

# Computer algebra independent integration tests

4-Trig-functions/4.2-Cosine/4.2.12-e-x^m-a+b-cos-c+d-x^n^p

Nasser M. Abbasi

May 18, 2020

Compiled on May 18, 2020 at 10:52am

## Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Listing of CAS systems tested . . . . .	3
1.2	Results . . . . .	3
1.3	Performance . . . . .	5
1.4	list of integrals that has no closed form antiderivative . . . . .	6
1.5	list of integrals solved by CAS but has no known antiderivative . . . . .	6
1.6	list of integrals solved by CAS but failed verification . . . . .	6
1.7	Timing . . . . .	7
1.8	Verification . . . . .	7
1.9	Important notes about some of the results . . . . .	7
1.10	Design of the test system . . . . .	8
<b>2</b>	<b>detailed summary tables of results</b>	<b>11</b>
2.1	List of integrals sorted by grade for each CAS . . . . .	11
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	12
2.3	Detailed conclusion table specific for Rubi results . . . . .	27
<b>3</b>	<b>Listing of integrals</b>	<b>31</b>
3.1	$\int x^3 \cos(a + bx^2) dx$ . . . . .	31
3.2	$\int x^2 \cos(a + bx^2) dx$ . . . . .	34
3.3	$\int x \cos(a + bx^2) dx$ . . . . .	37
3.4	$\int \cos(a + bx^2) dx$ . . . . .	40
3.5	$\int \frac{\cos(a+bx^2)}{x} dx$ . . . . .	43
3.6	$\int \frac{\cos(a+bx^2)}{x^2} dx$ . . . . .	46
3.7	$\int \frac{\cos(a+bx^2)}{x^3} dx$ . . . . .	49
3.8	$\int x^3 \cos^2(a + bx^2) dx$ . . . . .	52
3.9	$\int x^2 \cos^2(a + bx^2) dx$ . . . . .	55
3.10	$\int x \cos^2(a + bx^2) dx$ . . . . .	59
3.11	$\int \cos^2(a + bx^2) dx$ . . . . .	62
3.12	$\int \frac{\cos^2(a+bx^2)}{x} dx$ . . . . .	65
3.13	$\int \frac{\cos^2(a+bx^2)}{x^2} dx$ . . . . .	68
3.14	$\int \frac{\cos^2(a+bx^2)}{x^3} dx$ . . . . .	71

3.15	$\int x^3 \cos^3(a + bx^2) dx$	74
3.16	$\int x^2 \cos^3(a + bx^2) dx$	77
3.17	$\int x \cos^3(a + bx^2) dx$	81
3.18	$\int \cos^3(a + bx^2) dx$	84
3.19	$\int \frac{\cos^3(a+bx^2)}{x} dx$	88
3.20	$\int \frac{\cos^3(a+bx^2)}{x^2} dx$	91
3.21	$\int \frac{\cos^3(a+bx^2)}{x^3} dx$	95
3.22	$\int x \cos^7(a + bx^2) dx$	99
3.23	$\int x^{5/2} \cos(a + bx^2) dx$	102
3.24	$\int x^{3/2} \cos(a + bx^2) dx$	105
3.25	$\int \sqrt{x} \cos(a + bx^2) dx$	108
3.26	$\int \frac{\cos(a+bx^2)}{\sqrt{x}} dx$	111
3.27	$\int \frac{\cos(a+bx^2)}{x^{3/2}} dx$	114
3.28	$\int \frac{\cos(a+bx^2)}{x^{5/2}} dx$	117
3.29	$\int x^{5/2} \cos^2(a + bx^2) dx$	120
3.30	$\int x^{3/2} \cos^2(a + bx^2) dx$	123
3.31	$\int \sqrt{x} \cos^2(a + bx^2) dx$	126
3.32	$\int \frac{\cos^2(a+bx^2)}{\sqrt{x}} dx$	129
3.33	$\int \frac{\cos^2(a+bx^2)}{x^{3/2}} dx$	132
3.34	$\int \frac{\cos^2(a+bx^2)}{x^{5/2}} dx$	136
3.35	$\int \cos\left(a + \frac{b}{x}\right) dx$	140
3.36	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx$	143
3.37	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx$	146
3.38	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx$	149
3.39	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx$	152
3.40	$\int \cos\left(a + \frac{b}{x^2}\right) dx$	155
3.41	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$	158
3.42	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$	161
3.43	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx$	164
3.44	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$	167
3.45	$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx$	170
3.46	$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx$	173
3.47	$\int \cos(\sqrt{x}) dx$	175
3.48	$\int \cos^2(\sqrt{x}) dx$	178
3.49	$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx$	181
3.50	$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$	185
3.51	$\int \frac{\cos(a+b\sqrt[3]{x})}{\sqrt{x}} dx$	189
3.52	$\int \frac{\cos(a+b\sqrt[3]{x})}{x^{3/2}} dx$	193

3.53	$\int \frac{\cos(a+b\sqrt[3]{x})}{x^{5/2}} dx$	197
3.54	$\int \frac{\cos(a+b\sqrt[3]{x})}{x^{7/2}} dx$	201
3.55	$\int x^{3/2} \cos^2(a+b\sqrt[3]{x}) dx$	205
3.56	$\int \sqrt{x} \cos^2(a+b\sqrt[3]{x}) dx$	210
3.57	$\int \frac{\cos^2(a+b\sqrt[3]{x})}{\sqrt{x}} dx$	214
3.58	$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{3/2}} dx$	218
3.59	$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{5/2}} dx$	222
3.60	$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{7/2}} dx$	227
3.61	$\int \cos^3(\sqrt[3]{x}) dx$	232
3.62	$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx$	235
3.63	$\int (ex)^m (b \cos(c + dx^n))^p dx$	237
3.64	$\int (ex)^m (a + b \cos(c + dx^n))^p dx$	239
3.65	$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx$	241
3.66	$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$	244
3.67	$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx$	246
3.68	$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx$	249
3.69	$\int \frac{\cos(ax^n)}{x} dx$	251
3.70	$\int \frac{\cos^2(ax^n)}{x} dx$	254
3.71	$\int \frac{\cos^3(ax^n)}{x} dx$	257
3.72	$\int \frac{\cos^4(ax^n)}{x} dx$	260
3.73	$\int \cos(a + bx^n) dx$	263
3.74	$\int \cos^2(a + bx^n) dx$	266
3.75	$\int \cos^3(a + bx^n) dx$	269
3.76	$\int x^m \cos(a + bx^n) dx$	272
3.77	$\int x^m \cos^2(a + bx^n) dx$	275
3.78	$\int x^m \cos^3(a + bx^n) dx$	278
3.79	$\int x^{-1-n} \cos(a + bx^n) dx$	281
3.80	$\int x^{-1-n} \cos^2(a + bx^n) dx$	284
3.81	$\int x^{-1-n} \cos^3(a + bx^n) dx$	287
3.82	$\int x^{-1-2n} \cos(a + bx^n) dx$	290
3.83	$\int x^{-1-2n} \cos^2(a + bx^n) dx$	293
3.84	$\int x^{-1-2n} \cos^3(a + bx^n) dx$	296
3.85	$\int x^2 \cos((a + bx)^2) dx$	300
3.86	$\int x \cos((a + bx)^2) dx$	303
3.87	$\int \cos((a + bx)^2) dx$	306
3.88	$\int \frac{\cos((a+bx)^2)}{x} dx$	309
3.89	$\int \frac{\cos((a+bx)^2)}{x^2} dx$	311
3.90	$\int x^2 \cos(a + b\sqrt{c + dx}) dx$	313
3.91	$\int x \cos(a + b\sqrt{c + dx}) dx$	317
3.92	$\int \cos(a + b\sqrt{c + dx}) dx$	321
3.93	$\int \frac{\cos(a+b\sqrt{c+dx})}{x} dx$	324
3.94	$\int \frac{\cos(a+b\sqrt{c+dx})}{x^2} dx$	327
3.95	$\int x^2 \cos(a + b\sqrt[3]{c + dx}) dx$	331
3.96	$\int x \cos(a + b\sqrt[3]{c + dx}) dx$	337
3.97	$\int \cos(a + b\sqrt[3]{c + dx}) dx$	341
3.98	$\int \frac{\cos(a+b\sqrt[3]{c+dx})}{x} dx$	344

3.99  $\int \frac{\cos(a+b\sqrt[3]{c+dx})}{x^2} dx \dots\dots\dots 347$

**4 Listing of Grading functions 351**

# Chapter 1

## Introduction

This report gives the result of running the computer algebra independent integration problems. The listing of the problems are maintained by and can be downloaded from <https://rulebasedintegration.org>

The number of integrals in this report is [ 99 ]. This is test number [ 85 ].

### 1.1 Listing of CAS systems tested

The following systems were tested at this time.

1. Mathematica 12.1 (64 bit) on windows 10.
2. Rubi 4.16.1 in Mathematica 12 on windows 10.
3. Maple 2020 (64 bit) on windows 10.
4. Maxima 5.43 on Linux. (via sagemath 8.9)
5. Fricas 1.3.6 on Linux (via sagemath 9.0)
6. Sympy 1.5 under Python 3.7.3 using Anaconda distribution.
7. Giac/Xcas 1.5 on Linux. (via sagemath 8.9)

Maxima, Fricas and Giac/Xcas were called from inside SageMath. This was done using SageMath integrate command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly using Python.

### 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	solved	Failed
Rubi	% 100. ( 99 )	% 0. ( 0 )
Mathematica	% 100. ( 99 )	% 0. ( 0 )
Maple	% 87.88 ( 87 )	% 12.12 ( 12 )
Maxima	% 77.78 ( 77 )	% 22.22 ( 22 )
Fricas	% 91.92 ( 91 )	% 8.08 ( 8 )
Sympy	% 30.3 ( 30 )	% 69.7 ( 69 )
Giac	% 47.47 ( 47 )	% 52.53 ( 52 )


The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.

System	% A grade	% B grade	% C grade	% F grade
Rubi	100.	0.	0.	0.
Mathematica	93.94	0.	6.06	0.
Maple	69.7	5.05	13.13	12.12
Maxima	24.24	15.15	38.38	22.22
Fricas	87.88	0.	4.04	8.08
Sympy	25.25	5.05	0.	69.7
Giac	25.25	7.07	15.15	52.53

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



### 1.3 Performance

The table below summarizes the performance of each CAS system in terms of CPU time and leaf size of results.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.12	102.9	0.94	85.	1.
Mathematica	0.34	89.7	0.87	81.	0.95
Maple	0.11	144.4	1.17	64.	0.88
Maxima	1.45	273.45	2.51	338.	2.32
Fricas	1.64	224.51	2.42	180.	2.39
Sympy	2.41	95.4	1.32	53.5	1.13
Giac	1.03	174.81	1.49	78.	1.28

## 1.4 list of integrals that has no closed form antiderivative

{63, 64, 66, 68, 88, 89}

## 1.5 list of integrals solved by CAS but has no known antiderivative

**Rubi** {}

**Mathematica** {}

**Maple** {}

**Maxima** {}

**Fricas** {}

**Sympy** {}

**Giac** {}

## 1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

**Rubi** {}

**Mathematica** {67, 98}

**Maple** Verification phase not implemented yet.

**Maxima** Verification phase not implemented yet.

**Fricas** Verification phase not implemented yet.

**Sympy** Verification phase not implemented yet.

**Giac** Verification phase not implemented yet.



## 1.7 Timing

The command `AboluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of _int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call has completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out is not counted in the final statistics.

## 1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica. Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative produced was correct.

Verification phase has 3 minutes time out. An integral whose result was not verified could still be correct. Further investigation is needed on those integrals which failed verifications. Such integrals are marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.9 Important notes about some of the results

### 1.9.1 Important note about Maxima results

Since these integrals are run in a batch mode, using an automated script, and by using `sagemath` (SageMath uses Maxima), then any integral where Maxima needs an interactive response from the user to answer a question during evaluation of the integral in order to complete the integration, will fail and is counted as failed.

The exception raised is `ValueError`. Therefore Maxima result below is lower than what could result if Maxima was run directly and each question Maxima asks was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 30 such integrals out of total 705, or about 4 percent. This pecentage can be higher or lower depending on the specific input test file.

Such integrals can be indentified by looking at the output of the integration in each section for Maxima. If the output was an exception `ValueError` then this is most likely due to this reason.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath loading of Maxima `abs_integrate` was found to cause some problem. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

## 1.9.2 Important note about FriCAS and Giac/X-CAS results

There are Few integrals which failed due to SageMath not able to translate the result back to SageMath syntax and not because these CAS system were not able to do the integrations.

These will fail With error Exception raised: NotImplementedError

The number of such cases seems to be very small. About 1 or 2 percent of all integrals.

Hopefully the next version of SageMath will have complete translation of FriCAS and XCAS syntax and I will re-run all the tests again when this happens.

## 1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi and Maple, the builtin system function LeafSize is used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size is determined as follows.

For Fricas, Giac and Maxima (all called via sagemath) the following code is used

#see <https://stackoverflow.com/questions/25202346/how-to-obtain-leaf-count-expression-size-in->

```
def tree(expr):
    if expr.operator() is None:
        return expr
    else:
        return [expr.operator()+map(tree, expr.operands())
```

```
try:
    # 1.35 is a fudge factor since this estimate of leaf count is bit lower than
    #what it should be compared to Mathematica's
    leafCount = round(1.35*len(flatten(tree(anti))))
except Exception as ee:
    leafCount =1
```

For Sympy, called directly from Python, the following code is used

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

When these cas systems have a builtin function to find the leaf size of expressions, it will be used instead, and these tests run again.

## 1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. It contains 13 fields. This is description of each record (line)

1. integer, the problem number.
2. integer. 0 or 1 for failed or passed. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. The optimal antiderivative in CAS own syntax.

### High level overview of the CAS independent integration test build system

Nasser M. Abbasi  
June 22, 2018



# Chapter 2

## detailed summary tables of results

### 2.1 List of integrals sorted by grade for each CAS

#### 2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 }

B grade: { }

C grade: { }

F grade: { }

#### 2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 91, 92, 96, 97 }

B grade: { }

C grade: { 90, 93, 94, 95, 98, 99 }

F grade: { }

#### 2.1.3 Maple

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 20, 22, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 66, 68, 69, 70, 71, 72, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 91, 92, 97 }

B grade: { 90, 93, 94, 95, 96 }

C grade: { 14, 19, 21, 23, 24, 25, 26, 27, 28, 73, 76, 98, 99 }

F grade: { 29, 30, 31, 32, 33, 34, 65, 67, 74, 75, 77, 78 }

#### 2.1.4 Maxima

A grade: { 1, 3, 8, 10, 15, 17, 22, 37, 43, 45, 46, 47, 48, 61, 62, 63, 64, 66, 68, 88, 89, 91, 92, 97 }

B grade: { 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 90, 95, 96 }

C grade: { 2, 4, 5, 6, 7, 9, 11, 12, 13, 14, 16, 18, 19, 20, 21, 35, 36, 38, 39, 40, 41, 42, 44, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 85, 86, 87 }

F grade: { 65, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 93, 94, 98, 99 }

## 2.1.5 FriCAS

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 66, 68, 69, 70, 71, 72, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 95, 96, 97 }

B grade: { }

C grade: { 93, 94, 98, 99 }

F grade: { 65, 67, 73, 74, 75, 76, 77, 78 }

## 2.1.6 Sympy

A grade: { 1, 3, 4, 8, 10, 11, 15, 17, 18, 22, 36, 37, 38, 39, 43, 46, 47, 48, 63, 88, 89, 90, 91, 92, 97 }

B grade: { 2, 9, 16, 45, 61 }

C grade: { }

F grade: { 5, 6, 7, 12, 13, 14, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 40, 41, 42, 44, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 62, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 93, 94, 95, 96, 98, 99 }

## 2.1.7 Giac

A grade: { 1, 3, 5, 8, 10, 12, 15, 17, 19, 22, 37, 45, 46, 47, 48, 61, 62, 63, 64, 66, 68, 88, 89, 96, 97 }

B grade: { 7, 14, 21, 90, 91, 92, 95 }

C grade: { 2, 4, 9, 11, 16, 18, 49, 50, 51, 55, 56, 57, 85, 86, 87 }

F grade: { 6, 13, 20, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 38, 39, 40, 41, 42, 43, 44, 52, 53, 54, 58, 59, 60, 65, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 93, 94, 98, 99 }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column **normalized size** is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$

Problem 1	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	29	31	36	66	36	36
normalized size	1	1.	0.85	0.91	1.06	1.94	1.06	1.06
time (sec)	N/A	0.032	0.049	0.023	1.333	1.569	1.087	1.13

Problem 2	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	B	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	91	82	58	338	220	209	182
normalized size	1	1.	0.9	0.64	3.71	2.42	2.3	2.
time (sec)	N/A	0.069	0.145	0.025	2.102	1.643	2.13	1.152

Problem 3	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	18	30	19	18
normalized size	1	1.	1.	0.93	1.2	2.	1.27	1.2
time (sec)	N/A	0.014	0.003	0.023	1.354	1.571	0.208	1.159

Problem 4	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	A	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	57	44	308	185	61	128
normalized size	1	1.	0.81	0.63	4.4	2.64	0.87	1.83
time (sec)	N/A	0.021	0.09	0.025	2.25	1.593	0.565	1.162

Problem 5	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	24	22	58	122	0	28
normalized size	1	1.	0.96	0.88	2.32	4.88	0.	1.12
time (sec)	N/A	0.029	0.054	0.025	1.468	1.532	0.	1.129

Problem 6	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	80	81	57	355	209	0	0
normalized size	1	1.	1.01	0.71	4.44	2.61	0.	0.
time (sec)	N/A	0.045	0.202	0.024	1.739	1.687	0.	0.

Problem 7	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	42	39	65	178	0	117
normalized size	1	1.	1.	0.93	1.55	4.24	0.	2.79
time (sec)	N/A	0.089	0.068	0.024	1.545	1.593	0.	1.109

Problem 8	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	51	40	42	57	105	78	74
normalized size	1	1.	0.78	0.82	1.12	2.06	1.53	1.45
time (sec)	N/A	0.052	0.13	0.03	1.16	1.644	2.186	1.14

Problem 9	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	B	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	91	87	63	383	231	201	159
normalized size	1	1.	0.96	0.69	4.21	2.54	2.21	1.75
time (sec)	N/A	0.097	0.178	0.031	1.892	1.713	3.318	1.171

Problem 10	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	27	34	31	63	60	35
normalized size	1	1.	0.87	1.1	1.	2.03	1.94	1.13
time (sec)	N/A	0.029	0.045	0.026	1.178	1.618	0.518	1.116

Problem 11	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	A	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	67	45	338	163	56	111
normalized size	1	1.	0.96	0.64	4.83	2.33	0.8	1.59
time (sec)	N/A	0.044	0.061	0.035	1.913	1.633	1.047	1.162

Problem 12	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	34	32	69	153	0	47
normalized size	1	1.	0.92	0.86	1.86	4.14	0.	1.27
time (sec)	N/A	0.052	0.067	0.046	1.366	1.602	0.	1.168

Problem 13	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	76	62	369	180	0	0
normalized size	1	1.	1.	0.82	4.86	2.37	0.	0.
time (sec)	N/A	0.067	0.177	0.03	1.458	1.678	0.	0.

Problem 14	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	C	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	50	89	82	194	0	144
normalized size	1	1.	0.88	1.56	1.44	3.4	0.	2.53
time (sec)	N/A	0.118	0.128	0.138	1.266	1.645	0.	1.166



Problem 15	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	79	55	66	78	138	92	78
normalized size	1	1.	0.7	0.84	0.99	1.75	1.16	0.99
time (sec)	N/A	0.074	0.164	0.033	1.088	1.612	4.013	1.137

Problem 16	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	B	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	188	188	160	130	697	450	439	350
normalized size	1	1.	0.85	0.69	3.71	2.39	2.34	1.86
time (sec)	N/A	0.182	0.432	0.033	2.15	1.781	5.788	1.191

Problem 17	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	26	36	61	44	35
normalized size	1	1.	1.	0.79	1.09	1.85	1.33	1.06
time (sec)	N/A	0.028	0.017	0.029	1.175	1.565	0.97	1.162

Problem 18	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	A	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	153	153	116	101	653	373	129	250
normalized size	1	1.	0.76	0.66	4.27	2.44	0.84	1.63
time (sec)	N/A	0.089	0.237	0.033	2.22	1.623	1.564	1.166

Problem 19	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	C	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	50	125	120	261	0	63
normalized size	1	1.	0.91	2.27	2.18	4.75	0.	1.15
time (sec)	N/A	0.083	0.105	0.103	1.471	1.593	0.	1.167

Problem 20	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	168	166	128	721	412	0	0
normalized size	1	1.	0.99	0.76	4.29	2.45	0.	0.
time (sec)	N/A	0.144	0.644	0.033	1.39	1.828	0.	0.

Problem 21	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	C	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	91	90	162	132	340	0	250
normalized size	1	1.	0.99	1.78	1.45	3.74	0.	2.75
time (sec)	N/A	0.201	0.175	0.147	1.539	1.683	0.	1.178

Problem 22	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	54	50	74	123	94	70
normalized size	1	1.	0.81	0.75	1.1	1.84	1.4	1.04
time (sec)	N/A	0.054	0.088	0.028	1.178	1.702	17.455	1.274

Problem 23	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	B	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	111	111	113	229	397	176	0	0
normalized size	1	1.	1.02	2.06	3.58	1.59	0.	0.
time (sec)	N/A	0.081	0.201	0.142	1.474	1.702	0.	0.

Problem 24	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	111	111	111	290	390	170	0	0
normalized size	1	1.	1.	2.61	3.51	1.53	0.	0.
time (sec)	N/A	0.079	0.158	0.077	1.525	1.754	0.	0.

Problem 25	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	81	89	290	365	132	0	0
normalized size	1	1.	1.1	3.58	4.51	1.63	0.	0.
time (sec)	N/A	0.063	0.09	0.066	1.608	1.708	0.	0.

Problem 26	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	81	89	338	359	132	0	0
normalized size	1	1.	1.1	4.17	4.43	1.63	0.	0.
time (sec)	N/A	0.065	0.071	0.078	1.447	1.711	0.	0.

Problem 27	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	114	338	355	163	0	0
normalized size	1	1.	1.16	3.45	3.62	1.66	0.	0.
time (sec)	N/A	0.076	0.182	0.067	1.413	1.695	0.	0.

Problem 28	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	104	117	358	355	177	0	0
normalized size	1	1.	1.12	3.44	3.41	1.7	0.	0.
time (sec)	N/A	0.075	0.156	0.069	1.445	1.788	0.	0.

Problem 29	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	B	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	132	142	0	421	242	0	0
normalized size	1	1.	1.08	0.	3.19	1.83	0.	0.
time (sec)	N/A	0.175	0.458	0.094	1.495	1.81	0.	0.

Problem 30	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	132	142	0	424	236	0	0
normalized size	1	1.	1.08	0.	3.21	1.79	0.	0.
time (sec)	N/A	0.124	0.345	0.089	1.412	1.827	0.	0.

Problem 31	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	122	0	392	176	0	0
normalized size	1	1.	1.22	0.	3.92	1.76	0.	0.
time (sec)	N/A	0.13	0.226	0.085	1.422	1.744	0.	0.

Problem 32	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	96	96	120	0	393	170	0	0
normalized size	1	1.	1.25	0.	4.09	1.77	0.	0.
time (sec)	N/A	0.068	0.232	0.099	1.451	1.77	0.	0.

Problem 33	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	B	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	117	137	0	369	188	0	0
normalized size	1	1.	1.17	0.	3.15	1.61	0.	0.
time (sec)	N/A	0.163	0.34	0.089	1.347	1.766	0.	0.

Problem 34	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	B	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	116	137	0	373	196	0	0
normalized size	1	1.	1.18	0.	3.22	1.69	0.	0.
time (sec)	N/A	0.1	0.368	0.092	1.447	1.742	0.	0.

Problem 35	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	39	77	144	0	0
normalized size	1	1.	1.	1.26	2.48	4.65	0.	0.
time (sec)	N/A	0.072	0.031	0.037	1.287	1.595	0.	0.

Problem 36	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	21	58	109	15	0
normalized size	1	1.	1.	1.05	2.9	5.45	0.75	0.
time (sec)	N/A	0.027	0.047	0.033	1.421	1.573	1.915	0.

Problem 37	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	14	18	28	15	18
normalized size	1	1.	1.	1.08	1.38	2.15	1.15	1.38
time (sec)	N/A	0.014	0.003	0.026	1.055	1.511	1.759	1.133

Problem 38	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	29	42	69	70	31	0
normalized size	1	1.	0.97	1.4	2.3	2.33	1.03	0.
time (sec)	N/A	0.025	0.048	0.032	1.291	1.578	3.44	0.

Problem 39	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	46	92	68	95	46	0
normalized size	1	1.	1.	2.	1.48	2.07	1.	0.
time (sec)	N/A	0.047	0.004	0.033	1.368	1.515	6.182	0.

Problem 40	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	79	80	57	533	208	0	0
normalized size	1	1.	1.01	0.72	6.75	2.63	0.	0.
time (sec)	N/A	0.05	0.134	0.033	1.418	1.689	0.	0.

Problem 41	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	24	22	58	123	0	0
normalized size	1	1.	0.96	0.88	2.32	4.92	0.	0.
time (sec)	N/A	0.028	0.045	0.037	1.35	1.59	0.	0.

Problem 42	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	62	48	495	186	0	0
normalized size	1	1.	0.84	0.65	6.69	2.51	0.	0.
time (sec)	N/A	0.032	0.109	0.035	1.378	1.608	0.	0.

Problem 43	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	18	39	22	0
normalized size	1	1.	1.	0.93	1.2	2.6	1.47	0.
time (sec)	N/A	0.015	0.003	0.024	1.149	1.558	6.099	0.

Problem 44	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	88	64	359	235	0	0
normalized size	1	1.	0.91	0.66	3.7	2.42	0.	0.
time (sec)	N/A	0.053	0.156	0.036	1.332	1.67	0.	0.

Problem 45	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	B	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	18	14	16	50	39	16
normalized size	1	1.	0.95	0.74	0.84	2.63	2.05	0.84
time (sec)	N/A	0.021	0.029	0.032	1.14	1.644	0.357	1.112

Problem 46	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	8	22	7	8
normalized size	1	1.	1.	0.88	1.	2.75	0.88	1.
time (sec)	N/A	0.009	0.002	0.023	1.142	1.622	0.42	1.143

Problem 47	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	17	22	55	20	22
normalized size	1	1.	1.	0.77	1.	2.5	0.91	1.
time (sec)	N/A	0.011	0.015	0.027	1.161	1.637	0.41	1.113

Problem 48	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	31	34	31	86	51	31
normalized size	1	1.	0.86	0.94	0.86	2.39	1.42	0.86
time (sec)	N/A	0.021	0.028	0.026	1.077	1.605	0.387	1.117

Problem 49	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	235	235	165	196	459	462	0	325
normalized size	1	1.	0.7	0.83	1.95	1.97	0.	1.38
time (sec)	N/A	0.348	0.568	0.033	2.769	1.784	0.	1.154

Problem 50	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F(-1)	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	169	141	131	419	366	0	261
normalized size	1	1.	0.83	0.78	2.48	2.17	0.	1.54
time (sec)	N/A	0.196	0.363	0.03	2.435	1.763	0.	1.155

Problem 51	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	99	94	64	346	250	0	193
normalized size	1	1.	0.95	0.65	3.49	2.53	0.	1.95
time (sec)	N/A	0.113	0.153	0.032	1.952	1.923	0.	1.16

Problem 52	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	110	110	78	358	300	0	0
normalized size	1	1.	1.	0.71	3.25	2.73	0.	0.
time (sec)	N/A	0.143	0.248	0.03	1.86	2.022	0.	0.

Problem 53	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	184	184	180	129	359	405	0	0
normalized size	1	1.	0.98	0.7	1.95	2.2	0.	0.
time (sec)	N/A	0.238	0.238	0.03	1.503	2.076	0.	0.

Problem 54	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	250	250	238	180	362	490	0	0
normalized size	1	1.	0.95	0.72	1.45	1.96	0.	0.
time (sec)	N/A	0.337	0.316	0.032	1.436	2.131	0.	0.

Problem 55	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	310	310	174	219	497	591	0	302
normalized size	1	1.	0.56	0.71	1.6	1.91	0.	0.97
time (sec)	N/A	0.36	0.659	0.04	2.148	2.255	0.	1.186

Problem 56	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F(-1)	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	218	218	148	145	455	444	0	238
normalized size	1	1.	0.68	0.67	2.09	2.04	0.	1.09
time (sec)	N/A	0.251	0.442	0.037	1.968	2.085	0.	1.18

Problem 57	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	102	103	67	392	271	0	167
normalized size	1	1.	1.01	0.66	3.84	2.66	0.	1.64
time (sec)	N/A	0.17	0.189	0.039	2.965	2.083	0.	1.173

Problem 58	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	116	116	87	379	296	0	0
normalized size	1	1.	1.	0.75	3.27	2.55	0.	0.
time (sec)	N/A	0.187	0.254	0.037	1.744	2.092	0.	0.

Problem 59	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	228	185	146	381	455	0	0
normalized size	1	1.	0.81	0.64	1.67	2.	0.	0.
time (sec)	N/A	0.25	0.268	0.04	1.572	2.184	0.	0.

Problem 60	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	328	328	249	207	383	581	0	0
normalized size	1	1.	0.76	0.63	1.17	1.77	0.	0.
time (sec)	N/A	0.351	0.375	0.039	2.034	2.265	0.	0.

Problem 61	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	B	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	86	66	58	63	166	513	63
normalized size	1	1.	0.77	0.67	0.73	1.93	5.97	0.73
time (sec)	N/A	0.068	0.099	0.029	1.79	1.906	5.291	1.15

Problem 62	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	F(-1)	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	8	22	0	8
normalized size	1	1.	1.	0.88	1.	2.75	0.	1.
time (sec)	N/A	0.01	0.002	0.023	1.744	1.828	0.	1.147

Problem 63	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.02	0.987	1.156	0.	0.	0.	0.

Problem 64	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	F(-1)	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.022	1.261	1.145	0.	0.	0.	0.

Problem 65	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	93	93	89	0	0	0	0	0
normalized size	1	1.	0.96	0.	0.	0.	0.	0.
time (sec)	N/A	0.102	0.205	1.874	0.	0.	0.	0.

Problem 66	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	F(-1)	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	38	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.049	0.934	1.342	0.	0.	0.	0.

Problem 67	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F(-1)	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD
size	131	131	149	0	0	0	0	0
normalized size	1	1.	1.14	0.	0.	0.	0.	0.
time (sec)	N/A	0.167	0.441	1.907	0.	0.	0.	0.

Problem 68	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	F(-1)	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	40	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.054	1.158	1.388	0.	0.	0.	0.

Problem 69	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	24	25	0	131	0	0
normalized size	1	1.	0.92	0.96	0.	5.04	0.	0.
time (sec)	N/A	0.034	0.065	0.034	0.	1.895	0.	0.

Problem 70	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	37	45	0	165	0	0
normalized size	1	1.	0.86	1.05	0.	3.84	0.	0.
time (sec)	N/A	0.058	0.089	0.036	0.	1.862	0.	0.



Problem 71	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	53	52	0	274	0	0
normalized size	1	1.	0.79	0.78	0.	4.09	0.	0.
time (sec)	N/A	0.086	0.121	0.036	0.	2.006	0.	0.

Problem 72	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	79	66	77	0	309	0	0
normalized size	1	1.	0.84	0.97	0.	3.91	0.	0.
time (sec)	N/A	0.094	0.141	0.039	0.	2.025	0.	0.

Problem 73	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	83	92	75	0	0	0	0
normalized size	1	1.	1.11	0.9	0.	0.	0.	0.
time (sec)	N/A	0.022	0.09	0.092	0.	0.	0.	0.

Problem 74	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	102	94	0	0	0	0	0
normalized size	1	1.	0.92	0.	0.	0.	0.	0.
time (sec)	N/A	0.069	0.247	0.165	0.	0.	0.	0.

Problem 75	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	179	179	173	0	0	0	0	0
normalized size	1	1.	0.97	0.	0.	0.	0.	0.
time (sec)	N/A	0.079	0.24	180.	0.	0.	0.	0.

Problem 76	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	C	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	105	115	111	0	0	0	0
normalized size	1	1.	1.1	1.06	0.	0.	0.	0.
time (sec)	N/A	0.075	0.2	0.162	0.	0.	0.	0.

Problem 77	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	141	129	0	0	0	0	0
normalized size	1	1.	0.91	0.	0.	0.	0.	0.
time (sec)	N/A	0.16	0.516	0.126	0.	0.	0.	0.

Problem 78	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	229	229	221	0	0	0	0	0
normalized size	1	1.	0.97	0.	0.	0.	0.	0.
time (sec)	N/A	0.212	0.562	0.178	0.	0.	0.	0.

Problem 79	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	45	45	0	190	0	0
normalized size	1	1.	0.96	0.96	0.	4.04	0.	0.
time (sec)	N/A	0.086	0.094	0.036	0.	1.667	0.	0.

Problem 80	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	53	65	0	209	0	0
normalized size	1	1.	0.77	0.94	0.	3.03	0.	0.
time (sec)	N/A	0.121	0.187	0.044	0.	1.736	0.	0.

Problem 81	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	113	95	101	0	366	0	0
normalized size	1	1.	0.84	0.89	0.	3.24	0.	0.
time (sec)	N/A	0.2	0.235	0.05	0.	1.851	0.	0.

Problem 82	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	70	65	0	254	0	0
normalized size	1	1.	0.9	0.83	0.	3.26	0.	0.
time (sec)	N/A	0.111	0.13	0.037	0.	1.669	0.	0.

Problem 83	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-1)	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	82	89	0	290	0	0
normalized size	1	1.	0.86	0.94	0.	3.05	0.	0.
time (sec)	N/A	0.147	0.212	0.043	0.	2.123	0.	0.

Problem 84	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F	A	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	165	165	141	144	0	481	0	0
normalized size	1	1.	0.85	0.87	0.	2.92	0.	0.
time (sec)	N/A	0.253	0.356	0.05	0.	2.055	0.	0.

Problem 85	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	99	76	131	346	278	0	215
normalized size	1	1.	0.77	1.32	3.49	2.81	0.	2.17
time (sec)	N/A	0.068	0.265	0.029	2.781	1.667	0.	1.13

Problem 86	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	42	63	266	158	0	161
normalized size	1	1.	0.89	1.34	5.66	3.36	0.	3.43
time (sec)	N/A	0.032	0.047	0.028	2.371	1.641	0.	1.142

Problem 87	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	C	A	F	C
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	36	113	107	0	74
normalized size	1	1.	1.	1.24	3.9	3.69	0.	2.55
time (sec)	N/A	0.006	0.008	0.028	1.763	1.542	0.	1.09

Problem 88	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	14	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.008	2.662	0.124	0.	0.	0.	0.

Problem 89	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	14	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.009	5.288	0.116	0.	0.	0.	0.

Problem 90	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	346	346	224	825	907	248	274	1134
normalized size	1	1.	0.65	2.38	2.62	0.72	0.79	3.28
time (sec)	N/A	0.305	0.841	0.089	1.242	1.626	2.361	1.394

Problem 91	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	167	167	71	299	355	166	151	479
normalized size	1	1.	0.43	1.79	2.13	0.99	0.9	2.87
time (sec)	N/A	0.136	0.276	0.031	1.254	1.631	0.684	1.249

Problem 92	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	48	61	81	109	66	227
normalized size	1	1.	0.89	1.13	1.5	2.02	1.22	4.2
time (sec)	N/A	0.027	0.073	0.033	1.154	1.575	0.536	1.182

Problem 93	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	B	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	145	271	0	360	0	0
normalized size	1	1.	1.15	2.15	0.	2.86	0.	0.
time (sec)	N/A	0.239	0.669	0.058	0.	1.725	0.	0.

Problem 94	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	B	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	184	184	240	714	0	489	0	0
normalized size	1	1.	1.3	3.88	0.	2.66	0.	0.
time (sec)	N/A	0.349	1.225	0.181	0.	1.823	0.	0.

Problem 95	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	B	B	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	537	537	382	1809	1821	464	0	1490
normalized size	1	1.	0.71	3.37	3.39	0.86	0.	2.77
time (sec)	N/A	0.511	1.107	0.102	1.479	1.704	0.	1.613

Problem 96	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	B	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	261	117	655	706	293	0	500
normalized size	1	1.	0.45	2.51	2.7	1.12	0.	1.92
time (sec)	N/A	0.231	0.417	0.035	1.254	1.638	0.	1.238

Problem 97	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	A	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	65	131	159	155	94	109
normalized size	1	1.	0.76	1.54	1.87	1.82	1.11	1.28
time (sec)	N/A	0.055	0.111	0.034	1.118	1.625	1.678	1.263

Problem 98	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	C	F	C	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD
size	234	234	243	279	0	833	0	0
normalized size	1	1.	1.04	1.19	0.	3.56	0.	0.
time (sec)	N/A	0.468	0.433	0.149	0.	1.771	0.	0.

Problem 99	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	C	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	332	332	138	933	0	1137	0	0
normalized size	1	1.	0.42	2.81	0.	3.42	0.	0.
time (sec)	N/A	0.741	0.621	0.32	0.	2.06	0.	0.

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [35] had the largest ratio of [ 0.625 ]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	1.	12	0.25
2	A	4	4	1.	12	0.333
3	A	2	2	1.	10	0.2
4	A	3	3	1.	8	0.375
5	A	3	3	1.	12	0.25
6	A	4	4	1.	12	0.333
7	A	5	5	1.	12	0.417
8	A	3	3	1.	14	0.214
9	A	6	5	1.	14	0.357
10	A	3	3	1.	12	0.25
11	A	5	4	1.	10	0.4
12	A	5	4	1.	14	0.286
13	A	6	6	1.	14	0.429
14	A	7	6	1.	14	0.429
15	A	4	4	1.	14	0.286
16	A	10	5	1.	14	0.357
17	A	3	2	1.	12	0.167
18	A	8	4	1.	10	0.4
19	A	8	4	1.	14	0.286
20	A	9	5	1.	14	0.357
21	A	12	6	1.	14	0.429
22	A	3	2	1.	12	0.167
23	A	4	3	1.	14	0.214
24	A	4	3	1.	14	0.214
25	A	3	2	1.	14	0.143
26	A	3	2	1.	14	0.143
27	A	4	3	1.	14	0.214

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
28	A	4	3	1.	14	0.214
29	A	7	5	1.	16	0.312
30	A	7	5	1.	16	0.312
31	A	6	4	1.	16	0.25
32	A	6	4	1.	16	0.25
33	A	7	5	1.	16	0.312
34	A	7	6	1.	16	0.375
35	A	5	5	1.	8	0.625
36	A	3	3	1.	12	0.25
37	A	2	2	1.	12	0.167
38	A	3	3	1.	12	0.25
39	A	4	3	1.	12	0.25
40	A	5	5	1.	8	0.625
41	A	3	3	1.	12	0.25
42	A	4	4	1.	12	0.333
43	A	2	2	1.	12	0.167
44	A	5	5	1.	12	0.417
45	A	3	3	1.	14	0.214
46	A	2	2	1.	12	0.167
47	A	3	3	1.	6	0.5
48	A	3	3	1.	8	0.375
49	A	13	7	1.	16	0.438
50	A	10	7	1.	16	0.438
51	A	7	7	1.	16	0.438
52	A	8	7	1.	16	0.438
53	A	11	7	1.	16	0.438
54	A	14	7	1.	16	0.438
55	A	15	10	1.	18	0.556
56	A	12	9	1.	18	0.5
57	A	9	8	1.	18	0.444
58	A	10	9	1.	18	0.5
59	A	12	10	1.	18	0.556
60	A	16	9	1.	18	0.5
61	A	7	5	1.	8	0.625
62	A	2	2	1.	12	0.167
63	A	0	0	0.	0	0.
64	A	0	0	0.	0	0.
65	A	3	3	1.	20	0.15
66	A	0	0	0.	0	0.
67	A	5	5	1.	22	0.227
68	A	0	0	0.	0	0.
69	A	3	3	1.	12	0.25
70	A	5	4	1.	14	0.286
71	A	8	4	1.	14	0.286

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
72	A	8	4	1.	14	0.286
73	A	3	2	1.	8	0.25
74	A	5	3	1.	10	0.3
75	A	8	3	1.	10	0.3
76	A	3	2	1.	12	0.167
77	A	5	3	1.	14	0.214
78	A	8	3	1.	14	0.214
79	A	5	5	1.	16	0.312
80	A	7	6	1.	18	0.333
81	A	12	6	1.	18	0.333
82	A	6	5	1.	16	0.312
83	A	8	6	1.	18	0.333
84	A	14	6	1.	18	0.333
85	A	7	6	1.	12	0.5
86	A	5	4	1.	10	0.4
87	A	1	1	1.	8	0.125
88	A	0	0	0.	0	0.
89	A	0	0	0.	0	0.
90	A	14	3	1.	18	0.167
91	A	8	3	1.	16	0.188
92	A	3	3	1.	14	0.214
93	A	8	4	1.	18	0.222
94	A	10	6	1.	18	0.333
95	A	20	4	1.	18	0.222
96	A	11	4	1.	16	0.25
97	A	4	3	1.	14	0.214
98	A	11	4	1.	18	0.222
99	A	13	6	1.	18	0.333





# Chapter 3

## Listing of integrals

### 3.1 $\int x^3 \cos(a + bx^2) dx$

**Optimal.** Leaf size=34

$$\frac{\cos(a + bx^2)}{2b^2} + \frac{x^2 \sin(a + bx^2)}{2b}$$

```
[Out] Cos[a + b*x^2]/(2*b^2) + (x^2*Sin[a + b*x^2])/(2*b)
```

---

**Rubi [A]** time = 0.0317191, antiderivative size = 34, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3380, 3296, 2638}

$$\frac{\cos(a + bx^2)}{2b^2} + \frac{x^2 \sin(a + bx^2)}{2b}$$

Antiderivative was successfully verified.

```
[In] Int[x^3*Cos[a + b*x^2],x]
```

```
[Out] Cos[a + b*x^2]/(2*b^2) + (x^2*Sin[a + b*x^2])/(2*b)
```

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
:> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p,
x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
&& (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 3296

```
Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol]
:> -Simp[(c + d*x)^m*Cos[e + f*x]/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[e + f*x],
x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

#### Rule 2638

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol]
:> -Simp[Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int x^3 \cos(a + bx^2) dx &= \frac{1}{2} \text{Subst} \left( \int x \cos(a + bx) dx, x, x^2 \right) \\
&= \frac{x^2 \sin(a + bx^2)}{2b} - \frac{\text{Subst} \left( \int \sin(a + bx) dx, x, x^2 \right)}{2b} \\
&= \frac{\cos(a + bx^2)}{2b^2} + \frac{x^2 \sin(a + bx^2)}{2b}
\end{aligned}$$

**Mathematica [A]** time = 0.0492611, size = 29, normalized size = 0.85

$$\frac{bx^2 \sin(a + bx^2) + \cos(a + bx^2)}{2b^2}$$

Antiderivative was successfully verified.

[In] Integrate[x^3\*Cos[a + b\*x^2],x]

[Out] (Cos[a + b\*x^2] + b\*x^2\*Sin[a + b\*x^2])/(2\*b^2)

**Maple [A]** time = 0.023, size = 31, normalized size = 0.9

$$\frac{\cos(bx^2 + a)}{2b^2} + \frac{x^2 \sin(bx^2 + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3\*cos(b\*x^2+a),x)

[Out] 1/2\*cos(b\*x^2+a)/b^2+1/2\*x^2\*sin(b\*x^2+a)/b

**Maxima [A]** time = 1.33306, size = 36, normalized size = 1.06

$$\frac{bx^2 \sin(bx^2 + a) + \cos(bx^2 + a)}{2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3\*cos(b\*x^2+a),x, algorithm="maxima")

[Out] 1/2\*(b\*x^2\*sin(b\*x^2 + a) + cos(b\*x^2 + a))/b^2

**Fricas [A]** time = 1.56881, size = 66, normalized size = 1.94

$$\frac{bx^2 \sin(bx^2 + a) + \cos(bx^2 + a)}{2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3*cos(b*x^2+a),x, algorithm="fricas")
```

```
[Out] 1/2*(b*x^2*sin(b*x^2 + a) + cos(b*x^2 + a))/b^2
```

**Sympy [A]** time = 1.08675, size = 36, normalized size = 1.06

$$\begin{cases} \frac{x^2 \sin(a+bx^2)}{2b} + \frac{\cos(a+bx^2)}{2b^2} & \text{for } b \neq 0 \\ \frac{x^4 \cos(a)}{4} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**3*cos(b*x**2+a),x)
```

```
[Out] Piecewise((x**2*sin(a + b*x**2)/(2*b) + cos(a + b*x**2)/(2*b**2), Ne(b, 0))
, (x**4*cos(a)/4, True))
```

**Giac [A]** time = 1.12968, size = 36, normalized size = 1.06

$$\frac{bx^2 \sin(bx^2 + a) + \cos(bx^2 + a)}{2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3*cos(b*x^2+a),x, algorithm="giac")
```

```
[Out] 1/2*(b*x^2*sin(b*x^2 + a) + cos(b*x^2 + a))/b^2
```

## 3.2 $\int x^2 \cos(a + bx^2) dx$

**Optimal.** Leaf size=91

$$-\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right)}{2b^{3/2}} - \frac{\sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{2b^{3/2}} + \frac{x \sin(a + bx^2)}{2b}$$

[Out]  $-(\operatorname{Sqrt}[\pi/2] \operatorname{Cos}[a] \operatorname{FresnelS}[\operatorname{Sqrt}[b] \operatorname{Sqrt}[2/\pi] x]) / (2b^{3/2}) - (\operatorname{Sqrt}[\pi/2] \operatorname{FresnelC}[\operatorname{Sqrt}[b] \operatorname{Sqrt}[2/\pi] x] \operatorname{Sin}[a]) / (2b^{3/2}) + (x \operatorname{Sin}[a + bx^2]) / (2b)$

**Rubi [A]** time = 0.0694239, antiderivative size = 91, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3386, 3353, 3352, 3351}

$$-\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right)}{2b^{3/2}} - \frac{\sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{2b^{3/2}} + \frac{x \sin(a + bx^2)}{2b}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[x^2 \operatorname{Cos}[a + bx^2], x]$

[Out]  $-(\operatorname{Sqrt}[\pi/2] \operatorname{Cos}[a] \operatorname{FresnelS}[\operatorname{Sqrt}[b] \operatorname{Sqrt}[2/\pi] x]) / (2b^{3/2}) - (\operatorname{Sqrt}[\pi/2] \operatorname{FresnelC}[\operatorname{Sqrt}[b] \operatorname{Sqrt}[2/\pi] x] \operatorname{Sin}[a]) / (2b^{3/2}) + (x \operatorname{Sin}[a + bx^2]) / (2b)$

### Rule 3386

$\operatorname{Int}[\operatorname{Cos}[(c_.) + (d_.) (x_.)^{(n_)}] * ((e_.) (x_.)^{(m_)}), x\_Symbol] \rightarrow \operatorname{Simp}[(e^{(n-1)} (e^x)^{(m-n+1)} \operatorname{Sin}[c + d x^n]) / (d^n), x] - \operatorname{Dist}[(e^n (m-n+1)) / (d^n), \operatorname{Int}[(e^x)^{(m-n)} \operatorname{Sin}[c + d x^n], x], x] /;$   $\operatorname{FreeQ}\{c, d, e\}, x$  &&  $\operatorname{IGtQ}[n, 0]$  &&  $\operatorname{LtQ}[n, m + 1]$

### Rule 3353

$\operatorname{Int}[\operatorname{Sin}[(c_.) + (d_.) * ((e_.) + (f_.) (x_.)^2)], x\_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Sin}[c], \operatorname{Int}[\operatorname{Cos}[d * (e + f x)^2], x], x] + \operatorname{Dist}[\operatorname{Cos}[c], \operatorname{Int}[\operatorname{Sin}[d * (e + f x)^2], x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x$

### Rule 3352

$\operatorname{Int}[\operatorname{Cos}[(d_.) * ((e_.) + (f_.) (x_.)^2)], x\_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[\pi/2] \operatorname{FresnelC}[\operatorname{Sqrt}[2/\pi] \operatorname{Rt}[d, 2] * (e + f x)]) / (f \operatorname{Rt}[d, 2]), x] /;$   $\operatorname{FreeQ}\{d, e, f\}, x$

### Rule 3351

$\operatorname{Int}[\operatorname{Sin}[(d_.) * ((e_.) + (f_.) (x_.)^2)], x\_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[\pi/2] \operatorname{FresnelS}[\operatorname{Sqrt}[2/\pi] \operatorname{Rt}[d, 2] * (e + f x)]) / (f \operatorname{Rt}[d, 2]), x] /;$   $\operatorname{FreeQ}\{d, e, f\}, x$

### Rubi steps

$$\begin{aligned}
\int x^2 \cos(a + bx^2) dx &= \frac{x \sin(a + bx^2)}{2b} - \frac{\int \sin(a + bx^2) dx}{2b} \\
&= \frac{x \sin(a + bx^2)}{2b} - \frac{\cos(a) \int \sin(bx^2) dx}{2b} - \frac{\sin(a) \int \cos(bx^2) dx}{2b} \\
&= -\frac{\sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{2b^{3/2}} - \frac{\sqrt{\frac{\pi}{2}} C\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right) \sin(a)}{2b^{3/2}} + \frac{x \sin(a + bx^2)}{2b}
\end{aligned}$$

**Mathematica [A]** time = 0.144657, size = 82, normalized size = 0.9

$$\frac{-\sqrt{2\pi} \sin(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right) - \sqrt{2\pi} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right) + 2\sqrt{bx} \sin(a + bx^2)}{4b^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[x^2\*Cos[a + b\*x^2],x]

[Out]  $(-\text{Sqrt}[2*\text{Pi}]*\text{Cos}[a]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x]) - \text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x]*\text{Sin}[a] + 2*\text{Sqrt}[b]*x*\text{Sin}[a + b*x^2])/(4*b^(3/2))$

**Maple [A]** time = 0.025, size = 58, normalized size = 0.6

$$\frac{x \sin(bx^2 + a)}{2b} - \frac{\sqrt{2}\sqrt{\pi}}{4} \left( \cos(a) \text{FresnelS}\left(\frac{x\sqrt{2}}{\sqrt{\pi}} \sqrt{b}\right) + \sin(a) \text{FresnelC}\left(\frac{x\sqrt{2}}{\sqrt{\pi}} \sqrt{b}\right) \right) b^{-\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2\*cos(b\*x^2+a),x)

[Out]  $1/2*x*\sin(b*x^2+a)/b-1/4/b^(3/2)*2^(1/2)*\text{Pi}^(1/2)*(\cos(a)*\text{FresnelS}(x*b^(1/2)*2^(1/2)/\text{Pi}^(1/2))+\sin(a)*\text{FresnelC}(x*b^(1/2)*2^(1/2)/\text{Pi}^(1/2)))$

**Maxima [C]** time = 2.1017, size = 338, normalized size = 3.71

$$8x|b|\sin(bx^2 + a) + \sqrt{\pi} \left( \left( -i \cos\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) - i \cos\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) - \sin\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(b\*x^2+a),x, algorithm="maxima")

[Out]  $1/16*(8*x*\text{abs}(b)*\sin(b*x^2 + a) + \text{sqrt}(\text{pi})*((( -I*\cos(1/4*\text{pi} + 1/2*\arctan2(0, b)) - I*\cos(-1/4*\text{pi} + 1/2*\arctan2(0, b)) - \sin(1/4*\text{pi} + 1/2*\arctan2(0, b)) + \sin(-1/4*\text{pi} + 1/2*\arctan2(0, b))))*\cos(a) - (\cos(1/4*\text{pi} + 1/2*\arctan2(0, b)) + \cos(-1/4*\text{pi} + 1/2*\arctan2(0, b)) - I*\sin(1/4*\text{pi} + 1/2*\arctan2(0, b)) + I*\sin(-1/4*\text{pi} + 1/2*\arctan2(0, b))))*\sin(a)*\text{erf}(\text{sqrt}(I*b)*x) + ((I*\cos(1/4*\text{pi} + 1/2*\arctan2(0, b)) + I*\cos(-1/4*\text{pi} + 1/2*\arctan2(0, b)) - \sin(1/4*\text{pi} + 1/2*\arctan2(0, b)) + \sin(-1/4*\text{pi} + 1/2*\arctan2(0, b))))*\cos(a) - (\cos(1/4*\text{pi} + 1/2*\arctan2(0, b)) + \cos(-1/4*\text{pi} + 1/2*\arctan2(0, b)) + I*\sin(1/4*\text{pi}$

+ 1/2\*arctan2(0, b)) - I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))\*sin(a))\*erf(sqrt(-I\*b\*x))\*sqrt(abs(b)))/(b\*abs(b))

**Fricas [A]** time = 1.64268, size = 220, normalized size = 2.42

$$\frac{\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) + \sqrt{2}\pi\sqrt{\frac{b}{\pi}}C\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)\sin(a) - 2bx\sin(bx^2 + a)}{4b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(b\*x^2+a),x, algorithm="fricas")

[Out] -1/4\*(sqrt(2)\*pi\*sqrt(b/pi)\*cos(a)\*fresnel\_sin(sqrt(2)\*x\*sqrt(b/pi)) + sqrt(2)\*pi\*sqrt(b/pi)\*fresnel\_cos(sqrt(2)\*x\*sqrt(b/pi))\*sin(a) - 2\*b\*x\*sin(b\*x^2 + a))/b^2

**Sympy [B]** time = 2.13041, size = 209, normalized size = 2.3

$$\frac{b^{\frac{3}{2}}x^5\sqrt{\frac{1}{b}}\sin(a)\Gamma\left(\frac{3}{4}\right)\Gamma\left(\frac{5}{4}\right){}_2F_3\left(\frac{3}{4}, \frac{5}{4} \middle| \frac{3}{2}, \frac{7}{4}, \frac{9}{4} \right) - \sqrt{b}x^3\sqrt{\frac{1}{b}}\cos(a)\Gamma\left(\frac{1}{4}\right)\Gamma\left(\frac{3}{4}\right){}_2F_3\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{1}{2}, \frac{5}{4}, \frac{7}{4} \right) - \sqrt{2}\sqrt{\pi}x^2\sqrt{\frac{1}{b}}\sin(a)S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)}{8\Gamma\left(\frac{7}{4}\right)\Gamma\left(\frac{9}{4}\right) - 8\Gamma\left(\frac{5}{4}\right)\Gamma\left(\frac{7}{4}\right) - 2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*2\*cos(b\*x\*\*2+a),x)

[Out] b\*\*(3/2)\*x\*\*5\*sqrt(1/b)\*sin(a)\*gamma(3/4)\*gamma(5/4)\*hyper((3/4, 5/4), (3/2, 7/4, 9/4), -b\*\*2\*x\*\*4/4)/(8\*gamma(7/4)\*gamma(9/4)) - sqrt(b)\*x\*\*3\*sqrt(1/b)\*cos(a)\*gamma(1/4)\*gamma(3/4)\*hyper((1/4, 3/4), (1/2, 5/4, 7/4), -b\*\*2\*x\*\*4/4)/(8\*gamma(5/4)\*gamma(7/4)) - sqrt(2)\*sqrt(pi)\*x\*\*2\*sqrt(1/b)\*sin(a)\*fresnel\_s(sqrt(2)\*sqrt(b)\*x/sqrt(pi))/2 + sqrt(2)\*sqrt(pi)\*x\*\*2\*sqrt(1/b)\*cos(a)\*fresnel\_c(sqrt(2)\*sqrt(b)\*x/sqrt(pi))/2

**Giac [C]** time = 1.15215, size = 182, normalized size = 2.

$$\frac{ixe^{(ibx^2+ia)}}{4b} + \frac{ixe^{(-ibx^2-ia)}}{4b} - \frac{i\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{(ia)}}{8b\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}} + \frac{i\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{(-ia)}}{8b\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(b\*x^2+a),x, algorithm="giac")

[Out] -1/4\*I\*x\*e^(I\*b\*x^2 + I\*a)/b + 1/4\*I\*x\*e^(-I\*b\*x^2 - I\*a)/b - 1/8\*I\*sqrt(2)\*sqrt(pi)\*erf(-1/2\*sqrt(2)\*x\*(-I\*b/abs(b) + 1)\*sqrt(abs(b)))\*e^(I\*a)/(b\*(-I\*b/abs(b) + 1)\*sqrt(abs(b))) + 1/8\*I\*sqrt(2)\*sqrt(pi)\*erf(-1/2\*sqrt(2)\*x\*(I\*b/abs(b) + 1)\*sqrt(abs(b)))\*e^(-I\*a)/(b\*(I\*b/abs(b) + 1)\*sqrt(abs(b)))

### 3.3 $\int x \cos(a + bx^2) dx$

**Optimal.** Leaf size=15

$$\frac{\sin(a + bx^2)}{2b}$$

[Out] Sin[a + b\*x^2]/(2\*b)

**Rubi [A]** time = 0.0143389, antiderivative size = 15, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.2$ , Rules used = {3380, 2637}

$$\frac{\sin(a + bx^2)}{2b}$$

Antiderivative was successfully verified.

[In] Int[x\*Cos[a + b\*x^2],x]

[Out] Sin[a + b\*x^2]/(2\*b)

#### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

#### Rule 2637

Int[sin[Pi/2 + (c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Simp[Sin[c + d\*x]/d, x] /; FreeQ[{c, d}, x]

#### Rubi steps

$$\begin{aligned} \int x \cos(a + bx^2) dx &= \frac{1}{2} \text{Subst} \left( \int \cos(a + bx) dx, x, x^2 \right) \\ &= \frac{\sin(a + bx^2)}{2b} \end{aligned}$$

**Mathematica [A]** time = 0.0027367, size = 15, normalized size = 1.

$$\frac{\sin(a + bx^2)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[x\*Cos[a + b\*x^2],x]

[Out] Sin[a + b\*x^2]/(2\*b)

---

**Maple [A]** time = 0.023, size = 14, normalized size = 0.9

$$\frac{\sin(bx^2 + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x\*cos(b\*x^2+a),x)

[Out] 1/2\*sin(b\*x^2+a)/b

---

**Maxima [A]** time = 1.35392, size = 18, normalized size = 1.2

$$\frac{\sin(bx^2 + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x^2+a),x, algorithm="maxima")

[Out] 1/2\*sin(b\*x^2 + a)/b

---

**Fricas [A]** time = 1.57076, size = 30, normalized size = 2.

$$\frac{\sin(bx^2 + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x^2+a),x, algorithm="fricas")

[Out] 1/2\*sin(b\*x^2 + a)/b

---

**Sympy [A]** time = 0.208326, size = 19, normalized size = 1.27

$$\begin{cases} \frac{\sin(a+bx^2)}{2b} & \text{for } b \neq 0 \\ \frac{x^2 \cos(a)}{2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x\*\*2+a),x)

[Out] Piecewise((sin(a + b\*x\*\*2)/(2\*b), Ne(b, 0)), (x\*\*2\*cos(a)/2, True))

---

**Giac [A]** time = 1.15867, size = 18, normalized size = 1.2

$$\frac{\sin(bx^2 + a)}{2b}$$



Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x*cos(b*x^2+a),x, algorithm="giac")
```

```
[Out] 1/2*sin(b*x^2 + a)/b
```

### 3.4 $\int \cos(a + bx^2) dx$

**Optimal.** Leaf size=70

$$\frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right)}{\sqrt{b}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{\sqrt{b}}$$

[Out] (Sqrt [Pi/2]\*Cos [a]\*FresnelC[Sqrt [b]\*Sqrt [2/Pi]\*x])/Sqrt [b] - (Sqrt [Pi/2]\*FresnelS[Sqrt [b]\*Sqrt [2/Pi]\*x]\*Sin[a])/Sqrt [b]

**Rubi [A]** time = 0.0214534, antiderivative size = 70, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3354, 3352, 3351}

$$\frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right)}{\sqrt{b}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2], x]

[Out] (Sqrt [Pi/2]\*Cos [a]\*FresnelC[Sqrt [b]\*Sqrt [2/Pi]\*x])/Sqrt [b] - (Sqrt [Pi/2]\*FresnelS[Sqrt [b]\*Sqrt [2/Pi]\*x]\*Sin[a])/Sqrt [b]

#### Rule 3354

Int[Cos[(c\_) + (d\_)\*((e\_) + (f\_)\*(x\_))<sup>2</sup>], x\_Symbol] := Dist[Cos[c], Int[Cos[d\*(e + f\*x)<sup>2</sup>], x], x] - Dist[Sin[c], Int[Sin[d\*(e + f\*x)<sup>2</sup>], x], x] /; FreeQ[{c, d, e, f}, x]

#### Rule 3352

Int[Cos[(d\_)\*((e\_) + (f\_)\*(x\_))<sup>2</sup>], x\_Symbol] := Simp[(Sqrt [Pi/2]\*FresnelC[Sqrt [2/Pi]\*Rt[d, 2]\*(e + f\*x)])/ (f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rule 3351

Int[Sin[(d\_)\*((e\_) + (f\_)\*(x\_))<sup>2</sup>], x\_Symbol] := Simp[(Sqrt [Pi/2]\*FresnelS[Sqrt [2/Pi]\*Rt[d, 2]\*(e + f\*x)])/ (f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rubi steps

$$\begin{aligned} \int \cos(a + bx^2) dx &= \cos(a) \int \cos(bx^2) dx - \sin(a) \int \sin(bx^2) dx \\ &= \frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{C}\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{\sqrt{b}} - \frac{\sqrt{\frac{\pi}{2}} S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right) \sin(a)}{\sqrt{b}} \end{aligned}$$

**Mathematica [A]** time = 0.090349, size = 57, normalized size = 0.81

$$\frac{\sqrt{\frac{\pi}{2}} \left( \cos(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right) - \sin(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right) \right)}{\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2], x]

[Out] (Sqrt[Pi/2]\*(Cos[a]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x] - FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x]\*Sin[a]))/Sqrt[b]

**Maple [A]** time = 0.025, size = 44, normalized size = 0.6

$$\frac{\sqrt{2}\sqrt{\pi}}{2} \left( \cos(a) \operatorname{FresnelC} \left( \frac{x\sqrt{2}}{\sqrt{\pi}} \sqrt{b} \right) - \sin(a) \operatorname{FresnelS} \left( \frac{x\sqrt{2}}{\sqrt{\pi}} \sqrt{b} \right) \right) \frac{1}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a), x)

[Out] 1/2\*2^(1/2)\*Pi^(1/2)/b^(1/2)\*(cos(a)\*FresnelC(x\*b^(1/2)\*2^(1/2)/Pi^(1/2))-sin(a)\*FresnelS(x\*b^(1/2)\*2^(1/2)/Pi^(1/2)))

**Maxima [C]** time = 2.25014, size = 308, normalized size = 4.4

$$\sqrt{\pi} \left( \left( \cos \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + \cos \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) - i \sin \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + i \sin \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a), x, algorithm="maxima")

[Out] 1/8\*sqrt(pi)\*(((cos(1/4\*pi + 1/2\*arctan2(0, b)) + cos(-1/4\*pi + 1/2\*arctan2(0, b)) - I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(a) - (I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + sin(1/4\*pi + 1/2\*arctan2(0, b)) - sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(a)) \* erf(sqrt(I\*b)\*x) + (((cos(1/4\*pi + 1/2\*arctan2(0, b)) + cos(-1/4\*pi + 1/2\*arctan2(0, b)) + I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(a) - (-I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + sin(1/4\*pi + 1/2\*arctan2(0, b)) - sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(a)) \* erf(sqrt(-I\*b)\*x)) / sqrt(abs(b))

**Fricas [A]** time = 1.59267, size = 185, normalized size = 2.64

$$\frac{\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)\operatorname{C}\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)-\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\operatorname{S}\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)\sin(a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a), x, algorithm="fricas")

[Out] 1/2\*(sqrt(2)\*pi\*sqrt(b/pi)\*cos(a)\*fresnel\_cos(sqrt(2)\*x\*sqrt(b/pi)) - sqrt(2)\*pi\*sqrt(b/pi)\*fresnel\_sin(sqrt(2)\*x\*sqrt(b/pi))\*sin(a))/b

---

**Sympy [A]** time = 0.564621, size = 61, normalized size = 0.87

$$\frac{\sqrt{2}\sqrt{\pi}\left(-\sin(a)S\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right)+\cos(a)C\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right)\right)\sqrt{\frac{1}{b}}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a),x)

[Out] sqrt(2)\*sqrt(pi)\*(-sin(a)\*fresnels(sqrt(2)\*sqrt(b)\*x/sqrt(pi)) + cos(a)\*fresnelc(sqrt(2)\*sqrt(b)\*x/sqrt(pi)))\*sqrt(1/b)/2

---

**Giac [C]** time = 1.16209, size = 128, normalized size = 1.83

$$-\frac{\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{(ia)}}{4\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}}-\frac{\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{(-ia)}}{4\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a),x, algorithm="giac")

[Out] -1/4\*sqrt(2)\*sqrt(pi)\*erf(-1/2\*sqrt(2)\*x\*(-I\*b/abs(b) + 1)\*sqrt(abs(b)))\*e^(I\*a)/((-I\*b/abs(b) + 1)\*sqrt(abs(b))) - 1/4\*sqrt(2)\*sqrt(pi)\*erf(-1/2\*sqrt(2)\*x\*(I\*b/abs(b) + 1)\*sqrt(abs(b)))\*e^(-I\*a)/((I\*b/abs(b) + 1)\*sqrt(abs(b)))

$$3.5 \quad \int \frac{\cos(a+bx^2)}{x} dx$$

**Optimal.** Leaf size=25

$$\frac{1}{2} \cos(a) \operatorname{CosIntegral}(bx^2) - \frac{1}{2} \sin(a) \operatorname{Si}(bx^2)$$

[Out] (Cos[a]\*CosIntegral[b\*x^2])/2 - (Sin[a]\*SinIntegral[b\*x^2])/2

**Rubi [A]** time = 0.0289689, antiderivative size = 25, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3378, 3376, 3375}

$$\frac{1}{2} \cos(a) \operatorname{CosIntegral}(bx^2) - \frac{1}{2} \sin(a) \operatorname{Si}(bx^2)$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]/x,x]

[Out] (Cos[a]\*CosIntegral[b\*x^2])/2 - (Sin[a]\*SinIntegral[b\*x^2])/2

**Rule 3378**

Int[Cos[(c\_) + (d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Dist[Cos[c], Int[Cos[d\*x^n]/x, x], x] - Dist[Sin[c], Int[Sin[d\*x^n]/x, x], x] /; FreeQ[{c, d, n}, x]

**Rule 3376**

Int[Cos[(d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[CosIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

**Rule 3375**

Int[Sin[(d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[SinIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

**Rubi steps**

$$\begin{aligned} \int \frac{\cos(a+bx^2)}{x} dx &= \cos(a) \int \frac{\cos(bx^2)}{x} dx - \sin(a) \int \frac{\sin(bx^2)}{x} dx \\ &= \frac{1}{2} \cos(a) \operatorname{Ci}(bx^2) - \frac{1}{2} \sin(a) \operatorname{Si}(bx^2) \end{aligned}$$

**Mathematica [A]** time = 0.0544486, size = 24, normalized size = 0.96

$$\frac{1}{2} (\cos(a) \operatorname{CosIntegral}(bx^2) - \sin(a) \operatorname{Si}(bx^2))$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]/x,x]

[Out]  $(\text{Cos}[a] * \text{CosIntegral}[b*x^2] - \text{Sin}[a] * \text{SinIntegral}[b*x^2])/2$

---

**Maple [A]** time = 0.025, size = 22, normalized size = 0.9

$$\frac{\text{Ci}(bx^2) \cos(a)}{2} - \frac{\text{Si}(bx^2) \sin(a)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x^2+a)/x,x)`

[Out]  $1/2 * \text{Ci}(b*x^2) * \cos(a) - 1/2 * \text{Si}(b*x^2) * \sin(a)$

---

**Maxima [C]** time = 1.46817, size = 58, normalized size = 2.32

$$\frac{1}{4} (\text{Ei}(ibx^2) + \text{Ei}(-ibx^2)) \cos(a) + \frac{1}{4} (i \text{Ei}(ibx^2) - i \text{Ei}(-ibx^2)) \sin(a)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x^2+a)/x,x, algorithm="maxima")`

[Out]  $1/4 * (\text{Ei}(I*b*x^2) + \text{Ei}(-I*b*x^2)) * \cos(a) + 1/4 * (I*\text{Ei}(I*b*x^2) - I*\text{Ei}(-I*b*x^2)) * \sin(a)$

---

**Fricas [A]** time = 1.53231, size = 122, normalized size = 4.88

$$\frac{1}{4} (\text{Ci}(bx^2) + \text{Ci}(-bx^2)) \cos(a) - \frac{1}{2} \sin(a) \text{Si}(bx^2)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x^2+a)/x,x, algorithm="fricas")`

[Out]  $1/4 * (\cos\_integral(b*x^2) + \cos\_integral(-b*x^2)) * \cos(a) - 1/2 * \sin(a) * \sin\_integral(b*x^2)$

---

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + bx^2)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x**2+a)/x,x)`

[Out] `Integral(cos(a + b*x**2)/x, x)`

---

**Giac [A]** time = 1.1287, size = 28, normalized size = 1.12

$$\frac{1}{2} \cos(a) \operatorname{Ci}(bx^2) - \frac{1}{2} \sin(a) \operatorname{Si}(bx^2)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x^2+a)/x,x, algorithm="giac")
```

```
[Out] 1/2*cos(a)*cos_integral(b*x^2) - 1/2*sin(a)*sin_integral(b*x^2)
```

### 3.6 $\int \frac{\cos(a+bx^2)}{x^2} dx$

**Optimal.** Leaf size=80

$$-\sqrt{2\pi}\sqrt{b}\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{bx}\right) - \sqrt{2\pi}\sqrt{b}\cos(a)\text{S}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \frac{\cos(a+bx^2)}{x}$$

[Out] -(Cos[a + b\*x^2]/x) - Sqrt[b]\*Sqrt[2\*Pi]\*Cos[a]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x] - Sqrt[b]\*Sqrt[2\*Pi]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x]\*Sin[a]

**Rubi [A]** time = 0.0445045, antiderivative size = 80, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3388, 3353, 3352, 3351}

$$-\sqrt{2\pi}\sqrt{b}\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{bx}\right) - \sqrt{2\pi}\sqrt{b}\cos(a)\text{S}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \frac{\cos(a+bx^2)}{x}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]/x^2,x]

[Out] -(Cos[a + b\*x^2]/x) - Sqrt[b]\*Sqrt[2\*Pi]\*Cos[a]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x] - Sqrt[b]\*Sqrt[2\*Pi]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x]\*Sin[a]

#### Rule 3388

```
Int[Cos[(c_.) + (d_.)*(x_)^(n_.)]*((e_.)*(x_))^(m_), x_Symbol] := Simp[((e*x)^(m + 1)*Cos[c + d*x^n])/(e^(m + 1)), x] + Dist[(d*n)/(e^n*(m + 1)), Int[(e*x)^(m + n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[m, -1]
```

#### Rule 3353

```
Int[Sin[(c_) + (d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Dist[Sin[c], Int[Cos[d*(e + f*x)^2], x], x] + Dist[Cos[c], Int[Sin[d*(e + f*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]
```

#### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

#### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

#### Rubi steps



$$\begin{aligned} \int \frac{\cos(a + bx^2)}{x^2} dx &= -\frac{\cos(a + bx^2)}{x} - (2b) \int \sin(a + bx^2) dx \\ &= -\frac{\cos(a + bx^2)}{x} - (2b \cos(a)) \int \sin(bx^2) dx - (2b \sin(a)) \int \cos(bx^2) dx \\ &= -\frac{\cos(a + bx^2)}{x} - \sqrt{b}\sqrt{2\pi} \cos(a) S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \sqrt{b}\sqrt{2\pi} C\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a) \end{aligned}$$

**Mathematica [A]** time = 0.201662, size = 81, normalized size = 1.01

$$-\sqrt{2\pi}\sqrt{b}\left(\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{bx}\right) + \cos(a)S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)\right) + \frac{\sin(a)\sin(bx^2)}{x} - \frac{\cos(a)\cos(bx^2)}{x}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]/x^2,x]

[Out] -((Cos[a]\*Cos[b\*x^2])/x) - Sqrt[b]\*Sqrt[2\*Pi]\*(Cos[a]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x] + FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x]\*Sin[a]) + (Sin[a]\*Sin[b\*x^2])/x

**Maple [A]** time = 0.024, size = 57, normalized size = 0.7

$$-\frac{\cos(bx^2 + a)}{x} - \sqrt{b}\sqrt{2}\sqrt{\pi}\left(\cos(a)\text{FresnelS}\left(\frac{x\sqrt{2}}{\sqrt{\pi}}\sqrt{b}\right) + \sin(a)\text{FresnelC}\left(\frac{x\sqrt{2}}{\sqrt{\pi}}\sqrt{b}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)/x^2,x)

[Out] -cos(b\*x^2+a)/x-b^(1/2)\*2^(1/2)\*Pi^(1/2)\*(cos(a)\*FresnelS(x\*b^(1/2)\*2^(1/2)/Pi^(1/2))+sin(a)\*FresnelC(x\*b^(1/2)\*2^(1/2)/Pi^(1/2)))

**Maxima [C]** time = 1.73907, size = 355, normalized size = 4.44

$$\sqrt{x^2|b|\left(\left(\Gamma\left(-\frac{1}{2}, ibx^2\right) + \Gamma\left(-\frac{1}{2}, -ibx^2\right)\right)\cos\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right) + \left(\Gamma\left(-\frac{1}{2}, ibx^2\right) + \Gamma\left(-\frac{1}{2}, -ibx^2\right)\right)\cos\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right) + \left(\Gamma\left(-\frac{1}{2}, ibx^2\right) - \Gamma\left(-\frac{1}{2}, -ibx^2\right)\right)\sin\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right) + \left(-\Gamma\left(-\frac{1}{2}, ibx^2\right) + \Gamma\left(-\frac{1}{2}, -ibx^2\right)\right)\sin\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right)\right)\cos(a) + \left(-\Gamma\left(-\frac{1}{2}, ibx^2\right) + \Gamma\left(-\frac{1}{2}, -ibx^2\right)\right)\cos\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right) + \left(-\Gamma\left(-\frac{1}{2}, ibx^2\right) + \Gamma\left(-\frac{1}{2}, -ibx^2\right)\right)\cos\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right) + \left(\Gamma\left(-\frac{1}{2}, ibx^2\right) + \Gamma\left(-\frac{1}{2}, -ibx^2\right)\right)\sin\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right) - \left(\Gamma\left(-\frac{1}{2}, ibx^2\right) + \Gamma\left(-\frac{1}{2}, -ibx^2\right)\right)\sin\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^2,x, algorithm="maxima")

[Out] -1/8\*sqrt(x^2\*abs(b))\*(((gamma(-1/2, I\*b\*x^2) + gamma(-1/2, -I\*b\*x^2))\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + (gamma(-1/2, I\*b\*x^2) + gamma(-1/2, -I\*b\*x^2))\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + (I\*gamma(-1/2, I\*b\*x^2) - I\*gamma(-1/2, -I\*b\*x^2))\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + (-I\*gamma(-1/2, I\*b\*x^2) + I\*gamma(-1/2, -I\*b\*x^2))\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*cos(a) + ((-I\*gamma(-1/2, I\*b\*x^2) + I\*gamma(-1/2, -I\*b\*x^2))\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + (-I\*gamma(-1/2, I\*b\*x^2) + I\*gamma(-1/2, -I\*b\*x^2))\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + (gamma(-1/2, I\*b\*x^2) + gamma(-1/2, -I\*b\*x^2))\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - (gamma(-1/2, I\*b\*x^2) + gamma(-1/2, -I\*b\*x^2))\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))

$/4\pi + 1/2\arctan2(0, b))\sin(a))/x$

---

**Fricas [A]** time = 1.68656, size = 209, normalized size = 2.61

$$\frac{\sqrt{2}\pi x \sqrt{\frac{b}{\pi}} \cos(a) S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) + \sqrt{2}\pi x \sqrt{\frac{b}{\pi}} C\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) \sin(a) + \cos(bx^2 + a)}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^2,x, algorithm="fricas")

[Out]  $-(\sqrt{2}\pi x \sqrt{b/\pi} \cos(a) \operatorname{fresnel\_sin}(\sqrt{2}x\sqrt{b/\pi}) + \sqrt{2}\pi x \sqrt{b/\pi} \operatorname{fresnel\_cos}(\sqrt{2}x\sqrt{b/\pi}) \sin(a) + \cos(bx^2 + a))/x$

---

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + bx^2)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)/x\*\*2,x)

[Out] Integral(cos(a + b\*x\*\*2)/x\*\*2, x)

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^2 + a)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^2,x, algorithm="giac")

[Out] integrate(cos(b\*x^2 + a)/x^2, x)

### 3.7 $\int \frac{\cos(a+bx^2)}{x^3} dx$

**Optimal.** Leaf size=42

$$-\frac{1}{2}b \sin(a) \operatorname{CosIntegral}(bx^2) - \frac{1}{2}b \cos(a) \operatorname{Si}(bx^2) - \frac{\cos(a+bx^2)}{2x^2}$$

[Out]  $-\operatorname{Cos}[a + b*x^2]/(2*x^2) - (b*\operatorname{CosIntegral}[b*x^2]*\operatorname{Sin}[a])/2 - (b*\operatorname{Cos}[a]*\operatorname{SinIntegral}[b*x^2])/2$

**Rubi [A]** time = 0.0892469, antiderivative size = 42, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3380, 3297, 3303, 3299, 3302}

$$-\frac{1}{2}b \sin(a) \operatorname{CosIntegral}(bx^2) - \frac{1}{2}b \cos(a) \operatorname{Si}(bx^2) - \frac{\cos(a+bx^2)}{2x^2}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[\operatorname{Cos}[a + b*x^2]/x^3, x]$

[Out]  $-\operatorname{Cos}[a + b*x^2]/(2*x^2) - (b*\operatorname{CosIntegral}[b*x^2]*\operatorname{Sin}[a])/2 - (b*\operatorname{Cos}[a]*\operatorname{SinIntegral}[b*x^2])/2$

#### Rule 3380

$\operatorname{Int}[(a + \operatorname{Cos}(c + d*x) + (d*x)^n) * (b*x)^p * (x^m), x\_Symbol] \rightarrow \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n] - 1) * (a + b*\operatorname{Cos}[c + d*x])^p}, x], x, x^n], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, m, n, p\}, x \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]] \&\& (\operatorname{EqQ}[p, 1] \mid \mid \operatorname{EqQ}[m, n-1] \mid \mid (\operatorname{IntegerQ}[p] \&\& \operatorname{GtQ}[\operatorname{Simplify}[(m+1)/n], 0]))$

#### Rule 3297

$\operatorname{Int}[(c + d*x + (d*x)^m) * \sin(e + f*x), x\_Symbol] \rightarrow \operatorname{Simp}[(c + d*x)^{m+1} * \operatorname{Sin}[e + f*x] / (d*(m+1)), x] - \operatorname{Dist}[f / (d*(m+1)), \operatorname{Int}[(c + d*x)^{m+1} * \operatorname{Cos}[e + f*x], x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \&\& \operatorname{LtQ}[m, -1]$

#### Rule 3303

$\operatorname{Int}[\sin(e + f*x) / (c + d*x), x\_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Cos}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Sin}[(c*f)/d + f*x] / (c + d*x), x], x] + \operatorname{Dist}[\operatorname{Sin}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Cos}[(c*f)/d + f*x] / (c + d*x), x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \&\& \operatorname{NeQ}[d*e - c*f, 0]$

#### Rule 3299

$\operatorname{Int}[\sin(e + f*x) / (c + d*x), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{SinIntegral}[e + f*x] / d, x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \&\& \operatorname{EqQ}[d*e - c*f, 0]$

#### Rule 3302

$\operatorname{Int}[\sin(e + f*x) / (c + d*x), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{CosIntegral}[e - \operatorname{Pi}/2 + f*x] / d, x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \&\& \operatorname{EqQ}[d*(e - \operatorname{Pi}/2) -$

c\*f, 0]

### Rubi steps

$$\begin{aligned} \int \frac{\cos(a + bx^2)}{x^3} dx &= \frac{1}{2} \text{Subst} \left( \int \frac{\cos(a + bx)}{x^2} dx, x, x^2 \right) \\ &= -\frac{\cos(a + bx^2)}{2x^2} - \frac{1}{2}b \text{Subst} \left( \int \frac{\sin(a + bx)}{x} dx, x, x^2 \right) \\ &= -\frac{\cos(a + bx^2)}{2x^2} - \frac{1}{2}(b \cos(a)) \text{Subst} \left( \int \frac{\sin(bx)}{x} dx, x, x^2 \right) - \frac{1}{2}(b \sin(a)) \text{Subst} \left( \int \frac{\cos(bx)}{x} dx, x, x^2 \right) \\ &= -\frac{\cos(a + bx^2)}{2x^2} - \frac{1}{2}b \text{Ci}(bx^2) \sin(a) - \frac{1}{2}b \cos(a) \text{Si}(bx^2) \end{aligned}$$

**Mathematica [A]** time = 0.0684203, size = 42, normalized size = 1.

$$-\frac{bx^2 \sin(a) \text{CosIntegral}(bx^2) + bx^2 \cos(a) \text{Si}(bx^2) + \cos(a + bx^2)}{2x^2}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]/x^3, x]

[Out] -(Cos[a + b\*x^2] + b\*x^2\*CosIntegral[b\*x^2]\*Sin[a] + b\*x^2\*Cos[a]\*SinIntegral[b\*x^2])/(2\*x^2)

**Maple [A]** time = 0.024, size = 39, normalized size = 0.9

$$-\frac{\cos(bx^2 + a)}{2x^2} - b \left( \frac{\cos(a) \text{Si}(bx^2)}{2} + \frac{\text{Ci}(bx^2) \sin(a)}{2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)/x^3, x)

[Out] -1/2\*cos(b\*x^2+a)/x^2 - b\*(1/2\*cos(a)\*Si(b\*x^2) + 1/2\*Ci(b\*x^2)\*sin(a))

**Maxima [C]** time = 1.54482, size = 65, normalized size = 1.55

$$-\frac{1}{4} \left( (i\Gamma(-1, ibx^2) - i\Gamma(-1, -ibx^2)) \cos(a) + (\Gamma(-1, ibx^2) + \Gamma(-1, -ibx^2)) \sin(a) \right) b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^3, x, algorithm="maxima")

[Out] -1/4\*((I\*gamma(-1, I\*b\*x^2) - I\*gamma(-1, -I\*b\*x^2))\*cos(a) + (gamma(-1, I\*b\*x^2) + gamma(-1, -I\*b\*x^2))\*sin(a))\*b

**Fricas [A]** time = 1.59318, size = 178, normalized size = 4.24

$$\frac{2bx^2 \cos(a) \operatorname{Si}(bx^2) + (bx^2 \operatorname{Ci}(bx^2) + bx^2 \operatorname{Ci}(-bx^2)) \sin(a) + 2 \cos(bx^2 + a)}{4x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^3,x, algorithm="fricas")

[Out] -1/4\*(2\*b\*x^2\*cos(a)\*sin\_integral(b\*x^2) + (b\*x^2\*cos\_integral(b\*x^2) + b\*x^2\*cos\_integral(-b\*x^2))\*sin(a) + 2\*cos(b\*x^2 + a))/x^2

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + bx^2)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)/x\*\*3,x)

[Out] Integral(cos(a + b\*x\*\*2)/x\*\*3, x)

**Giac [B]** time = 1.10915, size = 117, normalized size = 2.79

$$\frac{(bx^2 + a)b^2 \operatorname{Ci}(bx^2) \sin(a) - ab^2 \operatorname{Ci}(bx^2) \sin(a) + (bx^2 + a)b^2 \cos(a) \operatorname{Si}(bx^2) - ab^2 \cos(a) \operatorname{Si}(bx^2) + b^2 \cos(bx^2 + a)}{2b^2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^3,x, algorithm="giac")

[Out] -1/2\*((b\*x^2 + a)\*b^2\*cos\_integral(b\*x^2)\*sin(a) - a\*b^2\*cos\_integral(b\*x^2)\*sin(a) + (b\*x^2 + a)\*b^2\*cos(a)\*sin\_integral(b\*x^2) - a\*b^2\*cos(a)\*sin\_integral(b\*x^2) + b^2\*cos(b\*x^2 + a))/(b^2\*x^2)

### 3.8 $\int x^3 \cos^2(a + bx^2) dx$

**Optimal.** Leaf size=51

$$\frac{\cos^2(a + bx^2)}{8b^2} + \frac{x^2 \sin(a + bx^2) \cos(a + bx^2)}{4b} + \frac{x^4}{8}$$

[Out]  $x^4/8 + \text{Cos}[a + b*x^2]^2/(8*b^2) + (x^2*\text{Cos}[a + b*x^2]*\text{Sin}[a + b*x^2])/(4*b)$

**Rubi [A]** time = 0.0519296, antiderivative size = 51, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3380, 3310, 30}

$$\frac{\cos^2(a + bx^2)}{8b^2} + \frac{x^2 \sin(a + bx^2) \cos(a + bx^2)}{4b} + \frac{x^4}{8}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[x^3*\text{Cos}[a + b*x^2]^2,x]$

[Out]  $x^4/8 + \text{Cos}[a + b*x^2]^2/(8*b^2) + (x^2*\text{Cos}[a + b*x^2]*\text{Sin}[a + b*x^2])/(4*b)$

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
  := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p,
    x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
  && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 3310

```
Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
  := Simp[(d*(b*Sine[e + f*x])^n)/(f^2*n^2), x] + (Dist[(b^2*(n - 1))/n, Int[(c + d*x)*
    (b*Sine[e + f*x])^(n - 2), x], x] - Simp[(b*(c + d*x)*Cos[e + f*x]*(b*Sine[e + f*x])^(n - 1))/
    (f*n), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]
```

#### Rule 30

```
Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]
```

#### Rubi steps

$$\begin{aligned} \int x^3 \cos^2(a + bx^2) dx &= \frac{1}{2} \text{Subst} \left( \int x \cos^2(a + bx) dx, x, x^2 \right) \\ &= \frac{\cos^2(a + bx^2)}{8b^2} + \frac{x^2 \cos(a + bx^2) \sin(a + bx^2)}{4b} + \frac{1}{4} \text{Subst} \left( \int x dx, x, x^2 \right) \\ &= \frac{x^4}{8} + \frac{\cos^2(a + bx^2)}{8b^2} + \frac{x^2 \cos(a + bx^2) \sin(a + bx^2)}{4b} \end{aligned}$$

**Mathematica [A]** time = 0.129541, size = 40, normalized size = 0.78

$$\frac{2bx^2 \left( \sin(2(a + bx^2)) + bx^2 \right) + \cos(2(a + bx^2))}{16b^2}$$

Antiderivative was successfully verified.

[In] Integrate[x^3\*Cos[a + b\*x^2]^2,x]

[Out] (Cos[2\*(a + b\*x^2)] + 2\*b\*x^2\*(b\*x^2 + Sin[2\*(a + b\*x^2)]))/(16\*b^2)

**Maple [A]** time = 0.03, size = 42, normalized size = 0.8

$$\frac{x^4}{8} + \frac{x^2 \sin(2bx^2 + 2a)}{8b} + \frac{\cos(2bx^2 + 2a)}{16b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3\*cos(b\*x^2+a)^2,x)

[Out] 1/8\*x^4+1/8/b\*x^2\*sin(2\*b\*x^2+2\*a)+1/16/b^2\*cos(2\*b\*x^2+2\*a)

**Maxima [A]** time = 1.1601, size = 57, normalized size = 1.12

$$\frac{2b^2x^4 + 2bx^2 \sin(2bx^2 + 2a) + \cos(2bx^2 + 2a)}{16b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3\*cos(b\*x^2+a)^2,x, algorithm="maxima")

[Out] 1/16\*(2\*b^2\*x^4 + 2\*b\*x^2\*sin(2\*b\*x^2 + 2\*a) + cos(2\*b\*x^2 + 2\*a))/b^2

**Fricas [A]** time = 1.64402, size = 105, normalized size = 2.06

$$\frac{b^2x^4 + 2bx^2 \cos(bx^2 + a) \sin(bx^2 + a) + \cos(bx^2 + a)^2}{8b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3\*cos(b\*x^2+a)^2,x, algorithm="fricas")

[Out] 1/8\*(b^2\*x^4 + 2\*b\*x^2\*cos(b\*x^2 + a)\*sin(b\*x^2 + a) + cos(b\*x^2 + a)^2)/b^2

**Sympy [A]** time = 2.18639, size = 78, normalized size = 1.53

$$\begin{cases} \frac{x^4 \sin^2(a+bx^2)}{8} + \frac{x^4 \cos^2(a+bx^2)}{8} + \frac{x^2 \sin(a+bx^2) \cos(a+bx^2)}{4b} - \frac{\sin^2(a+bx^2)}{8b^2} & \text{for } b \neq 0 \\ \frac{x^4 \cos^2(a)}{4} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*3\*cos(b\*x\*\*2+a)\*\*2,x)

[Out] Piecewise((x\*\*4\*sin(a + b\*x\*\*2)\*\*2/8 + x\*\*4\*cos(a + b\*x\*\*2)\*\*2/8 + x\*\*2\*sin(a + b\*x\*\*2)\*cos(a + b\*x\*\*2)/(4\*b) - sin(a + b\*x\*\*2)\*\*2/(8\*b\*\*2), Ne(b, 0)), (x\*\*4\*cos(a)\*\*2/4, True))

**Giac [A]** time = 1.14041, size = 74, normalized size = 1.45

$$\frac{2bx^2 \sin(2bx^2 + 2a) + 2(bx^2 + a)^2 - 4(bx^2 + a)a + \cos(2bx^2 + 2a)}{16b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3\*cos(b\*x^2+a)^2,x, algorithm="giac")

[Out] 1/16\*(2\*b\*x^2\*sin(2\*b\*x^2 + 2\*a) + 2\*(b\*x^2 + a)^2 - 4\*(b\*x^2 + a)\*a + cos(2\*b\*x^2 + 2\*a))/b^2



### 3.9 $\int x^2 \cos^2(a + bx^2) dx$

**Optimal.** Leaf size=91

$$-\frac{\sqrt{\pi} \sin(2a) \text{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{16b^{3/2}} - \frac{\sqrt{\pi} \cos(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{16b^{3/2}} + \frac{x \sin(2a + 2bx^2)}{8b} + \frac{x^3}{6}$$

```
[Out] x^3/6 - (Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x)/Sqrt[Pi]])/(16*b^(3/2)) -
(Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x)/Sqrt[Pi]]*Sin[2*a])/(16*b^(3/2)) + (x*Sin
[2*a + 2*b*x^2])/(8*b)
```

**Rubi [A]** time = 0.0973785, antiderivative size = 91, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 5, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {3404, 3386, 3353, 3352, 3351}

$$-\frac{\sqrt{\pi} \sin(2a) \text{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{16b^{3/2}} - \frac{\sqrt{\pi} \cos(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{16b^{3/2}} + \frac{x \sin(2a + 2bx^2)}{8b} + \frac{x^3}{6}$$

Antiderivative was successfully verified.

```
[In] Int[x^2*Cos[a + b*x^2]^2,x]
```

```
[Out] x^3/6 - (Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x)/Sqrt[Pi]])/(16*b^(3/2)) -
(Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x)/Sqrt[Pi]]*Sin[2*a])/(16*b^(3/2)) + (x*Sin
[2*a + 2*b*x^2])/(8*b)
```

#### Rule 3404

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_)*((e_.)*(x_)^(m_.), x
_Symbol] :> Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x]
/; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]
```

#### Rule 3386

```
Int[Cos[(c_.) + (d_.)*(x_)^(n_)])*((e_.)*(x_)^(m_.), x_Symbol] :> Simp[(e^(
n - 1)*(e*x)^(m - n + 1)*Sin[c + d*x^n])/(d*n), x] - Dist[(e^n*(m - n + 1))
/(d*n), Int[(e*x)^(m - n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] &&
IGtQ[n, 0] && LtQ[n, m + 1]
```

#### Rule 3353

```
Int[Sin[(c_) + (d_.)*((e_.) + (f_.)*(x_)^2)], x_Symbol] :> Dist[Sin[c], Int
[Cos[d*(e + f*x)^2], x], x] + Dist[Cos[c], Int[Sin[d*(e + f*x)^2], x], x] /
; FreeQ[{c, d, e, f}, x]
```

#### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_)^2)], x_Symbol] :> Simp[(Sqrt[Pi/2]*Fresne
lC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

#### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_)^2)], x_Symbol] :> Simp[(Sqrt[Pi/2]*Fresne
lS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

Rubi steps

$$\begin{aligned}
\int x^2 \cos^2(a + bx^2) dx &= \int \left( \frac{x^2}{2} + \frac{1}{2} x^2 \cos(2a + 2bx^2) \right) dx \\
&= \frac{x^3}{6} + \frac{1}{2} \int x^2 \cos(2a + 2bx^2) dx \\
&= \frac{x^3}{6} + \frac{x \sin(2a + 2bx^2)}{8b} - \frac{\int \sin(2a + 2bx^2) dx}{8b} \\
&= \frac{x^3}{6} + \frac{x \sin(2a + 2bx^2)}{8b} - \frac{\cos(2a) \int \sin(2bx^2) dx}{8b} - \frac{\sin(2a) \int \cos(2bx^2) dx}{8b} \\
&= \frac{x^3}{6} - \frac{\sqrt{\pi} \cos(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{16b^{3/2}} - \frac{\sqrt{\pi} C\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \sin(2a)}{16b^{3/2}} + \frac{x \sin(2a + 2bx^2)}{8b}
\end{aligned}$$

**Mathematica [A]** time = 0.178047, size = 87, normalized size = 0.96

$$\frac{-3\sqrt{\pi} \sin(2a) \text{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - 3\sqrt{\pi} \cos(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) + 2\sqrt{bx} (3 \sin(2(a + bx^2)) + 4bx^2)}{48b^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[x^2\*Cos[a + b\*x^2]^2,x]

[Out] (-3\*Sqrt[Pi]\*Cos[2\*a]\*FresnelS[(2\*Sqrt[b]\*x)/Sqrt[Pi]] - 3\*Sqrt[Pi]\*FresnelC[(2\*Sqrt[b]\*x)/Sqrt[Pi]]\*Sin[2\*a] + 2\*Sqrt[b]\*x\*(4\*b\*x^2 + 3\*Sin[2\*(a + b\*x^2)]))/(48\*b^(3/2))

**Maple [A]** time = 0.031, size = 63, normalized size = 0.7

$$\frac{x^3}{6} + \frac{x \sin(2bx^2 + 2a)}{8b} - \frac{\sqrt{\pi}}{16} \left( \cos(2a) \text{FresnelS}\left(2 \frac{x\sqrt{b}}{\sqrt{\pi}}\right) + \sin(2a) \text{FresnelC}\left(2 \frac{x\sqrt{b}}{\sqrt{\pi}}\right) \right) b^{-\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2\*cos(b\*x^2+a)^2,x)

[Out] 1/6\*x^3+1/8\*x\*sin(2\*b\*x^2+2\*a)/b-1/16/b^(3/2)\*Pi^(1/2)\*(cos(2\*a)\*FresnelS(2\*x\*b^(1/2)/Pi^(1/2))+sin(2\*a)\*FresnelC(2\*x\*b^(1/2)/Pi^(1/2)))

**Maxima [C]** time = 1.89165, size = 383, normalized size = 4.21

$$64bx^3|b| + 48x|b| \sin(2bx^2 + 2a) + \sqrt{2}\sqrt{\pi} \left( \left( -3i \cos\left(\frac{1}{4}\pi + \frac{1}{2} \arctan(0, b)\right) - 3i \cos\left(-\frac{1}{4}\pi + \frac{1}{2} \arctan(0, b)\right) - 3 \sin\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(b\*x^2+a)^2,x, algorithm="maxima")

[Out] 1/384\*(64\*b\*x^3\*abs(b) + 48\*x\*abs(b)\*sin(2\*b\*x^2 + 2\*a) + sqrt(2)\*sqrt(pi)\*((-3\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - 3\*I\*cos(-1/4\*pi + 1/2\*arctan2(0,

b)) - 3\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*cos(2\*a) - (3\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 3\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*sin(2\*a))\*erf(sqrt(2\*I\*b)\*x) + ((3\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 3\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*cos(2\*a) - (3\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + 3\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - 3\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*sin(2\*a))\*erf(sqrt(-2\*I\*b)\*x))\*sqrt(abs(b)))/(b\*abs(b))

**Fricas [A]** time = 1.7135, size = 231, normalized size = 2.54

$$\frac{8b^2x^3 + 12bx \cos(bx^2 + a) \sin(bx^2 + a) - 3\pi\sqrt{\frac{b}{\pi}} \cos(2a) S\left(2x\sqrt{\frac{b}{\pi}}\right) - 3\pi\sqrt{\frac{b}{\pi}} C\left(2x\sqrt{\frac{b}{\pi}}\right) \sin(2a)}{48b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(b\*x^2+a)^2,x, algorithm="fricas")

[Out] 1/48\*(8\*b^2\*x^3 + 12\*b\*x\*cos(b\*x^2 + a)\*sin(b\*x^2 + a) - 3\*pi\*sqrt(b/pi)\*cos(2\*a)\*fresnel\_sin(2\*x\*sqrt(b/pi)) - 3\*pi\*sqrt(b/pi)\*fresnel\_cos(2\*x\*sqrt(b/pi))\*sin(2\*a))/b^2

**Sympy [B]** time = 3.31795, size = 201, normalized size = 2.21

$$\frac{b^{\frac{3}{2}}x^5\sqrt{\frac{1}{b}}\sin(2a)\Gamma\left(\frac{3}{4}\right)\Gamma\left(\frac{5}{4}\right){}_2F_3\left(\frac{3}{4}, \frac{5}{4} \middle| \frac{3}{2}, \frac{7}{4}, \frac{9}{4} \right) - b^2x^4}{8\Gamma\left(\frac{7}{4}\right)\Gamma\left(\frac{9}{4}\right)} - \frac{\sqrt{b}x^3\sqrt{\frac{1}{b}}\cos(2a)\Gamma\left(\frac{1}{4}\right)\Gamma\left(\frac{3}{4}\right){}_2F_3\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{1}{2}, \frac{5}{4}, \frac{7}{4} \right) - b^2x^4}{16\Gamma\left(\frac{5}{4}\right)\Gamma\left(\frac{7}{4}\right)} + \frac{x^3}{6} - \frac{\sqrt{\pi}x^2\sqrt{\frac{1}{b}}\sin(2a)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*2\*cos(b\*x\*\*2+a)\*\*2,x)

[Out] b\*\*(3/2)\*x\*\*5\*sqrt(1/b)\*sin(2\*a)\*gamma(3/4)\*gamma(5/4)\*hyper((3/4, 5/4), (3/2, 7/4, 9/4), -b\*\*2\*x\*\*4)/(8\*gamma(7/4)\*gamma(9/4)) - sqrt(b)\*x\*\*3\*sqrt(1/b)\*cos(2\*a)\*gamma(1/4)\*gamma(3/4)\*hyper((1/4, 3/4), (1/2, 5/4, 7/4), -b\*\*2\*x\*\*4)/(16\*gamma(5/4)\*gamma(7/4)) + x\*\*3/6 - sqrt(pi)\*x\*\*2\*sqrt(1/b)\*sin(2\*a)\*fresnels(2\*sqrt(b)\*x/sqrt(pi))/4 + sqrt(pi)\*x\*\*2\*sqrt(1/b)\*cos(2\*a)\*fresnelc(2\*sqrt(b)\*x/sqrt(pi))/4

**Giac [C]** time = 1.17095, size = 159, normalized size = 1.75

$$\frac{1}{6}x^3 - \frac{ixe^{(2ibx^2+2ia)}}{16b} + \frac{ixe^{(-2ibx^2-2ia)}}{16b} - \frac{i\sqrt{\pi}\operatorname{erf}\left(-\sqrt{bx}\left(\frac{ib}{|b|}+1\right)\right)e^{(2ia)}}{32b^{\frac{3}{2}}\left(\frac{ib}{|b|}+1\right)} + \frac{i\sqrt{\pi}\operatorname{erf}\left(-\sqrt{bx}\left(\frac{ib}{|b|}+1\right)\right)e^{(-2ia)}}{32b^{\frac{3}{2}}\left(\frac{ib}{|b|}+1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(b\*x^2+a)^2,x, algorithm="giac")

```
[Out] 1/6*x^3 - 1/16*I*x*e^(2*I*b*x^2 + 2*I*a)/b + 1/16*I*x*e^(-2*I*b*x^2 - 2*I*a)
)/b - 1/32*I*sqrt(pi)*erf(-sqrt(b)*x*(-I*b/abs(b) + 1))*e^(2*I*a)/(b^(3/2)*
(-I*b/abs(b) + 1)) + 1/32*I*sqrt(pi)*erf(-sqrt(b)*x*(I*b/abs(b) + 1))*e^(-2
*I*a)/(b^(3/2)*(I*b/abs(b) + 1))
```

### 3.10 $\int x \cos^2(a + bx^2) dx$

**Optimal.** Leaf size=31

$$\frac{\sin(a + bx^2) \cos(a + bx^2)}{4b} + \frac{x^2}{4}$$

[Out]  $x^2/4 + (\text{Cos}[a + b*x^2]*\text{Sin}[a + b*x^2])/(4*b)$

**Rubi [A]** time = 0.028877, antiderivative size = 31, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3380, 2635, 8}

$$\frac{\sin(a + bx^2) \cos(a + bx^2)}{4b} + \frac{x^2}{4}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[x*\text{Cos}[a + b*x^2]^2,x]$

[Out]  $x^2/4 + (\text{Cos}[a + b*x^2]*\text{Sin}[a + b*x^2])/(4*b)$

#### Rule 3380

$\text{Int}[(a + \text{Cos}[c + d*x])*(b + d*x)^n*(x + e)^m, x\_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*\text{Cos}[c + d*x])^p, x}], x, x^n], x] /; \text{FreeQ}\{a, b, c, d, m, n, p\}, x] \&\& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]] \&\& (\text{EqQ}[p, 1] \parallel \text{EqQ}[m, n - 1] \parallel (\text{IntegerQ}[p] \&\& \text{GtQ}[\text{Simplify}[(m + 1)/n], 0]))$

#### Rule 2635

$\text{Int}[(b + \sin[c + d*x])*(x + e)^n, x\_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c + d*x])*(b*\text{Sin}[c + d*x])^{n-1}]/(d*n), x] + \text{Dist}[(b^2*(n - 1))/n, \text{Int}[(b*\text{Sin}[c + d*x])^{n-2}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

#### Rule 8

$\text{Int}[a, x\_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

#### Rubi steps

$$\begin{aligned} \int x \cos^2(a + bx^2) dx &= \frac{1}{2} \text{Subst} \left( \int \cos^2(a + bx) dx, x, x^2 \right) \\ &= \frac{\cos(a + bx^2) \sin(a + bx^2)}{4b} + \frac{1}{4} \text{Subst} \left( \int 1 dx, x, x^2 \right) \\ &= \frac{x^2}{4} + \frac{\cos(a + bx^2) \sin(a + bx^2)}{4b} \end{aligned}$$

**Mathematica [A]** time = 0.0451212, size = 27, normalized size = 0.87

$$\frac{2(a + bx^2) + \sin(2(a + bx^2))}{8b}$$

Antiderivative was successfully verified.

[In] Integrate[x\*Cos[a + b\*x^2]^2,x]

[Out] (2\*(a + b\*x^2) + Sin[2\*(a + b\*x^2)])/(8\*b)

**Maple [A]** time = 0.026, size = 34, normalized size = 1.1

$$\frac{1}{2b} \left( \frac{\cos(bx^2 + a) \sin(bx^2 + a)}{2} + \frac{bx^2}{2} + \frac{a}{2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x\*cos(b\*x^2+a)^2,x)

[Out] 1/2/b\*(1/2\*cos(b\*x^2+a)\*sin(b\*x^2+a)+1/2\*b\*x^2+1/2\*a)

**Maxima [A]** time = 1.17817, size = 31, normalized size = 1.

$$\frac{2bx^2 + \sin(2bx^2 + 2a)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x^2+a)^2,x, algorithm="maxima")

[Out] 1/8\*(2\*b\*x^2 + sin(2\*b\*x^2 + 2\*a))/b

**Fricas [A]** time = 1.61826, size = 63, normalized size = 2.03

$$\frac{bx^2 + \cos(bx^2 + a) \sin(bx^2 + a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x^2+a)^2,x, algorithm="fricas")

[Out] 1/4\*(b\*x^2 + cos(b\*x^2 + a)\*sin(b\*x^2 + a))/b

**Sympy [A]** time = 0.517633, size = 60, normalized size = 1.94

$$\begin{cases} \frac{x^2 \sin^2(a+bx^2)}{2} + \frac{x^2 \cos^2(a+bx^2)}{4} + \frac{\sin(a+bx^2) \cos(a+bx^2)}{4b} & \text{for } b \neq 0 \\ \frac{x^2 \cos^2(a)}{2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x\*\*2+a)\*\*2,x)

```
[Out] Piecewise((x**2*sin(a + b*x**2)**2/4 + x**2*cos(a + b*x**2)**2/4 + sin(a +
b*x**2)*cos(a + b*x**2)/(4*b), Ne(b, 0)), (x**2*cos(a)**2/2, True))
```

**Giac [A]** time = 1.11567, size = 35, normalized size = 1.13

$$\frac{2bx^2 + 2a + \sin(2bx^2 + 2a)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x*cos(b*x^2+a)^2,x, algorithm="giac")
```

```
[Out] 1/8*(2*b*x^2 + 2*a + sin(2*b*x^2 + 2*a))/b
```

### 3.11 $\int \cos^2(a + bx^2) dx$

**Optimal.** Leaf size=70

$$\frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4\sqrt{b}} - \frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4\sqrt{b}} + \frac{x}{2}$$

[Out]  $x/2 + (\operatorname{Sqrt}[\pi] \operatorname{Cos}[2*a] \operatorname{FresnelC}[(2*\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[\pi]])/(4*\operatorname{Sqrt}[b]) - (\operatorname{Sqrt}[\pi] \operatorname{FresnelS}[(2*\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[\pi]]*\operatorname{Sin}[2*a])/(4*\operatorname{Sqrt}[b])$

**Rubi [A]** time = 0.0437522, antiderivative size = 70, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 4, integrand size = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.4$ , Rules used = {3358, 3354, 3352, 3351}

$$\frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4\sqrt{b}} - \frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4\sqrt{b}} + \frac{x}{2}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[\operatorname{Cos}[a + b*x^2]^2, x]$

[Out]  $x/2 + (\operatorname{Sqrt}[\pi] \operatorname{Cos}[2*a] \operatorname{FresnelC}[(2*\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[\pi]])/(4*\operatorname{Sqrt}[b]) - (\operatorname{Sqrt}[\pi] \operatorname{FresnelS}[(2*\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[\pi]]*\operatorname{Sin}[2*a])/(4*\operatorname{Sqrt}[b])$

#### Rule 3358

$\operatorname{Int}[(a + \operatorname{Cos}(c + d*(e + f*x)^n))^p, x] \rightarrow \operatorname{Int}[\operatorname{ExpandTrigReduce}[(a + b*\operatorname{Cos}[c + d*(e + f*x)^n])^p, x], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \operatorname{IGtQ}[p, 1] \ \&\& \ \operatorname{IGtQ}[n, 1]$

#### Rule 3354

$\operatorname{Int}[\operatorname{Cos}(c + d*(e + f*x)^2), x] \rightarrow \operatorname{Dist}[\operatorname{Cos}[c], \operatorname{Int}[\operatorname{Cos}[d*(e + f*x)^2], x], x] - \operatorname{Dist}[\operatorname{Sin}[c], \operatorname{Int}[\operatorname{Sin}[d*(e + f*x)^2], x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f, x\}$

#### Rule 3352

$\operatorname{Int}[\operatorname{Cos}(d*(e + f*x)^2), x] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[\pi/2] \operatorname{FresnelC}[\operatorname{Sqrt}[2/\pi]*\operatorname{Rt}[d, 2]*(e + f*x)])/(f*\operatorname{Rt}[d, 2]), x] /;$   $\operatorname{FreeQ}\{d, e, f, x\}$

#### Rule 3351

$\operatorname{Int}[\operatorname{Sin}(d*(e + f*x)^2), x] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[\pi/2] \operatorname{FresnelS}[\operatorname{Sqrt}[2/\pi]*\operatorname{Rt}[d, 2]*(e + f*x)])/(f*\operatorname{Rt}[d, 2]), x] /;$   $\operatorname{FreeQ}\{d, e, f, x\}$

#### Rubi steps



$$\begin{aligned}
\int \cos^2(a + bx^2) dx &= \int \left( \frac{1}{2} + \frac{1}{2} \cos(2a + 2bx^2) \right) dx \\
&= \frac{x}{2} + \frac{1}{2} \int \cos(2a + 2bx^2) dx \\
&= \frac{x}{2} + \frac{1}{2} \cos(2a) \int \cos(2bx^2) dx - \frac{1}{2} \sin(2a) \int \sin(2bx^2) dx \\
&= \frac{x}{2} + \frac{\sqrt{\pi} \cos(2a) C\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \sqrt{\pi} \sin(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4\sqrt{b}}
\end{aligned}$$

**Mathematica [A]** time = 0.0608273, size = 67, normalized size = 0.96

$$\frac{\sqrt{\pi} \cos(2a) \text{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \sqrt{\pi} \sin(2a) \text{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) + 2\sqrt{bx}}{4\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]^2,x]

[Out] (2\*Sqrt[b]\*x + Sqrt[Pi]\*Cos[2\*a]\*FresnelC[(2\*Sqrt[b]\*x)/Sqrt[Pi]] - Sqrt[Pi]\*FresnelS[(2\*Sqrt[b]\*x)/Sqrt[Pi]]\*Sin[2\*a])/(4\*Sqrt[b])

**Maple [A]** time = 0.035, size = 45, normalized size = 0.6

$$\frac{x}{2} + \frac{\sqrt{\pi}}{4} \left( \cos(2a) \text{FresnelC}\left(2 \frac{x\sqrt{b}}{\sqrt{\pi}}\right) - \sin(2a) \text{FresnelS}\left(2 \frac{x\sqrt{b}}{\sqrt{\pi}}\right) \right) \frac{1}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)^2,x)

[Out] 1/2\*x+1/4\*Pi^(1/2)/b^(1/2)\*(cos(2\*a)\*FresnelC(2\*x\*b^(1/2)/Pi^(1/2))-sin(2\*a)\*FresnelS(2\*x\*b^(1/2)/Pi^(1/2)))

**Maxima [C]** time = 1.91315, size = 338, normalized size = 4.83

$$\sqrt{2}\sqrt{\pi} \left( \left( \cos\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) + \cos\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) - i \sin\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) + i \sin\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2,x, algorithm="maxima")

[Out] 1/32\*(sqrt(2)\*sqrt(pi))\*(((cos(1/4\*pi + 1/2\*arctan2(0, b)) + cos(-1/4\*pi + 1/2\*arctan2(0, b)) - I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + I\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*cos(2\*a) - (I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + sin(1/4\*pi + 1/2\*arctan2(0, b)) - sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*sin(2\*a))\*erf(sqrt(2\*I\*b)\*x) + ((cos(1/4\*pi + 1/2\*arctan2(0, b)) + cos(-1/4\*pi + 1/2\*arctan2(0, b)) + I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - I\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*sin(2\*a))

, b)) - I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))\*cos(2\*a) - (-I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + sin(1/4\*pi + 1/2\*arctan2(0, b)) - sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*sin(2\*a))\*erf(sqrt(-2\*I\*b\*x))\*sqrt(abs(b)) + 16\*x\*abs(b))/abs(b)

**Fricas [A]** time = 1.63341, size = 163, normalized size = 2.33

$$\frac{\pi\sqrt{\frac{b}{\pi}}\cos(2a)C\left(2x\sqrt{\frac{b}{\pi}}\right) - \pi\sqrt{\frac{b}{\pi}}S\left(2x\sqrt{\frac{b}{\pi}}\right)\sin(2a) + 2bx}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2,x, algorithm="fricas")

[Out] 1/4\*(pi\*sqrt(b/pi)\*cos(2\*a)\*fresnel\_cos(2\*x\*sqrt(b/pi)) - pi\*sqrt(b/pi)\*fresnel\_sin(2\*x\*sqrt(b/pi))\*sin(2\*a) + 2\*b\*x)/b

**Sympy [A]** time = 1.04674, size = 56, normalized size = 0.8

$$\frac{x}{2} + \frac{\sqrt{\pi}\left(-\sin(2a)S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) + \cos(2a)C\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)\right)\sqrt{\frac{1}{b}}}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*2,x)

[Out] x/2 + sqrt(pi)\*(-sin(2\*a)\*fresnels(2\*sqrt(b)\*x/sqrt(pi)) + cos(2\*a)\*fresnelc(2\*sqrt(b)\*x/sqrt(pi)))\*sqrt(1/b)/4

**Giac [C]** time = 1.16218, size = 111, normalized size = 1.59

$$\frac{1}{2}x - \frac{\sqrt{\pi}\operatorname{erf}\left(-\sqrt{bx}\left(-\frac{ib}{|b|} + 1\right)\right)e^{(2ia)}}{8\sqrt{b}\left(-\frac{ib}{|b|} + 1\right)} - \frac{\sqrt{\pi}\operatorname{erf}\left(-\sqrt{bx}\left(\frac{ib}{|b|} + 1\right)\right)e^{(-2ia)}}{8\sqrt{b}\left(\frac{ib}{|b|} + 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2,x, algorithm="giac")

[Out] 1/2\*x - 1/8\*sqrt(pi)\*erf(-sqrt(b)\*x\*(-I\*b/abs(b) + 1))\*e^(2\*I\*a)/(sqrt(b)\*(-I\*b/abs(b) + 1)) - 1/8\*sqrt(pi)\*erf(-sqrt(b)\*x\*(I\*b/abs(b) + 1))\*e^(-2\*I\*a)/(sqrt(b)\*(I\*b/abs(b) + 1))

$$3.12 \quad \int \frac{\cos^2(a+bx^2)}{x} dx$$

**Optimal.** Leaf size=37

$$\frac{1}{4} \cos(2a) \operatorname{CosIntegral}(2bx^2) - \frac{1}{4} \sin(2a) \operatorname{Si}(2bx^2) + \frac{\log(x)}{2}$$

[Out] (Cos[2\*a]\*CosIntegral[2\*b\*x^2])/4 + Log[x]/2 - (Sin[2\*a]\*SinIntegral[2\*b\*x^2])/4

**Rubi [A]** time = 0.0518501, antiderivative size = 37, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 4, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3404, 3378, 3376, 3375}

$$\frac{1}{4} \cos(2a) \operatorname{CosIntegral}(2bx^2) - \frac{1}{4} \sin(2a) \operatorname{Si}(2bx^2) + \frac{\log(x)}{2}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]^2/x,x]

[Out] (Cos[2\*a]\*CosIntegral[2\*b\*x^2])/4 + Log[x]/2 - (Sin[2\*a]\*SinIntegral[2\*b\*x^2])/4

#### Rule 3404

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] := Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]

#### Rule 3378

Int[Cos[(c\_) + (d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] := Dist[Cos[c], Int[Cos[d\*x^n]/x, x], x] - Dist[Sin[c], Int[Sin[d\*x^n]/x, x], x] /; FreeQ[{c, d, n}, x]

#### Rule 3376

Int[Cos[(d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] := Simp[CosIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

#### Rule 3375

Int[Sin[(d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] := Simp[SinIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(a + bx^2)}{x} dx &= \int \left( \frac{1}{2x} + \frac{\cos(2a + 2bx^2)}{2x} \right) dx \\
&= \frac{\log(x)}{2} + \frac{1}{2} \int \frac{\cos(2a + 2bx^2)}{x} dx \\
&= \frac{\log(x)}{2} + \frac{1}{2} \cos(2a) \int \frac{\cos(2bx^2)}{x} dx - \frac{1}{2} \sin(2a) \int \frac{\sin(2bx^2)}{x} dx \\
&= \frac{1}{4} \cos(2a) \text{Ci}(2bx^2) + \frac{\log(x)}{2} - \frac{1}{4} \sin(2a) \text{Si}(2bx^2)
\end{aligned}$$

**Mathematica [A]** time = 0.0672536, size = 34, normalized size = 0.92

$$\frac{1}{4} (\cos(2a) \text{CosIntegral}(2bx^2) - \sin(2a) \text{Si}(2bx^2) + 2 \log(x))$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]^2/x, x]

[Out] (Cos[2\*a]\*CosIntegral[2\*b\*x^2] + 2\*Log[x] - Sin[2\*a]\*SinIntegral[2\*b\*x^2])/4

**Maple [A]** time = 0.046, size = 32, normalized size = 0.9

$$\frac{\text{Ci}(2bx^2) \cos(2a)}{4} + \frac{\ln(x)}{2} - \frac{\text{Si}(2bx^2) \sin(2a)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)^2/x, x)

[Out] 1/4\*Ci(2\*b\*x^2)\*cos(2\*a)+1/2\*ln(x)-1/4\*Si(2\*b\*x^2)\*sin(2\*a)

**Maxima [C]** time = 1.36611, size = 69, normalized size = 1.86

$$\frac{1}{8} (\text{Ei}(2ibx^2) + \text{Ei}(-2ibx^2)) \cos(2a) + \frac{1}{8} (i \text{Ei}(2ibx^2) - i \text{Ei}(-2ibx^2)) \sin(2a) + \frac{1}{2} \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x, x, algorithm="maxima")

[Out] 1/8\*(Ei(2\*I\*b\*x^2) + Ei(-2\*I\*b\*x^2))\*cos(2\*a) + 1/8\*(I\*Ei(2\*I\*b\*x^2) - I\*Ei(-2\*I\*b\*x^2))\*sin(2\*a) + 1/2\*log(x)

**Fricas [A]** time = 1.60196, size = 153, normalized size = 4.14

$$\frac{1}{8} (\text{Ci}(2bx^2) + \text{Ci}(-2bx^2)) \cos(2a) - \frac{1}{4} \sin(2a) \text{Si}(2bx^2) + \frac{1}{2} \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x,x, algorithm="fricas")

[Out] 1/8\*(cos\_integral(2\*b\*x^2) + cos\_integral(-2\*b\*x^2))\*cos(2\*a) - 1/4\*sin(2\*a)\*sin\_integral(2\*b\*x^2) + 1/2\*log(x)

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + bx^2)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*2/x,x)

[Out] Integral(cos(a + b\*x\*\*2)\*\*2/x, x)

**Giac [A]** time = 1.16839, size = 47, normalized size = 1.27

$$\frac{1}{4} \cos(2a) \operatorname{Ci}(2bx^2) + \frac{1}{4} \sin(2a) \operatorname{Si}(-2bx^2) + \frac{1}{4} \log(bx^2)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x,x, algorithm="giac")

[Out] 1/4\*cos(2\*a)\*cos\_integral(2\*b\*x^2) + 1/4\*sin(2\*a)\*sin\_integral(-2\*b\*x^2) + 1/4\*log(b\*x^2)

### 3.13 $\int \frac{\cos^2(a+bx^2)}{x^2} dx$

**Optimal.** Leaf size=76

$$-\sqrt{\pi}\sqrt{b}\sin(2a)\text{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \sqrt{\pi}\sqrt{b}\cos(2a)\text{S}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \frac{\cos^2(a+bx^2)}{x}$$

[Out]  $-(\text{Cos}[a + b*x^2]^2/x) - \text{Sqrt}[b]*\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a]*\text{FresnelS}[(2*\text{Sqrt}[b]*x)/\text{Sqrt}[\text{Pi}]] - \text{Sqrt}[b]*\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(2*\text{Sqrt}[b]*x)/\text{Sqrt}[\text{Pi}]]*\text{Sin}[2*a]$

**Rubi [A]** time = 0.0673088, antiderivative size = 76, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 6, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3394, 4573, 3373, 3353, 3352, 3351}

$$-\sqrt{\pi}\sqrt{b}\sin(2a)\text{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \sqrt{\pi}\sqrt{b}\cos(2a)\text{S}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \frac{\cos^2(a+bx^2)}{x}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[\text{Cos}[a + b*x^2]^2/x^2, x]$

[Out]  $-(\text{Cos}[a + b*x^2]^2/x) - \text{Sqrt}[b]*\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a]*\text{FresnelS}[(2*\text{Sqrt}[b]*x)/\text{Sqrt}[\text{Pi}]] - \text{Sqrt}[b]*\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(2*\text{Sqrt}[b]*x)/\text{Sqrt}[\text{Pi}]]*\text{Sin}[2*a]$

#### Rule 3394

$\text{Int}[\text{Cos}[(a_.) + (b_.)*(x_.)^{n_.}]^{p_.}*(x_.)^{m_.}, x\_Symbol] \rightarrow \text{Simp}[(x^{m+1}*\text{Cos}[a + b*x^n]^p)/(m+1), x] + \text{Dist}[(b*n*p)/(m+1), \text{Int}[\text{Cos}[a + b*x^n]^{p-1}*\text{Sin}[a + b*x^n], x], x] /;$  FreeQ[{a, b}, x] && IGtQ[p, 1] && EqQ[m + n, 0] && NeQ[n, 1] && IntegerQ[n]

#### Rule 4573

$\text{Int}[\text{Cos}[w_.]^{p_.}*(u_.)*\text{Sin}[v_.]^{p_.}, x\_Symbol] \rightarrow \text{Dist}[1/2^p, \text{Int}[u*\text{Sin}[2*v]^p, x], x] /;$  EqQ[w, v] && IntegerQ[p]

#### Rule 3373

$\text{Int}[(a_. + (b_.)*\text{Sin}[u_.])^{p_.}, x\_Symbol] \rightarrow \text{Int}[(a + b*\text{Sin}[\text{ExpandToSum}[u, x]])^p, x] /;$  FreeQ[{a, b, p}, x] && BinomialQ[u, x] && !BinomialMatchQ[u, x]

#### Rule 3353

$\text{Int}[(\text{Sin}[(c_.) + (d_.)*((e_.) + (f_.)*(x_.))^2], x\_Symbol] \rightarrow \text{Dist}[\text{Sin}[c], \text{Int}[\text{Cos}[d*(e + f*x)^2], x], x] + \text{Dist}[\text{Cos}[c], \text{Int}[\text{Sin}[d*(e + f*x)^2], x], x] /;$  FreeQ[{c, d, e, f}, x]

#### Rule 3352

$\text{Int}[\text{Cos}[(d_.)*((e_.) + (f_.)*(x_.))^2], x\_Symbol] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)])/(f*\text{Rt}[d, 2]), x] /;$  FreeQ[{d, e, f}, x]

#### Rule 3351

Int[Sin[(d\_.)\*((e\_.) + (f\_.)\*(x\_.))^2], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)]]/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

### Rubi steps

$$\begin{aligned} \int \frac{\cos^2(a + bx^2)}{x^2} dx &= -\frac{\cos^2(a + bx^2)}{x} - (4b) \int \cos(a + bx^2) \sin(a + bx^2) dx \\ &= -\frac{\cos^2(a + bx^2)}{x} - (2b) \int \sin(2(a + bx^2)) dx \\ &= -\frac{\cos^2(a + bx^2)}{x} - (2b) \int \sin(2a + 2bx^2) dx \\ &= -\frac{\cos^2(a + bx^2)}{x} - (2b \cos(2a)) \int \sin(2bx^2) dx - (2b \sin(2a)) \int \cos(2bx^2) dx \\ &= -\frac{\cos^2(a + bx^2)}{x} - \sqrt{b}\sqrt{\pi} \cos(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \sqrt{b}\sqrt{\pi} C\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \sin(2a) \end{aligned}$$

**Mathematica [A]** time = 0.176776, size = 76, normalized size = 1.

$$\frac{\sqrt{\pi}\sqrt{bx} \sin(2a) \text{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) + \sqrt{\pi}\sqrt{bx} \cos(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) + \cos^2(a + bx^2)}{x}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]^2/x^2,x]

[Out] -((Cos[a + b\*x^2]^2 + Sqrt[b]\*Sqrt[Pi]\*x\*Cos[2\*a]\*FresnelS[(2\*Sqrt[b]\*x)/Sqrt[Pi]] + Sqrt[b]\*Sqrt[Pi]\*x\*FresnelC[(2\*Sqrt[b]\*x)/Sqrt[Pi]]\*Sin[2\*a])/x)

**Maple [A]** time = 0.03, size = 62, normalized size = 0.8

$$-\frac{1}{2x} - \frac{\cos(2bx^2 + 2a)}{2x} - \sqrt{b}\sqrt{\pi} \left( \cos(2a) \text{FresnelS}\left(2\frac{x\sqrt{b}}{\sqrt{\pi}}\right) + \sin(2a) \text{FresnelC}\left(2\frac{x\sqrt{b}}{\sqrt{\pi}}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)^2/x^2,x)

[Out] -1/2/x-1/2/x\*cos(2\*b\*x^2+2\*a)-b^(1/2)\*Pi^(1/2)\*(cos(2\*a)\*FresnelS(2\*x\*b^(1/2)/Pi^(1/2))+sin(2\*a)\*FresnelC(2\*x\*b^(1/2)/Pi^(1/2)))

**Maxima [C]** time = 1.45817, size = 369, normalized size = 4.86

$$\sqrt{2}\sqrt{x^2|b|} \left( \left( \Gamma\left(-\frac{1}{2}, 2ibx^2\right) + \Gamma\left(-\frac{1}{2}, -2ibx^2\right) \right) \cos\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right) + \left( \Gamma\left(-\frac{1}{2}, 2ibx^2\right) + \Gamma\left(-\frac{1}{2}, -2ibx^2\right) \right) \cos\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0, b)\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^2,x, algorithm="maxima")

```
[Out] -1/16*(sqrt(2)*sqrt(x^2*abs(b))*((gamma(-1/2, 2*I*b*x^2) + gamma(-1/2, -2*I*b*x^2))*cos(1/4*pi + 1/2*arctan2(0, b)) + (gamma(-1/2, 2*I*b*x^2) + gamma(-1/2, -2*I*b*x^2))*cos(-1/4*pi + 1/2*arctan2(0, b)) + (I*gamma(-1/2, 2*I*b*x^2) - I*gamma(-1/2, -2*I*b*x^2))*sin(1/4*pi + 1/2*arctan2(0, b)) + (-I*gamma(-1/2, 2*I*b*x^2) + I*gamma(-1/2, -2*I*b*x^2))*sin(-1/4*pi + 1/2*arctan2(0, b)))*cos(2*a) + ((-I*gamma(-1/2, 2*I*b*x^2) + I*gamma(-1/2, -2*I*b*x^2))*cos(1/4*pi + 1/2*arctan2(0, b)) + (-I*gamma(-1/2, 2*I*b*x^2) + I*gamma(-1/2, -2*I*b*x^2))*cos(-1/4*pi + 1/2*arctan2(0, b)) + (gamma(-1/2, 2*I*b*x^2) + gamma(-1/2, -2*I*b*x^2))*sin(1/4*pi + 1/2*arctan2(0, b)) - (gamma(-1/2, 2*I*b*x^2) + gamma(-1/2, -2*I*b*x^2))*sin(-1/4*pi + 1/2*arctan2(0, b)))*sin(2*a)) + 8)/x
```

**Fricas [A]** time = 1.67794, size = 180, normalized size = 2.37

$$\frac{\pi x \sqrt{\frac{b}{\pi}} \cos(2a) S\left(2x\sqrt{\frac{b}{\pi}}\right) + \pi x \sqrt{\frac{b}{\pi}} C\left(2x\sqrt{\frac{b}{\pi}}\right) \sin(2a) + \cos(bx^2 + a)^2}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x^2+a)^2/x^2,x, algorithm="fricas")
```

```
[Out] -(pi*x*sqrt(b/pi)*cos(2*a)*fresnel_sin(2*x*sqrt(b/pi)) + pi*x*sqrt(b/pi)*fresnel_cos(2*x*sqrt(b/pi))*sin(2*a) + cos(b*x^2 + a)^2)/x
```

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + bx^2)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x**2+a)**2/x**2,x)
```

```
[Out] Integral(cos(a + b*x**2)**2/x**2, x)
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^2 + a)^2}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x^2+a)^2/x^2,x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^2 + a)^2/x^2, x)
```



### 3.14 $\int \frac{\cos^2(a+bx^2)}{x^3} dx$

**Optimal.** Leaf size=57

$$-\frac{1}{2}b \sin(2a) \operatorname{CosIntegral}(2bx^2) - \frac{1}{2}b \cos(2a) \operatorname{Si}(2bx^2) - \frac{\cos(2(a+bx^2))}{4x^2} - \frac{1}{4x^2}$$

[Out]  $-1/(4*x^2) - \operatorname{Cos}[2*(a + b*x^2)]/(4*x^2) - (b*\operatorname{CosIntegral}[2*b*x^2]*\operatorname{Sin}[2*a])/2 - (b*\operatorname{Cos}[2*a]*\operatorname{SinIntegral}[2*b*x^2])/2$

**Rubi [A]** time = 0.11751, antiderivative size = 57, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 6, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3404, 3380, 3297, 3303, 3299, 3302}

$$-\frac{1}{2}b \sin(2a) \operatorname{CosIntegral}(2bx^2) - \frac{1}{2}b \cos(2a) \operatorname{Si}(2bx^2) - \frac{\cos(2(a+bx^2))}{4x^2} - \frac{1}{4x^2}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[\operatorname{Cos}[a + b*x^2]^2/x^3, x]$

[Out]  $-1/(4*x^2) - \operatorname{Cos}[2*(a + b*x^2)]/(4*x^2) - (b*\operatorname{CosIntegral}[2*b*x^2]*\operatorname{Sin}[2*a])/2 - (b*\operatorname{Cos}[2*a]*\operatorname{SinIntegral}[2*b*x^2])/2$

#### Rule 3404

$\operatorname{Int}[(a_. + \operatorname{Cos}(c_. + (d_.)*(x_.)^{n_.})*(b_.))^{p_.}*((e_.)*(x_.))^{m_.}, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandTrigReduce}[(e*x)^m, (a + b*\operatorname{Cos}[c + d*x^n])^p, x], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, e, m\}, x \ \&\& \operatorname{IGtQ}[p, 1] \ \&\& \operatorname{IGtQ}[n, 0]$

#### Rule 3380

$\operatorname{Int}[(a_. + \operatorname{Cos}(c_. + (d_.)*(x_.)^{n_.})*(b_.))^{p_.}*(x_.)^{m_.}, x\_Symbol] \rightarrow \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n] - 1)*(a + b*\operatorname{Cos}[c + d*x])^p}, x], x, x^n], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, m, n, p\}, x \ \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]] \ \&\& (\operatorname{EqQ}[p, 1] \ \|\ \operatorname{EqQ}[m, n-1] \ \|\ (\operatorname{IntegerQ}[p] \ \&\& \operatorname{GtQ}[\operatorname{Simplify}[(m+1)/n], 0]))$

#### Rule 3297

$\operatorname{Int}[(c_. + (d_.)*(x_.))^{m_.}*\operatorname{sin}[(e_.) + (f_.)*(x_.)], x\_Symbol] \rightarrow \operatorname{Simp}[(c + d*x)^{m+1}*\operatorname{Sin}[e + f*x]/(d*(m+1)), x] - \operatorname{Dist}[f/(d*(m+1)), \operatorname{Int}[(c + d*x)^{m+1}*\operatorname{Cos}[e + f*x], x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \ \&\& \operatorname{LtQ}[m, -1]$

#### Rule 3303

$\operatorname{Int}[\operatorname{sin}[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x\_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Cos}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Sin}[(c*f)/d + f*x]/(c + d*x), x], x] + \operatorname{Dist}[\operatorname{Sin}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Cos}[(c*f)/d + f*x]/(c + d*x), x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \ \&\& \operatorname{NeQ}[d*e - c*f, 0]$

#### Rule 3299

$\operatorname{Int}[\operatorname{sin}[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{SinIntegral}[e + f*x]/d, x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \ \&\& \operatorname{EqQ}[d*e - c*f, 0]$

**Rule 3302**

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

**Rubi steps**

$$\begin{aligned}
\int \frac{\cos^2(a + bx^2)}{x^3} dx &= \int \left( \frac{1}{2x^3} + \frac{\cos(2a + 2bx^2)}{2x^3} \right) dx \\
&= -\frac{1}{4x^2} + \frac{1}{2} \int \frac{\cos(2a + 2bx^2)}{x^3} dx \\
&= -\frac{1}{4x^2} + \frac{1}{4} \text{Subst} \left( \int \frac{\cos(2a + 2bx)}{x^2} dx, x, x^2 \right) \\
&= -\frac{1}{4x^2} - \frac{\cos(2(a + bx^2))}{4x^2} - \frac{1}{2} b \text{Subst} \left( \int \frac{\sin(2a + 2bx)}{x} dx, x, x^2 \right) \\
&= -\frac{1}{4x^2} - \frac{\cos(2(a + bx^2))}{4x^2} - \frac{1}{2} (b \cos(2a)) \text{Subst} \left( \int \frac{\sin(2bx)}{x} dx, x, x^2 \right) - \frac{1}{2} (b \sin(2a)) \text{Subst} \left( \int \frac{\cos(2bx)}{x} dx, x, x^2 \right) \\
&= -\frac{1}{4x^2} - \frac{\cos(2(a + bx^2))}{4x^2} - \frac{1}{2} b \text{Ci}(2bx^2) \sin(2a) - \frac{1}{2} b \cos(2a) \text{Si}(2bx^2)
\end{aligned}$$

**Mathematica [A]** time = 0.127861, size = 50, normalized size = 0.88

$$\frac{bx^2 \sin(2a) \text{CosIntegral}(2bx^2) + bx^2 \cos(2a) \text{Si}(2bx^2) + \cos^2(a + bx^2)}{2x^2}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x^2]^2/x^3, x]
```

```
[Out] -(Cos[a + b*x^2]^2 + b*x^2*CosIntegral[2*b*x^2]*Sin[2*a] + b*x^2*Cos[2*a]*SinIntegral[2*b*x^2])/(2*x^2)
```

**Maple [C]** time = 0.138, size = 89, normalized size = 1.6

$$-\frac{1}{4x^2} + \frac{\pi e^{-2ia} \text{csgn}(bx^2) b}{4} - \frac{e^{-2ia} \text{Si}(2bx^2) b}{2} + \frac{i e^{-2ia} \text{Ei}(1, -2ibx^2) b}{4} - \frac{i e^{2ia} b \text{Ei}(1, -2ibx^2)}{4} - \frac{\cos(2bx^2 + 2a)}{4x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(b*x^2+a)^2/x^3, x)
```

```
[Out] -1/4/x^2+1/4*Pi*exp(-2*I*a)*csgn(b*x^2)*b-1/2*exp(-2*I*a)*Si(2*b*x^2)*b+1/4*I*exp(-2*I*a)*Ei(1,-2*I*b*x^2)*b-1/4*I*exp(2*I*a)*b*Ei(1,-2*I*b*x^2)-1/4*cos(2*b*x^2+2*a)/x^2
```

**Maxima [C]** time = 1.26632, size = 82, normalized size = 1.44

$$\frac{\left( (i\Gamma(-1, 2ibx^2) - i\Gamma(-1, -2ibx^2)) \cos(2a) + (\Gamma(-1, 2ibx^2) + \Gamma(-1, -2ibx^2)) \sin(2a) \right) bx^2 + 1}{4x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^3,x, algorithm="maxima")

[Out]  $-1/4 * ((I * \gamma(-1, 2 * I * b * x^2) - I * \gamma(-1, -2 * I * b * x^2)) * \cos(2 * a) + (\gamma(-1, 2 * I * b * x^2) + \gamma(-1, -2 * I * b * x^2)) * \sin(2 * a)) * b * x^2 + 1) / x^2$

**Fricas [A]** time = 1.64515, size = 194, normalized size = 3.4

$$\frac{2 b x^2 \cos(2 a) \operatorname{Si}(2 b x^2) + 2 \cos(b x^2 + a)^2 + (b x^2 \operatorname{Ci}(2 b x^2) + b x^2 \operatorname{Ci}(-2 b x^2)) \sin(2 a)}{4 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^3,x, algorithm="fricas")

[Out]  $-1/4 * (2 * b * x^2 * \cos(2 * a) * \sin\_integral(2 * b * x^2) + 2 * \cos(b * x^2 + a)^2 + (b * x^2 * \cos\_integral(2 * b * x^2) + b * x^2 * \cos\_integral(-2 * b * x^2)) * \sin(2 * a)) / x^2$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + b x^2)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*2/x\*\*3,x)

[Out] Integral(cos(a + b\*x\*\*2)\*\*2/x\*\*3, x)

**Giac [B]** time = 1.16631, size = 144, normalized size = 2.53

$$\frac{2 (b x^2 + a) b^2 \operatorname{Ci}(2 b x^2) \sin(2 a) - 2 a b^2 \operatorname{Ci}(2 b x^2) \sin(2 a) - 2 (b x^2 + a) b^2 \cos(2 a) \operatorname{Si}(-2 b x^2) + 2 a b^2 \cos(2 a) \operatorname{Si}(-2 b x^2)}{4 b^2 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^3,x, algorithm="giac")

[Out]  $-1/4 * (2 * (b * x^2 + a) * b^2 * \cos\_integral(2 * b * x^2) * \sin(2 * a) - 2 * a * b^2 * \cos\_integral(2 * b * x^2) * \sin(2 * a) - 2 * (b * x^2 + a) * b^2 * \cos(2 * a) * \sin\_integral(-2 * b * x^2) + 2 * a * b^2 * \cos(2 * a) * \sin\_integral(-2 * b * x^2) + b^2 * \cos(2 * b * x^2 + 2 * a) + b^2) / (b^2 * x^2)$

### 3.15 $\int x^3 \cos^3(a + bx^2) dx$

**Optimal.** Leaf size=79

$$\frac{\cos^3(a + bx^2)}{18b^2} + \frac{\cos(a + bx^2)}{3b^2} + \frac{x^2 \sin(a + bx^2)}{3b} + \frac{x^2 \sin(a + bx^2) \cos^2(a + bx^2)}{6b}$$

[Out] Cos[a + b\*x^2]/(3\*b^2) + Cos[a + b\*x^2]^3/(18\*b^2) + (x^2\*Sin[a + b\*x^2])/(3\*b) + (x^2\*Cos[a + b\*x^2]^2\*Sin[a + b\*x^2])/(6\*b)

**Rubi [A]** time = 0.0738608, antiderivative size = 79, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3380, 3310, 3296, 2638}

$$\frac{\cos^3(a + bx^2)}{18b^2} + \frac{\cos(a + bx^2)}{3b^2} + \frac{x^2 \sin(a + bx^2)}{3b} + \frac{x^2 \sin(a + bx^2) \cos^2(a + bx^2)}{6b}$$

Antiderivative was successfully verified.

[In] Int[x^3\*Cos[a + b\*x^2]^3,x]

[Out] Cos[a + b\*x^2]/(3\*b^2) + Cos[a + b\*x^2]^3/(18\*b^2) + (x^2\*Sin[a + b\*x^2])/(3\*b) + (x^2\*Cos[a + b\*x^2]^2\*Sin[a + b\*x^2])/(6\*b)

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
  := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p,
    x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
  && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 3310

```
Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :=
  Simp[(d*(b*Sin[e + f*x])^n)/(f^2*n^2), x] + (Dist[(b^2*(n - 1))/n, Int[(c + d*x)*
    (b*Sin[e + f*x])^(n - 2), x], x] - Simp[(b*(c + d*x)*Cos[e + f*x]*(b*Sin[e + f*x])^(n - 1))/
    (f*n), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]
```

#### Rule 3296

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := -Simp[
  ((c + d*x)^m*Cos[e + f*x])/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[e + f*x],
  x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

#### Rule 2638

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

#### Rubi steps

$$\begin{aligned}
\int x^3 \cos^3(a + bx^2) dx &= \frac{1}{2} \text{Subst} \left( \int x \cos^3(a + bx) dx, x, x^2 \right) \\
&= \frac{\cos^3(a + bx^2)}{18b^2} + \frac{x^2 \cos^2(a + bx^2) \sin(a + bx^2)}{6b} + \frac{1}{3} \text{Subst} \left( \int x \cos(a + bx) dx, x, x^2 \right) \\
&= \frac{\cos^3(a + bx^2)}{18b^2} + \frac{x^2 \sin(a + bx^2)}{3b} + \frac{x^2 \cos^2(a + bx^2) \sin(a + bx^2)}{6b} - \frac{\text{Subst} \left( \int \sin(a + bx) dx, x, x^2 \right)}{3b} \\
&= \frac{\cos(a + bx^2)}{3b^2} + \frac{\cos^3(a + bx^2)}{18b^2} + \frac{x^2 \sin(a + bx^2)}{3b} + \frac{x^2 \cos^2(a + bx^2) \sin(a + bx^2)}{6b}
\end{aligned}$$

**Mathematica [A]** time = 0.163611, size = 55, normalized size = 0.7

$$\frac{3bx^2(9\sin(a+bx^2)+\sin(3(a+bx^2)))+27\cos(a+bx^2)+\cos(3(a+bx^2))}{72b^2}$$

Antiderivative was successfully verified.

[In] Integrate[x^3\*Cos[a + b\*x^2]^3,x]

[Out] (27\*Cos[a + b\*x^2] + Cos[3\*(a + b\*x^2)] + 3\*b\*x^2\*(9\*Sin[a + b\*x^2] + Sin[3\*(a + b\*x^2)]))/(72\*b^2)

**Maple [A]** time = 0.033, size = 66, normalized size = 0.8

$$\frac{3x^2 \sin(bx^2 + a)}{8b} + \frac{3 \cos(bx^2 + a)}{8b^2} + \frac{x^2 \sin(3bx^2 + 3a)}{24b} + \frac{\cos(3bx^2 + 3a)}{72b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3\*cos(b\*x^2+a)^3,x)

[Out] 3/8\*x^2\*sin(b\*x^2+a)/b+3/8\*cos(b\*x^2+a)/b^2+1/24/b\*x^2\*sin(3\*b\*x^2+3\*a)+1/72/b^2\*cos(3\*b\*x^2+3\*a)

**Maxima [A]** time = 1.08759, size = 78, normalized size = 0.99

$$\frac{3bx^2 \sin(3bx^2 + 3a) + 27bx^2 \sin(bx^2 + a) + \cos(3bx^2 + 3a) + 27 \cos(bx^2 + a)}{72b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3\*cos(b\*x^2+a)^3,x, algorithm="maxima")

[Out] 1/72\*(3\*b\*x^2\*sin(3\*b\*x^2 + 3\*a) + 27\*b\*x^2\*sin(b\*x^2 + a) + cos(3\*b\*x^2 + 3\*a) + 27\*cos(b\*x^2 + a))/b^2

**Fricas [A]** time = 1.61153, size = 138, normalized size = 1.75

$$\frac{\cos(bx^2 + a)^3 + 3(bx^2 \cos(bx^2 + a)^2 + 2bx^2) \sin(bx^2 + a) + 6 \cos(bx^2 + a)}{18b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3\*cos(b\*x^2+a)^3,x, algorithm="fricas")

[Out]  $1/18*(\cos(b*x^2 + a)^3 + 3*(b*x^2*\cos(b*x^2 + a)^2 + 2*b*x^2)*\sin(b*x^2 + a) + 6*\cos(b*x^2 + a))/b^2$

**Sympy [A]** time = 4.01279, size = 92, normalized size = 1.16

$$\begin{cases} \frac{x^2 \sin^3(a+bx^2)}{3b} + \frac{x^2 \sin(a+bx^2) \cos^2(a+bx^2)}{2b} + \frac{\sin^2(a+bx^2) \cos(a+bx^2)}{3b^2} + \frac{7 \cos^3(a+bx^2)}{18b^2} & \text{for } b \neq 0 \\ \frac{x^4 \cos^3(a)}{4} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*3\*cos(b\*x\*\*2+a)\*\*3,x)

[Out] Piecewise((x\*\*2\*sin(a + b\*x\*\*2)\*\*3/(3\*b) + x\*\*2\*sin(a + b\*x\*\*2)\*cos(a + b\*x\*\*2)\*\*2/(2\*b) + sin(a + b\*x\*\*2)\*\*2\*cos(a + b\*x\*\*2)/(3\*b\*\*2) + 7\*cos(a + b\*x\*\*2)\*\*3/(18\*b\*\*2), Ne(b, 0)), (x\*\*4\*cos(a)\*\*3/4, True))

**Giac [A]** time = 1.1368, size = 78, normalized size = 0.99

$$\frac{3bx^2 \sin(3bx^2 + 3a) + 27bx^2 \sin(bx^2 + a) + \cos(3bx^2 + 3a) + 27 \cos(bx^2 + a)}{72b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3\*cos(b\*x^2+a)^3,x, algorithm="giac")

[Out]  $1/72*(3*b*x^2*\sin(3*b*x^2 + 3*a) + 27*b*x^2*\sin(b*x^2 + a) + \cos(3*b*x^2 + 3*a) + 27*\cos(b*x^2 + a))/b^2$

### 3.16 $\int x^2 \cos^3(a + bx^2) dx$

**Optimal.** Leaf size=188

$$\frac{3\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \sin(3a) \operatorname{FresnelC}\left(\sqrt{\frac{6}{\pi}} \sqrt{bx}\right)}{24b^{3/2}} - \frac{3\sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \cos(3a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{24b^{3/2}}$$

[Out]  $(-3\sqrt{\pi/2} \cos[a] \operatorname{FresnelS}[\sqrt{b} \sqrt{2/\pi} x]) / (8b^{3/2}) - (\sqrt{\pi/6} \cos[3a] \operatorname{FresnelS}[\sqrt{b} \sqrt{6/\pi} x]) / (24b^{3/2}) - (3\sqrt{\pi/2} \operatorname{FresnelC}[\sqrt{b} \sqrt{2/\pi} x] \sin[a]) / (8b^{3/2}) - (\sqrt{\pi/6} \operatorname{FresnelC}[\sqrt{b} \sqrt{6/\pi} x] \sin[3a]) / (24b^{3/2}) + (3x \sin[a + bx^2]) / (8b) + (x \sin[3a + 3bx^2]) / (24b)$

**Rubi [A]** time = 0.182, antiderivative size = 188, normalized size of antiderivative = 1., number of steps used = 10, number of rules used = 5, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {3404, 3386, 3353, 3352, 3351}

$$\frac{3\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \sin(3a) \operatorname{FresnelC}\left(\sqrt{\frac{6}{\pi}} \sqrt{bx}\right)}{24b^{3/2}} - \frac{3\sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \cos(3a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{24b^{3/2}}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[x^2 \cos[a + bx^2]^3, x]$

[Out]  $(-3\sqrt{\pi/2} \cos[a] \operatorname{FresnelS}[\sqrt{b} \sqrt{2/\pi} x]) / (8b^{3/2}) - (\sqrt{\pi/6} \cos[3a] \operatorname{FresnelS}[\sqrt{b} \sqrt{6/\pi} x]) / (24b^{3/2}) - (3\sqrt{\pi/2} \operatorname{FresnelC}[\sqrt{b} \sqrt{2/\pi} x] \sin[a]) / (8b^{3/2}) - (\sqrt{\pi/6} \operatorname{FresnelC}[\sqrt{b} \sqrt{6/\pi} x] \sin[3a]) / (24b^{3/2}) + (3x \sin[a + bx^2]) / (8b) + (x \sin[3a + 3bx^2]) / (24b)$

#### Rule 3404

$\operatorname{Int}[(a + \cos(c + d(x)^n)) \cdot (b + e(x)^m)]^p, x] \rightarrow \operatorname{Int}[\operatorname{ExpandTrigReduce}[(e(x)^m) \cdot (a + b \cos[c + d(x)^n])]^p, x] /; \operatorname{FreeQ}\{a, b, c, d, e, m\}, x] \ \&\& \operatorname{IGtQ}[p, 1] \ \&\& \operatorname{IGtQ}[n, 0]$

#### Rule 3386

$\operatorname{Int}[\cos(c + d(x)^n) \cdot (e + f(x)^m)]^p, x] \rightarrow \operatorname{Simp}[(e^{n-1} \cdot (e(x)^m)^{m-n+1} \cdot \sin[c + d(x)^n]) / (d \cdot n), x] - \operatorname{Dist}[(e^{n-1} \cdot (e(x)^m)^{m-n+1}) / (d \cdot n), \operatorname{Int}[(e(x)^m)^{m-n} \cdot \sin[c + d(x)^n], x], x] /; \operatorname{FreeQ}\{c, d, e\}, x] \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{LtQ}[n, m + 1]$

#### Rule 3353

$\operatorname{Int}[\sin(c + d(x)^n) \cdot (e + f(x)^2)]^p, x] \rightarrow \operatorname{Dist}[\sin[c], \operatorname{Int}[\cos[d \cdot (e + f(x)^2)], x], x] + \operatorname{Dist}[\cos[c], \operatorname{Int}[\sin[d \cdot (e + f(x)^2)], x], x] /; \operatorname{FreeQ}\{c, d, e, f\}, x]$

#### Rule 3352

$\operatorname{Int}[\cos(d \cdot (e + f(x)^2))]^p, x] \rightarrow \operatorname{Simp}[(\sqrt{\pi/2} \operatorname{FresnelC}[\sqrt{2/\pi} \operatorname{Rt}[d, 2] \cdot (e + f(x))]) / (f \cdot \operatorname{Rt}[d, 2]), x] /; \operatorname{FreeQ}\{d, e, f\}, x]$

#### Rule 3351

Int[Sin[(d\_.)\*((e\_.) + (f\_.)\*(x\_))<sup>2</sup>], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)]]/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

### Rubi steps

$$\begin{aligned} \int x^2 \cos^3(a + bx^2) dx &= \int \left( \frac{3}{4}x^2 \cos(a + bx^2) + \frac{1}{4}x^2 \cos(3a + 3bx^2) \right) dx \\ &= \frac{1}{4} \int x^2 \cos(3a + 3bx^2) dx + \frac{3}{4} \int x^2 \cos(a + bx^2) dx \\ &= \frac{3x \sin(a + bx^2)}{8b} + \frac{x \sin(3a + 3bx^2)}{24b} - \frac{\int \sin(3a + 3bx^2) dx}{24b} - \frac{3 \int \sin(a + bx^2) dx}{8b} \\ &= \frac{3x \sin(a + bx^2)}{8b} + \frac{x \sin(3a + 3bx^2)}{24b} - \frac{(3 \cos(a)) \int \sin(bx^2) dx}{8b} - \frac{\cos(3a) \int \sin(3bx^2) dx}{24b} \\ &= -\frac{3\sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \cos(3a) S\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{24b^{3/2}} - \frac{3\sqrt{\frac{\pi}{2}} C\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} C\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right) \cos(3a)}{24b^{3/2}} \end{aligned}$$

**Mathematica [A]** time = 0.431522, size = 160, normalized size = 0.85

$$\frac{-27\sqrt{2\pi} \sin(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{bx}\right) - \sqrt{6\pi} \sin(3a) \text{FresnelC}\left(\sqrt{\frac{6}{\pi}}\sqrt{bx}\right) - 27\sqrt{2\pi} \cos(a) S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \sqrt{6\pi} \cos(3a) S\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{144b^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[x<sup>2</sup>\*Cos[a + b\*x<sup>2</sup>]<sup>3</sup>,x]

[Out] (-27\*Sqrt[2\*Pi]\*Cos[a]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x] - Sqrt[6\*Pi]\*Cos[3\*a]\*FresnelS[Sqrt[b]\*Sqrt[6/Pi]\*x] - 27\*Sqrt[2\*Pi]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x]\*Sin[a] - Sqrt[6\*Pi]\*FresnelC[Sqrt[b]\*Sqrt[6/Pi]\*x]\*Sin[3\*a] + 54\*Sqrt[b]\*x\*Sin[a + b\*x<sup>2</sup>] + 6\*Sqrt[b]\*x\*Sin[3\*(a + b\*x<sup>2</sup>)])/(144\*b<sup>(3/2)</sup>)

**Maple [A]** time = 0.033, size = 130, normalized size = 0.7

$$\frac{3x \sin(bx^2 + a)}{8b} - \frac{3\sqrt{2}\sqrt{\pi}}{16} \left( \cos(a) \text{FresnelS}\left(\frac{x\sqrt{2}}{\sqrt{\pi}}\sqrt{b}\right) + \sin(a) \text{FresnelC}\left(\frac{x\sqrt{2}}{\sqrt{\pi}}\sqrt{b}\right) \right) b^{-\frac{3}{2}} + \frac{x \sin(3bx^2 + 3a)}{24b} - \frac{\sqrt{2}\sqrt{\pi}}{16} \left( \cos(3a) \text{FresnelS}\left(\frac{x\sqrt{2}}{\sqrt{\pi}}\sqrt{b}\right) + \sin(3a) \text{FresnelC}\left(\frac{x\sqrt{2}}{\sqrt{\pi}}\sqrt{b}\right) \right) b^{-\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x<sup>2</sup>\*cos(b\*x<sup>2</sup>+a)<sup>3</sup>,x)

[Out] 3/8\*x\*sin(b\*x<sup>2</sup>+a)/b-3/16/b<sup>(3/2)</sup>\*2<sup>(1/2)</sup>\*Pi<sup>(1/2)</sup>\*(cos(a)\*FresnelS(x\*b<sup>(1/2)</sup>\*2<sup>(1/2)</sup>/Pi<sup>(1/2)</sup>)+sin(a)\*FresnelC(x\*b<sup>(1/2)</sup>\*2<sup>(1/2)</sup>/Pi<sup>(1/2)</sup>))+1/24\*x\*sin(3\*b\*x<sup>2</sup>+3\*a)/b-1/144/b<sup>(3/2)</sup>\*2<sup>(1/2)</sup>\*Pi<sup>(1/2)</sup>\*3<sup>(1/2)</sup>\*(cos(3\*a)\*FresnelS(2<sup>(1/2)</sup>/Pi<sup>(1/2)</sup>\*3<sup>(1/2)</sup>\*b<sup>(1/2)</sup>\*x)+sin(3\*a)\*FresnelC(2<sup>(1/2)</sup>/Pi<sup>(1/2)</sup>\*3<sup>(1/2)</sup>\*b<sup>(1/2)</sup>\*x))

**Maxima [C]** time = 2.1503, size = 697, normalized size = 3.71

result too large to display

Verification of antiderivative is not currently implemented for this CAS.



[In] integrate(x^2\*cos(b\*x^2+a)^3,x, algorithm="maxima")

[Out] 1/576\*(24\*x\*abs(b)\*sin(3\*b\*x^2 + 3\*a) + 216\*x\*abs(b)\*sin(b\*x^2 + a) + sqrt(3)\*sqrt(pi)\*(((-I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - sin(1/4\*pi + 1/2\*arctan2(0, b)) + sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(3\*a) - (cos(1/4\*pi + 1/2\*arctan2(0, b)) + cos(-1/4\*pi + 1/2\*arctan2(0, b)) - I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(3\*a)) \* erf(sqrt(3\*I\*b)\*x) + ((I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - sin(1/4\*pi + 1/2\*arctan2(0, b)) + sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(3\*a) - (cos(1/4\*pi + 1/2\*arctan2(0, b)) + cos(-1/4\*pi + 1/2\*arctan2(0, b)) + I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(3\*a)) \* erf(sqrt(-3\*I\*b)\*x)) \* sqrt(abs(b)) + sqrt(pi)\*((( -27\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - 27\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 27\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 27\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(a) - (27\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 27\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 27\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 27\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(a)) \* erf(sqrt(I\*b)\*x) + ((27\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 27\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 27\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 27\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(a) - (27\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 27\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + 27\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - 27\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(a)) \* erf(sqrt(-I\*b)\*x)) \* sqrt(abs(b)))/(b\*abs(b))

**Fricas [A]** time = 1.78099, size = 450, normalized size = 2.39

$$\frac{\sqrt{6}\pi\sqrt{\frac{b}{\pi}}\cos(3a)S\left(\sqrt{6}x\sqrt{\frac{b}{\pi}}\right) + 27\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) + \sqrt{6}\pi\sqrt{\frac{b}{\pi}}C\left(\sqrt{6}x\sqrt{\frac{b}{\pi}}\right)\sin(3a) + 27\sqrt{2}\pi\sqrt{\frac{b}{\pi}}C\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)\sin(a)}{144b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(b\*x^2+a)^3,x, algorithm="fricas")

[Out] -1/144\*(sqrt(6)\*pi\*sqrt(b/pi)\*cos(3\*a)\*fresnel\_sin(sqrt(6)\*x\*sqrt(b/pi)) + 27\*sqrt(2)\*pi\*sqrt(b/pi)\*cos(a)\*fresnel\_sin(sqrt(2)\*x\*sqrt(b/pi)) + sqrt(6)\*pi\*sqrt(b/pi)\*fresnel\_cos(sqrt(6)\*x\*sqrt(b/pi))\*sin(3\*a) + 27\*sqrt(2)\*pi\*sqrt(b/pi)\*fresnel\_cos(sqrt(2)\*x\*sqrt(b/pi))\*sin(a) - 24\*(b\*x\*cos(b\*x^2 + a)^2 + 2\*b\*x)\*sin(b\*x^2 + a))/b^2

**Sympy [B]** time = 5.78754, size = 439, normalized size = 2.34

$$\frac{3b^{\frac{3}{2}}x^5\sqrt{\frac{1}{b}}\sin(a)\Gamma\left(\frac{3}{4}\right)\Gamma\left(\frac{5}{4}\right){}_2F_3\left(\frac{3}{4}, \frac{5}{4} \middle| \frac{b^2x^4}{4}\right)}{32\Gamma\left(\frac{7}{4}\right)\Gamma\left(\frac{9}{4}\right)} + \frac{3b^{\frac{3}{2}}x^5\sqrt{\frac{1}{b}}\sin(3a)\Gamma\left(\frac{3}{4}\right)\Gamma\left(\frac{5}{4}\right){}_2F_3\left(\frac{3}{4}, \frac{5}{4} \middle| \frac{9b^2x^4}{4}\right)}{32\Gamma\left(\frac{7}{4}\right)\Gamma\left(\frac{9}{4}\right)} - \frac{3\sqrt{b}x^3\sqrt{\frac{1}{b}}\cos(a)}{144b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*2\*cos(b\*x\*\*2+a)\*\*3,x)

[Out] 3\*b\*\*(3/2)\*x\*\*5\*sqrt(1/b)\*sin(a)\*gamma(3/4)\*gamma(5/4)\*hyper((3/4, 5/4), (3/2, 7/4, 9/4), -b\*\*2\*x\*\*4/4)/(32\*gamma(7/4)\*gamma(9/4)) + 3\*b\*\*(3/2)\*x\*\*5\*sqrt(1/b)\*sin(3\*a)\*gamma(3/4)\*gamma(5/4)\*hyper((3/4, 5/4), (3/2, 7/4, 9/4), -9\*b\*\*2\*x\*\*4/4)/(32\*gamma(7/4)\*gamma(9/4)) - 3\*sqrt(b)\*x\*\*3\*sqrt(1/b)\*cos(a)

) $\gamma(1/4)\gamma(3/4)\text{hyper}((1/4, 3/4), (1/2, 5/4, 7/4), -b^{**2}x^{**4}/4)/(32\gamma(5/4)\gamma(7/4)) - \sqrt{b}x^{**3}\sqrt{1/b}\cos(3a)\gamma(1/4)\gamma(3/4)\text{hyper}((1/4, 3/4), (1/2, 5/4, 7/4), -9b^{**2}x^{**4}/4)/(32\gamma(5/4)\gamma(7/4)) - 3\sqrt{2}\sqrt{\pi}x^{**2}\sqrt{1/b}\sin(a)\text{fresnels}(\sqrt{2}\sqrt{b}x/\sqrt{\pi})/8 - \sqrt{6}\sqrt{\pi}x^{**2}\sqrt{1/b}\sin(3a)\text{fresnels}(\sqrt{6}\sqrt{b}x/\sqrt{\pi})/24 + 3\sqrt{2}\sqrt{\pi}x^{**2}\sqrt{1/b}\cos(a)\text{fresnelc}(\sqrt{2}\sqrt{b}x/\sqrt{\pi})/8 + \sqrt{6}\sqrt{\pi}x^{**2}\sqrt{1/b}\cos(3a)\text{fresnelc}(\sqrt{6}\sqrt{b}x/\sqrt{\pi})/24$

**Giac [C]** time = 1.19145, size = 350, normalized size = 1.86

$$-\frac{ixe^{(3ibx^2+3ia)}}{48b} - \frac{3ixe^{(ibx^2+ia)}}{16b} + \frac{3ixe^{(-ibx^2-ia)}}{16b} + \frac{ixe^{(-3ibx^2-3ia)}}{48b} - \frac{i\sqrt{6}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{6}\sqrt{bx}\left(-\frac{ib}{|b|}+1\right)\right)e^{(3ia)}}{288b^{\frac{3}{2}}\left(-\frac{ib}{|b|}+1\right)} - \frac{3i\sqrt{2}\sqrt{\pi}e^{(3ia)}}{288b^{\frac{3}{2}}\left(-\frac{ib}{|b|}+1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(b\*x^2+a)^3,x, algorithm="giac")

[Out]  $-1/48Ix^2e^{(3Ib^2x^2 + 3Ia)}/b - 3/16Ix^2e^{(Ib^2x^2 + Ia)}/b + 3/16Ix^2e^{(-Ib^2x^2 - Ia)}/b + 1/48Ix^2e^{(-3Ib^2x^2 - 3Ia)}/b - 1/288I\sqrt{6}\sqrt{\pi}\operatorname{erf}(-1/2\sqrt{6}\sqrt{bx}(-Ib/abs(b) + 1))e^{(3Ia)}/(b^{(3/2)}(-Ib/abs(b) + 1)) - 3/32I\sqrt{2}\sqrt{\pi}\operatorname{erf}(-1/2\sqrt{2}\sqrt{bx}(-Ib/abs(b) + 1))\sqrt{abs(b)}e^{(Ia)}/(b(-Ib/abs(b) + 1)\sqrt{abs(b)}) + 3/32I\sqrt{2}\sqrt{\pi}\operatorname{erf}(-1/2\sqrt{2}\sqrt{bx}(Ib/abs(b) + 1))\sqrt{abs(b)}e^{(-Ia)}/(b(Ib/abs(b) + 1)\sqrt{abs(b)}) + 1/288I\sqrt{6}\sqrt{\pi}\operatorname{erf}(-1/2\sqrt{6}\sqrt{bx}(Ib/abs(b) + 1))e^{(-3Ia)}/(b^{(3/2)}(Ib/abs(b) + 1))$

### 3.17 $\int x \cos^3(a + bx^2) dx$

**Optimal.** Leaf size=33

$$\frac{\sin(a + bx^2)}{2b} - \frac{\sin^3(a + bx^2)}{6b}$$

[Out] Sin[a + b\*x^2]/(2\*b) - Sin[a + b\*x^2]^3/(6\*b)

**Rubi [A]** time = 0.0283865, antiderivative size = 33, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 2, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3380, 2633}

$$\frac{\sin(a + bx^2)}{2b} - \frac{\sin^3(a + bx^2)}{6b}$$

Antiderivative was successfully verified.

[In] Int[x\*Cos[a + b\*x^2]^3,x]

[Out] Sin[a + b\*x^2]/(2\*b) - Sin[a + b\*x^2]^3/(6\*b)

#### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

#### Rule 2633

Int[sin[(c\_.) + (d\_.)\*(x\_)^(n\_.)], x\_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d\*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

#### Rubi steps

$$\begin{aligned} \int x \cos^3(a + bx^2) dx &= \frac{1}{2} \text{Subst} \left( \int \cos^3(a + bx) dx, x, x^2 \right) \\ &= -\frac{\text{Subst} \left( \int (1 - x^2) dx, x, -\sin(a + bx^2) \right)}{2b} \\ &= \frac{\sin(a + bx^2)}{2b} - \frac{\sin^3(a + bx^2)}{6b} \end{aligned}$$

**Mathematica [A]** time = 0.0170718, size = 33, normalized size = 1.

$$\frac{\sin(a + bx^2)}{2b} - \frac{\sin^3(a + bx^2)}{6b}$$

Antiderivative was successfully verified.

[In] Integrate[x\*Cos[a + b\*x^2]^3,x]

[Out]  $\text{Sin}[a + b*x^2]/(2*b) - \text{Sin}[a + b*x^2]^3/(6*b)$

**Maple [A]** time = 0.029, size = 26, normalized size = 0.8

$$\frac{\left(2 + \left(\cos\left(bx^2 + a\right)\right)^2\right) \sin\left(bx^2 + a\right)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*cos(b*x^2+a)^3,x)`

[Out]  $1/6/b*(2+\cos(b*x^2+a)^2)*\sin(b*x^2+a)$

**Maxima [A]** time = 1.17494, size = 36, normalized size = 1.09

$$\frac{\sin\left(3bx^2 + 3a\right) + 9\sin\left(bx^2 + a\right)}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*cos(b*x^2+a)^3,x, algorithm="maxima")`

[Out]  $1/24*(\sin(3*b*x^2 + 3*a) + 9*\sin(b*x^2 + a))/b$

**Fricas [A]** time = 1.56497, size = 61, normalized size = 1.85

$$\frac{\left(\cos\left(bx^2 + a\right)^2 + 2\right) \sin\left(bx^2 + a\right)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*cos(b*x^2+a)^3,x, algorithm="fricas")`

[Out]  $1/6*(\cos(b*x^2 + a)^2 + 2)*\sin(b*x^2 + a)/b$

**Sympy [A]** time = 0.970186, size = 44, normalized size = 1.33

$$\begin{cases} \frac{\sin^3(a+bx^2)}{3b} + \frac{\sin(a+bx^2)\cos^2(a+bx^2)}{2b} & \text{for } b \neq 0 \\ \frac{x^2 \cos^3(a)}{2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*cos(b*x**2+a)**3,x)`

[Out] `Piecewise((sin(a + b*x**2)**3/(3*b) + sin(a + b*x**2)*cos(a + b*x**2)**2/(2*b), Ne(b, 0)), (x**2*cos(a)**3/2, True))`

---

**Giac [A]** time = 1.1619, size = 35, normalized size = 1.06

$$\frac{\sin(bx^2 + a)^3 - 3 \sin(bx^2 + a)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x*cos(b*x^2+a)^3,x, algorithm="giac")
```

```
[Out] -1/6*(sin(b*x^2 + a)^3 - 3*sin(b*x^2 + a))/b
```

### 3.18 $\int \cos^3(a + bx^2) dx$

**Optimal.** Leaf size=153

$$\frac{3\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right)}{4\sqrt{b}} + \frac{\sqrt{\frac{\pi}{6}} \cos(3a) \operatorname{FresnelC}\left(\sqrt{\frac{6}{\pi}} \sqrt{bx}\right)}{4\sqrt{b}} - \frac{3\sqrt{\frac{\pi}{2}} \sin(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{4\sqrt{b}} - \frac{\sqrt{\frac{\pi}{6}} \sin(3a) S\left(\sqrt{b} \sqrt{\frac{6}{\pi}} x\right)}{4\sqrt{b}}$$

[Out] (3\*Sqrt [Pi/2]\*Cos [a]\*FresnelC[Sqrt [b]\*Sqrt [2/Pi]\*x])/(4\*Sqrt [b]) + (Sqrt [Pi/6]\*Cos [3\*a]\*FresnelC[Sqrt [b]\*Sqrt [6/Pi]\*x])/(4\*Sqrt [b]) - (3\*Sqrt [Pi/2]\*FresnelS[Sqrt [b]\*Sqrt [2/Pi]\*x]\*Sin[a])/(4\*Sqrt [b]) - (Sqrt [Pi/6]\*FresnelS[Sqrt [b]\*Sqrt [6/Pi]\*x]\*Sin[3\*a])/(4\*Sqrt [b])

**Rubi [A]** time = 0.0890601, antiderivative size = 153, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 4, integrand size = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.4$ , Rules used = {3358, 3354, 3352, 3351}

$$\frac{3\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{bx}\right)}{4\sqrt{b}} + \frac{\sqrt{\frac{\pi}{6}} \cos(3a) \operatorname{FresnelC}\left(\sqrt{\frac{6}{\pi}} \sqrt{bx}\right)}{4\sqrt{b}} - \frac{3\sqrt{\frac{\pi}{2}} \sin(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right)}{4\sqrt{b}} - \frac{\sqrt{\frac{\pi}{6}} \sin(3a) S\left(\sqrt{b} \sqrt{\frac{6}{\pi}} x\right)}{4\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]^3, x]

[Out] (3\*Sqrt [Pi/2]\*Cos [a]\*FresnelC[Sqrt [b]\*Sqrt [2/Pi]\*x])/(4\*Sqrt [b]) + (Sqrt [Pi/6]\*Cos [3\*a]\*FresnelC[Sqrt [b]\*Sqrt [6/Pi]\*x])/(4\*Sqrt [b]) - (3\*Sqrt [Pi/2]\*FresnelS[Sqrt [b]\*Sqrt [2/Pi]\*x]\*Sin[a])/(4\*Sqrt [b]) - (Sqrt [Pi/6]\*FresnelS[Sqrt [b]\*Sqrt [6/Pi]\*x]\*Sin[3\*a])/(4\*Sqrt [b])

#### Rule 3358

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)])\*(b\_.))^(p\_), x\_Symbol] := Int[ExpandTrigReduce[(a + b\*Cos[c + d\*(e + f\*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 1] && IGtQ[n, 1]

#### Rule 3354

Int[Cos[(c\_) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^2], x\_Symbol] := Dist[Cos[c], Int[Cos[d\*(e + f\*x)^2], x], x] - Dist[Sin[c], Int[Sin[d\*(e + f\*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]

#### Rule 3352

Int[Cos[(d\_.)\*((e\_.) + (f\_.)\*(x\_))^2], x\_Symbol] := Simp[(Sqrt [Pi/2]\*FresnelC[Sqrt [2/Pi]\*Rt [d, 2]\*(e + f\*x)])/(f\*Rt [d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rule 3351

Int[Sin[(d\_.)\*((e\_.) + (f\_.)\*(x\_))^2], x\_Symbol] := Simp[(Sqrt [Pi/2]\*FresnelS[Sqrt [2/Pi]\*Rt [d, 2]\*(e + f\*x)])/(f\*Rt [d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rubi steps

$$\begin{aligned}
\int \cos^3(a + bx^2) dx &= \int \left( \frac{3}{4} \cos(a + bx^2) + \frac{1}{4} \cos(3a + 3bx^2) \right) dx \\
&= \frac{1}{4} \int \cos(3a + 3bx^2) dx + \frac{3}{4} \int \cos(a + bx^2) dx \\
&= \frac{1}{4} (3 \cos(a)) \int \cos(bx^2) dx + \frac{1}{4} \cos(3a) \int \cos(3bx^2) dx - \frac{1}{4} (3 \sin(a)) \int \sin(bx^2) dx - \frac{1}{4} \sin(3a) \int \sin(3bx^2) dx \\
&= \frac{3\sqrt{\frac{\pi}{2}} \cos(a) C\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{4\sqrt{b}} + \frac{\sqrt{\frac{\pi}{6}} \cos(3a) C\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{4\sqrt{b}} - \frac{3\sqrt{\frac{\pi}{2}} S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a)}{4\sqrt{b}} - \frac{\sqrt{\frac{\pi}{6}} S\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right) \sin(3a)}{4\sqrt{b}}
\end{aligned}$$

**Mathematica [A]** time = 0.237082, size = 116, normalized size = 0.76

$$\frac{\sqrt{\frac{\pi}{6}} \left( 3\sqrt{3} \cos(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{bx}\right) + \cos(3a) \text{FresnelC}\left(\sqrt{\frac{6}{\pi}}\sqrt{bx}\right) - 3\sqrt{3} \sin(a) S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \sin(3a) S\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right) \right)}{4\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]^3, x]

[Out] (Sqrt[Pi/6]\*(3\*Sqrt[3]\*Cos[a]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x] + Cos[3\*a]\*FresnelC[Sqrt[b]\*Sqrt[6/Pi]\*x] - 3\*Sqrt[3]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x]\*Sin[a] - FresnelS[Sqrt[b]\*Sqrt[6/Pi]\*x]\*Sin[3\*a]))/(4\*Sqrt[b])

**Maple [A]** time = 0.033, size = 101, normalized size = 0.7

$$\frac{3\sqrt{2}\sqrt{\pi}}{8} \left( \cos(a) \text{FresnelC}\left(\frac{x\sqrt{2}}{\sqrt{\pi}}\sqrt{b}\right) - \sin(a) \text{FresnelS}\left(\frac{x\sqrt{2}}{\sqrt{\pi}}\sqrt{b}\right) \right) \frac{1}{\sqrt{b}} + \frac{\sqrt{2}\sqrt{\pi}\sqrt{3}}{24} \left( \cos(3a) \text{FresnelC}\left(\frac{\sqrt{2}\sqrt{3}x}{\sqrt{\pi}}\sqrt{b}\right) - \sin(3a) \text{FresnelS}\left(\frac{\sqrt{2}\sqrt{3}x}{\sqrt{\pi}}\sqrt{b}\right) \right) \frac{1}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)^3, x)

[Out] 3/8\*2^(1/2)\*Pi^(1/2)/b^(1/2)\*(cos(a)\*FresnelC(x\*b^(1/2)\*2^(1/2)/Pi^(1/2))-sin(a)\*FresnelS(x\*b^(1/2)\*2^(1/2)/Pi^(1/2)))+1/24\*2^(1/2)\*Pi^(1/2)\*3^(1/2)/b^(1/2)\*(cos(3\*a)\*FresnelC(2^(1/2)/Pi^(1/2)\*3^(1/2)\*b^(1/2)\*x)-sin(3\*a)\*FresnelS(2^(1/2)/Pi^(1/2)\*3^(1/2)\*b^(1/2)\*x))

**Maxima [C]** time = 2.21973, size = 653, normalized size = 4.27

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3, x, algorithm="maxima")

[Out] 1/96\*(sqrt(3)\*sqrt(pi)\*(((cos(1/4\*pi + 1/2\*arctan2(0, b)) + cos(-1/4\*pi + 1/2\*arctan2(0, b)) - I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*cos(3\*a) - (I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + sin(1/4\*pi + 1/2\*arctan2(0, b)) - sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*sin(3\*a))\*erf(sqrt(3\*I\*b)\*x) + ((cos(1/4\*pi + 1/2\*arctan2(0, b)) + cos(-1/4\*pi + 1/2\*arctan2(0, b)) - I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*sin(3\*a))

$n2(0, b)) + \cos(-1/4*\pi + 1/2*\arctan2(0, b)) + I*\sin(1/4*\pi + 1/2*\arctan2(0, b)) - I*\sin(-1/4*\pi + 1/2*\arctan2(0, b))*\cos(3*a) - (-I*\cos(1/4*\pi + 1/2*\arctan2(0, b)) - I*\cos(-1/4*\pi + 1/2*\arctan2(0, b)) + \sin(1/4*\pi + 1/2*\arctan2(0, b)) - \sin(-1/4*\pi + 1/2*\arctan2(0, b)))*\sin(3*a))*\operatorname{erf}(\sqrt{-3*I*b}*x))*\sqrt{\operatorname{abs}(b)} + \sqrt{\pi}*(((9*\cos(1/4*\pi + 1/2*\arctan2(0, b)) + 9*\cos(-1/4*\pi + 1/2*\arctan2(0, b)) - 9*I*\sin(1/4*\pi + 1/2*\arctan2(0, b)) + 9*I*\sin(-1/4*\pi + 1/2*\arctan2(0, b)))*\cos(a) - (9*I*\cos(1/4*\pi + 1/2*\arctan2(0, b)) + 9*I*\cos(-1/4*\pi + 1/2*\arctan2(0, b)) + 9*\sin(1/4*\pi + 1/2*\arctan2(0, b)) - 9*\sin(-1/4*\pi + 1/2*\arctan2(0, b)))*\sin(a))*\operatorname{erf}(\sqrt{I*b}*x) + ((9*\cos(1/4*\pi + 1/2*\arctan2(0, b)) + 9*\cos(-1/4*\pi + 1/2*\arctan2(0, b)) + 9*I*\sin(1/4*\pi + 1/2*\arctan2(0, b)) - 9*I*\sin(-1/4*\pi + 1/2*\arctan2(0, b)))*\cos(a) - (9*I*\cos(1/4*\pi + 1/2*\arctan2(0, b)) - 9*I*\cos(-1/4*\pi + 1/2*\arctan2(0, b)) + 9*\sin(1/4*\pi + 1/2*\arctan2(0, b)) - 9*\sin(-1/4*\pi + 1/2*\arctan2(0, b)))*\sin(a))*\operatorname{erf}(\sqrt{-I*b}*x))*\sqrt{\operatorname{abs}(b)))/\operatorname{abs}(b)$

**Fricas [A]** time = 1.62326, size = 373, normalized size = 2.44

$$\frac{\sqrt{6}\pi\sqrt{\frac{b}{\pi}}\cos(3a)C\left(\sqrt{6}x\sqrt{\frac{b}{\pi}}\right) + 9\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)C\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) - \sqrt{6}\pi\sqrt{\frac{b}{\pi}}S\left(\sqrt{6}x\sqrt{\frac{b}{\pi}}\right)\sin(3a) - 9\sqrt{2}\pi\sqrt{\frac{b}{\pi}}S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)\sin(a)}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3,x, algorithm="fricas")

[Out]  $\frac{1}{24}*(\sqrt{6}*\pi*\sqrt{b/\pi}*\cos(3*a)*\operatorname{fresnel\_cos}(\sqrt{6}*x*\sqrt{b/\pi})) + 9*\sqrt{2}*\pi*\sqrt{b/\pi}*\cos(a)*\operatorname{fresnel\_cos}(\sqrt{2}*x*\sqrt{b/\pi}) - \sqrt{6}*\pi*\sqrt{b/\pi}*\operatorname{fresnel\_sin}(\sqrt{6}*x*\sqrt{b/\pi})*\sin(3*a) - 9*\sqrt{2}*\pi*\sqrt{b/\pi}*\operatorname{fresnel\_sin}(\sqrt{2}*x*\sqrt{b/\pi})*\sin(a))/b$

**Sympy [A]** time = 1.56403, size = 129, normalized size = 0.84

$$\frac{3\sqrt{2}\sqrt{\pi}\left(-\sin(a)S\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right) + \cos(a)C\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right)\right)\sqrt{\frac{1}{b}}}{8} + \frac{\sqrt{6}\sqrt{\pi}\left(-\sin(3a)S\left(\frac{\sqrt{6}\sqrt{bx}}{\sqrt{\pi}}\right) + \cos(3a)C\left(\frac{\sqrt{6}\sqrt{bx}}{\sqrt{\pi}}\right)\right)\sqrt{\frac{1}{b}}}{24}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*3,x)

[Out]  $3*\sqrt{2}*\sqrt{\pi}*(-\sin(a)*\operatorname{fresnelc}(\sqrt{2}*\sqrt{b}*x/\sqrt{\pi})) + \cos(a)*\operatorname{fresnelc}(\sqrt{2}*\sqrt{b}*x/\sqrt{\pi}))*\sqrt{1/b}/8 + \sqrt{6}*\sqrt{\pi}*(-\sin(3*a)*\operatorname{fresnelc}(\sqrt{6}*\sqrt{b}*x/\sqrt{\pi})) + \cos(3*a)*\operatorname{fresnelc}(\sqrt{6}*\sqrt{b}*x/\sqrt{\pi}))*\sqrt{1/b}/24$

**Giac [C]** time = 1.1657, size = 250, normalized size = 1.63

$$\frac{\sqrt{6}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{6}\sqrt{bx}\left(-\frac{ib}{|b|}+1\right)\right)e^{(3ia)}}{48\sqrt{b}\left(-\frac{ib}{|b|}+1\right)} - \frac{3\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{(ia)}}{16\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}} - \frac{3\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)}{16\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}}$$

Verification of antiderivative is not currently implemented for this CAS.



[In] integrate(cos(b\*x^2+a)^3,x, algorithm="giac")

[Out] 
$$\begin{aligned} & -1/48*\sqrt{6}*\sqrt{\pi}*\operatorname{erf}(-1/2*\sqrt{6}*\sqrt{b}*x*(-I*b/\operatorname{abs}(b) + 1))*e^{(3*I*a)/(\sqrt{b}*(-I*b/\operatorname{abs}(b) + 1))} - 3/16*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-1/2*\sqrt{2}*x*(-I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})*e^{(I*a)/((-I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})} - \\ & 3/16*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-1/2*\sqrt{2}*x*(I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})*e^{(-I*a)/((I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})} - 1/48*\sqrt{6}*\sqrt{\pi}*\operatorname{erf}(-1/2*\sqrt{6}*\sqrt{b}*x*(I*b/\operatorname{abs}(b) + 1))*e^{(-3*I*a)/(\sqrt{b}*(I*b/\operatorname{abs}(b) + 1))} \end{aligned}$$

### 3.19 $\int \frac{\cos^3(a+bx^2)}{x} dx$

**Optimal.** Leaf size=55

$$\frac{3}{8} \cos(a) \text{CosIntegral}(bx^2) + \frac{1}{8} \cos(3a) \text{CosIntegral}(3bx^2) - \frac{3}{8} \sin(a) \text{Si}(bx^2) - \frac{1}{8} \sin(3a) \text{Si}(3bx^2)$$

[Out] (3\*Cos[a]\*CosIntegral[b\*x^2])/8 + (Cos[3\*a]\*CosIntegral[3\*b\*x^2])/8 - (3\*Sin[a]\*SinIntegral[b\*x^2])/8 - (Sin[3\*a]\*SinIntegral[3\*b\*x^2])/8

**Rubi [A]** time = 0.0832228, antiderivative size = 55, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 4, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3404, 3378, 3376, 3375}

$$\frac{3}{8} \cos(a) \text{CosIntegral}(bx^2) + \frac{1}{8} \cos(3a) \text{CosIntegral}(3bx^2) - \frac{3}{8} \sin(a) \text{Si}(bx^2) - \frac{1}{8} \sin(3a) \text{Si}(3bx^2)$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]^3/x,x]

[Out] (3\*Cos[a]\*CosIntegral[b\*x^2])/8 + (Cos[3\*a]\*CosIntegral[3\*b\*x^2])/8 - (3\*Sin[a]\*SinIntegral[b\*x^2])/8 - (Sin[3\*a]\*SinIntegral[3\*b\*x^2])/8

#### Rule 3404

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]

#### Rule 3378

Int[Cos[(c\_) + (d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Dist[Cos[c], Int[Cos[d\*x^n]/x, x], x] - Dist[Sin[c], Int[Sin[d\*x^n]/x, x], x] /; FreeQ[{c, d, n}, x]

#### Rule 3376

Int[Cos[(d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[CosIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

#### Rule 3375

Int[Sin[(d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[SinIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos^3(a + bx^2)}{x} dx &= \int \left( \frac{3 \cos(a + bx^2)}{4x} + \frac{\cos(3a + 3bx^2)}{4x} \right) dx \\
&= \frac{1}{4} \int \frac{\cos(3a + 3bx^2)}{x} dx + \frac{3}{4} \int \frac{\cos(a + bx^2)}{x} dx \\
&= \frac{1}{4} (3 \cos(a)) \int \frac{\cos(bx^2)}{x} dx + \frac{1}{4} \cos(3a) \int \frac{\cos(3bx^2)}{x} dx - \frac{1}{4} (3 \sin(a)) \int \frac{\sin(bx^2)}{x} dx - \frac{1}{4} \sin(3a) \int \frac{\sin(3bx^2)}{x} dx \\
&= \frac{3}{8} \cos(a) \text{Ci}(bx^2) + \frac{1}{8} \cos(3a) \text{Ci}(3bx^2) - \frac{3}{8} \sin(a) \text{Si}(bx^2) - \frac{1}{8} \sin(3a) \text{Si}(3bx^2)
\end{aligned}$$

**Mathematica [A]** time = 0.104799, size = 50, normalized size = 0.91

$$\frac{1}{8} \left( 3 \cos(a) \text{CosIntegral}(bx^2) + \cos(3a) \text{CosIntegral}(3bx^2) - 3 \sin(a) \text{Si}(bx^2) - \sin(3a) \text{Si}(3bx^2) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]^3/x,x]

[Out] (3\*Cos[a]\*CosIntegral[b\*x^2] + Cos[3\*a]\*CosIntegral[3\*b\*x^2] - 3\*Sin[a]\*SinIntegral[b\*x^2] - Sin[3\*a]\*SinIntegral[3\*b\*x^2])/8

**Maple [C]** time = 0.103, size = 125, normalized size = 2.3

$$\frac{i}{16} e^{-3ia} \pi \text{csgn}(bx^2) - \frac{i}{8} e^{-3ia} \text{Si}(3bx^2) - \frac{e^{-3ia} \text{Ei}(1, -3ibx^2)}{16} + \frac{3i}{16} \pi \text{csgn}(bx^2) e^{-ia} - \frac{3i}{8} e^{-ia} \text{Si}(bx^2) - \frac{3e^{-ia} \text{Ei}(1, -ibx^2)}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)^3/x,x)

[Out] 1/16\*I\*exp(-3\*I\*a)\*Pi\*csgn(b\*x^2)-1/8\*I\*exp(-3\*I\*a)\*Si(3\*b\*x^2)-1/16\*exp(-3\*I\*a)\*Ei(1,-3\*I\*b\*x^2)+3/16\*I\*Pi\*csgn(b\*x^2)\*exp(-I\*a)-3/8\*I\*exp(-I\*a)\*Si(b\*x^2)-3/16\*exp(-I\*a)\*Ei(1,-I\*b\*x^2)-3/16\*exp(I\*a)\*Ei(1,-I\*b\*x^2)-1/16\*exp(3\*I\*a)\*Ei(1,-3\*I\*b\*x^2)

**Maxima [C]** time = 1.47111, size = 120, normalized size = 2.18

$$\frac{1}{16} \left( \text{Ei}(3i bx^2) + \text{Ei}(-3i bx^2) \right) \cos(3a) + \frac{3}{16} \left( \text{Ei}(i bx^2) + \text{Ei}(-i bx^2) \right) \cos(a) + \frac{1}{16} \left( i \text{Ei}(3i bx^2) - i \text{Ei}(-3i bx^2) \right) \sin(3a) + \frac{3}{16} \left( i \text{Ei}(i bx^2) - i \text{Ei}(-i bx^2) \right) \sin(a)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3/x,x, algorithm="maxima")

[Out] 1/16\*(Ei(3\*I\*b\*x^2) + Ei(-3\*I\*b\*x^2))\*cos(3\*a) + 3/16\*(Ei(I\*b\*x^2) + Ei(-I\*b\*x^2))\*cos(a) + 1/16\*(I\*Ei(3\*I\*b\*x^2) - I\*Ei(-3\*I\*b\*x^2))\*sin(3\*a) + 1/16\*(3\*I\*Ei(I\*b\*x^2) - 3\*I\*Ei(-I\*b\*x^2))\*sin(a)

**Fricas [A]** time = 1.59335, size = 261, normalized size = 4.75

$$\frac{1}{16} (\text{Ci}(3bx^2) + \text{Ci}(-3bx^2)) \cos(3a) + \frac{3}{16} (\text{Ci}(bx^2) + \text{Ci}(-bx^2)) \cos(a) - \frac{1}{8} \sin(3a) \text{Si}(3bx^2) - \frac{3}{8} \sin(a) \text{Si}(bx^2)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3/x,x, algorithm="fricas")

[Out] 1/16\*(cos\_integral(3\*b\*x^2) + cos\_integral(-3\*b\*x^2))\*cos(3\*a) + 3/16\*(cos\_integral(b\*x^2) + cos\_integral(-b\*x^2))\*cos(a) - 1/8\*sin(3\*a)\*sin\_integral(3\*b\*x^2) - 3/8\*sin(a)\*sin\_integral(b\*x^2)

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^3(a + bx^2)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*3/x,x)

[Out] Integral(cos(a + b\*x\*\*2)\*\*3/x, x)

**Giac [A]** time = 1.16666, size = 63, normalized size = 1.15

$$\frac{1}{8} \cos(3a) \text{Ci}(3bx^2) + \frac{3}{8} \cos(a) \text{Ci}(bx^2) - \frac{3}{8} \sin(a) \text{Si}(bx^2) + \frac{1}{8} \sin(3a) \text{Si}(-3bx^2)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3/x,x, algorithm="giac")

[Out] 1/8\*cos(3\*a)\*cos\_integral(3\*b\*x^2) + 3/8\*cos(a)\*cos\_integral(b\*x^2) - 3/8\*sin(a)\*sin\_integral(b\*x^2) + 1/8\*sin(3\*a)\*sin\_integral(-3\*b\*x^2)

$$3.20 \quad \int \frac{\cos^3(a+bx^2)}{x^2} dx$$

**Optimal.** Leaf size=168

$$-\frac{3}{2}\sqrt{\frac{\pi}{2}}\sqrt{b}\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{bx}\right) - \frac{1}{2}\sqrt{\frac{3\pi}{2}}\sqrt{b}\sin(3a)\text{FresnelC}\left(\sqrt{\frac{6}{\pi}}\sqrt{bx}\right) - \frac{3}{2}\sqrt{\frac{\pi}{2}}\sqrt{b}\cos(a)\text{S}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \frac{1}{2}$$

[Out]  $-(\text{Cos}[a + b*x^2]^3/x) - (3*\text{Sqrt}[b]*\text{Sqrt}[\text{Pi}/2]*\text{Cos}[a]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x])/2 - (\text{Sqrt}[b]*\text{Sqrt}[(3*\text{Pi})/2]*\text{Cos}[3*a]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[6/\text{Pi}]*x])/2 - (3*\text{Sqrt}[b]*\text{Sqrt}[\text{Pi}/2]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x]*\text{Sin}[a])/2 - (\text{Sqrt}[b]*\text{Sqrt}[(3*\text{Pi})/2]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[6/\text{Pi}]*x]*\text{Sin}[3*a])/2$

**Rubi [A]** time = 0.143579, antiderivative size = 168, normalized size of antiderivative = 1., number of steps used = 9, number of rules used = 5, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {3394, 4574, 3353, 3352, 3351}

$$-\frac{3}{2}\sqrt{\frac{\pi}{2}}\sqrt{b}\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{bx}\right) - \frac{1}{2}\sqrt{\frac{3\pi}{2}}\sqrt{b}\sin(3a)\text{FresnelC}\left(\sqrt{\frac{6}{\pi}}\sqrt{bx}\right) - \frac{3}{2}\sqrt{\frac{\pi}{2}}\sqrt{b}\cos(a)\text{S}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \frac{1}{2}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]^3/x^2, x]

[Out]  $-(\text{Cos}[a + b*x^2]^3/x) - (3*\text{Sqrt}[b]*\text{Sqrt}[\text{Pi}/2]*\text{Cos}[a]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x])/2 - (\text{Sqrt}[b]*\text{Sqrt}[(3*\text{Pi})/2]*\text{Cos}[3*a]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[6/\text{Pi}]*x])/2 - (3*\text{Sqrt}[b]*\text{Sqrt}[\text{Pi}/2]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x]*\text{Sin}[a])/2 - (\text{Sqrt}[b]*\text{Sqrt}[(3*\text{Pi})/2]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[6/\text{Pi}]*x]*\text{Sin}[3*a])/2$

#### Rule 3394

Int[Cos[(a\_.) + (b\_.)\*(x\_)^(n\_)^(p\_)]\*(x\_)^(m\_), x\_Symbol] := Simp[(x^(m + 1)\*Cos[a + b\*x^n]^p)/(m + 1), x] + Dist[(b\*n\*p)/(m + 1), Int[Cos[a + b\*x^n]^(p - 1)\*Sin[a + b\*x^n], x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 1] && EqQ[m + n, 0] && NeQ[n, 1] && IntegerQ[n]

#### Rule 4574

Int[Cos[w\_]^(q\_.)\*Sin[v\_]^(p\_.), x\_Symbol] := Int[ExpandTrigReduce[Sin[v]^p \* Cos[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))

#### Rule 3353

Int[Sin[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^2], x\_Symbol] := Dist[Sin[c], Int[Cos[d\*(e + f\*x)^2], x], x] + Dist[Cos[c], Int[Sin[d\*(e + f\*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]

#### Rule 3352

Int[Cos[(d\_.)\*((e\_.) + (f\_.)\*(x\_))^2], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)])/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rule 3351

Int[Sin[(d\_.)\*((e\_.) + (f\_.)\*(x\_))<sup>2</sup>], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)]]/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

### Rubi steps

$$\begin{aligned} \int \frac{\cos^3(a + bx^2)}{x^2} dx &= -\frac{\cos^3(a + bx^2)}{x} - (6b) \int \cos^2(a + bx^2) \sin(a + bx^2) dx \\ &= -\frac{\cos^3(a + bx^2)}{x} - (6b) \int \left( \frac{1}{4} \sin(a + bx^2) + \frac{1}{4} \sin(3a + 3bx^2) \right) dx \\ &= -\frac{\cos^3(a + bx^2)}{x} - \frac{1}{2}(3b) \int \sin(a + bx^2) dx - \frac{1}{2}(3b) \int \sin(3a + 3bx^2) dx \\ &= -\frac{\cos^3(a + bx^2)}{x} - \frac{1}{2}(3b \cos(a)) \int \sin(bx^2) dx - \frac{1}{2}(3b \cos(3a)) \int \sin(3bx^2) dx - \frac{1}{2}(3b \sin(a)) \int \sin(3bx^2) dx \\ &= -\frac{\cos^3(a + bx^2)}{x} - \frac{3}{2} \sqrt{b} \sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right) - \frac{1}{2} \sqrt{b} \sqrt{\frac{3\pi}{2}} \cos(3a) S\left(\sqrt{b} \sqrt{\frac{6}{\pi}} x\right) - \frac{3}{2} \sqrt{b} \sqrt{\frac{\pi}{2}} C\left(\sqrt{b} \sqrt{\frac{2}{\pi}} x\right) \end{aligned}$$

**Mathematica [A]** time = 0.643549, size = 166, normalized size = 0.99

$$\frac{3\sqrt{2\pi}\sqrt{bx} \sin(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{bx}\right) + \sqrt{6\pi}\sqrt{bx} \sin(3a) \text{FresnelC}\left(\sqrt{\frac{6}{\pi}}\sqrt{bx}\right) + 3\sqrt{2\pi}\sqrt{bx} \cos(a) S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) + \sqrt{6\pi}\sqrt{bx} \cos(3a) S\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{4x}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]^3/x^2, x]

[Out]  $-(3 \cos[a + b x^2] + \cos[3(a + b x^2)] + 3 \sqrt{b} \sqrt{2 \pi} x \cos[a] \text{FresnelS}[\sqrt{b} \sqrt{2/\pi} x] + \sqrt{b} \sqrt{6 \pi} x \cos[3 a] \text{FresnelS}[\sqrt{b} \sqrt{6/\pi} x] + 3 \sqrt{b} \sqrt{2 \pi} x \text{FresnelC}[\sqrt{b} \sqrt{2/\pi} x] \sin[a] + \sqrt{b} \sqrt{6 \pi} x \text{FresnelC}[\sqrt{b} \sqrt{6/\pi} x] \sin[3 a]) / (4 x)$

**Maple [A]** time = 0.033, size = 128, normalized size = 0.8

$$-\frac{3 \cos(bx^2 + a)}{4x} - \frac{3 \sqrt{2} \sqrt{\pi}}{4} \sqrt{b} \left( \cos(a) \text{FresnelS}\left(\frac{x \sqrt{2}}{\sqrt{\pi}} \sqrt{b}\right) + \sin(a) \text{FresnelC}\left(\frac{x \sqrt{2}}{\sqrt{\pi}} \sqrt{b}\right) \right) - \frac{\cos(3bx^2 + 3a)}{4x} - \frac{\sqrt{2} \sqrt{\pi}}{4} \sqrt{b} \left( \cos(3a) \text{FresnelS}\left(\frac{x \sqrt{2}}{\sqrt{\pi}} \sqrt{b}\right) + \sin(3a) \text{FresnelC}\left(\frac{x \sqrt{2}}{\sqrt{\pi}} \sqrt{b}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)^3/x^2, x)

[Out]  $-3/4 \cos(b x^2 + a) / x - 3/4 b^{1/2} 2^{1/2} \pi^{1/2} (\cos(a) \text{FresnelS}(x b^{1/2} 2^{1/2} / \pi^{1/2}) + \sin(a) \text{FresnelC}(x b^{1/2} 2^{1/2} / \pi^{1/2})) - 1/4 x \cos(3 b x^2 + 3 a) - 1/4 b^{1/2} 2^{1/2} \pi^{1/2} 3^{1/2} (\cos(3 a) \text{FresnelS}(2^{1/2} / \pi^{1/2} 3^{1/2} b^{1/2} x) + \sin(3 a) \text{FresnelC}(2^{1/2} / \pi^{1/2} 3^{1/2} b^{1/2} x))$

**Maxima [C]** time = 1.38955, size = 721, normalized size = 4.29

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3/x^2,x, algorithm="maxima")

[Out] 
$$\begin{aligned} & -1/32*(\sqrt{3}*\sqrt{x^2*\text{abs}(b)})*((\text{gamma}(-1/2, 3*I*b*x^2) + \text{gamma}(-1/2, -3*I*b*x^2))*\cos(1/4*\pi + 1/2*\arctan2(0, b)) + (\text{gamma}(-1/2, 3*I*b*x^2) + \text{gamma}(-1/2, -3*I*b*x^2))*\cos(-1/4*\pi + 1/2*\arctan2(0, b)) + (I*\text{gamma}(-1/2, 3*I*b*x^2) - I*\text{gamma}(-1/2, -3*I*b*x^2))*\sin(1/4*\pi + 1/2*\arctan2(0, b)) + (-I*\text{gamma}(-1/2, 3*I*b*x^2) + I*\text{gamma}(-1/2, -3*I*b*x^2))*\sin(-1/4*\pi + 1/2*\arctan2(0, b))) * \cos(3*a) + ((-I*\text{gamma}(-1/2, 3*I*b*x^2) + I*\text{gamma}(-1/2, -3*I*b*x^2)) * \cos(1/4*\pi + 1/2*\arctan2(0, b)) + (-I*\text{gamma}(-1/2, 3*I*b*x^2) + I*\text{gamma}(-1/2, -3*I*b*x^2)) * \cos(-1/4*\pi + 1/2*\arctan2(0, b)) + (\text{gamma}(-1/2, 3*I*b*x^2) + \text{gamma}(-1/2, -3*I*b*x^2)) * \sin(1/4*\pi + 1/2*\arctan2(0, b)) - (\text{gamma}(-1/2, 3*I*b*x^2) + \text{gamma}(-1/2, -3*I*b*x^2)) * \sin(-1/4*\pi + 1/2*\arctan2(0, b))) * \sin(3*a) + \sqrt{x^2*\text{abs}(b)} * ((3*(\text{gamma}(-1/2, I*b*x^2) + \text{gamma}(-1/2, -I*b*x^2)) * \cos(1/4*\pi + 1/2*\arctan2(0, b)) + 3*(\text{gamma}(-1/2, I*b*x^2) + \text{gamma}(-1/2, -I*b*x^2)) * \cos(-1/4*\pi + 1/2*\arctan2(0, b)) + (3*I*\text{gamma}(-1/2, I*b*x^2) - 3*I*\text{gamma}(-1/2, -I*b*x^2)) * \sin(1/4*\pi + 1/2*\arctan2(0, b)) + (-3*I*\text{gamma}(-1/2, I*b*x^2) + 3*I*\text{gamma}(-1/2, -I*b*x^2)) * \sin(-1/4*\pi + 1/2*\arctan2(0, b))) * \cos(a) + ((-3*I*\text{gamma}(-1/2, I*b*x^2) + 3*I*\text{gamma}(-1/2, -I*b*x^2)) * \cos(1/4*\pi + 1/2*\arctan2(0, b)) + (-3*I*\text{gamma}(-1/2, I*b*x^2) + 3*I*\text{gamma}(-1/2, -I*b*x^2)) * \cos(-1/4*\pi + 1/2*\arctan2(0, b)) + 3*(\text{gamma}(-1/2, I*b*x^2) + \text{gamma}(-1/2, -I*b*x^2)) * \sin(1/4*\pi + 1/2*\arctan2(0, b)) - 3*(\text{gamma}(-1/2, I*b*x^2) + \text{gamma}(-1/2, -I*b*x^2)) * \sin(-1/4*\pi + 1/2*\arctan2(0, b))) * \sin(a)))/x \end{aligned}$$

---

**Fricas [A]** time = 1.82808, size = 412, normalized size = 2.45

$$\frac{\sqrt{6}\pi x \sqrt{\frac{b}{\pi}} \cos(3a) S\left(\sqrt{6}x \sqrt{\frac{b}{\pi}}\right) + 3\sqrt{2}\pi x \sqrt{\frac{b}{\pi}} \cos(a) S\left(\sqrt{2}x \sqrt{\frac{b}{\pi}}\right) + \sqrt{6}\pi x \sqrt{\frac{b}{\pi}} C\left(\sqrt{6}x \sqrt{\frac{b}{\pi}}\right) \sin(3a) + 3\sqrt{2}\pi x \sqrt{\frac{b}{\pi}} C\left(\sqrt{2}x \sqrt{\frac{b}{\pi}}\right) \sin(a)}{4x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3/x^2,x, algorithm="fricas")

[Out] 
$$\begin{aligned} & -1/4*(\sqrt{6}*\pi*x*\sqrt{b/\pi})*\cos(3*a)*\text{fresnel\_sin}(\sqrt{6}*x*\sqrt{b/\pi}) + 3*\sqrt{2}*\pi*x*\sqrt{b/\pi}*\cos(a)*\text{fresnel\_sin}(\sqrt{2}*x*\sqrt{b/\pi}) + \sqrt{6}*\pi*x*\sqrt{b/\pi}*\text{fresnel\_cos}(\sqrt{6}*x*\sqrt{b/\pi})*\sin(3*a) + 3*\sqrt{2}*\pi*x*\sqrt{b/\pi}*\text{fresnel\_cos}(\sqrt{2}*x*\sqrt{b/\pi})*\sin(a) + 4*\cos(b*x^2 + a)^3)/x \end{aligned}$$

---

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^3(a + bx^2)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*3/x\*\*2,x)

[Out] Integral(cos(a + b\*x\*\*2)\*\*3/x\*\*2, x)

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^2 + a)^3}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x^2+a)^3/x^2,x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^2 + a)^3/x^2, x)
```



## 3.21 $\int \frac{\cos^3(a+bx^2)}{x^3} dx$

**Optimal.** Leaf size=91

$$-\frac{3}{8}b \sin(a) \operatorname{CosIntegral}(bx^2) - \frac{3}{8}b \sin(3a) \operatorname{CosIntegral}(3bx^2) - \frac{3}{8}b \cos(a) \operatorname{Si}(bx^2) - \frac{3}{8}b \cos(3a) \operatorname{Si}(3bx^2) - \frac{3 \cos(a)}{8x}$$

[Out] (-3\*Cos[a + b\*x^2])/(8\*x^2) - Cos[3\*(a + b\*x^2)]/(8\*x^2) - (3\*b\*CosIntegral[b\*x^2]\*Sin[a])/8 - (3\*b\*CosIntegral[3\*b\*x^2]\*Sin[3\*a])/8 - (3\*b\*Cos[a]\*SinIntegral[b\*x^2])/8 - (3\*b\*Cos[3\*a]\*SinIntegral[3\*b\*x^2])/8

**Rubi [A]** time = 0.201348, antiderivative size = 91, normalized size of antiderivative = 1., number of steps used = 12, number of rules used = 6, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3404, 3380, 3297, 3303, 3299, 3302}

$$-\frac{3}{8}b \sin(a) \operatorname{CosIntegral}(bx^2) - \frac{3}{8}b \sin(3a) \operatorname{CosIntegral}(3bx^2) - \frac{3}{8}b \cos(a) \operatorname{Si}(bx^2) - \frac{3}{8}b \cos(3a) \operatorname{Si}(3bx^2) - \frac{3 \cos(a)}{8x}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]^3/x^3, x]

[Out] (-3\*Cos[a + b\*x^2])/(8\*x^2) - Cos[3\*(a + b\*x^2)]/(8\*x^2) - (3\*b\*CosIntegral[b\*x^2]\*Sin[a])/8 - (3\*b\*CosIntegral[3\*b\*x^2]\*Sin[3\*a])/8 - (3\*b\*Cos[a]\*SinIntegral[b\*x^2])/8 - (3\*b\*Cos[3\*a]\*SinIntegral[3\*b\*x^2])/8

### Rule 3404

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_)\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] :> Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]

### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

### Rule 3297

Int[((c\_.) + (d\_.)\*(x\_))^(m\_)\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] :> Simp[((c + d\*x)^(m + 1)\*Sin[e + f\*x])/(d\*(m + 1)), x] - Dist[f/(d\*(m + 1)), Int[(c + d\*x)^(m + 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

### Rule 3303

Int[sin[(e\_.) + (f\_.)\*(x\_)]/((c\_.) + (d\_.)\*(x\_)), x\_Symbol] :> Dist[Cos[(d\*e - c\*f)/d], Int[Sin[(c\*f)/d + f\*x]/(c + d\*x), x], x] + Dist[Sin[(d\*e - c\*f)/d], Int[Cos[(c\*f)/d + f\*x]/(c + d\*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d\*e - c\*f, 0]

### Rule 3299

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinInte
gral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

### Rule 3302

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosInte
gral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) -
c*f, 0]
```

### Rubi steps

$$\begin{aligned} \int \frac{\cos^3(a + bx^2)}{x^3} dx &= \int \left( \frac{3 \cos(a + bx^2)}{4x^3} + \frac{\cos(3a + 3bx^2)}{4x^3} \right) dx \\ &= \frac{1}{4} \int \frac{\cos(3a + 3bx^2)}{x^3} dx + \frac{3}{4} \int \frac{\cos(a + bx^2)}{x^3} dx \\ &= \frac{1}{8} \text{Subst} \left( \int \frac{\cos(3a + 3bx)}{x^2} dx, x, x^2 \right) + \frac{3}{8} \text{Subst} \left( \int \frac{\cos(a + bx)}{x^2} dx, x, x^2 \right) \\ &= -\frac{3 \cos(a + bx^2)}{8x^2} - \frac{\cos(3(a + bx^2))}{8x^2} - \frac{1}{8}(3b) \text{Subst} \left( \int \frac{\sin(a + bx)}{x} dx, x, x^2 \right) - \frac{1}{8}(3b) \text{Subst} \left( \int \frac{\sin(3a + 3bx)}{x} dx, x, x^2 \right) \\ &= -\frac{3 \cos(a + bx^2)}{8x^2} - \frac{\cos(3(a + bx^2))}{8x^2} - \frac{1}{8}(3b \cos(a)) \text{Subst} \left( \int \frac{\sin(bx)}{x} dx, x, x^2 \right) - \frac{1}{8}(3b \cos(3a)) \text{Subst} \left( \int \frac{\sin(3bx)}{x} dx, x, x^2 \right) \\ &= -\frac{3 \cos(a + bx^2)}{8x^2} - \frac{\cos(3(a + bx^2))}{8x^2} - \frac{3}{8} b \text{Ci}(bx^2) \sin(a) - \frac{3}{8} b \text{Ci}(3bx^2) \sin(3a) - \frac{3}{8} b \cos(a) \text{Si}(bx^2) - \frac{3}{8} b \cos(3a) \text{Si}(3bx^2) \end{aligned}$$

**Mathematica [A]** time = 0.175254, size = 90, normalized size = 0.99

$$\frac{3bx^2 \sin(a) \text{CosIntegral}(bx^2) + 3bx^2 \sin(3a) \text{CosIntegral}(3bx^2) + 3bx^2 \cos(a) \text{Si}(bx^2) + 3bx^2 \cos(3a) \text{Si}(3bx^2) + 3 \cos(a) \text{Si}(bx^2) + 3 \cos(3a) \text{Si}(3bx^2)}{8x^2}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x^2]^3/x^3, x]
```

```
[Out] -(3*Cos[a + b*x^2] + Cos[3*(a + b*x^2)] + 3*b*x^2*CosIntegral[b*x^2]*Sin[a]
+ 3*b*x^2*CosIntegral[3*b*x^2]*Sin[3*a] + 3*b*x^2*Cos[a]*SinIntegral[b*x^2]
] + 3*b*x^2*Cos[3*a]*SinIntegral[3*b*x^2])/(8*x^2)
```

**Maple [C]** time = 0.147, size = 162, normalized size = 1.8

$$\frac{3e^{-3ia} \pi \text{csgn}(bx^2) b}{16} - \frac{3e^{-3ia} \text{Si}(3bx^2) b}{8} + \frac{3i}{16} e^{-3ia} \text{Ei}(1, -3ibx^2) b + \frac{3\pi \text{csgn}(bx^2) e^{-ia} b}{16} - \frac{3e^{-ia} \text{Si}(bx^2) b}{8} + \frac{3i}{16} e^{-ia} \text{Ei}(1, -ibx^2) b$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(b*x^2+a)^3/x^3, x)
```

```
[Out] 3/16*exp(-3*I*a)*Pi*csgn(b*x^2)*b-3/8*exp(-3*I*a)*Si(3*b*x^2)*b+3/16*I*exp(-
3*I*a)*Ei(1, -3*I*b*x^2)*b+3/16*Pi*csgn(b*x^2)*exp(-I*a)*b-3/8*exp(-I*a)*Si
(b*x^2)*b+3/16*I*exp(-I*a)*Ei(1, -I*b*x^2)*b-3/16*I*exp(I*a)*b*Ei(1, -I*b*x^2
)-3/16*I*exp(3*I*a)*b*Ei(1, -3*I*b*x^2)-3/8*cos(b*x^2+a)/x^2-1/8*cos(3*b*x^2
```

$+3*a)/x^2$

**Maxima [C]** time = 1.53945, size = 132, normalized size = 1.45

$$-\frac{1}{16} \left( (3i\Gamma(-1, 3i bx^2) - 3i\Gamma(-1, -3i bx^2)) \cos(3a) + (3i\Gamma(-1, i bx^2) - 3i\Gamma(-1, -i bx^2)) \cos(a) + 3(\Gamma(-1, 3i bx^2) + \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3/x^3,x, algorithm="maxima")

[Out]  $-1/16*((3*I*\gamma(-1, 3*I*b*x^2) - 3*I*\gamma(-1, -3*I*b*x^2))*\cos(3*a) + (3*I*\gamma(-1, I*b*x^2) - 3*I*\gamma(-1, -I*b*x^2))*\cos(a) + 3*(\gamma(-1, 3*I*b*x^2) + \gamma(-1, -3*I*b*x^2))*\sin(3*a) + 3*(\gamma(-1, I*b*x^2) + \gamma(-1, -I*b*x^2))*\sin(a))*b$

**Fricas [A]** time = 1.68307, size = 340, normalized size = 3.74

$$\frac{6bx^2 \cos(3a) \operatorname{Si}(3bx^2) + 6bx^2 \cos(a) \operatorname{Si}(bx^2) + 8 \cos(bx^2 + a)^3 + 3(bx^2 \operatorname{Ci}(3bx^2) + bx^2 \operatorname{Ci}(-3bx^2)) \sin(3a) + 3(bx^2 \operatorname{Ci}(bx^2) + bx^2 \operatorname{Ci}(-bx^2)) \sin(a)}{16x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3/x^3,x, algorithm="fricas")

[Out]  $-1/16*(6*b*x^2*\cos(3*a)*\sin\_integral(3*b*x^2) + 6*b*x^2*\cos(a)*\sin\_integral(b*x^2) + 8*\cos(b*x^2 + a)^3 + 3*(b*x^2*\cos\_integral(3*b*x^2) + b*x^2*\cos\_integral(-3*b*x^2))*\sin(3*a) + 3*(b*x^2*\cos\_integral(b*x^2) + b*x^2*\cos\_integral(-b*x^2))*\sin(a))/x^2$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^3(a + bx^2)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*3/x\*\*3,x)

[Out] Integral(cos(a + b\*x\*\*2)\*\*3/x\*\*3, x)

**Giac [B]** time = 1.17793, size = 250, normalized size = 2.75

$$\frac{3(bx^2 + a)b^2 \operatorname{Ci}(3bx^2) \sin(3a) - 3ab^2 \operatorname{Ci}(3bx^2) \sin(3a) + 3(bx^2 + a)b^2 \operatorname{Ci}(bx^2) \sin(a) - 3ab^2 \operatorname{Ci}(bx^2) \sin(a) + 3(bx^2 + a)b^2 \operatorname{Ci}(-3bx^2) \sin(3a) - 3ab^2 \operatorname{Ci}(-3bx^2) \sin(3a) + 3(bx^2 + a)b^2 \operatorname{Ci}(-bx^2) \sin(a) - 3ab^2 \operatorname{Ci}(-bx^2) \sin(a)}{16x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^3/x^3,x, algorithm="giac")

```
[Out] -1/8*(3*(b*x^2 + a)*b^2*cos_integral(3*b*x^2)*sin(3*a) - 3*a*b^2*cos_integr  
al(3*b*x^2)*sin(3*a) + 3*(b*x^2 + a)*b^2*cos_integral(b*x^2)*sin(a) - 3*a*b  
^2*cos_integral(b*x^2)*sin(a) + 3*(b*x^2 + a)*b^2*cos(a)*sin_integral(b*x^2  
) - 3*a*b^2*cos(a)*sin_integral(b*x^2) - 3*(b*x^2 + a)*b^2*cos(3*a)*sin_int  
egral(-3*b*x^2) + 3*a*b^2*cos(3*a)*sin_integral(-3*b*x^2) + b^2*cos(3*b*x^2  
+ 3*a) + 3*b^2*cos(b*x^2 + a))/(b^2*x^2)
```

### 3.22 $\int x \cos^7(a + bx^2) dx$

**Optimal.** Leaf size=67

$$-\frac{\sin^7(a + bx^2)}{14b} + \frac{3 \sin^5(a + bx^2)}{10b} - \frac{\sin^3(a + bx^2)}{2b} + \frac{\sin(a + bx^2)}{2b}$$

[Out] Sin[a + b\*x^2]/(2\*b) - Sin[a + b\*x^2]^3/(2\*b) + (3\*Sin[a + b\*x^2]^5)/(10\*b) - Sin[a + b\*x^2]^7/(14\*b)

**Rubi [A]** time = 0.0537244, antiderivative size = 67, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 2, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3380, 2633}

$$-\frac{\sin^7(a + bx^2)}{14b} + \frac{3 \sin^5(a + bx^2)}{10b} - \frac{\sin^3(a + bx^2)}{2b} + \frac{\sin(a + bx^2)}{2b}$$

Antiderivative was successfully verified.

[In] Int[x\*Cos[a + b\*x^2]^7,x]

[Out] Sin[a + b\*x^2]/(2\*b) - Sin[a + b\*x^2]^3/(2\*b) + (3\*Sin[a + b\*x^2]^5)/(10\*b) - Sin[a + b\*x^2]^7/(14\*b)

#### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

#### Rule 2633

Int[sin[(c\_.) + (d\_.)\*(x\_)]^(n\_), x\_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d\*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

#### Rubi steps

$$\begin{aligned} \int x \cos^7(a + bx^2) dx &= \frac{1}{2} \text{Subst} \left( \int \cos^7(a + bx) dx, x, x^2 \right) \\ &= -\frac{\text{Subst} \left( \int (1 - 3x^2 + 3x^4 - x^6) dx, x, -\sin(a + bx^2) \right)}{2b} \\ &= \frac{\sin(a + bx^2)}{2b} - \frac{\sin^3(a + bx^2)}{2b} + \frac{3 \sin^5(a + bx^2)}{10b} - \frac{\sin^7(a + bx^2)}{14b} \end{aligned}$$

**Mathematica [A]** time = 0.0878097, size = 54, normalized size = 0.81

$$\frac{-5 \sin^7(a + bx^2) + 21 \sin^5(a + bx^2) - 35 \sin^3(a + bx^2) + 35 \sin(a + bx^2)}{70b}$$

Antiderivative was successfully verified.

[In] Integrate[x\*Cos[a + b\*x^2]^7,x]

[Out] (35\*Sin[a + b\*x^2] - 35\*Sin[a + b\*x^2]^3 + 21\*Sin[a + b\*x^2]^5 - 5\*Sin[a + b\*x^2]^7)/(70\*b)

**Maple [A]** time = 0.028, size = 50, normalized size = 0.8

$$\frac{\sin(bx^2 + a)}{14b} \left( \frac{16}{5} + (\cos(bx^2 + a))^6 + \frac{6(\cos(bx^2 + a))^4}{5} + \frac{8(\cos(bx^2 + a))^2}{5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x\*cos(b\*x^2+a)^7,x)

[Out] 1/14/b\*(16/5\*cos(b\*x^2+a)^6+6/5\*cos(b\*x^2+a)^4+8/5\*cos(b\*x^2+a)^2)\*sin(b\*x^2+a)

**Maxima [A]** time = 1.17826, size = 74, normalized size = 1.1

$$\frac{5 \sin(7bx^2 + 7a) + 49 \sin(5bx^2 + 5a) + 245 \sin(3bx^2 + 3a) + 1225 \sin(bx^2 + a)}{4480b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x^2+a)^7,x, algorithm="maxima")

[Out] 1/4480\*(5\*sin(7\*b\*x^2 + 7\*a) + 49\*sin(5\*b\*x^2 + 5\*a) + 245\*sin(3\*b\*x^2 + 3\*a) + 1225\*sin(b\*x^2 + a))/b

**Fricas [A]** time = 1.70206, size = 123, normalized size = 1.84

$$\frac{(5 \cos(bx^2 + a))^6 + 6 \cos(bx^2 + a)^4 + 8 \cos(bx^2 + a)^2 + 16) \sin(bx^2 + a)}{70b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x^2+a)^7,x, algorithm="fricas")

[Out] 1/70\*(5\*cos(b\*x^2 + a)^6 + 6\*cos(b\*x^2 + a)^4 + 8\*cos(b\*x^2 + a)^2 + 16)\*sin(b\*x^2 + a)/b

**Sympy [A]** time = 17.4554, size = 94, normalized size = 1.4

$$\begin{cases} \frac{8 \sin^7(a+bx^2)}{35b} + \frac{4 \sin^5(a+bx^2) \cos^2(a+bx^2)}{5b} + \frac{\sin^3(a+bx^2) \cos^4(a+bx^2)}{b} + \frac{\sin(a+bx^2) \cos^6(a+bx^2)}{2b} & \text{for } b \neq 0 \\ \frac{x^2 \cos^7(a)}{2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x\*\*2+a)\*\*7,x)

[Out] Piecewise((8\*sin(a + b\*x\*\*2)\*\*7/(35\*b) + 4\*sin(a + b\*x\*\*2)\*\*5\*cos(a + b\*x\*\*2)\*\*2/(5\*b) + sin(a + b\*x\*\*2)\*\*3\*cos(a + b\*x\*\*2)\*\*4/b + sin(a + b\*x\*\*2)\*cos(a + b\*x\*\*2)\*\*6/(2\*b), Ne(b, 0)), (x\*\*2\*cos(a)\*\*7/2, True))

**Giac [A]** time = 1.27403, size = 70, normalized size = 1.04

$$\frac{5 \sin(bx^2 + a)^7 - 21 \sin(bx^2 + a)^5 + 35 \sin(bx^2 + a)^3 - 35 \sin(bx^2 + a)}{70b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(b\*x^2+a)^7,x, algorithm="giac")

[Out] -1/70\*(5\*sin(b\*x^2 + a)^7 - 21\*sin(b\*x^2 + a)^5 + 35\*sin(b\*x^2 + a)^3 - 35\*sin(b\*x^2 + a))/b

### 3.23 $\int x^{5/2} \cos(a + bx^2) dx$

**Optimal.** Leaf size=111

$$-\frac{3ie^{ia}x^{3/2}\Gamma\left(\frac{3}{4}, -ibx^2\right)}{16b(-ibx^2)^{3/4}} + \frac{3ie^{-ia}x^{3/2}\Gamma\left(\frac{3}{4}, ibx^2\right)}{16b(ibx^2)^{3/4}} + \frac{x^{3/2} \sin(a + bx^2)}{2b}$$

[Out] (((-3\*I)/16)\*E^(I\*a)\*x^(3/2)\*Gamma[3/4, (-I)\*b\*x^2])/(b\*((-I)\*b\*x^2)^(3/4)) + (((3\*I)/16)\*x^(3/2)\*Gamma[3/4, I\*b\*x^2])/(b\*E^(I\*a)\*(I\*b\*x^2)^(3/4)) + (x^(3/2)\*Sin[a + b\*x^2])/(2\*b)

**Rubi [A]** time = 0.0811548, antiderivative size = 111, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3386, 3389, 2218}

$$-\frac{3ie^{ia}x^{3/2}\Gamma\left(\frac{3}{4}, -ibx^2\right)}{16b(-ibx^2)^{3/4}} + \frac{3ie^{-ia}x^{3/2}\Gamma\left(\frac{3}{4}, ibx^2\right)}{16b(ibx^2)^{3/4}} + \frac{x^{3/2} \sin(a + bx^2)}{2b}$$

Antiderivative was successfully verified.

[In] Int[x^(5/2)\*Cos[a + b\*x^2], x]

[Out] (((-3\*I)/16)\*E^(I\*a)\*x^(3/2)\*Gamma[3/4, (-I)\*b\*x^2])/(b\*((-I)\*b\*x^2)^(3/4)) + (((3\*I)/16)\*x^(3/2)\*Gamma[3/4, I\*b\*x^2])/(b\*E^(I\*a)\*(I\*b\*x^2)^(3/4)) + (x^(3/2)\*Sin[a + b\*x^2])/(2\*b)

#### Rule 3386

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)]\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] := Simp[(e^(n - 1)\*(e\*x)^(m - n + 1)\*Sin[c + d\*x^n])/(d\*n), x] - Dist[(e^n\*(m - n + 1))/(d\*n), Int[(e\*x)^(m - n)\*Sin[c + d\*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[n, m + 1]

#### Rule 3389

Int[((e\_.)\*(x\_))^(m\_.)\*Sin[(c\_.) + (d\_.)\*(x\_)^(n\_)], x\_Symbol] := Dist[I/2, Int[(e\*x)^m\*E^(-(c\*I) - d\*I\*x^n), x], x] - Dist[I/2, Int[(e\*x)^m\*E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]

#### Rule 2218

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_))^(n\_))\*((e\_.) + (f\_.)\*(x\_))^(m\_.), x\_Symbol] := -Simp[(F^a\*(e + f\*x)^(m + 1)\*Gamma[(m + 1)/n, -(b\*(c + d\*x)^n\*Log[F])])/(f\*n\*(-(b\*(c + d\*x)^n\*Log[F]))^((m + 1)/n)), x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d\*e - c\*f, 0]

#### Rubi steps



$$\begin{aligned}
\int x^{5/2} \cos(a + bx^2) dx &= \frac{x^{3/2} \sin(a + bx^2)}{2b} - \frac{3 \int \sqrt{x} \sin(a + bx^2) dx}{4b} \\
&= \frac{x^{3/2} \sin(a + bx^2)}{2b} - \frac{(3i) \int e^{-ia-ibx^2} \sqrt{x} dx}{8b} + \frac{(3i) \int e^{ia+ibx^2} \sqrt{x} dx}{8b} \\
&= -\frac{3ie^{ia} x^{3/2} \Gamma\left(\frac{3}{4}, -ibx^2\right)}{16b(-ibx^2)^{3/4}} + \frac{3ie^{-ia} x^{3/2} \Gamma\left(\frac{3}{4}, ibx^2\right)}{16b(ibx^2)^{3/4}} + \frac{x^{3/2} \sin(a + bx^2)}{2b}
\end{aligned}$$

**Mathematica [A]** time = 0.201149, size = 113, normalized size = 1.02

$$\frac{bx^{11/2} \left( 3 (ibx^2)^{3/4} (\sin(a) - i \cos(a)) \Gamma\left(\frac{3}{4}, -ibx^2\right) + 3 (-ibx^2)^{3/4} (\sin(a) + i \cos(a)) \Gamma\left(\frac{3}{4}, ibx^2\right) + 8 (b^2 x^4)^{7/4} \right)}{16 (b^2 x^4)^{7/4}}$$

Antiderivative was successfully verified.

[In] Integrate[x^(5/2)\*Cos[a + b\*x^2], x]

[Out] (b\*x^(11/2)\*(3\*(I\*b\*x^2)^(3/4)\*Gamma[3/4, (-I)\*b\*x^2]\*((-I)\*Cos[a] + Sin[a]) + 3\*((-I)\*b\*x^2)^(3/4)\*Gamma[3/4, I\*b\*x^2]\*(I\*Cos[a] + Sin[a]) + 8\*(b^2\*x^4)^(3/4)\*Sin[a + b\*x^2]))/(16\*(b^2\*x^4)^(7/4))

**Maple [C]** time = 0.142, size = 229, normalized size = 2.1

$$\frac{2^{\frac{3}{4}} \cos(a) \sqrt{\pi} \left( \frac{2 \sqrt[4]{2} \sin(bx^2)}{7 \sqrt{\pi} b} x^{\frac{3}{2}} (b^2)^{\frac{7}{8}} + \frac{3 \sqrt[4]{2} \sin(bx^2)}{14 \sqrt{\pi}} x^{\frac{7}{2}} (b^2)^{\frac{7}{8}} \text{LommelS1}\left(\frac{5}{4}, \frac{3}{2}, bx^2\right) (bx^2)^{-\frac{5}{4}} + \frac{3 \sqrt[4]{2} (\cos(bx^2) b)}{8 \sqrt{\pi}} \right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(5/2)\*cos(b\*x^2+a), x)

[Out] 1/2\*2^(3/4)/(b^2)^(7/8)\*cos(a)\*Pi^(1/2)\*(2/7/Pi^(1/2)\*x^(3/2)\*2^(1/4)\*(b^2)^(7/8)\*sin(b\*x^2)/b+3/14/Pi^(1/2)\*x^(7/2)\*(b^2)^(7/8)\*2^(1/4)/(b\*x^2)^(5/4)\*sin(b\*x^2)\*LommelS1(5/4, 3/2, b\*x^2)+3/8/Pi^(1/2)\*x^(7/2)\*(b^2)^(7/8)\*2^(1/4)/(b\*x^2)^(9/4)\*(cos(b\*x^2)\*b\*x^2-sin(b\*x^2))\*LommelS1(1/4, 1/2, b\*x^2))-1/2\*2^(3/4)/b^(7/4)\*sin(a)\*Pi^(1/2)\*(-1/8/Pi^(1/2)\*x^(7/2)\*b^(7/4)\*2^(1/4)/(b\*x^2)^(5/4)\*sin(b\*x^2)\*LommelS1(1/4, 3/2, b\*x^2)-1/2/Pi^(1/2)\*x^(7/2)\*b^(7/4)\*2^(1/4)/(b\*x^2)^(9/4)\*(cos(b\*x^2)\*b\*x^2-sin(b\*x^2))\*LommelS1(5/4, 1/2, b\*x^2))

**Maxima [B]** time = 1.47386, size = 397, normalized size = 3.58

$$\frac{16 x^2 |b| \sin(bx^2 + a) - (x^2 |b|)^{\frac{1}{4}} \left( \left( -3i \Gamma\left(\frac{3}{4}, ibx^2\right) + 3i \Gamma\left(\frac{3}{4}, -ibx^2\right) \right) \cos\left(\frac{3}{8} \pi + \frac{3}{4} \arctan(0, b)\right) + \left( -3i \Gamma\left(\frac{3}{4}, ibx^2\right) + \right) \right)}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(5/2)\*cos(b\*x^2+a), x, algorithm="maxima")

```
[Out] 1/32*(16*x^2*abs(b)*sin(b*x^2 + a) - (x^2*abs(b))^(1/4)*((-3*I*gamma(3/4,
I*b*x^2) + 3*I*gamma(3/4, -I*b*x^2))*cos(3/8*pi + 3/4*arctan2(0, b)) + (-3*
I*gamma(3/4, I*b*x^2) + 3*I*gamma(3/4, -I*b*x^2))*cos(-3/8*pi + 3/4*arctan2
(0, b)) - 3*(gamma(3/4, I*b*x^2) + gamma(3/4, -I*b*x^2))*sin(3/8*pi + 3/4*a
rctan2(0, b)) + 3*(gamma(3/4, I*b*x^2) + gamma(3/4, -I*b*x^2))*sin(-3/8*pi
+ 3/4*arctan2(0, b)))*cos(a) - (3*(gamma(3/4, I*b*x^2) + gamma(3/4, -I*b*x^
2))*cos(3/8*pi + 3/4*arctan2(0, b)) + 3*(gamma(3/4, I*b*x^2) + gamma(3/4, -
I*b*x^2))*cos(-3/8*pi + 3/4*arctan2(0, b)) - (3*I*gamma(3/4, I*b*x^2) - 3*I
*gamma(3/4, -I*b*x^2))*sin(3/8*pi + 3/4*arctan2(0, b)) - (-3*I*gamma(3/4, I
*b*x^2) + 3*I*gamma(3/4, -I*b*x^2))*sin(-3/8*pi + 3/4*arctan2(0, b)))*sin(a
))/ (b*sqrt(x)*abs(b))
```

**Fricas [A]** time = 1.70188, size = 176, normalized size = 1.59

$$\frac{8bx^{\frac{3}{2}}\sin(bx^2 + a) + 3(ib)^{\frac{1}{4}}e^{(-ia)}\Gamma\left(\frac{3}{4}, ibx^2\right) + 3(-ib)^{\frac{1}{4}}e^{(ia)}\Gamma\left(\frac{3}{4}, -ibx^2\right)}{16b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^(5/2)*cos(b*x^2+a),x, algorithm="fricas")
```

```
[Out] 1/16*(8*b*x^(3/2)*sin(b*x^2 + a) + 3*(I*b)^(1/4)*e^(-I*a)*gamma(3/4, I*b*x^
2) + 3*(-I*b)^(1/4)*e^(I*a)*gamma(3/4, -I*b*x^2))/b^2
```

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**(5/2)*cos(b*x**2+a),x)
```

```
[Out] Timed out
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{\frac{5}{2}} \cos(bx^2 + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^(5/2)*cos(b*x^2+a),x, algorithm="giac")
```

```
[Out] integrate(x^(5/2)*cos(b*x^2 + a), x)
```

### 3.24 $\int x^{3/2} \cos(a + bx^2) dx$

**Optimal.** Leaf size=111

$$-\frac{ie^{ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -ibx^2\right)}{16b^4\sqrt{-ibx^2}} + \frac{ie^{-ia}\sqrt{x}\Gamma\left(\frac{1}{4}, ibx^2\right)}{16b^4\sqrt{ibx^2}} + \frac{\sqrt{x}\sin(a + bx^2)}{2b}$$

[Out]  $((-I/16)*E^{(I*a)}*\text{Sqrt}[x]*\Gamma[1/4, (-I)*b*x^2])/(b*((-I)*b*x^2)^{(1/4)}) + ((I/16)*\text{Sqrt}[x]*\Gamma[1/4, I*b*x^2])/(b*E^{(I*a)}*(I*b*x^2)^{(1/4)}) + (\text{Sqrt}[x]*\text{Sin}[a + b*x^2])/(2*b)$

**Rubi [A]** time = 0.0786466, antiderivative size = 111, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3386, 3389, 2218}

$$-\frac{ie^{ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -ibx^2\right)}{16b^4\sqrt{-ibx^2}} + \frac{ie^{-ia}\sqrt{x}\Gamma\left(\frac{1}{4}, ibx^2\right)}{16b^4\sqrt{ibx^2}} + \frac{\sqrt{x}\sin(a + bx^2)}{2b}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[x^{(3/2)}*\text{Cos}[a + b*x^2], x]$

[Out]  $((-I/16)*E^{(I*a)}*\text{Sqrt}[x]*\Gamma[1/4, (-I)*b*x^2])/(b*((-I)*b*x^2)^{(1/4)}) + ((I/16)*\text{Sqrt}[x]*\Gamma[1/4, I*b*x^2])/(b*E^{(I*a)}*(I*b*x^2)^{(1/4)}) + (\text{Sqrt}[x]*\text{Sin}[a + b*x^2])/(2*b)$

#### Rule 3386

$\text{Int}[\text{Cos}[(c_.) + (d_.)*(x_.)^{(n_.)}]*((e_.)*(x_.))^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[(e^{(n-1)}*(e*x)^{(m-n+1)}*\text{Sin}[c + d*x^n])/(d*n), x] - \text{Dist}[(e^n*(m-n+1))/(d*n), \text{Int}[(e*x)^{(m-n)}*\text{Sin}[c + d*x^n], x], x] /; \text{FreeQ}\{c, d, e\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{LtQ}[n, m + 1]$

#### Rule 3389

$\text{Int}[(e_.)*(x_.)^{(m_.)}*\text{Sin}[(c_.) + (d_.)*(x_.)^{(n_.)}], x\_Symbol] \rightarrow \text{Dist}[I/2, \text{Int}[(e*x)^m * E^{-(c*I) - d*I*x^n}, x], x] - \text{Dist}[I/2, \text{Int}[(e*x)^m * E^{(c*I) + d*I*x^n}, x], x] /; \text{FreeQ}\{c, d, e, m\}, x] \&\& \text{IGtQ}[n, 0]$

#### Rule 2218

$\text{Int}[(F_.)^{((a_.) + (b_.)*((c_.) + (d_.)*(x_.))^{(n_.)})}*((e_.) + (f_.)*(x_.))^{(m_.)}, x\_Symbol] \rightarrow -\text{Simp}[(F^a*(e + f*x)^{(m+1)}*\Gamma[(m+1)/n, -(b*(c + d*x)^n*\text{Log}[F])])/(f*n*(-(b*(c + d*x)^n*\text{Log}[F]))^{((m+1)/n)}), x] /; \text{FreeQ}\{F, a, b, c, d, e, f, m, n\}, x] \&\& \text{EqQ}[d*e - c*f, 0]$

#### Rubi steps

$$\begin{aligned} \int x^{3/2} \cos(a + bx^2) dx &= \frac{\sqrt{x} \sin(a + bx^2)}{2b} - \frac{\int \frac{\sin(a+bx^2)}{\sqrt{x}} dx}{4b} \\ &= \frac{\sqrt{x} \sin(a + bx^2)}{2b} - \frac{i \int \frac{e^{-ia-ibx^2}}{\sqrt{x}} dx}{8b} + \frac{i \int \frac{e^{ia+ibx^2}}{\sqrt{x}} dx}{8b} \\ &= -\frac{ie^{ia} \sqrt{x} \Gamma\left(\frac{1}{4}, -ibx^2\right)}{16b \sqrt[4]{-ibx^2}} + \frac{ie^{-ia} \sqrt{x} \Gamma\left(\frac{1}{4}, ibx^2\right)}{16b \sqrt[4]{ibx^2}} + \frac{\sqrt{x} \sin(a + bx^2)}{2b} \end{aligned}$$

**Mathematica [A]** time = 0.157765, size = 111, normalized size = 1.

$$\frac{bx^{9/2} \left( \sqrt[4]{ibx^2} (\sin(a) - i \cos(a)) \Gamma\left(\frac{1}{4}, -ibx^2\right) + \sqrt[4]{-ibx^2} (\sin(a) + i \cos(a)) \Gamma\left(\frac{1}{4}, ibx^2\right) + 8 \sqrt[4]{b^2 x^4} \sin(a + bx^2) \right)}{16 (b^2 x^4)^{5/4}}$$

Antiderivative was successfully verified.

[In] Integrate[x^(3/2)\*Cos[a + b\*x^2],x]

[Out] (b\*x^(9/2)\*((I\*b\*x^2)^(1/4)\*Gamma[1/4, (-I)\*b\*x^2]\*((-I)\*Cos[a] + Sin[a]) + ((-I)\*b\*x^2)^(1/4)\*Gamma[1/4, I\*b\*x^2]\*(I\*Cos[a] + Sin[a]) + 8\*(b^2\*x^4)^(1/4)\*Sin[a + b\*x^2]))/(16\*(b^2\*x^4)^(5/4))

**Maple [C]** time = 0.077, size = 290, normalized size = 2.6

$$\frac{\sqrt[4]{2} \cos(a) \sqrt{\pi} \left( \frac{2 \cdot 2^{3/4} \sin(bx^2)}{5 \sqrt{\pi} b} \sqrt{x} (b^2)^{5/8} + \frac{2 \cdot 2^{3/4} (\cos(bx^2) bx^2 - \sin(bx^2))}{5 \sqrt{\pi} b} \sqrt{x} (b^2)^{5/8} + \frac{2^{3/4} b \sin(bx^2)}{10 \sqrt{\pi}} x^{9/2} (b^2)^{5/8} \right)}{2} \text{LommelS1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(3/2)\*cos(b\*x^2+a),x)

[Out] 1/2\*2^(1/4)/(b^2)^(5/8)\*cos(a)\*Pi^(1/2)\*(2/5/Pi^(1/2)\*x^(1/2)\*2^(3/4)\*(b^2)^(5/8)\*sin(b\*x^2)/b+2/5/Pi^(1/2)\*x^(1/2)\*2^(3/4)\*(b^2)^(5/8)/b\*(cos(b\*x^2)\*b\*x^2-sin(b\*x^2))+1/10/Pi^(1/2)\*x^(9/2)\*(b^2)^(5/8)\*2^(3/4)\*b/(b\*x^2)^(7/4)\*sin(b\*x^2)\*LommelS1(3/4,3/2,b\*x^2)-2/5/Pi^(1/2)\*x^(9/2)\*(b^2)^(5/8)\*2^(3/4)\*b/(b\*x^2)^(11/4)\*(cos(b\*x^2)\*b\*x^2-sin(b\*x^2))\*LommelS1(7/4,1/2,b\*x^2))-1/2\*2^(1/4)/b^(5/4)\*sin(a)\*Pi^(1/2)\*(2/9/Pi^(1/2)\*x^(5/2)\*2^(3/4)\*b^(5/4)\*sin(b\*x^2)-2/9/Pi^(1/2)\*x^(9/2)\*b^(9/4)\*2^(3/4)/(b\*x^2)^(7/4)\*sin(b\*x^2)\*LommelS1(7/4,3/2,b\*x^2)-1/2/Pi^(1/2)\*x^(9/2)\*b^(9/4)\*2^(3/4)/(b\*x^2)^(11/4)\*(cos(b\*x^2)\*b\*x^2-sin(b\*x^2))\*LommelS1(3/4,1/2,b\*x^2))

**Maxima [B]** time = 1.52483, size = 390, normalized size = 3.51

$$16 (x^2|b|)^{1/4} \sqrt{x} \sin(bx^2 + a) + \left( \left( i \Gamma\left(\frac{1}{4}, ibx^2\right) - i \Gamma\left(\frac{1}{4}, -ibx^2\right) \right) \cos\left(\frac{1}{8} \pi + \frac{1}{4} \arctan(0, b)\right) + \left( i \Gamma\left(\frac{1}{4}, ibx^2\right) - i \Gamma\left(\frac{1}{4}, -ibx^2\right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(b\*x^2+a),x, algorithm="maxima")

[Out]  $\frac{1}{32} \cdot (16 \cdot (x^2 \cdot \text{abs}(b))^{1/4}) \cdot \sqrt{x} \cdot \sin(b \cdot x^2 + a) + (((I \cdot \text{gamma}(1/4, I \cdot b \cdot x^2) - I \cdot \text{gamma}(1/4, -I \cdot b \cdot x^2)) \cdot \cos(1/8 \cdot \pi + 1/4 \cdot \arctan2(0, b)) + (I \cdot \text{gamma}(1/4, I \cdot b \cdot x^2) - I \cdot \text{gamma}(1/4, -I \cdot b \cdot x^2)) \cdot \cos(-1/8 \cdot \pi + 1/4 \cdot \arctan2(0, b)) + (\text{gamma}(1/4, I \cdot b \cdot x^2) + \text{gamma}(1/4, -I \cdot b \cdot x^2)) \cdot \sin(1/8 \cdot \pi + 1/4 \cdot \arctan2(0, b)) - (\text{gamma}(1/4, I \cdot b \cdot x^2) + \text{gamma}(1/4, -I \cdot b \cdot x^2)) \cdot \sin(-1/8 \cdot \pi + 1/4 \cdot \arctan2(0, b))) \cdot \cos(a) + ((\text{gamma}(1/4, I \cdot b \cdot x^2) + \text{gamma}(1/4, -I \cdot b \cdot x^2)) \cdot \cos(1/8 \cdot \pi + 1/4 \cdot \arctan2(0, b)) + (\text{gamma}(1/4, I \cdot b \cdot x^2) + \text{gamma}(1/4, -I \cdot b \cdot x^2)) \cdot \cos(-1/8 \cdot \pi + 1/4 \cdot \arctan2(0, b)) + (-I \cdot \text{gamma}(1/4, I \cdot b \cdot x^2) + I \cdot \text{gamma}(1/4, -I \cdot b \cdot x^2)) \cdot \sin(1/8 \cdot \pi + 1/4 \cdot \arctan2(0, b)) + (I \cdot \text{gamma}(1/4, I \cdot b \cdot x^2) - I \cdot \text{gamma}(1/4, -I \cdot b \cdot x^2)) \cdot \sin(-1/8 \cdot \pi + 1/4 \cdot \arctan2(0, b))) \cdot \sin(a) \cdot \sqrt{x}) / ((x^2 \cdot \text{abs}(b))^{1/4} \cdot b)$

**Fricas [A]** time = 1.75403, size = 170, normalized size = 1.53

$$\frac{(ib)^{\frac{3}{4}} e^{(-ia)} \Gamma\left(\frac{1}{4}, ibx^2\right) + (-ib)^{\frac{3}{4}} e^{(ia)} \Gamma\left(\frac{1}{4}, -ibx^2\right) + 8b\sqrt{x} \sin(bx^2 + a)}{16b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(b\*x^2+a),x, algorithm="fricas")

[Out]  $\frac{1}{16} \cdot ((I \cdot b)^{3/4} \cdot e^{-I \cdot a} \cdot \text{gamma}(1/4, I \cdot b \cdot x^2) + (-I \cdot b)^{3/4} \cdot e^{I \cdot a} \cdot \text{gamma}(1/4, -I \cdot b \cdot x^2) + 8 \cdot b \cdot \sqrt{x} \cdot \sin(b \cdot x^2 + a)) / b^2$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{\frac{3}{2}} \cos(a + bx^2) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*(3/2)\*cos(b\*x\*\*2+a),x)

[Out] Integral(x\*\*(3/2)\*cos(a + b\*x\*\*2), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{\frac{3}{2}} \cos(bx^2 + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(b\*x^2+a),x, algorithm="giac")

[Out] integrate(x^(3/2)\*cos(b\*x^2 + a), x)

### 3.25 $\int \sqrt{x} \cos(a + bx^2) dx$

**Optimal.** Leaf size=81

$$-\frac{e^{ia}x^{3/2}\Gamma\left(\frac{3}{4}, -ibx^2\right)}{4(-ibx^2)^{3/4}} - \frac{e^{-ia}x^{3/2}\Gamma\left(\frac{3}{4}, ibx^2\right)}{4(ibx^2)^{3/4}}$$

[Out]  $-(E^{(I*a)}*x^{(3/2)}*\Gamma[3/4, (-I)*b*x^2])/(4*((-I)*b*x^2)^{(3/4)}) - (x^{(3/2)}*\Gamma[3/4, I*b*x^2])/(4*E^{(I*a)}*(I*b*x^2)^{(3/4)})$

**Rubi [A]** time = 0.0629466, antiderivative size = 81, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 2, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {3390, 2218}

$$-\frac{e^{ia}x^{3/2}\Gamma\left(\frac{3}{4}, -ibx^2\right)}{4(-ibx^2)^{3/4}} - \frac{e^{-ia}x^{3/2}\Gamma\left(\frac{3}{4}, ibx^2\right)}{4(ibx^2)^{3/4}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[x]\*Cos[a + b\*x^2], x]

[Out]  $-(E^{(I*a)}*x^{(3/2)}*\Gamma[3/4, (-I)*b*x^2])/(4*((-I)*b*x^2)^{(3/4)}) - (x^{(3/2)}*\Gamma[3/4, I*b*x^2])/(4*E^{(I*a)}*(I*b*x^2)^{(3/4)})$

#### Rule 3390

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_.)]\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] := Dist[1/2, Int[(e\*x)^m\*E^(-c\*I - d\*I\*x^n), x], x] + Dist[1/2, Int[(e\*x)^m\*E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]

#### Rule 2218

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_))^(n\_.))\*((e\_.) + (f\_.)\*(x\_))^(m\_.), x\_Symbol] := -Simp[(F^a\*(e + f\*x)^(m + 1)\*Gamma[(m + 1)/n, -(b\*(c + d\*x)^n\*Log[F])])/(f\*n\*(-(b\*(c + d\*x)^n\*Log[F]))^(m + 1)/n), x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d\*e - c\*f, 0]

#### Rubi steps

$$\begin{aligned} \int \sqrt{x} \cos(a + bx^2) dx &= \frac{1}{2} \int e^{-ia-ibx^2} \sqrt{x} dx + \frac{1}{2} \int e^{ia+ibx^2} \sqrt{x} dx \\ &= -\frac{e^{ia}x^{3/2}\Gamma\left(\frac{3}{4}, -ibx^2\right)}{4(-ibx^2)^{3/4}} - \frac{e^{-ia}x^{3/2}\Gamma\left(\frac{3}{4}, ibx^2\right)}{4(ibx^2)^{3/4}} \end{aligned}$$

**Mathematica [A]** time = 0.0898811, size = 89, normalized size = 1.1

$$\frac{x^{3/2} \left( (-ibx^2)^{3/4} (\cos(a) - i \sin(a)) \Gamma\left(\frac{3}{4}, ibx^2\right) + (ibx^2)^{3/4} (\cos(a) + i \sin(a)) \Gamma\left(\frac{3}{4}, -ibx^2\right) \right)}{4(b^2x^4)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[x]\*Cos[a + b\*x^2], x]

[Out]  $-(x^{3/2} * (((-1) * b * x^2)^{3/4} * \Gamma[3/4, I * b * x^2] * (\cos[a] - I * \sin[a]) + (I * b * x^2)^{3/4} * \Gamma[3/4, (-1) * b * x^2] * (\cos[a] + I * \sin[a]))) / (4 * (b^2 * x^4)^{3/4})$

**Maple [C]** time = 0.066, size = 290, normalized size = 3.6

$$\frac{2^{\frac{3}{4}} \cos(a) \sqrt{\pi}}{4} \left( \frac{4 \sqrt[4]{2} \sin(bx^2)}{3 \sqrt{\pi} b} (b^2)^{\frac{3}{8}} \frac{1}{\sqrt{x}} + \frac{4 \sqrt[4]{2} (\cos(bx^2) bx^2 - \sin(bx^2))}{3 \sqrt{\pi} b} (b^2)^{\frac{3}{8}} \frac{1}{\sqrt{x}} - \frac{\sqrt[4]{2} b \sin(bx^2)}{3 \sqrt{\pi}} x^{\frac{7}{2}} (b^2)^{\frac{3}{8}} \text{LommelS1} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(1/2)\*cos(b\*x^2+a), x)

[Out]  $\frac{1}{4} * 2^{3/4} / (b^2)^{3/8} * \cos(a) * \pi^{1/2} * (4/3 / \pi^{1/2} / x^{1/2} * 2^{1/4} * (b^2)^{3/8} * \sin(bx^2) / b + 4/3 / \pi^{1/2} / x^{1/2} * 2^{1/4} * (b^2)^{3/8} / b * (\cos(bx^2) * bx^2 - \sin(bx^2)) - 1/3 / \pi^{1/2} * x^{7/2} * (b^2)^{3/8} * 2^{1/4} * b / (bx^2)^{5/4} * \sin(bx^2) * \text{LommelS1}(1/4, 3/2, bx^2) - 4/3 / \pi^{1/2} * x^{7/2} * (b^2)^{3/8} * 2^{1/4} * b / (bx^2)^{9/4} * (\cos(bx^2) * bx^2 - \sin(bx^2)) * \text{LommelS1}(5/4, 1/2, bx^2)) - 1/4 * 2^{3/4} / b^{3/4} * \sin(a) * \pi^{1/2} * (4/7 / \pi^{1/2} * x^{3/2} * 2^{1/4} * b^{3/4} * \sin(bx^2) - 4/7 / \pi^{1/2} * x^{7/2} * b^{7/4} * 2^{1/4} / (bx^2)^{5/4} * \sin(bx^2) * \text{LommelS1}(5/4, 3/2, bx^2) - 1/\pi^{1/2} * x^{7/2} * b^{7/4} * 2^{1/4} / (bx^2)^{9/4} * (\cos(bx^2) * bx^2 - \sin(bx^2)) * \text{LommelS1}(1/4, 1/2, bx^2))$

**Maxima [B]** time = 1.6082, size = 365, normalized size = 4.51

$$(x^2|b|)^{\frac{1}{4}} \left( \left( \Gamma\left(\frac{3}{4}, ibx^2\right) + \Gamma\left(\frac{3}{4}, -ibx^2\right) \right) \cos\left(\frac{3}{8}\pi + \frac{3}{4}\arctan(0, b)\right) + \left( \Gamma\left(\frac{3}{4}, ibx^2\right) + \Gamma\left(\frac{3}{4}, -ibx^2\right) \right) \cos\left(-\frac{3}{8}\pi + \frac{3}{4}\arctan(0, b)\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(1/2)\*cos(b\*x^2+a), x, algorithm="maxima")

[Out]  $-1/8 * (x^2 * \text{abs}(b))^{1/4} * (((\gamma(3/4, I * b * x^2) + \gamma(3/4, -I * b * x^2)) * \cos(3/8 * \pi + 3/4 * \arctan2(0, b)) + (\gamma(3/4, I * b * x^2) + \gamma(3/4, -I * b * x^2)) * \cos(-3/8 * \pi + 3/4 * \arctan2(0, b)) - (I * \gamma(3/4, I * b * x^2) - I * \gamma(3/4, -I * b * x^2)) * \sin(3/8 * \pi + 3/4 * \arctan2(0, b)) - (-I * \gamma(3/4, I * b * x^2) + I * \gamma(3/4, -I * b * x^2)) * \sin(-3/8 * \pi + 3/4 * \arctan2(0, b))) * \cos(a) - ((I * \gamma(3/4, I * b * x^2) - I * \gamma(3/4, -I * b * x^2)) * \cos(3/8 * \pi + 3/4 * \arctan2(0, b)) + (I * \gamma(3/4, I * b * x^2) + \gamma(3/4, -I * b * x^2)) * \cos(-3/8 * \pi + 3/4 * \arctan2(0, b)) + (\gamma(3/4, I * b * x^2) + \gamma(3/4, -I * b * x^2)) * \sin(3/8 * \pi + 3/4 * \arctan2(0, b)) - (\gamma(3/4, I * b * x^2) + \gamma(3/4, -I * b * x^2)) * \sin(-3/8 * \pi + 3/4 * \arctan2(0, b))) * \sin(a)) / (\text{sqrt}(x) * \text{abs}(b))$

**Fricas [A]** time = 1.7082, size = 132, normalized size = 1.63

$$\frac{i (ib)^{\frac{1}{4}} e^{(-ia)} \Gamma\left(\frac{3}{4}, ibx^2\right) - i (-ib)^{\frac{1}{4}} e^{(ia)} \Gamma\left(\frac{3}{4}, -ibx^2\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^(1/2)*cos(b*x^2+a),x, algorithm="fricas")`

[Out]  $\frac{1}{4} * (I * (I * b)^{(1/4)} * e^{(-I * a)} * \text{gamma}(3/4, I * b * x^2) - I * (-I * b)^{(1/4)} * e^{(I * a)} * \text{gamma}(3/4, -I * b * x^2)) / b$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \sqrt{x} \cos(a + bx^2) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**(1/2)*cos(b*x**2+a),x)`

[Out] `Integral(sqrt(x)*cos(a + b*x**2), x)`

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \sqrt{x} \cos(bx^2 + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^(1/2)*cos(b*x^2+a),x, algorithm="giac")`

[Out] `integrate(sqrt(x)*cos(b*x^2 + a), x)`



$$3.26 \quad \int \frac{\cos(a+bx^2)}{\sqrt{x}} dx$$

**Optimal.** Leaf size=81

$$\frac{e^{ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -ibx^2\right)}{4\sqrt[4]{-ibx^2}} - \frac{e^{-ia}\sqrt{x}\Gamma\left(\frac{1}{4}, ibx^2\right)}{4\sqrt[4]{ibx^2}}$$

[Out]  $-(E^{(I*a)}*\text{Sqrt}[x]*\text{Gamma}[1/4, (-I)*b*x^2])/(4*((-I)*b*x^2)^{(1/4)}) - (\text{Sqrt}[x]*\text{Gamma}[1/4, I*b*x^2])/(4*E^{(I*a)}*(I*b*x^2)^{(1/4)})$

**Rubi [A]** time = 0.0653032, antiderivative size = 81, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 2, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {3390, 2218}

$$\frac{e^{ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -ibx^2\right)}{4\sqrt[4]{-ibx^2}} - \frac{e^{-ia}\sqrt{x}\Gamma\left(\frac{1}{4}, ibx^2\right)}{4\sqrt[4]{ibx^2}}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[\text{Cos}[a + b*x^2]/\text{Sqrt}[x], x]$

[Out]  $-(E^{(I*a)}*\text{Sqrt}[x]*\text{Gamma}[1/4, (-I)*b*x^2])/(4*((-I)*b*x^2)^{(1/4)}) - (\text{Sqrt}[x]*\text{Gamma}[1/4, I*b*x^2])/(4*E^{(I*a)}*(I*b*x^2)^{(1/4)})$

#### Rule 3390

$\text{Int}[\text{Cos}[(c_.) + (d_.)*(x_)^{(n_)}]*((e_.)*(x_))^{(m_.)}, x\_Symbol] \rightarrow \text{Dist}[1/2, \text{Int}[(e*x)^m * E^{-(c*I) - d*I*x^n}, x], x] + \text{Dist}[1/2, \text{Int}[(e*x)^m * E^{(c*I) + d*I*x^n}, x], x] /;$  FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]

#### Rule 2218

$\text{Int}[(F_)^{((a_.) + (b_.)*((c_.) + (d_.)*(x_))^{(n_)})*((e_.) + (f_.)*(x_))^{(m_.)}, x\_Symbol] \rightarrow -\text{Simp}[(F^a*(e + f*x)^{(m+1)}*\text{Gamma}[(m+1)/n, -(b*(c + d*x)^n*\text{Log}[F])])]/(f*n*(-(b*(c + d*x)^n*\text{Log}[F]))^{((m+1)/n)}, x] /;$  FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d\*e - c\*f, 0]

#### Rubi steps

$$\begin{aligned} \int \frac{\cos(a+bx^2)}{\sqrt{x}} dx &= \frac{1}{2} \int \frac{e^{-ia-ibx^2}}{\sqrt{x}} dx + \frac{1}{2} \int \frac{e^{ia+ibx^2}}{\sqrt{x}} dx \\ &= -\frac{e^{ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -ibx^2\right)}{4\sqrt[4]{-ibx^2}} - \frac{e^{-ia}\sqrt{x}\Gamma\left(\frac{1}{4}, ibx^2\right)}{4\sqrt[4]{ibx^2}} \end{aligned}$$

**Mathematica [A]** time = 0.0714116, size = 89, normalized size = 1.1

$$\frac{\sqrt{x}\left(\sqrt[4]{-ibx^2}(\cos(a) - i\sin(a))\Gamma\left(\frac{1}{4}, ibx^2\right) + \sqrt[4]{ibx^2}(\cos(a) + i\sin(a))\Gamma\left(\frac{1}{4}, -ibx^2\right)\right)}{4\sqrt[4]{b^2x^4}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]/Sqrt[x], x]

[Out]  $-(\text{Sqrt}[x]*((( -I)*b*x^2)^{1/4}*\text{Gamma}[1/4, I*b*x^2]*(\text{Cos}[a] - I*\text{Sin}[a]) + (I*b*x^2)^{1/4}*\text{Gamma}[1/4, (-I)*b*x^2]*(\text{Cos}[a] + I*\text{Sin}[a])))/(4*(b^2*x^4)^{1/4})$

**Maple [C]** time = 0.078, size = 338, normalized size = 4.2

$$\frac{\cos(a)\sqrt{\pi}\sqrt[4]{2}}{4} \left( 6 \frac{2^{3/4}\sqrt[8]{b^2} \sin(bx^2)}{\sqrt{\pi}x^{3/2}b} \left( \frac{8b^2x^4}{27} + 2/3 \right) + 4 \frac{2^{3/4}\sqrt[8]{b^2} (\cos(bx^2)bx^2 - \sin(bx^2))}{\sqrt{\pi}x^{3/2}b} - \frac{16b^22^{3/4} \sin(bx^2)}{9\sqrt{\pi}} x^{9/2} \sqrt[8]{b^2} \text{L} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)/x^(1/2), x)

[Out]  $1/4*\cos(a)*\text{Pi}^{1/2}*2^{1/4}/(b^2)^{1/8}*(6/\text{Pi}^{1/2}/x^{3/2}*2^{3/4}*(b^2)^{1/8}*(8/27*b^2*x^4+2/3)*\sin(b*x^2)/b+4/\text{Pi}^{1/2}/x^{3/2}*2^{3/4}*(b^2)^{1/8}/b*(\cos(b*x^2)*b*x^2-\sin(b*x^2))-16/9/\text{Pi}^{1/2}*x^{9/2}*(b^2)^{1/8}*b^2*2^{3/4}/(b*x^2)^{7/4}*\sin(b*x^2)*\text{LommelS1}(7/4, 3/2, b*x^2)-4/\text{Pi}^{1/2}*x^{9/2}*(b^2)^{1/8}*b^2*2^{3/4}/(b*x^2)^{11/4}*(\cos(b*x^2)*b*x^2-\sin(b*x^2))*\text{LommelS1}(3/4, 1/2, b*x^2))-1/4*\sin(a)*\text{Pi}^{1/2}*2^{1/4}/b^{1/4}*(4/5/\text{Pi}^{1/2}*x^{1/2}*2^{3/4}*b^{1/4}*\sin(b*x^2)-16/5/\text{Pi}^{1/2}*x^{1/2}*2^{3/4}*b^{1/4}*(\cos(b*x^2)*b*x^2-\sin(b*x^2))-4/5/\text{Pi}^{1/2}*x^{9/2}*b^{9/4}*2^{3/4}/(b*x^2)^{7/4}*\sin(b*x^2)*\text{LommelS1}(3/4, 3/2, b*x^2)+16/5/\text{Pi}^{1/2}*x^{9/2}*b^{9/4}*2^{3/4}/(b*x^2)^{11/4}*(\cos(b*x^2)*b*x^2-\sin(b*x^2))*\text{LommelS1}(7/4, 1/2, b*x^2))$

**Maxima [B]** time = 1.44696, size = 359, normalized size = 4.43

$$\left( \left( \Gamma\left(\frac{1}{4}, ibx^2\right) + \Gamma\left(\frac{1}{4}, -ibx^2\right) \right) \cos\left(\frac{1}{8}\pi + \frac{1}{4}\arctan(0, b)\right) + \left( \Gamma\left(\frac{1}{4}, ibx^2\right) + \Gamma\left(\frac{1}{4}, -ibx^2\right) \right) \cos\left(-\frac{1}{8}\pi + \frac{1}{4}\arctan(0, b)\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^(1/2), x, algorithm="maxima")

[Out]  $-1/8*((\text{gamma}(1/4, I*b*x^2) + \text{gamma}(1/4, -I*b*x^2))*\cos(1/8*\text{pi} + 1/4*\arctan(0, b)) + (\text{gamma}(1/4, I*b*x^2) + \text{gamma}(1/4, -I*b*x^2))*\cos(-1/8*\text{pi} + 1/4*\arctan(0, b)) - (I*\text{gamma}(1/4, I*b*x^2) - I*\text{gamma}(1/4, -I*b*x^2))*\sin(1/8*\text{pi} + 1/4*\arctan(0, b)) - (-I*\text{gamma}(1/4, I*b*x^2) + I*\text{gamma}(1/4, -I*b*x^2))*\sin(-1/8*\text{pi} + 1/4*\arctan(0, b)))*\cos(a) - ((I*\text{gamma}(1/4, I*b*x^2) - I*\text{gamma}(1/4, -I*b*x^2))*\cos(1/8*\text{pi} + 1/4*\arctan(0, b)) + (I*\text{gamma}(1/4, I*b*x^2) - I*\text{gamma}(1/4, -I*b*x^2))*\cos(-1/8*\text{pi} + 1/4*\arctan(0, b)) + (\text{gamma}(1/4, I*b*x^2) + \text{gamma}(1/4, -I*b*x^2))*\sin(1/8*\text{pi} + 1/4*\arctan(0, b)) - (\text{gamma}(1/4, I*b*x^2) + \text{gamma}(1/4, -I*b*x^2))*\sin(-1/8*\text{pi} + 1/4*\arctan(0, b)))*\sin(a))*\text{sqrt}(x)/(x^2*\text{abs}(b))^{1/4}$

**Fricas [A]** time = 1.71051, size = 132, normalized size = 1.63

$$\frac{i (i b)^{\frac{3}{4}} e^{(-i a)} \Gamma\left(\frac{1}{4}, i b x^2\right) - i (-i b)^{\frac{3}{4}} e^{(i a)} \Gamma\left(\frac{1}{4}, -i b x^2\right)}{4 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^(1/2), x, algorithm="fricas")

[Out] 1/4\*(I\*(I\*b)^(3/4)\*e^(-I\*a)\*gamma(1/4, I\*b\*x^2) - I\*(-I\*b)^(3/4)\*e^(I\*a)\*gamma(1/4, -I\*b\*x^2))/b

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + b x^2)}{\sqrt{x}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)/x\*\*(1/2), x)

[Out] Integral(cos(a + b\*x\*\*2)/sqrt(x), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(b x^2 + a)}{\sqrt{x}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^(1/2), x, algorithm="giac")

[Out] integrate(cos(b\*x^2 + a)/sqrt(x), x)

$$3.27 \quad \int \frac{\cos(a+bx^2)}{x^{3/2}} dx$$

**Optimal.** Leaf size=98

$$-\frac{ie^{ia}bx^{3/2}\Gamma\left(\frac{3}{4}, -ibx^2\right)}{(-ibx^2)^{3/4}} + \frac{ie^{-ia}bx^{3/2}\Gamma\left(\frac{3}{4}, ibx^2\right)}{(ibx^2)^{3/4}} - \frac{2\cos(a+bx^2)}{\sqrt{x}}$$

[Out] (-2\*Cos[a + b\*x^2])/Sqrt[x] - (I\*b\*E^(I\*a)\*x^(3/2)\*Gamma[3/4, (-I)\*b\*x^2])/((-I)\*b\*x^2)^(3/4) + (I\*b\*x^(3/2)\*Gamma[3/4, I\*b\*x^2])/(E^(I\*a)\*(I\*b\*x^2)^(3/4))

**Rubi [A]** time = 0.0758502, antiderivative size = 98, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3388, 3389, 2218}

$$-\frac{ie^{ia}bx^{3/2}\Gamma\left(\frac{3}{4}, -ibx^2\right)}{(-ibx^2)^{3/4}} + \frac{ie^{-ia}bx^{3/2}\Gamma\left(\frac{3}{4}, ibx^2\right)}{(ibx^2)^{3/4}} - \frac{2\cos(a+bx^2)}{\sqrt{x}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]/x^(3/2), x]

[Out] (-2\*Cos[a + b\*x^2])/Sqrt[x] - (I\*b\*E^(I\*a)\*x^(3/2)\*Gamma[3/4, (-I)\*b\*x^2])/((-I)\*b\*x^2)^(3/4) + (I\*b\*x^(3/2)\*Gamma[3/4, I\*b\*x^2])/(E^(I\*a)\*(I\*b\*x^2)^(3/4))

#### Rule 3388

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)]\*((e\_.)\*(x\_))^(m\_), x\_Symbol] := Simp[((e\*x)^(m + 1)\*Cos[c + d\*x^n])/(e\*(m + 1)), x] + Dist[(d\*n)/(e^n\*(m + 1)), Int[(e\*x)^(m + n)\*Sin[c + d\*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[m, -1]

#### Rule 3389

Int[((e\_.)\*(x\_))^(m\_.)\*Sin[(c\_.) + (d\_.)\*(x\_)^(n\_)], x\_Symbol] := Dist[I/2, Int[(e\*x)^m\*E^(-(c\*I) - d\*I\*x^n), x], x] - Dist[I/2, Int[(e\*x)^m\*E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]

#### Rule 2218

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_)^(n\_)))\*((e\_.) + (f\_.)\*(x\_))^(m\_.), x\_Symbol] := -Simp[(F^a\*(e + f\*x)^(m + 1)\*Gamma[(m + 1)/n, -(b\*(c + d\*x)^n\*Log[F])])/(f\*n\*(-(b\*(c + d\*x)^n\*Log[F]))^((m + 1)/n)), x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d\*e - c\*f, 0]

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos(a + bx^2)}{x^{3/2}} dx &= -\frac{2 \cos(a + bx^2)}{\sqrt{x}} - (4b) \int \sqrt{x} \sin(a + bx^2) dx \\
&= -\frac{2 \cos(a + bx^2)}{\sqrt{x}} - (2ib) \int e^{-ia-ibx^2} \sqrt{x} dx + (2ib) \int e^{ia+ibx^2} \sqrt{x} dx \\
&= -\frac{2 \cos(a + bx^2)}{\sqrt{x}} - \frac{ibe^{ia} x^{3/2} \Gamma\left(\frac{3}{4}, -ibx^2\right)}{(-ibx^2)^{3/4}} + \frac{ibe^{-ia} x^{3/2} \Gamma\left(\frac{3}{4}, ibx^2\right)}{(ibx^2)^{3/4}}
\end{aligned}$$

**Mathematica [A]** time = 0.182243, size = 114, normalized size = 1.16

$$\frac{bx^2 (ibx^2)^{3/4} (\sin(a) - i \cos(a)) \Gamma\left(\frac{3}{4}, -ibx^2\right) + i (-ibx^2)^{7/4} (\sin(a) + i \cos(a)) \Gamma\left(\frac{3}{4}, ibx^2\right) - 2 (b^2 x^4)^{3/4} \cos(a)}{\sqrt{x} (b^2 x^4)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]/x^(3/2), x]

[Out]  $(-2*(b^2*x^4)^{(3/4)}*\cos[a + b*x^2] + b*x^2*(I*b*x^2)^{(3/4)}*\Gamma[3/4, (-I)*b*x^2]*((-I)*\cos[a] + \sin[a]) + I*((-I)*b*x^2)^{(7/4)}*\Gamma[3/4, I*b*x^2]*(I*\cos[a] + \sin[a]))/(\text{Sqrt}[x]*(b^2*x^4)^{(3/4)})$

**Maple [C]** time = 0.067, size = 338, normalized size = 3.5

$$\frac{\cos(a) \sqrt{\pi} 2^{\frac{3}{4}} \sqrt[8]{b^2} \left( -12 \frac{\sqrt[4]{2} \sin(bx^2)}{\sqrt{\pi} x^{5/2} \sqrt[8]{b^2} b} \left( \frac{8 b^2 x^4}{21} + 2/3 \right) - 8 \frac{\sqrt[4]{2} (\cos(bx^2) bx^2 - \sin(bx^2))}{\sqrt{\pi} x^{5/2} \sqrt[8]{b^2} b} + \frac{32 b^2 \sqrt[4]{2} \sin(bx^2)}{7 \sqrt{\pi}} x^{\frac{7}{2}} \text{LommelS1} \right)}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)/x^(3/2), x)

[Out]  $1/8*\cos(a)*\pi^{(1/2)}*2^{(3/4)}*(b^2)^{(1/8)}*(-12/\pi^{(1/2)}/x^{(5/2)}*2^{(1/4)}/(b^2)^{(1/8)}*(8/21*b^2*x^4+2/3)*\sin(b*x^2)/b-8/\pi^{(1/2)}/x^{(5/2)}*2^{(1/4)}/(b^2)^{(1/8)}/b*(\cos(b*x^2)*b*x^2-\sin(b*x^2))+32/7/\pi^{(1/2)}*x^{(7/2)}/(b^2)^{(1/8)}*b^2*2^{(1/4)}/(b*x^2)^{(5/4)}*\sin(b*x^2)*\text{LommelS1}(5/4,3/2,b*x^2)+8/\pi^{(1/2)}*x^{(7/2)}/(b^2)^{(1/8)}*b^2*2^{(1/4)}/(b*x^2)^{(9/4)}*(\cos(b*x^2)*b*x^2-\sin(b*x^2))*\text{LommelS1}(1/4,1/2,b*x^2)-1/8*\sin(a)*\pi^{(1/2)}*2^{(3/4)}*b^{(1/4)}*(8/3/\pi^{(1/2)}/x^{(1/2)}*2^{(1/4)}/b^{(1/4)}*\sin(b*x^2)+32/3/\pi^{(1/2)}/x^{(1/2)}*2^{(1/4)}/b^{(1/4)}*(\cos(b*x^2)*b*x^2-\sin(b*x^2))-8/3/\pi^{(1/2)}*x^{(7/2)}*b^{(7/4)}*2^{(1/4)}/(b*x^2)^{(5/4)}*\sin(b*x^2)*\text{LommelS1}(1/4,3/2,b*x^2)-32/3/\pi^{(1/2)}*x^{(7/2)}*b^{(7/4)}*2^{(1/4)}/(b*x^2)^{(9/4)}*(\cos(b*x^2)*b*x^2-\sin(b*x^2))*\text{LommelS1}(5/4,1/2,b*x^2))$

**Maxima [B]** time = 1.41277, size = 355, normalized size = 3.62

$$(x^2|b|)^{\frac{1}{4}} \left( \left( \Gamma\left(-\frac{1}{4}, ibx^2\right) + \Gamma\left(-\frac{1}{4}, -ibx^2\right) \right) \cos\left(\frac{1}{8}\pi + \frac{1}{4}\arctan(0, b)\right) + \left( \Gamma\left(-\frac{1}{4}, ibx^2\right) + \Gamma\left(-\frac{1}{4}, -ibx^2\right) \right) \cos\left(-\frac{1}{8}\pi + \frac{1}{4}\arctan(0, b)\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^(3/2),x, algorithm="maxima")

[Out] 
$$\frac{-1/8*(x^2*\text{abs}(b))^{1/4}*((\text{gamma}(-1/4, I*b*x^2) + \text{gamma}(-1/4, -I*b*x^2))*\cos(1/8*\text{pi} + 1/4*\text{arctan2}(0, b)) + (\text{gamma}(-1/4, I*b*x^2) + \text{gamma}(-1/4, -I*b*x^2))*\cos(-1/8*\text{pi} + 1/4*\text{arctan2}(0, b)) + (I*\text{gamma}(-1/4, I*b*x^2) - I*\text{gamma}(-1/4, -I*b*x^2))*\sin(1/8*\text{pi} + 1/4*\text{arctan2}(0, b)) + (-I*\text{gamma}(-1/4, I*b*x^2) + I*\text{gamma}(-1/4, -I*b*x^2))*\sin(-1/8*\text{pi} + 1/4*\text{arctan2}(0, b)))*\cos(a) + ((-I*\text{gamma}(-1/4, I*b*x^2) + I*\text{gamma}(-1/4, -I*b*x^2))*\cos(1/8*\text{pi} + 1/4*\text{arctan2}(0, b)) + (-I*\text{gamma}(-1/4, I*b*x^2) + I*\text{gamma}(-1/4, -I*b*x^2))*\cos(-1/8*\text{pi} + 1/4*\text{arctan2}(0, b)) + (\text{gamma}(-1/4, I*b*x^2) + \text{gamma}(-1/4, -I*b*x^2))*\sin(1/8*\text{pi} + 1/4*\text{arctan2}(0, b)) - (\text{gamma}(-1/4, I*b*x^2) + \text{gamma}(-1/4, -I*b*x^2))*\sin(-1/8*\text{pi} + 1/4*\text{arctan2}(0, b)))*\sin(a))/\text{sqrt}(x)}$$

**Fricas [A]** time = 1.69499, size = 163, normalized size = 1.66

$$\frac{(ib)^{\frac{1}{4}} x e^{(-ia)} \Gamma\left(\frac{3}{4}, ibx^2\right) + (-ib)^{\frac{1}{4}} x e^{(ia)} \Gamma\left(\frac{3}{4}, -ibx^2\right) - 2\sqrt{x} \cos(bx^2 + a)}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^(3/2),x, algorithm="fricas")

[Out] 
$$\frac{((I*b)^{1/4}*x*e^{(-I*a)}*\text{gamma}(3/4, I*b*x^2) + (-I*b)^{1/4}*x*e^{(I*a)}*\text{gamma}(3/4, -I*b*x^2) - 2*\text{sqrt}(x)*\cos(b*x^2 + a))/x}$$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + bx^2)}{x^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)/x\*\*(3/2),x)

[Out] Integral(cos(a + b\*x\*\*2)/x\*\*(3/2), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^2 + a)}{x^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^(3/2),x, algorithm="giac")

[Out] integrate(cos(b\*x^2 + a)/x^(3/2), x)

$$3.28 \quad \int \frac{\cos(a+bx^2)}{x^{5/2}} dx$$

**Optimal.** Leaf size=104

$$-\frac{ie^{iab}\sqrt{x}\Gamma\left(\frac{1}{4}, -ibx^2\right)}{3\sqrt[4]{-ibx^2}} + \frac{ie^{-ia}b\sqrt{x}\Gamma\left(\frac{1}{4}, ibx^2\right)}{3\sqrt[4]{ibx^2}} - \frac{2\cos(a+bx^2)}{3x^{3/2}}$$

[Out]  $(-2*\text{Cos}[a + b*x^2])/(3*x^{(3/2)}) - ((I/3)*b*E^{(I*a)}*\text{Sqrt}[x]*\Gamma[1/4, (-I)*b*x^2])/((-I)*b*x^2)^{(1/4)} + ((I/3)*b*\text{Sqrt}[x]*\Gamma[1/4, I*b*x^2])/(E^{(I*a)}*(I*b*x^2)^{(1/4)})$

**Rubi [A]** time = 0.0748721, antiderivative size = 104, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3388, 3389, 2218}

$$-\frac{ie^{iab}\sqrt{x}\Gamma\left(\frac{1}{4}, -ibx^2\right)}{3\sqrt[4]{-ibx^2}} + \frac{ie^{-ia}b\sqrt{x}\Gamma\left(\frac{1}{4}, ibx^2\right)}{3\sqrt[4]{ibx^2}} - \frac{2\cos(a+bx^2)}{3x^{3/2}}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[\text{Cos}[a + b*x^2]/x^{(5/2)}, x]$

[Out]  $(-2*\text{Cos}[a + b*x^2])/(3*x^{(3/2)}) - ((I/3)*b*E^{(I*a)}*\text{Sqrt}[x]*\Gamma[1/4, (-I)*b*x^2])/((-I)*b*x^2)^{(1/4)} + ((I/3)*b*\text{Sqrt}[x]*\Gamma[1/4, I*b*x^2])/(E^{(I*a)}*(I*b*x^2)^{(1/4)})$

#### Rule 3388

$\text{Int}[\text{Cos}[(c_.) + (d_.)*(x_)^{(n_)}]*((e_.)*(x_))^{(m_)}, x\_Symbol] \rightarrow \text{Simp}[(e*x)^{(m+1)}*\text{Cos}[c + d*x^n]/(e*(m+1)), x] + \text{Dist}[(d*n)/(e^n*(m+1)), \text{Int}[(e*x)^{(m+n)}*\text{Sin}[c + d*x^n], x], x] /; \text{FreeQ}\{c, d, e\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m, -1]$

#### Rule 3389

$\text{Int}[(e_.)*(x_)^{(m_.)}*\text{Sin}[(c_.) + (d_.)*(x_)^{(n_)}], x\_Symbol] \rightarrow \text{Dist}[I/2, \text{Int}[(e*x)^m*E^{-(c*I) - d*I*x^n}, x], x] - \text{Dist}[I/2, \text{Int}[(e*x)^m*E^{(c*I) + d*I*x^n}, x], x] /; \text{FreeQ}\{c, d, e, m\}, x \ \&\& \ \text{IGtQ}[n, 0]$

#### Rule 2218

$\text{Int}[(F_)^{((a_.) + (b_.)*((c_.) + (d_.)*(x_))^{(n_)})*((e_.) + (f_.)*(x_))^{(m_.)}], x\_Symbol] \rightarrow -\text{Simp}[(F^a*(e + f*x)^{(m+1)}*\Gamma[(m+1)/n, -(b*(c + d*x)^n*\text{Log}[F])])]/(f*n*(-(b*(c + d*x)^n*\text{Log}[F]))^{((m+1)/n)}), x] /; \text{FreeQ}\{F, a, b, c, d, e, f, m, n\}, x \ \&\& \ \text{EqQ}[d*e - c*f, 0]$

#### Rubi steps

$$\begin{aligned} \int \frac{\cos(a + bx^2)}{x^{5/2}} dx &= -\frac{2 \cos(a + bx^2)}{3x^{3/2}} - \frac{1}{3}(4b) \int \frac{\sin(a + bx^2)}{\sqrt{x}} dx \\ &= -\frac{2 \cos(a + bx^2)}{3x^{3/2}} - \frac{1}{3}(2ib) \int \frac{e^{-ia-ibx^2}}{\sqrt{x}} dx + \frac{1}{3}(2ib) \int \frac{e^{ia+ibx^2}}{\sqrt{x}} dx \\ &= -\frac{2 \cos(a + bx^2)}{3x^{3/2}} - \frac{ibe^{ia} \sqrt{x} \Gamma\left(\frac{1}{4}, -ibx^2\right)}{3\sqrt[4]{-ibx^2}} + \frac{ibe^{-ia} \sqrt{x} \Gamma\left(\frac{1}{4}, ibx^2\right)}{3\sqrt[4]{ibx^2}} \end{aligned}$$

**Mathematica [A]** time = 0.156479, size = 117, normalized size = 1.12

$$\frac{bx^2 \sqrt[4]{ibx^2} (\sin(a) - i \cos(a)) \Gamma\left(\frac{1}{4}, -ibx^2\right) + i(-ibx^2)^{5/4} (\sin(a) + i \cos(a)) \Gamma\left(\frac{1}{4}, ibx^2\right) - 2\sqrt[4]{b^2 x^4} \cos(a + bx^2)}{3x^{3/2} \sqrt[4]{b^2 x^4}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]/x^(5/2), x]

[Out]  $(-2*(b^2*x^4)^{(1/4)}*\text{Cos}[a + b*x^2] + b*x^2*(I*b*x^2)^{(1/4)}*\text{Gamma}[1/4, (-I)*b*x^2]*((-I)*\text{Cos}[a] + \text{Sin}[a]) + I*((-I)*b*x^2)^{(5/4)}*\text{Gamma}[1/4, I*b*x^2]*(I*\text{Cos}[a] + \text{Sin}[a]))/(3*x^{(3/2)}*(b^2*x^4)^{(1/4)})$

**Maple [C]** time = 0.069, size = 358, normalized size = 3.4

$$\frac{\cos(a) \sqrt{\pi} \sqrt[4]{2}}{8} (b^2)^{\frac{3}{8}} \left( -4 \frac{2^{3/4} \sin(bx^2)}{\sqrt{\pi} x^{7/2} (b^2)^{3/8} b} \left( \frac{8b^2 x^4}{15} + 2/3 \right) - \frac{8 \cdot 2^{3/4} (-16b^2 x^4 + 5) (\cos(bx^2) bx^2 - \sin(bx^2))}{15 \sqrt{\pi} b} x^{-7/2} (b^2)^{-3/8} + \dots \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)/x^(5/2), x)

[Out]  $1/8*\text{cos}(a)*\text{Pi}^{(1/2)}*2^{(1/4)}*(b^2)^{(3/8)}*(-4/\text{Pi}^{(1/2)}/x^{(7/2)}*2^{(3/4)}/(b^2)^{(3/8)}*(8/15*b^2*x^4+2/3)*\text{sin}(b*x^2)/b-8/15/\text{Pi}^{(1/2)}/x^{(7/2)}*2^{(3/4)}/(b^2)^{(3/8)}/b*(-16*b^2*x^4+5)*( \text{cos}(b*x^2)*b*x^2-\text{sin}(b*x^2) )+32/15/\text{Pi}^{(1/2)}*x^{(9/2)}/(b^2)^{(3/8)}*2^{(3/4)}*b^3/(b*x^2)^{(7/4)}*\text{sin}(b*x^2)*\text{LommelS1}(3/4, 3/2, b*x^2)-128/15/\text{Pi}^{(1/2)}*x^{(9/2)}/(b^2)^{(3/8)}*2^{(3/4)}*b^3/(b*x^2)^{(11/4)}*( \text{cos}(b*x^2)*b*x^2-\text{sin}(b*x^2) )*\text{LommelS1}(7/4, 1/2, b*x^2))-1/8*\text{sin}(a)*\text{Pi}^{(1/2)}*2^{(1/4)}*b^{(3/4)}*(12/\text{Pi}^{(1/2)}/x^{(3/2)}*2^{(3/4)}/b^{(3/4)}*(32/81*b^2*x^4+2/3)*\text{sin}(b*x^2)+32/3/\text{Pi}^{(1/2)}/x^{(3/2)}*2^{(3/4)}/b^{(3/4)}*( \text{cos}(b*x^2)*b*x^2-\text{sin}(b*x^2) )-128/27/\text{Pi}^{(1/2)}*x^{(9/2)}*b^{(9/4)}*2^{(3/4)}/(b*x^2)^{(7/4)}*\text{sin}(b*x^2)*\text{LommelS1}(7/4, 3/2, b*x^2)-32/3/\text{Pi}^{(1/2)}*x^{(9/2)}*b^{(9/4)}*2^{(3/4)}/(b*x^2)^{(11/4)}*( \text{cos}(b*x^2)*b*x^2-\text{sin}(b*x^2) )*\text{LommelS1}(3/4, 1/2, b*x^2))$

**Maxima [B]** time = 1.44541, size = 355, normalized size = 3.41

$$(x^2|b|)^{\frac{3}{4}} \left( \left( \Gamma\left(-\frac{3}{4}, ibx^2\right) + \Gamma\left(-\frac{3}{4}, -ibx^2\right) \right) \cos\left(\frac{3}{8}\pi + \frac{3}{4}\arctan(0, b)\right) + \left( \Gamma\left(-\frac{3}{4}, ibx^2\right) + \Gamma\left(-\frac{3}{4}, -ibx^2\right) \right) \cos\left(-\frac{3}{8}\pi + \frac{3}{4}\right)$$

Verification of antiderivative is not currently implemented for this CAS.



[In] integrate(cos(b\*x^2+a)/x^(5/2),x, algorithm="maxima")

[Out] 
$$\frac{-1/8*(x^2*\text{abs}(b))^{3/4}*(((\text{gamma}(-3/4, I*b*x^2) + \text{gamma}(-3/4, -I*b*x^2))*\cos(3/8*\text{pi} + 3/4*\text{arctan2}(0, b)) + (\text{gamma}(-3/4, I*b*x^2) + \text{gamma}(-3/4, -I*b*x^2))*\cos(-3/8*\text{pi} + 3/4*\text{arctan2}(0, b)) + (I*\text{gamma}(-3/4, I*b*x^2) - I*\text{gamma}(-3/4, -I*b*x^2))*\sin(3/8*\text{pi} + 3/4*\text{arctan2}(0, b)) + (-I*\text{gamma}(-3/4, I*b*x^2) + I*\text{gamma}(-3/4, -I*b*x^2))*\sin(-3/8*\text{pi} + 3/4*\text{arctan2}(0, b)))*\cos(a) + ((-I*\text{gamma}(-3/4, I*b*x^2) + I*\text{gamma}(-3/4, -I*b*x^2))*\cos(3/8*\text{pi} + 3/4*\text{arctan2}(0, b)) + (-I*\text{gamma}(-3/4, I*b*x^2) + I*\text{gamma}(-3/4, -I*b*x^2))*\cos(-3/8*\text{pi} + 3/4*\text{arctan2}(0, b)) + (\text{gamma}(-3/4, I*b*x^2) + \text{gamma}(-3/4, -I*b*x^2))*\sin(3/8*\text{pi} + 3/4*\text{arctan2}(0, b)) - (\text{gamma}(-3/4, I*b*x^2) + \text{gamma}(-3/4, -I*b*x^2))*\sin(-3/8*\text{pi} + 3/4*\text{arctan2}(0, b)))*\sin(a))}{x^{3/2}}$$

**Fricas [A]** time = 1.78771, size = 177, normalized size = 1.7

$$\frac{(ib)^{\frac{3}{4}}x^2e^{-ia}\Gamma\left(\frac{1}{4}, ibx^2\right) + (-ib)^{\frac{3}{4}}x^2e^{ia}\Gamma\left(\frac{1}{4}, -ibx^2\right) - 2\sqrt{x}\cos(bx^2 + a)}{3x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^(5/2),x, algorithm="fricas")

[Out] 
$$\frac{1/3*((I*b)^{3/4}*x^2*e^{-I*a}*\text{gamma}(1/4, I*b*x^2) + (-I*b)^{3/4}*x^2*e^{I*a}*\text{gamma}(1/4, -I*b*x^2) - 2*\text{sqrt}(x)*\cos(b*x^2 + a))}{x^2}$$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + bx^2)}{x^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)/x\*\*(5/2),x)

[Out] Integral(cos(a + b\*x\*\*2)/x\*\*(5/2), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^2 + a)}{x^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)/x^(5/2),x, algorithm="giac")

[Out] integrate(cos(b\*x^2 + a)/x^(5/2), x)

### 3.29 $\int x^{5/2} \cos^2(a + bx^2) dx$

**Optimal.** Leaf size=132

$$-\frac{3ie^{2ia}x^{3/2}\Gamma\left(\frac{3}{4}, -2ibx^2\right)}{64 \cdot 2^{3/4}b(-ibx^2)^{3/4}} + \frac{3ie^{-2ia}x^{3/2}\Gamma\left(\frac{3}{4}, 2ibx^2\right)}{64 \cdot 2^{3/4}b(ibx^2)^{3/4}} + \frac{x^{3/2} \sin(2(a + bx^2))}{8b} + \frac{x^{7/2}}{7}$$

[Out]  $x^{(7/2)}/7 - (((3*I)/64)*E^{((2*I)*a)}*x^{(3/2)}*\Gamma[3/4, (-2*I)*b*x^2])/(2^{(3/4)}*b*((-I)*b*x^2)^{(3/4)}) + (((3*I)/64)*x^{(3/2)}*\Gamma[3/4, (2*I)*b*x^2])/(2^{(3/4)}*b)*E^{((2*I)*a)}*(I*b*x^2)^{(3/4)} + (x^{(3/2)}*\text{Sin}[2*(a + b*x^2)])/(8*b)$

**Rubi [A]** time = 0.174658, antiderivative size = 132, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 5, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$ , Rules used = {3402, 3404, 3386, 3389, 2218}

$$-\frac{3ie^{2ia}x^{3/2}\Gamma\left(\frac{3}{4}, -2ibx^2\right)}{64 \cdot 2^{3/4}b(-ibx^2)^{3/4}} + \frac{3ie^{-2ia}x^{3/2}\Gamma\left(\frac{3}{4}, 2ibx^2\right)}{64 \cdot 2^{3/4}b(ibx^2)^{3/4}} + \frac{x^{3/2} \sin(2(a + bx^2))}{8b} + \frac{x^{7/2}}{7}$$

Antiderivative was successfully verified.

[In] Int[x^(5/2)\*Cos[a + b\*x^2]^2,x]

[Out]  $x^{(7/2)}/7 - (((3*I)/64)*E^{((2*I)*a)}*x^{(3/2)}*\Gamma[3/4, (-2*I)*b*x^2])/(2^{(3/4)}*b*((-I)*b*x^2)^{(3/4)}) + (((3*I)/64)*x^{(3/2)}*\Gamma[3/4, (2*I)*b*x^2])/(2^{(3/4)}*b)*E^{((2*I)*a)}*(I*b*x^2)^{(3/4)} + (x^{(3/2)}*\text{Sin}[2*(a + b*x^2)])/(8*b)$

#### Rule 3402

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> With[{k = Denominator[m]}, Dist[k/e, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + (d\*x^(k\*n))]/e^n)]^p, x], x, (e\*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]

#### Rule 3404

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]

#### Rule 3386

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> Simp[(e^(n - 1)\*(e\*x)^(m - n + 1)\*Sin[c + d\*x^n])/(d\*n), x] - Dist[(e^n\*(m - n + 1))/(d\*n), Int[(e\*x)^(m - n)\*Sin[c + d\*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[n, m + 1]

#### Rule 3389

Int[((e\_.)\*(x\_)^(m\_.)\*Sin[(c\_.) + (d\_.)\*(x\_)^(n\_)], x\_Symbol] :> Dist[I/2, Int[(e\*x)^m\*E^(-(c\*I) - d\*I\*x^n), x], x] - Dist[I/2, Int[(e\*x)^m\*E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]

#### Rule 2218

```
Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> -Simp[(F^a*(e + f*x)^(m + 1)*Gamma[(m + 1)/n, -(b*(c + d*x)^n*Log[F])])/(f*n*(-(b*(c + d*x)^n*Log[F]))^((m + 1)/n)), x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

### Rubi steps

$$\begin{aligned} \int x^{5/2} \cos^2(a + bx^2) dx &= 2 \operatorname{Subst}\left(\int x^6 \cos^2(a + bx^4) dx, x, \sqrt{x}\right) \\ &= 2 \operatorname{Subst}\left(\int \left(\frac{x^6}{2} + \frac{1}{2}x^6 \cos(2a + 2bx^4)\right) dx, x, \sqrt{x}\right) \\ &= \frac{x^{7/2}}{7} + \operatorname{Subst}\left(\int x^6 \cos(2a + 2bx^4) dx, x, \sqrt{x}\right) \\ &= \frac{x^{7/2}}{7} + \frac{x^{3/2} \sin(2(a + bx^2))}{8b} - \frac{3 \operatorname{Subst}\left(\int x^2 \sin(2a + 2bx^4) dx, x, \sqrt{x}\right)}{8b} \\ &= \frac{x^{7/2}}{7} + \frac{x^{3/2} \sin(2(a + bx^2))}{8b} - \frac{(3i) \operatorname{Subst}\left(\int e^{-2ia-2ibx^4} x^2 dx, x, \sqrt{x}\right)}{16b} + \frac{(3i) \operatorname{Subst}\left(\int e^{2ia+2ibx^4} x^2 dx, x, \sqrt{x}\right)}{16b} \\ &= \frac{x^{7/2}}{7} - \frac{3ie^{2ia}x^{3/2}\Gamma\left(\frac{3}{4}, -2ibx^2\right)}{64 \cdot 2^{3/4}b(-ibx^2)^{3/4}} + \frac{3ie^{-2ia}x^{3/2}\Gamma\left(\frac{3}{4}, 2ibx^2\right)}{64 \cdot 2^{3/4}b(ibx^2)^{3/4}} + \frac{x^{3/2} \sin(2(a + bx^2))}{8b} \end{aligned}$$

**Mathematica [A]** time = 0.458192, size = 142, normalized size = 1.08

$$\frac{bx^{11/2} \left( 21 \sqrt[4]{2} (ibx^2)^{3/4} (\sin(2a) - i \cos(2a)) \Gamma\left(\frac{3}{4}, -2ibx^2\right) + 21 \sqrt[4]{2} (-ibx^2)^{3/4} (\sin(2a) + i \cos(2a)) \Gamma\left(\frac{3}{4}, 2ibx^2\right) \right)}{896 (b^2x^4)^{7/4}}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^(5/2)*Cos[a + b*x^2]^2, x]
```

```
[Out] (b*x^(11/2)*(21*2^(1/4)*(I*b*x^2)^(3/4)*Gamma[3/4, (-2*I)*b*x^2]*((-I)*Cos[2*a] + Sin[2*a]) + 21*2^(1/4)*((-I)*b*x^2)^(3/4)*Gamma[3/4, (2*I)*b*x^2]*(I*Cos[2*a] + Sin[2*a]) + 16*(b^2*x^4)^(3/4)*(8*b*x^2 + 7*Sin[2*(a + b*x^2)]))/ (896*(b^2*x^4)^(7/4))
```

**Maple [F]** time = 0.094, size = 0, normalized size = 0.

$$\int x^{\frac{5}{2}} (\cos(bx^2 + a))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^(5/2)*cos(b*x^2+a)^2, x)
```

```
[Out] int(x^(5/2)*cos(b*x^2+a)^2, x)
```

**Maxima [B]** time = 1.49486, size = 421, normalized size = 3.19

$$256 bx^4|b| + 224 x^2|b| \sin(2bx^2 + 2a) - 2^{\frac{1}{4}} (x^2|b|)^{\frac{1}{4}} \left( \left( (-21i\Gamma\left(\frac{3}{4}, 2ibx^2\right) + 21i\Gamma\left(\frac{3}{4}, -2ibx^2\right)) \cos\left(\frac{3}{8}\pi + \frac{3}{4}\arctan\left(\frac{bx^2}{a}\right)\right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(5/2)\*cos(b\*x^2+a)^2,x, algorithm="maxima")

[Out]  $\frac{1}{1792} \cdot (256 \cdot b \cdot x^4 \cdot \text{abs}(b) + 224 \cdot x^2 \cdot \text{abs}(b) \cdot \sin(2 \cdot b \cdot x^2 + 2 \cdot a) - 2^{(1/4)} \cdot (x^2 \cdot \text{abs}(b))^{(1/4)} \cdot (((-21 \cdot I \cdot \text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) + 21 \cdot I \cdot \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) \cdot \cos(3/8 \cdot \pi + 3/4 \cdot \arctan(0, b)) + (-21 \cdot I \cdot \text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) + 21 \cdot I \cdot \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) \cdot \cos(-3/8 \cdot \pi + 3/4 \cdot \arctan(0, b)) - 21 \cdot (\text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) + \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) \cdot \sin(3/8 \cdot \pi + 3/4 \cdot \arctan(0, b)) + 21 \cdot (\text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) + \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) \cdot \sin(-3/8 \cdot \pi + 3/4 \cdot \arctan(0, b))) \cdot \cos(2 \cdot a) - (21 \cdot (\text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) + \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) \cdot \cos(3/8 \cdot \pi + 3/4 \cdot \arctan(0, b)) + 21 \cdot (\text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) + \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) \cdot \cos(-3/8 \cdot \pi + 3/4 \cdot \arctan(0, b)) - (21 \cdot I \cdot \text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) - 21 \cdot I \cdot \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) \cdot \sin(3/8 \cdot \pi + 3/4 \cdot \arctan(0, b)) - (-21 \cdot I \cdot \text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) + 21 \cdot I \cdot \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) \cdot \sin(-3/8 \cdot \pi + 3/4 \cdot \arctan(0, b))) \cdot \sin(2 \cdot a)) / (b \cdot \sqrt{x} \cdot \text{abs}(b))$

**Fricas [A]** time = 1.81022, size = 242, normalized size = 1.83

$$\frac{21 (2i b)^{\frac{1}{4}} e^{(-2ia)} \Gamma\left(\frac{3}{4}, 2i b x^2\right) + 21 (-2i b)^{\frac{1}{4}} e^{(2ia)} \Gamma\left(\frac{3}{4}, -2i b x^2\right) + 32 (4b^2 x^3 + 7bx \cos(bx^2 + a) \sin(bx^2 + a)) \sqrt{x}}{896 b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(5/2)\*cos(b\*x^2+a)^2,x, algorithm="fricas")

[Out]  $\frac{1}{896} \cdot (21 \cdot (2 \cdot I \cdot b)^{(1/4)} \cdot e^{(-2 \cdot I \cdot a)} \cdot \text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) + 21 \cdot (-2 \cdot I \cdot b)^{(1/4)} \cdot e^{(2 \cdot I \cdot a)} \cdot \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2) + 32 \cdot (4 \cdot b^2 \cdot x^3 + 7 \cdot b \cdot x \cdot \cos(b \cdot x^2 + a) \cdot \sin(b \cdot x^2 + a)) \cdot \sqrt{x}) / b^2$

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*(5/2)\*cos(b\*x\*\*2+a)\*\*2,x)

[Out] Timed out

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{\frac{5}{2}} \cos(bx^2 + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(5/2)\*cos(b\*x^2+a)^2,x, algorithm="giac")

[Out] integrate(x^(5/2)\*cos(b\*x^2 + a)^2, x)

### 3.30 $\int x^{3/2} \cos^2(a + bx^2) dx$

**Optimal.** Leaf size=132

$$-\frac{ie^{2ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -2ibx^2\right)}{64\sqrt[4]{2b}\sqrt[4]{-ibx^2}} + \frac{ie^{-2ia}\sqrt{x}\Gamma\left(\frac{1}{4}, 2ibx^2\right)}{64\sqrt[4]{2b}\sqrt[4]{ibx^2}} + \frac{\sqrt{x}\sin(2(a + bx^2))}{8b} + \frac{x^{5/2}}{5}$$

[Out]  $x^{(5/2)}/5 - ((I/64)*E^{((2*I)*a)}*Sqrt[x]*Gamma[1/4, (-2*I)*b*x^2])/(2^{(1/4)}*b*((-I)*b*x^2)^{(1/4)}) + ((I/64)*Sqrt[x]*Gamma[1/4, (2*I)*b*x^2])/(2^{(1/4)}*b)*E^{((2*I)*a)}*(I*b*x^2)^{(1/4)} + (Sqrt[x]*Sin[2*(a + b*x^2)])/(8*b)$

**Rubi [A]** time = 0.123891, antiderivative size = 132, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 5, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$ , Rules used = {3402, 3404, 3386, 3355, 2208}

$$-\frac{ie^{2ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -2ibx^2\right)}{64\sqrt[4]{2b}\sqrt[4]{-ibx^2}} + \frac{ie^{-2ia}\sqrt{x}\Gamma\left(\frac{1}{4}, 2ibx^2\right)}{64\sqrt[4]{2b}\sqrt[4]{ibx^2}} + \frac{\sqrt{x}\sin(2(a + bx^2))}{8b} + \frac{x^{5/2}}{5}$$

Antiderivative was successfully verified.

[In] Int[x^(3/2)\*Cos[a + b\*x^2]^2,x]

[Out]  $x^{(5/2)}/5 - ((I/64)*E^{((2*I)*a)}*Sqrt[x]*Gamma[1/4, (-2*I)*b*x^2])/(2^{(1/4)}*b*((-I)*b*x^2)^{(1/4)}) + ((I/64)*Sqrt[x]*Gamma[1/4, (2*I)*b*x^2])/(2^{(1/4)}*b)*E^{((2*I)*a)}*(I*b*x^2)^{(1/4)} + (Sqrt[x]*Sin[2*(a + b*x^2)])/(8*b)$

#### Rule 3402

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] :> With[{k = Denominator[m]}, Dist[k/e, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + (d\*x^(k\*n))/e^n])^p, x], x, (e\*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]

#### Rule 3404

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] :> Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]

#### Rule 3386

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] :> Simp[(e^(n - 1)\*(e\*x)^(m - n + 1)\*Sin[c + d\*x^n])/(d\*n), x] - Dist[(e^n\*(m - n + 1))/(d\*n), Int[(e\*x)^(m - n)\*Sin[c + d\*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[n, m + 1]

#### Rule 3355

Int[Sin[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_)^(n\_))], x\_Symbol] :> Dist[I/2, Int[E^(-c\*I) - d\*I\*(e + f\*x)^n], x], x] - Dist[I/2, Int[E^(c\*I + d\*I\*(e + f\*x)^n), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[n, 2]

#### Rule 2208

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_))^(n\_)), x\_Symbol] :> -Simp[(F^a\*(c + d\*x)\*Gamma[1/n, -(b\*(c + d\*x)^n\*Log[F]])]/(d\*n\*(-(b\*(c + d\*x)^n\*Log[F

]))^(1/n)), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]

### Rubi steps

$$\begin{aligned}
 \int x^{3/2} \cos^2(a + bx^2) dx &= 2 \operatorname{Subst} \left( \int x^4 \cos^2(a + bx^4) dx, x, \sqrt{x} \right) \\
 &= 2 \operatorname{Subst} \left( \int \left( \frac{x^4}{2} + \frac{1}{2} x^4 \cos(2a + 2bx^4) \right) dx, x, \sqrt{x} \right) \\
 &= \frac{x^{5/2}}{5} + \operatorname{Subst} \left( \int x^4 \cos(2a + 2bx^4) dx, x, \sqrt{x} \right) \\
 &= \frac{x^{5/2}}{5} + \frac{\sqrt{x} \sin(2(a + bx^2))}{8b} - \frac{\operatorname{Subst} \left( \int \sin(2a + 2bx^4) dx, x, \sqrt{x} \right)}{8b} \\
 &= \frac{x^{5/2}}{5} + \frac{\sqrt{x} \sin(2(a + bx^2))}{8b} - \frac{i \operatorname{Subst} \left( \int e^{-2ia - 2ibx^4} dx, x, \sqrt{x} \right)}{16b} + \frac{i \operatorname{Subst} \left( \int e^{2ia + 2ibx^4} dx, x, \sqrt{x} \right)}{16b} \\
 &= \frac{x^{5/2}}{5} - \frac{ie^{2ia} \sqrt{x} \Gamma\left(\frac{1}{4}, -2ibx^2\right)}{64 \sqrt[4]{2b} \sqrt[4]{-ibx^2}} + \frac{ie^{-2ia} \sqrt{x} \Gamma\left(\frac{1}{4}, 2ibx^2\right)}{64 \sqrt[4]{2b} \sqrt[4]{ibx^2}} + \frac{\sqrt{x} \sin(2(a + bx^2))}{8b}
 \end{aligned}$$

**Mathematica [A]** time = 0.344793, size = 142, normalized size = 1.08

$$\frac{bx^{9/2} \left( 5 \cdot 2^{3/4} \sqrt[4]{ibx^2} (\sin(2a) - i \cos(2a)) \Gamma\left(\frac{1}{4}, -2ibx^2\right) + 5 \cdot 2^{3/4} \sqrt[4]{-ibx^2} (\sin(2a) + i \cos(2a)) \Gamma\left(\frac{1}{4}, 2ibx^2\right) + 16 \right)}{640 (b^2 x^4)^{5/4}}$$

Antiderivative was successfully verified.

[In] Integrate[x^(3/2)\*Cos[a + b\*x^2]^2, x]

[Out] (b\*x^(9/2)\*(5\*2^(3/4)\*(I\*b\*x^2)^(1/4)\*Gamma[1/4, (-2\*I)\*b\*x^2]\*((-I)\*Cos[2\*a] + Sin[2\*a]) + 5\*2^(3/4)\*((-I)\*b\*x^2)^(1/4)\*Gamma[1/4, (2\*I)\*b\*x^2]\*(I\*Cos[2\*a] + Sin[2\*a]) + 16\*(b^2\*x^4)^(1/4)\*(8\*b\*x^2 + 5\*Sin[2\*(a + b\*x^2)])))/(640\*(b^2\*x^4)^(5/4))

**Maple [F]** time = 0.089, size = 0, normalized size = 0.

$$\int x^{\frac{3}{2}} (\cos(bx^2 + a))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(3/2)\*cos(b\*x^2+a)^2, x)

[Out] int(x^(3/2)\*cos(b\*x^2+a)^2, x)

**Maxima [B]** time = 1.41165, size = 424, normalized size = 3.21

$$2^{\frac{3}{4}} \left( 16 \cdot 2^{\frac{1}{4}} \left( 8bx^{\frac{5}{2}} + 5\sqrt{x} \sin(2bx^2 + 2a) \right) (x^2|b|)^{\frac{1}{4}} + \left( \left( 5i\Gamma\left(\frac{1}{4}, 2ibx^2\right) - 5i\Gamma\left(\frac{1}{4}, -2ibx^2\right) \right) \cos\left(\frac{1}{8}\pi + \frac{1}{4}\arctan(0, b)\right) + \right. \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(b\*x^2+a)^2,x, algorithm="maxima")

[Out]  $\frac{1}{1280} 2^{3/4} (16 \cdot 2^{1/4} (8 b x^{5/2} + 5 \sqrt{x} \sin(2 b x^2 + 2 a)) (x^2 \operatorname{abs}(b))^{1/4} + ((5 I \operatorname{gamma}(1/4, 2 I b x^2) - 5 I \operatorname{gamma}(1/4, -2 I b x^2)) \cos(1/8 \pi + 1/4 \arctan(0, b)) + (5 I \operatorname{gamma}(1/4, 2 I b x^2) - 5 I \operatorname{gamma}(1/4, -2 I b x^2)) \cos(-1/8 \pi + 1/4 \arctan(0, b)) + 5 (\operatorname{gamma}(1/4, 2 I b x^2) + \operatorname{gamma}(1/4, -2 I b x^2)) \sin(1/8 \pi + 1/4 \arctan(0, b)) - 5 (\operatorname{gamma}(1/4, 2 I b x^2) + \operatorname{gamma}(1/4, -2 I b x^2)) \sin(-1/8 \pi + 1/4 \arctan(0, b))) \cos(2 a) + (5 (\operatorname{gamma}(1/4, 2 I b x^2) + \operatorname{gamma}(1/4, -2 I b x^2)) \cos(1/8 \pi + 1/4 \arctan(0, b)) + 5 (\operatorname{gamma}(1/4, 2 I b x^2) + \operatorname{gamma}(1/4, -2 I b x^2)) \cos(-1/8 \pi + 1/4 \arctan(0, b)) + (-5 I \operatorname{gamma}(1/4, 2 I b x^2) + 5 I \operatorname{gamma}(1/4, -2 I b x^2)) \sin(1/8 \pi + 1/4 \arctan(0, b)) + (5 I \operatorname{gamma}(1/4, 2 I b x^2) - 5 I \operatorname{gamma}(1/4, -2 I b x^2)) \sin(-1/8 \pi + 1/4 \arctan(0, b))) \sin(2 a)) \sqrt{x} / ((x^2 \operatorname{abs}(b))^{1/4} b)$

**Fricas [A]** time = 1.82749, size = 236, normalized size = 1.79

$$\frac{5 (2i b)^{\frac{3}{4}} e^{(-2ia)} \Gamma\left(\frac{1}{4}, 2i b x^2\right) + 5 (-2i b)^{\frac{3}{4}} e^{(2ia)} \Gamma\left(\frac{1}{4}, -2i b x^2\right) + 32 (4 b^2 x^2 + 5 b \cos(b x^2 + a) \sin(b x^2 + a)) \sqrt{x}}{640 b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(b\*x^2+a)^2,x, algorithm="fricas")

[Out]  $\frac{1}{640} (5 (2 I b)^{3/4} e^{-2 I a} \operatorname{gamma}(1/4, 2 I b x^2) + 5 (-2 I b)^{3/4} e^{2 I a} \operatorname{gamma}(1/4, -2 I b x^2) + 32 (4 b^2 x^2 + 5 b \cos(b x^2 + a) \sin(b x^2 + a)) \sqrt{x}) / b^2$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{\frac{3}{2}} \cos^2(a + b x^2) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*(3/2)\*cos(b\*x\*\*2+a)\*\*2,x)

[Out] Integral(x\*\*(3/2)\*cos(a + b\*x\*\*2)\*\*2, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{\frac{3}{2}} \cos(b x^2 + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(b\*x^2+a)^2,x, algorithm="giac")

[Out] integrate(x^(3/2)\*cos(b\*x^2 + a)^2, x)

### 3.31 $\int \sqrt{x} \cos^2(a + bx^2) dx$

**Optimal.** Leaf size=100

$$-\frac{e^{2ia}x^{3/2}\Gamma\left(\frac{3}{4}, -2ibx^2\right)}{8 \cdot 2^{3/4}(-ibx^2)^{3/4}} - \frac{e^{-2ia}x^{3/2}\Gamma\left(\frac{3}{4}, 2ibx^2\right)}{8 \cdot 2^{3/4}(ibx^2)^{3/4}} + \frac{x^{3/2}}{3}$$

[Out]  $x^{(3/2)}/3 - (E^{((2*I)*a)}*x^{(3/2)}*\Gamma[3/4, (-2*I)*b*x^2])/(8*2^{(3/4)}*((-I)*b*x^2)^{(3/4)}) - (x^{(3/2)}*\Gamma[3/4, (2*I)*b*x^2])/(8*2^{(3/4)}*E^{((2*I)*a)}*(I*b*x^2)^{(3/4)})$

**Rubi [A]** time = 0.130385, antiderivative size = 100, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 4, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3402, 3404, 3390, 2218}

$$-\frac{e^{2ia}x^{3/2}\Gamma\left(\frac{3}{4}, -2ibx^2\right)}{8 \cdot 2^{3/4}(-ibx^2)^{3/4}} - \frac{e^{-2ia}x^{3/2}\Gamma\left(\frac{3}{4}, 2ibx^2\right)}{8 \cdot 2^{3/4}(ibx^2)^{3/4}} + \frac{x^{3/2}}{3}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[x]\*Cos[a + b\*x^2]^2,x]

[Out]  $x^{(3/2)}/3 - (E^{((2*I)*a)}*x^{(3/2)}*\Gamma[3/4, (-2*I)*b*x^2])/(8*2^{(3/4)}*((-I)*b*x^2)^{(3/4)}) - (x^{(3/2)}*\Gamma[3/4, (2*I)*b*x^2])/(8*2^{(3/4)}*E^{((2*I)*a)}*(I*b*x^2)^{(3/4)})$

#### Rule 3402

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> With[{k = Denominator[m]}, Dist[k/e, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + (d\*x^(k\*n))]/e^n)]^p, x], x, (e\*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]

#### Rule 3404

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]

#### Rule 3390

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> Dist[1/2, Int[(e\*x)^m\*E^(-(c\*I) - d\*I\*x^n), x], x] + Dist[1/2, Int[(e\*x)^m\*E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]

#### Rule 2218

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_)^(n\_)))\*((e\_.) + (f\_.)\*(x\_)^(m\_.), x\_Symbol] :> -Simp[(F^a\*(e + f\*x)^(m + 1)\*Gamma[(m + 1)/n, -(b\*(c + d\*x)^n\*Log[F])])/(f\*n\*(-(b\*(c + d\*x)^n\*Log[F]))^(m + 1)/n), x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d\*e - c\*f, 0]

#### Rubi steps



$$\begin{aligned}
\int \sqrt{x} \cos^2(a + bx^2) dx &= 2 \operatorname{Subst} \left( \int x^2 \cos^2(a + bx^4) dx, x, \sqrt{x} \right) \\
&= 2 \operatorname{Subst} \left( \int \left( \frac{x^2}{2} + \frac{1}{2} x^2 \cos(2a + 2bx^4) \right) dx, x, \sqrt{x} \right) \\
&= \frac{x^{3/2}}{3} + \operatorname{Subst} \left( \int x^2 \cos(2a + 2bx^4) dx, x, \sqrt{x} \right) \\
&= \frac{x^{3/2}}{3} + \frac{1}{2} \operatorname{Subst} \left( \int e^{-2ia - 2ibx^4} x^2 dx, x, \sqrt{x} \right) + \frac{1}{2} \operatorname{Subst} \left( \int e^{2ia + 2ibx^4} x^2 dx, x, \sqrt{x} \right) \\
&= \frac{x^{3/2}}{3} - \frac{e^{2ia} x^{3/2} \Gamma\left(\frac{3}{4}, -2ibx^2\right)}{8 \cdot 2^{3/4} (-ibx^2)^{3/4}} - \frac{e^{-2ia} x^{3/2} \Gamma\left(\frac{3}{4}, 2ibx^2\right)}{8 \cdot 2^{3/4} (ibx^2)^{3/4}}
\end{aligned}$$

**Mathematica [A]** time = 0.226213, size = 122, normalized size = 1.22

$$\frac{x^{3/2} \left( -3\sqrt[4]{2} (-ibx^2)^{3/4} (\cos(2a) - i \sin(2a)) \Gamma\left(\frac{3}{4}, 2ibx^2\right) - 3\sqrt[4]{2} (ibx^2)^{3/4} (\cos(2a) + i \sin(2a)) \Gamma\left(\frac{3}{4}, -2ibx^2\right) \right)}{48 (b^2 x^4)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[x]\*Cos[a + b\*x^2]^2,x]

[Out] (x^(3/2)\*(16\*(b^2\*x^4)^(3/4) - 3\*2^(1/4)\*((-I)\*b\*x^2)^(3/4)\*Gamma[3/4, (2\*I)\*b\*x^2]\*(Cos[2\*a] - I\*Sin[2\*a]) - 3\*2^(1/4)\*(I\*b\*x^2)^(3/4)\*Gamma[3/4, (-2\*I)\*b\*x^2]\*(Cos[2\*a] + I\*Sin[2\*a]))/(48\*(b^2\*x^4)^(3/4))

**Maple [F]** time = 0.085, size = 0, normalized size = 0.

$$\int \sqrt{x} (\cos(bx^2 + a))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(1/2)\*cos(b\*x^2+a)^2,x)

[Out] int(x^(1/2)\*cos(b\*x^2+a)^2,x)

**Maxima [B]** time = 1.4219, size = 392, normalized size = 3.92

$$32 x^2 |b| - 2^{\frac{1}{4}} (x^2 |b|)^{\frac{1}{4}} \left( \left( 3 \left( \Gamma\left(\frac{3}{4}, 2i b x^2\right) + \Gamma\left(\frac{3}{4}, -2i b x^2\right) \right) \cos\left(\frac{3}{8} \pi + \frac{3}{4} \arctan(0, b)\right) + 3 \left( \Gamma\left(\frac{3}{4}, 2i b x^2\right) + \Gamma\left(\frac{3}{4}, -2i b x^2\right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(1/2)\*cos(b\*x^2+a)^2,x, algorithm="maxima")

[Out] 1/96\*(32\*x^2\*abs(b) - 2^(1/4)\*(x^2\*abs(b))^(1/4))\*((3\*(gamma(3/4, 2\*I\*b\*x^2) + gamma(3/4, -2\*I\*b\*x^2))\*cos(3/8\*pi + 3/4\*arctan2(0, b)) + 3\*(gamma(3/4, 2\*I\*b\*x^2) + gamma(3/4, -2\*I\*b\*x^2))\*cos(-3/8\*pi + 3/4\*arctan2(0, b)) - (3\*I\*gamma(3/4, 2\*I\*b\*x^2) - 3\*I\*gamma(3/4, -2\*I\*b\*x^2))\*sin(3/8\*pi + 3/4\*arct

$$\begin{aligned} & \operatorname{atan2}(0, b) - (-3I\gamma(3/4, 2Ibx^2) + 3I\gamma(3/4, -2Ibx^2))\sin(-3/8\pi + 3/4\arctan2(0, b))\cos(2a) - ((3I\gamma(3/4, 2Ibx^2) - 3I\gamma(3/4, -2Ibx^2))\cos(3/8\pi + 3/4\arctan2(0, b)) + (3I\gamma(3/4, 2Ibx^2) - 3I\gamma(3/4, -2Ibx^2))\cos(-3/8\pi + 3/4\arctan2(0, b)) + 3(\gamma(3/4, 2Ibx^2) + \gamma(3/4, -2Ibx^2))\sin(3/8\pi + 3/4\arctan2(0, b)) - 3(\gamma(3/4, 2Ibx^2) + \gamma(3/4, -2Ibx^2))\sin(-3/8\pi + 3/4\arctan2(0, b)))\sin(2a))/(\sqrt{x}\operatorname{abs}(b)) \end{aligned}$$

**Fricas [A]** time = 1.7438, size = 176, normalized size = 1.76

$$\frac{16bx^{\frac{3}{2}} + 3i(2ib)^{\frac{1}{4}}e^{(-2ia)}\Gamma\left(\frac{3}{4}, 2ibx^2\right) - 3i(-2ib)^{\frac{1}{4}}e^{(2ia)}\Gamma\left(\frac{3}{4}, -2ibx^2\right)}{48b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(1/2)\*cos(b\*x^2+a)^2,x, algorithm="fricas")

[Out] 1/48\*(16\*b\*x^(3/2) + 3\*I\*(2\*I\*b)^(1/4)\*e^(-2\*I\*a)\*gamma(3/4, 2\*I\*b\*x^2) - 3\*I\*(-2\*I\*b)^(1/4)\*e^(2\*I\*a)\*gamma(3/4, -2\*I\*b\*x^2))/b

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \sqrt{x} \cos^2(a + bx^2) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*(1/2)\*cos(b\*x\*\*2+a)\*\*2,x)

[Out] Integral(sqrt(x)\*cos(a + b\*x\*\*2)\*\*2, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \sqrt{x} \cos(bx^2 + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(1/2)\*cos(b\*x^2+a)^2,x, algorithm="giac")

[Out] integrate(sqrt(x)\*cos(b\*x^2 + a)^2, x)

$$3.32 \quad \int \frac{\cos^2(a+bx^2)}{\sqrt{x}} dx$$

**Optimal.** Leaf size=96

$$-\frac{e^{2ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -2ibx^2\right)}{8\sqrt[4]{2}\sqrt[4]{-ibx^2}} - \frac{e^{-2ia}\sqrt{x}\Gamma\left(\frac{1}{4}, 2ibx^2\right)}{8\sqrt[4]{2}\sqrt[4]{ibx^2}} + \sqrt{x}$$

[Out] Sqrt[x] - (E^((2\*I)\*a)\*Sqrt[x]\*Gamma[1/4, (-2\*I)\*b\*x^2])/(8\*2^(1/4)\*((-I)\*b\*x^2)^(1/4)) - (Sqrt[x]\*Gamma[1/4, (2\*I)\*b\*x^2])/(8\*2^(1/4)\*E^((2\*I)\*a)\*(I\*b\*x^2)^(1/4))

**Rubi [A]** time = 0.0681866, antiderivative size = 96, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 4, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3402, 3358, 3356, 2208}

$$-\frac{e^{2ia}\sqrt{x}\Gamma\left(\frac{1}{4}, -2ibx^2\right)}{8\sqrt[4]{2}\sqrt[4]{-ibx^2}} - \frac{e^{-2ia}\sqrt{x}\Gamma\left(\frac{1}{4}, 2ibx^2\right)}{8\sqrt[4]{2}\sqrt[4]{ibx^2}} + \sqrt{x}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]^2/Sqrt[x], x]

[Out] Sqrt[x] - (E^((2\*I)\*a)\*Sqrt[x]\*Gamma[1/4, (-2\*I)\*b\*x^2])/(8\*2^(1/4)\*((-I)\*b\*x^2)^(1/4)) - (Sqrt[x]\*Gamma[1/4, (2\*I)\*b\*x^2])/(8\*2^(1/4)\*E^((2\*I)\*a)\*(I\*b\*x^2)^(1/4))

#### Rule 3402

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> With[{k = Denominator[m]}, Dist[k/e, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + (d\*x^(k\*n))/e^n])^p, x], x, (e\*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]

#### Rule 3358

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_)^(n\_))])\*(b\_.))^(p\_), x\_Symbol] :> Int[ExpandTrigReduce[(a + b\*Cos[c + d\*(e + f\*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 1] && IGtQ[n, 1]

#### Rule 3356

Int[Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_)^(n\_))], x\_Symbol] :> Dist[1/2, Int[E^(-c\*I) - d\*I\*(e + f\*x)^n], x], x] + Dist[1/2, Int[E^(c\*I + d\*I\*(e + f\*x)^n), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[n, 2]

#### Rule 2208

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_)^(n\_))), x\_Symbol] :> -Simp[(F^a\*(c + d\*x)\*Gamma[1/n, -(b\*(c + d\*x)^n\*Log[F]])]/(d\*n\*(-(b\*(c + d\*x)^n\*Log[F]))^(1/n)), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx &= 2 \operatorname{Subst} \left( \int \cos^2(a + bx^4) dx, x, \sqrt{x} \right) \\
&= 2 \operatorname{Subst} \left( \int \left( \frac{1}{2} + \frac{1}{2} \cos(2a + 2bx^4) \right) dx, x, \sqrt{x} \right) \\
&= \sqrt{x} + \operatorname{Subst} \left( \int \cos(2a + 2bx^4) dx, x, \sqrt{x} \right) \\
&= \sqrt{x} + \frac{1}{2} \operatorname{Subst} \left( \int e^{-2ia - 2ibx^4} dx, x, \sqrt{x} \right) + \frac{1}{2} \operatorname{Subst} \left( \int e^{2ia + 2ibx^4} dx, x, \sqrt{x} \right) \\
&= \sqrt{x} - \frac{e^{2ia} \sqrt{x} \Gamma\left(\frac{1}{4}, -2ibx^2\right)}{8\sqrt[4]{2}\sqrt[4]{-ibx^2}} - \frac{e^{-2ia} \sqrt{x} \Gamma\left(\frac{1}{4}, 2ibx^2\right)}{8\sqrt[4]{2}\sqrt[4]{ibx^2}}
\end{aligned}$$

**Mathematica [A]** time = 0.232448, size = 120, normalized size = 1.25

$$\frac{\sqrt{x} \left( 2^{3/4} \sqrt[4]{-ibx^2} (\cos(2a) - i \sin(2a)) \Gamma\left(\frac{1}{4}, 2ibx^2\right) + 2^{3/4} \sqrt[4]{ibx^2} (\cos(2a) + i \sin(2a)) \Gamma\left(\frac{1}{4}, -2ibx^2\right) - 16 \sqrt[4]{b^2} \right)}{16 \sqrt[4]{b^2} x^4}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]^2/Sqrt[x], x]

[Out]  $-\left(\sqrt{x} \left(-16 \left(b^2 x^4\right)^{1/4} + 2^{3/4} \left((-1) b x^2\right)^{1/4} \Gamma\left[\frac{1}{4}, (2 I) b x^2\right] (\cos[2 a] - I \sin[2 a]) + 2^{3/4} \left(I b x^2\right)^{1/4} \Gamma\left[\frac{1}{4}, (-2 I) b x^2\right] (\cos[2 a] + I \sin[2 a])\right)\right) / \left(16 \left(b^2 x^4\right)^{1/4}\right)$

**Maple [F]** time = 0.099, size = 0, normalized size = 0.

$$\int (\cos(bx^2 + a))^2 \frac{1}{\sqrt{x}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)^2/x^(1/2), x)

[Out] int(cos(b\*x^2+a)^2/x^(1/2), x)

**Maxima [B]** time = 1.45108, size = 393, normalized size = 4.09

$$2^{\frac{3}{4}} \left( \left( \left( \Gamma\left(\frac{1}{4}, 2i bx^2\right) + \Gamma\left(\frac{1}{4}, -2i bx^2\right) \right) \cos\left(\frac{1}{8} \pi + \frac{1}{4} \arctan(0, b)\right) + \left( \Gamma\left(\frac{1}{4}, 2i bx^2\right) + \Gamma\left(\frac{1}{4}, -2i bx^2\right) \right) \cos\left(-\frac{1}{8} \pi + \frac{1}{4} \arctan(0, b)\right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^(1/2), x, algorithm="maxima")

[Out]  $-1/32 \cdot 2^{3/4} \cdot \left( \left( \left( \gamma\left(\frac{1}{4}, 2I b x^2\right) + \gamma\left(\frac{1}{4}, -2I b x^2\right) \right) \cdot \cos\left(\frac{1}{8} \pi + \frac{1}{4} \arctan(0, b)\right) + \left( \gamma\left(\frac{1}{4}, 2I b x^2\right) + \gamma\left(\frac{1}{4}, -2I b x^2\right) \right) \cdot \cos\left(-\frac{1}{8} \pi + \frac{1}{4} \arctan(0, b)\right) \right) - \left( I \gamma\left(\frac{1}{4}, 2I b x^2\right) - I \gamma\left(\frac{1}{4}, -2I b x^2\right) \right) \cdot \sqrt{x}$

$$2*I*b*x^2))*\sin(1/8*\pi + 1/4*\arctan2(0, b)) - (-I*\gamma(1/4, 2*I*b*x^2) + I*\gamma(1/4, -2*I*b*x^2))*\sin(-1/8*\pi + 1/4*\arctan2(0, b)))*\cos(2*a) - ((I*\gamma(1/4, 2*I*b*x^2) - I*\gamma(1/4, -2*I*b*x^2))*\cos(1/8*\pi + 1/4*\arctan2(0, b)) + (I*\gamma(1/4, 2*I*b*x^2) - I*\gamma(1/4, -2*I*b*x^2))*\cos(-1/8*\pi + 1/4*\arctan2(0, b)) + (\gamma(1/4, 2*I*b*x^2) + \gamma(1/4, -2*I*b*x^2))*\sin(1/8*\pi + 1/4*\arctan2(0, b)) - (\gamma(1/4, 2*I*b*x^2) + \gamma(1/4, -2*I*b*x^2))*\sin(-1/8*\pi + 1/4*\arctan2(0, b)))*\sin(2*a))*\sqrt{x} - 16*2^{1/4}*(x^2*abs(b))^{1/4}*\sqrt{x})/(x^2*abs(b))^{1/4}$$

**Fricas [A]** time = 1.76987, size = 170, normalized size = 1.77

$$\frac{i(2ib)^{\frac{3}{4}}e^{(-2ia)}\Gamma\left(\frac{1}{4}, 2ibx^2\right) - i(-2ib)^{\frac{3}{4}}e^{(2ia)}\Gamma\left(\frac{1}{4}, -2ibx^2\right) + 16b\sqrt{x}}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^(1/2), x, algorithm="fricas")

[Out] 1/16\*(I\*(2\*I\*b)^(3/4)\*e^(-2\*I\*a)\*gamma(1/4, 2\*I\*b\*x^2) - I\*(-2\*I\*b)^(3/4)\*e^(2\*I\*a)\*gamma(1/4, -2\*I\*b\*x^2) + 16\*b\*sqrt(x))/b

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*2/x\*\*(1/2), x)

[Out] Integral(cos(a + b\*x\*\*2)\*\*2/sqrt(x), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^2 + a)^2}{\sqrt{x}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^(1/2), x, algorithm="giac")

[Out] integrate(cos(b\*x^2 + a)^2/sqrt(x), x)

$$3.33 \quad \int \frac{\cos^2(a+bx^2)}{x^{3/2}} dx$$

**Optimal.** Leaf size=117

$$-\frac{ie^{2ia}bx^{3/2}\Gamma\left(\frac{3}{4}, -2ibx^2\right)}{2^{3/4}(-ibx^2)^{3/4}} + \frac{ie^{-2ia}bx^{3/2}\Gamma\left(\frac{3}{4}, 2ibx^2\right)}{2^{3/4}(ibx^2)^{3/4}} - \frac{\cos(2(a+bx^2))}{\sqrt{x}} - \frac{1}{\sqrt{x}}$$

[Out] -(1/Sqrt[x]) - Cos[2\*(a + b\*x^2)]/Sqrt[x] - (I\*b\*E^((2\*I)\*a)\*x^(3/2)\*Gamma[3/4, (-2\*I)\*b\*x^2])/(2^(3/4)\*((-I)\*b\*x^2)^(3/4)) + (I\*b\*x^(3/2)\*Gamma[3/4, (2\*I)\*b\*x^2])/(2^(3/4)\*E^((2\*I)\*a)\*(I\*b\*x^2)^(3/4))

**Rubi [A]** time = 0.163031, antiderivative size = 117, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 5, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$ , Rules used = {3402, 3404, 3388, 3389, 2218}

$$-\frac{ie^{2ia}bx^{3/2}\Gamma\left(\frac{3}{4}, -2ibx^2\right)}{2^{3/4}(-ibx^2)^{3/4}} + \frac{ie^{-2ia}bx^{3/2}\Gamma\left(\frac{3}{4}, 2ibx^2\right)}{2^{3/4}(ibx^2)^{3/4}} - \frac{\cos(2(a+bx^2))}{\sqrt{x}} - \frac{1}{\sqrt{x}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^2]^2/x^(3/2), x]

[Out] -(1/Sqrt[x]) - Cos[2\*(a + b\*x^2)]/Sqrt[x] - (I\*b\*E^((2\*I)\*a)\*x^(3/2)\*Gamma[3/4, (-2\*I)\*b\*x^2])/(2^(3/4)\*((-I)\*b\*x^2)^(3/4)) + (I\*b\*x^(3/2)\*Gamma[3/4, (2\*I)\*b\*x^2])/(2^(3/4)\*E^((2\*I)\*a)\*(I\*b\*x^2)^(3/4))

#### Rule 3402

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*((e\_.)\*(x\_))^(m\_), x\_Symbol] :> With[{k = Denominator[m]}, Dist[k/e, Subst[Int[x^(k\*(m + 1) - 1)]\*(a + b\*Cos[c + d\*x^(k\*n)]/e^n)^p, x], x, (e\*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]

#### Rule 3404

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] :> Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]

#### Rule 3388

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*((e\_.)\*(x\_))^(m\_), x\_Symbol] :> Simp[((e\*x)^(m + 1)\*Cos[c + d\*x^n])/(e\*(m + 1)), x] + Dist[(d\*n)/(e^n\*(m + 1)), Int[(e\*x)^(m + n)\*Sin[c + d\*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[m, -1]

#### Rule 3389

Int[((e\_.)\*(x\_))^(m\_.)\*Sin[(c\_.) + (d\_.)\*(x\_)^(n\_)], x\_Symbol] :> Dist[I/2, Int[(e\*x)^m\*E^(-c\*I - d\*I\*x^n), x], x] - Dist[I/2, Int[(e\*x)^m\*E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]

#### Rule 2218

```
Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(m_
.), x_Symbol] :> -Simp[(F^a*(e + f*x)^(m + 1)*Gamma[(m + 1)/n, -(b*(c + d*x
)^n*Log[F]])]/(f*n*(-(b*(c + d*x)^n*Log[F]))^((m + 1)/n)), x] /; FreeQ[{F,
a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

### Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(a + bx^2)}{x^{3/2}} dx &= 2 \operatorname{Subst} \left( \int \frac{\cos^2(a + bx^4)}{x^2} dx, x, \sqrt{x} \right) \\
&= 2 \operatorname{Subst} \left( \int \left( \frac{1}{2x^2} + \frac{\cos(2a + 2bx^4)}{2x^2} \right) dx, x, \sqrt{x} \right) \\
&= -\frac{1}{\sqrt{x}} + \operatorname{Subst} \left( \int \frac{\cos(2a + 2bx^4)}{x^2} dx, x, \sqrt{x} \right) \\
&= -\frac{1}{\sqrt{x}} - \frac{\cos(2(a + bx^2))}{\sqrt{x}} - (8b) \operatorname{Subst} \left( \int x^2 \sin(2a + 2bx^4) dx, x, \sqrt{x} \right) \\
&= -\frac{1}{\sqrt{x}} - \frac{\cos(2(a + bx^2))}{\sqrt{x}} - (4ib) \operatorname{Subst} \left( \int e^{-2ia - 2ibx^4} x^2 dx, x, \sqrt{x} \right) + (4ib) \operatorname{Subst} \left( \int e^{2ia + 2ibx^4} x^2 dx, x, \sqrt{x} \right) \\
&= -\frac{1}{\sqrt{x}} - \frac{\cos(2(a + bx^2))}{\sqrt{x}} - \frac{ibe^{2ia} x^{3/2} \Gamma\left(\frac{3}{4}, -2ibx^2\right)}{2^{3/4} (-ibx^2)^{3/4}} + \frac{ibe^{-2ia} x^{3/2} \Gamma\left(\frac{3}{4}, 2ibx^2\right)}{2^{3/4} (ibx^2)^{3/4}}
\end{aligned}$$

**Mathematica [A]** time = 0.339617, size = 137, normalized size = 1.17

$$\frac{\sqrt[4]{2} b x^2 (i b x^2)^{3/4} (\sin(2a) - i \cos(2a)) \Gamma\left(\frac{3}{4}, -2i b x^2\right) + i \sqrt[4]{2} (-i b x^2)^{7/4} (\sin(2a) + i \cos(2a)) \Gamma\left(\frac{3}{4}, 2i b x^2\right)}{2 \sqrt{x} (b^2 x^4)^{3/4}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x^2]^2/x^(3/2), x]
```

```
[Out] (-4*(b^2*x^4)^(3/4)*Cos[a + b*x^2]^2 + 2^(1/4)*b*x^2*(I*b*x^2)^(3/4)*Gamma[
3/4, (-2*I)*b*x^2]*((-I)*Cos[2*a] + Sin[2*a]) + I*2^(1/4)*((-I)*b*x^2)^(7/4
)*Gamma[3/4, (2*I)*b*x^2]*(I*Cos[2*a] + Sin[2*a])/(2*Sqrt[x]*(b^2*x^4)^(3/
4))
```

**Maple [F]** time = 0.089, size = 0, normalized size = 0.

$$\int (\cos(bx^2 + a))^2 x^{-\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(b*x^2+a)^2/x^(3/2), x)
```

```
[Out] int(cos(b*x^2+a)^2/x^(3/2), x)
```

**Maxima [B]** time = 1.34728, size = 369, normalized size = 3.15

$$2^{\frac{1}{4}} (x^2|b|)^{\frac{1}{4}} \left( \left( \Gamma\left(-\frac{1}{4}, 2i bx^2\right) + \Gamma\left(-\frac{1}{4}, -2i bx^2\right) \right) \cos\left(\frac{1}{8}\pi + \frac{1}{4}\arctan(0, b)\right) + \left( \Gamma\left(-\frac{1}{4}, 2i bx^2\right) + \Gamma\left(-\frac{1}{4}, -2i bx^2\right) \right) \cos\left(-\frac{1}{8}\pi + \frac{1}{4}\arctan(0, b)\right) + (I\gamma(-1/4, 2I*b*x^2) - I\gamma(-1/4, -2I*b*x^2))*\sin(1/8\pi + 1/4*\arctan2(0, b)) + (-I\gamma(-1/4, 2I*b*x^2) + I\gamma(-1/4, -2I*b*x^2))*\sin(-1/8\pi + 1/4*\arctan2(0, b)) \right) \cos(2*a) + ((-I\gamma(-1/4, 2I*b*x^2) + I\gamma(-1/4, -2I*b*x^2))*\cos(1/8\pi + 1/4*\arctan2(0, b)) + (-I\gamma(-1/4, 2I*b*x^2) + I\gamma(-1/4, -2I*b*x^2))*\cos(-1/8\pi + 1/4*\arctan2(0, b)) + (\gamma(-1/4, 2I*b*x^2) + \gamma(-1/4, -2I*b*x^2))*\sin(1/8\pi + 1/4*\arctan2(0, b)) - (\gamma(-1/4, 2I*b*x^2) + \gamma(-1/4, -2I*b*x^2))*\sin(-1/8\pi + 1/4*\arctan2(0, b))) * \sin(2*a) + 16) / \sqrt{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^(3/2),x, algorithm="maxima")

[Out] -1/16\*(2^(1/4)\*(x^2\*abs(b))^(1/4)\*(((gamma(-1/4, 2\*I\*b\*x^2) + gamma(-1/4, -2\*I\*b\*x^2))\*cos(1/8\*pi + 1/4\*arctan2(0, b)) + (gamma(-1/4, 2\*I\*b\*x^2) + gamma(-1/4, -2\*I\*b\*x^2))\*cos(-1/8\*pi + 1/4\*arctan2(0, b)) + (I\*gamma(-1/4, 2\*I\*b\*x^2) - I\*gamma(-1/4, -2\*I\*b\*x^2))\*sin(1/8\*pi + 1/4\*arctan2(0, b)) + (-I\*gamma(-1/4, 2\*I\*b\*x^2) + I\*gamma(-1/4, -2\*I\*b\*x^2))\*sin(-1/8\*pi + 1/4\*arctan2(0, b))))\*cos(2\*a) + ((-I\*gamma(-1/4, 2\*I\*b\*x^2) + I\*gamma(-1/4, -2\*I\*b\*x^2))\*cos(1/8\*pi + 1/4\*arctan2(0, b)) + (-I\*gamma(-1/4, 2\*I\*b\*x^2) + I\*gamma(-1/4, -2\*I\*b\*x^2))\*cos(-1/8\*pi + 1/4\*arctan2(0, b)) + (gamma(-1/4, 2\*I\*b\*x^2) + gamma(-1/4, -2\*I\*b\*x^2))\*sin(1/8\*pi + 1/4\*arctan2(0, b)) - (gamma(-1/4, 2\*I\*b\*x^2) + gamma(-1/4, -2\*I\*b\*x^2))\*sin(-1/8\*pi + 1/4\*arctan2(0, b))))\*sin(2\*a) + 16)/sqrt(x)

**Fricas [A]** time = 1.76632, size = 188, normalized size = 1.61

$$\frac{(2ib)^{\frac{1}{4}} x e^{-2ia} \Gamma\left(\frac{3}{4}, 2i bx^2\right) + (-2ib)^{\frac{1}{4}} x e^{2ia} \Gamma\left(\frac{3}{4}, -2i bx^2\right) - 4\sqrt{x} \cos(bx^2 + a)^2}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^(3/2),x, algorithm="fricas")

[Out] 1/2\*((2\*I\*b)^(1/4)\*x\*e^(-2\*I\*a)\*gamma(3/4, 2\*I\*b\*x^2) + (-2\*I\*b)^(1/4)\*x\*e^(2\*I\*a)\*gamma(3/4, -2\*I\*b\*x^2) - 4\*sqrt(x)\*cos(b\*x^2 + a)^2)/x

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + bx^2)}{x^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*2/x\*\*(3/2),x)

[Out] Integral(cos(a + b\*x\*\*2)\*\*2/x\*\*(3/2), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(bx^2 + a)}{x^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.



```
[In] integrate(cos(b*x^2+a)^2/x^(3/2),x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^2 + a)^2/x^(3/2), x)
```

$$3.34 \quad \int \frac{\cos^2(a+bx^2)}{x^{5/2}} dx$$

**Optimal.** Leaf size=116

$$-\frac{ie^{2ia}b\sqrt{x}\Gamma\left(\frac{1}{4}, -2ibx^2\right)}{3\sqrt[4]{2}\sqrt[4]{-ibx^2}} + \frac{ie^{-2ia}b\sqrt{x}\Gamma\left(\frac{1}{4}, 2ibx^2\right)}{3\sqrt[4]{2}\sqrt[4]{ibx^2}} - \frac{2\cos^2(a+bx^2)}{3x^{3/2}}$$

[Out]  $(-2*\text{Cos}[a + b*x^2]^2)/(3*x^{(3/2)}) - ((I/3)*b*E^{((2*I)*a)*\text{Sqrt}[x]*\Gamma[1/4, (-2*I)*b*x^2]})/(2^{(1/4)}*((-I)*b*x^2)^{(1/4)}) + ((I/3)*b*\text{Sqrt}[x]*\Gamma[1/4, (2*I)*b*x^2])/(2^{(1/4)}*E^{((2*I)*a)*(I*b*x^2)^{(1/4)})}$

**Rubi [A]** time = 0.0997398, antiderivative size = 116, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 6, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3402, 3394, 4573, 3373, 3355, 2208}

$$-\frac{ie^{2ia}b\sqrt{x}\Gamma\left(\frac{1}{4}, -2ibx^2\right)}{3\sqrt[4]{2}\sqrt[4]{-ibx^2}} + \frac{ie^{-2ia}b\sqrt{x}\Gamma\left(\frac{1}{4}, 2ibx^2\right)}{3\sqrt[4]{2}\sqrt[4]{ibx^2}} - \frac{2\cos^2(a+bx^2)}{3x^{3/2}}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[\text{Cos}[a + b*x^2]^2/x^{(5/2)}, x]$

[Out]  $(-2*\text{Cos}[a + b*x^2]^2)/(3*x^{(3/2)}) - ((I/3)*b*E^{((2*I)*a)*\text{Sqrt}[x]*\Gamma[1/4, (-2*I)*b*x^2]})/(2^{(1/4)}*((-I)*b*x^2)^{(1/4)}) + ((I/3)*b*\text{Sqrt}[x]*\Gamma[1/4, (2*I)*b*x^2])/(2^{(1/4)}*E^{((2*I)*a)*(I*b*x^2)^{(1/4)})}$

#### Rule 3402

$\text{Int}[(a_. + \text{Cos}[c_. + (d_.)*(x_)^{(n_)}])*(b_.)^{(p_.)*((e_.)*(x_))^{(m_)}}, x\_Symbol] :> \text{With}[\{k = \text{Denominator}[m]\}, \text{Dist}[k/e, \text{Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*\text{Cos}[c + (d*x^{(k*n)})/e^n])^p, x], x, (e*x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{IntegerQ}[p] \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m]$

#### Rule 3394

$\text{Int}[\text{Cos}[(a_. + (b_.)*(x_)^{(n_)}])^{(p_)}*(x_)^{(m_.)}, x\_Symbol] :> \text{Simp}[(x^{(m+1)}*\text{Cos}[a + b*x^n]^p)/(m+1), x] + \text{Dist}[(b*n*p)/(m+1), \text{Int}[\text{Cos}[a + b*x^n]^{(p-1)}*\text{Sin}[a + b*x^n], x], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{IGtQ}[p, 1] \&\& \text{EqQ}[m+n, 0] \&\& \text{NeQ}[n, 1] \&\& \text{IntegerQ}[n]$

#### Rule 4573

$\text{Int}[\text{Cos}[w_]^{(p_.)*}(u_.)*\text{Sin}[v_]^{(p_.)}, x\_Symbol] :> \text{Dist}[1/2^p, \text{Int}[u*\text{Sin}[2*v]^p, x], x] /; \text{EqQ}[w, v] \&\& \text{IntegerQ}[p]$

#### Rule 3373

$\text{Int}[(a_. + (b_.)*\text{Sin}[u_])^{(p_.)}, x\_Symbol] :> \text{Int}[(a + b*\text{Sin}[\text{ExpandToSum}[u, x]])^p, x] /; \text{FreeQ}[\{a, b, p\}, x] \&\& \text{BinomialQ}[u, x] \&\& !\text{BinomialMatchQ}[u, x]$

#### Rule 3355

$\text{Int}[\text{Sin}[(c_. + (d_.)*((e_. + (f_.)*(x_))^{(n_)}], x\_Symbol] :> \text{Dist}[I/2, \text{Int}[E^{-(c*I) - d*I*(e + f*x)^n}, x], x] - \text{Dist}[I/2, \text{Int}[E^{(c*I + d*I*(e + f*x)^n}, x], x]$

$x)^n), x], x] /; \text{FreeQ}\{c, d, e, f\}, x] \&\& \text{IGtQ}[n, 2]$

### Rule 2208

$\text{Int}[(F\_)^{(a\_)} + (b\_)*((c\_)+ (d\_)*(x\_))^n], x\_Symbol] \text{:} > -\text{Simp}[(F^a * (c + d*x)*\text{Gamma}[1/n, -(b*(c + d*x)^n*\text{Log}[F])]) / (d*n*(-(b*(c + d*x)^n*\text{Log}[F]))^{(1/n)}), x] /; \text{FreeQ}\{F, a, b, c, d, n\}, x] \&\& !\text{IntegerQ}[2/n]$

### Rubi steps

$$\begin{aligned} \int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx &= 2 \text{Subst} \left( \int \frac{\cos^2(a + bx^4)}{x^4} dx, x, \sqrt{x} \right) \\ &= -\frac{2 \cos^2(a + bx^2)}{3x^{3/2}} - \frac{1}{3}(16b) \text{Subst} \left( \int \cos(a + bx^4) \sin(a + bx^4) dx, x, \sqrt{x} \right) \\ &= -\frac{2 \cos^2(a + bx^2)}{3x^{3/2}} - \frac{1}{3}(8b) \text{Subst} \left( \int \sin(2(a + bx^4)) dx, x, \sqrt{x} \right) \\ &= -\frac{2 \cos^2(a + bx^2)}{3x^{3/2}} - \frac{1}{3}(8b) \text{Subst} \left( \int \sin(2a + 2bx^4) dx, x, \sqrt{x} \right) \\ &= -\frac{2 \cos^2(a + bx^2)}{3x^{3/2}} - \frac{1}{3}(4ib) \text{Subst} \left( \int e^{-2ia-2ibx^4} dx, x, \sqrt{x} \right) + \frac{1}{3}(4ib) \text{Subst} \left( \int e^{2ia+2ibx^4} dx, x, \sqrt{x} \right) \\ &= -\frac{2 \cos^2(a + bx^2)}{3x^{3/2}} - \frac{ibe^{2ia} \sqrt{x} \Gamma\left(\frac{1}{4}, -2ibx^2\right)}{3\sqrt[4]{2}\sqrt[4]{-ibx^2}} + \frac{ibe^{-2ia} \sqrt{x} \Gamma\left(\frac{1}{4}, 2ibx^2\right)}{3\sqrt[4]{2}\sqrt[4]{ibx^2}} \end{aligned}$$

**Mathematica [A]** time = 0.368087, size = 137, normalized size = 1.18

$$\frac{2^{3/4}bx^2\sqrt[4]{ibx^2}(\sin(2a) - i\cos(2a))\text{Gamma}\left(\frac{1}{4}, -2ibx^2\right) + i2^{3/4}(-ibx^2)^{5/4}(\sin(2a) + i\cos(2a))\text{Gamma}\left(\frac{1}{4}, 2ibx^2\right) - 4\sqrt[4]{b^2x^4}}{6x^{3/2}\sqrt[4]{b^2x^4}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^2]^2/x^(5/2), x]

[Out]  $(-4*(b^2*x^4)^{(1/4)}*\text{Cos}[a + b*x^2]^2 + 2^{(3/4)}*b*x^2*(I*b*x^2)^{(1/4)}*\text{Gamma}[1/4, (-2*I)*b*x^2]*((-I)*\text{Cos}[2*a] + \text{Sin}[2*a]) + I*2^{(3/4)}*((-I)*b*x^2)^{(5/4)}*\text{Gamma}[1/4, (2*I)*b*x^2]*(I*\text{Cos}[2*a] + \text{Sin}[2*a]))/(6*x^{(3/2)}*(b^2*x^4)^{(1/4)})$

**Maple [F]** time = 0.092, size = 0, normalized size = 0.

$$\int (\cos(bx^2 + a))^2 x^{-5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b\*x^2+a)^2/x^(5/2), x)

[Out] int(cos(b\*x^2+a)^2/x^(5/2), x)

**Maxima [B]** time = 1.44658, size = 373, normalized size = 3.22

$$\frac{2^{\frac{3}{4}} (x^2|b|)^{\frac{3}{4}} \left( \left( 3 \Gamma\left(-\frac{3}{4}, 2i bx^2\right) + \Gamma\left(-\frac{3}{4}, -2i bx^2\right) \right) \cos\left(\frac{3}{8}\pi + \frac{3}{4}\arctan(0, b)\right) + 3 \left( \Gamma\left(-\frac{3}{4}, 2i bx^2\right) + \Gamma\left(-\frac{3}{4}, -2i bx^2\right) \right) \cos\right)}{}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^(5/2),x, algorithm="maxima")

[Out] 
$$\begin{aligned} & -1/48*(2^{(3/4)}*(x^2*\text{abs}(b))^{(3/4)}*((3*(\text{gamma}(-3/4, 2*I*b*x^2) + \text{gamma}(-3/4, \\ & -2*I*b*x^2))*\cos(3/8*\text{pi} + 3/4*\arctan2(0, b)) + 3*(\text{gamma}(-3/4, 2*I*b*x^2) + \\ & \text{gamma}(-3/4, -2*I*b*x^2))*\cos(-3/8*\text{pi} + 3/4*\arctan2(0, b)) + (3*I*\text{gamma}(-3/ \\ & 4, 2*I*b*x^2) - 3*I*\text{gamma}(-3/4, -2*I*b*x^2))*\sin(3/8*\text{pi} + 3/4*\arctan2(0, b) \\ & ) + (-3*I*\text{gamma}(-3/4, 2*I*b*x^2) + 3*I*\text{gamma}(-3/4, -2*I*b*x^2))*\sin(-3/8*\text{pi} \\ & + 3/4*\arctan2(0, b)))*\cos(2*a) + ((-3*I*\text{gamma}(-3/4, 2*I*b*x^2) + 3*I*\text{gamma} \\ & (-3/4, -2*I*b*x^2))*\cos(3/8*\text{pi} + 3/4*\arctan2(0, b)) + (-3*I*\text{gamma}(-3/4, 2*I \\ & *b*x^2) + 3*I*\text{gamma}(-3/4, -2*I*b*x^2))*\cos(-3/8*\text{pi} + 3/4*\arctan2(0, b)) + 3 \\ & *(\text{gamma}(-3/4, 2*I*b*x^2) + \text{gamma}(-3/4, -2*I*b*x^2))*\sin(3/8*\text{pi} + 3/4*\arctan \\ & 2(0, b)) - 3*(\text{gamma}(-3/4, 2*I*b*x^2) + \text{gamma}(-3/4, -2*I*b*x^2))*\sin(-3/8*\text{pi} \\ & + 3/4*\arctan2(0, b)))*\sin(2*a) + 16)/x^{(3/2)} \end{aligned}$$

**Fricas [A]** time = 1.74185, size = 196, normalized size = 1.69

$$\frac{(2i b)^{\frac{3}{4}} x^2 e^{-2ia} \Gamma\left(\frac{1}{4}, 2i bx^2\right) + (-2i b)^{\frac{3}{4}} x^2 e^{2ia} \Gamma\left(\frac{1}{4}, -2i bx^2\right) - 4 \sqrt{x} \cos(bx^2 + a)^2}{6x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x^2+a)^2/x^(5/2),x, algorithm="fricas")

[Out] 
$$\frac{1}{6} * ((2*I*b)^{(3/4)} * x^2 * e^{(-2*I*a)} * \text{gamma}(1/4, 2*I*b*x^2) + (-2*I*b)^{(3/4)} * x^2 * e^{(2*I*a)} * \text{gamma}(1/4, -2*I*b*x^2) - 4 * \text{sqrt}(x) * \cos(b*x^2 + a)^2) / x^2$$

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b\*x\*\*2+a)\*\*2/x\*\*(5/2),x)

[Out] Timed out

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^2 + a)^2}{x^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x^2+a)^2/x^(5/2),x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^2 + a)^2/x^(5/2), x)
```

### 3.35 $\int \cos\left(a + \frac{b}{x}\right) dx$

**Optimal.** Leaf size=31

$$b \sin(a) \operatorname{CosIntegral}\left(\frac{b}{x}\right) + b \cos(a) \operatorname{Si}\left(\frac{b}{x}\right) + x \cos\left(a + \frac{b}{x}\right)$$

[Out] x\*Cos[a + b/x] + b\*CosIntegral[b/x]\*Sin[a] + b\*Cos[a]\*SinIntegral[b/x]

**Rubi [A]** time = 0.0723927, antiderivative size = 31, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3362, 3297, 3303, 3299, 3302}

$$b \sin(a) \operatorname{CosIntegral}\left(\frac{b}{x}\right) + b \cos(a) \operatorname{Si}\left(\frac{b}{x}\right) + x \cos\left(a + \frac{b}{x}\right)$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b/x], x]

[Out] x\*Cos[a + b/x] + b\*CosIntegral[b/x]\*Sin[a] + b\*Cos[a]\*SinIntegral[b/x]

#### Rule 3362

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])*(b_.))^(p_.), x_Symbol]
:> Dist[1/(n*f), Subst[Int[x^(1/n - 1)*(a + b*Cos[c + d*x])^p, x], x, (e + f*x)^n], x]
;/; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

#### Rule 3297

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol]
:> Simp[(c + d*x)^(m + 1)*Sin[e + f*x]/(d*(m + 1)), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x]
;/; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]
```

#### Rule 3303

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Dist[Cos[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/(c + d*x), x], x]
;/; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

#### Rule 3299

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[SinIntegral[e + f*x]/d, x]
;/; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

#### Rule 3302

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[CosIntegral[e - Pi/2 + f*x]/d, x]
;/; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

#### Rubi steps

$$\begin{aligned}
\int \cos\left(a + \frac{b}{x}\right) dx &= -\text{Subst}\left(\int \frac{\cos(a + bx)}{x^2} dx, x, \frac{1}{x}\right) \\
&= x \cos\left(a + \frac{b}{x}\right) + b \text{Subst}\left(\int \frac{\sin(a + bx)}{x} dx, x, \frac{1}{x}\right) \\
&= x \cos\left(a + \frac{b}{x}\right) + (b \cos(a)) \text{Subst}\left(\int \frac{\sin(bx)}{x} dx, x, \frac{1}{x}\right) + (b \sin(a)) \text{Subst}\left(\int \frac{\cos(bx)}{x} dx, x, \frac{1}{x}\right) \\
&= x \cos\left(a + \frac{b}{x}\right) + b \text{Ci}\left(\frac{b}{x}\right) \sin(a) + b \cos(a) \text{Si}\left(\frac{b}{x}\right)
\end{aligned}$$

**Mathematica [A]** time = 0.030921, size = 31, normalized size = 1.

$$b \sin(a) \text{CosIntegral}\left(\frac{b}{x}\right) + b \cos(a) \text{Si}\left(\frac{b}{x}\right) + x \cos\left(a + \frac{b}{x}\right)$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b/x], x]

[Out] x\*Cos[a + b/x] + b\*CosIntegral[b/x]\*Sin[a] + b\*Cos[a]\*SinIntegral[b/x]

**Maple [A]** time = 0.037, size = 39, normalized size = 1.3

$$-b \left( -\frac{x}{b} \cos\left(a + \frac{b}{x}\right) - \text{Si}\left(\frac{b}{x}\right) \cos(a) - \text{Ci}\left(\frac{b}{x}\right) \sin(a) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x), x)

[Out] -b\*(-cos(a+b/x)\*x/b - Si(b/x)\*cos(a) - Ci(b/x)\*sin(a))

**Maxima [C]** time = 1.28664, size = 77, normalized size = 2.48

$$\frac{1}{2} \left( \left( -i \text{Ei}\left(\frac{ib}{x}\right) + i \text{Ei}\left(-\frac{ib}{x}\right) \right) \cos(a) + \left( \text{Ei}\left(\frac{ib}{x}\right) + \text{Ei}\left(-\frac{ib}{x}\right) \right) \sin(a) \right) b + x \cos\left(\frac{ax + b}{x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x), x, algorithm="maxima")

[Out] 1/2\*((-I\*Ei(I\*b/x) + I\*Ei(-I\*b/x))\*cos(a) + (Ei(I\*b/x) + Ei(-I\*b/x))\*sin(a))\*b + x\*cos((a\*x + b)/x)

**Fricas [A]** time = 1.59511, size = 144, normalized size = 4.65

$$b \cos(a) \text{Si}\left(\frac{b}{x}\right) + x \cos\left(\frac{ax + b}{x}\right) + \frac{1}{2} \left( b \text{Ci}\left(\frac{b}{x}\right) + b \text{Ci}\left(-\frac{b}{x}\right) \right) \sin(a)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b/x),x, algorithm="fricas")
```

```
[Out] b*cos(a)*sin_integral(b/x) + x*cos((a*x + b)/x) + 1/2*(b*cos_integral(b/x)
+ b*cos_integral(-b/x))*sin(a)
```

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos\left(a + \frac{b}{x}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b/x),x)
```

```
[Out] Integral(cos(a + b/x), x)
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos\left(a + \frac{b}{x}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b/x),x, algorithm="giac")
```

```
[Out] integrate(cos(a + b/x), x)
```



$$3.36 \quad \int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx$$

**Optimal.** Leaf size=20

$$\sin(a)\text{Si}\left(\frac{b}{x}\right) - \cos(a)\text{CosIntegral}\left(\frac{b}{x}\right)$$

[Out] -(Cos[a]\*CosIntegral[b/x]) + Sin[a]\*SinIntegral[b/x]

**Rubi [A]** time = 0.0273317, antiderivative size = 20, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3378, 3376, 3375}

$$\sin(a)\text{Si}\left(\frac{b}{x}\right) - \cos(a)\text{CosIntegral}\left(\frac{b}{x}\right)$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b/x]/x, x]

[Out] -(Cos[a]\*CosIntegral[b/x]) + Sin[a]\*SinIntegral[b/x]

#### Rule 3378

Int[Cos[(c\_) + (d\_)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Dist[Cos[c], Int[Cos[d\*x^n]/x, x], x] - Dist[Sin[c], Int[Sin[d\*x^n]/x, x], x] /; FreeQ[{c, d, n}, x]

#### Rule 3376

Int[Cos[(d\_)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[CosIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

#### Rule 3375

Int[Sin[(d\_)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[SinIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

#### Rubi steps

$$\begin{aligned} \int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx &= \cos(a) \int \frac{\cos\left(\frac{b}{x}\right)}{x} dx - \sin(a) \int \frac{\sin\left(\frac{b}{x}\right)}{x} dx \\ &= -\cos(a)\text{Ci}\left(\frac{b}{x}\right) + \sin(a)\text{Si}\left(\frac{b}{x}\right) \end{aligned}$$

**Mathematica [A]** time = 0.04694, size = 20, normalized size = 1.

$$\sin(a)\text{Si}\left(\frac{b}{x}\right) - \cos(a)\text{CosIntegral}\left(\frac{b}{x}\right)$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b/x]/x,x]

[Out] -(Cos[a]\*CosIntegral[b/x]) + Sin[a]\*SinIntegral[b/x]

**Maple [A]** time = 0.033, size = 21, normalized size = 1.1

$$-\text{Ci}\left(\frac{b}{x}\right)\cos(a) + \text{Si}\left(\frac{b}{x}\right)\sin(a)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x)/x,x)

[Out] -Ci(b/x)\*cos(a)+Si(b/x)\*sin(a)

**Maxima [C]** time = 1.4209, size = 58, normalized size = 2.9

$$-\frac{1}{2}\left(\text{Ei}\left(\frac{ib}{x}\right) + \text{Ei}\left(-\frac{ib}{x}\right)\right)\cos(a) - \frac{1}{2}\left(i\text{Ei}\left(\frac{ib}{x}\right) - i\text{Ei}\left(-\frac{ib}{x}\right)\right)\sin(a)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x,x, algorithm="maxima")

[Out] -1/2\*(Ei(I\*b/x) + Ei(-I\*b/x))\*cos(a) - 1/2\*(I\*Ei(I\*b/x) - I\*Ei(-I\*b/x))\*sin(a)

**Fricas [A]** time = 1.57274, size = 109, normalized size = 5.45

$$-\frac{1}{2}\left(\text{Ci}\left(\frac{b}{x}\right) + \text{Ci}\left(-\frac{b}{x}\right)\right)\cos(a) + \sin(a)\text{Si}\left(\frac{b}{x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x,x, algorithm="fricas")

[Out] -1/2\*(cos\_integral(b/x) + cos\_integral(-b/x))\*cos(a) + sin(a)\*sin\_integral(b/x)

**Sympy [A]** time = 1.9153, size = 15, normalized size = 0.75

$$\sin(a)\text{Si}\left(\frac{b}{x}\right) - \cos(a)\text{Ci}\left(\frac{b}{x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x,x)

[Out]  $\sin(a)*\text{Si}(b/x) - \cos(a)*\text{Ci}(b/x)$

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b/x)/x,x, algorithm="giac")`

[Out] `integrate(cos(a + b/x)/x, x)`

$$3.37 \quad \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx$$

**Optimal.** Leaf size=13

$$-\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

[Out] -(Sin[a + b/x]/b)

**Rubi [A]** time = 0.0144733, antiderivative size = 13, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3380, 2637}

$$-\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b/x]/x^2,x]

[Out] -(Sin[a + b/x]/b)

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
  := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p,
    x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
  && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 2637

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
  FreeQ[{c, d}, x]
```

#### Rubi steps

$$\begin{aligned} \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx &= -\text{Subst}\left(\int \cos(a + bx) dx, x, \frac{1}{x}\right) \\ &= -\frac{\sin\left(a + \frac{b}{x}\right)}{b} \end{aligned}$$

**Mathematica [A]** time = 0.0028499, size = 13, normalized size = 1.

$$-\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b/x]/x^2,x]

[Out] -(Sin[a + b/x]/b)

**Maple [A]** time = 0.026, size = 14, normalized size = 1.1

$$-\frac{1}{b} \sin\left(a + \frac{b}{x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x)/x^2,x)

[Out] -sin(a+b/x)/b

**Maxima [A]** time = 1.055, size = 18, normalized size = 1.38

$$\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x^2,x, algorithm="maxima")

[Out] -sin(a + b/x)/b

**Fricas [A]** time = 1.51144, size = 28, normalized size = 2.15

$$-\frac{\sin\left(\frac{ax+b}{x}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x^2,x, algorithm="fricas")

[Out] -sin((a\*x + b)/x)/b

**Sympy [A]** time = 1.7594, size = 15, normalized size = 1.15

$$\begin{cases} -\frac{\sin\left(a + \frac{b}{x}\right)}{b} & \text{for } b \neq 0 \\ -\frac{\cos(a)}{x} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x\*\*2,x)

```
[Out] Piecewise((-sin(a + b/x)/b, Ne(b, 0)), (-cos(a)/x, True))
```

---

**Giac [A]** time = 1.13324, size = 18, normalized size = 1.38

$$-\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b/x)/x^2,x, algorithm="giac")
```

```
[Out] -sin(a + b/x)/b
```

$$3.38 \quad \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx$$

**Optimal.** Leaf size=30

$$-\frac{\cos\left(a + \frac{b}{x}\right)}{b^2} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx}$$

[Out] -(Cos[a + b/x]/b^2) - Sin[a + b/x]/(b\*x)

**Rubi [A]** time = 0.0249832, antiderivative size = 30, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3380, 3296, 2638}

$$-\frac{\cos\left(a + \frac{b}{x}\right)}{b^2} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b/x]/x^3,x]

[Out] -(Cos[a + b/x]/b^2) - Sin[a + b/x]/(b\*x)

#### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

#### Rule 3296

Int[((c\_.) + (d\_.)\*(x\_)^(m\_.))\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] := -Simp[(c + d\*x)^m\*Cos[e + f\*x]/f, x] + Dist[(d\*m)/f, Int[(c + d\*x)^(m - 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

#### Rule 2638

Int[sin[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] := -Simp[Cos[c + d\*x]/d, x] /; FreeQ[{c, d}, x]

#### Rubi steps

$$\begin{aligned} \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx &= -\text{Subst}\left(\int x \cos(a + bx) dx, x, \frac{1}{x}\right) \\ &= -\frac{\sin\left(a + \frac{b}{x}\right)}{bx} + \frac{\text{Subst}\left(\int \sin(a + bx) dx, x, \frac{1}{x}\right)}{b} \\ &= -\frac{\cos\left(a + \frac{b}{x}\right)}{b^2} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx} \end{aligned}$$

**Mathematica [A]** time = 0.047531, size = 29, normalized size = 0.97

$$-\frac{b \sin\left(a + \frac{b}{x}\right) + x \cos\left(a + \frac{b}{x}\right)}{b^2 x}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b/x]/x^3,x]

[Out] -((x\*cos[a + b/x] + b\*sin[a + b/x])/(b^2\*x))

**Maple [A]** time = 0.032, size = 42, normalized size = 1.4

$$-\frac{1}{b^2} \left( \cos\left(a + \frac{b}{x}\right) + \left(a + \frac{b}{x}\right) \sin\left(a + \frac{b}{x}\right) - a \sin\left(a + \frac{b}{x}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x)/x^3,x)

[Out] -1/b^2\*(cos(a+b/x)+(a+b/x)\*sin(a+b/x)-a\*sin(a+b/x))

**Maxima [C]** time = 1.29148, size = 69, normalized size = 2.3

$$-\frac{\left(\Gamma\left(2, \frac{ib}{x}\right) + \Gamma\left(2, -\frac{ib}{x}\right)\right) \cos(a) - \left(i\Gamma\left(2, \frac{ib}{x}\right) - i\Gamma\left(2, -\frac{ib}{x}\right)\right) \sin(a)}{2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x^3,x, algorithm="maxima")

[Out] -1/2\*((gamma(2, I\*b/x) + gamma(2, -I\*b/x))\*cos(a) - (I\*gamma(2, I\*b/x) - I\*gamma(2, -I\*b/x))\*sin(a))/b^2

**Fricas [A]** time = 1.57816, size = 70, normalized size = 2.33

$$-\frac{x \cos\left(\frac{ax+b}{x}\right) + b \sin\left(\frac{ax+b}{x}\right)}{b^2 x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x^3,x, algorithm="fricas")

[Out] -(x\*cos((a\*x + b)/x) + b\*sin((a\*x + b)/x))/(b^2\*x)

**Sympy [A]** time = 3.43995, size = 31, normalized size = 1.03

$$\begin{cases} -\frac{\sin\left(a + \frac{b}{x}\right)}{bx} - \frac{\cos\left(a + \frac{b}{x}\right)}{b^2} & \text{for } b \neq 0 \\ -\frac{\cos(a)}{2x^2} & \text{otherwise} \end{cases}$$



Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b/x)/x**3,x)
```

```
[Out] Piecewise((-sin(a + b/x)/(b*x) - cos(a + b/x)/b**2, Ne(b, 0)), (-cos(a)/(2*x**2), True))
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b/x)/x^3,x, algorithm="giac")
```

```
[Out] integrate(cos(a + b/x)/x^3, x)
```

$$3.39 \quad \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx$$

**Optimal.** Leaf size=46

$$\frac{2 \sin\left(a + \frac{b}{x}\right)}{b^3} - \frac{2 \cos\left(a + \frac{b}{x}\right)}{b^2 x} - \frac{\sin\left(a + \frac{b}{x}\right)}{b x^2}$$

[Out]  $(-2*\text{Cos}[a + b/x])/(b^2*x) + (2*\text{Sin}[a + b/x])/b^3 - \text{Sin}[a + b/x]/(b*x^2)$

**Rubi [A]** time = 0.0467186, antiderivative size = 46, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 3, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3380, 3296, 2637}

$$\frac{2 \sin\left(a + \frac{b}{x}\right)}{b^3} - \frac{2 \cos\left(a + \frac{b}{x}\right)}{b^2 x} - \frac{\sin\left(a + \frac{b}{x}\right)}{b x^2}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[\text{Cos}[a + b/x]/x^4, x]$

[Out]  $(-2*\text{Cos}[a + b/x])/(b^2*x) + (2*\text{Sin}[a + b/x])/b^3 - \text{Sin}[a + b/x]/(b*x^2)$

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
 := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p,
x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
 && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 3296

```
Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol]
 := -Simp[(c + d*x)^m*Cos[e + f*x]/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[e + f*x],
x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

#### Rule 2637

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol]
 := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx &= -\text{Subst}\left(\int x^2 \cos(a + bx) dx, x, \frac{1}{x}\right) \\
&= -\frac{\sin\left(a + \frac{b}{x}\right)}{bx^2} + \frac{2 \text{Subst}\left(\int x \sin(a + bx) dx, x, \frac{1}{x}\right)}{b} \\
&= -\frac{2 \cos\left(a + \frac{b}{x}\right)}{b^2 x} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2} + \frac{2 \text{Subst}\left(\int \cos(a + bx) dx, x, \frac{1}{x}\right)}{b^2} \\
&= -\frac{2 \cos\left(a + \frac{b}{x}\right)}{b^2 x} + \frac{2 \sin\left(a + \frac{b}{x}\right)}{b^3} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2}
\end{aligned}$$

**Mathematica [A]** time = 0.0041019, size = 46, normalized size = 1.

$$\frac{2 \sin\left(a + \frac{b}{x}\right)}{b^3} - \frac{2 \cos\left(a + \frac{b}{x}\right)}{b^2 x} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b/x]/x^4,x]

[Out] (-2\*Cos[a + b/x])/(b^2\*x) + (2\*Sin[a + b/x])/b^3 - Sin[a + b/x]/(b\*x^2)

**Maple [A]** time = 0.033, size = 92, normalized size = 2.

$$-\frac{1}{b^3} \left( \left( a + \frac{b}{x} \right)^2 \sin\left(a + \frac{b}{x}\right) - 2 \sin\left(a + \frac{b}{x}\right) + 2 \left( a + \frac{b}{x} \right) \cos\left(a + \frac{b}{x}\right) - 2a \left( \cos\left(a + \frac{b}{x}\right) + \left( a + \frac{b}{x} \right) \sin\left(a + \frac{b}{x}\right) \right) + a^2 \sin\left(a + \frac{b}{x}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x)/x^4,x)

[Out] -1/b^3\*((a+b/x)^2\*sin(a+b/x)-2\*sin(a+b/x)+2\*(a+b/x)\*cos(a+b/x)-2\*a\*(cos(a+b/x)+(a+b/x)\*sin(a+b/x))+a^2\*sin(a+b/x))

**Maxima [C]** time = 1.36795, size = 68, normalized size = 1.48

$$\frac{\left(i\Gamma\left(3, \frac{ib}{x}\right) - i\Gamma\left(3, -\frac{ib}{x}\right)\right) \cos(a) + \left(\Gamma\left(3, \frac{ib}{x}\right) + \Gamma\left(3, -\frac{ib}{x}\right)\right) \sin(a)}{2b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x^4,x, algorithm="maxima")

[Out] 1/2\*((I\*gamma(3, I\*b/x) - I\*gamma(3, -I\*b/x))\*cos(a) + (gamma(3, I\*b/x) + gamma(3, -I\*b/x))\*sin(a))/b^3

**Fricas [A]** time = 1.51528, size = 95, normalized size = 2.07

$$\frac{2bx \cos\left(\frac{ax+b}{x}\right) + (b^2 - 2x^2) \sin\left(\frac{ax+b}{x}\right)}{b^3x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x^4,x, algorithm="fricas")

[Out] -(2\*b\*x\*cos((a\*x + b)/x) + (b^2 - 2\*x^2)\*sin((a\*x + b)/x))/(b^3\*x^2)

**Sympy [A]** time = 6.18221, size = 46, normalized size = 1.

$$\begin{cases} \frac{\sin\left(a+\frac{b}{x}\right)}{bx^2} - \frac{2\cos\left(a+\frac{b}{x}\right)}{b^2x} + \frac{2\sin\left(a+\frac{b}{x}\right)}{b^3} & \text{for } b \neq 0 \\ -\frac{\cos(a)}{3x^3} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x\*\*4,x)

[Out] Piecewise((-sin(a + b/x)/(b\*x\*\*2) - 2\*cos(a + b/x)/(b\*\*2\*x) + 2\*sin(a + b/x)/b\*\*3, Ne(b, 0)), (-cos(a)/(3\*x\*\*3), True))

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x)/x^4,x, algorithm="giac")

[Out] integrate(cos(a + b/x)/x^4, x)

### 3.40 $\int \cos\left(a + \frac{b}{x^2}\right) dx$

**Optimal.** Leaf size=79

$$\sqrt{2\pi}\sqrt{b}\sin(a)\text{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}}\sqrt{b}}{x}\right) + \sqrt{2\pi}\sqrt{b}\cos(a)S\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) + x\cos\left(a + \frac{b}{x^2}\right)$$

[Out] x\*Cos[a + b/x^2] + Sqrt[b]\*Sqrt[2\*Pi]\*Cos[a]\*FresnelS[(Sqrt[b]\*Sqrt[2/Pi])/x] + Sqrt[b]\*Sqrt[2\*Pi]\*FresnelC[(Sqrt[b]\*Sqrt[2/Pi])/x]\*Sin[a]

**Rubi [A]** time = 0.0495695, antiderivative size = 79, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3360, 3388, 3353, 3352, 3351}

$$\sqrt{2\pi}\sqrt{b}\sin(a)\text{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}}\sqrt{b}}{x}\right) + \sqrt{2\pi}\sqrt{b}\cos(a)S\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) + x\cos\left(a + \frac{b}{x^2}\right)$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b/x^2], x]

[Out] x\*Cos[a + b/x^2] + Sqrt[b]\*Sqrt[2\*Pi]\*Cos[a]\*FresnelS[(Sqrt[b]\*Sqrt[2/Pi])/x] + Sqrt[b]\*Sqrt[2\*Pi]\*FresnelC[(Sqrt[b]\*Sqrt[2/Pi])/x]\*Sin[a]

#### Rule 3360

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)]\*(b\_.))^(p\_.), x\_Symbol] :> -Dist[f^(-1), Subst[Int[(a + b\*Cos[c + d/x^n])^p/x^2, x], x, 1/(e + f\*x)], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[n, 0] && EqQ[n, -2]

#### Rule 3388

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)]\*((e\_.)\*(x\_))^(m\_), x\_Symbol] :> Simp[((e\*x)^(m + 1)\*Cos[c + d\*x^n])/(e\*(m + 1)), x] + Dist[(d\*n)/(e^(m + 1)), Int[(e\*x)^(m + n)\*Sin[c + d\*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[m, -1]

#### Rule 3353

Int[Sin[(c\_) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(2)], x\_Symbol] :> Dist[Sin[c], Int[Cos[d\*(e + f\*x)^2], x], x] + Dist[Cos[c], Int[Sin[d\*(e + f\*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]

#### Rule 3352

Int[Cos[(d\_.)\*((e\_.) + (f\_.)\*(x\_))^(2)], x\_Symbol] :> Simp[(Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)])/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rule 3351

Int[Sin[(d\_.)\*((e\_.) + (f\_.)\*(x\_))^(2)], x\_Symbol] :> Simp[(Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)])/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

Rubi steps

$$\begin{aligned}
\int \cos\left(a + \frac{b}{x^2}\right) dx &= -\text{Subst}\left(\int \frac{\cos(a + bx^2)}{x^2} dx, x, \frac{1}{x}\right) \\
&= x \cos\left(a + \frac{b}{x^2}\right) + (2b) \text{Subst}\left(\int \sin(a + bx^2) dx, x, \frac{1}{x}\right) \\
&= x \cos\left(a + \frac{b}{x^2}\right) + (2b \cos(a)) \text{Subst}\left(\int \sin(bx^2) dx, x, \frac{1}{x}\right) + (2b \sin(a)) \text{Subst}\left(\int \cos(bx^2) dx, x, \frac{1}{x}\right) \\
&= x \cos\left(a + \frac{b}{x^2}\right) + \sqrt{b}\sqrt{2\pi} \cos(a) S\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) + \sqrt{b}\sqrt{2\pi} C\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \sin(a)
\end{aligned}$$

**Mathematica [A]** time = 0.133651, size = 80, normalized size = 1.01

$$\sqrt{2\pi}\sqrt{b}\left(\sin(a)\text{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}}\sqrt{b}}{x}\right) + \cos(a)S\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)\right) - x \sin(a) \sin\left(\frac{b}{x^2}\right) + x \cos(a) \cos\left(\frac{b}{x^2}\right)$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b/x^2], x]

[Out] x\*Cos[a]\*Cos[b/x^2] + Sqrt[b]\*Sqrt[2\*Pi]\*(Cos[a]\*FresnelS[(Sqrt[b]\*Sqrt[2/Pi])/x] + FresnelC[(Sqrt[b]\*Sqrt[2/Pi])/x]\*Sin[a]) - x\*Sin[a]\*Sin[b/x^2]

**Maple [A]** time = 0.033, size = 57, normalized size = 0.7

$$x \cos\left(a + \frac{b}{x^2}\right) + \sqrt{b}\sqrt{2}\sqrt{\pi}\left(\cos(a) \text{FresnelS}\left(\frac{\sqrt{2}}{\sqrt{\pi x}}\sqrt{b}\right) + \sin(a) \text{FresnelC}\left(\frac{\sqrt{2}}{\sqrt{\pi x}}\sqrt{b}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x^2), x)

[Out] x\*cos(a+b/x^2)+b^(1/2)\*2^(1/2)\*Pi^(1/2)\*(cos(a)\*FresnelS(b^(1/2)\*2^(1/2)/Pi^(1/2)/x)+sin(a)\*FresnelC(b^(1/2)\*2^(1/2)/Pi^(1/2)/x)

**Maxima [C]** time = 1.41774, size = 533, normalized size = 6.75

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2), x, algorithm="maxima")

[Out] 1/4\*(4\*x^2\*sqrt(abs(b)/x^2)\*cos((a\*x^2 + b)/x^2) + (((I\*sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) - I\*sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + (I\*sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) - I\*sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + (sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) + sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*sin(1/4\*pi + 1/2\*arctan2(0

, b)) - (sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) + sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*sin(-1/4\*pi + 1/2\*arctan2(0, b))\*cos(a) + ((sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) + sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + (sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) + sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + (-I\*sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) + I\*sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + (I\*sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) - I\*sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*sin(-1/4\*pi + 1/2\*arctan2(0, b))\*sin(a))\*b)/(x\*sqrt(abs(b)/x^2))

**Fricas [A]** time = 1.68905, size = 208, normalized size = 2.63

$$\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)S\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) + \sqrt{2}\pi\sqrt{\frac{b}{\pi}}C\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right)\sin(a) + x\cos\left(\frac{ax^2 + b}{x^2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2),x, algorithm="fricas")

[Out] sqrt(2)\*pi\*sqrt(b/pi)\*cos(a)\*fresnel\_sin(sqrt(2)\*sqrt(b/pi)/x) + sqrt(2)\*pi\*sqrt(b/pi)\*fresnel\_cos(sqrt(2)\*sqrt(b/pi)/x)\*sin(a) + x\*cos((a\*x^2 + b)/x^2)

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos\left(a + \frac{b}{x^2}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x\*\*2),x)

[Out] Integral(cos(a + b/x\*\*2), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos\left(a + \frac{b}{x^2}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2),x, algorithm="giac")

[Out] integrate(cos(a + b/x^2), x)

$$3.41 \quad \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$$

**Optimal.** Leaf size=25

$$\frac{1}{2} \sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right) - \frac{1}{2} \cos(a) \operatorname{CosIntegral}\left(\frac{b}{x^2}\right)$$

[Out]  $-(\operatorname{Cos}[a] * \operatorname{CosIntegral}[b/x^2])/2 + (\operatorname{Sin}[a] * \operatorname{SinIntegral}[b/x^2])/2$

**Rubi [A]** time = 0.0277989, antiderivative size = 25, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3378, 3376, 3375}

$$\frac{1}{2} \sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right) - \frac{1}{2} \cos(a) \operatorname{CosIntegral}\left(\frac{b}{x^2}\right)$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[\operatorname{Cos}[a + b/x^2]/x, x]$

[Out]  $-(\operatorname{Cos}[a] * \operatorname{CosIntegral}[b/x^2])/2 + (\operatorname{Sin}[a] * \operatorname{SinIntegral}[b/x^2])/2$

#### Rule 3378

$\operatorname{Int}[\operatorname{Cos}[(c_.) + (d_.) * (x_.)^{(n_)}]/(x_), x\_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Cos}[c], \operatorname{Int}[\operatorname{Cos}[d * x^{(n)}]/x, x], x] - \operatorname{Dist}[\operatorname{Sin}[c], \operatorname{Int}[\operatorname{Sin}[d * x^{(n)}]/x, x], x] /; \operatorname{FreeQ}\{c, d, n\}, x]$

#### Rule 3376

$\operatorname{Int}[\operatorname{Cos}[(d_.) * (x_.)^{(n_)}]/(x_), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{CosIntegral}[d * x^{(n)}]/n, x] /; \operatorname{FreeQ}\{d, n\}, x]$

#### Rule 3375

$\operatorname{Int}[\operatorname{Sin}[(d_.) * (x_.)^{(n_)}]/(x_), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{SinIntegral}[d * x^{(n)}]/n, x] /; \operatorname{FreeQ}\{d, n\}, x]$

#### Rubi steps

$$\begin{aligned} \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx &= \cos(a) \int \frac{\cos\left(\frac{b}{x^2}\right)}{x} dx - \sin(a) \int \frac{\sin\left(\frac{b}{x^2}\right)}{x} dx \\ &= -\frac{1}{2} \cos(a) \operatorname{Ci}\left(\frac{b}{x^2}\right) + \frac{1}{2} \sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right) \end{aligned}$$

**Mathematica [A]** time = 0.0452284, size = 24, normalized size = 0.96

$$\frac{1}{2} \left( \sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right) - \cos(a) \operatorname{CosIntegral}\left(\frac{b}{x^2}\right) \right)$$

Antiderivative was successfully verified.



[In] Integrate[Cos[a + b/x^2]/x,x]

[Out]  $(-\text{Cos}[a]*\text{CosIntegral}[b/x^2]) + \text{Sin}[a]*\text{SinIntegral}[b/x^2])/2$

**Maple [A]** time = 0.037, size = 22, normalized size = 0.9

$$-\frac{\cos(a)}{2}\text{Ci}\left(\frac{b}{x^2}\right) + \frac{\sin(a)}{2}\text{Si}\left(\frac{b}{x^2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x^2)/x,x)

[Out]  $-1/2*\text{Ci}(b/x^2)*\cos(a)+1/2*\text{Si}(b/x^2)*\sin(a)$

**Maxima [C]** time = 1.35032, size = 58, normalized size = 2.32

$$-\frac{1}{4}\left(\text{Ei}\left(\frac{ib}{x^2}\right) + \text{Ei}\left(-\frac{ib}{x^2}\right)\right)\cos(a) - \frac{1}{4}\left(i\text{Ei}\left(\frac{ib}{x^2}\right) - i\text{Ei}\left(-\frac{ib}{x^2}\right)\right)\sin(a)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x,x, algorithm="maxima")

[Out]  $-1/4*(\text{Ei}(I*b/x^2) + \text{Ei}(-I*b/x^2))*\cos(a) - 1/4*(I*\text{Ei}(I*b/x^2) - I*\text{Ei}(-I*b/x^2))*\sin(a)$

**Fricas [A]** time = 1.58963, size = 123, normalized size = 4.92

$$-\frac{1}{4}\left(\text{Ci}\left(\frac{b}{x^2}\right) + \text{Ci}\left(-\frac{b}{x^2}\right)\right)\cos(a) + \frac{1}{2}\sin(a)\text{Si}\left(\frac{b}{x^2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x,x, algorithm="fricas")

[Out]  $-1/4*(\cos\_integral(b/x^2) + \cos\_integral(-b/x^2))*\cos(a) + 1/2*\sin(a)*\sin\_integral(b/x^2)$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x\*\*2)/x,x)

[Out] Integral(cos(a + b/x\*\*2)/x, x)

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x,x, algorithm="giac")

[Out] integrate(cos(a + b/x^2)/x, x)

$$3.42 \quad \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$$

**Optimal.** Leaf size=74

$$\frac{\sqrt{\frac{\pi}{2}} \sin(a) S\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}} - \frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}}\sqrt{b}}{x}\right)}{\sqrt{b}}$$

[Out]  $-\left(\frac{\sqrt{\pi/2} \cos[a] \text{FresnelC}[(\sqrt{b} \sqrt{2/\pi})/x]}{\sqrt{b}}\right) + \left(\frac{\sqrt{\pi/2} \text{FresnelS}[(\sqrt{b} \sqrt{2/\pi})/x] \sin[a]}{\sqrt{b}}\right)$

**Rubi [A]** time = 0.0323861, antiderivative size = 74, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3384, 3354, 3352, 3351}

$$\frac{\sqrt{\frac{\pi}{2}} \sin(a) S\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}} - \frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}}\sqrt{b}}{x}\right)}{\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b/x^2]/x^2, x]

[Out]  $-\left(\frac{\sqrt{\pi/2} \cos[a] \text{FresnelC}[(\sqrt{b} \sqrt{2/\pi})/x]}{\sqrt{b}}\right) + \left(\frac{\sqrt{\pi/2} \text{FresnelS}[(\sqrt{b} \sqrt{2/\pi})/x] \sin[a]}{\sqrt{b}}\right)$

#### Rule 3384

Int[Cos[(a\_.) + (b\_.)\*(x\_)^(n\_.)]\*(x\_)^(m\_.), x\_Symbol] :> Dist[2/n, Subst[Int[Cos[a + b\*x^2], x], x, x^(n/2)], x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n/2 - 1]

#### Rule 3354

Int[Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_)^2)], x\_Symbol] :> Dist[Cos[c], Int[Cos[d\*(e + f\*x)^2], x], x] - Dist[Sin[c], Int[Sin[d\*(e + f\*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]

#### Rule 3352

Int[Cos[(d\_.)\*((e\_.) + (f\_.)\*(x\_)^2)], x\_Symbol] :> Simp[(Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)])/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rule 3351

Int[Sin[(d\_.)\*((e\_.) + (f\_.)\*(x\_)^2)], x\_Symbol] :> Simp[(Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)])/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx &= -\text{Subst}\left(\int \cos(a + bx^2) dx, x, \frac{1}{x}\right) \\
&= -\left(\cos(a) \text{Subst}\left(\int \cos(bx^2) dx, x, \frac{1}{x}\right)\right) + \sin(a) \text{Subst}\left(\int \sin(bx^2) dx, x, \frac{1}{x}\right) \\
&= -\frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{C}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}} + \frac{\sqrt{\frac{\pi}{2}} \text{S}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \sin(a)}{\sqrt{b}}
\end{aligned}$$

**Mathematica [A]** time = 0.109059, size = 62, normalized size = 0.84

$$-\frac{\sqrt{\frac{\pi}{2}} \left( \cos(a) \text{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}}\sqrt{b}}{x}\right) - \sin(a) \text{S}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \right)}{\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b/x^2]/x^2,x]

[Out] -((Sqrt[Pi/2]\*(Cos[a]\*FresnelC[(Sqrt[b]\*Sqrt[2/Pi])/x] - FresnelS[(Sqrt[b]\*Sqrt[2/Pi])/x]\*Sin[a]))/Sqrt[b])

**Maple [A]** time = 0.035, size = 48, normalized size = 0.7

$$-\frac{\sqrt{2}\sqrt{\pi}}{2} \left( \cos(a) \text{FresnelC}\left(\frac{\sqrt{2}}{\sqrt{\pi x}}\sqrt{b}\right) - \sin(a) \text{FresnelS}\left(\frac{\sqrt{2}}{\sqrt{\pi x}}\sqrt{b}\right) \right) \frac{1}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x^2)/x^2,x)

[Out] -1/2\*2^(1/2)\*Pi^(1/2)/b^(1/2)\*(cos(a)\*FresnelC(b^(1/2)\*2^(1/2)/Pi^(1/2)/x)-sin(a)\*FresnelS(b^(1/2)\*2^(1/2)/Pi^(1/2)/x))

**Maxima [C]** time = 1.37842, size = 495, normalized size = 6.69

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x^2,x, algorithm="maxima")

[Out] -1/8\*(((sqrt(pi)\*(erf(sqrt(I\*b/x^2))) - 1) + sqrt(pi)\*(erf(sqrt(-I\*b/x^2))) - 1))\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + (sqrt(pi)\*(erf(sqrt(I\*b/x^2))) - 1) + sqrt(pi)\*(erf(sqrt(-I\*b/x^2))) - 1))\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + (-I\*sqrt(pi)\*(erf(sqrt(I\*b/x^2))) - 1) + I\*sqrt(pi)\*(erf(sqrt(-I\*b/x^2))) - 1))\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + (I\*sqrt(pi)\*(erf(sqrt(I\*b/x^2))) - 1) - I\*sqrt(pi)\*(erf(sqrt(-I\*b/x^2))) - 1))\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*cos(a) + ((-I\*sqrt(pi)\*(erf(sqrt(I\*b/x^2))) - 1) + I\*sqrt(pi)\*(erf(sqrt(-I\*b/x^2)))

) - 1))\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + (-I\*sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) + I\*sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - (sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) + sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + (sqrt(pi)\*(erf(sqrt(I\*b/x^2)) - 1) + sqrt(pi)\*(erf(sqrt(-I\*b/x^2)) - 1))\*sin(-1/4\*pi + 1/2\*arctan2(0, b)))\*sin(a))/(x\*sqrt(abs(b)/x^2))

**Fricas [A]** time = 1.60756, size = 186, normalized size = 2.51

$$\frac{\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)C\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) - \sqrt{2}\pi\sqrt{\frac{b}{\pi}}S\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right)\sin(a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x^2,x, algorithm="fricas")

[Out] -1/2\*(sqrt(2)\*pi\*sqrt(b/pi)\*cos(a)\*fresnel\_cos(sqrt(2)\*sqrt(b/pi)/x) - sqrt(2)\*pi\*sqrt(b/pi)\*fresnel\_sin(sqrt(2)\*sqrt(b/pi)/x)\*sin(a))/b

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x\*\*2)/x\*\*2,x)

[Out] Integral(cos(a + b/x\*\*2)/x\*\*2, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x^2,x, algorithm="giac")

[Out] integrate(cos(a + b/x^2)/x^2, x)

$$3.43 \quad \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx$$

**Optimal.** Leaf size=15

$$-\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b}$$

[Out] -Sin[a + b/x^2]/(2\*b)

**Rubi [A]** time = 0.0150678, antiderivative size = 15, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3380, 2637}

$$-\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b/x^2]/x^3, x]

[Out] -Sin[a + b/x^2]/(2\*b)

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
  := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p,
    x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
  && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 2637

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
  FreeQ[{c, d}, x]
```

#### Rubi steps

$$\begin{aligned} \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx &= -\left(\frac{1}{2} \text{Subst}\left(\int \cos(a + bx) dx, x, \frac{1}{x^2}\right)\right) \\ &= -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b} \end{aligned}$$

**Mathematica [A]** time = 0.0031249, size = 15, normalized size = 1.

$$-\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b/x^2]/x^3,x]

[Out] -Sin[a + b/x^2]/(2\*b)

**Maple [A]** time = 0.024, size = 14, normalized size = 0.9

$$-\frac{1}{2b} \sin\left(a + \frac{b}{x^2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b/x^2)/x^3,x)

[Out] -1/2\*sin(a+b/x^2)/b

**Maxima [A]** time = 1.14863, size = 18, normalized size = 1.2

$$-\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x^3,x, algorithm="maxima")

[Out] -1/2\*sin(a + b/x^2)/b

**Fricas [A]** time = 1.55821, size = 39, normalized size = 2.6

$$-\frac{\sin\left(\frac{ax^2+b}{x^2}\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x^3,x, algorithm="fricas")

[Out] -1/2\*sin((a\*x^2 + b)/x^2)/b

**Sympy [A]** time = 6.09938, size = 22, normalized size = 1.47

$$\begin{cases} -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b} & \text{for } b \neq 0 \\ -\frac{\cos(a)}{2x^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x\*\*2)/x\*\*3,x)

```
[Out] Piecewise((-sin(a + b/x**2)/(2*b), Ne(b, 0)), (-cos(a)/(2*x**2), True))
```

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b/x^2)/x^3,x, algorithm="giac")
```

```
[Out] integrate(cos(a + b/x^2)/x^3, x)
```



$$3.44 \quad \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$$

**Optimal.** Leaf size=97

$$\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}} \sqrt{b}}{x}\right)}{2b^{3/2}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) S\left(\frac{\sqrt{b} \sqrt{\frac{2}{\pi}}}{x}\right)}{2b^{3/2}} - \frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx}$$

[Out] (Sqrt[Pi/2]\*Cos[a]\*FresnelS[(Sqrt[b]\*Sqrt[2/Pi])/x])/(2\*b^(3/2)) + (Sqrt[Pi/2]\*FresnelC[(Sqrt[b]\*Sqrt[2/Pi])/x]\*Sin[a])/(2\*b^(3/2)) - Sin[a + b/x^2]/(2\*b\*x)

**Rubi [A]** time = 0.0532183, antiderivative size = 97, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3410, 3386, 3353, 3352, 3351}

$$\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}} \sqrt{b}}{x}\right)}{2b^{3/2}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) S\left(\frac{\sqrt{b} \sqrt{\frac{2}{\pi}}}{x}\right)}{2b^{3/2}} - \frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b/x^2]/x^4, x]

[Out] (Sqrt[Pi/2]\*Cos[a]\*FresnelS[(Sqrt[b]\*Sqrt[2/Pi])/x])/(2\*b^(3/2)) + (Sqrt[Pi/2]\*FresnelC[(Sqrt[b]\*Sqrt[2/Pi])/x]\*Sin[a])/(2\*b^(3/2)) - Sin[a + b/x^2]/(2\*b\*x)

#### Rule 3410

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> -Subst[Int[(a + b\*Cos[c + d\*x^n])^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m] && EqQ[n, -2]

#### Rule 3386

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)]\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> Simp[(e^(n - 1)\*(e\*x)^(m - n + 1)\*Sin[c + d\*x^n])/(d\*n), x] - Dist[(e^n\*(m - n + 1))/(d\*n), Int[(e\*x)^(m - n)\*Sin[c + d\*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[n, m + 1]

#### Rule 3353

Int[Sin[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(2)], x\_Symbol] :> Dist[Sin[c], Int[Cos[d\*(e + f\*x)^2], x], x] + Dist[Cos[c], Int[Sin[d\*(e + f\*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]

#### Rule 3352

Int[Cos[(d\_.)\*((e\_.) + (f\_.)\*(x\_))^(2)], x\_Symbol] :> Simp[(Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)])/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)]]/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rubi steps

$$\begin{aligned} \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx &= -\text{Subst}\left(\int x^2 \cos(a + bx^2) dx, x, \frac{1}{x}\right) \\ &= -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx} + \frac{\text{Subst}\left(\int \sin(a + bx^2) dx, x, \frac{1}{x}\right)}{2b} \\ &= -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx} + \frac{\cos(a) \text{Subst}\left(\int \sin(bx^2) dx, x, \frac{1}{x}\right)}{2b} + \frac{\sin(a) \text{Subst}\left(\int \cos(bx^2) dx, x, \frac{1}{x}\right)}{2b} \\ &= \frac{\sqrt{\frac{\pi}{2}} \cos(a) S\left(\frac{\sqrt{\frac{2}{\pi}} \sqrt{b}}{x}\right)}{2b^{3/2}} + \frac{\sqrt{\frac{\pi}{2}} C\left(\frac{\sqrt{\frac{2}{\pi}} \sqrt{b}}{x}\right) \sin(a)}{2b^{3/2}} - \frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx} \end{aligned}$$

**Mathematica [A]** time = 0.155535, size = 88, normalized size = 0.91

$$\frac{\sqrt{2\pi x} \sin(a) \text{FresnelC}\left(\frac{\sqrt{\frac{2}{\pi}} \sqrt{b}}{x}\right) + \sqrt{2\pi x} \cos(a) S\left(\frac{\sqrt{\frac{2}{\pi}} \sqrt{b}}{x}\right) - 2\sqrt{b} \sin\left(a + \frac{b}{x^2}\right)}{4b^{3/2}x}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b/x^2]/x^4, x]
```

```
[Out] (Sqrt[2*Pi]*x*Cos[a]*FresnelS[(Sqrt[b]*Sqrt[2/Pi])/x] + Sqrt[2*Pi]*x*FresnelC[(Sqrt[b]*Sqrt[2/Pi])/x]*Sin[a] - 2*Sqrt[b]*Sin[a + b/x^2])/(4*b^(3/2)*x)
```

**Maple [A]** time = 0.036, size = 64, normalized size = 0.7

$$-\frac{1}{2bx} \sin\left(a + \frac{b}{x^2}\right) + \frac{\sqrt{2}\sqrt{\pi}}{4} \left( \cos(a) \text{FresnelS}\left(\frac{\sqrt{2}}{\sqrt{\pi x}} \sqrt{b}\right) + \sin(a) \text{FresnelC}\left(\frac{\sqrt{2}}{\sqrt{\pi x}} \sqrt{b}\right) \right) b^{-\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a+b/x^2)/x^4, x)
```

```
[Out] -1/2*sin(a+b/x^2)/b/x+1/4/b^(3/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)/x)+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)/x)
```

**Maxima [C]** time = 1.33186, size = 359, normalized size = 3.7

$$\left( \left( \Gamma\left(\frac{3}{2}, \frac{ib}{x^2}\right) + \Gamma\left(\frac{3}{2}, -\frac{ib}{x^2}\right) \right) \cos\left(\frac{3}{4}\pi + \frac{3}{2} \arctan(0, b)\right) + \left( \Gamma\left(\frac{3}{2}, \frac{ib}{x^2}\right) + \Gamma\left(\frac{3}{2}, -\frac{ib}{x^2}\right) \right) \cos\left(-\frac{3}{4}\pi + \frac{3}{2} \arctan(0, b)\right) - i \Gamma\left(\frac{3}{2}, \frac{ib}{x^2}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x^4,x, algorithm="maxima")

[Out]  $\frac{1}{8} * ((\text{gamma}(3/2, I*b/x^2) + \text{gamma}(3/2, -I*b/x^2)) * \cos(3/4*\pi + 3/2*\arctan2(0, b)) + (\text{gamma}(3/2, I*b/x^2) + \text{gamma}(3/2, -I*b/x^2)) * \cos(-3/4*\pi + 3/2*\arctan2(0, b)) - (I*\text{gamma}(3/2, I*b/x^2) - I*\text{gamma}(3/2, -I*b/x^2)) * \sin(3/4*\pi + 3/2*\arctan2(0, b)) - (-I*\text{gamma}(3/2, I*b/x^2) + I*\text{gamma}(3/2, -I*b/x^2)) * \sin(-3/4*\pi + 3/2*\arctan2(0, b))) * \cos(a) - ((I*\text{gamma}(3/2, I*b/x^2) - I*\text{gamma}(3/2, -I*b/x^2)) * \cos(3/4*\pi + 3/2*\arctan2(0, b)) + (I*\text{gamma}(3/2, I*b/x^2) - I*\text{gamma}(3/2, -I*b/x^2)) * \cos(-3/4*\pi + 3/2*\arctan2(0, b)) + (\text{gamma}(3/2, I*b/x^2) + \text{gamma}(3/2, -I*b/x^2)) * \sin(3/4*\pi + 3/2*\arctan2(0, b)) - (\text{gamma}(3/2, I*b/x^2) + \text{gamma}(3/2, -I*b/x^2)) * \sin(-3/4*\pi + 3/2*\arctan2(0, b))) * \sin(a)) / (x^3 * (\text{abs}(b)/x^2)^(3/2))$

**Fricas [A]** time = 1.66973, size = 235, normalized size = 2.42

$$\frac{\sqrt{2}\pi x \sqrt{\frac{b}{\pi}} \cos(a) S\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) + \sqrt{2}\pi x \sqrt{\frac{b}{\pi}} C\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) \sin(a) - 2b \sin\left(\frac{ax^2+b}{x^2}\right)}{4b^2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x^4,x, algorithm="fricas")

[Out]  $\frac{1}{4} * (\text{sqrt}(2) * \pi * x * \text{sqrt}(b/\pi) * \cos(a) * \text{fresnel\_sin}(\text{sqrt}(2) * \text{sqrt}(b/\pi)/x) + \text{sqrt}(2) * \pi * x * \text{sqrt}(b/\pi) * \text{fresnel\_cos}(\text{sqrt}(2) * \text{sqrt}(b/\pi)/x) * \sin(a) - 2 * b * \sin((a * x^2 + b)/x^2)) / (b^2 * x)$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x\*\*2)/x\*\*4,x)

[Out] Integral(cos(a + b/x\*\*2)/x\*\*4, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b/x^2)/x^4,x, algorithm="giac")

[Out] integrate(cos(a + b/x^2)/x^4, x)

$$3.45 \quad \int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx$$

**Optimal.** Leaf size=19

$$\sqrt{x} + \sin(\sqrt{x}) \cos(\sqrt{x})$$

[Out] Sqrt[x] + Cos[Sqrt[x]]\*Sin[Sqrt[x]]

**Rubi [A]** time = 0.0214578, antiderivative size = 19, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3380, 2635, 8}

$$\sqrt{x} + \sin(\sqrt{x}) \cos(\sqrt{x})$$

Antiderivative was successfully verified.

[In] Int[Cos[Sqrt[x]]^2/Sqrt[x], x]

[Out] Sqrt[x] + Cos[Sqrt[x]]\*Sin[Sqrt[x]]

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
  := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p,
    x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
  && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 2635

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]
  *(b*Ssin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Ssin[c + d*x])^(n - 2),
  x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

#### Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

#### Rubi steps

$$\begin{aligned} \int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx &= 2 \operatorname{Subst} \left( \int \cos^2(x) dx, x, \sqrt{x} \right) \\ &= \cos(\sqrt{x}) \sin(\sqrt{x}) + \operatorname{Subst} \left( \int 1 dx, x, \sqrt{x} \right) \\ &= \sqrt{x} + \cos(\sqrt{x}) \sin(\sqrt{x}) \end{aligned}$$

**Mathematica [A]** time = 0.0286949, size = 18, normalized size = 0.95

$$\sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[Sqrt[x]]^2/Sqrt[x], x]
```

```
[Out] Sqrt[x] + Sin[2*Sqrt[x]]/2
```

**Maple [A]** time = 0.032, size = 14, normalized size = 0.7

$$\cos(\sqrt{x})\sin(\sqrt{x}) + \sqrt{x}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(x^(1/2))^2/x^(1/2), x)
```

```
[Out] cos(x^(1/2))*sin(x^(1/2))+x^(1/2)
```

**Maxima [A]** time = 1.14036, size = 16, normalized size = 0.84

$$\sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(x^(1/2))^2/x^(1/2), x, algorithm="maxima")
```

```
[Out] sqrt(x) + 1/2*sin(2*sqrt(x))
```

**Fricas [A]** time = 1.64442, size = 50, normalized size = 2.63

$$\cos(\sqrt{x})\sin(\sqrt{x}) + \sqrt{x}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(x^(1/2))^2/x^(1/2), x, algorithm="fricas")
```

```
[Out] cos(sqrt(x))*sin(sqrt(x)) + sqrt(x)
```

**Sympy [B]** time = 0.356543, size = 39, normalized size = 2.05

$$\sqrt{x} \sin^2(\sqrt{x}) + \sqrt{x} \cos^2(\sqrt{x}) + \sin(\sqrt{x}) \cos(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(x**(1/2))**2/x**(1/2), x)
```

```
[Out] sqrt(x)*sin(sqrt(x))**2 + sqrt(x)*cos(sqrt(x))**2 + sin(sqrt(x))*cos(sqrt(x))
```

---

**Giac [A]** time = 1.11242, size = 16, normalized size = 0.84

$$\sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x^(1/2))^2/x^(1/2),x, algorithm="giac")

[Out] sqrt(x) + 1/2\*sin(2\*sqrt(x))

$$3.46 \quad \int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx$$

**Optimal.** Leaf size=8

$$2 \sin(\sqrt{x})$$

[Out] 2\*Sin[Sqrt[x]]

**Rubi [A]** time = 0.0088485, antiderivative size = 8, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3380, 2637}

$$2 \sin(\sqrt{x})$$

Antiderivative was successfully verified.

[In] Int[Cos[Sqrt[x]]/Sqrt[x], x]

[Out] 2\*Sin[Sqrt[x]]

#### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

#### Rule 2637

Int[sin[Pi/2 + (c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Simp[Sin[c + d\*x]/d, x] /; FreeQ[{c, d}, x]

#### Rubi steps

$$\begin{aligned} \int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx &= 2 \text{Subst} \left( \int \cos(x) dx, x, \sqrt{x} \right) \\ &= 2 \sin(\sqrt{x}) \end{aligned}$$

**Mathematica [A]** time = 0.0019517, size = 8, normalized size = 1.

$$2 \sin(\sqrt{x})$$

Antiderivative was successfully verified.

[In] Integrate[Cos[Sqrt[x]]/Sqrt[x], x]

[Out] 2\*Sin[Sqrt[x]]

**Maple [A]** time = 0.023, size = 7, normalized size = 0.9

$$2 \sin(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(x^(1/2))/x^(1/2),x)`

[Out] `2*sin(x^(1/2))`

---

**Maxima [A]** time = 1.14184, size = 8, normalized size = 1.

$$2 \sin(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x^(1/2))/x^(1/2),x, algorithm="maxima")`

[Out] `2*sin(sqrt(x))`

---

**Fricas [A]** time = 1.62154, size = 22, normalized size = 2.75

$$2 \sin(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x^(1/2))/x^(1/2),x, algorithm="fricas")`

[Out] `2*sin(sqrt(x))`

---

**Sympy [A]** time = 0.419689, size = 7, normalized size = 0.88

$$2 \sin(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x**(1/2))/x**(1/2),x)`

[Out] `2*sin(sqrt(x))`

---

**Giac [A]** time = 1.14308, size = 8, normalized size = 1.

$$2 \sin(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x^(1/2))/x^(1/2),x, algorithm="giac")`

[Out] `2*sin(sqrt(x))`



### 3.47 $\int \cos(\sqrt{x}) dx$

**Optimal.** Leaf size=22

$$2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

[Out] 2\*Cos[Sqrt[x]] + 2\*Sqrt[x]\*Sin[Sqrt[x]]

**Rubi [A]** time = 0.0110084, antiderivative size = 22, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.5$ , Rules used = {3362, 3296, 2638}

$$2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

Antiderivative was successfully verified.

[In] Int[Cos[Sqrt[x]], x]

[Out] 2\*Cos[Sqrt[x]] + 2\*Sqrt[x]\*Sin[Sqrt[x]]

#### Rule 3362

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)]\*(b\_.))^(p\_.), x\_Symbol] :> Dist[1/(n\*f), Subst[Int[x^(1/n - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, (e + f\*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]

#### Rule 3296

Int[((c\_.) + (d\_.)\*(x\_))^(m\_.)\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] :> -Simp[(c + d\*x)^m\*Cos[e + f\*x]/f, x] + Dist[(d\*m)/f, Int[(c + d\*x)^(m - 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

#### Rule 2638

Int[sin[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> -Simp[Cos[c + d\*x]/d, x] /; FreeQ[{c, d}, x]

#### Rubi steps

$$\begin{aligned} \int \cos(\sqrt{x}) dx &= 2 \text{Subst} \left( \int x \cos(x) dx, x, \sqrt{x} \right) \\ &= 2\sqrt{x} \sin(\sqrt{x}) - 2 \text{Subst} \left( \int \sin(x) dx, x, \sqrt{x} \right) \\ &= 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x}) \end{aligned}$$

**Mathematica [A]** time = 0.0146263, size = 22, normalized size = 1.

$$2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

Antiderivative was successfully verified.

[In] Integrate[Cos[Sqrt[x]], x]

[Out]  $2*\text{Cos}[\text{Sqrt}[x]] + 2*\text{Sqrt}[x]*\text{Sin}[\text{Sqrt}[x]]$

**Maple [A]** time = 0.027, size = 17, normalized size = 0.8

$$2 \cos(\sqrt{x}) + 2 \sin(\sqrt{x}) \sqrt{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(x^(1/2)),x)`

[Out]  $2*\cos(x^{(1/2)})+2*\sin(x^{(1/2)})*x^{(1/2)}$

**Maxima [A]** time = 1.16089, size = 22, normalized size = 1.

$$2 \sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x^(1/2)),x, algorithm="maxima")`

[Out]  $2*\text{sqrt}(x)*\text{sin}(\text{sqrt}(x)) + 2*\text{cos}(\text{sqrt}(x))$

**Fricas [A]** time = 1.63744, size = 55, normalized size = 2.5

$$2 \sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x^(1/2)),x, algorithm="fricas")`

[Out]  $2*\text{sqrt}(x)*\text{sin}(\text{sqrt}(x)) + 2*\text{cos}(\text{sqrt}(x))$

**Sympy [A]** time = 0.409983, size = 20, normalized size = 0.91

$$2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x**(1/2)),x)`

[Out]  $2*\text{sqrt}(x)*\text{sin}(\text{sqrt}(x)) + 2*\text{cos}(\text{sqrt}(x))$

**Giac [A]** time = 1.11337, size = 22, normalized size = 1.

$$2 \sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(x^(1/2)),x, algorithm="giac")
```

```
[Out] 2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))
```

### 3.48 $\int \cos^2(\sqrt{x}) dx$

**Optimal.** Leaf size=36

$$\frac{x}{2} + \frac{1}{2} \cos^2(\sqrt{x}) + \sqrt{x} \sin(\sqrt{x}) \cos(\sqrt{x})$$

[Out] x/2 + Cos[Sqrt[x]]^2/2 + Sqrt[x]\*Cos[Sqrt[x]]\*Sin[Sqrt[x]]

**Rubi [A]** time = 0.0213152, antiderivative size = 36, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3362, 3310, 30}

$$\frac{x}{2} + \frac{1}{2} \cos^2(\sqrt{x}) + \sqrt{x} \sin(\sqrt{x}) \cos(\sqrt{x})$$

Antiderivative was successfully verified.

[In] Int[Cos[Sqrt[x]]^2,x]

[Out] x/2 + Cos[Sqrt[x]]^2/2 + Sqrt[x]\*Cos[Sqrt[x]]\*Sin[Sqrt[x]]

#### Rule 3362

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)]\*(b\_.))^(p\_.), x\_Symbol] :> Dist[1/(n\*f), Subst[Int[x^(1/n - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, (e + f\*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]

#### Rule 3310

Int[((c\_.) + (d\_.)\*(x\_))\*((b\_.)\*sin[(e\_.) + (f\_.)\*(x\_)])^(n\_), x\_Symbol] :> Simp[(d\*(b\*Ssin[e + f\*x])^n)/(f^2\*n^2), x] + (Dist[(b^2\*(n - 1))/n, Int[(c + d\*x)\*(b\*Ssin[e + f\*x])^(n - 2), x], x] - Simp[(b\*(c + d\*x)\*Cos[e + f\*x]\*(b\*Ssin[e + f\*x])^(n - 1))/(f\*n), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]

#### Rule 30

Int[(x\_)^(m\_.), x\_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

#### Rubi steps

$$\begin{aligned} \int \cos^2(\sqrt{x}) dx &= 2 \text{Subst} \left( \int x \cos^2(x) dx, x, \sqrt{x} \right) \\ &= \frac{1}{2} \cos^2(\sqrt{x}) + \sqrt{x} \cos(\sqrt{x}) \sin(\sqrt{x}) + \text{Subst} \left( \int x dx, x, \sqrt{x} \right) \\ &= \frac{x}{2} + \frac{1}{2} \cos^2(\sqrt{x}) + \sqrt{x} \cos(\sqrt{x}) \sin(\sqrt{x}) \end{aligned}$$

**Mathematica [A]** time = 0.0278696, size = 31, normalized size = 0.86

$$\frac{1}{4} (2(x + \sqrt{x} \sin(2\sqrt{x})) + \cos(2\sqrt{x}))$$

Antiderivative was successfully verified.

[In] Integrate[Cos[Sqrt[x]]^2,x]

[Out] (Cos[2\*Sqrt[x]] + 2\*(x + Sqrt[x]\*Sin[2\*Sqrt[x]]))/4

**Maple [A]** time = 0.026, size = 34, normalized size = 0.9

$$2\sqrt{x}\left(\frac{1}{2}\cos(\sqrt{x})\sin(\sqrt{x}) + \frac{1}{2}\sqrt{x}\right) - \frac{x}{2} - \frac{1}{2}\left(\sin(\sqrt{x})\right)^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x^(1/2))^2,x)

[Out] 2\*x^(1/2)\*(1/2\*cos(x^(1/2))\*sin(x^(1/2))+1/2\*x^(1/2))-1/2\*x-1/2\*sin(x^(1/2))^2

**Maxima [A]** time = 1.07688, size = 31, normalized size = 0.86

$$\frac{1}{2}\sqrt{x}\sin(2\sqrt{x}) + \frac{1}{2}x + \frac{1}{4}\cos(2\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x^(1/2))^2,x, algorithm="maxima")

[Out] 1/2\*sqrt(x)\*sin(2\*sqrt(x)) + 1/2\*x + 1/4\*cos(2\*sqrt(x))

**Fricas [A]** time = 1.60506, size = 86, normalized size = 2.39

$$\sqrt{x}\cos(\sqrt{x})\sin(\sqrt{x}) + \frac{1}{2}\cos(\sqrt{x})^2 + \frac{1}{2}x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x^(1/2))^2,x, algorithm="fricas")

[Out] sqrt(x)\*cos(sqrt(x))\*sin(sqrt(x)) + 1/2\*cos(sqrt(x))^2 + 1/2\*x

**Sympy [A]** time = 0.386695, size = 51, normalized size = 1.42

$$\sqrt{x}\sin(\sqrt{x})\cos(\sqrt{x}) + \frac{x\sin^2(\sqrt{x})}{2} + \frac{x\cos^2(\sqrt{x})}{2} + \frac{\cos^2(\sqrt{x})}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x\*\*(1/2))\*\*2,x)

```
[Out] sqrt(x)*sin(sqrt(x))*cos(sqrt(x)) + x*sin(sqrt(x))**2/2 + x*cos(sqrt(x))**2/2 + cos(sqrt(x))**2/2
```

---

**Giac [A]** time = 1.11741, size = 31, normalized size = 0.86

$$\frac{1}{2} \sqrt{x} \sin(2\sqrt{x}) + \frac{1}{2} x + \frac{1}{4} \cos(2\sqrt{x})$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(x^(1/2)))^2,x, algorithm="giac"
```

```
[Out] 1/2*sqrt(x)*sin(2*sqrt(x)) + 1/2*x + 1/4*cos(2*sqrt(x))
```

### 3.49 $\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx$

**Optimal.** Leaf size=235

$$\frac{405405\sqrt{\frac{\pi}{2}}\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right)}{64b^{15/2}} + \frac{405405\sqrt{\frac{\pi}{2}}\cos(a)S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{64b^{15/2}} - \frac{429x^{3/2}\sin(a + b\sqrt[3]{x})}{4b^3} + \frac{27027x^{5/6}\sin(a + b\sqrt[3]{x})}{16b}$$

```
[Out] (135135*Sqrt[x]*Cos[a + b*x^(1/3)])/(32*b^6) - (3861*x^(7/6)*Cos[a + b*x^(1/3)])/(8*b^4) + (39*x^(11/6)*Cos[a + b*x^(1/3)])/(2*b^2) + (405405*Sqrt[Pi/2]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)])/(64*b^(15/2)) + (405405*Sqrt[Pi/2]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a])/(64*b^(15/2)) - (405405*x^(1/6)*Sin[a + b*x^(1/3)])/(64*b^7) + (27027*x^(5/6)*Sin[a + b*x^(1/3)])/(16*b^5) - (429*x^(3/2)*Sin[a + b*x^(1/3)])/(4*b^3) + (3*x^(13/6)*Sin[a + b*x^(1/3)])/b
```

**Rubi [A]** time = 0.348002, antiderivative size = 235, normalized size of antiderivative = 1., number of steps used = 13, number of rules used = 7, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$ , Rules used = {3416, 3296, 3306, 3305, 3351, 3304, 3352}

$$\frac{405405\sqrt{\frac{\pi}{2}}\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right)}{64b^{15/2}} + \frac{405405\sqrt{\frac{\pi}{2}}\cos(a)S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{64b^{15/2}} - \frac{429x^{3/2}\sin(a + b\sqrt[3]{x})}{4b^3} + \frac{27027x^{5/6}\sin(a + b\sqrt[3]{x})}{16b}$$

Antiderivative was successfully verified.

```
[In] Int[x^(3/2)*Cos[a + b*x^(1/3)], x]
```

```
[Out] (135135*Sqrt[x]*Cos[a + b*x^(1/3)])/(32*b^6) - (3861*x^(7/6)*Cos[a + b*x^(1/3)])/(8*b^4) + (39*x^(11/6)*Cos[a + b*x^(1/3)])/(2*b^2) + (405405*Sqrt[Pi/2]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)])/(64*b^(15/2)) + (405405*Sqrt[Pi/2]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a])/(64*b^(15/2)) - (405405*x^(1/6)*Sin[a + b*x^(1/3)])/(64*b^7) + (27027*x^(5/6)*Sin[a + b*x^(1/3)])/(16*b^5) - (429*x^(3/2)*Sin[a + b*x^(1/3)])/(4*b^3) + (3*x^(13/6)*Sin[a + b*x^(1/3)])/b
```

#### Rule 3416

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
:> Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*x^(k*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x]
&& IntegerQ[p] && FractionQ[n]
```

#### Rule 3296

```
Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol]
:> -Simp[((c + d*x)^m*Cos[e + f*x])/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

#### Rule 3306

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol]
:> Dist[Cos[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]
```

#### Rule 3305

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d
, Subst[Int[Sin[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}
, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*Fresne
lS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rule 3304

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := D
ist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d
, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*Fresne
lC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rubi steps

$$\begin{aligned}
\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx &= 3 \operatorname{Subst}\left(\int x^{13/2} \cos(a + bx) dx, x, \sqrt[3]{x}\right) \\
&= \frac{3x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{39 \operatorname{Subst}\left(\int x^{11/2} \sin(a + bx) dx, x, \sqrt[3]{x}\right)}{2b} \\
&= \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} + \frac{3x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{429 \operatorname{Subst}\left(\int x^{9/2} \cos(a + bx) dx, x, \sqrt[3]{x}\right)}{4b^2} \\
&= \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{429x^{3/2} \sin(a + b\sqrt[3]{x})}{4b^3} + \frac{3x^{13/6} \sin(a + b\sqrt[3]{x})}{b} + \frac{3861 \operatorname{Subst}\left(\int x^{7/2} \cos(a + bx) dx, x, \sqrt[3]{x}\right)}{8b^2} \\
&= -\frac{3861x^{7/6} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{429x^{3/2} \sin(a + b\sqrt[3]{x})}{4b^3} + \frac{3x^{13/6} \sin(a + b\sqrt[3]{x})}{b} \\
&= -\frac{3861x^{7/6} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} + \frac{27027x^{5/6} \sin(a + b\sqrt[3]{x})}{16b^5} - \frac{429x^{3/2} \sin(a + b\sqrt[3]{x})}{4b^3} \\
&= \frac{135135\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^6} - \frac{3861x^{7/6} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} + \frac{27027x^{5/6} \sin(a + b\sqrt[3]{x})}{16b^5} \\
&= \frac{135135\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^6} - \frac{3861x^{7/6} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{405405\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{64b^5} \\
&= \frac{135135\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^6} - \frac{3861x^{7/6} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{405405\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{64b^5} \\
&= \frac{135135\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^6} - \frac{3861x^{7/6} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{405405\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{64b^5} \\
&= \frac{135135\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^6} - \frac{3861x^{7/6} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} + \frac{405405\sqrt{\frac{\pi}{2}} \cos(a + b\sqrt[3]{x})}{64b^5}
\end{aligned}$$

**Mathematica [A]** time = 0.568362, size = 165, normalized size = 0.7

$$6\sqrt{b}\sqrt[6]{x} \left( (64b^6x^2 - 2288b^4x^{4/3} + 36036b^2x^{2/3} - 135135) \sin(a + b\sqrt[3]{x}) + 26(16b^5x^{5/3} - 396b^3x + 3465b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x}) \right)$$

128b<sup>15/2</sup>

Antiderivative was successfully verified.



[In] Integrate[x^(3/2)\*Cos[a + b\*x^(1/3)],x]

[Out] (405405\*Sqrt[2\*Pi]\*Cos[a]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)] + 405405\*Sqrt[2\*Pi]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)]\*Sin[a] + 6\*Sqrt[b]\*x^(1/6)\*(26\*(3465\*b\*x^(1/3) - 396\*b^3\*x + 16\*b^5\*x^(5/3))\*Cos[a + b\*x^(1/3)] + (-135135 + 36036\*b^2\*x^(2/3) - 2288\*b^4\*x^(4/3) + 64\*b^6\*x^2)\*Sin[a + b\*x^(1/3)])) / (128\*b^(15/2))

**Maple [A]** time = 0.033, size = 196, normalized size = 0.8

$$3 \frac{\sin(a + b\sqrt[3]{x})}{b} x^{\frac{13}{6}} - 39 \frac{1}{b} \left( -1/2 \frac{\cos(a + b\sqrt[3]{x})}{b} x^{\frac{11}{6}} + 11/2 \frac{1}{b} \left( 1/2 \frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} - 9/2 \frac{1}{b} \left( -1/2 \frac{x^{7/6} \cos(a + b\sqrt[3]{x})}{b} \right. \right. \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(3/2)\*cos(a+b\*x^(1/3)),x)

[Out] 3\*x^(13/6)\*sin(a+b\*x^(1/3))/b-39/b\*(-1/2/b\*x^(11/6)\*cos(a+b\*x^(1/3))+11/2/b\*(1/2/b\*x^(3/2)\*sin(a+b\*x^(1/3))-9/2/b\*(-1/2/b\*x^(7/6)\*cos(a+b\*x^(1/3))+7/2/b\*(1/2/b\*x^(5/6)\*sin(a+b\*x^(1/3))-5/2/b\*(-1/2/b\*x^(1/2)\*cos(a+b\*x^(1/3))+3/2/b\*(1/2\*x^(1/6)\*sin(a+b\*x^(1/3)))/b-1/4/b^(3/2)\*2^(1/2)\*Pi^(1/2)\*(cos(a)\*FresnelS(x^(1/6)\*b^(1/2)\*2^(1/2)/Pi^(1/2))+sin(a)\*FresnelC(x^(1/6)\*b^(1/2)\*2^(1/2)/Pi^(1/2))))))

**Maxima [C]** time = 2.76905, size = 459, normalized size = 1.95

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(a+b\*x^(1/3)),x, algorithm="maxima")

[Out] -3/512\*(sqrt(pi)\*((( -135135\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - 135135\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 135135\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 135135\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*cos(a) - (135135\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 135135\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 135135\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 135135\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*sin(a))\*erf(sqrt(I\*b)\*x^(1/6)) + (((135135\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 135135\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 135135\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 135135\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*cos(a) - (135135\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 135135\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + 135135\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - 135135\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*sin(a))\*erf(sqrt(-I\*b)\*x^(1/6)))\*sqrt(abs(b)) - 208\*(16\*b^5\*x^(11/6)\*abs(b) - 396\*b^3\*x^(7/6)\*abs(b) + 3465\*b\*sqrt(x)\*abs(b))\*cos(b\*x^(1/3) + a) - 8\*(64\*b^6\*x^(13/6)\*abs(b) - 2288\*b^4\*x^(3/2)\*abs(b) + 36036\*b^2\*x^(5/6)\*abs(b) - 135135\*x^(1/6)\*abs(b))\*sin(b\*x^(1/3) + a)/(b^7\*abs(b))

**Fricas [A]** time = 1.78436, size = 462, normalized size = 1.97

$$3 \left( 135135 \sqrt{2\pi} \sqrt{\frac{b}{\pi}} \cos(a) S \left( \sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) + 135135 \sqrt{2\pi} \sqrt{\frac{b}{\pi}} C \left( \sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) \sin(a) + 52 \left( 16 b^6 x^{\frac{11}{6}} - 396 b^4 x^{\frac{7}{6}} + 3465 b^2 \right. \right.$$

128 b^8

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(a+b\*x^(1/3)),x, algorithm="fricas")

[Out] 3/128\*(135135\*sqrt(2)\*pi\*sqrt(b/pi)\*cos(a)\*fresnel\_sin(sqrt(2)\*x^(1/6)\*sqrt(b/pi)) + 135135\*sqrt(2)\*pi\*sqrt(b/pi)\*fresnel\_cos(sqrt(2)\*x^(1/6)\*sqrt(b/pi))\*sin(a) + 52\*(16\*b^6\*x^(11/6) - 396\*b^4\*x^(7/6) + 3465\*b^2\*sqrt(x))\*cos(b\*x^(1/3) + a) - 2\*(2288\*b^5\*x^(3/2) - 36036\*b^3\*x^(5/6) - (64\*b^7\*x^2 - 135135\*b)\*x^(1/6))\*sin(b\*x^(1/3) + a)/b^8

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{\frac{3}{2}} \cos(a + b\sqrt[3]{x}) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*(3/2)\*cos(a+b\*x\*\*(1/3)),x)

[Out] Integral(x\*\*(3/2)\*cos(a + b\*x\*\*(1/3)), x)

**Giac [C]** time = 1.15408, size = 325, normalized size = 1.38

$$\frac{3 \left( 64i b^6 x^{\frac{13}{6}} - 416 b^5 x^{\frac{11}{6}} - 2288i b^4 x^{\frac{9}{6}} + 10296 b^3 x^{\frac{7}{6}} + 36036i b^2 x^{\frac{5}{6}} - 90090 b \sqrt{x} - 135135i x^{\frac{1}{6}} \right) e^{\left( i b x^{\frac{1}{3}} + i a \right)} - 3 \left( -64i b^6 x^{\frac{13}{6}} + 416 b^5 x^{\frac{11}{6}} + 2288i b^4 x^{\frac{9}{6}} - 10296 b^3 x^{\frac{7}{6}} - 36036i b^2 x^{\frac{5}{6}} + 90090 b \sqrt{x} + 135135i x^{\frac{1}{6}} \right) e^{\left( -i b x^{\frac{1}{3}} - i a \right)}}{128 b^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(a+b\*x^(1/3)),x, algorithm="giac")

[Out] -3/128\*(64\*I\*b^6\*x^(13/6) - 416\*b^5\*x^(11/6) - 2288\*I\*b^4\*x^(9/6) + 10296\*b^3\*x^(7/6) + 36036\*I\*b^2\*x^(5/6) - 90090\*b\*sqrt(x) - 135135\*I\*x^(1/6))\*e^(I\*b\*x^(1/3) + I\*a)/b^7 - 3/128\*(-64\*I\*b^6\*x^(13/6) - 416\*b^5\*x^(11/6) + 2288\*I\*b^4\*x^(9/6) + 10296\*b^3\*x^(7/6) - 36036\*I\*b^2\*x^(5/6) - 90090\*b\*sqrt(x) + 135135\*I\*x^(1/6))\*e^(-I\*b\*x^(1/3) - I\*a)/b^7 + 405405/256\*I\*sqrt(2)\*sqrt(pi)\*erf(-1/2\*sqrt(2)\*x^(1/6)\*(-I\*b/abs(b) + 1)\*sqrt(abs(b)))\*e^(I\*a)/(b^7\*(-I\*b/abs(b) + 1)\*sqrt(abs(b))) - 405405/256\*I\*sqrt(2)\*sqrt(pi)\*erf(-1/2\*sqrt(2)\*x^(1/6)\*(I\*b/abs(b) + 1)\*sqrt(abs(b)))\*e^(-I\*a)/(b^7\*(I\*b/abs(b) + 1)\*sqrt(abs(b)))

### 3.50 $\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$

**Optimal.** Leaf size=169

$$\frac{315\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{b} \sqrt[6]{x}\right)}{8b^{9/2}} - \frac{315\sqrt{\frac{\pi}{2}} \sin(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right)}{8b^{9/2}} + \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{105\sqrt{x} \sin(a + b\sqrt[3]{x})}{4b^3}$$

[Out]  $(-315x^{(1/6)} \text{Cos}[a + b x^{(1/3)}]) / (8b^4) + (21x^{(5/6)} \text{Cos}[a + b x^{(1/3)}]) / (2b^2) + (315 \text{Sqrt}[\text{Pi}/2] \text{Cos}[a] \text{FresnelC}[\text{Sqrt}[b] \text{Sqrt}[2/\text{Pi}] x^{(1/6)}]) / (8b^{(9/2)}) - (315 \text{Sqrt}[\text{Pi}/2] \text{FresnelS}[\text{Sqrt}[b] \text{Sqrt}[2/\text{Pi}] x^{(1/6)}] \text{Sin}[a]) / (8b^{(9/2)}) - (105 \text{Sqrt}[x] \text{Sin}[a + b x^{(1/3)}]) / (4b^3) + (3x^{(7/6)} \text{Sin}[a + b x^{(1/3)}]) / b$

**Rubi [A]** time = 0.196254, antiderivative size = 169, normalized size of antiderivative = 1., number of steps used = 10, number of rules used = 7, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$ , Rules used = {3416, 3296, 3306, 3305, 3351, 3304, 3352}

$$\frac{315\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{b} \sqrt[6]{x}\right)}{8b^{9/2}} - \frac{315\sqrt{\frac{\pi}{2}} \sin(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right)}{8b^{9/2}} + \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{105\sqrt{x} \sin(a + b\sqrt[3]{x})}{4b^3}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[\text{Sqrt}[x] \text{Cos}[a + b x^{(1/3)}], x]$

[Out]  $(-315x^{(1/6)} \text{Cos}[a + b x^{(1/3)}]) / (8b^4) + (21x^{(5/6)} \text{Cos}[a + b x^{(1/3)}]) / (2b^2) + (315 \text{Sqrt}[\text{Pi}/2] \text{Cos}[a] \text{FresnelC}[\text{Sqrt}[b] \text{Sqrt}[2/\text{Pi}] x^{(1/6)}]) / (8b^{(9/2)}) - (315 \text{Sqrt}[\text{Pi}/2] \text{FresnelS}[\text{Sqrt}[b] \text{Sqrt}[2/\text{Pi}] x^{(1/6)}] \text{Sin}[a]) / (8b^{(9/2)}) - (105 \text{Sqrt}[x] \text{Sin}[a + b x^{(1/3)}]) / (4b^3) + (3x^{(7/6)} \text{Sin}[a + b x^{(1/3)}]) / b$

#### Rule 3416

$\text{Int}[(a_.) + \text{Cos}[(c_.) + (d_.)(x_)^{(n_)}] (b_.)^{(p_.)} (x_)^{(m_.)}, x\_Symbol] \rightarrow \text{Module}[\{k = \text{Denominator}[n]\}, \text{Dist}[k, \text{Subst}[\text{Int}[x^{(k(m+1)-1)} (a + b \text{Cos}[c + d x^{(k n)}])^p, x], x, x^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \&\& \text{IntegerQ}[p] \&\& \text{FractionQ}[n]$

#### Rule 3296

$\text{Int}[(c_.) + (d_.)(x_)^{(m_.)} \sin[(e_.) + (f_.)(x_)], x\_Symbol] \rightarrow -\text{Simp}[(c + d x)^m \text{Cos}[e + f x] / f, x] + \text{Dist}[(d m) / f, \text{Int}[(c + d x)^{(m-1)} \text{Cos}[e + f x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&\& \text{GtQ}[m, 0]$

#### Rule 3306

$\text{Int}[\sin[(e_.) + (f_.)(x_)] / \text{Sqrt}[(c_.) + (d_.)(x_)], x\_Symbol] \rightarrow \text{Dist}[\text{Cos}[(d e - c f) / d], \text{Int}[\text{Sin}[(c f) / d + f x] / \text{Sqrt}[c + d x], x], x] + \text{Dist}[\text{Sin}[(d e - c f) / d], \text{Int}[\text{Cos}[(c f) / d + f x] / \text{Sqrt}[c + d x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&\& \text{ComplexFreeQ}[f] \&\& \text{NeQ}[d e - c f, 0]$

#### Rule 3305

$\text{Int}[\sin[(e_.) + (f_.)(x_)] / \text{Sqrt}[(c_.) + (d_.)(x_)], x\_Symbol] \rightarrow \text{Dist}[2/d, \text{Subst}[\text{Int}[\text{Sin}[(f x^2) / d], x], x, \text{Sqrt}[c + d x]], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&\& \text{ComplexFreeQ}[f] \&\& \text{EqQ}[d e - c f, 0]$

**Rule 3351**

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

**Rule 3304**

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

**Rule 3352**

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

**Rubi steps**

$$\begin{aligned}
\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx &= 3 \operatorname{Subst}\left(\int x^{7/2} \cos(a + bx) dx, x, \sqrt[3]{x}\right) \\
&= \frac{3x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{21 \operatorname{Subst}\left(\int x^{5/2} \sin(a + bx) dx, x, \sqrt[3]{x}\right)}{2b} \\
&= \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2} + \frac{3x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{105 \operatorname{Subst}\left(\int x^{3/2} \cos(a + bx) dx, x, \sqrt[3]{x}\right)}{4b^2} \\
&= \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{105\sqrt{x} \sin(a + b\sqrt[3]{x})}{4b^3} + \frac{3x^{7/6} \sin(a + b\sqrt[3]{x})}{b} + \frac{315 \operatorname{Subst}\left(\int \sqrt{x} \sin(a + bx) dx, x, \sqrt[3]{x}\right)}{8b^3} \\
&= -\frac{315\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{105\sqrt{x} \sin(a + b\sqrt[3]{x})}{4b^3} + \frac{3x^{7/6} \sin(a + b\sqrt[3]{x})}{b} \\
&= -\frac{315\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{105\sqrt{x} \sin(a + b\sqrt[3]{x})}{4b^3} + \frac{3x^{7/6} \sin(a + b\sqrt[3]{x})}{b} \\
&= -\frac{315\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2} - \frac{105\sqrt{x} \sin(a + b\sqrt[3]{x})}{4b^3} + \frac{3x^{7/6} \sin(a + b\sqrt[3]{x})}{b} \\
&= -\frac{315\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2} + \frac{315\sqrt{\frac{\pi}{2}} \cos(a) C\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{8b^{9/2}} - \frac{315\sqrt{\frac{\pi}{2}} S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{8b^{9/2}}
\end{aligned}$$

**Mathematica [A]** time = 0.362793, size = 141, normalized size = 0.83

$$\frac{6\sqrt{b}\sqrt[6]{x}(2b\sqrt[3]{x}(4b^2x^{2/3} - 35)\sin(a + b\sqrt[3]{x}) + 7(4b^2x^{2/3} - 15)\cos(a + b\sqrt[3]{x})) + 315\sqrt{2\pi}\cos(a)\operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right) - 315\sqrt{2\pi}\sin(a)\operatorname{FresnelS}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right)}{16b^{9/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[x]*Cos[a + b*x^(1/3)], x]
```

```
[Out] (315*Sqrt[2*Pi]*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] - 315*Sqrt[2*Pi]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] + 6*Sqrt[b]*x^(1/6)*(7*(-15 + 4*b^2*x^(2/3))*Cos[a + b*x^(1/3)] + 2*b*(-35 + 4*b^2*x^(2/3))*x^(1/3)*Sin[a + b*x^(1/3)]))/(16*b^(9/2))
```

**Maple [A]** time = 0.03, size = 131, normalized size = 0.8

$$3 \frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - 21 \frac{1}{b} \left( -1/2 \frac{x^{5/6} \cos(a + b\sqrt[3]{x})}{b} + 5/2 \frac{1}{b} \left( 1/2 \frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - 3/2 \frac{1}{b} \left( -1/2 \frac{\cos(a + b\sqrt[3]{x}) \sqrt[6]{x}}{b} + 1/2 \frac{\sin(a + b\sqrt[3]{x}) \sqrt[6]{x}}{b} \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^(1/2)*cos(a+b*x^(1/3)),x)`

[Out]  $3*x^{7/6}*sin(a+b*x^{1/3})/b-21/b*(-1/2/b*x^{5/6}*cos(a+b*x^{1/3}))+5/2/b*(1/2/b*x^{1/2}*sin(a+b*x^{1/3}))-3/2/b*(-1/2/b*x^{1/6}*cos(a+b*x^{1/3}))+1/4/b^{3/2}*2^{1/2}*Pi^{1/2}*(cos(a)*FresnelC(x^{1/6}*b^{1/2}*2^{1/2}/Pi^{1/2}))-sin(a)*FresnelS(x^{1/6}*b^{1/2}*2^{1/2}/Pi^{1/2}))$

**Maxima [C]** time = 2.43454, size = 419, normalized size = 2.48

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^(1/2)*cos(a+b*x^(1/3)),x, algorithm="maxima")`

[Out]  $3/64*(sqrt(pi)*(((105*cos(1/4*pi + 1/2*arctan2(0, b)) + 105*cos(-1/4*pi + 1/2*arctan2(0, b)) - 105*I*sin(1/4*pi + 1/2*arctan2(0, b)) + 105*I*sin(-1/4*pi + 1/2*arctan2(0, b)))*cos(a) - (105*I*cos(1/4*pi + 1/2*arctan2(0, b)) + 105*I*cos(-1/4*pi + 1/2*arctan2(0, b)) + 105*sin(1/4*pi + 1/2*arctan2(0, b)) - 105*sin(-1/4*pi + 1/2*arctan2(0, b)))*sin(a))*erf(sqrt(I*b)*x^{1/6}) + ((105*cos(1/4*pi + 1/2*arctan2(0, b)) + 105*cos(-1/4*pi + 1/2*arctan2(0, b)) + 105*I*sin(1/4*pi + 1/2*arctan2(0, b)) - 105*I*sin(-1/4*pi + 1/2*arctan2(0, b)))*cos(a) - (-105*I*cos(1/4*pi + 1/2*arctan2(0, b)) - 105*I*cos(-1/4*pi + 1/2*arctan2(0, b)) + 105*sin(1/4*pi + 1/2*arctan2(0, b)) - 105*sin(-1/4*pi + 1/2*arctan2(0, b)))*sin(a))*erf(sqrt(-I*b)*x^{1/6}))*sqrt(abs(b)) + 56*(4*b^2*x^{5/6}*abs(b) - 15*x^{1/6}*abs(b))*cos(b*x^{1/3} + a) + 16*(4*b^3*x^{7/6}*abs(b) - 35*b*sqrt(x)*abs(b))*sin(b*x^{1/3} + a))/(b^4*abs(b))$

**Fricas [A]** time = 1.7632, size = 366, normalized size = 2.17

$$\frac{3 \left( 105 \sqrt{2} \pi \sqrt{\frac{b}{\pi}} \cos(a) C \left( \sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) - 105 \sqrt{2} \pi \sqrt{\frac{b}{\pi}} S \left( \sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) \sin(a) + 14 \left( 4 b^3 x^{\frac{5}{6}} - 15 b x^{\frac{1}{6}} \right) \cos \left( b x^{\frac{1}{3}} + a \right) + 4 \left( 4 b^3 x^{\frac{5}{6}} - 15 b x^{\frac{1}{6}} \right) \sin \left( b x^{\frac{1}{3}} + a \right) \right)}{16 b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^(1/2)*cos(a+b*x^(1/3)),x, algorithm="fricas")`

[Out]  $3/16*(105*sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel\_cos(sqrt(2)*x^{1/6}*sqrt(b/pi)) - 105*sqrt(2)*pi*sqrt(b/pi)*fresnel\_sin(sqrt(2)*x^{1/6}*sqrt(b/pi))*sin(a) + 14*(4*b^3*x^{5/6} - 15*b*x^{1/6})*cos(b*x^{1/3} + a) + 4*(4*b^3*x^{5/6} - 15*b*x^{1/6})*sin(b*x^{1/3} + a))/b^5$

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**(1/2)*cos(a+b*x**(1/3)),x)`

[Out] Timed out

---

**Giac [C]** time = 1.15468, size = 261, normalized size = 1.54

$$\frac{3 \left( 8i b^3 x^{\frac{7}{6}} - 28 b^2 x^{\frac{5}{6}} - 70i b \sqrt{x} + 105 x^{\frac{1}{6}} \right) e^{\left( i b x^{\frac{1}{3}} + i a \right)}}{16 b^4} - \frac{3 \left( -8i b^3 x^{\frac{7}{6}} - 28 b^2 x^{\frac{5}{6}} + 70i b \sqrt{x} + 105 x^{\frac{1}{6}} \right) e^{\left( -i b x^{\frac{1}{3}} - i a \right)}}{16 b^4} - \frac{315 \sqrt{2} \sqrt{\pi}}{16 b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(1/2)\*cos(a+b\*x^(1/3)),x, algorithm="giac")

[Out] 
$$-3/16*(8*I*b^3*x^{(7/6)} - 28*b^2*x^{(5/6)} - 70*I*b*\sqrt{x} + 105*x^{(1/6)})*e^{(I*b*x^{(1/3)} + I*a)/b^4} - 3/16*(-8*I*b^3*x^{(7/6)} - 28*b^2*x^{(5/6)} + 70*I*b*\sqrt{x} + 105*x^{(1/6)})*e^{(-I*b*x^{(1/3)} - I*a)/b^4} - 315/32*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-1/2*\sqrt{2}*x^{(1/6)}*(-I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})}*e^{(I*a)/(b^4*(-I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})} - 315/32*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-1/2*\sqrt{2}*x^{(1/6)}*(I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})}*e^{(-I*a)/(b^4*(I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})}$$

$$3.51 \quad \int \frac{\cos(a+b\sqrt[3]{x})}{\sqrt{x}} dx$$

**Optimal.** Leaf size=99

$$-\frac{3\sqrt{\frac{\pi}{2}} \sin(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{b} \sqrt[6]{x}\right)}{b^{3/2}} - \frac{3\sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right)}{b^{3/2}} + \frac{3\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b}$$

[Out]  $(-3\sqrt{\text{Pi}/2} \text{Cos}[a] \text{FresnelS}[\text{Sqrt}[b] \text{Sqrt}[2/\text{Pi}] x^{1/6}])/b^{3/2} - (3\sqrt{\text{Pi}/2} \text{FresnelC}[\text{Sqrt}[b] \text{Sqrt}[2/\text{Pi}] x^{1/6}] \text{Sin}[a])/b^{3/2} + (3x^{1/6} \text{Sin}[a + b x^{1/3}])/b$

**Rubi [A]** time = 0.113156, antiderivative size = 99, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 7, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$ , Rules used = {3416, 3296, 3306, 3305, 3351, 3304, 3352}

$$-\frac{3\sqrt{\frac{\pi}{2}} \sin(a) \text{FresnelC}\left(\sqrt{\frac{2}{\pi}} \sqrt{b} \sqrt[6]{x}\right)}{b^{3/2}} - \frac{3\sqrt{\frac{\pi}{2}} \cos(a) S\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right)}{b^{3/2}} + \frac{3\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b}$$

Antiderivative was successfully verified.

[In]  $\text{Int}[\text{Cos}[a + b x^{1/3}]/\text{Sqrt}[x], x]$

[Out]  $(-3\sqrt{\text{Pi}/2} \text{Cos}[a] \text{FresnelS}[\text{Sqrt}[b] \text{Sqrt}[2/\text{Pi}] x^{1/6}])/b^{3/2} - (3\sqrt{\text{Pi}/2} \text{FresnelC}[\text{Sqrt}[b] \text{Sqrt}[2/\text{Pi}] x^{1/6}] \text{Sin}[a])/b^{3/2} + (3x^{1/6} \text{Sin}[a + b x^{1/3}])/b$

#### Rule 3416

$\text{Int}[(a + \text{Cos}[c + (d \cdot x)^n] \cdot b)^p \cdot x^m, x\_Symbol] \rightarrow \text{Module}[\{k = \text{Denominator}[n]\}, \text{Dist}[k, \text{Subst}[\text{Int}[x^{k(m+1)-1} \cdot (a + b \cdot \text{Cos}[c + d \cdot x^{k \cdot n}])^p, x], x, x^{1/k}], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \&\& \text{IntegerQ}[p] \&\& \text{FractionQ}[n]$

#### Rule 3296

$\text{Int}[(c + (d \cdot x)^m) \cdot \sin[e + f \cdot x], x\_Symbol] \rightarrow -\text{Simp}[(c + d \cdot x)^m \cdot \text{Cos}[e + f \cdot x]/f, x] + \text{Dist}[(d \cdot m)/f, \text{Int}[(c + d \cdot x)^{m-1} \cdot \text{Cos}[e + f \cdot x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&\& \text{GtQ}[m, 0]$

#### Rule 3306

$\text{Int}[\sin[e + f \cdot x]/\text{Sqrt}[c + (d \cdot x)], x\_Symbol] \rightarrow \text{Dist}[\text{Cos}[(d \cdot e - c \cdot f)/d], \text{Int}[\text{Sin}[c \cdot f/d + f \cdot x]/\text{Sqrt}[c + d \cdot x], x], x] + \text{Dist}[\text{Sin}[(d \cdot e - c \cdot f)/d], \text{Int}[\text{Cos}[c \cdot f/d + f \cdot x]/\text{Sqrt}[c + d \cdot x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&\& \text{ComplexFreeQ}[f] \&\& \text{NeQ}[d \cdot e - c \cdot f, 0]$

#### Rule 3305

$\text{Int}[\sin[e + f \cdot x]/\text{Sqrt}[c + (d \cdot x)^2], x\_Symbol] \rightarrow \text{Dist}[2/d, \text{Subst}[\text{Int}[\text{Sin}[f \cdot x^2/d], x], x, \text{Sqrt}[c + d \cdot x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&\& \text{ComplexFreeQ}[f] \&\& \text{EqQ}[d \cdot e - c \cdot f, 0]$

#### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)]]/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rule 3304

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)]]/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rubi steps

$$\begin{aligned} \int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx &= 3 \operatorname{Subst} \left( \int \sqrt{x} \cos(a + bx) dx, x, \sqrt[3]{x} \right) \\ &= \frac{3\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{3 \operatorname{Subst} \left( \int \frac{\sin(a+bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right)}{2b} \\ &= \frac{3\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{(3 \cos(a)) \operatorname{Subst} \left( \int \frac{\sin(bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right)}{2b} - \frac{(3 \sin(a)) \operatorname{Subst} \left( \int \frac{\cos(bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right)}{2b} \\ &= \frac{3\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{(3 \cos(a)) \operatorname{Subst} \left( \int \sin(bx^2) dx, x, \sqrt[6]{x} \right)}{b} - \frac{(3 \sin(a)) \operatorname{Subst} \left( \int \cos(bx^2) dx, x, \sqrt[6]{x} \right)}{b} \\ &= -\frac{3\sqrt{\frac{\pi}{2}} \cos(a) S \left( \sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{b^{3/2}} - \frac{3\sqrt{\frac{\pi}{2}} C \left( \sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right) \sin(a)}{b^{3/2}} + \frac{3\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} \end{aligned}$$

**Mathematica [A]** time = 0.152974, size = 94, normalized size = 0.95

$$\frac{3 \left( \sqrt{2\pi} \sin(a) \operatorname{FresnelC} \left( \sqrt{\frac{2}{\pi}} \sqrt{b} \sqrt[6]{x} \right) + \sqrt{2\pi} \cos(a) S \left( \sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right) - 2\sqrt{b} \sqrt[6]{x} \sin(a + b\sqrt[3]{x}) \right)}{2b^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x^(1/3)]/Sqrt[x], x]
```

```
[Out] (-3*(Sqrt[2*Pi]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] + Sqrt[2*Pi]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] - 2*Sqrt[b]*x^(1/6)*Sin[a + b*x^(1/3)])/(2*b^(3/2))
```

**Maple [A]** time = 0.032, size = 64, normalized size = 0.7

$$3 \frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{3\sqrt{2}\sqrt{\pi}}{2} \left( \cos(a) \operatorname{FresnelS} \left( \frac{\sqrt{2}}{\sqrt{\pi}} \sqrt[6]{x} \sqrt{b} \right) + \sin(a) \operatorname{FresnelC} \left( \frac{\sqrt{2}}{\sqrt{\pi}} \sqrt[6]{x} \sqrt{b} \right) \right) b^{-3/2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a+b*x^(1/3))/x^(1/2), x)
```



[Out]  $3x^{1/6} \sin(a+bx^{1/3})/b - 3/2/b^{3/2} * 2^{1/2} * \pi^{1/2} * (\cos(a) * \text{FresnelS}(x^{1/6} * b^{1/2} * 2^{1/2} / \pi^{1/2}) + \sin(a) * \text{FresnelC}(x^{1/6} * b^{1/2} * 2^{1/2} / \pi^{1/2}))$

**Maxima [C]** time = 1.95197, size = 346, normalized size = 3.49

$3 \left( 8x^{1/6} |b| \sin \left( bx^{1/3} + a \right) + \sqrt{\pi} \left( \left( -i \cos \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) - i \cos \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) - \sin \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + \sin \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) \right) \cos(a) - \left( \cos \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + \cos \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) - I \sin \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + I \sin \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) \right) \sin(a) * \text{erf}(\sqrt{I*b} * x^{1/6}) + \left( (I \cos \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + I \cos \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) - \sin \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + \sin \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right)) \cos(a) - \left( \cos \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + \cos \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) + I \sin \left( \frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) - I \sin \left( -\frac{1}{4} \pi + \frac{1}{2} \arctan(0, b) \right) \right) \sin(a) * \text{erf}(\sqrt{-I*b} * x^{1/6}) \right) * \sqrt{\text{abs}(b)} \right) / (b * \text{abs}(b))$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+bx^(1/3))/x^(1/2), x, algorithm="maxima")

[Out]  $3/8 * (8 * x^{1/6} * \text{abs}(b) * \sin(b * x^{1/3} + a) + \sqrt{\pi} * (((-I * \cos(1/4 * \pi + 1/2 * \arctan(0, b)) - I * \cos(-1/4 * \pi + 1/2 * \arctan(0, b)) - \sin(1/4 * \pi + 1/2 * \arctan(0, b)) + \sin(-1/4 * \pi + 1/2 * \arctan(0, b))) * \cos(a) - (\cos(1/4 * \pi + 1/2 * \arctan(0, b)) + \cos(-1/4 * \pi + 1/2 * \arctan(0, b)) - I * \sin(1/4 * \pi + 1/2 * \arctan(0, b)) + I * \sin(-1/4 * \pi + 1/2 * \arctan(0, b))) * \sin(a) * \text{erf}(\sqrt{I * b} * x^{1/6}) + ((I * \cos(1/4 * \pi + 1/2 * \arctan(0, b)) + I * \cos(-1/4 * \pi + 1/2 * \arctan(0, b)) - \sin(1/4 * \pi + 1/2 * \arctan(0, b)) + \sin(-1/4 * \pi + 1/2 * \arctan(0, b))) * \cos(a) - (\cos(1/4 * \pi + 1/2 * \arctan(0, b)) + \cos(-1/4 * \pi + 1/2 * \arctan(0, b)) + I * \sin(1/4 * \pi + 1/2 * \arctan(0, b)) - I * \sin(-1/4 * \pi + 1/2 * \arctan(0, b))) * \sin(a) * \text{erf}(\sqrt{-I * b} * x^{1/6})) * \sqrt{\text{abs}(b)}) / (b * \text{abs}(b))$

**Fricas [A]** time = 1.92261, size = 250, normalized size = 2.53

$$\frac{3 \left( \sqrt{2} \pi \sqrt{\frac{b}{\pi}} \cos(a) S \left( \sqrt{2} x^{1/6} \sqrt{\frac{b}{\pi}} \right) + \sqrt{2} \pi \sqrt{\frac{b}{\pi}} C \left( \sqrt{2} x^{1/6} \sqrt{\frac{b}{\pi}} \right) \sin(a) - 2 b x^{1/6} \sin \left( b x^{1/3} + a \right) \right)}{2 b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+bx^(1/3))/x^(1/2), x, algorithm="fricas")

[Out]  $-3/2 * (\sqrt{2} * \pi * \sqrt{b/\pi} * \cos(a) * \text{fresnel\_sin}(\sqrt{2} * x^{1/6} * \sqrt{b/\pi})) + \sqrt{2} * \pi * \sqrt{b/\pi} * \text{fresnel\_cos}(\sqrt{2} * x^{1/6} * \sqrt{b/\pi}) * \sin(a) - 2 * b * x^{1/6} * \sin(b * x^{1/3} + a) / b^2$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos \left( a + b \sqrt[3]{x} \right)}{\sqrt{x}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+bx\*\*(1/3))/x\*\*(1/2), x)

[Out] Integral(cos(a + b\*x\*\*(1/3))/sqrt(x), x)

**Giac [C]** time = 1.16036, size = 193, normalized size = 1.95

$$\frac{3i\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x^{\frac{1}{6}}\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{(ia)}}{4b\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}} + \frac{3i\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x^{\frac{1}{6}}\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{(-ia)}}{4b\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}} - \frac{3ix^{\frac{1}{6}}e^{\left(ibx^{\frac{1}{3}}+ia\right)}}{2b} + \frac{3ix^{\frac{1}{6}}e^{\left(-ibx^{\frac{1}{3}}-ia\right)}}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))/x^(1/2),x, algorithm="giac")

[Out]  $-3/4*I*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}\left(-1/2*\sqrt{2}*x^{1/6}\right)*\left(-I*b/\operatorname{abs}(b)+1\right)*\sqrt{\operatorname{abs}(b)}*e^{(I*a)}/\left(b*\left(-I*b/\operatorname{abs}(b)+1\right)*\sqrt{\operatorname{abs}(b)}\right) + 3/4*I*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}\left(-1/2*\sqrt{2}*x^{1/6}\right)*\left(I*b/\operatorname{abs}(b)+1\right)*\sqrt{\operatorname{abs}(b)}*e^{(-I*a)}/\left(b*\left(I*b/\operatorname{abs}(b)+1\right)*\sqrt{\operatorname{abs}(b)}\right) - 3/2*I*x^{1/6}*e^{(I*b*x^{1/3}+I*a)}/b + 3/2*I*x^{1/6}*e^{(-I*b*x^{1/3}-I*a)}/b$

$$3.52 \quad \int \frac{\cos(a+b\sqrt[3]{x})}{x^{3/2}} dx$$

**Optimal.** Leaf size=110

$$-4\sqrt{2\pi}b^{3/2} \cos(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right) + 4\sqrt{2\pi}b^{3/2} \sin(a)\text{S}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) + \frac{4b \sin(a+b\sqrt[3]{x})}{\sqrt[6]{x}} - \frac{2 \cos(a+b\sqrt[3]{x})}{\sqrt{x}}$$

[Out] (-2\*Cos[a + b\*x^(1/3)])/Sqrt[x] - 4\*b^(3/2)\*Sqrt[2\*Pi]\*Cos[a]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)] + 4\*b^(3/2)\*Sqrt[2\*Pi]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)]\*Sin[a] + (4\*b\*Sin[a + b\*x^(1/3)])/x^(1/6)

**Rubi [A]** time = 0.142877, antiderivative size = 110, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 7, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$ , Rules used = {3416, 3297, 3306, 3305, 3351, 3304, 3352}

$$-4\sqrt{2\pi}b^{3/2} \cos(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right) + 4\sqrt{2\pi}b^{3/2} \sin(a)\text{S}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) + \frac{4b \sin(a+b\sqrt[3]{x})}{\sqrt[6]{x}} - \frac{2 \cos(a+b\sqrt[3]{x})}{\sqrt{x}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^(1/3)]/x^(3/2), x]

[Out] (-2\*Cos[a + b\*x^(1/3)])/Sqrt[x] - 4\*b^(3/2)\*Sqrt[2\*Pi]\*Cos[a]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)] + 4\*b^(3/2)\*Sqrt[2\*Pi]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)]\*Sin[a] + (4\*b\*Sin[a + b\*x^(1/3)])/x^(1/6)

#### Rule 3416

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + d\*x^(k\*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]

#### Rule 3297

Int[((c\_.) + (d\_.)\*(x\_)^(m\_))\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] :> Simp[((c + d\*x)^(m + 1)\*Sin[e + f\*x])/(d\*(m + 1)), x] - Dist[f/(d\*(m + 1)), Int[(c + d\*x)^(m + 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

#### Rule 3306

Int[sin[(e\_.) + (f\_.)\*(x\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Dist[Cos[(d\*e - c\*f)/d], Int[Sin[(c\*f)/d + f\*x]/Sqrt[c + d\*x], x], x] + Dist[Sin[(d\*e - c\*f)/d], Int[Cos[(c\*f)/d + f\*x]/Sqrt[c + d\*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d\*e - c\*f, 0]

#### Rule 3305

Int[sin[(e\_.) + (f\_.)\*(x\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Dist[2/d, Subst[Int[Sin[(f\*x^2)/d], x], x, Sqrt[c + d\*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d\*e - c\*f, 0]

#### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)]]/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rule 3304

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)]]/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rubi steps

$$\begin{aligned} \int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx &= 3 \operatorname{Subst} \left( \int \frac{\cos(a + bx)}{x^{5/2}} dx, x, \sqrt[3]{x} \right) \\ &= -\frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt{x}} - (2b) \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{x^{3/2}} dx, x, \sqrt[3]{x} \right) \\ &= -\frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} - (4b^2) \operatorname{Subst} \left( \int \frac{\cos(a + bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right) \\ &= -\frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} - (4b^2 \cos(a)) \operatorname{Subst} \left( \int \frac{\cos(bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right) + (4b^2 \sin(a)) \\ &= -\frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} - (8b^2 \cos(a)) \operatorname{Subst} \left( \int \cos(bx^2) dx, x, \sqrt[6]{x} \right) + (8b^2 \sin(a)) \\ &= -\frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt{x}} - 4b^{3/2} \sqrt{2\pi} \cos(a) C \left( \sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right) + 4b^{3/2} \sqrt{2\pi} S \left( \sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right) \sin(a) + \frac{4b \sin(a)}{\sqrt[6]{x}} \end{aligned}$$

**Mathematica [A]** time = 0.248194, size = 110, normalized size = 1.

$$-4\sqrt{2\pi}b^{3/2} \cos(a) \operatorname{FresnelC} \left( \sqrt{\frac{2}{\pi}} \sqrt{b} \sqrt[6]{x} \right) + 4\sqrt{2\pi}b^{3/2} \sin(a) S \left( \sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right) + \frac{4b \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} - \frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt{x}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x^(1/3)]/x^(3/2), x]
```

```
[Out] (-2*Cos[a + b*x^(1/3)])/Sqrt[x] - 4*b^(3/2)*Sqrt[2*Pi]*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] + 4*b^(3/2)*Sqrt[2*Pi]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] + (4*b*Sin[a + b*x^(1/3)])/x^(1/6)
```

**Maple [A]** time = 0.03, size = 78, normalized size = 0.7

$$-2 \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} - 4b \left( -\frac{\sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} + \sqrt{b} \sqrt{2} \sqrt{\pi} \left( \cos(a) \operatorname{FresnelC} \left( \frac{\sqrt[6]{x} \sqrt{b} \sqrt{2}}{\sqrt{\pi}} \right) - \sin(a) \operatorname{FresnelS} \left( \frac{\sqrt[6]{x} \sqrt{b} \sqrt{2}}{\sqrt{\pi}} \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a+b*x^(1/3))/x^(3/2), x)
```

[Out]  $-2\cos(a+bx^{1/3})/x^{1/2}-4b*(-1/x^{1/6}*\sin(a+bx^{1/3})+b^{1/2}*2^{1/2})*\text{Pi}^{1/2}*(\cos(a)*\text{FresnelC}(x^{1/6}*b^{1/2}*2^{1/2}/\text{Pi}^{1/2})-\sin(a)*\text{FresnelS}(x^{1/6}*b^{1/2}*2^{1/2}/\text{Pi}^{1/2}))$

**Maxima [C]** time = 1.85966, size = 358, normalized size = 3.25

$$3\left(\left(\Gamma\left(-\frac{3}{2}, ibx^{\frac{1}{3}}\right) + \Gamma\left(-\frac{3}{2}, -ibx^{\frac{1}{3}}\right)\right) \cos\left(\frac{3}{4}\pi + \frac{3}{2}\arctan(0, b)\right) + \left(\Gamma\left(-\frac{3}{2}, ibx^{\frac{1}{3}}\right) + \Gamma\left(-\frac{3}{2}, -ibx^{\frac{1}{3}}\right)\right) \cos\left(-\frac{3}{4}\pi + \frac{3}{2}\arctan(0, b)\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^(1/3))/x^(3/2), x, algorithm="maxima")`

[Out]  $-3/4*((\text{gamma}(-3/2, I*b*x^{1/3}) + \text{gamma}(-3/2, -I*b*x^{1/3}))*\cos(3/4*\text{pi} + 3/2*\arctan2(0, b)) + (\text{gamma}(-3/2, I*b*x^{1/3}) + \text{gamma}(-3/2, -I*b*x^{1/3}))*\cos(-3/4*\text{pi} + 3/2*\arctan2(0, b)) + (I*\text{gamma}(-3/2, I*b*x^{1/3}) - I*\text{gamma}(-3/2, -I*b*x^{1/3}))*\sin(3/4*\text{pi} + 3/2*\arctan2(0, b)) + (-I*\text{gamma}(-3/2, I*b*x^{1/3}) + I*\text{gamma}(-3/2, -I*b*x^{1/3}))*\sin(-3/4*\text{pi} + 3/2*\arctan2(0, b)))*\cos(a) + ((-I*\text{gamma}(-3/2, I*b*x^{1/3}) + I*\text{gamma}(-3/2, -I*b*x^{1/3}))*\cos(3/4*\text{pi} + 3/2*\arctan2(0, b)) + (-I*\text{gamma}(-3/2, I*b*x^{1/3}) + I*\text{gamma}(-3/2, -I*b*x^{1/3}))*\cos(-3/4*\text{pi} + 3/2*\arctan2(0, b)) + (\text{gamma}(-3/2, I*b*x^{1/3}) + \text{gamma}(-3/2, -I*b*x^{1/3}))*\sin(3/4*\text{pi} + 3/2*\arctan2(0, b)) - (\text{gamma}(-3/2, I*b*x^{1/3}) + \text{gamma}(-3/2, -I*b*x^{1/3}))*\sin(-3/4*\text{pi} + 3/2*\arctan2(0, b)))*\sin(a))*\text{sqrt}(x^{1/3}*\text{abs}(b))*\text{abs}(b)/x^{1/6}$

**Fricas [A]** time = 2.02239, size = 300, normalized size = 2.73

$$\frac{2\left(2\sqrt{2}\pi b x \sqrt{\frac{b}{\pi}} \cos(a) C\left(\sqrt{2}x^{\frac{1}{6}}\sqrt{\frac{b}{\pi}}\right) - 2\sqrt{2}\pi b x \sqrt{\frac{b}{\pi}} S\left(\sqrt{2}x^{\frac{1}{6}}\sqrt{\frac{b}{\pi}}\right) \sin(a) - 2bx^{\frac{5}{6}} \sin\left(bx^{\frac{1}{3}} + a\right) + \sqrt{x} \cos\left(bx^{\frac{1}{3}} + a\right)\right)}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^(1/3))/x^(3/2), x, algorithm="fricas")`

[Out]  $-2*(2*\text{sqrt}(2)*\text{pi}*b*x*\text{sqrt}(b/\text{pi}))*\cos(a)*\text{fresnel\_cos}(\text{sqrt}(2)*x^{1/6}*\text{sqrt}(b/\text{pi})) - 2*\text{sqrt}(2)*\text{pi}*b*x*\text{sqrt}(b/\text{pi}))*\text{fresnel\_sin}(\text{sqrt}(2)*x^{1/6}*\text{sqrt}(b/\text{pi}))*\sin(a) - 2*b*x^{5/6}*\sin(b*x^{1/3} + a) + \text{sqrt}(x)*\cos(b*x^{1/3} + a))/x$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x**(1/3))/x**(3/2), x)`

[Out] `Integral(cos(a + b*x**(1/3))/x**(3/2), x)`

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)}{x^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^(1/3))/x^(3/2),x, algorithm="giac")`

[Out] `integrate(cos(b*x^(1/3) + a)/x^(3/2), x)`

$$3.53 \quad \int \frac{\cos(a+b\sqrt[3]{x})}{x^{5/2}} dx$$

**Optimal.** Leaf size=184

$$-\frac{32}{315}\sqrt{2\pi}b^{9/2}\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right) - \frac{32}{315}\sqrt{2\pi}b^{9/2}\cos(a)S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) + \frac{8b^2\cos(a+b\sqrt[3]{x})}{105x^{5/6}} - \frac{16b^3\sin(a+b\sqrt[3]{x})}{315\sqrt{x}}$$

[Out]  $(-2*\text{Cos}[a + b*x^{(1/3)}])/(3*x^{(3/2)}) + (8*b^2*\text{Cos}[a + b*x^{(1/3)}])/(105*x^{(5/6)}) - (32*b^4*\text{Cos}[a + b*x^{(1/3)}])/(315*x^{(1/6)}) - (32*b^{(9/2)}*\text{Sqrt}[2*\text{Pi}]*\text{Cos}[a]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}])/315 - (32*b^{(9/2)}*\text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}]*\text{Sin}[a])/315 + (4*b*\text{Sin}[a + b*x^{(1/3)}])/(2*1*x^{(7/6)}) - (16*b^3*\text{Sin}[a + b*x^{(1/3)}])/(315*\text{Sqrt}[x])$

**Rubi [A]** time = 0.237778, antiderivative size = 184, normalized size of antiderivative = 1., number of steps used = 11, number of rules used = 7, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$ , Rules used = {3416, 3297, 3306, 3305, 3351, 3304, 3352}

$$-\frac{32}{315}\sqrt{2\pi}b^{9/2}\sin(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right) - \frac{32}{315}\sqrt{2\pi}b^{9/2}\cos(a)S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) + \frac{8b^2\cos(a+b\sqrt[3]{x})}{105x^{5/6}} - \frac{16b^3\sin(a+b\sqrt[3]{x})}{315\sqrt{x}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^(1/3)]/x^(5/2), x]

[Out]  $(-2*\text{Cos}[a + b*x^{(1/3)}])/(3*x^{(3/2)}) + (8*b^2*\text{Cos}[a + b*x^{(1/3)}])/(105*x^{(5/6)}) - (32*b^4*\text{Cos}[a + b*x^{(1/3)}])/(315*x^{(1/6)}) - (32*b^{(9/2)}*\text{Sqrt}[2*\text{Pi}]*\text{Cos}[a]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}])/315 - (32*b^{(9/2)}*\text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}]*\text{Sin}[a])/315 + (4*b*\text{Sin}[a + b*x^{(1/3)}])/(2*1*x^{(7/6)}) - (16*b^3*\text{Sin}[a + b*x^{(1/3)}])/(315*\text{Sqrt}[x])$

#### Rule 3416

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + d\*x^(k\*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]

#### Rule 3297

Int[((c\_.) + (d\_.)\*(x\_)^(m\_))\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] :> Simp[((c + d\*x)^(m + 1)\*Sin[e + f\*x])/(d\*(m + 1)), x] - Dist[f/(d\*(m + 1)), Int[(c + d\*x)^(m + 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

#### Rule 3306

Int[sin[(e\_.) + (f\_.)\*(x\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Dist[Cos[(d\*e - c\*f)/d], Int[Sin[(c\*f)/d + f\*x]/Sqrt[c + d\*x], x], x] + Dist[Sin[(d\*e - c\*f)/d], Int[Cos[(c\*f)/d + f\*x]/Sqrt[c + d\*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d\*e - c\*f, 0]

#### Rule 3305

Int[sin[(e\_.) + (f\_.)\*(x\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Dist[2/d, Subst[Int[Sin[(f\*x^2)/d], x], x, Sqrt[c + d\*x]], x] /; FreeQ[{c, d, e, f}

, x] && ComplexFreeQ[f] && EqQ[d\*e - c\*f, 0]

### Rule 3351

Int[Sin[(d\_.)\*((e\_.) + (f\_.)\*(x\_))<sup>2</sup>], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)]]/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

### Rule 3304

Int[sin[Pi/2 + (e\_.) + (f\_.)\*(x\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] := Dist[2/d, Subst[Int[Cos[(f\*x<sup>2</sup>)/d], x], x, Sqrt[c + d\*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d\*e - c\*f, 0]

### Rule 3352

Int[Cos[(d\_.)\*((e\_.) + (f\_.)\*(x\_))<sup>2</sup>], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)]]/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

### Rubi steps

$$\begin{aligned}
 \int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx &= 3 \operatorname{Subst} \left( \int \frac{\cos(a + bx)}{x^{11/2}} dx, x, \sqrt[3]{x} \right) \\
 &= -\frac{2 \cos(a + b\sqrt[3]{x})}{3x^{3/2}} - \frac{1}{3}(2b) \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{x^{9/2}} dx, x, \sqrt[3]{x} \right) \\
 &= -\frac{2 \cos(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{4b \sin(a + b\sqrt[3]{x})}{21x^{7/6}} - \frac{1}{21}(4b^2) \operatorname{Subst} \left( \int \frac{\cos(a + bx)}{x^{7/2}} dx, x, \sqrt[3]{x} \right) \\
 &= -\frac{2 \cos(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{105x^{5/6}} + \frac{4b \sin(a + b\sqrt[3]{x})}{21x^{7/6}} + \frac{1}{105}(8b^3) \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{x^{5/2}} dx, x, \sqrt[3]{x} \right) \\
 &= -\frac{2 \cos(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{105x^{5/6}} + \frac{4b \sin(a + b\sqrt[3]{x})}{21x^{7/6}} - \frac{16b^3 \sin(a + b\sqrt[3]{x})}{315\sqrt{x}} + \frac{1}{315}(16b^4) \operatorname{Subst} \left( \int \frac{\cos(a + bx)}{x^{3/2}} dx, x, \sqrt[3]{x} \right) \\
 &= -\frac{2 \cos(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{21x^{7/6}} - \frac{16b^3 \sin(a + b\sqrt[3]{x})}{315\sqrt{x}} \\
 &= -\frac{2 \cos(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{21x^{7/6}} - \frac{16b^3 \sin(a + b\sqrt[3]{x})}{315\sqrt{x}} \\
 &= -\frac{2 \cos(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{21x^{7/6}} - \frac{16b^3 \sin(a + b\sqrt[3]{x})}{315\sqrt{x}} \\
 &= -\frac{2 \cos(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} - \frac{32}{315}b^{9/2}\sqrt{2\pi} \cos(a)S \left( \sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x} \right)
 \end{aligned}$$

**Mathematica [A]** time = 0.238466, size = 180, normalized size = 0.98

$$\frac{2 \left( 16\sqrt{2\pi}b^{9/2}x^{3/2} \sin(a) \operatorname{FresnelC} \left( \sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x} \right) + 16\sqrt{2\pi}b^{9/2}x^{3/2} \cos(a)S \left( \sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x} \right) + 16b^4x^{4/3} \cos(a + b\sqrt[3]{x}) - 12b^2x^{2/3} \cos(a + b\sqrt[3]{x}) + 12b^{4/3} \cos(a + b\sqrt[3]{x}) \right)}{315x^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^(1/3)]/x^(5/2), x]

[Out] (-2\*(105\*Cos[a + b\*x^(1/3)] - 12\*b^2\*x^(2/3)\*Cos[a + b\*x^(1/3)] + 16\*b^4\*x^(4/3)\*Cos[a + b\*x^(1/3)] + 16\*b^(9/2)\*Sqrt[2\*Pi]\*x^(3/2)\*Cos[a]\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)] + 16\*b^(9/2)\*Sqrt[2\*Pi]\*x^(3/2)\*FresnelC[Sqrt[b]\*



$$\frac{\sqrt{2/\pi} x^{1/6} \sin[a] - 30 b x^{1/3} \sin[a + b x^{1/3}] + 8 b^3 x \sin[a + b x^{1/3}]}{315 x^{3/2}}$$

**Maple [A]** time = 0.03, size = 129, normalized size = 0.7

$$-\frac{2}{3} \cos(a + b\sqrt[3]{x}) x^{-\frac{3}{2}} - \frac{4b}{3} \left( -\frac{1}{7} \sin(a + b\sqrt[3]{x}) x^{-\frac{7}{6}} + \frac{2b}{7} \left( -\frac{1}{5} \cos(a + b\sqrt[3]{x}) x^{-\frac{5}{6}} - \frac{2b}{5} \left( -\frac{1}{3} \sin(a + b\sqrt[3]{x}) \frac{1}{\sqrt{x}} + \frac{2b}{3} \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^(1/3))/x^(5/2),x)

[Out]  $-2/3 \cos(a + b x^{1/3}) / x^{3/2} - 4/3 b \left( -1/7 x^{7/6} \sin(a + b x^{1/3}) + 2/7 b \left( -1/5 x^{5/6} \cos(a + b x^{1/3}) - 2/5 b \left( -1/3 x^{1/2} \sin(a + b x^{1/3}) + 2/3 b \left( -1/x^{1/6} \cos(a + b x^{1/3}) - b^{1/2} 2^{1/2} \pi^{1/2} (\cos(a) \operatorname{FresnelS}(x^{1/6}) * b^{1/2} 2^{1/2} / \pi^{1/2}) + \sin(a) \operatorname{FresnelC}(x^{1/6}) * b^{1/2} 2^{1/2} / \pi^{1/2}) \right) \right) \right)$

**Maxima [C]** time = 1.50286, size = 359, normalized size = 1.95

$$3 \left( \left( \Gamma\left(-\frac{9}{2}, i b x^{\frac{1}{3}}\right) + \Gamma\left(-\frac{9}{2}, -i b x^{\frac{1}{3}}\right) \right) \cos\left(\frac{9}{4} \pi + \frac{9}{2} \arctan(0, b)\right) + \left( \Gamma\left(-\frac{9}{2}, i b x^{\frac{1}{3}}\right) + \Gamma\left(-\frac{9}{2}, -i b x^{\frac{1}{3}}\right) \right) \cos\left(-\frac{9}{4} \pi + \frac{9}{2} \arctan(0, b)\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))/x^(5/2),x, algorithm="maxima")

[Out]  $-3/4 * \left( \left( \Gamma(-9/2, I * b * x^{1/3}) + \Gamma(-9/2, -I * b * x^{1/3}) \right) * \cos(9/4 * \pi + 9/2 * \arctan(0, b)) + \left( \Gamma(-9/2, I * b * x^{1/3}) + \Gamma(-9/2, -I * b * x^{1/3}) \right) * \cos(-9/4 * \pi + 9/2 * \arctan(0, b)) + \left( \Gamma(-9/2, I * b * x^{1/3}) - \Gamma(-9/2, -I * b * x^{1/3}) \right) * \sin(9/4 * \pi + 9/2 * \arctan(0, b)) + \left( -\Gamma(-9/2, I * b * x^{1/3}) + \Gamma(-9/2, -I * b * x^{1/3}) \right) * \sin(-9/4 * \pi + 9/2 * \arctan(0, b)) \right) * \cos(a) + \left( \left( -\Gamma(-9/2, I * b * x^{1/3}) + \Gamma(-9/2, -I * b * x^{1/3}) \right) * \cos(9/4 * \pi + 9/2 * \arctan(0, b)) + \left( -\Gamma(-9/2, I * b * x^{1/3}) + \Gamma(-9/2, -I * b * x^{1/3}) \right) * \cos(-9/4 * \pi + 9/2 * \arctan(0, b)) + \left( \Gamma(-9/2, I * b * x^{1/3}) + \Gamma(-9/2, -I * b * x^{1/3}) \right) * \sin(9/4 * \pi + 9/2 * \arctan(0, b)) - \left( \Gamma(-9/2, I * b * x^{1/3}) + \Gamma(-9/2, -I * b * x^{1/3}) \right) * \sin(-9/4 * \pi + 9/2 * \arctan(0, b)) \right) * \sin(a) * \sqrt{x^{1/3} * \text{abs}(b)} * b^4 / x^{1/6}$

**Fricas [A]** time = 2.07561, size = 405, normalized size = 2.2

$$\frac{2 \left( 16 \sqrt{2} \pi b^4 x^2 \sqrt{\frac{b}{\pi}} \cos(a) S\left(\sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) + 16 \sqrt{2} \pi b^4 x^2 \sqrt{\frac{b}{\pi}} C\left(\sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) \sin(a) + \left( 16 b^4 x^{\frac{11}{6}} - 12 b^2 x^{\frac{7}{6}} + 105 \sqrt{x} \right) \cos(a) \right)}{315 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))/x^(5/2),x, algorithm="fricas")

[Out]  $-2/315 * \left( 16 * \sqrt{2} * \pi * b^4 * x^2 * \sqrt{b/\pi} * \cos(a) * \operatorname{fresnel\_sin}(\sqrt{2} * x^{1/6} * \sqrt{b/\pi}) + 16 * \sqrt{2} * \pi * b^4 * x^2 * \sqrt{b/\pi} * \operatorname{fresnel\_cos}(\sqrt{2} * x^{1/6} * \sqrt{b/\pi}) \right)$

```
*sqrt(b/pi))*sin(a) + (16*b^4*x^(11/6) - 12*b^2*x^(7/6) + 105*sqrt(x))*cos(
b*x^(1/3) + a) + 2*(4*b^3*x^(3/2) - 15*b*x^(5/6))*sin(b*x^(1/3) + a))/x^2
```

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + b\sqrt[3]{x}\right)}{x^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x**(1/3))/x**(5/2),x)
```

```
[Out] Integral(cos(a + b*x**(1/3))/x**(5/2), x)
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)}{x^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^(1/3))/x^(5/2),x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^(1/3) + a)/x^(5/2), x)
```

$$3.54 \quad \int \frac{\cos(a+b\sqrt[3]{x})}{x^{7/2}} dx$$

**Optimal.** Leaf size=250

$$\frac{256\sqrt{2\pi}b^{15/2} \cos(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right)}{675675} - \frac{256\sqrt{2\pi}b^{15/2} \sin(a)S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{675675} + \frac{64b^5 \sin(a+b\sqrt[3]{x})}{225225x^{5/6}} - \frac{16b^3 \sin(a+b\sqrt[3]{x})}{6435x^{3/2}}$$

[Out]  $(-2*\text{Cos}[a + b*x^{(1/3)}])/(5*x^{(5/2)}) + (8*b^2*\text{Cos}[a + b*x^{(1/3)}])/(715*x^{(11/6)}) - (32*b^4*\text{Cos}[a + b*x^{(1/3)}])/(45045*x^{(7/6)}) + (128*b^6*\text{Cos}[a + b*x^{(1/3)}])/(675675*\text{Sqrt}[x]) + (256*b^{(15/2)}*\text{Sqrt}[2*\text{Pi}]*\text{Cos}[a]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}])/675675 - (256*b^{(15/2)}*\text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}]*\text{Sin}[a])/675675 + (4*b*\text{Sin}[a + b*x^{(1/3)}])/(65*x^{(13/6)}) - (16*b^3*\text{Sin}[a + b*x^{(1/3)}])/(6435*x^{(3/2)}) + (64*b^5*\text{Sin}[a + b*x^{(1/3)}])/(225225*x^{(5/6)}) - (256*b^7*\text{Sin}[a + b*x^{(1/3)}])/(675675*x^{(1/6)})$

**Rubi [A]** time = 0.336934, antiderivative size = 250, normalized size of antiderivative = 1., number of steps used = 14, number of rules used = 7, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$ , Rules used = {3416, 3297, 3306, 3305, 3351, 3304, 3352}

$$\frac{256\sqrt{2\pi}b^{15/2} \cos(a)\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}\sqrt{b}\sqrt[6]{x}\right)}{675675} - \frac{256\sqrt{2\pi}b^{15/2} \sin(a)S\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{675675} + \frac{64b^5 \sin(a+b\sqrt[3]{x})}{225225x^{5/6}} - \frac{16b^3 \sin(a+b\sqrt[3]{x})}{6435x^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^(1/3)]/x^(7/2), x]

[Out]  $(-2*\text{Cos}[a + b*x^{(1/3)}])/(5*x^{(5/2)}) + (8*b^2*\text{Cos}[a + b*x^{(1/3)}])/(715*x^{(11/6)}) - (32*b^4*\text{Cos}[a + b*x^{(1/3)}])/(45045*x^{(7/6)}) + (128*b^6*\text{Cos}[a + b*x^{(1/3)}])/(675675*\text{Sqrt}[x]) + (256*b^{(15/2)}*\text{Sqrt}[2*\text{Pi}]*\text{Cos}[a]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}])/675675 - (256*b^{(15/2)}*\text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}]*\text{Sin}[a])/675675 + (4*b*\text{Sin}[a + b*x^{(1/3)}])/(65*x^{(13/6)}) - (16*b^3*\text{Sin}[a + b*x^{(1/3)}])/(6435*x^{(3/2)}) + (64*b^5*\text{Sin}[a + b*x^{(1/3)}])/(225225*x^{(5/6)}) - (256*b^7*\text{Sin}[a + b*x^{(1/3)}])/(675675*x^{(1/6)})$

#### Rule 3416

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + d\*x^(k\*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]

#### Rule 3297

Int[((c\_.) + (d\_.)\*(x\_)^(m\_))\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] :> Simp[(c + d\*x)^(m + 1)\*Sin[e + f\*x]/(d\*(m + 1)), x] - Dist[f/(d\*(m + 1)), Int[(c + d\*x)^(m + 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

#### Rule 3306

Int[sin[(e\_.) + (f\_.)\*(x\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Dist[Cos[(d\*e - c\*f)/d], Int[Sin[(c\*f)/d + f\*x]/Sqrt[c + d\*x], x], x] + Dist[Sin[(d\*e - c\*f)/d], Int[Cos[(c\*f)/d + f\*x]/Sqrt[c + d\*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d\*e - c\*f, 0]

Rule 3305

Int[sin[(e\_.) + (f\_.)\*(x\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] := Dist[2/d, Subst[Int[Sin[(f\*x^2)/d], x], x, Sqrt[c + d\*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d\*e - c\*f, 0]

Rule 3351

Int[Sin[(d\_.)\*((e\_.) + (f\_.)\*(x\_))^2], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)]]/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

Rule 3304

Int[sin[Pi/2 + (e\_.) + (f\_.)\*(x\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] := Dist[2/d, Subst[Int[Cos[(f\*x^2)/d], x], x, Sqrt[c + d\*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d\*e - c\*f, 0]

Rule 3352

Int[Cos[(d\_.)\*((e\_.) + (f\_.)\*(x\_))^2], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)]]/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx &= 3 \operatorname{Subst} \left( \int \frac{\cos(a + bx)}{x^{17/2}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} - \frac{1}{5}(2b) \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{x^{15/2}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{4b \sin(a + b\sqrt[3]{x})}{65x^{13/6}} - \frac{1}{65}(4b^2) \operatorname{Subst} \left( \int \frac{\cos(a + bx)}{x^{13/2}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{715x^{11/6}} + \frac{4b \sin(a + b\sqrt[3]{x})}{65x^{13/6}} + \frac{1}{715}(8b^3) \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{x^{11/2}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{715x^{11/6}} + \frac{4b \sin(a + b\sqrt[3]{x})}{65x^{13/6}} - \frac{16b^3 \sin(a + b\sqrt[3]{x})}{6435x^{3/2}} + \frac{(16b^4) \operatorname{Subst} \left( \int \frac{\cos(a + bx)}{x^{9/2}} dx, x, \sqrt[3]{x} \right)}{6435x^{3/2}} \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{45045x^{7/6}} + \frac{4b \sin(a + b\sqrt[3]{x})}{65x^{13/6}} - \frac{16b^3 \sin(a + b\sqrt[3]{x})}{6435x^{3/2}} \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{45045x^{7/6}} + \frac{4b \sin(a + b\sqrt[3]{x})}{65x^{13/6}} - \frac{16b^3 \sin(a + b\sqrt[3]{x})}{6435x^{3/2}} \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{45045x^{7/6}} + \frac{128b^6 \cos(a + b\sqrt[3]{x})}{675675\sqrt{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{65x^{13/6}} \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{45045x^{7/6}} + \frac{128b^6 \cos(a + b\sqrt[3]{x})}{675675\sqrt{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{65x^{13/6}} \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{45045x^{7/6}} + \frac{128b^6 \cos(a + b\sqrt[3]{x})}{675675\sqrt{x}} + \frac{4b \sin(a + b\sqrt[3]{x})}{65x^{13/6}} \\
&= -\frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b^2 \cos(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{32b^4 \cos(a + b\sqrt[3]{x})}{45045x^{7/6}} + \frac{128b^6 \cos(a + b\sqrt[3]{x})}{675675\sqrt{x}} + \frac{256b^{15/2}}{675675\sqrt{x}}
\end{aligned}$$

**Mathematica [A]** time = 0.316166, size = 238, normalized size = 0.95

$$2 \left( 128 \sqrt{2\pi} b^{15/2} x^{5/2} \cos(a) \operatorname{FresnelC} \left( \sqrt{\frac{2}{\pi}} \sqrt{b} \sqrt[3]{x} \right) - 128 \sqrt{2\pi} b^{15/2} x^{5/2} \sin(a) S \left( \sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[3]{x} \right) - 128 b^7 x^{7/3} \sin(a + b \sqrt[3]{x}) + \right.$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^(1/3)]/x^(7/2), x]

[Out] (2\*(-135135\*Cos[a + b\*x^(1/3)] + 3780\*b^2\*x^(2/3)\*Cos[a + b\*x^(1/3)] - 240\*b^4\*x^(4/3)\*Cos[a + b\*x^(1/3)] + 64\*b^6\*x^2\*Cos[a + b\*x^(1/3)] + 128\*b^(15/2)\*Sqrt[2\*Pi]\*x^(5/2)\*Cos[a]\*FresnelC[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)] - 128\*b^(15/2)\*Sqrt[2\*Pi]\*x^(5/2)\*FresnelS[Sqrt[b]\*Sqrt[2/Pi]\*x^(1/6)]\*Sin[a] + 20790\*b\*x^(1/3)\*Sin[a + b\*x^(1/3)] - 840\*b^3\*x\*Sin[a + b\*x^(1/3)] + 96\*b^5\*x^(5/3)\*Sin[a + b\*x^(1/3)] - 128\*b^7\*x^(7/3)\*Sin[a + b\*x^(1/3)])/(675675\*x^(5/2))

**Maple [A]** time = 0.032, size = 180, normalized size = 0.7

$$-\frac{2}{5} \cos(a + b \sqrt[3]{x}) x^{-\frac{5}{2}} - \frac{4b}{5} \left( -\frac{1}{13} \sin(a + b \sqrt[3]{x}) x^{-\frac{13}{6}} + \frac{2b}{13} \left( -\frac{1}{11} \cos(a + b \sqrt[3]{x}) x^{-\frac{11}{6}} - \frac{2b}{11} \left( -\frac{1}{9} \sin(a + b \sqrt[3]{x}) x^{-\frac{3}{2}} + \frac{2}{9} \cos(a + b \sqrt[3]{x}) x^{-\frac{5}{2}} \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^(1/3))/x^(7/2), x)

[Out] -2/5\*cos(a+b\*x^(1/3))/x^(5/2)-4/5\*b\*(-1/13/x^(13/6)\*sin(a+b\*x^(1/3))+2/13\*b\*(-1/11/x^(11/6)\*cos(a+b\*x^(1/3))-2/11\*b\*(-1/9/x^(3/2)\*sin(a+b\*x^(1/3))+2/9\*b\*(-1/7/x^(7/6)\*cos(a+b\*x^(1/3))-2/7\*b\*(-1/5/x^(5/6)\*sin(a+b\*x^(1/3))+2/5\*b\*(-1/3\*cos(a+b\*x^(1/3))/x^(1/2)-2/3\*b\*(-1/x^(1/6)\*sin(a+b\*x^(1/3))+b^(1/2)\*2^(1/2)\*Pi^(1/2)\*(cos(a)\*FresnelC(x^(1/6)\*b^(1/2)\*2^(1/2)/Pi^(1/2))-sin(a)\*FresnelS(x^(1/6)\*b^(1/2)\*2^(1/2)/Pi^(1/2))))))

**Maxima [C]** time = 1.43609, size = 362, normalized size = 1.45

$$3 \left( \left( \Gamma \left( -\frac{15}{2}, i b x^{\frac{1}{3}} \right) + \Gamma \left( -\frac{15}{2}, -i b x^{\frac{1}{3}} \right) \right) \cos \left( \frac{15}{4} \pi + \frac{15}{2} \arctan(0, b) \right) + \left( \Gamma \left( -\frac{15}{2}, i b x^{\frac{1}{3}} \right) + \Gamma \left( -\frac{15}{2}, -i b x^{\frac{1}{3}} \right) \right) \cos \left( -\frac{15}{4} \pi \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))/x^(7/2), x, algorithm="maxima")

[Out] -3/4\*(((gamma(-15/2, I\*b\*x^(1/3)) + gamma(-15/2, -I\*b\*x^(1/3)))\*cos(15/4\*pi + 15/2\*arctan2(0, b)) + (gamma(-15/2, I\*b\*x^(1/3)) + gamma(-15/2, -I\*b\*x^(1/3)))\*cos(-15/4\*pi + 15/2\*arctan2(0, b)) + (I\*gamma(-15/2, I\*b\*x^(1/3)) - I\*gamma(-15/2, -I\*b\*x^(1/3)))\*sin(15/4\*pi + 15/2\*arctan2(0, b)) + (-I\*gamma(-15/2, I\*b\*x^(1/3)) + I\*gamma(-15/2, -I\*b\*x^(1/3)))\*sin(-15/4\*pi + 15/2\*arctan2(0, b)))\*cos(a) + ((-I\*gamma(-15/2, I\*b\*x^(1/3)) + I\*gamma(-15/2, -I\*b\*x^(1/3)))\*cos(15/4\*pi + 15/2\*arctan2(0, b)) + (-I\*gamma(-15/2, I\*b\*x^(1/3)) + I\*gamma(-15/2, -I\*b\*x^(1/3)))\*cos(-15/4\*pi + 15/2\*arctan2(0, b)) + (gamma(-15/2, I\*b\*x^(1/3)) + gamma(-15/2, -I\*b\*x^(1/3)))\*sin(15/4\*pi + 15/2\*arctan2(0, b)) - (gamma(-15/2, I\*b\*x^(1/3)) + gamma(-15/2, -I\*b\*x^(1/3)))\*sin(

$-15/4\pi + 15/2\arctan2(0, b))\sin(a)\sqrt{x^{1/3}\text{abs}(b)}b^6\text{abs}(b)/x^{1/6}$

**Fricas [A]** time = 2.13131, size = 490, normalized size = 1.96

$$\frac{2\left(128\sqrt{2}\pi b^7 x^3 \sqrt{\frac{b}{\pi}} \cos(a) C\left(\sqrt{2}x^{\frac{1}{6}}\sqrt{\frac{b}{\pi}}\right) - 128\sqrt{2}\pi b^7 x^3 \sqrt{\frac{b}{\pi}} S\left(\sqrt{2}x^{\frac{1}{6}}\sqrt{\frac{b}{\pi}}\right) \sin(a) - \left(240b^4 x^{\frac{11}{6}} - 3780b^2 x^{\frac{7}{6}} - (64b^6 x^2 - 135135)\sqrt{x}\right)\cos(bx^{1/3} + a) + 2(48b^5 x^{13/6} - 420b^3 x^{3/2} - (64b^7 x^2 - 10395b)x^{5/6})\sin(bx^{1/3} + a)\right)}{675675x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))/x^(7/2),x, algorithm="fricas")

[Out]  $2/675675*(128*\sqrt{2}*\pi*b^7*x^3*\sqrt{b/\pi}*\cos(a)*\text{fresnel\_cos}(\sqrt{2}*x^{1/6}*\sqrt{b/\pi}) - 128*\sqrt{2}*\pi*b^7*x^3*\sqrt{b/\pi}*\text{fresnel\_sin}(\sqrt{2}*x^{1/6}*\sqrt{b/\pi})*\sin(a) - (240*b^4*x^{11/6} - 3780*b^2*x^{7/6} - (64*b^6*x^2 - 135135)*\sqrt{x})*\cos(b*x^{1/3} + a) + 2*(48*b^5*x^{13/6} - 420*b^3*x^{3/2} - (64*b^7*x^2 - 10395*b)*x^{5/6})*\sin(b*x^{1/3} + a))/x^3$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + b\sqrt[3]{x}\right)}{x^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x\*\*(1/3))/x\*\*(7/2),x)

[Out] Integral(cos(a + b\*x\*\*(1/3))/x\*\*(7/2), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)}{x^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))/x^(7/2),x, algorithm="giac")

[Out] integrate(cos(b\*x^(1/3) + a)/x^(7/2), x)

### 3.55 $\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx$

**Optimal.** Leaf size=310

$$\frac{405405\sqrt{\pi} \sin(2a)\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{32768b^{15/2}} + \frac{405405\sqrt{\pi} \cos(2a)\text{S}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{32768b^{15/2}} + \frac{39x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4}$$

```
[Out] (-135135*Sqrt[x])/(4096*b^6) + (3861*x^(7/6))/(256*b^4) - (39*x^(11/6))/(16
*b^2) + x^(5/2)/5 + (135135*Sqrt[x]*Cos[a + b*x^(1/3)]^2)/(2048*b^6) - (386
1*x^(7/6)*Cos[a + b*x^(1/3)]^2)/(128*b^4) + (39*x^(11/6)*Cos[a + b*x^(1/3)]
^2)/(8*b^2) + (405405*Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[P
i]])/(32768*b^(15/2)) + (405405*Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[
Pi]]*Sin[2*a])/(32768*b^(15/2)) + (27027*x^(5/6)*Cos[a + b*x^(1/3)]*Sin[a +
b*x^(1/3)])/(512*b^5) - (429*x^(3/2)*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)]
)/(32*b^3) + (3*x^(13/6)*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(2*b) - (40
5405*x^(1/6)*Sin[2*(a + b*x^(1/3))])/(16384*b^7)
```

**Rubi [A]** time = 0.359734, antiderivative size = 310, normalized size of antiderivative = 1., number of steps used = 15, number of rules used = 10, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$ , Rules used = {3416, 3311, 30, 3312, 3296, 3306, 3305, 3351, 3304, 3352}

$$\frac{405405\sqrt{\pi} \sin(2a)\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{32768b^{15/2}} + \frac{405405\sqrt{\pi} \cos(2a)\text{S}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{32768b^{15/2}} + \frac{39x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4}$$

Antiderivative was successfully verified.

```
[In] Int[x^(3/2)*Cos[a + b*x^(1/3)]^2,x]
```

```
[Out] (-135135*Sqrt[x])/(4096*b^6) + (3861*x^(7/6))/(256*b^4) - (39*x^(11/6))/(16
*b^2) + x^(5/2)/5 + (135135*Sqrt[x]*Cos[a + b*x^(1/3)]^2)/(2048*b^6) - (386
1*x^(7/6)*Cos[a + b*x^(1/3)]^2)/(128*b^4) + (39*x^(11/6)*Cos[a + b*x^(1/3)]
^2)/(8*b^2) + (405405*Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[P
i]])/(32768*b^(15/2)) + (405405*Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[
Pi]]*Sin[2*a])/(32768*b^(15/2)) + (27027*x^(5/6)*Cos[a + b*x^(1/3)]*Sin[a +
b*x^(1/3)])/(512*b^5) - (429*x^(3/2)*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)]
)/(32*b^3) + (3*x^(13/6)*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(2*b) - (40
5405*x^(1/6)*Sin[2*(a + b*x^(1/3))])/(16384*b^7)
```

#### Rule 3416

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol
] :> Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k*(m + 1) - 1)*(a +
b*Cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x]
&& IntegerQ[p] && FractionQ[n]
```

#### Rule 3311

```
Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbo
l] :> Simp[(d*m*(c + d*x)^(m - 1)*(b*Ssin[e + f*x])^n)/(f^2*n^2), x] + (Dist
[(b^2*(n - 1))/n, Int[(c + d*x)^(m)*(b*Ssin[e + f*x])^(n - 2), x], x] - Dist[
(d^2*m*(m - 1))/(f^2*n^2), Int[(c + d*x)^(m - 2)*(b*Ssin[e + f*x])^n, x]
- Simp[(b*(c + d*x)^(m)*Cos[e + f*x]*(b*Ssin[e + f*x])^(n - 1))/(f*n), x]) /;
FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]
```

#### Rule 30

```
Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]
```

### Rule 3312

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

### Rule 3296

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := -Simp[(c + d*x)^m*Cos[e + f*x]/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

### Rule 3306

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]
```

### Rule 3305

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d, Subst[Int[Sin[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rule 3304

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rubi steps



$$\begin{aligned}
\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx &= 3 \operatorname{Subst}\left(\int x^{13/2} \cos^2(a + bx) dx, x, \sqrt[3]{x}\right) \\
&= \frac{39x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{3x^{13/6} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{2b} + \frac{3}{2} \operatorname{Subst}\left(\int x^{13/2} dx, x, \sqrt[3]{x}\right) \\
&= \frac{x^{5/2}}{5} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{39x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{429x^{3/2} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{32b^3} \\
&= -\frac{39x^{11/6}}{16b^2} + \frac{x^{5/2}}{5} + \frac{135135\sqrt{x} \cos^2(a + b\sqrt[3]{x})}{2048b^6} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{39x^{3/2} \cos^2(a + b\sqrt[3]{x})}{8b^2} \\
&= \frac{3861x^{7/6}}{256b^4} - \frac{39x^{11/6}}{16b^2} + \frac{x^{5/2}}{5} + \frac{135135\sqrt{x} \cos^2(a + b\sqrt[3]{x})}{2048b^6} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{39x^{3/2} \cos^2(a + b\sqrt[3]{x})}{8b^2} \\
&= -\frac{135135\sqrt{x}}{4096b^6} + \frac{3861x^{7/6}}{256b^4} - \frac{39x^{11/6}}{16b^2} + \frac{x^{5/2}}{5} + \frac{135135\sqrt{x} \cos^2(a + b\sqrt[3]{x})}{2048b^6} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4} \\
&= -\frac{135135\sqrt{x}}{4096b^6} + \frac{3861x^{7/6}}{256b^4} - \frac{39x^{11/6}}{16b^2} + \frac{x^{5/2}}{5} + \frac{135135\sqrt{x} \cos^2(a + b\sqrt[3]{x})}{2048b^6} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4} \\
&= -\frac{135135\sqrt{x}}{4096b^6} + \frac{3861x^{7/6}}{256b^4} - \frac{39x^{11/6}}{16b^2} + \frac{x^{5/2}}{5} + \frac{135135\sqrt{x} \cos^2(a + b\sqrt[3]{x})}{2048b^6} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4} \\
&= -\frac{135135\sqrt{x}}{4096b^6} + \frac{3861x^{7/6}}{256b^4} - \frac{39x^{11/6}}{16b^2} + \frac{x^{5/2}}{5} + \frac{135135\sqrt{x} \cos^2(a + b\sqrt[3]{x})}{2048b^6} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4}
\end{aligned}$$

**Mathematica [A]** time = 0.658827, size = 174, normalized size = 0.56

$$\frac{2\sqrt{b}\sqrt[6]{x} \left(15(4096b^6x^2 - 36608b^4x^{4/3} + 144144b^2x^{2/3} - 135135) \sin(2(a + b\sqrt[3]{x})) + 780(256b^5x^{5/3} - 1584b^3x + 3465)\right)}{163840}$$

Antiderivative was successfully verified.

[In] Integrate[x^(3/2)\*Cos[a + b\*x^(1/3)]^2,x]

[Out] (2027025\*Sqrt[Pi]\*Cos[2\*a]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]] + 2027025\*Sqrt[Pi]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a] + 2\*Sqrt[b]\*x^(1/6)\*(16384\*b^7\*x^(7/3) + 780\*(3465\*b\*x^(1/3) - 1584\*b^3\*x + 256\*b^5\*x^(5/3))\*Cos[2\*(a + b\*x^(1/3))] + 15\*(-135135 + 144144\*b^2\*x^(2/3) - 36608\*b^4\*x^(4/3) + 4096\*b^6\*x^2)\*Sin[2\*(a + b\*x^(1/3))])/ (163840\*b^(15/2))

**Maple [A]** time = 0.04, size = 219, normalized size = 0.7

$$\frac{1}{5}x^{5/2} + \frac{3}{4b}x^{13/6} \sin(2a + 2b\sqrt[3]{x}) - \frac{39}{4b} \left( -\frac{1}{4b}x^{11/6} \cos(2a + 2b\sqrt[3]{x}) + \frac{11}{4b} \left( \frac{1}{4b}x^{3/2} \sin(2a + 2b\sqrt[3]{x}) - \frac{9}{4b} \left( -\frac{1}{4b}x^{7/6} \cos(2a + 2b\sqrt[3]{x}) \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(3/2)\*cos(a+b\*x^(1/3))^2,x)

[Out] 1/5\*x^(5/2)+3/4/b\*x^(13/6)\*sin(2\*a+2\*b\*x^(1/3))-39/4/b\*(-1/4/b\*x^(11/6)\*cos(2\*a+2\*b\*x^(1/3))+11/4/b\*(1/4/b\*x^(3/2)\*sin(2\*a+2\*b\*x^(1/3))-9/4/b\*(-1/4/b\*x^(7/6)\*cos(2\*a+2\*b\*x^(1/3))+7/4/b\*(1/4/b\*x^(5/6)\*sin(2\*a+2\*b\*x^(1/3)))-5/4/

$$b*(-1/4/b*x^{(1/2)}*\cos(2*a+2*b*x^{(1/3)})+3/4/b*(1/4*x^{(1/6)}*\sin(2*a+2*b*x^{(1/3)}))/b-1/8/b^{(3/2)}*Pi^{(1/2)}*(\cos(2*a)*\text{FresnelS}(2*x^{(1/6)}*b^{(1/2)}/Pi^{(1/2)})+\sin(2*a)*\text{FresnelC}(2*x^{(1/6)}*b^{(1/2)}/Pi^{(1/2)}))))))$$

**Maxima [C]** time = 2.14783, size = 497, normalized size = 1.6

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(a+b\*x^(1/3))^2,x, algorithm="maxima")

[Out] 1/1310720\*(262144\*b^7\*x^(5/2)\*abs(b) - sqrt(2)\*sqrt(pi)\*((( -2027025\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - 2027025\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 2027025\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 2027025\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(2\*a) - (2027025\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 2027025\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 2027025\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 2027025\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(2\*a)) \* erf(sqrt(2\*I\*b)\*x^(1/6)) + ((2027025\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 2027025\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 2027025\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 2027025\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(2\*a) - (2027025\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 2027025\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + 2027025\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - 2027025\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(2\*a)) \* erf(sqrt(-2\*I\*b)\*x^(1/6))) \* sqrt(abs(b)) + 12480\*(256\*b^5\*x^(11/6)\*abs(b) - 1584\*b^3\*x^(7/6)\*abs(b) + 3465\*b\*sqrt(x)\*abs(b)) \* cos(2\*b\*x^(1/3) + 2\*a) + 240\*(4096\*b^6\*x^(13/6)\*abs(b) - 36608\*b^4\*x^(3/2)\*abs(b) + 144144\*b^2\*x^(5/6)\*abs(b) - 135135\*x^(1/6)\*abs(b)) \* sin(2\*b\*x^(1/3) + 2\*a))/(b^7\*abs(b))

**Fricas [A]** time = 2.25505, size = 591, normalized size = 1.91

$$399360 b^6 x^{\frac{11}{6}} - 2471040 b^4 x^{\frac{7}{6}} - 2027025 \pi \sqrt{\frac{b}{\pi}} \cos(2a) S\left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) - 2027025 \pi \sqrt{\frac{b}{\pi}} C\left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) \sin(2a) - 3120 \left(256 b^6 x^{\frac{11}{6}} - 1584 b^4 x^{\frac{7}{6}} + 3465 b^2 \sqrt{x}\right) \cos(b x^{\frac{1}{3}} + a)^2 + 60 \left(36608 b^5 x^{\frac{3}{2}} - 144144 b^3 x^{\frac{5}{6}} - (4096 b^7 x^2 - 135135 b) x^{\frac{1}{6}}\right) \cos(b x^{\frac{1}{3}} + a) \sin(b x^{\frac{1}{3}} + a) - 8 \left(4096 b^8 x^2 - 675675 b^2\right) \sqrt{x} / b^8$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(a+b\*x^(1/3))^2,x, algorithm="fricas")

[Out] -1/163840\*(399360\*b^6\*x^(11/6) - 2471040\*b^4\*x^(7/6) - 2027025\*pi\*sqrt(b/pi)\*cos(2\*a)\*fresnel\_sin(2\*x^(1/6)\*sqrt(b/pi)) - 2027025\*pi\*sqrt(b/pi)\*fresnel\_cos(2\*x^(1/6)\*sqrt(b/pi))\*sin(2\*a) - 3120\*(256\*b^6\*x^(11/6) - 1584\*b^4\*x^(7/6) + 3465\*b^2\*sqrt(x))\*cos(b\*x^(1/3) + a)^2 + 60\*(36608\*b^5\*x^(3/2) - 144144\*b^3\*x^(5/6) - (4096\*b^7\*x^2 - 135135\*b)\*x^(1/6))\*cos(b\*x^(1/3) + a)\*sin(b\*x^(1/3) + a) - 8\*(4096\*b^8\*x^2 - 675675\*b^2)\*sqrt(x))/b^8

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{\frac{3}{2}} \cos^2(a + b\sqrt[3]{x}) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*(3/2)\*cos(a+b\*x\*\*(1/3))\*\*2,x)

[Out] Integral(x\*\*(3/2)\*cos(a + b\*x\*\*(1/3))\*\*2, x)

**Giac [C]** time = 1.18605, size = 302, normalized size = 0.97

$$\frac{1}{5}x^{\frac{5}{2}} - \frac{3\left(4096ib^6x^{\frac{13}{6}} - 13312b^5x^{\frac{11}{6}} - 36608ib^4x^{\frac{9}{2}} + 82368b^3x^{\frac{7}{6}} + 144144ib^2x^{\frac{5}{6}} - 180180b\sqrt{x} - 135135ix^{\frac{1}{6}}\right)e^{2ibx^{\frac{1}{3}}}}{32768b^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(3/2)\*cos(a+b\*x^(1/3))^2,x, algorithm="giac")

[Out]  $\frac{1}{5}x^{5/2} - \frac{3}{32768}(4096Ib^6x^{13/6} - 13312b^5x^{11/6} - 36608Ib^4x^{9/2} + 82368b^3x^{7/6} + 144144Ib^2x^{5/6} - 180180b\sqrt{x} - 135135Ix^{1/6})e^{2Ibx^{1/3}}/b^7 - \frac{3}{32768}(-4096Ib^6x^{13/6} - 13312b^5x^{11/6} + 36608Ib^4x^{9/2} + 82368b^3x^{7/6} - 144144Ib^2x^{5/6} - 180180b\sqrt{x} + 135135Ix^{1/6})e^{-2Ibx^{1/3}} - \frac{2Ia}{b^7} + \frac{405405}{65536}I\sqrt{\pi}\operatorname{erf}(-\sqrt{b}x^{1/6})(-Ib/\operatorname{abs}(b) + 1)e^{2Ia}/(b^{15/2})(-Ib/\operatorname{abs}(b) + 1)) - \frac{405405}{65536}I\sqrt{\pi}\operatorname{erf}(-\sqrt{b}x^{1/6})(Ib/\operatorname{abs}(b) + 1)e^{-2Ia}/(b^{15/2})(Ib/\operatorname{abs}(b) + 1))$

### 3.56 $\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$

**Optimal.** Leaf size=218

$$\frac{315\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{512b^{9/2}} - \frac{315\sqrt{\pi} \sin(2a) S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{512b^{9/2}} + \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4} - \frac{105\sqrt{x}}{128b^4}$$

[Out] (315\*x^(1/6))/(256\*b^4) - (21\*x^(5/6))/(16\*b^2) + x^(3/2)/3 - (315\*x^(1/6)\*Cos[a + b\*x^(1/3)]^2)/(128\*b^4) + (21\*x^(5/6)\*Cos[a + b\*x^(1/3)]^2)/(8\*b^2) + (315\*Sqrt[Pi]\*Cos[2\*a]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]])/(512\*b^(9/2)) - (315\*Sqrt[Pi]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a])/(512\*b^(9/2)) - (105\*Sqrt[x]\*Cos[a + b\*x^(1/3)]\*Sin[a + b\*x^(1/3)])/(32\*b^3) + (3\*x^(7/6)\*Cos[a + b\*x^(1/3)]\*Sin[a + b\*x^(1/3)])/(2\*b)

**Rubi [A]** time = 0.250679, antiderivative size = 218, normalized size of antiderivative = 1., number of steps used = 12, number of rules used = 9, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.5$ , Rules used = {3416, 3311, 30, 3312, 3306, 3305, 3351, 3304, 3352}

$$\frac{315\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{512b^{9/2}} - \frac{315\sqrt{\pi} \sin(2a) S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{512b^{9/2}} + \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4} - \frac{105\sqrt{x}}{128b^4}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[x]\*Cos[a + b\*x^(1/3)]^2,x]

[Out] (315\*x^(1/6))/(256\*b^4) - (21\*x^(5/6))/(16\*b^2) + x^(3/2)/3 - (315\*x^(1/6)\*Cos[a + b\*x^(1/3)]^2)/(128\*b^4) + (21\*x^(5/6)\*Cos[a + b\*x^(1/3)]^2)/(8\*b^2) + (315\*Sqrt[Pi]\*Cos[2\*a]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]])/(512\*b^(9/2)) - (315\*Sqrt[Pi]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a])/(512\*b^(9/2)) - (105\*Sqrt[x]\*Cos[a + b\*x^(1/3)]\*Sin[a + b\*x^(1/3)])/(32\*b^3) + (3\*x^(7/6)\*Cos[a + b\*x^(1/3)]\*Sin[a + b\*x^(1/3)])/(2\*b)

#### Rule 3416

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*(x\_)^(m\_.), x\_Symbol] := Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + d\*x^(k\*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]

#### Rule 3311

Int[((c\_.) + (d\_.)\*(x\_)^(m\_.))\*((b\_.)\*sin[(e\_.) + (f\_.)\*(x\_)])^(n\_.), x\_Symbol] := Simp[(d\*m\*(c + d\*x)^(m - 1)\*(b\*Ssin[e + f\*x])^n)/(f^2\*n^2), x] + (Dist[(b^2\*(n - 1))/n, Int[(c + d\*x)^m\*(b\*Ssin[e + f\*x])^(n - 2), x], x] - Dist[(d^2\*m\*(m - 1))/(f^2\*n^2), Int[(c + d\*x)^(m - 2)\*(b\*Ssin[e + f\*x])^n, x], x] - Simp[(b\*(c + d\*x)^m\*Cos[e + f\*x]\*(b\*Ssin[e + f\*x])^(n - 1))/(f\*n), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]

#### Rule 30

Int[(x\_)^(m\_.), x\_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

#### Rule 3312

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

#### Rule 3306

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[Cos
[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] + Dist[Sin[(d
*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d,
e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]
```

#### Rule 3305

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d
, Subst[Int[Sin[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}
, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

#### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*Fresne
lS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

#### Rule 3304

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := D
ist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d
, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

#### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*Fresne
lC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

#### Rubi steps

$$\begin{aligned}
\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx &= 3 \operatorname{Subst} \left( \int x^{7/2} \cos^2(a + bx) dx, x, \sqrt[3]{x} \right) \\
&= \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{3x^{7/6} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{2b} + \frac{3}{2} \operatorname{Subst} \left( \int x^{7/2} dx, x, \sqrt[3]{x} \right) \\
&= \frac{x^{3/2}}{3} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{105\sqrt{x} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{32b^3} \\
&= -\frac{21x^{5/6}}{16b^2} + \frac{x^{3/2}}{3} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{105\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^3} \\
&= \frac{315\sqrt[6]{x}}{256b^4} - \frac{21x^{5/6}}{16b^2} + \frac{x^{3/2}}{3} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{105\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^3} \\
&= \frac{315\sqrt[6]{x}}{256b^4} - \frac{21x^{5/6}}{16b^2} + \frac{x^{3/2}}{3} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{105\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^3} \\
&= \frac{315\sqrt[6]{x}}{256b^4} - \frac{21x^{5/6}}{16b^2} + \frac{x^{3/2}}{3} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \frac{105\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^3} \\
&= \frac{315\sqrt[6]{x}}{256b^4} - \frac{21x^{5/6}}{16b^2} + \frac{x^{3/2}}{3} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4} + \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{315\sqrt{\pi} \cos(a + b\sqrt[3]{x})}{512b^3}
\end{aligned}$$

**Mathematica [A]** time = 0.442387, size = 148, normalized size = 0.68

$$\frac{2\sqrt{b}\sqrt[6]{x}\left(4b\sqrt[3]{x}\left(9\left(16b^2x^{2/3}-35\right)\sin\left(2\left(a+b\sqrt[3]{x}\right)\right)+64b^3x\right)+63\left(16b^2x^{2/3}-15\right)\cos\left(2\left(a+b\sqrt[3]{x}\right)\right)\right)+945\sqrt{\pi}\cos(2a)\operatorname{FresnelC}\left(\frac{2\sqrt{b}x^{1/6}}{\sqrt{\pi}}\right)}{1536b^{9/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[x]\*Cos[a + b\*x^(1/3)]^2,x]

[Out] (945\*Sqrt[Pi]\*Cos[2\*a]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]] - 945\*Sqrt[Pi]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a] + 2\*Sqrt[b]\*x^(1/6)\*(63\*(-15 + 16\*b^2\*x^(2/3))\*Cos[2\*(a + b\*x^(1/3))] + 4\*b\*x^(1/3)\*(64\*b^3\*x + 9\*(-35 + 16\*b^2\*x^(2/3))\*Sin[2\*(a + b\*x^(1/3))])))/(1536\*b^(9/2))

**Maple [A]** time = 0.037, size = 145, normalized size = 0.7

$$\frac{1}{3}x^{\frac{3}{2}} + \frac{3}{4b}x^{\frac{7}{6}}\sin(2a + 2b\sqrt[3]{x}) - \frac{21}{4b}\left(-\frac{1}{4b}x^{\frac{5}{6}}\cos(2a + 2b\sqrt[3]{x}) + \frac{5}{4b}\left(\frac{1}{4b}\sqrt{x}\sin(2a + 2b\sqrt[3]{x}) - \frac{3}{4b}\left(-\frac{1}{4b}\sqrt[6]{x}\cos(2a + 2b\sqrt[3]{x})\right)\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(1/2)\*cos(a+b\*x^(1/3))^2,x)

[Out] 1/3\*x^(3/2)+3/4/b\*x^(7/6)\*sin(2\*a+2\*b\*x^(1/3))-21/4/b\*(-1/4/b\*x^(5/6)\*cos(2\*a+2\*b\*x^(1/3))+5/4/b\*(1/4/b\*x^(1/2)\*sin(2\*a+2\*b\*x^(1/3))-3/4/b\*(-1/4/b\*x^(1/6)\*cos(2\*a+2\*b\*x^(1/3))+1/8/b^(3/2)\*Pi^(1/2)\*(cos(2\*a)\*FresnelC(2\*x^(1/6)\*b^(1/2)/Pi^(1/2))-sin(2\*a)\*FresnelS(2\*x^(1/6)\*b^(1/2)/Pi^(1/2))))

**Maxima [C]** time = 1.96806, size = 455, normalized size = 2.09

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(1/2)\*cos(a+b\*x^(1/3))^2,x, algorithm="maxima")

[Out] 1/12288\*(4096\*b^4\*x^(3/2)\*abs(b) + sqrt(2)\*sqrt(pi)\*(((945\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 945\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 945\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 945\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*cos(2\*a) - (945\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 945\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + 945\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - 945\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*sin(2\*a))\*erf(sqrt(2\*I\*b)\*x^(1/6)) + ((945\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 945\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + 945\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - 945\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*cos(2\*a) - (-945\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - 945\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + 945\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - 945\*sin(-1/4\*pi + 1/2\*arctan2(0, b))))\*sin(2\*a))\*erf(sqrt(-2\*I\*b)\*x^(1/6))\*sqrt(abs(b)) + 1008\*(16\*b^2\*x^(5/6)\*abs(b) - 15\*x^(1/6)\*abs(b))\*cos(2\*b\*x^(1/3) + 2\*a) + 576\*(16\*b^3\*x^(7/6)\*abs(b) - 35\*b\*sqrt(x)\*abs(b))\*sin(2\*b\*x^(1/3) + 2\*a))/(b^4\*abs(b))

**Fricas [A]** time = 2.08488, size = 444, normalized size = 2.04

$$\frac{512 b^5 x^{\frac{3}{2}} - 2016 b^3 x^{\frac{5}{6}} + 945 \pi \sqrt{\frac{b}{\pi}} \cos(2a) C\left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) - 945 \pi \sqrt{\frac{b}{\pi}} S\left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) \sin(2a) + 252 \left(16 b^3 x^{\frac{5}{6}} - 15 b x^{\frac{1}{6}}\right) \cos(a)}{1536 b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(1/2)\*cos(a+b\*x^(1/3))^2,x, algorithm="fricas")

[Out] 1/1536\*(512\*b^5\*x^(3/2) - 2016\*b^3\*x^(5/6) + 945\*pi\*sqrt(b/pi)\*cos(2\*a)\*fresnel\_cos(2\*x^(1/6)\*sqrt(b/pi)) - 945\*pi\*sqrt(b/pi)\*fresnel\_sin(2\*x^(1/6)\*sqrt(b/pi))\*sin(2\*a) + 252\*(16\*b^3\*x^(5/6) - 15\*b\*x^(1/6))\*cos(b\*x^(1/3) + a)^2 + 144\*(16\*b^4\*x^(7/6) - 35\*b^2\*sqrt(x))\*cos(b\*x^(1/3) + a)\*sin(b\*x^(1/3) + a) + 1890\*b\*x^(1/6))/b^5

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*(1/2)\*cos(a+b\*x\*\*(1/3))\*\*2,x)

[Out] Timed out

**Giac [C]** time = 1.17979, size = 238, normalized size = 1.09

$$\frac{1}{3} x^{\frac{3}{2}} - \frac{3 \left( 64i b^3 x^{\frac{7}{6}} - 112 b^2 x^{\frac{5}{6}} - 140i b \sqrt{x} + 105 x^{\frac{1}{6}} \right) e^{\left( 2i b x^{\frac{1}{3}} + 2i a \right)}}{512 b^4} - \frac{3 \left( -64i b^3 x^{\frac{7}{6}} - 112 b^2 x^{\frac{5}{6}} + 140i b \sqrt{x} + 105 x^{\frac{1}{6}} \right) e^{\left( -2i b x^{\frac{1}{3}} - 2i a \right)}}{512 b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(1/2)\*cos(a+b\*x^(1/3))^2,x, algorithm="giac")

[Out] 1/3\*x^(3/2) - 3/512\*(64\*I\*b^3\*x^(7/6) - 112\*b^2\*x^(5/6) - 140\*I\*b\*sqrt(x) + 105\*x^(1/6))\*e^(2\*I\*b\*x^(1/3) + 2\*I\*a)/b^4 - 3/512\*(-64\*I\*b^3\*x^(7/6) - 112\*b^2\*x^(5/6) + 140\*I\*b\*sqrt(x) + 105\*x^(1/6))\*e^(-2\*I\*b\*x^(1/3) - 2\*I\*a)/b^4 - 315/1024\*sqrt(pi)\*erf(-sqrt(b)\*x^(1/6)\*(-I\*b/abs(b) + 1))\*e^(2\*I\*a)/(b^(9/2)\*(-I\*b/abs(b) + 1)) - 315/1024\*sqrt(pi)\*erf(-sqrt(b)\*x^(1/6)\*(I\*b/abs(b) + 1))\*e^(-2\*I\*a)/(b^(9/2)\*(I\*b/abs(b) + 1))

$$3.57 \quad \int \frac{\cos^2(a+b\sqrt[3]{x})}{\sqrt{x}} dx$$

**Optimal.** Leaf size=102

$$-\frac{3\sqrt{\pi} \sin(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{8b^{3/2}} - \frac{3\sqrt{\pi} \cos(2a) S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{8b^{3/2}} + \frac{3\sqrt[6]{x} \sin\left(2\left(a+b\sqrt[3]{x}\right)\right)}{4b} + \sqrt{x}$$

[Out] Sqrt[x] - (3\*Sqrt[Pi]\*Cos[2\*a]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]])/(8\*b^(3/2)) - (3\*Sqrt[Pi]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a])/(8\*b^(3/2)) + (3\*x^(1/6)\*Sin[2\*(a + b\*x^(1/3))])/(4\*b)

**Rubi [A]** time = 0.169637, antiderivative size = 102, normalized size of antiderivative = 1., number of steps used = 9, number of rules used = 8, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$ , Rules used = {3416, 3312, 3296, 3306, 3305, 3351, 3304, 3352}

$$-\frac{3\sqrt{\pi} \sin(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{8b^{3/2}} - \frac{3\sqrt{\pi} \cos(2a) S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{8b^{3/2}} + \frac{3\sqrt[6]{x} \sin\left(2\left(a+b\sqrt[3]{x}\right)\right)}{4b} + \sqrt{x}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^(1/3)]^2/Sqrt[x], x]

[Out] Sqrt[x] - (3\*Sqrt[Pi]\*Cos[2\*a]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]])/(8\*b^(3/2)) - (3\*Sqrt[Pi]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a])/(8\*b^(3/2)) + (3\*x^(1/6)\*Sin[2\*(a + b\*x^(1/3))])/(4\*b)

#### Rule 3416

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*(x\_)^(m\_.), x\_Symbol] := Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + d\*x^(k\*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]

#### Rule 3312

Int[((c\_.) + (d\_.)\*(x\_)^(m\_))\*sin[(e\_.) + (f\_.)\*(x\_)^(n\_)], x\_Symbol] := Int[ExpandTrigReduce[(c + d\*x)^m, Sin[e + f\*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

#### Rule 3296

Int[((c\_.) + (d\_.)\*(x\_)^(m\_))\*sin[(e\_.) + (f\_.)\*(x\_)^(n\_)], x\_Symbol] := -Simp[(c + d\*x)^m\*Cos[e + f\*x]/f, x] + Dist[(d\*m)/f, Int[(c + d\*x)^(m - 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

#### Rule 3306

Int[sin[(e\_.) + (f\_.)\*(x\_)^(n\_)]/Sqrt[(c\_.) + (d\_.)\*(x\_)^(m\_)], x\_Symbol] := Dist[Cos[(d\*e - c\*f)/d], Int[Sin[(c\*f)/d + f\*x]/Sqrt[c + d\*x], x], x] + Dist[Sin[(d\*e - c\*f)/d], Int[Cos[(c\*f)/d + f\*x]/Sqrt[c + d\*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d\*e - c\*f, 0]

#### Rule 3305



```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d
, Subst[Int[Sin[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}
, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

Rule 3304

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx &= 3 \operatorname{Subst} \left( \int \sqrt{x} \cos^2(a + bx) dx, x, \sqrt[3]{x} \right) \\
&= 3 \operatorname{Subst} \left( \int \left( \frac{\sqrt{x}}{2} + \frac{1}{2} \sqrt{x} \cos(2a + 2bx) \right) dx, x, \sqrt[3]{x} \right) \\
&= \sqrt{x} + \frac{3}{2} \operatorname{Subst} \left( \int \sqrt{x} \cos(2a + 2bx) dx, x, \sqrt[3]{x} \right) \\
&= \sqrt{x} + \frac{3\sqrt[6]{x} \sin(2(a + b\sqrt[3]{x}))}{4b} - \frac{3 \operatorname{Subst} \left( \int \frac{\sin(2a + 2bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right)}{8b} \\
&= \sqrt{x} + \frac{3\sqrt[6]{x} \sin(2(a + b\sqrt[3]{x}))}{4b} - \frac{(3 \cos(2a)) \operatorname{Subst} \left( \int \frac{\sin(2bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right)}{8b} - \frac{(3 \sin(2a)) \operatorname{Subst} \left( \int \frac{1}{\sqrt{x}} dx, x, \sqrt[3]{x} \right)}{8b} \\
&= \sqrt{x} + \frac{3\sqrt[6]{x} \sin(2(a + b\sqrt[3]{x}))}{4b} - \frac{(3 \cos(2a)) \operatorname{Subst} \left( \int \sin(2bx^2) dx, x, \sqrt[6]{x} \right)}{4b} - \frac{(3 \sin(2a)) \operatorname{Subst} \left( \int \frac{1}{\sqrt{x}} dx, x, \sqrt[3]{x} \right)}{8b} \\
&= \sqrt{x} - \frac{3\sqrt{\pi} \cos(2a) S \left( \frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}} \right)}{8b^{3/2}} - \frac{3\sqrt{\pi} C \left( \frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}} \right) \sin(2a)}{8b^{3/2}} + \frac{3\sqrt[6]{x} \sin(2(a + b\sqrt[3]{x}))}{4b}
\end{aligned}$$

**Mathematica [A]** time = 0.189478, size = 103, normalized size = 1.01

$$\frac{-3\sqrt{\pi} \sin(2a) \operatorname{FresnelC} \left( \frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}} \right) - 3\sqrt{\pi} \cos(2a) S \left( \frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}} \right) + 2\sqrt{b}\sqrt[6]{x} (3 \sin(2(a + b\sqrt[3]{x})) + 4b\sqrt[3]{x})}{8b^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x^(1/3)]^2/Sqrt[x], x]
```

```
[Out] (-3*Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]] - 3*Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a] + 2*Sqrt[b]*x^(1/6)*(4*b*x^(1/3) + 3*Sin[2*(a + b*x^(1/3))]))/(8*b^(3/2))
```

**Maple [A]** time = 0.039, size = 67, normalized size = 0.7

$$\sqrt{x} + \frac{3}{4b} \sqrt[6]{x} \sin(2a + 2b\sqrt[3]{x}) - \frac{3\sqrt{\pi}}{8} \left( \cos(2a) \operatorname{FresnelS} \left( 2 \frac{\sqrt[6]{x}\sqrt{b}}{\sqrt{\pi}} \right) + \sin(2a) \operatorname{FresnelC} \left( 2 \frac{\sqrt[6]{x}\sqrt{b}}{\sqrt{\pi}} \right) \right) b^{-\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^(1/3))^2/x^(1/2),x)

[Out] x^(1/2)+3/4\*x^(1/6)\*sin(2\*a+2\*b\*x^(1/3))/b-3/8/b^(3/2)\*Pi^(1/2)\*(cos(2\*a)\*FresnelS(2\*x^(1/6)\*b^(1/2)/Pi^(1/2))+sin(2\*a)\*FresnelC(2\*x^(1/6)\*b^(1/2)/Pi^(1/2)))

**Maxima [C]** time = 2.96517, size = 392, normalized size = 3.84

$$\sqrt{2}\sqrt{\pi} \left( \left( -3i \cos\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) - 3i \cos\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) - 3 \sin\left(\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) + 3 \sin\left(-\frac{1}{4}\pi + \frac{1}{2}\arctan(0,b)\right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))^2/x^(1/2),x, algorithm="maxima")

[Out] 1/64\*(sqrt(2)\*sqrt(pi)\*(((3\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) - 3\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 3\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(2\*a) - (3\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 3\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(2\*a)) \* erf(sqrt(2\*I\*b)\*x^(1/6)) + ((3\*I\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*I\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) - 3\*sin(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* cos(2\*a) - (3\*cos(1/4\*pi + 1/2\*arctan2(0, b)) + 3\*cos(-1/4\*pi + 1/2\*arctan2(0, b)) + 3\*I\*sin(1/4\*pi + 1/2\*arctan2(0, b)) - 3\*I\*sin(-1/4\*pi + 1/2\*arctan2(0, b))) \* sin(2\*a)) \* erf(sqrt(-2\*I\*b)\*x^(1/6))) \* sqrt(abs(b)) + 64\*b\*sqrt(x)\*abs(b) + 48\*x^(1/6)\*abs(b)\*sin(2\*b\*x^(1/3) + 2\*a))/(b\*abs(b))

**Fricas [A]** time = 2.08258, size = 271, normalized size = 2.66

$$\frac{3\pi\sqrt{\frac{b}{\pi}}\cos(2a)\operatorname{S}\left(2x^{\frac{1}{6}}\sqrt{\frac{b}{\pi}}\right)+3\pi\sqrt{\frac{b}{\pi}}\operatorname{C}\left(2x^{\frac{1}{6}}\sqrt{\frac{b}{\pi}}\right)\sin(2a)-12bx^{\frac{1}{6}}\cos\left(bx^{\frac{1}{3}}+a\right)\sin\left(bx^{\frac{1}{3}}+a\right)-8b^2\sqrt{x}}{8b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))^2/x^(1/2),x, algorithm="fricas")

[Out] -1/8\*(3\*pi\*sqrt(b/pi)\*cos(2\*a)\*fresnel\_sin(2\*x^(1/6)\*sqrt(b/pi)) + 3\*pi\*sqrt(b/pi)\*fresnel\_cos(2\*x^(1/6)\*sqrt(b/pi))\*sin(2\*a) - 12\*b\*x^(1/6)\*cos(b\*x^(1/3) + a)\*sin(b\*x^(1/3) + a) - 8\*b^2\*sqrt(x))/b^2

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x\*\*(1/3))\*\*2/x\*\*(1/2), x)

[Out] Integral(cos(a + b\*x\*\*(1/3))\*\*2/sqrt(x), x)

**Giac [C]** time = 1.17301, size = 167, normalized size = 1.64

$$\sqrt{x} - \frac{3ix^{\frac{1}{6}}e^{\left(2ibx^{\frac{1}{3}}+2ia\right)}}{8b} + \frac{3ix^{\frac{1}{6}}e^{\left(-2ibx^{\frac{1}{3}}-2ia\right)}}{8b} - \frac{3i\sqrt{\pi}\operatorname{erf}\left(-\sqrt{bx^{\frac{1}{6}}}\left(-\frac{ib}{|b|}+1\right)\right)e^{(2ia)}}{16b^{\frac{3}{2}}\left(-\frac{ib}{|b|}+1\right)} + \frac{3i\sqrt{\pi}\operatorname{erf}\left(-\sqrt{bx^{\frac{1}{6}}}\left(\frac{ib}{|b|}+1\right)\right)e^{(-2ia)}}{16b^{\frac{3}{2}}\left(\frac{ib}{|b|}+1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))^2/x^(1/2), x, algorithm="giac")

[Out] sqrt(x) - 3/8\*I\*x^(1/6)\*e^(2\*I\*b\*x^(1/3) + 2\*I\*a)/b + 3/8\*I\*x^(1/6)\*e^(-2\*I\*b\*x^(1/3) - 2\*I\*a)/b - 3/16\*I\*sqrt(pi)\*erf(-sqrt(b)\*x^(1/6)\*(-I\*b/abs(b) + 1))\*e^(2\*I\*a)/(b^(3/2)\*(-I\*b/abs(b) + 1)) + 3/16\*I\*sqrt(pi)\*erf(-sqrt(b)\*x^(1/6)\*(I\*b/abs(b) + 1))\*e^(-2\*I\*a)/(b^(3/2)\*(I\*b/abs(b) + 1))

$$3.58 \quad \int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{3/2}} dx$$

**Optimal.** Leaf size=116

$$-8\sqrt{\pi}b^{3/2}\cos(2a)\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)+8\sqrt{\pi}b^{3/2}\sin(2a)\text{S}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)-\frac{2\cos^2(a+b\sqrt[3]{x})}{\sqrt{x}}+\frac{8b\sin(a+b\sqrt[3]{x})\cos(a+b\sqrt[3]{x})}{\sqrt[6]{x}}$$

[Out] (-2\*Cos[a + b\*x^(1/3)]^2)/Sqrt[x] - 8\*b^(3/2)\*Sqrt[Pi]\*Cos[2\*a]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]] + 8\*b^(3/2)\*Sqrt[Pi]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a] + (8\*b\*Cos[a + b\*x^(1/3)]\*Sin[a + b\*x^(1/3)])/x^(1/6)

**Rubi [A]** time = 0.18694, antiderivative size = 116, normalized size of antiderivative = 1., number of steps used = 10, number of rules used = 9, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.5$ , Rules used = {3416, 3314, 30, 3312, 3306, 3305, 3351, 3304, 3352}

$$-8\sqrt{\pi}b^{3/2}\cos(2a)\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)+8\sqrt{\pi}b^{3/2}\sin(2a)\text{S}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)-\frac{2\cos^2(a+b\sqrt[3]{x})}{\sqrt{x}}+\frac{8b\sin(a+b\sqrt[3]{x})\cos(a+b\sqrt[3]{x})}{\sqrt[6]{x}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^(1/3)]^2/x^(3/2), x]

[Out] (-2\*Cos[a + b\*x^(1/3)]^2)/Sqrt[x] - 8\*b^(3/2)\*Sqrt[Pi]\*Cos[2\*a]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]] + 8\*b^(3/2)\*Sqrt[Pi]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a] + (8\*b\*Cos[a + b\*x^(1/3)]\*Sin[a + b\*x^(1/3)])/x^(1/6)

#### Rule 3416

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_.)\*(x\_)^(m\_.), x\_Symbol] := Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k\*(m + 1) - 1)\*(a + b\*Cos[c + d\*x^(k\*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]

#### Rule 3314

Int[((c\_.) + (d\_.)\*(x\_)^(m\_))\*((b\_.)\*sin[(e\_.) + (f\_.)\*(x\_)])^(n\_), x\_Symbol] := Simp[((c + d\*x)^(m + 1)\*(b\*Ssin[e + f\*x])^n)/(d\*(m + 1)), x] + (Dist[(b^2\*f^2\*n\*(n - 1))/(d^2\*(m + 1)\*(m + 2)), Int[(c + d\*x)^(m + 2)\*(b\*Ssin[e + f\*x])^(n - 2), x], x] - Dist[(f^2\*n^2)/(d^2\*(m + 1)\*(m + 2)), Int[(c + d\*x)^(m + 2)\*(b\*Ssin[e + f\*x])^n, x], x] - Simp[(b\*f\*n\*(c + d\*x)^(m + 2)\*Cos[e + f\*x]\*(b\*Ssin[e + f\*x])^(n - 1))/(d^2\*(m + 1)\*(m + 2)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && LtQ[m, -2]

#### Rule 30

Int[(x\_)^(m\_.), x\_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

#### Rule 3312

Int[((c\_.) + (d\_.)\*(x\_)^(m\_))\*sin[(e\_.) + (f\_.)\*(x\_)])^(n\_), x\_Symbol] := Int[ExpandTrigReduce[(c + d\*x)^m, Sin[e + f\*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

#### Rule 3306

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[Cos
[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] + Dist[Sin[(d
*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d,
e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]
```

### Rule 3305

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d
, Subst[Int[Sin[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}
, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

### Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*Fresne
lS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rule 3304

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := D
ist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d
, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*Fresne
lC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

### Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx &= 3 \operatorname{Subst} \left( \int \frac{\cos^2(a + bx)}{x^{5/2}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} + (8b^2) \operatorname{Subst} \left( \int \frac{1}{\sqrt{x}} dx, x, \sqrt[3]{x} \right) - (16b^2) \operatorname{Subst} \left( \int \left( \frac{1}{2\sqrt{x}} + \frac{c}{\sqrt{x}} \right) dx, x, \sqrt[3]{x} \right) \\
&= 16b^2 \sqrt[6]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} - (16b^2) \operatorname{Subst} \left( \int \left( \frac{1}{2\sqrt{x}} + \frac{c}{\sqrt{x}} \right) dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} - (8b^2) \operatorname{Subst} \left( \int \frac{\cos(2a + 2bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} - (8b^2 \cos(2a)) \operatorname{Subst} \left( \int \frac{\cos(2bx)}{\sqrt{x}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} - (16b^2 \cos(2a)) \operatorname{Subst} \left( \int \cos(2bx^2) dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} - 8b^{3/2} \sqrt{\pi} \cos(2a) C \left( \frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}} \right) + 8b^{3/2} \sqrt{\pi} S \left( \frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}} \right) \sin(2a) + \frac{8b \cos(a)}{\sqrt{x}}
\end{aligned}$$

**Mathematica [A]** time = 0.254318, size = 116, normalized size = 1.

$$\frac{8\sqrt{\pi}b^{3/2}\sqrt{x}\sin(2a)S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) + 4b\sqrt[3]{x}\sin\left(2\left(a + b\sqrt[3]{x}\right)\right) - \cos\left(2\left(a + b\sqrt[3]{x}\right)\right) - 1}{\sqrt{x}} - 8\sqrt{\pi}b^{3/2}\cos(2a)\operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x^(1/3)]^2/x^(3/2), x]
```

[Out]  $-8*b^{(3/2)}*\text{Sqrt}[Pi]*\text{Cos}[2*a]*\text{FresnelC}[(2*\text{Sqrt}[b]*x^{(1/6)})/\text{Sqrt}[Pi]] + (-1 - \text{Cos}[2*(a + b*x^{(1/3)})] + 8*b^{(3/2)}*\text{Sqrt}[Pi]*\text{Sqrt}[x]*\text{FresnelS}[(2*\text{Sqrt}[b]*x^{(1/6)})/\text{Sqrt}[Pi]]*\text{Sin}[2*a] + 4*b*x^{(1/3)}*\text{Sin}[2*(a + b*x^{(1/3)})])/\text{Sqrt}[x]$

**Maple [A]** time = 0.037, size = 87, normalized size = 0.8

$$-\frac{1}{\sqrt{x}} - \cos(2a + 2b\sqrt[3]{x}) \frac{1}{\sqrt{x}} - 4b \left( -\frac{\sin(2a + 2b\sqrt[3]{x})}{\sqrt[6]{x}} + 2\sqrt{b}\sqrt{\pi} \left( \cos(2a) \text{FresnelC} \left( 2 \frac{\sqrt[6]{x}\sqrt{b}}{\sqrt{\pi}} \right) - \sin(2a) \text{FresnelS} \left( 2 \frac{\sqrt[6]{x}\sqrt{b}}{\sqrt{\pi}} \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a+b*x^(1/3))^2/x^(3/2),x)`

[Out]  $-1/x^{(1/2)} - 1/x^{(1/2)}*\text{cos}(2*a + 2*b*x^{(1/3)}) - 4*b*(-1/x^{(1/6)}*\text{sin}(2*a + 2*b*x^{(1/3)}) + 2*b^{(1/2)}*Pi^{(1/2)}*(\text{cos}(2*a)*\text{FresnelC}(2*x^{(1/6)}*b^{(1/2)}/Pi^{(1/2)}) - \text{sin}(2*a)*\text{FresnelS}(2*x^{(1/6)}*b^{(1/2)}/Pi^{(1/2)})))$

**Maxima [C]** time = 1.74449, size = 379, normalized size = 3.27

$$\sqrt{2} \left( \left( 3 \left( \Gamma \left( -\frac{3}{2}, 2i b x^{\frac{1}{3}} \right) + \Gamma \left( -\frac{3}{2}, -2i b x^{\frac{1}{3}} \right) \right) \cos \left( \frac{3}{4} \pi + \frac{3}{2} \arctan(0, b) \right) + 3 \left( \Gamma \left( -\frac{3}{2}, 2i b x^{\frac{1}{3}} \right) + \Gamma \left( -\frac{3}{2}, -2i b x^{\frac{1}{3}} \right) \right) \cos \left( -\frac{3}{4} \pi \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^(1/3))^2/x^(3/2),x, algorithm="maxima")`

[Out]  $-1/4*(\text{sqrt}(2)*((3*(\text{gamma}(-3/2, 2*I*b*x^{(1/3)})) + \text{gamma}(-3/2, -2*I*b*x^{(1/3)})) * \text{cos}(3/4*\text{pi} + 3/2*\text{arctan2}(0, b)) + 3*(\text{gamma}(-3/2, 2*I*b*x^{(1/3)})) + \text{gamma}(-3/2, -2*I*b*x^{(1/3)})) * \text{cos}(-3/4*\text{pi} + 3/2*\text{arctan2}(0, b)) + (3*I*\text{gamma}(-3/2, 2*I*b*x^{(1/3)}) - 3*I*\text{gamma}(-3/2, -2*I*b*x^{(1/3)})) * \text{sin}(3/4*\text{pi} + 3/2*\text{arctan2}(0, b)) + (-3*I*\text{gamma}(-3/2, 2*I*b*x^{(1/3)}) + 3*I*\text{gamma}(-3/2, -2*I*b*x^{(1/3)})) * \text{sin}(-3/4*\text{pi} + 3/2*\text{arctan2}(0, b))) * \text{cos}(2*a) + ((-3*I*\text{gamma}(-3/2, 2*I*b*x^{(1/3)}) + 3*I*\text{gamma}(-3/2, -2*I*b*x^{(1/3)})) * \text{cos}(3/4*\text{pi} + 3/2*\text{arctan2}(0, b)) + (-3*I*\text{gamma}(-3/2, 2*I*b*x^{(1/3)}) + 3*I*\text{gamma}(-3/2, -2*I*b*x^{(1/3)})) * \text{cos}(-3/4*\text{pi} + 3/2*\text{arctan2}(0, b)) + 3*(\text{gamma}(-3/2, 2*I*b*x^{(1/3)})) + \text{gamma}(-3/2, -2*I*b*x^{(1/3)})) * \text{sin}(3/4*\text{pi} + 3/2*\text{arctan2}(0, b)) - 3*(\text{gamma}(-3/2, 2*I*b*x^{(1/3)}) + \text{gamma}(-3/2, -2*I*b*x^{(1/3)})) * \text{sin}(-3/4*\text{pi} + 3/2*\text{arctan2}(0, b))) * \text{sin}(2*a) * \text{sqrt}(x^{(1/3)}*\text{abs}(b)) * x^{(1/3)}*\text{abs}(b) + 4)/\text{sqrt}(x)$

**Fricas [A]** time = 2.09213, size = 296, normalized size = 2.55

$$2 \left( 4 \pi b x \sqrt{\frac{b}{\pi}} \cos(2a) C \left( 2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) - 4 \pi b x \sqrt{\frac{b}{\pi}} S \left( 2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) \sin(2a) - 4 b x^{\frac{5}{6}} \cos \left( b x^{\frac{1}{3}} + a \right) \sin \left( b x^{\frac{1}{3}} + a \right) + \sqrt{x} \cos \left( b x^{\frac{1}{3}} + a \right) \right) x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^(1/3))^2/x^(3/2),x, algorithm="fricas")`

[Out]  $-2*(4*\text{pi}*b*x*\text{sqrt}(b/\text{pi})*\text{cos}(2*a)*\text{fresnel\_cos}(2*x^{(1/6)}*\text{sqrt}(b/\text{pi})) - 4*\text{pi}*b*x*\text{sqrt}(b/\text{pi})*\text{fresnel\_sin}(2*x^{(1/6)}*\text{sqrt}(b/\text{pi}))*\text{sin}(2*a) - 4*b*x^{(5/6)}*\text{cos}(b*x^{(1/3)} + a) * \text{sin}(b*x^{(1/3)} + a) + \text{sqrt}(x) * \text{cos}(b*x^{(1/3)} + a))$

$b*x^{(1/3)} + a)*\sin(b*x^{(1/3)} + a) + \sqrt{x}*\cos(b*x^{(1/3)} + a)^2/x$

---

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x\*\*(1/3))\*\*2/x\*\*(3/2), x)

[Out] Integral(cos(a + b\*x\*\*(1/3))\*\*2/x\*\*(3/2), x)

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)^2}{x^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))^2/x^(3/2), x, algorithm="giac")

[Out] integrate(cos(b\*x^(1/3) + a)^2/x^(3/2), x)

$$3.59 \quad \int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{5/2}} dx$$

**Optimal.** Leaf size=228

$$-\frac{512}{315}\sqrt{\pi}b^{9/2}\sin(2a)\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)-\frac{512}{315}\sqrt{\pi}b^{9/2}\cos(2a)S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)+\frac{32b^2\cos^2(a+b\sqrt[3]{x})}{105x^{5/6}}-\frac{512b^4\cos^2(a+b\sqrt[3]{x})}{315\sqrt[6]{x}}$$

```
[Out] (-16*b^2)/(105*x^(5/6)) + (256*b^4)/(315*x^(1/6)) - (2*Cos[a + b*x^(1/3)]^2)/(3*x^(3/2)) + (32*b^2*Cos[a + b*x^(1/3)]^2)/(105*x^(5/6)) - (512*b^4*Cos[a + b*x^(1/3)]^2)/(315*x^(1/6)) - (512*b^(9/2)*Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]])/315 - (512*b^(9/2)*Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a])/315 + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(21*x^(7/6)) - (128*b^3*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(315*Sqrt[x])
```

**Rubi [A]** time = 0.250116, antiderivative size = 228, normalized size of antiderivative = 1., number of steps used = 12, number of rules used = 10, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$ , Rules used = {3416, 3314, 30, 3313, 12, 3306, 3305, 3351, 3304, 3352}

$$-\frac{512}{315}\sqrt{\pi}b^{9/2}\sin(2a)\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)-\frac{512}{315}\sqrt{\pi}b^{9/2}\cos(2a)S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)+\frac{32b^2\cos^2(a+b\sqrt[3]{x})}{105x^{5/6}}-\frac{512b^4\cos^2(a+b\sqrt[3]{x})}{315\sqrt[6]{x}}$$

Antiderivative was successfully verified.

```
[In] Int[Cos[a + b*x^(1/3)]^2/x^(5/2), x]
```

```
[Out] (-16*b^2)/(105*x^(5/6)) + (256*b^4)/(315*x^(1/6)) - (2*Cos[a + b*x^(1/3)]^2)/(3*x^(3/2)) + (32*b^2*Cos[a + b*x^(1/3)]^2)/(105*x^(5/6)) - (512*b^4*Cos[a + b*x^(1/3)]^2)/(315*x^(1/6)) - (512*b^(9/2)*Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]])/315 - (512*b^(9/2)*Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a])/315 + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(21*x^(7/6)) - (128*b^3*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(315*Sqrt[x])
```

#### Rule 3416

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
:> Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*x^(k*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x]
&& IntegerQ[p] && FractionQ[n]
```

#### Rule 3314

```
Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
:> Simp[((c + d*x)^(m + 1)*(b*Ssin[e + f*x])^n)/(d*(m + 1)), x] + (Dist[(b^2*f^2*n*(n - 1))/(d^2*(m + 1)*(m + 2)), Int[(c + d*x)^(m + 2)*(b*Ssin[e + f*x])^(n - 2), x], x] - Dist[(f^2*n^2)/(d^2*(m + 1)*(m + 2)), Int[(c + d*x)^(m + 2)*(b*Ssin[e + f*x])^n, x], x] - Simp[(b*f*n*(c + d*x)^(m + 2)*Cos[e + f*x]*(b*Ssin[e + f*x])^(n - 1))/(d^2*(m + 1)*(m + 2)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && LtQ[m, -2]
```

#### Rule 30

```
Int[(x_)^(m_.), x_Symbol]
:> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]
```



Rule 3313

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Si
mp[((c + d*x)^(m + 1)*Sin[e + f*x]^n)/(d*(m + 1)), x] - Dist[(f*n)/(d*(m +
1)), Int[ExpandTrigReduce[(c + d*x)^(m + 1), Cos[e + f*x]*Sin[e + f*x]^(n -
1), x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && GeQ[m, -2] &&
LtQ[m, -1]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 3306

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[Cos
[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] + Dist[Sin[(d
*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d,
e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]
```

Rule 3305

```
Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Dist[2/d
, Subst[Int[Sin[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}
, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

Rule 3351

```
Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*Fresne
lS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

Rule 3304

```
Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := D
ist[2/d, Subst[Int[Cos[(f*x^2)/d], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d
, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]
```

Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]*Fresne
lC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx &= 3 \operatorname{Subst} \left( \int \frac{\cos^2(a + bx)}{x^{11/2}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos^2(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{21x^{7/6}} + \frac{1}{21} (8b^2) \operatorname{Subst} \left( \int \frac{1}{x^{7/2}} dx, x, \sqrt[3]{x} \right) - \frac{1}{21} \\
&= -\frac{16b^2}{105x^{5/6}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{105x^{5/6}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{21x^{7/6}} - \frac{128b^3}{105x^{5/6}} \\
&= -\frac{16b^2}{105x^{5/6}} + \frac{256b^4}{315\sqrt[6]{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{21x^{7/6}} \\
&= -\frac{16b^2}{105x^{5/6}} + \frac{256b^4}{315\sqrt[6]{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{21x^{7/6}} \\
&= -\frac{16b^2}{105x^{5/6}} + \frac{256b^4}{315\sqrt[6]{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{21x^{7/6}} \\
&= -\frac{16b^2}{105x^{5/6}} + \frac{256b^4}{315\sqrt[6]{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{21x^{7/6}} \\
&= -\frac{16b^2}{105x^{5/6}} + \frac{256b^4}{315\sqrt[6]{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3x^{3/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{105x^{5/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{315\sqrt[6]{x}} - \frac{512b^3}{315}
\end{aligned}$$

**Mathematica [A]** time = 0.268309, size = 185, normalized size = 0.81

$$\frac{-512\sqrt{\pi}b^{9/2}x^{3/2} \sin(2a)\operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - 512\sqrt{\pi}b^{9/2}x^{3/2} \cos(2a)\operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - 256b^4x^{4/3} \cos(2(a + b\sqrt[3]{x})) + 48b^2x^{2/3} \cos(2(a + b\sqrt[3]{x}))}{315x^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^(1/3)]^2/x^(5/2), x]

[Out] (-105 - 105\*Cos[2\*(a + b\*x^(1/3))] + 48\*b^2\*x^(2/3)\*Cos[2\*(a + b\*x^(1/3))] - 256\*b^4\*x^(4/3)\*Cos[2\*(a + b\*x^(1/3))] - 512\*b^(9/2)\*Sqrt[Pi]\*x^(3/2)\*Cos[2\*a]\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]] - 512\*b^(9/2)\*Sqrt[Pi]\*x^(3/2)\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a] + 60\*b\*x^(1/3)\*Sin[2\*(a + b\*x^(1/3))] - 64\*b^3\*x\*Sin[2\*(a + b\*x^(1/3))])/(315\*x^(3/2))

**Maple [A]** time = 0.04, size = 146, normalized size = 0.6

$$-\frac{1}{3}x^{-\frac{3}{2}} - \frac{1}{3} \cos(2a + 2b\sqrt[3]{x})x^{-\frac{3}{2}} - \frac{4b}{3} \left( -\frac{1}{7} \sin(2a + 2b\sqrt[3]{x})x^{-\frac{7}{6}} + \frac{4b}{7} \left( -\frac{1}{5} \cos(2a + 2b\sqrt[3]{x})x^{-\frac{5}{6}} - \frac{4b}{5} \left( -\frac{1}{3} \sin(2a + 2b\sqrt[3]{x})x^{-\frac{1}{2}} + \frac{4b}{3} \cos(2a + 2b\sqrt[3]{x})x^{-\frac{1}{2}} \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^(1/3))^2/x^(5/2), x)

[Out] -1/3/x^(3/2)-1/3/x^(3/2)\*cos(2\*a+2\*b\*x^(1/3))-4/3\*b\*(-1/7/x^(7/6)\*sin(2\*a+2\*b\*x^(1/3))+4/7\*b\*(-1/5/x^(5/6)\*cos(2\*a+2\*b\*x^(1/3))-4/5\*b\*(-1/3/x^(1/2)\*sin(2\*a+2\*b\*x^(1/3))+4/3\*b\*(-1/x^(1/6)\*cos(2\*a+2\*b\*x^(1/3))-2\*b^(1/2)\*Pi^(1/2))\*(cos(2\*a)\*FresnelS(2\*x^(1/6)\*b^(1/2)/Pi^(1/2))+sin(2\*a)\*FresnelC(2\*x^(1/6)\*b^(1/2)/Pi^(1/2))))))

**Maxima [C]** time = 1.57209, size = 381, normalized size = 1.67

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))^2/x^(5/2),x, algorithm="maxima")

[Out] 
$$-1/3*(\sqrt{2}*((18*(\gamma(-9/2, 2*I*b*x^{1/3})) + \gamma(-9/2, -2*I*b*x^{1/3}))\cos(9/4*\pi + 9/2*\arctan2(0, b)) + 18*(\gamma(-9/2, 2*I*b*x^{1/3})) + \gamma(-9/2, -2*I*b*x^{1/3}))\cos(-9/4*\pi + 9/2*\arctan2(0, b)) + (18*I*\gamma(-9/2, 2*I*b*x^{1/3}) - 18*I*\gamma(-9/2, -2*I*b*x^{1/3}))\sin(9/4*\pi + 9/2*\arctan2(0, b)) + (-18*I*\gamma(-9/2, 2*I*b*x^{1/3}) + 18*I*\gamma(-9/2, -2*I*b*x^{1/3}))\sin(-9/4*\pi + 9/2*\arctan2(0, b)))\cos(2*a) + ((-18*I*\gamma(-9/2, 2*I*b*x^{1/3}) + 18*I*\gamma(-9/2, -2*I*b*x^{1/3}))\cos(9/4*\pi + 9/2*\arctan2(0, b)) + (-18*I*\gamma(-9/2, 2*I*b*x^{1/3}) + 18*I*\gamma(-9/2, -2*I*b*x^{1/3}))\cos(-9/4*\pi + 9/2*\arctan2(0, b)) + 18*(\gamma(-9/2, 2*I*b*x^{1/3})) + \gamma(-9/2, -2*I*b*x^{1/3}))\sin(9/4*\pi + 9/2*\arctan2(0, b)) - 18*(\gamma(-9/2, 2*I*b*x^{1/3}) + \gamma(-9/2, -2*I*b*x^{1/3}))\sin(-9/4*\pi + 9/2*\arctan2(0, b)))\sin(2*a))*\sqrt{x^{1/3}*abs(b)}*b^4*x^{4/3} + 1/x^{3/2}$$

**Fricas [A]** time = 2.18392, size = 455, normalized size = 2.

$$\frac{2\left(256\pi b^4 x^2 \sqrt{\frac{b}{\pi}} \cos(2a) S\left(2x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) + 256\pi b^4 x^2 \sqrt{\frac{b}{\pi}} C\left(2x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) \sin(2a) - 128b^4 x^{\frac{11}{6}} + 24b^2 x^{\frac{7}{6}} + \left(256b^4 x^{\frac{11}{6}} - \dots\right)\right)}{315x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))^2/x^(5/2),x, algorithm="fricas")

[Out] 
$$-2/315*(256*\pi*b^4*x^2*\sqrt{b/\pi}*\cos(2*a)*\text{fresnel\_sin}(2*x^{1/6}*\sqrt{b/\pi}) + 256*\pi*b^4*x^2*\sqrt{b/\pi}*\text{fresnel\_cos}(2*x^{1/6}*\sqrt{b/\pi}))*\sin(2*a) - 128*b^4*x^{11/6} + 24*b^2*x^{7/6} + (256*b^4*x^{11/6} - 48*b^2*x^{7/6} + 105*\sqrt{x})*\cos(b*x^{1/3} + a)^2 + 4*(16*b^3*x^{3/2} - 15*b*x^{5/6})*\cos(b*x^{1/3} + a)*\sin(b*x^{1/3} + a)/x^2$$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x\*\*(1/3))\*\*2/x\*\*(5/2),x)

[Out] Integral(cos(a + b\*x\*\*(1/3))\*\*2/x\*\*(5/2), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2\left(bx^{\frac{1}{3}} + a\right)}{x^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^(1/3))^2/x^(5/2),x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^(1/3) + a)^2/x^(5/2), x)
```

$$3.60 \quad \int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{7/2}} dx$$

**Optimal.** Leaf size=328

$$\frac{32768\sqrt{\pi}b^{15/2}\cos(2a)\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{675675} - \frac{32768\sqrt{\pi}b^{15/2}\sin(2a)S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{675675} - \frac{512b^4\cos^2(a+b\sqrt[3]{x})}{45045x^{7/6}} + \frac{32b^2\cos^2(a+b\sqrt[3]{x})}{715x^{11/6}}$$

```
[Out] (-16*b^2)/(715*x^(11/6)) + (256*b^4)/(45045*x^(7/6)) - (4096*b^6)/(675675*Sqrt[x]) - (2*Cos[a + b*x^(1/3)]^2)/(5*x^(5/2)) + (32*b^2*Cos[a + b*x^(1/3)]^2)/(715*x^(11/6)) - (512*b^4*Cos[a + b*x^(1/3)]^2)/(45045*x^(7/6)) + (8192*b^6*Cos[a + b*x^(1/3)]^2)/(675675*Sqrt[x]) + (32768*b^(15/2)*Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]])/675675 - (32768*b^(15/2)*Sqrt[Pi]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a])/675675 + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(65*x^(13/6)) - (128*b^3*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(6435*x^(3/2)) + (2048*b^5*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(225225*x^(5/6)) - (32768*b^7*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(675675*x^(1/6))
```

**Rubi [A]** time = 0.350694, antiderivative size = 328, normalized size of antiderivative = 1., number of steps used = 16, number of rules used = 9, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.5$ , Rules used = {3416, 3314, 30, 3312, 3306, 3305, 3351, 3304, 3352}

$$\frac{32768\sqrt{\pi}b^{15/2}\cos(2a)\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{675675} - \frac{32768\sqrt{\pi}b^{15/2}\sin(2a)S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{675675} - \frac{512b^4\cos^2(a+b\sqrt[3]{x})}{45045x^{7/6}} + \frac{32b^2\cos^2(a+b\sqrt[3]{x})}{715x^{11/6}}$$

Antiderivative was successfully verified.

```
[In] Int[Cos[a + b*x^(1/3)]^2/x^(7/2), x]
```

```
[Out] (-16*b^2)/(715*x^(11/6)) + (256*b^4)/(45045*x^(7/6)) - (4096*b^6)/(675675*Sqrt[x]) - (2*Cos[a + b*x^(1/3)]^2)/(5*x^(5/2)) + (32*b^2*Cos[a + b*x^(1/3)]^2)/(715*x^(11/6)) - (512*b^4*Cos[a + b*x^(1/3)]^2)/(45045*x^(7/6)) + (8192*b^6*Cos[a + b*x^(1/3)]^2)/(675675*Sqrt[x]) + (32768*b^(15/2)*Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]])/675675 - (32768*b^(15/2)*Sqrt[Pi]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a])/675675 + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(65*x^(13/6)) - (128*b^3*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(6435*x^(3/2)) + (2048*b^5*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(225225*x^(5/6)) - (32768*b^7*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(675675*x^(1/6))
```

#### Rule 3416

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
:> Module[{k = Denominator[n]}, Dist[k, Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*x^(k*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x]
&& IntegerQ[p] && FractionQ[n]
```

#### Rule 3314

```
Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
:> Simp[((c + d*x)^(m + 1)*(b*Sin[e + f*x])^n)/(d*(m + 1)), x] + (Dist[(b^2*f^2*n*(n - 1))/(d^2*(m + 1)*(m + 2)), Int[(c + d*x)^(m + 2)*(b*Sin[e + f*x])^(n - 2), x], x] - Dist[(f^2*n^2)/(d^2*(m + 1)*(m + 2)), Int[(c + d*x)^(m + 2)*(b*Sin[e + f*x])^n, x], x] - Simp[(b*f*n*(c + d*x)^(m + 2)*Cos[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegerQ[m] && IntegerQ[n] && IntegerQ[n]
```

$f*x*(b*\sin[e + f*x])^{(n - 1)}/(d^2*(m + 1)*(m + 2)), x] /; \text{FreeQ}\{b, c, d, e, f\}, x] \ \&\& \text{GtQ}[n, 1] \ \&\& \text{LtQ}[m, -2]$

### Rule 30

$\text{Int}[(x_)^{(m_.)}, x\_Symbol] \text{ :> } \text{Simp}[x^{(m + 1)}/(m + 1), x] /; \text{FreeQ}[m, x] \ \&\& \text{NeQ}[m, -1]$

### Rule 3312

$\text{Int}[(c_.) + (d_.)*(x_)^{(m_.)}*\sin[(e_.) + (f_.)*(x_)]^{(n_.)}, x\_Symbol] \text{ :> } \text{Int}[\text{ExpandTrigReduce}[(c + d*x)^m, \sin[e + f*x]^n, x], x] /; \text{FreeQ}\{c, d, e, f, m\}, x] \ \&\& \text{IGtQ}[n, 1] \ \&\& (\text{!RationalQ}[m] \ || \ (\text{GeQ}[m, -1] \ \&\& \text{LtQ}[m, 1]))$

### Rule 3306

$\text{Int}[\sin[(e_.) + (f_.)*(x_)]/\text{Sqrt}[(c_.) + (d_.)*(x_)], x\_Symbol] \text{ :> } \text{Dist}[\text{Cos}[(d*e - c*f)/d], \text{Int}[\sin[(c*f)/d + f*x]/\text{Sqrt}[c + d*x], x], x] + \text{Dist}[\sin[(d*e - c*f)/d], \text{Int}[\text{Cos}[(c*f)/d + f*x]/\text{Sqrt}[c + d*x], x], x] /; \text{FreeQ}\{c, d, e, f\}, x] \ \&\& \text{ComplexFreeQ}[f] \ \&\& \text{NeQ}[d*e - c*f, 0]$

### Rule 3305

$\text{Int}[\sin[(e_.) + (f_.)*(x_)]/\text{Sqrt}[(c_.) + (d_.)*(x_)], x\_Symbol] \text{ :> } \text{Dist}[2/d, \text{Subst}[\text{Int}[\sin[(f*x^2)/d], x], x, \text{Sqrt}[c + d*x]], x] /; \text{FreeQ}\{c, d, e, f\}, x] \ \&\& \text{ComplexFreeQ}[f] \ \&\& \text{EqQ}[d*e - c*f, 0]$

### Rule 3351

$\text{Int}[\sin[(d_.)*((e_.) + (f_.)*(x_))^{2}], x\_Symbol] \text{ :> } \text{Simp}[(\text{Sqrt}[\text{Pi}/2]*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)])/(f*\text{Rt}[d, 2]), x] /; \text{FreeQ}\{d, e, f\}, x]$

### Rule 3304

$\text{Int}[\sin[\text{Pi}/2 + (e_.) + (f_.)*(x_)]/\text{Sqrt}[(c_.) + (d_.)*(x_)], x\_Symbol] \text{ :> } \text{Dist}[2/d, \text{Subst}[\text{Int}[\text{Cos}[(f*x^2)/d], x], x, \text{Sqrt}[c + d*x]], x] /; \text{FreeQ}\{c, d, e, f\}, x] \ \&\& \text{ComplexFreeQ}[f] \ \&\& \text{EqQ}[d*e - c*f, 0]$

### Rule 3352

$\text{Int}[\text{Cos}[(d_.)*((e_.) + (f_.)*(x_))^{2}], x\_Symbol] \text{ :> } \text{Simp}[(\text{Sqrt}[\text{Pi}/2]*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)])/(f*\text{Rt}[d, 2]), x] /; \text{FreeQ}\{d, e, f\}, x]$

### Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx &= 3 \operatorname{Subst} \left( \int \frac{\cos^2(a + bx)}{x^{17/2}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{65x^{13/6}} + \frac{1}{65} (8b^2) \operatorname{Subst} \left( \int \frac{1}{x^{13/2}} dx, x, \sqrt[3]{x} \right) \\
&= -\frac{16b^2}{715x^{11/6}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{715x^{11/6}} + \frac{8b \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{65x^{13/6}} - \frac{1}{65} \\
&= -\frac{16b^2}{715x^{11/6}} + \frac{256b^4}{45045x^{7/6}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{45045x^{7/6}} \\
&= -\frac{16b^2}{715x^{11/6}} + \frac{256b^4}{45045x^{7/6}} - \frac{4096b^6}{675675\sqrt{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{45045x^{7/6}} \\
&= -\frac{16b^2}{715x^{11/6}} + \frac{256b^4}{45045x^{7/6}} - \frac{4096b^6}{675675\sqrt{x}} - \frac{65536b^8 \sqrt[6]{x}}{675675} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{715x^{11/6}} \\
&= -\frac{16b^2}{715x^{11/6}} + \frac{256b^4}{45045x^{7/6}} - \frac{4096b^6}{675675\sqrt{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{45045x^{7/6}} \\
&= -\frac{16b^2}{715x^{11/6}} + \frac{256b^4}{45045x^{7/6}} - \frac{4096b^6}{675675\sqrt{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{45045x^{7/6}} \\
&= -\frac{16b^2}{715x^{11/6}} + \frac{256b^4}{45045x^{7/6}} - \frac{4096b^6}{675675\sqrt{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{45045x^{7/6}} \\
&= -\frac{16b^2}{715x^{11/6}} + \frac{256b^4}{45045x^{7/6}} - \frac{4096b^6}{675675\sqrt{x}} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2 \cos^2(a + b\sqrt[3]{x})}{715x^{11/6}} - \frac{512b^4 \cos^2(a + b\sqrt[3]{x})}{45045x^{7/6}}
\end{aligned}$$

**Mathematica [A]** time = 0.375443, size = 249, normalized size = 0.76

$$32768\sqrt{\pi}b^{15/2}x^{5/2} \cos(2a)\operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - 32768\sqrt{\pi}b^{15/2}x^{5/2} \sin(2a)S\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - 16384b^7x^{7/3} \sin\left(2(a + b\sqrt[3]{x})\right) +$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^(1/3)]^2/x^(7/2), x]

[Out] (-135135 - 135135\*Cos[2\*(a + b\*x^(1/3))] + 15120\*b^2\*x^(2/3)\*Cos[2\*(a + b\*x^(1/3))] - 3840\*b^4\*x^(4/3)\*Cos[2\*(a + b\*x^(1/3))] + 4096\*b^6\*x^2\*Cos[2\*(a + b\*x^(1/3))] + 32768\*b^(15/2)\*Sqrt[Pi]\*x^(5/2)\*Cos[2\*a]\*FresnelC[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]] - 32768\*b^(15/2)\*Sqrt[Pi]\*x^(5/2)\*FresnelS[(2\*Sqrt[b]\*x^(1/6))/Sqrt[Pi]]\*Sin[2\*a] + 41580\*b\*x^(1/3)\*Sin[2\*(a + b\*x^(1/3))] - 6720\*b^3\*x\*Sin[2\*(a + b\*x^(1/3))] + 3072\*b^5\*x^(5/3)\*Sin[2\*(a + b\*x^(1/3))] - 16384\*b^7\*x^(7/3)\*Sin[2\*(a + b\*x^(1/3))])/(675675\*x^(5/2))

**Maple [A]** time = 0.039, size = 207, normalized size = 0.6

$$-\frac{1}{5}x^{-\frac{5}{2}} - \frac{1}{5}\cos(2a + 2b\sqrt[3]{x})x^{-\frac{5}{2}} - \frac{4b}{5}\left(-\frac{1}{13}\sin(2a + 2b\sqrt[3]{x})x^{-\frac{13}{6}} + \frac{4b}{13}\left(-\frac{1}{11}\cos(2a + 2b\sqrt[3]{x})x^{-\frac{11}{6}} - \frac{4b}{11}\left(-\frac{1}{9}\sin\right.\right.\right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a+b*x^(1/3))^2/x^(7/2),x)`

[Out] `-1/5/x^(5/2)-1/5/x^(5/2)*cos(2*a+2*b*x^(1/3))-4/5*b*(-1/13/x^(13/6)*sin(2*a+2*b*x^(1/3))+4/13*b*(-1/11/x^(11/6)*cos(2*a+2*b*x^(1/3))-4/11*b*(-1/9/x^(3/2)*sin(2*a+2*b*x^(1/3))+4/9*b*(-1/7/x^(7/6)*cos(2*a+2*b*x^(1/3))-4/7*b*(-1/5/x^(5/6)*sin(2*a+2*b*x^(1/3))+4/5*b*(-1/3/x^(1/2)*cos(2*a+2*b*x^(1/3))-4/3*b*(-1/x^(1/6)*sin(2*a+2*b*x^(1/3))+2*b^(1/2)*Pi^(1/2)*(cos(2*a)*FresnelC(2*x^(1/6)*b^(1/2)/Pi^(1/2))-sin(2*a)*FresnelS(2*x^(1/6)*b^(1/2)/Pi^(1/2))))))`

**Maxima [C]** time = 2.03386, size = 383, normalized size = 1.17

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^(1/3))^2/x^(7/2),x, algorithm="maxima")`

[Out] `-1/5*(sqrt(2)*((240*(gamma(-15/2, 2*I*b*x^(1/3)) + gamma(-15/2, -2*I*b*x^(1/3))) * cos(15/4*pi + 15/2*arctan2(0, b)) + 240*(gamma(-15/2, 2*I*b*x^(1/3)) + gamma(-15/2, -2*I*b*x^(1/3))) * cos(-15/4*pi + 15/2*arctan2(0, b)) + (240*I*gamma(-15/2, 2*I*b*x^(1/3)) - 240*I*gamma(-15/2, -2*I*b*x^(1/3))) * sin(15/4*pi + 15/2*arctan2(0, b)) + (-240*I*gamma(-15/2, 2*I*b*x^(1/3)) + 240*I*gamma(-15/2, -2*I*b*x^(1/3))) * sin(-15/4*pi + 15/2*arctan2(0, b))) * cos(2*a) + ((-240*I*gamma(-15/2, 2*I*b*x^(1/3)) + 240*I*gamma(-15/2, -2*I*b*x^(1/3))) * cos(15/4*pi + 15/2*arctan2(0, b)) + (-240*I*gamma(-15/2, 2*I*b*x^(1/3)) + 240*I*gamma(-15/2, -2*I*b*x^(1/3))) * cos(-15/4*pi + 15/2*arctan2(0, b)) + 240*(gamma(-15/2, 2*I*b*x^(1/3)) + gamma(-15/2, -2*I*b*x^(1/3))) * sin(15/4*pi + 15/2*arctan2(0, b)) - 240*(gamma(-15/2, 2*I*b*x^(1/3)) + gamma(-15/2, -2*I*b*x^(1/3))) * sin(-15/4*pi + 15/2*arctan2(0, b))) * sin(2*a)) * sqrt(x^(1/3)*abs(b)) * b^6 * x^(7/3) * abs(b) + 1) / x^(5/2)`

**Fricas [A]** time = 2.26457, size = 581, normalized size = 1.77

$$2 \left( 16384 \pi b^7 x^3 \sqrt{\frac{b}{\pi}} \cos(2a) C \left( 2x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) - 16384 \pi b^7 x^3 \sqrt{\frac{b}{\pi}} S \left( 2x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) \sin(2a) - 2048 b^6 x^{\frac{5}{2}} + 1920 b^4 x^{\frac{11}{6}} - 7560 b^2 x^{\frac{7}{6}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^(1/3))^2/x^(7/2),x, algorithm="fricas")`

[Out] `2/675675*(16384*pi*b^7*x^3*sqrt(b/pi)*cos(2*a)*fresnel_cos(2*x^(1/6)*sqrt(b/pi)) - 16384*pi*b^7*x^3*sqrt(b/pi)*fresnel_sin(2*x^(1/6)*sqrt(b/pi))*sin(2*a) - 2048*b^6*x^(5/2) + 1920*b^4*x^(11/6) - 7560*b^2*x^(7/6) - (3840*b^4*x^(11/6) - 15120*b^2*x^(7/6) - (4096*b^6*x^2 - 135135)*sqrt(x))*cos(b*x^(1/3) + a)^2 + 4*(768*b^5*x^(13/6) - 1680*b^3*x^(3/2) - (4096*b^7*x^2 - 10395*b*x^(5/6))*cos(b*x^(1/3) + a)*sin(b*x^(1/3) + a))/x^3`

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{\frac{7}{2}}} dx$$



Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x\*\*(1/3))\*\*2/x\*\*(7/2), x)

[Out] Integral(cos(a + b\*x\*\*(1/3))\*\*2/x\*\*(7/2), x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)^2}{x^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^(1/3))^2/x^(7/2), x, algorithm="giac")

[Out] integrate(cos(b\*x^(1/3) + a)^2/x^(7/2), x)

### 3.61 $\int \cos^3(\sqrt[3]{x}) dx$

**Optimal.** Leaf size=86

$$2x^{2/3} \sin(\sqrt[3]{x}) + x^{2/3} \sin(\sqrt[3]{x}) \cos^2(\sqrt[3]{x}) + \frac{2}{9} \sin^3(\sqrt[3]{x}) - \frac{14}{3} \sin(\sqrt[3]{x}) + \frac{2}{3} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) + 4\sqrt[3]{x} \cos(\sqrt[3]{x})$$

[Out]  $4x^{1/3} \cos[x^{1/3}] + (2x^{1/3} \cos[x^{1/3}]^3)/3 - (14 \sin[x^{1/3}])/3 + 2x^{2/3} \sin[x^{1/3}] + x^{2/3} \cos[x^{1/3}]^2 \sin[x^{1/3}] + (2 \sin[x^{1/3}]^3)/9$

**Rubi [A]** time = 0.0682848, antiderivative size = 86, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 5, integrand size = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3362, 3311, 3296, 2637, 2633}

$$2x^{2/3} \sin(\sqrt[3]{x}) + x^{2/3} \sin(\sqrt[3]{x}) \cos^2(\sqrt[3]{x}) + \frac{2}{9} \sin^3(\sqrt[3]{x}) - \frac{14}{3} \sin(\sqrt[3]{x}) + \frac{2}{3} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) + 4\sqrt[3]{x} \cos(\sqrt[3]{x})$$

Antiderivative was successfully verified.

[In] Int[Cos[x^(1/3)]^3,x]

[Out]  $4x^{1/3} \cos[x^{1/3}] + (2x^{1/3} \cos[x^{1/3}]^3)/3 - (14 \sin[x^{1/3}])/3 + 2x^{2/3} \sin[x^{1/3}] + x^{2/3} \cos[x^{1/3}]^2 \sin[x^{1/3}] + (2 \sin[x^{1/3}]^3)/9$

#### Rule 3362

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)])\*(b\_.))^(p\_.), x\_Symbol] :> Dist[1/(n\*f), Subst[Int[x^(1/n - 1)\*(a + b\*cos[c + d\*x])^p, x], x, (e + f\*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]

#### Rule 3311

Int[((c\_.) + (d\_.)\*(x\_))^(m\_)\*((b\_.)\*sin[(e\_.) + (f\_.)\*(x\_)])^(n\_), x\_Symbol] :> Simp[(d\*m\*(c + d\*x)^(m - 1)\*(b\*sin[e + f\*x])^n)/(f^2\*n^2), x] + (Dist[(b^2\*(n - 1))/n, Int[(c + d\*x)^m\*(b\*sin[e + f\*x])^(n - 2), x], x] - Dist[(d^2\*m\*(m - 1))/(f^2\*n^2), Int[(c + d\*x)^(m - 2)\*(b\*sin[e + f\*x])^n, x], x] - Simp[(b\*(c + d\*x)^m\*cos[e + f\*x]\*(b\*sin[e + f\*x])^(n - 1))/(f\*n), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]

#### Rule 3296

Int[((c\_.) + (d\_.)\*(x\_))^(m\_)\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] :> -Simp[(c + d\*x)^m\*cos[e + f\*x]/f, x] + Dist[(d\*m)/f, Int[(c + d\*x)^(m - 1)\*cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

#### Rule 2637

Int[sin[Pi/2 + (c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Simp[Sin[c + d\*x]/d, x] /; FreeQ[{c, d}, x]

#### Rule 2633

Int[sin[(c\_.) + (d\_.)\*(x\_)]^(n\_), x\_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d\*x]], x] /; FreeQ[{c, d}, x]

&& IGtQ[(n - 1)/2, 0]

### Rubi steps

$$\begin{aligned}
 \int \cos^3(\sqrt[3]{x}) dx &= 3 \operatorname{Subst}\left(\int x^2 \cos^3(x) dx, x, \sqrt[3]{x}\right) \\
 &= \frac{2}{3} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) + x^{2/3} \cos^2(\sqrt[3]{x}) \sin(\sqrt[3]{x}) - \frac{2}{3} \operatorname{Subst}\left(\int \cos^3(x) dx, x, \sqrt[3]{x}\right) + 2 \operatorname{Subst}\left(\int x^2 \cos(x) dx, x, \sqrt[3]{x}\right) \\
 &= \frac{2}{3} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) + 2x^{2/3} \sin(\sqrt[3]{x}) + x^{2/3} \cos^2(\sqrt[3]{x}) \sin(\sqrt[3]{x}) + \frac{2}{3} \operatorname{Subst}\left(\int (1-x^2) dx, x, -\sin(\sqrt[3]{x})\right) \\
 &= 4\sqrt[3]{x} \cos(\sqrt[3]{x}) + \frac{2}{3} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) - \frac{2}{3} \sin(\sqrt[3]{x}) + 2x^{2/3} \sin(\sqrt[3]{x}) + x^{2/3} \cos^2(\sqrt[3]{x}) \sin(\sqrt[3]{x}) + \frac{2}{9} \sin^3(\sqrt[3]{x}) \\
 &= 4\sqrt[3]{x} \cos(\sqrt[3]{x}) + \frac{2}{3} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) - \frac{14}{3} \sin(\sqrt[3]{x}) + 2x^{2/3} \sin(\sqrt[3]{x}) + x^{2/3} \cos^2(\sqrt[3]{x}) \sin(\sqrt[3]{x}) + \frac{2}{9} \sin^3(\sqrt[3]{x})
 \end{aligned}$$

**Mathematica [A]** time = 0.0992082, size = 66, normalized size = 0.77

$$\frac{1}{36} (81 (x^{2/3} - 2) \sin(\sqrt[3]{x}) + (9x^{2/3} - 2) \sin(3\sqrt[3]{x}) + 162\sqrt[3]{x} \cos(\sqrt[3]{x}) + 6\sqrt[3]{x} \cos(3\sqrt[3]{x}))$$

Antiderivative was successfully verified.

[In] Integrate[Cos[x^(1/3)]^3, x]

[Out] (162\*x^(1/3)\*Cos[x^(1/3)] + 6\*x^(1/3)\*Cos[3\*x^(1/3)] + 81\*(-2 + x^(2/3))\*Sin[x^(1/3)] + (-2 + 9\*x^(2/3))\*Sin[3\*x^(1/3)])/36

**Maple [A]** time = 0.029, size = 58, normalized size = 0.7

$$x^{\frac{2}{3}} \left( 2 + (\cos(\sqrt[3]{x}))^2 \right) \sin(\sqrt[3]{x}) - 4 \sin(\sqrt[3]{x}) + 4 \sqrt[3]{x} \cos(\sqrt[3]{x}) + \frac{2}{3} \sqrt[3]{x} (\cos(\sqrt[3]{x}))^3 - \frac{2}{9} \left( 2 + (\cos(\sqrt[3]{x}))^2 \right) \sin(\sqrt[3]{x})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x^(1/3))^3, x)

[Out] x^(2/3)\*(2+cos(x^(1/3))^2)\*sin(x^(1/3))-4\*sin(x^(1/3))+4\*x^(1/3)\*cos(x^(1/3))+2/3\*x^(1/3)\*cos(x^(1/3))^3-2/9\*(2+cos(x^(1/3))^2)\*sin(x^(1/3))

**Maxima [A]** time = 1.78985, size = 63, normalized size = 0.73

$$\frac{1}{36} \left( 9x^{\frac{2}{3}} - 2 \right) \sin\left( 3x^{\frac{1}{3}} \right) + \frac{9}{4} \left( x^{\frac{2}{3}} - 2 \right) \sin\left( x^{\frac{1}{3}} \right) + \frac{1}{6} x^{\frac{1}{3}} \cos\left( 3x^{\frac{1}{3}} \right) + \frac{9}{2} x^{\frac{1}{3}} \cos\left( x^{\frac{1}{3}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x^(1/3))^3, x, algorithm="maxima")

[Out] 1/36\*(9\*x^(2/3) - 2)\*sin(3\*x^(1/3)) + 9/4\*(x^(2/3) - 2)\*sin(x^(1/3)) + 1/6\*x^(1/3)\*cos(3\*x^(1/3)) + 9/2\*x^(1/3)\*cos(x^(1/3))

**Fricas [A]** time = 1.90552, size = 166, normalized size = 1.93

$$\frac{2}{3}x^{\frac{1}{3}}\cos\left(x^{\frac{1}{3}}\right)^3 + \frac{1}{9}\left(\left(9x^{\frac{2}{3}} - 2\right)\cos\left(x^{\frac{1}{3}}\right)^2 + 18x^{\frac{2}{3}} - 40\right)\sin\left(x^{\frac{1}{3}}\right) + 4x^{\frac{1}{3}}\cos\left(x^{\frac{1}{3}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x^(1/3))^3,x, algorithm="fricas")

[Out] 2/3\*x^(1/3)\*cos(x^(1/3))^3 + 1/9\*((9\*x^(2/3) - 2)\*cos(x^(1/3))^2 + 18\*x^(2/3) - 40)\*sin(x^(1/3)) + 4\*x^(1/3)\*cos(x^(1/3))

**Sympy [B]** time = 5.29064, size = 513, normalized size = 5.97

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x\*\*(1/3))\*\*3,x)

[Out] 54\*x\*\*(2/3)\*tan(x\*\*(1/3)/2)\*\*5/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) + 36\*x\*\*(2/3)\*tan(x\*\*(1/3)/2)\*\*3/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) + 54\*x\*\*(2/3)\*tan(x\*\*(1/3)/2)/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) - 42\*x\*\*(1/3)\*tan(x\*\*(1/3)/2)\*\*6/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) - 18\*x\*\*(1/3)\*tan(x\*\*(1/3)/2)\*\*4/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) + 18\*x\*\*(1/3)\*tan(x\*\*(1/3)/2)\*\*2/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) + 42\*x\*\*(1/3)/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) - 84\*tan(x\*\*(1/3)/2)\*\*5/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) - 152\*tan(x\*\*(1/3)/2)\*\*3/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9) - 84\*tan(x\*\*(1/3)/2)/(9\*tan(x\*\*(1/3)/2)\*\*6 + 27\*tan(x\*\*(1/3)/2)\*\*4 + 27\*tan(x\*\*(1/3)/2)\*\*2 + 9)

**Giac [A]** time = 1.15009, size = 63, normalized size = 0.73

$$\frac{1}{36}\left(9x^{\frac{2}{3}} - 2\right)\sin\left(3x^{\frac{1}{3}}\right) + \frac{9}{4}\left(x^{\frac{2}{3}} - 2\right)\sin\left(x^{\frac{1}{3}}\right) + \frac{1}{6}x^{\frac{1}{3}}\cos\left(3x^{\frac{1}{3}}\right) + \frac{9}{2}x^{\frac{1}{3}}\cos\left(x^{\frac{1}{3}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x^(1/3))^3,x, algorithm="giac")

[Out] 1/36\*(9\*x^(2/3) - 2)\*sin(3\*x^(1/3)) + 9/4\*(x^(2/3) - 2)\*sin(x^(1/3)) + 1/6\*x^(1/3)\*cos(3\*x^(1/3)) + 9/2\*x^(1/3)\*cos(x^(1/3))

$$3.62 \quad \int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx$$

**Optimal.** Leaf size=8

$$6 \sin(\sqrt[6]{x})$$

[Out] 6\*Sin[x^(1/6)]

**Rubi [A]** time = 0.0097518, antiderivative size = 8, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3380, 2637}

$$6 \sin(\sqrt[6]{x})$$

Antiderivative was successfully verified.

[In] Int[Cos[x^(1/6)]/x^(5/6), x]

[Out] 6\*Sin[x^(1/6)]

#### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

#### Rule 2637

Int[sin[Pi/2 + (c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Simp[Sin[c + d\*x]/d, x] /; FreeQ[{c, d}, x]

#### Rubi steps

$$\begin{aligned} \int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx &= 6 \text{Subst} \left( \int \cos(x) dx, x, \sqrt[6]{x} \right) \\ &= 6 \sin(\sqrt[6]{x}) \end{aligned}$$

**Mathematica [A]** time = 0.0022308, size = 8, normalized size = 1.

$$6 \sin(\sqrt[6]{x})$$

Antiderivative was successfully verified.

[In] Integrate[Cos[x^(1/6)]/x^(5/6), x]

[Out] 6\*Sin[x^(1/6)]

**Maple [A]** time = 0.023, size = 7, normalized size = 0.9

$$6 \sin\left(\sqrt[6]{x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(x^(1/6))/x^(5/6),x)`

[Out] `6*sin(x^(1/6))`

---

**Maxima [A]** time = 1.74407, size = 8, normalized size = 1.

$$6 \sin\left(x^{\frac{1}{6}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x^(1/6))/x^(5/6),x, algorithm="maxima")`

[Out] `6*sin(x^(1/6))`

---

**Fricas [A]** time = 1.8284, size = 22, normalized size = 2.75

$$6 \sin\left(x^{\frac{1}{6}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x^(1/6))/x^(5/6),x, algorithm="fricas")`

[Out] `6*sin(x^(1/6))`

---

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x**(1/6))/x**(5/6),x)`

[Out] Timed out

---

**Giac [A]** time = 1.14654, size = 8, normalized size = 1.

$$6 \sin\left(x^{\frac{1}{6}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x^(1/6))/x^(5/6),x, algorithm="giac")`

[Out] `6*sin(x^(1/6))`

### 3.63 $\int (ex)^m (b \cos(c + dx^n))^p dx$

**Optimal.** Leaf size=20

$$\text{Unintegrable}((ex)^m (b \cos(c + dx^n))^p, x)$$

[Out] Unintegrable[(e\*x)^m\*(b\*cos[c + d\*x^n])^p, x]

**Rubi [A]** time = 0.0196607, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.$ , Rules used = {}

$$\int (ex)^m (b \cos(c + dx^n))^p dx$$

Verification is Not applicable to the result.

[In] Int[(e\*x)^m\*(b\*cos[c + d\*x^n])^p, x]

[Out] Defer[Int] [(e\*x)^m\*(b\*cos[c + d\*x^n])^p, x]

Rubi steps

$$\int (ex)^m (b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(c + dx^n))^p dx$$

**Mathematica [A]** time = 0.987134, size = 0, normalized size = 0.

$$\int (ex)^m (b \cos(c + dx^n))^p dx$$

Verification is Not applicable to the result.

[In] Integrate[(e\*x)^m\*(b\*cos[c + d\*x^n])^p, x]

[Out] Integrate[(e\*x)^m\*(b\*cos[c + d\*x^n])^p, x]

**Maple [A]** time = 1.156, size = 0, normalized size = 0.

$$\int (ex)^m (b \cos(c + dx^n))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e\*x)^m\*(b\*cos(c+d\*x^n))^p, x)

[Out] int((e\*x)^m\*(b\*cos(c+d\*x^n))^p, x)

**Maxima [A]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^m (b \cos(dx^n + c))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^m\*(b\*cos(c+d\*x^n))^p,x, algorithm="maxima")

[Out] integrate((e\*x)^m\*(b\*cos(d\*x^n + c))^p, x)

**Fricas [A]** time = 0., size = 0, normalized size = 0.

$$\text{integral} \left( (ex)^m (b \cos(dx^n + c))^p, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^m\*(b\*cos(c+d\*x^n))^p,x, algorithm="fricas")

[Out] integral((e\*x)^m\*(b\*cos(d\*x^n + c))^p, x)

**Sympy [A]** time = 0., size = 0, normalized size = 0.

$$\int (b \cos(c + dx^n))^p (ex)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)\*\*m\*(b\*cos(c+d\*x\*\*n))\*\*p,x)

[Out] Integral((b\*cos(c + d\*x\*\*n))\*\*p\*(e\*x)\*\*m, x)

**Giac [A]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^m (b \cos(dx^n + c))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^m\*(b\*cos(c+d\*x^n))^p,x, algorithm="giac")

[Out] integrate((e\*x)^m\*(b\*cos(d\*x^n + c))^p, x)



### 3.64 $\int (ex)^m (a + b \cos(c + dx^n))^p dx$

**Optimal.** Leaf size=22

$$\text{Unintegrable}((ex)^m (a + b \cos(c + dx^n))^p, x)$$

[Out] Unintegrable[(e\*x)^m\*(a + b\*cos[c + d\*x^n])^p, x]

**Rubi [A]** time = 0.0224955, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.$ , Rules used = {}

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx$$

Verification is Not applicable to the result.

[In] Int[(e\*x)^m\*(a + b\*cos[c + d\*x^n])^p, x]

[Out] Defer[Int] [(e\*x)^m\*(a + b\*cos[c + d\*x^n])^p, x]

Rubi steps

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = \int (ex)^m (a + b \cos(c + dx^n))^p dx$$

**Mathematica [A]** time = 1.26057, size = 0, normalized size = 0.

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx$$

Verification is Not applicable to the result.

[In] Integrate[(e\*x)^m\*(a + b\*cos[c + d\*x^n])^p, x]

[Out] Integrate[(e\*x)^m\*(a + b\*cos[c + d\*x^n])^p, x]

**Maple [A]** time = 1.145, size = 0, normalized size = 0.

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e\*x)^m\*(a+b\*cos(c+d\*x^n))^p, x)

[Out] int((e\*x)^m\*(a+b\*cos(c+d\*x^n))^p, x)

**Maxima [A]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^m (b \cos(dx^n + c) + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^m\*(a+b\*cos(c+d\*x^n))^p,x, algorithm="maxima")

[Out] integrate((e\*x)^m\*(b\*cos(d\*x^n + c) + a)^p, x)

**Fricas [A]** time = 0., size = 0, normalized size = 0.

$$\text{integral}\left((ex)^m (b \cos(dx^n + c) + a)^p, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^m\*(a+b\*cos(c+d\*x^n))^p,x, algorithm="fricas")

[Out] integral((e\*x)^m\*(b\*cos(d\*x^n + c) + a)^p, x)

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)\*\*m\*(a+b\*cos(c+d\*x\*\*n))\*\*p,x)

[Out] Timed out

**Giac [A]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^m (b \cos(dx^n + c) + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^m\*(a+b\*cos(c+d\*x^n))^p,x, algorithm="giac")

[Out] integrate((e\*x)^m\*(b\*cos(d\*x^n + c) + a)^p, x)

### 3.65 $\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx$

**Optimal.** Leaf size=93

$$\frac{x^{-n}(ex)^n \sin(c + dx^n) (b \cos(c + dx^n))^{p+1} {}_2F_1\left(\frac{1}{2}, \frac{p+1}{2}; \frac{p+3}{2}; \cos^2(dx^n + c)\right)}{b \operatorname{den}(p+1) \sqrt{\sin^2(c + dx^n)}}$$

[Out] -(((e\*x)^n\*(b\*Cos[c + d\*x^n])^(1 + p)\*Hypergeometric2F1[1/2, (1 + p)/2, (3 + p)/2, Cos[c + d\*x^n]^2]\*Sin[c + d\*x^n])/(b\*d\*e\*n\*(1 + p)\*x^n\*Sqrt[Sin[c + d\*x^n]^2]))

**Rubi [A]** time = 0.101958, antiderivative size = 93, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 20,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.15$ , Rules used = {3382, 3380, 2643}

$$\frac{x^{-n}(ex)^n \sin(c + dx^n) (b \cos(c + dx^n))^{p+1} {}_2F_1\left(\frac{1}{2}, \frac{p+1}{2}; \frac{p+3}{2}; \cos^2(dx^n + c)\right)}{b \operatorname{den}(p+1) \sqrt{\sin^2(c + dx^n)}}$$

Antiderivative was successfully verified.

[In] Int[(e\*x)^(-1 + n)\*(b\*Cos[c + d\*x^n])^p,x]

[Out] -(((e\*x)^n\*(b\*Cos[c + d\*x^n])^(1 + p)\*Hypergeometric2F1[1/2, (1 + p)/2, (3 + p)/2, Cos[c + d\*x^n]^2]\*Sin[c + d\*x^n])/(b\*d\*e\*n\*(1 + p)\*x^n\*Sqrt[Sin[c + d\*x^n]^2]))

#### Rule 3382

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*((e\_.)\*(x\_.))^(m\_.), x\_Symbol] :> Dist[(e^IntPart[m]\*(e\*x)^FracPart[m])/x^FracPart[m], Int[x^m\*(a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

#### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

#### Rule 2643

Int[((b\_.)\*sin[(c\_.) + (d\_.)\*(x\_)])^(n\_), x\_Symbol] :> Simp[(Cos[c + d\*x]\*(b\*Sin[c + d\*x])^(n + 1)\*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d\*x]^2])/(b\*d\*(n + 1)\*Sqrt[Cos[c + d\*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2\*n]

#### Rubi steps

$$\begin{aligned} \int (ex)^{-1+n} (b \cos(c + dx^n))^p dx &= \frac{(x^{-n}(ex)^n) \int x^{-1+n} (b \cos(c + dx^n))^p dx}{e} \\ &= \frac{(x^{-n}(ex)^n) \text{Subst}\left(\int (b \cos(c + dx))^p dx, x, x^n\right)}{en} \\ &= -\frac{x^{-n}(ex)^n (b \cos(c + dx^n))^{1+p} {}_2F_1\left(\frac{1}{2}, \frac{1+p}{2}; \frac{3+p}{2}; \cos^2(c + dx^n)\right) \sin(c + dx^n)}{bden(1+p)\sqrt{\sin^2(c + dx^n)}} \end{aligned}$$

**Mathematica [A]** time = 0.204899, size = 89, normalized size = 0.96

$$\frac{x^{1-n}(ex)^{n-1}\sqrt{\sin^2(c + dx^n)} \cot(c + dx^n) (b \cos(c + dx^n))^p {}_2F_1\left(\frac{1}{2}, \frac{p+1}{2}; \frac{p+3}{2}; \cos^2(dx^n + c)\right)}{dn(p+1)}$$

Antiderivative was successfully verified.

[In] Integrate[(e\*x)^(-1 + n)\*(b\*Cos[c + d\*x^n])^p, x]

[Out] -((x^(1 - n)\*(e\*x)^(-1 + n)\*(b\*Cos[c + d\*x^n])^p\*Cot[c + d\*x^n]\*Hypergeometric2F1[1/2, (1 + p)/2, (3 + p)/2, Cos[c + d\*x^n]^2]\*Sqrt[Sin[c + d\*x^n]^2])/(d\*n\*(1 + p))

**Maple [F]** time = 1.874, size = 0, normalized size = 0.

$$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e\*x)^(-1+n)\*(b\*cos(c+d\*x^n))^p, x)

[Out] int((e\*x)^(-1+n)\*(b\*cos(c+d\*x^n))^p, x)

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^{n-1} (b \cos(dx^n + c))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+n)\*(b\*cos(c+d\*x^n))^p, x, algorithm="maxima")

[Out] integrate((e\*x)^(n - 1)\*(b\*cos(d\*x^n + c))^p, x)

**Fricas [F]** time = 0., size = 0, normalized size = 0.

$$\text{integral}\left((ex)^{n-1} (b \cos(dx^n + c))^p, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+n)\*(b\*cos(c+d\*x^n))^p,x, algorithm="fricas")

[Out] integral((e\*x)^(n - 1)\*(b\*cos(d\*x^n + c))^p, x)

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)\*\*(-1+n)\*(b\*cos(c+d\*x\*\*n))\*\*p,x)

[Out] Timed out

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^{n-1} (b \cos(dx^n + c))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+n)\*(b\*cos(c+d\*x^n))^p,x, algorithm="giac")

[Out] integrate((e\*x)^(n - 1)\*(b\*cos(d\*x^n + c))^p, x)

### 3.66 $\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$

**Optimal.** Leaf size=38

$$\frac{x^{-2n}(ex)^{2n}\text{Unintegrable}(x^{2n-1}(b \cos(c + dx^n))^p, x)}{e}$$

[Out] ((e\*x)^(2\*n)\*Unintegrable[x^(-1 + 2\*n)\*(b\*Cos[c + d\*x^n])^p, x])/(e\*x^(2\*n))

**Rubi [A]** time = 0.0486985, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.$ , Rules used = {}

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$$

Verification is Not applicable to the result.

[In] Int[(e\*x)^(-1 + 2\*n)\*(b\*Cos[c + d\*x^n])^p, x]

[Out] ((e\*x)^(2\*n)\*Defer[Int][x^(-1 + 2\*n)\*(b\*Cos[c + d\*x^n])^p, x])/(e\*x^(2\*n))

Rubi steps

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \frac{(x^{-2n}(ex)^{2n}) \int x^{-1+2n} (b \cos(c + dx^n))^p dx}{e}$$

**Mathematica [A]** time = 0.934475, size = 0, normalized size = 0.

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$$

Verification is Not applicable to the result.

[In] Integrate[(e\*x)^(-1 + 2\*n)\*(b\*Cos[c + d\*x^n])^p, x]

[Out] Integrate[(e\*x)^(-1 + 2\*n)\*(b\*Cos[c + d\*x^n])^p, x]

**Maple [A]** time = 1.342, size = 0, normalized size = 0.

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e\*x)^(-1+2\*n)\*(b\*cos(c+d\*x^n))^p, x)

[Out] int((e\*x)^(-1+2\*n)\*(b\*cos(c+d\*x^n))^p, x)

**Maxima [A]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^{2n-1} (b \cos(dx^n + c))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+2\*n)\*(b\*cos(c+d\*x^n))^p,x, algorithm="maxima")

[Out] integrate((e\*x)^(2\*n - 1)\*(b\*cos(d\*x^n + c))^p, x)

---

**Fricas [A]** time = 0., size = 0, normalized size = 0.

$$\text{integral}\left((ex)^{2n-1} (b \cos(dx^n + c))^p, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+2\*n)\*(b\*cos(c+d\*x^n))^p,x, algorithm="fricas")

[Out] integral((e\*x)^(2\*n - 1)\*(b\*cos(d\*x^n + c))^p, x)

---

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)\*\*(-1+2\*n)\*(b\*cos(c+d\*x\*\*n))\*\*p,x)

[Out] Timed out

---

**Giac [A]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^{2n-1} (b \cos(dx^n + c))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+2\*n)\*(b\*cos(c+d\*x^n))^p,x, algorithm="giac")

[Out] integrate((e\*x)^(2\*n - 1)\*(b\*cos(d\*x^n + c))^p, x)

### 3.67 $\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx$

**Optimal.** Leaf size=131

$$\frac{\sqrt{2}x^{-n}(ex)^n \sin(c + dx^n) (a + b \cos(c + dx^n))^p \left(\frac{a+b \cos(c+dx^n)}{a+b}\right)^{-p} F_1\left(\frac{1}{2}; \frac{1}{2}, -p; \frac{3}{2}; \frac{1}{2}(1 - \cos(dx^n + c)), \frac{b(1-\cos(dx^n+c))}{a+b}\right)}{\text{den} \sqrt{\cos(c + dx^n) + 1}}$$

[Out] (Sqrt[2]\*(e\*x)^n\*AppellF1[1/2, 1/2, -p, 3/2, (1 - Cos[c + d\*x^n])/2, (b\*(1 - Cos[c + d\*x^n]))/(a + b)]\*(a + b\*Cos[c + d\*x^n])^p\*Sin[c + d\*x^n])/(d\*e\*n\*x^n\*Sqrt[1 + Cos[c + d\*x^n]]\*((a + b\*Cos[c + d\*x^n])/(a + b))^p)

**Rubi [A]** time = 0.166583, antiderivative size = 131, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 22,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$ , Rules used = {3382, 3380, 2665, 139, 138}

$$\frac{\sqrt{2}x^{-n}(ex)^n \sin(c + dx^n) (a + b \cos(c + dx^n))^p \left(\frac{a+b \cos(c+dx^n)}{a+b}\right)^{-p} F_1\left(\frac{1}{2}; \frac{1}{2}, -p; \frac{3}{2}; \frac{1}{2}(1 - \cos(dx^n + c)), \frac{b(1-\cos(dx^n+c))}{a+b}\right)}{\text{den} \sqrt{\cos(c + dx^n) + 1}}$$

Antiderivative was successfully verified.

[In] Int[(e\*x)^(-1 + n)\*(a + b\*Cos[c + d\*x^n])^p,x]

[Out] (Sqrt[2]\*(e\*x)^n\*AppellF1[1/2, 1/2, -p, 3/2, (1 - Cos[c + d\*x^n])/2, (b\*(1 - Cos[c + d\*x^n]))/(a + b)]\*(a + b\*Cos[c + d\*x^n])^p\*Sin[c + d\*x^n])/(d\*e\*n\*x^n\*Sqrt[1 + Cos[c + d\*x^n]]\*((a + b\*Cos[c + d\*x^n])/(a + b))^p)

#### Rule 3382

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] := Dist[(e^IntPart[m]\*(e\*x)^FracPart[m])/x^FracPart[m], Int[x^m\*(a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

#### Rule 3380

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_.)\*(x\_)^(m\_.), x\_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))

#### Rule 2665

Int[((a\_) + (b\_.)\*sin[(c\_.) + (d\_.)\*(x\_)])^(n\_), x\_Symbol] := Dist[Cos[c + d\*x]/(d\*Sqrt[1 + Sin[c + d\*x]]\*Sqrt[1 - Sin[c + d\*x]]), Subst[Int[(a + b\*x)^n/(Sqrt[1 + x]\*Sqrt[1 - x]), x], x, Sin[c + d\*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2\*n]

#### Rule 139

Int[((a\_) + (b\_.)\*(x\_)^(m\_))\*((c\_.) + (d\_.)\*(x\_)^(n\_))\*((e\_.) + (f\_.)\*(x\_)^(p\_)), x\_Symbol] := Dist[(e + f\*x)^FracPart[p]/((b/(b\*e - a\*f))^IntPart[p]\*((b\*(e + f\*x))/(b\*e - a\*f))^FracPart[p]), Int[(a + b\*x)^m\*(c + d\*x)^n\*((b\*e)/(b\*e - a\*f) + (b\*f\*x)/(b\*e - a\*f))^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[b/(b



\*c - a\*d), 0] && !GtQ[b/(b\*e - a\*f), 0]

### Rule 138

Int[((a\_) + (b\_)\*(x\_))^(m\_)\*((c\_) + (d\_)\*(x\_))^(n\_)\*((e\_) + (f\_)\*(x\_))^(p\_), x\_Symbol] := Simp[((a + b\*x)^(m + 1)\*AppellF1[m + 1, -n, -p, m + 2, -((d\*(a + b\*x))/(b\*c - a\*d)), -((f\*(a + b\*x))/(b\*e - a\*f))])/(b\*(m + 1)\*(b/(b\*c - a\*d))^n\*(b/(b\*e - a\*f))^p), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[b/(b\*c - a\*d), 0] && GtQ[b/(b\*e - a\*f), 0] && !(GtQ[d/(d\*a - c\*b), 0] && GtQ[d/(d\*e - c\*f), 0]) && SimplrQ[c + d\*x, a + b\*x] && !(GtQ[f/(f\*a - e\*b), 0] && GtQ[f/(f\*c - e\*d), 0]) && SimplrQ[e + f\*x, a + b\*x]

### Rubi steps

$$\begin{aligned} \int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx &= \frac{(x^{-n}(ex)^n) \int x^{-1+n} (a + b \cos(c + dx^n))^p dx}{e} \\ &= \frac{(x^{-n}(ex)^n) \text{Subst}\left(\int (a + b \cos(c + dx))^p dx, x, x^n\right)}{en} \\ &= -\frac{(x^{-n}(ex)^n \sin(c + dx^n)) \text{Subst}\left(\int \frac{(a+bx)^p}{\sqrt{1-x}\sqrt{1+x}} dx, x, \cos(c + dx^n)\right)}{den\sqrt{1 - \cos(c + dx^n)}\sqrt{1 + \cos(c + dx^n)}} \\ &= -\frac{\left(x^{-n}(ex)^n (a + b \cos(c + dx^n))^p \left(-\frac{a+b \cos(c+dx^n)}{-a-b}\right)^{-p} \sin(c + dx^n)\right) \text{Subst}\left(\int \frac{(-x)^p}{\sqrt{1-x}\sqrt{1+x}} dx, x, \cos(c + dx^n)\right)}{den\sqrt{1 - \cos(c + dx^n)}\sqrt{1 + \cos(c + dx^n)}} \\ &= \frac{\sqrt{2}x^{-n}(ex)^n F_1\left(\frac{1}{2}; \frac{1}{2}, -p; \frac{3}{2}; \frac{1}{2}(1 - \cos(c + dx^n)), \frac{b(1 - \cos(c+dx^n))}{a+b}\right) (a + b \cos(c + dx^n))^p}{den\sqrt{1 + \cos(c + dx^n)}} \end{aligned}$$

**Mathematica [A]** time = 0.440903, size = 149, normalized size = 1.14

$$\frac{x^{-n}(ex)^n \csc(c + dx^n) \sqrt{-\frac{b(\cos(c+dx^n)-1)}{a+b}} \sqrt{\frac{b(\cos(c+dx^n)+1)}{b-a}} (a + b \cos(c + dx^n))^{p+1} F_1\left(p + 1; \frac{1}{2}, \frac{1}{2}; p + 2; \frac{a+b \cos(dx^n+c)}{a-b}, \frac{a+b \cos(dx^n+c)}{a-b}\right)}{bden(p + 1)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(e\*x)^(-1+n)\*(a + b\*Cos[c + d\*x^n])^p,x]

[Out] -(((e\*x)^n\*AppellF1[1 + p, 1/2, 1/2, 2 + p, (a + b\*Cos[c + d\*x^n])/(a - b), (a + b\*Cos[c + d\*x^n])/(a + b)]\*Sqrt[-((b\*(-1 + Cos[c + d\*x^n]))/(a + b))]\*Sqrt[(b\*(1 + Cos[c + d\*x^n]))/(-a + b)]\*(a + b\*Cos[c + d\*x^n])^(1 + p)\*Csc[c + d\*x^n])/(b\*d\*e\*n\*(1 + p)\*x^n)

**Maple [F]** time = 1.907, size = 0, normalized size = 0.

$$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e\*x)^(-1+n)\*(a+b\*cos(c+d\*x^n))^p,x)

[Out] `int((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x)`

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^{n-1} (b \cos(dx^n + c) + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x, algorithm="maxima")`

[Out] `integrate((e*x)^(n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

**Fricas [F]** time = 0., size = 0, normalized size = 0.

$$\text{integral}((ex)^{n-1} (b \cos(dx^n + c) + a)^p, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x, algorithm="fricas")`

[Out] `integral((e*x)^(n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x)**(-1+n)*(a+b*cos(c+d*x**n))**p,x)`

[Out] Timed out

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^{n-1} (b \cos(dx^n + c) + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x, algorithm="giac")`

[Out] `integrate((e*x)^(n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

### 3.68 $\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx$

**Optimal.** Leaf size=40

$$\frac{x^{-2n}(ex)^{2n}\text{Unintegrable}(x^{2n-1}(a + b \cos(c + dx^n))^p, x)}{e}$$

[Out]  $((e*x)^{(2*n)}*\text{Unintegrable}[x^{(-1 + 2*n)}*(a + b*\text{Cos}[c + d*x^n])^p, x])/(e*x^{(2*n)})$

**Rubi [A]** time = 0.054143, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.$ , Rules used = {}

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx$$

Verification is Not applicable to the result.

[In]  $\text{Int}[(e*x)^{(-1 + 2*n)}*(a + b*\text{Cos}[c + d*x^n])^p, x]$

[Out]  $((e*x)^{(2*n)}*\text{Defer}[\text{Int}[x^{(-1 + 2*n)}*(a + b*\text{Cos}[c + d*x^n])^p, x])/(e*x^{(2*n)})$

Rubi steps

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx = \frac{(x^{-2n}(ex)^{2n}) \int x^{-1+2n} (a + b \cos(c + dx^n))^p dx}{e}$$

**Mathematica [A]** time = 1.15776, size = 0, normalized size = 0.

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx$$

Verification is Not applicable to the result.

[In]  $\text{Integrate}[(e*x)^{(-1 + 2*n)}*(a + b*\text{Cos}[c + d*x^n])^p, x]$

[Out]  $\text{Integrate}[(e*x)^{(-1 + 2*n)}*(a + b*\text{Cos}[c + d*x^n])^p, x]$

**Maple [A]** time = 1.388, size = 0, normalized size = 0.

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In]  $\text{int}((e*x)^{(-1+2*n)}*(a+b*\text{cos}(c+d*x^n))^p, x)$

[Out]  $\text{int}((e*x)^{(-1+2*n)}*(a+b*\text{cos}(c+d*x^n))^p, x)$

---

**Maxima [A]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^{2n-1} (b \cos(dx^n + c) + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+2\*n)\*(a+b\*cos(c+d\*x^n))^p,x, algorithm="maxima")

[Out] integrate((e\*x)^(2\*n - 1)\*(b\*cos(d\*x^n + c) + a)^p, x)

---

**Fricas [A]** time = 0., size = 0, normalized size = 0.

$$\text{integral}((ex)^{2n-1} (b \cos(dx^n + c) + a)^p, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+2\*n)\*(a+b\*cos(c+d\*x^n))^p,x, algorithm="fricas")

[Out] integral((e\*x)^(2\*n - 1)\*(b\*cos(d\*x^n + c) + a)^p, x)

---

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)\*\*(-1+2\*n)\*(a+b\*cos(c+d\*x\*\*n))\*\*p,x)

[Out] Timed out

---

**Giac [A]** time = 0., size = 0, normalized size = 0.

$$\int (ex)^{2n-1} (b \cos(dx^n + c) + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e\*x)^(-1+2\*n)\*(a+b\*cos(c+d\*x^n))^p,x, algorithm="giac")

[Out] integrate((e\*x)^(2\*n - 1)\*(b\*cos(d\*x^n + c) + a)^p, x)

$$3.69 \quad \int \frac{\cos(a+bx^n)}{x} dx$$

**Optimal.** Leaf size=26

$$\frac{\cos(a)\text{CosIntegral}(bx^n)}{n} - \frac{\sin(a)\text{Si}(bx^n)}{n}$$

[Out] (Cos[a]\*CosIntegral[b\*x^n])/n - (Sin[a]\*SinIntegral[b\*x^n])/n

**Rubi [A]** time = 0.0339921, antiderivative size = 26, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3378, 3376, 3375}

$$\frac{\cos(a)\text{CosIntegral}(bx^n)}{n} - \frac{\sin(a)\text{Si}(bx^n)}{n}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^n]/x,x]

[Out] (Cos[a]\*CosIntegral[b\*x^n])/n - (Sin[a]\*SinIntegral[b\*x^n])/n

**Rule 3378**

Int[Cos[(c\_) + (d\_)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Dist[Cos[c], Int[Cos[d\*x^n]/x, x], x] - Dist[Sin[c], Int[Sin[d\*x^n]/x, x], x] /; FreeQ[{c, d, n}, x]

**Rule 3376**

Int[Cos[(d\_)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[CosIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

**Rule 3375**

Int[Sin[(d\_)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[SinIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

**Rubi steps**

$$\begin{aligned} \int \frac{\cos(a+bx^n)}{x} dx &= \cos(a) \int \frac{\cos(bx^n)}{x} dx - \sin(a) \int \frac{\sin(bx^n)}{x} dx \\ &= \frac{\cos(a)\text{Ci}(bx^n)}{n} - \frac{\sin(a)\text{Si}(bx^n)}{n} \end{aligned}$$

**Mathematica [A]** time = 0.0645611, size = 24, normalized size = 0.92

$$\frac{\cos(a)\text{CosIntegral}(bx^n) - \sin(a)\text{Si}(bx^n)}{n}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^n]/x,x]

[Out]  $(\cos(a) \cdot \text{CosIntegral}[b \cdot x^n] - \sin(a) \cdot \text{SinIntegral}[b \cdot x^n]) / n$

---

**Maple [A]** time = 0.034, size = 25, normalized size = 1.

$$\frac{-\text{Si}(bx^n) \sin(a) + \text{Ci}(bx^n) \cos(a)}{n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a+b*x^n)/x,x)`

[Out] `1/n*(-Si(b*x^n)*sin(a)+Ci(b*x^n)*cos(a))`

---

**Maxima [F(-2)]** time = 0., size = 0, normalized size = 0.

Exception raised: IndexError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^n)/x,x, algorithm="maxima")`

[Out] Exception raised: IndexError

---

**Fricas [A]** time = 1.89521, size = 131, normalized size = 5.04

$$\frac{\cos(a) \text{Ci}(bx^n) + \cos(a) \text{Ci}(-bx^n) - 2 \sin(a) \text{Si}(bx^n)}{2n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^n)/x,x, algorithm="fricas")`

[Out] `1/2*(cos(a)*cos_integral(b*x^n) + cos(a)*cos_integral(-b*x^n) - 2*sin(a)*sin_integral(b*x^n))/n`

---

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + bx^n)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x**n)/x,x)`

[Out] `Integral(cos(a + b*x**n)/x, x)`

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^n + a)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n)/x,x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^n + a)/x, x)
```

### 3.70 $\int \frac{\cos^2(a+bx^n)}{x} dx$

**Optimal.** Leaf size=43

$$\frac{\cos(2a)\text{CosIntegral}(2bx^n)}{2n} - \frac{\sin(2a)\text{Si}(2bx^n)}{2n} + \frac{\log(x)}{2}$$

[Out] (Cos[2\*a]\*CosIntegral[2\*b\*x^n])/(2\*n) + Log[x]/2 - (Sin[2\*a]\*SinIntegral[2\*b\*x^n])/(2\*n)

**Rubi [A]** time = 0.0579186, antiderivative size = 43, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 4, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3426, 3378, 3376, 3375}

$$\frac{\cos(2a)\text{CosIntegral}(2bx^n)}{2n} - \frac{\sin(2a)\text{Si}(2bx^n)}{2n} + \frac{\log(x)}{2}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^n]^2/x,x]

[Out] (Cos[2\*a]\*CosIntegral[2\*b\*x^n])/(2\*n) + Log[x]/2 - (Sin[2\*a]\*SinIntegral[2\*b\*x^n])/(2\*n)

#### Rule 3426

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_)\*((e\_.)\*(x\_)^(m\_.), x\_Symbol] :> Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]

#### Rule 3378

Int[Cos[(c\_) + (d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Dist[Cos[c], Int[Cos[d\*x^n]/x, x], x] - Dist[Sin[c], Int[Sin[d\*x^n]/x, x], x] /; FreeQ[{c, d, n}, x]

#### Rule 3376

Int[Cos[(d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[CosIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

#### Rule 3375

Int[Sin[(d\_.)\*(x\_)^(n\_)]/(x\_), x\_Symbol] :> Simp[SinIntegral[d\*x^n]/n, x] /; FreeQ[{d, n}, x]

#### Rubi steps

$$\begin{aligned} \int \frac{\cos^2(a+bx^n)}{x} dx &= \int \left( \frac{1}{2x} + \frac{\cos(2a+2bx^n)}{2x} \right) dx \\ &= \frac{\log(x)}{2} + \frac{1}{2} \int \frac{\cos(2a+2bx^n)}{x} dx \\ &= \frac{\log(x)}{2} + \frac{1}{2} \cos(2a) \int \frac{\cos(2bx^n)}{x} dx - \frac{1}{2} \sin(2a) \int \frac{\sin(2bx^n)}{x} dx \\ &= \frac{\cos(2a)\text{Ci}(2bx^n)}{2n} + \frac{\log(x)}{2} - \frac{\sin(2a)\text{Si}(2bx^n)}{2n} \end{aligned}$$



**Mathematica [A]** time = 0.0890335, size = 37, normalized size = 0.86

$$\frac{\cos(2a)\text{CosIntegral}(2bx^n) - \sin(2a)\text{Si}(2bx^n) + n \log(x)}{2n}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^n]^2/x,x]

[Out] (Cos[2\*a]\*CosIntegral[2\*b\*x^n] + n\*Log[x] - Sin[2\*a]\*SinIntegral[2\*b\*x^n])/(2\*n)

**Maple [A]** time = 0.036, size = 45, normalized size = 1.1

$$-\frac{\text{Si}(2bx^n)\sin(2a)}{2n} + \frac{\text{Ci}(2bx^n)\cos(2a)}{2n} + \frac{\ln(bx^n)}{2n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^n)^2/x,x)

[Out] -1/2\*Si(2\*b\*x^n)\*sin(2\*a)/n+1/2\*Ci(2\*b\*x^n)\*cos(2\*a)/n+1/2/n\*ln(b\*x^n)

**Maxima [F(-2)]** time = 0., size = 0, normalized size = 0.

Exception raised: IndexError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^n)^2/x,x, algorithm="maxima")

[Out] Exception raised: IndexError

**Fricas [A]** time = 1.86181, size = 165, normalized size = 3.84

$$\frac{\cos(2a)\text{Ci}(2bx^n) + \cos(2a)\text{Ci}(-2bx^n) + 2n \log(x) - 2\sin(2a)\text{Si}(2bx^n)}{4n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^n)^2/x,x, algorithm="fricas")

[Out] 1/4\*(cos(2\*a)\*cos\_integral(2\*b\*x^n) + cos(2\*a)\*cos\_integral(-2\*b\*x^n) + 2\*n\*log(x) - 2\*sin(2\*a)\*sin\_integral(2\*b\*x^n))/n

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^2(a + bx^n)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x\*\*n)\*\*2/x,x)

[Out] Integral(cos(a + b\*x\*\*n)\*\*2/x, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^n + a)^2}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^n)^2/x,x, algorithm="giac")

[Out] integrate(cos(b\*x^n + a)^2/x, x)

### 3.71 $\int \frac{\cos^3(a+bx^n)}{x} dx$

**Optimal.** Leaf size=67

$$\frac{3 \cos(a) \operatorname{CosIntegral}(bx^n)}{4n} + \frac{\cos(3a) \operatorname{CosIntegral}(3bx^n)}{4n} - \frac{3 \sin(a) \operatorname{Si}(bx^n)}{4n} - \frac{\sin(3a) \operatorname{Si}(3bx^n)}{4n}$$

```
[Out] (3*Cos[a]*CosIntegral[b*x^n])/(4*n) + (Cos[3*a]*CosIntegral[3*b*x^n])/(4*n)
- (3*Sin[a]*SinIntegral[b*x^n])/(4*n) - (Sin[3*a]*SinIntegral[3*b*x^n])/(4
*n)
```

**Rubi [A]** time = 0.0860536, antiderivative size = 67, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 4, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3426, 3378, 3376, 3375}

$$\frac{3 \cos(a) \operatorname{CosIntegral}(bx^n)}{4n} + \frac{\cos(3a) \operatorname{CosIntegral}(3bx^n)}{4n} - \frac{3 \sin(a) \operatorname{Si}(bx^n)}{4n} - \frac{\sin(3a) \operatorname{Si}(3bx^n)}{4n}$$

Antiderivative was successfully verified.

```
[In] Int[Cos[a + b*x^n]^3/x, x]
```

```
[Out] (3*Cos[a]*CosIntegral[b*x^n])/(4*n) + (Cos[3*a]*CosIntegral[3*b*x^n])/(4*n)
- (3*Sin[a]*SinIntegral[b*x^n])/(4*n) - (Sin[3*a]*SinIntegral[3*b*x^n])/(4
*n)
```

#### Rule 3426

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_)*((e_.)*(x_)^(m_.), x
_Symbol] :> Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x]
/; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]
```

#### Rule 3378

```
Int[Cos[(c_) + (d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Dist[Cos[c], Int[Cos[d*x
^n]/x, x], x] - Dist[Sin[c], Int[Sin[d*x^n]/x, x], x] /; FreeQ[{c, d, n}, x
]
```

#### Rule 3376

```
Int[Cos[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[CosIntegral[d*x^n]/n, x] /
; FreeQ[{d, n}, x]
```

#### Rule 3375

```
Int[Sin[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[SinIntegral[d*x^n]/n, x] /
; FreeQ[{d, n}, x]
```

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos^3(a + bx^n)}{x} dx &= \int \left( \frac{3 \cos(a + bx^n)}{4x} + \frac{\cos(3a + 3bx^n)}{4x} \right) dx \\
&= \frac{1}{4} \int \frac{\cos(3a + 3bx^n)}{x} dx + \frac{3}{4} \int \frac{\cos(a + bx^n)}{x} dx \\
&= \frac{1}{4} (3 \cos(a)) \int \frac{\cos(bx^n)}{x} dx + \frac{1}{4} \cos(3a) \int \frac{\cos(3bx^n)}{x} dx - \frac{1}{4} (3 \sin(a)) \int \frac{\sin(bx^n)}{x} dx - \frac{1}{4} \sin(3a) \int \frac{\sin(3bx^n)}{x} dx \\
&= \frac{3 \cos(a) \text{Ci}(bx^n)}{4n} + \frac{\cos(3a) \text{Ci}(3bx^n)}{4n} - \frac{3 \sin(a) \text{Si}(bx^n)}{4n} - \frac{\sin(3a) \text{Si}(3bx^n)}{4n}
\end{aligned}$$

**Mathematica [A]** time = 0.121043, size = 53, normalized size = 0.79

$$\frac{3 \cos(a) \text{CosIntegral}(bx^n) + \cos(3a) \text{CosIntegral}(3bx^n) - 3 \sin(a) \text{Si}(bx^n) - \sin(3a) \text{Si}(3bx^n)}{4n}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^n]^3/x, x]

[Out] (3\*Cos[a]\*CosIntegral[b\*x^n] + Cos[3\*a]\*CosIntegral[3\*b\*x^n] - 3\*Sin[a]\*SinIntegral[b\*x^n] - Sin[3\*a]\*SinIntegral[3\*b\*x^n])/(4\*n)

**Maple [A]** time = 0.036, size = 52, normalized size = 0.8

$$\frac{1}{n} \left( -\frac{\text{Si}(3bx^n) \sin(3a)}{4} + \frac{\text{Ci}(3bx^n) \cos(3a)}{4} - \frac{3 \text{Si}(bx^n) \sin(a)}{4} + \frac{3 \text{Ci}(bx^n) \cos(a)}{4} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^n)^3/x, x)

[Out] 1/n\*(-1/4\*Si(3\*b\*x^n)\*sin(3\*a)+1/4\*Ci(3\*b\*x^n)\*cos(3\*a)-3/4\*Si(b\*x^n)\*sin(a)+3/4\*Ci(b\*x^n)\*cos(a))

**Maxima [F(-2)]** time = 0., size = 0, normalized size = 0.

Exception raised: IndexError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^n)^3/x, x, algorithm="maxima")

[Out] Exception raised: IndexError

**Fricas [A]** time = 2.00628, size = 274, normalized size = 4.09

$$\frac{\cos(3a) \text{Ci}(3bx^n) + 3 \cos(a) \text{Ci}(bx^n) + 3 \cos(a) \text{Ci}(-bx^n) + \cos(3a) \text{Ci}(-3bx^n) - 2 \sin(3a) \text{Si}(3bx^n) - 6 \sin(a) \text{Si}(bx^n)}{8n}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n)^3/x,x, algorithm="fricas")
```

```
[Out] 1/8*(cos(3*a)*cos_integral(3*b*x^n) + 3*cos(a)*cos_integral(b*x^n) + 3*cos(a)*cos_integral(-b*x^n) + cos(3*a)*cos_integral(-3*b*x^n) - 2*sin(3*a)*sin_integral(3*b*x^n) - 6*sin(a)*sin_integral(b*x^n))/n
```

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^3(a + bx^n)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x**n)**3/x,x)
```

```
[Out] Integral(cos(a + b*x**n)**3/x, x)
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^n + a)^3}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n)^3/x,x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^n + a)^3/x, x)
```

### 3.72 $\int \frac{\cos^4(a+bx^n)}{x} dx$

**Optimal.** Leaf size=79

$$\frac{\cos(2a)\text{CosIntegral}(2bx^n)}{2n} + \frac{\cos(4a)\text{CosIntegral}(4bx^n)}{8n} - \frac{\sin(2a)\text{Si}(2bx^n)}{2n} - \frac{\sin(4a)\text{Si}(4bx^n)}{8n} + \frac{3\log(x)}{8}$$

[Out] (Cos[2\*a]\*CosIntegral[2\*b\*x^n])/(2\*n) + (Cos[4\*a]\*CosIntegral[4\*b\*x^n])/(8\*n) + (3\*Log[x])/8 - (Sin[2\*a]\*SinIntegral[2\*b\*x^n])/(2\*n) - (Sin[4\*a]\*SinIntegral[4\*b\*x^n])/(8\*n)

**Rubi [A]** time = 0.0940672, antiderivative size = 79, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 4, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3426, 3378, 3376, 3375}

$$\frac{\cos(2a)\text{CosIntegral}(2bx^n)}{2n} + \frac{\cos(4a)\text{CosIntegral}(4bx^n)}{8n} - \frac{\sin(2a)\text{Si}(2bx^n)}{2n} - \frac{\sin(4a)\text{Si}(4bx^n)}{8n} + \frac{3\log(x)}{8}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^n]^4/x,x]

[Out] (Cos[2\*a]\*CosIntegral[2\*b\*x^n])/(2\*n) + (Cos[4\*a]\*CosIntegral[4\*b\*x^n])/(8\*n) + (3\*Log[x])/8 - (Sin[2\*a]\*SinIntegral[2\*b\*x^n])/(2\*n) - (Sin[4\*a]\*SinIntegral[4\*b\*x^n])/(8\*n)

#### Rule 3426

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x
_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x]
/; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]
```

#### Rule 3378

```
Int[Cos[(c_) + (d_.)*(x_)^(n_)]/(x_), x_Symbol] := Dist[Cos[c], Int[Cos[d*x
^n]/x, x], x] - Dist[Sin[c], Int[Sin[d*x^n]/x, x], x] /; FreeQ[{c, d, n}, x
]
```

#### Rule 3376

```
Int[Cos[(d_.)*(x_)^(n_)]/(x_), x_Symbol] := Simp[CosIntegral[d*x^n]/n, x] /
; FreeQ[{d, n}, x]
```

#### Rule 3375

```
Int[Sin[(d_.)*(x_)^(n_)]/(x_), x_Symbol] := Simp[SinIntegral[d*x^n]/n, x] /
; FreeQ[{d, n}, x]
```

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos^4(a + bx^n)}{x} dx &= \int \left( \frac{3}{8x} + \frac{\cos(2a + 2bx^n)}{2x} + \frac{\cos(4a + 4bx^n)}{8x} \right) dx \\
&= \frac{3 \log(x)}{8} + \frac{1}{8} \int \frac{\cos(4a + 4bx^n)}{x} dx + \frac{1}{2} \int \frac{\cos(2a + 2bx^n)}{x} dx \\
&= \frac{3 \log(x)}{8} + \frac{1}{2} \cos(2a) \int \frac{\cos(2bx^n)}{x} dx + \frac{1}{8} \cos(4a) \int \frac{\cos(4bx^n)}{x} dx - \frac{1}{2} \sin(2a) \int \frac{\sin(2bx^n)}{x} dx \\
&= \frac{\cos(2a)\text{Ci}(2bx^n)}{2n} + \frac{\cos(4a)\text{Ci}(4bx^n)}{8n} + \frac{3 \log(x)}{8} - \frac{\sin(2a)\text{Si}(2bx^n)}{2n} - \frac{\sin(4a)\text{Si}(4bx^n)}{8n}
\end{aligned}$$

**Mathematica [A]** time = 0.14085, size = 66, normalized size = 0.84

$$\frac{4 \cos(2a)\text{CosIntegral}(2bx^n) + \cos(4a)\text{CosIntegral}(4bx^n) - 4 \sin(2a)\text{Si}(2bx^n) - \sin(4a)\text{Si}(4bx^n)}{8n} + \frac{3 \log(x)}{8}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^n]^4/x,x]

[Out] (3\*Log[x])/8 + (4\*Cos[2\*a]\*CosIntegral[2\*b\*x^n] + Cos[4\*a]\*CosIntegral[4\*b\*x^n] - 4\*Sin[2\*a]\*SinIntegral[2\*b\*x^n] - Sin[4\*a]\*SinIntegral[4\*b\*x^n])/(8\*n)

**Maple [A]** time = 0.039, size = 77, normalized size = 1.

$$-\frac{\text{Si}(4bx^n)\sin(4a)}{8n} + \frac{\text{Ci}(4bx^n)\cos(4a)}{8n} - \frac{\text{Si}(2bx^n)\sin(2a)}{2n} + \frac{\text{Ci}(2bx^n)\cos(2a)}{2n} + \frac{3 \ln(bx^n)}{8n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^n)^4/x,x)

[Out] -1/8\*Si(4\*b\*x^n)\*sin(4\*a)/n+1/8\*Ci(4\*b\*x^n)\*cos(4\*a)/n-1/2\*Si(2\*b\*x^n)\*sin(2\*a)/n+1/2\*Ci(2\*b\*x^n)\*cos(2\*a)/n+3/8/n\*ln(b\*x^n)

**Maxima [F(-2)]** time = 0., size = 0, normalized size = 0.

Exception raised: IndexError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^n)^4/x,x, algorithm="maxima")

[Out] Exception raised: IndexError

**Fricas [A]** time = 2.02452, size = 309, normalized size = 3.91

$$\frac{\cos(4a)\text{Ci}(4bx^n) + 4 \cos(2a)\text{Ci}(2bx^n) + 4 \cos(2a)\text{Ci}(-2bx^n) + \cos(4a)\text{Ci}(-4bx^n) + 6n \log(x) - 2 \sin(4a)\text{Si}(4bx^n)}{16n}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n)^4/x,x, algorithm="fricas")
```

```
[Out] 1/16*(cos(4*a)*cos_integral(4*b*x^n) + 4*cos(2*a)*cos_integral(2*b*x^n) + 4
*cos(2*a)*cos_integral(-2*b*x^n) + cos(4*a)*cos_integral(-4*b*x^n) + 6*n*log
g(x) - 2*sin(4*a)*sin_integral(4*b*x^n) - 8*sin(2*a)*sin_integral(2*b*x^n))
/n
```

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos^4(a + bx^n)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x**n)**4/x,x)
```

```
[Out] Integral(cos(a + b*x**n)**4/x, x)
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(bx^n + a)^4}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n)^4/x,x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^n + a)^4/x, x)
```



### 3.73 $\int \cos(a + bx^n) dx$

**Optimal.** Leaf size=83

$$\frac{e^{ia}x(-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia}x(ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, ibx^n\right)}{2n}$$

[Out]  $-(E^{I*a}*x*\Gamma[n^{(-1)}, (-I)*b*x^n])/(2*n*((-I)*b*x^n)^n^{(-1)}) - (x*\Gamma[n^{(-1)}, I*b*x^n])/(2*E^{I*a}*n*(I*b*x^n)^n^{(-1)})$

**Rubi [A]** time = 0.0216957, antiderivative size = 83, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 2, integrand size = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3366, 2208}

$$\frac{e^{ia}x(-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia}x(ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, ibx^n\right)}{2n}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^n], x]

[Out]  $-(E^{I*a}*x*\Gamma[n^{(-1)}, (-I)*b*x^n])/(2*n*((-I)*b*x^n)^n^{(-1)}) - (x*\Gamma[n^{(-1)}, I*b*x^n])/(2*E^{I*a}*n*(I*b*x^n)^n^{(-1)})$

#### Rule 3366

Int[Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_)^n)], x\_Symbol] := Dist[1/2, Int[E^(-(c\*I) - d\*I\*(e + f\*x)^n), x], x] + Dist[1/2, Int[E^(c\*I + d\*I\*(e + f\*x)^n), x], x] /; FreeQ[{c, d, e, f, n}, x]

#### Rule 2208

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_)^n)), x\_Symbol] := -Simp[(F^a\*(c + d\*x)\*Gamma[1/n, -(b\*(c + d\*x)^n\*Log[F]])]/(d\*n\*(-(b\*(c + d\*x)^n\*Log[F]))^(1/n)), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]

#### Rubi steps

$$\begin{aligned} \int \cos(a + bx^n) dx &= \frac{1}{2} \int e^{-ia-ibx^n} dx + \frac{1}{2} \int e^{ia+ibx^n} dx \\ &= -\frac{e^{ia}x(-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia}x(ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, ibx^n\right)}{2n} \end{aligned}$$

**Mathematica [A]** time = 0.0900201, size = 92, normalized size = 1.11

$$\frac{x(b^2x^{2n})^{-1/n} \left( (\cos(a) - i \sin(a)) (-ibx^n)^{\frac{1}{n}} \Gamma\left(\frac{1}{n}, ibx^n\right) + (\cos(a) + i \sin(a)) (ibx^n)^{\frac{1}{n}} \Gamma\left(\frac{1}{n}, -ibx^n\right) \right)}{2n}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^n], x]

[Out]  $-(x*(((-I)*b*x^n)^n)^{-1}*\Gamma[n^{-1}], I*b*x^n*(\cos[a] - I*\sin[a]) + (I*b*x^n)^n)^{-1}*\Gamma[n^{-1}], (-I)*b*x^n*(\cos[a] + I*\sin[a]))/(2*n*(b^2*x^{(2*n)})^n)^{-1})$

**Maple [C]** time = 0.092, size = 75, normalized size = 0.9

$$x {}_1F_2\left(\frac{1}{2n}; \frac{1}{2}, 1 + \frac{1}{2n}; -\frac{x^{2n}b^2}{4}\right) \cos(a) - \frac{bx^{1+n} \sin(a)}{1+n} {}_1F_2\left(\frac{1}{2} + \frac{1}{2n}; \frac{3}{2}, \frac{3}{2} + \frac{1}{2n}; -\frac{x^{2n}b^2}{4}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a+b*x^n),x)`

[Out] `x*hypergeom([1/2/n],[1/2,1+1/2/n],-1/4*x^(2*n)*b^2)*cos(a)-b/(1+n)*x^(1+n)*hypergeom([1/2+1/2/n],[3/2,3/2+1/2/n],-1/4*x^(2*n)*b^2)*sin(a)`

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos(bx^n + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^n),x, algorithm="maxima")`

[Out] `integrate(cos(b*x^n + a), x)`

**Fricas [F]** time = 0., size = 0, normalized size = 0.

$$\text{integral}(\cos(bx^n + a), x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x^n),x, algorithm="fricas")`

[Out] `integral(cos(b*x^n + a), x)`

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos(a + bx^n) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(a+b*x**n),x)`

[Out] `Integral(cos(a + b*x**n), x)`

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos(bx^n + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n),x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^n + a), x)
```

### 3.74 $\int \cos^2(a + bx^n) dx$

**Optimal.** Leaf size=102

$$\frac{e^{2ia} 2^{-\frac{1}{n}-2} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -2ibx^n\right)}{n} - \frac{e^{-2ia} 2^{-\frac{1}{n}-2} x (ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, 2ibx^n\right)}{n} + \frac{x}{2}$$

[Out]  $x/2 - (2^{(-2 - n^{(-1)})} E^{((2*I)*a)} * x * \Gamma[n^{(-1)}, (-2*I)*b*x^n]) / (n * ((-I)*b*x^n)^{n^{(-1)}}) - (2^{(-2 - n^{(-1)})} * x * \Gamma[n^{(-1)}, (2*I)*b*x^n]) / (E^{((2*I)*a)} * n * (I*b*x^n)^{n^{(-1)}})$

**Rubi [A]** time = 0.0690008, antiderivative size = 102, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 3, integrand size = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.3$ , Rules used = {3368, 3366, 2208}

$$\frac{e^{2ia} 2^{-\frac{1}{n}-2} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -2ibx^n\right)}{n} - \frac{e^{-2ia} 2^{-\frac{1}{n}-2} x (ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, 2ibx^n\right)}{n} + \frac{x}{2}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^n]^2, x]

[Out]  $x/2 - (2^{(-2 - n^{(-1)})} E^{((2*I)*a)} * x * \Gamma[n^{(-1)}, (-2*I)*b*x^n]) / (n * ((-I)*b*x^n)^{n^{(-1)}}) - (2^{(-2 - n^{(-1)})} * x * \Gamma[n^{(-1)}, (2*I)*b*x^n]) / (E^{((2*I)*a)} * n * (I*b*x^n)^{n^{(-1)}})$

#### Rule 3368

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)])\*(b\_.))^(p\_), x\_Symbol] :> Int[ExpandTrigReduce[(a + b\*Cos[c + d\*(e + f\*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 1]

#### Rule 3366

Int[Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)], x\_Symbol] :> Dist[1/2, Int[E^(-(c\*I) - d\*I\*(e + f\*x)^n), x], x] + Dist[1/2, Int[E^(c\*I + d\*I\*(e + f\*x)^n), x], x] /; FreeQ[{c, d, e, f, n}, x]

#### Rule 2208

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_))^(n\_)), x\_Symbol] :> -Simp[(F^a \* (c + d\*x) \* Gamma[1/n, -(b\*(c + d\*x)^n \* Log[F]])] / (d \* n \* (-(b\*(c + d\*x)^n \* Log[F]))^(1/n)), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]

#### Rubi steps

$$\begin{aligned} \int \cos^2(a + bx^n) dx &= \int \left( \frac{1}{2} + \frac{1}{2} \cos(2a + 2bx^n) \right) dx \\ &= \frac{x}{2} + \frac{1}{2} \int \cos(2a + 2bx^n) dx \\ &= \frac{x}{2} + \frac{1}{4} \int e^{-2ia - 2ibx^n} dx + \frac{1}{4} \int e^{2ia + 2ibx^n} dx \\ &= \frac{x}{2} - \frac{2^{-2-\frac{1}{n}} e^{2ia} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -2ibx^n\right)}{n} - \frac{2^{-2-\frac{1}{n}} e^{-2ia} x (ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, 2ibx^n\right)}{n} \end{aligned}$$

**Mathematica [A]** time = 0.246937, size = 94, normalized size = 0.92

$$\frac{x \left( e^{2ia} 2^{-1/n} (-ibx^n)^{-1/n} \text{Gamma} \left( \frac{1}{n}, -2ibx^n \right) + e^{-2ia} 2^{-1/n} (ibx^n)^{-1/n} \text{Gamma} \left( \frac{1}{n}, 2ibx^n \right) - 2n \right)}{4n}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^n]^2,x]

[Out]  $-(x*(-2*n + (E^{((2*I)*a)}*Gamma[n^{(-1)}, (-2*I)*b*x^n])/(2^n^{(-1)}*((-I)*b*x^n)^n^{(-1)) + Gamma[n^{(-1)}, (2*I)*b*x^n]/(2^n^{(-1)}*E^{((2*I)*a)}*(I*b*x^n)^n^{(-1)})))/(4*n)$

**Maple [F]** time = 0.165, size = 0, normalized size = 0.

$$\int (\cos(a + bx^n))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^n)^2,x)

[Out] int(cos(a+b\*x^n)^2,x)

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\frac{1}{2}x + \frac{1}{2} \int \cos(2bx^n + 2a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^n)^2,x, algorithm="maxima")

[Out] 1/2\*x + 1/2\*integrate(cos(2\*b\*x^n + 2\*a), x)

**Fricas [F]** time = 0., size = 0, normalized size = 0.

$$\text{integral}(\cos(bx^n + a)^2, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^n)^2,x, algorithm="fricas")

[Out] integral(cos(b\*x^n + a)^2, x)

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos^2(a + bx^n) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x**n)**2,x)
```

```
[Out] Integral(cos(a + b*x**n)**2, x)
```

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos(bx^n + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n)^2,x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^n + a)^2, x)
```

### 3.75 $\int \cos^3(a + bx^n) dx$

**Optimal.** Leaf size=179

$$\frac{3e^{ia}x(-ibx^n)^{-1/n}\Gamma\left(\frac{1}{n}, -ibx^n\right)}{8n} - \frac{e^{3ia}3^{-1/n}x(-ibx^n)^{-1/n}\Gamma\left(\frac{1}{n}, -3ibx^n\right)}{8n} - \frac{3e^{-ia}x(ibx^n)^{-1/n}\Gamma\left(\frac{1}{n}, ibx^n\right)}{8n}$$

[Out]  $(-3E^{(I*a)}*x*\Gamma[n^{(-1)}, (-I)*b*x^n])/(8*n*((-I)*b*x^n)^{n^{(-1)}}) - (3*x*\Gamma[n^{(-1)}, I*b*x^n])/(8*E^{(I*a)}*n*(I*b*x^n)^{n^{(-1)}}) - (E^{((3*I)*a)}*x*\Gamma[a[n^{(-1)}, (-3*I)*b*x^n])/(8*3^n^{(-1)}*n*((-I)*b*x^n)^{n^{(-1)}}) - (x*\Gamma[n^{(-1)}, (3*I)*b*x^n])/(8*3^n^{(-1)}*E^{((3*I)*a)}*n*(I*b*x^n)^{n^{(-1)}})$

**Rubi [A]** time = 0.0789607, antiderivative size = 179, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 3, integrand size = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.3$ , Rules used = {3368, 3366, 2208}

$$\frac{3e^{ia}x(-ibx^n)^{-1/n}\Gamma\left(\frac{1}{n}, -ibx^n\right)}{8n} - \frac{e^{3ia}3^{-1/n}x(-ibx^n)^{-1/n}\Gamma\left(\frac{1}{n}, -3ibx^n\right)}{8n} - \frac{3e^{-ia}x(ibx^n)^{-1/n}\Gamma\left(\frac{1}{n}, ibx^n\right)}{8n}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*x^n]^3, x]

[Out]  $(-3E^{(I*a)}*x*\Gamma[n^{(-1)}, (-I)*b*x^n])/(8*n*((-I)*b*x^n)^{n^{(-1)}}) - (3*x*\Gamma[n^{(-1)}, I*b*x^n])/(8*E^{(I*a)}*n*(I*b*x^n)^{n^{(-1)}}) - (E^{((3*I)*a)}*x*\Gamma[a[n^{(-1)}, (-3*I)*b*x^n])/(8*3^n^{(-1)}*n*((-I)*b*x^n)^{n^{(-1)}}) - (x*\Gamma[n^{(-1)}, (3*I)*b*x^n])/(8*3^n^{(-1)}*E^{((3*I)*a)}*n*(I*b*x^n)^{n^{(-1)}})$

#### Rule 3368

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)])\*(b\_.))^(p\_), x\_Symbol] := Int[ExpandTrigReduce[(a + b\*Cos[c + d\*(e + f\*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 1]

#### Rule 3366

Int[Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)], x\_Symbol] := Dist[1/2, Int[E^(-c\*I) - d\*I\*(e + f\*x)^n], x] + Dist[1/2, Int[E^(c\*I + d\*I\*(e + f\*x)^n), x], x] /; FreeQ[{c, d, e, f, n}, x]

#### Rule 2208

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_))^(n\_)), x\_Symbol] := -Simp[(F^a\*(c + d\*x)\*Gamma[1/n, -(b\*(c + d\*x)^n\*Log[F]])]/(d\*n\*(-(b\*(c + d\*x)^n\*Log[F]))^(1/n)), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]

#### Rubi steps

$$\begin{aligned}
\int \cos^3(a + bx^n) dx &= \int \left( \frac{3}{4} \cos(a + bx^n) + \frac{1}{4} \cos(3a + 3bx^n) \right) dx \\
&= \frac{1}{4} \int \cos(3a + 3bx^n) dx + \frac{3}{4} \int \cos(a + bx^n) dx \\
&= \frac{1}{8} \int e^{-3ia-3ibx^n} dx + \frac{1}{8} \int e^{3ia+3ibx^n} dx + \frac{3}{8} \int e^{-ia-ibx^n} dx + \frac{3}{8} \int e^{ia+ibx^n} dx \\
&= \frac{3e^{ia} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -ibx^n\right)}{8n} - \frac{3e^{-ia} x (ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, ibx^n\right)}{8n} - \frac{3^{-1/n} e^{3ia} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -3ibx^n\right)}{8n}
\end{aligned}$$

**Mathematica [A]** time = 0.239687, size = 173, normalized size = 0.97

$$\frac{e^{-3ia} 3^{-1/n} x (b^2 x^{2n})^{-1/n} \left( e^{2ia} 3^{\frac{1}{n}+1} (-ibx^n)^{\frac{1}{n}} \text{Gamma}\left(\frac{1}{n}, ibx^n\right) + e^{4ia} 3^{\frac{1}{n}+1} (ibx^n)^{\frac{1}{n}} \text{Gamma}\left(\frac{1}{n}, -ibx^n\right) + e^{6ia} (ibx^n)^{\frac{1}{n}} \text{Gamma}\left(\frac{1}{n}, -3ibx^n\right) \right)}{8n}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*x^n]^3, x]

[Out]  $-(x*(3^{(1+n^{-1})})*E^{((4*I)*a)*(I*b*x^n)^{n^{-1}}*\text{Gamma}[n^{-1}, (-I)*b*x^n]} + 3^{(1+n^{-1})}*E^{((2*I)*a)*((-I)*b*x^n)^{n^{-1}}*\text{Gamma}[n^{-1}, I*b*x^n]} + E^{((6*I)*a)*(I*b*x^n)^{n^{-1}}*\text{Gamma}[n^{-1}, (-3*I)*b*x^n]} + ((-I)*b*x^n)^{n^{-1}}*\text{Gamma}[n^{-1}, (3*I)*b*x^n]))/(8*3^{n^{-1}}*E^{((3*I)*a)*n*(b^2*x^{(2*n)})^{n^{-1}}})$

**Maple [F]** time = 180., size = 0, normalized size = 0.

$$\int (\cos(a + bx^n))^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*x^n)^3, x)

[Out] int(cos(a+b\*x^n)^3, x)

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos(bx^n + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*x^n)^3, x, algorithm="maxima")

[Out] integrate(cos(b\*x^n + a)^3, x)

**Fricas [F]** time = 0., size = 0, normalized size = 0.

$$\text{integral}(\cos(bx^n + a)^3, x)$$



Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n)^3,x, algorithm="fricas")
```

```
[Out] integral(cos(b*x^n + a)^3, x)
```

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos^3(a + bx^n) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x**n)**3,x)
```

```
[Out] Integral(cos(a + b*x**n)**3, x)
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos(bx^n + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(a+b*x^n)^3,x, algorithm="giac")
```

```
[Out] integrate(cos(b*x^n + a)^3, x)
```

### 3.76 $\int x^m \cos(a + bx^n) dx$

**Optimal.** Leaf size=105

$$-\frac{e^{ia}x^{m+1}(-ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia}x^{m+1}(ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, ibx^n\right)}{2n}$$

[Out]  $-(E^{(I*a)}*x^{(1+m)}*\Gamma[(1+m)/n, (-I)*b*x^n])/(2*n*((-I)*b*x^n)^{((1+m)/n)}) - (x^{(1+m)}*\Gamma[(1+m)/n, I*b*x^n])/(2*E^{(I*a)}*n*(I*b*x^n)^{((1+m)/n)})$

**Rubi [A]** time = 0.0748536, antiderivative size = 105, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 2, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3424, 2218}

$$-\frac{e^{ia}x^{m+1}(-ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia}x^{m+1}(ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, ibx^n\right)}{2n}$$

Antiderivative was successfully verified.

[In] Int[x^m\*Cos[a + b\*x^n], x]

[Out]  $-(E^{(I*a)}*x^{(1+m)}*\Gamma[(1+m)/n, (-I)*b*x^n])/(2*n*((-I)*b*x^n)^{((1+m)/n)}) - (x^{(1+m)}*\Gamma[(1+m)/n, I*b*x^n])/(2*E^{(I*a)}*n*(I*b*x^n)^{((1+m)/n)})$

#### Rule 3424

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_.)]\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] :> Dist[1/2, Int[(e\*x)^m\*E^(-(c\*I) - d\*I\*x^n), x], x] + Dist[1/2, Int[(e\*x)^m\*E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m, n}, x]

#### Rule 2218

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_))^(n\_.))\*((e\_.) + (f\_.)\*(x\_))^(m\_.), x\_Symbol] :> -Simp[(F^a\*(e + f\*x)^(m + 1)\*Gamma[(m + 1)/n, -(b\*(c + d\*x)^(n)\*Log[F])])/(f\*n\*(-(b\*(c + d\*x)^(n)\*Log[F]))^((m + 1)/n)), x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d\*e - c\*f, 0]

#### Rubi steps

$$\begin{aligned} \int x^m \cos(a + bx^n) dx &= \frac{1}{2} \int e^{-ia-ibx^n} x^m dx + \frac{1}{2} \int e^{ia+ibx^n} x^m dx \\ &= -\frac{e^{ia}x^{1+m}(-ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia}x^{1+m}(ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, ibx^n\right)}{2n} \end{aligned}$$

**Mathematica [A]** time = 0.200067, size = 115, normalized size = 1.1

$$\frac{x^{m+1} (b^2 x^{2n})^{-\frac{m+1}{n}} \left( (\cos(a) - i \sin(a)) (-ibx^n)^{\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, ibx^n\right) + (\cos(a) + i \sin(a)) (ibx^n)^{\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -ibx^n\right) \right)}{2n}$$

Antiderivative was successfully verified.

[In] Integrate[x^m\*cos[a + b\*x^n],x]

[Out]  $-(x^{(1+m)}*((-I)*b*x^n)^{((1+m)/n)}*\Gamma[(1+m)/n, I*b*x^n]*(\cos[a] - I*\sin[a]) + (I*b*x^n)^{((1+m)/n)}*\Gamma[(1+m)/n, (-I)*b*x^n]*(\cos[a] + I*\sin[a]))/(2*n*(b^2*x^{(2*n)})^{((1+m)/n)})$

**Maple [C]** time = 0.162, size = 111, normalized size = 1.1

$$\frac{x^{1+m} \cos(a)}{1+m} {}_1F_2\left(\frac{m}{2n} + \frac{1}{2n}; \frac{1}{2}, 1 + \frac{m}{2n} + \frac{1}{2n}; -\frac{x^{2n}b^2}{4}\right) - \frac{bx^{m+n+1} \sin(a)}{m+n+1} {}_1F_2\left(\frac{1}{2} + \frac{m}{2n} + \frac{1}{2n}; \frac{3}{2}, \frac{3}{2} + \frac{m}{2n} + \frac{1}{2n}; -\frac{x^{2n}b^2}{4}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m\*cos(a+b\*x^n),x)

[Out]  $1/(1+m)*x^{(1+m)}*\text{hypergeom}([1/2/n*m+1/2/n], [1/2, 1+1/2/n*m+1/2/n], -1/4*x^{(2*n)}*b^2)*\cos(a) - b/(m+n+1)*x^{(m+n+1)}*\text{hypergeom}([1/2+1/2/n*m+1/2/n], [3/2, 3/2+1/2/n*m+1/2/n], -1/4*x^{(2*n)}*b^2)*\sin(a)$

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int x^m \cos(bx^n + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m\*cos(a+b\*x^n),x, algorithm="maxima")

[Out] integrate(x^m\*cos(b\*x^n + a), x)

**Fricas [F]** time = 0., size = 0, normalized size = 0.

$$\text{integral}(x^m \cos(bx^n + a), x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m\*cos(a+b\*x^n),x, algorithm="fricas")

[Out] integral(x^m\*cos(b\*x^n + a), x)

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^m \cos(a + bx^n) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*m\*cos(a+b\*x\*\*n),x)

[Out] Integral( $x^m \cos(a + b x^n)$ , x)

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^m \cos(bx^n + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate( $x^m \cos(a + b x^n)$ , x, algorithm="giac")

[Out] integrate( $x^m \cos(b x^n + a)$ , x)

### 3.77 $\int x^m \cos^2(a + bx^n) dx$

**Optimal.** Leaf size=141

$$\frac{e^{2ia} 2^{-\frac{m+2n+1}{n}} x^{m+1} (-ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -2ibx^n\right)}{n} - \frac{e^{-2ia} 2^{-\frac{m+2n+1}{n}} x^{m+1} (ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, 2ibx^n\right)}{n} + \frac{x^m}{2(m+1)}$$

[Out]  $x^{(1+m)/(2*(1+m))} - (E^{((2*I)*a)} * x^{(1+m)} * \Gamma[(1+m)/n, (-2*I)*b*x^n]) / (2^{((1+m+2*n)/n)} * n * ((-I)*b*x^n)^{((1+m)/n)}) - (x^{(1+m)} * \Gamma[(1+m)/n, (2*I)*b*x^n]) / (2^{((1+m+2*n)/n)} * E^{((2*I)*a)} * n * (I*b*x^n)^{((1+m)/n)})$

**Rubi [A]** time = 0.160436, antiderivative size = 141, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3426, 3424, 2218}

$$\frac{e^{2ia} 2^{-\frac{m+2n+1}{n}} x^{m+1} (-ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -2ibx^n\right)}{n} - \frac{e^{-2ia} 2^{-\frac{m+2n+1}{n}} x^{m+1} (ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, 2ibx^n\right)}{n} + \frac{x^m}{2(m+1)}$$

Antiderivative was successfully verified.

[In] Int[x^m \* Cos[a + b\*x^n]^2, x]

[Out]  $x^{(1+m)/(2*(1+m))} - (E^{((2*I)*a)} * x^{(1+m)} * \Gamma[(1+m)/n, (-2*I)*b*x^n]) / (2^{((1+m+2*n)/n)} * n * ((-I)*b*x^n)^{((1+m)/n)}) - (x^{(1+m)} * \Gamma[(1+m)/n, (2*I)*b*x^n]) / (2^{((1+m+2*n)/n)} * E^{((2*I)*a)} * n * (I*b*x^n)^{((1+m)/n)})$

#### Rule 3426

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.))^(p\_)\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] := Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]

#### Rule 3424

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)]\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] := Dist[1/2, Int[(e\*x)^m \* E^(-(c\*I) - d\*I\*x^n), x], x] + Dist[1/2, Int[(e\*x)^m \* E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m, n}, x]

#### Rule 2218

Int[(F\_)^(a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_)^(n\_))\*((e\_.) + (f\_.)\*(x\_))^(m\_.), x\_Symbol] := -Simp[(F^a\*(e + f\*x)^(m+1)\*Gamma[(m+1)/n, -(b\*(c + d\*x)^n \* Log[F]])]/(f\*n\*(-(b\*(c + d\*x)^n \* Log[F]))^((m+1)/n)), x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d\*e - c\*f, 0]

#### Rubi steps

$$\begin{aligned}
\int x^m \cos^2(a + bx^n) dx &= \int \left( \frac{x^m}{2} + \frac{1}{2} x^m \cos(2a + 2bx^n) \right) dx \\
&= \frac{x^{1+m}}{2(1+m)} + \frac{1}{2} \int x^m \cos(2a + 2bx^n) dx \\
&= \frac{x^{1+m}}{2(1+m)} + \frac{1}{4} \int e^{-2ia-2ibx^n} x^m dx + \frac{1}{4} \int e^{2ia+2ibx^n} x^m dx \\
&= \frac{x^{1+m}}{2(1+m)} - \frac{2^{-\frac{1+m+2n}{n}} e^{2ia} x^{1+m} (-ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -2ibx^n\right)}{n} - \frac{2^{-\frac{1+m+2n}{n}} e^{-2ia} x^{1+m} (ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, 2ibx^n\right)}{n}
\end{aligned}$$

**Mathematica [A]** time = 0.51575, size = 129, normalized size = 0.91

$$\frac{x^{m+1} \left( e^{2ia} (m+1) 2^{-\frac{m+1}{n}} (-ibx^n)^{-\frac{m+1}{n}} \text{Gamma}\left(\frac{m+1}{n}, -2ibx^n\right) + e^{-2ia} (m+1) 2^{-\frac{m+1}{n}} (ibx^n)^{-\frac{m+1}{n}} \text{Gamma}\left(\frac{m+1}{n}, 2ibx^n\right) - 2n \right)}{4(m+1)n}$$

Antiderivative was successfully verified.

[In] Integrate[x^m \* Cos[a + b \* x^n]^2, x]

[Out]  $-(x^{(1+m)} * (-2*n + (E^{((2*I)*a)} * (1+m) * \text{Gamma}[(1+m)/n, (-2*I)*b*x^n])) / (2^{((1+m)/n)} * ((-I)*b*x^n)^{((1+m)/n)}) + ((1+m) * \text{Gamma}[(1+m)/n, (2*I)*b*x^n]) / (2^{((1+m)/n)} * E^{((2*I)*a)} * (I*b*x^n)^{((1+m)/n)}) / (4*(1+m)*n)$

**Maple [F]** time = 0.126, size = 0, normalized size = 0.

$$\int x^m (\cos(a + bx^n))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m\*cos(a+b\*x^n)^2,x)

[Out] int(x^m\*cos(a+b\*x^n)^2,x)

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\frac{xx^m + (m+1) \int x^m \cos(2bx^n + 2a) dx}{2(m+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m\*cos(a+b\*x^n)^2,x, algorithm="maxima")

[Out]  $1/2*(x*x^m + (m+1)*integrate(x^m*cos(2*b*x^n + 2*a), x)) / (m+1)$

**Fricas [F]** time = 0., size = 0, normalized size = 0.

$$\text{integral}(x^m \cos(bx^n + a)^2, x)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^m*cos(a+b*x^n)^2,x, algorithm="fricas")
```

```
[Out] integral(x^m*cos(b*x^n + a)^2, x)
```

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^m \cos^2(a + bx^n) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**m*cos(a+b*x**n)**2,x)
```

```
[Out] Integral(x**m*cos(a + b*x**n)**2, x)
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^m \cos(bx^n + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^m*cos(a+b*x^n)^2,x, algorithm="giac")
```

```
[Out] integrate(x^m*cos(b*x^n + a)^2, x)
```

### 3.78 $\int x^m \cos^3(a + bx^n) dx$

**Optimal.** Leaf size=229

$$\frac{3e^{ia}x^{m+1}(-ibx^n)^{-\frac{m+1}{n}}\Gamma\left(\frac{m+1}{n}, -ibx^n\right)}{8n} - \frac{3e^{-ia}x^{m+1}(ibx^n)^{-\frac{m+1}{n}}\Gamma\left(\frac{m+1}{n}, ibx^n\right)}{8n} - \frac{e^{3ia}3^{-\frac{m+1}{n}}x^{m+1}(-ibx^n)^{-\frac{m+1}{n}}}{8n}$$

[Out]  $(-3E^{(I*a)}*x^{(1+m)}*\Gamma[(1+m)/n, (-I)*b*x^n])/(8*n*((-I)*b*x^n)^{((1+m)/n)}) - (3*x^{(1+m)}*\Gamma[(1+m)/n, I*b*x^n])/(8*E^{(I*a)}*n*(I*b*x^n)^{((1+m)/n)}) - (E^{((3*I)*a)}*x^{(1+m)}*\Gamma[(1+m)/n, (-3*I)*b*x^n])/(8*3^{((1+m)/n)}*n*((-I)*b*x^n)^{((1+m)/n)}) - (x^{(1+m)}*\Gamma[(1+m)/n, (3*I)*b*x^n])/(8*3^{((1+m)/n)}*E^{((3*I)*a)}*n*(I*b*x^n)^{((1+m)/n)})$

**Rubi [A]** time = 0.211757, antiderivative size = 229, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3426, 3424, 2218}

$$\frac{3e^{ia}x^{m+1}(-ibx^n)^{-\frac{m+1}{n}}\Gamma\left(\frac{m+1}{n}, -ibx^n\right)}{8n} - \frac{3e^{-ia}x^{m+1}(ibx^n)^{-\frac{m+1}{n}}\Gamma\left(\frac{m+1}{n}, ibx^n\right)}{8n} - \frac{e^{3ia}3^{-\frac{m+1}{n}}x^{m+1}(-ibx^n)^{-\frac{m+1}{n}}}{8n}$$

Antiderivative was successfully verified.

[In] Int[x^m\*Cos[a + b\*x^n]^3, x]

[Out]  $(-3E^{(I*a)}*x^{(1+m)}*\Gamma[(1+m)/n, (-I)*b*x^n])/(8*n*((-I)*b*x^n)^{((1+m)/n)}) - (3*x^{(1+m)}*\Gamma[(1+m)/n, I*b*x^n])/(8*E^{(I*a)}*n*(I*b*x^n)^{((1+m)/n)}) - (E^{((3*I)*a)}*x^{(1+m)}*\Gamma[(1+m)/n, (-3*I)*b*x^n])/(8*3^{((1+m)/n)}*n*((-I)*b*x^n)^{((1+m)/n)}) - (x^{(1+m)}*\Gamma[(1+m)/n, (3*I)*b*x^n])/(8*3^{((1+m)/n)}*E^{((3*I)*a)}*n*(I*b*x^n)^{((1+m)/n)})$

#### Rule 3426

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)])\*(b\_.)^(p\_)\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] :> Int[ExpandTrigReduce[(e\*x)^m, (a + b\*Cos[c + d\*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]

#### Rule 3424

Int[Cos[(c\_.) + (d\_.)\*(x\_)^(n\_)]\*((e\_.)\*(x\_))^(m\_.), x\_Symbol] :> Dist[1/2, Int[(e\*x)^m\*E^(-(c\*I) - d\*I\*x^n), x], x] + Dist[1/2, Int[(e\*x)^m\*E^(c\*I + d\*I\*x^n), x], x] /; FreeQ[{c, d, e, m, n}, x]

#### Rule 2218

Int[(F\_)^((a\_.) + (b\_.)\*((c\_.) + (d\_.)\*(x\_))^(n\_))\*((e\_.) + (f\_.)\*(x\_))^(m\_.), x\_Symbol] :> -Simp[(F^a\*(e + f\*x)^(m + 1)\*Gamma[(m + 1)/n, -(b\*(c + d\*x))^n\*Log[F]])/(f\*n\*(-(b\*(c + d\*x))^n\*Log[F]))^((m + 1)/n), x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d\*e - c\*f, 0]

#### Rubi steps



$$\begin{aligned}
\int x^m \cos^3(a + bx^n) dx &= \int \left( \frac{3}{4} x^m \cos(a + bx^n) + \frac{1}{4} x^m \cos(3a + 3bx^n) \right) dx \\
&= \frac{1}{4} \int x^m \cos(3a + 3bx^n) dx + \frac{3}{4} \int x^m \cos(a + bx^n) dx \\
&= \frac{1}{8} \int e^{-3ia-3ibx^n} x^m dx + \frac{1}{8} \int e^{3ia+3ibx^n} x^m dx + \frac{3}{8} \int e^{-ia-ibx^n} x^m dx + \frac{3}{8} \int e^{ia+ibx^n} x^m dx \\
&= \frac{3e^{ia} x^{1+m} (-ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -ibx^n\right)}{8n} - \frac{3e^{-ia} x^{1+m} (ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, ibx^n\right)}{8n} - \frac{3^{-\frac{1+m}{n}} e^{3ia} x^{1+m}}{8n}
\end{aligned}$$

**Mathematica [A]** time = 0.561795, size = 221, normalized size = 0.97

$$\frac{e^{-3ia} 3^{-\frac{m+1}{n}} x^{m+1} (b^2 x^{2n})^{-\frac{m+1}{n}} \left( e^{2ia} 3^{\frac{m+n+1}{n}} (-ibx^n)^{\frac{m+1}{n}} \text{Gamma}\left(\frac{m+1}{n}, ibx^n\right) + e^{4ia} 3^{\frac{m+n+1}{n}} (ibx^n)^{\frac{m+1}{n}} \text{Gamma}\left(\frac{m+1}{n}, -ibx^n\right) \right)}{8n}$$

Antiderivative was successfully verified.

[In] Integrate[x^m \* Cos[a + b\*x^n]^3, x]

[Out]  $-(x^{(1+m)} (3^{((1+m+n)/n)} E^{((4I)a)} (I b x^n)^{((1+m)/n)} \text{Gamma}[(1+m)/n, (-I) b x^n] + 3^{((1+m+n)/n)} E^{((2I)a)} ((-I) b x^n)^{((1+m)/n)} \text{Gamma}[(1+m)/n, I b x^n] + E^{((6I)a)} (I b x^n)^{((1+m)/n)} \text{Gamma}[(1+m)/n, (-3I) b x^n] + ((-I) b x^n)^{((1+m)/n)} \text{Gamma}[(1+m)/n, (3I) b x^n]) / (8 3^{((1+m)/n)} E^{((3I)a)} n (b^2 x^{(2n)})^{((1+m)/n)})$

**Maple [F]** time = 0.178, size = 0, normalized size = 0.

$$\int x^m (\cos(a + bx^n))^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m\*cos(a+b\*x^n)^3,x)

[Out] int(x^m\*cos(a+b\*x^n)^3,x)

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int x^m \cos(bx^n + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m\*cos(a+b\*x^n)^3,x, algorithm="maxima")

[Out] integrate(x^m\*cos(b\*x^n + a)^3, x)

**Fricas [F]** time = 0., size = 0, normalized size = 0.

$$\text{integral}(x^m \cos(bx^n + a)^3, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>m</sup>\*cos(a+b\*x<sup>n</sup>)<sup>3</sup>,x, algorithm="fricas")

[Out] integral(x<sup>m</sup>\*cos(b\*x<sup>n</sup> + a)<sup>3</sup>, x)

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^m \cos^3(a + bx^n) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*m\*cos(a+b\*x\*\*n)\*\*3,x)

[Out] Integral(x\*\*m\*cos(a + b\*x\*\*n)\*\*3, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^m \cos(bx^n + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>m</sup>\*cos(a+b\*x<sup>n</sup>)<sup>3</sup>,x, algorithm="giac")

[Out] integrate(x<sup>m</sup>\*cos(b\*x<sup>n</sup> + a)<sup>3</sup>, x)

### 3.79 $\int x^{-1-n} \cos(a + bx^n) dx$

**Optimal.** Leaf size=47

$$-\frac{b \sin(a) \operatorname{CosIntegral}(bx^n)}{n} - \frac{b \cos(a) \operatorname{Si}(bx^n)}{n} - \frac{x^{-n} \cos(a + bx^n)}{n}$$

[Out]  $-(\operatorname{Cos}[a + b*x^n]/(n*x^n)) - (b*\operatorname{CosIntegral}[b*x^n]*\operatorname{Sin}[a])/n - (b*\operatorname{Cos}[a]*\operatorname{SinIntegral}[b*x^n])/n$

**Rubi [A]** time = 0.0864871, antiderivative size = 47, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$ , Rules used = {3380, 3297, 3303, 3299, 3302}

$$-\frac{b \sin(a) \operatorname{CosIntegral}(bx^n)}{n} - \frac{b \cos(a) \operatorname{Si}(bx^n)}{n} - \frac{x^{-n} \cos(a + bx^n)}{n}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[x^{(-1 - n)}*\operatorname{Cos}[a + b*x^n], x]$

[Out]  $-(\operatorname{Cos}[a + b*x^n]/(n*x^n)) - (b*\operatorname{CosIntegral}[b*x^n]*\operatorname{Sin}[a])/n - (b*\operatorname{Cos}[a]*\operatorname{SinIntegral}[b*x^n])/n$

#### Rule 3380

$\operatorname{Int}[(a + \operatorname{Cos}(c + (d*x)^n)*(b*x)^p)*(x)^m, x\_Symbol] \rightarrow \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n) - 1]}*(a + b*\operatorname{Cos}[c + d*x])^p, x], x, x^n], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, m, n, p\}, x \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]] \&\& (\operatorname{EqQ}[p, 1] \mid \mid \operatorname{EqQ}[m, n-1] \mid \mid (\operatorname{IntegerQ}[p] \&\& \operatorname{GtQ}[\operatorname{Simplify}[(m+1)/n], 0]))$

#### Rule 3297

$\operatorname{Int}[(c + (d*x)^m)*\operatorname{sin}[e + (f*x)], x\_Symbol] \rightarrow \operatorname{Simp}[(c + d*x)^{m+1}*\operatorname{Sin}[e + f*x]/(d*(m+1)), x] - \operatorname{Dist}[f/(d*(m+1)), \operatorname{Int}[(c + d*x)^{m+1}*\operatorname{Cos}[e + f*x], x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \&\& \operatorname{LtQ}[m, -1]$

#### Rule 3303

$\operatorname{Int}[\operatorname{sin}[(e + (f*x))/(c + (d*x))], x\_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Cos}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Sin}[(c*f)/d + f*x]/(c + d*x), x], x] + \operatorname{Dist}[\operatorname{Sin}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Cos}[(c*f)/d + f*x]/(c + d*x), x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \&\& \operatorname{NeQ}[d*e - c*f, 0]$

#### Rule 3299

$\operatorname{Int}[\operatorname{sin}[(e + (f*x))/(c + (d*x))], x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{SinIntegral}[e + f*x]/d, x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \&\& \operatorname{EqQ}[d*e - c*f, 0]$

#### Rule 3302

$\operatorname{Int}[\operatorname{sin}[(e + (f*x))/(c + (d*x))], x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{CosIntegral}[e - \operatorname{Pi}/2 + f*x]/d, x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x \&\& \operatorname{EqQ}[d*(e - \operatorname{Pi}/2) - c*f, 0]$

Rubi steps

$$\begin{aligned}
\int x^{-1-n} \cos(a + bx^n) dx &= \frac{\text{Subst}\left(\int \frac{\cos(a+bx)}{x^2} dx, x, x^n\right)}{n} \\
&= -\frac{x^{-n} \cos(a + bx^n)}{n} - \frac{b \text{Subst}\left(\int \frac{\sin(a+bx)}{x} dx, x, x^n\right)}{n} \\
&= -\frac{x^{-n} \cos(a + bx^n)}{n} - \frac{(b \cos(a)) \text{Subst}\left(\int \frac{\sin(bx)}{x} dx, x, x^n\right)}{n} - \frac{(b \sin(a)) \text{Subst}\left(\int \frac{\cos(bx)}{x} dx, x, x^n\right)}{n} \\
&= -\frac{x^{-n} \cos(a + bx^n)}{n} - \frac{b \text{Ci}(bx^n) \sin(a)}{n} - \frac{b \cos(a) \text{Si}(bx^n)}{n}
\end{aligned}$$

**Mathematica [A]** time = 0.0938222, size = 45, normalized size = 0.96

$$-\frac{x^{-n} (b \sin(a) x^n \text{CosIntegral}(bx^n) + b \cos(a) x^n \text{Si}(bx^n) + \cos(a + bx^n))}{n}$$

Antiderivative was successfully verified.

[In] Integrate[x<sup>^</sup>(-1 - n)\*Cos[a + b\*x<sup>^</sup>n], x]

[Out] -((Cos[a + b\*x<sup>^</sup>n] + b\*x<sup>^</sup>n\*CosIntegral[b\*x<sup>^</sup>n]\*Sin[a] + b\*x<sup>^</sup>n\*Cos[a]\*SinIntegral[b\*x<sup>^</sup>n])/(n\*x<sup>^</sup>n))

**Maple [A]** time = 0.036, size = 45, normalized size = 1.

$$\frac{b}{n} \left( -\frac{\cos(a + bx^n)}{bx^n} - \text{Si}(bx^n) \cos(a) - \text{Ci}(bx^n) \sin(a) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x<sup>^</sup>(-n-1)\*cos(a+b\*x<sup>^</sup>n), x)

[Out] 1/n\*b\*(-cos(a+b\*x<sup>^</sup>n)/(x<sup>^</sup>n)/b-Si(b\*x<sup>^</sup>n)\*cos(a)-Ci(b\*x<sup>^</sup>n)\*sin(a))

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-n-1} \cos(bx^n + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>^</sup>(-1-n)\*cos(a+b\*x<sup>^</sup>n), x, algorithm="maxima")

[Out] integrate(x<sup>^</sup>(-n - 1)\*cos(b\*x<sup>^</sup>n + a), x)

**Fricas [A]** time = 1.66697, size = 190, normalized size = 4.04

$$-\frac{bx^n \text{Ci}(bx^n) \sin(a) + bx^n \text{Ci}(-bx^n) \sin(a) + 2bx^n \cos(a) \text{Si}(bx^n) + 2 \cos(bx^n + a)}{2nx^n}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^(-1-n)*cos(a+b*x^n),x, algorithm="fricas")
```

```
[Out] -1/2*(b*x^n*cos_integral(b*x^n)*sin(a) + b*x^n*cos_integral(-b*x^n)*sin(a)
+ 2*b*x^n*cos(a)*sin_integral(b*x^n) + 2*cos(b*x^n + a))/(n*x^n)
```

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**(-1-n)*cos(a+b*x**n),x)
```

```
[Out] Timed out
```

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-n-1} \cos(bx^n + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^(-1-n)*cos(a+b*x^n),x, algorithm="giac")
```

```
[Out] integrate(x^(-n - 1)*cos(b*x^n + a), x)
```

### 3.80 $\int x^{-1-n} \cos^2(a + bx^n) dx$

**Optimal.** Leaf size=69

$$-\frac{b \sin(2a) \operatorname{CosIntegral}(2bx^n)}{n} - \frac{b \cos(2a) \operatorname{Si}(2bx^n)}{n} - \frac{x^{-n} \cos(2(a + bx^n))}{2n} - \frac{x^{-n}}{2n}$$

[Out]  $-1/(2*n*x^n) - \operatorname{Cos}[2*(a + b*x^n)]/(2*n*x^n) - (b*\operatorname{CosIntegral}[2*b*x^n]*\operatorname{Sin}[2*a])/n - (b*\operatorname{Cos}[2*a]*\operatorname{SinIntegral}[2*b*x^n])/n$

**Rubi [A]** time = 0.120562, antiderivative size = 69, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 6, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3426, 3380, 3297, 3303, 3299, 3302}

$$-\frac{b \sin(2a) \operatorname{CosIntegral}(2bx^n)}{n} - \frac{b \cos(2a) \operatorname{Si}(2bx^n)}{n} - \frac{x^{-n} \cos(2(a + bx^n))}{2n} - \frac{x^{-n}}{2n}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[x^{(-1 - n)}*\operatorname{Cos}[a + b*x^n]^2, x]$

[Out]  $-1/(2*n*x^n) - \operatorname{Cos}[2*(a + b*x^n)]/(2*n*x^n) - (b*\operatorname{CosIntegral}[2*b*x^n]*\operatorname{Sin}[2*a])/n - (b*\operatorname{Cos}[2*a]*\operatorname{SinIntegral}[2*b*x^n])/n$

#### Rule 3426

$\operatorname{Int}[(a + \operatorname{Cos}[c + (d*x)^n])*(b*x)^p*((e*x)^m), x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandTrigReduce}[(e*x)^m, (a + b*\operatorname{Cos}[c + d*x^n])^p, x], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, e, m, n\}, x\} \ \&\& \ \operatorname{IGtQ}[p, 0]$

#### Rule 3380

$\operatorname{Int}[(a + \operatorname{Cos}[c + (d*x)^n])*(b*x)^p*(x)^m, x\_Symbol] \rightarrow \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m + 1)/n] - 1)*(a + b*\operatorname{Cos}[c + d*x])^p}], x, x^n], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, m, n, p\}, x\} \ \&\& \ \operatorname{IntegerQ}[\operatorname{Simplify}[(m + 1)/n]] \ \&\& \ (\operatorname{EqQ}[p, 1] \ \|\ \operatorname{EqQ}[m, n - 1] \ \|\ (\operatorname{IntegerQ}[p] \ \&\& \ \operatorname{GtQ}[\operatorname{Simplify}[(m + 1)/n], 0]))$

#### Rule 3297

$\operatorname{Int}[(c + (d*x)^m)*\operatorname{sin}[e + f*x], x\_Symbol] \rightarrow \operatorname{Simp}[(c + d*x)^{m+1}*\operatorname{Sin}[e + f*x]/(d*(m + 1)), x] - \operatorname{Dist}[f/(d*(m + 1)), \operatorname{Int}[(c + d*x)^{m+1}*\operatorname{Cos}[e + f*x], x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x\} \ \&\& \ \operatorname{LtQ}[m, -1]$

#### Rule 3303

$\operatorname{Int}[\operatorname{sin}[e + f*x]/((c + (d*x)^m)), x\_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Cos}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Sin}[(c*f)/d + f*x]/(c + d*x), x], x] + \operatorname{Dist}[\operatorname{Sin}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Cos}[(c*f)/d + f*x]/(c + d*x), x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x\} \ \&\& \ \operatorname{NeQ}[d*e - c*f, 0]$

#### Rule 3299

$\operatorname{Int}[\operatorname{sin}[e + f*x]/((c + (d*x)^m)), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{SinIntegral}[e + f*x]/d, x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x\} \ \&\& \ \operatorname{EqQ}[d*e - c*f, 0]$

Rule 3302

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

Rubi steps

$$\begin{aligned}
\int x^{-1-n} \cos^2(a + bx^n) dx &= \int \left( \frac{x^{-1-n}}{2} + \frac{1}{2} x^{-1-n} \cos(2a + 2bx^n) \right) dx \\
&= -\frac{x^{-n}}{2n} + \frac{1}{2} \int x^{-1-n} \cos(2a + 2bx^n) dx \\
&= -\frac{x^{-n}}{2n} + \frac{\text{Subst}\left(\int \frac{\cos(2a+2bx)}{x^2} dx, x, x^n\right)}{2n} \\
&= -\frac{x^{-n}}{2n} - \frac{x^{-n} \cos(2(a + bx^n))}{2n} - \frac{b \text{Subst}\left(\int \frac{\sin(2a+2bx)}{x} dx, x, x^n\right)}{n} \\
&= -\frac{x^{-n}}{2n} - \frac{x^{-n} \cos(2(a + bx^n))}{2n} - \frac{(b \cos(2a)) \text{Subst}\left(\int \frac{\sin(2bx)}{x} dx, x, x^n\right)}{n} - \frac{(b \sin(2a)) \text{Subst}\left(\int \frac{\cos(2bx)}{x} dx, x, x^n\right)}{n} \\
&= -\frac{x^{-n}}{2n} - \frac{x^{-n} \cos(2(a + bx^n))}{2n} - \frac{b \text{Ci}(2bx^n) \sin(2a)}{n} - \frac{b \cos(2a) \text{Si}(2bx^n)}{n}
\end{aligned}$$

**Mathematica [A]** time = 0.186908, size = 53, normalized size = 0.77

$$\frac{x^{-n} \left( b \sin(2a) x^n \text{CosIntegral}(2bx^n) + b \cos(2a) x^n \text{Si}(2bx^n) + \cos^2(a + bx^n) \right)}{n}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^(-1 - n)*Cos[a + b*x^n]^2, x]
```

```
[Out] -((Cos[a + b*x^n]^2 + b*x^n*CosIntegral[2*b*x^n]*Sin[2*a] + b*x^n*Cos[2*a]*SinIntegral[2*b*x^n])/(n*x^n))
```

**Maple [A]** time = 0.044, size = 65, normalized size = 0.9

$$-\frac{1}{2nx^n} + \frac{b}{n} \left( -\frac{\cos(2a + 2bx^n)}{2bx^n} - \text{Si}(2bx^n) \cos(2a) - \text{Ci}(2bx^n) \sin(2a) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^(-n-1)*cos(a+b*x^n)^2, x)
```

```
[Out] -1/2/n/(x^n)+1/n*b*(-1/2*cos(2*a+2*b*x^n)/(x^n)/b-Si(2*b*x^n)*cos(2*a)-Ci(2*b*x^n)*sin(2*a))
```

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\frac{nx^n \int \frac{\cos(2bx^n+2a)}{xx^n} dx - 1}{2nx^n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>(-1-n)</sup>\*cos(a+b\*x<sup>n</sup>)<sup>2</sup>,x, algorithm="maxima")

[Out] 1/2\*(n\*x<sup>n</sup>\*integrate(cos(2\*b\*x<sup>n</sup> + 2\*a)/(x\*x<sup>n</sup>), x) - 1)/(n\*x<sup>n</sup>)

**Fricas [A]** time = 1.73607, size = 209, normalized size = 3.03

$$\frac{bx^n \operatorname{Ci}(2bx^n) \sin(2a) + bx^n \operatorname{Ci}(-2bx^n) \sin(2a) + 2bx^n \cos(2a) \operatorname{Si}(2bx^n) + 2 \cos(bx^n + a)^2}{2nx^n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>(-1-n)</sup>\*cos(a+b\*x<sup>n</sup>)<sup>2</sup>,x, algorithm="fricas")

[Out] -1/2\*(b\*x<sup>n</sup>\*cos\_integral(2\*b\*x<sup>n</sup>)\*sin(2\*a) + b\*x<sup>n</sup>\*cos\_integral(-2\*b\*x<sup>n</sup>)\*sin(2\*a) + 2\*b\*x<sup>n</sup>\*cos(2\*a)\*sin\_integral(2\*b\*x<sup>n</sup>) + 2\*cos(b\*x<sup>n</sup> + a)<sup>2</sup>)/(n\*x<sup>n</sup>)

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>\*\*(-1-n)</sup>\*cos(a+b\*x<sup>\*\*n</sup>)<sup>\*\*2</sup>,x)

[Out] Timed out

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-n-1} \cos(bx^n + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>(-1-n)</sup>\*cos(a+b\*x<sup>n</sup>)<sup>2</sup>,x, algorithm="giac")

[Out] integrate(x<sup>(-n - 1)</sup>\*cos(b\*x<sup>n</sup> + a)<sup>2</sup>, x)



### 3.81 $\int x^{-1-n} \cos^3(a + bx^n) dx$

**Optimal.** Leaf size=113

$$\frac{3b \sin(a) \operatorname{CosIntegral}(bx^n)}{4n} - \frac{3b \sin(3a) \operatorname{CosIntegral}(3bx^n)}{4n} - \frac{3b \cos(a) \operatorname{Si}(bx^n)}{4n} - \frac{3b \cos(3a) \operatorname{Si}(3bx^n)}{4n} - \frac{3x^{-n} \cos(a + bx^n)}{4n}$$

[Out]  $(-3 \operatorname{Cos}[a + b x^n]) / (4 n x^n) - \operatorname{Cos}[3(a + b x^n)] / (4 n x^n) - (3 b \operatorname{CosIntegral}[b x^n] \operatorname{Sin}[a]) / (4 n) - (3 b \operatorname{CosIntegral}[3 b x^n] \operatorname{Sin}[3 a]) / (4 n) - (3 b \operatorname{Cos}[a] \operatorname{SinIntegral}[b x^n]) / (4 n) - (3 b \operatorname{Cos}[3 a] \operatorname{SinIntegral}[3 b x^n]) / (4 n)$

**Rubi [A]** time = 0.199548, antiderivative size = 113, normalized size of antiderivative = 1., number of steps used = 12, number of rules used = 6, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3426, 3380, 3297, 3303, 3299, 3302}

$$\frac{3b \sin(a) \operatorname{CosIntegral}(bx^n)}{4n} - \frac{3b \sin(3a) \operatorname{CosIntegral}(3bx^n)}{4n} - \frac{3b \cos(a) \operatorname{Si}(bx^n)}{4n} - \frac{3b \cos(3a) \operatorname{Si}(3bx^n)}{4n} - \frac{3x^{-n} \cos(a + bx^n)}{4n}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[x^{(-1 - n)} \operatorname{Cos}[a + b x^n]^3, x]$

[Out]  $(-3 \operatorname{Cos}[a + b x^n]) / (4 n x^n) - \operatorname{Cos}[3(a + b x^n)] / (4 n x^n) - (3 b \operatorname{CosIntegral}[b x^n] \operatorname{Sin}[a]) / (4 n) - (3 b \operatorname{CosIntegral}[3 b x^n] \operatorname{Sin}[3 a]) / (4 n) - (3 b \operatorname{Cos}[a] \operatorname{SinIntegral}[b x^n]) / (4 n) - (3 b \operatorname{Cos}[3 a] \operatorname{SinIntegral}[3 b x^n]) / (4 n)$

#### Rule 3426

$\operatorname{Int}[(a + \operatorname{Cos}(c + d x^n) + (d x^n)^m (b + e x^n)^p], x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandTrigReduce}[(e x^n)^m, (a + b \operatorname{Cos}[c + d x^n])^p, x], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, e, m, n\}, x$  &&  $\operatorname{IGtQ}[p, 0]$

#### Rule 3380

$\operatorname{Int}[(a + \operatorname{Cos}(c + d x^n) + (d x^n)^m (b + e x^n)^p] x^m, x\_Symbol] \rightarrow \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m + 1)/n] - 1)} (a + b \operatorname{Cos}[c + d x])^p], x, x^n], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, m, n, p\}, x$  &&  $\operatorname{IntegerQ}[\operatorname{Simplify}[(m + 1)/n]]$  &&  $(\operatorname{EqQ}[p, 1] \mid \mid \operatorname{EqQ}[m, n - 1] \mid \mid (\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[\operatorname{Simplify}[(m + 1)/n], 0]))$

#### Rule 3297

$\operatorname{Int}[(c + d x^n)^m \sin(e + f x^n), x\_Symbol] \rightarrow \operatorname{Simp}[(c + d x^n)^{m + 1} \operatorname{Sin}[e + f x^n] / (d(m + 1)), x] - \operatorname{Dist}[f / (d(m + 1)), \operatorname{Int}[(c + d x^n)^{m + 1} \operatorname{Cos}[e + f x^n], x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x$  &&  $\operatorname{LtQ}[m, -1]$

#### Rule 3303

$\operatorname{Int}[\sin(e + f x^n) / (c + d x^n), x\_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Cos}[(d e - c f) / d], \operatorname{Int}[\operatorname{Sin}[(c f) / d + f x^n] / (c + d x^n), x], x] + \operatorname{Dist}[\operatorname{Sin}[(d e - c f) / d], \operatorname{Int}[\operatorname{Cos}[(c f) / d + f x^n] / (c + d x^n), x], x] /;$   $\operatorname{FreeQ}\{c, d, e, f\}, x$  &&  $\operatorname{NeQ}[d e - c f, 0]$

#### Rule 3299

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinInte
gral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

### Rule 3302

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosInte
gral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) -
c*f, 0]
```

### Rubi steps

$$\begin{aligned}
\int x^{-1-n} \cos^3(a + bx^n) dx &= \int \left( \frac{3}{4} x^{-1-n} \cos(a + bx^n) + \frac{1}{4} x^{-1-n} \cos(3a + 3bx^n) \right) dx \\
&= \frac{1}{4} \int x^{-1-n} \cos(3a + 3bx^n) dx + \frac{3}{4} \int x^{-1-n} \cos(a + bx^n) dx \\
&= \frac{\text{Subst}\left(\int \frac{\cos(3a+3bx)}{x^2} dx, x, x^n\right)}{4n} + \frac{3 \text{Subst}\left(\int \frac{\cos(a+bx)}{x^2} dx, x, x^n\right)}{4n} \\
&= -\frac{3x^{-n} \cos(a + bx^n)}{4n} - \frac{x^{-n} \cos(3(a + bx^n))}{4n} - \frac{(3b) \text{Subst}\left(\int \frac{\sin(a+bx)}{x} dx, x, x^n\right)}{4n} - \frac{(3b) \text{Subst}\left(\int \frac{\sin(3a+3bx)}{x} dx, x, x^n\right)}{4n} \\
&= -\frac{3x^{-n} \cos(a + bx^n)}{4n} - \frac{x^{-n} \cos(3(a + bx^n))}{4n} - \frac{(3b \cos(a)) \text{Subst}\left(\int \frac{\sin(bx)}{x} dx, x, x^n\right)}{4n} - \frac{(3b \cos(3a)) \text{Subst}\left(\int \frac{\sin(3bx)}{x} dx, x, x^n\right)}{4n} \\
&= -\frac{3x^{-n} \cos(a + bx^n)}{4n} - \frac{x^{-n} \cos(3(a + bx^n))}{4n} - \frac{3b \text{Ci}(bx^n) \sin(a)}{4n} - \frac{3b \text{Ci}(3bx^n) \sin(3a)}{4n} - \frac{3b \cos(a) \text{Si}(bx^n)}{4n} - \frac{3b \cos(3a) \text{Si}(3bx^n)}{4n}
\end{aligned}$$

**Mathematica [A]** time = 0.234899, size = 95, normalized size = 0.84

$$\frac{x^{-n} \left( 3b \sin(a) x^n \text{CosIntegral}(bx^n) + 3b \sin(3a) x^n \text{CosIntegral}(3bx^n) + 3b \cos(a) x^n \text{Si}(bx^n) + 3b \cos(3a) x^n \text{Si}(3bx^n) \right)}{4n}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^(-1 - n)*Cos[a + b*x^n]^3, x]
```

```
[Out] -(3*Cos[a + b*x^n] + Cos[3*(a + b*x^n)] + 3*b*x^n*CosIntegral[b*x^n]*Sin[a]
+ 3*b*x^n*CosIntegral[3*b*x^n]*Sin[3*a] + 3*b*x^n*Cos[a]*SinIntegral[b*x^n]
] + 3*b*x^n*Cos[3*a]*SinIntegral[3*b*x^n])/(4*n*x^n)
```

**Maple [A]** time = 0.05, size = 101, normalized size = 0.9

$$\frac{3b}{4n} \left( -\frac{\cos(a + bx^n)}{bx^n} - \text{Si}(bx^n) \cos(a) - \text{Ci}(bx^n) \sin(a) \right) + \frac{3b}{4n} \left( -\frac{\cos(3a + 3bx^n)}{3bx^n} - \text{Si}(3bx^n) \cos(3a) - \text{Ci}(3bx^n) \sin(3a) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^(-n-1)*cos(a+b*x^n)^3, x)
```

```
[Out] 3/4/n*b*(-cos(a+b*x^n)/(x^n)/b-Si(b*x^n)*cos(a)-Ci(b*x^n)*sin(a))+3/4/n*b*(
-1/3*cos(3*a+3*b*x^n)/(x^n)/b-Si(3*b*x^n)*cos(3*a)-Ci(3*b*x^n)*sin(3*a))
```

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-n-1} \cos(bx^n + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(-1-n)\*cos(a+b\*x^n)^3,x, algorithm="maxima")

[Out] integrate(x^(-n - 1)\*cos(b\*x^n + a)^3, x)

**Fricas [A]** time = 1.85088, size = 366, normalized size = 3.24

$$\frac{3bx^n \operatorname{Ci}(3bx^n) \sin(3a) + 3bx^n \operatorname{Ci}(-3bx^n) \sin(3a) + 3bx^n \operatorname{Ci}(bx^n) \sin(a) + 3bx^n \operatorname{Ci}(-bx^n) \sin(a) + 6bx^n \cos(3a)}{8nx^n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(-1-n)\*cos(a+b\*x^n)^3,x, algorithm="fricas")

[Out] -1/8\*(3\*b\*x^n\*cos\_integral(3\*b\*x^n)\*sin(3\*a) + 3\*b\*x^n\*cos\_integral(-3\*b\*x^n)\*sin(3\*a) + 3\*b\*x^n\*cos\_integral(b\*x^n)\*sin(a) + 3\*b\*x^n\*cos\_integral(-b\*x^n)\*sin(a) + 6\*b\*x^n\*cos(3\*a)\*sin\_integral(3\*b\*x^n) + 6\*b\*x^n\*cos(a)\*sin\_integral(b\*x^n) + 8\*cos(b\*x^n + a)^3)/(n\*x^n)

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*(-1-n)\*cos(a+b\*x\*\*n)\*\*3,x)

[Out] Timed out

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-n-1} \cos(bx^n + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(-1-n)\*cos(a+b\*x^n)^3,x, algorithm="giac")

[Out] integrate(x^(-n - 1)\*cos(b\*x^n + a)^3, x)

### 3.82 $\int x^{-1-2n} \cos(a + bx^n) dx$

**Optimal.** Leaf size=78

$$-\frac{b^2 \cos(a) \operatorname{CosIntegral}(bx^n)}{2n} + \frac{b^2 \sin(a) \operatorname{Si}(bx^n)}{2n} + \frac{bx^{-n} \sin(a + bx^n)}{2n} - \frac{x^{-2n} \cos(a + bx^n)}{2n}$$

[Out]  $-\operatorname{Cos}[a + b*x^n]/(2*n*x^{(2*n)}) - (b^2*\operatorname{Cos}[a]*\operatorname{CosIntegral}[b*x^n])/(2*n) + (b*\operatorname{Sin}[a + b*x^n])/(2*n*x^n) + (b^2*\operatorname{Sin}[a]*\operatorname{SinIntegral}[b*x^n])/(2*n)$

**Rubi [A]** time = 0.111042, antiderivative size = 78, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 5, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$ , Rules used = {3380, 3297, 3303, 3299, 3302}

$$-\frac{b^2 \cos(a) \operatorname{CosIntegral}(bx^n)}{2n} + \frac{b^2 \sin(a) \operatorname{Si}(bx^n)}{2n} + \frac{bx^{-n} \sin(a + bx^n)}{2n} - \frac{x^{-2n} \cos(a + bx^n)}{2n}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[x^{(-1 - 2*n)}*\operatorname{Cos}[a + b*x^n], x]$

[Out]  $-\operatorname{Cos}[a + b*x^n]/(2*n*x^{(2*n)}) - (b^2*\operatorname{Cos}[a]*\operatorname{CosIntegral}[b*x^n])/(2*n) + (b*\operatorname{Sin}[a + b*x^n])/(2*n*x^n) + (b^2*\operatorname{Sin}[a]*\operatorname{SinIntegral}[b*x^n])/(2*n)$

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
  := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p,
    x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
  && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 3297

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(((c + d*x)^(m + 1)*Sin[e + f*x])/(d*(m + 1)), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]
```

#### Rule 3303

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/(c + d*x), x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

#### Rule 3299

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

#### Rule 3302

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

Rubi steps

$$\begin{aligned}
\int x^{-1-2n} \cos(a + bx^n) dx &= \frac{\text{Subst}\left(\int \frac{\cos(a+bx)}{x^3} dx, x, x^n\right)}{n} \\
&= -\frac{x^{-2n} \cos(a + bx^n)}{2n} - \frac{b \text{Subst}\left(\int \frac{\sin(a+bx)}{x^2} dx, x, x^n\right)}{2n} \\
&= -\frac{x^{-2n} \cos(a + bx^n)}{2n} + \frac{bx^{-n} \sin(a + bx^n)}{2n} - \frac{b^2 \text{Subst}\left(\int \frac{\cos(a+bx)}{x} dx, x, x^n\right)}{2n} \\
&= -\frac{x^{-2n} \cos(a + bx^n)}{2n} + \frac{bx^{-n} \sin(a + bx^n)}{2n} - \frac{(b^2 \cos(a)) \text{Subst}\left(\int \frac{\cos(bx)}{x} dx, x, x^n\right)}{2n} + \frac{(b^2 \sin(a)) \text{Subst}\left(\int \frac{\sin(bx)}{x} dx, x, x^n\right)}{2n} \\
&= -\frac{x^{-2n} \cos(a + bx^n)}{2n} - \frac{b^2 \cos(a) \text{Ci}(bx^n)}{2n} + \frac{bx^{-n} \sin(a + bx^n)}{2n} + \frac{b^2 \sin(a) \text{Si}(bx^n)}{2n}
\end{aligned}$$

**Mathematica [A]** time = 0.129637, size = 70, normalized size = 0.9

$$\frac{x^{-2n} \left( b^2 \cos(a) x^{2n} \text{CosIntegral}(bx^n) - b^2 \sin(a) x^{2n} \text{Si}(bx^n) - bx^n \sin(a + bx^n) + \cos(a + bx^n) \right)}{2n}$$

Antiderivative was successfully verified.

[In] Integrate[x^(-1 - 2\*n)\*Cos[a + b\*x^n], x]

[Out] -(Cos[a + b\*x^n] + b^2\*x^(2\*n)\*Cos[a]\*CosIntegral[b\*x^n] - b\*x^n\*Sin[a + b\*x^n] - b^2\*x^(2\*n)\*Sin[a]\*SinIntegral[b\*x^n])/(2\*n\*x^(2\*n))

**Maple [A]** time = 0.037, size = 65, normalized size = 0.8

$$\frac{b^2}{n} \left( -\frac{\cos(a + bx^n)}{2(x^n)^2 b^2} + \frac{\sin(a + bx^n)}{2bx^n} + \frac{\text{Si}(bx^n) \sin(a)}{2} - \frac{\text{Ci}(bx^n) \cos(a)}{2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(-1-2\*n)\*cos(a+b\*x^n), x)

[Out] 1/n\*b^2\*(-1/2/(x^n)^2/b^2\*cos(a+b\*x^n)+1/2\*sin(a+b\*x^n)/(x^n)/b+1/2\*Si(b\*x^n)\*sin(a)-1/2\*Ci(b\*x^n)\*cos(a))

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-2n-1} \cos(bx^n + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(-1-2\*n)\*cos(a+b\*x^n), x, algorithm="maxima")

[Out] integrate(x^(-2\*n - 1)\*cos(b\*x^n + a), x)

**Fricas [A]** time = 1.6691, size = 254, normalized size = 3.26

$$\frac{b^2 x^{2n} \cos(a) \operatorname{Ci}(bx^n) + b^2 x^{2n} \cos(a) \operatorname{Ci}(-bx^n) - 2 b^2 x^{2n} \sin(a) \operatorname{Si}(bx^n) - 2 bx^n \sin(bx^n + a) + 2 \cos(bx^n + a)}{4 n x^{2n}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>^</sup>(-1-2\*n)\*cos(a+b\*x<sup>^</sup>n),x, algorithm="fricas")

[Out] -1/4\*(b<sup>^</sup>2\*x<sup>^</sup>(2\*n)\*cos(a)\*cos\_integral(b\*x<sup>^</sup>n) + b<sup>^</sup>2\*x<sup>^</sup>(2\*n)\*cos(a)\*cos\_integral(-b\*x<sup>^</sup>n) - 2\*b<sup>^</sup>2\*x<sup>^</sup>(2\*n)\*sin(a)\*sin\_integral(b\*x<sup>^</sup>n) - 2\*b\*x<sup>^</sup>n\*sin(b\*x<sup>^</sup>n + a) + 2\*cos(b\*x<sup>^</sup>n + a))/(n\*x<sup>^</sup>(2\*n))

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>^</sup>(-1-2\*n)\*cos(a+b\*x<sup>^</sup>n),x)

[Out] Timed out

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-2n-1} \cos(bx^n + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>^</sup>(-1-2\*n)\*cos(a+b\*x<sup>^</sup>n),x, algorithm="giac")

[Out] integrate(x<sup>^</sup>(-2\*n - 1)\*cos(b\*x<sup>^</sup>n + a), x)

### 3.83 $\int x^{-1-2n} \cos^2(a + bx^n) dx$

**Optimal.** Leaf size=95

$$-\frac{b^2 \cos(2a) \operatorname{CosIntegral}(2bx^n)}{n} + \frac{b^2 \sin(2a) \operatorname{Si}(2bx^n)}{n} + \frac{bx^{-n} \sin(2(a + bx^n))}{2n} - \frac{x^{-2n} \cos(2(a + bx^n))}{4n} - \frac{x^{-2n}}{4n}$$

[Out]  $-1/(4*n*x^{(2*n)}) - \operatorname{Cos}[2*(a + b*x^n)]/(4*n*x^{(2*n)}) - (b^2*\operatorname{Cos}[2*a]*\operatorname{CosIntegral}[2*b*x^n])/n + (b*\operatorname{Sin}[2*(a + b*x^n)])/(2*n*x^n) + (b^2*\operatorname{Sin}[2*a]*\operatorname{SinIntegral}[2*b*x^n])/n$

**Rubi [A]** time = 0.14742, antiderivative size = 95, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 6, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3426, 3380, 3297, 3303, 3299, 3302}

$$-\frac{b^2 \cos(2a) \operatorname{CosIntegral}(2bx^n)}{n} + \frac{b^2 \sin(2a) \operatorname{Si}(2bx^n)}{n} + \frac{bx^{-n} \sin(2(a + bx^n))}{2n} - \frac{x^{-2n} \cos(2(a + bx^n))}{4n} - \frac{x^{-2n}}{4n}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[x^{(-1 - 2*n)}*\operatorname{Cos}[a + b*x^n]^2, x]$

[Out]  $-1/(4*n*x^{(2*n)}) - \operatorname{Cos}[2*(a + b*x^n)]/(4*n*x^{(2*n)}) - (b^2*\operatorname{Cos}[2*a]*\operatorname{CosIntegral}[2*b*x^n])/n + (b*\operatorname{Sin}[2*(a + b*x^n)])/(2*n*x^n) + (b^2*\operatorname{Sin}[2*a]*\operatorname{SinIntegral}[2*b*x^n])/n$

#### Rule 3426

$\operatorname{Int}[(a_.) + \operatorname{Cos}(c_.) + (d_.)*(x_.)^{(n_.)}*(b_.)]^{(p_.)}*((e_.)*(x_.))^{(m_.)}, x\_Symbol] :> \operatorname{Int}[\operatorname{ExpandTrigReduce}[(e*x)^m, (a + b*\operatorname{Cos}[c + d*x^n])^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, m, n\}, x \ \&\& \operatorname{IGtQ}[p, 0]$

#### Rule 3380

$\operatorname{Int}[(a_.) + \operatorname{Cos}(c_.) + (d_.)*(x_.)^{(n_.)}*(b_.)]^{(p_.)}*(x_.)^{(m_.)}, x\_Symbol] :> \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m + 1)/n] - 1)*(a + b*\operatorname{Cos}[c + d*x])^p}, x], x, x^n], x] /; \operatorname{FreeQ}\{a, b, c, d, m, n, p\}, x \ \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m + 1)/n]] \ \&\& (\operatorname{EqQ}[p, 1] \ \|\ \operatorname{EqQ}[m, n - 1] \ \|\ (\operatorname{IntegerQ}[p] \ \&\& \operatorname{GtQ}[\operatorname{Simplify}[(m + 1)/n], 0]))$

#### Rule 3297

$\operatorname{Int}[(c_.) + (d_.)*(x_.)]^{(m_.)}*\operatorname{sin}[(e_.) + (f_.)*(x_.)], x\_Symbol] :> \operatorname{Simp}[(c + d*x)^{(m + 1)}*\operatorname{Sin}[e + f*x]/(d*(m + 1)), x] - \operatorname{Dist}[f/(d*(m + 1)), \operatorname{Int}[(c + d*x)^{(m + 1)}*\operatorname{Cos}[e + f*x], x], x] /; \operatorname{FreeQ}\{c, d, e, f\}, x \ \&\& \operatorname{LtQ}[m, -1]$

#### Rule 3303

$\operatorname{Int}[\operatorname{sin}[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x\_Symbol] :> \operatorname{Dist}[\operatorname{Cos}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Sin}[(c*f)/d + f*x]/(c + d*x), x], x] + \operatorname{Dist}[\operatorname{Sin}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Cos}[(c*f)/d + f*x]/(c + d*x), x], x] /; \operatorname{FreeQ}\{c, d, e, f\}, x \ \&\& \operatorname{NeQ}[d*e - c*f, 0]$

#### Rule 3299

`Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

### Rule 3302

`Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

### Rubi steps

$$\begin{aligned}
 \int x^{-1-2n} \cos^2(a + bx^n) dx &= \int \left( \frac{1}{2} x^{-1-2n} + \frac{1}{2} x^{-1-2n} \cos(2a + 2bx^n) \right) dx \\
 &= -\frac{x^{-2n}}{4n} + \frac{1}{2} \int x^{-1-2n} \cos(2a + 2bx^n) dx \\
 &= -\frac{x^{-2n}}{4n} + \frac{\text{Subst}\left(\int \frac{\cos(2a+2bx)}{x^3} dx, x, x^n\right)}{2n} \\
 &= -\frac{x^{-2n}}{4n} - \frac{x^{-2n} \cos(2(a + bx^n))}{4n} - \frac{b \text{Subst}\left(\int \frac{\sin(2a+2bx)}{x^2} dx, x, x^n\right)}{2n} \\
 &= -\frac{x^{-2n}}{4n} - \frac{x^{-2n} \cos(2(a + bx^n))}{4n} + \frac{bx^{-n} \sin(2(a + bx^n))}{2n} - \frac{b^2 \text{Subst}\left(\int \frac{\cos(2a+2bx)}{x} dx, x, x^n\right)}{n} \\
 &= -\frac{x^{-2n}}{4n} - \frac{x^{-2n} \cos(2(a + bx^n))}{4n} + \frac{bx^{-n} \sin(2(a + bx^n))}{2n} - \frac{(b^2 \cos(2a)) \text{Subst}\left(\int \frac{\cos(2bx)}{x} dx, x, x^n\right)}{n} \\
 &= -\frac{x^{-2n}}{4n} - \frac{x^{-2n} \cos(2(a + bx^n))}{4n} - \frac{b^2 \cos(2a) \text{Ci}(2bx^n)}{n} + \frac{bx^{-n} \sin(2(a + bx^n))}{2n} + \frac{b^2 \sin(2a)}{n}
 \end{aligned}$$

**Mathematica [A]** time = 0.21181, size = 82, normalized size = 0.86

$$\frac{x^{-2n} \left( 4b^2 \cos(2a)x^{2n} \text{CosIntegral}(2bx^n) - 4b^2 \sin(2a)x^{2n} \text{Si}(2bx^n) - 2bx^n \sin(2(a + bx^n)) + \cos(2(a + bx^n)) + 1 \right)}{4n}$$

Antiderivative was successfully verified.

[In] Integrate[x^(-1 - 2\*n)\*Cos[a + b\*x^n]^2, x]

[Out] -(1 + Cos[2\*(a + b\*x^n)] + 4\*b^2\*x^(2\*n)\*Cos[2\*a]\*CosIntegral[2\*b\*x^n] - 2\*b\*x^n\*Sin[2\*(a + b\*x^n)] - 4\*b^2\*x^(2\*n)\*Sin[2\*a]\*SinIntegral[2\*b\*x^n])/(4\*n\*x^(2\*n))

**Maple [A]** time = 0.043, size = 89, normalized size = 0.9

$$-\frac{1}{4n(x^n)^2} + 2\frac{b^2}{n} \left( -\frac{1}{8} \frac{\cos(2a + 2bx^n)}{(x^n)^2 b^2} + \frac{1}{4} \frac{\sin(2a + 2bx^n)}{bx^n} + \frac{1}{2} \text{Si}(2bx^n) \sin(2a) - \frac{1}{2} \text{Ci}(2bx^n) \cos(2a) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(-1-2\*n)\*cos(a+b\*x^n)^2, x)

[Out] -1/4/n/(x^n)^2+2/n\*b^2\*(-1/8\*cos(2\*a+2\*b\*x^n)/(x^n)^2/b^2+1/4\*sin(2\*a+2\*b\*x^n)/(x^n)/b+1/2\*Si(2\*b\*x^n)\*sin(2\*a)-1/2\*Ci(2\*b\*x^n)\*cos(2\*a))



---

**Maxima [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>(-1-2\*n)</sup>\*cos(a+b\*x<sup>n</sup>)<sup>2</sup>,x, algorithm="maxima")

[Out] Timed out

---

**Fricas [A]** time = 2.12339, size = 290, normalized size = 3.05

$$\frac{b^2 x^{2n} \cos(2a) \operatorname{Ci}(2bx^n) + b^2 x^{2n} \cos(2a) \operatorname{Ci}(-2bx^n) - 2b^2 x^{2n} \sin(2a) \operatorname{Si}(2bx^n) - 2bx^n \cos(bx^n + a) \sin(bx^n + a)}{2nx^{2n}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>(-1-2\*n)</sup>\*cos(a+b\*x<sup>n</sup>)<sup>2</sup>,x, algorithm="fricas")

[Out]  $-1/2*(b^2*x^{(2*n)}*\cos(2*a)*\cos\_integral(2*b*x^n) + b^2*x^{(2*n)}*\cos(2*a)*\cos\_integral(-2*b*x^n) - 2*b^2*x^{(2*n)}*\sin(2*a)*\sin\_integral(2*b*x^n) - 2*b*x^n*\cos(b*x^n + a)*\sin(b*x^n + a) + \cos(b*x^n + a)^2)/(n*x^{(2*n)})$

---

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>\*\*(-1-2\*n)</sup>\*cos(a+b\*x<sup>\*\*n</sup>)<sup>\*\*2</sup>,x)

[Out] Timed out

---

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-2n-1} \cos(bx^n + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x<sup>(-1-2\*n)</sup>\*cos(a+b\*x<sup>n</sup>)<sup>2</sup>,x, algorithm="giac")

[Out] integrate(x<sup>(-2\*n - 1)</sup>\*cos(b\*x<sup>n</sup> + a)<sup>2</sup>, x)

### 3.84 $\int x^{-1-2n} \cos^3(a + bx^n) dx$

**Optimal.** Leaf size=165

$$-\frac{3b^2 \cos(a) \operatorname{CosIntegral}(bx^n)}{8n} - \frac{9b^2 \cos(3a) \operatorname{CosIntegral}(3bx^n)}{8n} + \frac{3b^2 \sin(a) \operatorname{Si}(bx^n)}{8n} + \frac{9b^2 \sin(3a) \operatorname{Si}(3bx^n)}{8n} + \frac{3bx^{-n}}{8n}$$

```
[Out] (-3*Cos[a + b*x^n])/(8*n*x^(2*n)) - Cos[3*(a + b*x^n)]/(8*n*x^(2*n)) - (3*b^2*Cos[a]*CosIntegral[b*x^n])/(8*n) - (9*b^2*Cos[3*a]*CosIntegral[3*b*x^n])/(8*n) + (3*b*Sin[a + b*x^n])/(8*n*x^n) + (3*b*Sin[3*(a + b*x^n)])/(8*n*x^n) + (3*b^2*Sin[a]*SinIntegral[b*x^n])/(8*n) + (9*b^2*Sin[3*a]*SinIntegral[3*b*x^n])/(8*n)
```

**Rubi [A]** time = 0.252791, antiderivative size = 165, normalized size of antiderivative = 1., number of steps used = 14, number of rules used = 6, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3426, 3380, 3297, 3303, 3299, 3302}

$$-\frac{3b^2 \cos(a) \operatorname{CosIntegral}(bx^n)}{8n} - \frac{9b^2 \cos(3a) \operatorname{CosIntegral}(3bx^n)}{8n} + \frac{3b^2 \sin(a) \operatorname{Si}(bx^n)}{8n} + \frac{9b^2 \sin(3a) \operatorname{Si}(3bx^n)}{8n} + \frac{3bx^{-n}}{8n}$$

Antiderivative was successfully verified.

```
[In] Int[x^(-1 - 2*n)*Cos[a + b*x^n]^3,x]
```

```
[Out] (-3*Cos[a + b*x^n])/(8*n*x^(2*n)) - Cos[3*(a + b*x^n)]/(8*n*x^(2*n)) - (3*b^2*Cos[a]*CosIntegral[b*x^n])/(8*n) - (9*b^2*Cos[3*a]*CosIntegral[3*b*x^n])/(8*n) + (3*b*Sin[a + b*x^n])/(8*n*x^n) + (3*b*Sin[3*(a + b*x^n)])/(8*n*x^n) + (3*b^2*Sin[a]*SinIntegral[b*x^n])/(8*n) + (9*b^2*Sin[3*a]*SinIntegral[3*b*x^n])/(8*n)
```

#### Rule 3426

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*((e_.)*(x_)^(m_.), x_Symbol]
:> Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x]
;/; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]
```

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
:> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x]
;/; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 3297

```
Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol]
:> Simp[((c + d*x)^(m + 1)*Sin[e + f*x])/(d*(m + 1)), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x]
;/; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]
```

#### Rule 3303

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Dist[Cos[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/(c + d*x), x], x]
;/; FreeQ[{c, d, e, f}, x] &&
```

NeQ[d\*e - c\*f, 0]

### Rule 3299

Int[sin[(e\_.) + (f\_.)\*(x\_.)]/((c\_.) + (d\_.)\*(x\_.)), x\_Symbol] := Simp[SinIntegral[e + f\*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d\*e - c\*f, 0]

### Rule 3302

Int[sin[(e\_.) + (f\_.)\*(x\_.)]/((c\_.) + (d\_.)\*(x\_.)), x\_Symbol] := Simp[CosIntegral[e - Pi/2 + f\*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d\*(e - Pi/2) - c\*f, 0]

### Rubi steps

$$\begin{aligned}
 \int x^{-1-2n} \cos^3(a + bx^n) dx &= \int \left( \frac{3}{4} x^{-1-2n} \cos(a + bx^n) + \frac{1}{4} x^{-1-2n} \cos(3a + 3bx^n) \right) dx \\
 &= \frac{1}{4} \int x^{-1-2n} \cos(3a + 3bx^n) dx + \frac{3}{4} \int x^{-1-2n} \cos(a + bx^n) dx \\
 &= \frac{\text{Subst}\left(\int \frac{\cos(3a+3bx)}{x^3} dx, x, x^n\right)}{4n} + \frac{3 \text{Subst}\left(\int \frac{\cos(a+bx)}{x^3} dx, x, x^n\right)}{4n} \\
 &= -\frac{3x^{-2n} \cos(a + bx^n)}{8n} - \frac{x^{-2n} \cos(3(a + bx^n))}{8n} - \frac{(3b) \text{Subst}\left(\int \frac{\sin(a+bx)}{x^2} dx, x, x^n\right)}{8n} \quad (3b) \\
 &= -\frac{3x^{-2n} \cos(a + bx^n)}{8n} - \frac{x^{-2n} \cos(3(a + bx^n))}{8n} + \frac{3bx^{-n} \sin(a + bx^n)}{8n} + \frac{3bx^{-n} \sin(3(a + bx^n))}{8n} \\
 &= -\frac{3x^{-2n} \cos(a + bx^n)}{8n} - \frac{x^{-2n} \cos(3(a + bx^n))}{8n} + \frac{3bx^{-n} \sin(a + bx^n)}{8n} + \frac{3bx^{-n} \sin(3(a + bx^n))}{8n} \\
 &= -\frac{3x^{-2n} \cos(a + bx^n)}{8n} - \frac{x^{-2n} \cos(3(a + bx^n))}{8n} - \frac{3b^2 \cos(a) \text{Ci}(bx^n)}{8n} - \frac{9b^2 \cos(3a) \text{Ci}(3bx^n)}{8n}
 \end{aligned}$$

**Mathematica [A]** time = 0.355798, size = 141, normalized size = 0.85

$$\frac{x^{-2n} (3b^2 \cos(a)x^{2n} \text{CosIntegral}(bx^n) + 9b^2 \cos(3a)x^{2n} \text{CosIntegral}(3bx^n) - 3b^2 \sin(a)x^{2n} \text{Si}(bx^n) - 9b^2 \sin(3a)x^{2n} \text{Si}(3bx^n))}{8n}$$

Antiderivative was successfully verified.

[In] Integrate[x^(-1 - 2\*n)\*Cos[a + b\*x^n]^3, x]

[Out] -(3\*Cos[a + b\*x^n] + Cos[3\*(a + b\*x^n)] + 3\*b^2\*x^(2\*n)\*Cos[a]\*CosIntegral[b\*x^n] + 9\*b^2\*x^(2\*n)\*Cos[3\*a]\*CosIntegral[3\*b\*x^n] - 3\*b\*x^n\*Sin[a + b\*x^n] - 3\*b\*x^n\*Sin[3\*(a + b\*x^n)] - 3\*b^2\*x^(2\*n)\*Sin[a]\*SinIntegral[b\*x^n] - 9\*b^2\*x^(2\*n)\*Sin[3\*a]\*SinIntegral[3\*b\*x^n])/(8\*n\*x^(2\*n))

**Maple [A]** time = 0.05, size = 144, normalized size = 0.9

$$\frac{3b^2}{4n} \left( -\frac{\cos(a + bx^n)}{2(x^n)^2 b^2} + \frac{\sin(a + bx^n)}{2bx^n} + \frac{\text{Si}(bx^n) \sin(a)}{2} - \frac{\text{Ci}(bx^n) \cos(a)}{2} \right) + \frac{9b^2}{4n} \left( -\frac{\cos(3a + 3bx^n)}{18(x^n)^2 b^2} + \frac{\sin(3a + 3bx^n)}{6bx^n} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^(-1-2*n)*cos(a+b*x^n)^3,x)`

[Out]  $\frac{3}{4} \frac{1}{n} b^2 \left( -\frac{1}{2} \frac{1}{(x^n)^2} \frac{1}{b^2} \cos(a+b*x^n) + \frac{1}{2} \frac{\sin(a+b*x^n)}{(x^n)} \frac{1}{b} + \frac{1}{2} \text{Si}(b*x^n) \sin(a) - \frac{1}{2} \text{Ci}(b*x^n) \cos(a) \right) + \frac{9}{4} \frac{1}{n} b^2 \left( -\frac{1}{18} \frac{\cos(3*a+3*b*x^n)}{(x^n)^2} \frac{1}{b^2} + \frac{1}{6} \frac{\sin(3*a+3*b*x^n)}{(x^n)} \frac{1}{b} + \frac{1}{2} \text{Si}(3*b*x^n) \sin(3*a) - \frac{1}{2} \text{Ci}(3*b*x^n) \cos(3*a) \right)$

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-2n-1} \cos(bx^n + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^(-1-2*n)*cos(a+b*x^n)^3,x, algorithm="maxima")`

[Out] `integrate(x^(-2*n - 1)*cos(b*x^n + a)^3, x)`

**Fricas [A]** time = 2.05471, size = 481, normalized size = 2.92

$$\frac{9 b^2 x^{2n} \cos(3 a) \text{Ci}(3 b x^n) + 3 b^2 x^{2n} \cos(a) \text{Ci}(b x^n) + 3 b^2 x^{2n} \cos(a) \text{Ci}(-b x^n) + 9 b^2 x^{2n} \cos(3 a) \text{Ci}(-3 b x^n) - 24 b x^{2n}}{16 n x^{2n}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^(-1-2*n)*cos(a+b*x^n)^3,x, algorithm="fricas")`

[Out]  $-1/16 * (9 * b^2 * x^{(2*n)} * \cos(3*a) * \cos\_integral(3*b*x^n) + 3 * b^2 * x^{(2*n)} * \cos(a) * \cos\_integral(b*x^n) + 3 * b^2 * x^{(2*n)} * \cos(a) * \cos\_integral(-b*x^n) + 9 * b^2 * x^{(2*n)} * \cos(3*a) * \cos\_integral(-3*b*x^n) - 24 * b * x^n * \cos(b*x^n + a)^2 * \sin(b*x^n + a) - 18 * b^2 * x^{(2*n)} * \sin(3*a) * \sin\_integral(3*b*x^n) - 6 * b^2 * x^{(2*n)} * \sin(a) * \sin\_integral(b*x^n) + 8 * \cos(b*x^n + a)^3) / (n * x^{(2*n)})$

**Sympy [F(-1)]** time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**(-1-2*n)*cos(a+b*x**n)**3,x)`

[Out] Timed out

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int x^{-2n-1} \cos(bx^n + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^(-1-2*n)*cos(a+b*x^n)^3,x, algorithm="giac")
```

```
[Out] integrate(x^(-2*n - 1)*cos(b*x^n + a)^3, x)
```

### 3.85 $\int x^2 \cos((a + bx)^2) dx$

**Optimal.** Leaf size=99

$$\frac{\sqrt{\frac{\pi}{2}}a^2 \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^3} - \frac{\sqrt{\frac{\pi}{2}}S\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{2b^3} - \frac{a \sin((a + bx)^2)}{b^3} + \frac{(a + bx) \sin((a + bx)^2)}{2b^3}$$

[Out] (a^2\*Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*(a + b\*x)])/b^3 - (Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*(a + b\*x)])/(2\*b^3) - (a\*Sin[(a + b\*x)^2])/b^3 + ((a + b\*x)\*Sin[(a + b\*x)^2])/(2\*b^3)

**Rubi [A]** time = 0.0675234, antiderivative size = 99, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 6, integrand size = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.5$ , Rules used = {3434, 3352, 3380, 2637, 3386, 3351}

$$\frac{\sqrt{\frac{\pi}{2}}a^2 \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^3} - \frac{\sqrt{\frac{\pi}{2}}S\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{2b^3} - \frac{a \sin((a + bx)^2)}{b^3} + \frac{(a + bx) \sin((a + bx)^2)}{2b^3}$$

Antiderivative was successfully verified.

[In] Int[x^2\*Cos[(a + b\*x)^2],x]

[Out] (a^2\*Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*(a + b\*x)])/b^3 - (Sqrt[Pi/2]\*FresnelS[Sqrt[2/Pi]\*(a + b\*x)])/(2\*b^3) - (a\*Sin[(a + b\*x)^2])/b^3 + ((a + b\*x)\*Sin[(a + b\*x)^2])/(2\*b^3)

#### Rule 3434

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] :> Module[{k = If[FractionQ[n], Denominator[n], 1]}, Dist[k/f^(m + 1), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x^(k*n)])^p, x^(k - 1)*(f*g - e*h + h*x^k)^m, x], x], x, (e + f*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && IGtQ[p, 0] && IGtQ[m, 0]
```

#### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] :> Simp[(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/ (f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 2637

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] :> Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

#### Rule 3386

```
Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_))^(m_.), x_Symbol] :> Simp[(e^(n - 1)*(e*x)^(m - n + 1)*Sin[c + d*x^n])/(d*n), x] - Dist[(e^n*(m - n + 1))
```

$/(d*n), \text{Int}[(e*x)^{(m-n)}*\text{Sin}[c + d*x^n], x], x] /; \text{FreeQ}\{c, d, e\}, x\} \&\& \text{IGtQ}[n, 0] \&\& \text{LtQ}[n, m + 1]$

### Rule 3351

$\text{Int}[\text{Sin}[(d_.)*((e_.) + (f_.)*(x_.))^2], x\_Symbol] :> \text{Simp}[(\text{Sqrt}[\text{Pi}/2]*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)])/(f*\text{Rt}[d, 2]), x] /; \text{FreeQ}\{d, e, f\}, x]$

### Rubi steps

$$\begin{aligned} \int x^2 \cos((a + bx)^2) dx &= \frac{\text{Subst}\left(\int (a^2 \cos(x^2) - 2ax \cos(x^2) + x^2 \cos(x^2)) dx, x, a + bx\right)}{b^3} \\ &= \frac{\text{Subst}\left(\int x^2 \cos(x^2) dx, x, a + bx\right)}{b^3} - \frac{(2a) \text{Subst}\left(\int x \cos(x^2) dx, x, a + bx\right)}{b^3} + \frac{a^2 \text{Subst}\left(\int \cos(x^2) dx, x, a + bx\right)}{b^3} \\ &= \frac{a^2 \sqrt{\frac{\pi}{2}} C\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^3} + \frac{(a + bx) \sin((a + bx)^2)}{2b^3} - \frac{\text{Subst}\left(\int \sin(x^2) dx, x, a + bx\right)}{2b^3} - \frac{a \text{Subst}\left(\int \cos(x^2) dx, x, a + bx\right)}{b^3} \\ &= \frac{a^2 \sqrt{\frac{\pi}{2}} C\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^3} - \frac{\sqrt{\frac{\pi}{2}} S\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{2b^3} - \frac{a \sin((a + bx)^2)}{b^3} + \frac{(a + bx) \sin((a + bx)^2)}{2b^3} \end{aligned}$$

**Mathematica [A]** time = 0.265072, size = 76, normalized size = 0.77

$$\frac{-2\sqrt{2\pi}a^2\text{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right) + \sqrt{2\pi}S\left(\sqrt{\frac{2}{\pi}}(a + bx)\right) + 2(a - bx)\sin((a + bx)^2)}{4b^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2\*cos[(a + b\*x)^2], x]

[Out]  $-( -2*a^2*\text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*(a + b*x)] + \text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*(a + b*x)] + 2*(a - b*x)*\text{Sin}[(a + b*x)^2])/(4*b^3)$

**Maple [A]** time = 0.029, size = 131, normalized size = 1.3

$$\frac{x \sin(x^2 b^2 + 2 a b x + a^2)}{2 b^2} - \frac{a}{b} \left( \frac{\sin(x^2 b^2 + 2 a b x + a^2)}{2 b^2} - \frac{\sqrt{2} a \sqrt{\pi}}{2 b} \text{FresnelC}\left(\frac{\sqrt{2}(b^2 x + a b)}{\sqrt{\pi}} \frac{1}{\sqrt{b^2}}\right) \frac{1}{\sqrt{b^2}} \right) - \frac{\sqrt{2} \sqrt{\pi}}{4 b^2} \text{FresnelS}\left(\frac{\sqrt{2}(b^2 x + a b)}{\sqrt{\pi}} \frac{1}{\sqrt{b^2}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2\*cos((b\*x+a)^2), x)

[Out]  $1/2/b^2*x*\sin(b^2*x^2+2*a*b*x+a^2) - a/b*(1/2/b^2*\sin(b^2*x^2+2*a*b*x+a^2) - 1/2*a/b*2^{(1/2)}*Pi^{(1/2)}/(b^2)^{(1/2)}*\text{FresnelC}(2^{(1/2)}/Pi^{(1/2)}/(b^2)^{(1/2)}*(b^2*x+a*b)) - 1/4/b^2*2^{(1/2)}*Pi^{(1/2)}/(b^2)^{(1/2)}*\text{FresnelS}(2^{(1/2)}/Pi^{(1/2)}/(b^2)^{(1/2)}*(b^2*x+a*b))$

**Maxima [C]** time = 2.78108, size = 346, normalized size = 3.49

$$\frac{abx\left(8ie^{(ib^2x^2+2iabx+ia^2)} - 8ie^{(-ib^2x^2-2iabx-ia^2)}\right) + a^2\left(8ie^{(ib^2x^2+2iabx+ia^2)} - 8ie^{(-ib^2x^2-2iabx-ia^2)}\right) + 2\sqrt{b^2x^2 + 2abx + a^2}}{4b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos((b\*x+a)^2),x, algorithm="maxima")

[Out]  $\frac{1}{16}*(a*b*x*(8*I*e^{(I*b^2*x^2 + 2*I*a*b*x + I*a^2)} - 8*I*e^{(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)}) + a^2*(8*I*e^{(I*b^2*x^2 + 2*I*a*b*x + I*a^2)} - 8*I*e^{(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)}) + 2*sqrt(b^2*x^2 + 2*a*b*x + a^2)*((-I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(I*b^2*x^2 + 2*I*a*b*x + I*a^2)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)) - 1))*a^2 + (I + 1)*sqrt(2)*gamma(3/2, I*b^2*x^2 + 2*I*a*b*x + I*a^2) - (I - 1)*sqrt(2)*gamma(3/2, -I*b^2*x^2 - 2*I*a*b*x - I*a^2))/(b^4*x + a*b^3)$

**Fricas [A]** time = 1.66701, size = 278, normalized size = 2.81

$$\frac{2\sqrt{2}\pi a^2\sqrt{\frac{b^2}{\pi}}C\left(\frac{\sqrt{2}(bx+a)\sqrt{\frac{b^2}{\pi}}}{b}\right) - \sqrt{2}\pi\sqrt{\frac{b^2}{\pi}}S\left(\frac{\sqrt{2}(bx+a)\sqrt{\frac{b^2}{\pi}}}{b}\right) + 2(b^2x - ab)\sin(b^2x^2 + 2abx + a^2)}{4b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos((b\*x+a)^2),x, algorithm="fricas")

[Out]  $\frac{1}{4}*(2*sqrt(2)*pi*a^2*sqrt(b^2/pi)*fresnel\_cos(sqrt(2)*(b*x + a)*sqrt(b^2/pi)/b) - sqrt(2)*pi*sqrt(b^2/pi)*fresnel\_sin(sqrt(2)*(b*x + a)*sqrt(b^2/pi)/b) + 2*(b^2*x - a*b)*sin(b^2*x^2 + 2*a*b*x + a^2))/b^4$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^2 \cos(a^2 + 2abx + b^2x^2) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*2\*cos((b\*x+a)\*\*2),x)

[Out] Integral(x\*\*2\*cos(a\*\*2 + 2\*a\*b\*x + b\*\*2\*x\*\*2), x)

**Giac [C]** time = 1.13045, size = 215, normalized size = 2.17

$$\frac{(i+1)\sqrt{2}\sqrt{\pi}(2a^2+i)\operatorname{erf}\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\left(x+\frac{a}{b}\right)|b|}{|b|} + \frac{4\left(ib\left(x+\frac{a}{b}\right)-2ia\right)e^{(ib^2x^2+2iabx+ia^2)}}{b} - \frac{(i-1)\sqrt{2}\sqrt{\pi}(2a^2-i)\operatorname{erf}\left(-\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\left(x+\frac{a}{b}\right)|b|}{|b|} + \frac{4\left(-ib\left(x+\frac{a}{b}\right)+2ia\right)e^{(-ib^2x^2-2iabx-ia^2)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos((b\*x+a)^2),x, algorithm="giac")

[Out]  $-1/16*((I + 1)*sqrt(2)*sqrt(pi)*(2*a^2 + I)*erf((1/2*I - 1/2)*sqrt(2)*(x + a/b)*abs(b))/abs(b) + 4*(I*b*(x + a/b) - 2*I*a)*e^{(I*b^2*x^2 + 2*I*a*b*x + I*a^2)/b}/b^2 - 1/16*(-(I - 1)*sqrt(2)*sqrt(pi)*(2*a^2 - I)*erf(-(1/2*I + 1/2)*sqrt(2)*(x + a/b)*abs(b))/abs(b) + 4*(-I*b*(x + a/b) + 2*I*a)*e^{(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)/b}/b^2$



### 3.86 $\int x \cos((a + bx)^2) dx$

**Optimal.** Leaf size=47

$$\frac{\sin((a + bx)^2)}{2b^2} - \frac{\sqrt{\frac{\pi}{2}} a \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^2}$$

[Out]  $-\left(\frac{a \sqrt{\pi/2} \operatorname{FresnelC}\left[\sqrt{2/\pi}(a + b*x)\right]}{b^2}\right) + \frac{\sin[(a + b*x)^2]}{2*b^2}$

**Rubi [A]** time = 0.0322894, antiderivative size = 47, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 4, integrand size = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.4$ , Rules used = {3434, 3352, 3380, 2637}

$$\frac{\sin((a + bx)^2)}{2b^2} - \frac{\sqrt{\frac{\pi}{2}} a \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^2}$$

Antiderivative was successfully verified.

[In] Int[x\*Cos[(a + b\*x)^2], x]

[Out]  $-\left(\frac{a \sqrt{\pi/2} \operatorname{FresnelC}\left[\sqrt{2/\pi}(a + b*x)\right]}{b^2}\right) + \frac{\sin[(a + b*x)^2]}{2*b^2}$

#### Rule 3434

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] :> Module[{k = If[FractionQ[n], Denominator[n], 1]}, Dist[k/f^(m + 1), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x^(k*n)])^p, x^(k - 1)*(f*g - e*h + h*x^k)^m, x], x], x, (e + f*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && IGtQ[p, 0] && IGtQ[m, 0]
```

#### Rule 3352

```
Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] :> Simp[(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)])/(f*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]
```

#### Rule 3380

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

#### Rule 2637

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] :> Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

#### Rubi steps

$$\begin{aligned}
\int x \cos((a + bx)^2) dx &= \frac{\text{Subst}\left(\int (-a \cos(x^2) + x \cos(x^2)) dx, x, a + bx\right)}{b^2} \\
&= \frac{\text{Subst}\left(\int x \cos(x^2) dx, x, a + bx\right)}{b^2} - \frac{a \text{Subst}\left(\int \cos(x^2) dx, x, a + bx\right)}{b^2} \\
&= -\frac{a\sqrt{\frac{\pi}{2}}C\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^2} + \frac{\text{Subst}\left(\int \cos(x) dx, x, (a + bx)^2\right)}{2b^2} \\
&= -\frac{a\sqrt{\frac{\pi}{2}}C\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^2} + \frac{\sin((a + bx)^2)}{2b^2}
\end{aligned}$$

**Mathematica [A]** time = 0.0469997, size = 42, normalized size = 0.89

$$\frac{\sin((a + bx)^2) - \sqrt{2\pi}a \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{2b^2}$$

Antiderivative was successfully verified.

[In] Integrate[x\*Cos[(a + b\*x)^2], x]

[Out]  $(-(a*\text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*(a + b*x)]) + \text{Sin}[(a + b*x)^2])/(2*b^2)$

**Maple [A]** time = 0.028, size = 63, normalized size = 1.3

$$\frac{\sin(x^2b^2 + 2abx + a^2)}{2b^2} - \frac{\sqrt{2}a\sqrt{\pi}}{2b} \text{FresnelC}\left(\frac{\sqrt{2}(b^2x + ab)}{\sqrt{\pi}} \frac{1}{\sqrt{b^2}}\right) \frac{1}{\sqrt{b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x\*cos((b\*x+a)^2), x)

[Out]  $1/2/b^2*\sin(b^2*x^2+2*a*b*x+a^2)-1/2*a/b*2^{(1/2)}*Pi^{(1/2)}/(b^2)^{(1/2)}*\text{FresnelC}(2^{(1/2)}/Pi^{(1/2)}/(b^2)^{(1/2)}*(b^2*x+a*b))$

**Maxima [C]** time = 2.37108, size = 266, normalized size = 5.66

$$\frac{bx\left(4ie^{(ib^2x^2+2iabx+ia^2)} - 4ie^{(-ib^2x^2-2iabx-ia^2)}\right) + 2\sqrt{b^2x^2 + 2abx + a^2}\left(- (i-1)\sqrt{2}\sqrt{\pi}\left(\text{erf}\left(\sqrt{ib^2x^2 + 2iabx + ia^2}\right) - 1\right)\right)}{16(b^3x + ab^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos((b\*x+a)^2), x, algorithm="maxima")

[Out]  $-1/16*(b*x*(4*I*e^{(I*b^2*x^2 + 2*I*a*b*x + I*a^2)} - 4*I*e^{(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)}) + 2*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2)*(-(I - 1)*\text{sqrt}(2)*\text{sqrt}(pi))*(\text{erf}(\text{sqrt}(I*b^2*x^2 + 2*I*a*b*x + I*a^2))) - 1) + (I + 1)*\text{sqrt}(2)*\text{sqrt}(pi)*(\text{erf}(\text{sqrt}(-I*b^2*x^2 - 2*I*a*b*x - I*a^2))) - 1)*a + a*(4*I*e^{(I*b^2*x^2 + 2*I*a*b*x + I*a^2)} - 4*I*e^{(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)})/(b^3*x + a$

$b^2$ )

**Fricas [A]** time = 1.64091, size = 158, normalized size = 3.36

$$\frac{\sqrt{2}\pi a\sqrt{\frac{b^2}{\pi}} C\left(\frac{\sqrt{2}(bx+a)\sqrt{\frac{b^2}{\pi}}}{b}\right) - b \sin(b^2x^2 + 2abx + a^2)}{2b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos((b\*x+a)^2),x, algorithm="fricas")

[Out]  $-1/2*(\sqrt{2}*\pi*a*\sqrt{b^2/\pi}*\text{fresnel\_cos}(\sqrt{2}*(b*x + a)*\sqrt{b^2/\pi})/b - b*\sin(b^2*x^2 + 2*a*b*x + a^2))/b^3$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x \cos(a^2 + 2abx + b^2x^2) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos((b\*x+a)\*\*2),x)

[Out] Integral(x\*cos(a\*\*2 + 2\*a\*b\*x + b\*\*2\*x\*\*2), x)

**Giac [C]** time = 1.14234, size = 161, normalized size = 3.43

$$\frac{\frac{(i+1)\sqrt{2}\sqrt{\pi}a \operatorname{erf}\left(\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\left(x+\frac{a}{b}\right)|b\right)}{|b|} + \frac{2ie^{(ib^2x^2+2iabx+ia^2)}}{b}}{8b} - \frac{\frac{(i-1)\sqrt{2}\sqrt{\pi}a \operatorname{erf}\left(-\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\left(x+\frac{a}{b}\right)|b\right)}{|b|} - \frac{2ie^{(-ib^2x^2-2iabx-ia^2)}}{b}}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos((b\*x+a)^2),x, algorithm="giac")

[Out]  $-1/8*(-(I + 1)*\sqrt{2}*\sqrt{\pi})*a*\operatorname{erf}((1/2*I - 1/2)*\sqrt{2}*(x + a/b)*\operatorname{abs}(b))/\operatorname{abs}(b) + 2*I*e^{(I*b^2*x^2 + 2*I*a*b*x + I*a^2)/b}/b - 1/8*((I - 1)*\sqrt{2}*\sqrt{\pi})*a*\operatorname{erf}(-(1/2*I + 1/2)*\sqrt{2}*(x + a/b)*\operatorname{abs}(b))/\operatorname{abs}(b) - 2*I*e^{(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)/b}/b$

### 3.87 $\int \cos((a + bx)^2) dx$

**Optimal.** Leaf size=29

$$\frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b}$$

[Out] (Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*(a + b\*x)])/b

**Rubi [A]** time = 0.0058369, antiderivative size = 29, normalized size of antiderivative = 1., number of steps used = 1, number of rules used = 1, integrand size = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {3352}

$$\frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[(a + b\*x)^2], x]

[Out] (Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*(a + b\*x)])/b

#### Rule 3352

Int[Cos[(d\_.)\*((e\_.) + (f\_.)\*(x\_)) ^2], x\_Symbol] := Simp[(Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*Rt[d, 2]\*(e + f\*x)]]/(f\*Rt[d, 2]), x] /; FreeQ[{d, e, f}, x]

#### Rubi steps

$$\int \cos((a + bx)^2) dx = \frac{\sqrt{\frac{\pi}{2}} \operatorname{C}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b}$$

**Mathematica [A]** time = 0.0080303, size = 29, normalized size = 1.

$$\frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[(a + b\*x)^2], x]

[Out] (Sqrt[Pi/2]\*FresnelC[Sqrt[2/Pi]\*(a + b\*x)])/b

**Maple [A]** time = 0.028, size = 36, normalized size = 1.2

$$\frac{\sqrt{2}\sqrt{\pi}}{2} \operatorname{FresnelC}\left(\frac{\sqrt{2}(b^2x + ab)}{\sqrt{\pi}} \frac{1}{\sqrt{b^2}}\right) \frac{1}{\sqrt{b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos((b\*x+a)^2),x)

[Out]  $\frac{1}{2} \cdot 2^{(1/2)} \cdot \pi^{(1/2)} / (b^2)^{(1/2)} \cdot \text{FresnelC}(2^{(1/2)} / \pi^{(1/2)} / (b^2)^{(1/2)} \cdot (b^2 \cdot x + a \cdot b))$

**Maxima [C]** time = 1.76279, size = 113, normalized size = 3.9

$$\frac{\sqrt{\pi} \left( (i-1) \sqrt{2} \operatorname{erf} \left( -(-1)^{\frac{3}{4}} (i b x + i a) \right) + (i-1) \sqrt{2} \operatorname{erf} \left( -\left( \frac{1}{4} i - \frac{1}{4} \right) \sqrt{2} (2 i b x + 2 i a) \right) - (i+1) \sqrt{2} \operatorname{erf} \left( -\left( \frac{1}{4} i + \frac{1}{4} \right) \sqrt{2} (2 i b x + 2 i a) \right) + (i+1) \sqrt{2} \operatorname{erf} \left( (i b x + i a) / \sqrt{-1} \right) \right)}{16 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)^2),x, algorithm="maxima")

[Out]  $-1/16 \cdot \sqrt{\pi} \cdot ((I - 1) \cdot \sqrt{2} \cdot \operatorname{erf}(-(-1)^{(3/4)} \cdot (I \cdot b \cdot x + I \cdot a)) + (I - 1) \cdot \sqrt{2} \cdot \operatorname{erf}(-\left(\frac{1}{4} \cdot I - \frac{1}{4}\right) \cdot \sqrt{2} \cdot (2 \cdot I \cdot b \cdot x + 2 \cdot I \cdot a)) - (I + 1) \cdot \sqrt{2} \cdot \operatorname{erf}(-\left(\frac{1}{4} \cdot I + \frac{1}{4}\right) \cdot \sqrt{2} \cdot (2 \cdot I \cdot b \cdot x + 2 \cdot I \cdot a)) + (I + 1) \cdot \sqrt{2} \cdot \operatorname{erf}((I \cdot b \cdot x + I \cdot a) / \sqrt{-1})) / b$

**Fricas [A]** time = 1.54214, size = 107, normalized size = 3.69

$$\frac{\sqrt{2} \pi \sqrt{\frac{b^2}{\pi}} C \left( \frac{\sqrt{2} (b x + a) \sqrt{\frac{b^2}{\pi}}}{b} \right)}{2 b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)^2),x, algorithm="fricas")

[Out]  $\frac{1}{2} \cdot \sqrt{2} \cdot \sqrt{\pi} \cdot \sqrt{b^2 / \pi} \cdot \text{fresnel\_cos}(\sqrt{2} \cdot (b \cdot x + a) \cdot \sqrt{b^2 / \pi}) / b / b^2$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \cos((a + b x)^2) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)\*\*2),x)

[Out] Integral(cos((a + b\*x)\*\*2), x)

**Giac [C]** time = 1.09037, size = 74, normalized size = 2.55

$$-\frac{(i+1) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left( \left( \frac{1}{2} i - \frac{1}{2} \right) \sqrt{2} \left( x + \frac{a}{b} \right) |b| \right)}{8 |b|} + \frac{(i-1) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left( -\left( \frac{1}{2} i + \frac{1}{2} \right) \sqrt{2} \left( x + \frac{a}{b} \right) |b| \right)}{8 |b|}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos((b*x+a)^2),x, algorithm="giac")
```

```
[Out] -(1/8*I + 1/8)*sqrt(2)*sqrt(pi)*erf((1/2*I - 1/2)*sqrt(2)*(x + a/b)*abs(b))  
/abs(b) + (1/8*I - 1/8)*sqrt(2)*sqrt(pi)*erf(-(1/2*I + 1/2)*sqrt(2)*(x + a/  
b)*abs(b))/abs(b)
```

$$3.88 \quad \int \frac{\cos((a+bx)^2)}{x} dx$$

**Optimal.** Leaf size=14

$$\text{Unintegrable}\left(\frac{\cos((a+bx)^2)}{x}, x\right)$$

[Out] Unintegrable[Cos[(a + b\*x)^2]/x, x]

**Rubi [A]** time = 0.0084172, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.$ , Rules used = {}

$$\int \frac{\cos((a+bx)^2)}{x} dx$$

Verification is Not applicable to the result.

[In] Int[Cos[(a + b\*x)^2]/x,x]

[Out] Defer[Int][Cos[(a + b\*x)^2]/x, x]

Rubi steps

$$\int \frac{\cos((a+bx)^2)}{x} dx = \int \frac{\cos((a+bx)^2)}{x} dx$$

**Mathematica [A]** time = 2.66219, size = 0, normalized size = 0.

$$\int \frac{\cos((a+bx)^2)}{x} dx$$

Verification is Not applicable to the result.

[In] Integrate[Cos[(a + b\*x)^2]/x,x]

[Out] Integrate[Cos[(a + b\*x)^2]/x, x]

**Maple [A]** time = 0.124, size = 0, normalized size = 0.

$$\int \frac{\cos((bx+a)^2)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos((b\*x+a)^2)/x,x)

[Out] int(cos((b\*x+a)^2)/x,x)

---

**Maxima [A]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos((bx + a)^2)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)^2)/x,x, algorithm="maxima")

[Out] integrate(cos((b\*x + a)^2)/x, x)

---

**Fricas [A]** time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\frac{\cos(b^2x^2 + 2abx + a^2)}{x}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)^2)/x,x, algorithm="fricas")

[Out] integral(cos(b^2\*x^2 + 2\*a\*b\*x + a^2)/x, x)

---

**Sympy [A]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a^2 + 2abx + b^2x^2)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)\*\*2)/x,x)

[Out] Integral(cos(a\*\*2 + 2\*a\*b\*x + b\*\*2\*x\*\*2)/x, x)

---

**Giac [A]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos((bx + a)^2)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)^2)/x,x, algorithm="giac")

[Out] integrate(cos((b\*x + a)^2)/x, x)



$$3.89 \quad \int \frac{\cos((a+bx)^2)}{x^2} dx$$

**Optimal.** Leaf size=14

$$\text{Unintegrable}\left(\frac{\cos((a+bx)^2)}{x^2}, x\right)$$

[Out] Unintegrable[Cos[(a + b\*x)^2]/x^2, x]

**Rubi [A]** time = 0.0085267, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.$ , Rules used = {}

$$\int \frac{\cos((a+bx)^2)}{x^2} dx$$

Verification is Not applicable to the result.

[In] Int[Cos[(a + b\*x)^2]/x^2,x]

[Out] Defer[Int][Cos[(a + b\*x)^2]/x^2, x]

Rubi steps

$$\int \frac{\cos((a+bx)^2)}{x^2} dx = \int \frac{\cos((a+bx)^2)}{x^2} dx$$

**Mathematica [A]** time = 5.28803, size = 0, normalized size = 0.

$$\int \frac{\cos((a+bx)^2)}{x^2} dx$$

Verification is Not applicable to the result.

[In] Integrate[Cos[(a + b\*x)^2]/x^2,x]

[Out] Integrate[Cos[(a + b\*x)^2]/x^2, x]

**Maple [A]** time = 0.116, size = 0, normalized size = 0.

$$\int \frac{\cos((bx+a)^2)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos((b\*x+a)^2)/x^2,x)

[Out] int(cos((b\*x+a)^2)/x^2,x)

---

**Maxima [A]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos((bx + a)^2)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)^2)/x^2,x, algorithm="maxima")

[Out] integrate(cos((b\*x + a)^2)/x^2, x)

---

**Fricas [A]** time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\frac{\cos(b^2x^2 + 2abx + a^2)}{x^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)^2)/x^2,x, algorithm="fricas")

[Out] integral(cos(b^2\*x^2 + 2\*a\*b\*x + a^2)/x^2, x)

---

**Sympy [A]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a^2 + 2abx + b^2x^2)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)\*\*2)/x\*\*2,x)

[Out] Integral(cos(a\*\*2 + 2\*a\*b\*x + b\*\*2\*x\*\*2)/x\*\*2, x)

---

**Giac [A]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos((bx + a)^2)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos((b\*x+a)^2)/x^2,x, algorithm="giac")

[Out] integrate(cos((b\*x + a)^2)/x^2, x)

### 3.90 $\int x^2 \cos(a + b\sqrt{c + dx}) dx$

**Optimal.** Leaf size=346

$$\frac{2c^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} - \frac{40(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{b^3 d^3} + \frac{24c\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^3 d^3} + \frac{240\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^5 d^3}$$

```
[Out] (240*Cos[a + b*Sqrt[c + d*x]])/(b^6*d^3) + (24*c*Cos[a + b*Sqrt[c + d*x]])/(b^4*d^3) + (2*c^2*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^3) - (120*(c + d*x)*Cos[a + b*Sqrt[c + d*x]])/(b^4*d^3) - (12*c*(c + d*x)*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^3) + (10*(c + d*x)^2*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^3) + (240*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b^5*d^3) + (24*c*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b^3*d^3) + (2*c^2*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b*d^3) - (40*(c + d*x)^(3/2)*Sin[a + b*Sqrt[c + d*x]])/(b^3*d^3) - (4*c*(c + d*x)^(3/2)*Sin[a + b*Sqrt[c + d*x]])/(b*d^3) + (2*(c + d*x)^(5/2)*Sin[a + b*Sqrt[c + d*x]])/(b*d^3)
```

**Rubi [A]** time = 0.304552, antiderivative size = 346, normalized size of antiderivative = 1., number of steps used = 14, number of rules used = 3, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3432, 3296, 2638}

$$\frac{2c^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} - \frac{40(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{b^3 d^3} + \frac{24c\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^3 d^3} + \frac{240\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^5 d^3}$$

Antiderivative was successfully verified.

```
[In] Int[x^2*Cos[a + b*Sqrt[c + d*x]],x]
```

```
[Out] (240*Cos[a + b*Sqrt[c + d*x]])/(b^6*d^3) + (24*c*Cos[a + b*Sqrt[c + d*x]])/(b^4*d^3) + (2*c^2*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^3) - (120*(c + d*x)*Cos[a + b*Sqrt[c + d*x]])/(b^4*d^3) - (12*c*(c + d*x)*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^3) + (10*(c + d*x)^2*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^3) + (240*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b^5*d^3) + (24*c*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b^3*d^3) + (2*c^2*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b*d^3) - (40*(c + d*x)^(3/2)*Sin[a + b*Sqrt[c + d*x]])/(b^3*d^3) - (4*c*(c + d*x)^(3/2)*Sin[a + b*Sqrt[c + d*x]])/(b*d^3) + (2*(c + d*x)^(5/2)*Sin[a + b*Sqrt[c + d*x]])/(b*d^3)
```

#### Rule 3432

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Dist[1/(n*f), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - (e*h)/f + (h*x^(1/n))/f)^m, x], x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

#### Rule 3296

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := -Simp[(c + d*x)^m*Cos[e + f*x]/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

#### Rule 2638

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int x^2 \cos(a + b\sqrt{c + dx}) dx &= \frac{2 \operatorname{Subst}\left(\int \left(\frac{c^2 x \cos(a+bx)}{d^2} - \frac{2cx^3 \cos(a+bx)}{d^2} + \frac{x^5 \cos(a+bx)}{d^2}\right) dx, x, \sqrt{c + dx}\right)}{d} \\
&= \frac{2 \operatorname{Subst}\left(\int x^5 \cos(a + bx) dx, x, \sqrt{c + dx}\right)}{d^3} - \frac{(4c) \operatorname{Subst}\left(\int x^3 \cos(a + bx) dx, x, \sqrt{c + dx}\right)}{d^3} \\
&= \frac{2c^2 \sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd^3} - \frac{4c(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{bd^3} + \frac{2(c + dx)^{5/2} \sin(a + b\sqrt{c + dx})}{bd^3} \\
&= \frac{2c^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} - \frac{12c(c + dx) \cos(a + b\sqrt{c + dx})}{b^2 d^3} + \frac{10(c + dx)^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} \\
&= \frac{2c^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} - \frac{12c(c + dx) \cos(a + b\sqrt{c + dx})}{b^2 d^3} + \frac{10(c + dx)^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} \\
&= \frac{24c \cos(a + b\sqrt{c + dx})}{b^4 d^3} + \frac{2c^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} - \frac{120(c + dx) \cos(a + b\sqrt{c + dx})}{b^4 d^3} - \frac{120(c + dx)^2 \cos(a + b\sqrt{c + dx})}{b^4 d^3} \\
&= \frac{24c \cos(a + b\sqrt{c + dx})}{b^4 d^3} + \frac{2c^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} - \frac{120(c + dx) \cos(a + b\sqrt{c + dx})}{b^4 d^3} - \frac{120(c + dx)^2 \cos(a + b\sqrt{c + dx})}{b^4 d^3} \\
&= \frac{240 \cos(a + b\sqrt{c + dx})}{b^6 d^3} + \frac{24c \cos(a + b\sqrt{c + dx})}{b^4 d^3} + \frac{2c^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} - \frac{120(c + dx) \cos(a + b\sqrt{c + dx})}{b^4 d^3} - \frac{120(c + dx)^2 \cos(a + b\sqrt{c + dx})}{b^4 d^3}
\end{aligned}$$

**Mathematica [C]** time = 0.841162, size = 224, normalized size = 0.65

$$\frac{e^{-i(a+b\sqrt{c+dx})} \left( -ib^5 d^2 x^2 \sqrt{c+dx} + b^4 dx(4c+5dx) + 4ib^3 \sqrt{c+dx}(2c+5dx) - 12b^2(4c+5dx) - 120ib\sqrt{c+dx} + 120 \right) e^{2i(a+b\sqrt{c+dx})}}{b^6 d^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2\*Cos[a + b\*Sqrt[c + d\*x]],x]

[Out] (120 + (120\*I)\*b\*Sqrt[c + d\*x] + I\*b^5\*d^2\*x^2\*Sqrt[c + d\*x] - (4\*I)\*b^3\*Sqrt[c + d\*x]\*(2\*c + 5\*d\*x) - 12\*b^2\*(4\*c + 5\*d\*x) + b^4\*d\*x\*(4\*c + 5\*d\*x) + E^((2\*I)\*(a + b\*Sqrt[c + d\*x]))\*(120 - (120\*I)\*b\*Sqrt[c + d\*x] - I\*b^5\*d^2\*x^2\*Sqrt[c + d\*x] + (4\*I)\*b^3\*Sqrt[c + d\*x]\*(2\*c + 5\*d\*x) - 12\*b^2\*(4\*c + 5\*d\*x) + b^4\*d\*x\*(4\*c + 5\*d\*x)))/(b^6\*d^3\*E^(I\*(a + b\*Sqrt[c + d\*x])))

**Maple [B]** time = 0.089, size = 825, normalized size = 2.4

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2\*cos(a+b\*(d\*x+c)^(1/2)),x)

[Out] 2/d^3/b^2\*(c^2\*(cos(a+b\*(d\*x+c)^(1/2))+(a+b\*(d\*x+c)^(1/2))\*sin(a+b\*(d\*x+c)^(1/2)))-a\*c^2\*sin(a+b\*(d\*x+c)^(1/2))-2/b^2\*c\*((a+b\*(d\*x+c)^(1/2))^3\*sin(a+b\*(d\*x+c)^(1/2))+3\*(a+b\*(d\*x+c)^(1/2))^2\*cos(a+b\*(d\*x+c)^(1/2))-6\*cos(a+b\*(d\*x+c)^(1/2))-6\*(a+b\*(d\*x+c)^(1/2))\*sin(a+b\*(d\*x+c)^(1/2)))+6/b^2\*a\*c\*((a+b\*(d\*x+c)^(1/2))^2\*sin(a+b\*(d\*x+c)^(1/2))-2\*sin(a+b\*(d\*x+c)^(1/2))+2\*(a+b\*(d\*x+c)^(1/2))\*cos(a+b\*(d\*x+c)^(1/2)))-6/b^2\*a^2\*c\*(cos(a+b\*(d\*x+c)^(1/2))+(a+b\*(d\*x+c)^(1/2))\*sin(a+b\*(d\*x+c)^(1/2)))+2/b^2\*a^3\*c\*sin(a+b\*(d\*x+c)^(1/2))+1/b^4\*((a+b\*(d\*x+c)^(1/2))^5\*sin(a+b\*(d\*x+c)^(1/2))+5\*(a+b\*(d\*x+c)^(1/2))^4\*cos(a+b\*(d\*x+c)^(1/2))-20\*(a+b\*(d\*x+c)^(1/2))^3\*sin(a+b\*(d\*x+c)^(1/2))-60

$$\begin{aligned} &*(a+b*(d*x+c)^{(1/2)})^2*\cos(a+b*(d*x+c)^{(1/2)})+120*\cos(a+b*(d*x+c)^{(1/2)})+12 \\ &0*(a+b*(d*x+c)^{(1/2)})*\sin(a+b*(d*x+c)^{(1/2)})-5/b^4*a*((a+b*(d*x+c)^{(1/2)})^4 \\ &4*\sin(a+b*(d*x+c)^{(1/2)})+4*(a+b*(d*x+c)^{(1/2)})^3*\cos(a+b*(d*x+c)^{(1/2)})-12* \\ &(a+b*(d*x+c)^{(1/2)})^2*\sin(a+b*(d*x+c)^{(1/2)})+24*\sin(a+b*(d*x+c)^{(1/2)})-24*( \\ &a+b*(d*x+c)^{(1/2)})*\cos(a+b*(d*x+c)^{(1/2)}))+10/b^4*a^2*((a+b*(d*x+c)^{(1/2)})^3 \\ &3*\sin(a+b*(d*x+c)^{(1/2)})+3*(a+b*(d*x+c)^{(1/2)})^2*\cos(a+b*(d*x+c)^{(1/2)})-6*\cos \\ &(a+b*(d*x+c)^{(1/2)})-6*(a+b*(d*x+c)^{(1/2)})*\sin(a+b*(d*x+c)^{(1/2)}))-10/b^4*a \\ &a^3*((a+b*(d*x+c)^{(1/2)})^2*\sin(a+b*(d*x+c)^{(1/2)})-2*\sin(a+b*(d*x+c)^{(1/2)})+ \\ &2*(a+b*(d*x+c)^{(1/2)})*\cos(a+b*(d*x+c)^{(1/2)}))+5/b^4*a^4*(\cos(a+b*(d*x+c)^{(1/2)}) \\ &)+(a+b*(d*x+c)^{(1/2)})*\sin(a+b*(d*x+c)^{(1/2)}))-1/b^4*a^5*\sin(a+b*(d*x+c)^{(1/2)}) \end{aligned}$$

**Maxima [B]** time = 1.24188, size = 907, normalized size = 2.62

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="maxima")

[Out] 
$$\begin{aligned} &-2*(a*c^2*\sin(\sqrt{d*x+c})*b+a)-((\sqrt{d*x+c})*b+a)*\sin(\sqrt{d*x+c}) \\ &+ \cos(\sqrt{d*x+c})*b+a)+\cos(\sqrt{d*x+c})*b+a)*c^2-2*a^3*c*\sin(\sqrt{d*x+c})*b+a \\ &)/b^2+6*((\sqrt{d*x+c})*b+a)*\sin(\sqrt{d*x+c})*b+a)+\cos(\sqrt{d*x+c})*b+a) \\ &)*a^2*c/b^2+a^5*\sin(\sqrt{d*x+c})*b+a)/b^4-5*((\sqrt{d*x+c})*b+a) \\ &)*\sin(\sqrt{d*x+c})*b+a)+\cos(\sqrt{d*x+c})*b+a)*a^4/b^4-6*(2 \\ &*(\sqrt{d*x+c})*b+a)*\cos(\sqrt{d*x+c})*b+a)+((\sqrt{d*x+c})*b+a)^2 \\ &-2)*\sin(\sqrt{d*x+c})*b+a)*a*c/b^2+10*(2*(\sqrt{d*x+c})*b+a)*\cos(\sqrt{d*x+c}) \\ &+((\sqrt{d*x+c})*b+a)^2-2)*\sin(\sqrt{d*x+c})*b+a) \\ &)*a^3/b^4+2*(3*((\sqrt{d*x+c})*b+a)^2-2)*\cos(\sqrt{d*x+c})*b+a)+ \\ &((\sqrt{d*x+c})*b+a)^3-6*\sqrt{d*x+c}*b-6*a)*\sin(\sqrt{d*x+c})*b+a) \\ &)*c/b^2-10*(3*((\sqrt{d*x+c})*b+a)^2-2)*\cos(\sqrt{d*x+c})*b+a)+ \\ &((\sqrt{d*x+c})*b+a)^3-6*\sqrt{d*x+c}*b-6*a)*\sin(\sqrt{d*x+c})*b+a) \\ &)*a^2/b^4+5*(4*((\sqrt{d*x+c})*b+a)^3-6*\sqrt{d*x+c}*b-6*a)*\cos(\sqrt{d*x+c}) \\ &+((\sqrt{d*x+c})*b+a)^4-12*(\sqrt{d*x+c})*b+a)^2+ \\ &24)*\sin(\sqrt{d*x+c})*b+a)*a/b^4-5*((\sqrt{d*x+c})*b+a)^4-12*(\sqrt{d*x+c}) \\ &+24)*\cos(\sqrt{d*x+c})*b+a)+((\sqrt{d*x+c})*b+a)^5-20*(\sqrt{d*x+c})*b+a)^3 \\ &+120*\sqrt{d*x+c}*b+120*a)*\sin(\sqrt{d*x+c})*b+a)/b^4)/(b^2*d^3) \end{aligned}$$

**Fricas [A]** time = 1.62633, size = 248, normalized size = 0.72

$$\frac{2\left(\left(b^5 d^2 x^2 - 20 b^3 d x - 8 b^3 c + 120 b\right) \sqrt{d x + c} \sin\left(\sqrt{d x + c} b + a\right) + \left(5 b^4 d^2 x^2 - 48 b^2 c + 4\left(b^4 c - 15 b^2\right) d x + 120\right) \cos\left(\sqrt{d x + c} b + a\right)\right)}{b^6 d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="fricas")

[Out] 
$$\begin{aligned} &2*((b^5*d^2*x^2-20*b^3*d*x-8*b^3*c+120*b)*\sqrt{d*x+c}*\sin(\sqrt{d*x+c})*b+a) \\ &+(5*b^4*d^2*x^2-48*b^2*c+4*(b^4*c-15*b^2)*d*x+120)*\cos(\sqrt{d*x+c})*b+a) \\ &)/(b^6*d^3) \end{aligned}$$

**Sympy [A]** time = 2.36067, size = 274, normalized size = 0.79

$$\left\{ \begin{array}{l} \frac{x^3 \cos(a)}{3} \\ \frac{x^3 \cos(a+b\sqrt{c})}{3} \\ \frac{x^3 \cos(a)}{3} \\ \frac{2x^2\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{bd} + \frac{8cx \cos(a+b\sqrt{c+dx})}{b^2d^2} + \frac{10x^2 \cos(a+b\sqrt{c+dx})}{b^2d} - \frac{16c\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3d^3} - \frac{40x\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3d^2} - \frac{96c \cos(a+b\sqrt{c+dx})}{b^4d^3} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*2\*cos(a+b\*(d\*x+c)\*\*(1/2)),x)

[Out] Piecewise((x\*\*3\*cos(a)/3, Eq(b, 0) & Eq(d, 0)), (x\*\*3\*cos(a + b\*sqrt(c))/3, Eq(d, 0)), (x\*\*3\*cos(a)/3, Eq(b, 0)), (2\*x\*\*2\*sqrt(c + d\*x)\*sin(a + b\*sqrt(c + d\*x))/(b\*d) + 8\*c\*x\*cos(a + b\*sqrt(c + d\*x))/(b\*\*2\*d\*\*2) + 10\*x\*\*2\*cos(a + b\*sqrt(c + d\*x))/(b\*\*2\*d) - 16\*c\*sqrt(c + d\*x)\*sin(a + b\*sqrt(c + d\*x))/(b\*\*3\*d\*\*3) - 40\*x\*sqrt(c + d\*x)\*sin(a + b\*sqrt(c + d\*x))/(b\*\*3\*d\*\*2) - 96\*c\*cos(a + b\*sqrt(c + d\*x))/(b\*\*4\*d\*\*3) - 120\*x\*cos(a + b\*sqrt(c + d\*x))/(b\*\*4\*d\*\*2) + 240\*sqrt(c + d\*x)\*sin(a + b\*sqrt(c + d\*x))/(b\*\*5\*d\*\*3) + 240\*cos(a + b\*sqrt(c + d\*x))/(b\*\*6\*d\*\*3), True))

**Giac [B]** time = 1.39436, size = 1134, normalized size = 3.28

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="giac")

[Out] 2\*((b^4\*c^2\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - 6\*(sqrt(d\*x + c)\*b + a)^2\*b^2\*c\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + 12\*(sqrt(d\*x + c)\*b + a)\*a\*b^2\*c\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - 6\*a^2\*b^2\*c\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + 5\*(sqrt(d\*x + c)\*b + a)^4\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - 20\*(sqrt(d\*x + c)\*b + a)^3\*a\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + 30\*(sqrt(d\*x + c)\*b + a)^2\*a^2\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - 20\*(sqrt(d\*x + c)\*b + a)\*a^3\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + 5\*a^4\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + 12\*b^2\*c\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - 60\*(sqrt(d\*x + c)\*b + a)^2\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + 120\*(sqrt(d\*x + c)\*b + a)\*a\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - 60\*a^2\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + 120\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b))\*cos(-(sqrt(d\*x + c)\*b + a)\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + a\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - a)/(b^3\*d^2) - ((sqrt(d\*x + c)\*b + a)\*b^4\*c^2 - a\*b^4\*c^2 - 2\*(sqrt(d\*x + c)\*b + a)^3\*b^2\*c + 6\*(sqrt(d\*x + c)\*b + a)^2\*a\*b^2\*c - 6\*(sqrt(d\*x + c)\*b + a)\*a^2\*b^2\*c + 2\*a^3\*b^2\*c + (sqrt(d\*x + c)\*b + a)^5 - 5\*(sqrt(d\*x + c)\*b + a)^4\*a + 10\*(sqrt(d\*x + c)\*b + a)^3\*a^2 - 10\*(sqrt(d\*x + c)\*b + a)^2\*a^3 + 5\*(sqrt(d\*x + c)\*b + a)\*a^4 - a^5 + 12\*(sqrt(d\*x + c)\*b + a)\*b^2\*c - 12\*a\*b^2\*c - 20\*(sqrt(d\*x + c)\*b + a)^3 + 60\*(sqrt(d\*x + c)\*b + a)^2\*a - 60\*(sqrt(d\*x + c)\*b + a)\*a^2 + 20\*a^3 + 120\*sqrt(d\*x + c)\*b)\*sin(-(sqrt(d\*x + c)\*b + a)\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + a\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - a)/(b^3\*d^2))/(b^3\*d)

### 3.91 $\int x \cos(a + b\sqrt{c + dx}) dx$

**Optimal.** Leaf size=167

$$-\frac{12\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3d^2} + \frac{6(c+dx) \cos(a+b\sqrt{c+dx})}{b^2d^2} - \frac{2c \cos(a+b\sqrt{c+dx})}{b^2d^2} - \frac{12 \cos(a+b\sqrt{c+dx})}{b^4d^2} + \dots$$

```
[Out] (-12*Cos[a + b*Sqrt[c + d*x]])/(b^4*d^2) - (2*c*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^2) + (6*(c + d*x)*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^2) - (12*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b^3*d^2) - (2*c*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b*d^2) + (2*(c + d*x)^(3/2)*Sin[a + b*Sqrt[c + d*x]])/(b*d^2)
```

**Rubi [A]** time = 0.13634, antiderivative size = 167, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 3, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {3432, 3296, 2638}

$$-\frac{12\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3d^2} + \frac{6(c+dx) \cos(a+b\sqrt{c+dx})}{b^2d^2} - \frac{2c \cos(a+b\sqrt{c+dx})}{b^2d^2} - \frac{12 \cos(a+b\sqrt{c+dx})}{b^4d^2} + \dots$$

Antiderivative was successfully verified.

```
[In] Int[x*Cos[a + b*Sqrt[c + d*x]], x]
```

```
[Out] (-12*Cos[a + b*Sqrt[c + d*x]])/(b^4*d^2) - (2*c*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^2) + (6*(c + d*x)*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^2) - (12*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b^3*d^2) - (2*c*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b*d^2) + (2*(c + d*x)^(3/2)*Sin[a + b*Sqrt[c + d*x]])/(b*d^2)
```

#### Rule 3432

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])*(b_.)^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Dist[1/(n*f), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - (e*h)/f + (h*x^(1/n))/f)^m, x], x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

#### Rule 3296

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := -Simp[((c + d*x)^m*Cos[e + f*x])/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

#### Rule 2638

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

#### Rubi steps

$$\begin{aligned}
\int x \cos(a + b\sqrt{c + dx}) dx &= \frac{2 \operatorname{Subst}\left(\int \left(-\frac{cx \cos(a+bx)}{d} + \frac{x^3 \cos(a+bx)}{d}\right) dx, x, \sqrt{c + dx}\right)}{d} \\
&= \frac{2 \operatorname{Subst}\left(\int x^3 \cos(a + bx) dx, x, \sqrt{c + dx}\right)}{d^2} - \frac{(2c) \operatorname{Subst}\left(\int x \cos(a + bx) dx, x, \sqrt{c + dx}\right)}{d^2} \\
&= -\frac{2c\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd^2} + \frac{2(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{bd^2} - \frac{6 \operatorname{Subst}\left(\int x^2 \sin(a + bx) dx, x, \sqrt{c + dx}\right)}{bd^2} \\
&= -\frac{2c \cos(a + b\sqrt{c + dx})}{b^2 d^2} + \frac{6(c + dx) \cos(a + b\sqrt{c + dx})}{b^2 d^2} - \frac{2c\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd^2} + \frac{6(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{bd^2} \\
&= -\frac{2c \cos(a + b\sqrt{c + dx})}{b^2 d^2} + \frac{6(c + dx) \cos(a + b\sqrt{c + dx})}{b^2 d^2} - \frac{12\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^3 d^2} + \frac{12(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{b^3 d^2} \\
&= -\frac{12 \cos(a + b\sqrt{c + dx})}{b^4 d^2} - \frac{2c \cos(a + b\sqrt{c + dx})}{b^2 d^2} + \frac{6(c + dx) \cos(a + b\sqrt{c + dx})}{b^2 d^2} - \frac{12\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^3 d^2} + \frac{12(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{b^3 d^2}
\end{aligned}$$

**Mathematica [A]** time = 0.275562, size = 71, normalized size = 0.43

$$\frac{2(b(b^2 dx - 6)\sqrt{c + dx} \sin(a + b\sqrt{c + dx}) + (b^2(2c + 3dx) - 6) \cos(a + b\sqrt{c + dx}))}{b^4 d^2}$$

Antiderivative was successfully verified.

[In] Integrate[x\*Cos[a + b\*Sqrt[c + d\*x]],x]

[Out] (2\*((-6 + b^2\*(2\*c + 3\*d\*x))\*Cos[a + b\*Sqrt[c + d\*x]] + b\*Sqrt[c + d\*x]\*(-6 + b^2\*d\*x)\*Sin[a + b\*Sqrt[c + d\*x]])/(b^4\*d^2)

**Maple [A]** time = 0.031, size = 299, normalized size = 1.8

$$2 \frac{1}{b^2 d^2} \left( -c \left( \cos(a + b\sqrt{dx + c}) + (a + b\sqrt{dx + c}) \sin(a + b\sqrt{dx + c}) \right) + ca \sin(a + b\sqrt{dx + c}) + \frac{(a + b\sqrt{dx + c})^3 \sin(a + b\sqrt{dx + c})}{b^2 d^2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x\*cos(a+b\*(d\*x+c)^(1/2)),x)

[Out] 2/d^2/b^2\*(-c\*(cos(a+b\*(d\*x+c)^(1/2))+(a+b\*(d\*x+c)^(1/2))\*sin(a+b\*(d\*x+c)^(1/2)))+c\*a\*sin(a+b\*(d\*x+c)^(1/2))+1/b^2\*((a+b\*(d\*x+c)^(1/2))^3\*sin(a+b\*(d\*x+c)^(1/2))+3\*(a+b\*(d\*x+c)^(1/2))^2\*cos(a+b\*(d\*x+c)^(1/2))-6\*cos(a+b\*(d\*x+c)^(1/2))-6\*(a+b\*(d\*x+c)^(1/2))\*sin(a+b\*(d\*x+c)^(1/2)))-3/b^2\*a\*((a+b\*(d\*x+c)^(1/2))^2\*sin(a+b\*(d\*x+c)^(1/2))-2\*sin(a+b\*(d\*x+c)^(1/2))+2\*(a+b\*(d\*x+c)^(1/2))\*cos(a+b\*(d\*x+c)^(1/2)))+3\*a^2/b^2\*(cos(a+b\*(d\*x+c)^(1/2))+(a+b\*(d\*x+c)^(1/2))\*sin(a+b\*(d\*x+c)^(1/2)))-1/b^2\*a^3\*sin(a+b\*(d\*x+c)^(1/2))

**Maxima [A]** time = 1.25414, size = 355, normalized size = 2.13

$$2 \left( ac \sin(\sqrt{dx + cb} + a) - ((\sqrt{dx + cb} + a) \sin(\sqrt{dx + cb} + a) + \cos(\sqrt{dx + cb} + a))c - \frac{a^3 \sin(\sqrt{dx + cb} + a)}{b^2} + \frac{3((\sqrt{dx + cb} + a) \sin(\sqrt{dx + cb} + a) + \cos(\sqrt{dx + cb} + a))a^2}{b^2} \right)$$



Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="maxima")

[Out]  $2*(a*c*\sin(\sqrt{d*x + c}*b + a) - ((\sqrt{d*x + c}*b + a)*\sin(\sqrt{d*x + c})*b + a) + \cos(\sqrt{d*x + c}*b + a))*c - a^3*\sin(\sqrt{d*x + c}*b + a)/b^2 + 3*((\sqrt{d*x + c}*b + a)*\sin(\sqrt{d*x + c}*b + a) + \cos(\sqrt{d*x + c}*b + a))*a^2/b^2 - 3*(2*(\sqrt{d*x + c}*b + a)*\cos(\sqrt{d*x + c}*b + a) + ((\sqrt{d*x + c}*b + a)^2 - 2)*\sin(\sqrt{d*x + c}*b + a))*a/b^2 + (3*((\sqrt{d*x + c}*b + a)^2 - 2)*\cos(\sqrt{d*x + c}*b + a) + ((\sqrt{d*x + c}*b + a)^3 - 6*\sqrt{d*x + c}*b - 6*a)*\sin(\sqrt{d*x + c}*b + a))/b^2)/(b^2*d^2)$

**Fricas [A]** time = 1.63106, size = 166, normalized size = 0.99

$$\frac{2\left(\left(b^3 dx - 6b\right)\sqrt{dx + c} \sin\left(\sqrt{dx + cb} + a\right) + \left(3b^2 dx + 2b^2 c - 6\right) \cos\left(\sqrt{dx + cb} + a\right)\right)}{b^4 d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="fricas")

[Out]  $2*((b^3*d*x - 6*b)*\sqrt{d*x + c}*\sin(\sqrt{d*x + c}*b + a) + (3*b^2*d*x + 2*b^2*c - 6)*\cos(\sqrt{d*x + c}*b + a))/(b^4*d^2)$

**Sympy [A]** time = 0.684471, size = 151, normalized size = 0.9

$$\begin{cases} \frac{x^2 \cos(a)}{2} & \text{for } b = 0 \wedge (b = \\ \frac{x^2 \cos(a+b\sqrt{c})}{2} & \text{for } d = 0 \\ \frac{2x\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{bd} + \frac{4c \cos(a+b\sqrt{c+dx})}{b^2 d^2} + \frac{6x \cos(a+b\sqrt{c+dx})}{b^2 d} - \frac{12\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3 d^2} - \frac{12 \cos(a+b\sqrt{c+dx})}{b^4 d^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(a+b\*(d\*x+c)\*\*(1/2)),x)

[Out] Piecewise((x\*\*2\*cos(a)/2, Eq(b, 0) & (Eq(b, 0) | Eq(d, 0))), (x\*\*2\*cos(a + b\*sqrt(c))/2, Eq(d, 0)), (2\*x\*sqrt(c + d\*x)\*sin(a + b\*sqrt(c + d\*x))/(b\*d) + 4\*c\*cos(a + b\*sqrt(c + d\*x))/(b\*\*2\*d\*\*2) + 6\*x\*cos(a + b\*sqrt(c + d\*x))/(b\*\*2\*d) - 12\*sqrt(c + d\*x)\*sin(a + b\*sqrt(c + d\*x))/(b\*\*3\*d\*\*2) - 12\*cos(a + b\*sqrt(c + d\*x))/(b\*\*4\*d\*\*2), True))

**Giac [B]** time = 1.24896, size = 479, normalized size = 2.87

$$2 \left( \frac{\left(b^2 \operatorname{csgn}(\sqrt{dx+cb+a})b-ab\right) - 3\left(\sqrt{dx+cb+a}\right)^2 \operatorname{sgn}(\sqrt{dx+cb+a})b-ab + 6\left(\sqrt{dx+cb+a}\right) \operatorname{asgn}(\sqrt{dx+cb+a})b-ab - 3a^2 \operatorname{sgn}(\sqrt{dx+cb+a})b-ab + 6 \operatorname{sgn}(\sqrt{dx+cb+a})b-ab}{b} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="giac")

```
[Out] -2*((b^2*c*sgn((sqrt(d*x + c)*b + a)*b - a*b) - 3*(sqrt(d*x + c)*b + a)^2*sgn((sqrt(d*x + c)*b + a)*b - a*b) + 6*(sqrt(d*x + c)*b + a)*a*sgn((sqrt(d*x + c)*b + a)*b - a*b) - 3*a^2*sgn((sqrt(d*x + c)*b + a)*b - a*b) + 6*sgn((sqrt(d*x + c)*b + a)*b - a*b))*cos(-(sqrt(d*x + c)*b + a)*sgn((sqrt(d*x + c)*b + a)*b - a*b) + a*sgn((sqrt(d*x + c)*b + a)*b - a*b) - a)/b - ((sqrt(d*x + c)*b + a)*b^2*c - a*b^2*c - (sqrt(d*x + c)*b + a)^3 + 3*(sqrt(d*x + c)*b + a)^2*a - 3*(sqrt(d*x + c)*b + a)*a^2 + a^3 + 6*sqrt(d*x + c)*b)*sin(-(sqrt(d*x + c)*b + a)*sgn((sqrt(d*x + c)*b + a)*b - a*b) + a*sgn((sqrt(d*x + c)*b + a)*b - a*b) - a)/b)/(b^3*d^2)
```

### 3.92 $\int \cos(a + b\sqrt{c + dx}) dx$

**Optimal.** Leaf size=54

$$\frac{2 \cos(a + b\sqrt{c + dx})}{b^2 d} + \frac{2\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd}$$

[Out] (2\*Cos[a + b\*Sqrt[c + d\*x]])/(b^2\*d) + (2\*Sqrt[c + d\*x]\*Sin[a + b\*Sqrt[c + d\*x]])/(b\*d)

**Rubi [A]** time = 0.0274606, antiderivative size = 54, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3362, 3296, 2638}

$$\frac{2 \cos(a + b\sqrt{c + dx})}{b^2 d} + \frac{2\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*Sqrt[c + d\*x]],x]

[Out] (2\*Cos[a + b\*Sqrt[c + d\*x]])/(b^2\*d) + (2\*Sqrt[c + d\*x]\*Sin[a + b\*Sqrt[c + d\*x]])/(b\*d)

#### Rule 3362

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)])\*(b\_.)^(p\_.), x\_Symbol] := Dist[1/(n\*f), Subst[Int[x^(1/n - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, (e + f\*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]

#### Rule 3296

Int[((c\_.) + (d\_.)\*(x\_))^(m\_.)\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] := -Simp[((c + d\*x)^m\*Cos[e + f\*x])/f, x] + Dist[(d\*m)/f, Int[(c + d\*x)^(m - 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

#### Rule 2638

Int[sin[(c\_.) + (d\_.)\*(x\_)], x\_Symbol] := -Simp[Cos[c + d\*x]/d, x] /; FreeQ[{c, d}, x]

#### Rubi steps

$$\begin{aligned} \int \cos(a + b\sqrt{c + dx}) dx &= \frac{2 \text{Subst}\left(\int x \cos(a + bx) dx, x, \sqrt{c + dx}\right)}{d} \\ &= \frac{2\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd} - \frac{2 \text{Subst}\left(\int \sin(a + bx) dx, x, \sqrt{c + dx}\right)}{bd} \\ &= \frac{2 \cos(a + b\sqrt{c + dx})}{b^2 d} + \frac{2\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd} \end{aligned}$$

**Mathematica [A]** time = 0.0734567, size = 48, normalized size = 0.89

$$\frac{2(b\sqrt{c+dx}\sin(a+b\sqrt{c+dx})+\cos(a+b\sqrt{c+dx}))}{b^2d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*Sqrt[c + d\*x]],x]

[Out] (2\*(Cos[a + b\*Sqrt[c + d\*x]] + b\*Sqrt[c + d\*x]\*Sin[a + b\*Sqrt[c + d\*x]]))/(b^2\*d)

**Maple [A]** time = 0.033, size = 61, normalized size = 1.1

$$2\frac{\cos(a+b\sqrt{dx+c})+(a+b\sqrt{dx+c})\sin(a+b\sqrt{dx+c})-a\sin(a+b\sqrt{dx+c})}{b^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*(d\*x+c)^(1/2)),x)

[Out] 2/d/b^2\*(cos(a+b\*(d\*x+c)^(1/2))+(a+b\*(d\*x+c)^(1/2))\*sin(a+b\*(d\*x+c)^(1/2))-a\*sin(a+b\*(d\*x+c)^(1/2)))

**Maxima [A]** time = 1.15409, size = 81, normalized size = 1.5

$$\frac{2((\sqrt{dx+cb}+a)\sin(\sqrt{dx+cb}+a)-a\sin(\sqrt{dx+cb}+a)+\cos(\sqrt{dx+cb}+a))}{b^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="maxima")

[Out] 2\*((sqrt(d\*x + c)\*b + a)\*sin(sqrt(d\*x + c)\*b + a) - a\*sin(sqrt(d\*x + c)\*b + a) + cos(sqrt(d\*x + c)\*b + a))/(b^2\*d)

**Fricas [A]** time = 1.57493, size = 109, normalized size = 2.02

$$\frac{2(\sqrt{dx+cb}\sin(\sqrt{dx+cb}+a)+\cos(\sqrt{dx+cb}+a))}{b^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="fricas")

[Out] 2\*(sqrt(d\*x + c)\*b\*sin(sqrt(d\*x + c)\*b + a) + cos(sqrt(d\*x + c)\*b + a))/(b^2\*d)

**Sympy [A]** time = 0.535975, size = 66, normalized size = 1.22

$$\begin{cases} x \cos(a) & \text{for } b = 0 \wedge d = 0 \\ x \cos(a + b\sqrt{c}) & \text{for } d = 0 \\ x \cos(a) & \text{for } b = 0 \\ \frac{2\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{bd} + \frac{2 \cos(a+b\sqrt{c+dx})}{b^2d} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)\*\*(1/2)),x)

[Out] Piecewise((x\*cos(a), Eq(b, 0) & Eq(d, 0)), (x\*cos(a + b\*sqrt(c)), Eq(d, 0)), (x\*cos(a), Eq(b, 0)), (2\*sqrt(c + d\*x)\*sin(a + b\*sqrt(c + d\*x))/(b\*d) + 2\*cos(a + b\*sqrt(c + d\*x))/(b\*\*2\*d), True))

**Giac [B]** time = 1.18209, size = 227, normalized size = 4.2

$$\frac{2 \left( \left( \sqrt{dx+cb+a} \right) b - ab \right) \sin \left( - \left( \sqrt{dx+cb+a} \right) \operatorname{sgn} \left( \left( \sqrt{dx+cb+a} \right) b - ab \right) + a \operatorname{sgn} \left( \left( \sqrt{dx+cb+a} \right) b - ab \right) - a \right) - b \cos \left( - \left( \sqrt{dx+cb+a} \right) \operatorname{sgn} \left( \left( \sqrt{dx+cb+a} \right) b - ab \right) + a \operatorname{sgn} \left( \left( \sqrt{dx+cb+a} \right) b - ab \right) - a \right)}{b^3 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/2)),x, algorithm="giac")

[Out] -2\*(((sqrt(d\*x + c)\*b + a)\*b - a\*b)\*sin(-(sqrt(d\*x + c)\*b + a)\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + a\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - a) - b\*cos(-(sqrt(d\*x + c)\*b + a)\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) + a\*sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b) - a)/sgn((sqrt(d\*x + c)\*b + a)\*b - a\*b))/(b^3\*d)

### 3.93 $\int \frac{\cos(a+b\sqrt{c+dx})}{x} dx$

**Optimal.** Leaf size=126

$$\cos(a - b\sqrt{c}) \operatorname{CosIntegral}\left(b\left(\sqrt{c+dx} + \sqrt{c}\right)\right) + \cos(a + b\sqrt{c}) \operatorname{CosIntegral}\left(b\sqrt{c} - b\sqrt{c+dx}\right) - \sin(a - b\sqrt{c}) \operatorname{Si}\left(b\left(\sqrt{c+dx} + \sqrt{c}\right)\right) + \sin(a + b\sqrt{c}) \operatorname{Si}\left(b\sqrt{c} - b\sqrt{c+dx}\right)$$

```
[Out] Cos[a - b*Sqrt[c]]*CosIntegral[b*(Sqrt[c] + Sqrt[c + d*x])] + Cos[a + b*Sqrt[c]]*CosIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]] - Sin[a - b*Sqrt[c]]*SinIntegral[b*(Sqrt[c] + Sqrt[c + d*x])] + Sin[a + b*Sqrt[c]]*SinIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]]
```

**Rubi [A]** time = 0.238869, antiderivative size = 126, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 4, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {3432, 3303, 3299, 3302}

$$\cos(a - b\sqrt{c}) \operatorname{CosIntegral}\left(b\left(\sqrt{c+dx} + \sqrt{c}\right)\right) + \cos(a + b\sqrt{c}) \operatorname{CosIntegral}\left(b\sqrt{c} - b\sqrt{c+dx}\right) - \sin(a - b\sqrt{c}) \operatorname{Si}\left(b\left(\sqrt{c+dx} + \sqrt{c}\right)\right) + \sin(a + b\sqrt{c}) \operatorname{Si}\left(b\sqrt{c} - b\sqrt{c+dx}\right)$$

Antiderivative was successfully verified.

```
[In] Int[Cos[a + b*Sqrt[c + d*x]]/x,x]
```

```
[Out] Cos[a - b*Sqrt[c]]*CosIntegral[b*(Sqrt[c] + Sqrt[c + d*x])] + Cos[a + b*Sqrt[c]]*CosIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]] - Sin[a - b*Sqrt[c]]*SinIntegral[b*(Sqrt[c] + Sqrt[c + d*x])] + Sin[a + b*Sqrt[c]]*SinIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]]
```

#### Rule 3432

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])*(b_.)^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] :> Dist[1/(n*f), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - (e*h)/f + (h*x^(1/n))/f)^m, x], x], x, (e + f*x)^n, x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

#### Rule 3303

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

#### Rule 3299

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

#### Rule 3302

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

#### Rubi steps

$$\begin{aligned}
\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx &= \frac{2 \operatorname{Subst}\left(\int \left(-\frac{d \cos(a+bx)}{2(\sqrt{c-x})} + \frac{d \cos(a+bx)}{2(\sqrt{c+x})}\right) dx, x, \sqrt{c + dx}\right)}{d} \\
&= -\operatorname{Subst}\left(\int \frac{\cos(a + bx)}{\sqrt{c - x}} dx, x, \sqrt{c + dx}\right) + \operatorname{Subst}\left(\int \frac{\cos(a + bx)}{\sqrt{c + x}} dx, x, \sqrt{c + dx}\right) \\
&= \cos(a - b\sqrt{c}) \operatorname{Subst}\left(\int \frac{\cos(b\sqrt{c} + bx)}{\sqrt{c + x}} dx, x, \sqrt{c + dx}\right) - \cos(a + b\sqrt{c}) \operatorname{Subst}\left(\int \frac{\cos(b\sqrt{c} + bx)}{\sqrt{c + x}} dx, x, \sqrt{c + dx}\right) \\
&= \cos(a - b\sqrt{c}) \operatorname{Ci}\left(b\left(\sqrt{c} + \sqrt{c + dx}\right)\right) + \cos(a + b\sqrt{c}) \operatorname{Ci}\left(b\sqrt{c} - b\sqrt{c + dx}\right) - \sin(a - b\sqrt{c}) \operatorname{Si}\left(b\left(\sqrt{c} + \sqrt{c + dx}\right)\right) - \sin(a + b\sqrt{c}) \operatorname{Si}\left(b\sqrt{c} - b\sqrt{c + dx}\right)
\end{aligned}$$

**Mathematica [C]** time = 0.66874, size = 145, normalized size = 1.15

$$\frac{1}{2} e^{-i(a+b\sqrt{c})} \left( e^{2i(a+b\sqrt{c})} \operatorname{Ei}\left(ib\left(\sqrt{c+dx} - \sqrt{c}\right)\right) + e^{2ia} \operatorname{Ei}\left(ib\left(\sqrt{c} + \sqrt{c+dx}\right)\right) + \operatorname{Ei}\left(-ib\left(\sqrt{c+dx} - \sqrt{c}\right)\right) + e^{2ib\sqrt{c}} \operatorname{Ei}\left(-ib\left(\sqrt{c} + \sqrt{c+dx}\right)\right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*Sqrt[c + d\*x]]/x,x]

[Out] (ExpIntegralEi[(-I)\*b\*(-Sqrt[c] + Sqrt[c + d\*x])] + E^((2\*I)\*(a + b\*Sqrt[c]))\*ExpIntegralEi[I\*b\*(-Sqrt[c] + Sqrt[c + d\*x])] + E^((2\*I)\*b\*Sqrt[c])\*ExpIntegralEi[(-I)\*b\*(Sqrt[c] + Sqrt[c + d\*x])] + E^((2\*I)\*a)\*ExpIntegralEi[I\*b\*(Sqrt[c] + Sqrt[c + d\*x])])/(2\*E^(I\*(a + b\*Sqrt[c])))

**Maple [B]** time = 0.058, size = 271, normalized size = 2.2

$$2 \frac{1}{b^2} \left( \frac{1}{2} \frac{b(a + b\sqrt{c}) \left( \operatorname{Si}(b\sqrt{c} - b\sqrt{dx + c}) \sin(a + b\sqrt{c}) + \operatorname{Ci}(b\sqrt{dx + c} - b\sqrt{c}) \cos(a + b\sqrt{c}) \right)}{\sqrt{c}} - \frac{1}{2} \frac{b(a - b\sqrt{c}) \left( \operatorname{Si}(b\sqrt{c} + b\sqrt{dx + c}) \sin(a - b\sqrt{c}) + \operatorname{Ci}(b\sqrt{c} + b\sqrt{dx + c}) \cos(a - b\sqrt{c}) \right)}{\sqrt{c}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*(d\*x+c)^(1/2))/x,x)

[Out] 2/b^2\*(1/2\*b\*(a+b\*c^(1/2))/c^(1/2)\*(Si(b\*c^(1/2)-b\*(d\*x+c)^(1/2))\*sin(a+b\*c^(1/2))+Ci(b\*(d\*x+c)^(1/2)-b\*c^(1/2))\*cos(a+b\*c^(1/2)))-1/2\*b\*(a-b\*c^(1/2))/c^(1/2)\*(-Si(b\*(d\*x+c)^(1/2)+b\*c^(1/2))\*sin(a-b\*c^(1/2))+Ci(b\*(d\*x+c)^(1/2)+b\*c^(1/2))\*cos(a-b\*c^(1/2)))-a\*b^2\*(1/2/b/c^(1/2)\*(Si(b\*c^(1/2)-b\*(d\*x+c)^(1/2))\*sin(a+b\*c^(1/2))+Ci(b\*(d\*x+c)^(1/2)-b\*c^(1/2))\*cos(a+b\*c^(1/2)))-1/2/b/c^(1/2)\*(-Si(b\*(d\*x+c)^(1/2)+b\*c^(1/2))\*sin(a-b\*c^(1/2))+Ci(b\*(d\*x+c)^(1/2)+b\*c^(1/2))\*cos(a-b\*c^(1/2))))

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(\sqrt{dx + cb} + a)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/2))/x,x, algorithm="maxima")

[Out] integrate(cos(sqrt(d\*x + c)\*b + a)/x, x)

**Fricas [C]** time = 1.72537, size = 360, normalized size = 2.86

$$\frac{1}{2} \operatorname{Ei}\left(i\sqrt{dx+cb}-\sqrt{-b^2c}\right)e^{(ia+\sqrt{-b^2c})} + \frac{1}{2} \operatorname{Ei}\left(i\sqrt{dx+cb}+\sqrt{-b^2c}\right)e^{(ia-\sqrt{-b^2c})} + \frac{1}{2} \operatorname{Ei}\left(-i\sqrt{dx+cb}-\sqrt{-b^2c}\right)e^{(-ia+\sqrt{-b^2c})}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/2))/x,x, algorithm="fricas")

[Out] 1/2\*Ei(I\*sqrt(d\*x + c)\*b - sqrt(-b^2\*c))\*e^(I\*a + sqrt(-b^2\*c)) + 1/2\*Ei(I\*sqrt(d\*x + c)\*b + sqrt(-b^2\*c))\*e^(I\*a - sqrt(-b^2\*c)) + 1/2\*Ei(-I\*sqrt(d\*x + c)\*b - sqrt(-b^2\*c))\*e^(-I\*a + sqrt(-b^2\*c)) + 1/2\*Ei(-I\*sqrt(d\*x + c)\*b + sqrt(-b^2\*c))\*e^(-I\*a - sqrt(-b^2\*c))

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + b\sqrt{c + dx}\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)\*\*(1/2))/x,x)

[Out] Integral(cos(a + b\*sqrt(c + d\*x))/x, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(\sqrt{dx+cb}+a\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/2))/x,x, algorithm="giac")

[Out] integrate(cos(sqrt(d\*x + c)\*b + a)/x, x)



$$3.94 \quad \int \frac{\cos(a+b\sqrt{c+dx})}{x^2} dx$$

**Optimal.** Leaf size=184

$$\frac{bd \sin(a - b\sqrt{c}) \operatorname{CosIntegral}(b(\sqrt{c+dx} + \sqrt{c}))}{2\sqrt{c}} - \frac{bd \sin(a + b\sqrt{c}) \operatorname{CosIntegral}(b\sqrt{c} - b\sqrt{c+dx})}{2\sqrt{c}} + \frac{bd \cos(a - b\sqrt{c})}{2\sqrt{c}}$$

```
[Out] -(Cos[a + b*Sqrt[c + d*x]]/x) + (b*d*CosIntegral[b*(Sqrt[c] + Sqrt[c + d*x])]*Sin[a - b*Sqrt[c]])/(2*Sqrt[c]) - (b*d*CosIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]]*Sin[a + b*Sqrt[c]])/(2*Sqrt[c]) + (b*d*Cos[a - b*Sqrt[c]]*SinIntegral[b*(Sqrt[c] + Sqrt[c + d*x])])/(2*Sqrt[c]) + (b*d*Cos[a + b*Sqrt[c]]*SinIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]])/(2*Sqrt[c])
```

**Rubi [A]** time = 0.349114, antiderivative size = 184, normalized size of antiderivative = 1., number of steps used = 10, number of rules used = 6, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3432, 3342, 3333, 3303, 3299, 3302}

$$\frac{bd \sin(a - b\sqrt{c}) \operatorname{CosIntegral}(b(\sqrt{c+dx} + \sqrt{c}))}{2\sqrt{c}} - \frac{bd \sin(a + b\sqrt{c}) \operatorname{CosIntegral}(b\sqrt{c} - b\sqrt{c+dx})}{2\sqrt{c}} + \frac{bd \cos(a - b\sqrt{c})}{2\sqrt{c}}$$

Antiderivative was successfully verified.

```
[In] Int[Cos[a + b*Sqrt[c + d*x]]/x^2,x]
```

```
[Out] -(Cos[a + b*Sqrt[c + d*x]]/x) + (b*d*CosIntegral[b*(Sqrt[c] + Sqrt[c + d*x])]*Sin[a - b*Sqrt[c]])/(2*Sqrt[c]) - (b*d*CosIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]]*Sin[a + b*Sqrt[c]])/(2*Sqrt[c]) + (b*d*Cos[a - b*Sqrt[c]]*SinIntegral[b*(Sqrt[c] + Sqrt[c + d*x])])/(2*Sqrt[c]) + (b*d*Cos[a + b*Sqrt[c]]*SinIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]])/(2*Sqrt[c])
```

#### Rule 3432

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Dist[1/(n*f), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - (e*h)/f + (h*x^(1/n))/f)^m, x], x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

#### Rule 3342

```
Int[Cos[(c_.) + (d_.)*(x_)]*((e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_))^(n_.)^(p_.), x_Symbol] := Simp[(e^m*(a + b*x^n)^(p + 1)*Cos[c + d*x])/(b*n*(p + 1)), x] + Dist[(d*e^m)/(b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*Sin[c + d*x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && ILtQ[p, -1] && EqQ[m, n - 1] && (IntegerQ[n] || GtQ[e, 0])
```

#### Rule 3333

```
Int[((a_.) + (b_.)*(x_))^(n_.)^(p_.)*Sin[(c_.) + (d_.)*(x_)], x_Symbol] := Int[ExpandIntegrand[Sin[c + d*x], (a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IGtQ[n, 0] && (EqQ[n, 2] || EqQ[p, -1])
```

#### Rule 3303

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)
```

)/d], Int[Cos[(c\*f)/d + f\*x]/(c + d\*x), x], x] /; FreeQ[{c, d, e, f}, x] &&  
NeQ[d\*e - c\*f, 0]

### Rule 3299

Int[sin[(e\_.) + (f\_.)\*(x\_.)]/((c\_.) + (d\_.)\*(x\_.)), x\_Symbol] :> Simp[SinInte  
gral[e + f\*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d\*e - c\*f, 0]

### Rule 3302

Int[sin[(e\_.) + (f\_.)\*(x\_.)]/((c\_.) + (d\_.)\*(x\_.)), x\_Symbol] :> Simp[CosInte  
gral[e - Pi/2 + f\*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d\*(e - Pi/2) -  
c\*f, 0]

### Rubi steps

$$\begin{aligned} \int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx &= \frac{2 \operatorname{Subst} \left( \int \frac{x \cos(a+bx)}{\left(-\frac{c}{d} + \frac{x^2}{d}\right)^2} dx, x, \sqrt{c + dx} \right)}{d} \\ &= -\frac{\cos(a + b\sqrt{c + dx})}{x} - b \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{-\frac{c}{d} + \frac{x^2}{d}} dx, x, \sqrt{c + dx} \right) \\ &= -\frac{\cos(a + b\sqrt{c + dx})}{x} - b \operatorname{Subst} \left( \int \left( -\frac{d \sin(a + bx)}{2\sqrt{c}(\sqrt{c} - x)} - \frac{d \sin(a + bx)}{2\sqrt{c}(\sqrt{c} + x)} \right) dx, x, \sqrt{c + dx} \right) \\ &= -\frac{\cos(a + b\sqrt{c + dx})}{x} + \frac{(bd) \operatorname{Subst} \left( \int \frac{\sin(a+bx)}{\sqrt{c}-x} dx, x, \sqrt{c + dx} \right)}{2\sqrt{c}} + \frac{(bd) \operatorname{Subst} \left( \int \frac{\sin(a+bx)}{\sqrt{c}+x} dx, x, \sqrt{c + dx} \right)}{2\sqrt{c}} \\ &= -\frac{\cos(a + b\sqrt{c + dx})}{x} + \frac{(bd \cos(a - b\sqrt{c})) \operatorname{Subst} \left( \int \frac{\sin(b\sqrt{c}+bx)}{\sqrt{c}+x} dx, x, \sqrt{c + dx} \right)}{2\sqrt{c}} - \frac{(bd \cos(a + b\sqrt{c})) \operatorname{Subst} \left( \int \frac{\sin(b\sqrt{c}+bx)}{\sqrt{c}-x} dx, x, \sqrt{c + dx} \right)}{2\sqrt{c}} \\ &= -\frac{\cos(a + b\sqrt{c + dx})}{x} + \frac{bd \operatorname{Ci}(b(\sqrt{c} + \sqrt{c + dx})) \sin(a - b\sqrt{c})}{2\sqrt{c}} - \frac{bd \operatorname{Ci}(b\sqrt{c} - b\sqrt{c + dx}) \sin(a + b\sqrt{c})}{2\sqrt{c}} \end{aligned}$$

**Mathematica [C]** time = 1.22471, size = 240, normalized size = 1.3

$$\frac{i \left( e^{-ia} \left( -bdxe^{-ib\sqrt{c}} \operatorname{Ei}(-ib(\sqrt{c + dx} - \sqrt{c})) + bdx e^{ib\sqrt{c}} \operatorname{Ei}(-ib(\sqrt{c} + \sqrt{c + dx})) + 2i\sqrt{c} e^{-ib\sqrt{c + dx}} \right) + e^{i(a - b\sqrt{c})} \left( bdx e^{2ib\sqrt{c}} \operatorname{Ei}(b(\sqrt{c} + \sqrt{c + dx})) - bdx e^{-2ib\sqrt{c}} \operatorname{Ei}(b\sqrt{c} - b\sqrt{c + dx}) \right) \right)}{4\sqrt{cx}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*Sqrt[c + d\*x]]/x^2,x]

[Out] ((I/4)\*(((2\*I)\*Sqrt[c])/E^(I\*b\*Sqrt[c + d\*x]) - (b\*d\*x\*ExpIntegralEi[(-I)\*  
b\*(-Sqrt[c] + Sqrt[c + d\*x])])/E^(I\*b\*Sqrt[c]) + b\*d\*E^(I\*b\*Sqrt[c])\*x\*ExpI  
ntegralEi[(-I)\*b\*(Sqrt[c] + Sqrt[c + d\*x])])/E^(I\*a) + E^(I\*(a - b\*Sqrt[c])  
)\*((2\*I)\*Sqrt[c]\*E^(I\*b\*(Sqrt[c] + Sqrt[c + d\*x])) + b\*d\*E^((2\*I)\*b\*Sqrt[c]  
) \*x\*ExpIntegralEi[I\*b\*(-Sqrt[c] + Sqrt[c + d\*x])]) - b\*d\*x\*ExpIntegralEi[I\*b  
(Sqrt[c] + Sqrt[c + d\*x])]))/(Sqrt[c]\*x)

**Maple [B]** time = 0.181, size = 714, normalized size = 3.9

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In]  $\int \cos(a+b*(d*x+c)^{(1/2)})/x^2, x$

[Out]  $2*d/b^2*(\cos(a+b*(d*x+c)^{(1/2)})*(-1/2*a*b^2/c*(a+b*(d*x+c)^{(1/2)})+1/2*b^2*(-b^2*c+a^2)/c)/(-b^2*c+(a+b*(d*x+c)^{(1/2)})^2-2*(a+b*(d*x+c)^{(1/2)})*a+a^2)-1/4*a*b/c^{(3/2)}*(\text{Si}(b*c^{(1/2)}-b*(d*x+c)^{(1/2)})*\sin(a+b*c^{(1/2)})+\text{Ci}(b*(d*x+c)^{(1/2)}-b*c^{(1/2)})*\cos(a+b*c^{(1/2)}))+1/4*a*b/c^{(3/2)}*(-\text{Si}(b*(d*x+c)^{(1/2)}+b*c^{(1/2)})*\sin(a-b*c^{(1/2)})+\text{Ci}(b*(d*x+c)^{(1/2)}+b*c^{(1/2)})*\cos(a-b*c^{(1/2)}))-1/4*b*(b^2*c+(a+b*c^{(1/2)})*a-a^2)/c^{(3/2)}*(-\text{Si}(b*c^{(1/2)}-b*(d*x+c)^{(1/2)})*\cos(a+b*c^{(1/2)})+\text{Ci}(b*(d*x+c)^{(1/2)}-b*c^{(1/2)})*\sin(a+b*c^{(1/2)}))+1/4*b*(b^2*c+(a-b*c^{(1/2)})*a-a^2)/c^{(3/2)}*(\text{Si}(b*(d*x+c)^{(1/2)}+b*c^{(1/2)})*\cos(a-b*c^{(1/2)})+\text{Ci}(b*(d*x+c)^{(1/2)}+b*c^{(1/2)})*\sin(a-b*c^{(1/2)}))-a*b^4*(\cos(a+b*(d*x+c)^{(1/2)})*(-1/2/b^2/c*(a+b*(d*x+c)^{(1/2)})+1/2*a/b^2/c)/(-b^2*c+(a+b*(d*x+c)^{(1/2)})^2-2*(a+b*(d*x+c)^{(1/2)})*a+a^2)-1/4/b^3/c^{(3/2)}*(\text{Si}(b*c^{(1/2)}-b*(d*x+c)^{(1/2)})*\sin(a+b*c^{(1/2)})+\text{Ci}(b*(d*x+c)^{(1/2)}-b*c^{(1/2)})*\cos(a+b*c^{(1/2)}))+1/4/b^3/c^{(3/2)}*(-\text{Si}(b*(d*x+c)^{(1/2)}+b*c^{(1/2)})*\sin(a-b*c^{(1/2)})+\text{Ci}(b*(d*x+c)^{(1/2)}+b*c^{(1/2)})*\cos(a-b*c^{(1/2)}))-1/4/b^2/c*(-\text{Si}(b*c^{(1/2)}-b*(d*x+c)^{(1/2)})*\cos(a+b*c^{(1/2)})+\text{Ci}(b*(d*x+c)^{(1/2)}-b*c^{(1/2)})*\sin(a+b*c^{(1/2)}))-1/4/b^2/c*(\text{Si}(b*(d*x+c)^{(1/2)}+b*c^{(1/2)})*\cos(a-b*c^{(1/2)})+\text{Ci}(b*(d*x+c)^{(1/2)}+b*c^{(1/2)})*\sin(a-b*c^{(1/2)}))$

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(\sqrt{dx+cb+a})}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In]  $\text{integrate}(\cos(a+b*(d*x+c)^{(1/2)})/x^2, x, \text{algorithm}="maxima")$

[Out]  $\text{integrate}(\cos(\text{sqrt}(d*x + c)*b + a)/x^2, x)$

**Fricas [C]** time = 1.82255, size = 489, normalized size = 2.66

$$\frac{\sqrt{-b^2c}dx\text{Ei}\left(i\sqrt{dx+cb}-\sqrt{-b^2c}\right)e^{(ia+\sqrt{-b^2c})}-\sqrt{-b^2c}dx\text{Ei}\left(i\sqrt{dx+cb}+\sqrt{-b^2c}\right)e^{(ia-\sqrt{-b^2c})}+\sqrt{-b^2c}dx\text{Ei}\left(-i\sqrt{dx+cb}-\sqrt{-b^2c}\right)e^{(-ia-\sqrt{-b^2c})}-\sqrt{-b^2c}dx\text{Ei}\left(-i\sqrt{dx+cb}+\sqrt{-b^2c}\right)e^{(-ia+\sqrt{-b^2c})}}{4cx}$$

Verification of antiderivative is not currently implemented for this CAS.

[In]  $\text{integrate}(\cos(a+b*(d*x+c)^{(1/2)})/x^2, x, \text{algorithm}="fricas")$

[Out]  $1/4*(\text{sqrt}(-b^2*c)*d*x*\text{Ei}(I*\text{sqrt}(d*x + c)*b - \text{sqrt}(-b^2*c))*e^{(I*a + \text{sqrt}(-b^2*c))} - \text{sqrt}(-b^2*c)*d*x*\text{Ei}(I*\text{sqrt}(d*x + c)*b + \text{sqrt}(-b^2*c))*e^{(I*a - \text{sqrt}(-b^2*c))} + \text{sqrt}(-b^2*c)*d*x*\text{Ei}(-I*\text{sqrt}(d*x + c)*b - \text{sqrt}(-b^2*c))*e^{(-I*a + \text{sqrt}(-b^2*c))} - \text{sqrt}(-b^2*c)*d*x*\text{Ei}(-I*\text{sqrt}(d*x + c)*b + \text{sqrt}(-b^2*c))*e^{(-I*a - \text{sqrt}(-b^2*c))} - 4*c*\cos(\text{sqrt}(d*x + c)*b + a))/(c*x)$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)\*\*(1/2))/x\*\*2,x)

[Out] Integral(cos(a + b\*sqrt(c + d\*x))/x\*\*2, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos(\sqrt{dx + cb + a})}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/2))/x^2,x, algorithm="giac")

[Out] integrate(cos(sqrt(d\*x + c)\*b + a)/x^2, x)

### 3.95 $\int x^2 \cos\left(a + b\sqrt[3]{c + dx}\right) dx$

**Optimal.** Leaf size=537

$$\frac{6c^2 \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^3 d^3} + \frac{6c^2 \sqrt[3]{c + dx} \cos\left(a + b\sqrt[3]{c + dx}\right)}{b^2 d^3} - \frac{168(c + dx)^2 \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^3 d^3} + \frac{5040(c + dx)^{4/3} \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^5 d^3}$$

```
[Out] (-720*c*Cos[a + b*(c + d*x)^(1/3)]/(b^6*d^3) - (120960*(c + d*x)^(1/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^8*d^3) + (6*c^2*(c + d*x)^(1/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^2*d^3) + (360*c*(c + d*x)^(2/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^4*d^3) + (20160*(c + d*x)*Cos[a + b*(c + d*x)^(1/3)]/(b^6*d^3) - (30*c*(c + d*x)^(4/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^2*d^3) - (1008*(c + d*x)^(5/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^4*d^3) + (24*(c + d*x)^(7/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^2*d^3) + (120960*Sin[a + b*(c + d*x)^(1/3)]/(b^9*d^3) - (6*c^2*Sin[a + b*(c + d*x)^(1/3)]/(b^3*d^3) - (720*c*(c + d*x)^(1/3)*Sin[a + b*(c + d*x)^(1/3)]/(b^5*d^3) - (60480*(c + d*x)^(2/3)*Sin[a + b*(c + d*x)^(1/3)]/(b^7*d^3) + (3*c^2*(c + d*x)^(2/3)*Sin[a + b*(c + d*x)^(1/3)]/(b*d^3) + (120*c*(c + d*x)*Sin[a + b*(c + d*x)^(1/3)]/(b^3*d^3) + (5040*(c + d*x)^(4/3)*Sin[a + b*(c + d*x)^(1/3)]/(b^5*d^3) - (6*c*(c + d*x)^(5/3)*Sin[a + b*(c + d*x)^(1/3)]/(b*d^3) - (168*(c + d*x)^2*Sin[a + b*(c + d*x)^(1/3)]/(b^3*d^3) + (3*(c + d*x)^(8/3)*Sin[a + b*(c + d*x)^(1/3)]/(b*d^3)
```

**Rubi [A]** time = 0.51131, antiderivative size = 537, normalized size of antiderivative = 1., number of steps used = 20, number of rules used = 4, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {3432, 3296, 2637, 2638}

$$\frac{6c^2 \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^3 d^3} + \frac{6c^2 \sqrt[3]{c + dx} \cos\left(a + b\sqrt[3]{c + dx}\right)}{b^2 d^3} - \frac{168(c + dx)^2 \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^3 d^3} + \frac{5040(c + dx)^{4/3} \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^5 d^3}$$

Antiderivative was successfully verified.

```
[In] Int[x^2*Cos[a + b*(c + d*x)^(1/3)], x]
```

```
[Out] (-720*c*Cos[a + b*(c + d*x)^(1/3)]/(b^6*d^3) - (120960*(c + d*x)^(1/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^8*d^3) + (6*c^2*(c + d*x)^(1/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^2*d^3) + (360*c*(c + d*x)^(2/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^4*d^3) + (20160*(c + d*x)*Cos[a + b*(c + d*x)^(1/3)]/(b^6*d^3) - (30*c*(c + d*x)^(4/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^2*d^3) - (1008*(c + d*x)^(5/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^4*d^3) + (24*(c + d*x)^(7/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^2*d^3) + (120960*Sin[a + b*(c + d*x)^(1/3)]/(b^9*d^3) - (6*c^2*Sin[a + b*(c + d*x)^(1/3)]/(b^3*d^3) - (720*c*(c + d*x)^(1/3)*Sin[a + b*(c + d*x)^(1/3)]/(b^5*d^3) - (60480*(c + d*x)^(2/3)*Sin[a + b*(c + d*x)^(1/3)]/(b^7*d^3) + (3*c^2*(c + d*x)^(2/3)*Sin[a + b*(c + d*x)^(1/3)]/(b*d^3) + (120*c*(c + d*x)*Sin[a + b*(c + d*x)^(1/3)]/(b^3*d^3) + (5040*(c + d*x)^(4/3)*Sin[a + b*(c + d*x)^(1/3)]/(b^5*d^3) - (6*c*(c + d*x)^(5/3)*Sin[a + b*(c + d*x)^(1/3)]/(b*d^3) - (168*(c + d*x)^2*Sin[a + b*(c + d*x)^(1/3)]/(b^3*d^3) + (3*(c + d*x)^(8/3)*Sin[a + b*(c + d*x)^(1/3)]/(b*d^3)
```

**Rule 3432**

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_.)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] :> Dist[1/(n*f), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - (e*h)/f + (h*x^(1/n))/f)^m, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

Rule 3296

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := -Simp[
((c + d*x)^m*cos[e + f*x])/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[
e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 2637

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

Rule 2638

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int x^2 \cos(a + b\sqrt[3]{c + dx}) dx &= \frac{3 \operatorname{Subst}\left(\int \left(\frac{c^2 x^2 \cos(ax + bx)}{d^2} - \frac{2cx^5 \cos(ax + bx)}{d^2} + \frac{x^8 \cos(ax + bx)}{d^2}\right) dx, x, \sqrt[3]{c + dx}\right)}{d} \\
&= \frac{3 \operatorname{Subst}\left(\int x^8 \cos(ax + bx) dx, x, \sqrt[3]{c + dx}\right)}{d^3} - \frac{(6c) \operatorname{Subst}\left(\int x^5 \cos(ax + bx) dx, x, \sqrt[3]{c + dx}\right)}{d^3} \\
&= \frac{3c^2(c + dx)^{2/3} \sin(a + b\sqrt[3]{c + dx})}{bd^3} - \frac{6c(c + dx)^{5/3} \sin(a + b\sqrt[3]{c + dx})}{bd^3} + \frac{3(c + dx)^{8/3} \sin(a + b\sqrt[3]{c + dx})}{bd^3} \\
&= \frac{6c^2 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} - \frac{30c(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} + \frac{24(c + dx)^{7/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} \\
&= \frac{6c^2 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} - \frac{30c(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} + \frac{24(c + dx)^{7/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} \\
&= \frac{6c^2 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} + \frac{360c(c + dx)^{2/3} \cos(a + b\sqrt[3]{c + dx})}{b^4 d^3} - \frac{30c(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} \\
&= \frac{6c^2 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} + \frac{360c(c + dx)^{2/3} \cos(a + b\sqrt[3]{c + dx})}{b^4 d^3} - \frac{30c(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} \\
&= -\frac{720c \cos(a + b\sqrt[3]{c + dx})}{b^6 d^3} + \frac{6c^2 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} + \frac{360c(c + dx)^{2/3} \cos(a + b\sqrt[3]{c + dx})}{b^4 d^3} \\
&= -\frac{720c \cos(a + b\sqrt[3]{c + dx})}{b^6 d^3} + \frac{6c^2 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} + \frac{360c(c + dx)^{2/3} \cos(a + b\sqrt[3]{c + dx})}{b^4 d^3} \\
&= -\frac{720c \cos(a + b\sqrt[3]{c + dx})}{b^6 d^3} - \frac{120960 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^8 d^3} + \frac{6c^2 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} \\
&= -\frac{720c \cos(a + b\sqrt[3]{c + dx})}{b^6 d^3} - \frac{120960 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^8 d^3} + \frac{6c^2 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3}
\end{aligned}$$

**Mathematica [C]** time = 1.10729, size = 382, normalized size = 0.71

$$3e^{-i(a+b\sqrt[3]{c+dx})} \left( 2ib^6 (9c^2 + 36cdx + 28d^2x^2) \left( -1 + e^{2i(a+b\sqrt[3]{c+dx})} \right) - ib^8 d^2 x^2 (c + dx)^{2/3} \left( -1 + e^{2i(a+b\sqrt[3]{c+dx})} \right) + 2b^7 dx \sqrt[3]{c + dx} \right)$$

Antiderivative was successfully verified.

```
[In] Integrate[x^2*Cos[a + b*(c + d*x)^(1/3)],x]
```

```
[Out] (3*((-40320*I)*(-1 + E^((2*I)*(a + b*(c + d*x)^(1/3)))) - 40320*b*(1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*(c + d*x)^(1/3) + (20160*I)*b^2*(-1 + E^((2*
```

$$I*(a + b*(c + d*x)^{(1/3)}))*(c + d*x)^{(2/3)} - I*b^8*d^2*(-1 + E^{((2*I)*(a + b*(c + d*x)^{(1/3)}))})*x^2*(c + d*x)^{(2/3)} + 2*b^7*d*(1 + E^{((2*I)*(a + b*(c + d*x)^{(1/3)}))})*x*(c + d*x)^{(1/3)}*(3*c + 4*d*x) - (240*I)*b^4*(-1 + E^{((2*I)*(a + b*(c + d*x)^{(1/3)}))})*(c + d*x)^{(1/3)}*(6*c + 7*d*x) - 24*b^5*(1 + E^{((2*I)*(a + b*(c + d*x)^{(1/3)}))})*(c + d*x)^{(2/3)}*(9*c + 14*d*x) + 240*b^3*(1 + E^{((2*I)*(a + b*(c + d*x)^{(1/3)}))})*(27*c + 28*d*x) + (2*I)*b^6*(-1 + E^{((2*I)*(a + b*(c + d*x)^{(1/3)}))})*(9*c^2 + 36*c*d*x + 28*d^2*x^2))/(2*b^9*d^3*E^{(I*(a + b*(c + d*x)^{(1/3)}))})$$

**Maple [B]** time = 0.102, size = 1809, normalized size = 3.4

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In]  $\text{int}(x^2 \cos(a+b*(d*x+c)^{(1/3)}), x)$

[Out] 
$$\begin{aligned} & 3/d^3/b^3*(c^2*((a+b*(d*x+c)^{(1/3)})^2*\sin(a+b*(d*x+c)^{(1/3)})-2*\sin(a+b*(d*x+c)^{(1/3)})+2*(a+b*(d*x+c)^{(1/3)})*\cos(a+b*(d*x+c)^{(1/3)}))-2*a*c^2*(\cos(a+b*(d*x+c)^{(1/3)})+(a+b*(d*x+c)^{(1/3)})*\sin(a+b*(d*x+c)^{(1/3)}))+c^2*a^2*\sin(a+b*(d*x+c)^{(1/3)})-2/b^3*c*((a+b*(d*x+c)^{(1/3)})^5*\sin(a+b*(d*x+c)^{(1/3)})+5*(a+b*(d*x+c)^{(1/3)})^4*\cos(a+b*(d*x+c)^{(1/3)})-20*(a+b*(d*x+c)^{(1/3)})^3*\sin(a+b*(d*x+c)^{(1/3)})-60*(a+b*(d*x+c)^{(1/3)})^2*\cos(a+b*(d*x+c)^{(1/3)})+120*\cos(a+b*(d*x+c)^{(1/3)})+120*(a+b*(d*x+c)^{(1/3)})*\sin(a+b*(d*x+c)^{(1/3)}))+10/b^3*a*c*((a+b*(d*x+c)^{(1/3)})^4*\sin(a+b*(d*x+c)^{(1/3)})+4*(a+b*(d*x+c)^{(1/3)})^3*\cos(a+b*(d*x+c)^{(1/3)})-12*(a+b*(d*x+c)^{(1/3)})^2*\sin(a+b*(d*x+c)^{(1/3)})+24*\sin(a+b*(d*x+c)^{(1/3)})-24*(a+b*(d*x+c)^{(1/3)})*\cos(a+b*(d*x+c)^{(1/3)}))-20/b^3*a^2*c*((a+b*(d*x+c)^{(1/3)})^3*\sin(a+b*(d*x+c)^{(1/3)})+3*(a+b*(d*x+c)^{(1/3)})^2*\cos(a+b*(d*x+c)^{(1/3)})-6*\cos(a+b*(d*x+c)^{(1/3)})-6*(a+b*(d*x+c)^{(1/3)})*\sin(a+b*(d*x+c)^{(1/3)}))+20/b^3*a^3*c*((a+b*(d*x+c)^{(1/3)})^2*\sin(a+b*(d*x+c)^{(1/3)})-2*\sin(a+b*(d*x+c)^{(1/3)})+2*(a+b*(d*x+c)^{(1/3)})*\cos(a+b*(d*x+c)^{(1/3)}))-10/b^3*a^4*c*(\cos(a+b*(d*x+c)^{(1/3)})+(a+b*(d*x+c)^{(1/3)})*\sin(a+b*(d*x+c)^{(1/3)}))+2/b^3*a^5*c*\sin(a+b*(d*x+c)^{(1/3)})+1/b^6*((a+b*(d*x+c)^{(1/3)})^8*\sin(a+b*(d*x+c)^{(1/3)})+8*(a+b*(d*x+c)^{(1/3)})^7*\cos(a+b*(d*x+c)^{(1/3)})-56*(a+b*(d*x+c)^{(1/3)})^6*\sin(a+b*(d*x+c)^{(1/3)})-336*(a+b*(d*x+c)^{(1/3)})^5*\cos(a+b*(d*x+c)^{(1/3)})+1680*(a+b*(d*x+c)^{(1/3)})^4*\sin(a+b*(d*x+c)^{(1/3)})+6720*(a+b*(d*x+c)^{(1/3)})^3*\cos(a+b*(d*x+c)^{(1/3)})-20160*(a+b*(d*x+c)^{(1/3)})^2*\sin(a+b*(d*x+c)^{(1/3)})+40320*\sin(a+b*(d*x+c)^{(1/3)})-40320*(a+b*(d*x+c)^{(1/3)})*\cos(a+b*(d*x+c)^{(1/3)}))-8/b^6*a*((a+b*(d*x+c)^{(1/3)})^7*\sin(a+b*(d*x+c)^{(1/3)})+7*(a+b*(d*x+c)^{(1/3)})^6*\cos(a+b*(d*x+c)^{(1/3)})-42*(a+b*(d*x+c)^{(1/3)})^5*\sin(a+b*(d*x+c)^{(1/3)})-210*(a+b*(d*x+c)^{(1/3)})^4*\cos(a+b*(d*x+c)^{(1/3)})+840*(a+b*(d*x+c)^{(1/3)})^3*\sin(a+b*(d*x+c)^{(1/3)})+2520*(a+b*(d*x+c)^{(1/3)})^2*\cos(a+b*(d*x+c)^{(1/3)})-5040*\cos(a+b*(d*x+c)^{(1/3)})-5040*(a+b*(d*x+c)^{(1/3)})*\sin(a+b*(d*x+c)^{(1/3)}))+28/b^6*a^2*((a+b*(d*x+c)^{(1/3)})^6*\sin(a+b*(d*x+c)^{(1/3)})+6*(a+b*(d*x+c)^{(1/3)})^5*\cos(a+b*(d*x+c)^{(1/3)})-30*(a+b*(d*x+c)^{(1/3)})^4*\sin(a+b*(d*x+c)^{(1/3)})-120*(a+b*(d*x+c)^{(1/3)})^3*\cos(a+b*(d*x+c)^{(1/3)})+360*(a+b*(d*x+c)^{(1/3)})^2*\sin(a+b*(d*x+c)^{(1/3)})-720*\sin(a+b*(d*x+c)^{(1/3)})+720*(a+b*(d*x+c)^{(1/3)})*\cos(a+b*(d*x+c)^{(1/3)}))-56/b^6*a^3*((a+b*(d*x+c)^{(1/3)})^5*\sin(a+b*(d*x+c)^{(1/3)})+5*(a+b*(d*x+c)^{(1/3)})^4*\cos(a+b*(d*x+c)^{(1/3)})-20*(a+b*(d*x+c)^{(1/3)})^3*\sin(a+b*(d*x+c)^{(1/3)})-60*(a+b*(d*x+c)^{(1/3)})^2*\cos(a+b*(d*x+c)^{(1/3)})+120*\cos(a+b*(d*x+c)^{(1/3)})+120*(a+b*(d*x+c)^{(1/3)})*\sin(a+b*(d*x+c)^{(1/3)}))+70/b^6*a^4*((a+b*(d*x+c)^{(1/3)})^4*\sin(a+b*(d*x+c)^{(1/3)})+4*(a+b*(d*x+c)^{(1/3)})^3*\cos(a+b*(d*x+c)^{(1/3)})-12*(a+b*(d*x+c)^{(1/3)})^2*\sin(a+b*(d*x+c)^{(1/3)})+24*\sin(a+b*(d*x+c)^{(1/3)})-24*(a+b*(d*x+c)^{(1/3)})*\cos(a+b*(d*x+c)^{(1/3)}))-56/b^6*a^5*((a+b*(d*x+c)^{(1/3)})^3*\sin(a+b*(d*x+c)^{(1/3)})+3*(a+b*(d*x+c)^{(1/3)})^2*\cos(a+b*(d*x+c)^{(1/3)})-6*\cos(a+b*(d*x+c)^{(1/3)})-6*(a+b*(d*x+c)^{(1/3)})*\sin(a+b*(d*x+c)^{(1/3)}))+28/b^6*a^6*((a+b*(d*x+c)^{(1/3)})^2*\sin(a+b$$

```
*(d*x+c)^(1/3))-2*sin(a+b*(d*x+c)^(1/3))+2*(a+b*(d*x+c)^(1/3))*cos(a+b*(d*x+c)^(1/3))-8/b^6*a^7*(cos(a+b*(d*x+c)^(1/3))+(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))+1/b^6*a^8*sin(a+b*(d*x+c)^(1/3))
```

**Maxima [B]** time = 1.47874, size = 1821, normalized size = 3.39

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^2*cos(a+b*(d*x+c)^(1/3)),x, algorithm="maxima")
```

```
[Out] 3*(a^2*c^2*sin((d*x + c)^(1/3)*b + a) - 2*(((d*x + c)^(1/3)*b + a)*sin((d*x + c)^(1/3)*b + a) + cos((d*x + c)^(1/3)*b + a))*a*c^2 + 2*a^5*c*sin((d*x + c)^(1/3)*b + a)/b^3 - 10*(((d*x + c)^(1/3)*b + a)*sin((d*x + c)^(1/3)*b + a) + cos((d*x + c)^(1/3)*b + a))*a^4*c/b^3 + (2*(((d*x + c)^(1/3)*b + a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2 - 2)*sin((d*x + c)^(1/3)*b + a))*c^2 + a^8*sin((d*x + c)^(1/3)*b + a)/b^6 - 8*(((d*x + c)^(1/3)*b + a)*sin((d*x + c)^(1/3)*b + a) + cos((d*x + c)^(1/3)*b + a))*a^7/b^6 + 20*(2*(((d*x + c)^(1/3)*b + a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2 - 2)*sin((d*x + c)^(1/3)*b + a))*a^3*c/b^3 + 28*(2*(((d*x + c)^(1/3)*b + a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2 - 2)*sin((d*x + c)^(1/3)*b + a))*a^6/b^6 - 20*(3*(((d*x + c)^(1/3)*b + a)^2 - 2)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*sin((d*x + c)^(1/3)*b + a))*a^2*c/b^3 - 56*(3*(((d*x + c)^(1/3)*b + a)^2 - 2)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*sin((d*x + c)^(1/3)*b + a))*a^5/b^6 + 10*(4*(((d*x + c)^(1/3)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^4 - 12*((d*x + c)^(1/3)*b + a)^2 + 24)*sin((d*x + c)^(1/3)*b + a))*a*c/b^3 + 70*(4*(((d*x + c)^(1/3)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^4 - 12*((d*x + c)^(1/3)*b + a)^2 + 24)*sin((d*x + c)^(1/3)*b + a))*a^4/b^6 - 2*(5*((d*x + c)^(1/3)*b + a)^4 - 12*((d*x + c)^(1/3)*b + a)^2 + 24)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^5 - 20*((d*x + c)^(1/3)*b + a)^3 + 120*(d*x + c)^(1/3)*b + 120*a)*sin((d*x + c)^(1/3)*b + a))*c/b^3 - 56*(5*((d*x + c)^(1/3)*b + a)^4 - 12*((d*x + c)^(1/3)*b + a)^2 + 24)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^5 - 20*((d*x + c)^(1/3)*b + a)^3 + 120*(d*x + c)^(1/3)*b + 120*a)*sin((d*x + c)^(1/3)*b + a))*a^3/b^6 + 28*(6*(((d*x + c)^(1/3)*b + a)^5 - 20*((d*x + c)^(1/3)*b + a)^3 + 120*(d*x + c)^(1/3)*b + 120*a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^6 - 30*((d*x + c)^(1/3)*b + a)^4 + 360*((d*x + c)^(1/3)*b + a)^2 - 720)*sin((d*x + c)^(1/3)*b + a))*a^2/b^6 - 8*(7*(((d*x + c)^(1/3)*b + a)^6 - 30*((d*x + c)^(1/3)*b + a)^4 + 360*((d*x + c)^(1/3)*b + a)^2 - 720)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^7 - 42*((d*x + c)^(1/3)*b + a)^5 + 840*((d*x + c)^(1/3)*b + a)^3 - 5040*(d*x + c)^(1/3)*b - 5040*a)*sin((d*x + c)^(1/3)*b + a))*a/b^6 + (8*(((d*x + c)^(1/3)*b + a)^7 - 42*((d*x + c)^(1/3)*b + a)^5 + 840*((d*x + c)^(1/3)*b + a)^3 - 5040*(d*x + c)^(1/3)*b - 5040*a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^8 - 56*((d*x + c)^(1/3)*b + a)^6 + 1680*((d*x + c)^(1/3)*b + a)^4 - 20160*((d*x + c)^(1/3)*b + a)^2 + 40320)*sin((d*x + c)^(1/3)*b + a))/b^6)/(b^3*d^3)
```

**Fricas [A]** time = 1.70366, size = 464, normalized size = 0.86

$$3 \left( 2 \left( 3360 b^3 dx + 3240 b^3 c - 12 (14 b^5 dx + 9 b^5 c) (dx + c)^{\frac{2}{3}} + (4 b^7 d^2 x^2 + 3 b^7 c dx - 20160 b) (dx + c)^{\frac{1}{3}} \right) \cos \left( (dx + c)^{\frac{1}{3}} b + a \right) \right)$$



Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(a+b\*(d\*x+c)^(1/3)),x, algorithm="fricas")

[Out] 3\*(2\*(3360\*b^3\*d\*x + 3240\*b^3\*c - 12\*(14\*b^5\*d\*x + 9\*b^5\*c)\*(d\*x + c)^(2/3) + (4\*b^7\*d^2\*x^2 + 3\*b^7\*c\*d\*x - 20160\*b)\*(d\*x + c)^(1/3))\*cos((d\*x + c)^(1/3)\*b + a) - (56\*b^6\*d^2\*x^2 + 72\*b^6\*c\*d\*x + 18\*b^6\*c^2 - (b^8\*d^2\*x^2 - 20160\*b^2)\*(d\*x + c)^(2/3) - 240\*(7\*b^4\*d\*x + 6\*b^4\*c)\*(d\*x + c)^(1/3) - 40320)\*sin((d\*x + c)^(1/3)\*b + a))/(b^9\*d^3)

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x^2 \cos\left(a + b\sqrt[3]{c + dx}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*\*2\*cos(a+b\*(d\*x+c)\*\*(1/3)),x)

[Out] Integral(x\*\*2\*cos(a + b\*(c + d\*x)\*\*(1/3)), x)

**Giac [B]** time = 1.61316, size = 1490, normalized size = 2.77

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2\*cos(a+b\*(d\*x+c)^(1/3)),x, algorithm="giac")

[Out] 3\*(2\*((d\*x + c)^(1/3)\*b + a)\*b^6\*c^2 - a\*b^6\*c^2 - 5\*((d\*x + c)^(1/3)\*b + a)^4\*b^3\*c + 20\*((d\*x + c)^(1/3)\*b + a)^3\*a\*b^3\*c - 30\*((d\*x + c)^(1/3)\*b + a)^2\*a^2\*b^3\*c + 20\*((d\*x + c)^(1/3)\*b + a)\*a^3\*b^3\*c - 5\*a^4\*b^3\*c + 4\*((d\*x + c)^(1/3)\*b + a)^7 - 28\*((d\*x + c)^(1/3)\*b + a)^6\*a + 84\*((d\*x + c)^(1/3)\*b + a)^5\*a^2 - 140\*((d\*x + c)^(1/3)\*b + a)^4\*a^3 + 140\*((d\*x + c)^(1/3)\*b + a)^3\*a^4 - 84\*((d\*x + c)^(1/3)\*b + a)^2\*a^5 + 28\*((d\*x + c)^(1/3)\*b + a)\*a^6 - 4\*a^7 + 60\*((d\*x + c)^(1/3)\*b + a)^2\*b^3\*c - 120\*((d\*x + c)^(1/3)\*b + a)\*a\*b^3\*c + 60\*a^2\*b^3\*c - 168\*((d\*x + c)^(1/3)\*b + a)^5 + 840\*((d\*x + c)^(1/3)\*b + a)^4\*a - 1680\*((d\*x + c)^(1/3)\*b + a)^3\*a^2 + 1680\*((d\*x + c)^(1/3)\*b + a)^2\*a^3 - 840\*((d\*x + c)^(1/3)\*b + a)\*a^4 + 168\*a^5 - 120\*b^3\*c + 3360\*((d\*x + c)^(1/3)\*b + a)^3 - 10080\*((d\*x + c)^(1/3)\*b + a)^2\*a + 10080\*((d\*x + c)^(1/3)\*b + a)\*a^2 - 3360\*a^3 - 20160\*(d\*x + c)^(1/3)\*b\*cos((d\*x + c)^(1/3)\*b + a)/(b^8\*d^2) + (((d\*x + c)^(1/3)\*b + a)^2\*b^6\*c^2 - 2\*((d\*x + c)^(1/3)\*b + a)\*a\*b^6\*c^2 + a^2\*b^6\*c^2 - 2\*((d\*x + c)^(1/3)\*b + a)^5\*b^3\*c + 10\*((d\*x + c)^(1/3)\*b + a)^4\*a\*b^3\*c - 20\*((d\*x + c)^(1/3)\*b + a)^3\*a^2\*b^3\*c + 20\*((d\*x + c)^(1/3)\*b + a)^2\*a^3\*b^3\*c - 10\*((d\*x + c)^(1/3)\*b + a)\*a^4\*b^3\*c + 2\*a^5\*b^3\*c + ((d\*x + c)^(1/3)\*b + a)^8 - 8\*((d\*x + c)^(1/3)\*b + a)^7\*a + 28\*((d\*x + c)^(1/3)\*b + a)^6\*a^2 - 56\*((d\*x + c)^(1/3)\*b + a)^5\*a^3 + 70\*((d\*x + c)^(1/3)\*b + a)^4\*a^4 - 56\*((d\*x + c)^(1/3)\*b + a)^3\*a^5 + 28\*((d\*x + c)^(1/3)\*b + a)^2\*a^6 - 8\*((d\*x + c)^(1/3)\*b + a)\*a^7 + a^8 - 2\*b^6\*c^2 + 40\*((d\*x + c)^(1/3)\*b + a)^3\*b^3\*c - 120\*((d\*x + c)^(1/3)\*b + a)^2\*a\*b^3\*c + 120\*((d\*x + c)^(1/3)\*b + a)\*a^2\*b^3\*c - 40\*a^3\*b^3\*c - 56\*((d\*x + c)^(1/3)\*b + a)^6 + 336\*((d\*x + c)^(1/3)\*b + a)^5\*a - 840\*((d\*x + c)^(1/3)\*b + a)^4\*a^2 + 1120\*((d\*x + c)^(1/3)\*b + a)^3\*a^3 - 840\*((d\*x + c)^(1/3)\*b + a)^2\*a^4 + 336\*((d\*x + c)^(1/3)\*b + a)\*a^5 - 56\*a^6 - 240\*((d\*x

$$\begin{aligned}
& + c)^{(1/3)} * b + a) * b^3 * c + 240 * a * b^3 * c + 1680 * ((d * x + c)^{(1/3)} * b + a)^4 - 6 \\
& 720 * ((d * x + c)^{(1/3)} * b + a)^3 * a + 10080 * ((d * x + c)^{(1/3)} * b + a)^2 * a^2 - 672 \\
& 0 * ((d * x + c)^{(1/3)} * b + a) * a^3 + 1680 * a^4 - 20160 * ((d * x + c)^{(1/3)} * b + a)^2 \\
& + 40320 * ((d * x + c)^{(1/3)} * b + a) * a - 20160 * a^2 + 40320) * \sin((d * x + c)^{(1/3)} * \\
& b + a) / (b^8 * d^2) / (b * d)
\end{aligned}$$

### 3.96 $\int x \cos(a + b\sqrt[3]{c + dx}) dx$

**Optimal.** Leaf size=261

$$\frac{60(c + dx) \sin(a + b\sqrt[3]{c + dx})}{b^3 d^2} + \frac{360\sqrt[3]{c + dx} \sin(a + b\sqrt[3]{c + dx})}{b^5 d^2} + \frac{6c \sin(a + b\sqrt[3]{c + dx})}{b^3 d^2} + \frac{15(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^2}$$

```
[Out] (360*Cos[a + b*(c + d*x)^(1/3)])/(b^6*d^2) - (6*c*(c + d*x)^(1/3)*Cos[a + b*(c + d*x)^(1/3)])/(b^2*d^2) - (180*(c + d*x)^(2/3)*Cos[a + b*(c + d*x)^(1/3)])/(b^4*d^2) + (15*(c + d*x)^(4/3)*Cos[a + b*(c + d*x)^(1/3)])/(b^2*d^2) + (6*c*Sin[a + b*(c + d*x)^(1/3)])/(b^3*d^2) + (360*(c + d*x)^(1/3)*Sin[a + b*(c + d*x)^(1/3)])/(b^5*d^2) - (3*c*(c + d*x)^(2/3)*Sin[a + b*(c + d*x)^(1/3)])/(b*d^2) - (60*(c + d*x)*Sin[a + b*(c + d*x)^(1/3)])/(b^3*d^2) + (3*(c + d*x)^(5/3)*Sin[a + b*(c + d*x)^(1/3)])/(b*d^2)
```

**Rubi [A]** time = 0.231053, antiderivative size = 261, normalized size of antiderivative = 1., number of steps used = 11, number of rules used = 4, integrand size = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$ , Rules used = {3432, 3296, 2637, 2638}

$$\frac{60(c + dx) \sin(a + b\sqrt[3]{c + dx})}{b^3 d^2} + \frac{360\sqrt[3]{c + dx} \sin(a + b\sqrt[3]{c + dx})}{b^5 d^2} + \frac{6c \sin(a + b\sqrt[3]{c + dx})}{b^3 d^2} + \frac{15(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^2}$$

Antiderivative was successfully verified.

```
[In] Int[x*Cos[a + b*(c + d*x)^(1/3)], x]
```

```
[Out] (360*Cos[a + b*(c + d*x)^(1/3)])/(b^6*d^2) - (6*c*(c + d*x)^(1/3)*Cos[a + b*(c + d*x)^(1/3)])/(b^2*d^2) - (180*(c + d*x)^(2/3)*Cos[a + b*(c + d*x)^(1/3)])/(b^4*d^2) + (15*(c + d*x)^(4/3)*Cos[a + b*(c + d*x)^(1/3)])/(b^2*d^2) + (6*c*Sin[a + b*(c + d*x)^(1/3)])/(b^3*d^2) + (360*(c + d*x)^(1/3)*Sin[a + b*(c + d*x)^(1/3)])/(b^5*d^2) - (3*c*(c + d*x)^(2/3)*Sin[a + b*(c + d*x)^(1/3)])/(b*d^2) - (60*(c + d*x)*Sin[a + b*(c + d*x)^(1/3)])/(b^3*d^2) + (3*(c + d*x)^(5/3)*Sin[a + b*(c + d*x)^(1/3)])/(b*d^2)
```

#### Rule 3432

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])*(b_.)^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Dist[1/(n*f), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - (e*h)/f + (h*x^(1/n))/f)^m, x], x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

#### Rule 3296

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := -Simp[(c + d*x)^m*Cos[e + f*x]/f, x] + Dist[(d*m)/f, Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

#### Rule 2637

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

#### Rule 2638

`Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

### Rubi steps

$$\begin{aligned}
 \int x \cos(a + b\sqrt[3]{c + dx}) dx &= \frac{3 \operatorname{Subst}\left(\int \left(-\frac{cx^2 \cos(a+bx)}{d} + \frac{x^5 \cos(a+bx)}{d}\right) dx, x, \sqrt[3]{c + dx}\right)}{d} \\
 &= \frac{3 \operatorname{Subst}\left(\int x^5 \cos(a + bx) dx, x, \sqrt[3]{c + dx}\right)}{d^2} - \frac{(3c) \operatorname{Subst}\left(\int x^2 \cos(a + bx) dx, x, \sqrt[3]{c + dx}\right)}{d^2} \\
 &= -\frac{3c(c + dx)^{2/3} \sin(a + b\sqrt[3]{c + dx})}{bd^2} + \frac{3(c + dx)^{5/3} \sin(a + b\sqrt[3]{c + dx})}{bd^2} - \frac{15 \operatorname{Subst}\left(\int x^4 \sin(a + bx) dx, x, \sqrt[3]{c + dx}\right)}{bd^2} \\
 &= -\frac{6c\sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} + \frac{15(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} - \frac{3c(c + dx)^{2/3} \sin(a + b\sqrt[3]{c + dx})}{bd^2} \\
 &= -\frac{6c\sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} + \frac{15(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} + \frac{6c \sin(a + b\sqrt[3]{c + dx})}{b^3d^2} \\
 &= -\frac{6c\sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} - \frac{180(c + dx)^{2/3} \cos(a + b\sqrt[3]{c + dx})}{b^4d^2} + \frac{15(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} \\
 &= -\frac{6c\sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} - \frac{180(c + dx)^{2/3} \cos(a + b\sqrt[3]{c + dx})}{b^4d^2} + \frac{15(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} \\
 &= \frac{360 \cos(a + b\sqrt[3]{c + dx})}{b^6d^2} - \frac{6c\sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2d^2} - \frac{180(c + dx)^{2/3} \cos(a + b\sqrt[3]{c + dx})}{b^4d^2}
 \end{aligned}$$

**Mathematica [A]** time = 0.417346, size = 117, normalized size = 0.45

$$\frac{3(b(b^4 dx(c + dx)^{2/3} - 2b^2(9c + 10dx) + 120\sqrt[3]{c + dx}) \sin(a + b\sqrt[3]{c + dx}) + (b^4\sqrt[3]{c + dx}(3c + 5dx) - 60b^2(c + dx)^{2/3} + 120c\sqrt[3]{c + dx}) \cos(a + b\sqrt[3]{c + dx})}{b^6d^2}$$

Antiderivative was successfully verified.

[In] `Integrate[x*Cos[a + b*(c + d*x)^(1/3)], x]`

[Out]  $(3*((120 - 60*b^2*(c + d*x)^{(2/3)} + b^4*(c + d*x)^{(1/3)}*(3*c + 5*d*x))*\cos[a + b*(c + d*x)^{(1/3)}] + b*(120*(c + d*x)^{(1/3)} + b^4*d*x*(c + d*x)^{(2/3)} - 2*b^2*(9*c + 10*d*x))*\sin[a + b*(c + d*x)^{(1/3)}])/(b^6*d^2)$

**Maple [B]** time = 0.035, size = 655, normalized size = 2.5

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*cos(a+b*(d*x+c)^(1/3)), x)`

[Out]  $3/d^2/b^3*(-c*((a+b*(d*x+c)^{(1/3)})^2*\sin(a+b*(d*x+c)^{(1/3)})-2*\sin(a+b*(d*x+c)^{(1/3}))+2*(a+b*(d*x+c)^{(1/3}))*\cos(a+b*(d*x+c)^{(1/3}))+2*c*a*(\cos(a+b*(d*x+c)^{(1/3}))+2*(a+b*(d*x+c)^{(1/3}))*\sin(a+b*(d*x+c)^{(1/3}))-a^2*c*\sin(a+b*(d*x+c)^{(1/3}))+1/b^3*((a+b*(d*x+c)^{(1/3)})^5*\sin(a+b*(d*x+c)^{(1/3}))+5*(a+b*(d*x+c)^{(1/3)})^4*\cos(a+b*(d*x+c)^{(1/3}))-20*(a+b*(d*x+c)^{(1/3)})^3*\sin(a+b*(d*x+c)^{(1/3}))-60*(a+b*(d*x+c)^{(1/3)})^2*\cos(a+b*(d*x+c)^{(1/3}))+120*\cos(a+b*(d*x+c)^{(1/3}))+120*(a+b*(d*x+c)^{(1/3}))*\sin(a+b*(d*x+c)^{(1/3}))-5/b^3*a*((a+b*(d*x+c)$

$$\begin{aligned} & \wedge(1/3))^4*\sin(a+b*(d*x+c)^(1/3))+4*(a+b*(d*x+c)^(1/3))^3*\cos(a+b*(d*x+c)^(1/3))-12*(a+b*(d*x+c)^(1/3))^2*\sin(a+b*(d*x+c)^(1/3))+24*\sin(a+b*(d*x+c)^(1/3))-24*(a+b*(d*x+c)^(1/3))*\cos(a+b*(d*x+c)^(1/3))+10/b^3*a^2*((a+b*(d*x+c)^(1/3))^3*\sin(a+b*(d*x+c)^(1/3))+3*(a+b*(d*x+c)^(1/3))^2*\cos(a+b*(d*x+c)^(1/3))-6*\cos(a+b*(d*x+c)^(1/3))-6*(a+b*(d*x+c)^(1/3))*\sin(a+b*(d*x+c)^(1/3))) \\ & -10/b^3*a^3*((a+b*(d*x+c)^(1/3))^2*\sin(a+b*(d*x+c)^(1/3))-2*\sin(a+b*(d*x+c)^(1/3))+2*(a+b*(d*x+c)^(1/3))*\cos(a+b*(d*x+c)^(1/3)))+5/b^3*a^4*(\cos(a+b*(d*x+c)^(1/3))+(a+b*(d*x+c)^(1/3))*\sin(a+b*(d*x+c)^(1/3)))-1/b^3*a^5*\sin(a+b*(d*x+c)^(1/3))) \end{aligned}$$

**Maxima [B]** time = 1.25364, size = 706, normalized size = 2.7

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(a+b\*(d\*x+c)^(1/3)),x, algorithm="maxima")

[Out] 
$$\begin{aligned} & -3*(a^2*c*\sin((d*x + c)^(1/3)*b + a) - 2*(((d*x + c)^(1/3)*b + a)*\sin((d*x + c)^(1/3)*b + a) + \cos((d*x + c)^(1/3)*b + a))*a*c + a^5*\sin((d*x + c)^(1/3)*b + a)/b^3 - 5*(((d*x + c)^(1/3)*b + a)*\sin((d*x + c)^(1/3)*b + a) + \cos((d*x + c)^(1/3)*b + a))*a^4/b^3 + (2*(((d*x + c)^(1/3)*b + a)*\cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2 - 2)*\sin((d*x + c)^(1/3)*b + a))*c + 10*(2*(((d*x + c)^(1/3)*b + a)*\cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2 - 2)*\sin((d*x + c)^(1/3)*b + a))*a^3/b^3 - 10*(3*(((d*x + c)^(1/3)*b + a)^2 - 2)*\cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*\sin((d*x + c)^(1/3)*b + a))*a^2/b^3 + 5*(4*(((d*x + c)^(1/3)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*\cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^4 - 12*(((d*x + c)^(1/3)*b + a)^2 + 24)*\sin((d*x + c)^(1/3)*b + a))*a/b^3 - (5*(((d*x + c)^(1/3)*b + a)^4 - 12*(((d*x + c)^(1/3)*b + a)^2 + 24)*\cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^5 - 20*(((d*x + c)^(1/3)*b + a)^3 + 120*(d*x + c)^(1/3)*b + 120*a)*\sin((d*x + c)^(1/3)*b + a))/b^3)/(b^3*d^2) \end{aligned}$$

**Fricas [A]** time = 1.63794, size = 293, normalized size = 1.12

$$\frac{3 \left( \left( 60(dx+c)^{\frac{2}{3}}b^2 - (5b^4dx + 3b^4c)(dx+c)^{\frac{1}{3}} - 120 \right) \cos \left( (dx+c)^{\frac{1}{3}}b + a \right) - \left( (dx+c)^{\frac{2}{3}}b^5dx - 20b^3dx - 18b^3c + 120 \right) \sin \left( (dx+c)^{\frac{1}{3}}b + a \right) \right)}{b^6d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(a+b\*(d\*x+c)^(1/3)),x, algorithm="fricas")

[Out] 
$$\begin{aligned} & -3*((60*(d*x + c)^(2/3)*b^2 - (5*b^4*d*x + 3*b^4*c)*(d*x + c)^(1/3) - 120)*\cos((d*x + c)^(1/3)*b + a) - ((d*x + c)^(2/3)*b^5*d*x - 20*b^3*d*x - 18*b^3*c + 120*(d*x + c)^(1/3)*b)*\sin((d*x + c)^(1/3)*b + a))/(b^6*d^2) \end{aligned}$$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int x \cos \left( a + b\sqrt[3]{c + dx} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(a+b\*(d\*x+c)\*\*(1/3)),x)

[Out] Integral(x\*cos(a + b\*(c + d\*x)\*\*(1/3)), x)

**Giac [A]** time = 1.23773, size = 500, normalized size = 1.92

$$3 \left( \frac{2 \left( (dx+c)^{\frac{1}{3}} b+a \right) b^3 c - 2 a b^3 c - 5 \left( (dx+c)^{\frac{1}{3}} b+a \right)^4 + 20 \left( (dx+c)^{\frac{1}{3}} b+a \right)^3 a - 30 \left( (dx+c)^{\frac{1}{3}} b+a \right)^2 a^2 + 20 \left( (dx+c)^{\frac{1}{3}} b+a \right) a^3 - 5 a^4 + 60 \left( (dx+c)^{\frac{1}{3}} b+a \right)^2 - 120 \left( (dx+c)^{\frac{1}{3}} b+a \right)}{b^5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x\*cos(a+b\*(d\*x+c)^(1/3)),x, algorithm="giac")

[Out]  $-3 \left( (2 \left( (dx+c)^{\frac{1}{3}} b+a \right) b^3 c - 2 a b^3 c - 5 \left( (dx+c)^{\frac{1}{3}} b+a \right)^4 + 20 \left( (dx+c)^{\frac{1}{3}} b+a \right)^3 a - 30 \left( (dx+c)^{\frac{1}{3}} b+a \right)^2 a^2 + 20 \left( (dx+c)^{\frac{1}{3}} b+a \right) a^3 - 5 a^4 + 60 \left( (dx+c)^{\frac{1}{3}} b+a \right)^2 - 120 \left( (dx+c)^{\frac{1}{3}} b+a \right) a + 60 a^2 - 120 \right) \cos \left( (dx+c)^{\frac{1}{3}} b+a \right) / b^5 + \left( (dx+c)^{\frac{1}{3}} b+a \right)^2 b^3 c - 2 \left( (dx+c)^{\frac{1}{3}} b+a \right) a b^3 c + a^2 b^3 c - \left( (dx+c)^{\frac{1}{3}} b+a \right)^5 + 5 \left( (dx+c)^{\frac{1}{3}} b+a \right)^4 a - 10 \left( (dx+c)^{\frac{1}{3}} b+a \right)^3 a^2 + 10 \left( (dx+c)^{\frac{1}{3}} b+a \right)^2 a^3 - 5 \left( (dx+c)^{\frac{1}{3}} b+a \right) a^4 + a^5 - 2 b^3 c + 20 \left( (dx+c)^{\frac{1}{3}} b+a \right)^3 - 60 \left( (dx+c)^{\frac{1}{3}} b+a \right)^2 a + 60 \left( (dx+c)^{\frac{1}{3}} b+a \right) a^2 - 20 a^3 - 120 \left( (dx+c)^{\frac{1}{3}} b \right) \sin \left( (dx+c)^{\frac{1}{3}} b+a \right) / b^5 \right) / (b d^2)$

### 3.97 $\int \cos\left(a + b\sqrt[3]{c + dx}\right) dx$

**Optimal.** Leaf size=85

$$-\frac{6 \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^3 d} + \frac{6\sqrt[3]{c + dx} \cos\left(a + b\sqrt[3]{c + dx}\right)}{b^2 d} + \frac{3(c + dx)^{2/3} \sin\left(a + b\sqrt[3]{c + dx}\right)}{bd}$$

[Out]  $(6*(c + d*x)^{(1/3)}*Cos[a + b*(c + d*x)^{(1/3)})/(b^2*d) - (6*Sin[a + b*(c + d*x)^{(1/3)})/(b^3*d) + (3*(c + d*x)^{(2/3)}*Sin[a + b*(c + d*x)^{(1/3)})/(b*d)$

**Rubi [A]** time = 0.0551264, antiderivative size = 85, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 3, integrand size = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3362, 3296, 2637}

$$-\frac{6 \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^3 d} + \frac{6\sqrt[3]{c + dx} \cos\left(a + b\sqrt[3]{c + dx}\right)}{b^2 d} + \frac{3(c + dx)^{2/3} \sin\left(a + b\sqrt[3]{c + dx}\right)}{bd}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b\*(c + d\*x)^(1/3)], x]

[Out]  $(6*(c + d*x)^{(1/3)}*Cos[a + b*(c + d*x)^{(1/3)})/(b^2*d) - (6*Sin[a + b*(c + d*x)^{(1/3)})/(b^3*d) + (3*(c + d*x)^{(2/3)}*Sin[a + b*(c + d*x)^{(1/3)})/(b*d)$

#### Rule 3362

Int[((a\_.) + Cos[(c\_.) + (d\_.)\*((e\_.) + (f\_.)\*(x\_))^(n\_)])\*(b\_.)^(p\_.), x\_Symbol] :> Dist[1/(n\*f), Subst[Int[x^(1/n - 1)\*(a + b\*Cos[c + d\*x])^p, x], x, (e + f\*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]

#### Rule 3296

Int[((c\_.) + (d\_.)\*(x\_))^(m\_.)\*sin[(e\_.) + (f\_.)\*(x\_)], x\_Symbol] :> -Simp[(c + d\*x)^m\*Cos[e + f\*x]/f, x] + Dist[(d\*m)/f, Int[(c + d\*x)^(m - 1)\*Cos[e + f\*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

#### Rule 2637

Int[sin[Pi/2 + (c\_.) + (d\_.)\*(x\_)], x\_Symbol] :> Simp[Sin[c + d\*x]/d, x] /; FreeQ[{c, d}, x]

#### Rubi steps

$$\begin{aligned} \int \cos\left(a + b\sqrt[3]{c + dx}\right) dx &= \frac{3 \text{Subst}\left(\int x^2 \cos(a + bx) dx, x, \sqrt[3]{c + dx}\right)}{d} \\ &= \frac{3(c + dx)^{2/3} \sin\left(a + b\sqrt[3]{c + dx}\right)}{bd} - \frac{6 \text{Subst}\left(\int x \sin(a + bx) dx, x, \sqrt[3]{c + dx}\right)}{bd} \\ &= \frac{6\sqrt[3]{c + dx} \cos\left(a + b\sqrt[3]{c + dx}\right)}{b^2 d} + \frac{3(c + dx)^{2/3} \sin\left(a + b\sqrt[3]{c + dx}\right)}{bd} - \frac{6 \text{Subst}\left(\int \cos(a + bx) dx, x, \sqrt[3]{c + dx}\right)}{b^2 d} \\ &= \frac{6\sqrt[3]{c + dx} \cos\left(a + b\sqrt[3]{c + dx}\right)}{b^2 d} - \frac{6 \sin\left(a + b\sqrt[3]{c + dx}\right)}{b^3 d} + \frac{3(c + dx)^{2/3} \sin\left(a + b\sqrt[3]{c + dx}\right)}{bd} \end{aligned}$$

**Mathematica [A]** time = 0.111096, size = 65, normalized size = 0.76

$$\frac{3 \left( b^2 (c + dx)^{2/3} - 2 \right) \sin \left( a + b \sqrt[3]{c + dx} \right) + 6b \sqrt[3]{c + dx} \cos \left( a + b \sqrt[3]{c + dx} \right)}{b^3 d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b\*(c + d\*x)^(1/3)],x]

[Out] (6\*b\*(c + d\*x)^(1/3)\*Cos[a + b\*(c + d\*x)^(1/3)] + 3\*(-2 + b^2\*(c + d\*x)^(2/3))\*Sin[a + b\*(c + d\*x)^(1/3)]/(b^3\*d)

**Maple [A]** time = 0.034, size = 131, normalized size = 1.5

$$\frac{3 \left( a + b \sqrt[3]{dx + c} \right)^2 \sin \left( a + b \sqrt[3]{dx + c} \right) - 2 \sin \left( a + b \sqrt[3]{dx + c} \right) + 2 \left( a + b \sqrt[3]{dx + c} \right) \cos \left( a + b \sqrt[3]{dx + c} \right) - 2a \left( \cos \left( a + b \sqrt[3]{dx + c} \right) \right)}{db^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*(d\*x+c)^(1/3)),x)

[Out] 3/d/b^3\*((a+b\*(d\*x+c)^(1/3))^2\*sin(a+b\*(d\*x+c)^(1/3))-2\*sin(a+b\*(d\*x+c)^(1/3))+2\*(a+b\*(d\*x+c)^(1/3))\*cos(a+b\*(d\*x+c)^(1/3))-2\*a\*(cos(a+b\*(d\*x+c)^(1/3)))+(a+b\*(d\*x+c)^(1/3))\*sin(a+b\*(d\*x+c)^(1/3)))+a^2\*sin(a+b\*(d\*x+c)^(1/3)))

**Maxima [A]** time = 1.11794, size = 159, normalized size = 1.87

$$\frac{3 \left( a^2 \sin \left( (dx + c)^{1/3} b + a \right) - 2 \left( \left( (dx + c)^{1/3} b + a \right) \sin \left( (dx + c)^{1/3} b + a \right) + \cos \left( (dx + c)^{1/3} b + a \right) \right) a + 2 \left( (dx + c)^{1/3} b + a \right) \cos \left( (dx + c)^{1/3} b + a \right) \right)}{b^3 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3)),x, algorithm="maxima")

[Out] 3\*(a^2\*sin((d\*x + c)^(1/3)\*b + a) - 2\*((d\*x + c)^(1/3)\*b + a)\*sin((d\*x + c)^(1/3)\*b + a) + cos((d\*x + c)^(1/3)\*b + a)\*a + 2\*((d\*x + c)^(1/3)\*b + a)\*cos((d\*x + c)^(1/3)\*b + a) + (((d\*x + c)^(1/3)\*b + a)^2 - 2)\*sin((d\*x + c)^(1/3)\*b + a))/(b^3\*d)

**Fricas [A]** time = 1.62542, size = 155, normalized size = 1.82

$$\frac{3 \left( 2 (dx + c)^{1/3} b \cos \left( (dx + c)^{1/3} b + a \right) + \left( (dx + c)^{2/3} b^2 - 2 \right) \sin \left( (dx + c)^{1/3} b + a \right) \right)}{b^3 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3)),x, algorithm="fricas")

[Out] 3\*(2\*(d\*x + c)^(1/3)\*b\*cos((d\*x + c)^(1/3)\*b + a) + ((d\*x + c)^(2/3)\*b^2 - 2)\*sin((d\*x + c)^(1/3)\*b + a))/(b^3\*d)



---

**Sympy [A]** time = 1.67818, size = 94, normalized size = 1.11

$$\begin{cases} x \cos(a) & \text{for } b = 0 \wedge (b = 0 \vee d = 0) \\ x \cos\left(a + b\sqrt[3]{c}\right) & \text{for } d = 0 \\ \frac{3(c+dx)^{\frac{2}{3}} \sin\left(a+b\sqrt[3]{c+dx}\right)}{bd} + \frac{6\sqrt[3]{c+dx} \cos\left(a+b\sqrt[3]{c+dx}\right)}{b^2d} - \frac{6 \sin\left(a+b\sqrt[3]{c+dx}\right)}{b^3d} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)\*\*(1/3)),x)

[Out] Piecewise((x\*cos(a), Eq(b, 0) & (Eq(b, 0) | Eq(d, 0))), (x\*cos(a + b\*c\*\*(1/3)), Eq(d, 0)), (3\*(c + d\*x)\*\*(2/3)\*sin(a + b\*(c + d\*x)\*\*(1/3))/(b\*d) + 6\*(c + d\*x)\*\*(1/3)\*cos(a + b\*(c + d\*x)\*\*(1/3))/(b\*\*2\*d) - 6\*sin(a + b\*(c + d\*x)\*\*(1/3))/(b\*\*3\*d), True))

---

**Giac [A]** time = 1.26277, size = 109, normalized size = 1.28

$$\frac{3 \left( \frac{2(dx+c)^{\frac{1}{3}} \cos\left((dx+c)^{\frac{1}{3}}b+a\right)}{b} + \frac{\left( \left( (dx+c)^{\frac{1}{3}}b+a \right)^2 - 2 \left( (dx+c)^{\frac{1}{3}}b+a \right) a + a^2 - 2 \right) \sin\left( (dx+c)^{\frac{1}{3}}b+a \right)}{b^2} \right)}{bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3)),x, algorithm="giac")

[Out] 3\*(2\*(d\*x + c)^(1/3)\*cos((d\*x + c)^(1/3)\*b + a)/b + (((d\*x + c)^(1/3)\*b + a)^2 - 2\*((d\*x + c)^(1/3)\*b + a)\*a + a^2 - 2)\*sin((d\*x + c)^(1/3)\*b + a)/b^2)/(b\*d)

$$3.98 \quad \int \frac{\cos\left(a+b\sqrt[3]{c+dx}\right)}{x} dx$$

**Optimal.** Leaf size=234

$$\cos\left(a+b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) + \cos\left(a+(-1)^{2/3}b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left((-1)^{2/3}b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) + \cos\left(a-(-1)^{1/3}b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(-b\sqrt[3]{c+dx}\right)$$

```
[Out] Cos[a + b*c^(1/3)]*CosIntegral[b*c^(1/3) - b*(c + d*x)^(1/3)] + Cos[a + (-1)^(2/3)*b*c^(1/3)]*CosIntegral[(-1)^(2/3)*b*c^(1/3) - b*(c + d*x)^(1/3)] + Cos[a - (-1)^(1/3)*b*c^(1/3)]*CosIntegral[(-1)^(1/3)*b*c^(1/3) + b*(c + d*x)^(1/3)] + Sin[a + b*c^(1/3)]*SinIntegral[b*c^(1/3) - b*(c + d*x)^(1/3)] + Sin[a + (-1)^(2/3)*b*c^(1/3)]*SinIntegral[(-1)^(2/3)*b*c^(1/3) - b*(c + d*x)^(1/3)] - Sin[a - (-1)^(1/3)*b*c^(1/3)]*SinIntegral[(-1)^(1/3)*b*c^(1/3) + b*(c + d*x)^(1/3)]
```

**Rubi [A]** time = 0.468404, antiderivative size = 234, normalized size of antiderivative = 1., number of steps used = 11, number of rules used = 4, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {3432, 3303, 3299, 3302}

$$\cos\left(a+b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) + \cos\left(a+(-1)^{2/3}b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left((-1)^{2/3}b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) + \cos\left(a-(-1)^{1/3}b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(-b\sqrt[3]{c+dx}\right)$$

Antiderivative was successfully verified.

```
[In] Int[Cos[a + b*(c + d*x)^(1/3)]/x,x]
```

```
[Out] Cos[a + b*c^(1/3)]*CosIntegral[b*c^(1/3) - b*(c + d*x)^(1/3)] + Cos[a + (-1)^(2/3)*b*c^(1/3)]*CosIntegral[(-1)^(2/3)*b*c^(1/3) - b*(c + d*x)^(1/3)] + Cos[a - (-1)^(1/3)*b*c^(1/3)]*CosIntegral[(-1)^(1/3)*b*c^(1/3) + b*(c + d*x)^(1/3)] + Sin[a + b*c^(1/3)]*SinIntegral[b*c^(1/3) - b*(c + d*x)^(1/3)] + Sin[a + (-1)^(2/3)*b*c^(1/3)]*SinIntegral[(-1)^(2/3)*b*c^(1/3) - b*(c + d*x)^(1/3)] - Sin[a - (-1)^(1/3)*b*c^(1/3)]*SinIntegral[(-1)^(1/3)*b*c^(1/3) + b*(c + d*x)^(1/3)]
```

#### Rule 3432

```
Int[((a_.) + Cos[(c_.) + (d_.)*(e_.) + (f_.)*(x_)])^(n_)]*(b_.)^(p_.)*((g_.) + (h_.)*(x_)^(m_.), x_Symbol] := Dist[1/(n*f), Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - (e*h)/f + (h*x^(1/n))/f)^m, x], x], x, (e + f*x)^n, x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

#### Rule 3303

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[(c*f)/d + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[(c*f)/d + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

#### Rule 3299

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

#### Rule 3302

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

### Rubi steps

$$\begin{aligned} \int \frac{\cos(a + b\sqrt[3]{c + dx})}{x} dx &= \frac{3 \operatorname{Subst}\left(\int \left(\frac{d \cos(ax)}{3(\sqrt[3]{c-x})} - \frac{d \cos(ax)}{3(-\sqrt[3]{-1}\sqrt[3]{c-x})} - \frac{d \cos(ax)}{3((-1)^{2/3}\sqrt[3]{c-x})}\right) dx, x, \sqrt[3]{c + dx}\right)}{d} \\ &= -\operatorname{Subst}\left(\int \frac{\cos(a + bx)}{\sqrt[3]{c-x}} dx, x, \sqrt[3]{c + dx}\right) - \operatorname{Subst}\left(\int \frac{\cos(a + bx)}{-\sqrt[3]{-1}\sqrt[3]{c-x}} dx, x, \sqrt[3]{c + dx}\right) - \operatorname{Subst}\left(\int \frac{\cos(a + bx)}{\sqrt[3]{(-1)^{2/3}\sqrt[3]{c-x}}}\right) \\ &= -\left(\cos(a + b\sqrt[3]{c}) \operatorname{Subst}\left(\int \frac{\cos(b\sqrt[3]{c} - bx)}{\sqrt[3]{c-x}} dx, x, \sqrt[3]{c + dx}\right)\right) - \cos\left(a - \sqrt[3]{-1}b\sqrt[3]{c}\right) \operatorname{Subst}\left(\int \frac{\cos(b\sqrt[3]{c} - bx)}{\sqrt[3]{c-x}} dx, x, \sqrt[3]{c + dx}\right) \\ &= \cos(a + b\sqrt[3]{c}) \operatorname{Ci}\left(b\sqrt[3]{c} - b\sqrt[3]{c + dx}\right) + \cos\left(a + (-1)^{2/3}b\sqrt[3]{c}\right) \operatorname{Ci}\left((-1)^{2/3}b\sqrt[3]{c} - b\sqrt[3]{c + dx}\right) \end{aligned}$$

**Mathematica [C]** time = 0.433259, size = 243, normalized size = 1.04

$$\frac{1}{2} \left( \operatorname{RootSum}\left[c - \#1^3 \&, -i \sin(\#1b + a) \operatorname{CosIntegral}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) + \cos(\#1b + a) \operatorname{CosIntegral}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) \right)$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[Cos[a + b*(c + d*x)^(1/3)]/x, x]
```

```
[Out] (RootSum[c - #1^3 &, Cos[a + b*#1]*CosIntegral[b*((c + d*x)^(1/3) - #1]] - I*CosIntegral[b*((c + d*x)^(1/3) - #1)]*Sin[a + b*#1] - I*Cos[a + b*#1]*SinIntegral[b*((c + d*x)^(1/3) - #1]] - Sin[a + b*#1]*SinIntegral[b*((c + d*x)^(1/3) - #1)] & ] + RootSum[c - #1^3 &, Cos[a + b*#1]*CosIntegral[b*((c + d*x)^(1/3) - #1]] + I*CosIntegral[b*((c + d*x)^(1/3) - #1)]*Sin[a + b*#1] + I*Cos[a + b*#1]*SinIntegral[b*((c + d*x)^(1/3) - #1]] - Sin[a + b*#1]*SinIntegral[b*((c + d*x)^(1/3) - #1)] & ])/2
```

**Maple [C]** time = 0.149, size = 279, normalized size = 1.2

$$3 \frac{1}{b^3} \left( \frac{1}{3} b^3 \sum_{_R1=\operatorname{RootOf}(-b^3c+_Z^3-3a_Z^2+3a^2_Z-a^3)} \frac{_R1^2 \left( \operatorname{Si}\left(-b\sqrt[3]{dx+c}+_R1-a\right) \sin(_R1) + \operatorname{Ci}\left(b\sqrt[3]{dx+c}-_R1\right) \right)}{_R1^2 - 2_R1 a + a^2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a+b*(d*x+c)^(1/3))/x, x)
```

```
[Out] 3/b^3*(1/3*b^3*sum(_R1^2/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^(1/3)+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^(1/3)-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))-2/3*a*b^3*sum(_R1/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^(1/3)+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^(1/3)-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))+1/3*a^2*b^3*sum(1/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^(1/3)+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^(1/3)-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3)))
```

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(\frac{1}{3}b(dx+c) + a\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3))/x,x, algorithm="maxima")

[Out] integrate(cos((d\*x + c)^(1/3)\*b + a)/x, x)

**Fricas [C]** time = 1.77086, size = 833, normalized size = 3.56

$$\frac{1}{2} \operatorname{Ei}\left(i(dx+c)^{\frac{1}{3}}b + \frac{1}{2}(ib^3c)^{\frac{1}{3}}(-i\sqrt{3}-1)\right)e^{\frac{1}{2}(ib^3c)^{\frac{1}{3}}(i\sqrt{3}+1)+ia} + \frac{1}{2} \operatorname{Ei}\left(-i(dx+c)^{\frac{1}{3}}b + \frac{1}{2}(-ib^3c)^{\frac{1}{3}}(-i\sqrt{3}-1)\right)e^{\frac{1}{2}(-i$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3))/x,x, algorithm="fricas")

[Out] 1/2\*Ei(I\*(d\*x + c)^(1/3)\*b + 1/2\*(I\*b^3\*c)^(1/3)\*(-I\*sqrt(3) - 1))\*e^(1/2\*(I\*b^3\*c)^(1/3)\*(I\*sqrt(3) + 1) + I\*a) + 1/2\*Ei(-I\*(d\*x + c)^(1/3)\*b + 1/2\*(-I\*b^3\*c)^(1/3)\*(-I\*sqrt(3) - 1))\*e^(1/2\*(-I\*b^3\*c)^(1/3)\*(I\*sqrt(3) + 1) - I\*a) + 1/2\*Ei(I\*(d\*x + c)^(1/3)\*b + 1/2\*(I\*b^3\*c)^(1/3)\*(I\*sqrt(3) - 1))\*e^(1/2\*(I\*b^3\*c)^(1/3)\*(-I\*sqrt(3) + 1) + I\*a) + 1/2\*Ei(-I\*(d\*x + c)^(1/3)\*b + 1/2\*(-I\*b^3\*c)^(1/3)\*(I\*sqrt(3) - 1))\*e^(1/2\*(-I\*b^3\*c)^(1/3)\*(-I\*sqrt(3) + 1) - I\*a) + 1/2\*Ei(I\*(d\*x + c)^(1/3)\*b + (I\*b^3\*c)^(1/3))\*e^(I\*a - (I\*b^3\*c)^(1/3)) + 1/2\*Ei(-I\*(d\*x + c)^(1/3)\*b + (-I\*b^3\*c)^(1/3))\*e^(-I\*a - (-I\*b^3\*c)^(1/3))

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)\*\*(1/3))/x,x)

[Out] Integral(cos(a + b\*(c + d\*x)\*\*(1/3))/x, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(\frac{1}{3}b(dx+c) + a\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3))/x,x, algorithm="giac")

[Out] integrate(cos((d\*x + c)^(1/3)\*b + a)/x, x)

$$3.99 \quad \int \frac{\cos\left(a+b\sqrt[3]{c+dx}\right)}{x^2} dx$$

**Optimal.** Leaf size=332

$$\frac{bd \sin\left(a+b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right)}{3c^{2/3}} + \frac{\sqrt[3]{-1}bd \sin\left(a-\sqrt[3]{-1}b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(b\sqrt[3]{c+dx}+\sqrt[3]{-1}b\sqrt[3]{c}\right)}{3c^{2/3}}$$

[Out]  $-(\operatorname{Cos}[a + b*(c + d*x)^{(1/3)}]/x) - (b*d*\operatorname{CosIntegral}[b*c^{(1/3)} - b*(c + d*x)^{(1/3)}]*\operatorname{Sin}[a + b*c^{(1/3)}])/(3*c^{(2/3)}) + ((-1)^{(1/3)}*b*d*\operatorname{CosIntegral}[(-1)^{(1/3)}*b*c^{(1/3)} + b*(c + d*x)^{(1/3)}]*\operatorname{Sin}[a - (-1)^{(1/3)}*b*c^{(1/3)}])/(3*c^{(2/3)}) - ((-1)^{(2/3)}*b*d*\operatorname{CosIntegral}[(-1)^{(2/3)}*b*c^{(1/3)} - b*(c + d*x)^{(1/3)}]*\operatorname{Sin}[a + (-1)^{(2/3)}*b*c^{(1/3)}])/(3*c^{(2/3)}) + (b*d*\operatorname{Cos}[a + b*c^{(1/3)}]*\operatorname{SinIntegral}[b*c^{(1/3)} - b*(c + d*x)^{(1/3)}])/(3*c^{(2/3)}) + ((-1)^{(2/3)}*b*d*\operatorname{Cos}[a + (-1)^{(2/3)}*b*c^{(1/3)}]*\operatorname{SinIntegral}[(-1)^{(2/3)}*b*c^{(1/3)} - b*(c + d*x)^{(1/3)}])/(3*c^{(2/3)}) + ((-1)^{(1/3)}*b*d*\operatorname{Cos}[a - (-1)^{(1/3)}*b*c^{(1/3)}]*\operatorname{SinIntegral}[(-1)^{(1/3)}*b*c^{(1/3)} + b*(c + d*x)^{(1/3)}])/(3*c^{(2/3)})$

**Rubi [A]** time = 0.741003, antiderivative size = 332, normalized size of antiderivative = 1., number of steps used = 13, number of rules used = 6, integrand size = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3432, 3342, 3333, 3303, 3299, 3302}

$$\frac{bd \sin\left(a+b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right)}{3c^{2/3}} + \frac{\sqrt[3]{-1}bd \sin\left(a-\sqrt[3]{-1}b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(b\sqrt[3]{c+dx}+\sqrt[3]{-1}b\sqrt[3]{c}\right)}{3c^{2/3}}$$

Antiderivative was successfully verified.

[In]  $\operatorname{Int}[\operatorname{Cos}[a + b*(c + d*x)^{(1/3)}]/x^2, x]$

[Out]  $-(\operatorname{Cos}[a + b*(c + d*x)^{(1/3)}]/x) - (b*d*\operatorname{CosIntegral}[b*c^{(1/3)} - b*(c + d*x)^{(1/3)}]*\operatorname{Sin}[a + b*c^{(1/3)}])/(3*c^{(2/3)}) + ((-1)^{(1/3)}*b*d*\operatorname{CosIntegral}[(-1)^{(1/3)}*b*c^{(1/3)} + b*(c + d*x)^{(1/3)}]*\operatorname{Sin}[a - (-1)^{(1/3)}*b*c^{(1/3)}])/(3*c^{(2/3)}) - ((-1)^{(2/3)}*b*d*\operatorname{CosIntegral}[(-1)^{(2/3)}*b*c^{(1/3)} - b*(c + d*x)^{(1/3)}]*\operatorname{Sin}[a + (-1)^{(2/3)}*b*c^{(1/3)}])/(3*c^{(2/3)}) + (b*d*\operatorname{Cos}[a + b*c^{(1/3)}]*\operatorname{SinIntegral}[b*c^{(1/3)} - b*(c + d*x)^{(1/3)}])/(3*c^{(2/3)}) + ((-1)^{(2/3)}*b*d*\operatorname{Cos}[a + (-1)^{(2/3)}*b*c^{(1/3)}]*\operatorname{SinIntegral}[(-1)^{(2/3)}*b*c^{(1/3)} - b*(c + d*x)^{(1/3)}])/(3*c^{(2/3)}) + ((-1)^{(1/3)}*b*d*\operatorname{Cos}[a - (-1)^{(1/3)}*b*c^{(1/3)}]*\operatorname{SinIntegral}[(-1)^{(1/3)}*b*c^{(1/3)} + b*(c + d*x)^{(1/3)}])/(3*c^{(2/3)})$

**Rule 3432**

$\operatorname{Int}[(a_.) + \operatorname{Cos}[(c_.) + (d_.)*(e_.) + (f_.)*(x_.)]^{(n_.)}]^{(p_.)}*(g_.) + (h_.)*(x_.)]^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Dist}[1/(n*f), \operatorname{Subst}[\operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b*\operatorname{Cos}[c + d*x])^p, x^{(1/n - 1)}*(g - (e*h)/f + (h*x^{(1/n)})/f)^m, x], x, (e + f*x)^n], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, g, h, m\}, x\} \&\& \operatorname{IGtQ}[p, 0] \&\& \operatorname{IntegerQ}[1/n]$

**Rule 3342**

$\operatorname{Int}[\operatorname{Cos}[(c_.) + (d_.)*(x_.)]^{(m_.)}*(e_.)*(x_.)]^{(p_.)}*(a_.) + (b_.)*(x_.)]^{(n_.)}]^{(p_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[(e^m*(a + b*x^n)^{(p + 1)}*\operatorname{Cos}[c + d*x])/(b*n*(p + 1)), x] + \operatorname{Dist}[(d*e^m)/(b*n*(p + 1)), \operatorname{Int}[(a + b*x^n)^{(p + 1)}*\operatorname{Sin}[c + d*x], x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, m, n\}, x\} \&\& \operatorname{ILtQ}[p, -1] \&\& \operatorname{EqQ}[m, n - 1] \&\& (\operatorname{IntegerQ}[n] \parallel \operatorname{GtQ}[e, 0])$

**Rule 3333**

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*Sin[(c_.) + (d_.)*(x_)], x_Symbol] := Int
[ExpandIntegrand[Sin[c + d*x], (a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d},
x] && ILtQ[p, 0] && IGtQ[n, 0] && (EqQ[n, 2] || EqQ[p, -1])
```

### Rule 3303

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*
e - c*f)/d], Int[Sin[(c*f)/d + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f
)/d], Int[Cos[(c*f)/d + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] &&
NeQ[d*e - c*f, 0]
```

### Rule 3299

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinInte
gral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

### Rule 3302

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosInte
gral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) -
c*f, 0]
```

### Rubi steps

$$\begin{aligned}
\int \frac{\cos(a + b\sqrt[3]{c + dx})}{x^2} dx &= \frac{3 \operatorname{Subst} \left( \int \frac{x^2 \cos(a + bx)}{\left(-\frac{c}{d} + \frac{x^3}{d}\right)^2} dx, x, \sqrt[3]{c + dx} \right)}{d} \\
&= -\frac{\cos(a + b\sqrt[3]{c + dx})}{x} - b \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{-\frac{c}{d} + \frac{x^3}{d}} dx, x, \sqrt[3]{c + dx} \right) \\
&= -\frac{\cos(a + b\sqrt[3]{c + dx})}{x} - b \operatorname{Subst} \left( \int \left( -\frac{d \sin(a + bx)}{3c^{2/3}(\sqrt[3]{c} - x)} - \frac{d \sin(a + bx)}{3c^{2/3}(\sqrt[3]{c} + \sqrt[3]{-1}x)} - \frac{d \sin(a + bx)}{3c^{2/3}(\sqrt[3]{c} - \sqrt[3]{-1}x)} \right) dx, x, \sqrt[3]{c + dx} \right) \\
&= -\frac{\cos(a + b\sqrt[3]{c + dx})}{x} + \frac{(bd) \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{\sqrt[3]{c} - x} dx, x, \sqrt[3]{c + dx} \right)}{3c^{2/3}} + \frac{(bd) \operatorname{Subst} \left( \int \frac{\sin(a + bx)}{\sqrt[3]{c} + \sqrt[3]{-1}x} dx, x, \sqrt[3]{c + dx} \right)}{3c^{2/3}} \\
&= -\frac{\cos(a + b\sqrt[3]{c + dx})}{x} - \frac{(bd \cos(a + b\sqrt[3]{c})) \operatorname{Subst} \left( \int \frac{\sin(b\sqrt[3]{c} - bx)}{\sqrt[3]{c} - x} dx, x, \sqrt[3]{c + dx} \right)}{3c^{2/3}} + \frac{(bd \cos(a + b\sqrt[3]{c})) \operatorname{Subst} \left( \int \frac{\sin(b\sqrt[3]{c} - bx)}{\sqrt[3]{c} + \sqrt[3]{-1}x} dx, x, \sqrt[3]{c + dx} \right)}{3c^{2/3}} \\
&= -\frac{\cos(a + b\sqrt[3]{c + dx})}{x} - \frac{bd \operatorname{Ci}(b\sqrt[3]{c} - b\sqrt[3]{c + dx}) \sin(a + b\sqrt[3]{c})}{3c^{2/3}} + \frac{\sqrt[3]{-1}bd \operatorname{Ci}(\sqrt[3]{-1}b\sqrt[3]{c} + b\sqrt[3]{c + dx}) \sin(a + b\sqrt[3]{c})}{3c^{2/3}}
\end{aligned}$$

**Mathematica [C]** time = 0.620772, size = 138, normalized size = 0.42

$$-\frac{1}{6} ibd \operatorname{RootSum} \left[ c - \#1^3 \&, \frac{e^{-i\#1b - ia} \operatorname{Ei} \left( -ib \left( \sqrt[3]{c + dx} - \#1 \right) \right)}{\#1^2} \& \right] + \frac{1}{6} ibd \operatorname{RootSum} \left[ c - \#1^3 \&, \frac{e^{i\#1b + ia} \operatorname{Ei} \left( ib \left( \sqrt[3]{c + dx} - \#1 \right) \right)}{\#1^2} \& \right]$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*(c + d*x)^(1/3)]/x^2, x]
```

```
[Out] -(Cos[a + b*(c + d*x)^(1/3)]/x) - (I/6)*b*d*RootSum[c - #1^3 &, (E^((-I)*a
- I*b*#1)*ExpIntegralEi[(-I)*b*((c + d*x)^(1/3) - #1)]/#1^2 & ] + (I/6)*b
```

\*d\*RootSum[c - #1^3 & , (E^(I\*a + I\*b\*#1)\*ExpIntegralEi[I\*b\*((c + d\*x)^(1/3) - #1))/#1^2 & ]

**Maple [C]** time = 0.32, size = 933, normalized size = 2.8

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a+b\*(d\*x+c)^(1/3))/x^2,x)

[Out]  $3*d/b^3*(\cos(a+b*(d*x+c)^{1/3}))*(-2/3*a*b^3/c*(a+b*(d*x+c)^{1/3})^2+a^2*b^3/c*(a+b*(d*x+c)^{1/3})-1/3*b^3*(b^3*c+a^3)/c)/(-b^3*c+(a+b*(d*x+c)^{1/3})^3-3*(a+b*(d*x+c)^{1/3})^2*a+3*a^2*(a+b*(d*x+c)^{1/3})-a^3)-2/9*a*b^3/c*\sum(_R1/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^{1/3}+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^{1/3}-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))-1/9*b^3/c*\sum((b^3*c+2*_RR1^2*a-3*_RR1*a^2+a^3)/(_RR1^2-2*_RR1*a+a^2)*(-Si(-b*(d*x+c)^{1/3}+_RR1-a)*cos(_RR1)+Ci(b*(d*x+c)^{1/3}-_RR1+a)*sin(_RR1)),_RR1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))+cos(a+b*(d*x+c)^{1/3})*(2/3*a*b^3/c*(a+b*(d*x+c)^{1/3})^2-2/3*a^2*b^3/c*(a+b*(d*x+c)^{1/3}))/(-b^3*c+(a+b*(d*x+c)^{1/3})^3-3*(a+b*(d*x+c)^{1/3})^2*a+3*a^2*(a+b*(d*x+c)^{1/3})-a^3)+2/9*a*b^3/c*\sum((_R1+a)/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^{1/3}+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^{1/3}-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))+2/9*a*b^3/c*\sum(_RR1/(_RR1-a)*(-Si(-b*(d*x+c)^{1/3}+_RR1-a)*cos(_RR1)+Ci(b*(d*x+c)^{1/3}-_RR1+a)*sin(_RR1)),_RR1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))+a^2*b^6*(cos(a+b*(d*x+c)^{1/3})*(-1/3/b^3/c*(a+b*(d*x+c)^{1/3}))+1/3*a/b^3/c)/(-b^3*c+(a+b*(d*x+c)^{1/3})^3-3*(a+b*(d*x+c)^{1/3})^2*a+3*a^2*(a+b*(d*x+c)^{1/3})-a^3)-2/9/b^3/c*\sum(1/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^{1/3}+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^{1/3}-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))-1/9/b^3/c*\sum(1/(_RR1-a)*(-Si(-b*(d*x+c)^{1/3}+_RR1-a)*cos(_RR1)+Ci(b*(d*x+c)^{1/3}-_RR1+a)*sin(_RR1)),_RR1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3)))))$

**Maxima [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left((dx+c)^{\frac{1}{3}}b+a\right)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3))/x^2,x, algorithm="maxima")

[Out] integrate(cos((d\*x + c)^(1/3)\*b + a)/x^2, x)

**Fricas [C]** time = 2.05983, size = 1137, normalized size = 3.42

$$2 (ib^3c)^{\frac{1}{3}} dx Ei\left(i(dx+c)^{\frac{1}{3}}b + (ib^3c)^{\frac{1}{3}}\right) e^{\left(ia-(ib^3c)^{\frac{1}{3}}\right)} + 2 (-ib^3c)^{\frac{1}{3}} dx Ei\left(-i(dx+c)^{\frac{1}{3}}b + (-ib^3c)^{\frac{1}{3}}\right) e^{\left(-ia-(ib^3c)^{\frac{1}{3}}\right)} -$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3))/x^2,x, algorithm="fricas")

[Out] 
$$-1/12*(2*(I*b^3*c)^(1/3)*d*x*Ei(I*(d*x + c)^(1/3)*b + (I*b^3*c)^(1/3))*e^{(I*a - (I*b^3*c)^(1/3))} + 2*(-I*b^3*c)^(1/3)*d*x*Ei(-I*(d*x + c)^(1/3)*b + (-I*b^3*c)^(1/3))*e^{(-I*a - (-I*b^3*c)^(1/3))} - (I*b^3*c)^(1/3)*(I*\sqrt{3}*d*x + d*x)*Ei(I*(d*x + c)^(1/3)*b + 1/2*(I*b^3*c)^(1/3)*(-I*\sqrt{3} - 1))*e^{(1/2*(I*b^3*c)^(1/3)*(I*\sqrt{3} + 1) + I*a)} - (-I*b^3*c)^(1/3)*(I*\sqrt{3}*d*x + d*x)*Ei(-I*(d*x + c)^(1/3)*b + 1/2*(-I*b^3*c)^(1/3)*(-I*\sqrt{3} - 1))*e^{(1/2*(-I*b^3*c)^(1/3)*(I*\sqrt{3} + 1) - I*a)} - (I*b^3*c)^(1/3)*(-I*\sqrt{3})*d*x + d*x)*Ei(I*(d*x + c)^(1/3)*b + 1/2*(I*b^3*c)^(1/3)*(I*\sqrt{3} - 1))*e^{(1/2*(I*b^3*c)^(1/3)*(-I*\sqrt{3} + 1) + I*a)} - (-I*b^3*c)^(1/3)*(-I*\sqrt{3})*d*x + d*x)*Ei(-I*(d*x + c)^(1/3)*b + 1/2*(-I*b^3*c)^(1/3)*(I*\sqrt{3} - 1))*e^{(1/2*(-I*b^3*c)^(1/3)*(-I*\sqrt{3} + 1) - I*a)} + 12*c*cos((d*x + c)^(1/3)*b + a))/(c*x)$$

**Sympy [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)\*\*(1/3))/x\*\*2,x)

[Out] Integral(cos(a + b\*(c + d\*x)\*\*(1/3))/x\*\*2, x)

**Giac [F]** time = 0., size = 0, normalized size = 0.

$$\int \frac{\cos\left((dx + c)^{\frac{1}{3}}b + a\right)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(a+b\*(d\*x+c)^(1/3))/x^2,x, algorithm="giac")

[Out] integrate(cos((d\*x + c)^(1/3)\*b + a)/x^2, x)



# Chapter 4

## Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

### 4.0.1 Mathematica and Rubi grading function

```
1 (* Original version thanks to Albert Rich emailed on 03/21/2017 *)
2 (* ::Package:: *)
3
4 (* ::Subsection:: *)
5 (*GradeAntiderivative[result,optimal]*)
6
7
8 (* ::Text:: *)
9 (*If result and optimal are mathematical expressions, *)
10 (*      GradeAntiderivative[result,optimal] returns*)
11 (* "F" if the result fails to integrate an expression that*)
12 (*   is integrable*)
13 (* "C" if result involves higher level functions than necessary*)
14 (* "B" if result is more than twice the size of the optimal*)
15 (*   antiderivative*)
16 (* "A" if result can be considered optimal*)
17
18
19 GradeAntiderivative[result_,optimal_] :=
20   If[ExpnType[result]<=ExpnType[optimal],
21     If[FreeQ[result,Complex] || Not[FreeQ[optimal,Complex]],
22       If[LeafCount[result]<=2*LeafCount[optimal],
23         "A",
24         "B"],
25       "C"],
26     If[FreeQ[result,Integrate] && FreeQ[result,Int],
27       "C",
28       "F"]]
29
30
31 (* ::Text:: *)
32 (*The following summarizes the type number assigned an *)
33 (*expression based on the functions it involves*)
34 (*1 = rational function*)
35 (*2 = algebraic function*)
36 (*3 = elementary function*)
37 (*4 = special function*)
```

```

38 (*5 = hyperpergeometric function*)
39 (*6 = appell function*)
40 (*7 = rootsum function*)
41 (*8 = integrate function*)
42 (*9 = unknown function*)
43
44
45 ExpnType[expn_] :=
46   If[AtomQ[expn],
47     1,
48     If[ListQ[expn],
49       Max[Map[ExpnType,expn]],
50       If[Head[expn]===Power,
51         If[IntegerQ[expn[[2]]],
52           ExpnType[expn[[1]]],
53           If[Head[expn[[2]]]===Rational,
54             If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
55               1,
56               Max[ExpnType[expn[[1]],2]],
57             Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
58             If[Head[expn]===Plus || Head[expn]===Times,
59               Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
60             If[ElementaryFunctionQ[Head[expn]],
61               Max[3,ExpnType[expn[[1]]],
62             If[SpecialFunctionQ[Head[expn]],
63               Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
64             If[HypergeometricFunctionQ[Head[expn]],
65               Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
66             If[AppellFunctionQ[Head[expn]],
67               Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
68             If[Head[expn]===RootSum,
69               Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
70             If[Head[expn]===Integrate || Head[expn]===Int,
71               Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
72             9]]]]]]]]]]
73
74
75 ElementaryFunctionQ[func_] :=
76   MemberQ[{
77     Exp,Log,
78     Sin,Cos,Tan,Cot,Sec,Csc,
79     ArcSin,ArcCos,ArcTan,ArcCot,ArcSec,ArcCsc,
80     Sinh,Cosh,Tanh,Coth,Sech,Csch,
81     ArcSinh,ArcCosh,ArcTanh,ArcCoth,ArcSech,ArcCsch
82   },func]
83
84
85 SpecialFunctionQ[func_] :=
86   MemberQ[{
87     Erf, Erfc, Erfi,
88     FresnelS, FresnelC,
89     ExpIntegralE, ExpIntegralEi, LogIntegral,
90     SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
91     Gamma, LogGamma, PolyGamma,
92     Zeta, PolyLog, ProductLog,
93     EllipticF, EllipticE, EllipticPi
94   },func]
95
96
97 HypergeometricFunctionQ[func_] :=
98   MemberQ[{Hypergeometric1F1,Hypergeometric2F1,HypergeometricPFQ},func]
99
100

```

```

101 AppellFunctionQ[func_] :=
102   MemberQ[{AppellF1},func]

```

## 4.0.2 Maple grading function

```

1 # File: GradeAntiderivative.mpl
2 # Original version thanks to Albert Rich emailed on 03/21/2017
3
4 #Nasser 03/22/2017 Use Maple leaf count instead since buildin
5 #Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
6 #Nasser 03/24/2017 corrected the check for complex result
7 #Nasser 10/27/2017 check for leafsize and do not call ExpnType()
8 # if leaf size is "too large". Set at 500,000
9 #Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
10 # see problem 156, file Apostol_Problems
11
12 GradeAntiderivative := proc(result,optimal)
13 local leaf_count_result, leaf_count_optimal,ExpnType_result,ExpnType_optimal,
14     debug:=false;
15
16     leaf_count_result:=leafcount(result);
17     #do NOT call ExpnType() if leaf size is too large. Recursion problem
18     if leaf_count_result > 500000 then
19         return "B";
20     fi;
21
22     leaf_count_optimal:=leafcount(optimal);
23
24     ExpnType_result:=ExpnType(result);
25     ExpnType_optimal:=ExpnType(optimal);
26
27     if debug then
28         print("ExpnType_result",ExpnType_result," ExpnType_optimal=",
29             ExpnType_optimal);
30     fi;
31
32 # If result and optimal are mathematical expressions,
33 # GradeAntiderivative[result,optimal] returns
34 # "F" if the result fails to integrate an expression that
35 # is integrable
36 # "C" if result involves higher level functions than necessary
37 # "B" if result is more than twice the size of the optimal
38 # antiderivative
39 # "A" if result can be considered optimal
40
41 #This check below actually is not needed, since I only
42 #call this grading only for passed integrals. i.e. I check
43 #for "F" before calling this. But no harm of keeping it here.
44 #just in case.
45
46 if not type(result,freeof('int')) then
47     return "F";
48 end if;
49
50 if ExpnType_result<=ExpnType_optimal then
51     if debug then
52         print("ExpnType_result<=ExpnType_optimal");
53     fi;
54     if is_contains_complex(result) then
55         if is_contains_complex(optimal) then
56             if debug then

```

```

57         print("both result and optimal complex");
58         fi;
59         #both result and optimal complex
60         if leaf_count_result<=2*leaf_count_optimal then
61             return "A";
62         else
63             return "B";
64         end if
65     else #result contains complex but optimal is not
66         if debug then
67             print("result contains complex but optimal is not");
68         fi;
69         return "C";
70     end if
71 else # result do not contain complex
72     # this assumes optimal do not as well
73     if debug then
74         print("result do not contain complex, this assumes optimal do
not as well");
75     fi;
76     if leaf_count_result<=2*leaf_count_optimal then
77         if debug then
78             print("leaf_count_result<=2*leaf_count_optimal");
79         fi;
80         return "A";
81     else
82         if debug then
83             print("leaf_count_result>2*leaf_count_optimal");
84         fi;
85         return "B";
86     end if
87 end if
88 else #ExpnType(result) > ExpnType(optimal)
89     if debug then
90         print("ExpnType(result) > ExpnType(optimal)");
91     fi;
92     return "C";
93 end if
94
95 end proc:
96
97 #
98 # is_contains_complex(result)
99 # takes expressions and returns true if it contains "I" else false
100 #
101 #Nasser 032417
102 is_contains_complex:= proc(expression)
103     return (has(expression,I));
104 end proc:
105
106 # The following summarizes the type number assigned an expression
107 # based on the functions it involves
108 # 1 = rational function
109 # 2 = algebraic function
110 # 3 = elementary function
111 # 4 = special function
112 # 5 = hyperpergeometric function
113 # 6 = appell function
114 # 7 = rootsum function
115 # 8 = integrate function
116 # 9 = unknown function
117
118 ExpnType := proc(expn)

```

```

119   if type(expn,'atomic') then
120     1
121   elif type(expn,'list') then
122     apply(max,map(ExpnType,expn))
123   elif type(expn,'sqrt') then
124     if type(op(1,expn),'rational') then
125       1
126     else
127       max(2,ExpnType(op(1,expn)))
128     end if
129   elif type(expn,'`^`') then
130     if type(op(2,expn),'integer') then
131       ExpnType(op(1,expn))
132     elif type(op(2,expn),'rational') then
133       if type(op(1,expn),'rational') then
134         1
135       else
136         max(2,ExpnType(op(1,expn)))
137       end if
138     else
139       max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
140     end if
141   elif type(expn,'`+`') or type(expn,'`*`') then
142     max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
143   elif ElementaryFunctionQ(op(0,expn)) then
144     max(3,ExpnType(op(1,expn)))
145   elif SpecialFunctionQ(op(0,expn)) then
146     max(4,apply(max,map(ExpnType,[op(expn)])))
147   elif HypergeometricFunctionQ(op(0,expn)) then
148     max(5,apply(max,map(ExpnType,[op(expn)])))
149   elif AppellFunctionQ(op(0,expn)) then
150     max(6,apply(max,map(ExpnType,[op(expn)])))
151   elif op(0,expn)='int' then
152     max(8,apply(max,map(ExpnType,[op(expn)]))) else
153     9
154   end if
155 end proc:
156
157
158 ElementaryFunctionQ := proc(func)
159   member(func,[
160     exp,log,ln,
161     sin,cos,tan,cot,sec,csc,
162     arcsin,arccos,arctan,arccot,arcsec,arccsc,
163     sinh,cosh,tanh,coth,sech,csch,
164     arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
165 end proc:
166
167 SpecialFunctionQ := proc(func)
168   member(func,[
169     erf,erfc,erfi,
170     FresnelS,FresnelC,
171     Ei,Ei,Li,Si,Ci,Shi,Chi,
172     GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
173     EllipticF,EllipticE,EllipticPi])
174 end proc:
175
176 HypergeometricFunctionQ := proc(func)
177   member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
178 end proc:
179
180 AppellFunctionQ := proc(func)
181   member(func,[AppellF1])

```

```

182 end proc:
183
184 # u is a sum or product. rest(u) returns all but the
185 # first term or factor of u.
186 rest := proc(u) local v;
187     if nops(u)=2 then
188         op(2,u)
189     else
190         apply(op(0,u),op(2..nops(u),u))
191     end if
192 end proc:
193
194 #leafcount(u) returns the number of nodes in u.
195 #Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
196 leafcount := proc(u)
197     MmaTranslator[Mma][LeafCount](u);
198 end proc:

```

### 4.0.3 Sympy grading function

```

1 #Dec 24, 2019. Nasser M. Abbasi:
2 #           Port of original Maple grading function by
3 #           Albert Rich to use with Sympy/Python
4 #Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
5 #           added 'exp_polar'
6 from sympy import *
7
8 def leaf_count(expr):
9     #sympy do not have leaf count function. This is approximation
10    return round(1.7*count_ops(expr))
11
12 def is_sqrt(expr):
13     if isinstance(expr,Pow):
14         if expr.args[1] == Rational(1,2):
15             return True
16         else:
17             return False
18     else:
19         return False
20
21 def is_elementary_function(func):
22     return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
23                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
24                    asinh,acosh,atanh,acoth,asech,acsch
25                    ]
26
27 def is_special_function(func):
28     return func in [ erf,erfc,erfi,
29                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
30                    gamma,loggamma,digamma,zeta,polylog,LambertW,
31                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
32                    ]
33
34 def is_hypergeometric_function(func):
35     return func in [hyper]
36
37 def is_appell_function(func):
38     return func in [appellf1]
39
40 def is_atom(expn):
41     try:
42         if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
43             return True

```

```

44     else:
45         return False
46
47     except AttributeError as error:
48         return False
49
50 def expnType(expn):
51     debug=False
52     if debug:
53         print("expn=",expn,"type(expn)=",type(expn))
54
55     if is_atom(expn):
56         return 1
57     elif isinstance(expn,list):
58         return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
59     elif is_sqrt(expn):
60         if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
61             return 1
62         else:
63             return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
64     elif isinstance(expn,Pow): #type(expn,``^`)
65         if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
66             return expnType(expn.args[0]) #ExpnType(op(1,expn))
67         elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
68             if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
69                 return 1
70             else:
71                 return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn
72 )))
73     else:
74         return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,
75 ExpnType(op(1,expn)),ExpnType(op(2,expn)))
76     elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or
77 type(expn,``*`)
78     m1 = expnType(expn.args[0])
79     m2 = expnType(list(expn.args[1:]))
80     return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
81     elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
82     return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
83     elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
84     m1 = max(map(expnType, list(expn.args)))
85     return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
86     elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,
87 expn))
88     m1 = max(map(expnType, list(expn.args)))
89     return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
90     elif is_appell_function(expn.func):
91     m1 = max(map(expnType, list(expn.args)))
92     return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
93     elif isinstance(expn,RootSum):
94     m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType
95 ,Apply[List,expn]],7]],
96     return max(7,m1)
97     elif str(expn).find("Integral") != -1:
98     m1 = max(map(expnType, list(expn.args)))
99     return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
100     else:
101     return 9
102
103 #main function
104 def grade_antiderivative(result,optimal):
105
106     leaf_count_result = leaf_count(result)

```

```

102 leaf_count_optimal = leaf_count(optimal)
103
104 expnType_result = expnType(result)
105 expnType_optimal = expnType(optimal)
106
107 if str(result).find("Integral") != -1:
108     return "F"
109
110 if expnType_result <= expnType_optimal:
111     if result.has(I):
112         if optimal.has(I): #both result and optimal complex
113             if leaf_count_result <= 2*leaf_count_optimal:
114                 return "A"
115             else:
116                 return "B"
117         else: #result contains complex but optimal is not
118             return "C"
119     else: # result do not contain complex, this assumes optimal do not as
well
120         if leaf_count_result <= 2*leaf_count_optimal:
121             return "A"
122         else:
123             return "B"
124 else:
125     return "C"

```

#### 4.0.4 SageMath grading function

```

1 #Dec 24, 2019. Nasser: Ported original Maple grading function by
2 #     Albert Rich to use with Sagemath. This is used to
3 #     grade Fracas, Giac and Maxima results.
4 #Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
5 #     'arctan2','floor','abs','log_integral'
6
7 from sage.all import *
8 from sage.symbolic.operators import add_vararg, mul_vararg
9
10 def tree(expr):
11     debug=False;
12     if debug:
13         print ("Enter tree(expr), expr=",expr)
14         print ("expr.operator()=",expr.operator())
15         print ("expr.operands()=",expr.operands())
16         print ("map(tree, expr.operands()=",map(tree, expr.operands()))
17
18     if expr.operator() is None:
19         return expr
20     else:
21         return [expr.operator()+list(map(tree, expr.operands()))
22
23 def leaf_count(anti):
24     debug=False;
25
26     if debug: print ("Enter leaf_count, anti=", anti, " len(anti)=", len(anti))
27
28     if len(anti) == 0: #special check for optimal being 0 for some test cases.
29         if debug: print ("len(anti) == 0")
30         return 1
31     else:
32         if debug: print ("round(1.35*len(flatten(tree(anti))))=",round(1.35*len
(flatten(tree(anti))))
33         return round(1.35*len(flatten(tree(anti)))) #fudge factor
34             #since this estimate of leaf count is bit lower than

```



```

35         #what it should be compared to Mathematica's
36
37 def is_sqrt(expr):
38     debug=False;
39     if expr.operator() == operator.pow: #isinstance(expr,Pow):
40         if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
41             if debug: print ("expr is sqrt")
42             return True
43         else:
44             return False
45     else:
46         return False
47
48 def is_elementary_function(func):
49     debug = False
50
51     m = func.name() in ['exp','log','ln',
52         'sin','cos','tan','cot','sec','csc',
53         'arcsin','arccos','arctan','arccot','arcsec','arccsc',
54         'sinh','cosh','tanh','coth','sech','csch',
55         'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
56         'arctan2','floor','abs'
57     ]
58     if debug:
59         if m:
60             print ("func ", func , " is elementary_function")
61         else:
62             print ("func ", func , " is NOT elementary_function")
63
64
65     return m
66
67 def is_special_function(func):
68     debug = False
69
70     if debug: print ("type(func)=", type(func))
71
72     m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
73         'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','
74     sinh_integral'
75         'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
76         'polylog','lambert_w','elliptic_f','elliptic_e',
77         'elliptic_pi','exp_integral_e','log_integral']
78
79     if debug:
80         print ("m=",m)
81         if m:
82             print ("func ", func , " is special_function")
83         else:
84             print ("func ", func , " is NOT special_function")
85
86     return m
87
88
89 def is_hypergeometric_function(func):
90     return func.name() in ['hypergeometric','hypergeometric_M','
91     hypergeometric_U']
92
93 def is_appell_function(func):
94     return func.name() in ['hypergeometric'] #[appellf1] can't find this in
95     sagemath

```

```

95 def is_atom(expn):
96
97     #thanks to answer at https://ask.sagemath.org/question/49179/what-is-
sagemath-equivalent-to-atomic-type-in-maple/
98     try:
99         if expn.parent() is SR:
100             return expn.operator() is None
101         if expn.parent() in (ZZ, QQ, AA, QQbar):
102             return expn in expn.parent() # Should always return True
103         if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens")
:
104             return expn in expn.parent().base_ring() or expn in expn.parent().
gens()
105         return False
106
107     except AttributeError as error:
108         return False
109
110
111 def expnType(expn):
112     debug=False
113
114     if debug:
115         print(">>>>Enter expnType, expn=", expn)
116         print(">>>>is_atom(expn)=", is_atom(expn))
117
118     if is_atom(expn):
119         return 1
120     elif type(expn)==list: #instance(expn,list):
121         return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
122     elif is_sqrt(expn):
123         if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],
Rational):
124             return 1
125         else:
126             return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
127     elif expn.operator() == operator.pow: #instance(expn,Pow)
128         if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer
)
129             return expnType(expn.operands()[0]) #expnType(expn.args[0])
130         elif type(expn.operands()[1])==Rational: #instance(expn.args[1],
Rational)
131             if type(expn.operands()[0])==Rational: #instance(expn.args[0],
Rational)
132                 return 1
133             else:
134                 return max(2,expnType(expn.operands()[0])) #max(2,expnType(
expn.args[0]))
135         else:
136             return max(3,expnType(expn.operands()[0]),expnType(expn.operands()
[1])) #max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))
137     elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #
instance(expn,Add) or instance(expn,Mul)
138         m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
139         m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
140         return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn)))
141     elif is_elementary_function(expn.operator()): #is_elementary_function(expn
.func)
142         return max(3,expnType(expn.operands()[0]))
143     elif is_special_function(expn.operator()): #is_special_function(expn.func)
144         m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))

```

```

145     return max(4,m1)    #max(4,m1)
146     elif is_hypergeometric_function(expn.operator()): #
is_hypergeometric_function(expn.func)
147         m1 = max(map(expnType, expn.operands()))    #max(map(expnType, list(
expn.args)))
148         return max(5,m1)    #max(5,m1)
149     elif is_appell_function(expn.operator()):
150         m1 = max(map(expnType, expn.operands()))    #max(map(expnType, list(
expn.args)))
151         return max(6,m1)    #max(6,m1)
152     elif str(expn).find("Integral") != -1: #this will never happen, since it
153         #is checked before calling the grading function that is passed.
154         #but kept it here.
155         m1 = max(map(expnType, expn.operands()))    #max(map(expnType, list(
expn.args)))
156         return max(8,m1)    #max(5,apply(max,map(ExpnType,[op(expn)])))
157     else:
158         return 9
159
160 #main function
161 def grade_antiderivative(result,optimal):
162     debug = False;
163
164     if debug: print ("Enter grade_antiderivative for sagemath")
165
166     leaf_count_result = leaf_count(result)
167     leaf_count_optimal = leaf_count(optimal)
168
169     if debug: print ("leaf_count_result=", leaf_count_result, "
leaf_count_optimal=",leaf_count_optimal)
170
171
172     expnType_result = expnType(result)
173     expnType_optimal = expnType(optimal)
174
175     if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",
expnType_optimal)
176
177     if expnType_result <= expnType_optimal:
178         if result.has(I):
179             if optimal.has(I): #both result and optimal complex
180                 if leaf_count_result <= 2*leaf_count_optimal:
181                     return "A"
182                 else:
183                     return "B"
184             else: #result contains complex but optimal is not
185                 return "C"
186         else: # result do not contain complex, this assumes optimal do not as
well
187             if leaf_count_result <= 2*leaf_count_optimal:
188                 return "A"
189             else:
190                 return "B"
191     else:
192         return "C"

```