


A Solution Manual For

Elementary Differential Equations,
Martin, Reissner, 2nd ed, 1961



Nasser M. Abbasi

May 15, 2024

Contents

1 Exercis 2, page 5

2

1 Exercis 2, page 5

1.1	problem 2(a)	3
1.2	problem 2(b)	6
1.3	problem 2(c)	9
1.4	problem 2(d)	12
1.5	problem 2(e)	15
1.6	problem 2(a)	18
1.7	problem 3(a)	21
1.8	problem 3(b)	36
1.9	problem 3(c)	50
1.10	problem 3(d)	63
1.11	problem 3(e)	76
1.12	problem 3(f)	88
1.13	problem 3(g)	91
1.14	problem 3(h)	94
1.15	problem 3(i)	97

1.1 problem 2(a)

1.1.1 Solving as quadrature ode	3
1.1.2 Maple step by step solution	4

Internal problem ID [2432]

Internal file name [OUTPUT/1924_Sunday_June_05_2022_02_39_42_AM_58612681/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = 2$$

1.1.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int 2 \, dx \\ &= 2x + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = 2x + c_1 \tag{1}$$




Figure 1: Slope field plot

Verification of solutions

$$y = 2x + c_1$$

Verified OK.

1.1.2 Maple step by step solution

Let's solve

$$y' = 2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int 2 dx + c_1$$

- Evaluate integral

$$y = 2x + c_1$$

- Solve for y

$$y = 2x + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=2,y(x), singsol=all)
```

$$y(x) = 2x + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 11

```
DSolve[y'[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x + c_1$$

1.2 problem 2(b)

1.2.1 Solving as quadrature ode	6
1.2.2 Maple step by step solution	7

Internal problem ID [2433]

Internal file name [OUTPUT/1925_Sunday_June_05_2022_02_39_43_AM_28780283/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = 2 e^{3x}$$

1.2.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int 2 e^{3x} dx \\ &= \frac{2 e^{3x}}{3} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{2 e^{3x}}{3} + c_1 \tag{1}$$




Figure 2: Slope field plot

Verification of solutions

$$y = \frac{2e^{3x}}{3} + c_1$$

Verified OK.

1.2.2 Maple step by step solution

Let's solve

$$y' = 2e^{3x}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int 2e^{3x} dx + c_1$$

- Evaluate integral

$$y = \frac{2e^{3x}}{3} + c_1$$

- Solve for y

$$y = \frac{2e^{3x}}{3} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=2*exp(3*x),y(x), singsol=all)
```

$$y(x) = \frac{2e^{3x}}{3} + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 17

```
DSolve[y'[x]==2*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2e^{3x}}{3} + c_1$$

1.3 problem 2(c)

1.3.1 Solving as quadrature ode	9
1.3.2 Maple step by step solution	10

Internal problem ID [2434]

Internal file name [OUTPUT/1926_Sunday_June_05_2022_02_39_45_AM_20320997/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = \frac{2}{\sqrt{-x^2 + 1}}$$

1.3.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{2}{\sqrt{-x^2 + 1}} dx \\ &= 2 \arcsin(x) + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = 2 \arcsin(x) + c_1 \tag{1}$$




Figure 3: Slope field plot

Verification of solutions

$$y = 2 \arcsin(x) + c_1$$

Verified OK.

1.3.2 Maple step by step solution

Let's solve

$$y' = \frac{2}{\sqrt{-x^2+1}}$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{2}{\sqrt{-x^2+1}} dx + c_1$$

- Evaluate integral

$$y = 2 \arcsin(x) + c_1$$

- Solve for y

$$y = 2 \arcsin(x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=2/sqrt(1-x^2),y(x), singsol=all)
```

$$y(x) = 2 \arcsin(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 28

```
DSolve[y'[x]==2/Sqrt[1-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -4 \arctan\left(\frac{\sqrt{1-x^2}}{x+1}\right) + c_1$$

1.4 problem 2(d)

1.4.1 Solving as quadrature ode	12
1.4.2 Maple step by step solution	13

Internal problem ID [2435]

Internal file name [OUTPUT/1927_Sunday_June_05_2022_02_39_47_AM_51050310/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = e^{x^2}$$

1.4.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int e^{x^2} dx \\ &= \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2} + c_1 \tag{1}$$




Figure 4: Slope field plot

Verification of solutions

$$y = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2} + c_1$$

Verified OK.

1.4.2 Maple step by step solution

Let's solve

$$y' = e^{x^2}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int e^{x^2} dx + c_1$$

- Evaluate integral

$$y = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2} + c_1$$

- Solve for y

$$y = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)=exp(x^2),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{\pi} \operatorname{erfi}(x)}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 19

```
DSolve[y'[x]==Exp[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}\sqrt{\pi}\operatorname{erfi}(x) + c_1$$

1.5 problem 2(e)

1.5.1 Solving as quadrature ode	15
1.5.2 Maple step by step solution	16

Internal problem ID [2436]

Internal file name [OUTPUT/1928_Sunday_June_05_2022_02_39_49_AM_13964111/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = x e^{x^2}$$

1.5.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int x e^{x^2} dx \\ &= \frac{e^{x^2}}{2} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{x^2}}{2} + c_1 \tag{1}$$




Figure 5: Slope field plot

Verification of solutions

$$y = \frac{e^{x^2}}{2} + c_1$$

Verified OK.

1.5.2 Maple step by step solution

Let's solve

$$y' = x e^{x^2}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int x e^{x^2} dx + c_1$$

- Evaluate integral

$$y = \frac{e^{x^2}}{2} + c_1$$

- Solve for y

$$y = \frac{e^{x^2}}{2} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x*exp(x^2),y(x), singsol=all)
```

$$y(x) = \frac{e^{x^2}}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 17

```
DSolve[y'[x]==x*Exp[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{x^2}}{2} + c_1$$

1.6 problem 2(a)

1.6.1 Solving as quadrature ode	18
1.6.2 Maple step by step solution	19

Internal problem ID [2437]

Internal file name [OUTPUT/1929_Sunday_June_05_2022_02_39_50_AM_71357439/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = \arcsin(x)$$

1.6.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \arcsin(x) \, dx \\ &= x \arcsin(x) + \sqrt{-x^2 + 1} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1 \tag{1}$$




Figure 6: Slope field plot

Verification of solutions

$$y = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

Verified OK.

1.6.2 Maple step by step solution

Let's solve

$$y' = \arcsin(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int \arcsin(x) dx + c_1$$

- Evaluate integral

$$y = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

- Solve for y

$$y = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=arcsin(x),y(x), singsol=all)
```

$$y(x) = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 23

```
DSolve[y'[x]==ArcSin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \arcsin(x) + \sqrt{1 - x^2} + c_1$$

1.7 problem 3(a)

1.7.1	Solving as separable ode	21
1.7.2	Solving as linear ode	23
1.7.3	Solving as homogeneousTypeD2 ode	24
1.7.4	Solving as first order ode lie symmetry lookup ode	26
1.7.5	Solving as exact ode	30
1.7.6	Maple step by step solution	34

Internal problem ID [2438]

Internal file name [OUTPUT/1930_Sunday_June_05_2022_02_39_52_AM_98288231/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$y' - yx = 0$$

1.7.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= xy\end{aligned}$$

Where $f(x) = x$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= x dx \\ \int \frac{1}{y} dy &= \int x dx \\ \ln(y) &= \frac{x^2}{2} + c_1 \\ y &= e^{\frac{x^2}{2} + c_1} \\ &= c_1 e^{\frac{x^2}{2}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x^2}{2}} \tag{1}$$




Figure 7: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x^2}{2}}$$

Verified OK.

1.7.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -x$$

$$q(x) = 0$$

Hence the ode is

$$y' - yx = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -x dx} \\ &= e^{-\frac{x^2}{2}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu y &= 0 \\ \frac{d}{dx}\left(e^{-\frac{x^2}{2}}y\right) &= 0\end{aligned}$$

Integrating gives

$$e^{-\frac{x^2}{2}}y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{-\frac{x^2}{2}}$ results in

$$y = c_1 e^{\frac{x^2}{2}}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x^2}{2}} \tag{1}$$




Figure 8: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x^2}{2}}$$

Verified OK.

1.7.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - u(x)x^2 = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(x^2 - 1)}{x} \end{aligned}$$

Where $f(x) = \frac{x^2-1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{x^2-1}{x} dx \\ \int \frac{1}{u} du &= \int \frac{x^2-1}{x} dx \\ \ln(u) &= \frac{x^2}{2} - \ln(x) + c_2 \\ u &= e^{\frac{x^2}{2} - \ln(x) + c_2} \\ &= c_2 e^{\frac{x^2}{2} - \ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_2 e^{\frac{x^2}{2}}}{x}$$

Therefore the solution y is

$$\begin{aligned}y &= xu \\ &= c_2 e^{\frac{x^2}{2}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 e^{\frac{x^2}{2}} \tag{1}$$




Figure 9: Slope field plot

Verification of solutions

$$y = c_2 e^{\frac{x^2}{2}}$$

Verified OK.

1.7.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} y' &= xy \\ y' &= \omega(x, y) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 7: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{\frac{x^2}{2}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{\frac{x^2}{2}}} dy \end{aligned}$$

Which results in

$$S = e^{-\frac{x^2}{2}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = xy$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -x e^{-\frac{x^2}{2}} y \\ S_y &= e^{-\frac{x^2}{2}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-\frac{x^2}{2}} y = c_1$$



Which simplifies to

$$e^{-\frac{x^2}{2}} y = c_1$$

Which gives

$$y = c_1 e^{\frac{x^2}{2}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = xy$ 	$R = x$ $S = e^{-\frac{x^2}{2}} y$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x^2}{2}} \quad (1)$$




Figure 10: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x^2}{2}}$$

Verified OK.

1.7.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{y}\right) dy &= (x) dx \\ (-x) dx + \left(\frac{1}{y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x \\ N(x, y) &= \frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -x dx \\ \phi &= -\frac{x^2}{2} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y} \right) dy \\ f(y) &= \ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} + \ln(y)$$

The solution becomes

$$y = e^{\frac{x^2}{2} + c_1}$$

Summary

The solution(s) found are the following

$$y = e^{\frac{x^2}{2} + c_1} \tag{1}$$




Figure 11: Slope field plot

Verification of solutions

$$y = e^{\frac{x^2}{2} + c_1}$$

Verified OK.

1.7.6 Maple step by step solution

Let's solve

$$y' - yx = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int x dx + c_1$$

- Evaluate integral

$$\ln(y) = \frac{x^2}{2} + c_1$$

- Solve for y

$$y = e^{\frac{x^2}{2} + c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=y(x)*x,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^2}{2}} c_1$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 22

```
DSolve[y'[x]==y[x]*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow 0$$

1.8 problem 3(b)

1.8.1	Solving as separable ode	36
1.8.2	Solving as first order ode lie symmetry lookup ode	38
1.8.3	Solving as exact ode	42
1.8.4	Solving as riccati ode	46
1.8.5	Maple step by step solution	48

Internal problem ID [2439]

Internal file name [OUTPUT/1931_Sunday_June_05_2022_02_39_53_AM_57998689/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "riccati", "separable",
"first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - y^2x^2 = 0$$

1.8.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= y^2x^2\end{aligned}$$

Where $f(x) = x^2$ and $g(y) = y^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y^2} dy &= x^2 dx \\ \int \frac{1}{y^2} dy &= \int x^2 dx \\ -\frac{1}{y} &= \frac{x^3}{3} + c_1\end{aligned}$$

Which results in

$$y = -\frac{3}{x^3 + 3c_1}$$

Summary

The solution(s) found are the following

$$y = -\frac{3}{x^3 + 3c_1} \tag{1}$$




Figure 12: Slope field plot

Verification of solutions

$$y = -\frac{3}{x^3 + 3c_1}$$

Verified OK.

1.8.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = y^2 x^2$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 10: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{1}{x^2} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{1}{x^2}} dx\end{aligned}$$

Which results in

$$S = \frac{x^3}{3}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y^2 x^2$$

Evaluating all the partial derivatives gives

$$R_x = 0$$

$$R_y = 1$$

$$S_x = x^2$$

$$S_y = 0$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{y^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{R^2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{1}{R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x^3}{3} = -\frac{1}{y} + c_1$$



Which simplifies to

$$\frac{x^3}{3} = -\frac{1}{y} + c_1$$

Which gives

$$y = \frac{3}{-x^3 + 3c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = y^2 x^2$ 	$R = y$ $S = \frac{x^3}{3}$	$\frac{dS}{dR} = \frac{1}{R^2}$ 

Summary

The solution(s) found are the following

$$y = \frac{3}{-x^3 + 3c_1} \tag{1}$$




Figure 13: Slope field plot

Verification of solutions

$$y = \frac{3}{-x^3 + 3c_1}$$

Verified OK.

1.8.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{y^2}\right) dy &= (x^2) dx \\ (-x^2) dx + \left(\frac{1}{y^2}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x^2 \\ N(x, y) &= \frac{1}{y^2}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y^2} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -x^2 dx \\ \phi &= -\frac{x^3}{3} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y^2}$. Therefore equation (4) becomes

$$\frac{1}{y^2} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y^2}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{y^2} \right) dy$$
$$f(y) = -\frac{1}{y} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^3}{3} - \frac{1}{y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^3}{3} - \frac{1}{y}$$

The solution becomes

$$y = -\frac{3}{x^3 + 3c_1}$$

Summary

The solution(s) found are the following

$$y = -\frac{3}{x^3 + 3c_1} \tag{1}$$




Figure 14: Slope field plot

Verification of solutions

$$y = -\frac{3}{x^3 + 3c_1}$$

Verified OK.

1.8.4 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= y^2 x^2 \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = y^2 x^2$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = 0$, $f_1(x) = 0$ and $f_2(x) = x^2$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{x^2 u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= 2x \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= 0\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$x^2 u''(x) - 2x u'(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_2 x^3 + c_1$$

The above shows that

$$u'(x) = 3c_2 x^2$$

Using the above in (1) gives the solution

$$y = -\frac{3c_2}{c_2 x^3 + c_1}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = -\frac{3}{x^3 + c_3}$$

Summary

The solution(s) found are the following

$$y = -\frac{3}{x^3 + c_3} \quad (1)$$




Figure 15: Slope field plot

Verification of solutions

$$y = -\frac{3}{x^3 + c_3}$$

Verified OK.

1.8.5 Maple step by step solution

Let's solve

$$y' - y^2 x^2 = 0$$

- Highest derivative means the order of the ODE is 1
- y'
- Separate variables

$$\frac{y'}{y^2} = x^2$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y^2} dx = \int x^2 dx + c_1$$

- Evaluate integral

$$-\frac{1}{y} = \frac{x^3}{3} + c_1$$

- Solve for y

$$y = -\frac{3}{x^3 + 3c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=y(x)^2*x^2,y(x), singsol=all)
```

$$y(x) = -\frac{3}{x^3 - 3c_1}$$

✓ Solution by Mathematica

Time used: 0.107 (sec). Leaf size: 22

```
DSolve[y'[x]==y[x]^2*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{3}{x^3 + 3c_1}$$
$$y(x) \rightarrow 0$$

1.9 problem 3(c)

1.9.1	Solving as separable ode	50
1.9.2	Solving as first order special form ID 1 ode	52
1.9.3	Solving as first order ode lie symmetry lookup ode	53
1.9.4	Solving as exact ode	57
1.9.5	Maple step by step solution	61

Internal problem ID [2440]

Internal file name [OUTPUT/1932_Sunday_June_05_2022_02_39_55_AM_49856130/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "separable", "first order special form ID 1", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_separable]`

$$y' + x e^y = 0$$

1.9.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -x e^y\end{aligned}$$

Where $f(x) = -x$ and $g(y) = e^y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{e^y} dy &= -x dx \\ \int \frac{1}{e^y} dy &= \int -x dx \\ -e^{-y} &= -\frac{x^2}{2} + c_1\end{aligned}$$

Which results in

$$y = \ln \left(-\frac{2}{-x^2 + 2c_1} \right)$$

Summary

The solution(s) found are the following

$$y = \ln \left(-\frac{2}{-x^2 + 2c_1} \right) \tag{1}$$




Figure 16: Slope field plot

Verification of solutions

$$y = \ln \left(-\frac{2}{-x^2 + 2c_1} \right)$$

Verified OK.

1.9.2 Solving as first order special form ID 1 ode

Writing the ode as

$$y' = -x e^y \quad (1)$$

And using the substitution $u = e^{-y}$ then

$$u' = -y' e^{-y}$$

The above shows that

$$\begin{aligned} y' &= -u'(x) e^y \\ &= -\frac{u'(x)}{u} \end{aligned}$$

Substituting this in (1) gives

$$-\frac{u'(x)}{u} = -\frac{x}{u}$$

The above simplifies to

$$u'(x) = x \quad (2)$$

Now ode (2) is solved for $u(x)$ Integrating both sides gives

$$\begin{aligned} u(x) &= \int x \, dx \\ &= \frac{x^2}{2} + c_1 \end{aligned}$$

Substituting the solution found for $u(x)$ in $u = e^{-y}$ gives

$$\begin{aligned} y &= -\ln(u(x)) \\ &= -\ln\left(\frac{x^2}{2} + c_1\right) \\ &= \ln(2) - \ln(x^2 + 2c_1) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(2) - \ln(x^2 + 2c_1) \quad (1)$$




Figure 17: Slope field plot

Verification of solutions

$$y = \ln(2) - \ln(x^2 + 2c_1)$$

Verified OK.

1.9.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -x e^y$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 13: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= -\frac{1}{x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{-\frac{1}{x}} dx \end{aligned}$$

Which results in

$$S = -\frac{x^2}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -x e^y$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 0 \\ R_y &= 1 \\ S_x &= -x \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^{-y} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^{-R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -e^{-R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{x^2}{2} = -e^{-y} + c_1$$



Which simplifies to

$$-\frac{x^2}{2} = -e^{-y} + c_1$$

Which gives

$$y = -\ln\left(\frac{x^2}{2} + c_1\right)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -x e^y$ 	$R = y$ $S = -\frac{x^2}{2}$	$\frac{dS}{dR} = e^{-R}$ 

Summary

The solution(s) found are the following

$$y = -\ln\left(\frac{x^2}{2} + c_1\right) \quad (1)$$




Figure 18: Slope field plot

Verification of solutions

$$y = -\ln\left(\frac{x^2}{2} + c_1\right)$$

Verified OK.

1.9.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(-e^{-y}) dy &= (x) dx \\ (-x) dx + (-e^{-y}) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x \\ N(x, y) &= -e^{-y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-e^{-y}) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x dx$$

$$\phi = -\frac{x^2}{2} + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -e^{-y}$. Therefore equation (4) becomes

$$-e^{-y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -e^{-y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (-e^{-y}) dy$$

$$f(y) = e^{-y} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} + e^{-y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} + e^{-y}$$

The solution becomes

$$y = -\ln\left(\frac{x^2}{2} + c_1\right)$$

Summary

The solution(s) found are the following

$$y = -\ln\left(\frac{x^2}{2} + c_1\right) \tag{1}$$




Figure 19: Slope field plot

Verification of solutions

$$y = -\ln\left(\frac{x^2}{2} + c_1\right)$$

Verified OK.

1.9.5 Maple step by step solution

Let's solve

$$y' + x e^y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{e^y} = -x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{e^y} dx = \int -x dx + c_1$$

- Evaluate integral

$$-\frac{1}{e^y} = -\frac{x^2}{2} + c_1$$

- Solve for y

$$y = \ln\left(-\frac{2}{-x^2+2c_1}\right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=-x*exp(y(x)),y(x), singsol=all)
```

$$y(x) = \ln(2) + \ln\left(\frac{1}{x^2 + 2c_1}\right)$$

✓ Solution by Mathematica

Time used: 0.307 (sec). Leaf size: 19

```
DSolve[y'[x]==-x*Exp[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(2) - \log(x^2 - 2c_1)$$

1.10 problem 3(d)

1.10.1 Solving as separable ode	63
1.10.2 Solving as differentialType ode	65
1.10.3 Solving as first order ode lie symmetry lookup ode	66
1.10.4 Solving as exact ode	70
1.10.5 Maple step by step solution	74

Internal problem ID [2441]

Internal file name [OUTPUT/1933_Sunday_June_05_2022_02_39_57_AM_21333843/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "differential-Type", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' \sin(y) = x^2$$

1.10.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{x^2}{\sin(y)}\end{aligned}$$

Where $f(x) = x^2$ and $g(y) = \frac{1}{\sin(y)}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\sin(y)} dy &= x^2 dx \\ \int \frac{1}{\sin(y)} dy &= \int x^2 dx\end{aligned}$$

$$-\cos(y) = \frac{x^3}{3} + c_1$$

Which results in

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right)$$

Summary

The solution(s) found are the following

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right) \tag{1}$$




Figure 20: Slope field plot

Verification of solutions

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right)$$

Verified OK.

1.10.2 Solving as differential Type ode

Writing the ode as

$$y' = \frac{x^2}{\sin(y)} \quad (1)$$

Which becomes

$$(\sin(y)) dy = (x^2) dx \quad (2)$$

But the RHS is complete differential because

$$(x^2) dx = d\left(\frac{x^3}{3}\right)$$

Hence (2) becomes

$$(\sin(y)) dy = d\left(\frac{x^3}{3}\right)$$

Integrating both sides gives gives these solutions

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right) + c_1$$

Summary

The solution(s) found are the following

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right) + c_1 \quad (1)$$




Figure 21: Slope field plot

Verification of solutions

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right) + c_1$$

Verified OK.

1.10.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{x^2}{\sin(y)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 16: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{1}{x^2} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{1}{x^2}} dx \end{aligned}$$

Which results in

$$S = \frac{x^3}{3}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x^2}{\sin(y)}$$

Evaluating all the partial derivatives gives

$$R_x = 0$$

$$R_y = 1$$

$$S_x = x^2$$

$$S_y = 0$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \sin(y) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \sin(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\cos(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x^3}{3} = -\cos(y) + c_1$$



Which simplifies to

$$\frac{x^3}{3} = -\cos(y) + c_1$$

Which gives

$$y = \arccos\left(-\frac{x^3}{3} + c_1\right)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x^2}{\sin(y)}$ 	$R = y$ $S = \frac{x^3}{3}$	$\frac{dS}{dR} = \sin(R)$ 

Summary

The solution(s) found are the following

$$y = \arccos\left(-\frac{x^3}{3} + c_1\right) \quad (1)$$




Figure 22: Slope field plot

Verification of solutions

$$y = \arccos\left(-\frac{x^3}{3} + c_1\right)$$

Verified OK.

1.10.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the

ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (\sin(y)) dy &= (x^2) dx \\ (-x^2) dx + (\sin(y)) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x^2 \\ N(x, y) &= \sin(y) \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(\sin(y)) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x^2 dx$$

$$\phi = -\frac{x^3}{3} + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \sin(y)$. Therefore equation (4) becomes

$$\sin(y) = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \sin(y)$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (\sin(y)) dy$$

$$f(y) = -\cos(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^3}{3} - \cos(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^3}{3} - \cos(y)$$

Summary

The solution(s) found are the following

$$-\frac{x^3}{3} - \cos(y) = c_1 \tag{1}$$




Figure 23: Slope field plot

Verification of solutions

$$-\frac{x^3}{3} - \cos(y) = c_1$$

Verified OK.

1.10.5 Maple step by step solution

Let's solve

$$y' \sin(y) = x^2$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int y' \sin(y) dx = \int x^2 dx + c_1$$

- Evaluate integral

$$-\cos(y) = \frac{x^3}{3} + c_1$$

- Solve for y

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)*sin(y(x))=x^2,y(x), singsol=all)
```

$$y(x) = \frac{\pi}{2} + \arcsin\left(\frac{x^3}{3} + c_1\right)$$

✓ Solution by Mathematica

Time used: 0.509 (sec). Leaf size: 37

```
DSolve[y'[x]*Sin[y[x]]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos\left(-\frac{x^3}{3} - c_1\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{x^3}{3} - c_1\right)$$

1.11 problem 3(e)

1.11.1 Solving as separable ode	76
1.11.2 Solving as first order ode lie symmetry lookup ode	78
1.11.3 Solving as exact ode	82
1.11.4 Maple step by step solution	86

Internal problem ID [2442]

Internal file name [OUTPUT/1934_Sunday_June_05_2022_02_39_59_AM_77840185/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$xy' - \sqrt{1 - y^2} = 0$$

1.11.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\sqrt{-y^2 + 1}}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = \sqrt{-y^2 + 1}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\sqrt{-y^2 + 1}} dy &= \frac{1}{x} dx \\ \int \frac{1}{\sqrt{-y^2 + 1}} dy &= \int \frac{1}{x} dx \\ \arcsin(y) &= \ln(x) + c_1\end{aligned}$$

Which results in

$$y = \sin(\ln(x) + c_1)$$

Summary

The solution(s) found are the following

$$y = \sin(\ln(x) + c_1) \tag{1}$$




Figure 24: Slope field plot

Verification of solutions

$$y = \sin(\ln(x) + c_1)$$

Verified OK.

1.11.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\sqrt{-y^2 + 1}}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 19: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= x \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{x} dx\end{aligned}$$

Which results in

$$S = \ln(x)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{\sqrt{-y^2 + 1}}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 0 \\R_y &= 1 \\S_x &= \frac{1}{x} \\S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{\sqrt{-y^2 + 1}} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{\sqrt{-R^2 + 1}}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \arcsin(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(x) = \arcsin(y) + c_1$$



Which simplifies to

$$\ln(x) = \arcsin(y) + c_1$$

Which gives

$$y = -\sin(-\ln(x) + c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{\sqrt{-y^2+1}}{x}$ 	$R = y$ $S = \ln(x)$	$\frac{dS}{dR} = \frac{1}{\sqrt{-R^2+1}}$ 

Summary

The solution(s) found are the following

$$y = -\sin(-\ln(x) + c_1) \tag{1}$$




Figure 25: Slope field plot

Verification of solutions

$$y = -\sin(-\ln(x) + c_1)$$

Verified OK.

1.11.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{1}{\sqrt{-y^2+1}}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{1}{\sqrt{-y^2+1}}\right) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x} \\ N(x, y) &= \frac{1}{\sqrt{-y^2+1}}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{\sqrt{-y^2+1}} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\sqrt{-y^2+1}}$. Therefore equation (4) becomes

$$\frac{1}{\sqrt{-y^2+1}} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{\sqrt{-y^2+1}}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{\sqrt{-y^2+1}} \right) dy \\ f(y) &= \arcsin(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \arcsin(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \arcsin(y)$$

The solution becomes

$$y = \sin(\ln(x) + c_1)$$

Summary

The solution(s) found are the following

$$y = \sin(\ln(x) + c_1) \tag{1}$$




Figure 26: Slope field plot

Verification of solutions

$$y = \sin(\ln(x) + c_1)$$

Verified OK.

1.11.4 Maple step by step solution

Let's solve

$$xy' - \sqrt{1-y^2} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{\sqrt{1-y^2}} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{1-y^2}} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$\arcsin(y) = \ln(x) + c_1$$

- Solve for y

$$y = \sin(\ln(x) + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)*x=sqrt(1-y(x)^2),y(x), singsol=all)
```

$$y(x) = \sin(\ln(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.217 (sec). Leaf size: 29

```
DSolve[y'[x]*x==Sqrt[1-y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(\log(x) + c_1)$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow \text{Interval}[\{-1, 1\}]$$

1.12 problem 3(f)

1.12.1 Maple step by step solution 89

Internal problem ID [2443]

Internal file name [OUTPUT/1935_Sunday_June_05_2022_02_40_01_AM_72065445/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(f).

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[`_quadrature`]

$$y'^2 - y^2 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = -y \tag{1}$$

$$y' = y \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int -\frac{1}{y} dy = \int dx$$
$$-\ln(y) = x + c_1$$

Raising both side to exponential gives

$$\frac{1}{y} = e^{x+c_1}$$

Which simplifies to

$$\frac{1}{y} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-x}}{c_2} \quad (1)$$

Verification of solutions

$$y = \frac{e^{-x}}{c_2}$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\int \frac{1}{y} dy = x + c_3$$

$$\ln(y) = x + c_3$$

$$y = e^{x+c_3}$$

$$y = c_3 e^x$$

Summary

The solution(s) found are the following

$$y = c_3 e^x \quad (1)$$

Verification of solutions

$$y = c_3 e^x$$

Verified OK.

1.12.1 Maple step by step solution

Let's solve

$$y'^2 - y^2 = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\ln(y) = x + c_1$$

- Solve for y

$$y = e^{x+c_1}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful
Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((diff(y(x),x))^2-y(x)^2=0,y(x), singsol=all)
```

$$y(x) = e^x c_1$$

$$y(x) = e^{-x} c_1$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 28

```
DSolve[(y'[x])^2-y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x}$$

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow 0$$

1.13 problem 3(g)

1.13.1 Maple step by step solution 92

Internal problem ID [2444]

Internal file name [OUTPUT/1936_Sunday_June_05_2022_02_40_04_AM_47833826/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(g).

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y'^2 - 3y' = -2$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = 2 \tag{1}$$

$$y' = 1 \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} y &= \int 2 \, dx \\ &= 2x + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = 2x + c_1 \tag{1}$$

Verification of solutions

$$y = 2x + c_1$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\begin{aligned}y &= \int 1 \, dx \\ &= x + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = x + c_2 \tag{1}$$

Verification of solutions

$$y = x + c_2$$

Verified OK.

1.13.1 Maple step by step solution

Let's solve

$$y'^2 - 3y' = -2$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int (y'^2 - 3y') \, dx = \int (-2) \, dx + c_1$$

- Cannot compute integral

$$\int (y'^2 - 3y') \, dx = -2x + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful  
Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)^2-3*diff(y(x),x)+2=0,y(x), singsol=all)
```

$$y(x) = 2x + c_1$$

$$y(x) = c_1 + x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 19

```
DSolve[(y'[x])^2-3*y'[x]+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + c_1$$

$$y(x) \rightarrow 2x + c_1$$

1.14 problem 3(h)

1.14.1 Solving as quadrature ode	94
1.14.2 Maple step by step solution	95

Internal problem ID [2445]

Internal file name [OUTPUT/1937_Sunday_June_05_2022_02_40_05_AM_55553494/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$(x^2 + 1) y' = 1$$

1.14.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{1}{x^2 + 1} dx \\ &= \arctan(x) + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \arctan(x) + c_1 \tag{1}$$




Figure 27: Slope field plot

Verification of solutions

$$y = \arctan(x) + c_1$$

Verified OK.

1.14.2 Maple step by step solution

Let's solve

$$(x^2 + 1)y' = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{1}{x^2+1}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{x^2+1} dx + c_1$$

- Evaluate integral

- $y = \arctan(x) + c_1$
Solve for y
 $y = \arctan(x) + c_1$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve((1+x^2)*diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = \arctan(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 10

```
DSolve[(1+x^2)*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arctan(x) + c_1$$

1.15 problem 3(i)

1.15.1 Solving as quadrature ode	97
1.15.2 Maple step by step solution	98

Internal problem ID [2446]

Internal file name [OUTPUT/1938_Sunday_June_05_2022_02_40_07_AM_86851588/index.tex]

Book: Elementary Differential Equations, Martin, Reissner, 2nd ed, 1961

Section: Exercis 2, page 5

Problem number: 3(i).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' \sin(x) = 1$$

1.15.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{1}{\sin(x)} dx \\ &= \ln(\csc(x) - \cot(x)) + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(\csc(x) - \cot(x)) + c_1 \tag{1}$$




Figure 28: Slope field plot

Verification of solutions

$$y = \ln(\csc(x) - \cot(x)) + c_1$$

Verified OK.

1.15.2 Maple step by step solution

Let's solve

$$y' \sin(x) = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{1}{\sin(x)}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{\sin(x)} dx + c_1$$

- Evaluate integral

$$y = \ln(\csc(x) - \cot(x)) + c_1$$

- Solve for y

$$y = \ln(\csc(x) - \cot(x)) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)*sin(x)=1,y(x), singsol=all)
```

$$y(x) = -\ln(\csc(x) + \cot(x)) + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 13

```
DSolve[y'[x]*Sin[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\operatorname{arctanh}(\cos(x)) + c_1$$