

A Solution Manual For

**Own collection of miscellaneous
problems**

Nasser M. Abbasi

May 16, 2024

Contents

1	section 1.0	2
2	section 2.0	735
3	section 3.0	1143
4	section 4.0	1472
5	section 5.0	2341

1 section 1.0

1.1	problem 1	5
1.2	problem 2	14
1.3	problem 3	25
1.4	problem 4	34
1.5	problem 5	44
1.6	problem 6	47
1.7	problem 7	50
1.8	problem 8	53
1.9	problem 9	56
1.10	problem 10	59
1.11	problem 11	62
1.12	problem 12	73
1.13	problem 13	86
1.14	problem 14	101
1.15	problem 15	110
1.16	problem 16	113
1.17	problem 17	126
1.18	problem 18	130
1.19	problem 19	135
1.20	problem 20	137
1.21	problem 21	147
1.22	problem 23	155
1.23	problem 24	169
1.24	problem 25	173
1.25	problem 26	176
1.26	problem 27	179
1.27	problem 28	182
1.28	problem 29	185
1.29	problem 30	188
1.30	problem 31	193
1.31	problem 32	205
1.32	problem 33	216
1.33	problem 34	221
1.34	problem 35	225
1.35	problem 36	239
1.36	problem 37	243
1.37	problem 38	247

1.38	problem 39	252
1.39	problem 40	256
1.40	problem 41	265
1.41	problem 41	274
1.42	problem 42	285
1.43	problem 43	297
1.44	problem 44	309
1.45	problem 45	315
1.46	problem 46	324
1.47	problem 47	327
1.48	problem 48	349
1.49	problem 49	366
1.50	problem 50	383
1.51	problem 51	397
1.52	problem 52	408
1.53	problem 53	410
1.54	problem 54	426
1.55	problem 55	440
1.56	problem 56	458
1.57	problem 57	473
1.58	problem 58	492
1.59	problem 59	500
1.60	problem 60	508
1.61	problem 61	524
1.62	problem 62	528
1.63	problem 63	532
1.64	problem 64	535
1.65	problem 65	538
1.66	problem 66	542
1.67	problem 67	545
1.68	problem 68	550
1.69	problem 69	555
1.70	problem 70	560
1.71	problem 71	564
1.72	problem 72	576
1.73	problem 73	586
1.74	problem 74	596
1.75	problem 75	606
1.76	problem 76	616

1.77	problem 77	625
1.78	problem 78	628
1.79	problem 78	633
1.80	problem 79	638
1.81	problem 80	651
1.82	problem 81	655
1.83	problem 82	659
1.84	problem 83	669
1.85	problem 84	672
1.86	problem 85	677
1.87	problem 86	687
1.88	problem 87	690
1.89	problem 88	699
1.90	problem 88	708
1.91	problem 89	719
1.92	problem 90	729

1.1 problem 1

1.1.1	Solving as separable ode	5
1.1.2	Solving as first order ode lie symmetry lookup ode	7
1.1.3	Solving as exact ode	8
1.1.4	Maple step by step solution	12

Internal problem ID [7045]

Internal file name [OUTPUT/6031_Sunday_June_05_2022_04_14_34_PM_36171372/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 1.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - \frac{\cos(y) \sec(x)}{x} = 0$$

1.1.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\cos(y) \sec(x)}{x}\end{aligned}$$

Where $f(x) = \frac{\sec(x)}{x}$ and $g(y) = \cos(y)$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\cos(y)} dy &= \frac{\sec(x)}{x} dx \\ \int \frac{1}{\cos(y)} dy &= \int \frac{\sec(x)}{x} dx\end{aligned}$$

$$\ln(\sec(y) + \tan(y)) = \int \frac{\sec(x)}{x} dx + c_1$$

Raising both side to exponential gives

$$\sec(y) + \tan(y) = e^{\int \frac{\sec(x)}{x} dx + c_1}$$

Which simplifies to

$$\sec(y) + \tan(y) = c_2 e^{\int \frac{\sec(x)}{x} dx}$$

Summary

The solution(s) found are the following

$$y = \arctan \left(\frac{e^{\int \frac{2\sec(x)}{x} dx + 2c_1} c_2^2 - 1}{e^{\int \frac{2\sec(x)}{x} dx + 2c_1} c_2^2 + 1}, \frac{2c_2 e^{\int \frac{\sec(x)}{x} dx + c_1}}{e^{\int \frac{2\sec(x)}{x} dx + 2c_1} c_2^2 + 1} \right) \quad (1)$$




Figure 1: Slope field plot

Verification of solutions

$$y = \arctan \left(\frac{e^{\int \frac{2\sec(x)}{x} dx + 2c_1} c_2^2 - 1}{e^{\int \frac{2\sec(x)}{x} dx + 2c_1} c_2^2 + 1}, \frac{2c_2 e^{\int \frac{\sec(x)}{x} dx + c_1}}{e^{\int \frac{2\sec(x)}{x} dx + 2c_1} c_2^2 + 1} \right)$$

Verified OK.

1.1.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\cos(y) \sec(x)}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 1: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{x}{\sec(x)} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{x}{\sec(x)}} dx\end{aligned}$$

Which results in

$$S = \int \frac{\sec(x)}{x} dx$$

1.1.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0\tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{1}{\cos(y)} \right) dy &= \left(\frac{\sec(x)}{x} \right) dx \\ \left(-\frac{\sec(x)}{x} \right) dx + \left(\frac{1}{\cos(y)} \right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{\sec(x)}{x} \\ N(x, y) &= \frac{1}{\cos(y)} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{\sec(x)}{x} \right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{\cos(y)} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{\sec(x)}{x} dx \\ \phi &= \int^x -\frac{\sec(-a)}{-a} d_{-a} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\cos(y)}$. Therefore equation (4) becomes

$$\frac{1}{\cos(y)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= \frac{1}{\cos(y)} \\ &= \sec(y)\end{aligned}$$

Integrating the above w.r.t y results in

$$\int f'(y) dy = \int (\sec(y)) dy$$

$$f(y) = \ln(\sec(y) + \tan(y)) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x -\frac{\sec(a)}{a} da + \ln(\sec(y) + \tan(y)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x -\frac{\sec(a)}{a} da + \ln(\sec(y) + \tan(y))$$

Summary

The solution(s) found are the following

$$\int^x -\frac{\sec(a)}{a} da + \ln(\sec(y) + \tan(y)) = c_1 \quad (1)$$




Figure 2: Slope field plot

Verification of solutions

$$\int^x -\frac{\sec(a)}{a} da + \ln(\sec(y) + \tan(y)) = c_1$$

Verified OK.

1.1.4 Maple step by step solution

Let's solve

$$y' - \frac{\cos(y)\sec(x)}{x} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{\cos(y)} = \frac{\sec(x)}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\cos(y)} dx = \int \frac{\sec(x)}{x} dx + c_1$$

- Evaluate integral

$$\ln(\sec(y) + \tan(y)) = \int \frac{\sec(x)}{x} dx + c_1$$

- Solve for y

$$y = \arctan \left(\frac{\left(e^{\int \frac{\sec(x)}{x} dx + c_1} \right)^2 - 1}{\left(e^{\int \frac{\sec(x)}{x} dx + c_1} \right)^2 + 1}, \frac{2 e^{\int \frac{\sec(x)}{x} dx + c_1}}{\left(e^{\int \frac{\sec(x)}{x} dx + c_1} \right)^2 + 1} \right)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 73

```
dsolve(diff(y(x),x) = cos(y(x))*sec(x)/x,y(x), singsol=all)
```

$$y(x) = \arctan \left(\frac{e^{2\left(\int \frac{\sec(x)}{x} dx\right)} c_1^2 - 1}{e^{2\left(\int \frac{\sec(x)}{x} dx\right)} c_1^2 + 1}, \frac{2 e^{\int \frac{\sec(x)}{x} dx} c_1}{e^{2\left(\int \frac{\sec(x)}{x} dx\right)} c_1^2 + 1} \right)$$

✓ Solution by Mathematica

Time used: 5.307 (sec). Leaf size: 49

```
DSolve[y'[x]== Cos[y[x]]*Sec[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \arctan \left(\tanh \left(\frac{1}{2} \left(\int_1^x \frac{\sec(K[1])}{K[1]} dK[1] + c_1 \right) \right) \right)$$

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

1.2 problem 2

1.2.1	Solving as separable ode	14
1.2.2	Solving as first order ode lie symmetry lookup ode	16
1.2.3	Solving as exact ode	19
1.2.4	Maple step by step solution	23

Internal problem ID [7046]

Internal file name [OUTPUT/6032_Sunday_June_05_2022_04_14_37_PM_48637153/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 2.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - x(\cos(y) + y) = 0$$

1.2.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= x(\cos(y) + y)\end{aligned}$$

Where $f(x) = x$ and $g(y) = \cos(y) + y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\cos(y) + y} dy &= x dx \\ \int \frac{1}{\cos(y) + y} dy &= \int x dx \\ \int^y \frac{1}{\cos(_a) + _a} d_a &= \frac{x^2}{2} + c_1\end{aligned}$$

Which results in

$$\int^y \frac{1}{\cos(_a) + _a} d_a = \frac{x^2}{2} + c_1$$

The solution is

$$\int^y \frac{1}{\cos(_a) + _a} d_a - \frac{x^2}{2} - c_1 = 0$$

Summary

The solution(s) found are the following

$$\int^y \frac{1}{\cos(_a) + _a} d_a - \frac{x^2}{2} - c_1 = 0 \tag{1}$$




Figure 3: Slope field plot

Verification of solutions

$$\int^y \frac{1}{\cos(_a) + _a} d_a - \frac{x^2}{2} - c_1 = 0$$

Verified OK.

1.2.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = x(\cos(y) + y)$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 4: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{1}{x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{1}{x}} dx\end{aligned}$$

Which results in

$$S = \frac{x^2}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = x(\cos(y) + y)$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 0 \\R_y &= 1 \\S_x &= x \\S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{\cos(y) + y} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{\cos(R) + R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \int \frac{1}{\cos(R) + R} dR + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x^2}{2} = \int^y \frac{1}{\cos(_a) + _a} d_a + c_1$$

Which simplifies to

$$\frac{x^2}{2} = \int^y \frac{1}{\cos(_a) + _a} d_a + c_1$$

This results in

$$\frac{x^2}{2} = \int^y \frac{1}{\cos(_a) + _a} d_a + c_1$$

Summary

The solution(s) found are the following

$$\frac{x^2}{2} = \int^y \frac{1}{\cos(_a) + _a} d_a + c_1 \quad (1)$$




Figure 4: Slope field plot

Verification of solutions

$$\frac{x^2}{2} = \int \frac{1}{\cos(a) + a} da + c_1$$

Verified OK.

1.2.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{\cos(y) + y}\right) dy &= (x) dx \\ (-x) dx + \left(\frac{1}{\cos(y) + y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x \\ N(x, y) &= \frac{1}{\cos(y) + y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{\cos(y) + y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -x dx \\ \phi &= -\frac{x^2}{2} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\cos(y) + y}$. Therefore equation (4) becomes

$$\frac{1}{\cos(y) + y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{\cos(y) + y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{\cos(y) + y} \right) dy$$

$$f(y) = \int_0^y \frac{1}{\cos(a) + a} da + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} + \int_0^y \frac{1}{\cos(a) + a} da + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} + \int_0^y \frac{1}{\cos(a) + a} da$$

Summary

The solution(s) found are the following

$$-\frac{x^2}{2} + \int_0^y \frac{1}{\cos(a) + a} da = c_1 \tag{1}$$




Figure 5: Slope field plot

Verification of solutions

$$-\frac{x^2}{2} + \int_0^y \frac{1}{\cos(a) + a} da = c_1$$

Verified OK.

1.2.4 Maple step by step solution

Let's solve

$$y' - x(\cos(y) + y) = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{\cos(y)+y} = x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\cos(y)+y} dx = \int x dx + c_1$$

- Cannot compute integral

$$\int \frac{y'}{\cos(y)+y} dx = \frac{x^2}{2} + c_1$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x) = x*(cos(y(x))+y(x)),y(x), singsol=all)
```

$$\frac{x^2}{2} - \left(\int^{y(x)} \frac{1}{\cos(a) + a} da \right) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.71 (sec). Leaf size: 33

```
DSolve[y'[x] == x*(Cos[y[x]]+y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{1}{\cos(K[1]) + K[1]} dK[1] \& \right] \left[\frac{x^2}{2} + c_1 \right]$$

1.3 problem 3

1.3.1	Solving as separable ode	25
1.3.2	Solving as first order ode lie symmetry lookup ode	27
1.3.3	Solving as exact ode	28
1.3.4	Maple step by step solution	32

Internal problem ID [7047]

Internal file name [OUTPUT/6033_Sunday_June_05_2022_04_14_40_PM_97928902/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 3.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - \frac{\sec(x)(\sin(y) + y)}{x} = 0$$

1.3.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\sec(x)(\sin(y) + y)}{x}\end{aligned}$$

Where $f(x) = \frac{\sec(x)}{x}$ and $g(y) = \sin(y) + y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\sin(y) + y} dy &= \frac{\sec(x)}{x} dx \\ \int \frac{1}{\sin(y) + y} dy &= \int \frac{\sec(x)}{x} dx\end{aligned}$$

$$\int^y \frac{1}{\sin(_a) + _a} d_a = \int \frac{\sec(x)}{x} dx + c_1$$

Which results in

$$\int^y \frac{1}{\sin(_a) + _a} d_a = \int \frac{\sec(x)}{x} dx + c_1$$

The solution is

$$\int^y \frac{1}{\sin(_a) + _a} d_a - \left(\int \frac{\sec(x)}{x} dx \right) - c_1 = 0$$

Summary

The solution(s) found are the following

$$\int^y \frac{1}{\sin(_a) + _a} d_a - \left(\int \frac{\sec(x)}{x} dx \right) - c_1 = 0 \quad (1)$$




Figure 6: Slope field plot

Verification of solutions

$$\int^y \frac{1}{\sin(_a) + _a} d_a - \left(\int \frac{\sec(x)}{x} dx \right) - c_1 = 0$$

Verified OK.

1.3.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\sec(x) (\sin(y) + y)}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 7: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{x}{\sec(x)} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{x}{\sec(x)}} dx\end{aligned}$$

Which results in

$$S = \int \frac{\sec(x)}{x} dx$$

1.3.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0\tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{1}{\sin(y) + y} \right) dy &= \left(\frac{\sec(x)}{x} \right) dx \\ \left(-\frac{\sec(x)}{x} \right) dx &+ \left(\frac{1}{\sin(y) + y} \right) dy = 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{\sec(x)}{x} \\ N(x, y) &= \frac{1}{\sin(y) + y} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{\sec(x)}{x} \right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{\sin(y) + y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{\sec(x)}{x} dx \\ \phi &= \int^x -\frac{\sec(x)}{x} dx + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\sin(y) + y}$. Therefore equation (4) becomes

$$\frac{1}{\sin(y) + y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{\sin(y) + y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{\sin(y) + y} \right) dy$$

$$f(y) = \int_0^y \frac{1}{\sin(_a) + _a} d_a + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x -\frac{\sec(_a)}{_a} d_a + \int_0^y \frac{1}{\sin(_a) + _a} d_a + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x -\frac{\sec(_a)}{_a} d_a + \int_0^y \frac{1}{\sin(_a) + _a} d_a$$

Summary

The solution(s) found are the following

$$\int^x -\frac{\sec(_a)}{_a} d_a + \int_0^y \frac{1}{\sin(_a) + _a} d_a = c_1 \quad (1)$$




Figure 7: Slope field plot

Verification of solutions

$$\int^x -\frac{\sec(_a)}{_a} d_a + \int_0^y \frac{1}{\sin(_a) + _a} d_a = c_1$$

Verified OK.

1.3.4 Maple step by step solution

Let's solve

$$y' - \frac{\sec(x)(\sin(y)+y)}{x} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{\sin(y)+y} = \frac{\sec(x)}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sin(y)+y} dx = \int \frac{\sec(x)}{x} dx + c_1$$

- Cannot compute integral

$$\int \frac{y'}{\sin(y)+y} dx = \int \frac{\sec(x)}{x} dx + c_1$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x) = sec(x)*(sin(y(x))+y(x))/x,y(x), singsol=all)
```

$$\int \frac{\sec(x)}{x} dx - \left(\int^{y(x)} \frac{1}{\sin(a)+a} da \right) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 1.312 (sec). Leaf size: 41

```
DSolve[y'[x]== Sec[x]*(Sin[y[x]]+y[x])/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{1}{K[1] + \sin(K[1])} dK[1] \& \right] \left[\int_1^x \frac{\sec(K[2])}{K[2]} dK[2] + c_1 \right]$$

1.4 problem 4

1.4.1	Solving as separable ode	34
1.4.2	Solving as first order ode lie symmetry lookup ode	36
1.4.3	Solving as exact ode	38
1.4.4	Maple step by step solution	42

Internal problem ID [7048]

Internal file name [OUTPUT/6034_Sunday_June_05_2022_04_14_43_PM_72189390/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - \left(5 + \frac{\sec(x)}{x}\right) (\sin(y) + y) = 0$$

1.4.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{(\sin(y) + y)(\sec(x) + 5x)}{x}\end{aligned}$$

Where $f(x) = \frac{\sec(x)+5x}{x}$ and $g(y) = \sin(y) + y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\sin(y) + y} dy &= \frac{\sec(x) + 5x}{x} dx \\ \int \frac{1}{\sin(y) + y} dy &= \int \frac{\sec(x) + 5x}{x} dx\end{aligned}$$

$$\int^y \frac{1}{\sin(_a) + _a} d_a = \int \frac{\sec(x) + 5x}{x} dx + c_1$$

Which results in

$$\int^y \frac{1}{\sin(_a) + _a} d_a = \int \frac{\sec(x) + 5x}{x} dx + c_1$$

The solution is

$$\int^y \frac{1}{\sin(_a) + _a} d_a - \left(\int \frac{\sec(x) + 5x}{x} dx \right) - c_1 = 0$$

Summary

The solution(s) found are the following

$$\int^y \frac{1}{\sin(_a) + _a} d_a - \left(\int \frac{\sec(x) + 5x}{x} dx \right) - c_1 = 0 \tag{1}$$




Figure 8: Slope field plot

Verification of solutions

$$\int^y \frac{1}{\sin(a) + a} da - \left(\int \frac{\sec(x) + 5x}{x} dx \right) - c_1 = 0$$

Verified OK.

1.4.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{(\sin(y) + y)(\sec(x) + 5x)}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 10: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{x}{\sec(x) + 5x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{x}{\sec(x)+5x}} dx \end{aligned}$$

Which results in

$$S = \int \frac{\sec(x) + 5x}{x} dx$$

1.4.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition

$\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\left(\frac{1}{\sin(y) + y} \right) dy = \left(\frac{\sec(x) + 5x}{x} \right) dx$$

$$\left(-\frac{\sec(x) + 5x}{x} \right) dx + \left(\frac{1}{\sin(y) + y} \right) dy = 0 \quad (2A)$$

Comparing (1A) and (2A) shows that

$$M(x, y) = -\frac{\sec(x) + 5x}{x}$$

$$N(x, y) = \frac{1}{\sin(y) + y}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y} \left(-\frac{\sec(x) + 5x}{x} \right)$$

$$= 0$$

And

$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x} \left(\frac{1}{\sin(y) + y} \right)$$

$$= 0$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{\sec(x) + 5x}{x} dx \\ \phi &= \int^x -\frac{\sec(_a) + 5_a}{_a} d_a + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\sin(y)+y}$. Therefore equation (4) becomes

$$\frac{1}{\sin(y) + y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{\sin(y) + y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{\sin(y) + y} \right) dy \\ f(y) &= \int_0^y \frac{1}{\sin(_a) + _a} d_a + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x -\frac{\sec(_a) + 5_a}{_a} d_a + \int_0^y \frac{1}{\sin(_a) + _a} d_a + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x -\frac{\sec(-a) + 5-a}{-a} d-a + \int_0^y \frac{1}{\sin(-a) + -a} d-a$$

Summary

The solution(s) found are the following

$$\int^x -\frac{\sec(-a) + 5-a}{-a} d-a + \int_0^y \frac{1}{\sin(-a) + -a} d-a = c_1 \quad (1)$$




Figure 9: Slope field plot

Verification of solutions

$$\int^x -\frac{\sec(-a) + 5-a}{-a} d-a + \int_0^y \frac{1}{\sin(-a) + -a} d-a = c_1$$

Verified OK.

1.4.4 Maple step by step solution

Let's solve

$$y' - \left(5 + \frac{\sec(x)}{x}\right) (\sin(y) + y) = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{\sin(y)+y} = 5 + \frac{\sec(x)}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sin(y)+y} dx = \int \left(5 + \frac{\sec(x)}{x}\right) dx + c_1$$

- Cannot compute integral

$$\int \frac{y'}{\sin(y)+y} dx = 5x + \int \frac{2e^{Ix}}{(e^{Ix})^2+1} dx + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x) = (5+sec(x)/x)*(sin(y(x))+y(x)),y(x), singsol=all)
```

$$\int \frac{5x + \sec(x)}{x} dx - \left(\int^{y(x)} \frac{1}{\sin(_a) + _a} d_a \right) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 19.938 (sec). Leaf size: 168

```
DSolve[y'[x] == (5+Sec[x]/x)*(Sin[y[x]]+y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} & \text{Solve} \left[\int_1^x \left(-\frac{2 \sec(K[1])}{K[1]} \right. \right. \\ & \left. \left. - \frac{5(-\sec(K[1]) \sin(K[1] - y(x)) + \sec(K[1]) \sin(K[1] + y(x)) + 2y(x))}{\sin(y(x)) + y(x)} \right) dK[1] \right. \\ & \left. + \int_1^{y(x)} \left(\frac{2}{K[2] + \sin(K[2])} \right. \right. \\ & \left. \left. - \int_1^x \left(\frac{5(\cos(K[2]) + 1)(2K[2] - \sec(K[1]) \sin(K[1] - K[2]) + \sec(K[1]) \sin(K[1] + K[2]))}{(K[2] + \sin(K[2]))^2} - \frac{5(\cos(K[1])}{\sin(y(x)) + y(x)} \right) dK[1] \right) dK[2] \right] \end{aligned}$$

1.5 problem 5

1.5.1 Solving as quadrature ode	44
1.5.2 Maple step by step solution	45

Internal problem ID [7049]

Internal file name [OUTPUT/6035_Sunday_June_05_2022_04_14_46_PM_80367263/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 5.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - y = 1$$

1.5.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{y+1} dy = \int dx$$
$$\ln(y+1) = x + c_1$$

Raising both side to exponential gives

$$y + 1 = e^{x+c_1}$$

Which simplifies to

$$y + 1 = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = c_2 e^x - 1 \tag{1}$$




Figure 10: Slope field plot

Verification of solutions

$$y = c_2 e^x - 1$$

Verified OK.

1.5.2 Maple step by step solution

Let's solve

$$y' - y = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y+1} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y+1} dx = \int 1 dx + c_1$$

- Evaluate integral

- $\ln(y + 1) = x + c_1$
Solve for y
 $y = e^{x+c_1} - 1$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x) = y(x)+1,y(x), singsol=all)
```

$$y(x) = -1 + e^x c_1$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 18

```
DSolve[y'[x] == y[x]+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1 + c_1 e^x$$

$$y(x) \rightarrow -1$$

1.6 problem 6

1.6.1 Solving as quadrature ode	47
1.6.2 Maple step by step solution	48

Internal problem ID [7050]

Internal file name [OUTPUT/6036_Sunday_June_05_2022_04_14_49_PM_28954144/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 6.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = 1 + x$$

1.6.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int 1 + x \, dx \\ &= x + \frac{1}{2}x^2 + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x + \frac{1}{2}x^2 + c_1 \tag{1}$$




Figure 11: Slope field plot

Verification of solutions

$$y = x + \frac{1}{2}x^2 + c_1$$

Verified OK.

1.6.2 Maple step by step solution

Let's solve

$$y' = 1 + x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int (1 + x) dx + c_1$$

- Evaluate integral

$$y = x + \frac{1}{2}x^2 + c_1$$

- Solve for y

$$y = x + \frac{1}{2}x^2 + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x) = 1+x,y(x), singsol=all)
```

$$y(x) = \frac{1}{2}x^2 + x + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 16

```
DSolve[y'[x]== 1+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{2} + x + c_1$$

1.7 problem 7

1.7.1 Solving as quadrature ode	50
1.7.2 Maple step by step solution	51

Internal problem ID [7051]

Internal file name [OUTPUT/6037_Sunday_June_05_2022_04_14_51_PM_54705140/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 7.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = x$$

1.7.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int x \, dx \\ &= \frac{x^2}{2} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2}{2} + c_1 \tag{1}$$




Figure 12: Slope field plot

Verification of solutions

$$y = \frac{x^2}{2} + c_1$$

Verified OK.

1.7.2 Maple step by step solution

Let's solve

$$y' = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int x dx + c_1$$

- Evaluate integral

$$y = \frac{x^2}{2} + c_1$$

- Solve for y

$$y = \frac{x^2}{2} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x) = x,y(x), singsol=all)
```

$$y(x) = \frac{x^2}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 15

```
DSolve[y'[x] == x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{2} + c_1$$

1.8 problem 8

1.8.1 Solving as quadrature ode	53
1.8.2 Maple step by step solution	54

Internal problem ID [7052]

Internal file name [OUTPUT/6038_Sunday_June_05_2022_04_14_53_PM_75126448/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 8.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - y = 0$$

1.8.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{y} dy = x + c_1$$

$$\ln(y) = x + c_1$$

$$y = e^{x+c_1}$$

$$y = c_1 e^x$$

Summary

The solution(s) found are the following

$$y = c_1 e^x \tag{1}$$




Figure 13: Slope field plot

Verification of solutions

$$y = c_1 e^x$$

Verified OK.

1.8.2 Maple step by step solution

Let's solve

$$y' - y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int 1 dx + c_1$$

- Evaluate integral

- $\ln(y) = x + c_1$
Solve for y
 $y = e^{x+c_1}$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(diff(y(x),x) = y(x),y(x), singsol=all)
```

$$y(x) = e^x c_1$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 16

```
DSolve[y'[x] == y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow 0$$

1.9 problem 9

1.9.1 Solving as quadrature ode	56
1.9.2 Maple step by step solution	57

Internal problem ID [7053]

Internal file name [OUTPUT/6039_Sunday_June_05_2022_04_14_55_PM_42731964/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 9.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = 0$$

1.9.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int 0 \, dx \\ &= c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \tag{1}$$




Figure 14: Slope field plot

Verification of solutions

$$y = c_1$$

Verified OK.

1.9.2 Maple step by step solution

Let's solve

$$y' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int 0 dx + c_1$$

- Evaluate integral

$$y = c_1$$

- Solve for y

$$y = c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(diff(y(x),x) = 0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[y'[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.10 problem 10

1.10.1 Solving as quadrature ode	59
1.10.2 Maple step by step solution	60

Internal problem ID [7054]

Internal file name [OUTPUT/6040_Sunday_June_05_2022_04_14_56_PM_58924344/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 10.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = 1 + \frac{\sec(x)}{x}$$

1.10.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{x + \sec(x)}{x} dx \\ &= x + \int \frac{2e^{ix}}{(e^{2ix} + 1)x} dx + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x + \int \frac{2e^{ix}}{(e^{2ix} + 1)x} dx + c_1 \tag{1}$$




Figure 15: Slope field plot

Verification of solutions

$$y = x + \int \frac{2e^{ix}}{(e^{2ix} + 1)x} dx + c_1$$

Verified OK.

1.10.2 Maple step by step solution

Let's solve

$$y' = 1 + \frac{\sec(x)}{x}$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int y' dx = \int \left(1 + \frac{\sec(x)}{x} \right) dx + c_1$$

- Evaluate integral

$$y = x + \int \frac{2e^{ix}}{(e^{ix})^2 + 1)x} dx + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(x),x) = 1+sec(x)/x,y(x), singsol=all)
```

$$y(x) = \int \frac{\sec(x)}{x} dx + x + c_1$$

✓ Solution by Mathematica

Time used: 0.833 (sec). Leaf size: 25

```
DSolve[y'[x] == 1+Sec[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \int_1^x \left(\frac{\sec(K[1])}{K[1]} + 1 \right) dK[1] + c_1$$

1.11 problem 11

1.11.1 Solving as linear ode	62
1.11.2 Solving as first order ode lie symmetry lookup ode	64
1.11.3 Solving as exact ode	66
1.11.4 Maple step by step solution	71

Internal problem ID [7055]

Internal file name [OUTPUT/6041_Sunday_June_05_2022_04_14_59_PM_80466540/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 11.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$y' - \frac{\sec(x)y}{x} = x$$

1.11.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{\sec(x)}{x}$$

$$q(x) = x$$

Hence the ode is

$$y' - \frac{\sec(x)y}{x} = x$$

The integrating factor μ is

$$\mu = e^{\int -\frac{\sec(x)}{x} dx}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x) \\ \frac{d}{dx}\left(e^{\int -\frac{\sec(x)}{x} dx} y\right) &= \left(e^{\int -\frac{\sec(x)}{x} dx}\right)(x) \\ d\left(e^{\int -\frac{\sec(x)}{x} dx} y\right) &= \left(x e^{-\left(\int \frac{\sec(x)}{x} dx\right)}\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{\int -\frac{\sec(x)}{x} dx} y &= \int x e^{-\left(\int \frac{\sec(x)}{x} dx\right)} dx \\ e^{\int -\frac{\sec(x)}{x} dx} y &= \int x e^{-\left(\int \frac{\sec(x)}{x} dx\right)} dx + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{\int -\frac{\sec(x)}{x} dx}$ results in

$$y = e^{\int \frac{\sec(x)}{x} dx} \left(\int x e^{-\left(\int \frac{\sec(x)}{x} dx\right)} dx \right) + c_1 e^{\int \frac{\sec(x)}{x} dx}$$

which simplifies to

$$y = e^{\int \frac{\sec(x)}{x} dx} \left(\int x e^{-\left(\int \frac{\sec(x)}{x} dx\right)} dx + c_1 \right)$$

Summary

The solution(s) found are the following

$$y = e^{\int \frac{\sec(x)}{x} dx} \left(\int x e^{-\left(\int \frac{\sec(x)}{x} dx\right)} dx + c_1 \right) \quad (1)$$




Figure 16: Slope field plot

Verification of solutions

$$y = e^{\int \frac{\sec(x)}{x} dx} \left(\int x e^{-\left(\int \frac{\sec(x)}{x} dx\right)} dx + c_1 \right)$$

Verified OK.

1.11.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\sec(x)y + x^2}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 19: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{\int \frac{\sec(x)}{x} dx}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{\int \frac{\sec(x)}{x} dx}} dy \end{aligned}$$

1.11.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might

or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} dy &= \left(x + \frac{\sec(x)y}{x} \right) dx \\ \left(-x - \frac{\sec(x)y}{x} \right) dx + dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x - \frac{\sec(x)y}{x} \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-x - \frac{\sec(x)y}{x} \right) \\ &= -\frac{\sec(x)}{x} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (1) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1 \left(\left(-\frac{\sec(x)}{x} \right) - (0) \right) \\ &= -\frac{\sec(x)}{x} \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int -\frac{\sec(x)}{x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\int -\frac{\sec(x)}{x} dx} \\ &= e^{-\left(\int \frac{\sec(x)}{x} dx\right)}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\overline{M} &= \mu M \\ &= e^{-\left(\int \frac{\sec(x)}{x} dx\right)} \left(-x - \frac{\sec(x) y}{x}\right) \\ &= -\frac{(\sec(x) y + x^2) e^{-\left(\int \frac{\sec(x)}{x} dx\right)}}{x}\end{aligned}$$

And

$$\begin{aligned}\overline{N} &= \mu N \\ &= e^{-\left(\int \frac{\sec(x)}{x} dx\right)} (1) \\ &= e^{-\left(\int \frac{\sec(x)}{x} dx\right)}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(-\frac{(\sec(x) y + x^2) e^{-\left(\int \frac{\sec(x)}{x} dx\right)}}{x}\right) + \left(e^{-\left(\int \frac{\sec(x)}{x} dx\right)}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{(\sec(x)y + x^2) e^{-\left(\int \frac{\sec(x)}{x} dx\right)}}{x} dx \\ \phi &= \int^x -\frac{(\sec(a)y + a^2) e^{-\left(\int \frac{\sec(a)}{a} da\right)}}{a} da + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-\left(\int^x \frac{\sec(a)}{a} da\right)} + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-\left(\int \frac{\sec(x)}{x} dx\right)}$. Therefore equation (4) becomes

$$e^{-\left(\int \frac{\sec(x)}{x} dx\right)} = e^{-\left(\int^x \frac{\sec(a)}{a} da\right)} + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -e^{-\left(\int^x \frac{\sec(a)}{a} da\right)} + e^{-\left(\int \frac{\sec(x)}{x} dx\right)}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(-e^{-\left(\int^x \frac{\sec(a)}{a} da\right)} + e^{-\left(\int \frac{\sec(x)}{x} dx\right)} \right) dy \\ f(y) &= \int_0^y \left(-e^{-\left(\int^x \frac{\sec(a)}{a} da\right)} + e^{-\left(\int \frac{\sec(x)}{x} dx\right)} \right) da + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\begin{aligned}\phi &= \int^x -\frac{(\sec(a)y + a^2) e^{-\left(\int \frac{\sec(a)}{a} da\right)}}{a} da \\ &\quad + \int_0^y \left(-e^{-\left(\int^x \frac{\sec(a)}{a} da\right)} + e^{-\left(\int \frac{\sec(x)}{x} dx\right)} \right) da + c_1\end{aligned}$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x \frac{(\sec(a)y + a^2) e^{-\left(\int \frac{\sec(a)}{a} da\right)}}{a} da + \int_0^y \left(-e^{-\left(\int^x \frac{\sec(a)}{a} da\right)} + e^{-\left(\int \frac{\sec(x)}{x} dx\right)} \right) da$$

Summary

The solution(s) found are the following

$$\int^x \frac{(\sec(a)y + a^2) e^{-\left(\int \frac{\sec(a)}{a} da\right)}}{a} da + \int_0^y \left(-e^{-\left(\int^x \frac{\sec(a)}{a} da\right)} + e^{-\left(\int \frac{\sec(x)}{x} dx\right)} \right) da = c_1 \quad (1)$$




Figure 17: Slope field plot

Verification of solutions

$$\int^x \frac{(\sec(a)y + a^2) e^{-\left(\int \frac{\sec(a)}{a} da\right)}}{a} da + \int_0^y \left(-e^{-\left(\int x \frac{\sec(a)}{a} da\right)} + e^{-\left(\int \frac{\sec(x)}{x} dx\right)} \right) da = c_1$$

Verified OK.

1.11.4 Maple step by step solution

Let's solve

$$y' - \frac{\sec(x)y}{x} = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = x + \frac{\sec(x)y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{\sec(x)y}{x} = x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{\sec(x)y}{x} \right) = \mu(x) x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' - \frac{\sec(x)y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{\mu(x)\sec(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = e^{\int -\frac{\sec(x)}{x} dx}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) x dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) x dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{\int -\frac{\sec(x)}{x} dx}$

$$y = \frac{\int x e^{\int -\frac{\sec(x)}{x} dx} dx + c_1}{e^{\int -\frac{\sec(x)}{x} dx}}$$

- Simplify

$$y = e^{\int \frac{\sec(x)}{x} dx} \left(\int x e^{-\left(\int \frac{\sec(x)}{x} dx\right)} dx + c_1 \right)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x) = x+sec(x)*y(x)/x,y(x), singsol=all)
```

$$y(x) = \left(\int x e^{-\left(\int \frac{\sec(x)}{x} dx\right)} dx + c_1 \right) e^{\int \frac{\sec(x)}{x} dx}$$

✓ Solution by Mathematica

Time used: 0.483 (sec). Leaf size: 56

```
DSolve[y'[x] == x+Sec[x]*y[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \exp\left(\int_1^x \frac{\sec(K[1])}{K[1]} dK[1]\right) \left(\int_1^x \exp\left(-\int_1^{K[2]} \frac{\sec(K[1])}{K[1]} dK[1]\right) K[2] dK[2] + c_1 \right)$$

1.12 problem 12

1.12.1 Existence and uniqueness analysis	73
1.12.2 Solving as separable ode	74
1.12.3 Solving as linear ode	75
1.12.4 Solving as homogeneousTypeD2 ode	76
1.12.5 Solving as first order ode lie symmetry lookup ode	77
1.12.6 Solving as exact ode	81
1.12.7 Maple step by step solution	84

Internal problem ID [7056]

Internal file name [OUTPUT/6042_Sunday_June_05_2022_04_15_02_PM_84488611/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 12.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - \frac{2y}{x} = 0$$

With initial conditions

$$[y(0) = 0]$$

1.12.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{2}{x}$$

$$q(x) = 0$$

Hence the ode is

$$y' - \frac{2y}{x} = 0$$

The domain of $p(x) = -\frac{2}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

But the point $x_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

1.12.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{2y}{x}\end{aligned}$$

Where $f(x) = \frac{2}{x}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{2}{x} dx \\ \int \frac{1}{y} dy &= \int \frac{2}{x} dx \\ \ln(y) &= 2 \ln(x) + c_1 \\ y &= e^{2 \ln(x) + c_1} \\ &= c_1 x^2\end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = 0$$

This solution is valid for any c_1 . Hence there are infinite number of solutions.

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.12.3 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{2}{x} dx} \\ &= \frac{1}{x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{y}{x^2} \right) &= 0\end{aligned}$$

Integrating gives

$$\frac{y}{x^2} = c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^2}$ results in

$$y = c_1 x^2$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = 0$$

This solution is valid for any c_1 . Hence there are infinite number of solutions.

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.12.4 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x - u(x) = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{1}{x} dx \\ \int \frac{1}{u} du &= \int \frac{1}{x} dx \\ \ln(u) &= \ln(x) + c_2 \\ u &= e^{\ln(x)+c_2} \\ &= c_2 x\end{aligned}$$

Therefore the solution y is

$$\begin{aligned}y &= xu \\ &= c_2 x^2\end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = 0$$

This solution is valid for any c_2 . Hence there are infinite number of solutions.

Summary

The solution(s) found are the following

$$y = c_2 x^2 \tag{1}$$

Verification of solutions

$$y = c_2 x^2$$

Verified OK.

1.12.5 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{2y}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 22: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x^2\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x^2} dy\end{aligned}$$

Which results in

$$S = \frac{y}{x^2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\R_y &= 0 \\S_x &= -\frac{2y}{x^3} \\S_y &= \frac{1}{x^2}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x^2} = c_1$$



Which simplifies to

$$\frac{y}{x^2} = c_1$$

Which gives

$$y = c_1 x^2$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2y}{x}$ 	$R = x$ $S = \frac{y}{x^2}$	$\frac{dS}{dR} = 0$ 

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = 0$$

This solution is valid for any c_1 . Hence there are infinite number of solutions.

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.12.6 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{1}{2y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{1}{2y}\right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$M(x, y) = -\frac{1}{x}$$
$$N(x, y) = \frac{1}{2y}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y} \left(-\frac{1}{x} \right)$$
$$= 0$$

And

$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x} \left(\frac{1}{2y} \right)$$
$$= 0$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$
$$\int \frac{\partial \phi}{\partial x} dx = \int -\frac{1}{x} dx$$
$$\phi = -\ln(x) + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{2y}$. Therefore equation (4) becomes

$$\frac{1}{2y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{2y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{2y} \right) dy$$
$$f(y) = \frac{\ln(y)}{2} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \frac{\ln(y)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \frac{\ln(y)}{2}$$

The solution becomes

$$y = e^{2c_1} x^2$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = 0$$

This solution is valid for any c_1 . Hence there are infinite number of solutions.

Summary

The solution(s) found are the following

$$y = e^{2c_1} x^2 \quad (1)$$

Verification of solutions

$$y = e^{2c_1} x^2$$

Verified OK.

1.12.7 Maple step by step solution

Let's solve

$$[y' - \frac{2y}{x} = 0, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{2}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{2}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = 2 \ln(x) + c_1$$

- Solve for y

$$y = e^{c_1} x^2$$

- Use initial condition $y(0) = 0$

$$0 = 0$$

- Solve for c_1

$$c_1 = c_1$$

- Substitute $c_1 = c_1$ into general solution and simplify

$$y = e^{c_1} x^2$$

- Solution to the IVP

$$y = e^{c_1} x^2$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve([diff(y(x),x) = 2*y(x)/x,y(0) = 0],y(x), singsol=all)
```

$$y(x) = c_1 x^2$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{y'[x] == 2*y[x]/x,y[0]==0},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

1.13 problem 13

1.13.1 Solving as separable ode	86
1.13.2 Solving as linear ode	88
1.13.3 Solving as homogeneousTypeD2 ode	89
1.13.4 Solving as first order ode lie symmetry lookup ode	91
1.13.5 Solving as exact ode	95
1.13.6 Maple step by step solution	99

Internal problem ID [7057]

Internal file name [OUTPUT/6043_Sunday_June_05_2022_04_15_04_PM_71324988/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 13.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$y' - \frac{2y}{x} = 0$$

1.13.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{2y}{x}\end{aligned}$$

Where $f(x) = \frac{2}{x}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{2}{x} dx \\ \int \frac{1}{y} dy &= \int \frac{2}{x} dx \\ \ln(y) &= 2 \ln(x) + c_1 \\ y &= e^{2 \ln(x) + c_1} \\ &= c_1 x^2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$




Figure 18: Slope field plot

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.13.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{2}{x}$$

$$q(x) = 0$$

Hence the ode is

$$y' - \frac{2y}{x} = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{2}{x} dx} \\ &= \frac{1}{x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{y}{x^2} \right) &= 0\end{aligned}$$

Integrating gives

$$\frac{y}{x^2} = c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^2}$ results in

$$y = c_1 x^2$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$




Figure 19: Slope field plot

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.13.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x - u(x) = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{1}{x} dx \\ \int \frac{1}{u} du &= \int \frac{1}{x} dx \\ \ln(u) &= \ln(x) + c_2 \\ u &= e^{\ln(x)+c_2} \\ &= c_2 x\end{aligned}$$

Therefore the solution y is

$$\begin{aligned}y &= xu \\ &= c_2 x^2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 x^2 \tag{1}$$




Figure 20: Slope field plot

Verification of solutions

$$y = c_2 x^2$$

Verified OK.

1.13.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{2y}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 25: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x^2\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x^2} dy \end{aligned}$$

Which results in

$$S = \frac{y}{x^2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{2y}{x^3} \\ S_y &= \frac{1}{x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x^2} = c_1$$



Which simplifies to

$$\frac{y}{x^2} = c_1$$

Which gives

$$y = c_1 x^2$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2y}{x}$ 	$R = x$ $S = \frac{y}{x^2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$




Figure 21: Slope field plot

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.13.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{2y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{1}{2y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x} \\ N(x, y) &= \frac{1}{2y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{2y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{2y}$. Therefore equation (4) becomes

$$\frac{1}{2y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{2y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{2y} \right) dy \\ f(y) &= \frac{\ln(y)}{2} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \frac{\ln(y)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \frac{\ln(y)}{2}$$

The solution becomes

$$y = e^{2c_1} x^2$$

Summary

The solution(s) found are the following

$$y = e^{2c_1} x^2 \tag{1}$$




Figure 22: Slope field plot

Verification of solutions

$$y = e^{2c_1} x^2$$

Verified OK.

1.13.6 Maple step by step solution

Let's solve

$$y' - \frac{2y}{x} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{2}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{2}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = 2 \ln(x) + c_1$$

- Solve for y

$$y = e^{c_1} x^2$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x) = 2*y(x)/x,y(x), singsol=all)
```

$$y(x) = c_1 x^2$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 16

```
DSolve[y'[x] == 2*y[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x^2$$

$$y(x) \rightarrow 0$$

1.14 problem 14

1.14.1 Solving as separable ode	101
1.14.2 Solving as first order ode lie symmetry lookup ode	103
1.14.3 Solving as exact ode	104
1.14.4 Maple step by step solution	108

Internal problem ID [7058]

Internal file name [OUTPUT/6044_Sunday_June_05_2022_04_15_06_PM_69283102/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 14.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - \frac{\ln(1+y^2)}{\ln(x^2+1)} = 0$$

1.14.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\ln(y^2+1)}{\ln(x^2+1)}\end{aligned}$$

Where $f(x) = \frac{1}{\ln(x^2+1)}$ and $g(y) = \ln(y^2+1)$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\ln(y^2+1)} dy &= \frac{1}{\ln(x^2+1)} dx \\ \int \frac{1}{\ln(y^2+1)} dy &= \int \frac{1}{\ln(x^2+1)} dx\end{aligned}$$

$$\int^y \frac{1}{\ln(_a^2 + 1)} d_a = \int \frac{1}{\ln(x^2 + 1)} dx + c_1$$

Which results in

$$\int^y \frac{1}{\ln(_a^2 + 1)} d_a = \int \frac{1}{\ln(x^2 + 1)} dx + c_1$$

The solution is

$$\int^y \frac{1}{\ln(_a^2 + 1)} d_a - \left(\int \frac{1}{\ln(x^2 + 1)} dx \right) - c_1 = 0$$

Summary

The solution(s) found are the following

$$\int^y \frac{1}{\ln(_a^2 + 1)} d_a - \left(\int \frac{1}{\ln(x^2 + 1)} dx \right) - c_1 = 0 \tag{1}$$




Figure 23: Slope field plot

Verification of solutions

$$\int^y \frac{1}{\ln(_a^2 + 1)} d_a - \left(\int \frac{1}{\ln(x^2 + 1)} dx \right) - c_1 = 0$$

Verified OK.

1.14.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\ln(y^2 + 1)}{\ln(x^2 + 1)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 28: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \ln(x^2 + 1) \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\ln(x^2 + 1)} dx\end{aligned}$$

Which results in

$$S = \int \frac{1}{\ln(x^2 + 1)} dx$$

1.14.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0\tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{1}{\ln(y^2 + 1)} \right) dy &= \left(\frac{1}{\ln(x^2 + 1)} \right) dx \\ \left(-\frac{1}{\ln(x^2 + 1)} \right) dx + \left(\frac{1}{\ln(y^2 + 1)} \right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{1}{\ln(x^2 + 1)} \\ N(x, y) &= \frac{1}{\ln(y^2 + 1)} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{\ln(x^2 + 1)} \right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{\ln(y^2 + 1)} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{\ln(x^2 + 1)} dx \\ \phi &= \int -\frac{1}{\ln(x^2 + 1)} dx + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\ln(y^2 + 1)}$. Therefore equation (4) becomes

$$\frac{1}{\ln(y^2 + 1)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{\ln(y^2 + 1)}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{\ln(y^2 + 1)} \right) dy$$

$$f(y) = \int_0^y \frac{1}{\ln(a^2 + 1)} da + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x -\frac{1}{\ln(a^2 + 1)} da + \int_0^y \frac{1}{\ln(a^2 + 1)} da + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x -\frac{1}{\ln(a^2 + 1)} da + \int_0^y \frac{1}{\ln(a^2 + 1)} da$$

Summary

The solution(s) found are the following

$$\int^x -\frac{1}{\ln(a^2 + 1)} da + \int_0^y \frac{1}{\ln(a^2 + 1)} da = c_1 \quad (1)$$




Figure 24: Slope field plot

Verification of solutions

$$\int^x -\frac{1}{\ln(-a^2 + 1)} d_a + \int_0^y \frac{1}{\ln(-a^2 + 1)} d_a = c_1$$

Verified OK.

1.14.4 Maple step by step solution

Let's solve

$$y' - \frac{\ln(1+y^2)}{\ln(x^2+1)} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{\ln(1+y^2)} = \frac{1}{\ln(x^2+1)}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\ln(1+y^2)} dx = \int \frac{1}{\ln(x^2+1)} dx + c_1$$

- Cannot compute integral

$$\int \frac{y'}{\ln(1+y^2)} dx = \int \frac{1}{\ln(x^2+1)} dx + c_1$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(diff(y(x),x)=ln(y(x)^2+1)/ln(x^2+1),y(x), singsol=all)
```

$$\int \frac{1}{\ln(x^2+1)} dx - \left(\int^{y(x)} \frac{1}{\ln(a^2+1)} da \right) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.64 (sec). Leaf size: 48

```
DSolve[y'[x] == Log[1+y[x]^2]/Log[1+x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{1}{\log(K[1]^2+1)} dK[1] \& \right] \left[\int_1^x \frac{1}{\log(K[2]^2+1)} dK[2] + c_1 \right]$$

$$y(x) \rightarrow 0$$

1.15 problem 15

1.15.1 Solving as quadrature ode	110
1.15.2 Maple step by step solution	111

Internal problem ID [7059]

Internal file name [OUTPUT/6045_Sunday_June_05_2022_04_15_09_PM_85665527/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 15.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = \frac{1}{x}$$

1.15.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{1}{x} dx \\ &= \ln(x) + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(x) + c_1 \tag{1}$$




Figure 25: Slope field plot

Verification of solutions

$$y = \ln(x) + c_1$$

Verified OK.

1.15.2 Maple step by step solution

Let's solve

$$y' = \frac{1}{x}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$y = \ln(x) + c_1$$

- Solve for y

$$y = \ln(x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(diff(y(x),x)=1/x,y(x), singsol=all)
```

$$y(x) = \ln(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 10

```
DSolve[y'[x] == 1/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(x) + c_1$$

1.16 problem 16

- 1.16.1 Solving as first order ode lie symmetry calculated ode 113
- 1.16.2 Solving as exact ode 119

Internal problem ID [7060]

Internal file name [OUTPUT/6046_Sunday_June_05_2022_04_15_11_PM_93014366/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 16.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exactWithIntegrationFactor",
"first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, [_Abel, `2nd type`, `class B`]]
```

$$y' - \frac{-yx - 1}{4yx^3 - 2x^2} = 0$$

1.16.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{xy + 1}{2x^2(2xy - 1)}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{(xy+1)(b_3-a_2)}{2x^2(2xy-1)} - \frac{(xy+1)^2 a_3}{4x^4(2xy-1)^2} \\ - \left(-\frac{y}{2x^2(2xy-1)} + \frac{xy+1}{x^3(2xy-1)} + \frac{(xy+1)y}{x^2(2xy-1)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{1}{2x(2xy-1)} + \frac{xy+1}{x(2xy-1)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{16x^6y^2b_2 - 16x^5yb_2 - 4x^4y^2a_2 - 4x^4y^2b_3 - 8x^3y^3a_3 - 8x^3y^2a_1 - 2b_2x^4 - 8x^3ya_2 - 8x^3yb_3 - 11x^2y^2a_3 - 6x^3b_1 - 10x^2ya_1 + 2x^2a_2 + 2x^2b_3 + 2xya_3 + 4xa_1 - a_3}{4x^4(2xy-1)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} 16x^6y^2b_2 - 16x^5yb_2 - 4x^4y^2a_2 - 4x^4y^2b_3 - 8x^3y^3a_3 - 8x^3y^2a_1 - 2b_2x^4 - 8x^3ya_2 \\ - 8x^3yb_3 - 11x^2y^2a_3 - 6x^3b_1 - 10x^2ya_1 + 2x^2a_2 + 2x^2b_3 + 2xya_3 + 4xa_1 - a_3 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} 16b_2v_1^6v_2^2 - 4a_2v_1^4v_2^2 - 8a_3v_1^3v_2^3 - 16b_2v_1^5v_2 - 4b_3v_1^4v_2^2 - 8a_1v_1^3v_2^2 \\ - 8a_2v_1^3v_2 - 11a_3v_1^2v_2^2 - 2b_2v_1^4 - 8b_3v_1^3v_2 - 10a_1v_1^2v_2 \\ - 6b_1v_1^3 + 2a_2v_1^2 + 2a_3v_1v_2 + 2b_3v_1^2 + 4a_1v_1 - a_3 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} &16b_2v_1^6v_2^2 - 16b_2v_1^5v_2 + (-4a_2 - 4b_3)v_1^4v_2^2 - 2b_2v_1^4 - 8a_3v_1^3v_2^3 \\ &- 8a_1v_1^3v_2^2 + (-8a_2 - 8b_3)v_1^3v_2 - 6b_1v_1^3 - 11a_3v_1^2v_2^2 \\ &- 10a_1v_1^2v_2 + (2a_2 + 2b_3)v_1^2 + 2a_3v_1v_2 + 4a_1v_1 - a_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -10a_1 &= 0 \\ -8a_1 &= 0 \\ 4a_1 &= 0 \\ -11a_3 &= 0 \\ -8a_3 &= 0 \\ -a_3 &= 0 \\ 2a_3 &= 0 \\ -6b_1 &= 0 \\ -16b_2 &= 0 \\ -2b_2 &= 0 \\ 16b_2 &= 0 \\ -8a_2 - 8b_3 &= 0 \\ -4a_2 - 4b_3 &= 0 \\ 2a_2 + 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= -b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}\xi &= -x \\ \eta &= y\end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(-\frac{xy + 1}{2x^2(2xy - 1)} \right) (-x) \\ &= \frac{4x^2y^2 - 3xy - 1}{4yx^2 - 2x} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{4x^2y^2 - 3xy - 1}{4yx^2 - 2x}} dy\end{aligned}$$

Which results in

$$S = \frac{3 \ln(4xy + 1)}{5} + \frac{2 \ln(xy - 1)}{5}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{xy + 1}{2x^2(2xy - 1)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{4x y^2 - 2y}{4x^2 y^2 - 3xy - 1} \\ S_y &= \frac{4y x^2 - 2x}{4x^2 y^2 - 3xy - 1} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{x} \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \ln(R) + c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{3 \ln(1 + 4yx)}{5} + \frac{2 \ln(yx - 1)}{5} = \ln(x) + c_1$$

Which simplifies to

$$\frac{3 \ln(1 + 4yx)}{5} + \frac{2 \ln(yx - 1)}{5} = \ln(x) + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{xy+1}{2x^2(2xy-1)}$ 	$R = x$ $S = \frac{3 \ln(4xy + 1)}{5} + \frac{2 \ln}{5}$	$\frac{dS}{dR} = \frac{1}{R}$ 

Summary

The solution(s) found are the following

$$\frac{3 \ln(1 + 4yx)}{5} + \frac{2 \ln(yx - 1)}{5} = \ln(x) + c_1 \quad (1)$$




Figure 26: Slope field plot

Verification of solutions

$$\frac{3 \ln(1 + 4yx)}{5} + \frac{2 \ln(yx - 1)}{5} = \ln(x) + c_1$$

Verified OK.

1.16.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}dy &= \left(\frac{-xy - 1}{4y x^3 - 2x^2} \right) dx \\ \left(-\frac{-xy - 1}{4y x^3 - 2x^2} \right) dx + dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{-xy - 1}{4y x^3 - 2x^2} \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{-xy - 1}{4y x^3 - 2x^2} \right) \\ &= -\frac{3}{2x(2xy - 1)^2}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1 \left(\left(\frac{x}{4y x^3 - 2x^2} + \frac{4(-xy - 1)x^3}{(4y x^3 - 2x^2)^2} \right) - (0) \right) \\ &= -\frac{3}{2x(2xy - 1)^2}\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{4y x^3 - 2x^2}{xy + 1} \left((0) - \left(\frac{x}{4y x^3 - 2x^2} + \frac{4(-xy - 1)x^3}{(4y x^3 - 2x^2)^2} \right) \right) \\ &= \frac{3x}{2x^2y^2 + xy - 1}\end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned}R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(0) - \left(\frac{x}{4y x^3 - 2x^2} + \frac{4(-xy - 1)x^3}{(4y x^3 - 2x^2)^2} \right)}{x \left(-\frac{-xy - 1}{4y x^3 - 2x^2} \right) - y(1)} \\ &= -\frac{3}{8x^3y^3 - 10x^2y^2 + xy + 1}\end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = -\frac{3}{8t^3 - 10t^2 + t + 1}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned}\mu &= e^{\int R dt} \\ &= e^{\int \left(-\frac{3}{8t^3-10t^2+t+1}\right) dt}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\ln(2t-1) - \frac{2\ln(4t+1)}{5} - \frac{3\ln(t-1)}{5}} \\ &= \frac{2t-1}{(4t+1)^{\frac{2}{5}}(t-1)^{\frac{3}{5}}}\end{aligned}$$

Now t is replaced back with xy giving

$$\mu = \frac{2xy-1}{(4xy+1)^{\frac{2}{5}}(xy-1)^{\frac{3}{5}}}$$

Multiplying M and N by this integrating factor gives new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{2xy-1}{(4xy+1)^{\frac{2}{5}}(xy-1)^{\frac{3}{5}}} \left(-\frac{-xy-1}{4yx^3-2x^2} \right) \\ &= \frac{xy+1}{2(xy-1)^{\frac{3}{5}}(4xy+1)^{\frac{2}{5}}x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{2xy-1}{(4xy+1)^{\frac{2}{5}}(xy-1)^{\frac{3}{5}}} (1) \\ &= \frac{2xy-1}{(4xy+1)^{\frac{2}{5}}(xy-1)^{\frac{3}{5}}}\end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{xy+1}{2(xy-1)^{\frac{3}{5}}(4xy+1)^{\frac{2}{5}}x^2} \right) + \left(\frac{2xy-1}{(4xy+1)^{\frac{2}{5}}(xy-1)^{\frac{3}{5}}} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{xy + 1}{2(xy - 1)^{\frac{3}{5}}(4xy + 1)^{\frac{2}{5}}x^2} dx \\ \phi &= \frac{(4xy + 1)^{\frac{3}{5}}(xy - 1)^{\frac{2}{5}}}{2x} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned} \frac{\partial \phi}{\partial y} &= \frac{6(xy - 1)^{\frac{2}{5}}}{5(4xy + 1)^{\frac{2}{5}}} + \frac{(4xy + 1)^{\frac{3}{5}}}{5(xy - 1)^{\frac{3}{5}}} + f'(y) \\ &= \frac{2xy - 1}{(4xy + 1)^{\frac{2}{5}}(xy - 1)^{\frac{3}{5}}} + f'(y) \end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{2xy-1}{(4xy+1)^{\frac{2}{5}}(xy-1)^{\frac{3}{5}}}$. Therefore equation (4) becomes

$$\frac{2xy - 1}{(4xy + 1)^{\frac{2}{5}}(xy - 1)^{\frac{3}{5}}} = \frac{2xy - 1}{(4xy + 1)^{\frac{2}{5}}(xy - 1)^{\frac{3}{5}}} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(4xy + 1)^{\frac{3}{5}}(xy - 1)^{\frac{2}{5}}}{2x} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(4xy + 1)^{\frac{3}{5}} (xy - 1)^{\frac{2}{5}}}{2x}$$

Summary

The solution(s) found are the following

$$\frac{(1 + 4yx)^{\frac{3}{5}} (yx - 1)^{\frac{2}{5}}}{2x} = c_1 \quad (1)$$




Figure 27: Slope field plot

Verification of solutions

$$\frac{(1 + 4yx)^{\frac{3}{5}} (yx - 1)^{\frac{2}{5}}}{2x} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous G  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.562 (sec). Leaf size: 37

```
dsolve(diff(y(x),x)=(-x*y(x)-1)/(4*x^3*y(x)-2*x^2),y(x), singsol=all)
```

$$y(x) = \frac{\text{RootOf}(_Z^{25}c_1 - 10_Z^{20}c_1 + 25_Z^{15}c_1 - 16x^5)^5 - 1}{4x}$$

✓ Solution by Mathematica

Time used: 15.76 (sec). Leaf size: 391

```
DSolve[y'[x] == (-x*y[x]-1)/(4*x^3*y[x]-2*x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{Root}[64\#1^5c_1^5x^5 - 80\#1^4c_1^5x^4 - 20\#1^3c_1^5x^3 + 25\#1^2c_1^5x^2 + 10\#1c_1^5x - x^5 + c_1^5\&, 1]$$

$$y(x) \rightarrow \text{Root}[64\#1^5c_1^5x^5 - 80\#1^4c_1^5x^4 - 20\#1^3c_1^5x^3 + 25\#1^2c_1^5x^2 + 10\#1c_1^5x - x^5 + c_1^5\&, 2]$$

$$y(x) \rightarrow \text{Root}[64\#1^5c_1^5x^5 - 80\#1^4c_1^5x^4 - 20\#1^3c_1^5x^3 + 25\#1^2c_1^5x^2 + 10\#1c_1^5x - x^5 + c_1^5\&, 3]$$

$$y(x) \rightarrow \text{Root}[64\#1^5c_1^5x^5 - 80\#1^4c_1^5x^4 - 20\#1^3c_1^5x^3 + 25\#1^2c_1^5x^2 + 10\#1c_1^5x - x^5 + c_1^5\&, 4]$$

$$y(x) \rightarrow \text{Root}[64\#1^5c_1^5x^5 - 80\#1^4c_1^5x^4 - 20\#1^3c_1^5x^3 + 25\#1^2c_1^5x^2 + 10\#1c_1^5x - x^5 + c_1^5\&, 5]$$

1.17 problem 17

1.17.1 Solving as clairaut ode 126

Internal problem ID [7061]

Internal file name [OUTPUT/6047_Sunday_June_05_2022_04_15_15_PM_17424678/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 17.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**clairaut**"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries], _Clairaut]
```

$$\frac{y'^2}{4} - xy' + y = 0$$

1.17.1 Solving as clairaut ode

This is Clairaut ODE. It has the form

$$y = xy' + g(y')$$

Where g is function of $y'(x)$. Let $p = y'$ the ode becomes

$$\frac{1}{4}p^2 - xp + y = 0$$

Solving for y from the above results in

$$y = -\frac{1}{4}p^2 + xp \tag{1A}$$

The above ode is a Clairaut ode which is now solved. We start by replacing y' by p which gives

$$\begin{aligned} y &= -\frac{1}{4}p^2 + xp \\ &= -\frac{1}{4}p^2 + xp \end{aligned}$$

Writing the ode as

$$y = xp + g(p)$$

We now write $g \equiv g(p)$ to make notation simpler but we should always remember that g is function of p which in turn is function of x . Hence the above becomes

$$y = xp + g \tag{1}$$

Then we see that

$$g = -\frac{p^2}{4}$$

Taking derivative of (1) w.r.t. x gives

$$\begin{aligned} p &= \frac{d}{dx}(xp + g) \\ p &= \left(p + x \frac{dp}{dx} \right) + \left(g' \frac{dp}{dx} \right) \\ p &= p + (x + g') \frac{dp}{dx} \\ 0 &= (x + g') \frac{dp}{dx} \end{aligned}$$

Where g' is derivative of $g(p)$ w.r.t. p . The general solution is given by

$$\begin{aligned} \frac{dp}{dx} &= 0 \\ p &= c_1 \end{aligned}$$

Substituting this in (1) gives the general solution as

$$y = c_1 x - \frac{1}{4} c_1^2$$

The singular solution is found from solving for p from

$$x + g'(p) = 0$$

And substituting the result back in (1). Since we found above that $g = -\frac{p^2}{4}$, then the above equation becomes

$$\begin{aligned} x + g'(p) &= x - \frac{p}{2} \\ &= 0 \end{aligned}$$

Solving the above for p results in

$$p_1 = 2x$$

Substituting the above back in (1) results in

$$y_1 = x^2$$

Summary

The solution(s) found are the following

$$y = c_1x - \frac{1}{4}c_1^2 \quad (1)$$

$$y = x^2 \quad (2)$$

Verification of solutions

$$y = c_1x - \frac{1}{4}c_1^2$$

Verified OK.

$$y = x^2$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
*** Sublevel 2 ***
Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
trying dAlembert
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 18

```
dsolve((1/4)*diff(y(x),x)^2-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = x^2$$
$$y(x) = -\frac{c_1(c_1 - 4x)}{4}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 25

```
DSolve[(1/4)*(y'[x])^2-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x - \frac{c_1^2}{4}$$
$$y(x) \rightarrow x^2$$

1.18 problem 18

1.18.1 Existence and uniqueness analysis	130
1.18.2 Solving as quadrature ode	131
1.18.3 Maple step by step solution	132

Internal problem ID [7062]

Internal file name [OUTPUT/6048_Sunday_June_05_2022_04_15_19_PM_15217227/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 18.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"quadrature"**

Maple gives the following as the ode type

`[_quadrature]`

$$y' - \sqrt{\frac{y+1}{y^2}} = 0$$

With initial conditions

$$[y(0) = 1]$$

1.18.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= \sqrt{\frac{y+1}{y^2}}\end{aligned}$$

The y domain of $f(x, y)$ when $x = 0$ is

$$\{-1 \leq y < 0, 0 < y \leq \infty\}$$

And the point $y_0 = 1$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(\sqrt{\frac{y+1}{y^2}} \right) \\ &= \frac{\frac{1}{y^2} - \frac{2(y+1)}{y^3}}{2\sqrt{\frac{y+1}{y^2}}}\end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{-\infty \leq y < -1, -1 < y < 0, 0 < y \leq \infty\}$$

And the point $y_0 = 1$ is inside this domain. Therefore solution exists and is unique.

1.18.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned}\int \frac{1}{\sqrt{\frac{y+1}{y^2}}} dy &= \int dx \\ \frac{2(y+1)(y-2)}{3y\sqrt{\frac{y+1}{y^2}}} &= x + c_1\end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$-\frac{2\sqrt{2}}{3} = c_1$$

$$c_1 = -\frac{2\sqrt{2}}{3}$$

Substituting c_1 found above in the general solution gives

$$\frac{2(y+1)(y-2)}{3y\sqrt{\frac{y+1}{y^2}}} = x - \frac{2\sqrt{2}}{3}$$

The above simplifies to

$$2\sqrt{2}y\sqrt{\frac{y+1}{y^2}} - 3xy\sqrt{\frac{y+1}{y^2}} + 2y^2 - 2y - 4 = 0$$

Summary

The solution(s) found are the following

$$-3\left(x - \frac{2\sqrt{2}}{3}\right)y\sqrt{\frac{y+1}{y^2}} + 2y^2 - 2y - 4 = 0 \quad (1)$$

Verification of solutions

$$-3\left(x - \frac{2\sqrt{2}}{3}\right)y\sqrt{\frac{y+1}{y^2}} + 2y^2 - 2y - 4 = 0$$

Verified OK.

1.18.3 Maple step by step solution

Let's solve

$$\left[y' - \sqrt{\frac{y+1}{y^2}} = 0, y(0) = 1 \right]$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{\sqrt{\frac{y+1}{y^2}}} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{\frac{y+1}{y^2}}} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\frac{2(y+1)(y-2)}{3y\sqrt{\frac{y+1}{y^2}}} = x + c_1$$

- Solve for y

$$y = \frac{\left(-8+9c_1^2+18c_1x+9x^2+3\sqrt{9c_1^4+36c_1^3x+54c_1^2x^2+36c_1x^3+9x^4-16c_1^2-32c_1x-16x^2}\right)^{\frac{1}{3}}}{2} + \frac{\left(-8+9c_1^2+18c_1x+9x^2+3\sqrt{9c_1^4+36c_1^3}\right)^{\frac{1}{3}}}{\left(-8+9c_1^2+3\sqrt{9c_1^4-16c_1^2}\right)^{\frac{1}{3}}}$$

- Use initial condition $y(0) = 1$

$$1 = \frac{\left(-8+9c_1^2+3\sqrt{9c_1^4-16c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{2}{\left(-8+9c_1^2+3\sqrt{9c_1^4-16c_1^2}\right)^{\frac{1}{3}}} + 1$$

- Solve for c_1

$$c_1 = \text{RootOf}\left(\left(-8 + 9Z^2 + 3\sqrt{9Z^4 - 16Z^2}\right)^{\frac{2}{3}} + 4\right)$$

- Substitute $c_1 = \text{RootOf}\left(\left(-8 + 9Z^2 + 3\sqrt{9Z^4 - 16Z^2}\right)^{\frac{2}{3}} + 4\right)$ into general solution and simplify

$$y = \frac{\left(-8 + 9\text{RootOf}\left(\left(-8 + 9Z^2 + 3\sqrt{9Z^4 - 16Z^2}\right)^{\frac{2}{3}} + 4\right)\right)^2 + 18\text{RootOf}\left(\left(-8 + 9Z^2 + 3\sqrt{9Z^4 - 16Z^2}\right)^{\frac{2}{3}} + 4\right)}{x + 9}$$

- Solution to the IVP

$$y = \frac{\left(-8 + 9\text{RootOf}\left(\left(-8 + 9Z^2 + 3\sqrt{9Z^4 - 16Z^2}\right)^{\frac{2}{3}} + 4\right)\right)^2 + 18\text{RootOf}\left(\left(-8 + 9Z^2 + 3\sqrt{9Z^4 - 16Z^2}\right)^{\frac{2}{3}} + 4\right)}{x + 9}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.469 (sec). Leaf size: 148

```
dsolve([diff(y(x),x)=sqrt((1+y(x))/y(x)^2),y(0) = 1],y(x), singsol=all)
```

$y(x) =$

$$\frac{(1 + i\sqrt{3}) \left(-12\sqrt{2}x + 9x^2 + \sqrt{(-12\sqrt{2}x + 9x^2 - 8)(3x - 2\sqrt{2})^2}\right)^{\frac{2}{3}} - 4i\sqrt{3} - 4 \left(-12\sqrt{2}x + 9x^2 + \sqrt{(-12\sqrt{2}x + 9x^2 - 8)(3x - 2\sqrt{2})^2}\right)^{\frac{2}{3}}}{4 \left(-12\sqrt{2}x + 9x^2 + \sqrt{(-12\sqrt{2}x + 9x^2 - 8)(3x - 2\sqrt{2})^2}\right)^{\frac{2}{3}}}$$

✓ Solution by Mathematica

Time used: 0.097 (sec). Leaf size: 123

```
DSolve[{y'[x]==Sqrt[(1+y[x])/y[x]^2],y[0]==1},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{4} \left(1 + i\sqrt{3}\right) \sqrt[3]{9x^2 + \sqrt{81x^4 - 216\sqrt{2}x^3 + 288x^2 - 64} - 12\sqrt{2}x} \\ + \frac{i(\sqrt{3} + i)}{\sqrt[3]{9x^2 + \sqrt{81x^4 - 216\sqrt{2}x^3 + 288x^2 - 64} - 12\sqrt{2}x}} + 1$$

1.19 problem 19

Internal problem ID [7063]

Internal file name [OUTPUT/6049_Sunday_June_05_2022_04_15_22_PM_6405240/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 19.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

[`y=_G(x,y)']

Unable to solve or complete the solution.

$$y' - \sqrt{1 - x^2 - y^2} = 0$$

Unable to determine ODE type.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying homogeneous types:
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying an equivalence to an Abel ODE
trying 1st order ODE linearizable_by_differentiation
--- Trying Lie symmetry methods, 1st order ---
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 5
trying symmetry patterns for 1st order ODEs
-> trying a symmetry pattern of the form [F(x)*G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)*G(y)]
-> trying symmetry patterns of the forms [F(x),G(y)] and [G(y),F(x)]
-> trying a symmetry pattern of the form [F(x),G(x)]
-> trying a symmetry pattern of the form [F(y),G(y)]
-> trying a symmetry pattern of the form [F(x)+G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)+G(y)]
-> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying a symmetry pattern of conformal type`
```

X Solution by Maple

```
dsolve(diff(y(x),x)=sqrt( 1-x^2-y(x)^2),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==Sqrt[ 1-x^2-y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

1.20 problem 20

- 1.20.1 Solving as first order ode lie symmetry lookup ode 137
- 1.20.2 Solving as bernoulli ode 141

Internal problem ID [7064]

Internal file name [OUTPUT/6050_Sunday_June_05_2022_04_15_25_PM_21854630/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 20.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**bernoulli**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

`[_Bernoulli]`

$$y' + \frac{y}{3} - \frac{(1-2x)y^4}{3} = 0$$

1.20.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{1}{3}y - \frac{2}{3}y^4x + \frac{1}{3}y^4$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \tag{A}$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 33: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= y^4 e^x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{y^4 e^x} dy \end{aligned}$$

Which results in

$$S = -\frac{e^{-x}}{3y^3}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{1}{3}y - \frac{2}{3}y^4x + \frac{1}{3}y^4$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{e^{-x}}{3y^3} \\ S_y &= \frac{e^{-x}}{y^4} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{e^{-x}(2x-1)}{3} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{e^{-R}(2R-1)}{3}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{(2R + 1)e^{-R}}{3} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{e^{-x}}{3y^3} = \frac{(2x + 1)e^{-x}}{3} + c_1$$

Which simplifies to

$$-\frac{e^{-x}}{3y^3} = \frac{(2x + 1)e^{-x}}{3} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{1}{3}y - \frac{2}{3}y^4x + \frac{1}{3}y^4$ 	$R = x$ $S = -\frac{e^{-x}}{3y^3}$	$\frac{dS}{dR} = -\frac{e^{-R}(2R-1)}{3}$ 

Summary

The solution(s) found are the following

$$-\frac{e^{-x}}{3y^3} = \frac{(2x + 1)e^{-x}}{3} + c_1 \quad (1)$$




Figure 28: Slope field plot

Verification of solutions

$$-\frac{e^{-x}}{3y^3} = \frac{(2x+1)e^{-x}}{3} + c_1$$

Verified OK.

1.20.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{1}{3}y - \frac{2}{3}y^4x + \frac{1}{3}y^4 \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -\frac{1}{3}y - \frac{2x}{3} + \frac{1}{3}y^4 \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= -\frac{1}{3} \\ f_1(x) &= -\frac{2x}{3} + \frac{1}{3} \\ n &= 4 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^4$ gives

$$y' \frac{1}{y^4} = -\frac{1}{3y^3} - \frac{2x}{3} + \frac{1}{3} \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= \frac{1}{y^3} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{3}{y^4} y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -\frac{w'(x)}{3} &= -\frac{w(x)}{3} - \frac{2x}{3} + \frac{1}{3} \\ w' &= w + 2x - 1 \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned} p(x) &= -1 \\ q(x) &= 2x - 1 \end{aligned}$$

Hence the ode is

$$w'(x) - w(x) = 2x - 1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int(-1)dx} \\ &= e^{-x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu)(2x - 1) \\ \frac{d}{dx}(e^{-x}w) &= (e^{-x})(2x - 1) \\ d(e^{-x}w) &= (e^{-x}(2x - 1)) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-x}w &= \int e^{-x}(2x - 1) dx \\ e^{-x}w &= -(2x + 1)e^{-x} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-x}$ results in

$$w(x) = -e^x(2x + 1)e^{-x} + c_1e^x$$

which simplifies to

$$w(x) = -2x - 1 + c_1e^x$$

Replacing w in the above by $\frac{1}{y^3}$ using equation (5) gives the final solution.

$$\frac{1}{y^3} = -2x - 1 + c_1e^x$$

Solving for y gives

$$\begin{aligned}y(x) &= \frac{1}{(-2x - 1 + c_1e^x)^{\frac{1}{3}}} \\ y(x) &= \frac{i\sqrt{3} - 1}{2(-2x - 1 + c_1e^x)^{\frac{1}{3}}} \\ y(x) &= -\frac{1 + i\sqrt{3}}{2(-2x - 1 + c_1e^x)^{\frac{1}{3}}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{(-2x - 1 + c_1 e^x)^{\frac{1}{3}}} \quad (1)$$

$$y = \frac{i\sqrt{3} - 1}{2(-2x - 1 + c_1 e^x)^{\frac{1}{3}}} \quad (2)$$

$$y = -\frac{1 + i\sqrt{3}}{2(-2x - 1 + c_1 e^x)^{\frac{1}{3}}} \quad (3)$$




Figure 29: Slope field plot

Verification of solutions

$$y = \frac{1}{(-2x - 1 + c_1 e^x)^{\frac{1}{3}}}$$

Verified OK.

$$y = \frac{i\sqrt{3} - 1}{2(-2x - 1 + c_1 e^x)^{\frac{1}{3}}}$$

Verified OK.

$$y = -\frac{1 + i\sqrt{3}}{2(-2x - 1 + c_1 e^x)^{\frac{1}{3}}}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

```
dsolve(diff(y(x),x)+y(x)/3=(1-2*x)/3*y(x)^4,y(x), singsol=all)
```

$$y(x) = \frac{1}{(e^x c_1 - 2x - 1)^{\frac{1}{3}}}$$

$$y(x) = -\frac{1 + i\sqrt{3}}{2(e^x c_1 - 2x - 1)^{\frac{1}{3}}}$$

$$y(x) = \frac{i\sqrt{3} - 1}{2(e^x c_1 - 2x - 1)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 4.53 (sec). Leaf size: 76

```
DSolve[y'[x]+y[x]/3==(1-2*x)/3*y[x]^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{\sqrt[3]{-2x + c_1 e^x - 1}}$$
$$y(x) \rightarrow -\frac{\sqrt[3]{-1}}{\sqrt[3]{-2x + c_1 e^x - 1}}$$
$$y(x) \rightarrow \frac{(-1)^{2/3}}{\sqrt[3]{-2x + c_1 e^x - 1}}$$
$$y(x) \rightarrow 0$$

1.21 problem 21

1.21.1 Solving as first order ode lie symmetry calculated ode 147

Internal problem ID [7065]

Internal file name [OUTPUT/6051_Sunday_June_05_2022_04_15_29_PM_29923346/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 21.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries] , _Chini]
```

$$y' - \sqrt{y} = x$$

1.21.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \sqrt{y} + x$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + (\sqrt{y} + x)(b_3 - a_2) - (\sqrt{y} + x)^2 a_3 - xa_2 - ya_3 - a_1 - \frac{xb_2 + yb_3 + b_1}{2\sqrt{y}} = 0 \quad (5E)$$

Putting the above in normal form gives

$$\frac{4y^{\frac{3}{2}}a_3 + 4yxa_3 + 2\sqrt{y}x^2a_3 + 2ya_2 - yb_3 + 4xa_2\sqrt{y} - 2\sqrt{y}xb_3 + 2a_1\sqrt{y} - 2b_2\sqrt{y} + xb_2 + b_1}{2\sqrt{y}} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -4y^{\frac{3}{2}}a_3 - 2\sqrt{y}x^2a_3 - 4xa_2\sqrt{y} + 2\sqrt{y}xb_3 - 4yxa_3 \\ - 2a_1\sqrt{y} + 2b_2\sqrt{y} - xb_2 - 2ya_2 + yb_3 - b_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \sqrt{y}, y^{\frac{3}{2}}\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \sqrt{y} = v_3, y^{\frac{3}{2}} = v_4\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -2v_3v_1^2a_3 - 4v_1a_2v_3 - 4v_2v_1a_3 + 2v_3v_1b_3 - 2a_1v_3 \\ - 2v_2a_2 - 4v_4a_3 - v_1b_2 + 2b_2v_3 + v_2b_3 - b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4\}$$

Equation (7E) now becomes

$$\begin{aligned} -2v_3v_1^2a_3 - 4v_2v_1a_3 + (-4a_2 + 2b_3)v_1v_3 - v_1b_2 \\ + (-2a_2 + b_3)v_2 + (-2a_1 + 2b_2)v_3 - 4v_4a_3 - b_1 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 -4a_3 &= 0 \\
 -2a_3 &= 0 \\
 -b_1 &= 0 \\
 -b_2 &= 0 \\
 -2a_1 + 2b_2 &= 0 \\
 -4a_2 + 2b_3 &= 0 \\
 -2a_2 + b_3 &= 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= 0 \\
 a_2 &= a_2 \\
 a_3 &= 0 \\
 b_1 &= 0 \\
 b_2 &= 0 \\
 b_3 &= 2a_2
 \end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= x \\
 \eta &= 2y
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= 2y - (\sqrt{y} + x)(x) \\
 &= -x\sqrt{y} - x^2 + 2y \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-x\sqrt{y} - x^2 + 2y} dy \end{aligned}$$

Which results in

$$S = \frac{\ln(x + 2\sqrt{y})}{6} - \frac{\ln(\sqrt{y} + x)}{3} - \frac{\ln(-x + 2\sqrt{y})}{6} + \frac{\ln(\sqrt{y} - x)}{3} + \frac{\ln(-x^2 + y)}{3} + \frac{\ln(-x^2 + 4y)}{6}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \sqrt{y} + x$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{x^3 - 3xy - 2y^{\frac{3}{2}}}{(x^2 - 4y)(x^2 - y)} \\ S_y &= \frac{-x^2 + x\sqrt{y} + 2y}{(x^2 - 4y)(x^2 - y)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(x + 2\sqrt{y})}{6} - \frac{\ln(\sqrt{y} + x)}{3} - \frac{\ln(-x + 2\sqrt{y})}{6} + \frac{\ln(\sqrt{y} - x)}{3} + \frac{\ln(-x^2 + y)}{3} + \frac{\ln(-x^2 + 4y)}{6} = c_1$$

Which simplifies to

$$\frac{\ln(x + 2\sqrt{y})}{6} - \frac{\ln(\sqrt{y} + x)}{3} - \frac{\ln(-x + 2\sqrt{y})}{6} + \frac{\ln(\sqrt{y} - x)}{3} + \frac{\ln(-x^2 + y)}{3} + \frac{\ln(-x^2 + 4y)}{6} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \sqrt{y} + x$ 	$R = x$ $S = \frac{\ln(x + 2\sqrt{y})}{6} - \ln(\sqrt{y} - x)$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{\ln(x + 2\sqrt{y})}{6} - \frac{\ln(\sqrt{y} + x)}{3} - \frac{\ln(-x + 2\sqrt{y})}{6} + \frac{\ln(\sqrt{y} - x)}{3} + \frac{\ln(-x^2 + y)}{3} + \frac{\ln(-x^2 + 4y)}{6} = c_1 \tag{1}$$




Figure 30: Slope field plot

Verification of solutions

$$\frac{\ln(x + 2\sqrt{y})}{6} - \frac{\ln(\sqrt{y} + x)}{3} - \frac{\ln(-x + 2\sqrt{y})}{6} + \frac{\ln(\sqrt{y} - x)}{3} + \frac{\ln(-x^2 + y)}{3} + \frac{\ln(-x^2 + 4y)}{6} = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
<- Chini successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 69

```
dsolve(diff(y(x),x)=sqrt(y(x))+x,y(x), singsol=all)
```

$$\frac{4 \operatorname{arctanh}\left(\sqrt{\frac{y(x)}{x^2}}\right)}{3} - \frac{2 \operatorname{arctanh}\left(2\sqrt{\frac{y(x)}{x^2}}\right)}{3} - \frac{\ln\left(\frac{-x^2+4y(x)}{x^2}\right)}{3} - \frac{2 \ln(2)}{3} - \frac{2 \ln\left(\frac{y(x)-x^2}{x^2}\right)}{3} - 2 \ln(x) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 47.265 (sec). Leaf size: 716

`DSolve[y'[x]==Sqrt[y[x]]+x,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{4} \left(3x^2 + \frac{e^{3c_1} x (8 + e^{3c_1} x^3)}{\sqrt[3]{-e^{18c_1} x^6 + 20e^{15c_1} x^3 + 8\sqrt{-e^{24c_1} (-1 + e^{3c_1} x^3)^3 + 8e^{12c_1}}}} + e^{-6c_1} \sqrt[3]{-e^{18c_1} x^6 + 20e^{15c_1} x^3 + 8\sqrt{-e^{24c_1} (-1 + e^{3c_1} x^3)^3 + 8e^{12c_1}}} \right)$$

$$y(x) \rightarrow \frac{1}{72} \left(54x^2 - \frac{9i(\sqrt{3} - i) e^{3c_1} x (8 + e^{3c_1} x^3)}{\sqrt[3]{-e^{18c_1} x^6 + 20e^{15c_1} x^3 + 8\sqrt{-e^{24c_1} (-1 + e^{3c_1} x^3)^3 + 8e^{12c_1}}}} + 9i(\sqrt{3} + i) e^{-6c_1} \sqrt[3]{-e^{18c_1} x^6 + 20e^{15c_1} x^3 + 8\sqrt{-e^{24c_1} (-1 + e^{3c_1} x^3)^3 + 8e^{12c_1}}} \right)$$

$$y(x) \rightarrow \frac{1}{72} \left(54x^2 + \frac{9i(\sqrt{3} + i) e^{3c_1} x (8 + e^{3c_1} x^3)}{\sqrt[3]{-e^{18c_1} x^6 + 20e^{15c_1} x^3 + 8\sqrt{-e^{24c_1} (-1 + e^{3c_1} x^3)^3 + 8e^{12c_1}}}} - 9 \left(1 + i\sqrt{3} \right) e^{-6c_1} \sqrt[3]{-e^{18c_1} x^6 + 20e^{15c_1} x^3 + 8\sqrt{-e^{24c_1} (-1 + e^{3c_1} x^3)^3 + 8e^{12c_1}}} \right)$$

$$y(x) \rightarrow \frac{-(-x^6)^{2/3} + 3x^4 + \sqrt[3]{-x^6} x^2}{4x^2}$$

$$y(x) \rightarrow \frac{(1 + i\sqrt{3}) (-x^6)^{2/3} + 6x^4 + i(\sqrt{3} + i) \sqrt[3]{-x^6} x^2}{8x^2}$$

$$y(x) \rightarrow \frac{1}{8} x^2 \left(\frac{(1 + i\sqrt{3}) x^4}{(-x^6)^{2/3}} + \frac{i(\sqrt{3} + i) x^2}{\sqrt[3]{-x^6}} + 6 \right)$$

1.22 problem 23

1.22.1 Solving as homogeneousTypeD2 ode	155
1.22.2 Solving as first order ode lie symmetry calculated ode	157
1.22.3 Solving as exact ode	162

Internal problem ID [7066]

Internal file name [OUTPUT/6052_Sunday_June_05_2022_04_15_33_PM_89745658/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 23.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactByInspection", "homogeneousTypeD2", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, [_Abel, `2nd type`, `class B`]]
```

$$x^2y' + y^2 - xyy' = 0$$

1.22.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$x^2(u'(x)x + u(x)) + u(x)^2x^2 - x^2u(x)(u'(x)x + u(x)) = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u}{x(u-1)}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = \frac{u}{u-1}$. Integrating both sides gives

$$\frac{1}{\frac{u}{u-1}} du = \frac{1}{x} dx$$

$$\int \frac{1}{\frac{u}{u-1}} du = \int \frac{1}{x} dx$$

$$u - \ln(u) = \ln(x) + c_2$$

The solution is

$$u(x) - \ln(u(x)) - \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\frac{y}{x} - \ln\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

$$\frac{y}{x} - \ln\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{y}{x} - \ln\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0 \tag{1}$$




Figure 31: Slope field plot

Verification of solutions

$$\frac{y}{x} - \ln\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

Verified OK.

1.22.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y^2}{x(y-x)}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{y^2(b_3 - a_2)}{x(y-x)} - \frac{y^4 a_3}{x^2(y-x)^2} - \left(-\frac{y^2}{x^2(y-x)} + \frac{y^2}{x(y-x)^2} \right) (xa_2 + ya_3 + a_1) \quad (\text{5E})$$
$$- \left(\frac{2y}{x(y-x)} - \frac{y^2}{x(y-x)^2} \right) (xb_2 + yb_3 + b_1) = 0$$

Putting the above in normal form gives

$$\frac{x^4 b_2 - x^2 y^2 a_2 + x^2 y^2 b_3 - 2x y^3 a_3 + 2x^2 y b_1 - 2x y^2 a_1 - x y^2 b_1 + y^3 a_1}{x^2 (x-y)^2} = 0$$

Setting the numerator to zero gives

$$x^4 b_2 - x^2 y^2 a_2 + x^2 y^2 b_3 - 2x y^3 a_3 + 2x^2 y b_1 - 2x y^2 a_1 - x y^2 b_1 + y^3 a_1 = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$-a_2 v_1^2 v_2^2 - 2a_3 v_1 v_2^3 + b_2 v_1^4 + b_3 v_1^2 v_2^2 - 2a_1 v_1 v_2^2 + a_1 v_2^3 + 2b_1 v_1^2 v_2 - b_1 v_1 v_2^2 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$b_2 v_1^4 + (b_3 - a_2) v_1^2 v_2^2 + 2b_1 v_1^2 v_2 - 2a_3 v_1 v_2^3 + (-2a_1 - b_1) v_1 v_2^2 + a_1 v_2^3 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ b_2 &= 0 \\ -2a_3 &= 0 \\ 2b_1 &= 0 \\ -2a_1 - b_1 &= 0 \\ b_3 - a_2 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}\xi &= x \\ \eta &= y\end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(\frac{y^2}{x(y-x)} \right) (x) \\ &= \frac{yx}{x-y} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{yx}{x-y}} dy\end{aligned}$$

Which results in

$$S = \ln(y) - \frac{y}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y^2}{x(y-x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{y}{x^2} \\ S_y &= \frac{x-y}{xy} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(y)x - y}{x} = c_1$$



Which simplifies to

$$\frac{\ln(y)x - y}{x} = c_1$$

Which gives

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y^2}{x(y-x)}$ 	$R = x$ $S = \frac{\ln(y) x - y}{x}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1} \quad (1)$$




Figure 32: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1}$$

Verified OK.

1.22.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(x^2 - xy) dy &= (-y^2) dx \\ (y^2) dx + (x^2 - xy) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= y^2 \\ N(x, y) &= x^2 - xy\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y^2) \\ &= 2y\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x^2 - xy) \\ &= 2x - y\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. By inspection $\frac{1}{x^2y}$ is an integrating factor. Therefore by multiplying $M = y^2$ and $N = -yx + x^2$ by this integrating factor the ode becomes exact. The new M, N are

$$M = \frac{y}{x^2}$$

$$N = \frac{-yx + x^2}{x^2y}$$

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\frac{\partial \phi}{\partial x} = M$$

$$\frac{\partial \phi}{\partial y} = N$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} \left(\frac{x^2 - xy}{x^2y}\right) dy &= \left(-\frac{y}{x^2}\right) dx \\ \left(\frac{y}{x^2}\right) dx + \left(\frac{x^2 - xy}{x^2y}\right) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= \frac{y}{x^2} \\ N(x, y) &= \frac{x^2 - xy}{x^2y} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{y}{x^2}\right) \\ &= \frac{1}{x^2} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{x^2 - xy}{x^2y}\right) \\ &= \frac{1}{x^2} \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{y}{x^2} dx \\ \phi &= -\frac{y}{x} + f(y) \end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -\frac{1}{x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{x^2 - xy}{x^2 y}$. Therefore equation (4) becomes

$$\frac{x^2 - xy}{x^2 y} = -\frac{1}{x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{y}\right) dy$$

$$f(y) = \ln(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \ln(y) - \frac{y}{x} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \ln(y) - \frac{y}{x}$$

The solution becomes

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1}$$

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1} \quad (1)$$




Figure 33: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```


✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x)+y(x)^2=x*y(x)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = -x \operatorname{LambertW}\left(-\frac{e^{-c_1}}{x}\right)$$

✓ Solution by Mathematica

Time used: 2.396 (sec). Leaf size: 25

```
DSolve[x^2*y'[x]+y[x]^2==x*y[x]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -xW\left(-\frac{e^{-c_1}}{x}\right)$$

$$y(x) \rightarrow 0$$

1.23 problem 24

1.23.1 Maple step by step solution 171

Internal problem ID [7067]

Internal file name [OUTPUT/6053_Sunday_June_05_2022_04_15_36_PM_49751359/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 24.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y - xy' - x^2y'^2 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{-\frac{1}{2} + \frac{\sqrt{1+4y}}{2}}{x} \quad (1)$$

$$y' = \frac{-\frac{1}{2} - \frac{\sqrt{1+4y}}{2}}{x} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{-\frac{1}{2} + \frac{\sqrt{1+4y}}{2}}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = -\frac{1}{2} + \frac{\sqrt{1+4y}}{2}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{-\frac{1}{2} + \frac{\sqrt{1+4y}}{2}} dy &= \frac{1}{x} dx \\ \int \frac{1}{-\frac{1}{2} + \frac{\sqrt{1+4y}}{2}} dy &= \int \frac{1}{x} dx \\ \sqrt{1+4y} + \frac{\ln(-1 + \sqrt{1+4y})}{2} - \frac{\ln(\sqrt{1+4y} + 1)}{2} + \frac{\ln(y)}{2} &= \ln(x) + c_1 \end{aligned}$$

The solution is

$$\sqrt{1+4y} + \frac{\ln(-1 + \sqrt{1+4y})}{2} - \frac{\ln(\sqrt{1+4y} + 1)}{2} + \frac{\ln(y)}{2} - \ln(x) - c_1 = 0$$

Summary

The solution(s) found are the following

$$\sqrt{1+4y} + \frac{\ln(-1 + \sqrt{1+4y})}{2} - \frac{\ln(\sqrt{1+4y} + 1)}{2} + \frac{\ln(y)}{2} - \ln(x) - c_1 = 0 \quad (1)$$

Verification of solutions

$$\sqrt{1+4y} + \frac{\ln(-1 + \sqrt{1+4y})}{2} - \frac{\ln(\sqrt{1+4y} + 1)}{2} + \frac{\ln(y)}{2} - \ln(x) - c_1 = 0$$

Verified OK.

Solving equation (2)

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{-\frac{1}{2} - \frac{\sqrt{1+4y}}{2}}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = -\frac{1}{2} - \frac{\sqrt{1+4y}}{2}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{-\frac{1}{2} - \frac{\sqrt{1+4y}}{2}} dy &= \frac{1}{x} dx \\ \int \frac{1}{-\frac{1}{2} - \frac{\sqrt{1+4y}}{2}} dy &= \int \frac{1}{x} dx \\ -\sqrt{1+4y} - \frac{\ln(-1 + \sqrt{1+4y})}{2} + \frac{\ln(\sqrt{1+4y} + 1)}{2} + \frac{\ln(y)}{2} &= \ln(x) + c_2 \end{aligned}$$

The solution is

$$-\sqrt{1+4y} - \frac{\ln(-1 + \sqrt{1+4y})}{2} + \frac{\ln(\sqrt{1+4y} + 1)}{2} + \frac{\ln(y)}{2} - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$-\sqrt{1+4y} - \frac{\ln(-1 + \sqrt{1+4y})}{2} + \frac{\ln(\sqrt{1+4y} + 1)}{2} + \frac{\ln(y)}{2} - \ln(x) - c_2 = 0 \quad (1)$$

Verification of solutions

$$-\sqrt{1+4y} - \frac{\ln(-1 + \sqrt{1+4y})}{2} + \frac{\ln(\sqrt{1+4y} + 1)}{2} + \frac{\ln(y)}{2} - \ln(x) - c_2 = 0$$

Verified OK.

1.23.1 Maple step by step solution

Let's solve

$$y - xy' - x^2y'^2 = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{-\frac{1}{2} + \frac{\sqrt{1+4y}}{2}} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{-\frac{1}{2} + \frac{\sqrt{1+4y}}{2}} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$\frac{\ln(y)}{2} + \sqrt{1+4y} + \frac{\ln(-1 + \sqrt{1+4y})}{2} - \frac{\ln(\sqrt{1+4y} + 1)}{2} = \ln(x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
trying dAlembert  
trying simple symmetries for implicit equations  
<- symmetries for implicit equations successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 97

```
dsolve(y(x)=x*diff(y(x),x)+x^2*diff(y(x),x)^2,y(x), singsol=all)
```

$$\begin{aligned} \ln(x) - \sqrt{4y(x)+1} - \frac{\ln(-1 + \sqrt{4y(x)+1})}{2} \\ + \frac{\ln(1 + \sqrt{4y(x)+1})}{2} - \frac{\ln(y(x))}{2} - c_1 = 0 \\ \ln(x) + \sqrt{4y(x)+1} + \frac{\ln(-1 + \sqrt{4y(x)+1})}{2} \\ - \frac{\ln(1 + \sqrt{4y(x)+1})}{2} - \frac{\ln(y(x))}{2} - c_1 = 0 \end{aligned}$$

✓ Solution by Mathematica

Time used: 22.779 (sec). Leaf size: 72

```
DSolve[y[x]==x*y'[x]+x^2*(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} y(x) &\rightarrow \frac{1}{4}W(-e^{-1-2c_1}x)(2+W(-e^{-1-2c_1}x)) \\ y(x) &\rightarrow \frac{1}{4}W(e^{-1+2c_1}x)(2+W(e^{-1+2c_1}x)) \\ y(x) &\rightarrow 0 \end{aligned}$$

1.24 problem 25

1.24.1 Solving as quadrature ode	173
1.24.2 Maple step by step solution	174

Internal problem ID [7068]

Internal file name [OUTPUT/6054_Sunday_June_05_2022_04_15_43_PM_10654432/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 25.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$(x + y) y' = 0$$

1.24.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int 0 \, dx \\ &= c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \tag{1}$$




Figure 34: Slope field plot

Verification of solutions

$$y = c_1$$

Verified OK.

1.24.2 Maple step by step solution

Let's solve

$$(x + y) y' = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int (x + y) y' dx = \int 0 dx + c_1$$

- Cannot compute integral

$$\int (x + y) y' dx = c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve((x+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x$$
$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 14

```
DSolve[(x+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x$$
$$y(x) \rightarrow c_1$$

1.25 problem 26

1.25.1 Solving as quadrature ode	176
1.25.2 Maple step by step solution	177

Internal problem ID [7069]

Internal file name [OUTPUT/6055_Sunday_June_05_2022_04_15_45_PM_52012227/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 26.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$xy' = 0$$

1.25.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned}y &= \int 0 \, dx \\ &= c_1\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \tag{1}$$




Figure 35: Slope field plot

Verification of solutions

$$y = c_1$$

Verified OK.

1.25.2 Maple step by step solution

Let's solve

$$xy' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int xy'dx = \int 0dx + c_1$$

- Cannot compute integral

$$\int xy'dx = c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.26 problem 27

1.26.1 Solving as quadrature ode	179
1.26.2 Maple step by step solution	180

Internal problem ID [7070]

Internal file name [OUTPUT/6056_Sunday_June_05_2022_04_15_47_PM_11684547/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 27.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$\frac{y'}{x+y} = 0$$

1.26.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned}y &= \int 0 \, dx \\ &= c_1\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \tag{1}$$




Figure 36: Slope field plot

Verification of solutions

$$y = c_1$$

Verified OK.

1.26.2 Maple step by step solution

Let's solve

$$\frac{y'}{x+y} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int \frac{y'}{x+y} dx = \int 0 dx + c_1$$

- Cannot compute integral

$$\int \frac{y'}{x+y} dx = c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(1/(x+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[1/(x+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.27 problem 28

1.27.1 Solving as quadrature ode	182
1.27.2 Maple step by step solution	183

Internal problem ID [7071]

Internal file name [OUTPUT/6057_Sunday_June_05_2022_04_15_49_PM_43898222/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 28.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$\boxed{\frac{y'}{x} = 0}$$

1.27.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned}y &= \int 0 \, dx \\ &= c_1\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \tag{1}$$




Figure 37: Slope field plot

Verification of solutions

$$y = c_1$$

Verified OK.

1.27.2 Maple step by step solution

Let's solve

$$\frac{y'}{x} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int \frac{y'}{x} dx = \int 0 dx + c_1$$

- Cannot compute integral

$$\int \frac{y'}{x} dx = c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(1/x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 7

```
DSolve[1/x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.28 problem 29

1.28.1 Solving as quadrature ode	185
1.28.2 Maple step by step solution	186

Internal problem ID [7072]

Internal file name [OUTPUT/6058_Sunday_June_05_2022_04_15_51_PM_23005375/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 29.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = 0$$

1.28.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int 0 \, dx \\ &= c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \tag{1}$$




Figure 38: Slope field plot

Verification of solutions

$$y = c_1$$

Verified OK.

1.28.2 Maple step by step solution

Let's solve

$$y' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int 0 dx + c_1$$

- Evaluate integral

$$y = c_1$$

- Solve for y

$$y = c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve(diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 7

```
DSolve[y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

1.29 problem 30

1.29.1 Solving as dAlembert ode 188

Internal problem ID [7073]

Internal file name [OUTPUT/6059_Sunday_June_05_2022_04_15_53_PM_89480368/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 30.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**dAlembert**"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, _dAlembert]
```

$$y - xy'^2 - y'^2 = 0$$

1.29.1 Solving as dAlembert ode

Let $p = y'$ the ode becomes

$$-xp^2 - p^2 + y = 0$$

Solving for y from the above results in

$$y = xp^2 + p^2 \tag{1A}$$

This has the form

$$y = xf(p) + g(p) \tag{*}$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved. Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g')\frac{dp}{dx} \\ p - f &= (xf' + g')\frac{dp}{dx} \end{aligned} \tag{2}$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned}f &= p^2 \\g &= p^2\end{aligned}$$

Hence (2) becomes

$$-p^2 + p = (2xp + 2p)p'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$-p^2 + p = 0$$

Solving for p from the above gives

$$\begin{aligned}p &= 0 \\p &= 1\end{aligned}$$

Substituting these in (1A) gives

$$\begin{aligned}y &= 0 \\y &= 1 + x\end{aligned}$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = \frac{-p(x)^2 + p(x)}{2p(x)x + 2p(x)} \quad (3)$$

This ODE is now solved for $p(x)$.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(x) + p(x)p(x) = q(x)$$

Where here

$$\begin{aligned}p(x) &= \frac{1}{2x + 2} \\q(x) &= \frac{1}{2x + 2}\end{aligned}$$

Hence the ode is

$$p'(x) + \frac{p(x)}{2x + 2} = \frac{1}{2x + 2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{2x+2} dx} \\ &= \sqrt{1+x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= (\mu) \left(\frac{1}{2x+2} \right) \\ \frac{d}{dx}(\sqrt{1+x} p) &= (\sqrt{1+x}) \left(\frac{1}{2x+2} \right) \\ d(\sqrt{1+x} p) &= \left(\frac{1}{2\sqrt{1+x}} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\sqrt{1+x} p &= \int \frac{1}{2\sqrt{1+x}} dx \\ \sqrt{1+x} p &= \sqrt{1+x} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \sqrt{1+x}$ results in

$$p(x) = 1 + \frac{c_1}{\sqrt{1+x}}$$

Substituting the above solution for p in (2A) gives

$$y = x \left(1 + \frac{c_1}{\sqrt{1+x}} \right)^2 + \left(1 + \frac{c_1}{\sqrt{1+x}} \right)^2$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$

$$y = 1 + x \tag{2}$$

$$y = x \left(1 + \frac{c_1}{\sqrt{1+x}} \right)^2 + \left(1 + \frac{c_1}{\sqrt{1+x}} \right)^2 \tag{3}$$

Verification of solutions

$$y = 0$$

Verified OK.

$$y = 1 + x$$

Verified OK.

$$y = x \left(1 + \frac{c_1}{\sqrt{1+x}} \right)^2 + \left(1 + \frac{c_1}{\sqrt{1+x}} \right)^2$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
*** Sublevel 2 ***  
Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
trying dAlembert  
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 53

```
dsolve(y(x)=x*diff(y(x),x)^2+diff(y(x),x)^2,y(x), singsol=all)
```

$$y(x) = 0$$
$$y(x) = \frac{\left(x + 1 + \sqrt{(x + 1)(c_1 + 1)} \right)^2}{x + 1}$$
$$y(x) = \frac{\left(-x - 1 + \sqrt{(x + 1)(c_1 + 1)} \right)^2}{x + 1}$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 57

```
DSolve[y[x]==x*(y'[x])^2+(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - c_1\sqrt{x+1} + 1 + \frac{c_1^2}{4}$$

$$y(x) \rightarrow x + c_1\sqrt{x+1} + 1 + \frac{c_1^2}{4}$$

$$y(x) \rightarrow 0$$

1.30 problem 31

1.30.1 Solving as homogeneousTypeD2 ode	193
1.30.2 Solving as first order ode lie symmetry calculated ode	195
1.30.3 Solving as riccati ode	201

Internal problem ID [7074]

Internal file name [OUTPUT/6060_Sunday_June_05_2022_04_16_29_PM_79850988/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 31.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**", "**homogeneousTypeD2**", "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Riccati]
```

$$y' - \frac{5x^2 - yx + y^2}{x^2} = 0$$

1.30.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - \frac{5x^2 - u(x)x^2 + u(x)^2x^2}{x^2} = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u^2 - 2u + 5}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u^2 - 2u + 5$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u^2 - 2u + 5} du &= \frac{1}{x} dx \\ \int \frac{1}{u^2 - 2u + 5} du &= \int \frac{1}{x} dx \\ \frac{\arctan\left(\frac{u}{2} - \frac{1}{2}\right)}{2} &= \ln(x) + c_2\end{aligned}$$

The solution is

$$\frac{\arctan\left(\frac{u(x)}{2} - \frac{1}{2}\right)}{2} - \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned}-\frac{\arctan\left(-\frac{y}{2x} + \frac{1}{2}\right)}{2} - \ln(x) - c_2 &= 0 \\ \frac{\arctan\left(\frac{y-x}{2x}\right)}{2} - \ln(x) - c_2 &= 0\end{aligned}$$

Summary

The solution(s) found are the following

$$\frac{\arctan\left(\frac{y-x}{2x}\right)}{2} - \ln(x) - c_2 = 0 \tag{1}$$




Figure 39: Slope field plot

Verification of solutions

$$\frac{\arctan\left(\frac{y-x}{2x}\right)}{2} - \ln(x) - c_2 = 0$$

Verified OK.

1.30.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{5x^2 - xy + y^2}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(5x^2 - xy + y^2)(b_3 - a_2)}{x^2} - \frac{(5x^2 - xy + y^2)^2 a_3}{x^4} \\ - \left(\frac{10x - y}{x^2} - \frac{2(5x^2 - xy + y^2)}{x^3} \right) (xa_2 + ya_3 + a_1) \\ - \frac{(-x + 2y)(xb_2 + yb_3 + b_1)}{x^2} = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{5x^4 a_2 + 25x^4 a_3 - 2b_2 x^4 - 5x^4 b_3 - 10x^3 y a_3 + 2x^3 y b_2 - x^2 y^2 a_2 + 12x^2 y^2 a_3 + x^2 y^2 b_3 - 4x y^3 a_3 + y^4 a_3 - x^3 b_1 - x^2 y a_1 - 2x^2 y b_1 + 2x y^2 a_1}{x^4} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -5x^4 a_2 - 25x^4 a_3 + 2b_2 x^4 + 5x^4 b_3 + 10x^3 y a_3 - 2x^3 y b_2 + x^2 y^2 a_2 - 12x^2 y^2 a_3 \\ - x^2 y^2 b_3 + 4x y^3 a_3 - y^4 a_3 + x^3 b_1 - x^2 y a_1 - 2x^2 y b_1 + 2x y^2 a_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -5a_2 v_1^4 + a_2 v_1^2 v_2^2 - 25a_3 v_1^4 + 10a_3 v_1^3 v_2 - 12a_3 v_1^2 v_2^2 + 4a_3 v_1 v_2^3 - a_3 v_2^4 + 2b_2 v_1^4 \\ - 2b_2 v_1^3 v_2 + 5b_3 v_1^4 - b_3 v_1^2 v_2^2 - a_1 v_1^2 v_2 + 2a_1 v_1 v_2^2 + b_1 v_1^3 - 2b_1 v_1^2 v_2 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} &(-5a_2 - 25a_3 + 2b_2 + 5b_3)v_1^4 + (10a_3 - 2b_2)v_1^3v_2 + b_1v_1^3 \\ &+ (a_2 - 12a_3 - b_3)v_1^2v_2^2 + (-a_1 - 2b_1)v_1^2v_2 + 4a_3v_1v_2^3 + 2a_1v_1v_2^2 - a_3v_2^4 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} b_1 &= 0 \\ 2a_1 &= 0 \\ -a_3 &= 0 \\ 4a_3 &= 0 \\ -a_1 - 2b_1 &= 0 \\ 10a_3 - 2b_2 &= 0 \\ a_2 - 12a_3 - b_3 &= 0 \\ -5a_2 - 25a_3 + 2b_2 + 5b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(\frac{5x^2 - xy + y^2}{x^2} \right) (x) \\ &= \frac{-5x^2 + 2xy - y^2}{x} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-5x^2 + 2xy - y^2}{x}} dy\end{aligned}$$

Which results in

$$S = -\frac{\arctan\left(\frac{2y-2x}{4x}\right)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{5x^2 - xy + y^2}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{y}{5x^2 - 2xy + y^2} \\ S_y &= -\frac{x}{5x^2 - 2xy + y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\ln(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\arctan\left(\frac{x-y}{2x}\right)}{2} = -\ln(x) + c_1$$



Which simplifies to

$$\frac{\arctan\left(\frac{x-y}{2x}\right)}{2} = -\ln(x) + c_1$$

Which gives

$$y = -2 \tan(-2 \ln(x) + 2c_1) x + x$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{5x^2 - xy + y^2}{x^2}$ 	$R = x$ $S = \frac{\arctan\left(\frac{x-y}{2x}\right)}{2}$	$\frac{dS}{dR} = -\frac{1}{R}$ 

Summary

The solution(s) found are the following

$$y = -2 \tan(-2 \ln(x) + 2c_1) x + x \quad (1)$$




Figure 40: Slope field plot

Verification of solutions

$$y = -2 \tan(-2 \ln(x) + 2c_1) x + x$$

Verified OK.

1.30.3 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{5x^2 - xy + y^2}{x^2} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = 5 - \frac{y}{x} + \frac{y^2}{x^2}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = 5$, $f_1(x) = -\frac{1}{x}$ and $f_2(x) = \frac{1}{x^2}$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{u}{x^2}} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \tag{2}$$

But

$$\begin{aligned} f_2' &= -\frac{2}{x^3} \\ f_1 f_2 &= -\frac{1}{x^3} \\ f_2^2 f_0 &= \frac{5}{x^4} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{u''(x)}{x^2} + \frac{3u'(x)}{x^3} + \frac{5u(x)}{x^4} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \frac{c_1 \sin(2 \ln(x)) + c_2 \cos(2 \ln(x))}{x}$$

The above shows that

$$u'(x) = \frac{(2c_1 - c_2) \cos(2 \ln(x)) - \sin(2 \ln(x)) (c_1 + 2c_2)}{x^2}$$

Using the above in (1) gives the solution

$$y = -\frac{((2c_1 - c_2) \cos(2 \ln(x)) - \sin(2 \ln(x)) (c_1 + 2c_2)) x}{c_1 \sin(2 \ln(x)) + c_2 \cos(2 \ln(x))}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = -\frac{((2c_3 - 1) \cos(2 \ln(x)) - \sin(2 \ln(x)) (c_3 + 2)) x}{c_3 \sin(2 \ln(x)) + \cos(2 \ln(x))}$$

Summary

The solution(s) found are the following

$$y = -\frac{((2c_3 - 1) \cos(2 \ln(x)) - \sin(2 \ln(x)) (c_3 + 2)) x}{c_3 \sin(2 \ln(x)) + \cos(2 \ln(x))} \quad (1)$$




Figure 41: Slope field plot

Verification of solutions

$$y = -\frac{((2c_3 - 1) \cos(2 \ln(x)) - \sin(2 \ln(x)) (c_3 + 2)) x}{c_3 \sin(2 \ln(x)) + \cos(2 \ln(x))}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=(5*x^2-x*y(x)+y(x)^2)/x^2,y(x), singsol=all)
```

$$y(x) = x(1 + 2 \tan(2 \ln(x) + 2c_1))$$

✓ Solution by Mathematica

Time used: 0.789 (sec). Leaf size: 18

```
DSolve[y'[x]==(5*x^2-x*y[x]+y[x]^2)/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + 2x \tan(2(\log(x) + c_1))$$

1.31 problem 32

- 1.31.1 Solving as homogeneousTypeMapleC ode 205
- 1.31.2 Solving as first order ode lie symmetry calculated ode 209

Internal problem ID [7075]

Internal file name [OUTPUT/6061_Sunday_June_05_2022_04_16_32_PM_51526962/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 32.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "homogeneousTypeMapleC",
"first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$3x + (x + 2)x' = -2t$$

1.31.1 Solving as homogeneousTypeMapleC ode

Let $Y = x + y_0$ and $X = t + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = -\frac{2X + 2x_0 + 3Y(X) + 3y_0}{Y(X) + y_0 + 2}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = 3$$

$$y_0 = -2$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = -\frac{2X + 3Y(X)}{Y(X)}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= -\frac{2X + 3Y}{Y} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = -2X - 3Y$ and $N = Y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= -\frac{2}{u} - 3 \\ \frac{du}{dX} &= \frac{-\frac{2}{u(X)} - 3 - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{-\frac{2}{u(X)} - 3 - u(X)}{X} = 0$$

Or

$$\left(\frac{d}{dX}u(X) \right) u(X) X + u(X)^2 + 3u(X) + 2 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{u^2 + 3u + 2}{uX} \end{aligned}$$

Where $f(X) = -\frac{1}{X}$ and $g(u) = \frac{u^2+3u+2}{u}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u^2+3u+2}{u}} du &= -\frac{1}{X} dX \\ \int \frac{1}{\frac{u^2+3u+2}{u}} du &= \int -\frac{1}{X} dX \\ -\ln(u+1) + 2\ln(u+2) &= -\ln(X) + c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{-\ln(u+1)+2\ln(u+2)} = e^{-\ln(X)+c_2}$$

Which simplifies to

$$\frac{(u+2)^2}{u+1} = \frac{c_3}{X}$$

The solution is

$$\frac{(u(X)+2)^2}{u(X)+1} = \frac{c_3}{X}$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$\frac{\left(\frac{Y(X)}{X} + 2\right)^2}{\frac{Y(X)}{X} + 1} = \frac{c_3}{X}$$

Which simplifies to

$$\frac{(Y(X) + 2X)^2}{Y(X) + X} = c_3$$

Using the solution for $Y(X)$

$$\frac{(Y(X) + 2X)^2}{Y(X) + X} = c_3$$

And replacing back terms in the above solution using

$$\begin{aligned}Y &= x + y_0 \\ X &= t + x_0\end{aligned}$$

Or

$$Y = x - 2$$

$$X = t + 3$$

Then the solution in x becomes

$$\frac{(x - 4 + 2t)^2}{x - 1 + t} = c_3$$

Summary

The solution(s) found are the following

$$\frac{(x - 4 + 2t)^2}{x - 1 + t} = c_3 \tag{1}$$




Figure 42: Slope field plot

Verification of solutions

$$\frac{(x - 4 + 2t)^2}{x - 1 + t} = c_3$$

Verified OK.

1.31.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$x' = -\frac{2t + 3x}{x + 2}$$

$$x' = \omega(t, x)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_t + \omega(\eta_x - \xi_t) - \omega^2 \xi_x - \omega_t \xi - \omega_x \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = ta_2 + xa_3 + a_1 \quad (\text{1E})$$

$$\eta = tb_2 + xb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 - \frac{(2t + 3x)(b_3 - a_2)}{x + 2} - \frac{(2t + 3x)^2 a_3}{(x + 2)^2} + \frac{2ta_2 + 2xa_3 + 2a_1}{x + 2} \quad (\text{5E})$$

$$- \left(-\frac{3}{x + 2} + \frac{2t + 3x}{(x + 2)^2} \right) (tb_2 + xb_3 + b_1) = 0$$

Putting the above in normal form gives

$$\frac{4t^2 a_3 + 2t^2 b_2 - 4txa_2 + 12txa_3 + 4txb_3 - 3x^2 a_2 + 7x^2 a_3 - x^2 b_2 + 3x^2 b_3 - 8ta_2 + 2tb_1 - 6tb_2 + 4tb_3 - 2a_1}{(x + 2)^2}$$

$$= 0$$

Setting the numerator to zero gives

$$-4t^2 a_3 - 2t^2 b_2 + 4txa_2 - 12txa_3 - 4txb_3 + 3x^2 a_2 - 7x^2 a_3 + x^2 b_2 - 3x^2 b_3 \quad (\text{6E})$$

$$+ 8ta_2 - 2tb_1 + 6tb_2 - 4tb_3 + 2xa_1 + 6xa_2 + 4xa_3 + 4xb_2 + 4a_1 + 6b_1 + 4b_2 = 0$$

Looking at the above PDE shows the following are all the terms with $\{t, x\}$ in them.

$$\{t, x\}$$

The following substitution is now made to be able to collect on all terms with $\{t, x\}$ in them

$$\{t = v_1, x = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} 4a_2v_1v_2 + 3a_2v_2^2 - 4a_3v_1^2 - 12a_3v_1v_2 - 7a_3v_2^2 - 2b_2v_1^2 + b_2v_2^2 \\ - 4b_3v_1v_2 - 3b_3v_2^2 + 2a_1v_2 + 8a_2v_1 + 6a_2v_2 + 4a_3v_2 \\ - 2b_1v_1 + 6b_2v_1 + 4b_2v_2 - 4b_3v_1 + 4a_1 + 6b_1 + 4b_2 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} (-4a_3 - 2b_2)v_1^2 + (4a_2 - 12a_3 - 4b_3)v_1v_2 + (8a_2 - 2b_1 + 6b_2 - 4b_3)v_1 \\ + (3a_2 - 7a_3 + b_2 - 3b_3)v_2^2 + (2a_1 + 6a_2 + 4a_3 + 4b_2)v_2 + 4a_1 + 6b_1 + 4b_2 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -4a_3 - 2b_2 &= 0 \\ 4a_1 + 6b_1 + 4b_2 &= 0 \\ 4a_2 - 12a_3 - 4b_3 &= 0 \\ 2a_1 + 6a_2 + 4a_3 + 4b_2 &= 0 \\ 3a_2 - 7a_3 + b_2 - 3b_3 &= 0 \\ 8a_2 - 2b_1 + 6b_2 - 4b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= -7a_3 - 3b_3 \\ a_2 &= 3a_3 + b_3 \\ a_3 &= a_3 \\ b_1 &= 6a_3 + 2b_3 \\ b_2 &= -2a_3 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\xi = t - 3$$

$$\eta = x + 2$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(t, x) \xi \\ &= x + 2 - \left(-\frac{2t + 3x}{x + 2} \right) (t - 3) \\ &= \frac{2t^2 + 3tx + x^2 - 6t - 5x + 4}{x + 2} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(t, x) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dt}{\xi} = \frac{dx}{\eta} = dS \quad (1)$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial t} + \eta \frac{\partial}{\partial x}) S(t, x) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = t$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{2t^2 + 3tx + x^2 - 6t - 5x + 4}{x + 2}} dy \end{aligned}$$

Which results in

$$S = -\ln(t + x - 1) + 2 \ln(2t + x - 4)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_t + \omega(t, x) S_x}{R_t + \omega(t, x) R_x} \quad (2)$$

Where in the above R_t, R_x, S_t, S_x are all partial derivatives and $\omega(t, x)$ is the right hand side of the original ode given by

$$\omega(t, x) = -\frac{2t + 3x}{x + 2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_t &= 1 \\ R_x &= 0 \\ S_t &= -\frac{1}{t + x - 1} + \frac{4}{2t + x - 4} \\ S_x &= \frac{x + 2}{(t + x - 1)(2t + x - 4)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for t, x in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to t, x coordinates. This results in

$$-\ln(x - 1 + t) + 2 \ln(x - 4 + 2t) = c_1$$

Which simplifies to

$$-\ln(x - 1 + t) + 2 \ln(x - 4 + 2t) = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in t, x coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dx}{dt} = -\frac{2t+3x}{x+2}$ 	$R = t$ $S = -\ln(t + x - 1) + 2t$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$-\ln(x - 1 + t) + 2 \ln(x - 4 + 2t) = c_1 \tag{1}$$




Figure 43: Slope field plot

Verification of solutions

$$-\ln(x - 1 + t) + 2 \ln(x - 4 + 2t) = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous C
trying homogeneous types:
trying homogeneous D
<- homogeneous successful
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 2.813 (sec). Leaf size: 30

`dsolve(2*t+3*x(t)+(x(t)+2)*diff(x(t),t)=0,x(t), singsol=all)`

$$x(t) = \frac{-\sqrt{4(t-3)c_1+1}-1+(-4t+8)c_1}{2c_1}$$

✓ Solution by Mathematica

Time used: 60.104 (sec). Leaf size: 1165

`DSolve[2*t+3*x[t]+(x[t]+2)*x'[t]==0,x[t],t,IncludeSingularSolutions -> True]`

$x(t) \rightarrow -2$

$$-\frac{t\sqrt{\frac{3}{(t-3)^2}-\frac{3(t-3)^2\cosh\left(\frac{4c_1}{9}\right)+3(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+2}{(t-3)^2\left((t-3)^2\cosh\left(\frac{4c_1}{9}\right)+(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+1\right)}-\sqrt{-\frac{\cosh\left(\frac{4c_1}{9}\right)+\sinh\left(\frac{4c_1}{9}\right)}{(t-3)^2\left((t-3)^2\cosh\left(\frac{4c_1}{9}\right)+(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+1\right)^2}}-3\sqrt{2(t-3)}}{2(t-3)}$$

$x(t) \rightarrow -2$

$$+\frac{t\sqrt{\frac{3}{(t-3)^2}-\frac{3(t-3)^2\cosh\left(\frac{4c_1}{9}\right)+3(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+2}{(t-3)^2\left((t-3)^2\cosh\left(\frac{4c_1}{9}\right)+(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+1\right)}-\sqrt{-\frac{\cosh\left(\frac{4c_1}{9}\right)+\sinh\left(\frac{4c_1}{9}\right)}{(t-3)^2\left((t-3)^2\cosh\left(\frac{4c_1}{9}\right)+(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+1\right)^2}}-3\sqrt{2(t-3)}}{2(t-3)}$$

$x(t) \rightarrow -2$

$$-\frac{t\sqrt{\frac{3}{(t-3)^2}-\frac{3(t-3)^2\cosh\left(\frac{4c_1}{9}\right)+3(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+2}{(t-3)^2\left((t-3)^2\cosh\left(\frac{4c_1}{9}\right)+(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+1\right)}+\sqrt{-\frac{\cosh\left(\frac{4c_1}{9}\right)+\sinh\left(\frac{4c_1}{9}\right)}{(t-3)^2\left((t-3)^2\cosh\left(\frac{4c_1}{9}\right)+(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+1\right)^2}}-3\sqrt{2(t-3)}}{2(t-3)}$$

$x(t) \rightarrow -2$

$$+\frac{t\sqrt{\frac{3}{(t-3)^2}-\frac{3(t-3)^2\cosh\left(\frac{4c_1}{9}\right)+3(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+2}{(t-3)^2\left((t-3)^2\cosh\left(\frac{4c_1}{9}\right)+(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+1\right)}+\sqrt{-\frac{\cosh\left(\frac{4c_1}{9}\right)+\sinh\left(\frac{4c_1}{9}\right)}{(t-3)^2\left((t-3)^2\cosh\left(\frac{4c_1}{9}\right)+(t-3)^2\sinh\left(\frac{4c_1}{9}\right)+1\right)^2}}-3\sqrt{2(t-3)}}{2(t-3)}$$

1.32 problem 33

1.32.1 Existence and uniqueness analysis	216
1.32.2 Solving as quadrature ode	217
1.32.3 Maple step by step solution	218

Internal problem ID [7076]

Internal file name [OUTPUT/6062_Sunday_June_05_2022_04_16_50_PM_96181182/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 33.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' - \frac{1}{1-y} = 0$$

With initial conditions

$$[y(0) = 2]$$

1.32.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(t, y) \\ &= -\frac{1}{-1+y} \end{aligned}$$

The y domain of $f(t, y)$ when $t = 0$ is

$$\{y < 1 \vee 1 < y\}$$

And the point $y_0 = 2$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{-1+y} \right) \\ &= \frac{1}{(-1+y)^2}\end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $t = 0$ is

$$\{y < 1 \vee 1 < y\}$$

And the point $y_0 = 2$ is inside this domain. Therefore solution exists and is unique.

1.32.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned}\int (1-y) dy &= t + c_1 \\ y - \frac{1}{2}y^2 &= t + c_1\end{aligned}$$

Solving for y gives these solutions

$$\begin{aligned}y_1 &= 1 - \sqrt{1 - 2c_1 - 2t} \\ y_2 &= 1 + \sqrt{1 - 2c_1 - 2t}\end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $t = 0$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 1 + \sqrt{1 - 2c_1}$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$y = 1 + \sqrt{1 - 2t}$$


Initial conditions are used to solve for c_1 . Substituting $t = 0$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 1 - \sqrt{1 - 2c_1}$$


Summary

Warning: Unable to solve for constant of integration. The solution(s) found are the following

$$y = 1 + \sqrt{1 - 2t}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 1 + \sqrt{1 - 2t}$$

Verified OK.

1.32.3 Maple step by step solution

Let's solve

$$\left[y' - \frac{1}{1-y} = 0, y(0) = 2 \right]$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$y'(1 - y) = 1$$

- Integrate both sides with respect to t

$$\int y'(1 - y) dt = \int 1 dt + c_1$$

- Evaluate integral

$$-\frac{y^2}{2} + y = t + c_1$$

- Solve for y
 $\{y = 1 - \sqrt{1 - 2c_1 - 2t}, y = 1 + \sqrt{1 - 2c_1 - 2t}\}$
- Use initial condition $y(0) = 2$
 $2 = 1 - \sqrt{1 - 2c_1}$
- Solution does not satisfy initial condition
- Use initial condition $y(0) = 2$
 $2 = 1 + \sqrt{1 - 2c_1}$
- Solve for c_1
 $c_1 = 0$
- Substitute $c_1 = 0$ into general solution and simplify
 $y = 1 + \sqrt{1 - 2t}$
- Solution to the IVP
 $y = 1 + \sqrt{1 - 2t}$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 13

```
dsolve([diff(y(t),t)=1/(1-y(t)),y(0) = 2],y(t), singsol=all)
```

$$y(t) = 1 + \sqrt{1 - 2t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 16

```
DSolve[{y'[t]==1/(1-y[t]),y[0]==2},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \sqrt{1 - 2t} + 1$$

1.33 problem 34

1.33.1 Existence and uniqueness analysis	221
1.33.2 Solving as quadrature ode	222
1.33.3 Maple step by step solution	223

Internal problem ID [7077]

Internal file name [OUTPUT/6063_Sunday_June_05_2022_04_16_52_PM_44126501/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 34.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$p' - ap + bp^2 = 0$$

With initial conditions

$$[p(t_0) = p_0]$$

1.33.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} p' &= f(t, p) \\ &= -bp^2 + ap \end{aligned}$$

The p domain of $f(t, p)$ when $t = t_0$ is

$$\{-\infty < p < \infty\}$$

But the point $p_0 = p_0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

1.33.2 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{-bp^2 + ap} dp = \int dt$$
$$\frac{\ln(p)}{a} - \frac{\ln(bp - a)}{a} = t + c_1$$

The above can be written as

$$\left(\frac{1}{a}\right) (\ln(p) - \ln(bp - a)) = t + c_1$$
$$\ln(p) - \ln(bp - a) = (a)(t + c_1)$$
$$= a(t + c_1)$$

Raising both side to exponential gives

$$e^{\ln(p) - \ln(bp - a)} = ac_1 e^{at}$$

Which simplifies to

$$-\frac{p}{-bp + a} = c_2 e^{at}$$

Initial conditions are used to solve for c_2 . Substituting $t = t_0$ and $p = p_0$ in the above solution gives an equation to solve for the constant of integration.

$$p_0 = \frac{e^{at_0} c_2 a}{e^{at_0} c_2 b - 1}$$

$$c_2 = -\frac{p_0 e^{-at_0}}{-b p_0 + a}$$

Substituting c_2 found above in the general solution gives

$$p = \frac{a p_0 e^{a(t-t_0)}}{b p_0 e^{a(t-t_0)} - b p_0 + a}$$

Summary

The solution(s) found are the following

$$p = \frac{a p_0 e^{a(t-t_0)}}{b p_0 e^{a(t-t_0)} - b p_0 + a} \quad (1)$$

Verification of solutions

$$p = \frac{a p_0 e^{a(t-t_0)}}{b p_0 e^{a(t-t_0)} - b p_0 + a}$$

Verified OK.

1.33.3 Maple step by step solution

Let's solve

$$[p' - ap + bp^2 = 0, p(t_0) = p_0]$$

- Highest derivative means the order of the ODE is 1

$$p'$$

- Separate variables

$$\frac{p'}{ap - bp^2} = 1$$

- Integrate both sides with respect to t

$$\int \frac{p'}{ap - bp^2} dt = \int 1 dt + c_1$$

- Evaluate integral

$$\frac{\ln(p)}{a} - \frac{\ln(bp - a)}{a} = t + c_1$$

- Solve for p

$$p = \frac{e^{c_1 a + at} a}{-1 + b e^{c_1 a + at}}$$

- Use initial condition $p(t_0) = p_0$

$$p_0 = \frac{e^{c_1 a + at_0} a}{-1 + b e^{c_1 a + at_0}}$$

- Solve for c_1

$$c_1 = \frac{-at_0 + \ln\left(-\frac{p_0}{-bp_0 + a}\right)}{a}$$

- Substitute $c_1 = \frac{-at_0 + \ln\left(-\frac{p_0}{-bp_0 + a}\right)}{a}$ into general solution and simplify

$$p = \frac{ap_0 e^{a(t-t_0)}}{bp_0 e^{a(t-t_0)} - bp_0 + a}$$

- Solution to the IVP

$$p = \frac{ap_0 e^{a(t-t_0)}}{bp_0 e^{a(t-t_0)} - bp_0 + a}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 29

```
dsolve([diff(p(t),t)=a*p(t)-b*p(t)^2,p(t0) = p0],p(t), singsol=all)
```

$$p(t) = \frac{a p_0}{(-p_0 b + a) e^{-a(t-t_0)} + p_0 b}$$

✓ Solution by Mathematica

Time used: 0.865 (sec). Leaf size: 39

```
DSolve[{p'[t]==a*p[t]-b*p[t]^2,p[t0]==p0},p[t],t,IncludeSingularSolutions -> True]
```

$$p(t) \rightarrow \frac{a p_0 e^{at}}{b p_0 (e^{at} - e^{at_0}) + a e^{at_0}}$$

1.34 problem 35

1.34.1 Solving as first order ode lie symmetry lookup ode	225
1.34.2 Solving as bernoulli ode	229
1.34.3 Solving as exact ode	233
1.34.4 Maple step by step solution	236

Internal problem ID [7078]

Internal file name [OUTPUT/6064_Sunday_June_05_2022_04_16_55_PM_47732975/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 35.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "bernoulli", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _exact, _rational, _Bernoulli]
```

$$y^2 + 2xyy' = -\frac{2}{x}$$

1.34.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{xy^2 + 2}{2yx^2}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type `Bernoulli`. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 43: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{xy}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{xy}} dy \end{aligned}$$

Which results in

$$S = \frac{x y^2}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{x y^2 + 2}{2y x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{y^2}{2} \\ S_y &= xy \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\ln(R) + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{xy^2}{2} = -\ln(x) + c_1$$

Which simplifies to

$$\frac{xy^2}{2} = -\ln(x) + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{xy^2+2}{2yx^2}$ 	$R = x$ $S = \frac{xy^2}{2}$	$\frac{dS}{dR} = -\frac{1}{R}$ 

Summary

The solution(s) found are the following

$$\frac{xy^2}{2} = -\ln(x) + c_1 \quad (1)$$




Figure 45: Slope field plot

Verification of solutions

$$\frac{xy^2}{2} = -\ln(x) + c_1$$

Verified OK.

1.34.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{xy^2 + 2}{2yx^2} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -\frac{1}{2x}y - \frac{1}{x^2}\frac{1}{y} \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= -\frac{1}{2x} \\ f_1(x) &= -\frac{1}{x^2} \\ n &= -1 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = \frac{1}{y}$ gives

$$y'y = -\frac{y^2}{2x} - \frac{1}{x^2} \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= y^2 \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = 2yy' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} \frac{w'(x)}{2} &= -\frac{w(x)}{2x} - \frac{1}{x^2} \\ w' &= -\frac{w}{x} - \frac{2}{x^2} \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned} p(x) &= \frac{1}{x} \\ q(x) &= -\frac{2}{x^2} \end{aligned}$$

Hence the ode is

$$w'(x) + \frac{w(x)}{x} = -\frac{2}{x^2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu) \left(-\frac{2}{x^2}\right) \\ \frac{d}{dx}(wx) &= (x) \left(-\frac{2}{x^2}\right) \\ d(wx) &= \left(-\frac{2}{x}\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}wx &= \int -\frac{2}{x} dx \\ wx &= -2 \ln(x) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$w(x) = -\frac{2 \ln(x)}{x} + \frac{c_1}{x}$$

which simplifies to

$$w(x) = \frac{-2 \ln(x) + c_1}{x}$$

Replacing w in the above by y^2 using equation (5) gives the final solution.

$$y^2 = \frac{-2 \ln(x) + c_1}{x}$$

Solving for y gives

$$\begin{aligned}y(x) &= \frac{\sqrt{x(-2 \ln(x) + c_1)}}{x} \\ y(x) &= -\frac{\sqrt{x(-2 \ln(x) + c_1)}}{x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\sqrt{x(-2\ln(x) + c_1)}}{x} \quad (1)$$

$$y = -\frac{\sqrt{x(-2\ln(x) + c_1)}}{x} \quad (2)$$




Figure 46: Slope field plot

Verification of solutions

$$y = \frac{\sqrt{x(-2\ln(x) + c_1)}}{x}$$

Verified OK.

$$y = -\frac{\sqrt{x(-2\ln(x) + c_1)}}{x}$$

Verified OK.

1.34.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (2xy) dy &= \left(-y^2 - \frac{2}{x}\right) dx \\ \left(y^2 + \frac{2}{x}\right) dx + (2xy) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$M(x, y) = y^2 + \frac{2}{x}$$
$$N(x, y) = 2xy$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y} \left(y^2 + \frac{2}{x} \right)$$
$$= 2y$$

And

$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x} (2xy)$$
$$= 2y$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$
$$\int \frac{\partial \phi}{\partial x} dx = \int y^2 + \frac{2}{x} dx$$
$$\phi = x y^2 + 2 \ln(x) + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 2xy + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 2xy$. Therefore equation (4) becomes

$$2xy = 2xy + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = x y^2 + 2 \ln(x) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x y^2 + 2 \ln(x)$$

Summary

The solution(s) found are the following

$$x y^2 + 2 \ln(x) = c_1 \tag{1}$$




Figure 47: Slope field plot

Verification of solutions

$$xy^2 + 2 \ln(x) = c_1$$

Verified OK.

1.34.4 Maple step by step solution

Let's solve

$$y^2 + 2xyy' = -\frac{2}{x}$$

- Highest derivative means the order of the ODE is 1
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 - $F'(x, y) = 0$
 - Compute derivative of lhs
 - $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$
 - Evaluate derivatives
 - $2y = 2y$
 - Condition met, ODE is exact
- Exact ODE implies solution will be of this form
- $\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x
- $F(x, y) = \int \left(y^2 + \frac{2}{x} \right) dx + f_1(y)$
- Evaluate integral
- $F(x, y) = x y^2 + 2 \ln(x) + f_1(y)$
- Take derivative of $F(x, y)$ with respect to y
- $N(x, y) = \frac{\partial}{\partial y} F(x, y)$
- Compute derivative
- $2xy = 2xy + \frac{d}{dy} f_1(y)$
- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 0$$

- Solve for $f_1(y)$

$$f_1(y) = 0$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = x y^2 + 2 \ln(x)$$

- Substitute $F(x, y)$ into the solution of the ODE

$$x y^2 + 2 \ln(x) = c_1$$

- Solve for y

$$\left\{ y = \frac{\sqrt{-x(2 \ln(x) - c_1)}}{x}, y = -\frac{\sqrt{-x(2 \ln(x) - c_1)}}{x} \right\}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 36

```
dsolve((y(x)^2+2/x)+2*y(x)*x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{x(-2 \ln(x) + c_1)}}{x}$$

$$y(x) = -\frac{\sqrt{x(-2 \ln(x) + c_1)}}{x}$$

✓ Solution by Mathematica

Time used: 0.207 (sec). Leaf size: 44

```
DSolve[(y[x]^2+2/x)+2*y[x]*x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-2\log(x) + c_1}}{\sqrt{x}}$$

$$y(x) \rightarrow \frac{\sqrt{-2\log(x) + c_1}}{\sqrt{x}}$$

1.35 problem 36

1.35.1 Solving as Clairaut ode 239

Internal problem ID [7079]

Internal file name [OUTPUT/6065_Sunday_June_05_2022_04_16_59_PM_89045391/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 36.

ODE order: 1.

ODE degree: 0.

The type(s) of ODE detected by this program : "**clairaut**"

Maple gives the following as the ode type

[_Clairaut]

$$xf' - f - \frac{f'^2(1 - f'^\lambda)^2}{\lambda^2} = 0$$

1.35.1 Solving as Clairaut ode

This is Clairaut ODE. It has the form

$$f = xf' + g(f')$$

Where g is function of $f'(x)$. Let $p = f'$ the ode becomes

$$xp - f - \frac{p^2(1 - p^\lambda)^2}{\lambda^2} = 0$$

Solving for f from the above results in

$$f = -\frac{p(p^{2\lambda}p - x\lambda^2 - 2p^\lambda p + p)}{\lambda^2} \tag{1A}$$

The above ode is a Clairaut ode which is now solved. We start by replacing f' by p which gives

$$\begin{aligned} f &= xp - \frac{p^2(p^{2\lambda} - 2p^\lambda + 1)}{\lambda^2} \\ &= xp - \frac{p^2(p^{2\lambda} - 2p^\lambda + 1)}{\lambda^2} \end{aligned}$$

Writing the ode as

$$f = xp + g(p)$$

We now write $g \equiv g(p)$ to make notation simpler but we should always remember that g is function of p which in turn is function of x . Hence the above becomes

$$f = xp + g \tag{1}$$

Then we see that

$$g = -\frac{p^2(p^{2\lambda} - 2p^\lambda + 1)}{\lambda^2}$$

Taking derivative of (1) w.r.t. x gives

$$\begin{aligned} p &= \frac{d}{dx}(xp + g) \\ p &= \left(p + x \frac{dp}{dx}\right) + \left(g' \frac{dp}{dx}\right) \\ p &= p + (x + g') \frac{dp}{dx} \\ 0 &= (x + g') \frac{dp}{dx} \end{aligned}$$

Where g' is derivative of $g(p)$ w.r.t. p . The general solution is given by

$$\begin{aligned} \frac{dp}{dx} &= 0 \\ p &= c_1 \end{aligned}$$

Substituting this in (1) gives the general solution as

$$f = c_1 x - \frac{c_1^2(c_1^{2\lambda} - 2c_1^\lambda + 1)}{\lambda^2}$$

The singular solution is found from solving for p from

$$x + g'(p) = 0$$

And substituting the result back in (1). Since we found above that $g = -\frac{p^2(p^{2\lambda} - 2p^\lambda + 1)}{\lambda^2}$, then the above equation becomes

$$\begin{aligned} x + g'(p) &= x - \frac{2p(p^{2\lambda} - 2p^\lambda + 1)}{\lambda^2} - \frac{p^2 \left(\frac{2p^{2\lambda}\lambda}{p} - \frac{2p^\lambda\lambda}{p} \right)}{\lambda^2} \\ &= 0 \end{aligned}$$

Solving the above for p results in

$$p_1 = \text{RootOf}(2_Z^{2\lambda} Z\lambda + 2_Z^{2\lambda} Z - 2_Z^\lambda Z\lambda - x\lambda^2 - 4_Z^\lambda Z + 2_Z)$$

Substituting the above back in (1) results in

$$f_1 = \frac{\text{RootOf}(2_Z^{1+2\lambda}\lambda + 2_Z^{1+2\lambda} - 2_Z^{\lambda+1}\lambda - x\lambda^2 - 4_Z^{\lambda+1} + 2_Z) x\lambda^2 + 2\text{RootOf}(2_Z^{1+2\lambda}\lambda + 2_Z)}{2}$$

Summary

The solution(s) found are the following

$$f = c_1 x - \frac{c_1^2(c_1^{2\lambda} - 2c_1^\lambda + 1)}{\lambda^2} \quad (1)$$

$$f = \frac{\text{RootOf}(2_Z^{1+2\lambda}\lambda + 2_Z^{1+2\lambda} - 2_Z^{\lambda+1}\lambda - x\lambda^2 - 4_Z^{\lambda+1} + 2_Z) x\lambda^2 + 2\text{RootOf}(2_Z^{1+2\lambda}\lambda + 2_Z)}{2} \quad (2)$$

Verification of solutions

$$f = c_1 x - \frac{c_1^2(c_1^{2\lambda} - 2c_1^\lambda + 1)}{\lambda^2}$$

Verified OK.

$$f = \frac{\text{RootOf}(2_Z^{1+2\lambda}\lambda + 2_Z^{1+2\lambda} - 2_Z^{\lambda+1}\lambda - x\lambda^2 - 4_Z^{\lambda+1} + 2_Z) x\lambda^2 + 2\text{RootOf}(2_Z^{1+2\lambda}\lambda + 2_Z)}{2}$$

Warning, solution could not be verified

Maple trace

```

`Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
trying dAlembert
<- 1st order, parametric methods successful
<- dAlembert successful`

```

✓ Solution by Maple

Time used: 0.468 (sec). Leaf size: 318

```
dsolve(x*diff(f(x),x)-f(x)=diff(f(x),x)^2/lambda^2*(1-diff(f(x),x)^lambda)^2,f(x), singsol=a
```

$$f(x) = 0$$

$$f(x)$$

$$= \frac{\lambda^2 x^2 \left(2\lambda e^{\text{RootOf}(2\lambda e^{-Z(2\lambda+1)} + 2e^{-Z(2\lambda+1)} - 2\lambda e^{-Z(\lambda+1)} - x\lambda^2 - 4e^{-Z(\lambda+1)} + 2e^{-Z})\lambda} + e^{\text{RootOf}(2\lambda e^{-Z(2\lambda+1)} + 2e^{-Z(2\lambda+1)} - 2\lambda e^{-Z(\lambda+1)} - x\lambda^2 - 4e^{-Z(\lambda+1)} + 2e^{-Z})\lambda} \right)}{4 \left(\lambda e^{\text{RootOf}(2\lambda e^{-Z(2\lambda+1)} + 2e^{-Z(2\lambda+1)} - 2\lambda e^{-Z(\lambda+1)} - x\lambda^2 - 4e^{-Z(\lambda+1)} + 2e^{-Z})\lambda} + e^{\text{RootOf}(2\lambda e^{-Z(2\lambda+1)} + 2e^{-Z(2\lambda+1)} - 2\lambda e^{-Z(\lambda+1)} - x\lambda^2 - 4e^{-Z(\lambda+1)} + 2e^{-Z})\lambda} \right)}$$

$$f(x) = c_1 x - \frac{c_1^2 (-1 + c_1^\lambda)^2}{\lambda^2}$$

✓ Solution by Mathematica

Time used: 15.811 (sec). Leaf size: 30

```
DSolve[x*f'[x]-f[x]==f'[x]^2/\[Lambda]^2*(1-f'[x]^\[Lambda])^2,f[x],x,IncludeSingularSolutio
```

$$f(x) \rightarrow c_1 \left(x - \frac{c_1 (-1 + c_1^\lambda)^2}{\lambda^2} \right)$$

$$f(x) \rightarrow 0$$

1.36 problem 37

1.36.1 Solving as riccati ode 243

Internal problem ID [7080]

Internal file name [OUTPUT/6066_Sunday_June_05_2022_04_17_16_PM_84389658/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 37.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**"

Maple gives the following as the ode type

`[_rational, _Riccati]`

$$xy' - 2y + by^2 = cx^4$$

1.36.1 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= -\frac{-cx^4 + by^2 - 2y}{x}\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = x^3c - \frac{by^2}{x} + \frac{2y}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x^3c$, $f_1(x) = \frac{2}{x}$ and $f_2(x) = -\frac{b}{x}$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2u} \\ &= \frac{-u'}{-\frac{bu}{x}}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= \frac{b}{x^2} \\ f_1 f_2 &= -\frac{2b}{x^2} \\ f_2^2 f_0 &= b^2 x c \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-\frac{b u''(x)}{x} + \frac{b u'(x)}{x^2} + b^2 x c u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 \sinh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) + c_2 \cosh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right)$$

The above shows that

$$u'(x) = x \sqrt{b} \sqrt{c} \left(c_1 \cosh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) + c_2 \sinh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) \right)$$

Using the above in (1) gives the solution

$$y = \frac{x^2 \sqrt{c} \left(c_1 \cosh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) + c_2 \sinh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) \right)}{\sqrt{b} \left(c_1 \sinh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) + c_2 \cosh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) \right)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{x^2 \sqrt{c} \left(c_3 \cosh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) + \sinh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) \right)}{\sqrt{b} \left(c_3 \sinh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) + \cosh\left(\frac{x^2 \sqrt{b} \sqrt{c}}{2}\right) \right)}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2\sqrt{c} \left(c_3 \cosh\left(\frac{x^2\sqrt{b}\sqrt{c}}{2}\right) + \sinh\left(\frac{x^2\sqrt{b}\sqrt{c}}{2}\right) \right)}{\sqrt{b} \left(c_3 \sinh\left(\frac{x^2\sqrt{b}\sqrt{c}}{2}\right) + \cosh\left(\frac{x^2\sqrt{b}\sqrt{c}}{2}\right) \right)} \quad (1)$$

Verification of solutions

$$y = \frac{x^2\sqrt{c} \left(c_3 \cosh\left(\frac{x^2\sqrt{b}\sqrt{c}}{2}\right) + \sinh\left(\frac{x^2\sqrt{b}\sqrt{c}}{2}\right) \right)}{\sqrt{b} \left(c_3 \sinh\left(\frac{x^2\sqrt{b}\sqrt{c}}{2}\right) + \cosh\left(\frac{x^2\sqrt{b}\sqrt{c}}{2}\right) \right)}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
<- Chini successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 31

```
dsolve(x*diff(y(x),x)-2*y(x)+b*y(x)^2=c*x^4,y(x), singsol=all)
```

$$y(x) = \frac{i \tan\left(-\frac{ix^2\sqrt{b}\sqrt{c}}{2} + c_1\right) x^2\sqrt{c}}{\sqrt{b}}$$

✓ Solution by Mathematica

Time used: 0.251 (sec). Leaf size: 153

```
DSolve[x*y'[x]-2*y[x]+b*y[x]^2==c*x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{cx^2} \left(-\cos\left(\frac{1}{2}\sqrt{-b}\sqrt{cx^2}\right) + c_1 \sin\left(\frac{1}{2}\sqrt{-b}\sqrt{cx^2}\right) \right)}{\sqrt{-b} \left(\sin\left(\frac{1}{2}\sqrt{-b}\sqrt{cx^2}\right) + c_1 \cos\left(\frac{1}{2}\sqrt{-b}\sqrt{cx^2}\right) \right)}$$

$$y(x) \rightarrow \frac{\sqrt{cx^2} \tan\left(\frac{1}{2}\sqrt{-b}\sqrt{cx^2}\right)}{\sqrt{-b}}$$

1.37 problem 38

1.37.1 Solving as riccati ode 247

Internal problem ID [7081]

Internal file name [OUTPUT/6067_Sunday_June_05_2022_04_17_18_PM_83377121/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 38.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**"

Maple gives the following as the ode type

`[_rational, _Riccati]`

$$xy' - y + y^2 = x^{\frac{2}{3}}$$

1.37.1 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y - y^2 + x^{\frac{2}{3}}}{x} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = \frac{1}{x^{\frac{1}{3}}} - \frac{y^2}{x} + \frac{y}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = \frac{1}{x^{\frac{1}{3}}}$, $f_1(x) = \frac{1}{x}$ and $f_2(x) = -\frac{1}{x}$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-\frac{u}{x}} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= \frac{1}{x^2} \\ f_1 f_2 &= -\frac{1}{x^2} \\ f_2^2 f_0 &= \frac{1}{x^{\frac{7}{3}}} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-\frac{u''(x)}{x} + \frac{u(x)}{x^{\frac{7}{3}}} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_2 e^{3x^{\frac{1}{3}}} \left(3x^{\frac{1}{3}} - 1 \right) + 3 e^{-3x^{\frac{1}{3}}} c_1 \left(x^{\frac{1}{3}} + \frac{1}{3} \right)$$

The above shows that

$$u'(x) = -\frac{3 \left(c_1 e^{-3x^{\frac{1}{3}}} - c_2 e^{3x^{\frac{1}{3}}} \right)}{x^{\frac{1}{3}}}$$

Using the above in (1) gives the solution

$$y = -\frac{3x^{\frac{2}{3}} \left(c_1 e^{-3x^{\frac{1}{3}}} - c_2 e^{3x^{\frac{1}{3}}} \right)}{c_2 e^{3x^{\frac{1}{3}}} \left(3x^{\frac{1}{3}} - 1 \right) + 3 e^{-3x^{\frac{1}{3}}} c_1 \left(x^{\frac{1}{3}} + \frac{1}{3} \right)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = -\frac{3x^{\frac{2}{3}} \left(-e^{6x^{\frac{1}{3}}} + c_3 \right)}{\left(3x^{\frac{1}{3}} - 1 \right) e^{6x^{\frac{1}{3}}} + 3x^{\frac{1}{3}} c_3 + c_3}$$

Summary

The solution(s) found are the following

$$y = -\frac{3x^{\frac{2}{3}}(-e^{6x^{\frac{1}{3}}} + c_3)}{(3x^{\frac{1}{3}} - 1)e^{6x^{\frac{1}{3}}} + 3x^{\frac{1}{3}}c_3 + c_3} \quad (1)$$




Figure 48: Slope field plot

Verification of solutions

$$y = -\frac{3x^{\frac{2}{3}}(-e^{6x^{\frac{1}{3}}} + c_3)}{(3x^{\frac{1}{3}} - 1)e^{6x^{\frac{1}{3}}} + 3x^{\frac{1}{3}}c_3 + c_3}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying Riccati
trying Riccati sub-methods:
  trying Riccati_symmetries
  trying Riccati to 2nd Order
  -> Calling odsolve with the ODE`, diff(diff(y(x), x), x) = y(x)/x^(4/3), y(x)`
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a symmetry of the form [xi=0, eta=F(x)]
  checking if the LODE is missing y
  -> Trying an equivalence, under non-integer power transformations,
  to LODEs admitting Liouvillian solutions.
  -> Trying a Liouvillian solution using Kovacics algorithm
  A Liouvillian solution exists
  Group is reducible or imprimitive
  <- Kovacics algorithm successful
  <- Equivalence, under non-integer power transformations successful
  <- Riccati to 2nd Order successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 72

```
dsolve(x*diff(y(x),x)-y(x)+y(x)^2=x^(2/3),y(x), singsol=all)
```

$$y(x) = \frac{x^{\frac{1}{3}} \left(c_1 e^{6x^{\frac{1}{3}}} \operatorname{abs} \left(1, 3x^{\frac{1}{3}} - 1 \right) + c_1 e^{6x^{\frac{1}{3}}} |3x^{\frac{1}{3}} - 1| - 3x^{\frac{1}{3}} \right)}{c_1 e^{6x^{\frac{1}{3}}} |3x^{\frac{1}{3}} - 1| + 3x^{\frac{1}{3}} + 1}$$

✓ Solution by Mathematica

Time used: 0.221 (sec). Leaf size: 131

```
DSolve[x*y'[x]-y[x]+y[x]^2==x^(2/3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x^{2/3}(c_1 \cosh(3\sqrt[3]{x}) - i \sinh(3\sqrt[3]{x}))}{(-3i\sqrt[3]{x} - c_1) \cosh(3\sqrt[3]{x}) + (3c_1\sqrt[3]{x} + i) \sinh(3\sqrt[3]{x})}$$

$$y(x) \rightarrow \frac{3x^{2/3} \cosh(3\sqrt[3]{x})}{3\sqrt[3]{x} \sinh(3\sqrt[3]{x}) - \cosh(3\sqrt[3]{x})}$$

1.38 problem 39

1.38.1 Solving as riccati ode 252

Internal problem ID [7082]

Internal file name [OUTPUT/6068_Sunday_June_05_2022_04_17_23_PM_32546509/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 39.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_rational, _Riccati]`

$$u' + u^2 = \frac{1}{x^{\frac{4}{5}}}$$

1.38.1 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= -\frac{u^2 x^{\frac{4}{5}} - 1}{x^{\frac{4}{5}}} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$u' = -u^2 + \frac{1}{x^{\frac{4}{5}}}$$

With Riccati ODE standard form

$$u' = f_0(x) + f_1(x)u + f_2(x)u^2$$

Shows that $f_0(x) = \frac{1}{x^{\frac{4}{5}}}$, $f_1(x) = 0$ and $f_2(x) = -1$. Let

$$\begin{aligned} u &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-u} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= \frac{1}{x^{\frac{4}{5}}} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-u''(x) + \frac{u(x)}{x^{\frac{4}{5}}} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \left(\text{BesselK} \left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_2 + \text{BesselI} \left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_1 \right) \sqrt{x}$$

The above shows that

$$u'(x) = \left(-\text{BesselK} \left(\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_2 + \text{BesselI} \left(-\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_1 \right) x^{\frac{1}{10}}$$

Using the above in (1) gives the solution

$$u = \frac{-\text{BesselK} \left(\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_2 + \text{BesselI} \left(-\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_1}{x^{\frac{2}{5}} \left(\text{BesselK} \left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_2 + \text{BesselI} \left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_1 \right)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$u = \frac{-\text{BesselK} \left(\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) + \text{BesselI} \left(-\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_3}{x^{\frac{2}{5}} \left(\text{BesselK} \left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) + \text{BesselI} \left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3} \right) c_3 \right)}$$

Summary

The solution(s) found are the following

$$u = \frac{-\text{BesselK}\left(\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3}\right) + \text{BesselI}\left(-\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3}\right) c_3}{x^{\frac{2}{5}} \left(\text{BesselK}\left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3}\right) + \text{BesselI}\left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3}\right) c_3\right)} \quad (1)$$




Figure 49: Slope field plot

Verification of solutions

$$u = \frac{-\text{BesselK}\left(\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3}\right) + \text{BesselI}\left(-\frac{1}{6}, \frac{5x^{\frac{3}{5}}}{3}\right) c_3}{x^{\frac{2}{5}} \left(\text{BesselK}\left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3}\right) + \text{BesselI}\left(\frac{5}{6}, \frac{5x^{\frac{3}{5}}}{3}\right) c_3\right)}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying Riccati
trying Riccati Special
<- Riccati Special successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 46

```
dsolve(diff(u(x),x)+u(x)^2=x^(-4/5),u(x), singsol=all)
```

$$u(x) = \frac{\text{BesselI}\left(-\frac{1}{6}, \frac{5x^{3/5}}{3}\right) c_1 - \text{BesselK}\left(\frac{1}{6}, \frac{5x^{3/5}}{3}\right)}{x^{2/5} \left(c_1 \text{BesselI}\left(\frac{5}{6}, \frac{5x^{3/5}}{3}\right) + \text{BesselK}\left(\frac{5}{6}, \frac{5x^{3/5}}{3}\right) \right)}$$

✓ Solution by Mathematica

Time used: 0.293 (sec). Leaf size: 286

```
DSolve[u'[x]+u[x]^2==x^(-4/5),u[x],x,IncludeSingularSolutions -> True]
```

$$u(x) \rightarrow \frac{(-1)^{5/6} x^{3/5} \text{Gamma}\left(\frac{11}{6}\right) \text{BesselI}\left(-\frac{1}{6}, \frac{5x^{3/5}}{3}\right) + (-1)^{5/6} \text{Gamma}\left(\frac{11}{6}\right) \text{BesselI}\left(\frac{5}{6}, \frac{5x^{3/5}}{3}\right) + (-1)^{5/6} x^{3/5} \text{Gamma}\left(\frac{11}{6}\right) \text{BesselK}\left(-\frac{1}{6}, \frac{5x^{3/5}}{3}\right) + (-1)^{5/6} x^{3/5} \text{Gamma}\left(\frac{11}{6}\right) \text{BesselK}\left(\frac{5}{6}, \frac{5x^{3/5}}{3}\right)}{2x \left((-1)^{5/6} \text{Gamma}\left(\frac{11}{6}\right) \text{BesselI}\left(\frac{5}{6}, \frac{5x^{3/5}}{3}\right) + (-1)^{5/6} \text{Gamma}\left(\frac{11}{6}\right) \text{BesselI}\left(-\frac{1}{6}, \frac{5x^{3/5}}{3}\right) + (-1)^{5/6} x^{3/5} \text{Gamma}\left(\frac{11}{6}\right) \text{BesselK}\left(-\frac{1}{6}, \frac{5x^{3/5}}{3}\right) + (-1)^{5/6} x^{3/5} \text{Gamma}\left(\frac{11}{6}\right) \text{BesselK}\left(\frac{5}{6}, \frac{5x^{3/5}}{3}\right) \right)}$$
$$u(x) \rightarrow \frac{x^{3/5} \text{BesselI}\left(-\frac{11}{6}, \frac{5x^{3/5}}{3}\right) + \text{BesselI}\left(-\frac{5}{6}, \frac{5x^{3/5}}{3}\right) + x^{3/5} \text{BesselI}\left(\frac{1}{6}, \frac{5x^{3/5}}{3}\right)}{2x \text{BesselI}\left(-\frac{5}{6}, \frac{5x^{3/5}}{3}\right)}$$

1.39 problem 40

- 1.39.1 Solving as homogeneousTypeD2 ode 256
1.39.2 Solving as first order ode lie symmetry calculated ode 258

Internal problem ID [7083]

Internal file name [OUTPUT/6069_Sunday_June_05_2022_04_17_26_PM_31417552/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 40.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**homogeneousTypeD2**", "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$yy' - y = x$$

1.39.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u(x)x(u'(x)x + u(x)) - u(x)x = x$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^2 - u - 1}{ux}\end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^2 - u - 1}{u}$. Integrating both sides gives

$$\frac{1}{\frac{u^2 - u - 1}{u}} du = -\frac{1}{x} dx$$

$$\int \frac{1}{\frac{u^2-u-1}{u}} du = \int -\frac{1}{x} dx$$

$$\frac{\ln(u^2 - u - 1)}{2} - \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(2u-1)\sqrt{5}}{5}\right)}{5} = -\ln(x) + c_2$$

The solution is

$$\frac{\ln(u(x)^2 - u(x) - 1)}{2} - \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(2u(x)-1)\sqrt{5}}{5}\right)}{5} + \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned} \frac{\ln\left(\frac{y^2}{x^2} - \frac{y}{x} - 1\right)}{2} - \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{\left(\frac{2y}{x}-1\right)\sqrt{5}}{5}\right)}{5} + \ln(x) - c_2 &= 0 \\ \frac{\ln\left(\frac{y^2}{x^2} - \frac{y}{x} - 1\right)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(x-2y)\sqrt{5}}{5x}\right)}{5} + \ln(x) - c_2 &= 0 \end{aligned}$$

Summary

The solution(s) found are the following

$$\frac{\ln\left(\frac{y^2}{x^2} - \frac{y}{x} - 1\right)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(x-2y)\sqrt{5}}{5x}\right)}{5} + \ln(x) - c_2 = 0 \quad (1)$$




Figure 50: Slope field plot

Verification of solutions

$$\frac{\ln\left(\frac{y^2}{x^2} - \frac{y}{x} - 1\right)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(x-2y)\sqrt{5}}{5x}\right)}{5} + \ln(x) - c_2 = 0$$

Verified OK.

1.39.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{x+y}{y}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{(x+y)(b_3 - a_2)}{y} - \frac{(x+y)^2 a_3}{y^2} - \frac{xa_2 + ya_3 + a_1}{y} - \left(\frac{1}{y} - \frac{x+y}{y^2}\right)(xb_2 + yb_3 + b_1) = 0 \quad (5E)$$

Putting the above in normal form gives

$$\frac{-x^2 a_3 - x^2 b_2 + 2xy a_2 + 2xy a_3 - 2xy b_3 + y^2 a_2 + 2y^2 a_3 - b_2 y^2 - y^2 b_3 - x b_1 + y a_1}{y^2} = 0$$

Setting the numerator to zero gives

$$-x^2 a_3 + x^2 b_2 - 2xy a_2 - 2xy a_3 + 2xy b_3 - y^2 a_2 - 2y^2 a_3 + b_2 y^2 + y^2 b_3 + x b_1 - y a_1 = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$-2a_2 v_1 v_2 - a_2 v_2^2 - a_3 v_1^2 - 2a_3 v_1 v_2 - 2a_3 v_2^2 + b_2 v_1^2 + b_2 v_2^2 + 2b_3 v_1 v_2 + b_3 v_2^2 - a_1 v_2 + b_1 v_1 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$(-a_3 + b_2) v_1^2 + (-2a_2 - 2a_3 + 2b_3) v_1 v_2 + b_1 v_1 + (-a_2 - 2a_3 + b_2 + b_3) v_2^2 - a_1 v_2 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 b_1 &= 0 \\
 -a_1 &= 0 \\
 -a_3 + b_2 &= 0 \\
 -2a_2 - 2a_3 + 2b_3 &= 0 \\
 -a_2 - 2a_3 + b_2 + b_3 &= 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= 0 \\
 a_2 &= -b_2 + b_3 \\
 a_3 &= b_2 \\
 b_1 &= 0 \\
 b_2 &= b_2 \\
 b_3 &= b_3
 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= x \\
 \eta &= y
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= y - \left(\frac{x + y}{y} \right) (x) \\
 &= \frac{-x^2 - xy + y^2}{y} \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x^2 - xy + y^2}{y}} dy \end{aligned}$$

Which results in

$$S = \frac{\ln(-x^2 - xy + y^2)}{2} - \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(-x+2y)\sqrt{5}}{5x}\right)}{5}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x + y}{y}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{x + y}{x^2 + xy - y^2} \\ S_y &= -\frac{y}{x^2 + xy - y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(y^2 - yx - x^2)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(x-2y)\sqrt{5}}{5x}\right)}{5} = c_1$$

Which simplifies to

$$\frac{\ln(y^2 - yx - x^2)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(x-2y)\sqrt{5}}{5x}\right)}{5} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x+y}{y}$ 	$R = x$ $S = \frac{\ln(-x^2 - xy + y^2)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{\ln(y^2 - yx - x^2)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(x-2y)\sqrt{5}}{5x}\right)}{5} = c_1 \quad (1)$$




Figure 51: Slope field plot

Verification of solutions

$$\frac{\ln(y^2 - yx - x^2)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(x-2y)\sqrt{5}}{5x}\right)}{5} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.359 (sec). Leaf size: 53

```
dsolve(y(x)*diff(y(x),x)-y(x)=x,y(x), singsol=all)
```

$$-\frac{\ln\left(\frac{-x^2-xy(x)+y(x)^2}{x^2}\right)}{2} - \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(-2y(x)+x)\sqrt{5}}{5x}\right)}{5} - \ln(x) - c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 63

```
DSolve[y[x]*y'[x] - y[x] == x,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[\frac{1}{10}\left(\left(5+\sqrt{5}\right)\log\left(-\frac{2y(x)}{x}+\sqrt{5}+1\right)-\left(\sqrt{5}-5\right)\log\left(\frac{2y(x)}{x}+\sqrt{5}-1\right)\right)=-\log(x)+c_1,y(x)\right]$$

1.40 problem 41

1.40.1 Solving as second order linear constant coeff ode	265
1.40.2 Solving as linear second order ode solved by an integrating factor ode	267
1.40.3 Solving using Kovacic algorithm	268
1.40.4 Maple step by step solution	272

Internal problem ID [7084]

Internal file name [OUTPUT/6070_Sunday_June_05_2022_04_17_31_PM_81036611/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 41.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"**

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + 2y' + y = 0$$

1.40.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(2)^2 - (4)(1)(1)} \\ &= -1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 1$. Therefore the solution is

$$y = c_1 e^{-x} + c_2 x e^{-x} \tag{1}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 x e^{-x} \tag{1}$$




Figure 52: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + c_2 x e^{-x}$$

Verified OK.

1.40.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = 2$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int 2 dx} \\ &= e^x\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= 0 \\ (e^x y)'' &= 0\end{aligned}$$

Integrating once gives

$$(e^x y)' = c_1$$

Integrating again gives

$$(e^x y) = c_1 x + c_2$$

Hence the solution is

$$y = \frac{c_1 x + c_2}{e^x}$$

Or

$$y = c_1 x e^{-x} + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = c_1 x e^{-x} + c_2 e^{-x} \quad (1)$$




Figure 53: Slope field plot

Verification of solutions

$$y = c_1 x e^{-x} + c_2 e^{-x}$$

Verified OK.

1.40.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 2 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 46: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \\ &= z_1 e^{-x} \\ &= z_1 (e^{-x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(x))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 x e^{-x} \quad (1)$$




Figure 54: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + c_2 x e^{-x}$$

Verified OK.

1.40.4 Maple step by step solution

Let's solve

$$y'' + 2y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r + 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = -1$$

- 1st solution of the ODE

$$y_1(x) = e^{-x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{-x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-x} + c_2 x e^{-x}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 18

```
DSolve[y''[x]+2*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_2x + c_1)$$

1.41 problem 41

1.41.1 Existence and uniqueness analysis	274
1.41.2 Solving as second order linear constant coeff ode	275
1.41.3 Solving using Kovacic algorithm	278
1.41.4 Maple step by step solution	282

Internal problem ID [7085]

Internal file name [OUTPUT/6071_Sunday_June_05_2022_04_17_33_PM_53577441/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 41.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$5y'' + 2y' + 4y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 5]$$

1.41.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = \frac{2}{5}$$
$$q(x) = \frac{4}{5}$$
$$F = 0$$

Hence the ode is

$$y'' + \frac{2y'}{5} + \frac{4y}{5} = 0$$

The domain of $p(x) = \frac{2}{5}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = \frac{4}{5}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

1.41.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 5, B = 2, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$5\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + 4e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$5\lambda^2 + 2\lambda + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 5, B = 2, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(5)} \pm \frac{1}{(2)(5)} \sqrt{2^2 - (4)(5)(4)} \\ &= -\frac{1}{5} \pm \frac{i\sqrt{19}}{5} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{5} + \frac{i\sqrt{19}}{5} \\ \lambda_2 &= -\frac{1}{5} - \frac{i\sqrt{19}}{5} \end{aligned}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{5} + \frac{i\sqrt{19}}{5}$$

$$\lambda_2 = -\frac{1}{5} - \frac{i\sqrt{19}}{5}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{5}$ and $\beta = \frac{\sqrt{19}}{5}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{5}} \left(c_1 \cos \left(\frac{\sqrt{19}x}{5} \right) + c_2 \sin \left(\frac{\sqrt{19}x}{5} \right) \right)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-\frac{x}{5}} \left(c_1 \cos \left(\frac{\sqrt{19}x}{5} \right) + c_2 \sin \left(\frac{\sqrt{19}x}{5} \right) \right) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{e^{-\frac{x}{5}} \left(c_1 \cos \left(\frac{\sqrt{19}x}{5} \right) + c_2 \sin \left(\frac{\sqrt{19}x}{5} \right) \right)}{5} + e^{-\frac{x}{5}} \left(-\frac{c_1 \sqrt{19} \sin \left(\frac{\sqrt{19}x}{5} \right)}{5} + \frac{c_2 \sqrt{19} \cos \left(\frac{\sqrt{19}x}{5} \right)}{5} \right)$$

substituting $y' = 5$ and $x = 0$ in the above gives

$$5 = -\frac{c_1}{5} + \frac{\sqrt{19}c_2}{5} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = \frac{25\sqrt{19}}{19}$$


Substituting these values back in above solution results in

$$y = \frac{25\sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19}$$


Summary

The solution(s) found are the following

$$y = \frac{25\sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{25\sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19}$$

Verified OK.

1.41.3 Solving using Kovacic algorithm

Writing the ode as

$$5y'' + 2y' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 5$$

$$B = 2 \quad (3)$$

$$C = 4$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-19}{25} \quad (6)$$

Comparing the above to (5) shows that

$$s = -19$$

$$t = 25$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{19z(x)}{25} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 48: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{19}{25}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{19}x}{5}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{2}{5} dx} \\&= z_1 e^{-\frac{x}{5}} \\&= z_1 \left(e^{-\frac{x}{5}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{5}} \cos \left(\frac{\sqrt{19}x}{5} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{2}{5} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\frac{2x}{5}}}{(y_1)^2} dx \\&= y_1 \left(\frac{5\sqrt{19} \tan \left(\frac{\sqrt{19}x}{5} \right)}{19} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{x}{5}} \cos \left(\frac{\sqrt{19}x}{5} \right) \right) + c_2 \left(e^{-\frac{x}{5}} \cos \left(\frac{\sqrt{19}x}{5} \right) \left(\frac{5\sqrt{19} \tan \left(\frac{\sqrt{19}x}{5} \right)}{19} \right) \right)\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{x}{5}} \cos\left(\frac{\sqrt{19}x}{5}\right) + \frac{5c_2 \sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{c_1 e^{-\frac{x}{5}} \cos\left(\frac{\sqrt{19}x}{5}\right)}{5} - \frac{c_1 e^{-\frac{x}{5}} \sqrt{19} \sin\left(\frac{\sqrt{19}x}{5}\right)}{5} - \frac{c_2 \sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19} + c_2 e^{-\frac{x}{5}} \cos\left(\frac{\sqrt{19}x}{5}\right)$$

substituting $y' = 5$ and $x = 0$ in the above gives

$$5 = -\frac{c_1}{5} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 5$$


Substituting these values back in above solution results in

$$y = \frac{25\sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19}$$


Summary

The solution(s) found are the following

$$y = \frac{25\sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{25\sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19}$$

Verified OK.

1.41.4 Maple step by step solution

Let's solve

$$\left[5y'' + 2y' + 4y = 0, y(0) = 0, y'|_{\{x=0\}} = 5 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y'}{5} - \frac{4y}{5}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y'}{5} + \frac{4y}{5} = 0$$

- Characteristic polynomial of ODE

$$r^2 + \frac{2}{5}r + \frac{4}{5} = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-\frac{2}{5}) \pm \left(\sqrt{-\frac{76}{25}}\right)}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{5} - \frac{i\sqrt{19}}{5}, -\frac{1}{5} + \frac{i\sqrt{19}}{5}\right)$$

- 1st solution of the ODE

$$y_1(x) = e^{-\frac{x}{5}} \cos\left(\frac{\sqrt{19}x}{5}\right)$$

- 2nd solution of the ODE

$$y_2(x) = e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-\frac{x}{5}} \cos\left(\frac{\sqrt{19}x}{5}\right) + c_2 \sin\left(\frac{\sqrt{19}x}{5}\right) e^{-\frac{x}{5}}$$

- Check validity of solution $y = c_1 e^{-\frac{x}{5}} \cos\left(\frac{\sqrt{19}x}{5}\right) + c_2 \sin\left(\frac{\sqrt{19}x}{5}\right) e^{-\frac{x}{5}}$

- Use initial condition $y(0) = 0$

$$0 = c_1$$

- Compute derivative of the solution

$$y' = -\frac{c_1 e^{-\frac{x}{5}} \cos\left(\frac{\sqrt{19}x}{5}\right)}{5} - \frac{c_1 e^{-\frac{x}{5}} \sqrt{19} \sin\left(\frac{\sqrt{19}x}{5}\right)}{5} + \frac{c_2 \sqrt{19} \cos\left(\frac{\sqrt{19}x}{5}\right) e^{-\frac{x}{5}}}{5} - \frac{c_2 \sin\left(\frac{\sqrt{19}x}{5}\right) e^{-\frac{x}{5}}}{5}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 5$

$$5 = -\frac{c_1}{5} + \frac{\sqrt{19}c_2}{5}$$

- Solve for c_1 and c_2

$$\left\{c_1 = 0, c_2 = \frac{25\sqrt{19}}{19}\right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{25\sqrt{19}e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19}$$

- Solution to the IVP

$$y = \frac{25\sqrt{19}e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 20

```
dsolve([5*diff(y(x),x$2)+2*diff(y(x),x)+4*y(x)=0,y(0) = 0, D(y)(0) = 5],y(x), singsol=all)
```

$$y(x) = \frac{25\sqrt{19} e^{-\frac{x}{5}} \sin\left(\frac{\sqrt{19}x}{5}\right)}{19}$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 6

```
DSolve[{5*y''[x]+2*y'[x]+4*y[x]==0,{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

1.42 problem 42

1.42.1 Solving as second order linear constant coeff ode	285
1.42.2 Solving using Kovacic algorithm	289
1.42.3 Maple step by step solution	294

Internal problem ID [7086]

Internal file name [OUTPUT/6072_Sunday_June_05_2022_04_17_36_PM_92423913/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 42.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y' + 4y = 1$$

1.42.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 4, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(4)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{15}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{15}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{15}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{15}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{15}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{15}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{15} x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{15} x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right) \right) + \left(\frac{1}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right) + \frac{1}{4} \quad (1)$$




Figure 57: Slope field plot

Verification of solutions

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right) + \frac{1}{4}$$

Verified OK.

1.42.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \quad (3)$$

$$C = 4$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-15}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = -15$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{15z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 50: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{15}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{15}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{15} x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{15} \tan \left(\frac{\sqrt{15} x}{2} \right)}{15} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{15} x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{15} x}{2} \right) \left(\frac{2\sqrt{15} \tan \left(\frac{\sqrt{15} x}{2} \right)}{15} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{2c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right), \frac{2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{2c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15} \right) + \left(\frac{1}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{2c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15} + \frac{1}{4} \quad (1)$$




Figure 58: Slope field plot

Verification of solutions

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{2c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15} + \frac{1}{4}$$

Verified OK.

1.42.3 Maple step by step solution

Let's solve

$$y'' + y' + 4y = 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-15})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{15}}{2}, -\frac{1}{2} + \frac{i\sqrt{15}}{2}\right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \sqrt{15} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{15}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2e^{-\frac{x}{2}}\sqrt{15}\left(\cos\left(\frac{\sqrt{15}x}{2}\right)\left(\int e^{\frac{x}{2}}\sin\left(\frac{\sqrt{15}x}{2}\right)dx\right) - \sin\left(\frac{\sqrt{15}x}{2}\right)\left(\int e^{\frac{x}{2}}\cos\left(\frac{\sqrt{15}x}{2}\right)dx\right)\right)}{15}$$

- Compute integrals

$$y_p(x) = \frac{1}{4}$$

- Substitute particular solution into general solution to ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) c_2 + \frac{1}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)+diff(y(x),x)+4*y(x)=1,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) c_2 + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{1}{4}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 51

```
DSolve[y''[x]+y'[x]+4*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{-x/2} \cos\left(\frac{\sqrt{15}x}{2}\right) + c_1 e^{-x/2} \sin\left(\frac{\sqrt{15}x}{2}\right) + \frac{1}{4}$$

1.43 problem 43

1.43.1 Solving as second order linear constant coeff ode	297
1.43.2 Solving using Kovacic algorithm	301
1.43.3 Maple step by step solution	306

Internal problem ID [7087]

Internal file name [OUTPUT/6073_Sunday_June_05_2022_04_17_39_PM_84374832/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 43.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y' + 4y = \sin(x)$$

1.43.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 4, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(4)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{15}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{15}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{15}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{15}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{15}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{15}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{15} x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{15} x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) - A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{10}, A_2 = \frac{3}{10} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{10} + \frac{3 \sin(x)}{10}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right) \right) + \left(-\frac{\cos(x)}{10} + \frac{3 \sin(x)}{10} \right)$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right) - \frac{\cos(x)}{10} + \frac{3 \sin(x)}{10} \quad (1)$$




Figure 59: Slope field plot

Verification of solutions

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right) - \frac{\cos(x)}{10} + \frac{3 \sin(x)}{10}$$

Verified OK.

1.43.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \quad (3)$$

$$C = 4$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-15}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = -15$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{15z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 52: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{15}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{15}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\&= z_1 e^{-\frac{x}{2}} \\&= z_1 \left(e^{-\frac{x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{15} \tan\left(\frac{\sqrt{15}x}{2}\right)}{15} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) \left(\frac{2\sqrt{15} \tan\left(\frac{\sqrt{15}x}{2}\right)}{15} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{2c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right), \frac{2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) - A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{10}, A_2 = \frac{3}{10} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{10} + \frac{3 \sin(x)}{10}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{2c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15} \right) + \left(-\frac{\cos(x)}{10} + \frac{3 \sin(x)}{10} \right)$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{2c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15} - \frac{\cos(x)}{10} + \frac{3 \sin(x)}{10} \quad (1)$$




Figure 60: Slope field plot

Verification of solutions

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{2c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{15} - \frac{\cos(x)}{10} + \frac{3 \sin(x)}{10}$$

Verified OK.

1.43.3 Maple step by step solution

Let's solve

$$y'' + y' + 4y = \sin(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-15})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{15}}{2}, -\frac{1}{2} + \frac{i\sqrt{15}}{2}\right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{15}x}{2}\right) e^{-\frac{x}{2}} \sqrt{15}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \sqrt{15} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{15}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{15}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{15}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin(x) \sin\left(\frac{\sqrt{15}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin(x) \cos\left(\frac{\sqrt{15}x}{2}\right) dx \right) \right)}{15}$$

- Compute integrals

$$y_p(x) = -\frac{\cos(x)}{10} + \frac{3\sin(x)}{10}$$

- Substitute particular solution into general solution to ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) c_2 + \frac{3\sin(x)}{10} - \frac{\cos(x)}{10}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 39

```
dsolve(diff(y(x),x$2)+diff(y(x),x)+4*y(x)=sin(x),y(x), singsol=all)
```

$$y(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) c_2 + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) c_1 + \frac{3\sin(x)}{10} - \frac{\cos(x)}{10}$$

✓ Solution by Mathematica

Time used: 1.949 (sec). Leaf size: 60

```
DSolve[y''[x]+y'[x]+4*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3 \sin(x)}{10} - \frac{\cos(x)}{10} + c_2 e^{-x/2} \cos\left(\frac{\sqrt{15}x}{2}\right) + c_1 e^{-x/2} \sin\left(\frac{\sqrt{15}x}{2}\right)$$

1.44 problem 44

- 1.44.1 Solving as first order nonlinear p but separable ode 309
- 1.44.2 Solving as dAlembert ode 311

Internal problem ID [7088]

Internal file name [OUTPUT/6074_Sunday_June_05_2022_04_17_42_PM_34052322/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 44.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "dAlembert", "first_order_non-linear_p_but_separable"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _dAlembert]
```

$$y - xy'^2 = 0$$

1.44.1 Solving as first order nonlinear p but separable ode

The ode has the form

$$(y')^{\frac{n}{m}} = f(x)g(y) \tag{1}$$

Where $n = 2, m = 1, f = \frac{1}{x}, g = y$. Hence the ode is

$$(y')^2 = \frac{y}{x}$$

Solving for y' from (1) gives

$$y' = \sqrt{fg}$$
$$y' = -\sqrt{fg}$$

To be able to solve as separable ode, we have to now assume that $f > 0, g > 0$.

$$\frac{1}{x} > 0$$
$$y > 0$$

Under the above assumption the differential equations become separable and can be written as

$$y' = \sqrt{f} \sqrt{g}$$

$$y' = -\sqrt{f} \sqrt{g}$$

Therefore

$$\frac{1}{\sqrt{g}} dy = (\sqrt{f}) dx$$

$$-\frac{1}{\sqrt{g}} dy = (\sqrt{f}) dx$$

Replacing $f(x), g(y)$ by their values gives

$$\frac{1}{\sqrt{y}} dy = \left(\sqrt{\frac{1}{x}} \right) dx$$

$$-\frac{1}{\sqrt{y}} dy = \left(\sqrt{\frac{1}{x}} \right) dx$$

Integrating now gives the solutions.

$$\int \frac{1}{\sqrt{y}} dy = \int \sqrt{\frac{1}{x}} dx + c_1$$

$$\int -\frac{1}{\sqrt{y}} dy = \int \sqrt{\frac{1}{x}} dx + c_1$$

Integrating gives

$$2\sqrt{y} = 2x\sqrt{\frac{1}{x}} + c_1$$

$$-2\sqrt{y} = 2x\sqrt{\frac{1}{x}} + c_1$$

Therefore

$$y = x\sqrt{\frac{1}{x}} c_1 + \frac{c_1^2}{4} + x$$

$$y = x\sqrt{\frac{1}{x}} c_1 + \frac{c_1^2}{4} + x$$

Summary

The solution(s) found are the following

$$y = x\sqrt{\frac{1}{x}}c_1 + \frac{c_1^2}{4} + x \quad (1)$$

$$y = x\sqrt{\frac{1}{x}}c_1 + \frac{c_1^2}{4} + x \quad (2)$$

Verification of solutions

$$y = x\sqrt{\frac{1}{x}}c_1 + \frac{c_1^2}{4} + x$$

Verified OK. $\{0 < y, 0 < 1/x\}$

$$y = x\sqrt{\frac{1}{x}}c_1 + \frac{c_1^2}{4} + x$$

Verified OK. $\{0 < y, 0 < 1/x\}$

1.44.2 Solving as dAlembert ode

Let $p = y'$ the ode becomes

$$-xp^2 + y = 0$$

Solving for y from the above results in

$$y = xp^2 \quad (1A)$$

This has the form

$$y = xf(p) + g(p) \quad (*)$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved. Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g')\frac{dp}{dx} \\ p - f &= (xf' + g')\frac{dp}{dx} \end{aligned} \quad (2)$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned} f &= p^2 \\ g &= 0 \end{aligned}$$

Hence (2) becomes

$$-p^2 + p = 2xpp'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$-p^2 + p = 0$$

Solving for p from the above gives

$$p = 0$$

$$p = 1$$

Substituting these in (1A) gives

$$y = 0$$

$$y = x$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = \frac{-p(x)^2 + p(x)}{2xp(x)} \quad (3)$$

This ODE is now solved for $p(x)$.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(x) + p(x)p(x) = q(x)$$

Where here

$$p(x) = \frac{1}{2x}$$

$$q(x) = \frac{1}{2x}$$

Hence the ode is

$$p'(x) + \frac{p(x)}{2x} = \frac{1}{2x}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int \frac{1}{2x} dx} \\ &= \sqrt{x} \end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= (\mu) \left(\frac{1}{2x} \right) \\ \frac{d}{dx}(\sqrt{x} p) &= (\sqrt{x}) \left(\frac{1}{2x} \right) \\ d(\sqrt{x} p) &= \left(\frac{1}{2\sqrt{x}} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\sqrt{x} p &= \int \frac{1}{2\sqrt{x}} dx \\ \sqrt{x} p &= \sqrt{x} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \sqrt{x}$ results in

$$p(x) = 1 + \frac{c_1}{\sqrt{x}}$$

Substituting the above solution for p in (2A) gives

$$y = x \left(1 + \frac{c_1}{\sqrt{x}} \right)^2$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$

$$y = x \tag{2}$$

$$y = x \left(1 + \frac{c_1}{\sqrt{x}} \right)^2 \tag{3}$$

Verification of solutions

$$y = 0$$

Verified OK. $\{0 < y, 0 < 1/x\}$

$$y = x$$

Verified OK. $\{0 < y, 0 < 1/x\}$

$$y = x \left(1 + \frac{c_1}{\sqrt{x}} \right)^2$$

Verified OK. $\{0 < y, 0 < 1/x\}$

Maple trace

```
`Methods for first order ODEs:  
*** Sublevel 2 ***  
Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
trying dAlembert  
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 39

```
dsolve(y(x)=x*(diff(y(x),x))^2,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \frac{(x + \sqrt{c_1 x})^2}{x}$$

$$y(x) = \frac{(-x + \sqrt{c_1 x})^2}{x}$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 46

```
DSolve[y[x]==x*(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(-2\sqrt{x} + c_1)^2$$

$$y(x) \rightarrow \frac{1}{4}(2\sqrt{x} + c_1)^2$$

$$y(x) \rightarrow 0$$

1.45 problem 45

1.45.1 Solving as dAlembert ode 315

Internal problem ID [7089]

Internal file name [OUTPUT/6075_Sunday_June_05_2022_04_17_47_PM_71869709/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 45.

ODE order: 1.

ODE degree: 3.

The type(s) of ODE detected by this program : "**dAlembert**"

Maple gives the following as the ode type

[_dAlembert]

$$y'y + xy'^3 = 1$$

1.45.1 Solving as dAlembert ode

Let $p = y'$ the ode becomes

$$x p^3 + p y = 1$$

Solving for y from the above results in

$$y = -p^2 x + \frac{1}{p} \tag{1A}$$

This has the form

$$y = x f(p) + g(p) \tag{*}$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved. Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (x f' + g') \frac{dp}{dx} \\ p - f &= (x f' + g') \frac{dp}{dx} \end{aligned} \tag{2}$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned}f &= -p^2 \\g &= \frac{1}{p}\end{aligned}$$

Hence (2) becomes

$$p^2 + p = \left(-2xp - \frac{1}{p^2}\right) p'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$p^2 + p = 0$$

Solving for p from the above gives

$$\begin{aligned}p &= -1 \\p &= 0\end{aligned}$$

Removing solutions for p which leads to undefined results and substituting these in (1A) gives

$$y = -x - 1$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = \frac{p(x)^2 + p(x)}{-2p(x)x - \frac{1}{p(x)^2}} \quad (3)$$

This ODE is now solved for $p(x)$.

Inverting the above ode gives

$$\frac{d}{dp}x(p) = \frac{-2x(p)p - \frac{1}{p^2}}{p^2 + p} \quad (4)$$

This ODE is now solved for $x(p)$.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$\frac{d}{dp}x(p) + p(p)x(p) = q(p)$$

Where here

$$p(p) = \frac{2}{p+1}$$
$$q(p) = -\frac{1}{p^3(p+1)}$$

Hence the ode is

$$\frac{d}{dp}x(p) + \frac{2x(p)}{p+1} = -\frac{1}{p^3(p+1)}$$

The integrating factor μ is

$$\mu = e^{\int \frac{2}{p+1} dp}$$
$$= (p+1)^2$$

The ode becomes

$$\frac{d}{dp}(\mu x) = (\mu) \left(-\frac{1}{p^3(p+1)} \right)$$
$$\frac{d}{dp}((p+1)^2 x) = ((p+1)^2) \left(-\frac{1}{p^3(p+1)} \right)$$
$$d((p+1)^2 x) = \left(\frac{-p-1}{p^3} \right) dp$$

Integrating gives

$$(p+1)^2 x = \int \frac{-p-1}{p^3} dp$$
$$(p+1)^2 x = \frac{1}{2p^2} + \frac{1}{p} + c_1$$

Dividing both sides by the integrating factor $\mu = (p+1)^2$ results in

$$x(p) = \frac{\frac{1}{2p^2} + \frac{1}{p}}{(p+1)^2} + \frac{c_1}{(p+1)^2}$$

which simplifies to

$$x(p) = \frac{2c_1 p^2 + 2p + 1}{2(p+1)^2 p^2}$$

Now we need to eliminate p between the above and (1A). One way to do this is by solving (1) for p . This results in

$$p = \frac{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)x^2\right)^{\frac{1}{3}}}{6x} - \frac{2y}{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)x^2\right)^{\frac{1}{3}}}$$

$$p = -\frac{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)x^2\right)^{\frac{1}{3}}}{12x} + \frac{y}{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)x^2\right)^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(\frac{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)\right)^{\frac{1}{3}}}{6x}\right)}{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)x^2\right)^{\frac{1}{3}}}$$

$$p = -\frac{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)x^2\right)^{\frac{1}{3}}}{12x} + \frac{y}{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)x^2\right)^{\frac{1}{3}}} - \frac{i\sqrt{3}\left(\frac{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)\right)^{\frac{1}{3}}}{6x}\right)}{\left(\left(12\sqrt{3}\sqrt{\frac{4y^3+27x}{x}} + 108\right)x^2\right)^{\frac{1}{3}}}$$

Substituting the above in the solution for x found above gives

$$x = \frac{54x^3 2^{\frac{2}{3}} 3^{\frac{1}{3}} \left(\frac{x \left(\sqrt{\frac{4y^3+27x}{x}} c_1 3^{\frac{1}{6}} - 2 \left(y - \frac{3c_1}{2} \right) 3^{\frac{2}{3}} \right) 2^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}}}{3} + \frac{2^{\frac{2}{3}} 3^{\frac{5}{6}} x^2 \sqrt{\frac{4y^3+27x}{x}}}{3} + 3x \left(\frac{2y^2 c_1}{9} + x \right) 3^{\frac{1}{3}} 2^{\frac{2}{3}} \right)}{\left(-3^{\frac{1}{3}} 2^{\frac{2}{3}} xy + \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \right)^2 \left(3^{\frac{1}{3}} 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right) \right)}$$

$$x = \frac{36x^3 2^{\frac{2}{3}} 3^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \left(\left(-\frac{8yc_1}{9} + \frac{2x}{3} \right) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + x \left(-\frac{\left((i3^{\frac{2}{3}} + 3^{\frac{1}{6}}) c_1 \right)}{\left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}}} \right) \right)}{\left(\frac{2^{\frac{2}{3}} (i3^{\frac{5}{6}} - 3^{\frac{1}{3}}) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}}}{6} + x \left(2 \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}} + y \right) \right)}$$

$$\begin{aligned}
& x \\
& 36x^3 \left(\left(\frac{8yc_1}{9} - \frac{2x}{3} \right) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + x \left(- \frac{\left(c_1 \left(i3^{\frac{2}{3}} - 3^{\frac{1}{6}} \right) \sqrt{\frac{4y^3+27x}{x}} - 6 \left(y - \frac{3c_1}{2} \right) \left(i3^{\frac{1}{6}} - 3^{\frac{2}{3}} \right) \right)}{9} \right) 2^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \right) \\
= & \frac{\left((-i + \sqrt{3}) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + yx2^{\frac{2}{3}} \left(3^{\frac{5}{6}} + i3^{\frac{1}{3}} \right) \right)^2 \left(\frac{\left(3^{\frac{1}{3}} + i3^{\frac{5}{6}} \right) 2^{\frac{2}{3}}}{3} \right)}{\left(\frac{\left(3^{\frac{1}{3}} + i3^{\frac{5}{6}} \right) 2^{\frac{2}{3}}}{3} \right)}
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = -x - 1 \tag{1}$$

$$x \tag{2}$$

$$\begin{aligned}
& 54x^3 2^{\frac{2}{3}} 3^{\frac{1}{3}} \left(\frac{x \left(\sqrt{\frac{4y^3+27x}{x}} c_1 3^{\frac{1}{6}} - 2 \left(y - \frac{3c_1}{2} \right) 3^{\frac{2}{3}} \right) 2^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}}}{3} + \frac{2^{\frac{2}{3}} 3^{\frac{5}{6}} x^2 \sqrt{\frac{4y^3+27x}{x}}}{3} + 3x \left(\frac{2y^2 c_1}{9} + x \right) 3^{\frac{1}{3}} 2^{\frac{2}{3}} \right) \\
= & \frac{\left(-3^{\frac{1}{3}} 2^{\frac{2}{3}} xy + \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \right)^2 \left(3^{\frac{1}{3}} 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \right)}{\left(3^{\frac{1}{3}} 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \right)}
\end{aligned}$$

$$x = \tag{3}$$

$$\begin{aligned}
& 36x^3 2^{\frac{2}{3}} 3^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \left(\left(-\frac{8yc_1}{9} + \frac{2x}{3} \right) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + x \left(- \frac{\left(\left(i3^{\frac{2}{3}} + 3^{\frac{1}{6}} \right) c_1 \right)}{9} \right) 2^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \right) \\
= & \frac{\left(\frac{2^{\frac{2}{3}} \left(i3^{\frac{5}{6}} - 3^{\frac{1}{3}} \right) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}}}{6} + x \left(2 \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}} + y \right) \right)^2 \left(\frac{\left(3^{\frac{1}{3}} + i3^{\frac{5}{6}} \right) 2^{\frac{2}{3}}}{3} \right)}{\left(\frac{\left(3^{\frac{1}{3}} + i3^{\frac{5}{6}} \right) 2^{\frac{2}{3}}}{3} \right)}
\end{aligned}$$

$$x \tag{4}$$

$$\begin{aligned}
& 36x^3 \left(\left(\frac{8yc_1}{9} - \frac{2x}{3} \right) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + x \left(- \frac{\left(c_1 \left(i3^{\frac{2}{3}} - 3^{\frac{1}{6}} \right) \sqrt{\frac{4y^3+27x}{x}} - 6 \left(y - \frac{3c_1}{2} \right) \left(i3^{\frac{1}{6}} - 3^{\frac{2}{3}} \right) \right)}{9} \right) 2^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \right) \\
= & \frac{\left((-i + \sqrt{3}) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + yx2^{\frac{2}{3}} \left(3^{\frac{5}{6}} + i3^{\frac{1}{3}} \right) \right)^2 \left(\frac{\left(3^{\frac{1}{3}} + i3^{\frac{5}{6}} \right) 2^{\frac{2}{3}}}{3} \right)}{\left(\frac{\left(3^{\frac{1}{3}} + i3^{\frac{5}{6}} \right) 2^{\frac{2}{3}}}{3} \right)}
\end{aligned}$$

Verification of solutions

$$y = -x - 1$$

Verified OK.

x

$$54x^3 2^{\frac{2}{3}} 3^{\frac{1}{3}} \left(\frac{x \left(\sqrt{\frac{4y^3+27x}{x}} c_1 3^{\frac{1}{6}} - 2 \left(y - \frac{3c_1}{2} \right) 3^{\frac{2}{3}} \right) 2^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}}}{3} + \frac{2^{\frac{2}{3}} 3^{\frac{5}{6}} x^2 \sqrt{\frac{4y^3+27x}{x}}}{3} + 3x \left(\frac{2y^2 c_1}{9} + x \right) 3^{\frac{1}{3}} 2^{\frac{2}{3}} \right) -$$

$$\left(-3^{\frac{1}{3}} 2^{\frac{2}{3}} xy + \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \right)^2 \left(3^{\frac{1}{3}} 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right) \right)$$

Warning, solution could not be verified

$x =$

$$36x^3 2^{\frac{2}{3}} 3^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \left(\left(-\frac{8yc_1}{9} + \frac{2x}{3} \right) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + x \left(-\frac{\left((i3^{\frac{2}{3}} + 3^{\frac{1}{6}}) c_1 \right)}{\left((i3^{\frac{2}{3}} + 3^{\frac{1}{6}}) c_1 \right)} \right) \right) -$$

$$\left(\frac{2^{\frac{2}{3}} (i3^{\frac{5}{6}} - 3^{\frac{1}{3}}) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}}}{6} + x \left(2 \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}} + y \right) \right)$$

Warning, solution could not be verified

x

$$36x^3 \left(\left(\frac{8yc_1}{9} - \frac{2x}{3} \right) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + x \left(-\frac{\left(c_1 (i3^{\frac{2}{3}} - 3^{\frac{1}{6}}) \sqrt{\frac{4y^3+27x}{x}} - 6 \left(y - \frac{3c_1}{2} \right) \left(i3^{\frac{1}{6}} - \frac{3^{\frac{2}{3}}}{3} \right) \right) 2^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}}}{9} \right) \right) -$$

$$\left((-i + \sqrt{3}) \left(\left(\sqrt{3} \sqrt{\frac{4y^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + yx 2^{\frac{2}{3}} \left(3^{\frac{5}{6}} + i3^{\frac{1}{3}} \right) \right)^2 \left(\frac{\left((3^{\frac{1}{3}} + i3^{\frac{5}{6}}) 2^{\frac{2}{3}} \right)}{\left((3^{\frac{1}{3}} + i3^{\frac{5}{6}}) 2^{\frac{2}{3}} \right)} \right)$$

Warning, solution could not be verified

Maple trace

```
`Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
trying dAlembert  
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 1817

```
dsolve(diff(y(x),x)*y(x)=1-x*(diff(y(x),x))^3,y(x), singsol=all)
```

$$\begin{aligned} & 12 \left(-2 \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} y(x) + \left(\frac{3^{\frac{2}{3}} 2^{\frac{1}{3}} \left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}}}{6} + 2^{\frac{2}{3}} 3^{\frac{1}{3}} y(x) \right)^2 \right. \\ & \left. + x \left(2^{\frac{2}{3}} 3^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right)^2 x^4 \right)^{\frac{1}{3}} - 2x \left(y(x) 3^{\frac{2}{3}} 2^{\frac{1}{3}} - 3 \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}} \right) \right) \right)^2 \left(y(x) 2^{\frac{2}{3}} \right. \\ & \left. + 18x^4 \left(\sqrt{\frac{4y(x)^3+27x}{x}} 2^{\frac{2}{3}} 3^{\frac{5}{6}} x - 2 \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}} 3^{\frac{2}{3}} 2^{\frac{1}{3}} y(x) + 9 3^{\frac{1}{3}} 2^{\frac{2}{3}} x + 3 \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} \right. \right. \right. \right. \\ & \left. \left. \left. - 2y(x) 3^{\frac{2}{3}} 2^{\frac{1}{3}} x + 2^{\frac{2}{3}} 3^{\frac{1}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right)^2 x^4 \right)^{\frac{1}{3}} + 6x \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}} \right) \right)^2 \left(-y \right. \right. \\ & \left. \left. = 0 \right. \right. \\ & \left. \left. \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} \left(-8 \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} y(x) + x \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) 2^{\frac{1}{3}} \right) \right. \right. \right. \\ & \left. \left. \left. + 6 \left(\frac{\left(i3^{\frac{5}{6}} + 3^{\frac{1}{3}} \right) 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right)^2 x^4 \right)^{\frac{1}{3}}}{6} + \left(-2 \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{1}{3}} + y(x) 2^{\frac{1}{3}} \left(i3^{\frac{1}{6}} - \frac{3^{\frac{2}{3}}}{3} \right) \right) \right) x \right. \right. \right. \\ & \left. \left. + x \right. \right. \\ & \left. \left. 24 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right)^2 x^4 \right)^{\frac{1}{3}} x^4 3^{\frac{1}{3}} \left(- \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + y(x) \left(i3^{\frac{1}{6}} - \frac{3^{\frac{2}{3}}}{3} \right) 2^{\frac{1}{3}} \right) \right. \right. \right. \\ & \left. \left. + \right. \right. \\ & \left. \left. \left((-i\sqrt{3} - 1) \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + (-i3^{\frac{5}{6}} + 3^{\frac{1}{3}}) 2^{\frac{2}{3}} y(x) x \right)^2 \left(\frac{\left(i3^{\frac{5}{6}} + 3^{\frac{1}{3}} \right) 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right)^2 x^4 \right)^{\frac{1}{3}}}{6} \right. \right. \right. \\ & \left. \left. = 0 \right. \right. \\ & \left. \left. \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} c_1 x^3 \left(8 \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} y(x) + \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) 2^{\frac{1}{3}} \right) \right. \right. \right. \\ & \left. \left. \left. + 6 \left((i\sqrt{3} - 1) \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + y(x) \left(i3^{\frac{5}{6}} + 3^{\frac{1}{3}} \right) x 2^{\frac{2}{3}} \right)^2 \left(\frac{\left(-3^{\frac{1}{3}} + i3^{\frac{5}{6}} \right) 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right)^2 x^4 \right)^{\frac{1}{3}}}{6} \right. \right. \right. \\ & \left. \left. + x \right. \right. \\ & \left. \left. 24 2^{\frac{2}{3}} \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right)^2 x^4 \right)^{\frac{1}{3}} x^4 3^{\frac{1}{3}} 2^{\frac{2}{3}} \left(\left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} + 9 \right) x^2 \right)^{\frac{2}{3}} + y(x) \left(i3^{\frac{1}{6}} + \frac{3^{\frac{2}{3}}}{3} \right) 2^{\frac{1}{3}} \right) \left(\left(\sqrt{3} \sqrt{\frac{4y(x)^3+27x}{x}} \right. \right. \right. \\ & \left. \left. + \right. \right. \end{aligned}$$

✓ Solution by Mathematica

Time used: 89.497 (sec). Leaf size: 20717

```
DSolve[y'[x]*y[x]==1-x*(y'[x])^3,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

1.46 problem 46

1.46.1 Solving as quadrature ode	324
1.46.2 Maple step by step solution	325

Internal problem ID [7090]

Internal file name [OUTPUT/6076_Sunday_June_05_2022_04_18_45_PM_61799293/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 46.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$f' - \frac{1}{f} = 0$$

1.46.1 Solving as quadrature ode

Integrating both sides gives

$$\int f df = x + c_1$$
$$\frac{f^2}{2} = x + c_1$$

Solving for f gives these solutions

$$f_1 = \sqrt{2c_1 + 2x}$$
$$f_2 = -\sqrt{2c_1 + 2x}$$

Summary

The solution(s) found are the following

$$f = \sqrt{2c_1 + 2x} \tag{1}$$

$$f = -\sqrt{2c_1 + 2x} \tag{2}$$




Figure 61: Slope field plot

Verification of solutions

$$f = \sqrt{2c_1 + 2x}$$

Verified OK.

$$f = -\sqrt{2c_1 + 2x}$$

Verified OK.

1.46.2 Maple step by step solution

Let's solve

$$f' - \frac{1}{f} = 0$$

- Highest derivative means the order of the ODE is 1

$$f'$$

- Separate variables

$$f'f = 1$$

- Integrate both sides with respect to x

$$\int f' f dx = \int 1 dx + c_1$$

- Evaluate integral

$$\frac{f^2}{2} = x + c_1$$

- Solve for f

$$\{f = \sqrt{2c_1 + 2x}, f = -\sqrt{2c_1 + 2x}\}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(f(x),x)=f(x)^(-1),f(x), singsol=all)
```

$$f(x) = \sqrt{c_1 + 2x}$$

$$f(x) = -\sqrt{c_1 + 2x}$$

✓ Solution by Mathematica

Time used: 0.078 (sec). Leaf size: 38

```
DSolve[f'[x]==f[x]^(-1),f[x],x,IncludeSingularSolutions -> True]
```

$$f(x) \rightarrow -\sqrt{2}\sqrt{x + c_1}$$

$$f(x) \rightarrow \sqrt{2}\sqrt{x + c_1}$$

1.47 problem 47

1.47.1 Solving as second order integrable as is ode	327
1.47.2 Solving as second order ode missing y ode	329
1.47.3 Solving as second order ode non constant coeff transformation on B ode	330
1.47.4 Solving as type second_order_integrable_as_is (not using ABC version)	335
1.47.5 Solving using Kovacic algorithm	336
1.47.6 Solving as exact linear second order ode ode	343
1.47.7 Maple step by step solution	346

Internal problem ID [7091]

Internal file name [OUTPUT/6077_Sunday_June_05_2022_04_18_47_PM_44648207/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 47.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$ty'' + 4y' = t^2$$

1.47.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t t gives

$$\int (ty'' + 4y') dt = \int t^2 dt$$
$$ty' + 3y = \frac{t^3}{3} + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(t)y = q(t)$$

Where here

$$p(t) = \frac{3}{t}$$
$$q(t) = \frac{t^3 + 3c_1}{3t}$$

Hence the ode is

$$y' + \frac{3y}{t} = \frac{t^3 + 3c_1}{3t}$$

The integrating factor μ is

$$\mu = e^{\int \frac{3}{t} dt}$$
$$= t^3$$

The ode becomes

$$\frac{d}{dt}(\mu y) = (\mu) \left(\frac{t^3 + 3c_1}{3t} \right)$$
$$\frac{d}{dt}(t^3 y) = (t^3) \left(\frac{t^3 + 3c_1}{3t} \right)$$
$$d(t^3 y) = \left(\frac{(t^3 + 3c_1)t^2}{3} \right) dt$$

Integrating gives

$$t^3 y = \int \frac{(t^3 + 3c_1)t^2}{3} dt$$
$$t^3 y = \frac{(t^3 + 3c_1)^2}{18} + c_2$$

Dividing both sides by the integrating factor $\mu = t^3$ results in

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3}$$

Summary

The solution(s) found are the following

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3} \quad (1)$$

Verification of solutions

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3}$$

Verified OK.

1.47.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(t) = y'$$

Then

$$p'(t) = y''$$

Hence the ode becomes

$$tp'(t) + 4p(t) - t^2 = 0$$

Which is now solve for $p(t)$ as first order ode.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(t) + p(t)p(t) = q(t)$$

Where here

$$p(t) = \frac{4}{t}$$

$$q(t) = t$$

Hence the ode is

$$p'(t) + \frac{4p(t)}{t} = t$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{4}{t} dt} \\ &= t^4\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dt}(\mu p) &= (\mu)(t) \\ \frac{d}{dt}(t^4 p) &= (t^4)(t) \\ d(t^4 p) &= t^5 dt\end{aligned}$$

Integrating gives

$$\begin{aligned}t^4 p &= \int t^5 dt \\ t^4 p &= \frac{t^6}{6} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = t^4$ results in

$$p(t) = \frac{t^2}{6} + \frac{c_1}{t^4}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{t^2}{6} + \frac{c_1}{t^4}$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{t^6 + 6c_1}{6t^4} dt \\ &= \frac{t^3}{18} - \frac{c_1}{3t^3} + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{t^3}{18} - \frac{c_1}{3t^3} + c_2 \quad (1)$$

Verification of solutions

$$y = \frac{t^3}{18} - \frac{c_1}{3t^3} + c_2$$

Verified OK.

1.47.3 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(t)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$A = t$$

$$B = 4$$

$$C = 0$$

$$F = t^2$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (t)(0) + (4)(0) + (0)(4) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$4tv'' + (16)v' = 0$$

Now by applying $v' = u$ the above becomes

$$4tu'(t) + 16u(t) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(t, u) \\ &= f(t)g(u) \\ &= -\frac{4u}{t} \end{aligned}$$

Where $f(t) = -\frac{4}{t}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{4}{t} dt \\ \int \frac{1}{u} du &= \int -\frac{4}{t} dt \\ \ln(u) &= -4 \ln(t) + c_1 \\ u &= e^{-4 \ln(t) + c_1} \\ &= \frac{c_1}{t^4}\end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= \frac{c_1}{t^4}\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(t) &= \int \frac{c_1}{t^4} dt \\ &= -\frac{c_1}{3t^3} + c_2\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(t) &= Bv \\ &= (4) \left(-\frac{c_1}{3t^3} + c_2 \right) \\ &= -\frac{4c_1}{3t^3} + 4c_2\end{aligned}$$

And now the particular solution $y_p(t)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$y_p(t) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= 4 \\ y_2 &= \frac{1}{t^3}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 4 & \frac{1}{t^3} \\ \frac{d}{dt}(4) & \frac{d}{dt}\left(\frac{1}{t^3}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 4 & \frac{1}{t^3} \\ 0 & -\frac{3}{t^4} \end{vmatrix}$$

Therefore

$$W = (4) \left(-\frac{3}{t^4}\right) - \left(\frac{1}{t^3}\right) (0)$$

Which simplifies to

$$W = -\frac{12}{t^4}$$

Which simplifies to

$$W = -\frac{12}{t^4}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{1}{t}}{-\frac{12}{t^3}} dt$$

Which simplifies to

$$u_1 = - \int -\frac{t^2}{12} dt$$

Hence

$$u_1 = \frac{t^3}{36}$$

And Eq. (3) becomes

$$u_2 = \int \frac{4t^2}{-\frac{12}{t^3}} dt$$

Which simplifies to

$$u_2 = \int -\frac{t^5}{3} dt$$

Hence

$$u_2 = -\frac{t^6}{18}$$

Therefore the particular solution, from equation (1) is

$$y_p(t) = \frac{t^3}{18}$$

Hence the complete solution is

$$\begin{aligned} y(t) &= y_h + y_p \\ &= \left(-\frac{4c_1}{3t^3} + 4c_2 \right) + \left(\frac{t^3}{18} \right) \\ &= -\frac{4c_1}{3t^3} + 4c_2 + \frac{t^3}{18} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{4c_1}{3t^3} + 4c_2 + \frac{t^3}{18} \quad (1)$$

Verification of solutions

$$y = -\frac{4c_1}{3t^3} + 4c_2 + \frac{t^3}{18}$$

Verified OK.

1.47.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$ty'' + 4y' = t^2$$

Integrating both sides of the ODE w.r.t t gives

$$\int (ty'' + 4y') dt = \int t^2 dt$$
$$ty' + 3y = \frac{t^3}{3} + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(t)y = q(t)$$

Where here

$$p(t) = \frac{3}{t}$$
$$q(t) = \frac{t^3 + 3c_1}{3t}$$

Hence the ode is

$$y' + \frac{3y}{t} = \frac{t^3 + 3c_1}{3t}$$

The integrating factor μ is

$$\mu = e^{\int \frac{3}{t} dt}$$
$$= t^3$$

The ode becomes

$$\frac{d}{dt}(\mu y) = (\mu) \left(\frac{t^3 + 3c_1}{3t} \right)$$
$$\frac{d}{dt}(t^3 y) = (t^3) \left(\frac{t^3 + 3c_1}{3t} \right)$$
$$d(t^3 y) = \left(\frac{(t^3 + 3c_1)t^2}{3} \right) dt$$

Integrating gives

$$t^3 y = \int \frac{(t^3 + 3c_1)t^2}{3} dt$$
$$t^3 y = \frac{(t^3 + 3c_1)^2}{18} + c_2$$

Dividing both sides by the integrating factor $\mu = t^3$ results in

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3}$$

Summary

The solution(s) found are the following

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3} \quad (1)$$

Verification of solutions

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3}$$

Verified OK.

1.47.5 Solving using Kovacic algorithm

Writing the ode as

$$ty'' + 4y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = t$$
$$B = 4$$
$$C = 0 \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{t^2} \tag{6}$$

Comparing the above to (5) shows that

$$s = 2$$

$$t = t^2$$

Therefore eq. (4) becomes

$$z''(t) = \left(\frac{2}{t^2}\right) z(t) \tag{7}$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 55: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = t^2$. There is a pole at $t = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{t^2}$$

For the pole at $t = 0$ let b be the coefficient of $\frac{1}{t^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{t^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{t^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{t^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{t-c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{t-c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= -\frac{1}{t} + (-)(0) \\ &= -\frac{1}{t} \\ &= -\frac{1}{t} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(t)$ of degree $d = 0$ to solve the ode. The polynomial $p(t)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(t) = 1 \tag{2A}$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{t}\right)(0) + \left(\left(\frac{1}{t^2}\right) + \left(-\frac{1}{t}\right)^2 - \left(\frac{2}{t^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(t) = pe^{\int \omega dt}$$
$$= e^{\int -\frac{1}{t} dt}$$
$$= \frac{1}{t}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$
$$= z_1 e^{-\int \frac{1}{2} \frac{4}{t} dt}$$
$$= z_1 e^{-2 \ln(t)}$$
$$= z_1 \left(\frac{1}{t^2}\right)$$

Which simplifies to

$$y_1 = \frac{1}{t^3}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{4}{t} dt}}{(y_1)^2} dt$$
$$= y_1 \int \frac{e^{-4 \ln(t)}}{(y_1)^2} dt$$
$$= y_1 \left(\frac{t^3}{3}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{t^3} \right) + c_2 \left(\frac{1}{t^3} \left(\frac{t^3}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$ty'' + 4y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1}{t^3} + \frac{c_2}{3}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$y_p(t) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{t^3}$$

$$y_2 = \frac{1}{3}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(t)}{aW(t)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(t)}{aW(t)} \tag{3}$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{t^3} & \frac{1}{3} \\ \frac{d}{dt}\left(\frac{1}{t^3}\right) & \frac{d}{dt}\left(\frac{1}{3}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{t^3} & \frac{1}{3} \\ -\frac{3}{t^4} & 0 \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{t^3}\right)(0) - \left(\frac{1}{3}\right)\left(-\frac{3}{t^4}\right)$$

Which simplifies to

$$W = \frac{1}{t^4}$$

Which simplifies to

$$W = \frac{1}{t^4}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{t^2}{\frac{3}{\frac{1}{t^3}}} dt$$

Which simplifies to

$$u_1 = - \int \frac{t^5}{3} dt$$

Hence

$$u_1 = -\frac{t^6}{18}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{t}}{\frac{1}{t^3}} dt$$

Which simplifies to

$$u_2 = \int t^2 dt$$

Hence

$$u_2 = \frac{t^3}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(t) = \frac{t^3}{18}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1}{t^3} + \frac{c_2}{3} \right) + \left(\frac{t^3}{18} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{t^3} + \frac{c_2}{3} + \frac{t^3}{18} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{t^3} + \frac{c_2}{3} + \frac{t^3}{18}$$

Verified OK.

1.47.6 Solving as exact linear second order ode ode

An ode of the form

$$p(t) y'' + q(t) y' + r(t) y = s(t)$$

is exact if

$$p''(t) - q'(t) + r(t) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned}p(x) &= t \\q(x) &= 4 \\r(x) &= 0 \\s(x) &= t^2\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(t) y' + (q(t) - p'(t)) y)' = s(x)$$

Integrating gives

$$p(t) y' + (q(t) - p'(t)) y = \int s(t) dt$$

Substituting the above values for p, q, r, s gives

$$ty' + 3y = \int t^2 dt$$

We now have a first order ode to solve which is

$$ty' + 3y = \frac{t^3}{3} + c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(t)y = q(t)$$

Where here

$$\begin{aligned}p(t) &= \frac{3}{t} \\q(t) &= \frac{t^3 + 3c_1}{3t}\end{aligned}$$

Hence the ode is

$$y' + \frac{3y}{t} = \frac{t^3 + 3c_1}{3t}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{3}{t} dt} \\ &= t^3\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dt}(\mu y) &= (\mu) \left(\frac{t^3 + 3c_1}{3t} \right) \\ \frac{d}{dt}(t^3 y) &= (t^3) \left(\frac{t^3 + 3c_1}{3t} \right) \\ d(t^3 y) &= \left(\frac{(t^3 + 3c_1)t^2}{3} \right) dt\end{aligned}$$

Integrating gives

$$\begin{aligned}t^3 y &= \int \frac{(t^3 + 3c_1)t^2}{3} dt \\ t^3 y &= \frac{(t^3 + 3c_1)^2}{18} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = t^3$ results in

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3}$$

Summary

The solution(s) found are the following

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3} \quad (1)$$

Verification of solutions

$$y = \frac{(t^3 + 3c_1)^2}{18t^3} + \frac{c_2}{t^3}$$

Verified OK.

1.47.7 Maple step by step solution

Let's solve

$$ty'' + 4y' = t^2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$tu'(t) + 4u(t) = t^2$$

- Isolate the derivative

$$u'(t) = -\frac{4u(t)}{t} + t$$

- Group terms with $u(t)$ on the lhs of the ODE and the rest on the rhs of the ODE

$$u'(t) + \frac{4u(t)}{t} = t$$

- The ODE is linear; multiply by an integrating factor $\mu(t)$

$$\mu(t) \left(u'(t) + \frac{4u(t)}{t} \right) = \mu(t) t$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dt}(\mu(t) u(t))$

$$\mu(t) \left(u'(t) + \frac{4u(t)}{t} \right) = \mu'(t) u(t) + \mu(t) u'(t)$$

- Isolate $\mu'(t)$

$$\mu'(t) = \frac{4\mu(t)}{t}$$

- Solve to find the integrating factor

$$\mu(t) = t^4$$

- Integrate both sides with respect to t

$$\int \left(\frac{d}{dt}(\mu(t) u(t)) \right) dt = \int \mu(t) t dt + c_1$$

- Evaluate the integral on the lhs

$$\mu(t) u(t) = \int \mu(t) t dt + c_1$$

- Solve for $u(t)$

$$u(t) = \frac{\int \mu(t) t dt + c_1}{\mu(t)}$$

- Substitute $\mu(t) = t^4$

$$u(t) = \frac{\int t^5 dt + c_1}{t^4}$$

- Evaluate the integrals on the rhs

$$u(t) = \frac{t^6 + c_1}{t^4}$$

- Simplify

$$u(t) = \frac{t^6 + 6c_1}{6t^4}$$

- Solve 1st ODE for $u(t)$

$$u(t) = \frac{t^6 + 6c_1}{6t^4}$$

- Make substitution $u = y'$

$$y' = \frac{t^6 + 6c_1}{6t^4}$$

- Integrate both sides to solve for y

$$\int y' dt = \int \frac{t^6 + 6c_1}{6t^4} dt + c_2$$

- Compute integrals

$$y = \frac{t^3}{18} - \frac{c_1}{3t^3} + c_2$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(-_a^2+4*_b(_a))/_a, _b(_a)` *** Sub
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(t*diff(y(t),t$2)+4*diff(y(t),t)=t^2,y(t), singsol=all)
```

$$y(t) = \frac{t^3}{18} - \frac{c_1}{3t^3} + c_2$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 24

```
DSolve[t*y''[t]+4*y'[t]==t^2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{t^3}{18} - \frac{c_1}{3t^3} + c_2$$

1.48 problem 48

1.48.1 Existence and uniqueness analysis	349
1.48.2 Solving as second order integrable as is ode	350
1.48.3 Solving as second order ode missing y ode	352
1.48.4 Solving as type second_order_integrable_as_is (not using ABC version)	354
1.48.5 Solving using Kovacic algorithm	356
1.48.6 Solving as exact linear second order ode ode	362

Internal problem ID [7092]

Internal file name [OUTPUT/6078_Sunday_June_05_2022_04_18_49_PM_32004638/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 48.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$(t^2 + 9) y'' + 2ty' = 0$$

With initial conditions

$$\left[y(3) = 2\pi, y'(3) = \frac{2}{3} \right]$$

1.48.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = \frac{2t}{t^2 + 9}$$

$$q(t) = 0$$

$$F = 0$$

Hence the ode is

$$y'' + \frac{2ty'}{t^2 + 9} = 0$$

The domain of $p(t) = \frac{2t}{t^2+9}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 3$ is inside this domain. Hence solution exists and is unique.

1.48.2 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t t gives

$$\begin{aligned} \int ((t^2 + 9) y'' + 2ty') dt &= 0 \\ (t^2 + 9) y' &= c_1 \end{aligned}$$

Which is now solved for y . Integrating both sides gives

$$\begin{aligned} y &= \int \frac{c_1}{t^2 + 9} dt \\ &= \frac{c_1 \arctan\left(\frac{t}{3}\right)}{3} + c_2 \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1 \arctan\left(\frac{t}{3}\right)}{3} + c_2 \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2\pi$ and $t = 3$ in the above gives

$$2\pi = \frac{\pi c_1}{12} + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{t^2 + 9}$$

substituting $y' = \frac{2}{3}$ and $t = 3$ in the above gives

$$\frac{2}{3} = \frac{c_1}{18} \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 12$$

$$c_2 = \pi$$

Substituting these values back in above solution results in

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Summary

The solution(s) found are the following

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi \tag{1}$$




Figure 62: Solution plot

Verification of solutions

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Verified OK.

1.48.3 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(t) = y'$$

Then

$$p'(t) = y''$$

Hence the ode becomes

$$(t^2 + 9)p'(t) + 2tp(t) = 0$$

Which is now solve for $p(t)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(t, p) \\ &= f(t)g(p) \\ &= -\frac{2tp}{t^2 + 9} \end{aligned}$$

Where $f(t) = -\frac{2t}{t^2+9}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\frac{2t}{t^2 + 9} dt \\ \int \frac{1}{p} dp &= \int -\frac{2t}{t^2 + 9} dt \\ \ln(p) &= -\ln(t^2 + 9) + c_1 \\ p &= e^{-\ln(t^2+9)+c_1} \\ &= \frac{c_1}{t^2 + 9} \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $t = 3$ and $p = \frac{2}{3}$ in the above solution gives an equation to solve for the constant of integration.

$$\frac{2}{3} = \frac{c_1}{18}$$

$$c_1 = 12$$

Substituting c_1 found above in the general solution gives

$$p(t) = \frac{12}{t^2 + 9}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{12}{t^2 + 9}$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{12}{t^2 + 9} dt \\ &= 4 \arctan\left(\frac{t}{3}\right) + c_2 \end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $t = 3$ and $y = 2\pi$ in the above solution gives an equation to solve for the constant of integration.

$$2\pi = \pi + c_2$$

$$c_2 = \pi$$

Substituting c_2 found above in the general solution gives

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi \tag{1}$$




Figure 63: Solution plot

Verification of solutions

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Verified OK.

1.48.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$(t^2 + 9) y'' + 2ty' = 0$$

Integrating both sides of the ODE w.r.t t gives

$$\int ((t^2 + 9) y'' + 2ty') dt = 0$$
$$(t^2 + 9) y' = c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int \frac{c_1}{t^2 + 9} dt$$
$$= \frac{c_1 \arctan\left(\frac{t}{3}\right)}{3} + c_2$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1 \arctan\left(\frac{t}{3}\right)}{3} + c_2 \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2\pi$ and $t = 3$ in the above gives

$$2\pi = \frac{\pi c_1}{12} + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{t^2 + 9}$$

substituting $y' = \frac{2}{3}$ and $t = 3$ in the above gives

$$\frac{2}{3} = \frac{c_1}{18} \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 12$$

$$c_2 = \pi$$

Substituting these values back in above solution results in

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Summary

The solution(s) found are the following

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi \tag{1}$$




Figure 64: Solution plot

Verification of solutions

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Verified OK.

1.48.5 Solving using Kovacic algorithm

Writing the ode as

$$(t^2 + 9) y'' + 2ty' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= t^2 + 9 \\ B &= 2t \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{(t^2 + 9)^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= (t^2 + 9)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = \left(\frac{9}{(t^2 + 9)^2} \right) z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 57: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 4 - 0 \\ &= 4 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = (t^2 + 9)^2$. There is a pole at $t = 3i$ of order 2. There is a pole at $t = -3i$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 4 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 4 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4(t-3i)^2} - \frac{1}{4(t+3i)^2} - \frac{i}{12(t-3i)} + \frac{i}{12t+36i}$$

For the pole at $t = 3i$ let b be the coefficient of $\frac{1}{(t-3i)^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

For the pole at $t = -3i$ let b be the coefficient of $\frac{1}{(t+3i)^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $4 > 2$ then

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= 0 \\ \alpha_\infty^- &= 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{9}{(t^2 + 9)^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
$3i$	2	0	$\frac{1}{2}$	$\frac{1}{2}$
$-3i$	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
4	0	0	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^+ + \alpha_{c_2}^+) \\ &= 1 - (1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{t - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((+)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^+}{t - c_1} \right) + \left((+)[\sqrt{r}]_{c_2} + \frac{\alpha_{c_2}^+}{t - c_2} \right) + (-)[\sqrt{r}]_\infty \\ &= \frac{1}{2t - 6i} + \frac{1}{2t + 6i} + (-)(0) \\ &= \frac{1}{2t - 6i} + \frac{1}{2t + 6i} \\ &= \frac{t}{t^2 + 9} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(t)$ of degree $d = 0$ to solve the ode. The polynomial $p(t)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(t) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2 \left(\frac{1}{2t - 6i} + \frac{1}{2t + 6i} \right) (0) + \left(\left(-\frac{1}{2(t - 3i)^2} - \frac{1}{2(t + 3i)^2} \right) + \left(\frac{1}{2t - 6i} + \frac{1}{2t + 6i} \right)^2 - \left(\frac{9}{(t^2 + 9)^2} \right) \right) 1 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(t) &= p e^{\int \omega dt} \\ &= e^{\int \left(\frac{1}{2t - 6i} + \frac{1}{2t + 6i} \right) dt} \\ &= \sqrt{t^2 + 9} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\&= z_1 e^{-\int \frac{1}{2} \frac{2t}{t^2+9} dt} \\&= z_1 e^{-\frac{\ln(t^2+9)}{2}} \\&= z_1 \left(\frac{1}{\sqrt{t^2+9}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{2t}{t^2+9} dt}}{(y_1)^2} dt \\&= y_1 \int \frac{e^{-\ln(t^2+9)}}{(y_1)^2} dt \\&= y_1 \left(\frac{\arctan\left(\frac{t}{3}\right)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 (1) + c_2 \left(1 \left(\frac{\arctan\left(\frac{t}{3}\right)}{3} \right) \right)\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 + \frac{c_2 \arctan\left(\frac{t}{3}\right)}{3} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2\pi$ and $t = 3$ in the above gives

$$2\pi = c_1 + \frac{\pi c_2}{12} \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_2}{t^2 + 9}$$

substituting $y' = \frac{2}{3}$ and $t = 3$ in the above gives

$$\frac{2}{3} = \frac{c_2}{18} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \pi$$

$$c_2 = 12$$

Substituting these values back in above solution results in

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Summary

The solution(s) found are the following

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi \quad (1)$$




Figure 65: Solution plot

Verification of solutions

$$y = 4 \arctan \left(\frac{t}{3} \right) + \pi$$

Verified OK.

1.48.6 Solving as exact linear second order ode

An ode of the form

$$p(t) y'' + q(t) y' + r(t) y = s(t)$$

is exact if

$$p''(t) - q'(t) + r(t) = 0 \tag{1}$$

For the given ode we have

$$p(x) = t^2 + 9$$

$$q(x) = 2t$$

$$r(x) = 0$$

$$s(x) = 0$$

Hence

$$p''(x) = 2$$

$$q'(x) = 2$$

Therefore (1) becomes

$$2 - (2) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(t) y' + (q(t) - p'(t)) y)' = s(x)$$

Integrating gives

$$p(t) y' + (q(t) - p'(t)) y = \int s(t) dt$$

Substituting the above values for p, q, r, s gives

$$(t^2 + 9) y' = c_1$$

We now have a first order ode to solve which is

$$(t^2 + 9) y' = c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{c_1}{t^2 + 9} dt \\ &= \frac{c_1 \arctan\left(\frac{t}{3}\right)}{3} + c_2 \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1 \arctan\left(\frac{t}{3}\right)}{3} + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2\pi$ and $t = 3$ in the above gives

$$2\pi = \frac{\pi c_1}{12} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{t^2 + 9}$$

substituting $y' = \frac{2}{3}$ and $t = 3$ in the above gives

$$\frac{2}{3} = \frac{c_1}{18} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 12$$

$$c_2 = \pi$$

Substituting these values back in above solution results in

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Summary

The solution(s) found are the following

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi \quad (1)$$




Figure 66: Solution plot

Verification of solutions

$$y = 4 \arctan\left(\frac{t}{3}\right) + \pi$$

Verified OK.

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
<- LODE missing y successful`

```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 12

```
dsolve([(t^2+9)*diff(y(t),t)+2*t*diff(y(t),t)=0,y(3) = 2*Pi, D(y)(3) = 2/3],y(t), singsol=
```

$$y(t) = \pi + 4 \arctan\left(\frac{t}{3}\right)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 15

```
DSolve[{(t^2+9)*y'[t]+2*t*y'[t]==0,{y[3]==2*Pi,y'[3]==2/3}},y[t],t,IncludeSingularSolutions
```

$$y(t) \rightarrow 4 \arctan\left(\frac{t}{3}\right) + \pi$$

1.49 problem 49

1.49.1 Solving as second order euler ode ode	366
1.49.2 Solving as second order change of variable on x method 2 ode .	368
1.49.3 Solving as second order change of variable on x method 1 ode .	371
1.49.4 Solving as second order change of variable on y method 2 ode .	373
1.49.5 Solving using Kovacic algorithm	375
1.49.6 Maple step by step solution	380

Internal problem ID [7093]

Internal file name [OUTPUT/6079_Sunday_June_05_2022_04_18_52_PM_86132628/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 49.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$t^2 y'' - 3ty' + 5y = 0$$

1.49.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = t^r$, then $y' = rt^{r-1}$ and $y'' = r(r-1)t^{r-2}$. Substituting these back into the given ODE gives

$$t^2(r(r-1))t^{r-2} - 3trt^{r-1} + 5t^r = 0$$

Simplifying gives

$$r(r-1)t^r - 3rt^r + 5t^r = 0$$

Since $t^r \neq 0$ then dividing throughout by t^r gives

$$r(r-1) - 3r + 5 = 0$$

Or

$$r^2 - 4r + 5 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 2 - i$$

$$r_2 = 2 + i$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$

$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = 2$ and $\beta = -1$. Hence the solution becomes

$$\begin{aligned} y &= c_1 t^{r_1} + c_2 t^{r_2} \\ &= c_1 t^{\alpha+i\beta} + c_2 t^{\alpha-i\beta} \\ &= t^\alpha (c_1 t^{i\beta} + c_2 t^{-i\beta}) \\ &= t^\alpha (c_1 e^{\ln(t^{i\beta})} + c_2 e^{\ln(t^{-i\beta})}) \\ &= t^\alpha (c_1 e^{i(\beta \ln t)} + c_2 e^{-i(\beta \ln t)}) \end{aligned}$$

Using the values for $\alpha = 2, \beta = -1$, the above becomes

$$y = t^2 (c_1 e^{-i \ln(t)} + c_2 e^{i \ln(t)})$$

Using Euler relation, the expression $c_1 e^{iA} + c_2 e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y = t^2 (c_1 \cos (\ln (t)) + c_2 \sin (\ln (t)))$$

Summary

The solution(s) found are the following

$$y = t^2 (c_1 \cos (\ln (t)) + c_2 \sin (\ln (t))) \quad (1)$$

Verification of solutions

$$y = t^2 (c_1 \cos (\ln (t)) + c_2 \sin (\ln (t)))$$

Verified OK.

1.49.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$t^2 y'' - 3ty' + 5y = 0 \quad (1)$$

Becomes

$$y'' + p(t)y' + q(t)y = 0 \quad (2)$$

Where

$$p(t) = -\frac{3}{t}$$
$$q(t) = \frac{5}{t^2}$$

Applying change of variables $\tau = g(t)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(t) + p(t)\tau'(t)}{\tau'(t)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(t)}{\tau'(t)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(t) + p(t)\tau'(t) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(t)dt)} dt \\ &= \int e^{-(\int -\frac{3}{t}dt)} dt \\ &= \int e^{3\ln(t)} dt \\ &= \int t^3 dt \\ &= \frac{t^4}{4} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(t)}{\tau'(t)^2} \\ &= \frac{5}{t^2} \\ &= \frac{5}{t^8} \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{5y(\tau)}{t^8} &= 0 \end{aligned}$$

But in terms of τ

$$\frac{5}{t^8} = \frac{5}{16\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{5y(\tau)}{16\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$16 \left(\frac{d^2}{d\tau^2}y(\tau) \right) \tau^2 + 5y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$16\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + 5\tau^r = 0$$

Simplifying gives

$$16r(r-1)\tau^r + 0\tau^r + 5\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$16r(r-1) + 0 + 5 = 0$$

Or

$$16r^2 - 16r + 5 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2} - \frac{i}{4}$$

$$r_2 = \frac{1}{2} + \frac{i}{4}$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$

$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = \frac{1}{2}$ and $\beta = -\frac{1}{4}$. Hence the solution becomes

$$\begin{aligned} y(\tau) &= c_1\tau^{r_1} + c_2\tau^{r_2} \\ &= c_1\tau^{\alpha+i\beta} + c_2\tau^{\alpha-i\beta} \\ &= \tau^\alpha(c_1\tau^{i\beta} + c_2\tau^{-i\beta}) \\ &= \tau^\alpha(c_1e^{\ln(\tau^{i\beta})} + c_2e^{\ln(\tau^{-i\beta})}) \\ &= \tau^\alpha(c_1e^{i(\beta \ln \tau)} + c_2e^{-i(\beta \ln \tau)}) \end{aligned}$$

Using the values for $\alpha = \frac{1}{2}, \beta = -\frac{1}{4}$, the above becomes

$$y(\tau) = \tau^{\frac{1}{2}} \left(c_1 e^{-\frac{i \ln(\tau)}{4}} + c_2 e^{\frac{i \ln(\tau)}{4}} \right)$$

Using Euler relation, the expression $c_1 e^{iA} + c_2 e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y(\tau) = \sqrt{\tau} \left(c_1 \cos \left(\frac{\ln(\tau)}{4} \right) + c_2 \sin \left(\frac{\ln(\tau)}{4} \right) \right)$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{\left(c_1 \cos \left(-\frac{\ln(2)}{2} + \ln(t) \right) + c_2 \sin \left(-\frac{\ln(2)}{2} + \ln(t) \right) \right) t^2}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\left(c_1 \cos \left(-\frac{\ln(2)}{2} + \ln(t) \right) + c_2 \sin \left(-\frac{\ln(2)}{2} + \ln(t) \right) \right) t^2}{2} \quad (1)$$

Verification of solutions

$$y = \frac{\left(c_1 \cos \left(-\frac{\ln(2)}{2} + \ln(t) \right) + c_2 \sin \left(-\frac{\ln(2)}{2} + \ln(t) \right) \right) t^2}{2}$$

Verified OK.

1.49.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$t^2 y'' - 3ty' + 5y = 0 \quad (1)$$

Becomes

$$y'' + p(t)y' + q(t)y = 0 \quad (2)$$

Where

$$p(t) = -\frac{3}{t}$$

$$q(t) = \frac{5}{t^2}$$

Applying change of variables $\tau = g(t)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1 \left(\frac{d}{d\tau}y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(t) + p(t)\tau'(t)}{\tau'(t)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(t)}{\tau'(t)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c} \sqrt{q} \\ &= \frac{\sqrt{5} \sqrt{\frac{1}{t^2}}}{c} \\ \tau'' &= -\frac{\sqrt{5}}{c \sqrt{\frac{1}{t^2}} t^3} \end{aligned} \quad (6)$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(t) + p(t) \tau'(t)}{\tau'(t)^2} \\
 &= \frac{-\frac{\sqrt{5}}{c\sqrt{\frac{1}{t^2}}t^3} - \frac{3}{t} \frac{\sqrt{5}\sqrt{\frac{1}{t^2}}}{c}}{\left(\frac{\sqrt{5}\sqrt{\frac{1}{t^2}}}{c}\right)^2} \\
 &= -\frac{4c\sqrt{5}}{5}
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2} y(\tau) - \frac{4c\sqrt{5}}{5} \left(\frac{d}{d\tau} y(\tau)\right) + c^2 y(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{\frac{2\sqrt{5}c\tau}{5}} \left(c_1 \cos\left(\frac{\sqrt{5}c\tau}{5}\right) + c_2 \sin\left(\frac{\sqrt{5}c\tau}{5}\right) \right)$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c} \sqrt{q} dt \\
 &= \frac{\int \sqrt{5} \sqrt{\frac{1}{t^2}} dt}{c} \\
 &= \frac{\sqrt{5} \sqrt{\frac{1}{t^2}} t \ln(t)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = t^2 (c_1 \cos(\ln(t)) + c_2 \sin(\ln(t)))$$

Summary

The solution(s) found are the following

$$y = t^2 (c_1 \cos(\ln(t)) + c_2 \sin(\ln(t))) \tag{1}$$

Verification of solutions

$$y = t^2 (c_1 \cos(\ln(t)) + c_2 \sin(\ln(t)))$$

Verified OK.

1.49.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$t^2 y'' - 3ty' + 5y = 0 \quad (1)$$

Becomes

$$y'' + p(t)y' + q(t)y = 0 \quad (2)$$

Where

$$p(t) = -\frac{3}{t}$$
$$q(t) = \frac{5}{t^2}$$

Applying change of variables on the dependent variable $y = v(t)t^n$ to (2) gives the following ode where the dependent variables is $v(t)$ and not y .

$$v''(t) + \left(\frac{2n}{t} + p\right)v'(t) + \left(\frac{n(n-1)}{t^2} + \frac{np}{t} + q\right)v(t) = 0 \quad (3)$$

Let the coefficient of $v(t)$ above be zero. Hence

$$\frac{n(n-1)}{t^2} + \frac{np}{t} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(t)$ and $q(t)$ into (4) gives

$$\frac{n(n-1)}{t^2} - \frac{3n}{t} + \frac{5}{t^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 + i \quad (6)$$

Substituting this value in (3) gives

$$v''(t) + \left(\frac{4+2i}{t} - \frac{3}{t}\right)v'(t) = 0$$
$$v''(t) + \frac{(1+2i)v'(t)}{t} = 0 \quad (7)$$

Using the substitution

$$u(t) = v'(t)$$

Then (7) becomes

$$u'(t) + \frac{(1 + 2i)u(t)}{t} = 0 \quad (8)$$

The above is now solved for $u(t)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(t, u) \\ &= f(t)g(u) \\ &= \frac{(-1 - 2i)u}{t} \end{aligned}$$

Where $f(t) = \frac{-1-2i}{t}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{-1 - 2i}{t} dt \\ \int \frac{1}{u} du &= \int \frac{-1 - 2i}{t} dt \\ \ln(u) &= (-1 - 2i) \ln(t) + c_1 \\ u &= e^{(-1-2i)\ln(t)+c_1} \\ &= c_1 e^{(-1-2i)\ln(t)} \end{aligned}$$

Which simplifies to

$$u(t) = \frac{c_1 t^{-2i}}{t}$$

Now that $u(t)$ is known, then

$$\begin{aligned} v'(t) &= u(t) \\ v(t) &= \int u(t) dt + c_2 \\ &= \frac{ic_1 t^{-2i}}{2} + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(t) t^n \\ &= \left(\frac{ic_1 t^{-2i}}{2} + c_2 \right) t^{2+i} \\ &= c_2 t^{2+i} + \frac{ic_1 t^{2-i}}{2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{ic_1 t^{-2i}}{2} + c_2 \right) t^{2+i} \quad (1)$$

Verification of solutions

$$y = \left(\frac{ic_1 t^{-2i}}{2} + c_2 \right) t^{2+i}$$

Verified OK.

1.49.5 Solving using Kovacic algorithm

Writing the ode as

$$t^2 y'' - 3ty' + 5y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= t^2 \\ B &= -3t \\ C &= 5 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-5}{4t^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -5 \\ t &= 4t^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = \left(-\frac{5}{4t^2}\right) z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 58: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4t^2$. There is a pole at $t = 0$ of order 2. Since there is no odd order pole

larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{5}{4t^2}$$

For the pole at $t = 0$ let b be the coefficient of $\frac{1}{t^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{5}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + i \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - i \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{t^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{5}{4t^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{5}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + i \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - i \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{5}{4t^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + i$	$\frac{1}{2} - i$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + i$	$\frac{1}{2} - i$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{1}{2} - i$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{1}{2} - i - \left(\frac{1}{2} - i \right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{t - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{t - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= \frac{\frac{1}{2} - i}{t} + (-)(0) \\ &= \frac{\frac{1}{2} - i}{t} \\ &= \frac{\frac{1}{2} - i}{t} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(t)$ of degree $d = 0$ to solve the ode. The polynomial $p(t)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(t) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(\frac{\frac{1}{2} - i}{t}\right)(0) + \left(\left(\frac{-\frac{1}{2} + i}{t^2}\right) + \left(\frac{\frac{1}{2} - i}{t}\right)^2 - \left(-\frac{5}{4t^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(t) = pe^{\int \omega dt}$$

$$= e^{\int \frac{\frac{1}{2} - i}{t} dt}$$

$$= t^{\frac{1}{2} - i}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$

$$= z_1 e^{-\int \frac{1}{2} \frac{-3t}{t^2} dt}$$

$$= z_1 e^{\frac{3 \ln(t)}{2}}$$

$$= z_1 \left(t^{\frac{3}{2}}\right)$$

Which simplifies to

$$y_1 = t^{2-i}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{-3t}{t^2} dt}}{(y_1)^2} dt$$

$$= y_1 \int \frac{e^{3 \ln(t)}}{(y_1)^2} dt$$

$$= y_1 \left(-\frac{it^{2i}}{2}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (t^{2-i}) + c_2 \left(t^{2-i} \left(-\frac{it^{2i}}{2} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 t^{2-i} - \frac{ic_2 t^{2+i}}{2} \quad (1)$$

Verification of solutions

$$y = c_1 t^{2-i} - \frac{ic_2 t^{2+i}}{2}$$

Verified OK.

1.49.6 Maple step by step solution

Let's solve

$$y''t^2 - 3ty' + 5y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{3y'}{t} - \frac{5y}{t^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{3y'}{t} + \frac{5y}{t^2} = 0$$

- Multiply by denominators of the ODE

$$y''t^2 - 3ty' + 5y = 0$$

- Make a change of variables

$$s = \ln(t)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to t , using the chain rule

$$y' = \left(\frac{d}{ds} y(s) \right) s'(t)$$

- Compute derivative

$$y' = \frac{\frac{d}{ds}y(s)}{t}$$

- Calculate the 2nd derivative of y with respect to t , using the chain rule

$$y'' = \left(\frac{d^2}{ds^2}y(s) \right) s'(t)^2 + s''(t) \left(\frac{d}{ds}y(s) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{ds^2}y(s)}{t^2} - \frac{\frac{d}{ds}y(s)}{t^2}$$

Substitute the change of variables back into the ODE

$$\left(\frac{\frac{d^2}{ds^2}y(s)}{t^2} - \frac{\frac{d}{ds}y(s)}{t^2} \right) t^2 - 3 \frac{d}{ds}y(s) + 5y(s) = 0$$

- Simplify

$$\frac{d^2}{ds^2}y(s) - 4 \frac{d}{ds}y(s) + 5y(s) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 4r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{4 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (2 - I, 2 + I)$$

- 1st solution of the ODE

$$y_1(s) = e^{2s} \cos(s)$$

- 2nd solution of the ODE

$$y_2(s) = e^{2s} \sin(s)$$

- General solution of the ODE

$$y(s) = c_1 y_1(s) + c_2 y_2(s)$$

- Substitute in solutions

$$y(s) = c_1 e^{2s} \cos(s) + c_2 e^{2s} \sin(s)$$

- Change variables back using $s = \ln(t)$

$$y = c_1 t^2 \cos(\ln(t)) + c_2 t^2 \sin(\ln(t))$$

- Simplify

$$y = t^2(c_1 \cos(\ln(t)) + c_2 \sin(\ln(t)))$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(t^2*diff(y(t),t$2)-3*t*diff(y(t),t)+5*y(t)=0,y(t), singsol=all)
```

$$y(t) = t^2(c_1 \sin(\ln(t)) + c_2 \cos(\ln(t)))$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 22

```
DSolve[t^2*y''[t]-3*t*y'[t]+5*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow t^2(c_2 \cos(\log(t)) + c_1 \sin(\log(t)))$$

1.50 problem 50

1.50.1 Solving as second order integrable as is ode	383
1.50.2 Solving as second order ode missing y ode	384
1.50.3 Solving as second order ode non constant coeff transformation on B ode	385
1.50.4 Solving as type second_order_integrable_as_is (not using ABC version)	387
1.50.5 Solving using Kovacic algorithm	388
1.50.6 Solving as exact linear second order ode ode	393
1.50.7 Maple step by step solution	394

Internal problem ID [7094]

Internal file name [OUTPUT/6080_Sunday_June_05_2022_04_18_53_PM_46488860/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 50.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$ty'' + y' = 0$$

1.50.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t t gives

$$\int (ty'' + y') dt = 0$$
$$ty' = c_1$$

Which is now solved for y . Integrating both sides gives

$$\begin{aligned}y &= \int \frac{c_1}{t} dt \\ &= c_1 \ln(t) + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \ln(t) + c_2 \quad (1)$$

Verification of solutions

$$y = c_1 \ln(t) + c_2$$

Verified OK.

1.50.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(t) = y'$$

Then

$$p'(t) = y''$$

Hence the ode becomes

$$tp'(t) + p(t) = 0$$

Which is now solve for $p(t)$ as first order ode. In canonical form the ODE is

$$\begin{aligned}p' &= F(t, p) \\ &= f(t)g(p) \\ &= -\frac{p}{t}\end{aligned}$$

Where $f(t) = -\frac{1}{t}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned}\frac{1}{p} dp &= -\frac{1}{t} dt \\ \int \frac{1}{p} dp &= \int -\frac{1}{t} dt \\ \ln(p) &= -\ln(t) + c_1 \\ p &= e^{-\ln(t)+c_1} \\ &= \frac{c_1}{t}\end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{c_1}{t}$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{c_1}{t} dt \\ &= c_1 \ln(t) + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \ln(t) + c_2 \quad (1)$$

Verification of solutions

$$y = c_1 \ln(t) + c_2$$

Verified OK.

1.50.3 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(t)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2) u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$A = t$$

$$B = 1$$

$$C = 0$$

$$F = 0$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (t)(0) + (1)(0) + (0)(1) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$tv'' + (1)v' = 0$$

Now by applying $v' = u$ the above becomes

$$tu'(t) + u(t) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(t, u) \\ &= f(t)g(u) \\ &= -\frac{u}{t} \end{aligned}$$

Where $f(t) = -\frac{1}{t}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{1}{t} dt \\ \int \frac{1}{u} du &= \int -\frac{1}{t} dt \\ \ln(u) &= -\ln(t) + c_1 \\ u &= e^{-\ln(t)+c_1} \\ &= \frac{c_1}{t} \end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= \frac{c_1}{t}\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(t) &= \int \frac{c_1}{t} dt \\ &= c_1 \ln(t) + c_2\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y(t) &= Bv \\ &= (1)(c_1 \ln(t) + c_2) \\ &= c_1 \ln(t) + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \ln(t) + c_2 \tag{1}$$

Verification of solutions

$$y = c_1 \ln(t) + c_2$$

Verified OK.

1.50.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$ty'' + y' = 0$$

Integrating both sides of the ODE w.r.t t gives

$$\begin{aligned}\int (ty'' + y') dt &= 0 \\ ty' &= c_1\end{aligned}$$

Which is now solved for y . Integrating both sides gives

$$\begin{aligned}y &= \int \frac{c_1}{t} dt \\ &= c_1 \ln(t) + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \ln(t) + c_2 \quad (1)$$

Verification of solutions

$$y = c_1 \ln(t) + c_2$$

Verified OK.

1.50.5 Solving using Kovacic algorithm

Writing the ode as

$$ty'' + y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= t \\ B &= 1 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{4t^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 4t^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = \left(-\frac{1}{4t^2}\right) z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 60: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4t^2$. There is a pole at $t = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4t^2}$$

For the pole at $t = 0$ let b be the coefficient of $\frac{1}{t^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{t^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{4t^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{4t^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to

determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_{∞}^{\pm} . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_{\infty}^{-} = \frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \left(\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{t - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{t - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{1}{2t} + (-)(0) \\ &= \frac{1}{2t} \\ &= \frac{1}{2t} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(t)$ of degree $d = 0$ to solve the ode. The polynomial $p(t)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(t) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{2t}\right)(0) + \left(\left(-\frac{1}{2t^2}\right) + \left(\frac{1}{2t}\right)^2 - \left(-\frac{1}{4t^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned}z_1(t) &= pe^{\int \omega dt} \\ &= e^{\int \frac{1}{2t} dt} \\ &= \sqrt{t}\end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{t} dt} \\ &= z_1 e^{-\frac{\ln(t)}{2}} \\ &= z_1 \left(\frac{1}{\sqrt{t}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{t} dt}}{(y_1)^2} dt \\ &= y_1 \int \frac{e^{-\ln(t)}}{(y_1)^2} dt \\ &= y_1(\ln(t))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(\ln(t)))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2 \ln(t) \quad (1)$$

Verification of solutions

$$y = c_1 + c_2 \ln(t)$$

Verified OK.

1.50.6 Solving as exact linear second order ode

An ode of the form

$$p(t) y'' + q(t) y' + r(t) y = s(t)$$

is exact if

$$p''(t) - q'(t) + r(t) = 0 \quad (1)$$

For the given ode we have

$$p(x) = t$$

$$q(x) = 1$$

$$r(x) = 0$$

$$s(x) = 0$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(t) y' + (q(t) - p'(t)) y)' = s(x)$$

Integrating gives

$$p(t) y' + (q(t) - p'(t)) y = \int s(t) dt$$

Substituting the above values for p, q, r, s gives

$$ty' = c_1$$

We now have a first order ode to solve which is

$$ty' = c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{c_1}{t} dt \\ &= c_1 \ln(t) + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \ln(t) + c_2 \tag{1}$$

Verification of solutions

$$y = c_1 \ln(t) + c_2$$

Verified OK.

1.50.7 Maple step by step solution

Let's solve

$$ty'' + y' = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{t}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{t} = 0$$

- Multiply by denominators of the ODE

$$ty'' + y' = 0$$

- Make a change of variables

$$s = \ln(t)$$

- Substitute the change of variables back into the ODE
 - Calculate the 1st derivative of y with respect to t , using the chain rule

$$y' = \left(\frac{d}{ds}y(s)\right) s'(t)$$
 - Compute derivative

$$y' = \frac{\frac{d}{ds}y(s)}{t}$$
 - Calculate the 2nd derivative of y with respect to t , using the chain rule

$$y'' = \left(\frac{d^2}{ds^2}y(s)\right) s'(t)^2 + s''(t) \left(\frac{d}{ds}y(s)\right)$$
 - Compute derivative

$$y'' = \frac{\frac{d^2}{ds^2}y(s)}{t^2} - \frac{\frac{d}{ds}y(s)}{t^2}$$

Substitute the change of variables back into the ODE

$$t \left(\frac{\frac{d^2}{ds^2}y(s)}{t^2} - \frac{\frac{d}{ds}y(s)}{t^2} \right) + \frac{\frac{d}{ds}y(s)}{t} = 0$$

- Simplify

$$\frac{\frac{d^2}{ds^2}y(s)}{t} = 0$$
- Isolate 2nd derivative

$$\frac{d^2}{ds^2}y(s) = 0$$
- Characteristic polynomial of ODE

$$r^2 = 0$$
- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$
- Roots of the characteristic polynomial

$$r = 0$$
- 1st solution of the ODE

$$y_1(s) = 1$$
- Repeated root, multiply $y_1(s)$ by s to ensure linear independence

$$y_2(s) = s$$
- General solution of the ODE

$$y(s) = c_1 y_1(s) + c_2 y_2(s)$$

- Substitute in solutions
 $y(s) = c_2 s + c_1$
- Change variables back using $s = \ln(t)$
 $y = c_1 + c_2 \ln(t)$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(t*diff(y(t),t$2)+diff(y(t),t)=0,y(t), singsol=all)
```

$$y(t) = c_2 \ln(t) + c_1$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 13

```
DSolve[t*y''[t]+y'[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_1 \log(t) + c_2$$

1.51 problem 51

1.51.1 Solving as second order ode missing y ode	397
1.51.2 Solving as second order ode non constant coeff transformation on B ode	398
1.51.3 Solving using Kovacic algorithm	401

Internal problem ID [7095]

Internal file name [OUTPUT/6081_Sunday_June_05_2022_04_18_55_PM_45043879/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 51.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_ode_missing_y", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$t^2 y'' - 2y' = 0$$

1.51.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(t) = y'$$

Then

$$p'(t) = y''$$

Hence the ode becomes

$$p'(t) t^2 - 2p(t) = 0$$

Which is now solve for $p(t)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(t, p) \\ &= f(t)g(p) \\ &= \frac{2p}{t^2} \end{aligned}$$

Where $f(t) = \frac{2}{t^2}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned}\frac{1}{p} dp &= \frac{2}{t^2} dt \\ \int \frac{1}{p} dp &= \int \frac{2}{t^2} dt \\ \ln(p) &= -\frac{2}{t} + c_1 \\ p &= e^{-\frac{2}{t} + c_1} \\ &= c_1 e^{-\frac{2}{t}}\end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = c_1 e^{-\frac{2}{t}}$$

Integrating both sides gives

$$\begin{aligned}y &= \int c_1 e^{-\frac{2}{t}} dt \\ &= c_1 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right) + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right) + c_2 \quad (1)$$

Verification of solutions

$$y = c_1 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right) + c_2$$

Verified OK.

1.51.2 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(t)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned}y' &= B'v + v'B \\y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v\end{aligned}$$

And now the original ode becomes

$$\begin{aligned}A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0\end{aligned}\tag{1}$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned}A &= t^2 \\B &= -2 \\C &= 0 \\F &= 0\end{aligned}$$

The above shows that for this ode

$$\begin{aligned}AB'' + BB' + CB &= (t^2)(0) + (-2)(0) + (0)(-2) \\ &= 0\end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-2t^2v'' + (4)v' = 0$$

Now by applying $v' = u$ the above becomes

$$-2t^2u'(t) + 4u(t) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(t, u) \\ &= f(t)g(u) \\ &= \frac{2u}{t^2} \end{aligned}$$

Where $f(t) = \frac{2}{t^2}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{2}{t^2} dt \\ \int \frac{1}{u} du &= \int \frac{2}{t^2} dt \\ \ln(u) &= -\frac{2}{t} + c_1 \\ u &= e^{-\frac{2}{t} + c_1} \\ &= c_1 e^{-\frac{2}{t}} \end{aligned}$$

The ode for v now becomes

$$\begin{aligned} v' &= u \\ &= c_1 e^{-\frac{2}{t}} \end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned} v(t) &= \int c_1 e^{-\frac{2}{t}} dt \\ &= c_1 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right) + c_2 \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y(t) &= Bv \\ &= (-2) \left(c_1 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right) + c_2 \right) \\ &= -2tc_1 e^{-\frac{2}{t}} + 4 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) c_1 - 2c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -2tc_1 e^{-\frac{2}{t}} + 4 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) c_1 - 2c_2 \quad (1)$$

Verification of solutions

$$y = -2tc_1e^{-\frac{2}{t}} + 4 \exp\text{Integral}_1\left(\frac{2}{t}\right)c_1 - 2c_2$$

Verified OK.

1.51.3 Solving using Kovacic algorithm

Writing the ode as

$$t^2y'' - 2y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = t^2$$

$$B = -2 \quad (3)$$

$$C = 0$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2t + 1}{t^4} \quad (6)$$

Comparing the above to (5) shows that

$$s = 2t + 1$$

$$t = t^4$$

Therefore eq. (4) becomes

$$z''(t) = \left(\frac{2t+1}{t^4} \right) z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 62: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 4 - 1 \\ &= 3 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = t^4$. There is a pole at $t = 0$ of order 4. Since there is no odd order pole larger than 2 and the order at ∞ is 3 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Looking at higher order poles of order $2v \geq 4$ (must be even order for case one). Then for each pole c , $[\sqrt{r}]_c$ is the sum of terms $\frac{1}{(t-c)^i}$ for $2 \leq i \leq v$ in the Laurent series expansion of \sqrt{r} expanded around each pole c . Hence

$$[\sqrt{r}]_c = \sum_2^v \frac{a_i}{(t-c)^i} \quad (1B)$$

Let a be the coefficient of the term $\frac{1}{(t-c)^v}$ in the above where v is the pole order divided by 2. Let b be the coefficient of $\frac{1}{(t-c)^{v+1}}$ in r minus the coefficient of $\frac{1}{(t-c)^{v+1}}$ in $[\sqrt{r}]_c$. Then

$$\alpha_c^+ = \frac{1}{2} \left(\frac{b}{a} + v \right)$$

$$\alpha_c^- = \frac{1}{2} \left(-\frac{b}{a} + v \right)$$

The partial fraction decomposition of r is

$$r = \frac{1}{t^4} + \frac{2}{t^3}$$

There is pole in r at $t = 0$ of order 4, hence $v = 2$. Expanding \sqrt{r} as Laurent series about this pole $c = 0$ gives

$$[\sqrt{r}]_c \approx \frac{1}{t^2} + \frac{1}{t} - \frac{1}{2} + \frac{t}{2} - \frac{5t^2}{8} + \frac{7t^3}{8} + \dots \quad (2B)$$

Using eq. (1B), taking the sum up to $v = 2$ the above becomes

$$[\sqrt{r}]_c = \frac{1}{t^2} \quad (3B)$$

The above shows that the coefficient of $\frac{1}{(t-0)^2}$ is

$$a = 1$$

Now we need to find b . let b be the coefficient of the term $\frac{1}{(t-c)^{v+1}}$ in r minus the coefficient of the same term but in the sum $[\sqrt{r}]_c$ found in eq. (3B). Here c is current pole which is $c = 0$. This term becomes $\frac{1}{t^3}$. The coefficient of this term in the sum $[\sqrt{r}]_c$ is seen to be 0 and the coefficient of this term r is found from the partial fraction decomposition from above to be 2. Therefore

$$b = (2) - (0)$$

$$= 2$$

Hence

$$\begin{aligned}
[\sqrt{r}]_c &= \frac{1}{t^2} \\
\alpha_c^+ &= \frac{1}{2} \left(\frac{b}{a} + v \right) = \frac{1}{2} \left(\frac{2}{1} + 2 \right) = 2 \\
\alpha_c^- &= \frac{1}{2} \left(-\frac{b}{a} + v \right) = \frac{1}{2} \left(-\frac{2}{1} + 2 \right) = 0
\end{aligned}$$

Since the order of r at ∞ is $3 > 2$ then

$$\begin{aligned}
[\sqrt{r}]_\infty &= 0 \\
\alpha_\infty^+ &= 0 \\
\alpha_\infty^- &= 1
\end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2t + 1}{t^4}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	4	$\frac{1}{t^2}$	2	0

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
3	0	0	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^+ = 0$ then

$$\begin{aligned}
d &= \alpha_\infty^+ - (\alpha_{c_1}^-) \\
&= 0 - (0) \\
&= 0
\end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{t - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

Substituting the above values in the above results in

$$\begin{aligned}
 \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{t - c_1} \right) + (+)[\sqrt{r}]_{\infty} \\
 &= -\frac{1}{t^2} + (0) \\
 &= -\frac{1}{t^2} \\
 &= -\frac{1}{t^2}
 \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(t)$ of degree $d = 0$ to solve the ode. The polynomial $p(t)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(t) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned}
 (0) + 2\left(-\frac{1}{t^2}\right)(0) + \left(\left(\frac{2}{t^3}\right) + \left(-\frac{1}{t^2}\right)^2 - \left(\frac{2t+1}{t^4}\right)\right) &= 0 \\
 0 &= 0
 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned}
 z_1(t) &= pe^{\int \omega dt} \\
 &= e^{\int -\frac{1}{t^2} dt} \\
 &= e^{\frac{1}{t}}
 \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-2}{t^2} dt} \\
 &= z_1 e^{-\frac{1}{t}} \\
 &= z_1 \left(e^{-\frac{1}{t}} \right)
 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{t^2} dt}}{(y_1)^2} dt \\ &= y_1 \int \frac{e^{-\frac{2}{t}}}{(y_1)^2} dt \\ &= y_1 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2 \left(1 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right) \quad (1)$$

Verification of solutions

$$y = c_1 + c_2 \left(t e^{-\frac{2}{t}} - 2 \operatorname{expIntegral}_1 \left(\frac{2}{t} \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
<- LODE missing y successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(t^2*diff(y(t),t$2)-2*diff(y(t),t)=0,y(t), singsol=all)
```

$$y(t) = e^{-\frac{2}{t}}c_2t - 2 \operatorname{expIntegral}_1\left(\frac{2}{t}\right)c_2 + c_1$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 29

```
DSolve[t^2*y''[t]-2*y'[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow 2c_1 \operatorname{ExpIntegralEi}\left(-\frac{2}{t}\right) + c_1 e^{-2/t} + c_2$$

1.52 problem 52

Internal problem ID [7096]

Internal file name [OUTPUT/6082_Sunday_June_05_2022_04_18_57_PM_26022958/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 52.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' + \frac{(t^2 - 1)y'}{t} + \frac{t^2 y}{\left(1 + e^{\frac{t^2}{2}}\right)^2} = 0$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
<- unable to find a useful change of variables
    trying a symmetry of the form [xi=0, eta=F(x)]
    trying 2nd order exact linear
    trying symmetries linear in x and y(x)
    trying to convert to a linear ODE with constant coefficients
    <- to_const_coeffs successful: conversion to a linear ODE with constant coefficients was
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 84

```
dsolve(diff(y(t),t$2)+(t^2-1)/t*diff(y(t),t)+t^2/(1 + exp(t^2/2))^2*y(t)=0,y(t), singsol=all
```

$$y(t) = \frac{\left(c_1 \left(1 + e^{\frac{t^2}{2}} \right)^{-\frac{i\sqrt{3}}{2}} \left(e^{\frac{t^2}{2}} \right)^{\frac{i\sqrt{3}}{2}} + c_2 \left(1 + e^{\frac{t^2}{2}} \right)^{\frac{i\sqrt{3}}{2}} \left(e^{\frac{t^2}{2}} \right)^{-\frac{i\sqrt{3}}{2}} \right) \sqrt{1 + e^{\frac{t^2}{2}}}}{\sqrt{e^{\frac{t^2}{2}}}}$$

✓ Solution by Mathematica

Time used: 0.116 (sec). Leaf size: 72

```
DSolve[y''[t]+(t^2-1)/t*y'[t]+t^2/(1 + Exp[t^2/2])^2*y[t]==0,y[t],t,IncludeSingularSolutions
```

$$y(t) \rightarrow e^{\operatorname{arctanh}\left(2e^{\frac{t^2}{2}}+1\right)} \left(c_2 \cos\left(\sqrt{3}\operatorname{arctanh}\left(2e^{\frac{t^2}{2}}+1\right)\right) - c_1 \sin\left(\sqrt{3}\operatorname{arctanh}\left(2e^{\frac{t^2}{2}}+1\right)\right) \right)$$

1.53 problem 53

- 1.53.1 Solving as second order change of variable on x method 2 ode . 410
- 1.53.2 Solving as second order change of variable on x method 1 ode . 413
- 1.53.3 Solving as second order bessele ode 415
- 1.53.4 Solving using Kovacic algorithm 416
- 1.53.5 Maple step by step solution 422

Internal problem ID [7097]

Internal file name [OUTPUT/6083_Sunday_June_05_2022_04_19_00_PM_68918213/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 53.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_bessel_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$ty'' - y' + 4t^3y = 0$$

1.53.1 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$ty'' - y' + 4t^3y = 0 \tag{1}$$

Becomes

$$y'' + p(t)y' + q(t)y = 0 \tag{2}$$

Where

$$p(t) = -\frac{1}{t}$$
$$q(t) = 4t^2$$

Applying change of variables $\tau = g(t)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(t) + p(t)\tau'(t)}{\tau'(t)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(t)}{\tau'(t)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(t) + p(t)\tau'(t) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(t)dt)} dt \\ &= \int e^{-(\int -\frac{1}{t}dt)} dt \\ &= \int e^{\ln(t)} dt \\ &= \int t dt \\ &= \frac{t^2}{2} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(t)}{\tau'(t)^2} \\ &= \frac{4t^2}{t^2} \\ &= 4 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + 4y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} + 4 e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y(\tau) = e^{\alpha\tau} (c_1 \cos(\beta\tau) + c_2 \sin(\beta\tau))$$

Which becomes

$$y(\tau) = e^0 (c_1 \cos(2\tau) + c_2 \sin(2\tau))$$

Or

$$y(\tau) = c_1 \cos(2\tau) + c_2 \sin(2\tau)$$

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \cos(t^2) + c_2 \sin(t^2)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(t^2) + c_2 \sin(t^2) \quad (1)$$

Verification of solutions

$$y = c_1 \cos(t^2) + c_2 \sin(t^2)$$

Verified OK.

1.53.2 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$ty'' - y' + 4t^3y = 0 \quad (1)$$

Becomes

$$y'' + p(t)y' + q(t)y = 0 \quad (2)$$

Where

$$p(t) = -\frac{1}{t}$$
$$q(t) = 4t^2$$

Applying change of variables $\tau = g(t)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(t) + p(t)\tau'(t)}{\tau'(t)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(t)}{\tau'(t)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{2\sqrt{t^2}}{c} \\ \tau'' &= \frac{2t}{c\sqrt{t^2}}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(t) + p(t)\tau'(t)}{\tau'(t)^2} \\ &= \frac{\frac{2t}{c\sqrt{t^2}} - \frac{1}{t}\frac{2\sqrt{t^2}}{c}}{\left(\frac{2\sqrt{t^2}}{c}\right)^2} \\ &= 0\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dt \\ &= \frac{\int 2\sqrt{t^2} dt}{c} \\ &= \frac{t\sqrt{t^2}}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos(t^2) + c_2 \sin(t^2)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(t^2) + c_2 \sin(t^2)\tag{1}$$

Verification of solutions

$$y = c_1 \cos(t^2) + c_2 \sin(t^2)$$

Verified OK.

1.53.3 Solving as second order Bessel ODE

Writing the ODE as

$$y''t^2 - ty' + 4t^4y = 0 \quad (1)$$

Bessel ODE has the form

$$y''t^2 + ty' + (-n^2 + t^2)y = 0 \quad (2)$$

The generalized form of Bessel ODE is given by Bowman (1958) as the following

$$y''t^2 + (1 - 2\alpha)ty' + (\beta^2\gamma^2t^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \quad (3)$$

With the standard solution

$$y = t^\alpha(c_1 \text{BesselJ}(n, \beta t^\gamma) + c_2 \text{BesselY}(n, \beta t^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\alpha = 1$$

$$\beta = 1$$

$$n = \frac{1}{2}$$

$$\gamma = 2$$

Substituting all the above into (4) gives the solution as

$$y = \frac{c_1 t \sqrt{2} \sin(t^2)}{\sqrt{\pi} \sqrt{t^2}} - \frac{c_2 t \sqrt{2} \cos(t^2)}{\sqrt{\pi} \sqrt{t^2}}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 t \sqrt{2} \sin(t^2)}{\sqrt{\pi} \sqrt{t^2}} - \frac{c_2 t \sqrt{2} \cos(t^2)}{\sqrt{\pi} \sqrt{t^2}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 t \sqrt{2} \sin(t^2)}{\sqrt{\pi} \sqrt{t^2}} - \frac{c_2 t \sqrt{2} \cos(t^2)}{\sqrt{\pi} \sqrt{t^2}}$$

Verified OK.

1.53.4 Solving using Kovacic algorithm

Writing the ode as

$$ty'' - y' + 4t^3y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= t \\ B &= -1 \\ C &= 4t^3 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-16t^4 + 3}{4t^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -16t^4 + 3 \\ t &= 4t^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = \left(\frac{-16t^4 + 3}{4t^2} \right) z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 63: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 4 \\ &= -2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4t^2$. There is a pole at $t = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Therefore

$$L = [1, 2]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -4t^2 + \frac{3}{4t^2}$$

For the pole at $t = 0$ let b be the coefficient of $\frac{1}{t^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving t^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i t^i \\ &= \sum_{i=0}^1 a_i t^i \end{aligned} \tag{8}$$

Let a be the coefficient of $t^v = t^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx 2it - \frac{3i}{16t^3} - \frac{9i}{1024t^7} - \frac{27i}{32768t^{11}} - \frac{405i}{4194304t^{15}} - \frac{1701i}{134217728t^{19}} - \frac{15309i}{8589934592t^{23}} - \frac{72171i}{274877906944t^{27}} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = 2i$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i t^i \\ &= 2it \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $t^{v-1} = t^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = -4t^2$$

This shows that the coefficient of 1 in the above is 0. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned}
 r &= \frac{s}{t} \\
 &= \frac{-16t^4 + 3}{4t^2} \\
 &= Q + \frac{R}{4t^2} \\
 &= (-4t^2) + \left(\frac{3}{4t^2}\right) \\
 &= -4t^2 + \frac{3}{4t^2}
 \end{aligned}$$

We see that the coefficient of the term t in the quotient is 0. Now b can be found.

$$\begin{aligned}
 b &= (0) - (0) \\
 &= 0
 \end{aligned}$$

Hence

$$\begin{aligned}
 [\sqrt{r}]_{\infty} &= 2it \\
 \alpha_{\infty}^{+} &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{0}{2i} - 1 \right) = -\frac{1}{2} \\
 \alpha_{\infty}^{-} &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{0}{2i} - 1 \right) = -\frac{1}{2}
 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{-16t^4 + 3}{4t^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^{+}	α_c^{-}
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_{\infty}$	α_{∞}^{+}	α_{∞}^{-}
-2	$2it$	$-\frac{1}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^{\pm} have been determined for all the poles in the set Γ and $[\sqrt{r}]_{\infty}$ and its associated α_{∞}^{\pm} have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{t-c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{t-c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2t} + (-)(2it) \\ &= -\frac{1}{2t} - 2it \\ &= -\frac{1}{2t} - 2it \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(t)$ of degree $d = 0$ to solve the ode. The polynomial $p(t)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(t) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{1}{2t} - 2it\right)(0) + \left(\left(\frac{1}{2t^2} - 2i\right) + \left(-\frac{1}{2t} - 2it\right)^2 - \left(\frac{-16t^4 + 3}{4t^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(t) &= pe^{\int \omega dt} \\ &= e^{\int \left(-\frac{1}{2t} - 2it\right) dt} \\ &= \frac{e^{-it^2}}{\sqrt{t}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\&= z_1 e^{-\int \frac{1}{2} \frac{-1}{t} dt} \\&= z_1 e^{\frac{\ln(t)}{2}} \\&= z_1 (\sqrt{t})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-it^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{t} dt}}{(y_1)^2} dt \\&= y_1 \int \frac{e^{\ln(t)}}{(y_1)^2} dt \\&= y_1 \left(-\frac{ie^{2it^2}}{4} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 (e^{-it^2}) + c_2 \left(e^{-it^2} \left(-\frac{ie^{2it^2}}{4} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-it^2} - \frac{ic_2 e^{it^2}}{4} \quad (1)$$

Verification of solutions

$$y = c_1 e^{-it^2} - \frac{ic_2 e^{it^2}}{4}$$

Verified OK.

1.53.5 Maple step by step solution

Let's solve

$$ty'' - y' + 4t^3y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y'}{t} - 4t^2y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{t} + 4t^2y = 0$$

- Check to see if $t_0 = 0$ is a regular singular point

- Define functions

$$[P_2(t) = -\frac{1}{t}, P_3(t) = 4t^2]$$

- $t \cdot P_2(t)$ is analytic at $t = 0$

$$(t \cdot P_2(t)) \Big|_{t=0} = -1$$

- $t^2 \cdot P_3(t)$ is analytic at $t = 0$

$$(t^2 \cdot P_3(t)) \Big|_{t=0} = 0$$

- $t = 0$ is a regular singular point

Check to see if $t_0 = 0$ is a regular singular point

$$t_0 = 0$$

- Multiply by denominators

$$ty'' - y' + 4t^3y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k t^{k+r}$$

□ Rewrite ODE with series expansions

- Convert $t^3 \cdot y$ to series expansion

$$t^3 \cdot y = \sum_{k=0}^{\infty} a_k t^{k+r+3}$$

- Shift index using $k \rightarrow k - 3$

$$t^3 \cdot y = \sum_{k=3}^{\infty} a_{k-3} t^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) t^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) t^{k+r}$$

- Convert $t \cdot y''$ to series expansion

$$t \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r) (k+r-1) t^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$t \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) (k+r) t^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-2+r) t^{-1+r} + a_1 (1+r) (-1+r) t^r + a_2 (2+r) r t^{1+r} + a_3 (3+r) (1+r) t^{2+r} + \left(\sum_{k=3}^{\infty} a_{k+1} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-2+r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \{0, 2\}$$
- The coefficients of each power of t must be 0

$$[a_1(1+r)(-1+r) = 0, a_2(2+r)r = 0, a_3(3+r)(1+r) = 0]$$
- Solve for the dependent coefficient(s)

$$\{a_1 = 0, a_2 = 0, a_3 = 0\}$$
- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1+r)(k+r-1) + 4a_{k-3} = 0$$

- Shift index using $k \rightarrow k + 3$

$$a_{k+4}(k+4+r)(k+2+r) + 4a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+4} = -\frac{4a_k}{(k+4+r)(k+2+r)}$$

- Recursion relation for $r = 0$

$$a_{k+4} = -\frac{4a_k}{(k+4)(k+2)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k t^k, a_{k+4} = -\frac{4a_k}{(k+4)(k+2)}, a_1 = 0, a_2 = 0, a_3 = 0 \right]$$

- Recursion relation for $r = 2$

$$a_{k+4} = -\frac{4a_k}{(k+6)(k+4)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k t^{k+2}, a_{k+4} = -\frac{4a_k}{(k+6)(k+4)}, a_1 = 0, a_2 = 0, a_3 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k t^k \right) + \left(\sum_{k=0}^{\infty} b_k t^{k+2} \right), a_{k+4} = -\frac{4a_k}{(k+4)(k+2)}, a_1 = 0, a_2 = 0, a_3 = 0, b_{k+4} = -\frac{4b_k}{(k+6)(k+4)}, \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(t*diff(y(t),t$2)-diff(y(t),t)+4*t^3*y(t)=0,y(t), singsol=all)
```

$$y(t) = c_1 \sin(t^2) + c_2 \cos(t^2)$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 20

```
DSolve[t*y''[t]-y'[t]+4*t^3*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_1 \cos(t^2) + c_2 \sin(t^2)$$

1.54 problem 54

1.54.1 Solving as second order ode quadrature ode	426
1.54.2 Solving as second order linear constant coeff ode	427
1.54.3 Solving as second order ode can be made integrable ode	429
1.54.4 Solving as second order integrable as is ode	430
1.54.5 Solving as second order ode missing y ode	431
1.54.6 Solving using Kovacic algorithm	433
1.54.7 Solving as exact linear second order ode ode	436
1.54.8 Maple step by step solution	438

Internal problem ID [7098]

Internal file name [OUTPUT/6084_Sunday_June_05_2022_04_19_02_PM_96915118/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 54.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = 0$$

1.54.1 Solving as second order ode quadrature ode

Integrating twice gives the solution

$$y = c_1 t + c_2$$

Summary

The solution(s) found are the following

$$y = c_1 t + c_2 \tag{1}$$




Figure 67: Slope field plot

Verification of solutions

$$y = c_1t + c_2$$

Verified OK.

1.54.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0\end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 t \quad (1)$$

Summary

The solution(s) found are the following

$$y = c_2 t + c_1 \quad (1)$$




Figure 68: Slope field plot

Verification of solutions

$$y = c_2 t + c_1$$

Verified OK.

1.54.3 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' = 0$$

Integrating the above w.r.t t gives

$$\int y'y'' dt = 0$$
$$\frac{y'^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{c_1} \sqrt{2} \quad (1)$$

$$y' = -\sqrt{c_1} \sqrt{2} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$y = \int \sqrt{c_1} \sqrt{2} dt$$
$$= t\sqrt{c_1} \sqrt{2} + c_2$$

Solving equation (2)

Integrating both sides gives

$$y = \int -\sqrt{c_1} \sqrt{2} dt$$
$$= -t\sqrt{c_1} \sqrt{2} + c_3$$

Summary

The solution(s) found are the following

$$y = t\sqrt{c_1} \sqrt{2} + c_2 \quad (1)$$

$$y = -t\sqrt{c_1} \sqrt{2} + c_3 \quad (2)$$




Figure 69: Slope field plot

Verification of solutions

$$y = t\sqrt{c_1} \sqrt{2} + c_2$$

Verified OK.

$$y = -t\sqrt{c_1} \sqrt{2} + c_3$$

Verified OK.

1.54.4 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t t gives

$$\int y'' dt = 0$$

$$y' = c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int c_1 dt$$

$$= c_1 t + c_2$$

Summary

The solution(s) found are the following

$$y = c_1 t + c_2 \quad (1)$$




Figure 70: Slope field plot

Verification of solutions

$$y = c_1 t + c_2$$

Verified OK.

1.54.5 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(t) = y'$$

Then

$$p'(t) = y''$$

Hence the ode becomes

$$p'(t) = 0$$

Which is now solve for $p(t)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(t) &= \int 0 \, dt \\ &= c_1 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int c_1 \, dt \\ &= c_1 t + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 t + c_2 \tag{1}$$




Figure 71: Slope field plot

Verification of solutions

$$y = c_1 t + c_2$$

Verified OK.

1.54.6 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 65: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= 1\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dt \\ &= 1 \int \frac{1}{1} dt \\ &= 1(t)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(t))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 t + c_1 \tag{1}$$




Figure 72: Slope field plot

Verification of solutions

$$y = c_2 t + c_1$$

Verified OK.

1.54.7 Solving as exact linear second order ode ode

An ode of the form

$$p(t) y'' + q(t) y' + r(t) y = s(t)$$

is exact if

$$p''(t) - q'(t) + r(t) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \\ s(x) &= 0 \end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(t) y' + (q(t) - p'(t)) y)' = s(x)$$

Integrating gives

$$p(t) y' + (q(t) - p'(t)) y = \int s(t) dt$$

Substituting the above values for p, q, r, s gives

$$y' = c_1$$

We now have a first order ode to solve which is

$$y' = c_1$$

Integrating both sides gives

$$\begin{aligned}y &= \int c_1 dt \\&= c_1 t + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 t + c_2 \tag{1}$$




Figure 73: Slope field plot

Verification of solutions

$$y = c_1 t + c_2$$

Verified OK.

1.54.8 Maple step by step solution

Let's solve

$$y'' = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

- $r = 0$
- 1st solution of the ODE
 $y_1(t) = 1$
- Repeated root, multiply $y_1(t)$ by t to ensure linear independence
 $y_2(t) = t$
- General solution of the ODE
 $y = c_1 y_1(t) + c_2 y_2(t)$
- Substitute in solutions
 $y = c_2 t + c_1$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(t),t$2)=0,y(t), singsol=all)
```

$$y(t) = c_1 t + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
DSolve[y''[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_2 t + c_1$$

1.55 problem 55

1.55.1 Solving as second order ode quadrature ode	440
1.55.2 Solving as second order linear constant coeff ode	441
1.55.3 Solving as second order ode can be made integrable ode	444
1.55.4 Solving as second order integrable as is ode	446
1.55.5 Solving as second order ode missing y ode	447
1.55.6 Solving using Kovacic algorithm	449
1.55.7 Solving as exact linear second order ode ode	454
1.55.8 Maple step by step solution	456

Internal problem ID [7099]

Internal file name [OUTPUT/6085_Sunday_June_05_2022_04_19_03_PM_25081960/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 55.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = 1$$

1.55.1 Solving as second order ode quadrature ode

The ODE can be written as

$$y'' = 1$$

Integrating once gives

$$y' = t + c_1$$

Integrating again gives

$$y = \frac{t^2}{2} + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}t^2 + c_1t + c_2 \quad (1)$$




Figure 74: Slope field plot

Verification of solutions

$$y = \frac{1}{2}t^2 + c_1t + c_2$$

Verified OK.

1.55.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = f(t)$$

Where $A = 1, B = 0, C = 0, f(t) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$.

y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 t \tag{1}$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 t + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, t\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t\}]$$

Since t is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 t^2$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{t^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2 t + c_1) + \left(\frac{t^2}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 t + c_1 + \frac{1}{2} t^2 \tag{1}$$




Figure 75: Slope field plot

Verification of solutions

$$y = c_2 t + c_1 + \frac{1}{2} t^2$$

Verified OK.

1.55.3 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' - y' = 0$$

Integrating the above w.r.t t gives

$$\int (y' y'' - y') dt = 0$$

$$\frac{y'^2}{2} - y = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{2y + 2c_1} \tag{1}$$

$$y' = -\sqrt{2y + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{2y + 2c_1}} dy = \int dt$$
$$\sqrt{2y + 2c_1} = c_2 + t$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{2y + 2c_1}} dy = \int dt$$
$$-\sqrt{2y + 2c_1} = t + c_3$$

Summary

The solution(s) found are the following

$$\sqrt{2y + 2c_1} = c_2 + t \quad (1)$$

$$-\sqrt{2y + 2c_1} = t + c_3 \quad (2)$$




Figure 76: Slope field plot

Verification of solutions

$$\sqrt{2y + 2c_1} = c_2 + t$$

Verified OK.

$$-\sqrt{2y + 2c_1} = t + c_3$$

Verified OK.

1.55.4 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t t gives

$$\int y'' dt = \int 1 dt$$

$$y' = t + c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int t + c_1 dt$$

$$= \frac{1}{2}t^2 + c_1t + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}t^2 + c_1t + c_2 \quad (1)$$




Figure 77: Slope field plot

Verification of solutions

$$y = \frac{1}{2}t^2 + c_1t + c_2$$

Verified OK.

1.55.5 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(t) = y'$$

Then

$$p'(t) = y''$$

Hence the ode becomes

$$p'(t) - 1 = 0$$

Which is now solve for $p(t)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(t) &= \int 1 \, dt \\ &= t + c_1 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = t + c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int t + c_1 \, dt \\ &= \frac{1}{2}t^2 + c_1t + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}t^2 + c_1t + c_2 \tag{1}$$




Figure 78: Slope field plot

Verification of solutions

$$y = \frac{1}{2}t^2 + c_1t + c_2$$

Verified OK.

1.55.6 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 67: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dt \\ &= 1 \int \frac{1}{1} dt \\ &= 1(t) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(t)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2 t + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, t\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t\}]$$

Since t is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 t^2$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{t^2}{2}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_2 t + c_1) + \left(\frac{t^2}{2}\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 t + c_1 + \frac{1}{2} t^2 \tag{1}$$




Figure 79: Slope field plot

Verification of solutions

$$y = c_2 t + c_1 + \frac{1}{2} t^2$$

Verified OK.

1.55.7 Solving as exact linear second order ode

An ode of the form

$$p(t) y'' + q(t) y' + r(t) y = s(t)$$

is exact if

$$p''(t) - q'(t) + r(t) = 0 \tag{1}$$

For the given ode we have

$$p(x) = 1$$

$$q(x) = 0$$

$$r(x) = 0$$

$$s(x) = 1$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(t) y' + (q(t) - p'(t)) y)' = s(x)$$

Integrating gives

$$p(t) y' + (q(t) - p'(t)) y = \int s(t) dt$$

Substituting the above values for p, q, r, s gives

$$y' = \int 1 dt$$

We now have a first order ode to solve which is

$$y' = t + c_1$$

Integrating both sides gives

$$\begin{aligned}y &= \int t + c_1 dt \\ &= \frac{1}{2}t^2 + c_1t + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}t^2 + c_1t + c_2 \tag{1}$$




Figure 80: Slope field plot

Verification of solutions

$$y = \frac{1}{2}t^2 + c_1t + c_2$$

Verified OK.

1.55.8 Maple step by step solution

Let's solve

$$y'' = 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(t) = 1$$

- Repeated root, multiply $y_1(t)$ by t to ensure linear independence

$$y_2(t) = t$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 t + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = 1$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -(\int t dt) + t(\int 1 dt)$$

- Compute integrals

$$y_p(t) = \frac{t^2}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_2 t + c_1 + \frac{1}{2} t^2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(t),t$2)=1,y(t), singsol=all)
```

$$y(t) = \frac{1}{2} t^2 + c_1 t + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 19

```
DSolve[y''[t]==1,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{t^2}{2} + c_2 t + c_1$$

1.56 problem 56

1.56.1 Solving as second order ode quadrature ode	458
1.56.2 Solving as second order linear constant coeff ode	459
1.56.3 Solving as second order integrable as is ode	462
1.56.4 Solving as second order ode missing y ode	463
1.56.5 Solving using Kovacic algorithm	464
1.56.6 Solving as exact linear second order ode ode	469
1.56.7 Maple step by step solution	470

Internal problem ID [7100]

Internal file name [OUTPUT/6086_Sunday_June_05_2022_04_19_04_PM_45789474/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 56.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = f(t)$$

1.56.1 Solving as second order ode quadrature ode

Integrating once gives

$$y' = \int f(t) dt + c_1$$

Integrating again gives

$$y = \int \left(\int f(t) dt \right) dt + c_1 x + c_2$$

Summary

The solution(s) found are the following

$$y = \int \int f(t) dt dt + c_1 t + c_2 \tag{1}$$

Verification of solutions

$$y = \int \int f(t) dt dt + c_1 t + c_2$$

Verified OK.

1.56.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = f(t)$$

Where $A = 1, B = 0, C = 0, f(t) = f(t)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 t \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 t + c_1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$y_p(t) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = 1$$

$$y_2 = t$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & t \\ \frac{d}{dt}(1) & \frac{d}{dt}(t) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & t \\ 0 & 1 \end{vmatrix}$$

Therefore

$$W = (1) (1) - (t) (0)$$

Which simplifies to

$$W = 1$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{f(t) t}{1} dt$$

Which simplifies to

$$u_1 = - \int f(t) t dt$$

Hence

$$u_1 = - \left(\int_0^t f(\alpha) \alpha d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{f(t)}{1} dt$$

Which simplifies to

$$u_2 = \int f(t) dt$$

Hence

$$u_2 = \int_0^t f(\alpha) d\alpha$$

Therefore the particular solution, from equation (1) is

$$y_p(t) = - \left(\int_0^t f(\alpha) \alpha d\alpha \right) + \left(\int_0^t f(\alpha) d\alpha \right) t$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_2 t + c_1) + \left(- \left(\int_0^t f(\alpha) \alpha d\alpha \right) + \left(\int_0^t f(\alpha) d\alpha \right) t \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 t + c_1 - \left(\int_0^t f(\alpha) \alpha d\alpha \right) + \left(\int_0^t f(\alpha) d\alpha \right) t \quad (1)$$

Verification of solutions

$$y = c_2 t + c_1 - \left(\int_0^t f(\alpha) \alpha d\alpha \right) + \left(\int_0^t f(\alpha) d\alpha \right) t$$

Verified OK.

1.56.3 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t t gives

$$\begin{aligned}\int y'' dt &= \int f(t) dt \\ y' &= \int f(t) dt + c_1\end{aligned}$$

Which is now solved for y . Integrating both sides gives

$$\begin{aligned}y &= \int \int f(t) dt + c_1 dt \\ &= \int \left(\int f(t) dt + c_1 \right) dt + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \left(\int f(t) dt + c_1 \right) dt + c_2 \quad (1)$$

Verification of solutions

$$y = \int \left(\int f(t) dt + c_1 \right) dt + c_2$$

Verified OK.

1.56.4 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(t) = y'$$

Then

$$p'(t) = y''$$

Hence the ode becomes

$$p'(t) - f(t) = 0$$

Which is now solve for $p(t)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(t) &= \int f(t) dt \\ &= \int f(t) dt + c_1 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \int f(t) dt + c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int \int f(t) dt + c_1 dt \\ &= \int \left(\int f(t) dt + c_1 \right) dt + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \left(\int f(t) dt + c_1 \right) dt + c_2 \quad (1)$$

Verification of solutions

$$y = \int \left(\int f(t) dt + c_1 \right) dt + c_2$$

Verified OK.

1.56.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 69: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= 1\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dt \\ &= 1 \int \frac{1}{1} dt \\ &= 1(t)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(t))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2 t + c_1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$y_p(t) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = 1$$

$$y_2 = t$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & t \\ \frac{d}{dt}(1) & \frac{d}{dt}(t) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & t \\ 0 & 1 \end{vmatrix}$$

Therefore

$$W = (1)(1) - (t)(0)$$

Which simplifies to

$$W = 1$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{f(t) t}{1} dt$$

Which simplifies to

$$u_1 = - \int f(t) t dt$$

Hence

$$u_1 = - \left(\int_0^t f(\alpha) \alpha d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{f(t)}{1} dt$$

Which simplifies to

$$u_2 = \int f(t) dt$$

Hence

$$u_2 = \int_0^t f(\alpha) d\alpha$$

Therefore the particular solution, from equation (1) is

$$y_p(t) = - \left(\int_0^t f(\alpha) \alpha d\alpha \right) + \left(\int_0^t f(\alpha) d\alpha \right) t$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2 t + c_1) + \left(- \left(\int_0^t f(\alpha) \alpha d\alpha \right) + \left(\int_0^t f(\alpha) d\alpha \right) t \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 t + c_1 - \left(\int_0^t f(\alpha) \alpha d\alpha \right) + \left(\int_0^t f(\alpha) d\alpha \right) t \quad (1)$$

Verification of solutions

$$y = c_2 t + c_1 - \left(\int_0^t f(\alpha) \alpha d\alpha \right) + \left(\int_0^t f(\alpha) d\alpha \right) t$$

Verified OK.

1.56.6 Solving as exact linear second order ode ode

An ode of the form

$$p(t) y'' + q(t) y' + r(t) y = s(t)$$

is exact if

$$p''(t) - q'(t) + r(t) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \\ s(x) &= f(t) \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(t) y' + (q(t) - p'(t)) y)' = s(x)$$

Integrating gives

$$p(t) y' + (q(t) - p'(t)) y = \int s(t) dt$$

Substituting the above values for p, q, r, s gives

$$y' = \int f(t) dt$$

We now have a first order ode to solve which is

$$y' = \int f(t) dt + c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int \int f(t) dt + c_1 dt \\ &= \int \left(\int f(t) dt + c_1 \right) dt + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \left(\int f(t) dt + c_1 \right) dt + c_2 \quad (1)$$

Verification of solutions

$$y = \int \left(\int f(t) dt + c_1 \right) dt + c_2$$

Verified OK.

1.56.7 Maple step by step solution

Let's solve

$$y'' = f(t)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(t) = 1$$

- Repeated root, multiply $y_1(t)$ by t to ensure linear independence

$$y_2(t) = t$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 t + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t),y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t),y_2(t))} dt \right), f(t) = f(t) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = 1$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -\left(\int f(t) t dt \right) + t \left(\int f(t) dt \right)$$

- Compute integrals

$$y_p(t) = -\left(\int f(t) t dt \right) + t \left(\int f(t) dt \right)$$

- Substitute particular solution into general solution to ODE

$$y = c_1 + c_2 t - \left(\int f(t) t dt \right) + t \left(\int f(t) dt \right)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(t),t$2)=f(t),y(t), singsol=all)
```

$$y(t) = \int \int f(t) dt dt + c_1 t + c_2$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 30

```
DSolve[y''[t]==f[t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \int_1^t \int_1^{K[2]} f(K[1]) dK[1] dK[2] + c_2 t + c_1$$

1.57 problem 57

1.57.1 Solving as second order ode quadrature ode	473
1.57.2 Solving as second order linear constant coeff ode	474
1.57.3 Solving as second order ode can be made integrable ode	477
1.57.4 Solving as second order integrable as is ode	479
1.57.5 Solving as second order ode missing y ode	481
1.57.6 Solving using Kovacic algorithm	482
1.57.7 Solving as exact linear second order ode ode	487
1.57.8 Maple step by step solution	489

Internal problem ID [7101]

Internal file name [OUTPUT/6087_Sunday_June_05_2022_04_19_05_PM_3354666/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 57.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = k$$

1.57.1 Solving as second order ode quadrature ode

The ODE can be written as

$$y'' = k$$

Integrating once gives

$$y' = kt + c_1$$

Integrating again gives

$$y = \frac{k t^2}{2} + c_1 x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}k t^2 + c_1 t + c_2 \quad (1)$$




Figure 81: Slope field plot

Verification of solutions

$$y = \frac{1}{2}k t^2 + c_1 t + c_2$$

Verified OK.

1.57.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = f(t)$$

Where $A = 1, B = 0, C = 0, f(t) = k$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$.

y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 t \tag{1}$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 t + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, t\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t\}]$$

Since t is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 t^2$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 = k$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{k}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{k t^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2 t + c_1) + \left(\frac{k t^2}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 t + c_1 + \frac{1}{2} k t^2 \quad (1)$$




Figure 82: Slope field plot

Verification of solutions

$$y = c_2 t + c_1 + \frac{1}{2} k t^2$$

Verified OK.

1.57.3 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' - y' k = 0$$

Integrating the above w.r.t t gives

$$\int (y' y'' - y' k) dt = 0$$

$$\frac{y'^2}{2} - ky = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{2ky + 2c_1} \tag{1}$$

$$y' = -\sqrt{2ky + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{2ky + 2c_1}} dy = \int dt$$
$$\frac{\sqrt{2ky + 2c_1}}{k} = t + c_2$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{2ky + 2c_1}} dy = \int dt$$
$$-\frac{\sqrt{2ky + 2c_1}}{k} = t + c_3$$

Summary

The solution(s) found are the following

$$\frac{\sqrt{2ky + 2c_1}}{k} = t + c_2 \tag{1}$$

$$-\frac{\sqrt{2ky + 2c_1}}{k} = t + c_3 \tag{2}$$




Figure 83: Slope field plot

Verification of solutions

$$\frac{\sqrt{2ky + 2c_1}}{k} = t + c_2$$

Verified OK.

$$-\frac{\sqrt{2ky + 2c_1}}{k} = t + c_3$$

Verified OK.

1.57.4 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t t gives

$$\int y'' dt = \int k dt$$

$$y' = kt + c_1$$

Which is now solved for y . Integrating both sides gives

$$\begin{aligned}y &= \int kt + c_1 dt \\ &= \frac{1}{2}kt^2 + c_1t + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}kt^2 + c_1t + c_2 \quad (1)$$




Figure 84: Slope field plot

Verification of solutions

$$y = \frac{1}{2}kt^2 + c_1t + c_2$$

Verified OK.

1.57.5 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(t) = y'$$

Then

$$p'(t) = y''$$

Hence the ode becomes

$$p'(t) - k = 0$$

Which is now solve for $p(t)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(t) &= \int k \, dt \\ &= kt + c_1 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = kt + c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int kt + c_1 \, dt \\ &= \frac{1}{2}k t^2 + c_1 t + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}k t^2 + c_1 t + c_2 \quad (1)$$




Figure 85: Slope field plot

Verification of solutions

$$y = \frac{1}{2}k t^2 + c_1 t + c_2$$

Verified OK.

1.57.6 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = 0$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(t) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 71: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dt \\ &= 1 \int \frac{1}{1} dt \\ &= 1(t) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(t))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2 t + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, t\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t\}]$$

Since t is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 t^2$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 = k$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{k}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{k t^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2 t + c_1) + \left(\frac{k t^2}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 t + c_1 + \frac{1}{2} k t^2 \tag{1}$$




Figure 86: Slope field plot

Verification of solutions

$$y = c_2 t + c_1 + \frac{1}{2} k t^2$$

Verified OK.

1.57.7 Solving as exact linear second order ode ode

An ode of the form

$$p(t) y'' + q(t) y' + r(t) y = s(t)$$

is exact if

$$p''(t) - q'(t) + r(t) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \\ s(x) &= k \end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(t) y' + (q(t) - p'(t)) y)' = s(x)$$

Integrating gives

$$p(t) y' + (q(t) - p'(t)) y = \int s(t) dt$$

Substituting the above values for p, q, r, s gives

$$y' = \int k dt$$

We now have a first order ode to solve which is

$$y' = kt + c_1$$

Integrating both sides gives

$$\begin{aligned}y &= \int kt + c_1 dt \\&= \frac{1}{2}k t^2 + c_1 t + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}k t^2 + c_1 t + c_2 \quad (1)$$




Figure 87: Slope field plot

Verification of solutions

$$y = \frac{1}{2}k t^2 + c_1 t + c_2$$

Verified OK.

1.57.8 Maple step by step solution

Let's solve

$$y'' = k$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(t) = 1$$
 - Repeated root, multiply $y_1(t)$ by t to ensure linear independence

$$y_2(t) = t$$
 - General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$
 - Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 t + y_p(t)$$
- Find a particular solution $y_p(t)$ of the ODE
- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = k \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(t), y_2(t)) = 1$$
 - Substitute functions into equation for $y_p(t)$

$$y_p(t) = k(-(\int t dt) + t(\int 1 dt))$$
 - Compute integrals

$$y_p(t) = \frac{k t^2}{2}$$
 - Substitute particular solution into general solution to ODE

$$y = c_2 t + c_1 + \frac{1}{2} k t^2$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(t),t$2)=k,y(t), singsol=all)
```

$$y(t) = \frac{1}{2}kt^2 + c_1t + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 20

```
DSolve[y''[t]==k,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{kt^2}{2} + c_2t + c_1$$

1.58 problem 58

1.58.1 Solving as first order ode lie symmetry calculated ode 492

Internal problem ID [7102]

Internal file name [OUTPUT/6088_Sunday_June_05_2022_04_19_07_PM_97370974/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 58.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _dAlembert]
```

$$y' + 4 \sin(x - y) = -4$$

1.58.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -4 \sin(x - y) - 4$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + (-4 \sin(x - y) - 4)(b_3 - a_2) - (-4 \sin(x - y) - 4)^2 a_3 + 4 \cos(x - y)(xa_2 + ya_3 + a_1) - 4 \cos(x - y)(xb_2 + yb_3 + b_1) = 0 \quad (5E)$$

Putting the above in normal form gives

$$-16 \sin(x - y)^2 a_3 + 4 \cos(x - y)xa_2 - 4 \cos(x - y)xb_2 + 4 \cos(x - y)ya_3 - 4 \cos(x - y)yb_3 + 4 \sin(x - y)a_2 - 32 \sin(x - y)a_3 - 4 \sin(x - y)b_3 + 4 \cos(x - y)a_1 - 4 \cos(x - y)b_1 + 4a_2 - 16a_3 + b_2 - 4b_3 = 0$$

Setting the numerator to zero gives

$$-16 \sin(x - y)^2 a_3 + 4 \cos(x - y)xa_2 - 4 \cos(x - y)xb_2 + 4 \cos(x - y)ya_3 - 4 \cos(x - y)yb_3 + 4 \sin(x - y)a_2 - 32 \sin(x - y)a_3 - 4 \sin(x - y)b_3 + 4 \cos(x - y)a_1 - 4 \cos(x - y)b_1 + 4a_2 - 16a_3 + b_2 - 4b_3 = 0 \quad (6E)$$

Simplifying the above gives

$$4a_2 - 24a_3 + b_2 - 4b_3 + 4 \cos(x - y)xa_2 - 4 \cos(x - y)xb_2 + 4 \cos(x - y)ya_3 - 4 \cos(x - y)yb_3 + 4 \sin(x - y)a_2 - 32 \sin(x - y)a_3 - 4 \sin(x - y)b_3 + 4 \cos(x - y)a_1 - 4 \cos(x - y)b_1 + 8a_3 \cos(2x - 2y) = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \cos(x - y), \cos(2x - 2y), \sin(x - y)\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \cos(x - y) = v_3, \cos(2x - 2y) = v_4, \sin(x - y) = v_5\}$$

The above PDE (6E) now becomes

$$4v_3v_1a_2 + 4v_3v_2a_3 - 4v_3v_1b_2 - 4v_3v_2b_3 + 4v_3a_1 + 4v_5a_2 + 8a_3v_4 - 32v_5a_3 - 4v_3b_1 - 4v_5b_3 + 4a_2 - 24a_3 + b_2 - 4b_3 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4, v_5\}$$

Equation (7E) now becomes

$$(4a_2 - 4b_2)v_1v_3 + (4a_3 - 4b_3)v_2v_3 + (4a_1 - 4b_1)v_3 + 8a_3v_4 + (4a_2 - 32a_3 - 4b_3)v_5 + 4a_2 - 24a_3 + b_2 - 4b_3 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} 8a_3 &= 0 \\ 4a_1 - 4b_1 &= 0 \\ 4a_2 - 4b_2 &= 0 \\ 4a_3 - 4b_3 &= 0 \\ 4a_2 - 32a_3 - 4b_3 &= 0 \\ 4a_2 - 24a_3 + b_2 - 4b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= b_1 \\ a_2 &= 0 \\ a_3 &= 0 \\ b_1 &= b_1 \\ b_2 &= 0 \\ b_3 &= 0 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 1 \\ \eta &= 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 1 - (-4 \sin(x - y) - 4) (1) \\ &= 5 + 4 \sin(x) \cos(y) - 4 \cos(x) \sin(y) \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{5 + 4 \sin(x) \cos(y) - 4 \cos(x) \sin(y)} dy \end{aligned}$$

Which results in

$$S = -\frac{2 \arctan\left(\frac{2(4 \sin(x)-5) \tan(\frac{y}{2})+8 \cos(x)}{2\sqrt{25-16 \sin(x)^2-16 \cos(x)^2}}\right)}{\sqrt{25-16 \sin(x)^2-16 \cos(x)^2}}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -4 \sin(x - y) - 4$$

Evaluating all the partial derivatives gives

$$R_x = 1$$

$$R_y = 0$$

$$S_x = \frac{-8 \cos(x) \tan\left(\frac{y}{2}\right) + 8 \sin(x)}{16 \left(\sin(x) - \frac{5}{4}\right)^2 \tan^2\left(\frac{y}{2}\right) + (32 \sin(x) - 40) \cos(x) \tan\left(\frac{y}{2}\right) + 16 \cos(x)^2 + 9}$$

$$S_y = \frac{1}{5 + 4 \sin(x) \cos(y) - 4 \cos(x) \sin(y)}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{-8 \cos(x) \tan\left(\frac{y}{2}\right) + 8 \sin(x)}{16 \left(\sin(x) - \frac{5}{4}\right)^2 \tan\left(\frac{y}{2}\right)^2 + (32 \sin(x) - 40) \cos(x) \tan\left(\frac{y}{2}\right) + 16 \cos(x)^2 + 9} + \frac{4 + 4 \sin(x) \cos\left(\frac{y}{2}\right)}{-5 - 4 \sin(x) \cos\left(\frac{y}{2}\right)} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{-4 \sin(R) + 4}{4 \sin(R) - 5}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{2 \arctan\left(\frac{5 \tan\left(\frac{R}{2}\right) - \frac{4}{3}}{3}\right)}{3} - R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{2 \arctan\left(\frac{(4 \sin(x) - 5) \tan\left(\frac{y}{2}\right) + \frac{4 \cos(x)}{3}}{3}\right)}{3} = \frac{2 \arctan\left(\frac{5 \tan\left(\frac{x}{2}\right) - \frac{4}{3}}{3}\right)}{3} - x + c_1$$



Which simplifies to

$$-\frac{2 \arctan\left(\frac{(4 \sin(x) - 5) \tan\left(\frac{y}{2}\right) + \frac{4 \cos(x)}{3}}{3}\right)}{3} = \frac{2 \arctan\left(\frac{5 \tan\left(\frac{x}{2}\right) - \frac{4}{3}}{3}\right)}{3} - x + c_1$$

Which gives

Expression too large to display

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -4 \sin(x - y) - 4$ 	$R = x$ $S = -\frac{2 \arctan\left(\frac{(4 \sin(x) - 5)}{3}\right)}{3}$	$\frac{dS}{dR} = \frac{-4 \sin(R) + 4}{4 \sin(R) - 5}$ 

Summary

The solution(s) found are the following

Expression too large to display (1)




Figure 88: Slope field plot

Verification of solutions

Expression too large to display

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous C
1st order, trying the canonical coordinates of the invariance group
<- 1st order, canonical coordinates successful
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)=4*sin(y(x)-x)-4,y(x), singsol=all)
```

$$y(x) = x + 2 \arctan \left(\frac{3 \tan \left(-\frac{3x}{2} + \frac{3c_1}{2} \right)}{5} + \frac{4}{5} \right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==4*Sin[y[x]-x]-4,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

1.59 problem 59

1.59.1 Solving as first order ode lie symmetry calculated ode 500

Internal problem ID [7103]

Internal file name [OUTPUT/6089_Sunday_June_05_2022_04_21_23_PM_58036799/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 59.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _dAlembert]
```

$$y' + \sin(x - y) = 0$$

1.59.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$\begin{aligned}y' &= -\sin(x - y) \\y' &= \omega(x, y)\end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 - \sin(x-y)(b_3 - a_2) - \sin(x-y)^2 a_3 + \cos(x-y)(xa_2 + ya_3 + a_1) - \cos(x-y)(xb_2 + yb_3 + b_1) = 0 \quad (5E)$$

Putting the above in normal form gives

$$-\sin(x-y)^2 a_3 + \cos(x-y)xa_2 - \cos(x-y)xb_2 + \cos(x-y)ya_3 - \cos(x-y)yb_3 + \sin(x-y)a_2 - \sin(x-y)b_3 + \cos(x-y)a_1 - \cos(x-y)b_1 + b_2 = 0$$

Setting the numerator to zero gives

$$-\sin(x-y)^2 a_3 + \cos(x-y)xa_2 - \cos(x-y)xb_2 + \cos(x-y)ya_3 - \cos(x-y)yb_3 + \sin(x-y)a_2 - \sin(x-y)b_3 + \cos(x-y)a_1 - \cos(x-y)b_1 + b_2 = 0 \quad (6E)$$

Simplifying the above gives

$$b_2 - \frac{a_3}{2} + \frac{a_3 \cos(2x-2y)}{2} + \cos(x-y)xa_2 - \cos(x-y)xb_2 + \cos(x-y)ya_3 - \cos(x-y)yb_3 + \sin(x-y)a_2 - \sin(x-y)b_3 + \cos(x-y)a_1 - \cos(x-y)b_1 = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \cos(x-y), \cos(2x-2y), \sin(x-y)\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \cos(x-y) = v_3, \cos(2x-2y) = v_4, \sin(x-y) = v_5\}$$

The above PDE (6E) now becomes

$$b_2 - \frac{1}{2}a_3 + \frac{1}{2}a_3v_4 + v_3v_1a_2 - v_3v_1b_2 + v_3v_2a_3 - v_3v_2b_3 + v_5a_2 - v_5b_3 + v_3a_1 - v_3b_1 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4, v_5\}$$

Equation (7E) now becomes

$$b_2 - \frac{a_3}{2} + (a_1 - b_1) v_3 + \frac{a_3 v_4}{2} + (a_2 - b_3) v_5 + (a_2 - b_2) v_1 v_3 + (a_3 - b_3) v_2 v_3 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} \frac{a_3}{2} &= 0 \\ a_1 - b_1 &= 0 \\ a_2 - b_2 &= 0 \\ a_2 - b_3 &= 0 \\ a_3 - b_3 &= 0 \\ b_2 - \frac{a_3}{2} &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= b_1 \\ a_2 &= 0 \\ a_3 &= 0 \\ b_1 &= b_1 \\ b_2 &= 0 \\ b_3 &= 0 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 1 \\ \eta &= 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 1 - (-\sin(x - y)) (1) \\ &= \sin(x) \cos(y) - \cos(x) \sin(y) + 1 \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\sin(x) \cos(y) - \cos(x) \sin(y) + 1} dy \end{aligned}$$

Which results in

$$S = -\frac{2}{-\tan\left(\frac{x}{2} - \frac{y}{2}\right) - 1}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\sin(x - y)$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{1}{\sin(x - y) + 1} \\ S_y &= \frac{1}{\sin(x - y) + 1} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -1 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -R + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{2}{\tan\left(\frac{x}{2} - \frac{y}{2}\right) + 1} = -x + c_1$$



Which simplifies to

$$\frac{2}{\tan\left(\frac{x}{2} - \frac{y}{2}\right) + 1} = -x + c_1$$

Which gives

$$y = x + 2 \arctan\left(\frac{c_1 - x - 2}{-x + c_1}\right)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\sin(x - y)$ 	$R = x$ $S = \frac{2}{\tan\left(\frac{x}{2} - \frac{y}{2}\right) + 1}$	$\frac{dS}{dR} = -1$ 

Summary

The solution(s) found are the following

$$y = x + 2 \arctan \left(\frac{c_1 - x - 2}{-x + c_1} \right) \quad (1)$$




Figure 89: Slope field plot

Verification of solutions

$$y = x + 2 \arctan \left(\frac{c_1 - x - 2}{-x + c_1} \right)$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
1st order, trying the canonical coordinates of the invariance group  
<- 1st order, canonical coordinates successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 23

```
dsolve(diff(y(x),x)-sin(y(x)-x)=0,y(x), singsol=all)
```

$$y(x) = x + 2 \arctan \left(\frac{c_1 - x - 2}{-x + c_1} \right)$$

✓ Solution by Mathematica

Time used: 37.233 (sec). Leaf size: 553

`DSolve[y'[x]-Sin[y[x]-x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow -2 \arccos \left(\frac{(-x + 2 + c_1) \cos \left(\frac{x}{2}\right) + (x - c_1) \sin \left(\frac{x}{2}\right)}{\sqrt{2} \sqrt{x^2 - 2(1 + c_1)x + 2 + c_1^2 + 2c_1}} \right)$$

$$y(x) \rightarrow 2 \arccos \left(\frac{(-x + 2 + c_1) \cos \left(\frac{x}{2}\right) + (x - c_1) \sin \left(\frac{x}{2}\right)}{\sqrt{2} \sqrt{x^2 - 2(1 + c_1)x + 2 + c_1^2 + 2c_1}} \right)$$

$$y(x) \rightarrow -2 \arccos \left(\frac{(x - 2 - c_1) \cos \left(\frac{x}{2}\right) + (-x + c_1) \sin \left(\frac{x}{2}\right)}{\sqrt{2} \sqrt{x^2 - 2(1 + c_1)x + 2 + c_1^2 + 2c_1}} \right)$$

$$y(x) \rightarrow 2 \arccos \left(\frac{(x - 2 - c_1) \cos \left(\frac{x}{2}\right) + (-x + c_1) \sin \left(\frac{x}{2}\right)}{\sqrt{2} \sqrt{x^2 - 2(1 + c_1)x + 2 + c_1^2 + 2c_1}} \right)$$

$$y(x) \rightarrow -2 \arccos \left(\frac{\cos \left(\frac{x}{2}\right) - \sin \left(\frac{x}{2}\right)}{\sqrt{2}} \right)$$

$$y(x) \rightarrow 2 \arccos \left(\frac{\cos \left(\frac{x}{2}\right) - \sin \left(\frac{x}{2}\right)}{\sqrt{2}} \right)$$

$$y(x) \rightarrow -2 \arccos \left(\frac{\sin \left(\frac{x}{2}\right) - \cos \left(\frac{x}{2}\right)}{\sqrt{2}} \right)$$

$$y(x) \rightarrow 2 \arccos \left(\frac{\sin \left(\frac{x}{2}\right) - \cos \left(\frac{x}{2}\right)}{\sqrt{2}} \right)$$

$$y(x) \rightarrow -2 \arccos \left(\frac{(x - 2) \cos \left(\frac{x}{2}\right) - x \sin \left(\frac{x}{2}\right)}{\sqrt{2} \sqrt{x^2 - 2x + 2}} \right)$$

$$y(x) \rightarrow 2 \arccos \left(\frac{(x - 2) \cos \left(\frac{x}{2}\right) - x \sin \left(\frac{x}{2}\right)}{\sqrt{2} \sqrt{x^2 - 2x + 2}} \right)$$

$$y(x) \rightarrow -2 \arccos \left(\frac{x \sin \left(\frac{x}{2}\right) - (x - 2) \cos \left(\frac{x}{2}\right)}{\sqrt{2} \sqrt{x^2 - 2x + 2}} \right)$$

$$y(x) \rightarrow 2 \arccos \left(\frac{x \sin \left(\frac{x}{2}\right) - (x - 2) \cos \left(\frac{x}{2}\right)}{\sqrt{2} \sqrt{x^2 - 2x + 2}} \right)$$

1.60 problem 60

1.60.1 Solving as second order ode quadrature ode	508
1.60.2 Solving as second order linear constant coeff ode	509
1.60.3 Solving as second order integrable as is ode	512
1.60.4 Solving as second order ode missing y ode	513
1.60.5 Solving using Kovacic algorithm	515
1.60.6 Solving as exact linear second order ode ode	519
1.60.7 Maple step by step solution	521

Internal problem ID [7104]

Internal file name [OUTPUT/6090_Sunday_June_05_2022_04_21_28_PM_90558027/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 60.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = 4 \sin(x) - 4$$

1.60.1 Solving as second order ode quadrature ode

Integrating once gives

$$y' = -4x - 4 \cos(x) + c_1$$

Integrating again gives

$$y = -2x^2 - 4 \sin(x) + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = -2x^2 - 4 \sin(x) + c_1x + c_2 \tag{1}$$




Figure 90: Slope field plot

Verification of solutions

$$y = -2x^2 - 4 \sin(x) + c_1x + c_2$$

Verified OK.

1.60.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 0, f(x) = 4 \sin(x) - 4$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$4 \sin(x) - 4$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}, \{\cos(x), \sin(x)\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2\}, \{\cos(x), \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 + A_2 \cos(x) + A_3 \sin(x)$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 - A_2 \cos(x) - A_3 \sin(x) = 4 \sin(x) - 4$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -2, A_2 = 0, A_3 = -4]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2x^2 - 4 \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2 x + c_1) + (-2x^2 - 4 \sin(x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 x + c_1 - 2x^2 - 4 \sin(x) \tag{1}$$




Figure 91: Slope field plot

Verification of solutions

$$y = c_2x + c_1 - 2x^2 - 4 \sin(x)$$

Verified OK.

1.60.3 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = \int (4 \sin(x) - 4) dx$$

$$y' = -4x - 4 \cos(x) + c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int -4x - 4 \cos(x) + c_1 dx$$

$$= -2x^2 - 4 \sin(x) + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = -2x^2 - 4 \sin(x) + c_1x + c_2 \tag{1}$$




Figure 92: Slope field plot

Verification of solutions

$$y = -2x^2 - 4 \sin(x) + c_1 x + c_2$$

Verified OK.

1.60.4 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 4 \sin(x) + 4 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(x) &= \int 4 \sin(x) - 4 \, dx \\ &= -4x - 4 \cos(x) + c_1 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = -4x - 4 \cos(x) + c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int -4x - 4 \cos(x) + c_1 \, dx \\ &= -2x^2 - 4 \sin(x) + c_1x + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -2x^2 - 4 \sin(x) + c_1x + c_2 \tag{1}$$




Figure 93: Slope field plot

Verification of solutions

$$y = -2x^2 - 4 \sin(x) + c_1x + c_2$$

Verified OK.

1.60.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 73: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= 1\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2 x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$4 \sin(x) - 4$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}, \{\cos(x), \sin(x)\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2\}, \{\cos(x), \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 + A_2 \cos(x) + A_3 \sin(x)$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 - A_2 \cos(x) - A_3 \sin(x) = 4 \sin(x) - 4$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -2, A_2 = 0, A_3 = -4]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2x^2 - 4 \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2 x + c_1) + (-2x^2 - 4 \sin(x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2x + c_1 - 2x^2 - 4 \sin(x) \quad (1)$$




Figure 94: Slope field plot

Verification of solutions

$$y = c_2x + c_1 - 2x^2 - 4 \sin(x)$$

Verified OK.

1.60.6 Solving as exact linear second order ode ode

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned}p(x) &= 1 \\q(x) &= 0 \\r(x) &= 0 \\s(x) &= 4 \sin(x) - 4\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = \int 4 \sin(x) - 4 dx$$

We now have a first order ode to solve which is

$$y' = -4x - 4 \cos(x) + c_1$$

Integrating both sides gives

$$\begin{aligned}y &= \int -4x - 4 \cos(x) + c_1 dx \\&= -2x^2 - 4 \sin(x) + c_1x + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = -2x^2 - 4 \sin(x) + c_1x + c_2 \tag{1}$$




Figure 95: Slope field plot

Verification of solutions

$$y = -2x^2 - 4 \sin(x) + c_1x + c_2$$

Verified OK.

1.60.7 Maple step by step solution

Let's solve

$$y'' = 4 \sin(x) - 4$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 4 \sin(x) - 4 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -4 \left(\int x(\sin(x) - 1) dx \right) + 4x \left(\int (\sin(x) - 1) dx \right)$$

- Compute integrals

$$y_p(x) = -2x^2 - 4 \sin(x)$$

- Substitute particular solution into general solution to ODE

$$y = c_2 x + c_1 - 2x^2 - 4 \sin(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)=4*sin(x)-4,y(x), singsol=all)
```

$$y(x) = -2x^2 - 4 \sin(x) + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 21

```
DSolve[y''[x]==4*Sin[x]-4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x^2 - 4 \sin(x) + c_2x + c_1$$

1.61 problem 61

1.61.1 Maple step by step solution 525

Internal problem ID [7105]

Internal file name [OUTPUT/6091_Sunday_June_05_2022_04_21_29_PM_82852476/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 61.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "algebraic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$yy'' = 0$$

The ode

$$yy'' = 0$$

Gives the following equations

$$y = 0 \tag{1}$$

$$y'' = 0 \tag{2}$$

Each of the above equations is now solved.

Solving ODE (1) Since $y = 0$, is missing derivative in y then it is an algebraic equation.

Solving for y .

$$y = 0$$

Summary

The solution(s) found are the following

$$y = 0 \quad (1)$$

Verification of solutions

$$y = 0$$

Verified OK.

Solving ODE (2) Integrating twice gives the solution

$$y = c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = c_1x + c_2 \quad (1)$$

Verification of solutions

$$y = c_1x + c_2$$

Verified OK.

Summary

The solution(s) found are the following

$$y = c_1x + c_2 \quad (1)$$

Verification of solutions

$$y = c_1x + c_2$$

Verified OK.

1.61.1 Maple step by step solution

Let's solve

$$yy'' = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = 0$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_2 x + c_1$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(y(x)*diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = 0$$
$$y(x) = c_1 x + c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 17

```
DSolve[y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_2x + c_1$$

1.62 problem 62

1.62.1 Solving as second order ode missing x ode 528

Internal problem ID [7106]

Internal file name [OUTPUT/6092_Sunday_June_05_2022_04_21_31_PM_77928761/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 62.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]

$$yy'' = 1$$

1.62.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable.

Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) = 1$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{1}{yp} \end{aligned}$$

Where $f(y) = \frac{1}{y}$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= \frac{1}{y} dy \\ \int \frac{1}{p} dp &= \int \frac{1}{y} dy \\ \frac{p^2}{2} &= \ln(y) + c_1 \end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} - \ln(y) - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} - \ln(y) - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{2 \ln(y) + 2c_1} \quad (1)$$

$$y' = -\sqrt{2 \ln(y) + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} \int \frac{1}{\sqrt{2 \ln(y) + 2c_1}} dy &= \int dx \\ \int \frac{1}{\sqrt{2 \ln(a) + 2c_1}} da &= x + c_2 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{2 \ln (y)+2 c_1}} d y = \int d x$$
$$-\left(\int^y \frac{1}{\sqrt{2 \ln (a)+2 c_1}} d a\right)=x+c_3$$

Summary

The solution(s) found are the following

$$\int^y \frac{1}{\sqrt{2 \ln (a)+2 c_1}} d a = x+c_2 \quad (1)$$

$$-\left(\int^y \frac{1}{\sqrt{2 \ln (a)+2 c_1}} d a\right)=x+c_3 \quad (2)$$

Verification of solutions

$$\int^y \frac{1}{\sqrt{2 \ln (a)+2 c_1}} d a = x+c_2$$

Verified OK.

$$-\left(\int^y \frac{1}{\sqrt{2 \ln (a)+2 c_1}} d a\right)=x+c_3$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
trying differential order: 2; missing variables  
, `-> Computing symmetries using: way = 3  
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-1/_a = 0, _b(_a), HINT = [[_a, 0  
    symmetry methods on request  
, `1st order, trying reduction of order with given symmetries: `_[_a, 0]
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 51

```
dsolve(y(x)*diff(y(x),x$2)=1,y(x), singsol=all)
```

$$\int^{y(x)} \frac{1}{\sqrt{2 \ln(_a) - c_1}} d_a - x - c_2 = 0$$
$$- \left(\int^{y(x)} \frac{1}{\sqrt{2 \ln(_a) - c_1}} d_a \right) - x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 60.072 (sec). Leaf size: 93

```
DSolve[y[x]*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \exp \left(-\operatorname{erf}^{-1} \left(-i \sqrt{\frac{2}{\pi}} \sqrt{e^{c_1(x+c_2)^2}} \right)^2 - \frac{c_1}{2} \right)$$
$$y(x) \rightarrow \exp \left(-\operatorname{erf}^{-1} \left(i \sqrt{\frac{2}{\pi}} \sqrt{e^{c_1(x+c_2)^2}} \right)^2 - \frac{c_1}{2} \right)$$

1.63 problem 63

Internal problem ID [7107]

Internal file name [OUTPUT/6093_Sunday_June_05_2022_04_21_33_PM_4614970/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 63.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$yy'' = x$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integrating f
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular cases
trying symmetries linear in x and y(x)
trying differential order: 2; exact nonlinear
trying 2nd order, integrating factor of the form mu(y)
trying 2nd order, integrating factor of the form mu(x,y)
trying 2nd order, integrating factor of the form mu(x,y)/(y)^n, only the general case
trying 2nd order, integrating factor of the form mu(y,y)
-> Calling odsolve with the ODE`, -(_y1^3*x-1)*y(x)/(x*_y1^3)+(1/3)*(3*(diff(y(x), x))*x+2*_
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
trying differential order: 2; mu polynomial in y
trying 2nd order, integrating factor of the form mu(x,y)
differential order: 2; looking for linear symmetries
differential order: 2; found: 1 linear symmetries. Trying reduction of order
`, `2nd order, trying reduction of order with given symmetries: `[x, 3/2*y]
```

X Solution by Maple

```
dsolve(y(x)*diff(y(x),x$2)=x,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

1.64 problem 64

Internal problem ID [7108]

Internal file name [OUTPUT/6094_Sunday_June_05_2022_04_21_36_PM_24970600/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 64.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y^2 y'' = x$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integrating f
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular cases
trying symmetries linear in x and y(x)
trying differential order: 2; exact nonlinear
trying 2nd order, integrating factor of the form mu(y)
trying 2nd order, integrating factor of the form mu(x,y)
trying 2nd order, integrating factor of the form mu(x,y)/(y)^n, only the general case
trying 2nd order, integrating factor of the form mu(y,y)
-> Calling odsolve with the ODE`, -(_y1^3-4)*y(x)/_y1^3+2*((diff(y(x), x))*x+_y1)/_y1^3, y(x)
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
trying differential order: 2; mu polynomial in y
trying 2nd order, integrating factor of the form mu(x,y)
differential order: 2; looking for linear symmetries
differential order: 2; found: 1 linear symmetries. Trying reduction of order
`, `2nd order, trying reduction of order with given symmetries: `[x, y]
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 106

```
dsolve(y(x)^2*diff(y(x),x$2)=x,y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(\ln(x) \right. \\ \left. + 2^{\frac{1}{3}} \left(\int^{-Z} \frac{1}{2^{\frac{1}{3}} \sqrt[3]{f} + 2 \text{RootOf} \left(\text{AiryBi} \left(\frac{2-Z^2 \sqrt[3]{f} + 2^{\frac{2}{3}}}{2 \sqrt[3]{f}} \right) c_1 - Z \text{AiryAi} \left(\frac{2-Z^2 \sqrt[3]{f} + 2^{\frac{2}{3}}}{2 \sqrt[3]{f}} \right) + \text{AiryBi} \left(1, \frac{2-Z^2 \sqrt[3]{f} + 2^{\frac{2}{3}}}{2 \sqrt[3]{f}} \right) - c_2 \right) x} \right) \right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]^2*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

1.65 problem 65

1.65.1 Maple step by step solution 539

Internal problem ID [7109]

Internal file name [OUTPUT/6095_Sunday_June_05_2022_04_21_37_PM_79446194/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 65.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "algebraic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y^2 y'' = 0$$

The ode

$$y^2 y'' = 0$$

Gives the following equations

$$y^2 = 0 \tag{1}$$

$$y'' = 0 \tag{2}$$

Each of the above equations is now solved.

Solving ODE (1) Since $y^2 = 0$, is missing derivative in y then it is an algebraic equation.

Solving for y .

$$y = 0$$

Summary

The solution(s) found are the following

$$y = 0 \quad (1)$$

Verification of solutions

$$y = 0$$

Verified OK.

Solving ODE (2) Integrating twice gives the solution

$$y = c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = c_1x + c_2 \quad (1)$$

Verification of solutions

$$y = c_1x + c_2$$

Verified OK.

Summary

The solution(s) found are the following

$$y = c_1x + c_2 \quad (1)$$

Verification of solutions

$$y = c_1x + c_2$$

Verified OK.

1.65.1 Maple step by step solution

Let's solve

$$y^2y'' = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = 0$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_2 x + c_1$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(y(x)^2*diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = 0$$
$$y(x) = c_1 x + c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 17

```
DSolve[y[x]^2*y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_2x + c_1$$

1.66 problem 66

Internal problem ID [7110]

Internal file name [OUTPUT/6096_Sunday_June_05_2022_04_21_39_PM_99630569/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 66.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

[NONE]

Unable to solve or complete the solution.

$$3yy'' = \sin(x)$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integrating f
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular cases
trying symmetries linear in x and y(x)
trying differential order: 2; exact nonlinear
trying 2nd order, integrating factor of the form mu(y)
trying 2nd order, integrating factor of the form mu(x,y)
trying 2nd order, integrating factor of the form mu(x,y)/(y)^n, only the general case
trying 2nd order, integrating factor of the form mu(y,y)
trying differential order: 2; mu polynomial in y
trying 2nd order, integrating factor of the form mu(x,y)
differential order: 2; looking for linear symmetries
-> trying 2nd order, the S-function method
  -> trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for the S-
  -> trying 2nd order, the S-function method
  -> trying 2nd order, No Point Symmetries Class V
  -> trying 2nd order, No Point Symmetries Class V
  -> trying 2nd order, No Point Symmetries Class V
trying 2nd order, integrating factor of the form mu(x,y)/(y)^n, only the general case
-> trying 2nd order, dynamical_symmetries, only a reduction of order through one integrating
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 5
`, `-> Computing symmetries using: way = formal`
```

X Solution by Maple

```
dsolve(3*y(x)*diff(y(x),x$2)=sin(x),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[3*y[x]*y''[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

1.67 problem 67

1.67.1 Solving as second order ode missing x ode 545

Internal problem ID [7111]

Internal file name [OUTPUT/6097_Sunday_June_05_2022_04_21_41_PM_30241354/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 67.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]
```

$$3yy'' + y = 5$$

1.67.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$3yp(y) \left(\frac{d}{dy} p(y) \right) + y = 5$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{y-5}{3yp} \end{aligned}$$

Where $f(y) = -\frac{y-5}{3y}$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{1}{p}} dp &= -\frac{y-5}{3y} dy \\ \int \frac{1}{\frac{1}{p}} dp &= \int -\frac{y-5}{3y} dy \\ \frac{p^2}{2} &= -\frac{y}{3} + \frac{5 \ln(y)}{3} + c_1 \end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} + \frac{y}{3} - \frac{5 \ln(y)}{3} - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} + \frac{y}{3} - \frac{5 \ln(y)}{3} - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{-6y + 30 \ln(y) + 18c_1}}{3} \quad (1)$$

$$y' = -\frac{\sqrt{-6y + 30 \ln(y) + 18c_1}}{3} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} \int \frac{3}{\sqrt{-6y + 30 \ln(y) + 18c_1}} dy &= \int dx \\ 3 \left(\int^y \frac{1}{\sqrt{-6_a + 30 \ln(_a) + 18c_1}} d_a \right) &= x + c_2 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{3}{\sqrt{-6y + 30 \ln(y) + 18c_1}} dy = \int dx$$
$$-3 \left(\int^y \frac{1}{\sqrt{-6_a + 30 \ln(_a) + 18c_1}} d_a \right) = x + c_3$$

Summary

The solution(s) found are the following

$$3 \left(\int^y \frac{1}{\sqrt{-6_a + 30 \ln(_a) + 18c_1}} d_a \right) = x + c_2 \quad (1)$$

$$-3 \left(\int^y \frac{1}{\sqrt{-6_a + 30 \ln(_a) + 18c_1}} d_a \right) = x + c_3 \quad (2)$$

Verification of solutions

$$3 \left(\int^y \frac{1}{\sqrt{-6_a + 30 \ln(_a) + 18c_1}} d_a \right) = x + c_2$$

Verified OK.

$$-3 \left(\int^y \frac{1}{\sqrt{-6_a + 30 \ln(_a) + 18c_1}} d_a \right) = x + c_3$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+(1/3)*(_a-5)/_a = 0, _b(_a)` *
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 59

```
dsolve(3*y(x)*diff(y(x),x$2)+y(x)=5,y(x), singsol=all)
```

$$-3 \left(\int^{y(x)} \frac{1}{\sqrt{30 \ln(_a) + 9c_1 - 6_a}} d_a \right) - x - c_2 = 0$$
$$3 \left(\int^{y(x)} \frac{1}{\sqrt{30 \ln(_a) + 9c_1 - 6_a}} d_a \right) - x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 0.333 (sec). Leaf size: 41

```
DSolve[3*y[x]*y'[x]+y[x]==5,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\int_1^{y(x)} \frac{1}{\sqrt{c_1 + \frac{2}{3}(5 \log(K[1]) - K[1])}} dK[1]^2 = (x + c_2)^2, y(x) \right]$$

1.68 problem 68

1.68.1 Solving as second order ode missing x ode 550

Internal problem ID [7112]

Internal file name [OUTPUT/6098_Sunday_June_05_2022_04_21_44_PM_6504858/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 68.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]
```

$$ayy'' + by = c$$

1.68.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable.

Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$ayp(y) \left(\frac{d}{dy} p(y) \right) + by = c$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{by-c}{ayp} \end{aligned}$$

Where $f(y) = -\frac{by-c}{ay}$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{1}{p}} dp &= -\frac{by-c}{ay} dy \\ \int \frac{1}{\frac{1}{p}} dp &= \int -\frac{by-c}{ay} dy \\ \frac{p^2}{2} &= \frac{c \ln(y)}{a} - \frac{yb}{a} + c_1 \end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} - \frac{c \ln(y)}{a} + \frac{yb}{a} - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} - \frac{c \ln(y)}{a} + \frac{yb}{a} - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{-2a(by - c \ln(y) - c_1a)}}{a} \quad (1)$$

$$y' = -\frac{\sqrt{-2a(by - c \ln(y) - c_1a)}}{a} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} \int \frac{a}{\sqrt{-2a(by - c \ln(y) - c_1a)}} dy &= \int dx \\ \int^y \frac{a}{\sqrt{-2a(b_a - c \ln(_a) - c_1a)}} d_a &= x + c_2 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{a}{\sqrt{-2a(by - c \ln(y) - c_1a)}} dy = \int dx$$
$$-\left(\int^y \frac{a}{\sqrt{-2a(b_a - c \ln(_a) - c_1a)}} d_a\right) = x + c_3$$

Summary

The solution(s) found are the following

$$\int^y \frac{a}{\sqrt{-2a(b_a - c \ln(_a) - c_1a)}} d_a = x + c_2 \quad (1)$$

$$-\left(\int^y \frac{a}{\sqrt{-2a(b_a - c \ln(_a) - c_1a)}} d_a\right) = x + c_3 \quad (2)$$

Verification of solutions

$$\int^y \frac{a}{\sqrt{-2a(b_a - c \ln(_a) - c_1a)}} d_a = x + c_2$$

Verified OK.

$$-\left(\int^y \frac{a}{\sqrt{-2a(b_a - c \ln(_a) - c_1a)}} d_a\right) = x + c_3$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+(_a*b-c)/(_a*a) = 0, _b(_a)` *
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 68

```
dsolve(a*y(x)*diff(y(x),x$2)+b*y(x)=c,y(x), singsol=all)
```

$$a \left(\int^{y(x)} \frac{1}{\sqrt{a(2c \ln(_a) + c_1 a - 2_ab)}} d_a \right) - x - c_2 = 0$$
$$-a \left(\int^{y(x)} \frac{1}{\sqrt{a(2c \ln(_a) + c_1 a - 2_ab)}} d_a \right) - x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 0.43 (sec). Leaf size: 43

```
DSolve[a*y[x]*y'[x]+b*y[x]==c,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\int_1^{y(x)} \frac{1}{\sqrt{c_1 + \frac{2(c \log(K[1]) - bK[1])}{a}}} dK[1]^2 = (x + c_2)^2, y(x) \right]$$

1.69 problem 69

1.69.1 Solving as second order ode missing x ode 555

Internal problem ID [7113]

Internal file name [OUTPUT/6099_Sunday_June_05_2022_04_21_48_PM_25779835/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 69.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]
```

$$ay^2y'' + by^2 = c$$

1.69.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable.

Using

$$y' = p(y)$$

Then

$$\begin{aligned}y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy}\end{aligned}$$

Hence the ode becomes

$$ay^2p(y) \left(\frac{d}{dy}p(y) \right) + by^2 = c$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{by^2 - c}{ay^2p} \end{aligned}$$

Where $f(y) = -\frac{by^2 - c}{ay^2}$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\frac{by^2 - c}{ay^2} dy \\ \int \frac{1}{p} dp &= \int -\frac{by^2 - c}{ay^2} dy \\ \frac{p^2}{2} &= -\frac{by + \frac{c}{y}}{a} + c_1 \end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} + \frac{by + \frac{c}{y}}{a} - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} + \frac{by + \frac{c}{y}}{a} - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{-2ay(by^2 - c_1ay + c)}}{ay} \tag{1}$$

$$y' = -\frac{\sqrt{-2ay(by^2 - c_1ay + c)}}{ay} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} \int \frac{ay}{\sqrt{-2ay(-ac_1y + by^2 + c)}} dy &= \int dx \\ \int^y \frac{a_a}{\sqrt{-2a_a(-a^2b - _aac_1 + c)}} d_a &= x + c_2 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{ay}{\sqrt{-2ay(-ac_1y + by^2 + c)}} dy = \int dx$$
$$\int^y -\frac{a_a}{\sqrt{-2a_a(a^2b - aac_1 + c)}} d_a = x + c_3$$

Summary

The solution(s) found are the following

$$\int^y \frac{a_a}{\sqrt{-2a_a(a^2b - aac_1 + c)}} d_a = x + c_2 \quad (1)$$

$$\int^y -\frac{a_a}{\sqrt{-2a_a(a^2b - aac_1 + c)}} d_a = x + c_3 \quad (2)$$

Verification of solutions

$$\int^y \frac{a_a}{\sqrt{-2a_a(a^2b - aac_1 + c)}} d_a = x + c_2$$

Verified OK.

$$\int^y -\frac{a_a}{\sqrt{-2a_a(a^2b - aac_1 + c)}} d_a = x + c_3$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+(_a^2*b-c)/(_a^2*a) = 0, _b(_a)`
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 76

```
dsolve(a*y(x)^2*diff(y(x),x$2)+b*y(x)^2=c,y(x), singsol=all)
```

$$a \left(\int^{y(x)} \frac{-a}{\sqrt{-aa(-2b_a^2 + _aac_1 - 2c)}} d_a \right) - x - c_2 = 0$$
$$-a \left(\int^{y(x)} \frac{-a}{\sqrt{-aa(-2b_a^2 + _aac_1 - 2c)}} d_a \right) - x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 0.801 (sec). Leaf size: 346

```
DSolve[a*y[x]^2*y'[x]+b*y[x]^2==c,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[-\frac{(\sqrt{-16bc + a^2c_1^2} - ac_1) (\sqrt{-16bc + a^2c_1^2} + ac_1)^2 \left(1 + \frac{4by(x)}{\sqrt{-16bc + a^2c_1^2} - ac_1}\right) \left(1 - \frac{4by(x)}{\sqrt{-16bc + a^2c_1^2} + ac_1}\right)}{\dots} \right]$$

1.70 problem 70

1.70.1 Maple step by step solution 562

Internal problem ID [7114]

Internal file name [OUTPUT/6100_Sunday_June_05_2022_04_21_54_PM_34240908/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 70.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "algebraic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$ayy'' + by = 0$$

The ode

$$ayy'' + by = 0$$

is factored to

$$y(ay'' + b) = 0$$

Which gives the following equations

$$y = 0 \tag{1}$$

$$ay'' + b = 0 \tag{2}$$

Each of the above equations is now solved.

Solving ODE (1) Since $y = 0$, is missing derivative in y then it is an algebraic equation.
Solving for y .

$$y = 0$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$

Verification of solutions

$$y = 0$$

Verified OK.

Solving ODE (2) The ODE can be written as

$$y'' = -\frac{b}{a}$$

Integrating once gives

$$y' = -\frac{bx}{a} + c_1$$

Integrating again gives

$$y = -\frac{bx^2}{2a} + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = -\frac{bx^2}{2a} + c_1x + c_2 \tag{1}$$

Verification of solutions

$$y = -\frac{bx^2}{2a} + c_1x + c_2$$

Verified OK.

Summary

The solution(s) found are the following

$$y = -\frac{bx^2}{2a} + c_1x + c_2 \tag{1}$$

Verification of solutions

$$y = -\frac{bx^2}{2a} + c_1x + c_2$$

Verified OK.

1.70.1 Maple step by step solution

Let's solve

$$ayy'' + by = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{b}{a}$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = -\frac{b}{a} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{b(\int 1 dx)x - (\int x dx)}{a}$$

- Compute integrals

$$y_p(x) = -\frac{bx^2}{2a}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 + c_2x - \frac{bx^2}{2a}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(a*y(x)*diff(y(x),x$2)+b*y(x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = -\frac{bx^2}{2a} + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 28

```
DSolve[a*y[x]*y'[x]+b*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\frac{bx^2}{2a} + c_2x + c_1$$

1.71 problem 71

1.71.1 Solution using Matrix exponential method	564
1.71.2 Solution using explicit Eigenvalue and Eigenvector method . . .	565
1.71.3 Maple step by step solution	572

Internal problem ID [7115]

Internal file name [OUTPUT/6101_Sunday_June_05_2022_04_21_56_PM_67912897/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 71.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"system of linear ODEs"**

Solve

$$\begin{aligned}x'(t) &= 9x(t) + 4y(t) \\y'(t) &= -6x(t) - y(t) \\z'(t) &= 6x(t) + 4y(t) + 3z(t)\end{aligned}$$

1.71.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} -2e^{3t} + 3e^{5t} & -2e^{3t} + 2e^{5t} & 0 \\ -3e^{5t} + 3e^{3t} & 3e^{3t} - 2e^{5t} & 0 \\ 3e^{5t} - 3e^{3t} & -2e^{3t} + 2e^{5t} & e^{3t} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} -2e^{3t} + 3e^{5t} & -2e^{3t} + 2e^{5t} & 0 \\ -3e^{5t} + 3e^{3t} & 3e^{3t} - 2e^{5t} & 0 \\ 3e^{5t} - 3e^{3t} & -2e^{3t} + 2e^{5t} & e^{3t} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \\
 &= \begin{bmatrix} (-2e^{3t} + 3e^{5t})c_1 + (-2e^{3t} + 2e^{5t})c_2 \\ (-3e^{5t} + 3e^{3t})c_1 + (3e^{3t} - 2e^{5t})c_2 \\ (3e^{5t} - 3e^{3t})c_1 + (-2e^{3t} + 2e^{5t})c_2 + e^{3t}c_3 \end{bmatrix} \\
 &= \begin{bmatrix} (-2c_1 - 2c_2)e^{3t} + 3(c_1 + \frac{2c_2}{3})e^{5t} \\ (3c_1 + 3c_2)e^{3t} - 3(c_1 + \frac{2c_2}{3})e^{5t} \\ (-3c_1 - 2c_2 + c_3)e^{3t} + 3(c_1 + \frac{2c_2}{3})e^{5t} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

1.71.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 9 - \lambda & 4 & 0 \\ -6 & -1 - \lambda & 0 \\ 6 & 4 & 3 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^3 - 11\lambda^2 + 39\lambda - 45 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 3$$

$$\lambda_2 = 5$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
3	1	real eigenvalue
5	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 6 & 4 & 0 \\ -6 & -4 & 0 \\ 6 & 4 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} 6 & 4 & 0 & 0 \\ -6 & -4 & 0 & 0 \\ 6 & 4 & 0 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{ccc|c} 6 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 6 & 4 & 0 & 0 \end{array} \right]$$

$$R_3 = R_3 - R_1 \implies \left[\begin{array}{ccc|c} 6 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 6 & 4 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2, v_3\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Let $v_3 = s$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -\frac{2t}{3}\}$

Hence the solution is

$$\begin{bmatrix} -\frac{2t}{3} \\ t \\ s \end{bmatrix} = \begin{bmatrix} -\frac{2t}{3} \\ t \\ s \end{bmatrix}$$

Since there are two free Variable, we have found two eigenvectors associated with this eigenvalue. The above can be written as

$$\begin{aligned} \begin{bmatrix} -\frac{2t}{3} \\ t \\ s \end{bmatrix} &= \begin{bmatrix} -\frac{2t}{3} \\ t \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ s \end{bmatrix} \\ &= t \begin{bmatrix} -\frac{2}{3} \\ 1 \\ 0 \end{bmatrix} + s \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \end{aligned}$$

By letting $t = 1$ and $s = 1$ then the above becomes

$$\begin{bmatrix} -\frac{2t}{3} \\ t \\ s \end{bmatrix} = \begin{bmatrix} -\frac{2}{3} \\ 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Hence the two eigenvectors associated with this eigenvalue are

$$\left(\begin{bmatrix} -\frac{2}{3} \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)$$

Which are normalized to

$$\left(\begin{bmatrix} -2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)$$

Considering the eigenvalue $\lambda_2 = 5$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} - (5) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 4 & 0 \\ -6 & -6 & 0 \\ 6 & 4 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} 4 & 4 & 0 & 0 \\ -6 & -6 & 0 & 0 \\ 6 & 4 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{3R_1}{2} \implies \left[\begin{array}{ccc|c} 4 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 6 & 4 & -2 & 0 \end{array} \right]$$

$$R_3 = R_3 - \frac{3R_1}{2} \implies \left[\begin{array}{ccc|c} 4 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & -2 & -2 & 0 \end{array} \right]$$

Since the current pivot $A(2,2)$ is zero, then the current pivot row is replaced with a row with a non-zero pivot. Swapping row 2 and row 3 gives

$$\left[\begin{array}{ccc|c} 4 & 4 & 0 & 0 \\ 0 & -2 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 4 & 4 & 0 \\ 0 & -2 & -2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = t, v_2 = -t\}$

Hence the solution is

$$\begin{bmatrix} t \\ -t \\ t \end{bmatrix} = \begin{bmatrix} t \\ -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ -t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ -t \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
3	2	2	No	$\begin{bmatrix} 0 & -\frac{2}{3} \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$
5	1	1	No	$\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 3 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 96: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2 which is the same as its geometric

multiplicity 2, then it is complete eigenvalue and this falls into case 1 shown above. Hence the corresponding eigenvector basis are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{3t} \\ &= \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} e^{3t}\end{aligned}$$

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{3t} \\ &= \begin{bmatrix} -\frac{2}{3} \\ 1 \\ 0 \end{bmatrix} e^{3t}\end{aligned}$$

Since eigenvalue 5 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_3(t) &= \vec{v}_3 e^{5t} \\ &= \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} e^{5t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + c_3 \vec{x}_3(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = c_1 \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} -\frac{2e^{3t}}{3} \\ e^{3t} \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} e^{5t} \\ -e^{5t} \\ e^{5t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} -\frac{2c_2 e^{3t}}{3} + c_3 e^{5t} \\ c_2 e^{3t} - c_3 e^{5t} \\ c_1 e^{3t} + c_3 e^{5t} \end{bmatrix}$$

1.71.3 Maple step by step solution

Let's solve

$$[x'(t) = 9x(t) + 4y(t), y'(t) = -6x(t) - y(t), z'(t) = 6x(t) + 4y(t) + 3z(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \right], \left[\begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix} \right], \left[\begin{bmatrix} 5 \\ -1 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[3, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 3

$$\vec{x}_1(t) = e^{3t} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 3$ is the eigenvalue, and

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained
- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 3

$$\left(\begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} - 3 \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 3

$$\vec{x}_2(t) = e^{3t} \cdot \left(t \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right)$$

- Consider eigenpair

$$\left[5, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_3 = e^{5t} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + c_3 \vec{x}_3$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{3t} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + c_2 e^{3t} \cdot \left(t \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right) + c_3 e^{5t} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} c_3 e^{5t} \\ -c_3 e^{5t} \\ (c_2 t + c_1) e^{3t} + c_3 e^{5t} \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = c_3 e^{5t}, y(t) = -c_3 e^{5t}, z(t) = (c_2 t + c_1) e^{3t} + c_3 e^{5t}\}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 58

```
dsolve([diff(x(t),t)=9*x(t)+4*y(t),diff(y(t),t)=-6*x(t)-y(t),diff(z(t),t)=6*x(t)+4*y(t)+3*z(t))
```

$$\begin{aligned}x(t) &= c_2 e^{3t} + c_3 e^{5t} \\y(t) &= -\frac{3c_2 e^{3t}}{2} - c_3 e^{5t} \\z(t) &= c_2 e^{3t} + c_3 e^{5t} + c_1 e^{3t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 103

```
DSolve[{x'[t]==9*x[t]+4*y[t],y'[t]==-6*x[t]-y[t],z'[t]==6*x[t]+4*y[t]+3*z[t]},{x[t],y[t],z[t]
```

$$\begin{aligned}x(t) &\rightarrow e^{3t}(c_1(3e^{2t} - 2) + 2c_2(e^{2t} - 1)) \\y(t) &\rightarrow -e^{3t}(3c_1(e^{2t} - 1) + c_2(2e^{2t} - 3)) \\z(t) &\rightarrow e^{3t}(3c_1(e^{2t} - 1) + 2c_2(e^{2t} - 1) + c_3)\end{aligned}$$

1.72 problem 72

1.72.1 Solution using Matrix exponential method	576
1.72.2 Solution using explicit Eigenvalue and Eigenvector method . . .	577
1.72.3 Maple step by step solution	582

Internal problem ID [7116]

Internal file name [OUTPUT/6102_Sunday_June_05_2022_04_21_58_PM_12651686/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 72.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"system of linear ODEs"**

Solve

$$\begin{aligned}x'(t) &= x(t) - 3y(t) \\y'(t) &= 3x(t) + 7y(t)\end{aligned}$$

1.72.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{4t}(1 - 3t) & -3t e^{4t} \\ 3t e^{4t} & e^{4t}(1 + 3t) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^{4t}(1-3t) & -3te^{4t} \\ 3te^{4t} & e^{4t}(1+3t) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^{4t}(1-3t)c_1 - 3te^{4t}c_2 \\ 3te^{4t}c_1 + e^{4t}(1+3t)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} (c_1(1-3t) - 3c_2t)e^{4t} \\ e^{4t}(3tc_1 + 3c_2t + c_2) \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

1.72.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1-\lambda & -3 \\ 3 & 7-\lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 8\lambda + 16 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 4$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
4	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 4$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix} - (4) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -3 & -3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -3 & -3 & 0 \\ 3 & 3 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} -3 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 & -3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
4	2	1	Yes	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 4 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 97: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix} - (4) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} -3 & -3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} -\frac{2}{3} \\ 1 \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue 4. Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{4t} \\ &= \begin{bmatrix} -e^{4t} \\ e^{4t} \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= (\vec{v}_1 t + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} -1 \\ 1 \end{bmatrix} t + \begin{bmatrix} -\frac{2}{3} \\ 1 \end{bmatrix} \right) e^{4t} \\ &= \begin{bmatrix} -\frac{e^{4t}(3t+2)}{3} \\ e^{4t}(t+1) \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -e^{4t} \\ e^{4t} \end{bmatrix} + c_2 \begin{bmatrix} e^{4t}(-t - \frac{2}{3}) \\ e^{4t}(t+1) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^{4t}(-c_1 - c_2 t - \frac{2}{3}c_2) \\ e^{4t}(c_2 t + c_1 + c_2) \end{bmatrix}$$

The following is the phase plot of the system.




Figure 98: Phase plot

1.72.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) - 3y(t), y'(t) = 3x(t) + 7y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[4, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right], \left[4, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[4, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 4

$$\vec{x}_1(t) = e^{4t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 4$ is the eigenvalue, and

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained
- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 4

$$\left(\begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix} - 4 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} \frac{1}{3} \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 4

$$\vec{x}_2(t) = e^{4t} \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} \frac{1}{3} \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{4t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + c_2 e^{4t} \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} \frac{1}{3} \\ 0 \end{bmatrix} \right)$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^{4t}(-c_1 - c_2 t + \frac{1}{3}c_2) \\ e^{4t}(c_2 t + c_1) \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = e^{4t}(-c_1 - c_2 t + \frac{1}{3}c_2), y(t) = e^{4t}(c_2 t + c_1)\}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve([diff(x(t),t)=x(t)-3*y(t),diff(y(t),t)=3*x(t)+7*y(t)],singsol=all)
```

$$x(t) = e^{4t}(c_2 t + c_1)$$

$$y(t) = -\frac{e^{4t}(3c_2 t + 3c_1 + c_2)}{3}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{x'[t]==x[t]-3*y[t],y'[t]==3*x[t]+7*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> T
```

$$x(t) \rightarrow -e^{4t}(c_1(3t - 1) + 3c_2t)$$

$$y(t) \rightarrow e^{4t}(3(c_1 + c_2)t + c_2)$$

1.73 problem 73

1.73.1 Solution using Matrix exponential method	586
1.73.2 Solution using explicit Eigenvalue and Eigenvector method . . .	587
1.73.3 Maple step by step solution	592

Internal problem ID [7117]

Internal file name [OUTPUT/6103_Sunday_June_05_2022_04_22_00_PM_4714900/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 73.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= x(t) - 2y(t) \\y'(t) &= 2x(t) + 5y(t)\end{aligned}$$

1.73.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{3t}(1 - 2t) & -2t e^{3t} \\ 2t e^{3t} & e^{3t}(2t + 1) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^{3t}(1-2t) & -2t e^{3t} \\ 2t e^{3t} & e^{3t}(2t+1) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^{3t}(1-2t)c_1 - 2t e^{3t}c_2 \\ 2t e^{3t}c_1 + e^{3t}(2t+1)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} (c_1(1-2t) - 2c_2t) e^{3t} \\ e^{3t}(2tc_1 + 2c_2t + c_2) \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

1.73.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1 - \lambda & -2 \\ 2 & 5 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 6\lambda + 9 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 3$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
3	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} -2 & -2 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -2 & -2 & 0 \\ 2 & 2 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} -2 & -2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -2 & -2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
3	2	1	Yes	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 3 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 99: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} -2 & -2 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue 3. Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{3t} \\ &= \begin{bmatrix} -e^{3t} \\ e^{3t} \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= (\vec{v}_1 t + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} -1 \\ 1 \end{bmatrix} t + \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right) e^{3t} \\ &= \begin{bmatrix} -\frac{e^{3t}(2t+1)}{2} \\ e^{3t}(t+1) \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -e^{3t} \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} e^{3t}(-t - \frac{1}{2}) \\ e^{3t}(t+1) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^{3t}(-c_1 - c_2 t - \frac{1}{2}c_2) \\ e^{3t}(c_2 t + c_1 + c_2) \end{bmatrix}$$

The following is the phase plot of the system.




Figure 100: Phase plot

1.73.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) - 2y(t), y'(t) = 2x(t) + 5y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[3, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right], \left[3, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[3, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 3

$$\vec{x}_1(t) = e^{3t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 3$ is the eigenvalue, and

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 3

$$\left(\begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} - 3 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} \frac{1}{2} \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 3

$$\vec{x}_2(t) = e^{3t} \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} \frac{1}{2} \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{3t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + c_2 e^{3t} \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} \frac{1}{2} \\ 0 \end{bmatrix} \right)$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^{3t}(-c_1 + \frac{1}{2}c_2 - c_2 t) \\ e^{3t}(c_2 t + c_1) \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = e^{3t}(-c_1 + \frac{1}{2}c_2 - c_2 t), y(t) = e^{3t}(c_2 t + c_1)\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve([diff(x(t),t) = x(t)-2*y(t), diff(y(t),t) = 2*x(t)+5*y(t)],singsol=all)
```

$$x(t) = e^{3t}(c_2 t + c_1)$$

$$y(t) = -\frac{e^{3t}(2c_2 t + 2c_1 + c_2)}{2}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 46

```
DSolve[{x'[t]== x[t]-2*y[t],y'[t] == 2*x[t]+5*y[t]},{x[t],y[t]},t,IncludeSingularSolutions
```

$$x(t) \rightarrow -e^{3t}(c_1(2t - 1) + 2c_2t)$$

$$y(t) \rightarrow e^{3t}(2(c_1 + c_2)t + c_2)$$

1.74 problem 74

1.74.1 Solution using Matrix exponential method	596
1.74.2 Solution using explicit Eigenvalue and Eigenvector method . . .	597
1.74.3 Maple step by step solution	602

Internal problem ID [7118]

Internal file name [OUTPUT/6104_Sunday_June_05_2022_04_22_01_PM_80773381/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 74.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"system of linear ODEs"**

Solve

$$\begin{aligned}x'(t) &= 7x(t) + y(t) \\y'(t) &= -4x(t) + 3y(t)\end{aligned}$$

1.74.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{5t}(2t + 1) & t e^{5t} \\ -4t e^{5t} & e^{5t}(1 - 2t) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^{5t}(2t+1) & t e^{5t} \\ -4t e^{5t} & e^{5t}(1-2t) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^{5t}(2t+1)c_1 + t e^{5t}c_2 \\ -4t e^{5t}c_1 + e^{5t}(1-2t)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^{5t}(2tc_1 + c_2t + c_1) \\ (c_2(1-2t) - 4tc_1) e^{5t} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

1.74.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 7 - \lambda & 1 \\ -4 & 3 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 10\lambda + 25 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 5$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
5	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 5$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} - (5) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ -4 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2 & 1 & 0 \\ -4 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 + 2R_1 \implies \left[\begin{array}{cc|c} 2 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -\frac{t}{2}\}$

Hence the solution is

$$\begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
5	2	1	Yes	$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 5 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 101: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} - (5) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ -4 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} 1 \\ -\frac{5}{2} \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue 5. Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} e^{5t} \\ &= \begin{bmatrix} -\frac{e^{5t}}{2} \\ e^{5t} \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= (\vec{v}_1 t + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} t + \begin{bmatrix} 1 \\ -\frac{5}{2} \end{bmatrix} \right) e^{5t} \\ &= \begin{bmatrix} -\frac{e^{5t}(t-2)}{2} \\ \frac{e^{5t}(2t-5)}{2} \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -\frac{e^{5t}}{2} \\ e^{5t} \end{bmatrix} + c_2 \begin{bmatrix} e^{5t}(-\frac{t}{2} + 1) \\ e^{5t}(t - \frac{5}{2}) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{((t-2)c_2 + c_1)e^{5t}}{2} \\ e^{5t}(c_1 + c_2 t - \frac{5}{2}c_2) \end{bmatrix}$$

The following is the phase plot of the system.




Figure 102: Phase plot

1.74.3 Maple step by step solution

Let's solve

$$[x'(t) = 7x(t) + y(t), y'(t) = -4x(t) + 3y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[5, \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right], \left[5, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[5, \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 5

$$\vec{x}_1(t) = e^{5t} \cdot \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 5$ is the eigenvalue, and

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained
- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 5

$$\left(\begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} - 5 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -\frac{1}{4} \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 5

$$\vec{x}_2(t) = e^{5t} \cdot \left(t \cdot \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} -\frac{1}{4} \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{5t} \cdot \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} + c_2 e^{5t} \cdot \left(t \cdot \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} -\frac{1}{4} \\ 0 \end{bmatrix} \right)$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{e^{5t}(2c_2t+2c_1+c_2)}{4} \\ e^{5t}(c_2t+c_1) \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = -\frac{e^{5t}(2c_2t+2c_1+c_2)}{4}, y(t) = e^{5t}(c_2t+c_1) \right\}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve([diff(x(t),t) = 7*x(t)+y(t), diff(y(t),t) = -4*x(t)+3*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= e^{5t}(c_2t+c_1) \\ y(t) &= -e^{5t}(2c_2t+2c_1-c_2) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 45

```
DSolve[{x'[t]== 7*x[t]+y[t],y'[t] == -4*x[t]+3*y[t]},{x[t],y[t]},t,IncludeSingularSolutions
```

$$x(t) \rightarrow e^{5t}(2c_1t + c_2t + c_1)$$

$$y(t) \rightarrow e^{5t}(c_2 - 2(2c_1 + c_2)t)$$

1.75 problem 75

1.75.1 Solution using Matrix exponential method 606

1.75.2 Solution using explicit Eigenvalue and Eigenvector method . . . 607

Internal problem ID [7119]

Internal file name [OUTPUT/6105_Sunday_June_05_2022_04_22_03_PM_33672749/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 75.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = x(t) + y(t)$$

$$y'(t) = y(t)$$

$$z'(t) = z(t)$$

1.75.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^t & e^t t & 0 \\ 0 & e^t & 0 \\ 0 & 0 & e^t \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At}\vec{c} \\
 &= \begin{bmatrix} e^t & e^t t & 0 \\ 0 & e^t & 0 \\ 0 & 0 & e^t \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \\
 &= \begin{bmatrix} e^t c_1 + e^t t c_2 \\ e^t c_2 \\ e^t c_3 \end{bmatrix} \\
 &= \begin{bmatrix} e^t (c_2 t + c_1) \\ e^t c_2 \\ e^t c_3 \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

1.75.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1 - \lambda & 1 & 0 \\ 0 & 1 - \lambda & 0 \\ 0 & 0 & 1 - \lambda \end{bmatrix} \right) = 0$$

Since the matrix A is triangular matrix, then the determinant is the product of the elements along the diagonal. Therefore the above becomes

$$(1 - \lambda)(1 - \lambda)(1 - \lambda) = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
1	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_1, v_3\}$ and the leading variables are $\{v_2\}$. Let $v_1 = t$. Let $v_3 = s$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_2 = 0\}$

Hence the solution is

$$\begin{bmatrix} t \\ 0 \\ s \end{bmatrix} = \begin{bmatrix} t \\ 0 \\ s \end{bmatrix}$$

Since there are two free Variable, we have found two eigenvectors associated with this eigenvalue. The above can be written as

$$\begin{aligned} \begin{bmatrix} t \\ 0 \\ s \end{bmatrix} &= \begin{bmatrix} t \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ s \end{bmatrix} \\ &= t \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \end{aligned}$$

By letting $t = 1$ and $s = 1$ then the above becomes

$$\begin{bmatrix} t \\ 0 \\ s \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Hence the two eigenvectors associated with this eigenvalue are

$$\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated

with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
1	3	2	Yes	$\begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 1 & 0 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 1 is real and repeated eigenvalue of multiplicity 3. There are three possible cases that can happen. This is illustrated in this diagram




Figure 103: Possible case for repeated λ of multiplicity 3

This eigenvalue has algebraic multiplicity of 3, and geometric multiplicity 2, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to find rank-2 eigenvector \vec{v}_3 . This eigenvector must therefore satisfy $(A - \lambda I)^2 \vec{v}_3 = \vec{0}$.

But

$$(A - \lambda I)^2 = \left(\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - 1 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right)^2$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Therefore \vec{v}_3 could be any eigenvector vector we want (but not the zero vector). Let

$$\vec{v}_3 = \begin{bmatrix} \eta_1 \\ \eta_2 \\ \eta_3 \end{bmatrix}$$

To determine the actual \vec{v}_3 we need now to enforce the condition that \vec{v}_3 satisfies

$$(A - \lambda I) \vec{v}_3 = \vec{u} \tag{1}$$

Where \vec{u} is linear combination of \vec{v}_1, \vec{v}_2 . Hence

$$\vec{u} = \alpha \vec{v}_1 + \beta \vec{v}_2$$

Where α, β are arbitrary constants (not both zero). Eq. (1) becomes

$$(A - \lambda I) \begin{bmatrix} \eta_1 \\ \eta_2 \\ \eta_3 \end{bmatrix} = \alpha \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + \beta \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \eta_1 \\ \eta_2 \\ \eta_3 \end{bmatrix} = \alpha \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + \beta \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} \eta_2 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} \beta \\ 0 \\ \alpha \end{bmatrix}$$

Expanding the above gives the following equations equations

$$\eta_2 = \beta$$

$$0 = \alpha$$

solving for α, β from the above gives

$$\begin{aligned}\eta_2 &= \beta \\ 0 &= \alpha\end{aligned}$$

Since α, β are not both zero, then we just need to determine η_i values, not all zero, which satisfy the above equations for α, β not both zero. By inspection we see that the following values satisfy this condition

$$[\eta_2 = -1]$$

Hence we found the missing generalized eigenvector

$$\vec{v}_3 = \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}$$

Which implies that

$$\begin{aligned}\alpha &= 0 \\ \beta &= -1\end{aligned}$$

Therefore

$$\begin{aligned}\vec{u} &= \alpha\vec{v}_1 + \beta\vec{v}_2 \\ &= 0 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + (-1) \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \\ &= \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix}\end{aligned}$$

Therefore the missing generalized eigenvector is now found. We have found three generalized eigenvectors for eigenvalue 1. Therefore the three basis solutions associated

with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} e^t \\ &= \begin{bmatrix} 0 \\ 0 \\ e^t \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{\lambda t} \\ &= \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} e^t \\ &= \begin{bmatrix} e^t \\ 0 \\ 0 \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_3(t) &= (\vec{u}t + \vec{v}_3) e^{\lambda t} \\ &= \left(\begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix} t + \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix} \right) e^t\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + c_3 \vec{x}_3(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = c_1 \begin{bmatrix} 0 \\ 0 \\ e^t \end{bmatrix} + c_2 \begin{bmatrix} e^t \\ 0 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} -e^t t \\ -e^t \\ 0 \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} e^t(-tc_3 + c_2) \\ -c_3e^t \\ c_1e^t \end{bmatrix}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve([diff(x(t),t)=x(t)+y(t),diff(y(t),t)=y(t),diff(z(t),t)=z(t)],singsol=all)
```

$$\begin{aligned} x(t) &= e^t(c_2t + c_1) \\ y(t) &= c_2e^t \\ z(t) &= c_3e^t \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 62

```
DSolve[{x'[t]== x[t]+y[t],y'[t] == y[t],z'[t]==z[t]},{x[t],y[t],z[t]},t,IncludeSingularSolut
```

$$\begin{aligned} x(t) &\rightarrow e^t(c_2t + c_1) \\ y(t) &\rightarrow c_2e^t \\ z(t) &\rightarrow c_3e^t \\ x(t) &\rightarrow e^t(c_2t + c_1) \\ y(t) &\rightarrow c_2e^t \\ z(t) &\rightarrow 0 \end{aligned}$$

1.76 problem 76

1.76.1 Solution using Matrix exponential method 616

1.76.2 Solution using explicit Eigenvalue and Eigenvector method . . . 617

Internal problem ID [7120]

Internal file name [OUTPUT/6106_Sunday_June_05_2022_04_22_04_PM_11878266/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 76.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = 2x(t) + y(t) - z(t)$$

$$y'(t) = -x(t) + 2z(t)$$

$$z'(t) = -x(t) - 2y(t) + 4z(t)$$

1.76.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 2 & 1 & -1 \\ -1 & 0 & 2 \\ -1 & -2 & 4 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{2t} & t e^{2t} & -t e^{2t} \\ -t e^{2t} & e^{2t} \left(1 - \frac{1}{2}t^2 - 2t\right) & \frac{e^{2t}t(t+4)}{2} \\ -t e^{2t} & -\frac{e^{2t}t(t+4)}{2} & e^{2t} \left(1 + \frac{1}{2}t^2 + 2t\right) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^{2t} & t e^{2t} & -t e^{2t} \\ -t e^{2t} & e^{2t} \left(1 - \frac{1}{2} t^2 - 2t\right) & \frac{e^{2t} t(t+4)}{2} \\ -t e^{2t} & -\frac{e^{2t} t(t+4)}{2} & e^{2t} \left(1 + \frac{1}{2} t^2 + 2t\right) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \\
 &= \begin{bmatrix} e^{2t} c_1 + t e^{2t} c_2 - t e^{2t} c_3 \\ -t e^{2t} c_1 + e^{2t} \left(1 - \frac{1}{2} t^2 - 2t\right) c_2 + \frac{e^{2t} t(t+4) c_3}{2} \\ -t e^{2t} c_1 - \frac{e^{2t} t(t+4) c_2}{2} + e^{2t} \left(1 + \frac{1}{2} t^2 + 2t\right) c_3 \end{bmatrix} \\
 &= \begin{bmatrix} ((-c_3 + c_2) t + c_1) e^{2t} \\ -\frac{((-c_3 + c_2) t^2 + (2c_1 + 4c_2 - 4c_3) t - 2c_2) e^{2t}}{2} \\ -\frac{((-c_3 + c_2) t^2 + (2c_1 + 4c_2 - 4c_3) t - 2c_3) e^{2t}}{2} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

1.76.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 2 & 1 & -1 \\ -1 & 0 & 2 \\ -1 & -2 & 4 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 2 & 1 & -1 \\ -1 & 0 & 2 \\ -1 & -2 & 4 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 2 - \lambda & 1 & -1 \\ -1 & -\lambda & 2 \\ -1 & -2 & 4 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^3 - 6\lambda^2 + 12\lambda - 8 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 2$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
2	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 2 & 1 & -1 \\ -1 & 0 & 2 \\ -1 & -2 & 4 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & -1 \\ -1 & -2 & 2 \\ -1 & -2 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} 0 & 1 & -1 & 0 \\ -1 & -2 & 2 & 0 \\ -1 & -2 & 2 & 0 \end{array} \right]$$

Since the current pivot $A(1,1)$ is zero, then the current pivot row is replaced with a row with a non-zero pivot. Swapping row 1 and row 2 gives

$$\left[\begin{array}{ccc|c} -1 & -2 & 2 & 0 \\ 0 & 1 & -1 & 0 \\ -1 & -2 & 2 & 0 \end{array} \right]$$

$$R_3 = R_3 - R_1 \implies \left[\begin{array}{ccc|c} -1 & -2 & 2 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{ccc} -1 & -2 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = 0, v_2 = t\}$

Hence the solution is

$$\begin{bmatrix} 0 \\ t \\ t \end{bmatrix} = \begin{bmatrix} 0 \\ t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} 0 \\ t \\ t \end{bmatrix} = t \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} 0 \\ t \\ t \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
2	3	1	Yes	$\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 2 is real and repeated eigenvalue of multiplicity 3. There are three possible cases that can happen. This is illustrated in this diagram




Figure 104: Possible case for repeated λ of multiplicity 3

This eigenvalue has algebraic multiplicity of 3, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 2. This falls into case 3 shown above. First we find generalized eigenvector \vec{v}_2 of rank 2 and then use this to find generalized eigenvector \vec{v}_3 of rank 3. \vec{v}_2 is found by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence

$$\left(\begin{bmatrix} 2 & 1 & -1 \\ -1 & 0 & 2 \\ -1 & -2 & 4 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & -1 \\ -1 & -2 & 2 \\ -1 & -2 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$$

Now \vec{v}_3 is found by solving

$$(A - \lambda I) \vec{v}_3 = \vec{v}_2$$

Where \vec{v}_2 is the (rank 2) generalized eigenvector found above. Hence

$$\left(\begin{bmatrix} 2 & 1 & -1 \\ -1 & 0 & 2 \\ -1 & -2 & 4 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & -1 \\ -1 & -2 & 2 \\ -1 & -2 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_3 gives

$$\vec{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

We have found three generalized eigenvectors for eigenvalue 2. Therefore the three basis

solutions associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} e^{2t} \\ &= \begin{bmatrix} 0 \\ e^{2t} \\ e^{2t} \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= e^{\lambda t}(\vec{v}_1 t + \vec{v}_2) \\ &= e^{2t} \left(\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} t + \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix} \right) \\ &= \begin{bmatrix} -e^{2t} \\ e^{2t}(t+1) \\ e^{2t}(t+1) \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_3(t) &= \left(\vec{v}_1 \frac{t^2}{2} + \vec{v}_2 t + \vec{v}_3 \right) e^{\lambda t} \\ &= \left(\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \frac{t^2}{2} + \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix} t + \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right) e^{2t} \\ &= \begin{bmatrix} -e^{2t}(t-1) \\ \frac{e^{2t}t(t+2)}{2} \\ \frac{e^{2t}(t^2+2t+2)}{2} \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + c_3 \vec{x}_3(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = c_1 \begin{bmatrix} 0 \\ e^{2t} \\ e^{2t} \end{bmatrix} + c_2 \begin{bmatrix} -e^{2t} \\ e^{2t}(t+1) \\ e^{2t}(t+1) \end{bmatrix} + c_3 \begin{bmatrix} e^{2t}(-t+1) \\ e^{2t}(\frac{1}{2}t^2+t) \\ e^{2t}(\frac{1}{2}t^2+t+1) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} -e^{2t}((t-1)c_3 + c_2) \\ \frac{(c_3t^2 + (2c_2 + 2c_3)t + 2c_1 + 2c_2)e^{2t}}{2} \\ \frac{((t^2 + 2t + 2)c_3 + 2c_2t + 2c_1 + 2c_2)e^{2t}}{2} \end{bmatrix}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 59

```
dsolve([diff(x(t),t)=2*x(t)+y(t)-z(t),diff(y(t),t)=-x(t)+2*z(t),diff(z(t),t)=-x(t)-2*y(t)+4*z(t)},{x(t),y(t),z(t)});
```

$$\begin{aligned} x(t) &= -e^{2t}(2c_3t + c_2 - 4c_3) \\ y(t) &= e^{2t}(c_3t^2 + c_2t + c_1) \\ z(t) &= e^{2t}(c_3t^2 + c_2t + c_1 + 2c_3) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 107

```
DSolve[{x'[t]== 2*x[t]+y[t]-z[t],y'[t]== -x[t]+2*z[t],z'[t]==-x[t]-2*y[t]+4*z[t]},{x[t],y[t],z[t]}];
```

$$\begin{aligned} x(t) &\rightarrow e^{2t}((c_2 - c_3)t + c_1) \\ y(t) &\rightarrow -\frac{1}{2}e^{2t}((c_2 - c_3)t^2 + 2(c_1 + 2c_2 - 2c_3)t - 2c_2) \\ z(t) &\rightarrow -\frac{1}{2}e^{2t}((c_2 - c_3)t^2 + 2(c_1 + 2c_2 - 2c_3)t - 2c_3) \end{aligned}$$

1.77 problem 77

1.77.1 Solving as quadrature ode	625
1.77.2 Maple step by step solution	626

Internal problem ID [7121]

Internal file name [OUTPUT/6107_Sunday_June_05_2022_04_22_06_PM_97271531/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 77.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"quadrature"**

Maple gives the following as the ode type

[_quadrature]

$$x' - 4Ak \left(\frac{x}{A}\right)^{\frac{3}{4}} + 3kx = 0$$

1.77.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{4Ak \left(\frac{x}{A}\right)^{\frac{3}{4}} - 3kx} dx = \int dt$$
$$\frac{-\frac{\ln(256A-81x)}{3} + \frac{\ln(9\sqrt{\frac{x}{A}}+16)}{3} - \frac{\ln(9\sqrt{\frac{x}{A}}-16)}{3} - \frac{2\ln\left(3\left(\frac{x}{A}\right)^{\frac{1}{4}}-4\right)}{3} + \frac{2\ln\left(4+3\left(\frac{x}{A}\right)^{\frac{1}{4}}\right)}{3}}{k} = t + c_1$$

Raising both side to exponential gives

$$e^{-\frac{\ln(256A-81x)}{3} + \frac{\ln(9\sqrt{\frac{x}{A}}+16)}{3} - \frac{\ln(9\sqrt{\frac{x}{A}}-16)}{3} - \frac{2\ln\left(3\left(\frac{x}{A}\right)^{\frac{1}{4}}-4\right)}{3} + \frac{2\ln\left(4+3\left(\frac{x}{A}\right)^{\frac{1}{4}}\right)}{3}}{k} = e^{t+c_1}$$

Which simplifies to

$$e^{-\frac{\ln(256A-81x) - \ln(9\sqrt{\frac{x}{A}}+16) + \ln(9\sqrt{\frac{x}{A}}-16) + 2\ln\left(3\left(\frac{x}{A}\right)^{\frac{1}{4}}-4\right) - 2\ln\left(4+3\left(\frac{x}{A}\right)^{\frac{1}{4}}\right)}{3k}} = c_2 e^t$$

Summary

The solution(s) found are the following

$$e^{-\frac{\ln(256A-81x)-\ln(9\sqrt{\frac{x}{A}}+16)+\ln(9\sqrt{\frac{x}{A}}-16)+2\ln\left(3\left(\frac{x}{A}\right)^{\frac{1}{4}}-4\right)-2\ln\left(4+3\left(\frac{x}{A}\right)^{\frac{1}{4}}\right)}{3k}} = c_2 e^t \quad (1)$$

Verification of solutions

$$e^{-\frac{\ln(256A-81x)-\ln(9\sqrt{\frac{x}{A}}+16)+\ln(9\sqrt{\frac{x}{A}}-16)+2\ln\left(3\left(\frac{x}{A}\right)^{\frac{1}{4}}-4\right)-2\ln\left(4+3\left(\frac{x}{A}\right)^{\frac{1}{4}}\right)}{3k}} = c_2 e^t$$

Verified OK.

1.77.2 Maple step by step solution

Let's solve

$$x' - 4Ak\left(\frac{x}{A}\right)^{\frac{3}{4}} + 3kx = 0$$

- Highest derivative means the order of the ODE is 1

$$x'$$

- Separate variables

$$\frac{x'}{4Ak\left(\frac{x}{A}\right)^{\frac{3}{4}} - 3kx} = 1$$

- Integrate both sides with respect to t

$$\int \frac{x'}{4Ak\left(\frac{x}{A}\right)^{\frac{3}{4}} - 3kx} dt = \int 1 dt + c_1$$

- Evaluate integral

$$\frac{-\frac{\ln(256A-81x)}{3} + \frac{\ln(9\sqrt{\frac{x}{A}}+16)}{3} - \frac{\ln(9\sqrt{\frac{x}{A}}-16)}{3} - \frac{2\ln\left(3\left(\frac{x}{A}\right)^{\frac{1}{4}}-4\right)}{3} + \frac{2\ln\left(4+3\left(\frac{x}{A}\right)^{\frac{1}{4}}\right)}{3}}{k} = t + c_1$$

- Solve for x

$$\left\{ x = \frac{\frac{16\left(8Ae^{c_1k}e^{tk} - (-A^3e^{c_1k}e^{tk})^{\frac{1}{4}}\right)^3 e^{3(t+c_1)k}}{A^2(e^{c_1k})^3(e^{tk})^3} - \frac{288e^{3(t+c_1)k}\left(8Ae^{c_1k}e^{tk} - (-A^3e^{c_1k}e^{tk})^{\frac{1}{4}}\right)^2}{A(e^{c_1k})^2(e^{tk})^2} + \frac{1792e^{3(t+c_1)k}\left(8Ae^{c_1k}e^{tk} - (-A^3e^{c_1k}e^{tk})^{\frac{1}{4}}\right)}{e^{c_1k}e^{tk}}}{81e^{3(t+c_1)k}} \right.$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 85

```
dsolve(diff(x(t),t)=4*A*k*(x(t)/A)^(3/4)-3*k*x(t),x(t), singsol=all)
```

$$\frac{\ln\left(9\sqrt{\frac{x(t)}{A}} - 16\right) - \ln\left(9\sqrt{\frac{x(t)}{A}} + 16\right) + 2\ln\left(3\left(\frac{x(t)}{A}\right)^{\frac{1}{4}} - 4\right) - 2\ln\left(3\left(\frac{x(t)}{A}\right)^{\frac{1}{4}} + 4\right) + \ln(256A - 81x(t))}{3k} = 0$$

✓ Solution by Mathematica

Time used: 0.409 (sec). Leaf size: 51

```
DSolve[x'[t]==4*A*k*(x[t]/A)^(3/4)-3*k*x[t],x[t],t,IncludeSingularSolutions -> True]
```

$$\begin{aligned}x(t) &\rightarrow \frac{1}{81}Ae^{-3kt}\left(4e^{\frac{3kt}{4}} + e^{\frac{3c_1}{4}}\right)^4 \\x(t) &\rightarrow 0 \\x(t) &\rightarrow \frac{256A}{81}\end{aligned}$$

1.78 problem 78

1.78.1 Solving as dAlembert ode 628

Internal problem ID [7122]

Internal file name [OUTPUT/6108_Sunday_June_05_2022_04_22_10_PM_61546553/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 78.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**dAlembert**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _dAlembert]
```

$$\frac{y'y}{1 + \frac{\sqrt{1+y'^2}}{2}} = -x$$

1.78.1 Solving as dAlembert ode

Let $p = y'$ the ode becomes

$$\frac{py}{1 + \frac{\sqrt{p^2+1}}{2}} = -x$$

Solving for y from the above results in

$$y = -\frac{x(2 + \sqrt{p^2 + 1})}{2p} \quad (1A)$$

This has the form

$$y = xf(p) + g(p) \quad (*)$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved. Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g')\frac{dp}{dx} \\ p - f &= (xf' + g')\frac{dp}{dx} \end{aligned} \quad (2)$$

Comparing the form $y = xf + g$ to (1A) shows that

$$f = \frac{-2 - \sqrt{p^2 + 1}}{2p}$$

$$g = 0$$

Hence (2) becomes

$$p - \frac{-2 - \sqrt{p^2 + 1}}{2p} = x \left(-\frac{1}{2\sqrt{p^2 + 1}} - \frac{-2 - \sqrt{p^2 + 1}}{2p^2} \right) p'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$p - \frac{-2 - \sqrt{p^2 + 1}}{2p} = 0$$

Solving for p from the above gives

$$p = i$$

$$p = -i$$

Substituting these in (1A) gives

$$y = -ix$$

$$y = ix$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = \frac{p(x) - \frac{-2 - \sqrt{p(x)^2 + 1}}{2p(x)}}{x \left(-\frac{1}{2\sqrt{p(x)^2 + 1}} - \frac{-2 - \sqrt{p(x)^2 + 1}}{2p(x)^2} \right)} \quad (3)$$

This ODE is now solved for $p(x)$. In canonical form the ODE is

$$p' = F(x, p)$$

$$= f(x)g(p)$$

$$= \frac{\sqrt{p^2 + 1} p (2p^2 + \sqrt{p^2 + 1} + 2)}{x (1 + 2\sqrt{p^2 + 1})}$$

Where $f(x) = \frac{1}{x}$ and $g(p) = \frac{\sqrt{p^2+1}p(2p^2+\sqrt{p^2+1}+2)}{1+2\sqrt{p^2+1}}$. Integrating both sides gives

$$\frac{1}{\frac{\sqrt{p^2+1}p(2p^2+\sqrt{p^2+1}+2)}{1+2\sqrt{p^2+1}}} dp = \frac{1}{x} dx$$

$$\int \frac{1}{\frac{\sqrt{p^2+1}p(2p^2+\sqrt{p^2+1}+2)}{1+2\sqrt{p^2+1}}} dp = \int \frac{1}{x} dx$$

$$\ln(p) - \frac{\ln(p^2+1)}{2} = \ln(x) + c_1$$

Raising both side to exponential gives

$$e^{\ln(p) - \frac{\ln(p^2+1)}{2}} = e^{\ln(x) + c_1}$$

Which simplifies to

$$\frac{p}{\sqrt{p^2+1}} = c_2 x$$

Substituing the above solution for p in (2A) gives

$$y = \frac{-2 - \sqrt{-\frac{c_2^2 x^2}{c_2^2 x^2 - 1} + 1}}{2c_2 \sqrt{-\frac{1}{c_2^2 x^2 - 1}}}$$

Summary

The solution(s) found are the following

$$y = -ix \tag{1}$$

$$y = ix \tag{2}$$

$$y = \frac{-2 - \sqrt{-\frac{c_2^2 x^2}{c_2^2 x^2 - 1} + 1}}{2c_2 \sqrt{-\frac{1}{c_2^2 x^2 - 1}}} \tag{3}$$

Verification of solutions

$$y = -ix$$

Verified OK.

$$y = ix$$

Verified OK.

$$y = \frac{-2 - \sqrt{-\frac{c_2^2 x^2}{c_2^2 x^2 - 1} + 1}}{2c_2 \sqrt{-\frac{1}{c_2^2 x^2 - 1}}}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
trying dAlembert  
<- dAlembert successful`
```


✓ Solution by Maple

Time used: 1.891 (sec). Leaf size: 187

```
dsolve(diff(y(x),x)*y(x)/(1+1/2*sqrt(1+diff(y(x),x)^2))=-x,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{-x^2 + c_1} \left(2 + \sqrt{\frac{c_1}{-x^2 + c_1}}\right)}{2}$$

$$y(x) = \frac{\sqrt{-x^2 + c_1} \left(2 + \sqrt{\frac{c_1}{-x^2 + c_1}}\right)}{2}$$

$$y(x) = -\frac{\sqrt{-9x^2 + 15c_1 - 6\sqrt{-3c_1x^2 + 4c_1^2}}}{3}$$

$$y(x) = \frac{\sqrt{-9x^2 + 15c_1 - 6\sqrt{-3c_1x^2 + 4c_1^2}}}{3}$$

$$y(x) = -\frac{\sqrt{-9x^2 + 15c_1 + 6\sqrt{-3c_1x^2 + 4c_1^2}}}{3}$$

$$y(x) = \frac{\sqrt{-9x^2 + 15c_1 + 6\sqrt{-3c_1x^2 + 4c_1^2}}}{3}$$

✓ Solution by Mathematica

Time used: 2.255 (sec). Leaf size: 153

```
DSolve[y'[x]*y[x]/(1+1/2*Sqrt[1+(y'[x])^2])==-x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \left(e^{c_1} - \sqrt{-9x^2 + 4e^{2c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{3} \left(\sqrt{-9x^2 + 4e^{2c_1}} + e^{c_1} \right)$$

$$y(x) \rightarrow -\sqrt{-x^2 + 4e^{2c_1}} - e^{c_1}$$

$$y(x) \rightarrow \sqrt{-x^2 + 4e^{2c_1}} - e^{c_1}$$

$$y(x) \rightarrow -\sqrt{-x^2}$$

$$y(x) \rightarrow \sqrt{-x^2}$$

1.79 problem 78

1.79.1 Solving as dAlembert ode 633

Internal problem ID [7123]

Internal file name [OUTPUT/6109_Sunday_June_05_2022_04_22_25_PM_59366813/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 78.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**dAlembert**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _dAlembert]
```

$$\frac{y'y}{1 + \frac{\sqrt{1+y'^2}}{2}} = -x$$

With initial conditions

$$[y(0) = 3]$$

1.79.1 Solving as dAlembert ode

Let $p = y'$ the ode becomes

$$\frac{py}{1 + \frac{\sqrt{p^2+1}}{2}} = -x$$

Solving for y from the above results in

$$y = -\frac{x(2 + \sqrt{p^2+1})}{2p} \tag{1A}$$

This has the form

$$y = xf(p) + g(p) \tag{*}$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved. Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g') \frac{dp}{dx} \\ p - f &= (xf' + g') \frac{dp}{dx} \end{aligned} \quad (2)$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned} f &= \frac{-2 - \sqrt{p^2 + 1}}{2p} \\ g &= 0 \end{aligned}$$

Hence (2) becomes

$$p - \frac{-2 - \sqrt{p^2 + 1}}{2p} = x \left(-\frac{1}{2\sqrt{p^2 + 1}} - \frac{-2 - \sqrt{p^2 + 1}}{2p^2} \right) p'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$p - \frac{-2 - \sqrt{p^2 + 1}}{2p} = 0$$

Solving for p from the above gives

$$\begin{aligned} p &= i \\ p &= -i \end{aligned}$$

Substituting these in (1A) gives

$$\begin{aligned} y &= -ix \\ y &= ix \end{aligned}$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = \frac{p(x) - \frac{-2 - \sqrt{p(x)^2 + 1}}{2p(x)}}{x \left(-\frac{1}{2\sqrt{p(x)^2 + 1}} - \frac{-2 - \sqrt{p(x)^2 + 1}}{2p(x)^2} \right)} \quad (3)$$

This ODE is now solved for $p(x)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= \frac{\sqrt{p^2 + 1} p (2p^2 + \sqrt{p^2 + 1} + 2)}{x (1 + 2\sqrt{p^2 + 1})} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(p) = \frac{\sqrt{p^2+1}p(2p^2+\sqrt{p^2+1}+2)}{1+2\sqrt{p^2+1}}$. Integrating both sides gives

$$\frac{1}{\frac{\sqrt{p^2+1}p(2p^2+\sqrt{p^2+1}+2)}{1+2\sqrt{p^2+1}}} dp = \frac{1}{x} dx$$

$$\int \frac{1}{\frac{\sqrt{p^2+1}p(2p^2+\sqrt{p^2+1}+2)}{1+2\sqrt{p^2+1}}} dp = \int \frac{1}{x} dx$$

$$\ln(p) - \frac{\ln(p^2+1)}{2} = \ln(x) + c_1$$

Raising both side to exponential gives

$$e^{\ln(p) - \frac{\ln(p^2+1)}{2}} = e^{\ln(x) + c_1}$$

Which simplifies to

$$\frac{p}{\sqrt{p^2+1}} = c_2 x$$

Substituing the above solution for p in (2A) gives

$$y = \frac{-2 - \sqrt{-\frac{c_2^2 x^2}{c_2^2 x^2 - 1} + 1}}{2c_2 \sqrt{-\frac{1}{c_2^2 x^2 - 1}}}$$

Initial conditions are used to solve for c_2 . Substituing $x = 0$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = -\frac{3}{2c_2}$$

$$c_2 = -\frac{1}{2}$$

Substituing c_2 found above in the general solution gives

$$y = \frac{1 + \sqrt{-\frac{1}{x^2-4}}}{\sqrt{-\frac{1}{x^2-4}}}$$

Summary

The solution(s) found are the following

$$y = -ix \quad (1)$$

$$y = ix \quad (2)$$

$$y = \frac{1 + \sqrt{-\frac{1}{x^2-4}}}{\sqrt{-\frac{1}{x^2-4}}} \quad (3)$$




Figure 105: Solution plot

Verification of solutions

$$y = -ix$$

Warning, solution could not be verified

$$y = ix$$

Warning, solution could not be verified

$$y = \frac{1 + \sqrt{-\frac{1}{x^2-4}}}{\sqrt{-\frac{1}{x^2-4}}}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
trying dAlembert  
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 5.672 (sec). Leaf size: 33

```
dsolve([diff(y(x),x)*y(x)/(1+1/2*sqrt(1+diff(y(x),x)^2))=-x,y(0) = 3],y(x), singsol=all)
```

$$y(x) = -3 + \sqrt{-x^2 + 36}$$

$$y(x) = 1 + \sqrt{-x^2 + 4}$$

✓ Solution by Mathematica

Time used: 0.55 (sec). Leaf size: 35

```
DSolve[{y'[x]*y[x]/(1+1/2*Sqrt[1+(y'[x])^2])==-x,y[0]==3},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \sqrt{4 - x^2} + 1$$

$$y(x) \rightarrow \sqrt{36 - x^2} - 3$$

1.80 problem 79

1.80.1 Solving as separable ode	638
1.80.2 Solving as linear ode	639
1.80.3 Solving as homogeneousTypeD2 ode	641
1.80.4 Solving as first order ode lie symmetry lookup ode	642
1.80.5 Solving as exact ode	646
1.80.6 Maple step by step solution	649

Internal problem ID [7124]

Internal file name [OUTPUT/6110_Sunday_June_05_2022_04_23_08_PM_54080598/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 79.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$y' - \frac{y \left(1 + \frac{a^2 x}{\sqrt{a^2(x^2+1)}} \right)}{\sqrt{a^2(x^2+1)}} = 0$$

1.80.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y \left(a^2 x + \sqrt{a^2(x^2+1)} \right)}{a^2(x^2+1)} \end{aligned}$$

Where $f(x) = \frac{a^2x + \sqrt{a^2(x^2+1)}}{a^2(x^2+1)}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{a^2x + \sqrt{a^2(x^2+1)}}{a^2(x^2+1)} dx \\ \int \frac{1}{y} dy &= \int \frac{a^2x + \sqrt{a^2(x^2+1)}}{a^2(x^2+1)} dx \\ \ln(y) &= \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2}\right)}{\sqrt{a^2}} + \frac{\ln(x^2+1)}{2} + c_1 \\ y &= e^{\frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2}\right)}{\sqrt{a^2}} + \frac{\ln(x^2+1)}{2} + c_1} \\ &= c_1 e^{\frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2}\right)}{\sqrt{a^2}} + \frac{\ln(x^2+1)}{2}}\end{aligned}$$

Which simplifies to

$$y = c_1 \left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2} \right)^{\frac{1}{\sqrt{a^2}}} \sqrt{x^2 + 1}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2} \right)^{\frac{1}{\sqrt{a^2}}} \sqrt{x^2 + 1} \quad (1)$$

Verification of solutions

$$y = c_1 \left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2} \right)^{\frac{1}{\sqrt{a^2}}} \sqrt{x^2 + 1}$$

Verified OK.

1.80.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -\frac{a^2x + \sqrt{a^2(x^2+1)}}{a^2(x^2+1)} \\ q(x) &= 0\end{aligned}$$

Hence the ode is

$$y' - \frac{y(a^2x + \sqrt{a^2(x^2+1)})}{a^2(x^2+1)} = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{a^2x + \sqrt{a^2(x^2+1)}}{a^2(x^2+1)} dx} \\ &= e^{-\frac{\ln\left(\frac{a^2x + \sqrt{a^2x^2+a^2}}{\sqrt{a^2}}\right) - \ln(x^2+1)}{2}}\end{aligned}$$

Which simplifies to

$$\mu = \frac{\left(ax \operatorname{csgn}(a) + \sqrt{a^2(x^2+1)}\right)^{-\frac{\operatorname{csgn}(a)}{a}}}{\sqrt{x^2+1}}$$

Which assuming all positive simplifies to

$$\mu = \frac{\left(xa + \sqrt{a^2(x^2+1)}\right)^{-\frac{1}{a}}}{\sqrt{x^2+1}}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{\left(xa + \sqrt{a^2(x^2+1)}\right)^{-\frac{1}{a}} y}{\sqrt{x^2+1}} \right) &= 0\end{aligned}$$

Integrating gives

$$\frac{\left(xa + \sqrt{a^2(x^2+1)}\right)^{-\frac{1}{a}} y}{\sqrt{x^2+1}} = c_1$$

Dividing both sides by the integrating factor $\mu = \frac{\left(xa + \sqrt{a^2(x^2+1)}\right)^{-\frac{1}{a}}}{\sqrt{x^2+1}}$ results in

$$y = c_1 \left(xa + \sqrt{a^2(x^2+1)}\right)^{\frac{1}{a}} \sqrt{x^2+1}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(xa + \sqrt{a^2(x^2 + 1)} \right)^{\frac{1}{a}} \sqrt{x^2 + 1} \quad (1)$$

Verification of solutions

$$y = c_1 \left(xa + \sqrt{a^2(x^2 + 1)} \right)^{\frac{1}{a}} \sqrt{x^2 + 1}$$

Verified OK. {positive}

1.80.3 Solving as homogeneous TypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - \frac{u(x)x \left(1 + \frac{a^2x}{\sqrt{a^2(x^2+1)}} \right)}{\sqrt{a^2(x^2+1)}} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u \left(x\sqrt{a^2(x^2+1)} - a^2 \right)}{a^2(x^2+1)x} \end{aligned}$$

Where $f(x) = \frac{x\sqrt{a^2(x^2+1)} - a^2}{a^2(x^2+1)x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{x\sqrt{a^2(x^2+1)} - a^2}{a^2(x^2+1)x} dx \\ \int \frac{1}{u} du &= \int \frac{x\sqrt{a^2(x^2+1)} - a^2}{a^2(x^2+1)x} dx \\ \ln(u) &= \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2}\right)}{\sqrt{a^2}} - \ln(x) + \frac{\ln(x^2+1)}{2} + c_2 \\ u &= e^{\frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2}\right)}{\sqrt{a^2}} - \ln(x) + \frac{\ln(x^2+1)}{2} + c_2} \\ &= c_2 e^{\frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2}\right)}{\sqrt{a^2}} - \ln(x) + \frac{\ln(x^2+1)}{2}} \end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_2 \left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + a^2} \right)^{\frac{1}{\sqrt{a^2}}}}{x} \sqrt{x^2 + 1}$$

Therefore the solution y is

$$\begin{aligned} y &= xu \\ &= c_2 \left(\frac{a^2 x}{\sqrt{a^2}} + \sqrt{a^2 x^2 + a^2} \right)^{\frac{1}{\sqrt{a^2}}} \sqrt{x^2 + 1} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 \left(\frac{a^2 x}{\sqrt{a^2}} + \sqrt{a^2 x^2 + a^2} \right)^{\frac{1}{\sqrt{a^2}}} \sqrt{x^2 + 1} \quad (1)$$

Verification of solutions

$$y = c_2 \left(\frac{a^2 x}{\sqrt{a^2}} + \sqrt{a^2 x^2 + a^2} \right)^{\frac{1}{\sqrt{a^2}}} \sqrt{x^2 + 1}$$

Verified OK. {positive}

1.80.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} y' &= \frac{y \left(a^2 x + \sqrt{a^2 (x^2 + 1)} \right)}{a^2 (x^2 + 1)} \\ y' &= \omega(x, y) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (A)$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 83: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\xi(x, y) = 0$$

$$\eta(x, y) = e^{\frac{\ln\left(\frac{a^2x}{\sqrt{a^2} + \sqrt{a^2x^2+a^2}}\right) + \ln(x^2+1)}{\sqrt{a^2}}} \quad (A1)$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{\frac{\ln\left(\frac{a^2x}{\sqrt{a^2} + \sqrt{a^2x^2+a^2}}\right)}{\sqrt{a^2}} + \frac{\ln(x^2+1)}{2}}} dy \end{aligned}$$

Which results in

$$S = e^{-\frac{\ln\left(\frac{a^2x}{\sqrt{a^2} + \sqrt{a^2x^2+a^2}}\right)}{\sqrt{a^2}} + \ln\left(\frac{1}{\sqrt{x^2+1}}\right)} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y\left(a^2x + \sqrt{a^2(x^2 + 1)}\right)}{a^2(x^2 + 1)}$$

Evaluating all the partial derivatives gives

$$R_x = 1$$

$$R_y = 0$$

$$S_x = -\frac{y a^{-\frac{a-1}{a}} (xa + \sqrt{x^2 + 1}) (\sqrt{x^2 + 1} + x)^{-\frac{1}{a}}}{(x^2 + 1)^{\frac{3}{2}}}$$

$$S_y = \frac{a^{-\frac{1}{a}} (\sqrt{x^2 + 1} + x)^{-\frac{1}{a}}}{\sqrt{x^2 + 1}}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{\left(-a^{\frac{a-1}{a}} \sqrt{x^2 + 1} + \sqrt{a^2(x^2 + 1)} a^{-\frac{1}{a}}\right) y (\sqrt{x^2 + 1} + x)^{-\frac{1}{a}}}{(x^2 + 1)^{\frac{3}{2}} a^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{ya^{-\frac{1}{a}}(\sqrt{x^2+1}+x)^{-\frac{1}{a}}}{\sqrt{x^2+1}} = c_1$$

Which simplifies to

$$\frac{ya^{-\frac{1}{a}}(\sqrt{x^2+1}+x)^{-\frac{1}{a}}}{\sqrt{x^2+1}} = c_1$$

Which gives

$$y = c_1\sqrt{x^2+1}a^{\frac{1}{a}}\left(\sqrt{x^2+1}+x\right)^{\frac{1}{a}}$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x^2+1}a^{\frac{1}{a}}\left(\sqrt{x^2+1}+x\right)^{\frac{1}{a}} \tag{1}$$

Verification of solutions

$$y = c_1\sqrt{x^2+1}a^{\frac{1}{a}}\left(\sqrt{x^2+1}+x\right)^{\frac{1}{a}}$$

Verified OK. {positive}

1.80.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{a^2}{y}\right) dy &= \left(\frac{a^2x + \sqrt{a^2(x^2 + 1)}}{x^2 + 1}\right) dx \\ \left(-\frac{a^2x + \sqrt{a^2(x^2 + 1)}}{x^2 + 1}\right) dx &+ \left(\frac{a^2}{y}\right) dy = 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$M(x, y) = -\frac{a^2x + \sqrt{a^2(x^2 + 1)}}{x^2 + 1}$$

$$N(x, y) = \frac{a^2}{y}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y} \left(-\frac{a^2x + \sqrt{a^2(x^2 + 1)}}{x^2 + 1} \right)$$

$$= 0$$

And

$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x} \left(\frac{a^2}{y} \right)$$

$$= 0$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -\frac{a^2x + \sqrt{a^2(x^2 + 1)}}{x^2 + 1} dx$$

$$\phi = -a \ln \left(ax \operatorname{csgn}(a) + \sqrt{a^2(x^2 + 1)} \right) \operatorname{csgn}(a) - \frac{a^2 \ln(x^2 + 1)}{2} + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{a^2}{y}$. Therefore equation (4) becomes

$$\frac{a^2}{y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{a^2}{y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{a^2}{y} \right) dy$$

$$f(y) = a^2 \ln(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -a \ln \left(ax \operatorname{csgn}(a) + \sqrt{a^2(x^2+1)} \right) \operatorname{csgn}(a) - \frac{a^2 \ln(x^2+1)}{2} + a^2 \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -a \ln \left(ax \operatorname{csgn}(a) + \sqrt{a^2(x^2+1)} \right) \operatorname{csgn}(a) - \frac{a^2 \ln(x^2+1)}{2} + a^2 \ln(y)$$

The solution becomes

$$y = e^{\frac{2a \ln \left(ax \operatorname{csgn}(a) + \sqrt{a^2(x^2+1)} \right) \operatorname{csgn}(a) + a^2 \ln(x^2+1) + 2c_1}{2a^2}}$$

Simplifying the solution $y = e^{\frac{2a \ln(ax \operatorname{csgn}(a) + \sqrt{a^2(x^2+1)}) \operatorname{csgn}(a) + a^2 \ln(x^2+1) + 2c_1}{2a^2}}$ to $y = e^{\frac{2a \ln(xa + \sqrt{a^2(x^2+1)}) + a^2 \ln(x^2+1) + 2c_1}{2a^2}}$
Summary

The solution(s) found are the following

$$y = e^{\frac{2a \ln(xa + \sqrt{a^2(x^2+1)}) + a^2 \ln(x^2+1) + 2c_1}{2a^2}} \quad (1)$$

Verification of solutions

$$y = e^{\frac{2a \ln(xa + \sqrt{a^2(x^2+1)}) + a^2 \ln(x^2+1) + 2c_1}{2a^2}}$$

Verified OK. {positive}

1.80.6 Maple step by step solution

Let's solve

$$y' - \frac{y \left(1 + \frac{a^2 x}{\sqrt{a^2(x^2+1)}} \right)}{\sqrt{a^2(x^2+1)}} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{y} = \frac{1 + \frac{a^2 x}{\sqrt{a^2(x^2+1)}}}{\sqrt{a^2(x^2+1)}}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1 + \frac{a^2 x}{\sqrt{a^2(x^2+1)}}}{\sqrt{a^2(x^2+1)}} dx + c_1$$

- Evaluate integral

$$\ln(y) = \frac{\ln\left(\frac{a^2 x}{\sqrt{a^2} + \sqrt{a^2 x^2 + a^2}}\right)}{\sqrt{a^2}} + \frac{\ln(x^2+1)}{2} + c_1$$

- Solve for y

$$y = e^{\frac{\ln(x^2+1) \sqrt{a^2} + 2c_1 \sqrt{a^2} + 2 \ln(a^2 x + \sqrt{a^2 x^2 + a^2} \sqrt{a^2}) - \ln(a^2)}{2\sqrt{a^2}}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x) = y(x)*(1+ a^2*x/sqrt(a^2*(x^2+1)))/sqrt(a^2*(x^2+1)),y(x), singsol=all)
```

$$y(x) = c_1 \left(ax \operatorname{csgn}(a) + \sqrt{a^2(x^2+1)} \right)^{\frac{1}{\sqrt{a^2}}} \sqrt{x^2+1}$$

✓ Solution by Mathematica

Time used: 0.365 (sec). Leaf size: 116

```
DSolve[y'[x]== y[x]*(1+ a^2*x/Sqrt[a^2*(x^2+1)]/Sqrt[a^2*(x^2+1)],y[x],x,IncludeSingularSol
```

$$y(x) \rightarrow c_1 \left(a \left(-\sqrt{a^2(x^2+1)} + \sqrt{a^2+ax} \right) \right)^{-\frac{a+1}{a}} \left(a \left(\sqrt{a^2(x^2+1)} - \sqrt{a^2+ax} \right) \right)^{\frac{1}{a}-1} \left(\sqrt{a^2} \sqrt{a^2(x^2+1)} - a^2(x^2+1) \right)$$

$$y(x) \rightarrow 0$$

1.81 problem 80

1.81.1 Solving as riccati ode 651

Internal problem ID [7125]

Internal file name [OUTPUT/6111_Sunday_June_05_2022_04_23_11_PM_2402927/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 80.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**"

Maple gives the following as the ode type

`[[_Riccati, _special]]`

$$y' - y^2 = x^2$$

1.81.1 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= x^2 + y^2\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = x^2 + y^2$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x^2$, $f_1(x) = 0$ and $f_2(x) = 1$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= x^2 \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$u''(x) + x^2 u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \left(\text{BesselJ} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_1 + \text{BesselY} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_2 \right) \sqrt{x}$$

The above shows that

$$u'(x) = x^{\frac{3}{2}} \left(\text{BesselJ} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_1 + \text{BesselY} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_2 \right)$$

Using the above in (1) gives the solution

$$y = - \frac{x \left(\text{BesselJ} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_1 + \text{BesselY} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_2 \right)}{\text{BesselJ} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_1 + \text{BesselY} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_2}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = - \frac{x \left(\text{BesselJ} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselY} \left(-\frac{3}{4}, \frac{x^2}{2} \right) \right)}{\text{BesselJ} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselY} \left(\frac{1}{4}, \frac{x^2}{2} \right)}$$

Summary

The solution(s) found are the following

$$y = - \frac{x \left(\text{BesselJ} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselY} \left(-\frac{3}{4}, \frac{x^2}{2} \right) \right)}{\text{BesselJ} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselY} \left(\frac{1}{4}, \frac{x^2}{2} \right)} \quad (1)$$




Figure 106: Slope field plot

Verification of solutions

$$y = -\frac{x \left(\text{BesselJ} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselY} \left(-\frac{3}{4}, \frac{x^2}{2} \right) \right)}{\text{BesselJ} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselY} \left(\frac{1}{4}, \frac{x^2}{2} \right)}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
Looking for potential symmetries  
trying Riccati  
trying Riccati Special  
<- Riccati Special successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
dsolve(diff(y(x),x)=x^2+y(x)^2,y(x), singsol=all)
```

$$y(x) = -\frac{x \left(\text{BesselJ} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_1 + \text{BesselY} \left(-\frac{3}{4}, \frac{x^2}{2} \right) \right)}{c_1 \text{BesselJ} \left(\frac{1}{4}, \frac{x^2}{2} \right) + \text{BesselY} \left(\frac{1}{4}, \frac{x^2}{2} \right)}$$

✓ Solution by Mathematica

Time used: 0.127 (sec). Leaf size: 169

```
DSolve[y'[x]==x^2+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2 \left(-2 \text{BesselJ} \left(-\frac{3}{4}, \frac{x^2}{2} \right) + c_1 \left(\text{BesselJ} \left(\frac{3}{4}, \frac{x^2}{2} \right) - \text{BesselJ} \left(-\frac{5}{4}, \frac{x^2}{2} \right) \right) \right) - c_1 \text{BesselJ} \left(-\frac{1}{4}, \frac{x^2}{2} \right)}{2x \left(\text{BesselJ} \left(\frac{1}{4}, \frac{x^2}{2} \right) + c_1 \text{BesselJ} \left(-\frac{1}{4}, \frac{x^2}{2} \right) \right)}$$
$$y(x) \rightarrow -\frac{x^2 \text{BesselJ} \left(-\frac{5}{4}, \frac{x^2}{2} \right) - x^2 \text{BesselJ} \left(\frac{3}{4}, \frac{x^2}{2} \right) + \text{BesselJ} \left(-\frac{1}{4}, \frac{x^2}{2} \right)}{2x \text{BesselJ} \left(-\frac{1}{4}, \frac{x^2}{2} \right)}$$

1.82 problem 81

1.82.1 Existence and uniqueness analysis	655
1.82.2 Solving as quadrature ode	656
1.82.3 Maple step by step solution	657

Internal problem ID [7126]

Internal file name [OUTPUT/6112_Sunday_June_05_2022_04_23_13_PM_22840556/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 81.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' - 2\sqrt{y} = 0$$

With initial conditions

$$[y(0) = 0]$$

1.82.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= 2\sqrt{y}\end{aligned}$$

The y domain of $f(x, y)$ when $x = 0$ is

$$\{0 \leq y\}$$

And the point $y_0 = 0$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(2\sqrt{y}) \\ &= \frac{1}{\sqrt{y}}\end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{0 < y\}$$

But the point $y_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. Solution exists but no guarantee that unique solution exists.

1.82.2 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{2\sqrt{y}} dy = \int dx$$
$$\sqrt{y} = x + c_1$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_1$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$\sqrt{y} = x$$


Solving for y from the above gives

$$y = x^2$$


Summary

The solution(s) found are the following

$$y = x^2 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = x^2$$

Verified OK.

1.82.3 Maple step by step solution

Let's solve

$$[y' - 2\sqrt{y} = 0, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{\sqrt{y}} = 2$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{y}} dx = \int 2 dx + c_1$$

- Evaluate integral

$$2\sqrt{y} = 2x + c_1$$

- Solve for y

$$y = x^2 + c_1 x + \frac{1}{4}c_1^2$$

- Use initial condition $y(0) = 0$
 $0 = \frac{c_1^2}{4}$
- Solve for c_1
 $c_1 = (0, 0)$
- Substitute $c_1 = (0, 0)$ into general solution and simplify
 $y = x^2$
- Solution to the IVP
 $y = x^2$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x) = 2*sqrt(y(x)),y(0) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 8

```
DSolve[{y'[x]==2*Sqrt[y[x]],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2$$

1.83 problem 82

1.83.1 Solving as second order linear constant coeff ode	659
1.83.2 Solving using Kovacic algorithm	662
1.83.3 Maple step by step solution	667

Internal problem ID [7127]

Internal file name [OUTPUT/6113_Sunday_June_05_2022_04_23_16_PM_18109688/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 82.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$z'' + 3z' + 2z = 24e^{-3t} - 24e^{-4t}$$

1.83.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Az''(t) + Bz'(t) + Cz(t) = f(t)$$

Where $A = 1, B = 3, C = 2, f(t) = 24e^{-3t} - 24e^{-4t}$. Let the solution be

$$z = z_h + z_p$$

Where z_h is the solution to the homogeneous ODE $Az''(t) + Bz'(t) + Cz(t) = 0$, and z_p is a particular solution to the non-homogeneous ODE $Az''(t) + Bz'(t) + Cz(t) = f(t)$. z_h is the solution to

$$z'' + 3z' + 2z = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Az''(t) + Bz'(t) + Cz(t) = 0$$

Where in the above $A = 1, B = 3, C = 2$. Let the solution be $z = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} + 3\lambda e^{\lambda t} + 2e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 + 3\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 3, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-3}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{3^2 - (4)(1)(2)} \\ &= -\frac{3}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{3}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{3}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -1 \\ \lambda_2 &= -2 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} z &= c_1 e^{\lambda_1 t} + c_2 e^{\lambda_2 t} \\ z &= c_1 e^{(-1)t} + c_2 e^{(-2)t} \end{aligned}$$

Or

$$z = c_1 e^{-t} + c_2 e^{-2t}$$

Therefore the homogeneous solution z_h is

$$z_h = c_1 e^{-t} + c_2 e^{-2t}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$24e^{-3t} - 24e^{-4t}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-4t}\}, \{e^{-3t}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-2t}, e^{-t}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$z_p = A_1e^{-4t} + A_2e^{-3t}$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution z_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_1e^{-4t} + 2A_2e^{-3t} = 24e^{-3t} - 24e^{-4t}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -4, A_2 = 12]$$

Substituting the above back in the above trial solution z_p , gives the particular solution

$$z_p = -4e^{-4t} + 12e^{-3t}$$

Therefore the general solution is

$$\begin{aligned} z &= z_h + z_p \\ &= (c_1e^{-t} + c_2e^{-2t}) + (-4e^{-4t} + 12e^{-3t}) \end{aligned}$$

Summary

The solution(s) found are the following

$$z = c_1e^{-t} + c_2e^{-2t} - 4e^{-4t} + 12e^{-3t} \quad (1)$$




Figure 108: Slope field plot

Verification of solutions

$$z = c_1 e^{-t} + c_2 e^{-2t} - 4 e^{-4t} + 12 e^{-3t}$$

Verified OK.

1.83.2 Solving using Kovacic algorithm

Writing the ode as

$$z'' + 3z' + 2z = 0 \tag{1}$$

$$Az'' + Bz' + Cz = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 3 \\ C &= 2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ze^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = \frac{z}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(t)$ then z is found using the inverse transformation

$$z = ze^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 87: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = e^{-\frac{t}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in z is found from

$$\begin{aligned} z_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\ &= z_1 e^{-\int \frac{1}{2} \frac{3}{1} dt} \\ &= z_1 e^{-\frac{3t}{2}} \\ &= z_1 \left(e^{-\frac{3t}{2}} \right) \end{aligned}$$

Which simplifies to

$$z_1 = e^{-2t}$$

The second solution z_2 to the original ode is found using reduction of order

$$z_2 = z_1 \int \frac{e^{\int -\frac{B}{A} dt}}{z_1^2} dt$$

Substituting gives

$$\begin{aligned} z_2 &= z_1 \int \frac{e^{\int -\frac{3}{1} dt}}{(z_1)^2} dt \\ &= z_1 \int \frac{e^{-3t}}{(z_1)^2} dt \\ &= z_1 (e^t) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}z &= c_1 z_1 + c_2 z_2 \\ &= c_1 (e^{-2t}) + c_2 (e^{-2t} (e^t))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$z = z_h + z_p$$

Where z_h is the solution to the homogeneous ODE $Az''(t) + Bz'(t) + Cz(t) = 0$, and z_p is a particular solution to the nonhomogeneous ODE $Az''(t) + Bz'(t) + Cz(t) = f(t)$. z_h is the solution to

$$z'' + 3z' + 2z = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$z_h = c_1 e^{-2t} + e^{-t} c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$24 e^{-3t} - 24 e^{-4t}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-4t}\}, \{e^{-3t}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-2t}, e^{-t}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$z_p = A_1 e^{-4t} + A_2 e^{-3t}$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution z_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_1 e^{-4t} + 2A_2 e^{-3t} = 24 e^{-3t} - 24 e^{-4t}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -4, A_2 = 12]$$

Substituting the above back in the above trial solution z_p , gives the particular solution

$$z_p = -4e^{-4t} + 12e^{-3t}$$

Therefore the general solution is

$$\begin{aligned} z &= z_h + z_p \\ &= (c_1e^{-2t} + e^{-t}c_2) + (-4e^{-4t} + 12e^{-3t}) \end{aligned}$$

Summary

The solution(s) found are the following

$$z = c_1e^{-2t} + e^{-t}c_2 - 4e^{-4t} + 12e^{-3t} \quad (1)$$




Figure 109: Slope field plot

Verification of solutions

$$z = c_1e^{-2t} + e^{-t}c_2 - 4e^{-4t} + 12e^{-3t}$$

Verified OK.

1.83.3 Maple step by step solution

Let's solve

$$z'' + 3z' + 2z = 24e^{-3t} - 24e^{-4t}$$

- Highest derivative means the order of the ODE is 2

$$z''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 3r + 2 = 0$$

- Factor the characteristic polynomial

$$(r + 2)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-2, -1)$$

- 1st solution of the homogeneous ODE

$$z_1(t) = e^{-2t}$$

- 2nd solution of the homogeneous ODE

$$z_2(t) = e^{-t}$$

- General solution of the ODE

$$z = c_1 z_1(t) + c_2 z_2(t) + z_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$z = c_1 e^{-2t} + e^{-t} c_2 + z_p(t)$$

- Find a particular solution $z_p(t)$ of the ODE

- Use variation of parameters to find z_p here $f(t)$ is the forcing function

$$\left[z_p(t) = -z_1(t) \left(\int \frac{z_2(t)f(t)}{W(z_1(t), z_2(t))} dt \right) + z_2(t) \left(\int \frac{z_1(t)f(t)}{W(z_1(t), z_2(t))} dt \right), f(t) = 24e^{-3t} - 24e^{-4t} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(z_1(t), z_2(t)) = \begin{bmatrix} e^{-2t} & e^{-t} \\ -2e^{-2t} & -e^{-t} \end{bmatrix}$$

- Compute Wronskian

$$W(z_1(t), z_2(t)) = e^{-3t}$$

- Substitute functions into equation for $z_p(t)$

$$z_p(t) = -24e^{-2t} \left(\int (e^t - 1)e^{-2t} dt \right) + 24e^{-t} \left(\int (e^t - 1)e^{-3t} dt \right)$$

- Compute integrals

$$z_p(t) = -4e^{-4t} + 12e^{-3t}$$

- Substitute particular solution into general solution to ODE

$$z = c_1e^{-2t} + e^{-t}c_2 - 4e^{-4t} + 12e^{-3t}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
<- double symmetry of the form [xi=0, eta=F(x)] successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 30

```
dsolve(diff(z(t),t$2)+3*diff(z(t),t)+2*z(t)=24*(exp(-3*t)-exp(-4*t)),z(t), singsol=all)
```

$$z(t) = (-e^{-t}c_1 - 4e^{-3t} + 12e^{-2t} + c_2)e^{-t}$$

✓ Solution by Mathematica

Time used: 0.084 (sec). Leaf size: 34

```
DSolve[z''[t]+3*z'[t]+2*z[t]==24*(Exp[-3*t]-Exp[-4*t]),z[t],t,IncludeSingularSolutions -> Tr
```

$$z(t) \rightarrow e^{-4t}(12e^t + c_1e^{2t} + c_2e^{3t} - 4)$$

1.84 problem 83

1.84.1 Solving as separable ode	669
1.84.2 Maple step by step solution	670

Internal problem ID [7128]

Internal file name [OUTPUT/6114_Sunday_June_05_2022_04_23_18_PM_8884940/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 83.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

[_quadrature]

$$y' - \sqrt{1 - y^2} = 0$$

1.84.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \sqrt{-y^2 + 1}\end{aligned}$$

Where $f(x) = 1$ and $g(y) = \sqrt{-y^2 + 1}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\sqrt{-y^2 + 1}} dy &= 1 dx \\ \int \frac{1}{\sqrt{-y^2 + 1}} dy &= \int 1 dx \\ \arcsin(y) &= x + c_1\end{aligned}$$

Which results in

$$y = \sin(x + c_1)$$

Summary

The solution(s) found are the following

$$y = \sin(x + c_1) \quad (1)$$




Figure 110: Slope field plot

Verification of solutions

$$y = \sin(x + c_1)$$

Verified OK.

1.84.2 Maple step by step solution

Let's solve

$$y' - \sqrt{1 - y^2} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{\sqrt{1 - y^2}} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{1-y^2}} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\arcsin(y) = x + c_1$$

- Solve for y

$$y = \sin(x + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(diff(y(x),x)=sqrt(1-y(x)^2),y(x), singsol=all)
```

$$y(x) = \sin(x + c_1)$$

✓ Solution by Mathematica

Time used: 0.228 (sec). Leaf size: 28

```
DSolve[y'[x]==Sqrt[1-y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(x + c_1)$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow \text{Interval}[\{-1, 1\}]$$

1.85 problem 84

1.85.1 Solving as riccati ode 672

Internal problem ID [7129]

Internal file name [OUTPUT/6115_Sunday_June_05_2022_04_23_19_PM_88734211/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 84.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**"

Maple gives the following as the ode type

[_Riccati]

$$y' - y^2 = x^2 - 1$$

1.85.1 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= x^2 + y^2 - 1\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = x^2 + y^2 - 1$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x^2 - 1$, $f_1(x) = 0$ and $f_2(x) = 1$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= x^2 - 1 \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$u''(x) + (x^2 - 1) u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \frac{c_1 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + c_2 \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)}{\sqrt{x}}$$

The above shows that

$$\begin{aligned} u'(x) &= \frac{\left(\frac{3}{2} + \frac{i}{2}\right) c_1 \text{WhittakerM}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) - 2 \text{WhittakerW}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) c_2 + \left(c_1 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + c_2 \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)\right) x^{-\frac{3}{2}}}{x^{\frac{3}{2}}} \end{aligned}$$

Using the above in (1) gives the solution

$$y = \frac{\left(\frac{3}{2} + \frac{i}{2}\right) c_1 \text{WhittakerM}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) - 2 \text{WhittakerW}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) c_2 + \left(c_1 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + c_2 \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)\right) x^{-\frac{3}{2}}}{x \left(c_1 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + c_2 \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)\right)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant

$\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{(-3 - i) c_3 \text{WhittakerM}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) + 4 \text{WhittakerW}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) + (-2ix^2 + i + 1) c_3 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + c_3 \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)}{2x \left(c_3 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)\right)}$$

Summary

The solution(s) found are the following

$$y = \frac{(-3 - i) c_3 \text{WhittakerM}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) + 4 \text{WhittakerW}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) + (-2ix^2 + i + 1) c_3 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)}{2x \left(c_3 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)\right)} \quad (1)$$




Figure 111: Slope field plot

Verification of solutions

$$y = \frac{(-3 - i) c_3 \text{WhittakerM}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) + 4 \text{WhittakerW}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) + (-2ix^2 + i + 1) c_3 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)}{2x \left(c_3 \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)\right)}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying Riccati
trying Riccati Special
trying Riccati sub-methods:
  trying Riccati to 2nd Order
  -> Calling odsolve with the ODE`, diff(diff(y(x), x), x) = (-x^2+1)*y(x), y(x)`
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  checking if the LODE has constant coefficients
  checking if the LODE is of Euler type
  trying a symmetry of the form [xi=0, eta=F(x)]
  checking if the LODE is missing y
  -> Trying a Liouvillian solution using Kovacic's algorithm
  <- No Liouvillian solutions exist
  -> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Whittaker
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
      <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
      <- Whittaker successful
    <- special function solution successful
  <- Riccati to 2nd Order successful`
***
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 107

```
dsolve(diff(y(x),x)=x^2+y(x)^2-1,y(x), singsol=all)
```

$$y(x) = \frac{(-3 - i) \text{WhittakerM}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) + 4 \text{WhittakerW}\left(1 + \frac{i}{4}, \frac{1}{4}, ix^2\right) c_1 + (-2ix^2 + i + 1) \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)}{2x \left(c_1 \text{WhittakerW}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right) + \text{WhittakerM}\left(\frac{i}{4}, \frac{1}{4}, ix^2\right)\right)}$$

✓ Solution by Mathematica

Time used: 0.236 (sec). Leaf size: 153

```
DSolve[y'[x]==x^2+y[x]^2-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{i \left(x \text{ParabolicCylinderD}\left(-\frac{1}{2} - \frac{i}{2}, (-1 + i)x\right) + (1 + i) \text{ParabolicCylinderD}\left(\frac{1}{2} - \frac{i}{2}, (-1 + i)x\right) - c_1 x\right)}{\text{ParabolicCylinderD}\left(-\frac{1}{2} - \frac{i}{2}, (-1 + i)x\right) + c_1}$$

$$y(x) \rightarrow \frac{(1 + i) \text{ParabolicCylinderD}\left(\frac{1}{2} + \frac{i}{2}, (1 + i)x\right)}{\text{ParabolicCylinderD}\left(-\frac{1}{2} + \frac{i}{2}, (1 + i)x\right)} - ix$$

1.86 problem 85

1.86.1 Existence and uniqueness analysis	677
1.86.2 Solving as first order ode lie symmetry lookup ode	678
1.86.3 Solving as bernoulli ode	683

Internal problem ID [7130]

Internal file name [OUTPUT/6116_Sunday_June_05_2022_04_23_22_PM_35750931/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 85.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**bernoulli**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

[_Bernoulli]

$$y' - 2y(x\sqrt{y} - 1) = 0$$

With initial conditions

$$[y(0) = 1]$$

1.86.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= 2y(x\sqrt{y} - 1)\end{aligned}$$

The x domain of $f(x, y)$ when $y = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The y domain of $f(x, y)$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 1$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(2y(x\sqrt{y} - 1)) \\ &= -2 + 3x\sqrt{y}\end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 1$ is inside this domain. Therefore solution exists and is unique.

1.86.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned}y' &= 2y(x\sqrt{y} - 1) \\ y' &= \omega(x, y)\end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type Bernoulli. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 90: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int(n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= y^{\frac{3}{2}}e^x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{y^{\frac{3}{2}} e^x} dy \end{aligned}$$

Which results in

$$S = -\frac{2e^{-x}}{\sqrt{y}}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = 2y(x\sqrt{y} - 1)$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{2e^{-x}}{\sqrt{y}} \\ S_y &= \frac{e^{-x}}{y^{\frac{3}{2}}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 2x e^{-x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 2R e^{-R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -2(R + 1)e^{-R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{2e^{-x}}{\sqrt{y}} = -2(1+x)e^{-x} + c_1$$



Which simplifies to

$$-\frac{2e^{-x}}{\sqrt{y}} = -2(1+x)e^{-x} + c_1$$

Which gives

$$y = \frac{4}{(-2 + c_1 e^x - 2x)^2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = 2y(x\sqrt{y} - 1)$ 	$R = x$ $S = -\frac{2e^{-x}}{\sqrt{y}}$	$\frac{dS}{dR} = 2Re^{-R}$ 

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = \frac{4}{c_1^2 - 4c_1 + 4}$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = \frac{1}{x^2 + 2x + 1}$$


Summary

The solution(s) found are the following

$$y = \frac{1}{x^2 + 2x + 1} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{1}{x^2 + 2x + 1}$$

Verified OK.

1.86.3 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= 2y(x\sqrt{y} - 1)\end{aligned}$$

This is a Bernoulli ODE.

$$y' = -2y + 2xy^{\frac{3}{2}} \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned}f_0(x) &= -2 \\ f_1(x) &= 2x \\ n &= \frac{3}{2}\end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^{\frac{3}{2}}$ gives

$$y' \frac{1}{y^{\frac{3}{2}}} = -\frac{2}{\sqrt{y}} + 2x \quad (4)$$

Let

$$\begin{aligned}w &= y^{1-n} \\ &= \frac{1}{\sqrt{y}}\end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{1}{2y^{\frac{3}{2}}}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -2w'(x) &= -2w(x) + 2x \\ w' &= w - x \end{aligned} \tag{7}$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned} p(x) &= -1 \\ q(x) &= -x \end{aligned}$$

Hence the ode is

$$w'(x) - w(x) = -x$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int (-1)dx} \\ &= e^{-x} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu w) &= (\mu)(-x) \\ \frac{d}{dx}(e^{-x}w) &= (e^{-x})(-x) \\ d(e^{-x}w) &= (-x e^{-x}) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} e^{-x}w &= \int -x e^{-x} dx \\ e^{-x}w &= (1+x)e^{-x} + c_1 \end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-x}$ results in

$$w(x) = e^x(1+x)e^{-x} + c_1e^x$$

which simplifies to

$$w(x) = 1 + x + c_1e^x$$

Replacing w in the above by $\frac{1}{\sqrt{y}}$ using equation (5) gives the final solution.

$$\frac{1}{\sqrt{y}} = 1 + x + c_1 e^x$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = c_1 + 1$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$\frac{1}{\sqrt{y}} = 1 + x$$

The above simplifies to

$$-x\sqrt{y} - \sqrt{y} + 1 = 0$$


Solving for y from the above gives

$$y = \frac{1}{(1+x)^2}$$


Summary

The solution(s) found are the following

$$y = \frac{1}{(1+x)^2} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{1}{(1+x)^2}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 9

```
dsolve([diff(y(x),x)= 2*y(x)*(x*sqrt(y(x)) - 1),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{1}{(x+1)^2}$$

✓ Solution by Mathematica

Time used: 0.684 (sec). Leaf size: 20

```
DSolve[{y'[x]==2*y[x]*(x*Sqrt[y[x]-1]),{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1$$
$$y(x) \rightarrow \sec^2\left(\frac{x^2}{2}\right)$$

1.87 problem 86

Internal problem ID [7131]

Internal file name [OUTPUT/6117_Sunday_June_05_2022_04_23_26_PM_23415603/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 86.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order, _exact, _nonlinear], [_2nd_order,
  _with_linear_symmetries], [_2nd_order, _reducible, _mu_x_y1],
  [_2nd_order, _reducible, _mu_y_y1], [_2nd_order, _reducible,
  _mu_xy]]
```

Unable to solve or complete the solution.

$$y'' - \frac{1}{y} + \frac{xy'}{y^2} = 0$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integrating f
    --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dynam
    -> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integratin
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular cases
trying symmetries linear in x and y(x)
trying differential order: 2; exact nonlinear
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(c__1*_b(_a)-_a)/_b(_a), _b(_a)` ***
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  trying separable
  trying inverse linear
  trying homogeneous types:
  trying homogeneous D
  <- homogeneous successful
<- differential order: 2; exact nonlinear successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 56

```
dsolve(diff(y(x),x$2)=1/y(x)-x/y(x)^2*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(_Z^2 - e^{\text{RootOf} \left(x^2 \left(4e^{-Z \cosh \left(\frac{\sqrt{c_1^2+4} (2c_2 + _Z + 2 \ln(x))}{2c_1} \right)^2 + c_1^2 + 4 \right)} \right) - 1 + _Z c_1} \right) x$$

✓ Solution by Mathematica

Time used: 0.199 (sec). Leaf size: 77

```
DSolve[y''[x]==1/y[x]-x/y[x]^2*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{2} \log \left(-\frac{y(x)^2}{x^2} - \frac{c_1 y(x)}{x} + 1 \right) - \frac{c_1 \arctan \left(\frac{\frac{2y(x)}{x} + c_1}{\sqrt{-4 - c_1^2}} \right)}{\sqrt{-4 - c_1^2}} = -\log(x) + c_2, y(x) \right]$$

1.88 problem 87

1.88.1 Existence and uniqueness analysis	690
1.88.2 Solving as second order linear constant coeff ode	691
1.88.3 Solving using Kovacic algorithm	693
1.88.4 Maple step by step solution	696

Internal problem ID [7132]

Internal file name [OUTPUT/6118_Sunday_June_05_2022_04_23_29_PM_130275/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 87.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y' + y = 0$$

With initial conditions

$$[y(0) = 0]$$

1.88.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = 1$$

$$F = 0$$

Hence the ode is

$$y'' + y' + y = 0$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

1.88.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$
$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}}$$

Summary

The solution(s) found are the following

$$y = c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \quad (1)$$

Verification of solutions

$$y = c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}$$

Verified OK.

1.88.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 92: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

Summary

The solution(s) found are the following

$$y = \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} \quad (1)$$

Verification of solutions

$$y = \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

Verified OK.

1.88.4 Maple step by step solution

Let's solve

$$[y'' + y' + y = 0, y(0) = 0]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of ODE
 $r^2 + r + 1 = 0$
- Use quadratic formula to solve for r
 $r = \frac{(-1) \pm (\sqrt{-3})}{2}$
- Roots of the characteristic polynomial
 $r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)$
- 1st solution of the ODE
 $y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$
- 2nd solution of the ODE
 $y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$
- General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
- Substitute in solutions
 $y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+diff(y(x),x)+y(x)=0,y(0) = 0],y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 26

```
DSolve[{y''[x]+y'[x]+y[x]==0,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x/2} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

1.89 problem 88

1.89.1 Existence and uniqueness analysis	699
1.89.2 Solving as second order linear constant coeff ode	700
1.89.3 Solving using Kovacic algorithm	702
1.89.4 Maple step by step solution	706

Internal problem ID [7133]

Internal file name [OUTPUT/6119_Sunday_June_05_2022_04_23_31_PM_60836407/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 88.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y' + y = 0$$

With initial conditions

$$[y'(0) = 0]$$

1.89.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = 1$$

$$F = 0$$

Hence the ode is

$$y'' + y' + y = 0$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

1.89.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$
$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. Taking derivative of the solution gives

$$y' = -\frac{e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)}{2} + e^{-\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} + \frac{c_2 \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right)}{2} \right)$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -\frac{c_1}{2} + \frac{\sqrt{3}c_2}{2} \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \sqrt{3}c_2$$

Substituting these values back in above solution results in

$$y = e^{-\frac{x}{2}} \sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right) c_2 + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}$$

Which simplifies to

$$y = \left(\sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \right) c_2 e^{-\frac{x}{2}}$$

Summary

The solution(s) found are the following

$$y = \left(\sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \right) c_2 e^{-\frac{x}{2}} \quad (1)$$

Verification of solutions

$$y = \left(\sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \right) c_2 e^{-\frac{x}{2}}$$

Verified OK.

1.89.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = -3$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 94: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
 y_2 &= y_1 \int \frac{e^{\int -\frac{1}{x} dx}}{(y_1)^2} dx \\
 &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\
 &= y_1 \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right)
 \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
 y &= c_1 y_1 + c_2 y_2 \\
 &= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \right)
 \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. Taking derivative of the solution gives

$$y' = -\frac{c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{c_1 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} - \frac{c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -\frac{c_1}{2} + c_2 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 2c_2$$

Substituting these values back in above solution results in

$$y = \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} + 2c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}$$

Which simplifies to

$$y = \frac{2c_2\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3}$$

Summary

The solution(s) found are the following

$$y = \frac{2c_2\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3} \quad (1)$$

Verification of solutions

$$y = \frac{2c_2\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3}$$

Verified OK.

1.89.4 Maple step by step solution

Let's solve

$$\left[y'' + y' + y = 0, y' \Big|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve([diff(y(x),x$2)+diff(y(x),x)+y(x)=0,D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{x}{2}} \left(\sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 44

```
DSolve[{y''[x]+y'[x]+y[x]==0,{y'[0]==0}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x/2} \left(\sin\left(\frac{\sqrt{3}x}{2}\right) + \sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right) \right)$$

1.90 problem 88

1.90.1 Existence and uniqueness analysis	708
1.90.2 Solving as second order linear constant coeff ode	709
1.90.3 Solving using Kovacic algorithm	712
1.90.4 Maple step by step solution	716

Internal problem ID [7134]

Internal file name [OUTPUT/6120_Sunday_June_05_2022_04_23_33_PM_34490047/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 88.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y' + y = 0$$

With initial conditions

$$[y'(0) = 0, y(0) = 1]$$

1.90.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = 1$$

$$F = 0$$

Hence the ode is

$$y'' + y' + y = 0$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

1.90.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)}{2} + e^{-\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} + \frac{c_2 \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right)}{2} \right)$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -\frac{c_1}{2} + \frac{\sqrt{3}c_2}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = \frac{\sqrt{3}}{3}$$

Substituting these values back in above solution results in

$$y = \frac{\sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$


Which simplifies to

$$y = \frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3}$$


Summary

The solution(s) found are the following

$$y = \frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3}$$

Verified OK.

1.90.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \quad (3)$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = -3$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 96: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\&= z_1 e^{-\frac{x}{2}} \\&= z_1 \left(e^{-\frac{x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3}x}{2} \right)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3}x}{2} \right)}{3} \right) \right)\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{c_1 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} - \frac{c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -\frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= 1 \\ c_2 &= \frac{1}{2} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{\sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$


Which simplifies to

$$y = \frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3}$$


Summary

The solution(s) found are the following

$$y = \frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{x}{2}}}{3}$$

Verified OK.

1.90.4 Maple step by step solution

Let's solve

$$\left[y'' + y' + y = 0, y'|_{\{x=0\}} = 0, y(0) = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)$$

- 1st solution of the ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}$$

- Check validity of solution $y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}$

- Use initial condition $y(0) = 1$

$$1 = c_1$$

- Compute derivative of the solution

$$y' = -\frac{c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{c_1 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right) c_2}{2} - \frac{c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}}{2}$$

- Use the initial condition $y'|_{\{x=0\}} = 0$

$$0 = -\frac{c_1}{2} + \frac{\sqrt{3}c_2}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = 1, c_2 = \frac{\sqrt{3}}{3} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)) e^{-\frac{x}{2}}}{3}$$

- Solution to the IVP

$$y = \frac{(\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right) + 3 \cos\left(\frac{\sqrt{3}x}{2}\right)) e^{-\frac{x}{2}}}{3}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 31

```
dsolve([diff(y(x),x$2)+diff(y(x),x)+y(x)=0,D(y)(0) = 0, y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{e^{-\frac{x}{2}} \left(\sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right) + 3 \cos \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 47

```
DSolve[{y''[x]+y'[x]+y[x]==0,{y'[0]==0,y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} e^{-x/2} \left(\sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right) + 3 \cos \left(\frac{\sqrt{3}x}{2} \right) \right)$$

1.91 problem 89

1.91.1 Solving as second order integrable as is ode	719
1.91.2 Solving as type second_order_integrable_as_is (not using ABC version)	721
1.91.3 Solving as exact nonlinear second order ode ode	724

Internal problem ID [7135]

Internal file name [OUTPUT/6121_Sunday_June_05_2022_04_23_34_PM_68750402/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 89.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_integrable_as_is**",
"**exact nonlinear second order ode**"

Maple gives the following as the ode type

```
[[_2nd_order, _exact, _nonlinear], [_2nd_order, _reducible,  
_mu_x_y1], [_2nd_order, _reducible, _mu_y_y1], [_2nd_order,  
_reducible, _mu_xy]]
```

$$y'' - yy' = 2x$$

1.91.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - yy') dx = \int 2x dx$$
$$-\frac{y^2}{2} + y' = x^2 + c_1$$

Which is now solved for y . In canonical form the ODE is

$$y' = F(x, y)$$
$$= \frac{y^2}{2} + x^2 + c_1$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = \frac{y^2}{2} + x^2 + c_1$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x^2 + c_1$, $f_1(x) = 0$ and $f_2(x) = \frac{1}{2}$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{u}{2}} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \tag{2}$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= \frac{x^2}{4} + \frac{c_1}{4} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{u''(x)}{2} + \left(\frac{x^2}{4} + \frac{c_1}{4} \right) u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \frac{c_3 \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + c_3 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{\sqrt{x}}$$

The above shows that

$$\begin{aligned} &u'(x) \\ &= \frac{\left(\frac{(-ic_2\sqrt{2}+6) \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2} - 4 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + (-1 + i(x^2 + \frac{c_2}{2})\sqrt{2}) \right)}{2x^{\frac{3}{2}}} \end{aligned}$$

Using the above in (1) gives the solution

$$y = \frac{\left(\frac{(-ic_2\sqrt{2}+6) \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2} - 4 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + (-1+i(x^2+\frac{c_2}{2})) \right)}{x \left(c_3 \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + c_3 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)}$$

Dividing both numerator and denominator by c_2 gives, after renaming the constant

$\frac{c_3}{c_2} = c_4$ the following solution

$$y = \frac{(ic_4\sqrt{2}-6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)}$$

Summary

The solution(s) found are the following

$$y = \frac{(ic_4\sqrt{2}-6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)} \quad (1)$$

Verification of solutions

$$y = \frac{(ic_4\sqrt{2}-6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)}$$

Verified OK.

1.91.2 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$y'' - yy' = 2x$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - yy') dx = \int 2x dx$$

$$-\frac{y^2}{2} + y' = x^2 + c_1$$

Which is now solved for y . In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y^2}{2} + x^2 + c_1 \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = \frac{y^2}{2} + x^2 + c_1$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x^2 + c_1$, $f_1(x) = 0$ and $f_2(x) = \frac{1}{2}$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{u}{2}} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \tag{2}$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= \frac{x^2}{4} + \frac{c_1}{4} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{u''(x)}{2} + \left(\frac{x^2}{4} + \frac{c_1}{4} \right) u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \frac{c_3 \text{WhittakerM} \left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right) + c_3 \text{WhittakerW} \left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right)}{\sqrt{x}}$$

The above shows that

$$u'(x) = \frac{\left(\frac{(-ic_2\sqrt{2}+6) \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2} - 4 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + (-1+i(x^2+\frac{c_2}{2})\sqrt{2}) \right)}{2x^{\frac{3}{2}}}$$

Using the above in (1) gives the solution

$$y = \frac{\left(\frac{(-ic_2\sqrt{2}+6) \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2} - 4 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + (-1+i(x^2+\frac{c_2}{2})\sqrt{2}) \right)}{x \left(c_3 \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + c_3 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)}$$

Dividing both numerator and denominator by c_2 gives, after renaming the constant $\frac{c_3}{c_2} = c_4$ the following solution

$$y = \frac{(ic_4\sqrt{2}-6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)}$$

Summary

The solution(s) found are the following

$$y = \frac{(ic_4\sqrt{2}-6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)} \quad (1)$$

Verification of solutions

$$y = \frac{(ic_4\sqrt{2}-6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)}$$

Verified OK.

1.91.3 Solving as exact nonlinear second order ode

An exact non-linear second order ode has the form

$$a_2(x, y, y') y'' + a_1(x, y, y') y' + a_0(x, y, y') = 0$$

Where the following conditions are satisfied

$$\begin{aligned}\frac{\partial a_2}{\partial y} &= \frac{\partial a_1}{\partial y'} \\ \frac{\partial a_2}{\partial x} &= \frac{\partial a_0}{\partial y'} \\ \frac{\partial a_1}{\partial x} &= \frac{\partial a_0}{\partial y}\end{aligned}$$

Looking at the the ode given we see that

$$\begin{aligned}a_2 &= 1 \\ a_1 &= -y \\ a_0 &= -2x\end{aligned}$$

Applying the conditions to the above shows this is a nonlinear exact second order ode. Therefore it can be reduced to first order ode given by

$$\begin{aligned}\int a_2 dy' + \int a_1 dy + \int a_0 dx &= c_1 \\ \int 1 dy' + \int -y dy + \int -2x dx &= c_1\end{aligned}$$

Which results in

$$y' - \frac{y^2}{2} - x^2 = c_1$$

Which is now solved In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= \frac{y^2}{2} + x^2 + c_1\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = \frac{y^2}{2} + x^2 + c_1$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x^2 + c_1$, $f_1(x) = 0$ and $f_2(x) = \frac{1}{2}$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{u}{2}} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \tag{2}$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= \frac{x^2}{4} + \frac{c_1}{4} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{u''(x)}{2} + \left(\frac{x^2}{4} + \frac{c_1}{4} \right) u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \frac{c_3 \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + c_3 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{\sqrt{x}}$$

The above shows that

$$\begin{aligned} &u'(x) \\ &= \frac{\left(\frac{(-ic_2\sqrt{2}+6) \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2} - 4 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + (-1 + i(x^2 + \frac{c_2}{2})\sqrt{2}) \right)}{2x^{\frac{3}{2}}} \end{aligned}$$

Using the above in (1) gives the solution

$$y = \frac{\left(\frac{(-ic_2\sqrt{2}+6) \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2} - 4 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}+1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + (-1 + i(x^2 + \frac{c_2}{2})\sqrt{2}) \right)}{x \left(c_3 \text{WhittakerM}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + c_3 \text{WhittakerW}\left(-\frac{ic_2\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right)}$$

Dividing both numerator and denominator by c_2 gives, after renaming the constant $\frac{c_3}{c_2} = c_4$ the following solution

$$y = \frac{(ic_4\sqrt{2} - 6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8} + 1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8} + 1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right)}$$

Summary

The solution(s) found are the following

$$y = \frac{(ic_4\sqrt{2} - 6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8} + 1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8} + 1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right)} \quad (1)$$

Verification of solutions

$$y = \frac{(ic_4\sqrt{2} - 6) \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8} + 1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 8 \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8} + 1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2x \left(\text{WhittakerM}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \text{WhittakerW}\left(-\frac{ic_4\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right)}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integrating f
    --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dynam
    -> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integratin
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular cases
trying symmetries linear in x and y(x)
trying differential order: 2; exact nonlinear
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = (1/2)*_b(_a)^2+_a^2-c__1, _b(_a)` ***
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  trying separable
  trying inverse linear
  trying homogeneous types:
  trying Chini
  differential order: 1; looking for linear symmetries
  trying exact
  Looking for potential symmetries
  trying Riccati
  trying Riccati Special
  trying Riccati sub-methods:
    trying Riccati to 2nd Order
    -> Calling odsolve with the ODE`, diff(diff(y(x), x), x) = (-(1/2)*x^2+(1/2)*c__1)*y(x)
      Methods for second order ODEs:
      --- Trying classification methods ---
      trying a quadrature
      checking if the LODE has constant coefficients
      checking if the LODE is of Euler type
      trying a symmetry of the form [xi=0, eta=F(x)]
      checking if the LODE is missing y
      -> Trying a Liouvillian solution using Kovacic's algorithm
      <- No Liouvillian solutions exist
      -> Trying a solution in terms of special functions:
        -> Bessel
        -> elliptic
```


✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 161

```
dsolve(diff(y(x),x$2)-diff(y(x),x)*y(x)=2*x,y(x), singsol=all)
```

$$y(x) = \frac{-\text{WhittakerM}\left(\frac{ic_1\sqrt{2}}{8} + 1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) (6 + ic_1\sqrt{2}) + 8c_2 \text{WhittakerW}\left(\frac{ic_1\sqrt{2}}{8} + 1, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + 2(1 - i(x^2))}{2x \left(c_2 \text{WhittakerW}\left(\frac{ic_1\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \text{WhittakerM}\left(\frac{ic_1\sqrt{2}}{8}, \frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right)}$$

✓ Solution by Mathematica

Time used: 42.411 (sec). Leaf size: 318

```
DSolve[y''[x]+y'[x]*y[x]==2*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[4]{2} \left(\sqrt[4]{2} x \text{ParabolicCylinderD}\left(\frac{1}{4}(-\sqrt{2}c_1 - 2), i\sqrt[4]{2}x\right) + 2i \text{ParabolicCylinderD}\left(\frac{1}{4}(2 - \sqrt{2}c_1), i\sqrt[4]{2}x\right) \right)}{\text{ParabolicCylinderD}\left(\frac{1}{4}(-\sqrt{2}c_1 - 2), i\sqrt[4]{2}x\right)}$$

$$y(x) \rightarrow \sqrt{2}x - \frac{2\sqrt[4]{2} \text{ParabolicCylinderD}\left(\frac{1}{4}(\sqrt{2}c_1 + 2), \sqrt[4]{2}x\right)}{\text{ParabolicCylinderD}\left(\frac{1}{4}(\sqrt{2}c_1 - 2), \sqrt[4]{2}x\right)}$$

$$y(x) \rightarrow \sqrt{2}x - \frac{2\sqrt[4]{2} \text{ParabolicCylinderD}\left(\frac{1}{4}(\sqrt{2}c_1 + 2), \sqrt[4]{2}x\right)}{\text{ParabolicCylinderD}\left(\frac{1}{4}(\sqrt{2}c_1 - 2), \sqrt[4]{2}x\right)}$$

1.92 problem 90

1.92.1 Solving as riccati ode 729

Internal problem ID [7136]

Internal file name [OUTPUT/6122_Sunday_June_05_2022_04_23_38_PM_44636970/index.tex]

Book: Own collection of miscellaneous problems

Section: section 1.0

Problem number: 90.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**"

Maple gives the following as the ode type

[_Riccati]

$$y' - y^2 = x^2 + x$$

1.92.1 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= x^2 + y^2 + x\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = x^2 + y^2 + x$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x^2 + x$, $f_1(x) = 0$ and $f_2(x) = 1$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= x^2 + x \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$u''(x) + (x^2 + x) u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$\begin{aligned} u(x) = 2 e^{-\frac{ix(1+x)}{2}} & \left(c_2 \left(x + \frac{1}{2} \right) \text{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) \right. \\ & \left. + \frac{\text{hypergeom} \left(\left[\frac{1}{4} - \frac{i}{16} \right], \left[\frac{1}{2} \right], \frac{i(2x+1)^2}{4} \right) c_1}{2} \right) \end{aligned}$$

The above shows that

$$\begin{aligned} u'(x) = -2 e^{-\frac{ix(1+x)}{2}} & \left(\left(-\frac{1}{12} - i \right) \left(x + \frac{1}{2} \right)^2 c_2 \text{hypergeom} \left(\left[\frac{7}{4} - \frac{i}{16} \right], \left[\frac{5}{2} \right], \frac{i(2x+1)^2}{4} \right) \right. \\ & + \left(ix^2 + ix - 1 + \frac{1}{4}i \right) c_2 \text{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) \\ & \left. + \frac{\left(x + \frac{1}{2} \right) \left(\left(-\frac{1}{4} - i \right) \text{hypergeom} \left(\left[\frac{5}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) + i \text{hypergeom} \left(\left[\frac{1}{4} - \frac{i}{16} \right], \left[\frac{1}{2} \right], \frac{i(2x+1)^2}{4} \right) \right) c_1}{2} \right) \end{aligned}$$

Using the above in (1) gives the solution

y

$$\begin{aligned} & \frac{\left(-\frac{1}{12} - i \right) \left(x + \frac{1}{2} \right)^2 c_2 \text{hypergeom} \left(\left[\frac{7}{4} - \frac{i}{16} \right], \left[\frac{5}{2} \right], \frac{i(2x+1)^2}{4} \right) + \left(ix^2 + ix - 1 + \frac{1}{4}i \right) c_2 \text{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right)}{=} \\ & c_2 \left(x + \frac{1}{2} \right) \text{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) \end{aligned}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

y

$$= \frac{(4ix^2 + 4ix + i - 4) \operatorname{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) + 4 \left(\left(-\frac{1}{12} - i \right) \left(x + \frac{1}{2} \right) \operatorname{hypergeom} \left(\left[\frac{7}{4} - \frac{i}{16} \right] \right)}{2(2x+1) \operatorname{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) + \dots}$$

Summary

The solution(s) found are the following

y

(1)

$$= \frac{(4ix^2 + 4ix + i - 4) \operatorname{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) + 4 \left(\left(-\frac{1}{12} - i \right) \left(x + \frac{1}{2} \right) \operatorname{hypergeom} \left(\left[\frac{7}{4} - \frac{i}{16} \right] \right)}{2(2x+1) \operatorname{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) + \dots}$$




Figure 116: Slope field plot

Verification of solutions

y

$$(4ix^2 + 4ix + i - 4) \operatorname{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) + 4 \left(\left(-\frac{1}{12} - i \right) \left(x + \frac{1}{2} \right) \operatorname{hypergeom} \left(\left[\frac{7}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) + 2(2x+1) \operatorname{hypergeom} \left(\left[\frac{3}{4} - \frac{i}{16} \right], \left[\frac{3}{2} \right], \frac{i(2x+1)^2}{4} \right) + \dots \right)$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying Riccati
trying Riccati Special
trying Riccati sub-methods:
  trying Riccati to 2nd Order
  -> Calling odsolve with the ODE`, diff(diff(y(x), x), x) = (-x^2-x)*y(x), y(x)`
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  checking if the LODE has constant coefficients
  checking if the LODE is of Euler type
  trying a symmetry of the form [xi=0, eta=F(x)]
  checking if the LODE is missing y
  -> Trying a Liouvillian solution using Kovacic's algorithm
  <- No Liouvillian solutions exists
  -> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Whittaker
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
      -> heuristic approach
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
        <- hyper3 successful: indirect Equivalence to 0F1 under \\\`^ @ Moebius\\` is r
        <- hypergeometric successful
      <- special function solution successful
    <- Riccati to 2nd Order successful`
***
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 155

```
dsolve(diff(y(x),x)-y(x)^2-x-x^2=0,y(x), singsol=all)
```

$y(x)$

$$2\left(ix^2 + ix - 1 + \frac{1}{4}i\right) c_1 \operatorname{hypergeom}\left(\left[\frac{3}{4} - \frac{i}{16}\right], \left[\frac{3}{2}\right], \frac{i(2x+1)^2}{4}\right) + 2\left(-\frac{1}{12} - i\right) c_1 \left(x + \frac{1}{2}\right) \operatorname{hypergeom}\left(\left[\frac{7}{4} - \frac{i}{16}\right], \left[\frac{3}{2}\right], \frac{i(2x+1)^2}{4}\right) \\ = \frac{(2x+1) c_1 \operatorname{hypergeom}\left(\left[\frac{3}{4} - \frac{i}{16}\right], \left[\frac{3}{2}\right], \frac{i(2x+1)^2}{4}\right)}{(2x+1) c_1 \operatorname{hypergeom}\left(\left[\frac{3}{4} - \frac{i}{16}\right], \left[\frac{3}{2}\right], \frac{i(2x+1)^2}{4}\right)}$$

✓ Solution by Mathematica

Time used: 0.306 (sec). Leaf size: 298

```
DSolve[y'[x]-y[x]^2-x-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{i((2x+1) \operatorname{ParabolicCylinderD}\left(-\frac{1}{2} - \frac{i}{8}, \left(-\frac{1}{2} + \frac{i}{2}\right)(2x+1)\right) - c_1(2x+1) \operatorname{ParabolicCylinderD}\left(-\frac{1}{2} + \frac{i}{8}, \left(-\frac{1}{2} + \frac{i}{2}\right)(2x+1)\right))}{2(\operatorname{ParabolicCylinderD}\left(-\frac{1}{2} - \frac{i}{8}, \left(-\frac{1}{2} + \frac{i}{2}\right)(2x+1)\right) - c_1(2x+1) \operatorname{ParabolicCylinderD}\left(-\frac{1}{2} + \frac{i}{8}, \left(-\frac{1}{2} + \frac{i}{2}\right)(2x+1)\right))} \\ y(x) \rightarrow \frac{(1+i) \operatorname{ParabolicCylinderD}\left(\frac{1}{2} + \frac{i}{8}, (1+i)x + \left(\frac{1}{2} + \frac{i}{2}\right)\right)}{\operatorname{ParabolicCylinderD}\left(-\frac{1}{2} + \frac{i}{8}, (1+i)x + \left(\frac{1}{2} + \frac{i}{2}\right)\right)} - \frac{1}{2}i(2x+1) \\ y(x) \rightarrow \frac{(1+i) \operatorname{ParabolicCylinderD}\left(\frac{1}{2} + \frac{i}{8}, (1+i)x + \left(\frac{1}{2} + \frac{i}{2}\right)\right)}{\operatorname{ParabolicCylinderD}\left(-\frac{1}{2} + \frac{i}{8}, (1+i)x + \left(\frac{1}{2} + \frac{i}{2}\right)\right)} - \frac{1}{2}i(2x+1)$$

2 section 2.0

2.1	problem 1	737
2.2	problem 2	749
2.3	problem 3	761
2.4	problem 4	773
2.5	problem 5	785
2.6	problem 6	797
2.7	problem 7	809
2.8	problem 8	821
2.9	problem 9	833
2.10	problem 10	846
2.11	problem 11	859
2.12	problem 12	862
2.13	problem 13	865
2.14	problem 14	868
2.15	problem 15	874
2.16	problem 16	880
2.17	problem 16	886
2.18	problem 17	892
2.19	problem 18	898
2.20	problem 19	904
2.21	problem 20	910
2.22	problem 21	916
2.23	problem 22	922
2.24	problem 23	928
2.25	problem 24	934
2.26	problem 25	940
2.27	problem 26	946
2.28	problem 27	952
2.29	problem 28	958
2.30	problem 29	968
2.31	problem 30	978
2.32	problem 31	989
2.33	problem 32	999
2.34	problem 33	1010
2.35	problem 34	1020
2.36	problem 35	1027
2.37	problem 36	1034

2.38	problem 37	1041
2.39	problem 38	1048
2.40	problem 39	1054
2.41	problem 40	1061
2.42	problem 41	1068
2.43	problem 42	1071
2.44	problem 43	1074
2.45	problem 44	1086
2.46	problem 45	1088
2.47	problem 46	1091
2.48	problem 47	1094
2.49	problem 48	1101
2.50	problem 49	1125
2.51	problem 50	1132
2.52	problem 51	1135
2.53	problem 52	1138
2.54	problem 50	1141

2.1 problem 1

2.1.1 Solving using Kovacic algorithm 737

Internal problem ID [7137]

Internal file name [OUTPUT/6123_Sunday_June_05_2022_04_23_42_PM_34541086/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 1.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xy' - yx = x$$

2.1.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -x \\ C &= -x \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 98: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2) e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2) e^{-x - \frac{1}{4}x^2} \\ &= (x + 2) e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2) e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2 e^{\frac{x(4+x)}{2}}}{2x + 4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\frac{x(x+2)}{2}} x \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2} dx$$

Hence

$$u_1 = - \frac{i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x}}{2} + \frac{i e^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2})}{2} - e^x + 1$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int x(x+2) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -(1+x) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(- \frac{i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x}}{2} + \frac{i e^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2})}{2} - e^x + 1 \right) (x+2) e^{-x} + \frac{(1+x) e^{-\frac{x(x+2)}{2}} e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2}$$

Which simplifies to

$$y_p(x) = -1 - \frac{\sqrt{2} \sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1(x+2)e^{-x} - \frac{c_2e^{-x} \left(ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \\ &\quad + \left(-1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 55

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x=0,y(x), singsol=all)
```

$$y(x) = -\pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + i\sqrt{\pi}\sqrt{2} e^{\frac{x(x+2)}{2}} c_1 - 1 + e^{-x}(x+2) c_2$$

✓ Solution by Mathematica

Time used: 4.759 (sec). Leaf size: 216

`DSolve[y''[x]-x*y'[x]-x*y[x]-x==0,y[x],x,IncludeSingularSolutions -> True]`

$$\begin{aligned}
 y(x) \rightarrow & \frac{1}{2}e^{-\frac{1}{2}(x+2)^2} \left(2\sqrt{2}e^{\frac{x^2}{2}+x+2}(x+2) \int_1^x \left(\frac{e^{K[1]}K[1]}{\sqrt{2}} \right. \right. \\
 & - \left. \frac{1}{2}e^{-\frac{1}{2}K[1]^2-K[1]-2}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{(K[1]+2)^2}}{\sqrt{2}}\right) K[1]\sqrt{(K[1]+2)^2} \right) dK[1] \\
 & - \sqrt{2\pi}\sqrt{(x+2)^2} \left(c_2e^{\frac{x^2}{2}+x+2} + x + 1 \right) \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) \\
 & \left. + 2e^{\frac{x^2}{2}+x+2} \left(e^x(x+1) + \sqrt{2}c_1(x+2) + c_2e^{\frac{1}{2}(x+2)^2} \right) \right)
 \end{aligned}$$

2.2 problem 2

2.2.1 Solving using Kovacic algorithm 749

Internal problem ID [7138]

Internal file name [OUTPUT/6124_Sunday_June_05_2022_04_23_45_PM_72233922/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xy' - yx = 2x$$

2.2.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -x \tag{3}$$

$$C = -x$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 99: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int -e^{-\frac{x(x+2)}{2}} x \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) dx$$

Hence

$$u_1 = -i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x} + i e^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2}) - 2e^x + 2$$

And Eq. (3) becomes

$$u_2 = \int \frac{2(x+2) e^{-x} x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int 2x(x+2) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -2(1+x) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x} + i e^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2}) - 2e^x + 2 \right) (x+2) e^{-x} + (1+x) e^{-\frac{x(x+2)}{2}} e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)$$

Which simplifies to

$$y_p(x) = -\sqrt{2} \sqrt{\pi} \operatorname{erfi}(\sqrt{2}) (x+2) e^{-x-2} - 2 + (2x+4) e^{-x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(i e^{-2} (x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2} \right) \\ &\quad + \left(-\sqrt{2} \sqrt{\pi} \operatorname{erfi}(\sqrt{2}) (x+2) e^{-x-2} - 2 + (2x+4) e^{-x} \right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} y &= -\frac{i c_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \\ &\quad - \sqrt{2} \sqrt{\pi} \operatorname{erfi}(\sqrt{2}) (x+2) e^{-x-2} - 2 + (2x+4) e^{-x} \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= -\frac{i c_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \\ &\quad - \sqrt{2} \sqrt{\pi} \operatorname{erfi}(\sqrt{2}) (x+2) e^{-x-2} - 2 + (2x+4) e^{-x} \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y &= -\frac{i c_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \\ &\quad - \sqrt{2} \sqrt{\pi} \operatorname{erfi}(\sqrt{2}) (x+2) e^{-x-2} - 2 + (2x+4) e^{-x} \end{aligned}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    trying a symmetry of the form [xi=0, eta=F(x)]  
    checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
    A Liouvillian solution exists  
    Reducible group (found an exponential solution)  
    Group is reducible, not completely reducible  
    <- Kovacics algorithm successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 54

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-2*x=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 - 2 + e^{-x} (x+2) c_2$$

✓ Solution by Mathematica

Time used: 1.745 (sec). Leaf size: 217

`DSolve[y''[x]-x*y'[x]-x*y[x]-2*x==0,y[x],x,IncludeSingularSolutions -> True]`

$$\begin{aligned}
 y(x) \rightarrow & \frac{1}{2}e^{-\frac{1}{2}(x+2)^2} \left(2\sqrt{2}e^{\frac{x^2}{2}+x+2}(x+2) \int_1^x \left(\sqrt{2}e^{K[1]}K[1] \right. \right. \\
 & - e^{-\frac{1}{2}K[1]^2-K[1]-2}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{(K[1]+2)^2}}{\sqrt{2}}\right) K[1]\sqrt{(K[1]+2)^2} \left. \right) dK[1] \\
 & - \sqrt{2\pi}\sqrt{(x+2)^2} \left(c_2e^{\frac{x^2}{2}+x+2} + 2x + 2 \right) \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) \\
 & \left. + 2e^{\frac{x^2}{2}+x+2} \left(2e^x(x+1) + \sqrt{2}c_1(x+2) + c_2e^{\frac{1}{2}(x+2)^2} \right) \right)
 \end{aligned}$$

2.3 problem 3

2.3.1 Solving using Kovacic algorithm 761

Internal problem ID [7139]

Internal file name [OUTPUT/6125_Sunday_June_05_2022_04_23_48_PM_28311767/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xy' - yx = 3x$$

2.3.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -x \tag{3}$$

$$C = -x$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 100: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{3e^{-x} \left(ie^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right) x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{3e^{-\frac{x(x+2)}{2}} x \left(ie^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} dx$$

Hence

$$u_1 = - \frac{3i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x}}{2} + \frac{3ie^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2})}{2} - 3e^x + 3$$

And Eq. (3) becomes

$$u_2 = \int \frac{3(x+2)e^{-x}x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int 3x(x+2)e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -3(1+x)e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(- \frac{3i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x}}{2} + \frac{3ie^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2})}{2} - 3e^x \right. \\ \left. + 3 \right) (x+2)e^{-x} \\ + \frac{3(1+x)e^{-\frac{x(x+2)}{2}} e^{-x} \left(ie^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

Which simplifies to

$$y_p(x) = -3 - \frac{3\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + 3(x+2)e^{-x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1(x+2)e^{-x} - \frac{c_2e^{-x}\left(ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}}\right)}{2} \right) \\ &\quad + \left(-3 - \frac{3\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + 3(x+2)e^{-x} \right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 3 - \frac{3\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + 3(x+2)e^{-x} \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 3 - \frac{3\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + 3(x+2)e^{-x} \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 3 - \frac{3\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + 3(x+2)e^{-x} \end{aligned}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 54

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-3*x=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 - 3 + e^{-x} (x+2) c_2$$

✓ Solution by Mathematica

Time used: 2.238 (sec). Leaf size: 220

`DSolve[y''[x]-x*y'[x]-x*y[x]-3*x==0,y[x],x,IncludeSingularSolutions -> True]`

$$\begin{aligned}
 y(x) \rightarrow & \frac{1}{2}e^{-\frac{1}{2}(x+2)^2} \left(2\sqrt{2}e^{\frac{x^2}{2}+x+2}(x+2) \int_1^x \left(\frac{3e^{K[1]}K[1]}{\sqrt{2}} \right. \right. \\
 & - \frac{3}{2}e^{-\frac{1}{2}K[1]^2-K[1]-2}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{(K[1]+2)^2}}{\sqrt{2}}\right) K[1]\sqrt{(K[1]+2)^2} \left. \right) dK[1] \\
 & - \sqrt{2}\pi\sqrt{(x+2)^2} \left(c_2e^{\frac{x^2}{2}+x+2} + 3x + 3 \right) \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) \\
 & \left. + 2e^{\frac{x^2}{2}+x+2} \left(3e^x(x+1) + \sqrt{2}c_1(x+2) + c_2e^{\frac{1}{2}(x+2)^2} \right) \right)
 \end{aligned}$$

2.4 problem 4

2.4.1 Solving using Kovacic algorithm 773

Internal problem ID [7140]

Internal file name [OUTPUT/6126_Sunday_June_05_2022_04_23_51_PM_90967048/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xy' - yx = x^2 + x$$

2.4.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -x \\ C &= -x \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 101: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) (x^2+x)}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\frac{x(x+2)}{2}} (1+x) \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) x}{2} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{e^{-\frac{\alpha(\alpha+2)}{2}} (1+\alpha) \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) \alpha}{2} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} (x^2+x)}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int (x+2) x(1+x) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -(x^2 + 2x + 2) e^{-\frac{x(x+2)}{2}}$$

Which simplifies to

$$u_1 = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (1+\alpha) \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) \alpha d\alpha \right)}{2}$$

$$u_2 = -(x^2 + 2x + 2) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (1+\alpha) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) \alpha d\alpha \right) (x+2) e^{-x}}{2} + \frac{(x^2 + 2x + 2) e^{-\frac{x(x+2)}{2}} e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

Which simplifies to

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (1+\alpha) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) \alpha d\alpha \right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + 2x + 2)}{2}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) + \left(\frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (1+\alpha) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) \alpha d\alpha \right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + 2x + 2)}{2} \right)$$

Which simplifies to

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (1+\alpha) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) \alpha d\alpha \right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + 2x + 2)}{2}$$

Summary

The solution(s) found are the following

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \quad (1)$$
$$+ \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (1+\alpha) \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) \alpha d\alpha\right) (x+2) e^{-x}}{2}$$
$$+ \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + 2x + 2)}{2}$$

Verification of solutions

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x}$$
$$+ \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (1+\alpha) \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) \alpha d\alpha\right) (x+2) e^{-x}}{2}$$
$$+ \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + 2x + 2)}{2}$$

Verified OK.

Maple trace **Kovacic algorithm successful**

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 56

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x^2-x=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 + e^{-x}(x+2) c_2 - x$$

✓ Solution by Mathematica

Time used: 2.153 (sec). Leaf size: 84

```
DSolve[y''[x]-x*y'[x]-x*y[x]-x^2-x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-x} \left(-\sqrt{2\pi} c_2 \sqrt{(x+2)^2} \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) - 2e^x x + 2\sqrt{2} c_1 (x+2) + 2c_2 e^{\frac{1}{2}(x+2)^2} \right)$$

2.5 problem 5

2.5.1 Solving using Kovacic algorithm 785

Internal problem ID [7141]

Internal file name [OUTPUT/6127_Sunday_June_05_2022_04_23_57_PM_16311551/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

`[[_2nd_order , _linear , _nonhomogeneous]]`

$$y'' - xy' - yx = x^3 - 2$$

2.5.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -x \tag{3}$$

$$C = -x$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 102: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) (x^3 - 2)}{2 e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\frac{x(x+2)}{2}} (x^3 - 2) \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^3 - 2) \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right)}{2} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} (x^3 - 2)}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int (x+2) (x^3 - 2) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -(x^3 + x^2 + 2x - 2) e^{-\frac{x(x+2)}{2}}$$

Which simplifies to

$$u_1 = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^3 - 2) \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right)}{2}$$

$$u_2 = -(x^3 + x^2 + 2x - 2) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}}(\alpha^3 - 2) \left(ie^{-2}(\alpha + 2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2} + \frac{(x^3 + x^2 + 2x - 2) e^{-\frac{x(x+2)}{2}} e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}}\right)}{2}$$

Which simplifies to

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}}(\alpha^3 - 2) \left(ie^{-2}(\alpha + 2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}}}{2} (x^3 + x^2 + 2x - 2)$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}}\right)}{2}\right) + \left(\frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}}(\alpha^3 - 2) \left(ie^{-2}(\alpha + 2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}}}{2} (x^3 + x^2 + 2x - 2)\right)$$

Which simplifies to

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}}(\alpha^3 - 2) \left(ie^{-2}(\alpha + 2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}}}{2} (x^3 + x^2 + 2x - 2)$$

Summary

The solution(s) found are the following

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \quad (1)$$
$$+ \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^3 - 2) \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2}$$
$$+ \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^3 + x^2 + 2x - 2)}{2}$$

Verification of solutions

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x}$$
$$+ \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^3 - 2) \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2}$$
$$+ \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^3 + x^2 + 2x - 2)}{2}$$

Verified OK.

Maple trace **Kovacic algorithm successful**

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 62

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x^3+2=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 + e^{-x}(x+2) c_2 - x^2 + 2x - 2$$

✓ Solution by Mathematica

Time used: 5.186 (sec). Leaf size: 91

```
DSolve[y''[x]-x*y'[x]-x*y[x]-x^3+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-x} \left(-\sqrt{2\pi} c_2 \sqrt{(x+2)^2} \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) - 2e^x(x^2 - 2x + 2) + 2\sqrt{2} c_1 (x+2) + 2c_2 e^{\frac{1}{2}(x+2)^2} \right)$$

2.6 problem 6

2.6.1 Solving using Kovacic algorithm 797

Internal problem ID [7142]

Internal file name [OUTPUT/6128_Sunday_June_05_2022_04_24_02_PM_33158656/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

`[[_2nd_order , _linear , _nonhomogeneous]]`

$$y'' - xy' - yx = x^4 + 6$$

2.6.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -x \tag{3}$$

$$C = -x$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 103: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x + 2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x + 4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
 y &= c_1 y_1 + c_2 y_2 \\
 &= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2 e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)
 \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}
 y_1 &= (x+2) e^{-x} \\
 y_2 &= - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2}
 \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)\sqrt{2}} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) (x^4+6)}{2 e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\frac{x(x+2)}{2}} \left(i e^{-2(x+2)\sqrt{2}} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) (x^4+6)}{2} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{e^{-\frac{\alpha(\alpha+2)}{2}} \left(i e^{-2(\alpha+2)\sqrt{2}} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) (\alpha^4+6)}{2} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} (x^4+6)}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int (x+2) (x^4+6) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -(x^4 + x^3 + 3x^2 + 12) e^{-\frac{x(x+2)}{2}}$$

Which simplifies to

$$u_1 = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(i e^{-2(\alpha+2)\sqrt{2}} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) (\alpha^4+6) d\alpha \right)}{2}$$

$$u_2 = -(x^4 + x^3 + 3x^2 + 12) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) (\alpha^4 + 6) d\alpha \right) (x+2) e^{-x}}{2} + \frac{(x^4 + x^3 + 3x^2 + 12) e^{-\frac{x(x+2)}{2}} e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2}$$

Which simplifies to

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) (\alpha^4 + 6) d\alpha \right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^4 + x^3 + 3x^2 + 12)}{2}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2} \right) + \left(\frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) (\alpha^4 + 6) d\alpha \right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^4 + x^3 + 3x^2 + 12)}{2} \right)$$

Which simplifies to

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1(x+2) e^{-x} + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) (\alpha^4 + 6) d\alpha \right) (x+2) e^{-x}}{2} + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^4 + x^3 + 3x^2 + 12)}{2}$$

Summary

The solution(s) found are the following

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \quad (1)$$
$$+ \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) (\alpha^4 + 6) d\alpha\right) (x+2) e^{-x}}{2}$$
$$+ \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^4 + x^3 + 3x^2 + 12)}{2}$$

Verification of solutions

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x}$$
$$+ \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) (\alpha^4 + 6) d\alpha\right) (x+2) e^{-x}}{2}$$
$$+ \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^4 + x^3 + 3x^2 + 12)}{2}$$

Verified OK.

Maple trace **Kovacic algorithm successful**

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```


✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 66

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x^4-6=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 + e^{-x} (x+2) c_2 - x^3 + 3x^2 - 6x$$

✓ Solution by Mathematica

Time used: 7.359 (sec). Leaf size: 92

```
DSolve[y''[x]-x*y'[x]-x*y[x]-x^4-6==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-x} \left(-\sqrt{2\pi} c_2 \sqrt{(x+2)^2} \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) - 2e^x x (x^2 - 3x + 6) + 2\sqrt{2} c_1 (x+2) + 2c_2 e^{\frac{1}{2}(x+2)^2} \right)$$

2.7 problem 7

2.7.1 Solving using Kovacic algorithm 809

Internal problem ID [7143]

Internal file name [OUTPUT/6129_Sunday_June_05_2022_04_24_07_PM_53103097/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 7.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - xy' - yx = x^5 - 24$$

2.7.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -x \tag{3}$$

$$C = -x$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 104: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2) e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2) e^{-x - \frac{1}{4}x^2} \\ &= (x + 2) e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2) e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2 e^{\frac{x(4+x)}{2}}}{2x + 4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) (x^5 - 24)}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int - \frac{e^{-\frac{x(x+2)}{2}} (x^5 - 24) \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2} dx$$

Hence

$$u_1 = - \left(\int_0^x - \frac{e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^5 - 24) \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right)}{2} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} (x^5 - 24)}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int (x+2) e^{-\frac{x(x+2)}{2}} (x^5 - 24) dx$$

Hence

$$u_2 = -(x^5 + x^4 + 4x^3 + 12x - 36) e^{-\frac{x(x+2)}{2}}$$

Which simplifies to

$$u_1 = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^5 - 24) \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right)}{2}$$

$$u_2 = -(x^5 + x^4 + 4x^3 + 12x - 36) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^5 - 24) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right) (x+2) e^{-x}}{2} + \frac{(x^5 + x^4 + 4x^3 + 12x - 36) e^{-\frac{x(x+2)}{2}} e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

Which simplifies to

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^5 - 24) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right) (x+2) e^{-x}}{2} + \frac{(x^5 + x^4 + 4x^3 + 12x - 36) \left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}}}{2}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) + \left(\frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^5 - 24) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right) (x+2) e^{-x}}{2} + \frac{(x^5 + x^4 + 4x^3 + 12x - 36) \left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}}}{2} \right)$$

Which simplifies to

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1(x+2) e^{-x} + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^5 - 24) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right) (x+2) e^{-x}}{2} + \frac{(x^5 + x^4 + 4x^3 + 12x - 36) \left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}}}{2}$$

Summary

The solution(s) found are the following

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \quad (1)$$
$$+ \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^5 - 24) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2}$$
$$+ \frac{(x^5 + x^4 + 4x^3 + 12x - 36) \left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}}}{2}$$

Verification of solutions

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x}$$
$$+ \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} (\alpha^5 - 24) \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2}$$
$$+ \frac{(x^5 + x^4 + 4x^3 + 12x - 36) \left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}}}{2}$$

Verified OK.

Maple trace **Kovacic algorithm successful**

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 72

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x^5+24=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 \\ + e^{-x} (x+2) c_2 - x^4 + 4x^3 - 12x^2 + 12x + 12$$

✓ Solution by Mathematica

Time used: 3.222 (sec). Leaf size: 102

```
DSolve[y''[x]-x*y'[x]-x*y[x]-x^5+24==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-x} \left(-\sqrt{2\pi} c_2 \sqrt{(x+2)^2} \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) \right. \\ \left. + e^x (-2x^4 + 8x^3 - 24x^2 + 24x + 24) + 2\sqrt{2} c_1 (x+2) + 2c_2 e^{\frac{1}{2}(x+2)^2} \right)$$

2.8 problem 8

2.8.1 Solving using Kovacic algorithm 821

Internal problem ID [7144]

Internal file name [OUTPUT/6130_Sunday_June_05_2022_04_24_13_PM_87287391/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xy' - yx = x$$

2.8.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -x \tag{3}$$

$$C = -x$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 105: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\frac{x(x+2)}{2}} x \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2} dx$$

Hence

$$u_1 = - \frac{i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x}}{2} + \frac{i e^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2})}{2} - e^x + 1$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int x(x+2) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -(1+x) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(- \frac{i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x}}{2} + \frac{i e^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2})}{2} - e^x + 1 \right) (x+2) e^{-x} + \frac{(1+x) e^{-\frac{x(x+2)}{2}} e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2}$$

Which simplifies to

$$y_p(x) = -1 - \frac{\sqrt{2} \sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1(x+2)e^{-x} - \frac{c_2e^{-x} \left(ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \\ &\quad + \left(-1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 - 1 + e^{-x} (x+2) c_2$$

✓ Solution by Mathematica

Time used: 0.689 (sec). Leaf size: 216

`DSolve[y''[x]-x*y'[x]-x*y[x]-x==0,y[x],x,IncludeSingularSolutions -> True]`

$$\begin{aligned}
 y(x) \rightarrow & \frac{1}{2}e^{-\frac{1}{2}(x+2)^2} \left(2\sqrt{2}e^{\frac{x^2}{2}+x+2}(x+2) \int_1^x \left(\frac{e^{K[1]}K[1]}{\sqrt{2}} \right. \right. \\
 & - \frac{1}{2}e^{-\frac{1}{2}K[1]^2-K[1]-2}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{(K[1]+2)^2}}{\sqrt{2}}\right) K[1]\sqrt{(K[1]+2)^2} \left. \right) dK[1] \\
 & - \sqrt{2\pi}\sqrt{(x+2)^2} \left(c_2e^{\frac{x^2}{2}+x+2} + x + 1 \right) \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) \\
 & \left. + 2e^{\frac{x^2}{2}+x+2} \left(e^x(x+1) + \sqrt{2}c_1(x+2) + c_2e^{\frac{1}{2}(x+2)^2} \right) \right)
 \end{aligned}$$

2.9 problem 9

2.9.1 Solving using Kovacic algorithm 833

Internal problem ID [7145]

Internal file name [OUTPUT/6131_Sunday_June_05_2022_04_24_16_PM_84183808/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 9.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xy' - yx = x^2$$

2.9.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -x \\ C &= -x \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 106: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) x^2}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\frac{x(x+2)}{2}} x^2 \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{e^{-\frac{\alpha(\alpha+2)}{2}} \alpha^2 \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right)}{2} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} x^2}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int x^2(x+2) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -(x^2 + x + 1) e^{-\frac{x(x+2)}{2}}$$

Which simplifies to

$$u_1 = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \alpha^2 \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right)}{2}$$

$$u_2 = -(x^2 + x + 1) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \alpha^2 \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right) (x+2) e^{-x}}{2} \\ + \frac{(x^2 + x + 1) e^{-\frac{x(x+2)}{2}} e^{-x} \left(ie^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

Which simplifies to

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \alpha^2 \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right) (x+2) e^{-x}}{2} \\ + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + x + 1)}{2}$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \\ + \left(\frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \alpha^2 \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right) (x+2) e^{-x}}{2} \right. \\ \left. + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + x + 1)}{2} \right)$$

Which simplifies to

$$y = -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \\ + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \alpha^2 \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2e^{\frac{\alpha(4+\alpha)}{2}} \right) d\alpha \right) (x+2) e^{-x}}{2} \\ + \frac{\left(2 + i\sqrt{2} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + x + 1)}{2}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = & -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \\ & + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \alpha^2 \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2} \\ & + \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + x + 1)}{2} \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y = & -\frac{ic_2 e^{-x-2} \sqrt{\pi} (x+2) \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2 e^{\frac{x(x+2)}{2}} + c_1 (x+2) e^{-x} \\ & + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \alpha^2 \left(ie^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) d\alpha\right) (x+2) e^{-x}}{2} \\ & + \frac{\left(2 + i\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)\right) \sqrt{\pi} (x+2) e^{-\frac{(x+2)^2}{2}} (x^2 + x + 1)}{2} \end{aligned}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 57

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x^2=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 + e^{-x} (x+2) c_2 - x + 1$$

✓ Solution by Mathematica

Time used: 4.289 (sec). Leaf size: 226

`DSolve[y''[x]-x*y'[x]-x*y[x]-x^2==0,y[x],x,IncludeSingularSolutions -> True]`

$$\begin{aligned}
 y(x) \rightarrow & \frac{1}{2}e^{-\frac{1}{2}(x+2)^2} \left(2\sqrt{2}e^{\frac{x^2}{2}+x+2}(x+2) \int_1^x \left(\frac{e^{K[1]}K[1]^2}{\sqrt{2}} \right. \right. \\
 & - \left. \frac{1}{2}e^{-\frac{1}{2}K[1]^2-K[1]-2}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{(K[1]+2)^2}}{\sqrt{2}}\right) K[1]^2\sqrt{(K[1]+2)^2} \right) dK[1] \\
 & - \sqrt{2\pi}\sqrt{(x+2)^2} \left(x^2 + c_2e^{\frac{x^2}{2}+x+2} + x + 1 \right) \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) \\
 & \left. + 2e^{\frac{x^2}{2}+x+2} \left(e^x(x^2+x+1) + \sqrt{2}c_1(x+2) + c_2e^{\frac{1}{2}(x+2)^2} \right) \right)
 \end{aligned}$$

2.10 problem 10

2.10.1 Solving using Kovacic algorithm 846

Internal problem ID [7146]

Internal file name [OUTPUT/6132_Sunday_June_05_2022_04_24_21_PM_29009897/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 10.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

`[[_2nd_order , _linear , _nonhomogeneous]]`

$$y'' - xy' - yx = x^3$$

2.10.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -x \\ C &= -x \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 107: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
 y &= c_1 y_1 + c_2 y_2 \\
 &= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)
 \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}
 y_1 &= (x+2) e^{-x} \\
 y_2 &= - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}
 \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) x^3}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\frac{x(x+2)}{2}} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) x^3}{2} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{e^{-\frac{\alpha(\alpha+2)}{2}} \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) \alpha^3}{2} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} x^3}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int x^3 (x+2) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -x^3 e^{-x-\frac{1}{2}x^2} - x^2 e^{-x-\frac{1}{2}x^2} - 2x e^{-x-\frac{1}{2}x^2} + \sqrt{\pi} e^{\frac{1}{2}} \sqrt{2} \operatorname{erf} \left(\frac{\sqrt{2}x}{2} + \frac{\sqrt{2}}{2} \right)$$

Which simplifies to

$$u_1 = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(i e^{-2(\alpha+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(\alpha+2)}{2} \right) + 2 e^{\frac{\alpha(4+\alpha)}{2}} \right) \alpha^3 d\alpha \right)}{2}$$

$$u_2 = \sqrt{\pi} e^{\frac{1}{2}} \sqrt{2} \operatorname{erf} \left(\frac{\sqrt{2}(1+x)}{2} \right) - e^{-\frac{x(x+2)}{2}} x (x^2 + x + 2)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) \alpha^3 d\alpha\right) (x+2) e^{-x}}{2} - \frac{\left(\sqrt{\pi} e^{\frac{1}{2}} \sqrt{2} \operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right) - e^{-\frac{x(x+2)}{2}} x(x^2+x+2)\right) e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}}\right)}{2}$$

Which simplifies to

$$y_p(x) = \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) \alpha^3 d\alpha\right) (x+2) e^{-x}}{2} + \frac{i\sqrt{2} \sqrt{\pi} (x+2) x(x^2+x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) e^{-\frac{(x+2)^2}{2}}}{2} - \sqrt{2} \sqrt{\pi} e^{\frac{(1+x)^2}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right) - ie^{-\frac{3}{2}-x}(x+2) \operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right) \pi \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + x^3 + x^2 + 2x$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}}\right)}{2}\right) + \left(\frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(ie^{-2}(\alpha+2) \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right) \alpha^3 d\alpha\right) (x+2) e^{-x}}{2} + \frac{i\sqrt{2} \sqrt{\pi} (x+2) x(x^2+x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) e^{-\frac{(x+2)^2}{2}}}{2} - \sqrt{2} \sqrt{\pi} e^{\frac{(1+x)^2}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right) - ie^{-\frac{3}{2}-x}(x+2) \operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right) \pi \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + x^3 + x^2 + 2x\right)$$

Which simplifies to

$$\begin{aligned}
 y = & -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2}\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\
 & + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(ie^{-2(\alpha+2)}\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right)\alpha^3d\alpha\right)(x+2)e^{-x}}{2} \\
 & + \frac{i\sqrt{2}\sqrt{\pi}(x+2)x(x^2+x+2)\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)e^{-\frac{(x+2)^2}{2}}}{2} \\
 & - \sqrt{2}\sqrt{\pi}e^{\frac{(1+x)^2}{2}}\operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right) \\
 & - ie^{-\frac{3}{2}-x}(x+2)\operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right)\pi\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + x^3 + x^2 + 2x
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y = & -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2}\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\
 & + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(ie^{-2(\alpha+2)}\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right)\alpha^3d\alpha\right)(x+2)e^{-x}}{2} \\
 & + \frac{i\sqrt{2}\sqrt{\pi}(x+2)x(x^2+x+2)\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)e^{-\frac{(x+2)^2}{2}}}{2} \\
 & - \sqrt{2}\sqrt{\pi}e^{\frac{(1+x)^2}{2}}\operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right) \\
 & - ie^{-\frac{3}{2}-x}(x+2)\operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right)\pi\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + x^3 + x^2 + 2x
 \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned} y = & -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2}\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ & + \frac{\left(\int_0^x e^{-\frac{\alpha(\alpha+2)}{2}} \left(ie^{-2(\alpha+2)}\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(\frac{i\sqrt{2}(\alpha+2)}{2}\right) + 2e^{\frac{\alpha(4+\alpha)}{2}}\right)\alpha^3d\alpha\right)(x+2)e^{-x}}{2} \\ & + \frac{i\sqrt{2}\sqrt{\pi}(x+2)x(x^2+x+2)\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)e^{-\frac{(x+2)^2}{2}}}{2} \\ & - \sqrt{2}\sqrt{\pi}e^{\frac{(1+x)^2}{2}}\operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right) \\ & - ie^{-\frac{3}{2}-x}(x+2)\operatorname{erf}\left(\frac{\sqrt{2}(1+x)}{2}\right)\pi\operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + x^3 + x^2 + 2x \end{aligned}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 211

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x^3=0,y(x), singsol=all)
```

$y(x)$

$$= \frac{\sqrt{2} e^{-x} (x+2) \left(\int x^3 e^{-\frac{x(x+2)}{2}} \left(i\pi e^{-2(x+2)} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + \sqrt{2} \sqrt{\pi} e^{\frac{x(x+4)}{2}} \right) dx \right) + i\sqrt{2} (x+2) x \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right)}{1}$$

✓ Solution by Mathematica

Time used: 6.619 (sec). Leaf size: 453

```
DSolve[y''[x]-x*y'[x]-x*y[x]-x^3==0,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-\frac{1}{2}(x+2)^2} \left(2\sqrt{2} e^{\frac{x^2}{2}+x+2} (x+2) \int_1^x \left(\frac{e^{K[1]} K[1]^3}{\sqrt{2}} - \frac{1}{2} e^{-\frac{1}{2}K[1]^2-K[1]-2} \sqrt{\pi} \operatorname{erfi} \left(\frac{\sqrt{(K[1]+2)^2}}{\sqrt{2}} \right) K[1]^3 \sqrt{(K[1]+2)^2} \right) dK[1] - 2\operatorname{erf} \left(\frac{x+1}{\sqrt{2}} \right) \left(\sqrt{2\pi} e^{x^2+3x+\frac{5}{2}} - \pi e^{\frac{1}{2}(x+1)^2} \sqrt{(x+2)^2} \operatorname{erfi} \left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}} \right) \right) - \sqrt{2\pi} \sqrt{(x+2)^2} x^3 \operatorname{erfi} \left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}} \right) - \sqrt{2\pi} \sqrt{(x+2)^2} x^2 \operatorname{erfi} \left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}} \right) - \sqrt{2\pi} c_2 e^{\frac{x^2}{2}+x+2} \sqrt{(x+2)^2} \operatorname{erfi} \left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}} \right) - 2\sqrt{2\pi} \sqrt{(x+2)^2} x \operatorname{erfi} \left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}} \right) + 2e^{\frac{1}{2}(x+2)^2} x^3 + 2e^{\frac{1}{2}(x+2)^2} x^2 + 2\sqrt{2} c_1 e^{\frac{x^2}{2}+x+2} x + 4\sqrt{2} c_1 e^{\frac{x^2}{2}+x+2} + 2c_2 e^{x^2+3x+4} + 4e^{\frac{1}{2}(x+2)^2} x \right)$$

2.11 problem 11

Internal problem ID [7147]

Internal file name [OUTPUT/6133_Sunday_June_05_2022_04_24_26_PM_92145129/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 11.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - axy' - bxy = cx$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        <- heuristic approach successful
    <- hypergeometric successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 86

```
dsolve(diff(y(x),x$2)-a*x*diff(y(x),x)-b*x*y(x)-c*x=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-\frac{bx}{a}} \text{KummerU}\left(-\frac{b^2}{2a^3}, \frac{1}{2}, \frac{(a^2x+2b)^2}{2a^3}\right) c_1 b + e^{-\frac{bx}{a}} \text{KummerM}\left(-\frac{b^2}{2a^3}, \frac{1}{2}, \frac{(a^2x+2b)^2}{2a^3}\right) c_2 b - c}{b}$$

✓ Solution by Mathematica

Time used: 5.384 (sec). Leaf size: 565

`DSolve[y''[x]-a*x*y'[x]-b*x*y[x]-c*x==0,y[x],x,IncludeSingularSolutions -> True]`

$y(x)$

$$\rightarrow e^{-\frac{bx}{a}} \left(\text{HermiteH} \left(\frac{b^2}{a^3}, \frac{xa^2 + 2b}{\sqrt{2a^{3/2}}} \right) \int_1^x \frac{a^4 c e^{\frac{bK}{a}}}{b^2 \left(\sqrt{2} \text{HermiteH} \left(\frac{b^2}{a^3} - 1, \frac{K[1]a^2 + 2b}{\sqrt{2a^{3/2}}} \right) \text{Hypergeometric1F1} \left(-\frac{b^2}{2a^3}, \right. \right. \right.$$

2.12 problem 12

Internal problem ID [7148]

Internal file name [OUTPUT/6134_Sunday_June_05_2022_04_24_29_PM_62312471/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 12.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - axy' - bxy = cx^2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
            <- heuristic approach successful
        <- hypergeometric successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 95

```
dsolve(diff(y(x),x$2)-a*x*diff(y(x),x)-b*x*y(x)-c*x^2=0,y(x), singsol=all)
```

$$y(x) = \frac{e^{-\frac{bx}{a}} \text{KummerM}\left(-\frac{b^2}{2a^3}, \frac{1}{2}, \frac{(a^2x+2b)^2}{2a^3}\right) c_2 b^2 + e^{-\frac{bx}{a}} \text{KummerU}\left(-\frac{b^2}{2a^3}, \frac{1}{2}, \frac{(a^2x+2b)^2}{2a^3}\right) c_1 b^2 + c(-bx + a)}{b^2}$$

✓ Solution by Mathematica

Time used: 2.978 (sec). Leaf size: 569

```
DSolve[y''[x]-a*x*y'[x]-b*x*y[x]-c*x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{bx}{a}} \left(\text{HermiteH} \left(\frac{b^2}{a^3}, \frac{xa^2 + 2b}{\sqrt{2a^{3/2}}} \right) \int_1^x \frac{a^4 c e^{\frac{bK[1]}{a}}}{b^2 \left(\sqrt{2} \text{HermiteH} \left(\frac{b^2}{a^3} - 1, \frac{K[1]a^2 + 2b}{\sqrt{2a^{3/2}}} \right) \text{Hypergeometric1F1} \left(-\frac{b^2}{2a^3}, \right. \right. \right.$$

2.13 problem 13

Internal problem ID [7149]

Internal file name [OUTPUT/6135_Sunday_June_05_2022_04_24_31_PM_59286259/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 13.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$y'' - axy' - bxy = cx^3$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
            <- heuristic approach successful
        <- hypergeometric successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 517

```
dsolve(diff(y(x),x$2)-a*x*diff(y(x),x)-b*x*y(x)-c*x^3=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{bx}{a}} \left(2 \operatorname{KummerU} \left(-\frac{b^2}{2a^3}, \frac{1}{2}, \frac{(a^2x+2b)^2}{2a^3} \right) \left(\int -\frac{(a^2x+2b)x^3 \operatorname{KummerM} \left(-\frac{b^2}{2a^3}, \frac{1}{2}, \frac{(a^2x+2b)^2}{2a^3} \right)}{\operatorname{KummerM} \left(-\frac{b^2}{2a^3}, \frac{1}{2}, \frac{(a^2x+2b)^2}{2a^3} \right)} dx \right) + (a^3 - b^2) \operatorname{KummerU} \left(\frac{2a^3 - b^2}{2a^3}, \frac{1}{2}, \frac{(a^2x+2b)^2}{2a^3} \right) \right)$$

✓ Solution by Mathematica

Time used: 3.085 (sec). Leaf size: 569

```
DSolve[y''[x]-a*x*y'[x]-b*x*y[x]-c*x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{bx}{a}} \left(\text{HermiteH} \left(\frac{b^2}{a^3}, \frac{xa^2 + 2b}{\sqrt{2a^{3/2}}} \right) \int_1^x \frac{a^4 c e^{\frac{bK[1]}{a}}}{b^2 \left(\sqrt{2} \text{HermiteH} \left(\frac{b^2}{a^3} - 1, \frac{K[1]a^2 + 2b}{\sqrt{2a^{3/2}}} \right) \text{Hypergeometric1F1} \left(-\frac{b^2}{2a^3}, \right. \right. \right.$$

2.14 problem 14

2.14.1 Solving as second order airy ode 868

Internal problem ID [7150]

Internal file name [OUTPUT/6136_Sunday_June_05_2022_04_24_34_PM_92301162/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 14.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - yx = x$$

2.14.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -1$$

$$c = -1$$

$$F = x$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi} \left(\frac{1}{4} + x \right)$$

$$y_2 = \text{AiryBi} \left(\frac{1}{4} + x \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \frac{d}{dx} (\text{AiryAi} \left(\frac{1}{4} + x \right)) & \frac{d}{dx} (\text{AiryBi} \left(\frac{1}{4} + x \right)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \text{AiryAi} \left(1, \frac{1}{4} + x \right) & \text{AiryBi} \left(1, \frac{1}{4} + x \right) \end{vmatrix}$$

Therefore

$$W = \left(\text{AiryAi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryBi} \left(1, \frac{1}{4} + x \right) \right) - \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryAi} \left(1, \frac{1}{4} + x \right) \right)$$

Which simplifies to

$$W = \text{AiryAi}\left(\frac{1}{4} + x\right) \text{AiryBi}\left(1, \frac{1}{4} + x\right) - \text{AiryBi}\left(\frac{1}{4} + x\right) \text{AiryAi}\left(1, \frac{1}{4} + x\right)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}\left(\frac{1}{4} + x\right) x}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}\left(\frac{1}{4} + x\right) x \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}\left(\frac{1}{4} + \alpha\right) \alpha \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}\left(\frac{1}{4} + x\right) x}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}\left(\frac{1}{4} + x\right) x \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}\left(\frac{1}{4} + \alpha\right) \alpha \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}\left(\frac{1}{4} + \alpha\right) \alpha d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}\left(\frac{1}{4} + \alpha\right) \alpha d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ + \left(-\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Summary

The solution(s) found are the following

$$y = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \quad (1)$$

Verification of solutions

$$y = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-x*y(x)-x=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \text{AiryAi} \left(\frac{1}{4} + x \right) c_2 + e^{\frac{x}{2}} \text{AiryBi} \left(\frac{1}{4} + x \right) c_1 - 1$$

✓ Solution by Mathematica

Time used: 13.6 (sec). Leaf size: 99

```
DSolve[y''[x]-y'[x]-x*y[x]-x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} y(x) \rightarrow e^{x/2} & \left(\text{AiryAi} \left(x + \frac{1}{4} \right) \int_1^x -e^{-\frac{K[1]}{2}} \pi \text{AiryBi} \left(K[1] + \frac{1}{4} \right) K[1] dK[1] \right. \\ & + \text{AiryBi} \left(x + \frac{1}{4} \right) \int_1^x e^{-\frac{K[2]}{2}} \pi \text{AiryAi} \left(K[2] + \frac{1}{4} \right) K[2] dK[2] \\ & \left. + c_1 \text{AiryAi} \left(x + \frac{1}{4} \right) + c_2 \text{AiryBi} \left(x + \frac{1}{4} \right) \right) \end{aligned}$$

2.15 problem 15

2.15.1 Solving as second order airy ode 874

Internal problem ID [7151]

Internal file name [OUTPUT/6137_Sunday_June_05_2022_04_24_38_PM_67771200/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 15.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_airy**"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - y' - yx = x^2$$

2.15.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -1$$

$$c = -1$$

$$F = x^2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi} \left(\frac{1}{4} + x \right)$$

$$y_2 = \text{AiryBi} \left(\frac{1}{4} + x \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \frac{d}{dx} (\text{AiryAi} \left(\frac{1}{4} + x \right)) & \frac{d}{dx} (\text{AiryBi} \left(\frac{1}{4} + x \right)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \text{AiryAi} \left(1, \frac{1}{4} + x \right) & \text{AiryBi} \left(1, \frac{1}{4} + x \right) \end{vmatrix}$$

Therefore

$$W = \left(\text{AiryAi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryBi} \left(1, \frac{1}{4} + x \right) \right) - \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryAi} \left(1, \frac{1}{4} + x \right) \right)$$

Which simplifies to

$$W = \text{AiryAi}\left(\frac{1}{4} + x\right) \text{AiryBi}\left(1, \frac{1}{4} + x\right) - \text{AiryBi}\left(\frac{1}{4} + x\right) \text{AiryAi}\left(1, \frac{1}{4} + x\right)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}\left(\frac{1}{4} + x\right) x^2}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}\left(\frac{1}{4} + x\right) x^2 \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}\left(\frac{1}{4} + \alpha\right) \alpha^2 \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}\left(\frac{1}{4} + x\right) x^2}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}\left(\frac{1}{4} + x\right) x^2 \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}\left(\frac{1}{4} + \alpha\right) \alpha^2 \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}\left(\frac{1}{4} + \alpha\right) \alpha^2 d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}\left(\frac{1}{4} + \alpha\right) \alpha^2 d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^2 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
            <- Bessel successful
        <- special function solution successful
    <- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 63

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-x*y(x)-x^2=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \pi \left(\int x^2 \text{AiryAi} \left(\frac{1}{4} + x \right) e^{-\frac{x}{2}} dx \right) \right. \\ \left. - \text{AiryAi} \left(\frac{1}{4} + x \right) \pi \left(\int x^2 \text{AiryBi} \left(\frac{1}{4} + x \right) e^{-\frac{x}{2}} dx \right) + c_1 \text{AiryBi} \left(\frac{1}{4} + x \right) \right. \\ \left. + c_2 \text{AiryAi} \left(\frac{1}{4} + x \right) \right)$$

✓ Solution by Mathematica

Time used: 9.743 (sec). Leaf size: 103

```
DSolve[y''[x]-y'[x]-x*y[x]-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left(\text{AiryAi} \left(x + \frac{1}{4} \right) \int_1^x -e^{-\frac{K[1]}{2}} \pi \text{AiryBi} \left(K[1] + \frac{1}{4} \right) K[1]^2 dK[1] \right. \\ \left. + \text{AiryBi} \left(x + \frac{1}{4} \right) \int_1^x e^{-\frac{K[2]}{2}} \pi \text{AiryAi} \left(K[2] + \frac{1}{4} \right) K[2]^2 dK[2] \right. \\ \left. + c_1 \text{AiryAi} \left(x + \frac{1}{4} \right) + c_2 \text{AiryBi} \left(x + \frac{1}{4} \right) \right)$$

2.16 problem 16

2.16.1 Solving as second order airy ode 880

Internal problem ID [7152]

Internal file name [OUTPUT/6138_Sunday_June_05_2022_04_24_42_PM_19515171/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 16.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_airy**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - yx = x^2 + 1$$

2.16.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -1$$

$$c = -1$$

$$F = x^2 + 1$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi} \left(\frac{1}{4} + x \right)$$

$$y_2 = \text{AiryBi} \left(\frac{1}{4} + x \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \frac{d}{dx} (\text{AiryAi} \left(\frac{1}{4} + x \right)) & \frac{d}{dx} (\text{AiryBi} \left(\frac{1}{4} + x \right)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \text{AiryAi} \left(1, \frac{1}{4} + x \right) & \text{AiryBi} \left(1, \frac{1}{4} + x \right) \end{vmatrix}$$

Therefore

$$W = \left(\text{AiryAi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryBi} \left(1, \frac{1}{4} + x \right) \right) - \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryAi} \left(1, \frac{1}{4} + x \right) \right)$$

Which simplifies to

$$W = \text{AiryAi}\left(\frac{1}{4} + x\right) \text{AiryBi}\left(1, \frac{1}{4} + x\right) - \text{AiryBi}\left(\frac{1}{4} + x\right) \text{AiryAi}\left(1, \frac{1}{4} + x\right)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}\left(\frac{1}{4} + x\right) (x^2 + 1)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}\left(\frac{1}{4} + x\right) (x^2 + 1) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}\left(\frac{1}{4} + \alpha\right) (\alpha^2 + 1) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}\left(\frac{1}{4} + x\right) (x^2 + 1)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}\left(\frac{1}{4} + x\right) (x^2 + 1) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}\left(\frac{1}{4} + \alpha\right) (\alpha^2 + 1) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}\left(\frac{1}{4} + \alpha\right) (\alpha^2 + 1) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}\left(\frac{1}{4} + \alpha\right) (\alpha^2 + 1) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-x*y(x)-x^2-1=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \text{AiryAi} \left(\frac{1}{4} + x \right) c_2 + e^{\frac{x}{2}} \text{AiryBi} \left(\frac{1}{4} + x \right) c_1 - x$$

✓ Solution by Mathematica

Time used: 4.468 (sec). Leaf size: 107

```
DSolve[y''[x]-y'[x]-x*y[x]-x^2-1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left(\text{AiryAi} \left(x + \frac{1}{4} \right) \int_1^x -e^{-\frac{K[1]}{2}} \pi \text{AiryBi} \left(K[1] + \frac{1}{4} \right) (K[1]^2 + 1) dK[1] \right. \\ \left. + \text{AiryBi} \left(x + \frac{1}{4} \right) \int_1^x e^{-\frac{K[2]}{2}} \pi \text{AiryAi} \left(K[2] + \frac{1}{4} \right) (K[2]^2 + 1) dK[2] \right. \\ \left. + c_1 \text{AiryAi} \left(x + \frac{1}{4} \right) + c_2 \text{AiryBi} \left(x + \frac{1}{4} \right) \right)$$

2.17 problem 16

2.17.1 Solving as second order airy ode 886

Internal problem ID [7153]

Internal file name [OUTPUT/6139_Sunday_June_05_2022_04_24_47_PM_70450727/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 16.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - yx = x^2 + 1$$

2.17.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -1$$

$$c = -1$$

$$F = x^2 + 1$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi} \left(\frac{1}{4} + x \right)$$

$$y_2 = \text{AiryBi} \left(\frac{1}{4} + x \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \frac{d}{dx} (\text{AiryAi} \left(\frac{1}{4} + x \right)) & \frac{d}{dx} (\text{AiryBi} \left(\frac{1}{4} + x \right)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \text{AiryAi} \left(1, \frac{1}{4} + x \right) & \text{AiryBi} \left(1, \frac{1}{4} + x \right) \end{vmatrix}$$

Therefore

$$W = \left(\text{AiryAi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryBi} \left(1, \frac{1}{4} + x \right) \right) - \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryAi} \left(1, \frac{1}{4} + x \right) \right)$$

Which simplifies to

$$W = \text{AiryAi} \left(\frac{1}{4} + x \right) \text{AiryBi} \left(1, \frac{1}{4} + x \right) - \text{AiryBi} \left(\frac{1}{4} + x \right) \text{AiryAi} \left(1, \frac{1}{4} + x \right)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi} \left(\frac{1}{4} + x \right) (x^2 + 1)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi} \left(\frac{1}{4} + x \right) (x^2 + 1) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi} \left(\frac{1}{4} + x \right) (x^2 + 1)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi} \left(\frac{1}{4} + x \right) (x^2 + 1) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^2 + 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-x*y(x)-x^2-1=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \text{AiryAi} \left(\frac{1}{4} + x \right) c_2 + e^{\frac{x}{2}} \text{AiryBi} \left(\frac{1}{4} + x \right) c_1 - x$$

✓ Solution by Mathematica

Time used: 1.289 (sec). Leaf size: 107

```
DSolve[y''[x]-y'[x]-x*y[x]-x^2-1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left(\text{AiryAi} \left(x + \frac{1}{4} \right) \int_1^x -e^{-\frac{K[1]}{2}} \pi \text{AiryBi} \left(K[1] + \frac{1}{4} \right) (K[1]^2 + 1) dK[1] \right. \\ \left. + \text{AiryBi} \left(x + \frac{1}{4} \right) \int_1^x e^{-\frac{K[2]}{2}} \pi \text{AiryAi} \left(K[2] + \frac{1}{4} \right) (K[2]^2 + 1) dK[2] \right. \\ \left. + c_1 \text{AiryAi} \left(x + \frac{1}{4} \right) + c_2 \text{AiryBi} \left(x + \frac{1}{4} \right) \right)$$

2.18 problem 17

2.18.1 Solving as second order airy ode 892

Internal problem ID [7154]

Internal file name [OUTPUT/6140_Sunday_June_05_2022_04_24_51_PM_93637378/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 17.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_airy**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2y' - yx = x^2 + 2$$

2.18.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -2$$

$$c = -1$$

$$F = x^2 + 2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x))$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(1+x)$$

$$y_2 = \text{AiryBi}(1+x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(1+x) & \text{AiryBi}(1+x) \\ \frac{d}{dx}(\text{AiryAi}(1+x)) & \frac{d}{dx}(\text{AiryBi}(1+x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(1+x) & \text{AiryBi}(1+x) \\ \text{AiryAi}(1, 1+x) & \text{AiryBi}(1, 1+x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(1+x))(\text{AiryBi}(1, 1+x)) - (\text{AiryBi}(1+x))(\text{AiryAi}(1, 1+x))$$

Which simplifies to

$$W = \text{AiryAi}(1+x)\text{AiryBi}(1, 1+x) - \text{AiryBi}(1+x)\text{AiryAi}(1, 1+x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(1+x)(x^2+2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(1+x)(x^2+2) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(1+\alpha)(\alpha^2+2) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(1+x)(x^2+2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(1+x)(x^2+2) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}(1+\alpha)(\alpha^2+2) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(1+\alpha)(\alpha^2+2) d\alpha \right)$$
$$u_2 = \pi \left(\int_0^x \text{AiryAi}(1+\alpha)(\alpha^2+2) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryAi}(1 + x) \\ + \pi \left(\int_0^x \text{AiryAi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryBi}(1 + x)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryAi}(1 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryBi}(1 + x) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = (e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x))) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryAi}(1 + x) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryBi}(1 + x) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryAi}(1 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryBi}(1 + x) \right) \\ + e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x))$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryAi}(1 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryBi}(1 + x) \right) \\ + e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x)) \quad (1)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryAi}(1 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1 + \alpha) (\alpha^2 + 2) d\alpha \right) \text{AiryBi}(1 + x) \right) \\ + e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x))$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x), x$2) - 2*diff(y(x), x) - x*y(x) - x^2 - 2 = 0, y(x), singsol=all)
```

$$y(x) = e^x \text{AiryAi}(x + 1) c_2 + e^x \text{AiryBi}(x + 1) c_1 - x$$

✓ Solution by Mathematica

Time used: 5.71 (sec). Leaf size: 87

```
DSolve[y''[x]-2*y'[x]-x*y[x]-x^2-2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x \left(\text{AiryAi}(x+1) \int_1^x -e^{-K[1]} \pi \text{AiryBi}(K[1]+1) (K[1]^2+2) dK[1] \right. \\ \left. + \text{AiryBi}(x+1) \int_1^x e^{-K[2]} \pi \text{AiryAi}(K[2]+1) (K[2]^2+2) dK[2] \right. \\ \left. + c_1 \text{AiryAi}(x+1) + c_2 \text{AiryBi}(x+1) \right)$$

2.19 problem 18

2.19.1 Solving as second order airy ode 898

Internal problem ID [7155]

Internal file name [OUTPUT/6141_Sunday_June_05_2022_04_24_55_PM_36421319/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 18.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 4y' - yx = x^2 + 4$$

2.19.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -4$$

$$c = -1$$

$$F = x^2 + 4$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{2x} (c_1 \text{AiryAi} (4 + x) + c_2 \text{AiryBi} (4 + x))$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(4 + x)$$

$$y_2 = \text{AiryBi}(4 + x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(4 + x) & \text{AiryBi}(4 + x) \\ \frac{d}{dx}(\text{AiryAi}(4 + x)) & \frac{d}{dx}(\text{AiryBi}(4 + x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(4 + x) & \text{AiryBi}(4 + x) \\ \text{AiryAi}(1, 4 + x) & \text{AiryBi}(1, 4 + x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(4 + x))(\text{AiryBi}(1, 4 + x)) - (\text{AiryBi}(4 + x))(\text{AiryAi}(1, 4 + x))$$

Which simplifies to

$$W = \text{AiryAi}(4 + x) \text{AiryBi}(1, 4 + x) - \text{AiryBi}(4 + x) \text{AiryAi}(1, 4 + x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(4+x)(x^2+4)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(4+x)(x^2+4) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(4+\alpha)(\alpha^2+4) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(4+x)(x^2+4)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(4+x)(x^2+4) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}(4+\alpha)(\alpha^2+4) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(4+\alpha)(\alpha^2+4) d\alpha \right)$$
$$u_2 = \pi \left(\int_0^x \text{AiryAi}(4+\alpha)(\alpha^2+4) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryAi}(4 + x) \\ + \pi \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryBi}(4 + x)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryAi}(4 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryBi}(4 + x) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = (e^{2x}(c_1 \text{AiryAi}(4 + x) + c_2 \text{AiryBi}(4 + x))) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryAi}(4 + x) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryBi}(4 + x) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryAi}(4 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryBi}(4 + x) \right) \\ + e^{2x}(c_1 \text{AiryAi}(4 + x) + c_2 \text{AiryBi}(4 + x))$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryAi}(4 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryBi}(4 + x) \right) \\ + e^{2x}(c_1 \text{AiryAi}(4 + x) + c_2 \text{AiryBi}(4 + x)) \quad (1)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryAi}(4 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^2 + 4) d\alpha \right) \text{AiryBi}(4 + x) \right) \\ + e^{2x} (c_1 \text{AiryAi}(4 + x) + c_2 \text{AiryBi}(4 + x))$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x), x$2) - 4*diff(y(x), x) - x*y(x) - x^2 - 4 = 0, y(x), singsol=all)
```

$$y(x) = e^{2x} \text{AiryAi}(x + 4) c_2 + e^{2x} \text{AiryBi}(x + 4) c_1 - x$$

✓ Solution by Mathematica

Time used: 6.139 (sec). Leaf size: 89

```
DSolve[y''[x]-4*y'[x]-x*y[x]-x^2-4==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x} \left(\text{AiryAi}(x+4) \int_1^x -e^{-2K[1]} \pi \text{AiryBi}(K[1]+4) (K[1]^2+4) dK[1] \right. \\ \left. + \text{AiryBi}(x+4) \int_1^x e^{-2K[2]} \pi \text{AiryAi}(K[2]+4) (K[2]^2+4) dK[2] \right. \\ \left. + c_1 \text{AiryAi}(x+4) + c_2 \text{AiryBi}(x+4) \right)$$

2.20 problem 19

2.20.1 Solving as second order airy ode 904

Internal problem ID [7156]

Internal file name [OUTPUT/6142_Sunday_June_05_2022_04_24_59_PM_5678011/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 19.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - y' - yx = x^3 - 1$$

2.20.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -1$$

$$c = -1$$

$$F = x^3 - 1$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi} \left(\frac{1}{4} + x \right)$$

$$y_2 = \text{AiryBi} \left(\frac{1}{4} + x \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \frac{d}{dx} (\text{AiryAi} \left(\frac{1}{4} + x \right)) & \frac{d}{dx} (\text{AiryBi} \left(\frac{1}{4} + x \right)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \text{AiryAi} \left(1, \frac{1}{4} + x \right) & \text{AiryBi} \left(1, \frac{1}{4} + x \right) \end{vmatrix}$$

Therefore

$$W = \left(\text{AiryAi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryBi} \left(1, \frac{1}{4} + x \right) \right) - \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryAi} \left(1, \frac{1}{4} + x \right) \right)$$

Which simplifies to

$$W = \text{AiryAi} \left(\frac{1}{4} + x \right) \text{AiryBi} \left(1, \frac{1}{4} + x \right) - \text{AiryBi} \left(\frac{1}{4} + x \right) \text{AiryAi} \left(1, \frac{1}{4} + x \right)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi} \left(\frac{1}{4} + x \right) (x^3 - 1)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi} \left(\frac{1}{4} + x \right) (x^3 - 1) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi} \left(\frac{1}{4} + x \right) (x^3 - 1)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi} \left(\frac{1}{4} + x \right) (x^3 - 1) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 1) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
            <- Bessel successful
        <- special function solution successful
    <- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 67

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-x*y(x)-x^3+1=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \left(-\text{AiryAi} \left(\frac{1}{4} + x \right) \pi \left(\int (x^3 - 1) \text{AiryBi} \left(\frac{1}{4} + x \right) e^{-\frac{x}{2}} dx \right) \right. \\ \left. + \text{AiryBi} \left(\frac{1}{4} + x \right) \pi \left(\int (x^3 - 1) \text{AiryAi} \left(\frac{1}{4} + x \right) e^{-\frac{x}{2}} dx \right) \right. \\ \left. + c_2 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_1 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

✓ Solution by Mathematica

Time used: 3.972 (sec). Leaf size: 107

```
DSolve[y''[x]-y'[x]-x*y[x]-x^3+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left(\text{AiryAi} \left(x + \frac{1}{4} \right) \int_1^x -e^{-\frac{K[1]}{2}} \pi \text{AiryBi} \left(K[1] + \frac{1}{4} \right) (K[1]^3 - 1) dK[1] \right. \\ \left. + \text{AiryBi} \left(x + \frac{1}{4} \right) \int_1^x e^{-\frac{K[2]}{2}} \pi \text{AiryAi} \left(K[2] + \frac{1}{4} \right) (K[2]^3 - 1) dK[2] \right. \\ \left. + c_1 \text{AiryAi} \left(x + \frac{1}{4} \right) + c_2 \text{AiryBi} \left(x + \frac{1}{4} \right) \right)$$

2.21 problem 20

2.21.1 Solving as second order airy ode 910

Internal problem ID [7157]

Internal file name [OUTPUT/6143_Sunday_June_05_2022_04_25_04_PM_69291286/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 20.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - 2y' - yx = x^3 + x^2$$

2.21.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -2$$

$$c = -1$$

$$F = x^2(1 + x)$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x))$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(1+x)$$

$$y_2 = \text{AiryBi}(1+x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(1+x) & \text{AiryBi}(1+x) \\ \frac{d}{dx}(\text{AiryAi}(1+x)) & \frac{d}{dx}(\text{AiryBi}(1+x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(1+x) & \text{AiryBi}(1+x) \\ \text{AiryAi}(1, 1+x) & \text{AiryBi}(1, 1+x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(1+x))(\text{AiryBi}(1, 1+x)) - (\text{AiryBi}(1+x))(\text{AiryAi}(1, 1+x))$$

Which simplifies to

$$W = \text{AiryAi}(1+x)\text{AiryBi}(1, 1+x) - \text{AiryBi}(1+x)\text{AiryAi}(1, 1+x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(1+x) x^2(1+x)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(1+x) x^2(1+x) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(1+\alpha) \alpha^2(1+\alpha) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(1+x) x^2(1+x)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(1+x) x^2(1+x) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}(1+\alpha) \alpha^2(1+\alpha) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryAi}(1+x) \\ + \pi \left(\int_0^x \text{AiryAi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryBi}(1+x)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryAi}(1+x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryBi}(1+x) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = (e^x(c_1 \text{AiryAi}(1+x) + c_2 \text{AiryBi}(1+x))) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryAi}(1+x) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryBi}(1+x) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryAi}(1+x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryBi}(1+x) \right) \\ + e^x(c_1 \text{AiryAi}(1+x) + c_2 \text{AiryBi}(1+x))$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryAi}(1+x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1+\alpha) \alpha^2(1+\alpha) d\alpha \right) \text{AiryBi}(1+x) \right) \\ + e^x(c_1 \text{AiryAi}(1+x) + c_2 \text{AiryBi}(1+x)) \quad (1)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(1 + \alpha) \alpha^2 (1 + \alpha) d\alpha \right) \text{AiryAi}(1 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1 + \alpha) \alpha^2 (1 + \alpha) d\alpha \right) \text{AiryBi}(1 + x) \right) \\ + e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x))$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)-x*y(x)-x^3-x^2=0,y(x), singsol=all)
```

$$y(x) = e^x \text{AiryAi}(x + 1) c_2 + e^x \text{AiryBi}(x + 1) c_1 - x^2 - x + 4$$

✓ Solution by Mathematica

Time used: 8.466 (sec). Leaf size: 91

```
DSolve[y''[x]-2*y'[x]-x*y[x]-x^3-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x \left(\text{AiryAi}(x+1) \int_1^x -e^{-K[1]} \pi \text{AiryBi}(K[1]+1) K[1]^2 (K[1]+1) dK[1] \right. \\ \left. + \text{AiryBi}(x+1) \int_1^x e^{-K[2]} \pi \text{AiryAi}(K[2]+1) K[2]^2 (K[2]+1) dK[2] \right. \\ \left. + c_1 \text{AiryAi}(x+1) + c_2 \text{AiryBi}(x+1) \right)$$

2.22 problem 21

2.22.1 Solving as second order airy ode 916

Internal problem ID [7158]

Internal file name [OUTPUT/6144_Sunday_June_05_2022_04_25_08_PM_25142294/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 21.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - y' - yx = x^3 - 2$$

2.22.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -1$$

$$c = -1$$

$$F = x^3 - 2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi} \left(\frac{1}{4} + x \right)$$

$$y_2 = \text{AiryBi} \left(\frac{1}{4} + x \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \frac{d}{dx} (\text{AiryAi} \left(\frac{1}{4} + x \right)) & \frac{d}{dx} (\text{AiryBi} \left(\frac{1}{4} + x \right)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \text{AiryAi} \left(1, \frac{1}{4} + x \right) & \text{AiryBi} \left(1, \frac{1}{4} + x \right) \end{vmatrix}$$

Therefore

$$W = \left(\text{AiryAi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryBi} \left(1, \frac{1}{4} + x \right) \right) - \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryAi} \left(1, \frac{1}{4} + x \right) \right)$$

Which simplifies to

$$W = \text{AiryAi} \left(\frac{1}{4} + x \right) \text{AiryBi} \left(1, \frac{1}{4} + x \right) - \text{AiryBi} \left(\frac{1}{4} + x \right) \text{AiryAi} \left(1, \frac{1}{4} + x \right)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi} \left(\frac{1}{4} + x \right) (x^3 - 2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi} \left(\frac{1}{4} + x \right) (x^3 - 2) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi} \left(\frac{1}{4} + x \right) (x^3 - 2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi} \left(\frac{1}{4} + x \right) (x^3 - 2) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^3 - 2) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-x*y(x)-x^3+2=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \text{AiryAi} \left(\frac{1}{4} + x \right) c_2 + e^{\frac{x}{2}} \text{AiryBi} \left(\frac{1}{4} + x \right) c_1 - x^2 + 2$$

✓ Solution by Mathematica

Time used: 3.963 (sec). Leaf size: 107

```
DSolve[y''[x]-y'[x]-x*y[x]-x^3+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left(\text{AiryAi} \left(x + \frac{1}{4} \right) \int_1^x -e^{-\frac{K[1]}{2}} \pi \text{AiryBi} \left(K[1] + \frac{1}{4} \right) (K[1]^3 - 2) dK[1] \right. \\ \left. + \text{AiryBi} \left(x + \frac{1}{4} \right) \int_1^x e^{-\frac{K[2]}{2}} \pi \text{AiryAi} \left(K[2] + \frac{1}{4} \right) (K[2]^3 - 2) dK[2] \right. \\ \left. + c_1 \text{AiryAi} \left(x + \frac{1}{4} \right) + c_2 \text{AiryBi} \left(x + \frac{1}{4} \right) \right)$$

2.23 problem 22

2.23.1 Solving as second order airy ode 922

Internal problem ID [7159]

Internal file name [OUTPUT/6145_Sunday_June_05_2022_04_25_12_PM_11559479/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 22.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - 2y' - yx = x^3 - 2$$

2.23.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -2$$

$$c = -1$$

$$F = x^3 - 2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x))$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(1+x)$$

$$y_2 = \text{AiryBi}(1+x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(1+x) & \text{AiryBi}(1+x) \\ \frac{d}{dx}(\text{AiryAi}(1+x)) & \frac{d}{dx}(\text{AiryBi}(1+x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(1+x) & \text{AiryBi}(1+x) \\ \text{AiryAi}(1, 1+x) & \text{AiryBi}(1, 1+x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(1+x))(\text{AiryBi}(1, 1+x)) - (\text{AiryBi}(1+x))(\text{AiryAi}(1, 1+x))$$

Which simplifies to

$$W = \text{AiryAi}(1+x)\text{AiryBi}(1, 1+x) - \text{AiryBi}(1+x)\text{AiryAi}(1, 1+x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(1+x)(x^3-2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(1+x)(x^3-2) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(1+\alpha)(\alpha^3-2) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(1+x)(x^3-2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(1+x)(x^3-2) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}(1+\alpha)(\alpha^3-2) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(1+\alpha)(\alpha^3-2) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}(1+\alpha)(\alpha^3-2) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(1+x) \\ + \pi \left(\int_0^x \text{AiryAi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(1+x)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(1+x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(1+x) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = (e^x (c_1 \text{AiryAi}(1+x) + c_2 \text{AiryBi}(1+x))) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(1+x) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(1+x) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(1+x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(1+x) \right) \\ + e^x (c_1 \text{AiryAi}(1+x) + c_2 \text{AiryBi}(1+x))$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(1+x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1+\alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(1+x) \right) \\ + e^x (c_1 \text{AiryAi}(1+x) + c_2 \text{AiryBi}(1+x)) \quad (1)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(1 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(1 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(1 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(1 + x) \right) \\ + e^x (c_1 \text{AiryAi}(1 + x) + c_2 \text{AiryBi}(1 + x))$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
  checking if the LODE has constant coefficients  
  checking if the LODE is of Euler type  
  trying a symmetry of the form [xi=0, eta=F(x)]  
  checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
  -> Bessel  
    <- Bessel successful  
  <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)-x*y(x)-x^3+2=0,y(x), singsol=all)
```

$$y(x) = e^x \text{AiryAi}(x + 1) c_2 + e^x \text{AiryBi}(x + 1) c_1 - x^2 + 4$$

✓ Solution by Mathematica

Time used: 2.673 (sec). Leaf size: 87

```
DSolve[y''[x]-2*y'[x]-x*y[x]-x^3+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x \left(\text{AiryAi}(x+1) \int_1^x -e^{-K[1]} \pi \text{AiryBi}(K[1]+1) (K[1]^3 - 2) dK[1] \right. \\ \left. + \text{AiryBi}(x+1) \int_1^x e^{-K[2]} \pi \text{AiryAi}(K[2]+1) (K[2]^3 - 2) dK[2] \right. \\ \left. + c_1 \text{AiryAi}(x+1) + c_2 \text{AiryBi}(x+1) \right)$$

2.24 problem 23

2.24.1 Solving as second order airy ode 928

Internal problem ID [7160]

Internal file name [OUTPUT/6146_Sunday_June_05_2022_04_25_16_PM_96489219/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 23.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - 4y' - yx = x^3 - 2$$

2.24.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -4$$

$$c = -1$$

$$F = x^3 - 2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{2x} (c_1 \text{AiryAi} (4 + x) + c_2 \text{AiryBi} (4 + x))$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(4 + x)$$

$$y_2 = \text{AiryBi}(4 + x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(4 + x) & \text{AiryBi}(4 + x) \\ \frac{d}{dx}(\text{AiryAi}(4 + x)) & \frac{d}{dx}(\text{AiryBi}(4 + x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(4 + x) & \text{AiryBi}(4 + x) \\ \text{AiryAi}(1, 4 + x) & \text{AiryBi}(1, 4 + x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(4 + x))(\text{AiryBi}(1, 4 + x)) - (\text{AiryBi}(4 + x))(\text{AiryAi}(1, 4 + x))$$

Which simplifies to

$$W = \text{AiryAi}(4 + x) \text{AiryBi}(1, 4 + x) - \text{AiryBi}(4 + x) \text{AiryAi}(1, 4 + x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(4+x)(x^3-2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(4+x)(x^3-2) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(4+\alpha)(\alpha^3-2) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(4+x)(x^3-2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(4+x)(x^3-2) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}(4+\alpha)(\alpha^3-2) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(4+\alpha)(\alpha^3-2) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}(4+\alpha)(\alpha^3-2) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(4 + x) \\ + \pi \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(4 + x)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(4 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(4 + x) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = (e^{2x}(c_1 \text{AiryAi}(4 + x) + c_2 \text{AiryBi}(4 + x))) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(4 + x) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(4 + x) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(4 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(4 + x) \right) \\ + e^{2x}(c_1 \text{AiryAi}(4 + x) + c_2 \text{AiryBi}(4 + x))$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(4 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(4 + x) \right) \\ + e^{2x}(c_1 \text{AiryAi}(4 + x) + c_2 \text{AiryBi}(4 + x))$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(4 + x) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(4 + \alpha) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(4 + x) \right) \\ + e^{2x} (c_1 \text{AiryAi}(4 + x) + c_2 \text{AiryBi}(4 + x))$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x), x$2) - 4*diff(y(x), x) - x*y(x) - x^3 + 2 = 0, y(x), singsol=all)
```

$$y(x) = e^{2x} \text{AiryAi}(x + 4) c_2 + e^{2x} \text{AiryBi}(x + 4) c_1 - x^2 + 8$$

✓ Solution by Mathematica

Time used: 2.795 (sec). Leaf size: 89

```
DSolve[y''[x]-4*y'[x]-x*y[x]-x^3+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x} \left(\text{AiryAi}(x+4) \int_1^x -e^{-2K[1]} \pi \text{AiryBi}(K[1]+4) (K[1]^3 - 2) dK[1] \right. \\ \left. + \text{AiryBi}(x+4) \int_1^x e^{-2K[2]} \pi \text{AiryAi}(K[2]+4) (K[2]^3 - 2) dK[2] \right. \\ \left. + c_1 \text{AiryAi}(x+4) + c_2 \text{AiryBi}(x+4) \right)$$

2.25 problem 24

2.25.1 Solving as second order airy ode 934

Internal problem ID [7161]

Internal file name [OUTPUT/6147_Sunday_June_05_2022_04_25_20_PM_89844936/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 24.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - 6y' - yx = x^3 - 2$$

2.25.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -6$$

$$c = -1$$

$$F = x^3 - 2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{3x} (c_1 \text{AiryAi} (x + 9) + c_2 \text{AiryBi} (x + 9))$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x + 9)$$

$$y_2 = \text{AiryBi}(x + 9)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x + 9) & \text{AiryBi}(x + 9) \\ \frac{d}{dx}(\text{AiryAi}(x + 9)) & \frac{d}{dx}(\text{AiryBi}(x + 9)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x + 9) & \text{AiryBi}(x + 9) \\ \text{AiryAi}(1, x + 9) & \text{AiryBi}(1, x + 9) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x + 9))(\text{AiryBi}(1, x + 9)) - (\text{AiryBi}(x + 9))(\text{AiryAi}(1, x + 9))$$

Which simplifies to

$$W = \text{AiryAi}(x + 9) \text{AiryBi}(1, x + 9) - \text{AiryBi}(x + 9) \text{AiryAi}(1, x + 9)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x+9)(x^3-2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x+9)(x^3-2) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(\alpha+9)(\alpha^3-2) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x+9)(x^3-2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x+9)(x^3-2) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}(\alpha+9)(\alpha^3-2) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(\alpha+9)(\alpha^3-2) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}(\alpha+9)(\alpha^3-2) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 9) \\ + \pi \left(\int_0^x \text{AiryAi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 9)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 9) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 9) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = (e^{3x}(c_1 \text{AiryAi}(x + 9) + c_2 \text{AiryBi}(x + 9))) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 9) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 9) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 9) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 9) \right) \\ + e^{3x}(c_1 \text{AiryAi}(x + 9) + c_2 \text{AiryBi}(x + 9))$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 9) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 9) \right) \\ + e^{3x}(c_1 \text{AiryAi}(x + 9) + c_2 \text{AiryBi}(x + 9))$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 9) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(\alpha + 9) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 9) \right) \\ + e^{3x} (c_1 \text{AiryAi}(x + 9) + c_2 \text{AiryBi}(x + 9))$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve(diff(y(x), x$2) - 6*diff(y(x), x) - x*y(x) - x^3 + 2 = 0, y(x), singsol=all)
```

$$y(x) = e^{3x} \text{AiryAi}(9 + x) c_2 + e^{3x} \text{AiryBi}(9 + x) c_1 - x^2 + 12$$

✓ Solution by Mathematica

Time used: 6.656 (sec). Leaf size: 89

```
DSolve[y''[x]-6*y'[x]-x*y[x]-x^3+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x} \left(\text{AiryAi}(x+9) \int_1^x -e^{-3K[1]} \pi \text{AiryBi}(K[1]+9) (K[1]^3 - 2) dK[1] \right. \\ \left. + \text{AiryBi}(x+9) \int_1^x e^{-3K[2]} \pi \text{AiryAi}(K[2]+9) (K[2]^3 - 2) dK[2] \right. \\ \left. + c_1 \text{AiryAi}(x+9) + c_2 \text{AiryBi}(x+9) \right)$$

2.26 problem 25

2.26.1 Solving as second order airy ode 940

Internal problem ID [7162]

Internal file name [OUTPUT/6148_Sunday_June_05_2022_04_25_24_PM_69506047/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 25.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - 8y' - yx = x^3 - 2$$

2.26.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -8$$

$$c = -1$$

$$F = x^3 - 2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{4x} (c_1 \text{AiryAi}(x + 16) + c_2 \text{AiryBi}(x + 16))$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x + 16)$$

$$y_2 = \text{AiryBi}(x + 16)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x + 16) & \text{AiryBi}(x + 16) \\ \frac{d}{dx}(\text{AiryAi}(x + 16)) & \frac{d}{dx}(\text{AiryBi}(x + 16)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x + 16) & \text{AiryBi}(x + 16) \\ \text{AiryAi}(1, x + 16) & \text{AiryBi}(1, x + 16) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x + 16)) (\text{AiryBi}(1, x + 16)) - (\text{AiryBi}(x + 16)) (\text{AiryAi}(1, x + 16))$$

Which simplifies to

$$W = \text{AiryAi}(x + 16) \text{AiryBi}(1, x + 16) - \text{AiryBi}(x + 16) \text{AiryAi}(1, x + 16)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x + 16) (x^3 - 2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x + 16) (x^3 - 2) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(\alpha + 16) (\alpha^3 - 2) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x + 16) (x^3 - 2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x + 16) (x^3 - 2) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}(\alpha + 16) (\alpha^3 - 2) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 16) \\ + \pi \left(\int_0^x \text{AiryAi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 16)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 16) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 16) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = (e^{4x}(c_1 \text{AiryAi}(x + 16) + c_2 \text{AiryBi}(x + 16))) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 16) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 16) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 16) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 16) \right) \\ + e^{4x}(c_1 \text{AiryAi}(x + 16) + c_2 \text{AiryBi}(x + 16))$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 16) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 16) \right) \\ + e^{4x}(c_1 \text{AiryAi}(x + 16) + c_2 \text{AiryBi}(x + 16)) \quad (1)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryAi}(x + 16) \right. \\ \left. + \left(\int_0^x \text{AiryAi}(\alpha + 16) (\alpha^3 - 2) d\alpha \right) \text{AiryBi}(x + 16) \right) \\ + e^{4x} (c_1 \text{AiryAi}(x + 16) + c_2 \text{AiryBi}(x + 16))$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$2)-8*diff(y(x),x)-x*y(x)-x^3+2=0,y(x), singsol=all)
```

$$y(x) = e^{4x} \text{AiryAi}(16 + x) c_2 + e^{4x} \text{AiryBi}(16 + x) c_1 - x^2 + 16$$

✓ Solution by Mathematica

Time used: 6.555 (sec). Leaf size: 89

```
DSolve[y''[x]-8*y'[x]-x*y[x]-x^3+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{4x} \left(\text{AiryAi}(x + 16) \int_1^x -e^{-4K[1]} \pi \text{AiryBi}(K[1] + 16) (K[1]^3 - 2) dK[1] \right. \\ \left. + \text{AiryBi}(x + 16) \int_1^x e^{-4K[2]} \pi \text{AiryAi}(K[2] + 16) (K[2]^3 - 2) dK[2] \right. \\ \left. + c_1 \text{AiryAi}(x + 16) + c_2 \text{AiryBi}(x + 16) \right)$$

2.27 problem 26

2.27.1 Solving as second order airy ode 946

Internal problem ID [7163]

Internal file name [OUTPUT/6149_Sunday_June_05_2022_04_25_28_PM_78560358/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 26.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - y' - yx = x^4 - 3$$

2.27.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -1$$

$$c = -1$$

$$F = x^4 - 3$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi} \left(\frac{1}{4} + x \right)$$

$$y_2 = \text{AiryBi} \left(\frac{1}{4} + x \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \frac{d}{dx} (\text{AiryAi} \left(\frac{1}{4} + x \right)) & \frac{d}{dx} (\text{AiryBi} \left(\frac{1}{4} + x \right)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \text{AiryAi} \left(1, \frac{1}{4} + x \right) & \text{AiryBi} \left(1, \frac{1}{4} + x \right) \end{vmatrix}$$

Therefore

$$W = \left(\text{AiryAi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryBi} \left(1, \frac{1}{4} + x \right) \right) - \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryAi} \left(1, \frac{1}{4} + x \right) \right)$$

Which simplifies to

$$W = \text{AiryAi} \left(\frac{1}{4} + x \right) \text{AiryBi} \left(1, \frac{1}{4} + x \right) - \text{AiryBi} \left(\frac{1}{4} + x \right) \text{AiryAi} \left(1, \frac{1}{4} + x \right)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi} \left(\frac{1}{4} + x \right) (x^4 - 3)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi} \left(\frac{1}{4} + x \right) (x^4 - 3) \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi} \left(\frac{1}{4} + x \right) (x^4 - 3)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi} \left(\frac{1}{4} + x \right) (x^4 - 3) \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) (\alpha^4 - 3) d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-x*y(x)-x^4+3=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \text{AiryAi} \left(\frac{1}{4} + x \right) c_2 + e^{\frac{x}{2}} \text{AiryBi} \left(\frac{1}{4} + x \right) c_1 - x^3 + 3x - 6$$

✓ Solution by Mathematica

Time used: 4.059 (sec). Leaf size: 107

```
DSolve[y''[x]-y'[x]-x*y[x]-x^4+3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left(\text{AiryAi} \left(x + \frac{1}{4} \right) \int_1^x -e^{-\frac{K[1]}{2}} \pi \text{AiryBi} \left(K[1] + \frac{1}{4} \right) (K[1]^4 - 3) dK[1] \right. \\ \left. + \text{AiryBi} \left(x + \frac{1}{4} \right) \int_1^x e^{-\frac{K[2]}{2}} \pi \text{AiryAi} \left(K[2] + \frac{1}{4} \right) (K[2]^4 - 3) dK[2] \right. \\ \left. + c_1 \text{AiryAi} \left(x + \frac{1}{4} \right) + c_2 \text{AiryBi} \left(x + \frac{1}{4} \right) \right)$$

2.28 problem 27

2.28.1 Solving as second order airy ode 952

Internal problem ID [7164]

Internal file name [OUTPUT/6150_Sunday_June_05_2022_04_25_34_PM_99044401/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 27.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - y' - yx = x^3$$

2.28.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = -1$$

$$c = -1$$

$$F = x^3$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi} \left(\frac{1}{4} + x \right)$$

$$y_2 = \text{AiryBi} \left(\frac{1}{4} + x \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \frac{d}{dx} (\text{AiryAi} \left(\frac{1}{4} + x \right)) & \frac{d}{dx} (\text{AiryBi} \left(\frac{1}{4} + x \right)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi} \left(\frac{1}{4} + x \right) & \text{AiryBi} \left(\frac{1}{4} + x \right) \\ \text{AiryAi} \left(1, \frac{1}{4} + x \right) & \text{AiryBi} \left(1, \frac{1}{4} + x \right) \end{vmatrix}$$

Therefore

$$W = \left(\text{AiryAi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryBi} \left(1, \frac{1}{4} + x \right) \right) - \left(\text{AiryBi} \left(\frac{1}{4} + x \right) \right) \left(\text{AiryAi} \left(1, \frac{1}{4} + x \right) \right)$$

Which simplifies to

$$W = \text{AiryAi}\left(\frac{1}{4} + x\right) \text{AiryBi}\left(1, \frac{1}{4} + x\right) - \text{AiryBi}\left(\frac{1}{4} + x\right) \text{AiryAi}\left(1, \frac{1}{4} + x\right)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}\left(\frac{1}{4} + x\right) x^3}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}\left(\frac{1}{4} + x\right) x^3 \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}\left(\frac{1}{4} + \alpha\right) \alpha^3 \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}\left(\frac{1}{4} + x\right) x^3}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}\left(\frac{1}{4} + x\right) x^3 \pi dx$$

Hence

$$u_2 = \int_0^x \text{AiryAi}\left(\frac{1}{4} + \alpha\right) \alpha^3 \pi d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}\left(\frac{1}{4} + \alpha\right) \alpha^3 d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \text{AiryAi}\left(\frac{1}{4} + \alpha\right) \alpha^3 d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \\ + \pi \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right)$$

Which simplifies to

$$y_p(x) = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ + \left(\pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \right. \\ \left. \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \right) \\ = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Summary

The solution(s) found are the following

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verification of solutions

$$y = \pi \left(- \left(\int_0^x \text{AiryBi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + \left(\int_0^x \text{AiryAi} \left(\frac{1}{4} + \alpha \right) \alpha^3 d\alpha \right) \text{AiryBi} \left(\frac{1}{4} + x \right) \right) \\ + e^{\frac{x}{2}} \left(c_1 \text{AiryAi} \left(\frac{1}{4} + x \right) + c_2 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
            <- Bessel successful
        <- special function solution successful
    <- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 63

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-x*y(x)-x^3=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \left(-\text{AiryAi} \left(\frac{1}{4} + x \right) \pi \left(\int x^3 \text{AiryBi} \left(\frac{1}{4} + x \right) e^{-\frac{x}{2}} dx \right) \right. \\ \left. + \text{AiryBi} \left(\frac{1}{4} + x \right) \pi \left(\int x^3 \text{AiryAi} \left(\frac{1}{4} + x \right) e^{-\frac{x}{2}} dx \right) + c_2 \text{AiryAi} \left(\frac{1}{4} + x \right) \right. \\ \left. + c_1 \text{AiryBi} \left(\frac{1}{4} + x \right) \right)$$

✓ Solution by Mathematica

Time used: 10.277 (sec). Leaf size: 103

```
DSolve[y''[x]-y'[x]-x*y[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left(\text{AiryAi} \left(x + \frac{1}{4} \right) \int_1^x -e^{-\frac{K[1]}{2}} \pi \text{AiryBi} \left(K[1] + \frac{1}{4} \right) K[1]^3 dK[1] \right. \\ \left. + \text{AiryBi} \left(x + \frac{1}{4} \right) \int_1^x e^{-\frac{K[2]}{2}} \pi \text{AiryAi} \left(K[2] + \frac{1}{4} \right) K[2]^3 dK[2] \right. \\ \left. + c_1 \text{AiryAi} \left(x + \frac{1}{4} \right) + c_2 \text{AiryBi} \left(x + \frac{1}{4} \right) \right)$$

2.29 problem 28

2.29.1 Solving as second order airy ode 958

2.29.2 Solving as second order besel ode ode 962

Internal problem ID [7165]

Internal file name [OUTPUT/6151_Sunday_June_05_2022_04_25_38_PM_22911779/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 28.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_bessel_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - yx = x^3 - 2$$

2.29.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = 0$$

$$c = -1$$

$$F = x^3 - 2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) (x^3 - 2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) (x^3 - 2) \pi dx$$

Hence

$$u_1 = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) x^4 - 5 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) x + \frac{5\pi 3^{\frac{5}{6}} (\text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right))}{10\Gamma\left(\frac{2}{3}\right)}}{10\Gamma\left(\frac{2}{3}\right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) (x^3 - 2)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) (x^3 - 2) \pi dx$$

Hence

$$u_2 = \frac{x \left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) x^4 - 5 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) - \frac{5\pi 3^{\frac{1}{3}} (\text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right))}{10\Gamma\left(\frac{2}{3}\right)}}{10\Gamma\left(\frac{2}{3}\right)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) x^4 - 5 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) x + \frac{5\pi 3^{\frac{5}{6}} (\text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right))}{10 \Gamma\left(\frac{2}{3}\right)}}{10 \Gamma\left(\frac{2}{3}\right)} - \frac{x \left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) x^4 - 5 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) - \frac{5\pi 3^{\frac{1}{3}} (\text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right))}{10 \Gamma\left(\frac{2}{3}\right)}}{10 \Gamma\left(\frac{2}{3}\right)}$$

Which simplifies to

$$y_p(x) = \frac{\left(\frac{20\pi (\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right)}{3} + x \left(-5 \Gamma\left(\frac{2}{3}\right)^2 (\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}}) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right) \right)}{3} \right)}{3}$$

Therefore the general solution is

$$y = y_h + y_p = (c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)) + \frac{\left(\frac{20\pi (\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right)}{3} + x \left(-5 \Gamma\left(\frac{2}{3}\right)^2 (\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}}) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right) \right)}{3} \right)}{3} = \frac{\left(\frac{20\pi (\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right)}{3} + x \left(-5 \Gamma\left(\frac{2}{3}\right)^2 (\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}}) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right) \right)}{3} \right)}{3} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Summary

The solution(s) found are the following

$$y = \frac{\left(\frac{20\pi (\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right)}{3} + x \left(-5 \Gamma\left(\frac{2}{3}\right)^2 (\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}}) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right) \right)}{3} \right)}{3} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x) \tag{1}$$

Verification of solutions

$$y = \frac{20\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom}\left(\left[\frac{1}{3}\right], \left[\frac{2}{3}, \frac{4}{3}\right], \frac{x^3}{9}\right)}{3} + x \left(-5\Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}} \right) \text{hypergeom}\left(\left[\frac{1}{3}\right], \left[\frac{2}{3}, \frac{4}{3}\right], \frac{x^3}{9}\right) \right)$$

$$+ c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Verified OK.

2.29.2 Solving as second order bessel ode

Writing the ode as

$$x^2 y'' - y x^3 = x^2 (x^3 - 2) \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + x y' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) x y' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\alpha = \frac{1}{2}$$

$$\beta = \frac{2i}{3}$$

$$n = \frac{1}{3}$$

$$\gamma = \frac{3}{2}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + c_2 \sqrt{x} \text{BesselY}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1\sqrt{x} \operatorname{BesselJ}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + c_2\sqrt{x} \operatorname{BesselY}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \operatorname{AiryAi}(x)$$

$$y_2 = \operatorname{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \operatorname{AiryAi}(x) & \operatorname{AiryBi}(x) \\ \frac{d}{dx}(\operatorname{AiryAi}(x)) & \frac{d}{dx}(\operatorname{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \operatorname{AiryAi}(x) & \operatorname{AiryBi}(x) \\ \operatorname{AiryAi}(1, x) & \operatorname{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x)) (\text{AiryBi}(1, x)) - (\text{AiryBi}(x)) (\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) x^2 (x^3 - 2)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) (x^3 - 2) \pi dx$$

Hence

$$u_1 = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) x^4 - 5 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) x + \frac{5\pi 3^{\frac{5}{6}} (\text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right))}{10 \Gamma\left(\frac{2}{3}\right)} \right)}{10 \Gamma\left(\frac{2}{3}\right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) x^2 (x^3 - 2)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) (x^3 - 2) \pi dx$$

Hence

$$u_2 = \frac{x \left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) x^4 - 5 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) - \frac{5\pi 3^{\frac{1}{3}} (\text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right))}{10 \Gamma\left(\frac{2}{3}\right)} \right)}{10 \Gamma\left(\frac{2}{3}\right)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) x^4 - 5\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) x + \frac{5\pi 3^{\frac{5}{6}} (\text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right))}{10\Gamma\left(\frac{2}{3}\right)} \right)}{10\Gamma\left(\frac{2}{3}\right)} - \frac{x \left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) x^4 - 5 \cdot 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) - \frac{5\pi 3^{\frac{1}{3}} (\text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right))}{10\Gamma\left(\frac{2}{3}\right)} \right)}{10\Gamma\left(\frac{2}{3}\right)}$$

Which simplifies to

$$y_p(x) = \frac{\left(\frac{20\pi (\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right)}{3} + x \left(-5\Gamma\left(\frac{2}{3}\right)^2 (\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}}) \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) \right) \right)}{10\Gamma\left(\frac{2}{3}\right)}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \right) + \frac{\left(\frac{20\pi (\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right)}{3} + x \left(-5\Gamma\left(\frac{2}{3}\right)^2 (\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}}) \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) \right) \right)}{10\Gamma\left(\frac{2}{3}\right)}$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + \frac{\left(\frac{20\pi (\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)) \text{hypergeom} \left(\left[\frac{1}{3} \right], \left[\frac{2}{3}, \frac{4}{3} \right], \frac{x^3}{9} \right)}{3} + x \left(-5\Gamma\left(\frac{2}{3}\right)^2 (\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}}) \text{hypergeom} \left(\left[\frac{2}{3} \right], \left[\frac{4}{3}, \frac{5}{3} \right], \frac{x^3}{9} \right) \right) \right)}{10\Gamma\left(\frac{2}{3}\right)} \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \operatorname{BesselJ}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + c_2 \sqrt{x} \operatorname{BesselY}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) \\ + \frac{20\pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x)\right) \operatorname{hypergeom}\left(\left[\frac{1}{3}\right], \left[\frac{2}{3}, \frac{4}{3}\right], \frac{x^3}{9}\right)}{3} + x \left(-5\Gamma\left(\frac{2}{3}\right)^2 \left(\operatorname{AiryAi}(x) 3^{\frac{2}{3}} + \operatorname{AiryBi}(x) 3^{\frac{1}{6}}\right) \operatorname{hype}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(diff(y(x), x$2) - x*y(x) - x^3 + 2 = 0, y(x), singsol=all)
```

$$y(x) = \operatorname{AiryAi}(x) c_2 + \operatorname{AiryBi}(x) c_1 - x^2$$

✓ Solution by Mathematica

Time used: 0.458 (sec). Leaf size: 290

```
DSolve[y''[x]-x*y[x]-x^3+2==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$\rightarrow \frac{6\sqrt[3]{3}\pi x \Gamma\left(\frac{1}{3}\right) \Gamma\left(\frac{5}{3}\right) \Gamma\left(\frac{7}{3}\right) \Gamma\left(\frac{8}{3}\right) (\sqrt{3} \text{AiryAi}(x) - \text{AiryBi}(x)) {}_1F_2\left(\frac{1}{3}; \frac{2}{3}, \frac{4}{3}; \frac{x^3}{9}\right)}{\dots}$

2.30 problem 29

- 2.30.1 Solving as second order airy ode 968
2.30.2 Solving as second order besseel ode ode 972

Internal problem ID [7166]

Internal file name [OUTPUT/6152_Sunday_June_05_2022_04_25_39_PM_88887002/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 29.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_besseel_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - yx = x^6 - 64$$

2.30.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$\begin{aligned} a &= 1 \\ b &= 0 \\ c &= -1 \\ F &= x^6 - 64 \end{aligned}$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) (x^6 - 64)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) (x^6 - 64) \pi dx$$

Hence

$$u_1 = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7 + \frac{16\pi \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) 3^{\frac{5}{6}} x^6}{21} - 256 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7}{16 \Gamma\left(\frac{2}{3}\right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) (x^6 - 64)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) (x^6 - 64) \pi dx$$

Hence

$$u_2 = \frac{\left(\Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) 3^{\frac{1}{6}} x^7 - \frac{16\pi 3^{\frac{1}{3}} \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) x^6}{21} - 256 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7}{16 \Gamma\left(\frac{2}{3}\right)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7 + \frac{16\pi \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) 3^{\frac{5}{6}} x^6 - 256 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7}{16 \Gamma\left(\frac{2}{3}\right)} - \frac{\left(\Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) 3^{\frac{1}{6}} x^7 - \frac{16\pi 3^{\frac{1}{3}} \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) x^6 - 256 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7}{16 \Gamma\left(\frac{2}{3}\right)}}$$

Which simplifies to

$$y_p(x) = \left(-\frac{16x^6\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}} \right) \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) \right)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)) \\ &\quad + \left(-\frac{16x^6\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}} \right) \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) \right) \\ &= \left(-\frac{16x^6\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}} \right) \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) \right) \\ &\quad + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{16x^6\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}} \right) \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) \right) + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x) \tag{1}$$

Verification of solutions

$y =$

$$\left(-\frac{16x^6\pi(\text{AiryBi}(x)3^{\frac{1}{3}} - 3^{\frac{5}{6}}\text{AiryAi}(x))\text{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right)}{21} + x^7\Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x)3^{\frac{2}{3}} + \text{AiryBi}(x)3^{\frac{1}{6}}\right) \text{hyp}$$

$$+ c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Verified OK.

2.30.2 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' - yx^3 = x^2(x^6 - 64) \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= \frac{1}{2} \\ \beta &= \frac{2i}{3} \\ n &= \frac{1}{3} \\ \gamma &= \frac{3}{2} \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + c_2 \sqrt{x} \text{BesselY}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x)) (\text{AiryBi}(1, x)) - (\text{AiryBi}(x)) (\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) x^2 (x^6 - 64)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) (x^6 - 64) \pi dx$$

Hence

$$u_1 = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7 + \frac{16\pi \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) 3^{\frac{5}{6}} x^6 - 256 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7}{16 \Gamma\left(\frac{2}{3}\right)^2}}{16 \Gamma\left(\frac{2}{3}\right)^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) x^2 (x^6 - 64)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) (x^6 - 64) \pi dx$$

Hence

$$u_2 = \frac{\left(\Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) 3^{\frac{1}{6}} x^7 - \frac{16\pi 3^{\frac{1}{3}} \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) x^6 - 256 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7}{16 \Gamma\left(\frac{2}{3}\right)^2}}{16 \Gamma\left(\frac{2}{3}\right)^2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7 + \frac{16\pi \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) 3^{\frac{5}{6}} x^6 - 256 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) x^7}{16 \Gamma\left(\frac{2}{3}\right)} - \frac{\left(\Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) 3^{\frac{1}{6}} x^7 - \frac{16\pi 3^{\frac{1}{3}} \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) x^6 - 256 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right)}{16 \Gamma\left(\frac{2}{3}\right)}$$

Which simplifies to

$$y_p(x) = \left(-\frac{16x^6 \pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}} \right) \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) \right)$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \right) + \left(-\frac{16x^6 \pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}} \right) \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + \left(-\frac{16x^6 \pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}} \right) \text{hypergeom} \left(\left[\frac{8}{3} \right], \left[\frac{4}{3}, \frac{11}{3} \right], \frac{x^3}{9} \right) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \operatorname{BesselJ}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + c_2 \sqrt{x} \operatorname{BesselY}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)$$

$$\left(-\frac{16x^6\pi\left(\operatorname{AiryBi}(x)3^{\frac{1}{3}}-3^{\frac{5}{6}}\operatorname{AiryAi}(x)\right)\operatorname{hypergeom}\left(\left[\frac{7}{3}\right],\left[\frac{2}{3},\frac{10}{3}\right],\frac{x^3}{9}\right)}{21} + x^7\Gamma\left(\frac{2}{3}\right)^2\left(\operatorname{AiryAi}(x)3^{\frac{2}{3}} + \operatorname{AiryBi}(x)3^{\frac{1}{6}}\right)\operatorname{hyp}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 149

```
dsolve(diff(y(x),x$2)-x*y(x)-x^6+64=0,y(x), singsol=all)
```

$y(x)$

$$= \frac{16x^7\pi\left(\operatorname{AiryBi}(x)3^{\frac{1}{3}} - 3^{\frac{5}{6}}\operatorname{AiryAi}(x)\right)\operatorname{hypergeom}\left(\left[\frac{7}{3}\right],\left[\frac{2}{3},\frac{10}{3}\right],\frac{x^3}{9}\right) - 21x^8\Gamma\left(\frac{2}{3}\right)^2\left(3^{\frac{1}{6}}\operatorname{AiryBi}(x) + 3^{\frac{2}{3}}\right)}{21}$$

✓ Solution by Mathematica

Time used: 0.493 (sec). Leaf size: 256

```
DSolve[y''[x]-x*y[x]-x^6+64==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{192\sqrt[3]{3}\pi x \Gamma\left(\frac{1}{3}\right) \left(\sqrt{3} \operatorname{AiryAi}(x) - \operatorname{AiryBi}(x)\right) {}_1F_2\left(\frac{1}{3}; \frac{2}{3}, \frac{4}{3}; \frac{x^3}{9}\right) - \sqrt[6]{3}\pi x^8 \Gamma\left(\frac{2}{3}\right) \Gamma\left(\frac{8}{3}\right) \left(3 \operatorname{AiryAi}(x) - \operatorname{AiryBi}(x)\right)}{\Gamma\left(\frac{1}{3}\right) \Gamma\left(\frac{2}{3}\right) \Gamma\left(\frac{8}{3}\right)}$$

2.31 problem 30

- 2.31.1 Solving as second order airy ode 978
- 2.31.2 Solving as second order besseel ode ode 982

Internal problem ID [7167]

Internal file name [OUTPUT/6153_Sunday_June_05_2022_04_25_41_PM_9233294/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 30.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_besseel_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - yx = x$$

2.31.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = 0$$

$$c = -1$$

$$F = x$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) x}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) x \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(\alpha) \alpha \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x \text{AiryAi}(x)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) x \pi dx$$

Hence

$$u_2 = - \frac{x^3 3^{\frac{1}{6}} \text{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)}{6} + \frac{\sqrt{x} \left(x^{\frac{3}{2}} \right)^{\frac{1}{3}} \pi \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3}$$

Which simplifies to

$$u_1 = - \pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right)$$

$$u_2 = - \frac{x^3 3^{\frac{1}{6}} \text{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)}{6} + \frac{\sqrt{x} \left(x^{\frac{3}{2}} \right)^{\frac{1}{3}} \pi \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right) \text{AiryAi}(x) + \left(-\frac{x^3 3^{\frac{1}{6}} \text{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3} \right) \text{AiryBi}(x)$$

Which simplifies to

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right) \text{AiryAi}(x) - \frac{\text{AiryBi}(x) x^3 3^{\frac{1}{6}} \text{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\text{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)) + \left(-\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right) \text{AiryAi}(x) - \frac{\text{AiryBi}(x) x^3 3^{\frac{1}{6}} \text{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\text{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3} \right) \\ &= -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right) \text{AiryAi}(x) - \frac{\text{AiryBi}(x) x^3 3^{\frac{1}{6}} \text{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\text{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right) \text{AiryAi}(x) - \frac{\text{AiryBi}(x) x^3 3^{\frac{1}{6}} \text{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\text{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x) \quad (1)$$

Verification of solutions

$$y = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right) \text{AiryAi}(x) - \frac{\text{AiryBi}(x) x^3 3^{\frac{1}{6}} \text{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\text{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Verified OK.

2.31.2 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' - yx^3 = x^3 \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{2i}{3} \\ n &= \frac{1}{3} \\ \gamma &= \frac{3}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) x^3}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) x \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(\alpha) \alpha \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) x^3}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) x \pi dx$$

Hence

$$u_2 = -\frac{x^3 3^{\frac{1}{6}} \text{hypergeom}\left(\left[1\right], \left[\frac{4}{3}, 2\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI}\left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3}\right)}{3}$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right)$$

$$u_2 = -\frac{x^3 3^{\frac{1}{6}} \text{hypergeom}\left(\left[1\right], \left[\frac{4}{3}, 2\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI}\left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3}\right)}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right) \text{AiryAi}(x)$$

$$+ \left(-\frac{x^3 3^{\frac{1}{6}} \text{hypergeom}\left(\left[1\right], \left[\frac{4}{3}, 2\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI}\left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3}\right)}{3} \right) \text{AiryBi}(x)$$

Which simplifies to

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha d\alpha \right) \text{AiryAi}(x)$$

$$- \frac{\text{AiryBi}(x) x^3 3^{\frac{1}{6}} \text{hypergeom}\left(\left[1\right], \left[\frac{4}{3}, 2\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)}{6}$$

$$+ \frac{\text{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \text{BesselI}\left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3}\right)}{3}$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \right) \\
 &\quad + \left(-\pi \left(\int_0^x \operatorname{AiryBi}(\alpha) \alpha d\alpha \right) \operatorname{AiryAi}(x) \right. \\
 &\quad \left. - \frac{\operatorname{AiryBi}(x) x^3 3^{\frac{1}{6}} \operatorname{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)}{6} \right. \\
 &\quad \left. + \frac{\operatorname{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}} \right)^{\frac{1}{3}} \pi \operatorname{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3} \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \\
 &\quad - \pi \left(\int_0^x \operatorname{AiryBi}(\alpha) \alpha d\alpha \right) \operatorname{AiryAi}(x) \\
 &\quad - \frac{\operatorname{AiryBi}(x) x^3 3^{\frac{1}{6}} \operatorname{hypergeom} \left([1], \left[\frac{4}{3}, 2 \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)}{6} \\
 &\quad + \frac{\operatorname{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}} \right)^{\frac{1}{3}} \pi \operatorname{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right)}{3}
 \end{aligned} \tag{1}$$

Verification of solutions

$$y = c_1 \sqrt{x} \operatorname{BesselJ}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + c_2 \sqrt{x} \operatorname{BesselY}\left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) - \pi \left(\int_0^x \operatorname{AiryBi}(\alpha) \alpha d\alpha \right) \operatorname{AiryAi}(x) - \frac{\operatorname{AiryBi}(x) x^3 3^{\frac{1}{6}} \operatorname{hypergeom}\left([1], \left[\frac{4}{3}, 2\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)}{6} + \frac{\operatorname{AiryBi}(x) \sqrt{x} \left(x^{\frac{3}{2}}\right)^{\frac{1}{3}} \pi \operatorname{BesselI}\left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3}\right)}{3}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x), x$2) - x*y(x) - x = 0, y(x), singsol=all)
```

$$y(x) = \operatorname{AiryAi}(x) c_2 + \operatorname{AiryBi}(x) c_1 - 1$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 28

```
DSolve[y''[x]-x*y[x]-x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \pi \text{AiryAiPrime}(x) \text{AiryBi}(x) + c_2 \text{AiryBi}(x) \\ + \text{AiryAi}(x)(-\pi \text{AiryBiPrime}(x) + c_1)$$

2.32 problem 31

2.32.1 Solving as second order airy ode 989

2.32.2 Solving as second order besseel ode ode 993

Internal problem ID [7168]

Internal file name [OUTPUT/6154_Sunday_June_05_2022_04_25_43_PM_21708165/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 31.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_besseel_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - yx = x^2$$

2.32.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = 0$$

$$c = -1$$

$$F = x^2$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^2 \text{AiryBi}(x)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) x^2 \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) x^2}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) x^2 \pi dx$$

Hence

u_2

$$= \frac{\pi \left(-3 \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) x^{\frac{3}{2}} - \text{BesselI} \left(\frac{1}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) \right) \Gamma \left(\frac{2}{3} \right) + x^{\frac{7}{2}} 3^{\frac{1}{3}} \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right)$$

$$u_2 = \frac{\pi \left(-3 \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) x^{\frac{3}{2}} - \text{BesselI} \left(\frac{1}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) \right) \Gamma \left(\frac{2}{3} \right) + x^{\frac{7}{2}} 3^{\frac{1}{3}} \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right) \text{AiryAi}(x)$$

$$+ \frac{\pi \left(-3 \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) x^{\frac{3}{2}} - \text{BesselI} \left(\frac{1}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) \right) \Gamma \left(\frac{2}{3} \right) + x^{\frac{7}{2}} 3^{\frac{1}{3}} \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

Which simplifies to

$$y_p(x) = \frac{\pi \left(\text{AiryBi}(x) \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3 \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right) \sqrt{x} \text{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \text{AiryAi}(x) \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x))$$

$$+ \left(\frac{\pi \left(\text{AiryBi}(x) \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3 \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right) \sqrt{x} \text{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \text{AiryAi}(x) \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)} \right)$$

$$= \frac{\pi \left(\text{AiryBi}(x) \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3 \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right) \sqrt{x} \text{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \text{AiryAi}(x) \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

$$+ c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Summary

The solution(s) found are the following

$$y = \frac{\pi \left(\text{AiryBi}(x) \text{ hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3 \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right) \sqrt{x} \text{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \text{AiryAi}(x) \right) \right)}{9\sqrt{x} \Gamma\left(\frac{2}{3}\right)} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x) \quad (1)$$

Verification of solutions

$$y = \frac{\pi \left(\text{AiryBi}(x) \text{ hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3 \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right) \sqrt{x} \text{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \text{AiryAi}(x) \right) \right)}{9\sqrt{x} \Gamma\left(\frac{2}{3}\right)} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Verified OK.

2.32.2 Solving as second order bessel ode

Writing the ode as

$$x^2 y'' - y x^3 = x^4 \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + x y' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) x y' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{2i}{3} \\ n &= \frac{1}{3} \\ \gamma &= \frac{3}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^4 \text{AiryBi}(x)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) x^2 \pi dx$$

Hence

$$u_1 = - \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 \pi d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) x^4}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) x^2 \pi dx$$

Hence

$$u_2 = \frac{\pi \left(-3 \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) x^{\frac{3}{2}} - \text{BesselI} \left(\frac{1}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) \right) \Gamma \left(\frac{2}{3} \right) + x^{\frac{7}{2}} 3^{\frac{1}{3}} \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right)$$

$$u_2 = \frac{\pi \left(-3 \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) x^{\frac{3}{2}} - \text{BesselI} \left(\frac{1}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) \right) \Gamma \left(\frac{2}{3} \right) + x^{\frac{7}{2}} 3^{\frac{1}{3}} \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\pi \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right) \text{AiryAi}(x) + \frac{\pi \left(-3 \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) x^{\frac{3}{2}} - \text{BesselI} \left(\frac{1}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) \right) \Gamma \left(\frac{2}{3} \right) + x^{\frac{7}{2}} 3^{\frac{1}{3}} \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

Which simplifies to

$$y_p(x) = \frac{\pi \left(\text{AiryBi}(x) \text{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3\sqrt{x} \left(\int_0^x \text{AiryBi}(\alpha) \alpha^2 d\alpha \right) \text{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \text{Ai} \right) \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \right) \\
 &\quad + \frac{\pi \left(\operatorname{AiryBi}(x) \operatorname{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3\sqrt{x} \left(\int_0^x \operatorname{AiryBi}(\alpha) \alpha^2 d\alpha \right) \operatorname{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \tag{1} \\
 &\quad + \frac{\pi \left(\operatorname{AiryBi}(x) \operatorname{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3\sqrt{x} \left(\int_0^x \operatorname{AiryBi}(\alpha) \alpha^2 d\alpha \right) \operatorname{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \\
 &\quad + \frac{\pi \left(\operatorname{AiryBi}(x) \operatorname{hypergeom} \left([1], \left[\frac{2}{3}, 2 \right], \frac{x^3}{9} \right) x^{\frac{7}{2}} 3^{\frac{1}{3}} - 3 \left(3\sqrt{x} \left(\int_0^x \operatorname{AiryBi}(\alpha) \alpha^2 d\alpha \right) \operatorname{AiryAi}(x) + \left(x^{\frac{3}{2}} \right)^{\frac{2}{3}} \right)}{9\sqrt{x} \Gamma \left(\frac{2}{3} \right)}
 \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x$2)-x*y(x)-x^2=0,y(x), singsol=all)
```

$$y(x) = \text{AiryAi}(x) c_2 + \text{AiryBi}(x) c_1 - x$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 30

```
DSolve[y''[x]-x*y[x]-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \pi x \text{AiryAiPrime}(x) \text{AiryBi}(x) + c_2 \text{AiryBi}(x) \\ + \text{AiryAi}(x)(-\pi x \text{AiryBiPrime}(x) + c_1)$$

2.33 problem 32

2.33.1 Solving as second order airy ode 999

2.33.2 Solving as second order besseel ode ode 1003

Internal problem ID [7169]

Internal file name [OUTPUT/6155_Sunday_June_05_2022_04_25_45_PM_56980621/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 32.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_besseel_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - yx = x^3$$

2.33.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = 0$$

$$c = -1$$

$$F = x^3$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) x^3}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) x^3 \pi dx$$

Hence

$$u_1 = - \frac{6x^5 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) + 5 3^{\frac{5}{6}} x^4 \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{60 \Gamma\left(\frac{2}{3}\right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) x^3}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) x^3 \pi dx$$

Hence

$$u_2 = - \frac{\left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) - \frac{5 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6}\right) x^4}{10 \Gamma\left(\frac{2}{3}\right)}$$

Which simplifies to

$$u_1 = -\frac{\left(\Gamma\left(\frac{2}{3}\right)^2 x 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) + \frac{5 3^{\frac{5}{6}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6}\right) x^4}{10\Gamma\left(\frac{2}{3}\right)}$$

$$u_2 = -\frac{\left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) - \frac{5 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6}\right) x^4}{10\Gamma\left(\frac{2}{3}\right)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) =$$

$$\frac{\left(\Gamma\left(\frac{2}{3}\right)^2 x 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) + \frac{5 3^{\frac{5}{6}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6}\right) x^4 \text{AiryAi}(x)}{10\Gamma\left(\frac{2}{3}\right)}$$

$$-\frac{\left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) - \frac{5 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6}\right) x^4 \text{AiryBi}(x)}{10\Gamma\left(\frac{2}{3}\right)}$$

Which simplifies to

$$y_p(x) =$$

$$\frac{x^4 \left(-\frac{5\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)\right) \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right)}{6} + x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) - \text{AiryBi}(x)\right) \right)}{10\Gamma\left(\frac{2}{3}\right)}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x))$$

$$+ \left(\frac{x^4 \left(-\frac{5\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)\right) \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right)}{6} + x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) - \text{AiryBi}(x)\right) \right)}{10\Gamma\left(\frac{2}{3}\right)} \right)$$

$$=$$

$$\frac{x^4 \left(-\frac{5\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)\right) \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right)}{6} + x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) - \text{AiryBi}(x)\right) \right)}{10\Gamma\left(\frac{2}{3}\right)}$$

$$+ c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Summary

The solution(s) found are the following

$$y = \frac{x^4 \left(-\frac{5\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{4}{3} \right], \left[\frac{2}{3}, \frac{7}{3} \right], \frac{x^3}{9} \right)}{6} + x \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)^2 \left(\text{AiryAi}(x) \right)}{10\Gamma \left(\frac{2}{3} \right)} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x) \tag{1}$$

Verification of solutions

$$y = \frac{x^4 \left(-\frac{5\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom} \left(\left[\frac{4}{3} \right], \left[\frac{2}{3}, \frac{7}{3} \right], \frac{x^3}{9} \right)}{6} + x \text{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)^2 \left(\text{AiryAi}(x) \right)}{10\Gamma \left(\frac{2}{3} \right)} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Verified OK.

2.33.2 Solving as second order Bessel ODE

Writing the ODE as

$$x^2 y'' - y x^3 = x^5 \tag{1}$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ODE has the form

$$x^2 y'' + x y' + (-n^2 + x^2) y = 0 \tag{2}$$

The generalized form of Bessel ODE is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) x y' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{2i}{3} \\ n &= \frac{1}{3} \\ \gamma &= \frac{3}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^5 \text{AiryBi}(x)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) x^3 \pi dx$$

Hence

$$u_1 = - \frac{6x^5 \Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) + 5 \cdot 3^{\frac{5}{6}} x^4 \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{60 \Gamma\left(\frac{2}{3}\right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) x^5}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) x^3 \pi dx$$

Hence

$$u_2 = - \frac{\left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) - \frac{5 \cdot 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6} \right) x^4}{10 \Gamma\left(\frac{2}{3}\right)}$$

Which simplifies to

$$u_1 = - \frac{\left(\Gamma\left(\frac{2}{3}\right)^2 x 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) + \frac{5 \cdot 3^{\frac{5}{6}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6} \right) x^4}{10 \Gamma\left(\frac{2}{3}\right)}$$

$$u_2 = - \frac{\left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) - \frac{5 \cdot 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6} \right) x^4}{10 \Gamma\left(\frac{2}{3}\right)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\Gamma\left(\frac{2}{3}\right)^2 x 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) + \frac{5 \cdot 3^{\frac{5}{6}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6} \right) x^4 \text{AiryAi}(x)}{10 \Gamma\left(\frac{2}{3}\right)} - \frac{\left(3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) - \frac{5 \cdot 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \pi}{6} \right) x^4 \text{AiryBi}(x)}{10 \Gamma\left(\frac{2}{3}\right)}$$

Which simplifies to

$$y_p(x) = \frac{x^4 \left(- \frac{5 \pi (\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)) \text{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right)}{6} + x \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) \Gamma\left(\frac{2}{3}\right)^2 (\text{AiryAi}(x) - \text{AiryBi}(x)) \right)}{10 \Gamma\left(\frac{2}{3}\right)}$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \right) \\
 &\quad + \left(\frac{x^4 \left(-\frac{5\pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom} \left(\left[\frac{4}{3} \right], \left[\frac{2}{3}, \frac{7}{3} \right], \frac{x^3}{9} \right)}{6} + x \operatorname{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)^2 \left(\operatorname{AiryAi}(x) \right)}{10\Gamma \left(\frac{2}{3} \right)} \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \tag{1} \\
 &\quad + \frac{x^4 \left(-\frac{5\pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom} \left(\left[\frac{4}{3} \right], \left[\frac{2}{3}, \frac{7}{3} \right], \frac{x^3}{9} \right)}{6} + x \operatorname{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)^2 \left(\operatorname{AiryAi}(x) \right)}{10\Gamma \left(\frac{2}{3} \right)}
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \\
 &\quad + \frac{x^4 \left(-\frac{5\pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom} \left(\left[\frac{4}{3} \right], \left[\frac{2}{3}, \frac{7}{3} \right], \frac{x^3}{9} \right)}{6} + x \operatorname{hypergeom} \left(\left[\frac{5}{3} \right], \left[\frac{4}{3}, \frac{8}{3} \right], \frac{x^3}{9} \right) \Gamma \left(\frac{2}{3} \right)^2 \left(\operatorname{AiryAi}(x) \right)}{10\Gamma \left(\frac{2}{3} \right)}
 \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 87

```
dsolve(diff(y(x),x$2)-x*y(x)-x^3=0,y(x), singsol=all)
```

$y(x)$

$$= \frac{5x^4\pi \operatorname{hypergeom}\left(\left[\frac{4}{3}\right], \left[\frac{2}{3}, \frac{7}{3}\right], \frac{x^3}{9}\right) \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x)\right) - 6\Gamma\left(\frac{2}{3}\right) \left(x^5 \operatorname{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right)\right)}{60\Gamma\left(\frac{2}{3}\right)}$$

✓ Solution by Mathematica

Time used: 0.093 (sec). Leaf size: 137

```
DSolve[y''[x]-x*y[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\pi x^5 \Gamma\left(\frac{5}{3}\right) (3 \operatorname{AiryAi}(x) + \sqrt{3} \operatorname{AiryBi}(x)) {}_1F_2\left(\frac{5}{3}; \frac{4}{3}, \frac{8}{3}; \frac{x^3}{9}\right)}{9 \cdot 3^{5/6} \Gamma\left(\frac{4}{3}\right) \Gamma\left(\frac{8}{3}\right)} \\ + \frac{\pi x^4 \Gamma\left(\frac{4}{3}\right) (\operatorname{AiryBi}(x) - \sqrt{3} \operatorname{AiryAi}(x)) {}_1F_2\left(\frac{4}{3}; \frac{2}{3}, \frac{7}{3}; \frac{x^3}{9}\right)}{3 \cdot 3^{2/3} \Gamma\left(\frac{2}{3}\right) \Gamma\left(\frac{7}{3}\right)} \\ + c_1 \operatorname{AiryAi}(x) + c_2 \operatorname{AiryBi}(x)$$

2.34 problem 33

- 2.34.1 Solving as second order airy ode 1010
- 2.34.2 Solving as second order bessel ode ode 1014

Internal problem ID [7170]

Internal file name [OUTPUT/6156_Sunday_June_05_2022_04_25_47_PM_37607482/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 33.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_bessel_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - yx = x^6 + x^3 - 42$$

2.34.1 Solving as second order airy ode

This is Airy ODE. It has the general form

$$ay'' + by' + cyx = F(x)$$

Where in this case

$$a = 1$$

$$b = 0$$

$$c = -1$$

$$F = x^6 + x^3 - 42$$

Therefore the solution to the homogeneous Airy ODE becomes

$$y = e^{-\frac{bx}{2a}} \left(c_1 \text{AiryAi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) + c_2 \text{AiryBi} \left(\frac{\left(-\frac{c}{a}\right)^{\frac{1}{3}} (4cxa + b^2)}{4ca} \right) \right)$$

Substituting the values for a, b, c gives

$$y = c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Since this is inhomogeneous Airy ODE, then we need to find the particular solution and add that to the homogeneous above. The particular solution is found using variation of parameters. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x))(\text{AiryBi}(1, x)) - (\text{AiryBi}(x))(\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) (x^6 + x^3 - 42)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) (x^6 + x^3 - 42) \pi dx$$

Hence

$$u_1 =$$

$$\frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) x^7 + \frac{16\pi \text{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right) 3^{\frac{5}{6}} x^6}{21} + \frac{8\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) x^5}{5}}{1}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) (x^6 + x^3 - 42)}{\frac{1}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) (x^6 + x^3 - 42) \pi dx$$

Hence

$$u_2 =$$

$$\frac{\left(\Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) 3^{\frac{1}{6}} x^7 - \frac{16\pi 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right) x^6}{21} + \frac{8 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) x^5}{5}}{1}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \operatorname{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) x^7 + \frac{16\pi \operatorname{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right) 3^{\frac{5}{6}} x^6}{21} + \frac{8\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \operatorname{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) x^5}{5} \right)}{\left(\Gamma\left(\frac{2}{3}\right)^2 \operatorname{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) 3^{\frac{1}{6}} x^7 - \frac{16\pi 3^{\frac{1}{3}} \operatorname{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right) x^6}{21} + \frac{8 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 \operatorname{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) x^5}{5} \right)}$$

Which simplifies to

$$y_p(x) = \left(-\frac{16x^6\pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\operatorname{AiryAi}(x) 3^{\frac{2}{3}} + \operatorname{AiryBi}(x) 3^{\frac{1}{6}} \right) \operatorname{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) \right)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \operatorname{AiryAi}(x) + c_2 \operatorname{AiryBi}(x)) \\ &+ \left(-\frac{16x^6\pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\operatorname{AiryAi}(x) 3^{\frac{2}{3}} + \operatorname{AiryBi}(x) 3^{\frac{1}{6}} \right) \operatorname{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) \right) \\ &= \left(-\frac{16x^6\pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\operatorname{AiryAi}(x) 3^{\frac{2}{3}} + \operatorname{AiryBi}(x) 3^{\frac{1}{6}} \right) \operatorname{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) \right) \\ &+ c_1 \operatorname{AiryAi}(x) + c_2 \operatorname{AiryBi}(x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{16x^6\pi \left(\text{AiryBi}(x)3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom}\left(\left[\frac{7}{3}, \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right)\right)}{21} + x^7\Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x)3^{\frac{2}{3}} + \text{AiryBi}(x)3^{\frac{1}{6}} \right) \right) \text{hyp} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x) \quad (1)$$

Verification of solutions

$$y = \left(-\frac{16x^6\pi \left(\text{AiryBi}(x)3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x) \right) \text{hypergeom}\left(\left[\frac{7}{3}, \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right)\right)}{21} + x^7\Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x)3^{\frac{2}{3}} + \text{AiryBi}(x)3^{\frac{1}{6}} \right) \right) \text{hyp} + c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

Verified OK.

2.34.2 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' - yx^3 = x^2(x^6 + x^3 - 42) \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{2i}{3} \\ n &= \frac{1}{3} \\ \gamma &= \frac{3}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \text{AiryAi}(x)$$

$$y_2 = \text{AiryBi}(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \frac{d}{dx}(\text{AiryAi}(x)) & \frac{d}{dx}(\text{AiryBi}(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \text{AiryAi}(x) & \text{AiryBi}(x) \\ \text{AiryAi}(1, x) & \text{AiryBi}(1, x) \end{vmatrix}$$

Therefore

$$W = (\text{AiryAi}(x)) (\text{AiryBi}(1, x)) - (\text{AiryBi}(x)) (\text{AiryAi}(1, x))$$

Which simplifies to

$$W = \text{AiryAi}(x) \text{AiryBi}(1, x) - \text{AiryBi}(x) \text{AiryAi}(1, x)$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\text{AiryBi}(x) x^2 (x^6 + x^3 - 42)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \text{AiryBi}(x) (x^6 + x^3 - 42) \pi dx$$

Hence

$$u_1 = \frac{x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) x^7 + \frac{16\pi \text{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right) 3^{\frac{5}{3}} x^6}{21} + \frac{8\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}\right], \frac{x^3}{9}\right) x^3}{5}}{\pi}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\text{AiryAi}(x) x^2 (x^6 + x^3 - 42)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \text{AiryAi}(x) (x^6 + x^3 - 42) \pi dx$$

Hence

$$u_2 = \left(\Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) 3^{\frac{1}{6}} x^7 - \frac{16\pi 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right) x^6}{21} + \frac{8 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) x^5}{5} \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x \left(\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) x^7 + \frac{16\pi \text{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right) 3^{\frac{5}{6}} x^6}{21} + \frac{8\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) x^5}{5} \right)$$

$$\left(\Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) 3^{\frac{1}{6}} x^7 - \frac{16\pi 3^{\frac{1}{3}} \text{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right) x^6}{21} + \frac{8 3^{\frac{1}{6}} \Gamma\left(\frac{2}{3}\right)^2 \text{hypergeom}\left(\left[\frac{5}{3}\right], \left[\frac{4}{3}, \frac{8}{3}\right], \frac{x^3}{9}\right) x^5}{5} \right)$$

Which simplifies to

$$y_p(x) = \left(-\frac{16x^6\pi \left(\text{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \text{AiryAi}(x)\right) \text{hypergeom}\left(\left[\frac{7}{3}\right], \left[\frac{2}{3}, \frac{10}{3}\right], \frac{x^3}{9}\right)}{21} + x^7 \Gamma\left(\frac{2}{3}\right)^2 \left(\text{AiryAi}(x) 3^{\frac{2}{3}} + \text{AiryBi}(x) 3^{\frac{1}{6}}\right) \text{hypergeom}\left(\left[\frac{8}{3}\right], \left[\frac{4}{3}, \frac{11}{3}\right], \frac{x^3}{9}\right) \right)$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \right) + \left(- \frac{16x^6 \pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) + x^7 \Gamma \left(\frac{2}{3} \right)^2 \left(\operatorname{AiryAi}(x) 3^{\frac{2}{3}} + \operatorname{AiryBi}(x) 3^{\frac{1}{6}} \right)}{21} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \quad (1)$$

$$\left(- \frac{16x^6 \pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) + x^7 \Gamma \left(\frac{2}{3} \right)^2 \left(\operatorname{AiryAi}(x) 3^{\frac{2}{3}} + \operatorname{AiryBi}(x) 3^{\frac{1}{6}} \right) \operatorname{hyp} \right)$$

Verification of solutions

$$y = c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)$$

$$\left(- \frac{16x^6 \pi \left(\operatorname{AiryBi}(x) 3^{\frac{1}{3}} - 3^{\frac{5}{6}} \operatorname{AiryAi}(x) \right) \operatorname{hypergeom} \left(\left[\frac{7}{3} \right], \left[\frac{2}{3}, \frac{10}{3} \right], \frac{x^3}{9} \right) + x^7 \Gamma \left(\frac{2}{3} \right)^2 \left(\operatorname{AiryAi}(x) 3^{\frac{2}{3}} + \operatorname{AiryBi}(x) 3^{\frac{1}{6}} \right) \operatorname{hyp} \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-x*y(x)-x^6-x^3+42=0,y(x), singsol=all)
```

$$y(x) = \text{AiryAi}(x) c_2 + \text{AiryBi}(x) c_1 - x^5 - 21x^2$$

✓ Solution by Mathematica

Time used: 1.142 (sec). Leaf size: 367

```
DSolve[y''[x]-x*y[x]-x^6-x^3+42==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x) \rightarrow$

$$\frac{-126\sqrt[3]{3}\pi x \Gamma\left(\frac{1}{3}\right) \left(\sqrt{3} \text{AiryAi}(x) - \text{AiryBi}(x)\right) {}_1F_2\left(\frac{1}{3}; \frac{2}{3}, \frac{4}{3}; \frac{x^3}{9}\right) + \frac{\sqrt[6]{3}\pi x^8 \Gamma\left(\frac{2}{3}\right) \Gamma\left(\frac{8}{3}\right) (3 \text{Ai}(x) - \text{Bi}(x))}{\Gamma\left(\frac{1}{3}\right) \Gamma\left(\frac{2}{3}\right) \Gamma\left(\frac{4}{3}\right) \Gamma\left(\frac{5}{3}\right) \Gamma\left(\frac{8}{3}\right)}}{\Gamma\left(\frac{1}{3}\right) \Gamma\left(\frac{2}{3}\right) \Gamma\left(\frac{4}{3}\right) \Gamma\left(\frac{5}{3}\right) \Gamma\left(\frac{8}{3}\right)}$$

2.35 problem 34

2.35.1 Solving as second order bessel ode ode 1020

Internal problem ID [7171]

Internal file name [OUTPUT/6157_Sunday_June_05_2022_04_25_48_PM_4074723/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 34.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - x^2y = x^2$$

2.35.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' - yx^4 = x^4 \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{i}{2} \\ n &= \frac{1}{4} \\ \gamma &= 2\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

$$y_2 = \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ \frac{d}{dx}\left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) & \frac{d}{dx}\left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ \frac{\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) & \frac{\text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \left(\frac{\text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}}\left(-\text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \\ &\quad - \left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \left(\frac{\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}}\left(-\text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \end{aligned}$$

Which simplifies to

$$W = -ix^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) \right)$$

Which simplifies to

$$W = \frac{4}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{\frac{5}{2}} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \pi}{4} dx$$

Hence

$$u_1 = - \frac{(-1)^{\frac{1}{8}} x^4 \text{hypergeom} \left([1], \left[\frac{5}{4}, 2 \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right)}{8} - \frac{\sqrt{2} (-1)^{\frac{7}{8}} x^3 \text{BesselI} \left(\frac{3}{4}, \frac{x^2}{2} \right) \pi}{4 (x^2)^{\frac{3}{4}}}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{\frac{9}{2}} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{\frac{5}{2}} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \pi}{4} dx$$

Hence

$$u_2 = \frac{(-1)^{\frac{1}{8}} x^4 \text{hypergeom} \left([1], \left[\frac{5}{4}, 2 \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right)}{8}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(- \frac{(-1)^{\frac{1}{8}} x^4 \text{hypergeom} \left([1], \left[\frac{5}{4}, 2 \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right)}{8} - \frac{\sqrt{2} (-1)^{\frac{7}{8}} x^3 \text{BesselI} \left(\frac{3}{4}, \frac{x^2}{2} \right) \pi}{4 (x^2)^{\frac{3}{4}}} \right) \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + \frac{(-1)^{\frac{1}{8}} x^{\frac{9}{2}} \text{hypergeom} \left([1], \left[\frac{5}{4}, 2 \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right) \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{8}$$

Which simplifies to

$$y_p(x) = \frac{x^{\frac{7}{2}}(-1)^{\frac{1}{8}} \left(\Gamma\left(\frac{3}{4}\right) x(x^2)^{\frac{3}{4}} \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) \text{hypergeom}\left([1], \left[\frac{5}{4}, 2\right], \frac{x^4}{16}\right) + 2\pi \text{Besse}}{8(x^2)^{\frac{3}{4}}}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + c_2 \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) + \left(-\frac{x^{\frac{7}{2}}(-1)^{\frac{1}{8}} \left(\Gamma\left(\frac{3}{4}\right) x(x^2)^{\frac{3}{4}} \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) \text{hypergeom}\left([1], \left[\frac{5}{4}, 2\right], \frac{x^4}{16}\right) + 2\pi \text{Besse}}{8(x^2)^{\frac{3}{4}}}$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + c_2 \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + \frac{x^{\frac{7}{2}}(-1)^{\frac{1}{8}} \left(\Gamma\left(\frac{3}{4}\right) x(x^2)^{\frac{3}{4}} \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) \text{hypergeom}\left([1], \left[\frac{5}{4}, 2\right], \frac{x^4}{16}\right) + 2\pi \text{Besse}}{8(x^2)^{\frac{3}{4}}} \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + c_2 \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + \frac{x^{\frac{7}{2}}(-1)^{\frac{1}{8}} \left(\Gamma\left(\frac{3}{4}\right) x(x^2)^{\frac{3}{4}} \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) \text{hypergeom}\left([1], \left[\frac{5}{4}, 2\right], \frac{x^4}{16}\right) + 2\pi \text{Besse}}{8(x^2)^{\frac{3}{4}}}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
            <- Bessel successful
        <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)-x^2*y(x)-x^2=0,y(x), singsol=all)
```

$$y(x) = \sqrt{x} \operatorname{BesselI}\left(\frac{1}{4}, \frac{x^2}{2}\right) c_2 + \sqrt{x} \operatorname{BesselK}\left(\frac{1}{4}, \frac{x^2}{2}\right) c_1 - 1$$

✓ Solution by Mathematica

Time used: 6.053 (sec). Leaf size: 213

```
DSolve[y''[x]-x^2*y[x]-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\begin{aligned} \rightarrow & \text{ParabolicCylinderD} \left(-\frac{1}{2}, \sqrt{2}x \right) \left(\int_1^x \frac{K[1]^2 \text{ParabolicCyl}}{\sqrt{2} (\text{HermiteH}(-\frac{1}{2}, K[1]) (i \text{HermiteH}(\frac{1}{2}, iK[1]) + 2 \text{HermiteH} \right.} \\ & \left. + c_1) \right) \\ & + \text{ParabolicCylinderD} \left(-\frac{1}{2}, i\sqrt{2}x \right) \left(\int_1^x \frac{K[2]^2 \text{ParabolicCyl}}{\sqrt{2} (\text{HermiteH}(-\frac{1}{2}, iK[2]) \text{HermiteH}(\frac{1}{2}, K[2]) + \text{HermiteH} \right. \\ & \left. + c_2) \right) \end{aligned}$$

2.36 problem 35

2.36.1 Solving as second order bessel ode ode 1027

Internal problem ID [7172]

Internal file name [OUTPUT/6158_Sunday_June_05_2022_04_25_52_PM_37822212/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 35.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - x^2y = x^3$$

2.36.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' - yx^4 = x^5 \tag{1}$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{i}{2} \\ n &= \frac{1}{4} \\ \gamma &= 2\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

$$y_2 = \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ \frac{d}{dx}\left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) & \frac{d}{dx}\left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ \frac{\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) & \frac{\text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \left(\frac{\text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}}\left(-\text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \\ &\quad - \left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \left(\frac{\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}}\left(-\text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \end{aligned}$$

Which simplifies to

$$W = -ix^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) \right)$$

Which simplifies to

$$W = \frac{4}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^{\frac{11}{2}} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{\frac{7}{2}} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \pi}{4} dx$$

Hence

$$u_1 = \frac{x \left((-1)^{\frac{3}{4}} x^3 \text{hypergeom} \left([1], \left[\frac{3}{4}, 2 \right], \frac{x^4}{16} \right) (x^2)^{\frac{1}{4}} + 2\Gamma\left(\frac{3}{4}\right) \left(\text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) x^2 - \text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) \right) \right) (-1)}{8 (x^2)^{\frac{1}{4}} \Gamma \left(\frac{3}{4} \right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{\frac{11}{2}} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{\frac{7}{2}} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \pi}{4} dx$$

Hence

$$u_2 = \frac{(-1)^{\frac{1}{8}} \pi x^3 \text{BesselI} \left(\frac{5}{4}, \frac{x^2}{2} \right)}{4 (x^2)^{\frac{1}{4}}}$$

Which simplifies to

$$u_1 = \frac{x \left((-1)^{\frac{3}{4}} x^3 \text{hypergeom} \left([1], \left[\frac{3}{4}, 2 \right], \frac{x^4}{16} \right) (x^2)^{\frac{1}{4}} + 2\Gamma\left(\frac{3}{4}\right) \left(\text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) x^2 - \text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) \right) \right) (-1)}{8 (x^2)^{\frac{1}{4}} \Gamma \left(\frac{3}{4} \right)}$$

$$u_2 = - \frac{(-1)^{\frac{1}{8}} \pi x \left(- \text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) x^2 + \text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) \right)}{4 (x^2)^{\frac{1}{4}}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x^{\frac{3}{2}} \left((-1)^{\frac{3}{4}} x^3 \operatorname{hypergeom} \left([1], \left[\frac{3}{4}, 2 \right], \frac{x^4}{16} \right) (x^2)^{\frac{1}{4}} + 2\Gamma\left(\frac{3}{4}\right) \left(\operatorname{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) x^2 - \operatorname{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) \right) \right)}{8 (x^2)^{\frac{1}{4}} \Gamma\left(\frac{3}{4}\right)} - \frac{(-1)^{\frac{1}{8}} \pi x^{\frac{3}{2}} \left(-\operatorname{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) x^2 + \operatorname{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) \right) \operatorname{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{4 (x^2)^{\frac{1}{4}}}$$

Which simplifies to

$$y_p(x) = \frac{(-1)^{\frac{1}{8}} \left(\operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \operatorname{hypergeom} \left([1], \left[\frac{3}{4}, 2 \right], \frac{x^4}{16} \right) (-1)^{\frac{3}{4}} (x^2)^{\frac{1}{4}} x^3 + 2\Gamma\left(\frac{3}{4}\right) \left(\operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) - \operatorname{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) \right)}{8 (x^2)^{\frac{1}{4}} \Gamma\left(\frac{3}{4}\right)}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) + \left(-\frac{(-1)^{\frac{1}{8}} \left(\operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \operatorname{hypergeom} \left([1], \left[\frac{3}{4}, 2 \right], \frac{x^4}{16} \right) (-1)^{\frac{3}{4}} (x^2)^{\frac{1}{4}} x^3 + 2\Gamma\left(\frac{3}{4}\right) \left(\operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) - \operatorname{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) \right)}{8 (x^2)^{\frac{1}{4}} \Gamma\left(\frac{3}{4}\right)} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + \frac{(-1)^{\frac{1}{8}} \left(\operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \operatorname{hypergeom} \left([1], \left[\frac{3}{4}, 2 \right], \frac{x^4}{16} \right) (-1)^{\frac{3}{4}} (x^2)^{\frac{1}{4}} x^3 + 2\Gamma\left(\frac{3}{4}\right) \left(\operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) - \operatorname{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) \right)}{8 (x^2)^{\frac{1}{4}} \Gamma\left(\frac{3}{4}\right)} \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + \frac{(-1)^{\frac{1}{8}} \left(\operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \operatorname{hypergeom} \left([1], \left[\frac{3}{4}, 2 \right], \frac{x^4}{16} \right) (-1)^{\frac{3}{4}} (x^2)^{\frac{1}{4}} x^3 + 2\Gamma\left(\frac{3}{4}\right) \left(\operatorname{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) - \operatorname{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) \right)}{8 (x^2)^{\frac{1}{4}} \Gamma\left(\frac{3}{4}\right)}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    trying a symmetry of the form [xi=0, eta=F(x)]  
    checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
    -> Bessel  
        <- Bessel successful  
    <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)-x^2*y(x)-x^3=0,y(x), singsol=all)
```

$$y(x) = \sqrt{x} \operatorname{BesselI}\left(\frac{1}{4}, \frac{x^2}{2}\right) c_2 + \sqrt{x} \operatorname{BesselK}\left(\frac{1}{4}, \frac{x^2}{2}\right) c_1 - x$$

✓ Solution by Mathematica

Time used: 4.871 (sec). Leaf size: 213

```
DSolve[y''[x]-x^2*y[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\begin{aligned} \rightarrow & \text{ParabolicCylinderD} \left(-\frac{1}{2}, \sqrt{2}x \right) \left(\int_1^x \frac{K[1]^3 \text{ParabolicCyl}}{\sqrt{2} (\text{HermiteH}(-\frac{1}{2}, K[1]) (i \text{HermiteH}(\frac{1}{2}, iK[1]) + 2 \text{HermiteH} \right.} \\ & \left. + c_1) \right) \\ & + \text{ParabolicCylinderD} \left(-\frac{1}{2}, i\sqrt{2}x \right) \left(\int_1^x \frac{K[2]^3 \text{ParabolicCyl}}{\sqrt{2} (\text{HermiteH}(-\frac{1}{2}, iK[2]) \text{HermiteH}(\frac{1}{2}, K[2]) + \text{HermiteH} \right. \\ & \left. + c_2) \right) \end{aligned}$$

2.37 problem 36

2.37.1 Solving as second order bessel ode ode 1034

Internal problem ID [7173]

Internal file name [OUTPUT/6159_Sunday_June_05_2022_04_25_55_PM_90936483/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 36.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - x^2y = x^4$$

2.37.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' - yx^4 = x^6 \tag{1}$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{i}{2} \\ n &= \frac{1}{4} \\ \gamma &= 2\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

$$y_2 = \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ \frac{d}{dx}\left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) & \frac{d}{dx}\left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ \frac{\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) & \frac{\text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \left(\frac{\text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}} \left(-\text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \\ &\quad - \left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \left(\frac{\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}} \left(-\text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \end{aligned}$$

Which simplifies to

$$W = -ix^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right)\right)$$

Which simplifies to

$$W = \frac{4}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^{\frac{13}{2}} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \pi}{4} dx$$

Hence

$$u_1 = \frac{(-1)^{\frac{1}{8}} \left(5x^6 \Gamma \left(\frac{3}{4} \right)^2 \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right) + 6x^5 (-1)^{\frac{3}{4}} \text{hypergeom} \left(\left[\frac{5}{4} \right], \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{16} \right) \pi \right)}{60 \Gamma \left(\frac{3}{4} \right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{\frac{13}{2}} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{\frac{9}{2}} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \pi}{4} dx$$

Hence

$$u_2 = \frac{(-1)^{\frac{1}{8}} x^6 \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right)}{12}$$

Which simplifies to

$$u_1 = - \frac{x^5 \left((-1)^{\frac{3}{4}} \text{hypergeom} \left(\left[\frac{5}{4} \right], \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{16} \right) \pi + \frac{5 \Gamma \left(\frac{3}{4} \right)^2 x \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right)}{6} \right) (-1)^{\frac{1}{8}}}{10 \Gamma \left(\frac{3}{4} \right)}$$

$$u_2 = \frac{(-1)^{\frac{1}{8}} x^6 \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right)}{12}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x^{\frac{11}{2}} \left((-1)^{\frac{3}{4}} \text{hypergeom} \left(\left[\frac{5}{4} \right], \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{16} \right) \pi + \frac{5\Gamma(\frac{3}{4})^2 x \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right)}{6} \right) (-1)^{\frac{1}{8}} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{10\Gamma \left(\frac{3}{4} \right)} + \frac{(-1)^{\frac{1}{8}} x^{\frac{13}{2}} \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right) \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{12}$$

Which simplifies to

$$y_p(x) = \frac{\left(\frac{5x\Gamma(\frac{3}{4})^2 \left(\text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) - \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right)}{6} + \pi \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \text{hypergeom} \left(\left[\frac{5}{4} \right], \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{16} \right) \right)}{10\Gamma \left(\frac{3}{4} \right)}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) + \frac{\left(\frac{5x\Gamma(\frac{3}{4})^2 \left(\text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) - \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right)}{6} + \pi \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \text{hypergeom} \left(\left[\frac{5}{4} \right], \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{16} \right) \right)}{10\Gamma \left(\frac{3}{4} \right)}$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + \frac{\left(\frac{5x\Gamma(\frac{3}{4})^2 \left(\text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) - \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \right) \text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right)}{6} + \pi \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) \text{hypergeom} \left(\left[\frac{5}{4} \right], \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{16} \right) \right)}{10\Gamma \left(\frac{3}{4} \right)} \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \operatorname{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + c_2 \sqrt{x} \operatorname{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ - \frac{\left(\frac{5x\Gamma\left(\frac{3}{4}\right)^2 \left(\operatorname{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \operatorname{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \operatorname{hypergeom}\left(\left[\frac{3}{2}\right], \left[\frac{5}{4}, \frac{5}{2}\right], \frac{x^4}{16}\right)\right)}{6} + \pi \operatorname{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \operatorname{hypergeom}\left(\left[\frac{5}{4}\right], \left[\frac{3}{4}, \frac{9}{4}\right]\right)}{10\Gamma\left(\frac{3}{4}\right)}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 124

```
dsolve(diff(y(x), x$2) - x^2*y(x) - x^4=0, y(x), singsol=all)
```

$$y(x) = \frac{\left(-\frac{6x^5\pi^2 \operatorname{csgn}(x) \operatorname{hypergeom}\left(\left[\frac{5}{4}\right], \left[\frac{3}{4}, \frac{5}{2}\right], \frac{x^4}{16}\right) \operatorname{BesselI}\left(\frac{1}{4}, \frac{x^2}{2}\right)\right)}{5} + \Gamma\left(\frac{3}{4}\right) \left(2x^6\Gamma\left(\frac{3}{4}\right) \operatorname{BesselK}\left(\frac{1}{4}, \frac{x^2}{2}\right) \operatorname{hypergeom}\left(\left[\frac{3}{2}\right], \left[\frac{5}{4}, \frac{9}{4}\right]\right)\right)$$

✓ Solution by Mathematica

Time used: 3.699 (sec). Leaf size: 213

```
DSolve[y''[x]-x^2*y[x]-x^4==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\begin{aligned} \rightarrow & \text{ParabolicCylinderD} \left(-\frac{1}{2}, \sqrt{2}x \right) \left(\int_1^x \frac{K[1]^4 \text{ParabolicCyl}}{\sqrt{2} (\text{HermiteH}(-\frac{1}{2}, K[1]) (i \text{HermiteH}(\frac{1}{2}, iK[1]) + 2 \text{HermiteH} \right.} \\ & \left. + c_1) \right) \\ & + \text{ParabolicCylinderD} \left(-\frac{1}{2}, i\sqrt{2}x \right) \left(\int_1^x \frac{K[2]^4 \text{ParabolicCyl}}{\sqrt{2} (\text{HermiteH}(-\frac{1}{2}, iK[2]) \text{HermiteH}(\frac{1}{2}, K[2]) + \text{HermiteH} \right. \\ & \left. + c_2) \right) \end{aligned}$$

2.38 problem 37

2.38.1 Solving as second order bessel ode ode 1041

Internal problem ID [7174]

Internal file name [OUTPUT/6160_Sunday_June_05_2022_04_25_58_PM_28309459/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 37.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - x^2y = x^4 - 2$$

2.38.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' - yx^4 = x^2(x^4 - 2) \tag{1}$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{i}{2} \\ n &= \frac{1}{4} \\ \gamma &= 2\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

$$y_2 = \sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ \frac{d}{dx}\left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) & \frac{d}{dx}\left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \\ \frac{\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) & \frac{\text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \left(\frac{\text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}}\left(-\text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \\ &\quad - \left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right)\right) \left(\frac{\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}}\left(-\text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \frac{i \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right)}{2x^2}\right) \end{aligned}$$

Which simplifies to

$$W = -ix^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \text{BesselY}\left(\frac{5}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \text{BesselJ}\left(\frac{5}{4}, \frac{ix^2}{2}\right) \right)$$

Which simplifies to

$$W = \frac{4}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^{\frac{5}{2}} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) (x^4 - 2)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right) (x^4 - 2) \pi}{4} dx$$

Hence

$$u_1 = \frac{\left(\text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right)^2 x^5 + \frac{6 \text{hypergeom} \left(\left[\frac{5}{4}, \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{16} \right) (-1)^{\frac{3}{4}} \pi x^4}{5} - 6 \text{hypergeom} \left(\left[\frac{1}{2} \right], \left[\frac{5}{4}, \frac{3}{2} \right] \right) \Gamma \left(\frac{3}{4} \right) \right)}{12 \Gamma \left(\frac{3}{4} \right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{\frac{5}{2}} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) (x^4 - 2)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{\sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) (x^4 - 2) \pi}{4} dx$$

Hence

$$u_2 = \frac{(-1)^{\frac{1}{8}} x^2 \left(\text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right) x^4 - 6 \text{hypergeom} \left(\left[\frac{1}{2} \right], \left[\frac{5}{4}, \frac{3}{2} \right], \frac{x^4}{16} \right) \right) \Gamma \left(\frac{3}{4} \right)}{12}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right) \Gamma \left(\frac{3}{4} \right)^2 x^5 + \frac{6 \text{hypergeom} \left(\left[\frac{5}{4}, \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{16} \right) (-1)^{\frac{3}{4}} \pi x^4}{5} - 6 \text{hypergeom} \left(\left[\frac{1}{2} \right], \left[\frac{5}{4}, \frac{3}{2} \right] \right) \Gamma \left(\frac{3}{4} \right) \right)}{12 \Gamma \left(\frac{3}{4} \right)} + \frac{(-1)^{\frac{1}{8}} x^{\frac{5}{2}} \left(\text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{16} \right) x^4 - 6 \text{hypergeom} \left(\left[\frac{1}{2} \right], \left[\frac{5}{4}, \frac{3}{2} \right], \frac{x^4}{16} \right) \right) \Gamma \left(\frac{3}{4} \right) \text{BesselY} \left(\frac{1}{4}, \frac{ix^2}{2} \right)}{12}$$

Which simplifies to

$$y_p(x) = \frac{(-1)^{\frac{1}{8}} x^{\frac{3}{2}} \left(-6x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) \text{hypergeom}\left(\left[\frac{1}{2}\right], \left[\frac{5}{4}, \frac{3}{2}\right], \frac{x^4}{16}\right) + x^5\Gamma\left(\frac{3}{4}\right)^2}{1}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + c_2\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) + \frac{(-1)^{\frac{1}{8}} x^{\frac{3}{2}} \left(-6x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) \text{hypergeom}\left(\left[\frac{1}{2}\right], \left[\frac{5}{4}, \frac{3}{2}\right], \frac{x^4}{16}\right) + x^5\Gamma\left(\frac{3}{4}\right)^2}{1}$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + c_2\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + \frac{(-1)^{\frac{1}{8}} x^{\frac{3}{2}} \left(-6x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) \text{hypergeom}\left(\left[\frac{1}{2}\right], \left[\frac{5}{4}, \frac{3}{2}\right], \frac{x^4}{16}\right) + x^5\Gamma\left(\frac{3}{4}\right)^2}{1} \quad (1)$$

Verification of solutions

$$y = c_1\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + c_2\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) + \frac{(-1)^{\frac{1}{8}} x^{\frac{3}{2}} \left(-6x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{ix^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{ix^2}{2}\right) \right) \text{hypergeom}\left(\left[\frac{1}{2}\right], \left[\frac{5}{4}, \frac{3}{2}\right], \frac{x^4}{16}\right) + x^5\Gamma\left(\frac{3}{4}\right)^2}{1}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    trying a symmetry of the form [xi=0, eta=F(x)]  
    checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
    -> Bessel  
        <- Bessel successful  
    <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)-x^2*y(x)-x^4+2=0,y(x), singsol=all)
```

$$y(x) = \sqrt{x} \operatorname{BesselI}\left(\frac{1}{4}, \frac{x^2}{2}\right) c_2 + \sqrt{x} \operatorname{BesselK}\left(\frac{1}{4}, \frac{x^2}{2}\right) c_1 - x^2$$

✓ Solution by Mathematica

Time used: 4.998 (sec). Leaf size: 217

`DSolve[y''[x]-x^2*y[x]-x^4+2==0,y[x],x,IncludeSingularSolutions -> True]`

$$\begin{aligned}
 y(x) \rightarrow & \text{ParabolicCylinderD}\left(-\frac{1}{2}, \sqrt{2}x\right) \left(\int_1^x \frac{(K[1]^4 - 2) \text{ParabolicCylinderD}\left(-\frac{1}{2}, i\sqrt{2}K[1]\right)}{\sqrt{2} \left(\text{HermiteH}\left(-\frac{1}{2}, iK[1]\right) \text{HermiteH}\left(\frac{1}{2}, K[1]\right) + \text{HermiteH}\left(-\frac{1}{2}, K[1]\right) \left(-i \text{HermiteH}\left(\frac{1}{2}, iK[1]\right) - 2\right) + c_1} \right) \\
 & + \text{ParabolicCylinderD}\left(-\frac{1}{2}, i\sqrt{2}x\right) \left(\int_1^x \frac{(K[2]^4 - 2) \text{ParabolicCylinderD}\left(-\frac{1}{2}, i\sqrt{2}K[2]\right)}{\sqrt{2} \left(\text{HermiteH}\left(-\frac{1}{2}, iK[2]\right) \text{HermiteH}\left(\frac{1}{2}, K[2]\right) + \text{HermiteH}\left(-\frac{1}{2}, K[2]\right) \left(-i \text{HermiteH}\left(\frac{1}{2}, iK[2]\right) - 2\right) + c_2} \right)
 \end{aligned}$$

2.39 problem 38

2.39.1 Solving as second order bessel ode ode 1048

Internal problem ID [7175]

Internal file name [OUTPUT/6161_Sunday_June_05_2022_04_26_03_PM_89021356/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 38.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$y'' - 2x^2y = x^4 - 1$$

2.39.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' - 2yx^4 = x^2(x^4 - 1) \tag{1}$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{i\sqrt{2}}{2} \\ n &= \frac{1}{4} \\ \gamma &= 2\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1\sqrt{x} \text{ BesselJ} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right) + c_2\sqrt{x} \text{ BesselY} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1\sqrt{x} \text{ BesselJ} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right) + c_2\sqrt{x} \text{ BesselY} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= \sqrt{x} \text{ BesselJ} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right) \\ y_2 &= \sqrt{x} \text{ BesselY} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right)\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \\ \frac{d}{dx}\left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right) & \frac{d}{dx}\left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \\ \frac{\text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselJ}\left(\frac{5}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \frac{i\sqrt{2} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{4x^2}\right) & \sqrt{2} \frac{\text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2\sqrt{x}} + ix^{\frac{3}{2}}\left(-\text{BesselY}\left(\frac{5}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \frac{i\sqrt{2} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{4x^2}\right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right) \left(\frac{\text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}} \left(-\text{BesselY}\left(\frac{5}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \frac{i\sqrt{2} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{4x^2}\right) \sqrt{2} \\ &\quad - \left(\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right) \left(\frac{\text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{2\sqrt{x}}\right) \\ &\quad + ix^{\frac{3}{2}} \left(-\text{BesselJ}\left(\frac{5}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \frac{i\sqrt{2} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right)}{4x^2}\right) \sqrt{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} W &= -ix^2\sqrt{2} \left(\text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \text{BesselY}\left(\frac{5}{4}, \frac{i\sqrt{2}x^2}{2}\right)\right) \\ &\quad - \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \text{BesselJ}\left(\frac{5}{4}, \frac{i\sqrt{2}x^2}{2}\right) \end{aligned}$$

Which simplifies to

$$W = \frac{4}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^{\frac{5}{2}} \text{BesselY} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right) (x^4 - 1)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sqrt{x} \text{BesselY} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right) (x^4 - 1) \pi}{4} dx$$

Hence

$$u_1 = \frac{\left(\frac{3\pi(-1)^{\frac{3}{4}} 2^{\frac{3}{4}} \text{hypergeom} \left(\left[\frac{5}{4}, \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{8} \right) x^4}{5} - 3\pi(-1)^{\frac{3}{4}} 2^{\frac{3}{4}} \text{hypergeom} \left(\left[\frac{1}{4} \right], \left[\frac{3}{4}, \frac{5}{4} \right], \frac{x^4}{8} \right) + x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{8} \right) x^4 - 3 \text{hypergeom} \left(\left[\frac{1}{2} \right], \left[\frac{5}{4}, \frac{3}{2} \right], \frac{x^4}{8} \right) \right) \Gamma\left(\frac{3}{4}\right)}{12\Gamma\left(\frac{3}{4}\right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{\frac{5}{2}} \text{BesselJ} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right) (x^4 - 1)}{\frac{4x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{\sqrt{x} \text{BesselJ} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right) (x^4 - 1) \pi}{4} dx$$

Hence

$$u_2 = \frac{2^{\frac{1}{8}}(-1)^{\frac{1}{8}} x^2 \left(\text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{8} \right) x^4 - 3 \text{hypergeom} \left(\left[\frac{1}{2} \right], \left[\frac{5}{4}, \frac{3}{2} \right], \frac{x^4}{8} \right) \right) \Gamma\left(\frac{3}{4}\right)}{12}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\frac{3\pi(-1)^{\frac{3}{4}} 2^{\frac{3}{4}} \text{hypergeom} \left(\left[\frac{5}{4}, \left[\frac{3}{4}, \frac{9}{4} \right], \frac{x^4}{8} \right) x^4}{5} - 3\pi(-1)^{\frac{3}{4}} 2^{\frac{3}{4}} \text{hypergeom} \left(\left[\frac{1}{4} \right], \left[\frac{3}{4}, \frac{5}{4} \right], \frac{x^4}{8} \right) + x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{8} \right) x^4 - 3 \text{hypergeom} \left(\left[\frac{1}{2} \right], \left[\frac{5}{4}, \frac{3}{2} \right], \frac{x^4}{8} \right) \right) \Gamma\left(\frac{3}{4}\right)}{12\Gamma\left(\frac{3}{4}\right)} + \frac{2^{\frac{1}{8}}(-1)^{\frac{1}{8}} x^{\frac{5}{2}} \left(\text{hypergeom} \left(\left[\frac{3}{2} \right], \left[\frac{5}{4}, \frac{5}{2} \right], \frac{x^4}{8} \right) x^4 - 3 \text{hypergeom} \left(\left[\frac{1}{2} \right], \left[\frac{5}{4}, \frac{3}{2} \right], \frac{x^4}{8} \right) \right) \Gamma\left(\frac{3}{4}\right) \text{BesselY} \left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2} \right)}{12}$$

Which simplifies to

$$y_p(x) = \frac{x^{\frac{3}{2}} 2^{\frac{1}{8}} (-1)^{\frac{1}{8}} \left(-3x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right) \text{hypergeom}\left(\left[\frac{1}{2}\right], \left[\frac{5}{4}, \frac{3}{2}\right], \frac{x^4}{8}\right) + x}{\dots}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + c_2\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right) + \left(\frac{x^{\frac{3}{2}} 2^{\frac{1}{8}} (-1)^{\frac{1}{8}} \left(-3x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right) \text{hypergeom}\left(\left[\frac{1}{2}\right], \left[\frac{5}{4}, \frac{3}{2}\right], \frac{x^4}{8}\right) + x}{\dots} \right)$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + c_2\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \frac{x^{\frac{3}{2}} 2^{\frac{1}{8}} (-1)^{\frac{1}{8}} \left(-3x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right) \text{hypergeom}\left(\left[\frac{1}{2}\right], \left[\frac{5}{4}, \frac{3}{2}\right], \frac{x^4}{8}\right) + x}{\dots} \quad (1)$$

Verification of solutions

$$y = c_1\sqrt{x} \text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + c_2\sqrt{x} \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) + \frac{x^{\frac{3}{2}} 2^{\frac{1}{8}} (-1)^{\frac{1}{8}} \left(-3x\Gamma\left(\frac{3}{4}\right)^2 \left(\text{BesselJ}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) - \text{BesselY}\left(\frac{1}{4}, \frac{i\sqrt{2}x^2}{2}\right) \right) \text{hypergeom}\left(\left[\frac{1}{2}\right], \left[\frac{5}{4}, \frac{3}{2}\right], \frac{x^4}{8}\right) + x}{\dots}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    trying a symmetry of the form [xi=0, eta=F(x)]  
    checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
    -> Bessel  
        <- Bessel successful  
    <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(diff(y(x), x$2) - 2*x^2*y(x) - x^4 + 1 = 0, y(x), singsol=all)
```

$$y(x) = \sqrt{x} \operatorname{BesselI}\left(\frac{1}{4}, \frac{\sqrt{2}x^2}{2}\right) c_2 + \sqrt{x} \operatorname{BesselK}\left(\frac{1}{4}, \frac{\sqrt{2}x^2}{2}\right) c_1 - \frac{x^2}{2}$$

✓ Solution by Mathematica

Time used: 3.94 (sec). Leaf size: 288

```
DSolve[y''[x] - 2*x^2*y[x] - x^4 + 1 == 0, y[x], x, IncludeSingularSolutions -> True]
```

$y(x)$

→ $\operatorname{ParabolicCylinderD}\left(-\frac{1}{2}, 2^{3/4}x\right) \left(\int_1^x \frac{1}{i^{23/4} \operatorname{HermiteH}\left(-\frac{1}{2}, \sqrt[4]{2}K[1]\right) \operatorname{HermiteH}\left(\frac{1}{2}, i\sqrt[4]{2}K[1]\right)} dx \right) + \operatorname{ParabolicCylinderD}\left(\frac{1}{2}, 2^{3/4}x\right) \left(\int_1^x \frac{1}{i^{23/4} \operatorname{HermiteH}\left(-\frac{1}{2}, \sqrt[4]{2}K[1]\right) \operatorname{HermiteH}\left(\frac{1}{2}, i\sqrt[4]{2}K[1]\right)} dx \right) + \dots$

2.40 problem 39

2.40.1 Solving as second order bessel ode ode 1054

Internal problem ID [7176]

Internal file name [OUTPUT/6162_Sunday_June_05_2022_04_26_06_PM_27513699/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 39.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - yx^3 = x^3$$

2.40.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' - yx^5 = x^5 \tag{1}$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{2i}{5} \\ n &= \frac{1}{5} \\ \gamma &= \frac{5}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

$$y_2 = \sqrt{x} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \\ \frac{d}{dx}\left(\sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) & \frac{d}{dx}\left(\sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \\ \frac{\text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2\sqrt{x}} + ix^2 \left(-\text{BesselJ}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2x^{\frac{5}{2}}} \right) & \frac{\text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2\sqrt{x}} + ix^2 \left(-\text{BesselY}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2x^{\frac{5}{2}}} \right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(\sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right) \left(\frac{\text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2\sqrt{x}} \right. \\ &\quad \left. + ix^2 \left(-\text{BesselY}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2x^{\frac{5}{2}}} \right) \right) \\ &\quad - \left(\sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right) \left(\frac{\text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2\sqrt{x}} \right. \\ &\quad \left. + ix^2 \left(-\text{BesselJ}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2x^{\frac{5}{2}}} \right) \right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} W &= -ix^{\frac{5}{2}} \left(\text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \text{BesselY}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right. \\ &\quad \left. - \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \text{BesselJ}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right) \end{aligned}$$

Which simplifies to

$$W = \frac{5}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^{\frac{11}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{\frac{5x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{\frac{7}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \pi}{5} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{\alpha^{\frac{7}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) \pi}{5} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{\frac{11}{2}} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{\frac{5x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{\frac{7}{2}} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \pi}{5} dx$$

Hence

$$u_2 = \frac{5^{\frac{4}{5}} (-1)^{\frac{1}{10}} \sin \left(\frac{\pi}{5} \right) \Gamma \left(\frac{4}{5} \right) x^5 \text{hypergeom} \left([1], \left[\frac{6}{5}, 2 \right], \frac{x^5}{25} \right)}{25}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \left(\int_0^x \frac{\alpha^{\frac{7}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) \pi}{5} d\alpha \right) \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + \frac{5^{\frac{4}{5}} (-1)^{\frac{1}{10}} \sin \left(\frac{\pi}{5} \right) \Gamma \left(\frac{4}{5} \right) x^{\frac{11}{2}} \text{hypergeom} \left([1], \left[\frac{6}{5}, 2 \right], \frac{x^5}{25} \right) \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{25}$$

Which simplifies to

$$y_p(x) = \frac{\sqrt{x} \left(5^{\frac{4}{5}} (-1)^{\frac{1}{10}} \sin\left(\frac{\pi}{5}\right) \Gamma\left(\frac{4}{5}\right) x^5 \text{hypergeom}\left([1], \left[\frac{6}{5}, 2\right], \frac{x^5}{25}\right) \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - 5\pi \left(\int_0^x \alpha^{\frac{7}{2}} \text{BesselY}\left(\frac{1}{5}, \frac{2\alpha^{\frac{5}{2}}}{5}\right) d\alpha\right) \right)}{25}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 \sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + c_2 \sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right) + \left(\frac{\sqrt{x} \left(5^{\frac{4}{5}} (-1)^{\frac{1}{10}} \sin\left(\frac{\pi}{5}\right) \Gamma\left(\frac{4}{5}\right) x^5 \text{hypergeom}\left([1], \left[\frac{6}{5}, 2\right], \frac{x^5}{25}\right) \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - 5\pi \left(\int_0^x \alpha^{\frac{7}{2}} \text{BesselY}\left(\frac{1}{5}, \frac{2\alpha^{\frac{5}{2}}}{5}\right) d\alpha\right) \right)}{25} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + c_2 \sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + \frac{\sqrt{x} \left(5^{\frac{4}{5}} (-1)^{\frac{1}{10}} \sin\left(\frac{\pi}{5}\right) \Gamma\left(\frac{4}{5}\right) x^5 \text{hypergeom}\left([1], \left[\frac{6}{5}, 2\right], \frac{x^5}{25}\right) \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - 5\pi \left(\int_0^x \alpha^{\frac{7}{2}} \text{BesselY}\left(\frac{1}{5}, \frac{2\alpha^{\frac{5}{2}}}{5}\right) d\alpha\right) \right)}{25} \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + c_2 \sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + \frac{\sqrt{x} \left(5^{\frac{4}{5}} (-1)^{\frac{1}{10}} \sin\left(\frac{\pi}{5}\right) \Gamma\left(\frac{4}{5}\right) x^5 \text{hypergeom}\left([1], \left[\frac{6}{5}, 2\right], \frac{x^5}{25}\right) \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - 5\pi \left(\int_0^x \alpha^{\frac{7}{2}} \text{BesselY}\left(\frac{1}{5}, \frac{2\alpha^{\frac{5}{2}}}{5}\right) d\alpha\right) \right)}{25}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    trying a symmetry of the form [xi=0, eta=F(x)]  
    checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
    -> Bessel  
        <- Bessel successful  
    <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)-x^3*y(x)-x^3=0,y(x), singsol=all)
```

$$y(x) = \sqrt{x} \operatorname{BesselI}\left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5}\right) c_2 + \sqrt{x} \operatorname{BesselK}\left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5}\right) c_1 - 1$$

✓ Solution by Mathematica

Time used: 0.275 (sec). Leaf size: 217

```
DSolve[y''[x]-x^3*y[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{\sqrt[5]{-1} \Gamma\left(\frac{4}{5}\right) \left(5^{4/5} x^5 \Gamma\left(\frac{6}{5}\right) \text{Hypergeometric0F1Regularized}\left(\frac{9}{5}, \frac{x^5}{25}\right) \text{BesselI}\left(\frac{1}{5}, \frac{2x^{5/2}}{5}\right) + 5 \sqrt[5]{25} \sqrt[5]{x^{5/2}} \text{Root}\left[25x^5 - 1, 1\right]\right)}{25 \sqrt[5]{x^{5/2}} \text{Root}\left[25x^5 - 1, 1\right]}$$

$$+ \frac{c_1 \sqrt{x} \Gamma\left(\frac{4}{5}\right) \text{BesselI}\left(-\frac{1}{5}, \frac{2x^{5/2}}{5}\right)}{\sqrt[5]{5}}$$

$$+ \sqrt[5]{-\frac{1}{5}} c_2 \sqrt{x} \Gamma\left(\frac{6}{5}\right) \text{BesselI}\left(\frac{1}{5}, \frac{2x^{5/2}}{5}\right)$$

2.41 problem 40

2.41.1 Solving as second order bessel ode ode 1061

Internal problem ID [7177]

Internal file name [OUTPUT/6163_Sunday_June_05_2022_04_26_07_PM_78234632/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 40.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - yx^3 = x^4$$

2.41.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' - yx^5 = x^6 \tag{1}$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= \frac{2i}{5} \\ n &= \frac{1}{5} \\ \gamma &= \frac{5}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

$$y_2 = \sqrt{x} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \\ \frac{d}{dx}\left(\sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) & \frac{d}{dx}\left(\sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) & \sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \\ \frac{\text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2\sqrt{x}} + ix^2 \left(-\text{BesselJ}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2x^{\frac{5}{2}}} \right) & \frac{\text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2\sqrt{x}} + ix^2 \left(-\text{BesselY}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2x^{\frac{5}{2}}} \right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(\sqrt{x} \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right) \left(\frac{\text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2\sqrt{x}} \right. \\ &\quad \left. + ix^2 \left(-\text{BesselY}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2x^{\frac{5}{2}}} \right) \right) \\ &\quad - \left(\sqrt{x} \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right) \left(\frac{\text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2\sqrt{x}} \right. \\ &\quad \left. + ix^2 \left(-\text{BesselJ}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{2x^{\frac{5}{2}}} \right) \right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} W &= -ix^{\frac{5}{2}} \left(\text{BesselJ}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \text{BesselY}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right. \\ &\quad \left. - \text{BesselY}\left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \text{BesselJ}\left(\frac{6}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \right) \end{aligned}$$

Which simplifies to

$$W = \frac{5}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^{\frac{13}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{\frac{5x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \pi}{5} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{\alpha^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) \pi}{5} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{\frac{13}{2}} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{\frac{5x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{\frac{9}{2}} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \pi}{5} dx$$

Hence

$$u_2 = \frac{(-1)^{\frac{1}{10}} \pi x^{\frac{7}{2}} \text{BesselI} \left(\frac{6}{5}, \frac{2x^{\frac{5}{2}}}{5} \right)}{5 \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}}}$$

Which simplifies to

$$u_1 = - \frac{\pi \left(\int_0^x \alpha^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) d\alpha \right)}{5}$$

$$u_2 = \frac{(-1)^{\frac{1}{10}} \pi x \left(\text{BesselI} \left(-\frac{4}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) x^{\frac{5}{2}} - \text{BesselI} \left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) \right)}{5 \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\pi \left(\int_0^x \alpha^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) d\alpha \right) \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{5} + \frac{(-1)^{\frac{1}{10}} \pi x^{\frac{3}{2}} \left(\text{BesselI} \left(-\frac{4}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) x^{\frac{5}{2}} - \text{BesselI} \left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) \right) \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{5 \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}}}$$

Which simplifies to

$$y_p(x) = \frac{\sqrt{x} \left(\left(\int_0^x \alpha^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) d\alpha \right) \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}} + (-1)^{\frac{1}{10}} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x \text{BesselI} \left(-\frac{4}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) x^{\frac{5}{2}} - \text{BesselI} \left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) \right) \right)}{5 \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}}}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \right) + \frac{\sqrt{x} \left(\left(\int_0^x \alpha^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) d\alpha \right) \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}} + (-1)^{\frac{1}{10}} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x \text{BesselI} \left(-\frac{4}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) x^{\frac{5}{2}} - \text{BesselI} \left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) \right) \right)}{5 \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}}}$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 \sqrt{x} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + \frac{\sqrt{x} \left(\left(\int_0^x \alpha^{\frac{9}{2}} \text{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) d\alpha \right) \text{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}} + (-1)^{\frac{1}{10}} \text{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x \text{BesselI} \left(-\frac{4}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) x^{\frac{5}{2}} - \text{BesselI} \left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) \right) \right)}{5 \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}}} \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \operatorname{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 \sqrt{x} \operatorname{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)$$

$$\frac{\sqrt{x} \left(\int_0^x \alpha^{\frac{9}{2}} \operatorname{BesselY} \left(\frac{1}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) d\alpha \right) \operatorname{BesselJ} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}} + (-1)^{\frac{1}{10}} \operatorname{BesselY} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x \operatorname{BesselI} \left(\frac{1}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \right)^{\frac{1}{5}}}{5 \left(x^{\frac{5}{2}} \right)^{\frac{1}{5}}}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x), x$2) - x^3*y(x) - x^4 = 0, y(x), singsol=all)
```

$$y(x) = \sqrt{x} \operatorname{BesselI} \left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) c_2 + \sqrt{x} \operatorname{BesselK} \left(\frac{1}{5}, \frac{2x^{\frac{5}{2}}}{5} \right) c_1 - x$$

✓ Solution by Mathematica

Time used: 0.182 (sec). Leaf size: 219

```
DSolve[y''[x]-x^3*y[x]-x^4==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{\sqrt[5]{-1} \Gamma\left(\frac{6}{5}\right) \left(-5^{2/5} \sqrt[5]{x^{5/2}} x^{15/2} \Gamma\left(\frac{4}{5}\right) \text{Hypergeometric0F1Regularized}\left(\frac{11}{5}, \frac{x^5}{25}\right) \text{BesselI}\left(-\frac{1}{5}, \frac{2x^{5/2}}{5}\right)\right)}{25x^{3/2} \text{Root}\left[5x^5 - 1, 1\right]} + \frac{c_1 \sqrt{x} \Gamma\left(\frac{4}{5}\right) \text{BesselI}\left(-\frac{1}{5}, \frac{2x^{5/2}}{5}\right)}{\sqrt[5]{5}} + \sqrt[5]{-\frac{1}{5}} c_2 \sqrt{x} \Gamma\left(\frac{6}{5}\right) \text{BesselI}\left(\frac{1}{5}, \frac{2x^{5/2}}{5}\right)$$

2.42 problem 41

Internal problem ID [7178]

Internal file name [OUTPUT/6164_Sunday_June_05_2022_04_26_09_PM_19571536/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 41.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - x^2 y' - x^2 y = x^2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
        -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunT ODE, case c = 0
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 55

```
dsolve(diff(y(x),x$2)-x^2*diff(y(x),x)-x^2*y(x)-x^2=0,y(x), singsol=all)
```

$$y(x) = \text{HeunT} \left(3^{\frac{2}{3}}, 3, 2 \cdot 3^{\frac{1}{3}}, \frac{3^{\frac{2}{3}} x}{3} \right) e^{-x} c_2 + \text{HeunT} \left(3^{\frac{2}{3}}, -3, 2 \cdot 3^{\frac{1}{3}}, -\frac{3^{\frac{2}{3}} x}{3} \right) e^{\frac{x(x^2+3)}{3}} c_1 - 1$$

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]-x^2*y'[x]-x^2*y[x]-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

2.43 problem 42

Internal problem ID [7179]

Internal file name [OUTPUT/6165_Sunday_June_05_2022_04_26_12_PM_13843688/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 42.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - y'x^3 - yx^3 = x^3$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
        -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * {}_2F_1$ 
    trying a symmetry of the form [xi=0, eta=F(x)]
    trying symmetries linear in x and y(x)
    trying to convert to a linear ODE with constant coefficients
    trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
        -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
```

X Solution by Maple

```
dsolve(diff(y(x),x$2)-x^3*diff(y(x),x)-x^3*y(x)-x^3=0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]-x^3*y'[x]-x^3*y[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

2.44 problem 43

2.44.1 Solving using Kovacic algorithm 1074

Internal problem ID [7180]

Internal file name [OUTPUT/6166_Sunday_June_05_2022_04_26_16_PM_7748534/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 43.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xy' - yx = x$$

2.44.1 Solving using Kovacic algorithm

Writing the ode as

$$y'' - xy' - yx = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -x \\ C &= -x \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x - 2}{4} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 4x - 2$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 108: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x}{2} + 1 - \frac{3}{2x} + \frac{3}{x^2} - \frac{33}{4x^3} + \frac{51}{2x^4} - \frac{339}{4x^5} + \frac{591}{2x^6} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= \frac{x}{2} + 1 \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}x^2 + x + 1$$

This shows that the coefficient of 1 in the above is 1. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x - 2}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) + (0) \\ &= \frac{1}{4}x^2 + x - \frac{1}{2} \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2} \right) - (1) \\ &= -\frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x}{2} + 1 \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = -2 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-\frac{3}{2}}{\frac{1}{2}} - 1 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{1}{4}x^2 + x - \frac{1}{2}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$\frac{x}{2} + 1$	-2	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either + or - and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x}{2} + 1 \right) \\ &= -1 - \frac{x}{2} \\ &= -1 - \frac{x}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-1 - \frac{x}{2} \right) (1) + \left(\left(-\frac{1}{2} \right) + \left(-1 - \frac{x}{2} \right)^2 - \left(\frac{1}{4}x^2 + x - \frac{1}{2} \right) \right) &= 0 \\ -2 + a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 2\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x + 2$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x + 2)e^{\int (-1 - \frac{x}{2}) dx} \\ &= (x + 2)e^{-x - \frac{1}{4}x^2} \\ &= (x + 2)e^{-\frac{x(4+x)}{4}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{1} dx} \\ &= z_1 e^{\frac{x^2}{4}} \\ &= z_1 \left(e^{\frac{x^2}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (x + 2)e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^2}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{-ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 ((x+2) e^{-x}) + c_2 \left((x+2) e^{-x} \left(\frac{-ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) - 2e^{\frac{x(4+x)}{2}}}{2x+4} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - xy' - yx = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x+2) e^{-x} - \frac{c_2 e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x+2) e^{-x}$$

$$y_2 = - \frac{e^{-x} \left(ie^{-2}(x+2) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2e^{\frac{x(4+x)}{2}} \right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ \frac{d}{dx}((x+2)e^{-x}) & \frac{d}{dx} \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+2)e^{-x} & -\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \\ e^{-x} - (x+2)e^{-x} & \frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} \right)}{2} \end{vmatrix}$$

Therefore

$$W = ((x+2)e^{-x}) \left(\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} - \frac{e^{-x} \left(i\sqrt{\pi}e^{-2}\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - 2e^{-2(x+2)}e^{\frac{(x+2)^2}{2}} + 2(x+2)e^{\frac{x(4+x)}{2}} \right)}{2} \right) - \left(-\frac{e^{-x} \left(ie^{-2(x+2)}\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) (e^{-x} - (x+2)e^{-x})$$

Which simplifies to

$$W = e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} x - e^{-2x} e^{\frac{x(4+x)}{2}} x^2 + 4e^{\frac{(x+2)^2}{2}} e^{-2} e^{-2x} - 4e^{-2x} e^{\frac{x(4+x)}{2}} x - 3e^{-2x} e^{\frac{x(4+x)}{2}}$$

Which simplifies to

$$W = e^{\frac{x^2}{2}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right) x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\frac{x(x+2)}{2}} x \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2} dx$$

Hence

$$u_1 = - \frac{i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x}}{2} + \frac{i e^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2})}{2} - e^x + 1$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x+2) e^{-x} x}{e^{\frac{x^2}{2}}} dx$$

Which simplifies to

$$u_2 = \int x(x+2) e^{-\frac{x(x+2)}{2}} dx$$

Hence

$$u_2 = -(1+x) e^{-\frac{x(x+2)}{2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(- \frac{i \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) \sqrt{\pi} (1+x) \sqrt{2} e^{-2-\frac{1}{2}x^2-x}}{2} + \frac{i e^{-2} \sqrt{2} \sqrt{\pi} \operatorname{erf} (i\sqrt{2})}{2} - e^x + 1 \right) (x+2) e^{-x} + \frac{(1+x) e^{-\frac{x(x+2)}{2}} e^{-x} \left(i e^{-2(x+2)} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{i\sqrt{2}(x+2)}{2} \right) + 2 e^{\frac{x(4+x)}{2}} \right)}{2}$$

Which simplifies to

$$y_p(x) = -1 - \frac{\sqrt{2} \sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1(x+2)e^{-x} - \frac{c_2e^{-x} \left(ie^{-2}(x+2)\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) + 2e^{\frac{x(4+x)}{2}} \right)}{2} \right) \\ &\quad + \left(-1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y &= -\frac{ic_2e^{-x-2}\sqrt{\pi}(x+2)\sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right)}{2} - c_2e^{\frac{x(x+2)}{2}} + c_1(x+2)e^{-x} \\ &\quad - 1 - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erfi}(\sqrt{2})(x+2)e^{-x-2}}{2} + (x+2)e^{-x} \end{aligned}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 54

```
dsolve(diff(y(x),x$2)-x*diff(y(x),x)-x*y(x)-x=0,y(x), singsol=all)
```

$$y(x) = \pi e^{-2-x} c_1 (x+2) \operatorname{erf}\left(\frac{i\sqrt{2}(x+2)}{2}\right) - i\sqrt{\pi} \sqrt{2} e^{\frac{x(x+2)}{2}} c_1 - 1 + e^{-x} (x+2) c_2$$

✓ Solution by Mathematica

Time used: 0.809 (sec). Leaf size: 216

`DSolve[y''[x]-x*y'[x]-x*y[x]-x==0,y[x],x,IncludeSingularSolutions -> True]`

$$\begin{aligned}
 y(x) \rightarrow & \frac{1}{2}e^{-\frac{1}{2}(x+2)^2} \left(2\sqrt{2}e^{\frac{x^2}{2}+x+2}(x+2) \int_1^x \left(\frac{e^{K[1]}K[1]}{\sqrt{2}} \right. \right. \\
 & - \left. \frac{1}{2}e^{-\frac{1}{2}K[1]^2-K[1]-2}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{(K[1]+2)^2}}{\sqrt{2}}\right) K[1]\sqrt{(K[1]+2)^2} \right) dK[1] \\
 & - \sqrt{2\pi}\sqrt{(x+2)^2} \left(c_2e^{\frac{x^2}{2}+x+2} + x + 1 \right) \operatorname{erfi}\left(\frac{\sqrt{(x+2)^2}}{\sqrt{2}}\right) \\
 & \left. + 2e^{\frac{x^2}{2}+x+2} \left(e^x(x+1) + \sqrt{2}c_1(x+2) + c_2e^{\frac{1}{2}(x+2)^2} \right) \right)
 \end{aligned}$$

2.45 problem 44

Internal problem ID [7181]

Internal file name [OUTPUT/6167_Sunday_June_05_2022_04_26_19_PM_22928568/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 44.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - x^2y' - yx = x^2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
dsolve(diff(y(x),x$2)-x^2*diff(y(x),x)-x*y(x)-x^2=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^3}{6}} \sqrt{x} \operatorname{BesselI}\left(\frac{1}{6}, \frac{x^3}{6}\right) c_2 + e^{\frac{x^3}{6}} \sqrt{x} \operatorname{BesselK}\left(\frac{1}{6}, \frac{x^3}{6}\right) c_1 - \frac{x}{2}$$

✓ Solution by Mathematica

Time used: 0.344 (sec). Leaf size: 224

```
DSolve[y''[x]-x^2*y'[x]-x*y[x]-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x) \rightarrow$

$$e^{\frac{x^3}{6}} \left(12(x^3)^{5/6} \Gamma\left(\frac{1}{6}\right) \Gamma\left(\frac{7}{6}\right) \operatorname{BesselI}\left(\frac{1}{6}, \frac{x^3}{6}\right) {}_1F_1\left(-\frac{2}{3}; -\frac{1}{3}; -\frac{x^3}{6}\right) + \sqrt[3]{23}^{2/3} \sqrt[6]{x^3 x^6} \Gamma\left(\frac{1}{6}\right) \Gamma\left(\frac{7}{6}\right) \operatorname{BesselK}\left(\frac{1}{6}, \frac{x^3}{6}\right) \right) c_1 + \frac{x}{2}$$

2.46 problem 45

Internal problem ID [7182]

Internal file name [OUTPUT/6168_Sunday_June_05_2022_04_26_24_PM_50542001/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 45.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - x^2y' - x^2y = x^3 + x^2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
        -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunT ODE, case c = 0
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 57

```
dsolve(diff(y(x),x$2)-x^2*diff(y(x),x)-x^2*y(x)-x^3-x^2=0,y(x), singsol=all)
```

$$y(x) = \text{HeunT}\left(3^{\frac{2}{3}}, 3, 2 \cdot 3^{\frac{1}{3}}, \frac{3^{\frac{2}{3}}x}{3}\right) e^{-x} c_2 + \text{HeunT}\left(3^{\frac{2}{3}}, -3, 2 \cdot 3^{\frac{1}{3}}, -\frac{3^{\frac{2}{3}}x}{3}\right) e^{\frac{x(x^2+3)}{3}} c_1 - x$$

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]-x^2*y'[x]-x^2*y[x]-x^3-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

2.47 problem 46

Internal problem ID [7183]

Internal file name [OUTPUT/6169_Sunday_June_05_2022_04_26_27_PM_63707127/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 46.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - x^2y' - yx^3 = x^4 + x^2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
        -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunT ODE, case c = 0
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 74

```
dsolve(diff(y(x),x$2)-x^2*diff(y(x),x)-x^3*y(x)-x^4-x^2=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x(x-2)}{2}} \operatorname{HeunT}\left(2\sqrt[3]{3}, -3, -3\sqrt[3]{3}, \frac{3^{\frac{2}{3}}(x+1)}{3}\right) c_2 \\ + e^{\frac{1}{3}x^3 + \frac{1}{2}x^2 - x} \operatorname{HeunT}\left(2\sqrt[3]{3}, 3, -3\sqrt[3]{3}, -\frac{3^{\frac{2}{3}}(x+1)}{3}\right) c_1 - x$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]-x^2*y'[x]-x^3*y[x]-x^4-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

2.48 problem 47

2.48.1 Solving as second order bessel ode ode 1094

Internal problem ID [7184]

Internal file name [OUTPUT/6170_Sunday_June_05_2022_04_26_30_PM_73972892/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 47.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - \frac{y'}{x} - yx = x^2 + \frac{1}{x}$$

2.48.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' - xy' - yx^3 = x^2 \left(x^2 + \frac{1}{x} \right) \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= 1 \\ \beta &= \frac{2i}{3} \\ n &= \frac{2}{3} \\ \gamma &= \frac{3}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 x \text{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + c_2 x \text{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 x \text{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + c_2 x \text{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x \text{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)$$

$$y_2 = x \text{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x \operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) & x \operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) \\ \frac{d}{dx}\left(x \operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)\right) & \frac{d}{dx}\left(x \operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x \operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) & x \operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) \\ \operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + ix^{\frac{3}{2}}\left(\operatorname{BesselJ}\left(-\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + \frac{i \operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)}{x^{\frac{3}{2}}}\right) & \operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + ix^{\frac{3}{2}}\left(\operatorname{BesselY}\left(-\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + \frac{i \operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)}{x^{\frac{3}{2}}}\right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(x \operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)\right) \left(\operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)\right) \\ &\quad + ix^{\frac{3}{2}} \left(\operatorname{BesselY}\left(-\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + \frac{i \operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)}{x^{\frac{3}{2}}}\right) \\ &\quad - \left(x \operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)\right) \left(\operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)\right) \\ &\quad + ix^{\frac{3}{2}} \left(\operatorname{BesselJ}\left(-\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) + \frac{i \operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)}{x^{\frac{3}{2}}}\right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} W &= ix^{\frac{5}{2}} \left(\operatorname{BesselJ}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) \operatorname{BesselY}\left(-\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right)\right) \\ &\quad - \operatorname{BesselY}\left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) \operatorname{BesselJ}\left(-\frac{1}{3}, \frac{2ix^{\frac{3}{2}}}{3}\right) \end{aligned}$$

Which simplifies to

$$W = \frac{3x}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 \text{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \left(x^2 + \frac{1}{x} \right)}{\frac{3x^3}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\text{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) (x^3 + 1) \pi}{3x} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{\text{BesselY} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3 + 1) \pi}{3\alpha} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 \text{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \left(x^2 + \frac{1}{x} \right)}{\frac{3x^3}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{\text{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) (x^3 + 1) \pi}{3x} dx$$

Hence

$$u_2 = \int_0^x \frac{\text{BesselJ} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3 + 1) \pi}{3\alpha} d\alpha$$

Which simplifies to

$$u_1 = - \frac{\pi \left(\int_0^x \frac{\text{BesselY} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3 + 1)}{\alpha} d\alpha \right)}{3}$$

$$u_2 = - \frac{\pi \left(\int_0^x \frac{\text{BesselJ} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3 + 1)}{\alpha} d\alpha \right)}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{\pi \left(\int_0^x \frac{\text{BesselY} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3 + 1)}{\alpha} d\alpha \right) x \text{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)}{3} \\ + \frac{\pi \left(\int_0^x \frac{\text{BesselJ} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3 + 1)}{\alpha} d\alpha \right) x \text{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)}{3}$$

Which simplifies to

$$y_p(x) = \frac{\pi x \left(\left(\int_0^x \frac{\text{BesselY} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3 + 1)}{\alpha} d\alpha \right) \text{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) - \left(\int_0^x \frac{\text{BesselJ} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3 + 1)}{\alpha} d\alpha \right) \text{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \right)}{3}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(c_1 x \operatorname{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 x \operatorname{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \right) \\ + \left(\frac{\pi x \left(\left(\int_0^x \frac{\operatorname{BesselY} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3+1)}{\alpha} d\alpha \right) \operatorname{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) - \left(\int_0^x \frac{\operatorname{BesselJ} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3+1)}{\alpha} d\alpha \right) \operatorname{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)}{3} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 x \operatorname{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 x \operatorname{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \quad (1) \\ \frac{\pi x \left(\left(\int_0^x \frac{\operatorname{BesselY} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3+1)}{\alpha} d\alpha \right) \operatorname{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) - \left(\int_0^x \frac{\operatorname{BesselJ} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3+1)}{\alpha} d\alpha \right) \operatorname{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)}{3}$$

Verification of solutions

$$y = c_1 x \operatorname{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) + c_2 x \operatorname{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) \\ \frac{\pi x \left(\left(\int_0^x \frac{\operatorname{BesselY} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3+1)}{\alpha} d\alpha \right) \operatorname{BesselJ} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right) - \left(\int_0^x \frac{\operatorname{BesselJ} \left(\frac{2}{3}, \frac{2i\alpha^{\frac{3}{2}}}{3} \right) (\alpha^3+1)}{\alpha} d\alpha \right) \operatorname{BesselY} \left(\frac{2}{3}, \frac{2ix^{\frac{3}{2}}}{3} \right)}{3}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-1/x*diff(y(x),x)-x*y(x)-x^2-1/x=0,y(x), singsol=all)
```

$$y(x) = x \left(-1 + \text{BesselI} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) c_2 + \text{BesselK} \left(\frac{2}{3}, \frac{2x^{\frac{3}{2}}}{3} \right) c_1 \right)$$

✓ Solution by Mathematica

Time used: 0.487 (sec). Leaf size: 253

```
DSolve[y''[x]-1/x*y'[x]-x*y[x]-x^2-1/x==0,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{3 \sqrt[6]{3} \pi \Gamma\left(-\frac{1}{3}\right) \left(3 \text{AiryAiPrime}(x) + \sqrt{3} \text{AiryBiPrime}(x)\right) {}_1F_2\left(-\frac{1}{3}, \frac{1}{3}, \frac{2}{3}, \frac{x^3}{9}\right)}{x \Gamma\left(\frac{2}{3}\right)} + \frac{\sqrt[3]{3} \pi x \Gamma\left(\frac{1}{3}\right)^2 \left(\sqrt{3} \text{AiryAiPrime}(x) - \text{AiryBiPrime}(x)\right)}{\Gamma\left(\frac{4}{3}\right)}$$

2.49 problem 48

- 2.49.1 Solving as second order change of variable on x method 2 ode . 1101
- 2.49.2 Solving as second order change of variable on x method 1 ode . 1106
- 2.49.3 Solving as second order bessel ode ode 1111
- 2.49.4 Solving using Kovacic algorithm 1114

Internal problem ID [7185]

Internal file name [OUTPUT/6171_Sunday_June_05_2022_04_26_33_PM_98289116/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 48.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_bessel_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - \frac{y'}{x} - x^2y = x^3 + \frac{1}{x}$$

2.49.1 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - \frac{y'}{x} - x^2y = 0$$

In normal form the ode

$$y'' - \frac{y'}{x} - x^2y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{1}{x}$$
$$q(x) = -x^2$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int -\frac{1}{x}dx)} dx \\ &= \int e^{\ln(x)} dx \\ &= \int x dx \\ &= \frac{x^2}{2} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-x^2}{x^2} \\ &= -1 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - y(\tau) &= 0\end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1\end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(1)\tau} + c_2 e^{(-1)\tau}$$

Or

$$y(\tau) = c_1 e^\tau + c_2 e^{-\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = c_1 e^{\frac{x^2}{2}} + c_2 e^{-\frac{x^2}{2}}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{\frac{x^2}{2}} + c_2 e^{-\frac{x^2}{2}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-\frac{x^2}{2}}$$

$$y_2 = e^{\frac{x^2}{2}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-\frac{x^2}{2}} & e^{\frac{x^2}{2}} \\ \frac{d}{dx} \left(e^{-\frac{x^2}{2}} \right) & \frac{d}{dx} \left(e^{\frac{x^2}{2}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-\frac{x^2}{2}} & e^{\frac{x^2}{2}} \\ -x e^{-\frac{x^2}{2}} & x e^{\frac{x^2}{2}} \end{vmatrix}$$

Therefore

$$W = \left(e^{-\frac{x^2}{2}} \right) \left(x e^{\frac{x^2}{2}} \right) - \left(e^{\frac{x^2}{2}} \right) \left(-x e^{-\frac{x^2}{2}} \right)$$

Which simplifies to

$$W = 2 e^{-\frac{x^2}{2}} x e^{\frac{x^2}{2}}$$

Which simplifies to

$$W = 2x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{\frac{x^2}{2}} \left(x^3 + \frac{1}{x} \right)}{2x} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{\frac{x^2}{2}} (x^4 + 1)}{2x^2} dx$$

Hence

$$u_1 = - \frac{(x^2 - 1) e^{\frac{x^2}{2}}}{2x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-\frac{x^2}{2}} \left(x^3 + \frac{1}{x} \right)}{2x} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-\frac{x^2}{2}} (x^4 + 1)}{2x^2} dx$$

Hence

$$u_2 = - \frac{(x^2 + 1) e^{-\frac{x^2}{2}}}{2x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{(x^2 - 1)e^{\frac{x^2}{2}}e^{-\frac{x^2}{2}}}{2x} - \frac{(x^2 + 1)e^{-\frac{x^2}{2}}e^{\frac{x^2}{2}}}{2x}$$

Which simplifies to

$$y_p(x) = -x$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= \left(c_1e^{\frac{x^2}{2}} + c_2e^{-\frac{x^2}{2}}\right) + (-x)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{\frac{x^2}{2}} + c_2e^{-\frac{x^2}{2}} - x \quad (1)$$

Verification of solutions

$$y = c_1e^{\frac{x^2}{2}} + c_2e^{-\frac{x^2}{2}} - x$$

Verified OK.

2.49.2 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -\frac{1}{x}, C = -x^2, f(x) = x^3 + \frac{1}{x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$y'' - \frac{y'}{x} - x^2y = 0$$

In normal form the ode

$$y'' - \frac{y'}{x} - x^2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{1}{x}$$
$$q(x) = -x^2$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$
$$= \frac{\sqrt{-x^2}}{c} \quad (6)$$
$$\tau'' = -\frac{x}{c\sqrt{-x^2}}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2}$$
$$= \frac{-\frac{x}{c\sqrt{-x^2}} - \frac{1}{x}\frac{\sqrt{-x^2}}{c}}{\left(\frac{\sqrt{-x^2}}{c}\right)^2}$$
$$= 0$$

Therefore ode (3) now becomes

$$y(\tau)'' + p_1y(\tau)' + q_1y(\tau) = 0$$
$$\frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) = 0 \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{-x^2} dx}{c} \\ &= \frac{x\sqrt{-x^2}}{2c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cosh\left(\frac{x^2}{2}\right) + ic_2 \sinh\left(\frac{x^2}{2}\right)$$

Now the particular solution to this ODE is found

$$y'' - \frac{y'}{x} - x^2 y = x^3 + \frac{1}{x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{-\frac{x^2}{2}} \\ y_2 &= e^{\frac{x^2}{2}} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-\frac{x^2}{2}} & e^{\frac{x^2}{2}} \\ \frac{d}{dx}\left(e^{-\frac{x^2}{2}}\right) & \frac{d}{dx}\left(e^{\frac{x^2}{2}}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-\frac{x^2}{2}} & e^{\frac{x^2}{2}} \\ -x e^{-\frac{x^2}{2}} & x e^{\frac{x^2}{2}} \end{vmatrix}$$

Therefore

$$W = \left(e^{-\frac{x^2}{2}}\right) \left(x e^{\frac{x^2}{2}}\right) - \left(e^{\frac{x^2}{2}}\right) \left(-x e^{-\frac{x^2}{2}}\right)$$

Which simplifies to

$$W = 2e^{-\frac{x^2}{2}} x e^{\frac{x^2}{2}}$$

Which simplifies to

$$W = 2x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{\frac{x^2}{2}} \left(x^3 + \frac{1}{x}\right)}{2x} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{\frac{x^2}{2}} (x^4 + 1)}{2x^2} dx$$

Hence

$$u_1 = - \frac{(x^2 - 1) e^{\frac{x^2}{2}}}{2x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-\frac{x^2}{2}} \left(x^3 + \frac{1}{x}\right)}{2x} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-\frac{x^2}{2}}(x^4 + 1)}{2x^2} dx$$

Hence

$$u_2 = -\frac{(x^2 + 1)e^{-\frac{x^2}{2}}}{2x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{(x^2 - 1)e^{\frac{x^2}{2}}e^{-\frac{x^2}{2}}}{2x} - \frac{(x^2 + 1)e^{-\frac{x^2}{2}}e^{\frac{x^2}{2}}}{2x}$$

Which simplifies to

$$y_p(x) = -x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cosh\left(\frac{x^2}{2}\right) + ic_2 \sinh\left(\frac{x^2}{2}\right) \right) + (-x) \\ &= -x + c_1 \cosh\left(\frac{x^2}{2}\right) + ic_2 \sinh\left(\frac{x^2}{2}\right) \end{aligned}$$

Which simplifies to

$$y = -x + c_1 \cosh\left(\frac{x^2}{2}\right) + ic_2 \sinh\left(\frac{x^2}{2}\right)$$

Summary

The solution(s) found are the following

$$y = -x + c_1 \cosh\left(\frac{x^2}{2}\right) + ic_2 \sinh\left(\frac{x^2}{2}\right) \quad (1)$$

Verification of solutions

$$y = -x + c_1 \cosh\left(\frac{x^2}{2}\right) + ic_2 \sinh\left(\frac{x^2}{2}\right)$$

Verified OK.

2.49.3 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' - xy' - yx^4 = x^2 \left(x^3 + \frac{1}{x} \right) \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= 1 \\ \beta &= \frac{i}{2} \\ n &= \frac{1}{2} \\ \gamma &= 2 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = \frac{2ic_1 x \sinh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}} - \frac{2c_2 x \cosh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{2ic_1 x \sinh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}} - \frac{2c_2 x \cosh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-\frac{x^2}{2}}$$

$$y_2 = e^{\frac{x^2}{2}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-\frac{x^2}{2}} & e^{\frac{x^2}{2}} \\ \frac{d}{dx} \left(e^{-\frac{x^2}{2}} \right) & \frac{d}{dx} \left(e^{\frac{x^2}{2}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-\frac{x^2}{2}} & e^{\frac{x^2}{2}} \\ -x e^{-\frac{x^2}{2}} & x e^{\frac{x^2}{2}} \end{vmatrix}$$

Therefore

$$W = \left(e^{-\frac{x^2}{2}} \right) \left(x e^{\frac{x^2}{2}} \right) - \left(e^{\frac{x^2}{2}} \right) \left(-x e^{-\frac{x^2}{2}} \right)$$

Which simplifies to

$$W = 2 e^{-\frac{x^2}{2}} x e^{\frac{x^2}{2}}$$

Which simplifies to

$$W = 2x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{\frac{x^2}{2}} x^2 (x^3 + \frac{1}{x})}{2x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{\frac{x^2}{2}} (x^4 + 1)}{2x^2} dx$$

Hence

$$u_1 = - \frac{(x^2 - 1) e^{\frac{x^2}{2}}}{2x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-\frac{x^2}{2}} x^2 (x^3 + \frac{1}{x})}{2x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-\frac{x^2}{2}} (x^4 + 1)}{2x^2} dx$$

Hence

$$u_2 = - \frac{(x^2 + 1) e^{-\frac{x^2}{2}}}{2x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{(x^2 - 1) e^{\frac{x^2}{2}} e^{-\frac{x^2}{2}}}{2x} - \frac{(x^2 + 1) e^{-\frac{x^2}{2}} e^{\frac{x^2}{2}}}{2x}$$

Which simplifies to

$$y_p(x) = -x$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(\frac{2ic_1x \sinh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}} - \frac{2c_2x \cosh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}} \right) + (-x)$$

Summary

The solution(s) found are the following

$$y = \frac{2ic_1x \sinh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}} - \frac{2c_2x \cosh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}} - x \quad (1)$$

Verification of solutions

$$y = \frac{2ic_1x \sinh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}} - \frac{2c_2x \cosh\left(\frac{x^2}{2}\right)}{\sqrt{\pi} \sqrt{ix^2}} - x$$

Verified OK.

2.49.4 Solving using Kovacic algorithm

Writing the ode as

$$y'' - \frac{y'}{x} - x^2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -\frac{1}{x} \quad (3)$$

$$C = -x^2$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4x^4 + 3}{4x^2} \tag{6}$$

Comparing the above to (5) shows that

$$s = 4x^4 + 3$$

$$t = 4x^2$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{4x^4 + 3}{4x^2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 109: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 4 \\ &= -2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Therefore

$$L = [1, 2]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4x^2} + x^2$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx x + \frac{3}{8x^3} - \frac{9}{128x^7} + \frac{27}{1024x^{11}} - \frac{405}{32768x^{15}} + \frac{1701}{262144x^{19}} - \frac{15309}{4194304x^{23}} + \frac{72171}{33554432x^{27}} + \dots \quad (9)$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = 1$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_{\infty} &= \sum_{i=0}^1 a_i x^i \\ &= x \end{aligned} \quad (10)$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_{\infty})^2$ where $[\sqrt{r}]_{\infty}$ was found above in Eq (10). Hence

$$([\sqrt{r}]_{\infty})^2 = x^2$$

This shows that the coefficient of 1 in the above is 0. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{4x^4 + 3}{4x^2} \\ &= Q + \frac{R}{4x^2} \\ &= (x^2) + \left(\frac{3}{4x^2}\right) \\ &= \frac{3}{4x^2} + x^2 \end{aligned}$$

We see that the coefficient of the term x in the quotient is 0. Now b can be found.

$$\begin{aligned} b &= (0) - (0) \\ &= 0 \end{aligned}$$

Hence

$$\begin{aligned}
 [\sqrt{r}]_\infty &= x \\
 \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{0}{1} - 1 \right) = -\frac{1}{2} \\
 \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{0}{1} - 1 \right) = -\frac{1}{2}
 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{4x^4 + 3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	x	$-\frac{1}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned}
 d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\
 &= -\frac{1}{2} - \left(-\frac{1}{2} \right) \\
 &= 0
 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned}
 \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\
 &= -\frac{1}{2x} + (-)(x) \\
 &= -\frac{1}{2x} - x \\
 &= -\frac{1}{2x} - x
 \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned}
 (0) + 2\left(-\frac{1}{2x} - x\right)(0) + \left(\left(\frac{1}{2x^2} - 1\right) + \left(-\frac{1}{2x} - x\right)^2 - \left(\frac{4x^4 + 3}{4x^2}\right)\right) &= 0 \\
 0 &= 0
 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned}
 z_1(x) &= p e^{\int \omega dx} \\
 &= e^{\int \left(-\frac{1}{2x} - x\right) dx} \\
 &= \frac{e^{-\frac{x^2}{2}}}{\sqrt{x}}
 \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\
 &= z_1 e^{\frac{\ln(x)}{2}} \\
 &= z_1 (\sqrt{x})
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x^2}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{x^2}}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x^2}{2}} \right) + c_2 \left(e^{-\frac{x^2}{2}} \left(\frac{e^{x^2}}{2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - \frac{y'}{x} - x^2 y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x^2}{2}} + \frac{c_2 e^{\frac{x^2}{2}}}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-\frac{x^2}{2}}$$

$$y_2 = \frac{e^{\frac{x^2}{2}}}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-\frac{x^2}{2}} & \frac{e^{\frac{x^2}{2}}}{2} \\ \frac{d}{dx} \left(e^{-\frac{x^2}{2}} \right) & \frac{d}{dx} \left(\frac{e^{\frac{x^2}{2}}}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-\frac{x^2}{2}} & \frac{e^{\frac{x^2}{2}}}{2} \\ -x e^{-\frac{x^2}{2}} & \frac{x e^{\frac{x^2}{2}}}{2} \end{vmatrix}$$

Therefore

$$W = \left(e^{-\frac{x^2}{2}} \right) \left(\frac{x e^{\frac{x^2}{2}}}{2} \right) - \left(\frac{e^{\frac{x^2}{2}}}{2} \right) \left(-x e^{-\frac{x^2}{2}} \right)$$

Which simplifies to

$$W = e^{-\frac{x^2}{2}} x e^{\frac{x^2}{2}}$$

Which simplifies to

$$W = x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{\frac{x^2}{2}} (x^3 + \frac{1}{x})}{x} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{\frac{x^2}{2}} (x^4 + 1)}{2x^2} dx$$

Hence

$$u_1 = - \frac{(x^2 - 1) e^{\frac{x^2}{2}}}{2x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-\frac{x^2}{2}} (x^3 + \frac{1}{x})}{x} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-\frac{x^2}{2}} (x^4 + 1)}{x^2} dx$$

Hence

$$u_2 = - \frac{(x^2 + 1) e^{-\frac{x^2}{2}}}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{(x^2 - 1) e^{\frac{x^2}{2}} e^{-\frac{x^2}{2}}}{2x} - \frac{(x^2 + 1) e^{-\frac{x^2}{2}} e^{\frac{x^2}{2}}}{2x}$$

Which simplifies to

$$y_p(x) = -x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{x^2}{2}} + \frac{c_2 e^{\frac{x^2}{2}}}{2} \right) + (-x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{x^2}{2}} + \frac{c_2 e^{\frac{x^2}{2}}}{2} - x \quad (1)$$

Verification of solutions

$$y = c_1 e^{-\frac{x^2}{2}} + \frac{c_2 e^{\frac{x^2}{2}}}{2} - x$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    <- linear_1 successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)-1/x*diff(y(x),x)-x^2*y(x)-x^3-1/x=0,y(x), singsol=all)
```

$$y(x) = \sinh\left(\frac{x^2}{2}\right) c_2 + \cosh\left(\frac{x^2}{2}\right) c_1 - x$$

✓ Solution by Mathematica

Time used: 0.094 (sec). Leaf size: 34

```
DSolve[y''[x]-1/x*y'[x]-x^2*y[x]-x^3-1/x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cosh\left(\frac{x^2}{2}\right) + ic_2 \sinh\left(\frac{x^2}{2}\right) - x$$

2.50 problem 49

2.50.1 Solving as second order bessel ode ode 1125

Internal problem ID [7186]

Internal file name [OUTPUT/6172_Sunday_June_05_2022_04_26_34_PM_47666754/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 49.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

`[[_2nd_order , _with_linear_symmetries]]`

$$y'' - \frac{y'}{x} - yx^3 = x^4 + \frac{1}{x}$$

2.50.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' - xy' - yx^5 = x^2 \left(x^4 + \frac{1}{x} \right) \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= 1 \\ \beta &= \frac{2i}{5} \\ n &= \frac{2}{5} \\ \gamma &= \frac{5}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 x \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + c_2 x \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 x \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + c_2 x \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)$$

$$y_2 = x \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) & x \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \\ \frac{d}{dx}\left(x \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) & \frac{d}{dx}\left(x \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) & x \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \\ \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + ix^{\frac{5}{2}}\left(-\text{BesselJ}\left(\frac{7}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{x^{\frac{5}{2}}}\right) & \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) + ix^{\frac{5}{2}}\left(-\text{BesselY}\left(\frac{7}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{x^{\frac{5}{2}}}\right) \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(x \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) \left(\text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) \\ &\quad + ix^{\frac{5}{2}}\left(-\text{BesselY}\left(\frac{7}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{x^{\frac{5}{2}}}\right) \\ &\quad - \left(x \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) \left(\text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) \\ &\quad + ix^{\frac{5}{2}}\left(-\text{BesselJ}\left(\frac{7}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) - \frac{i \text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)}{x^{\frac{5}{2}}}\right) \end{aligned}$$

Which simplifies to

$$\begin{aligned} W &= -ix^{\frac{7}{2}}\left(\text{BesselJ}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\text{BesselY}\left(\frac{7}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\right) \\ &\quad - \text{BesselY}\left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right)\text{BesselJ}\left(\frac{7}{5}, \frac{2ix^{\frac{5}{2}}}{5}\right) \end{aligned}$$

Which simplifies to

$$W = \frac{5x}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 \text{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x^4 + \frac{1}{x} \right)}{\frac{5x^3}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\text{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) (x^5 + 1) \pi}{5x} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{\text{BesselY} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1) \pi}{5\alpha} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 \text{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \left(x^4 + \frac{1}{x} \right)}{\frac{5x^3}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{\text{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) (x^5 + 1) \pi}{5x} dx$$

Hence

$$u_2 = \int_0^x \frac{\text{BesselJ} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1) \pi}{5\alpha} d\alpha$$

Which simplifies to

$$u_1 = -\frac{\pi \left(\int_0^x \frac{\text{BesselY} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right)}{5}$$

$$u_2 = -\frac{\pi \left(\int_0^x \frac{\text{BesselJ} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right)}{5}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\pi \left(\int_0^x \frac{\text{BesselY} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) x \text{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{5} \\ + \frac{\pi \left(\int_0^x \frac{\text{BesselJ} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) x \text{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{5}$$

Which simplifies to

$$y_p(x) \\ = \frac{\pi x \left(-\left(\int_0^x \frac{\text{BesselY} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) \text{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + \left(\int_0^x \frac{\text{BesselJ} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) \text{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \right)}{5}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$\begin{aligned}
&= \left(c_1 x \operatorname{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 x \operatorname{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \right) \\
&+ \left(\frac{\pi x \left(- \left(\int_0^x \frac{\operatorname{BesselY} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) \operatorname{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + \left(\int_0^x \frac{\operatorname{BesselJ} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) \operatorname{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{5} \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
y &= c_1 x \operatorname{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 x \operatorname{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \tag{1} \\
&+ \left(\frac{\pi x \left(- \left(\int_0^x \frac{\operatorname{BesselY} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) \operatorname{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + \left(\int_0^x \frac{\operatorname{BesselJ} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) \operatorname{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{5} \right)
\end{aligned}$$

Verification of solutions

$$\begin{aligned}
y &= c_1 x \operatorname{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + c_2 x \operatorname{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) \\
&+ \left(\frac{\pi x \left(- \left(\int_0^x \frac{\operatorname{BesselY} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) \operatorname{BesselJ} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right) + \left(\int_0^x \frac{\operatorname{BesselJ} \left(\frac{2}{5}, \frac{2i\alpha^{\frac{5}{2}}}{5} \right) (\alpha^5 + 1)}{\alpha} d\alpha \right) \operatorname{BesselY} \left(\frac{2}{5}, \frac{2ix^{\frac{5}{2}}}{5} \right)}{5} \right)
\end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
            <- Bessel successful
        <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x), x$2)-1/x*diff(y(x), x)-x^3*y(x)-x^4-1/x=0, y(x), singsol=all)
```

$$y(x) = x \left(-1 + \text{BesselI} \left(\frac{2}{5}, \frac{2x^{5/2}}{5} \right) c_2 + \text{BesselK} \left(\frac{2}{5}, \frac{2x^{5/2}}{5} \right) c_1 \right)$$

✓ Solution by Mathematica

Time used: 0.364 (sec). Leaf size: 316

```
DSolve[y''[x]-1/x*y'[x]-x^3*y[x]-x^4-1/x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) = \frac{5(x^{5/2})^{13/5} \Gamma(\frac{4}{5}) \Gamma(\frac{7}{5}) \text{BesselI}(\frac{2}{5}, \frac{2x^{5/2}}{5}) {}_1F_2(\frac{4}{5}; \frac{3}{5}, \frac{9}{5}; \frac{x^5}{25})}{\Gamma(\frac{9}{5})} - \frac{\sqrt[5]{5}(x^{5/2})^{7/5} \Gamma(\frac{1}{5}) \Gamma(\frac{3}{5}) \text{BesselI}(-\frac{2}{5}, \frac{2x^{5/2}}{5}) {}_1F_2(\frac{1}{5}; \frac{3}{5}, \frac{9}{5}; \frac{x^5}{25})}{\Gamma(\frac{6}{5})}$$

→

2.51 problem 50

Internal problem ID [7187]

Internal file name [OUTPUT/6173_Sunday_June_05_2022_04_26_37_PM_44292969/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 50.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - y'x^3 - yx = x^3 + x^2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
        -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1$ 
    trying a symmetry of the form [xi=0, eta=F(x)]
    trying symmetries linear in x and y(x)
    trying to convert to a linear ODE with constant coefficients
    trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
        -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
```

X Solution by Maple

```
dsolve(diff(y(x),x$2)-x^3*diff(y(x),x)-x*y(x)-x^3-x^2=0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]-x^3*y'[x]-x*y[x]-x^3-x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

2.52 problem 51

Internal problem ID [7188]

Internal file name [OUTPUT/6174_Sunday_June_05_2022_04_26_41_PM_38669530/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 51.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - y'x^3 - x^2y = x^3$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
        <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
    <- Kummer successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-x^3*diff(y(x),x)-x^2*y(x)-x^3=0,y(x), singsol=all)
```

$$y(x) = x \left(\text{KummerU} \left(\frac{1}{2}, \frac{5}{4}, \frac{x^4}{4} \right) c_1 + \text{KummerM} \left(\frac{1}{2}, \frac{5}{4}, \frac{x^4}{4} \right) c_2 - \frac{1}{2} \right)$$

✓ Solution by Mathematica

Time used: 1.216 (sec). Leaf size: 337

`DSolve[y''[x]-x^3*y'[x]-x^2*y[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]`

$y(x)$

$$\begin{aligned} &\rightarrow \text{Hypergeometric1F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{x^4}{4}\right) \int_1^x \frac{1}{5 \text{Hypergeometric1F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{K[1]^4}{4}\right) \text{Hypergeometric1F1}\left(\frac{5}{4}, \frac{7}{4}, \frac{K[1]^4}{4}\right)} \\ &+ \frac{\sqrt[4]{-1}x \text{Hypergeometric1F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{x^4}{4}\right) \int_1^x \frac{1}{3 \text{Hypergeometric1F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{K[2]^4}{4}\right) \left(2 \text{Hypergeometric1F1}\left(\frac{3}{2}, \frac{9}{4}, \frac{K[2]^4}{4}\right) K[2]^4 + 5 \text{Hypergeometric1F1}\left(\frac{5}{2}, \frac{11}{4}, \frac{K[2]^4}{4}\right) K[2]^4\right)} dx}{\sqrt{2}} \\ &+ c_1 \text{Hypergeometric1F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{x^4}{4}\right) + \left(\frac{1}{2} + \frac{i}{2}\right) c_2 x \text{Hypergeometric1F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{x^4}{4}\right) \end{aligned}$$

2.53 problem 52

Internal problem ID [7189]

Internal file name [OUTPUT/6175_Sunday_June_05_2022_04_26_44_PM_29136993/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 52.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - y'x^3 - yx^3 = x^4 + x^3$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
        -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
        -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1$ 
    trying a symmetry of the form [xi=0, eta=F(x)]
    trying symmetries linear in x and y(x)
    trying to convert to a linear ODE with constant coefficients
    trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
    trying to convert to an ODE of Bessel type
    -> trying reduction of order to Riccati
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 3` [0, x+y]
```

X Solution by Maple

```
dsolve(diff(y(x),x$2)-x^3*diff(y(x),x)-x^3*y(x)-x^4-x^3=0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]-x^3*y'[x]-x^3*y[x]-x^4-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

2.54 problem 50

2.54.1 Maple step by step solution 1141

Internal problem ID [7190]

Internal file name [OUTPUT/6176_Sunday_June_05_2022_04_26_49_PM_70486816/index.tex]

Book: Own collection of miscellaneous problems

Section: section 2.0

Problem number: 50.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y''' - y'x^3 - x^2y = x^3$$

Unable to solve this ODE.

2.54.1 Maple step by step solution

Let's solve

$$y''' - y'x^3 - x^2y = x^3$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 3; missing the dependent variable
checking if the LODE is of Euler type
trying Louvillian solutions for 3rd order ODEs, imprimitive case
-> pFq: Equivalence to the 3F2 or one of its 3 confluent cases under a power @ Moebius
<- pFq successful: received ODE is equivalent to the 1F2 ODE, case c = 0`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 51

```
dsolve(diff(y(x),x$3)-x^3*diff(y(x),x)-x^2*y(x)-x^3=0,y(x), singsol=all)
```

$$y(x) = -\frac{x}{2} + c_1 \operatorname{hypergeom} \left(\left[\frac{1}{5} \right], \left[\frac{3}{5}, \frac{4}{5} \right], \frac{x^5}{25} \right) \\ + c_2 x \operatorname{hypergeom} \left(\left[\frac{2}{5} \right], \left[\frac{4}{5}, \frac{6}{5} \right], \frac{x^5}{25} \right) + c_3 x^2 \operatorname{hypergeom} \left(\left[\frac{3}{5} \right], \left[\frac{6}{5}, \frac{7}{5} \right], \frac{x^5}{25} \right)$$

✓ Solution by Mathematica

Time used: 12.206 (sec). Leaf size: 2548

```
DSolve[y'''[x]-x^3*y'[x]-x^2*y[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

3 section 3.0

3.1	problem 1	1144
3.2	problem 2	1151
3.3	problem 3	1155
3.4	problem 4	1167
3.5	problem 5	1179
3.6	problem 6	1193
3.7	problem 7	1205
3.8	problem 8	1217
3.9	problem 9	1230
3.10	problem 10	1243
3.11	problem 11	1256
3.12	problem 12	1270
3.13	problem 13	1283
3.14	problem 14	1297
3.15	problem 15	1304
3.16	problem 16	1328
3.17	problem 17	1352
3.18	problem 18	1379
3.19	problem 19	1390
3.20	problem 20	1392
3.21	problem 21	1397
3.22	problem 22	1401
3.23	problem 23	1406
3.24	problem 24	1410
3.25	problem 25	1413
3.26	problem 26	1417
3.27	problem 27	1420
3.28	problem 28	1426
3.29	problem 29	1435
3.30	problem 30	1447
3.31	problem 31	1460

3.1 problem 1

3.1.1 Solving as second order linear constant coeff ode	1144
3.1.2 Solving using Kovacic algorithm	1146
3.1.3 Maple step by step solution	1149

Internal problem ID [7191]

Internal file name [OUTPUT/6177_Sunday_June_05_2022_04_26_50_PM_81515482/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 1.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y'c + ky = 0$$

3.1.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = c, C = k$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + c\lambda e^{\lambda x} + k e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$c\lambda + \lambda^2 + k = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = c, C = k$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-c}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{c^2 - (4)(1)(k)} \\ &= -\frac{c}{2} \pm \frac{\sqrt{c^2 - 4k}}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2} \\ \lambda_2 &= -\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= -\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2} \\ \lambda_2 &= -\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2}\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{\left(-\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2}\right)x} + c_2 e^{\left(-\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2}\right)x}\end{aligned}$$

Or

$$y = c_1 e^{\left(-\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2}\right)x} + c_2 e^{\left(-\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2}\right)x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\left(-\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2}\right)x} + c_2 e^{\left(-\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2}\right)x} \quad (1)$$

Verification of solutions

$$y = c_1 e^{\left(-\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2}\right)x} + c_2 e^{\left(-\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2}\right)x}$$

Verified OK.

3.1.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y'c + ky = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= c \\ C &= k \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{c^2 - 4k}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= c^2 - 4k \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{c^2}{4} - k \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 111: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{c^2}{4} - k$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\frac{x\sqrt{c^2-4k}}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

$$\begin{aligned}
&= z_1 e^{-\int \frac{1}{2} \frac{c}{1} dx} \\
&= z_1 e^{-\frac{cx}{2}} \\
&= z_1 \left(e^{-\frac{cx}{2}} \right)
\end{aligned}$$

Which simplifies to

$$y_1 = e^{\frac{(-c + \sqrt{c^2 - 4k})x}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
y_2 &= y_1 \int \frac{e^{\int -\frac{c}{1} dx}}{(y_1)^2} dx \\
&= y_1 \int \frac{e^{-cx}}{(y_1)^2} dx \\
&= y_1 \left(-\frac{e^{-x\sqrt{c^2 - 4k}}}{\sqrt{c^2 - 4k}} \right)
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 \left(e^{\frac{(-c + \sqrt{c^2 - 4k})x}{2}} \right) + c_2 \left(e^{\frac{(-c + \sqrt{c^2 - 4k})x}{2}} \left(-\frac{e^{-x\sqrt{c^2 - 4k}}}{\sqrt{c^2 - 4k}} \right) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{(-c + \sqrt{c^2 - 4k})x}{2}} - \frac{c_2 e^{-\frac{(c + \sqrt{c^2 - 4k})x}{2}}}{\sqrt{c^2 - 4k}} \quad (1)$$

Verification of solutions

$$y = c_1 e^{\frac{(-c + \sqrt{c^2 - 4k})x}{2}} - \frac{c_2 e^{-\frac{(c + \sqrt{c^2 - 4k})x}{2}}}{\sqrt{c^2 - 4k}}$$

Verified OK.

3.1.3 Maple step by step solution

Let's solve

$$y'' + y'c + ky = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$rc + r^2 + k = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-c) \pm (\sqrt{c^2 - 4k})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2}, -\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2} \right)$$

- 1st solution of the ODE

$$y_1(x) = e^{\left(-\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2} \right)x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{\left(-\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2} \right)x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{\left(-\frac{c}{2} - \frac{\sqrt{c^2 - 4k}}{2} \right)x} + c_2 e^{\left(-\frac{c}{2} + \frac{\sqrt{c^2 - 4k}}{2} \right)x}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
dsolve(diff(y(x),x$2)+c*diff(y(x),x)+k*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{(-c + \sqrt{c^2 - 4k})x}{2}} + c_2 e^{-\frac{(c + \sqrt{c^2 - 4k})x}{2}}$$

✓ Solution by Mathematica

Time used: 8.987 (sec). Leaf size: 2548

```
DSolve[y'''[x]-x^3*y'[x]-x^2*y[x]-x^3==0,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

3.2 problem 2

3.2.1	Existence and uniqueness analysis	1151
3.2.2	Solving as quadrature ode	1152
3.2.3	Maple step by step solution	1153

Internal problem ID [7192]

Internal file name [OUTPUT/6178_Sunday_June_05_2022_04_26_52_PM_3626605/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 2.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$w' + \frac{\sqrt{1-12w}}{2} = -\frac{1}{2}$$

With initial conditions

$$[w(1) = -1]$$

3.2.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} w' &= f(z, w) \\ &= -\frac{1}{2} - \frac{\sqrt{1-12w}}{2} \end{aligned}$$

The w domain of $f(z, w)$ when $z = 1$ is

$$\left\{ w \leq \frac{1}{12} \right\}$$

And the point $w_0 = -1$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial w} &= \frac{\partial}{\partial w} \left(-\frac{1}{2} - \frac{\sqrt{1-12w}}{2} \right) \\ &= \frac{3}{\sqrt{1-12w}}\end{aligned}$$

The w domain of $\frac{\partial f}{\partial w}$ when $z = 1$ is

$$\left\{ w < \frac{1}{12} \right\}$$

And the point $w_0 = -1$ is inside this domain. Therefore solution exists and is unique.

3.2.2 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{-\frac{1}{2} - \frac{\sqrt{1-12w}}{2}} dw = \int dz$$

$$-\frac{\ln(w)}{6} + \frac{\sqrt{1-12w}}{3} + \frac{\ln(-1 + \sqrt{1-12w})}{6} - \frac{\ln(1 + \sqrt{1-12w})}{6} = z + c_1$$

Initial conditions are used to solve for c_1 . Substituting $z = 1$ and $w = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-\frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1 + \sqrt{13})}{6} - \frac{\ln(1 + \sqrt{13})}{6} = c_1 + 1$$

$$c_1 = -1 - \frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1 + \sqrt{13})}{6} - \frac{\ln(1 + \sqrt{13})}{6}$$

Substituting c_1 found above in the general solution gives

$$-\frac{\ln(w)}{6} + \frac{\sqrt{1-12w}}{3} + \frac{\ln(-1 + \sqrt{1-12w})}{6} - \frac{\ln(1 + \sqrt{1-12w})}{6} = z - 1 - \frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1 + \sqrt{13})}{6}$$

Summary

The solution(s) found are the following

$$\begin{aligned}&-\frac{\ln(w)}{6} + \frac{\sqrt{1-12w}}{3} + \frac{\ln(-1 + \sqrt{1-12w})}{6} - \frac{\ln(1 + \sqrt{1-12w})}{6} \\ &= z - 1 - \frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1 + \sqrt{13})}{6} - \frac{\ln(1 + \sqrt{13})}{6}\end{aligned}\tag{1}$$

Verification of solutions

$$\begin{aligned} & -\frac{\ln(w)}{6} + \frac{\sqrt{1-12w}}{3} + \frac{\ln(-1+\sqrt{1-12w})}{6} - \frac{\ln(1+\sqrt{1-12w})}{6} \\ & = z - 1 - \frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1+\sqrt{13})}{6} - \frac{\ln(1+\sqrt{13})}{6} \end{aligned}$$

Verified OK.

3.2.3 Maple step by step solution

Let's solve

$$\left[w' + \frac{\sqrt{1-12w}}{2} = -\frac{1}{2}, w(1) = -1 \right]$$

- Highest derivative means the order of the ODE is 1

w'

- Separate variables

$$\frac{w'}{-\frac{1}{2} - \frac{\sqrt{1-12w}}{2}} = 1$$

- Integrate both sides with respect to z

$$\int \frac{w'}{-\frac{1}{2} - \frac{\sqrt{1-12w}}{2}} dz = \int 1 dz + c_1$$

- Evaluate integral

$$-\frac{\ln(w)}{6} + \frac{\sqrt{1-12w}}{3} + \frac{\ln(-1+\sqrt{1-12w})}{6} - \frac{\ln(1+\sqrt{1-12w})}{6} = z + c_1$$

- Use initial condition $w(1) = -1$

$$-\frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1+\sqrt{13})}{6} - \frac{\ln(1+\sqrt{13})}{6} = c_1 + 1$$

- Solve for c_1

$$c_1 = -1 - \frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1+\sqrt{13})}{6} - \frac{\ln(1+\sqrt{13})}{6}$$

- Substitute $c_1 = -1 - \frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1+\sqrt{13})}{6} - \frac{\ln(1+\sqrt{13})}{6}$ into general solution and simplify

$$-\frac{\ln(w)}{6} + \frac{\sqrt{1-12w}}{3} + \frac{\ln(-1+\sqrt{1-12w})}{6} - \frac{\ln(1+\sqrt{1-12w})}{6} = z - 1 - \frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1+\sqrt{13})}{6} - \frac{\ln(1+\sqrt{13})}{6}$$

- Solution to the IVP

$$-\frac{\ln(w)}{6} + \frac{\sqrt{1-12w}}{3} + \frac{\ln(-1+\sqrt{1-12w})}{6} - \frac{\ln(1+\sqrt{1-12w})}{6} = z - 1 - \frac{i\pi}{6} + \frac{\sqrt{13}}{3} + \frac{\ln(-1+\sqrt{13})}{6} - \frac{\ln(1+\sqrt{13})}{6}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.64 (sec). Leaf size: 66

```
dsolve([diff(w(z),z) = -1/2 - sqrt(1/4 - 3*w(z)),w(1) = -1],w(z), singsol=all)
```

$$w(z) = \text{RootOf} \left(-i\pi + 2\sqrt{13} - 2\sqrt{1 - 12_Z} + \ln(_Z) - \ln(-1 + \sqrt{1 - 12_Z}) \right. \\ \left. + \ln(1 + \sqrt{1 - 12_Z}) - \ln(1 + \sqrt{13}) + \ln(-1 + \sqrt{13}) + 6z - 6 \right)$$

✓ Solution by Mathematica

Time used: 14.307 (sec). Leaf size: 105

```
DSolve[{w'[z] == -1/2 - Sqrt[1/4 - 3*w[z]],{w[1] == -1}},w[z],z,IncludeSingularSolutions ->
```

$$w(z) \rightarrow -\frac{1}{12}W\left(\left(\sqrt{13}-1\right)e^{-3z+\sqrt{13}+2}\right)\left(W\left(\left(\sqrt{13}-1\right)e^{-3z+\sqrt{13}+2}\right)+2\right) \\ w(z) \rightarrow -\frac{1}{12}W\left(\left(\sqrt{13}-1\right)e^{-3z+\sqrt{13}+2}\right)\left(W\left(\left(\sqrt{13}-1\right)e^{-3z+\sqrt{13}+2}\right)+2\right)$$

3.3 problem 3

3.3.1	Existence and uniqueness analysis	1155
3.3.2	Solving as second order linear constant coeff ode	1156
3.3.3	Solving using Kovacic algorithm	1159
3.3.4	Maple step by step solution	1164

Internal problem ID [7193]

Internal file name [OUTPUT/6179_Sunday_June_05_2022_04_26_56_PM_2515758/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y(0) = 1]$$

3.3.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 1$$

$$F = \sin(x)$$

Hence the ode is

$$y'' + y = \sin(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

3.3.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

Substituting these values back in above solution results in

$$y = \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2}$$

Summary

The solution(s) found are the following

$$y = \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Verification of solutions

$$y = \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2}$$

Verified OK.

3.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 114: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

Substituting these values back in above solution results in

$$y = \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2}$$

Summary

The solution(s) found are the following

$$y = \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Verification of solutions

$$y = \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2}$$

Verified OK.

3.3.4 Maple step by step solution

Let's solve

$$[y'' + y = \sin(x), y(0) = 1]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{\sin(x)(2c_2 + 1)}{2} - \frac{\cos(x)(x - 2)}{2}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 20

```
DSolve[{y''[x]+y[x]==Sin[x],{y[0] == 1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}x \cos(x) + \cos(x) + c_2 \sin(x)$$

3.4 problem 4

3.4.1	Existence and uniqueness analysis	1167
3.4.2	Solving as second order linear constant coeff ode	1168
3.4.3	Solving using Kovacic algorithm	1171
3.4.4	Maple step by step solution	1176

Internal problem ID [7194]

Internal file name [OUTPUT/6180_Sunday_June_05_2022_04_26_58_PM_46413825/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y'(0) = 1]$$

3.4.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 1$$

$$F = \sin(x)$$

Hence the ode is

$$y'' + y = \sin(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

3.4.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -\frac{1}{2} + c_2 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_2 = \frac{3}{2}$$

Substituting these values back in above solution results in

$$y = c_1 \cos(x) + \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2} \quad (1)$$

Verification of solutions

$$y = c_1 \cos(x) + \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Verified OK.

3.4.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 116: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -\frac{1}{2} + c_2 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_2 = \frac{3}{2}$$

Substituting these values back in above solution results in

$$y = c_1 \cos(x) + \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2} \quad (1)$$

Verification of solutions

$$y = c_1 \cos(x) + \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Verified OK.

3.4.4 Maple step by step solution

Let's solve

$$\left[y'' + y = \sin(x), y' \Big|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(-x + 2c_1) \cos(x)}{2} + \frac{3 \sin(x)}{2}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 23

```
DSolve[{y''[x]+y[x]==Sin[x],{y'[0] == 1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3 \sin(x)}{2} + \left(-\frac{x}{2} + c_1\right) \cos(x)$$

3.5 problem 5

3.5.1	Existence and uniqueness analysis	1179
3.5.2	Solving as second order linear constant coeff ode	1180
3.5.3	Solving using Kovacic algorithm	1184
3.5.4	Maple step by step solution	1190

Internal problem ID [7195]

Internal file name [OUTPUT/6181_Sunday_June_05_2022_04_27_00_PM_10273806/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y'(0) = 1, y(0) = 0]$$

3.5.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 1$$

$$F = \sin(x)$$

Hence the ode is

$$y'' + y = \sin(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

3.5.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -\frac{1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= 0 \\ c_2 &= \frac{3}{2} \end{aligned}$$


Substituting these values back in above solution results in

$$y = \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Verified OK.

3.5.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 118: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x)(\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -\frac{1}{2} + c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$
$$c_2 = \frac{3}{2}$$


Substituting these values back in above solution results in

$$y = \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Verified OK.

3.5.4 Maple step by step solution

Let's solve

$$\left[y'' + y = \sin(x), y' \Big|_{\{x=0\}} = 1, y(0) = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Check validity of solution $y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$

- Use initial condition $y(0) = 0$

$$0 = c_1$$

- Compute derivative of the solution

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{4} + \frac{x \sin(x)}{2}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 1$

$$1 = -\frac{1}{4} + c_2$$

- Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = \frac{5}{4}\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$

- Solution to the IVP

$$y = \frac{3 \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
    <- solving first the homogeneous part of the ODE successful`

```


✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),D(y)(0) = 1, y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{3 \sin(x)}{2} - \frac{\cos(x) x}{2}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 19

```
DSolve[{y'[x]+y[x]==Sin[x],{y'[0] == 1,y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(3 \sin(x) - x \cos(x))$$

3.6 problem 6

3.6.1	Existence and uniqueness analysis	1193
3.6.2	Solving as second order linear constant coeff ode	1194
3.6.3	Solving using Kovacic algorithm	1197
3.6.4	Maple step by step solution	1202

Internal problem ID [7196]

Internal file name [OUTPUT/6182_Sunday_June_05_2022_04_27_02_PM_45975686/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y(1) = 0]$$

3.6.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 1$$

$$F = \sin(x)$$

Hence the ode is

$$y'' + y = \sin(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

3.6.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = c_1 \cos(1) + c_2 \sin(1) - \frac{\cos(1)}{2} \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\tan(1) c_2 + \frac{1}{2}$$

Substituting these values back in above solution results in

$$y = -\cos(x) \tan(1) c_2 + \frac{\cos(x)}{2} + c_2 \sin(x) - \frac{x \cos(x)}{2}$$

Which simplifies to

$$y = \frac{(-2 \tan(1) c_2 - x + 1) \cos(x)}{2} + c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = \frac{(-2 \tan(1) c_2 - x + 1) \cos(x)}{2} + c_2 \sin(x) \quad (1)$$

Verification of solutions

$$y = \frac{(-2 \tan(1) c_2 - x + 1) \cos(x)}{2} + c_2 \sin(x)$$

Verified OK.

3.6.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 1\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 120: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = c_1 \cos(1) + c_2 \sin(1) - \frac{\cos(1)}{2} \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\tan(1) c_2 + \frac{1}{2}$$

Substituting these values back in above solution results in

$$y = -\cos(x) \tan(1) c_2 + \frac{\cos(x)}{2} + c_2 \sin(x) - \frac{x \cos(x)}{2}$$

Which simplifies to

$$y = \frac{(-2 \tan(1) c_2 - x + 1) \cos(x)}{2} + c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = \frac{(-2 \tan(1) c_2 - x + 1) \cos(x)}{2} + c_2 \sin(x) \quad (1)$$

Verification of solutions

$$y = \frac{(-2 \tan(1) c_2 - x + 1) \cos(x)}{2} + c_2 \sin(x)$$

Verified OK.

3.6.4 Maple step by step solution

Let's solve

$$[y'' + y = \sin(x), y(1) = 0]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 32

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),y(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{((-2c_2 - 1) \tan(1) - x + 1) \cos(x)}{2} + \frac{\sin(x)(2c_2 + 1)}{2}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 18

```
DSolve[{y''[x]+y[x]==Sin[x],{y[0] == 0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}x \cos(x) + c_2 \sin(x)$$

3.7 problem 7

3.7.1	Existence and uniqueness analysis	1205
3.7.2	Solving as second order linear constant coeff ode	1206
3.7.3	Solving using Kovacic algorithm	1210
3.7.4	Maple step by step solution	1215

Internal problem ID [7197]

Internal file name [OUTPUT/6183_Sunday_June_05_2022_04_27_05_PM_9550656/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 7.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y'(1) = 0]$$

3.7.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 1$$

$$F = \sin(x)$$

Hence the ode is

$$y'' + y = \sin(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

3.7.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(2c_2 - 1) \cos(1)}{2} + \frac{(1 - 2c_1) \sin(1)}{2} \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \cot(1) c_2 + \frac{1}{2} - \frac{\cot(1)}{2}$$

Substituting these values back in above solution results in

$$y = \cos(x) \cot(1) c_2 + \frac{\cos(x)}{2} - \frac{\cos(x) \cot(1)}{2} + c_2 \sin(x) - \frac{x \cos(x)}{2}$$

Which simplifies to

$$y = \frac{((2c_2 - 1) \cot(1) - x + 1) \cos(x)}{2} + c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = \frac{((2c_2 - 1) \cot(1) - x + 1) \cos(x)}{2} + c_2 \sin(x) \quad (1)$$

Verification of solutions

$$y = \frac{((2c_2 - 1) \cot(1) - x + 1) \cos(x)}{2} + c_2 \sin(x)$$

Verified OK.

3.7.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 122: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(2c_2 - 1) \cos(1)}{2} + \frac{(1 - 2c_1) \sin(1)}{2} \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \cot(1) c_2 + \frac{1}{2} - \frac{\cot(1)}{2}$$

Substituting these values back in above solution results in

$$y = \cos(x) \cot(1) c_2 + \frac{\cos(x)}{2} - \frac{\cos(x) \cot(1)}{2} + c_2 \sin(x) - \frac{x \cos(x)}{2}$$

Which simplifies to

$$y = \frac{((2c_2 - 1) \cot(1) - x + 1) \cos(x)}{2} + c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = \frac{((2c_2 - 1) \cot(1) - x + 1) \cos(x)}{2} + c_2 \sin(x) \quad (1)$$

Verification of solutions

$$y = \frac{((2c_2 - 1) \cot(1) - x + 1) \cos(x)}{2} + c_2 \sin(x)$$

Verified OK.

3.7.4 Maple step by step solution

Let's solve

$$\left[y'' + y = \sin(x), y' \Big|_{\{x=1\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 27

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),D(y)(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(2 \cot(1) c_2 - x + 1) \cos(x)}{2} + \frac{\sin(x) (2c_2 + 1)}{2}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 35

```
DSolve[{y''[x]+y[x]==Sin[x],{y'[1] == 0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}((1 - \tan(1) + 2c_1 \tan(1)) \sin(x) - (x - 2c_1) \cos(x))$$

3.8 problem 8

3.8.1 Solving as second order linear constant coeff ode	1217
3.8.2 Solving using Kovacic algorithm	1221
3.8.3 Maple step by step solution	1227

Internal problem ID [7198]

Internal file name [OUTPUT/6184_Sunday_June_05_2022_04_27_07_PM_37680447/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y'(1) = 0, y(0) = 0]$$

3.8.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2}\right)\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(2c_2 - 1) \cos(1)}{2} + \frac{(1 - 2c_1) \sin(1)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= 0 \\ c_2 &= \frac{1}{2} - \frac{\tan(1)}{2}\end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{\sin(x)}{2} - \frac{\sin(x) \tan(1)}{2} - \frac{x \cos(x)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\sin(x)}{2} - \frac{\sin(x) \tan(1)}{2} - \frac{x \cos(x)}{2} \quad (1)$$




Figure 119: Solution plot

Verification of solutions

$$y = \frac{\sin(x)}{2} - \frac{\sin(x) \tan(1)}{2} - \frac{x \cos(x)}{2}$$

Verified OK.

3.8.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 124: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x)(\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(2c_2 - 1) \cos(1)}{2} + \frac{(1 - 2c_1) \sin(1)}{2} \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$
$$c_2 = \frac{1}{2} - \frac{\tan(1)}{2}$$

Substituting these values back in above solution results in

$$y = \frac{\sin(x)}{2} - \frac{\sin(x)\tan(1)}{2} - \frac{x\cos(x)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\sin(x)}{2} - \frac{\sin(x)\tan(1)}{2} - \frac{x\cos(x)}{2} \quad (1)$$




Figure 120: Solution plot

Verification of solutions

$$y = \frac{\sin(x)}{2} - \frac{\sin(x)\tan(1)}{2} - \frac{x\cos(x)}{2}$$

Verified OK.

3.8.3 Maple step by step solution

Let's solve

$$\left[y'' + y = \sin(x), y' \Big|_{\{x=1\}} = 0, y(0) = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Check validity of solution $y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$

- Use initial condition $y(0) = 0$

$$0 = c_1$$

- Compute derivative of the solution

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{4} + \frac{x \sin(x)}{2}$$

- Use the initial condition $y' \Big|_{\{x=1\}} = 0$

$$0 = -c_1 \sin(1) + c_2 \cos(1) - \frac{\cos(1)}{4} + \frac{\sin(1)}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = 0, c_2 = \frac{\cos(1) - 2 \sin(1)}{4 \cos(1)} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(1 - \tan(1)) \sin(x)}{2} - \frac{x \cos(x)}{2}$$

- Solution to the IVP

$$y = \frac{(1 - \tan(1)) \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),D(y)(1) = 0, y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(-\tan(1) + 1)\sin(x)}{2} - \frac{\cos(x)x}{2}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 23

```
DSolve[{y'[x]+y[x]==Sin[x],{y'[1] == 0,y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(\sin(x) - x \cos(x) - \tan(1) \sin(x))$$

3.9 problem 9

3.9.1 Solving as second order linear constant coeff ode	1230
3.9.2 Solving using Kovacic algorithm	1234
3.9.3 Maple step by step solution	1240

Internal problem ID [7199]

Internal file name [OUTPUT/6185_Sunday_June_05_2022_04_27_09_PM_39315846/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 9.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y'(1) = 0, y(2) = 0]$$

3.9.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2}\right)
 \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 2$ in the above gives

$$0 = (c_1 - 1) \cos(2) + c_2 \sin(2) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(2c_2 - 1) \cos(1)}{2} + \frac{(1 - 2c_1) \sin(1)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}
 c_1 &= \frac{\cos(2)}{2} + \frac{1}{2} - \frac{\sin(2)}{2} \\
 c_2 &= \frac{\sec(1) \cos(2) (\sin(1) + \cos(1))}{2}
 \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{\cos(x) \cos(2)}{2} + \frac{\cos(x)}{2} - \frac{\cos(x) \sin(2)}{2} + \frac{\sin(x) \sec(1) \cos(2) \cos(1)}{2} + \frac{\sin(x) \sec(1) \cos(2) \sin(1)}{2}$$

Which simplifies to

$$y = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x) (\sin(2) - \tan(1) + \cos(2))}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x) (\sin(2) - \tan(1) + \cos(2))}{2} \quad (1)$$




Figure 121: Solution plot

Verification of solutions

$$y = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x) (\sin(2) - \tan(1) + \cos(2))}{2}$$

Verified OK.

3.9.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 126: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 2$ in the above gives

$$0 = (c_1 - 1) \cos(2) + c_2 \sin(2) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(2c_2 - 1) \cos(1)}{2} + \frac{(1 - 2c_1) \sin(1)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{\cos(2)}{2} + \frac{1}{2} - \frac{\sin(2)}{2}$$

$$c_2 = \frac{\sec(1) \cos(2) (\sin(1) + \cos(1))}{2}$$

Substituting these values back in above solution results in

$$y = \frac{\cos(x) \cos(2)}{2} + \frac{\cos(x)}{2} - \frac{\cos(x) \sin(2)}{2} + \frac{\sin(x) \sec(1) \cos(2) \cos(1)}{2} + \frac{\sin(x) \sec(1) \cos(2) \sin(1)}{2}$$

Which simplifies to

$$y = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x) (\sin(2) - \tan(1) + \cos(2))}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x) (\sin(2) - \tan(1) + \cos(2))}{2} \quad (1)$$




Figure 122: Solution plot

Verification of solutions

$$y = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x) (\sin(2) - \tan(1) + \cos(2))}{2}$$

Verified OK.

3.9.3 Maple step by step solution

Let's solve

$$\left[y'' + y = \sin(x), y' \Big|_{\{x=1\}} = 0, y(2) = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE
 $y_1(x) = \cos(x)$
 - 2nd solution of the homogeneous ODE
 $y_2(x) = \sin(x)$
 - General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$
 - Substitute in solutions of the homogeneous ODE
 $y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$
- Find a particular solution $y_p(x)$ of the ODE
- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
 - Compute Wronskian
 $W(y_1(x), y_2(x)) = 1$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$
 - Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$
 - Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$
- Check validity of solution $y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$
- Use initial condition $y(2) = 0$
 $0 = c_1 \cos(2) + c_2 \sin(2) + \frac{\sin(2)}{4} - \cos(2)$
 - Compute derivative of the solution

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{4} + \frac{x \sin(x)}{2}$$
 - Use the initial condition $y' \Big|_{\{x=1\}} = 0$

$$0 = -c_1 \sin(1) + c_2 \cos(1) - \frac{\cos(1)}{4} + \frac{\sin(1)}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{2 \cos(1) \cos(2) - \sin(2) \cos(1) + \sin(1) \sin(2)}{2(\cos(1) \cos(2) + \sin(1) \sin(2))}, c_2 = \frac{\cos(1) \cos(2) + 2 \cos(2) \sin(1) - \sin(1) \sin(2)}{4(\cos(1) \cos(2) + \sin(1) \sin(2))} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x)(\sin(2) - \tan(1) + \cos(2))}{2}$$

- Solution to the IVP

$$y = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x)(\sin(2) - \tan(1) + \cos(2))}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 33

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),D(y)(1) = 0, y(2) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(-x + \cos(2) - \sin(2) + 1) \cos(x)}{2} + \frac{\sin(x)(\sin(2) - \tan(1) + \cos(2))}{2}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 39

```
DSolve[{y''[x]+y[x]==Sin[x],{y'[1] == 0,y[2]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(\sec(1) \sin(x)(-\sin(1) + \sin(3) + \cos(1) + \cos(3)) - 2 \cos(x)(x - 1 + \sin(2) - \cos(2)))$$

3.10 problem 10

3.10.1 Solving as second order linear constant coeff ode	1243
3.10.2 Solving using Kovacic algorithm	1247
3.10.3 Maple step by step solution	1253

Internal problem ID [7200]

Internal file name [OUTPUT/6186_Sunday_June_05_2022_04_27_11_PM_45422128/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 10.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y'(1) = 0, y(0) = 0]$$

3.10.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2}\right)\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(2c_2 - 1) \cos(1)}{2} + \frac{(1 - 2c_1) \sin(1)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= 0 \\ c_2 &= \frac{1}{2} - \frac{\tan(1)}{2}\end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{\sin(x)}{2} - \frac{\sin(x) \tan(1)}{2} - \frac{x \cos(x)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\sin(x)}{2} - \frac{\sin(x) \tan(1)}{2} - \frac{x \cos(x)}{2} \quad (1)$$




Figure 123: Solution plot

Verification of solutions

$$y = \frac{\sin(x)}{2} - \frac{\sin(x) \tan(1)}{2} - \frac{x \cos(x)}{2}$$

Verified OK.

3.10.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 128: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x)(\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(x), x \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(x) + A_2 x \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{x \cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{x \cos(x)}{2} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{2} + \frac{x \sin(x)}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(2c_2 - 1) \cos(1)}{2} + \frac{(1 - 2c_1) \sin(1)}{2} \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$
$$c_2 = \frac{1}{2} - \frac{\tan(1)}{2}$$

Substituting these values back in above solution results in

$$y = \frac{\sin(x)}{2} - \frac{\sin(x)\tan(1)}{2} - \frac{x\cos(x)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\sin(x)}{2} - \frac{\sin(x)\tan(1)}{2} - \frac{x\cos(x)}{2} \tag{1}$$




Figure 124: Solution plot

Verification of solutions

$$y = \frac{\sin(x)}{2} - \frac{\sin(x)\tan(1)}{2} - \frac{x\cos(x)}{2}$$

Verified OK.

3.10.3 Maple step by step solution

Let's solve

$$\left[y'' + y = \sin(x), y' \Big|_{\{x=1\}} = 0, y(0) = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

- Check validity of solution $y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$

- Use initial condition $y(0) = 0$

$$0 = c_1$$

- Compute derivative of the solution

$$y' = -c_1 \sin(x) + c_2 \cos(x) - \frac{\cos(x)}{4} + \frac{x \sin(x)}{2}$$

- Use the initial condition $y' \Big|_{\{x=1\}} = 0$

$$0 = -c_1 \sin(1) + c_2 \cos(1) - \frac{\cos(1)}{4} + \frac{\sin(1)}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = 0, c_2 = \frac{\cos(1) - 2 \sin(1)}{4 \cos(1)} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(-\tan(1)+1) \sin(x)}{2} - \frac{x \cos(x)}{2}$$

- Solution to the IVP

$$y = \frac{(-\tan(1)+1) \sin(x)}{2} - \frac{x \cos(x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),D(y)(1) = 0, y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(-\tan(1) + 1)\sin(x)}{2} - \frac{\cos(x)x}{2}$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 23

```
DSolve[{y'[x]+y[x]==Sin[x],{y'[1] == 0,y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(\sin(x) - x \cos(x) - \tan(1) \sin(x))$$

3.11 problem 11

3.11.1 Solving as second order linear constant coeff ode	1256
3.11.2 Solving using Kovacic algorithm	1261
3.11.3 Maple step by step solution	1267

Internal problem ID [7201]

Internal file name [OUTPUT/6187_Sunday_June_05_2022_04_27_13_PM_30427453/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 11.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y' + y = \sin(x)$$

With initial conditions

$$[y'(1) = 0, y(2) = 0]$$

3.11.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \right) + (-\cos(x))$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) - \cos(x) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 2$ in the above gives

$$0 = e^{-1} \cos(\sqrt{3}) c_1 + e^{-1} \sin(\sqrt{3}) c_2 - \cos(2) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)}{2} + e^{-\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} + \frac{c_2 \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right)}{2} \right) + \sin(x)$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = -\frac{e^{-\frac{1}{2}} (-\sqrt{3} c_2 + c_1) \cos \left(\frac{\sqrt{3}}{2} \right)}{2} - \frac{e^{-\frac{1}{2}} (c_1 \sqrt{3} + c_2) \sin \left(\frac{\sqrt{3}}{2} \right)}{2} + \sin(1) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{\left(\cos \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{1}{2}} \cos(2) - \sin \left(\frac{\sqrt{3}}{2} \right) e^{\frac{1}{2}} \cos(2) + 2 \sin(\sqrt{3}) \sin(1) \right) e^{\frac{1}{2}}}{\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right)}$$

$$c_2 = \frac{\left(\sin \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{1}{2}} \cos(2) + \cos \left(\frac{\sqrt{3}}{2} \right) e^{\frac{1}{2}} \cos(2) - 2 \cos(\sqrt{3}) \sin(1) \right) e^{\frac{1}{2}}}{\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right)}$$

Substituting these values back in above solution results in

$$y = \frac{-2 \sin \left(\frac{\sqrt{3}x}{2} \right) \cos \left(\frac{\sqrt{3}}{2} \right) \sin(1) \sqrt{3} e^{-\frac{x}{2} + \frac{1}{2}} \cos(\sqrt{3}) + 2 \cos \left(\frac{\sqrt{3}}{2} \right) \sin(1) \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2} + \frac{1}{2}} \sin(\sqrt{3})}{\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right)}$$

Which simplifies to

y

$$-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1+\sqrt{3}\sin(\sqrt{3})-\cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right)+\sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right)+\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)\right)$$

Summary

The solution(s) found are the following

y

(1)

$$-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1+\sqrt{3}\sin(\sqrt{3})-\cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right)+\sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right)+\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)\right)$$




Figure 125: Solution plot

Verification of solutions

y

$$-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1+\sqrt{3}\sin(\sqrt{3})-\cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right)+\sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right)+\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)\right)$$

Verified OK.

3.11.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \quad (3)$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = -3$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 130: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\&= z_1 e^{-\frac{x}{2}} \\&= z_1 \left(e^{-\frac{x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3}x}{2} \right)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3}x}{2} \right)}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) + \frac{2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}} \sqrt{3}}{3} \right) + (-\cos(x))$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) + \frac{2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}} \sqrt{3}}{3} - \cos(x) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 2$ in the above gives

$$0 = e^{-1} \cos(\sqrt{3}) c_1 + \frac{2e^{-1} \sin(\sqrt{3}) c_2 \sqrt{3}}{3} - \cos(2) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right)}{2} - \frac{c_1 e^{-\frac{x}{2}} \sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} + c_2 \cos \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}} - \frac{c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}} \sqrt{3}}{3} + \sin(x)$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = -\frac{e^{-\frac{1}{2}}(c_1 - 2c_2) \cos \left(\frac{\sqrt{3}}{2} \right)}{2} - \frac{\sqrt{3} (c_1 + \frac{2c_2}{3}) e^{-\frac{1}{2}} \sin \left(\frac{\sqrt{3}}{2} \right)}{2} + \sin(1) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{\left(-\sin \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{1}{2}} \cos(2) + 3 \cos \left(\frac{\sqrt{3}}{2} \right) e^{\frac{1}{2}} \cos(2) + 2 \sin(1) \sqrt{3} \sin(\sqrt{3}) \right) e^{\frac{1}{2}}}{\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + 3 \cos \left(\frac{\sqrt{3}}{2} \right)}$$

$$c_2 = \frac{3 \left(\sin \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{1}{2}} \cos(2) + \cos \left(\frac{\sqrt{3}}{2} \right) e^{\frac{1}{2}} \cos(2) - 2 \cos(\sqrt{3}) \sin(1) \right) e^{\frac{1}{2}}}{2 \left(\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + 3 \cos \left(\frac{\sqrt{3}}{2} \right) \right)}$$

Substituting these values back in above solution results in

$$y = \frac{-2 \sin \left(\frac{\sqrt{3}x}{2} \right) \cos \left(\frac{\sqrt{3}}{2} \right) \sin(1) \sqrt{3} e^{-\frac{x}{2} + \frac{1}{2}} \cos(\sqrt{3}) + 2 \cos \left(\frac{\sqrt{3}}{2} \right) \sin(1) \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2} + \frac{1}{2}} \sin(\sqrt{3})}{2 \left(\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + 3 \cos \left(\frac{\sqrt{3}}{2} \right) \right)}$$

Which simplifies to

$$y = -2 \left(-\cos \left(\frac{\sqrt{3}}{2} \right) (1 + \sqrt{3} \sin(\sqrt{3}) - \cos(\sqrt{3})) \cos \left(\frac{\sqrt{3}x}{2} \right) + \sin \left(\frac{\sqrt{3}x}{2} \right) \cos(\sqrt{3}) \left(\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right) \right) \right)$$

Summary

The solution(s) found are the following

$$y = -2 \left(-\cos \left(\frac{\sqrt{3}}{2} \right) (1 + \sqrt{3} \sin(\sqrt{3}) - \cos(\sqrt{3})) \cos \left(\frac{\sqrt{3}x}{2} \right) + \sin \left(\frac{\sqrt{3}x}{2} \right) \cos(\sqrt{3}) \left(\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right) \right) \right) \tag{1}$$




Figure 126: Solution plot

Verification of solutions

$$y = -2 \left(-\cos \left(\frac{\sqrt{3}}{2} \right) (1 + \sqrt{3} \sin(\sqrt{3}) - \cos(\sqrt{3})) \cos \left(\frac{\sqrt{3}x}{2} \right) + \sin \left(\frac{\sqrt{3}x}{2} \right) \cos(\sqrt{3}) \left(\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right) \right) \right)$$

Verified OK.

3.11.3 Maple step by step solution

Let's solve

$$\left[y'' + y' + y = \sin(x), y'|_{\{x=1\}} = 0, y(2) = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{2e^{-\frac{x}{2}}\sqrt{3}\left(-\cos\left(\frac{\sqrt{3}x}{2}\right)\left(\int e^{\frac{x}{2}}\sin(x)\sin\left(\frac{\sqrt{3}x}{2}\right)dx\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)\left(\int e^{\frac{x}{2}}\sin(x)\cos\left(\frac{\sqrt{3}x}{2}\right)dx\right)\right)}{3}$$

- Compute integrals

$$y_p(x) = -\cos(x)$$

- Substitute particular solution into general solution to ODE

$$y = c_1e^{-\frac{x}{2}}\cos\left(\frac{\sqrt{3}x}{2}\right) + c_2\sin\left(\frac{\sqrt{3}x}{2}\right)e^{-\frac{x}{2}} - \cos(x)$$

- Check validity of solution $y = c_1e^{-\frac{x}{2}}\cos\left(\frac{\sqrt{3}x}{2}\right) + c_2\sin\left(\frac{\sqrt{3}x}{2}\right)e^{-\frac{x}{2}} - \cos(x)$

- Use initial condition $y(2) = 0$

$$0 = e^{-1}\cos(\sqrt{3})c_1 + e^{-1}\sin(\sqrt{3})c_2 - \cos(2)$$

- Compute derivative of the solution

$$y' = -\frac{c_1e^{-\frac{x}{2}}\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{c_1e^{-\frac{x}{2}}\sqrt{3}\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}}\sqrt{3}\cos\left(\frac{\sqrt{3}x}{2}\right)c_2}{2} - \frac{c_2\sin\left(\frac{\sqrt{3}x}{2}\right)e^{-\frac{x}{2}}}{2} + \sin(x)$$

- Use the initial condition $y'|_{\{x=1\}} = 0$

$$0 = -\frac{e^{-\frac{1}{2}}c_1\sin\left(\frac{\sqrt{3}}{2}\right)\sqrt{3}}{2} - \frac{e^{-\frac{1}{2}}c_1\cos\left(\frac{\sqrt{3}}{2}\right)}{2} + \frac{c_2e^{-\frac{1}{2}}\cos\left(\frac{\sqrt{3}}{2}\right)\sqrt{3}}{2} - \frac{c_2e^{-\frac{1}{2}}\sin\left(\frac{\sqrt{3}}{2}\right)}{2} + \sin(1)$$

- Solve for c_1 and c_2

$$\begin{cases} c_1 = \frac{\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)e^{-\frac{1}{2}}\cos(2) + 2\sin(1)e^{-1}\sin(\sqrt{3}) - \sin\left(\frac{\sqrt{3}}{2}\right)e^{-\frac{1}{2}}\cos(2)}{e^{-\frac{1}{2}}\left(\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\sqrt{3}\sin\left(\frac{\sqrt{3}}{2}\right) - \sin\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\cos\left(\frac{\sqrt{3}}{2}\right)\right)e^{-1}}, c_2 = -\frac{e^{-\frac{1}{2}}\left(\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\sqrt{3}\sin\left(\frac{\sqrt{3}}{2}\right) - \sin\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\cos\left(\frac{\sqrt{3}}{2}\right)\right)e^{-1}}{e^{-\frac{1}{2}}\left(\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\sqrt{3}\sin\left(\frac{\sqrt{3}}{2}\right) - \sin\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\cos\left(\frac{\sqrt{3}}{2}\right)\right)e^{-1}} \end{cases}$$

- Substitute constant values into general solution and simplify

$$y = \frac{2\sin(1)\left(\cos\left(\frac{\sqrt{3}x}{2}\right)\sin(\sqrt{3}) - \sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\right)e^{-\frac{x}{2} + \frac{1}{2}} - \cos(2)\left(\cos\left(\frac{\sqrt{3}x}{2}\right)\left(-\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right)\right) - \sin\left(\frac{\sqrt{3}x}{2}\right)\left(\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right)\right)\right)e^{-\frac{x}{2}}}{\sin\left(\frac{\sqrt{3}}{2}\right) + \sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)}$$

- Solution to the IVP

$$y = \frac{2\sin(1)\left(\cos\left(\frac{\sqrt{3}x}{2}\right)\sin(\sqrt{3}) - \sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\right)e^{-\frac{x}{2} + \frac{1}{2}} - \cos(2)\left(\cos\left(\frac{\sqrt{3}x}{2}\right)\left(-\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right)\right) - \sin\left(\frac{\sqrt{3}x}{2}\right)\left(\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right)\right)\right)e^{-\frac{x}{2}}}{\sin\left(\frac{\sqrt{3}}{2}\right) + \sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.469 (sec). Leaf size: 144

```
dsolve([diff(y(x),x$2)+diff(y(x),x)+y(x)=sin(x),D(y)(1) = 0, y(2) = 0],y(x), singsol=all)
```

$$y(x) = \frac{2 \sin(1) \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \sin(\sqrt{3}) - \sin\left(\frac{\sqrt{3}x}{2}\right) \cos(\sqrt{3}) \right) e^{\frac{1}{2} - \frac{x}{2}} - \cos(2) \left(\left(-\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right) \right) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \right)}{\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right)}$$

✓ Solution by Mathematica

Time used: 1.065 (sec). Leaf size: 12765

```
DSolve[{y'''[x]+y'[x]+y[x]==Sin[x],{y'[1] == 0,y[2]==0}},y[x],x,IncludeSingularSolutions ->
```

Too large to display

3.12 problem 12

3.12.1 Existence and uniqueness analysis	1270
3.12.2 Solving as second order linear constant coeff ode	1271
3.12.3 Solving using Kovacic algorithm	1275
3.12.4 Maple step by step solution	1280

Internal problem ID [7202]

Internal file name [OUTPUT/6188_Sunday_June_05_2022_04_27_17_PM_16711888/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 12.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y' + y = \sin(x)$$

With initial conditions

$$[y'(1) = 0]$$

3.12.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = 1$$

$$F = \sin(x)$$

Hence the ode is

$$y'' + y' + y = \sin(x)$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

3.12.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \right) + (-\cos(x)) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) - \cos(x) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. Taking derivative of the solution gives

$$y' = -\frac{e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)}{2} + e^{-\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} + \frac{c_2 \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right)}{2} \right) + \sin(x)$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = -\frac{e^{-\frac{1}{2}} (-\sqrt{3}c_2 + c_1) \cos \left(\frac{\sqrt{3}}{2} \right)}{2} - \frac{e^{-\frac{1}{2}} (c_1 \sqrt{3} + c_2) \sin \left(\frac{\sqrt{3}}{2} \right)}{2} + \sin(1) \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{c_2 \cos \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} + 2 \sin(1) e^{\frac{1}{2}} - c_2 \sin \left(\frac{\sqrt{3}}{2} \right)}{\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + \cos \left(\frac{\sqrt{3}}{2} \right)}$$

Substituting these values back in above solution results in

$$y = \frac{\sin \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) c_2 + \cos \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) c_2 - \sin \left(\frac{\sqrt{3}}{2} \right) e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) c_2 + \cos \left(\frac{\sqrt{3}}{2} \right) e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) c_2}{\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + \cos \left(\frac{\sqrt{3}}{2} \right)} - \cos(x)$$

Which simplifies to

$$y = \frac{2 \sin(1) \cos \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2} + \frac{1}{2}} + \left(\sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right) - \sin \left(\frac{\sqrt{3}}{2} \right) \right) e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) c_2 + \left(\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + \cos \left(\frac{\sqrt{3}}{2} \right) \right) e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) c_2}{\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + \cos \left(\frac{\sqrt{3}}{2} \right)} - \cos(x)$$

Summary

The solution(s) found are the following

$$y = \frac{2 \sin(1) \cos \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2} + \frac{1}{2}} + \left(\sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right) - \sin \left(\frac{\sqrt{3}}{2} \right) \right) e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) c_2 + \left(\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + \cos \left(\frac{\sqrt{3}}{2} \right) \right) e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) c_2}{\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + \cos \left(\frac{\sqrt{3}}{2} \right)} - \cos(x) \quad (1)$$

Verification of solutions

$$y = \frac{2 \sin(1) \cos\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2} + \frac{1}{2}} + \left(\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) - \sin\left(\frac{\sqrt{3}}{2}\right)\right) e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_2 + \left(\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) + \cos\left(\frac{\sqrt{3}}{2}\right)\right) e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) + \cos\left(\frac{\sqrt{3}}{2}\right)}$$

Verified OK.

3.12.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \quad (3)$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = -3$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 132: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} \right) + (-\cos(x)) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} - \cos(x) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. Taking derivative of the solution gives

$$y' = -\frac{c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{c_1 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} - \frac{c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} + \sin(x)$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = -\frac{e^{-\frac{1}{2}}(c_1 - 2c_2) \cos\left(\frac{\sqrt{3}}{2}\right)}{2} - \frac{\sqrt{3}(c_1 + \frac{2c_2}{3}) e^{-\frac{1}{2}} \sin\left(\frac{\sqrt{3}}{2}\right)}{2} + \sin(1) \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{2\left(c_2 \sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) - 3 \sin(1) e^{\frac{1}{2}} - 3 \cos\left(\frac{\sqrt{3}}{2}\right) c_2\right)}{3\left(\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) + \cos\left(\frac{\sqrt{3}}{2}\right)\right)}$$

Substituting these values back in above solution results in

$$y = \frac{-2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \sin\left(\frac{\sqrt{3}}{2}\right) \sqrt{3}c_2 + 2 \sin\left(\frac{\sqrt{3}x}{2}\right) \cos\left(\frac{\sqrt{3}}{2}\right) \sqrt{3}e^{-\frac{x}{2}}c_2 + 6e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \cos\left(\frac{\sqrt{3}}{2}\right)c_2 + \dots}{3\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right)}$$

Which simplifies to

$$y = \frac{6 \sin(1) \cos\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2} + \frac{1}{2}} - 2c_2 e^{-\frac{x}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) - 3 \cos\left(\frac{\sqrt{3}}{2}\right)\right) \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 e^{-\frac{x}{2}} \left(\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \dots\right)}{3\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) + 3 \cos\left(\frac{\sqrt{3}}{2}\right)}$$

Summary

The solution(s) found are the following

$$y = \frac{6 \sin(1) \cos\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2} + \frac{1}{2}} - 2c_2 e^{-\frac{x}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) - 3 \cos\left(\frac{\sqrt{3}}{2}\right)\right) \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 e^{-\frac{x}{2}} \left(\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \dots\right)}{3\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) + 3 \cos\left(\frac{\sqrt{3}}{2}\right)} \quad (1)$$

Verification of solutions

$$y = \frac{6 \sin(1) \cos\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2} + \frac{1}{2}} - 2c_2 e^{-\frac{x}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) - 3 \cos\left(\frac{\sqrt{3}}{2}\right)\right) \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 e^{-\frac{x}{2}} \left(\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \dots\right)}{3\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) + 3 \cos\left(\frac{\sqrt{3}}{2}\right)}$$

Verified OK.

3.12.4 Maple step by step solution

Let's solve

$$\left[y'' + y' + y = \sin(x), y' \Big|_{\{x=1\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{2e^{-\frac{x}{2}}\sqrt{3} \left(-\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin(x) \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin(x) \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = -\cos(x)$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} - \cos(x)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 80

```
dsolve([diff(y(x),x$2)+diff(y(x),x)+y(x)=sin(x),D(y)(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{2 \sin(1) e^{\frac{1}{2} - \frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 e^{-\frac{x}{2}} \left(\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) - \sin\left(\frac{\sqrt{3}}{2}\right)\right) \cos\left(\frac{\sqrt{3}x}{2}\right) + \left(\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) + \cos\left(\frac{\sqrt{3}}{2}\right)\right)}{\sqrt{3} \sin\left(\frac{\sqrt{3}}{2}\right) + \cos\left(\frac{\sqrt{3}}{2}\right)}$$

✓ Solution by Mathematica

Time used: 0.346 (sec). Leaf size: 4176

```
DSolve[{y'''[x]+y'[x]+y[x]==Sin[x],{y'[1] == 0}},y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

3.13 problem 13

3.13.1 Solving as second order linear constant coeff ode	1283
3.13.2 Solving using Kovacic algorithm	1288
3.13.3 Maple step by step solution	1294

Internal problem ID [7203]

Internal file name [OUTPUT/6189_Sunday_June_05_2022_04_27_23_PM_60745131/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 13.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y' + y = \sin(x)$$

With initial conditions

$$[y'(1) = 0, y(2) = 0]$$

3.13.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \right) + (-\cos(x))
 \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) - \cos(x) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 2$ in the above gives

$$0 = e^{-1} \cos(\sqrt{3}) c_1 + e^{-1} \sin(\sqrt{3}) c_2 - \cos(2) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)}{2} + e^{-\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} + \frac{c_2 \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right)}{2} \right) + \sin(x)$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = -\frac{e^{-\frac{1}{2}} (-\sqrt{3} c_2 + c_1) \cos \left(\frac{\sqrt{3}}{2} \right)}{2} - \frac{e^{-\frac{1}{2}} (c_1 \sqrt{3} + c_2) \sin \left(\frac{\sqrt{3}}{2} \right)}{2} + \sin(1) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}
 c_1 &= \frac{\left(\cos \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{1}{2}} \cos(2) - \sin \left(\frac{\sqrt{3}}{2} \right) e^{\frac{1}{2}} \cos(2) + 2 \sin(\sqrt{3}) \sin(1) \right) e^{\frac{1}{2}}}{\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right)} \\
 c_2 &= \frac{\left(\sin \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{1}{2}} \cos(2) + \cos \left(\frac{\sqrt{3}}{2} \right) e^{\frac{1}{2}} \cos(2) - 2 \cos(\sqrt{3}) \sin(1) \right) e^{\frac{1}{2}}}{\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right)}
 \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{-2 \sin \left(\frac{\sqrt{3}x}{2} \right) \cos \left(\frac{\sqrt{3}}{2} \right) \sin(1) \sqrt{3} e^{-\frac{x}{2} + \frac{1}{2}} \cos(\sqrt{3}) + 2 \cos \left(\frac{\sqrt{3}}{2} \right) \sin(1) \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2} + \frac{1}{2}} \sin(\sqrt{3})}{\sin \left(\frac{\sqrt{3}}{2} \right) + \sqrt{3} \cos \left(\frac{\sqrt{3}}{2} \right)}$$

Which simplifies to

y

$$-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1+\sqrt{3}\sin(\sqrt{3})-\cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right)+\sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right)+\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)\right)$$

Summary

The solution(s) found are the following

y

(1)

$$-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1+\sqrt{3}\sin(\sqrt{3})-\cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right)+\sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right)+\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)\right)$$




Figure 127: Solution plot

Verification of solutions

y

$$-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1+\sqrt{3}\sin(\sqrt{3})-\cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right)+\sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right)+\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)\right)$$

Verified OK.

3.13.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \quad (3)$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = -3$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 134: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\&= z_1 e^{-\frac{x}{2}} \\&= z_1 \left(e^{-\frac{x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3}x}{2} \right)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3}x}{2} \right)}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) + \frac{2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}} \sqrt{3}}{3} \right) + (-\cos(x))$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) + \frac{2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}} \sqrt{3}}{3} - \cos(x) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 2$ in the above gives

$$0 = e^{-1} \cos(\sqrt{3}) c_1 + \frac{2e^{-1} \sin(\sqrt{3}) c_2 \sqrt{3}}{3} - \cos(2) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right)}{2} - \frac{c_1 e^{-\frac{x}{2}} \sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} + c_2 \cos \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}} - \frac{c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2}} \sqrt{3}}{3} + \sin(x)$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = -\frac{e^{-\frac{1}{2}}(c_1 - 2c_2) \cos \left(\frac{\sqrt{3}}{2} \right)}{2} - \frac{\sqrt{3} (c_1 + \frac{2c_2}{3}) e^{-\frac{1}{2}} \sin \left(\frac{\sqrt{3}}{2} \right)}{2} + \sin(1) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{\left(-\sin \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{1}{2}} \cos(2) + 3 \cos \left(\frac{\sqrt{3}}{2} \right) e^{\frac{1}{2}} \cos(2) + 2 \sin(1) \sqrt{3} \sin(\sqrt{3}) \right) e^{\frac{1}{2}}}{\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + 3 \cos \left(\frac{\sqrt{3}}{2} \right)}$$

$$c_2 = \frac{3 \left(\sin \left(\frac{\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{1}{2}} \cos(2) + \cos \left(\frac{\sqrt{3}}{2} \right) e^{\frac{1}{2}} \cos(2) - 2 \cos(\sqrt{3}) \sin(1) \right) e^{\frac{1}{2}}}{2 \left(\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + 3 \cos \left(\frac{\sqrt{3}}{2} \right) \right)}$$

Substituting these values back in above solution results in

$$y = \frac{-2 \sin \left(\frac{\sqrt{3}x}{2} \right) \cos \left(\frac{\sqrt{3}}{2} \right) \sin(1) \sqrt{3} e^{-\frac{x}{2} + \frac{1}{2}} \cos(\sqrt{3}) + 2 \cos \left(\frac{\sqrt{3}}{2} \right) \sin(1) \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right) e^{-\frac{x}{2} + \frac{1}{2}} \sin(\sqrt{3})}{2 \left(\sqrt{3} \sin \left(\frac{\sqrt{3}}{2} \right) + 3 \cos \left(\frac{\sqrt{3}}{2} \right) \right)}$$

Which simplifies to

$$y = \frac{-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1 + \sqrt{3}\sin(\sqrt{3}) - \cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right) + \sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)}{=}$$

Summary

The solution(s) found are the following

$$y = \frac{-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1 + \sqrt{3}\sin(\sqrt{3}) - \cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right) + \sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)}{=} \tag{1}$$




Figure 128: Solution plot

Verification of solutions

$$y = \frac{-2\left(-\cos\left(\frac{\sqrt{3}}{2}\right)\left(1 + \sqrt{3}\sin(\sqrt{3}) - \cos(\sqrt{3})\right)\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\left(\sin\left(\frac{\sqrt{3}}{2}\right) + \sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\right)}{=}$$

Verified OK.

3.13.3 Maple step by step solution

Let's solve

$$\left[y'' + y' + y = \sin(x), y'|_{\{x=1\}} = 0, y(2) = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{2e^{-\frac{x}{2}}\sqrt{3}\left(-\cos\left(\frac{\sqrt{3}x}{2}\right)\left(\int e^{\frac{x}{2}}\sin(x)\sin\left(\frac{\sqrt{3}x}{2}\right)dx\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)\left(\int e^{\frac{x}{2}}\sin(x)\cos\left(\frac{\sqrt{3}x}{2}\right)dx\right)\right)}{3}$$

- Compute integrals

$$y_p(x) = -\cos(x)$$

- Substitute particular solution into general solution to ODE

$$y = c_1e^{-\frac{x}{2}}\cos\left(\frac{\sqrt{3}x}{2}\right) + c_2\sin\left(\frac{\sqrt{3}x}{2}\right)e^{-\frac{x}{2}} - \cos(x)$$

- Check validity of solution $y = c_1e^{-\frac{x}{2}}\cos\left(\frac{\sqrt{3}x}{2}\right) + c_2\sin\left(\frac{\sqrt{3}x}{2}\right)e^{-\frac{x}{2}} - \cos(x)$

- Use initial condition $y(2) = 0$

$$0 = e^{-1}\cos(\sqrt{3})c_1 + e^{-1}\sin(\sqrt{3})c_2 - \cos(2)$$

- Compute derivative of the solution

$$y' = -\frac{c_1e^{-\frac{x}{2}}\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{c_1e^{-\frac{x}{2}}\sqrt{3}\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}}\sqrt{3}\cos\left(\frac{\sqrt{3}x}{2}\right)c_2}{2} - \frac{c_2\sin\left(\frac{\sqrt{3}x}{2}\right)e^{-\frac{x}{2}}}{2} + \sin(x)$$

- Use the initial condition $y'|_{\{x=1\}} = 0$

$$0 = -\frac{e^{-\frac{1}{2}}c_1\sin\left(\frac{\sqrt{3}}{2}\right)\sqrt{3}}{2} - \frac{e^{-\frac{1}{2}}c_1\cos\left(\frac{\sqrt{3}}{2}\right)}{2} + \frac{c_2e^{-\frac{1}{2}}\cos\left(\frac{\sqrt{3}}{2}\right)\sqrt{3}}{2} - \frac{c_2e^{-\frac{1}{2}}\sin\left(\frac{\sqrt{3}}{2}\right)}{2} + \sin(1)$$

- Solve for c_1 and c_2

$$\begin{cases} c_1 = \frac{\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)e^{-\frac{1}{2}}\cos(2) + 2\sin(1)e^{-1}\sin(\sqrt{3}) - \sin\left(\frac{\sqrt{3}}{2}\right)e^{-\frac{1}{2}}\cos(2)}{e^{-\frac{1}{2}}\left(\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\sqrt{3}\sin\left(\frac{\sqrt{3}}{2}\right) - \sin\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\cos\left(\frac{\sqrt{3}}{2}\right)\right)e^{-1}}, c_2 = -\frac{e^{-\frac{1}{2}}\left(\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\sqrt{3}\sin\left(\frac{\sqrt{3}}{2}\right) - \sin\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\cos\left(\frac{\sqrt{3}}{2}\right)\right)e^{-1}}{e^{-\frac{1}{2}}\left(\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\sqrt{3}\sin\left(\frac{\sqrt{3}}{2}\right) - \sin\left(\frac{\sqrt{3}}{2}\right)\cos(\sqrt{3}) + \sin(\sqrt{3})\cos\left(\frac{\sqrt{3}}{2}\right)\right)e^{-1}} \end{cases}$$

- Substitute constant values into general solution and simplify

$$y = \frac{2\sin(1)\left(\cos\left(\frac{\sqrt{3}x}{2}\right)\sin(\sqrt{3}) - \sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\right)e^{-\frac{x}{2} + \frac{1}{2}} - \left(\left(-\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right)\right)\cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right)\left(\sqrt{3}\sin\left(\frac{\sqrt{3}}{2}\right) - \cos\left(\frac{\sqrt{3}}{2}\right)\right)\right)e^{-\frac{x}{2}}}{\sin\left(\frac{\sqrt{3}}{2}\right) + \sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)}$$

- Solution to the IVP

$$y = \frac{2\sin(1)\left(\cos\left(\frac{\sqrt{3}x}{2}\right)\sin(\sqrt{3}) - \sin\left(\frac{\sqrt{3}x}{2}\right)\cos(\sqrt{3})\right)e^{-\frac{x}{2} + \frac{1}{2}} - \left(\left(-\sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right)\right)\cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right)\left(\sqrt{3}\sin\left(\frac{\sqrt{3}}{2}\right) - \cos\left(\frac{\sqrt{3}}{2}\right)\right)\right)e^{-\frac{x}{2}}}{\sin\left(\frac{\sqrt{3}}{2}\right) + \sqrt{3}\cos\left(\frac{\sqrt{3}}{2}\right)}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 144

```
dsolve([diff(y(x),x$2)+diff(y(x),x)+y(x)=sin(x),D(y)(1) = 0, y(2) = 0],y(x), singsol=all)
```

$$y(x) = \frac{2 \sin(1) \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \sin(\sqrt{3}) - \sin\left(\frac{\sqrt{3}x}{2}\right) \cos(\sqrt{3}) \right) e^{\frac{1}{2} - \frac{x}{2}} - \cos(2) \left(\left(-\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right) \right) \cos(\sqrt{3}x) + \sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right) \right)}{\sqrt{3} \cos\left(\frac{\sqrt{3}}{2}\right) + \sin\left(\frac{\sqrt{3}}{2}\right)}$$

✓ Solution by Mathematica

Time used: 0.786 (sec). Leaf size: 12765

```
DSolve[{y'''[x]+y'[x]+y[x]==Sin[x],{y'[1] == 0,y[2]==0}},y[x],x,IncludeSingularSolutions ->
```

Too large to display

3.14 problem 14

Internal problem ID [7204]

Internal file name [OUTPUT/6190_Sunday_June_05_2022_04_27_25_PM_95321338/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 14.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

$$y''' + y' + y = x$$

With initial conditions

$$[y'(0) = 0, y(0) = 0, y''(0) = 1]$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' + y' + y = 0$$

The characteristic equation is

$$\lambda^3 + \lambda + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = -\frac{(108 + 12\sqrt{93})^{\frac{1}{3}}}{6} + \frac{2}{(108 + 12\sqrt{93})^{\frac{1}{3}}}$$

$$\lambda_2 = \frac{(108 + 12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108 + 12\sqrt{93})^{\frac{1}{3}}} + \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2}$$

$$\lambda_3 = \frac{(108 + 12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108 + 12\sqrt{93})^{\frac{1}{3}}} - \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2}$$

Therefore the homogeneous solution is

$$y_h(x) = e^{\left(\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108+12\sqrt{93})^{\frac{1}{3}}} + \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2} \right) x} c_1 + e^{\left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} + \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right) x} c_2 + e^{\left(\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108+12\sqrt{93})^{\frac{1}{3}}} - \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2} \right) x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{\left(\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108+12\sqrt{93})^{\frac{1}{3}}} + \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2} \right) x}$$

$$y_2 = e^{\left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} + \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right) x}$$

$$y_3 = e^{\left(\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108+12\sqrt{93})^{\frac{1}{3}}} - \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2} \right) x}$$

Now the particular solution to the given ODE is found

$$y''' + y' + y = x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{\left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} + \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right) x}, e^{\left(\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108+12\sqrt{93})^{\frac{1}{3}}} - \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2} \right) x}, e^{\left(\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{12} \right) x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2x + A_1 + A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x - 1$$

Therefore the general solution is

$$y = y_h + y_p$$

$$\begin{aligned}
&= \left(e^{\left(\frac{\left(\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108+12\sqrt{93})^{\frac{1}{3}}} + \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2} \right) x}{\right)} c_1 \right. \\
&\quad \left. + e^{\left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} + \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right) x} c_2 \right. \\
&\quad \left. + e^{\left(\frac{\left(\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{12} - \frac{1}{(108+12\sqrt{93})^{\frac{1}{3}}} - \frac{i\sqrt{3} \left(-\frac{(108+12\sqrt{93})^{\frac{1}{3}}}{6} - \frac{2}{(108+12\sqrt{93})^{\frac{1}{3}}} \right)}{2} \right) x}{\right)} c_3 \right) + (x-1)
\end{aligned}$$

Which simplifies to

$$\begin{aligned}
y = e & \frac{x \left((i\sqrt{3}-1) \frac{(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}} + 12i\sqrt{3}+12}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}} \right)}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}} c_1 + e^{\left(\frac{((108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}-12)x}{6(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}} \right)} c_2 \\
& + e^{\left(\frac{(i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3} + (108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}} + 12i\sqrt{3}-12)x}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}} \right)} c_3 + x - 1
\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{\left(\frac{x \left((i\sqrt{3}-1) \frac{(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}} + 12i\sqrt{3}+12}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}} \right)}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}} \right)} c_1 + e^{\left(\frac{((108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}-12)x}{6(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}} \right)} c_2 + e^{\left(\frac{(i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3} + (108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}} + 12i\sqrt{3}-12)x}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}} \right)} c_3 + x - 1 \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 + c_3 - 1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -\frac{\left((i\sqrt{3}-1)(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}+12i\sqrt{3}+12\right)e^{-\frac{x\left((i\sqrt{3}-1)(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}+12i\sqrt{3}+12\right)}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}}}}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}}c_1 - \frac{\left((108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3}\sqrt{31}-3i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{31}-19i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3}-38(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\right)}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = \frac{(108+12\sqrt{93})^{\frac{2}{3}}(-i(c_1-c_3)\sqrt{3}+c_1-2c_2+c_3)+12(108+12\sqrt{93})^{\frac{1}{3}}-12i(c_1-c_3)\sqrt{3}-12c_1+2c_2-c_3}{12(108+12\sqrt{93})^{\frac{1}{3}}}$$

(2A)

Taking two derivatives of the solution gives

$$y'' = \frac{\left((i\sqrt{3}-1)(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}+12i\sqrt{3}+12\right)^2 e^{-\frac{x\left((i\sqrt{3}-1)(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}+12i\sqrt{3}+12\right)}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}}}}{144(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}}c_1 + \frac{\left((108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3}\sqrt{31}-3i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{31}-19i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3}-38(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\right)}{144(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}}$$

substituting $y'' = 1$ and $x = 0$ in the above gives

$$1 = \frac{3\left(-i(c_1-c_3)\sqrt{3}-\frac{(c_1-2c_2+c_3)\sqrt{93}}{9}-\frac{i(c_1-c_3)\sqrt{31}}{3}-c_1+2c_2-c_3\right)(108+12\sqrt{93})^{\frac{1}{3}}-\frac{2(c_1+c_2+c_3)(108+12\sqrt{93})^{\frac{2}{3}}}{3}-2i(-c_1+c_3)\sqrt{3}}{(108+12\sqrt{93})^{\frac{2}{3}}}$$

(3A)

Equations {1A,2A,3A} are now solved for $\{c_1, c_2, c_3\}$. Solving for the constants gives

$$c_1 = \frac{\left(5(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3}\sqrt{31}-3i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{31}-19i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3}-38(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\right)}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}}$$

$$c_2 = \frac{30\sqrt{31}(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}+3(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{31}+19(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3}+78(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}}{1116\sqrt{3}+324\sqrt{31}}$$

$$c_3 = \frac{\left(5(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{3}\sqrt{31}-38(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}\sqrt{3}\sqrt{31}-96\sqrt{3}\sqrt{31}+3i(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{31}\right)}{12(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}}$$

Substituting these values back in above solution results in

$$y = \text{Expression too large to display}$$

Summary

The solution(s) found are the following

$$y \tag{1}$$

$$= \frac{\left((-3i\sqrt{93} - 57i + 43\sqrt{3} + 15\sqrt{31}) (108 + 12\sqrt{93})^{\frac{2}{3}} + (-54i\sqrt{93} - 558i - 342\sqrt{3} - 114\sqrt{31}) (108 + \dots\right)}{\dots}$$

Verification of solutions

$$y$$

$$= \frac{\left((-3i\sqrt{93} - 57i + 43\sqrt{3} + 15\sqrt{31}) (108 + 12\sqrt{93})^{\frac{2}{3}} + (-54i\sqrt{93} - 558i - 342\sqrt{3} - 114\sqrt{31}) (108 + \dots\right)}{\dots}$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 3; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.609 (sec). Leaf size: 359

```
dsolve([diff(y(x),x$3)+diff(y(x),x)+y(x)=x,D(y)(0) = 0, y(0) = 0, (D@@2)(y)(0) = 1],y(x), si
```

$$y(x) = \frac{10 e^{-\frac{x(108+12\sqrt{93})^{\frac{1}{3}}(-12+(\sqrt{93}-9)(108+12\sqrt{93})^{\frac{1}{3}})}{144}} \left((108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}\sqrt{3}\sqrt{31} + \frac{3\sqrt{3}(108+12\sqrt{3}\sqrt{31})^{\frac{2}{3}}\sqrt{31}}{5} - \frac{6\sqrt{3}\sqrt{31}}{5} - \frac{39(108+12\sqrt{3}\sqrt{31})^{\frac{1}{3}}}{5} \right)}{3}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 1546

```
DSolve[{y'''[x]+y'[x]+y[x]==x,{y'[1] == 0,y[0]==0,y''[0]==1}},y[x],x,IncludeSingularSolution
```

Too large to display

3.15 problem 15

- 3.15.1 Solving as second order change of variable on x method 2 ode . 1304
- 3.15.2 Solving as second order change of variable on x method 1 ode . 1309
- 3.15.3 Solving as second order change of variable on y method 2 ode . 1314
- 3.15.4 Solving using Kovacic algorithm 1319

Internal problem ID [7205]

Internal file name [OUTPUT/6191_Sunday_June_05_2022_04_27_29_PM_70179798/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 15.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^4y'' + y'x^3 - 4x^2y = 1$$

3.15.1 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^4y'' + y'x^3 - 4x^2y = 0$$

In normal form the ode

$$x^4y'' + y'x^3 - 4x^2y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x} dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{4}{x^2}}{\frac{1}{x^2}} \\ &= -4 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 4y(\tau) &= 0\end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 4e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)} \\ &= \pm 2\end{aligned}$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(2)\tau} + c_2 e^{(-2)\tau}$$

Or

$$y(\tau) = c_1 e^{2\tau} + c_2 e^{-2\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 x^4 + c_2}{x^2}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 x^4 + c_2}{x^2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx}\left(\frac{1}{x^2}\right) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)(2x) - (x^2)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^2}{4x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{4x} dx$$

Hence

$$u_1 = -\frac{\ln(x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x^2}}{4x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^5} dx$$

Hence

$$u_2 = -\frac{1}{16x^4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\ln(x)}{4x^2} - \frac{1}{16x^2}$$

Which simplifies to

$$y_p(x) = \frac{-1 - 4 \ln(x)}{16x^2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1 x^4 + c_2}{x^2} \right) + \left(\frac{-1 - 4 \ln(x)}{16x^2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^4 + c_2}{x^2} + \frac{-1 - 4 \ln(x)}{16x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 x^4 + c_2}{x^2} + \frac{-1 - 4 \ln(x)}{16x^2}$$

Verified OK.

3.15.2 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^4$, $B = x^3$, $C = -4x^2$, $f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^4 y'' + y' x^3 - 4x^2 y = 0$$

In normal form the ode

$$x^4 y'' + y' x^3 - 4x^2 y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c} \sqrt{q}$$

$$= \frac{2\sqrt{-\frac{1}{x^2}}}{c} \quad (6)$$

$$\tau'' = \frac{2}{c\sqrt{-\frac{1}{x^2}} x^3}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2}$$

$$= \frac{\frac{2}{c\sqrt{-\frac{1}{x^2}} x^3} + \frac{1}{x} \frac{2\sqrt{-\frac{1}{x^2}}}{c}}{\left(\frac{2\sqrt{-\frac{1}{x^2}}}{c} \right)^2}$$

$$= 0$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int 2\sqrt{-\frac{1}{x^2}} dx}{c} \\ &= \frac{2\sqrt{-\frac{1}{x^2}} x \ln(x)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))$$

Now the particular solution to this ODE is found

$$x^4 y'' + y' x^3 - 4x^2 y = 1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \frac{1}{x^2} \\ y_2 &= x^2 \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx} \left(\frac{1}{x^2} \right) & \frac{d}{dx} (x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2} \right) (2x) - (x^2) \left(-\frac{2}{x^3} \right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^2}{4x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{4x} dx$$

Hence

$$u_1 = -\frac{\ln(x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x^2}}{4x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^5} dx$$

Hence

$$u_2 = -\frac{1}{16x^4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\ln(x)}{4x^2} - \frac{1}{16x^2}$$

Which simplifies to

$$y_p(x) = \frac{-1 - 4 \ln(x)}{16x^2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))) + \left(\frac{-1 - 4 \ln(x)}{16x^2} \right) \\ &= \frac{-1 - 4 \ln(x)}{16x^2} + c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x)) \end{aligned}$$

Which simplifies to

$$y = \frac{-1 - 4 \ln(x)}{16x^2} + c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))$$

Summary

The solution(s) found are the following

$$y = \frac{-1 - 4 \ln(x)}{16x^2} + c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x)) \quad (1)$$

Verification of solutions

$$y = \frac{-1 - 4 \ln(x)}{16x^2} + c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))$$

Verified OK.

3.15.3 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^4$, $B = x^3$, $C = -4x^2$, $f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^4 y'' + y' x^3 - 4x^2 y = 0$$

In normal form the ode

$$x^4 y'' + y' x^3 - 4x^2 y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right) v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right) v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{4}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \frac{5v'(x)}{x} &= 0 \\ v''(x) + \frac{5v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{5u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{5u}{x} \end{aligned}$$

Where $f(x) = -\frac{5}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{5}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{5}{x} dx \\ \ln(u) &= -5 \ln(x) + c_1 \\ u &= e^{-5 \ln(x) + c_1} \\ &= \frac{c_1}{x^5} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= -\frac{c_1}{4x^4} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(-\frac{c_1}{4x^4} + c_2\right) x^2 \\&= \frac{4c_2x^4 - c_1}{4x^2}\end{aligned}$$

Now the particular solution to this ODE is found

$$x^4y'' + y'x^3 - 4x^2y = 1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= \frac{1}{x^2} \\y_2 &= x^2\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx}\left(\frac{1}{x^2}\right) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)(2x) - (x^2)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^2}{4x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{4x} dx$$

Hence

$$u_1 = -\frac{\ln(x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x^2}}{4x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^5} dx$$

Hence

$$u_2 = -\frac{1}{16x^4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\ln(x)}{4x^2} - \frac{1}{16x^2}$$

Which simplifies to

$$y_p(x) = \frac{-1 - 4 \ln(x)}{16x^2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\left(-\frac{c_1}{4x^4} + c_2 \right) x^2 \right) + \left(\frac{-1 - 4 \ln(x)}{16x^2} \right) \\ &= \frac{-1 - 4 \ln(x)}{16x^2} + \left(-\frac{c_1}{4x^4} + c_2 \right) x^2 \end{aligned}$$

Which simplifies to

$$y = -\frac{-16c_2x^4 + 4 \ln(x) + 4c_1 + 1}{16x^2}$$

Summary

The solution(s) found are the following

$$y = -\frac{-16c_2x^4 + 4 \ln(x) + 4c_1 + 1}{16x^2} \tag{1}$$

Verification of solutions

$$y = -\frac{-16c_2x^4 + 4 \ln(x) + 4c_1 + 1}{16x^2}$$

Verified OK.

3.15.4 Solving using Kovacic algorithm

Writing the ode as

$$x^4 y'' + y' x^3 - 4x^2 y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^4 \\ B &= x^3 \\ C &= -4x^2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{15}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 15 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{15}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 136: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{15}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{15}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{15}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = -\frac{3}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -\frac{3}{2} - \left(-\frac{3}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= -\frac{3}{2x} + (-)(0) \\ &= -\frac{3}{2x} \\ &= -\frac{3}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{3}{2x}\right)(0) + \left(\left(\frac{3}{2x^2}\right) + \left(-\frac{3}{2x}\right)^2 - \left(\frac{15}{4x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{3}{2x} dx} \\ &= \frac{1}{x^{\frac{3}{2}}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{x^3}{x^4} dx} \\&= z_1 e^{-\frac{\ln(x)}{2}} \\&= z_1 \left(\frac{1}{\sqrt{x}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{x^3}{x^4} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{x^4}{4} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(\frac{1}{x^2} \right) + c_2 \left(\frac{1}{x^2} \left(\frac{x^4}{4} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^4y'' + y'x^3 - 4x^2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1}{x^2} + \frac{c_2x^2}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = \frac{x^2}{4}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & \frac{x^2}{4} \\ \frac{d}{dx} \left(\frac{1}{x^2} \right) & \frac{d}{dx} \left(\frac{x^2}{4} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & \frac{x^2}{4} \\ -\frac{2}{x^3} & \frac{x}{2} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)\left(\frac{x}{2}\right) - \left(\frac{x^2}{4}\right)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{1}{x}$$

Which simplifies to

$$W = \frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{x^2}{4}}{x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{4x} dx$$

Hence

$$u_1 = -\frac{\ln(x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x^2}}{x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x^5} dx$$

Hence

$$u_2 = -\frac{1}{4x^4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\ln(x)}{4x^2} - \frac{1}{16x^2}$$

Which simplifies to

$$y_p(x) = \frac{-1 - 4 \ln(x)}{16x^2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1}{x^2} + \frac{c_2 x^2}{4} \right) + \left(\frac{-1 - 4 \ln(x)}{16x^2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} + \frac{c_2 x^2}{4} + \frac{-1 - 4 \ln(x)}{16x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^2} + \frac{c_2 x^2}{4} + \frac{-1 - 4 \ln(x)}{16x^2}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    <- LODE of Euler type successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(x^4*diff(y(x),x$2)+x^3*diff(y(x),x)-4*x^2*y(x)=1,y(x), singsol=all)
```

$$y(x) = \frac{16c_2x^4 - 4\ln(x) + 16c_1 - 1}{16x^2}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 29

```
DSolve[x^4*y''[x]+x^3*y'[x]-4*x^2*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{16c_2x^4 - 4\log(x) - 1 + 16c_1}{16x^2}$$

3.16 problem 16

- 3.16.1 Solving as second order change of variable on x method 2 ode . 1328
- 3.16.2 Solving as second order change of variable on x method 1 ode . 1333
- 3.16.3 Solving as second order change of variable on y method 2 ode . 1338
- 3.16.4 Solving using Kovacic algorithm 1342

Internal problem ID [7206]

Internal file name [OUTPUT/6192_Sunday_June_05_2022_04_27_31_PM_6887828/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 16.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^4y'' + y'x^3 - 4x^2y = x$$

3.16.1 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^4y'' + y'x^3 - 4x^2y = 0$$

In normal form the ode

$$x^4y'' + y'x^3 - 4x^2y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x} dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{4}{x^2}}{\frac{1}{x^2}} \\ &= -4 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) = 0$$

$$\frac{d^2}{d\tau^2}y(\tau) - 4y(\tau) = 0$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 4e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\lambda_{1,2} = \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)}$$

$$= \pm 2$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(2)\tau} + c_2 e^{(-2)\tau}$$

Or

$$y(\tau) = c_1 e^{2\tau} + c_2 e^{-2\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 x^4 + c_2}{x^2}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 x^4 + c_2}{x^2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx}\left(\frac{1}{x^2}\right) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)(2x) - (x^2)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3}{4x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{4} dx$$

Hence

$$u_1 = -\frac{x}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x}}{4x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^4} dx$$

Hence

$$u_2 = -\frac{1}{12x^3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{1}{3x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1 x^4 + c_2}{x^2} \right) + \left(-\frac{1}{3x} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^4 + c_2}{x^2} - \frac{1}{3x} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 x^4 + c_2}{x^2} - \frac{1}{3x}$$

Verified OK.

3.16.2 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^4$, $B = x^3$, $C = -4x^2$, $f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^4 y'' + y' x^3 - 4x^2 y = 0$$

In normal form the ode

$$x^4 y'' + y' x^3 - 4x^2 y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{2\sqrt{-\frac{1}{x^2}}}{c} \quad (6)$$

$$\tau'' = \frac{2}{c\sqrt{-\frac{1}{x^2}}x^3}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2}$$

$$= \frac{\frac{2}{c\sqrt{-\frac{1}{x^2}}x^3} + \frac{1}{x}\frac{2\sqrt{-\frac{1}{x^2}}}{c}}{\left(\frac{2\sqrt{-\frac{1}{x^2}}}{c}\right)^2}$$

$$= 0$$

Therefore ode (3) now becomes

$$y(\tau)'' + p_1y(\tau)' + q_1y(\tau) = 0$$

$$\frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) = 0 \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int 2\sqrt{-\frac{1}{x^2}} dx}{c} \\ &= \frac{2\sqrt{-\frac{1}{x^2}} x \ln(x)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))$$

Now the particular solution to this ODE is found

$$x^4 y'' + y' x^3 - 4x^2 y = x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \frac{1}{x^2} \\ y_2 &= x^2 \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx}\left(\frac{1}{x^2}\right) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)(2x) - (x^2)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3}{4x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{4} dx$$

Hence

$$u_1 = -\frac{x}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x}}{4x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^4} dx$$

Hence

$$u_2 = -\frac{1}{12x^3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{1}{3x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))) + \left(-\frac{1}{3x}\right) \\ &= -\frac{1}{3x} + c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x)) \end{aligned}$$

Which simplifies to

$$y = \frac{(3ic_2 + 3c_1)x^4 - 2x - 3ic_2 + 3c_1}{6x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{(3ic_2 + 3c_1)x^4 - 2x - 3ic_2 + 3c_1}{6x^2} \quad (1)$$

Verification of solutions

$$y = \frac{(3ic_2 + 3c_1)x^4 - 2x - 3ic_2 + 3c_1}{6x^2}$$

Verified OK.

3.16.3 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^4$, $B = x^3$, $C = -4x^2$, $f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^4y'' + y'x^3 - 4x^2y = 0$$

In normal form the ode

$$x^4y'' + y'x^3 - 4x^2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variable is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{4}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \tag{6}$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \frac{5v'(x)}{x} &= 0 \\ v''(x) + \frac{5v'(x)}{x} &= 0 \end{aligned} \tag{7}$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{5u(x)}{x} = 0 \tag{8}$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{5u}{x} \end{aligned}$$

Where $f(x) = -\frac{5}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{5}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{5}{x} dx \\ \ln(u) &= -5 \ln(x) + c_1 \\ u &= e^{-5 \ln(x) + c_1} \\ &= \frac{c_1}{x^5} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{4x^4} + c_2 \end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{4x^4} + c_2\right) x^2 \\ &= \frac{4c_2x^4 - c_1}{4x^2}\end{aligned}$$

Now the particular solution to this ODE is found

$$x^4y'' + y'x^3 - 4x^2y = x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx}\left(\frac{1}{x^2}\right) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)(2x) - (x^2)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3}{4x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{4} dx$$

Hence

$$u_1 = -\frac{x}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x}}{4x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^4} dx$$

Hence

$$u_2 = -\frac{1}{12x^3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{1}{3x}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\&= \left(\left(-\frac{c_1}{4x^4} + c_2 \right) x^2 \right) + \left(-\frac{1}{3x} \right) \\&= -\frac{1}{3x} + \left(-\frac{c_1}{4x^4} + c_2 \right) x^2\end{aligned}$$

Which simplifies to

$$y = -\frac{-12c_2x^4 + 3c_1 + 4x}{12x^2}$$

Summary

The solution(s) found are the following

$$y = -\frac{-12c_2x^4 + 3c_1 + 4x}{12x^2} \quad (1)$$

Verification of solutions

$$y = -\frac{-12c_2x^4 + 3c_1 + 4x}{12x^2}$$

Verified OK.

3.16.4 Solving using Kovacic algorithm

Writing the ode as

$$x^4y'' + y'x^3 - 4x^2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^4 \\B &= x^3 \\C &= -4x^2\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{15}{4x^2} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 15 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{15}{4x^2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 137: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 2 - 0 \\
 &= 2
 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{15}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition

of r given above. Therefore $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{15}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{15}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^- = -\frac{3}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^- - (\alpha_{c_1}^-) \\ &= -\frac{3}{2} - \left(-\frac{3}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= -\frac{3}{2x} + (-)(0) \\ &= -\frac{3}{2x} \\ &= -\frac{3}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{3}{2x}\right)(0) + \left(\left(\frac{3}{2x^2}\right) + \left(-\frac{3}{2x}\right)^2 - \left(\frac{15}{4x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{3}{2x} dx} \\ &= \frac{1}{x^{\frac{3}{2}}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{x^3}{x^4} dx} \\&= z_1 e^{-\frac{\ln(x)}{2}} \\&= z_1 \left(\frac{1}{\sqrt{x}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{x^3}{x^4} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{x^4}{4} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(\frac{1}{x^2} \right) + c_2 \left(\frac{1}{x^2} \left(\frac{x^4}{4} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^4y'' + y'x^3 - 4x^2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1}{x^2} + \frac{c_2x^2}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = \frac{x^2}{4}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & \frac{x^2}{4} \\ \frac{d}{dx} \left(\frac{1}{x^2} \right) & \frac{d}{dx} \left(\frac{x^2}{4} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & \frac{x^2}{4} \\ -\frac{2}{x^3} & \frac{x}{2} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)\left(\frac{x}{2}\right) - \left(\frac{x^2}{4}\right)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{1}{x}$$

Which simplifies to

$$W = \frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{x^3}{4}}{x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{4} dx$$

Hence

$$u_1 = -\frac{x}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x}}{x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x^4} dx$$

Hence

$$u_2 = -\frac{1}{3x^3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{1}{3x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1}{x^2} + \frac{c_2 x^2}{4} \right) + \left(-\frac{1}{3x} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} + \frac{c_2 x^2}{4} - \frac{1}{3x} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^2} + \frac{c_2 x^2}{4} - \frac{1}{3x}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    <- LODE of Euler type successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(x^4*diff(y(x),x$2)+x^3*diff(y(x),x)-4*x^2*y(x)=x,y(x), singsol=all)
```

$$y(x) = \frac{3c_2x^4 + 3c_1 - x}{3x^2}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 25

```
DSolve[x^4*y''[x]+x^3*y'[x]-4*x^2*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^2 + \frac{c_1}{x^2} - \frac{1}{3x}$$

3.17 problem 17

3.17.1 Solving as second order euler ode ode	1352
3.17.2 Solving as second order change of variable on x method 2 ode .	1356
3.17.3 Solving as second order change of variable on x method 1 ode .	1361
3.17.4 Solving as second order change of variable on y method 2 ode .	1365
3.17.5 Solving using Kovacic algorithm	1370

Internal problem ID [7207]

Internal file name [OUTPUT/6193_Sunday_June_05_2022_04_27_33_PM_88434747/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 17.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + xy' - 4y = x$$

3.17.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = x$, $C = -4$, $f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + xy' - 4y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + xrx^{r-1} - 4x^r = 0$$

Simplifying gives

$$r(r-1)x^r + rx^r - 4x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + r - 4 = 0$$

Or

$$r^2 - 4 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -2$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x^2} + c_2x^2$$

Next, we find the particular solution to the ODE

$$x^2y'' + xy' - 4y = x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx}\left(\frac{1}{x^2}\right) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)(2x) - (x^2)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3}{4x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^2}{4} dx$$

Hence

$$u_1 = -\frac{x^3}{12}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x}}{4x} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^2} dx$$

Hence

$$u_2 = -\frac{1}{4x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= -\frac{x}{3} + \frac{c_1}{x^2} + c_2x^2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{x}{3} + \frac{c_1}{x^2} + c_2x^2 \quad (1)$$

Verification of solutions

$$y = -\frac{x}{3} + \frac{c_1}{x^2} + c_2x^2$$

Verified OK.

3.17.2 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2y'' + xy' - 4y = 0$$

In normal form the ode

$$x^2y'' + xy' - 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned}
 \tau &= \int e^{-(\int p(x)dx)} dx \\
 &= \int e^{-(\int \frac{1}{x} dx)} dx \\
 &= \int e^{-\ln(x)} dx \\
 &= \int \frac{1}{x} dx \\
 &= \ln(x)
 \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned}
 q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\
 &= \frac{-\frac{4}{x^2}}{\frac{1}{x^2}} \\
 &= -4
 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}
 \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) - 4y(\tau) &= 0
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 4 e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)} \\ &= \pm 2\end{aligned}$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(2)\tau} + c_2 e^{(-2)\tau}$$

Or

$$y(\tau) = c_1 e^{2\tau} + c_2 e^{-2\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 x^4 + c_2}{x^2}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 x^4 + c_2}{x^2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx} \left(\frac{1}{x^2} \right) & \frac{d}{dx} (x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2} \right) (2x) - (x^2) \left(-\frac{2}{x^3} \right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3}{4x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^2}{4} dx$$

Hence

$$u_1 = - \frac{x^3}{12}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x}}{4x} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^2} dx$$

Hence

$$u_2 = - \frac{1}{4x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{x}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1 x^4 + c_2}{x^2} \right) + \left(- \frac{x}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^4 + c_2}{x^2} - \frac{x}{3} \tag{1}$$

Verification of solutions

$$y = \frac{c_1 x^4 + c_2}{x^2} - \frac{x}{3}$$

Verified OK.

3.17.3 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = x, C = -4, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' + xy' - 4y = 0$$

In normal form the ode

$$x^2 y'' + xy' - 4y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{2\sqrt{-\frac{1}{x^2}}}{c} \\ \tau'' &= \frac{2}{c\sqrt{-\frac{1}{x^2}}x^3}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{\frac{2}{c\sqrt{-\frac{1}{x^2}}x^3} + \frac{1}{x}\frac{2\sqrt{-\frac{1}{x^2}}}{c}}{\left(\frac{2\sqrt{-\frac{1}{x^2}}}{c}\right)^2} \\ &= 0\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int 2\sqrt{-\frac{1}{x^2}} dx}{c} \\ &= \frac{2\sqrt{-\frac{1}{x^2}}x \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))$$

Now the particular solution to this ODE is found

$$x^2y'' + xy' - 4y = x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx}\left(\frac{1}{x^2}\right) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)(2x) - (x^2)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3}{4x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^2}{4} dx$$

Hence

$$u_1 = -\frac{x^3}{12}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x}}{4x} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^2} dx$$

Hence

$$u_2 = -\frac{1}{4x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x}{3}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\&= (c_1 \cosh (2 \ln (x)) + ic_2 \sinh (2 \ln (x))) + \left(-\frac{x}{3}\right) \\&= -\frac{x}{3} + c_1 \cosh (2 \ln (x)) + ic_2 \sinh (2 \ln (x))\end{aligned}$$

Which simplifies to

$$y = -\frac{x}{3} + c_1 \cosh (2 \ln (x)) + ic_2 \sinh (2 \ln (x))$$

Summary

The solution(s) found are the following

$$y = -\frac{x}{3} + c_1 \cosh (2 \ln (x)) + ic_2 \sinh (2 \ln (x)) \quad (1)$$

Verification of solutions

$$y = -\frac{x}{3} + c_1 \cosh (2 \ln (x)) + ic_2 \sinh (2 \ln (x))$$

Verified OK.

3.17.4 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = x$, $C = -4$, $f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + xy' - 4y = 0$$

In normal form the ode

$$x^2y'' + xy' - 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{4}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \frac{5v'(x)}{x} = 0$$
$$v''(x) + \frac{5v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{5u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{5u}{x}\end{aligned}$$

Where $f(x) = -\frac{5}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{5}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{5}{x} dx \\ \ln(u) &= -5 \ln(x) + c_1 \\ u &= e^{-5 \ln(x) + c_1} \\ &= \frac{c_1}{x^5}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{4x^4} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{4x^4} + c_2\right) x^2 \\ &= \frac{4c_2x^4 - c_1}{4x^2}\end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' + xy' - 4y = x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ \frac{d}{dx} \left(\frac{1}{x^2} \right) & \frac{d}{dx} (x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & x^2 \\ -\frac{2}{x^3} & 2x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2} \right) (2x) - (x^2) \left(-\frac{2}{x^3} \right)$$

Which simplifies to

$$W = \frac{4}{x}$$

Which simplifies to

$$W = \frac{4}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3}{4x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^2}{4} dx$$

Hence

$$u_1 = - \frac{x^3}{12}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{1}{x}}{4x} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{4x^2} dx$$

Hence

$$u_2 = - \frac{1}{4x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{x}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\left(-\frac{c_1}{4x^4} + c_2 \right) x^2 \right) + \left(-\frac{x}{3} \right) \\ &= -\frac{x}{3} + \left(-\frac{c_1}{4x^4} + c_2 \right) x^2 \end{aligned}$$

Which simplifies to

$$y = -\frac{x}{3} + \left(-\frac{c_1}{4x^4} + c_2 \right) x^2$$

Summary

The solution(s) found are the following

$$y = -\frac{x}{3} + \left(-\frac{c_1}{4x^4} + c_2\right) x^2 \quad (1)$$

Verification of solutions

$$y = -\frac{x}{3} + \left(-\frac{c_1}{4x^4} + c_2\right) x^2$$

Verified OK.

3.17.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + xy' - 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= x \\ C &= -4 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{15}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 15 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{15}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 138: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{15}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{15}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{15}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to

determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_{∞}^{\pm} . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_{\infty}^{-} = -\frac{3}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -\frac{3}{2} - \left(-\frac{3}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= -\frac{3}{2x} + (-)(0) \\ &= -\frac{3}{2x} \\ &= -\frac{3}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{3}{2x}\right)(0) + \left(\left(\frac{3}{2x^2}\right) + \left(-\frac{3}{2x}\right)^2 - \left(\frac{15}{4x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned}z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{3}{2x} dx} \\ &= \frac{1}{x^{\frac{3}{2}}}\end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx} \\ &= z_1 e^{-\frac{\ln(x)}{2}} \\ &= z_1 \left(\frac{1}{\sqrt{x}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(\frac{x^4}{4} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x^2} \right) + c_2 \left(\frac{1}{x^2} \left(\frac{x^4}{4} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2y'' + xy' - 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1}{x^2} + \frac{c_2x^2}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x^2}$$
$$y_2 = \frac{x^2}{4}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x^2} & \frac{x^2}{4} \\ \frac{d}{dx}\left(\frac{1}{x^2}\right) & \frac{d}{dx}\left(\frac{x^2}{4}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x^2} & \frac{x^2}{4} \\ -\frac{2}{x^3} & \frac{x}{2} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x^2}\right)\left(\frac{x}{2}\right) - \left(\frac{x^2}{4}\right)\left(-\frac{2}{x^3}\right)$$

Which simplifies to

$$W = \frac{1}{x}$$

Which simplifies to

$$W = \frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3}{4} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^2}{4} dx$$

Hence

$$u_1 = -\frac{x^3}{12}$$

And Eq. (3) becomes

$$u_2 = \int \frac{1}{x} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x^2} dx$$

Hence

$$u_2 = -\frac{1}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1}{x^2} + \frac{c_2 x^2}{4} \right) + \left(-\frac{x}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} + \frac{c_2 x^2}{4} - \frac{x}{3} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^2} + \frac{c_2 x^2}{4} - \frac{x}{3}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    <- LODE of Euler type successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(x^2*diff(diff(y(x),x),x)+x*diff(y(x),x)-4*y(x) = x,y(x), singsol=all)
```

$$y(x) = c_2x^2 + \frac{c_1}{x^2} - \frac{x}{3}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 23

```
DSolve[x^2*y''[x]+x*y'[x]-4*y[x] == x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^2 + \frac{c_1}{x^2} - \frac{x}{3}$$

3.18 problem 18

3.18.1 Maple step by step solution 1382

Internal problem ID [7208]

Internal file name [OUTPUT/6194_Sunday_June_05_2022_04_27_35_PM_24677674/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 18.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : **"higher_order_ODE_non_constant_coefficients_of_type_Euler"**

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

$$x^4 y''' + x^3 y'' + x^2 y' + yx = 0$$

This is Euler ODE of higher order. Let $y = x^\lambda$. Hence

$$\begin{aligned}y' &= \lambda x^{\lambda-1} \\y'' &= \lambda(\lambda-1) x^{\lambda-2} \\y''' &= \lambda(\lambda-1)(\lambda-2) x^{\lambda-3}\end{aligned}$$

Substituting these back into

$$x^3 y''' + x^2 y'' + x y' + y = 0$$

gives

$$x\lambda x^{\lambda-1} + x^2\lambda(\lambda-1) x^{\lambda-2} + x^3\lambda(\lambda-1)(\lambda-2) x^{\lambda-3} + x^\lambda = 0$$

Which simplifies to

$$\lambda x^\lambda + \lambda(\lambda-1) x^\lambda + \lambda(\lambda-1)(\lambda-2) x^\lambda + x^\lambda = 0$$

And since $x^\lambda \neq 0$ then dividing through by x^λ , the above becomes

$$\lambda + \lambda(\lambda - 1) + \lambda(\lambda - 1)(\lambda - 2) + 1 = 0$$

Simplifying gives the characteristic equation as

$$\lambda^3 - 2\lambda^2 + 2\lambda + 1 = 0$$

Solving the above gives the following roots

$$\lambda_1 = -\frac{(188 + 12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188 + 12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}$$

$$\lambda_2 = \frac{(188 + 12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188 + 12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3} + \frac{i\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right)}{2}$$

$$\lambda_3 = \frac{(188 + 12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188 + 12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3} - \frac{i\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right)}{2}$$

This table summarises the result

root	multiplicity	type of root
$\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3} \pm \frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right)}{2} i$	1	complex conjugate
$-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}$	1	real root

The solution is generated by going over the above table. For each real root λ of multiplicity one generates a $c_1 x^\lambda$ basis solution. Each real root of multiplicity two, generates $c_1 x^\lambda$ and $c_2 x^\lambda \ln(x)$ basis solutions. Each real root of multiplicity three, generates $c_1 x^\lambda$ and $c_2 x^\lambda \ln(x)$ and $c_3 x^\lambda \ln(x)^2$ basis solutions, and so on. Each complex root $\alpha \pm i\beta$ of multiplicity one generates $x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity two generates $\ln(x) x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity three generates $\ln(x)^2 x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And so on. Using the above show that the solution is

$$y = x^{\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}} + \frac{2}{3}}} \left(c_1 \cos \left(\frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right) \ln(x)}{2} \right) - c_2 \sin \left(\frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right) \ln(x)}{2} \right) \right)$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = x^{\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}} + \frac{2}{3}}} \cos \left(\frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right) \ln(x)}{2} \right)$$

$$y_2 = -x^{\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}} + \frac{2}{3}}} \sin \left(\frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right) \ln(x)}{2} \right)$$

$$y_3 = x^{-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}} + \frac{2}{3}}}$$

Summary

The solution(s) found are the following

$$y = x^{\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}} + \frac{2}{3}}} \left(c_1 \cos \left(\frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right) \ln(x)}{2} \right) - c_2 \sin \left(\frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right) \ln(x)}{2} \right) \right) + c_3 x^{-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}} + \frac{2}{3}}} \quad (1)$$

Verification of solutions

y

$$= x \left(c_1 \cos \left(\frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right) \ln(x)}{2} \right) - c_2 \sin \left(\frac{\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right) \ln(x)}{2} \right) \right) + c_3 x - \frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}$$

Verified OK.

3.18.1 Maple step by step solution

Let's solve

$$x^3 y''' + x^2 y'' + x y' + y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Isolate 3rd derivative

$$y''' = -\frac{y}{x^3} - \frac{x y'' + y'}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y''' + \frac{y''}{x} + \frac{y'}{x^2} + \frac{y}{x^3} = 0$$

- Multiply by denominators of the ODE

$$x^3 y''' + x^2 y'' + x y' + y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

- Calculate the 3rd derivative of y with respect to x , using the chain rule

$$y''' = \left(\frac{d^3}{dt^3}y(t) \right) t'(x)^3 + 3t'(x)t''(x) \left(\frac{d^2}{dt^2}y(t) \right) + t'''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y''' = \frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3}$$

Substitute the change of variables back into the ODE

$$x^3 \left(\frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3} \right) + x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + \frac{d}{dt}y(t) + y(t) = 0$$

- Simplify

$$\frac{d^3}{dt^3}y(t) - 2\frac{d^2}{dt^2}y(t) + 2\frac{d}{dt}y(t) + y(t) = 0$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(t)$

$$y_1(t) = y(t)$$

- Define new variable $y_2(t)$

$$y_2(t) = \frac{d}{dt}y(t)$$

- Define new variable $y_3(t)$

$$y_3(t) = \frac{d^2}{dt^2}y(t)$$

- Isolate for $\frac{d}{dt}y_3(t)$ using original ODE

$$\frac{d}{dt}y_3(t) = 2y_3(t) - 2y_2(t) - y_1(t)$$

Convert linear ODE into a system of first order ODEs

$$\left[y_2(t) = \frac{d}{dt}y_1(t), y_3(t) = \frac{d}{dt}y_2(t), \frac{d}{dt}y_3(t) = 2y_3(t) - 2y_2(t) - y_1(t) \right]$$

- Define vector

$$\vec{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \end{bmatrix}$$

- System to solve

$$\frac{d}{dt}\vec{y}(t) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & -2 & 2 \end{bmatrix} \cdot \vec{y}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & -2 & 2 \end{bmatrix}$$

- Rewrite the system as

$$\frac{d}{dt}\vec{y}(t) = A \cdot \vec{y}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\begin{bmatrix} -\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}, \right. \\ \left. \begin{bmatrix} \frac{1}{\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right)^2} \\ \frac{1}{-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}} \\ 1 \end{bmatrix}, \right. \\ \left. \left. \begin{bmatrix} \frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[\begin{array}{c} -\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}, \\ \left(\frac{1}{-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}} \right)^2 \\ \frac{1}{-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}} \\ 1 \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3} \right) t} \cdot \left[\begin{array}{c} \frac{1}{\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3} \right)^2} \\ \frac{1}{-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}} \\ 1 \end{array} \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[\begin{array}{c} \frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3} - \frac{I\sqrt{3} \left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} \right)}{2}, \\ \frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3} \\ \frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3} \end{array} \right]$$

- Solution from eigenpair

$$e^{\left(\frac{\left(\frac{188+12\sqrt{249}}{12} \right)^{\frac{1}{3}} - \frac{2}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}}} + \frac{2}{3} - \frac{i\sqrt{3} \left(-\frac{\left(\frac{188+12\sqrt{249}}{6} \right)^{\frac{1}{3}} - \frac{4}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}}}{2} \right) t \right)} \cdot \left[\begin{array}{l} \left(\frac{\left(\frac{188+12\sqrt{249}}{12} \right)^{\frac{1}{3}} - \frac{2}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}}}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}} \right) \\ \left(\frac{\left(\frac{188+12\sqrt{249}}{12} \right)^{\frac{1}{3}} - \frac{2}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}}}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}} \right) \end{array} \right]$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{\left(\frac{\left(\frac{188+12\sqrt{249}}{12} \right)^{\frac{1}{3}} - \frac{2}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}}} + \frac{2}{3} \right) t} \cdot \left(\cos \left(\frac{\sqrt{3} \left(-\frac{\left(\frac{188+12\sqrt{249}}{6} \right)^{\frac{1}{3}} - \frac{4}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}}}{2} \right) t}{2} \right) - i \sin \left(\frac{\sqrt{3} \left(-\frac{\left(\frac{188+12\sqrt{249}}{6} \right)^{\frac{1}{3}} - \frac{4}{3\left(188+12\sqrt{249}\right)^{\frac{1}{3}}}}{2} \right) t}{2} \right) \right)$$

- Simplify expression

$$e^{\left(\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right)t} \begin{bmatrix} \cos\left(\frac{\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}\right) - I \sin\left(\frac{\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}\right)}{\left(\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right) - \frac{I\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}} \\ \cos\left(\frac{\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}\right) - I \sin\left(\frac{\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}\right)}{\left(\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right) - \frac{I\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}} \\ \cos\left(\frac{\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}\right) - I \sin\left(\frac{\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}\right)}{\left(\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right) - \frac{I\sqrt{3}\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} - \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}} \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_2(t) = e^{\left(\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right)t} \begin{bmatrix} 18(188+12\sqrt{249})^{\frac{2}{3}} \left((188+12\sqrt{249})^{\frac{4}{3}} \sqrt{3} \sin\left(\frac{\sqrt{3}\left(\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{12} - \frac{2}{3(188+12\sqrt{249})^{\frac{1}{3}}}\right)t}{2}\right) \right) \\ \dots \end{bmatrix}$$

- General solution to the system of ODEs
 $\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(t) + c_3 \vec{y}_3(t)$
- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right)t} \begin{bmatrix} \frac{1}{\left(-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right)^2} \\ \frac{1}{-\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}} \\ 1 \end{bmatrix} + c_2 e^{\left(\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right)t} \begin{bmatrix} \frac{1}{\left(\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}\right)^2} \\ \frac{1}{\frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{4}{3(188+12\sqrt{249})^{\frac{1}{3}}} + \frac{2}{3}} \\ 1 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y(t) = \frac{96 \left(\left(\left(\left(-\frac{7c_3}{32} - \frac{7\sqrt{3}c_2}{96} \right) \sqrt{83} - \frac{115c_3\sqrt{3}}{96} - \frac{115c_2}{96} \right) (188+12\sqrt{3}\sqrt{83})^{\frac{2}{3}} + c_2 \left(\sqrt{3}\sqrt{83} + \frac{47}{3} \right) (188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}} + \left(-\frac{10\sqrt{3}c_2}{3} + 10 \right) \right)}{1}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{96 \left(\left(\left(\left(-\frac{7c_3}{32} - \frac{7\sqrt{3}c_2}{96} \right) \sqrt{83} - \frac{115c_3\sqrt{3}}{96} - \frac{115c_2}{96} \right) (188+12\sqrt{3}\sqrt{83})^{\frac{2}{3}} + c_2 \left(\sqrt{3}\sqrt{83} + \frac{47}{3} \right) (188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}} + \left(-\frac{10\sqrt{3}c_2}{3} + 10 \right) \right)}{1}$$

- Simplify

$$y = \frac{7 \left(\left(\left(\left(\sqrt{3}c_2 + 3c_3 \right) \sqrt{83} + \frac{115c_3\sqrt{3}}{7} + \frac{115c_2}{7} \right) (188+12\sqrt{3}\sqrt{83})^{\frac{2}{3}} - \frac{96c_2 \left(\sqrt{3}\sqrt{83} + \frac{47}{3} \right) (188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}}}{7} + \frac{320(\sqrt{3}c_2 - 3c_3)\sqrt{83}}{7} + 10 \right)}{1}$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 184

```
dsolve(x^4*diff(y(x),x$3)+x^3*diff(y(x),x$2)+x^2*diff(y(x),x)+x*y(x)= 0,y(x), singsol=all)
```

$$y(x) = c_1 x \frac{-(188+12\sqrt{249})^{\frac{2}{3}} - 4(188+12\sqrt{249})^{\frac{1}{3}} - 8}{6(188+12\sqrt{249})^{\frac{1}{3}}} + c_2 x \frac{-8 + (188+12\sqrt{249})^{\frac{2}{3}} + 8(188+12\sqrt{249})^{\frac{1}{3}}}{12(188+12\sqrt{249})^{\frac{1}{3}}} \sin\left(\frac{\sqrt{3}\left((188+12\sqrt{3}\sqrt{83})^{\frac{2}{3}} + 8\right)\ln(x)}{12(188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}}}\right) + c_3 x \frac{-8 + (188+12\sqrt{249})^{\frac{2}{3}} + 8(188+12\sqrt{249})^{\frac{1}{3}}}{12(188+12\sqrt{249})^{\frac{1}{3}}} \cos\left(\frac{\sqrt{3}\left((188+12\sqrt{3}\sqrt{83})^{\frac{2}{3}} + 8\right)\ln(x)}{12(188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}}}\right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 81

```
DSolve[x^4*y'''[x]+x^3*y''[x]+x^2*y'[x]+x*y[x]== 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x^{\text{Root}[\#1^3 - 2\#1^2 + 2\#1 + 1 \&, 1]} + c_3 x^{\text{Root}[\#1^3 - 2\#1^2 + 2\#1 + 1 \&, 3]} + c_2 x^{\text{Root}[\#1^3 - 2\#1^2 + 2\#1 + 1 \&, 2]}$$

3.19 problem 19

3.19.1 Maple step by step solution 1390

Internal problem ID [7209]

Internal file name [OUTPUT/6195_Sunday_June_05_2022_04_27_37_PM_96743745/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 19.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$x^4 y''' + x^3 y'' + x^2 y' + yx = x$$

Unable to solve this ODE.

3.19.1 Maple step by step solution

Let's solve

$$x^4 y''' + x^3 y'' + x^2 y' + yx = x$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 3; missing the dependent variable  
checking if the LODE is of Euler type  
<- LODE of Euler type successful  
Euler equation successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 223

```
dsolve(x^4*diff(y(x),x$3)+x^3*diff(y(x),x$2)+x^2*diff(y(x),x)+x*y(x)= x,y(x), singsol=all)
```

$$y(x) = c_2 x \frac{(47-3\sqrt{249})(188+12\sqrt{249})^{\frac{2}{3}} + (188+12\sqrt{249})^{\frac{1}{3}}}{192} + \frac{2}{3} \cos \left(\frac{(188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}} \sqrt{3} \left(3(188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}} \sqrt{3}\sqrt{83} - \right)}{192} \right) + c_3 x \frac{(47-3\sqrt{249})(188+12\sqrt{249})^{\frac{2}{3}} + (188+12\sqrt{249})^{\frac{1}{3}}}{192} + \frac{2}{3} \sin \left(\frac{(188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}} \sqrt{3} \left(3(188+12\sqrt{3}\sqrt{83})^{\frac{1}{3}} \sqrt{3}\sqrt{83} - \right)}{192} \right) + x \frac{(188+12\sqrt{249})^{\frac{2}{3}} (-47+3\sqrt{249}) - (188+12\sqrt{249})^{\frac{1}{3}}}{96} - \frac{(188+12\sqrt{249})^{\frac{1}{3}}}{6} + \frac{2}{3} c_1 + 1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 82

```
DSolve[x^4*y'''[x]+x^3*y''[x]+x^2*y'[x]+x*y[x]== x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x^{\text{Root}[\#1^3-2\#1^2+2\#1+1\&,1]} + c_3 x^{\text{Root}[\#1^3-2\#1^2+2\#1+1\&,3]} + c_2 x^{\text{Root}[\#1^3-2\#1^2+2\#1+1\&,2]} + 1$$

3.20 problem 20

Internal problem ID [7210]

Internal file name [OUTPUT/6196_Sunday_June_05_2022_04_27_39_PM_75557255/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 20.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_ODE_non_constant_coefficients_of_type_Euler**"

Maple gives the following as the ode type

```
[[_high_order , _with_linear_symmetries]]
```

$$5x^5y'''' + 4x^4y'''' + x^2y' + xy = 0$$

This is Euler ODE of higher order. Let $y = x^\lambda$. Hence

$$\begin{aligned}y' &= \lambda x^{\lambda-1} \\y'' &= \lambda(\lambda-1) x^{\lambda-2} \\y''' &= \lambda(\lambda-1)(\lambda-2) x^{\lambda-3} \\y'''' &= \lambda(\lambda-1)(\lambda-2)(\lambda-3) x^{\lambda-4}\end{aligned}$$

Substituting these back into

$$5y''''x^4 + 4y''''x^3 + xy' + y = 0$$

gives

$$x\lambda x^{\lambda-1} + 4x^3\lambda(\lambda-1)(\lambda-2)x^{\lambda-3} + 5x^4\lambda(\lambda-1)(\lambda-2)(\lambda-3)x^{\lambda-4} + x^\lambda = 0$$

Which simplifies to

$$\lambda x^\lambda + 4\lambda(\lambda-1)(\lambda-2)x^\lambda + 5\lambda(\lambda-1)(\lambda-2)(\lambda-3)x^\lambda + x^\lambda = 0$$

And since $x^\lambda \neq 0$ then dividing through by x^λ , the above becomes

$$\lambda + 4\lambda(\lambda - 1)(\lambda - 2) + 5\lambda(\lambda - 1)(\lambda - 2)(\lambda - 3) + 1 = 0$$

Simplifying gives the characteristic equation as

$$5\lambda^4 - 26\lambda^3 + 43\lambda^2 - 21\lambda + 1 = 0$$

Solving the above gives the following roots

$$\lambda_1 = \frac{13}{10} + \frac{\sqrt{6} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}{60} + \frac{i\sqrt{6} \sqrt{\frac{5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}{5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}}}{60}$$

$$\lambda_2 = \frac{13}{10} + \frac{\sqrt{6} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}{60} - \frac{i\sqrt{6} \sqrt{\frac{5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}{5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}}}{60}$$

$$\lambda_3 = \frac{13}{10} - \frac{\sqrt{6} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}{60} + \frac{\sqrt{6} \sqrt{\frac{-5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}{-5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}}}{60}$$

$$\lambda_4 = \frac{13}{10} - \frac{\sqrt{6} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}{60} - \frac{\sqrt{6} \sqrt{\frac{-5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}{-5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}} + 154(40076+12\sqrt{2307813})^{\frac{1}{3}} + 5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}}}}{60}$$

This table summarises the result

root	
$\frac{13}{10} - \frac{\sqrt{6} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}{60}}$	$+ \sqrt{6} \sqrt{\frac{-5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} (40076+12\sqrt{2307813})^{\frac{1}{3}}$
$\frac{13}{10} + \frac{\sqrt{6} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}{60}}$	$\pm \sqrt{6} \sqrt{\frac{5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} (40076+12\sqrt{2307813})^{\frac{1}{3}}$
$\frac{13}{10} - \frac{\sqrt{6} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}{60}}$	$- \sqrt{6} \sqrt{\frac{-5 \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} (40076+12\sqrt{2307813})^{\frac{1}{3}}$

The solution is generated by going over the above table. For each real root λ of multiplicity one generates a $c_1 x^\lambda$ basis solution. Each real root of multiplicity two, generates $c_1 x^\lambda$ and $c_2 x^\lambda \ln(x)$ basis solutions. Each real root of multiplicity three, generates $c_1 x^\lambda$ and $c_2 x^\lambda \ln(x)$ and $c_3 x^\lambda \ln(x)^2$ basis solutions, and so on. Each complex root $\alpha \pm i\beta$ of multiplicity one generates $x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity two generates $\ln(x) x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity three generates $\ln(x)^2 x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And so on. Using the above show that the solution is

Expression too large to display

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}
 y_1 &= x^{\frac{13}{10}} \left[\frac{\sqrt{6}}{60} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} + \sqrt{\frac{-5 \sqrt{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} \right] \\
 y_2 &= x^{\frac{13}{10}} \left[\frac{\sqrt{6}}{60} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} \right] \cos \left[\sqrt{6} \sqrt{\frac{5 \sqrt{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} \right] \\
 y_3 &= x^{\frac{13}{10}} \left[\frac{\sqrt{6}}{60} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} \right] \sin \left[\sqrt{6} \sqrt{\frac{5 \sqrt{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} \right] \\
 y_4 &= x^{\frac{13}{10}} \left[\frac{\sqrt{6}}{60} \sqrt{\frac{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} - \sqrt{\frac{-5 \sqrt{5(40076+12\sqrt{2307813})^{\frac{2}{3}}+154(40076+12\sqrt{2307813})^{\frac{1}{3}}+5420}}{(40076+12\sqrt{2307813})^{\frac{1}{3}}}} \right]
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\text{Expression too large to display} \tag{1}$$

Verification of solutions

$$\text{Expression too large to display}$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve(5*x^5*diff(y(x),x$4)+4*x^4*diff(y(x),x$3)+x^2*diff(y(x),x)+x*y(x)= 0,y(x), singsol=all
```

$$y(x) = \sum_{a=1}^4 x^{\text{RootOf}(5_Z^4-26_Z^3+43_Z^2-21_Z+1, \text{index}=_a)} C_a$$

✓ Solution by Mathematica

Time used: 1.114 (sec). Leaf size: 1931

```
DSolve[5*x^5*y''''[x]+4*x^4*y''''[x]+x^2*y'[x]+x*y[x]== Sin[x],y[x],x,IncludeSingularSolution
```

Too large to display

3.21 problem 21

3.21.1 Solving as second order ode missing y ode 1397

3.21.2 Maple step by step solution 1399

Internal problem ID [7211]

Internal file name [OUTPUT/6197_Sunday_June_05_2022_04_27_45_PM_68194267/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 21.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]
```

$$(x^2 + 1) y'' + y'^2 = -1$$

3.21.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$(x^2 + 1) p'(x) + 1 + p(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= \frac{-p^2 - 1}{x^2 + 1} \end{aligned}$$

Where $f(x) = \frac{1}{x^2+1}$ and $g(p) = -p^2 - 1$. Integrating both sides gives

$$\begin{aligned}\frac{1}{-p^2-1} dp &= \frac{1}{x^2+1} dx \\ \int \frac{1}{-p^2-1} dp &= \int \frac{1}{x^2+1} dx \\ -\arctan(p) &= \arctan(x) + c_1\end{aligned}$$

The solution is

$$-\arctan(p(x)) - \arctan(x) - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$-\arctan(y') - \arctan(x) - c_1 = 0$$

Integrating both sides gives

$$\begin{aligned}y &= \int -\tan(\arctan(x) + c_1) dx \\ &= \frac{ie^{4ic_1}x}{(e^{2ic_1}-1)^2} - \frac{ix}{(e^{2ic_1}-1)^2} - \frac{4e^{2ic_1} \ln((-e^{2ic_1}+1)x + ie^{2ic_1}+i)}{(e^{2ic_1}-1)^2} + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{ie^{4ic_1}x}{(e^{2ic_1}-1)^2} - \frac{ix}{(e^{2ic_1}-1)^2} - \frac{4e^{2ic_1} \ln((-e^{2ic_1}+1)x + ie^{2ic_1}+i)}{(e^{2ic_1}-1)^2} + c_2 \quad (1)$$

Verification of solutions

$$y = \frac{ie^{4ic_1}x}{(e^{2ic_1}-1)^2} - \frac{ix}{(e^{2ic_1}-1)^2} - \frac{4e^{2ic_1} \ln((-e^{2ic_1}+1)x + ie^{2ic_1}+i)}{(e^{2ic_1}-1)^2} + c_2$$

Verified OK.

3.21.2 Maple step by step solution

Let's solve

$$(x^2 + 1) y'' + y'^2 = -1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$(x^2 + 1) u'(x) + u(x)^2 = -1$$

- Separate variables

$$\frac{u'(x)}{-u(x)^2 - 1} = \frac{1}{x^2 + 1}$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{-u(x)^2 - 1} dx = \int \frac{1}{x^2 + 1} dx + c_1$$

- Evaluate integral

$$-\arctan(u(x)) = \arctan(x) + c_1$$

- Solve for $u(x)$

$$u(x) = -\tan(\arctan(x) + c_1)$$

- Solve 1st ODE for $u(x)$

$$u(x) = -\tan(\arctan(x) + c_1)$$

- Make substitution $u = y'$

$$y' = -\tan(\arctan(x) + c_1)$$

- Integrate both sides to solve for y

$$\int y' dx = \int -\tan(\arctan(x) + c_1) dx + c_2$$

- Compute integrals

$$y = \frac{1 e^{4 I c_1} x}{(e^{2 I c_1} - 1)^2} - \frac{I x}{(e^{2 I c_1} - 1)^2} - \frac{4 e^{2 I c_1} \ln((-e^{2 I c_1} + 1)x + I e^{2 I c_1} + 1)}{(e^{2 I c_1} - 1)^2} + c_2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(_b(_a)^2+1)/(_a^2+1), _b(_a)` *** S
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  trying separable
  <- separable successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve((1+x^2)*diff(y(x),x$2)+1+diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = \frac{\ln(c_1x - 1)c_1^2 + c_2c_1^2 + c_1x + \ln(c_1x - 1)}{c_1^2}$$

✓ Solution by Mathematica

Time used: 8.017 (sec). Leaf size: 33

```
DSolve[(1+x^2)*y''[x]+1+(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \cot(c_1) + \csc^2(c_1) \log(-x \sin(c_1) - \cos(c_1)) + c_2$$

3.22 problem 22

3.22.1 Solving as second order ode missing y ode 1401

Internal problem ID [7212]

Internal file name [OUTPUT/6198_Sunday_June_05_2022_04_27_49_PM_82611728/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 22.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_y**"

Maple gives the following as the ode type

`[[_2nd_order, _missing_y]]`

$$(x^2 + 1)y'' + y'^2 = x - 1$$

3.22.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$(x^2 + 1)p'(x) + 1 + p(x)^2 - x = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= -\frac{p^2 - x + 1}{x^2 + 1} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$p' = -\frac{p^2}{x^2 + 1} + \frac{x}{x^2 + 1} - \frac{1}{x^2 + 1}$$

With Riccati ODE standard form

$$p' = f_0(x) + f_1(x)p + f_2(x)p^2$$

Shows that $f_0(x) = -\frac{1-x}{x^2+1}$, $f_1(x) = 0$ and $f_2(x) = -\frac{1}{x^2+1}$. Let

$$\begin{aligned} p &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-\frac{u}{x^2+1}} \end{aligned} \quad (1)$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= \frac{2x}{(x^2+1)^2} \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= -\frac{1-x}{(x^2+1)^3} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-\frac{u''(x)}{x^2+1} - \frac{2xu'(x)}{(x^2+1)^2} - \frac{(1-x)u(x)}{(x^2+1)^3} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$\begin{aligned} u(x) = & \left(\text{hypergeom} \left(\left[\left[-\frac{i\sqrt{-2+2\sqrt{2}}}{2}, \frac{\sqrt{1-i}}{2} + 1 - \frac{\sqrt{1+i}}{2} \right], [1 - \sqrt{1+i}], \frac{1}{2} \right. \right. \right. \\ & \left. \left. \left. - \frac{ix}{2} \right) (x+i)^{-\frac{\sqrt{1+i}}{2}} c_1 \right. \right. \\ & \left. \left. + \text{hypergeom} \left(\left[\left[\frac{\sqrt{2+2\sqrt{2}}}{2}, \frac{\sqrt{2+2\sqrt{2}}}{2} + 1 \right], [1 + \sqrt{1+i}], \frac{1}{2} - \frac{ix}{2} \right] \right. \right. \right. \\ & \left. \left. \left. + i \right)^{\frac{\sqrt{1+i}}{2}} c_2 \right) (x-i)^{\frac{\sqrt{1-i}}{2}} \right. \end{aligned}$$

The above shows that

$$u'(x) = \frac{\left((x+i)^{-\frac{\sqrt{1+i}}{2}} \sqrt{-2+2\sqrt{2}} (i\sqrt{2} + i\sqrt{1-i} + i\sqrt{1+i} + 1+i) (x^2+1) c_1 \text{ hypergeom} \left(\left[1 - \frac{i\sqrt{-2+2\sqrt{2}}}{2} \right] \right)}{\dots}$$

Using the above in (1) gives the solution

$$p(x) = \frac{\left((x+i)^{-\frac{\sqrt{1+i}}{2}} \sqrt{-2+2\sqrt{2}} (i\sqrt{2} + i\sqrt{1-i} + i\sqrt{1+i} + 1+i) (x^2+1) c_1 \text{ hypergeom} \left(\left[1 - \frac{i\sqrt{-2+2\sqrt{2}}}{2} \right] \right)}{\dots}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$p(x) = \frac{-(x+i)^{-\frac{\sqrt{1+i}}{2}} \sqrt{-2+2\sqrt{2}} (i\sqrt{2} + i\sqrt{1-i} + i\sqrt{1+i} + 1+i) (x^2+1) c_3 \text{ hypergeom} \left(\left[1 - \frac{i\sqrt{-2+2\sqrt{2}}}{2} \right] \right)}{\dots}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{-(x+i)^{-\frac{\sqrt{1+i}}{2}} \sqrt{-2+2\sqrt{2}} (i\sqrt{2} + i\sqrt{1-i} + i\sqrt{1+i} + 1+i) (x^2+1) c_3 \text{ hypergeom} \left(\left[1 - \frac{i\sqrt{-2+2\sqrt{2}}}{2} \right] \right)}{\dots}$$

Integrating both sides gives

$$y = \int \text{Expression too large to display } dx$$

$$= \text{Expression too large to display}$$

Summary

The solution(s) found are the following

$$\text{Expression too large to display} \tag{1}$$

Verification of solutions

Expression too large to display

Warning, solution could not be verified

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(_b(_a)^2-_a+1)/(_a^2+1), _b(_a)` **
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  trying separable
  trying inverse linear
  trying homogeneous types:
  trying Chini
  differential order: 1; looking for linear symmetries
  trying exact
  Looking for potential symmetries
  trying Riccati
  trying Riccati sub-methods:
    <- Abel AIR successful: ODE belongs to the 2F1 2-parameter class
  <- differential order: 2; canonical coordinates successful
  <- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 460

```
dsolve((1+x^2)*diff(y(x),x^2)+1+diff(y(x),x)^2=x,y(x), singsol=all)
```

$$y(x) = - \left(\int \frac{-\left(\frac{1}{2} - \frac{ix}{2}\right)^{\frac{i\sqrt{-2+2\sqrt{2}}}{2}} (x+i) \left(\frac{1}{2} + \frac{ix}{2}\right)^{i\sqrt{-1+i}} \sqrt{-1+i} \operatorname{hypergeom}\left(\left[\frac{i\sqrt{-2+2\sqrt{2}}}{2}, \frac{i\sqrt{-1+i}}{2} + \frac{\sqrt{1+i}}{2} + 1\right], \left(4\left(\frac{1}{2} - \frac{ix}{2}\right)^{\frac{\sqrt{2+2\sqrt{2}}}{2}} c_1\right.\right.}{\left.4\left(\frac{1}{2} - \frac{ix}{2}\right)^{\frac{\sqrt{2+2\sqrt{2}}}{2}} c_1\right)} dx \right) + c_2$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(1+x^2)*y'[x]+1+(y'[x])^2==x,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

3.23 problem 23

3.23.1 Solving as second order ode missing y ode	1406
3.23.2 Maple step by step solution	1408

Internal problem ID [7213]

Internal file name [OUTPUT/6199_Sunday_June_05_2022_04_31_48_PM_39722752/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 23.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]
```

$$(x^2 + 1)y'' + xy'^2 = 0$$

3.23.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$(x^2 + 1)p'(x) + xp(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= -\frac{xp^2}{x^2 + 1} \end{aligned}$$

Where $f(x) = -\frac{x}{x^2+1}$ and $g(p) = p^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{p^2} dp &= -\frac{x}{x^2+1} dx \\ \int \frac{1}{p^2} dp &= \int -\frac{x}{x^2+1} dx \\ -\frac{1}{p} &= -\frac{\ln(x^2+1)}{2} + c_1\end{aligned}$$

The solution is

$$-\frac{1}{p(x)} + \frac{\ln(x^2+1)}{2} - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$-\frac{1}{y'} + \frac{\ln(x^2+1)}{2} - c_1 = 0$$

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{2}{\ln(x^2+1) - 2c_1} dx \\ &= \int \frac{2}{\ln(x^2+1) - 2c_1} dx + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \frac{2}{\ln(x^2+1) - 2c_1} dx + c_2 \tag{1}$$

Verification of solutions

$$y = \int \frac{2}{\ln(x^2+1) - 2c_1} dx + c_2$$

Verified OK.

3.23.2 Maple step by step solution

Let's solve

$$(x^2 + 1)y'' + xy'^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$(x^2 + 1)u'(x) + xu(x)^2 = 0$$

- Separate variables

$$\frac{u'(x)}{u(x)^2} = -\frac{x}{x^2+1}$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{u(x)^2} dx = \int -\frac{x}{x^2+1} dx + c_1$$

- Evaluate integral

$$-\frac{1}{u(x)} = -\frac{\ln(x^2+1)}{2} + c_1$$

- Solve for $u(x)$

$$u(x) = \frac{2}{\ln(x^2+1)-2c_1}$$

- Solve 1st ODE for $u(x)$

$$u(x) = \frac{2}{\ln(x^2+1)-2c_1}$$

- Make substitution $u = y'$

$$y' = \frac{2}{\ln(x^2+1)-2c_1}$$

- Integrate both sides to solve for y

$$\int y' dx = \int \frac{2}{\ln(x^2+1)-2c_1} dx + c_2$$

- Compute integrals

$$y = \int \frac{2}{\ln(x^2+1)-2c_1} dx + c_2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)^2*_a/(_a^2+1), _b(_a)` *** Su
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve((1+x^2)*diff(y(x),x$2)+1+x*diff(y(x),x)^2=1,y(x), singsol=all)
```

$$y(x) = 2 \left(\int \frac{1}{\ln(x^2 + 1) + 2c_1} dx \right) + c_2$$

✓ Solution by Mathematica

Time used: 60.288 (sec). Leaf size: 33

```
DSolve[(1+x^2)*y'[x]+1+x*(y'[x])^2==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \int_1^x -\frac{2}{2c_1 - \log(K[1]^2 + 1)} dK[1] + c_2$$

3.24 problem 24

Internal problem ID [7214]

Internal file name [OUTPUT/6200_Sunday_June_05_2022_04_31_53_PM_23928755/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 24.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

[NONE]

Unable to solve or complete the solution.

$$(x^2 + 1) y'' + yy'^2 = 0$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integrating f
    --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dynam
    -> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integratin
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular cases
trying symmetries linear in x and y(x)
trying differential order: 2; exact nonlinear
trying 2nd order, integrating factor of the form mu(y)
trying 2nd order, integrating factor of the form mu(x,y)
-> Calling odsolve with the ODE`, -(_y1^2*x^2+_y1^2-4*x^2)*y(x)/((x^2+1)*_y1^2)+(2*(diff(y(x)
    Methods for first order ODEs:
    --- Trying classification methods ---
    trying a quadrature
    trying 1st order linear
    <- 1st order linear successful
trying 2nd order, integrating factor of the form mu(x,y)/(y)^n, only the general case
trying 2nd order, integrating factor of the form mu(y,y)
trying differential order: 2; mu polynomial in y
trying 2nd order, integrating factor of the form mu(x,y)
differential order: 2; looking for linear symmetries
-> trying 2nd order, the S-function method
    -> trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for the S-
    -> trying 2nd order, the S-function method
    -> trying 2nd order, No Point Symmetries Class V
        --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dy
        -> trying 2nd order, No Point Symmetries Class V
    -> trying 2nd order, No Point Symmetries Class V
        --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dy
        -> trying 2nd order, No Point Symmetries Class V
    -> trying 2nd order, No Point Symmetries Class V
        --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dy
        -> trying 2nd order, No Point Symmetries Class V
trying 2nd order, integrating factor of the form mu(x,y)/(y)^n, only the general case
-> trying 2nd order, dynamical_symmetries, only a reduction of order through one integrating
    --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dynam
    -> trying 2nd order, dynamical_symmetries, only a reduction of order through one integrat
    --- Trying Lie symmetry methods, 2nd order ---
```

X Solution by Maple

```
dsolve((1+x^2)*diff(y(x),x$2)+y(x)*diff(y(x),x)^2=0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(1+x^2)*y'[x]+y[x]*(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

3.25 problem 25

3.25.1 Solving as second order ode missing y ode 1413

3.25.2 Maple step by step solution 1415

Internal problem ID [7215]

Internal file name [OUTPUT/6201_Sunday_June_05_2022_04_31_56_PM_84200508/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 25.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]
```

$$(x^2 + 1)y'' + y'^2 = 0$$

3.25.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$(x^2 + 1)p'(x) + p(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= -\frac{p^2}{x^2 + 1} \end{aligned}$$

Where $f(x) = -\frac{1}{x^2+1}$ and $g(p) = p^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{p^2} dp &= -\frac{1}{x^2+1} dx \\ \int \frac{1}{p^2} dp &= \int -\frac{1}{x^2+1} dx \\ -\frac{1}{p} &= -\arctan(x) + c_1\end{aligned}$$

The solution is

$$-\frac{1}{p(x)} + \arctan(x) - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$-\frac{1}{y'} + \arctan(x) - c_1 = 0$$

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{1}{\arctan(x) - c_1} dx \\ &= \int \frac{1}{\arctan(x) - c_1} dx + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \frac{1}{\arctan(x) - c_1} dx + c_2 \quad (1)$$

Verification of solutions

$$y = \int \frac{1}{\arctan(x) - c_1} dx + c_2$$

Verified OK.

3.25.2 Maple step by step solution

Let's solve

$$(x^2 + 1) y'' + y'^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$(x^2 + 1) u'(x) + u(x)^2 = 0$$

- Separate variables

$$\frac{u'(x)}{u(x)^2} = -\frac{1}{x^2+1}$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{u(x)^2} dx = \int -\frac{1}{x^2+1} dx + c_1$$

- Evaluate integral

$$-\frac{1}{u(x)} = -\arctan(x) + c_1$$

- Solve for $u(x)$

$$u(x) = \frac{1}{\arctan(x) - c_1}$$

- Solve 1st ODE for $u(x)$

$$u(x) = \frac{1}{\arctan(x) - c_1}$$

- Make substitution $u = y'$

$$y' = \frac{1}{\arctan(x) - c_1}$$

- Integrate both sides to solve for y

$$\int y' dx = \int \frac{1}{\arctan(x) - c_1} dx + c_2$$

- Compute integrals

$$y = \int \frac{1}{\arctan(x) - c_1} dx + c_2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)^2/(_a^2+1), _b(_a)` *** Suble
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve((1+x^2)*diff(y(x),x$2)+diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = \int \frac{1}{\arctan(x) + c_1} dx + c_2$$

✓ Solution by Mathematica

Time used: 60.278 (sec). Leaf size: 25

```
DSolve[(1+x^2)*y'[x]+(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \int_1^x \frac{1}{\arctan(K[1]) - c_1} dK[1] + c_2$$

3.26 problem 26

3.26.1 Solving as second order ode missing x ode 1417

Internal problem ID [7216]

Internal file name [OUTPUT/6202_Sunday_June_05_2022_04_32_00_PM_14548145/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 26.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible,
    _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]
```

$$y'' + \sin(y) y'^2 = 0$$

3.26.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) + \sin(y) p(y)^2 = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\sin(y)p \end{aligned}$$

Where $f(y) = -\sin(y)$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\sin(y) dy \\ \int \frac{1}{p} dp &= \int -\sin(y) dy \\ \ln(p) &= \cos(y) + c_1 \\ p &= e^{\cos(y)+c_1} \\ &= c_1 e^{\cos(y)} \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1 e^{\cos(y)}$$

Integrating both sides gives

$$\begin{aligned} \int \frac{e^{-\cos(y)}}{c_1} dy &= \int dx \\ \int^y \frac{e^{-\cos(a)}}{c_1} da &= x + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$\int^y \frac{e^{-\cos(a)}}{c_1} da = x + c_2 \quad (1)$$

Verification of solutions

$$\int^y \frac{e^{-\cos(a)}}{c_1} da = x + c_2$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
<- 2nd_order Liouville successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)+sin(y(x))*diff(y(x),x)^2=0,y(x), singsol=all)
```

$$\int^{y(x)} e^{-\cos(-a)} d_a - c_1 x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 1.584 (sec). Leaf size: 111

```
DSolve[y''[x]+y[x]*Sin[y[x]](y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{e^{\sin(K[1]) - \cos(K[1])K[1]}}{c_1} dK[1] \& \right] [x + c_2]$$
$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} -\frac{e^{\sin(K[1]) - \cos(K[1])K[1]}}{c_1} dK[1] \& \right] [x + c_2]$$
$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{e^{\sin(K[1]) - \cos(K[1])K[1]}}{c_1} dK[1] \& \right] [x + c_2]$$

3.27 problem 27

- 3.27.1 Solving as second order ode missing y ode 1420
- 3.27.2 Maple step by step solution 1422

Internal problem ID [7217]

Internal file name [OUTPUT/6203_Sunday_June_05_2022_04_32_03_PM_48103369/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 27.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]
```

$$(x^2 + 1)y'' + y'^3 = 0$$

3.27.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$(x^2 + 1)p'(x) + p(x)^3 = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= -\frac{p^3}{x^2 + 1} \end{aligned}$$

Where $f(x) = -\frac{1}{x^2+1}$ and $g(p) = p^3$. Integrating both sides gives

$$\begin{aligned}\frac{1}{p^3} dp &= -\frac{1}{x^2+1} dx \\ \int \frac{1}{p^3} dp &= \int -\frac{1}{x^2+1} dx \\ -\frac{1}{2p^2} &= -\arctan(x) + c_1\end{aligned}$$

The solution is

$$-\frac{1}{2p(x)^2} + \arctan(x) - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$-\frac{1}{2y'^2} + \arctan(x) - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = -\frac{1}{\sqrt{-2c_1 + 2\arctan(x)}} \quad (1)$$

$$y' = \frac{1}{\sqrt{-2c_1 + 2\arctan(x)}} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned}y &= \int -\frac{1}{\sqrt{-2c_1 + 2\arctan(x)}} dx \\ &= \int -\frac{1}{\sqrt{-2c_1 + 2\arctan(x)}} dx + c_2\end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{1}{\sqrt{-2c_1 + 2\arctan(x)}} dx \\ &= \int \frac{1}{\sqrt{-2c_1 + 2\arctan(x)}} dx + c_3\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int -\frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} dx + c_2 \quad (1)$$

$$y = \int \frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} dx + c_3 \quad (2)$$

Verification of solutions

$$y = \int -\frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} dx + c_2$$

Verified OK.

$$y = \int \frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} dx + c_3$$

Verified OK.

3.27.2 Maple step by step solution

Let's solve

$$(x^2 + 1) y'' + y'^3 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$(x^2 + 1) u'(x) + u(x)^3 = 0$$

- Separate variables

$$\frac{u'(x)}{u(x)^3} = -\frac{1}{x^2+1}$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{u(x)^3} dx = \int -\frac{1}{x^2+1} dx + c_1$$

- Evaluate integral

$$-\frac{1}{2u(x)^2} = -\arctan(x) + c_1$$

- Solve for $u(x)$

$$\left\{ u(x) = \frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}}, u(x) = -\frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} \right\}$$

- Solve 1st ODE for $u(x)$

$$u(x) = \frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}}$$

- Make substitution $u = y'$

$$y' = \frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}}$$

- Integrate both sides to solve for y

$$\int y' dx = \int \frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} dx + c_2$$

- Compute integrals

$$y = \int \frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} dx + c_2$$

- Solve 2nd ODE for $u(x)$

$$u(x) = -\frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}}$$

- Make substitution $u = y'$

$$y' = -\frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}}$$

- Integrate both sides to solve for y

$$\int y' dx = \int -\frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} dx + c_2$$

- Compute integrals

$$y = \int -\frac{1}{\sqrt{-2c_1 + 2 \arctan(x)}} dx + c_2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)^3/(_a^2+1), _b(_a)` *** Suble
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 33

```
dsolve((1+x^2)*diff(y(x),x$2)+diff(y(x),x)^3=0,y(x), singsol=all)
```

$$y(x) = \int \frac{1}{\sqrt{c_1 + 2 \arctan(x)}} dx + c_2$$
$$y(x) = - \left(\int \frac{1}{\sqrt{c_1 + 2 \arctan(x)}} dx \right) + c_2$$

✓ Solution by Mathematica

Time used: 62.161 (sec). Leaf size: 59

```
DSolve[(1+x^2)*y'[x]+y[x]^3==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \int_1^x -\frac{1}{\sqrt{2 \arctan(K[1]) - 2c_1}} dK[1] + c_2$$

$$y(x) \rightarrow \int_1^x \frac{1}{\sqrt{2 \arctan(K[2]) - 2c_1}} dK[2] + c_2$$

3.28 problem 28

3.28.1 Solving as homogeneousTypeD2 ode 1426

3.28.2 Solving as first order ode lie symmetry calculated ode 1428

Internal problem ID [7218]

Internal file name [OUTPUT/6204_Sunday_June_05_2022_04_32_07_PM_6154067/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 28.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**homogeneousTypeD2**", "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _dAlembert]
```

$$y' - e^{-\frac{y}{x}} = 0$$

3.28.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - e^{-u(x)} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{-u + e^{-u}}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = -u + e^{-u}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{-u + e^{-u}} du &= \frac{1}{x} dx \\ \int \frac{1}{-u + e^{-u}} du &= \int \frac{1}{x} dx \\ \int \frac{1}{-a + e^{-a}} da &= \ln(x) + c_2 \end{aligned}$$

Which results in

$$\int^u \frac{1}{-a + e^{-a}} da = \ln(x) + c_2$$

The solution is

$$\int^{u(x)} \frac{1}{-a + e^{-a}} da - \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\int^{\frac{y}{x}} \frac{1}{-a + e^{-a}} da - \ln(x) - c_2 = 0$$

$$\int^{\frac{y}{x}} \frac{1}{-a + e^{-a}} da - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\int^{\frac{y}{x}} \frac{1}{-a + e^{-a}} da - \ln(x) - c_2 = 0 \tag{1}$$




Figure 129: Slope field plot

Verification of solutions

$$\int^{\frac{y}{x}} \frac{1}{-a + e^{-a}} d(-a - \ln(x) - c_2) = 0$$

Verified OK.

3.28.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = e^{-\frac{y}{x}}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (A)$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as ansatz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1E)$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2E)$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + e^{-\frac{y}{x}}(b_3 - a_2) - e^{-\frac{2y}{x}}a_3 - \frac{ye^{-\frac{y}{x}}(xa_2 + ya_3 + a_1)}{x^2} + \frac{e^{-\frac{y}{x}}(xb_2 + yb_3 + b_1)}{x} = 0 \quad (5E)$$

Putting the above in normal form gives

$$\frac{e^{-\frac{2y}{x}}a_3x^2 + e^{-\frac{y}{x}}x^2a_2 - e^{-\frac{y}{x}}x^2b_2 - e^{-\frac{y}{x}}x^2b_3 + e^{-\frac{y}{x}}xya_2 - e^{-\frac{y}{x}}xyb_3 + e^{-\frac{y}{x}}y^2a_3 - e^{-\frac{y}{x}}xb_1 + e^{-\frac{y}{x}}ya_1 - b_2x^2}{x^2} = 0$$

Setting the numerator to zero gives

$$-e^{-\frac{2y}{x}}a_3x^2 - e^{-\frac{y}{x}}x^2a_2 + e^{-\frac{y}{x}}x^2b_2 + e^{-\frac{y}{x}}x^2b_3 - e^{-\frac{y}{x}}xya_2 + e^{-\frac{y}{x}}xyb_3 - e^{-\frac{y}{x}}y^2a_3 + e^{-\frac{y}{x}}xb_1 - e^{-\frac{y}{x}}ya_1 + b_2x^2 = 0 \quad (6E)$$

Simplifying the above gives

$$\begin{aligned} & -e^{-\frac{2y}{x}} a_3 x^2 - e^{-\frac{y}{x}} x^2 a_2 + e^{-\frac{y}{x}} x^2 b_2 + e^{-\frac{y}{x}} x^2 b_3 - e^{-\frac{y}{x}} x y a_2 \\ & + e^{-\frac{y}{x}} x y b_3 - e^{-\frac{y}{x}} y^2 a_3 + e^{-\frac{y}{x}} x b_1 - e^{-\frac{y}{x}} y a_1 + b_2 x^2 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\left\{ x, y, e^{-\frac{2y}{x}}, e^{-\frac{y}{x}} \right\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\left\{ x = v_1, y = v_2, e^{-\frac{2y}{x}} = v_3, e^{-\frac{y}{x}} = v_4 \right\}$$

The above PDE (6E) now becomes

$$\begin{aligned} & -v_4 v_1^2 a_2 - v_4 v_1 v_2 a_2 - v_3 a_3 v_1^2 - v_4 v_2^2 a_3 + v_4 v_1^2 b_2 \\ & + v_4 v_1^2 b_3 + v_4 v_1 v_2 b_3 - v_4 v_2 a_1 + v_4 v_1 b_1 + b_2 v_1^2 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4\}$$

Equation (7E) now becomes

$$-v_3 a_3 v_1^2 + (-a_2 + b_2 + b_3) v_1^2 v_4 + b_2 v_1^2 + (b_3 - a_2) v_1 v_2 v_4 + v_4 v_1 b_1 - v_4 v_2^2 a_3 - v_4 v_2 a_1 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} b_1 &= 0 \\ b_2 &= 0 \\ -a_1 &= 0 \\ -a_3 &= 0 \\ b_3 - a_2 &= 0 \\ -a_2 + b_2 + b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$a_1 = 0$$

$$a_2 = b_3$$

$$a_3 = 0$$

$$b_1 = 0$$

$$b_2 = 0$$

$$b_3 = b_3$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\xi = x$$

$$\eta = y$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Therefore

$$\begin{aligned} \frac{dy}{dx} &= \frac{\eta}{\xi} \\ &= \frac{y}{x} \\ &= \frac{y}{x} \end{aligned}$$

This is easily solved to give

$$y = c_1 x$$

Where now the coordinate R is taken as the constant of integration. Hence

$$R = \frac{y}{x}$$

And S is found from

$$\begin{aligned} dS &= \frac{dx}{\xi} \\ &= \frac{dx}{x} \end{aligned}$$

Integrating gives

$$\begin{aligned} S &= \int \frac{dx}{T} \\ &= \ln(x) \end{aligned}$$

Where the constant of integration is set to zero as we just need one solution. Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = e^{-\frac{y}{x}}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= -\frac{y}{x^2} \\ R_y &= \frac{1}{x} \\ S_x &= \frac{1}{x} \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{x}{x e^{-\frac{y}{x}} - y} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{e^{-R} - R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \int \frac{1}{e^{-R} - R} dR + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(x) = \int \frac{y}{x} \frac{1}{e^{-y/x} - y/x} d_{y/x} + c_1$$

Which simplifies to

$$\ln(x) = \int \frac{y}{x} \frac{1}{e^{-y/x} - y/x} d_{y/x} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = e^{-\frac{y}{x}}$ 	$R = \frac{y}{x}$ $S = \ln(x)$	$\frac{dS}{dR} = \frac{1}{e^{-R} - R}$ 

Summary

The solution(s) found are the following

$$\ln(x) = \int \frac{y}{x} \frac{1}{e^{-y/x} - y/x} d_{y/x} + c_1 \quad (1)$$




Figure 130: Slope field plot

Verification of solutions

$$\ln(x) = \int^x \frac{1}{e^{-a} - a} da + c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)=exp(-y(x)/x),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(- \left(\int^{-z} - \frac{1}{-e^{-a} + a} da \right) + \ln(x) + c_1 \right) x$$

✓ Solution by Mathematica

Time used: 0.166 (sec). Leaf size: 39

```
DSolve[y'[x]==Exp[-y[x]/x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\int_1^{\frac{y(x)}{x}} \frac{e^{K[1]}}{e^{K[1]}K[1] - 1} dK[1] = -\log(x) + c_1, y(x) \right]$$

3.29 problem 29

- 3.29.1 Solving as homogeneousTypeD ode 1435
- 3.29.2 Solving as homogeneousTypeD2 ode 1437
- 3.29.3 Solving as first order ode lie symmetry lookup ode 1439

Internal problem ID [7219]

Internal file name [OUTPUT/6205_Sunday_June_05_2022_04_32_08_PM_80973960/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 29.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"homogeneousTypeD", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_homogeneous , `class D`]]
```

$$y' - 2x^2 \sin\left(\frac{y}{x}\right)^2 - \frac{y}{x} = 0$$

3.29.1 Solving as homogeneousTypeD ode

Writing the ode as

$$y' = 2x^2 \sin\left(\frac{y}{x}\right)^2 + \frac{y}{x} \tag{A}$$

The given ode has the form

$$y' = \frac{y}{x} + g(x) f\left(b\frac{y}{x}\right)^{\frac{n}{m}} \tag{1}$$

Where b is scalar and $g(x)$ is function of x and n, m are integers. The solution is given in Kamke page 20. Using the substitution $y(x) = u(x)x$ then

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Hence the given ode becomes

$$\begin{aligned}\frac{du}{dx}x + u &= u + g(x) f(bu)^{\frac{n}{m}} \\ u' &= \frac{1}{x}g(x) f(bu)^{\frac{n}{m}}\end{aligned}\quad (2)$$

The above ode is always separable. This is easily solved for u assuming the integration can be resolved, and then the solution to the original ode becomes $y = ux$. Comparing the given ode (A) with the form (1) shows that

$$\begin{aligned}g(x) &= 2x^2 \\ b &= 1 \\ f\left(\frac{bx}{y}\right) &= \sin\left(\frac{y}{x}\right)\end{aligned}$$

Substituting the above in (2) results in the $u(x)$ ode as

$$u'(x) = 2x \sin(u(x))^2$$

Which is now solved as separable In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= 2x \sin(u)^2\end{aligned}$$

Where $f(x) = 2x$ and $g(u) = \sin(u)^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\sin(u)^2} du &= 2x dx \\ \int \frac{1}{\sin(u)^2} du &= \int 2x dx \\ -\cot(u) &= x^2 + c_1\end{aligned}$$

The solution is

$$-\cot(u(x)) - x^2 - c_1 = 0$$

Therefore the solution is found using $y = ux$. Hence

$$-\cot\left(\frac{y}{x}\right) - x^2 - c_1 = 0$$

Summary

The solution(s) found are the following

$$-\cot\left(\frac{y}{x}\right) - x^2 - c_1 = 0 \quad (1)$$




Figure 131: Slope field plot

Verification of solutions

$$-\cot\left(\frac{y}{x}\right) - x^2 - c_1 = 0$$

Verified OK.

3.29.2 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x - 2x^2 \sin(u(x))^2 = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= 2 \sin(u)^2 x \end{aligned}$$

Where $f(x) = 2x$ and $g(u) = \sin(u)^2$. Integrating both sides gives

$$\frac{1}{\sin(u)^2} du = 2x dx$$

$$\int \frac{1}{\sin(u)^2} du = \int 2x dx$$

$$-\cot(u) = x^2 + c_2$$

The solution is

$$-\cot(u(x)) - x^2 - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$-\cot\left(\frac{y}{x}\right) - x^2 - c_2 = 0$$

$$-\cot\left(\frac{y}{x}\right) - x^2 - c_2 = 0$$

Summary

The solution(s) found are the following

$$-\cot\left(\frac{y}{x}\right) - x^2 - c_2 = 0 \tag{1}$$




Figure 132: Slope field plot

Verification of solutions

$$-\cot\left(\frac{y}{x}\right) - x^2 - c_2 = 0$$

Verified OK.

3.29.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{2x^3 \sin\left(\frac{y}{x}\right)^2 + y}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **homogeneous Type D**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 145: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= x^2 \\ \eta(x, y) &= xy\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Therefore

$$\begin{aligned}\frac{dy}{dx} &= \frac{\eta}{\xi} \\ &= \frac{xy}{x^2} \\ &= \frac{y}{x}\end{aligned}$$

This is easily solved to give

$$y = c_1 x$$

Where now the coordinate R is taken as the constant of integration. Hence

$$R = \frac{y}{x}$$

And S is found from

$$\begin{aligned}dS &= \frac{dx}{\xi} \\ &= \frac{dx}{x^2}\end{aligned}$$

Integrating gives

$$\begin{aligned}S &= \int \frac{dx}{T} \\ &= -\frac{1}{x}\end{aligned}$$

Where the constant of integration is set to zero as we just need one solution. Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2x^3 \sin\left(\frac{y}{x}\right)^2 + y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= -\frac{y}{x^2} \\ R_y &= \frac{1}{x} \\ S_x &= \frac{1}{x^2} \\ S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{\csc\left(\frac{y}{x}\right)^2}{2x^3} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{\csc(R)^2 S(R)^3}{2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives these solutions

$$S(R) = \frac{\sqrt{(c_1 \tan(R) - 1) \tan(R)}}{c_1 \tan(R) - 1} \quad (4)$$

$$S(R) = -\frac{\sqrt{(c_1 \tan(R) - 1) \tan(R)}}{c_1 \tan(R) - 1}$$

Each will now be processed. To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{1}{x} = \frac{\sqrt{(c_1 \tan\left(\frac{y}{x}\right) - 1) \tan\left(\frac{y}{x}\right)}}{c_1 \tan\left(\frac{y}{x}\right) - 1}$$



Which simplifies to

$$-\frac{1}{x} = \frac{\sqrt{(c_1 \tan\left(\frac{y}{x}\right) - 1) \tan\left(\frac{y}{x}\right)}}{c_1 \tan\left(\frac{y}{x}\right) - 1}$$

Which gives

$$y = \arctan\left(\frac{1}{-x^2 + c_1}\right) x$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2x^3 \sin\left(\frac{y}{x}\right)^2 + y}{x}$ 	$R = \frac{y}{x}$ $S = -\frac{1}{x}$	$\frac{dS}{dR} = -\frac{\csc(R)^2 S(R)^3}{2}$ 

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{1}{x} = -\frac{\sqrt{(c_1 \tan\left(\frac{y}{x}\right) - 1) \tan\left(\frac{y}{x}\right)}}{c_1 \tan\left(\frac{y}{x}\right) - 1}$$



Which simplifies to

$$-\frac{1}{x} = -\frac{\sqrt{(c_1 \tan\left(\frac{y}{x}\right) - 1) \tan\left(\frac{y}{x}\right)}}{c_1 \tan\left(\frac{y}{x}\right) - 1}$$

Which gives

$$y = \arctan\left(\frac{1}{-x^2 + c_1}\right) x$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2x^3 \sin\left(\frac{y}{x}\right)^2 + y}{x}$ 	$R = \frac{y}{x}$ $S = -\frac{1}{x}$	$\frac{dS}{dR} = -\frac{\csc(R)^2 S(R)^3}{2}$ 

Summary

The solution(s) found are the following

$$y = \arctan\left(\frac{1}{-x^2 + c_1}\right) x \quad (1)$$

$$y = \arctan\left(\frac{1}{-x^2 + c_1}\right) x \quad (2)$$




Figure 133: Slope field plot

Verification of solutions

$$y = \arctan \left(\frac{1}{-x^2 + c_1} \right) x$$

Verified OK.

$$y = \arctan \left(\frac{1}{-x^2 + c_1} \right) x$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)= 2*x^2 * sin(y(x)/x)^2 + y(x)/x,y(x), singsol=all)
```

$$y(x) = \left(\frac{\pi}{2} + \arctan(x^2 + 2c_1) \right) x$$

✓ Solution by Mathematica

Time used: 0.341 (sec). Leaf size: 22

```
DSolve[y'[x]== 2*x^2 * Sin[y[x]/x]^2 + y[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \cot^{-1}(x^2 - 2c_1)$$
$$y(x) \rightarrow 0$$

3.30 problem 30

3.30.1 Solving as second order euler ode ode	1447
3.30.2 Solving using Kovacic algorithm	1451

Internal problem ID [7220]

Internal file name [OUTPUT/6206_Sunday_June_05_2022_04_32_13_PM_27191802/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 30.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$4x^2y'' + y = 8\sqrt{x}(1 + \ln(x))$$

3.30.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 4x^2$, $B = 0$, $C = 1$, $f(x) = 8\sqrt{x}(1 + \ln(x))$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$4x^2y'' + y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$4x^2(r(r-1))x^{r-2} + 0rx^{r-1} + x^r = 0$$

Simplifying gives

$$4r(r-1)x^r + 0x^r + x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$4r(r-1) + 0 + 1 = 0$$

Or

$$4r^2 - 4r + 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2}$$
$$r_2 = \frac{1}{2}$$

Since the roots are equal, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^r$ and $y_2 = x^r \ln(x)$. Hence

$$y = c_1\sqrt{x} + c_2\sqrt{x} \ln(x)$$

Next, we find the particular solution to the ODE

$$4x^2y'' + y = 8\sqrt{x}(1 + \ln(x))$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x}$$
$$y_2 = \sqrt{x} \ln(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} & \sqrt{x} \ln(x) \\ \frac{d}{dx}(\sqrt{x}) & \frac{d}{dx}(\sqrt{x} \ln(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} & \sqrt{x} \ln(x) \\ \frac{1}{2\sqrt{x}} & \frac{\ln(x)}{2\sqrt{x}} + \frac{1}{\sqrt{x}} \end{vmatrix}$$

Therefore

$$W = (\sqrt{x}) \left(\frac{\ln(x)}{2\sqrt{x}} + \frac{1}{\sqrt{x}} \right) - (\sqrt{x} \ln(x)) \left(\frac{1}{2\sqrt{x}} \right)$$

Which simplifies to

$$W = 1$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{8x \ln(x) (1 + \ln(x))}{4x^2} dx$$

Which simplifies to

$$u_1 = - \int \frac{2 \ln(x) (1 + \ln(x))}{x} dx$$

Hence

$$u_1 = -\frac{2 \ln(x)^3}{3} - \ln(x)^2$$

And Eq. (3) becomes

$$u_2 = \int \frac{8x(1 + \ln(x))}{4x^2} dx$$

Which simplifies to

$$u_2 = \int \frac{2 + 2 \ln(x)}{x} dx$$

Hence

$$u_2 = \ln(x)^2 + 2 \ln(x)$$

Which simplifies to

$$u_1 = -\frac{2 \ln(x)^3}{3} - \ln(x)^2$$
$$u_2 = \ln(x)(\ln(x) + 2)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{2 \ln(x)^3}{3} - \ln(x)^2 \right) \sqrt{x} + \ln(x)^2 (\ln(x) + 2) \sqrt{x}$$

Which simplifies to

$$y_p(x) = \frac{\ln(x)^2 (\ln(x) + 3) \sqrt{x}}{3}$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= \left(\frac{\ln(x)^3}{3} + \ln(x)^2 + c_1 + c_2 \ln(x) \right) \sqrt{x}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{\ln(x)^3}{3} + \ln(x)^2 + c_1 + c_2 \ln(x) \right) \sqrt{x} \quad (1)$$

Verification of solutions

$$y = \left(\frac{\ln(x)^3}{3} + \ln(x)^2 + c_1 + c_2 \ln(x) \right) \sqrt{x}$$

Verified OK.

3.30.2 Solving using Kovacic algorithm

Writing the ode as

$$4x^2y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 4x^2 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 147: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to

determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_{∞}^{\pm} . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_{\infty}^{-} = \frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \left(\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{1}{2x} + (-)(0) \\ &= \frac{1}{2x} \\ &= \frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{2x}\right)(0) + \left(\left(-\frac{1}{2x^2}\right) + \left(\frac{1}{2x}\right)^2 - \left(-\frac{1}{4x^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned}z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{1}{2x} dx} \\ &= \sqrt{x}\end{aligned}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \sqrt{x}\end{aligned}$$

Which simplifies to

$$y_1 = \sqrt{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \sqrt{x} \int \frac{1}{x} dx \\ &= \sqrt{x}(\ln(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\sqrt{x}) + c_2(\sqrt{x}(\ln(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$4x^2y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1\sqrt{x} + c_2\sqrt{x} \ln(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x}$$

$$y_2 = \sqrt{x} \ln(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} & \sqrt{x} \ln(x) \\ \frac{d}{dx}(\sqrt{x}) & \frac{d}{dx}(\sqrt{x} \ln(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} & \sqrt{x} \ln(x) \\ \frac{1}{2\sqrt{x}} & \frac{\ln(x)}{2\sqrt{x}} + \frac{1}{\sqrt{x}} \end{vmatrix}$$

Therefore

$$W = (\sqrt{x}) \left(\frac{\ln(x)}{2\sqrt{x}} + \frac{1}{\sqrt{x}} \right) - (\sqrt{x} \ln(x)) \left(\frac{1}{2\sqrt{x}} \right)$$

Which simplifies to

$$W = 1$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{8x \ln(x) (1 + \ln(x))}{4x^2} dx$$

Which simplifies to

$$u_1 = - \int \frac{2 \ln(x) (1 + \ln(x))}{x} dx$$

Hence

$$u_1 = - \frac{2 \ln(x)^3}{3} - \ln(x)^2$$

And Eq. (3) becomes

$$u_2 = \int \frac{8x(1 + \ln(x))}{4x^2} dx$$

Which simplifies to

$$u_2 = \int \frac{2 + 2 \ln(x)}{x} dx$$

Hence

$$u_2 = \ln(x)^2 + 2 \ln(x)$$

Which simplifies to

$$u_1 = -\frac{2 \ln(x)^3}{3} - \ln(x)^2$$

$$u_2 = \ln(x) (\ln(x) + 2)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{2 \ln(x)^3}{3} - \ln(x)^2 \right) \sqrt{x} + \ln(x)^2 (\ln(x) + 2) \sqrt{x}$$

Which simplifies to

$$y_p(x) = \frac{\ln(x)^2 (\ln(x) + 3) \sqrt{x}}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \sqrt{x} + c_2 \sqrt{x} \ln(x)) + \left(\frac{\ln(x)^2 (\ln(x) + 3) \sqrt{x}}{3} \right) \end{aligned}$$

Which simplifies to

$$y = (c_2 \ln(x) + c_1) \sqrt{x} + \frac{\ln(x)^2 (\ln(x) + 3) \sqrt{x}}{3}$$

Summary

The solution(s) found are the following

$$y = (c_2 \ln(x) + c_1) \sqrt{x} + \frac{\ln(x)^2 (\ln(x) + 3) \sqrt{x}}{3} \quad (1)$$

Verification of solutions

$$y = (c_2 \ln(x) + c_1) \sqrt{x} + \frac{\ln(x)^2 (\ln(x) + 3) \sqrt{x}}{3}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    <- LODE of Euler type successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(4*x^2*diff(y(x),x$2)+ y(x) = 8*sqrt(x)*(1+ln(x)),y(x), singsol=all)
```

$$y(x) = \left(c_2 + \ln(x) c_1 + \frac{\ln(x)^3}{3} + \ln(x)^2 \right) \sqrt{x}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 37

```
DSolve[4*x^2*y''[x]+y[x] == 8*Sqrt[x]*(1+Log[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6} \sqrt{x} (2 \log^3(x) + 6 \log^2(x) + 3 c_2 \log(x) + 6 c_1)$$

3.31 problem 31

- 3.31.1 Solving as first order ode lie symmetry lookup ode 1460
- 3.31.2 Solving as bernoulli ode 1463
- 3.31.3 Solving as exact ode 1466

Internal problem ID [7221]

Internal file name [OUTPUT/6207_Sunday_June_05_2022_04_32_15_PM_33122274/index.tex]

Book: Own collection of miscellaneous problems

Section: section 3.0

Problem number: 31.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "bernoulli", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_rational, _Bernoulli]`

$$vv' - \frac{2v^2}{r^3} = \frac{\lambda r}{3}$$

3.31.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$v' = \frac{\lambda r^4 + 6v^2}{3r^3v}$$
$$v' = \omega(r, v)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_r + \omega(\eta_v - \xi_r) - \omega^2 \xi_v - \omega_r \xi - \omega_v \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 148: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(r, v) &= 0 \\ \eta(r, v) &= \frac{e^{-\frac{2}{r^2}}}{v}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(r, v) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dr}{\xi} = \frac{dv}{\eta} = dS\tag{1}$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial r} + \eta \frac{\partial}{\partial v}) S(r, v) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = r$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{e^{-\frac{2}{r^2}}}{v}} dy \end{aligned}$$

Which results in

$$S = \frac{v^2 e^{\frac{2}{r^2}}}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_r + \omega(r, v)S_v}{R_r + \omega(r, v)R_v} \quad (2)$$

Where in the above R_r, R_v, S_r, S_v are all partial derivatives and $\omega(r, v)$ is the right hand side of the original ode given by

$$\omega(r, v) = \frac{\lambda r^4 + 6v^2}{3r^3v}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_r &= 1 \\ R_v &= 0 \\ S_r &= -\frac{2v^2 e^{\frac{2}{r^2}}}{r^3} \\ S_v &= v e^{\frac{2}{r^2}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{e^{\frac{2}{R^2}} \lambda R}{3} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for r, v in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{e^{\frac{2}{R^2}} \lambda R}{3}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{\lambda \left(\frac{e^{\frac{2}{R^2}} R^2}{2} + \text{expIntegral}_1 \left(-\frac{2}{R^2} \right) \right)}{3} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to r, v coordinates. This results in

$$\frac{v^2 e^{\frac{2}{r^2}}}{2} = \frac{\lambda \left(\frac{e^{\frac{2}{r^2}} r^2}{2} + \text{expIntegral}_1 \left(-\frac{2}{r^2} \right) \right)}{3} + c_1$$

Which simplifies to

$$\frac{v^2 e^{\frac{2}{r^2}}}{2} = \frac{\lambda \left(\frac{e^{\frac{2}{r^2}} r^2}{2} + \text{expIntegral}_1 \left(-\frac{2}{r^2} \right) \right)}{3} + c_1$$

Summary

The solution(s) found are the following

$$\frac{v^2 e^{\frac{2}{r^2}}}{2} = \frac{\lambda \left(\frac{e^{\frac{2}{r^2}} r^2}{2} + \text{expIntegral}_1 \left(-\frac{2}{r^2} \right) \right)}{3} + c_1 \quad (1)$$

Verification of solutions

$$\frac{v^2 e^{\frac{2}{r^2}}}{2} = \frac{\lambda \left(\frac{e^{\frac{2}{r^2}} r^2}{2} + \text{expIntegral}_1 \left(-\frac{2}{r^2} \right) \right)}{3} + c_1$$

Verified OK.

3.31.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} v' &= F(r, v) \\ &= \frac{\lambda r^4 + 6v^2}{3r^3 v} \end{aligned}$$

This is a Bernoulli ODE.

$$v' = \frac{2}{r^3} v + \frac{\lambda r}{3} \frac{1}{v} \quad (1)$$

The standard Bernoulli ODE has the form

$$v' = f_0(r)v + f_1(r)v^n \quad (2)$$

The first step is to divide the above equation by v^n which gives

$$\frac{v'}{v^n} = f_0(r)v^{1-n} + f_1(r) \quad (3)$$

The next step is use the substitution $w = v^{1-n}$ in equation (3) which generates a new ODE in $w(r)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $v(r)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(r) &= \frac{2}{r^3} \\ f_1(r) &= \frac{\lambda r}{3} \\ n &= -1 \end{aligned}$$

Dividing both sides of ODE (1) by $v^n = \frac{1}{v}$ gives

$$v'v = \frac{2v^2}{r^3} + \frac{\lambda r}{3} \quad (4)$$

Let

$$\begin{aligned} w &= v^{1-n} \\ &= v^2 \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t r gives

$$w' = 2vv' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} \frac{w'(r)}{2} &= \frac{2w(r)}{r^3} + \frac{\lambda r}{3} \\ w' &= \frac{4w}{r^3} + \frac{2\lambda r}{3} \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(r)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(r) + p(r)w(r) = q(r)$$

Where here

$$p(r) = -\frac{4}{r^3}$$
$$q(r) = \frac{2\lambda r}{3}$$

Hence the ode is

$$w'(r) - \frac{4w(r)}{r^3} = \frac{2\lambda r}{3}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{4}{r^3} dr}$$
$$= e^{\frac{2}{r^2}}$$

The ode becomes

$$\frac{d}{dr}(\mu w) = (\mu) \left(\frac{2\lambda r}{3} \right)$$
$$\frac{d}{dr} \left(e^{\frac{2}{r^2}} w \right) = \left(e^{\frac{2}{r^2}} \right) \left(\frac{2\lambda r}{3} \right)$$
$$d \left(e^{\frac{2}{r^2}} w \right) = \left(\frac{2 e^{\frac{2}{r^2}} \lambda r}{3} \right) dr$$

Integrating gives

$$e^{\frac{2}{r^2}} w = \int \frac{2 e^{\frac{2}{r^2}} \lambda r}{3} dr$$
$$e^{\frac{2}{r^2}} w = \frac{2\lambda \left(\frac{e^{\frac{2}{r^2}} r^2}{2} + \text{expIntegral}_1 \left(-\frac{2}{r^2} \right) \right)}{3} + c_1$$

Dividing both sides by the integrating factor $\mu = e^{\frac{2}{r^2}}$ results in

$$w(r) = \frac{2 e^{-\frac{2}{r^2}} \lambda \left(\frac{e^{\frac{2}{r^2}} r^2}{2} + \text{expIntegral}_1 \left(-\frac{2}{r^2} \right) \right)}{3} + c_1 e^{-\frac{2}{r^2}}$$

which simplifies to

$$w(r) = \frac{\lambda r^2}{3} + \frac{2 e^{-\frac{2}{r^2}} \lambda \text{expIntegral}_1 \left(-\frac{2}{r^2} \right)}{3} + c_1 e^{-\frac{2}{r^2}}$$

Replacing w in the above by v^2 using equation (5) gives the final solution.

$$v^2 = \frac{\lambda r^2}{3} + \frac{2 e^{-\frac{2}{r^2}} \lambda \exp \text{Integral}_1 \left(-\frac{2}{r^2}\right)}{3} + c_1 e^{-\frac{2}{r^2}}$$

Solving for v gives

$$v(r) = \frac{\sqrt{6 e^{-\frac{2}{r^2}} \lambda \exp \text{Integral}_1 \left(-\frac{2}{r^2}\right) + 3 \lambda r^2 + 9 c_1 e^{-\frac{2}{r^2}}}}{3}$$

$$v(r) = -\frac{\sqrt{6 e^{-\frac{2}{r^2}} \lambda \exp \text{Integral}_1 \left(-\frac{2}{r^2}\right) + 3 \lambda r^2 + 9 c_1 e^{-\frac{2}{r^2}}}}{3}$$

Summary

The solution(s) found are the following

$$v = \frac{\sqrt{6 e^{-\frac{2}{r^2}} \lambda \exp \text{Integral}_1 \left(-\frac{2}{r^2}\right) + 3 \lambda r^2 + 9 c_1 e^{-\frac{2}{r^2}}}}{3} \quad (1)$$

$$v = -\frac{\sqrt{6 e^{-\frac{2}{r^2}} \lambda \exp \text{Integral}_1 \left(-\frac{2}{r^2}\right) + 3 \lambda r^2 + 9 c_1 e^{-\frac{2}{r^2}}}}{3} \quad (2)$$

Verification of solutions

$$v = \frac{\sqrt{6 e^{-\frac{2}{r^2}} \lambda \exp \text{Integral}_1 \left(-\frac{2}{r^2}\right) + 3 \lambda r^2 + 9 c_1 e^{-\frac{2}{r^2}}}}{3}$$

Verified OK.

$$v = -\frac{\sqrt{6 e^{-\frac{2}{r^2}} \lambda \exp \text{Integral}_1 \left(-\frac{2}{r^2}\right) + 3 \lambda r^2 + 9 c_1 e^{-\frac{2}{r^2}}}}{3}$$

Verified OK.

3.31.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(r, v) dr + N(r, v) dv = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (v) dv &= \left(\frac{2v^2}{r^3} + \frac{\lambda r}{3} \right) dr \\ \left(-\frac{2v^2}{r^3} - \frac{\lambda r}{3} \right) dr + (v) dv &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(r, v) &= -\frac{2v^2}{r^3} - \frac{\lambda r}{3} \\ N(r, v) &= v \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial v} = \frac{\partial N}{\partial r}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial v} &= \frac{\partial}{\partial v} \left(-\frac{2v^2}{r^3} - \frac{\lambda r}{3} \right) \\ &= -\frac{4v}{r^3} \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial r} &= \frac{\partial}{\partial r}(v) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial v} \neq \frac{\partial N}{\partial r}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial v} - \frac{\partial N}{\partial r} \right) \\ &= \frac{1}{v} \left(\left(-\frac{4v}{r^3} \right) - (0) \right) \\ &= -\frac{4}{r^3}\end{aligned}$$

Since A does not depend on v , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A \, dr} \\ &= e^{\int -\frac{4}{r^3} \, dr}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\frac{2}{r^2}} \\ &= e^{\frac{2}{r^2}}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^{\frac{2}{r^2}} \left(-\frac{2v^2}{r^3} - \frac{\lambda r}{3} \right) \\ &= -\frac{e^{\frac{2}{r^2}} (\lambda r^4 + 6v^2)}{3r^3}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{\frac{2}{r^2}}(v) \\ &= v e^{\frac{2}{r^2}}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dv}{dr} &= 0 \\ \left(-\frac{e^{\frac{2}{r^2}}(\lambda r^4 + 6v^2)}{3r^3} \right) + \left(v e^{\frac{2}{r^2}} \right) \frac{dv}{dr} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(r, v)$

$$\frac{\partial \phi}{\partial r} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial v} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. r gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial r} dr &= \int \overline{M} dr \\ \int \frac{\partial \phi}{\partial r} dr &= \int -\frac{e^{\frac{2}{r^2}}(\lambda r^4 + 6v^2)}{3r^3} dr \\ \phi &= -\frac{\lambda \exp \text{Integral}_1 \left(-\frac{2}{r^2} \right)}{3} - \frac{e^{\frac{2}{r^2}}(\lambda r^2 - 3v^2)}{6} + f(v) \end{aligned} \quad (3)$$

Where $f(v)$ is used for the constant of integration since ϕ is a function of both r and v . Taking derivative of equation (3) w.r.t v gives

$$\frac{\partial \phi}{\partial v} = v e^{\frac{2}{r^2}} + f'(v) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial v} = v e^{\frac{2}{r^2}}$. Therefore equation (4) becomes

$$v e^{\frac{2}{r^2}} = v e^{\frac{2}{r^2}} + f'(v) \quad (5)$$

Solving equation (5) for $f'(v)$ gives

$$f'(v) = 0$$

Therefore

$$f(v) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(v)$ into equation (3) gives ϕ

$$\phi = -\frac{\lambda \exp\text{Integral}_1\left(-\frac{2}{r^2}\right)}{3} - \frac{e^{\frac{2}{r^2}}(\lambda r^2 - 3v^2)}{6} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{\lambda \exp\text{Integral}_1\left(-\frac{2}{r^2}\right)}{3} - \frac{e^{\frac{2}{r^2}}(\lambda r^2 - 3v^2)}{6}$$

Summary

The solution(s) found are the following

$$-\frac{\lambda \exp\text{Integral}_1\left(-\frac{2}{r^2}\right)}{3} - \frac{e^{\frac{2}{r^2}}(\lambda r^2 - 3v^2)}{6} = c_1 \quad (1)$$

Verification of solutions

$$-\frac{\lambda \exp\text{Integral}_1\left(-\frac{2}{r^2}\right)}{3} - \frac{e^{\frac{2}{r^2}}(\lambda r^2 - 3v^2)}{6} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 97

```
dsolve(v(r)*diff(v(r),r)=2*v(r)^2/r^3+1/3*lambda*r,v(r), singsol=all)
```

$$v(r) = -\frac{\sqrt{3} \sqrt{e^{-\frac{2}{r^2}} \left(\lambda e^{\frac{2}{r^2}} r^2 + 2\lambda \operatorname{expIntegral}_1 \left(-\frac{2}{r^2} \right) + 3c_1 \right) e^{-\frac{2}{r^2}}}}{3}$$
$$v(r) = \frac{\sqrt{3} \sqrt{e^{-\frac{2}{r^2}} \left(\lambda e^{\frac{2}{r^2}} r^2 + 2\lambda \operatorname{expIntegral}_1 \left(-\frac{2}{r^2} \right) + 3c_1 \right) e^{-\frac{2}{r^2}}}}{3}$$

✓ Solution by Mathematica

Time used: 10.758 (sec). Leaf size: 98

```
DSolve[v[r]*v'[r]==2*v[r]^2/r^3+1/3*\[Lambda]*r,v[r],r,IncludeSingularSolutions -> True]
```

$$v(r) \rightarrow -\frac{\sqrt{e^{-\frac{2}{r^2}} \left(-2\lambda \operatorname{ExpIntegralEi} \left(\frac{2}{r^2} \right) + \lambda e^{\frac{2}{r^2}} r^2 + 3c_1 \right)}}{\sqrt{3}}$$
$$v(r) \rightarrow \frac{\sqrt{e^{-\frac{2}{r^2}} \left(-2\lambda \operatorname{ExpIntegralEi} \left(\frac{2}{r^2} \right) + \lambda e^{\frac{2}{r^2}} r^2 + 3c_1 \right)}}{\sqrt{3}}$$

4 section 4.0

4.1	problem 1	1474
4.2	problem 2	1486
4.3	problem 3	1499
4.4	problem 4	1511
4.5	problem 5	1522
4.6	problem 6	1534
4.7	problem 7	1547
4.8	problem 8	1561
4.9	problem 9	1574
4.10	problem 10	1585
4.11	problem 11	1597
4.12	problem 12	1611
4.13	problem 13	1623
4.14	problem 14	1634
4.15	problem 15	1648
4.16	problem 16	1663
4.17	problem 17	1675
4.18	problem 18	1689
4.19	problem 19	1704
4.20	problem 20	1717
4.21	problem 21	1727
4.22	problem 22	1737
4.23	problem 23	1751
4.24	problem 24	1766
4.25	problem 24	1781
4.26	problem 24	1798
4.27	problem 24	1812
4.28	problem 24	1829
4.29	problem 25	1841
4.30	problem 26	1856
4.31	problem 27	1870
4.32	problem 28	1884
4.33	problem 29	1887
4.34	problem 31	1898
4.35	problem 32	1907
4.36	problem 33	1918
4.37	problem 34	1931

4.38	problem 35	1942
4.39	problem 36	1954
4.40	problem 37	1970
4.41	problem 38	1982
4.42	problem 39	1995
4.43	problem 40	2010
4.44	problem 41	2025
4.45	problem 42	2039
4.46	problem 43	2054
4.47	problem 44	2069
4.48	problem 45	2079
4.49	problem 46	2092
4.50	problem 47	2103
4.51	problem 48	2116
4.52	problem 49	2130
4.53	problem 50	2145
4.54	problem 51	2161
4.55	problem 52	2171
4.56	problem 53	2184
4.57	problem 54	2197
4.58	problem 55	2211
4.59	problem 56	2222
4.60	problem 57	2231
4.61	problem 58	2243
4.62	problem 59	2258
4.63	problem 60	2268
4.64	problem 61	2272
4.65	problem 62	2277
4.66	problem 63	2281
4.67	problem 64	2288
4.68	problem 65	2292
4.69	problem 66	2302
4.70	problem 67	2312
4.71	problem 68	2324
4.72	problem 69	2335

4.1 problem 1

4.1.1 Maple step by step solution 1482

Internal problem ID [7222]

Internal file name [OUTPUT/6208_Sunday_June_05_2022_04_32_18_PM_40583659/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 1.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 150: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \quad (1)$$

Verification of solutions

$$y = c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

4.1.1 Maple step by step solution

Let's solve

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{(x^2-1)y}{2x^2} + \frac{y'}{2x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{2x} - \frac{(x^2-1)y}{2x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{1}{2x}, P_3(x) = -\frac{x^2-1}{2x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -\frac{1}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = \frac{1}{2}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-1+2r)(-1+r)x^r + a_1(1+2r)rx^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r-1)(k+r-1) - a_{k-2})x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(-1+2r)(-1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 1, \frac{1}{2} \right\}$$

- Each term must be 0

$$a_1(1+2r)r = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$2(k+r-1)\left(k+r-\frac{1}{2}\right)a_k - a_{k-2} = 0$$

- Shift index using $k \rightarrow k+2$

$$2(k+1+r)\left(k+\frac{3}{2}+r\right)a_{k+2} - a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = \frac{a_k}{(k+1+r)(2k+3+2r)}$$

- Recursion relation for $r = 1$

$$a_{k+2} = \frac{a_k}{(k+2)(2k+5)}$$

- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+2} = \frac{a_k}{(k+2)(2k+5)}, a_1 = 0 \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = \frac{a_k}{\left(k+\frac{3}{2}\right)(2k+4)}$$

- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = \frac{a_k}{\left(k+\frac{3}{2}\right)(2k+4)}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = \frac{a_k}{(k+2)(2k+5)}, a_1 = 0, b_{k+2} = \frac{b_k}{\left(k+\frac{3}{2}\right)(2k+4)}, b_1 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
Order:=6;  
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 48

```
AsymptoticDSolveValue[2*x^2*y'[x] - x*y'[x] + (1-x^2)*y[x] == 0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1x \left(\frac{x^4}{360} + \frac{x^2}{10} + 1 \right) + c_2\sqrt{x} \left(\frac{x^4}{168} + \frac{x^2}{6} + 1 \right)$$

4.2 problem 2

Internal problem ID [7223]

Internal file name [OUTPUT/6209_Sunday_June_05_2022_04_32_21_PM_99689811/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = 1$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 152: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = 1$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= 1 \\ m &= 0 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+0} \end{aligned}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = 1$
$c_1 = 0$
$c_2 = \frac{1}{3}$
$c_3 = 0$
$c_4 = \frac{1}{63}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= 1 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\ &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4\end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = 1 + \frac{x^2}{3} + \frac{x^4}{63} + O(x^6)$$

Hence the final solution is

$$\begin{aligned}y &= y_h + y_p \\ &= 1 + \frac{x^2}{3} + \frac{x^4}{63} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= 1 + \frac{x^2}{3} + \frac{x^4}{63} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ &\quad + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)\end{aligned}\tag{1}$$

Verification of solutions

$$y = 1 + \frac{x^2}{3} + \frac{x^4}{63} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
  checking if the LODE has constant coefficients  
  checking if the LODE is of Euler type  
  trying a symmetry of the form [xi=0, eta=F(x)]  
  checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
  -> Bessel  
    <- Bessel successful  
  <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
Order:=6;  
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = 1,y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) \\ + c_2x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 176

```
AsymptoticDSolveValue[2*x^2*y''[x] - x*y'[x] + (1-x^2)*y[x] ==1,y[x],{x,0,5}]
```

$y(x)$

$$\begin{aligned} &\rightarrow c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ &+ c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{154440} - \frac{x^{7/2}}{1260} - \frac{x^{3/2}}{15} \right. \\ &\left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^5}{55440} + \frac{x^3}{504} + \frac{x}{6} - \frac{1}{x} \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \end{aligned}$$

4.3 problem 3

Internal problem ID [7224]

Internal file name [OUTPUT/6210_Sunday_June_05_2022_04_32_23_PM_43528752/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$2x^2y'' - xy' + (1 - x^2)y = 1 + x$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 153: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = 1 + x$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r)(n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r(-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r(-1 + r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1 + x$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. In order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$ $a_2 = \frac{a_0}{2r^2+5r+3}$ $a_3 = 0$ $a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$ $a_5 = 0$
--

Since the $F = 1 + x$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$c_0 = 1$$

$$m = 0$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+0}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$$\begin{array}{l}
 c_0 = 1 \\
 c_1 = 0 \\
 c_2 = \frac{1}{3} \\
 c_3 = 0 \\
 c_4 = \frac{1}{63} \\
 c_5 = 0
 \end{array}$$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= 1 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\
 &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4
 \end{aligned}$$

Unable to solve the balance equation $(2x^m m(-1 + m) - x^m m + x^m) c_0$ for c_0 and x . No particular solution exists.

Failed to convert RHS $1 + x$ to series in order to find particular solution. Unable to solve. Terminating Unable to find the particular solution or no solution exists.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✗ Solution by Maple

```
Order:=6;
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = 1+x,y(x),type='series',x=0)
```

No solution found

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 224

```
AsymptoticDSolveValue[2*x^2*y'[x] - x*y'[x] + (1-x^2)*y[x] ==1+x,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ + \sqrt{x} \left(-\frac{x^{11/2}}{154440} - \frac{x^{9/2}}{1620} - \frac{x^{7/2}}{1260} - \frac{x^{5/2}}{25} - \frac{x^{3/2}}{15} - 2\sqrt{x} \right. \\ \left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \left(\frac{x^6}{66528} + \frac{x^5}{55440} + \frac{x^4}{672} + \frac{x^3}{504} + \frac{x^2}{12} + \frac{x}{6} - \frac{1}{x} + \dots \right)$$

4.4 problem 4

Internal problem ID [7225]

Internal file name [OUTPUT/6211_Sunday_June_05_2022_04_32_24_PM_91532832/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$2x^2y'' - xy' + (1 - x^2)y = x$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 154: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = x$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r)(n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r(-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r(-1 + r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. In order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$$\begin{array}{l} a_1 = 0 \\ a_2 = \frac{a_0}{2r^2+5r+3} \\ a_3 = 0 \\ a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)} \\ a_5 = 0 \end{array}$$

Unable to solve the balance equation $(2x^m m(-1+m) - x^m m + x^m) c_0$ for c_0 and x . No particular solution exists.

Adding all the above particular solution(s) gives

$$y_p = \text{FAIL}$$

Unable to find the particular solution or no solution exists.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✗ Solution by Maple

```
Order:=6;
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = x,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 166

```
AsymptoticDSolveValue[2*x^2*y'[x] - x*y'[x] + (1-x^2)*y[x] ==x,y[x],{x,0,5}]
```

$$y(x) \rightarrow x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \left(\frac{x^6}{66528} + \frac{x^4}{672} + \frac{x^2}{12} + \log(x) \right) \\ + c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ + \sqrt{x} \left(-\frac{x^{9/2}}{1620} - \frac{x^{5/2}}{25} - 2\sqrt{x} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right)$$

4.5 problem 5

Internal problem ID [7226]

Internal file name [OUTPUT/6212_Sunday_June_05_2022_04_32_26_PM_81686768/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$2x^2y'' - xy' + (1 - x^2)y = x^2 + x + 1$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 155: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = x^2 + x + 1$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r)(n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r(-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r(-1 + r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^2 + x + 1$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. In order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$ $a_2 = \frac{a_0}{2r^2+5r+3}$ $a_3 = 0$ $a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$ $a_5 = 0$
--

Since the $F = x^2 + x + 1$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x^2$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^2$$

For c_0 and x . This results in

$$c_0 = \frac{1}{3}$$

$$m = 2$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+2}$$

Where in the above $c_0 = \frac{1}{3}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{3}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$$\begin{array}{l}
c_0 = \frac{1}{3} \\
c_1 = 0 \\
c_2 = \frac{1}{63} \\
c_3 = 0 \\
c_4 = \frac{1}{3465} \\
c_5 = 0
\end{array}$$

The particular solution is now found using

$$\begin{aligned}
y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
&= x^2 \sum_{n=0}^{\infty} c_n x^n
\end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
y_p &= x^2 \left(\frac{1}{3} + \frac{1}{63} x^2 + \frac{1}{3465} x^4 \right) \\
&= \frac{1}{3} x^2 + \frac{1}{63} x^4 + \frac{1}{3465} x^6
\end{aligned}$$

Unable to solve the balance equation $(2x^m m(-1 + m) - x^m m + x^m) c_0$ for c_0 and x .
No particular solution exists.

Failed to convert RHS $x^2 + x + 1$ to series in order to find particular solution. Unable to solve. Terminating Unable to find the particular solution or no solution exists.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✗ Solution by Maple

```
Order:=6;
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = 1+x+x^2,y(x),type='series',
```

No solution found

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 224

```
AsymptoticDSolveValue[2*x^2*y'[x] - x*y'[x] + (1-x^2)*y[x] == 1+x+x^2,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ + \sqrt{x} \left(-\frac{79x^{11/2}}{154440} - \frac{x^{9/2}}{1620} - \frac{37x^{7/2}}{1260} - \frac{x^{5/2}}{25} - \frac{11x^{3/2}}{15} - 2\sqrt{x} \right. \\ \left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \left(\frac{x^6}{66528} + \frac{67x^5}{55440} + \frac{x^4}{672} + \frac{29x^3}{504} + \frac{x^2}{12} + \frac{7x}{6} - 1 \right)$$

4.6 problem 6

Internal problem ID [7227]

Internal file name [OUTPUT/6213_Sunday_June_05_2022_04_32_27_PM_17721390/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = x^2$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 156: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	"regular"

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	"regular"

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = x^2$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^2$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Now we determine the particular solution y_p associated with $F = x^2$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^2$$

For c_0 and x . This results in

$$c_0 = \frac{1}{3}$$

$$m = 2$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+2}$$

Where in the above $c_0 = \frac{1}{3}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{3}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{3}$
$c_1 = 0$
$c_2 = \frac{1}{63}$
$c_3 = 0$
$c_4 = \frac{1}{3465}$
$c_5 = 0$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^2 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}y_p &= x^2 \left(\frac{1}{3} + \frac{1}{63}x^2 + \frac{1}{3465}x^4 \right) \\ &= \frac{1}{3}x^2 + \frac{1}{63}x^4 + \frac{1}{3465}x^6\end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = \frac{x^2}{3} + \frac{x^4}{63} + \frac{x^6}{3465} + O(x^6)$$

Hence the final solution is

$$y = y_h + y_p$$

$$= \frac{x^2}{3} + \frac{x^4}{63} + \frac{x^6}{3465} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= \frac{x^2}{3} + \frac{x^4}{63} + \frac{x^6}{3465} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ &\quad + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)\end{aligned}\tag{1}$$

Verification of solutions

$$\begin{aligned}y &= \frac{x^2}{3} + \frac{x^4}{63} + \frac{x^6}{3465} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ &\quad + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)\end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
  checking if the LODE has constant coefficients  
  checking if the LODE is of Euler type  
  trying a symmetry of the form [xi=0, eta=F(x)]  
  checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
  -> Bessel  
    <- Bessel successful  
  <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
Order:=6;  
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = x^2,y(x),type='series',x=0)
```

$$y(x) = c_1 \sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2 x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + x^2 \left(\frac{1}{3} + \frac{1}{63}x^2 + O(x^4) \right)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 160

```
AsymptoticDSolveValue[2*x^2*y'[x] - x*y'[x] + (1-x^2)*y[x] ==x^2,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) + c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{1980} - \frac{x^{7/2}}{35} - \frac{2x^{3/2}}{3} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^5}{840} + \frac{x^3}{18} + x \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right)$$

4.7 problem 7

Internal problem ID [7228]

Internal file name [OUTPUT/6214_Sunday_June_05_2022_04_32_30_PM_7253412/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 7.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = x^2 + 1$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 157: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = x^2 + 1$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & 2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^2 + 1$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Since the $F = x^2 + 1$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x^2$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^2$$

For c_0 and x . This results in

$$c_0 = \frac{1}{3}$$

$$m = 2$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+2}$$

Where in the above $c_0 = \frac{1}{3}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{3}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{3}$
$c_1 = 0$
$c_2 = \frac{1}{63}$
$c_3 = 0$
$c_4 = \frac{1}{3465}$
$c_5 = 0$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^2 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x^2 \left(\frac{1}{3} + \frac{1}{63}x^2 + \frac{1}{3465}x^4 \right) \\ &= \frac{1}{3}x^2 + \frac{1}{63}x^4 + \frac{1}{3465}x^6 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= 1 \\ m &= 0 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+0} \end{aligned}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = 1$
$c_1 = 0$
$c_2 = \frac{1}{3}$
$c_3 = 0$
$c_4 = \frac{1}{63}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= 1 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\ &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = 1 + \frac{2x^2}{3} + \frac{2x^4}{63} + \frac{x^6}{3465} + O(x^6)$$

Hence the final solution is

$$\begin{aligned} y &= y_h + y_p \\ &= 1 + \frac{2x^2}{3} + \frac{2x^4}{63} + \frac{x^6}{3465} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ &\quad + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= 1 + \frac{2x^2}{3} + \frac{2x^4}{63} + \frac{x^6}{3465} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ &\quad + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned} y &= 1 + \frac{2x^2}{3} + \frac{2x^4}{63} + \frac{x^6}{3465} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ &\quad + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

```
Order:=6;
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = 1+x^2,y(x),type='series',x=
```

$$y(x) = c_1\sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) \\ + c_2x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + \left(1 + \frac{2}{3}x^2 + \frac{2}{63}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 176

```
AsymptoticDSolveValue[2*x^2*y''[x] - x*y'[x] + (1-x^2)*y[x] ==1+x^2,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ & + c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{79x^{11/2}}{154440} - \frac{37x^{7/2}}{1260} - \frac{11x^{3/2}}{15} \right. \\ & \left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{67x^5}{55440} + \frac{29x^3}{504} + \frac{7x}{6} - \frac{1}{x} \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \end{aligned}$$

4.8 problem 8

Internal problem ID [7229]

Internal file name [OUTPUT/6215_Sunday_June_05_2022_04_32_32_PM_8454786/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = x^4$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 158: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2 - 1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = x^4$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^4$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Now we determine the particular solution y_p associated with $F = x^4$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^4$$

For c_0 and x . This results in

$$c_0 = \frac{1}{21}$$

$$m = 4$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+4}$$

Where in the above $c_0 = \frac{1}{21}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 4$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{21}$ and $r = m$ or $r = 4$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{21}$
$c_1 = 0$
$c_2 = \frac{1}{1155}$
$c_3 = 0$
$c_4 = \frac{1}{121275}$
$c_5 = 0$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^4 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}y_p &= x^4 \left(\frac{1}{21} + \frac{1}{1155}x^2 + \frac{1}{121275}x^4 \right) \\ &= \frac{1}{21}x^4 + \frac{1}{1155}x^6 + \frac{1}{121275}x^8\end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = \frac{x^4}{21} + \frac{x^6}{1155} + \frac{x^8}{121275} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{x^4}{21} + O(x^6)$$

Hence the final solution is

$$\begin{aligned}y &= y_h + y_p \\ &= \frac{x^4}{21} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{x^4}{21} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verification of solutions

$$y = \frac{x^4}{21} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 43

```
Order:=6;
```

```
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = x^4,y(x),type='series',x=0)
```

$$y(x) = c_1 \sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2 x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + x^4 \left(\frac{1}{21} + O(x^2) \right)$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 150

```
AsymptoticDSolveValue[2*x^2*y''[x] - x*y'[x] + (1-x^2)*y[x] ==x^4,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) + c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{55} - \frac{2x^{7/2}}{7} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^5}{30} + \frac{x^3}{3} \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right)$$

4.9 problem 9

Internal problem ID [7230]

Internal file name [OUTPUT/6216_Sunday_June_05_2022_04_32_35_PM_55173922/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 9.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$2x^2y'' - xy' + (1 - x^2)y = \sin(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 159: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = \sin(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r)(n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r(-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r(-1 + r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \sin(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. In order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$$\begin{array}{l} a_1 = 0 \\ a_2 = \frac{a_0}{2r^2+5r+3} \\ a_3 = 0 \\ a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)} \\ a_5 = 0 \end{array}$$

Expanding the rhs of the ode $\sin(x)$ in series gives

$$\sin(x) = x - \frac{1}{6}x^3 + \frac{1}{120}x^5$$

Since the $F = x - \frac{1}{6}x^3 + \frac{1}{120}x^5$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Unable to solve the balance equation $(2x^m m(-1 + m) - x^m m + x^m) c_0$ for c_0 and x . No particular solution exists.

Failed to convert RHS $\sin(x)$ to series in order to find particular solution. Unable to solve. Terminating Unable to find the particular solution or no solution exists.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

Solution by Maple

```
Order:=6;
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = sin(x),y(x),type='series',x
```

No solution found

Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 159

```
AsymptoticDSolveValue[2*x^2*y''[x] - x*y'[x] + (1-x^2)*y[x] ==Sin[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \left(\frac{x^6}{20790} - \frac{17x^4}{5040} + \log(x) \right) \\ + c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ + \sqrt{x} \left(\frac{x^{9/2}}{810} + \frac{2x^{5/2}}{75} - 2\sqrt{x} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right)$$

4.10 problem 10

Internal problem ID [7231]

Internal file name [OUTPUT/6217_Sunday_June_05_2022_04_32_36_PM_90296483/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 10.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$2x^2y'' - xy' + (1 - x^2)y = 1 + \sin(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 160: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = 1 + \sin(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r)(n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r(-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r(-1 + r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1 + \sin(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. In order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$ $a_2 = \frac{a_0}{2r^2+5r+3}$ $a_3 = 0$ $a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$ $a_5 = 0$
--

Expanding the rhs of the ode $1 + \sin(x)$ in series gives

$$1 + \sin(x) = 1 + x - \frac{1}{6}x^3 + \frac{1}{120}x^5$$

Since the $F = 1 + x - \frac{1}{6}x^3 + \frac{1}{120}x^5$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$c_0 = 1$$

$$m = 0$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+0}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and

using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = 1$
$c_1 = 0$
$c_2 = \frac{1}{3}$
$c_3 = 0$
$c_4 = \frac{1}{63}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= 1 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\
 &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4
 \end{aligned}$$

Unable to solve the balance equation $(2x^m m(-1 + m) - x^m m + x^m) c_0$ for c_0 and x . No particular solution exists.

Failed to convert RHS $1 + \sin(x)$ to series in order to find particular solution. Unable to solve. Terminating Unable to find the particular solution or no solution exists.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✗ Solution by Maple

```
Order:=6;
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = 1+sin(x),y(x),type='series'
```

No solution found

✓ Solution by Mathematica

Time used: 0.142 (sec). Leaf size: 217

```
AsymptoticDSolveValue[2*x^2*y'[x] - x*y'[x] + (1-x^2)*y[x] ==1+Sin[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ + \sqrt{x} \left(-\frac{x^{11/2}}{154440} + \frac{x^{9/2}}{810} - \frac{x^{7/2}}{1260} + \frac{2x^{5/2}}{75} - \frac{x^{3/2}}{15} - 2\sqrt{x} \right. \\ \left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \left(\frac{x^6}{20790} + \frac{x^5}{55440} - \frac{17x^4}{5040} + \frac{x^3}{504} + \frac{x}{6} - \frac{1}{x} + \log \right)$$

4.11 problem 11

Internal problem ID [7232]

Internal file name [OUTPUT/6218_Sunday_June_05_2022_04_32_38_PM_63408434/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 11.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = x \sin(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 161: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	"regular"

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	"regular"

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = x \sin(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x \sin(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Expanding the rhs of the ode $x \sin(x)$ in series gives

$$x \sin(x) = x^2 - \frac{1}{6}x^4$$

Since the $F = x^2 - \frac{1}{6}x^4$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x^2$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^2$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= \frac{1}{3} \\ m &= 2 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+2} \end{aligned}$$

Where in the above $c_0 = \frac{1}{3}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{3}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{3}$
$c_1 = 0$
$c_2 = \frac{1}{63}$
$c_3 = 0$
$c_4 = \frac{1}{3465}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= x^2 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x^2 \left(\frac{1}{3} + \frac{1}{63} x^2 + \frac{1}{3465} x^4 \right) \\ &= \frac{1}{3} x^2 + \frac{1}{63} x^4 + \frac{1}{3465} x^6 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{x^4}{6}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = -\frac{x^4}{6}$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= -\frac{1}{126} \\ m &= 4 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+4} \end{aligned}$$

Where in the above $c_0 = -\frac{1}{126}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 4$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{126}$ and $r = m$ or $r = 4$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = -\frac{1}{126}$ $c_1 = 0$ $c_2 = -\frac{1}{6930}$ $c_3 = 0$ $c_4 = -\frac{1}{727650}$ $c_5 = 0$
--

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^4 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$y_p = x^4 \left(-\frac{1}{126} - \frac{1}{6930} x^2 - \frac{1}{727650} x^4 \right)$$

$$= -\frac{1}{126} x^4 - \frac{1}{6930} x^6 - \frac{1}{727650} x^8$$

Adding all the above particular solution(s) gives

$$y_p = \frac{x^2}{3} + \frac{x^4}{126} + \frac{x^6}{6930} - \frac{x^8}{727650} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{x^2}{3} + \frac{x^4}{126} + O(x^6)$$

Hence the final solution is

$$y = y_h + y_p$$

$$= \frac{x^2}{3} + \frac{x^4}{126} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Summary

The solution(s) found are the following

$$y = \frac{x^2}{3} + \frac{x^4}{126} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \quad (1)$$

Verification of solutions

$$y = \frac{x^2}{3} + \frac{x^4}{126} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
  checking if the LODE has constant coefficients
  checking if the LODE is of Euler type
  trying a symmetry of the form [xi=0, eta=F(x)]
  checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  <- Bessel successful
  <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

dsolve(2*x^2*diff(y(x), x\$2) - x*diff(y(x), x) + (1-x^2)*y(x) = x*sin(x),y(x),type='series')

$$y(x) = c_1\sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + x^2 \left(\frac{1}{3} + \frac{1}{126}x^2 + O(x^4) \right)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 167

AsymptoticDSolveValue[2*x^2*y'[x] - x*y'[x] + (1-x^2)*y[x] ==x*sin(x),y[x],{x,0,5}]

$$y(x) \rightarrow c_2x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) + c_1\sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) \left(-\frac{x^{11/2} \sin}{1980} - \frac{1}{35}x^{7/2} \sin - \frac{2}{3}x^{3/2} \sin \right) + x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \left(\frac{x^5 \sin}{840} + \frac{x^3 \sin}{18} + x \sin \right)$$

4.12 problem 12

Internal problem ID [7233]

Internal file name [OUTPUT/6219_Sunday_June_05_2022_04_32_41_PM_63660793/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 12.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$2x^2y'' - xy' + (1 - x^2)y = \cos(x) + \sin(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 162: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = \cos(x) + \sin(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r)(n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r(-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r(-1 + r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \cos(x) + \sin(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$$\begin{array}{l} a_1 = 0 \\ a_2 = \frac{a_0}{2r^2+5r+3} \\ a_3 = 0 \\ a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)} \\ a_5 = 0 \end{array}$$

Expanding the rhs of the ode $\cos(x) + \sin(x)$ in series gives

$$\cos(x) + \sin(x) = 1 + x - \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5$$

Since the $F = 1 + x - \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$\begin{array}{l} c_0 = 1 \\ m = 0 \end{array}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+0} \end{aligned}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = 1$
$c_1 = 0$
$c_2 = \frac{1}{3}$
$c_3 = 0$
$c_4 = \frac{1}{63}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= 1 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\
 &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4
 \end{aligned}$$

Unable to solve the balance equation $(2x^m m(-1 + m) - x^m m + x^m) c_0$ for c_0 and x . No particular solution exists.

Failed to convert RHS $\cos(x) + \sin(x)$ to series in order to find particular solution. Unable to solve. Terminating Unable to find the particular solution or no solution exists.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

Solution by Maple

```
Order:=6;
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (1-x^2)*y(x) = sin(x)+cos(x), y(x), type='se
```

No solution found

Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 217

```
AsymptoticDSolveValue[2*x^2*y'[x] - x*y'[x] + (1-x^2)*y[x] ==Sin[x]+Cos[x], y[x], {x, 0, 5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ + \sqrt{x} \left(-\frac{x^{11/2}}{3861} + \frac{x^{9/2}}{810} + \frac{x^{7/2}}{630} + \frac{2x^{5/2}}{75} + \frac{4x^{3/2}}{15} - 2\sqrt{x} \right. \\ \left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \left(\frac{x^6}{20790} + \frac{37x^5}{69300} - \frac{17x^4}{5040} - \frac{x^3}{84} - \frac{x}{3} - \frac{1}{x} + \log(x) \right)$$

4.13 problem 13

Internal problem ID [7234]

Internal file name [OUTPUT/6220_Sunday_June_05_2022_04_32_45_PM_5690383/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 13.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' + (\cos(x) - 1) y' + e^x y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2 y'' + (\cos(x) - 1) y' + e^x y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{\cos(x) - 1}{x^2}$$
$$q(x) = \frac{e^x}{x^2}$$

Table 163: Table $p(x), q(x)$ singularities.

$p(x) = \frac{\cos(x)-1}{x^2}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{e^x}{x^2}$	
singularity	type
$x = 0$	“regular”
$x = \infty$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + (\cos(x) - 1) y' + e^x y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + (\cos(x) - 1) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + e^x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Expanding $\cos(x) - 1$ as Taylor series around $x = 0$ and keeping only the first 6 terms gives

$$\begin{aligned}\cos(x) - 1 &= -\frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6 + \dots \\ &= -\frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6\end{aligned}$$

Expanding e^x as Taylor series around $x = 0$ and keeping only the first 6 terms gives

$$\begin{aligned}e^x &= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \dots \\ &= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6\end{aligned}$$

Which simplifies to

$$\begin{aligned}&\left(\sum_{n=0}^{\infty} x^{n+r} a_n(n+r)(n+r-1)\right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+5} a_n(n+r)}{720}\right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n(n+r)}{24}\right) + \sum_{n=0}^{\infty} \left(-\frac{x^{1+n+r} a_n(n+r)}{2}\right) \\ &+ \left(\sum_{n=0}^{\infty} a_n x^{n+r}\right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n\right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+2} a_n}{2}\right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n}{6}\right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{24}\right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120}\right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+6} a_n}{720}\right) = 0\end{aligned}\tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=0}^{\infty} \left(-\frac{x^{n+r+5} a_n(n+r)}{720}\right) &= \sum_{n=5}^{\infty} \left(-\frac{a_{n-5}(n-5+r) x^{n+r}}{720}\right) \\ \sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n(n+r)}{24} &= \sum_{n=3}^{\infty} \frac{a_{n-3}(n-3+r) x^{n+r}}{24} \\ \sum_{n=0}^{\infty} \left(-\frac{x^{1+n+r} a_n(n+r)}{2}\right) &= \sum_{n=1}^{\infty} \left(-\frac{a_{n-1}(n+r-1) x^{n+r}}{2}\right)\end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{\infty} x^{1+n+r} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+2} a_n}{2} &= \sum_{n=2}^{\infty} \frac{a_{n-2} x^{n+r}}{2} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n}{6} &= \sum_{n=3}^{\infty} \frac{a_{n-3} x^{n+r}}{6} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{24} &= \sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r}}{24} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} &= \sum_{n=5}^{\infty} \frac{a_{n-5} x^{n+r}}{120} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+6} a_n}{720} &= \sum_{n=6}^{\infty} \frac{a_{n-6} x^{n+r}}{720}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\begin{aligned}
&\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=5}^{\infty} \left(-\frac{a_{n-5} (n-5+r) x^{n+r}}{720} \right) \\
&+ \left(\sum_{n=3}^{\infty} \frac{a_{n-3} (n-3+r) x^{n+r}}{24} \right) + \sum_{n=1}^{\infty} \left(-\frac{a_{n-1} (n+r-1) x^{n+r}}{2} \right) \\
&+ \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) + \left(\sum_{n=2}^{\infty} \frac{a_{n-2} x^{n+r}}{2} \right) \\
&+ \left(\sum_{n=3}^{\infty} \frac{a_{n-3} x^{n+r}}{6} \right) + \left(\sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r}}{24} \right) \\
&+ \left(\sum_{n=5}^{\infty} \frac{a_{n-5} x^{n+r}}{120} \right) + \left(\sum_{n=6}^{\infty} \frac{a_{n-6} x^{n+r}}{720} \right) = 0
\end{aligned} \tag{2B}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r (-1+r) + a_0 x^r = 0$$

Or

$$(x^r r(-1+r) + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2} + \frac{i\sqrt{3}}{2}$$
$$r_2 = \frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$
$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}+\frac{i\sqrt{3}}{2}}$$
$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}-\frac{i\sqrt{3}}{2}}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{r-2}{2r^2+2r+2}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = -\frac{r(r+5)}{4(r^2+r+1)(r^2+3r+3)}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = \frac{-r^5 - 8r^4 - 32r^3 - 55r^2 - 9r + 24}{24(r^2+r+1)(r^2+3r+3)(r^2+5r+7)}$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{-4r^6 - 38r^5 - 141r^4 - 196r^3 + 85r^2 + 408r + 192}{48(r^2+r+1)(r^2+3r+3)(r^2+5r+7)(r^2+7r+13)}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = \frac{2r^9 + 20r^8 - 17r^7 - 757r^6 - 1964r^5 + 7667r^4 + 50216r^3 + 98979r^2 + 73140r + 9504}{1440(r^2+r+1)(r^2+3r+3)(r^2+5r+7)(r^2+7r+13)(r^2+9r+21)}$$

For $6 \leq n$ the recursive equation is

$$\begin{aligned} a_n(n+r)(n+r-1) - \frac{a_{n-5}(n-5+r)}{720} + \frac{a_{n-3}(n-3+r)}{24} \\ - \frac{a_{n-1}(n+r-1)}{2} + a_n + a_{n-1} + \frac{a_{n-2}}{2} + \frac{a_{n-3}}{6} + \frac{a_{n-4}}{24} + \frac{a_{n-5}}{120} + \frac{a_{n-6}}{720} = 0 \end{aligned} \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{na_{n-5} - 30na_{n-3} + 360na_{n-1} + ra_{n-5} - 30ra_{n-3} + 360ra_{n-1} - a_{n-6} - 11a_{n-5} - 30a_{n-4} - 30a_{n-3} - \dots}{720n^2 + 1440nr + 720r^2 - 720n - 720r + 720} \quad (4)$$

Which for the root $r = \frac{1}{2} + \frac{i\sqrt{3}}{2}$ becomes

$$a_n = \frac{i(a_{n-5} - 30a_{n-3} + 360a_{n-1})\sqrt{3} + 2(a_{n-5} - 30a_{n-3} + 360a_{n-1})n - 2a_{n-6} - 21a_{n-5} - 60a_{n-4} - 90a_{n-3} - \dots}{1440n(i\sqrt{3} + n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2} + \frac{i\sqrt{3}}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r-2}{2r^2+2r+2}$	$\frac{i\sqrt{3}}{4}$
a_2	$-\frac{r(r+5)}{4(r^2+r+1)(r^2+3r+3)}$	$\frac{-i\sqrt{3}-11}{32i\sqrt{3}+64}$
a_3	$\frac{-r^5-8r^4-32r^3-55r^2-9r+24}{24(r^2+r+1)(r^2+3r+3)(r^2+5r+7)}$	$\frac{\frac{55\sqrt{3}}{288} + \frac{55i}{96}}{(i-\sqrt{3})(i\sqrt{3}+2)(i\sqrt{3}+3)}$
a_4	$\frac{-4r^6-38r^5-141r^4-196r^3+85r^2+408r+192}{48(r^2+r+1)(r^2+3r+3)(r^2+5r+7)(r^2+7r+13)}$	$\frac{112i\sqrt{3}+199}{384(-\sqrt{3}+2i)(-i+\sqrt{3})(i\sqrt{3}+3)(i\sqrt{3}+4)}$
a_5	$\frac{2r^9+20r^8-17r^7-757r^6-1964r^5+7667r^4+50216r^3+98979r^2+73140r+9504}{1440(r^2+r+1)(r^2+3r+3)(r^2+5r+7)(r^2+7r+13)(r^2+9r+21)}$	$\frac{\frac{18491\sqrt{3}}{38400} + \frac{4387i}{12800}}{(-i+\sqrt{3})(i\sqrt{3}+2)(i\sqrt{3}+3)(i\sqrt{3}+4)(i\sqrt{3}+5)}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
y_1(x) &= x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
&= x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(1 + \frac{i\sqrt{3}x}{4} + \frac{(-i\sqrt{3}-11)x^2}{32i\sqrt{3}+64} + \frac{55(\sqrt{3}+3i)x^3}{288(i-\sqrt{3})(i\sqrt{3}+2)(i\sqrt{3}+3)} \right. \\
&\quad \left. + \frac{(112i\sqrt{3}+199)x^4}{384(-\sqrt{3}+2i)(-i+\sqrt{3})(i\sqrt{3}+3)(i\sqrt{3}+4)} \right. \\
&\quad \left. + \frac{41(451\sqrt{3}+321i)x^5}{38400(-i+\sqrt{3})(i\sqrt{3}+2)(i\sqrt{3}+3)(i\sqrt{3}+4)(i\sqrt{3}+5)} + O(x^6) \right)
\end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned}
y_2(x) &= x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(1 - \frac{i\sqrt{3}x}{4} + \frac{(i\sqrt{3}-11)x^2}{-32i\sqrt{3}+64} + \frac{55(\sqrt{3}-3i)x^3}{288(-i-\sqrt{3})(-i\sqrt{3}+2)(-i\sqrt{3}+3)} \right. \\
&\quad \left. + \frac{(-112i\sqrt{3}+199)x^4}{384(-\sqrt{3}-2i)(\sqrt{3}+i)(-i\sqrt{3}+3)(-i\sqrt{3}+4)} \right. \\
&\quad \left. + \frac{41(451\sqrt{3}-321i)x^5}{38400(\sqrt{3}+i)(-i\sqrt{3}+2)(-i\sqrt{3}+3)(-i\sqrt{3}+4)(-i\sqrt{3}+5)} + O(x^6) \right)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
&= c_1 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(1 + \frac{i\sqrt{3}x}{4} + \frac{(-i\sqrt{3} - 11)x^2}{32i\sqrt{3} + 64} + \frac{55(\sqrt{3} + 3i)x^3}{288(i - \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)} \right. \\
&\quad \left. + \frac{(112i\sqrt{3} + 199)x^4}{384(-\sqrt{3} + 2i)(-i + \sqrt{3})(i\sqrt{3} + 3)(i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{41(451\sqrt{3} + 321i)x^5}{38400(-i + \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)(i\sqrt{3} + 4)(i\sqrt{3} + 5)} + O(x^6) \right) \\
&\quad + c_2 x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(1 - \frac{i\sqrt{3}x}{4} + \frac{(i\sqrt{3} - 11)x^2}{-32i\sqrt{3} + 64} \right. \\
&\quad \left. + \frac{55(\sqrt{3} - 3i)x^3}{288(-i - \sqrt{3})(-i\sqrt{3} + 2)(-i\sqrt{3} + 3)} \right. \\
&\quad \left. + \frac{(-112i\sqrt{3} + 199)x^4}{384(-\sqrt{3} - 2i)(\sqrt{3} + i)(-i\sqrt{3} + 3)(-i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{41(451\sqrt{3} - 321i)x^5}{38400(\sqrt{3} + i)(-i\sqrt{3} + 2)(-i\sqrt{3} + 3)(-i\sqrt{3} + 4)(-i\sqrt{3} + 5)} + O(x^6) \right)
\end{aligned}$$

Hence the final solution is

$$\begin{aligned}
y &= y_h \\
&= c_1 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(1 + \frac{i\sqrt{3}x}{4} + \frac{(-i\sqrt{3} - 11)x^2}{32i\sqrt{3} + 64} + \frac{55(\sqrt{3} + 3i)x^3}{288(i - \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)} \right. \\
&\quad \left. + \frac{(112i\sqrt{3} + 199)x^4}{384(-\sqrt{3} + 2i)(-i + \sqrt{3})(i\sqrt{3} + 3)(i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{41(451\sqrt{3} + 321i)x^5}{38400(-i + \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)(i\sqrt{3} + 4)(i\sqrt{3} + 5)} + O(x^6) \right) \\
&\quad + c_2 x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(1 - \frac{i\sqrt{3}x}{4} + \frac{(i\sqrt{3} - 11)x^2}{-32i\sqrt{3} + 64} + \frac{55(\sqrt{3} - 3i)x^3}{288(-i - \sqrt{3})(-i\sqrt{3} + 2)(-i\sqrt{3} + 3)} \right. \\
&\quad \left. + \frac{(-112i\sqrt{3} + 199)x^4}{384(-\sqrt{3} - 2i)(\sqrt{3} + i)(-i\sqrt{3} + 3)(-i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{41(451\sqrt{3} - 321i)x^5}{38400(\sqrt{3} + i)(-i\sqrt{3} + 2)(-i\sqrt{3} + 3)(-i\sqrt{3} + 4)(-i\sqrt{3} + 5)} + O(x^6) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = c_1 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} & \left(1 + \frac{i\sqrt{3}x}{4} + \frac{(-i\sqrt{3} - 11)x^2}{32i\sqrt{3} + 64} + \frac{55(\sqrt{3} + 3i)x^3}{288(i - \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)} \right. \\ & + \frac{(112i\sqrt{3} + 199)x^4}{384(-\sqrt{3} + 2i)(-i + \sqrt{3})(i\sqrt{3} + 3)(i\sqrt{3} + 4)} \\ & \left. + \frac{41(451\sqrt{3} + 321i)x^5}{38400(-i + \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)(i\sqrt{3} + 4)(i\sqrt{3} + 5)} + O(x^6) \right) \\ + c_2 x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} & \left(1 - \frac{i\sqrt{3}x}{4} + \frac{(i\sqrt{3} - 11)x^2}{-32i\sqrt{3} + 64} \right. \\ & + \frac{55(\sqrt{3} - 3i)x^3}{288(-i - \sqrt{3})(-i\sqrt{3} + 2)(-i\sqrt{3} + 3)} \\ & + \frac{(-112i\sqrt{3} + 199)x^4}{384(-\sqrt{3} - 2i)(\sqrt{3} + i)(-i\sqrt{3} + 3)(-i\sqrt{3} + 4)} \\ & \left. + \frac{41(451\sqrt{3} - 321i)x^5}{38400(\sqrt{3} + i)(-i\sqrt{3} + 2)(-i\sqrt{3} + 3)(-i\sqrt{3} + 4)(-i\sqrt{3} + 5)} + O(x^6) \right) \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y = c_1 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} & \left(1 + \frac{i\sqrt{3}x}{4} + \frac{(-i\sqrt{3} - 11)x^2}{32i\sqrt{3} + 64} + \frac{55(\sqrt{3} + 3i)x^3}{288(i - \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)} \right. \\ & + \frac{(112i\sqrt{3} + 199)x^4}{384(-\sqrt{3} + 2i)(-i + \sqrt{3})(i\sqrt{3} + 3)(i\sqrt{3} + 4)} \\ & \left. + \frac{41(451\sqrt{3} + 321i)x^5}{38400(-i + \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)(i\sqrt{3} + 4)(i\sqrt{3} + 5)} + O(x^6) \right) \\ + c_2 x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} & \left(1 - \frac{i\sqrt{3}x}{4} + \frac{(i\sqrt{3} - 11)x^2}{-32i\sqrt{3} + 64} + \frac{55(\sqrt{3} - 3i)x^3}{288(-i - \sqrt{3})(-i\sqrt{3} + 2)(-i\sqrt{3} + 3)} \right. \\ & + \frac{(-112i\sqrt{3} + 199)x^4}{384(-\sqrt{3} - 2i)(\sqrt{3} + i)(-i\sqrt{3} + 3)(-i\sqrt{3} + 4)} \\ & \left. + \frac{41(451\sqrt{3} - 321i)x^5}{38400(\sqrt{3} + i)(-i\sqrt{3} + 2)(-i\sqrt{3} + 3)(-i\sqrt{3} + 4)(-i\sqrt{3} + 5)} + O(x^6) \right) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 5`[0, u]
```

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 1171

Order:=6;

dsolve(x^2*diff(y(x), x\$2) + (cos(x)-1)*diff(y(x), x) + exp(x)*y(x) = 0,y(x),type='series',x

$$\begin{aligned}
 y(x) = & \sqrt{x} \left(c_2 x^{\frac{i\sqrt{3}}{2}} \left(1 + \frac{1}{4}i\sqrt{3}x + \frac{-i\sqrt{3} - 11}{32i\sqrt{3} + 64}x^2 + \frac{\frac{55\sqrt{3}}{288} + \frac{55i}{96}}{(i - \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)}x^3 \right. \right. \\
 & + \frac{1}{384} \frac{112i\sqrt{3} + 199}{(-\sqrt{3} + 2i)(-i + \sqrt{3})(i\sqrt{3} + 3)(i\sqrt{3} + 4)}x^4 \\
 & \left. \left. + \frac{\frac{18491\sqrt{3}}{38400} + \frac{4387i}{12800}}{(-i + \sqrt{3})(i\sqrt{3} + 2)(i\sqrt{3} + 3)(i\sqrt{3} + 4)(i\sqrt{3} + 5)}x^5 + O(x^6) \right) \right. \\
 & + c_1 x^{-\frac{i\sqrt{3}}{2}} \left(1 - \frac{1}{4}i\sqrt{3}x + \frac{-\sqrt{3} - 11i}{32\sqrt{3} + 64i}x^2 + \frac{55\sqrt{3} - 165i}{3456i - 2304\sqrt{3}}x^3 \right. \\
 & + \frac{199i + 112\sqrt{3}}{-27648i + 7680\sqrt{3}}x^4 \\
 & \left. \left. + \frac{\frac{18491\sqrt{3}}{38400} - \frac{4387i}{12800}}{(\sqrt{3} + i)(\sqrt{3} + 2i)(\sqrt{3} + 3i)(\sqrt{3} + 4i)(\sqrt{3} + 5i)}x^5 + O(x^6) \right) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 2502

AsymptoticDSolveValue[x^2*y''[x] + (Cos[x]-1)*y'[x] + Exp[x]*y[x] ==0,y[x],{x,0,5}]

Too large to display

4.14 problem 14

4.14.1 Maple step by step solution 1643

Internal problem ID [7235]

Internal file name [OUTPUT/6221_Sunday_June_05_2022_04_32_52_PM_9038864/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 14.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x - 2)y'' + \frac{y'}{x} + (1 + x)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x - 2)y'' + \frac{y'}{x} + (1 + x)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x(x - 2)}$$
$$q(x) = \frac{1 + x}{x - 2}$$

Table 164: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x(x-2)}$	
singularity	type
$x = 0$	“regular”
$x = 2$	“regular”

$q(x) = \frac{1+x}{x-2}$	
singularity	type
$x = 2$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, 2]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$y''x(x-2) + y' + (1+x)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x(x-2) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1+x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-2x^{n+r-1} a_n (n+r) (n+r-1)) \quad (2A) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0 \end{aligned}$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \\ \sum_{n=0}^{\infty} x^{n+r+2} a_n &= \sum_{n=3}^{\infty} a_{n-3} x^{n+r-1} \\ \sum_{n=0}^{\infty} x^{1+n+r} a_n &= \sum_{n=2}^{\infty} a_{n-2} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-2x^{n+r-1} a_n (n+r) (n+r-1)) \quad (2B) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=3}^{\infty} a_{n-3} x^{n+r-1} \right) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r-1} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$-2x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$-2x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(-2x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(-2r^2 + 3r)x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation becomes

$$-2r^2 + 3r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= \frac{3}{2} \\ r_2 &= 0 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(-2r^2 + 3r)x^{-1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{3}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+\frac{3}{2}} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^n \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{r(-1+r)}{2r^2+r-1}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{r^3 - r^2 + 2r - 1}{4r^3 + 8r^2 - r - 2}$$

For $3 \leq n$ the recursive equation is

$$a_{n-1}(n+r-1)(n+r-2) - 2a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-3} + a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{n^2 a_{n-1} + 2nr a_{n-1} + r^2 a_{n-1} - 3n a_{n-1} - 3r a_{n-1} + a_{n-3} + a_{n-2} + 2a_{n-1}}{2n^2 + 4nr + 2r^2 - 3n - 3r} \quad (4)$$

Which for the root $r = \frac{3}{2}$ becomes

$$a_n = \frac{4n^2 a_{n-1} + 4a_{n-3} + 4a_{n-2} - a_{n-1}}{8n^2 + 12n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{3}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r(-1+r)}{2r^2+r-1}$	$\frac{3}{20}$
a_2	$\frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$	$\frac{25}{224}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{r^5 + r^4 + 7r^3 + 5r^2 - 2r - 2}{8r^5 + 44r^4 + 70r^3 + 25r^2 - 18r - 9}$$

Which for the root $r = \frac{3}{2}$ becomes

$$a_3 = \frac{1361}{17280}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r(-1+r)}{2r^2+r-1}$	$\frac{3}{20}$
a_2	$\frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$	$\frac{25}{224}$
a_3	$\frac{r^5+r^4+7r^3+5r^2-2r-2}{8r^5+44r^4+70r^3+25r^2-18r-9}$	$\frac{1361}{17280}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r^7 + 5r^6 + 21r^5 + 52r^4 + 51r^3 + 2r^2 - 21r - 11}{(4r^3 + 8r^2 - r - 2)(1+r)(2r+3)(2r^2 + 13r + 20)}$$

Which for the root $r = \frac{3}{2}$ becomes

$$a_4 = \frac{80753}{2365440}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r(-1+r)}{2r^2+r-1}$	$\frac{3}{20}$
a_2	$\frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$	$\frac{25}{224}$
a_3	$\frac{r^5+r^4+7r^3+5r^2-2r-2}{8r^5+44r^4+70r^3+25r^2-18r-9}$	$\frac{1361}{17280}$
a_4	$\frac{r^7+5r^6+21r^5+52r^4+51r^3+2r^2-21r-11}{(4r^3+8r^2-r-2)(1+r)(2r+3)(2r^2+13r+20)}$	$\frac{80753}{2365440}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{r^9 + 11r^8 + 66r^7 + 262r^6 + 652r^5 + 936r^4 + 648r^3 - 11r^2 - 311r - 164}{(8r^5 + 44r^4 + 70r^3 + 25r^2 - 18r - 9)(2+r)(2r+5)(2r^2 + 17r + 35)}$$

Which for the root $r = \frac{3}{2}$ becomes

$$a_5 = \frac{616517}{38707200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r(-1+r)}{2r^2+r-1}$	$\frac{3}{20}$
a_2	$\frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$	$\frac{25}{224}$
a_3	$\frac{r^5+r^4+7r^3+5r^2-2r-2}{8r^5+44r^4+70r^3+25r^2-18r-9}$	$\frac{1361}{17280}$
a_4	$\frac{r^7+5r^6+21r^5+52r^4+51r^3+2r^2-21r-11}{(4r^3+8r^2-r-2)(1+r)(2r+3)(2r^2+13r+20)}$	$\frac{80753}{2365440}$
a_5	$\frac{r^9+11r^8+66r^7+262r^6+652r^5+936r^4+648r^3-11r^2-311r-164}{(8r^5+44r^4+70r^3+25r^2-18r-9)(2+r)(2r+5)(2r^2+17r+35)}$	$\frac{616517}{38707200}$

Using the above table, then the solution $y_1(x)$ is

$$y_1(x) = x^{\frac{3}{2}}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots)$$

$$= x^{\frac{3}{2}}\left(1 + \frac{3x}{20} + \frac{25x^2}{224} + \frac{1361x^3}{17280} + \frac{80753x^4}{2365440} + \frac{616517x^5}{38707200} + O(x^6)\right)$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = \frac{r(-1+r)}{2r^2+r-1}$$

Substituting $n = 2$ in Eq. (2B) gives

$$b_2 = \frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$$

For $3 \leq n$ the recursive equation is

$$b_{n-1}(n+r-1)(n+r-2) - 2b_n(n+r)(n+r-1) + (n+r)b_n + b_{n-3} + b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{n^2b_{n-1} + 2nr b_{n-1} + r^2b_{n-1} - 3nb_{n-1} - 3rb_{n-1} + b_{n-3} + b_{n-2} + 2b_{n-1}}{2n^2 + 4nr + 2r^2 - 3n - 3r} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = \frac{(n^2 - 3n + 2)b_{n-1} + b_{n-3} + b_{n-2}}{2n^2 - 3n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r(-1+r)}{2r^2+r-1}$	0
b_2	$\frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{r^5 + r^4 + 7r^3 + 5r^2 - 2r - 2}{8r^5 + 44r^4 + 70r^3 + 25r^2 - 18r - 9}$$

Which for the root $r = 0$ becomes

$$b_3 = \frac{2}{9}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r(-1+r)}{2r^2+r-1}$	0
b_2	$\frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$	$\frac{1}{2}$
b_3	$\frac{r^5+r^4+7r^3+5r^2-2r-2}{8r^5+44r^4+70r^3+25r^2-18r-9}$	$\frac{2}{9}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{r^7 + 5r^6 + 21r^5 + 52r^4 + 51r^3 + 2r^2 - 21r - 11}{(4r^3 + 8r^2 - r - 2)(1+r)(2r+3)(2r^2+13r+20)}$$

Which for the root $r = 0$ becomes

$$b_4 = \frac{11}{120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r(-1+r)}{2r^2+r-1}$	0
b_2	$\frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$	$\frac{1}{2}$
b_3	$\frac{r^5+r^4+7r^3+5r^2-2r-2}{8r^5+44r^4+70r^3+25r^2-18r-9}$	$\frac{2}{9}$
b_4	$\frac{r^7+5r^6+21r^5+52r^4+51r^3+2r^2-21r-11}{(4r^3+8r^2-r-2)(1+r)(2r+3)(2r^2+13r+20)}$	$\frac{11}{120}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{r^9 + 11r^8 + 66r^7 + 262r^6 + 652r^5 + 936r^4 + 648r^3 - 11r^2 - 311r - 164}{(8r^5 + 44r^4 + 70r^3 + 25r^2 - 18r - 9)(2+r)(2r+5)(2r^2+17r+35)}$$

Which for the root $r = 0$ becomes

$$b_5 = \frac{82}{1575}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r(-1+r)}{2r^2+r-1}$	0
b_2	$\frac{r^3-r^2+2r-1}{4r^3+8r^2-r-2}$	$\frac{1}{2}$
b_3	$\frac{r^5+r^4+7r^3+5r^2-2r-2}{8r^5+44r^4+70r^3+25r^2-18r-9}$	$\frac{2}{9}$
b_4	$\frac{r^7+5r^6+21r^5+52r^4+51r^3+2r^2-21r-11}{(4r^3+8r^2-r-2)(1+r)(2r+3)(2r^2+13r+20)}$	$\frac{11}{120}$
b_5	$\frac{r^9+11r^8+66r^7+262r^6+652r^5+936r^4+648r^3-11r^2-311r-164}{(8r^5+44r^4+70r^3+25r^2-18r-9)(2+r)(2r+5)(2r^2+17r+35)}$	$\frac{82}{1575}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= 1 + \frac{x^2}{2} + \frac{2x^3}{9} + \frac{11x^4}{120} + \frac{82x^5}{1575} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^{\frac{3}{2}} \left(1 + \frac{3x}{20} + \frac{25x^2}{224} + \frac{1361x^3}{17280} + \frac{80753x^4}{2365440} + \frac{616517x^5}{38707200} + O(x^6) \right) \\ &\quad + c_2 \left(1 + \frac{x^2}{2} + \frac{2x^3}{9} + \frac{11x^4}{120} + \frac{82x^5}{1575} + O(x^6) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x^{\frac{3}{2}} \left(1 + \frac{3x}{20} + \frac{25x^2}{224} + \frac{1361x^3}{17280} + \frac{80753x^4}{2365440} + \frac{616517x^5}{38707200} + O(x^6) \right) \\ &\quad + c_2 \left(1 + \frac{x^2}{2} + \frac{2x^3}{9} + \frac{11x^4}{120} + \frac{82x^5}{1575} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1x^{\frac{3}{2}} \left(1 + \frac{3x}{20} + \frac{25x^2}{224} + \frac{1361x^3}{17280} + \frac{80753x^4}{2365440} + \frac{616517x^5}{38707200} + O(x^6) \right) \\ &\quad + c_2 \left(1 + \frac{x^2}{2} + \frac{2x^3}{9} + \frac{11x^4}{120} + \frac{82x^5}{1575} + O(x^6) \right) \end{aligned} \tag{1}$$

Verification of solutions

$$y = c_1 x^{\frac{3}{2}} \left(1 + \frac{3x}{20} + \frac{25x^2}{224} + \frac{1361x^3}{17280} + \frac{80753x^4}{2365440} + \frac{616517x^5}{38707200} + O(x^6) \right) \\ + c_2 \left(1 + \frac{x^2}{2} + \frac{2x^3}{9} + \frac{11x^4}{120} + \frac{82x^5}{1575} + O(x^6) \right)$$

Verified OK.

4.14.1 Maple step by step solution

Let's solve

$$y''x(x-2) + y' + (1+x)yx = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x(x-2)} - \frac{(1+x)y}{x-2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x(x-2)} + \frac{(1+x)y}{x-2} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x(x-2)}, P_3(x) = \frac{1+x}{x-2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -\frac{1}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x(x-2) + y' + (1+x)yx = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 1..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k- > k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) x^{k+r}$$

- Convert $x^m \cdot y''$ to series expansion for $m = 1..2$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r) (k+r-1) x^{k+r-2+m}$$

- Shift index using $k- > k + 2 - m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r) (k+1-m+r) x^{k+r}$$

Rewrite ODE with series expansions

$$-a_0 r(-3+2r) x^{-1+r} + (-a_1(1+r)(-1+2r) + a_0 r(-1+r)) x^r + (-a_2(2+r)(1+2r) + a_1(1+r)r + a_0) x^{1+r} + \dots = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(-3+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{0, \frac{3}{2}\right\}$$

- The coefficients of each power of x must be 0

$$[-a_1(1+r)(-1+2r) + a_0 r(-1+r) = 0, -a_2(2+r)(1+2r) + a_1(1+r)r + a_0 = 0]$$

- Solve for the dependent coefficient(s)

$$\left\{ a_1 = \frac{a_0 r(-1+r)}{2r^2+r-1}, a_2 = \frac{a_0(r^3-r^2+2r-1)}{4r^3+8r^2-r-2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$-2\left(k - \frac{1}{2} + r\right)(k+1+r)a_{k+1} + a_k(k+r)(k+r-1) + a_{k-1} + a_{k-2} = 0$$

- Shift index using $k \rightarrow k+2$

$$-2\left(k + \frac{3}{2} + r\right)(k+3+r)a_{k+3} + a_{k+2}(k+2+r)(k+1+r) + a_{k+1} + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+3} = \frac{k^2 a_{k+2} + 2kr a_{k+2} + r^2 a_{k+2} + 3ka_{k+2} + 3ra_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{(2k+3+2r)(k+3+r)}$$

- Recursion relation for $r = 0$

$$a_{k+3} = \frac{k^2 a_{k+2} + 3ka_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{(2k+3)(k+3)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = \frac{k^2 a_{k+2} + 3ka_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{(2k+3)(k+3)}, a_1 = 0, a_2 = \frac{a_0}{2} \right]$$

- Recursion relation for $r = \frac{3}{2}$

$$a_{k+3} = \frac{k^2 a_{k+2} + 6ka_{k+2} + a_k + a_{k+1} + \frac{35}{4} a_{k+2}}{(2k+6)(k+\frac{9}{2})}$$

- Solution for $r = \frac{3}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{3}{2}}, a_{k+3} = \frac{k^2 a_{k+2} + 6ka_{k+2} + a_k + a_{k+1} + \frac{35}{4} a_{k+2}}{(2k+6)(k+\frac{9}{2})}, a_1 = \frac{3a_0}{20}, a_2 = \frac{25a_0}{224} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{3}{2}} \right), a_{k+3} = \frac{k^2 a_{k+2} + 3ka_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{(2k+3)(k+3)}, a_1 = 0, a_2 = \frac{a_0}{2}, b_{k+3} = \frac{k^2}{20} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunC ODE, case a <> 0, e <> 0, c =
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
Order:=6;
dsolve((x-2)*diff(y(x), x$2) + 1/x*diff(y(x), x) + (x+1)*y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{3}{2}} \left(1 + \frac{3}{20}x + \frac{25}{224}x^2 + \frac{1361}{17280}x^3 + \frac{80753}{2365440}x^4 + \frac{616517}{38707200}x^5 + O(x^6) \right) \\ + c_2 \left(1 + \frac{1}{2}x^2 + \frac{2}{9}x^3 + \frac{11}{120}x^4 + \frac{82}{1575}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 80

```
AsymptoticDSolveValue[(x-2)*y''[x] + 1/x*y'[x] + (x+1)*y[x] ==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{82x^5}{1575} + \frac{11x^4}{120} + \frac{2x^3}{9} + \frac{x^2}{2} + 1 \right) + c_1 \left(\frac{616517x^5}{38707200} + \frac{80753x^4}{2365440} + \frac{1361x^3}{17280} + \frac{25x^2}{224} + \frac{3x}{20} + 1 \right) x^{3/2}$$

4.15 problem 15

4.15.1 Maple step by step solution 1658

Internal problem ID [7236]

Internal file name [OUTPUT/6222_Sunday_June_05_2022_04_32_56_PM_85136051/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 15.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x - 2)y'' + \frac{y'}{x} + (1 + x)y = 0$$

With the expansion point for the power series method at $x = 2$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = x - 2$$

The ode is converted to be in terms of the new independent variable t . This results in

$$t \left(\frac{d^2}{dt^2} y(t) \right) + \frac{d}{dt} y(t) + (3 + t) y(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$t \left(\frac{d^2}{dt^2} y(t) \right) + \frac{d}{dt} y(t) + (3 + t) y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2}y(t) + p(t)\frac{d}{dt}y(t) + q(t)y(t) = 0$$

Where

$$p(t) = \frac{1}{t(t+2)}$$

$$q(t) = \frac{3+t}{t}$$

Table 166: Table $p(t), q(t)$ singularities.

$p(t) = \frac{1}{t(t+2)}$	
singularity	type
$t = -2$	“regular”
$t = 0$	“regular”

$q(t) = \frac{3+t}{t}$	
singularity	type
$t = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-2, 0]$

Irregular singular points : $[\infty]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$\left(\frac{d^2}{dt^2}y(t)\right)t(t+2) + \frac{d}{dt}y(t) + (3+t)y(t)(t+2) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+r)(n+r-1)a_n t^{n+r-2} \right) t(t+2) \\ & + \left(\sum_{n=0}^{\infty} (n+r)a_n t^{n+r-1} \right) + (3+t) \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) (t+2) = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} t^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2t^{n+r-1} a_n (n+r)(n+r-1) \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r)a_n t^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} t^{n+r+2} a_n \right) \\ & + \left(\sum_{n=0}^{\infty} 5t^{1+n+r} a_n \right) + \left(\sum_{n=0}^{\infty} 6a_n t^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of t be $n+r-1$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} t^{n+r} a_n (n+r)(n+r-1) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1)(n+r-2) t^{n+r-1} \\ \sum_{n=0}^{\infty} t^{n+r+2} a_n &= \sum_{n=3}^{\infty} a_{n-3} t^{n+r-1} \\ \sum_{n=0}^{\infty} 5t^{1+n+r} a_n &= \sum_{n=2}^{\infty} 5a_{n-2} t^{n+r-1} \\ \sum_{n=0}^{\infty} 6a_n t^{n+r} &= \sum_{n=1}^{\infty} 6a_{n-1} t^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers

of t are the same and equal to $n + r - 1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) t^{n+r-1} \right) \\ & + \left(\sum_{n=0}^{\infty} 2t^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) \\ & + \left(\sum_{n=3}^{\infty} a_{n-3} t^{n+r-1} \right) + \left(\sum_{n=2}^{\infty} 5a_{n-2} t^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} 6a_{n-1} t^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2t^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n t^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$2t^{-1+r} a_0 r (-1+r) + r a_0 t^{-1+r} = 0$$

Or

$$(2t^{-1+r} r (-1+r) + r t^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r t^{-1+r} (2r - 1) = 0$$

Since the above is true for all t then the indicial equation becomes

$$2r^2 - r = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r t^{-1+r} (2r - 1) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(t) = t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right)$$

$$y_2(t) = t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right)$$

Or

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^{n+\frac{1}{2}}$$

$$y_2(t) = \sum_{n=0}^{\infty} b_n t^n$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{-r^2 + r - 6}{2r^2 + 3r + 1}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{r^4 + r^2 - 15r + 31}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$$

For $3 \leq n$ the recursive equation is

$$a_{n-1}(n+r-1)(n+r-2) + 2a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-3} + 5a_{n-2} + 6a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{n^2 a_{n-1} + 2nra_{n-1} + r^2 a_{n-1} - 3na_{n-1} - 3ra_{n-1} + a_{n-3} + 5a_{n-2} + 8a_{n-1}}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = \frac{-4n^2 a_{n-1} + 8na_{n-1} - 4a_{n-3} - 20a_{n-2} - 27a_{n-1}}{8n^2 + 4n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2+r-6}{2r^2+3r+1}$	$-\frac{23}{12}$
a_2	$\frac{r^4+r^2-15r+31}{4r^4+20r^3+35r^2+25r+6}$	$\frac{127}{160}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-r^6 - 3r^5 - 3r^4 + 17r^3 + 26r^2 + 182r - 74}{(2r^2 + 11r + 15)(2r^2 + 3r + 1)(2r^2 + 7r + 6)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = \frac{1621}{40320}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2+r-6}{2r^2+3r+1}$	$-\frac{23}{12}$
a_2	$\frac{r^4+r^2-15r+31}{4r^4+20r^3+35r^2+25r+6}$	$\frac{127}{160}$
a_3	$\frac{-r^6-3r^5-3r^4+17r^3+26r^2+182r-74}{(2r^2+11r+15)(2r^2+3r+1)(2r^2+7r+6)}$	$\frac{1621}{40320}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r^8 + 8r^7 + 24r^6 + 11r^5 - 53r^4 - 153r^3 - 75r^2 - 1458r - 897}{(2r^2 + 15r + 28)(4r^4 + 20r^3 + 35r^2 + 25r + 6)(2r^2 + 11r + 15)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = -\frac{426599}{5806080}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2+r-6}{2r^2+3r+1}$	$-\frac{23}{12}$
a_2	$\frac{r^4+r^2-15r+31}{4r^4+20r^3+35r^2+25r+6}$	$\frac{127}{160}$
a_3	$\frac{-r^6-3r^5-3r^4+17r^3+26r^2+182r-74}{(2r^2+11r+15)(2r^2+3r+1)(2r^2+7r+6)}$	$\frac{1621}{40320}$
a_4	$\frac{r^8+8r^7+24r^6+11r^5-53r^4-153r^3-75r^2-1458r-897}{(2r^2+15r+28)(4r^4+20r^3+35r^2+25r+6)(2r^2+11r+15)}$	$-\frac{426599}{5806080}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-r^{10} - 15r^9 - 92r^8 - 270r^7 - 316r^6 + 276r^5 + 970r^4 + 207r^3 - 4303r^2 + 2370r + 13486}{(2r^2 + 19r + 45)(2r^2 + 15r + 28)(4r^4 + 20r^3 + 35r^2 + 25r + 6)(2r^2 + 11r + 15)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = \frac{4670443}{425779200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2+r-6}{2r^2+3r+1}$	$-\frac{23}{12}$
a_2	$\frac{r^4+r^2-15r+31}{4r^4+20r^3+35r^2+25r+6}$	$\frac{127}{160}$
a_3	$\frac{-r^6-3r^5-3r^4+17r^3+26r^2+182r-74}{(2r^2+11r+15)(2r^2+3r+1)(2r^2+7r+6)}$	$\frac{1621}{40320}$
a_4	$\frac{r^8+8r^7+24r^6+11r^5-53r^4-153r^3-75r^2-1458r-897}{(2r^2+15r+28)(4r^4+20r^3+35r^2+25r+6)(2r^2+11r+15)}$	$-\frac{426599}{5806080}$
a_5	$\frac{-r^{10}-15r^9-92r^8-270r^7-316r^6+276r^5+970r^4+207r^3-4303r^2+2370r+13486}{(2r^2+19r+45)(2r^2+15r+28)(4r^4+20r^3+35r^2+25r+6)(2r^2+11r+15)}$	$\frac{4670443}{425779200}$

Using the above table, then the solution $y_1(t)$ is

$$\begin{aligned} y_1(t) &= \sqrt{t}(a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 \dots) \\ &= \sqrt{t} \left(1 - \frac{23t}{12} + \frac{127t^2}{160} + \frac{1621t^3}{40320} - \frac{426599t^4}{5806080} + \frac{4670443t^5}{425779200} + O(t^6) \right) \end{aligned}$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = \frac{-r^2 + r - 6}{2r^2 + 3r + 1}$$

Substituting $n = 2$ in Eq. (2B) gives

$$b_2 = \frac{r^4 + r^2 - 15r + 31}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$$

For $3 \leq n$ the recursive equation is

$$\begin{aligned} &b_{n-1}(n+r-1)(n+r-2) + 2b_n(n+r)(n+r-1) \\ &+ (n+r)b_n + b_{n-3} + 5b_{n-2} + 6b_{n-1} = 0 \end{aligned} \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{n^2b_{n-1} + 2nrb_{n-1} + r^2b_{n-1} - 3nb_{n-1} - 3rb_{n-1} + b_{n-3} + 5b_{n-2} + 8b_{n-1}}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = \frac{-n^2 b_{n-1} + 3n b_{n-1} - b_{n-3} - 5b_{n-2} - 8b_{n-1}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r^2+r-6}{2r^2+3r+1}$	-6
b_2	$\frac{r^4+r^2-15r+31}{4r^4+20r^3+35r^2+25r+6}$	$\frac{31}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{-r^6 - 3r^5 - 3r^4 + 17r^3 + 26r^2 + 182r - 74}{(2r^2 + 11r + 15)(2r^2 + 3r + 1)(2r^2 + 7r + 6)}$$

Which for the root $r = 0$ becomes

$$b_3 = -\frac{37}{45}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r^2+r-6}{2r^2+3r+1}$	-6
b_2	$\frac{r^4+r^2-15r+31}{4r^4+20r^3+35r^2+25r+6}$	$\frac{31}{6}$
b_3	$\frac{-r^6-3r^5-3r^4+17r^3+26r^2+182r-74}{(2r^2+11r+15)(2r^2+3r+1)(2r^2+7r+6)}$	$-\frac{37}{45}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{r^8 + 8r^7 + 24r^6 + 11r^5 - 53r^4 - 153r^3 - 75r^2 - 1458r - 897}{(2r^2 + 15r + 28)(4r^4 + 20r^3 + 35r^2 + 25r + 6)(2r^2 + 11r + 15)}$$

Which for the root $r = 0$ becomes

$$b_4 = -\frac{299}{840}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r^2+r-6}{2r^2+3r+1}$	-6
b_2	$\frac{r^4+r^2-15r+31}{4r^4+20r^3+35r^2+25r+6}$	$\frac{31}{6}$
b_3	$\frac{-r^6-3r^5-3r^4+17r^3+26r^2+182r-74}{(2r^2+11r+15)(2r^2+3r+1)(2r^2+7r+6)}$	$-\frac{37}{45}$
b_4	$\frac{r^8+8r^7+24r^6+11r^5-53r^4-153r^3-75r^2-1458r-897}{(2r^2+15r+28)(4r^4+20r^3+35r^2+25r+6)(2r^2+11r+15)}$	$-\frac{299}{840}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{-r^{10} - 15r^9 - 92r^8 - 270r^7 - 316r^6 + 276r^5 + 970r^4 + 207r^3 - 4303r^2 + 2370r + 13486}{(2r^2 + 19r + 45)(2r^2 + 15r + 28)(4r^4 + 20r^3 + 35r^2 + 25r + 6)(2r^2 + 11r + 15)}$$

Which for the root $r = 0$ becomes

$$b_5 = \frac{6743}{56700}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r^2+r-6}{2r^2+3r+1}$	-6
b_2	$\frac{r^4+r^2-15r+31}{4r^4+20r^3+35r^2+25r+6}$	$\frac{31}{6}$
b_3	$\frac{-r^6-3r^5-3r^4+17r^3+26r^2+182r-74}{(2r^2+11r+15)(2r^2+3r+1)(2r^2+7r+6)}$	$-\frac{37}{45}$
b_4	$\frac{r^8+8r^7+24r^6+11r^5-53r^4-153r^3-75r^2-1458r-897}{(2r^2+15r+28)(4r^4+20r^3+35r^2+25r+6)(2r^2+11r+15)}$	$-\frac{299}{840}$
b_5	$\frac{-r^{10}-15r^9-92r^8-270r^7-316r^6+276r^5+970r^4+207r^3-4303r^2+2370r+13486}{(2r^2+19r+45)(2r^2+15r+28)(4r^4+20r^3+35r^2+25r+6)(2r^2+11r+15)}$	$\frac{6743}{56700}$

Using the above table, then the solution $y_2(t)$ is

$$\begin{aligned} y_2(t) &= b_0 + b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_5t^5 + b_6t^6 \dots \\ &= 1 - 6t + \frac{31t^2}{6} - \frac{37t^3}{45} - \frac{299t^4}{840} + \frac{6743t^5}{56700} + O(t^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(t) &= c_1 y_1(t) + c_2 y_2(t) \\
 &= c_1 \sqrt{t} \left(1 - \frac{23t}{12} + \frac{127t^2}{160} + \frac{1621t^3}{40320} - \frac{426599t^4}{5806080} + \frac{4670443t^5}{425779200} + O(t^6) \right) \\
 &\quad + c_2 \left(1 - 6t + \frac{31t^2}{6} - \frac{37t^3}{45} - \frac{299t^4}{840} + \frac{6743t^5}{56700} + O(t^6) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y(t) &= y_h \\
 &= c_1 \sqrt{t} \left(1 - \frac{23t}{12} + \frac{127t^2}{160} + \frac{1621t^3}{40320} - \frac{426599t^4}{5806080} + \frac{4670443t^5}{425779200} + O(t^6) \right) \\
 &\quad + c_2 \left(1 - 6t + \frac{31t^2}{6} - \frac{37t^3}{45} - \frac{299t^4}{840} + \frac{6743t^5}{56700} + O(t^6) \right)
 \end{aligned}$$

Replacing t in the above with the original independent variable x using $t = x - 2$ results in

$$\begin{aligned}
 y &= c_1 \sqrt{x-2} \left(\frac{29}{6} - \frac{23x}{12} + \frac{127(x-2)^2}{160} + \frac{1621(x-2)^3}{40320} - \frac{426599(x-2)^4}{5806080} \right. \\
 &\quad \left. + \frac{4670443(x-2)^5}{425779200} + O((x-2)^6) \right) + c_2 \left(13 - 6x + \frac{31(x-2)^2}{6} - \frac{37(x-2)^3}{45} \right. \\
 &\quad \left. - \frac{299(x-2)^4}{840} + \frac{6743(x-2)^5}{56700} + O((x-2)^6) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 \sqrt{x-2} \left(\frac{29}{6} - \frac{23x}{12} + \frac{127(x-2)^2}{160} + \frac{1621(x-2)^3}{40320} - \frac{426599(x-2)^4}{5806080} \right. \\
 &\quad \left. + \frac{4670443(x-2)^5}{425779200} + O((x-2)^6) \right) + c_2 \left(13 - 6x + \frac{31(x-2)^2}{6} - \frac{37(x-2)^3}{45} \right. \\
 &\quad \left. - \frac{299(x-2)^4}{840} + \frac{6743(x-2)^5}{56700} + O((x-2)^6) \right)
 \end{aligned}$$

Verification of solutions

$$y = c_1 \sqrt{x-2} \left(\frac{29}{6} - \frac{23x}{12} + \frac{127(x-2)^2}{160} + \frac{1621(x-2)^3}{40320} - \frac{426599(x-2)^4}{5806080} \right. \\ \left. + \frac{4670443(x-2)^5}{425779200} + O((x-2)^6) \right) + c_2 \left(13 - 6x + \frac{31(x-2)^2}{6} - \frac{37(x-2)^3}{45} \right. \\ \left. - \frac{299(x-2)^4}{840} + \frac{6743(x-2)^5}{56700} + O((x-2)^6) \right)$$

Verified OK.

4.15.1 Maple step by step solution

Let's solve

$$(x-2)y'' + \frac{y'}{x} + (1+x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x(x-2)} - \frac{(1+x)y}{x-2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x(x-2)} + \frac{(1+x)y}{x-2} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x(x-2)}, P_3(x) = \frac{1+x}{x-2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -\frac{1}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x(x-2) + y' + (1+x)yx = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 1..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k- > k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) x^{k+r}$$

- Convert $x^m \cdot y''$ to series expansion for $m = 1..2$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-2+m}$$

- Shift index using $k- > k + 2 - m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) x^{k+r}$$

Rewrite ODE with series expansions

$$-a_0 r(-3+2r) x^{-1+r} + (-a_1(1+r)(-1+2r) + a_0 r(-1+r)) x^r + (-a_2(2+r)(1+2r) + a_1(1+r)r + a_0) x^{1+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(-3+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{0, \frac{3}{2}\right\}$$

- The coefficients of each power of x must be 0

$$[-a_1(1+r)(-1+2r) + a_0 r(-1+r) = 0, -a_2(2+r)(1+2r) + a_1(1+r)r + a_0 = 0]$$

- Solve for the dependent coefficient(s)

$$\left\{ a_1 = \frac{a_0 r(-1+r)}{2r^2+r-1}, a_2 = \frac{a_0(r^3-r^2+2r-1)}{4r^3+8r^2-r-2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$-2\left(k - \frac{1}{2} + r\right)(k+1+r)a_{k+1} + a_k(k+r)(k+r-1) + a_{k-1} + a_{k-2} = 0$$

- Shift index using $k \rightarrow k+2$

$$-2\left(k + \frac{3}{2} + r\right)(k+3+r)a_{k+3} + a_{k+2}(k+2+r)(k+1+r) + a_{k+1} + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+3} = \frac{k^2 a_{k+2} + 2k r a_{k+2} + r^2 a_{k+2} + 3k a_{k+2} + 3r a_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{(2k+3+2r)(k+3+r)}$$

- Recursion relation for $r = 0$

$$a_{k+3} = \frac{k^2 a_{k+2} + 3k a_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{(2k+3)(k+3)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = \frac{k^2 a_{k+2} + 3k a_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{(2k+3)(k+3)}, a_1 = 0, a_2 = \frac{a_0}{2} \right]$$

- Recursion relation for $r = \frac{3}{2}$

$$a_{k+3} = \frac{k^2 a_{k+2} + 6k a_{k+2} + a_k + a_{k+1} + \frac{35}{4} a_{k+2}}{(2k+6)(k+\frac{9}{2})}$$

- Solution for $r = \frac{3}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{3}{2}}, a_{k+3} = \frac{k^2 a_{k+2} + 6k a_{k+2} + a_k + a_{k+1} + \frac{35}{4} a_{k+2}}{(2k+6)(k+\frac{9}{2})}, a_1 = \frac{3a_0}{20}, a_2 = \frac{25a_0}{224} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{3}{2}} \right), a_{k+3} = \frac{k^2 a_{k+2} + 3k a_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{(2k+3)(k+3)}, a_1 = 0, a_2 = \frac{a_0}{2}, b_{k+3} = \frac{k^2}{20} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
      -> heuristic approach
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
      -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunC ODE, case a <> 0, e <> 0, c =
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 46

```
Order:=6;
dsolve((x-2)*diff(y(x), x$2) + 1/x*diff(y(x), x) + (x+1)*y(x) = 0,y(x),type='series',x=2);
```

$$y(x) = c_1 \sqrt{x-2} \left(1 - \frac{23}{12}(x-2) + \frac{127}{160}(x-2)^2 + \frac{1621}{40320}(x-2)^3 - \frac{426599}{5806080}(x-2)^4 + \frac{4670443}{425779200}(x-2)^5 + O((x-2)^6) \right) + c_2 \left(1 - 6(x-2) + \frac{31}{6}(x-2)^2 - \frac{37}{45}(x-2)^3 - \frac{299}{840}(x-2)^4 + \frac{6743}{56700}(x-2)^5 + O((x-2)^6) \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 105

```
AsymptoticDSolveValue[(x-2)*y''[x] + 1/x*y'[x] + (x+1)*y[x] ==0,y[x],{x,2,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{4670443(x-2)^5}{425779200} - \frac{426599(x-2)^4}{5806080} + \frac{1621(x-2)^3}{40320} + \frac{127}{160}(x-2)^2 - \frac{23(x-2)}{12} + 1 \right) \sqrt{x-2} + c_2 \left(\frac{6743(x-2)^5}{56700} - \frac{299}{840}(x-2)^4 - \frac{37}{45}(x-2)^3 + \frac{31}{6}(x-2)^2 - 6(x-2) + 1 \right)$$

4.16 problem 16

Internal problem ID [7237]

Internal file name [OUTPUT/6223_Sunday_June_05_2022_04_33_01_PM_45818337/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 16.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(1 + x)(3x - 1)y'' + y' \cos(x) - 3yx = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (298)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (299)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{y' \cos(x) - 3yx}{3x^2 + 2x - 1}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{\left(\cos(x)^2 + \cos(x)(6x+2) + 9(1+x)\left(x - \frac{1}{3}\right)\left(x + \frac{\sin(x)}{3}\right) \right) y' - 9x^2 y - 3 \cos(x) yx - 3y}{(3x^2 + 2x - 1)^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{(-18 \cos(x)^2 x + (\sin(x)^2 + (-9x^2 - 6x + 3) \sin(x) + 9x^4 - 6x^3 - 68x^2 - 34x - 14) \cos(x) + 6 \sin(x)) y' - 18x^2 y - 6 \cos(x) yx - 3y}{(3x^2 + 2x - 1)^3} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(\cos(x)^4 + (36x + 12) \cos(x)^3 + ((18x^2 + 12x - 6) \sin(x) - 63x^4 - 57x^3 + 356x^2 + 235x + 61) \cos(x) + 6 \sin(x)^2 x) y' - 18x^3 y - 6 \cos(x) yx - 3y}{(3x^2 + 2x - 1)^4} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{243(1+x)\left(x - \frac{1}{3}\right) \left(\left(\frac{\cos(x)^5}{81} + \left(\frac{20}{81} + \frac{20x}{27} \right) \cos(x)^4 + \left(\left(-\frac{10}{81} + \frac{10}{27}x^2 + \frac{20}{81}x \right) \sin(x) - \frac{25x^4}{9} + \frac{175}{81} + \frac{80x}{27} \right) \cos(x) + 6 \sin(x)^2 x \right) y' - 18x^4 y - 6 \cos(x) yx - 3y}{(3x^2 + 2x - 1)^5} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= y'(0) \\ F_1 &= 3y'(0) - 3y(0) \\ F_2 &= -15y(0) + 14y'(0) \\ F_3 &= -159y(0) + 140y'(0) \\ F_4 &= -1917y(0) + 1711y'(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^3 - \frac{5}{8}x^4 - \frac{53}{40}x^5 - \frac{213}{80}x^6\right) y(0) + \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{7}{12}x^4 + \frac{7}{6}x^5 + \frac{1711}{720}x^6\right) y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. The ode is normalized to be

$$y''(3x^2 + 2x - 1) + y' \cos(x) - 3yx = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}\right) (3x^2 + 2x - 1) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1}\right) \cos(x) - 3\left(\sum_{n=0}^{\infty} a_n x^n\right) x = 0 \quad (1)$$

Expanding $\cos(x)$ as Taylor series around $x = 0$ and keeping only the first 6 terms gives

$$\begin{aligned} \cos(x) &= -\frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 + \dots \\ &= -\frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\begin{aligned} &\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}\right) (3x^2 + 2x - 1) \\ &+ \left(-\frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2\right) \left(\sum_{n=1}^{\infty} n a_n x^{n-1}\right) - 3\left(\sum_{n=0}^{\infty} a_n x^n\right) x = 0 \end{aligned}$$

Expanding the second term in (1) gives

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) (3x^2 + 2x - 1) + -\frac{x^6}{720} \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \frac{x^4}{24} \\ & \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + 1 \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \frac{x^2}{2} \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - 3 \left(\sum_{n=0}^{\infty} a_n x^n \right) x = 0 \end{aligned}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} 3x^n a_n n(n-1) \right) + \left(\sum_{n=2}^{\infty} 2n x^{n-1} a_n (n-1) \right) \\ & + \sum_{n=2}^{\infty} (-n(n-1) a_n x^{n-2}) + \sum_{n=1}^{\infty} \left(-\frac{n x^{n+5} a_n}{720} \right) + \left(\sum_{n=1}^{\infty} \frac{n x^{n+3} a_n}{24} \right) \\ & + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=1}^{\infty} \left(-\frac{n x^{1+n} a_n}{2} \right) + \sum_{n=0}^{\infty} (-3x^{1+n} a_n) = 0 \end{aligned} \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} 2n x^{n-1} a_n (n-1) &= \sum_{n=1}^{\infty} 2(1+n) a_{1+n} n x^n \\ \sum_{n=2}^{\infty} (-n(n-1) a_n x^{n-2}) &= \sum_{n=0}^{\infty} (-(n+2) a_{n+2} (1+n) x^n) \\ \sum_{n=1}^{\infty} \left(-\frac{n x^{n+5} a_n}{720} \right) &= \sum_{n=6}^{\infty} \left(-\frac{(n-5) a_{n-5} x^n}{720} \right) \\ \sum_{n=1}^{\infty} \frac{n x^{n+3} a_n}{24} &= \sum_{n=4}^{\infty} \frac{(n-3) a_{n-3} x^n}{24} \\ \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \\ \sum_{n=1}^{\infty} \left(-\frac{n x^{1+n} a_n}{2} \right) &= \sum_{n=2}^{\infty} \left(-\frac{(n-1) a_{n-1} x^n}{2} \right) \end{aligned}$$

$$\sum_{n=0}^{\infty} (-3x^{1+n}a_n) = \sum_{n=1}^{\infty} (-3a_{n-1}x^n)$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} 3x^n a_n n(n-1) \right) + \left(\sum_{n=1}^{\infty} 2(1+n) a_{1+n} n x^n \right) \\ & + \sum_{n=0}^{\infty} (-(n+2) a_{n+2} (1+n) x^n) + \sum_{n=6}^{\infty} \left(-\frac{(n-5) a_{n-5} x^n}{720} \right) \\ & + \left(\sum_{n=4}^{\infty} \frac{(n-3) a_{n-3} x^n}{24} \right) + \left(\sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \right) \\ & + \sum_{n=2}^{\infty} \left(-\frac{(n-1) a_{n-1} x^n}{2} \right) + \sum_{n=1}^{\infty} (-3a_{n-1} x^n) = 0 \end{aligned} \quad (3)$$

$n = 0$ gives

$$-2a_2 + a_1 = 0$$

$$a_2 = \frac{a_1}{2}$$

$n = 1$ gives

$$6a_2 - 6a_3 - 3a_0 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{a_0}{2} + \frac{a_1}{2}$$

$n = 2$ gives

$$6a_2 + 15a_3 - 12a_4 - \frac{7a_1}{2} = 0$$

Which after substituting earlier equations, simplifies to

$$a_4 = -\frac{5a_0}{8} + \frac{7a_1}{12}$$

$n = 3$ gives

$$18a_3 + 28a_4 - 20a_5 - 4a_2 = 0$$

Which after substituting earlier equations, simplifies to

$$-\frac{53a_0}{2} + \frac{70a_1}{3} - 20a_5 = 0$$

Or

$$a_5 = -\frac{53a_0}{40} + \frac{7a_1}{6}$$

$n = 4$ gives

$$36a_4 + 45a_5 - 30a_6 + \frac{a_1}{24} - \frac{9a_3}{2} = 0$$

Which after substituting earlier equations, simplifies to

$$a_6 = -\frac{213a_0}{80} + \frac{1711a_1}{720}$$

$n = 5$ gives

$$60a_5 + 66a_6 - 42a_7 + \frac{a_2}{12} - 5a_4 = 0$$

Which after substituting earlier equations, simplifies to

$$-\frac{2521a_0}{10} + \frac{6719a_1}{30} - 42a_7 = 0$$

Or

$$a_7 = -\frac{2521a_0}{420} + \frac{6719a_1}{1260}$$

For $6 \leq n$, the recurrence equation is

$$\begin{aligned} & 3na_n(n-1) + 2(1+n)a_{1+n}n - (n+2)a_{n+2}(1+n) - \frac{(n-5)a_{n-5}}{720} \\ & + \frac{(n-3)a_{n-3}}{24} + (1+n)a_{1+n} - \frac{(n-1)a_{n-1}}{2} - 3a_{n-1} = 0 \end{aligned} \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned} & a_{n+2} \\ &= \frac{2160n^2 a_n + 1440n^2 a_{1+n} - 2160n a_n + 2160n a_{1+n} - n a_{n-5} + 30n a_{n-3} - 360n a_{n-1} + 720 a_{1+n} + 5 a_{n-5}}{720(n+2)(1+n)} \\ (5) \quad &= \frac{(2160n^2 - 2160n) a_n}{720(n+2)(1+n)} + \frac{(1440n^2 + 2160n + 720) a_{1+n}}{720(n+2)(1+n)} \\ &+ \frac{(-n+5) a_{n-5}}{720(n+2)(1+n)} + \frac{(30n-90) a_{n-3}}{720(n+2)(1+n)} + \frac{(-360n-1800) a_{n-1}}{720(n+2)(1+n)} \end{aligned}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + \frac{a_1 x^2}{2} + \left(-\frac{a_0}{2} + \frac{a_1}{2}\right) x^3 + \left(-\frac{5a_0}{8} + \frac{7a_1}{12}\right) x^4 + \left(-\frac{53a_0}{40} + \frac{7a_1}{6}\right) x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^3 - \frac{5}{8}x^4 - \frac{53}{40}x^5\right) a_0 + \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{7}{12}x^4 + \frac{7}{6}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^3 - \frac{5}{8}x^4 - \frac{53}{40}x^5\right) c_1 + \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{7}{12}x^4 + \frac{7}{6}x^5\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^3 - \frac{5}{8}x^4 - \frac{53}{40}x^5 - \frac{213}{80}x^6\right) y(0) \\ &+ \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{7}{12}x^4 + \frac{7}{6}x^5 + \frac{1711}{720}x^6\right) y'(0) + O(x^6) \end{aligned} \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^3 - \frac{5}{8}x^4 - \frac{53}{40}x^5\right) c_1 + \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{7}{12}x^4 + \frac{7}{6}x^5\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^3 - \frac{5}{8}x^4 - \frac{53}{40}x^5 - \frac{213}{80}x^6\right) y(0) \\ + \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{7}{12}x^4 + \frac{7}{6}x^5 + \frac{1711}{720}x^6\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^3 - \frac{5}{8}x^4 - \frac{53}{40}x^5\right) c_1 + \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{7}{12}x^4 + \frac{7}{6}x^5\right) c_2 + O(x^6)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
trying a symmetry of the form [xi=0, eta=F(x)]
trying differential order: 2; exact nonlinear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying 2nd order, integrating factor of the form mu(x,y)
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) *
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

Order:=6;

```
dsolve((x+1)*(3*x-1)*diff(y(x),x$2)+cos(x)*diff(y(x),x)-3*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^3 - \frac{5}{8}x^4 - \frac{53}{40}x^5\right) y(0) + \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{7}{12}x^4 + \frac{7}{6}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[(x+1)*(3*x-1)*y'[x]+Cos[x]*y'[x]-3*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{53x^5}{40} - \frac{5x^4}{8} - \frac{x^3}{2} + 1 \right) + c_2 \left(\frac{7x^5}{6} + \frac{7x^4}{12} + \frac{x^3}{2} + \frac{x^2}{2} + x \right)$$

4.17 problem 17

4.17.1 Existence and uniqueness analysis	1675
4.17.2 Maple step by step solution	1685

Internal problem ID [7238]

Internal file name [OUTPUT/6224_Sunday_June_05_2022_04_33_07_PM_33923488/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 17.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

[_Lienard]

$$xy'' + 2y' + yx = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

4.17.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = \frac{2}{x}$$

$$q(x) = 1$$

$$F = 0$$

Hence the ode is

$$y'' + \frac{2y'}{x} + y = 0$$

The domain of $p(x) = \frac{2}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

But the point $x_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$xy'' + 2y' + yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{2}{x}$$

$$q(x) = 1$$

Table 168: Table $p(x), q(x)$ singularities.

$p(x) = \frac{2}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = 1$	
singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$xy'' + 2y' + yx = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2(n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{1+n+r} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2(n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + 2(n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + 2r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + 2r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} (1+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(1+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = -1$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} (1+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 2a_n(n+r) + a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2 + n + r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-2}}{n(1+n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{r^2 + 5r + 6}$$

Which for the root $r = 0$ becomes

$$a_2 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{r^4 + 14r^3 + 71r^2 + 154r + 120}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{120}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{120}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -1} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-1} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + 2(n+r)b_n + b_{n-2} = 0 \quad (4)$$

Which for the root $r = -1$ becomes

$$b_n(n-1)(n-2) + 2(n-1)b_n + b_{n-2} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{b_{n-2}}{n^2 + 2nr + r^2 + n + r} \quad (5)$$

Which for the root $r = -1$ becomes

$$b_n = -\frac{b_{n-2}}{n^2 - n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{1}{r^2 + 5r + 6}$$

Which for the root $r = -1$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{(r^2 + 5r + 6)(r^2 + 9r + 20)}$$

Which for the root $r = -1$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{24}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)}{x} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)\right)}{x} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)\right)}{x} \end{aligned}$$

$$y = 1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)$$

Summary

The solution(s) found are the following

$$y = 1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \tag{1}$$

Verification of solutions

$$y = 1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)$$

Verified OK.

4.17.2 Maple step by step solution

Let's solve

$$\left[xy'' + 2y' + yx = 0, y(0) = 1, y'|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y'}{x} - y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y'}{x} + y = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{2}{x}, P_3(x) = 1]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$xy'' + 2y' + yx = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(1+r) x^{-1+r} + a_1 (1+r)(2+r) x^r + \left(\sum_{k=1}^{\infty} (a_{k+1} (k+r+1)(k+2+r) + a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(1+r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \{-1, 0\}$$
- Each term must be 0

$$a_1 (1+r)(2+r) = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_{k+1} (k+r+1)(k+2+r) + a_{k-1} = 0$$
- Shift index using $k \rightarrow k + 1$

$$a_{k+2}(k+2+r)(k+3+r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+2+r)(k+3+r)}$$

- Recursion relation for $r = -1$

$$a_{k+2} = -\frac{a_k}{(k+1)(k+2)}$$

- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+2} = -\frac{a_k}{(k+1)(k+2)}, 0 = 0 \right]$$

- Recursion relation for $r = 0$

$$a_{k+2} = -\frac{a_k}{(k+2)(k+3)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{(k+2)(k+3)}, 2a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k x^k \right), a_{k+2} = -\frac{a_k}{(k+1)(k+2)}, 0 = 0, b_{k+2} = -\frac{b_k}{(k+2)(k+3)}, 2b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

Order:=6;

```
dsolve([x*diff(y(x),x$2)+2*diff(y(x),x)+x*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series',x
```

$$y(x) = 1 - \frac{1}{6}x^2 + \frac{1}{120}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{x*y'[x]+2*y'[x]+x*y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^4}{120} - \frac{x^2}{6} + 1$$

4.18 problem 18

Internal problem ID [7239]

Internal file name [OUTPUT/6225_Sunday_June_05_2022_04_33_09_PM_21549085/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 18.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2x^2y'' + 3xy' - yx = x^2 + 2x$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + 3xy' - yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3}{2x}$$
$$q(x) = -\frac{1}{2x}$$

Table 170: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + 3xy' - yx = x^2 + 2x$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' + 3xy' - yx = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & 2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + 3x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + 3x^{n+r} a_n (n+r) = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r (-1+r) + 3x^r a_0 r = 0$$

Or

$$(2x^r r (-1+r) + 3x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 + r) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 + r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -\frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m (-1+m) + 3x^m m) c_0 = x^2 + 2x$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 + r)x^r = 0$$

Solving for r gives the roots of the indicial equation as $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + 3a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{2n^2 + 4nr + 2r^2 + n + r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 0$ becomes

$$a_1 = \frac{1}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2r^2+5r+3}$	$\frac{1}{3}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4r^4 + 28r^3 + 71r^2 + 77r + 30}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{30}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2r^2+5r+3}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^4+28r^3+71r^2+77r+30}$	$\frac{1}{30}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{8r^6 + 108r^5 + 590r^4 + 1665r^3 + 2552r^2 + 2007r + 630}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{630}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2r^2+5r+3}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^4+28r^3+71r^2+77r+30}$	$\frac{1}{30}$
a_3	$\frac{1}{8r^6+108r^5+590r^4+1665r^3+2552r^2+2007r+630}$	$\frac{1}{630}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16r^8 + 352r^7 + 3304r^6 + 17248r^5 + 54649r^4 + 107338r^3 + 127251r^2 + 82962r + 22680}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{22680}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2r^2+5r+3}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^4+28r^3+71r^2+77r+30}$	$\frac{1}{30}$
a_3	$\frac{1}{8r^6+108r^5+590r^4+1665r^3+2552r^2+2007r+630}$	$\frac{1}{630}$
a_4	$\frac{1}{16r^8+352r^7+3304r^6+17248r^5+54649r^4+107338r^3+127251r^2+82962r+22680}$	$\frac{1}{22680}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{32r^{10} + 1040r^9 + 14880r^8 + 123240r^7 + 653226r^6 + 2310945r^5 + 5514295r^4 + 8741785r^3 + 8786367r^2 + 4443183r + 1102200}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{1102200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2r^2+5r+3}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^4+28r^3+71r^2+77r+30}$	$\frac{1}{30}$
a_3	$\frac{1}{8r^6+108r^5+590r^4+1665r^3+2552r^2+2007r+630}$	$\frac{1}{630}$
a_4	$\frac{1}{16r^8+352r^7+3304r^6+17248r^5+54649r^4+107338r^3+127251r^2+82962r+22680}$	$\frac{1}{22680}$
a_5	$\frac{1}{32r^{10}+1040r^9+14880r^8+123240r^7+653226r^6+2310945r^5+5514295r^4+8741785r^3+8786367r^2+5039190r+1247400}$	$\frac{1}{1247400}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\
 &= 1 + \frac{x}{3} + \frac{x^2}{30} + \frac{x^3}{630} + \frac{x^4}{22680} + \frac{x^5}{1247400} + O(x^6)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) + 3b_n(n+r) - b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-1}}{2n^2 + 4nr + 2r^2 + n + r} \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = \frac{b_{n-1}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_1 = 1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2r^2+5r+3}$	1

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{4r^4 + 28r^3 + 71r^2 + 77r + 30}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2r^2+5r+3}$	1
b_2	$\frac{1}{4r^4+28r^3+71r^2+77r+30}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{1}{8r^6 + 108r^5 + 590r^4 + 1665r^3 + 2552r^2 + 2007r + 630}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_3 = \frac{1}{90}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2r^2+5r+3}$	1
b_2	$\frac{1}{4r^4+28r^3+71r^2+77r+30}$	$\frac{1}{6}$
b_3	$\frac{1}{8r^6+108r^5+590r^4+1665r^3+2552r^2+2007r+630}$	$\frac{1}{90}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{16r^8 + 352r^7 + 3304r^6 + 17248r^5 + 54649r^4 + 107338r^3 + 127251r^2 + 82962r + 22680}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = \frac{1}{2520}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2r^2+5r+3}$	1
b_2	$\frac{1}{4r^4+28r^3+71r^2+77r+30}$	$\frac{1}{6}$
b_3	$\frac{1}{8r^6+108r^5+590r^4+1665r^3+2552r^2+2007r+630}$	$\frac{1}{90}$
b_4	$\frac{1}{16r^8+352r^7+3304r^6+17248r^5+54649r^4+107338r^3+127251r^2+82962r+22680}$	$\frac{1}{2520}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{1}{32r^{10} + 1040r^9 + 14880r^8 + 123240r^7 + 653226r^6 + 2310945r^5 + 5514295r^4 + 8741785r^3 + 8786367r^2 + 5039190r + 1247400}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_5 = \frac{1}{113400}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2r^2+5r+3}$	1
b_2	$\frac{1}{4r^4+28r^3+71r^2+77r+30}$	$\frac{1}{6}$
b_3	$\frac{1}{8r^6+108r^5+590r^4+1665r^3+2552r^2+2007r+630}$	$\frac{1}{90}$
b_4	$\frac{1}{16r^8+352r^7+3304r^6+17248r^5+54649r^4+107338r^3+127251r^2+82962r+22680}$	$\frac{1}{2520}$
b_5	$\frac{1}{32r^{10}+1040r^9+14880r^8+123240r^7+653226r^6+2310945r^5+5514295r^4+8741785r^3+8786367r^2+5039190r+1247400}$	$\frac{1}{113400}$

Using the above table, then the solution $y_2(x)$ is

$$y_2(x) = 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots)$$

$$= \frac{1 + x + \frac{x^2}{6} + \frac{x^3}{90} + \frac{x^4}{2520} + \frac{x^5}{113400} + O(x^6)}{\sqrt{x}}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1y_1(x) + c_2y_2(x)$$

$$= c_1\left(1 + \frac{x}{3} + \frac{x^2}{30} + \frac{x^3}{630} + \frac{x^4}{22680} + \frac{x^5}{1247400} + O(x^6)\right)$$

$$+ \frac{c_2\left(1 + x + \frac{x^2}{6} + \frac{x^3}{90} + \frac{x^4}{2520} + \frac{x^5}{113400} + O(x^6)\right)}{\sqrt{x}}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1 + m) + 3x^m m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$$a_1 = \frac{a_0}{2r^2+5r+3}$$

$$a_2 = \frac{a_0}{4r^4+28r^3+71r^2+77r+30}$$

$$a_3 = \frac{a_0}{8r^6+108r^5+590r^4+1665r^3+2552r^2+2007r+630}$$

$$a_4 = \frac{a_0}{16r^8+352r^7+3304r^6+17248r^5+54649r^4+107338r^3+127251r^2+82962r+22680}$$

$$a_5 = \frac{a_0}{32r^{10}+1040r^9+14880r^8+123240r^7+653226r^6+2310945r^5+5514295r^4+8741785r^3+8786367r^2+5039190r+1247400}$$

Since the $F = x^2 + 2x$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x^2$ by solving the balance equation

$$(2x^m m(-1 + m) + 3x^m m) c_0 = x^2$$

For c_0 and x . This results in

$$c_0 = \frac{1}{10}$$

$$m = 2$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+2}$$

Where in the above $c_0 = \frac{1}{10}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{10}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{10}$
$c_1 = \frac{1}{210}$
$c_2 = \frac{1}{7560}$
$c_3 = \frac{1}{415800}$
$c_4 = \frac{1}{32432400}$
$c_5 = \frac{1}{3405402000}$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^2 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$y_p = x^2 \left(\frac{1}{10} + \frac{1}{210}x + \frac{1}{7560}x^2 + \frac{1}{415800}x^3 + \frac{1}{32432400}x^4 + \frac{1}{3405402000}x^5 \right)$$

$$= \frac{1}{10}x^2 + \frac{1}{210}x^3 + \frac{1}{7560}x^4 + \frac{1}{415800}x^5 + \frac{1}{32432400}x^6 + \frac{1}{3405402000}x^7$$

Now we determine the particular solution y_p associated with $F = 2x$ by solving the balance equation

$$(2x^m m(-1 + m) + 3x^m m) c_0 = 2x$$

For c_0 and x . This results in

$$c_0 = \frac{2}{3}$$

$$m = 1$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+1}$$

Where in the above $c_0 = \frac{2}{3}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 1$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{2}{3}$ and $r = m$ or $r = 1$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{2}{3}$
$c_1 = \frac{1}{15}$
$c_2 = \frac{1}{315}$
$c_3 = \frac{1}{11340}$
$c_4 = \frac{1}{623700}$
$c_5 = \frac{1}{48648600}$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x \left(\frac{2}{3} + \frac{1}{15}x + \frac{1}{315}x^2 + \frac{1}{11340}x^3 + \frac{1}{623700}x^4 + \frac{1}{48648600}x^5 \right) \\ &= \frac{2}{3}x + \frac{1}{15}x^2 + \frac{1}{315}x^3 + \frac{1}{11340}x^4 + \frac{1}{623700}x^5 + \frac{1}{48648600}x^6 \end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = \frac{2x}{3} + \frac{x^2}{6} + \frac{x^3}{126} + \frac{x^4}{4536} + \frac{x^5}{249480} + \frac{x^6}{19459440} + \frac{x^7}{3405402000} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{2x}{3} + \frac{x^2}{6} + \frac{x^3}{126} + \frac{x^4}{4536} + \frac{x^5}{249480} + O(x^6)$$

Hence the final solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \frac{2x}{3} + \frac{x^2}{6} + \frac{x^3}{126} + \frac{x^4}{4536} + \frac{x^5}{249480} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{3} + \frac{x^2}{30} + \frac{x^3}{630} + \frac{x^4}{22680} + \frac{x^5}{1247400} + O(x^6) \right) \\ &\quad + \frac{c_2 \left(1 + x + \frac{x^2}{6} + \frac{x^3}{90} + \frac{x^4}{2520} + \frac{x^5}{113400} + O(x^6) \right)}{\sqrt{x}} \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \frac{2x}{3} + \frac{x^2}{6} + \frac{x^3}{126} + \frac{x^4}{4536} + \frac{x^5}{249480} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{3} + \frac{x^2}{30} + \frac{x^3}{630} + \frac{x^4}{22680} + \frac{x^5}{1247400} + O(x^6) \right) \\ &\quad + \frac{c_2 \left(1 + x + \frac{x^2}{6} + \frac{x^3}{90} + \frac{x^4}{2520} + \frac{x^5}{113400} + O(x^6) \right)}{\sqrt{x}} \end{aligned} \tag{1}$$

Verification of solutions

$$y = \frac{2x}{3} + \frac{x^2}{6} + \frac{x^3}{126} + \frac{x^4}{4536} + \frac{x^5}{249480} + O(x^6) \\ + c_1 \left(1 + \frac{x}{3} + \frac{x^2}{30} + \frac{x^3}{630} + \frac{x^4}{22680} + \frac{x^5}{1247400} + O(x^6) \right) \\ + \frac{c_2 \left(1 + x + \frac{x^2}{6} + \frac{x^3}{90} + \frac{x^4}{2520} + \frac{x^5}{113400} + O(x^6) \right)}{\sqrt{x}}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 60

Order:=6;

dsolve(2*x^2*diff(y(x),x\$2)+3*x*diff(y(x),x)-x*y(x)=x^2+2*x,y(x),type='series',x=0);

$$y(x) = \frac{c_1 \left(1 + x + \frac{1}{6}x^2 + \frac{1}{90}x^3 + \frac{1}{2520}x^4 + \frac{1}{113400}x^5 + O(x^6)\right)}{\sqrt{x}} + c_2 \left(1 + \frac{1}{3}x + \frac{1}{30}x^2 + \frac{1}{630}x^3 + \frac{1}{22680}x^4 + \frac{1}{1247400}x^5 + O(x^6)\right) + x \left(\frac{2}{3} + \frac{1}{6}x + \frac{1}{126}x^2 + \frac{1}{4536}x^3 + \frac{1}{249480}x^4 + O(x^5)\right)$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 239

AsymptoticDSolveValue[2*x^2*y''[x]+3*x*y'[x]-x*y[x]==x^2+2*x,y[x],{x,0,5}]

$y(x)$

$$\rightarrow c_1 \left(\frac{x^5}{1247400} + \frac{x^4}{22680} + \frac{x^3}{630} + \frac{x^2}{30} + \frac{x}{3} + 1 \right) + \frac{c_2 \left(\frac{x^5}{113400} + \frac{x^4}{2520} + \frac{x^3}{90} + \frac{x^2}{6} + x + 1 \right)}{\sqrt{x}} + \frac{\left(\frac{x^5}{113400} + \frac{x^4}{2520} + \frac{x^3}{90} + \frac{x^2}{6} + x + 1 \right) \left(-\frac{19x^{11/2}}{62370} - \frac{23x^{9/2}}{2835} - \frac{4x^{7/2}}{35} - \frac{2x^{5/2}}{3} - \frac{4x^{3/2}}{3} \right)}{\sqrt{x}} + \left(\frac{x^5}{1247400} + \frac{x^4}{22680} + \frac{x^3}{630} + \frac{x^2}{30} + \frac{x}{3} + 1 \right) \left(\frac{47x^6}{680400} + \frac{x^5}{420} + \frac{17x^4}{360} + \frac{4x^3}{9} + \frac{3x^2}{2} + 2x \right)$$

4.19 problem 19

Internal problem ID [7240]

Internal file name [OUTPUT/6226_Sunday_June_05_2022_04_33_11_PM_25930002/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 19.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = 1$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 171: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	"regular"

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	"regular"

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : [0]

Irregular singular points : [∞]

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = 1$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= 1 \\ m &= 0 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+0} \end{aligned}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = 1$
$c_1 = 0$
$c_2 = \frac{1}{3}$
$c_3 = 0$
$c_4 = \frac{1}{63}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= 1 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\ &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4\end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = 1 + \frac{x^2}{3} + \frac{x^4}{63} + O(x^6)$$

Hence the final solution is

$$\begin{aligned}y &= y_h + y_p \\ &= 1 + \frac{x^2}{3} + \frac{x^4}{63} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= 1 + \frac{x^2}{3} + \frac{x^4}{63} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ &\quad + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)\end{aligned}\tag{1}$$

Verification of solutions

$$y = 1 + \frac{x^2}{3} + \frac{x^4}{63} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 43

```
Order:=6;
dsolve(2*x^2*diff(y(x), x$2) - x*diff(y(x), x) + (-x^2 + 1)*y(x) = 1,y(x),type='series',x=0)
```

$$y(x) = c_1\sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 176

```
AsymptoticDSolveValue[2*x^2*y'[x]-x*y'[x]+(1-x^2)*y[x]==1,y[x],{x,0,5}]
```

$y(x)$

$$\begin{aligned} &\rightarrow c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ &+ c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{154440} - \frac{x^{7/2}}{1260} - \frac{x^{3/2}}{15} \right. \\ &\left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^5}{55440} + \frac{x^3}{504} + \frac{x}{6} - \frac{1}{x} \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \end{aligned}$$

4.20 problem 20

Internal problem ID [7241]

Internal file name [OUTPUT/6227_Sunday_June_05_2022_04_33_12_PM_60709388/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 20.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$2x^2y'' + 2xy' - yx = 1$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + 2xy' - yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{2x}$$

Table 172: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + 2xy' - yx = 1$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' + 2xy' - yx = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 2x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r)(n+r-1) + 2x^{n+r} a_n (n+r) = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r(-1+r) + 2x^r a_0 r = 0$$

Or

$$(2x^r r(-1+r) + 2x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$2x^r r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= 0 \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) + 2x^m m) c_0 = 1$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$2x^r r^2 = 0$$

Solving for r gives the roots of the indicial equation as. Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + 2a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{2n^2 + 4nr + 2r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{2n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{2(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4(r+1)^2(2+r)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{16}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{288}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{9216}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{460800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$
a_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r=0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$	$-\frac{1}{(r+1)^3}$	-1
b_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$	$\frac{-3-2r}{2(r+1)^3(2+r)^3}$	$-\frac{3}{16}$
b_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$	$\frac{-3r^2-12r-11}{4(r+1)^3(2+r)^3(r+3)^3}$	$-\frac{11}{864}$
b_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$	$\frac{-2r^3-15r^2-35r-25}{4(r+1)^3(2+r)^3(r+3)^3(r+4)^3}$	$-\frac{25}{55296}$
b_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$	$\frac{-5r^4-60r^3-255r^2-450r-274}{16(r+1)^3(2+r)^3(r+3)^3(r+4)^3(r+5)^3}$	$-\frac{137}{13824000}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned}
y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\
&= \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) \\
&\quad - x - \frac{3x^2}{16} - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
&= c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\
&\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\
&\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right)
\end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) + 2x^m m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$$a_1 = \frac{a_0}{2(r+1)^2}$$

$$a_2 = \frac{a_0}{4(r+1)^2(2+r)^2}$$

$$a_3 = \frac{a_0}{8(r+1)^2(2+r)^2(r+3)^2}$$

$$a_4 = \frac{a_0}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

$$a_5 = \frac{a_0}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Unable to solve the balance equation $(2x^m m(-1 + m) + 2x^m m) c_0$ for c_0 and x . No particular solution exists.

Adding all the above particular solution(s) gives

$$y_p = \text{FAIL}$$

Unable to find the particular solution or no solution exists.

Verification of solutions N/A

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`

```

✗ Solution by Maple

```
Order:=6;  
dsolve(2*x^2*diff(y(x), x, x) + 2*x*diff(y(x), x) - x*y(x) = 1,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.148 (sec). Leaf size: 360

```
AsymptoticDSolveValue[2*x^2*y''[x]+2*x*y'[x]-x*y[x]==1,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_2 \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \\ & + c_1 \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) \right. \\ & \quad \left. + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \right) \\ & + \left(-\frac{137x^6}{1990656000} + \frac{x^5}{4608000} + \frac{x^4}{73728} + \frac{x^3}{1728} + \frac{x^2}{64} + \frac{x}{4} \right. \\ & \quad \left. + \frac{\log(x)}{2} \right) \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) \right. \\ & \quad \left. + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \right) \\ & + \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \left(\frac{137x^6(6 \log(x) + 5)}{11943936000} \right. \\ & \quad \left. + \frac{x^5(113 - 30 \log(x))}{138240000} + \frac{x^4(41 - 12 \log(x))}{884736} + \frac{x^3(3 - \log(x))}{1728} \right. \\ & \quad \left. + \frac{1}{128}x^2(5 - 2 \log(x)) + \frac{1}{4}x(2 - \log(x)) - \frac{1}{4} \log(x)(\log(x) + 2) \right) \end{aligned}$$

4.21 problem 21

4.21.1 Maple step by step solution 1734

Internal problem ID [7242]

Internal file name [OUTPUT/6228_Sunday_June_05_2022_04_33_14_PM_93469576/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 21.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + (x - 6)y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (304)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (305)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -(x - 6)y \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= (-x + 6)y' - y \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= -2y' + (x^2 - 12x + 36)y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= (x - 6)((x - 6)y' + 4y) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= -(x - 6)^3y + 6(x - 6)y' + 4y
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 6y(0) \\
 F_1 &= 6y'(0) - y(0) \\
 F_2 &= -2y'(0) + 36y(0) \\
 F_3 &= 36y'(0) - 24y(0) \\
 F_4 &= 220y(0) - 36y'(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned}
 y &= \left(1 + 3x^2 - \frac{1}{6}x^3 + \frac{3}{2}x^4 - \frac{1}{5}x^5 + \frac{11}{36}x^6\right)y(0) \\
 &\quad + \left(x + x^3 - \frac{1}{12}x^4 + \frac{3}{10}x^5 - \frac{1}{20}x^6\right)y'(0) + O(x^6)
 \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -(x-6) \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} x^{1+n} a_n \right) + \sum_{n=0}^{\infty} (-6a_n x^n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=0}^{\infty} x^{1+n} a_n = \sum_{n=1}^{\infty} a_{n-1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^n \right) + \sum_{n=0}^{\infty} (-6a_n x^n) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 - 6a_0 = 0$$

$$a_2 = 3a_0$$

For $1 \leq n$, the recurrence equation is

$$(n + 2) a_{n+2}(1 + n) + a_{n-1} - 6a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$(5) \quad \begin{aligned} a_{n+2} &= \frac{-a_{n-1} + 6a_n}{(n + 2)(1 + n)} \\ &= \frac{6a_n}{(n + 2)(1 + n)} - \frac{a_{n-1}}{(n + 2)(1 + n)} \end{aligned}$$

For $n = 1$ the recurrence equation gives

$$6a_3 + a_0 - 6a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{6} + a_1$$

For $n = 2$ the recurrence equation gives

$$12a_4 + a_1 - 6a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_1}{12} + \frac{3a_0}{2}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + a_2 - 6a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_0}{5} + \frac{3a_1}{10}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + a_3 - 6a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{11a_0}{36} - \frac{a_1}{20}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + a_4 - 6a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{113a_1}{2520} - \frac{9a_0}{140}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + 3a_0 x^2 + \left(-\frac{a_0}{6} + a_1\right) x^3 + \left(-\frac{a_1}{12} + \frac{3a_0}{2}\right) x^4 + \left(-\frac{a_0}{5} + \frac{3a_1}{10}\right) x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + 3x^2 - \frac{1}{6}x^3 + \frac{3}{2}x^4 - \frac{1}{5}x^5\right) a_0 + \left(x + x^3 - \frac{1}{12}x^4 + \frac{3}{10}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + 3x^2 - \frac{1}{6}x^3 + \frac{3}{2}x^4 - \frac{1}{5}x^5\right) c_1 + \left(x + x^3 - \frac{1}{12}x^4 + \frac{3}{10}x^5\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 + 3x^2 - \frac{1}{6}x^3 + \frac{3}{2}x^4 - \frac{1}{5}x^5 + \frac{11}{36}x^6\right) y(0) \\ + \left(x + x^3 - \frac{1}{12}x^4 + \frac{3}{10}x^5 - \frac{1}{20}x^6\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 + 3x^2 - \frac{1}{6}x^3 + \frac{3}{2}x^4 - \frac{1}{5}x^5\right) c_1 + \left(x + x^3 - \frac{1}{12}x^4 + \frac{3}{10}x^5\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 + 3x^2 - \frac{1}{6}x^3 + \frac{3}{2}x^4 - \frac{1}{5}x^5 + \frac{11}{36}x^6\right) y(0) \\ + \left(x + x^3 - \frac{1}{12}x^4 + \frac{3}{10}x^5 - \frac{1}{20}x^6\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 + 3x^2 - \frac{1}{6}x^3 + \frac{3}{2}x^4 - \frac{1}{5}x^5\right) c_1 + \left(x + x^3 - \frac{1}{12}x^4 + \frac{3}{10}x^5\right) c_2 + O(x^6)$$

Verified OK.

4.21.1 Maple step by step solution

Let's solve

$$y'' = -(x - 6)y$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = (-x + 6)y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + (x - 6)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=\max(0,-m)}^{\infty} a_k x^{k+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=\max(0,-m)+m}^{\infty} a_{k-m} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 - 6a_0 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) - 6a_k + a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 - 6a_0 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - 6a_k + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} - 6a_{k+1} + a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{-6a_{k+1} + a_k}{k^2 + 5k + 6}, 2a_2 - 6a_0 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
    -> Bessel  
    <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
Order:=6;  
dsolve(diff(y(x), x, x) + (x-6)*y(x) = 0, y(x), type='series', x=0);
```

$$y(x) = \left(1 + 3x^2 - \frac{1}{6}x^3 + \frac{3}{2}x^4 - \frac{1}{5}x^5\right) y(0) + \left(x + x^3 - \frac{1}{12}x^4 + \frac{3}{10}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 57

```
AsymptoticDSolveValue[y''[x] + (x-6)*y[x] == 0, y[x], {x, 0, 5}]
```

$$y(x) \rightarrow c_2 \left(\frac{3x^5}{10} - \frac{x^4}{12} + x^3 + x \right) + c_1 \left(-\frac{x^5}{5} + \frac{3x^4}{2} - \frac{x^3}{6} + 3x^2 + 1 \right)$$

4.22 problem 22

4.22.1 Maple step by step solution 1748

Internal problem ID [7243]

Internal file name [OUTPUT/6229_Sunday_June_05_2022_04_33_16_PM_45710257/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 22.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + (3x^2 + 2x)y' - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + (3x^2 + 2x)y' - 2y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3x + 2}{x}$$
$$q(x) = -\frac{2}{x^2}$$

Table 174: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3x+2}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{2}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + (3x^2 + 2x) y' - 2y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + (3x^2 + 2x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{1+n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 3x^{1+n+r} a_n (n+r) = \sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) x^{n+r} \right) \\ & + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + 2x^{n+r} a_n (n+r) - 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r (-1+r) + 2x^r a_0 r - 2a_0 x^r = 0$$

Or

$$(x^r r (-1+r) + 2x^r r - 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 + r - 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + r - 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= -2 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 + r - 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 3$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= x \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x^2} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{1+n} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-2} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 3a_{n-1}(n+r-1) + 2a_n(n+r) - 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{3a_{n-1}}{n+r+2} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = -\frac{3a_{n-1}}{n+3} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{3}{3+r}$$

Which for the root $r = 1$ becomes

$$a_1 = -\frac{3}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3+r}$	$-\frac{3}{4}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{9}{(3+r)(4+r)}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{9}{20}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3+r}$	$-\frac{3}{4}$
a_2	$\frac{9}{(3+r)(4+r)}$	$\frac{9}{20}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{27}{(3+r)(4+r)(5+r)}$$

Which for the root $r = 1$ becomes

$$a_3 = -\frac{9}{40}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3+r}$	$-\frac{3}{4}$
a_2	$\frac{9}{(3+r)(4+r)}$	$\frac{9}{20}$
a_3	$-\frac{27}{(3+r)(4+r)(5+r)}$	$-\frac{9}{40}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{81}{(4+r)(5+r)(3+r)(6+r)}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{27}{280}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3+r}$	$-\frac{3}{4}$
a_2	$\frac{9}{(3+r)(4+r)}$	$\frac{9}{20}$
a_3	$-\frac{27}{(3+r)(4+r)(5+r)}$	$-\frac{9}{40}$
a_4	$\frac{81}{(4+r)(5+r)(3+r)(6+r)}$	$\frac{27}{280}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{243}{(5+r)(3+r)(6+r)(7+r)(4+r)}$$

Which for the root $r = 1$ becomes

$$a_5 = -\frac{81}{2240}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3+r}$	$-\frac{3}{4}$
a_2	$\frac{9}{(3+r)(4+r)}$	$\frac{9}{20}$
a_3	$-\frac{27}{(3+r)(4+r)(5+r)}$	$-\frac{9}{40}$
a_4	$\frac{81}{(4+r)(5+r)(3+r)(6+r)}$	$\frac{27}{280}$
a_5	$-\frac{243}{(5+r)(3+r)(6+r)(7+r)(4+r)}$	$-\frac{81}{2240}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 - \frac{3x}{4} + \frac{9x^2}{20} - \frac{9x^3}{40} + \frac{27x^4}{280} - \frac{81x^5}{2240} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 3$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_3(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_3 \\ &= -\frac{27}{(3+r)(4+r)(5+r)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} -\frac{27}{(3+r)(4+r)(5+r)} &= \lim_{r \rightarrow -2} -\frac{27}{(3+r)(4+r)(5+r)} \\ &= -\frac{9}{2} \end{aligned}$$

The limit is $-\frac{9}{2}$. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-2} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + 3b_{n-1}(n+r-1) + 2b_n(n+r) - 2b_n = 0 \quad (4)$$

Which for for the root $r = -2$ becomes

$$b_n(n-2)(n-3) + 3b_{n-1}(n-3) + 2b_n(n-2) - 2b_n = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{3b_{n-1}}{n+r+2} \quad (5)$$

Which for the root $r = -2$ becomes

$$b_n = -\frac{3b_{n-1}}{n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -2$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -\frac{3}{3+r}$$

Which for the root $r = -2$ becomes

$$b_1 = -3$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3+r}$	-3

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{9}{(3+r)(4+r)}$$

Which for the root $r = -2$ becomes

$$b_2 = \frac{9}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3+r}$	-3
b_2	$\frac{9}{(3+r)(4+r)}$	$\frac{9}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{27}{(3+r)(4+r)(5+r)}$$

Which for the root $r = -2$ becomes

$$b_3 = -\frac{9}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3+r}$	-3
b_2	$\frac{9}{(3+r)(4+r)}$	$\frac{9}{2}$
b_3	$-\frac{27}{(3+r)(4+r)(5+r)}$	$-\frac{9}{2}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{81}{(4+r)(5+r)(3+r)(6+r)}$$

Which for the root $r = -2$ becomes

$$b_4 = \frac{27}{8}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3+r}$	-3
b_2	$\frac{9}{(3+r)(4+r)}$	$\frac{9}{2}$
b_3	$-\frac{27}{(3+r)(4+r)(5+r)}$	$-\frac{9}{2}$
b_4	$\frac{81}{(4+r)(5+r)(3+r)(6+r)}$	$\frac{27}{8}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{243}{(5+r)(3+r)(6+r)(7+r)(4+r)}$$

Which for the root $r = -2$ becomes

$$b_5 = -\frac{81}{40}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3+r}$	-3
b_2	$\frac{9}{(3+r)(4+r)}$	$\frac{9}{2}$
b_3	$-\frac{27}{(3+r)(4+r)(5+r)}$	$-\frac{9}{2}$
b_4	$\frac{81}{(4+r)(5+r)(3+r)(6+r)}$	$\frac{27}{8}$
b_5	$-\frac{243}{(5+r)(3+r)(6+r)(7+r)(4+r)}$	$-\frac{81}{40}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 - 3x + \frac{9x^2}{2} - \frac{9x^3}{2} + \frac{27x^4}{8} - \frac{81x^5}{40} + O(x^6)}{x^2} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x \left(1 - \frac{3x}{4} + \frac{9x^2}{20} - \frac{9x^3}{40} + \frac{27x^4}{280} - \frac{81x^5}{2240} + O(x^6) \right) \\ &\quad + \frac{c_2 \left(1 - 3x + \frac{9x^2}{2} - \frac{9x^3}{2} + \frac{27x^4}{8} - \frac{81x^5}{40} + O(x^6) \right)}{x^2} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x \left(1 - \frac{3x}{4} + \frac{9x^2}{20} - \frac{9x^3}{40} + \frac{27x^4}{280} - \frac{81x^5}{2240} + O(x^6) \right) \\ &\quad + \frac{c_2 \left(1 - 3x + \frac{9x^2}{2} - \frac{9x^3}{2} + \frac{27x^4}{8} - \frac{81x^5}{40} + O(x^6) \right)}{x^2} \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1x \left(1 - \frac{3x}{4} + \frac{9x^2}{20} - \frac{9x^3}{40} + \frac{27x^4}{280} - \frac{81x^5}{2240} + O(x^6) \right) \\ &\quad + \frac{c_2 \left(1 - 3x + \frac{9x^2}{2} - \frac{9x^3}{2} + \frac{27x^4}{8} - \frac{81x^5}{40} + O(x^6) \right)}{x^2} \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned} y &= c_1x \left(1 - \frac{3x}{4} + \frac{9x^2}{20} - \frac{9x^3}{40} + \frac{27x^4}{280} - \frac{81x^5}{2240} + O(x^6) \right) \\ &\quad + \frac{c_2 \left(1 - 3x + \frac{9x^2}{2} - \frac{9x^3}{2} + \frac{27x^4}{8} - \frac{81x^5}{40} + O(x^6) \right)}{x^2} \end{aligned}$$

Verified OK.

4.22.1 Maple step by step solution

Let's solve

$$x^2 y'' + (3x^2 + 2x)y' - 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{2y}{x^2} - \frac{(3x+2)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(3x+2)y'}{x} - \frac{2y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{3x+2}{x}, P_3(x) = -\frac{2}{x^2}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + x(3x + 2)y' - 2y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 1..2$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k- > k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m}(k+1-m+r)x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(2+r)(-1+r)x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r+2)(k+r-1) + 3a_{k-1}(k+r-1))x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(2+r)(-1+r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \{-2, 1\}$$
- Each term in the series must be 0, giving the recursion relation

$$(k+r-1)(a_k(k+r+2) + 3a_{k-1}) = 0$$
- Shift index using $k \rightarrow k+1$

$$(k+r)(a_{k+1}(k+3+r) + 3a_k) = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{3a_k}{k+3+r}$$
- Recursion relation for $r = -2$

$$a_{k+1} = -\frac{3a_k}{k+1}$$
- Solution for $r = -2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-2}, a_{k+1} = -\frac{3a_k}{k+1} \right]$$
- Recursion relation for $r = 1$

$$a_{k+1} = -\frac{3a_k}{k+4}$$
- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = -\frac{3a_k}{k+4} \right]$$
- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-2} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1} \right), a_{k+1} = -\frac{3a_k}{k+1}, b_{k+1} = -\frac{3b_k}{k+4} \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 45

```
Order:=6;
dsolve(x^2*diff(y(x), x, x) + (2*x+3*x^2)*diff(y(x),x)-2*y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = c_1 x \left(1 - \frac{3}{4}x + \frac{9}{20}x^2 - \frac{9}{40}x^3 + \frac{27}{280}x^4 - \frac{81}{2240}x^5 + O(x^6) \right) \\ + \frac{c_2 (12 - 36x + 54x^2 - 54x^3 + \frac{81}{2}x^4 - \frac{243}{10}x^5 + O(x^6))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 64

```
AsymptoticDSolveValue[x^2*y''[x]+(2*x+3*x^2)*y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{27x^2}{8} + \frac{1}{x^2} - \frac{9x}{2} - \frac{3}{x} + \frac{9}{2} \right) + c_2 \left(\frac{27x^5}{280} - \frac{9x^4}{40} + \frac{9x^3}{20} - \frac{3x^2}{4} + x \right)$$

4.23 problem 23

Internal problem ID [7244]

Internal file name [OUTPUT/6230_Sunday_June_05_2022_04_33_19_PM_19504862/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 23.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = x^2 + \cos(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 176: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	"regular"

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	"regular"

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : [0]

Irregular singular points : [∞]

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = x^2 + \cos(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^2 + \cos(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Expanding the rhs of the ode $x^2 + \cos(x)$ in series gives

$$x^2 + \cos(x) = 1 + \frac{1}{2}x^2 + \frac{1}{24}x^4$$

Since the $F = 1 + \frac{1}{2}x^2 + \frac{1}{24}x^4$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$c_0 = 1$$

$$m = 0$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+0} \end{aligned}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = 1$
$c_1 = 0$
$c_2 = \frac{1}{3}$
$c_3 = 0$
$c_4 = \frac{1}{63}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= 1 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\ &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = \frac{x^2}{2}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \frac{x^2}{2}$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= \frac{1}{6} \\ m &= 2 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+2} \end{aligned}$$

Where in the above $c_0 = \frac{1}{6}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{6}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{6}$
$c_1 = 0$
$c_2 = \frac{1}{126}$
$c_3 = 0$
$c_4 = \frac{1}{6930}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= x^2 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= x^2 \left(\frac{1}{6} + \frac{1}{126} x^2 + \frac{1}{6930} x^4 \right) \\
 &= \frac{1}{6} x^2 + \frac{1}{126} x^4 + \frac{1}{6930} x^6
 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = \frac{x^4}{24}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \frac{x^4}{24}$$

For c_0 and x . This results in

$$\begin{aligned}
 c_0 &= \frac{1}{504} \\
 m &= 4
 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
 y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
 &= \sum_{n=0}^{\infty} c_n x^{n+4}
 \end{aligned}$$

Where in the above $c_0 = \frac{1}{504}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 4$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{504}$ and $r = m$ or $r = 4$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{504}$
$c_1 = 0$
$c_2 = \frac{1}{27720}$
$c_3 = 0$
$c_4 = \frac{1}{2910600}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= x^4 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= x^4 \left(\frac{1}{504} + \frac{1}{27720} x^2 + \frac{1}{2910600} x^4 \right) \\
 &= \frac{1}{504} x^4 + \frac{1}{27720} x^6 + \frac{1}{2910600} x^8
 \end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = 1 + \frac{x^2}{2} + \frac{13x^4}{504} + \frac{x^6}{5544} + \frac{x^8}{2910600} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = 1 + \frac{x^2}{2} + \frac{13x^4}{504} + O(x^6)$$

Hence the final solution is

$$y = y_h + y_p \\ = 1 + \frac{x^2}{2} + \frac{13x^4}{504} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Summary

The solution(s) found are the following

$$y = 1 + \frac{x^2}{2} + \frac{13x^4}{504} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \quad (1)$$

Verification of solutions

$$y = 1 + \frac{x^2}{2} + \frac{13x^4}{504} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 43

Order:=6;

```
dsolve(2*x^2*diff(y(x), x, x) - x*diff(y(x), x) + (-x^2 + 1)*y(x) = x^2+cos(x), y(x), type='se
```

$$y(x) = c_1 \sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2 x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + \left(1 + \frac{1}{2}x^2 + \frac{13}{504}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.107 (sec). Leaf size: 176

```
AsymptoticDSolveValue[2*x^2*y'[x]-x*y'[x]+(1-x^2)*y[x]==x^2+Cos[x], y[x], {x, 0, 5}]
```

$$y(x) \rightarrow c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) + c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{59x^{11/2}}{77220} - \frac{17x^{7/2}}{630} - \frac{2x^{3/2}}{5} + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{239x^5}{138600} + \frac{11x^3}{252} + \frac{2x}{3} - \frac{1}{x} \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right)$$

4.24 problem 24

Internal problem ID [7245]

Internal file name [OUTPUT/6231_Sunday_June_05_2022_04_33_20_PM_65496285/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 24.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = \cos(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 177: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = \cos(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \cos(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Expanding the rhs of the ode $\cos(x)$ in series gives

$$\cos(x) = \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2$$

Since the $F = \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = \frac{x^4}{24}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \frac{x^4}{24}$$

For c_0 and x . This results in

$$c_0 = \frac{1}{504}$$

$$m = 4$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+4}$$

Where in the above $c_0 = \frac{1}{504}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 4$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{504}$ and $r = m$ or $r = 4$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{504}$
$c_1 = 0$
$c_2 = \frac{1}{27720}$
$c_3 = 0$
$c_4 = \frac{1}{2910600}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= x^4 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x^4 \left(\frac{1}{504} + \frac{1}{27720} x^2 + \frac{1}{2910600} x^4 \right) \\ &= \frac{1}{504} x^4 + \frac{1}{27720} x^6 + \frac{1}{2910600} x^8 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= 1 \\ m &= 0 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+0} \end{aligned}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$$\begin{array}{l}
 c_0 = 1 \\
 c_1 = 0 \\
 c_2 = \frac{1}{3} \\
 c_3 = 0 \\
 c_4 = \frac{1}{63} \\
 c_5 = 0
 \end{array}$$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= 1 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\
 &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4
 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{x^2}{2}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = -\frac{x^2}{2}$$

For c_0 and x . This results in

$$\begin{aligned}
 c_0 &= -\frac{1}{6} \\
 m &= 2
 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
 y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
 &= \sum_{n=0}^{\infty} c_n x^{n+2}
 \end{aligned}$$

Where in the above $c_0 = -\frac{1}{6}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{6}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = -\frac{1}{6}$
$c_1 = 0$
$c_2 = -\frac{1}{126}$
$c_3 = 0$
$c_4 = -\frac{1}{6930}$
$c_5 = 0$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^2 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$y_p = x^2 \left(-\frac{1}{6} - \frac{1}{126}x^2 - \frac{1}{6930}x^4 \right)$$

$$= -\frac{1}{6}x^2 - \frac{1}{126}x^4 - \frac{1}{6930}x^6$$

Adding all the above particular solution(s) gives

$$y_p = 1 + \frac{x^2}{6} + \frac{5x^4}{504} - \frac{x^6}{9240} + \frac{x^8}{2910600} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = 1 + \frac{x^2}{6} + \frac{5x^4}{504} + O(x^6)$$

Hence the final solution is

$$y = y_h + y_p \\ = 1 + \frac{x^2}{6} + \frac{5x^4}{504} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Summary

The solution(s) found are the following

$$y = 1 + \frac{x^2}{6} + \frac{5x^4}{504} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \quad (1)$$

Verification of solutions

$$y = 1 + \frac{x^2}{6} + \frac{5x^4}{504} + O(x^6) + c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 43

Order:=6;

```
dsolve(2*x^2*diff(y(x), x, x) - x*diff(y(x), x) + (-x^2 + 1)*y(x) = cos(x), y(x), type='series
```

$$y(x) = c_1 \sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2 x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + \left(1 + \frac{1}{6}x^2 + \frac{5}{504}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 176

```
AsymptoticDSolveValue[2*x^2*y'[x]-x*y'[x]+(1-x^2)*y[x]==Cos[x], y[x], {x, 0, 5}]
```

$y(x)$

$$\begin{aligned} &\rightarrow c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ &+ c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{3861} + \frac{x^{7/2}}{630} + \frac{4x^{3/2}}{15} \right. \\ &\left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{37x^5}{69300} - \frac{x^3}{84} - \frac{x}{3} - \frac{1}{x} \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \end{aligned}$$

4.25 problem 24

Internal problem ID [7246]

Internal file name [OUTPUT/6232_Sunday_June_05_2022_04_33_22_PM_21405477/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 24.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = x^3 + \cos(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 178: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = x^3 + \cos(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & 2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^3 + \cos(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Expanding the rhs of the ode $x^3 + \cos(x)$ in series gives

$$x^3 + \cos(x) = 1 - \frac{1}{2}x^2 + x^3 + \frac{1}{24}x^4$$

Since the $F = 1 - \frac{1}{2}x^2 + x^3 + \frac{1}{24}x^4$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = 1$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = 1$$

For c_0 and x . This results in

$$c_0 = 1$$

$$m = 0$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+0} \end{aligned}$$

Where in the above $c_0 = 1$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 0$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = 1$ and $r = m$ or $r = 0$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = 1$
$c_1 = 0$
$c_2 = \frac{1}{3}$
$c_3 = 0$
$c_4 = \frac{1}{63}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= 1 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= 1 \left(1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \right) \\ &= 1 + \frac{1}{3}x^2 + \frac{1}{63}x^4 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{x^2}{2}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = -\frac{x^2}{2}$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= -\frac{1}{6} \\ m &= 2 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+2} \end{aligned}$$

Where in the above $c_0 = -\frac{1}{6}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{6}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$$\begin{array}{l}
c_0 = -\frac{1}{6} \\
c_1 = 0 \\
c_2 = -\frac{1}{126} \\
c_3 = 0 \\
c_4 = -\frac{1}{6930} \\
c_5 = 0
\end{array}$$

The particular solution is now found using

$$\begin{aligned}
y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
&= x^2 \sum_{n=0}^{\infty} c_n x^n
\end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
y_p &= x^2 \left(-\frac{1}{6} - \frac{1}{126} x^2 - \frac{1}{6930} x^4 \right) \\
&= -\frac{1}{6} x^2 - \frac{1}{126} x^4 - \frac{1}{6930} x^6
\end{aligned}$$

Now we determine the particular solution y_p associated with $F = x^3$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^3$$

For c_0 and x . This results in

$$\begin{aligned}
c_0 &= \frac{1}{10} \\
m &= 3
\end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
&= \sum_{n=0}^{\infty} c_n x^{n+3}
\end{aligned}$$

Where in the above $c_0 = \frac{1}{10}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 3$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{10}$ and $r = m$ or $r = 3$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{10}$
$c_1 = 0$
$c_2 = \frac{1}{360}$
$c_3 = 0$
$c_4 = \frac{1}{28080}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= x^3 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= x^3 \left(\frac{1}{10} + \frac{1}{360} x^2 + \frac{1}{28080} x^4 \right) \\
 &= \frac{1}{10} x^3 + \frac{1}{360} x^5 + \frac{1}{28080} x^7
 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = \frac{x^4}{24}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \frac{x^4}{24}$$

For c_0 and x . This results in

$$\begin{aligned}
 c_0 &= \frac{1}{504} \\
 m &= 4
 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
 y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
 &= \sum_{n=0}^{\infty} c_n x^{n+4}
 \end{aligned}$$

Where in the above $c_0 = \frac{1}{504}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 4$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{504}$ and $r = m$ or $r = 4$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{504}$
$c_1 = 0$
$c_2 = \frac{1}{27720}$
$c_3 = 0$
$c_4 = \frac{1}{2910600}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= x^4 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= x^4 \left(\frac{1}{504} + \frac{1}{27720} x^2 + \frac{1}{2910600} x^4 \right) \\
 &= \frac{1}{504} x^4 + \frac{1}{27720} x^6 + \frac{1}{2910600} x^8
 \end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = 1 + \frac{x^2}{6} + \frac{x^3}{10} + \frac{5x^4}{504} + \frac{x^5}{360} - \frac{x^6}{9240} + \frac{x^7}{28080} + \frac{x^8}{2910600} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = 1 + \frac{x^2}{6} + \frac{x^3}{10} + \frac{5x^4}{504} + \frac{x^5}{360} + O(x^6)$$

Hence the final solution is

$$\begin{aligned} y &= y_h + y_p \\ &= 1 + \frac{x^2}{6} + \frac{x^3}{10} + \frac{5x^4}{504} + \frac{x^5}{360} + O(x^6) \\ &\quad + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= 1 + \frac{x^2}{6} + \frac{x^3}{10} + \frac{5x^4}{504} + \frac{x^5}{360} + O(x^6) \\ &\quad + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned} y &= 1 + \frac{x^2}{6} + \frac{x^3}{10} + \frac{5x^4}{504} + \frac{x^5}{360} + O(x^6) \\ &\quad + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
Order:=6;
```

```
dsolve(2*x^2*diff(y(x), x, x) - x*diff(y(x), x) + (-x^2 + 1)*y(x) = x^3+cos(x), y(x), type='se
```

$$y(x) = c_1 \sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2 x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) \\ + \left(1 + \frac{1}{6}x^2 + \frac{1}{10}x^3 + \frac{5}{504}x^4 + \frac{1}{360}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 176

```
AsymptoticDSolveValue[2*x^2*y'[x]-x*y'[x]+(1-x^2)*y[x]==Cos[x],y[x],{x,0,5}]
```

$y(x)$

$$\begin{aligned} &\rightarrow c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \\ &+ c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{3861} + \frac{x^{7/2}}{630} + \frac{4x^{3/2}}{15} \right. \\ &\left. + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{37x^5}{69300} - \frac{x^3}{84} - \frac{x}{3} - \frac{1}{x} \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \end{aligned}$$

4.26 problem 24

Internal problem ID [7247]

Internal file name [OUTPUT/6233_Sunday_June_05_2022_04_33_24_PM_76326205/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 24.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = \cos(x)x^3$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 179: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = \cos(x)x^3$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \cos(x) x^3$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Expanding the rhs of the ode $\cos(x)x^3$ in series gives

$$\cos(x)x^3 = x^3 - \frac{1}{2}x^5$$

Since the $F = x^3 - \frac{1}{2}x^5$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x^3$ by solving the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = x^3$$

For c_0 and x . This results in

$$c_0 = \frac{1}{10}$$

$$m = 3$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+3}$$

Where in the above $c_0 = \frac{1}{10}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 3$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{10}$ and $r = m$ or $r = 3$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{10}$
$c_1 = 0$
$c_2 = \frac{1}{360}$
$c_3 = 0$
$c_4 = \frac{1}{28080}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= x^3 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x^3 \left(\frac{1}{10} + \frac{1}{360} x^2 + \frac{1}{28080} x^4 \right) \\ &= \frac{1}{10} x^3 + \frac{1}{360} x^5 + \frac{1}{28080} x^7 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{x^5}{2}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = -\frac{x^5}{2}$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= -\frac{1}{72} \\ m &= 5 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+5} \end{aligned}$$

Where in the above $c_0 = -\frac{1}{72}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 5$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{72}$ and $r = m$ or $r = 5$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = -\frac{1}{72}$ $c_1 = 0$ $c_2 = -\frac{1}{5616}$ $c_3 = 0$ $c_4 = -\frac{1}{763776}$ $c_5 = 0$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^5 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$y_p = x^5 \left(-\frac{1}{72} - \frac{1}{5616} x^2 - \frac{1}{763776} x^4 \right)$$

$$= -\frac{1}{72} x^5 - \frac{1}{5616} x^7 - \frac{1}{763776} x^9$$

Adding all the above particular solution(s) gives

$$y_p = \frac{x^3}{10} - \frac{x^5}{90} - \frac{x^7}{7020} - \frac{x^9}{763776} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{x^3}{10} - \frac{x^5}{90} + O(x^6)$$

Hence the final solution is

$$y = y_h + y_p$$

$$= \frac{x^3}{10} - \frac{x^5}{90} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Summary

The solution(s) found are the following

$$y = \frac{x^3}{10} - \frac{x^5}{90} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verification of solutions

$$y = \frac{x^3}{10} - \frac{x^5}{90} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 45

Order:=6;

dsolve(2*x^2*diff(y(x), x, x) - x*diff(y(x), x) + (-x^2 + 1)*y(x) = x^3*cos(x), y(x), type='se

$$y(x) = c_1\sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + x^3 \left(\frac{1}{10} - \frac{1}{90}x^2 + O(x^4) \right)$$

✓ Solution by Mathematica

Time used: 0.138 (sec). Leaf size: 215

AsymptoticDSolveValue[2*x^2*y'[x]-x*y'[x]+(1-x^2)*y[x]==x^3+Cos[x], y[x], {x, 0, 5}]

$$y(x) \rightarrow c_1\sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + c_2x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{3861} - \frac{x^{9/2}}{45} + \frac{x^{7/2}}{630} - \frac{2x^{5/2}}{5} + \frac{4x^{3/2}}{15} + \frac{2}{\sqrt{x}} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) \left(\frac{x^6}{1008} + \frac{37x^5}{69300} + \frac{x^4}{24} - \frac{x^3}{84} + \frac{x^2}{2} - \frac{x}{3} - \frac{1}{x} \right)$$

4.27 problem 24

Internal problem ID [7248]

Internal file name [OUTPUT/6234_Sunday_June_05_2022_04_33_27_PM_99187992/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 24.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = \cos(x)x^3 + \sin(x)^2$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' - xy' + (1 - x^2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2x}$$
$$q(x) = -\frac{x^2 - 1}{2x^2}$$

Table 180: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	"regular"

$q(x) = -\frac{x^2-1}{2x^2}$	
singularity	type
$x = 0$	"regular"

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' - xy' + (1 - x^2)y = \cos(x)x^3 + \sin(x)^2$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' - xy' + (1 - x^2)y = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1-x^2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) - x^r a_0 r + a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) - x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 3r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 3r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \frac{1}{2} \end{aligned}$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = \cos(x) x^3 + \sin(x)^2$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 3r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_n(n+r) + a_n - a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-2}}{2n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{360}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x\left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_n(n+r) + b_n - b_{n-2} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}}{2n^2 + 4nr + 2r^2 - 3n - 3r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{b_{n-2}}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{2r^2 + 5r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{4r^4 + 36r^3 + 113r^2 + 144r + 63}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{1}{168}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{1}{2r^2+5r+3}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{4r^4+36r^3+113r^2+144r+63}$	$\frac{1}{168}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)
 \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = 0$
$a_2 = \frac{a_0}{2r^2+5r+3}$
$a_3 = 0$
$a_4 = \frac{a_0}{(2r^2+5r+3)(2r^2+13r+21)}$
$a_5 = 0$

Expanding the rhs of the ode $\cos(x)x^3 + \sin(x)^2$ in series gives

$$\cos(x)x^3 + \sin(x)^2 = x^2 + x^3 - \frac{1}{3}x^4 - \frac{1}{2}x^5$$

Since the $F = x^2 + x^3 - \frac{1}{3}x^4 - \frac{1}{2}x^5$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x^2$ by solving the balance equation

$$(2x^m m(-1+m) - x^m m + x^m) c_0 = x^2$$

For c_0 and x . This results in

$$c_0 = \frac{1}{3}$$

$$m = 2$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+2}$$

Where in the above $c_0 = \frac{1}{3}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{3}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{3}$
$c_1 = 0$
$c_2 = \frac{1}{63}$
$c_3 = 0$
$c_4 = \frac{1}{3465}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= x^2 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x^2 \left(\frac{1}{3} + \frac{1}{63} x^2 + \frac{1}{3465} x^4 \right) \\ &= \frac{1}{3} x^2 + \frac{1}{63} x^4 + \frac{1}{3465} x^6 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = x^3$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = x^3$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= \frac{1}{10} \\ m &= 3 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+3} \end{aligned}$$

Where in the above $c_0 = \frac{1}{10}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 3$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{10}$ and $r = m$ or $r = 3$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{10}$
$c_1 = 0$
$c_2 = \frac{1}{360}$
$c_3 = 0$
$c_4 = \frac{1}{28080}$
$c_5 = 0$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^3 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$y_p = x^3 \left(\frac{1}{10} + \frac{1}{360} x^2 + \frac{1}{28080} x^4 \right)$$

$$= \frac{1}{10} x^3 + \frac{1}{360} x^5 + \frac{1}{28080} x^7$$

Now we determine the particular solution y_p associated with $F = -\frac{x^4}{3}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = -\frac{x^4}{3}$$

For c_0 and x . This results in

$$c_0 = -\frac{1}{63}$$

$$m = 4$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+4}$$

Where in the above $c_0 = -\frac{1}{63}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 4$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{63}$ and $r = m$ or $r = 4$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = -\frac{1}{63}$
$c_1 = 0$
$c_2 = -\frac{1}{3465}$
$c_3 = 0$
$c_4 = -\frac{1}{363825}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= x^4 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= x^4 \left(-\frac{1}{63} - \frac{1}{3465}x^2 - \frac{1}{363825}x^4 \right) \\
 &= -\frac{1}{63}x^4 - \frac{1}{3465}x^6 - \frac{1}{363825}x^8
 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{x^5}{2}$ by solving the balance equation

$$(2x^m m(-1 + m) - x^m m + x^m) c_0 = -\frac{x^5}{2}$$

For c_0 and x . This results in

$$\begin{aligned}
 c_0 &= -\frac{1}{72} \\
 m &= 5
 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+5} \end{aligned}$$

Where in the above $c_0 = -\frac{1}{72}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 5$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{72}$ and $r = m$ or $r = 5$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = -\frac{1}{72}$
$c_1 = 0$
$c_2 = -\frac{1}{5616}$
$c_3 = 0$
$c_4 = -\frac{1}{763776}$
$c_5 = 0$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= x^5 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x^5 \left(-\frac{1}{72} - \frac{1}{5616}x^2 - \frac{1}{763776}x^4 \right) \\ &= -\frac{1}{72}x^5 - \frac{1}{5616}x^7 - \frac{1}{763776}x^9 \end{aligned}$$

Adding all the above particular solution(s) gives

$$y_p = \frac{x^2}{3} + \frac{x^3}{10} - \frac{x^5}{90} - \frac{x^7}{7020} - \frac{x^8}{363825} - \frac{x^9}{763776} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{x^2}{3} + \frac{x^3}{10} - \frac{x^5}{90} + O(x^6)$$

Hence the final solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \frac{x^2}{3} + \frac{x^3}{10} - \frac{x^5}{90} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \frac{x^2}{3} + \frac{x^3}{10} - \frac{x^5}{90} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) \\ &\quad + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right) \end{aligned} \tag{1}$$

Verification of solutions

$$y = \frac{x^2}{3} + \frac{x^3}{10} - \frac{x^5}{90} + O(x^6) + c_1 x \left(1 + \frac{x^2}{10} + \frac{x^4}{360} + O(x^6) \right) + c_2 \sqrt{x} \left(1 + \frac{x^2}{6} + \frac{x^4}{168} + O(x^6) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
Order:=6;
```

```
dsolve(2*x^2*diff(y(x), x, x) - x*diff(y(x), x) + (-x^2 + 1)*y(x) = x^3*cos(x)+sin(x)^2,y(x))
```

$$y(x) = c_1 \sqrt{x} \left(1 + \frac{1}{6}x^2 + \frac{1}{168}x^4 + O(x^6) \right) + c_2 x \left(1 + \frac{1}{10}x^2 + \frac{1}{360}x^4 + O(x^6) \right) + x^2 \left(\frac{1}{3} + \frac{1}{10}x - \frac{1}{90}x^3 + O(x^4) \right)$$

✓ Solution by Mathematica

Time used: 0.514 (sec). Leaf size: 199

```
AsymptoticDSolveValue[2*x^2*y'[x]-x*y'[x]+(1-x^2)*y[x]==x^3*Cos[x]+Sin[x]^2,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 x \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right) + c_1 \sqrt{x} \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + \sqrt{x} \left(-\frac{x^{11/2}}{396} + \frac{4x^{9/2}}{45} + \frac{x^{7/2}}{15} - \frac{2x^{5/2}}{5} - \frac{2x^{3/2}}{3} \right) \left(\frac{x^6}{11088} + \frac{x^4}{168} + \frac{x^2}{6} + 1 \right) + x \left(-\frac{x^6}{168} - \frac{13x^5}{12600} - \frac{x^4}{12} - \frac{x^3}{18} + \frac{x^2}{2} + x \right) \left(\frac{x^6}{28080} + \frac{x^4}{360} + \frac{x^2}{10} + 1 \right)$$

4.28 problem 24

Internal problem ID [7249]

Internal file name [OUTPUT/6235_Sunday_June_05_2022_04_33_32_PM_64206728/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 24.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' - xy' + (1 - x^2)y = \ln(x)$$

With the expansion point for the power series method at $x = 1$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = x - 1$$

The ode is converted to be in terms of the new independent variable t . This results in

$$2(t+1)^2 \left(\frac{d^2}{dt^2} y(t) \right) - (t+1) \left(\frac{d}{dt} y(t) \right) + (1 - (t+1)^2) y(t) = \ln(t+1)$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \dots \\ &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \dots \\ &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{313}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{314}$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \tag{2}$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = \frac{y(t)t^2 + 2y(t)t + t\left(\frac{d}{dt}y(t)\right) + \frac{d}{dt}y(t) + \ln(t+1)}{2(t+1)^2}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dt} \\ &= \frac{\partial F_0}{\partial t} + \frac{\partial F_0}{\partial y} \frac{d}{dt}y(t) + \frac{\partial F_0}{\partial \frac{d}{dt}y(t)} F_0 \\ &= \frac{-3\ln(t+1) + (2t^3 + 6t^2 + 3t - 1)\left(\frac{d}{dt}y(t)\right) + 2 + (t^2 + 2t + 4)y(t)}{4(t+1)^3} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dt} \\ &= \frac{\partial F_1}{\partial t} + \frac{\partial F_1}{\partial y} \frac{d}{dt}y(t) + \frac{\partial F_1}{\partial \frac{d}{dt}y(t)} F_1 \\ &= \frac{(2t^2 + 4t + 17)\ln(t+1) + (4t^3 + 12t^2 + 27t + 19)\left(\frac{d}{dt}y(t)\right) - 18 + (2t^4 + 8t^3 + 5t^2 - 6t - 20)y(t)}{8(t+1)^4} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dt} \\ &= \frac{\partial F_2}{\partial t} + \frac{\partial F_2}{\partial y} \frac{d}{dt}y(t) + \frac{\partial F_2}{\partial \frac{d}{dt}y(t)} F_2 \\ &= \frac{(-4t^2 - 8t - 109)\ln(t+1) + (4t^5 + 20t^4 + 22t^3 - 14t^2 - 139t - 119)\left(\frac{d}{dt}y(t)\right) + (4t^4 + 16t^3 + 63t^2)}{16(t+1)^5} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dt} \\ &= \frac{\partial F_3}{\partial t} + \frac{\partial F_3}{\partial y} \frac{d}{dt}y(t) + \frac{\partial F_3}{\partial \frac{d}{dt}y(t)} F_3 \\ &= \frac{(4t^4 + 16t^3 + 30t^2 + 28t + 955)\ln(t+1) + (12t^5 + 60t^4 + 252t^3 + 516t^2 + 1401t + 1089)\left(\frac{d}{dt}y(t)\right) +}{32(t+1)^6} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $t = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= \frac{y'(0)}{2} \\
 F_1 &= y(0) - \frac{y'(0)}{4} + \frac{1}{2} \\
 F_2 &= -\frac{5y(0)}{2} + \frac{19y'(0)}{8} - \frac{9}{4} \\
 F_3 &= \frac{37y(0)}{4} - \frac{119y'(0)}{16} + \frac{89}{8} \\
 F_4 &= -\frac{323y(0)}{8} + \frac{1089y'(0)}{32} - \frac{991}{16}
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned}
 y(t) &= \left(1 + \frac{1}{6}t^3 - \frac{5}{48}t^4 + \frac{37}{480}t^5 - \frac{323}{5760}t^6\right) y(0) \\
 &+ \left(t + \frac{1}{4}t^2 - \frac{1}{24}t^3 + \frac{19}{192}t^4 - \frac{119}{1920}t^5 + \frac{121}{2560}t^6\right) y'(0) \\
 &+ \frac{t^3}{12} - \frac{3t^4}{32} + \frac{89t^5}{960} - \frac{991t^6}{11520} + O(t^6)
 \end{aligned}$$

Since the expansion point $t = 0$ is an ordinary, we can also solve this using standard power series. The ode is normalized to be

$$(2t^2 + 4t + 2) \left(\frac{d^2}{dt^2}y(t)\right) + (-t - 1) \left(\frac{d}{dt}y(t)\right) + (-t^2 - 2t)y(t) = \ln(t + 1)$$

Let the solution be represented as power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^n$$

Then

$$\begin{aligned}
 \frac{d}{dt}y(t) &= \sum_{n=1}^{\infty} n a_n t^{n-1} \\
 \frac{d^2}{dt^2}y(t) &= \sum_{n=2}^{\infty} n(n-1) a_n t^{n-2}
 \end{aligned}$$

Substituting the above back into the ode gives

$$(2t^2 + 4t + 2) \left(\sum_{n=2}^{\infty} n(n-1) a_n t^{n-2}\right) + (-t - 1) \left(\sum_{n=1}^{\infty} n a_n t^{n-1}\right) + (-t^2 - 2t) \left(\sum_{n=0}^{\infty} a_n t^n\right) = \ln(t + 1) \quad (1)$$

Expanding $\ln(t + 1)$ as Taylor series around $t = 0$ and keeping only the first 6 terms gives

$$\begin{aligned}\ln(t + 1) &= t - \frac{1}{2}t^2 + \frac{1}{3}t^3 - \frac{1}{4}t^4 + \frac{1}{5}t^5 + \dots \\ &= t - \frac{1}{2}t^2 + \frac{1}{3}t^3 - \frac{1}{4}t^4 + \frac{1}{5}t^5\end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\begin{aligned}(2t^2 + 4t + 2) \left(\sum_{n=2}^{\infty} n(n-1) a_n t^{n-2} \right) + (-t-1) \left(\sum_{n=1}^{\infty} n a_n t^{n-1} \right) \\ + (-t^2 - 2t) \left(\sum_{n=0}^{\infty} a_n t^n \right) = t - \frac{1}{2}t^2 + \frac{1}{3}t^3 - \frac{1}{4}t^4 + \frac{1}{5}t^5\end{aligned}$$

Which simplifies to

$$\begin{aligned}\left(\sum_{n=2}^{\infty} 2t^n a_n n(n-1) \right) + \left(\sum_{n=2}^{\infty} 4n t^{n-1} a_n (n-1) \right) + \left(\sum_{n=2}^{\infty} 2n(n-1) a_n t^{n-2} \right) \\ + \sum_{n=1}^{\infty} (-n a_n t^n) + \sum_{n=1}^{\infty} (-n a_n t^{n-1}) + \sum_{n=0}^{\infty} (-t^{n+2} a_n) \\ + \sum_{n=0}^{\infty} (-2t^{1+n} a_n) = t - \frac{1}{2}t^2 + \frac{1}{3}t^3 - \frac{1}{4}t^4 + \frac{1}{5}t^5\end{aligned} \quad (2)$$

The next step is to make all powers of t be n in each summation term. Going over each summation term above with power of t in it which is not already t^n and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=2}^{\infty} 4n t^{n-1} a_n (n-1) &= \sum_{n=1}^{\infty} 4(1+n) a_{1+n} n t^n \\ \sum_{n=2}^{\infty} 2n(n-1) a_n t^{n-2} &= \sum_{n=0}^{\infty} 2(n+2) a_{n+2} (1+n) t^n \\ \sum_{n=1}^{\infty} (-n a_n t^{n-1}) &= \sum_{n=0}^{\infty} (-(1+n) a_{1+n} t^n) \\ \sum_{n=0}^{\infty} (-t^{n+2} a_n) &= \sum_{n=2}^{\infty} (-a_{n-2} t^n)\end{aligned}$$

$$\sum_{n=0}^{\infty} (-2t^{1+n}a_n) = \sum_{n=1}^{\infty} (-2a_{n-1}t^n)$$

Substituting all the above in Eq (2) gives the following equation where now all powers of t are the same and equal to n .

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} 2t^n a_n n(n-1) \right) + \left(\sum_{n=1}^{\infty} 4(1+n) a_{1+n} n t^n \right) \\ & + \left(\sum_{n=0}^{\infty} 2(n+2) a_{n+2} (1+n) t^n \right) + \sum_{n=1}^{\infty} (-n a_n t^n) + \sum_{n=0}^{\infty} (-(1+n) a_{1+n} t^n) \\ & + \sum_{n=2}^{\infty} (-a_{n-2} t^n) + \sum_{n=1}^{\infty} (-2a_{n-1} t^n) = t - \frac{1}{2}t^2 + \frac{1}{3}t^3 - \frac{1}{4}t^4 + \frac{1}{5}t^5 \end{aligned} \quad (3)$$

$n = 0$ gives

$$4a_2 - a_1 = 0$$

$$a_2 = \frac{a_1}{4}$$

$n = 1$ gives

$$\begin{aligned} (6a_2 + 12a_3 - a_1 - 2a_0)t &= t \\ 6a_2 + 12a_3 - a_1 - 2a_0 &= 1 \end{aligned}$$

Which after substituting earlier equations, simplifies to

$$a_3 = \frac{a_0}{6} - \frac{a_1}{24} + \frac{1}{12}$$

For $2 \leq n$, the recurrence equation is

$$\begin{aligned} (2na_n(n-1) + 4(1+n) a_{1+n} n + 2(n+2) a_{n+2} (1+n) - n a_n \\ - (1+n) a_{1+n} - a_{n-2} - 2a_{n-1}) t^n = t - \frac{1}{2}t^2 + \frac{1}{3}t^3 - \frac{1}{4}t^4 + \frac{1}{5}t^5 \end{aligned} \quad (4)$$

For $n = 2$ the recurrence equation gives

$$(2a_2 + 21a_3 + 24a_4 - a_0 - 2a_1)t^2 = -\frac{t^2}{2}$$

$$2a_2 + 21a_3 + 24a_4 - a_0 - 2a_1 = -\frac{1}{2}$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{3}{32} + \frac{19a_1}{192} - \frac{5a_0}{48}$$

For $n = 3$ the recurrence equation gives

$$(9a_3 + 44a_4 + 40a_5 - a_1 - 2a_2)t^3 = \frac{t^3}{3}$$

$$9a_3 + 44a_4 + 40a_5 - a_1 - 2a_2 = \frac{1}{3}$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{89}{960} + \frac{37a_0}{480} - \frac{119a_1}{1920}$$

For $n = 4$ the recurrence equation gives

$$(20a_4 + 75a_5 + 60a_6 - a_2 - 2a_3)t^4 = -\frac{t^4}{4}$$

$$20a_4 + 75a_5 + 60a_6 - a_2 - 2a_3 = -\frac{1}{4}$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{991}{11520} + \frac{121a_1}{2560} - \frac{323a_0}{5760}$$

For $n = 5$ the recurrence equation gives

$$(35a_5 + 114a_6 + 84a_7 - a_3 - 2a_4)t^5 = \frac{t^5}{5}$$

$$35a_5 + 114a_6 + 84a_7 - a_3 - 2a_4 = \frac{1}{5}$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{4261}{53760} + \frac{167a_0}{3840} - \frac{11761a_1}{322560}$$

And so on. Therefore the solution is

$$\begin{aligned} y(t) &= \sum_{n=0}^{\infty} a_n t^n \\ &= a_3 t^3 + a_2 t^2 + a_1 t + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y(t) &= a_0 + a_1 t + \frac{a_1 t^2}{4} + \left(\frac{a_0}{6} - \frac{a_1}{24} + \frac{1}{12} \right) t^3 \\ &\quad + \left(-\frac{3}{32} + \frac{19a_1}{192} - \frac{5a_0}{48} \right) t^4 + \left(\frac{89}{960} + \frac{37a_0}{480} - \frac{119a_1}{1920} \right) t^5 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y(t) &= \left(1 + \frac{1}{6}t^3 - \frac{5}{48}t^4 + \frac{37}{480}t^5 \right) a_0 \\ &\quad + \left(t + \frac{1}{4}t^2 - \frac{1}{24}t^3 + \frac{19}{192}t^4 - \frac{119}{1920}t^5 \right) a_1 + \frac{t^3}{12} - \frac{3t^4}{32} + \frac{89t^5}{960} + O(t^6) \end{aligned} \quad (3)$$

At $t = 0$ the solution above becomes

$$\begin{aligned} y(t) &= \left(1 + \frac{1}{6}t^3 - \frac{5}{48}t^4 + \frac{37}{480}t^5 \right) c_1 + \left(t + \frac{1}{4}t^2 - \frac{1}{24}t^3 + \frac{19}{192}t^4 - \frac{119}{1920}t^5 \right) c_2 \\ &\quad + \frac{t^3}{12} - \frac{3t^4}{32} + \frac{89t^5}{960} + O(t^6) \end{aligned}$$

Replacing t in the above with the original independent variable x using $t = x - 1$ results in

$$\begin{aligned} y &= \left(1 + \frac{(x-1)^3}{6} - \frac{5(x-1)^4}{48} + \frac{37(x-1)^5}{480} - \frac{323(x-1)^6}{5760} \right) y(1) \\ &\quad + \left(x-1 + \frac{(x-1)^2}{4} - \frac{(x-1)^3}{24} + \frac{19(x-1)^4}{192} - \frac{119(x-1)^5}{1920} + \frac{121(x-1)^6}{2560} \right) y'(1) \\ &\quad + \frac{(x-1)^3}{12} - \frac{3(x-1)^4}{32} + \frac{89(x-1)^5}{960} - \frac{991(x-1)^6}{11520} + O((x-1)^6) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = & \left(1 + \frac{(x-1)^3}{6} - \frac{5(x-1)^4}{48} + \frac{37(x-1)^5}{480} - \frac{323(x-1)^6}{5760} \right) y(1) \\ & + \left(x-1 + \frac{(x-1)^2}{4} - \frac{(x-1)^3}{24} + \frac{19(x-1)^4}{192} - \frac{119(x-1)^5}{1920} + \frac{121(x-1)^6}{2560} \right) y'(1) \\ & + \frac{(x-1)^3}{12} - \frac{3(x-1)^4}{32} + \frac{89(x-1)^5}{960} - \frac{991(x-1)^6}{11520} + O((x-1)^6) \end{aligned}$$

Verification of solutions

$$\begin{aligned} y = & \left(1 + \frac{(x-1)^3}{6} - \frac{5(x-1)^4}{48} + \frac{37(x-1)^5}{480} - \frac{323(x-1)^6}{5760} \right) y(1) \\ & + \left(x-1 + \frac{(x-1)^2}{4} - \frac{(x-1)^3}{24} + \frac{19(x-1)^4}{192} - \frac{119(x-1)^5}{1920} + \frac{121(x-1)^6}{2560} \right) y'(1) \\ & + \frac{(x-1)^3}{12} - \frac{3(x-1)^4}{32} + \frac{89(x-1)^5}{960} - \frac{991(x-1)^6}{11520} + O((x-1)^6) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 52

```
Order:=6;
```

```
dsolve(2*x^2*diff(y(x), x, x) - x*diff(y(x), x) + (-x^2 + 1)*y(x) = ln(x),y(x),type='series')
```

$$\begin{aligned} y(x) = & \left(1 + \frac{(x-1)^3}{6} - \frac{5(x-1)^4}{48} + \frac{37(x-1)^5}{480} \right) y(1) \\ & + \left(x-1 + \frac{(x-1)^2}{4} - \frac{(x-1)^3}{24} + \frac{19(x-1)^4}{192} - \frac{119(x-1)^5}{1920} \right) D(y)(1) \\ & + \frac{(x-1)^3}{12} - \frac{3(x-1)^4}{32} + \frac{89(x-1)^5}{960} + O(x^6) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 105

```
AsymptoticDSolveValue[2*x^2*y'[x]-x*y'[x]+(1-x^2)*y[x]==Log[x],y[x],{x,1,5}]
```

$$\begin{aligned} y(x) \rightarrow & \frac{89}{960}(x-1)^5 - \frac{3}{32}(x-1)^4 + \frac{1}{12}(x-1)^3 \\ & + c_1 \left(\frac{37}{480}(x-1)^5 - \frac{5}{48}(x-1)^4 + \frac{1}{6}(x-1)^3 + 1 \right) \\ & + c_2 \left(-\frac{119(x-1)^5}{1920} + \frac{19}{192}(x-1)^4 - \frac{1}{24}(x-1)^3 + \frac{1}{4}(x-1)^2 + x - 1 \right) \end{aligned}$$

4.29 problem 25

4.29.1 Maple step by step solution 1851

Internal problem ID [7250]

Internal file name [OUTPUT/6236_Sunday_June_05_2022_04_33_36_PM_43509992/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 25.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2x^2(x^2 + x + 1)y'' + x(11x^2 + 11x + 9)y' + (7x^2 + 10x + 6)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(2x^4 + 2x^3 + 2x^2)y'' + (11x^3 + 11x^2 + 9x)y' + (7x^2 + 10x + 6)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{11x^2 + 11x + 9}{2x(x^2 + x + 1)}$$
$$q(x) = \frac{7x^2 + 10x + 6}{2x^2(x^2 + x + 1)}$$

Table 181: Table $p(x), q(x)$ singularities.

$p(x) = \frac{11x^2+11x+9}{2x(x^2+x+1)}$		$q(x) = \frac{7x^2+10x+6}{2x^2(x^2+x+1)}$	
singularity	type	singularity	type
$x = 0$	“regular”	$x = 0$	“regular”
$x = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$	“regular”	$x = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$	“regular”
$x = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$	“regular”	$x = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $\left[0, -\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2}, \infty\right]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2(x^2 + x + 1)y'' + (11x^3 + 11x^2 + 9x)y' + (7x^2 + 10x + 6)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2(x^2 + x + 1) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + (11x^3 + 11x^2 + 9x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (7x^2 + 10x + 6) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned}
& \left(\sum_{n=0}^{\infty} 2x^{n+r+2} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{1+n+r} a_n (n+r) (n+r-1) \right) \\
& + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 11x^{n+r+2} a_n (n+r) \right) \\
& + \left(\sum_{n=0}^{\infty} 11x^{1+n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 9x^{n+r} a_n (n+r) \right) \\
& + \left(\sum_{n=0}^{\infty} 7x^{n+r+2} a_n \right) + \left(\sum_{n=0}^{\infty} 10x^{1+n+r} a_n \right) + \left(\sum_{n=0}^{\infty} 6a_n x^{n+r} \right) = 0
\end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=0}^{\infty} 2x^{n+r+2} a_n (n+r) (n+r-1) &= \sum_{n=2}^{\infty} 2a_{n-2} (n+r-2) (n-3+r) x^{n+r} \\
\sum_{n=0}^{\infty} 2x^{1+n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} 2a_{n-1} (n+r-1) (n+r-2) x^{n+r} \\
\sum_{n=0}^{\infty} 11x^{n+r+2} a_n (n+r) &= \sum_{n=2}^{\infty} 11a_{n-2} (n+r-2) x^{n+r} \\
\sum_{n=0}^{\infty} 11x^{1+n+r} a_n (n+r) &= \sum_{n=1}^{\infty} 11a_{n-1} (n+r-1) x^{n+r} \\
\sum_{n=0}^{\infty} 7x^{n+r+2} a_n &= \sum_{n=2}^{\infty} 7a_{n-2} x^{n+r} \\
\sum_{n=0}^{\infty} 10x^{1+n+r} a_n &= \sum_{n=1}^{\infty} 10a_{n-1} x^{n+r}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers

of x are the same and equal to $n + r$.

$$\begin{aligned}
& \left(\sum_{n=2}^{\infty} 2a_{n-2}(n+r-2)(n-3+r)x^{n+r} \right) \\
& + \left(\sum_{n=1}^{\infty} 2a_{n-1}(n+r-1)(n+r-2)x^{n+r} \right) \\
& + \left(\sum_{n=0}^{\infty} 2x^{n+r}a_n(n+r)(n+r-1) \right) + \left(\sum_{n=2}^{\infty} 11a_{n-2}(n+r-2)x^{n+r} \right) \quad (2B) \\
& + \left(\sum_{n=1}^{\infty} 11a_{n-1}(n+r-1)x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 9x^{n+r}a_n(n+r) \right) \\
& + \left(\sum_{n=2}^{\infty} 7a_{n-2}x^{n+r} \right) + \left(\sum_{n=1}^{\infty} 10a_{n-1}x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 6a_nx^{n+r} \right) = 0
\end{aligned}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2x^{n+r}a_n(n+r)(n+r-1) + 9x^{n+r}a_n(n+r) + 6a_nx^{n+r} = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r(-1+r) + 9x^r a_0 r + 6a_0 x^r = 0$$

Or

$$(2x^r r(-1+r) + 9x^r r + 6x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 + 7r + 6) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 + 7r + 6 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned}
r_1 &= -\frac{3}{2} \\
r_2 &= -2
\end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 + 7r + 6) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n-\frac{3}{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n-2}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{-r - 2}{r + 3}$$

For $2 \leq n$ the recursive equation is

$$2a_{n-2}(n+r-2)(n-3+r) + 2a_{n-1}(n+r-1)(n+r-2) + 2a_n(n+r)(n+r-1) + 11a_{n-2}(n+r-2) + 11a_{n-1}(n+r-1) + 9a_n(n+r) + 7a_{n-2} + 10a_{n-1} + 6a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{na_{n-2} + na_{n-1} + ra_{n-2} + ra_{n-1} - a_{n-2} + a_{n-1}}{n+r+2} \quad (4)$$

Which for the root $r = -\frac{3}{2}$ becomes

$$a_n = \frac{(-2a_{n-2} - 2a_{n-1})n + 5a_{n-2} + a_{n-1}}{2n+1} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = -\frac{3}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r-2}{r+3}$	$-\frac{1}{3}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4+r}$$

Which for the root $r = -\frac{3}{2}$ becomes

$$a_2 = \frac{2}{5}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r-2}{r+3}$	$-\frac{1}{3}$
a_2	$\frac{1}{4+r}$	$\frac{2}{5}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{r^2 + 3r + 1}{(r+3)(5+r)}$$

Which for the root $r = -\frac{3}{2}$ becomes

$$a_3 = -\frac{5}{21}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r-2}{r+3}$	$-\frac{1}{3}$
a_2	$\frac{1}{4+r}$	$\frac{2}{5}$
a_3	$\frac{r^2+3r+1}{(r+3)(5+r)}$	$-\frac{5}{21}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{-r^3 - 8r^2 - 19r - 13}{(6+r)(r+3)(4+r)}$$

Which for the root $r = -\frac{3}{2}$ becomes

$$a_4 = \frac{7}{135}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r-2}{r+3}$	$-\frac{1}{3}$
a_2	$\frac{1}{4+r}$	$\frac{2}{5}$
a_3	$\frac{r^2+3r+1}{(r+3)(5+r)}$	$-\frac{5}{21}$
a_4	$\frac{-r^3-8r^2-19r-13}{(6+r)(r+3)(4+r)}$	$\frac{7}{135}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{2r^3 + 18r^2 + 52r + 49}{(r+3)(4+r)(5+r)(7+r)}$$

Which for the root $r = -\frac{3}{2}$ becomes

$$a_5 = \frac{76}{1155}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r-2}{r+3}$	$-\frac{1}{3}$
a_2	$\frac{1}{4+r}$	$\frac{2}{5}$
a_3	$\frac{r^2+3r+1}{(r+3)(5+r)}$	$-\frac{5}{21}$
a_4	$\frac{-r^3-8r^2-19r-13}{(6+r)(r+3)(4+r)}$	$\frac{7}{135}$
a_5	$\frac{2r^3+18r^2+52r+49}{(r+3)(4+r)(5+r)(7+r)}$	$\frac{76}{1155}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \frac{1}{x^{\frac{3}{2}}} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= \frac{1 - \frac{x}{3} + \frac{2x^2}{5} - \frac{5x^3}{21} + \frac{7x^4}{135} + \frac{76x^5}{1155} + O(x^6)}{x^{\frac{3}{2}}} \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = \frac{-r-2}{r+3}$$

For $2 \leq n$ the recursive equation is

$$2b_{n-2}(n+r-2)(n-3+r) + 2b_{n-1}(n+r-1)(n+r-2) + 2b_n(n+r)(n+r-1) + 11b_{n-2}(n+r-2) + 11b_{n-1}(n+r-1) + 9b_n(n+r) + 7b_{n-2} + 10b_{n-1} + 6b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{nb_{n-2} + nb_{n-1} + rb_{n-2} + rb_{n-1} - b_{n-2} + b_{n-1}}{n+r+2} \quad (4)$$

Which for the root $r = -2$ becomes

$$b_n = \frac{(-b_{n-2} - b_{n-1})n + 3b_{n-2} + b_{n-1}}{n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -2$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r-2}{r+3}$	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{4+r}$$

Which for the root $r = -2$ becomes

$$b_2 = \frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r-2}{r+3}$	0
b_2	$\frac{1}{4+r}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{r^2 + 3r + 1}{(r + 3)(5 + r)}$$

Which for the root $r = -2$ becomes

$$b_3 = -\frac{1}{3}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r-2}{r+3}$	0
b_2	$\frac{1}{4+r}$	$\frac{1}{2}$
b_3	$\frac{r^2+3r+1}{(r+3)(5+r)}$	$-\frac{1}{3}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{-r^3 - 8r^2 - 19r - 13}{(6 + r)(r + 3)(4 + r)}$$

Which for the root $r = -2$ becomes

$$b_4 = \frac{1}{8}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r-2}{r+3}$	0
b_2	$\frac{1}{4+r}$	$\frac{1}{2}$
b_3	$\frac{r^2+3r+1}{(r+3)(5+r)}$	$-\frac{1}{3}$
b_4	$\frac{-r^3-8r^2-19r-13}{(6+r)(r+3)(4+r)}$	$\frac{1}{8}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{2r^3 + 18r^2 + 52r + 49}{(r + 3)(4 + r)(5 + r)(7 + r)}$$

Which for the root $r = -2$ becomes

$$b_5 = \frac{1}{30}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r-2}{r+3}$	0
b_2	$\frac{1}{4+r}$	$\frac{1}{2}$
b_3	$\frac{r^2+3r+1}{(r+3)(5+r)}$	$-\frac{1}{3}$
b_4	$\frac{-r^3-8r^2-19r-13}{(6+r)(r+3)(4+r)}$	$\frac{1}{8}$
b_5	$\frac{2r^3+18r^2+52r+49}{(r+3)(4+r)(5+r)(7+r)}$	$\frac{1}{30}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= \frac{1}{x^{\frac{3}{2}}} (b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 + \frac{x^2}{2} - \frac{x^3}{3} + \frac{x^4}{8} + \frac{x^5}{30} + O(x^6)}{x^2} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= \frac{c_1 \left(1 - \frac{x}{3} + \frac{2x^2}{5} - \frac{5x^3}{21} + \frac{7x^4}{135} + \frac{76x^5}{1155} + O(x^6) \right)}{x^{\frac{3}{2}}} + \frac{c_2 \left(1 + \frac{x^2}{2} - \frac{x^3}{3} + \frac{x^4}{8} + \frac{x^5}{30} + O(x^6) \right)}{x^2} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= \frac{c_1 \left(1 - \frac{x}{3} + \frac{2x^2}{5} - \frac{5x^3}{21} + \frac{7x^4}{135} + \frac{76x^5}{1155} + O(x^6) \right)}{x^{\frac{3}{2}}} + \frac{c_2 \left(1 + \frac{x^2}{2} - \frac{x^3}{3} + \frac{x^4}{8} + \frac{x^5}{30} + O(x^6) \right)}{x^2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \left(1 - \frac{x}{3} + \frac{2x^2}{5} - \frac{5x^3}{21} + \frac{7x^4}{135} + \frac{76x^5}{1155} + O(x^6) \right)}{x^{\frac{3}{2}}} + \frac{c_2 \left(1 + \frac{x^2}{2} - \frac{x^3}{3} + \frac{x^4}{8} + \frac{x^5}{30} + O(x^6) \right)}{x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \left(1 - \frac{x}{3} + \frac{2x^2}{5} - \frac{5x^3}{21} + \frac{7x^4}{135} + \frac{76x^5}{1155} + O(x^6) \right)}{x^{\frac{3}{2}}} + \frac{c_2 \left(1 + \frac{x^2}{2} - \frac{x^3}{3} + \frac{x^4}{8} + \frac{x^5}{30} + O(x^6) \right)}{x^2}$$

Verified OK.

4.29.1 Maple step by step solution

Let's solve

$$2x^2(x^2 + x + 1)y'' + (11x^3 + 11x^2 + 9x)y' + (7x^2 + 10x + 6)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(7x^2+10x+6)y}{2x^2(x^2+x+1)} - \frac{(11x^2+11x+9)y'}{2x(x^2+x+1)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(11x^2+11x+9)y'}{2x(x^2+x+1)} + \frac{(7x^2+10x+6)y}{2x^2(x^2+x+1)} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{11x^2+11x+9}{2x(x^2+x+1)}, P_3(x) = \frac{7x^2+10x+6}{2x^2(x^2+x+1)} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{9}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 3$$

- $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$2x^2(x^2 + x + 1)y'' + x(11x^2 + 11x + 9)y' + (7x^2 + 10x + 6)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^m \cdot y'$ to series expansion for $m = 1..3$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^m \cdot y''$ to series expansion for $m = 2..4$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(2+r)(3+2r)x^r + (a_1(3+r)(5+2r) + a_0(5+2r)(2+r))x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r+2)(2k+r) + a_{k-1}(k+r+1)(k+r)) \right) x^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(2+r)(3+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ -2, -\frac{3}{2} \right\}$$

- Each term must be 0

$$a_1(3+r)(5+2r) + a_0(5+2r)(2+r) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = -\frac{(2+r)a_0}{3+r}$$

- Each term in the series must be 0, giving the recursion relation

$$2((a_k + a_{k-2} + a_{k-1})k + (a_k + a_{k-2} + a_{k-1})r + 2a_k - a_{k-2} + a_{k-1})(k+r+\frac{3}{2}) = 0$$

- Shift index using $k \rightarrow k+2$

$$2((a_{k+2} + a_k + a_{k+1})(k+2) + (a_{k+2} + a_k + a_{k+1})r + 2a_{k+2} - a_k + a_{k+1})(k+\frac{7}{2}+r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{ka_k + ka_{k+1} + ra_k + ra_{k+1} + a_k + 3a_{k+1}}{k+4+r}$$

- Recursion relation for $r = -2$

$$a_{k+2} = -\frac{ka_k + ka_{k+1} - a_k + a_{k+1}}{k+2}$$

- Solution for $r = -2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-2}, a_{k+2} = -\frac{ka_k + ka_{k+1} - a_k + a_{k+1}}{k+2}, a_1 = 0 \right]$$

- Recursion relation for $r = -\frac{3}{2}$

$$a_{k+2} = -\frac{ka_k + ka_{k+1} - \frac{1}{2}a_k + \frac{3}{2}a_{k+1}}{k+\frac{5}{2}}$$

- Solution for $r = -\frac{3}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{3}{2}}, a_{k+2} = -\frac{ka_k + ka_{k+1} - \frac{1}{2}a_k + \frac{3}{2}a_{k+1}}{k+\frac{5}{2}}, a_1 = -\frac{a_0}{3} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-2} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-\frac{3}{2}} \right), a_{k+2} = -\frac{ka_k + ka_{k+1} - a_k + a_{k+1}}{k+2}, a_1 = 0, b_{k+2} = -\frac{kb_k + kb_{k+1} - \frac{1}{2}b_k + \frac{3}{2}b_{k+1}}{k+\frac{5}{2}} \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
  A Liouvillian solution exists
  Reducible group (found an exponential solution)
  Group is reducible, not completely reducible
  Solution has integrals. Trying a special function solution free of integrals...
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
  -> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
  <- Heun successful: received ODE is equivalent to the HeunG ODE, case a <> 0, e <>
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 45

```
Order:=6;
dsolve(2*x^2*(1+x+x^2)*diff(y(x), x$2) + x*(9+11*x+11*x^2)*diff(y(x), x) + (6+10*x+7*x^2)*y(x), x)
```

$$y(x) = \frac{c_1 \left(1 + \frac{1}{2}x^2 - \frac{1}{3}x^3 + \frac{1}{8}x^4 + \frac{1}{30}x^5 + O(x^6)\right)}{x^2} + \frac{c_2 \left(1 - \frac{1}{3}x + \frac{2}{5}x^2 - \frac{5}{21}x^3 + \frac{7}{135}x^4 + \frac{76}{1155}x^5 + O(x^6)\right)}{x^{\frac{3}{2}}}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 83

```
AsymptoticDSolveValue[2*x^2*(1+x+x^2)*y'[x] + x*(9+11*x+11*x^2)*y'[x] + (6+10*x+7*x^2)*y[x]
```

$$y(x) \rightarrow \frac{c_2 \left(\frac{x^5}{30} + \frac{x^4}{8} - \frac{x^3}{3} + \frac{x^2}{2} + 1 \right)}{x^2} + \frac{c_1 \left(\frac{76x^5}{1155} + \frac{7x^4}{135} - \frac{5x^3}{21} + \frac{2x^2}{5} - \frac{x}{3} + 1 \right)}{x^{3/2}}$$

4.30 problem 26

4.30.1 Maple step by step solution 1866

Internal problem ID [7251]

Internal file name [OUTPUT/6237_Sunday_June_05_2022_04_33_42_PM_25973204/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 26.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$x^2(x+3)y'' + 5x(1+x)y' - (1-4x)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x^3 + 3x^2)y'' + (5x^2 + 5x)y' + (4x - 1)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{5x + 5}{x(x + 3)}$$
$$q(x) = \frac{4x - 1}{x^2(x + 3)}$$

Table 183: Table $p(x), q(x)$ singularities.

$p(x) = \frac{5x+5}{x(x+3)}$		$q(x) = \frac{4x-1}{x^2(x+3)}$	
singularity	type	singularity	type
$x = -3$	“regular”	$x = -3$	“regular”
$x = 0$	“regular”	$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-3, 0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2(x+3)y'' + (5x^2 + 5x)y' + (4x - 1)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2(x+3) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + (5x^2 + 5x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (4x - 1) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned}
 & \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) (n+r-1) \right) \\
 & + \left(\sum_{n=0}^{\infty} 5x^{1+n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) \\
 & + \left(\sum_{n=0}^{\infty} 4x^{1+n+r} a_n \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0
 \end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}
 \sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r} \\
 \sum_{n=0}^{\infty} 5x^{1+n+r} a_n (n+r) &= \sum_{n=1}^{\infty} 5a_{n-1} (n+r-1) x^{n+r} \\
 \sum_{n=0}^{\infty} 4x^{1+n+r} a_n &= \sum_{n=1}^{\infty} 4a_{n-1} x^{n+r}
 \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned}
 & \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r} \right) \\
 & + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} 5a_{n-1} (n+r-1) x^{n+r} \right) \\
 & + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) + \left(\sum_{n=1}^{\infty} 4a_{n-1} x^{n+r} \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0
 \end{aligned} \tag{2B}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$3x^{n+r} a_n (n+r) (n+r-1) + 5x^{n+r} a_n (n+r) - a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$3x^r a_0 r(-1 + r) + 5x^r a_0 r - a_0 x^r = 0$$

Or

$$(3x^r r(-1 + r) + 5x^r r - x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(3r^2 + 2r - 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$3r^2 + 2r - 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{3}$$

$$r_2 = -1$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(3r^2 + 2r - 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{4}{3}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{3}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n-1}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_{n-1}(n+r-1)(n+r-2) + 3a_n(n+r)(n+r-1) + 5a_{n-1}(n+r-1) + 5a_n(n+r) + 4a_{n-1} - a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{(1+n+r)a_{n-1}}{3n+3r-1} \quad (4)$$

Which for the root $r = \frac{1}{3}$ becomes

$$a_n = -\frac{(4+3n)a_{n-1}}{9n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{3}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-2-r}{2+3r}$$

Which for the root $r = \frac{1}{3}$ becomes

$$a_1 = -\frac{7}{9}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{2+3r}$	$-\frac{7}{9}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r^2 + 5r + 6}{9r^2 + 21r + 10}$$

Which for the root $r = \frac{1}{3}$ becomes

$$a_2 = \frac{35}{81}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{2+3r}$	$-\frac{7}{9}$
a_2	$\frac{r^2+5r+6}{9r^2+21r+10}$	$\frac{35}{81}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-r^3 - 9r^2 - 26r - 24}{27r^3 + 135r^2 + 198r + 80}$$

Which for the root $r = \frac{1}{3}$ becomes

$$a_3 = -\frac{455}{2187}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{2+3r}$	$-\frac{7}{9}$
a_2	$\frac{r^2+5r+6}{9r^2+21r+10}$	$\frac{35}{81}$
a_3	$\frac{-r^3-9r^2-26r-24}{27r^3+135r^2+198r+80}$	$-\frac{455}{2187}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r^4 + 14r^3 + 71r^2 + 154r + 120}{81r^4 + 702r^3 + 2079r^2 + 2418r + 880}$$

Which for the root $r = \frac{1}{3}$ becomes

$$a_4 = \frac{1820}{19683}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{2+3r}$	$-\frac{7}{9}$
a_2	$\frac{r^2+5r+6}{9r^2+21r+10}$	$\frac{35}{81}$
a_3	$\frac{-r^3-9r^2-26r-24}{27r^3+135r^2+198r+80}$	$-\frac{455}{2187}$
a_4	$\frac{r^4+14r^3+71r^2+154r+120}{81r^4+702r^3+2079r^2+2418r+880}$	$\frac{1820}{19683}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-r^5 - 20r^4 - 155r^3 - 580r^2 - 1044r - 720}{243r^5 + 3240r^4 + 16065r^3 + 36360r^2 + 36492r + 12320}$$

Which for the root $r = \frac{1}{3}$ becomes

$$a_5 = -\frac{6916}{177147}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{2+3r}$	$-\frac{7}{9}$
a_2	$\frac{r^2+5r+6}{9r^2+21r+10}$	$\frac{35}{81}$
a_3	$\frac{-r^3-9r^2-26r-24}{27r^3+135r^2+198r+80}$	$-\frac{455}{2187}$
a_4	$\frac{r^4+14r^3+71r^2+154r+120}{81r^4+702r^3+2079r^2+2418r+880}$	$\frac{1820}{19683}$
a_5	$\frac{-r^5-20r^4-155r^3-580r^2-1044r-720}{243r^5+3240r^4+16065r^3+36360r^2+36492r+12320}$	$-\frac{6916}{177147}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^{\frac{1}{3}}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^{\frac{1}{3}}\left(1 - \frac{7x}{9} + \frac{35x^2}{81} - \frac{455x^3}{2187} + \frac{1820x^4}{19683} - \frac{6916x^5}{177147} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$\begin{aligned} b_{n-1}(n+r-1)(n+r-2) + 3b_n(n+r)(n+r-1) \\ + 5b_{n-1}(n+r-1) + 5b_n(n+r) + 4b_{n-1} - b_n = 0 \end{aligned} \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{(1+n+r)b_{n-1}}{3n+3r-1} \quad (4)$$

Which for the root $r = -1$ becomes

$$b_n = -\frac{nb_{n-1}}{3n-4} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{-2 - r}{2 + 3r}$$

Which for the root $r = -1$ becomes

$$b_1 = 1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-2-r}{2+3r}$	1

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{r^2 + 5r + 6}{9r^2 + 21r + 10}$$

Which for the root $r = -1$ becomes

$$b_2 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-2-r}{2+3r}$	1
b_2	$\frac{r^2+5r+6}{9r^2+21r+10}$	-1

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{-r^3 - 9r^2 - 26r - 24}{27r^3 + 135r^2 + 198r + 80}$$

Which for the root $r = -1$ becomes

$$b_3 = \frac{3}{5}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-2-r}{2+3r}$	1
b_2	$\frac{r^2+5r+6}{9r^2+21r+10}$	-1
b_3	$\frac{-r^3-9r^2-26r-24}{27r^3+135r^2+198r+80}$	$\frac{3}{5}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{r^4 + 14r^3 + 71r^2 + 154r + 120}{81r^4 + 702r^3 + 2079r^2 + 2418r + 880}$$

Which for the root $r = -1$ becomes

$$b_4 = -\frac{3}{10}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-2-r}{2+3r}$	1
b_2	$\frac{r^2+5r+6}{9r^2+21r+10}$	-1
b_3	$\frac{-r^3-9r^2-26r-24}{27r^3+135r^2+198r+80}$	$\frac{3}{5}$
b_4	$\frac{r^4+14r^3+71r^2+154r+120}{81r^4+702r^3+2079r^2+2418r+880}$	$-\frac{3}{10}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{-r^5 - 20r^4 - 155r^3 - 580r^2 - 1044r - 720}{243r^5 + 3240r^4 + 16065r^3 + 36360r^2 + 36492r + 12320}$$

Which for the root $r = -1$ becomes

$$b_5 = \frac{3}{22}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-2-r}{2+3r}$	1
b_2	$\frac{r^2+5r+6}{9r^2+21r+10}$	-1
b_3	$\frac{-r^3-9r^2-26r-24}{27r^3+135r^2+198r+80}$	$\frac{3}{5}$
b_4	$\frac{r^4+14r^3+71r^2+154r+120}{81r^4+702r^3+2079r^2+2418r+880}$	$-\frac{3}{10}$
b_5	$\frac{-r^5-20r^4-155r^3-580r^2-1044r-720}{243r^5+3240r^4+16065r^3+36360r^2+36492r+12320}$	$\frac{3}{22}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x^{\frac{1}{3}}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \frac{1 + x - x^2 + \frac{3x^3}{5} - \frac{3x^4}{10} + \frac{3x^5}{22} + O(x^6)}{x}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x^{\frac{1}{3}}\left(1 - \frac{7x}{9} + \frac{35x^2}{81} - \frac{455x^3}{2187} + \frac{1820x^4}{19683} - \frac{6916x^5}{177147} + O(x^6)\right) \\
 &\quad + \frac{c_2\left(1 + x - x^2 + \frac{3x^3}{5} - \frac{3x^4}{10} + \frac{3x^5}{22} + O(x^6)\right)}{x}
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1x^{\frac{1}{3}}\left(1 - \frac{7x}{9} + \frac{35x^2}{81} - \frac{455x^3}{2187} + \frac{1820x^4}{19683} - \frac{6916x^5}{177147} + O(x^6)\right) \\
 &\quad + \frac{c_2\left(1 + x - x^2 + \frac{3x^3}{5} - \frac{3x^4}{10} + \frac{3x^5}{22} + O(x^6)\right)}{x}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{\frac{1}{3}} \left(1 - \frac{7x}{9} + \frac{35x^2}{81} - \frac{455x^3}{2187} + \frac{1820x^4}{19683} - \frac{6916x^5}{177147} + O(x^6) \right) + \frac{c_2 \left(1 + x - x^2 + \frac{3x^3}{5} - \frac{3x^4}{10} + \frac{3x^5}{22} + O(x^6) \right)}{x} \quad (1)$$

Verification of solutions

$$y = c_1 x^{\frac{1}{3}} \left(1 - \frac{7x}{9} + \frac{35x^2}{81} - \frac{455x^3}{2187} + \frac{1820x^4}{19683} - \frac{6916x^5}{177147} + O(x^6) \right) + \frac{c_2 \left(1 + x - x^2 + \frac{3x^3}{5} - \frac{3x^4}{10} + \frac{3x^5}{22} + O(x^6) \right)}{x}$$

Verified OK.

4.30.1 Maple step by step solution

Let's solve

$$x^2(x+3)y'' + (5x^2 + 5x)y' + (4x-1)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(4x-1)y}{x^2(x+3)} - \frac{5(1+x)y'}{x(x+3)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{5(1+x)y'}{x(x+3)} + \frac{(4x-1)y}{x^2(x+3)} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{5(1+x)}{x(x+3)}, P_3(x) = \frac{4x-1}{x^2(x+3)} \right]$$

- $(x+3) \cdot P_2(x)$ is analytic at $x = -3$

$$\left. ((x+3) \cdot P_2(x)) \right|_{x=-3} = \frac{10}{3}$$

- $(x+3)^2 \cdot P_3(x)$ is analytic at $x = -3$

$$\left. ((x+3)^2 \cdot P_3(x)) \right|_{x=-3} = 0$$

- $x = -3$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -3$$

- Multiply by denominators

$$x^2(x+3)y'' + 5x(1+x)y' + (4x-1)y = 0$$

- Change variables using $x = u - 3$ so that the regular singular point is at $u = 0$

$$(u^3 - 6u^2 + 9u) \left(\frac{d^2}{du^2} y(u) \right) + (5u^2 - 25u + 30) \left(\frac{d}{du} y(u) \right) + (4u - 13) y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot y(u)$ to series expansion for $m = 0..1$

$$u^m \cdot y(u) = \sum_{k=0}^{\infty} a_k u^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$u^m \cdot y(u) = \sum_{k=m}^{\infty} a_{k-m} u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..2$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..3$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$3a_0 r(7+3r) u^{-1+r} + (3a_1(1+r)(10+3r) - a_0(13+6r)(1+r)) u^r + \left(\sum_{k=1}^{\infty} (3a_{k+1}(k+r+1) \right) (3$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$3r(7 + 3r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{0, -\frac{7}{3}\right\}$$

- Each term must be 0

$$3a_1(1 + r)(10 + 3r) - a_0(13 + 6r)(1 + r) = 0$$

- Each term in the series must be 0, giving the recursion relation

$$-6(k + r + 1) \left(\left(a_k - \frac{a_{k-1}}{6} - \frac{3a_{k+1}}{2} \right) k + \left(a_k - \frac{a_{k-1}}{6} - \frac{3a_{k+1}}{2} \right) r + \frac{13a_k}{6} - \frac{a_{k-1}}{6} - 5a_{k+1} \right) = 0$$

- Shift index using $k \rightarrow k + 1$

$$-6(k + r + 2) \left(\left(a_{k+1} - \frac{a_k}{6} - \frac{3a_{k+2}}{2} \right) (k + 1) + \left(a_{k+1} - \frac{a_k}{6} - \frac{3a_{k+2}}{2} \right) r + \frac{13a_{k+1}}{6} - \frac{a_k}{6} - 5a_{k+2} \right) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{ka_k - 6ka_{k+1} + ra_k - 6ra_{k+1} + 2a_k - 19a_{k+1}}{3(3k + 13 + 3r)}$$

- Recursion relation for $r = 0$

$$a_{k+2} = -\frac{ka_k - 6ka_{k+1} + 2a_k - 19a_{k+1}}{3(3k + 13)}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+2} = -\frac{ka_k - 6ka_{k+1} + 2a_k - 19a_{k+1}}{3(3k + 13)}, 30a_1 - 13a_0 = 0 \right]$$

- Revert the change of variables $u = x + 3$

$$\left[y = \sum_{k=0}^{\infty} a_k (x + 3)^k, a_{k+2} = -\frac{ka_k - 6ka_{k+1} + 2a_k - 19a_{k+1}}{3(3k + 13)}, 30a_1 - 13a_0 = 0 \right]$$

- Recursion relation for $r = -\frac{7}{3}$

$$a_{k+2} = -\frac{ka_k - 6ka_{k+1} - \frac{1}{3}a_k - 5a_{k+1}}{3(3k + 6)}$$

- Solution for $r = -\frac{7}{3}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k - \frac{7}{3}}, a_{k+2} = -\frac{ka_k - 6ka_{k+1} - \frac{1}{3}a_k - 5a_{k+1}}{3(3k + 6)}, -12a_1 - \frac{4a_0}{3} = 0 \right]$$

- Revert the change of variables $u = x + 3$

$$\left[y = \sum_{k=0}^{\infty} a_k (x + 3)^{k - \frac{7}{3}}, a_{k+2} = -\frac{ka_k - 6ka_{k+1} - \frac{1}{3}a_k - 5a_{k+1}}{3(3k + 6)}, -12a_1 - \frac{4a_0}{3} = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (x+3)^k \right) + \left(\sum_{k=0}^{\infty} b_k (x+3)^{k-\frac{7}{3}} \right), a_{k+2} = -\frac{ka_k - 6ka_{k+1} + 2a_k - 19a_{k+1}}{3(3k+13)}, 30a_1 - 13a_0 = 0, \dots \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
  One independent solution has integrals. Trying a hypergeometric solution free of integral
  -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  <- hyper3 successful: received ODE is equivalent to the 2F1 ODE
  -> Trying to convert hypergeometric functions to elementary form...
  <- elementary form for at least one hypergeometric solution is achieved - returning with
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 47

```

Order:=6;
dsolve(x^2*(3+x)*diff(y(x), x$2) + 5*x*(1+x)*diff(y(x), x) - (1-4*x)*y(x) = 0, y(x), type='series')

```

$$y(x) = \frac{c_2 x^{\frac{4}{3}} \left(1 - \frac{7}{9}x + \frac{35}{81}x^2 - \frac{455}{2187}x^3 + \frac{1820}{19683}x^4 - \frac{6916}{177147}x^5 + O(x^6) \right) + c_1 \left(1 + x - x^2 + \frac{3}{5}x^3 - \frac{3}{10}x^4 + \frac{3}{22}x^5 + O(x^6) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 82

```

AsymptoticDSolveValue[x^2*(3+x)*y'[x] + 5*x*(1+x)*y'[x] - (1-4*x)*y[x] == 0, y[x], {x, 0, 5}]

```

$$y(x) \rightarrow c_1 \sqrt[3]{x} \left(-\frac{6916x^5}{177147} + \frac{1820x^4}{19683} - \frac{455x^3}{2187} + \frac{35x^2}{81} - \frac{7x}{9} + 1 \right) + \frac{c_2 \left(\frac{3x^5}{22} - \frac{3x^4}{10} + \frac{3x^3}{5} - x^2 + x + 1 \right)}{x}$$

4.31 problem 27

4.31.1 Maple step by step solution 1879

Internal problem ID [7252]

Internal file name [OUTPUT/6238_Sunday_June_05_2022_04_33_46_PM_98768854/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 27.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2(-x^2 + 2)y'' - x(4x^2 + 3)y' + (-2x^2 + 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(-x^4 + 2x^2)y'' + (-4x^3 - 3x)y' + (-2x^2 + 2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{4x^2 + 3}{x(x^2 - 2)}$$
$$q(x) = \frac{2x^2 - 2}{x^2(x^2 - 2)}$$

Table 185: Table $p(x), q(x)$ singularities.

$p(x) = \frac{4x^2+3}{x(x^2-2)}$	
singularity	type
$x = 0$	“regular”
$x = \sqrt{2}$	“regular”
$x = -\sqrt{2}$	“regular”

$q(x) = \frac{2x^2-2}{x^2(x^2-2)}$	
singularity	type
$x = 0$	“regular”
$x = \sqrt{2}$	“regular”
$x = -\sqrt{2}$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \sqrt{2}, -\sqrt{2}, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$-y''x^2(x^2 - 2) + (-4x^3 - 3x)y' + (-2x^2 + 2)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & - \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x^2(x^2 - 2) \\
 & + (-4x^3 - 3x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (-2x^2 + 2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned}
& \sum_{n=0}^{\infty} (-x^{n+r+2} a_n (n+r) (n+r-1)) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) \\
& + \sum_{n=0}^{\infty} (-4x^{n+r+2} a_n (n+r)) + \sum_{n=0}^{\infty} (-3x^{n+r} a_n (n+r)) \\
& + \sum_{n=0}^{\infty} (-2x^{n+r+2} a_n) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0
\end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=0}^{\infty} (-x^{n+r+2} a_n (n+r) (n+r-1)) &= \sum_{n=2}^{\infty} (-a_{n-2} (n+r-2) (n-3+r) x^{n+r}) \\
\sum_{n=0}^{\infty} (-4x^{n+r+2} a_n (n+r)) &= \sum_{n=2}^{\infty} (-4a_{n-2} (n+r-2) x^{n+r}) \\
\sum_{n=0}^{\infty} (-2x^{n+r+2} a_n) &= \sum_{n=2}^{\infty} (-2a_{n-2} x^{n+r})
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned}
& \sum_{n=2}^{\infty} (-a_{n-2} (n+r-2) (n-3+r) x^{n+r}) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) \\
& + \sum_{n=2}^{\infty} (-4a_{n-2} (n+r-2) x^{n+r}) + \sum_{n=0}^{\infty} (-3x^{n+r} a_n (n+r)) \\
& + \sum_{n=2}^{\infty} (-2a_{n-2} x^{n+r}) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0
\end{aligned} \tag{2B}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) - 3x^{n+r} a_n (n+r) + 2a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r(-1 + r) - 3x^r a_0 r + 2a_0 x^r = 0$$

Or

$$(2x^r r(-1 + r) - 3x^r r + 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - 5r + 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - 5r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2$$

$$r_2 = \frac{1}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - 5r + 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{3}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+2}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$\begin{aligned} -a_{n-2}(n+r-2)(n-3+r) + 2a_n(n+r)(n+r-1) \\ - 4a_{n-2}(n+r-2) - 3a_n(n+r) - 2a_{n-2} + 2a_n = 0 \end{aligned} \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}(n^2 + 2nr + r^2 - n - r)}{2n^2 + 4nr + 2r^2 - 5n - 5r + 2} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = \frac{a_{n-2}(n^2 + 3n + 2)}{2n^2 + 3n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r^2 + 3r + 2}{2r^2 + 3r}$$

Which for the root $r = 2$ becomes

$$a_2 = \frac{6}{7}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{r^2+3r+2}{2r^2+3r}$	$\frac{6}{7}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{r^2+3r+2}{2r^2+3r}$	$\frac{6}{7}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r^3 + 8r^2 + 19r + 12}{4r^3 + 20r^2 + 21r}$$

Which for the root $r = 2$ becomes

$$a_4 = \frac{45}{77}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{r^2+3r+2}{2r^2+3r}$	$\frac{6}{7}$
a_3	0	0
a_4	$\frac{r^3+8r^2+19r+12}{4r^3+20r^2+21r}$	$\frac{45}{77}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{r^2+3r+2}{2r^2+3r}$	$\frac{6}{7}$
a_3	0	0
a_4	$\frac{r^3+8r^2+19r+12}{4r^3+20r^2+21r}$	$\frac{45}{77}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^2\left(1 + \frac{6x^2}{7} + \frac{45x^4}{77} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$\begin{aligned} -b_{n-2}(n+r-2)(n-3+r) + 2b_n(n+r)(n+r-1) \\ - 4b_{n-2}(n+r-2) - 3b_n(n+r) - 2b_{n-2} + 2b_n = 0 \end{aligned} \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}(n^2 + 2nr + r^2 - n - r)}{2n^2 + 4nr + 2r^2 - 5n - 5r + 2} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_n = \frac{4n^2b_{n-2} - b_{n-2}}{8n^2 - 12n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{r^2 + 3r + 2}{2r^2 + 3r}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_2 = \frac{15}{8}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{r^2+3r+2}{2r^2+3r}$	$\frac{15}{8}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{r^2+3r+2}{2r^2+3r}$	$\frac{15}{8}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{r^3 + 8r^2 + 19r + 12}{4r^3 + 20r^2 + 21r}$$

Which for the root $r = \frac{1}{2}$ becomes

$$b_4 = \frac{189}{128}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{r^2+3r+2}{2r^2+3r}$	$\frac{15}{8}$
b_3	0	0
b_4	$\frac{r^3+8r^2+19r+12}{4r^3+20r^2+21r}$	$\frac{189}{128}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{r^2+3r+2}{2r^2+3r}$	$\frac{15}{8}$
b_3	0	0
b_4	$\frac{r^3+8r^2+19r+12}{4r^3+20r^2+21r}$	$\frac{189}{128}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^2(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \sqrt{x} \left(1 + \frac{15x^2}{8} + \frac{189x^4}{128} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^2 \left(1 + \frac{6x^2}{7} + \frac{45x^4}{77} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{15x^2}{8} + \frac{189x^4}{128} + O(x^6) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x^2 \left(1 + \frac{6x^2}{7} + \frac{45x^4}{77} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{15x^2}{8} + \frac{189x^4}{128} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1x^2 \left(1 + \frac{6x^2}{7} + \frac{45x^4}{77} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{15x^2}{8} + \frac{189x^4}{128} + O(x^6) \right) \quad (1)$$

Verification of solutions

$$y = c_1x^2 \left(1 + \frac{6x^2}{7} + \frac{45x^4}{77} + O(x^6) \right) + c_2\sqrt{x} \left(1 + \frac{15x^2}{8} + \frac{189x^4}{128} + O(x^6) \right)$$

Verified OK.

4.31.1 Maple step by step solution

Let's solve

$$-y''x^2(x^2 - 2) + (-4x^3 - 3x)y' + (-2x^2 + 2)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2(x^2-1)y}{x^2(x^2-2)} - \frac{(4x^2+3)y'}{x(x^2-2)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(4x^2+3)y'}{x(x^2-2)} + \frac{2(x^2-1)y}{x^2(x^2-2)} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{4x^2+3}{x(x^2-2)}, P_3(x) = \frac{2(x^2-1)}{x^2(x^2-2)} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -\frac{3}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 1$$

- $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x^2(x^2 - 2) + x(4x^2 + 3)y' + (2x^2 - 2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^m \cdot y'$ to series expansion for $m = 1..3$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^m \cdot y''$ to series expansion for $m = 2..4$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) x^{k+r}$$

Rewrite ODE with series expansions

$$-a_0(-1+2r)(-2+r)x^r - a_1(1+2r)(-1+r)x^{1+r} + \left(\sum_{k=2}^{\infty} (-a_k(2k+2r-1)(k+r-2) + a_{k-1}(k+r-1)(k+r-2)) \right) x^{k+r}$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-(-1+2r)(-2+r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \left\{ 2, \frac{1}{2} \right\}$$
- Each term must be 0

$$-a_1(1+2r)(-1+r) = 0$$
- Solve for the dependent coefficient(s)

$$a_1 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$-2\left(k+r-\frac{1}{2}\right)(k+r-2)a_k + a_{k-2}(k+r)(k+r-1) = 0$$
- Shift index using $k \rightarrow k + 2$

$$-2\left(k+\frac{3}{2}+r\right)(k+r)a_{k+2} + a_k(k+r+2)(k+r+1) = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = \frac{a_k(k+r+2)(k+r+1)}{(2k+3+2r)(k+r)}$$

- Recursion relation for $r = 2$

$$a_{k+2} = \frac{a_k(k+4)(k+3)}{(2k+7)(k+2)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+2} = \frac{a_k(k+4)(k+3)}{(2k+7)(k+2)}, a_1 = 0 \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = \frac{a_k(k+\frac{5}{2})(k+\frac{3}{2})}{(2k+4)(k+\frac{1}{2})}$$

- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = \frac{a_k(k+\frac{5}{2})(k+\frac{3}{2})}{(2k+4)(k+\frac{1}{2})}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+2} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = \frac{a_k(k+4)(k+3)}{(2k+7)(k+2)}, a_1 = 0, b_{k+2} = \frac{b_k(k+\frac{5}{2})(k+\frac{3}{2})}{(2k+4)(k+\frac{1}{2})}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
  A Liouvillian solution exists
  Reducible group (found an exponential solution)
  Group is reducible, not completely reducible
  Solution has integrals. Trying a special function solution free of integrals...
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
      <- heuristic approach successful
    <- hypergeometric successful
  <- special function solution successful
    -> Trying to convert hypergeometric functions to elementary form...
      <- elementary form for at least one hypergeometric solution is achieved - returning wi
  <- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
Order:=6;
```

```
dsolve(x^2*(2-x^2)*diff(y(x), x$2) - x*(3+4*x^2)*diff(y(x), x) + (2-2*x^2)*y(x) = 0,y(x),typ
```

$$y(x) = c_1\sqrt{x} \left(1 + \frac{15}{8}x^2 + \frac{189}{128}x^4 + O(x^6) \right) + c_2x^2 \left(1 + \frac{6}{7}x^2 + \frac{45}{77}x^4 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 50

```
AsymptoticDSolveValue[x^2*(2-x^2)*y'[x] - x*(3+4*x^2)*y'[x] + (2-2*x^2)*y[x] == 0,y[x],{x,0
```

$$y(x) \rightarrow c_1 \left(\frac{45x^4}{77} + \frac{6x^2}{7} + 1 \right) x^2 + c_2 \left(\frac{189x^4}{128} + \frac{15x^2}{8} + 1 \right) \sqrt{x}$$

4.32 problem 28

Internal problem ID [7253]

Internal file name [OUTPUT/6239_Sunday_June_05_2022_04_33_49_PM_22094621/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 28.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : **"unknown"**

Maple gives the following as the ode type

[`y=_G(x,y')`]

Unable to solve or complete the solution.

$$y'^2 + y^2 = \sec(x)^4$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{1 - y^2 \cos(x)^4}}{\cos(x)^2} \quad (1)$$

$$y' = -\frac{\sqrt{1 - y^2 \cos(x)^4}}{\cos(x)^2} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Unable to determine ODE type.

Unable to determine ODE type.

Solving equation (2)

Unable to determine ODE type.

Unable to determine ODE type.

Maple trace

```
`Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
trying simple symmetries for implicit equations
Successful isolation of dy/dx: 2 solutions were found. Trying to solve each resulting ODE.
  *** Sublevel 2 ***
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying homogeneous types:
  trying exact
  Looking for potential symmetries
  trying an equivalence to an Abel ODE
  trying 1st order ODE linearizable_by_differentiation
-> Solving 1st order ODE of high degree, Lie methods, 1st trial
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 4
`, `-> Computing symmetries using: way = 2
`, `-> Computing symmetries using: way = 2
-> Solving 1st order ODE of high degree, 2nd attempt. Trying parametric methods
trying dAlembert
-> Calling odsolve with the ODE`,  $\text{diff}(y(x), x) = x^2/(2*(x^2+y(x)^2)^{(5/4)}*((x^2+y(x)^2)^{(5/4)}))$ 
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying homogeneous types:
  trying exact
  Looking for potential symmetries
  trying an equivalence to an Abel ODE
  trying 1st order ODE linearizable_by_differentiation
-> Calling odsolve with the ODE`,  $\text{diff}(y(x), x) = -x*(10+15*\cos(2*y(x))+\cos(6*y(x))+6*\cos(4*y(x)))$ 
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying homogeneous types:
  trying exact
  Looking for potential symmetries
  trying an equivalence to an Abel ODE
  trying 1st order ODE linearizable_by_differentiation
-> Solving 1st order ODE of high degree, Lie methods, 2nd trial
`, `-> Computing symmetries using: way = 4
`, `-> Computing symmetries using: way = 5
`, `-> Computing symmetries using: way = 5`
```

X Solution by Maple

```
dsolve(diff(y(x),x)^2+y(x)^2=sec(x)^4,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]^2+y[x]^2==Sec[x]^4,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.33 problem 29

4.33.1 Solving as dAlembert ode 1887

Internal problem ID [7254]

Internal file name [OUTPUT/6240_Sunday_June_05_2022_04_33_59_PM_32030808/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 29.

ODE order: 1.

ODE degree: 3.

The type(s) of ODE detected by this program : "**dAlembert**"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries] , _dAlembert]
```

$$(y - 2xy')^2 - y'^3 = 0$$

4.33.1 Solving as dAlembert ode

Let $p = y'$ the ode becomes

$$(-2xp + y)^2 - p^3 = 0$$

Solving for y from the above results in

$$y = 2xp + p^{\frac{3}{2}} \tag{1A}$$

$$y = 2xp - p^{\frac{3}{2}} \tag{2A}$$

This has the form

$$y = xf(p) + g(p) \tag{*}$$

Where f, g are functions of $p = y'(x)$. Each of the above ode's is dAlembert ode which is now solved. Solving ode 1A Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g')\frac{dp}{dx} \\ p - f &= (xf' + g')\frac{dp}{dx} \end{aligned} \tag{2}$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned}f &= 2p \\g &= p^{\frac{3}{2}}\end{aligned}$$

Hence (2) becomes

$$-p = \left(2x + \frac{3\sqrt{p}}{2}\right) p'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$-p = 0$$

Solving for p from the above gives

$$p = 0$$

Substituting these in (1A) gives

$$y = 0$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = -\frac{p(x)}{2x + \frac{3\sqrt{p(x)}}{2}} \quad (3)$$

This ODE is now solved for $p(x)$.

Inverting the above ode gives

$$\frac{d}{dp}x(p) = -\frac{2x(p) + \frac{3\sqrt{p}}{2}}{p} \quad (4)$$

This ODE is now solved for $x(p)$.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$\frac{d}{dp}x(p) + p(p)x(p) = q(p)$$

Where here

$$\begin{aligned}p(p) &= \frac{2}{p} \\q(p) &= -\frac{3}{2\sqrt{p}}\end{aligned}$$

Hence the ode is

$$\frac{d}{dp}x(p) + \frac{2x(p)}{p} = -\frac{3}{2\sqrt{p}}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{2}{p} dp} \\ &= p^2\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dp}(\mu x) &= (\mu) \left(-\frac{3}{2\sqrt{p}} \right) \\ \frac{d}{dp}(p^2 x) &= (p^2) \left(-\frac{3}{2\sqrt{p}} \right) \\ d(p^2 x) &= \left(-\frac{3p^{\frac{3}{2}}}{2} \right) dp\end{aligned}$$

Integrating gives

$$\begin{aligned}p^2 x &= \int -\frac{3p^{\frac{3}{2}}}{2} dp \\ p^2 x &= -\frac{3p^{\frac{5}{2}}}{5} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = p^2$ results in

$$x(p) = -\frac{3\sqrt{p}}{5} + \frac{c_1}{p^2}$$

Now we need to eliminate p between the above and (1A). One way to do this is by

solving (1) for p . This results in

$$p = \left(\frac{(108y - 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}}{6} + \frac{8x^2}{3(108y - 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}} - \frac{2x}{3} \right)^2$$

$$p = \left(-\frac{(108y - 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}}{12} - \frac{4x^2}{3(108y - 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}} - \frac{2x}{3} + \frac{i\sqrt{3}}{3} \right)^2$$

$$p = \left(-\frac{(108y - 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}}{12} - \frac{4x^2}{3(108y - 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}} - \frac{2x}{3} - \frac{i\sqrt{3}}{3} \right)^2$$

Substituting the above in the solution for x found above gives

x

$$432 \left((16\sqrt{3} \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} - 128x^6 + 160yx^3 - 27y^2) (108y - 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2})^{\frac{2}{3}} - 2048 \left(x^3 - \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16} \right) \right)$$

x

$$\left((82944 \left(i - \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} - 221184 \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) (i\sqrt{3} - 1) \right) (108y - 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2})^{\frac{2}{3}} - 2048 \left(x^3 - \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16} \right) \right)$$

x

$$\left((-82944 \left(i + \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} + 221184 (1 + i\sqrt{3}) \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) \right) (108y - 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2})^{\frac{2}{3}} - 2048 \left(x^3 - \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16} \right) \right)$$

Solving ode 2A Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g') \frac{dp}{dx} \\ p - f &= (xf' + g') \frac{dp}{dx} \end{aligned} \quad (2)$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned} f &= 2p \\ g &= -p^{\frac{3}{2}} \end{aligned}$$

Hence (2) becomes

$$-p = \left(2x - \frac{3\sqrt{p}}{2} \right) p'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$-p = 0$$

Solving for p from the above gives

$$p = 0$$

Substituting these in (1A) gives

$$y = 0$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = -\frac{p(x)}{2x - \frac{3\sqrt{p(x)}}{2}} \quad (3)$$

This ODE is now solved for $p(x)$.

Inverting the above ode gives

$$\frac{d}{dp} x(p) = -\frac{2x(p) - \frac{3\sqrt{p}}{2}}{p} \quad (4)$$

This ODE is now solved for $x(p)$.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$\frac{d}{dp} x(p) + p(p)x(p) = q(p)$$

Where here

$$p(p) = \frac{2}{p}$$
$$q(p) = \frac{3}{2\sqrt{p}}$$

Hence the ode is

$$\frac{d}{dp}x(p) + \frac{2x(p)}{p} = \frac{3}{2\sqrt{p}}$$

The integrating factor μ is

$$\mu = e^{\int \frac{2}{p} dp}$$
$$= p^2$$

The ode becomes

$$\frac{d}{dp}(\mu x) = (\mu) \left(\frac{3}{2\sqrt{p}} \right)$$
$$\frac{d}{dp}(p^2 x) = (p^2) \left(\frac{3}{2\sqrt{p}} \right)$$
$$d(p^2 x) = \left(\frac{3p^{\frac{3}{2}}}{2} \right) dp$$

Integrating gives

$$p^2 x = \int \frac{3p^{\frac{3}{2}}}{2} dp$$
$$p^2 x = \frac{3p^{\frac{5}{2}}}{5} + c_2$$

Dividing both sides by the integrating factor $\mu = p^2$ results in

$$x(p) = \frac{3\sqrt{p}}{5} + \frac{c_2}{p^2}$$

Now we need to eliminate p between the above and (1A). One way to do this is by

solving (1) for p . This results in

$$p = \left(\frac{(-108y + 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}}{6} + \frac{8x^2}{3(-108y + 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}} + \frac{2x}{3} \right)^2$$

$$p = \left(-\frac{(-108y + 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}}{12} - \frac{4x^2}{3(-108y + 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}} + \frac{2x}{3} + \frac{i\sqrt{3}}{3} \right)^2$$

$$p = \left(-\frac{(-108y + 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}}{12} - \frac{4x^2}{3(-108y + 64x^3 + 12\sqrt{-96yx^3 + 81y^2})^{\frac{1}{3}}} + \frac{2x}{3} - \frac{i\sqrt{3}}{3} \right)^2$$

Substituting the above in the solution for x found above gives

x

$$432 \left((16\sqrt{3} \left(x^3 - \frac{3y}{16}\right) \sqrt{-32yx^3 + 27y^2} + 128x^6 - 160yx^3 + 27y^2) (-108y + 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2})^{\frac{2}{3}} + 2048 \left(\frac{x^3 + \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16}}{\dots} \right) \right)$$

x

$$\left((82944 \left(i - \frac{\sqrt{3}}{3}\right) \left(x^3 - \frac{3y}{16}\right) \sqrt{-32yx^3 + 27y^2} + 221184 \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128}\right) (i\sqrt{3} - 1)) (-108y + 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2})^{\frac{2}{3}} + 2048 \left(\frac{x^3 + \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16}}{\dots} \right) \right)$$

x

$$\left((-82944 \left(i + \frac{\sqrt{3}}{3}\right) \left(x^3 - \frac{3y}{16}\right) \sqrt{-32yx^3 + 27y^2} - 221184 (1 + i\sqrt{3}) \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128}\right)) (-108y + 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2})^{\frac{2}{3}} + 2048 \left(\frac{x^3 + \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16}}{\dots} \right) \right)$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$

$$x \tag{2}$$

$$432 \left(\left(16\sqrt{3} \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} - 128x^6 + 160yx^3 - 27y^2 \right) \left(108y - 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2} \right)^{\frac{2}{3}} - 2048 \left(\frac{x^3 - \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16}}{\dots} \right) \right)$$

$$x \tag{3}$$

$$\left(\left(82944 \left(i - \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} - 221184 \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) \left(i\sqrt{3} - 1 \right) \right) \left(108y - 64x^3 + \dots \right) \right)$$

$$x \tag{4}$$

$$\left(\left(-82944 \left(i + \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} + 221184 \left(1 + i\sqrt{3} \right) \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) \right) \left(108y - 64x^3 + \dots \right) \right)$$

$$y = 0 \tag{5}$$

$$x \tag{6}$$

$$432 \left(\left(16\sqrt{3} \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} + 128x^6 - 160yx^3 + 27y^2 \right) \left(-108y + 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2} \right)^{\frac{2}{3}} + 2048 \left(\frac{x^3 + \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16}}{\dots} \right) \right)$$

$$x \tag{7}$$

$$\left(\left(82944 \left(i - \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} + 221184 \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) \left(i\sqrt{3} - 1 \right) \right) \left(-108y + 64x^3 + \dots \right) \right)$$

$$x \tag{8}$$

$$\left(\left(-82944 \left(i + \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} - 221184 \left(1 + i\sqrt{3} \right) \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) \right) \left(-108y + 64x^3 + \dots \right) \right)$$

Verification of solutions

$$y = 0$$

Verified OK.

x

$$432 \left(\left(16\sqrt{3} \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} - 128x^6 + 160yx^3 - 27y^2 \right) (108y - 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2})^{\frac{2}{3}} - 2048 \left(- \frac{\left(x^3 - \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16} \right)}{\dots} \right) \right)$$

Warning, solution could not be verified

x

$$\left(\left(82944 \left(i - \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} - 221184 \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) (i\sqrt{3} - 1) \right) (108y - 64x^3 + \dots) \right)$$

Warning, solution could not be verified

x

$$\left(\left(-82944 \left(i + \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} + 221184 (1 + i\sqrt{3}) \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) \right) (108y - 64x^3 + \dots) \right)$$

Warning, solution could not be verified

$$y = 0$$

Verified OK.

x

$$432 \left(\left(16\sqrt{3} \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} + 128x^6 - 160yx^3 + 27y^2 \right) (-108y + 64x^3 + 12\sqrt{3} \sqrt{-32yx^3 + 27y^2})^{\frac{2}{3}} + 2048 \left(\frac{\left(x^3 + \frac{3\sqrt{3} \sqrt{-32yx^3 + 27y^2}}{16} \right)}{\dots} \right) \right)$$

Warning, solution could not be verified

x

$$\left(\left(82944 \left(i - \frac{\sqrt{3}}{3} \right) \left(x^3 - \frac{3y}{16} \right) \sqrt{-32yx^3 + 27y^2} + 221184 \left(x^6 - \frac{5yx^3}{4} + \frac{27y^2}{128} \right) (i\sqrt{3} - 1) \right) (-108y + 64x^3 + \dots) \right)$$

Maple trace

```
`Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
trying simple symmetries for implicit equations
Successful isolation of dy/dx: 3 solutions were found. Trying to solve each resulting ODE.
  *** Sublevel 2 ***
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying homogeneous types:
  trying exact
  Looking for potential symmetries
  trying an equivalence to an Abel ODE
  trying 1st order ODE linearizable_by_differentiation
-> Solving 1st order ODE of high degree, Lie methods, 1st trial
`, `-> Computing symmetries using: way = 2
`, `-> Computing symmetries using: way = 2
-> Solving 1st order ODE of high degree, 2nd attempt. Trying parametric methods
trying dAlembert
<- dAlembert successful
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 73

```
dsolve((y(x)-2*x*diff(y(x),x))^2= diff(y(x),x)^3,y(x), singsol=all)
```

$$y(x) = 0$$
$$\left[x(-T) = \frac{3-T^{\frac{5}{2}} + 5c_1}{5-T^2}, y(-T) = \frac{-T^{\frac{5}{2}} + 10c_1}{5-T} \right]$$
$$\left[x(-T) = \frac{-3-T^{\frac{5}{2}} + 5c_1}{5-T^2}, y(-T) = \frac{-T^{\frac{5}{2}} + 10c_1}{5-T} \right]$$

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(y[x]-2*x*y'[x])^2== y'[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

4.34 problem 31

4.34.1 Maple step by step solution 1904

Internal problem ID [7255]

Internal file name [OUTPUT/6241_Sunday_June_05_2022_04_35_07_PM_49233843/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 31.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

`[[_Emden, _Fowler]]`

$$x^2y'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$

$$q(x) = \frac{1}{x^2}$$

Table 187: Table $p(x), q(x)$ singularities.

$p(x) = 0$	
singularity	type

$q(x) = \frac{1}{x^2}$	
singularity	type
$x = 0$	"regular"

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r (-1+r) + a_0 x^r = 0$$

Or

$$(x^r r (-1+r) + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - r + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$r_2 = \frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - r + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}+\frac{i\sqrt{3}}{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}-\frac{i\sqrt{3}}{2}}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $0 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = 0 \quad (4)$$

Which for the root $r = \frac{1}{2} + \frac{i\sqrt{3}}{2}$ becomes

$$a_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2} + \frac{i\sqrt{3}}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}y_1(x) &= x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} (1 + O(x^6))\end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$y_2(x) = x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} (1 + O(x^6))$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} (1 + O(x^6)) + c_2x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} (1 + O(x^6))\end{aligned}$$

Hence the final solution is

$$\begin{aligned}y &= y_h \\ &= c_1x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} (1 + O(x^6)) + c_2x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} (1 + O(x^6))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} (1 + O(x^6)) + c_2x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} (1 + O(x^6)) \quad (1)$$

Verification of solutions

$$y = c_1x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} (1 + O(x^6)) + c_2x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} (1 + O(x^6))$$

Verified OK.

4.34.1 Maple step by step solution

Let's solve

$$x^2 y'' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}\right) + y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) - \frac{d}{dt}y(t) + y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{1 \pm (\sqrt{-3})}{2}$$
- Roots of the characteristic polynomial

$$r = \left(\frac{1}{2} - \frac{I\sqrt{3}}{2}, \frac{1}{2} + \frac{I\sqrt{3}}{2} \right)$$
- 1st solution of the ODE

$$y_1(t) = e^{\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)$$
- 2nd solution of the ODE

$$y_2(t) = e^{\frac{t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)$$
- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$
- Substitute in solutions

$$y(t) = c_1 e^{\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) + c_2 e^{\frac{t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)$$
- Change variables back using $t = \ln(x)$

$$y = c_1 \sqrt{x} \cos\left(\frac{\sqrt{3} \ln(x)}{2}\right) + c_2 \sqrt{x} \sin\left(\frac{\sqrt{3} \ln(x)}{2}\right)$$
- Simplify

$$y = \sqrt{x} \left(c_1 \cos\left(\frac{\sqrt{3} \ln(x)}{2}\right) + c_2 \sin\left(\frac{\sqrt{3} \ln(x)}{2}\right) \right)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
Order:=6;  
dsolve(x^2*diff(y(x), x$2) +y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left(c_1 x^{-\frac{i\sqrt{3}}{2}} + c_2 x^{\frac{i\sqrt{3}}{2}} \right) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 26

```
AsymptoticDSolveValue[x^2*y''[x] +y[x] == 0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x^{-(-1)^{2/3}} + c_2 x^{\sqrt[3]{-1}}$$

4.35 problem 32

4.35.1 Maple step by step solution 1914

Internal problem ID [7256]

Internal file name [OUTPUT/6242_Sunday_June_05_2022_04_35_09_PM_47896713/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 32.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

`[[_Emden , _Fowler]]`

$$xy'' + y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$xy'' + y' - y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{1}{x}$$

Table 189: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$xy'' + y' - y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-a_n x^{n+r}) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r-1})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r-1}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^{-1+r} r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^{-1+r} r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = 1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(r+1)^2(2+r)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{36}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{576}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{14400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$
a_5	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{14400}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{1}{(r+1)^2}$	1	$-\frac{2}{(r+1)^3}$	-2
b_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$	$\frac{-6-4r}{(r+1)^3(2+r)^3}$	$-\frac{3}{4}$
b_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$	$\frac{-6r^2-24r-22}{(r+1)^3(2+r)^3(r+3)^3}$	$-\frac{11}{108}$
b_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$	$\frac{-8r^3-60r^2-140r-100}{(r+1)^3(2+r)^3(r+3)^3(r+4)^3}$	$-\frac{25}{3456}$
b_5	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{14400}$	$\frac{-10r^4-120r^3-510r^2-900r-548}{(r+1)^3(2+r)^3(r+3)^3(r+4)^3(r+5)^3}$	$-\frac{137}{432000}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) \\ &\quad - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\ &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\
 &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\
 &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\
 &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right)
 \end{aligned}$$

Verified OK.

4.35.1 Maple step by step solution

Let's solve

$$xy'' + y' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{x} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1}{x}, P_3(x) = -\frac{1}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$xy'' + y' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r^2 x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1}(k+1+r)^2 - a_k) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $r^2 = 0$
- Values of r that satisfy the indicial equation
 $r = 0$
- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1)^2 - a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k}{(k+1)^2}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k}{(k+1)^2}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = \frac{a_k}{(k+1)^2} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 59

```
Order:=6;  
dsolve(x*diff(y(x), x$2) +diff(y(x),x)-y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 + x + \frac{1}{4}x^2 + \frac{1}{36}x^3 + \frac{1}{576}x^4 + \frac{1}{14400}x^5 + O(x^6) \right) \\ + \left((-2)x - \frac{3}{4}x^2 - \frac{11}{108}x^3 - \frac{25}{3456}x^4 - \frac{137}{432000}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 107

```
AsymptoticDSolveValue[x*y'[x] +y'[x]-y[x] == 0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) + c_2 \left(-\frac{137x^5}{432000} - \frac{25x^4}{3456} - \frac{11x^3}{108} - \frac{3x^2}{4} \right. \\ \left. + \left(\frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) \log(x) - 2x \right)$$

4.36 problem 33

4.36.1 Maple step by step solution 1928

Internal problem ID [7257]

Internal file name [OUTPUT/6243_Sunday_June_05_2022_04_35_12_PM_864337/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 33.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$4xy'' + 2y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$4xy'' + 2y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{2x}$$
$$q(x) = \frac{1}{4x}$$

Table 191: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1}{4x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$4xy'' + 2y' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$4x \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 4x^{n+r-1} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2(n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} a_n x^{n+r} = \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} 4x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2(n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$4x^{n+r-1} a_n (n+r) (n+r-1) + 2(n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$4x^{-1+r} a_0 r (-1+r) + 2r a_0 x^{-1+r} = 0$$

Or

$$(4x^{-1+r} r (-1+r) + 2r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(4r^2 - 2r) x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation becomes

$$4r^2 - 2r = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(4r^2 - 2r) x^{-1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^n$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$4a_n(n+r)(n+r-1) + 2a_n(n+r) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{2(2n^2 + 4nr + 2r^2 - n - r)} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = -\frac{a_{n-1}}{4n^2 + 2n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{4r^2 + 6r + 2}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_1 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{6}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{16r^4 + 80r^3 + 140r^2 + 100r + 24}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{6}$
a_2	$\frac{1}{16r^4+80r^3+140r^2+100r+24}$	$\frac{1}{120}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{64r^6 + 672r^5 + 2800r^4 + 5880r^3 + 6496r^2 + 3528r + 720}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = -\frac{1}{5040}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{6}$
a_2	$\frac{1}{16r^4+80r^3+140r^2+100r+24}$	$\frac{1}{120}$
a_3	$-\frac{1}{64r^6+672r^5+2800r^4+5880r^3+6496r^2+3528r+720}$	$-\frac{1}{5040}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r^2 + 15r + 28)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{1}{362880}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{6}$
a_2	$\frac{1}{16r^4+80r^3+140r^2+100r+24}$	$\frac{1}{120}$
a_3	$-\frac{1}{64r^6+672r^5+2800r^4+5880r^3+6496r^2+3528r+720}$	$-\frac{1}{5040}$
a_4	$\frac{1}{16(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r^2+15r+28)}$	$\frac{1}{362880}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{32(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r^2 + 15r + 28)(2r^2 + 19r + 45)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = -\frac{1}{39916800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{6}$
a_2	$\frac{1}{16r^4+80r^3+140r^2+100r+24}$	$\frac{1}{120}$
a_3	$-\frac{1}{64r^6+672r^5+2800r^4+5880r^3+6496r^2+3528r+720}$	$-\frac{1}{5040}$
a_4	$\frac{1}{16(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r^2+15r+28)}$	$\frac{1}{362880}$
a_5	$-\frac{1}{32(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r^2+15r+28)(2r^2+19r+45)}$	$-\frac{1}{39916800}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= \sqrt{x} \left(1 - \frac{x}{6} + \frac{x^2}{120} - \frac{x^3}{5040} + \frac{x^4}{362880} - \frac{x^5}{39916800} + O(x^6) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$4b_n(n+r)(n+r-1) + 2(n+r)b_n + b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{b_{n-1}}{2(2n^2 + 4nr + 2r^2 - n - r)} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = -\frac{b_{n-1}}{4n^2 - 2n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -\frac{1}{4r^2 + 6r + 2}$$

Which for the root $r = 0$ becomes

$$b_1 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{16r^4 + 80r^3 + 140r^2 + 100r + 24}$$

Which for the root $r = 0$ becomes

$$b_2 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{2}$
b_2	$\frac{1}{16r^4+80r^3+140r^2+100r+24}$	$\frac{1}{24}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{1}{64r^6 + 672r^5 + 2800r^4 + 5880r^3 + 6496r^2 + 3528r + 720}$$

Which for the root $r = 0$ becomes

$$b_3 = -\frac{1}{720}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{2}$
b_2	$\frac{1}{16r^4+80r^3+140r^2+100r+24}$	$\frac{1}{24}$
b_3	$-\frac{1}{64r^6+672r^5+2800r^4+5880r^3+6496r^2+3528r+720}$	$-\frac{1}{720}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{16(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r^2 + 15r + 28)}$$

Which for the root $r = 0$ becomes

$$b_4 = \frac{1}{40320}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{2}$
b_2	$\frac{1}{16r^4+80r^3+140r^2+100r+24}$	$\frac{1}{24}$
b_3	$-\frac{1}{64r^6+672r^5+2800r^4+5880r^3+6496r^2+3528r+720}$	$-\frac{1}{720}$
b_4	$\frac{1}{16(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r^2+15r+28)}$	$\frac{1}{40320}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{1}{32(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r^2 + 15r + 28)(2r^2 + 19r + 45)}$$

Which for the root $r = 0$ becomes

$$b_5 = -\frac{1}{3628800}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+6r+2}$	$-\frac{1}{2}$
b_2	$\frac{1}{16r^4+80r^3+140r^2+100r+24}$	$\frac{1}{24}$
b_3	$-\frac{1}{64r^6+672r^5+2800r^4+5880r^3+6496r^2+3528r+720}$	$-\frac{1}{720}$
b_4	$\frac{1}{16(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r^2+15r+28)}$	$\frac{1}{40320}$
b_5	$-\frac{1}{32(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r^2+15r+28)(2r^2+19r+45)}$	$-\frac{1}{3628800}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= 1 - \frac{x}{2} + \frac{x^2}{24} - \frac{x^3}{720} + \frac{x^4}{40320} - \frac{x^5}{3628800} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\&= c_1 \sqrt{x} \left(1 - \frac{x}{6} + \frac{x^2}{120} - \frac{x^3}{5040} + \frac{x^4}{362880} - \frac{x^5}{39916800} + O(x^6) \right) \\&\quad + c_2 \left(1 - \frac{x}{2} + \frac{x^2}{24} - \frac{x^3}{720} + \frac{x^4}{40320} - \frac{x^5}{3628800} + O(x^6) \right)\end{aligned}$$

Hence the final solution is

$$\begin{aligned}y &= y_h \\&= c_1 \sqrt{x} \left(1 - \frac{x}{6} + \frac{x^2}{120} - \frac{x^3}{5040} + \frac{x^4}{362880} - \frac{x^5}{39916800} + O(x^6) \right) \\&\quad + c_2 \left(1 - \frac{x}{2} + \frac{x^2}{24} - \frac{x^3}{720} + \frac{x^4}{40320} - \frac{x^5}{3628800} + O(x^6) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= c_1 \sqrt{x} \left(1 - \frac{x}{6} + \frac{x^2}{120} - \frac{x^3}{5040} + \frac{x^4}{362880} - \frac{x^5}{39916800} + O(x^6) \right) \\&\quad + c_2 \left(1 - \frac{x}{2} + \frac{x^2}{24} - \frac{x^3}{720} + \frac{x^4}{40320} - \frac{x^5}{3628800} + O(x^6) \right)\end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned}y &= c_1 \sqrt{x} \left(1 - \frac{x}{6} + \frac{x^2}{120} - \frac{x^3}{5040} + \frac{x^4}{362880} - \frac{x^5}{39916800} + O(x^6) \right) \\&\quad + c_2 \left(1 - \frac{x}{2} + \frac{x^2}{24} - \frac{x^3}{720} + \frac{x^4}{40320} - \frac{x^5}{3628800} + O(x^6) \right)\end{aligned}$$

Verified OK.

4.36.1 Maple step by step solution

Let's solve

$$4xy'' + 2y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{4x} - \frac{y'}{2x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{2x} + \frac{y}{4x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1}{2x}, P_3(x) = \frac{1}{4x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{1}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4xy'' + 2y' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k- > k+1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r-1}$$

- Shift index using $k- > k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)(k+r)x^{k+r}$$

Rewrite ODE with series expansions

$$2a_0r(-1+2r)x^{-1+r} + \left(\sum_{k=0}^{\infty} (2a_{k+1}(k+1+r)(2k+1+2r) + a_k)x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$2r(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, \frac{1}{2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$4(k+1+r)\left(k+\frac{1}{2}+r\right)a_{k+1} + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{2(k+1+r)(2k+1+2r)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{a_k}{2(k+1)(2k+1)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{a_k}{2(k+1)(2k+1)} \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+1} = -\frac{a_k}{2\left(k+\frac{3}{2}\right)(2k+2)}$$

- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+1} = -\frac{a_k}{2\left(k+\frac{3}{2}\right)(2k+2)} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+1} = -\frac{a_k}{2(k+1)(2k+1)}, b_{k+1} = -\frac{b_k}{2(k+\frac{3}{2})(2k+2)} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 44

```

Order:=6;
dsolve(4*x*diff(y(x), x$2) +2*diff(y(x),x)+y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = c_1 \sqrt{x} \left(1 - \frac{1}{6}x + \frac{1}{120}x^2 - \frac{1}{5040}x^3 + \frac{1}{362880}x^4 - \frac{1}{39916800}x^5 + O(x^6) \right) \\ + c_2 \left(1 - \frac{1}{2}x + \frac{1}{24}x^2 - \frac{1}{720}x^3 + \frac{1}{40320}x^4 - \frac{1}{3628800}x^5 + O(x^6) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 85

```

AsymptoticDSolveValue[4*x*y''[x] +2*y'[x]+y[x] == 0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_1 \sqrt{x} \left(-\frac{x^5}{39916800} + \frac{x^4}{362880} - \frac{x^3}{5040} + \frac{x^2}{120} - \frac{x}{6} + 1 \right) \\ + c_2 \left(-\frac{x^5}{3628800} + \frac{x^4}{40320} - \frac{x^3}{720} + \frac{x^2}{24} - \frac{x}{2} + 1 \right)$$

4.37 problem 34

4.37.1 Maple step by step solution 1938

Internal problem ID [7258]

Internal file name [OUTPUT/6244_Sunday_June_05_2022_04_35_15_PM_27026346/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 34.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

`[[_Emden , _Fowler]]`

$$xy'' + y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$xy'' + y' - y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{1}{x}$$

Table 193: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	"regular"

$q(x) = -\frac{1}{x}$	
singularity	type
$x = 0$	"regular"

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$xy'' + y' - y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-a_n x^{n+r}) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r-1})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r-1}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^{-1+r} r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^{-1+r} r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = 1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(r+1)^2(2+r)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{36}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{576}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{14400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$
a_5	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{14400}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{1}{(r+1)^2}$	1	$-\frac{2}{(r+1)^3}$	-2
b_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$	$\frac{-6-4r}{(r+1)^3(2+r)^3}$	$-\frac{3}{4}$
b_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$	$\frac{-6r^2-24r-22}{(r+1)^3(2+r)^3(r+3)^3}$	$-\frac{11}{108}$
b_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$	$\frac{-8r^3-60r^2-140r-100}{(r+1)^3(2+r)^3(r+3)^3(r+4)^3}$	$-\frac{25}{3456}$
b_5	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{14400}$	$\frac{-10r^4-120r^3-510r^2-900r-548}{(r+1)^3(2+r)^3(r+3)^3(r+4)^3(r+5)^3}$	$-\frac{137}{432000}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) \\ &\quad - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\ &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\
 &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\
 &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\
 &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right)
 \end{aligned}$$

Verified OK.

4.37.1 Maple step by step solution

Let's solve

$$xy'' + y' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{x} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1}{x}, P_3(x) = -\frac{1}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$xy'' + y' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r^2 x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1}(k+1+r)^2 - a_k) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $r^2 = 0$
- Values of r that satisfy the indicial equation
 $r = 0$
- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1)^2 - a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k}{(k+1)^2}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k}{(k+1)^2}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = \frac{a_k}{(k+1)^2} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

```
Order:=6;  
dsolve(x*diff(y(x), x$2) +diff(y(x),x)-y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 + x + \frac{1}{4}x^2 + \frac{1}{36}x^3 + \frac{1}{576}x^4 + \frac{1}{14400}x^5 + O(x^6) \right) + \left((-2)x - \frac{3}{4}x^2 - \frac{11}{108}x^3 - \frac{25}{3456}x^4 - \frac{137}{432000}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 107

```
AsymptoticDSolveValue[x*y''[x] +y'[x]-y[x] == 0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) + c_2 \left(-\frac{137x^5}{432000} - \frac{25x^4}{3456} - \frac{11x^3}{108} - \frac{3x^2}{4} + \left(\frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) \log(x) - 2x \right)$$

4.38 problem 35

4.38.1 Maple step by step solution 1950

Internal problem ID [7259]

Internal file name [OUTPUT/6245_Sunday_June_05_2022_04_35_18_PM_25840892/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 35.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$xy'' + (1 + x)y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$xy'' + (1 + x)y' + 2y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1 + x}{x}$$
$$q(x) = \frac{2}{x}$$

Table 195: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1+x}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{2}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$xy'' + (1+x)y' + 2y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + (1+x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r} a_n (n+r) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r-1} \\ \sum_{n=0}^{\infty} 2a_n x^{n+r} &= \sum_{n=1}^{\infty} 2a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r-1} \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} 2a_{n-1} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^{-1+r} r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^{-1+r}r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_{n-1}(n+r-1) + a_n(n+r) + 2a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}(n+r+1)}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-1}(n+1)}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-2-r}{(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = -2$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{(r+1)^2}$	-2

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{3+r}{(2+r)(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{3}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{(r+1)^2}$	-2
a_2	$\frac{3+r}{(2+r)(r+1)^2}$	$\frac{3}{2}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-4 - r}{(3 + r)(2 + r)(r + 1)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{2}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{(r+1)^2}$	-2
a_2	$\frac{3+r}{(2+r)(r+1)^2}$	$\frac{3}{2}$
a_3	$\frac{-4-r}{(3+r)(2+r)(r+1)^2}$	$-\frac{2}{3}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{5 + r}{(4 + r)(3 + r)(2 + r)(r + 1)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{5}{24}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{(r+1)^2}$	-2
a_2	$\frac{3+r}{(2+r)(r+1)^2}$	$\frac{3}{2}$
a_3	$\frac{-4-r}{(3+r)(2+r)(r+1)^2}$	$-\frac{2}{3}$
a_4	$\frac{5+r}{(4+r)(3+r)(2+r)(r+1)^2}$	$\frac{5}{24}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-6 - r}{(5 + r)(4 + r)(3 + r)(2 + r)(r + 1)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{1}{20}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2-r}{(r+1)^2}$	-2
a_2	$\frac{3+r}{(2+r)(r+1)^2}$	$\frac{3}{2}$
a_3	$\frac{-4-r}{(3+r)(2+r)(r+1)^2}$	$-\frac{2}{3}$
a_4	$\frac{5+r}{(4+r)(3+r)(2+r)(r+1)^2}$	$\frac{5}{24}$
a_5	$\frac{-6-r}{(5+r)(4+r)(3+r)(2+r)(r+1)^2}$	$-\frac{1}{20}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r=0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{-2-r}{(r+1)^2}$	-2	$\frac{3+r}{(r+1)^3}$	3
b_2	$\frac{3+r}{(2+r)(r+1)^2}$	$\frac{3}{2}$	$\frac{-2r^2-11r-13}{(2+r)^2(r+1)^3}$	$-\frac{13}{4}$
b_3	$\frac{-4-r}{(3+r)(2+r)(r+1)^2}$	$-\frac{2}{3}$	$\frac{3r^3+27r^2+74r+62}{(3+r)^2(2+r)^2(r+1)^3}$	$\frac{31}{18}$
b_4	$\frac{5+r}{(4+r)(3+r)(2+r)(r+1)^2}$	$\frac{5}{24}$	$\frac{-4r^4-54r^3-256r^2-504r-346}{(4+r)^2(3+r)^2(2+r)^2(r+1)^3}$	$-\frac{173}{288}$
b_5	$\frac{-6-r}{(5+r)(4+r)(3+r)(2+r)(r+1)^2}$	$-\frac{1}{20}$	$\frac{5r^5+95r^4+685r^3+2335r^2+3744r+2244}{(5+r)^2(4+r)^2(3+r)^2(2+r)^2(r+1)^3}$	$\frac{187}{1200}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned}
y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\
&= \left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \ln(x) \\
&\quad + 3x - \frac{13x^2}{4} + \frac{31x^3}{18} - \frac{173x^4}{288} + \frac{187x^5}{1200} + O(x^6)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
&= c_1 \left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \\
&\quad + c_2 \left(\left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \ln(x) + 3x - \frac{13x^2}{4} + \frac{31x^3}{18} \right. \\
&\quad \left. - \frac{173x^4}{288} + \frac{187x^5}{1200} + O(x^6) \right)
\end{aligned}$$

Hence the final solution is

$$\begin{aligned}
y &= y_h \\
&= c_1 \left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \\
&\quad + c_2 \left(\left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \ln(x) + 3x - \frac{13x^2}{4} + \frac{31x^3}{18} \right. \\
&\quad \left. - \frac{173x^4}{288} + \frac{187x^5}{1200} + O(x^6) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \\ + c_2 \left(\left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \ln(x) + 3x - \frac{13x^2}{4} + \frac{31x^3}{18} \right. \\ \left. - \frac{173x^4}{288} + \frac{187x^5}{1200} + O(x^6) \right)$$

Verification of solutions

$$y = c_1 \left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \\ + c_2 \left(\left(1 - 2x + \frac{3x^2}{2} - \frac{2x^3}{3} + \frac{5x^4}{24} - \frac{x^5}{20} + O(x^6) \right) \ln(x) + 3x - \frac{13x^2}{4} + \frac{31x^3}{18} \right. \\ \left. - \frac{173x^4}{288} + \frac{187x^5}{1200} + O(x^6) \right)$$

Verified OK.

4.38.1 Maple step by step solution

Let's solve

$$xy'' + (1+x)y' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(1+x)y'}{x} - \frac{2y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(1+x)y'}{x} + \frac{2y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1+x}{x}, P_3(x) = \frac{2}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$xy'' + (1 + x)y' + 2y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r^2 x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)^2 + a_k (k+r+2)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1)^2 + a_k(k+2) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k(k+2)}{(k+1)^2}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{a_k(k+2)}{(k+1)^2}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{a_k(k+2)}{(k+1)^2} \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 59

```

Order:=6;
dsolve(x*dif(y(x), x$2) +(1+x)*dif(y(x),x)+2*y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - 2x + \frac{3}{2}x^2 - \frac{2}{3}x^3 + \frac{5}{24}x^4 - \frac{1}{20}x^5 + O(x^6) \right) + \left(3x - \frac{13}{4}x^2 + \frac{31}{18}x^3 - \frac{173}{288}x^4 + \frac{187}{1200}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 111

```
AsymptoticDSolveValue[x*y''[x] + (1+x)*y'[x] + 2*y[x] == 0, y[x], {x, 0, 5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{20} + \frac{5x^4}{24} - \frac{2x^3}{3} + \frac{3x^2}{2} - 2x + 1 \right) + c_2 \left(\frac{187x^5}{1200} - \frac{173x^4}{288} + \frac{31x^3}{18} - \frac{13x^2}{4} + \left(-\frac{x^5}{20} + \frac{5x^4}{24} - \frac{2x^3}{3} + \frac{3x^2}{2} - 2x + 1 \right) \log(x) + 3x \right)$$

4.39 problem 36

4.39.1 Maple step by step solution 1966

Internal problem ID [7260]

Internal file name [OUTPUT/6246_Sunday_June_05_2022_04_35_21_PM_68480318/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 36.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$x(x - 1)y'' + 3xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x^2 - x)y'' + 3xy' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3}{x - 1}$$
$$q(x) = \frac{1}{x(x - 1)}$$

Table 197: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3}{x-1}$	
singularity	type
$x = 1$	“regular”

$q(x) = \frac{1}{x(x-1)}$	
singularity	type
$x = 0$	“regular”
$x = 1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[1, 0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x(x-1)y'' + 3xy' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x(x-1) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + 3x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r-1} a_n (n+r) (n+r-1)) \quad (2A) \\ & + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned}$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \\ \sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) &= \sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) x^{n+r-1} \\ \sum_{n=0}^{\infty} a_n x^{n+r} &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-x^{n+r-1} a_n (n+r) (n+r-1)) \quad (2B) \\ & + \left(\sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$-x^{n+r-1} a_n (n+r) (n+r-1) = 0$$

When $n=0$ the above becomes

$$-x^{-1+r} a_0 r (-1+r) = 0$$

Or

$$-x^{-1+r} a_0 r (-1+r) = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$-x^{-1+r}r(-1+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$-r(-1+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 1$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$-x^{-1+r}r(-1+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+1}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots

of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_{n-1}(n+r-1)(n+r-2) - a_n(n+r)(n+r-1) + 3a_{n-1}(n+r-1) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{(n+r)a_{n-1}}{n+r-1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{(n+1)a_{n-1}}{n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1+r}{r}$$

Which for the root $r = 1$ becomes

$$a_1 = 2$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1+r}{r}$	2

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{2+r}{r}$$

Which for the root $r = 1$ becomes

$$a_2 = 3$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1+r}{r}$	2
a_2	$\frac{2+r}{r}$	3

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{3+r}{r}$$

Which for the root $r = 1$ becomes

$$a_3 = 4$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1+r}{r}$	2
a_2	$\frac{2+r}{r}$	3
a_3	$\frac{3+r}{r}$	4

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{4+r}{r}$$

Which for the root $r = 1$ becomes

$$a_4 = 5$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1+r}{r}$	2
a_2	$\frac{2+r}{r}$	3
a_3	$\frac{3+r}{r}$	4
a_4	$\frac{4+r}{r}$	5

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{5 + r}{r}$$

Which for the root $r = 1$ becomes

$$a_5 = 6$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1+r}{r}$	2
a_2	$\frac{2+r}{r}$	3
a_3	$\frac{3+r}{r}$	4
a_4	$\frac{4+r}{r}$	5
a_5	$\frac{5+r}{r}$	6

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= \frac{1 + r}{r} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} \frac{1 + r}{r} &= \lim_{r \rightarrow 0} \frac{1 + r}{r} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x(x-1)y'' + 3xy' + y = 0$ gives

$$\begin{aligned} &x(x-1) \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ &\quad + 3x \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \right) \\ &\quad + Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned}$$

Which can be written as

$$\begin{aligned} & \left((x(x-1)y_1''(x) + 3y_1'(x)x + y_1(x)) \ln(x) + x(x-1) \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) \right. \\ & \left. + 3y_1(x) \right) C + x(x-1) \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\ & + 3x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (7)$$

But since $y_1(x)$ is a solution to the ode, then

$$x(x-1)y_1''(x) + 3y_1'(x)x + y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned} & \left(x(x-1) \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + 3y_1(x) \right) C \\ & + x(x-1) \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\ & + 3x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \frac{\left(2x(x-1) \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) + (2x+1) \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C}{x} \\ & + \frac{x^2(x-1) \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) + 3 \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) x^2 + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) x}{x} \\ & = 0 \end{aligned} \quad (9)$$

Since $r_1 = 1$ and $r_2 = 0$ then the above becomes

$$\begin{aligned} & \frac{\left(2x(x-1) \left(\sum_{n=0}^{\infty} x^n a_n (n+1) \right) + (2x+1) \left(\sum_{n=0}^{\infty} a_n x^{n+1} \right) \right) C}{x} \\ & + \frac{x^2(x-1) \left(\sum_{n=0}^{\infty} x^{-2+n} b_n n (n-1) \right) + 3 \left(\sum_{n=0}^{\infty} x^{n-1} b_n n \right) x^2 + \left(\sum_{n=0}^{\infty} b_n x^n \right) x}{x} = 0 \end{aligned} \quad (10)$$

Which simplifies to

$$\begin{aligned}
& \left(\sum_{n=0}^{\infty} 2C x^{n+1} a_n (n+1) \right) + \sum_{n=0}^{\infty} (-2C x^n a_n (n+1)) \\
& + \left(\sum_{n=0}^{\infty} 2C x^{n+1} a_n \right) + \left(\sum_{n=0}^{\infty} C a_n x^n \right) + \left(\sum_{n=0}^{\infty} x^n b_n n (n-1) \right) \\
& + \sum_{n=0}^{\infty} (-n x^{n-1} b_n (n-1)) + \left(\sum_{n=0}^{\infty} 3x^n b_n n \right) + \left(\sum_{n=0}^{\infty} b_n x^n \right) = 0
\end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-1} and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=0}^{\infty} 2C x^{n+1} a_n (n+1) &= \sum_{n=2}^{\infty} 2C a_{-2+n} (n-1) x^{n-1} \\
\sum_{n=0}^{\infty} (-2C x^n a_n (n+1)) &= \sum_{n=1}^{\infty} (-2C a_{n-1} n x^{n-1}) \\
\sum_{n=0}^{\infty} 2C x^{n+1} a_n &= \sum_{n=2}^{\infty} 2C a_{-2+n} x^{n-1} \\
\sum_{n=0}^{\infty} C a_n x^n &= \sum_{n=1}^{\infty} C a_{n-1} x^{n-1} \\
\sum_{n=0}^{\infty} x^n b_n n (n-1) &= \sum_{n=1}^{\infty} (n-1) b_{n-1} (-2+n) x^{n-1} \\
\sum_{n=0}^{\infty} 3x^n b_n n &= \sum_{n=1}^{\infty} 3(n-1) b_{n-1} x^{n-1} \\
\sum_{n=0}^{\infty} b_n x^n &= \sum_{n=1}^{\infty} b_{n-1} x^{n-1}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers

of x are the same and equal to $n - 1$.

$$\begin{aligned}
& \left(\sum_{n=2}^{\infty} 2Ca_{-2+n}(n-1)x^{n-1} \right) + \sum_{n=1}^{\infty} (-2Ca_{n-1}nx^{n-1}) \\
& + \left(\sum_{n=2}^{\infty} 2Ca_{-2+n}x^{n-1} \right) + \left(\sum_{n=1}^{\infty} Ca_{n-1}x^{n-1} \right) \\
& + \left(\sum_{n=1}^{\infty} (n-1)b_{n-1}(-2+n)x^{n-1} \right) + \sum_{n=0}^{\infty} (-nx^{n-1}b_n(n-1)) \\
& + \left(\sum_{n=1}^{\infty} 3(n-1)b_{n-1}x^{n-1} \right) + \left(\sum_{n=1}^{\infty} b_{n-1}x^{n-1} \right) = 0
\end{aligned} \tag{2B}$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = N$, where $N = 1$ which is the difference between the two roots, we are free to choose $b_1 = 0$. Hence for $n = 1$, Eq (2B) gives

$$-C + 1 = 0$$

Which is solved for C . Solving for C gives

$$C = 1$$

For $n = 2$, Eq (2B) gives

$$(4a_0 - 3a_1)C + 4b_1 - 2b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-2 - 2b_2 = 0$$

Solving the above for b_2 gives

$$b_2 = -1$$

For $n = 3$, Eq (2B) gives

$$(6a_1 - 5a_2)C + 9b_2 - 6b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-12 - 6b_3 = 0$$

Solving the above for b_3 gives

$$b_3 = -2$$

For $n = 4$, Eq (2B) gives

$$(8a_2 - 7a_3)C + 16b_3 - 12b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-36 - 12b_4 = 0$$

Solving the above for b_4 gives

$$b_4 = -3$$

For $n = 5$, Eq (2B) gives

$$(10a_3 - 9a_4)C + 25b_4 - 20b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-80 - 20b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = -4$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = 1$ and all b_n , then the second solution becomes

$$y_2(x) = 1(x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6))) \ln(x) + 1 - x^2 - 2x^3 - 3x^4 - 4x^5 + O(x^6)$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \\ &\quad + c_2 (1(x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6))) \ln(x) + 1 - x^2 - 2x^3 \\ &\quad \quad \quad - 3x^4 - 4x^5 + O(x^6)) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \\ &\quad + c_2 (x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6))) \ln(x) + 1 - x^2 - 2x^3 - 3x^4 - 4x^5 \\ &\quad \quad \quad + O(x^6) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \\ + c_2(x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \ln(x) + 1 - x^2 - 2x^3 - 3x^4(1) \\ - 4x^5 + O(x^6))$$

Verification of solutions

$$y = c_1x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \\ + c_2(x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \ln(x) + 1 - x^2 - 2x^3 - 3x^4 - 4x^5 \\ + O(x^6))$$

Verified OK.

4.39.1 Maple step by step solution

Let's solve

$$x(x-1)y'' + 3xy' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{x(x-1)} - \frac{3y'}{x-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{x-1} + \frac{y}{x(x-1)} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{3}{x-1}, P_3(x) = \frac{1}{x(x-1)} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x(x-1)y'' + 3xy' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^m \cdot y''$ to series expansion for $m = 1..2$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-2+m}$$

- Shift index using $k- > k+2-m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) x^{k+r}$$

Rewrite ODE with series expansions

$$-a_0 r(-1+r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (-a_{k+1}(k+r+1)(k+r) + a_k(k+r+1)^2) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(-1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{0, 1\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k+r+1)(-a_{k+1}(k+r) + a_k(k+r+1)) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k(k+r+1)}{k+r}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k(k+1)}{k}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = \frac{a_k(k+1)}{k} \right]$$

- Recursion relation for $r = 1$

$$a_{k+1} = \frac{a_k(k+2)}{k+1}$$

- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = \frac{a_k(k+2)}{k+1} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1} \right), a_{k+1} = \frac{a_k(k+1)}{k}, b_{k+1} = \frac{b_k(k+2)}{k+1} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 60

```

Order:=6;
dsolve(x*(x-1)*diff(y(x), x$2) +3*x*diff(y(x),x)+y(x) = 0,y(x),type='series',x=0);

```

$$\begin{aligned}
 y(x) = & c_1 x(1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + O(x^6)) \\
 & + (x + 2x^2 + 3x^3 + 4x^4 + 5x^5 + O(x^6)) \ln(x) c_2 \\
 & + (1 + 3x + 5x^2 + 7x^3 + 9x^4 + 11x^5 + O(x^6)) c_2
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 63

```
AsymptoticDSolveValue[x*(x-1)*y''[x] + 3*x*y'[x] + y[x] == 0, y[x], {x, 0, 5}]
```

$$y(x) \rightarrow c_1(x^4 + x^3 + x^2 + (4x^3 + 3x^2 + 2x + 1)x \log(x) + x + 1) \\ + c_2(5x^5 + 4x^4 + 3x^3 + 2x^2 + x)$$

4.40 problem 37

4.40.1 Maple step by step solution 1978

Internal problem ID [7261]

Internal file name [OUTPUT/6247_Sunday_June_05_2022_04_35_27_PM_89333441/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 37.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2(x^2 - 2x + 1)y'' - x(x + 3)y' + (4 + x)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x^4 - 2x^3 + x^2)y'' + (-x^2 - 3x)y' + (4 + x)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{x + 3}{x(x - 1)^2}$$
$$q(x) = \frac{4 + x}{x^2(x - 1)^2}$$

Table 199: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{x+3}{x(x-1)^2}$		$q(x) = \frac{4+x}{x^2(x-1)^2}$	
singularity	type	singularity	type
$x = 0$	“regular”	$x = 0$	“regular”
$x = 1$	“irregular”	$x = 1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[1]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2(x^2 - 2x + 1)y'' + (-x^2 - 3x)y' + (4 + x)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2(x^2 - 2x + 1) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + (-x^2 - 3x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (4+x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\begin{aligned}
& \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-2x^{1+n+r} a_n (n+r) (n+r-1)) \\
& + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n (n+r)) \\
& + \sum_{n=0}^{\infty} (-3x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 4a_n x^{n+r} \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0
\end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=0}^{\infty} x^{n+r+2} a_n (n+r) (n+r-1) &= \sum_{n=2}^{\infty} a_{n-2} (n+r-2) (n-3+r) x^{n+r} \\
\sum_{n=0}^{\infty} (-2x^{1+n+r} a_n (n+r) (n+r-1)) &= \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) (n+r-2) x^{n+r}) \\
\sum_{n=0}^{\infty} (-x^{1+n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r}) \\
\sum_{n=0}^{\infty} x^{1+n+r} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned}
& \left(\sum_{n=2}^{\infty} a_{n-2} (n+r-2) (n-3+r) x^{n+r} \right) \\
& + \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) (n+r-2) x^{n+r}) \\
& + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r}) \\
& + \sum_{n=0}^{\infty} (-3x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 4a_n x^{n+r} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) = 0
\end{aligned} \tag{2B}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - 3x^{n+r} a_n (n+r) + 4a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r (-1+r) - 3x^r a_0 r + 4a_0 x^r = 0$$

Or

$$(x^r r (-1+r) - 3x^r r + 4x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(-2+r)^2 x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$(-2+r)^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2$$

$$r_2 = 2$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(-2+r)^2 x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \tag{1A}$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \tag{1B}$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitray constants of

integration which can be found from initial conditions. Using the value of the indicial root found earlier, $r = 2$, Eqs (1A,1B) become

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+2}$$

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+2} \right)$$

We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{2r + 1}{-1 + r}$$

For $2 \leq n$ the recursive equation is

$$a_{n-2}(n+r-2)(n-3+r) - 2a_{n-1}(n+r-1)(n+r-2) + a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) - 3a_n(n+r) + 4a_n + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{na_{n-2} - 2na_{n-1} + ra_{n-2} - 2ra_{n-1} - 3a_{n-2} + a_{n-1}}{n+r-2} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = \frac{(-a_{n-2} + 2a_{n-1})n + a_{n-2} + 3a_{n-1}}{n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r+1}{-1+r}$	5

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{3r^2 + 10r + 2}{r(-1+r)}$$

Which for the root $r = 2$ becomes

$$a_2 = 17$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r+1}{-1+r}$	5
a_2	$\frac{3r^2+10r+2}{r(-1+r)}$	17

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{4r^3 + 34r^2 + 54r + 10}{r^3 - r}$$

Which for the root $r = 2$ becomes

$$a_3 = \frac{143}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r+1}{-1+r}$	5
a_2	$\frac{3r^2+10r+2}{r(-1+r)}$	17
a_3	$\frac{4r^3+34r^2+54r+10}{r^3-r}$	$\frac{143}{3}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{5r^4 + 80r^3 + 321r^2 + 384r + 68}{(2+r)r(r^2-1)}$$

Which for the root $r = 2$ becomes

$$a_4 = \frac{355}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r+1}{-1+r}$	5
a_2	$\frac{3r^2+10r+2}{r(-1+r)}$	17
a_3	$\frac{4r^3+34r^2+54r+10}{r^3-r}$	$\frac{143}{3}$
a_4	$\frac{5r^4+80r^3+321r^2+384r+68}{(2+r)r(r^2-1)}$	$\frac{355}{3}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{6r^5 + 155r^4 + 1156r^3 + 3295r^2 + 3336r + 572}{r^5 + 5r^4 + 5r^3 - 5r^2 - 6r}$$

Which for the root $r = 2$ becomes

$$a_5 = \frac{4043}{15}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r+1}{-1+r}$	5
a_2	$\frac{3r^2+10r+2}{r(-1+r)}$	17
a_3	$\frac{4r^3+34r^2+54r+10}{r^3-r}$	$\frac{143}{3}$
a_4	$\frac{5r^4+80r^3+321r^2+384r+68}{(2+r)r(r^2-1)}$	$\frac{355}{3}$
a_5	$\frac{6r^5+155r^4+1156r^3+3295r^2+3336r+572}{r^5+5r^4+5r^3-5r^2-6r}$	$\frac{4043}{15}$

Using the above table, then the first solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^2\left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6)\right) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r}\right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 2$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$
b_0	1	1	N/A since b_n starts from 1
b_1	$\frac{2r+1}{-1+r}$	5	$-\frac{3}{(-1+r)^2}$
b_2	$\frac{3r^2+10r+2}{r(-1+r)}$	17	$\frac{-13r^2-4r+2}{r^2(-1+r)^2}$
b_3	$\frac{4r^3+34r^2+54r+10}{r^3-r}$	$\frac{143}{3}$	$\frac{-34r^4-116r^3-64r^2+10}{r^2(r^2-1)^2}$
b_4	$\frac{5r^4+80r^3+321r^2+384r+68}{(2+r)r(r^2-1)}$	$\frac{355}{3}$	$\frac{-70r^6-652r^5-1904r^4-2128r^3-666r^2+136r+136}{(2+r)^2 r^2 (r^2-1)^2}$
b_5	$\frac{6r^5+155r^4+1156r^3+3295r^2+3336r+572}{r^5+5r^4+5r^3-5r^2-6r}$	$\frac{4043}{15}$	$\frac{-125r^8-2252r^7-14980r^6-47988r^5-77945r^4-58672r^3-11670r^2+5720r+3432}{r^2(r^4+5r^3+5r^2-5r-6)^2}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \ln(x) \\ &\quad + x^2 \left(-3x - \frac{29x^2}{2} - \frac{859x^3}{18} - \frac{4693x^4}{36} - \frac{285181x^5}{900} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \\ &\quad + c_2 \left(x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \ln(x) \right. \\ &\quad \left. + x^2 \left(-3x - \frac{29x^2}{2} - \frac{859x^3}{18} - \frac{4693x^4}{36} - \frac{285181x^5}{900} + O(x^6) \right) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \\
 &\quad + c_2 \left(x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \ln(x) \right. \\
 &\quad \left. + x^2 \left(-3x - \frac{29x^2}{2} - \frac{859x^3}{18} - \frac{4693x^4}{36} - \frac{285181x^5}{900} + O(x^6) \right) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \\
 &\quad + c_2 \left(x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \ln(x) \right. \\
 &\quad \left. + x^2 \left(-3x - \frac{29x^2}{2} - \frac{859x^3}{18} - \frac{4693x^4}{36} - \frac{285181x^5}{900} + O(x^6) \right) \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \\
 &\quad + c_2 \left(x^2 \left(17x^2 + 5x + 1 + \frac{143x^3}{3} + \frac{355x^4}{3} + \frac{4043x^5}{15} + O(x^6) \right) \ln(x) \right. \\
 &\quad \left. + x^2 \left(-3x - \frac{29x^2}{2} - \frac{859x^3}{18} - \frac{4693x^4}{36} - \frac{285181x^5}{900} + O(x^6) \right) \right)
 \end{aligned}$$

Verified OK.

4.40.1 Maple step by step solution

Let's solve

$$x^2(x^2 - 2x + 1)y'' + (-x^2 - 3x)y' + (4 + x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(4+x)y}{x^2(x^2-2x+1)} + \frac{(x+3)y'}{x(x^2-2x+1)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(x+3)y'}{x(x^2-2x+1)} + \frac{(4+x)y}{x^2(x^2-2x+1)} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{x+3}{x(x^2-2x+1)}, P_3(x) = \frac{4+x}{x^2(x^2-2x+1)} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -3$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 4$$

- $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2(x^2 - 2x + 1)y'' - x(x + 3)y' + (4 + x)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^m \cdot y'$ to series expansion for $m = 1..2$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^m \cdot y''$ to series expansion for $m = 2.4$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r-2+m}$$

- Shift index using $k \rightarrow k+2-m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m}(k+2-m+r)(k+1-m+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-2+r)^2 x^r + (a_1(-1+r)^2 - a_0(1+2r)(-1+r))x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r-2)^2 - a_{k-1}(2k-2+r)(k+r-2)) \right) x^{k+r}$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(-2+r)^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 2$$

- Each term must be 0

$$a_1(-1+r)^2 - a_0(1+2r)(-1+r) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = \frac{a_0(1+2r)}{-1+r}$$

- Each term in the series must be 0, giving the recursion relation

$$((a_k + a_{k-2} - 2a_{k-1})k + (a_k + a_{k-2} - 2a_{k-1})r - 2a_k - 3a_{k-2} + a_{k-1})(k+r-2) = 0$$

- Shift index using $k \rightarrow k+2$

$$((a_{k+2} + a_k - 2a_{k+1})(k+2) + (a_{k+2} + a_k - 2a_{k+1})r - 2a_{k+2} - 3a_k + a_{k+1})(k+r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{ka_k - 2ka_{k+1} + ra_k - 2ra_{k+1} - a_k - 3a_{k+1}}{k+r}$$

- Recursion relation for $r = 2$

$$a_{k+2} = -\frac{ka_k - 2ka_{k+1} + a_k - 7a_{k+1}}{k+2}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+2} = -\frac{ka_k - 2ka_{k+1} + a_k - 7a_{k+1}}{k+2}, a_1 = 5a_0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=6;
```

```
dsolve(x^2*(1-2*x+x^2)*diff(y(x), x$2) -x*(3+x)*diff(y(x),x)+(4+x)*y(x) = 0,y(x),type='series')
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 + 5x + 17x^2 + \frac{143}{3}x^3 + \frac{355}{3}x^4 + \frac{4043}{15}x^5 + O(x^6) \right) + \left((-3)x - \frac{29}{2}x^2 - \frac{859}{18}x^3 - \frac{4693}{36}x^4 - \frac{285181}{900}x^5 + O(x^6) \right) c_2 \right) x^2$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 118

```
AsymptoticDSolveValue[x^2*(1-2*x+x^2)*y'[x] -x*(3+x)*y'[x]+(4+x)*y[x] == 0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{4043x^5}{15} + \frac{355x^4}{3} + \frac{143x^3}{3} + 17x^2 + 5x + 1 \right) x^2 + c_2 \left(\left(-\frac{285181x^5}{900} - \frac{4693x^4}{36} - \frac{859x^3}{18} - \frac{29x^2}{2} - 3x \right) x^2 + \left(\frac{4043x^5}{15} + \frac{355x^4}{3} + \frac{143x^3}{3} + 17x^2 + 5x + 1 \right) x^2 \log(x) \right)$$

4.41 problem 38

4.41.1 Maple step by step solution 1990

Internal problem ID [7262]

Internal file name [OUTPUT/6248_Sunday_June_05_2022_04_35_29_PM_70306277/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 38.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2x^2(x+2)y'' + 5x^2y' + (1+x)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(2x^3 + 4x^2)y'' + 5x^2y' + (1+x)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{5}{2(x+2)}$$
$$q(x) = \frac{1+x}{2x^2(x+2)}$$

Table 201: Table $p(x), q(x)$ singularities.

$p(x) = \frac{5}{2(x+2)}$	
singularity	type
$x = -2$	“regular”

$q(x) = \frac{1+x}{2x^2(x+2)}$	
singularity	type
$x = -2$	“regular”
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-2, 0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2(x+2)y'' + 5x^2y' + (1+x)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2(x+2) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 5x^2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1+x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{1+n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) \quad (2A) \\ & + \left(\sum_{n=0}^{\infty} 5x^{1+n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0 \end{aligned}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 2x^{1+n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} 2a_{n-1} (n+r-1) (n+r-2) x^{n+r} \\ \sum_{n=0}^{\infty} 5x^{1+n+r} a_n (n+r) &= \sum_{n=1}^{\infty} 5a_{n-1} (n+r-1) x^{n+r} \\ \sum_{n=0}^{\infty} x^{1+n+r} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} 2a_{n-1} (n+r-1) (n+r-2) x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) \quad (2B) \\ & + \left(\sum_{n=1}^{\infty} 5a_{n-1} (n+r-1) x^{n+r} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$4x^{n+r} a_n (n+r) (n+r-1) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$4x^r a_0 r(-1+r) + a_0 x^r = 0$$

Or

$$(4x^r r(-1+r) + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r - 1)^2 x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$(2r - 1)^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = \frac{1}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r - 1)^2 x^r = 0$$

Solving for r gives the roots of the indicial equation as. Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. Using the value of the indicial root found earlier, $r = \frac{1}{2}$, Eqs (1A,1B) become

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+\frac{1}{2}} \right)$$

We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_{n-1}(n+r-1)(n+r-2) + 4a_n(n+r)(n+r-1) + 5a_{n-1}(n+r-1) + a_n + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{(n+r)a_{n-1}}{-1+2n+2r} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = -\frac{(2n+1)a_{n-1}}{4n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-1-r}{1+2r}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_1 = -\frac{3}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{1+2r}$	$-\frac{3}{4}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r^2 + 3r + 2}{4r^2 + 8r + 3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = \frac{15}{32}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{1+2r}$	$-\frac{3}{4}$
a_2	$\frac{r^2+3r+2}{4r^2+8r+3}$	$\frac{15}{32}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-r^3 - 6r^2 - 11r - 6}{8r^3 + 36r^2 + 46r + 15}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = -\frac{35}{128}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{1+2r}$	$-\frac{3}{4}$
a_2	$\frac{r^2+3r+2}{4r^2+8r+3}$	$\frac{15}{32}$
a_3	$\frac{-r^3-6r^2-11r-6}{8r^3+36r^2+46r+15}$	$-\frac{35}{128}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r^4 + 10r^3 + 35r^2 + 50r + 24}{16r^4 + 128r^3 + 344r^2 + 352r + 105}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{315}{2048}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{1+2r}$	$-\frac{3}{4}$
a_2	$\frac{r^2+3r+2}{4r^2+8r+3}$	$\frac{15}{32}$
a_3	$\frac{-r^3-6r^2-11r-6}{8r^3+36r^2+46r+15}$	$-\frac{35}{128}$
a_4	$\frac{r^4+10r^3+35r^2+50r+24}{16r^4+128r^3+344r^2+352r+105}$	$\frac{315}{2048}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-r^5 - 15r^4 - 85r^3 - 225r^2 - 274r - 120}{32r^5 + 400r^4 + 1840r^3 + 3800r^2 + 3378r + 945}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = -\frac{693}{8192}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{1+2r}$	$-\frac{3}{4}$
a_2	$\frac{r^2+3r+2}{4r^2+8r+3}$	$\frac{15}{32}$
a_3	$\frac{-r^3-6r^2-11r-6}{8r^3+36r^2+46r+15}$	$-\frac{35}{128}$
a_4	$\frac{r^4+10r^3+35r^2+50r+24}{16r^4+128r^3+344r^2+352r+105}$	$\frac{315}{2048}$
a_5	$\frac{-r^5-15r^4-85r^3-225r^2-274r-120}{32r^5+400r^4+1840r^3+3800r^2+3378r+945}$	$-\frac{693}{8192}$

Using the above table, then the first solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= \sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = \frac{1}{2}$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$
b_0	1	1	N/A since b_n starts from 1
b_1	$\frac{-1-r}{1+2r}$	$-\frac{3}{4}$	$\frac{1}{(1+2r)^2}$
b_2	$\frac{r^2+3r+2}{4r^2+8r+3}$	$\frac{15}{32}$	$\frac{-4r^2-10r-7}{(4r^2+8r+3)^2}$
b_3	$\frac{-r^3-6r^2-11r-6}{8r^3+36r^2+46r+15}$	$-\frac{35}{128}$	$\frac{12r^4+84r^3+219r^2+252r+111}{(8r^3+36r^2+46r+15)^2}$
b_4	$\frac{r^4+10r^3+35r^2+50r+24}{16r^4+128r^3+344r^2+352r+105}$	$\frac{315}{2048}$	$\frac{-32r^6-432r^5-2384r^4-6876r^3-10946r^2-9162r-3198}{(16r^4+128r^3+344r^2+352r+105)^2}$
b_5	$\frac{-r^5-15r^4-85r^3-225r^2-274r-120}{32r^5+400r^4+1840r^3+3800r^2+3378r+945}$	$-\frac{693}{8192}$	$\frac{80r^8+1760r^7+16600r^6+87560r^5+282265r^4+569360r^3+702575r^2+486750r+111111}{(32r^5+400r^4+1840r^3+3800r^2+3378r+945)^2}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \ln(x) \\ &\quad + \sqrt{x} \left(\frac{x}{4} - \frac{13x^2}{64} + \frac{101x^3}{768} - \frac{641x^4}{8192} + \frac{7303x^5}{163840} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 \sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \\ &\quad + c_2 \left(\sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \ln(x) \right. \\ &\quad \left. + \sqrt{x} \left(\frac{x}{4} - \frac{13x^2}{64} + \frac{101x^3}{768} - \frac{641x^4}{8192} + \frac{7303x^5}{163840} + O(x^6) \right) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 \sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \\
 &\quad + c_2 \left(\sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \ln(x) \right. \\
 &\quad \left. + \sqrt{x} \left(\frac{x}{4} - \frac{13x^2}{64} + \frac{101x^3}{768} - \frac{641x^4}{8192} + \frac{7303x^5}{163840} + O(x^6) \right) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 \sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \\
 &\quad + c_2 \left(\sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \ln(x) \right. \\
 &\quad \left. + \sqrt{x} \left(\frac{x}{4} - \frac{13x^2}{64} + \frac{101x^3}{768} - \frac{641x^4}{8192} + \frac{7303x^5}{163840} + O(x^6) \right) \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 \sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \\
 &\quad + c_2 \left(\sqrt{x} \left(1 - \frac{3x}{4} + \frac{15x^2}{32} - \frac{35x^3}{128} + \frac{315x^4}{2048} - \frac{693x^5}{8192} + O(x^6) \right) \ln(x) \right. \\
 &\quad \left. + \sqrt{x} \left(\frac{x}{4} - \frac{13x^2}{64} + \frac{101x^3}{768} - \frac{641x^4}{8192} + \frac{7303x^5}{163840} + O(x^6) \right) \right)
 \end{aligned}$$

Verified OK.

4.41.1 Maple step by step solution

Let's solve

$$2x^2(x+2)y'' + 5x^2y' + (1+x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(1+x)y}{2x^2(x+2)} - \frac{5y'}{2(x+2)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{5y'}{2(x+2)} + \frac{(1+x)y}{2x^2(x+2)} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{5}{2(x+2)}, P_3(x) = \frac{1+x}{2x^2(x+2)} \right]$$

- $(x+2) \cdot P_2(x)$ is analytic at $x = -2$

$$\left. ((x+2) \cdot P_2(x)) \right|_{x=-2} = \frac{5}{2}$$

- $(x+2)^2 \cdot P_3(x)$ is analytic at $x = -2$

$$\left. ((x+2)^2 \cdot P_3(x)) \right|_{x=-2} = 0$$

- $x = -2$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -2$$

- Multiply by denominators

$$2x^2(x+2)y'' + 5x^2y' + (1+x)y = 0$$

- Change variables using $x = u - 2$ so that the regular singular point is at $u = 0$

$$(2u^3 - 8u^2 + 8u) \left(\frac{d^2}{du^2} y(u) \right) + (5u^2 - 20u + 20) \left(\frac{d}{du} y(u) \right) + (-1 + u) y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot y(u)$ to series expansion for $m = 0..1$

$$u^m \cdot y(u) = \sum_{k=0}^{\infty} a_k u^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$u^m \cdot y(u) = \sum_{k=m}^{\infty} a_{k-m} u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..2$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k + 1 - m + r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..3$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k + r) (k + r - 1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k + 2 - m + r) (k + 1 - m + r) u^{k+r}$$

Rewrite ODE with series expansions

$$4a_0 r (3 + 2r) u^{-1+r} + (4a_1 (1 + r) (5 + 2r) - a_0 (8r^2 + 12r + 1)) u^r + \left(\sum_{k=1}^{\infty} (4a_{k+1} (k + r + 1) (2k + r + 1) - a_k (k + r) (k + r - 1)) u^{k+r} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$4r(3 + 2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, -\frac{3}{2} \right\}$$

- Each term must be 0

$$4a_1 (1 + r) (5 + 2r) - a_0 (8r^2 + 12r + 1) = 0$$

- Each term in the series must be 0, giving the recursion relation

$$2(-4a_k + a_{k-1} + 4a_{k+1}) k^2 + (4(-4a_k + a_{k-1} + 4a_{k+1}) r - 12a_k - a_{k-1} + 28a_{k+1}) k + 2(-4a_k + a_{k-1} + 4a_{k+1}) = 0$$

- Shift index using $k \rightarrow k + 1$

$$2(-4a_{k+1} + a_k + 4a_{k+2}) (k + 1)^2 + (4(-4a_{k+1} + a_k + 4a_{k+2}) r - 12a_{k+1} - a_k + 28a_{k+2}) (k + 1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{2k^2 a_k - 8k^2 a_{k+1} + 4k r a_k - 16k r a_{k+1} + 2r^2 a_k - 8r^2 a_{k+1} + 3k a_k - 28k a_{k+1} + 3r a_k - 28r a_{k+1} + a_k - 21a_{k+1}}{4(2k^2 + 4kr + 2r^2 + 11k + 11r + 14)}$$

- Recursion relation for $r = 0$

$$a_{k+2} = -\frac{2k^2 a_k - 8k^2 a_{k+1} + 3k a_k - 28k a_{k+1} + a_k - 21a_{k+1}}{4(2k^2 + 11k + 14)}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+2} = -\frac{2k^2 a_k - 8k^2 a_{k+1} + 3k a_k - 28k a_{k+1} + a_k - 21a_{k+1}}{4(2k^2 + 11k + 14)}, 20a_1 - a_0 = 0 \right]$$

- Revert the change of variables $u = x + 2$

$$\left[y = \sum_{k=0}^{\infty} a_k (x+2)^k, a_{k+2} = -\frac{2k^2 a_k - 8k^2 a_{k+1} + 3ka_k - 28ka_{k+1} + a_k - 21a_{k+1}}{4(2k^2 + 11k + 14)}, 20a_1 - a_0 = 0 \right]$$

- Recursion relation for $r = -\frac{3}{2}$

$$a_{k+2} = -\frac{2k^2 a_k - 8k^2 a_{k+1} - 3ka_k - 4ka_{k+1} + a_k + 3a_{k+1}}{4(2k^2 + 5k + 2)}$$

- Solution for $r = -\frac{3}{2}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k-\frac{3}{2}}, a_{k+2} = -\frac{2k^2 a_k - 8k^2 a_{k+1} - 3ka_k - 4ka_{k+1} + a_k + 3a_{k+1}}{4(2k^2 + 5k + 2)}, -4a_1 - a_0 = 0 \right]$$

- Revert the change of variables $u = x + 2$

$$\left[y = \sum_{k=0}^{\infty} a_k (x+2)^{k-\frac{3}{2}}, a_{k+2} = -\frac{2k^2 a_k - 8k^2 a_{k+1} - 3ka_k - 4ka_{k+1} + a_k + 3a_{k+1}}{4(2k^2 + 5k + 2)}, -4a_1 - a_0 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (x+2)^k \right) + \left(\sum_{k=0}^{\infty} b_k (x+2)^{k-\frac{3}{2}} \right), a_{k+2} = -\frac{2k^2 a_k - 8k^2 a_{k+1} + 3ka_k - 28ka_{k+1} + a_k - 21a_{k+1}}{4(2k^2 + 11k + 14)}, 20a_1 - a_0 = 0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

Order:=6;

```
dsolve(2*x^2*(2+x)*diff(y(x), x$2) +5*x^2*diff(y(x),x)+(1+x)*y(x) = 0,y(x),type='series',x=0
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 - \frac{3}{4}x + \frac{15}{32}x^2 - \frac{35}{128}x^3 + \frac{315}{2048}x^4 - \frac{693}{8192}x^5 + O(x^6) \right) + \left(\frac{1}{4}x - \frac{13}{64}x^2 + \frac{101}{768}x^3 - \frac{641}{8192}x^4 + \frac{7303}{163840}x^5 + O(x^6) \right) c_2 \right) \sqrt{x}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 134

```
AsymptoticDSolveValue[2*x^2*(2+x)*y'[x] +5*x^2*y'[x]+(1+x)*y[x] == 0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(-\frac{693x^5}{8192} + \frac{315x^4}{2048} - \frac{35x^3}{128} + \frac{15x^2}{32} - \frac{3x}{4} + 1 \right) + c_2 \left(\sqrt{x} \left(\frac{7303x^5}{163840} - \frac{641x^4}{8192} + \frac{101x^3}{768} - \frac{13x^2}{64} + \frac{x}{4} \right) + \sqrt{x} \left(-\frac{693x^5}{8192} + \frac{315x^4}{2048} - \frac{35x^3}{128} + \frac{15x^2}{32} - \frac{3x}{4} + 1 \right) \log(x) \right)$$

4.42 problem 39

4.42.1 Maple step by step solution 2006

Internal problem ID [7263]

Internal file name [OUTPUT/6249_Sunday_June_05_2022_04_35_32_PM_87536086/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 39.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2x^2y'' + xy' + (x - 5)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + xy' + (x - 5)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{2x}$$
$$q(x) = \frac{x - 5}{2x^2}$$

Table 203: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{x-5}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + xy' + (x - 5)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x-5) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) + \sum_{n=0}^{\infty} (-5a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{1+n+r} a_n = \sum_{n=1}^{\infty} a_{n-1} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) + \sum_{n=0}^{\infty} (-5a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - 5a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) + x^r a_0 r - 5a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) + x^r r - 5x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - r - 5) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - r - 5 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{4} + \frac{\sqrt{41}}{4}$$

$$r_2 = \frac{1}{4} - \frac{\sqrt{41}}{4}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - r - 5) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{\sqrt{41}}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{4}+\frac{\sqrt{41}}{4}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{4}-\frac{\sqrt{41}}{4}}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-1} - 5a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{2n^2 + 4nr + 2r^2 - n - r - 5} \quad (4)$$

Which for the root $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$ becomes

$$a_n = -\frac{a_{n-1}}{n(\sqrt{41} + 2n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{2r^2 + 3r - 4}$$

Which for the root $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$ becomes

$$a_1 = \frac{1}{-2 - \sqrt{41}}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2-\sqrt{41}}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4r^4 + 20r^3 + 15r^2 - 25r - 4}$$

Which for the root $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$ becomes

$$a_2 = \frac{1}{2(2 + \sqrt{41})(4 + \sqrt{41})}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2-\sqrt{41}}$
a_2	$\frac{1}{4r^4+20r^3+15r^2-25r-4}$	$\frac{1}{2(2+\sqrt{41})(4+\sqrt{41})}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{8r^6 + 84r^5 + 290r^4 + 315r^3 - 133r^2 - 294r - 40}$$

Which for the root $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$ becomes

$$a_3 = -\frac{1}{3240 + 510\sqrt{41}}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2-\sqrt{41}}$
a_2	$\frac{1}{4r^4+20r^3+15r^2-25r-4}$	$\frac{1}{2(2+\sqrt{41})(4+\sqrt{41})}$
a_3	$-\frac{1}{8r^6+84r^5+290r^4+315r^3-133r^2-294r-40}$	$-\frac{1}{3240+510\sqrt{41}}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16r^8 + 288r^7 + 2024r^6 + 6912r^5 + 11129r^4 + 4662r^3 - 7549r^2 - 7362r - 920}$$

Which for the root $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$ becomes

$$a_4 = \frac{1}{187320 + 29280\sqrt{41}}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2-\sqrt{41}}$
a_2	$\frac{1}{4r^4+20r^3+15r^2-25r-4}$	$\frac{1}{2(2+\sqrt{41})(4+\sqrt{41})}$
a_3	$-\frac{1}{8r^6+84r^5+290r^4+315r^3-133r^2-294r-40}$	$-\frac{1}{3240+510\sqrt{41}}$
a_4	$\frac{1}{16r^8+288r^7+2024r^6+6912r^5+11129r^4+4662r^3-7549r^2-7362r-920}$	$\frac{1}{187320+29280\sqrt{41}}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{32r^{10} + 880r^9 + 10160r^8 + 63800r^7 + 234546r^6 + 497255r^5 + 518640r^4 + 28325r^3 - 443678r^2 - 3}$$

Which for the root $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$ becomes

$$a_5 = -\frac{1}{600(1561 + 244\sqrt{41})(10 + \sqrt{41})}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2-\sqrt{41}}$
a_2	$\frac{1}{4r^4+20r^3+15r^2-25r-4}$	$\frac{1}{2(2+\sqrt{41})(4+\sqrt{41})}$
a_3	$-\frac{1}{8r^6+84r^5+290r^4+315r^3-133r^2-294r-40}$	$-\frac{1}{3240+510\sqrt{41}}$
a_4	$\frac{1}{16r^8+288r^7+2024r^6+6912r^5+11129r^4+4662r^3-7549r^2-7362r-920}$	$\frac{1}{187320+29280\sqrt{41}}$
a_5	$-\frac{1}{32r^{10}+880r^9+10160r^8+63800r^7+234546r^6+497255r^5+518640r^4+28325r^3-443678r^2-311960r-36800}$	$-\frac{1}{600(1561+244\sqrt{41})}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x^{\frac{1}{4} + \frac{\sqrt{41}}{4}} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= x^{\frac{1}{4} + \frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2 - \sqrt{41}} + \frac{x^2}{2(2 + \sqrt{41})(4 + \sqrt{41})} - \frac{x^3}{3240 + 510\sqrt{41}} + \frac{x^4}{187320 + 29280\sqrt{41}} - \dots \right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) + b_n(n+r) + b_{n-1} - 5b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{b_{n-1}}{2n^2 + 4nr + 2r^2 - n - r - 5} \quad (4)$$

Which for the root $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$ becomes

$$b_n = \frac{b_{n-1}}{n(\sqrt{41} - 2n)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -\frac{1}{2r^2 + 3r - 4}$$

Which for the root $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$ becomes

$$b_1 = \frac{1}{-2 + \sqrt{41}}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2+\sqrt{41}}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{4r^4 + 20r^3 + 15r^2 - 25r - 4}$$

Which for the root $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$ becomes

$$b_2 = \frac{1}{2(-2 + \sqrt{41})(-4 + \sqrt{41})}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2+\sqrt{41}}$
b_2	$\frac{1}{4r^4+20r^3+15r^2-25r-4}$	$\frac{1}{2(-2+\sqrt{41})(-4+\sqrt{41})}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{1}{8r^6 + 84r^5 + 290r^4 + 315r^3 - 133r^2 - 294r - 40}$$

Which for the root $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$ becomes

$$b_3 = \frac{1}{-3240 + 510\sqrt{41}}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2+\sqrt{41}}$
b_2	$\frac{1}{4r^4+20r^3+15r^2-25r-4}$	$\frac{1}{2(-2+\sqrt{41})(-4+\sqrt{41})}$
b_3	$-\frac{1}{8r^6+84r^5+290r^4+315r^3-133r^2-294r-40}$	$\frac{1}{-3240+510\sqrt{41}}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{16r^8 + 288r^7 + 2024r^6 + 6912r^5 + 11129r^4 + 4662r^3 - 7549r^2 - 7362r - 920}$$

Which for the root $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$ becomes

$$b_4 = \frac{1}{187320 - 29280\sqrt{41}}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2+\sqrt{41}}$
b_2	$\frac{1}{4r^4+20r^3+15r^2-25r-4}$	$\frac{1}{2(-2+\sqrt{41})(-4+\sqrt{41})}$
b_3	$-\frac{1}{8r^6+84r^5+290r^4+315r^3-133r^2-294r-40}$	$\frac{1}{-3240+510\sqrt{41}}$
b_4	$\frac{1}{16r^8+288r^7+2024r^6+6912r^5+11129r^4+4662r^3-7549r^2-7362r-920}$	$\frac{1}{187320-29280\sqrt{41}}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{1}{32r^{10} + 880r^9 + 10160r^8 + 63800r^7 + 234546r^6 + 497255r^5 + 518640r^4 + 28325r^3 - 443678r^2 - 3}$$

Which for the root $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$ becomes

$$b_5 = -\frac{1}{600(-1561 + 244\sqrt{41})(-10 + \sqrt{41})}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2r^2+3r-4}$	$\frac{1}{-2+\sqrt{41}}$
b_2	$\frac{1}{4r^4+20r^3+15r^2-25r-4}$	$\frac{1}{2(-2+\sqrt{41})(-4+\sqrt{41})}$
b_3	$-\frac{1}{8r^6+84r^5+290r^4+315r^3-133r^2-294r-40}$	$\frac{1}{-3240+510\sqrt{41}}$
b_4	$\frac{1}{16r^8+288r^7+2024r^6+6912r^5+11129r^4+4662r^3-7549r^2-7362r-920}$	$\frac{1}{187320-29280\sqrt{41}}$
b_5	$-\frac{1}{32r^{10}+880r^9+10160r^8+63800r^7+234546r^6+497255r^5+518640r^4+28325r^3-443678r^2-311960r-36800}$	$-\frac{1}{600(-1561+244\sqrt{41})}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
y_2(x) &= x^{\frac{1}{4}+\frac{\sqrt{41}}{4}} (b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
&= x^{\frac{1}{4}-\frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2+\sqrt{41}} + \frac{x^2}{2(-2+\sqrt{41})(-4+\sqrt{41})} + \frac{x^3}{-3240+510\sqrt{41}} + \frac{x^4}{187320-29280\sqrt{41}} \right)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
&= c_1x^{\frac{1}{4}+\frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2-\sqrt{41}} + \frac{x^2}{2(2+\sqrt{41})(4+\sqrt{41})} - \frac{x^3}{3240+510\sqrt{41}} \right. \\
&\quad \left. + \frac{x^4}{187320+29280\sqrt{41}} - \frac{x^5}{600(1561+244\sqrt{41})(10+\sqrt{41})} + O(x^6) \right) \\
&\quad + c_2x^{\frac{1}{4}-\frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2+\sqrt{41}} + \frac{x^2}{2(-2+\sqrt{41})(-4+\sqrt{41})} + \frac{x^3}{-3240+510\sqrt{41}} \right. \\
&\quad \left. + \frac{x^4}{187320-29280\sqrt{41}} - \frac{x^5}{600(-1561+244\sqrt{41})(-10+\sqrt{41})} + O(x^6) \right)
\end{aligned}$$

Hence the final solution is

$$y = y_h$$

$$\begin{aligned}
&= c_1 x^{\frac{1}{4} + \frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2 - \sqrt{41}} + \frac{x^2}{2(2 + \sqrt{41})(4 + \sqrt{41})} - \frac{x^3}{3240 + 510\sqrt{41}} \right. \\
&\quad \left. + \frac{x^4}{187320 + 29280\sqrt{41}} - \frac{x^5}{600(1561 + 244\sqrt{41})(10 + \sqrt{41})} + O(x^6) \right) \\
&+ c_2 x^{\frac{1}{4} - \frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2 + \sqrt{41}} + \frac{x^2}{2(-2 + \sqrt{41})(-4 + \sqrt{41})} + \frac{x^3}{-3240 + 510\sqrt{41}} \right. \\
&\quad \left. + \frac{x^4}{187320 - 29280\sqrt{41}} - \frac{x^5}{600(-1561 + 244\sqrt{41})(-10 + \sqrt{41})} + O(x^6) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
y &= c_1 x^{\frac{1}{4} + \frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2 - \sqrt{41}} + \frac{x^2}{2(2 + \sqrt{41})(4 + \sqrt{41})} - \frac{x^3}{3240 + 510\sqrt{41}} \right. \\
&\quad \left. + \frac{x^4}{187320 + 29280\sqrt{41}} - \frac{x^5}{600(1561 + 244\sqrt{41})(10 + \sqrt{41})} + O(x^6) \right) \\
&+ c_2 x^{\frac{1}{4} - \frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2 + \sqrt{41}} + \frac{x^2}{2(-2 + \sqrt{41})(-4 + \sqrt{41})} + \frac{x^3}{-3240 + 510\sqrt{41}} \right. \\
&\quad \left. + \frac{x^4}{187320 - 29280\sqrt{41}} - \frac{x^5}{600(-1561 + 244\sqrt{41})(-10 + \sqrt{41})} + O(x^6) \right)
\end{aligned}$$

Verification of solutions

$$\begin{aligned}
y &= c_1 x^{\frac{1}{4} + \frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2 - \sqrt{41}} + \frac{x^2}{2(2 + \sqrt{41})(4 + \sqrt{41})} - \frac{x^3}{3240 + 510\sqrt{41}} \right. \\
&\quad \left. + \frac{x^4}{187320 + 29280\sqrt{41}} - \frac{x^5}{600(1561 + 244\sqrt{41})(10 + \sqrt{41})} + O(x^6) \right) \\
&+ c_2 x^{\frac{1}{4} - \frac{\sqrt{41}}{4}} \left(1 + \frac{x}{-2 + \sqrt{41}} + \frac{x^2}{2(-2 + \sqrt{41})(-4 + \sqrt{41})} + \frac{x^3}{-3240 + 510\sqrt{41}} \right. \\
&\quad \left. + \frac{x^4}{187320 - 29280\sqrt{41}} - \frac{x^5}{600(-1561 + 244\sqrt{41})(-10 + \sqrt{41})} + O(x^6) \right)
\end{aligned}$$

Verified OK.

4.42.1 Maple step by step solution

Let's solve

$$2x^2y'' + xy' + (x - 5)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{2x} - \frac{(x-5)y}{2x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{2x} + \frac{(x-5)y}{2x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{2x}, P_3(x) = \frac{x-5}{2x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{1}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{5}{2}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$2x^2y'' + xy' + (x - 5)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(2r^2 - r - 5) x^r + \left(\sum_{k=1}^{\infty} (a_k(2k^2 + 4kr + 2r^2 - k - r - 5) + a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$2r^2 - r - 5 = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ \frac{1}{4} - \frac{\sqrt{41}}{4}, \frac{1}{4} + \frac{\sqrt{41}}{4} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$(2k^2 + (4r - 1)k + 2r^2 - r - 5) a_k + a_{k-1} = 0$$

- Shift index using $k- > k + 1$

$$(2(k+1)^2 + (4r - 1)(k+1) + 2r^2 - r - 5) a_{k+1} + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{2k^2 + 4kr + 2r^2 + 3k + 3r - 4}$$

- Recursion relation for $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$

$$a_{k+1} = -\frac{a_k}{2k^2 + 4k\left(\frac{1}{4} - \frac{\sqrt{41}}{4}\right) + 2\left(\frac{1}{4} - \frac{\sqrt{41}}{4}\right)^2 + 3k - \frac{13}{4} - \frac{3\sqrt{41}}{4}}$$

- Solution for $r = \frac{1}{4} - \frac{\sqrt{41}}{4}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k + \frac{1}{4} - \frac{\sqrt{41}}{4}}, a_{k+1} = -\frac{a_k}{2k^2 + 4k\left(\frac{1}{4} - \frac{\sqrt{41}}{4}\right) + 2\left(\frac{1}{4} - \frac{\sqrt{41}}{4}\right)^2 + 3k - \frac{13}{4} - \frac{3\sqrt{41}}{4}} \right]$$

- Recursion relation for $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$

$$a_{k+1} = -\frac{a_k}{2k^2 + 4k\left(\frac{1}{4} + \frac{\sqrt{41}}{4}\right) + 2\left(\frac{1}{4} + \frac{\sqrt{41}}{4}\right)^2 + 3k - \frac{13}{4} + \frac{3\sqrt{41}}{4}}$$

- Solution for $r = \frac{1}{4} + \frac{\sqrt{41}}{4}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{4}+\frac{\sqrt{41}}{4}}, a_{k+1} = -\frac{a_k}{2k^2+4k\left(\frac{1}{4}+\frac{\sqrt{41}}{4}\right)+2\left(\frac{1}{4}+\frac{\sqrt{41}}{4}\right)^2+3k-\frac{13}{4}+\frac{3\sqrt{41}}{4}} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+\frac{1}{4}+\frac{\sqrt{41}}{4}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{4}+\frac{\sqrt{41}}{4}} \right), a_{k+1} = -\frac{a_k}{2k^2+4k\left(\frac{1}{4}-\frac{\sqrt{41}}{4}\right)+2\left(\frac{1}{4}-\frac{\sqrt{41}}{4}\right)^2+3k-\frac{13}{4}-\frac{3\sqrt{41}}{4}}, b_{k+1} = -\frac{b_k}{2k^2+4k\left(\frac{1}{4}+\frac{\sqrt{41}}{4}\right)+2\left(\frac{1}{4}+\frac{\sqrt{41}}{4}\right)^2+3k-\frac{13}{4}+\frac{3\sqrt{41}}{4}} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 665

Order:=6;

dsolve(2*x^2*diff(y(x), x, x) +x*diff(y(x), x) +(x-5)*y(x) = 0,y(x),type='series',x=0);

$$\begin{aligned}
 y(x) = & x^{\frac{1}{4}} \left(c_1 x^{-\frac{\sqrt{41}}{4}} \left(1 + \frac{1}{-2 + \sqrt{41}} x + \frac{1}{2} \frac{1}{(-2 + \sqrt{41})(-4 + \sqrt{41})} x^2 \right. \right. \\
 & + \frac{1}{6} \frac{1}{(-2 + \sqrt{41})(-4 + \sqrt{41})(-6 + \sqrt{41})} x^3 \\
 & + \frac{1}{24} \frac{1}{(-2 + \sqrt{41})(-4 + \sqrt{41})(-6 + \sqrt{41})(-8 + \sqrt{41})} x^4 \\
 & + \frac{1}{120} \frac{1}{(-2 + \sqrt{41})(-4 + \sqrt{41})(-6 + \sqrt{41})(-8 + \sqrt{41})(-10 + \sqrt{41})} x^5 \\
 & \left. \left. + O(x^6) \right) + c_2 x^{\frac{\sqrt{41}}{4}} \left(1 + \frac{1}{-2 - \sqrt{41}} x + \frac{1}{2} \frac{1}{(2 + \sqrt{41})(4 + \sqrt{41})} x^2 \right. \right. \\
 & - \frac{1}{6} \frac{1}{(2 + \sqrt{41})(4 + \sqrt{41})(6 + \sqrt{41})} x^3 \\
 & + \frac{1}{24} \frac{1}{(2 + \sqrt{41})(4 + \sqrt{41})(6 + \sqrt{41})(8 + \sqrt{41})} x^4 \\
 & \left. \left. - \frac{1}{120} \frac{1}{(2 + \sqrt{41})(4 + \sqrt{41})(6 + \sqrt{41})(8 + \sqrt{41})(10 + \sqrt{41})} x^5 + O(x^6) \right) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 1668

AsymptoticDSolveValue[2*x^2*y''[x]+x*y'[x]+(x-5)*y[x]==0,y[x],{x,0,5}]

Too large to display

4.43 problem 40

Internal problem ID [7264]

Internal file name [OUTPUT/6250_Sunday_June_05_2022_04_35_36_PM_10600353/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 40.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' + 2xy' - yx = \sin(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + 2xy' - yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{2x}$$

Table 205: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + 2xy' - yx = \sin(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' + 2xy' - yx = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & 2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + 2x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + 2x^{n+r} a_n (n+r) = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r (-1+r) + 2x^r a_0 r = 0$$

Or

$$(2x^r r (-1+r) + 2x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$2x^r r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m (-1+m) + 2x^m m) c_0 = \sin(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$2x^r r^2 = 0$$

Solving for r gives the roots of the indicial equation as $r = 0$ and $r = -1$. Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + 2a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{2n^2 + 4nr + 2r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{2n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{2(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4(r+1)^2(2+r)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{16}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{288}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{9216}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{460800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$
a_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned}
y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\
&= 1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)
\end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$	$-\frac{1}{(r+1)^3}$	-1
b_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$	$\frac{-3-2r}{2(r+1)^3(2+r)^3}$	$-\frac{3}{16}$
b_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$	$\frac{-3r^2-12r-11}{4(r+1)^3(2+r)^3(r+3)^3}$	$-\frac{11}{864}$
b_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$	$\frac{-2r^3-15r^2-35r-25}{4(r+1)^3(2+r)^3(r+3)^3(r+4)^3}$	$-\frac{25}{55296}$
b_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$	$\frac{-5r^4-60r^3-255r^2-450r-274}{16(r+1)^3(2+r)^3(r+3)^3(r+4)^3(r+5)^3}$	$-\frac{137}{13824000}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \ln(x) \\ &\quad - x - \frac{3x^2}{16} - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = \frac{a_0}{2(r+1)^2}$
$a_2 = \frac{a_0}{4(r+1)^2(2+r)^2}$
$a_3 = \frac{a_0}{8(r+1)^2(2+r)^2(r+3)^2}$
$a_4 = \frac{a_0}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$
$a_5 = \frac{a_0}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$

Expanding the rhs of the ode $\sin(x)$ in series gives

$$\sin(x) = x - \frac{1}{6}x^3 + \frac{1}{120}x^5$$

Since the $F = x - \frac{1}{6}x^3 + \frac{1}{120}x^5$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = x$$

For c_0 and x . This results in

$$c_0 = \frac{1}{2}$$

$$m = 1$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+1}$$

Where in the above $c_0 = \frac{1}{2}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 1$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{2}$ and $r = m$ or $r = 1$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{2}$
$c_1 = \frac{1}{16}$
$c_2 = \frac{1}{288}$
$c_3 = \frac{1}{9216}$
$c_4 = \frac{1}{460800}$
$c_5 = \frac{1}{33177600}$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= x \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x \left(\frac{1}{2} + \frac{1}{16}x + \frac{1}{288}x^2 + \frac{1}{9216}x^3 + \frac{1}{460800}x^4 + \frac{1}{33177600}x^5 \right) \\ &= \frac{1}{2}x + \frac{1}{16}x^2 + \frac{1}{288}x^3 + \frac{1}{9216}x^4 + \frac{1}{460800}x^5 + \frac{1}{33177600}x^6 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{x^3}{6}$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = -\frac{x^3}{6}$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= -\frac{1}{108} \\ m &= 3 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+3} \end{aligned}$$

Where in the above $c_0 = -\frac{1}{108}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 3$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{108}$ and $r = m$ or $r = 3$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$$\begin{aligned}
c_0 &= -\frac{1}{108} \\
c_1 &= -\frac{1}{3456} \\
c_2 &= -\frac{1}{172800} \\
c_3 &= -\frac{1}{12441600} \\
c_4 &= -\frac{1}{1219276800} \\
c_5 &= -\frac{1}{156067430400}
\end{aligned}$$

The particular solution is now found using

$$\begin{aligned}
y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
&= x^3 \sum_{n=0}^{\infty} c_n x^n
\end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
y_p &= x^3 \left(-\frac{1}{108} - \frac{1}{3456}x - \frac{1}{172800}x^2 - \frac{1}{12441600}x^3 - \frac{1}{1219276800}x^4 - \frac{1}{156067430400}x^5 \right) \\
&= -\frac{1}{108}x^3 - \frac{1}{3456}x^4 - \frac{1}{172800}x^5 - \frac{1}{12441600}x^6 - \frac{1}{1219276800}x^7 - \frac{1}{156067430400}x^8
\end{aligned}$$

Now we determine the particular solution y_p associated with $F = \frac{x^5}{120}$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = \frac{x^5}{120}$$

For c_0 and x . This results in

$$\begin{aligned}
c_0 &= \frac{1}{6000} \\
m &= 5
\end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
&= \sum_{n=0}^{\infty} c_n x^{n+5}
\end{aligned}$$

Where in the above $c_0 = \frac{1}{6000}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 5$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{6000}$ and $r = m$ or $r = 5$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{6000}$
$c_1 = \frac{1}{432000}$
$c_2 = \frac{1}{42336000}$
$c_3 = \frac{1}{5419008000}$
$c_4 = \frac{1}{877879296000}$
$c_5 = \frac{1}{175575859200000}$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^5 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$y_p = x^5 \left(\frac{1}{6000} + \frac{1}{432000}x + \frac{1}{42336000}x^2 + \frac{1}{5419008000}x^3 + \frac{1}{877879296000}x^4 + \frac{1}{175575859200000}x^5 \right)$$

$$= \frac{1}{6000}x^5 + \frac{1}{432000}x^6 + \frac{1}{42336000}x^7 + \frac{1}{5419008000}x^8 + \frac{1}{877879296000}x^9 + \frac{1}{175575859200000}x^{10}$$

Adding all the above particular solution(s) gives

$$y_p = \frac{x}{2} + \frac{x^2}{16} - \frac{5x^3}{864} - \frac{5x^4}{27648} + \frac{1127x^5}{6912000} + \frac{1127x^6}{497664000} + \frac{139x^7}{6096384000}$$

$$+ \frac{139x^8}{780337152000} + \frac{x^9}{877879296000} + \frac{x^{10}}{175575859200000} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{x}{2} + \frac{x^2}{16} - \frac{5x^3}{864} - \frac{5x^4}{27648} + \frac{1127x^5}{6912000} + O(x^6)$$

Hence the final solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \frac{x}{2} + \frac{x^2}{16} - \frac{5x^3}{864} - \frac{5x^4}{27648} + \frac{1127x^5}{6912000} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \frac{x}{2} + \frac{x^2}{16} - \frac{5x^3}{864} - \frac{5x^4}{27648} + \frac{1127x^5}{6912000} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} \right. \\ &\quad \left. - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y &= \frac{x}{2} + \frac{x^2}{16} - \frac{5x^3}{864} - \frac{5x^4}{27648} + \frac{1127x^5}{6912000} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 77

```
Order:=6;
```

```
dsolve(2*x^2*diff(y(x), x, x) + 2*x*diff(y(x), x) - x*y(x) = sin(x),y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 + \frac{1}{2}x + \frac{1}{16}x^2 + \frac{1}{288}x^3 + \frac{1}{9216}x^4 + \frac{1}{460800}x^5 + O(x^6) \right) \\ + x \left(\frac{1}{2} + \frac{1}{16}x - \frac{5}{864}x^2 - \frac{5}{27648}x^3 + \frac{1127}{6912000}x^4 + \frac{1127}{497664000}x^5 + O(x^6) \right) \\ + \left(-x - \frac{3}{16}x^2 - \frac{11}{864}x^3 - \frac{25}{55296}x^4 - \frac{137}{13824000}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.156 (sec). Leaf size: 340

AsymptoticDSolveValue[2*x^2*y'[x]+2*x*y'[x]-x*y[x]==Sin[x],y[x],{x,0,5}]

$$\begin{aligned}
 y(x) \rightarrow & c_2 \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \\
 & + c_1 \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) \right. \\
 & \quad + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \left. \right) + \left(\frac{4963x^6}{16588800} - \frac{91x^5}{460800} \right. \\
 & \quad - \frac{23x^4}{2304} - \frac{5x^3}{288} + \frac{x^2}{8} + \frac{x}{2} \left. \right) \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) \right. \\
 & \quad \left. + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \right) \\
 & + \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \left(\frac{x^6(66968 - 74445 \log(x))}{248832000} \right. \\
 & \quad + \frac{13x^5(210 \log(x) - 3107)}{13824000} + \frac{x^4(276 \log(x) - 325)}{27648} + \frac{1}{864} x^3(15 \log(x) + 37) \\
 & \quad \left. + \frac{1}{16} x^2(3 - 2 \log(x)) - \frac{1}{2} x \log(x) \right)
 \end{aligned}$$

4.44 problem 41

Internal problem ID [7265]

Internal file name [OUTPUT/6251_Sunday_June_05_2022_04_35_37_PM_8662565/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 41.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' + 2xy' - yx = x \sin(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + 2xy' - yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{2x}$$

Table 206: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + 2xy' - yx = x \sin(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' + 2xy' - yx = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & 2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + 2x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + 2x^{n+r} a_n (n+r) = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r (-1+r) + 2x^r a_0 r = 0$$

Or

$$(2x^r r (-1+r) + 2x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$2x^r r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m (-1+m) + 2x^m m) c_0 = x \sin(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$2x^r r^2 = 0$$

Solving for r gives the roots of the indicial equation as $r = 0$. Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + 2a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{2n^2 + 4nr + 2r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{2n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{2(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4(r+1)^2(2+r)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{16}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{288}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{9216}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{460800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$
a_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned}
 y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\
 &= 1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)
 \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$	$-\frac{1}{(r+1)^3}$	-1
b_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$	$\frac{-3-2r}{2(r+1)^3(2+r)^3}$	$-\frac{3}{16}$
b_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$	$\frac{-3r^2-12r-11}{4(r+1)^3(2+r)^3(r+3)^3}$	$-\frac{11}{864}$
b_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$	$\frac{-2r^3-15r^2-35r-25}{4(r+1)^3(2+r)^3(r+3)^3(r+4)^3}$	$-\frac{25}{55296}$
b_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$	$\frac{-5r^4-60r^3-255r^2-450r-274}{16(r+1)^3(2+r)^3(r+3)^3(r+4)^3(r+5)^3}$	$-\frac{137}{13824000}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \ln(x) \\ &\quad - x - \frac{3x^2}{16} - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = \frac{a_0}{2(r+1)^2}$
$a_2 = \frac{a_0}{4(r+1)^2(2+r)^2}$
$a_3 = \frac{a_0}{8(r+1)^2(2+r)^2(r+3)^2}$
$a_4 = \frac{a_0}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$
$a_5 = \frac{a_0}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$

Expanding the rhs of the ode $x \sin(x)$ in series gives

$$x \sin(x) = x^2 - \frac{1}{6}x^4$$

Since the $F = x^2 - \frac{1}{6}x^4$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x^2$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = x^2$$

For c_0 and x . This results in

$$c_0 = \frac{1}{8}$$

$$m = 2$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+2}$$

Where in the above $c_0 = \frac{1}{8}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{8}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{8}$
$c_1 = \frac{1}{144}$
$c_2 = \frac{1}{4608}$
$c_3 = \frac{1}{230400}$
$c_4 = \frac{1}{16588800}$
$c_5 = \frac{1}{1625702400}$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= x^2 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= x^2 \left(\frac{1}{8} + \frac{1}{144}x + \frac{1}{4608}x^2 + \frac{1}{230400}x^3 + \frac{1}{16588800}x^4 + \frac{1}{1625702400}x^5 \right) \\
 &= \frac{1}{8}x^2 + \frac{1}{144}x^3 + \frac{1}{4608}x^4 + \frac{1}{230400}x^5 + \frac{1}{16588800}x^6 + \frac{1}{1625702400}x^7
 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{x^4}{6}$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = -\frac{x^4}{6}$$

For c_0 and x . This results in

$$\begin{aligned}
 c_0 &= -\frac{1}{192} \\
 m &= 4
 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
 y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
 &= \sum_{n=0}^{\infty} c_n x^{n+4}
 \end{aligned}$$

Where in the above $c_0 = -\frac{1}{192}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 4$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{192}$ and $r = m$ or $r = 4$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$$\begin{aligned}
c_0 &= -\frac{1}{192} \\
c_1 &= -\frac{1}{9600} \\
c_2 &= -\frac{1}{691200} \\
c_3 &= -\frac{1}{67737600} \\
c_4 &= -\frac{1}{8670412800} \\
c_5 &= -\frac{1}{1404606873600}
\end{aligned}$$

The particular solution is now found using

$$\begin{aligned}
y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
&= x^4 \sum_{n=0}^{\infty} c_n x^n
\end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
y_p &= x^4 \left(-\frac{1}{192} - \frac{1}{9600}x - \frac{1}{691200}x^2 - \frac{1}{67737600}x^3 - \frac{1}{8670412800}x^4 - \frac{1}{1404606873600}x^5 \right) \\
&= -\frac{1}{192}x^4 - \frac{1}{9600}x^5 - \frac{1}{691200}x^6 - \frac{1}{67737600}x^7 - \frac{1}{8670412800}x^8 - \frac{1}{1404606873600}x^9
\end{aligned}$$

Adding all the above particular solution(s) gives

$$\begin{aligned}
y_p &= \frac{x^2}{8} + \frac{x^3}{144} - \frac{23x^4}{4608} - \frac{23x^5}{230400} - \frac{23x^6}{16588800} - \frac{23x^7}{1625702400} \\
&\quad - \frac{1}{8670412800} - \frac{1}{1404606873600} + O(x^6)
\end{aligned}$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{x^2}{8} + \frac{x^3}{144} - \frac{23x^4}{4608} - \frac{23x^5}{230400} + O(x^6)$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \frac{x^2}{8} + \frac{x^3}{144} - \frac{23x^4}{4608} - \frac{23x^5}{230400} + O(x^6) + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\
 &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= \frac{x^2}{8} + \frac{x^3}{144} - \frac{23x^4}{4608} - \frac{23x^5}{230400} + O(x^6) \\
 &\quad + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} \right. \\
 &\quad \left. - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= \frac{x^2}{8} + \frac{x^3}{144} - \frac{23x^4}{4608} - \frac{23x^5}{230400} + O(x^6) + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\
 &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\
 &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right)
 \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    trying a symmetry of the form [xi=0, eta=F(x)]  
    checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
    -> Bessel  
        <- Bessel successful  
    <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 75

```
Order:=6;
```

```
dsolve(2*x^2*diff(y(x), x, x) + 2*x*diff(y(x), x) - x*y(x) = x*sin(x), y(x), type='series', x=0
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 + \frac{1}{2}x + \frac{1}{16}x^2 + \frac{1}{288}x^3 + \frac{1}{9216}x^4 + \frac{1}{460800}x^5 + O(x^6) \right) \\ + x^2 \left(\frac{1}{8} + \frac{1}{144}x - \frac{23}{4608}x^2 - \frac{23}{230400}x^3 + O(x^4) \right) \\ + \left(-x - \frac{3}{16}x^2 - \frac{11}{864}x^3 - \frac{25}{55296}x^4 - \frac{137}{13824000}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.186 (sec). Leaf size: 328

AsymptoticDSolveValue[2*x^2*y'[x]+2*x*y'[x]-x*y[x]==x*Sin[x],y[x],{x,0,5}]

$$\begin{aligned}
 y(x) \rightarrow & c_2 \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \\
 & + c_1 \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) \right. \\
 & + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \left. \right) + \left(-\frac{91x^6}{552960} - \frac{23x^5}{2880} \right. \\
 & - \frac{5x^4}{384} + \frac{x^3}{12} + \frac{x^2}{4} \left. \right) \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) \right. \\
 & + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \left. \right) \\
 & + \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \left(\frac{13x^6(21 \log(x) - 310)}{1658880} \right. \\
 & + \frac{x^5(345 \log(x) - 389)}{43200} + \frac{x^4(20 \log(x) + 51)}{1536} + \frac{1}{36}x^3(4 - 3 \log(x)) \\
 & \left. + \frac{1}{8}x^2(-2 \log(x) - 1) \right)
 \end{aligned}$$

4.45 problem 42

Internal problem ID [7266]

Internal file name [OUTPUT/6252_Sunday_June_05_2022_04_35_40_PM_59085868/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 42.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' + 2xy' - yx = \sin(x) \cos(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + 2xy' - yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{2x}$$

Table 207: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : [0]

Irregular singular points : [∞]

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + 2xy' - yx = \frac{\sin(2x)}{2}$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' + 2xy' - yx = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & 2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + 2x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + 2x^{n+r} a_n (n+r) = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r (-1+r) + 2x^r a_0 r = 0$$

Or

$$(2x^r r (-1+r) + 2x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$2x^r r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m (-1+m) + 2x^m m) c_0 = \frac{\sin(2x)}{2}$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$2x^r r^2 = 0$$

Solving for r gives the roots of the indicial equation as $r = 0$ and $r = -1$. Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + 2a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{2n^2 + 4nr + 2r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{2n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{2(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4(r+1)^2(2+r)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{16}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{288}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{9216}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{460800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$
a_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned}
 y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\
 &= 1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)
 \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$	$-\frac{1}{(r+1)^3}$	-1
b_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$	$\frac{-3-2r}{2(r+1)^3(2+r)^3}$	$-\frac{3}{16}$
b_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$	$\frac{-3r^2-12r-11}{4(r+1)^3(2+r)^3(r+3)^3}$	$-\frac{11}{864}$
b_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$	$\frac{-2r^3-15r^2-35r-25}{4(r+1)^3(2+r)^3(r+3)^3(r+4)^3}$	$-\frac{25}{55296}$
b_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$	$\frac{-5r^4-60r^3-255r^2-450r-274}{16(r+1)^3(2+r)^3(r+3)^3(r+4)^3(r+5)^3}$	$-\frac{137}{13824000}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \ln(x) \\ &\quad - x - \frac{3x^2}{16} - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = \frac{a_0}{2(r+1)^2}$
$a_2 = \frac{a_0}{4(r+1)^2(2+r)^2}$
$a_3 = \frac{a_0}{8(r+1)^2(2+r)^2(r+3)^2}$
$a_4 = \frac{a_0}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$
$a_5 = \frac{a_0}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$

Expanding the rhs of the ode $\frac{\sin(2x)}{2}$ in series gives

$$\frac{\sin(2x)}{2} = x - \frac{2}{3}x^3 + \frac{2}{15}x^5$$

Since the $F = x - \frac{2}{3}x^3 + \frac{2}{15}x^5$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = x$$

For c_0 and x . This results in

$$c_0 = \frac{1}{2}$$

$$m = 1$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+1}$$

Where in the above $c_0 = \frac{1}{2}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 1$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{2}$ and $r = m$ or $r = 1$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{2}$
$c_1 = \frac{1}{16}$
$c_2 = \frac{1}{288}$
$c_3 = \frac{1}{9216}$
$c_4 = \frac{1}{460800}$
$c_5 = \frac{1}{33177600}$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= x \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= x \left(\frac{1}{2} + \frac{1}{16}x + \frac{1}{288}x^2 + \frac{1}{9216}x^3 + \frac{1}{460800}x^4 + \frac{1}{33177600}x^5 \right) \\
 &= \frac{1}{2}x + \frac{1}{16}x^2 + \frac{1}{288}x^3 + \frac{1}{9216}x^4 + \frac{1}{460800}x^5 + \frac{1}{33177600}x^6
 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{2x^3}{3}$ by solving the balance equation

$$(2x^m m(-1+m) + 2x^m m) c_0 = -\frac{2x^3}{3}$$

For c_0 and x . This results in

$$\begin{aligned}
 c_0 &= -\frac{1}{27} \\
 m &= 3
 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
 y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
 &= \sum_{n=0}^{\infty} c_n x^{n+3}
 \end{aligned}$$

Where in the above $c_0 = -\frac{1}{27}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 3$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{27}$ and $r = m$ or $r = 3$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$$\begin{aligned}
c_0 &= -\frac{1}{27} \\
c_1 &= -\frac{1}{864} \\
c_2 &= -\frac{1}{43200} \\
c_3 &= -\frac{1}{3110400} \\
c_4 &= -\frac{1}{304819200} \\
c_5 &= -\frac{1}{39016857600}
\end{aligned}$$

The particular solution is now found using

$$\begin{aligned}
y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
&= x^3 \sum_{n=0}^{\infty} c_n x^n
\end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
y_p &= x^3 \left(-\frac{1}{27} - \frac{1}{864}x - \frac{1}{43200}x^2 - \frac{1}{3110400}x^3 - \frac{1}{304819200}x^4 - \frac{1}{39016857600}x^5 \right) \\
&= -\frac{1}{27}x^3 - \frac{1}{864}x^4 - \frac{1}{43200}x^5 - \frac{1}{3110400}x^6 - \frac{1}{304819200}x^7 - \frac{1}{39016857600}x^8
\end{aligned}$$

Now we determine the particular solution y_p associated with $F = \frac{2x^5}{15}$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = \frac{2x^5}{15}$$

For c_0 and x . This results in

$$\begin{aligned}
c_0 &= \frac{1}{375} \\
m &= 5
\end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
&= \sum_{n=0}^{\infty} c_n x^{n+5}
\end{aligned}$$

Where in the above $c_0 = \frac{1}{375}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 5$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{375}$ and $r = m$ or $r = 5$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{375}$
$c_1 = \frac{1}{27000}$
$c_2 = \frac{1}{2646000}$
$c_3 = \frac{1}{338688000}$
$c_4 = \frac{1}{54867456000}$
$c_5 = \frac{1}{10973491200000}$

The particular solution is now found using

$$\begin{aligned}
 y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
 &= x^5 \sum_{n=0}^{\infty} c_n x^n
 \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
 y_p &= x^5 \left(\frac{1}{375} + \frac{1}{27000}x + \frac{1}{2646000}x^2 + \frac{1}{338688000}x^3 + \frac{1}{54867456000}x^4 \right. \\
 &\quad \left. + \frac{1}{10973491200000}x^5 \right) \\
 &= \frac{1}{375}x^5 + \frac{1}{27000}x^6 + \frac{1}{2646000}x^7 + \frac{1}{338688000}x^8 + \frac{1}{54867456000}x^9 + \frac{1}{10973491200000}x^{10}
 \end{aligned}$$

Adding all the above particular solution(s) gives

$$\begin{aligned}
 y_p &= \frac{x}{2} + \frac{x^2}{16} - \frac{29x^3}{864} - \frac{29x^4}{27648} + \frac{18287x^5}{6912000} + \frac{18287x^6}{497664000} + \frac{571x^7}{1524096000} \\
 &\quad + \frac{571x^8}{195084288000} + \frac{x^9}{54867456000} + \frac{x^{10}}{10973491200000} + O(x^6)
 \end{aligned}$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{x}{2} + \frac{x^2}{16} - \frac{29x^3}{864} - \frac{29x^4}{27648} + \frac{18287x^5}{6912000} + O(x^6)$$

Hence the final solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \frac{x}{2} + \frac{x^2}{16} - \frac{29x^3}{864} - \frac{29x^4}{27648} + \frac{18287x^5}{6912000} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \frac{x}{2} + \frac{x^2}{16} - \frac{29x^3}{864} - \frac{29x^4}{27648} + \frac{18287x^5}{6912000} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} \right. \\ &\quad \left. - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y &= \frac{x}{2} + \frac{x^2}{16} - \frac{29x^3}{864} - \frac{29x^4}{27648} + \frac{18287x^5}{6912000} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
    <- No Liouvillian solutions exists
    -> Trying a solution in terms of special functions:
        -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 77

```
Order:=6;
dsolve(2*x^2*diff(y(x), x, x) + 2*x*diff(y(x), x) - x*y(x) = cos(x)*sin(x),y(x),type='series
```

$$\begin{aligned} y(x) = & (c_2 \ln(x) + c_1) \left(1 + \frac{1}{2}x + \frac{1}{16}x^2 + \frac{1}{288}x^3 + \frac{1}{9216}x^4 + \frac{1}{460800}x^5 + O(x^6) \right) \\ & + x \left(\frac{1}{2} + \frac{1}{16}x - \frac{29}{864}x^2 - \frac{29}{27648}x^3 + \frac{18287}{6912000}x^4 + \frac{18287}{497664000}x^5 + O(x^6) \right) \\ & + \left(-x - \frac{3}{16}x^2 - \frac{11}{864}x^3 - \frac{25}{55296}x^4 - \frac{137}{13824000}x^5 + O(x^6) \right) c_2 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.157 (sec). Leaf size: 340

AsymptoticDSolveValue[2*x^2*y'[x]+2*x*y'[x]-x*y[x]==Cos[x]*Sin[x],y[x],{x,0,5}]

$$\begin{aligned}
 y(x) \rightarrow & c_2 \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \\
 & + c_1 \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) \right. \\
 & \quad + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \left. \right) + \left(\frac{88963x^6}{16588800} + \frac{4229x^5}{460800} \right. \\
 & \quad - \frac{95x^4}{2304} - \frac{29x^3}{288} + \frac{x^2}{8} + \frac{x}{2} \left. \right) \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) \right. \\
 & \quad \left. + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \right) \\
 & + \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \left(\frac{x^6(1476968 - 1334445 \log(x))}{248832000} \right. \\
 & \quad + \frac{x^5(-126870 \log(x) - 273671)}{13824000} + \frac{5x^4(228 \log(x) - 281)}{27648} \\
 & \quad \left. + \frac{1}{864}x^3(87 \log(x) + 85) + \frac{1}{16}x^2(3 - 2 \log(x)) - \frac{1}{2}x \log(x) \right)
 \end{aligned}$$

4.46 problem 43

Internal problem ID [7267]

Internal file name [OUTPUT/6253_Sunday_June_05_2022_04_35_42_PM_10931439/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 43.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$2x^2y'' + 2xy' - yx = x^3 + x \sin(x)$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + 2xy' - yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{2x}$$

Table 208: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + 2xy' - yx = x^3 + x \sin(x)$$

Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $2x^2y'' + 2xy' - yx = 0$, and y_p is a particular solution to the inhomogeneous ode. which is found using the balance equation generated from indicial equation

First, we solve for y_h Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & 2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + 2x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + 2x^{n+r} a_n (n+r) = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r (-1+r) + 2x^r a_0 r = 0$$

Or

$$(2x^r r (-1+r) + 2x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$2x^r r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(2x^m m (-1+m) + 2x^m m) c_0 = x^3 + x \sin(x)$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$2x^r r^2 = 0$$

Solving for r gives the roots of the indicial equation as $r = 0$ and $r = -1$. Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + 2a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{2n^2 + 4nr + 2r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{2n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{2(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4(r+1)^2(2+r)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{16}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{288}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{9216}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{460800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$
a_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$
a_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$
a_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$
a_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned}
 y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\
 &= 1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)
 \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{1}{2(r+1)^2}$	$\frac{1}{2}$	$-\frac{1}{(r+1)^3}$	-1
b_2	$\frac{1}{4(r+1)^2(2+r)^2}$	$\frac{1}{16}$	$\frac{-3-2r}{2(r+1)^3(2+r)^3}$	$-\frac{3}{16}$
b_3	$\frac{1}{8(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{288}$	$\frac{-3r^2-12r-11}{4(r+1)^3(2+r)^3(r+3)^3}$	$-\frac{11}{864}$
b_4	$\frac{1}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{9216}$	$\frac{-2r^3-15r^2-35r-25}{4(r+1)^3(2+r)^3(r+3)^3(r+4)^3}$	$-\frac{25}{55296}$
b_5	$\frac{1}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{460800}$	$\frac{-5r^4-60r^3-255r^2-450r-274}{16(r+1)^3(2+r)^3(r+3)^3(r+4)^3(r+5)^3}$	$-\frac{137}{13824000}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \ln(x) \\ &\quad - x - \frac{3x^2}{16} - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6)\right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

The particular solution is found by solving for c, m the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = F$$

Where $F(x)$ is the RHS of the ode. If $F(x)$ has more than one term, then this is done for each term one at a time and then all the particular solutions are added. The function $F(x)$ will be converted to series if needed. in order to solve for c_n, m for each term, the same recursive relation used to find $y_h(x)$ is used to find c_n, m which is used to find the particular solution $\sum_{n=0} c_n x^{n+m}$ by replacing a_n by c_n and r by m .

The following are the values of a_n found in terms of the indicial root r .

$a_1 = \frac{a_0}{2(r+1)^2}$
$a_2 = \frac{a_0}{4(r+1)^2(2+r)^2}$
$a_3 = \frac{a_0}{8(r+1)^2(2+r)^2(r+3)^2}$
$a_4 = \frac{a_0}{16(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$
$a_5 = \frac{a_0}{32(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$

Expanding the rhs of the ode $x^3 + x \sin(x)$ in series gives

$$x^3 + x \sin(x) = x^2 + x^3 - \frac{1}{6}x^4$$

Since the $F = x^2 + x^3 - \frac{1}{6}x^4$ has more than one term then we find a particular solution for each term and add the result to find the particular solution to the ode.

Now we determine the particular solution y_p associated with $F = x^2$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = x^2$$

For c_0 and x . This results in

$$c_0 = \frac{1}{8}$$

$$m = 2$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+2}$$

Where in the above $c_0 = \frac{1}{8}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 2$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{8}$ and $r = m$ or $r = 2$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = \frac{1}{8}$
$c_1 = \frac{1}{144}$
$c_2 = \frac{1}{4608}$
$c_3 = \frac{1}{230400}$
$c_4 = \frac{1}{16588800}$
$c_5 = \frac{1}{1625702400}$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= x^2 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned} y_p &= x^2 \left(\frac{1}{8} + \frac{1}{144}x + \frac{1}{4608}x^2 + \frac{1}{230400}x^3 + \frac{1}{16588800}x^4 + \frac{1}{1625702400}x^5 \right) \\ &= \frac{1}{8}x^2 + \frac{1}{144}x^3 + \frac{1}{4608}x^4 + \frac{1}{230400}x^5 + \frac{1}{16588800}x^6 + \frac{1}{1625702400}x^7 \end{aligned}$$

Now we determine the particular solution y_p associated with $F = x^3$ by solving the balance equation

$$(2x^m m(-1 + m) + 2x^m m) c_0 = x^3$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= \frac{1}{18} \\ m &= 3 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+3} \end{aligned}$$

Where in the above $c_0 = \frac{1}{18}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 3$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = \frac{1}{18}$ and $r = m$ or $r = 3$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$$\begin{aligned}
c_0 &= \frac{1}{18} \\
c_1 &= \frac{1}{576} \\
c_2 &= \frac{1}{28800} \\
c_3 &= \frac{1}{2073600} \\
c_4 &= \frac{1}{203212800} \\
c_5 &= \frac{1}{26011238400}
\end{aligned}$$

The particular solution is now found using

$$\begin{aligned}
y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\
&= x^3 \sum_{n=0}^{\infty} c_n x^n
\end{aligned}$$

Using the values found above for c_n into the above sum gives

$$\begin{aligned}
y_p &= x^3 \left(\frac{1}{18} + \frac{1}{576}x + \frac{1}{28800}x^2 + \frac{1}{2073600}x^3 + \frac{1}{203212800}x^4 + \frac{1}{26011238400}x^5 \right) \\
&= \frac{1}{18}x^3 + \frac{1}{576}x^4 + \frac{1}{28800}x^5 + \frac{1}{2073600}x^6 + \frac{1}{203212800}x^7 + \frac{1}{26011238400}x^8
\end{aligned}$$

Now we determine the particular solution y_p associated with $F = -\frac{x^4}{6}$ by solving the balance equation

$$(2x^m m(-1+m) + 2x^m m) c_0 = -\frac{x^4}{6}$$

For c_0 and x . This results in

$$\begin{aligned}
c_0 &= -\frac{1}{192} \\
m &= 4
\end{aligned}$$

The particular solution is therefore

$$\begin{aligned}
y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\
&= \sum_{n=0}^{\infty} c_n x^{n+4}
\end{aligned}$$

Where in the above $c_0 = -\frac{1}{192}$.

The remaining c_n values are found using the same recurrence relation given in the earlier table which was used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 4$ in place of the root of the indicial equation used to find the homogeneous solution. By letting $a_0 = c_0$ or $a_0 = -\frac{1}{192}$ and $r = m$ or $r = 4$. The following table gives the resulting c_n values. These values will be used to find the particular solution. Values of c_n found not defined when doing the substitution will be discarded and not used

$c_0 = -\frac{1}{192}$
$c_1 = -\frac{1}{9600}$
$c_2 = -\frac{1}{691200}$
$c_3 = -\frac{1}{67737600}$
$c_4 = -\frac{1}{8670412800}$
$c_5 = -\frac{1}{1404606873600}$

The particular solution is now found using

$$y_p = x^m \sum_{n=0}^{\infty} c_n x^n$$

$$= x^4 \sum_{n=0}^{\infty} c_n x^n$$

Using the values found above for c_n into the above sum gives

$$y_p = x^4 \left(-\frac{1}{192} - \frac{1}{9600}x - \frac{1}{691200}x^2 - \frac{1}{67737600}x^3 - \frac{1}{8670412800}x^4 - \frac{1}{1404606873600}x^5 \right)$$

$$= -\frac{1}{192}x^4 - \frac{1}{9600}x^5 - \frac{1}{691200}x^6 - \frac{1}{67737600}x^7 - \frac{1}{8670412800}x^8 - \frac{1}{1404606873600}x^9$$

Adding all the above particular solution(s) gives

$$y_p = \frac{x^2}{8} + \frac{x^3}{16} - \frac{5x^4}{1536} - \frac{x^5}{15360} - \frac{x^6}{1105920} - \frac{x^7}{108380160}$$

$$- \frac{x^8}{13005619200} - \frac{x^9}{1404606873600} + O(x^6)$$

Truncating the particular solution to the order of series requested gives

$$y_p = \frac{x^2}{8} + \frac{x^3}{16} - \frac{5x^4}{1536} - \frac{x^5}{15360} + O(x^6)$$

Hence the final solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \frac{x^2}{8} + \frac{x^3}{16} - \frac{5x^4}{1536} - \frac{x^5}{15360} + O(x^6) + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \frac{x^2}{8} + \frac{x^3}{16} - \frac{5x^4}{1536} - \frac{x^5}{15360} + O(x^6) \\ &\quad + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} \right. \\ &\quad \left. - \frac{11x^3}{864} - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned} y &= \frac{x^2}{8} + \frac{x^3}{16} - \frac{5x^4}{1536} - \frac{x^5}{15360} + O(x^6) + c_1 \left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 + \frac{x}{2} + \frac{x^2}{16} + \frac{x^3}{288} + \frac{x^4}{9216} + \frac{x^5}{460800} + O(x^6) \right) \ln(x) - x - \frac{3x^2}{16} - \frac{11x^3}{864} \right. \\ &\quad \left. - \frac{25x^4}{55296} - \frac{137x^5}{13824000} + O(x^6) \right) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    trying a symmetry of the form [xi=0, eta=F(x)]  
    checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
    -> Bessel  
        <- Bessel successful  
    <- special function solution successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 75

```
Order:=6;
```

```
dsolve(2*x^2*diff(y(x), x, x) + 2*x*diff(y(x), x) - x*y(x) = x^3+x*sin(x),y(x),type='series')
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 + \frac{1}{2}x + \frac{1}{16}x^2 + \frac{1}{288}x^3 + \frac{1}{9216}x^4 + \frac{1}{460800}x^5 + O(x^6) \right) \\ + x^2 \left(\frac{1}{8} + \frac{1}{16}x - \frac{5}{1536}x^2 - \frac{1}{15360}x^3 + O(x^4) \right) \\ + \left(-x - \frac{3}{16}x^2 - \frac{11}{864}x^3 - \frac{25}{55296}x^4 - \frac{137}{13824000}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.296 (sec). Leaf size: 268

```
AsymptoticDSolveValue[2*x^2*y'[x]+2*x*y'[x]-x*y[x]==x^3*x*Sin[x],y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_2 \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \\ & + c_1 \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) \right. \\ & + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \left. \right) + \left(\frac{x^5}{460800} + \frac{x^4}{9216} + \frac{x^3}{288} \right. \\ & \quad \left. + \frac{x^2}{16} + \frac{x}{2} + 1 \right) \left(\frac{1}{144} x^6 (7 - 6 \log(x)) + \frac{1}{50} x^5 (-5 \log(x) - 4) \right) \\ & + \left(\frac{x^6}{24} + \frac{x^5}{10} \right) \left(x^5 \left(\frac{\log(x)}{460800} - \frac{107}{13824000} \right) + x^4 \left(\frac{\log(x)}{9216} - \frac{19}{55296} \right) \right. \\ & \quad \left. + x^3 \left(\frac{\log(x)}{288} - \frac{1}{108} \right) + x^2 \left(\frac{\log(x)}{16} - \frac{1}{8} \right) + x \left(\frac{\log(x)}{2} - \frac{1}{2} \right) + \log(x) + 1 \right) \end{aligned}$$

4.47 problem 44

Internal problem ID [7268]

Internal file name [OUTPUT/6254_Sunday_June_05_2022_04_35_47_PM_92925667/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 44.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' \cos(x) + 2xy' - yx = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (333)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (334)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{x(2y' - y)}{\cos(x)}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= (2 \sec(x) x^2 (2y' - y) + (-2 \tan(x) x + x - 2) y' + y(\tan(x) x + 1)) \sec(x) \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= 4 \sec(x)^3 \left(\left(\frac{\cos(x)^2 (1+x)}{2} + \left((-1 + \frac{x}{2}) \sin(x) - x^2 + 3x \right) \cos(x) - 2x^3 + 3 \sin(x) x^2 - x \right) y' - \right. \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= -8 \sec(x)^4 \left(\left(\left(\frac{3x}{8} - \frac{3}{4} \right) \cos(x)^3 + \left(\left(-\frac{x}{4} - \frac{3}{4} \right) \sin(x) + \frac{27x^2}{8} + \frac{9x}{4} - \frac{3}{2} \right) \cos(x)^2 + \left((-7x + \right. \right. \right. \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= 16 \sec(x)^5 \left(\left(\left((-3 - \frac{x}{2}) \cos(x)^4 + \frac{(-3 + (1 - \frac{x}{2}) \sin(x) + 7x^2 - \frac{87x}{4}) \cos(x)^3}{2} + \left(\left(-\frac{27}{8} x^2 + 5 - \right. \right. \right. \right. \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$F_0 = 0$$

$$F_1 = -2y'(0) + y(0)$$

$$F_2 = 2y'(0)$$

$$F_3 = 6y'(0) - 3y(0)$$

$$F_4 = -12y'(0) + 4y(0)$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 + \frac{1}{6}x^3 - \frac{1}{40}x^5 + \frac{1}{180}x^6\right)y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{20}x^5 - \frac{1}{60}x^6\right)y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -\frac{x \left(2 \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) \right)}{\cos(x)} \quad (1)$$

Expanding $\cos(x)$ as Taylor series around $x = 0$ and keeping only the first 6 terms gives

$$\begin{aligned} \cos(x) &= -\frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 + \dots \\ &= -\frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\begin{aligned} &\left(-\frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2\right) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}\right) \\ &+ 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1}\right) - x \left(\sum_{n=0}^{\infty} a_n x^n\right) = 0 \end{aligned}$$

Expanding the first term in (1) gives

$$\begin{aligned}
& -\frac{x^6}{720} \cdot \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \frac{x^4}{24} \cdot \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + 1 \\
& \cdot \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) - \frac{x^2}{2} \cdot \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) \\
& + 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - x \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0
\end{aligned}$$

Which simplifies to

$$\begin{aligned}
& \sum_{n=2}^{\infty} \left(-\frac{n x^{n+4} a_n (n-1)}{720} \right) + \left(\sum_{n=2}^{\infty} \frac{n x^{n+2} a_n (n-1)}{24} \right) + \sum_{n=2}^{\infty} \left(-\frac{n a_n x^n (n-1)}{2} \right) \quad (2) \\
& + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} 2n a_n x^n \right) + \sum_{n=0}^{\infty} (-x^{1+n} a_n) = 0
\end{aligned}$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=2}^{\infty} \left(-\frac{n x^{n+4} a_n (n-1)}{720} \right) &= \sum_{n=6}^{\infty} \left(-\frac{(n-4) a_{n-4} (n-5) x^n}{720} \right) \\
\sum_{n=2}^{\infty} \frac{n x^{n+2} a_n (n-1)}{24} &= \sum_{n=4}^{\infty} \frac{(n-2) a_{n-2} (n-3) x^n}{24} \\
\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \\
\sum_{n=0}^{\infty} (-x^{1+n} a_n) &= \sum_{n=1}^{\infty} (-a_{n-1} x^n)
\end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers

of x are the same and equal to n .

$$\begin{aligned} & \sum_{n=6}^{\infty} \left(-\frac{(n-4)a_{n-4}(n-5)x^n}{720} \right) + \left(\sum_{n=4}^{\infty} \frac{(n-2)a_{n-2}(n-3)x^n}{24} \right) \\ & + \sum_{n=2}^{\infty} \left(-\frac{na_n x^n (n-1)}{2} \right) + \left(\sum_{n=0}^{\infty} (n+2)a_{n+2}(1+n)x^n \right) \\ & + \left(\sum_{n=1}^{\infty} 2na_n x^n \right) + \sum_{n=1}^{\infty} (-a_{n-1}x^n) = 0 \end{aligned} \quad (3)$$

$n = 1$ gives

$$6a_3 + 2a_1 - a_0 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = \frac{a_0}{6} - \frac{a_1}{3}$$

$n = 2$ gives

$$3a_2 + 12a_4 - a_1 = 0$$

Which after substituting earlier equations, simplifies to

$$a_4 = \frac{a_1}{12}$$

$n = 3$ gives

$$3a_3 + 20a_5 - a_2 = 0$$

Which after substituting earlier equations, simplifies to

$$\frac{a_0}{2} - a_1 + 20a_5 = 0$$

Or

$$a_5 = -\frac{a_0}{40} + \frac{a_1}{20}$$

$n = 4$ gives

$$\frac{a_2}{12} + 2a_4 + 30a_6 - a_3 = 0$$

Which after substituting earlier equations, simplifies to

$$\frac{a_1}{2} + 30a_6 - \frac{a_0}{6} = 0$$

Or

$$a_6 = \frac{a_0}{180} - \frac{a_1}{60}$$

$n = 5$ gives

$$\frac{a_3}{4} + 42a_7 - a_4 = 0$$

Which after substituting earlier equations, simplifies to

$$\frac{a_0}{24} - \frac{a_1}{6} + 42a_7 = 0$$

Or

$$a_7 = -\frac{a_0}{1008} + \frac{a_1}{252}$$

For $6 \leq n$, the recurrence equation is

$$\begin{aligned} & -\frac{(n-4)a_{n-4}(n-5)}{720} + \frac{(n-2)a_{n-2}(n-3)}{24} \\ & -\frac{na_n(n-1)}{2} + (n+2)a_{n+2}(1+n) + 2na_n - a_{n-1} = 0 \end{aligned} \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{360n^2a_n + n^2a_{n-4} - 30n^2a_{n-2} - 1800na_n - 9na_{n-4} + 150na_{n-2} + 20a_{n-4} - 180a_{n-2} + 720a_{n-1}}{720(n+2)(1+n)}$$

$$\begin{aligned} (5) \quad & = \frac{(360n^2 - 1800n)a_n}{720(n+2)(1+n)} + \frac{(n^2 - 9n + 20)a_{n-4}}{720(n+2)(1+n)} \\ & + \frac{(-30n^2 + 150n - 180)a_{n-2}}{720(n+2)(1+n)} + \frac{a_{n-1}}{(n+2)(1+n)} \end{aligned}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1x + \left(\frac{a_0}{6} - \frac{a_1}{3}\right)x^3 + \frac{a_1x^4}{12} + \left(-\frac{a_0}{40} + \frac{a_1}{20}\right)x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + \frac{1}{6}x^3 - \frac{1}{40}x^5\right)a_0 + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{20}x^5\right)a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + \frac{1}{6}x^3 - \frac{1}{40}x^5\right)c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{20}x^5\right)c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 + \frac{1}{6}x^3 - \frac{1}{40}x^5 + \frac{1}{180}x^6\right)y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{20}x^5 - \frac{1}{60}x^6\right)y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 + \frac{1}{6}x^3 - \frac{1}{40}x^5\right)c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{20}x^5\right)c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{1}{6}x^3 - \frac{1}{40}x^5 + \frac{1}{180}x^6\right)y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{20}x^5 - \frac{1}{60}x^6\right)y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 + \frac{1}{6}x^3 - \frac{1}{40}x^5\right)c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{20}x^5\right)c_2 + O(x^6)$$

Verified OK.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 39

```
Order:=6;
dsolve(cos(x)*diff(y(x), x, x) + 2*x*diff(y(x), x) - x*y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{6}x^3 - \frac{1}{40}x^5\right)y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{12}x^4 + \frac{1}{20}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 49

```
AsymptoticDSolveValue[Cos[x]*y''[x]+2*x*y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{40} + \frac{x^3}{6} + 1 \right) + c_2 \left(\frac{x^5}{20} + \frac{x^4}{12} - \frac{x^3}{3} + x \right)$$

4.48 problem 45

4.48.1 Maple step by step solution 2088

Internal problem ID [7269]

Internal file name [OUTPUT/6255_Sunday_June_05_2022_04_35_50_PM_10751942/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 45.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + 4xy' + (x^2 + 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + 4xy' + (x^2 + 2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{4}{x}$$
$$q(x) = \frac{x^2 + 2}{x^2}$$

Table 209: Table $p(x), q(x)$ singularities.

$p(x) = \frac{4}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{x^2+2}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + 4xy' + (x^2 + 2)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + 4x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x^2 + 2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + 4x^{n+r} a_n (n+r) + 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + 4x^r a_0 r + 2a_0 x^r = 0$$

Or

$$(x^r r(-1+r) + 4x^r r + 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 + 3r + 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + 3r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= -1 \\ r_2 &= -2 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 + 3r + 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \frac{\sum_{n=0}^{\infty} a_n x^n}{x} \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x^2} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n-1} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-2} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 4a_n(n+r) + a_{n-2} + 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2 + 3n + 3r + 2} \quad (4)$$

Which for the root $r = -1$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+1)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{r^2 + 7r + 12}$$

Which for the root $r = -1$ becomes

$$a_2 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+7r+12}$	$-\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+7r+12}$	$-\frac{1}{6}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+4)(r+3)(6+r)(r+5)}$$

Which for the root $r = -1$ becomes

$$a_4 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+7r+12}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{1}{(r+4)(r+3)(6+r)(r+5)}$	$\frac{1}{120}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+7r+12}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{1}{(r+4)(r+3)(6+r)(r+5)}$	$\frac{1}{120}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \frac{1}{x} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= \frac{1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)}{x} \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -2} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-2} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + 4b_n(n+r) + b_{n-2} + 2b_n = 0 \quad (4)$$

Which for for the root $r = -2$ becomes

$$b_n(n-2)(n-3) + 4b_n(n-2) + b_{n-2} + 2b_n = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{b_{n-2}}{n^2 + 2nr + r^2 + 3n + 3r + 2} \quad (5)$$

Which for the root $r = -2$ becomes

$$b_n = -\frac{b_{n-2}}{n^2 - n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -2$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{1}{r^2 + 7r + 12}$$

Which for the root $r = -2$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+7r+12}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+7r+12}$	$-\frac{1}{2}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{(r^2 + 7r + 12)(r^2 + 11r + 30)}$$

Which for the root $r = -2$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+7r+12}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{1}{(r+4)(r+3)(6+r)(r+5)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+7r+12}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{1}{(r+4)(r+3)(6+r)(r+5)}$	$\frac{1}{24}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= \frac{1}{x} (b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)}{x^2} \end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 y_1(x) + c_2 y_2(x) \\ = \frac{c_1 \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)\right)}{x} + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)\right)}{x^2}$$

Hence the final solution is

$$y = y_h \\ = \frac{c_1 \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)\right)}{x} + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)\right)}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)\right)}{x} + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)\right)}{x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6)\right)}{x} + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)\right)}{x^2}$$

Verified OK.

4.48.1 Maple step by step solution

Let's solve

$$x^2 y'' + 4xy' + (x^2 + 2)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x^2+2)y}{x^2} - \frac{4y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{4y'}{x} + \frac{(x^2+2)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{4}{x}, P_3(x) = \frac{x^2+2}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 4$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + 4xy' + (x^2 + 2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(2+r)(1+r)x^r + a_1(3+r)(2+r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r+2)(k+r+1) + a_{k-2}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $(2 + r)(1 + r) = 0$
- Values of r that satisfy the indicial equation
 $r \in \{-2, -1\}$
- Each term must be 0
 $a_1(3 + r)(2 + r) = 0$
- Solve for the dependent coefficient(s)
 $a_1 = 0$
- Each term in the series must be 0, giving the recursion relation
 $a_k(k + r + 2)(k + r + 1) + a_{k-2} = 0$
- Shift index using $k- \rightarrow k + 2$
 $a_{k+2}(k + 4 + r)(k + 3 + r) + a_k = 0$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+4+r)(k+3+r)}$$
- Recursion relation for $r = -2$

$$a_{k+2} = -\frac{a_k}{(k+2)(k+1)}$$
- Solution for $r = -2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-2}, a_{k+2} = -\frac{a_k}{(k+2)(k+1)}, a_1 = 0 \right]$$
- Recursion relation for $r = -1$

$$a_{k+2} = -\frac{a_k}{(k+3)(k+2)}$$
- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+2} = -\frac{a_k}{(k+3)(k+2)}, a_1 = 0 \right]$$
- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-2} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-1} \right), a_{k+2} = -\frac{a_k}{(k+2)(k+1)}, a_1 = 0, b_{k+2} = -\frac{b_k}{(k+3)(k+2)}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Group is reducible or imprimitive  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
Order:=6;  
dsolve(x^2*diff(y(x), x, x) + 4*x*diff(y(x), x) + (x^2+2)*y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{6}x^2 + \frac{1}{120}x^4 + O(x^6)\right) x + c_2 \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 + O(x^6)\right)}{x^2}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 40

```
AsymptoticDSolveValue[x^2*y''[x]+4*x*y'[x]+(x^2+2)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^3}{120} - \frac{x}{6} + \frac{1}{x} \right) + c_1 \left(\frac{x^2}{24} + \frac{1}{x^2} - \frac{1}{2} \right)$$

4.49 problem 46

4.49.1 Maple step by step solution 2100

Internal problem ID [7270]

Internal file name [OUTPUT/6256_Sunday_June_05_2022_04_35_53_PM_8811729/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 46.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$x^2y'' + xy' - yx = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$x^2y'' + xy' - yx = 0$$

Or

$$x(xy'' + y' - y) = 0$$

For $x \neq 0$ the above simplifies to

$$xy'' + y' - y = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + xy' - yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{1}{x}$$

Table 211: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$		$q(x) = -\frac{1}{x}$	
singularity	type	singularity	type
$x = 0$	“regular”	$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + xy' - yx = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r = 0$$

Or

$$(x^r r(-1+r) + x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitray constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = 1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(r+1)^2(2+r)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{36}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{576}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{14400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(r+1)^2}$	1
a_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$
a_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$
a_5	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{14400}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned}
 y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\
 &= 1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6)
 \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{1}{(r+1)^2}$	1	$-\frac{2}{(r+1)^3}$	-2
b_2	$\frac{1}{(r+1)^2(2+r)^2}$	$\frac{1}{4}$	$\frac{-6-4r}{(r+1)^3(2+r)^3}$	$-\frac{3}{4}$
b_3	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2}$	$\frac{1}{36}$	$\frac{-6r^2-24r-22}{(r+1)^3(2+r)^3(r+3)^3}$	$-\frac{11}{108}$
b_4	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$	$\frac{-8r^3-60r^2-140r-100}{(r+1)^3(2+r)^3(r+3)^3(r+4)^3}$	$-\frac{25}{3456}$
b_5	$\frac{1}{(r+1)^2(2+r)^2(r+3)^2(r+4)^2(r+5)^2}$	$\frac{1}{14400}$	$\frac{-10r^4-120r^3-510r^2-900r-548}{(r+1)^3(2+r)^3(r+3)^3(r+4)^3(r+5)^3}$	$-\frac{137}{432000}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6)\right) \ln(x) \\ &\quad - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6)\right) \\ &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6)\right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\ &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6)\right) \\ &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6)\right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\ &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6)\right) \\ &\quad + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6)\right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\ &\quad \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right) \end{aligned} \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \\ + c_2 \left(\left(1 + x + \frac{x^2}{4} + \frac{x^3}{36} + \frac{x^4}{576} + \frac{x^5}{14400} + O(x^6) \right) \ln(x) - 2x - \frac{3x^2}{4} - \frac{11x^3}{108} \right. \\ \left. - \frac{25x^4}{3456} - \frac{137x^5}{432000} + O(x^6) \right)$$

Verified OK.

4.49.1 Maple step by step solution

Let's solve

$$x^2 y'' + xy' - yx = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{x} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1}{x}, P_3(x) = -\frac{1}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$xy'' + y' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r^2 x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)^2 - a_k) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1} (k+1)^2 - a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k}{(k+1)^2}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k}{(k+1)^2}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = \frac{a_k}{(k+1)^2} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

```

Order:=6;
dsolve(x^2*diff(y(x), x, x) + x*diff(y(x), x) - x*y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 + x + \frac{1}{4}x^2 + \frac{1}{36}x^3 + \frac{1}{576}x^4 + \frac{1}{14400}x^5 + O(x^6) \right) + \left((-2)x - \frac{3}{4}x^2 - \frac{11}{108}x^3 - \frac{25}{3456}x^4 - \frac{137}{432000}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 107

```

AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]-x*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) + c_2 \left(-\frac{137x^5}{432000} - \frac{25x^4}{3456} - \frac{11x^3}{108} - \frac{3x^2}{4} + \left(\frac{x^5}{14400} + \frac{x^4}{576} + \frac{x^3}{36} + \frac{x^2}{4} + x + 1 \right) \log(x) - 2x \right)$$

4.50 problem 47

4.50.1 Maple step by step solution 2112

Internal problem ID [7271]

Internal file name [OUTPUT/6257_Sunday_June_05_2022_04_35_55_PM_61307053/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 47.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{4x^2 - 1}{4x^2}$$

Table 213: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{4x^2-1}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + \left(x^2 - \frac{1}{4}\right) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(x^2 - \frac{1}{4} \right) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} \left(-\frac{a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} \left(-\frac{a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - \frac{a_n x^{n+r}}{4} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r - \frac{a_0 x^r}{4} = 0$$

Or

$$\left(x^r r(-1+r) + x^r r - \frac{x^r}{4} \right) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$\frac{(4r^2 - 1) x^r}{4} = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - \frac{1}{4} = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = -\frac{1}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$\frac{(4r^2 - 1)x^r}{4} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sqrt{x} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{\sqrt{x}}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} - \frac{a_n}{4} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-2}}{4n^2 + 8nr + 4r^2 - 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+1)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{4}{4r^2 + 16r + 15}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= \sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -\frac{1}{2}} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + b_n(n+r) + b_{n-2} - \frac{b_n}{4} = 0 \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n \left(n - \frac{1}{2} \right) \left(n - \frac{3}{2} \right) + b_n \left(n - \frac{1}{2} \right) + b_{n-2} - \frac{b_n}{4} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{4b_{n-2}}{4n^2 + 8nr + 4r^2 - 1} \quad (5)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = -\frac{4b_{n-2}}{4n^2 - 4n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{4}{4r^2 + 16r + 15}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2 + 16r + 15}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{16}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= \sqrt{x}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)}{\sqrt{x}} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6) \right)}{\sqrt{x}} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6) \right)}{\sqrt{x}} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6) \right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6) \right)}{\sqrt{x}}$$

Verified OK.

4.50.1 Maple step by step solution

Let's solve

$$x^2y'' + xy' + \left(x^2 - \frac{1}{4}\right)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(4x^2-1)y}{4x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(4x^2-1)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{4x^2-1}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{1}{4}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2y'' + 4xy' + (4x^2 - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+2r)(-1+2r)x^r + a_1(3+2r)(1+2r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r+1)(2k+2r-1) + 4a_{k-2}) \right) x^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+2r)(-1+2r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \left\{ -\frac{1}{2}, \frac{1}{2} \right\}$$
- Each term must be 0

$$a_1(3+2r)(1+2r) = 0$$
- Solve for the dependent coefficient(s)

$$a_1 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_k(4k^2 + 8kr + 4r^2 - 1) + 4a_{k-2} = 0$$
- Shift index using $k \rightarrow k+2$

$$a_{k+2}(4(k+2)^2 + 8(k+2)r + 4r^2 - 1) + 4a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{4a_k}{4k^2 + 8kr + 4r^2 + 16k + 16r + 15}$$
- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}$$
- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}, a_1 = 0 \right]$$
- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}$$
- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}, a_1 = 0 \right]$$
- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = -\frac{4a_k}{4k^2+12k+8}, a_1 = 0, b_{k+2} = -\frac{4b_k}{4k^2+20k+24}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```

Order:=6;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1/4)*y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{6}x^2 + \frac{1}{120}x^4 + O(x^6) \right) x + c_2 \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 + O(x^6) \right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 58

```

AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1/4)*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_1 \left(\frac{x^{7/2}}{24} - \frac{x^{3/2}}{2} + \frac{1}{\sqrt{x}} \right) + c_2 \left(\frac{x^{9/2}}{120} - \frac{x^{5/2}}{6} + \sqrt{x} \right)$$

4.51 problem 48

Internal problem ID [7272]

Internal file name [OUTPUT/6258_Sunday_June_05_2022_04_35_58_PM_44807529/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 48.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 - x)y'' - xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x^2 - x)y'' - xy' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{x-1}$$
$$q(x) = \frac{1}{x(x-1)}$$

Table 215: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{x-1}$	
singularity	type
$x = 1$	“regular”

$q(x) = \frac{1}{x(x-1)}$	
singularity	type
$x = 0$	“regular”
$x = 1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[1, 0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x(x-1)y'' - xy' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x(x-1) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r-1} a_n (n+r) (n+r-1)) \quad (2A) \\ & + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned}$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \\ \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ \sum_{n=0}^{\infty} a_n x^{n+r} &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-x^{n+r-1} a_n (n+r) (n+r-1)) \quad (2B) \\ & + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$-x^{n+r-1} a_n (n+r) (n+r-1) = 0$$

When $n=0$ the above becomes

$$-x^{-1+r} a_0 r (-1+r) = 0$$

Or

$$-x^{-1+r} a_0 r (-1+r) = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$-x^{-1+r}r(-1+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$-r(-1+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 1$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$-x^{-1+r}r(-1+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+1}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots

of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_{n-1}(n+r-1)(n+r-2) - a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}(n^2 + 2nr + r^2 - 4n - 4r + 4)}{(n+r)(n+r-1)} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-1}(n-1)^2}{n(n+1)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{(-1+r)^2}{r(1+r)}$$

Which for the root $r = 1$ becomes

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(-1+r)^2}{r(1+r)}$	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r(-1+r)^2}{(1+r)^2(2+r)}$$

Which for the root $r = 1$ becomes

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(-1+r)^2}{r(1+r)}$	0
a_2	$\frac{r(-1+r)^2}{(1+r)^2(2+r)}$	0

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{r(-1+r)^2}{(3+r)(2+r)^2}$$

Which for the root $r = 1$ becomes

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(-1+r)^2}{r(1+r)}$	0
a_2	$\frac{r(-1+r)^2}{(1+r)^2(2+r)}$	0
a_3	$\frac{r(-1+r)^2}{(3+r)(2+r)^2}$	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r(-1+r)^2}{(4+r)(3+r)^2}$$

Which for the root $r = 1$ becomes

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(-1+r)^2}{r(1+r)}$	0
a_2	$\frac{r(-1+r)^2}{(1+r)^2(2+r)}$	0
a_3	$\frac{r(-1+r)^2}{(3+r)(2+r)^2}$	0
a_4	$\frac{r(-1+r)^2}{(4+r)(3+r)^2}$	0

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{r(-1+r)^2}{(5+r)(4+r)^2}$$

Which for the root $r = 1$ becomes

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(-1+r)^2}{r(1+r)}$	0
a_2	$\frac{r(-1+r)^2}{(1+r)^2(2+r)}$	0
a_3	$\frac{r(-1+r)^2}{(3+r)(2+r)^2}$	0
a_4	$\frac{r(-1+r)^2}{(4+r)(3+r)^2}$	0
a_5	$\frac{r(-1+r)^2}{(5+r)(4+r)^2}$	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x(1 + O(x^6)) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= \frac{(-1+r)^2}{r(1+r)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} \frac{(-1+r)^2}{r(1+r)} &= \lim_{r \rightarrow 0} \frac{(-1+r)^2}{r(1+r)} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x(x-1)y'' - xy' + y = 0$ gives

$$\begin{aligned}
& x(x-1) \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\
& \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
& - x \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \right) \\
& + Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
\end{aligned}$$

Which can be written as

$$\begin{aligned}
& \left((x(x-1)y_1''(x) - y_1'(x)x + y_1(x)) \ln(x) + x(x-1) \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) \right. \\
& \left. - y_1(x) \right) C + x(x-1) \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
& - x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
\end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$x(x-1)y_1''(x) - y_1'(x)x + y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned}
& \left(x(x-1) \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) - y_1(x) \right) C \\
& + x(x-1) \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
& - x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
\end{aligned} \tag{8}$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \frac{\left(2x(x-1) \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1)\right) + (1-2x) \left(\sum_{n=0}^{\infty} a_n x^{n+r_1}\right)\right) C}{x} \\ & + \frac{x^2(x-1) \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2)\right) - \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2)\right) x^2 + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2}\right) x}{x} \\ & = 0 \end{aligned} \tag{9}$$

Since $r_1 = 1$ and $r_2 = 0$ then the above becomes

$$\begin{aligned} & \frac{\left(2x(x-1) \left(\sum_{n=0}^{\infty} x^n a_n (n+1)\right) + (1-2x) \left(\sum_{n=0}^{\infty} a_n x^{n+1}\right)\right) C}{x} \\ & + \frac{x^2(x-1) \left(\sum_{n=0}^{\infty} x^{-2+n} b_n n (n-1)\right) - \left(\sum_{n=0}^{\infty} x^{n-1} b_n n\right) x^2 + \left(\sum_{n=0}^{\infty} b_n x^n\right) x}{x} = 0 \end{aligned} \tag{10}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2C x^{n+1} a_n (n+1)\right) + \sum_{n=0}^{\infty} (-2C x^n a_n (n+1)) \\ & + \left(\sum_{n=0}^{\infty} C a_n x^n\right) + \sum_{n=0}^{\infty} (-2C x^{n+1} a_n) + \left(\sum_{n=0}^{\infty} x^n b_n n (n-1)\right) \\ & + \sum_{n=0}^{\infty} (-n x^{n-1} b_n (n-1)) + \sum_{n=0}^{\infty} (-x^n b_n n) + \left(\sum_{n=0}^{\infty} b_n x^n\right) = 0 \end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 2C x^{n+1} a_n (n+1) &= \sum_{n=2}^{\infty} 2C a_{-2+n} (n-1) x^{n-1} \\ \sum_{n=0}^{\infty} (-2C x^n a_n (n+1)) &= \sum_{n=1}^{\infty} (-2C a_{n-1} n x^{n-1}) \end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{\infty} C a_n x^n &= \sum_{n=1}^{\infty} C a_{n-1} x^{n-1} \\
\sum_{n=0}^{\infty} (-2C x^{n+1} a_n) &= \sum_{n=2}^{\infty} (-2C a_{-2+n} x^{n-1}) \\
\sum_{n=0}^{\infty} x^n b_n n(n-1) &= \sum_{n=1}^{\infty} (n-1) b_{n-1} (-2+n) x^{n-1} \\
\sum_{n=0}^{\infty} (-x^n b_n n) &= \sum_{n=1}^{\infty} (-(n-1) b_{n-1} x^{n-1}) \\
\sum_{n=0}^{\infty} b_n x^n &= \sum_{n=1}^{\infty} b_{n-1} x^{n-1}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n - 1$.

$$\begin{aligned}
&\left(\sum_{n=2}^{\infty} 2C a_{-2+n} (n-1) x^{n-1} \right) + \sum_{n=1}^{\infty} (-2C a_{n-1} n x^{n-1}) \\
&+ \left(\sum_{n=1}^{\infty} C a_{n-1} x^{n-1} \right) + \sum_{n=2}^{\infty} (-2C a_{-2+n} x^{n-1}) \\
&+ \left(\sum_{n=1}^{\infty} (n-1) b_{n-1} (-2+n) x^{n-1} \right) + \sum_{n=0}^{\infty} (-n x^{n-1} b_n (n-1)) \\
&+ \sum_{n=1}^{\infty} (-(n-1) b_{n-1} x^{n-1}) + \left(\sum_{n=1}^{\infty} b_{n-1} x^{n-1} \right) = 0
\end{aligned} \tag{2B}$$

For $n = 0$ in Eq. (2B), we choose arbitray value for b_0 as $b_0 = 1$. For $n = N$, where $N = 1$ which is the difference between the two roots, we are free to choose $b_1 = 0$. Hence for $n = 1$, Eq (2B) gives

$$-C + 1 = 0$$

Which is solved for C . Solving for C gives

$$C = 1$$

For $n = 2$, Eq (2B) gives

$$-3C a_1 - 2b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-2b_2 = 0$$

Solving the above for b_2 gives

$$b_2 = 0$$

For $n = 3$, Eq (2B) gives

$$(2a_1 - 5a_2)C + b_2 - 6b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-6b_3 = 0$$

Solving the above for b_3 gives

$$b_3 = 0$$

For $n = 4$, Eq (2B) gives

$$(4a_2 - 7a_3)C + 4b_3 - 12b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-12b_4 = 0$$

Solving the above for b_4 gives

$$b_4 = 0$$

For $n = 5$, Eq (2B) gives

$$(6a_3 - 9a_4)C + 9b_4 - 20b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-20b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = 0$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = 1$ and all b_n , then the second solution becomes

$$y_2(x) = 1(x(1 + O(x^6))) \ln(x) + 1 + O(x^6)$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x(1 + O(x^6)) + c_2 (1(x(1 + O(x^6))) \ln(x) + 1 + O(x^6)) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 x(1 + O(x^6)) + c_2 (x(1 + O(x^6)) \ln(x) + 1 + O(x^6)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x(1 + O(x^6)) + c_2 (x(1 + O(x^6)) \ln(x) + 1 + O(x^6)) \quad (1)$$

Verification of solutions

$$y = c_1 x(1 + O(x^6)) + c_2 (x(1 + O(x^6)) \ln(x) + 1 + O(x^6))$$

Verified OK.

Maple trace **Kovacic algorithm successful**

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
  A Liouvillian solution exists
  Reducible group (found an exponential solution)
  Group is reducible, not completely reducible
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((x^2-x)*diff(y(x), x)-x*diff(y(x), x)+y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = \ln(x) (x + O(x^6)) c_2 + c_1 x (1 + O(x^6)) + (1 - x + O(x^6)) c_2$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 20

```
AsymptoticDSolveValue[(x^2-x)*y'[x]-x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 x + c_1 (-3x + x \log(x) + 1)$$

4.52 problem 49

4.52.1 Maple step by step solution 2141

Internal problem ID [7273]

Internal file name [OUTPUT/6259_Sunday_June_05_2022_04_36_02_PM_79190898/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 49.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order, _with_linear_symmetries], [_2nd_order, _linear, `
  _with_symmetry_[0,F(x)]`]]
```

$$x^2y'' + (x^2 + 6x)y' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$x^2y'' + (x^2 + 6x)y' + yx = 0$$

Or

$$x(xy' + xy'' + y + 6y') = 0$$

For $x \neq 0$ the above simplifies to

$$xy'' + (x + 6)y' + y = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + (x^2 + 6x)y' + yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{x+6}{x}$$

$$q(x) = \frac{1}{x}$$

Table 216: Table $p(x), q(x)$ singularities.

$p(x) = \frac{x+6}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + (x^2 + 6x) y' + yx = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + (x^2 + 6x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned}
 & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) \right) \\
 & + \left(\sum_{n=0}^{\infty} 6x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0
 \end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}
 \sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r} \\
 \sum_{n=0}^{\infty} x^{1+n+r} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r}
 \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned}
 & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r} \right) \\
 & + \left(\sum_{n=0}^{\infty} 6x^{n+r} a_n (n+r) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) = 0
 \end{aligned} \tag{2B}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r)(n+r-1) + 6x^{n+r} a_n (n+r) = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r(-1 + r) + 6x^r a_0 r = 0$$

Or

$$(x^r r(-1 + r) + 6x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r(5 + r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(5 + r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = -5$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r(5 + r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 5$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x^5}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-5} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_{n-1}(n+r-1) + 6a_n(n+r) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{n+5+r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-1}}{n+5} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{6+r}$$

Which for the root $r = 0$ becomes

$$a_1 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{6+r}$	$-\frac{1}{6}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(6+r)(7+r)}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{42}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{6+r}$	$-\frac{1}{6}$
a_2	$\frac{1}{(6+r)(7+r)}$	$\frac{1}{42}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{(6+r)(7+r)(8+r)}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{1}{336}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{6+r}$	$-\frac{1}{6}$
a_2	$\frac{1}{(6+r)(7+r)}$	$\frac{1}{42}$
a_3	$-\frac{1}{(6+r)(7+r)(8+r)}$	$-\frac{1}{336}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(6+r)(7+r)(8+r)(9+r)}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{3024}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{6+r}$	$-\frac{1}{6}$
a_2	$\frac{1}{(6+r)(7+r)}$	$\frac{1}{42}$
a_3	$-\frac{1}{(6+r)(7+r)(8+r)}$	$-\frac{1}{336}$
a_4	$\frac{1}{(6+r)(7+r)(8+r)(9+r)}$	$\frac{1}{3024}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{(6+r)(7+r)(8+r)(9+r)(10+r)}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{1}{30240}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{6+r}$	$-\frac{1}{6}$
a_2	$\frac{1}{(6+r)(7+r)}$	$\frac{1}{42}$
a_3	$-\frac{1}{(6+r)(7+r)(8+r)}$	$-\frac{1}{336}$
a_4	$\frac{1}{(6+r)(7+r)(8+r)(9+r)}$	$\frac{1}{3024}$
a_5	$-\frac{1}{(6+r)(7+r)(8+r)(9+r)(10+r)}$	$-\frac{1}{30240}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 - \frac{x}{6} + \frac{x^2}{42} - \frac{x^3}{336} + \frac{x^4}{3024} - \frac{x^5}{30240} + O(x^6) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 5$. Now we need to determine if

C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_5(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_5 \\ &= -\frac{1}{(6+r)(7+r)(8+r)(9+r)(10+r)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} -\frac{1}{(6+r)(7+r)(8+r)(9+r)(10+r)} &= \lim_{r \rightarrow -5} -\frac{1}{(6+r)(7+r)(8+r)(9+r)(10+r)} \\ &= -\frac{1}{120} \end{aligned}$$

The limit is $-\frac{1}{120}$. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-5} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + b_{n-1}(n+r-1) + 6b_n(n+r) + b_{n-1} = 0 \quad (4)$$

Which for the root $r = -5$ becomes

$$b_n(n-5)(n-6) + b_{n-1}(n-6) + 6b_n(n-5) + b_{n-1} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{b_{n-1}}{n+5+r} \quad (5)$$

Which for the root $r = -5$ becomes

$$b_n = -\frac{b_{n-1}}{n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -5$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -\frac{1}{6+r}$$

Which for the root $r = -5$ becomes

$$b_1 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{6+r}$	-1

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{(6+r)(7+r)}$$

Which for the root $r = -5$ becomes

$$b_2 = \frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{6+r}$	-1
b_2	$\frac{1}{(6+r)(7+r)}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{1}{(6+r)(7+r)(8+r)}$$

Which for the root $r = -5$ becomes

$$b_3 = -\frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{6+r}$	-1
b_2	$\frac{1}{(6+r)(7+r)}$	$\frac{1}{2}$
b_3	$-\frac{1}{(6+r)(7+r)(8+r)}$	$-\frac{1}{6}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{(6+r)(7+r)(8+r)(9+r)}$$

Which for the root $r = -5$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{6+r}$	-1
b_2	$\frac{1}{(6+r)(7+r)}$	$\frac{1}{2}$
b_3	$-\frac{1}{(6+r)(7+r)(8+r)}$	$-\frac{1}{6}$
b_4	$\frac{1}{(6+r)(7+r)(8+r)(9+r)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{1}{(6+r)(7+r)(8+r)(9+r)(10+r)}$$

Which for the root $r = -5$ becomes

$$b_5 = -\frac{1}{120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{6+r}$	-1
b_2	$\frac{1}{(6+r)(7+r)}$	$\frac{1}{2}$
b_3	$-\frac{1}{(6+r)(7+r)(8+r)}$	$-\frac{1}{6}$
b_4	$\frac{1}{(6+r)(7+r)(8+r)(9+r)}$	$\frac{1}{24}$
b_5	$-\frac{1}{(6+r)(7+r)(8+r)(9+r)(10+r)}$	$-\frac{1}{120}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\
 &= \frac{1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6)}{x^5}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1 \left(1 - \frac{x}{6} + \frac{x^2}{42} - \frac{x^3}{336} + \frac{x^4}{3024} - \frac{x^5}{30240} + O(x^6) \right) \\
 &\quad + \frac{c_2 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6) \right)}{x^5}
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 \left(1 - \frac{x}{6} + \frac{x^2}{42} - \frac{x^3}{336} + \frac{x^4}{3024} - \frac{x^5}{30240} + O(x^6) \right) \\
 &\quad + \frac{c_2 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6) \right)}{x^5}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - \frac{x}{6} + \frac{x^2}{42} - \frac{x^3}{336} + \frac{x^4}{3024} - \frac{x^5}{30240} + O(x^6) \right) + \frac{c_2 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6) \right)}{x^5} \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 - \frac{x}{6} + \frac{x^2}{42} - \frac{x^3}{336} + \frac{x^4}{3024} - \frac{x^5}{30240} + O(x^6) \right) + \frac{c_2 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6) \right)}{x^5}$$

Verified OK.

4.52.1 Maple step by step solution

Let's solve

$$x^2 y'' + (x^2 + 6x) y' + yx = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{x} - \frac{(x+6)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(x+6)y'}{x} + \frac{y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{x+6}{x}, P_3(x) = \frac{1}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 6$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$xy'' + (x + 6)y' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(5+r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)(k+6+r) + a_k (k+1+r)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(5+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-5, 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k+1+r)(a_{k+1}(k+6+r) + a_k) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{k+6+r}$$

- Recursion relation for $r = -5$

$$a_{k+1} = -\frac{a_k}{k+1}$$

- Solution for $r = -5$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-5}, a_{k+1} = -\frac{a_k}{k+1} \right]$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{a_k}{k+6}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{a_k}{k+6} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-5} \right) + \left(\sum_{k=0}^{\infty} b_k x^k \right), a_{k+1} = -\frac{a_k}{k+1}, b_{k+1} = -\frac{b_k}{k+6} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=6;
dsolve(x^2*diff(y(x), x$2)+(6*x+x^2)*diff(y(x), x)+x*y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 - \frac{1}{6}x + \frac{1}{42}x^2 - \frac{1}{336}x^3 + \frac{1}{3024}x^4 - \frac{1}{30240}x^5 + O(x^6) \right) + \frac{c_2(2880 - 2880x + 1440x^2 - 480x^3 + 120x^4 - 24x^5 + O(x^6))}{x^5}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 68

```
AsymptoticDSolveValue[x^2*y''[x]+(6*x+x^2)*y'[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^4}{3024} - \frac{x^3}{336} + \frac{x^2}{42} - \frac{x}{6} + 1 \right) + c_1 \left(\frac{1}{x^5} - \frac{1}{x^4} + \frac{1}{2x^3} - \frac{1}{6x^2} + \frac{1}{24x} \right)$$

4.53 problem 50

4.53.1 Maple step by step solution 2157

Internal problem ID [7274]

Internal file name [OUTPUT/6260_Sunday_June_05_2022_04_36_07_PM_1175942/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 50.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' - xy' + (x^2 - 8)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' - xy' + (x^2 - 8)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{x}$$
$$q(x) = \frac{x^2 - 8}{x^2}$$

Table 218: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{x^2-8}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' - x y' + (x^2 - 8) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x^2 - 8) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} (-8a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} (-8a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) - 8a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) - x^r a_0 r - 8a_0 x^r = 0$$

Or

$$(x^r r(-1+r) - x^r r - 8x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 2r - 8) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 2r - 8 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 4 \\ r_2 &= -2 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 2r - 8) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 6$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= x^4 \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x^2} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+4} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-2} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_n(n+r) + a_{n-2} - 8a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2 - 2n - 2r - 8} \quad (4)$$

Which for the root $r = 4$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+6)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 4$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{r^2 + 2r - 8}$$

Which for the root $r = 4$ becomes

$$a_2 = -\frac{1}{16}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+2r-8}$	$-\frac{1}{16}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+2r-8}$	$-\frac{1}{16}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+4)(-2+r)r(r+6)}$$

Which for the root $r = 4$ becomes

$$a_4 = \frac{1}{640}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+2r-8}$	$-\frac{1}{16}$
a_3	0	0
a_4	$\frac{1}{(r+4)(-2+r)r(r+6)}$	$\frac{1}{640}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+2r-8}$	$-\frac{1}{16}$
a_3	0	0
a_4	$\frac{1}{(r+4)(-2+r)r(r+6)}$	$\frac{1}{640}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{1}{(r+4)(-2+r)r(r+6)(8+r)(2+r)}$$

Which for the root $r = 4$ becomes

$$a_6 = -\frac{1}{46080}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+2r-8}$	$-\frac{1}{16}$
a_3	0	0
a_4	$\frac{1}{(r+4)(-2+r)r(r+6)}$	$\frac{1}{640}$
a_5	0	0
a_6	$-\frac{1}{(r+4)(-2+r)r(r+6)(8+r)(2+r)}$	$-\frac{1}{46080}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^4(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 \dots) \\ &= x^4\left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 6$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_6(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_6 \\ &= -\frac{1}{(r+4)(-2+r)r(r+6)(8+r)(2+r)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} -\frac{1}{(r+4)(-2+r)r(r+6)(8+r)(2+r)} &= \lim_{r \rightarrow -2} -\frac{1}{(r+4)(-2+r)r(r+6)(8+r)(2+r)} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx}y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2}y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x^2y'' - xy' + (x^2 - 8)y = 0$ gives

$$\begin{aligned} &x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\ &\quad - x \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \right) \\ &\quad + (x^2 - 8) \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) = 0 \end{aligned}$$

Which can be written as

$$\begin{aligned} &\left((x^2y_1''(x) - y_1'(x)x + (x^2 - 8)y_1(x)) \ln(x) + x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) - y_1(x) \right) C \\ &\quad + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\ &\quad - x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) + (x^2 - 8) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$x^2 y_1''(x) - y_1'(x)x + (x^2 - 8)y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned} & \left(x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) - y_1(x) \right) C \\ & + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ & - x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + (x^2 - 8) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x - 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) x^2 \\ & - \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) x - 8 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (9)$$

Since $r_1 = 4$ and $r_2 = -2$ then the above becomes

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{3+n} a_n (n+4) \right) x - 2 \left(\sum_{n=0}^{\infty} a_n x^{n+4} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-4+n} b_n (n-2) (-3+n) \right) x^2 + \left(\sum_{n=0}^{\infty} b_n x^{n-2} \right) x^2 \\ & - \left(\sum_{n=0}^{\infty} x^{-3+n} b_n (n-2) \right) x - 8 \left(\sum_{n=0}^{\infty} b_n x^{n-2} \right) = 0 \end{aligned} \quad (10)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2C x^{n+4} a_n (n+4) \right) + \sum_{n=0}^{\infty} (-2C a_n x^{n+4}) + \left(\sum_{n=0}^{\infty} x^{n-2} b_n (n^2 - 5n + 6) \right) \\ & + \left(\sum_{n=0}^{\infty} b_n x^n \right) + \sum_{n=0}^{\infty} (-x^{n-2} b_n (n-2)) + \sum_{n=0}^{\infty} (-8b_n x^{n-2}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n - 2$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-2} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 2C x^{n+4} a_n (n+4) &= \sum_{n=6}^{\infty} 2C a_{n-6} (n-2) x^{n-2} \\ \sum_{n=0}^{\infty} (-2C a_n x^{n+4}) &= \sum_{n=6}^{\infty} (-2C a_{n-6} x^{n-2}) \\ \sum_{n=0}^{\infty} b_n x^n &= \sum_{n=2}^{\infty} b_{n-2} x^{n-2} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n - 2$.

$$\begin{aligned} & \left(\sum_{n=6}^{\infty} 2C a_{n-6} (n-2) x^{n-2} \right) + \sum_{n=6}^{\infty} (-2C a_{n-6} x^{n-2}) \\ & + \left(\sum_{n=0}^{\infty} x^{n-2} b_n (n^2 - 5n + 6) \right) + \left(\sum_{n=2}^{\infty} b_{n-2} x^{n-2} \right) \\ & + \sum_{n=0}^{\infty} (-x^{n-2} b_n (n-2)) + \sum_{n=0}^{\infty} (-8b_n x^{n-2}) = 0 \end{aligned} \quad (2B)$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = 1$, Eq (2B) gives

$$-5b_1 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-5b_1 = 0$$

Solving the above for b_1 gives

$$b_1 = 0$$

For $n = 2$, Eq (2B) gives

$$b_0 - 8b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$1 - 8b_2 = 0$$

Solving the above for b_2 gives

$$b_2 = \frac{1}{8}$$

For $n = 3$, Eq (2B) gives

$$b_1 - 9b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-9b_3 = 0$$

Solving the above for b_3 gives

$$b_3 = 0$$

For $n = 4$, Eq (2B) gives

$$b_2 - 8b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{1}{8} - 8b_4 = 0$$

Solving the above for b_4 gives

$$b_4 = \frac{1}{64}$$

For $n = 5$, Eq (2B) gives

$$-5b_5 + b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-5b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = 0$$

For $n = N$, where $N = 6$ which is the difference between the two roots, we are free to choose $b_6 = 0$. Hence for $n = 6$, Eq (2B) gives

$$6C + \frac{1}{64} = 0$$

Which is solved for C . Solving for C gives

$$C = -\frac{1}{384}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -\frac{1}{384}$ and all b_n , then the second solution becomes

$$y_2(x) = -\frac{1}{384} \left(x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \right) \ln(x) + \frac{1 + \frac{x^2}{8} + \frac{x^4}{64} + O(x^7)}{x^2}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \\ &\quad + c_2 \left(-\frac{1}{384} \left(x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \right) \ln(x) \right. \\ &\quad \left. + \frac{1 + \frac{x^2}{8} + \frac{x^4}{64} + O(x^7)}{x^2} \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \\ &\quad + c_2 \left(-\frac{x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \ln(x)}{384} + \frac{1 + \frac{x^2}{8} + \frac{x^4}{64} + O(x^7)}{x^2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1 x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \\ &\quad + c_2 \left(-\frac{x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \ln(x)}{384} + \frac{1 + \frac{x^2}{8} + \frac{x^4}{64} + O(x^7)}{x^2} \right) \quad (1) \end{aligned}$$

Verification of solutions

$$y = c_1 x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \\ + c_2 \left(-\frac{x^4 \left(1 - \frac{x^2}{16} + \frac{x^4}{640} - \frac{x^6}{46080} + O(x^7) \right) \ln(x)}{384} + \frac{1 + \frac{x^2}{8} + \frac{x^4}{64} + O(x^7)}{x^2} \right)$$

Verified OK.

4.53.1 Maple step by step solution

Let's solve

$$x^2 y'' - xy' + (x^2 - 8)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x^2-8)y}{x^2} + \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{x} + \frac{(x^2-8)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{1}{x}, P_3(x) = \frac{x^2-8}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -8$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' - xy' + (x^2 - 8)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(2+r)(-4+r)x^r + a_1(3+r)(-3+r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r+2)(k+r-4) + a_{k-2}) x^{k+r} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(2+r)(-4+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-2, 4\}$$

- Each term must be 0

$$a_1(3+r)(-3+r) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+2)(k+r-4) + a_{k-2} = 0$$

- Shift index using $k- > k + 2$

$$a_{k+2}(k+4+r)(k-2+r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+4+r)(k-2+r)}$$

- Recursion relation for $r = -2$

$$a_{k+2} = -\frac{a_k}{(k+2)(k-4)}$$

- Series not valid for $r = -2$, division by 0 in the recursion relation at $k = 4$

$$a_{k+2} = -\frac{a_k}{(k+2)(k-4)}$$

- Recursion relation for $r = 4$

$$a_{k+2} = -\frac{a_k}{(k+8)(k+2)}$$

- Solution for $r = 4$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+4}, a_{k+2} = -\frac{a_k}{(k+8)(k+2)}, a_1 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```

Order:=6;
dsolve(x^2*diff(y(x), x$2)-x*diff(y(x), x)+(x^2-8)*y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = c_1 x^4 \left(1 - \frac{1}{16} x^2 + \frac{1}{640} x^4 + O(x^6) \right) + \frac{c_2 (-86400 - 10800x^2 - 1350x^4 + O(x^6))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 42

```
AsymptoticDSolveValue[x^2*y''[x]-x*y'[x]+(x^2-8)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^2}{64} + \frac{1}{x^2} + \frac{1}{8} \right) + c_2 \left(\frac{x^8}{640} - \frac{x^6}{16} + x^4 \right)$$

4.54 problem 51

4.54.1 Maple step by step solution 2168

Internal problem ID [7275]

Internal file name [OUTPUT/6261_Sunday_June_05_2022_04_36_11_PM_97759864/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 51.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$x^2y'' - 9xy' + 25y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' - 9xy' + 25y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{9}{x}$$
$$q(x) = \frac{25}{x^2}$$

Table 220: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{9}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{25}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' - 9xy' + 25y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - 9x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 25 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-9x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 25a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-9x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 25a_n x^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - 9x^{n+r} a_n (n+r) + 25a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r (-1+r) - 9x^r a_0 r + 25a_0 x^r = 0$$

Or

$$(x^r r (-1+r) - 9x^r r + 25x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r-5)^2 x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$(r-5)^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 5 \\ r_2 &= 5 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r-5)^2 x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. Using the value of the indicial root found earlier, $r = 5$, Eqs (1A,1B) become

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+5}$$

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+5} \right)$$

We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $0 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - 9a_n(n+r) + 25a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = 0 \quad (4)$$

Which for the root $r = 5$ becomes

$$a_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 5$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0

Using the above table, then the first solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^5(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^5(1 + O(x^6)) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 5$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 5)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	0	0	0	0
b_2	0	0	0	0
b_3	0	0	0	0
b_4	0	0	0	0
b_5	0	0	0	0

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned}
 y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\
 &= x^5(1 + O(x^6)) \ln(x) + x^5O(x^6)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x^5(1 + O(x^6)) + c_2(x^5(1 + O(x^6)) \ln(x) + x^5O(x^6))
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1x^5(1 + O(x^6)) + c_2(x^5(1 + O(x^6)) \ln(x) + x^5O(x^6))
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1x^5(1 + O(x^6)) + c_2(x^5(1 + O(x^6)) \ln(x) + x^5O(x^6)) \quad (1)$$

Verification of solutions

$$y = c_1x^5(1 + O(x^6)) + c_2(x^5(1 + O(x^6)) \ln(x) + x^5O(x^6))$$

Verified OK.

4.54.1 Maple step by step solution

Let's solve

$$x^2y'' - 9xy' + 25y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{9y'}{x} - \frac{25y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{9y'}{x} + \frac{25y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2y'' - 9xy' + 25y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) - 9 \frac{d}{dt}y(t) + 25y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) - 10 \frac{d}{dt}y(t) + 25y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 10r + 25 = 0$$

- Factor the characteristic polynomial
 $(r - 5)^2 = 0$
- Root of the characteristic polynomial
 $r = 5$
- 1st solution of the ODE
 $y_1(t) = e^{5t}$
- Repeated root, multiply $y_1(t)$ by t to ensure linear independence
 $y_2(t) = t e^{5t}$
- General solution of the ODE
 $y(t) = c_1 y_1(t) + c_2 y_2(t)$
- Substitute in solutions
 $y(t) = c_1 e^{5t} + c_2 t e^{5t}$
- Change variables back using $t = \ln(x)$
 $y = c_2 \ln(x) x^5 + c_1 x^5$
- Simplify
 $y = x^5(c_2 \ln(x) + c_1)$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```

Order:=6;
dsolve(x^2*diff(y(x), x$2)-9*x*diff(y(x), x)+25*y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = x^5(c_2 \ln(x) + c_1) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 18

```
AsymptoticDSolveValue[x^2*y''[x]-9*x*y'[x]+25*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x^5 + c_2 x^5 \log(x)$$

4.55 problem 52

4.55.1 Maple step by step solution 2180

Internal problem ID [7276]

Internal file name [OUTPUT/6262_Sunday_June_05_2022_04_36_13_PM_23047735/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 52.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' - xy' - \left(x^2 + \frac{5}{4}\right) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2 y'' - xy' + \left(-x^2 - \frac{5}{4}\right) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{x}$$
$$q(x) = -\frac{4x^2 + 5}{4x^2}$$

Table 222: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{4x^2+5}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' - x y' + \left(-x^2 - \frac{5}{4}\right) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(-x^2 - \frac{5}{4} \right) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \sum_{n=0}^{\infty} (-x^{n+r+2} a_n) + \sum_{n=0}^{\infty} \left(-\frac{5a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \sum_{n=2}^{\infty} (-a_{n-2} x^{n+r}) + \sum_{n=0}^{\infty} \left(-\frac{5a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) - \frac{5a_n x^{n+r}}{4} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) - x^r a_0 r - \frac{5a_0 x^r}{4} = 0$$

Or

$$\left(x^r r(-1+r) - x^r r - \frac{5x^r}{4} \right) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$\frac{(4r^2 - 8r - 5) x^r}{4} = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 2r - \frac{5}{4} = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{5}{2}$$

$$r_2 = -\frac{1}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$\frac{(4r^2 - 8r - 5)x^r}{4} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 3$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x^{\frac{5}{2}} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{\sqrt{x}}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{5}{2}}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_n(n+r) - a_{n-2} - \frac{5a_n}{4} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{4a_{n-2}}{4n^2 + 8nr + 4r^2 - 8n - 8r - 5} \quad (4)$$

Which for the root $r = \frac{5}{2}$ becomes

$$a_n = \frac{a_{n-2}}{n(n+3)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{5}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{4}{4r^2 + 8r - 5}$$

Which for the root $r = \frac{5}{2}$ becomes

$$a_2 = \frac{1}{10}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{4}{4r^2+8r-5}$	$\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{4}{4r^2+8r-5}$	$\frac{1}{10}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16}{(4r^2 + 8r - 5)(4r^2 + 24r + 27)}$$

Which for the root $r = \frac{5}{2}$ becomes

$$a_4 = \frac{1}{280}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{4}{4r^2+8r-5}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{16}{(4r^2+8r-5)(4r^2+24r+27)}$	$\frac{1}{280}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{4}{4r^2+8r-5}$	$\frac{1}{10}$
a_3	0	0
a_4	$\frac{16}{(4r^2+8r-5)(4r^2+24r+27)}$	$\frac{1}{280}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^{\frac{5}{2}}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^{\frac{5}{2}}\left(1 + \frac{x^2}{10} + \frac{x^4}{280} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 3$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_3(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_3 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -\frac{1}{2}} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) - b_n(n+r) - b_{n-2} - \frac{5b_n}{4} = 0 \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n \left(n - \frac{1}{2} \right) \left(n - \frac{3}{2} \right) - b_n \left(n - \frac{1}{2} \right) - b_{n-2} - \frac{5b_n}{4} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = \frac{4b_{n-2}}{4n^2 + 8nr + 4r^2 - 8n - 8r - 5} \quad (5)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = \frac{4b_{n-2}}{4n^2 - 12n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{4}{4r^2 + 8r - 5}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{4}{4r^2 + 8r - 5}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{4}{4r^2+8r-5}$	$-\frac{1}{2}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{16}{(4r^2 + 8r - 5)(4r^2 + 24r + 27)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = -\frac{1}{8}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{4}{4r^2+8r-5}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+8r-5)(4r^2+24r+27)}$	$-\frac{1}{8}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$\frac{4}{4r^2+8r-5}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+8r-5)(4r^2+24r+27)}$	$-\frac{1}{8}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^{\frac{5}{2}}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 - \frac{x^2}{2} - \frac{x^4}{8} + O(x^6)}{\sqrt{x}} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^{\frac{5}{2}}\left(1 + \frac{x^2}{10} + \frac{x^4}{280} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^2}{2} - \frac{x^4}{8} + O(x^6)\right)}{\sqrt{x}} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x^{\frac{5}{2}}\left(1 + \frac{x^2}{10} + \frac{x^4}{280} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^2}{2} - \frac{x^4}{8} + O(x^6)\right)}{\sqrt{x}} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1x^{\frac{5}{2}}\left(1 + \frac{x^2}{10} + \frac{x^4}{280} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^2}{2} - \frac{x^4}{8} + O(x^6)\right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = c_1x^{\frac{5}{2}}\left(1 + \frac{x^2}{10} + \frac{x^4}{280} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^2}{2} - \frac{x^4}{8} + O(x^6)\right)}{\sqrt{x}}$$

Verified OK.

4.55.1 Maple step by step solution

Let's solve

$$x^2y'' - xy' + \left(-x^2 - \frac{5}{4}\right)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{(4x^2+5)y}{4x^2} + \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{x} - \frac{(4x^2+5)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{1}{x}, P_3(x) = -\frac{4x^2+5}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{5}{4}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2 y'' - 4xy' + (-4x^2 - 5)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+2r)(-5+2r)x^r + a_1(3+2r)(-3+2r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r+1)(2k+2r-5) - 4a_{k-2}) \right) x^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+2r)(-5+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ -\frac{1}{2}, \frac{5}{2} \right\}$$

- Each term must be 0

$$a_1(3+2r)(-3+2r) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$4\left(k+r-\frac{5}{2}\right)\left(k+r+\frac{1}{2}\right)a_k - 4a_{k-2} = 0$$

- Shift index using $k- > k+2$

$$4\left(k-\frac{1}{2}+r\right)\left(k+\frac{5}{2}+r\right)a_{k+2} - 4a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = \frac{4a_k}{(2k-1+2r)(2k+5+2r)}$$

- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+2} = \frac{4a_k}{(2k-2)(2k+4)}$$

- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+2} = \frac{4a_k}{(2k-2)(2k+4)}, a_1 = 0 \right]$$

- Recursion relation for $r = \frac{5}{2}$

$$a_{k+2} = \frac{4a_k}{(2k+4)(2k+10)}$$

- Solution for $r = \frac{5}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{5}{2}}, a_{k+2} = \frac{4a_k}{(2k+4)(2k+10)}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{5}{2}} \right), a_{k+2} = \frac{4a_k}{(2k-2)(2k+4)}, a_1 = 0, b_{k+2} = \frac{4b_k}{(2k+4)(2k+10)}, b_1 = 0 \right]$$

Maple trace **Kovacic algorithm successful**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```

Order:=6;
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)-(x^2+5/4)*y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = \frac{c_1 x^3 \left(1 + \frac{1}{10} x^2 + \frac{1}{280} x^4 + O(x^6) \right) + c_2 \left(12 - 6x^2 - \frac{3}{2} x^4 + O(x^6) \right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 58

```

AsymptoticDSolveValue[x^2*y''[x]-x*y'[x]-(x^2+5/4)*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_1 \left(-\frac{x^{7/2}}{8} - \frac{x^{3/2}}{2} + \frac{1}{\sqrt{x}} \right) + c_2 \left(\frac{x^{13/2}}{280} + \frac{x^{9/2}}{10} + x^{5/2} \right)$$

4.56 problem 53

4.56.1 Maple step by step solution 2193

Internal problem ID [7277]

Internal file name [OUTPUT/6263_Sunday_June_05_2022_04_36_16_PM_43181729/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 53.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{4x^2 - 1}{4x^2}$$

Table 224: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{4x^2-1}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + \left(x^2 - \frac{1}{4}\right) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(x^2 - \frac{1}{4}\right) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} \left(-\frac{a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} \left(-\frac{a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - \frac{a_n x^{n+r}}{4} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r - \frac{a_0 x^r}{4} = 0$$

Or

$$\left(x^r r(-1+r) + x^r r - \frac{x^r}{4} \right) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$\frac{(4r^2 - 1) x^r}{4} = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - \frac{1}{4} = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = -\frac{1}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$\frac{(4r^2 - 1)x^r}{4} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sqrt{x} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{\sqrt{x}}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} - \frac{a_n}{4} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-2}}{4n^2 + 8nr + 4r^2 - 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+1)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{4}{4r^2 + 16r + 15}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= \sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -\frac{1}{2}} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + b_n(n+r) + b_{n-2} - \frac{b_n}{4} = 0 \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n \left(n - \frac{1}{2} \right) \left(n - \frac{3}{2} \right) + b_n \left(n - \frac{1}{2} \right) + b_{n-2} - \frac{b_n}{4} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{4b_{n-2}}{4n^2 + 8nr + 4r^2 - 1} \quad (5)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = -\frac{4b_{n-2}}{4n^2 - 4n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{4}{4r^2 + 16r + 15}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2 + 16r + 15}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{16}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= \sqrt{x}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6)}{\sqrt{x}} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6) \right)}{\sqrt{x}} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6) \right)}{\sqrt{x}} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6) \right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} + O(x^6) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} + O(x^6) \right)}{\sqrt{x}}$$

Verified OK.

4.56.1 Maple step by step solution

Let's solve

$$x^2y'' + xy' + \left(x^2 - \frac{1}{4}\right)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(4x^2-1)y}{4x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(4x^2-1)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{4x^2-1}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{1}{4}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2y'' + 4xy' + (4x^2 - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+2r)(-1+2r)x^r + a_1(3+2r)(1+2r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r+1)(2k+2r-1) + 4a_{k-2}) \right) x^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+2r)(-1+2r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \left\{ -\frac{1}{2}, \frac{1}{2} \right\}$$
- Each term must be 0

$$a_1(3+2r)(1+2r) = 0$$
- Solve for the dependent coefficient(s)

$$a_1 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_k(4k^2 + 8kr + 4r^2 - 1) + 4a_{k-2} = 0$$
- Shift index using $k- > k+2$

$$a_{k+2}(4(k+2)^2 + 8(k+2)r + 4r^2 - 1) + 4a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{4a_k}{4k^2 + 8kr + 4r^2 + 16k + 16r + 15}$$
- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}$$
- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}, a_1 = 0 \right]$$
- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}$$
- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}, a_1 = 0 \right]$$
- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = -\frac{4a_k}{4k^2+12k+8}, a_1 = 0, b_{k+2} = -\frac{4b_k}{4k^2+20k+24}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```

Order:=6;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1/4)*y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{6}x^2 + \frac{1}{120}x^4 + O(x^6) \right) x + c_2 \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 + O(x^6) \right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 58

```

AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1/4)*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_1 \left(\frac{x^{7/2}}{24} - \frac{x^{3/2}}{2} + \frac{1}{\sqrt{x}} \right) + c_2 \left(\frac{x^{9/2}}{120} - \frac{x^{5/2}}{6} + \sqrt{x} \right)$$

4.57 problem 54

4.57.1 Maple step by step solution 2208

Internal problem ID [7278]

Internal file name [OUTPUT/6264_Sunday_June_05_2022_04_36_18_PM_46684790/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 54.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$xy'' + (-x + 2)y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$xy'' + (-x + 2)y' - y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{x-2}{x}$$
$$q(x) = -\frac{1}{x}$$

Table 226: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{x-2}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$xy'' + (-x + 2)y' - y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + (-x+2) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} 2(n+r) a_n x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ \sum_{n=0}^{\infty} (-a_n x^{n+r}) &= \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r-1}) \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ & + \left(\sum_{n=0}^{\infty} 2(n+r) a_n x^{n+r-1} \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r-1}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + 2(n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + 2r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + 2r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} (1+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(1+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -1 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r}(1+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) + 2a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{n + 1 + r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{n + 1} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{2 + r}$$

Which for the root $r = 0$ becomes

$$a_1 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2+r}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(2 + r)(3 + r)}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2+r}$	$\frac{1}{2}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{(2+r)(4+r)(3+r)}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{24}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2+r}$	$\frac{1}{2}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{6}$
a_3	$\frac{1}{(2+r)(4+r)(3+r)}$	$\frac{1}{24}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(2+r)(3+r)(5+r)(4+r)}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2+r}$	$\frac{1}{2}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{6}$
a_3	$\frac{1}{(2+r)(4+r)(3+r)}$	$\frac{1}{24}$
a_4	$\frac{1}{(2+r)(3+r)(5+r)(4+r)}$	$\frac{1}{120}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{(2+r)(4+r)(3+r)(6+r)(5+r)}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{720}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{2+r}$	$\frac{1}{2}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{6}$
a_3	$\frac{1}{(2+r)(4+r)(3+r)}$	$\frac{1}{24}$
a_4	$\frac{1}{(2+r)(3+r)(5+r)(4+r)}$	$\frac{1}{120}$
a_5	$\frac{1}{(2+r)(4+r)(3+r)(6+r)(5+r)}$	$\frac{1}{720}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 + \frac{x}{2} + \frac{x^2}{6} + \frac{x^3}{24} + \frac{x^4}{120} + \frac{x^5}{720} + O(x^6) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= \frac{1}{2+r} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} \frac{1}{2+r} &= \lim_{r \rightarrow -1} \frac{1}{2+r} \\ &= 1 \end{aligned}$$

The limit is 1. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-1} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) - b_{n-1}(n+r-1) + 2(n+r)b_n - b_{n-1} = 0 \quad (4)$$

Which for for the root $r = -1$ becomes

$$b_n(n-1)(n-2) - b_{n-1}(n-2) + 2(n-1)b_n - b_{n-1} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = \frac{b_{n-1}}{n+1+r} \quad (5)$$

Which for the root $r = -1$ becomes

$$b_n = \frac{b_{n-1}}{n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{1}{2+r}$$

Which for the root $r = -1$ becomes

$$b_1 = 1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2+r}$	1

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{(2+r)(3+r)}$$

Which for the root $r = -1$ becomes

$$b_2 = \frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2+r}$	1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{1}{(2+r)(4+r)(3+r)}$$

Which for the root $r = -1$ becomes

$$b_3 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2+r}$	1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$
b_3	$\frac{1}{(2+r)(4+r)(3+r)}$	$\frac{1}{6}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{(2+r)(3+r)(5+r)(4+r)}$$

Which for the root $r = -1$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2+r}$	1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$
b_3	$\frac{1}{(2+r)(4+r)(3+r)}$	$\frac{1}{6}$
b_4	$\frac{1}{(2+r)(3+r)(5+r)(4+r)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{1}{(2+r)(4+r)(3+r)(6+r)(5+r)}$$

Which for the root $r = -1$ becomes

$$b_5 = \frac{1}{120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{2+r}$	1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$
b_3	$\frac{1}{(2+r)(4+r)(3+r)}$	$\frac{1}{6}$
b_4	$\frac{1}{(2+r)(3+r)(5+r)(4+r)}$	$\frac{1}{24}$
b_5	$\frac{1}{(2+r)(4+r)(3+r)(6+r)(5+r)}$	$\frac{1}{120}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + O(x^6)}{x} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\left(1 + \frac{x}{2} + \frac{x^2}{6} + \frac{x^3}{24} + \frac{x^4}{120} + \frac{x^5}{720} + O(x^6)\right) + \frac{c_2\left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + O(x^6)\right)}{x} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\left(1 + \frac{x}{2} + \frac{x^2}{6} + \frac{x^3}{24} + \frac{x^4}{120} + \frac{x^5}{720} + O(x^6)\right) + \frac{c_2\left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + O(x^6)\right)}{x} \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1\left(1 + \frac{x}{2} + \frac{x^2}{6} + \frac{x^3}{24} + \frac{x^4}{120} + \frac{x^5}{720} + O(x^6)\right) \\ &\quad + \frac{c_2\left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + O(x^6)\right)}{x} \end{aligned} \tag{1}$$

Verification of solutions

$$y = c_1\left(1 + \frac{x}{2} + \frac{x^2}{6} + \frac{x^3}{24} + \frac{x^4}{120} + \frac{x^5}{720} + O(x^6)\right) + \frac{c_2\left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + O(x^6)\right)}{x}$$

Verified OK.

4.57.1 Maple step by step solution

Let's solve

$$xy'' + (-x + 2)y' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{x} + \frac{(x-2)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(x-2)y'}{x} - \frac{y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{x-2}{x}, P_3(x) = -\frac{1}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$xy'' + (-x + 2)y' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k- \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m}(k+1-m+r)x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r-1}$$

- Shift index using $k- > k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)(k+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0r(1+r)x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1}(k+1+r)(k+2+r) - a_k(k+1+r))x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
- $r(1+r) = 0$
- Values of r that satisfy the indicial equation
- $r \in \{-1, 0\}$
- Each term in the series must be 0, giving the recursion relation
- $(k+1+r)(a_{k+1}(k+2+r) - a_k) = 0$
- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k}{k+2+r}$$

- Recursion relation for $r = -1$

$$a_{k+1} = \frac{a_k}{k+1}$$

- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+1} = \frac{a_k}{k+1} \right]$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k}{k+2}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = \frac{a_k}{k+2} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k x^k \right), a_{k+1} = \frac{a_k}{k+1}, b_{k+1} = \frac{b_k}{k+2} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+(2-x)*diff(y(x),x)-y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 + \frac{1}{2}x + \frac{1}{6}x^2 + \frac{1}{24}x^3 + \frac{1}{120}x^4 + \frac{1}{720}x^5 + O(x^6) \right) + \frac{c_2 \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + O(x^6) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 62

```
AsymptoticDSolveValue[x*y'[x]+(2-x)*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{24} + \frac{x^2}{6} + \frac{x}{2} + \frac{1}{x} + 1 \right) + c_2 \left(\frac{x^4}{120} + \frac{x^3}{24} + \frac{x^2}{6} + \frac{x}{2} + 1 \right)$$

4.58 problem 55

4.58.1 Maple step by step solution 2219

Internal problem ID [7279]

Internal file name [OUTPUT/6265_Sunday_June_05_2022_04_36_21_PM_68300846/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 55.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$2x^2y'' + 3xy' - y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + 3xy' - y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3}{2x}$$
$$q(x) = -\frac{1}{2x^2}$$

Table 228: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + 3xy' - y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 3x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + 3x^{n+r} a_n (n+r) - a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r(-1+r) + 3x^r a_0 r - a_0 x^r = 0$$

Or

$$(2x^r r(-1+r) + 3x^r r - x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 + r - 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 + r - 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = -1$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 + r - 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{3}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$
$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n-1}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $0 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + 3a_n(n+r) - a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = 0 \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= \sqrt{x}(1 + O(x^6))
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $0 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) + 3b_n(n+r) - b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = 0 \quad (4)$$

Which for the root $r = -1$ becomes

$$b_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= \sqrt{x}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 + O(x^6)}{x} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\sqrt{x}(1 + O(x^6)) + \frac{c_2(1 + O(x^6))}{x} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\sqrt{x}(1 + O(x^6)) + \frac{c_2(1 + O(x^6))}{x} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x}(1 + O(x^6)) + \frac{c_2(1 + O(x^6))}{x} \tag{1}$$

Verification of solutions

$$y = c_1\sqrt{x}(1 + O(x^6)) + \frac{c_2(1 + O(x^6))}{x}$$

Verified OK.

4.58.1 Maple step by step solution

Let's solve

$$2x^2y'' + 3xy' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{3y'}{2x} + \frac{y}{2x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{2x} - \frac{y}{2x^2} = 0$$

- Multiply by denominators of the ODE

$$2x^2y'' + 3xy' - y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$2x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + 3 \frac{d}{dt}y(t) - y(t) = 0$$

- Simplify

$$2 \frac{d^2}{dt^2}y(t) + \frac{d}{dt}y(t) - y(t) = 0$$

- Isolate 2nd derivative

$$\frac{d^2}{dt^2}y(t) = -\frac{d}{dt}\frac{y(t)}{2} + \frac{y(t)}{2}$$

- Group terms with $y(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dt^2}y(t) + \frac{d}{dt}\frac{y(t)}{2} - \frac{y(t)}{2} = 0$$

- Characteristic polynomial of ODE

$$r^2 + \frac{1}{2}r - \frac{1}{2} = 0$$

- Factor the characteristic polynomial

$$\frac{(r+1)(2r-1)}{2} = 0$$

- Roots of the characteristic polynomial

$$r = \left(-1, \frac{1}{2}\right)$$

- 1st solution of the ODE

$$y_1(t) = e^{-t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{\frac{t}{2}}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-t} + c_2 e^{\frac{t}{2}}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x} + c_2 \sqrt{x}$$

- Simplify

$$y = \frac{c_1}{x} + c_2 \sqrt{x}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
Order:=6;  
dsolve(2*x^2*diff(y(x),x$2)+3*x*diff(y(x),x)-y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = \frac{x^{\frac{3}{2}}c_2 + c_1}{x} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 18

```
AsymptoticDSolveValue[2*x^2*y'[x]+3*x*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1\sqrt{x} + \frac{c_2}{x}$$

4.59 problem 56

4.59.1 Maple step by step solution 2228

Internal problem ID [7280]

Internal file name [OUTPUT/6266_Sunday_June_05_2022_04_36_23_PM_27616334/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 56.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$2x^2y'' + 5xy' + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + 5xy' + 4y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{5}{2x}$$
$$q(x) = \frac{2}{x^2}$$

Table 230: Table $p(x), q(x)$ singularities.

$p(x) = \frac{5}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{2}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + 5xy' + 4y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 5x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 4 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 4a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 4a_n x^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + 5x^{n+r} a_n (n+r) + 4a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$2x^r a_0 r (-1+r) + 5x^r a_0 r + 4a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) + 5x^r r + 4x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 + 3r + 4) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 + 3r + 4 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = -\frac{3}{4} + \frac{i\sqrt{23}}{4}$$

$$r_2 = -\frac{3}{4} - \frac{i\sqrt{23}}{4}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 + 3r + 4) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n - \frac{3}{4} + \frac{i\sqrt{23}}{4}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n - \frac{3}{4} - \frac{i\sqrt{23}}{4}}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $0 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + 5a_n(n+r) + 4a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = 0 \quad (4)$$

Which for the root $r = -\frac{3}{4} + \frac{i\sqrt{23}}{4}$ becomes

$$a_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = -\frac{3}{4} + \frac{i\sqrt{23}}{4}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^{-\frac{3}{4} + \frac{i\sqrt{23}}{4}} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^{-\frac{3}{4} + \frac{i\sqrt{23}}{4}} (1 + O(x^6)) \end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$y_2(x) = x^{-\frac{3}{4} - \frac{i\sqrt{23}}{4}} (1 + O(x^6))$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^{-\frac{3}{4} + \frac{i\sqrt{23}}{4}} (1 + O(x^6)) + c_2x^{-\frac{3}{4} - \frac{i\sqrt{23}}{4}} (1 + O(x^6)) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x^{-\frac{3}{4} + \frac{i\sqrt{23}}{4}} (1 + O(x^6)) + c_2x^{-\frac{3}{4} - \frac{i\sqrt{23}}{4}} (1 + O(x^6)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1x^{-\frac{3}{4} + \frac{i\sqrt{23}}{4}} (1 + O(x^6)) + c_2x^{-\frac{3}{4} - \frac{i\sqrt{23}}{4}} (1 + O(x^6)) \quad (1)$$

Verification of solutions

$$y = c_1x^{-\frac{3}{4} + \frac{i\sqrt{23}}{4}} (1 + O(x^6)) + c_2x^{-\frac{3}{4} - \frac{i\sqrt{23}}{4}} (1 + O(x^6))$$

Verified OK.

4.59.1 Maple step by step solution

Let's solve

$$2x^2y'' + 5xy' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{5y'}{2x} - \frac{2y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{5y'}{2x} + \frac{2y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$2x^2y'' + 5xy' + 4y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$2x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + 5 \frac{d}{dt}y(t) + 4y(t) = 0$$

- Simplify

$$2 \frac{d^2}{dt^2}y(t) + 3 \frac{d}{dt}y(t) + 4y(t) = 0$$

- Isolate 2nd derivative

$$\frac{d^2}{dt^2}y(t) = -\frac{3\frac{d}{dt}y(t)}{2} - 2y(t)$$

- Group terms with $y(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dt^2}y(t) + \frac{3\frac{d}{dt}y(t)}{2} + 2y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + \frac{3}{2}r + 2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-\frac{3}{2}) \pm \left(\sqrt{-\frac{23}{4}}\right)}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{3}{4} - \frac{i\sqrt{23}}{4}, -\frac{3}{4} + \frac{i\sqrt{23}}{4}\right)$$

- 1st solution of the ODE

$$y_1(t) = e^{-\frac{3t}{4}} \cos\left(\frac{\sqrt{23}t}{4}\right)$$

- 2nd solution of the ODE

$$y_2(t) = e^{-\frac{3t}{4}} \sin\left(\frac{\sqrt{23}t}{4}\right)$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-\frac{3t}{4}} \cos\left(\frac{\sqrt{23}t}{4}\right) + c_2 e^{-\frac{3t}{4}} \sin\left(\frac{\sqrt{23}t}{4}\right)$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1 \cos\left(\frac{\sqrt{23} \ln(x)}{4}\right)}{x^{\frac{3}{4}}} + \frac{c_2 \sin\left(\frac{\sqrt{23} \ln(x)}{4}\right)}{x^{\frac{3}{4}}}$$

- Simplify

$$y = \frac{c_1 \cos\left(\frac{\sqrt{23} \ln(x)}{4}\right)}{x^{\frac{3}{4}}} + \frac{c_2 \sin\left(\frac{\sqrt{23} \ln(x)}{4}\right)}{x^{\frac{3}{4}}}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
Order:=6;  
dsolve(2*x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+4*y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = \frac{x^{-\frac{i\sqrt{23}}{4}} c_1 + x^{\frac{i\sqrt{23}}{4}} c_2}{x^{\frac{3}{4}}} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 44

```
AsymptoticDSolveValue[2*x^2*y'[x]+5*x*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 x^{\frac{1}{4}(-3+i\sqrt{23})} + c_2 x^{\frac{1}{4}(-3-i\sqrt{23})}$$

4.60 problem 57

4.60.1 Maple step by step solution 2239

Internal problem ID [7281]

Internal file name [OUTPUT/6267_Sunday_June_05_2022_04_36_26_PM_67548285/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 57.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

[[_Emden , _Fowler]]

$$x^2y'' + 3xy' + 4yx^4 = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$x^2y'' + 3xy' + 4yx^4 = 0$$

Or

$$x(4yx^3 + xy'' + 3y') = 0$$

For $x \neq 0$ the above simplifies to

$$4yx^3 + xy'' + 3y' = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + 3xy' + 4yx^4 = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3}{x}$$

$$q(x) = 4x^2$$

Table 232: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = 4x^2$	
singularity	type
$x = \infty$	“regular”
$x = -\infty$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty, -\infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + 3xy' + 4yx^4 = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 3x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 4 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x^4 = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 4x^{4+n+r} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 4x^{4+n+r} a_n = \sum_{n=4}^{\infty} 4a_{n-4} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \left(\sum_{n=4}^{\infty} 4a_{n-4} x^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r)(n+r-1) + 3x^{n+r} a_n (n+r) = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + 3x^r a_0 r = 0$$

Or

$$(x^r r(-1+r) + 3x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r(2+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(2+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -2 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r(2+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x^2} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-2} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = 0$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = 0$$

For $4 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 3a_n(n+r) + 4a_{n-4} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-4}}{n^2 + 2nr + r^2 + 2n + 2r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{4a_{n-4}}{n(n+2)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = -\frac{4}{r^2 + 10r + 24}$$

Which for the root $r = 0$ becomes

$$a_4 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	$-\frac{4}{r^2+10r+24}$	$-\frac{1}{6}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	$-\frac{4}{r^2+10r+24}$	$-\frac{1}{6}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 - \frac{x^4}{6} + O(x^6) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 2$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_2(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_2 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned}\lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -2} 0 \\ &= 0\end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned}y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-2}\end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

Substituting $n = 2$ in Eq(3) gives

$$b_2 = 0$$

Substituting $n = 3$ in Eq(3) gives

$$b_3 = 0$$

For $4 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + 3b_n(n+r) + 4b_{n-4} = 0 \quad (4)$$

Which for the root $r = -2$ becomes

$$b_n(n-2)(n-3) + 3b_n(n-2) + 4b_{n-4} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{4b_{n-4}}{n^2 + 2nr + r^2 + 2n + 2r} \quad (5)$$

Which for the root $r = -2$ becomes

$$b_n = -\frac{4b_{n-4}}{n^2 - 2n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -2$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = -\frac{4}{r^2 + 10r + 24}$$

Which for the root $r = -2$ becomes

$$b_4 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	$-\frac{4}{r^2+10r+24}$	$-\frac{1}{2}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	$-\frac{4}{r^2+10r+24}$	$-\frac{1}{2}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}y_2(x) &= 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= \frac{1 - \frac{x^4}{2} + O(x^6)}{x^2}\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\left(1 - \frac{x^4}{6} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^4}{2} + O(x^6)\right)}{x^2}\end{aligned}$$

Hence the final solution is

$$\begin{aligned}y &= y_h \\ &= c_1\left(1 - \frac{x^4}{6} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^4}{2} + O(x^6)\right)}{x^2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1\left(1 - \frac{x^4}{6} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^4}{2} + O(x^6)\right)}{x^2} \quad (1)$$

Verification of solutions

$$y = c_1\left(1 - \frac{x^4}{6} + O(x^6)\right) + \frac{c_2\left(1 - \frac{x^4}{2} + O(x^6)\right)}{x^2}$$

Verified OK.

4.60.1 Maple step by step solution

Let's solve

$$x^2y'' + 3xy' + 4yx^4 = 0$$

- Highest derivative means the order of the ODE is 2
 y''
- Isolate 2nd derivative

$$y'' = -\frac{3y'}{x} - 4x^2y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{x} + 4x^2y = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{3}{x}, P_3(x) = 4x^2]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 3$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4yx^3 + xy'' + 3y' = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^3 \cdot y$ to series expansion

$$x^3 \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+3}$$

- Shift index using $k \rightarrow k - 3$

$$x^3 \cdot y = \sum_{k=3}^{\infty} a_{k-3} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r-1}$$

- Shift index using $k- > k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)(k+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0r(2+r)x^{-1+r} + a_1(1+r)(3+r)x^r + a_2(2+r)(4+r)x^{1+r} + a_3(3+r)(5+r)x^{2+r} + \left(\sum_{k=3}^{\infty} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-2, 0\}$$

- The coefficients of each power of x must be 0

$$[a_1(1+r)(3+r) = 0, a_2(2+r)(4+r) = 0, a_3(3+r)(5+r) = 0]$$

- Solve for the dependent coefficient(s)

$$\{a_1 = 0, a_2 = 0, a_3 = 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1+r)(k+r+3) + 4a_{k-3} = 0$$

- Shift index using $k- > k+3$

$$a_{k+4}(k+4+r)(k+6+r) + 4a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+4} = -\frac{4a_k}{(k+4+r)(k+6+r)}$$

- Recursion relation for $r = -2$

$$a_{k+4} = -\frac{4a_k}{(k+2)(k+4)}$$

- Solution for $r = -2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-2}, a_{k+4} = -\frac{4a_k}{(k+2)(k+4)}, a_1 = 0, a_2 = 0, a_3 = 0 \right]$$

- Recursion relation for $r = 0$

$$a_{k+4} = -\frac{4a_k}{(k+4)(k+6)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = -\frac{4a_k}{(k+4)(k+6)}, a_1 = 0, a_2 = 0, a_3 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-2} \right) + \left(\sum_{k=0}^{\infty} b_k x^k \right), a_{k+4} = -\frac{4a_k}{(k+2)(k+4)}, a_1 = 0, a_2 = 0, a_3 = 0, b_{k+4} = -\frac{4b_k}{(k+4)(k+6)} \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```

Order:=6;
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+4*x^4*y(x) = 0,y(x),type='series',x=0);

```

$$y(x) = c_1 \left(1 - \frac{1}{6}x^4 + O(x^6) \right) + \frac{c_2(-2 + x^4 + O(x^6))}{x^2}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 30

```

AsymptoticDSolveValue[x^2*y''[x]+3*x*y'[x]+4*x^4*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_2 \left(1 - \frac{x^4}{6} \right) + c_1 \left(\frac{1}{x^2} - \frac{x^2}{2} \right)$$

4.61 problem 58

4.61.1 Maple step by step solution 2255

Internal problem ID [7282]

Internal file name [OUTPUT/6268_Sunday_June_05_2022_04_36_28_PM_22985598/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 58.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

[[_Emden , _Fowler]]

$$x^2y'' - yx = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$x^2y'' - yx = 0$$

Or

$$x(xy'' - y) = 0$$

For $x \neq 0$ the above simplifies to

$$xy'' - y = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' - yx = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$

$$q(x) = -\frac{1}{x}$$

Table 234: Table $p(x), q(x)$ singularities.

$p(x) = 0$		$q(x) = -\frac{1}{x}$	
singularity	type	singularity	type
		$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' - yx = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) = 0$$

When $n=0$ the above becomes

$$x^r a_0 r (-1+r) = 0$$

Or

$$x^r a_0 r (-1+r) = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r (-1+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(-1+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 1$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r(-1+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{1+n}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_{n-1} = 0 \tag{3}$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{(n+r)(n+r-1)} \tag{4}$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-1}}{(1+n)n} \tag{5}$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{(1+r)r}$$

Which for the root $r = 1$ becomes

$$a_1 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(1+r)r}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(1+r)^2 r (2+r)}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{12}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(1+r)r}$	$\frac{1}{2}$
a_2	$\frac{1}{(1+r)^2 r (2+r)}$	$\frac{1}{12}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{(1+r)^2 r (2+r)^2 (3+r)}$$

Which for the root $r = 1$ becomes

$$a_3 = \frac{1}{144}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(1+r)r}$	$\frac{1}{2}$
a_2	$\frac{1}{(1+r)^2 r(2+r)}$	$\frac{1}{12}$
a_3	$\frac{1}{(1+r)^2 r(2+r)^2(3+r)}$	$\frac{1}{144}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(1+r)^2 r(2+r)^2(3+r)^2(4+r)}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{2880}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(1+r)r}$	$\frac{1}{2}$
a_2	$\frac{1}{(1+r)^2 r(2+r)}$	$\frac{1}{12}$
a_3	$\frac{1}{(1+r)^2 r(2+r)^2(3+r)}$	$\frac{1}{144}$
a_4	$\frac{1}{(1+r)^2 r(2+r)^2(3+r)^2(4+r)}$	$\frac{1}{2880}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{(1+r)^2 r(2+r)^2(3+r)^2(4+r)^2(5+r)}$$

Which for the root $r = 1$ becomes

$$a_5 = \frac{1}{86400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{(1+r)r}$	$\frac{1}{2}$
a_2	$\frac{1}{(1+r)^2 r(2+r)}$	$\frac{1}{12}$
a_3	$\frac{1}{(1+r)^2 r(2+r)^2(3+r)}$	$\frac{1}{144}$
a_4	$\frac{1}{(1+r)^2 r(2+r)^2(3+r)^2(4+r)}$	$\frac{1}{2880}$
a_5	$\frac{1}{(1+r)^2 r(2+r)^2(3+r)^2(4+r)^2(5+r)}$	$\frac{1}{86400}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\
 &= x\left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned}
 a_N &= a_1 \\
 &= \frac{1}{(1+r)r}
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \lim_{r \rightarrow r_2} \frac{1}{(1+r)r} &= \lim_{r \rightarrow 0} \frac{1}{(1+r)r} \\
 &= \text{undefined}
 \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2}\right)$$

Therefore

$$\begin{aligned} \frac{d}{dx}y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2}y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x^2y'' - yx = 0$ gives

$$\begin{aligned} &x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\ &\quad - \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) x = 0 \end{aligned}$$

Which can be written as

$$\begin{aligned} &\left((x^2y_1''(x) - y_1(x)x) \ln(x) + x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) \right) C \\ &\quad + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) - \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) x = 0 \end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$x^2y_1''(x) - y_1(x)x = 0$$

Eq (7) simplifies to

$$x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) C + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) - \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) x = 0 \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\left(2 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x - \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 - \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) x = 0 \quad (9)$$

Since $r_1 = 1$ and $r_2 = 0$ then the above becomes

$$\left(2 \left(\sum_{n=0}^{\infty} x^n a_n (1+n) \right) x - \left(\sum_{n=0}^{\infty} a_n x^{1+n} \right) \right) C + \left(\sum_{n=0}^{\infty} x^{-2+n} b_n n (n-1) \right) x^2 - \left(\sum_{n=0}^{\infty} b_n x^n \right) x = 0 \quad (10)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2C x^{1+n} a_n (1+n) \right) + \sum_{n=0}^{\infty} (-C a_n x^{1+n}) + \left(\sum_{n=0}^{\infty} n x^n b_n (n-1) \right) + \sum_{n=0}^{\infty} (-x^{1+n} b_n) = 0 \quad (2A)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the

power and the corresponding index gives

$$\begin{aligned}\sum_{n=0}^{\infty} 2C x^{1+n} a_n (1+n) &= \sum_{n=1}^{\infty} 2C a_{n-1} n x^n \\ \sum_{n=0}^{\infty} (-C a_n x^{1+n}) &= \sum_{n=1}^{\infty} (-C a_{n-1} x^n) \\ \sum_{n=0}^{\infty} (-x^{1+n} b_n) &= \sum_{n=1}^{\infty} (-b_{n-1} x^n)\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=1}^{\infty} 2C a_{n-1} n x^n \right) + \sum_{n=1}^{\infty} (-C a_{n-1} x^n) + \left(\sum_{n=0}^{\infty} n x^n b_n (n-1) \right) + \sum_{n=1}^{\infty} (-b_{n-1} x^n) = 0 \quad (2B)$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = N$, where $N = 1$ which is the difference between the two roots, we are free to choose $b_1 = 0$. Hence for $n = 1$, Eq (2B) gives

$$C - 1 = 0$$

Which is solved for C . Solving for C gives

$$C = 1$$

For $n = 2$, Eq (2B) gives

$$3C a_1 - b_1 + 2b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$2b_2 + \frac{3}{2} = 0$$

Solving the above for b_2 gives

$$b_2 = -\frac{3}{4}$$

For $n = 3$, Eq (2B) gives

$$5C a_2 - b_2 + 6b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$6b_3 + \frac{7}{6} = 0$$

Solving the above for b_3 gives

$$b_3 = -\frac{7}{36}$$

For $n = 4$, Eq (2B) gives

$$7Ca_3 - b_3 + 12b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$12b_4 + \frac{35}{144} = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{35}{1728}$$

For $n = 5$, Eq (2B) gives

$$9Ca_4 - b_4 + 20b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$20b_5 + \frac{101}{4320} = 0$$

Solving the above for b_5 gives

$$b_5 = -\frac{101}{86400}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = 1$ and all b_n , then the second solution becomes

$$y_2(x) = 1 \left(x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \right) \ln(x) \\ + 1 - \frac{3x^2}{4} - \frac{7x^3}{36} - \frac{35x^4}{1728} - \frac{101x^5}{86400} + O(x^6)$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \\
 &\quad + c_2 \left(1 \left(x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \right) \ln(x) + 1 - \frac{3x^2}{4} \right. \\
 &\quad \left. - \frac{7x^3}{36} - \frac{35x^4}{1728} - \frac{101x^5}{86400} + O(x^6) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \\
 &\quad + c_2 \left(x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \ln(x) + 1 - \frac{3x^2}{4} - \frac{7x^3}{36} \right. \\
 &\quad \left. - \frac{35x^4}{1728} - \frac{101x^5}{86400} + O(x^6) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \\
 &\quad + c_2 \left(x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \ln(x) + 1 - \frac{3x^2}{4} - \frac{7x^3}{36} \right. \\
 &\quad \left. - \frac{35x^4}{1728} - \frac{101x^5}{86400} + O(x^6) \right)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \\
 &\quad + c_2 \left(x \left(1 + \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} + \frac{x^4}{2880} + \frac{x^5}{86400} + O(x^6) \right) \ln(x) + 1 - \frac{3x^2}{4} - \frac{7x^3}{36} \right. \\
 &\quad \left. - \frac{35x^4}{1728} - \frac{101x^5}{86400} + O(x^6) \right)
 \end{aligned}$$

Verified OK.

4.61.1 Maple step by step solution

Let's solve

$$x^2 y'' - yx = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = 0, P_3(x) = -\frac{1}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$xy'' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k- > k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)(k+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-1+r)x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1}(k+1+r)(k+r) - a_k)x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-1+r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \{0, 1\}$$
- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1+r)(k+r) - a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k}{(k+1+r)(k+r)}$$
- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k}{(k+1)k}$$
- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = \frac{a_k}{(k+1)k} \right]$$
- Recursion relation for $r = 1$

$$a_{k+1} = \frac{a_k}{(k+2)(k+1)}$$
- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = \frac{a_k}{(k+2)(k+1)} \right]$$
- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1} \right), a_{k+1} = \frac{a_k}{(k+1)k}, b_{k+1} = \frac{b_k}{(k+2)(k+1)} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 58

```
Order:=6;  
dsolve(x^2*diff(y(x),x$2)-x*y(x) = 0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 + \frac{1}{2}x + \frac{1}{12}x^2 + \frac{1}{144}x^3 + \frac{1}{2880}x^4 + \frac{1}{86400}x^5 + O(x^6) \right) \\ + c_2 \left(\ln(x) \left(x + \frac{1}{2}x^2 + \frac{1}{12}x^3 + \frac{1}{144}x^4 + \frac{1}{2880}x^5 + O(x^6) \right) \right. \\ \left. + \left(1 - \frac{3}{4}x^2 - \frac{7}{36}x^3 - \frac{35}{1728}x^4 - \frac{101}{86400}x^5 + O(x^6) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 85

```
AsymptoticDSolveValue[x^2*y''[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{144}x(x^3 + 12x^2 + 72x + 144) \log(x) \right. \\ \left. + \frac{-47x^4 - 480x^3 - 2160x^2 - 1728x + 1728}{1728} \right) + c_2 \left(\frac{x^5}{2880} + \frac{x^4}{144} + \frac{x^3}{12} + \frac{x^2}{2} + x \right)$$

4.62 problem 59

Internal problem ID [7283]

Internal file name [OUTPUT/6269_Sunday_June_05_2022_04_36_33_PM_2867427/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 59.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$(1 - x^2) y'' + y' + y = x e^x$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \dots \\ &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \dots \\ &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (350)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (351)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{-y' - y + x e^x}{x^2 - 1}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{(x^2 - 2x) y' + (-x^3 + x^2 + 1) e^x + (1 - 2x) y}{(x^2 - 1)^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{(-4x^3 + 8x^2 - 2x + 1) y' + (-x^5 + 2x^4 - 2x^3 + 5x^2 - 6x - 1) e^x + y(7x^2 - 6x + 2)}{(x^2 - 1)^3} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(19x^4 - 42x^3 + 25x^2 - 18x + 1) y' + (-x^7 + 3x^6 - 5x^5 + 18x^4 - 40x^3 + 32x^2 + x + 7) e^x - 32(x^3 - 1)}{(x^2 - 1)^4} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{(-108x^5 + 267x^4 - 264x^3 + 246x^2 - 48x + 12) y' + (-x^9 + 4x^8 - 10x^7 + 37x^6 - 144x^5 + 281x^4 - 192x^3 + 64x^2 - 8x + 1) e^x - 32(7x^3 - 6x^2 + 2x - 1)}{(x^2 - 1)^5} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$F_0 = -y'(0) - y(0)$$

$$F_1 = y(0) + 1$$

$$F_2 = 1 - 2y(0) - y'(0)$$

$$F_3 = 7 + 7y(0) + y'(0)$$

$$F_4 = 8 - 29y(0) - 12y'(0)$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{7}{120}x^5 - \frac{29}{720}x^6\right) y(0) \\ + \left(x - \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{120}x^5 - \frac{1}{60}x^6\right) y'(0) + \frac{x^3}{6} + \frac{x^4}{24} + \frac{7x^5}{120} + \frac{x^6}{90} + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$(1 - x^2) y'' + y' + y = x e^x$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$(1 - x^2) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x e^x \quad (1)$$

Expanding $x e^x$ as Taylor series around $x = 0$ and keeping only the first 6 terms gives

$$x e^x = x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5 + \dots \\ = x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5$$

Hence the ODE in Eq (1) becomes

$$(1 - x^2) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ = x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5$$

Which simplifies to

$$\begin{aligned} \sum_{n=2}^{\infty} (-x^n a_n n(n-1)) + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (2) \\ = x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5 \end{aligned}$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \\ \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} \sum_{n=2}^{\infty} (-x^n a_n n(n-1)) + \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) \quad (3) \\ + \left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5 \end{aligned}$$

$n = 0$ gives

$$2a_2 + a_1 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2} - \frac{a_1}{2}$$

$n = 1$ gives

$$(6a_3 + 2a_2 + a_1) x = x$$

$$6a_3 + 2a_2 + a_1 = 1$$

Which after substituting earlier equations, simplifies to

$$a_3 = \frac{a_0}{6} + \frac{1}{6}$$

For $2 \leq n$, the recurrence equation is

$$(-na_n(n-1) + (n+2)a_{n+2}(n+1) + (n+1)a_{n+1} + a_n)x^n = x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5 \quad (4)$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned} (-a_2 + 12a_4 + 3a_3)x^2 &= x^2 \\ -a_2 + 12a_4 + 3a_3 &= 1 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{24} - \frac{a_0}{12} - \frac{a_1}{24}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned} (-5a_3 + 20a_5 + 4a_4)x^3 &= \frac{x^3}{2} \\ -5a_3 + 20a_5 + 4a_4 &= \frac{1}{2} \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{7}{120} + \frac{7a_0}{120} + \frac{a_1}{120}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned} (-11a_4 + 30a_6 + 5a_5)x^4 &= \frac{x^4}{6} \\ -11a_4 + 30a_6 + 5a_5 &= \frac{1}{6} \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{1}{90} - \frac{29a_0}{720} - \frac{a_1}{60}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned} (-19a_5 + 42a_7 + 6a_6) x^5 &= \frac{x^5}{24} \\ -19a_5 + 42a_7 + 6a_6 &= \frac{1}{24} \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{13}{504} + \frac{9a_0}{280} + \frac{31a_1}{5040}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x + \left(-\frac{a_0}{2} - \frac{a_1}{2}\right) x^2 + \left(\frac{a_0}{6} + \frac{1}{6}\right) x^3 \\ &\quad + \left(\frac{1}{24} - \frac{a_0}{12} - \frac{a_1}{24}\right) x^4 + \left(\frac{7}{120} + \frac{7a_0}{120} + \frac{a_1}{120}\right) x^5 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{7}{120}x^5\right) a_0 \\ &\quad + \left(x - \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{120}x^5\right) a_1 + \frac{x^3}{6} + \frac{x^4}{24} + \frac{7x^5}{120} + O(x^6) \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{7}{120}x^5\right) c_1 \\ &\quad + \left(x - \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{120}x^5\right) c_2 + \frac{x^3}{6} + \frac{x^4}{24} + \frac{7x^5}{120} + O(x^6) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{7}{120}x^5 - \frac{29}{720}x^6\right) y(0) + \left(x - \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{120}x^5 - \frac{1}{60}x^6\right) y'(0) + \frac{x^3}{6} + \frac{x^4}{24} + \frac{7x^5}{120} + \frac{x^6}{90} + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{7}{120}x^5\right) c_1 + \left(x - \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{120}x^5\right) c_2 + \frac{x^3}{6} + \frac{x^4}{24} + \frac{7x^5}{120} + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{7}{120}x^5 - \frac{29}{720}x^6\right) y(0) + \left(x - \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{120}x^5 - \frac{1}{60}x^6\right) y'(0) + \frac{x^3}{6} + \frac{x^4}{24} + \frac{7x^5}{120} + \frac{x^6}{90} + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{7}{120}x^5\right) c_1 + \left(x - \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{120}x^5\right) c_2 + \frac{x^3}{6} + \frac{x^4}{24} + \frac{7x^5}{120} + O(x^6)$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 53

```
Order:=6;
dsolve((1-x^2)*diff(y(x),x$2)+diff(y(x),x)+y(x)=x*exp(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{7}{120}x^5\right) y(0) \\ + \left(x - \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{120}x^5\right) D(y)(0) + \frac{x^3}{6} + \frac{x^4}{24} + \frac{7x^5}{120} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 63

```
AsymptoticDSolveValue[(1-x^2)*y'[x]+y'[x]+y[x]==x*Exp[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} - \frac{x^4}{24} - \frac{x^2}{2} + x \right) + c_1 \left(\frac{7x^5}{120} - \frac{x^4}{12} + \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

4.63 problem 60

4.63.1 Solving as quadrature ode 2268

4.63.2 Maple step by step solution 2269

Internal problem ID [7284]

Internal file name [OUTPUT/6270_Sunday_June_05_2022_04_36_36_PM_75428236/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 60.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "quadrature"

Maple gives the following as the ode type

[_quadrature]

$$y' - y(1 - y^2) = 0$$

4.63.1 Solving as quadrature ode

Integrating both sides gives

$$\int -\frac{1}{y(y^2 - 1)} dy = \int dx$$
$$\ln(y) - \frac{\ln(y+1)}{2} - \frac{\ln(-1+y)}{2} = x + c_1$$

Raising both side to exponential gives

$$e^{\ln(y) - \frac{\ln(y+1)}{2} - \frac{\ln(-1+y)}{2}} = e^{x+c_1}$$

Which simplifies to

$$\frac{y}{\sqrt{y+1}\sqrt{-1+y}} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{c_2^2 e^{2x} - \sqrt{e^{4x} c_2^4 - c_2^2 e^{2x} - 1}}{c_2^2 e^{2x} - 1} - 1 \quad (1)$$




Figure 134: Slope field plot

Verification of solutions

$$y = \frac{c_2^2 e^{2x} - \sqrt{e^{4x} c_2^4 - c_2^2 e^{2x} - 1}}{c_2^2 e^{2x} - 1} - 1$$

Verified OK.

4.63.2 Maple step by step solution

Let's solve

$$y' - y(1 - y^2) = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y(1-y^2)} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y(1-y^2)} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\ln(y) - \frac{\ln(y+1)}{2} - \frac{\ln(-1+y)}{2} = x + c_1$$

- Solve for y

$$\left\{ y = \frac{\sqrt{(e^{2c_1+2x}-1)e^{2c_1+2x}}}{e^{2c_1+2x}-1}, y = -\frac{\sqrt{(e^{2c_1+2x}-1)e^{2c_1+2x}}}{e^{2c_1+2x}-1} \right\}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)=y(x)*(1-y(x)^2),y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{e^{-2x}c_1 + 1}}$$
$$y(x) = -\frac{1}{\sqrt{e^{-2x}c_1 + 1}}$$

✓ Solution by Mathematica

Time used: 0.787 (sec). Leaf size: 100

```
DSolve[y'[x]==y[x]*(1-y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^x}{\sqrt{e^{2x} + e^{2c_1}}}$$
$$y(x) \rightarrow \frac{e^x}{\sqrt{e^{2x} + e^{2c_1}}}$$
$$y(x) \rightarrow -1$$
$$y(x) \rightarrow 0$$
$$y(x) \rightarrow 1$$
$$y(x) \rightarrow -\frac{e^x}{\sqrt{e^{2x}}}$$
$$y(x) \rightarrow \frac{e^x}{\sqrt{e^{2x}}}$$

4.64 problem 61

4.64.1 Solving as second order ode lagrange adjoint equation method od2272

Internal problem ID [7285]

Internal file name [OUTPUT/6271_Sunday_June_05_2022_04_36_40_PM_63226158/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 61.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$\frac{xy''}{1-x} + y = \frac{1}{1-x}$$

4.64.1 Solving as second order ode lagrange adjoint equation method ode

In normal form the ode

$$-\frac{xy''}{x-1} + y = \frac{1}{1-x} \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 0 \\ q(x) &= \frac{1-x}{x} \\ r(x) &= \frac{1}{x} \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned}\xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + \left(\frac{(1-x)\xi(x)}{x}\right) &= 0 \\ \xi''(x) + \frac{(1-x)\xi(x)}{x} &= 0\end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$x^2\xi''(x) + (-x^2 + x)\xi(x) = 0 \quad (1)$$

Bessel ode has the form

$$x^2\xi''(x) + \xi'(x)x + (-n^2 + x^2)\xi(x) = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2\xi''(x) + (1 - 2\alpha)x\xi'(x) + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)\xi(x) = 0 \quad (3)$$

With the standard solution

$$\xi(x) = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= 2 \\ n &= -1 \\ \gamma &= \frac{1}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$\xi(x) = -c_1\sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{BesselY}(1, 2\sqrt{x})$$

The original ode (2) now reduces to first order ode

$$\begin{aligned}y' - \frac{y\left(-\frac{c_3 \text{BesselJ}(1, 2\sqrt{x})}{2\sqrt{x}} - c_3\left(\text{BesselJ}(0, 2\sqrt{x}) - \frac{\text{BesselJ}(1, 2\sqrt{x})}{2\sqrt{x}}\right) - \frac{c_2 \text{BesselY}(1, 2\sqrt{x})}{2\sqrt{x}} - c_2\left(\text{BesselY}(0, 2\sqrt{x}) - \frac{\text{BesselY}(1, 2\sqrt{x})}{2\sqrt{x}}\right)\right)}{-c_3\sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{BesselY}(1, 2\sqrt{x})} &= \xi(x) y' - y \xi'(x) \\ &= y' + y \xi(x)\end{aligned}$$

Which is now a first order ode. This is now solved for y . In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{(-1 + y) (c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x}))}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} \end{aligned}$$

Where $f(x) = \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))}$ and $g(y) = -1 + y$. Integrating both sides gives

$$\begin{aligned} \frac{1}{-1 + y} dy &= \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} dx \\ \int \frac{1}{-1 + y} dy &= \int \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} dx \\ \ln(-1 + y) &= \int \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} dx + c_3 \end{aligned}$$

Raising both side to exponential gives

$$-1 + y = e^{\int \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} dx + c_3}$$

Which simplifies to

$$-1 + y = c_4 e^{\int \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} dx}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_4 e^{\int \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} dx + c_3} + 1$$

Summary

The solution(s) found are the following

$$y = c_4 e^{\int \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} dx + c_3} + 1 \quad (1)$$

Verification of solutions

$$y = c_4 e^{\int \frac{c_2 \text{BesselY}(0, 2\sqrt{x}) + c_3 \text{BesselJ}(0, 2\sqrt{x})}{\sqrt{x} (c_2 \text{BesselY}(1, 2\sqrt{x}) + c_3 \text{BesselJ}(1, 2\sqrt{x}))} dx + c_3} + 1$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    <- Kummer successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 167

```
dsolve(x/(1-x)*diff(y(x),x$2)+y(x)=1/(1-x),y(x), singsol=all)
```

$$y(x) = -x \left(\text{BesselK}(0, -x) - \text{BesselK}(1, -x) \left(\int \frac{-\text{BesselI}(0, -x) - \text{BesselI}(1, -x)}{x(\text{BesselI}(0, x)(x+1)\text{BesselK}(1, -x) + 1 - (x+1)\text{BesselK}(0, -x)\text{BesselI}(1, x) + (-\text{BesselI}(0, -x)))} dx \right) - \text{BesselI}(1, -x) \left(\int \frac{-\text{BesselK}(0, -x) + \text{BesselK}(1, -x)}{(\text{BesselI}(0, x)(x+1)\text{BesselK}(1, -x) + 1 - (x+1)\text{BesselK}(0, -x)\text{BesselI}(1, x))} dx \right) - \text{BesselK}(0, -x) c_1 + \text{BesselK}(1, -x) c_1 - \text{BesselI}(0, -x) c_2 - \text{BesselI}(1, -x) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.266 (sec). Leaf size: 136

```
DSolve[x/(1-x)*y'[x]+y[x]==1/(1-x),y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} y(x) \rightarrow e^{-x} x & \left(e^x \text{BesselI}(0, x) \right. \\ & - \text{BesselI}(1, x) \int_1^x 2e^{-K[1]} \sqrt{\pi} \text{HypergeometricU} \left(\frac{1}{2}, 2, 2K[1] \right) dK[1] \\ & - 2\sqrt{\pi} x \text{HypergeometricU} \left(\frac{1}{2}, 2, 2x \right) {}_1F_2 \left(\frac{1}{2}; 1, \frac{3}{2}; \frac{x^2}{4} \right) \\ & + 2\sqrt{\pi} \text{HypergeometricU} \left(\frac{1}{2}, 2, 2x \right) \text{BesselI}(0, x) \\ & \left. + c_1 \text{HypergeometricU} \left(\frac{1}{2}, 2, 2x \right) + c_2 e^x \text{BesselI}(0, x) - c_2 e^x \text{BesselI}(1, x) \right) \end{aligned}$$

4.65 problem 62

4.65.1 Solving as second order bessel ode ode 2277

4.65.2 Maple step by step solution 2278

Internal problem ID [7286]

Internal file name [OUTPUT/6272_Sunday_June_05_2022_04_36_42_PM_97420023/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 62.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_bessel_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$\frac{xy''}{1-x} + yx = 0$$

4.65.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' + (-x^3 + x^2)y = 0 \tag{1}$$

Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= 2 \\ n &= -1 \\ \gamma &= \frac{1}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = -c_1\sqrt{x} \text{ BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{ BesselY}(1, 2\sqrt{x})$$

Summary

The solution(s) found are the following

$$y = -c_1\sqrt{x} \text{ BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{ BesselY}(1, 2\sqrt{x}) \quad (1)$$

Verification of solutions

$$y = -c_1\sqrt{x} \text{ BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{ BesselY}(1, 2\sqrt{x})$$

Verified OK.

4.65.2 Maple step by step solution

Let's solve

$$y'' + (1 - x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=\max(0, -m)}^{\infty} a_k x^{k+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=\max(0,-m)+m}^{\infty} a_{k-m} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + a_0 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) + a_k - a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 + a_0 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} + a_k - a_{k-1} = 0$$

- Shift index using $k- > k+1$

$$((k+1)^2 + 3k + 5) a_{k+3} + a_{k+1} - a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = \frac{-a_{k+1} + a_k}{k^2 + 5k + 6}, 2a_2 + a_0 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x/(1-x)*diff(y(x),x$2)+x*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \text{AiryAi}(x - 1) + c_2 \text{AiryBi}(x - 1)$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 20

```
DSolve[x/(1-x)*y'[x]+x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \text{AiryAi}(x - 1) + c_2 \text{AiryBi}(x - 1)$$

4.66 problem 63

4.66.1 Solving as second order bessel ode ode 2281

Internal problem ID [7287]

Internal file name [OUTPUT/6273_Sunday_June_05_2022_04_36_43_PM_54839824/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 63.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

[[_2nd_order , _linear , _nonhomogeneous]]

$$\frac{xy''}{1-x} + y = \cos(x)$$

Multiplying the ode throughout by the denominator of the coefficient of y'' results in

$$-xy'' + (x-1)y = \cos(x)(x-1)$$

4.66.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' + (-x^2 + x)y = x \cos(x)(1-x) \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= \frac{1}{2} \\ \beta &= 2 \\ n &= -1 \\ \gamma &= \frac{1}{2} \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = -c_1 \sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) - c_2 \sqrt{x} \text{BesselY}(1, 2\sqrt{x})$$

Therefore the homogeneous solution y_h is

$$y_h = -c_1 \sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) - c_2 \sqrt{x} \text{BesselY}(1, 2\sqrt{x})$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= -\sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) \\ y_2 &= -\sqrt{x} \text{BesselY}(1, 2\sqrt{x}) \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} -\sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) & -\sqrt{x} \text{BesselY}(1, 2\sqrt{x}) \\ \frac{d}{dx}(-\sqrt{x} \text{BesselJ}(1, 2\sqrt{x})) & \frac{d}{dx}(-\sqrt{x} \text{BesselY}(1, 2\sqrt{x})) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} -\sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) & -\sqrt{x} \text{BesselY}(1, 2\sqrt{x}) \\ -\text{BesselJ}(0, 2\sqrt{x}) & -\text{BesselY}(0, 2\sqrt{x}) \end{vmatrix}$$

Therefore

$$W = (-\sqrt{x} \text{BesselJ}(1, 2\sqrt{x})) (-\text{BesselY}(0, 2\sqrt{x})) - (-\sqrt{x} \text{BesselY}(1, 2\sqrt{x})) (-\text{BesselJ}(0, 2\sqrt{x}))$$

Which simplifies to

$$W = \sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) \text{BesselY}(0, 2\sqrt{x}) - \sqrt{x} \text{BesselY}(1, 2\sqrt{x}) \text{BesselJ}(0, 2\sqrt{x})$$

Which simplifies to

$$W = \frac{1}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-x^{\frac{3}{2}} \text{BesselY}(1, 2\sqrt{x}) \cos(x) (1-x)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\text{BesselY}(1, 2\sqrt{x}) \cos(x) (x-1) \pi}{\sqrt{x}} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{\text{BesselY}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha-1) \pi}{\sqrt{\alpha}} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{-x^{\frac{3}{2}} \text{BesselJ}(1, 2\sqrt{x}) \cos(x) (1-x)}{\frac{x^2}{\pi}} dx$$

Which simplifies to

$$u_2 = \int \frac{\text{BesselJ}(1, 2\sqrt{x}) \cos(x) (x-1) \pi}{\sqrt{x}} dx$$

Hence

$$u_2 = \int_0^x \frac{\text{BesselJ}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha-1) \pi}{\sqrt{\alpha}} d\alpha$$

Which simplifies to

$$u_1 = -\pi \left(\int_0^x \frac{\text{BesselY}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha-1)}{\sqrt{\alpha}} d\alpha \right)$$

$$u_2 = \pi \left(\int_0^x \frac{\text{BesselJ}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha-1)}{\sqrt{\alpha}} d\alpha \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \pi \left(\int_0^x \frac{\text{BesselY}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha-1)}{\sqrt{\alpha}} d\alpha \right) \sqrt{x} \text{BesselJ}(1, 2\sqrt{x})$$

$$- \pi \left(\int_0^x \frac{\text{BesselJ}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha-1)}{\sqrt{\alpha}} d\alpha \right) \sqrt{x} \text{BesselY}(1, 2\sqrt{x})$$

Which simplifies to

$$y_p(x) = \pi \sqrt{x} \left(\left(\int_0^x \frac{\text{BesselY}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha-1)}{\sqrt{\alpha}} d\alpha \right) \text{BesselJ}(1, 2\sqrt{x}) \right.$$

$$\left. - \left(\int_0^x \frac{\text{BesselJ}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha-1)}{\sqrt{\alpha}} d\alpha \right) \text{BesselY}(1, 2\sqrt{x}) \right)$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= (-c_1\sqrt{x} \operatorname{BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \operatorname{BesselY}(1, 2\sqrt{x})) \\
 &\quad + \left(\pi\sqrt{x} \left(\left(\int_0^x \frac{\operatorname{BesselY}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha - 1)}{\sqrt{\alpha}} d\alpha \right) \operatorname{BesselJ}(1, 2\sqrt{x}) \right. \right. \\
 &\quad \left. \left. - \left(\int_0^x \frac{\operatorname{BesselJ}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha - 1)}{\sqrt{\alpha}} d\alpha \right) \operatorname{BesselY}(1, 2\sqrt{x}) \right) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= -c_1\sqrt{x} \operatorname{BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \operatorname{BesselY}(1, 2\sqrt{x}) \\
 &\quad + \pi\sqrt{x} \left(\left(\int_0^x \frac{\operatorname{BesselY}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha - 1)}{\sqrt{\alpha}} d\alpha \right) \operatorname{BesselJ}(1, 2\sqrt{x}) \right. \\
 &\quad \left. - \left(\int_0^x \frac{\operatorname{BesselJ}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha - 1)}{\sqrt{\alpha}} d\alpha \right) \operatorname{BesselY}(1, 2\sqrt{x}) \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= -c_1\sqrt{x} \operatorname{BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \operatorname{BesselY}(1, 2\sqrt{x}) \\
 &\quad + \pi\sqrt{x} \left(\left(\int_0^x \frac{\operatorname{BesselY}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha - 1)}{\sqrt{\alpha}} d\alpha \right) \operatorname{BesselJ}(1, 2\sqrt{x}) \right. \\
 &\quad \left. - \left(\int_0^x \frac{\operatorname{BesselJ}(1, 2\sqrt{\alpha}) \cos(\alpha) (\alpha - 1)}{\sqrt{\alpha}} d\alpha \right) \operatorname{BesselY}(1, 2\sqrt{x}) \right)
 \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    <- Kummer successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 169

```
dsolve(x/(1-x)*diff(y(x),x$2)+y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = - \left((\text{BesselI}(0, -x) + \text{BesselI}(1, -x)) \left(\int \frac{\cos(x) (\text{BesselK}(0, -x) - \text{BesselK}(1, -x)) (x - 1)}{x (\text{BesselI}(0, x) (x + 1) \text{BesselK}(1, -x) + 1 - (x + 1) \text{BesselK}(0, -x) \text{BesselI}(1, x))} dx \right) \right. \\ \left. + (-\text{BesselK}(0, -x) + \text{BesselK}(1, -x)) \left(\int \frac{\cos(x) (\text{BesselI}(0, x) - \text{BesselI}(1, x)) (x - 1)}{x (\text{BesselI}(0, x) (x + 1) \text{BesselK}(1, -x) + 1 - (x + 1) \text{BesselK}(0, -x) \text{BesselI}(1, x))} dx \right) \right) \\ + \text{BesselK}(1, -x) c_1 - \text{BesselK}(0, -x) c_1 - \text{BesselI}(0, -x) c_2 - \text{BesselI}(1, -x) c_2 \Big) x$$

✓ Solution by Mathematica

Time used: 8.805 (sec). Leaf size: 133

```
DSolve[x/(1-x)*y'[x]+y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} y(x) \rightarrow & e^{-x} x \left(\text{HypergeometricU} \left(\frac{1}{2}, 2, 2x \right) \int_1^x 2\sqrt{\pi} (\text{BesselI}(0, K[1]) \right. \\ & \left. - \text{BesselI}(1, K[1])) \cos(K[1]) (K[1] - 1) dK[1] \right. \\ & \left. + e^x (\text{BesselI}(0, x) - \text{BesselI}(1, x)) \int_1^x \right. \\ & \left. - 2e^{-K[2]} \sqrt{\pi} \cos(K[2]) \text{HypergeometricU} \left(\frac{1}{2}, 2, 2K[2] \right) (K[2] - 1) dK[2] \right. \\ & \left. + c_1 \text{HypergeometricU} \left(\frac{1}{2}, 2, 2x \right) + c_2 e^x \text{BesselI}(0, x) - c_2 e^x \text{BesselI}(1, x) \right) \end{aligned}$$

4.67 problem 64

4.67.1 Solving as second order bessel ode ode 2288

Internal problem ID [7288]

Internal file name [OUTPUT/6274_Sunday_June_05_2022_04_36_46_PM_19328234/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 64.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

[[_2nd_order , _with_linear_symmetries]]

$$\frac{xy''}{1-x^2} + y = 0$$

4.67.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' + (-x^3 + x)y = 0 \tag{1}$$

Bessel ode has the form

$$x^2y'' + xy' + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\alpha = \frac{1}{2}$$

$$\beta = 2$$

$$n = -1$$

$$\gamma = \frac{1}{2}$$

Substituting all the above into (4) gives the solution as

$$y = -c_1\sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{BesselY}(1, 2\sqrt{x})$$

Summary

The solution(s) found are the following

$$y = -c_1\sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{BesselY}(1, 2\sqrt{x}) \quad (1)$$

Verification of solutions

$$y = -c_1\sqrt{x} \text{BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{BesselY}(1, 2\sqrt{x})$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Whittaker
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * {}_2F_1([a$ 
  trying differential order: 2; exact nonlinear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying to convert to an ODE of Bessel type
  -> trying reduction of order to Riccati
    trying Riccati sub-methods:
      -> trying a symmetry pattern of the form  $[F(x)*G(y), 0]$ 
      -> trying a symmetry pattern of the form  $[0, F(x)*G(y)]$ 
      -> trying a symmetry pattern of the form  $[F(x), G(x)*y+H(x)]$ 
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 3` [0, y]
```

X Solution by Maple

```
dsolve(x/(1-x^2)*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[x/(1-x^2)*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.68 problem 65

- 4.68.1 Solving as second order bessel ode ode 2292
- 4.68.2 Solving using Kovacic algorithm 2293
- 4.68.3 Maple step by step solution 2299

Internal problem ID [7289]

Internal file name [OUTPUT/6275_Sunday_June_05_2022_04_36_48_PM_61369062/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 65.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_bessel_ode"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - (x^2 + 3)y = 0$$

4.68.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' + (-x^4 - 3x^2)y = 0 \tag{1}$$

Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha)xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= \frac{1}{2} \\ \beta &= 2 \\ n &= -1 \\ \gamma &= \frac{1}{2}\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = -c_1\sqrt{x} \text{ BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{ BesselY}(1, 2\sqrt{x})$$

Summary

The solution(s) found are the following

$$y = -c_1\sqrt{x} \text{ BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{ BesselY}(1, 2\sqrt{x}) \quad (1)$$

Verification of solutions

$$y = -c_1\sqrt{x} \text{ BesselJ}(1, 2\sqrt{x}) - c_2\sqrt{x} \text{ BesselY}(1, 2\sqrt{x})$$

Verified OK.

4.68.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + (-x^2 - 3)y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\ B &= 0 \\ C &= -x^2 - 3\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 3}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = x^2 + 3$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = (x^2 + 3) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 238: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 2 \\ &= -2 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx x + \frac{3}{2x} - \frac{9}{8x^3} + \frac{27}{16x^5} - \frac{405}{128x^7} + \frac{1701}{256x^9} - \frac{15309}{1024x^{11}} + \frac{72171}{2048x^{13}} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = 1$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= x \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = x^2$$

This shows that the coefficient of 1 in the above is 0. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 3}{1} \\ &= Q + \frac{R}{1} \\ &= (x^2 + 3) + (0) \\ &= x^2 + 3 \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is 3. Now b can be found.

$$\begin{aligned} b &= (3) - (0) \\ &= 3 \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_{\infty} &= x \\ \alpha_{\infty}^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{3}{1} - 1 \right) = 1 \\ \alpha_{\infty}^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{3}{1} - 1 \right) = -2 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = x^2 + 3$$

Order of r at ∞	$[\sqrt{r}]_{\infty}$	α_{∞}^+	α_{∞}^-
-2	x	1	-2

Now that the all $[\sqrt{r}]_c$ and its associated α_c^{\pm} have been determined for all the poles in the set Γ and $[\sqrt{r}]_{\infty}$ and its associated α_{∞}^{\pm} have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^+ = 1$, and since there are no poles, then

$$\begin{aligned} d &= \alpha_\infty^+ \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

Substituting the above values in the above results in

$$\begin{aligned} \omega &= (+) [\sqrt{r}]_\infty \\ &= 0 + (x) \\ &= x \\ &= x \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \tag{1A}$$

Let

$$p(x) = x + a_0 \tag{2A}$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2(x)(1) + ((1) + (x)^2 - (x^2 + 3)) &= 0 \\ -2a_0 &= 0 \end{aligned}$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 0\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= (x) e^{\int x dx} \\ &= (x) e^{\frac{x^2}{2}} \\ &= x e^{\frac{x^2}{2}} \end{aligned}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= x e^{\frac{x^2}{2}} \end{aligned}$$

Which simplifies to

$$y_1 = x e^{\frac{x^2}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= x e^{\frac{x^2}{2}} \int \frac{1}{x^2 e^{x^2}} dx \\ &= x e^{\frac{x^2}{2}} \left(\frac{-\sqrt{\pi} \operatorname{erf}(x) x - e^{-x^2}}{x} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(x e^{\frac{x^2}{2}} \right) + c_2 \left(x e^{\frac{x^2}{2}} \left(\frac{-\sqrt{\pi} \operatorname{erf}(x) x - e^{-x^2}}{x} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x e^{\frac{x^2}{2}} + c_2 \left(-\sqrt{\pi} \operatorname{erf}(x) x e^{\frac{x^2}{2}} - e^{-\frac{x^2}{2}} \right) \quad (1)$$

Verification of solutions

$$y = c_1 x e^{\frac{x^2}{2}} + c_2 \left(-\sqrt{\pi} \operatorname{erf}(x) x e^{\frac{x^2}{2}} - e^{-\frac{x^2}{2}} \right)$$

Verified OK.

4.68.3 Maple step by step solution

Let's solve

$$y'' + (-x^2 - 3)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=\max(0,-m)}^{\infty} a_k x^{k+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=\max(0,-m)+m}^{\infty} a_{k-m} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 - 3a_0 + (6a_3 - 3a_1)x + \left(\sum_{k=2}^{\infty} (a_{k+2}(k+2)(k+1) - 3a_k - a_{k-2}) x^k \right) = 0$$

- The coefficients of each power of x must be 0
 $[2a_2 - 3a_0 = 0, 6a_3 - 3a_1 = 0]$
- Solve for the dependent coefficient(s)
 $\{a_2 = \frac{3a_0}{2}, a_3 = \frac{a_1}{2}\}$
- Each term in the series must be 0, giving the recursion relation
 $(k^2 + 3k + 2) a_{k+2} - 3a_k - a_{k-2} = 0$
- Shift index using $k \rightarrow k + 2$
 $((k + 2)^2 + 3k + 8) a_{k+4} - 3a_{k+2} - a_k = 0$
- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = \frac{3a_{k+2} + a_k}{k^2 + 7k + 12}, a_2 = \frac{3a_0}{2}, a_3 = \frac{a_1}{2} \right]$$

Maple trace **Kovacic algorithm successful**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 30

```
dsolve(diff(y(x), x$2)=(x^2+3)*y(x), y(x), singsol=all)
```

$$y(x) = x(c_2\sqrt{\pi} \operatorname{erf}(x) + c_1) e^{\frac{x^2}{2}} + e^{-\frac{x^2}{2}} c_2$$

✓ Solution by Mathematica

Time used: 0.125 (sec). Leaf size: 46

```
DSolve[y''[x]==(x^2+3)*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{x^2}{2}} \left(-\sqrt{\pi} c_2 e^{x^2} \operatorname{erf}(x) + c_1 e^{x^2} x - c_2 \right)$$

4.69 problem 66

4.69.1 Maple step by step solution 2309

Internal problem ID [7290]

Internal file name [OUTPUT/6276_Sunday_June_05_2022_04_36_51_PM_22286391/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 66.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + (x - 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (360)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (361)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -(x - 1)y \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= -y - (x - 1)y' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= -2y' + (x - 1)^2y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= (x - 1)((x - 1)y' + 4y) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= -(x - 1)^3y + (6x - 6)y' + 4y
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= y(0) \\
 F_1 &= -y(0) + y'(0) \\
 F_2 &= -2y'(0) + y(0) \\
 F_3 &= y'(0) - 4y(0) \\
 F_4 &= 5y(0) - 6y'(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned}
 y &= \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + \frac{1}{144}x^6\right)y(0) \\
 &+ \left(x + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5 - \frac{1}{120}x^6\right)y'(0) + O(x^6)
 \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -(x-1) \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} x^{1+n} a_n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=0}^{\infty} x^{1+n} a_n = \sum_{n=1}^{\infty} a_{n-1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 - a_0 = 0$$

$$a_2 = \frac{a_0}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+2)a_{n+2}(1+n) + a_{n-1} - a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned} a_{n+2} &= \frac{-a_{n-1} + a_n}{(n+2)(1+n)} \\ (5) \qquad &= \frac{a_n}{(n+2)(1+n)} - \frac{a_{n-1}}{(n+2)(1+n)} \end{aligned}$$

For $n = 1$ the recurrence equation gives

$$6a_3 + a_0 - a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{6} + \frac{a_1}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + a_1 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_1}{12} + \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + a_2 - a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_0}{30} + \frac{a_1}{120}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + a_3 - a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{144} - \frac{a_1}{120}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + a_4 - a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{11a_1}{5040} - \frac{a_0}{560}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + \frac{a_0 x^2}{2} + \left(-\frac{a_0}{6} + \frac{a_1}{6}\right) x^3 + \left(-\frac{a_1}{12} + \frac{a_0}{24}\right) x^4 + \left(-\frac{a_0}{30} + \frac{a_1}{120}\right) x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5\right) a_0 + \left(x + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5\right) c_1 + \left(x + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + \frac{1}{144}x^6\right) y(0) \\ &+ \left(x + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5 - \frac{1}{120}x^6\right) y'(0) + O(x^6) \\ y &= \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5\right) c_1 + \left(x + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5\right) c_2 + O(x^6)\end{aligned}\tag{1}$$

Verification of solutions

$$\begin{aligned}y &= \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + \frac{1}{144}x^6\right) y(0) \\ &+ \left(x + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5 - \frac{1}{120}x^6\right) y'(0) + O(x^6)\end{aligned}$$

Verified OK.

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5\right) c_1 + \left(x + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5\right) c_2 + O(x^6)$$

Verified OK.

4.69.1 Maple step by step solution

Let's solve

$$y'' = -(x - 1)y$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = (1 - x)y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + (x - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=\max(0,-m)}^{\infty} a_k x^{k+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=\max(0,-m)+m}^{\infty} a_{k-m} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 - a_0 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) - a_k + a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 - a_0 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - a_k + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} - a_{k+1} + a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{-a_{k+1} + a_k}{k^2 + 5k + 6}, 2a_2 - a_0 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;  
dsolve(diff(y(x),x$2)+(x-1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5\right) y(0) \\ + \left(x + \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[y''[x]+(x-1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} - \frac{x^4}{12} + \frac{x^3}{6} + x \right) + c_1 \left(-\frac{x^5}{30} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} + 1 \right)$$

4.70 problem 67

- 4.70.1 Solution using Matrix exponential method 2312
- 4.70.2 Solution using explicit Eigenvalue and Eigenvector method . . . 2314
- 4.70.3 Maple step by step solution 2320

Internal problem ID [7291]

Internal file name [OUTPUT/6277_Sunday_June_05_2022_04_36_52_PM_56014866/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 67.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "system of linear ODEs"

Solve

$$\begin{aligned}x'(t) &= x(t) + 2y(t) + 2t + 1 \\y'(t) &= 5x(t) + y(t) + 3t - 1\end{aligned}$$

4.70.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 5 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} 2t + 1 \\ 3t - 1 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation

of parameters method applied to the fundamental matrix. For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} & -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{10} \\ -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{4} & \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned} \vec{x}_h(t) &= e^{At}\vec{c} \\ &= \begin{bmatrix} \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} & -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{10} \\ -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{4} & \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2}\right)c_1 - \frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}c_2}{10} \\ -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}c_1}{4} + \left(\frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2}\right)c_2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{(-c_2\sqrt{10}+5c_1)e^{-(-1+\sqrt{10})t}}{10} + \frac{\left(\frac{c_2\sqrt{10}}{5}+c_1\right)e^{(1+\sqrt{10})t}}{2} \\ \frac{(-c_1\sqrt{10}+2c_2)e^{-(-1+\sqrt{10})t}}{4} + \frac{e^{(1+\sqrt{10})t}(c_1\sqrt{10}+2c_2)}{4} \end{bmatrix} \end{aligned}$$

The particular solution given by

$$\vec{x}_p(t) = e^{At} \int e^{-At}\vec{G}(t) dt$$

But

$$\begin{aligned} e^{-At} &= (e^{At})^{-1} \\ &= \begin{bmatrix} \frac{e^{-2t}\left(e^{-(-1+\sqrt{10})t} + e^{(1+\sqrt{10})t}\right)}{2} & \frac{\sqrt{10}e^{-2t}\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)}{10} \\ \frac{\sqrt{10}e^{-2t}\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)}{4} & \frac{e^{-2t}\left(e^{-(-1+\sqrt{10})t} + e^{(1+\sqrt{10})t}\right)}{2} \end{bmatrix} \end{aligned}$$

Hence

$$\begin{aligned}
\vec{x}_p(t) &= \begin{bmatrix} \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} & -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{10} \\ -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{4} & \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} \end{bmatrix} \int \begin{bmatrix} \frac{e^{-2t}\left(e^{-(-1+\sqrt{10})t} + e^{(1+\sqrt{10})t}\right)}{2} \\ \frac{\sqrt{10}e^{-2t}\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)}{4} \end{bmatrix} \\
&= \begin{bmatrix} \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} & -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{10} \\ -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{4} & \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} \end{bmatrix} \begin{bmatrix} \frac{\left((-63t-67)\sqrt{10}-180t+85\right)e^{-(1+\sqrt{10})t}}{810} \\ \frac{\left((-36t+17)\sqrt{10}-126t-134\right)e^{-(1+\sqrt{10})t}}{324} \end{bmatrix} \\
&= \begin{bmatrix} -\frac{4t}{9} + \frac{17}{81} \\ -\frac{7t}{9} - \frac{67}{81} \end{bmatrix}
\end{aligned}$$

Hence the complete solution is

$$\begin{aligned}
\vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\
&= \begin{bmatrix} \frac{17}{81} + \frac{\left(-c_2\sqrt{10}+5c_1\right)e^{-(-1+\sqrt{10})t}}{10} + \frac{e^{(1+\sqrt{10})t}\left(c_2\sqrt{10}+5c_1\right)}{10} - \frac{4t}{9} \\ \frac{\left(-c_1\sqrt{10}+2c_2\right)e^{-(-1+\sqrt{10})t}}{4} + \frac{e^{(1+\sqrt{10})t}\left(c_1\sqrt{10}+2c_2\right)}{4} - \frac{7t}{9} - \frac{67}{81} \end{bmatrix}
\end{aligned}$$

4.70.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 5 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} 2t + 1 \\ 3t - 1 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix.

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 2 \\ 5 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 2 \\ 5 & 1 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda - 9 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1 + \sqrt{10}$$

$$\lambda_2 = 1 - \sqrt{10}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$1 - \sqrt{10}$	1	real eigenvalue
$1 + \sqrt{10}$	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1 - \sqrt{10}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ 5 & 1 \end{bmatrix} - (1 - \sqrt{10}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} \sqrt{10} & 2 \\ 5 & \sqrt{10} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} \sqrt{10} & 2 & 0 \\ 5 & \sqrt{10} & 0 \end{array} \right]$$

$$R_2 = R_2 - \frac{\sqrt{10} R_1}{2} \implies \left[\begin{array}{cc|c} \sqrt{10} & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{cc} \sqrt{10} & 2 \\ 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = -\frac{t\sqrt{10}}{5} \right\}$

Hence the solution is

$$\begin{bmatrix} -\frac{t\sqrt{10}}{5} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{t\sqrt{10}}{5} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{t\sqrt{10}}{5} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix} = \begin{bmatrix} -\sqrt{10} \\ 5 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 1 + \sqrt{10}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ 5 & 1 \end{bmatrix} - (1 + \sqrt{10}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -\sqrt{10} & 2 \\ 5 & -\sqrt{10} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -\sqrt{10} & 2 & 0 \\ 5 & -\sqrt{10} & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{\sqrt{10} R_1}{2} \implies \left[\begin{array}{cc|c} -\sqrt{10} & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -\sqrt{10} & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = \frac{t\sqrt{10}}{5} \right\}$

Hence the solution is

$$\begin{bmatrix} \frac{t\sqrt{10}}{5} \\ t \end{bmatrix} = \begin{bmatrix} \frac{t\sqrt{10}}{5} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{t\sqrt{10}}{5} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{\sqrt{10}}{5} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{t\sqrt{10}}{5} \\ t \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{10}}{5} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{t\sqrt{10}}{5} \\ t \end{bmatrix} = \begin{bmatrix} \sqrt{10} \\ 5 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$1 + \sqrt{10}$	1	1	No	$\begin{bmatrix} \frac{\sqrt{10}}{5} \\ 1 \end{bmatrix}$
$1 - \sqrt{10}$	1	1	No	$\begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue $1 + \sqrt{10}$ is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_1(t) &= \vec{v}_1 e^{(1+\sqrt{10})t} \\ &= \begin{bmatrix} \frac{\sqrt{10}}{5} \\ 1 \end{bmatrix} e^{(1+\sqrt{10})t} \end{aligned}$$

Since eigenvalue $1 - \sqrt{10}$ is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_2(t) &= \vec{v}_2 e^{(1-\sqrt{10})t} \\ &= \begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix} e^{(1-\sqrt{10})t} \end{aligned}$$

Therefore the homogeneous solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \frac{\sqrt{10}e^{(1+\sqrt{10})t}}{5} \\ e^{(1+\sqrt{10})t} \end{bmatrix} + c_2 \begin{bmatrix} -\frac{e^{(1-\sqrt{10})t}\sqrt{10}}{5} \\ e^{(1-\sqrt{10})t} \end{bmatrix}$$

Now that we found homogeneous solution above, we need to find a particular solution $\vec{x}_p(t)$. We will use Variation of parameters. The fundamental matrix is

$$\Phi = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 & \cdots \end{bmatrix}$$

Where \vec{x}_i are the solution basis found above. Therefore the fundamental matrix is

$$\Phi(t) = \begin{bmatrix} \frac{\sqrt{10}e^{(1+\sqrt{10})t}}{5} & -\frac{e^{(1-\sqrt{10})t}\sqrt{10}}{5} \\ e^{(1+\sqrt{10})t} & e^{(1-\sqrt{10})t} \end{bmatrix}$$

The particular solution is then given by

$$\vec{x}_p(t) = \Phi \int \Phi^{-1} \vec{G}(t) dt$$

But

$$\Phi^{-1} = \begin{bmatrix} \frac{\sqrt{10}e^{-(1+\sqrt{10})t}}{4} & \frac{e^{-(1+\sqrt{10})t}}{2} \\ -\frac{\sqrt{10}e^{-(1-\sqrt{10})t}}{4} & \frac{e^{-(1-\sqrt{10})t}}{2} \end{bmatrix}$$

Hence

$$\begin{aligned} \vec{x}_p(t) &= \begin{bmatrix} \frac{\sqrt{10}e^{(1+\sqrt{10})t}}{5} & -\frac{e^{(1-\sqrt{10})t}\sqrt{10}}{5} \\ e^{(1+\sqrt{10})t} & e^{(1-\sqrt{10})t} \end{bmatrix} \int \begin{bmatrix} \frac{\sqrt{10}e^{-(1+\sqrt{10})t}}{4} & \frac{e^{-(1+\sqrt{10})t}}{2} \\ -\frac{\sqrt{10}e^{-(1-\sqrt{10})t}}{4} & \frac{e^{-(1-\sqrt{10})t}}{2} \end{bmatrix} \begin{bmatrix} 2t+1 \\ 3t-1 \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{\sqrt{10}e^{(1+\sqrt{10})t}}{5} & -\frac{e^{(1-\sqrt{10})t}\sqrt{10}}{5} \\ e^{(1+\sqrt{10})t} & e^{(1-\sqrt{10})t} \end{bmatrix} \int \begin{bmatrix} \frac{e^{-(1+\sqrt{10})t}(2t\sqrt{10}+\sqrt{10}+6t-2)}{4} \\ -\frac{e^{-(1-\sqrt{10})t}(2t\sqrt{10}+\sqrt{10}-6t+2)}{4} \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{\sqrt{10}e^{(1+\sqrt{10})t}}{5} & -\frac{e^{(1-\sqrt{10})t}\sqrt{10}}{5} \\ e^{(1+\sqrt{10})t} & e^{(1-\sqrt{10})t} \end{bmatrix} \begin{bmatrix} \frac{(18t\sqrt{10}+185\sqrt{10}-18t-572)e^{-(1+\sqrt{10})t}(2t\sqrt{10}+\sqrt{10}+6t-2)}{-648t-5184+1620\sqrt{10}} \\ -\frac{(18t\sqrt{10}+185\sqrt{10}+18t+572)e^{-(1-\sqrt{10})t}(2t\sqrt{10}+\sqrt{10}-6t+2)}{324(2t+16+5\sqrt{10})} \end{bmatrix} \\ &= \begin{bmatrix} \frac{-72t^3-1118t^2+436t+51}{162t^2+2592t+243} \\ \frac{-126t^3-2150t^2-2333t-201}{162t^2+2592t+243} \end{bmatrix} \end{aligned}$$

Now that we found particular solution, the final solution is

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{c_1\sqrt{10}e^{(1+\sqrt{10})t}}{5} \\ c_1e^{(1+\sqrt{10})t} \end{bmatrix} + \begin{bmatrix} -\frac{c_2e^{(1-\sqrt{10})t}\sqrt{10}}{5} \\ c_2e^{(1-\sqrt{10})t} \end{bmatrix} + \begin{bmatrix} \frac{-72t^3-1118t^2+436t+51}{162t^2+2592t+243} \\ \frac{-126t^3-2150t^2-2333t-201}{162t^2+2592t+243} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{c_1\sqrt{10}e^{(1+\sqrt{10})t}}{5} - \frac{c_2e^{-(1+\sqrt{10})t}\sqrt{10}}{5} - \frac{4t}{9} + \frac{17}{81} \\ c_1e^{(1+\sqrt{10})t} + c_2e^{-(1+\sqrt{10})t} - \frac{7t}{9} - \frac{67}{81} \end{bmatrix}$$

4.70.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + 2y(t) + 2t + 1, y'(t) = 5x(t) + y(t) + 3t - 1]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ 5 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 2t + 1 \\ 3t - 1 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ 5 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 2t + 1 \\ 3t - 1 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(t) = \begin{bmatrix} 2t + 1 \\ 3t - 1 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 2 \\ 5 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[1 - \sqrt{10}, \begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix} \right], \left[1 + \sqrt{10}, \begin{bmatrix} \frac{\sqrt{10}}{5} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[1 - \sqrt{10}, \begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{(1-\sqrt{10})t} \cdot \begin{bmatrix} -\frac{\sqrt{10}}{5} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1 + \sqrt{10}, \begin{bmatrix} \frac{\sqrt{10}}{5} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{(1+\sqrt{10})t} \cdot \begin{bmatrix} \frac{\sqrt{10}}{5} \\ 1 \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{x}_p(t)$

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \vec{x}_p(t)$$

- Fundamental matrix

- Let $\phi(t)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(t) = \begin{bmatrix} -\frac{e^{(1-\sqrt{10})t}\sqrt{10}}{5} & \frac{\sqrt{10}e^{(1+\sqrt{10})t}}{5} \\ e^{(1-\sqrt{10})t} & e^{(1+\sqrt{10})t} \end{bmatrix}$$

- The fundamental matrix, $\Phi(t)$ is a normalized version of $\phi(t)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(t) = \phi(t) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(t)$ and $\phi(0)$

$$\Phi(t) = \begin{bmatrix} -\frac{e^{(1-\sqrt{10})t}\sqrt{10}}{5} & \frac{\sqrt{10}e^{(1+\sqrt{10})t}}{5} \\ e^{(1-\sqrt{10})t} & e^{(1+\sqrt{10})t} \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} -\frac{\sqrt{10}}{5} & \frac{\sqrt{10}}{5} \\ 1 & 1 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(t) = \begin{bmatrix} \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} & -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{10} \\ -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t}\right)\sqrt{10}}{4} & \frac{e^{-(-1+\sqrt{10})t}}{2} + \frac{e^{(1+\sqrt{10})t}}{2} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(t)$ and solve for $\vec{v}(t)$

$$\vec{x}_p(t) = \Phi(t) \cdot \vec{v}(t)$$

- Take the derivative of the particular solution

$$\vec{x}'_p(t) = \Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(t)$

$$A \cdot \Phi(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- Cancel like terms

$$\Phi(t) \cdot \vec{v}'(t) = \vec{f}(t)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(t) = \frac{1}{\Phi(t)} \cdot \vec{f}(t)$$

- Integrate to solve for $\vec{v}(t)$

$$\vec{v}(t) = \int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(t)$ into the equation for the particular solution

$$\vec{x}_p(t) = \Phi(t) \cdot \left(\int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{x}_p(t) = \begin{bmatrix} \frac{(-67\sqrt{10}-85)e^{-(1+\sqrt{10})t}}{810} + \frac{(67\sqrt{10}-85)e^{(1+\sqrt{10})t}}{810} - \frac{4t}{9} + \frac{17}{81} \\ \frac{(17\sqrt{10}+134)e^{-(1+\sqrt{10})t}}{324} + \frac{(-17\sqrt{10}+134)e^{(1+\sqrt{10})t}}{324} - \frac{7t}{9} - \frac{67}{81} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \begin{bmatrix} \frac{(-67\sqrt{10}-85)e^{-(1+\sqrt{10})t}}{810} + \frac{(67\sqrt{10}-85)e^{(1+\sqrt{10})t}}{810} - \frac{4t}{9} + \frac{17}{81} \\ \frac{(17\sqrt{10}+134)e^{-(1+\sqrt{10})t}}{324} + \frac{(-17\sqrt{10}+134)e^{(1+\sqrt{10})t}}{324} - \frac{7t}{9} - \frac{67}{81} \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{(-85+(-162c_1-67)\sqrt{10})e^{-(1+\sqrt{10})t}}{810} + \frac{(-85+(162c_2+67)\sqrt{10})e^{(1+\sqrt{10})t}}{810} - \frac{4t}{9} + \frac{17}{81} \\ \frac{(324c_1+17\sqrt{10}+134)e^{-(1+\sqrt{10})t}}{324} + \frac{(324c_2-17\sqrt{10}+134)e^{(1+\sqrt{10})t}}{324} - \frac{7t}{9} - \frac{67}{81} \end{bmatrix}$$

- Solution to the system of ODEs

$$\begin{cases} x(t) = \frac{(-85+(-162c_1-67)\sqrt{10})e^{-(1+\sqrt{10})t}}{810} + \frac{(-85+(162c_2+67)\sqrt{10})e^{(1+\sqrt{10})t}}{810} - \frac{4t}{9} + \frac{17}{81}, y(t) = \frac{(324c_1+17\sqrt{10}+134)e^{-(1+\sqrt{10})t}}{324} + \frac{(324c_2-17\sqrt{10}+134)e^{(1+\sqrt{10})t}}{324} - \frac{7t}{9} - \frac{67}{81} \end{cases}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 68

```
dsolve([diff(x(t),t)=x(t)+2*y(t)+2*t+1,diff(y(t),t)=5*x(t)+y(t)+3*t-1],singsol=all)
```

$$x(t) = e^{(1+\sqrt{10})t} c_2 + e^{-(1+\sqrt{10})t} c_1 - \frac{4t}{9} + \frac{17}{81}$$

$$y(t) = \frac{e^{(1+\sqrt{10})t} c_2 \sqrt{10}}{2} - \frac{e^{-(1+\sqrt{10})t} c_1 \sqrt{10}}{2} - \frac{7t}{9} - \frac{67}{81}$$

✓ Solution by Mathematica

Time used: 10.731 (sec). Leaf size: 158

```
DSolve[{x'[t]==x[t]+2*y[t]+2*t+1,y'[t]==5*x[t]+y[t]+3*t-1},{x[t],y[t]},t,IncludeSingularSolu
```

$$x(t) \rightarrow \frac{1}{810} e^{t-\sqrt{10}t} \left(e^{(\sqrt{10}-1)t} (170-360t) + 81(5c_1 + \sqrt{10}c_2) e^{2\sqrt{10}t} + 81(5c_1 - \sqrt{10}c_2) \right)$$

$$y(t) \rightarrow \frac{1}{324} e^{t-\sqrt{10}t} \left(-4e^{(\sqrt{10}-1)t} (63t+67) + 81(\sqrt{10}c_1 + 2c_2) e^{2\sqrt{10}t} - 81(\sqrt{10}c_1 - 2c_2) \right)$$

4.71 problem 68

- 4.71.1 Solving as second order linear constant coeff ode 2324
- 4.71.2 Solving using Kovacic algorithm 2327
- 4.71.3 Maple step by step solution 2332

Internal problem ID [7292]

Internal file name [OUTPUT/6278_Sunday_June_05_2022_04_36_56_PM_32947359/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 68.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 20y' + 500y = 100000 \cos(100x)$$

4.71.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 20, C = 500, f(x) = 100000 \cos(100x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 20y' + 500y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 20, C = 500$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 20\lambda e^{\lambda x} + 500 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 20\lambda + 500 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 20, C = 500$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-20}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{20^2 - (4)(1)(500)} \\ &= -10 \pm 20i \end{aligned}$$

Hence

$$\lambda_1 = -10 + 20i$$

$$\lambda_2 = -10 - 20i$$

Which simplifies to

$$\lambda_1 = -10 + 20i$$

$$\lambda_2 = -10 - 20i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -10$ and $\beta = 20$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-10x} (c_1 \cos(20x) + c_2 \sin(20x))$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-10x} (c_1 \cos(20x) + c_2 \sin(20x))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$100000 \cos(100x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(100x), \sin(100x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-10x} \cos(20x), e^{-10x} \sin(20x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(100x) + A_2 \sin(100x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} -9500A_1 \cos(100x) - 9500A_2 \sin(100x) - 2000A_1 \sin(100x) + 2000A_2 \cos(100x) \\ = 100000 \cos(100x) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{3800}{377}, A_2 = \frac{800}{377} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{-10x}(c_1 \cos(20x) + c_2 \sin(20x))) + \left(-\frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{-10x}(c_1 \cos(20x) + c_2 \sin(20x)) - \frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377} \quad (1)$$




Figure 135: Slope field plot

Verification of solutions

$$y = e^{-10x}(c_1 \cos(20x) + c_2 \sin(20x)) - \frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377}$$

Verified OK.

4.71.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 20y' + 500y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 20 \\C &= 500\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-400}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -400 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -400z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 242: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -400$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(20x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{20}{1} dx} \\
 &= z_1 e^{-10x} \\
 &= z_1 (e^{-10x})
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-10x} \cos(20x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{20}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-20x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\tan(20x)}{20} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-10x} \cos(20x)) + c_2 \left(e^{-10x} \cos(20x) \left(\frac{\tan(20x)}{20} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 20y' + 500y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(20x) e^{-10x} c_1 + \frac{\sin(20x) e^{-10x} c_2}{20}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$100000 \cos(100x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(100x), \sin(100x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-10x} \cos(20x), \frac{e^{-10x} \sin(20x)}{20} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(100x) + A_2 \sin(100x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} -9500A_1 \cos(100x) - 9500A_2 \sin(100x) - 2000A_1 \sin(100x) + 2000A_2 \cos(100x) \\ = 100000 \cos(100x) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{3800}{377}, A_2 = \frac{800}{377} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\cos(20x) e^{-10x} c_1 + \frac{\sin(20x) e^{-10x} c_2}{20} \right) + \left(-\frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(20x) e^{-10x} c_1 + \frac{\sin(20x) e^{-10x} c_2}{20} - \frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377} \quad (1)$$




Figure 136: Slope field plot

Verification of solutions

$$y = \cos(20x) e^{-10x} c_1 + \frac{\sin(20x) e^{-10x} c_2}{20} - \frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377}$$

Verified OK.

4.71.3 Maple step by step solution

Let's solve

$$y'' + 20y' + 500y = 100000 \cos(100x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 20r + 500 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-20) \pm (\sqrt{-1600})}{2}$$

- Roots of the characteristic polynomial

$$r = (-10 - 20I, -10 + 20I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-10x} \cos(20x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-10x} \sin(20x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(20x) e^{-10x} c_1 + \sin(20x) e^{-10x} c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 100000 \cos(100x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-10x} \cos(20x) & e^{-10x} \sin(20x) \\ -10 e^{-10x} \cos(20x) - 20 e^{-10x} \sin(20x) & -10 e^{-10x} \sin(20x) + 20 e^{-10x} \cos(20x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 20 e^{-20x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -5000 e^{-10x} (\cos(20x) \left(\int \sin(20x) \cos(100x) e^{10x} dx \right) - \sin(20x) \left(\int \cos(20x) \cos(100x) e^{10x} dx \right))$$

- Compute integrals

$$y_p(x) = -\frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377}$$

- Substitute particular solution into general solution to ODE

$$y = \cos(20x) e^{-10x} c_1 + \sin(20x) e^{-10x} c_2 - \frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 37

```
dsolve(diff(diff(y(x),x),x)+20*diff(y(x),x)+500*y(x) = 100000*cos(100*x),y(x), singsol=all)
```

$$y(x) = e^{-10x} \sin(20x) c_2 + e^{-10x} \cos(20x) c_1 - \frac{3800 \cos(100x)}{377} + \frac{800 \sin(100x)}{377}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 47

```
DSolve[y''[x]+20*y'[x]+500*y[x] == 100000*Cos[100*x],y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow -\frac{200}{377}(19 \cos(100x) - 4 \sin(100x)) + c_2 e^{-10x} \cos(20x) + c_1 e^{-10x} \sin(20x)$$

4.72 problem 69

4.72.1 Solving as second order change of variable on x method 2 ode . 2335

4.72.2 Solving as second order change of variable on x method 1 ode . 2338

Internal problem ID [7293]

Internal file name [OUTPUT/6279_Sunday_June_05_2022_04_36_59_PM_15964274/index.tex]

Book: Own collection of miscellaneous problems

Section: section 4.0

Problem number: 69.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2"

Maple gives the following as the ode type

```
[[_2nd_order, _with_linear_symmetries], [_2nd_order, _linear, `
  _with_symmetry_[0,F(x)]`]]
```

$$y'' \sin(2x)^2 + y' \sin(4x) - 4y = 0$$

4.72.1 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$y'' \sin(2x)^2 + y' \sin(4x) - 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{\sin(4x)}{\sin(2x)^2}$$
$$q(x) = -\frac{4}{\sin(2x)^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x) \tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-\left(\int p(x) dx\right)} dx \\ &= \int e^{-\left(\int \frac{\sin(4x)}{\sin(2x)^2} dx\right)} dx \\ &= \int e^{\frac{\ln(\csc(2x)^2)}{2}} dx \\ &= \int \operatorname{csgn}(\csc(2x)) \csc(2x) dx \\ &= -\frac{\operatorname{csgn}(\csc(2x)) \ln(\csc(2x) + \cot(2x))}{2} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{4}{\sin(2x)^2}}{\operatorname{csgn}(\csc(2x))^2 \csc(2x)^2} \\ &= -4 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2} y(\tau) + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) - 4y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 4 e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)} \\ &= \pm 2 \end{aligned}$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(2)\tau} + c_2 e^{(-2)\tau}$$

Or

$$y(\tau) = c_1 e^{2\tau} + c_2 e^{-2\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = c_1 (\csc(2x) + \cot(2x))^{-\text{signum}(\sin(2x))} + c_2 (\csc(2x) + \cot(2x))^{\text{signum}(\sin(2x))}$$

Summary

The solution(s) found are the following

$$y = c_1(\csc(2x) + \cot(2x))^{-\text{signum}(\sin(2x))} + c_2(\csc(2x) + \cot(2x))^{\text{signum}(\sin(2x))} \quad (1)$$

Verification of solutions

$$y = c_1(\csc(2x) + \cot(2x))^{-\text{signum}(\sin(2x))} + c_2(\csc(2x) + \cot(2x))^{\text{signum}(\sin(2x))}$$

Verified OK.

4.72.2 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$y'' \sin(2x)^2 + y' \sin(4x) - 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= 2 \cot(2x) \\ q(x) &= -4 \csc(2x)^2 \end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{2\sqrt{-\csc(2x)^2}}{c} \\ \tau'' &= \frac{4 \cot(2x) \csc(2x)^2}{c\sqrt{-\csc(2x)^2}} \end{aligned} \quad (6)$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\
 &= \frac{\frac{4 \cot(2x) \csc(2x)^2}{c\sqrt{-\csc(2x)^2}} + 2 \cot(2x) \frac{2\sqrt{-\csc(2x)^2}}{c}}{\left(\frac{2\sqrt{-\csc(2x)^2}}{c}\right)^2} \\
 &= 0
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c} \sqrt{q} dx \\
 &= \frac{\int 2\sqrt{-\csc(2x)^2} dx}{c} \\
 &= \frac{\sqrt{-\csc(2x)^2} \ln(-\cot(2x) + \csc(2x)) \sin(2x)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = -i \cot(2x) c_2 + c_1 \cosh(\ln(-\cot(2x) + \csc(2x)))$$

Summary

The solution(s) found are the following

$$y = -i \cot(2x) c_2 + c_1 \left(-\frac{\cot(2x)}{2} + \frac{\csc(2x)}{2} + \frac{1}{-2 \cot(2x) + 2 \csc(2x)} \right) \tag{1}$$

Verification of solutions

$$y = -i \cot(2x) c_2 + c_1 \left(-\frac{\cot(2x)}{2} + \frac{\csc(2x)}{2} + \frac{1}{-2 \cot(2x) + 2 \csc(2x)} \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
    trying a quadrature
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful
Change of variables used:
    [x = 1/4*arccos(t)]
Linear ODE actually solved:
    -u(t)+(3*t^2-2*t-1)*diff(u(t),t)+(2*t^3-2*t^2-2*t+2)*diff(diff(u(t),t),t) = 0
<- change of variables successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)*sin(2*x)^2+diff(y(x),x)*sin(4*x)-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \csc(2x) + \cot(2x) c_2$$

✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 29

```
DSolve[y''[x]*Sin[2*x]^2+y'[x]*Sin[4*x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 - i c_2 \cos(2x)}{\sqrt{\sin^2(2x)}}$$

5 section 5.0

5.1	problem 1	2342
5.2	problem 2	2348
5.3	problem 3	2357
5.4	problem 4	2364
5.5	problem 5	2375
5.6	problem 6	2388
5.7	problem 7	2397
5.8	problem 8	2404
5.9	problem 9	2412
5.10	problem 10	2418
5.11	problem 11	2422
5.12	problem 12	2426
5.13	problem 13	2430
5.14	problem 14	2434
5.15	problem 15	2438
5.16	problem 16	2442
5.17	problem 17	2455
5.18	problem 18	2462
5.19	problem 19	2475
5.20	problem 20	2487
5.21	problem 21	2496
5.22	problem 22	2509
5.23	problem 23	2516

5.1 problem 1

5.1.1 Solving as second order ode can be made integrable ode 2342

5.1.2 Solving as second order ode missing x ode 2344

Internal problem ID [7294]

Internal file name [OUTPUT/6280_Sunday_June_05_2022_04_37_01_PM_50628687/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 1.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**", "**second_order_ode_can_be_made_integrable**"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x], [_2nd_order , _reducible , _mu_x_y1]]
```

$$y'' - Ay^{\frac{2}{3}} = 0$$

5.1.1 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' - Ay^{\frac{2}{3}}y' = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' - Ay^{\frac{2}{3}}y') dx = 0$$
$$\frac{y'^2}{2} - \frac{3Ay^{\frac{5}{3}}}{5} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{30Ay^{\frac{5}{3}} + 50c_1}}{5} \tag{1}$$

$$y' = -\frac{\sqrt{30Ay^{\frac{5}{3}} + 50c_1}}{5} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{5}{\sqrt{30A y^{\frac{5}{3}} + 50c_1}} dy = \int dx$$
$$5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_2$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{5}{\sqrt{30A y^{\frac{5}{3}} + 50c_1}} dy = \int dx$$
$$-5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_3$$

Summary

The solution(s) found are the following

$$5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_2 \quad (1)$$

$$-5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_3 \quad (2)$$

Verification of solutions

$$5 \left(\int^y \frac{1}{\sqrt{30A - a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_2$$

Verified OK.

$$-5 \left(\int^y \frac{1}{\sqrt{30A - a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_3$$

Verified OK.

5.1.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) - A y^{\frac{2}{3}} = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{A y^{\frac{2}{3}}}{p} \end{aligned}$$

Where $f(y) = Ay^{\frac{2}{3}}$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\frac{1}{p} dp = Ay^{\frac{2}{3}} dy$$

$$\int \frac{1}{p} dp = \int Ay^{\frac{2}{3}} dy$$

$$\frac{p^2}{2} = \frac{3Ay^{\frac{5}{3}}}{5} + c_1$$

The solution is

$$\frac{p(y)^2}{2} - \frac{3Ay^{\frac{5}{3}}}{5} - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} - \frac{3Ay^{\frac{5}{3}}}{5} - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{30Ay^{\frac{5}{3}} + 50c_1}}{5} \tag{1}$$

$$y' = -\frac{\sqrt{30Ay^{\frac{5}{3}} + 50c_1}}{5} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{5}{\sqrt{30Ay^{\frac{5}{3}} + 50c_1}} dy = \int dx$$

$$5 \left(\int^y \frac{1}{\sqrt{30A_a^{\frac{5}{3}} + 50c_1}} d_a \right) = x + c_2$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{5}{\sqrt{30A y^{\frac{5}{3}} + 50c_1}} dy = \int dx$$
$$-5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_3$$

Summary

The solution(s) found are the following

$$5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_2 \quad (1)$$

$$-5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_3 \quad (2)$$

Verification of solutions

$$5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_2$$

Verified OK.

$$-5 \left(\int^y \frac{1}{\sqrt{30A a^{\frac{5}{3}} + 50c_1}} da \right) = x + c_3$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-A*_a^(2/3) = 0, _b(_a), HINT = [
    symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, 5/6*_b]
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 61

```
dsolve(diff(y(x),x$2)=A*y(x)^(2/3),y(x), singsol=all)
```

$$y(x) = 0$$
$$-5 \left(\int^{y(x)} \frac{1}{\sqrt{30_a^{\frac{5}{3}}A - 5c_1}} d_a \right) - x - c_2 = 0$$
$$5 \left(\int^{y(x)} \frac{1}{\sqrt{30_a^{\frac{5}{3}}A - 5c_1}} d_a \right) - x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 75

```
DSolve[y''[x]==A*y[x]^(2/3),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{y(x)^2 \left(1 + \frac{6Ay(x)^{5/3}}{5c_1} \right) \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{3}{5}, \frac{8}{5}, -\frac{6Ay(x)^{5/3}}{5c_1} \right)^2}{\frac{6}{5}Ay(x)^{5/3} + c_1} = (x+c_2)^2, y(x) \right]$$

5.2 problem 2

5.2.1 Solving as linear second order ode solved by an integrating factor ode	2348
5.2.2 Solving as second order change of variable on y method 1 ode .	2349
5.2.3 Solving using Kovacic algorithm	2351
5.2.4 Maple step by step solution	2354

Internal problem ID [7295]

Internal file name [OUTPUT/6281_Sunday_June_05_2022_04_37_05_PM_55717402/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_change_of_variable_on_y_method_1", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 2xy' + (x^2 + 1)y = 0$$

5.2.1 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x))^2 + p'(x)}{2}y = f(x)$$

Where $p(x) = 2x$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p \, dx} \\ &= e^{\int 2x \, dx} \\ &= e^{\frac{x^2}{2}} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = 0$$
$$\left(e^{\frac{x^2}{2}}y\right)'' = 0$$

Integrating once gives

$$\left(e^{\frac{x^2}{2}}y\right)' = c_1$$

Integrating again gives

$$\left(e^{\frac{x^2}{2}}y\right) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{e^{\frac{x^2}{2}}}$$

Or

$$y = c_1x e^{-\frac{x^2}{2}} + c_2e^{-\frac{x^2}{2}}$$

Summary

The solution(s) found are the following

$$y = c_1x e^{-\frac{x^2}{2}} + c_2e^{-\frac{x^2}{2}} \quad (1)$$

Verification of solutions

$$y = c_1x e^{-\frac{x^2}{2}} + c_2e^{-\frac{x^2}{2}}$$

Verified OK.

5.2.2 Solving as second order change of variable on y method 1 ode

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = 2x$$
$$q(x) = x^2 + 1$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= x^2 + 1 - \frac{(2x)'}{2} - \frac{(2x)^2}{4} \\ &= x^2 + 1 - \frac{(2)}{2} - \frac{(4x^2)}{4} \\ &= x^2 + 1 - (1) - x^2 \\ &= 0 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x) z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\left(\int \frac{p(x)}{2} dx\right)} \\ &= e^{-\int \frac{2x}{2}} \\ &= e^{-\frac{x^2}{2}} \end{aligned} \tag{5}$$

Hence (3) becomes

$$y = v(x) e^{-\frac{x^2}{2}} \tag{4}$$

Applying this change of variable to the original ode results in

$$v''(x) e^{-\frac{x^2}{2}} = 0$$

Which is now solved for $v(x)$ Integrating twice gives the solution

$$v(x) = c_1 x + c_2$$

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1 x + c_2) (z(x)) \end{aligned} \tag{7}$$

But from (5)

$$z(x) = e^{-\frac{x^2}{2}}$$

Hence (7) becomes

$$y = e^{-\frac{x^2}{2}}(c_1x + c_2)$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x^2}{2}}(c_1x + c_2) \quad (1)$$

Verification of solutions

$$y = e^{-\frac{x^2}{2}}(c_1x + c_2)$$

Verified OK.

5.2.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2xy' + (x^2 + 1)y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2x \\ C &= x^2 + 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 244: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2x}{1} dx} \\ &= z_1 e^{-\frac{x^2}{2}} \\ &= z_1 \left(e^{-\frac{x^2}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x^2}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x^2}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x^2}{2}} \right) + c_2 \left(e^{-\frac{x^2}{2}}(x) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{x^2}{2}} + c_2 x e^{-\frac{x^2}{2}} \quad (1)$$

Verification of solutions

$$y = c_1 e^{-\frac{x^2}{2}} + c_2 x e^{-\frac{x^2}{2}}$$

Verified OK.

5.2.4 Maple step by step solution

Let's solve

$$y'' + 2xy' + (x^2 + 1)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=\max(0,-m)}^{\infty} a_k x^{k+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=\max(0,-m)+m}^{\infty} a_{k-m} x^k$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + a_0 + (6a_3 + 3a_1)x + \left(\sum_{k=2}^{\infty} (a_{k+2}(k+2)(k+1) + a_k(2k+1) + a_{k-2}) x^k \right) = 0$$

- The coefficients of each power of x must be 0

$$[2a_2 + a_0 = 0, 6a_3 + 3a_1 = 0]$$

- Solve for the dependent coefficient(s)

$$\{a_2 = -\frac{a_0}{2}, a_3 = -\frac{a_1}{2}\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} + 2a_k k + a_k + a_{k-2} = 0$$

- Shift index using $k \rightarrow k + 2$

$$((k + 2)^2 + 3k + 8) a_{k+4} + 2a_{k+2}(k + 2) + a_{k+2} + a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = -\frac{2ka_{k+2} + a_k + 5a_{k+2}}{k^2 + 7k + 12}, a_2 = -\frac{a_0}{2}, a_3 = -\frac{a_1}{2} \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x), x$2) + 2*x*diff(y(x), x) + (x^2 + 1)*y(x) = 0, y(x), singsol=all)
```

$$y(x) = e^{-\frac{x^2}{2}} (c_2 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 22

```
DSolve[y''[x]+2*x*y'[x]+(x^2+1)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{x^2}{2}}(c_2x + c_1)$$

5.3 problem 3

5.3.1 Solving as linear second order ode solved by an integrating factor ode	2357
5.3.2 Solving as second order change of variable on y method 1 ode	2358
5.3.3 Solving using Kovacic algorithm	2360

Internal problem ID [7296]

Internal file name [OUTPUT/6282_Sunday_June_05_2022_04_37_06_PM_80670546/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**", "**second_order_change_of_variable_on_y_method_1**", "**linear_second_order_ode_solved_by_an_integrating_factor**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 2 \cot(x) y' - y = 0$$

5.3.1 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x) y' + \frac{(p(x))^2 + p'(x)}{2} y = f(x)$$

Where $p(x) = 2 \cot(x)$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p \, dx} \\ &= e^{\int 2 \cot(x) \, dx} \\ &= \sin(x) \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = 0$$

$$(\sin(x)y)'' = 0$$

Integrating once gives

$$(\sin(x)y)' = c_1$$

Integrating again gives

$$(\sin(x)y) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{\sin(x)}$$

Or

$$y = \frac{c_1x}{\sin(x)} + \frac{c_2}{\sin(x)}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1x}{\sin(x)} + \frac{c_2}{\sin(x)} \quad (1)$$

Verification of solutions

$$y = \frac{c_1x}{\sin(x)} + \frac{c_2}{\sin(x)}$$

Verified OK.

5.3.2 Solving as second order change of variable on y method 1 ode

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = 2 \cot(x)$$

$$q(x) = -1$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned}
 Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\
 &= -1 - \frac{(2 \cot(x))'}{2} - \frac{(2 \cot(x))^2}{4} \\
 &= -1 - \frac{(-2 - 2 \cot(x)^2)}{2} - \frac{(4 \cot(x)^2)}{4} \\
 &= -1 - (-1 - \cot(x)^2) - \cot(x)^2 \\
 &= 0
 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x) z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned}
 z(x) &= e^{-\left(\int \frac{p(x)}{2} dx\right)} \\
 &= e^{-\int \frac{2 \cot(x)}{2}} \\
 &= \csc(x)
 \end{aligned} \quad (5)$$

Hence (3) becomes

$$y = v(x) \csc(x) \quad (4)$$

Applying this change of variable to the original ode results in

$$v''(x) \csc(x) = 0$$

Which is now solved for $v(x)$ Integrating twice gives the solution

$$v(x) = c_1 x + c_2$$

Now that $v(x)$ is known, then

$$\begin{aligned}
 y &= v(x) z(x) \\
 &= (c_1 x + c_2) (z(x))
 \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \csc(x)$$

Hence (7) becomes

$$y = \csc(x) (c_1x + c_2)$$

Summary

The solution(s) found are the following

$$y = \csc(x) (c_1x + c_2) \quad (1)$$

Verification of solutions

$$y = \csc(x) (c_1x + c_2)$$

Verified OK.

5.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2 \cot(x) y' - y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2 \cot(x) \\ C &= -1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 246: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2 \cot(x)}{1} dx} \\ &= z_1 e^{-\ln(\sin(x))} \\ &= z_1 (\csc(x)) \end{aligned}$$

Which simplifies to

$$y_1 = \csc(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2 \cot(x)}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2 \ln(\sin(x))}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\csc(x)) + c_2 (\csc(x) (x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \csc(x) c_1 + c_2 x \csc(x) \tag{1}$$

Verification of solutions

$$y = \csc(x) c_1 + c_2 x \csc(x)$$

Verified OK.

Maple trace **Kovacic algorithm successful**

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x$2)+2*cot(x)*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \csc(x) (c_2 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.058 (sec). Leaf size: 15

```
DSolve[y''[x]+2*Cot[x]*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (c_2 x + c_1) \csc(x)$$

5.4 problem 4

- 5.4.1 Solving as second order change of variable on y method 1 ode . 2364
- 5.4.2 Solving as second order bessel ode ode 2367
- 5.4.3 Solving using Kovacic algorithm 2368
- 5.4.4 Maple step by step solution 2371

Internal problem ID [7297]

Internal file name [OUTPUT/6283_Sunday_June_05_2022_04_37_08_PM_91391385/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_bessel_ode", "second_order_change_of_variable_on_y_method_1"

Maple gives the following as the ode type

[[_2nd_order , _with_linear_symmetries]]

$$x^2 y'' + x y' + \left(x^2 - \frac{1}{4}\right) y = 0$$

5.4.1 Solving as second order change of variable on y method 1 ode

In normal form the given ode is written as

$$y'' + p(x) y' + q(x) y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{x^2 - \frac{1}{4}}{x^2}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned}
 Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\
 &= \frac{x^2 - \frac{1}{4}}{x^2} - \frac{\left(\frac{1}{x}\right)'}{2} - \frac{\left(\frac{1}{x}\right)^2}{4} \\
 &= \frac{x^2 - \frac{1}{4}}{x^2} - \frac{\left(-\frac{1}{x^2}\right)}{2} - \frac{\left(\frac{1}{x^2}\right)}{4} \\
 &= \frac{x^2 - \frac{1}{4}}{x^2} - \left(-\frac{1}{2x^2}\right) - \frac{1}{4x^2} \\
 &= 1
 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x) z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned}
 z(x) &= e^{-\left(\int \frac{p(x)}{2} dx\right)} \\
 &= e^{-\int \frac{1}{2}} \\
 &= \frac{1}{\sqrt{x}}
 \end{aligned} \tag{5}$$

Hence (3) becomes

$$y = \frac{v(x)}{\sqrt{x}} \tag{4}$$

Applying this change of variable to the original ode results in

$$x^{\frac{3}{2}}(v''(x) + v(x)) = 0$$

Which is now solved for $v(x)$ This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$v(x) = c_1 \cos(x) + c_2 \sin(x)$$

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1 \cos(x) + c_2 \sin(x)) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \frac{1}{\sqrt{x}}$$

Hence (7) becomes

$$y = \frac{c_1 \cos(x) + c_2 \sin(x)}{\sqrt{x}}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \cos(x) + c_2 \sin(x)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \cos(x) + c_2 \sin(x)}{\sqrt{x}}$$

Verified OK.

5.4.2 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0 \quad (1)$$

Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\alpha = 0$$

$$\beta = 1$$

$$n = -\frac{1}{2}$$

$$\gamma = 1$$

Substituting all the above into (4) gives the solution as

$$y = \frac{c_1\sqrt{2} \cos(x)}{\sqrt{\pi} \sqrt{x}} + \frac{c_2\sqrt{2} \sin(x)}{\sqrt{\pi} \sqrt{x}}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1\sqrt{2} \cos(x)}{\sqrt{\pi} \sqrt{x}} + \frac{c_2\sqrt{2} \sin(x)}{\sqrt{\pi} \sqrt{x}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1\sqrt{2} \cos(x)}{\sqrt{\pi} \sqrt{x}} + \frac{c_2\sqrt{2} \sin(x)}{\sqrt{\pi} \sqrt{x}}$$

Verified OK.

5.4.3 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + xy' + \left(x^2 - \frac{1}{4}\right)y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= x \\ C &= x^2 - \frac{1}{4} \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 247: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx} \\ &= z_1 e^{-\frac{\ln(x)}{2}} \\ &= z_1 \left(\frac{1}{\sqrt{x}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{\cos(x)}{\sqrt{x}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\ &= y_1 (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
 y &= c_1 y_1 + c_2 y_2 \\
 &= c_1 \left(\frac{\cos(x)}{\sqrt{x}} \right) + c_2 \left(\frac{\cos(x)}{\sqrt{x}} (\tan(x)) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \cos(x)}{\sqrt{x}} + \frac{c_2 \sin(x)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \cos(x)}{\sqrt{x}} + \frac{c_2 \sin(x)}{\sqrt{x}}$$

Verified OK.

5.4.4 Maple step by step solution

Let's solve

$$x^2 y'' + x y' + \left(x^2 - \frac{1}{4}\right) y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(4x^2-1)y}{4x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(4x^2-1)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{4x^2-1}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{1}{4}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2y'' + 4xy' + (4x^2 - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+2r)(-1+2r)x^r + a_1(3+2r)(1+2r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r+1)(2k+2r-1) + 4a_{k-1}) \right) x^{k+r}$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+2r)(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ -\frac{1}{2}, \frac{1}{2} \right\}$$

- Each term must be 0

$$a_1(3+2r)(1+2r) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(4k^2 + 8kr + 4r^2 - 1) + 4a_{k-2} = 0$$

- Shift index using $k \rightarrow k + 2$

$$a_{k+2}(4(k+2)^2 + 8(k+2)r + 4r^2 - 1) + 4a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{4a_k}{4k^2 + 8kr + 4r^2 + 16k + 16r + 15}$$

- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}$$

- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}, a_1 = 0 \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}$$

- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}, a_1 = 0, b_{k+2} = -\frac{4b_k}{4k^2 + 20k + 24}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(x^2*diff(diff(y(x),x),x)+x*diff(y(x),x)+(x^2-1/4)*y(x) = 0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \sin(x) + c_2 \cos(x)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 39

```
DSolve[x^2*y'[x]+x*y'[x]+(x^2-1/4)*y[x] == 0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-ix}(2c_1 - ic_2 e^{2ix})}{2\sqrt{x}}$$

5.5 problem 5

- 5.5.1 Solving as linear second order ode solved by an integrating factor ode 2375
- 5.5.2 Solving as second order change of variable on y method 1 ode . 2377
- 5.5.3 Solving using Kovacic algorithm 2381

Internal problem ID [7298]

Internal file name [OUTPUT/6284_Sunday_June_05_2022_04_37_11_PM_48763785/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**", "**second_order_change_of_variable_on_y_method_1**", "**linear_second_order_ode_solved_by_an_integrating_factor**"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$4x^2y'' + (-8x^2 + 4x)y' + (4x^2 - 4x - 1)y = 4\sqrt{x}e^x$$

5.5.1 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = \frac{1-2x}{x}$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p \, dx} \\ &= e^{\int \frac{1-2x}{x} \, dx} \\ &= \sqrt{x} e^{-x}\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = \frac{e^{-x}e^x}{x}$$

$$(\sqrt{x}e^{-x}y)'' = \frac{e^{-x}e^x}{x}$$

Integrating once gives

$$(\sqrt{x}e^{-x}y)' = \ln(x) + c_1$$

Integrating again gives

$$(\sqrt{x}e^{-x}y) = x(\ln(x) + c_1 - 1) + c_2$$

Hence the solution is

$$y = \frac{x(\ln(x) + c_1 - 1) + c_2}{\sqrt{x}e^{-x}}$$

Or

$$y = c_1\sqrt{x}e^x + \sqrt{x}e^x \ln(x) + \frac{c_2e^x}{\sqrt{x}} - \sqrt{x}e^x$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x}e^x + \sqrt{x}e^x \ln(x) + \frac{c_2e^x}{\sqrt{x}} - \sqrt{x}e^x \quad (1)$$

Verification of solutions

$$y = c_1\sqrt{x}e^x + \sqrt{x}e^x \ln(x) + \frac{c_2e^x}{\sqrt{x}} - \sqrt{x}e^x$$

Verified OK.

5.5.2 Solving as second order change of variable on y method 1 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$4x^2y'' + (-8x^2 + 4x)y' + (4x^2 - 4x - 1)y = 0$$

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{-8x^2 + 4x}{4x^2}$$

$$q(x) = \frac{4x^2 - 4x - 1}{4x^2}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= \frac{4x^2 - 4x - 1}{4x^2} - \frac{\left(\frac{-8x^2+4x}{4x^2}\right)'}{2} - \frac{\left(\frac{-8x^2+4x}{4x^2}\right)^2}{4} \\ &= \frac{4x^2 - 4x - 1}{4x^2} - \frac{\left(\frac{-16x+4}{4x^2} - \frac{-8x^2+4x}{2x^3}\right)}{2} - \frac{\left(\frac{(-8x^2+4x)^2}{16x^4}\right)}{4} \\ &= \frac{4x^2 - 4x - 1}{4x^2} - \left(\frac{-16x+4}{8x^2} - \frac{-8x^2+4x}{4x^3}\right) - \frac{(-8x^2+4x)^2}{64x^4} \\ &= 0 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\left(\int \frac{p(x)}{2} dx\right)} \\ &= e^{-\int \frac{-8x^2+4x}{4x^2} dx} \\ &= \frac{e^x}{\sqrt{x}} \end{aligned} \tag{5}$$

Hence (3) becomes

$$y = \frac{v(x) e^x}{\sqrt{x}} \tag{4}$$

Applying this change of variable to the original ode results in

$$xv''(x) = 1$$

Which is now solved for $v(x)$ Simplifying the ode gives

$$v''(x) = \frac{1}{x}$$

Integrating once gives

$$v'(x) = \ln(x) + c_1$$

Integrating again gives

$$v(x) = x \ln(x) - x + c_1x + c_2$$

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1x + x \ln(x) - x + c_2) (z(x)) \end{aligned} \tag{7}$$

But from (5)

$$z(x) = \frac{e^x}{\sqrt{x}}$$

Hence (7) becomes

$$y = \frac{e^x (c_1x + x \ln(x) - x + c_2)}{\sqrt{x}}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{e^x(c_1x + x \ln(x) - x + c_2)}{\sqrt{x}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x} e^x$$

$$y_2 = \frac{e^x}{\sqrt{x}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x} e^x & \frac{e^x}{\sqrt{x}} \\ \frac{d}{dx}(\sqrt{x} e^x) & \frac{d}{dx}\left(\frac{e^x}{\sqrt{x}}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x} e^x & \frac{e^x}{\sqrt{x}} \\ \frac{e^x}{2\sqrt{x}} + \sqrt{x} e^x & -\frac{e^x}{2x^{\frac{3}{2}}} + \frac{e^x}{\sqrt{x}} \end{vmatrix}$$

Therefore

$$W = (\sqrt{x} e^x) \left(-\frac{e^x}{2x^{\frac{3}{2}}} + \frac{e^x}{\sqrt{x}} \right) - \left(\frac{e^x}{\sqrt{x}} \right) \left(\frac{e^x}{2\sqrt{x}} + \sqrt{x} e^x \right)$$

Which simplifies to

$$W = -\frac{e^{2x}}{x}$$

Which simplifies to

$$W = -\frac{e^{2x}}{x}$$

Therefore Eq. (2) becomes

$$u_1 = -\int \frac{4(e^x)^2}{-4x e^{2x}} dx$$

Which simplifies to

$$u_1 = -\int -\frac{1}{x} dx$$

Hence

$$u_1 = \ln(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{4x(e^x)^2}{-4x e^{2x}} dx$$

Which simplifies to

$$u_2 = \int (-1) dx$$

Hence

$$u_2 = -x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \sqrt{x} e^x \ln(x) - \sqrt{x} e^x$$

Which simplifies to

$$y_p(x) = \sqrt{x} (\ln(x) - 1) e^x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{e^x (c_1 x + x \ln(x) - x + c_2)}{\sqrt{x}} \right) + (\sqrt{x} (\ln(x) - 1) e^x) \end{aligned}$$

Which simplifies to

$$y = \frac{e^x (x \ln(x) + (c_1 - 1)x + c_2)}{\sqrt{x}} + \sqrt{x} (\ln(x) - 1) e^x$$

Summary

The solution(s) found are the following

$$y = \frac{e^x (x \ln(x) + (c_1 - 1)x + c_2)}{\sqrt{x}} + \sqrt{x} (\ln(x) - 1) e^x \quad (1)$$

Verification of solutions

$$y = \frac{e^x (x \ln(x) + (c_1 - 1)x + c_2)}{\sqrt{x}} + \sqrt{x} (\ln(x) - 1) e^x$$

Verified OK.

5.5.3 Solving using Kovacic algorithm

Writing the ode as

$$4x^2 y'' + (-8x^2 + 4x) y' + (4x^2 - 4x - 1) y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 4x^2 \\ B &= -8x^2 + 4x \\ C &= 4x^2 - 4x - 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 249: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-8x^2+4x}{4x^2} dx} \\ &= z_1 e^{x - \frac{\ln(x)}{2}} \\ &= z_1 \left(\frac{e^x}{\sqrt{x}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{e^x}{\sqrt{x}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-8x^2+4x}{4x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x - \ln(x)}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{e^x}{\sqrt{x}} \right) + c_2 \left(\frac{e^x}{\sqrt{x}}(x) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$4x^2 y'' + (-8x^2 + 4x) y' + (4x^2 - 4x - 1) y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1 e^x}{\sqrt{x}} + \sqrt{x} e^x c_2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \frac{e^x}{\sqrt{x}} \\ y_2 &= \sqrt{x} e^x \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{e^x}{\sqrt{x}} & \sqrt{x} e^x \\ \frac{d}{dx} \left(\frac{e^x}{\sqrt{x}} \right) & \frac{d}{dx} (\sqrt{x} e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{e^x}{\sqrt{x}} & \sqrt{x} e^x \\ -\frac{e^x}{2x^{\frac{3}{2}}} + \frac{e^x}{\sqrt{x}} & \frac{e^x}{2\sqrt{x}} + \sqrt{x} e^x \end{vmatrix}$$

Therefore

$$W = \left(\frac{e^x}{\sqrt{x}} \right) \left(\frac{e^x}{2\sqrt{x}} + \sqrt{x} e^x \right) - (\sqrt{x} e^x) \left(-\frac{e^x}{2x^{\frac{3}{2}}} + \frac{e^x}{\sqrt{x}} \right)$$

Which simplifies to

$$W = \frac{e^{2x}}{x}$$

Which simplifies to

$$W = \frac{e^{2x}}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{4x e^{2x}}{4x e^{2x}} dx$$

Which simplifies to

$$u_1 = - \int 1 dx$$

Hence

$$u_1 = -x$$

And Eq. (3) becomes

$$u_2 = \int \frac{4 e^{2x}}{4x e^{2x}} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x} dx$$

Hence

$$u_2 = \ln(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \sqrt{x} e^x \ln(x) - \sqrt{x} e^x$$

Which simplifies to

$$y_p(x) = \sqrt{x} (\ln(x) - 1) e^x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1 e^x}{\sqrt{x}} + \sqrt{x} e^x c_2 \right) + (\sqrt{x} (\ln(x) - 1) e^x) \end{aligned}$$

Which simplifies to

$$y = \frac{e^x(c_2 x + c_1)}{\sqrt{x}} + \sqrt{x} (\ln(x) - 1) e^x$$

Summary

The solution(s) found are the following

$$y = \frac{e^x(c_2 x + c_1)}{\sqrt{x}} + \sqrt{x} (\ln(x) - 1) e^x \quad (1)$$

Verification of solutions

$$y = \frac{e^x(c_2 x + c_1)}{\sqrt{x}} + \sqrt{x} (\ln(x) - 1) e^x$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(4*x^2*diff(diff(y(x),x),x)+(-8*x^2+4*x)*diff(y(x),x)+(4*x^2-4*x-1)*y(x) = 4*x^(1/2)*e
```

$$y(x) = \frac{(x \ln(x) + (-1 + c_1)x + c_2)e^x}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 27

```
DSolve[4*x^2*y''[x]+(-8*x^2+4*x)*y'[x]+(4*x^2-4*x-1)*y[x] == 4*x^(1/2)*Exp[x],y[x],x,Include
```

$$y(x) \rightarrow \frac{e^x(x \log(x) + (-1 + c_2)x + c_1)}{\sqrt{x}}$$

5.6 problem 6

5.6.1 Solving using Kovacic algorithm 2388

Internal problem ID [7299]

Internal file name [OUTPUT/6285_Sunday_June_05_2022_04_37_14_PM_49765877/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

`[[_2nd_order , _linear , _nonhomogeneous]]`

$$xy'' - (2x + 2)y' + (x + 2)y = 6x^3e^x$$

5.6.1 Solving using Kovacic algorithm

Writing the ode as

$$xy'' + (-2x - 2)y' + (x + 2)y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x \\ B &= -2x - 2 \\ C &= x + 2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{x^2} \tag{6}$$

Comparing the above to (5) shows that

$$s = 2$$

$$t = x^2$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{2}{x^2}\right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 250: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= -\frac{1}{x} + (-) (0) \\ &= -\frac{1}{x} \\ &= -\frac{1}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{x}\right)(0) + \left(\left(\frac{1}{x^2}\right) + \left(-\frac{1}{x}\right)^2 - \left(\frac{2}{x^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(x) = pe^{\int \omega dx}$$
$$= e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$
$$= z_1 e^{-\int \frac{1}{2} \frac{-2x-2}{x} dx}$$
$$= z_1 e^{x+\ln(x)}$$
$$= z_1 (x e^x)$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{-2x-2}{x} dx}}{(y_1)^2} dx$$
$$= y_1 \int \frac{e^{2x+2\ln(x)}}{(y_1)^2} dx$$
$$= y_1 \left(\frac{x^3}{3}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^x) + c_2 \left(e^x \left(\frac{x^3}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$xy'' + (-2x - 2)y' + (x + 2)y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + \frac{c_2 x^3 e^x}{3}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= e^x \\ y_2 &= \frac{x^3 e^x}{3}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^x & \frac{x^3 e^x}{3} \\ \frac{d}{dx}(e^x) & \frac{d}{dx}\left(\frac{x^3 e^x}{3}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^x & \frac{x^3 e^x}{3} \\ e^x & x^2 e^x + \frac{x^3 e^x}{3} \end{vmatrix}$$

Therefore

$$W = (e^x) \left(x^2 e^x + \frac{x^3 e^x}{3} \right) - \left(\frac{x^3 e^x}{3} \right) (e^x)$$

Which simplifies to

$$W = x^2 e^{2x}$$

Which simplifies to

$$W = x^2 e^{2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2x^6 e^{2x}}{x^3 e^{2x}} dx$$

Which simplifies to

$$u_1 = - \int 2x^3 dx$$

Hence

$$u_1 = - \frac{x^4}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{6x^3 e^{2x}}{x^3 e^{2x}} dx$$

Which simplifies to

$$u_2 = \int 6dx$$

Hence

$$u_2 = 6x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{3x^4e^x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^x + \frac{c_2x^3e^x}{3} \right) + \left(\frac{3x^4e^x}{2} \right) \end{aligned}$$

Which simplifies to

$$y = e^x \left(c_1 + \frac{c_2x^3}{3} \right) + \frac{3x^4e^x}{2}$$

Summary

The solution(s) found are the following

$$y = e^x \left(c_1 + \frac{c_2x^3}{3} \right) + \frac{3x^4e^x}{2} \tag{1}$$

Verification of solutions

$$y = e^x \left(c_1 + \frac{c_2x^3}{3} \right) + \frac{3x^4e^x}{2}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x*diff(diff(y(x),x),x)-(2*x+2)*diff(y(x),x)+(2+x)*y(x) = 6*x^3*exp(x),y(x), singsol=a
```

$$y(x) = e^x \left(c_2 + c_1 x^3 + \frac{3}{2} x^4 \right)$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 29

```
DSolve[x*y'[x]-(2*x+2)*y'[x]+(2+x)*y[x] == 6*x^3*Exp[x],y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \frac{1}{6} e^x (9x^4 + 2c_2 x^3 + 6c_1)$$

5.7 problem 7

5.7.1 Solving as series ode	2397
5.7.2 Maple step by step solution	2402

Internal problem ID [7300]

Internal file name [OUTPUT/6286_Sunday_June_05_2022_04_37_17_PM_68930396/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 7.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

Unable to solve or complete the solution.

$$y' + y = \frac{1}{x}$$

With the expansion point for the power series method at $x = 0$.

5.7.1 Solving as series ode

Writing the ODE as

$$y' + q(x)y = p(x)$$
$$y' + y = \frac{1}{x}$$

Where

$$q(x) = 1$$
$$p(x) = \frac{1}{x}$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular

singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point if $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not.

Since $x = 0$ is not an ordinary point, we now check to see if it is a regular singular point. Since $x = 0$ is regular singular point, then Frobenius power series is used. Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $y' + y = 0$, and y_p is a particular solution to the inhomogeneous ode. First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} a_n x^{n+r} = \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2) this gives

$$(n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$r a_0 x^{-1+r} = 0$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$m c_0 x^{-1+m} = \frac{1}{x}$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation simplifies to

$$r = 0$$

Solving for r gives the root of the indicial equation as

$$r = 0$$

We start by finding y_n . For $1 \leq n$, the recurrence equation is

$$a_n(n+r) + a_{n-1} = 0 \quad (4)$$

For $n = 1$ the recurrence equation gives

$$a_1(1+r) + a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_1 = -\frac{a_0}{1+r}$$

For $n = 2$ the recurrence equation gives

$$a_2(2 + r) + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_2 = \frac{a_0}{(1 + r)(2 + r)}$$

For $n = 3$ the recurrence equation gives

$$a_3(3 + r) + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{(1 + r)(2 + r)(3 + r)}$$

For $n = 4$ the recurrence equation gives

$$a_4(4 + r) + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{(1 + r)(2 + r)(3 + r)(4 + r)}$$

For $n = 5$ the recurrence equation gives

$$a_5(5 + r) + a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_0}{(1 + r)(2 + r)(3 + r)(4 + r)(5 + r)}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^{n+r} \\ &= a_0 x^r + a_1 x^{1+r} + a_2 x^{2+r} + a_3 x^{3+r} + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 x^r - \frac{a_0 x^{1+r}}{1+r} + \frac{a_0 x^{2+r}}{(1+r)(2+r)} - \frac{a_0 x^{3+r}}{(1+r)(2+r)(3+r)} + \frac{a_0 x^{4+r}}{(1+r)(2+r)(3+r)(4+r)} - \frac{a_0 x^{5+r}}{(1+r)(2+r)(3+r)(4+r)(5+r)} + \dots$$

Which can be written as

$$y = x^r \left(a_0 - \frac{a_0 x}{1+r} + \frac{a_0 x^2}{(1+r)(2+r)} - \frac{a_0 x^3}{(1+r)(2+r)(3+r)} + \frac{a_0 x^4}{(1+r)(2+r)(3+r)(4+r)} - \frac{a_0 x^5}{(1+r)(2+r)(3+r)(4+r)(5+r)} + O(x^6) a_0 \right)$$

Collecting terms, the solution becomes

$$y = x^r \left(1 - \frac{x}{1+r} + \frac{x^2}{(1+r)(2+r)} - \frac{x^3}{(1+r)(2+r)(3+r)} + \frac{x^4}{(1+r)(2+r)(3+r)(4+r)} - \frac{x^5}{(1+r)(2+r)(3+r)(4+r)(5+r)} + \dots \right) \quad (3)$$

Finally, since $r = 0$, then the solution becomes

$$y = \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6) \right) a_0 \quad (3)$$

Therefore the homogeneous solution is

$$y_h(x) = \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6) \right) a_0$$

Unable to solve the balance equation $mc_0 x^{-1+m} = \frac{1}{x}$ for c_0 and x . No particular solution exists.

Unable to find the particular solution. No solution exist.

Verification of solutions N/A

5.7.2 Maple step by step solution

Let's solve

$$y' + y = \frac{1}{x}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -y + \frac{1}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + y = \frac{1}{x}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x)(y' + y) = \frac{\mu(x)}{x}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y' + y) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{\mu(x)}{x} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)}{x} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)}{x} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^x$

$$y = \frac{\int \frac{e^x}{x} dx + c_1}{e^x}$$

- Evaluate the integrals on the rhs

$$y = \frac{-\text{Ei}_1(-x) + c_1}{e^x}$$

- Simplify

$$y = e^{-x}(-\text{Ei}_1(-x) + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✗ Solution by Maple

```

Order:=6;
dsolve(diff(y(x),x)+y(x)=1/x,y(x),type='series',x=0);

```

No solution found

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 113

```

AsymptoticDSolveValue[y'[x]+y[x]==1/x,y[x],{x,0,5}]

```

$$y(x) \rightarrow \left(-\frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right) \left(\frac{x^6}{2160} + \frac{x^5}{600} + \frac{x^4}{96} + \frac{x^3}{18} + \frac{x^2}{4} + x + \log(x) \right) + c_1 \left(-\frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right)$$

5.8 problem 8

5.8.1 Solving as series ode	2404
5.8.2 Maple step by step solution	2409

Internal problem ID [7301]

Internal file name [OUTPUT/6287_Sunday_June_05_2022_04_37_19_PM_60245184/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 8.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

Unable to solve or complete the solution.

$$y' + y = \frac{1}{x^2}$$

With the expansion point for the power series method at $x = 0$.

5.8.1 Solving as series ode

Writing the ODE as

$$y' + q(x)y = p(x)$$
$$y' + y = \frac{1}{x^2}$$

Where

$$q(x) = 1$$
$$p(x) = \frac{1}{x^2}$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular

singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point if $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not.

Since $x = 0$ is not an ordinary point, we now check to see if it is a regular singular point. Since $x = 0$ is regular singular point, then Frobenius power series is used. Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $y' + y = 0$, and y_p is a particular solution to the inhomogeneous ode. First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} a_n x^{n+r} = \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2) this gives

$$(n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$r a_0 x^{-1+r} = 0$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$m c_0 x^{-1+m} = \frac{1}{x^2}$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation simplifies to

$$r = 0$$

Solving for r gives the root of the indicial equation as

$$r = 0$$

We start by finding y_n . For $1 \leq n$, the recurrence equation is

$$a_n(n+r) + a_{n-1} = 0 \quad (4)$$

For $n = 1$ the recurrence equation gives

$$a_1(1+r) + a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_1 = -\frac{a_0}{1+r}$$

For $n = 2$ the recurrence equation gives

$$a_2(2 + r) + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_2 = \frac{a_0}{(1 + r)(2 + r)}$$

For $n = 3$ the recurrence equation gives

$$a_3(3 + r) + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{(1 + r)(2 + r)(3 + r)}$$

For $n = 4$ the recurrence equation gives

$$a_4(4 + r) + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{(1 + r)(2 + r)(3 + r)(4 + r)}$$

For $n = 5$ the recurrence equation gives

$$a_5(5 + r) + a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_0}{(1 + r)(2 + r)(3 + r)(4 + r)(5 + r)}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^{n+r} \\ &= a_0 x^r + a_1 x^{1+r} + a_2 x^{2+r} + a_3 x^{3+r} + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 x^r - \frac{a_0 x^{1+r}}{1+r} + \frac{a_0 x^{2+r}}{(1+r)(2+r)} - \frac{a_0 x^{3+r}}{(1+r)(2+r)(3+r)} + \frac{a_0 x^{4+r}}{(1+r)(2+r)(3+r)(4+r)} - \frac{a_0 x^{5+r}}{(1+r)(2+r)(3+r)(4+r)(5+r)} + \dots$$

Which can be written as

$$y = x^r \left(a_0 - \frac{a_0 x}{1+r} + \frac{a_0 x^2}{(1+r)(2+r)} - \frac{a_0 x^3}{(1+r)(2+r)(3+r)} + \frac{a_0 x^4}{(1+r)(2+r)(3+r)(4+r)} - \frac{a_0 x^5}{(1+r)(2+r)(3+r)(4+r)(5+r)} + O(x^6) a_0 \right)$$

Collecting terms, the solution becomes

$$y = x^r \left(1 - \frac{x}{1+r} + \frac{x^2}{(1+r)(2+r)} - \frac{x^3}{(1+r)(2+r)(3+r)} + \frac{x^4}{(1+r)(2+r)(3+r)(4+r)} - \frac{x^5}{(1+r)(2+r)(3+r)(4+r)(5+r)} + \dots \right) \quad (3)$$

Finally, since $r = 0$, then the solution becomes

$$y = \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6) \right) a_0 \quad (3)$$

Therefore the homogeneous solution is

$$y_h(x) = \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + O(x^6) \right) a_0$$

Now we determine the particular solution y_p by solving the balance equation

$$m c_0 x^{-1+m} = \frac{1}{x^2}$$

For c_0 and x . This results in

$$c_0 = -1$$

$$m = -1$$

The particular solution is therefore

$$y_p = \sum_{n=0}^{\infty} c_n x^{n+m}$$

$$= \sum_{n=0}^{\infty} c_n x^{n+1}$$

Where in the above $c_0 = -1$. The remaining c_n values are found using the same recurrence relation used to find the homogeneous solution but using c_0 in place of a_0 and using $m = -1$ in place of the root of the indicial equation used to find the homogeneous solution. The following are the values of a_n found in terms of the indicial root r . These will be now used to find c_n by replacing $a_0 = -1$ and $r = -1$. The following table gives the a_n values found and the corresponding c_n values which will be used to find the particular solution

n	a_n	c_n
0	$a_0 = 1$	$c_0 = -1$

Unable to find particular solution .Unable to find the particular solution. No solution exist.

Verification of solutions N/A

5.8.2 Maple step by step solution

Let's solve

$$y' + y = \frac{1}{x^2}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -y + \frac{1}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + y = \frac{1}{x^2}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + y) = \frac{\mu(x)}{x^2}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' + y) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx} (\mu(x) y) \right) dx = \int \frac{\mu(x)}{x^2} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \frac{\mu(x)}{x^2} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)}{x^2} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^x$

$$y = \frac{\int \frac{e^x}{x^2} dx + c_1}{e^x}$$

- Evaluate the integrals on the rhs

$$y = \frac{-\frac{e^x}{x} - \text{Ei}_1(-x) + c_1}{e^x}$$

- Simplify

$$y = \frac{c_1 x e^{-x} - \text{Ei}_1(-x) x e^{-x} - 1}{x}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✗ Solution by Maple

```
Order:=6;
dsolve(diff(y(x),x)+y(x)=1/x^2,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 122

```
AsymptoticDSolveValue[y'[x]+y[x]==1/x^2,y[x],{x,0,5}]
```

$$y(x) \rightarrow \left(-\frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right) \left(\frac{x^6}{2160} + \frac{x^5}{1800} + \frac{x^4}{480} + \frac{x^3}{72} + \frac{x^2}{12} + \frac{x}{2} - \frac{1}{x} + \log(x) \right) \\ + c_1 \left(-\frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right)$$

5.9 problem 9

5.9.1 Solving as series ode	2412
5.9.2 Maple step by step solution	2416

Internal problem ID [7302]

Internal file name [OUTPUT/6288_Sunday_June_05_2022_04_37_21_PM_62041604/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 9.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method.**

Regular singular point"

Maple gives the following as the ode type

`[_separable]`

$$xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

5.9.1 Solving as series ode

Writing the ODE as

$$y' + q(x)y = p(x)$$
$$y' + \frac{y}{x} = 0$$

Where

$$q(x) = \frac{1}{x}$$
$$p(x) = 0$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a

regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not.

Since $x = 0$ is not an ordinary point, we now check to see if it is a regular singular point. $xq(x) = 1$ has a Taylor series around $x = 0$. Since $x = 0$ is regular singular point, then Frobenius power series is used. Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \frac{\sum_{n=0}^{\infty} a_n x^{n+r}}{x} = 0 \quad (1)$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \frac{\sum_{n=0}^{\infty} a_n x^{n+r}}{x} = 0 \quad (1)$$

Expanding the second term in (1) gives

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + x \cdot \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) - 1 \cdot \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2) this gives

$$(n+r) a_n x^{n+r-1} + x^{n+r-1} a_n = 0$$

When $n = 0$ the above becomes

$$r a_0 x^{-1+r} + x^{-1+r} a_0 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r+1) x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation simplifies to

$$r+1 = 0$$

Solving for r gives the root of the indicial equation as

$$r = -1$$

We start by finding y_h . Replacing $r = -1$ found above results in

$$\left(\sum_{n=0}^{\infty} (n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} x^{n-2} a_n \right) = 0$$

From the above we see that there is no recurrence relation since there is only one summation term. Therefore all a_n terms are zero except for a_0 . Hence

$$y_h = a_0 x^r$$

Therefore the homogeneous solution is

$$y_h(x) = a_0 \left(\frac{1}{x} + O(x^6) \right)$$

At $x = 0$ the solution above becomes

$$y = c_1 \left(\frac{1}{x} + O(x^6) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \left(\frac{1}{x} + O(x^6) \right) \tag{1}$$




Figure 137: Slope field plot

Verification of solutions

$$y = c_1 \left(\frac{1}{x} + O(x^6) \right)$$

Verified OK.

5.9.2 Maple step by step solution

Let's solve

$$y' + \frac{y}{x} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = -\frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int -\frac{1}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = -\ln(x) + c_1$$

- Solve for y

$$y = \frac{e^{c_1}}{x}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
Order:=6;  
dsolve(x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1}{x} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 9

```
AsymptoticDSolveValue[x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1}{x}$$

5.10 problem 10

5.10.1 Solving as series ode	2418
5.10.2 Maple step by step solution	2421

Internal problem ID [7303]

Internal file name [OUTPUT/6289_Sunday_June_05_2022_04_37_23_PM_28782687/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 10.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

[_quadrature]

Unable to solve or complete the solution.

$$y' = \frac{1}{x}$$

With the expansion point for the power series method at $x = 0$.

5.10.1 Solving as series ode

Writing the ODE as

$$y' + q(x)y = p(x)$$
$$y' = \frac{1}{x}$$

Where

$$q(x) = 0$$
$$p(x) = \frac{1}{x}$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular

singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point if $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not.

Since $x = 0$ is not an ordinary point, we now check to see if it is a regular singular point. Since $x = 0$ is regular singular point, then Frobenius power series is used. Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $y' = 0$, and y_p is a particular solution to the inhomogeneous ode. First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

Substituting the above back into the ode gives

$$\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} = 0 \tag{1}$$

Which simplifies to

$$\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} = 0 \tag{2A}$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives Substituting all the above in

Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2) this gives

$$(n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$r a_0 x^{-1+r} = 0$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$m c_0 x^{-1+m} = \frac{1}{x}$$

This equation will be used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation simplifies to

$$r = 0$$

Solving for r gives the root of the indicial equation as

$$r = 0$$

We start by finding y_h . From the above we see that there is no recurrence relation since there is only one summation term. Therefore all a_n terms are zero except for a_0 . Hence

$$y_h = a_0 x^r$$

Therefore the homogeneous solution is

$$y_h(x) = a_0 (1 + O(x^6))$$

Unable to solve the balance equation $m c_0 x^{-1+m} = \frac{1}{x}$ for c_0 and x . No particular solution exists.

Unable to find the particular solution. No solution exist.

Verification of solutions N/A

5.10.2 Maple step by step solution

Let's solve

$$y' = \frac{1}{x}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$y = \ln(x) + c_1$$

- Solve for y

$$y = \ln(x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

Solution by Maple

```
Order:=6;  
dsolve(diff(y(x),x)=1/x,y(x),type='series',x=0);
```

No solution found

Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 8

```
AsymptoticDSolveValue[y'[x]==1/x,y[x],{x,0,5}]
```

$$y(x) \rightarrow \log(x) + c_1$$

5.11 problem 11

5.11.1 Maple step by step solution 2423

Internal problem ID [7304]

Internal file name [OUTPUT/6290_Sunday_June_05_2022_04_37_25_PM_27329765/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 11.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

[[_2nd_order , _quadrature]]

Unable to solve or complete the solution.

$$y'' = \frac{1}{x}$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$y'' = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$

$$q(x) = 0$$

Table 255: Table $p(x), q(x)$ singularities.

$p(x) = 0$		$q(x) = 0$	
singularity	type	singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[\infty]$

Irregular singular points : $[\]$

Verification of solutions N/A

5.11.1 Maple step by step solution

Let's solve

$$y'' = \frac{1}{x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{1}{x} \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\left(\int 1 dx\right) + x\left(\int \frac{1}{x} dx\right)$$
 - Compute integrals

$$y_p(x) = x(\ln(x) - 1)$$
 - Substitute particular solution into general solution to ODE

$$y = c_1 + c_2x + x(\ln(x) - 1)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```

✗ Solution by Maple

```

Order:=6;
dsolve(diff(y(x),x$2)=1/x,y(x),type='series',x=0);

```

No solution found

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 17

```
AsymptoticDSolveValue[y''[x]==1/x,y[x],{x,0,5}]
```

$$y(x) \rightarrow -x + x \log(x) + c_2 x + c_1$$

5.12 problem 12

5.12.1 Maple step by step solution 2427

Internal problem ID [7305]

Internal file name [OUTPUT/6291_Sunday_June_05_2022_04_37_26_PM_50109572/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 12.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

Unable to solve or complete the solution.

$$y'' + y' = \frac{1}{x}$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$y'' + y' = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 1$$

$$q(x) = 0$$

Table 257: Table $p(x), q(x)$ singularities.

$p(x) = 1$		$q(x) = 0$	
singularity	type	singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : []

Irregular singular points : [∞]

Verification of solutions N/A

5.12.1 Maple step by step solution

Let's solve

$$y'' + y' = \frac{1}{x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{1}{x} \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int \frac{e^x}{x} dx \right) + \int \frac{1}{x} dx$$
 - Compute integrals

$$y_p(x) = e^{-x} \text{Ei}_1(-x) + \ln(x)$$
 - Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 + e^{-x} \text{Ei}_1(-x) + \ln(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(_b(_a)*_a-1)/_a, _b(_a)`
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

*** Sublev

X Solution by Maple

```
Order:=6;  
dsolve(diff(y(x),x$2)+diff(y(x),x)=1/x,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 159

```
AsymptoticDSolveValue[y''[x]+y'[x]==1/x,y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^6}{4320} - \frac{x^5}{600} - \frac{x^4}{96} - \frac{x^3}{18} - \frac{x^2}{4} + c_2 \left(-\frac{x^5}{720} + \frac{x^4}{120} - \frac{x^3}{24} + \frac{x^2}{6} - \frac{x}{2} + 1 \right) x \\ + \left(-\frac{x^5}{720} + \frac{x^4}{120} - \frac{x^3}{24} + \frac{x^2}{6} - \frac{x}{2} + 1 \right) x \left(\frac{x^6}{2160} + \frac{x^5}{600} + \frac{x^4}{96} + \frac{x^3}{18} + \frac{x^2}{4} + x + \log(x) \right) \\ - x + c_1$$

5.13 problem 13

5.13.1 Maple step by step solution 2431

Internal problem ID [7306]

Internal file name [OUTPUT/6292_Sunday_June_05_2022_04_37_27_PM_5680654/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 13.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_linear_constant_coeff**"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$y'' + y = \frac{1}{x}$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$y'' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$

$$q(x) = 1$$

Table 259: Table $p(x), q(x)$ singularities.

$p(x) = 0$		$q(x) = 1$	
singularity	type	singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : []

Irregular singular points : [∞]

Verification of solutions N/A

5.13.1 Maple step by step solution

Let's solve

$$y'' + y = \frac{1}{x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{1}{x} \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \frac{\sin(x)}{x} dx \right) + \sin(x) \left(\int \frac{\cos(x)}{x} dx \right)$$
 - Compute integrals

$$y_p(x) = -\cos(x) \text{Si}(x) + \sin(x) \text{Ci}(x)$$
 - Substitute particular solution into general solution to ODE


$$y = c_1 \cos(x) + c_2 \sin(x) - \cos(x) \text{Si}(x) + \sin(x) \text{Ci}(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

 Solution by Maple

```
Order:=6;  
dsolve(diff(y(x),x$2)+y(x)=1/x,y(x),type='series',x=0);
```

No solution found

 Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 148

```
AsymptoticDSolveValue[y''[x]+y[x]==1/x,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & x \left(-\frac{x^6}{5040} + \frac{x^4}{120} - \frac{x^2}{6} + 1 \right) \left(-\frac{x^6}{4320} + \frac{x^4}{96} - \frac{x^2}{4} + \log(x) \right) \\ & + c_1 \left(-\frac{x^6}{720} + \frac{x^4}{24} - \frac{x^2}{2} + 1 \right) + c_2 x \left(-\frac{x^6}{5040} + \frac{x^4}{120} - \frac{x^2}{6} + 1 \right) \\ & + \left(-\frac{x^5}{600} + \frac{x^3}{18} - x \right) \left(-\frac{x^6}{720} + \frac{x^4}{24} - \frac{x^2}{2} + 1 \right) \end{aligned}$$

5.14 problem 14

5.14.1 Maple step by step solution 2435

Internal problem ID [7307]

Internal file name [OUTPUT/6293_Sunday_June_05_2022_04_37_28_PM_25391719/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 14.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_linear_constant_coeff**"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

Unable to solve or complete the solution.

$$y'' + y' + y = \frac{1}{x}$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$y'' + y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 1$$

$$q(x) = 1$$

Table 261: Table $p(x), q(x)$ singularities.

$p(x) = 1$		$q(x) = 1$	
singularity	type	singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : []

Irregular singular points : [∞]

Verification of solutions N/A

5.14.1 Maple step by step solution

Let's solve

$$y'' + y' + y = \frac{1}{x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE
- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{1}{x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \sqrt{3} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3} e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3} e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int \frac{e^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{x} dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int \frac{e^{\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{x} dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = -\frac{\left(\left(\text{Icos}\left(\frac{\sqrt{3}x}{2}\right) + \text{sin}\left(\frac{\sqrt{3}x}{2}\right) \right) \text{Ei}_1\left(-\frac{x(1+I\sqrt{3})}{2}\right) - \left(\text{Icos}\left(\frac{\sqrt{3}x}{2}\right) - \text{sin}\left(\frac{\sqrt{3}x}{2}\right) \right) \text{Ei}_1\left(\frac{x(I\sqrt{3}-1)}{2}\right) \right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

- Substitute particular solution into general solution to ODE


$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} - \frac{\left(\left(\text{Icos}\left(\frac{\sqrt{3}x}{2}\right) + \text{sin}\left(\frac{\sqrt{3}x}{2}\right) \right) \text{Ei}_1\left(-\frac{x(1+I\sqrt{3})}{2}\right) - \left(\text{Icos}\left(\frac{\sqrt{3}x}{2}\right) - \text{sin}\left(\frac{\sqrt{3}x}{2}\right) \right) \text{Ei}_1\left(\frac{x(I\sqrt{3}-1)}{2}\right) \right) e^{-\frac{x}{2}} \sqrt{3}}{3}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

 Solution by Maple

```
Order:=6;  
dsolve(diff(y(x),x$2)+diff(y(x),x)+y(x)=1/x,y(x),type='series',x=0);
```

No solution found

 Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 152

```
AsymptoticDSolveValue[y''[x]+y'[x]+y[x]==1/x,y[x],{x,0,5}]
```

$$\begin{aligned} y(x) \rightarrow & c_2 x \left(-\frac{x^4}{120} + \frac{x^3}{24} - \frac{x}{2} + 1 \right) + c_1 \left(\frac{x^3}{6} - \frac{x^2}{2} + 1 \right) \\ & + x \left(-\frac{x^4}{120} + \frac{x^3}{24} - \frac{x}{2} + 1 \right) \left(\frac{41x^6}{4320} + \frac{x^5}{120} - \frac{x^4}{96} - \frac{x^3}{18} + x + \log(x) \right) \\ & + \left(\frac{x^3}{6} - \frac{x^2}{2} + 1 \right) \left(-\frac{x^6}{180} + \frac{x^5}{600} + \frac{x^4}{96} - \frac{x^2}{4} - x \right) \end{aligned}$$

5.15 problem 15

5.15.1 Maple step by step solution 2439

Internal problem ID [7308]

Internal file name [OUTPUT/6294_Sunday_June_05_2022_04_37_30_PM_76305912/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 15.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$h^2 + \frac{2ah}{\sqrt{1+h'^2}} = b^2$$

Solving the given ode for h' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$h' = -\frac{\sqrt{-h^4 + 4a^2h^2 + 2h^2b^2 - b^4}}{(h+b)(h-b)} \quad (1)$$

$$h' = \frac{\sqrt{-h^4 + 4a^2h^2 + 2h^2b^2 - b^4}}{(h+b)(h-b)} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int -\frac{(h+b)(h-b)}{\sqrt{4a^2h^2 - b^4 + 2h^2b^2 - h^4}} dh = \int du$$
$$\int -\frac{(_a + b)(_a - b)}{\sqrt{-_a^4 + 4_a^2a^2 + 2_a^2b^2 - b^4}} d_a = u + c_1$$

Summary

The solution(s) found are the following

$$\int^h -\frac{(a+b)(a-b)}{\sqrt{-a^4+4a^2a^2+2a^2b^2-b^4}} da = u + c_1 \quad (1)$$

Verification of solutions

$$\int^h -\frac{(a+b)(a-b)}{\sqrt{-a^4+4a^2a^2+2a^2b^2-b^4}} da = u + c_1$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\int \frac{(h+b)(h-b)}{\sqrt{4a^2h^2-b^4+2h^2b^2-h^4}} dh = \int du$$
$$\int^h \frac{(a+b)(a-b)}{\sqrt{-a^4+4a^2a^2+2a^2b^2-b^4}} da = u + c_2$$

Summary

The solution(s) found are the following

$$\int^h \frac{(a+b)(a-b)}{\sqrt{-a^4+4a^2a^2+2a^2b^2-b^4}} da = u + c_2 \quad (1)$$

Verification of solutions

$$\int^h \frac{(a+b)(a-b)}{\sqrt{-a^4+4a^2a^2+2a^2b^2-b^4}} da = u + c_2$$

Verified OK.

5.15.1 Maple step by step solution

Let's solve

$$h^2 + \frac{2ah}{\sqrt{1+h^2}} = b^2$$

- Highest derivative means the order of the ODE is 1

h'

- Separate variables

$$\frac{(h+b)(h-b)h'}{\sqrt{-h^4+4a^2h^2+2h^2b^2-b^4}} = 1$$

- Integrate both sides with respect to u

$$\int \frac{(h+b)(h-b)h'}{\sqrt{-h^4+4a^2h^2+2h^2b^2-b^4}} du = \int 1 du + c_1$$

- Evaluate integral

$$2b^4 \sqrt{1 + \frac{h^2(2a\sqrt{a^2+b^2}-2a^2-b^2)}{b^4}} \sqrt{1 - \frac{(2a\sqrt{a^2+b^2}+2a^2+b^2)h^2}{b^4}} \left(\text{EllipticF} \left(h \sqrt{-\frac{2a\sqrt{a^2+b^2}-2a^2-b^2}{b^4}}, \sqrt{-1 + \frac{(4a^2+2b^2)(2a\sqrt{a^2+b^2}+2a^2+b^2)}{b^4}} \right) \right) \sqrt{-\frac{2a\sqrt{a^2+b^2}-2a^2-b^2}{b^4}} \sqrt{-h^4+4a^2h^2+2h^2b^2-b^4} (4a^2+2b^2)$$

Maple trace

```

`Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
<- differential order: 1; missing x successful`

```

✓ Solution by Maple

Time used: 0.438 (sec). Leaf size: 103

```
dsolve(h(u)^2 + 2*a*h(u)/sqrt(1 + diff(h(u), u)^2) = b^2,h(u), singsol=all)
```

$$u - \left(\int^{h(u)} \frac{-a^2 - b^2}{\sqrt{-a^4 + (4a^2 + 2b^2)a^2 - b^4}} da \right) - c_1 = 0$$

$$u + \int^{h(u)} \frac{-a^2 - b^2}{\sqrt{-a^4 + (4a^2 + 2b^2)a^2 - b^4}} da - c_1 = 0$$

✓ Solution by Mathematica

Time used: 24.41 (sec). Leaf size: 913

```
DSolve[h[u]^2 + 2*a*h[u]/Sqrt[1 + (h'[u])^2] == b^2,h[u],u,IncludeSingularSolutions -> True]
```

$$h(u) \rightarrow \text{InverseFunction} \left[\frac{i\sqrt{(b^2 - \#1^2)^2} \sqrt{1 - \frac{\#1^2}{-2\sqrt{a^2(a^2+b^2)+2a^2+b^2}}} \sqrt{1 - \frac{\#1^2}{2\sqrt{a^2(a^2+b^2)+2a^2+b^2}}} \left((2\sqrt{a^2(a^2+b^2)} + c_1 \right)}{\dots} \right]$$

$$h(u) \rightarrow \text{InverseFunction} \left[\frac{i\sqrt{(b^2 - \#1^2)^2} \sqrt{1 - \frac{\#1^2}{-2\sqrt{a^2(a^2+b^2)+2a^2+b^2}}} \sqrt{1 - \frac{\#1^2}{2\sqrt{a^2(a^2+b^2)+2a^2+b^2}}} \left((2\sqrt{a^2(a^2+b^2)} + c_1 \right)}{\dots} \right]$$

$$h(u) \rightarrow -\sqrt{a^2 + b^2} - a$$

$$h(u) \rightarrow \sqrt{a^2 + b^2} - a$$

5.16 problem 16

5.16.1 Solving as second order linear constant coeff ode	2442
5.16.2 Solving using Kovacic algorithm	2445
5.16.3 Maple step by step solution	2452

Internal problem ID [7309]

Internal file name [OUTPUT/6295_Sunday_June_05_2022_04_39_02_PM_53865576/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 16.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 2y' - 24y = 16 - (x + 2)e^{4x}$$

5.16.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 2, C = -24, f(x) = 16 + (-x - 2)e^{4x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' - 24y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = -24$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} - 24 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda - 24 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = -24$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{2^2 - (4)(1)(-24)} \\ &= -1 \pm 5 \end{aligned}$$

Hence

$$\lambda_1 = -1 + 5$$

$$\lambda_2 = -1 - 5$$

Which simplifies to

$$\lambda_1 = 4$$

$$\lambda_2 = -6$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(4)x} + c_2 e^{(-6)x}$$

Or

$$y = c_1 e^{4x} + c_2 e^{-6x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{4x} + c_2 e^{-6x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$16 + (-x - 2)e^{4x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{e^{4x}x, e^{4x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-6x}, e^{4x}\}$$

Since e^{4x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{1\}, \{x^2e^{4x}, e^{4x}x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 + A_2x^2e^{4x} + A_3e^{4x}x$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_2e^{4x} + 20A_2xe^{4x} + 10A_3e^{4x} - 24A_1 = 16 + (-x - 2)e^{4x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{2}{3}, A_2 = -\frac{1}{20}, A_3 = -\frac{19}{100} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{2}{3} - \frac{x^2e^{4x}}{20} - \frac{19e^{4x}x}{100}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^{4x} + c_2e^{-6x}) + \left(-\frac{2}{3} - \frac{x^2e^{4x}}{20} - \frac{19e^{4x}x}{100} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{4x} + c_2 e^{-6x} - \frac{2}{3} - \frac{x^2 e^{4x}}{20} - \frac{19 e^{4x} x}{100} \quad (1)$$




Figure 138: Slope field plot

Verification of solutions

$$y = c_1 e^{4x} + c_2 e^{-6x} - \frac{2}{3} - \frac{x^2 e^{4x}}{20} - \frac{19 e^{4x} x}{100}$$

Verified OK.

5.16.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' - 24y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 2 \\C &= -24\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{25}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 25 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 25z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 264: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 25$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-5x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \\
 &= z_1 e^{-x} \\
 &= z_1 (e^{-x})
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-6x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{10x}}{10} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-6x}) + c_2 \left(e^{-6x} \left(\frac{e^{10x}}{10} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' - 24y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-6x} + \frac{c_2 e^{4x}}{10}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-6x}$$

$$y_2 = \frac{e^{4x}}{10}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-6x} & \frac{e^{4x}}{10} \\ \frac{d}{dx}(e^{-6x}) & \frac{d}{dx}\left(\frac{e^{4x}}{10}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-6x} & \frac{e^{4x}}{10} \\ -6e^{-6x} & \frac{2e^{4x}}{5} \end{vmatrix}$$

Therefore

$$W = (e^{-6x}) \left(\frac{2e^{4x}}{5} \right) - \left(\frac{e^{4x}}{10} \right) (-6e^{-6x})$$

Which simplifies to

$$W = e^{-6x} e^{4x}$$

Which simplifies to

$$W = e^{-2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{e^{4x}(16+(-x-2)e^{4x})}{10}}{e^{-2x}} dx$$

Which simplifies to

$$u_1 = - \int - \frac{e^{6x}(-16 + (x + 2) e^{4x})}{10} dx$$

Hence

$$u_1 = \frac{e^{10x}x}{100} + \frac{19e^{10x}}{1000} - \frac{4e^{6x}}{15}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-6x}(16 + (-x - 2) e^{4x})}{e^{-2x}} dx$$

Which simplifies to

$$u_2 = \int (16e^{-4x} - x - 2) dx$$

Hence

$$u_2 = -2x - \frac{x^2}{2} - 4e^{-4x}$$

Which simplifies to

$$u_1 = \frac{(30x + 57)e^{10x}}{3000} - \frac{4e^{6x}}{15}$$

$$u_2 = -2x - \frac{x^2}{2} - 4e^{-4x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{(30x + 57)e^{10x}}{3000} - \frac{4e^{6x}}{15} \right) e^{-6x} + \frac{\left(-2x - \frac{x^2}{2} - 4e^{-4x} \right) e^{4x}}{10}$$

Which simplifies to

$$y_p(x) = -\frac{2}{3} + \frac{(-50x^2 - 190x + 19)e^{4x}}{1000}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-6x} + \frac{c_2 e^{4x}}{10} \right) + \left(-\frac{2}{3} + \frac{(-50x^2 - 190x + 19)e^{4x}}{1000} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-6x} + \frac{c_2 e^{4x}}{10} - \frac{2}{3} + \frac{(-50x^2 - 190x + 19)e^{4x}}{1000} \quad (1)$$




Figure 139: Slope field plot

Verification of solutions

$$y = c_1 e^{-6x} + \frac{c_2 e^{4x}}{10} - \frac{2}{3} + \frac{(-50x^2 - 190x + 19)e^{4x}}{1000}$$

Verified OK.

5.16.3 Maple step by step solution

Let's solve

$$y'' + 2y' - 24y = 16 + (-x - 2)e^{4x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -e^{4x}x - 2e^{4x} + 24y - 2y' + 16$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + 2y' - 24y = 16 - e^{4x}x - 2e^{4x}$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 2r - 24 = 0$$

- Factor the characteristic polynomial

$$(r + 6)(r - 4) = 0$$

- Roots of the characteristic polynomial

$$r = (-6, 4)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-6x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{4x}$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1e^{-6x} + c_2e^{4x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 16 - e^{4x}x - 2e^{4x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-6x} & e^{4x} \\ -6e^{-6x} & 4e^{4x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 10 e^{-2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{(e^{10x} (\int (16 e^{-4x} - x - 2) dx) + \int e^{6x} (-16 + (x+2)e^{4x}) dx) e^{-6x}}{10}$$

- Compute integrals

$$y_p(x) = -\frac{2}{3} + \frac{(-50x^2 - 190x + 19)e^{4x}}{1000}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-6x} + c_2 e^{4x} - \frac{2}{3} + \frac{(-50x^2 - 190x + 19)e^{4x}}{1000}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)-24*y(x)=16-(x+2)*exp(4*x),y(x), singsol=all)
```

$$y(x) = -\frac{\left(\left(x^2 + \frac{19}{5}x - 20c_2 - \frac{19}{50} \right) e^{10x} - 20c_1 + \frac{40e^{6x}}{3} \right) e^{-6x}}{20}$$

✓ Solution by Mathematica

Time used: 0.186 (sec). Leaf size: 41

```
DSolve[y''[x]+2*y'[x]-24*y[x]==16-(x+2)*Exp[4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{4x} \left(-\frac{x^2}{20} - \frac{19x}{100} + \frac{19}{1000} + c_2 \right) + c_1 e^{-6x} - \frac{2}{3}$$

5.17 problem 17

- 5.17.1 Existence and uniqueness analysis 2455
- 5.17.2 Maple step by step solution 2458

Internal problem ID [7310]

Internal file name [OUTPUT/6296_Sunday_June_05_2022_04_39_04_PM_47141029/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 17.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 3y' - 4y = 6e^{2t-2}$$

With initial conditions

$$[y(1) = 4, y'(1) = 5]$$

5.17.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = 3$$

$$q(t) = -4$$

$$F = 6e^{2t-2}$$

Hence the ode is

$$y'' + 3y' - 4y = 6e^{2t-2}$$

The domain of $p(t) = 3$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 1$ is inside this domain. The domain of $q(t) = -4$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 1$ is also inside this domain. The domain of $F = 6e^{2t-2}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 1$ is also inside this domain. Hence solution exists and is unique.

Since both initial conditions are not at zero, then let

$$\begin{aligned}y(0) &= c_1 \\y'(0) &= c_2\end{aligned}$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 3sY(s) - 3y(0) - 4Y(s) = \frac{6e^{-2}}{s-2} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= c_1 \\y'(0) &= c_2\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - c_2 - sc_1 + 3sY(s) - 3c_1 - 4Y(s) = \frac{6e^{-2}}{s-2}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{s^2 c_1 + s c_1 + c_2 s + 6 e^{-2} - 6 c_1 - 2 c_2}{(s - 2)(s^2 + 3s - 4)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{e^{-2}}{s - 2} + \frac{\frac{c_1}{5} - \frac{c_2}{5} + \frac{e^{-2}}{5}}{s + 4} + \frac{\frac{4c_1}{5} + \frac{c_2}{5} - \frac{6e^{-2}}{5}}{s - 1}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(\frac{e^{-2}}{s - 2}\right) &= e^{2t-2} \\ \mathcal{L}^{-1}\left(\frac{\frac{c_1}{5} - \frac{c_2}{5} + \frac{e^{-2}}{5}}{s + 4}\right) &= \frac{(c_1 - c_2 + e^{-2}) e^{-4t}}{5} \\ \mathcal{L}^{-1}\left(\frac{\frac{4c_1}{5} + \frac{c_2}{5} - \frac{6e^{-2}}{5}}{s - 1}\right) &= \frac{e^t(4c_1 + c_2 - 6e^{-2})}{5}\end{aligned}$$

Adding the above results and simplifying gives

$$y = e^{2t-2} + \frac{e^t(4c_1 + c_2 - 6e^{-2})}{5} + \frac{(c_1 - c_2 + e^{-2}) e^{-4t}}{5}$$

Since both initial conditions given are not at zero, then we need to setup two equations to solve for c_1, c_1 . At $t = 1$ the first equation becomes, using the above solution

$$4 = 1 + \frac{e(4c_1 + c_2 - 6e^{-2})}{5} + \frac{(c_1 - c_2 + e^{-2}) e^{-4}}{5}$$

And taking derivative of the solution and evaluating at $t = 1$ gives the second equation as

$$5 = 2 + \frac{e(4c_1 + c_2 - 6e^{-2})}{5} - \frac{4(c_1 - c_2 + e^{-2}) e^{-4}}{5}$$

Solving gives

$$\begin{aligned}c_1 &= (e e^{-2} + 3) e^{-1} \\ c_2 &= e^{-1}(2 e e^{-2} + 3)\end{aligned}$$

Substituting these in the solution obtained above gives

$$\begin{aligned}y &= e^{2t-2} + \frac{e^t(4(e e^{-2} + 3) e^{-1} + e^{-1}(2 e e^{-2} + 3) - 6 e^{-2})}{5} + \frac{((e e^{-2} + 3) e^{-1} - e^{-1}(2 e e^{-2} + 3) + e^{-2}) e^{-4t}}{5} \\ &= e^{2t-2} + 3 e^{t-1}\end{aligned}$$


Simplifying the solution gives

$$y = e^{2t-2} + 3e^{t-1}$$


Summary

The solution(s) found are the following

$$y = e^{2t-2} + 3e^{t-1} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{2t-2} + 3e^{t-1}$$

Verified OK.

5.17.2 Maple step by step solution

Let's solve

$$\left[y'' + 3y' - 4y = 6e^{2t-2}, y(1) = 4, y'|_{\{t=1\}} = 5 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 3r - 4 = 0$$

- Factor the characteristic polynomial
 $(r + 4)(r - 1) = 0$
- Roots of the characteristic polynomial
 $r = (-4, 1)$
- 1st solution of the homogeneous ODE
 $y_1(t) = e^{-4t}$
- 2nd solution of the homogeneous ODE
 $y_2(t) = e^t$
- General solution of the ODE
 $y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$
- Substitute in solutions of the homogeneous ODE
 $y = c_1 e^{-4t} + c_2 e^t + y_p(t)$
- Find a particular solution $y_p(t)$ of the ODE
 - Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t),y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t),y_2(t))} dt \right), f(t) = 6 e^{2t-2} \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{-4t} & e^t \\ -4e^{-4t} & e^t \end{bmatrix}$$
 - Compute Wronskian
 $W(y_1(t), y_2(t)) = 5 e^{-3t}$
 - Substitute functions into equation for $y_p(t)$

$$y_p(t) = -\frac{6(-e^{5t}(\int e^{t-2} dt) + \int e^{6t-2} dt)e^{-4t}}{5}$$
 - Compute integrals
 $y_p(t) = e^{2t-2}$
- Substitute particular solution into general solution to ODE
 $y = c_1 e^{-4t} + c_2 e^t + e^{2t-2}$
- Check validity of solution $y = c_1 e^{-4t} + c_2 e^t + e^{2t-2}$
 - Use initial condition $y(1) = 4$
 $4 = c_1 e^{-4} + c_2 e + 1$

- Compute derivative of the solution

$$y' = -4c_1e^{-4t} + c_2e^t + 2e^{2t-2}$$

- Use the initial condition $y'|_{\{t=1\}} = 5$

$$5 = -4c_1e^{-4} + c_2e + 2$$

- Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = \frac{3}{e}\}$$

- Substitute constant values into general solution and simplify

$$y = e^{2t-2} + 3e^{t-1}$$

- Solution to the IVP

$$y = e^{2t-2} + 3e^{t-1}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.297 (sec). Leaf size: 17

```
dsolve([diff(y(t),t$2)+3*diff(y(t),t)-4*y(t)=6*exp(2*t-2),y(1) = 4, D(y)(1) = 5],y(t), sings
```

$$y(t) = e^{2t-2} + 3e^{t-1}$$

✓ Solution by Mathematica

Time used: 0.078 (sec). Leaf size: 18

```
DSolve[{y''[t]+3*y'[t]-4*y[t]==6*Exp[2*t-2],{y[1]==4,y'[1]==5}},y[t],t,IncludeSingularSoluti
```

$$y(t) \rightarrow e^{t-2}(e^t + 3e)$$

5.18 problem 18

5.18.1 Maple step by step solution 2472

Internal problem ID [7311]

Internal file name [OUTPUT/6297_Sunday_June_05_2022_04_39_06_PM_23353552/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 18.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_linear_constant_coeff**", "**second order series method. Ordinary point**", "**second order series method. Taylor series method**"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = e^{a \cos(x)}$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \cdots \\ &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (388)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (389)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -y + e^{a \cos(x)}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= -y' - a \sin(x) e^{a \cos(x)} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= e^{a \cos(x)} (a^2 \sin(x)^2 - a \cos(x) - 1) + y \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= -a \sin(x) (a^2 \sin(x)^2 - 3a \cos(x) - 2) e^{a \cos(x)} + y' \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= (8a^2 \cos(x)^2 + (-6a^3 \sin(x)^2 + 2a) \cos(x) + \sin(x)^4 a^4 - 5a^2 + 1) e^{a \cos(x)} - y \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= -a \sin(x) (\sin(x)^4 a^4 - 10 \cos(x) \sin(x)^2 a^3 + 26a^2 \cos(x)^2 + 18a \cos(x) - 11a^2 + 3) e^{a \cos(x)} - y' \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= (-96a^3 \cos(x)^3 + (66 \sin(x)^2 a^4 - 39a^2) \cos(x)^2 + (-15a^5 \sin(x)^4 + 81a^3 - 3a) \cos(x) + \sin(x)^6 a^4) e^{a \cos(x)} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= -y(0) + e^a \\
 F_1 &= -y'(0) \\
 F_2 &= -e^a a - e^a + y(0) \\
 F_3 &= y'(0) \\
 F_4 &= 3e^a a^2 + 2e^a a + e^a - y(0) \\
 F_5 &= -y'(0) \\
 F_6 &= -15e^a a^3 - 18e^a a^2 - 3e^a a - e^a + y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned}
 y &= \left(\frac{1}{40320}x^8 - \frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 \right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 \right) y'(0) \\
 &+ \frac{x^2 e^a}{2} - \frac{x^4 e^a a}{24} - \frac{x^4 e^a}{24} + \frac{x^6 e^a a^2}{240} + \frac{x^6 e^a a}{360} + \frac{x^6 e^a}{720} - \frac{x^8 e^a a^3}{2688} - \frac{x^8 e^a a^2}{2240} - \frac{x^8 e^a a}{13440} - \frac{x^8 e^a}{40320} \\
 &+ O(x^8)
 \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned}
 y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\
 y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}
 \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = - \left(\sum_{n=0}^{\infty} a_n x^n \right) + e^{a \cos(x)} \quad (1)$$

Expanding $e^{a \cos(x)}$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned}
 e^{a \cos(x)} &= e^a - \frac{e^a a x^2}{2} + e^a \left(\frac{1}{24} a + \frac{1}{8} a^2 \right) x^4 + e^a \left(-\frac{1}{720} a - \frac{1}{48} a^2 - \frac{1}{48} a^3 \right) x^6 + \dots \\
 &= e^a - \frac{e^a a x^2}{2} + e^a \left(\frac{1}{24} a + \frac{1}{8} a^2 \right) x^4 + e^a \left(-\frac{1}{720} a - \frac{1}{48} a^2 - \frac{1}{48} a^3 \right) x^6
 \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\begin{aligned} \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) &= e^a - \frac{e^a a x^2}{2} + e^a \left(\frac{1}{24} a + \frac{1}{8} a^2 \right) x^4 \\ &+ e^a \left(-\frac{1}{720} a - \frac{1}{48} a^2 - \frac{1}{48} a^3 \right) x^6 \end{aligned}$$

Which simplifies to

$$\begin{aligned} \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) &= e^a - \frac{e^a a x^2}{2} + e^a \left(\frac{1}{24} a + \frac{1}{8} a^2 \right) x^4 \quad (2) \\ &+ e^a \left(-\frac{1}{720} a - \frac{1}{48} a^2 - \frac{1}{48} a^3 \right) x^6 \end{aligned}$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) & \quad (3) \\ = e^a - \frac{e^a a x^2}{2} + e^a \left(\frac{1}{24} a + \frac{1}{8} a^2 \right) x^4 + e^a \left(-\frac{1}{720} a - \frac{1}{48} a^2 - \frac{1}{48} a^3 \right) x^6 \end{aligned}$$

For $0 \leq n$, the recurrence equation is

$$\begin{aligned} ((n+2) a_{n+2} (n+1) + a_n) x^n &= e^a - \frac{e^a a x^2}{2} + e^a \left(\frac{1}{24} a + \frac{1}{8} a^2 \right) x^4 \quad (4) \\ &+ e^a \left(-\frac{1}{720} a - \frac{1}{48} a^2 - \frac{1}{48} a^3 \right) x^6 \end{aligned}$$

For $n = 0$ the recurrence equation gives

$$(2a_2 + a_0) 1 = e^a$$
$$2a_2 + a_0 = e^a$$

Which after substituting the earlier terms found becomes

$$a_2 = \frac{e^a}{2} - \frac{a_0}{2}$$

For $n = 1$ the recurrence equation gives

$$(6a_3 + a_1) x = 0$$
$$6a_3 + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_1}{6}$$

For $n = 2$ the recurrence equation gives

$$(12a_4 + a_2) x^2 = -\frac{e^a a x^2}{2}$$
$$12a_4 + a_2 = -\frac{e^a a}{2}$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{e^a a}{24} - \frac{e^a}{24} + \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$(20a_5 + a_3) x^3 = 0$$
$$20a_5 + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{120}$$

For $n = 4$ the recurrence equation gives

$$(30a_6 + a_4)x^4 = e^a \left(\frac{1}{24}a + \frac{1}{8}a^2 \right) x^4$$
$$30a_6 + a_4 = e^a \left(\frac{1}{24}a + \frac{1}{8}a^2 \right)$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{e^a a}{360} + \frac{e^a a^2}{240} + \frac{e^a}{720} - \frac{a_0}{720}$$

For $n = 5$ the recurrence equation gives

$$(42a_7 + a_5)x^5 = 0$$
$$42a_7 + a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_1}{5040}$$

For $n = 6$ the recurrence equation gives

$$(56a_8 + a_6)x^6 = e^a \left(-\frac{1}{720}a - \frac{1}{48}a^2 - \frac{1}{48}a^3 \right) x^6$$
$$56a_8 + a_6 = e^a \left(-\frac{1}{720}a - \frac{1}{48}a^2 - \frac{1}{48}a^3 \right)$$

Which after substituting the earlier terms found becomes

$$a_8 = -\frac{e^a a}{13440} - \frac{e^a a^2}{2240} - \frac{e^a a^3}{2688} - \frac{e^a}{40320} + \frac{a_0}{40320}$$

For $n = 7$ the recurrence equation gives

$$(72a_9 + a_7)x^7 = 0$$
$$72a_9 + a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{a_1}{362880}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x + \left(\frac{e^a}{2} - \frac{a_0}{2} \right) x^2 - \frac{a_1 x^3}{6} + \left(-\frac{e^a a}{24} - \frac{e^a}{24} + \frac{a_0}{24} \right) x^4 \\ &\quad + \frac{a_1 x^5}{120} + \left(\frac{e^a a}{360} + \frac{e^a a^2}{240} + \frac{e^a}{720} - \frac{a_0}{720} \right) x^6 - \frac{a_1 x^7}{5040} + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(-\frac{1}{720} x^6 + \frac{1}{24} x^4 + 1 - \frac{1}{2} x^2 \right) a_0 + \left(x - \frac{1}{6} x^3 + \frac{1}{120} x^5 - \frac{1}{5040} x^7 \right) a_1 \quad (3) \\ &\quad + \frac{x^2 e^a}{2} + \left(-\frac{e^a a}{24} - \frac{e^a}{24} \right) x^4 + \left(\frac{e^a a}{360} + \frac{e^a a^2}{240} + \frac{e^a}{720} \right) x^6 + O(x^8) \end{aligned}$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y &= \left(-\frac{1}{720} x^6 + \frac{1}{24} x^4 + 1 - \frac{1}{2} x^2 \right) c_1 + \left(x - \frac{1}{6} x^3 + \frac{1}{120} x^5 - \frac{1}{5040} x^7 \right) c_2 \\ &\quad + \frac{x^2 e^a}{2} + \left(-\frac{e^a a}{24} - \frac{e^a}{24} \right) x^4 + \left(\frac{e^a a}{360} + \frac{e^a a^2}{240} + \frac{e^a}{720} \right) x^6 + O(x^8) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{1}{40320}x^8 - \frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 \right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 \right) y'(0) + \frac{x^2 e^a}{2} - \frac{x^4 e^a a}{24} - \frac{x^4 e^a}{24} + \frac{x^6 e^a a^2}{240} + \frac{x^6 e^a a}{360} + \frac{x^6 e^a}{720} - \frac{x^8 e^a a^3}{2688} - \frac{x^8 e^a a^2}{2240} - \frac{x^8 e^a a}{13440} - \frac{x^8 e^a}{40320} + O(x^8) \quad (1)$$

$$y = \left(-\frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 \right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 \right) c_2 + \frac{x^2 e^a}{2} + \left(-\frac{e^a a}{24} - \frac{e^a}{24} \right) x^4 + \left(\frac{e^a a}{360} + \frac{e^a a^2}{240} + \frac{e^a}{720} \right) x^6 + O(x^8) \quad (2)$$




Figure 141: Slope field plot

Verification of solutions

$$y = \left(\frac{1}{40320}x^8 - \frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 \right) y(0) \\ + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 \right) y'(0) + \frac{x^2 e^a}{2} - \frac{x^4 e^a a}{24} - \frac{x^4 e^a}{24} + \frac{x^6 e^a a^2}{240} \\ + \frac{x^6 e^a a}{360} + \frac{x^6 e^a}{720} - \frac{x^8 e^a a^3}{2688} - \frac{x^8 e^a a^2}{2240} - \frac{x^8 e^a a}{13440} - \frac{x^8 e^a}{40320} + O(x^8)$$

Verified OK.

$$y = \left(-\frac{1}{720}x^6 + \frac{1}{24}x^4 + 1 - \frac{1}{2}x^2 \right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 \right) c_2 \\ + \frac{x^2 e^a}{2} + \left(-\frac{e^a a}{24} - \frac{e^a}{24} \right) x^4 + \left(\frac{e^a a}{360} + \frac{e^a a^2}{240} + \frac{e^a}{720} \right) x^6 + O(x^8)$$

Verified OK.

5.18.1 Maple step by step solution

Let's solve

$$y'' = -y + e^{a \cos(x)}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + y = e^{a \cos(x)}$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{a \cos(x)} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) e^{a \cos(x)} dx \right) + \sin(x) \left(\int \cos(x) e^{a \cos(x)} dx \right)$$

- Compute integrals

$$y_p(x) = \frac{\sin(x) \left(\int \cos(x) e^{a \cos(x)} dx \right) a + \cos(x) e^{a \cos(x)}}{a}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\sin(x) \left(\int \cos(x) e^{a \cos(x)} dx \right) a + \cos(x) e^{a \cos(x)}}{a}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 76

```
Order:=8;
dsolve(diff(y(x),x$2)+y(x)=exp(a*cos(x)),y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7\right) D(y)(0) \\ + \frac{e^a x^2}{2} + \frac{(-a-1)e^a x^4}{24} + \frac{(3a^2+2a+1)e^a x^6}{720} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 239

```
AsymptoticDSolveValue[y''[x]+y[x]==Exp[a*Cos[x]],y[x],{x,0,7}]
```

$$y(x) \rightarrow \left(-\frac{x^7}{5040} + \frac{x^5}{120} - \frac{x^3}{6} + x\right) \left(\frac{1}{120}(3a^2+7a+1)e^a x^5 \right. \\ \left. - \frac{(15a^3+60a^2+31a+1)e^a x^7}{5040} - \frac{1}{6}(a+1)e^a x^3 + e^a x\right) \\ + \left(-\frac{x^6}{720} + \frac{x^4}{24} - \frac{x^2}{2} + 1\right) \left(-\frac{1}{720}(15a^2+15a+1)e^a x^6 \right. \\ \left. + \frac{(105a^3+210a^2+63a+1)e^a x^8}{40320} + \frac{1}{24}(3a+1)e^a x^4 - \frac{e^a x^2}{2}\right) \\ + c_2 \left(-\frac{x^7}{5040} + \frac{x^5}{120} - \frac{x^3}{6} + x\right) + c_1 \left(-\frac{x^6}{720} + \frac{x^4}{24} - \frac{x^2}{2} + 1\right)$$

5.19 problem 19

- 5.19.1 Solving as differentialType ode 2475
- 5.19.2 Solving as first order ode lie symmetry calculated ode 2477
- 5.19.3 Solving as exact ode 2482

Internal problem ID [7312]

Internal file name [OUTPUT/6311_Monday_July_25_2022_10_21_09_PM_9550685/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 19.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "differentialType", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries]]
```

$$y' - \frac{y}{2y \ln(y) + y - x} = 0$$

5.19.1 Solving as differentialType ode

Writing the ode as

$$y' = \frac{y}{2y \ln(y) + y - x} \tag{1}$$

Which becomes

$$(2y \ln(y) + y) dy = (x) dy + (y) dx \tag{2}$$

But the RHS is complete differential because

$$(x) dy + (y) dx = d(xy)$$

Hence (2) becomes

$$(2y \ln(y) + y) dy = d(xy)$$

Integrating both sides gives gives the solution as

$$\ln(y) y^2 = yx + c_1$$

Summary

The solution(s) found are the following

$$\ln(y) y^2 = yx + c_1 \tag{1}$$




Figure 142: Slope field plot

Verification of solutions

$$\ln(y) y^2 = yx + c_1$$

Verified OK.

5.19.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y}{2y \ln(y) + y - x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{y(b_3 - a_2)}{2y \ln(y) + y - x} - \frac{y^2 a_3}{(2y \ln(y) + y - x)^2} - \frac{y(xa_2 + ya_3 + a_1)}{(2y \ln(y) + y - x)^2} \quad (\text{5E})$$

$$- \left(\frac{1}{2y \ln(y) + y - x} - \frac{y(2 \ln(y) + 3)}{(2y \ln(y) + y - x)^2} \right) (xb_2 + yb_3 + b_1) = 0$$

Putting the above in normal form gives

$$\frac{4 \ln(y)^2 y^2 b_2 - 4 \ln(y) xyb_2 - 2 \ln(y) y^2 a_2 + 4 \ln(y) y^2 b_2 + 2 \ln(y) y^2 b_3 + 2x^2 b_2 - y^2 a_2 - 2y^2 a_3 + y^2 b_2 + \dots}{(2y \ln(y) + y - x)^2}$$

$$= 0$$

Setting the numerator to zero gives

$$4 \ln(y)^2 y^2 b_2 - 4 \ln(y) xyb_2 - 2 \ln(y) y^2 a_2 + 4 \ln(y) y^2 b_2 + 2 \ln(y) y^2 b_3 \quad (\text{6E})$$

$$+ 2x^2 b_2 - y^2 a_2 - 2y^2 a_3 + y^2 b_2 + 3y^2 b_3 + xb_1 - ya_1 + 2yb_1 = 0$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \ln(y)\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \ln(y) = v_3\}$$

The above PDE (6E) now becomes

$$\begin{aligned} 4v_3^2v_2^2b_2 - 2v_3v_2^2a_2 - 4v_3v_1v_2b_2 + 4v_3v_2^2b_2 + 2v_3v_2^2b_3 - v_2^2a_2 \\ - 2v_2^2a_3 + 2v_1^2b_2 + v_2^2b_2 + 3v_2^2b_3 - v_2a_1 + v_1b_1 + 2v_2b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$\begin{aligned} 2v_1^2b_2 - 4v_3v_1v_2b_2 + v_1b_1 + 4v_3^2v_2^2b_2 + (-2a_2 + 4b_2 + 2b_3)v_2^2v_3 \\ + (-a_2 - 2a_3 + b_2 + 3b_3)v_2^2 + (-a_1 + 2b_1)v_2 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} b_1 &= 0 \\ -4b_2 &= 0 \\ 2b_2 &= 0 \\ 4b_2 &= 0 \\ -a_1 + 2b_1 &= 0 \\ -2a_2 + 4b_2 + 2b_3 &= 0 \\ -a_2 - 2a_3 + b_2 + 3b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= b_3 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}\xi &= x + y \\ \eta &= y\end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(\frac{y}{2y \ln(y) + y - x} \right) (x + y) \\ &= \frac{2y^2 \ln(y) - 2xy}{2y \ln(y) + y - x} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{2y^2 \ln(y) - 2xy}{2y \ln(y) + y - x}} dy\end{aligned}$$

Which results in

$$S = \frac{\ln(y^2 \ln(y) - xy)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{2y \ln(y) + y - x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{-2y \ln(y) + 2x} \\ S_y &= \frac{2y \ln(y) + y - x}{2y(y \ln(y) - x)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(y)}{2} + \frac{\ln(y \ln(y) - x)}{2} = c_1$$

Which simplifies to

$$\frac{\ln(y)}{2} + \frac{\ln(y \ln(y) - x)}{2} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{2y \ln(y) + y - x}$ 	$R = x$ $S = \frac{\ln(y)}{2} + \frac{\ln(y \ln(y) - x)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{\ln(y)}{2} + \frac{\ln(y \ln(y) - x)}{2} = c_1 \tag{1}$$




Figure 143: Slope field plot

Verification of solutions

$$\frac{\ln(y)}{2} + \frac{\ln(y \ln(y) - x)}{2} = c_1$$

Verified OK.

5.19.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(2y \ln(y) + y - x) dy &= (y) dx \\ (-y) dx + (2y \ln(y) + y - x) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y \\ N(x, y) &= 2y \ln(y) + y - x\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(2y \ln(y) + y - x) \\ &= -1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -y dx \\ \phi &= -xy + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 2y \ln(y) + y - x$. Therefore equation (4) becomes

$$2y \ln(y) + y - x = -x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 2y \ln(y) + y$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int ((2 \ln(y) + 1) y) dy \\ f(y) &= y^2 \ln(y) + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = y^2 \ln(y) - xy + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = y^2 \ln(y) - xy$$

Summary

The solution(s) found are the following

$$\ln(y) y^2 - yx = c_1 \tag{1}$$




Figure 144: Slope field plot

Verification of solutions

$$\ln(y) y^2 - yx = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
<- 1st order linear successful  
<- inverse linear successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=y(x)/(2*y(x)*ln(y(x))+y(x)-x),y(x), singsol=all)
```

$$y(x) = e^{\text{RootOf}(-Ze^{2-Z} - xe^{-Z} + c_1)}$$

✓ Solution by Mathematica

Time used: 0.198 (sec). Leaf size: 19

```
DSolve[y'[x]==y[x]/(2*y[x]*Log[y[x]]+y[x]-x),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[x = y(x) \log(y(x)) + \frac{c_1}{y(x)}, y(x)\right]$$

5.20 problem 20

- 5.20.1 Solving using Kovacic algorithm 2487
- 5.20.2 Maple step by step solution 2492

Internal problem ID [7313]

Internal file name [OUTPUT/6312_Monday_July_25_2022_10_21_11_PM_88747455/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 20.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**kovacic**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$xy'' - (2x + 1)y' + (1 + x)y = 0$$

5.20.1 Solving using Kovacic algorithm

Writing the ode as

$$xy'' + (-2x - 1)y' + (1 + x)y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = x$$

$$B = -2x - 1 \tag{3}$$

$$C = 1 + x$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{4x^2} \tag{6}$$

Comparing the above to (5) shows that

$$s = 3$$

$$t = 4x^2$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{3}{4x^2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 268: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{3}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + (-)(0) \\ &= -\frac{1}{2x} \\ &= -\frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{2x}\right)(0) + \left(\left(\frac{1}{2x^2}\right) + \left(-\frac{1}{2x}\right)^2 - \left(\frac{3}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int -\frac{1}{2x} dx} \\ &= \frac{1}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2x-1}{x} dx} \\ &= z_1 e^{x + \frac{\ln(x)}{2}} \\ &= z_1 (\sqrt{x} e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-2x-1}{x} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{2x+\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{x^2}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 (e^x) + c_2 \left(e^x \left(\frac{x^2}{2} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + \frac{x^2 e^x c_2}{2} \quad (1)$$

Verification of solutions

$$y = c_1 e^x + \frac{x^2 e^x c_2}{2}$$

Verified OK.

5.20.2 Maple step by step solution

Let's solve

$$xy'' + (-2x - 1)y' + (1 + x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(1+x)y}{x} + \frac{(2x+1)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(2x+1)y'}{x} + \frac{(1+x)y}{x} = 0$$

□ Check to see if $x_0 = 0$ is a regular singular point

○ Define functions

$$[P_2(x) = -\frac{2x+1}{x}, P_3(x) = \frac{1+x}{x}]$$

○ $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -1$$

○ $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

○ $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

• Multiply by denominators

$$xy'' + (-2x - 1)y' + (1 + x)y = 0$$

• Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

○ Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

○ Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

○ Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

○ Shift index using $k- > k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

○ Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)(k+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0r(-2+r)x^{-1+r} + (a_1(1+r)(-1+r) - a_0(-1+2r))x^r + \left(\sum_{k=1}^{\infty} (a_{k+1}(k+1+r)(k+r-1) \right.$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{0, 2\}$$

- Each term must be 0

$$a_1(1+r)(-1+r) - a_0(-1+2r) = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1+r)(k+r-1) + a_k(-2k-2r+1) + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$a_{k+2}(k+2+r)(k+r) + a_{k+1}(-2k-1-2r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = \frac{2ka_{k+1} + 2ra_{k+1} - a_k + a_{k+1}}{(k+2+r)(k+r)}$$

- Recursion relation for $r = 0$

$$a_{k+2} = \frac{2ka_{k+1} - a_k + a_{k+1}}{(k+2)k}$$

- Series not valid for $r = 0$, division by 0 in the recursion relation at $k = 0$

$$a_{k+2} = \frac{2ka_{k+1} - a_k + a_{k+1}}{(k+2)k}$$

- Recursion relation for $r = 2$

$$a_{k+2} = \frac{2ka_{k+1} - a_k + 5a_{k+1}}{(k+4)(k+2)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+2} = \frac{2ka_{k+1} - a_k + 5a_{k+1}}{(k+4)(k+2)}, 3a_1 - 3a_0 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(x*diff(y(x),x$2)-(2*x+1)*diff(y(x),x)+(x+1)*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x (c_2 x^2 + c_1)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 23

```
DSolve[x*y''[x]-(2*x+1)*y'[x]+(x+1)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^x (c_2 x^2 + 2c_1)$$

5.21 problem 21

5.21.1 Solving as separable ode	2496
5.21.2 Solving as first order special form ID 1 ode	2498
5.21.3 Solving as first order ode lie symmetry lookup ode	2499
5.21.4 Solving as exact ode	2503
5.21.5 Maple step by step solution	2507

Internal problem ID [7314]

Internal file name [OUTPUT/6313_Monday_July_25_2022_10_21_12_PM_64086885/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 21.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "separable", "first order special form ID 1", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_separable]`

$$x^2y' + e^{-y} = 0$$

5.21.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{e^{-y}}{x^2}\end{aligned}$$

Where $f(x) = -\frac{1}{x^2}$ and $g(y) = e^{-y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{e^{-y}} dy &= -\frac{1}{x^2} dx \\ \int \frac{1}{e^{-y}} dy &= \int -\frac{1}{x^2} dx \\ e^y &= \frac{1}{x} + c_1\end{aligned}$$

Which results in

$$y = -\ln\left(\frac{x}{c_1x + 1}\right)$$

Summary

The solution(s) found are the following

$$y = -\ln\left(\frac{x}{c_1x + 1}\right) \tag{1}$$




Figure 145: Slope field plot

Verification of solutions

$$y = -\ln\left(\frac{x}{c_1x + 1}\right)$$

Verified OK.

5.21.2 Solving as first order special form ID 1 ode

Writing the ode as

$$y' = -\frac{e^{-y}}{x^2} \quad (1)$$

And using the substitution $u = e^y$ then

$$u' = y'e^y$$

The above shows that

$$\begin{aligned} y' &= u'(x) e^{-y} \\ &= \frac{u'(x)}{u} \end{aligned}$$

Substituting this in (1) gives

$$\frac{u'(x)}{u} = -\frac{1}{x^2 u}$$

The above simplifies to

$$u'(x) = -\frac{1}{x^2} \quad (2)$$

Now ode (2) is solved for $u(x)$ Integrating both sides gives

$$\begin{aligned} u(x) &= \int -\frac{1}{x^2} dx \\ &= \frac{1}{x} + c_1 \end{aligned}$$

Substituting the solution found for $u(x)$ in $u = e^y$ gives

$$\begin{aligned} y &= \ln(u(x)) \\ &= \ln\left(\frac{1}{x} + c_1\right) \\ &= \ln\left(\frac{1}{x} + c_1\right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln\left(\frac{1}{x} + c_1\right) \quad (1)$$




Figure 146: Slope field plot

Verification of solutions

$$y = \ln\left(\frac{1}{x} + c_1\right)$$

Verified OK.

5.21.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{e^{-y}}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 270: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= -x^2 \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{-x^2} dx \end{aligned}$$

Which results in

$$S = \frac{1}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{e^{-y}}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 0 \\ R_y &= 1 \\ S_x &= -\frac{1}{x^2} \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^y \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = e^R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{1}{x} = e^y + c_1$$



Which simplifies to

$$\frac{1}{x} = e^y + c_1$$

Which gives

$$y = \ln \left(-\frac{c_1 x - 1}{x} \right)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{e^{-y}}{x^2}$ 	$R = y$ $S = \frac{1}{x}$	$\frac{dS}{dR} = e^R$ 

Summary

The solution(s) found are the following

$$y = \ln\left(-\frac{c_1x - 1}{x}\right) \quad (1)$$




Figure 147: Slope field plot

Verification of solutions

$$y = \ln\left(-\frac{c_1x - 1}{x}\right)$$

Verified OK.

5.21.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the

ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (-e^y) dy &= \left(\frac{1}{x^2}\right) dx \\ \left(-\frac{1}{x^2}\right) dx + (-e^y) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{1}{x^2} \\ N(x, y) &= -e^y \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x^2} \right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (-e^y) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x^2} dx \\ \phi &= \frac{1}{x} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -e^y$. Therefore equation (4) becomes

$$-e^y = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= -e^y \\ &= -e^y\end{aligned}$$

Integrating the above w.r.t y results in

$$\int f'(y) dy = \int (-e^y) dy$$
$$f(y) = -e^y + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{1}{x} - e^y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{1}{x} - e^y$$

The solution becomes

$$y = \ln \left(-\frac{c_1 x - 1}{x} \right)$$

Summary

The solution(s) found are the following

$$y = \ln \left(-\frac{c_1 x - 1}{x} \right) \tag{1}$$




Figure 148: Slope field plot

Verification of solutions

$$y = \ln\left(-\frac{c_1x - 1}{x}\right)$$

Verified OK.

5.21.5 Maple step by step solution

Let's solve

$$x^2y' + e^{-y} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{e^{-y}} = -\frac{1}{x^2}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{e^{-y}} dx = \int -\frac{1}{x^2} dx + c_1$$

- Evaluate integral

- $$\frac{1}{e^{-y}} = \frac{1}{x} + c_1$$
 Solve for y

$$y = -\ln\left(\frac{x}{c_1x+1}\right)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x)+exp(-y(x))=0,y(x), singsol=all)
```

$$y(x) = \ln\left(\frac{-c_1x + 1}{x}\right)$$

✓ Solution by Mathematica

Time used: 0.441 (sec). Leaf size: 12

```
DSolve[x^2*y'[x]+Exp[-y[x]]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log\left(\frac{1}{x} + c_1\right)$$

5.22 problem 22

- 5.22.1 Solving as second order ode can be made integrable ode 2509
- 5.22.2 Solving as second order ode missing x ode 2511
- 5.22.3 Maple step by step solution 2513

Internal problem ID [7315]

Internal file name [OUTPUT/6562_Friday_October_14_2022_05_49_35_AM_9550685/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 22.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**", "**second_order_ode_can_be_made_integrable**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]
```

$$y'' + e^y = 0$$

5.22.1 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' + y' e^y = 0$$

Integrating the above w.r.t x gives

$$\int (y' y'' + y' e^y) dx = 0$$
$$\frac{y'^2}{2} + e^y = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-2e^y + 2c_1} \tag{1}$$

$$y' = -\sqrt{-2e^y + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-2e^y + 2c_1}} dy = \int dx$$
$$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_2$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-2e^y + 2c_1}} dy = \int dx$$
$$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_3$$

Summary

The solution(s) found are the following

$$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_2 \quad (1)$$

$$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_3 \quad (2)$$

Verification of solutions

$$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_2$$

Verified OK.

$$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_3$$

Verified OK.

5.22.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned}y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy}\end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) = -e^y$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned}p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{e^y}{p}\end{aligned}$$

Where $f(y) = -e^y$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{p} dp &= -e^y dy \\ \int \frac{1}{p} dp &= \int -e^y dy \\ \frac{p^2}{2} &= -e^y + c_1\end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} + e^y - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} + e^y - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-2e^y + 2c_1} \quad (1)$$

$$y' = -\sqrt{-2e^y + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-2e^y + 2c_1}} dy = \int dx$$

$$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_2$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-2e^y + 2c_1}} dy = \int dx$$

$$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_3$$

Summary

The solution(s) found are the following

$$y = \ln \left(-\tanh \left(\frac{\sqrt{c_1} (x + c_2) \sqrt{2}}{2} \right)^2 c_1 + c_1 \right) \quad (1)$$

$$y = \ln \left(-\tanh \left(\frac{\sqrt{c_1} (x + c_3) \sqrt{2}}{2} \right)^2 c_1 + c_1 \right) \quad (2)$$

Verification of solutions

$$y = \ln \left(-\tanh \left(\frac{\sqrt{c_1} (x + c_2) \sqrt{2}}{2} \right)^2 c_1 + c_1 \right)$$

Verified OK.

$$y = \ln \left(-\tanh \left(\frac{\sqrt{c_1} (x + c_3) \sqrt{2}}{2} \right)^2 c_1 + c_1 \right)$$

Verified OK.

5.22.3 Maple step by step solution

Let's solve

$$y'' = -e^y$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$u(y) \left(\frac{d}{dy} u(y) \right) = -e^y$$

- Integrate both sides with respect to y

$$\int u(y) \left(\frac{d}{dy} u(y) \right) dy = \int -e^y dy + c_1$$

- Evaluate integral

$$\frac{u(y)^2}{2} = -e^y + c_1$$

- Solve for $u(y)$
 $\{u(y) = \sqrt{-2e^y + 2c_1}, u(y) = -\sqrt{-2e^y + 2c_1}\}$
- Solve 1st ODE for $u(y)$
 $u(y) = \sqrt{-2e^y + 2c_1}$
- Revert to original variables with substitution $u(y) = y', y = y$
 $y' = \sqrt{-2e^y + 2c_1}$
- Separate variables
 $\frac{y'}{\sqrt{-2e^y + 2c_1}} = 1$
- Integrate both sides with respect to x
 $\int \frac{y'}{\sqrt{-2e^y + 2c_1}} dx = \int 1 dx + c_2$
- Evaluate integral
 $-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = x + c_2$
- Solve for y
 $y = \ln\left(-\tanh\left(\frac{\sqrt{c_1}(x+c_2)\sqrt{2}}{2}\right)^2 c_1 + c_1\right)$
- Solve 2nd ODE for $u(y)$
 $u(y) = -\sqrt{-2e^y + 2c_1}$
- Revert to original variables with substitution $u(y) = y', y = y$
 $y' = -\sqrt{-2e^y + 2c_1}$
- Separate variables
 $\frac{y'}{\sqrt{-2e^y + 2c_1}} = -1$
- Integrate both sides with respect to x
 $\int \frac{y'}{\sqrt{-2e^y + 2c_1}} dx = \int (-1) dx + c_2$
- Evaluate integral
 $-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{-2e^y + 2c_1} \sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = -x + c_2$
- Solve for y
 $y = \ln\left(-\tanh\left(\frac{\sqrt{c_1}(-x+c_2)\sqrt{2}}{2}\right)^2 c_1 + c_1\right)$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
trying differential order: 2; missing variables  
, ` -> Computing symmetries using: way = 3  
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+exp(_a) = 0, _b(_a), HINT = [[1,  
    symmetry methods on request  
, `1st order, trying reduction of order with given symmetries:` [1, 1/2*_b]
```

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+exp(y(x))=0,y(x), singsol=all)
```

$$y(x) = -\ln(2) + \ln\left(\frac{\operatorname{sech}\left(\frac{x+c_2}{2c_1}\right)^2}{c_1^2}\right)$$

✓ Solution by Mathematica

Time used: 29.642 (sec). Leaf size: 60

```
DSolve[y''[x]+Exp[y[x]]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log\left(\frac{1}{2}c_1\operatorname{sech}^2\left(\frac{1}{2}\sqrt{c_1}(x+c_2)^2\right)\right)$$

$$y(x) \rightarrow \log\left(\frac{1}{2}c_1\operatorname{sech}^2\left(\frac{\sqrt{c_1}x^2}{2}\right)\right)$$

5.23 problem 23

Internal problem ID [7316]

Internal file name [OUTPUT/6563_Wednesday_October_19_2022_08_36_58_PM_9550685/index.tex]

Book: Own collection of miscellaneous problems

Section: section 5.0

Problem number: 23.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[_rational, [_Abel, `2nd type`, `class B`]]
```

Unable to solve or complete the solution.

$$y' - \frac{yx + 3x - 2y + 6}{yx - 3x - 2y + 6} = 0$$

Unable to determine ODE type.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
trying Abel
Looking for potential symmetries
Looking for potential symmetries
Looking for potential symmetries
trying inverse_Riccati
trying an equivalence to an Abel ODE
differential order: 1; trying a linearization to 2nd order
--- trying a change of variables {x -> y(x), y(x) -> x}
differential order: 1; trying a linearization to 2nd order
trying 1st order ODE linearizable_by_differentiation
--- Trying Lie symmetry methods, 1st order ---
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 4
`, `-> Computing symmetries using: way = 2
trying symmetry patterns for 1st order ODEs
-> trying a symmetry pattern of the form [F(x)*G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)*G(y)]
-> trying symmetry patterns of the forms [F(x),G(y)] and [G(y),F(x)]
-> trying a symmetry pattern of the form [F(x),G(x)]
-> trying a symmetry pattern of the form [F(y),G(y)]
-> trying a symmetry pattern of the form [F(x)+G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)+G(y)]
-> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying a symmetry pattern of conformal type`
```

X Solution by Maple

```
dsolve(diff(y(x),x)=(x*y(x)+3*x-2*y(x)+6)/(x*y(x)-3*x-2*y(x)+6),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y'[x]==(x*y[x]+3*x-2*y[x]+6)/(x*y[x]-3*x-2*y[x]+6),y[x],x,IncludeSingularSolutions ->
```

Not solved