


A Solution Manual For

Schaums Outline Differential Equations,
4th edition. Bronson and Costa.

McGraw Hill 2014



Nasser M. Abbasi

May 16, 2024

Contents

1	Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95	2
2	Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101	143
3	Chapter 12. VARIATION OF PARAMETERS. page 104	250
4	Chapter 12. VARIATION OF PARAMETERS. Supplementary Prob- lems. page 109	357
5	Chapter 24. Solutions of linear DE by Laplace transforms. Supple- mentary Problems. page 248	496
6	Chapter 27. Power series solutions of linear DE with variable coef- ficients. Supplementary Problems. page 274	590

1 Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

1.1	problem Problem 11.1	3
1.2	problem Problem 11.2	14
1.3	problem Problem 11.3	25
1.4	problem Problem 11.4	36
1.5	problem Problem 11.5	47
1.6	problem Problem 11.6	58
1.7	problem Problem 11.7	69
1.8	problem Problem 11.8	77
1.9	problem Problem 11.10	93
1.10	problem Problem 11.12	104
1.11	problem Problem 11.13	117
1.12	problem Problem 11.14	130

1.1 problem Problem 11.1

1.1.1	Solving as second order linear constant coeff ode	3
1.1.2	Solving using Kovacic algorithm	6
1.1.3	Maple step by step solution	11

Internal problem ID [5163]

Internal file name [OUTPUT/4656_Sunday_June_05_2022_03_02_48_PM_29362966/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.1.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - 2y = 4x^2$$

1.1.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -2, f(x) = 4x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-2)} \\ &= \frac{1}{2} \pm \frac{3}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} + \frac{3}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{3}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3x^2 + A_2x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_3x^2 - 2A_2x - 2xA_3 - 2A_1 - A_2 + 2A_3 = 4x^2$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -3, A_2 = 2, A_3 = -2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2x^2 + 2x - 3$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^{2x} + c_2e^{-x}) + (-2x^2 + 2x - 3) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{2x} + c_2e^{-x} - 2x^2 + 2x - 3 \quad (1)$$




Figure 1: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-x} - 2x^2 + 2x - 3$$

Verified OK.

1.1.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -1 \\ C &= -2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 1: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\ &= z_1 e^{\frac{x}{2}} \\ &= z_1 \left(e^{\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{3x}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{3x}}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^{2x}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{2x}}{3}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_3x^2 - 2A_2x - 2xA_3 - 2A_1 - A_2 + 2A_3 = 4x^2$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -3, A_2 = 2, A_3 = -2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2x^2 + 2x - 3$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{-x} + \frac{c_2e^{2x}}{3} \right) + (-2x^2 + 2x - 3) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-x} + \frac{c_2e^{2x}}{3} - 2x^2 + 2x - 3 \quad (1)$$




Figure 2: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} - 2x^2 + 2x - 3$$

Verified OK.

1.1.3 Maple step by step solution

Let's solve

$$y'' - y' - 2y = 4x^2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 4x^2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{2x} \\ -e^{-x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{4e^{-x}(\int x^2 e^x dx)}{3} + \frac{4e^{2x}(\int x^2 e^{-2x} dx)}{3}$$

- Compute integrals

$$y_p(x) = -2x^2 + 2x - 3$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{2x} - 2x^2 + 2x - 3$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=4*x^2,y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{2x} c_1 - 2x^2 + 2x - 3$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 31

```
DSolve[y''[x]-y'[x]-2*y[x]==4*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2x^2 + 2x + c_1 e^{-x} + c_2 e^{2x} - 3$$

1.2 problem Problem 11.2

1.2.1	Solving as second order linear constant coeff ode	14
1.2.2	Solving using Kovacic algorithm	17
1.2.3	Maple step by step solution	22

Internal problem ID [5164]

Internal file name [OUTPUT/4657_Sunday_June_05_2022_03_02_49_PM_41519936/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - 2y = e^{3x}$$

1.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -2, f(x) = e^{3x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-2)} \\ &= \frac{1}{2} \pm \frac{3}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} + \frac{3}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{3}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1 e^{3x} = e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{3x}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^{-x}) + \left(\frac{e^{3x}}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-x} + \frac{e^{3x}}{4} \quad (1)$$




Figure 3: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-x} + \frac{e^{3x}}{4}$$

Verified OK.

1.2.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -1 \\ C &= -2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 3: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\ &= z_1 e^{\frac{x}{2}} \\ &= z_1 \left(e^{\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{3x}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{3x}}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^{2x}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{2x}}{3}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1e^{3x} = e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{3x}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{-x} + \frac{c_2e^{2x}}{3} \right) + \left(\frac{e^{3x}}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-x} + \frac{c_2e^{2x}}{3} + \frac{e^{3x}}{4} \quad (1)$$




Figure 4: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} + \frac{e^{3x}}{4}$$

Verified OK.

1.2.3 Maple step by step solution

Let's solve

$$y'' - y' - 2y = e^{3x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{3x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{2x} \\ -e^{-x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x}(\int e^{4x} dx)}{3} + \frac{e^{2x}(\int e^x dx)}{3}$$

- Compute integrals

$$y_p(x) = \frac{e^{3x}}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + \frac{e^{3x}}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=exp(3*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{2x} c_1 + \frac{e^{3x}}{4}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 31

```
DSolve[y''[x]-y'[x]-2*y[x]==Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{3x}}{4} + c_1 e^{-x} + c_2 e^{2x}$$

1.3 problem Problem 11.3

1.3.1	Solving as second order linear constant coeff ode	25
1.3.2	Solving using Kovacic algorithm	28
1.3.3	Maple step by step solution	33

Internal problem ID [5165]

Internal file name [OUTPUT/4658_Sunday_June_05_2022_03_02_50_PM_59966924/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - y' - 2y = \sin(2x)$$

1.3.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -2, f(x) = \sin(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-2)} \\ &= \frac{1}{2} \pm \frac{3}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} + \frac{3}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{3}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(2x) + A_2 \sin(2x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-6A_1 \cos(2x) - 6A_2 \sin(2x) + 2A_1 \sin(2x) - 2A_2 \cos(2x) = \sin(2x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{20}, A_2 = -\frac{3}{20} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^{-x}) + \left(\frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-x} + \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20} \quad (1)$$




Figure 5: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-x} + \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20}$$

Verified OK.

1.3.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= -1 \\C &= -2\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{4}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 9 \\t &= 4\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{4}\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 5: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
O(\infty) &= \deg(t) - \deg(s) \\
&= 0 - 0 \\
&= 0
\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
&= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\
&= z_1 e^{\frac{x}{2}} \\
&= z_1 \left(e^{\frac{x}{2}} \right)
\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{3x}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{3x}}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^{2x}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{2x}}{3}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(2x) + A_2 \sin(2x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-6A_1 \cos(2x) - 6A_2 \sin(2x) + 2A_1 \sin(2x) - 2A_2 \cos(2x) = \sin(2x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{20}, A_2 = -\frac{3}{20} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-x} + \frac{c_2 e^{2x}}{3} \right) + \left(\frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} + \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20} \quad (1)$$




Figure 6: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} + \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20}$$

Verified OK.

1.3.3 Maple step by step solution

Let's solve

$$y'' - y' - 2y = \sin(2x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{2x} \\ -e^{-x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x} \left(\int e^x \sin(2x) dx \right)}{3} + \frac{e^{2x} \left(\int e^{-2x} \sin(2x) dx \right)}{3}$$

- Compute integrals

$$y_p(x) = \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=sin(2*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{2x} c_1 + \frac{\cos(2x)}{20} - \frac{3 \sin(2x)}{20}$$

✓ Solution by Mathematica

Time used: 0.116 (sec). Leaf size: 37

```
DSolve[y''[x]-y'[x]-2*y[x]==Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x} + c_2 e^{2x} + \frac{1}{20}(\cos(2x) - 3 \sin(2x))$$

1.4 problem Problem 11.4

1.4.1	Solving as second order linear constant coeff ode	36
1.4.2	Solving using Kovacic algorithm	39
1.4.3	Maple step by step solution	44

Internal problem ID [5166]

Internal file name [OUTPUT/4659_Sunday_June_05_2022_03_02_51_PM_17805542/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 6y' + 25y = 2 \sin\left(\frac{t}{2}\right) - \cos\left(\frac{t}{2}\right)$$

1.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = f(t)$$

Where $A = 1, B = -6, C = 25, f(t) = 2 \sin\left(\frac{t}{2}\right) - \cos\left(\frac{t}{2}\right)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' - 6y' + 25y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = 0$$

Where in the above $A = 1, B = -6, C = 25$. Let the solution be $y = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} - 6\lambda e^{\lambda t} + 25 e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 - 6\lambda + 25 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -6, C = 25$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{6}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-6^2 - (4)(1)(25)} \\ &= 3 \pm 4i \end{aligned}$$

Hence

$$\lambda_1 = 3 + 4i$$

$$\lambda_2 = 3 - 4i$$

Which simplifies to

$$\lambda_1 = 3 + 4i$$

$$\lambda_2 = 3 - 4i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 3$ and $\beta = 4$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha t} (c_1 \cos(\beta t) + c_2 \sin(\beta t))$$

Which becomes

$$y = e^{3t} (c_1 \cos(4t) + c_2 \sin(4t))$$

Therefore the homogeneous solution y_h is

$$y_h = e^{3t} (c_1 \cos(4t) + c_2 \sin(4t))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2 \sin\left(\frac{t}{2}\right) - \cos\left(\frac{t}{2}\right)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$\left[\left\{ \cos\left(\frac{t}{2}\right), \sin\left(\frac{t}{2}\right) \right\} \right]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{3t} \cos(4t), e^{3t} \sin(4t)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos\left(\frac{t}{2}\right) + A_2 \sin\left(\frac{t}{2}\right)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\frac{99A_1 \cos\left(\frac{t}{2}\right)}{4} + \frac{99A_2 \sin\left(\frac{t}{2}\right)}{4} + 3A_1 \sin\left(\frac{t}{2}\right) - 3A_2 \cos\left(\frac{t}{2}\right) = 2 \sin\left(\frac{t}{2}\right) - \cos\left(\frac{t}{2}\right)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{20}{663}, A_2 = \frac{56}{663} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{20 \cos\left(\frac{t}{2}\right)}{663} + \frac{56 \sin\left(\frac{t}{2}\right)}{663}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{3t}(c_1 \cos(4t) + c_2 \sin(4t))) + \left(-\frac{20 \cos\left(\frac{t}{2}\right)}{663} + \frac{56 \sin\left(\frac{t}{2}\right)}{663} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3t}(c_1 \cos(4t) + c_2 \sin(4t)) - \frac{20 \cos\left(\frac{t}{2}\right)}{663} + \frac{56 \sin\left(\frac{t}{2}\right)}{663} \quad (1)$$




Figure 7: Slope field plot

Verification of solutions

$$y = e^{3t}(c_1 \cos(4t) + c_2 \sin(4t)) - \frac{20 \cos\left(\frac{t}{2}\right)}{663} + \frac{56 \sin\left(\frac{t}{2}\right)}{663}$$

Verified OK.

1.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 6y' + 25y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= -6 \\C &= 25\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-16}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -16 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = -16z(t)\tag{7}$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 7: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
O(\infty) &= \deg(t) - \deg(s) \\
&= 0 - 0 \\
&= 0
\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -16$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = \cos(4t)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\
&= z_1 e^{-\int \frac{1}{2} \frac{-6}{1} dt} \\
&= z_1 e^{3t} \\
&= z_1 (e^{3t})
\end{aligned}$$

Which simplifies to

$$y_1 = e^{3t} \cos(4t)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{6}{1} dt}}{(y_1)^2} dt \\ &= y_1 \int \frac{e^{6t}}{(y_1)^2} dt \\ &= y_1 \left(\frac{\tan(4t)}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{3t} \cos(4t)) + c_2 \left(e^{3t} \cos(4t) \left(\frac{\tan(4t)}{4} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' - 6y' + 25y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2 \sin\left(\frac{t}{2}\right) - \cos\left(\frac{t}{2}\right)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$\left[\left\{ \cos\left(\frac{t}{2}\right), \sin\left(\frac{t}{2}\right) \right\} \right]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{3t} \cos(4t), \frac{e^{3t} \sin(4t)}{4} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos\left(\frac{t}{2}\right) + A_2 \sin\left(\frac{t}{2}\right)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\frac{99A_1 \cos\left(\frac{t}{2}\right)}{4} + \frac{99A_2 \sin\left(\frac{t}{2}\right)}{4} + 3A_1 \sin\left(\frac{t}{2}\right) - 3A_2 \cos\left(\frac{t}{2}\right) = 2 \sin\left(\frac{t}{2}\right) - \cos\left(\frac{t}{2}\right)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{20}{663}, A_2 = \frac{56}{663} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{20 \cos\left(\frac{t}{2}\right)}{663} + \frac{56 \sin\left(\frac{t}{2}\right)}{663}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} \right) + \left(-\frac{20 \cos\left(\frac{t}{2}\right)}{663} + \frac{56 \sin\left(\frac{t}{2}\right)}{663} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} - \frac{20 \cos\left(\frac{t}{2}\right)}{663} + \frac{56 \sin\left(\frac{t}{2}\right)}{663} \quad (1)$$




Figure 8: Slope field plot

Verification of solutions

$$y = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} - \frac{20 \cos\left(\frac{t}{2}\right)}{663} + \frac{56 \sin\left(\frac{t}{2}\right)}{663}$$

Verified OK.

1.4.3 Maple step by step solution

Let's solve

$$y'' - 6y' + 25y = 2 \sin\left(\frac{t}{2}\right) - \cos\left(\frac{t}{2}\right)$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 - 6r + 25 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{6 \pm (\sqrt{-64})}{2}$$

- Roots of the characteristic polynomial

$$r = (3 - 4I, 3 + 4I)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{3t} \cos(4t)$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{3t} \sin(4t)$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{3t} \cos(4t) c_1 + e^{3t} \sin(4t) c_2 + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = 2 \sin\left(\frac{t}{2}\right) - \cos\left(\frac{t}{2}\right) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{3t} \cos(4t) & e^{3t} \sin(4t) \\ 3e^{3t} \cos(4t) - 4e^{3t} \sin(4t) & 3e^{3t} \sin(4t) + 4e^{3t} \cos(4t) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = 4e^{6t}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = \frac{e^{3t} (\cos(4t) (\int \sin(4t) (-2 \sin(\frac{t}{2}) + \cos(\frac{t}{2})) e^{-3t} dt) - \sin(4t) (\int \cos(4t) (-2 \sin(\frac{t}{2}) + \cos(\frac{t}{2})) e^{-3t} dt))}{4}$$

- Compute integrals

$$y_p(t) = -\frac{20 \cos(\frac{t}{2})}{663} + \frac{56 \sin(\frac{t}{2})}{663}$$

- Substitute particular solution into general solution to ODE

$$y = e^{3t} \cos(4t) c_1 + e^{3t} \sin(4t) c_2 - \frac{20 \cos(\frac{t}{2})}{663} + \frac{56 \sin(\frac{t}{2})}{663}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.578 (sec). Leaf size: 37

```
dsolve(diff(y(t),t$2)-6*diff(y(t),t)+25*y(t)=2*sin(t/2)-cos(t/2),y(t), singsol=all)
```

$$y(t) = e^{3t} \sin(4t) c_2 + e^{3t} \cos(4t) c_1 + \frac{56 \sin\left(\frac{t}{2}\right)}{663} - \frac{20 \cos\left(\frac{t}{2}\right)}{663}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 51

```
DSolve[y''[t]-6*y'[t]+25*y[t]==2*Sin[t/2]-Cos[t/2],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{663} \left(56 \sin\left(\frac{t}{2}\right) - 20 \cos\left(\frac{t}{2}\right) \right) + c_2 e^{3t} \cos(4t) + c_1 e^{3t} \sin(4t)$$

1.5 problem Problem 11.5

1.5.1	Solving as second order linear constant coeff ode	47
1.5.2	Solving using Kovacic algorithm	50
1.5.3	Maple step by step solution	55

Internal problem ID [5167]

Internal file name [OUTPUT/4660_Sunday_June_05_2022_03_02_53_PM_58783062/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 6y' + 25y = 64e^{-t}$$

1.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = f(t)$$

Where $A = 1, B = -6, C = 25, f(t) = 64e^{-t}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' - 6y' + 25y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = 0$$

Where in the above $A = 1, B = -6, C = 25$. Let the solution be $y = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} - 6\lambda e^{\lambda t} + 25 e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 - 6\lambda + 25 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -6, C = 25$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{6}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-6^2 - (4)(1)(25)} \\ &= 3 \pm 4i \end{aligned}$$

Hence

$$\lambda_1 = 3 + 4i$$

$$\lambda_2 = 3 - 4i$$

Which simplifies to

$$\lambda_1 = 3 + 4i$$

$$\lambda_2 = 3 - 4i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 3$ and $\beta = 4$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha t} (c_1 \cos(\beta t) + c_2 \sin(\beta t))$$

Which becomes

$$y = e^{3t} (c_1 \cos(4t) + c_2 \sin(4t))$$

Therefore the homogeneous solution y_h is

$$y_h = e^{3t} (c_1 \cos(4t) + c_2 \sin(4t))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$64 e^{-t}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-t}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{3t} \cos(4t), e^{3t} \sin(4t)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-t}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$32A_1 e^{-t} = 64 e^{-t}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2 e^{-t}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{3t}(c_1 \cos(4t) + c_2 \sin(4t))) + (2 e^{-t}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3t}(c_1 \cos(4t) + c_2 \sin(4t)) + 2 e^{-t} \quad (1)$$




Figure 9: Slope field plot

Verification of solutions

$$y = e^{3t}(c_1 \cos(4t) + c_2 \sin(4t)) + 2e^{-t}$$

Verified OK.

1.5.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 6y' + 25y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -6 \tag{3}$$

$$C = 25$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-16}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -16 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = -16z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 9: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -16$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = \cos(4t)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{B}{A} dt} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-6}{1} dt} \\ &= z_1 e^{3t} \\ &= z_1 (e^{3t}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{3t} \cos(4t)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-6}{1} dt}}{(y_1)^2} dt \\ &= y_1 \int \frac{e^{6t}}{(y_1)^2} dt \\ &= y_1 \left(\frac{\tan(4t)}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{3t} \cos(4t)) + c_2 \left(e^{3t} \cos(4t) \left(\frac{\tan(4t)}{4} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' - 6y' + 25y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$64e^{-t}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[e^{-t}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{3t} \cos(4t), \frac{e^{3t} \sin(4t)}{4} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-t}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$32A_1e^{-t} = 64e^{-t}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2e^{-t}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} \right) + (2e^{-t}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} + 2e^{-t} \quad (1)$$




Figure 10: Slope field plot

Verification of solutions

$$y = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} + 2e^{-t}$$

Verified OK.

1.5.3 Maple step by step solution

Let's solve

$$y'' - 6y' + 25y = 64e^{-t}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 6r + 25 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{6 \pm (\sqrt{-64})}{2}$$

- Roots of the characteristic polynomial

$$r = (3 - 4I, 3 + 4I)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{3t} \cos(4t)$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{3t} \sin(4t)$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{3t} \cos(4t) c_1 + e^{3t} \sin(4t) c_2 + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = 64 e^{-t} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{3t} \cos(4t) & e^{3t} \sin(4t) \\ 3e^{3t} \cos(4t) - 4e^{3t} \sin(4t) & 3e^{3t} \sin(4t) + 4e^{3t} \cos(4t) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = 4e^{6t}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = 16 e^{3t} (-\cos(4t) \left(\int \sin(4t) e^{-4t} dt \right) + \sin(4t) \left(\int \cos(4t) e^{-4t} dt \right))$$

- Compute integrals

$$y_p(t) = 2e^{-t}$$

- Substitute particular solution into general solution to ODE

$$y = e^{3t} \sin(4t) c_2 + e^{3t} \cos(4t) c_1 + 2e^{-t}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve(diff(y(t),t$2)-6*diff(y(t),t)+25*y(t)=64*exp(-t),y(t), singsol=all)
```

$$y(t) = e^{3t} \sin(4t) c_2 + e^{3t} \cos(4t) c_1 + 2e^{-t}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 37

```
DSolve[y''[t]-6*y'[t]+25*y[t]==64*Exp[-t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^{-t} (c_2 e^{4t} \cos(4t) + c_1 e^{4t} \sin(4t) + 2)$$

1.6 problem Problem 11.6

1.6.1	Solving as second order linear constant coeff ode	58
1.6.2	Solving using Kovacic algorithm	61
1.6.3	Maple step by step solution	66

Internal problem ID [5168]

Internal file name [OUTPUT/4661_Sunday_June_05_2022_03_02_54_PM_98534113/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 6y' + 25y = 50t^3 - 36t^2 - 63t + 18$$

1.6.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = f(t)$$

Where $A = 1, B = -6, C = 25, f(t) = 50t^3 - 36t^2 - 63t + 18$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' - 6y' + 25y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(t) + By'(t) + Cy(t) = 0$$

Where in the above $A = 1, B = -6, C = 25$. Let the solution be $y = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} - 6\lambda e^{\lambda t} + 25 e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 - 6\lambda + 25 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -6, C = 25$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{6}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-6^2 - (4)(1)(25)} \\ &= 3 \pm 4i \end{aligned}$$

Hence

$$\lambda_1 = 3 + 4i$$

$$\lambda_2 = 3 - 4i$$

Which simplifies to

$$\lambda_1 = 3 + 4i$$

$$\lambda_2 = 3 - 4i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 3$ and $\beta = 4$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha t} (c_1 \cos(\beta t) + c_2 \sin(\beta t))$$

Which becomes

$$y = e^{3t} (c_1 \cos(4t) + c_2 \sin(4t))$$

Therefore the homogeneous solution y_h is

$$y_h = e^{3t} (c_1 \cos(4t) + c_2 \sin(4t))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$t^3 + t^2 + t + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, t, t^2, t^3\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{3t} \cos(4t), e^{3t} \sin(4t)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_4 t^3 + A_3 t^2 + A_2 t + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} 25A_4 t^3 + 25A_3 t^2 - 18t^2 A_4 + 25A_2 t - 12t A_3 + 6t A_4 + 25A_1 - 6A_2 + 2A_3 \\ = 50t^3 - 36t^2 - 63t + 18 \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = -3, A_3 = 0, A_4 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2t^3 - 3t$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{3t}(c_1 \cos(4t) + c_2 \sin(4t))) + (2t^3 - 3t) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3t}(c_1 \cos(4t) + c_2 \sin(4t)) + 2t^3 - 3t \quad (1)$$




Figure 11: Slope field plot

Verification of solutions

$$y = e^{3t}(c_1 \cos(4t) + c_2 \sin(4t)) + 2t^3 - 3t$$

Verified OK.

1.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 6y' + 25y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -6 \\ C &= 25 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = ye^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-16}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -16 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = -16z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then y is found using the inverse transformation

$$y = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 11: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -16$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = \cos(4t)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{B}{A} dt} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-6}{1} dt} \\ &= z_1 e^{3t} \\ &= z_1 (e^{3t}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{3t} \cos(4t)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dt}}{y_1^2} dt$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-6}{1} dt}}{(y_1)^2} dt \\ &= y_1 \int \frac{e^{6t}}{(y_1)^2} dt \\ &= y_1 \left(\frac{\tan(4t)}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{3t} \cos(4t)) + c_2 \left(e^{3t} \cos(4t) \left(\frac{\tan(4t)}{4} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(t) + By'(t) + Cy(t) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(t) + By'(t) + Cy(t) = f(t)$. y_h is the solution to

$$y'' - 6y' + 25y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$t^3 + t^2 + t + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[1, t, t^2, t^3]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{3t} \cos(4t), \frac{e^{3t} \sin(4t)}{4} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_4 t^3 + A_3 t^2 + A_2 t + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} 25A_4t^3 + 25A_3t^2 - 18t^2A_4 + 25A_2t - 12tA_3 + 6tA_4 + 25A_1 - 6A_2 + 2A_3 \\ = 50t^3 - 36t^2 - 63t + 18 \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = -3, A_3 = 0, A_4 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2t^3 - 3t$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} \right) + (2t^3 - 3t) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} + 2t^3 - 3t \quad (1)$$




Figure 12: Slope field plot

Verification of solutions

$$y = e^{3t} \cos(4t) c_1 + \frac{e^{3t} \sin(4t) c_2}{4} + 2t^3 - 3t$$

Verified OK.

1.6.3 Maple step by step solution

Let's solve

$$y'' - 6y' + 25y = 50t^3 - 36t^2 - 63t + 18$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 6r + 25 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{6 \pm (\sqrt{-64})}{2}$$

- Roots of the characteristic polynomial

$$r = (3 - 4I, 3 + 4I)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{3t} \cos(4t)$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{3t} \sin(4t)$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{3t} \cos(4t) c_1 + e^{3t} \sin(4t) c_2 + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = 50t^3 - 36t^2 - 63t + 18 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{3t} \cos(4t) & e^{3t} \sin(4t) \\ 3e^{3t} \cos(4t) - 4e^{3t} \sin(4t) & 3e^{3t} \sin(4t) + 4e^{3t} \cos(4t) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = 4e^{6t}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -\frac{e^{3t}(\cos(4t)(\int e^{-3t} \sin(4t)(50t^3 - 36t^2 - 63t + 18) dt) - \sin(4t)(\int \cos(4t)e^{-3t}(50t^3 - 36t^2 - 63t + 18) dt))}{4}$$

- Compute integrals

$$y_p(t) = 2t^3 - 3t$$

- Substitute particular solution into general solution to ODE

$$y = e^{3t} \cos(4t) c_1 + e^{3t} \sin(4t) c_2 + 2t^3 - 3t$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(t),t$2)-6*diff(y(t),t)+25*y(t)=50*t^3-36*t^2-63*t+18,y(t), singsol=all)
```

$$y(t) = e^{3t} \sin(4t) c_2 + e^{3t} \cos(4t) c_1 + 2t^3 - 3t$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 38

```
DSolve[y''[t]-6*y'[t]+25*y[t]==50*t^3-36*t^2-63*t+18,y[t],t,IncludeSingularSolutions -> True
```

$$y(t) \rightarrow 2t^3 - 3t + c_2 e^{3t} \cos(4t) + c_1 e^{3t} \sin(4t)$$

1.7 problem Problem 11.7

1.7.1 Maple step by step solution 71

Internal problem ID [5169]

Internal file name [OUTPUT/4662_Sunday_June_05_2022_03_02_55_PM_22257302/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.7.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _linear , _nonhomogeneous]]
```

$$y''' - 6y'' + 11y' - 6y = 2x e^{-x}$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' - 6y'' + 11y' - 6y = 0$$

The characteristic equation is

$$\lambda^3 - 6\lambda^2 + 11\lambda - 6 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = 2$$

$$\lambda_3 = 3$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^x + c_2 e^{2x} + c_3 e^{3x}$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^x$$

$$y_2 = e^{2x}$$

$$y_3 = e^{3x}$$

Now the particular solution to the given ODE is found

$$y''' - 6y'' + 11y' - 6y = 2x e^{-x}$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2x e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{x e^{-x}, e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, e^{2x}, e^{3x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 x e^{-x} + A_2 e^{-x}$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$26A_1 e^{-x} - 24A_1 x e^{-x} - 24A_2 e^{-x} = 2x e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{12}, A_2 = -\frac{13}{144} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x e^{-x}}{12} - \frac{13 e^{-x}}{144}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 e^{2x} + c_3 e^{3x}) + \left(-\frac{x e^{-x}}{12} - \frac{13 e^{-x}}{144} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 e^{2x} + c_3 e^{3x} - \frac{x e^{-x}}{12} - \frac{13 e^{-x}}{144} \quad (1)$$

Verification of solutions

$$y = c_1 e^x + c_2 e^{2x} + c_3 e^{3x} - \frac{x e^{-x}}{12} - \frac{13 e^{-x}}{144}$$

Verified OK.

1.7.1 Maple step by step solution

Let's solve

$$y''' - 6y'' + 11y' - 6y = 2x e^{-x}$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 2x e^{-x} + 6y_3(x) - 11y_2(x) + 6y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 2x e^{-x} + 6y_3(x) - 11y_2(x) + 6y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 6 & -11 & 6 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ 2x e^{-x} \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ 2x e^{-x} \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 6 & -11 & 6 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[\begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[\begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[3, \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{3x} \cdot \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + \vec{y}_p(x)$$

- Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} e^x & \frac{e^{2x}}{4} & \frac{e^{3x}}{9} \\ e^x & \frac{e^{2x}}{2} & \frac{e^{3x}}{3} \\ e^x & e^{2x} & e^{3x} \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} e^x & \frac{e^{2x}}{4} & \frac{e^{3x}}{9} \\ e^x & \frac{e^{2x}}{2} & \frac{e^{3x}}{3} \\ e^x & e^{2x} & e^{3x} \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} 1 & \frac{1}{4} & \frac{1}{9} \\ 1 & \frac{1}{2} & \frac{1}{3} \\ 1 & 1 & 1 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} 3e^x - 3e^{2x} + e^{3x} & -\frac{5e^x}{2} + 4e^{2x} - \frac{3e^{3x}}{2} & \frac{e^x}{2} - e^{2x} + \frac{e^{3x}}{2} \\ 3e^x - 6e^{2x} + 3e^{3x} & -\frac{5e^x}{2} + 8e^{2x} - \frac{9e^{3x}}{2} & \frac{e^x}{2} - 2e^{2x} + \frac{3e^{3x}}{2} \\ 3e^x - 12e^{2x} + 9e^{3x} & -\frac{5e^x}{2} + 16e^{2x} - \frac{27e^{3x}}{2} & \frac{e^x}{2} - 4e^{2x} + \frac{9e^{3x}}{2} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} \frac{(-12x-13)e^{-x}}{144} + \frac{e^x}{4} - \frac{2e^{2x}}{9} + \frac{e^{3x}}{16} \\ \frac{(12x+1)e^{-x}}{144} + \frac{e^x}{4} - \frac{4e^{2x}}{9} + \frac{3e^{3x}}{16} \\ \frac{(-12x+11)e^{-x}}{144} + \frac{e^x}{4} - \frac{8e^{2x}}{9} + \frac{9e^{3x}}{16} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + \begin{bmatrix} \frac{(-12x-13)e^{-x}}{144} + \frac{e^x}{4} - \frac{2e^{2x}}{9} + \frac{e^{3x}}{16} \\ \frac{(12x+1)e^{-x}}{144} + \frac{e^x}{4} - \frac{4e^{2x}}{9} + \frac{3e^{3x}}{16} \\ \frac{(-12x+11)e^{-x}}{144} + \frac{e^x}{4} - \frac{8e^{2x}}{9} + \frac{9e^{3x}}{16} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{(-12x-13)e^{-x}}{144} + \frac{(-8+9c_2)e^{2x}}{36} + \frac{(16c_3+9)e^{3x}}{144} + \frac{e^x(1+4c_1)}{4}$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 3; missing the dependent variable
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$3)-6*diff(y(x),x$2)+11*diff(y(x),x)-6*y(x)=2*x*exp(-x),y(x), singsol=all)
```

$$y(x) = \frac{(-12x - 13)e^{-x}}{144} + e^x c_1 + c_2 e^{2x} + c_3 e^{3x}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 42

```
DSolve[y'''[x]-6*y''[x]+11*y'[x]-6*y[x]==2*x*Exp[-x],y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow -\frac{1}{144}e^{-x}(12x + 13) + c_1 e^x + c_2 e^{2x} + c_3 e^{3x}$$

1.8 problem Problem 11.8

1.8.1	Solving as second order ode quadrature ode	77
1.8.2	Solving as second order linear constant coeff ode	78
1.8.3	Solving as second order integrable as is ode	81
1.8.4	Solving as second order ode missing y ode	82
1.8.5	Solving using Kovacic algorithm	84
1.8.6	Solving as exact linear second order ode ode	89
1.8.7	Maple step by step solution	91

Internal problem ID [5170]

Internal file name [OUTPUT/4663_Sunday_June_05_2022_03_02_56_PM_99349420/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = 9x^2 + 2x - 1$$

1.8.1 Solving as second order ode quadrature ode

Integrating once gives

$$y' = 3x^3 + x^2 - x + c_1$$

Integrating again gives

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2 \quad (1)$$




Figure 13: Slope field plot

Verification of solutions

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2$$

Verified OK.

1.8.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 0, f(x) = 9x^2 + 2x - 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3, x^4\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_3x^4 + A_2x^3 + A_1x^2$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$12x^2A_3 + 6xA_2 + 2A_1 = 9x^2 + 2x - 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = \frac{1}{3}, A_3 = \frac{3}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2x + c_1 + \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 \quad (1)$$




Figure 14: Slope field plot

Verification of solutions

$$y = c_2x + c_1 + \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2$$

Verified OK.

1.8.3 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = \int (9x^2 + 2x - 1) dx$$

$$y' = 3x^3 + x^2 - x + c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int 3x^3 + x^2 + c_1 - x dx$$

$$= \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2 \quad (1)$$




Figure 15: Slope field plot

Verification of solutions

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2$$

Verified OK.

1.8.4 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 9x^2 - 2x + 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(x) &= \int 9x^2 + 2x - 1 \, dx \\ &= 3x^3 + x^2 + c_1 - x \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = 3x^3 + x^2 + c_1 - x$$

Integrating both sides gives

$$\begin{aligned} y &= \int 3x^3 + x^2 + c_1 - x \, dx \\ &= \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2 \tag{1}$$




Figure 16: Slope field plot

Verification of solutions

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2$$

Verified OK.

1.8.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 14: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3, x^4\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_3x^4 + A_2x^3 + A_1x^2$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$12x^2A_3 + 6xA_2 + 2A_1 = 9x^2 + 2x - 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = \frac{1}{3}, A_3 = \frac{3}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2x + c_1 + \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 \quad (1)$$




Figure 17: Slope field plot

Verification of solutions

$$y = c_2x + c_1 + \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2$$

Verified OK.

1.8.6 Solving as exact linear second order ode

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \\ s(x) &= 9x^2 + 2x - 1 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = \int 9x^2 + 2x - 1 dx$$

We now have a first order ode to solve which is

$$y' = 3x^3 + x^2 + c_1 - x$$

Integrating both sides gives

$$\begin{aligned}y &= \int 3x^3 + x^2 + c_1 - x \, dx \\ &= \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2 \quad (1)$$




Figure 18: Slope field plot

Verification of solutions

$$y = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2$$

Verified OK.

1.8.7 Maple step by step solution

Let's solve

$$y'' = 9x^2 + 2x - 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 9x^2 + 2x - 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\left(\int (9x^3 + 2x^2 - x) dx\right) + x\left(\int (9x^2 + 2x - 1) dx\right)$$

- Compute integrals

$$y_p(x) = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2$$

- Substitute particular solution into general solution to ODE

$$y = c_2x + c_1 + \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)=9*x^2+2*x-1,y(x), singsol=all)
```

$$y(x) = \frac{3}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 33

```
DSolve[y''[x]==9*x^2+2*x-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x^4}{4} + \frac{x^3}{3} - \frac{x^2}{2} + c_2x + c_1$$

1.9 problem Problem 11.10

1.9.1	Solving as second order linear constant coeff ode	93
1.9.2	Solving using Kovacic algorithm	96
1.9.3	Maple step by step solution	101

Internal problem ID [5171]

Internal file name [OUTPUT/4664_Sunday_June_05_2022_03_02_57_PM_17961132/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.10.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 5y = 2e^{5x}$$

1.9.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = -5, f(x) = 2e^{5x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 5y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 5 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 5 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -5$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-5)} \\ &= \pm \sqrt{5} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +\sqrt{5} \\ \lambda_2 &= -\sqrt{5} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= \sqrt{5} \\ \lambda_2 &= -\sqrt{5} \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(\sqrt{5})x} + c_2 e^{(-\sqrt{5})x} \end{aligned}$$

Or

$$y = c_1 e^{x\sqrt{5}} + c_2 e^{-x\sqrt{5}}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{x\sqrt{5}} + c_2 e^{-x\sqrt{5}}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2e^{5x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{5x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{x\sqrt{5}}, e^{-x\sqrt{5}}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{5x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$20A_1 e^{5x} = 2e^{5x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{10} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{5x}}{10}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{x\sqrt{5}} + c_2 e^{-x\sqrt{5}}) + \left(\frac{e^{5x}}{10} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{x\sqrt{5}} + c_2 e^{-x\sqrt{5}} + \frac{e^{5x}}{10} \quad (1)$$




Figure 19: Slope field plot

Verification of solutions

$$y = c_1 e^{x\sqrt{5}} + c_2 e^{-x\sqrt{5}} + \frac{e^{5x}}{10}$$

Verified OK.

1.9.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 5y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= -5 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{5}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 5 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 5z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 16: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 5$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x\sqrt{5}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= e^{-x\sqrt{5}} \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x\sqrt{5}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-x\sqrt{5}} \int \frac{1}{e^{-2x\sqrt{5}}} dx \\ &= e^{-x\sqrt{5}} \left(\frac{e^{2x\sqrt{5}} \sqrt{5}}{10} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-x\sqrt{5}} \right) + c_2 \left(e^{-x\sqrt{5}} \left(\frac{e^{2x\sqrt{5}} \sqrt{5}}{10} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 5y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x\sqrt{5}} + \frac{c_2 \sqrt{5} e^{x\sqrt{5}}}{10}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2e^{5x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{5x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sqrt{5} e^{x\sqrt{5}}}{10}, e^{-x\sqrt{5}} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{5x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$20A_1e^{5x} = 2e^{5x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{10} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{5x}}{10}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{-x\sqrt{5}} + \frac{c_2\sqrt{5}e^{x\sqrt{5}}}{10} \right) + \left(\frac{e^{5x}}{10} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-x\sqrt{5}} + \frac{c_2\sqrt{5}e^{x\sqrt{5}}}{10} + \frac{e^{5x}}{10} \quad (1)$$




Figure 20: Slope field plot

Verification of solutions

$$y = c_1 e^{-x\sqrt{5}} + \frac{c_2 \sqrt{5} e^{x\sqrt{5}}}{10} + \frac{e^{5x}}{10}$$

Verified OK.

1.9.3 Maple step by step solution

Let's solve

$$y'' - 5y = 2e^{5x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{20})}{2}$$

- Roots of the characteristic polynomial
 $r = (\sqrt{5}, -\sqrt{5})$
- 1st solution of the homogeneous ODE
 $y_1(x) = e^{x\sqrt{5}}$
- 2nd solution of the homogeneous ODE
 $y_2(x) = e^{-x\sqrt{5}}$
- General solution of the ODE
 $y = c_1y_1(x) + c_2y_2(x) + y_p(x)$
- Substitute in solutions of the homogeneous ODE
 $y = c_1e^{x\sqrt{5}} + c_2e^{-x\sqrt{5}} + y_p(x)$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 2e^{5x} \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{x\sqrt{5}} & e^{-x\sqrt{5}} \\ \sqrt{5}e^{x\sqrt{5}} & -\sqrt{5}e^{-x\sqrt{5}} \end{bmatrix}$$
 - Compute Wronskian
 $W(y_1(x), y_2(x)) = -2\sqrt{5}$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{\sqrt{5} \left(e^{x\sqrt{5}} \left(\int e^{-x(-5+\sqrt{5})} dx \right) - e^{-x\sqrt{5}} \left(\int e^{x(5+\sqrt{5})} dx \right) \right)}{5}$$
 - Compute integrals
 $y_p(x) = \frac{e^{5x}}{10}$
- Substitute particular solution into general solution to ODE
 $y = c_1e^{x\sqrt{5}} + c_2e^{-x\sqrt{5}} + \frac{e^{5x}}{10}$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-5*y(x)=2*exp(5*x),y(x), singsol=all)
```

$$y(x) = e^{\sqrt{5}x}c_2 + e^{-\sqrt{5}x}c_1 + \frac{e^{5x}}{10}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 40

```
DSolve[y''[x]-5*y[x]==2*Exp[5*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{5x}}{10} + c_1 e^{\sqrt{5}x} + c_2 e^{-\sqrt{5}x}$$

1.10 problem Problem 11.12

1.10.1 Solving as linear ode	104
1.10.2 Solving as first order ode lie symmetry lookup ode	106
1.10.3 Solving as exact ode	110
1.10.4 Maple step by step solution	115

Internal problem ID [5172]

Internal file name [OUTPUT/4665_Sunday_June_05_2022_03_02_58_PM_49852271/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.12.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - 5y = (x - 1) \sin(x) + (x + 1) \cos(x)$$

1.10.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -5$$

$$q(x) = (x - 1) \sin(x) + (x + 1) \cos(x)$$

Hence the ode is

$$y' - 5y = (x - 1) \sin(x) + (x + 1) \cos(x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-5) dx} \\ &= e^{-5x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) ((x-1) \sin(x) + (x+1) \cos(x)) \\ \frac{d}{dx}(e^{-5x} y) &= (e^{-5x}) ((x-1) \sin(x) + (x+1) \cos(x)) \\ d(e^{-5x} y) &= (((x-1) \sin(x) + (x+1) \cos(x)) e^{-5x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-5x} y &= \int ((x-1) \sin(x) + (x+1) \cos(x)) e^{-5x} dx \\ e^{-5x} y &= \left(-\frac{x}{26} - \frac{5}{338}\right) e^{-5x} \cos(x) + \left(-\frac{5x}{26} - \frac{6}{169}\right) e^{-5x} \sin(x) + \left(-\frac{5x}{26} - \frac{6}{169}\right) e^{-5x} \cos(x) - \left(-\frac{x}{26}\right)\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-5x}$ results in

$$y = e^{5x} \left(\left(-\frac{x}{26} - \frac{5}{338}\right) e^{-5x} \cos(x) + \left(-\frac{5x}{26} - \frac{6}{169}\right) e^{-5x} \sin(x) + \left(-\frac{5x}{26} - \frac{6}{169}\right) e^{-5x} \cos(x) - \left(-\frac{x}{26}\right) \right)$$

which simplifies to

$$y = c_1 e^{5x} + \frac{(-78x - 69) \cos(x)}{338} + \frac{(-52x + 71) \sin(x)}{338}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{5x} + \frac{(-78x - 69) \cos(x)}{338} + \frac{(-52x + 71) \sin(x)}{338} \quad (1)$$




Figure 21: Slope field plot

Verification of solutions

$$y = c_1 e^{5x} + \frac{(-78x - 69) \cos(x)}{338} + \frac{(-52x + 71) \sin(x)}{338}$$

Verified OK.

1.10.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \cos(x) x + \sin(x) x + \cos(x) - \sin(x) + 5y$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 18: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{5x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{5x}} dy \end{aligned}$$

Which results in

$$S = e^{-5x} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \cos(x)x + \sin(x)x + \cos(x) - \sin(x) + 5y$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -5e^{-5x}y \\ S_y &= e^{-5x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = ((x - 1) \sin(x) + (x + 1) \cos(x)) e^{-5x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = ((R - 1) \sin(R) + (R + 1) \cos(R)) e^{-5R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 - \frac{e^{-5R}(52 \sin(R) R + 78 \cos(R) R - 71 \sin(R) + 69 \cos(R))}{338} \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-5x} y = c_1 - \frac{e^{-5x}(52 \sin(x) x + 78 \cos(x) x - 71 \sin(x) + 69 \cos(x))}{338}$$



Which simplifies to

$$\frac{((78x + 69) \cos(x) + (52x - 71) \sin(x) + 338y) e^{-5x}}{338} - c_1 = 0$$

Which gives

$$y = -\frac{e^{5x}(52 e^{-5x} \sin(x) x + 78 e^{-5x} \cos(x) x - 71 e^{-5x} \sin(x) + 69 e^{-5x} \cos(x) - 338c_1)}{338}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \cos(x) x + \sin(x) x + \cos(x) - \sin(x) + 5y$ 	$R = x$ $S = e^{-5x} y$	$\frac{dS}{dR} = ((R - 1) \sin(R) + (R + 1) \cos(R)) e^{-5R}$ 

Summary

The solution(s) found are the following

$$y = -\frac{e^{5x}(52 e^{-5x} \sin(x) x + 78 e^{-5x} \cos(x) x - 71 e^{-5x} \sin(x) + 69 e^{-5x} \cos(x) - 338c_1)}{338} \quad (1)$$




Figure 22: Slope field plot

Verification of solutions

$$y = -\frac{e^{5x}(52 e^{-5x} \sin(x) x + 78 e^{-5x} \cos(x) x - 71 e^{-5x} \sin(x) + 69 e^{-5x} \cos(x) - 338c_1)}{338}$$

Verified OK.

1.10.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the

ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= (5y + (x - 1) \sin(x) + (x + 1) \cos(x)) dx \\ (-5y - (x - 1) \sin(x) - (x + 1) \cos(x)) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -5y - (x - 1) \sin(x) - (x + 1) \cos(x) \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-5y - (x-1)\sin(x) - (x+1)\cos(x)) \\ &= -5\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-5) - (0)) \\ &= -5\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int -5 dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-5x} \\ &= e^{-5x}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\overline{M} &= \mu M \\ &= e^{-5x}(-5y - (x-1)\sin(x) - (x+1)\cos(x)) \\ &= -(5y + (x-1)\sin(x) + (x+1)\cos(x))e^{-5x}\end{aligned}$$

And

$$\begin{aligned}\overline{N} &= \mu N \\ &= e^{-5x}(1) \\ &= e^{-5x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ (-5y + (x - 1) \sin(x) + (x + 1) \cos(x)) e^{-5x} + (e^{-5x}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -(5y + (x - 1) \sin(x) + (x + 1) \cos(x)) e^{-5x} dx \\ \phi &= \frac{((78x + 69) \cos(x) + (52x - 71) \sin(x) + 338y) e^{-5x}}{338} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-5x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-5x}$. Therefore equation (4) becomes

$$e^{-5x} = e^{-5x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{((78x + 69) \cos(x) + (52x - 71) \sin(x) + 338y) e^{-5x}}{338} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{((78x + 69) \cos(x) + (52x - 71) \sin(x) + 338y) e^{-5x}}{338}$$

The solution becomes

$$y = -\frac{e^{5x}(52 e^{-5x} \sin(x) x + 78 e^{-5x} \cos(x) x - 71 e^{-5x} \sin(x) + 69 e^{-5x} \cos(x) - 338c_1)}{338}$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{5x}(52 e^{-5x} \sin(x) x + 78 e^{-5x} \cos(x) x - 71 e^{-5x} \sin(x) + 69 e^{-5x} \cos(x) - 338c_1)}{338} \quad (1)$$




Figure 23: Slope field plot

Verification of solutions

$$y = -\frac{e^{5x}(52 e^{-5x} \sin(x) x + 78 e^{-5x} \cos(x) x - 71 e^{-5x} \sin(x) + 69 e^{-5x} \cos(x) - 338c_1)}{338}$$

Verified OK.

1.10.4 Maple step by step solution

Let's solve

$$y' - 5y = (x - 1) \sin(x) + (x + 1) \cos(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = \cos(x) x + \sin(x) x + \cos(x) - \sin(x) + 5y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - 5y = \cos(x) x + \sin(x) x + \cos(x) - \sin(x)$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - 5y) = \mu(x) (\cos(x) x + \sin(x) x + \cos(x) - \sin(x))$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' - 5y) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -5\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{-5x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) (\cos(x) x + \sin(x) x + \cos(x) - \sin(x)) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) (\cos(x) x + \sin(x) x + \cos(x) - \sin(x)) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) (\cos(x) x + \sin(x) x + \cos(x) - \sin(x)) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-5x}$

$$y = \frac{\int e^{-5x} (\cos(x) x + \sin(x) x + \cos(x) - \sin(x)) dx + c_1}{e^{-5x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{\left(-\frac{x}{26} - \frac{5}{338}\right)e^{-5x} \cos(x) + \left(-\frac{5x}{26} - \frac{6}{169}\right)e^{-5x} \sin(x) + \left(-\frac{5x}{26} - \frac{6}{169}\right)e^{-5x} \cos(x) - \left(-\frac{x}{26} - \frac{5}{338}\right)e^{-5x} \sin(x) - \frac{2e^{-5x} \cos(x)}{13} + \frac{3e^{-5x} \sin(x)}{13}}{e^{-5x}}$$

- Simplify

$$y = c_1 e^{5x} + \frac{(-78x-69) \cos(x)}{338} + \frac{(-52x+71) \sin(x)}{338}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x), x) - 5*y(x) = (x-1)*sin(x) + (x+1)*cos(x), y(x), singsol=all)
```

$$y(x) = c_1 e^{5x} + \frac{(-78x - 69) \cos(x)}{338} + \frac{(-52x + 71) \sin(x)}{338}$$

✓ Solution by Mathematica

Time used: 0.229 (sec). Leaf size: 36

```
DSolve[y' [x] - 5*y [x] == (x-1)*Sin [x] + (x+1)*Cos [x], y [x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{338}((71 - 52x) \sin(x) - 3(26x + 23) \cos(x)) + c_1 e^{5x}$$

1.11 problem Problem 11.13

1.11.1 Solving as linear ode	117
1.11.2 Solving as first order ode lie symmetry lookup ode	119
1.11.3 Solving as exact ode	123
1.11.4 Maple step by step solution	127

Internal problem ID [5173]

Internal file name [OUTPUT/4666_Sunday_June_05_2022_03_02_59_PM_68550176/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.13.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - 5y = 3e^x - 2x + 1$$

1.11.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -5$$

$$q(x) = 3e^x - 2x + 1$$

Hence the ode is

$$y' - 5y = 3e^x - 2x + 1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-5) dx} \\ &= e^{-5x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(3e^x - 2x + 1) \\ \frac{d}{dx}(e^{-5x}y) &= (e^{-5x})(3e^x - 2x + 1) \\ d(e^{-5x}y) &= ((3e^x - 2x + 1)e^{-5x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-5x}y &= \int (3e^x - 2x + 1)e^{-5x} dx \\ e^{-5x}y &= -\frac{3e^{-5x}}{25} - \frac{3e^{-4x}}{4} + \frac{2xe^{-5x}}{5} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-5x}$ results in

$$y = e^{5x} \left(-\frac{3e^{-5x}}{25} - \frac{3e^{-4x}}{4} + \frac{2xe^{-5x}}{5} \right) + c_1 e^{5x}$$

which simplifies to

$$y = -\frac{3}{25} - \frac{3e^x}{4} + \frac{2x}{5} + c_1 e^{5x}$$

Summary

The solution(s) found are the following

$$y = -\frac{3}{25} - \frac{3e^x}{4} + \frac{2x}{5} + c_1 e^{5x} \quad (1)$$




Figure 24: Slope field plot

Verification of solutions

$$y = -\frac{3}{25} - \frac{3e^x}{4} + \frac{2x}{5} + c_1e^{5x}$$

Verified OK.

1.11.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = 5y + 3e^x - 2x + 1$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 21: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{5x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{5x}} dy \end{aligned}$$

Which results in

$$S = e^{-5x} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = 5y + 3e^x - 2x + 1$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -5e^{-5x}y \\ S_y &= e^{-5x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 3e^{-4x} - 2xe^{-5x} + e^{-5x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 3e^{-4R} - 2Re^{-5R} + e^{-5R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{2R e^{-5R}}{5} - \frac{3 e^{-5R}}{25} - \frac{3 e^{-4R}}{4} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-5x} y = \frac{2x e^{-5x}}{5} - \frac{3 e^{-5x}}{25} - \frac{3 e^{-4x}}{4} + c_1$$



Which simplifies to

$$\frac{(3 - 10x + 25y) e^{-5x}}{25} - c_1 + \frac{3 e^{-4x}}{4} = 0$$

Which gives

$$y = -\frac{(-40x e^{-5x} + 75 e^{-4x} + 12 e^{-5x} - 100c_1) e^{5x}}{100}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = 5y + 3e^x - 2x + 1$ 	$R = x$ $S = e^{-5x} y$	$\frac{dS}{dR} = 3e^{-4R} - 2R e^{-5R} + e^{-5R}$ 

Summary

The solution(s) found are the following

$$y = -\frac{(-40x e^{-5x} + 75 e^{-4x} + 12 e^{-5x} - 100c_1) e^{5x}}{100} \quad (1)$$




Figure 25: Slope field plot

Verification of solutions

$$y = -\frac{(-40x e^{-5x} + 75 e^{-4x} + 12 e^{-5x} - 100c_1) e^{5x}}{100}$$

Verified OK.

1.11.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (5y + 3e^x - 2x + 1) dx \\ (-5y - 3e^x + 2x - 1) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -5y - 3e^x + 2x - 1 \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-5y - 3e^x + 2x - 1) \\ &= -5\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-5) - (0)) \\ &= -5 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int -5 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-5x} \\ &= e^{-5x} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{-5x}(-5y - 3e^x + 2x - 1) \\ &= (-5y - 3e^x + 2x - 1)e^{-5x} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{-5x}(1) \\ &= e^{-5x} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((-5y - 3e^x + 2x - 1)e^{-5x}) + (e^{-5x}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (-5y - 3e^x + 2x - 1)e^{-5x} dx \\ \phi &= \frac{(3 - 10x + 25y)e^{-5x}}{25} + \frac{3e^{-4x}}{4} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-5x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-5x}$. Therefore equation (4) becomes

$$e^{-5x} = e^{-5x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(3 - 10x + 25y)e^{-5x}}{25} + \frac{3e^{-4x}}{4} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(3 - 10x + 25y)e^{-5x}}{25} + \frac{3e^{-4x}}{4}$$

The solution becomes

$$y = -\frac{(-40xe^{-5x} + 75e^{-4x} + 12e^{-5x} - 100c_1)e^{5x}}{100}$$

Summary

The solution(s) found are the following

$$y = -\frac{(-40x e^{-5x} + 75 e^{-4x} + 12 e^{-5x} - 100c_1) e^{5x}}{100} \quad (1)$$




Figure 26: Slope field plot

Verification of solutions

$$y = -\frac{(-40x e^{-5x} + 75 e^{-4x} + 12 e^{-5x} - 100c_1) e^{5x}}{100}$$

Verified OK.

1.11.4 Maple step by step solution

Let's solve

$$y' - 5y = 3e^x - 2x + 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = 5y + 3e^x - 2x + 1$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - 5y = 3e^x - 2x + 1$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - 5y) = \mu(x) (3e^x - 2x + 1)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) (y' - 5y) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -5\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{-5x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) (3e^x - 2x + 1) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) (3e^x - 2x + 1) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x)(3e^x - 2x + 1) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-5x}$

$$y = \frac{\int (3e^x - 2x + 1)e^{-5x} dx + c_1}{e^{-5x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{2x}{5(e^x)^5} - \frac{3}{25(e^x)^5} - \frac{3}{4(e^x)^4} + c_1}{e^{-5x}}$$

- Simplify

$$y = -\frac{3}{25} - \frac{3e^x}{4} + \frac{2x}{5} + c_1e^{5x}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)-5*y(x)=3*exp(x)-2*x+1,y(x), singsol=all)
```

$$y(x) = \frac{2x}{5} - \frac{3}{25} - \frac{3e^x}{4} + c_1e^{5x}$$

✓ Solution by Mathematica

Time used: 0.112 (sec). Leaf size: 29

```
DSolve[y'[x]-5*y[x]==3*Exp[x]-2*x+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2x}{5} - \frac{3e^x}{4} + c_1e^{5x} - \frac{3}{25}$$

1.12 problem Problem 11.14

1.12.1 Solving as linear ode	130
1.12.2 Solving as first order ode lie symmetry lookup ode	132
1.12.3 Solving as exact ode	136
1.12.4 Maple step by step solution	140

Internal problem ID [5174]

Internal file name [OUTPUT/4667_Sunday_June_05_2022_03_03_00_PM_98372085/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. page 95

Problem number: Problem 11.14.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - 5y = x^2e^x - e^{5x}x$$

1.12.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -5$$
$$q(x) = e^x x(-e^{4x} + x)$$

Hence the ode is

$$y' - 5y = e^x x(-e^{4x} + x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-5) dx} \\ &= e^{-5x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (e^x x (-e^{4x} + x)) \\ \frac{d}{dx}(e^{-5x} y) &= (e^{-5x}) (e^x x (-e^{4x} + x)) \\ d(e^{-5x} y) &= (x(e^{-4x} x - 1)) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-5x} y &= \int x(e^{-4x} x - 1) dx \\ e^{-5x} y &= -\frac{x^2}{2} - \frac{e^{-4x} x^2}{4} - \frac{e^{-4x} x}{8} - \frac{e^{-4x}}{32} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-5x}$ results in

$$y = e^{5x} \left(-\frac{x^2}{2} - \frac{e^{-4x} x^2}{4} - \frac{e^{-4x} x}{8} - \frac{e^{-4x}}{32} \right) + c_1 e^{5x}$$

which simplifies to

$$y = \frac{(-x^2 + 2c_1) e^{5x}}{2} + \frac{(-8x^2 - 4x - 1) e^x}{32}$$

Summary

The solution(s) found are the following

$$y = \frac{(-x^2 + 2c_1) e^{5x}}{2} + \frac{(-8x^2 - 4x - 1) e^x}{32} \quad (1)$$




Figure 27: Slope field plot

Verification of solutions

$$y = \frac{(-x^2 + 2c_1)e^{5x}}{2} + \frac{(-8x^2 - 4x - 1)e^x}{32}$$

Verified OK.

1.12.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = 5y + x^2e^x - e^{5x}x$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 24: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{5x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{5x}} dy \end{aligned}$$

Which results in

$$S = e^{-5x}y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = 5y + x^2e^x - e^{5x}x$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -5e^{-5x}y \\ S_y &= e^{-5x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x(e^{-4x}x - 1) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R(e^{-4R}R - 1)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{R^2}{2} - \frac{e^{-4R}R^2}{4} - \frac{e^{-4R}R}{8} - \frac{e^{-4R}}{32} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-5x}y = -\frac{x^2}{2} - \frac{e^{-4x}x^2}{4} - \frac{e^{-4x}x}{8} - \frac{e^{-4x}}{32} + c_1$$



Which simplifies to

$$\frac{(8x^2 + 4x + 1)e^{-4x}}{32} + \frac{x^2}{2} + e^{-5x}y - c_1 = 0$$

Which gives

$$y = -\frac{(8e^{-4x}x^2 + 4e^{-4x}x + 16x^2 + e^{-4x} - 32c_1)e^{5x}}{32}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = 5y + x^2e^x - e^{5x}x$ 	$R = x$ $S = e^{-5x}y$	$\frac{dS}{dR} = R(e^{-4R}R - 1)$ 

Summary

The solution(s) found are the following

$$y = -\frac{(8e^{-4x}x^2 + 4e^{-4x}x + 16x^2 + e^{-4x} - 32c_1)e^{5x}}{32} \quad (1)$$




Figure 28: Slope field plot

Verification of solutions

$$y = -\frac{(8e^{-4x}x^2 + 4e^{-4x}x + 16x^2 + e^{-4x} - 32c_1)e^{5x}}{32}$$

Verified OK.

1.12.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (5y + x^2e^x - e^{5x}x) dx \\ (-5y - x^2e^x + e^{5x}x) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -5y - x^2e^x + e^{5x}x \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-5y - x^2e^x + e^{5x}x) \\ &= -5\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-5) - (0)) \\ &= -5 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int -5 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-5x} \\ &= e^{-5x} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{-5x} (-5y - x^2 e^x + e^{5x} x) \\ &= (-5y - x^2 e^x + e^{5x} x) e^{-5x} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{-5x} (1) \\ &= e^{-5x} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((-5y - x^2 e^x + e^{5x} x) e^{-5x}) + (e^{-5x}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (-5y - x^2 e^x + e^{5x} x) e^{-5x} dx \\ \phi &= \frac{(8x^2 + 4x + 1) e^{-4x}}{32} + \frac{x^2}{2} + e^{-5x} y + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-5x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-5x}$. Therefore equation (4) becomes

$$e^{-5x} = e^{-5x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(8x^2 + 4x + 1) e^{-4x}}{32} + \frac{x^2}{2} + e^{-5x} y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(8x^2 + 4x + 1) e^{-4x}}{32} + \frac{x^2}{2} + e^{-5x} y$$

The solution becomes

$$y = -\frac{(8e^{-4x}x^2 + 4e^{-4x}x + 16x^2 + e^{-4x} - 32c_1)e^{5x}}{32}$$

Summary

The solution(s) found are the following

$$y = -\frac{(8e^{-4x}x^2 + 4e^{-4x}x + 16x^2 + e^{-4x} - 32c_1)e^{5x}}{32} \quad (1)$$




Figure 29: Slope field plot

Verification of solutions

$$y = -\frac{(8e^{-4x}x^2 + 4e^{-4x}x + 16x^2 + e^{-4x} - 32c_1)e^{5x}}{32}$$

Verified OK.

1.12.4 Maple step by step solution

Let's solve

$$y' - 5y = x^2e^x - e^{5x}x$$

- Highest derivative means the order of the ODE is 1

y'

- Isolate the derivative

$$y' = 5y + x^2e^x - e^{5x}x$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - 5y = x^2e^x - e^{5x}x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - 5y) = \mu(x) (x^2e^x - e^{5x}x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) (y' - 5y) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -5\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{-5x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) (x^2e^x - e^{5x}x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) (x^2e^x - e^{5x}x) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x)(x^2e^x - e^{5x}x)dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-5x}$

$$y = \frac{\int e^{-5x}(x^2e^x - e^{5x}x)dx + c_1}{e^{-5x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{-\frac{x^2}{2} - \frac{x^2}{4(e^x)^4} - \frac{x}{8(e^x)^4} - \frac{1}{32(e^x)^4} + c_1}{e^{-5x}}$$

- Simplify

$$y = -\frac{e^x(x^2 - 2c_1)e^{4x}}{2} + \frac{(-8x^2 - 4x - 1)e^x}{32}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(y(x),x)-5*y(x)=x^2*exp(x)-x*exp(5*x),y(x), singsol=all)
```

$$y(x) = -\frac{(x^2 - 2c_1)e^x e^{4x}}{2} + \frac{(-8x^2 - 4x - 1)e^x}{32}$$

✓ Solution by Mathematica

Time used: 0.209 (sec). Leaf size: 39

```
DSolve[y'[x]-5*y[x]==x^2*Exp[x]-x*Exp[5*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{32}e^x(8x^2 + 4x + 1) + e^{5x}\left(-\frac{x^2}{2} + c_1\right)$$

2 Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS.

Supplementary Problems. page 101

2.1	problem Problem 11.44	144
2.2	problem Problem 11.45	156
2.3	problem Problem 11.46	168
2.4	problem Problem 11.47	180
2.5	problem Problem 11.48	193
2.6	problem Problem 11.49	206
2.7	problem Problem 11.50	219
2.8	problem Problem 11.51	232
2.9	problem Problem 11.52	246

2.1 problem Problem 11.44

2.1.1	Solving as second order linear constant coeff ode	144
2.1.2	Solving as linear second order ode solved by an integrating factor ode	147
2.1.3	Solving using Kovacic algorithm	149
2.1.4	Maple step by step solution	154

Internal problem ID [5175]

Internal file name [OUTPUT/4668_Sunday_June_05_2022_03_03_02_PM_73750163/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.44.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2y' + y = x^2 - 1$$

2.1.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = x^2 - 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 e^x x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3x^2 + A_2x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_3x^2 + A_2x - 4xA_3 + A_1 - 2A_2 + 2A_3 = x^2 - 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 5, A_2 = 4, A_3 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^2 + 4x + 5$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^x + c_2xe^x) + (x^2 + 4x + 5) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2x + c_1) + x^2 + 4x + 5$$

Summary

The solution(s) found are the following

$$y = e^x(c_2x + c_1) + x^2 + 4x + 5 \quad (1)$$




Figure 30: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + x^2 + 4x + 5$$

Verified OK.

2.1.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x))^2 + p'(x)}{2}y = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2 dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = e^{-x}(x^2 - 1)$$

$$(e^{-x}y)'' = e^{-x}(x^2 - 1)$$

Integrating once gives

$$(e^{-x}y)' = -(x+1)^2 e^{-x} + c_1$$

Integrating again gives

$$(e^{-x}y) = (x^2 + 4x + 5) e^{-x} + c_1x + c_2$$

Hence the solution is

$$y = \frac{(x^2 + 4x + 5) e^{-x} + c_1x + c_2}{e^{-x}}$$

Or

$$y = c_1x e^x + c_2e^x + x^2 + 4x + 5$$

Summary

The solution(s) found are the following

$$y = c_1x e^x + c_2e^x + x^2 + 4x + 5 \tag{1}$$




Figure 31: Slope field plot

Verification of solutions

$$y = c_1 x e^x + c_2 e^x + x^2 + 4x + 5$$

Verified OK.

2.1.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 27: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1(e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_3 x^2 + A_2 x - 4x A_3 + A_1 - 2A_2 + 2A_3 = x^2 - 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 5, A_2 = 4, A_3 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^2 + 4x + 5$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (x^2 + 4x + 5) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) + x^2 + 4x + 5$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) + x^2 + 4x + 5 \quad (1)$$




Figure 32: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + x^2 + 4x + 5$$

Verified OK.

2.1.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = x^2 - 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = e^x x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 x e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^2 - 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^x x \\ e^x & e^x x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$
- Substitute functions into equation for $y_p(x)$

$$y_p(x) = e^x \left(-\int (x^3 - x) e^{-x} dx \right) + \left(\int e^{-x} (x^2 - 1) dx \right) x$$
- Compute integrals

$$y_p(x) = x^2 + 4x + 5$$
- Substitute particular solution into general solution to ODE

$$y = c_2 x e^x + c_1 e^x + x^2 + 4x + 5$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=x^2-1,y(x), singsol=all)
```

$$y(x) = (c_1 x + c_2) e^x + x^2 + 4x + 5$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 26

```
DSolve[y''[x]-2*y'[x]+y[x]==x^2-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + x(4 + c_2 e^x) + c_1 e^x + 5$$

2.2 problem Problem 11.45

2.2.1	Solving as second order linear constant coeff ode	156
2.2.2	Solving as linear second order ode solved by an integrating factor ode	159
2.2.3	Solving using Kovacic algorithm	161
2.2.4	Maple step by step solution	166

Internal problem ID [5176]

Internal file name [OUTPUT/4669_Sunday_June_05_2022_03_03_03_PM_52958204/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.45.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2y' + y = 4e^{2x}$$

2.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = 4e^{2x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 e^x x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$4 e^{2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{2x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 e^{2x} = 4 e^{2x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 4]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 4 e^{2x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (4 e^{2x}) \end{aligned}$$

Which simplifies to

$$y = e^x (c_2 x + c_1) + 4 e^{2x}$$

Summary

The solution(s) found are the following

$$y = e^x (c_2 x + c_1) + 4 e^{2x} \tag{1}$$




Figure 33: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + 4e^{2x}$$

Verified OK.

2.2.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x))^2 + p'(x)}{2}y = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2 dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = 4 e^{-x} e^{2x}$$

$$(e^{-x}y)'' = 4 e^{-x} e^{2x}$$

Integrating once gives

$$(e^{-x}y)' = 4 e^x + c_1$$

Integrating again gives

$$(e^{-x}y) = c_1 x + 4 e^x + c_2$$

Hence the solution is

$$y = \frac{c_1 x + 4 e^x + c_2}{e^{-x}}$$

Or

$$y = c_1 x e^x + c_2 e^x + 4 e^{2x}$$

Summary

The solution(s) found are the following

$$y = c_1 x e^x + c_2 e^x + 4 e^{2x} \tag{1}$$




Figure 34: Slope field plot

Verification of solutions

$$y = c_1 x e^x + c_2 e^x + 4 e^{2x}$$

Verified OK.

2.2.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 29: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1 (e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^x) + c_2 (e^x(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$4 e^{2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, x e^x\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{2x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 e^{2x} = 4 e^{2x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 4]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 4 e^{2x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (4 e^{2x}) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2x + c_1) + 4e^{2x}$$

Summary

The solution(s) found are the following

$$y = e^x(c_2x + c_1) + 4e^{2x} \tag{1}$$




Figure 35: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + 4e^{2x}$$

Verified OK.

2.2.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = 4e^{2x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = e^x x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 x e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 4e^{2x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^x x \\ e^x & e^x x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -4e^x \left(\int e^x x dx - \left(\int e^x dx \right) x \right)$$

- Compute integrals

$$y_p(x) = 4e^{2x}$$

- Substitute particular solution into general solution to ODE

$$y = c_2 x e^x + c_1 e^x + 4e^{2x}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=4*exp(2*x),y(x), singsol=all)
```

$$y(x) = 4e^{2x} + (c_1 x + c_2) e^x$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 21

```
DSolve[y''[x]-2*y'[x]+y[x]==4*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(4e^x + c_2 x + c_1)$$

2.3 problem Problem 11.46

2.3.1	Solving as second order linear constant coeff ode	168
2.3.2	Solving as linear second order ode solved by an integrating factor ode	171
2.3.3	Solving using Kovacic algorithm	173
2.3.4	Maple step by step solution	178

Internal problem ID [5177]

Internal file name [OUTPUT/4670_Sunday_June_05_2022_03_03_04_PM_31127158/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.46.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 2y' + y = 4 \cos(x)$$

2.3.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = 4 \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 e^x x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$4 \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 \sin(x) - 2A_2 \cos(x) = 4 \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = -2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2 \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (-2 \sin(x)) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) - 2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) - 2 \sin(x) \tag{1}$$




Figure 36: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) - 2 \sin(x)$$

Verified OK.

2.3.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2 dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = 4e^{-x} \cos(x)$$

$$(e^{-x}y)'' = 4e^{-x} \cos(x)$$

Integrating once gives

$$(e^{-x}y)' = -2e^{-x}(-\sin(x) + \cos(x)) + c_1$$

Integrating again gives

$$(e^{-x}y) = c_1x - 2e^{-x} \sin(x) + c_2$$

Hence the solution is

$$y = \frac{c_1x - 2e^{-x} \sin(x) + c_2}{e^{-x}}$$

Or

$$y = c_1x e^x + c_2e^x - 2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = c_1x e^x + c_2e^x - 2 \sin(x) \tag{1}$$




Figure 37: Slope field plot

Verification of solutions

$$y = c_1 x e^x + c_2 e^x - 2 \sin(x)$$

Verified OK.

2.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -2 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 31: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1(e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$4 \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 \sin(x) - 2A_2 \cos(x) = 4 \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = -2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2 \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (-2 \sin(x)) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) - 2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) - 2 \sin(x) \tag{1}$$




Figure 38: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) - 2 \sin(x)$$

Verified OK.

2.3.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = 4 \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = e^x x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 x e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 4 \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^x x \\ e^x & e^x x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$
- Substitute functions into equation for $y_p(x)$

$$y_p(x) = 4e^x \left(-\int \cos(x) x e^{-x} dx + x \int e^{-x} \cos(x) dx \right)$$
- Compute integrals

$$y_p(x) = -2 \sin(x)$$
- Substitute particular solution into general solution to ODE

$$y = c_2 x e^x + c_1 e^x - 2 \sin(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=4*cos(x),y(x), singsol=all)
```

$$y(x) = (c_1 x + c_2) e^x - 2 \sin(x)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 21

```
DSolve[y''[x]-2*y'[x]+y[x]==4*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \sin(x) + e^x (c_2 x + c_1)$$

2.4 problem Problem 11.47

2.4.1	Solving as second order linear constant coeff ode	180
2.4.2	Solving as linear second order ode solved by an integrating factor ode	183
2.4.3	Solving using Kovacic algorithm	185
2.4.4	Maple step by step solution	190

Internal problem ID [5178]

Internal file name [OUTPUT/4671_Sunday_June_05_2022_03_03_05_PM_28672195/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.47.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2y' + y = 3e^x$$

2.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = 3e^x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 x e^x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$3 e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[e^x]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^x x\}]$$

Since $e^x x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2 e^x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^x = 3 e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{3}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{3x^2 e^x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + \left(\frac{3x^2 e^x}{2} \right) \end{aligned}$$

Which simplifies to

$$y = e^x (c_2 x + c_1) + \frac{3x^2 e^x}{2}$$

Summary

The solution(s) found are the following

$$y = e^x (c_2 x + c_1) + \frac{3x^2 e^x}{2} \quad (1)$$




Figure 39: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + \frac{3x^2e^x}{2}$$

Verified OK.

2.4.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2 dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = 3e^{-x}e^x$$

$$(e^{-x}y)'' = 3e^{-x}e^x$$

Integrating once gives

$$(e^{-x}y)' = 3x + c_1$$

Integrating again gives

$$(e^{-x}y) = \frac{x(3x + 2c_1)}{2} + c_2$$

Hence the solution is

$$y = \frac{\frac{x(3x+2c_1)}{2} + c_2}{e^{-x}}$$

Or

$$y = c_1x e^x + \frac{3x^2e^x}{2} + c_2e^x$$

Summary

The solution(s) found are the following

$$y = c_1x e^x + \frac{3x^2e^x}{2} + c_2e^x \quad (1)$$




Figure 40: Slope field plot

Verification of solutions

$$y = c_1 x e^x + \frac{3x^2 e^x}{2} + c_2 e^x$$

Verified OK.

2.4.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = y e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 33: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1(e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$3 e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^x x\}]$$

Since $e^x x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2 e^x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^x = 3 e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{3}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{3x^2 e^x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + \left(\frac{3x^2 e^x}{2} \right) \end{aligned}$$

Which simplifies to

$$y = e^x (c_2 x + c_1) + \frac{3x^2 e^x}{2}$$

Summary

The solution(s) found are the following

$$y = e^x (c_2 x + c_1) + \frac{3x^2 e^x}{2} \tag{1}$$




Figure 41: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + \frac{3x^2e^x}{2}$$

Verified OK.

2.4.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = 3e^x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = e^x x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 x e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 3 e^x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^x x \\ e^x & e^x x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -3 e^x \left(\int x dx - \left(\int 1 dx \right) x \right)$$

- Compute integrals

$$y_p(x) = \frac{3x^2 e^x}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_2 x e^x + c_1 e^x + \frac{3x^2 e^x}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=3*exp(x),y(x), singsol=all)
```

$$y(x) = e^x \left(c_2 + c_1 x + \frac{3}{2} x^2 \right)$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 27

```
DSolve[y''[x]-2*y'[x]+y[x]==3*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^x (3x^2 + 2c_2 x + 2c_1)$$

2.5 problem Problem 11.48

2.5.1	Solving as second order linear constant coeff ode	193
2.5.2	Solving as linear second order ode solved by an integrating factor ode	196
2.5.3	Solving using Kovacic algorithm	198
2.5.4	Maple step by step solution	203

Internal problem ID [5179]

Internal file name [OUTPUT/4672_Sunday_June_05_2022_03_03_06_PM_66551782/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.48.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 2y' + y = e^x x$$

2.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = e^x x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 e^x x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x x, e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^x x, x^2 e^x\}]$$

Since $e^x x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^3 e^x, x^2 e^x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^3 e^x + A_2 x^2 e^x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_1 x e^x + 2A_2 e^x = e^x x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{6}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^3 e^x}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + \left(\frac{x^3 e^x}{6} \right) \end{aligned}$$

Which simplifies to

$$y = e^x (c_2 x + c_1) + \frac{x^3 e^x}{6}$$

Summary

The solution(s) found are the following

$$y = e^x (c_2 x + c_1) + \frac{x^3 e^x}{6} \quad (1)$$




Figure 42: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + \frac{x^3e^x}{6}$$

Verified OK.

2.5.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2 dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = e^{-x}e^x x$$

$$(e^{-x}y)'' = e^{-x}e^x x$$

Integrating once gives

$$(e^{-x}y)' = \frac{x^2}{2} + c_1$$

Integrating again gives

$$(e^{-x}y) = \frac{1}{6}x^3 + c_1x + c_2$$

Hence the solution is

$$y = \frac{\frac{1}{6}x^3 + c_1x + c_2}{e^{-x}}$$

Or

$$y = \frac{x^3 e^x}{6} + c_1 x e^x + c_2 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{x^3 e^x}{6} + c_1 x e^x + c_2 e^x \quad (1)$$




Figure 43: Slope field plot

Verification of solutions

$$y = \frac{x^3 e^x}{6} + c_1 x e^x + c_2 e^x$$

Verified OK.

2.5.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 35: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1(e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[{\{e^x x, e^x\}}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, e^x\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{e^x x, x^2 e^x\}}]$$

Since $e^x x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x^3 e^x, x^2 e^x\}}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^3 e^x + A_2 x^2 e^x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_1 x e^x + 2A_2 e^x = e^x x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{6}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^3 e^x}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + \left(\frac{x^3 e^x}{6} \right) \end{aligned}$$

Which simplifies to

$$y = e^x (c_2 x + c_1) + \frac{x^3 e^x}{6}$$

Summary

The solution(s) found are the following

$$y = e^x (c_2 x + c_1) + \frac{x^3 e^x}{6} \tag{1}$$




Figure 44: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + \frac{x^3e^x}{6}$$

Verified OK.

2.5.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = e^x x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = e^x x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 x e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^x x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^x x \\ e^x & e^x x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = e^x \left(- \left(\int x^2 dx \right) + \left(\int x dx \right) x \right)$$

- Compute integrals

$$y_p(x) = \frac{x^3 e^x}{6}$$

- Substitute particular solution into general solution to ODE

$$y = c_2 x e^x + c_1 e^x + \frac{x^3 e^x}{6}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=x*exp(x),y(x), singsol=all)
```

$$y(x) = e^x \left(c_2 + c_1 x + \frac{1}{6} x^3 \right)$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 25

```
DSolve[y''[x]-2*y'[x]+y[x]==x*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6} e^x (x^3 + 6c_2 x + 6c_1)$$

2.6 problem Problem 11.49

2.6.1	Solving as linear ode	206
2.6.2	Solving as first order ode lie symmetry lookup ode	208
2.6.3	Solving as exact ode	212
2.6.4	Maple step by step solution	216

Internal problem ID [5180]

Internal file name [OUTPUT/4673_Sunday_June_05_2022_03_03_07_PM_56470516/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.49.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - y = e^x$$

2.6.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -1$$

$$q(x) = e^x$$

Hence the ode is

$$y' - y = e^x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-1)dx} \\ &= e^{-x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(e^x) \\ \frac{d}{dx}(e^{-x}y) &= (e^{-x})(e^x) \\ d(e^{-x}y) &= dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-x}y &= \int dx \\ e^{-x}y &= x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-x}$ results in

$$y = e^x x + c_1 e^x$$

which simplifies to

$$y = e^x(x + c_1)$$

Summary

The solution(s) found are the following

$$y = e^x(x + c_1) \tag{1}$$




Figure 45: Slope field plot

Verification of solutions

$$y = e^x(x + c_1)$$

Verified OK.

2.6.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = y + e^x$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 37: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^x} dy \end{aligned}$$

Which results in

$$S = e^{-x}y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y + e^x$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -e^{-x}y \\ S_y &= e^{-x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 1 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = R + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$y e^{-x} = x + c_1$$



Which simplifies to

$$y e^{-x} = x + c_1$$

Which gives

$$y = e^x(x + c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = y + e^x$ 	$R = x$ $S = e^{-x}y$	$\frac{dS}{dR} = 1$ 

Summary

The solution(s) found are the following

$$y = e^x(x + c_1) \tag{1}$$




Figure 46: Slope field plot

Verification of solutions

$$y = e^x(x + c_1)$$

Verified OK.

2.6.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}dy &= (y + e^x) dx \\ (-y - e^x) dx + dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y - e^x \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y - e^x) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-1) - (0)) \\ &= -1 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int -1 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-x} \\ &= e^{-x} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{-x}(-y - e^x) \\ &= -e^{-x}y - 1 \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{-x}(1) \\ &= e^{-x} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (-e^{-x}y - 1) + (e^{-x}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -e^{-x}y - 1 dx \\ \phi &= -x + e^{-x}y + f(y)\end{aligned}\tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-x} + f'(y)\tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-x}$. Therefore equation (4) becomes

$$e^{-x} = e^{-x} + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -x + e^{-x}y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -x + e^{-x}y$$

The solution becomes

$$y = e^x(x + c_1)$$

Summary

The solution(s) found are the following

$$y = e^x(x + c_1)\tag{1}$$




Figure 47: Slope field plot

Verification of solutions

$$y = e^x(x + c_1)$$

Verified OK.

2.6.4 Maple step by step solution

Let's solve

$$y' - y = e^x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = y + e^x$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - y = e^x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x)(y' - y) = \mu(x)e^x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y' - y) = \mu'(x)y + \mu(x)y'$$
- Isolate $\mu'(x)$

$$\mu'(x) = -\mu(x)$$
- Solve to find the integrating factor

$$\mu(x) = e^{-x}$$
- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x)e^x dx + c_1$$
- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x)e^x dx + c_1$$
- Solve for y

$$y = \frac{\int \mu(x)e^x dx + c_1}{\mu(x)}$$
- Substitute $\mu(x) = e^{-x}$

$$y = \frac{\int e^{-x}e^x dx + c_1}{e^{-x}}$$
- Evaluate the integrals on the rhs

$$y = \frac{x + c_1}{e^{-x}}$$
- Simplify

$$y = e^x(x + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)-y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = (x + c_1) e^x$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 13

```
DSolve[y'[x]-y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(x + c_1)$$

2.7 problem Problem 11.50

2.7.1	Solving as linear ode	219
2.7.2	Solving as first order ode lie symmetry lookup ode	221
2.7.3	Solving as exact ode	225
2.7.4	Maple step by step solution	229

Internal problem ID [5181]

Internal file name [OUTPUT/4674_Sunday_June_05_2022_03_08_PM_32285306/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.50.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - y = e^{2x}x + 1$$

2.7.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -1$$
$$q(x) = e^{2x}x + 1$$

Hence the ode is

$$y' - y = e^{2x}x + 1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-1)dx} \\ &= e^{-x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(e^{2x}x + 1) \\ \frac{d}{dx}(e^{-x}y) &= (e^{-x})(e^{2x}x + 1) \\ d(e^{-x}y) &= (e^x x + e^{-x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-x}y &= \int e^x x + e^{-x} dx \\ e^{-x}y &= e^x x - e^x - e^{-x} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-x}$ results in

$$y = e^x(e^x x - e^x - e^{-x}) + c_1 e^x$$

which simplifies to

$$y = (x - 1)e^{2x} + c_1 e^x - 1$$

Summary

The solution(s) found are the following

$$y = (x - 1)e^{2x} + c_1 e^x - 1 \tag{1}$$




Figure 48: Slope field plot

Verification of solutions

$$y = (x - 1)e^{2x} + c_1e^x - 1$$

Verified OK.

2.7.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = y + e^{2x}x + 1$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 40: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^x} dy \end{aligned}$$

Which results in

$$S = e^{-x}y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y + e^{2x}x + 1$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -e^{-x}y \\ S_y &= e^{-x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^x x + e^{-x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^R R + e^{-R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = e^R R - e^R - e^{-R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$y e^{-x} = e^x x - e^x - e^{-x} + c_1$$



Which simplifies to

$$y e^{-x} = e^x x - e^x - e^{-x} + c_1$$

Which gives

$$y = (e^x x - e^x - e^{-x} + c_1) e^x$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = y + e^{2x}x + 1$ 	$R = x$ $S = e^{-x}y$	$\frac{dS}{dR} = e^R R + e^{-R}$ 

Summary

The solution(s) found are the following

$$y = (e^x x - e^x - e^{-x} + c_1) e^x \quad (1)$$




Figure 49: Slope field plot

Verification of solutions

$$y = (e^x x - e^x - e^{-x} + c_1) e^x$$

Verified OK.

2.7.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (y + e^{2x}x + 1) dx \\ (-y - e^{2x}x - 1) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y - e^{2x}x - 1 \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y - e^{2x}x - 1) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-1) - (0)) \\ &= -1 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int -1 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-x} \\ &= e^{-x} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{-x}(-y - e^{2x}x - 1) \\ &= (-y - 1)e^{-x} - e^x x \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{-x}(1) \\ &= e^{-x} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((-y - 1)e^{-x} - e^x x) + (e^{-x}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (-y - 1)e^{-x} - e^x x dx \\ \phi &= -e^x x + e^{-x} y + e^x + e^{-x} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-x}$. Therefore equation (4) becomes

$$e^{-x} = e^{-x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -e^x x + e^{-x} y + e^x + e^{-x} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -e^x x + e^{-x} y + e^x + e^{-x}$$

The solution becomes

$$y = (e^x x - e^x - e^{-x} + c_1) e^x$$

Summary

The solution(s) found are the following

$$y = (e^x x - e^x - e^{-x} + c_1) e^x \quad (1)$$




Figure 50: Slope field plot

Verification of solutions

$$y = (e^x x - e^x - e^{-x} + c_1) e^x$$

Verified OK.

2.7.4 Maple step by step solution

Let's solve

$$y' - y = e^{2x} x + 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = y + e^{2x} x + 1$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - y = e^{2x} x + 1$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - y) = \mu(x) (e^{2x} x + 1)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y' - y) = \mu'(x)y + \mu(x)y'$$
- Isolate $\mu'(x)$

$$\mu'(x) = -\mu(x)$$
- Solve to find the integrating factor

$$\mu(x) = e^{-x}$$
- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x)(e^{2x}x + 1) dx + c_1$$
- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x)(e^{2x}x + 1) dx + c_1$$
- Solve for y

$$y = \frac{\int \mu(x)(e^{2x}x + 1) dx + c_1}{\mu(x)}$$
- Substitute $\mu(x) = e^{-x}$

$$y = \frac{\int (e^{2x}x + 1)e^{-x} dx + c_1}{e^{-x}}$$
- Evaluate the integrals on the rhs

$$y = \frac{e^x x - e^x - \frac{1}{e^x} + c_1}{e^{-x}}$$
- Simplify

$$y = (x - 1)e^{2x} + c_1 e^x - 1$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)-y(x)=x*exp(2*x)+1,y(x), singsol=all)
```

$$y(x) = (x - 1)e^{2x} + e^x c_1 - 1$$

✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 22

```
DSolve[y'[x]-y[x]==x*Exp[2*x]+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(x - 1) + c_1 e^x - 1$$

2.8 problem Problem 11.51

2.8.1	Solving as linear ode	232
2.8.2	Solving as first order ode lie symmetry lookup ode	234
2.8.3	Solving as exact ode	239
2.8.4	Maple step by step solution	243

Internal problem ID [5182]

Internal file name [OUTPUT/4675_Sunday_June_05_2022_03_03_09_PM_21862982/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.51.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - y = \sin(x) + \cos(2x)$$

2.8.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -1$$

$$q(x) = \sin(x) + \cos(2x)$$

Hence the ode is

$$y' - y = \sin(x) + \cos(2x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int(-1)dx} \\ &= e^{-x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (\sin(x) + \cos(2x)) \\ \frac{d}{dx}(e^{-x}y) &= (e^{-x}) (\sin(x) + \cos(2x)) \\ d(e^{-x}y) &= ((\sin(x) + \cos(2x)) e^{-x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-x}y &= \int (\sin(x) + \cos(2x)) e^{-x} dx \\ e^{-x}y &= -\frac{e^{-x} \cos(x)}{2} - \frac{e^{-x} \sin(x)}{2} + \frac{2(-\cos(x) + 2 \sin(x)) e^{-x} \cos(x)}{5} + \frac{e^{-x}}{5} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-x}$ results in

$$y = e^x \left(-\frac{e^{-x} \cos(x)}{2} - \frac{e^{-x} \sin(x)}{2} + \frac{2(-\cos(x) + 2 \sin(x)) e^{-x} \cos(x)}{5} + \frac{e^{-x}}{5} \right) + c_1 e^x$$

which simplifies to

$$y = -\frac{2 \cos(x)^2}{5} + \frac{(8 \sin(x) - 5) \cos(x)}{10} + c_1 e^x - \frac{\sin(x)}{2} + \frac{1}{5}$$

Summary

The solution(s) found are the following

$$y = -\frac{2 \cos(x)^2}{5} + \frac{(8 \sin(x) - 5) \cos(x)}{10} + c_1 e^x - \frac{\sin(x)}{2} + \frac{1}{5} \quad (1)$$




Figure 51: Slope field plot

Verification of solutions

$$y = -\frac{2 \cos(x)^2}{5} + \frac{(8 \sin(x) - 5) \cos(x)}{10} + c_1 e^x - \frac{\sin(x)}{2} + \frac{1}{5}$$

Verified OK.

2.8.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} y' &= y + \sin(x) + \cos(2x) \\ y' &= \omega(x, y) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 43: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^x} dy \end{aligned}$$

Which results in

$$S = e^{-x}y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y + \sin(x) + \cos(2x)$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -e^{-x}y \\ S_y &= e^{-x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = (\sin(x) + \cos(2x)) e^{-x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = (\sin(R) + \cos(2R)) e^{-R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{e^{-R}}{5} + c_1 - \frac{e^{-R}(5 \cos(R) + 5 \sin(R) + 2 \cos(2R) + 2 - 4 \sin(2R))}{10} \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$y e^{-x} = \frac{e^{-x}}{5} + c_1 - \frac{e^{-x}(-4 \sin(2x) + 2 \cos(2x) + 2 + 5 \sin(x) + 5 \cos(x))}{10}$$



Which simplifies to

$$\frac{(4 \cos(x))^2 + (-8 \sin(x) + 5) \cos(x) + 10y + 5 \sin(x) - 2) e^{-x}}{10} - c_1 = 0$$

Which gives

$$y = \frac{(8 e^{-x} \sin(x) \cos(x) - 4 e^{-x} \cos(x))^2 - 5 e^{-x} \sin(x) - 5 e^{-x} \cos(x) + 2 e^{-x} + 10c_1) e^x}{10}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = y + \sin(x) + \cos(2x)$ 	$R = x$ $S = e^{-x}y$	$\frac{dS}{dR} = (\sin(R) + \cos(2R)) e^{-R}$ 

Summary

The solution(s) found are the following

$$y = \frac{(8 e^{-x} \sin(x) \cos(x) - 4 e^{-x} \cos(x)^2 - 5 e^{-x} \sin(x) - 5 e^{-x} \cos(x) + 2 e^{-x} + 10c_1) e^x}{10} \quad (1)$$




Figure 52: Slope field plot

Verification of solutions

$$y = \frac{(8 e^{-x} \sin(x) \cos(x) - 4 e^{-x} \cos(x)^2 - 5 e^{-x} \sin(x) - 5 e^{-x} \cos(x) + 2 e^{-x} + 10c_1) e^x}{10}$$

Verified OK.

2.8.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= (y + \sin(x) + \cos(2x)) dx \\ (-y - \sin(x) - \cos(2x)) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -y - \sin(x) - \cos(2x) \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y - \sin(x) - \cos(2x)) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-1) - (0)) \\ &= -1\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int -1 dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-x} \\ &= e^{-x}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\overline{M} &= \mu M \\ &= e^{-x}(-y - \sin(x) - \cos(2x)) \\ &= -e^{-x}(y + \sin(x) + \cos(2x))\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{-x}(1) \\ &= e^{-x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (-e^{-x}(y + \sin(x) + \cos(2x))) + (e^{-x}) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int \bar{M} dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -e^{-x}(y + \sin(x) + \cos(2x)) dx$$

$$\phi = \frac{(4 \cos(x)^2 + (-8 \sin(x) + 5) \cos(x) + 10y + 5 \sin(x) - 2) e^{-x}}{10} + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-x} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-x}$. Therefore equation (4) becomes

$$e^{-x} = e^{-x} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(4 \cos(x)^2 + (-8 \sin(x) + 5) \cos(x) + 10y + 5 \sin(x) - 2) e^{-x}}{10} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(4 \cos(x)^2 + (-8 \sin(x) + 5) \cos(x) + 10y + 5 \sin(x) - 2) e^{-x}}{10}$$

The solution becomes

$$y = \frac{(8 e^{-x} \sin(x) \cos(x) - 4 e^{-x} \cos(x)^2 - 5 e^{-x} \sin(x) - 5 e^{-x} \cos(x) + 2 e^{-x} + 10c_1) e^x}{10}$$

Summary

The solution(s) found are the following

$$y = \frac{(8 e^{-x} \sin(x) \cos(x) - 4 e^{-x} \cos(x)^2 - 5 e^{-x} \sin(x) - 5 e^{-x} \cos(x) + 2 e^{-x} + 10c_1) e^x}{10} \quad (1)$$




Figure 53: Slope field plot

Verification of solutions

$$y = \frac{(8e^{-x} \sin(x) \cos(x) - 4e^{-x} \cos(x)^2 - 5e^{-x} \sin(x) - 5e^{-x} \cos(x) + 2e^{-x} + 10c_1) e^x}{10}$$

Verified OK.

2.8.4 Maple step by step solution

Let's solve

$$y' - y = \sin(x) + \cos(2x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = y + \sin(x) + \cos(2x)$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - y = \sin(x) + \cos(2x)$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - y) = \mu(x) (\sin(x) + \cos(2x))$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' - y) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{-x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) (\sin(x) + \cos(2x)) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) (\sin(x) + \cos(2x)) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x)(\sin(x)+\cos(2x))dx+c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-x}$

$$y = \frac{\int (\sin(x)+\cos(2x))e^{-x} dx+c_1}{e^{-x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{-\frac{e^{-x} \cos(x)}{2} - \frac{e^{-x} \sin(x)}{2} + \frac{2(-\cos(x)+2\sin(x))e^{-x} \cos(x)}{5} + \frac{1}{5e^x} + c_1}{e^{-x}}$$

- Simplify

$$y = -\frac{2\cos(x)^2}{5} + \frac{(8\sin(x)-5)\cos(x)}{10} + c_1 e^x - \frac{\sin(x)}{2} + \frac{1}{5}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x)-y(x)=sin(x)+cos(2*x),y(x), singsol=all)
```

$$y(x) = e^x c_1 - \frac{\cos(x)}{2} - \frac{\sin(x)}{2} + \frac{2 \sin(2x)}{5} - \frac{\cos(2x)}{5}$$

✓ Solution by Mathematica

Time used: 0.16 (sec). Leaf size: 37

```
DSolve[y'[x]-y[x]==Sin[x]+Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{10}(-5 \sin(x) + 4 \sin(2x) - 5 \cos(x) - 2 \cos(2x) + 10c_1 e^x)$$

2.9 problem Problem 11.52

Internal problem ID [5183]

Internal file name [OUTPUT/4676_Sunday_June_05_2022_03_03_10_PM_88926433/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 11. THE METHOD OF UNDETERMINED COEFFICIENTS. Supplementary Problems. page 101

Problem number: Problem 11.52.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

$$y''' - 3y'' + 3y' - y = 1 + e^x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' - 3y'' + 3y' - y = 0$$

The characteristic equation is

$$\lambda^3 - 3\lambda^2 + 3\lambda - 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = 1$$

$$\lambda_3 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^x + c_2 x e^x + x^2 e^x c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^x$$

$$y_2 = e^x x$$

$$y_3 = x^2 e^x$$

Now the particular solution to the given ODE is found

$$y''' - 3y'' + 3y' - y = 1 + e^x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x x, x^2 e^x, e^x\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{1\}, \{e^x x\}]$$

Since $e^x x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{1\}, \{x^2 e^x\}]$$

Since $x^2 e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{1\}, \{x^3 e^x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 + A_2 x^3 e^x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_2e^x - A_1 = 1 + e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -1, A_2 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -1 + \frac{x^3e^x}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^x + c_2xe^x + x^2e^xc_3) + \left(-1 + \frac{x^3e^x}{6} \right) \end{aligned}$$

Which simplifies to

$$y = e^x(c_3x^2 + c_2x + c_1) - 1 + \frac{x^3e^x}{6}$$

Summary

The solution(s) found are the following

$$y = e^x(c_3x^2 + c_2x + c_1) - 1 + \frac{x^3e^x}{6} \quad (1)$$

Verification of solutions

$$y = e^x(c_3x^2 + c_2x + c_1) - 1 + \frac{x^3e^x}{6}$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 3; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)+3*diff(y(x),x)-y(x)=exp(x)+1,y(x), singsol=all)
```

$$y(x) = -1 + \frac{(6c_3x^2 + x^3 + 6c_2x + 6c_1) e^x}{6}$$

✓ Solution by Mathematica

Time used: 0.067 (sec). Leaf size: 31

```
DSolve[y'''[x]-3*y''[x]+3*y'[x]-y[x]==Exp[x]+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1 + e^x \left(\frac{x^3}{6} + c_3x^2 + c_2x + c_1 \right)$$

3 Chapter 12. VARIATION OF PARAMETERS.

page 104

3.1	problem Problem 12.1	251
3.2	problem Problem 12.2	261
3.3	problem Problem 12.3	267
3.4	problem Problem 12.4	281
3.5	problem Problem 12.5	292
3.6	problem Problem 12.6	303
3.7	problem Problem 12.7	340
3.8	problem Problem 12.8	353

3.1 problem Problem 12.1

3.1.1 Maple step by step solution 255

Internal problem ID [5184]

Internal file name [OUTPUT/4677_Sunday_June_05_2022_03_03_11_PM_55752686/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. page 104

Problem number: Problem 12.1.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_y]]
```

$$y''' + y' = \sec(x)$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' + y' = 0$$

The characteristic equation is

$$\lambda^3 + \lambda = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = i$$

$$\lambda_3 = -i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 + e^{-ix}c_2 + e^{ix}c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= 1 \\y_2 &= e^{-ix} \\y_3 &= e^{ix}\end{aligned}$$

Now the particular solution to the given ODE is found

$$y''' + y' = \sec(x)$$

Let the particular solution be

$$y_p = U_1y_1 + U_2y_2 + U_3y_3$$

Where y_i are the basis solutions found above for the homogeneous solution y_h and $U_i(x)$ are functions to be determined as follows

$$U_i = (-1)^{n-i} \int \frac{F(x)W_i(x)}{aW(x)} dx$$

Where $W(x)$ is the Wronskian and $W_i(x)$ is the Wronskian that results after deleting the last row and the i -th column of the determinant and n is the order of the ODE or equivalently, the number of basis solutions, and a is the coefficient of the leading derivative in the ODE, and $F(x)$ is the RHS of the ODE. Therefore, the first step is to find the Wronskian $W(x)$. This is given by

$$W(x) = \begin{vmatrix} y_1 & y_2 & y_3 \\ y_1' & y_2' & y_3' \\ y_1'' & y_2'' & y_3'' \end{vmatrix}$$

Substituting the fundamental set of solutions y_i found above in the Wronskian gives

$$\begin{aligned}W &= \begin{bmatrix} 1 & e^{-ix} & e^{ix} \\ 0 & -ie^{-ix} & ie^{ix} \\ 0 & -e^{-ix} & -e^{ix} \end{bmatrix} \\|W| &= 2ie^{-ix}e^{ix}\end{aligned}$$

The determinant simplifies to

$$|W| = 2i$$

Now we determine W_i for each U_i .

$$\begin{aligned} W_1(x) &= \det \begin{bmatrix} e^{-ix} & e^{ix} \\ -ie^{-ix} & ie^{ix} \end{bmatrix} \\ &= 2i \end{aligned}$$

$$\begin{aligned} W_2(x) &= \det \begin{bmatrix} 1 & e^{ix} \\ 0 & ie^{ix} \end{bmatrix} \\ &= ie^{ix} \end{aligned}$$

$$\begin{aligned} W_3(x) &= \det \begin{bmatrix} 1 & e^{-ix} \\ 0 & -ie^{-ix} \end{bmatrix} \\ &= -ie^{-ix} \end{aligned}$$

Now we are ready to evaluate each $U_i(x)$.

$$\begin{aligned} U_1 &= (-1)^{3-1} \int \frac{F(x)W_1(x)}{aW(x)} dx \\ &= (-1)^2 \int \frac{(\sec(x))(2i)}{(1)(2i)} dx \\ &= \int \frac{2i \sec(x)}{2i} dx \\ &= \int (\sec(x)) dx \\ &= \ln(\sec(x) + \tan(x)) \end{aligned}$$

$$\begin{aligned} U_2 &= (-1)^{3-2} \int \frac{F(x)W_2(x)}{aW(x)} dx \\ &= (-1)^1 \int \frac{(\sec(x))(ie^{ix})}{(1)(2i)} dx \\ &= - \int \frac{i \sec(x) e^{ix}}{2i} dx \\ &= - \int \left(\frac{\sec(x) e^{ix}}{2} \right) dx \\ &= \frac{i \ln(e^{2ix} + 1)}{2} \end{aligned}$$

$$\begin{aligned}
U_3 &= (-1)^{3-3} \int \frac{F(x)W_3(x)}{aW(x)} dx \\
&= (-1)^0 \int \frac{(\sec(x))(-ie^{-ix})}{(1)(2i)} dx \\
&= \int \frac{-i \sec(x) e^{-ix}}{2i} dx \\
&= \int \left(-\frac{\sec(x) e^{-ix}}{2} \right) dx \\
&= -\frac{i \ln(e^{2ix} + 1)}{2} + i \ln(e^{ix})
\end{aligned}$$

Now that all the U_i functions have been determined, the particular solution is found from

$$y_p = U_1 y_1 + U_2 y_2 + U_3 y_3$$

Hence

$$\begin{aligned}
y_p &= (\ln(\sec(x) + \tan(x))) \\
&+ \left(\frac{i \ln(e^{2ix} + 1)}{2} \right) (e^{-ix}) \\
&+ \left(-\frac{i \ln(e^{2ix} + 1)}{2} + i \ln(e^{ix}) \right) (e^{ix})
\end{aligned}$$

Therefore the particular solution is

$$y_p = \frac{i(-e^{ix} + e^{-ix}) \ln(e^{2ix} + 1)}{2} + ie^{ix} \ln(e^{ix}) + \ln(\sec(x) + \tan(x))$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 + e^{-ix} c_2 + e^{ix} c_3) + \left(\frac{i(-e^{ix} + e^{-ix}) \ln(e^{2ix} + 1)}{2} + ie^{ix} \ln(e^{ix}) + \ln(\sec(x) + \tan(x)) \right)$$

Summary

The solution(s) found are the following

$$\begin{aligned}
y &= c_1 + e^{-ix} c_2 + e^{ix} c_3 + \frac{i(-e^{ix} + e^{-ix}) \ln(e^{2ix} + 1)}{2} \\
&+ ie^{ix} \ln(e^{ix}) + \ln(\sec(x) + \tan(x))
\end{aligned} \tag{1}$$

Verification of solutions

$$y = c_1 + e^{-ix} c_2 + e^{ix} c_3 + \frac{i(-e^{ix} + e^{-ix}) \ln(e^{2ix} + 1)}{2} + ie^{ix} \ln(e^{ix}) + \ln(\sec(x) + \tan(x))$$

Verified OK.

3.1.1 Maple step by step solution

Let's solve

$$y''' + y' = \sec(x)$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = \sec(x) - y_2(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = \sec(x) - y_2(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ \sec(x) \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ \sec(x) \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right], \left[-I, \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix} \right], \left[I, \begin{bmatrix} -1 \\ -I \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{y}_2(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_3(x) = \begin{bmatrix} \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix} \right]$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$
 $\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \vec{y}_p(x)$

□ Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} 1 & -\cos(x) & \sin(x) \\ 0 & \sin(x) & \cos(x) \\ 0 & \cos(x) & -\sin(x) \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix
 $\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} 1 & -\cos(x) & \sin(x) \\ 0 & \sin(x) & \cos(x) \\ 0 & \cos(x) & -\sin(x) \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} 1 & \sin(x) & 1 - \cos(x) \\ 0 & \cos(x) & \sin(x) \\ 0 & -\sin(x) & \cos(x) \end{bmatrix}$$

□ Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} -\sin(x) \left(\int_0^x \tan(s) ds \right) - \cos(x)x + \int_0^x (\sec(s) - 1) ds + x \\ -\cos(x) \left(\int_0^x \tan(s) ds \right) + \sin(x)x \\ \sin(x) \left(\int_0^x \tan(s) ds \right) + \cos(x)x \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \begin{bmatrix} -\sin(x) \left(\int_0^x \tan(s) ds \right) - \cos(x) x + \int_0^x (\sec(s) - 1) ds \\ -\cos(x) \left(\int_0^x \tan(s) ds \right) + \sin(x) x \\ \sin(x) \left(\int_0^x \tan(s) ds \right) + \cos(x) x \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \int_0^x (\sec(s) - 1) ds - \sin(x) \left(\int_0^x \tan(s) ds \right) + (-c_2 - x) \cos(x) + c_3 \sin(x) + x + c_1$$

Maple trace

```

Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
-> Calling odsolve with the ODE`, diff(diff(_b(_a), _a), _a) = -_b(_a)+sec(_a), _b(_a)`
Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
    <- solving first the homogeneous part of the ODE successful
<- differential order: 3; linear nonhomogeneous with symmetry [0,1] successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 83

```
dsolve(diff(y(x),x$3)+diff(y(x),x)=sec(x),y(x), singsol=all)
```

$$y(x) = \frac{i(e^{ix} - e^{-ix}) \ln\left(\frac{e^{ix}}{e^{2ix}+1}\right)}{2} - \frac{ie^{-ix}}{2} - 2i \arctan(e^{ix}) + \frac{ie^{ix}}{2} + (1 + c_1 - \ln(2)) \sin(x) + (-x - c_2) \cos(x) + c_3$$

✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 57

```
DSolve[y'''[x]+y'[x]==Sec[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -(x + c_2) \cos(x) - \log\left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right) \\ + \log\left(\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)\right) + \sin(x)(\log(\cos(x)) + c_1) + c_3$$

3.2 problem Problem 12.2

Internal problem ID [5185]

Internal file name [OUTPUT/4678_Sunday_June_05_2022_03_03_13_PM_87732597/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. page 104

Problem number: Problem 12.2.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_y]]
```

$$y''' - 3y'' + 2y' = \frac{e^x}{1 + e^{-x}}$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' - 3y'' + 2y' = 0$$

The characteristic equation is

$$\lambda^3 - 3\lambda^2 + 2\lambda = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 2$$

$$\lambda_3 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 + c_2 e^x + e^{2x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned} y_1 &= 1 \\ y_2 &= e^x \\ y_3 &= e^{2x} \end{aligned}$$

Now the particular solution to the given ODE is found

$$y''' - 3y'' + 2y' = \frac{e^x}{1 + e^{-x}}$$

Let the particular solution be

$$y_p = U_1 y_1 + U_2 y_2 + U_3 y_3$$

Where y_i are the basis solutions found above for the homogeneous solution y_h and $U_i(x)$ are functions to be determined as follows

$$U_i = (-1)^{n-i} \int \frac{F(x)W_i(x)}{aW(x)} dx$$

Where $W(x)$ is the Wronskian and $W_i(x)$ is the Wronskian that results after deleting the last row and the i -th column of the determinant and n is the order of the ODE or equivalently, the number of basis solutions, and a is the coefficient of the leading derivative in the ODE, and $F(x)$ is the RHS of the ODE. Therefore, the first step is to find the Wronskian $W(x)$. This is given by

$$W(x) = \begin{vmatrix} y_1 & y_2 & y_3 \\ y_1' & y_2' & y_3' \\ y_1'' & y_2'' & y_3'' \end{vmatrix}$$

Substituting the fundamental set of solutions y_i found above in the Wronskian gives

$$W = \begin{bmatrix} 1 & e^x & e^{2x} \\ 0 & e^x & 2e^{2x} \\ 0 & e^x & 4e^{2x} \end{bmatrix}$$

$$|W| = 2e^x e^{2x}$$

The determinant simplifies to

$$|W| = 2 e^{3x}$$

Now we determine W_i for each U_i .

$$\begin{aligned} W_1(x) &= \det \begin{bmatrix} e^x & e^{2x} \\ e^x & 2e^{2x} \end{bmatrix} \\ &= e^{3x} \end{aligned}$$

$$\begin{aligned} W_2(x) &= \det \begin{bmatrix} 1 & e^{2x} \\ 0 & 2e^{2x} \end{bmatrix} \\ &= 2 e^{2x} \end{aligned}$$

$$\begin{aligned} W_3(x) &= \det \begin{bmatrix} 1 & e^x \\ 0 & e^x \end{bmatrix} \\ &= e^x \end{aligned}$$

Now we are ready to evaluate each $U_i(x)$.

$$\begin{aligned} U_1 &= (-1)^{3-1} \int \frac{F(x)W_1(x)}{aW(x)} dx \\ &= (-1)^2 \int \frac{\left(\frac{e^x}{1+e^{-x}}\right) (e^{3x})}{(1)(2e^{3x})} dx \\ &= \int \frac{\frac{e^x e^{3x}}{1+e^{-x}}}{2e^{3x}} dx \\ &= \int \left(\frac{e^{2x}}{2e^x + 2} \right) dx \\ &= \frac{e^x}{2} - \frac{\ln(1+e^x)}{2} \end{aligned}$$

$$\begin{aligned}
U_2 &= (-1)^{3-2} \int \frac{F(x)W_2(x)}{aW(x)} dx \\
&= (-1)^1 \int \frac{\left(\frac{e^x}{1+e^{-x}}\right) (2e^{2x})}{(1)(2e^{3x})} dx \\
&= - \int \frac{\frac{2e^xe^{2x}}{1+e^{-x}}}{2e^{3x}} dx \\
&= - \int \left(\frac{1}{1+e^{-x}}\right) dx \\
&= -\ln(1+e^{-x}) + \ln(e^{-x})
\end{aligned}$$

$$\begin{aligned}
U_3 &= (-1)^{3-3} \int \frac{F(x)W_3(x)}{aW(x)} dx \\
&= (-1)^0 \int \frac{\left(\frac{e^x}{1+e^{-x}}\right) (e^x)}{(1)(2e^{3x})} dx \\
&= \int \frac{\frac{e^{2x}}{1+e^{-x}}}{2e^{3x}} dx \\
&= \int \left(\frac{e^{-x}}{2e^{-x}+2}\right) dx \\
&= -\frac{\ln(1+e^{-x})}{2}
\end{aligned}$$

Now that all the U_i functions have been determined, the particular solution is found from

$$y_p = U_1y_1 + U_2y_2 + U_3y_3$$

Hence

$$\begin{aligned}
y_p &= \left(\frac{e^x}{2} - \frac{\ln(1+e^x)}{2}\right) \\
&\quad + (-\ln(1+e^{-x}) + \ln(e^{-x})) (e^x) \\
&\quad + \left(-\frac{\ln(1+e^{-x})}{2}\right) (e^{2x})
\end{aligned}$$

Therefore the particular solution is

$$y_p = \frac{(-2e^x - e^{2x}) \ln(1+e^{-x})}{2} + e^x \ln(e^{-x}) + \frac{e^x}{2} - \frac{\ln(1+e^x)}{2}$$

Therefore the general solution is

$$\begin{aligned}
y &= y_h + y_p \\
&= (c_1 + c_2e^x + e^{2x}c_3) + \left(\frac{(-2e^x - e^{2x}) \ln(1+e^{-x})}{2} + e^x \ln(e^{-x}) + \frac{e^x}{2} - \frac{\ln(1+e^x)}{2}\right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2 e^x + e^{2x} c_3 + \frac{(-2e^x - e^{2x}) \ln(1 + e^{-x})}{2} + e^x \ln(e^{-x}) + \frac{e^x}{2} - \frac{\ln(1 + e^x)}{2} \quad (1)$$

Verification of solutions

$$y = c_1 + c_2 e^x + e^{2x} c_3 + \frac{(-2e^x - e^{2x}) \ln(1 + e^{-x})}{2} + e^x \ln(e^{-x}) + \frac{e^x}{2} - \frac{\ln(1 + e^x)}{2}$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
-> Calling odsolve with the ODE`, diff(diff(b(a), a), a) = -(2*b(a)*exp(-a)-3*(diff(
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying high order exact linear fully integrable
  trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
  trying a double symmetry of the form [xi=0, eta=F(x)]
  <- double symmetry of the form [xi=0, eta=F(x)] successful
  <- differential order: 3; linear nonhomogeneous with symmetry [0,1] successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 56

```
dsolve(diff(y(x), x$3)-3*diff(y(x), x$2)+2*diff(y(x), x)=exp(x)/(1+exp(-x)), y(x), singsol=all)
```

$$y(x) = \frac{(-2e^x - e^{2x} - 1) \ln(1 + e^{-x})}{2} + \frac{(2e^x + 1) \ln(e^{-x})}{2} + \frac{e^{2x} c_1}{2} + \frac{(2c_2 + 1) e^x}{2} + c_3$$

✓ Solution by Mathematica

Time used: 0.137 (sec). Leaf size: 59

```
DSolve[y'''[x]-3*y''[x]+2*y'[x]==Exp[x]/(1+Exp[-x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(-2e^{2x} \operatorname{arctanh}(2e^x + 1) - (2e^x + 1) \log(e^x + 1) + e^x(c_2 e^x + 1 + 2c_1)) + c_3$$

3.3 problem Problem 12.3

3.3.1	Solving as second order linear constant coeff ode	267
3.3.2	Solving as linear second order ode solved by an integrating factor ode	271
3.3.3	Solving using Kovacic algorithm	273
3.3.4	Maple step by step solution	279

Internal problem ID [5186]

Internal file name [OUTPUT/4679_Sunday_June_05_2022_03_03_14_PM_9585856/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. page 104

Problem number: Problem 12.3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 2y' + y = \frac{e^x}{x}$$

3.3.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = \frac{e^x}{x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 e^x x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^x \\ y_2 &= e^x x \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^x & e^x x \\ \frac{d}{dx}(e^x) & \frac{d}{dx}(e^x x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^x & e^x x \\ e^x & e^x x + e^x \end{vmatrix}$$

Therefore

$$W = (e^x)(e^x x + e^x) - (e^x x)(e^x)$$

Which simplifies to

$$W = e^{2x}$$

Which simplifies to

$$W = e^{2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{2x}}{e^{2x}} dx$$

Which simplifies to

$$u_1 = - \int 1 dx$$

Hence

$$u_1 = -x$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{2x}}{e^{2x}} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x} dx$$

Hence

$$u_2 = \ln(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -e^x x + \ln(x) e^x x$$

Which simplifies to

$$y_p(x) = e^x x(\ln(x) - 1)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (e^x x(\ln(x) - 1)) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) + e^x x(\ln(x) - 1)$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) + e^x x(\ln(x) - 1) \tag{1}$$




Figure 54: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + e^x x(\ln(x) - 1)$$

Verified OK.

3.3.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2 dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = \frac{e^{-x}e^x}{x}$$
$$(e^{-x}y)'' = \frac{e^{-x}e^x}{x}$$

Integrating once gives

$$(e^{-x}y)' = \ln(x) + c_1$$

Integrating again gives

$$(e^{-x}y) = x(\ln(x) + c_1 - 1) + c_2$$

Hence the solution is

$$y = \frac{x(\ln(x) + c_1 - 1) + c_2}{e^{-x}}$$

Or

$$y = c_1 x e^x + x e^x \ln(x) + c_2 e^x - x e^x$$

Summary

The solution(s) found are the following

$$y = c_1 x e^x + x e^x \ln(x) + c_2 e^x - x e^x \tag{1}$$




Figure 55: Slope field plot

Verification of solutions

$$y = c_1 x e^x + x e^x \ln(x) + c_2 e^x - x e^x$$

Verified OK.

3.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = y e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 47: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1(e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^x$$

$$y_2 = x e^x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^x & x e^x \\ \frac{d}{dx}(e^x) & \frac{d}{dx}(x e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^x & x e^x \\ e^x & x e^x + e^x \end{vmatrix}$$

Therefore

$$W = (e^x)(x e^x + e^x) - (x e^x)(e^x)$$

Which simplifies to

$$W = e^{2x}$$

Which simplifies to

$$W = e^{2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{2x}}{e^{2x}} dx$$

Which simplifies to

$$u_1 = - \int 1 dx$$

Hence

$$u_1 = -x$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{e^{2x}}{x}}{e^{2x}} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x} dx$$

Hence

$$u_2 = \ln(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -x e^x + x e^x \ln(x)$$

Which simplifies to

$$y_p(x) = e^x x (\ln(x) - 1)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (e^x x (\ln(x) - 1)) \end{aligned}$$

Which simplifies to

$$y = e^x (c_2 x + c_1) + e^x x (\ln(x) - 1)$$

Summary

The solution(s) found are the following

$$y = e^x (c_2 x + c_1) + e^x x (\ln(x) - 1) \tag{1}$$




Figure 56: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + e^x x(\ln(x) - 1)$$

Verified OK.

3.3.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = \frac{e^x}{x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 x e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{e^x}{x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & x e^x \\ e^x & x e^x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$
- Substitute functions into equation for $y_p(x)$

$$y_p(x) = e^x \left(-\int 1 dx + \left(\int \frac{1}{x} dx \right) x \right)$$
- Compute integrals

$$y_p(x) = e^x x (\ln(x) - 1)$$
- Substitute particular solution into general solution to ODE

$$y = c_1 e^x + c_2 x e^x + e^x x (\ln(x) - 1)$$

Maple trace

```

Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=exp(x)/x,y(x), singsol=all)
```

$$y(x) = (\ln(x) x + x(c_1 - 1) + c_2) e^x$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 22

```
DSolve[y''[x]-2*y'[x]+y[x]==Exp[x]/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x (x \log(x) + (-1 + c_2)x + c_1)$$

3.4 problem Problem 12.4

3.4.1	Solving as second order linear constant coeff ode	281
3.4.2	Solving using Kovacic algorithm	284
3.4.3	Maple step by step solution	289

Internal problem ID [5187]

Internal file name [OUTPUT/4680_Sunday_June_05_2022_03_03_15_PM_92094135/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. page 104

Problem number: Problem 12.4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - 2y = e^{3x}$$

3.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -2, f(x) = e^{3x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-2)} \\ &= \frac{1}{2} \pm \frac{3}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} + \frac{3}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{3}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1 e^{3x} = e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{3x}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^{-x}) + \left(\frac{e^{3x}}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-x} + \frac{e^{3x}}{4} \quad (1)$$




Figure 57: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-x} + \frac{e^{3x}}{4}$$

Verified OK.

3.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -1 \tag{3}$$

$$C = -2$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 49: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\ &= z_1 e^{\frac{x}{2}} \\ &= z_1 \left(e^{\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{3x}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{3x}}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^{2x}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{2x}}{3}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1e^{3x} = e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{3x}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{-x} + \frac{c_2e^{2x}}{3} \right) + \left(\frac{e^{3x}}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-x} + \frac{c_2e^{2x}}{3} + \frac{e^{3x}}{4} \quad (1)$$




Figure 58: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} + \frac{e^{3x}}{4}$$

Verified OK.

3.4.3 Maple step by step solution

Let's solve

$$y'' - y' - 2y = e^{3x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{3x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{2x} \\ -e^{-x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x}(\int e^{4x} dx)}{3} + \frac{e^{2x}(\int e^x dx)}{3}$$

- Compute integrals

$$y_p(x) = \frac{e^{3x}}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + \frac{e^{3x}}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=exp(3*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{2x} c_1 + \frac{e^{3x}}{4}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 31

```
DSolve[y''[x]-y'[x]-2*y[x]==Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{3x}}{4} + c_1 e^{-x} + c_2 e^{2x}$$

3.5 problem Problem 12.5

3.5.1	Solving as second order linear constant coeff ode	292
3.5.2	Solving using Kovacic algorithm	295
3.5.3	Maple step by step solution	300

Internal problem ID [5188]

Internal file name [OUTPUT/4681_Sunday_June_05_2022_03_03_16_PM_90074113/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. page 104

Problem number: Problem 12.5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$x'' + 4x = \sin(2t)^2$$

3.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = f(t)$$

Where $A = 1, B = 0, C = 4, f(t) = \sin(2t)^2$. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the non-homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' + 4x = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $x = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} + 4 e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$x = e^{\alpha t} (c_1 \cos(\beta t) + c_2 \sin(\beta t))$$

Which becomes

$$x = e^0 (c_1 \cos(2t) + c_2 \sin(2t))$$

Or

$$x = c_1 \cos(2t) + c_2 \sin(2t)$$

Therefore the homogeneous solution x_h is

$$x_h = c_1 \cos(2t) + c_2 \sin(2t)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(2t)^2$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{\cos(4t), \sin(4t)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(2t), \sin(2t)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$x_p = A_1 + A_2 \cos(4t) + A_3 \sin(4t)$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-12A_2 \cos(4t) - 12A_3 \sin(4t) + 4A_1 = \sin(2t)^2$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{8}, A_2 = \frac{1}{24}, A_3 = 0 \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = \frac{1}{8} + \frac{\cos(4t)}{24}$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= (c_1 \cos(2t) + c_2 \sin(2t)) + \left(\frac{1}{8} + \frac{\cos(4t)}{24} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$x = c_1 \cos(2t) + c_2 \sin(2t) + \frac{1}{8} + \frac{\cos(4t)}{24} \quad (1)$$




Figure 59: Slope field plot

Verification of solutions

$$x = c_1 \cos(2t) + c_2 \sin(2t) + \frac{1}{8} + \frac{\cos(4t)}{24}$$

Verified OK.

3.5.2 Solving using Kovacic algorithm

Writing the ode as

$$x'' + 4x = 0 \quad (1)$$

$$Ax'' + Bx' + Cx = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = xe^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = -4z(t)\tag{7}$$

Equation (7) is now solved. After finding $z(t)$ then x is found using the inverse transformation

$$x = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 51: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = \cos(2t)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in x is found from

$$x_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 x_1 &= z_1 \\
 &= \cos(2t)
 \end{aligned}$$

Which simplifies to

$$x_1 = \cos(2t)$$

The second solution x_2 to the original ode is found using reduction of order

$$x_2 = x_1 \int \frac{e^{\int -\frac{B}{A} dt}}{x_1^2} dt$$

Since $B = 0$ then the above becomes

$$\begin{aligned} x_2 &= x_1 \int \frac{1}{x_1^2} dt \\ &= \cos(2t) \int \frac{1}{\cos(2t)^2} dt \\ &= \cos(2t) \left(\frac{\tan(2t)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} x &= c_1 x_1 + c_2 x_2 \\ &= c_1(\cos(2t)) + c_2 \left(\cos(2t) \left(\frac{\tan(2t)}{2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the nonhomogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' + 4x = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$x_h = c_1 \cos(2t) + \frac{c_2 \sin(2t)}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(2t)^2$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{\cos(4t), \sin(4t)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(2t)}{2}, \cos(2t) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$x_p = A_1 + A_2 \cos(4t) + A_3 \sin(4t)$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-12A_2 \cos(4t) - 12A_3 \sin(4t) + 4A_1 = \sin(2t)^2$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{8}, A_2 = \frac{1}{24}, A_3 = 0 \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = \frac{1}{8} + \frac{\cos(4t)}{24}$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= \left(c_1 \cos(2t) + \frac{c_2 \sin(2t)}{2} \right) + \left(\frac{1}{8} + \frac{\cos(4t)}{24} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$x = c_1 \cos(2t) + \frac{c_2 \sin(2t)}{2} + \frac{1}{8} + \frac{\cos(4t)}{24} \quad (1)$$




Figure 60: Slope field plot

Verification of solutions

$$x = c_1 \cos(2t) + \frac{c_2 \sin(2t)}{2} + \frac{1}{8} + \frac{\cos(4t)}{24}$$

Verified OK.

3.5.3 Maple step by step solution

Let's solve

$$x'' + 4x = \sin(2t)^2$$

- Highest derivative means the order of the ODE is 2
 x''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2I, 2I)$$

- 1st solution of the homogeneous ODE

$$x_1(t) = \cos(2t)$$

- 2nd solution of the homogeneous ODE

$$x_2(t) = \sin(2t)$$

- General solution of the ODE

$$x = c_1 x_1(t) + c_2 x_2(t) + x_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$x = c_1 \cos(2t) + c_2 \sin(2t) + x_p(t)$$

- Find a particular solution $x_p(t)$ of the ODE

- Use variation of parameters to find x_p here $f(t)$ is the forcing function

$$\left[x_p(t) = -x_1(t) \left(\int \frac{x_2(t)f(t)}{W(x_1(t), x_2(t))} dt \right) + x_2(t) \left(\int \frac{x_1(t)f(t)}{W(x_1(t), x_2(t))} dt \right), f(t) = \sin(2t)^2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(x_1(t), x_2(t)) = \begin{bmatrix} \cos(2t) & \sin(2t) \\ -2\sin(2t) & 2\cos(2t) \end{bmatrix}$$

- Compute Wronskian

$$W(x_1(t), x_2(t)) = 2$$

- Substitute functions into equation for $x_p(t)$

$$x_p(t) = -\frac{\cos(2t) \left(\int \sin(2t)^3 dt \right)}{2} + \frac{\sin(2t) \left(\int \cos(2t) \sin(2t)^2 dt \right)}{2}$$

- Compute integrals

$$x_p(t) = \frac{1}{8} + \frac{\cos(4t)}{24}$$

- Substitute particular solution into general solution to ODE

$$x = c_1 \cos(2t) + c_2 \sin(2t) + \frac{1}{8} + \frac{\cos(4t)}{24}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(x(t),t$2)+4*x(t)=sin(2*t)^2,x(t), singsol=all)
```

$$x(t) = \sin(2t) c_2 + \cos(2t) c_1 + \frac{1}{8} + \frac{\cos(4t)}{24}$$

✓ Solution by Mathematica

Time used: 0.09 (sec). Leaf size: 31

```
DSolve[x''[t]+4*x[t]==Sin[2*t]^2,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{24} \cos(4t) + c_1 \cos(2t) + c_2 \sin(2t) + \frac{1}{8}$$

3.6 problem Problem 12.6

3.6.1	Solving as second order euler ode	304
3.6.2	Solving as linear second order ode solved by an integrating factor ode	307
3.6.3	Solving as second order change of variable on x method 2 ode .	308
3.6.4	Solving as second order change of variable on x method 1 ode .	314
3.6.5	Solving as second order change of variable on y method 1 ode .	318
3.6.6	Solving as second order change of variable on y method 2 ode .	323
3.6.7	Solving as second order ode non constant coeff transformation on B ode	328
3.6.8	Solving using Kovacic algorithm	333

Internal problem ID [5189]

Internal file name [OUTPUT/4682_Sunday_June_05_2022_03_03_17_PM_364858/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. page 104

Problem number: Problem 12.6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_1", "second_order_change_of_variable_on_y_method_2", "linear_second_order_ode_solved_by_an_integrating_factor", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$t^2 N'' - 2tN' + 2N = t \ln(t)$$

3.6.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$AN''(t) + BN'(t) + CN(t) = f(t)$$

Where $A = t^2, B = -2t, C = 2, f(t) = t \ln(t)$. Let the solution be

$$N = N_h + N_p$$

Where N_h is the solution to the homogeneous ODE $AN''(t) + BN'(t) + CN(t) = 0$, and N_p is a particular solution to the non-homogeneous ODE $AN''(t) + BN'(t) + CN(t) = f(t)$. Solving for N_h from

$$t^2N'' - 2tN' + 2N = 0$$

This is Euler second order ODE. Let the solution be $N = t^r$, then $N' = rt^{r-1}$ and $N'' = r(r-1)t^{r-2}$. Substituting these back into the given ODE gives

$$t^2(r(r-1))t^{r-2} - 2trt^{r-1} + 2t^r = 0$$

Simplifying gives

$$r(r-1)t^r - 2rt^r + 2t^r = 0$$

Since $t^r \neq 0$ then dividing throughout by t^r gives

$$r(r-1) - 2r + 2 = 0$$

Or

$$r^2 - 3r + 2 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 1$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$N = c_1N_1 + c_2N_2$$

Where $N_1 = t^{r_1}$ and $N_2 = t^{r_2}$. Hence

$$N = c_2t^2 + c_1t$$

Next, we find the particular solution to the ODE

$$t^2N'' - 2tN' + 2N = t \ln(t)$$

The particular solution N_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$N_p(t) = u_1 N_1 + u_2 N_2 \quad (1)$$

Where u_1, u_2 to be determined, and N_1, N_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$N_1 = t$$

$$N_2 = t^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{N_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{N_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of N'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} N_1 & N_2 \\ N_1' & N_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} t & t^2 \\ \frac{d}{dt}(t) & \frac{d}{dt}(t^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} t & t^2 \\ 1 & 2t \end{vmatrix}$$

Therefore

$$W = (t)(2t) - (t^2) \quad (1)$$

Which simplifies to

$$W = t^2$$

Which simplifies to

$$W = t^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{t^3 \ln(t)}{t^4} dt$$

Which simplifies to

$$u_1 = - \int \frac{\ln(t)}{t} dt$$

Hence

$$u_1 = - \frac{\ln(t)^2}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{t^2 \ln(t)}{t^4} dt$$

Which simplifies to

$$u_2 = \int \frac{\ln(t)}{t^2} dt$$

Hence

$$u_2 = - \frac{\ln(t)}{t} - \frac{1}{t}$$

Which simplifies to

$$u_1 = - \frac{\ln(t)^2}{2}$$
$$u_2 = \frac{-\ln(t) - 1}{t}$$

Therefore the particular solution, from equation (1) is

$$N_p(t) = - \frac{\ln(t)^2 t}{2} + (-\ln(t) - 1) t$$

Which simplifies to

$$N_p(t) = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Therefore the general solution is

$$\begin{aligned} N &= N_h + N_p \\ &= -\frac{t(\ln(t)^2 - 2c_2t + 2\ln(t) - 2c_1 + 2)}{2} \end{aligned}$$

Summary

The solution(s) found are the following

$$N = -\frac{t(\ln(t)^2 - 2c_2t + 2\ln(t) - 2c_1 + 2)}{2} \quad (1)$$

Verification of solutions

$$N = -\frac{t(\ln(t)^2 - 2c_2t + 2\ln(t) - 2c_1 + 2)}{2}$$

Verified OK.

3.6.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$N'' + p(t) N' + \frac{(p(t)^2 + p'(t)) N}{2} = f(t)$$

Where $p(t) = -\frac{2}{t}$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -\frac{2}{t} dx} \\ &= \frac{1}{t} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)N)'' &= \frac{\ln(t)}{t^2} \\ \left(\frac{N}{t}\right)'' &= \frac{\ln(t)}{t^2} \end{aligned}$$

Integrating once gives

$$\left(\frac{N}{t}\right)' = \frac{-\ln(t) - 1}{t} + c_1$$

Integrating again gives

$$\left(\frac{N}{t}\right) = c_1 t - \ln(t) - \frac{\ln(t)^2}{2} + c_2$$

Hence the solution is

$$N = \frac{c_1 t - \ln(t) - \frac{\ln(t)^2}{2} + c_2}{\frac{1}{t}}$$

Or

$$N = c_1 t^2 - \frac{t \ln(t)^2}{2} + c_2 t - t \ln(t)$$

Summary

The solution(s) found are the following

$$N = c_1 t^2 - \frac{t \ln(t)^2}{2} + c_2 t - t \ln(t) \quad (1)$$

Verification of solutions

$$N = c_1 t^2 - \frac{t \ln(t)^2}{2} + c_2 t - t \ln(t)$$

Verified OK.

3.6.3 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$N = N_h + N_p$$

Where N_h is the solution to the homogeneous ODE $AN''(t) + BN'(t) + CN(t) = 0$, and N_p is a particular solution to the non-homogeneous ODE $AN''(t) + BN'(t) + CN(t) = f(t)$. N_h is the solution to

$$t^2 N'' - 2t N' + 2N = 0$$

In normal form the ode

$$t^2 N'' - 2t N' + 2N = 0 \quad (1)$$

Becomes

$$N'' + p(t) N' + q(t) N = 0 \quad (2)$$

Where

$$p(t) = -\frac{2}{t}$$
$$q(t) = \frac{2}{t^2}$$

Applying change of variables $\tau = g(t)$ to (2) gives

$$\frac{d^2}{d\tau^2} N(\tau) + p_1 \left(\frac{d}{d\tau} N(\tau) \right) + q_1 N(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(t) + p(t) \tau'(t)}{\tau'(t)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(t)}{\tau'(t)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(t) + p(t) \tau'(t) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(t) dt)} dt \\ &= \int e^{-(\int -\frac{2}{t} dt)} dt \\ &= \int e^{2 \ln(t)} dt \\ &= \int t^2 dt \\ &= \frac{t^3}{3} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(t)}{\tau'(t)^2} \\ &= \frac{\frac{2}{t^2}}{t^4} \\ &= \frac{2}{t^6} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}N(\tau) + q_1N(\tau) &= 0 \\ \frac{d^2}{d\tau^2}N(\tau) + \frac{2N(\tau)}{t^6} &= 0\end{aligned}$$

But in terms of τ

$$\frac{2}{t^6} = \frac{2}{9\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}N(\tau) + \frac{2N(\tau)}{9\tau^2} = 0$$

The above ode is now solved for $N(\tau)$. The ode can be written as

$$9\left(\frac{d^2}{d\tau^2}N(\tau)\right)\tau^2 + 2N(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $N(\tau) = \tau^r$, then $N' = r\tau^{r-1}$ and $N'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$9\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + 2\tau^r = 0$$

Simplifying gives

$$9r(r-1)\tau^r + 0\tau^r + 2\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$9r(r-1) + 0 + 2 = 0$$

Or

$$9r^2 - 9r + 2 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$\begin{aligned}r_1 &= \frac{1}{3} \\ r_2 &= \frac{2}{3}\end{aligned}$$

Since the roots are real and distinct, then the general solution is

$$N(\tau) = c_1N_1 + c_2N_2$$

Where $N_1 = \tau^{r_1}$ and $N_2 = \tau^{r_2}$. Hence

$$N(\tau) = c_1\tau^{\frac{1}{3}} + c_2\tau^{\frac{2}{3}}$$

The above solution is now transformed back to N using (6) which results in

$$N = \frac{c_1 3^{\frac{2}{3}} (t^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}} (t^3)^{\frac{2}{3}}}{3}$$

Therefore the homogeneous solution N_h is

$$N_h = \frac{c_1 3^{\frac{2}{3}} (t^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}} (t^3)^{\frac{2}{3}}}{3}$$

The particular solution N_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$N_p(t) = u_1 N_1 + u_2 N_2 \quad (1)$$

Where u_1, u_2 to be determined, and N_1, N_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$N_1 = (t^3)^{\frac{1}{3}}$$

$$N_2 = (t^3)^{\frac{2}{3}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{N_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{N_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of N'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} N_1 & N_2 \\ N_1' & N_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (t^3)^{\frac{1}{3}} & (t^3)^{\frac{2}{3}} \\ \frac{d}{dt} \left((t^3)^{\frac{1}{3}} \right) & \frac{d}{dt} \left((t^3)^{\frac{2}{3}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (t^3)^{\frac{1}{3}} & (t^3)^{\frac{2}{3}} \\ \frac{t^2}{(t^3)^{\frac{2}{3}}} & \frac{2t^2}{(t^3)^{\frac{1}{3}}} \end{vmatrix}$$

Therefore

$$W = \left((t^3)^{\frac{1}{3}} \right) \left(\frac{2t^2}{(t^3)^{\frac{1}{3}}} \right) - \left((t^3)^{\frac{2}{3}} \right) \left(\frac{t^2}{(t^3)^{\frac{2}{3}}} \right)$$

Which simplifies to

$$W = t^2$$

Which simplifies to

$$W = t^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(t^3)^{\frac{2}{3}} t \ln(t)}{t^4} dt$$

Which simplifies to

$$u_1 = - \int \frac{(t^3)^{\frac{2}{3}} \ln(t)}{t^3} dt$$

Hence

$$u_1 = - \frac{(t^3)^{\frac{2}{3}} \ln(t)^2}{2t^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(t^3)^{\frac{1}{3}} t \ln(t)}{t^4} dt$$

Which simplifies to

$$u_2 = \int \frac{(t^3)^{\frac{1}{3}} \ln(t)}{t^3} dt$$

Hence

$$u_2 = -\frac{(t^3)^{\frac{1}{3}} \ln(t)}{t^2} - \frac{(t^3)^{\frac{1}{3}}}{t^2}$$

Which simplifies to

$$u_1 = -\frac{(t^3)^{\frac{2}{3}} \ln(t)^2}{2t^2}$$
$$u_2 = \frac{(-\ln(t) - 1)(t^3)^{\frac{1}{3}}}{t^2}$$

Therefore the particular solution, from equation (1) is

$$N_p(t) = -\frac{t \ln(t)^2}{2} + (-\ln(t) - 1)t$$

Which simplifies to

$$N_p(t) = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Therefore the general solution is

$$N = N_h + N_p$$
$$= \left(\frac{c_1 3^{\frac{2}{3}} (t^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}} (t^3)^{\frac{2}{3}}}{3} \right) + \left(t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \right)$$

Summary

The solution(s) found are the following

$$N = \frac{c_1 3^{\frac{2}{3}} (t^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}} (t^3)^{\frac{2}{3}}}{3} + t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \quad (1)$$

Verification of solutions

$$N = \frac{c_1 3^{\frac{2}{3}} (t^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}} (t^3)^{\frac{2}{3}}}{3} + t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Verified OK.

3.6.4 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$AN''(t) + BN'(t) + CN(t) = f(t)$$

Where $A = t^2, B = -2t, C = 2, f(t) = t \ln(t)$. Let the solution be

$$N = N_h + N_p$$

Where N_h is the solution to the homogeneous ODE $AN''(t) + BN'(t) + CN(t) = 0$, and N_p is a particular solution to the non-homogeneous ODE $AN''(t) + BN'(t) + CN(t) = f(t)$. Solving for N_h from

$$t^2N'' - 2tN' + 2N = 0$$

In normal form the ode

$$t^2N'' - 2tN' + 2N = 0 \quad (1)$$

Becomes

$$N'' + p(t)N' + q(t)N = 0 \quad (2)$$

Where

$$p(t) = -\frac{2}{t}$$

$$q(t) = \frac{2}{t^2}$$

Applying change of variables $\tau = g(t)$ to (2) results

$$\frac{d^2}{d\tau^2}N(\tau) + p_1\left(\frac{d}{d\tau}N(\tau)\right) + q_1N(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(t) + p(t)\tau'(t)}{\tau'(t)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(t)}{\tau'(t)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{\sqrt{2}\sqrt{\frac{1}{t^2}}}{c} \quad (6)$$

$$\tau'' = -\frac{\sqrt{2}}{c\sqrt{\frac{1}{t^2}}t^3}$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(t) + p(t) \tau'(t)}{\tau'(t)^2} \\
 &= \frac{-\frac{\sqrt{2}}{c\sqrt{\frac{1}{t^2}}t^3} - \frac{2}{t} \frac{\sqrt{2}}{c} \sqrt{\frac{1}{t^2}}}{\left(\frac{\sqrt{2}}{c} \sqrt{\frac{1}{t^2}}\right)^2} \\
 &= -\frac{3c\sqrt{2}}{2}
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 N(\tau)'' + p_1 N(\tau)' + q_1 N(\tau) &= 0 \\
 \frac{d^2}{d\tau^2} N(\tau) - \frac{3c\sqrt{2}}{2} \left(\frac{d}{d\tau} N(\tau)\right) + c^2 N(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $N(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$N(\tau) = e^{\frac{3\sqrt{2}c\tau}{4}} \left(c_1 \cosh\left(\frac{\sqrt{2}c\tau}{4}\right) + ic_2 \sinh\left(\frac{\sqrt{2}c\tau}{4}\right) \right)$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c} \sqrt{q} dt \\
 &= \frac{\int \sqrt{2} \sqrt{\frac{1}{t^2}} dt}{c} \\
 &= \frac{\sqrt{2} \sqrt{\frac{1}{t^2}} t \ln(t)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$N = t^{\frac{3}{2}} \left(c_1 \cosh\left(\frac{\ln(t)}{2}\right) + ic_2 \sinh\left(\frac{\ln(t)}{2}\right) \right)$$

Now the particular solution to this ODE is found

$$t^2 N'' - 2tN' + 2N = t \ln(t)$$

The particular solution N_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$N_p(t) = u_1 N_1 + u_2 N_2 \quad (1)$$

Where u_1, u_2 to be determined, and N_1, N_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$N_1 = (t^3)^{\frac{1}{3}}$$

$$N_2 = (t^3)^{\frac{2}{3}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{N_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{N_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of N'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} N_1 & N_2 \\ N_1' & N_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (t^3)^{\frac{1}{3}} & (t^3)^{\frac{2}{3}} \\ \frac{d}{dt} \left((t^3)^{\frac{1}{3}} \right) & \frac{d}{dt} \left((t^3)^{\frac{2}{3}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (t^3)^{\frac{1}{3}} & (t^3)^{\frac{2}{3}} \\ \frac{t^2}{(t^3)^{\frac{2}{3}}} & \frac{2t^2}{(t^3)^{\frac{1}{3}}} \end{vmatrix}$$

Therefore

$$W = \left((t^3)^{\frac{1}{3}} \right) \left(\frac{2t^2}{(t^3)^{\frac{1}{3}}} \right) - \left((t^3)^{\frac{2}{3}} \right) \left(\frac{t^2}{(t^3)^{\frac{2}{3}}} \right)$$

Which simplifies to

$$W = t^2$$

Which simplifies to

$$W = t^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(t^3)^{\frac{2}{3}} t \ln(t)}{t^4} dt$$

Which simplifies to

$$u_1 = - \int \frac{(t^3)^{\frac{2}{3}} \ln(t)}{t^3} dt$$

Hence

$$u_1 = - \frac{(t^3)^{\frac{2}{3}} \ln(t)^2}{2t^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(t^3)^{\frac{1}{3}} t \ln(t)}{t^4} dt$$

Which simplifies to

$$u_2 = \int \frac{(t^3)^{\frac{1}{3}} \ln(t)}{t^3} dt$$

Hence

$$u_2 = - \frac{(t^3)^{\frac{1}{3}} \ln(t)}{t^2} - \frac{(t^3)^{\frac{1}{3}}}{t^2}$$

Which simplifies to

$$u_1 = - \frac{(t^3)^{\frac{2}{3}} \ln(t)^2}{2t^2}$$
$$u_2 = \frac{(-\ln(t) - 1)(t^3)^{\frac{1}{3}}}{t^2}$$

Therefore the particular solution, from equation (1) is

$$N_p(t) = - \frac{t \ln(t)^2}{2} + (-\ln(t) - 1)t$$

Which simplifies to

$$N_p(t) = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Therefore the general solution is

$$\begin{aligned} N &= N_h + N_p \\ &= \left(t^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(t)}{2} \right) + i c_2 \sinh \left(\frac{\ln(t)}{2} \right) \right) \right) + \left(t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \right) \\ &= t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) + t^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(t)}{2} \right) + i c_2 \sinh \left(\frac{\ln(t)}{2} \right) \right) \end{aligned}$$

Which simplifies to

$$N = i t^{\frac{3}{2}} \sinh \left(\frac{\ln(t)}{2} \right) c_2 + t^{\frac{3}{2}} \cosh \left(\frac{\ln(t)}{2} \right) c_1 - \frac{t \ln(t)^2}{2} - t \ln(t) - t$$

Summary

The solution(s) found are the following

$$N = i t^{\frac{3}{2}} \sinh \left(\frac{\ln(t)}{2} \right) c_2 + t^{\frac{3}{2}} \cosh \left(\frac{\ln(t)}{2} \right) c_1 - \frac{t \ln(t)^2}{2} - t \ln(t) - t \quad (1)$$

Verification of solutions

$$N = i t^{\frac{3}{2}} \sinh \left(\frac{\ln(t)}{2} \right) c_2 + t^{\frac{3}{2}} \cosh \left(\frac{\ln(t)}{2} \right) c_1 - \frac{t \ln(t)^2}{2} - t \ln(t) - t$$

Verified OK.

3.6.5 Solving as second order change of variable on y method 1 ode

This is second order non-homogeneous ODE. Let the solution be

$$N = N_h + N_p$$

Where N_h is the solution to the homogeneous ODE $AN''(t) + BN'(t) + CN(t) = 0$, and N_p is a particular solution to the non-homogeneous ODE $AN''(t) + BN'(t) + CN(t) = f(t)$. N_h is the solution to

$$t^2 N'' - 2t N' + 2N = 0$$

In normal form the given ode is written as

$$N'' + p(t) N' + q(t) N = 0 \quad (2)$$

Where

$$p(t) = -\frac{2}{t}$$

$$q(t) = \frac{2}{t^2}$$

Calculating the Liouville ode invariant Q given by

$$Q = q - \frac{p'}{2} - \frac{p^2}{4}$$

$$= \frac{2}{t^2} - \frac{\left(-\frac{2}{t}\right)'}{2} - \frac{\left(-\frac{2}{t}\right)^2}{4}$$

$$= \frac{2}{t^2} - \frac{\left(\frac{2}{t^2}\right)}{2} - \frac{\left(\frac{4}{t^2}\right)}{4}$$

$$= \frac{2}{t^2} - \left(\frac{1}{t^2}\right) - \frac{1}{t^2}$$

$$= 0$$

Since the Liouville ode invariant does not depend on the independent variable t then the transformation

$$N = v(t) z(t) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(t)$ is given by

$$z(t) = e^{-\left(\int \frac{p(t)}{2} dt\right)}$$

$$= e^{-\int \frac{-2}{2} dt}$$

$$= t \quad (5)$$

Hence (3) becomes

$$N = v(t) t \quad (4)$$

Applying this change of variable to the original ode results in

$$t^2 v''(t) = \ln(t)$$

Which is now solved for $v(t)$ Simplifying the ode gives

$$v''(t) = \frac{\ln(t)}{t^2}$$

Integrating once gives

$$v'(t) = -\frac{\ln(t)}{t} - \frac{1}{t} + c_1$$

Integrating again gives

$$v(t) = -\frac{\ln(t)^2}{2} - \ln(t) + c_1x + c_2$$

Now that $v(t)$ is known, then

$$\begin{aligned} N &= v(t) z(t) \\ &= \left(c_1t - \ln(t) - \frac{\ln(t)^2}{2} + c_2 \right) (z(t)) \end{aligned} \tag{7}$$

But from (5)

$$z(t) = t$$

Hence (7) becomes

$$N = \left(c_1t - \ln(t) - \frac{\ln(t)^2}{2} + c_2 \right) t$$

Therefore the homogeneous solution N_h is

$$N_h = \left(c_1t - \ln(t) - \frac{\ln(t)^2}{2} + c_2 \right) t$$

The particular solution N_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$N_p(t) = u_1N_1 + u_2N_2 \tag{1}$$

Where u_1, u_2 to be determined, and N_1, N_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$N_1 = (t^3)^{\frac{1}{3}}$$

$$N_2 = (t^3)^{\frac{2}{3}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{N_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{N_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of N'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} N_1 & N_2 \\ N_1' & N_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (t^3)^{\frac{1}{3}} & (t^3)^{\frac{2}{3}} \\ \frac{d}{dt} \left((t^3)^{\frac{1}{3}} \right) & \frac{d}{dt} \left((t^3)^{\frac{2}{3}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (t^3)^{\frac{1}{3}} & (t^3)^{\frac{2}{3}} \\ \frac{t^2}{(t^3)^{\frac{2}{3}}} & \frac{2t^2}{(t^3)^{\frac{1}{3}}} \end{vmatrix}$$

Therefore

$$W = \left((t^3)^{\frac{1}{3}} \right) \left(\frac{2t^2}{(t^3)^{\frac{1}{3}}} \right) - \left((t^3)^{\frac{2}{3}} \right) \left(\frac{t^2}{(t^3)^{\frac{2}{3}}} \right)$$

Which simplifies to

$$W = t^2$$

Which simplifies to

$$W = t^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(t^3)^{\frac{2}{3}} t \ln(t)}{t^4} dt$$

Which simplifies to

$$u_1 = - \int \frac{(t^3)^{\frac{2}{3}} \ln(t)}{t^3} dt$$

Hence

$$u_1 = - \frac{(t^3)^{\frac{2}{3}} \ln(t)^2}{2t^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(t^3)^{\frac{1}{3}} t \ln(t)}{t^4} dt$$

Which simplifies to

$$u_2 = \int \frac{(t^3)^{\frac{1}{3}} \ln(t)}{t^3} dt$$

Hence

$$u_2 = - \frac{(t^3)^{\frac{1}{3}} \ln(t)}{t^2} - \frac{(t^3)^{\frac{1}{3}}}{t^2}$$

Which simplifies to

$$u_1 = - \frac{(t^3)^{\frac{2}{3}} \ln(t)^2}{2t^2}$$
$$u_2 = \frac{(-\ln(t) - 1)(t^3)^{\frac{1}{3}}}{t^2}$$

Therefore the particular solution, from equation (1) is

$$N_p(t) = - \frac{t \ln(t)^2}{2} + (-\ln(t) - 1)t$$

Which simplifies to

$$N_p(t) = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Therefore the general solution is

$$\begin{aligned} N &= N_h + N_p \\ &= \left(\left(c_1 t - \ln(t) - \frac{\ln(t)^2}{2} + c_2 \right) t \right) + \left(t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$N = \left(c_1 t - \ln(t) - \frac{\ln(t)^2}{2} + c_2 \right) t + t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \quad (1)$$

Verification of solutions

$$N = \left(c_1 t - \ln(t) - \frac{\ln(t)^2}{2} + c_2 \right) t + t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Verified OK.

3.6.6 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$AN''(t) + BN'(t) + CN(t) = f(t)$$

Where $A = t^2$, $B = -2t$, $C = 2$, $f(t) = t \ln(t)$. Let the solution be

$$N = N_h + N_p$$

Where N_h is the solution to the homogeneous ODE $AN''(t) + BN'(t) + CN(t) = 0$, and N_p is a particular solution to the non-homogeneous ODE $AN''(t) + BN'(t) + CN(t) = f(t)$. Solving for N_h from

$$t^2 N'' - 2t N' + 2N = 0$$

In normal form the ode

$$t^2 N'' - 2t N' + 2N = 0 \quad (1)$$

Becomes

$$N'' + p(t) N' + q(t) N = 0 \quad (2)$$

Where

$$p(t) = -\frac{2}{t}$$
$$q(t) = \frac{2}{t^2}$$

Applying change of variables on the dependent variable $N = v(t) t^n$ to (2) gives the following ode where the dependent variables is $v(t)$ and not N .

$$v''(t) + \left(\frac{2n}{t} + p \right) v'(t) + \left(\frac{n(n-1)}{t^2} + \frac{np}{t} + q \right) v(t) = 0 \quad (3)$$

Let the coefficient of $v(t)$ above be zero. Hence

$$\frac{n(n-1)}{t^2} + \frac{np}{t} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(t)$ and $q(t)$ into (4) gives

$$\frac{n(n-1)}{t^2} - \frac{2n}{t^2} + \frac{2}{t^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$v''(t) + \frac{2v'(t)}{t} = 0$$
$$v''(t) + \frac{2v'(t)}{t} = 0 \quad (7)$$

Using the substitution

$$u(t) = v'(t)$$

Then (7) becomes

$$u'(t) + \frac{2u(t)}{t} = 0 \quad (8)$$

The above is now solved for $u(t)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(t, u) \\ &= f(t)g(u) \\ &= -\frac{2u}{t}\end{aligned}$$

Where $f(t) = -\frac{2}{t}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{2}{t} dt \\ \int \frac{1}{u} du &= \int -\frac{2}{t} dt \\ \ln(u) &= -2 \ln(t) + c_1 \\ u &= e^{-2 \ln(t) + c_1} \\ &= \frac{c_1}{t^2}\end{aligned}$$

Now that $u(t)$ is known, then

$$\begin{aligned}v'(t) &= u(t) \\ v(t) &= \int u(t) dt + c_2 \\ &= -\frac{c_1}{t} + c_2\end{aligned}$$

Hence

$$\begin{aligned}N &= v(t) t^n \\ &= \left(-\frac{c_1}{t} + c_2\right) t^2 \\ &= (c_2 t - c_1) t\end{aligned}$$

Now the particular solution to this ODE is found

$$t^2 N'' - 2tN' + 2N = t \ln(t)$$

The particular solution N_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$N_p(t) = u_1 N_1 + u_2 N_2 \tag{1}$$

Where u_1, u_2 to be determined, and N_1, N_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$N_1 = t$$

$$N_2 = t^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{N_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{N_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of N'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} N_1 & N_2 \\ N_1' & N_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} t & t^2 \\ \frac{d}{dt}(t) & \frac{d}{dt}(t^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} t & t^2 \\ 1 & 2t \end{vmatrix}$$

Therefore

$$W = (t)(2t) - (t^2)(1)$$

Which simplifies to

$$W = t^2$$

Which simplifies to

$$W = t^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{t^3 \ln(t)}{t^4} dt$$

Which simplifies to

$$u_1 = - \int \frac{\ln(t)}{t} dt$$

Hence

$$u_1 = -\frac{\ln(t)^2}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{t^2 \ln(t)}{t^4} dt$$

Which simplifies to

$$u_2 = \int \frac{\ln(t)}{t^2} dt$$

Hence

$$u_2 = -\frac{\ln(t)}{t} - \frac{1}{t}$$

Which simplifies to

$$u_1 = -\frac{\ln(t)^2}{2}$$
$$u_2 = \frac{-\ln(t) - 1}{t}$$

Therefore the particular solution, from equation (1) is

$$N_p(t) = -\frac{t \ln(t)^2}{2} + (-\ln(t) - 1)t$$

Which simplifies to

$$N_p(t) = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Therefore the general solution is

$$\begin{aligned}
 N &= N_h + N_p \\
 &= \left(\left(-\frac{c_1}{t} + c_2 \right) t^2 \right) + \left(t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \right) \\
 &= t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) + \left(-\frac{c_1}{t} + c_2 \right) t^2
 \end{aligned}$$

Which simplifies to

$$N = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 + c_2 t - c_1 \right)$$

Summary

The solution(s) found are the following

$$N = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 + c_2 t - c_1 \right) \quad (1)$$

Verification of solutions

$$N = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 + c_2 t - c_1 \right)$$

Verified OK.

3.6.7 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$AN'' + BN' + CN = F(t)$$

This method reduces the order ode the ODE by one by applying the transformation

$$N = Bv$$

This results in

$$\begin{aligned}
 N' &= B'v + v'B \\
 N'' &= B''v + B'v' + v''B + v'B' \\
 &= v''B + 2v' + B' + B''v
 \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $N = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned} A &= t^2 \\ B &= -2t \\ C &= 2 \\ F &= t \ln(t) \end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (t^2)(0) + (-2t)(-2) + (2)(-2t) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-2t^3v'' + (0)v' = 0$$

Now by applying $v' = u$ the above becomes

$$-2t^3u'(t) = 0$$

Which is now solved for u . Integrating both sides gives

$$\begin{aligned} u(t) &= \int 0 dt \\ &= c_1 \end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= c_1\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(t) &= \int c_1 dt \\ &= c_1 t + c_2\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}N_h(t) &= Bv \\ &= (-2t)(c_1 t + c_2) \\ &= -2t(c_1 t + c_2)\end{aligned}$$

And now the particular solution $N_p(t)$ will be found. The particular solution N_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$N_p(t) = u_1 N_1 + u_2 N_2 \quad (1)$$

Where u_1, u_2 to be determined, and N_1, N_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}N_1 &= t \\ N_2 &= t^2\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{N_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{N_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of N'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} N_1 & N_2 \\ N_1' & N_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} t & t^2 \\ \frac{d}{dt}(t) & \frac{d}{dt}(t^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} t & t^2 \\ 1 & 2t \end{vmatrix}$$

Therefore

$$W = (t)(2t) - (t^2)(1)$$

Which simplifies to

$$W = t^2$$

Which simplifies to

$$W = t^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{t^3 \ln(t)}{t^4} dt$$

Which simplifies to

$$u_1 = - \int \frac{\ln(t)}{t} dt$$

Hence

$$u_1 = - \frac{\ln(t)^2}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{t^2 \ln(t)}{t^4} dt$$

Which simplifies to

$$u_2 = \int \frac{\ln(t)}{t^2} dt$$

Hence

$$u_2 = - \frac{\ln(t)}{t} - \frac{1}{t}$$

Which simplifies to

$$u_1 = -\frac{\ln(t)^2}{2}$$
$$u_2 = \frac{-\ln(t) - 1}{t}$$

Therefore the particular solution, from equation (1) is

$$N_p(t) = -\frac{t \ln(t)^2}{2} + (-\ln(t) - 1)t$$

Which simplifies to

$$N_p(t) = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Hence the complete solution is

$$\begin{aligned} N(t) &= N_h + N_p \\ &= (-2t(c_1t + c_2)) + \left(t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \right) \\ &= t \left(-2c_1t - 2c_2 - \frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$N = t \left(-2c_1t - 2c_2 - \frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \quad (1)$$

Verification of solutions

$$N = t \left(-2c_1t - 2c_2 - \frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Verified OK.

3.6.8 Solving using Kovacic algorithm

Writing the ode as

$$t^2 N'' - 2tN' + 2N = 0 \quad (1)$$

$$AN'' + BN' + CN = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= t^2 \\ B &= -2t \\ C &= 2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = Ne^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then N is found using the inverse transformation

$$N = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 53: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in N is found from

$$\begin{aligned} N_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2t}{t^2} dt} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^{\ln(t)} \\
&= z_1(t)
\end{aligned}$$

Which simplifies to

$$N_1 = t$$

The second solution N_2 to the original ode is found using reduction of order

$$N_2 = N_1 \int \frac{e^{\int -\frac{B}{A} dt}}{N_1^2} dt$$

Substituting gives

$$\begin{aligned}
N_2 &= N_1 \int \frac{e^{\int -\frac{-2t}{t^2} dt}}{(N_1)^2} dt \\
&= N_1 \int \frac{e^{2 \ln(t)}}{(N_1)^2} dt \\
&= N_1(t)
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
N &= c_1 N_1 + c_2 N_2 \\
&= c_1(t) + c_2(t(t))
\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$N = N_h + N_p$$

Where N_h is the solution to the homogeneous ODE $AN''(t) + BN'(t) + CN(t) = 0$, and N_p is a particular solution to the nonhomogeneous ODE $AN''(t) + BN'(t) + CN(t) = f(t)$. N_h is the solution to

$$t^2 N'' - 2t N' + 2N = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$N_h = c_2 t^2 + c_1 t$$

The particular solution N_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$N_p(t) = u_1 N_1 + u_2 N_2 \quad (1)$$

Where u_1, u_2 to be determined, and N_1, N_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$N_1 = t$$

$$N_2 = t^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{N_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{N_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of N'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} N_1 & N_2 \\ N_1' & N_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} t & t^2 \\ \frac{d}{dt}(t) & \frac{d}{dt}(t^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} t & t^2 \\ 1 & 2t \end{vmatrix}$$

Therefore

$$W = (t)(2t) - (t^2) \quad (1)$$

Which simplifies to

$$W = t^2$$

Which simplifies to

$$W = t^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{t^3 \ln(t)}{t^4} dt$$

Which simplifies to

$$u_1 = - \int \frac{\ln(t)}{t} dt$$

Hence

$$u_1 = - \frac{\ln(t)^2}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{t^2 \ln(t)}{t^4} dt$$

Which simplifies to

$$u_2 = \int \frac{\ln(t)}{t^2} dt$$

Hence

$$u_2 = - \frac{\ln(t)}{t} - \frac{1}{t}$$

Which simplifies to

$$u_1 = - \frac{\ln(t)^2}{2}$$
$$u_2 = \frac{-\ln(t) - 1}{t}$$

Therefore the particular solution, from equation (1) is

$$N_p(t) = - \frac{t \ln(t)^2}{2} + (-\ln(t) - 1)t$$

Which simplifies to

$$N_p(t) = t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Therefore the general solution is

$$\begin{aligned} N &= N_h + N_p \\ &= (c_2 t^2 + c_1 t) + \left(t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \right) \end{aligned}$$

Which simplifies to

$$N = t(c_2 t + c_1) + t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Summary

The solution(s) found are the following

$$N = t(c_2 t + c_1) + t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right) \quad (1)$$

Verification of solutions

$$N = t(c_2 t + c_1) + t \left(-\frac{\ln(t)^2}{2} - \ln(t) - 1 \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    <- LODE of Euler type successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(t^2*diff(N(t),t$2)-2*t*diff(N(t),t)+2*N(t)=t*ln(t),N(t), singsol=all)
```

$$N(t) = -\frac{t(\ln(t))^2 - 2c_1t + 2\ln(t) - 2c_2 + 2}{2}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 30

```
DSolve[t^2*n'[t]-2*t*n'[t]+2*n[t]==t*Log[t],n[t],t,IncludeSingularSolutions -> True]
```

$$n(t) \rightarrow -\frac{1}{2}t \log^2(t) - t \log(t) + t(c_2t - 1 + c_1)$$

3.7 problem Problem 12.7

3.7.1	Solving as linear ode	340
3.7.2	Solving as first order ode lie symmetry lookup ode	342
3.7.3	Solving as exact ode	346
3.7.4	Maple step by step solution	351

Internal problem ID [5190]

Internal file name [OUTPUT/4683_Sunday_June_05_2022_03_03_18_PM_63440513/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. page 104

Problem number: Problem 12.7.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$y' + \frac{4y}{x} = x^4$$

3.7.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{4}{x}$$
$$q(x) = x^4$$

Hence the ode is

$$y' + \frac{4y}{x} = x^4$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{4}{x} dx} \\ &= x^4\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (x^4) \\ \frac{d}{dx}(y x^4) &= (x^4) (x^4) \\ d(y x^4) &= x^8 dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y x^4 &= \int x^8 dx \\ y x^4 &= \frac{x^9}{9} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x^4$ results in

$$y = \frac{x^5}{9} + \frac{c_1}{x^4}$$

Summary

The solution(s) found are the following

$$y = \frac{x^5}{9} + \frac{c_1}{x^4} \tag{1}$$




Figure 61: Slope field plot

Verification of solutions

$$y = \frac{x^5}{9} + \frac{c_1}{x^4}$$

Verified OK.

3.7.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{-x^5 + 4y}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 54: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int(n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{x^4}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x^4} dy \end{aligned}$$

Which results in

$$S = y x^4$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{-x^5 + 4y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= 4y x^3 \\ S_y &= x^4 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x^8 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R^8$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^9}{9} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$yx^4 = \frac{x^9}{9} + c_1$$



Which simplifies to

$$yx^4 = \frac{x^9}{9} + c_1$$

Which gives

$$y = \frac{x^9 + 9c_1}{9x^4}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{-x^5+4y}{x}$ 	$R = x$ $S = yx^4$	$\frac{dS}{dR} = R^8$ 

Summary

The solution(s) found are the following

$$y = \frac{x^9 + 9c_1}{9x^4} \quad (1)$$




Figure 62: Slope field plot

Verification of solutions

$$y = \frac{x^9 + 9c_1}{9x^4}$$

Verified OK.

3.7.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= \left(-\frac{4y}{x} + x^4\right) dx \\ \left(-x^4 + \frac{4y}{x}\right) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x^4 + \frac{4y}{x} \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-x^4 + \frac{4y}{x}\right) \\ &= \frac{4}{x}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1 \left(\left(\frac{4}{x} \right) - (0) \right) \\ &= \frac{4}{x}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int \frac{4}{x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{4 \ln(x)} \\ &= x^4\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= x^4 \left(-x^4 + \frac{4y}{x} \right) \\ &= -(x^5 - 4y) x^3\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= x^4(1) \\ &= x^4\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ (-x^5 - 4y)x^3 + (x^4) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -(x^5 - 4y)x^3 dx \\ \phi &= -\frac{1}{9}x^9 + yx^4 + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x^4 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x^4$. Therefore equation (4) becomes

$$x^4 = x^4 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{1}{9}x^9 + yx^4 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{1}{9}x^9 + yx^4$$

The solution becomes

$$y = \frac{x^9 + 9c_1}{9x^4}$$

Summary

The solution(s) found are the following

$$y = \frac{x^9 + 9c_1}{9x^4} \tag{1}$$




Figure 63: Slope field plot

Verification of solutions

$$y = \frac{x^9 + 9c_1}{9x^4}$$

Verified OK.

3.7.4 Maple step by step solution

Let's solve

$$y' + \frac{4y}{x} = x^4$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{4y}{x} + x^4$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{4y}{x} = x^4$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{4y}{x} \right) = \mu(x) x^4$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{4y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{4\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x^4$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) x^4 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) x^4 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^4 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x^4$

$$y = \frac{\int x^8 dx + c_1}{x^4}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{x^9}{9} + c_1}{x^4}$$

- Simplify

$$y = \frac{x^9 + 9c_1}{9x^4}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)+(4/x)*y(x)=x^4,y(x), singsol=all)
```

$$y(x) = \frac{x^9 + 9c_1}{9x^4}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 19

```
DSolve[y'[x]+(4/x)*y[x]==x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^5}{9} + \frac{c_1}{x^4}$$

3.8 problem Problem 12.8

Internal problem ID [5191]

Internal file name [OUTPUT/4684_Sunday_June_05_2022_03_03_20_PM_22251152/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. page 104

Problem number: Problem 12.8.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _quadrature]]
```

$$y'''' = 5x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y'''' = 0$$

The characteristic equation is

$$\lambda^4 = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 0$$

$$\lambda_3 = 0$$

$$\lambda_4 = 0$$

Therefore the homogeneous solution is

$$y_h(x) = c_4x^3 + c_3x^2 + c_2x + c_1$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = x$$

$$y_3 = x^2$$

$$y_4 = x^3$$

Now the particular solution to the given ODE is found

$$y'''' = 5x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x, x^2, x^3\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3\}]$$

Since x^2 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^3, x^4\}]$$

Since x^3 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^4, x^5\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^5 + A_1x^4$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$120xA_2 + 24A_1 = 5x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{24} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^5}{24}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_4x^3 + c_3x^2 + c_2x + c_1) + \left(\frac{x^5}{24} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_4x^3 + c_3x^2 + c_2x + c_1 + \frac{1}{24}x^5 \quad (1)$$

Verification of solutions

$$y = c_4x^3 + c_3x^2 + c_2x + c_1 + \frac{1}{24}x^5$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$4)=5*x,y(x), singsol=all)
```

$$y(x) = \frac{x^5}{24} + \frac{c_1 x^3}{6} + \frac{c_2 x^2}{2} + \frac{(3c_1^2 + 10c_3)x}{10} + c_4$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 31

```
DSolve[y''''[x]==5*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^5}{24} + c_4 x^3 + c_3 x^2 + c_2 x + c_1$$

4 Chapter 12. VARIATION OF PARAMETERS.

Supplementary Problems. page 109

4.1	problem Problem 12.9	358
4.2	problem Problem 12.10	373
4.3	problem Problem 12.11	386
4.4	problem Problem 12.12	397
4.5	problem Problem 12.13	408
4.6	problem Problem 12.14	428
4.7	problem Problem 12.15	466
4.8	problem Problem 12.16	482

4.1 problem Problem 12.9

4.1.1	Solving as second order linear constant coeff ode	358
4.1.2	Solving as linear second order ode solved by an integrating factor ode	362
4.1.3	Solving using Kovacic algorithm	364
4.1.4	Maple step by step solution	370

Internal problem ID [5192]

Internal file name [OUTPUT/4685_Sunday_June_05_2022_03_03_21_PM_45980404/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. Supplementary Problems. page 109

Problem number: Problem 12.9.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 2y' + y = \frac{e^x}{x^5}$$

4.1.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = \frac{e^x}{x^5}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 x e^x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^x \\ y_2 &= x e^x \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^x & x e^x \\ \frac{d}{dx}(e^x) & \frac{d}{dx}(x e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^x & x e^x \\ e^x & x e^x + e^x \end{vmatrix}$$

Therefore

$$W = (e^x)(x e^x + e^x) - (x e^x)(e^x)$$

Which simplifies to

$$W = e^{2x}$$

Which simplifies to

$$W = e^{2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{2x}}{e^{2x}} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{x^4} dx$$

Hence

$$u_1 = \frac{1}{3x^3}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{2x}}{x^5 e^{2x}} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x^5} dx$$

Hence

$$u_2 = -\frac{1}{4x^4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{e^x}{12x^3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + \left(\frac{e^x}{12x^3} \right) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) + \frac{e^x}{12x^3}$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) + \frac{e^x}{12x^3} \quad (1)$$




Figure 64: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + \frac{e^x}{12x^3}$$

Verified OK.

4.1.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p \, dx} \\ &= e^{\int -2 \, dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = \frac{e^{-x}e^x}{x^5}$$
$$(e^{-x}y)'' = \frac{e^{-x}e^x}{x^5}$$

Integrating once gives

$$(e^{-x}y)' = -\frac{1}{4x^4} + c_1$$

Integrating again gives

$$(e^{-x}y) = c_1x + \frac{1}{12x^3} + c_2$$

Hence the solution is

$$y = \frac{c_1x + \frac{1}{12x^3} + c_2}{e^{-x}}$$

Or

$$y = c_1x e^x + c_2e^x + \frac{e^x}{12x^3}$$

Summary

The solution(s) found are the following

$$y = c_1x e^x + c_2e^x + \frac{e^x}{12x^3} \quad (1)$$




Figure 65: Slope field plot

Verification of solutions

$$y = c_1 x e^x + c_2 e^x + \frac{e^x}{12x^3}$$

Verified OK.

4.1.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0 \\ t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 57: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1(e^x)\end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^x$$

$$y_2 = x e^x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^x & x e^x \\ \frac{d}{dx}(e^x) & \frac{d}{dx}(x e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^x & x e^x \\ e^x & x e^x + e^x \end{vmatrix}$$

Therefore

$$W = (e^x)(x e^x + e^x) - (x e^x)(e^x)$$

Which simplifies to

$$W = e^{2x}$$

Which simplifies to

$$W = e^{2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{2x}}{e^{2x}} dx$$

Which simplifies to

$$u_1 = - \int \frac{1}{x^4} dx$$

Hence

$$u_1 = \frac{1}{3x^3}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{2x}}{e^{2x}} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x^5} dx$$

Hence

$$u_2 = -\frac{1}{4x^4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{e^x}{12x^3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + \left(\frac{e^x}{12x^3} \right) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) + \frac{e^x}{12x^3}$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) + \frac{e^x}{12x^3} \quad (1)$$




Figure 66: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + \frac{e^x}{12x^3}$$

Verified OK.

4.1.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = \frac{e^x}{x^5}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 x e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{e^x}{x^5} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & x e^x \\ e^x & x e^x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = e^x \left(- \left(\int \frac{1}{x^4} dx \right) + \left(\int \frac{1}{x^5} dx \right) x \right)$$

- Compute integrals

$$y_p(x) = \frac{e^x}{12x^3}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^x + c_2 x e^x + \frac{e^x}{12x^3}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=exp(x)/x^5,y(x), singsol=all)
```

$$y(x) = \frac{e^x(12c_1x^4 + 12c_2x^3 + 1)}{12x^3}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 25

```
DSolve[y''[x]-2*y'[x]+y[x]==Exp[x]/x^5,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{12}e^x \left(\frac{1}{x^3} + 12c_2x + 12c_1 \right)$$

4.2 problem Problem 12.10

4.2.1	Solving as second order linear constant coeff ode	373
4.2.2	Solving using Kovacic algorithm	377
4.2.3	Maple step by step solution	383

Internal problem ID [5193]

Internal file name [OUTPUT/4686_Sunday_June_05_2022_03_03_22_PM_44270153/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. Supplementary Problems. page 109

Problem number: Problem 12.10.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sec(x)$$

4.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sec(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \sec(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \tan(x) dx$$

Hence

$$u_1 = \ln(\cos(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sec(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int 1 dx$$

Hence

$$u_2 = x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (\ln(\cos(x)) \cos(x) + \sin(x) x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x \quad (1)$$




Figure 67: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Verified OK.

4.2.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \quad (3)$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 59: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \sec(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \tan(x) dx$$

Hence

$$u_1 = \ln(\cos(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sec(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int 1 dx$$

Hence

$$u_2 = x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (\ln(\cos(x)) \cos(x) + \sin(x) x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x \quad (1)$$




Figure 68: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Verified OK.

4.2.3 Maple step by step solution

Let's solve

$$y'' + y = \sec(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sec(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \tan(x) dx \right) + \sin(x) \left(\int 1 dx \right)$$

- Compute integrals

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)+y(x)=sec(x),y(x), singsol=all)
```

$$y(x) = -\ln(\sec(x)) \cos(x) + \cos(x) c_1 + \sin(x) (x + c_2)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 22

```
DSolve[y''[x]+y[x]==Sec[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_2) \sin(x) + \cos(x) (\log(\cos(x)) + c_1)$$

4.3 problem Problem 12.11

4.3.1 Solving as second order linear constant coeff ode	386
4.3.2 Solving using Kovacic algorithm	389
4.3.3 Maple step by step solution	394

Internal problem ID [5194]

Internal file name [OUTPUT/4687_Sunday_June_05_2022_03_03_23_PM_27460064/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. Supplementary Problems. page 109

Problem number: Problem 12.11.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - 2y = e^{3x}$$

4.3.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -2, f(x) = e^{3x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-2)} \\ &= \frac{1}{2} \pm \frac{3}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} + \frac{3}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{3}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1 e^{3x} = e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{3x}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^{-x}) + \left(\frac{e^{3x}}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-x} + \frac{e^{3x}}{4} \quad (1)$$




Figure 69: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-x} + \frac{e^{3x}}{4}$$

Verified OK.

4.3.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -1 \\ C &= -2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 61: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\ &= z_1 e^{\frac{x}{2}} \\ &= z_1 \left(e^{\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{3x}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{3x}}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^{2x}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{2x}}{3}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1e^{3x} = e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{3x}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{-x} + \frac{c_2e^{2x}}{3} \right) + \left(\frac{e^{3x}}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-x} + \frac{c_2e^{2x}}{3} + \frac{e^{3x}}{4} \quad (1)$$




Figure 70: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} + \frac{e^{3x}}{4}$$

Verified OK.

4.3.3 Maple step by step solution

Let's solve

$$y'' - y' - 2y = e^{3x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{3x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{2x} \\ -e^{-x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x}(\int e^{4x} dx)}{3} + \frac{e^{2x}(\int e^x dx)}{3}$$

- Compute integrals

$$y_p(x) = \frac{e^{3x}}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + \frac{e^{3x}}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=exp(3*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^{2x} c_1 + \frac{e^{3x}}{4}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 31

```
DSolve[y''[x]-y'[x]-2*y[x]==Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{3x}}{4} + c_1 e^{-x} + c_2 e^{2x}$$

4.4 problem Problem 12.12

4.4.1	Solving as second order linear constant coeff ode	397
4.4.2	Solving using Kovacic algorithm	400
4.4.3	Maple step by step solution	405

Internal problem ID [5195]

Internal file name [OUTPUT/4688_Sunday_June_05_2022_03_03_24_PM_30692838/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. Supplementary Problems. page 109

Problem number: Problem 12.12.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 60y' - 900y = 5e^{10x}$$

4.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -60, C = -900, f(x) = 5e^{10x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 60y' - 900y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -60, C = -900$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 60\lambda e^{\lambda x} - 900 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 60\lambda - 900 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -60, C = -900$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{60}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-60^2 - (4)(1)(-900)} \\ &= 30 \pm 30\sqrt{2} \end{aligned}$$

Hence

$$\lambda_1 = 30 + 30\sqrt{2}$$

$$\lambda_2 = 30 - 30\sqrt{2}$$

Which simplifies to

$$\lambda_1 = 30 + 30\sqrt{2}$$

$$\lambda_2 = 30 - 30\sqrt{2}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(30+30\sqrt{2})x} + c_2 e^{(30-30\sqrt{2})x}$$

Or

$$y = c_1 e^{(30+30\sqrt{2})x} + c_2 e^{(30-30\sqrt{2})x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{(30+30\sqrt{2})x} + c_2 e^{(30-30\sqrt{2})x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$5 e^{10x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{10x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{(30-30\sqrt{2})x}, e^{(30+30\sqrt{2})x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{10x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-1400A_1 e^{10x} = 5 e^{10x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{280} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{e^{10x}}{280}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{(30+30\sqrt{2})x} + c_2 e^{(30-30\sqrt{2})x} \right) + \left(-\frac{e^{10x}}{280} \right) \end{aligned}$$

Which simplifies to

$$y = c_1 e^{30(1+\sqrt{2})x} + c_2 e^{-30(\sqrt{2}-1)x} - \frac{e^{10x}}{280}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{30(1+\sqrt{2})x} + c_2 e^{-30(\sqrt{2}-1)x} - \frac{e^{10x}}{280} \quad (1)$$




Figure 71: Slope field plot

Verification of solutions

$$y = c_1 e^{30(1+\sqrt{2})x} + c_2 e^{-30(\sqrt{2}-1)x} - \frac{e^{10x}}{280}$$

Verified OK.

4.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 60y' - 900y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= -60 \\C &= -900\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1800}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 1800 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 1800z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 63: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1800$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-30x\sqrt{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-60}{1} dx} \\
 &= z_1 e^{30x} \\
 &= z_1 (e^{30x})
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-30(\sqrt{2}-1)x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{60}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{60x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{2} e^{60x\sqrt{2}}}{120} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-30(\sqrt{2}-1)x} \right) + c_2 \left(e^{-30(\sqrt{2}-1)x} \left(\frac{\sqrt{2} e^{60x\sqrt{2}}}{120} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 60y' - 900y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-30(\sqrt{2}-1)x} + \frac{c_2 \sqrt{2} e^{30(1+\sqrt{2})x}}{120}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$5e^{10x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{10x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sqrt{2} e^{30(1+\sqrt{2})x}}{120}, e^{-30(\sqrt{2}-1)x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{10x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-1400A_1 e^{10x} = 5e^{10x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{280} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{e^{10x}}{280}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-30(\sqrt{2}-1)x} + \frac{c_2 \sqrt{2} e^{30(1+\sqrt{2})x}}{120} \right) + \left(-\frac{e^{10x}}{280} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-30(\sqrt{2}-1)x} + \frac{c_2 \sqrt{2} e^{30(1+\sqrt{2})x}}{120} - \frac{e^{10x}}{280} \quad (1)$$




Figure 72: Slope field plot

Verification of solutions

$$y = c_1 e^{-30(\sqrt{2}-1)x} + \frac{c_2 \sqrt{2} e^{30(1+\sqrt{2})x}}{120} - \frac{e^{10x}}{280}$$

Verified OK.

4.4.3 Maple step by step solution

Let's solve

$$y'' - 60y' - 900y = 5e^{10x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 60r - 900 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{60 \pm (\sqrt{7200})}{2}$$

- Roots of the characteristic polynomial

$$r = (30 - 30\sqrt{2}, 30 + 30\sqrt{2})$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{(30-30\sqrt{2})x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{(30+30\sqrt{2})x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{(30-30\sqrt{2})x} + c_2 e^{(30+30\sqrt{2})x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 5 e^{10x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{(30-30\sqrt{2})x} & e^{(30+30\sqrt{2})x} \\ (30 - 30\sqrt{2}) e^{(30-30\sqrt{2})x} & (30 + 30\sqrt{2}) e^{(30+30\sqrt{2})x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 60\sqrt{2} e^{60x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{\sqrt{2} \left(-e^{-30(\sqrt{2}-1)x} \left(\int e^{30x\sqrt{2}-20x} dx \right) + e^{30(1+\sqrt{2})x} \left(\int e^{-30x\sqrt{2}-20x} dx \right) \right)}{24}$$

- Compute integrals

$$y_p(x) = -\frac{e^{10x}}{280}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{(30-30\sqrt{2})x} + c_2 e^{(30+30\sqrt{2})x} - \frac{e^{10x}}{280}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)-60*diff(y(x),x)-900*y(x)=5*exp(10*x),y(x), singsol=all)
```

$$y(x) = e^{30(1+\sqrt{2})x} c_2 + e^{-30(\sqrt{2}-1)x} c_1 - \frac{e^{10x}}{280}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 45

```
DSolve[y''[x]-60*y'[x]-900*y[x]==5*Exp[10*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^{10x}}{280} + c_1 e^{-30(\sqrt{2}-1)x} + c_2 e^{30(1+\sqrt{2})x}$$

4.5 problem Problem 12.13

4.5.1	Solving as second order linear constant coeff ode	408
4.5.2	Solving as second order integrable as is ode	412
4.5.3	Solving as second order ode missing y ode	414
4.5.4	Solving as type second_order_integrable_as_is (not using ABC version)	415
4.5.5	Solving using Kovacic algorithm	417
4.5.6	Solving as exact linear second order ode ode	422
4.5.7	Maple step by step solution	425

Internal problem ID [5196]

Internal file name [OUTPUT/4689_Sunday_June_05_2022_03_03_25_PM_72658977/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. Supplementary Problems. page 109

Problem number: Problem 12.13.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 7y' = -3$$

4.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -7, C = 0, f(x) = -3$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 7y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -7, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 7\lambda e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 7\lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -7, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{7}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-7^2 - (4)(1)(0)} \\ &= \frac{7}{2} \pm \frac{7}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{7}{2} + \frac{7}{2}$$

$$\lambda_2 = \frac{7}{2} - \frac{7}{2}$$

Which simplifies to

$$\lambda_1 = 7$$

$$\lambda_2 = 0$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(7)x} + c_2 e^{(0)x}$$

Or

$$y = c_1 e^{7x} + c_2$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{7x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{7x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-7A_1 = -3$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{3}{7} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{3x}{7}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 e^{7x} + c_2) + \left(\frac{3x}{7}\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{7x} + c_2 + \frac{3x}{7} \quad (1)$$




Figure 73: Slope field plot

Verification of solutions

$$y = c_1 e^{7x} + c_2 + \frac{3x}{7}$$

Verified OK.

4.5.2 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - 7y') dx = \int (-3) dx$$
$$-7y + y' = -3x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -7$$
$$q(x) = -3x + c_1$$

Hence the ode is

$$-7y + y' = -3x + c_1$$

The integrating factor μ is

$$\mu = e^{\int (-7) dx}$$
$$= e^{-7x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu)(-3x + c_1)$$
$$\frac{d}{dx}(e^{-7x}y) = (e^{-7x})(-3x + c_1)$$
$$d(e^{-7x}y) = ((-3x + c_1)e^{-7x}) dx$$

Integrating gives

$$e^{-7x}y = \int (-3x + c_1)e^{-7x} dx$$
$$e^{-7x}y = \frac{e^{-7x}(21x - 7c_1 + 3)}{49} + c_2$$

Dividing both sides by the integrating factor $\mu = e^{-7x}$ results in

$$y = \frac{e^{7x}e^{-7x}(21x - 7c_1 + 3)}{49} + c_2e^{7x}$$

which simplifies to

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2 e^{7x}$$

Summary

The solution(s) found are the following

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2 e^{7x} \tag{1}$$




Figure 74: Slope field plot

Verification of solutions

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2 e^{7x}$$

Verified OK.

4.5.3 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 7p(x) + 3 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\int \frac{1}{7p-3} dp = \int dx$$
$$\frac{\ln(7p-3)}{7} = x + c_1$$

Raising both side to exponential gives

$$(7p-3)^{\frac{1}{7}} = e^{x+c_1}$$

Which simplifies to

$$(7p-3)^{\frac{1}{7}} = c_2 e^x$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{c_2^7 e^{7x}}{7} + \frac{3}{7}$$

Integrating both sides gives

$$y = \int \frac{c_2^7 e^{7x}}{7} + \frac{3}{7} dx$$
$$= \frac{3x}{7} + \frac{c_2^7 e^{7x}}{49} + c_3$$

Summary

The solution(s) found are the following

$$y = \frac{3x}{7} + \frac{c_2^7 e^{7x}}{49} + c_3 \quad (1)$$




Figure 75: Slope field plot

Verification of solutions

$$y = \frac{3x}{7} + \frac{c_2^7 e^{7x}}{49} + c_3$$

Verified OK.

4.5.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$y'' - 7y' = -3$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - 7y') dx = \int (-3) dx$$

$$-7y + y' = -3x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -7 \\q(x) &= -3x + c_1\end{aligned}$$

Hence the ode is

$$-7y + y' = -3x + c_1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-7) dx} \\ &= e^{-7x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(-3x + c_1) \\ \frac{d}{dx}(e^{-7x}y) &= (e^{-7x})(-3x + c_1) \\ d(e^{-7x}y) &= ((-3x + c_1)e^{-7x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-7x}y &= \int (-3x + c_1)e^{-7x} dx \\ e^{-7x}y &= \frac{e^{-7x}(21x - 7c_1 + 3)}{49} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-7x}$ results in

$$y = \frac{e^{7x}e^{-7x}(21x - 7c_1 + 3)}{49} + c_2e^{7x}$$

which simplifies to

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2e^{7x}$$

Summary

The solution(s) found are the following

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2e^{7x} \tag{1}$$




Figure 76: Slope field plot

Verification of solutions

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2 e^{7x}$$

Verified OK.

4.5.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 7y' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -7 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{49}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 49 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{49z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 65: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{49}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{7x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-7}{1} dx} \\ &= z_1 e^{\frac{7x}{2}} \\ &= z_1 \left(e^{\frac{7x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-7}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{7x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{7x}}{7} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2 \left(1 \left(\frac{e^{7x}}{7} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 7y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 + \frac{c_2 e^{7x}}{7}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ 1, \frac{e^{7x}}{7} \right\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-7A_1 = -3$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{3}{7} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{3x}{7}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 + \frac{c_2 e^{7x}}{7} \right) + \left(\frac{3x}{7} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + \frac{c_2 e^{7x}}{7} + \frac{3x}{7} \tag{1}$$




Figure 77: Slope field plot

Verification of solutions

$$y = c_1 + \frac{c_2 e^{7x}}{7} + \frac{3x}{7}$$

Verified OK.

4.5.6 Solving as exact linear second order ode ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= -7 \\ r(x) &= 0 \\ s(x) &= -3 \end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$-7y + y' = \int -3 dx$$

We now have a first order ode to solve which is

$$-7y + y' = -3x + c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -7 \\q(x) &= -3x + c_1\end{aligned}$$

Hence the ode is

$$-7y + y' = -3x + c_1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-7)dx} \\&= e^{-7x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(-3x + c_1) \\ \frac{d}{dx}(e^{-7x}y) &= (e^{-7x})(-3x + c_1) \\ d(e^{-7x}y) &= ((-3x + c_1)e^{-7x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-7x}y &= \int (-3x + c_1)e^{-7x} dx \\ e^{-7x}y &= \frac{e^{-7x}(21x - 7c_1 + 3)}{49} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-7x}$ results in

$$y = \frac{e^{7x}e^{-7x}(21x - 7c_1 + 3)}{49} + c_2e^{7x}$$

which simplifies to

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2e^{7x}$$

Summary

The solution(s) found are the following

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2e^{7x} \tag{1}$$




Figure 78: Slope field plot

Verification of solutions

$$y = \frac{3x}{7} - \frac{c_1}{7} + \frac{3}{49} + c_2 e^{7x}$$

Verified OK.

4.5.7 Maple step by step solution

Let's solve

$$y'' - 7y' = -3$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 7r = 0$$

- Factor the characteristic polynomial

$$r(r - 7) = 0$$

- Roots of the characteristic polynomial

$$r = (0, 7)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{7x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 e^{7x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = -3 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & e^{7x} \\ 0 & 7e^{7x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 7e^{7x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{3(\int 1 dx)}{7} - \frac{3e^{7x}(\int e^{-7x} dx)}{7}$$

- Compute integrals

$$y_p(x) = \frac{3x}{7} + \frac{3}{49}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 + c_2 e^{7x} + \frac{3x}{7} + \frac{3}{49}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = 7*_b(_a)-3, _b(_a)` *** Sublevel 2 **  
  Methods for first order ODEs:  
    --- Trying classification methods ---  
    trying a quadrature  
    trying 1st order linear  
    <- 1st order linear successful  
  <- high order exact linear fully integrable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x$2)-7*diff(y(x),x)=-3,y(x), singsol=all)
```

$$y(x) = \frac{e^{7x}c_1}{7} + \frac{3x}{7} + c_2$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 24

```
DSolve[y''[x]-7*y'[x]==-3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x}{7} + \frac{1}{7}c_1e^{7x} + c_2$$

4.6 problem Problem 12.14

4.6.1	Solving as second order euler ode ode	429
4.6.2	Solving as second order change of variable on x method 2 ode .	432
4.6.3	Solving as second order change of variable on x method 1 ode .	438
4.6.4	Solving as second order change of variable on y method 2 ode .	442
4.6.5	Solving as second order integrable as is ode	447
4.6.6	Solving as second order ode non constant coeff transformation on B ode	449
4.6.7	Solving as type second_order_integrable_as_is (not using ABC version)	453
4.6.8	Solving using Kovacic algorithm	455
4.6.9	Solving as exact linear second order ode ode	463

Internal problem ID [5197]

Internal file name [OUTPUT/4690_Sunday_June_05_2022_03_03_26_PM_86935249/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. Supplementary Problems. page 109

Problem number: Problem 12.14.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "exact linear second order ode", "second_order_integrable_as_is", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _nonhomogeneous]]
```

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = \ln(x)$$

The ode can be written as

$$x^2 y'' + x y' - y = \ln(x) x^2$$

Which shows it is a Euler ODE.

4.6.1 Solving as second order euler ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = x, C = -1, f(x) = \ln(x) x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + xy' - y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + xx^{r-1} - x^r = 0$$

Simplifying gives

$$r(r-1)x^r + rx^r - x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + r - 1 = 0$$

Or

$$r^2 - 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -1$$

$$r_2 = 1$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x} + c_2x$$

Next, we find the particular solution to the ODE

$$x^2y'' + xy' - y = \ln(x) x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \frac{1}{x} \\ y_2 &= x \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x} & x \\ \frac{d}{dx}\left(\frac{1}{x}\right) & \frac{d}{dx}(x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x} & x \\ -\frac{1}{x^2} & 1 \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x}\right)(1) - (x)\left(-\frac{1}{x^2}\right)$$

Which simplifies to

$$W = \frac{2}{x}$$

Which simplifies to

$$W = \frac{2}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 \ln(x)}{2x} dx$$

Which simplifies to

$$u_1 = - \int \frac{\ln(x) x^2}{2} dx$$

Hence

$$u_1 = - \frac{x^3 \ln(x)}{6} + \frac{x^3}{18}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\ln(x) x}{2x} dx$$

Which simplifies to

$$u_2 = \int \frac{\ln(x)}{2} dx$$

Hence

$$u_2 = \frac{\ln(x) x}{2} - \frac{x}{2}$$

Which simplifies to

$$u_1 = - \frac{x^3(-1 + 3 \ln(x))}{18}$$
$$u_2 = \frac{x(\ln(x) - 1)}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{x^2(-1 + 3 \ln(x))}{18} + \frac{x^2(\ln(x) - 1)}{2}$$

Which simplifies to

$$y_p(x) = \frac{x^2(-4 + 3 \ln(x))}{9}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \frac{x^2(-4 + 3 \ln(x))}{9} + \frac{c_1}{x} + c_2x \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2(-4 + 3 \ln(x))}{9} + \frac{c_1}{x} + c_2x \quad (1)$$

Verification of solutions

$$y = \frac{x^2(-4 + 3 \ln(x))}{9} + \frac{c_1}{x} + c_2x$$

Verified OK.

4.6.2 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$x^2y'' + xy' - y = 0$$

In normal form the ode

$$x^2y'' + xy' - y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x} dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{1}{x^2}}{\frac{1}{x^2}} \\ &= -1 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - y(\tau) &= 0\end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1\end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(1)\tau} + c_2 e^{(-1)\tau}$$

Or

$$y(\tau) = c_1 e^\tau + c_2 e^{-\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 x^2 + c_2}{x}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 x^2 + c_2}{x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = \frac{1}{x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & \frac{1}{x} \\ \frac{d}{dx}(x) & \frac{d}{dx}\left(\frac{1}{x}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & \frac{1}{x} \\ 1 & -\frac{1}{x^2} \end{vmatrix}$$

Therefore

$$W = (x) \left(-\frac{1}{x^2} \right) - \left(\frac{1}{x} \right) \quad (1)$$

Which simplifies to

$$W = -\frac{2}{x}$$

Which simplifies to

$$W = -\frac{2}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\ln(x) x}{-2x} dx$$

Which simplifies to

$$u_1 = - \int -\frac{\ln(x)}{2} dx$$

Hence

$$u_1 = \frac{\ln(x) x}{2} - \frac{x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 \ln(x)}{-2x} dx$$

Which simplifies to

$$u_2 = \int -\frac{\ln(x) x^2}{2} dx$$

Hence

$$u_2 = -\frac{x^3 \ln(x)}{6} + \frac{x^3}{18}$$

Which simplifies to

$$u_1 = \frac{x(\ln(x) - 1)}{2}$$
$$u_2 = -\frac{x^3(-1 + 3 \ln(x))}{18}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2(-1 + 3 \ln(x))}{18} + \frac{x^2(\ln(x) - 1)}{2}$$

Which simplifies to

$$y_p(x) = \frac{x^2(-4 + 3 \ln(x))}{9}$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= \left(\frac{c_1 x^2 + c_2}{x} \right) + \left(\frac{x^2(-4 + 3 \ln(x))}{9} \right)$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^2 + c_2}{x} + \frac{x^2(-4 + 3 \ln(x))}{9} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 x^2 + c_2}{x} + \frac{x^2(-4 + 3 \ln(x))}{9}$$

Verified OK.

4.6.3 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = x, C = -1, f(x) = \ln(x) x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + xy' - y = 0$$

In normal form the ode

$$x^2y'' + xy' - y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{1}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{\sqrt{-\frac{1}{x^2}}}{c} \tag{6}$$

$$\tau'' = \frac{1}{c\sqrt{-\frac{1}{x^2}}x^3}$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \\
 &= \frac{\frac{1}{c\sqrt{-\frac{1}{x^2}}x^3} + \frac{1}{x} \frac{\sqrt{-\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{-\frac{1}{x^2}}}{c}\right)^2} \\
 &= 0
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c} \sqrt{q} dx \\
 &= \frac{\int \sqrt{-\frac{1}{x^2}} dx}{c} \\
 &= \frac{\sqrt{-\frac{1}{x^2}} x \ln(x)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = \frac{(ic_2 + c_1) x^2 - ic_2 + c_1}{2x}$$

Now the particular solution to this ODE is found

$$x^2 y'' + xy' - y = \ln(x) x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = \frac{1}{x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & \frac{1}{x} \\ \frac{d}{dx}(x) & \frac{d}{dx}\left(\frac{1}{x}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & \frac{1}{x} \\ 1 & -\frac{1}{x^2} \end{vmatrix}$$

Therefore

$$W = (x) \left(-\frac{1}{x^2}\right) - \left(\frac{1}{x}\right) \quad (1)$$

Which simplifies to

$$W = -\frac{2}{x}$$

Which simplifies to

$$W = -\frac{2}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\ln(x) x}{-2x} dx$$

Which simplifies to

$$u_1 = - \int -\frac{\ln(x)}{2} dx$$

Hence

$$u_1 = \frac{\ln(x) x}{2} - \frac{x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 \ln(x)}{-2x} dx$$

Which simplifies to

$$u_2 = \int -\frac{\ln(x) x^2}{2} dx$$

Hence

$$u_2 = -\frac{x^3 \ln(x)}{6} + \frac{x^3}{18}$$

Which simplifies to

$$u_1 = \frac{x(\ln(x) - 1)}{2}$$
$$u_2 = -\frac{x^3(-1 + 3 \ln(x))}{18}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2(-1 + 3 \ln(x))}{18} + \frac{x^2(\ln(x) - 1)}{2}$$

Which simplifies to

$$y_p(x) = \frac{x^2(-4 + 3 \ln(x))}{9}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{(ic_2 + c_1)x^2 - ic_2 + c_1}{2x} \right) + \left(\frac{x^2(-4 + 3 \ln(x))}{9} \right) \\ &= \frac{x^2(-4 + 3 \ln(x))}{9} + \frac{(ic_2 + c_1)x^2 - ic_2 + c_1}{2x} \end{aligned}$$

Which simplifies to

$$y = \frac{6x^3 \ln(x) + 9ic_2x^2 + 9c_1x^2 - 8x^3 - 9ic_2 + 9c_1}{18x}$$

Summary

The solution(s) found are the following

$$y = \frac{6x^3 \ln(x) + 9ic_2x^2 + 9c_1x^2 - 8x^3 - 9ic_2 + 9c_1}{18x} \quad (1)$$

Verification of solutions

$$y = \frac{6x^3 \ln(x) + 9ic_2x^2 + 9c_1x^2 - 8x^3 - 9ic_2 + 9c_1}{18x}$$

Verified OK.

4.6.4 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = x$, $C = -1$, $f(x) = \ln(x)x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + xy' - y = 0$$

In normal form the ode

$$x^2 y'' + xy' - y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{1}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 1 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \frac{3v'(x)}{x} = 0$$
$$v''(x) + \frac{3v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{3u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{3u}{x} \end{aligned}$$

Where $f(x) = -\frac{3}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{3}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{3}{x} dx \\ \ln(u) &= -3 \ln(x) + c_1 \\ u &= e^{-3 \ln(x) + c_1} \\ &= \frac{c_1}{x^3} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{2x^2} + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= \left(-\frac{c_1}{2x^2} + c_2 \right) x \\ &= \left(-\frac{c_1}{2x^2} + c_2 \right) x \end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' + xy' - y = \ln(x) x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = \frac{1}{x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & \frac{1}{x} \\ \frac{d}{dx}(x) & \frac{d}{dx}\left(\frac{1}{x}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & \frac{1}{x} \\ 1 & -\frac{1}{x^2} \end{vmatrix}$$

Therefore

$$W = (x) \left(-\frac{1}{x^2}\right) - \left(\frac{1}{x}\right) \quad (1)$$

Which simplifies to

$$W = -\frac{2}{x}$$

Which simplifies to

$$W = -\frac{2}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\ln(x) x}{-2x} dx$$

Which simplifies to

$$u_1 = - \int -\frac{\ln(x)}{2} dx$$

Hence

$$u_1 = \frac{\ln(x) x}{2} - \frac{x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 \ln(x)}{-2x} dx$$

Which simplifies to

$$u_2 = \int -\frac{\ln(x) x^2}{2} dx$$

Hence

$$u_2 = -\frac{x^3 \ln(x)}{6} + \frac{x^3}{18}$$

Which simplifies to

$$u_1 = \frac{x(\ln(x) - 1)}{2}$$
$$u_2 = -\frac{x^3(-1 + 3 \ln(x))}{18}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2(-1 + 3 \ln(x))}{18} + \frac{x^2(\ln(x) - 1)}{2}$$

Which simplifies to

$$y_p(x) = \frac{x^2(-4 + 3 \ln(x))}{9}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\left(-\frac{c_1}{2x^2} + c_2 \right) x \right) + \left(\frac{x^2(-4 + 3 \ln(x))}{9} \right) \\ &= \frac{x^2(-4 + 3 \ln(x))}{9} + \left(-\frac{c_1}{2x^2} + c_2 \right) x \end{aligned}$$

Which simplifies to

$$y = \frac{6x^3 \ln(x) + 18c_2x^2 - 8x^3 - 9c_1}{18x}$$

Summary

The solution(s) found are the following

$$y = \frac{6x^3 \ln(x) + 18c_2x^2 - 8x^3 - 9c_1}{18x} \tag{1}$$

Verification of solutions

$$y = \frac{6x^3 \ln(x) + 18c_2x^2 - 8x^3 - 9c_1}{18x}$$

Verified OK.

4.6.5 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned} \int (x^2y'' + xy' - y) dx &= \int \ln(x) x^2 dx \\ x^2y' - xy &= \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} + c_1 \end{aligned}$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x} \right) = \left(\frac{1}{x} \right) \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2} \right)$$
$$d \left(\frac{y}{x} \right) = \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^3} \right) dx$$

Integrating gives

$$\frac{y}{x} = \int \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^3} dx$$
$$\frac{y}{x} = \frac{\ln(x) x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = x \left(\frac{\ln(x) x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} \right) + c_2 x$$

which simplifies to

$$y = x \left(\frac{\ln(x) x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right)$$

Summary

The solution(s) found are the following

$$y = x \left(\frac{\ln(x) x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right) \quad (1)$$

Verification of solutions

$$y = x \left(\frac{\ln(x)x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right)$$

Verified OK.

4.6.6 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned} A &= x^2 \\ B &= x \\ C &= -1 \\ F &= \ln(x)x^2 \end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x^2)(0) + (x)(1) + (-1)(x) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$x^3 v'' + (3x^2) v' = 0$$

Now by applying $v' = u$ the above becomes

$$x^2(u'(x)x + 3u(x)) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{3u}{x} \end{aligned}$$

Where $f(x) = -\frac{3}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{3}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{3}{x} dx \\ \ln(u) &= -3 \ln(x) + c_1 \\ u &= e^{-3 \ln(x) + c_1} \\ &= \frac{c_1}{x^3} \end{aligned}$$

The ode for v now becomes

$$\begin{aligned} v' &= u \\ &= \frac{c_1}{x^3} \end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned} v(x) &= \int \frac{c_1}{x^3} dx \\ &= -\frac{c_1}{2x^2} + c_2 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= Bv \\ &= (x) \left(-\frac{c_1}{2x^2} + c_2 \right) \\ &= \left(-\frac{c_1}{2x^2} + c_2 \right) x \end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= x \\ y_2 &= \frac{1}{x} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & \frac{1}{x} \\ \frac{d}{dx}(x) & \frac{d}{dx}\left(\frac{1}{x}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & \frac{1}{x} \\ 1 & -\frac{1}{x^2} \end{vmatrix}$$

Therefore

$$W = (x) \left(-\frac{1}{x^2} \right) - \left(\frac{1}{x} \right) \quad (1)$$

Which simplifies to

$$W = -\frac{2}{x}$$

Which simplifies to

$$W = -\frac{2}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\ln(x) x}{-2x} dx$$

Which simplifies to

$$u_1 = - \int -\frac{\ln(x)}{2} dx$$

Hence

$$u_1 = \frac{\ln(x) x}{2} - \frac{x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 \ln(x)}{-2x} dx$$

Which simplifies to

$$u_2 = \int -\frac{\ln(x) x^2}{2} dx$$

Hence

$$u_2 = -\frac{x^3 \ln(x)}{6} + \frac{x^3}{18}$$

Which simplifies to

$$u_1 = \frac{x(\ln(x) - 1)}{2}$$
$$u_2 = -\frac{x^3(-1 + 3 \ln(x))}{18}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2(-1 + 3 \ln(x))}{18} + \frac{x^2(\ln(x) - 1)}{2}$$

Which simplifies to

$$y_p(x) = \frac{x^2(-4 + 3 \ln(x))}{9}$$

Hence the complete solution is

$$\begin{aligned} y(x) &= y_h + y_p \\ &= \left(\left(-\frac{c_1}{2x^2} + c_2 \right) x \right) + \left(\frac{x^2(-4 + 3 \ln(x))}{9} \right) \\ &= \frac{6x^3 \ln(x) + 18c_2x^2 - 8x^3 - 9c_1}{18x} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{6x^3 \ln(x) + 18c_2x^2 - 8x^3 - 9c_1}{18x} \quad (1)$$

Verification of solutions

$$y = \frac{6x^3 \ln(x) + 18c_2x^2 - 8x^3 - 9c_1}{18x}$$

Verified OK.

4.6.7 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$x^2y'' + xy' - y = \ln(x) x^2$$

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned} \int (x^2y'' + xy' - y) dx &= \int \ln(x) x^2 dx \\ x^2y' - xy &= \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} + c_1 \end{aligned}$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x} \right) = \left(\frac{1}{x} \right) \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2} \right)$$
$$d \left(\frac{y}{x} \right) = \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^3} \right) dx$$

Integrating gives

$$\frac{y}{x} = \int \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^3} dx$$
$$\frac{y}{x} = \frac{\ln(x)x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = x \left(\frac{\ln(x)x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} \right) + c_2 x$$

which simplifies to

$$y = x \left(\frac{\ln(x)x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right)$$

Summary

The solution(s) found are the following

$$y = x \left(\frac{\ln(x) x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right) \quad (1)$$

Verification of solutions

$$y = x \left(\frac{\ln(x) x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right)$$

Verified OK.

4.6.8 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + xy' - y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= x \\ C &= -1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 3 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{3}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 67: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole

larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{3}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + (-)(0) \\ &= -\frac{1}{2x} \\ &= -\frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{2x}\right)(0) + \left(\left(\frac{1}{2x^2}\right) + \left(-\frac{1}{2x}\right)^2 - \left(\frac{3}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(x) = pe^{\int \omega dx}$$

$$= e^{\int -\frac{1}{2x} dx}$$

$$= \frac{1}{\sqrt{x}}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

$$= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx}$$

$$= z_1 e^{-\frac{\ln(x)}{2}}$$

$$= z_1 \left(\frac{1}{\sqrt{x}}\right)$$

Which simplifies to

$$y_1 = \frac{1}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx$$

$$= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx$$

$$= y_1 \left(\frac{x^2}{2}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x} \right) + c_2 \left(\frac{1}{x} \left(\frac{x^2}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2 y'' + x y' - y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1}{x} + \frac{c_2 x}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= \frac{1}{x} \\ y_2 &= \frac{x}{2}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x} & \frac{x}{2} \\ \frac{d}{dx} \left(\frac{1}{x} \right) & \frac{d}{dx} \left(\frac{x}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x} & \frac{x}{2} \\ -\frac{1}{x^2} & \frac{1}{2} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x} \right) \left(\frac{1}{2} \right) - \left(\frac{x}{2} \right) \left(-\frac{1}{x^2} \right)$$

Which simplifies to

$$W = \frac{1}{x}$$

Which simplifies to

$$W = \frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{x^3 \ln(x)}{2}}{x} dx$$

Which simplifies to

$$u_1 = - \int \frac{\ln(x) x^2}{2} dx$$

Hence

$$u_1 = -\frac{x^3 \ln(x)}{6} + \frac{x^3}{18}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\ln(x) x}{x} dx$$

Which simplifies to

$$u_2 = \int \ln(x) dx$$

Hence

$$u_2 = \ln(x)x - x$$

Which simplifies to

$$u_1 = -\frac{x^3(-1 + 3 \ln(x))}{18}$$
$$u_2 = x(\ln(x) - 1)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2(-1 + 3 \ln(x))}{18} + \frac{x^2(\ln(x) - 1)}{2}$$

Which simplifies to

$$y_p(x) = \frac{x^2(-4 + 3 \ln(x))}{9}$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= \left(\frac{c_1}{x} + \frac{c_2 x}{2} \right) + \left(\frac{x^2(-4 + 3 \ln(x))}{9} \right)$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} + \frac{c_2 x}{2} + \frac{x^2(-4 + 3 \ln(x))}{9} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x} + \frac{c_2 x}{2} + \frac{x^2(-4 + 3 \ln(x))}{9}$$

Verified OK.

4.6.9 Solving as exact linear second order ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= x^2 \\ q(x) &= x \\ r(x) &= -1 \\ s(x) &= \ln(x) x^2 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 2 \\ q'(x) &= 1 \end{aligned}$$

Therefore (1) becomes

$$2 - (1) + (-1) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x) y' + (q(x) - p'(x)) y)' = s(x)$$

Integrating gives

$$p(x) y' + (q(x) - p'(x)) y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$x^2 y' - xy = \int \ln(x) x^2 dx$$

We now have a first order ode to solve which is

$$x^2 y' - xy = \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} + c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x} \right) = \left(\frac{1}{x} \right) \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^2} \right)$$
$$d \left(\frac{y}{x} \right) = \left(\frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^3} \right) dx$$

Integrating gives

$$\frac{y}{x} = \int \frac{3x^3 \ln(x) - x^3 + 9c_1}{9x^3} dx$$
$$\frac{y}{x} = \frac{\ln(x)x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = x \left(\frac{\ln(x)x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} \right) + c_2 x$$

which simplifies to

$$y = x \left(\frac{\ln(x)x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right)$$

Summary

The solution(s) found are the following

$$y = x \left(\frac{\ln(x) x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right) \quad (1)$$

Verification of solutions

$$y = x \left(\frac{\ln(x) x}{3} - \frac{4x}{9} - \frac{c_1}{2x^2} + c_2 \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
<- high order exact linear fully integrable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+1/x*diff(y(x),x)-1/x^2*y(x)=ln(x),y(x), singsol=all)
```

$$y(x) = c_1 x + \frac{c_2}{x} + \frac{x^2(3 \ln(x) - 4)}{9}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 32

```
DSolve[y''[x]+1/x*y'[x]-1/x^2*y[x]==Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{4x^2}{9} + \frac{1}{3}x^2 \log(x) + c_2 x + \frac{c_1}{x}$$

4.7 problem Problem 12.15

4.7.1	Solving as second order euler ode ode	466
4.7.2	Solving as second order ode missing y ode	470
4.7.3	Solving using Kovacic algorithm	471
4.7.4	Maple step by step solution	479

Internal problem ID [5198]

Internal file name [OUTPUT/4691_Sunday_June_05_2022_03_03_27_PM_42950579/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. Supplementary Problems. page 109

Problem number: Problem 12.15.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$x^2y'' - xy' = x^3e^x$$

4.7.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -x$, $C = 0$, $f(x) = x^3e^x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - xy' = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - xx^{r-1} + 0 = 0$$

Simplifying gives

$$r(r-1)x^r - rx^r + 0 = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) - r + 0 = 0$$

Or

$$r^2 - 2r = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 0$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_2x^2 + c_1$$

Next, we find the particular solution to the ODE

$$x^2y'' - xy' = x^3e^x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = 1$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & x^2 \\ \frac{d}{dx}(1) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & x^2 \\ 0 & 2x \end{vmatrix}$$

Therefore

$$W = (1)(2x) - (x^2)(0)$$

Which simplifies to

$$W = 2x$$

Which simplifies to

$$W = 2x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^5 e^x}{2x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^2 e^x}{2} dx$$

Hence

$$u_1 = - \frac{(x^2 - 2x + 2) e^x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 e^x}{2x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{e^x}{2} dx$$

Hence

$$u_2 = \frac{e^x}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{(x^2 - 2x + 2) e^x}{2} + \frac{x^2 e^x}{2}$$

Which simplifies to

$$y_p(x) = (x - 1) e^x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (x - 1) e^x + c_2 x^2 + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = (x - 1) e^x + c_2 x^2 + c_1 \quad (1)$$

Verification of solutions

$$y = (x - 1) e^x + c_2 x^2 + c_1$$

Verified OK.

4.7.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$x^2 p'(x) - xp(x) - x^3 e^x = 0$$

Which is now solve for $p(x)$ as first order ode.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(x) + p(x)p(x) = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = x e^x$$

Hence the ode is

$$p'(x) - \frac{p(x)}{x} = x e^x$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu p) = (\mu) (x e^x)$$
$$\frac{d}{dx}\left(\frac{p}{x}\right) = \left(\frac{1}{x}\right) (x e^x)$$
$$d\left(\frac{p}{x}\right) = e^x dx$$

Integrating gives

$$\frac{p}{x} = \int e^x dx$$
$$\frac{p}{x} = e^x + c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$p(x) = x e^x + c_1 x$$

which simplifies to

$$p(x) = x(e^x + c_1)$$

Since $p = y'$ then the new first order ode to solve is

$$y' = x(e^x + c_1)$$

Integrating both sides gives

$$y = \int x(e^x + c_1) dx$$
$$= \frac{c_1 x^2}{2} + x e^x - e^x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^2}{2} + x e^x - e^x + c_2 \quad (1)$$

Verification of solutions

$$y = \frac{c_1 x^2}{2} + x e^x - e^x + c_2$$

Verified OK.

4.7.3 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' - x y' = 0 \quad (1)$$

$$A y'' + B y' + C y = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\B &= -x \\C &= 0\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{4x^2}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 3 \\t &= 4x^2\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{3}{4x^2}\right) z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 68: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 2 - 0 \\
 &= 2
 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition

of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{3}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= -\frac{1}{2x} + (-)(0) \\ &= -\frac{1}{2x} \\ &= -\frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{1}{2x}\right)(0) + \left(\left(\frac{1}{2x^2}\right) + \left(-\frac{1}{2x}\right)^2 - \left(\frac{3}{4x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{2x} dx} \\ &= \frac{1}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{-x}{x^2} dx} \\&= z_1 e^{\frac{\ln(x)}{2}} \\&= z_1 (\sqrt{x})\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{x^2}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 (1) + c_2 \left(1 \left(\frac{x^2}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2 y'' - x y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 + \frac{c_2 x^2}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = 1$$

$$y_2 = \frac{x^2}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & \frac{x^2}{2} \\ \frac{d}{dx}(1) & \frac{d}{dx}\left(\frac{x^2}{2}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & \frac{x^2}{2} \\ 0 & x \end{vmatrix}$$

Therefore

$$W = (1)(x) - \left(\frac{x^2}{2}\right)(0)$$

Which simplifies to

$$W = x$$

Which simplifies to

$$W = x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{x^5 e^x}{2}}{x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^2 e^x}{2} dx$$

Hence

$$u_1 = - \frac{(x^2 - 2x + 2) e^x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 e^x}{x^3} dx$$

Which simplifies to

$$u_2 = \int e^x dx$$

Hence

$$u_2 = e^x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{(x^2 - 2x + 2) e^x}{2} + \frac{x^2 e^x}{2}$$

Which simplifies to

$$y_p(x) = (x - 1) e^x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 + \frac{c_2 x^2}{2} \right) + ((x - 1) e^x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + \frac{c_2 x^2}{2} + (x - 1) e^x \quad (1)$$

Verification of solutions

$$y = c_1 + \frac{c_2 x^2}{2} + (x - 1) e^x$$

Verified OK.

4.7.4 Maple step by step solution

Let's solve

$$x^2 y'' - x y' = x^3 e^x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$x^2 u'(x) - x u(x) = x^3 e^x$$

- Isolate the derivative

$$u'(x) = \frac{u(x)}{x} + x e^x$$

- Group terms with $u(x)$ on the lhs of the ODE and the rest on the rhs of the ODE

$$u'(x) - \frac{u(x)}{x} = x e^x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(u'(x) - \frac{u(x)}{x} \right) = \mu(x) x e^x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) u(x))$

$$\mu(x) \left(u'(x) - \frac{u(x)}{x} \right) = \mu'(x) u(x) + \mu(x) u'(x)$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx} (\mu(x) u(x)) \right) dx = \int \mu(x) x e^x dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) u(x) = \int \mu(x) x e^x dx + c_1$$

- Solve for $u(x)$

$$u(x) = \frac{\int \mu(x) x e^x dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{x}$

$$u(x) = x \left(\int e^x dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$u(x) = x(e^x + c_1)$$

- Solve 1st ODE for $u(x)$

$$u(x) = x(e^x + c_1)$$

- Make substitution $u = y'$

$$y' = x(e^x + c_1)$$

- Integrate both sides to solve for y

$$\int y' dx = \int x(e^x + c_1) dx + c_2$$

- Compute integrals

$$y = \frac{c_1 x^2}{2} + x e^x - e^x + c_2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = (exp(_a)*_a^2+_b(_a))/_a, _b(_a)` ***
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)-x*diff(y(x),x)=x^3*exp(x),y(x), singsol=all)
```

$$y(x) = (x - 1)e^x + \frac{c_1 x^2}{2} + c_2$$

✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 24

```
DSolve[x^2*y'[x]-x*y'[x]==x^3*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 x^2}{2} + e^x(x - 1) + c_2$$

4.8 problem Problem 12.16

4.8.1	Solving as linear ode	482
4.8.2	Solving as homogeneousTypeD2 ode	484
4.8.3	Solving as first order ode lie symmetry lookup ode	485
4.8.4	Solving as exact ode	489
4.8.5	Maple step by step solution	494

Internal problem ID [5199]

Internal file name [OUTPUT/4692_Sunday_June_05_2022_03_03_28_PM_94374798/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 12. VARIATION OF PARAMETERS. Supplementary Problems. page 109

Problem number: Problem 12.16.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**homogeneousTypeD2**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

[_linear]

$$y' - \frac{y}{x} = x^2$$

4.8.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = x^2$$

Hence the ode is

$$y' - \frac{y}{x} = x^2$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x^2) \\ \frac{d}{dx}\left(\frac{y}{x}\right) &= \left(\frac{1}{x}\right)(x^2) \\ d\left(\frac{y}{x}\right) &= x dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{x} &= \int x dx \\ \frac{y}{x} &= \frac{x^2}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = \frac{1}{2}x^3 + c_1x$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}x^3 + c_1x \tag{1}$$




Figure 79: Slope field plot

Verification of solutions

$$y = \frac{1}{2}x^3 + c_1x$$

Verified OK.

4.8.2 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x = x^2$$

Integrating both sides gives

$$\begin{aligned} u(x) &= \int x \, dx \\ &= \frac{x^2}{2} + c_2 \end{aligned}$$

Therefore the solution y is

$$\begin{aligned} y &= xu \\ &= x \left(\frac{x^2}{2} + c_2 \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x \left(\frac{x^2}{2} + c_2 \right) \quad (1)$$




Figure 80: Slope field plot

Verification of solutions

$$y = x \left(\frac{x^2}{2} + c_2 \right)$$

Verified OK.

4.8.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{x^3 + y}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 70: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x} dy \end{aligned}$$

Which results in

$$S = \frac{y}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x^3 + y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{y}{x^2} \\ S_y &= \frac{1}{x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^2}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x} = \frac{x^2}{2} + c_1$$



Which simplifies to

$$\frac{y}{x} = \frac{x^2}{2} + c_1$$

Which gives

$$y = \frac{x(x^2 + 2c_1)}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x^3 + y}{x}$ 	$R = x$ $S = \frac{y}{x}$	$\frac{dS}{dR} = R$ 

Summary

The solution(s) found are the following

$$y = \frac{x(x^2 + 2c_1)}{2} \quad (1)$$




Figure 81: Slope field plot

Verification of solutions

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Verified OK.

4.8.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= \left(\frac{y}{x} + x^2 \right) dx \\ \left(-\frac{y}{x} - x^2 \right) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{y}{x} - x^2 \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{y}{x} - x^2 \right) \\ &= -\frac{1}{x} \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1 \left(\left(-\frac{1}{x} \right) - (0) \right) \\ &= -\frac{1}{x}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int -\frac{1}{x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-\ln(x)} \\ &= \frac{1}{x}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{x} \left(-\frac{y}{x} - x^2 \right) \\ &= \frac{-x^3 - y}{x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{x}(1) \\ &= \frac{1}{x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(\frac{-x^3 - y}{x^2} \right) + \left(\frac{1}{x} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-x^3 - y}{x^2} dx \\ \phi &= \frac{-x^3 + 2y}{2x} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{1}{x} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{x}$. Therefore equation (4) becomes

$$\frac{1}{x} = \frac{1}{x} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{-x^3 + 2y}{2x} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{-x^3 + 2y}{2x}$$

The solution becomes

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{x(x^2 + 2c_1)}{2} \tag{1}$$




Figure 82: Slope field plot

Verification of solutions

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Verified OK.

4.8.5 Maple step by step solution

Let's solve

$$y' - \frac{y}{x} = x^2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = \frac{y}{x} + x^2$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{y}{x} = x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu(x) x^2$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x^2 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x^2 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^2 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{x}$

$$y = x(\int x dx + c_1)$$

- Evaluate the integrals on the rhs

$$y = x\left(\frac{x^2}{2} + c_1\right)$$

- Simplify

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)-1/x*y(x)=x^2,y(x), singsol=all)
```

$$y(x) = \frac{x(x^2 + 2c_1)}{2}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 17

```
DSolve[y'[x]-1/x*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{2} + c_1 x$$

5 Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

5.1	problem Problem 24.17	497
5.2	problem Problem 24.18	501
5.3	problem Problem 24.19	505
5.4	problem Problem 24.26	510
5.5	problem Problem 24.27	515
5.6	problem Problem 24.28	521
5.7	problem Problem 24.29	527
5.8	problem Problem 24.30	533
5.9	problem Problem 24.31	539
5.10	problem Problem 24.32	545
5.11	problem Problem 24.33	551
5.12	problem Problem 24.35	558
5.13	problem Problem 24.36	567
5.14	problem Problem 24.37	575
5.15	problem Problem 24.44	579
5.16	problem Problem 24.46	584

5.1 problem Problem 24.17

5.1.1	Existence and uniqueness analysis	497
5.1.2	Solving as laplace ode	498
5.1.3	Maple step by step solution	499

Internal problem ID [5200]

Internal file name [OUTPUT/4693_Sunday_June_05_2022_03_03_30_PM_75048576/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.17.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' + 2y = 0$$

With initial conditions

$$[y(0) = 1]$$

5.1.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 2$$

$$q(x) = 0$$

Hence the ode is

$$y' + 2y = 0$$

The domain of $p(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. Hence solution exists and is unique.

5.1.2 Solving as laplace ode

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$sY(s) - y(0) + 2Y(s) = 0 \tag{1}$$

Replacing initial condition gives

$$sY(s) - 1 + 2Y(s) = 0$$

Solving for $Y(s)$ gives

$$Y(s) = \frac{1}{s+2}$$


Taking the inverse Laplace transform gives

$$\begin{aligned} y &= \mathcal{L}^{-1}(Y(s)) \\ &= \mathcal{L}^{-1}\left(\frac{1}{s+2}\right) \\ &= e^{-2x} \end{aligned}$$


Summary

The solution(s) found are the following

$$y = e^{-2x} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{-2x}$$

Verified OK.

5.1.3 Maple step by step solution

Let's solve

$$[y' + 2y = 0, y(0) = 1]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = -2$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int (-2) dx + c_1$$

- Evaluate integral

$$\ln(y) = -2x + c_1$$

- Solve for y

$$y = e^{-2x+c_1}$$

- Use initial condition $y(0) = 1$
 $1 = e^{c_1}$
- Solve for c_1
 $c_1 = 0$
- Substitute $c_1 = 0$ into general solution and simplify
 $y = e^{-2x}$
- Solution to the IVP
 $y = e^{-2x}$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.391 (sec). Leaf size: 8

```
dsolve([diff(y(x),x)+2*y(x)=0,y(0) = 1],y(x), singsol=all)
```

$$y(x) = e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 10

```
DSolve[{y'[x]+2*y[x]==0,{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}$$

5.2 problem Problem 24.18

5.2.1	Existence and uniqueness analysis	501
5.2.2	Solving as laplace ode	502
5.2.3	Maple step by step solution	503

Internal problem ID [5201]

Internal file name [OUTPUT/4694_Sunday_June_05_2022_03_03_31_PM_6745626/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.18.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' + 2y = 2$$

With initial conditions

$$[y(0) = 1]$$

5.2.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 2$$

$$q(x) = 2$$

Hence the ode is

$$y' + 2y = 2$$

The domain of $p(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

5.2.2 Solving as laplace ode

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$sY(s) - y(0) + 2Y(s) = \frac{2}{s} \quad (1)$$

Replacing initial condition gives

$$sY(s) - 1 + 2Y(s) = \frac{2}{s}$$

Solving for $Y(s)$ gives

$$Y(s) = \frac{1}{s}$$


Taking the inverse Laplace transform gives

$$\begin{aligned} y &= \mathcal{L}^{-1}(Y(s)) \\ &= \mathcal{L}^{-1}\left(\frac{1}{s}\right) \\ &= 1 \end{aligned}$$


Summary

The solution(s) found are the following

$$y = 1 \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 1$$

Verified OK.

5.2.3 Maple step by step solution

Let's solve

$$[y' + 2y = 2, y(0) = 1]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{-2y+2} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{-2y+2} dx = \int 1 dx + c_1$$

- Evaluate integral

$$-\frac{\ln(1-y)}{2} = x + c_1$$

- Solve for y

$$y = -e^{-2c_1-2x} + 1$$

- Use initial condition $y(0) = 1$
 $1 = -e^{-2c_1} + 1$
- Solution does not satisfy initial condition

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.312 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)+2*y(x)=2,y(0) = 1],y(x), singsol=all)
```

$$y(x) = 1$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{y'[x]+2*y[x]==2,{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1$$

5.3 problem Problem 24.19

5.3.1	Existence and uniqueness analysis	505
5.3.2	Solving as laplace ode	506
5.3.3	Maple step by step solution	507

Internal problem ID [5202]

Internal file name [OUTPUT/4695_Sunday_June_05_2022_03_03_32_PM_35441377/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.19.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' + 2y = e^x$$

With initial conditions

$$[y(0) = 1]$$

5.3.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 2$$

$$q(x) = e^x$$

Hence the ode is

$$y' + 2y = e^x$$

The domain of $p(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = e^x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

5.3.2 Solving as laplace ode

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$sY(s) - y(0) + 2Y(s) = \frac{1}{s-1} \quad (1)$$

Replacing initial condition gives

$$sY(s) - 1 + 2Y(s) = \frac{1}{s-1}$$

Solving for $Y(s)$ gives

$$Y(s) = \frac{s}{(s-1)(s+2)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{1}{3s-3} + \frac{2}{3(s+2)}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned} \mathcal{L}^{-1}\left(\frac{1}{3s-3}\right) &= \frac{e^x}{3} \\ \mathcal{L}^{-1}\left(\frac{2}{3(s+2)}\right) &= \frac{2e^{-2x}}{3} \end{aligned}$$


Adding the above results and simplifying gives

$$y = \frac{e^x}{3} + \frac{2e^{-2x}}{3}$$


Summary

The solution(s) found are the following

$$y = \frac{e^x}{3} + \frac{2e^{-2x}}{3} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^x}{3} + \frac{2e^{-2x}}{3}$$

Verified OK.

5.3.3 Maple step by step solution

Let's solve

$$[y' + 2y = e^x, y(0) = 1]$$

- Highest derivative means the order of the ODE is 1

y'

- Isolate the derivative

$$y' = -2y + e^x$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + 2y = e^x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + 2y) = \mu(x) e^x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' + 2y) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = 2\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{2x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) e^x dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) e^x dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) e^x dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{2x}$

$$y = \frac{\int e^{2x} e^x dx + c_1}{e^{2x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{e^{3x}}{3} + c_1}{e^{2x}}$$

- Simplify

$$y = \frac{(e^{3x} + 3c_1)e^{-2x}}{3}$$

- Use initial condition $y(0) = 1$

$$1 = \frac{1}{3} + c_1$$

- Solve for c_1

$$c_1 = \frac{2}{3}$$

- Substitute $c_1 = \frac{2}{3}$ into general solution and simplify

$$y = \frac{(e^{3x} + 2)e^{-2x}}{3}$$

- Solution to the IVP

$$y = \frac{(e^{3x}+2)e^{-2x}}{3}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.343 (sec). Leaf size: 15

```
dsolve([diff(y(x),x)+2*y(x)=exp(x),y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(e^{3x} + 2)e^{-2x}}{3}$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 21

```
DSolve[{y'[x]+2*y[x]==Exp[x],{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^{-2x}(e^{3x} + 2)$$

5.4 problem Problem 24.26

5.4.1	Existence and uniqueness analysis	510
5.4.2	Maple step by step solution	512

Internal problem ID [5203]

Internal file name [OUTPUT/4696_Sunday_June_05_2022_03_03_33_PM_29252225/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.26.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 1]$$

5.4.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = -1$$

$$F = 0$$

Hence the ode is

$$y'' - y = 0$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) - Y(s) = 0 \tag{1}$$

But the initial conditions are

$$y(0) = 1$$

$$y'(0) = 1$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 1 - s - Y(s) = 0$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{1}{s-1}$$

Taking the inverse Laplace transform gives

$$\begin{aligned} y &= \mathcal{L}^{-1}(Y(s)) \\ &= \mathcal{L}^{-1}\left(\frac{1}{s-1}\right) \\ &= e^x \end{aligned}$$


Simplifying the solution gives

$$y = e^x$$


Summary

The solution(s) found are the following

$$y = e^x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^x$$

Verified OK.

5.4.2 Maple step by step solution

Let's solve

$$\left[y'' - y = 0, y(0) = 1, y' \Big|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial
 $(r - 1)(r + 1) = 0$
- Roots of the characteristic polynomial
 $r = (-1, 1)$
- 1st solution of the ODE
 $y_1(x) = e^{-x}$
- 2nd solution of the ODE
 $y_2(x) = e^x$
- General solution of the ODE
 $y = c_1y_1(x) + c_2y_2(x)$
- Substitute in solutions
 $y = c_1e^{-x} + c_2e^x$
- Check validity of solution $y = c_1e^{-x} + c_2e^x$
 - Use initial condition $y(0) = 1$
 $1 = c_1 + c_2$
 - Compute derivative of the solution
 $y' = -c_1e^{-x} + c_2e^x$
 - Use the initial condition $y' \Big|_{\{x=0\}} = 1$
 $1 = -c_1 + c_2$
 - Solve for c_1 and c_2
 $\{c_1 = 0, c_2 = 1\}$
 - Substitute constant values into general solution and simplify
 $y = e^x$
- Solution to the IVP
 $y = e^x$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.359 (sec). Leaf size: 6

```
dsolve([diff(y(x),x$2)-y(x)=0,y(0) = 1, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = e^x$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 26

```
DSolve[{y''[x]-y[x]==Sin[x],{y[0]==1,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(-e^{-x} + 5e^x - 2\sin(x))$$

5.5 problem Problem 24.27

5.5.1	Existence and uniqueness analysis	515
5.5.2	Maple step by step solution	518

Internal problem ID [5204]

Internal file name [OUTPUT/4697_Sunday_June_05_2022_03_03_34_PM_23039275/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.27.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - y = \sin(x)$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

5.5.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = -1$$

$$F = \sin(x)$$

Hence the ode is

$$y'' - y = \sin(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) - Y(s) = \frac{1}{s^2 + 1} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 0 \\ y'(0) &= 1\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 1 - Y(s) = \frac{1}{s^2 + 1}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{s^2 + 2}{(s^2 + 1)(s^2 - 1)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{i}{4s - 4i} - \frac{i}{4(s + i)} + \frac{3}{4(s - 1)} - \frac{3}{4(s + 1)}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(\frac{i}{4s - 4i}\right) &= \frac{ie^{ix}}{4} \\ \mathcal{L}^{-1}\left(-\frac{i}{4(s + i)}\right) &= -\frac{ie^{-ix}}{4} \\ \mathcal{L}^{-1}\left(\frac{3}{4(s - 1)}\right) &= \frac{3e^x}{4} \\ \mathcal{L}^{-1}\left(-\frac{3}{4(s + 1)}\right) &= -\frac{3e^{-x}}{4}\end{aligned}$$

Adding the above results and simplifying gives

$$y = -\frac{\sin(x)}{2} + \frac{3\sinh(x)}{2}$$


Simplifying the solution gives

$$y = -\frac{\sin(x)}{2} + \frac{3\sinh(x)}{2}$$


Summary

The solution(s) found are the following

$$y = -\frac{\sin(x)}{2} + \frac{3\sinh(x)}{2} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -\frac{\sin(x)}{2} + \frac{3 \sinh(x)}{2}$$

Verified OK.

5.5.2 Maple step by step solution

Let's solve

$$\left[y'' - y = \sin(x), y(0) = 0, y'|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE
- $r^2 - 1 = 0$
- Factor the characteristic polynomial
- $(r - 1)(r + 1) = 0$
- Roots of the characteristic polynomial
- $r = (-1, 1)$
- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^x \\ -e^{-x} & e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x} \left(\int \sin(x)e^x dx \right)}{2} + \frac{e^x \left(\int e^{-x} \sin(x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = -\frac{\sin(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^x - \frac{\sin(x)}{2}$$

- Check validity of solution $y = c_1 e^{-x} + c_2 e^x - \frac{\sin(x)}{2}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = -c_1 e^{-x} + c_2 e^x - \frac{\cos(x)}{2}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 1$

$$1 = -c_1 + c_2 - \frac{1}{2}$$

- Solve for c_1 and c_2

$$\left\{c_1 = -\frac{3}{4}, c_2 = \frac{3}{4}\right\}$$

- Substitute constant values into general solution and simplify

$$y = -\frac{3e^{-x}}{4} + \frac{3e^x}{4} - \frac{\sin(x)}{2}$$

- Solution to the IVP

$$y = -\frac{3e^{-x}}{4} + \frac{3e^x}{4} - \frac{\sin(x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.406 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)-y(x)=sin(x),y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{\sin(x)}{2} + \frac{3 \sinh(x)}{2}$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 26

```
DSolve[{y'[x]-y[x]==Sin[x],{y[0]==1,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(-e^{-x} + 5e^x - 2 \sin(x))$$

5.6 problem Problem 24.28

5.6.1	Existence and uniqueness analysis	521
5.6.2	Maple step by step solution	524

Internal problem ID [5205]

Internal file name [OUTPUT/4698_Sunday_June_05_2022_03_03_35_PM_85166985/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.28.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y = e^x$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

5.6.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$\begin{aligned} p(x) &= 0 \\ q(x) &= -1 \\ F &= e^x \end{aligned}$$

Hence the ode is

$$y'' - y = e^x$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = e^x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) - Y(s) = \frac{1}{s-1} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 1 \\ y'(0) &= 0\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - s - Y(s) = \frac{1}{s-1}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{s^2 - s + 1}{(s-1)(s^2-1)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{3}{4(s+1)} + \frac{1}{2(s-1)^2} + \frac{1}{4s-4}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(\frac{3}{4(s+1)}\right) &= \frac{3e^{-x}}{4} \\ \mathcal{L}^{-1}\left(\frac{1}{2(s-1)^2}\right) &= \frac{xe^x}{2} \\ \mathcal{L}^{-1}\left(\frac{1}{4s-4}\right) &= \frac{e^x}{4}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{3e^{-x}}{4} + \frac{e^x(1+2x)}{4}$$


Simplifying the solution gives

$$y = \frac{3e^{-x}}{4} + \frac{e^x(1+2x)}{4}$$


Summary

The solution(s) found are the following

$$y = \frac{3e^{-x}}{4} + \frac{e^x(1+2x)}{4} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{3e^{-x}}{4} + \frac{e^x(1+2x)}{4}$$

Verified OK.

5.6.2 Maple step by step solution

Let's solve

$$\left[y'' - y = e^x, y(0) = 1, y'|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 1)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^x \\ -e^{-x} & e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x}(\int e^{2x} dx)}{2} + \frac{e^x(\int 1 dx)}{2}$$

- Compute integrals

$$y_p(x) = \frac{(2x-1)e^x}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^x + \frac{(2x-1)e^x}{4}$$

- Check validity of solution $y = c_1 e^{-x} + c_2 e^x + \frac{(2x-1)e^x}{4}$

- Use initial condition $y(0) = 1$

$$1 = c_1 + c_2 - \frac{1}{4}$$

- Compute derivative of the solution

$$y' = -c_1 e^{-x} + c_2 e^x + \frac{e^x}{2} + \frac{(2x-1)e^x}{4}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 0$

$$0 = -c_1 + c_2 + \frac{1}{4}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{3}{4}, c_2 = \frac{1}{2} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{3e^{-x}}{4} + \frac{e^x(1+2x)}{4}$$

- Solution to the IVP

$$y = \frac{3e^{-x}}{4} + \frac{e^x(1+2x)}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.391 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$2)-y(x)=exp(x),y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{3e^{-x}}{4} + \frac{e^x(2x+1)}{4}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 27

```
DSolve[{y'[x]-y[x]==Exp[x],{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}e^{-x}(e^{2x}(2x+1) + 3)$$

5.7 problem Problem 24.29

5.7.1	Existence and uniqueness analysis	527
5.7.2	Maple step by step solution	530

Internal problem ID [5206]

Internal file name [OUTPUT/4699_Sunday_June_05_2022_03_03_36_PM_97048772/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.29.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 2y' - 3y = \sin(2x)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

5.7.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 2$$

$$q(x) = -3$$

$$F = \sin(2x)$$

Hence the ode is

$$y'' + 2y' - 3y = \sin(2x)$$

The domain of $p(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = \sin(2x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 2sY(s) - 2y(0) - 3Y(s) = \frac{2}{s^2 + 4} \quad (1)$$

But the initial conditions are

$$y(0) = 0$$

$$y'(0) = 0$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) + 2sY(s) - 3Y(s) = \frac{2}{s^2 + 4}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{2}{(s^2 + 4)(s^2 + 2s - 3)}$$

Applying partial fractions decomposition results in

$$Y(s) = -\frac{1}{26(s+3)} + \frac{-\frac{2}{65} + \frac{7i}{130}}{s-2i} + \frac{-\frac{2}{65} - \frac{7i}{130}}{s+2i} + \frac{1}{10s-10}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(-\frac{1}{26(s+3)}\right) &= -\frac{e^{-3x}}{26} \\ \mathcal{L}^{-1}\left(\frac{-\frac{2}{65} + \frac{7i}{130}}{s-2i}\right) &= \left(-\frac{2}{65} + \frac{7i}{130}\right)e^{2ix} \\ \mathcal{L}^{-1}\left(\frac{-\frac{2}{65} - \frac{7i}{130}}{s+2i}\right) &= \left(-\frac{2}{65} - \frac{7i}{130}\right)e^{-2ix} \\ \mathcal{L}^{-1}\left(\frac{1}{10s-10}\right) &= \frac{e^x}{10}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{e^x}{10} - \frac{e^{-3x}}{26} - \frac{4 \cos(2x)}{65} - \frac{7 \sin(2x)}{65}$$


Simplifying the solution gives

$$y = -\frac{4 e^{-3x} \left(\left(\cos(2x) + \frac{7 \sin(2x)}{4} \right) e^{3x} - \frac{13 e^{4x}}{8} + \frac{5}{8} \right)}{65}$$


Summary

The solution(s) found are the following

$$y = -\frac{4 e^{-3x} \left(\left(\cos(2x) + \frac{7 \sin(2x)}{4} \right) e^{3x} - \frac{13 e^{4x}}{8} + \frac{5}{8} \right)}{65} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -\frac{4 e^{-3x} \left(\cos(2x) + \frac{7 \sin(2x)}{4} \right) e^{3x} - \frac{13 e^{4x}}{8} + \frac{5}{8}}{65}$$

Verified OK.

5.7.2 Maple step by step solution

Let's solve

$$\left[y'' + 2y' - 3y = \sin(2x), y(0) = 0, y' \Big|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE
- $r^2 + 2r - 3 = 0$
- Factor the characteristic polynomial
- $(r + 3)(r - 1) = 0$
- Roots of the characteristic polynomial
- $r = (-3, 1)$
- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-3x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-3x} + c_2 e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-3x} & e^x \\ -3e^{-3x} & e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 4e^{-2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{(e^{4x} (\int e^{-x} \sin(2x) dx) - (\int \sin(2x) e^{3x} dx)) e^{-3x}}{4}$$

- Compute integrals

$$y_p(x) = -\frac{4 \cos(2x)}{65} - \frac{7 \sin(2x)}{65}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-3x} + c_2 e^x - \frac{4 \cos(2x)}{65} - \frac{7 \sin(2x)}{65}$$

- Check validity of solution $y = c_1 e^{-3x} + c_2 e^x - \frac{4 \cos(2x)}{65} - \frac{7 \sin(2x)}{65}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2 - \frac{4}{65}$$

- Compute derivative of the solution

$$y' = -3c_1 e^{-3x} + c_2 e^x + \frac{8 \sin(2x)}{65} - \frac{14 \cos(2x)}{65}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 0$

$$0 = -3c_1 + c_2 - \frac{14}{65}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{1}{26}, c_2 = \frac{1}{10} \right\}$$

- Substitute constant values into general solution and simplify

$$y = -\frac{4e^{-3x} \left(\cos(2x) + \frac{7\sin(2x)}{4} \right) e^{3x} - \frac{13e^{4x}}{8} + \frac{5}{8}}{65}$$

- Solution to the IVP

$$y = -\frac{4e^{-3x} \left(\cos(2x) + \frac{7\sin(2x)}{4} \right) e^{3x} - \frac{13e^{4x}}{8} + \frac{5}{8}}{65}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.454 (sec). Leaf size: 27

```
dsolve([diff(y(x),x$2)+2*diff(y(x),x)-3*y(x)=sin(2*x),y(0) = 0, D(y)(0) = 0],y(x), singsol=a
```

$$y(x) = -\frac{4e^{-3x} \left(\cos(2x) + \frac{7\sin(2x)}{4} \right) e^{3x} - \frac{13e^{4x}}{8} + \frac{5}{8}}{65}$$

✓ Solution by Mathematica

Time used: 0.109 (sec). Leaf size: 36

```
DSolve[{y''[x]-2*y'[x]-3*y[x]==Sin[2*x],{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{130} (-13e^{-x} + 5e^{3x} - 14\sin(2x) + 8\cos(2x))$$

5.8 problem Problem 24.30

5.8.1	Existence and uniqueness analysis	533
5.8.2	Maple step by step solution	536

Internal problem ID [5207]

Internal file name [OUTPUT/4700_Sunday_June_05_2022_03_03_37_PM_82796084/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.30.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(x)$$

With initial conditions

$$[y(0) = 0, y'(0) = 2]$$

5.8.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 1$$

$$F = \sin(x)$$

Hence the ode is

$$y'' + y = \sin(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + Y(s) = \frac{1}{s^2 + 1} \quad (1)$$

But the initial conditions are

$$y(0) = 0$$

$$y'(0) = 2$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 2 + Y(s) = \frac{1}{s^2 + 1}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{2s^2 + 3}{(s^2 + 1)^2}$$

Applying partial fractions decomposition results in

$$Y(s) = -\frac{1}{4(s-i)^2} - \frac{1}{4(s+i)^2} - \frac{5i}{4(s-i)} + \frac{5i}{4(s+i)}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(-\frac{1}{4(s-i)^2}\right) &= -\frac{x e^{ix}}{4} \\ \mathcal{L}^{-1}\left(-\frac{1}{4(s+i)^2}\right) &= -\frac{x e^{-ix}}{4} \\ \mathcal{L}^{-1}\left(-\frac{5i}{4(s-i)}\right) &= -\frac{5ie^{ix}}{4} \\ \mathcal{L}^{-1}\left(\frac{5i}{4(s+i)}\right) &= \frac{5ie^{-ix}}{4}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{5 \sin(x)}{2} - \frac{\cos(x) x}{2}$$


Simplifying the solution gives

$$y = \frac{5 \sin(x)}{2} - \frac{\cos(x) x}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{5 \sin(x)}{2} - \frac{\cos(x) x}{2} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{5 \sin(x)}{2} - \frac{\cos(x)x}{2}$$

Verified OK.

5.8.2 Maple step by step solution

Let's solve

$$\left[y'' + y = \sin(x), y(0) = 0, y'|_{\{x=0\}} = 2 \right]$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE
- $r^2 + 1 = 0$
- Use quadratic formula to solve for r
- $r = \frac{0 \pm (\sqrt{-4})}{2}$
- Roots of the characteristic polynomial
- $r = (-I, I)$
- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{\cos(x)x}{2}$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{\cos(x)x}{2}$$

- Check validity of solution $y = \cos(x) c_1 + c_2 \sin(x) + \frac{\sin(x)}{4} - \frac{\cos(x)x}{2}$

- Use initial condition $y(0) = 0$

$$0 = c_1$$

- Compute derivative of the solution

$$y' = -\sin(x) c_1 + c_2 \cos(x) - \frac{\cos(x)}{4} + \frac{\sin(x)x}{2}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 2$

$$2 = -\frac{1}{4} + c_2$$

- Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = \frac{9}{4}\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{5 \sin(x)}{2} - \frac{\cos(x)x}{2}$$

- Solution to the IVP

$$y = \frac{5 \sin(x)}{2} - \frac{\cos(x)x}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.375 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)+y(x)=sin(x),y(0) = 0, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = \frac{5 \sin(x)}{2} - \frac{\cos(x)x}{2}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 19

```
DSolve[{y'[x]+y[x]==Sin[x],{y[0]==0,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(5 \sin(x) - x \cos(x))$$

5.9 problem Problem 24.31

5.9.1	Existence and uniqueness analysis	539
5.9.2	Maple step by step solution	542

Internal problem ID [5208]

Internal file name [OUTPUT/4701_Sunday_June_05_2022_03_03_38_PM_30494857/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.31.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y' + y = 0$$

With initial conditions

$$[y(0) = 4, y'(0) = -3]$$

5.9.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = 1$$

$$F = 0$$

Hence the ode is

$$y'' + y' + y = 0$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + sY(s) - y(0) + Y(s) = 0 \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 4 \\ y'(0) &= -3\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 1 - 4s + sY(s) + Y(s) = 0$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{4s + 1}{s^2 + s + 1}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{2 + \frac{i\sqrt{3}}{3}}{s + \frac{1}{2} - \frac{i\sqrt{3}}{2}} + \frac{2 - \frac{i\sqrt{3}}{3}}{s + \frac{1}{2} + \frac{i\sqrt{3}}{2}}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{2 + \frac{i\sqrt{3}}{3}}{s + \frac{1}{2} - \frac{i\sqrt{3}}{2}}\right) = \frac{(i\sqrt{3} + 6) e^{-\frac{(1-i\sqrt{3})x}{2}}}{3}$$

$$\mathcal{L}^{-1}\left(\frac{2 - \frac{i\sqrt{3}}{3}}{s + \frac{1}{2} + \frac{i\sqrt{3}}{2}}\right) = \frac{(6 - i\sqrt{3}) e^{-\frac{(1+i\sqrt{3})x}{2}}}{3}$$

Adding the above results and simplifying gives

$$y = \frac{2e^{-\frac{x}{2}}\left(6\cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}\right)}{3}$$


Simplifying the solution gives

$$y = -\frac{2\left(\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} - 6\cos\left(\frac{\sqrt{3}x}{2}\right)\right)e^{-\frac{x}{2}}}{3}$$


Summary

The solution(s) found are the following

$$y = -\frac{2\left(\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} - 6\cos\left(\frac{\sqrt{3}x}{2}\right)\right)e^{-\frac{x}{2}}}{3} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -\frac{2\left(\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} - 6\cos\left(\frac{\sqrt{3}x}{2}\right)\right)e^{-\frac{x}{2}}}{3}$$

Verified OK.

5.9.2 Maple step by step solution

Let's solve

$$\left[y'' + y' + y = 0, y(0) = 4, y'|_{\{x=0\}} = -3 \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the ODE

$$y_2(x) = \sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2$$

- Check validity of solution $y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2$

- Use initial condition $y(0) = 4$

$$4 = c_1$$

- Compute derivative of the solution

$$y' = -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)c_1}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}c_1}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)c_2}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}c_2}{2}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = -3$

$$-3 = -\frac{c_1}{2} + \frac{\sqrt{3}c_2}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = 4, c_2 = -\frac{2\sqrt{3}}{3} \right\}$$

- Substitute constant values into general solution and simplify

$$y = -\frac{2\left(\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} - 6\cos\left(\frac{\sqrt{3}x}{2}\right)\right)e^{-\frac{x}{2}}}{3}$$

- Solution to the IVP

$$y = -\frac{2\left(\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} - 6\cos\left(\frac{\sqrt{3}x}{2}\right)\right)e^{-\frac{x}{2}}}{3}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.453 (sec). Leaf size: 32

```
dsolve([diff(y(x),x$2)+diff(y(x),x)+y(x)=0,y(0) = 4, D(y)(0) = -3],y(x), singsol=all)
```

$$y(x) = -\frac{2\left(\sqrt{3}\sin\left(\frac{\sqrt{3}x}{2}\right) - 6\cos\left(\frac{\sqrt{3}x}{2}\right)\right)e^{-\frac{x}{2}}}{3}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 47

```
DSolve[{y''[x]+y'[x]+y[x]==0,{y[0]==4,y'[0]==-3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2}{3}e^{-x/2} \left(\sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right) - 6 \cos \left(\frac{\sqrt{3}x}{2} \right) \right)$$

5.10 problem Problem 24.32

5.10.1 Existence and uniqueness analysis	545
5.10.2 Maple step by step solution	548

Internal problem ID [5209]

Internal file name [OUTPUT/4702_Sunday_June_05_2022_03_03_39_PM_85824528/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.32.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 2y' + 5y = 3e^{-2x}$$

With initial conditions

$$[y(0) = 1, y'(0) = 1]$$

5.10.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 2$$

$$q(x) = 5$$

$$F = 3e^{-2x}$$

Hence the ode is

$$y'' + 2y' + 5y = 3e^{-2x}$$

The domain of $p(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 5$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = 3e^{-2x}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 2sY(s) - 2y(0) + 5Y(s) = \frac{3}{s+2} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 1 \\ y'(0) &= 1\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 3 - s + 2sY(s) + 5Y(s) = \frac{3}{s+2}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{s^2 + 5s + 9}{(s+2)(s^2 + 2s + 5)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{\frac{1}{5} - \frac{13i}{20}}{s + 1 - 2i} + \frac{\frac{1}{5} + \frac{13i}{20}}{s + 1 + 2i} + \frac{3}{5(s + 2)}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{\frac{1}{5} - \frac{13i}{20}}{s + 1 - 2i}\right) = \left(\frac{1}{5} - \frac{13i}{20}\right) e^{(-1+2i)x}$$

$$\mathcal{L}^{-1}\left(\frac{\frac{1}{5} + \frac{13i}{20}}{s + 1 + 2i}\right) = \left(\frac{1}{5} + \frac{13i}{20}\right) e^{(-1-2i)x}$$

$$\mathcal{L}^{-1}\left(\frac{3}{5(s + 2)}\right) = \frac{3e^{-2x}}{5}$$

Adding the above results and simplifying gives

$$y = \frac{3e^{-2x}}{5} + \frac{e^{-x}(4\cos(2x) + 13\sin(2x))}{10}$$


Simplifying the solution gives

$$y = \frac{3e^{-2x}}{5} + \frac{e^{-x}(4\cos(2x) + 13\sin(2x))}{10}$$


Summary

The solution(s) found are the following

$$y = \frac{3e^{-2x}}{5} + \frac{e^{-x}(4\cos(2x) + 13\sin(2x))}{10} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{3 e^{-2x}}{5} + \frac{e^{-x}(4 \cos(2x) + 13 \sin(2x))}{10}$$

Verified OK.

5.10.2 Maple step by step solution

Let's solve

$$\left[y'' + 2y' + 5y = 3 e^{-2x}, y(0) = 1, y'|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 2r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2) \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - 2I, -1 + 2I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x} \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-x} \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} \cos(2x) + c_2 e^{-x} \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 3 e^{-2x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} \cos(2x) & e^{-x} \sin(2x) \\ -e^{-x} \cos(2x) - 2e^{-x} \sin(2x) & -e^{-x} \sin(2x) + 2e^{-x} \cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2e^{-2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{3e^{-x}(\cos(2x)(\int e^{-x} \sin(2x) dx) - \sin(2x)(\int e^{-x} \cos(2x) dx))}{2}$$

- Compute integrals

$$y_p(x) = \frac{3e^{-2x}}{5}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} \cos(2x) + c_2 e^{-x} \sin(2x) + \frac{3e^{-2x}}{5}$$

- Check validity of solution $y = c_1 e^{-x} \cos(2x) + c_2 e^{-x} \sin(2x) + \frac{3e^{-2x}}{5}$

- Use initial condition $y(0) = 1$

$$1 = c_1 + \frac{3}{5}$$

- Compute derivative of the solution

$$y' = -c_1 e^{-x} \cos(2x) - 2c_1 e^{-x} \sin(2x) - c_2 e^{-x} \sin(2x) + 2c_2 e^{-x} \cos(2x) - \frac{6e^{-2x}}{5}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 1$

$$1 = -c_1 - \frac{6}{5} + 2c_2$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{2}{5}, c_2 = \frac{13}{10} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{3e^{-2x}}{5} + \frac{e^{-x}(4 \cos(2x) + 13 \sin(2x))}{10}$$

- Solution to the IVP

$$y = \frac{3e^{-2x}}{5} + \frac{e^{-x}(4 \cos(2x) + 13 \sin(2x))}{10}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.469 (sec). Leaf size: 30

```
dsolve([diff(y(x),x$2)+2*diff(y(x),x)+5*y(x)=3*exp(-2*x),y(0) = 1, D(y)(0) = 1],y(x), singsol
```

$$y(x) = \frac{3e^{-2x}}{5} + \frac{e^{-x}(4\cos(2x) + 13\sin(2x))}{10}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 34

```
DSolve[{y''[x]+2*y'[x]+5*y[x]==3*Exp[-2*x]},{y[0]==1,y'[0]==1},y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow \frac{1}{10}e^{-2x}(13e^x \sin(2x) + 4e^x \cos(2x) + 6)$$

5.11 problem Problem 24.33

5.11.1 Existence and uniqueness analysis	551
5.11.2 Maple step by step solution	554

Internal problem ID [5210]

Internal file name [OUTPUT/4703_Sunday_June_05_2022_03_03_40_PM_78854413/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.33.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 5y' - 3y = \text{Heaviside}(-4 + x)$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

5.11.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 5$$

$$q(x) = -3$$

$$F = \text{Heaviside}(-4 + x)$$

Hence the ode is

$$y'' + 5y' - 3y = \text{Heaviside}(-4 + x)$$

The domain of $p(x) = 5$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = \text{Heaviside}(-4 + x)$ is

$$\{x < 4 \vee 4 < x\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 5sY(s) - 5y(0) - 3Y(s) = \frac{e^{-4s}}{s} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 0 \\ y'(0) &= 0\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) + 5sY(s) - 3Y(s) = \frac{e^{-4s}}{s}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{e^{-4s}}{s(s^2 + 5s - 3)}$$

Taking the inverse Laplace transform gives

$$\begin{aligned}
 y &= \mathcal{L}^{-1}(Y(s)) \\
 &= \mathcal{L}^{-1}\left(\frac{e^{-4s}}{s(s^2 + 5s - 3)}\right) \\
 &= \frac{\text{Heaviside}(-4 + x) \left(-37 + e^{10 - \frac{5x}{2}} \left(5 \sinh\left(\frac{\sqrt{37}(-4+x)}{2}\right) \sqrt{37} + 37 \cosh\left(\frac{\sqrt{37}(-4+x)}{2}\right)\right)\right)}{111}
 \end{aligned}$$

Hence the final solution is

$$y = \frac{\text{Heaviside}(-4 + x) \left(-37 + e^{10 - \frac{5x}{2}} \left(5 \sinh\left(\frac{\sqrt{37}(-4+x)}{2}\right) \sqrt{37} + 37 \cosh\left(\frac{\sqrt{37}(-4+x)}{2}\right)\right)\right)}{111}$$


Simplifying the solution gives

$$y = \frac{\text{Heaviside}(-4 + x) \left(-1 + \frac{5 e^{10 - \frac{5x}{2}} \sinh\left(\frac{\sqrt{37}(-4+x)}{2}\right) \sqrt{37}}{37} + e^{10 - \frac{5x}{2}} \cosh\left(\frac{\sqrt{37}(-4+x)}{2}\right)\right)}{3}$$


Summary

The solution(s) found are the following

$$y = \frac{\text{Heaviside}(-4 + x) \left(-1 + \frac{5 e^{10 - \frac{5x}{2}} \sinh\left(\frac{\sqrt{37}(-4+x)}{2}\right) \sqrt{37}}{37} + e^{10 - \frac{5x}{2}} \cosh\left(\frac{\sqrt{37}(-4+x)}{2}\right)\right)}{3} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\text{Heaviside}(-4+x) \left(-1 + \frac{5e^{10-\frac{5x}{2}} \sinh\left(\frac{\sqrt{37}(-4+x)}{2}\right) \sqrt{37}}{37} + e^{10-\frac{5x}{2}} \cosh\left(\frac{\sqrt{37}(-4+x)}{2}\right) \right)}{3}$$

Verified OK.

5.11.2 Maple step by step solution

Let's solve

$$\left[y'' + 5y' - 3y = \text{Heaviside}(-4+x), y(0) = 0, y' \Big|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of homogeneous ODE

$$r^2 + 5r - 3 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-5) \pm (\sqrt{37})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{5}{2} - \frac{\sqrt{37}}{2}, -\frac{5}{2} + \frac{\sqrt{37}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{\left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right)x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{\left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right)x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{\left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right)x} + c_2 e^{\left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right)x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \text{Heaviside}(-4+x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{\left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right)x} & e^{\left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right)x} \\ \left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right) e^{\left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right)x} & \left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right) e^{\left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right)x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \sqrt{37} e^{-5x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\sqrt{37} \left(e^{-\frac{(5+\sqrt{37})x}{2}} \left(\int e^{\frac{(5+\sqrt{37})x}{2}} Heaviside(-4+x) dx \right) - e^{-\frac{(-5+\sqrt{37})x}{2}} \left(\int e^{-\frac{(-5+\sqrt{37})x}{2}} Heaviside(-4+x) dx \right) \right)}{37}$$

- Compute integrals

$$y_p(x) = \frac{Heaviside(-4+x) \left(5\sqrt{37} e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} - 5\sqrt{37} e^{-\frac{(-4+x)(5+\sqrt{37})}{2}} + 37 e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} + 37 e^{-\frac{(-4+x)(5+\sqrt{37})}{2}} \right)}{222}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{\left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right)x} + c_2 e^{\left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right)x} + \frac{Heaviside(-4+x) \left(5\sqrt{37} e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} - 5\sqrt{37} e^{-\frac{(-4+x)(5+\sqrt{37})}{2}} + 37 e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} + 37 e^{-\frac{(-4+x)(5+\sqrt{37})}{2}} \right)}{222}$$

- Check validity of solution $y = c_1 e^{\left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right)x} + c_2 e^{\left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right)x} + \frac{Heaviside(-4+x) \left(5\sqrt{37} e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} - 5\sqrt{37} e^{-\frac{(-4+x)(5+\sqrt{37})}{2}} + 37 e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} + 37 e^{-\frac{(-4+x)(5+\sqrt{37})}{2}} \right)}{222}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = c_1 \left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right) e^{\left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right)x} + c_2 \left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right) e^{\left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right)x} + \frac{Dirac(-4+x) \left(5\sqrt{37} e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} - 5\sqrt{37} e^{-\frac{(-4+x)(5+\sqrt{37})}{2}} + 37 e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} + 37 e^{-\frac{(-4+x)(5+\sqrt{37})}{2}} \right)}{222}$$

- Use the initial condition $y'|_{\{x=0\}} = 0$

$$0 = c_1 \left(-\frac{5}{2} - \frac{\sqrt{37}}{2}\right) + c_2 \left(-\frac{5}{2} + \frac{\sqrt{37}}{2}\right)$$

- Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = 0\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{\text{Heaviside}(-4+x) \left(5\sqrt{37} e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} - 5\sqrt{37} e^{\frac{(-4+x)(5+\sqrt{37})}{2}} + 37 e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} + 37 e^{\frac{(-4+x)(5+\sqrt{37})}{2}} - 74 \right)}{222}$$

- Solution to the IVP

$$y = \frac{\text{Heaviside}(-4+x) \left(5\sqrt{37} e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} - 5\sqrt{37} e^{\frac{(-4+x)(5+\sqrt{37})}{2}} + 37 e^{\frac{(-4+x)(-5+\sqrt{37})}{2}} + 37 e^{\frac{(-4+x)(5+\sqrt{37})}{2}} - 74 \right)}{222}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.531 (sec). Leaf size: 45

```
dsolve([diff(y(x),x$2)+5*diff(y(x),x)-3*y(x)=Heaviside(x-4),y(0) = 0, D(y)(0) = 0],y(x), sin
```

$$y(x) = \frac{\text{Heaviside}(x-4) \left(-1 + \frac{5\sqrt{37} \sinh\left(\frac{(x-4)\sqrt{37}}{2}\right) e^{-\frac{5x}{2}+10}}{37} + \cosh\left(\frac{(x-4)\sqrt{37}}{2}\right) e^{-\frac{5x}{2}+10} \right)}{3}$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 70

```
DSolve[{y''[x]+5*y'[x]-3*y[x]==UnitStep[x-4],{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolut
```

$y(x)$

$$\rightarrow \left\{ \begin{array}{ll} \frac{1}{222} \left(-74 + (37 + 5\sqrt{37}) e^{\frac{1}{2}(-5+\sqrt{37})(x-4)} + (37 - 5\sqrt{37}) e^{-\frac{1}{2}(5+\sqrt{37})(x-4)} \right) & x > 4 \\ 0 & \text{True} \end{array} \right.$$

5.12 problem Problem 24.35

5.12.1 Maple step by step solution 560

Internal problem ID [5211]

Internal file name [OUTPUT/4704_Sunday_June_05_2022_03_03_43_PM_92798513/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.35.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : **"higher_order_laplace"**

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - y = 5$$

With initial conditions

$$[y(0) = 0, y'(0) = 0, y''(0) = 0]$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

$$\mathcal{L}(y''') = s^3Y(s) - y''(0) - sy'(0) - s^2y(0)$$

The given ode becomes an algebraic equation in the Laplace domain

$$s^3Y(s) - y''(0) - sy'(0) - s^2y(0) - Y(s) = \frac{5}{s} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 0 \\y'(0) &= 0 \\y''(0) &= 0\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^3Y(s) - Y(s) = \frac{5}{s}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{5}{s(s^3 - 1)}$$

Applying partial fractions decomposition results in

$$Y(s) = -\frac{5}{s} + \frac{5}{3(s-1)} + \frac{5}{3\left(s + \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)} + \frac{5}{3\left(s + \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(-\frac{5}{s}\right) &= -5 \\ \mathcal{L}^{-1}\left(\frac{5}{3(s-1)}\right) &= \frac{5e^x}{3} \\ \mathcal{L}^{-1}\left(\frac{5}{3\left(s + \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)}\right) &= \frac{5e^{-\frac{(1-i\sqrt{3})x}{2}}}{3} \\ \mathcal{L}^{-1}\left(\frac{5}{3\left(s + \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}\right) &= \frac{5e^{-\frac{(1+i\sqrt{3})x}{2}}}{3}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{10e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - 5 + \frac{5e^x}{3}$$

Summary

The solution(s) found are the following

$$y = \frac{10e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - 5 + \frac{5e^x}{3} \quad (1)$$




Figure 94: Solution plot

Verification of solutions

$$y = \frac{10 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - 5 + \frac{5 e^x}{3}$$

Verified OK.

5.12.1 Maple step by step solution

Let's solve

$$\left[y''' - y = 5, y(0) = 0, y'|_{\{x=0\}} = 0, y''|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 3
 y'''
- Convert linear ODE into a system of first order ODEs
 - Define new variable $y_1(x)$
 $y_1(x) = y$
 - Define new variable $y_2(x)$
 $y_2(x) = y'$
 - Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 5 + y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 5 + y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ 5 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ 5 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\begin{array}{c} \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \\ 1, \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \end{array} \right], \left[\begin{array}{c} -\frac{1}{2} - \frac{i\sqrt{3}}{2}, \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right] \end{array} \right], \left[\begin{array}{c} -\frac{1}{2} + \frac{i\sqrt{3}}{2}, \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right] \end{array} \right]$$

- Consider eigenpair

$$\left[\begin{array}{c} \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \\ 1, \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^x \cdot \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[\begin{array}{c} -\frac{1}{2} - \frac{i\sqrt{3}}{2}, \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right] \end{array} \right]$$

- Solution from eigenpair

$$e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} \cdot \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right]$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-\frac{x}{2}} \cdot \left(\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \cdot \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right]$$

- Simplify expression

$$e^{-\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - I \sin\left(\frac{\sqrt{3}x}{2}\right)}{\left(-\frac{1}{2} - \frac{I\sqrt{3}}{2}\right)^2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - I \sin\left(\frac{\sqrt{3}x}{2}\right)}{-\frac{1}{2} - \frac{I\sqrt{3}}{2}} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) - I \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_2(x) = e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}, \vec{y}_3(x) = e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$
 $\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \vec{y}_p(x)$

□ Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & -\sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & -\sin\left(\frac{\sqrt{3}x}{2}\right) e^{-\frac{x}{2}} \end{bmatrix} \cdot \begin{bmatrix} 1 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 1 & -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ 1 & 1 & 0 \end{bmatrix}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \\ \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \\ \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} + \end{bmatrix}$$

□

Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}_p'(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} \frac{10e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - 5 + \frac{5e^x}{3} \\ -\frac{5e^{-\frac{x}{2}} \left(-e^{\frac{3x}{2}} + \sin\left(\frac{\sqrt{3}x}{2}\right) \sqrt{3} + \cos\left(\frac{\sqrt{3}x}{2}\right) \right)}{3} \\ \frac{5e^{-\frac{x}{2}} \left(e^{\frac{3x}{2}} + \sin\left(\frac{\sqrt{3}x}{2}\right) \sqrt{3} - \cos\left(\frac{\sqrt{3}x}{2}\right) \right)}{3} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \begin{bmatrix} \frac{10e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - 5 + \frac{5e^x}{3} \\ \frac{5e^{-\frac{x}{2}} \left(-e^{\frac{3x}{2}} + \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} + \cos\left(\frac{\sqrt{3}x}{2}\right)\right)}{3} \\ \frac{5e^{-\frac{x}{2}} \left(e^{\frac{3x}{2}} + \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} - \cos\left(\frac{\sqrt{3}x}{2}\right)\right)}{3} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -5 - \frac{e^{-\frac{x}{2}} \left(c_3\sqrt{3} + c_2 - \frac{20}{3}\right) \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \left(\sqrt{3}c_2 - c_3\right) \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{(5+3c_1)e^x}{3}$$

- Use the initial condition $y(0) = 0$

$$0 = -\frac{c_3\sqrt{3}}{2} - \frac{c_2}{2} + c_1$$

- Calculate the 1st derivative of the solution

$$y' = \frac{e^{-\frac{x}{2}} \left(c_3\sqrt{3} + c_2 - \frac{20}{3}\right) \cos\left(\frac{\sqrt{3}x}{2}\right)}{4} + \frac{e^{-\frac{x}{2}} \left(c_3\sqrt{3} + c_2 - \frac{20}{3}\right) \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{4} + \frac{e^{-\frac{x}{2}} \left(\sqrt{3}c_2 - c_3\right) \sin\left(\frac{\sqrt{3}x}{2}\right)}{4} - \frac{e^{-\frac{x}{2}} \left(\sqrt{3}c_2 - c_3\right)}{4}$$

- Use the initial condition $y'|_{\{x=0\}} = 0$

$$0 = \frac{c_3\sqrt{3}}{4} + \frac{c_2}{4} - \frac{(\sqrt{3}c_2 - c_3)\sqrt{3}}{4} + c_1$$

- Calculate the 2nd derivative of the solution

$$y'' = \frac{e^{-\frac{x}{2}} \left(c_3\sqrt{3} + c_2 - \frac{20}{3}\right) \cos\left(\frac{\sqrt{3}x}{2}\right)}{4} - \frac{e^{-\frac{x}{2}} \left(c_3\sqrt{3} + c_2 - \frac{20}{3}\right) \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{4} + \frac{e^{-\frac{x}{2}} \left(\sqrt{3}c_2 - c_3\right) \sin\left(\frac{\sqrt{3}x}{2}\right)}{4} + \frac{e^{-\frac{x}{2}} \left(\sqrt{3}c_2 - c_3\right)}{4}$$

- Use the initial condition $y''|_{\{x=0\}} = 0$

$$0 = \frac{c_3\sqrt{3}}{4} + \frac{c_2}{4} + \frac{(\sqrt{3}c_2 - c_3)\sqrt{3}}{4} + c_1$$

- Solve for the unknown coefficients

$$\{c_1 = 0, c_2 = 0, c_3 = 0\}$$

- Solution to the IVP

$$y = \frac{10e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - 5 + \frac{5e^x}{3}$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 3; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.422 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$3)-y(x)=5,y(0) = 0, D(y)(0) = 0, (D@@2)(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = -5 + \frac{5e^x}{3} + \frac{10e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 34

```
DSolve[{y'''[x]-y[x]==5,{y[0]==0,y'[0]==0,y''[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{5}{3} \left(e^x + 2e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) - 3 \right)$$

5.13 problem Problem 24.36

5.13.1 Maple step by step solution 569

Internal problem ID [5212]

Internal file name [OUTPUT/4705_Sunday_June_05_2022_03_03_44_PM_45275139/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.36.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_laplace**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' - y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0, y''(0) = 0, y'''(0) = 0]$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

$$\mathcal{L}(y''') = s^3Y(s) - y''(0) - sy'(0) - s^2y(0)$$

$$\mathcal{L}(y'''') = s^4Y(s) - y'''(0) - sy''(0) - s^2y'(0) - s^3y(0)$$

The given ode becomes an algebraic equation in the Laplace domain

$$s^4Y(s) - y'''(0) - sy''(0) - s^2y'(0) - s^3y(0) - Y(s) = 0 \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 1 \\y'(0) &= 0 \\y''(0) &= 0 \\y'''(0) &= 0\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^4 Y(s) - s^3 - Y(s) = 0$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{s^3}{s^4 - 1}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{1}{4s + 4} + \frac{1}{4s - 4i} + \frac{1}{4s + 4i} + \frac{1}{4s - 4}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(\frac{1}{4s + 4}\right) &= \frac{e^{-x}}{4} \\ \mathcal{L}^{-1}\left(\frac{1}{4s - 4i}\right) &= \frac{e^{ix}}{4} \\ \mathcal{L}^{-1}\left(\frac{1}{4s + 4i}\right) &= \frac{e^{-ix}}{4} \\ \mathcal{L}^{-1}\left(\frac{1}{4s - 4}\right) &= \frac{e^x}{4}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{\cos(x)}{2} + \frac{\cosh(x)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\cos(x)}{2} + \frac{\cosh(x)}{2} \tag{1}$$




Figure 95: Solution plot

Verification of solutions

$$y = \frac{\cos(x)}{2} + \frac{\cosh(x)}{2}$$

Verified OK.

5.13.1 Maple step by step solution

Let's solve

$$\left[y'''' - y = 0, y(0) = 1, y'|_{\{x=0\}} = 0, y''|_{\{x=0\}} = 0, y'''|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 4
 - y''''
- Convert linear ODE into a system of first order ODEs
 - Define new variable $y_1(x)$

$$y_1(x) = y$$
 - Define new variable $y_2(x)$

$$y_2(x) = y'$$
 - Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{array}{c} -1 \\ -1 \\ 1 \end{array} \right], \left[\begin{array}{c} -1 \\ 1 \\ -1 \\ 1 \end{array} \right] \right], \left[\left[\begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \end{array} \right], \left[\begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \end{array} \right] \right], \left[\left[\begin{array}{c} -1 \\ -1 \\ 1 \\ 1 \end{array} \right], \left[\begin{array}{c} -I \\ -1 \\ I \\ 1 \end{array} \right] \right], \left[\left[\begin{array}{c} I \\ -1 \\ -1 \\ 1 \end{array} \right], \left[\begin{array}{c} I \\ -1 \\ -I \\ 1 \end{array} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -I(\cos(x) - I \sin(x)) \\ -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -c_3 \sin(x) - c_4 \cos(x) \\ -c_3 \cos(x) + c_4 \sin(x) \\ c_3 \sin(x) + c_4 \cos(x) \\ c_3 \cos(x) - c_4 \sin(x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -c_1 e^{-x} + c_2 e^x - c_4 \cos(x) - c_3 \sin(x)$$

- Use the initial condition $y(0) = 1$
 $1 = -c_1 + c_2 - c_4$
- Calculate the 1st derivative of the solution
 $y' = c_1e^{-x} + c_2e^x + c_4 \sin(x) - c_3 \cos(x)$
- Use the initial condition $y' \Big|_{\{x=0\}} = 0$
 $0 = c_1 + c_2 - c_3$
- Calculate the 2nd derivative of the solution
 $y'' = -c_1e^{-x} + c_2e^x + c_4 \cos(x) + c_3 \sin(x)$
- Use the initial condition $y'' \Big|_{\{x=0\}} = 0$
 $0 = -c_1 + c_2 + c_4$
- Calculate the 3rd derivative of the solution
 $y''' = c_1e^{-x} + c_2e^x - c_4 \sin(x) + c_3 \cos(x)$
- Use the initial condition $y''' \Big|_{\{x=0\}} = 0$
 $0 = c_1 + c_2 + c_3$
- Solve for the unknown coefficients
 $\{c_1 = -\frac{1}{4}, c_2 = \frac{1}{4}, c_3 = 0, c_4 = -\frac{1}{2}\}$
- Solution to the IVP
 $y = \frac{\cos(x)}{2} + \frac{e^{-x}}{4} + \frac{e^x}{4}$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.422 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$4)-y(x)=0,y(0) = 1, D(y)(0) = 0, (D@@2)(y)(0) = 0, (D@@3)(y)(0) = 0],y(x))
```

$$y(x) = \frac{\cos(x)}{2} + \frac{\cosh(x)}{2}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 22

```
DSolve[{y''''[x]-y[x]==0,{y[0]==1,y'[0]==0,y''[0]==0,y'''[0]==0}},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{4}(e^{-x} + e^x + 2 \cos(x))$$

5.14 problem Problem 24.37

Internal problem ID [5213]

Internal file name [OUTPUT/4706_Sunday_June_05_2022_03_03_45_PM_1165393/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.37.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_laplace**"

Maple gives the following as the ode type

```
[[_3rd_order , _linear , _nonhomogeneous]]
```

$$y''' - 3y'' + 3y' - y = x^2 e^x$$

With initial conditions

$$[y(0) = 1, y'(0) = 2, y''(0) = 3]$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

$$\mathcal{L}(y''') = s^3Y(s) - y''(0) - sy'(0) - s^2y(0)$$

The given ode becomes an algebraic equation in the Laplace domain

$$s^3Y(s) - y''(0) - sy'(0) - s^2y(0) - 3s^2Y(s) + 3y'(0) + 3sy(0) + 3sY(s) - 3y(0) - Y(s) = \frac{2}{(s-1)^3} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 1 \\y'(0) &= 2 \\y''(0) &= 3\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^3Y(s) + s - s^2 - 3s^2Y(s) + 3sY(s) - Y(s) = \frac{2}{(s-1)^3}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{s^5 - 4s^4 + 6s^3 - 4s^2 + s + 2}{(s-1)^3(s^3 - 3s^2 + 3s - 1)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{2}{(s-1)^6} + \frac{1}{(s-1)^2} + \frac{1}{s-1}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(\frac{2}{(s-1)^6}\right) &= \frac{x^5 e^x}{60} \\ \mathcal{L}^{-1}\left(\frac{1}{(s-1)^2}\right) &= x e^x \\ \mathcal{L}^{-1}\left(\frac{1}{s-1}\right) &= e^x\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{e^x(x^5 + 60x + 60)}{60}$$

Summary

The solution(s) found are the following

$$y = \frac{e^x(x^5 + 60x + 60)}{60} \tag{1}$$




Figure 96: Solution plot

Verification of solutions

$$y = \frac{e^x(x^5 + 60x + 60)}{60}$$

Verified OK.

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 3; missing the dependent variable
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.453 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$3)-3*diff(y(x),x$2)+3*diff(y(x),x)-y(x)=x^2*exp(x),y(0) = 1, D(y)(0) = 2
```

$$y(x) = \frac{e^x(x^5 + 60x + 60)}{60}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 20

```
DSolve[{y'''[x]-3*y''[x]+3*y'[x]-y[x]==x^2*Exp[x],{y[0]==1,y'[0]==2,y''[0]==3}},y[x],x,Inclu
```

$$y(x) \rightarrow \frac{1}{60}e^x(x^5 + 60x + 60)$$

5.15 problem Problem 24.44

5.15.1 Existence and uniqueness analysis	579
5.15.2 Maple step by step solution	582

Internal problem ID [5214]

Internal file name [OUTPUT/4707_Sunday_June_05_2022_03_03_46_PM_75360293/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.44.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$x'' + 4x' + 4x = 0$$

With initial conditions

$$[x(0) = 2, x'(0) = -2]$$

5.15.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$x'' + p(t)x' + q(t)x = F$$

Where here

$$p(t) = 4$$

$$q(t) = 4$$

$$F = 0$$

Hence the ode is

$$x'' + 4x' + 4x = 0$$

The domain of $p(t) = 4$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = 4$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(x) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(x') = sY(s) - x(0)$$

$$\mathcal{L}(x'') = s^2Y(s) - x'(0) - sx(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - x'(0) - sx(0) + 4sY(s) - 4x(0) + 4Y(s) = 0 \quad (1)$$

But the initial conditions are

$$x(0) = 2$$

$$x'(0) = -2$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 6 - 2s + 4sY(s) + 4Y(s) = 0$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{2s + 6}{s^2 + 4s + 4}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{2}{s + 2} + \frac{2}{(s + 2)^2}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{2}{s+2}\right) = 2e^{-2t}$$
$$\mathcal{L}^{-1}\left(\frac{2}{(s+2)^2}\right) = 2te^{-2t}$$

Adding the above results and simplifying gives

$$x = 2(t+1)e^{-2t}$$


Simplifying the solution gives

$$x = 2(t+1)e^{-2t}$$


Summary

The solution(s) found are the following

$$x = 2(t+1)e^{-2t} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = 2(t+1)e^{-2t}$$

Verified OK.

5.15.2 Maple step by step solution

Let's solve

$$\left[x'' + 4x' + 4x = 0, x(0) = 2, x' \Big|_{\{t=0\}} = -2 \right]$$

- Highest derivative means the order of the ODE is 2
 x''
- Characteristic polynomial of ODE
 $r^2 + 4r + 4 = 0$
- Factor the characteristic polynomial
 $(r + 2)^2 = 0$
- Root of the characteristic polynomial
 $r = -2$
- 1st solution of the ODE
 $x_1(t) = e^{-2t}$
- Repeated root, multiply $x_1(t)$ by t to ensure linear independence
 $x_2(t) = t e^{-2t}$
- General solution of the ODE
 $x = c_1 x_1(t) + c_2 x_2(t)$
- Substitute in solutions
 $x = c_1 e^{-2t} + c_2 t e^{-2t}$
- Check validity of solution $x = c_1 e^{-2t} + c_2 t e^{-2t}$
 - Use initial condition $x(0) = 2$
 $2 = c_1$
 - Compute derivative of the solution
 $x' = -2c_1 e^{-2t} + c_2 e^{-2t} - 2c_2 t e^{-2t}$
 - Use the initial condition $x' \Big|_{\{t=0\}} = -2$
 $-2 = -2c_1 + c_2$
 - Solve for c_1 and c_2
 $\{c_1 = 2, c_2 = 2\}$

- Substitute constant values into general solution and simplify

$$x = 2(t + 1)e^{-2t}$$

- Solution to the IVP

$$x = 2(t + 1)e^{-2t}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.453 (sec). Leaf size: 13

```
dsolve([diff(x(t),t$2)+4*diff(x(t),t)+4*x(t)=0,x(0) = 2, D(x)(0) = -2],x(t), singsol=all)
```

$$x(t) = 2(t + 1)e^{-2t}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 47

```
DSolve[{x''[t]+3*x'[t]+4*x[t]==0,{x[0]==2,x'[0]==-2}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{2}{7}e^{-3t/2} \left(\sqrt{7} \sin \left(\frac{\sqrt{7}t}{2} \right) + 7 \cos \left(\frac{\sqrt{7}t}{2} \right) \right)$$

5.16 problem Problem 24.46

5.16.1 Existence and uniqueness analysis	584
5.16.2 Maple step by step solution	587

Internal problem ID [5215]

Internal file name [OUTPUT/4708_Sunday_June_05_2022_03_03_47_PM_35789450/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 24. Solutions of linear DE by Laplace transforms. Supplementary Problems. page 248

Problem number: Problem 24.46.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$q'' + 9q' + 14q = \frac{\sin(t)}{2}$$

With initial conditions

$$[q(0) = 0, q'(0) = 1]$$

5.16.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$q'' + p(t)q' + q(t)q = F$$

Where here

$$p(t) = 9$$

$$q(t) = 14$$

$$F = \frac{\sin(t)}{2}$$

Hence the ode is

$$q'' + 9q' + 14q = \frac{\sin(t)}{2}$$

The domain of $p(t) = 9$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = 14$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. The domain of $F = \frac{\sin(t)}{2}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(q) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(q') &= sY(s) - q(0) \\ \mathcal{L}(q'') &= s^2Y(s) - q'(0) - sq(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - q'(0) - sq(0) + 9sY(s) - 9q(0) + 14Y(s) = \frac{1}{2s^2 + 2} \quad (1)$$

But the initial conditions are

$$\begin{aligned}q(0) &= 0 \\ q'(0) &= 1\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 1 + 9sY(s) + 14Y(s) = \frac{1}{2s^2 + 2}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{2s^2 + 3}{2(s^2 + 1)(s^2 + 9s + 14)}$$

Applying partial fractions decomposition results in

$$Y(s) = -\frac{101}{500(s+7)} + \frac{11}{50(s+2)} + \frac{-\frac{9}{1000} - \frac{13i}{1000}}{s-i} + \frac{-\frac{9}{1000} + \frac{13i}{1000}}{s+i}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(-\frac{101}{500(s+7)}\right) &= -\frac{101 e^{-7t}}{500} \\ \mathcal{L}^{-1}\left(\frac{11}{50(s+2)}\right) &= \frac{11 e^{-2t}}{50} \\ \mathcal{L}^{-1}\left(\frac{-\frac{9}{1000} - \frac{13i}{1000}}{s-i}\right) &= \left(-\frac{9}{1000} - \frac{13i}{1000}\right) e^{it} \\ \mathcal{L}^{-1}\left(\frac{-\frac{9}{1000} + \frac{13i}{1000}}{s+i}\right) &= \left(-\frac{9}{1000} + \frac{13i}{1000}\right) e^{-it}\end{aligned}$$

Adding the above results and simplifying gives

$$q = -\frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500} - \frac{101 e^{-7t}}{500} + \frac{11 e^{-2t}}{50}$$


Simplifying the solution gives

$$q = -\frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500} - \frac{101 e^{-7t}}{500} + \frac{11 e^{-2t}}{50}$$


Summary

The solution(s) found are the following

$$q = -\frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500} - \frac{101 e^{-7t}}{500} + \frac{11 e^{-2t}}{50} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$q = -\frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500} - \frac{101 e^{-7t}}{500} + \frac{11 e^{-2t}}{50}$$

Verified OK.

5.16.2 Maple step by step solution

Let's solve

$$\left[q'' + 9q' + 14q = \frac{\sin(t)}{2}, q(0) = 0, q' \Big|_{\{t=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2
 q''
- Characteristic polynomial of homogeneous ODE
 $r^2 + 9r + 14 = 0$
- Factor the characteristic polynomial
 $(r + 7)(r + 2) = 0$
- Roots of the characteristic polynomial
 $r = (-7, -2)$
- 1st solution of the homogeneous ODE

$$q_1(t) = e^{-7t}$$

- 2nd solution of the homogeneous ODE

$$q_2(t) = e^{-2t}$$

- General solution of the ODE

$$q = c_1 q_1(t) + c_2 q_2(t) + q_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$q = c_1 e^{-7t} + c_2 e^{-2t} + q_p(t)$$

- Find a particular solution $q_p(t)$ of the ODE

- Use variation of parameters to find q_p here $f(t)$ is the forcing function

$$\left[q_p(t) = -q_1(t) \left(\int \frac{q_2(t)f(t)}{W(q_1(t),q_2(t))} dt \right) + q_2(t) \left(\int \frac{q_1(t)f(t)}{W(q_1(t),q_2(t))} dt \right), f(t) = \frac{\sin(t)}{2} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(q_1(t), q_2(t)) = \begin{bmatrix} e^{-7t} & e^{-2t} \\ -7e^{-7t} & -2e^{-2t} \end{bmatrix}$$

- Compute Wronskian

$$W(q_1(t), q_2(t)) = 5e^{-9t}$$

- Substitute functions into equation for $q_p(t)$

$$q_p(t) = -\frac{e^{-7t} \left(\int \sin(t) e^{7t} dt \right)}{10} + \frac{e^{-2t} \left(\int \sin(t) e^{2t} dt \right)}{10}$$

- Compute integrals

$$q_p(t) = -\frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500}$$

- Substitute particular solution into general solution to ODE

$$q = c_1 e^{-7t} + c_2 e^{-2t} - \frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500}$$

- Check validity of solution $q = c_1 e^{-7t} + c_2 e^{-2t} - \frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500}$

- Use initial condition $q(0) = 0$

$$0 = c_1 + c_2 - \frac{9}{500}$$

- Compute derivative of the solution

$$q' = -7c_1 e^{-7t} - 2c_2 e^{-2t} + \frac{9 \sin(t)}{500} + \frac{13 \cos(t)}{500}$$

- Use the initial condition $q' \Big|_{\{t=0\}} = 1$

$$1 = -7c_1 - 2c_2 + \frac{13}{500}$$

- Solve for c_1 and c_2
- Substitute constant values into general solution and simplify

$$q = -\frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500} - \frac{101 e^{-7t}}{500} + \frac{11 e^{-2t}}{50}$$

- Solution to the IVP

$$q = -\frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500} - \frac{101 e^{-7t}}{500} + \frac{11 e^{-2t}}{50}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.531 (sec). Leaf size: 25

```
dsolve([diff(q(t),t$2)+9*diff(q(t),t)+14*q(t)=1/2*sin(t),q(0) = 0, D(q)(0) = 1],q(t), singsol
```

$$q(t) = -\frac{9 \cos(t)}{500} + \frac{13 \sin(t)}{500} - \frac{101 e^{-7t}}{500} + \frac{11 e^{-2t}}{50}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 32

```
DSolve[{q'[t]+9*q[t]+14*q[t]==1/2*Sin[t],{q[0]==0,q'[0]==1}},q[t],t,IncludeSingularSolutio
```

$$q(t) \rightarrow \frac{1}{500}(-101e^{-7t} + 110e^{-2t} + 13 \sin(t) - 9 \cos(t))$$

6 Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary

Problems. page 274

6.1	problem Problem 27.28	591
6.2	problem Problem 27.30	601
6.3	problem Problem 27.36	604
6.4	problem Problem 27.37	613
6.5	problem Problem 27.38	622
6.6	problem Problem 27.39	631
6.7	problem Problem 27.40	642
6.8	problem Problem 27.41	651
6.9	problem Problem 27.42	660
6.10	problem Problem 27.48	669

6.1 problem Problem 27.28

Internal problem ID [5216]

Internal file name [OUTPUT/4709_Sunday_June_05_2022_03_03_48_PM_60152543/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.28.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x + 1)y'' + \frac{y'}{x} + xy = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x + 1)y'' + \frac{y'}{x} + xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x(x+1)}$$
$$q(x) = \frac{x}{x+1}$$

Table 88: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x(x+1)}$	
singularity	type
$x = -1$	“regular”
$x = 0$	“regular”

$q(x) = \frac{x}{x+1}$	
singularity	type
$x = -1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-1, 0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$y''x(x+1) + y' + yx^2 = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x(x+1) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x^2 = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} x^{2+n+r} a_n \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \\ \sum_{n=0}^{\infty} x^{2+n+r} a_n &= \sum_{n=3}^{\infty} a_{n-3} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=3}^{\infty} a_{n-3} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^{-1+r} r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^{-1+r}r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = -\frac{r(-1+r)}{(1+r)^2}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{r^2(-1+r)}{(1+r)(2+r)^2}$$

For $3 \leq n$ the recursive equation is

$$a_{n-1}(n+r-1)(n+r-2) + a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-3} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{n^2 a_{n-1} + 2nra_{n-1} + r^2 a_{n-1} - 3na_{n-1} - 3ra_{n-1} + a_{n-3} + 2a_{n-1}}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{(-n^2 + 3n - 2)a_{n-1} - a_{n-3}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r(-1+r)}{(1+r)^2}$	0
a_2	$\frac{r^2(-1+r)}{(1+r)(2+r)^2}$	0

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-r^3 + r^2 - r - 2}{(r+3)^2(2+r)}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{1}{9}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r(-1+r)}{(1+r)^2}$	0
a_2	$\frac{r^2(-1+r)}{(1+r)(2+r)^2}$	0
a_3	$\frac{-r^3+r^2-r-2}{(r+3)^2(2+r)}$	$-\frac{1}{9}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r^5 + r^4 + r^3 + 5r^2 + 2r + 2}{(r + 4)^2 (1 + r)^2 (r + 3)}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{24}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r(-1+r)}{(1+r)^2}$	0
a_2	$\frac{r^2(-1+r)}{(1+r)(2+r)^2}$	0
a_3	$\frac{-r^3+r^2-r-2}{(r+3)^2(2+r)}$	$-\frac{1}{9}$
a_4	$\frac{r^5+r^4+r^3+5r^2+2r+2}{(r+4)^2(1+r)^2(r+3)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-r^7 - 5r^6 - 10r^5 - 17r^4 - 25r^3 - 26r^2 - 16r - 8}{(r + 5)^2 (2 + r)^2 (1 + r)^2 (r + 4)}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{1}{50}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r(-1+r)}{(1+r)^2}$	0
a_2	$\frac{r^2(-1+r)}{(1+r)(2+r)^2}$	0
a_3	$\frac{-r^3+r^2-r-2}{(r+3)^2(2+r)}$	$-\frac{1}{9}$
a_4	$\frac{r^5+r^4+r^3+5r^2+2r+2}{(r+4)^2(1+r)^2(r+3)}$	$\frac{1}{24}$
a_5	$\frac{-r^7-5r^6-10r^5-17r^4-25r^3-26r^2-16r-8}{(r+5)^2(2+r)^2(1+r)^2(r+4)}$	$-\frac{1}{50}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots \\ &= 1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$
b_0	1	1	N/A since b_n starts from 1
b_1	$-\frac{r(-1+r)}{(1+r)^2}$	0	$\frac{-3r+1}{(1+r)^3}$
b_2	$\frac{r^2(-1+r)}{(1+r)(2+r)^2}$	0	$\frac{6r^3+4r^2-4r}{(1+r)^2(2+r)^3}$
b_3	$\frac{-r^3+r^2-r-2}{(r+3)^2(2+r)}$	$-\frac{1}{9}$	$\frac{-9r^3-13r^2+20r+8}{(r+3)^3(2+r)^2}$
b_4	$\frac{r^5+r^4+r^3+5r^2+2r+2}{(r+4)^2(1+r)^2(r+3)}$	$\frac{1}{24}$	$\frac{12r^6+64r^5+84r^4+8r^3+8r^2+36r-44}{(r+4)^3(1+r)^3(r+3)^2}$
b_5	$\frac{-r^7-5r^6-10r^5-17r^4-25r^3-26r^2-16r-8}{(r+5)^2(2+r)^2(1+r)^2(r+4)}$	$-\frac{1}{50}$	$\frac{-15r^9-184r^8-884r^7-2178r^6-3099r^5-2886r^4-1926r^3-668r^2+440r+528}{(r+5)^3(2+r)^3(1+r)^3(r+4)^2}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots \\ &= \left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \ln(x) + x + \frac{2x^3}{27} - \frac{11x^4}{144} + \frac{33x^5}{1000} + O(x^6) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1 \left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \\ &\quad + c_2 \left(\left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \ln(x) + x + \frac{2x^3}{27} - \frac{11x^4}{144} + \frac{33x^5}{1000} + O(x^6) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}y &= y_h \\&= c_1 \left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \\&\quad + c_2 \left(\left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \ln(x) + x + \frac{2x^3}{27} - \frac{11x^4}{144} + \frac{33x^5}{1000} + O(x^6) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= c_1 \left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \\&\quad + c_2 \left(\left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \ln(x) + x + \frac{2x^3}{27} - \frac{11x^4}{144} + \frac{33x^5}{1000} + O(x^6) \right) \quad (1)\end{aligned}$$

Verification of solutions

$$\begin{aligned}y &= c_1 \left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \\&\quad + c_2 \left(\left(1 - \frac{x^3}{9} + \frac{x^4}{24} - \frac{x^5}{50} + O(x^6) \right) \ln(x) + x + \frac{2x^3}{27} - \frac{11x^4}{144} + \frac{33x^5}{1000} + O(x^6) \right)\end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunC ODE, case a <> 0, e <> 0, c =
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 49

```
Order:=6;
dsolve((x+1)*diff(y(x),x$2)+1/x*diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - \frac{1}{9}x^3 + \frac{1}{24}x^4 - \frac{1}{50}x^5 + O(x^6) \right) \\ + \left(x + \frac{2}{27}x^3 - \frac{11}{144}x^4 + \frac{33}{1000}x^5 + O(x^6) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 82

```
AsymptoticDSolveValue[(1+x)*y'[x]+1/x*y'[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{50} + \frac{x^4}{24} - \frac{x^3}{9} + 1 \right) + c_2 \left(\frac{33x^5}{1000} - \frac{11x^4}{144} + \frac{2x^3}{27} + \left(-\frac{x^5}{50} + \frac{x^4}{24} - \frac{x^3}{9} + 1 \right) \log(x) + x \right)$$

6.2 problem Problem 27.30

Internal problem ID [5217]

Internal file name [OUTPUT/4710_Sunday_June_05_2022_03_03_50_PM_5107071/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.30.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_bessel_ode", "second order series method. Irregular singular point"

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

Unable to solve or complete the solution.

$$x^3 y'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^3 y'' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$
$$q(x) = \frac{1}{x^3}$$

Table 89: Table $p(x), q(x)$ singularities.

$p(x) = 0$	
singularity	type

$q(x) = \frac{1}{x^3}$	
singularity	type
$x = 0$	“irregular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[\infty]$

Irregular singular points : $[0]$

Since $x = 0$ is not an ordinary point, then we will now check if it is a regular singular point. Unable to solve since $x = 0$ is not regular singular point. Terminating.

Verification of solutions N/A

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

X Solution by Maple

```

Order:=6;
dsolve(x^3*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);

```

No solution found

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 222

```
AsymptoticDSolveValue[x^3*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 e^{-\frac{2i}{\sqrt{x}}x^{3/4}} \left(-\frac{468131288625ix^{9/2}}{8796093022208} + \frac{66891825ix^{7/2}}{4294967296} - \frac{72765ix^{5/2}}{8388608} + \frac{105ix^{3/2}}{8192} \right. \\ \left. + \frac{33424574007825x^5}{281474976710656} - \frac{14783093325x^4}{549755813888} + \frac{2837835x^3}{268435456} - \frac{4725x^2}{524288} + \frac{15x}{512} - \frac{3i\sqrt{x}}{16} \right. \\ \left. + 1 \right) + c_2 e^{\frac{2i}{\sqrt{x}}x^{3/4}} \left(\frac{468131288625ix^{9/2}}{8796093022208} - \frac{66891825ix^{7/2}}{4294967296} + \frac{72765ix^{5/2}}{8388608} - \frac{105ix^{3/2}}{8192} + \frac{33424574007825x^5}{281474976710656} - \right.$$

6.3 problem Problem 27.36

6.3.1 Maple step by step solution 611

Internal problem ID [5218]

Internal file name [OUTPUT/4711_Sunday_June_05_2022_03_03_51_PM_32978055/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.36.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$y'' + xy = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{125}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{126}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -xy \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= -y - xy' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= -2y' + yx^2 \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= x(xy' + 4y) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= -yx^3 + 6xy' + 4y
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= -y(0) \\
 F_2 &= -2y'(0) \\
 F_3 &= 0 \\
 F_4 &= 4y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right)y(0) + \left(x - \frac{1}{12}x^4\right)y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -x \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} x^{1+n} a_n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=0}^{\infty} x^{1+n} a_n = \sum_{n=1}^{\infty} a_{n-1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^n \right) = 0 \quad (3)$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (1+n) + a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_{n-1}}{(n+2)(1+n)} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$6a_3 + a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_1}{12}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$30a_6 + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{180}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{a_1}{504}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{6} a_0 x^3 - \frac{1}{12} a_1 x^4 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{x^3}{6}\right) a_0 + \left(x - \frac{1}{12} x^4\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{x^3}{6}\right) c_1 + \left(x - \frac{1}{12} x^4\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{6} x^3 + \frac{1}{180} x^6\right) y(0) + \left(x - \frac{1}{12} x^4\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{x^3}{6}\right) c_1 + \left(x - \frac{1}{12} x^4\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{6} x^3 + \frac{1}{180} x^6\right) y(0) + \left(x - \frac{1}{12} x^4\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{x^3}{6}\right) c_1 + \left(x - \frac{1}{12} x^4\right) c_2 + O(x^6)$$

Verified OK.

6.3.1 Maple step by step solution

Let's solve

$$y'' = -xy$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) + a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} + a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{a_k}{k^2+5k+6}, 2a_2 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```

Order:=6;
dsolve(diff(y(x),x$2)+x*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 - \frac{x^3}{6}\right) y(0) + \left(x - \frac{1}{12}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```

AsymptoticDSolveValue[y'[x]+x*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{12}\right) + c_1 \left(1 - \frac{x^3}{6}\right)$$

6.4 problem Problem 27.37

6.4.1 Maple step by step solution 620

Internal problem ID [5219]

Internal file name [OUTPUT/4712_Sunday_June_05_2022_03_03_52_PM_83320885/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.37.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact linear second order ode", "second_order_integrable_as_is", "second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$y'' - 2xy' - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (128)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (129)$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \quad (2)
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \quad (3)
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= 2xy' + 2y \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= 4x^2y' + 4xy + 4y' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= 8y'x^3 + 8yx^2 + 20xy' + 12y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= (16x^4 + 72x^2 + 32)y' + (16x^3 + 56x)y \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= (32x^5 + 224x^3 + 264x)y' + 32y\left(x^4 + 6x^2 + \frac{15}{4}\right)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 2y(0) \\
 F_1 &= 4y'(0) \\
 F_2 &= 12y(0) \\
 F_3 &= 32y'(0) \\
 F_4 &= 120y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6\right)y(0) + \left(x + \frac{2}{3}x^3 + \frac{4}{15}x^5\right)y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-2n x^n a_n) + \sum_{n=0}^{\infty} (-2a_n x^n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \sum_{n=1}^{\infty} (-2n x^n a_n) + \sum_{n=0}^{\infty} (-2a_n x^n) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 - 2a_0 = 0$$

$$a_2 = a_0$$

For $1 \leq n$, the recurrence equation is

$$(n + 2) a_{n+2}(n + 1) - 2na_n - 2a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{2a_n}{n + 2} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$6a_3 - 4a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{2a_1}{3}$$

For $n = 2$ the recurrence equation gives

$$12a_4 - 6a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{2}$$

For $n = 3$ the recurrence equation gives

$$20a_5 - 8a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{4a_1}{15}$$

For $n = 4$ the recurrence equation gives

$$30a_6 - 10a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{6}$$

For $n = 5$ the recurrence equation gives

$$42a_7 - 12a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{8a_1}{105}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + a_0 x^2 + \frac{2}{3} a_1 x^3 + \frac{1}{2} a_0 x^4 + \frac{4}{15} a_1 x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + x^2 + \frac{1}{2}x^4\right) a_0 + \left(x + \frac{2}{3}x^3 + \frac{4}{15}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + x^2 + \frac{1}{2}x^4\right) c_1 + \left(x + \frac{2}{3}x^3 + \frac{4}{15}x^5\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6\right) y(0) + \left(x + \frac{2}{3}x^3 + \frac{4}{15}x^5\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 + x^2 + \frac{1}{2}x^4\right) c_1 + \left(x + \frac{2}{3}x^3 + \frac{4}{15}x^5\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6\right) y(0) + \left(x + \frac{2}{3}x^3 + \frac{4}{15}x^5\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 + x^2 + \frac{1}{2}x^4\right) c_1 + \left(x + \frac{2}{3}x^3 + \frac{4}{15}x^5\right) c_2 + O(x^6)$$

Verified OK.

6.4.1 Maple step by step solution

Let's solve

$$y'' = 2xy' + 2y$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - 2xy' - 2y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1) x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (a_{k+2}(k+2)(k+1) - 2a_k(k+1)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k+1)(a_{k+2}(k+2) - 2a_k) = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = \frac{2a_k}{k+2} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
Order:=6;  
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x^2 + \frac{1}{2}x^4\right) y(0) + \left(x + \frac{2}{3}x^3 + \frac{4}{15}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 38

```
AsymptoticDSolveValue[y'[x]-2*x*y'[x]-2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{4x^5}{15} + \frac{2x^3}{3} + x \right) + c_1 \left(\frac{x^4}{2} + x^2 + 1 \right)$$

6.5 problem Problem 27.38

6.5.1 Maple step by step solution 629

Internal problem ID [5220]

Internal file name [OUTPUT/4713_Sunday_June_05_2022_03_03_53_PM_98761641/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.38.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact linear second order ode", "second_order_integrable_as_is", "second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$y'' + x^2y' + 2xy = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{131}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{132}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -x^2 y' - 2xy \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= (x^4 - 4x) y' + (2x^3 - 2) y \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= (-x^6 + 10x^3 - 6) y' - 2yx^2(x^3 - 7) \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= x((x^7 - 18x^4 + 50x) y' + 2y(x^6 - 15x^3 + 20)) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= (-x^{10} + 28x^7 - 170x^4 + 140x) y' - 2y(x^9 - 25x^6 + 110x^3 - 20)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= -2y(0) \\
 F_2 &= -6y'(0) \\
 F_3 &= 0 \\
 F_4 &= 40y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{3}x^3 + \frac{1}{18}x^6\right) y(0) + \left(x - \frac{1}{4}x^4\right) y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -x^2 \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - 2x \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n x^{1+n} a_n \right) + \left(\sum_{n=0}^{\infty} 2x^{1+n} a_n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=1}^{\infty} n x^{1+n} a_n = \sum_{n=2}^{\infty} (n-1) a_{n-1} x^n$$

$$\sum_{n=0}^{\infty} 2x^{1+n} a_n = \sum_{n=1}^{\infty} 2a_{n-1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=2}^{\infty} (n-1) a_{n-1} x^n \right) + \left(\sum_{n=1}^{\infty} 2a_{n-1} x^n \right) = 0 \quad (3)$$

$n = 1$ gives

$$6a_3 + 2a_0 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{a_0}{3}$$

For $2 \leq n$, the recurrence equation is

$$(n + 2) a_{n+2}(1 + n) + (n - 1) a_{n-1} + 2a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_{n-1}}{n + 2} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 3a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_1}{4}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 4a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 5a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{18}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 6a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{a_1}{28}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{3} a_0 x^3 - \frac{1}{4} a_1 x^4 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{x^3}{3}\right) a_0 + \left(x - \frac{1}{4} x^4\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{x^3}{3}\right) c_1 + \left(x - \frac{1}{4} x^4\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{3} x^3 + \frac{1}{18} x^6\right) y(0) + \left(x - \frac{1}{4} x^4\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{x^3}{3}\right) c_1 + \left(x - \frac{1}{4} x^4\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{3} x^3 + \frac{1}{18} x^6\right) y(0) + \left(x - \frac{1}{4} x^4\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{x^3}{3}\right) c_1 + \left(x - \frac{1}{4} x^4\right) c_2 + O(x^6)$$

Verified OK.

6.5.1 Maple step by step solution

Let's solve

$$y'' = -x^2y' - 2xy$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + x^2y' + 2xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k- > k-1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k k x^{k+1}$$

- Shift index using $k- > k-1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) + a_{k-1}(k+1)) x^k \right) = 0$$

- Each term must be 0
 $2a_2 = 0$
- Each term in the series must be 0, giving the recursion relation
 $(k + 1)(a_{k+2}(k + 2) + a_{k-1}) = 0$
- Shift index using $k \rightarrow k + 1$
 $(k + 2)(a_{k+3}(k + 3) + a_k) = 0$
- Recursion relation that defines the series solution to the ODE
$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{a_k}{k+3}, 2a_2 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```

Order:=6;
dsolve(diff(y(x),x$2)+x^2*diff(y(x),x)+2*x*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 - \frac{x^3}{3}\right) y(0) + \left(x - \frac{1}{4}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```

AsymptoticDSolveValue[y'[x]+x^2*y'[x]+2*x*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{4}\right) + c_1 \left(1 - \frac{x^3}{3}\right)$$

6.6 problem Problem 27.39

6.6.1 Maple step by step solution 638

Internal problem ID [5221]

Internal file name [OUTPUT/4714_Sunday_June_05_2022_03_03_54_PM_95464487/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.39.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - x^2y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (134)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (135)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = x^2 y' + y$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= (x^4 + 2x + 1) y' + yx^2 \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= (x^6 + 6x^3 + 2x^2 + 2) y' + y(x^4 + 4x + 1) \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= (x^8 + 12x^5 + 3x^4 + 20x^2 + 8x + 1) y' + y(x^6 + 10x^3 + 2x^2 + 6) \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= (x^{10} + 20x^7 + 4x^6 + 80x^4 + 30x^3 + 3x^2 + 40x + 14) y' + y(x^8 + 18x^5 + 3x^4 + 50x^2 + 12x + 1) \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= y(0) \\ F_1 &= y'(0) \\ F_2 &= 2y'(0) + y(0) \\ F_3 &= y'(0) + 6y(0) \\ F_4 &= 14y'(0) + y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4 + \frac{1}{20}x^5 + \frac{1}{720}x^6 \right) y(0) \\ &\quad + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5 + \frac{7}{360}x^6 \right) y'(0) + O(x^6) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = x^2 \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-n x^{1+n} a_n) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=1}^{\infty} (-n x^{1+n} a_n) = \sum_{n=2}^{\infty} (-(n-1) a_{n-1} x^n)$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \sum_{n=2}^{\infty} (-(n-1) a_{n-1} x^n) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 - a_0 = 0$$

$$a_2 = \frac{a_0}{2}$$

$n = 1$ gives

$$6a_3 - a_1 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = \frac{a_1}{6}$$

For $2 \leq n$, the recurrence equation is

$$(n+2)a_{n+2}(1+n) - (n-1)a_{n-1} - a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned} a_{n+2} &= \frac{na_{n-1} + a_n - a_{n-1}}{(n+2)(1+n)} \\ (5) \quad &= \frac{a_n}{(n+2)(1+n)} + \frac{(n-1)a_{n-1}}{(n+2)(1+n)} \end{aligned}$$

For $n = 2$ the recurrence equation gives

$$12a_4 - a_1 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_1}{12} + \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$20a_5 - 2a_2 - a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_0}{20} + \frac{a_1}{120}$$

For $n = 4$ the recurrence equation gives

$$30a_6 - 3a_3 - a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{7a_1}{360} + \frac{a_0}{720}$$

For $n = 5$ the recurrence equation gives

$$42a_7 - 4a_4 - a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{41a_1}{5040} + \frac{13a_0}{2520}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + \frac{a_0 x^2}{2} + \frac{a_1 x^3}{6} + \left(\frac{a_1}{12} + \frac{a_0}{24} \right) x^4 + \left(\frac{a_0}{20} + \frac{a_1}{120} \right) x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4 + \frac{1}{20}x^5 \right) a_0 + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5 \right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4 + \frac{1}{20}x^5 \right) c_1 + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5 \right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4 + \frac{1}{20}x^5 + \frac{1}{720}x^6 \right) y(0) \\ &\quad + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5 + \frac{7}{360}x^6 \right) y'(0) + O(x^6) \end{aligned} \quad (1)$$

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4 + \frac{1}{20}x^5 \right) c_1 + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5 \right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4 + \frac{1}{20}x^5 + \frac{1}{720}x^6\right) y(0) \\ + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5 + \frac{7}{360}x^6\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4 + \frac{1}{20}x^5\right) c_1 + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5\right) c_2 + O(x^6)$$

Verified OK.

6.6.1 Maple step by step solution

Let's solve

$$y'' = x^2 y' + y$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - x^2 y' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k k x^{k+1}$$

- Shift index using $k- > k-1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1)x^k$$

Rewrite ODE with series expansions

$$2a_2 - a_0 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) - a_k - a_{k-1}(k-1))x^k \right) = 0$$

- Each term must be 0
 $2a_2 - a_0 = 0$
- Each term in the series must be 0, giving the recursion relation
 $(k^2 + 3k + 2)a_{k+2} - a_{k-1}k - a_k + a_{k-1} = 0$
- Shift index using $k \rightarrow k+1$
 $((k+1)^2 + 3k + 5)a_{k+3} - a_k(k+1) - a_{k+1} + a_k = 0$
- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = \frac{a_k k + a_{k+1}}{k^2 + 5k + 6}, 2a_2 - a_0 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
      -> heuristic approach
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
      -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunT ODE, case c = 0`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
Order:=6;
dsolve(diff(y(x),x$2)-x^2*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4 + \frac{1}{20}x^5\right) y(0) + \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[y''[x]-x^2*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} + \frac{x^4}{12} + \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^5}{20} + \frac{x^4}{24} + \frac{x^2}{2} + 1 \right)$$

6.7 problem Problem 27.40

6.7.1 Maple step by step solution 648

Internal problem ID [5222]

Internal file name [OUTPUT/4715_Sunday_June_05_2022_03_03_55_PM_79979292/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.40.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**", "**second order series method. Ordinary point**", "**second order series method. Taylor series method**"

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$y'' + 2yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \dots \\
 &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{137}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{138}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -2yx^2 \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= -2x(xy' + 2y) \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= 4yx^4 - 8xy' - 4y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= 4y'x^4 + 32yx^3 - 12y' \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= 48y'x^3 - 8x^2y(x^4 - 15)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= 0 \\
 F_2 &= -4y(0) \\
 F_3 &= -12y'(0) \\
 F_4 &= 0
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{x^4}{6}\right)y(0) + \left(x - \frac{1}{10}x^5\right)y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -2 \left(\sum_{n=0}^{\infty} a_n x^n \right) x^2 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} 2x^{n+2} a_n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

$$\sum_{n=0}^{\infty} 2x^{n+2} a_n = \sum_{n=2}^{\infty} 2a_{n-2} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=2}^{\infty} 2a_{n-2} x^n \right) = 0 \quad (3)$$

For $2 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (n+1) + 2a_{n-2} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{2a_{n-2}}{(n+2)(n+1)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 2a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_0}{6}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 2a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_1}{10}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 2a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = 0$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 2a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1x - \frac{1}{6}a_0x^4 - \frac{1}{10}a_1x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{x^4}{6}\right) a_0 + \left(x - \frac{1}{10}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{x^4}{6}\right) c_1 + \left(x - \frac{1}{10}x^5\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{x^4}{6}\right) y(0) + \left(x - \frac{1}{10}x^5\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{x^4}{6}\right) c_1 + \left(x - \frac{1}{10}x^5\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{x^4}{6}\right) y(0) + \left(x - \frac{1}{10}x^5\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{x^4}{6}\right) c_1 + \left(x - \frac{1}{10}x^5\right) c_2 + O(x^6)$$

Verified OK.

6.7.1 Maple step by step solution

Let's solve

$$y'' = -2yx^2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + 2yx^2 = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^2 \cdot y$ to series expansion

$$x^2 \cdot y = \sum_{k=0}^{\infty} a_k x^{k+2}$$

- Shift index using $k \rightarrow k - 2$

$$x^2 \cdot y = \sum_{k=2}^{\infty} a_{k-2} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$6a_3x + 2a_2 + \left(\sum_{k=2}^{\infty} (a_{k+2}(k+2)(k+1) + 2a_{k-2}) x^k \right) = 0$$

- The coefficients of each power of x must be 0

$$[2a_2 = 0, 6a_3 = 0]$$

- Solve for the dependent coefficient(s)

$$\{a_2 = 0, a_3 = 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} + 2a_{k-2} = 0$$

- Shift index using $k \rightarrow k + 2$

$$((k+2)^2 + 3k + 8) a_{k+4} + 2a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = -\frac{2a_k}{k^2 + 7k + 12}, a_2 = 0, a_3 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+2*x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^4}{6}\right) y(0) + \left(x - \frac{1}{10}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]+2*x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^5}{10}\right) + c_1 \left(1 - \frac{x^4}{6}\right)$$

6.8 problem Problem 27.41

6.8.1 Maple step by step solution 658

Internal problem ID [5223]

Internal file name [OUTPUT/4716_Sunday_June_05_2022_03_03_56_PM_64263502/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.41.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact linear second order ode", "second_order_integrable_as_is", "second order series method. Ordinary point", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2", "second order series method. Taylor series method", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$(x^2 - 1)y'' + xy' - y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \dots \\
 &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{140}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{141}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -\frac{-y + xy'}{x^2 - 1} \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= \frac{3(-y + xy')x}{(x^2 - 1)^2} \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= -\frac{3(-y + xy')(4x^2 + 1)}{(x^2 - 1)^3} \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= \frac{60(x^2 + \frac{3}{4})x(-y + xy')}{(x^2 - 1)^4} \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= -\frac{45(-y + xy')(8x^4 + 12x^2 + 1)}{(x^2 - 1)^5}
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= -y(0) \\
 F_1 &= 0 \\
 F_2 &= -3y(0) \\
 F_3 &= 0 \\
 F_4 &= -45y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6\right) y(0) + xy'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$(x^2 - 1) y'' + xy' - y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$(x^2 - 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \sum_{n=2}^{\infty} (-n(n-1) a_n x^{n-2}) + \left(\sum_{n=1}^{\infty} n a_n x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} (-n(n-1) a_n x^{n-2}) = \sum_{n=0}^{\infty} (-(n+2) a_{n+2} (n+1) x^n)$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \sum_{n=0}^{\infty} (-(n+2) a_{n+2} (n+1) x^n) + \left(\sum_{n=1}^{\infty} n a_n x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (3)$$

$n = 0$ gives

$$-2a_2 - a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

For $2 \leq n$, the recurrence equation is

$$na_n(n-1) - (n+2)a_{n+2}(n+1) + na_n - a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{(n-1)a_n}{n+2} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$3a_2 - 12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_0}{8}$$

For $n = 3$ the recurrence equation gives

$$8a_3 - 20a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$15a_4 - 30a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{16}$$

For $n = 5$ the recurrence equation gives

$$24a_5 - 42a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 - \frac{1}{8} a_0 x^4 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4\right) a_0 + a_1 x + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4\right) c_1 + c_2 x + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6\right) y(0) + xy'(0) + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4\right) c_1 + c_2 x + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6\right) y(0) + xy'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4\right) c_1 + c_2 x + O(x^6)$$

Verified OK.

6.8.1 Maple step by step solution

Let's solve

$$(x^2 - 1)y'' + xy' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{xy'}{x^2-1} + \frac{y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{xy'}{x^2-1} - \frac{y}{x^2-1} = 0$$

- Multiply by denominators of ODE

$$(-x^2 + 1)y'' - xy' + y = 0$$

- Make a change of variables

$$\theta = \arccos(x)$$

- Calculate y' with change of variables

$$y' = \left(\frac{d}{d\theta}y(\theta)\right)\theta'(x)$$

- Compute 1st derivative y'

$$y' = -\frac{\frac{d}{d\theta}y(\theta)}{\sqrt{-x^2+1}}$$

- Calculate y'' with change of variables

$$y'' = \left(\frac{d^2}{d\theta^2}y(\theta)\right)\theta'(x)^2 + \theta''(x)\left(\frac{d}{d\theta}y(\theta)\right)$$

- Compute 2nd derivative y''

$$y'' = \frac{\frac{d^2}{d\theta^2}y(\theta)}{-x^2+1} - \frac{x\left(\frac{d}{d\theta}y(\theta)\right)}{(-x^2+1)^{\frac{3}{2}}}$$

- Apply the change of variables to the ODE

$$(-x^2 + 1)\left(\frac{\frac{d^2}{d\theta^2}y(\theta)}{-x^2+1} - \frac{x\left(\frac{d}{d\theta}y(\theta)\right)}{(-x^2+1)^{\frac{3}{2}}}\right) + \frac{x\left(\frac{d}{d\theta}y(\theta)\right)}{\sqrt{-x^2+1}} + y = 0$$

- Multiply through

$$-\frac{\left(\frac{d^2}{d\theta^2}y(\theta)\right)x^2}{-x^2+1} + \frac{\frac{d^2}{d\theta^2}y(\theta)}{-x^2+1} + \frac{x^3\left(\frac{d}{d\theta}y(\theta)\right)}{(-x^2+1)^{\frac{3}{2}}} - \frac{x\left(\frac{d}{d\theta}y(\theta)\right)}{(-x^2+1)^{\frac{3}{2}}} + \frac{x\left(\frac{d}{d\theta}y(\theta)\right)}{\sqrt{-x^2+1}} + y = 0$$

- Simplify ODE

$$y + \frac{d^2}{d\theta^2}y(\theta) = 0$$

- ODE is that of a harmonic oscillator with given general solution

$$y(\theta) = c_1 \sin(\theta) + c_2 \cos(\theta)$$

- Revert back to x

$$y = c_1 \sin(\arccos(x)) + c_2 \cos(\arccos(x))$$

- Use trig identity to simplify $\sin(\arccos(x))$

$$\sin(\arccos(x)) = \sqrt{-x^2 + 1}$$

- Simplify solution to the ODE

$$y = c_1 \sqrt{-x^2 + 1} + c_2 x$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;
dsolve((x^2-1)*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4\right) y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 27

```
AsymptoticDSolveValue[(x^2-1)*y''[x]+x*y'[x]-y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^4}{8} - \frac{x^2}{2} + 1\right) + c_2 x$$

6.9 problem Problem 27.42

6.9.1 Maple step by step solution 667

Internal problem ID [5224]

Internal file name [OUTPUT/4717_Sunday_June_05_2022_03_03_57_PM_70489940/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.42.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$y'' - xy = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{143}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{144}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= xy \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= y + xy' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= 2y' + yx^2 \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= x(xy' + 4y) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= yx^3 + 6xy' + 4y
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= y(0) \\
 F_2 &= 2y'(0) \\
 F_3 &= 0 \\
 F_4 &= 4y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 + \frac{1}{6}x^3 + \frac{1}{180}x^6\right) y(0) + \left(x + \frac{1}{12}x^4\right) y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = x \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=0}^{\infty} (-x^{1+n} a_n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=0}^{\infty} (-x^{1+n} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^n)$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^n) = 0 \quad (3)$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (1+n) - a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{a_{n-1}}{(n+2)(1+n)} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$6a_3 - a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{a_0}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 - a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_1}{12}$$

For $n = 3$ the recurrence equation gives

$$20a_5 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$30a_6 - a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{180}$$

For $n = 5$ the recurrence equation gives

$$42a_7 - a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{a_1}{504}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + \frac{1}{6} a_0 x^3 + \frac{1}{12} a_1 x^4 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + \frac{x^3}{6}\right) a_0 + \left(x + \frac{1}{12} x^4\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + \frac{x^3}{6}\right) c_1 + \left(x + \frac{1}{12} x^4\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 + \frac{1}{6} x^3 + \frac{1}{180} x^6\right) y(0) + \left(x + \frac{1}{12} x^4\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 + \frac{x^3}{6}\right) c_1 + \left(x + \frac{1}{12} x^4\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{1}{6} x^3 + \frac{1}{180} x^6\right) y(0) + \left(x + \frac{1}{12} x^4\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 + \frac{x^3}{6}\right) c_1 + \left(x + \frac{1}{12} x^4\right) c_2 + O(x^6)$$

Verified OK.

6.9.1 Maple step by step solution

Let's solve

$$y'' = xy$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + \left(\sum_{k=1}^{\infty} (a_{k+2} (k+2)(k+1) - a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} - a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = \frac{a_k}{k^2 + 5k + 6}, 2a_2 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```

Order:=6;
dsolve(diff(y(x),x$2)-x*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 + \frac{x^3}{6}\right) y(0) + \left(x + \frac{1}{12}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```

AsymptoticDSolveValue[y''[x]-x*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_2 \left(\frac{x^4}{12} + x \right) + c_1 \left(\frac{x^3}{6} + 1 \right)$$

6.10 problem Problem 27.48

6.10.1 Existence and uniqueness analysis	669
6.10.2 Maple step by step solution	677

Internal problem ID [5225]

Internal file name [OUTPUT/4718_Sunday_June_05_2022_03_03_57_PM_82975089/index.tex]

Book: Schaums Outline Differential Equations, 4th edition. Bronson and Costa. McGraw Hill 2014

Section: Chapter 27. Power series solutions of linear DE with variable coefficients. Supplementary Problems. page 274

Problem number: Problem 27.48.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_y_method_1", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2xy' + yx^2 = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -1]$$

With the expansion point for the power series method at $x = 0$.

6.10.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = -2x$$

$$q(x) = x^2$$

$$F = 0$$

Hence the ode is

$$y'' - 2xy' + yx^2 = 0$$

The domain of $p(x) = -2x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = x^2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (146)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (147)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= 2xy' - yx^2 \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= -2yx^3 + 3x^2y' - 2xy + 2y' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= (4x^3 + 8x)y' + (-3x^4 - 8x^2 - 2)y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= (5x^4 + 20x^2 + 6)y' - 4xy(x^2 + 4)(x^2 + 1) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= (6x^5 + 40x^3 + 36x)y' - 5y\left(x^6 + 8x^4 + \frac{66}{5}x^2 + \frac{16}{5}\right)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = 1$ and $y'(0) = -1$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= -2 \\
 F_2 &= -2 \\
 F_3 &= -6 \\
 F_4 &= -16
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = 1 - x - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{20} - \frac{x^6}{45} + O(x^6)$$

$$y = 1 - x - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{20} - \frac{x^6}{45} + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) x^2 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-2n x^n a_n) + \left(\sum_{n=0}^{\infty} x^{n+2} a_n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

$$\sum_{n=0}^{\infty} x^{n+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \sum_{n=1}^{\infty} (-2n x^n a_n) + \left(\sum_{n=2}^{\infty} a_{n-2} x^n \right) = 0 \quad (3)$$

$n = 1$ gives

$$6a_3 - 2a_1 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = \frac{a_1}{3}$$

For $2 \leq n$, the recurrence equation is

$$(n+2)a_{n+2}(n+1) - 2na_n + a_{n-2} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned} a_{n+2} &= \frac{2na_n - a_{n-2}}{(n+2)(n+1)} \\ (5) \quad &= \frac{2na_n}{(n+2)(n+1)} - \frac{a_{n-2}}{(n+2)(n+1)} \end{aligned}$$

For $n = 2$ the recurrence equation gives

$$12a_4 - 4a_2 + a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_0}{12}$$

For $n = 3$ the recurrence equation gives

$$20a_5 - 6a_3 + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{20}$$

For $n = 4$ the recurrence equation gives

$$30a_6 - 8a_4 + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{45}$$

For $n = 5$ the recurrence equation gives

$$42a_7 - 10a_5 + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{a_1}{252}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + \frac{1}{3} a_1 x^3 - \frac{1}{12} a_0 x^4 + \frac{1}{20} a_1 x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{x^4}{12}\right) a_0 + \left(x + \frac{1}{3} x^3 + \frac{1}{20} x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{x^4}{12}\right) c_1 + \left(x + \frac{1}{3} x^3 + \frac{1}{20} x^5\right) c_2 + O(x^6)$$

$$y = 1 - \frac{x^4}{12} - x - \frac{x^3}{3} - \frac{x^5}{20} + O(x^6)$$

Summary

The solution(s) found are the following

$$y = 1 - x - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{20} - \frac{x^6}{45} + O(x^6) \quad (1)$$

$$y = 1 - \frac{x^4}{12} - x - \frac{x^3}{3} - \frac{x^5}{20} + O(x^6) \quad (2)$$

Verification of solutions

$$y = 1 - x - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{20} - \frac{x^6}{45} + O(x^6)$$

Verified OK.

$$y = 1 - \frac{x^4}{12} - x - \frac{x^3}{3} - \frac{x^5}{20} + O(x^6)$$

Verified OK.

6.10.2 Maple step by step solution

Let's solve

$$\left[y'' = 2xy' - yx^2, y(0) = 1, y'|_{\{x=0\}} = -1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - 2xy' + yx^2 = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^2 \cdot y$ to series expansion

$$x^2 \cdot y = \sum_{k=0}^{\infty} a_k x^{k+2}$$

- Shift index using $k- > k - 2$

$$x^2 \cdot y = \sum_{k=2}^{\infty} a_{k-2} x^k$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1)x^k$$

Rewrite ODE with series expansions

$$2a_2 + (6a_3 - 2a_1)x + \left(\sum_{k=2}^{\infty} (a_{k+2}(k+2)(k+1) - 2a_k k + a_{k-2})x^k \right) = 0$$

- The coefficients of each power of x must be 0
 $[2a_2 = 0, 6a_3 - 2a_1 = 0]$
- Solve for the dependent coefficient(s)
 $\{a_2 = 0, a_3 = \frac{a_1}{3}\}$
- Each term in the series must be 0, giving the recursion relation
 $(k^2 + 3k + 2)a_{k+2} - 2a_k k + a_{k-2} = 0$
- Shift index using $k \rightarrow k + 2$
 $((k+2)^2 + 3k + 8)a_{k+4} - 2a_{k+2}(k+2) + a_k = 0$
- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = \frac{2ka_{k+2} - a_k + 4a_{k-2}}{k^2 + 7k + 12}, a_2 = 0, a_3 = \frac{a_1}{3} \right]$$

Maple trace **Kovacic algorithm successful**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

Order:=6;

```
dsolve([diff(y(x),x$2)-2*x*diff(y(x),x)+x^2*y(x)=0,y(0) = 1, D(y)(0) = -1],y(x),type='series')
```

$$y(x) = 1 - x - \frac{1}{3}x^3 - \frac{1}{12}x^4 - \frac{1}{20}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 29

```
AsymptoticDSolveValue[{y'[x]-2*x*y'[x]+x^2*y[x]==0,{y[0]==1,y'[0]==-1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^5}{20} - \frac{x^4}{12} - \frac{x^3}{3} - x + 1$$