

HW 10  
MATH 4567 Applied Fourier Analysis  
Spring 2019  
University of Minnesota, Twin Cities

Nasser M. Abbasi

November 2, 2019

Compiled on November 2, 2019 at 9:52pm [public]

# Contents

<b>1</b>	<b>Section 69, Problem 1</b>	<b>2</b>
1.1	Part (a) . . . . .	2
1.2	Part (b) . . . . .	3
<b>2</b>	<b>Section 72, Problem 3</b>	<b>5</b>
<b>3</b>	<b>Section 72, Problem 6</b>	<b>7</b>
<b>4</b>	<b>Section 72, Problem 9</b>	<b>9</b>

# 1 Section 69, Problem 1

1. (a) After writing the differential equation in the regular Sturm-Liouville problem

$$[xX'(x)]' + \frac{\lambda}{x}X(x) = 0 \quad (1 < x < b),$$

$$X(1) = 0, \quad X(b) = 0$$

in Cauchy-Euler form (see Problem 1, Sec. 44), use the substitution  $x = \exp s$  to transform the problem into one consisting of the differential equation

$$\frac{d^2X}{ds^2} + \lambda X = 0 \quad (0 < s < \ln b)$$

and the boundary conditions

$$X = 0 \quad \text{when } s = 0 \quad \text{and} \quad X = 0 \quad \text{when } s = \ln b.$$

Then, by simply referring to the solutions of the Sturm-Liouville problem (4) in Sec. 35, show that the eigenvalues and eigenfunctions of the original problem here are

$$\lambda_n = \alpha_n^2, \quad X_n(x) = \sin(\alpha_n \ln x) \quad (n = 1, 2, \dots),$$

where  $\alpha_n = n\pi / \ln b$ .

(b) By making the substitution

$$s = \pi \frac{\ln x}{\ln b} \quad \text{p. 13}$$

in the integral involved and then referring to Problem 9, Sec. 5, give a direct verification that the set of eigenfunctions  $X_n(x)$  obtained in part (a) is orthogonal on the interval  $1 < x < b$  with weight function  $p(x) = 1/x$ , as ensured by Theorem 1 in Sec. 69.

Figure 1: Problem statement

## Solution

### 1.1 Part (a)

$$\begin{aligned} X'(x) + xX''(x) + \frac{\lambda}{x}X(x) &= 0 \\ x^2X''(x) + xX'(x) + \lambda X(x) &= 0 \end{aligned} \quad (1)$$

To transform the above to  $X''(s) + \lambda X(s) = 0$ , let  $x = e^s$ . Therefore  $\frac{dx}{ds} = e^s$  or  $\frac{ds}{dx} = e^{-s}$ .  
Now

$$\begin{aligned} \frac{dX}{dx} &= \frac{dX}{ds} \frac{ds}{dx} \\ &= \frac{dX}{ds} e^{-s} \end{aligned} \quad (2)$$

And

$$\begin{aligned} \frac{d^2X}{dx^2} &= \frac{d}{dx} \left( \frac{dX}{dx} \right) \\ &= \frac{d}{dx} \left( \frac{dX}{ds} e^{-s} \right) \end{aligned}$$

Hence, by product rule

$$\begin{aligned}
 \frac{d^2 X}{dx^2} &= \frac{d^2 X}{ds^2} \frac{ds}{dx} e^{-s} + \frac{dX}{ds} \frac{d}{dx} (e^{-s}) \\
 &= \frac{d^2 X}{ds^2} e^{-s} e^{-s} + \frac{dX}{ds} \frac{d}{ds} (e^{-s}) \frac{ds}{dx} \\
 &= \frac{d^2 X}{ds^2} e^{-2s} + \frac{dX}{ds} (-e^{-s}) (e^{-s}) \\
 &= e^{-2s} \frac{d^2 X}{ds^2} - e^{-2s} \frac{dX}{ds}
 \end{aligned} \tag{3}$$

Substituting (2,3) back into (1) gives

$$x^2 \left( e^{-2s} \frac{d^2 X}{ds^2} - e^{-2s} \frac{dX}{ds} \right) + x \left( \frac{dX}{ds} e^{-s} \right) + \lambda X = 0$$

But  $x = e^s$  and the above simplifies to

$$\begin{aligned}
 e^{2s} \left( e^{-2s} \frac{d^2 X}{ds^2} - e^{-2s} \frac{dX}{ds} \right) + e^s \left( \frac{dX}{ds} e^{-s} \right) + \lambda X &= 0 \\
 \frac{d^2 X}{ds^2} - \frac{dX}{ds} + \frac{dX}{ds} + \lambda X &= 0 \\
 \frac{d^2 X(s)}{ds^2} + \lambda X(s) &= 0
 \end{aligned}$$

When  $X(1) = 0$ , which means when  $x = 1$ , and since  $x = e^s$ , then when  $s = 0$ . Hence  $X(1) = 0$  becomes  $X(0) = 0$ . And when  $x = b$ , then  $s = \ln(b)$ . Hence the second condition becomes  $X(\ln(b)) = 0$ . Therefore the new B.C. are

$$\begin{aligned}
 X(0) &= 0 \\
 X(\ln(b)) &= 0
 \end{aligned}$$

By referring to problem (4) in section 35 we see that the eigenvalues are

$$\lambda_n = \left( \frac{n\pi}{c} \right)^2$$

Where here  $c = \ln(b)$ . Hence

$$\begin{aligned}
 \lambda_n &= \left( \frac{n\pi}{\ln(b)} \right)^2 \quad n = 1, 2, 3, \dots \\
 &= \alpha_n^2
 \end{aligned}$$

Where  $\alpha_n = \frac{n\pi}{\ln(b)}$ . And the eigenfunctions are, per section 35

$$X_n(s) = \sin(\alpha_n s)$$

In terms of  $x$ , the eigenfunctions become

$$X_n(s) = \sin(\alpha_n \ln x)$$

## 1.2 Part (b)

$$\langle X_n(x), X_m(x) \rangle = \int_1^b \sin(\alpha_n \ln x) \sin(\alpha_m \ln x) p(x) dx$$

But from  $(xX'(x))' + \frac{\lambda}{x}X(x) = 0$  and comparing this to  $(rX')' + (\lambda p + q)X = 0$ , we see that  $r(x) = x$  and  $q = 0$  and  $p = \frac{1}{x}$ . Hence the above integral becomes

$$\langle X_n(x), X_m(x) \rangle = \int_1^b \frac{1}{x} \sin(\alpha_n \ln x) \sin(\alpha_m \ln x) dx$$

Let  $s = \frac{\ln x}{\ln b} \pi$ . Then  $\frac{ds}{dx} = \frac{1}{x} \frac{\pi}{\ln b}$  or  $dx = \frac{x}{\pi} \ln(b) ds$ . When  $x = 1$  then  $s = 0$  and when  $x = b$  then  $s = \pi$ . Hence the above integral becomes

$$\begin{aligned} \langle X_n(x), X_m(x) \rangle &= \int_{s=0}^{s=\pi} \frac{1}{x} \sin\left(\alpha_n \frac{s \ln b}{\pi}\right) \sin\left(\alpha_m \frac{s \ln b}{\pi}\right) \left(\frac{x}{\pi} \ln(b) ds\right) \\ &= \frac{1}{\pi} \ln(b) \int_0^\pi \sin\left(\alpha_n \frac{s \ln b}{\pi}\right) \sin\left(\alpha_m \frac{s \ln b}{\pi}\right) ds \end{aligned}$$

But  $\alpha_n = \frac{n\pi}{\ln(b)}$  and  $\alpha_m = \frac{m\pi}{\ln(b)}$ , therefore the above becomes

$$\begin{aligned} \langle X_n(x), X_m(x) \rangle &= \frac{1}{\pi} \ln(b) \int_0^\pi \sin\left(\frac{n\pi}{\ln(b)} \frac{s \ln b}{\pi}\right) \sin\left(\frac{m\pi}{\ln(b)} \frac{s \ln b}{\pi}\right) ds \\ &= \frac{1}{\pi} \ln(b) \int_0^\pi \sin(ns) \sin(ms) ds \end{aligned} \tag{1}$$

Referring to Problem 9., section 5 which says that


$$\int_0^\pi \sin(nx) \sin(mx) dx = \begin{cases} 0 & n \neq m \\ \frac{\pi}{2} & n = 0 \end{cases}$$

Applying this to (1) shows that

$$\langle X_n(x), X_m(x) \rangle = \begin{cases} 0 & n \neq m \\ \frac{\pi}{2} & n = 0 \end{cases}$$

Hence  $X_n(x)$  and  $X_m(x)$  are orthogonal, since this is the definition of orthogonality.

## 2 Section 72, Problem 3



3.  $X'' + \lambda X = 0, \quad X'(0) = 0, \quad X(c) = 0.$   
 Answer:  $\lambda_n = \alpha_n^2, \quad \phi_n(x) = \sqrt{\frac{2}{c}} \cos \alpha_n x \quad (n = 1, 2, \dots); \quad \alpha_n = \frac{(2n-1)\pi}{2c}.$

Figure 2: Problem statement

### Solution

Solve for eigenvalues and normalized eigenfunctions.

$$\begin{aligned} X'' + \lambda X &= 0 \\ X'(0) &= 0 \\ X(c) &= 0 \end{aligned}$$

Writing the boundary conditions in SL standard form

$$\begin{aligned} a_1 X(0) + a_2 X'(0) &= 0 \\ b_1 X(c) + b_2 X'(c) &= 0 \end{aligned}$$

Shows that  $a_1 = 0, a_2 = 1$  and  $b_1 = 1, b_2 = 0$ . Therefore  $a_1 a_2 = 0$  and  $b_1 b_2 = 0$ . But we know that if  $a_1 a_2 \geq 0$  and  $b_1 b_2 \geq 0$ , then  $\lambda > 0$  is only possible eigenvalues. Let  $\lambda_n = \alpha_n^2, \alpha > 0$ . Hence the solution to the ODE is

$$\begin{aligned} X_n(x) &= A \cos(\alpha_n x) + B \sin(\alpha_n x) \\ X'_n(x) &= -A\alpha_n \sin(\alpha_n x) + B\alpha_n \cos(\alpha_n x) \end{aligned}$$

First B.C  $X'(0) = 0$  gives

$$0 = B\alpha_n$$

Which implies  $B = 0$ . Hence the solution now becomes  $X_n(x) = A \cos(\alpha_n x)$ . For the second BC

$$\begin{aligned} 0 &= A \cos(\alpha_n c) \\ 0 &= \cos(\alpha_n c) \end{aligned}$$

Which implies

$$\begin{aligned} \alpha_n c &= \frac{\pi}{2}, 3\frac{\pi}{2}, 5\frac{\pi}{2}, \dots \\ &= (2n-1)\frac{\pi}{2} \quad n = 1, 2, 3, \dots \end{aligned}$$

Hence

$$\alpha_n = \frac{(2n-1)\pi}{c} \quad n = 1, 2, 3, \dots$$

And the corresponding eigenfunctions are

$$\begin{aligned} X_n(x) &= \cos(\alpha_n x) \\ &= \cos\left(\frac{(2n-1)\pi}{c} x\right) \end{aligned}$$

To find the normalized  $X_n(x)$  which we call it  $\phi_n(x)$ , then by definition

$$\phi_n(x) = \frac{X_n(x)}{\|X_n(x)\|}$$

But

$$\|X_n(x)\|^2 = \int_0^c p(x) X_n^2(x) dx$$

Comparing the ODE  $X'' + \lambda X = 0$  to  $(rX')' + (\lambda p + q)X = 0$ , we see that  $r(x) = 1$  and

$q = 0$  and  $p = 1$ . Hence the above becomes

$$\begin{aligned}\|X_n(x)\|^2 &= \int_0^c \cos^2(\alpha_n x) dx \\ &= \frac{c}{2}\end{aligned}$$

Therefore  $\|X_n(x)\| = \sqrt{\frac{c}{2}}$  which shows that

$$\begin{aligned}\phi_n(x) &= \frac{X_n(x)}{\sqrt{\frac{c}{2}}} \\ &= \sqrt{\frac{2}{c}} \cos(\alpha_n x)\end{aligned}$$

where

$$\alpha_n = \frac{(2n-1)\pi}{c} \quad n = 1, 2, 3, \dots$$

Which is what required to show.

### 3 Section 72, Problem 6




Figure 3: Problem statement

#### Solution

$$X_n(x) = \sin(\alpha_n \ln x)$$

$$\alpha_n = \frac{n\pi}{\ln b} \quad n = 1, 2, 3, \dots$$

The normalized eigenfunction is given by

$$\phi_n(x) = \frac{X_n(x)}{\|X_n(x)\|}$$

But

$$\|X_n(x)\|^2 = \int_1^b p(x) X_n^2(x) dx$$

Comparing the ODE  $(xX') + \frac{\lambda}{x}X = 0$  to  $(rX') + (\lambda p + q)X = 0$ , we see that  $r(x) = x$  and  $q = 0$  and  $p = \frac{1}{x}$ . Hence the above becomes

$$\|X_n(x)\|^2 = \int_1^b \frac{1}{x} \sin^2(\alpha_n \ln x) dx$$

Let  $s = \frac{\ln x}{\ln b} \pi$ . Then  $\frac{ds}{dx} = \frac{1}{x \ln b} \pi$  or  $dx = \frac{x}{\pi} \ln(b) ds$ . When  $x = 1$  then  $s = 0$  and when  $x = b$  then  $s = \pi$ . Hence the above integral becomes

$$\begin{aligned} \|X_n(x)\|^2 &= \int_{s=0}^{s=\pi} \frac{1}{x} \sin^2\left(\alpha_n \frac{s \ln b}{\pi}\right) \left(\frac{x}{\pi} \ln(b) ds\right) \\ &= \frac{1}{\pi} \ln(b) \int_0^\pi \sin^2\left(\alpha_n \frac{s \ln b}{\pi}\right) ds \end{aligned}$$



But  $\alpha_n = \frac{n\pi}{\ln(b)}$  therefore the above becomes

$$\begin{aligned}
 \|X_n(x)\|^2 &= \frac{1}{\pi} \ln(b) \int_0^\pi \sin^2\left(\frac{n\pi}{\ln(b)} \frac{s \ln b}{\pi}\right) ds \\
 &= \frac{1}{\pi} \ln(b) \int_0^\pi \sin^2(ns) ds \\
 &= \frac{1}{\pi} \ln(b) \int_0^\pi \frac{1}{2} - \frac{1}{2} \cos(2ns) ds \\
 &= \frac{1}{\pi} \ln(b) \left( \frac{\pi}{2} - \frac{1}{2} \sin\left(\frac{2ns}{2n}\right)_0^\pi \right) \\
 &= \frac{1}{\pi} \ln(b) \left( \frac{\pi}{2} - \frac{1}{2} \sin(s)_0^\pi \right) \\
 &= \frac{1}{2} \ln(b)
 \end{aligned}$$

Hence

$$\begin{aligned}
 \phi_n(x) &= \frac{\sin(\alpha_n \ln x)}{\sqrt{\frac{1}{2} \ln(b)}} \\
 &= \sqrt{\frac{2}{\ln(b)}} \sin(\alpha_n \ln x)
 \end{aligned}$$

Which is what required to show.

## 4 Section 72, Problem 9




Figure 4: Problem description

### solution

From problem section 69 problem 1, we know that  $(xX'(x))' + \frac{\lambda}{x}X(x) = 0$  can be transformed to  $X''(s) + \lambda X(s) = 0$  using  $x = e^s$ . With boundary conditions in  $s$  found as follows. When  $x = 1$  then  $s = 0$  and when  $x = b$  then  $s = \ln b$ . Hence we obtain the SL problem

$$\begin{aligned} X''(s) + \lambda X(s) &= 0 \\ X'(0) &= 0 \\ X(\ln b) &= 0 \end{aligned} \tag{1}$$

But problem 3 is

$$\begin{aligned} X'' + \lambda X &= 0 \\ X'(0) &= 0 \\ X(c) &= 0 \end{aligned} \tag{2}$$

And it had the solution

$$\phi_n(x) = \sqrt{\frac{2}{c}} \cos(\alpha_n x)$$

where

$$\alpha_n = \frac{(2n-1)\pi}{c} \quad n = 1, 2, 3, \dots$$

By comparing (2) and (1) we see it is the same problem, except  $c \rightarrow \ln b$ . Hence the solution to (2) is the same as the solution in (1) but with  $c$  replaced by  $\ln b$ . Hence the solution is

$$\begin{aligned} \phi_n(s) &= \sqrt{\frac{2}{\ln b}} \cos(\alpha_n s) \\ \alpha_n &= \frac{(2n-1)\pi}{\ln b} \quad n = 1, 2, 3, \dots \end{aligned}$$

But  $s = \ln x$ , hence the above becomes

$$\begin{aligned} \phi_n(x) &= \sqrt{\frac{2}{\ln b}} \cos(\alpha_n \ln x) \\ \alpha_n &= \frac{(2n-1)\pi}{\ln b} \quad n = 1, 2, 3, \dots \end{aligned}$$

Which is what required to show.