


quizz 8, ME 240 Dynamics, Fall 2017

Nasser M. Abbasi

December 30, 2019

0.1 Problem 1



The particle A, which has mass 3 kg, is moving in the coordinate system shown above. A is currently at position $X=1.5$, $Y=5.8$, $Z=-1.2$ m relative to the origin O, and is moving with velocity $\vec{v}=4.1\hat{i}+4.4\hat{j}+5.3\hat{k}$ m/s.

What is the magnitude of the angular momentum of particle A about the origin O ($\text{kg}\cdot\text{m}^2/\text{s}$)?

Report your answer to the nearest whole number.

Let \vec{h}_O be the angular momentum of A w.r.t to O. Therefore apply the definition


$$\begin{aligned}\vec{h}_O &= \vec{r}_{A/O} \times m\vec{v}_A \\ &= 3 \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1.5 & 5.8 & -1.2 \\ 4.1 & 4.4 & 5.3 \end{vmatrix} \\ &= 3(36.02\hat{i} - 12.87\hat{j} - 17.18\hat{k}) \\ &= 108.06\hat{i} - 38.61\hat{j} - 51.54\hat{k}\end{aligned}$$

Hence

$$\begin{aligned}|\vec{h}_O| &= \sqrt{108^2 + 38.61^2 + 51.54^2} \\ &= 125.794 \text{ kg}\cdot\text{m}^2/\text{sec}\end{aligned}$$

0.2 Problem 2

Question 2 2 pts



Particle A with mass 3.6 kg is attached to the origin O by a string of viscoelastic putty (silly putty) that slowly stretches. Initially A travels at speed $V=9.87$ m/s and at radius $r = r_0$ m.

What is the speed of A (m/s) when the radius is $r = 2.29r_0$

Report your answer to two decimal places.

By conservation of angular momentum


$$\begin{aligned} r_0 m v_A &= 2.29 r_0 m v_2 \\ v_2 &= \frac{v_A}{2.29} \\ &= \frac{9.87}{2.29} \\ &= 4.31 \text{ m/s} \end{aligned}$$

0.3 Problem 3



Particles A (mass 9.8 kg) and B (mass 8.3 kg) are on a massless rod with length $L=5.0$ meters which spins about the origin O. There is a linear motor at the origin that can move the bar back and forth relative to the origin, without applying a moment to the system. Initially particle B is at the origin O and particle A is traveling with a speed of $V_A=3.70$ m/s (shown above).

The linear motor at O slowly moves the rod, so that A moves toward O and B moves away from O. There are no other forces or moments.



When the rod is centered at O (as seen above), what is the velocity of B (m/s)?

Report your answer to two decimal places.

Angular momentum initially

$$\begin{aligned}
 \vec{h}_1 &= \vec{r}_{A/O} \times m_A \vec{v}_A \\
 &= 9.8 \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5 & 0 & 0 \\ 0 & -3.7 & 0 \end{vmatrix} \\
 &= -181.3 \hat{k}
 \end{aligned} \tag{1}$$

In new state, we first note that $|\vec{v}_{A2}| = |\vec{v}_B|$ since both are the same radius from origin. This means $|v_{Ay}| = |v_{By}|$ since they move only in y direction. Then

$$\begin{aligned}
 \vec{h}_2 &= \vec{r}_{A/O} \times m_A \vec{v}_{A2} + \vec{r}_{B/O} \times m_B \vec{v}_B \\
 &= 9.8 \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2.5 & 0 & 0 \\ 0 & -v_{Ay} & 0 \end{vmatrix} + 8.3 \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -2.5 & 0 & 0 \\ 0 & v_{By} & 0 \end{vmatrix} \\
 &= 9.8 \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2.5 & 0 & 0 \\ 0 & -v_{By} & 0 \end{vmatrix} + 8.3 \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -2.5 & 0 & 0 \\ 0 & v_{By} & 0 \end{vmatrix} \\
 &= -46.5 v_{By} \hat{k}
 \end{aligned} \tag{2}$$

Since (1) and (2) are equal (conservation of angular momentum) then

$$v_{By} = \frac{-181.3}{-46.5} = 3.899 \text{ m/s}$$