

HW 1, Math 601
Fall 2018
University Of Wisconsin, Milwaukee

Nasser M. Abbasi

March 27, 2019 Compiled on March 27, 2019 at 3:44am

Contents

1 Problem set	1
2 key solution	4

1 Problem set

PROBLEM SET 13.1

1. (Powers of i) Show that $i^2 = -1$, $i^3 = -i$, $i^4 = 1$, $i^5 = i$, \dots and $1/i = -i$, $1/i^2 = -1$, $1/i^3 = i$, \dots .
2. (Rotation) Multiplication by i is geometrically a counterclockwise rotation through $\pi/2$ (90°). Verify this by graphing z and iz and the angle of rotation for $z = 2 + 2i$, $z = -1 - 5i$, $z = 4 - 3i$.
3. (Division) Verify the calculation in (7).
4. (Multiplication) If the product of two complex numbers is zero, show that at least one factor must be zero.
5. Show that $z = x + iy$ is pure imaginary if and only if $\bar{z} = -z$.
6. (Laws for conjugates) Verify (9) for $z_1 = 24 + 10i$, $z_2 = 4 + 6i$.

7-15 COMPLEX ARITHMETIC

Let $z_1 = 2 + 3i$ and $z_2 = 4 - 5i$. Showing the details of your work, find (in the form $x + iy$):

7. $(5z_1 + 3z_2)^2$ 8. $\bar{z}_1\bar{z}_2$
 9. $\operatorname{Re}(1/z_1^2)$ 10. $\operatorname{Re}(z_2^2)$, $(\operatorname{Re} z_2)^2$
 11. z_2/z_1 12. \bar{z}_1/\bar{z}_2 , (z_1/z_2)

13. $(4z_1 - z_2)^2$ 14. \bar{z}_1/z_1 , z_1/\bar{z}_1
 15. $(z_1 + z_2)/(z_1 - z_2)$

16-19 Let $z = x + iy$. Find:

16. $\operatorname{Im} z^3$, $(\operatorname{Im} z)^3$
 17. $\operatorname{Re}(1/\bar{z})$
 18. $\operatorname{Im} [(1 + i)^8 z^2]$
 19. $\operatorname{Re}(1/\bar{z}^2)$

20. (Laws of addition and multiplication) Derive the following laws for complex numbers from the corresponding laws for real numbers.

$$\begin{aligned} z_1 + z_2 &= z_2 + z_1, \quad z_1 z_2 = z_2 z_1 \quad (\text{Commutative laws}) \\ (z_1 + z_2) + z_3 &= z_1 + (z_2 + z_3), \quad (z_1 z_2) z_3 = z_1 (z_2 z_3) \quad (\text{Associative laws}) \\ z_1(z_2 + z_3) &= z_1 z_2 + z_1 z_3 \quad (\text{Distributive law}) \\ 0 + z &= z + 0 = z, \quad z \cdot 1 = z. \\ z + (-z) &= (-z) + z = 0, \quad z \cdot 0 = 0. \end{aligned}$$

Problem Set 13.2

7. $\frac{-6 + 5i}{3i}$

8. $\frac{2 + 3i}{5 + 4i}$

9-15 PRINCIPAL ARGUMENT

Determine the principal value of the argument.

9. $-1 - i$

10. $-20 + i; -20 - i$

11. $4 \pm 3i$

12. $-\pi^2$

13. $7 \pm 7i$

14. $(1 + i)^{12}$

15. $(9 + 9i)^3$

16-20 CONVERSION TO $x + iy$

Represent in the form $x + iy$ and graph it in the complex plane.

16. $\cos \frac{1}{2}\pi + i \sin(\pm \frac{1}{2}\pi)$

17. $3(\cos 0.2 + i \sin 0.2)$

18. $4(\cos \frac{1}{3}\pi \pm i \sin \frac{1}{3}\pi)$

19. $\cos(-1) + i \sin(-1)$

20. $12(\cos \frac{3}{2}\pi + i \sin \frac{3}{2}\pi)$

21-25 ROOTS

Find and graph all roots in the complex plane.

21. $\sqrt{-i}$

22. $\sqrt[8]{1}$

23. $\sqrt[4]{-1}$

24. $\sqrt[3]{3 + 4i}$

25. $\sqrt[5]{-1}$

26. TEAM PROJECT. Square Root. (a) Show that $w = \sqrt{z}$ has the values

$$w_1 = \sqrt{r} \left[\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} \right],$$

$$(18) w_2 = \sqrt{r} \left[\cos \left(\frac{\theta}{2} + \pi \right) + i \sin \left(\frac{\theta}{2} + \pi \right) \right]$$

$$= -w_1.$$

(b) Obtain from (18) the often more practical formula

$$(19) \sqrt{z} = \pm \left[\sqrt{\frac{1}{2}(|z| + x)} + (\text{sign } y)i\sqrt{\frac{1}{2}(|z| + x)} \right]$$

where $\text{sign } y = 1$ if $y \geq 0$, $\text{sign } y = -1$ if $y < 0$, and all square roots of positive numbers are taken with positive sign. Hint: Use (10) in App. A3.1 with $x = \theta/2$.

(c) Find the square roots of $4i$, $16 - 30i$, and $9 + 8\sqrt{7}i$ by both (18) and (19) and comment on the work involved.

(d) Do some further examples of your own and apply a method of checking your results.

27-30 EQUATIONS

Solve and graph all solutions, showing the details:

27. $z^2 - (8 - 5i)z + 40 - 20i = 0$ (Use (19).)

28. $z^4 + (5 - 14i)z^2 - (24 + 10i) = 0$

29. $8z^2 - (36 - 6i)z + 42 - 11i = 0$

30. $z^4 + 16 = 0$. Then use the solutions to factor $z^4 + 16$ into quadratic factors with *real* coefficients.

31. CAS PROJECT. Roots of Unity and Their Graphs. Write a program for calculating these roots and for graphing them as points on the unit circle. Apply the program to $z^n = 1$ with $n = 2, 3, \dots, 10$. Then extend the program to one for arbitrary roots, using an idea near the end of the text, and apply the program to examples of your choice.

32-35 INEQUALITIES AND AN EQUATION

Verify or prove as indicated.

32. (Re and Im) Prove $|\text{Re } z| \leq |z|$, $|\text{Im } z| \leq |z|$.

33. (Parallelogram equality) Prove

$$|z_1 + z_2|^2 + |z_1 - z_2|^2 = 2(|z_1|^2 + |z_2|^2).$$

Explain the name.

34. (Triangle inequality) Verify (6) for $z_1 = 4 + 7i$, $z_2 = 5 + 2i$.

35. (Triangle inequality) Prove (6).

PROBLEM SET 13.3**1-10 CURVES AND REGIONS OF PRACTICAL INTEREST**

Find and sketch or graph the sets in the complex plane given by

1. $|z - 3 - 2i| = \frac{4}{3}$

2. $1 \leq |z - 1 + 4i| \leq 5$

3. $0 < |z - 1| < 1$ 4. $-\pi < \text{Re } z < \pi$

5. $\text{Im } z^2 = 2$

6. $\text{Re } z > -1$

7. $|z + 1| = |z - 1|$

8. $|\text{Arg } z| \leq \frac{1}{4}\pi$

9. $\text{Re } z \leq \text{Im } z$

10. $\text{Re}(1/z) < 1$

PROBLEM SET 13.5

1. Using the Cauchy-Riemann equations, show that e^z is entire.

2-8 Values of e^z . Compute e^z in the form $u + iv$ and $|e^z|$, where z equals:

(2) $3 + \pi i$

(4) $\sqrt{2} - \frac{1}{2}\pi i$

(6) $(1 + i)\pi$

(8) $9\pi i/2$

3. $1 + 2i$

5. $7\pi i/2$

7. $0.8 - 5i$

9-12 Real and Imaginary Parts. Find Re and Im of:

9. e^{-2z}

11. e^{z^2}

10. e^{z^3}

12. $e^{1/z}$

13-17 Polar Form. Write in polar form:

13. \sqrt{i}

15. $\sqrt[n]{z}$

17. -9

14. $1 + i$

16. $3 + 4i$

18-21 Equations. Find all solutions and graph some of them in the complex plane.

18. $e^{3z} = 4$

20. $e^z = 0$

19. $e^z = -2$

21. $e^z = 4 - 3i$

22. TEAM PROJECT. Further Properties of the Exponential Function. (a) Analyticity. Show that e^z is entire. What about $e^{1/z}$? $e^{\bar{z}}$? $e^x(\cos ky + i \sin ky)$? (Use the Cauchy-Riemann equations.)

(b) Special values. Find all z such that (i) e^z is real, (ii) $|e^{-z}| < 1$, (iii) $e^z = \overline{e^z}$.

(c) Harmonic function. Show that

$u = e^{xy} \cos(x^2/2 - y^2/2)$ is harmonic and find a conjugate.

(d) Uniqueness. It is interesting that $f(z) = e^z$ is uniquely determined by the two properties $f(x + i0) = e^x$ and $f'(z) = f(z)$, where f is assumed to be entire. Prove this using the Cauchy-Riemann equations.

PROBLEM SET 13.6

1. Prove that $\cos z$, $\sin z$, $\cosh z$, $\sinh z$ are entire functions.

2. Verify by differentiation that $\operatorname{Re} \cos z$ and $\operatorname{Im} \sin z$ are harmonic.

3-6 FORMULAS FOR HYPERBOLIC FUNCTIONS

Show that

3. $\cosh z = \cosh x \cos y + i \sinh x \sin y$
 $\sinh z = \sinh x \cos y + i \cosh x \sin y$

4. $\cosh(z_1 + z_2) = \cosh z_1 \cosh z_2 + \sinh z_1 \sinh z_2$
 $\sinh(z_1 + z_2) = \sinh z_1 \cosh z_2 + \cosh z_1 \sinh z_2$

5. $\cosh^2 z - \sinh^2 z = 1$

6. $\cosh^2 z + \sinh^2 z = \cosh 2z$

7-15 Function Values. Compute (in the form $u + iv$)

7. $\cos(1 + i)$

8. $\sin(1 + i)$

9. $\sin 5i, \cos 5i$

10. $\cos 3\pi i$

11. $\cosh(-2 + 3i), \cos(-3 - 2i)$

12. $-i \sinh(-\pi + 2i), \sin(2 + \pi i)$

13. $\cosh(2n + 1)\pi i, n = 1, 2, \dots$

14. $\sinh(4 - 3i)$

15. $\cosh(4 - 6\pi i)$

16. (Real and imaginary parts) Show that

$$\operatorname{Re} \tan z = \frac{\sin x \cos x}{\cos^2 x + \sinh^2 y},$$

$$\operatorname{Im} \tan z = \frac{\sinh y \cosh y}{\cos^2 x + \sinh^2 y}.$$

17-21 Equations. Find all solutions of the following equations.

17. $\cosh z = 0$

18. $\sin z = 100$

19. $\cos z = 2i$

20. $\cosh z = -1$

21. $\sinh z = 0$

22. Find all z for which (a) $\cos z$, (b) $\sin z$ has real values.

23-25 Equations and Inequalities. Using the definitions, prove:

23. $\cos z$ is even, $\cos(-z) = \cos z$, and $\sin z$ is odd, $\sin(-z) = -\sin z$.

24. $|\sinh y| \leq |\cos z| \leq \cosh y, |\sinh y| \leq |\sin z| \leq \cosh y$. Conclude that the complex cosine and sine are not bounded in the whole complex plane.

25. $\sin z_1 \cos z_2 = \frac{1}{2}[\sin(z_1 + z_2) + \sin(z_1 - z_2)]$

PROBLEM SET 13.7

1-9 Principal Value $\operatorname{Ln} z$. Find $\operatorname{Ln} z$ when z equals:

1. -10

2. $2 + 2i$

3. $2 - 2i$

4. $-5 \pm 0.1i$

5. $-3 - 4i$

6. -100

7. $0.6 + 0.8i$

8. $-ei$

9. $1 - i$

10-16 All Values of $\operatorname{Ln} z$. Find all values and graph some of them in the complex plane.

10. $\ln 1$

11. $\ln(-1)$

12. $\ln e$

13. $\ln(-6)$

14. $\ln(4 + 3i)$

15. $\ln(-e^{-i})$

16. $\ln(e^{3i})$

17. Show that the set of values of $\ln(i^2)$ differs from the set of values of $2 \ln i$.

18-21 Equations. Solve for z :

18. $\ln z = (2 - \frac{1}{2}i)\pi$

19. $\ln z = 0.3 + 0.7i$

20. $\ln z = e - \pi i$

21. $\ln z = 2 + \frac{1}{4}\pi i$

22–28 General Powers. Showing the details of your work, find the principal value of:

22. i^{2i} , $(2i)^i$

24. $(1-i)^{1+i}$

26. $(-1)^{1-2i}$

28. $(3-4i)^{1/3}$

23. 4^{3+i}

25. $(1+i)^{1-i}$

27. $i^{1/2}$

29. How can you find the answer to Prob. 24 from the answer to Prob. 25?

30. TEAM PROJECT. Inverse Trigonometric and Hyperbolic Functions. By definition, the inverse sine $w = \arcsin z$ is the relation such that $\sin w = z$. The inverse cosine $w = \arccos z$ is the relation such that $\cos w = z$. The inverse tangent, inverse cotangent, inverse hyperbolic sine, etc., are defined and denoted in a similar fashion. (Note that all these relations are *multivalued*.) Using $\sin w = (e^{iw} - e^{-iw})/(2i)$ and similar representations of $\cos w$, etc., show that

(a) $\arccos z = -i \ln(z + \sqrt{z^2 - 1})$

(b) $\arcsin z = -i \ln(iz + \sqrt{1 - z^2})$

(c) $\text{arccosh } z = \ln(z + \sqrt{z^2 - 1})$

(d) $\text{arcsinh } z = \ln(z + \sqrt{z^2 + 1})$

(e) $\arctan z = \frac{i}{2} \ln \frac{i+z}{i-z}$

(f) $\text{arctanh } z = \frac{1}{2} \ln \frac{1+z}{1-z}$

(g) Show that $w = \arcsin z$ is infinitely many-valued, and if w_1 is one of these values, the others are of the form $w_1 \pm 2n\pi$ and $\pi - w_1 \pm 2n\pi$, $n = 0, 1, \dots$. (The *principal value* of $w = u + iv = \arcsin z$ is defined to be the value for which $-\pi/2 \leq u \leq \pi/2$ if $v \geq 0$ and $-\pi/2 < u < \pi/2$ if $v < 0$.)

2 key solution

Math 601

[13.1] By definition

[1]

the sequence

$\{i, -1, -i, 1\}$ repeats.

$$i^1 = i$$

$$i^2 = i \cdot i = -1$$

$$i^3 = i^2 \cdot i = -i$$

$$i^4 = i^2 \cdot i^2 = (-1)(-1) = 1$$

$$i^5 = i^4 \cdot i = i$$

$$i^6 = i^4 \cdot i^2 = -1$$

$$i^7 = i^4 \cdot i^3 = -i$$

$$i^8 = i^4 \cdot i^4 = 1$$

⋮

[7]

$$z_1 = 2+3i, z_2 = 4-5i$$

$$(5z_1 + 3z_2)^2 = (5(2+3i) + 3(4-5i))^2 =$$

$$= (10+15i+12-15i)^2 = 22^2 = 484$$

[10]

$$z_2 = 4-5i \quad (\operatorname{Re} z_2)^2 = 4^2 = 16$$

$$z_2^2 = (4-5i)(4-5i) \Rightarrow \operatorname{Re}(z_2^2) = -9$$

$$= 16-25-2 \cdot 4 \cdot 5i$$

$$= -9-40i$$

[12] $\frac{\bar{z}_1}{\bar{z}_2} = \frac{2-3i}{4+5i} = \frac{\bar{z}_1 z_2}{\bar{z}_2 z_2} = \frac{\bar{z}_1 z_2}{|z_2|^2} = \frac{-7-22i}{41} = -\frac{7}{41} - \frac{22}{41}i$

$$|z_2|^2 = 4^2 + 5^2 = 41$$

$$\bar{z}_1 z_2 = (2-3i)(4-5i) = 8-15-(10+12)i = -7-22i$$

We know that $\left(\frac{\bar{z}_1}{\bar{z}_2}\right) = \frac{\bar{z}_1}{\bar{z}_2} = -\frac{7}{41} - \frac{22}{41}i$

[13.2] 18 polar form of $z = \frac{2+3i}{5+4i} = r e^{i\theta} = r(\cos\theta + i\sin\theta)$

with $r = \sqrt{z \cdot \bar{z}}$

$\theta = \tan^{-1}\left(\frac{y}{x}\right)$

$(z = x+iy)$

$$z = \frac{2+3i}{5+4i} = \frac{(2+3i)(5-4i)}{(5+4i)(5-4i)} = \frac{10+12+(15-8)i}{25+16} = \frac{22}{41} + \frac{7}{41}i = x+iy$$

thus $r = \sqrt{\left(\frac{22}{41}\right)^2 + \left(\frac{7}{41}\right)^2}$

and $\theta = \tan^{-1} \frac{7/41}{22/41} = \tan^{-1}\left(\frac{7}{22}\right)$

III

$z = x+iy$ $z = 4+3i$ $-\pi < \operatorname{Arg} z \leq \pi$ and $\tan \operatorname{Arg} z = \frac{y}{x} = \frac{3}{4}$

$\bar{z} = x-iy$ $\bar{z} = 4-3i$

$\operatorname{Arg} z = \tan^{-1} \frac{3}{4} \approx 0.643$


$\tan^{-1} \frac{y}{x} = \operatorname{Arg} \bar{z} = -\operatorname{Arg} z = \tan^{-1} \frac{y}{x}$

(and \tan^{-1} is odd function)

[18] $z = 4 \cdot \left(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}\right) = \frac{4}{2} + i \frac{4\sqrt{3}}{2} = 2 + 2\sqrt{3}i$

$$\bar{z} = 4 \cdot \left(\cos \frac{\pi}{3} - i \sin \frac{\pi}{3}\right) = 2 - 2\sqrt{3}i$$

$\frac{\pi}{2}$ $\frac{\pi}{2}$




[22] $8\sqrt{1} = 8\sqrt{r} \left(\cos \frac{\theta + 2k\pi}{8} + i \sin \frac{\theta + 2k\pi}{8} \right) =$

$k=0, 1, 2, \dots, 7$

$r = 1 \cdot 1 = 1$

$\theta = \operatorname{Arg} 1 = 0$


$= \cos \frac{k\pi}{4} + i \sin \frac{k\pi}{4}$



$$\boxed{13.2} \quad \boxed{24} \quad \sqrt[3]{3+4i} = r^{\frac{1}{3}} \left(\cos \frac{\theta + 2k\pi}{3} + i \sin \frac{\theta + 2k\pi}{3} \right)$$

$$r = |3+4i| = \sqrt{9+16} = 5 \quad = \sqrt[3]{5} \left[\cos \left(\frac{\theta}{3} + \frac{2\pi k}{3} \right) + i \sin \left(\frac{\theta}{3} + \frac{2\pi k}{3} \right) \right]$$

$$\theta = \tan^{-1} \frac{4}{3} \approx 0.9273 \quad k=0, 1, 2$$



35 Show that $|z_1 + z_2| \leq |z_1| + |z_2|$
or equivalently

$$|z_1 + z_2|^2 \leq |z_1|^2 + 2|z_1||z_2| + |z_2|^2$$

thus we need to show:

$$\cancel{x_1^2 + 2x_1x_2 + x_2^2 + y_1^2 + 2y_1y_2 + y_2^2} \leq \\ \leq x_1^2 + y_1^2 + x_2^2 + y_2^2 + 2\sqrt{x_1^2 + y_1^2} \cdot \sqrt{x_2^2 + y_2^2}$$

or $x_1x_2 + y_1y_2 \leq \sqrt{x_1^2 + y_1^2} \sqrt{x_2^2 + y_2^2}$

(Note: RHS non-negative) to show this, square each sides

$$\cancel{x_1^2 x_2^2 + 2x_1 x_2 y_1 y_2 + y_1^2 y_2^2} \leq (x_1^2 + y_1^2)(x_2^2 + y_2^2)$$

$$= x_1^2 x_2^2 + x_1^2 y_2^2 + x_2^2 y_1^2 + y_1^2 y_2^2$$

$$|z_1| = \sqrt{x_1^2 + y_1^2}$$

$$|z_2| = \sqrt{x_2^2 + y_2^2}$$

$$z_1 + z_2 = x_1 + x_2 + i(y_1 + y_2)$$

$$|z_1 + z_2|^2 = \\ = (x_1 + x_2)^2 + (y_1 + y_2)^2 \\ = x_1^2 + 2x_1x_2 + x_2^2 \\ + y_1^2 + 2y_1y_2 + y_2^2$$

or

$$0 \leq x_1^2 y_2^2 + x_2^2 y_1^2 - 2x_1 x_2 y_1 y_2 =$$


$$= (x_1 y_2 - x_2 y_1)^2 \text{ which is true for any } x_1, x_2, y_1, y_2 \in \mathbb{R}.$$

13.3 2

$$1 \leq |z - (1-4i)| \leq 5 \quad \text{take } z_0 = 1-4i$$

$|z - z_0| = 1$: circle of radius 1
around z_0

$|z - z_0| = 5$ ——— radius 5




5 $\operatorname{Im} z^2 = 2$

$$z = r \cdot e^{i\theta} \quad z^2 = r^2 \cdot e^{i2\theta}$$



$$r^2 \cos 2\theta + i r^2 \sin 2\theta$$

$$\operatorname{Im} z^2 = r^2 \sin 2\theta = 2$$

$$\Rightarrow r^2 = 2 \csc 2\theta$$



$$r = \sqrt{2 \csc 2\theta}$$




8 $|\operatorname{Arg} z| \leq \frac{1}{4}\pi \Rightarrow -\frac{\pi}{4} \leq \operatorname{Arg} z \leq \frac{\pi}{4}$

$$z = r \cdot e^{\operatorname{Arg} z}$$

with

$$-\pi < \operatorname{Arg} z \leq \pi$$



13.5 (2) $e^{3+i\pi} = e^3 e^{i\pi} = e^3 (\underbrace{\cos \pi}_{-1} + i \underbrace{\sin \pi}_0) = -e^3$

12 $z = r \cdot e^{i\theta} \Rightarrow z^{-1} = r^{-1} e^{-i\theta} = \frac{1}{r} (\cos(-\theta) + i \sin(-\theta)) =$

$$r=|z|$$

$$\theta = \arg z$$

$$\text{if } z = x+iy$$

$$|z| = \sqrt{x^2+y^2}$$

$$\arg z = \tan^{-1} \frac{y}{x}$$

$$e^{\frac{1}{z}} = e^{\frac{1}{r} \cos \theta - \frac{1}{r} \sin \theta i} =$$

$$= e^{\frac{1}{r} \cos \theta} (\cos(-\frac{1}{r} \sin \theta) + i \sin(-\frac{1}{r} \sin \theta)) =$$

$$= e^{\frac{1}{r} \cos \theta} \cdot \cos(\frac{1}{r} \sin \theta)$$

$$+ i e^{\frac{1}{r} \cos \theta} \cos(\frac{1}{r} \sin \theta)$$

$$\text{Thus } \operatorname{Re} e^{\frac{1}{z}} = e^{\frac{1}{r} \cos \theta} \cdot \cos(\frac{1}{r} \sin \theta)$$

$$\operatorname{Im} e^{\frac{1}{z}} = e^{\frac{1}{r} \cos \theta} \cdot \cos(\frac{1}{r} \sin \theta)$$

13 \sqrt{i} :

$$i = e^{\frac{\pi}{2}i} = \cos \frac{\pi}{2} + i \sin \frac{\pi}{2}$$

$$|i|=1 \quad \text{thus} \quad \sqrt{i} = \sqrt{1} \left(\cos \frac{\frac{\pi}{2} + k \cdot 2\pi}{2} + i \sin \frac{\frac{\pi}{2} + k \cdot 2\pi}{2} \right)$$

$$\text{with } k=0, k=1$$


$$\begin{aligned} 1) & \text{ either } \sqrt{i} = \cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \\ & (k=0) \\ 2) & \text{ or } \sqrt{i} = \cos \frac{5\pi}{4} + i \sin \frac{5\pi}{4} \\ & (k=1) \end{aligned}$$

$$\text{i.e. } \sqrt{i} = \frac{\sqrt{2}}{2} + i \frac{\sqrt{2}}{2}$$

$$\text{or } \sqrt{i} = -\frac{\sqrt{2}}{2} - i \frac{\sqrt{2}}{2}$$

13.5 16 $z + 4i = r \cdot e^{i\theta} = 5 \cdot \left(\cos(\tan^{-1} \frac{4}{3}) + i \sin(\tan^{-1} \frac{4}{3}) \right)$

$$r = \sqrt{3^2 + 4^2} = 5 \quad / = 5 \left(\frac{3}{5} + i \frac{4}{5} \right) /$$



20 $e^z = e^{|z|} \cdot e^{i \operatorname{Arg} z} \Rightarrow |e^z| = |e^{|z|} \cdot e^{i \theta}| =$

$$r = |z|$$

$$\theta = \operatorname{Arg} z$$

$$|e^r| \cdot |\cos \theta + i \sin \theta| > 0 = 1$$

$$\text{thus } |e^{|z|}| = 0$$

so $e^{|z|} = 0$ has
no solution.

*HYP^O $(4-3i)$

21 $e^z = 3 - 4i \Rightarrow \ln e^z = z = \ln(3 - 4i) =$

$$= \ln|3 - 4i| + i \operatorname{Arg} z =$$

$$|3 - 4i| = 5$$

$$\operatorname{Arg} z = \tan^{-1} \frac{4}{3}$$

$$= \ln 5 - i \tan^{-1} \frac{4}{3}$$

13.6
18

Compute as $u+iv \rightarrow \sin(1+i)$

$$z = 1+i$$

$$iz = -1+i$$

$$-iz = 1-i$$

$$|iz| = |-iz| = |z| = \sqrt{2}$$

$$e^{iz} = e^{-1+i} = \frac{1}{e}(\cos 1 + i \sin 1)$$

$$e^{-iz} = e^{1-i} = e(\cos 1 - i \sin 1)$$

$$\sin z = \frac{1}{2i}(e^{iz} - e^{-iz})$$

$$\sin(1+i) = \frac{e^{iz} - e^{-iz}}{2i} =$$

$$= \frac{-i}{2} \left(\cos 1 \cdot \left(\frac{1}{e} - e \right) + i \sin 1 \cdot \left(\frac{1}{e} + e \right) \right)$$

$$= \underbrace{\frac{\sin 1}{2} \left(\frac{1}{e} + e \right)}_u - i \underbrace{\frac{\cos 1}{2} \left(\frac{1}{e} - e \right)}_v = u + iv$$

23 $\cos(-z) = \frac{1}{2}(e^{i(-z)} + e^{-i(-z)}) = \frac{1}{2}(e^{-iz} + e^{iz}) = \cos z$

$$\sin(-z) = \frac{1}{2i}(e^{i(-z)} - e^{-i(-z)}) = \frac{1}{2i}(e^{-iz} - e^{iz}) = -\frac{1}{2i}(e^{iz} - e^{-iz}) = -\sin z$$

thus $\cos(z)$ is even

$\sin(z)$ is odd

13.7 14 See 13.5 21

19 solve $\ln z = 0.3 + 0.7i \Rightarrow z = e^{\ln z} = e^{\frac{3}{10} + \frac{7}{10}i} = e^{\frac{3}{10}} \left(\cos \frac{7}{10} + i \sin \frac{7}{10} \right) = e^{0.3} \cos 0.7 + i e^{0.3} \sin 0.7$

$$\boxed{13.7} \quad \boxed{\frac{125}{29}} \quad (1+i)^{(1-i)} = e^{\ln(1+i)^{(1-i)}} = e^{(1-i)\ln(1+i)}$$

and $\boxed{\frac{29}{2}}$

$$\ln(1+i) = \ln\sqrt{2} + i\frac{\pi}{4} \quad \text{thus} \quad (1-i)(\ln\sqrt{2} + i\frac{\pi}{4}) =$$

$$|\ln(1+i)| = \sqrt{2}$$

$$\ln\sqrt{2} + i\left(\frac{\pi}{4} - \ln\sqrt{2}\right)$$

$$\operatorname{Arg}(1+i) = \tan^{-1} 1 = \frac{\pi}{4} \quad \text{So}$$

$$(1+i)^{1-i} = e^{\ln\sqrt{2} + i\frac{\pi}{4}} \cdot \cos\left(\frac{\pi}{4} - \ln\sqrt{2}\right)$$

$$+ i e^{\ln\sqrt{2} + i\frac{\pi}{4}} \cdot \sin\left(\frac{\pi}{4} - \ln\sqrt{2}\right)$$

consider

$$z^{\bar{z}} \quad \text{and} \quad \bar{z}^z \quad \bar{z}^z = \overline{z^{\bar{z}}} = \overline{e^{\bar{z} \cdot \ln z}} = e^{\bar{\bar{z}} \cdot \bar{\ln z}} = e^{\bar{z}} e^{\ln \bar{z}} =$$

① ②

(note) $e^{\bar{z}} = \overline{e^z} = \overline{e^{x(\cos y + i \sin y)}} =$

$z = x+iy \quad e^x(\cos y + i \sin y) = e^{\bar{z}}$

also $\overline{\ln z} = \overline{\ln|z| + i \operatorname{Arg} z}$

$$\begin{aligned} &= \ln|z| - i \operatorname{Arg} z \\ &= \ln|z| + i \operatorname{Arg} \bar{z} \\ &= \overline{\ln z} \end{aligned}$$

thus (as $\overline{i+i} = 1-i$)

taking $z = 1+i$

we have that

$$(1+i)^{1-i} = (1-i)^{1+i}$$