

Computer algebra independent integration tests

4-Trig-functions/4.6-Cosecant/4.6.1.2-d-cscⁿ-a+b-csc^m

Nasser M. Abbasi

May 23, 2020

Compiled on May 23, 2020 at 7:02am

Contents

1	Introduction	5
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Performance	9
1.4	list of integrals that has no closed form antiderivative	10
1.5	list of integrals solved by CAS but has no known antiderivative	10
1.6	list of integrals solved by CAS but failed verification	10
1.7	Timing	11
1.8	Verification	11
1.9	Important notes about some of the results	11
1.10	Design of the test system	13
2	detailed summary tables of results	15
2.1	List of integrals sorted by grade for each CAS	15
2.2	Detailed conclusion table per each integral for all CAS systems	17
2.3	Detailed conclusion table specific for Rubi results	29
3	Listing of integrals	33
3.1	$\int \frac{\csc^5(x)}{a+a \csc(x)} dx$	33
3.2	$\int \frac{\csc^4(x)}{a+a \csc(x)} dx$	38
3.3	$\int \frac{\csc^3(x)}{a+a \csc(x)} dx$	42

3.4	$\int \frac{\csc^2(x)}{a+a \csc(x)} dx$	46
3.5	$\int \frac{\csc(x)}{a+a \csc(x)} dx$	50
3.6	$\int \frac{1}{a+a \csc(c+dx)} dx$	53
3.7	$\int \frac{\sin(x)}{a+a \csc(x)} dx$	56
3.8	$\int \frac{\sin^2(x)}{a+a \csc(x)} dx$	60
3.9	$\int \frac{\sin^3(x)}{a+a \csc(x)} dx$	64
3.10	$\int \frac{\sin^4(x)}{a+a \csc(x)} dx$	68
3.11	$\int \frac{1}{(a+a \csc(c+dx))^2} dx$	72
3.12	$\int \frac{1}{(a+a \csc(c+dx))^3} dx$	76
3.13	$\int (a+a \csc(x))^{5/2} dx$	80
3.14	$\int (a+a \csc(x))^{3/2} dx$	85
3.15	$\int \sqrt{a+a \csc(x)} dx$	90
3.16	$\int \frac{1}{\sqrt{a+a \csc(x)}} dx$	94
3.17	$\int \frac{1}{(a+a \csc(x))^{3/2}} dx$	98
3.18	$\int \frac{1}{(a+a \csc(x))^{5/2}} dx$	104
3.19	$\int \sqrt{\csc(e+fx)} \sqrt{a+a \csc(e+fx)} dx$	111
3.20	$\int \sqrt{-\csc(e+fx)} \sqrt{a-a \csc(e+fx)} dx$	115
3.21	$\int \csc^{\frac{4}{3}}(c+dx) \sqrt{a+a \csc(c+dx)} dx$	119
3.22	$\int \sqrt[3]{\csc(c+dx)} \sqrt{a+a \csc(c+dx)} dx$	124
3.23	$\int \frac{\sqrt{a+a \csc(c+dx)}}{\csc^{\frac{2}{3}}(c+dx)} dx$	128
3.24	$\int \csc^{\frac{5}{3}}(c+dx) \sqrt{a+a \csc(c+dx)} dx$	133
3.25	$\int \csc^{\frac{2}{3}}(c+dx) \sqrt{a+a \csc(c+dx)} dx$	138
3.26	$\int \frac{\sqrt{a+a \csc(c+dx)}}{\sqrt[3]{\csc(c+dx)}} dx$	143
3.27	$\int \frac{\sqrt{a+a \csc(c+dx)}}{\csc^{\frac{4}{3}}(c+dx)} dx$	148
3.28	$\int \csc^n(c+dx) \sqrt{a+a \csc(c+dx)} dx$	153
3.29	$\int \csc^n(c+dx) \sqrt{a-a \csc(c+dx)} dx$	156
3.30	$\int \csc^3(e+fx)(a+a \csc(e+fx))^m dx$	160
3.31	$\int \csc^2(e+fx)(a+a \csc(e+fx))^m dx$	165
3.32	$\int \csc(e+fx)(a+a \csc(e+fx))^m dx$	169
3.33	$\int (a+a \csc(e+fx))^m dx$	173
3.34	$\int (a+a \csc(e+fx))^m \sin(e+fx) dx$	177
3.35	$\int (a+a \csc(e+fx))^m \sin^2(e+fx) dx$	181
3.36	$\int (a+b \csc(c+dx))^4 dx$	185

3.37	$\int (a + b \csc(c + dx))^3 dx$	190
3.38	$\int (a + b \csc(c + dx))^2 dx$	194
3.39	$\int \frac{\csc^5(x)}{a+b \csc(x)} dx$	198
3.40	$\int \frac{\csc^4(x)}{a+b \csc(x)} dx$	204
3.41	$\int \frac{\csc^3(x)}{a+b \csc(x)} dx$	209
3.42	$\int \frac{\csc^2(x)}{a+b \csc(x)} dx$	214
3.43	$\int \frac{\csc(x)}{a+b \csc(x)} dx$	219
3.44	$\int \frac{1}{a+b \csc(c+dx)} dx$	223
3.45	$\int \frac{\sin(x)}{a+b \csc(x)} dx$	227
3.46	$\int \frac{\sin^2(x)}{a+b \csc(x)} dx$	232
3.47	$\int \frac{\sin^3(x)}{a+b \csc(x)} dx$	237
3.48	$\int \frac{\sin^4(x)}{a+b \csc(x)} dx$	242
3.49	$\int \frac{1}{(a+b \csc(c+dx))^2} dx$	248
3.50	$\int \frac{1}{(a+b \csc(c+dx))^3} dx$	253
3.51	$\int \frac{1}{(a+b \csc(c+dx))^4} dx$	259
3.52	$\int \frac{1}{3+5 \csc(c+dx)} dx$	266
3.53	$\int \frac{1}{5+3 \csc(c+dx)} dx$	270
3.54	$\int \csc^3(e + fx)(a + b \csc(e + fx))^m dx$	274
3.55	$\int \csc^2(e + fx)(a + b \csc(e + fx))^m dx$	279
3.56	$\int \csc(e + fx)(a + b \csc(e + fx))^m dx$	283
3.57	$\int (a + b \csc(e + fx))^m dx$	287
3.58	$\int (a + b \csc(e + fx))^m \sin(e + fx) dx$	290
3.59	$\int (a + b \csc(e + fx))^m \sin^2(e + fx) dx$	293

4 Listing of Grading functions

297

Chapter 1

Introduction

This report gives the result of running the computer algebra independent integration problems. The listing of the problems are maintained by and can be downloaded from <https://rulebasedintegration.org>

The number of integrals in this report is [59]. This is test number [129].

1.1 Listing of CAS systems tested

The following systems were tested at this time.

1. Mathematica 12.1 (64 bit) on windows 10.
2. Rubi 4.16.1 in Mathematica 12 on windows 10.
3. Maple 2020 (64 bit) on windows 10.
4. Maxima 5.43 on Linux. (via sagemath 8.9)
5. Fricas 1.3.6 on Linux (via sagemath 9.0)
6. Sympy 1.5 under Python 3.7.3 using Anaconda distribution.
7. Giac/Xcas 1.5 on Linux. (via sagemath 8.9)

Maxima, Fricas and Giac/Xcas were called from inside SageMath. This was done using SageMath integrate command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly using Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	solved	Failed
Rubi	% 100. (59)	% 0. (0)
Mathematica	% 89.83 (53)	% 10.17 (6)
Maple	% 69.49 (41)	% 30.51 (18)
Maxima	% 42.37 (25)	% 57.63 (34)
Fricas	% 69.49 (41)	% 30.51 (18)
Sympy	% 3.39 (2)	% 96.61 (57)
Giac	% 64.41 (38)	% 35.59 (21)


The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.

System	% A grade	% B grade	% C grade	% F grade
Rubi	100.	0.	0.	0.
Mathematica	61.02	16.95	11.86	10.17
Maple	42.37	27.12	0.	30.51
Maxima	20.34	22.03	0.	57.63
Fricas	30.51	38.98	0.	30.51
Sympy	3.39	0.	0.	96.61
Giac	49.15	15.25	0.	35.59

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



1.3 Performance

The table below summarizes the performance of each CAS system in terms of CPU time and leaf size of results.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.14	113.1	0.95	69.	1.
Mathematica	0.8	95.75	1.29	76.	1.06
Maple	0.11	248.15	2.98	100.	1.53
Maxima	1.16	148.56	3.03	128.	2.51
Fricas	0.5	627.49	8.81	520.	9.
Sympy	0.	0.	0.	0.	0.
Giac	1.41	172.45	2.62	110.	2.05

1.4 list of integrals that has no closed form antiderivative

{57, 58, 59}

1.5 list of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

1.7 Timing

The command `AboluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of _int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call has completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out is not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica. Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative produced was correct.

Verification phase has 3 minutes time out. An integral whose result was not verified could still be correct. Further investigation is needed on those integrals which failed verifications. Such integrals are marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since these integrals are run in a batch mode, using an automated script, and by using `sagemath` (SageMath uses Maxima), then any integral where Maxima needs an interactive response from the user to answer a question during evaluation of the integral in order to complete the integration, will fail and is counted as failed.

The exception raised is `ValueError`. Therefore Maxima result below is lower than what could result if Maxima was run directly and each question Maxima asks was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 30 such integrals out of total 705, or about 4 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be indentified by looking at the output of the integration in each section for Maxima. If the output was an exception `ValueError` then this is most likely due to this reason.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath loading of Maxima `abs_integrate` was found to cause some problem. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS and Giac/X-CAS results

There are Few integrals which failed due to SageMath not able to translate the result back to SageMath syntax and not because these CAS system were not able to do the integrations.

These will fail With error `Exception raised: NotImplementedError`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals.

Hopefully the next version of SageMath will have complete translation of FriCAS and XCAS syntax and I will re-run all the tests again when this happens.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi and Maple, the buildin system function `LeafSize` is used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special buildin function for this purpose at this time. Therefore the leaf size is determined as follows.

For Fricas, Giac and Maxima (all called via `sagemath`) the following code is used

#see <https://stackoverflow.com/questions/25202346/how-to-obtain-leaf-count-expression-size-in>

```
def tree(expr):
    if expr.operator() is None:
        return expr
    else:
        return [expr.operator()+map(tree, expr.operands())

try:
    # 1.35 is a fudge factor since this estimate of leaf count is bit lower than
    #what it should be compared to Mathematica's
    leafCount = round(1.35*len(flatten(tree(anti))))
except Exception as ee:
    leafCount =1
```

For Sympy, called directly from Python, the following code is used


```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

When these cas systems have a builtin function to find the leaf size of expressions, it will be used instead, and these tests run again.

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. It contains 13 fields. This is description of each record (line)

1. integer, the problem number.
2. integer. 0 or 1 for failed or passed. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. The optimal antiderivative in CAS own syntax.

High level overview of the CAS independent integration test build system

Chapter 2

detailed summary tables of results

2.1 List of integrals sorted by grade for each CAS

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 2, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 28, 29, 30, 31, 32, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 53, 57, 58, 59 }

B grade: { 1, 3, 4, 5, 19, 20, 36, 37, 38, 52 }

C grade: { 21, 22, 23, 24, 25, 26, 27 }

F grade: { 33, 34, 35, 54, 55, 56 }

2.1.3 Maple

A grade: { 1, 2, 3, 4, 5, 6, 7, 11, 12, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 52, 53, 57, 58, 59 }

B grade: { 8, 9, 10, 13, 14, 15, 16, 17, 18, 19, 20, 47, 48, 49, 50, 51 }

C grade: { }

F grade: { 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 54, 55, 56 }

2.1.4 Maxima

A grade: { 4, 5, 6, 16, 36, 37, 38, 52, 53, 57, 58, 59 }

B grade: { 1, 2, 3, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17 }

C grade: { }

F grade: { 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 54, 55, 56 }

2.1.5 FriCAS

A grade: { 5, 6, 7, 8, 9, 10, 16, 43, 44, 45, 46, 47, 48, 52, 53, 57, 58, 59 }

B grade: { 1, 2, 3, 4, 11, 12, 13, 14, 15, 17, 18, 19, 20, 36, 37, 38, 39, 40, 41, 42, 49, 50, 51 }

C grade: { }

F grade: { 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 54, 55, 56 }

2.1.6 Sympy

A grade: { 57, 58 }

B grade: { }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 59 }

2.1.7 Giac

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 52, 53, 57, 58, 59 }

B grade: { 13, 14, 15, 17, 18, 36, 37, 38, 51 }

C grade: { }

F grade: { 16, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 54, 55, 56 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$

Problem 1	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	A	B	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	113	89	162	537	0	130
normalized size	1	1.	2.05	1.62	2.95	9.76	0.	2.36
time (sec)	N/A	0.071	0.765	0.033	0.994	0.497	0.	1.463

Problem 2	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	B	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	83	67	131	433	0	99
normalized size	1	1.	1.89	1.52	2.98	9.84	0.	2.25
time (sec)	N/A	0.066	0.325	0.033	0.991	0.488	0.	1.395

Problem 3	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	A	B	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	63	45	92	301	0	72
normalized size	1	1.	2.33	1.67	3.41	11.15	0.	2.67
time (sec)	N/A	0.087	0.164	0.028	1.031	0.487	0.	1.348

Problem 4	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	A	A	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	44	24	42	204	0	32
normalized size	1	1.	2.2	1.2	2.1	10.2	0.	1.6
time (sec)	N/A	0.056	0.053	0.027	0.97	0.481	0.	1.369

Problem 5	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	A	A	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	26	14	22	68	0	18
normalized size	1	1.	2.17	1.17	1.83	5.67	0.	1.5
time (sec)	N/A	0.02	0.024	0.024	1.02	0.444	0.	1.337

Problem 6	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	47	41	68	142	0	43
normalized size	1	1.	1.68	1.46	2.43	5.07	0.	1.54
time (sec)	N/A	0.013	0.089	0.042	1.471	0.467	0.	1.277

Problem 7	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	B	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	32	40	105	122	0	59
normalized size	1	1.	1.28	1.6	4.2	4.88	0.	2.36
time (sec)	N/A	0.047	0.078	0.043	1.458	0.473	0.	1.33

Problem 8	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	B	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	42	100	173	166	0	76
normalized size	1	1.	1.05	2.5	4.32	4.15	0.	1.9
time (sec)	N/A	0.061	0.13	0.042	1.459	0.474	0.	1.368

Problem 9	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	B	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	49	121	243	211	0	90
normalized size	1	1.	0.92	2.28	4.58	3.98	0.	1.7
time (sec)	N/A	0.067	0.164	0.047	1.464	0.483	0.	1.277

Problem 10	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	B	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	66	57	185	311	259	0	123
normalized size	1	1.	0.86	2.8	4.71	3.92	0.	1.86
time (sec)	N/A	0.073	0.209	0.046	1.472	0.485	0.	1.392

Problem 11	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	B	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	108	83	192	306	0	81
normalized size	1	1.	1.89	1.46	3.37	5.37	0.	1.42
time (sec)	N/A	0.066	0.325	0.063	1.47	0.481	0.	1.33

Problem 12	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	B	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	123	125	308	475	0	116
normalized size	1	1.	1.4	1.42	3.5	5.4	0.	1.32
time (sec)	N/A	0.11	0.944	0.079	1.509	0.486	0.	1.251

Problem 13	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	B	B	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	80	535	563	900	0	339
normalized size	1	1.	1.23	8.23	8.66	13.85	0.	5.22
time (sec)	N/A	0.091	1.63	0.21	1.583	0.515	0.	2.242

Problem 14	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	B	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	69	273	270	666	0	263
normalized size	1	1.	1.57	6.2	6.14	15.14	0.	5.98
time (sec)	N/A	0.03	0.085	0.16	1.543	0.502	0.	2.074

Problem 15	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	B	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	32	199	200	377	0	477
normalized size	1	1.	1.23	7.65	7.69	14.5	0.	18.35
time (sec)	N/A	0.016	0.052	0.151	1.582	0.489	0.	2.038

Problem 16	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	A	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	54	221	112	668	0	0
normalized size	1	1.	0.87	3.56	1.81	10.77	0.	0.
time (sec)	N/A	0.058	0.126	0.139	1.577	0.506	0.	0.

Problem 17	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	B	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	81	129	1141	203	1299	0	410
normalized size	1	1.	1.59	14.09	2.51	16.04	0.	5.06
time (sec)	N/A	0.102	0.411	0.184	1.544	0.532	0.	2.973

Problem 18	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	F	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	139	1961	0	1677	0	470
normalized size	1	1.	1.39	19.61	0.	16.77	0.	4.7
time (sec)	N/A	0.148	0.503	0.148	0.	0.546	0.	2.529

Problem 19	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	B	F	B	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	108	114	0	755	0	0
normalized size	1	1.	2.92	3.08	0.	20.41	0.	0.
time (sec)	N/A	0.059	0.408	0.359	0.	0.535	0.	0.

Problem 20	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	B	F	B	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	101	117	0	764	0	0
normalized size	1	1.	2.66	3.08	0.	20.11	0.	0.
time (sec)	N/A	0.071	0.775	0.333	0.	0.53	0.	0.

Problem 21	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	F	F	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	254	254	102	0	0	0	0	0
normalized size	1	1.	0.4	0.	0.	0.	0.	0.
time (sec)	N/A	0.281	0.382	0.384	0.	0.	0.	0.

Problem 22	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	213	46	0	0	0	0	0
normalized size	1	1.	0.22	0.	0.	0.	0.	0.
time (sec)	N/A	0.128	0.223	0.645	0.	0.	0.	0.

Problem 23	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	254	254	110	0	0	0	0	0
normalized size	1	1.	0.43	0.	0.	0.	0.	0.
time (sec)	N/A	0.145	0.447	0.448	0.	0.	0.	0.

Problem 24	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	F	F	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	514	514	120	0	0	0	0	0
normalized size	1	1.	0.23	0.	0.	0.	0.	0.
time (sec)	N/A	0.299	1.187	0.335	0.	0.	0.	0.

Problem 25	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	470	470	109	0	0	0	0	0
normalized size	1	1.	0.23	0.	0.	0.	0.	0.
time (sec)	N/A	0.253	0.985	0.325	0.	0.	0.	0.

Problem 26	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	508	508	46	0	0	0	0	0
normalized size	1	1.	0.09	0.	0.	0.	0.	0.
time (sec)	N/A	0.275	0.755	0.44	0.	0.	0.	0.

Problem 27	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	552	552	72	0	0	0	0	0
normalized size	1	1.	0.13	0.	0.	0.	0.	0.
time (sec)	N/A	0.307	1.201	0.323	0.	0.	0.	0.

Problem 28	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	48	0	0	0	0	0
normalized size	1	1.	1.	0.	0.	0.	0.	0.
time (sec)	N/A	0.058	0.169	0.401	0.	0.	0.	0.

Problem 29	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	73	0	0	0	0	0
normalized size	1	1.	1.06	0.	0.	0.	0.	0.
time (sec)	N/A	0.07	0.913	0.484	0.	0.	0.	0.

Problem 30	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	156	199	0	0	0	0	0
normalized size	1	1.	1.28	0.	0.	0.	0.	0.
time (sec)	N/A	0.19	0.726	0.609	0.	0.	0.	0.

Problem 31	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	109	131	0	0	0	0	0
normalized size	1	1.	1.2	0.	0.	0.	0.	0.
time (sec)	N/A	0.103	0.431	0.448	0.	0.	0.	0.

Problem 32	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	60	0	0	0	0	0
normalized size	1	1.	0.81	0.	0.	0.	0.	0.
time (sec)	N/A	0.056	0.202	0.429	0.	0.	0.	0.

Problem 33	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	84	84	0	0	0	0	0	0
normalized size	1	1.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.057	0.595	0.287	0.	0.	0.	0.

Problem 34	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	83	83	0	0	0	0	0	0
normalized size	1	1.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.09	3.647	0.678	0.	0.	0.	0.

Problem 35	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	F	F	F	F	F(-1)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	84	84	0	0	0	0	0	0
normalized size	1	1.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.105	7.648	0.882	0.	0.	0.	0.

Problem 36	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	A	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	568	139	169	524	0	277
normalized size	1	1.	5.31	1.3	1.58	4.9	0.	2.59
time (sec)	N/A	0.11	6.238	0.05	1.006	0.524	0.	1.598

Problem 37	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	A	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	73	73	152	99	128	383	0	181
normalized size	1	1.	2.08	1.36	1.75	5.25	0.	2.48
time (sec)	N/A	0.047	0.648	0.044	1.004	0.511	0.	1.473

Problem 38	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	A	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	76	52	58	209	0	100
normalized size	1	1.	2.24	1.53	1.71	6.15	0.	2.94
time (sec)	N/A	0.026	0.182	0.029	0.996	0.504	0.	1.428

Problem 39	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	112	125	162	0	1378	0	262
normalized size	1	1.	1.12	1.45	0.	12.3	0.	2.34
time (sec)	N/A	0.407	1.631	0.05	0.	0.898	0.	1.387

Problem 40	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	144	112	0	1146	0	190
normalized size	1	1.	1.71	1.33	0.	13.64	0.	2.26
time (sec)	N/A	0.252	0.487	0.045	0.	0.879	0.	1.445

Problem 41	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	106	77	0	768	0	132
normalized size	1	1.	1.71	1.24	0.	12.39	0.	2.13
time (sec)	N/A	0.155	0.204	0.049	0.	0.624	0.	1.285

Problem 42	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	62	53	0	595	0	85
normalized size	1	1.	1.17	1.	0.	11.23	0.	1.6
time (sec)	N/A	0.113	0.061	0.045	0.	0.612	0.	1.583

Problem 43	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	40	39	0	360	0	65
normalized size	1	1.	1.	0.98	0.	9.	0.	1.62
time (sec)	N/A	0.07	0.024	0.042	0.	0.497	0.	1.432

Problem 44	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	59	70	0	525	0	104
normalized size	1	1.	1.04	1.23	0.	9.21	0.	1.82
time (sec)	N/A	0.065	0.107	0.056	0.	0.521	0.	1.455

Problem 45	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	56	72	0	520	0	104
normalized size	1	1.	0.92	1.18	0.	8.52	0.	1.7
time (sec)	N/A	0.107	0.099	0.061	0.	0.532	0.	1.456

Problem 46	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	F(-2)	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	82	82	78	142	0	644	0	151
normalized size	1	1.	0.95	1.73	0.	7.85	0.	1.84
time (sec)	N/A	0.261	0.119	0.072	0.	0.54	0.	1.381

Problem 47	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	F(-2)	A	F(-1)	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	110	98	213	0	744	0	201
normalized size	1	1.	0.89	1.94	0.	6.76	0.	1.83
time (sec)	N/A	0.398	0.238	0.066	0.	0.552	0.	1.437

Problem 48	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	F(-2)	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	144	129	405	0	909	0	340
normalized size	1	1.	0.9	2.81	0.	6.31	0.	2.36
time (sec)	N/A	0.588	0.315	0.079	0.	0.581	0.	1.388

Problem 49	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	F(-2)	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	108	139	247	0	1076	0	213
normalized size	1	1.	1.29	2.29	0.	9.96	0.	1.97
time (sec)	N/A	0.171	0.456	0.093	0.	0.563	0.	1.384

Problem 50	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	F(-2)	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	170	216	796	0	1994	0	401
normalized size	1	1.	1.27	4.68	0.	11.73	0.	2.36
time (sec)	N/A	0.319	1.103	0.115	0.	0.645	0.	1.181

Problem 51	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	B	F(-2)	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	239	239	279	1912	0	3389	0	722
normalized size	1	1.	1.17	8.	0.	14.18	0.	3.02
time (sec)	N/A	0.504	2.095	0.147	0.	0.778	0.	1.39

Problem 52	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	B	A	A	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	66	36	66	89	0	66
normalized size	1	1.	2.13	1.16	2.13	2.87	0.	2.13
time (sec)	N/A	0.029	0.048	0.039	1.474	0.492	0.	1.443

Problem 53	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	A	A	A	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	67	53	96	144	0	61
normalized size	1	1.	0.99	0.78	1.41	2.12	0.	0.9
time (sec)	N/A	0.039	0.047	0.045	1.463	0.496	0.	1.374

Problem 54	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	274	274	0	0	0	0	0	0
normalized size	1	1.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.343	4.181	0.309	0.	0.	0.	0.

Problem 55	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	220	220	0	0	0	0	0	0
normalized size	1	1.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.226	2.718	0.247	0.	0.	0.	0.

Problem 56	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	A	A	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	104	104	0	0	0	0	0	0
normalized size	1	1.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.074	1.913	0.511	0.	0.	0.	0.

Problem 57	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	14	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.01	1.546	0.333	0.	0.	0.	0.

Problem 58	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	21	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.032	6.055	0.257	0.	0.	0.	0.

Problem 59	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac
grade	N/A	A	A	A	A	A	F(-1)	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	23	0	0	0	0	0	0	0
normalized size	1	0.	0.	0.	0.	0.	0.	0.
time (sec)	N/A	0.04	5.904	0.567	0.	0.	0.	0.

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [39] had the largest ratio of [0.6923]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	5	1.	13	0.385
2	A	6	6	1.	13	0.462
3	A	4	4	1.	13	0.308
4	A	3	3	1.	13	0.231
5	A	1	1	1.	11	0.091
6	A	2	2	1.	12	0.167
7	A	4	4	1.	11	0.364
8	A	5	5	1.	13	0.385
9	A	6	5	1.	13	0.385
10	A	7	5	1.	13	0.385
11	A	3	3	1.	12	0.25
12	A	4	4	1.	12	0.333
13	A	5	5	1.	10	0.5
14	A	4	4	1.	10	0.4
15	A	2	2	1.	10	0.2
16	A	5	4	1.	10	0.4
17	A	6	5	1.	10	0.5
18	A	7	6	1.	10	0.6
19	A	2	2	1.	25	0.08
20	A	2	2	1.	28	0.071
21	A	4	4	1.	25	0.16
22	A	3	3	1.	25	0.12
23	A	4	4	1.	25	0.16

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
24	A	6	6	1.	25	0.24
25	A	5	5	1.	25	0.2
26	A	6	6	1.	25	0.24
27	A	7	6	1.	25	0.24
28	A	2	2	1.	23	0.087
29	A	3	3	1.	24	0.125
30	A	5	5	1.	21	0.238
31	A	4	4	1.	21	0.19
32	A	3	3	1.	19	0.158
33	A	3	3	1.	12	0.25
34	A	3	3	1.	19	0.158
35	A	3	3	1.	21	0.143
36	A	6	5	1.	12	0.417
37	A	5	4	1.	12	0.333
38	A	4	4	1.	12	0.333
39	A	9	9	1.	13	0.692
40	A	8	8	1.	13	0.615
41	A	7	7	1.	13	0.538
42	A	6	6	1.	13	0.462
43	A	4	4	1.	11	0.364
44	A	4	4	1.	12	0.333
45	A	6	6	1.	11	0.546
46	A	7	7	1.	13	0.538
47	A	8	7	1.	13	0.538
48	A	9	7	1.	13	0.538
49	A	6	6	1.	12	0.5
50	A	7	7	1.	12	0.583
51	A	8	7	1.	12	0.583
52	A	2	2	1.	12	0.167
53	A	5	4	1.	12	0.333
54	A	8	5	1.	21	0.238
55	A	7	4	1.	21	0.19

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
56	A	3	3	1.	19	0.158
57	A	0	0	0.	0	0.
58	A	0	0	0.	0	0.
59	A	0	0	0.	0	0.

Chapter 3

Listing of integrals

3.1 $\int \frac{\csc^5(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=55

$$-\frac{4 \cot^3(x)}{3a} - \frac{4 \cot(x)}{a} + \frac{3 \tanh^{-1}(\cos(x))}{2a} + \frac{\cot(x) \csc^3(x)}{a \csc(x) + a} + \frac{3 \cot(x) \csc(x)}{2a}$$

[Out] (3*ArcTanh[Cos[x]])/(2*a) - (4*Cot[x])/a - (4*Cot[x]^3)/(3*a) + (3*Cot[x]*Csc[x])/(2*a) + (Cot[x]*Csc[x]^3)/(a + a*Csc[x])

Rubi [A] time = 0.0714401, antiderivative size = 55, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 5, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3818, 3787, 3768, 3770, 3767}

$$-\frac{4 \cot^3(x)}{3a} - \frac{4 \cot(x)}{a} + \frac{3 \tanh^{-1}(\cos(x))}{2a} + \frac{\cot(x) \csc^3(x)}{a \csc(x) + a} + \frac{3 \cot(x) \csc(x)}{2a}$$

Antiderivative was successfully verified.

[In] Int[Csc[x]^5/(a + a*Csc[x]),x]

[Out] (3*ArcTanh[Cos[x]])/(2*a) - (4*Cot[x])/a - (4*Cot[x]^3)/(3*a) + (3*Cot[x]*Csc[x])/(2*a) + (Cot[x]*Csc[x]^3)/(a + a*Csc[x])

Rule 3818

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[(d^2*Cot[e + f*x]*(d*Csc[e + f*x])^(n - 2))/(f*(a + b*Csc[e + f*x])), x] - Dist[d^2/(a*b), Int[(d*Csc[e + f*x])^(n - 2)*(b*(n - 2) - a*(n - 1)*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && GtQ[n, 1]
```

Rule 3787

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Dist[a, Int[(d*Csc[e + f*x])^n, x], x] + Dist[b/d, Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]
```

Rule 3768

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 3770

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3767

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned}
 \int \frac{\csc^5(x)}{a + a \csc(x)} dx &= \frac{\cot(x) \csc^3(x)}{a + a \csc(x)} - \frac{\int \csc^3(x)(3a - 4a \csc(x)) dx}{a^2} \\
 &= \frac{\cot(x) \csc^3(x)}{a + a \csc(x)} - \frac{3 \int \csc^3(x) dx}{a} + \frac{4 \int \csc^4(x) dx}{a} \\
 &= \frac{3 \cot(x) \csc(x)}{2a} + \frac{\cot(x) \csc^3(x)}{a + a \csc(x)} - \frac{3 \int \csc(x) dx}{2a} - \frac{4 \text{Subst}\left(\int (1 + x^2) dx, x, \cot(x)\right)}{a} \\
 &= \frac{3 \tanh^{-1}(\cos(x))}{2a} - \frac{4 \cot(x)}{a} - \frac{4 \cot^3(x)}{3a} + \frac{3 \cot(x) \csc(x)}{2a} + \frac{\cot(x) \csc^3(x)}{a + a \csc(x)}
 \end{aligned}$$

Mathematica [B] time = 0.764763, size = 113, normalized size = 2.05

$$\frac{20 \tan\left(\frac{x}{2}\right) - 20 \cot\left(\frac{x}{2}\right) + 3 \csc^2\left(\frac{x}{2}\right) - 3 \sec^2\left(\frac{x}{2}\right) - 36 \log\left(\sin\left(\frac{x}{2}\right)\right) + 36 \log\left(\cos\left(\frac{x}{2}\right)\right) + \frac{48 \sin\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)} + 8 \sin^4\left(\frac{x}{2}\right) \csc^3\left(\frac{x}{2}\right)}{24a}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[x]^5/(a + a*Csc[x]),x]

[Out] (-20*Cot[x/2] + 3*Csc[x/2]^2 + 36*Log[Cos[x/2]] - 36*Log[Sin[x/2]] - 3*Sec[x/2]^2 + 8*Csc[x]^3*Sin[x/2]^4 + (48*Sin[x/2])/(Cos[x/2] + Sin[x/2]) - (Csc[x/2]^4*Sin[x])/2 + 20*Tan[x/2])/(24*a)

Maple [A] time = 0.033, size = 89, normalized size = 1.6

$$\frac{1}{24a} \left(\tan\left(\frac{x}{2}\right) \right)^3 - \frac{1}{8a} \left(\tan\left(\frac{x}{2}\right) \right)^2 + \frac{7}{8a} \tan\left(\frac{x}{2}\right) - 2 \frac{1}{a(\tan(x/2) + 1)} - \frac{1}{24a} \left(\tan\left(\frac{x}{2}\right) \right)^{-3} + \frac{1}{8a} \left(\tan\left(\frac{x}{2}\right) \right)^{-2} - \frac{7}{8a} \left(\tan\left(\frac{x}{2}\right) \right)^{-1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(x)^5/(a+a*csc(x)),x)

[Out] 1/24/a*tan(1/2*x)^3-1/8/a*tan(1/2*x)^2+7/8/a*tan(1/2*x)-2/a/(tan(1/2*x)+1)-1/24/a/tan(1/2*x)^3+1/8/a/tan(1/2*x)^2-7/8/a/tan(1/2*x)-3/2/a*ln(tan(1/2*x))

Maxima [B] time = 0.993993, size = 162, normalized size = 2.95

$$\frac{\frac{21 \sin(x)}{\cos(x)+1} - \frac{3 \sin(x)^2}{(\cos(x)+1)^2} + \frac{\sin(x)^3}{(\cos(x)+1)^3}}{24a} + \frac{\frac{2 \sin(x)}{\cos(x)+1} - \frac{18 \sin(x)^2}{(\cos(x)+1)^2} - \frac{69 \sin(x)^3}{(\cos(x)+1)^3} - 1}{24 \left(\frac{a \sin(x)^3}{(\cos(x)+1)^3} + \frac{a \sin(x)^4}{(\cos(x)+1)^4} \right)} - \frac{3 \log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^5/(a+a*csc(x)),x, algorithm="maxima")

[Out] 1/24*(21*sin(x)/(cos(x) + 1) - 3*sin(x)^2/(cos(x) + 1)^2 + sin(x)^3/(cos(x) + 1)^3)/a + 1/24*(2*sin(x)/(cos(x) + 1) - 18*sin(x)^2/(cos(x) + 1)^2 - 69*

$$\frac{\sin(x)^3/(\cos(x) + 1)^3 - 1}{a \sin(x)^3/(\cos(x) + 1)^3 + a \sin(x)^4/(\cos(x) + 1)^4} - \frac{3}{2} \log(\sin(x)/(\cos(x) + 1))/a$$

Fricas [B] time = 0.497218, size = 537, normalized size = 9.76

$$\frac{32 \cos(x)^4 + 14 \cos(x)^3 - 48 \cos(x)^2 + 9(\cos(x)^4 - 2 \cos(x)^2 - (\cos(x)^3 + \cos(x)^2 - \cos(x) - 1) \sin(x) + 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - 9(\cos(x)^4 - 2 \cos(x)^2 - (\cos(x)^3 + \cos(x)^2 - \cos(x) - 1) \sin(x) + 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) + 2(16 \cos(x)^3 + 9 \cos(x)^2 - 15 \cos(x) - 6) \sin(x) - 18 \cos(x) + 12}{12(a \cos(x)^4 - 2a \cos(x)^2 - (a \cos(x)^3 + a \cos(x)^2 - a \cos(x) - a) \sin(x) + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^5/(a+a*csc(x)),x, algorithm="fricas")

[Out] $\frac{1}{12} * (32 * \cos(x)^4 + 14 * \cos(x)^3 - 48 * \cos(x)^2 + 9 * (\cos(x)^4 - 2 * \cos(x)^2 - (\cos(x)^3 + \cos(x)^2 - \cos(x) - 1) * \sin(x) + 1) * \log(1/2 * \cos(x) + 1/2) - 9 * (\cos(x)^4 - 2 * \cos(x)^2 - (\cos(x)^3 + \cos(x)^2 - \cos(x) - 1) * \sin(x) + 1) * \log(-1/2 * \cos(x) + 1/2) + 2 * (16 * \cos(x)^3 + 9 * \cos(x)^2 - 15 * \cos(x) - 6) * \sin(x) - 18 * \cos(x) + 12) / (a * \cos(x)^4 - 2 * a * \cos(x)^2 - (a * \cos(x)^3 + a * \cos(x)^2 - a * \cos(x) - a) * \sin(x) + a)$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\frac{\int \frac{\csc^5(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)**5/(a+a*csc(x)),x)

[Out] Integral(csc(x)**5/(csc(x) + 1), x)/a

Giac [A] time = 1.46273, size = 130, normalized size = 2.36

$$-\frac{3 \log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{2a} + \frac{a^2 \tan\left(\frac{1}{2}x\right)^3 - 3a^2 \tan\left(\frac{1}{2}x\right)^2 + 21a^2 \tan\left(\frac{1}{2}x\right)}{24a^3} - \frac{2}{a\left(\tan\left(\frac{1}{2}x\right) + 1\right)} + \frac{66 \tan\left(\frac{1}{2}x\right)^3 - 21 \tan\left(\frac{1}{2}x\right)}{24a \tan\left(\frac{1}{2}x\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(x)^5/(a+a*csc(x)),x, algorithm="giac")
```

```
[Out] -3/2*log(abs(tan(1/2*x)))/a + 1/24*(a^2*tan(1/2*x)^3 - 3*a^2*tan(1/2*x)^2 +  
21*a^2*tan(1/2*x))/a^3 - 2/(a*(tan(1/2*x) + 1)) + 1/24*(66*tan(1/2*x)^3 -  
21*tan(1/2*x)^2 + 3*tan(1/2*x) - 1)/(a*tan(1/2*x)^3)
```

3.2 $\int \frac{\csc^4(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=44

$$\frac{2 \cot(x)}{a} - \frac{3 \tanh^{-1}(\cos(x))}{2a} + \frac{\cot(x) \csc^2(x)}{a \csc(x) + a} - \frac{3 \cot(x) \csc(x)}{2a}$$

[Out] $(-3*\text{ArcTanh}[\text{Cos}[x]])/(2*a) + (2*\text{Cot}[x])/a - (3*\text{Cot}[x]*\text{Csc}[x])/(2*a) + (\text{Cot}[x]*\text{Csc}[x]^2)/(a + a*\text{Csc}[x])$

Rubi [A] time = 0.0659157, antiderivative size = 44, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 6, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3818, 3787, 3767, 8, 3768, 3770}

$$\frac{2 \cot(x)}{a} - \frac{3 \tanh^{-1}(\cos(x))}{2a} + \frac{\cot(x) \csc^2(x)}{a \csc(x) + a} - \frac{3 \cot(x) \csc(x)}{2a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[x]^4/(a + a*\text{Csc}[x]), x]$

[Out] $(-3*\text{ArcTanh}[\text{Cos}[x]])/(2*a) + (2*\text{Cot}[x])/a - (3*\text{Cot}[x]*\text{Csc}[x])/(2*a) + (\text{Cot}[x]*\text{Csc}[x]^2)/(a + a*\text{Csc}[x])$

Rule 3818

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}/(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[(d^2*\text{Cot}[e + f*x]*(d*\text{Csc}[e + f*x])^{(n - 2)})/(f*(a + b*\text{Csc}[e + f*x])), x] - \text{Dist}[d^2/(a*b), \text{Int}[(d*\text{Csc}[e + f*x])^{(n - 2)}*(b*(n - 2) - a*(n - 1)*\text{Csc}[e + f*x]), x], x] /;$ $\text{FreeQ}\{a, b, d, e, f\}, x \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[n, 1]$

Rule 3787

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Dist}[b/d, \text{Int}[(d*\text{Csc}[e + f*x])^{(n + 1)}, x], x] /;$ $\text{FreeQ}\{a, b, d, e, f, n\}, x]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] /;$ $\text{FreeQ}\{c,$

d}, x] && IGtQ[n/2, 0]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x] * (b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\csc^4(x)}{a + a \csc(x)} dx &= \frac{\cot(x) \csc^2(x)}{a + a \csc(x)} - \frac{\int \csc^2(x)(2a - 3a \csc(x)) dx}{a^2} \\ &= \frac{\cot(x) \csc^2(x)}{a + a \csc(x)} - \frac{2 \int \csc^2(x) dx}{a} + \frac{3 \int \csc^3(x) dx}{a} \\ &= -\frac{3 \cot(x) \csc(x)}{2a} + \frac{\cot(x) \csc^2(x)}{a + a \csc(x)} + \frac{3 \int \csc(x) dx}{2a} + \frac{2 \text{Subst}(\int 1 dx, x, \cot(x))}{a} \\ &= -\frac{3 \tanh^{-1}(\cos(x))}{2a} + \frac{2 \cot(x)}{a} - \frac{3 \cot(x) \csc(x)}{2a} + \frac{\cot(x) \csc^2(x)}{a + a \csc(x)} \end{aligned}$$

Mathematica [A] time = 0.324768, size = 83, normalized size = 1.89

$$\frac{-4 \tan\left(\frac{x}{2}\right) + 4 \cot\left(\frac{x}{2}\right) - \csc^2\left(\frac{x}{2}\right) + \sec^2\left(\frac{x}{2}\right) + 12 \log\left(\sin\left(\frac{x}{2}\right)\right) - 12 \log\left(\cos\left(\frac{x}{2}\right)\right) - \frac{16 \sin\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)}}{8a}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[x]^4/(a + a*Csc[x]), x]

[Out] $(4*\text{Cot}[x/2] - \text{Csc}[x/2]^2 - 12*\text{Log}[\text{Cos}[x/2]] + 12*\text{Log}[\text{Sin}[x/2]] + \text{Sec}[x/2]^2 - (16*\text{Sin}[x/2])/(\text{Cos}[x/2] + \text{Sin}[x/2]) - 4*\text{Tan}[x/2])/(8*a)$

Maple [A] time = 0.033, size = 67, normalized size = 1.5

$$\frac{1}{8a} \left(\tan\left(\frac{x}{2}\right) \right)^2 - \frac{1}{2a} \tan\left(\frac{x}{2}\right) + 2 \frac{1}{a(\tan(x/2) + 1)} - \frac{1}{8a} \left(\tan\left(\frac{x}{2}\right) \right)^{-2} + \frac{1}{2a} \left(\tan\left(\frac{x}{2}\right) \right)^{-1} + \frac{3}{2a} \ln\left(\tan\left(\frac{x}{2}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(x)^4/(a+a*csc(x)),x)`

[Out] $1/8/a*\tan(1/2*x)^2-1/2/a*\tan(1/2*x)+2/a/(\tan(1/2*x)+1)-1/8/a/\tan(1/2*x)^2+1/2/a/\tan(1/2*x)+3/2/a*\ln(\tan(1/2*x))$

Maxima [B] time = 0.991286, size = 131, normalized size = 2.98

$$-\frac{\frac{4 \sin(x)}{\cos(x)+1} - \frac{\sin(x)^2}{(\cos(x)+1)^2}}{8a} + \frac{\frac{3 \sin(x)}{\cos(x)+1} + \frac{20 \sin(x)^2}{(\cos(x)+1)^2} - 1}{8 \left(\frac{a \sin(x)^2}{(\cos(x)+1)^2} + \frac{a \sin(x)^3}{(\cos(x)+1)^3} \right)} + \frac{3 \log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(x)^4/(a+a*csc(x)),x, algorithm="maxima")`

[Out] $-1/8*(4*\sin(x)/(\cos(x) + 1) - \sin(x)^2/(\cos(x) + 1)^2)/a + 1/8*(3*\sin(x)/(\cos(x) + 1) + 20*\sin(x)^2/(\cos(x) + 1)^2 - 1)/(a*\sin(x)^2/(\cos(x) + 1)^2 + a*\sin(x)^3/(\cos(x) + 1)^3) + 3/2*\log(\sin(x)/(\cos(x) + 1))/a$

Fricas [B] time = 0.488282, size = 433, normalized size = 9.84

$$\frac{8 \cos(x)^3 + 6 \cos(x)^2 - 3(\cos(x)^3 + \cos(x)^2 + (\cos(x)^2 - 1)\sin(x) - \cos(x) - 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + 3(\cos(x)^3 + \cos(x)^2 - \cos(x) - 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right)}{4(a \cos(x)^3 + a \cos(x)^2 - a \cos(x) - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^4/(a+a*csc(x)),x, algorithm="fricas")

[Out] $\frac{1}{4}*(8*\cos(x)^3 + 6*\cos(x)^2 - 3*(\cos(x)^3 + \cos(x)^2 + (\cos(x)^2 - 1)*\sin(x) - \cos(x) - 1)*\log(1/2*\cos(x) + 1/2) + 3*(\cos(x)^3 + \cos(x)^2 + (\cos(x)^2 - 1)*\sin(x) - \cos(x) - 1)*\log(-1/2*\cos(x) + 1/2) - 2*(4*\cos(x)^2 + \cos(x) - 2)*\sin(x) - 6*\cos(x) - 4)/(a*\cos(x)^3 + a*\cos(x)^2 - a*\cos(x) + (a*\cos(x)^2 - a)*\sin(x) - a)$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\csc^4(x)}{\csc(x)+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)**4/(a+a*csc(x)),x)

[Out] Integral(csc(x)**4/(csc(x) + 1), x)/a

Giac [A] time = 1.39513, size = 99, normalized size = 2.25

$$\frac{3 \log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{2a} + \frac{a \tan\left(\frac{1}{2}x\right)^2 - 4a \tan\left(\frac{1}{2}x\right)}{8a^2} + \frac{2}{a\left(\tan\left(\frac{1}{2}x\right) + 1\right)} - \frac{18 \tan\left(\frac{1}{2}x\right)^2 - 4 \tan\left(\frac{1}{2}x\right) + 1}{8a \tan\left(\frac{1}{2}x\right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^4/(a+a*csc(x)),x, algorithm="giac")

[Out] $\frac{3}{2}*\log(\text{abs}(\tan(1/2*x)))/a + 1/8*(a*\tan(1/2*x)^2 - 4*a*\tan(1/2*x))/a^2 + 2/(a*(\tan(1/2*x) + 1)) - 1/8*(18*\tan(1/2*x)^2 - 4*\tan(1/2*x) + 1)/(a*\tan(1/2*x)^2)$

3.3 $\int \frac{\csc^3(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=27

$$-\frac{\cot(x)}{a} + \frac{\tanh^{-1}(\cos(x))}{a} - \frac{\cot(x)}{a \csc(x) + a}$$

[Out] ArcTanh[Cos[x]]/a - Cot[x]/a - Cot[x]/(a + a*Csc[x])

Rubi [A] time = 0.0867268, antiderivative size = 27, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3790, 3789, 3770, 3794}

$$-\frac{\cot(x)}{a} + \frac{\tanh^{-1}(\cos(x))}{a} - \frac{\cot(x)}{a \csc(x) + a}$$

Antiderivative was successfully verified.

[In] Int[Csc[x]^3/(a + a*Csc[x]),x]

[Out] ArcTanh[Cos[x]]/a - Cot[x]/a - Cot[x]/(a + a*Csc[x])

Rule 3790

Int[csc[(e_.) + (f_.)*(x_)]^3/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] :> -Simp[Cot[e + f*x]/(b*f), x] - Dist[a/b, Int[Csc[e + f*x]^2/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x]

Rule 3789

Int[csc[(e_.) + (f_.)*(x_)]^2/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] :> Dist[1/b, Int[Csc[e + f*x], x], x] - Dist[a/b, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3794

```
Int[csc[(e_.) + (f_.)*(x_.)]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol]
:= -Simp[Cot[e + f*x]/(f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x]
&& EqQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\csc^3(x)}{a + a \csc(x)} dx &= -\frac{\cot(x)}{a} - \int \frac{\csc^2(x)}{a + a \csc(x)} dx \\ &= -\frac{\cot(x)}{a} - \frac{\int \csc(x) dx}{a} + \int \frac{\csc(x)}{a + a \csc(x)} dx \\ &= \frac{\tanh^{-1}(\cos(x))}{a} - \frac{\cot(x)}{a} - \frac{\cot(x)}{a + a \csc(x)} \end{aligned}$$

Mathematica [B] time = 0.164203, size = 63, normalized size = 2.33

$$\frac{\tan\left(\frac{x}{2}\right) - \cot\left(\frac{x}{2}\right) - 2 \log\left(\sin\left(\frac{x}{2}\right)\right) + 2 \log\left(\cos\left(\frac{x}{2}\right)\right) + \frac{4 \sin\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)}}{2a}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[x]^3/(a + a*Csc[x]), x]
```

```
[Out] (-Cot[x/2] + 2*Log[Cos[x/2]] - 2*Log[Sin[x/2]] + (4*Sin[x/2]))/(Cos[x/2] + Sin[x/2]) + Tan[x/2]/(2*a)
```

Maple [A] time = 0.028, size = 45, normalized size = 1.7

$$\frac{1}{2a} \tan\left(\frac{x}{2}\right) - 2 \frac{1}{a(\tan(x/2) + 1)} - \frac{1}{2a} \left(\tan\left(\frac{x}{2}\right)\right)^{-1} - \frac{1}{a} \ln\left(\tan\left(\frac{x}{2}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(x)^3/(a+a*csc(x)), x)
```

```
[Out] 1/2/a*tan(1/2*x)-2/a/(tan(1/2*x)+1)-1/2/a/tan(1/2*x)-1/a*ln(tan(1/2*x))
```

Maxima [B] time = 1.0311, size = 92, normalized size = 3.41

$$-\frac{\frac{5 \sin(x)}{\cos(x)+1} + 1}{2 \left(\frac{a \sin(x)}{\cos(x)+1} + \frac{a \sin(x)^2}{(\cos(x)+1)^2} \right)} - \frac{\log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a} + \frac{\sin(x)}{2a(\cos(x)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^3/(a+a*csc(x)),x, algorithm="maxima")

[Out] -1/2*(5*sin(x)/(cos(x) + 1) + 1)/(a*sin(x)/(cos(x) + 1) + a*sin(x)^2/(cos(x) + 1)^2) - log(sin(x)/(cos(x) + 1))/a + 1/2*sin(x)/(a*(cos(x) + 1))

Fricas [B] time = 0.486788, size = 301, normalized size = 11.15

$$\frac{4 \cos(x)^2 + (\cos(x)^2 - (\cos(x) + 1) \sin(x) - 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x)^2 - (\cos(x) + 1) \sin(x) - 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)}{2(a \cos(x)^2 - (a \cos(x) + a) \sin(x) - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^3/(a+a*csc(x)),x, algorithm="fricas")

[Out] 1/2*(4*cos(x)^2 + (cos(x)^2 - (cos(x) + 1)*sin(x) - 1)*log(1/2*cos(x) + 1/2) - (cos(x)^2 - (cos(x) + 1)*sin(x) - 1)*log(-1/2*cos(x) + 1/2) + 2*(2*cos(x) + 1)*sin(x) + 2*cos(x) - 2)/(a*cos(x)^2 - (a*cos(x) + a)*sin(x) - a)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\frac{\int \frac{\csc^3(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)**3/(a+a*csc(x)),x)

[Out] Integral(csc(x)**3/(csc(x) + 1), x)/a

Giac [A] time = 1.34789, size = 72, normalized size = 2.67

$$-\frac{\log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{a} + \frac{\tan\left(\frac{1}{2}x\right)}{2a} + \frac{\tan\left(\frac{1}{2}x\right)^2 - 4\tan\left(\frac{1}{2}x\right) - 1}{2\left(\tan\left(\frac{1}{2}x\right)^2 + \tan\left(\frac{1}{2}x\right)\right)a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^3/(a+a*csc(x)),x, algorithm="giac")

[Out] -log(abs(tan(1/2*x)))/a + 1/2*tan(1/2*x)/a + 1/2*(tan(1/2*x)^2 - 4*tan(1/2*x) - 1)/((tan(1/2*x)^2 + tan(1/2*x))*a)

$$3.4 \quad \int \frac{\csc^2(x)}{a+a \csc(x)} dx$$

Optimal. Leaf size=20

$$\frac{\cot(x)}{a \csc(x) + a} - \frac{\tanh^{-1}(\cos(x))}{a}$$

[Out] -(ArcTanh[Cos[x]]/a) + Cot[x]/(a + a*Csc[x])

Rubi [A] time = 0.0560358, antiderivative size = 20, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3789, 3770, 3794}

$$\frac{\cot(x)}{a \csc(x) + a} - \frac{\tanh^{-1}(\cos(x))}{a}$$

Antiderivative was successfully verified.

[In] Int[Csc[x]^2/(a + a*Csc[x]),x]

[Out] -(ArcTanh[Cos[x]]/a) + Cot[x]/(a + a*Csc[x])

Rule 3789

Int[csc[(e_.) + (f_.)*(x_)]^2/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] :> Dist[1/b, Int[Csc[e + f*x], x], x] - Dist[a/b, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3794

Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] :> -Simp[Cot[e + f*x]/(f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]

Rubi steps

$$\int \frac{\csc^2(x)}{a + a \csc(x)} dx = \frac{\int \csc(x) dx}{a} - \int \frac{\csc(x)}{a + a \csc(x)} dx$$

$$= -\frac{\tanh^{-1}(\cos(x))}{a} + \frac{\cot(x)}{a + a \csc(x)}$$

Mathematica [B] time = 0.0528889, size = 44, normalized size = 2.2

$$\frac{\log\left(\sin\left(\frac{x}{2}\right)\right) - \log\left(\cos\left(\frac{x}{2}\right)\right) - \frac{2 \sin\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)}}{a}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[x]^2/(a + a*Csc[x]),x]

[Out] (-Log[Cos[x/2]] + Log[Sin[x/2]] - (2*Sin[x/2])/(Cos[x/2] + Sin[x/2]))/a

Maple [A] time = 0.027, size = 24, normalized size = 1.2

$$2 \frac{1}{a(\tan(x/2) + 1)} + \frac{1}{a} \ln\left(\tan\left(\frac{x}{2}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(x)^2/(a+a*csc(x)),x)

[Out] 2/a/(tan(1/2*x)+1)+1/a*ln(tan(1/2*x))

Maxima [A] time = 0.969735, size = 42, normalized size = 2.1

$$\frac{\log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a} + \frac{2}{a + \frac{a \sin(x)}{\cos(x)+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^2/(a+a*csc(x)),x, algorithm="maxima")

[Out] log(sin(x)/(cos(x) + 1))/a + 2/(a + a*sin(x)/(cos(x) + 1))

Fricas [B] time = 0.480746, size = 204, normalized size = 10.2

$$\frac{(\cos(x) + \sin(x) + 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x) + \sin(x) + 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) - 2 \cos(x) + 2 \sin(x) - 2}{2(a \cos(x) + a \sin(x) + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^2/(a+a*csc(x)),x, algorithm="fricas")

[Out] -1/2*((cos(x) + sin(x) + 1)*log(1/2*cos(x) + 1/2) - (cos(x) + sin(x) + 1)*log(-1/2*cos(x) + 1/2) - 2*cos(x) + 2*sin(x) - 2)/(a*cos(x) + a*sin(x) + a)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\frac{\int \frac{\csc^2(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)**2/(a+a*csc(x)),x)

[Out] Integral(csc(x)**2/(csc(x) + 1), x)/a

Giac [A] time = 1.36932, size = 32, normalized size = 1.6

$$\frac{\log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{a} + \frac{2}{a\left(\tan\left(\frac{1}{2}x\right) + 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^2/(a+a*csc(x)),x, algorithm="giac")


```
[Out] log(abs(tan(1/2*x)))/a + 2/(a*(tan(1/2*x) + 1))
```

$$3.5 \quad \int \frac{\csc(x)}{a+a \csc(x)} dx$$

Optimal. Leaf size=12

$$-\frac{\cot(x)}{a \csc(x) + a}$$

[Out] -(Cot[x]/(a + a*Csc[x]))

Rubi [A] time = 0.0198414, antiderivative size = 12, normalized size of antiderivative = 1., number of steps used = 1, number of rules used = 1, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {3794}

$$-\frac{\cot(x)}{a \csc(x) + a}$$

Antiderivative was successfully verified.

[In] Int[Csc[x]/(a + a*Csc[x]),x]

[Out] -(Cot[x]/(a + a*Csc[x]))

Rule 3794

Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] :> -Simp[Cot[e + f*x]/(f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]

Rubi steps

$$\int \frac{\csc(x)}{a+a \csc(x)} dx = -\frac{\cot(x)}{a+a \csc(x)}$$

Mathematica [B] time = 0.0239544, size = 26, normalized size = 2.17

$$\frac{2 \sin\left(\frac{x}{2}\right)}{a \left(\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[x]/(a + a*Csc[x]),x]

[Out] (2*Sin[x/2])/(a*(Cos[x/2] + Sin[x/2]))

Maple [A] time = 0.024, size = 14, normalized size = 1.2

$$-2 \frac{1}{a (\tan(x/2) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(x)/(a+a*csc(x)),x)

[Out] -2/a/(tan(1/2*x)+1)

Maxima [A] time = 1.01956, size = 22, normalized size = 1.83

$$-\frac{2}{a + \frac{a \sin(x)}{\cos(x)+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)/(a+a*csc(x)),x, algorithm="maxima")

[Out] -2/(a + a*sin(x)/(cos(x) + 1))

Fricas [A] time = 0.444167, size = 68, normalized size = 5.67

$$\frac{\cos(x) - \sin(x) + 1}{a \cos(x) + a \sin(x) + a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)/(a+a*csc(x)),x, algorithm="fricas")

[Out] $-(\cos(x) - \sin(x) + 1)/(a\cos(x) + a\sin(x) + a)$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\csc(x)}{\csc(x)+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(x)/(a+a*csc(x)),x)`

[Out] `Integral(csc(x)/(csc(x) + 1), x)/a`

Giac [A] time = 1.3368, size = 18, normalized size = 1.5

$$-\frac{2}{a\left(\tan\left(\frac{1}{2}x\right)+1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(x)/(a+a*csc(x)),x, algorithm="giac")`

[Out] `-2/(a*(tan(1/2*x) + 1))`

$$3.6 \quad \int \frac{1}{a+a \csc(c+dx)} dx$$

Optimal. Leaf size=28

$$\frac{\cot(c+dx)}{d(a \csc(c+dx)+a)} + \frac{x}{a}$$

[Out] x/a + Cot[c + d*x]/(d*(a + a*Csc[c + d*x]))

Rubi [A] time = 0.0126391, antiderivative size = 28, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3777, 8}

$$\frac{\cot(c+dx)}{d(a \csc(c+dx)+a)} + \frac{x}{a}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Csc[c + d*x])^(-1), x]

[Out] x/a + Cot[c + d*x]/(d*(a + a*Csc[c + d*x]))

Rule 3777

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^n], x_Symbol] :> -Simp[(Cot[c + d*x]*(a + b*Csc[c + d*x])^n)/(d*(2*n + 1)), x] + Dist[1/(a^2*(2*n + 1)), Int[(a + b*Csc[c + d*x])^(n + 1)*(a*(2*n + 1) - b*(n + 1)*Csc[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{a+a \csc(c+dx)} dx &= \frac{\cot(c+dx)}{d(a+a \csc(c+dx))} + \frac{\int a dx}{a^2} \\ &= \frac{x}{a} + \frac{\cot(c+dx)}{d(a+a \csc(c+dx))} \end{aligned}$$

Mathematica [A] time = 0.0894926, size = 47, normalized size = 1.68

$$\frac{\frac{2 \sin\left(\frac{1}{2}(c+dx)\right)}{\sin\left(\frac{1}{2}(c+dx)\right) + \cos\left(\frac{1}{2}(c+dx)\right)} + c + dx}{ad}$$

Antiderivative was successfully verified.

[In] Integrate[(a + a*Csc[c + d*x])^(-1), x]

[Out] (c + d*x - (2*Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])/(a*d)

Maple [A] time = 0.042, size = 41, normalized size = 1.5

$$2 \frac{\arctan(\tan(1/2 dx + c/2))}{ad} + 2 \frac{1}{ad(\tan(1/2 dx + c/2) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+a*csc(d*x+c)), x)

[Out] 2/a/d*arctan(tan(1/2*d*x+1/2*c))+2/a/d/(tan(1/2*d*x+1/2*c)+1)

Maxima [A] time = 1.47103, size = 68, normalized size = 2.43

$$\frac{2 \left(\frac{\arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a} + \frac{1}{a + \frac{a \sin(dx+c)}{\cos(dx+c)+1}} \right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c)), x, algorithm="maxima")

[Out] 2*(arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a + 1/(a + a*sin(d*x + c)/(cos(d*x + c) + 1)))/d

Fricas [A] time = 0.467268, size = 142, normalized size = 5.07

$$\frac{dx + (dx + 1) \cos(dx + c) + (dx - 1) \sin(dx + c) + 1}{ad \cos(dx + c) + ad \sin(dx + c) + ad}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c)),x, algorithm="fricas")

[Out] (d*x + (d*x + 1)*cos(d*x + c) + (d*x - 1)*sin(d*x + c) + 1)/(a*d*cos(d*x + c) + a*d*sin(d*x + c) + a*d)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{\csc(c+dx)+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c)),x)

[Out] Integral(1/(csc(c + d*x) + 1), x)/a

Giac [A] time = 1.2772, size = 43, normalized size = 1.54

$$\frac{\frac{dx+c}{a} + \frac{2}{a\left(\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)+1\right)}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c)),x, algorithm="giac")

[Out] ((d*x + c)/a + 2/(a*(tan(1/2*d*x + 1/2*c) + 1)))/d

$$3.7 \quad \int \frac{\sin(x)}{a+a \csc(x)} dx$$

Optimal. Leaf size=25

$$-\frac{x}{a} - \frac{2 \cos(x)}{a} + \frac{\cos(x)}{a \csc(x) + a}$$

[Out] $-(x/a) - (2*\text{Cos}[x])/a + \text{Cos}[x]/(a + a*\text{Csc}[x])$

Rubi [A] time = 0.0468814, antiderivative size = 25, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3819, 3787, 2638, 8}

$$-\frac{x}{a} - \frac{2 \cos(x)}{a} + \frac{\cos(x)}{a \csc(x) + a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[x]/(a + a*\text{Csc}[x]), x]$

[Out] $-(x/a) - (2*\text{Cos}[x])/a + \text{Cos}[x]/(a + a*\text{Csc}[x])$

Rule 3819

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(d_.))^n]/(\csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[(\text{Cot}[e + f*x]*(d*\text{Csc}[e + f*x])^n)/(f*(a + b*\text{Csc}[e + f*x])), x] - \text{Dist}[1/a^2, \text{Int}[(d*\text{Csc}[e + f*x])^n*(a*(n - 1) - b*n*\text{Csc}[e + f*x]), x], x] /; \text{FreeQ}\{a, b, d, e, f\}, x] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{LtQ}[n, 0]$

Rule 3787

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(d_.))^n*(\csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Dist}[b/d, \text{Int}[(d*\text{Csc}[e + f*x])^{n+1}, x], x] /; \text{FreeQ}\{a, b, d, e, f, n\}, x]$

Rule 2638

$\text{Int}[\sin[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow -\text{Simp}[\text{Cos}[c + d*x]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rubi steps

$$\begin{aligned} \int \frac{\sin(x)}{a + a \csc(x)} dx &= \frac{\cos(x)}{a + a \csc(x)} - \frac{\int (-2a + a \csc(x)) \sin(x) dx}{a^2} \\ &= \frac{\cos(x)}{a + a \csc(x)} - \frac{\int 1 dx}{a} + \frac{2 \int \sin(x) dx}{a} \\ &= -\frac{x}{a} - \frac{2 \cos(x)}{a} + \frac{\cos(x)}{a + a \csc(x)} \end{aligned}$$

Mathematica [A] time = 0.077861, size = 32, normalized size = 1.28

$$-\frac{x + \cos(x) - \frac{2 \sin\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)}}{a}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]/(a + a*Csc[x]),x]

[Out] -((x + Cos[x] - (2*Sin[x/2]))/(Cos[x/2] + Sin[x/2]))/a

Maple [A] time = 0.043, size = 40, normalized size = 1.6

$$-2 \frac{1}{a \left((\tan(x/2))^2 + 1 \right)} - 2 \frac{\arctan(\tan(x/2))}{a} - 2 \frac{1}{a (\tan(x/2) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)/(a+a*csc(x)),x)

[Out] -2/a/(tan(1/2*x)^2+1)-2/a*arctan(tan(1/2*x))-2/a/(tan(1/2*x)+1)

Maxima [B] time = 1.45752, size = 105, normalized size = 4.2

$$\frac{2 \left(\frac{\sin(x)}{\cos(x)+1} + \frac{\sin(x)^2}{(\cos(x)+1)^2} + 2 \right)}{a + \frac{a \sin(x)}{\cos(x)+1} + \frac{a \sin(x)^2}{(\cos(x)+1)^2} + \frac{a \sin(x)^3}{(\cos(x)+1)^3}} - \frac{2 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)/(a+a*csc(x)),x, algorithm="maxima")

[Out] -2*(sin(x)/(cos(x) + 1) + sin(x)^2/(cos(x) + 1)^2 + 2)/(a + a*sin(x)/(cos(x) + 1) + a*sin(x)^2/(cos(x) + 1)^2 + a*sin(x)^3/(cos(x) + 1)^3) - 2*arctan(sin(x)/(cos(x) + 1))/a

Fricas [A] time = 0.472957, size = 122, normalized size = 4.88

$$\frac{(x + 2) \cos(x) + \cos(x)^2 + (x + \cos(x) - 1) \sin(x) + x + 1}{a \cos(x) + a \sin(x) + a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)/(a+a*csc(x)),x, algorithm="fricas")

[Out] -((x + 2)*cos(x) + cos(x)^2 + (x + cos(x) - 1)*sin(x) + x + 1)/(a*cos(x) + a*sin(x) + a)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\frac{\int \frac{\sin(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)/(a+a*csc(x)),x)

[Out] Integral(sin(x)/(csc(x) + 1), x)/a

Giac [A] time = 1.33014, size = 59, normalized size = 2.36

$$-\frac{x}{a} - \frac{2 \left(\tan\left(\frac{1}{2}x\right)^2 + \tan\left(\frac{1}{2}x\right) + 2 \right)}{\left(\tan\left(\frac{1}{2}x\right)^3 + \tan\left(\frac{1}{2}x\right)^2 + \tan\left(\frac{1}{2}x\right) + 1 \right) a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)/(a+a*csc(x)),x, algorithm="giac")

[Out] -x/a - 2*(tan(1/2*x)^2 + tan(1/2*x) + 2)/((tan(1/2*x)^3 + tan(1/2*x)^2 + tan(1/2*x) + 1)*a)

$$3.8 \quad \int \frac{\sin^2(x)}{a+a \csc(x)} dx$$

Optimal. Leaf size=40

$$\frac{3x}{2a} + \frac{2 \cos(x)}{a} - \frac{3 \sin(x) \cos(x)}{2a} + \frac{\sin(x) \cos(x)}{a \csc(x) + a}$$

[Out] (3*x)/(2*a) + (2*Cos[x])/a - (3*Cos[x]*Sin[x])/(2*a) + (Cos[x]*Sin[x])/(a + a*Csc[x])

Rubi [A] time = 0.0612801, antiderivative size = 40, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3819, 3787, 2635, 8, 2638}

$$\frac{3x}{2a} + \frac{2 \cos(x)}{a} - \frac{3 \sin(x) \cos(x)}{2a} + \frac{\sin(x) \cos(x)}{a \csc(x) + a}$$

Antiderivative was successfully verified.

[In] Int[Sin[x]^2/(a + a*Csc[x]),x]

[Out] (3*x)/(2*a) + (2*Cos[x])/a - (3*Cos[x]*Sin[x])/(2*a) + (Cos[x]*Sin[x])/(a + a*Csc[x])

Rule 3819

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[(Cot[e + f*x]*(d*Csc[e + f*x])^n)/(f*(a + b*Csc[e + f*x])), x] - Dist[1/a^2, Int[(d*Csc[e + f*x])^n*(a*(n - 1) - b*n*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, 0]

Rule 3787

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Dist[a, Int[(d*Csc[e + f*x])^n, x], x] + Dist[b/d, Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^n], x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c

+ d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2638

Int[sin[(c_.) + (d_.)*(x_.)], x_Symbol] := -Simp[Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sin^2(x)}{a + a \csc(x)} dx &= \frac{\cos(x) \sin(x)}{a + a \csc(x)} - \frac{\int (-3a + 2a \csc(x)) \sin^2(x) dx}{a^2} \\ &= \frac{\cos(x) \sin(x)}{a + a \csc(x)} - \frac{2 \int \sin(x) dx}{a} + \frac{3 \int \sin^2(x) dx}{a} \\ &= \frac{2 \cos(x)}{a} - \frac{3 \cos(x) \sin(x)}{2a} + \frac{\cos(x) \sin(x)}{a + a \csc(x)} + \frac{3 \int 1 dx}{2a} \\ &= \frac{3x}{2a} + \frac{2 \cos(x)}{a} - \frac{3 \cos(x) \sin(x)}{2a} + \frac{\cos(x) \sin(x)}{a + a \csc(x)} \end{aligned}$$

Mathematica [A] time = 0.129952, size = 42, normalized size = 1.05

$$-\frac{-6x + \sin(2x) - 4 \cos(x) + \frac{8 \sin\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)}}{4a}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]^2/(a + a*Csc[x]),x]

[Out] -(-6*x - 4*Cos[x] + (8*Sin[x/2]))/(Cos[x/2] + Sin[x/2]) + Sin[2*x]/(4*a)

Maple [B] time = 0.042, size = 100, normalized size = 2.5

$$\frac{1}{a} \left(\tan\left(\frac{x}{2}\right) \right)^3 \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-2} + 2 \frac{(\tan(x/2))^2}{a \left((\tan(x/2))^2 + 1 \right)^2} - \frac{1}{a} \tan\left(\frac{x}{2}\right) \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-2} + 2 \frac{1}{a \left((\tan(x/2))^2 + 1 \right)^2} +$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(x)^2/(a+a*csc(x)),x)`

[Out] $1/a/(\tan(1/2*x)^2+1)^2*\tan(1/2*x)^3+2/a/(\tan(1/2*x)^2+1)^2*\tan(1/2*x)^2-1/a/(\tan(1/2*x)^2+1)^2*\tan(1/2*x)+2/a/(\tan(1/2*x)^2+1)^2+3/a*\arctan(\tan(1/2*x))+2/a/(\tan(1/2*x)+1)$

Maxima [B] time = 1.45891, size = 173, normalized size = 4.32

$$\frac{\frac{\sin(x)}{\cos(x)+1} + \frac{5 \sin(x)^2}{(\cos(x)+1)^2} + \frac{3 \sin(x)^3}{(\cos(x)+1)^3} + \frac{3 \sin(x)^4}{(\cos(x)+1)^4} + 4}{a + \frac{a \sin(x)}{\cos(x)+1} + \frac{2 a \sin(x)^2}{(\cos(x)+1)^2} + \frac{2 a \sin(x)^3}{(\cos(x)+1)^3} + \frac{a \sin(x)^4}{(\cos(x)+1)^4} + \frac{a \sin(x)^5}{(\cos(x)+1)^5}} + \frac{3 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^2/(a+a*csc(x)),x, algorithm="maxima")`

[Out] $(\sin(x)/(\cos(x) + 1) + 5*\sin(x)^2/(\cos(x) + 1)^2 + 3*\sin(x)^3/(\cos(x) + 1)^3 + 3*\sin(x)^4/(\cos(x) + 1)^4 + 4)/(a + a*\sin(x)/(\cos(x) + 1) + 2*a*\sin(x)^2/(\cos(x) + 1)^2 + 2*a*\sin(x)^3/(\cos(x) + 1)^3 + a*\sin(x)^4/(\cos(x) + 1)^4 + a*\sin(x)^5/(\cos(x) + 1)^5) + 3*\arctan(\sin(x)/(\cos(x) + 1)))/a$

Fricas [A] time = 0.474, size = 166, normalized size = 4.15

$$\frac{\cos(x)^3 + 3(x+1)\cos(x) + 2\cos(x)^2 - (\cos(x)^2 - 3x - \cos(x) + 2)\sin(x) + 3x + 2}{2(a\cos(x) + a\sin(x) + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^2/(a+a*csc(x)),x, algorithm="fricas")`

[Out] $1/2*(\cos(x)^3 + 3*(x + 1)*\cos(x) + 2*\cos(x)^2 - (\cos(x)^2 - 3*x - \cos(x) + 2)*\sin(x) + 3*x + 2)/(a*\cos(x) + a*\sin(x) + a)$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sin^2(x)}{\csc(x)+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)**2/(a+a*csc(x)),x)

[Out] Integral(sin(x)**2/(csc(x) + 1), x)/a

Giac [A] time = 1.36766, size = 76, normalized size = 1.9

$$\frac{3x}{2a} + \frac{\tan\left(\frac{1}{2}x\right)^3 + 2\tan\left(\frac{1}{2}x\right)^2 - \tan\left(\frac{1}{2}x\right) + 2}{\left(\tan\left(\frac{1}{2}x\right)^2 + 1\right)^2 a} + \frac{2}{a\left(\tan\left(\frac{1}{2}x\right) + 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^2/(a+a*csc(x)),x, algorithm="giac")

[Out] 3/2*x/a + (tan(1/2*x)^3 + 2*tan(1/2*x)^2 - tan(1/2*x) + 2)/((tan(1/2*x)^2 + 1)^2*a) + 2/(a*(tan(1/2*x) + 1))

3.9 $\int \frac{\sin^3(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=53

$$-\frac{3x}{2a} + \frac{4 \cos^3(x)}{3a} - \frac{4 \cos(x)}{a} + \frac{3 \sin(x) \cos(x)}{2a} + \frac{\sin^2(x) \cos(x)}{a \csc(x) + a}$$

[Out] $(-3*x)/(2*a) - (4*\text{Cos}[x])/a + (4*\text{Cos}[x]^3)/(3*a) + (3*\text{Cos}[x]*\text{Sin}[x])/(2*a) + (\text{Cos}[x]*\text{Sin}[x]^2)/(a + a*\text{Csc}[x])$

Rubi [A] time = 0.0670422, antiderivative size = 53, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 5, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3819, 3787, 2633, 2635, 8}

$$-\frac{3x}{2a} + \frac{4 \cos^3(x)}{3a} - \frac{4 \cos(x)}{a} + \frac{3 \sin(x) \cos(x)}{2a} + \frac{\sin^2(x) \cos(x)}{a \csc(x) + a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[x]^3/(a + a*\text{Csc}[x]), x]$

[Out] $(-3*x)/(2*a) - (4*\text{Cos}[x])/a + (4*\text{Cos}[x]^3)/(3*a) + (3*\text{Cos}[x]*\text{Sin}[x])/(2*a) + (\text{Cos}[x]*\text{Sin}[x]^2)/(a + a*\text{Csc}[x])$

Rule 3819

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^n/(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[(\text{Cot}[e + f*x]*(d*\text{Csc}[e + f*x])^n)/(f*(a + b*\text{Csc}[e + f*x])), x] - \text{Dist}[1/a^2, \text{Int}[(d*\text{Csc}[e + f*x])^n*(a*(n - 1) - b*n*\text{Csc}[e + f*x]), x], x] /;$ $\text{FreeQ}\{a, b, d, e, f\}, x \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, 0]$

Rule 3787

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^n*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Dist}[b/d, \text{Int}[(d*\text{Csc}[e + f*x])^{n+1}, x], x] /;$ $\text{FreeQ}\{a, b, d, e, f, n\}, x]$

Rule 2633

$\text{Int}[\text{sin}[(c_.) + (d_.)*(x_.)]^n, x_Symbol] \rightarrow -\text{Dist}[d^{-1}, \text{Subst}[\text{Int}[\text{Expand}[(1 - x^2)^{(n-1)/2}, x], x], x, \text{Cos}[c + d*x]], x] /;$ $\text{FreeQ}\{c, d\}, x]$

&& IGtQ[(n - 1)/2, 0]

Rule 2635

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]
)*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c
+ d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
]
```

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(x)}{a + a \csc(x)} dx &= \frac{\cos(x) \sin^2(x)}{a + a \csc(x)} - \frac{\int (-4a + 3a \csc(x)) \sin^3(x) dx}{a^2} \\ &= \frac{\cos(x) \sin^2(x)}{a + a \csc(x)} - \frac{3 \int \sin^2(x) dx}{a} + \frac{4 \int \sin^3(x) dx}{a} \\ &= \frac{3 \cos(x) \sin(x)}{2a} + \frac{\cos(x) \sin^2(x)}{a + a \csc(x)} - \frac{3 \int 1 dx}{2a} - \frac{4 \text{Subst}\left(\int (1 - x^2) dx, x, \cos(x)\right)}{a} \\ &= -\frac{3x}{2a} - \frac{4 \cos(x)}{a} + \frac{4 \cos^3(x)}{3a} + \frac{3 \cos(x) \sin(x)}{2a} + \frac{\cos(x) \sin^2(x)}{a + a \csc(x)} \end{aligned}$$

Mathematica [A] time = 0.16353, size = 49, normalized size = 0.92

$$\frac{-21 \cos(x) + \cos(3x) + 3 \left(-6x + \sin(2x) + \frac{8 \sin\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)} \right)}{12a}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[x]^3/(a + a*Csc[x]),x]
```

```
[Out] (-21*Cos[x] + Cos[3*x] + 3*(-6*x + (8*Sin[x/2])/(Cos[x/2] + Sin[x/2]) + Sin[2*x]))/(12*a)
```

Maple [B] time = 0.047, size = 121, normalized size = 2.3

$$-\frac{1}{a} \left(\tan\left(\frac{x}{2}\right) \right)^5 \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-3} - 2 \frac{(\tan(x/2))^4}{a \left((\tan(x/2))^2 + 1 \right)^3} - 8 \frac{(\tan(x/2))^2}{a \left((\tan(x/2))^2 + 1 \right)^3} + \frac{1}{a} \tan\left(\frac{x}{2}\right) \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-3} - \dots$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)^3/(a+a*csc(x)),x)

[Out] -1/a/(tan(1/2*x)^2+1)^3*tan(1/2*x)^5-2/a/(tan(1/2*x)^2+1)^3*tan(1/2*x)^4-8/a/(tan(1/2*x)^2+1)^3*tan(1/2*x)^2+1/a/(tan(1/2*x)^2+1)^3*tan(1/2*x)-10/3/a/(tan(1/2*x)^2+1)^3-3/a*arctan(tan(1/2*x))-2/a/(tan(1/2*x)+1)

Maxima [B] time = 1.46401, size = 243, normalized size = 4.58

$$\frac{\frac{7 \sin(x)}{\cos(x)+1} + \frac{39 \sin(x)^2}{(\cos(x)+1)^2} + \frac{24 \sin(x)^3}{(\cos(x)+1)^3} + \frac{24 \sin(x)^4}{(\cos(x)+1)^4} + \frac{9 \sin(x)^5}{(\cos(x)+1)^5} + \frac{9 \sin(x)^6}{(\cos(x)+1)^6} + 16}{3 \left(a + \frac{a \sin(x)}{\cos(x)+1} + \frac{3 a \sin(x)^2}{(\cos(x)+1)^2} + \frac{3 a \sin(x)^3}{(\cos(x)+1)^3} + \frac{3 a \sin(x)^4}{(\cos(x)+1)^4} + \frac{3 a \sin(x)^5}{(\cos(x)+1)^5} + \frac{a \sin(x)^6}{(\cos(x)+1)^6} + \frac{a \sin(x)^7}{(\cos(x)+1)^7} \right)} - \frac{3 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^3/(a+a*csc(x)),x, algorithm="maxima")

[Out] -1/3*(7*sin(x)/(cos(x) + 1) + 39*sin(x)^2/(cos(x) + 1)^2 + 24*sin(x)^3/(cos(x) + 1)^3 + 24*sin(x)^4/(cos(x) + 1)^4 + 9*sin(x)^5/(cos(x) + 1)^5 + 9*sin(x)^6/(cos(x) + 1)^6 + 16)/(a + a*sin(x)/(cos(x) + 1) + 3*a*sin(x)^2/(cos(x) + 1)^2 + 3*a*sin(x)^3/(cos(x) + 1)^3 + 3*a*sin(x)^4/(cos(x) + 1)^4 + 3*a*sin(x)^5/(cos(x) + 1)^5 + a*sin(x)^6/(cos(x) + 1)^6 + a*sin(x)^7/(cos(x) + 1)^7) - 3*arctan(sin(x)/(cos(x) + 1))/a

Fricas [A] time = 0.482549, size = 211, normalized size = 3.98

$$\frac{2 \cos(x)^4 - \cos(x)^3 - 3(3x + 5) \cos(x) - 12 \cos(x)^2 + (2 \cos(x)^3 + 3 \cos(x)^2 - 9x - 9 \cos(x) + 6) \sin(x) - 9x - 6}{6(a \cos(x) + a \sin(x) + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^3/(a+a*csc(x)),x, algorithm="fricas")

[Out] $\frac{1}{6}(2\cos(x)^4 - \cos(x)^3 - 3(3x + 5)\cos(x) - 12\cos(x)^2 + (2\cos(x)^3 + 3\cos(x)^2 - 9x - 9\cos(x) + 6)\sin(x) - 9x - 6)/(a\cos(x) + a\sin(x) + a)$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sin^3(x)}{\csc(x)+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)**3/(a+a*csc(x)),x)`

[Out] `Integral(sin(x)**3/(csc(x) + 1), x)/a`

Giac [A] time = 1.27658, size = 90, normalized size = 1.7

$$\frac{3x}{2a} - \frac{2}{a\left(\tan\left(\frac{1}{2}x\right) + 1\right)} - \frac{3\tan\left(\frac{1}{2}x\right)^5 + 6\tan\left(\frac{1}{2}x\right)^4 + 24\tan\left(\frac{1}{2}x\right)^2 - 3\tan\left(\frac{1}{2}x\right) + 10}{3\left(\tan\left(\frac{1}{2}x\right)^2 + 1\right)^3 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^3/(a+a*csc(x)),x, algorithm="giac")`

[Out] $-3/2*x/a - 2/(a*(\tan(1/2*x) + 1)) - 1/3*(3*\tan(1/2*x)^5 + 6*\tan(1/2*x)^4 + 24*\tan(1/2*x)^2 - 3*\tan(1/2*x) + 10)/((\tan(1/2*x)^2 + 1)^3*a)$

3.10 $\int \frac{\sin^4(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=66

$$\frac{15x}{8a} - \frac{4 \cos^3(x)}{3a} + \frac{4 \cos(x)}{a} - \frac{5 \sin^3(x) \cos(x)}{4a} - \frac{15 \sin(x) \cos(x)}{8a} + \frac{\sin^3(x) \cos(x)}{a \csc(x) + a}$$

[Out] (15*x)/(8*a) + (4*Cos[x])/a - (4*Cos[x]^3)/(3*a) - (15*Cos[x]*Sin[x])/(8*a) - (5*Cos[x]*Sin[x]^3)/(4*a) + (Cos[x]*Sin[x]^3)/(a + a*Csc[x])

Rubi [A] time = 0.0733532, antiderivative size = 66, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 5, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3819, 3787, 2635, 8, 2633}

$$\frac{15x}{8a} - \frac{4 \cos^3(x)}{3a} + \frac{4 \cos(x)}{a} - \frac{5 \sin^3(x) \cos(x)}{4a} - \frac{15 \sin(x) \cos(x)}{8a} + \frac{\sin^3(x) \cos(x)}{a \csc(x) + a}$$

Antiderivative was successfully verified.

[In] Int[Sin[x]^4/(a + a*Csc[x]),x]

[Out] (15*x)/(8*a) + (4*Cos[x])/a - (4*Cos[x]^3)/(3*a) - (15*Cos[x]*Sin[x])/(8*a) - (5*Cos[x]*Sin[x]^3)/(4*a) + (Cos[x]*Sin[x]^3)/(a + a*Csc[x])

Rule 3819

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n_]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[(Cot[e + f*x]*(d*Csc[e + f*x])^n)/(f*(a + b*Csc[e + f*x])), x] - Dist[1/a^2, Int[(d*Csc[e + f*x])^n*(a*(n - 1) - b*n*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, 0]

Rule 3787

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n_]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Dist[a, Int[(d*Csc[e + f*x])^n, x], x] + Dist[b/d, Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c

+ d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned}
 \int \frac{\sin^4(x)}{a + a \csc(x)} dx &= \frac{\cos(x) \sin^3(x)}{a + a \csc(x)} - \frac{\int (-5a + 4a \csc(x)) \sin^4(x) dx}{a^2} \\
 &= \frac{\cos(x) \sin^3(x)}{a + a \csc(x)} - \frac{4 \int \sin^3(x) dx}{a} + \frac{5 \int \sin^4(x) dx}{a} \\
 &= -\frac{5 \cos(x) \sin^3(x)}{4a} + \frac{\cos(x) \sin^3(x)}{a + a \csc(x)} + \frac{15 \int \sin^2(x) dx}{4a} + \frac{4 \text{Subst}\left(\int (1 - x^2) dx, x, \cos(x)\right)}{a} \\
 &= \frac{4 \cos(x)}{a} - \frac{4 \cos^3(x)}{3a} - \frac{15 \cos(x) \sin(x)}{8a} - \frac{5 \cos(x) \sin^3(x)}{4a} + \frac{\cos(x) \sin^3(x)}{a + a \csc(x)} + \frac{15 \int 1 dx}{8a} \\
 &= \frac{15x}{8a} + \frac{4 \cos(x)}{a} - \frac{4 \cos^3(x)}{3a} - \frac{15 \cos(x) \sin(x)}{8a} - \frac{5 \cos(x) \sin^3(x)}{4a} + \frac{\cos(x) \sin^3(x)}{a + a \csc(x)}
 \end{aligned}$$

Mathematica [A] time = 0.20905, size = 57, normalized size = 0.86

$$\frac{168 \cos(x) - 8 \cos(3x) + 3 \left(60x - 16 \sin(2x) + \sin(4x) - \frac{64 \sin\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)} \right)}{96a}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]^4/(a + a*Csc[x]),x]

[Out] (168*Cos[x] - 8*Cos[3*x] + 3*(60*x - (64*Sin[x/2])/(Cos[x/2] + Sin[x/2]) - 16*Sin[2*x] + Sin[4*x]))/(96*a)

Maple [B] time = 0.046, size = 185, normalized size = 2.8

$$\frac{7}{4a} \left(\tan\left(\frac{x}{2}\right) \right)^7 \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-4} + 2 \frac{(\tan(x/2))^6}{a \left((\tan(x/2))^2 + 1 \right)^4} + \frac{15}{4a} \left(\tan\left(\frac{x}{2}\right) \right)^5 \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-4} + 10 \frac{(\tan(x/2))^4}{a \left((\tan(x/2))^2 + 1 \right)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)^4/(a+a*csc(x)),x)

[Out] 7/4/a/(tan(1/2*x)^2+1)^4*tan(1/2*x)^7+2/a/(tan(1/2*x)^2+1)^4*tan(1/2*x)^6+15/4/a/(tan(1/2*x)^2+1)^4*tan(1/2*x)^5+10/a/(tan(1/2*x)^2+1)^4*tan(1/2*x)^4-15/4/a/(tan(1/2*x)^2+1)^4*tan(1/2*x)^3+34/3/a/(tan(1/2*x)^2+1)^4*tan(1/2*x)^2-7/4/a/(tan(1/2*x)^2+1)^4*tan(1/2*x)+10/3/a/(tan(1/2*x)^2+1)^4+15/4/a*arc tan(tan(1/2*x))+2/a/(tan(1/2*x)+1)

Maxima [B] time = 1.47195, size = 311, normalized size = 4.71

$$\frac{\frac{19 \sin(x)}{\cos(x)+1} + \frac{211 \sin(x)^2}{(\cos(x)+1)^2} + \frac{91 \sin(x)^3}{(\cos(x)+1)^3} + \frac{219 \sin(x)^4}{(\cos(x)+1)^4} + \frac{165 \sin(x)^5}{(\cos(x)+1)^5} + \frac{165 \sin(x)^6}{(\cos(x)+1)^6} + \frac{45 \sin(x)^7}{(\cos(x)+1)^7} + \frac{45 \sin(x)^8}{(\cos(x)+1)^8} + 64}{12 \left(a + \frac{a \sin(x)}{\cos(x)+1} + \frac{4a \sin(x)^2}{(\cos(x)+1)^2} + \frac{4a \sin(x)^3}{(\cos(x)+1)^3} + \frac{6a \sin(x)^4}{(\cos(x)+1)^4} + \frac{6a \sin(x)^5}{(\cos(x)+1)^5} + \frac{4a \sin(x)^6}{(\cos(x)+1)^6} + \frac{4a \sin(x)^7}{(\cos(x)+1)^7} + \frac{a \sin(x)^8}{(\cos(x)+1)^8} + \frac{a \sin(x)^9}{(\cos(x)+1)^9} \right)} + \frac{15}{\cos(x)+1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^4/(a+a*csc(x)),x, algorithm="maxima")

[Out] 1/12*(19*sin(x)/(cos(x) + 1) + 211*sin(x)^2/(cos(x) + 1)^2 + 91*sin(x)^3/(cos(x) + 1)^3 + 219*sin(x)^4/(cos(x) + 1)^4 + 165*sin(x)^5/(cos(x) + 1)^5 + 165*sin(x)^6/(cos(x) + 1)^6 + 45*sin(x)^7/(cos(x) + 1)^7 + 45*sin(x)^8/(cos(x) + 1)^8 + 64)/(a + a*sin(x)/(cos(x) + 1) + 4*a*sin(x)^2/(cos(x) + 1)^2 + 4*a*sin(x)^3/(cos(x) + 1)^3 + 6*a*sin(x)^4/(cos(x) + 1)^4 + 6*a*sin(x)^5/(cos(x) + 1)^5 + 4*a*sin(x)^6/(cos(x) + 1)^6 + 4*a*sin(x)^7/(cos(x) + 1)^7 + a*sin(x)^8/(cos(x) + 1)^8 + a*sin(x)^9/(cos(x) + 1)^9) + 15/4*arctan(sin(x)/(cos(x) + 1))/a

Fricas [A] time = 0.485014, size = 259, normalized size = 3.92

$$\frac{6 \cos(x)^5 + 8 \cos(x)^4 - 25 \cos(x)^3 - 45(x+1) \cos(x) - 48 \cos(x)^2 - (6 \cos(x)^4 - 2 \cos(x)^3 - 27 \cos(x)^2 + 45x + 15)}{24(a \cos(x) + a \sin(x) + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^4/(a+a*csc(x)),x, algorithm="fricas")

[Out]
$$\frac{-1/24*(6*\cos(x)^5 + 8*\cos(x)^4 - 25*\cos(x)^3 - 45*(x + 1)*\cos(x) - 48*\cos(x)^2 - (6*\cos(x)^4 - 2*\cos(x)^3 - 27*\cos(x)^2 + 45*x + 21*\cos(x) - 24)*\sin(x) - 45*x - 24)/(a*\cos(x) + a*\sin(x) + a)}$$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sin^4(x)}{\csc(x)+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)**4/(a+a*csc(x)),x)

[Out] Integral(sin(x)**4/(csc(x) + 1), x)/a

Giac [A] time = 1.39204, size = 123, normalized size = 1.86

$$\frac{15x}{8a} + \frac{2}{a\left(\tan\left(\frac{1}{2}x\right) + 1\right)} + \frac{21 \tan\left(\frac{1}{2}x\right)^7 + 24 \tan\left(\frac{1}{2}x\right)^6 + 45 \tan\left(\frac{1}{2}x\right)^5 + 120 \tan\left(\frac{1}{2}x\right)^4 - 45 \tan\left(\frac{1}{2}x\right)^3 + 136 \tan\left(\frac{1}{2}x\right)^2 - 21 \tan\left(\frac{1}{2}x\right) + 40}{12\left(\tan\left(\frac{1}{2}x\right)^2 + 1\right)^4 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^4/(a+a*csc(x)),x, algorithm="giac")

[Out]
$$15/8*x/a + 2/(a*(\tan(1/2*x) + 1)) + 1/12*(21*\tan(1/2*x)^7 + 24*\tan(1/2*x)^6 + 45*\tan(1/2*x)^5 + 120*\tan(1/2*x)^4 - 45*\tan(1/2*x)^3 + 136*\tan(1/2*x)^2 - 21*\tan(1/2*x) + 40)/((\tan(1/2*x)^2 + 1)^4*a)$$

$$3.11 \quad \int \frac{1}{(a+a \csc(c+dx))^2} dx$$

Optimal. Leaf size=57

$$\frac{4 \cot(c+dx)}{3a^2d(\csc(c+dx)+1)} + \frac{x}{a^2} + \frac{\cot(c+dx)}{3d(a \csc(c+dx)+a)^2}$$

[Out] x/a^2 + (4*Cot[c + d*x])/(3*a^2*d*(1 + Csc[c + d*x])) + Cot[c + d*x]/(3*d*(a + a*Csc[c + d*x])^2)

Rubi [A] time = 0.066061, antiderivative size = 57, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$, Rules used = {3777, 3919, 3794}

$$\frac{4 \cot(c+dx)}{3a^2d(\csc(c+dx)+1)} + \frac{x}{a^2} + \frac{\cot(c+dx)}{3d(a \csc(c+dx)+a)^2}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Csc[c + d*x])^(-2), x]

[Out] x/a^2 + (4*Cot[c + d*x])/(3*a^2*d*(1 + Csc[c + d*x])) + Cot[c + d*x]/(3*d*(a + a*Csc[c + d*x])^2)

Rule 3777

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^n, x_Symbol] :> -Simp[(Cot[c + d*x]*(a + b*Csc[c + d*x])^n)/(d*(2*n + 1)), x] + Dist[1/(a^2*(2*n + 1)), Int[(a + b*Csc[c + d*x])^(n + 1)*(a*(2*n + 1) - b*(n + 1)*Csc[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 3919

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Simp[(c*x)/a, x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]

Rule 3794


```
Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol]
:= -Simp[Cot[e + f*x]/(f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x]
&& EqQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + a \csc(c + dx))^2} dx &= \frac{\cot(c + dx)}{3d(a + a \csc(c + dx))^2} - \frac{\int \frac{-3a + a \csc(c + dx)}{a + a \csc(c + dx)} dx}{3a^2} \\ &= \frac{x}{a^2} + \frac{\cot(c + dx)}{3d(a + a \csc(c + dx))^2} - \frac{4 \int \frac{\csc(c + dx)}{a + a \csc(c + dx)} dx}{3a} \\ &= \frac{x}{a^2} + \frac{\cot(c + dx)}{3d(a + a \csc(c + dx))^2} + \frac{4 \cot(c + dx)}{3d(a^2 + a^2 \csc(c + dx))} \end{aligned}$$

Mathematica [A] time = 0.324959, size = 108, normalized size = 1.89

$$\frac{3(3c + 3dx - 4) \cos\left(\frac{1}{2}(c + dx)\right) + (-3c - 3dx + 10) \cos\left(\frac{3}{2}(c + dx)\right) + 6 \sin\left(\frac{1}{2}(c + dx)\right) ((c + dx) \cos(c + dx) + 2c + 2dx)}{6a^2d \left(\sin\left(\frac{1}{2}(c + dx)\right) + \cos\left(\frac{1}{2}(c + dx)\right)\right)^3}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + a*Csc[c + d*x])^(-2), x]
```

```
[Out] (3*(-4 + 3*c + 3*d*x)*Cos[(c + d*x)/2] + (10 - 3*c - 3*d*x)*Cos[(3*(c + d*x))/2] + 6*(-3 + 2*c + 2*d*x + (c + d*x)*Cos[c + d*x])*Sin[(c + d*x)/2])/(6*a^2*d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^3)
```

Maple [A] time = 0.063, size = 83, normalized size = 1.5

$$2 \frac{\arctan(\tan(1/2 dx + c/2))}{da^2} - \frac{4}{3da^2} \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)^{-3} + 2 \frac{1}{da^2 (\tan(1/2 dx + c/2) + 1)^2} + 2 \frac{1}{da^2 (\tan(1/2 dx + c/2) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(a+a*csc(d*x+c))^2,x)
```

[Out] $2/d/a^2 \arctan(\tan(1/2*d*x+1/2*c)) - 4/3/d/a^2/(\tan(1/2*d*x+1/2*c)+1)^3 + 2/d/a^2/(\tan(1/2*d*x+1/2*c)+1)^2 + 2/d/a^2/(\tan(1/2*d*x+1/2*c)+1)$

Maxima [B] time = 1.47041, size = 192, normalized size = 3.37

$$2 \left(\frac{\frac{9 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 4}{a^2 + \frac{3a^2 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a^2 \sin(dx+c)^3}{(\cos(dx+c)+1)^3}} + \frac{3 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^2} \right) / 3d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c))^2,x, algorithm="maxima")

[Out] $2/3 * ((9 * \sin(d*x + c) / (\cos(d*x + c) + 1) + 3 * \sin(d*x + c)^2 / (\cos(d*x + c) + 1)^2 + 4) / (a^2 + 3 * a^2 * \sin(d*x + c) / (\cos(d*x + c) + 1) + 3 * a^2 * \sin(d*x + c)^2 / (\cos(d*x + c) + 1)^2 + a^2 * \sin(d*x + c)^3 / (\cos(d*x + c) + 1)^3) + 3 * \arctan(\sin(d*x + c) / (\cos(d*x + c) + 1)) / a^2) / d$

Fricas [B] time = 0.480864, size = 306, normalized size = 5.37

$$\frac{(3dx - 5) \cos(dx + c)^2 - 6dx - (3dx + 4) \cos(dx + c) - (6dx + (3dx + 5) \cos(dx + c) + 1) \sin(dx + c) + 1}{3(a^2d \cos(dx + c)^2 - a^2d \cos(dx + c) - 2a^2d - (a^2d \cos(dx + c) + 2a^2d) \sin(dx + c))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c))^2,x, algorithm="fricas")

[Out] $1/3 * ((3*d*x - 5) * \cos(d*x + c)^2 - 6*d*x - (3*d*x + 4) * \cos(d*x + c) - (6*d*x + (3*d*x + 5) * \cos(d*x + c) + 1) * \sin(d*x + c) + 1) / (a^2*d*\cos(d*x + c)^2 - a^2*d*\cos(d*x + c) - 2*a^2*d - (a^2*d*\cos(d*x + c) + 2*a^2*d)*\sin(d*x + c))$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{\frac{\csc^2(c+dx)+2 \csc(c+dx)+1}{a^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c))**2,x)

[Out] Integral(1/(csc(c + d*x)**2 + 2*csc(c + d*x) + 1), x)/a**2

Giac [A] time = 1.33023, size = 81, normalized size = 1.42

$$\frac{\frac{3(dx+c)}{a^2} + \frac{2\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 9 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 4\right)}{a^2\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right)^3}}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c))^2,x, algorithm="giac")

[Out] 1/3*(3*(d*x + c)/a^2 + 2*(3*tan(1/2*d*x + 1/2*c)^2 + 9*tan(1/2*d*x + 1/2*c) + 4)/(a^2*(tan(1/2*d*x + 1/2*c) + 1)^3)/d

$$3.12 \quad \int \frac{1}{(a+a \csc(c+dx))^3} dx$$

Optimal. Leaf size=88

$$\frac{22 \cot(c+dx)}{15d(a^3 \csc(c+dx) + a^3)} + \frac{x}{a^3} + \frac{7 \cot(c+dx)}{15ad(a \csc(c+dx) + a)^2} + \frac{\cot(c+dx)}{5d(a \csc(c+dx) + a)^3}$$

[Out] x/a^3 + Cot[c + d*x]/(5*d*(a + a*Csc[c + d*x])^3) + (7*Cot[c + d*x])/(15*a*d*(a + a*Csc[c + d*x])^2) + (22*Cot[c + d*x])/(15*d*(a^3 + a^3*Csc[c + d*x]))

Rubi [A] time = 0.109679, antiderivative size = 88, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3777, 3922, 3919, 3794}

$$\frac{22 \cot(c+dx)}{15d(a^3 \csc(c+dx) + a^3)} + \frac{x}{a^3} + \frac{7 \cot(c+dx)}{15ad(a \csc(c+dx) + a)^2} + \frac{\cot(c+dx)}{5d(a \csc(c+dx) + a)^3}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Csc[c + d*x])^(-3), x]

[Out] x/a^3 + Cot[c + d*x]/(5*d*(a + a*Csc[c + d*x])^3) + (7*Cot[c + d*x])/(15*a*d*(a + a*Csc[c + d*x])^2) + (22*Cot[c + d*x])/(15*d*(a^3 + a^3*Csc[c + d*x]))

Rule 3777

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^ (n_.), x_Symbol] :> -Simp[(Cot[c + d*x]*(a + b*Csc[c + d*x])^n)/(d*(2*n + 1)), x] + Dist[1/(a^2*(2*n + 1)), Int[(a + b*Csc[c + d*x])^(n + 1)*(a*(2*n + 1) - b*(n + 1)*Csc[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 3922

Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^ (m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)), x_Symbol] :> -Simp[((b*c - a*d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b*f*(2*m + 1)), x] + Dist[1/(a^2*(2*m + 1)), Int[(a + b*Csc[e + f*x])^(m + 1)*Simp[a*c*(2*m + 1) - (b*c - a*d)*(m + 1)*Csc[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ

$[a^2 - b^2, 0] \ \&\& \ \text{IntegerQ}[2*m]$

Rule 3919

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.) + (c_))/(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \text{ :> } \text{Simp}[(c*x)/a, x] - \text{Dist}[(b*c - a*d)/a, \text{Int}[\text{Csc}[e + f*x]/(a + b*\text{Csc}[e + f*x]), x], x] \text{ /; } \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0]$

Rule 3794

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]/(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \text{ :> } -\text{Simp}[\text{Cot}[e + f*x]/(f*(b + a*\text{Csc}[e + f*x])), x] \text{ /; } \text{FreeQ}[\{a, b, e, f\}, x] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + a \csc(c + dx))^3} dx &= \frac{\cot(c + dx)}{5d(a + a \csc(c + dx))^3} - \frac{\int \frac{-5a + 2a \csc(c + dx)}{(a + a \csc(c + dx))^2} dx}{5a^2} \\ &= \frac{\cot(c + dx)}{5d(a + a \csc(c + dx))^3} + \frac{7 \cot(c + dx)}{15ad(a + a \csc(c + dx))^2} + \frac{\int \frac{15a^2 - 7a^2 \csc(c + dx)}{a + a \csc(c + dx)} dx}{15a^4} \\ &= \frac{x}{a^3} + \frac{\cot(c + dx)}{5d(a + a \csc(c + dx))^3} + \frac{7 \cot(c + dx)}{15ad(a + a \csc(c + dx))^2} - \frac{22 \int \frac{\csc(c + dx)}{a + a \csc(c + dx)} dx}{15a^2} \\ &= \frac{x}{a^3} + \frac{\cot(c + dx)}{5d(a + a \csc(c + dx))^3} + \frac{7 \cot(c + dx)}{15ad(a + a \csc(c + dx))^2} + \frac{22 \cot(c + dx)}{15d(a^3 + a^3 \csc(c + dx))} \end{aligned}$$

Mathematica [A] time = 0.94444, size = 123, normalized size = 1.4

$$\frac{2 \sin\left(\frac{1}{2}(c+dx)\right)(-51 \sin(c+dx)+16 \cos(2(c+dx))-38)}{\left(\sin\left(\frac{1}{2}(c+dx)\right)+\cos\left(\frac{1}{2}(c+dx)\right)\right)^5} - \frac{13}{\left(\sin\left(\frac{1}{2}(c+dx)\right)+\cos\left(\frac{1}{2}(c+dx)\right)\right)^2} + \frac{3}{\left(\sin\left(\frac{1}{2}(c+dx)\right)+\cos\left(\frac{1}{2}(c+dx)\right)\right)^4} + 15c + 15dx}{15a^3d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + a*Csc[c + d*x])^(-3), x]

[Out] (15*c + 15*d*x + 3/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^4 - 13/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 + (2*Sin[(c + d*x)/2]*(-38 + 16*Cos[2*(c + d*x

)] - 51*Sin[c + d*x]))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^5)/(15*a^3*d)

Maple [A] time = 0.079, size = 125, normalized size = 1.4

$$2 \frac{\arctan(\tan(1/2 dx + c/2))}{da^3} - 4 \frac{1}{da^3 (\tan(1/2 dx + c/2) + 1)^4} + \frac{8}{5 da^3} \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)^{-5} + \frac{4}{3 da^3} \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+a*csc(d*x+c))^3,x)

[Out] 2/d/a^3*arctan(tan(1/2*d*x+1/2*c))-4/d/a^3/(tan(1/2*d*x+1/2*c)+1)^4+8/5/d/a^3/(tan(1/2*d*x+1/2*c)+1)^5+4/3/d/a^3/(tan(1/2*d*x+1/2*c)+1)^3+2/d/a^3/(tan(1/2*d*x+1/2*c)+1)^2+2/d/a^3/(tan(1/2*d*x+1/2*c)+1)

Maxima [B] time = 1.50908, size = 308, normalized size = 3.5

$$2 \left(\frac{\frac{95 \sin(dx+c)}{\cos(dx+c)+1} + \frac{145 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{75 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{15 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 22}{a^3 + \frac{5a^3 \sin(dx+c)}{\cos(dx+c)+1} + \frac{10a^3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{10a^3 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{5a^3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{a^3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5}} + \frac{15 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^3} \right) / 15d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c))^3,x, algorithm="maxima")

[Out] 2/15*((95*sin(d*x + c)/(cos(d*x + c) + 1) + 145*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 75*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 15*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 22)/(a^3 + 5*a^3*sin(d*x + c)/(cos(d*x + c) + 1) + 10*a^3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 10*a^3*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 5*a^3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + a^3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5) + 15*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^3)/d

Fricas [B] time = 0.486009, size = 475, normalized size = 5.4

$$\frac{(15 dx + 32) \cos(dx + c)^3 + (45 dx - 19) \cos(dx + c)^2 - 60 dx - 6(5 dx + 9) \cos(dx + c) + ((15 dx - 32) \cos(dx + c)^2 - 60 dx + 32) \cos(dx + c) + 15}{15(a^3 d \cos(dx + c)^3 + 3 a^3 d \cos(dx + c)^2 - 2 a^3 d \cos(dx + c) - 4 a^3 d + (a^3 d \cos(dx + c)^2 - 2 a^3 d \cos(dx + c) + 15) \arctan(\frac{\sin(dx + c)}{\cos(dx + c) + 1})) / d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c))^3,x, algorithm="fricas")

[Out] 1/15*((15*d*x + 32)*cos(d*x + c)^3 + (45*d*x - 19)*cos(d*x + c)^2 - 60*d*x - 6*(5*d*x + 9)*cos(d*x + c) + ((15*d*x - 32)*cos(d*x + c)^2 - 60*d*x - 3*(10*d*x + 17)*cos(d*x + c) + 3)*sin(d*x + c) - 3)/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d + (a^3*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d)*sin(d*x + c))

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{\frac{\csc^3(c+dx)+3\csc^2(c+dx)+3\csc(c+dx)+1}{a^3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c))**3,x)

[Out] Integral(1/(csc(c + d*x)**3 + 3*csc(c + d*x)**2 + 3*csc(c + d*x) + 1), x)/a**3

Giac [A] time = 1.25117, size = 116, normalized size = 1.32

$$\frac{\frac{15(dx+c)}{a^3} + \frac{2\left(15 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 + 75 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 + 145 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 95 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 22\right)}{a^3\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right)^5}}{15d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(d*x+c))^3,x, algorithm="giac")

[Out] 1/15*(15*(d*x + c)/a^3 + 2*(15*tan(1/2*d*x + 1/2*c)^4 + 75*tan(1/2*d*x + 1/2*c)^3 + 145*tan(1/2*d*x + 1/2*c)^2 + 95*tan(1/2*d*x + 1/2*c) + 22)/(a^3*(tan(1/2*d*x + 1/2*c) + 1)^5))/d

3.13 $\int (a + a \csc(x))^{5/2} dx$

Optimal. Leaf size=65

$$-\frac{14a^3 \cot(x)}{3\sqrt{a \csc(x) + a}} - \frac{2}{3}a^2 \cot(x)\sqrt{a \csc(x) + a} - 2a^{5/2} \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x) + a}}\right)$$

[Out] $-2*a^{(5/2)}*ArcTan[(Sqrt[a]*Cot[x])/Sqrt[a + a*Csc[x]]] - (14*a^3*Cot[x])/(3*Sqrt[a + a*Csc[x]]) - (2*a^2*Cot[x]*Sqrt[a + a*Csc[x]])/3$

Rubi [A] time = 0.0913834, antiderivative size = 65, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.5$, Rules used = {3775, 3915, 3774, 203, 3792}

$$-\frac{14a^3 \cot(x)}{3\sqrt{a \csc(x) + a}} - \frac{2}{3}a^2 \cot(x)\sqrt{a \csc(x) + a} - 2a^{5/2} \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x) + a}}\right)$$

Antiderivative was successfully verified.

[In] Int[(a + a*Csc[x])^(5/2),x]

[Out] $-2*a^{(5/2)}*ArcTan[(Sqrt[a]*Cot[x])/Sqrt[a + a*Csc[x]]] - (14*a^3*Cot[x])/(3*Sqrt[a + a*Csc[x]]) - (2*a^2*Cot[x]*Sqrt[a + a*Csc[x]])/3$

Rule 3775

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] :> -Simp[(b^2*Cot[c + d*x]*(a + b*Csc[c + d*x])^(n - 2))/(d*(n - 1)), x] + Dist[a/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 2)*(a*(n - 1) + b*(3*n - 4)*Csc[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 3915

```
Int[Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)), x_Symbol] :> Dist[c, Int[Sqrt[a + b*Csc[e + f*x]], x], x] + Dist[d, Int[Sqrt[a + b*Csc[e + f*x]]*Csc[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0]
```

Rule 3774


```
Int[Sqrt[csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Dist[(-2*b)/d,
  Subst[Int[1/(a + x^2), x], x, (b*Cot[c + d*x])/Sqrt[a + b*Csc[c + d*x]]],
x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]
```

Rule 203

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt
[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 3792

```
Int[csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_S
ymbol] := Simp[(-2*b*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]]), x] /; Free
Q[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned}
\int (a + a \csc(x))^{5/2} dx &= -\frac{2}{3}a^2 \cot(x)\sqrt{a + a \csc(x)} + \frac{1}{3}(2a) \int \sqrt{a + a \csc(x)} \left(\frac{3a}{2} + \frac{7}{2}a \csc(x) \right) dx \\
&= -\frac{2}{3}a^2 \cot(x)\sqrt{a + a \csc(x)} + a^2 \int \sqrt{a + a \csc(x)} dx + \frac{1}{3}(7a^2) \int \csc(x)\sqrt{a + a \csc(x)} dx \\
&= -\frac{14a^3 \cot(x)}{3\sqrt{a + a \csc(x)}} - \frac{2}{3}a^2 \cot(x)\sqrt{a + a \csc(x)} - (2a^3) \text{Subst} \left(\int \frac{1}{a + x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}} \right) \\
&= -2a^{5/2} \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{a + a \csc(x)}} \right) - \frac{14a^3 \cot(x)}{3\sqrt{a + a \csc(x)}} - \frac{2}{3}a^2 \cot(x)\sqrt{a + a \csc(x)}
\end{aligned}$$

Mathematica [A] time = 1.62999, size = 80, normalized size = 1.23

$$-\frac{2a^2\sqrt{a(\csc(x)+1)}\left(\cos\left(\frac{x}{2}\right)-\sin\left(\frac{x}{2}\right)\right)\left(\sqrt{\csc(x)-1}(\csc(x)+8)+3\tan^{-1}\left(\sqrt{\csc(x)-1}\right)\right)}{3\sqrt{\csc(x)-1}\left(\sin\left(\frac{x}{2}\right)+\cos\left(\frac{x}{2}\right)\right)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + a*Csc[x])^(5/2),x]
```

```
[Out] (-2*a^2*Sqrt[a*(1 + Csc[x])]*(3*ArcTan[Sqrt[-1 + Csc[x]]] + Sqrt[-1 + Csc[x]]*(8 + Csc[x]))*(Cos[x/2] - Sin[x/2]))/(3*Sqrt[-1 + Csc[x]]*(Cos[x/2] + Sin[x/2]))
```

Maple [B] time = 0.21, size = 535, normalized size = 8.2

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+a*csc(x))^(5/2),x)`

[Out]
$$\begin{aligned} & \frac{1}{6}2^{1/2}*(-12*\sin(x)*\cos(x)*\arctan(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}-1) \\ &)*(-(-1+\cos(x))/\sin(x))^{3/2}-3*\sin(x)*\cos(x)*\ln(-(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2} \\ &)*\sin(x)-\sin(x)+\cos(x)-1)/(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}*\sin(x)+\sin(x)-\cos(x)+1)) \\ &)*(-(-1+\cos(x))/\sin(x))^{3/2}-3*\sin(x)*\cos(x)*\ln(-(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2} \\ &)*\sin(x)+\sin(x)-\cos(x)+1)/(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}*\sin(x)-\sin(x)+\cos(x)-1)) \\ &)*(-(-1+\cos(x))/\sin(x))^{3/2}-12*\sin(x)*\cos(x)*\arctan(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}+1) \\ &)*(-(-1+\cos(x))/\sin(x))^{3/2}-12*\sin(x)*\arctan(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}-1) \\ &)*(-(-1+\cos(x))/\sin(x))^{3/2}-3*\sin(x)*\ln(-(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}*\sin(x)-\sin(x)+\cos(x)-1) \\ &)/(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}*\sin(x)+\sin(x)-\cos(x)+1)) \\ &)*(-(-1+\cos(x))/\sin(x))^{3/2}-3*\sin(x)*\ln(-(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}*\sin(x)-\sin(x)+\cos(x)-1) \\ &)/(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}*\sin(x)+\sin(x)-\cos(x)+1)) \\ &)*(-(-1+\cos(x))/\sin(x))^{3/2}-12*\sin(x)*\arctan(2^{1/2}*(-(-1+\cos(x))/\sin(x))^{1/2}+1) \\ &)*(-(-1+\cos(x))/\sin(x))^{3/2}+16*\sin(x)*\cos(x)*2^{1/2}-16*\cos(x)^2*2^{1/2}-14*\sin(x)*2^{1/2}+2*\cos(x)*2^{1/2}+14*2^{1/2} \\ &) \\ &)*\sin(x)*(a*(\sin(x)+1)/\sin(x))^{5/2}/(\cos(x)^2*\sin(x)-\cos(x)^3+2*\cos(x)*\sin(x)+3*\cos(x)^2-4*\sin(x)+2*\cos(x)-4) \end{aligned}$$

Maxima [B] time = 1.58343, size = 563, normalized size = 8.66

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(x))^(5/2),x, algorithm="maxima")`

[Out]
$$\begin{aligned} & \frac{1}{22}\sqrt{2}*a^{5/2}*(\sin(x)/(\cos(x)+1))^{11/2} + \frac{5}{18}\sqrt{2}*a^{5/2}*(\sin(x)/(\cos(x)+1))^{9/2} \\ & + \frac{9}{14}\sqrt{2}*a^{5/2}*(\sin(x)/(\cos(x)+1))^{7/2} + \frac{1}{2}\sqrt{2}*a^{5/2}*(\sin(x)/(\cos(x)+1))^{5/2} - \frac{2}{3}\sqrt{2}*a^{5/2}*(\sin(x)/(\cos(x)+1))^{3/2} \\ & + \sqrt{2}*(\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2}+2*\sqrt{\sin(x)/(\cos(x)+1)}))) + \sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2}-2*\sqrt{\sin(x)/(\cos(x)+1)}))) \\ &)*a^{5/2} - 2*\sqrt{2}*a^{5/2}*\sqrt{\sin(x)/(\cos(x)+1)} \end{aligned}$$

+ 1)) - 1/1386*(693*sqrt(2)*a^(5/2)*sin(x)/(cos(x) + 1) + 1155*sqrt(2)*a^(5/2)*sin(x)^2/(cos(x) + 1)^2 + 1386*sqrt(2)*a^(5/2)*sin(x)^3/(cos(x) + 1)^3 + 990*sqrt(2)*a^(5/2)*sin(x)^4/(cos(x) + 1)^4 + 385*sqrt(2)*a^(5/2)*sin(x)^5/(cos(x) + 1)^5 + 63*sqrt(2)*a^(5/2)*sin(x)^6/(cos(x) + 1)^6)/sqrt(sin(x)/(cos(x) + 1)) - 1/42*(7*sqrt(2)*a^(5/2)*sin(x)/(cos(x) + 1) + 105*sqrt(2)*a^(5/2)*sin(x)^2/(cos(x) + 1)^2 - 210*sqrt(2)*a^(5/2)*sin(x)^3/(cos(x) + 1)^3 - 70*sqrt(2)*a^(5/2)*sin(x)^4/(cos(x) + 1)^4 - 21*sqrt(2)*a^(5/2)*sin(x)^5/(cos(x) + 1)^5 - 3*sqrt(2)*a^(5/2)*sin(x)^6/(cos(x) + 1)^6)/(sin(x)/(cos(x) + 1))^(5/2)

Fricas [B] time = 0.514679, size = 900, normalized size = 13.85

$$\left[\frac{3 \left(a^2 \cos(x)^2 - a^2 - (a^2 \cos(x) + a^2) \sin(x) \right) \sqrt{-a} \log \left(\frac{2a \cos(x)^2 - 2(\cos(x)^2 + (\cos(x)+1) \sin(x) - 1) \sqrt{-a} \sqrt{\frac{a \sin(x) + a}{\sin(x)}} + a \cos(x) - (2a \cos(x) + a) \sin(x) - a}{\cos(x) + \sin(x) + 1} \right)}{3 \left(\cos(x)^2 - (\cos(x) + 1) \sin(x) - 1 \right)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(x))^(5/2),x, algorithm="fricas")

[Out] [1/3*(3*(a^2*cos(x)^2 - a^2 - (a^2*cos(x) + a^2)*sin(x))*sqrt(-a)*log((2*a*cos(x)^2 - 2*(cos(x)^2 + (cos(x) + 1)*sin(x) - 1)*sqrt(-a)*sqrt((a*sin(x) + a)/sin(x)) + a*cos(x) - (2*a*cos(x) + a)*sin(x) - a)/(cos(x) + sin(x) + 1)) + 2*(8*a^2*cos(x)^2 + a^2*cos(x) - 7*a^2 + (8*a^2*cos(x) + 7*a^2)*sin(x))*sqrt((a*sin(x) + a)/sin(x)))/(cos(x)^2 - (cos(x) + 1)*sin(x) - 1), 2/3*(3*(a^2*cos(x)^2 - a^2 - (a^2*cos(x) + a^2)*sin(x))*sqrt(a)*arctan(-sqrt(a)*sqrt((a*sin(x) + a)/sin(x))*(cos(x) - sin(x) + 1)/(a*cos(x) + a*sin(x) + a)) + (8*a^2*cos(x)^2 + a^2*cos(x) - 7*a^2 + (8*a^2*cos(x) + 7*a^2)*sin(x))*sqrt((a*sin(x) + a)/sin(x)))/(cos(x)^2 - (cos(x) + 1)*sin(x) - 1)]

Sympy [F(-1)] time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(x))**(5/2),x)

[Out] Timed out

Giac [B] time = 2.24225, size = 339, normalized size = 5.22

$$\frac{1}{6} \sqrt{2} \sqrt{a \tan\left(\frac{1}{2}x\right)} a^2 \tan\left(\frac{1}{2}x\right) + \frac{5}{2} \sqrt{2} \sqrt{a \tan\left(\frac{1}{2}x\right)} a^2 + \left(a^2 \sqrt{|a|} + a|a|^{\frac{3}{2}}\right) \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\sqrt{|a|} + 2\sqrt{a \tan\left(\frac{1}{2}x\right)}\right)}{2\sqrt{|a|}}\right) + \left(a\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(x))^(5/2),x, algorithm="giac")

[Out] 1/6*sqrt(2)*sqrt(a*tan(1/2*x))*a^2*tan(1/2*x) + 5/2*sqrt(2)*sqrt(a*tan(1/2*x))*a^2 + (a^2*sqrt(abs(a)) + a*abs(a)^(3/2))*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(abs(a)) + 2*sqrt(a*tan(1/2*x)))/sqrt(abs(a))) + (a^2*sqrt(abs(a)) + a*abs(a)^(3/2))*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(abs(a)) - 2*sqrt(a*tan(1/2*x)))/sqrt(abs(a))) + 1/2*(a^2*sqrt(abs(a)) - a*abs(a)^(3/2))*log(a*tan(1/2*x) + sqrt(2)*sqrt(a*tan(1/2*x))*sqrt(abs(a)) + abs(a)) - 1/2*(a^2*sqrt(abs(a)) - a*abs(a)^(3/2))*log(a*tan(1/2*x) - sqrt(2)*sqrt(a*tan(1/2*x))*sqrt(abs(a)) + abs(a)) - 1/6*sqrt(2)*(15*a^4*tan(1/2*x) + a^4)/(sqrt(a*tan(1/2*x))*a*tan(1/2*x))

3.14 $\int (a + a \csc(x))^{3/2} dx$

Optimal. Leaf size=44

$$-\frac{2a^2 \cot(x)}{\sqrt{a \csc(x) + a}} - 2a^{3/2} \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x) + a}} \right)$$

[Out] $-2*a^{(3/2)}*ArcTan[(Sqrt[a]*Cot[x])/Sqrt[a + a*Csc[x]]] - (2*a^2*Cot[x])/Sqrt[a + a*Csc[x]]$

Rubi [A] time = 0.0304638, antiderivative size = 44, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.4$, Rules used = {3775, 21, 3774, 203}

$$-\frac{2a^2 \cot(x)}{\sqrt{a \csc(x) + a}} - 2a^{3/2} \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x) + a}} \right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + a*\text{Csc}[x])^{(3/2)}, x]$

[Out] $-2*a^{(3/2)}*ArcTan[(Sqrt[a]*Cot[x])/Sqrt[a + a*Csc[x]]] - (2*a^2*Cot[x])/Sqrt[a + a*Csc[x]]$

Rule 3775

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^{(n_)}, x_Symbol] := -\text{Simp}[(b^2*\text{Cot}[c + d*x]*(a + b*\text{Csc}[c + d*x])^{(n-2)})/(d*(n-1)), x] + \text{Dist}[a/(n-1), \text{Int}[(a + b*\text{Csc}[c + d*x])^{(n-2)}*(a*(n-1) + b*(3*n-4)*\text{Csc}[c + d*x]), x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 21

$\text{Int}[(u_.)*((a_.) + (b_.)*(v_))^{(m_.)}*((c_.) + (d_.)*(v_))^{(n_.)}, x_Symbol] := \text{Dist}[(b/d)^m, \text{Int}[u*(c + d*v)^{(m+n)}, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \text{EqQ}[b*c - a*d, 0] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (!\text{IntegerQ}[n] \ || \ \text{SimplerQ}[c + d*x, a + b*x])$

Rule 3774

```
Int[Sqrt[csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] := Dist[(-2*b)/d,
  Subst[Int[1/(a + x^2), x], x, (b*Cot[c + d*x])/Sqrt[a + b*Csc[c + d*x]]],
  x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]
```

Rule 203

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[Rt[b, 2]*x]/Rt
[a, 2])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rubi steps

$$\begin{aligned}
\int (a + a \csc(x))^{3/2} dx &= -\frac{2a^2 \cot(x)}{\sqrt{a + a \csc(x)}} + (2a) \int \frac{\frac{a}{2} + \frac{1}{2}a \csc(x)}{\sqrt{a + a \csc(x)}} dx \\
&= -\frac{2a^2 \cot(x)}{\sqrt{a + a \csc(x)}} + a \int \sqrt{a + a \csc(x)} dx \\
&= -\frac{2a^2 \cot(x)}{\sqrt{a + a \csc(x)}} - (2a^2) \text{Subst} \left(\int \frac{1}{a + x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}} \right) \\
&= -2a^{3/2} \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{a + a \csc(x)}} \right) - \frac{2a^2 \cot(x)}{\sqrt{a + a \csc(x)}}
\end{aligned}$$

Mathematica [A] time = 0.0853796, size = 69, normalized size = 1.57

$$-\frac{2a\sqrt{a(\csc(x)+1)}\left(\cos\left(\frac{x}{2}\right)-\sin\left(\frac{x}{2}\right)\right)\left(\sqrt{\csc(x)-1}+\tan^{-1}\left(\sqrt{\csc(x)-1}\right)\right)}{\sqrt{\csc(x)-1}\left(\sin\left(\frac{x}{2}\right)+\cos\left(\frac{x}{2}\right)\right)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + a*Csc[x])^(3/2), x]
```

```
[Out] (-2*a*(ArcTan[Sqrt[-1 + Csc[x]]] + Sqrt[-1 + Csc[x]])*Sqrt[a*(1 + Csc[x])]*
(Cos[x/2] - Sin[x/2]))/(Sqrt[-1 + Csc[x]]*(Cos[x/2] + Sin[x/2]))
```

Maple [B] time = 0.16, size = 273, normalized size = 6.2

$$-\frac{\sin(x)\sqrt{2}}{2\cos(x)\sin(x)+2(\cos(x))^2-4\sin(x)+2\cos(x)-4}\left(\sin(x)\sqrt{\frac{-1+\cos(x)}{\sin(x)}}\ln\left(-\left(\sqrt{2}\sqrt{\frac{-1+\cos(x)}{\sin(x)}}\sin(x)+\dots\right)\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+a*csc(x))^(3/2),x)`

[Out]
$$\begin{aligned} & -1/2*2^{(1/2)}*(\sin(x)*(-(-1+\cos(x))/\sin(x))^{(1/2)}*\ln(-(2^{(1/2)}*(-(-1+\cos(x)) \\ & / \sin(x))^{(1/2)}*\sin(x)+\sin(x)-\cos(x)+1)/(2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)} \\ & *\sin(x)-\sin(x)+\cos(x)-1))+4*\sin(x)*(-(-1+\cos(x))/\sin(x))^{(1/2)}*\arctan(2^{(1/2)} \\ & *(-(-1+\cos(x))/\sin(x))^{(1/2)}+1)+4*\sin(x)*(-(-1+\cos(x))/\sin(x))^{(1/2)}*\arctan \\ & (2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)}-1)+\sin(x)*(-(-1+\cos(x))/\sin(x))^{(1/2)} \\ &)*\ln(-(2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)}*\sin(x)-\sin(x)+\cos(x)-1)/(2^{(1/2)} \\ & *(-(-1+\cos(x))/\sin(x))^{(1/2)}*\sin(x)+\sin(x)-\cos(x)+1))-2*\sin(x)*2^{(1/2)}-2*\cos \\ & (x)*2^{(1/2)}+2*2^{(1/2)}*\sin(x)*(a*(\sin(x)+1)/\sin(x))^{(3/2)}/(\cos(x)*\sin(x)+\cos \\ & (x)^2-2*\sin(x)+\cos(x)-2) \end{aligned}$$

Maxima [B] time = 1.54284, size = 270, normalized size = 6.14

$$\sqrt{2} \left(\sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + 2 \sqrt{\frac{\sin(x)}{\cos(x)+1}} \right) \right) + \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} \left(\sqrt{2} - 2 \sqrt{\frac{\sin(x)}{\cos(x)+1}} \right) \right) \right) a^{\frac{3}{2}} - \frac{1}{5} \sqrt{2} \left(a^{\frac{3}{2}} \left(\frac{\sin(x)}{\cos(x)+1} \right)^{\frac{3}{2}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(x))^(3/2),x, algorithm="maxima")`

[Out]
$$\begin{aligned} & \sqrt{2}*(\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*\sqrt{\sin(x)/(\cos(x) + 1)})) \\ &) + \sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*\sqrt{\sin(x)/(\cos(x) + 1)}))) * a \\ & ^{(3/2)} - 1/5*\sqrt{2}*(a^{(3/2)}*(\sin(x)/(\cos(x) + 1))^{(5/2)} + 5*a^{(3/2)}*(\sin \\ & (x)/(\cos(x) + 1))^{(3/2)} + 10*a^{(3/2)}*\sqrt{\sin(x)/(\cos(x) + 1)}) - 1/5*(5*\sqrt{2} \\ & *a^{(3/2)}*\sin(x)/(\cos(x) + 1) - 15*\sqrt{2} *a^{(3/2)}*\sin(x)^2/(\cos(x) + 1) \\ & ^2 - 5*\sqrt{2} *a^{(3/2)}*\sin(x)^3/(\cos(x) + 1)^3 - \sqrt{2} *a^{(3/2)}*\sin(x)^4/(\cos(x) + 1)^4) / (\sin(x)/(\cos(x) + 1))^{(3/2)} \end{aligned}$$

Fricas [B] time = 0.501578, size = 666, normalized size = 15.14

$$\left[\frac{(a \cos(x) + a \sin(x) + a) \sqrt{-a} \log \left(\frac{2a \cos(x)^2 - 2(\cos(x)^2 + (\cos(x)+1) \sin(x) - 1) \sqrt{-a} \sqrt{\frac{a \sin(x)+a}{\sin(x)}} + a \cos(x) - (2a \cos(x)+a) \sin(x) - a}{\cos(x) + \sin(x) + 1} \right) - 2(a \cos(x) + a \sin(x) + a) \sqrt{a} \arctan \left(\frac{\sqrt{a} \sqrt{\frac{a \sin(x)+a}{\sin(x)}}}{\cos(x) - \sin(x) + 1} \right)}{\cos(x) + \sin(x) + 1} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(x))^(3/2),x, algorithm="fricas")

[Out] [((a*cos(x) + a*sin(x) + a)*sqrt(-a)*log((2*a*cos(x)^2 - 2*(cos(x)^2 + (cos(x) + 1)*sin(x) - 1)*sqrt(-a)*sqrt((a*sin(x) + a)/sin(x)) + a*cos(x) - (2*a*cos(x) + a)*sin(x) - a)/(cos(x) + sin(x) + 1)) - 2*(a*cos(x) - a*sin(x) + a)*sqrt((a*sin(x) + a)/sin(x)))/(cos(x) + sin(x) + 1), 2*((a*cos(x) + a*sin(x) + a)*sqrt(a)*arctan(-sqrt(a)*sqrt((a*sin(x) + a)/sin(x))*(cos(x) - sin(x) + 1)/(a*cos(x) + a*sin(x) + a)) - (a*cos(x) - a*sin(x) + a)*sqrt((a*sin(x) + a)/sin(x)))/(cos(x) + sin(x) + 1)]

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(x) + a)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(x))**(3/2),x)

[Out] Integral((a*csc(x) + a)**(3/2), x)

Giac [B] time = 2.07439, size = 263, normalized size = 5.98

$$\sqrt{2} \sqrt{a \tan\left(\frac{1}{2}x\right)} a - \frac{\sqrt{2}a^2}{\sqrt{a \tan\left(\frac{1}{2}x\right)}} + \left(a\sqrt{|a|} + |a|^{\frac{3}{2}}\right) \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\sqrt{|a|} + 2\sqrt{a \tan\left(\frac{1}{2}x\right)}\right)}{2\sqrt{|a|}}\right) + \left(a\sqrt{|a|} + |a|^{\frac{3}{2}}\right) \arctan\left(\frac{1}{\sqrt{|a|}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.


```
[In] integrate((a+a*csc(x))^(3/2),x, algorithm="giac")
```

```
[Out] sqrt(2)*sqrt(a*tan(1/2*x))*a - sqrt(2)*a^2/sqrt(a*tan(1/2*x)) + (a*sqrt(abs(a) + abs(a)^(3/2))*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(abs(a)) + 2*sqrt(a*tan(1/2*x)))/sqrt(abs(a))) + (a*sqrt(abs(a)) + abs(a)^(3/2))*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(abs(a)) - 2*sqrt(a*tan(1/2*x)))/sqrt(abs(a))) + 1/2*(a*sqrt(abs(a)) - abs(a)^(3/2))*log(a*tan(1/2*x) + sqrt(2)*sqrt(a*tan(1/2*x))*sqrt(abs(a) + abs(a)) - 1/2*(a*sqrt(abs(a)) - abs(a)^(3/2))*log(a*tan(1/2*x) - sqrt(2)*sqrt(a*tan(1/2*x))*sqrt(abs(a) + abs(a)))
```

3.15 $\int \sqrt{a + a \csc(x)} dx$

Optimal. Leaf size=26

$$-2\sqrt{a} \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x) + a}} \right)$$

[Out] $-2*\text{Sqrt}[a]*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])/\text{Sqrt}[a + a*\text{Csc}[x]]]$

Rubi [A] time = 0.0155253, antiderivative size = 26, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.2$, Rules used = {3774, 203}

$$-2\sqrt{a} \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x) + a}} \right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[a + a*\text{Csc}[x]], x]$

[Out] $-2*\text{Sqrt}[a]*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])/\text{Sqrt}[a + a*\text{Csc}[x]]]$

Rule 3774

$\text{Int}[\text{Sqrt}[\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] \rightarrow \text{Dist}[(-2*b)/d, \text{Subst}[\text{Int}[1/(a + x^2), x], x, (b*\text{Cot}[c + d*x])/\text{Sqrt}[a + b*\text{Csc}[c + d*x]]], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

Rule 203

$\text{Int}[(a_. + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1*\text{ArcTan}[(\text{Rt}[b, 2]*x)/\text{Rt}[a, 2]])/(\text{Rt}[a, 2]*\text{Rt}[b, 2]), x] /;$ $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rubi steps

$$\begin{aligned} \int \sqrt{a + a \csc(x)} dx &= - \left((2a) \text{Subst} \left(\int \frac{1}{a + x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}} \right) \right) \\ &= -2\sqrt{a} \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{a + a \csc(x)}} \right) \end{aligned}$$

Mathematica [A] time = 0.0518105, size = 32, normalized size = 1.23

$$-\frac{2a \cot(x) \tan^{-1}(\sqrt{\csc(x)-1})}{\sqrt{\csc(x)-1} \sqrt{a(\csc(x)+1)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + a*Csc[x]], x]

[Out] (-2*a*ArcTan[Sqrt[-1 + Csc[x]]]*Cot[x])/(Sqrt[-1 + Csc[x]]*Sqrt[a*(1 + Csc[x])])

Maple [B] time = 0.151, size = 199, normalized size = 7.7

$$\frac{\sin(x) \sqrt{2}}{2 - 2 \cos(x) + 2 \sin(x)} \sqrt{\frac{a(\sin(x)+1)}{\sin(x)}} \sqrt{\frac{-1 + \cos(x)}{\sin(x)}} \left(\ln \left(- \left(\sqrt{2} \sqrt{\frac{-1 + \cos(x)}{\sin(x)}} \sin(x) + \sin(x) - \cos(x) + 1 \right) \right) \right) \left(\sqrt{\frac{a(\sin(x)+1)}{\sin(x)}} \sqrt{\frac{-1 + \cos(x)}{\sin(x)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+a*csc(x))^(1/2), x)

[Out] 1/2*2^(1/2)*(a*(sin(x)+1)/sin(x))^(1/2)*sin(x)*(-(-1+cos(x))/sin(x))^(1/2)*(ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1))+4*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1)+4*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)-1)+ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1)))/(1-cos(x)+sin(x))

Maxima [B] time = 1.58212, size = 200, normalized size = 7.69

$$-\frac{2}{3} \sqrt{2} \sqrt{a} \left(\frac{\sin(x)}{\cos(x)+1} \right)^{\frac{3}{2}} + \sqrt{2} \left(\sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + 2 \sqrt{\frac{\sin(x)}{\cos(x)+1}} \right) \right) + \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} \left(\sqrt{2} - 2 \sqrt{\frac{\sin(x)}{\cos(x)+1}} \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(x))^(1/2), x, algorithm="maxima")

```
[Out] -2/3*sqrt(2)*sqrt(a)*(sin(x)/(cos(x) + 1))^(3/2) + sqrt(2)*(sqrt(2)*arctan(
1/2*sqrt(2)*(sqrt(2) + 2*sqrt(sin(x)/(cos(x) + 1)))) + sqrt(2)*arctan(-1/2*
sqrt(2)*(sqrt(2) - 2*sqrt(sin(x)/(cos(x) + 1)))))*sqrt(a) - 2*sqrt(2)*sqrt(
a)*sqrt(sin(x)/(cos(x) + 1)) + 2/3*(3*sqrt(2)*sqrt(a)*sin(x)/(cos(x) + 1) +
sqrt(2)*sqrt(a)*sin(x)^2/(cos(x) + 1)^2)/sqrt(sin(x)/(cos(x) + 1))
```

Fricas [B] time = 0.488618, size = 377, normalized size = 14.5

$$\left[\sqrt{-a} \log \left(\frac{2a \cos(x)^2 - 2(\cos(x)^2 + (\cos(x) + 1) \sin(x) - 1) \sqrt{-a} \sqrt{\frac{a \sin(x) + a}{\sin(x)}} + a \cos(x) - (2a \cos(x) + a) \sin(x) - a}{\cos(x) + \sin(x) + 1} \right) \right],$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+a*csc(x))^(1/2),x, algorithm="fricas")
```

```
[Out] [sqrt(-a)*log((2*a*cos(x)^2 - 2*(cos(x)^2 + (cos(x) + 1)*sin(x) - 1)*sqrt(-
a)*sqrt((a*sin(x) + a)/sin(x)) + a*cos(x) - (2*a*cos(x) + a)*sin(x) - a)/(c
os(x) + sin(x) + 1)), 2*sqrt(a)*arctan(-sqrt(a)*sqrt((a*sin(x) + a)/sin(x))
*(cos(x) - sin(x) + 1)/(a*cos(x) + a*sin(x) + a))]
```

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(x) + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+a*csc(x))**(1/2),x)
```

```
[Out] Integral(sqrt(a*csc(x) + a), x)
```

Giac [B] time = 2.0381, size = 477, normalized size = 18.35

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+a*csc(x))^(1/2),x, algorithm="giac")
```

```
[Out] 1/4*sqrt(2)*(2*sqrt(2)*(a*sqrt(abs(a))*sgn(tan(1/2*x)^3 + tan(1/2*x)^2 + ta
n(1/2*x) + 1) + abs(a)^(3/2)*sgn(tan(1/2*x)^3 + tan(1/2*x)^2 + tan(1/2*x) +
1))*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(abs(a)) + 2*sqrt(a*tan(1/2*x)))/sqrt(
abs(a)))/a + 2*sqrt(2)*(a*sqrt(abs(a))*sgn(tan(1/2*x)^3 + tan(1/2*x)^2 + ta
n(1/2*x) + 1) + abs(a)^(3/2)*sgn(tan(1/2*x)^3 + tan(1/2*x)^2 + tan(1/2*x) +
1))*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(abs(a)) - 2*sqrt(a*tan(1/2*x)))/sqrt
(abs(a)))/a + sqrt(2)*(a*sqrt(abs(a))*sgn(tan(1/2*x)^3 + tan(1/2*x)^2 + tan
(1/2*x) + 1) - abs(a)^(3/2)*sgn(tan(1/2*x)^3 + tan(1/2*x)^2 + tan(1/2*x) +
1))*log(a*tan(1/2*x) + sqrt(2)*sqrt(a*tan(1/2*x))*sqrt(abs(a)) + abs(a))/a
- sqrt(2)*(a*sqrt(abs(a))*sgn(tan(1/2*x)^3 + tan(1/2*x)^2 + tan(1/2*x) + 1)
- abs(a)^(3/2)*sgn(tan(1/2*x)^3 + tan(1/2*x)^2 + tan(1/2*x) + 1))*log(a*ta
n(1/2*x) - sqrt(2)*sqrt(a*tan(1/2*x))*sqrt(abs(a)) + abs(a))/a)*sgn(sin(x))
```

$$3.16 \quad \int \frac{1}{\sqrt{a+a \csc(x)}} dx$$

Optimal. Leaf size=62

$$\frac{\sqrt{2} \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{2}\sqrt{a \csc(x)+a}}\right)}{\sqrt{a}} - \frac{2 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x)+a}}\right)}{\sqrt{a}}$$

[Out] (-2*ArcTan[(Sqrt[a]*Cot[x])/Sqrt[a + a*Csc[x]]])/Sqrt[a] + (Sqrt[2]*ArcTan[(Sqrt[a]*Cot[x])/(Sqrt[2]*Sqrt[a + a*Csc[x]])])/Sqrt[a]

Rubi [A] time = 0.0575193, antiderivative size = 62, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.4$, Rules used = {3776, 3774, 203, 3795}

$$\frac{\sqrt{2} \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{2}\sqrt{a \csc(x)+a}}\right)}{\sqrt{a}} - \frac{2 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x)+a}}\right)}{\sqrt{a}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a + a*Csc[x]],x]

[Out] (-2*ArcTan[(Sqrt[a]*Cot[x])/Sqrt[a + a*Csc[x]]])/Sqrt[a] + (Sqrt[2]*ArcTan[(Sqrt[a]*Cot[x])/(Sqrt[2]*Sqrt[a + a*Csc[x]])])/Sqrt[a]

Rule 3776

Int[1/Sqrt[csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Dist[1/a, Int[Sqrt[a + b*Csc[c + d*x]], x], x] - Dist[b/a, Int[Csc[c + d*x]/Sqrt[a + b*Csc[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]

Rule 3774

Int[Sqrt[csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Dist[(-2*b)/d, Subst[Int[1/(a + x^2), x], x, (b*Cot[c + d*x])/Sqrt[a + b*Csc[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]

Rule 203

Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a

, 0] || GtQ[b, 0])

Rule 3795

Int[csc[(e_.) + (f_.)*(x_.)]/Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] :> Dist[-2/f, Subst[Int[1/(2*a + x^2), x], x, (b*Cot[e + f*x])/Sqrt[a + b*Csc[e + f*x]]], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a + a \csc(x)}} dx &= \frac{\int \sqrt{a + a \csc(x)} dx}{a} - \int \frac{\csc(x)}{\sqrt{a + a \csc(x)}} dx \\ &= -\left(2 \operatorname{Subst}\left(\int \frac{1}{a + x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}}\right)\right) + 2 \operatorname{Subst}\left(\int \frac{1}{2a + x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}}\right) \\ &= -\frac{2 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a + a \csc(x)}}\right)}{\sqrt{a}} + \frac{\sqrt{2} \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{2}\sqrt{a + a \csc(x)}}\right)}{\sqrt{a}} \end{aligned}$$

Mathematica [A] time = 0.12595, size = 54, normalized size = 0.87

$$\frac{\cot(x) \left(\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\csc(x)-1}}{\sqrt{2}} \right) - 2 \tan^{-1} \left(\sqrt{\csc(x)-1} \right) \right)}{\sqrt{\csc(x)-1} \sqrt{a(\csc(x)+1)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[a + a*Csc[x]], x]

[Out] ((-2*ArcTan[Sqrt[-1 + Csc[x]]] + Sqrt[2]*ArcTan[Sqrt[-1 + Csc[x]]/Sqrt[2]])*Cot[x])/(Sqrt[-1 + Csc[x]]*Sqrt[a*(1 + Csc[x])])

Maple [B] time = 0.139, size = 221, normalized size = 3.6

$$-\frac{\sqrt{2}(1 - \cos(x) + \sin(x))}{4 \sin(x)} \left(4 \sqrt{2} \arctan \left(\sqrt{-\frac{-1 + \cos(x)}{\sin(x)}} \right) - \ln \left(-\left(\sqrt{2} \sqrt{-\frac{-1 + \cos(x)}{\sin(x)}} \sin(x) + \sin(x) - \cos(x) + 1 \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+a*csc(x))^(1/2),x)

[Out]
$$-1/4*2^{(1/2)}*(4*2^{(1/2)}*\arctan((-(-1+\cos(x))/\sin(x))^{(1/2)})-\ln(-(2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)}*\sin(x)+\sin(x)-\cos(x)+1)/(2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)}*\sin(x)-\sin(x)+\cos(x)-1))-4*\arctan(2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)}+1)-4*\arctan(2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)}-1)-\ln(-(2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)}*\sin(x)-\sin(x)+\cos(x)-1)/(2^{(1/2)}*(-(-1+\cos(x))/\sin(x))^{(1/2)}*\sin(x)+\sin(x)-\cos(x)+1)))*(1-\cos(x)+\sin(x))/(a*(\sin(x)+1)/\sin(x))^{(1/2)}/\sin(x)/(-(-1+\cos(x))/\sin(x))^{(1/2)}$$

Maxima [A] time = 1.57662, size = 112, normalized size = 1.81

$$\frac{\sqrt{2}\left(\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+2\sqrt{\frac{\sin(x)}{\cos(x)+1}}\right)\right)+\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}-2\sqrt{\frac{\sin(x)}{\cos(x)+1}}\right)\right)\right)}{\sqrt{a}}-\frac{2\sqrt{2}\arctan\left(\sqrt{\frac{\sin(x)}{\cos(x)+1}}\right)}{\sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(x))^(1/2),x, algorithm="maxima")

[Out]
$$\frac{\sqrt{2}*\left(\sqrt{2}*\arctan\left(\frac{1}{2}*\sqrt{2}*\left(\sqrt{2}+2*\sqrt{\frac{\sin(x)}{\cos(x)+1}}\right)\right)+\sqrt{2}*\arctan\left(-\frac{1}{2}*\sqrt{2}*\left(\sqrt{2}-2*\sqrt{\frac{\sin(x)}{\cos(x)+1}}\right)\right)\right)}{\sqrt{a}}-\frac{2*\sqrt{2}*\arctan\left(\sqrt{\frac{\sin(x)}{\cos(x)+1}}\right)}{\sqrt{a}}$$

Fricas [A] time = 0.506052, size = 668, normalized size = 10.77

$$\frac{\sqrt{2a}\sqrt{-\frac{1}{a}}\log\left(\frac{\sqrt{2}\sqrt{\frac{a\sin(x)+a}{\sin(x)}}\sqrt{-\frac{1}{a}}\sin(x)+\cos(x)}{\sin(x)+1}\right)-\sqrt{-a}\log\left(\frac{2a\cos(x)^2+2(\cos(x)^2+(\cos(x)+1)\sin(x)-1)\sqrt{-a}\sqrt{\frac{a\sin(x)+a}{\sin(x)}}+a\cos(x)-(2a\cos(x)+a)}{\cos(x)+\sin(x)+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(x))^(1/2),x, algorithm="fricas")

[Out]
$$\left[\left(\sqrt{2}*a*\sqrt{-1/a}*\log\left(\sqrt{2}*\sqrt{\frac{a*\sin(x)+a}{\sin(x)}}*\sqrt{-1/a}*\sin(x)+\cos(x)\right)/\left(\sin(x)+1\right)\right)-\sqrt{-a}*\log\left(2*a*\cos(x)^2+2*\cos(x)^2+\right)$$


```
(cos(x) + 1)*sin(x) - 1)*sqrt(-a)*sqrt((a*sin(x) + a)/sin(x)) + a*cos(x) -
(2*a*cos(x) + a)*sin(x) - a)/(cos(x) + sin(x) + 1))/a, -2*(sqrt(2)*sqrt(a)
)*arctan(sqrt(2)*sqrt((a*sin(x) + a)/sin(x))*sin(x)/(sqrt(a)*(cos(x) + sin(
x) + 1))) - sqrt(a)*arctan(-sqrt(a)*sqrt((a*sin(x) + a)/sin(x))*(cos(x) - s
in(x) + 1)/(a*cos(x) + a*sin(x) + a)))/a]
```

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{\sqrt{a \csc(x) + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(x))**(1/2),x)

[Out] Integral(1/sqrt(a*csc(x) + a), x)

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{\sqrt{a \csc(x) + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(x))^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(a*csc(x) + a), x)

$$3.17 \quad \int \frac{1}{(a+a \csc(x))^{3/2}} dx$$

Optimal. Leaf size=81

$$-\frac{2 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x)+a}}\right)}{a^{3/2}} + \frac{5 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{2}\sqrt{a \csc(x)+a}}\right)}{2\sqrt{2}a^{3/2}} + \frac{\cot(x)}{2(a \csc(x) + a)^{3/2}}$$

[Out] $(-2*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])/\text{Sqrt}[a + a*\text{Csc}[x]])/a^{(3/2)} + (5*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])]/(\text{Sqrt}[2]*\text{Sqrt}[a + a*\text{Csc}[x]]))/((2*\text{Sqrt}[2]*a^{(3/2)}) + \text{Cot}[x]/(2*(a + a*\text{Csc}[x])^{(3/2)})$

Rubi [A] time = 0.101898, antiderivative size = 81, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 5, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.5$, Rules used = {3777, 3920, 3774, 203, 3795}

$$-\frac{2 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x)+a}}\right)}{a^{3/2}} + \frac{5 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{2}\sqrt{a \csc(x)+a}}\right)}{2\sqrt{2}a^{3/2}} + \frac{\cot(x)}{2(a \csc(x) + a)^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + a*\text{Csc}[x])^{(-3/2)}, x]$

[Out] $(-2*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])/\text{Sqrt}[a + a*\text{Csc}[x]])/a^{(3/2)} + (5*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])]/(\text{Sqrt}[2]*\text{Sqrt}[a + a*\text{Csc}[x]]))/((2*\text{Sqrt}[2]*a^{(3/2)}) + \text{Cot}[x]/(2*(a + a*\text{Csc}[x])^{(3/2)})$

Rule 3777

$\text{Int}[(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^{(n_.)}, x_Symbol] \text{ :> } -\text{Simp}[(\text{Cot}[c + d*x]*(a + b*\text{Csc}[c + d*x])^n)/(d*(2*n + 1)), x] + \text{Dist}[1/(a^2*(2*n + 1)), \text{Int}[(a + b*\text{Csc}[c + d*x])^{(n + 1)}*(a*(2*n + 1) - b*(n + 1)*\text{Csc}[c + d*x]), x], x] /;$ FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 3920

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))/\text{Sqrt}[\csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] \text{ :> } \text{Dist}[c/a, \text{Int}[\text{Sqrt}[a + b*\text{Csc}[e + f*x]], x], x] - \text{Dist}[(b*c - a*d)/a, \text{Int}[\text{Csc}[e + f*x]/\text{Sqrt}[a + b*\text{Csc}[e + f*x]], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

$eQ[\{a, b, c, d, e, f\}, x] \&\& NeQ[b*c - a*d, 0] \&\& EqQ[a^2 - b^2, 0]$

Rule 3774

$Int[Sqrt[csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] \rightarrow Dist[(-2*b)/d, Subst[Int[1/(a + x^2), x], x, (b*Cot[c + d*x])/Sqrt[a + b*Csc[c + d*x]]], x] /;$ FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]

Rule 203

$Int[((a_.) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /;$ FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 3795

$Int[csc[(e_.) + (f_.)*(x_.)]/Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] \rightarrow Dist[-2/f, Subst[Int[1/(2*a + x^2), x], x, (b*Cot[e + f*x])/Sqrt[a + b*Csc[e + f*x]]], x] /;$ FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + a \csc(x))^{3/2}} dx &= \frac{\cot(x)}{2(a + a \csc(x))^{3/2}} - \frac{\int \frac{-2a + \frac{1}{2}a \csc(x)}{\sqrt{a + a \csc(x)}} dx}{2a^2} \\ &= \frac{\cot(x)}{2(a + a \csc(x))^{3/2}} + \frac{\int \sqrt{a + a \csc(x)} dx}{a^2} - \frac{5 \int \frac{\csc(x)}{\sqrt{a + a \csc(x)}} dx}{4a} \\ &= \frac{\cot(x)}{2(a + a \csc(x))^{3/2}} - \frac{2 \operatorname{Subst}\left(\int \frac{1}{a+x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}}\right)}{a} + \frac{5 \operatorname{Subst}\left(\int \frac{1}{2a+x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}}\right)}{2a} \\ &= -\frac{2 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a + a \csc(x)}}\right)}{a^{3/2}} + \frac{5 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{2}\sqrt{a + a \csc(x)}}\right)}{2\sqrt{2}a^{3/2}} + \frac{\cot(x)}{2(a + a \csc(x))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.410785, size = 129, normalized size = 1.59

$$\frac{\left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right)\left(-2 \csc(x) + 8\sqrt{\csc(x) - 1}(\csc(x) + 1) \tan^{-1}\left(\sqrt{\csc(x) - 1}\right) - 5\sqrt{2}\sqrt{\csc(x) - 1} \csc(x)\left(\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)\right)\right)}{4a(\csc(x) - 1)\sqrt{a}(\csc(x) + 1)\left(\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)\right)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + a*Csc[x])^(-3/2), x]
```

```
[Out] -((Cos[x/2] - Sin[x/2])*(2 - 2*Csc[x] + 8*ArcTan[Sqrt[-1 + Csc[x]]]*Sqrt[-1
+ Csc[x]]*(1 + Csc[x]) - 5*Sqrt[2]*ArcTan[Sqrt[-1 + Csc[x]]/Sqrt[2]]*Sqrt[
-1 + Csc[x]]*Csc[x]*(Cos[x/2] + Sin[x/2])^2))/(4*a*(-1 + Csc[x])*Sqrt[a*(1
+ Csc[x])]*(Cos[x/2] + Sin[x/2]))
```

Maple [B] time = 0.184, size = 1141, normalized size = 14.1

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(a+a*csc(x))^(3/2), x)
```

```
[Out] 1/8*2^(1/2)*(-1+cos(x))*(-2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)
*cos(x)*(-(-1+cos(x))/sin(x))^(3/2)*2^(1/2)-sin(x)*cos(x)*(-(-1+cos(x))/sin
(x))^(1/2)*2^(1/2)-10*sin(x)*cos(x)*arctan((-(-1+cos(x))/sin(x))^(1/2))*2^(
1/2)-4*sin(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)
-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1))-4*sin(x)*
ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1)/(2^(1/2)*
(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1))-16*sin(x)*arctan(2^(1/2)
)*(-(-1+cos(x))/sin(x))^(1/2)+1)+8*cos(x)*arctan(2^(1/2)*(-(-1+cos(x))/sin(
x))^(1/2)-1)+2*cos(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)
)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1))+2
*cos(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1)/(2
^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1))+8*cos(x)*arctan
(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1)+(-(-1+cos(x))/sin(x))^(3/2)*2^(1/2)
-4*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1)/(2^(1/2)
)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1))-4*ln(-(2^(1/2)*(-(-1
+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x)
))^(1/2)*sin(x)-sin(x)+cos(x)-1))-16*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(
1/2)+1)-16*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)-1)+8*cos(x)^2*arctan(
2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)-1)+2*cos(x)^2*ln(-(2^(1/2)*(-(-1+cos(x)
)/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)
)*sin(x)+sin(x)-cos(x)+1))+2*cos(x)^2*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1
/2)*sin(x)+sin(x)-cos(x)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin
(x)+cos(x)-1))+8*cos(x)^2*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1)-16*
sin(x)*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)-1)-10*cos(x)^2*arctan((-(-
1+cos(x))/sin(x))^(1/2))*2^(1/2)+20*sin(x)*arctan((-(-1+cos(x))/sin(x))^(1
/2))*2^(1/2)-10*cos(x)*arctan((-(-1+cos(x))/sin(x))^(1/2))*2^(1/2)-2^(1/2)*
(-(-1+cos(x))/sin(x))^(1/2)+20*2^(1/2)*arctan((-(-1+cos(x))/sin(x))^(1/2))-
```

$$\cos(x)^2 * (-(-1 + \cos(x)) / \sin(x))^{3/2} * 2^{1/2} + \sin(x) * (-(-1 + \cos(x)) / \sin(x))^{3/2} * 2^{1/2} + \cos(x)^2 * (-(-1 + \cos(x)) / \sin(x))^{1/2} * 2^{1/2} + 8 * \sin(x) * \cos(x) * \arctan(2^{1/2} * (-(-1 + \cos(x)) / \sin(x))^{1/2} - 1) + 2 * \sin(x) * \cos(x) * \ln(-2^{1/2} * (-(-1 + \cos(x)) / \sin(x))^{1/2} * \sin(x) - \sin(x) + \cos(x) - 1) / (2^{1/2} * (-(-1 + \cos(x)) / \sin(x))^{1/2} * \sin(x) + \sin(x) - \cos(x) + 1)) + 2 * \sin(x) * \cos(x) * \ln(-2^{1/2} * (-(-1 + \cos(x)) / \sin(x))^{1/2} * \sin(x) + \sin(x) - \cos(x) + 1) / (2^{1/2} * (-(-1 + \cos(x)) / \sin(x))^{1/2} * \sin(x) - \sin(x) + \cos(x) - 1)) + 8 * \sin(x) * \cos(x) * \arctan(2^{1/2} * (-(-1 + \cos(x)) / \sin(x))^{1/2} + 1) / (a * (\sin(x) + 1) / \sin(x))^{3/2} / \sin(x)^3 / (-(-1 + \cos(x)) / \sin(x))^{3/2})$$

Maxima [B] time = 1.54373, size = 203, normalized size = 2.51

$$\frac{\sqrt{2} \left(\frac{\sin(x)}{\cos(x)+1} \right)^{\frac{3}{2}} - \sqrt{2} \sqrt{\frac{\sin(x)}{\cos(x)+1}} + \frac{\sqrt{2} \left(\sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + 2 \sqrt{\frac{\sin(x)}{\cos(x)+1}} \right) \right) + \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} \left(\sqrt{2} - 2 \sqrt{\frac{\sin(x)}{\cos(x)+1}} \right) \right) \right)}{2 \left(a^{\frac{3}{2}} + \frac{2a^{\frac{3}{2}} \sin(x)}{\cos(x)+1} + \frac{a^{\frac{3}{2}} \sin(x)^2}{(\cos(x)+1)^2} \right) + \frac{a^{\frac{3}{2}}}{a^{\frac{3}{2}}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(x))^(3/2),x, algorithm="maxima")

[Out] $-1/2 * (\sqrt{2} * (\sin(x) / (\cos(x) + 1))^{3/2} - \sqrt{2} * \sqrt{\sin(x) / (\cos(x) + 1)}) / (a^{3/2} + 2 * a^{3/2} * \sin(x) / (\cos(x) + 1) + a^{3/2} * \sin(x)^2 / (\cos(x) + 1)^2) + \sqrt{2} * (\sqrt{2} * \arctan(1/2 * \sqrt{2} * (\sqrt{2} + 2 * \sqrt{\sin(x) / (\cos(x) + 1)}))) + \sqrt{2} * \arctan(-1/2 * \sqrt{2} * (\sqrt{2} - 2 * \sqrt{\sin(x) / (\cos(x) + 1)}))) / a^{3/2} - 5/2 * \sqrt{2} * \arctan(\sqrt{\sin(x) / (\cos(x) + 1)}) / a^{3/2}$

Fricas [B] time = 0.532199, size = 1299, normalized size = 16.04

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(x))^(3/2),x, algorithm="fricas")

[Out] $[-1/4 * (5 * \sqrt{2} * (\cos(x)^2 - (\cos(x) + 2) * \sin(x) - \cos(x) - 2) * \sqrt{-a}) * \log(-(\sqrt{2} * \sqrt{-a} * \sqrt{(a * \sin(x) + a) / \sin(x)} * \sin(x) - a * \cos(x)) / (\sin(x) + 1)) + 4 * (\cos(x)^2 - (\cos(x) + 2) * \sin(x) - \cos(x) - 2) * \sqrt{-a}) * \log((2 * a * \cos(x)^2 + 2 * (\cos(x)^2 + (\cos(x) + 1) * \sin(x) - 1) * \sqrt{-a}) * \sqrt{(a * \sin(x) +$

$$a/\sin(x)) + a\cos(x) - (2a\cos(x) + a)\sin(x) - a)/(\cos(x) + \sin(x) + 1) + 2*(\cos(x)^2 + (\cos(x) + 1)\sin(x) - 1)*\sqrt{(a\sin(x) + a)/\sin(x))}/(a^2*\cos(x)^2 - a^2*\cos(x) - 2*a^2 - (a^2*\cos(x) + 2*a^2)*\sin(x)), 1/2*(5*\sqrt{2}*(\cos(x)^2 - (\cos(x) + 2)*\sin(x) - \cos(x) - 2)*\sqrt{a}*\arctan(\sqrt{2}*\sqrt{a}*\sqrt{(a\sin(x) + a)/\sin(x))}*(\cos(x) + 1)/(a*\cos(x) + a*\sin(x) + a)) + 4*(\cos(x)^2 - (\cos(x) + 2)*\sin(x) - \cos(x) - 2)*\sqrt{a}*\arctan(-\sqrt{a}*\sqrt{(a\sin(x) + a)/\sin(x))}*(\cos(x) - \sin(x) + 1)/(a*\cos(x) + a*\sin(x) + a)) - (\cos(x)^2 + (\cos(x) + 1)\sin(x) - 1)*\sqrt{(a\sin(x) + a)/\sin(x))}/(a^2*\cos(x)^2 - a^2*\cos(x) - 2*a^2 - (a^2*\cos(x) + 2*a^2)*\sin(x))]$$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{(a \csc(x) + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(x))**(3/2),x)

[Out] Integral((a*csc(x) + a)**(-3/2), x)

Giac [B] time = 2.9728, size = 410, normalized size = 5.06

$$-\frac{1}{2} \left(\frac{5\sqrt{2} \arctan\left(\frac{\sqrt{a \tan\left(\frac{1}{2}x\right)}}{\sqrt{a}}\right)}{a^{\frac{3}{2}} \operatorname{sgn}\left(\tan\left(\frac{1}{2}x\right) + 1\right)} - \frac{2\left(a\sqrt{|a|} + |a|^{\frac{3}{2}}\right) \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\sqrt{|a|} + 2\sqrt{a \tan\left(\frac{1}{2}x\right)}\right)}{2\sqrt{|a|}}\right)}{a^3 \operatorname{sgn}\left(\tan\left(\frac{1}{2}x\right) + 1\right)} - \frac{2\left(a\sqrt{|a|} + |a|^{\frac{3}{2}}\right) \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\sqrt{|a|} - 2\sqrt{a \tan\left(\frac{1}{2}x\right)}\right)}{2\sqrt{|a|}}\right)}{a^3 \operatorname{sgn}\left(\tan\left(\frac{1}{2}x\right) + 1\right)} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*csc(x))^(3/2),x, algorithm="giac")

[Out] -1/2*(5*sqrt(2)*arctan(sqrt(a*tan(1/2*x))/sqrt(a))/(a^(3/2)*sgn(tan(1/2*x) + 1)) - 2*(a*sqrt(abs(a)) + abs(a)^(3/2))*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(abs(a)) + 2*sqrt(a*tan(1/2*x)))/sqrt(abs(a)))/(a^3*sgn(tan(1/2*x) + 1)) - 2*(a*sqrt(abs(a)) + abs(a)^(3/2))*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(abs(a)) - 2*sqrt(a*tan(1/2*x)))/sqrt(abs(a)))/(a^3*sgn(tan(1/2*x) + 1)) - (a*sqrt(a

$$\begin{aligned} & \text{bs}(a) - \text{abs}(a)^{(3/2)} * \log(a * \tan(1/2 * x) + \text{sqrt}(2) * \text{sqrt}(a * \tan(1/2 * x)) * \text{sqrt}(a \\ & \text{bs}(a) + \text{abs}(a)) / (a^3 * \text{sgn}(\tan(1/2 * x) + 1)) + (a * \text{sqrt}(\text{abs}(a)) - \text{abs}(a)^{(3/2)} \\ &) * \log(a * \tan(1/2 * x) - \text{sqrt}(2) * \text{sqrt}(a * \tan(1/2 * x)) * \text{sqrt}(\text{abs}(a)) + \text{abs}(a)) / (a^3 \\ & * \text{sgn}(\tan(1/2 * x) + 1)) + \text{sqrt}(2) * (\text{sqrt}(a * \tan(1/2 * x)) * a * \tan(1/2 * x) - \text{sqrt}(a * \tan \\ & (1/2 * x)) * a) / ((a * \tan(1/2 * x) + a)^2 * a * \text{sgn}(\tan(1/2 * x) + 1)) * \text{sgn}(\tan(1/2 * x)) \end{aligned}$$

$$3.18 \quad \int \frac{1}{(a+a \csc(x))^{5/2}} dx$$

Optimal. Leaf size=100

$$-\frac{2 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x)+a}}\right)}{a^{5/2}} + \frac{43 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{2}\sqrt{a \csc(x)+a}}\right)}{16\sqrt{2}a^{5/2}} + \frac{11 \cot(x)}{16a(a \csc(x) + a)^{3/2}} + \frac{\cot(x)}{4(a \csc(x) + a)^{5/2}}$$

[Out] $(-2*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])/\text{Sqrt}[a + a*\text{Csc}[x]])/a^{(5/2)} + (43*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])/(\text{Sqrt}[2]*\text{Sqrt}[a + a*\text{Csc}[x]])]/(16*\text{Sqrt}[2]*a^{(5/2)}) + \text{Cot}[x]/(4*(a + a*\text{Csc}[x])^{(5/2)}) + (11*\text{Cot}[x])/(16*a*(a + a*\text{Csc}[x])^{(3/2)})$

Rubi [A] time = 0.148174, antiderivative size = 100, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.6$, Rules used = {3777, 3922, 3920, 3774, 203, 3795}

$$-\frac{2 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{a \csc(x)+a}}\right)}{a^{5/2}} + \frac{43 \tan^{-1}\left(\frac{\sqrt{a} \cot(x)}{\sqrt{2}\sqrt{a \csc(x)+a}}\right)}{16\sqrt{2}a^{5/2}} + \frac{11 \cot(x)}{16a(a \csc(x) + a)^{3/2}} + \frac{\cot(x)}{4(a \csc(x) + a)^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + a*\text{Csc}[x])^{(-5/2)}, x]$

[Out] $(-2*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])/\text{Sqrt}[a + a*\text{Csc}[x]])/a^{(5/2)} + (43*\text{ArcTan}[(\text{Sqrt}[a]*\text{Cot}[x])/(\text{Sqrt}[2]*\text{Sqrt}[a + a*\text{Csc}[x]])]/(16*\text{Sqrt}[2]*a^{(5/2)}) + \text{Cot}[x]/(4*(a + a*\text{Csc}[x])^{(5/2)}) + (11*\text{Cot}[x])/(16*a*(a + a*\text{Csc}[x])^{(3/2)})$

Rule 3777

$\text{Int}[(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(\text{Cot}[c + d*x]*(a + b*\text{Csc}[c + d*x])^n)/(d*(2*n + 1)), x] + \text{Dist}[1/(a^{2*(2*n + 1)}), \text{Int}[(a + b*\text{Csc}[c + d*x])^{(n + 1)}*(a*(2*n + 1) - b*(n + 1)*\text{Csc}[c + d*x]), x], x] /;$ FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 3922

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^{(m_.)}*(\csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)), x_Symbol] \rightarrow -\text{Simp}[(b*c - a*d)*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^m)/(b*f*(2*m + 1)), x] + \text{Dist}[1/(a^{2*(2*m + 1)}), \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m + 1)}*\text{Simp}[a*c*(2*m + 1) - (b*c - a*d)*(m + 1)*\text{Csc}[e + f*x], x], x], x]$

] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 - b^2, 0] && IntegerQ[2*m]

Rule 3920

Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Dist[c/a, Int[Sqrt[a + b*Csc[e + f*x]], x], x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/Sqrt[a + b*Csc[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0]

Rule 3774

Int[Sqrt[csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Dist[(-2*b)/d, Subst[Int[1/(a + x^2), x], x, (b*Cot[c + d*x])/Sqrt[a + b*Csc[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]

Rule 203

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 3795

Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Dist[-2/f, Subst[Int[1/(2*a + x^2), x], x, (b*Cot[e + f*x])/Sqrt[a + b*Csc[e + f*x]]], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a + a \csc(x))^{5/2}} dx &= \frac{\cot(x)}{4(a + a \csc(x))^{5/2}} - \frac{\int \frac{-4a + \frac{3}{2}a \csc(x)}{(a + a \csc(x))^{3/2}} dx}{4a^2} \\
&= \frac{\cot(x)}{4(a + a \csc(x))^{5/2}} + \frac{11 \cot(x)}{16a(a + a \csc(x))^{3/2}} + \frac{\int \frac{8a^2 - \frac{11}{4}a^2 \csc(x)}{\sqrt{a + a \csc(x)}} dx}{8a^4} \\
&= \frac{\cot(x)}{4(a + a \csc(x))^{5/2}} + \frac{11 \cot(x)}{16a(a + a \csc(x))^{3/2}} + \frac{\int \sqrt{a + a \csc(x)} dx}{a^3} - \frac{43 \int \frac{\csc(x)}{\sqrt{a + a \csc(x)}} dx}{32a^2} \\
&= \frac{\cot(x)}{4(a + a \csc(x))^{5/2}} + \frac{11 \cot(x)}{16a(a + a \csc(x))^{3/2}} - \frac{2 \text{Subst} \left(\int \frac{1}{a+x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}} \right)}{a^2} + \frac{43 \text{Subst} \left(\int \frac{1}{2a-x^2} dx, x, \frac{a \cot(x)}{\sqrt{a + a \csc(x)}} \right)}{32a^2} \\
&= -\frac{2 \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{a + a \csc(x)}} \right)}{a^{5/2}} + \frac{43 \tan^{-1} \left(\frac{\sqrt{a} \cot(x)}{\sqrt{2} \sqrt{a + a \csc(x)}} \right)}{16\sqrt{2}a^{5/2}} + \frac{\cot(x)}{4(a + a \csc(x))^{5/2}} + \frac{11 \cot(x)}{16a(a + a \csc(x))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.503013, size = 139, normalized size = 1.39

$$\frac{\csc^2(x) \left(\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right) \right) \left(8 \sin(x) + 15 \cos(2x) - 64 \sqrt{\csc(x) - 1} \left(\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right) \right)^4 \tan^{-1} \left(\sqrt{\csc(x) - 1} \right) + 43 \sqrt{2} \sqrt{\csc(x) - 1} \right)}{32(a(\csc(x) + 1))^{5/2} \left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right) \right)}$$

Antiderivative was successfully verified.

[In] Integrate[(a + a*Csc[x])^(-5/2), x]

[Out] (Csc[x]^2*(Cos[x/2] + Sin[x/2])*(7 + 15*Cos[2*x] - 64*ArcTan[Sqrt[-1 + Csc[x]]]*Sqrt[-1 + Csc[x]]*(Cos[x/2] + Sin[x/2])^4 + 43*Sqrt[2]*ArcTan[Sqrt[-1 + Csc[x]]/Sqrt[2]]*Sqrt[-1 + Csc[x]]*(Cos[x/2] + Sin[x/2])^4 + 8*Sin[x]))/(32*(a*(1 + Csc[x]))^(5/2)*(Cos[x/2] - Sin[x/2]))

Maple [B] time = 0.148, size = 1961, normalized size = 19.6

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+a*csc(x))^(5/2), x)

```
[Out] 1/128*2^(1/2)*(-1+cos(x))^2*(11*2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+
38*sin(x)*cos(x)*(-(-1+cos(x))/sin(x))^(3/2)*2^(1/2)+22*sin(x)*cos(x)*(-(-1
+cos(x))/sin(x))^(1/2)*2^(1/2)+344*sin(x)*cos(x)*arctan((-(-1+cos(x))/sin(x)
))^(1/2))*2^(1/2)+128*sin(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)
)-sin(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)
)+1))+128*sin(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(
x)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1))+512*si
n(x)*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1)-256*cos(x)*arctan(2^(1/2
))*(-(-1+cos(x))/sin(x))^(1/2)-1)-64*cos(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x)
))^(1/2)*sin(x)-sin(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)
)+sin(x)-cos(x)+1))-64*cos(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(
x)+sin(x)-cos(x)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(
x)-1))-256*cos(x)*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1)+19*(-(-1+co
s(x))/sin(x))^(3/2)*2^(1/2)+11*cos(x)^2*sin(x)*(-(-1+cos(x))/sin(x))^(1/2)*
2^(1/2)+172*cos(x)^2*sin(x)*2^(1/2)*arctan((-(-1+cos(x))/sin(x))^(1/2))-11*
cos(x)^2*sin(x)*(-(-1+cos(x))/sin(x))^(7/2)*2^(1/2)-19*cos(x)^2*sin(x)*(-(-
1+cos(x))/sin(x))^(5/2)*2^(1/2)+19*cos(x)^2*sin(x)*(-(-1+cos(x))/sin(x))^(3
/2)*2^(1/2)-38*cos(x)*sin(x)*(-(-1+cos(x))/sin(x))^(5/2)*2^(1/2)-22*cos(x)*
sin(x)*(-(-1+cos(x))/sin(x))^(7/2)*2^(1/2)+128*ln(-(2^(1/2)*(-(-1+cos(x))/s
in(x))^(1/2)*sin(x)-sin(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*s
in(x)+sin(x)-cos(x)+1))+128*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)
+sin(x)-cos(x)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)
-1))+512*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1)+512*arctan(2^(1/2)*(-
(-1+cos(x))/sin(x))^(1/2)-1)-384*cos(x)^2*arctan(2^(1/2)*(-(-1+cos(x))/sin
(x))^(1/2)-1)-96*cos(x)^2*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-s
in(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1
))-96*cos(x)^2*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)
)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1))-384*cos(
x)^2*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1)+512*sin(x)*arctan(2^(1/2
))*(-(-1+cos(x))/sin(x))^(1/2)-1)+32*cos(x)^3*ln(-(2^(1/2)*(-(-1+cos(x))/sin
(x))^(1/2)*sin(x)-sin(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin
(x)+sin(x)-cos(x)+1))+32*cos(x)^3*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*
sin(x)+sin(x)-cos(x)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+
cos(x)-1))+128*cos(x)^3*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1)+128*c
os(x)^3*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)-1)-19*(-(-1+cos(x))/sin(
x))^(5/2)*2^(1/2)-11*(-(-1+cos(x))/sin(x))^(7/2)*2^(1/2)+516*cos(x)^2*arcta
n((-(-1+cos(x))/sin(x))^(1/2))*2^(1/2)-688*sin(x)*arctan((-(-1+cos(x))/sin(
x))^(1/2))*2^(1/2)+344*cos(x)*arctan((-(-1+cos(x))/sin(x))^(1/2))*2^(1/2)-1
72*cos(x)^3*2^(1/2)*arctan((-(-1+cos(x))/sin(x))^(1/2))-11*cos(x)^3*(-(-1+c
os(x))/sin(x))^(1/2)*2^(1/2)+19*cos(x)*(-(-1+cos(x))/sin(x))^(3/2)*2^(1/2)+
11*cos(x)*(-(-1+cos(x))/sin(x))^(1/2)*2^(1/2)-32*cos(x)^2*sin(x)*ln(-(2^(1/
2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x)
))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1))-32*cos(x)^2*sin(x)*ln(-(2^(1/2)*(-
(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1)/(2^(1/2)*(-(-1+cos(x))/s
in(x))^(1/2)*sin(x)-sin(x)+cos(x)-1))-128*cos(x)^2*sin(x)*arctan(2^(1/2)*(-
```

```
(-1+cos(x))/sin(x))^(1/2)+1)-128*cos(x)^2*sin(x)*arctan(2^(1/2)*(-(-1+cos(x)))/sin(x))^(1/2)-1)+11*cos(x)^3*(-(-1+cos(x))/sin(x))^(7/2)*2^(1/2)+11*cos(x)^2*(-(-1+cos(x))/sin(x))^(7/2)*2^(1/2)+19*cos(x)^3*(-(-1+cos(x))/sin(x))^(5/2)*2^(1/2)-11*cos(x)*(-(-1+cos(x))/sin(x))^(7/2)*2^(1/2)-11*sin(x)*(-(-1+cos(x))/sin(x))^(7/2)*2^(1/2)+19*cos(x)^2*(-(-1+cos(x))/sin(x))^(5/2)*2^(1/2)-19*cos(x)^3*(-(-1+cos(x))/sin(x))^(3/2)*2^(1/2)-19*cos(x)*(-(-1+cos(x))/sin(x))^(5/2)*2^(1/2)-19*sin(x)*(-(-1+cos(x))/sin(x))^(5/2)*2^(1/2)+11*2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)-688*2^(1/2)*arctan((-(-1+cos(x))/sin(x))^(1/2))-19*cos(x)^2*(-(-1+cos(x))/sin(x))^(3/2)*2^(1/2)+19*sin(x)*(-(-1+cos(x))/sin(x))^(3/2)*2^(1/2)-11*cos(x)^2*(-(-1+cos(x))/sin(x))^(1/2)*2^(1/2)-256*sin(x)*cos(x)*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)-1)-64*sin(x)*cos(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1))-64*sin(x)*cos(x)*ln(-(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)+sin(x)-cos(x)+1)/(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)*sin(x)-sin(x)+cos(x)-1))-256*sin(x)*cos(x)*arctan(2^(1/2)*(-(-1+cos(x))/sin(x))^(1/2)+1))/(a*(sin(x)+1)/sin(x))^(5/2)/sin(x)^5/(-(-1+cos(x))/sin(x))^(5/2)
```

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{(a \csc(x) + a)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a+a*csc(x))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate((a*csc(x) + a)^(-5/2), x)
```

Fricas [B] time = 0.546279, size = 1677, normalized size = 16.77

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a+a*csc(x))^(5/2),x, algorithm="fricas")
```

```
[Out] [-1/32*(43*sqrt(2)*(cos(x)^3 + 3*cos(x)^2 + (cos(x)^2 - 2*cos(x) - 4)*sin(x) - 2*cos(x) - 4)*sqrt(-a)*log(-(sqrt(2)*sqrt(-a)*sqrt((a*sin(x) + a)/sin(x)))*sin(x) - a*cos(x))/(sin(x) + 1)) + 32*(cos(x)^3 + 3*cos(x)^2 + (cos(x)^2
```

```

- 2*cos(x) - 4)*sin(x) - 2*cos(x) - 4)*sqrt(-a)*log((2*a*cos(x)^2 + 2*(cos
(x)^2 + (cos(x) + 1)*sin(x) - 1)*sqrt(-a)*sqrt((a*sin(x) + a)/sin(x)) + a*c
os(x) - (2*a*cos(x) + a)*sin(x) - a)/(cos(x) + sin(x) + 1)) - 2*(15*cos(x)^
3 + 4*cos(x)^2 - (15*cos(x)^2 + 11*cos(x) - 4)*sin(x) - 15*cos(x) - 4)*sqrt
((a*sin(x) + a)/sin(x)))/(a^3*cos(x)^3 + 3*a^3*cos(x)^2 - 2*a^3*cos(x) - 4*
a^3 + (a^3*cos(x)^2 - 2*a^3*cos(x) - 4*a^3)*sin(x)), 1/16*(43*sqrt(2)*(cos(
x)^3 + 3*cos(x)^2 + (cos(x)^2 - 2*cos(x) - 4)*sin(x) - 2*cos(x) - 4)*sqrt(a
)*arctan(sqrt(2)*sqrt(a)*sqrt((a*sin(x) + a)/sin(x))*(cos(x) + 1)/(a*cos(x)
+ a*sin(x) + a)) + 32*(cos(x)^3 + 3*cos(x)^2 + (cos(x)^2 - 2*cos(x) - 4)*s
in(x) - 2*cos(x) - 4)*sqrt(a)*arctan(-sqrt(a)*sqrt((a*sin(x) + a)/sin(x))*
(cos(x) - sin(x) + 1)/(a*cos(x) + a*sin(x) + a)) + (15*cos(x)^3 + 4*cos(x)^2
- (15*cos(x)^2 + 11*cos(x) - 4)*sin(x) - 15*cos(x) - 4)*sqrt((a*sin(x) + a
)/sin(x)))/(a^3*cos(x)^3 + 3*a^3*cos(x)^2 - 2*a^3*cos(x) - 4*a^3 + (a^3*cos
(x)^2 - 2*a^3*cos(x) - 4*a^3)*sin(x))]

```

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{(a \csc(x) + a)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a+a*csc(x))**(5/2),x)
```

```
[Out] Integral((a*csc(x) + a)**(-5/2), x)
```

Giac [B] time = 2.52895, size = 470, normalized size = 4.7

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a+a*csc(x))^(5/2),x, algorithm="giac")
```

```
[Out] -1/16*(43*sqrt(2)*arctan(sqrt(a*tan(1/2*x))/sqrt(a))/sqrt(a))/(a^(5/2)*sgn(tan(1/2*x)
+ 1)) - 16*(a*sqrt(abs(a)) + abs(a)^(3/2))*arctan(1/2*sqrt(2)*(sqrt(2)*sq
rt(abs(a)) + 2*sqrt(a*tan(1/2*x)))/sqrt(abs(a)))/(a^4*sgn(tan(1/2*x) + 1))
- 16*(a*sqrt(abs(a)) + abs(a)^(3/2))*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(abs(
a)) - 2*sqrt(a*tan(1/2*x)))/sqrt(abs(a)))/(a^4*sgn(tan(1/2*x) + 1)) - 8*(a*
```

```

sqrt(abs(a)) - abs(a)^(3/2))*log(a*tan(1/2*x) + sqrt(2)*sqrt(a*tan(1/2*x))*
sqrt(abs(a)) + abs(a))/(a^4*sgn(tan(1/2*x) + 1)) + 8*(a*sqrt(abs(a)) - abs(
a)^(3/2))*log(a*tan(1/2*x) - sqrt(2)*sqrt(a*tan(1/2*x))*sqrt(abs(a)) + abs(
a))/(a^4*sgn(tan(1/2*x) + 1)) + sqrt(2)*(11*sqrt(a*tan(1/2*x))*a^3*tan(1/2*
x)^3 + 19*sqrt(a*tan(1/2*x))*a^3*tan(1/2*x)^2 - 19*sqrt(a*tan(1/2*x))*a^3*t
an(1/2*x) - 11*sqrt(a*tan(1/2*x))*a^3)/((a*tan(1/2*x) + a)^4*a^2*sgn(tan(1/
2*x) + 1))*sgn(tan(1/2*x))

```

$$3.19 \quad \int \sqrt{\csc(e + fx)} \sqrt{a + a \csc(e + fx)} dx$$

Optimal. Leaf size=37

$$-\frac{2\sqrt{a} \sinh^{-1}\left(\frac{\sqrt{a} \cot(e+fx)}{\sqrt{a \csc(e+fx)+a}}\right)}{f}$$

[Out] $(-2*\text{Sqrt}[a]*\text{ArcSinh}[(\text{Sqrt}[a]*\text{Cot}[e + f*x])/\text{Sqrt}[a + a*\text{Csc}[e + f*x]]])/f$

Rubi [A] time = 0.0585123, antiderivative size = 37, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.08$, Rules used = {3801, 215}

$$-\frac{2\sqrt{a} \sinh^{-1}\left(\frac{\sqrt{a} \cot(e+fx)}{\sqrt{a \csc(e+fx)+a}}\right)}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[\text{Csc}[e + f*x]]*\text{Sqrt}[a + a*\text{Csc}[e + f*x]], x]$

[Out] $(-2*\text{Sqrt}[a]*\text{ArcSinh}[(\text{Sqrt}[a]*\text{Cot}[e + f*x])/\text{Sqrt}[a + a*\text{Csc}[e + f*x]]])/f$

Rule 3801

$\text{Int}[\text{Sqrt}[\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.)]*\text{Sqrt}[\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] \rightarrow \text{Dist}[(-2*a*\text{Sqrt}[(a*d)/b])/(b*f), \text{Subst}[\text{Int}[1/\text{Sqrt}[1 + x^2/a], x], x, (b*\text{Cot}[e + f*x])/\text{Sqrt}[a + b*\text{Csc}[e + f*x]]], x] /; \text{FreeQ}\{a, b, d, e, f\}, x] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{GtQ}[(a*d)/b, 0]$

Rule 215

$\text{Int}[1/\text{Sqrt}[(a_.) + (b_.)*(x_.)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[(\text{Rt}[b, 2]*x)/\text{Sqrt}[a]]/\text{Rt}[b, 2], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{GtQ}[a, 0] \&\& \text{PosQ}[b]$

Rubi steps

$$\int \sqrt{\csc(e+fx)} \sqrt{a+a \csc(e+fx)} dx = -\frac{2 \operatorname{Subst} \left(\int \frac{1}{\sqrt{1+\frac{x^2}{a}}} dx, x, \frac{a \cot(e+fx)}{\sqrt{a+a \csc(e+fx)}} \right)}{f}$$

$$= -\frac{2\sqrt{a} \sinh^{-1} \left(\frac{\sqrt{a} \cot(e+fx)}{\sqrt{a+a \csc(e+fx)}} \right)}{f}$$

Mathematica [B] time = 0.408282, size = 108, normalized size = 2.92

$$\frac{2 \cot(e+fx) \sqrt{a(\csc(e+fx)+1)} \left(\log(\csc(e+fx)+1) - \log \left(\csc^3(e+fx) + \sqrt{\csc(e+fx)} + \sqrt{\cot^2(e+fx)} \sqrt{\csc(e+fx)} \right) \right)}{f \sqrt{\cot^2(e+fx)} \sqrt{\csc(e+fx)+1}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Csc[e + f*x]]*Sqrt[a + a*Csc[e + f*x]],x]

[Out] (2*Cot[e + f*x]*Sqrt[a*(1 + Csc[e + f*x])]*(Log[1 + Csc[e + f*x]] - Log[Sqrt[Csc[e + f*x]] + Csc[e + f*x]^(3/2) + Sqrt[Cot[e + f*x]^2]*Sqrt[1 + Csc[e + f*x]]]))/(f*Sqrt[Cot[e + f*x]^2]*Sqrt[1 + Csc[e + f*x]])

Maple [B] time = 0.359, size = 114, normalized size = 3.1

$$-\frac{\sqrt{2}(-1 + \cos(fx + e))}{f(-1 + \cos(fx + e) - \sin(fx + e))} \sqrt{(\sin(fx + e))^{-1}} \sqrt{\frac{a(\sin(fx + e) + 1)}{\sin(fx + e)}} \left(\operatorname{Arcsinh} \left(\frac{-1 + \cos(fx + e)}{\sin(fx + e)} \right) + \operatorname{Arctanh} \left(\frac{-1 + \cos(fx + e)}{\sin(fx + e)} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^(1/2)*(a+a*csc(f*x+e))^(1/2),x)

[Out] -1/f*2^(1/2)*(1/sin(f*x+e))^(1/2)*(-1+cos(f*x+e))*(a*(sin(f*x+e)+1)/sin(f*x+e))^(1/2)*(arcsinh((-1+cos(f*x+e))/sin(f*x+e))+arctanh(1/2*2^(1/2)/(1/(1+cos(f*x+e))))^(1/2))/(-1+cos(f*x+e)-sin(f*x+e))/(1/(1+cos(f*x+e)))^(1/2)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(fx + e) + a} \sqrt{\csc(fx + e)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^(1/2)*(a+a*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a*csc(f*x + e) + a)*sqrt(csc(f*x + e)), x)

Fricas [B] time = 0.534522, size = 755, normalized size = 20.41

$$\sqrt{a} \log \left(\frac{a \cos(fx+e)^3 - 7a \cos(fx+e)^2 - 9a \cos(fx+e) + (a \cos(fx+e)^2 + 8a \cos(fx+e) - a) \sin(fx+e) + \frac{4(\cos(fx+e)^3 + 3 \cos(fx+e)^2 - (\cos(fx+e)^2 - 2 \cos(fx+e) - 3) \sin(fx+e) - \cos(fx+e) - 3)}{\sqrt{\sin(fx+e)}}}{\cos(fx+e)^3 + \cos(fx+e)^2 + (\cos(fx+e)^2 - 1) \sin(fx+e) - \cos(fx+e) - 1} \right) \frac{1}{2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^(1/2)*(a+a*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] [1/2*sqrt(a)*log((a*cos(f*x + e)^3 - 7*a*cos(f*x + e)^2 - 9*a*cos(f*x + e) + (a*cos(f*x + e)^2 + 8*a*cos(f*x + e) - a)*sin(f*x + e) + 4*(cos(f*x + e)^3 + 3*cos(f*x + e)^2 - (cos(f*x + e)^2 - 2*cos(f*x + e) - 3)*sin(f*x + e) - cos(f*x + e) - 3)*sqrt(a)*sqrt((a*sin(f*x + e) + a)/sin(f*x + e))/sqrt(sin(f*x + e)) - a)/(cos(f*x + e)^3 + cos(f*x + e)^2 + (cos(f*x + e)^2 - 1)*sin(f*x + e) - cos(f*x + e) - 1))/f, sqrt(-a)*arctan(1/2*(cos(f*x + e)^2 + 2*sin(f*x + e) - 1)*sqrt(-a)*sqrt((a*sin(f*x + e) + a)/sin(f*x + e)))/(a*cos(f*x + e)*sqrt(sin(f*x + e))))/f]

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a(\csc(e + fx) + 1)} \sqrt{\csc(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**(1/2)*(a+a*csc(f*x+e))**(1/2),x)`

[Out] `Integral(sqrt(a*(csc(e + f*x) + 1))*sqrt(csc(e + f*x)), x)`

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(fx + e) + a} \sqrt{\csc(fx + e)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^(1/2)*(a+a*csc(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(a*csc(f*x + e) + a)*sqrt(csc(f*x + e)), x)`

$$3.20 \quad \int \sqrt{-\csc(e + fx)} \sqrt{a - a \csc(e + fx)} dx$$

Optimal. Leaf size=38

$$-\frac{2\sqrt{a} \sinh^{-1}\left(\frac{\sqrt{a} \cot(e+fx)}{\sqrt{a-a \csc(e+fx)}}\right)}{f}$$

[Out] $(-2*\text{Sqrt}[a]*\text{ArcSinh}[(\text{Sqrt}[a]*\text{Cot}[e + f*x])/\text{Sqrt}[a - a*\text{Csc}[e + f*x]]])/f$

Rubi [A] time = 0.0708915, antiderivative size = 38, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {3801, 215}

$$-\frac{2\sqrt{a} \sinh^{-1}\left(\frac{\sqrt{a} \cot(e+fx)}{\sqrt{a-a \csc(e+fx)}}\right)}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[-\text{Csc}[e + f*x]]*\text{Sqrt}[a - a*\text{Csc}[e + f*x]], x]$

[Out] $(-2*\text{Sqrt}[a]*\text{ArcSinh}[(\text{Sqrt}[a]*\text{Cot}[e + f*x])/\text{Sqrt}[a - a*\text{Csc}[e + f*x]]])/f$

Rule 3801

$\text{Int}[\text{Sqrt}[\csc[(e_.) + (f_.)*(x_.)]*(d_.)]*\text{Sqrt}[\csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] \rightarrow \text{Dist}[(-2*a*\text{Sqrt}[(a*d)/b])/(b*f), \text{Subst}[\text{Int}[1/\text{Sqrt}[1 + x^2/a], x], x, (b*\text{Cot}[e + f*x])/\text{Sqrt}[a + b*\text{Csc}[e + f*x]]], x] /; \text{FreeQ}\{a, b, d, e, f\}, x] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{GtQ}[(a*d)/b, 0]$

Rule 215

$\text{Int}[1/\text{Sqrt}[(a_.) + (b_.)*(x_.)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[(\text{Rt}[b, 2]*x)/\text{Sqrt}[a]]/\text{Rt}[b, 2], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{GtQ}[a, 0] \&\& \text{PosQ}[b]$

Rubi steps

$$\int \sqrt{-\csc(e+fx)} \sqrt{a-a\csc(e+fx)} dx = \frac{2 \operatorname{Subst} \left(\int \frac{1}{\sqrt{1+\frac{x^2}{a}}} dx, x, -\frac{a \cot(e+fx)}{\sqrt{a-a\csc(e+fx)}} \right)}{f}$$

$$= -\frac{2\sqrt{a} \sinh^{-1} \left(\frac{\sqrt{a} \cot(e+fx)}{\sqrt{a-a\csc(e+fx)}} \right)}{f}$$

Mathematica [B] time = 0.775479, size = 101, normalized size = 2.66

$$\frac{2 \tan\left(\frac{1}{2}(e+fx)\right) \sqrt{-\csc(e+fx)} \sqrt{a-a\csc(e+fx)} \left(\tanh^{-1} \left(\sqrt{\sec^2\left(\frac{1}{2}(e+fx)\right)} \right) + \sinh^{-1} \left(\tan\left(\frac{1}{2}(e+fx)\right) \right) \right)}{f \left(\tan\left(\frac{1}{2}(e+fx)\right) - 1 \right) \sqrt{\sec^2\left(\frac{1}{2}(e+fx)\right)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-Csc[e + f*x]]*Sqrt[a - a*Csc[e + f*x]],x]

[Out] (2*(ArcSinh[Tan[(e + f*x)/2]] + ArcTanh[Sqrt[Sec[(e + f*x)/2]^2]])*Sqrt[-Csc[e + f*x]]*Sqrt[a - a*Csc[e + f*x]]*Tan[(e + f*x)/2])/(f*Sqrt[Sec[(e + f*x)/2]^2]*(-1 + Tan[(e + f*x)/2]))

Maple [B] time = 0.333, size = 117, normalized size = 3.1

$$2 \frac{-1 + \cos(fx + e)}{f(-1 + \cos(fx + e) + \sin(fx + e))} \sqrt{-(\sin(fx + e))^{-1}} \sqrt{\frac{a(\sin(fx + e) - 1)}{\sin(fx + e)}} \left(\arctan\left(\frac{1}{2} \sin(fx + e)\right) \sqrt{-2(1 + \cos(fx + e))} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-csc(f*x+e))^(1/2)*(a-a*csc(f*x+e))^(1/2),x)

[Out] 2/f*(-1/sin(f*x+e))^(1/2)*(-1+cos(f*x+e))*(a*(sin(f*x+e)-1)/sin(f*x+e))^(1/2)*(arctan(1/2*sin(f*x+e)*(-2/(1+cos(f*x+e))))^(1/2))-arctan(1/(-2/(1+cos(f*x+e))))^(1/2))/(-1+cos(f*x+e)+sin(f*x+e))/(-2/(1+cos(f*x+e)))^(1/2)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{-a \csc(fx + e) + a} \sqrt{-\csc(fx + e)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-csc(f*x+e))^(1/2)*(a-a*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-a*csc(f*x + e) + a)*sqrt(-csc(f*x + e)), x)

Fricas [B] time = 0.52994, size = 764, normalized size = 20.11

$$\sqrt{a} \log \left(\frac{a \cos(fx+e)^3 - 7a \cos(fx+e)^2 - 4(\cos(fx+e)^3 + 3 \cos(fx+e)^2 + (\cos(fx+e)^2 - 2 \cos(fx+e) - 3) \sin(fx+e) - \cos(fx+e) - 3) \sqrt{a} \sqrt{\frac{a \sin(fx+e) - a}{\sin(fx+e)}} \sqrt{-\frac{1}{\sin(fx+e)}}}{\cos(fx+e)^3 + \cos(fx+e)^2 - (\cos(fx+e)^2 - 1) \sin(fx+e) - \cos(fx+e) - 1} \right) \frac{1}{2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-csc(f*x+e))^(1/2)*(a-a*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] [1/2*sqrt(a)*log((a*cos(f*x + e)^3 - 7*a*cos(f*x + e)^2 - 4*(cos(f*x + e)^3 + 3*cos(f*x + e)^2 + (cos(f*x + e)^2 - 2*cos(f*x + e) - 3)*sin(f*x + e) - cos(f*x + e) - 3)*sqrt(a)*sqrt((a*sin(f*x + e) - a)/sin(f*x + e))*sqrt(-1/sin(f*x + e)) - 9*a*cos(f*x + e) - (a*cos(f*x + e)^2 + 8*a*cos(f*x + e) - a)*sin(f*x + e) - a)/(cos(f*x + e)^3 + cos(f*x + e)^2 - (cos(f*x + e)^2 - 1)*sin(f*x + e) - cos(f*x + e) - 1))/f, sqrt(-a)*arctan(-1/2*(cos(f*x + e)^2 - 2*sin(f*x + e) - 1)*sqrt(-a)*sqrt((a*sin(f*x + e) - a)/sin(f*x + e))*sqrt(-1/sin(f*x + e)))/(a*cos(f*x + e)))/f]

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{-\csc(e + fx)} \sqrt{-a(\csc(e + fx) - 1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-csc(f*x+e))**(1/2)*(a-a*csc(f*x+e))**(1/2),x)
```

```
[Out] Integral(sqrt(-csc(e + f*x))*sqrt(-a*(csc(e + f*x) - 1)), x)
```

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{-a \csc(fx + e) + a} \sqrt{-\csc(fx + e)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-csc(f*x+e))^(1/2)*(a-a*csc(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(-a*csc(f*x + e) + a)*sqrt(-csc(f*x + e)), x)
```

3.21 $\int \csc^{\frac{4}{3}}(c + dx) \sqrt{a + a \csc(c + dx)} dx$

Optimal. Leaf size=254

$$\frac{4 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c + dx) + \sqrt[3]{\csc(c + dx)} + 1}{(-\sqrt[3]{\csc(c + dx)} + \sqrt{3} + 1)^2}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c + dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c + dx)} + \sqrt{3} + 1}\right), -7 - 4\sqrt{3}\right)}{5d \sqrt{\frac{1 - \sqrt[3]{\csc(c + dx)}}{(-\sqrt[3]{\csc(c + dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c + dx)) \sqrt{a \csc(c + dx) + a}}$$

[Out] $(-6*a*\cos[c + d*x]*\csc[c + d*x]^{(4/3)})/(5*d*\sqrt{a + a*\csc[c + d*x]}) - (4*3^{(3/4)}*\sqrt{2 + \sqrt{3}}*a^2*\cot[c + d*x]*(1 - \csc[c + d*x]^{(1/3)})*\sqrt{(1 + \csc[c + d*x]^{(1/3)} + \csc[c + d*x]^{(2/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})^2}*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \sqrt{3} - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})], -7 - 4*\sqrt{3}])/(5*d*\sqrt{(1 - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})^2}*(a - a*\csc[c + d*x])* \sqrt{a + a*\csc[c + d*x]})$

Rubi [A] time = 0.280558, antiderivative size = 254, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.16$, Rules used = {3806, 50, 63, 218}

$$\frac{4 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c + dx) + \sqrt[3]{\csc(c + dx)} + 1}{(-\sqrt[3]{\csc(c + dx)} + \sqrt{3} + 1)^2}} F\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c + dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c + dx)} + \sqrt{3} + 1}\right) \middle| -7 - 4\sqrt{3}\right)}{5d \sqrt{\frac{1 - \sqrt[3]{\csc(c + dx)}}{(-\sqrt[3]{\csc(c + dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c + dx)) \sqrt{a \csc(c + dx) + a}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\csc[c + d*x]^{(4/3)}*\sqrt{a + a*\csc[c + d*x]}, x]$

[Out] $(-6*a*\cos[c + d*x]*\csc[c + d*x]^{(4/3)})/(5*d*\sqrt{a + a*\csc[c + d*x]}) - (4*3^{(3/4)}*\sqrt{2 + \sqrt{3}}*a^2*\cot[c + d*x]*(1 - \csc[c + d*x]^{(1/3)})*\sqrt{(1 + \csc[c + d*x]^{(1/3)} + \csc[c + d*x]^{(2/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})^2}*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \sqrt{3} - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})], -7 - 4*\sqrt{3}])/(5*d*\sqrt{(1 - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})^2}*(a - a*\csc[c + d*x])* \sqrt{a + a*\csc[c + d*x]})$

Rule 3806

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.)
+ (a_.)], x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]
]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[(d*x)^(n - 1)/Sqrt[a - b*x], x], x,
Csc[e + f*x]], x] /; FreeQ[{a, b, d, e, f, n}, x] && EqQ[a^2 - b^2, 0]
```

Rule 50

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
((a + b*x)^(m + 1)*(c + d*x)^n)/(b*(m + n + 1)), x] + Dist[(n*(b*c - a*d))/
(b*(m + n + 1)), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 63

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - (a*d)/b +
(d*x^p)/b)^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 218

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 + Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 - Sqrt[3])*s
+ r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[(s*(s + r*x))/((1 + Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b}, x] &
& PosQ[a]
```

Rubi steps

$$\begin{aligned}
\int \csc^{\frac{4}{3}}(c+dx) \sqrt{a+a \csc(c+dx)} dx &= \frac{(a^2 \cot(c+dx)) \operatorname{Subst}\left(\int \frac{\sqrt[3]{x}}{\sqrt{a-ax}} dx, x, \csc(c+dx)\right)}{d \sqrt{a-a \csc(c+dx)} \sqrt{a+a \csc(c+dx)}} \\
&= -\frac{6a \cos(c+dx) \csc^{\frac{4}{3}}(c+dx)}{5d \sqrt{a+a \csc(c+dx)}} + \frac{(2a^2 \cot(c+dx)) \operatorname{Subst}\left(\int \frac{1}{x^{2/3} \sqrt{a-ax}} dx, x, \csc(c+dx)\right)}{5d \sqrt{a-a \csc(c+dx)} \sqrt{a+a \csc(c+dx)}} \\
&= -\frac{6a \cos(c+dx) \csc^{\frac{4}{3}}(c+dx)}{5d \sqrt{a+a \csc(c+dx)}} + \frac{(6a^2 \cot(c+dx)) \operatorname{Subst}\left(\int \frac{1}{\sqrt{a-ax^3}} dx, x, \sqrt[3]{\csc(c+dx)}\right)}{5d \sqrt{a-a \csc(c+dx)} \sqrt{a+a \csc(c+dx)}} \\
&= -\frac{6a \cos(c+dx) \csc^{\frac{4}{3}}(c+dx)}{5d \sqrt{a+a \csc(c+dx)}} - \frac{4 \cdot 3^{3/4} \sqrt{2+\sqrt{3}a^2} \cot(c+dx) \left(1 - \sqrt[3]{\csc(c+dx)}\right)}{5d \sqrt{\frac{1-\sqrt[3]{\csc(c+dx)}}{(1+\sqrt{3}-\sqrt[3]{\csc(c+dx)})}}}
\end{aligned}$$

Mathematica [C] time = 0.381845, size = 102, normalized size = 0.4

$$\frac{2\sqrt{a(\csc(c+dx)+1)} \left(\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right) \left(2\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{3}{2}, 1 - \csc(c+dx)\right) + 3\sqrt[3]{\csc(c+dx)}\right)}{5d \left(\sin\left(\frac{1}{2}(c+dx)\right) + \cos\left(\frac{1}{2}(c+dx)\right)\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[c + d*x]^(4/3)*Sqrt[a + a*Csc[c + d*x]],x]

[Out] (-2*Sqrt[a*(1 + Csc[c + d*x])]*(3*Csc[c + d*x]^(1/3) + 2*Hypergeometric2F1[1/2, 2/3, 3/2, 1 - Csc[c + d*x]])*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(5*d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))

Maple [F] time = 0.384, size = 0, normalized size = 0.

$$\int (\csc(dx+c))^{\frac{4}{3}} \sqrt{a+a \csc(dx+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(d*x+c)^(4/3)*(a+a*csc(d*x+c))^(1/2),x)

[Out] `int(csc(d*x+c)^(4/3)*(a+a*csc(d*x+c))^(1/2),x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a \csc(dx + c)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(d*x+c)^(4/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(4/3), x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\sqrt{a \csc(dx + c) + a \csc(dx + c)^{\frac{4}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(d*x+c)^(4/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="fricas")`

[Out] `integral(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(4/3), x)`

Sympy [F(-1)] time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(d*x+c)**(4/3)*(a+a*csc(d*x+c))**(1/2),x)`

[Out] Timed out

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a} \csc(dx + c)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(d*x+c)^(4/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(4/3), x)
```

3.22 $\int \sqrt[3]{\csc(c+dx)} \sqrt{a+a \csc(c+dx)} dx$

Optimal. Leaf size=213

$$\frac{2 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a^2 \cot(c+dx) \left(1 - \sqrt[3]{\csc(c+dx)}\right) \sqrt{\frac{\csc^2(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{\left(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1\right)^2}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right), -7 - 4\sqrt{3}\right)}{d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{\left(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1\right)^2}} (a - a \csc(c+dx)) \sqrt{a \csc(c+dx) + a}}$$

[Out] $(-2 \cdot 3^{3/4} \cdot \operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]] \cdot a^2 \cdot \operatorname{Cot}[c + d \cdot x] \cdot (1 - \operatorname{Csc}[c + d \cdot x]^{1/3}) \cdot \operatorname{Sqrt}[(1 + \operatorname{Csc}[c + d \cdot x]^{1/3} + \operatorname{Csc}[c + d \cdot x]^{2/3}) / (1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d \cdot x]^{1/3})^2] \cdot \operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d \cdot x]^{1/3}) / (1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d \cdot x]^{1/3})], -7 - 4 \cdot \operatorname{Sqrt}[3]]) / (d \cdot \operatorname{Sqrt}[(1 - \operatorname{Csc}[c + d \cdot x]^{1/3}) / (1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d \cdot x]^{1/3})^2] \cdot (a - a \cdot \operatorname{Csc}[c + d \cdot x]) \cdot \operatorname{Sqrt}[a + a \cdot \operatorname{Csc}[c + d \cdot x]])$

Rubi [A] time = 0.127755, antiderivative size = 213, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.12$, Rules used = {3806, 63, 218}

$$\frac{2 \cdot 3^{3/4} \sqrt{2 + \sqrt{3}} a^2 \cot(c+dx) \left(1 - \sqrt[3]{\csc(c+dx)}\right) \sqrt{\frac{\csc^2(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{\left(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1\right)^2}} F\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right) \mid -7 - 4\sqrt{3}\right)}{d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{\left(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1\right)^2}} (a - a \csc(c+dx)) \sqrt{a \csc(c+dx) + a}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[c + d \cdot x]^{1/3} \cdot \operatorname{Sqrt}[a + a \cdot \operatorname{Csc}[c + d \cdot x]], x]$

[Out] $(-2 \cdot 3^{3/4} \cdot \operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]] \cdot a^2 \cdot \operatorname{Cot}[c + d \cdot x] \cdot (1 - \operatorname{Csc}[c + d \cdot x]^{1/3}) \cdot \operatorname{Sqrt}[(1 + \operatorname{Csc}[c + d \cdot x]^{1/3} + \operatorname{Csc}[c + d \cdot x]^{2/3}) / (1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d \cdot x]^{1/3})^2] \cdot \operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d \cdot x]^{1/3}) / (1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d \cdot x]^{1/3})], -7 - 4 \cdot \operatorname{Sqrt}[3]]) / (d \cdot \operatorname{Sqrt}[(1 - \operatorname{Csc}[c + d \cdot x]^{1/3}) / (1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d \cdot x]^{1/3})^2] \cdot (a - a \cdot \operatorname{Csc}[c + d \cdot x]) \cdot \operatorname{Sqrt}[a + a \cdot \operatorname{Csc}[c + d \cdot x]])$

Rule 3806

$\operatorname{Int}[(\operatorname{csc}[(e_.) + (f_.) \cdot (x_.)] \cdot (d_.)^n) \cdot \operatorname{Sqrt}[\operatorname{csc}[(e_.) + (f_.) \cdot (x_.)] \cdot (b_.) + (a_.)], x_Symbol] \rightarrow \operatorname{Dist}[(a^2 \cdot d \cdot \operatorname{Cot}[e + f \cdot x]) / (f \cdot \operatorname{Sqrt}[a + b \cdot \operatorname{Csc}[e + f \cdot x])]$

] * Sqrt[a - b * Csc[e + f * x]], Subst[Int[(d * x)^(n - 1) / Sqrt[a - b * x], x], x, Csc[e + f * x]], x] /; FreeQ[{a, b, d, e, f, n}, x] && EqQ[a^2 - b^2, 0]

Rule 63

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - (a*d)/b + (d*x^p)/b)^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 218

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 + Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(s*(s + r*x))/((1 + Sqrt[3])*s + r*x)^2]), x] /; FreeQ[{a, b}, x] && PosQ[a]

Rubi steps

$$\begin{aligned} \int \sqrt[3]{\csc(c+dx)} \sqrt{a+a \csc(c+dx)} dx &= \frac{(a^2 \cot(c+dx)) \operatorname{Subst}\left(\int \frac{1}{x^{2/3} \sqrt{a-ax}} dx, x, \csc(c+dx)\right)}{d \sqrt{a-a \csc(c+dx)} \sqrt{a+a \csc(c+dx)}} \\ &= \frac{(3a^2 \cot(c+dx)) \operatorname{Subst}\left(\int \frac{1}{\sqrt{a-ax^3}} dx, x, \sqrt[3]{\csc(c+dx)}\right)}{d \sqrt{a-a \csc(c+dx)} \sqrt{a+a \csc(c+dx)}} \\ &= -\frac{2 \cdot 3^{3/4} \sqrt{2+\sqrt{3}} a^2 \cot(c+dx) \left(1-\sqrt[3]{\csc(c+dx)}\right) \sqrt{\frac{1+\sqrt[3]{\csc(c+dx)}+\csc^{\frac{2}{3}}(c+dx)}{(1+\sqrt{3}-\sqrt[3]{\csc(c+dx)})^2}} F\left(\arcsin\left(\frac{1-\sqrt[3]{\csc(c+dx)}}{1+\sqrt{3}-\sqrt[3]{\csc(c+dx)}}\right), -\frac{7-4\sqrt{3}}{3}\right)}{d \sqrt{\frac{1-\sqrt[3]{\csc(c+dx)}}{(1+\sqrt{3}-\sqrt[3]{\csc(c+dx)})^2}} (a-a \csc(c+dx)) \sqrt{a+a \csc(c+dx)}} \end{aligned}$$

Mathematica [C] time = 0.223329, size = 46, normalized size = 0.22

$$\frac{2a \cot(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{3}{2}, 1-\csc(c+dx)\right)}{d \sqrt{a(\csc(c+dx)+1)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[c + d*x]^(1/3)*Sqrt[a + a*Csc[c + d*x]],x]

[Out] $(-2*a*\cot[c + d*x]*\text{Hypergeometric2F1}[1/2, 2/3, 3/2, 1 - \text{Csc}[c + d*x]])/(d*\text{Sqrt}[a*(1 + \text{Csc}[c + d*x])])$

Maple [F] time = 0.645, size = 0, normalized size = 0.

$$\int \sqrt[3]{\csc(dx+c)} \sqrt{a+a \csc(dx+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(d*x+c)^(1/3)*(a+a*csc(d*x+c))^(1/2),x)

[Out] int(csc(d*x+c)^(1/3)*(a+a*csc(d*x+c))^(1/2),x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx+c) + a \csc(dx+c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^(1/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(1/3), x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\sqrt{a \csc(dx+c) + a \csc(dx+c)^{\frac{1}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^(1/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(1/3), x)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a(\csc(c + dx) + 1)} \sqrt[3]{\csc(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)**(1/3)*(a+a*csc(d*x+c))**(1/2),x)

[Out] Integral(sqrt(a*(csc(c + d*x) + 1))*csc(c + d*x)**(1/3), x)

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a \csc(dx + c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^(1/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(1/3), x)

$$3.23 \quad \int \frac{\sqrt{a+a \csc(c+dx)}}{\csc^{\frac{2}{3}}(c+dx)} dx$$

Optimal. Leaf size=254

$$\frac{3^{3/4} \sqrt{2 + \sqrt{3}} a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right), -7 - 4\sqrt{3}\right)}{2d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c + dx)) \sqrt{a \csc(c + dx) + a}}$$

[Out] $(-3*a*\cos[c + d*x]*\csc[c + d*x]^{(1/3)})/(2*d*\sqrt{a + a*\csc[c + d*x]}) - (3^{(3/4)}*\sqrt{2 + \sqrt{3}}*a^2*\cot[c + d*x]*(1 - \csc[c + d*x]^{(1/3)})*\sqrt{((1 + \csc[c + d*x]^{(1/3)} + \csc[c + d*x]^{(2/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})^2}*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \sqrt{3} - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})], -7 - 4*\sqrt{3}])/(2*d*\sqrt{((1 - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})^2}*(a - a*\csc[c + d*x])* \sqrt{a + a*\csc[c + d*x]})$

Rubi [A] time = 0.144931, antiderivative size = 254, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.16$, Rules used = {3806, 51, 63, 218}

$$\frac{3^{3/4} \sqrt{2 + \sqrt{3}} a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} F\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right) \mid -7 - 4\sqrt{3}\right)}{2d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c + dx)) \sqrt{a \csc(c + dx) + a}} - \frac{3a \csc(c + dx)}{2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\sqrt{a + a*\csc[c + d*x]}/\csc[c + d*x]^{(2/3)}, x]$

[Out] $(-3*a*\cos[c + d*x]*\csc[c + d*x]^{(1/3)})/(2*d*\sqrt{a + a*\csc[c + d*x]}) - (3^{(3/4)}*\sqrt{2 + \sqrt{3}}*a^2*\cot[c + d*x]*(1 - \csc[c + d*x]^{(1/3)})*\sqrt{((1 + \csc[c + d*x]^{(1/3)} + \csc[c + d*x]^{(2/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})^2}*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \sqrt{3} - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})], -7 - 4*\sqrt{3}])/(2*d*\sqrt{((1 - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})^2}*(a - a*\csc[c + d*x])* \sqrt{a + a*\csc[c + d*x]})$

Rule 3806

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.)
+ (a_.)], x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]
]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[(d*x)^(n - 1)/Sqrt[a - b*x], x], x,
Csc[e + f*x]], x] /; FreeQ[{a, b, d, e, f, n}, x] && EqQ[a^2 - b^2, 0]
```

Rule 51

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
((a + b*x)^(m + 1)*(c + d*x)^(n + 1))/((b*c - a*d)*(m + 1)), x] - Dist[(d*(
m + n + 2))/((b*c - a*d)*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ
[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]
```

Rule 63

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - (a*d)/b +
(d*x^p)/b)^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 218

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 + Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 - Sqrt[3])*s
+ r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[(s*(s + r*x))/((1 + Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b}, x] &
& PosQ[a]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{a + a \csc(c + dx)}}{\csc^{\frac{2}{3}}(c + dx)} dx &= \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{x^{5/3} \sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\
&= -\frac{3a \cos(c + dx) \sqrt[3]{\csc(c + dx)}}{2d \sqrt{a + a \csc(c + dx)}} + \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{x^{2/3} \sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{4d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\
&= -\frac{3a \cos(c + dx) \sqrt[3]{\csc(c + dx)}}{2d \sqrt{a + a \csc(c + dx)}} + \frac{(3a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{\sqrt{a-ax^3}} dx, x, \sqrt[3]{\csc(c + dx)}\right)}{4d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\
&= -\frac{3a \cos(c + dx) \sqrt[3]{\csc(c + dx)}}{2d \sqrt{a + a \csc(c + dx)}} - \frac{3^{3/4} \sqrt{2 + \sqrt{3}} a^2 \cot(c + dx) (1 - \sqrt[3]{\csc(c + dx)}) \sqrt{\frac{1 + \sqrt[3]{\csc(c + dx)}}{(1 + \sqrt{3} - \sqrt[3]{\csc(c + dx)})}}}{2d \sqrt{\frac{1 - \sqrt[3]{\csc(c + dx)}}{(1 + \sqrt{3} - \sqrt[3]{\csc(c + dx)})^2}} (a - a \csc(c + dx))}
\end{aligned}$$

Mathematica [C] time = 0.44718, size = 110, normalized size = 0.43

$$\frac{\sqrt{a(\csc(c + dx) + 1)} \left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right) \right) \left(\csc^{\frac{2}{3}}(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{3}{2}, 1 - \csc(c + dx)\right) \right)}{2d \csc^{\frac{2}{3}}(c + dx) \left(\sin\left(\frac{1}{2}(c + dx)\right) + \cos\left(\frac{1}{2}(c + dx)\right) \right)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + a*Csc[c + d*x]]/Csc[c + d*x]^(2/3), x]

[Out] -(Sqrt[a*(1 + Csc[c + d*x]])*(3 + Csc[c + d*x]^(2/3)*Hypergeometric2F1[1/2, 2/3, 3/2, 1 - Csc[c + d*x]])*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(2*d*sc[c + d*x]^(2/3)*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))

Maple [F] time = 0.448, size = 0, normalized size = 0.

$$\int \sqrt{a + a \csc(dx + c)} (\csc(dx + c))^{-\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(2/3), x)

[Out] `int((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(2/3),x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(2/3),x, algorithm="maxima")`

[Out] `integrate(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(2/3), x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{2}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(2/3),x, algorithm="fricas")`

[Out] `integral(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(2/3), x)`

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a (\csc(c + dx) + 1)}}{\csc^{\frac{2}{3}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(d*x+c))**(1/2)/csc(d*x+c)**(2/3),x)`

[Out] `Integral(sqrt(a*(csc(c + d*x) + 1))/csc(c + d*x)**(2/3), x)`

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(2/3),x, algorithm="giac")
```

```
[Out] integrate(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(2/3), x)
```

3.24 $\int \csc^{\frac{5}{3}}(c + dx) \sqrt{a + a \csc(c + dx)} dx$

Optimal. Leaf size=514

$$8\sqrt{2}3^{3/4}a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} \text{EllipticF}\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right), -7 - 4\sqrt{3}\right) - 12\sqrt{2}3^{3/4}a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c + dx)) \sqrt{a \csc(c + dx) + a}$$

[Out] (24*a*Cot[c + d*x])/(7*d*(1 + Sqrt[3] - Csc[c + d*x]^(1/3))*Sqrt[a + a*Csc[c + d*x]]) - (6*a*Cos[c + d*x]*Csc[c + d*x]^(5/3))/(7*d*Sqrt[a + a*Csc[c + d*x]]) - (12*3^(1/4)*Sqrt[2 - Sqrt[3]]*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*EllipticE[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(7*d*Sqrt[(1 - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a + a*Csc[c + d*x]]) + (8*Sqrt[2]*3^(3/4)*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*EllipticF[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(7*d*Sqrt[(1 - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a + a*Csc[c + d*x]])

Rubi [A] time = 0.298501, antiderivative size = 514, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 6, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.24$, Rules used = {3806, 50, 63, 303, 218, 1877}

$$8\sqrt{2}3^{3/4}a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} F\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right) \mid -7 - 4\sqrt{3}\right) - 12\sqrt{2}3^{3/4}a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c + dx)) \sqrt{a \csc(c + dx) + a}$$

Antiderivative was successfully verified.

[In] Int[Csc[c + d*x]^(5/3)*Sqrt[a + a*Csc[c + d*x]], x]

[Out] (24*a*Cot[c + d*x])/(7*d*(1 + Sqrt[3] - Csc[c + d*x]^(1/3))*Sqrt[a + a*Csc[c + d*x]]) - (6*a*Cos[c + d*x]*Csc[c + d*x]^(5/3))/(7*d*Sqrt[a + a*Csc[c + d*x]]) - (12*3^(1/4)*Sqrt[2 - Sqrt[3]]*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*EllipticE[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(7*d*Sqrt[(1 - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a + a*Csc[c + d*x]]) + (8*Sqrt[2]*3^(3/4)*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*EllipticF[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(7*d*Sqrt[(1 - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a + a*Csc[c + d*x]])

```

1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc
[c + d*x]^(1/3))^2]*EllipticE[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1
+ Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(7*d*Sqrt[(1 - Csc[c + d
*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[
a + a*Csc[c + d*x]]) + (8*Sqrt[2]*3^(3/4)*a^2*Cot[c + d*x]*(1 - Csc[c + d*x
]^(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] -
Csc[c + d*x]^(1/3))^2]*EllipticF[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/
(1 + Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(7*d*Sqrt[(1 - Csc[c
+ d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqr
t[a + a*Csc[c + d*x]])

```

Rule 3806

```

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.)
+ (a_.)], x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]
]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[(d*x)^(n - 1)/Sqrt[a - b*x], x], x,
Csc[e + f*x]], x] /; FreeQ[{a, b, d, e, f, n}, x] && EqQ[a^2 - b^2, 0]

```

Rule 50

```

Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := Simp[
((a + b*x)^(m + 1)*(c + d*x)^n)/(b*(m + n + 1)), x] + Dist[(n*(b*c - a*d))/
(b*(m + n + 1)), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

```

Rule 63

```

Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - (a*d)/b +
(d*x^p)/b)^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

```

Rule 303

```

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, Dist[(Sqrt[2]*s)/(Sqrt[2 + Sqrt[3]]*r), Int[1/Sqr
t[a + b*x^3], x], x] + Dist[1/r, Int[((1 - Sqrt[3])*s + r*x)/Sqrt[a + b*x^
3], x], x]] /; FreeQ[{a, b}, x] && PosQ[a]

```

Rule 218

```

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]]},

```

```
s = Denom[Rt[b/a, 3]], Simp[(2*Sqrt[2 + Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]]]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(s*(s + r*x))/((1 + Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b}, x] & & PosQ[a]
```

Rule 1877

```
Int[((c_) + (d_)*(x_))/Sqrt[(a_) + (b_)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 - Sqrt[3])*d)/c]], s = Denom[Simplify[((1 - Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 + Sqrt[3])*s + r*x)), x] - Simp[(3^(1/4)*Sqrt[2 - Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[(s*(s + r*x))/((1 + Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b, c, d}, x] && PosQ[a] && EqQ[b*c^3 - 2*(5 - 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
 \int \csc^{\frac{5}{3}}(c + dx) \sqrt{a + a \csc(c + dx)} dx &= \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{x^{2/3}}{\sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{d\sqrt{a - a \csc(c + dx)}\sqrt{a + a \csc(c + dx)}} \\
 &= -\frac{6a \cos(c + dx) \csc^{\frac{5}{3}}(c + dx)}{7d\sqrt{a + a \csc(c + dx)}} + \frac{(4a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{\sqrt[3]{x}\sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{7d\sqrt{a - a \csc(c + dx)}\sqrt{a + a \csc(c + dx)}} \\
 &= -\frac{6a \cos(c + dx) \csc^{\frac{5}{3}}(c + dx)}{7d\sqrt{a + a \csc(c + dx)}} + \frac{(12a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{x}{\sqrt{a-ax^3}} dx, x, \sqrt[3]{\csc(c + dx)}\right)}{7d\sqrt{a - a \csc(c + dx)}\sqrt{a + a \csc(c + dx)}} \\
 &= -\frac{6a \cos(c + dx) \csc^{\frac{5}{3}}(c + dx)}{7d\sqrt{a + a \csc(c + dx)}} - \frac{(12a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1-\sqrt{3}-x}{\sqrt{a-ax^3}} dx, x, \sqrt[3]{\csc(c + dx)}\right)}{7d\sqrt{a - a \csc(c + dx)}\sqrt{a + a \csc(c + dx)}} \\
 &= \frac{24a \cot(c + dx)}{7d(1 + \sqrt{3} - \sqrt[3]{\csc(c + dx)})\sqrt{a + a \csc(c + dx)}} - \frac{6a \cos(c + dx) \csc^{\frac{5}{3}}(c + dx)}{7d\sqrt{a + a \csc(c + dx)}}
 \end{aligned}$$

Mathematica [C] time = 1.18692, size = 120, normalized size = 0.23

$$\frac{2\sqrt{a(\csc(c + dx) + 1)}\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right)\left(3(\csc(c + dx) + 4) - 8\sqrt[3]{\csc(c + dx)}\right)\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{\csc(c + dx) + 1}{\csc(c + dx)}\right) - 7d\sqrt[3]{\csc(c + dx)}\left(\sin\left(\frac{1}{2}(c + dx)\right) + \cos\left(\frac{1}{2}(c + dx)\right)\right)}{7d\sqrt[3]{\csc(c + dx)}\left(\sin\left(\frac{1}{2}(c + dx)\right) + \cos\left(\frac{1}{2}(c + dx)\right)\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[c + d*x]^(5/3)*Sqrt[a + a*Csc[c + d*x]],x]

[Out] (-2*Sqrt[a*(1 + Csc[c + d*x])]*(3*(4 + Csc[c + d*x]) - 8*Csc[c + d*x]^(1/3)) *Hypergeometric2F1[1/2, 4/3, 3/2, 1 - Csc[c + d*x]])*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(7*d*Csc[c + d*x]^(1/3)*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))

Maple [F] time = 0.335, size = 0, normalized size = 0.

$$\int (\csc(dx + c))^{5/3} \sqrt{a + a \csc(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(d*x+c)^(5/3)*(a+a*csc(d*x+c))^(1/2),x)

[Out] int(csc(d*x+c)^(5/3)*(a+a*csc(d*x+c))^(1/2),x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a} \csc(dx + c)^{5/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^(5/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(5/3), x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\sqrt{a \csc(dx + c) + a} \csc(dx + c)^{5/3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.


```
[In] integrate(csc(d*x+c)^(5/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] integral(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(5/3), x)
```

Sympy [F(-1)] time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(d*x+c)**(5/3)*(a+a*csc(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a} \csc(dx + c)^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(d*x+c)^(5/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(5/3), x)
```

3.25 $\int \csc^{\frac{2}{3}}(c + dx) \sqrt{a + a \csc(c + dx)} dx$

Optimal. Leaf size=470

$$2\sqrt{2}3^{3/4}a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} \text{EllipticF}\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right), -7 - 4\sqrt{3}\right) - 3\sqrt[4]{2} \\ d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c + dx)) \sqrt{a \csc(c + dx) + a}$$

[Out] (6*a*Cot[c + d*x])/(d*(1 + Sqrt[3] - Csc[c + d*x]^(1/3))*Sqrt[a + a*Csc[c + d*x]]) - (3*3^(1/4)*Sqrt[2 - Sqrt[3]]*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*EllipticE[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(d*Sqrt[(1 - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a + a*Csc[c + d*x]]) + (2*Sqrt[2]*3^(3/4)*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*EllipticF[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(d*Sqrt[(1 - Csc[c + d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a + a*Csc[c + d*x]])

Rubi [A] time = 0.253267, antiderivative size = 470, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.2$, Rules used = {3806, 63, 303, 218, 1877}

$$2\sqrt{2}3^{3/4}a^2 \cot(c + dx) \left(1 - \sqrt[3]{\csc(c + dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} F\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right) \middle| -7 - 4\sqrt{3}\right) - 3\sqrt[4]{2}\sqrt{2} - \\ d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c + dx)) \sqrt{a \csc(c + dx) + a}$$

Antiderivative was successfully verified.

[In] Int[Csc[c + d*x]^(2/3)*Sqrt[a + a*Csc[c + d*x]],x]

[Out] (6*a*Cot[c + d*x])/(d*(1 + Sqrt[3] - Csc[c + d*x]^(1/3))*Sqrt[a + a*Csc[c + d*x]]) - (3*3^(1/4)*Sqrt[2 - Sqrt[3]]*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*EllipticE[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1

```
+ Sqrt[3] - Csc[c + d*x]^(1/3)], -7 - 4*Sqrt[3]]/(d*Sqrt[(1 - Csc[c + d*x]
]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2*(a - a*Csc[c + d*x])*Sqrt[a
+ a*Csc[c + d*x]]) + (2*Sqrt[2]*3^(3/4)*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^
(1/3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Cs
c[c + d*x]^(1/3))^2]*EllipticF[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1
+ Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]]/(d*Sqrt[(1 - Csc[c + d*
x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2*(a - a*Csc[c + d*x])*Sqrt[a
+ a*Csc[c + d*x]])
```

Rule 3806

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.)
+ (a_.)], x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]
]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[(d*x)^(n - 1)/Sqrt[a - b*x], x], x,
Csc[e + f*x], x] /; FreeQ[{a, b, d, e, f, n}, x] && EqQ[a^2 - b^2, 0]
```

Rule 63

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - (a*d)/b +
(d*x^p)/b)^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 303

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, Dist[(Sqrt[2]*s)/(Sqrt[2 + Sqrt[3]]*r), Int[1/Sq
rt[a + b*x^3], x], x] + Dist[1/r, Int[((1 - Sqrt[3])*s + r*x)/Sqrt[a + b*x^
3], x], x] /; FreeQ[{a, b}, x] && PosQ[a]
```

Rule 218

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 + Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 - Sqrt[3])*s
+ r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]]/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[(s*(s + r*x))/((1 + Sqrt[3])*s + r*x)^2]), x] /; FreeQ[{a, b}, x] &
& PosQ[a]
```

Rule 1877

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 - Sqrt[3])*d)/c]], s = Denom[Simplify[((1 - Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 + Sqrt[3])*s + r*x)), x] - S
```

```
imp[(3^(1/4)*Sqrt[2 - Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/
(1 + Sqrt[3])*s + r*x]^2]*EllipticE[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]]/(r^2*Sqrt[a + b*x^3]*Sqrt[(s*(s + r*x))/
((1 + Sqrt[3])*s + r*x)^2]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[a] && Eq
Q[b*c^3 - 2*(5 - 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned} \int \csc^{\frac{2}{3}}(c + dx) \sqrt{a + a \csc(c + dx)} dx &= \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{\sqrt[3]{x}\sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{d\sqrt{a - a \csc(c + dx)}\sqrt{a + a \csc(c + dx)}} \\ &= \frac{(3a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{x}{\sqrt{a-ax^3}} dx, x, \sqrt[3]{\csc(c + dx)}\right)}{d\sqrt{a - a \csc(c + dx)}\sqrt{a + a \csc(c + dx)}} \\ &= -\frac{(3a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1-\sqrt{3}-x}{\sqrt{a-ax^3}} dx, x, \sqrt[3]{\csc(c + dx)}\right)}{d\sqrt{a - a \csc(c + dx)}\sqrt{a + a \csc(c + dx)}} - \frac{(3\sqrt{2(2-\sqrt{3})}a^2 \cot(c + dx))}{d\sqrt{a - a \csc(c + dx)}} \\ &= \frac{6a \cot(c + dx)}{d(1 + \sqrt{3} - \sqrt[3]{\csc(c + dx)})\sqrt{a + a \csc(c + dx)}} - \frac{3^{\frac{4}{3}}\sqrt{2-\sqrt{3}}a^2 \cot(c + dx)}{d\sqrt{a - a \csc(c + dx)}} \end{aligned}$$

Mathematica [C] time = 0.985294, size = 109, normalized size = 0.23

$$\frac{2\sqrt{a(\csc(c + dx) + 1)}\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right)\left(2\sqrt[3]{\csc(c + dx)}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{3}{2}, 1 - \csc(c + dx)\right)\right)}{d\sqrt[3]{\csc(c + dx)}\left(\sin\left(\frac{1}{2}(c + dx)\right) + \cos\left(\frac{1}{2}(c + dx)\right)\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[c + d*x]^(2/3)*Sqrt[a + a*Csc[c + d*x]], x]

[Out] (2*Sqrt[a*(1 + Csc[c + d*x])]*(-3 + 2*Csc[c + d*x]^(1/3)*Hypergeometric2F1[1/2, 4/3, 3/2, 1 - Csc[c + d*x]]*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(d*Csc[c + d*x]^(1/3)*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))

Maple [F] time = 0.325, size = 0, normalized size = 0.

$$\int (\csc(dx + c))^{\frac{2}{3}} \sqrt{a + a \csc(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(d*x+c)^(2/3)*(a+a*csc(d*x+c))^(1/2), x)

[Out] int(csc(d*x+c)^(2/3)*(a+a*csc(d*x+c))^(1/2), x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a \csc(dx + c)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^(2/3)*(a+a*csc(d*x+c))^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(2/3), x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\sqrt{a \csc(dx + c) + a \csc(dx + c)^{\frac{2}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^(2/3)*(a+a*csc(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(2/3), x)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a (\csc(c + dx) + 1)} \csc^{\frac{2}{3}}(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(d*x+c)**(2/3)*(a+a*csc(d*x+c))**(1/2),x)
```

```
[Out] Integral(sqrt(a*(csc(c + d*x) + 1))*csc(c + d*x)**(2/3), x)
```

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a \csc(dx + c)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(d*x+c)^(2/3)*(a+a*csc(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^(2/3), x)
```

$$3.26 \quad \int \frac{\sqrt{a+a \csc(c+dx)}}{\sqrt[3]{\csc(c+dx)}} dx$$

Optimal. Leaf size=508

$$\frac{\sqrt{2}3^{3/4}a^2 \cot(c+dx) \left(1 - \sqrt[3]{\csc(c+dx)}\right) \sqrt{\frac{\csc^2(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right), -7 - 4\sqrt{3}\right)}{d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c+dx)) \sqrt{a \csc(c+dx) + a}}$$

[Out] $(-3*a*\cot[c + d*x])/(d*(1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})*\operatorname{Sqrt}[a + a*\operatorname{Csc}[c + d*x]]) - (3*a*\cos[c + d*x]*\operatorname{Csc}[c + d*x]^{(2/3)})/(d*\operatorname{Sqrt}[a + a*\operatorname{Csc}[c + d*x]]) + (3*3^{(1/4)}*\operatorname{Sqrt}[2 - \operatorname{Sqrt}[3]]*a^2*\cot[c + d*x]*(1 - \operatorname{Csc}[c + d*x]^{(1/3)}))*\operatorname{Sqrt}[(1 + \operatorname{Csc}[c + d*x]^{(1/3)} + \operatorname{Csc}[c + d*x]^{(2/3)})/(1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1 - \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})/(1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})], -7 - 4*\operatorname{Sqrt}[3]]/(2*d*\operatorname{Sqrt}[(1 - \operatorname{Csc}[c + d*x]^{(1/3)})/(1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})^2]*(a - a*\operatorname{Csc}[c + d*x])* \operatorname{Sqrt}[a + a*\operatorname{Csc}[c + d*x]]) - (\operatorname{Sqrt}[2]*3^{(3/4)}*a^2*\cot[c + d*x]*(1 - \operatorname{Csc}[c + d*x]^{(1/3)}))*\operatorname{Sqrt}[(1 + \operatorname{Csc}[c + d*x]^{(1/3)} + \operatorname{Csc}[c + d*x]^{(2/3)})/(1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})/(1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})], -7 - 4*\operatorname{Sqrt}[3]]/(d*\operatorname{Sqrt}[(1 - \operatorname{Csc}[c + d*x]^{(1/3)})/(1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})^2]*(a - a*\operatorname{Csc}[c + d*x])* \operatorname{Sqrt}[a + a*\operatorname{Csc}[c + d*x]])$

Rubi [A] time = 0.274672, antiderivative size = 508, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 6, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.24$, Rules used = {3806, 51, 63, 303, 218, 1877}

$$\frac{\sqrt{2}3^{3/4}a^2 \cot(c+dx) \left(1 - \sqrt[3]{\csc(c+dx)}\right) \sqrt{\frac{\csc^2(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} F\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right) \middle| -7 - 4\sqrt{3}\right)}{d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c+dx)) \sqrt{a \csc(c+dx) + a}} + 3\sqrt[4]{3}\sqrt{2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Sqrt}[a + a*\operatorname{Csc}[c + d*x]]/\operatorname{Csc}[c + d*x]^{(1/3)}, x]$

[Out] $(-3*a*\cot[c + d*x])/(d*(1 + \operatorname{Sqrt}[3] - \operatorname{Csc}[c + d*x]^{(1/3)})*\operatorname{Sqrt}[a + a*\operatorname{Csc}[c + d*x]]) - (3*a*\cos[c + d*x]*\operatorname{Csc}[c + d*x]^{(2/3)})/(d*\operatorname{Sqrt}[a + a*\operatorname{Csc}[c + d*x]])$

```

]) + (3*3^(1/4)*Sqrt[2 - Sqrt[3]]*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))
*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c +
d*x]^(1/3))^2]*EllipticE[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sqr
t[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(2*d*Sqrt[(1 - Csc[c + d*x]^(
1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a + a
*Csc[c + d*x]]) - (Sqrt[2]*3^(3/4)*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3)
)*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c +
d*x]^(1/3))^2]*EllipticF[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sq
rt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(d*Sqrt[(1 - Csc[c + d*x]^(1
/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a + a
Csc[c + d*x]])

```

Rule 3806

```

Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.)
+ (a_.)], x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]
]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[(d*x)^(n - 1)/Sqrt[a - b*x], x], x,
Csc[e + f*x]], x] /; FreeQ[{a, b, d, e, f, n}, x] && EqQ[a^2 - b^2, 0]

```

Rule 51

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
((a + b*x)^(m + 1)*(c + d*x)^(n + 1))/((b*c - a*d)*(m + 1)), x] - Dist[(d*(
m + n + 2))/((b*c - a*d)*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ
[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]

```

Rule 63

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - (a*d)/b +
(d*x^p)/b)^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

```

Rule 303

```

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, Dist[(Sqrt[2]*s)/(Sqrt[2 + Sqrt[3]]*r), Int[1/Sqr
t[a + b*x^3], x], x] + Dist[1/r, Int[((1 - Sqrt[3])*s + r*x)/Sqrt[a + b*x^
3], x], x]] /; FreeQ[{a, b}, x] && PosQ[a]

```

Rule 218


```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 + Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 - Sqrt[3])*s
+ r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[(s*(s + r*x))/((1 + Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b}, x] &
& PosQ[a]
```

Rule 1877

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 - Sqrt[3])*d)/c]], s = Denom[Simplify[((1 - Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 + Sqrt[3])*s + r*x)), x] - S
imp[(3^(1/4)*Sqrt[2 - Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/(
(1 + Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sq
rt[3])*s + r*x)], -7 - 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[(s*(s + r*x))/
((1 + Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b, c, d}, x] && PosQ[a] && Eq
Q[b*c^3 - 2*(5 - 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{a + a \csc(c + dx)}}{\sqrt[3]{\csc(c + dx)}} dx &= \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{x^{4/3} \sqrt{a - ax}} dx, x, \csc(c + dx)\right)}{d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\
&= -\frac{3a \cos(c + dx) \csc^{2/3}(c + dx)}{d \sqrt{a + a \csc(c + dx)}} - \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{\sqrt[3]{x} \sqrt{a - ax}} dx, x, \csc(c + dx)\right)}{2d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\
&= -\frac{3a \cos(c + dx) \csc^{2/3}(c + dx)}{d \sqrt{a + a \csc(c + dx)}} - \frac{(3a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{x}{\sqrt{a - ax^3}} dx, x, \sqrt[3]{\csc(c + dx)}\right)}{2d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\
&= -\frac{3a \cos(c + dx) \csc^{2/3}(c + dx)}{d \sqrt{a + a \csc(c + dx)}} + \frac{(3a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1 - \sqrt{3} - x}{\sqrt{a - ax^3}} dx, x, \sqrt[3]{\csc(c + dx)}\right)}{2d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} + \dots \\
&= -\frac{3a \cot(c + dx)}{d(1 + \sqrt{3} - \sqrt[3]{\csc(c + dx)}) \sqrt{a + a \csc(c + dx)}} - \frac{3a \cos(c + dx) \csc^{2/3}(c + dx)}{d \sqrt{a + a \csc(c + dx)}} + \frac{3^4 \sqrt{3} \sqrt{2 - \dots}}{\dots}
\end{aligned}$$

Mathematica [C] time = 0.754867, size = 46, normalized size = 0.09

$$\frac{2a \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{3}{2}, 1 - \csc(c + dx)\right)}{d\sqrt{a(\csc(c + dx) + 1)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + a*Csc[c + d*x]]/Csc[c + d*x]^(1/3), x]

[Out] (-2*a*Cot[c + d*x]*Hypergeometric2F1[1/2, 4/3, 3/2, 1 - Csc[c + d*x]])/(d*Sqrt[a*(1 + Csc[c + d*x])])

Maple [F] time = 0.44, size = 0, normalized size = 0.

$$\int \frac{\sqrt{a + a \csc(dx + c)}}{\sqrt[3]{\csc(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(1/3), x)

[Out] int((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(1/3), x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(1/3), x, algorithm="maxima")

[Out] integrate(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(1/3), x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{1}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(1/3),x, algorithm="fricas")

[Out] integral(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(1/3), x)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a(\csc(c + dx) + 1)}}{\sqrt[3]{\csc(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(d*x+c))**(1/2)/csc(d*x+c)**(1/3),x)

[Out] Integral(sqrt(a*(csc(c + d*x) + 1))/csc(c + d*x)**(1/3), x)

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(1/3),x, algorithm="giac")

[Out] integrate(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(1/3), x)

$$3.27 \quad \int \frac{\sqrt{a+a \csc(c+dx)}}{\csc^{\frac{4}{3}}(c+dx)} dx$$

Optimal. Leaf size=552

$$\frac{5 \cdot 3^{3/4} a^2 \cot(c+dx) \left(1 - \sqrt[3]{\csc(c+dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right), -7 - 4\sqrt{3}\right) + 15\sqrt[4]{3} \sqrt{2} d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c+dx)) \sqrt{a \csc(c+dx) + a}}{4\sqrt{2}d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c+dx)) \sqrt{a \csc(c+dx) + a}}$$

[Out] $(-15*a*\cot[c + d*x])/(8*d*(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})*\sqrt{a + a*\csc[c + d*x]}) - (3*a*\cos[c + d*x])/(4*d*\csc[c + d*x]^{(1/3)}*\sqrt{a + a*\csc[c + d*x]}) - (15*a*\cos[c + d*x]*\csc[c + d*x]^{(2/3)})/(8*d*\sqrt{a + a*\csc[c + d*x]}) + (15*3^{(1/4)}*\sqrt{2 - \sqrt{3}}*a^2*\cot[c + d*x]*(1 - \csc[c + d*x]^{(1/3)})*\sqrt{(1 + \csc[c + d*x]^{(1/3)} + \csc[c + d*x]^{(2/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3}))^2}*\operatorname{EllipticE}[\operatorname{ArcSin}[(1 - \sqrt{3} - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})], -7 - 4*\sqrt{3}])/(16*d*\sqrt{(1 - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3}))^2}*(a - a*\csc[c + d*x])*\sqrt{a + a*\csc[c + d*x]}) - (5*3^{(3/4)}*a^2*\cot[c + d*x]*(1 - \csc[c + d*x]^{(1/3)})*\sqrt{(1 + \csc[c + d*x]^{(1/3)} + \csc[c + d*x]^{(2/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3}))^2}*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \sqrt{3} - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})], -7 - 4*\sqrt{3}])/(4*\sqrt{2}*d*\sqrt{(1 - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3}))^2}*(a - a*\csc[c + d*x])*\sqrt{a + a*\csc[c + d*x]})$

Rubi [A] time = 0.306694, antiderivative size = 552, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 6, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.24$, Rules used = {3806, 51, 63, 303, 218, 1877}

$$\frac{5 \cdot 3^{3/4} a^2 \cot(c+dx) \left(1 - \sqrt[3]{\csc(c+dx)}\right) \sqrt{\frac{\csc^{\frac{2}{3}}(c+dx) + \sqrt[3]{\csc(c+dx)} + 1}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} F\left(\sin^{-1}\left(\frac{-\sqrt[3]{\csc(c+dx)} - \sqrt{3} + 1}{-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1}\right) \middle| -7 - 4\sqrt{3}\right) + 15\sqrt[4]{3} \sqrt{2} d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c+dx)) \sqrt{a \csc(c+dx) + a}}{4\sqrt{2}d \sqrt{\frac{1 - \sqrt[3]{\csc(c+dx)}}{(-\sqrt[3]{\csc(c+dx)} + \sqrt{3} + 1)^2}} (a - a \csc(c+dx)) \sqrt{a \csc(c+dx) + a}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\sqrt{a + a*\csc[c + d*x]}/\csc[c + d*x]^{(4/3)}, x]$

[Out] $(-15*a*\cot[c + d*x])/(8*d*(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})*\sqrt{a + a*\csc[c + d*x]}) - (3*a*\cos[c + d*x])/(4*d*\csc[c + d*x]^{(1/3)}*\sqrt{a + a*\csc[c + d*x]}) - (15*a*\cos[c + d*x]*\csc[c + d*x]^{(2/3)})/(8*d*\sqrt{a + a*\csc[c + d*x]}) + (15*3^{(1/4)}*\sqrt{2 - \sqrt{3}}*a^2*\cot[c + d*x]*(1 - \csc[c + d*x]^{(1/3)})*\sqrt{(1 + \csc[c + d*x]^{(1/3)} + \csc[c + d*x]^{(2/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3}))^2}*\operatorname{EllipticE}[\operatorname{ArcSin}[(1 - \sqrt{3} - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})], -7 - 4*\sqrt{3}])/(16*d*\sqrt{(1 - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3}))^2}*(a - a*\csc[c + d*x])*\sqrt{a + a*\csc[c + d*x]}) - (5*3^{(3/4)}*a^2*\cot[c + d*x]*(1 - \csc[c + d*x]^{(1/3)})*\sqrt{(1 + \csc[c + d*x]^{(1/3)} + \csc[c + d*x]^{(2/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3}))^2}*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 - \sqrt{3} - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3)})], -7 - 4*\sqrt{3}])/(4*\sqrt{2}*d*\sqrt{(1 - \csc[c + d*x]^{(1/3)})/(1 + \sqrt{3} - \csc[c + d*x]^{(1/3}))^2}*(a - a*\csc[c + d*x])*\sqrt{a + a*\csc[c + d*x]})$

```

d*x]]) - (15*a*cos[c + d*x]*Csc[c + d*x]^(2/3))/(8*d*Sqrt[a + a*Csc[c + d*
x]]) + (15*3^(1/4)*Sqrt[2 - Sqrt[3]]*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/
3))*Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c
+ d*x]^(1/3))^2]*EllipticE[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 +
Sqrt[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(16*d*Sqrt[(1 - Csc[c + d*
x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqrt[a
+ a*Csc[c + d*x]]) - (5*3^(3/4)*a^2*Cot[c + d*x]*(1 - Csc[c + d*x]^(1/3))*
Sqrt[(1 + Csc[c + d*x]^(1/3) + Csc[c + d*x]^(2/3))/(1 + Sqrt[3] - Csc[c + d
*x]^(1/3))^2]*EllipticF[ArcSin[(1 - Sqrt[3] - Csc[c + d*x]^(1/3))/(1 + Sqrt
[3] - Csc[c + d*x]^(1/3))], -7 - 4*Sqrt[3]])/(4*Sqrt[2]*d*Sqrt[(1 - Csc[c +
d*x]^(1/3))/(1 + Sqrt[3] - Csc[c + d*x]^(1/3))^2]*(a - a*Csc[c + d*x])*Sqr
t[a + a*Csc[c + d*x]])

```

Rule 3806

```

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.)
+ (a_.)], x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]
]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[(d*x)^(n - 1)/Sqrt[a - b*x], x], x,
Csc[e + f*x], x] /; FreeQ[{a, b, d, e, f, n}, x] && EqQ[a^2 - b^2, 0]

```

Rule 51

```

Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := Simp[
((a + b*x)^(m + 1)*(c + d*x)^(n + 1))/((b*c - a*d)*(m + 1)), x] - Dist[(d*(
m + n + 2))/((b*c - a*d)*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ
[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]

```

Rule 63

```

Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - (a*d)/b +
(d*x^p)/b)^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

```

Rule 303

```

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, Dist[(Sqrt[2]*s)/(Sqrt[2 + Sqrt[3]]*r), Int[1/Sq
rt[a + b*x^3], x], x] + Dist[1/r, Int[((1 - Sqrt[3])*s + r*x)/Sqrt[a + b*x^
3], x], x] /; FreeQ[{a, b}, x] && PosQ[a]

```

Rule 218

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 + Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 + Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 - Sqrt[3])*s
+ r*x)/((1 + Sqrt[3])*s + r*x)], -7 - 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[(s*(s + r*x))/((1 + Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b}, x] &
& PosQ[a]
```

Rule 1877

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 - Sqrt[3])*d)/c]], s = Denom[Simplify[((1 - Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 + Sqrt[3])*s + r*x)), x] - S
imp[(3^(1/4)*Sqrt[2 - Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/(
(1 + Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 - Sqrt[3])*s + r*x)/((1 + Sq
rt[3])*s + r*x)], -7 - 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[(s*(s + r*x))/
((1 + Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b, c, d}, x] && PosQ[a] && Eq
Q[b*c^3 - 2*(5 - 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\int \frac{\sqrt{a + a \csc(c + dx)}}{\csc^{\frac{4}{3}}(c + dx)} dx = \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{x^{7/3} \sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}}$$

$$= -\frac{3a \cos(c + dx)}{4d \sqrt[3]{\csc(c + dx)} \sqrt{a + a \csc(c + dx)}} + \frac{(5a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{x^{4/3} \sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{8d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}}$$

$$= -\frac{3a \cos(c + dx)}{4d \sqrt[3]{\csc(c + dx)} \sqrt{a + a \csc(c + dx)}} - \frac{15a \cos(c + dx) \csc^{\frac{2}{3}}(c + dx)}{8d \sqrt{a + a \csc(c + dx)}} - \frac{(5a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{x^{4/3} \sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{16d \sqrt{a - a \csc(c + dx)}}$$

$$= -\frac{3a \cos(c + dx)}{4d \sqrt[3]{\csc(c + dx)} \sqrt{a + a \csc(c + dx)}} - \frac{15a \cos(c + dx) \csc^{\frac{2}{3}}(c + dx)}{8d \sqrt{a + a \csc(c + dx)}} - \frac{(15a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{x^{4/3} \sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{16d \sqrt{a - a \csc(c + dx)}}$$

$$= -\frac{3a \cos(c + dx)}{4d \sqrt[3]{\csc(c + dx)} \sqrt{a + a \csc(c + dx)}} - \frac{15a \cos(c + dx) \csc^{\frac{2}{3}}(c + dx)}{8d \sqrt{a + a \csc(c + dx)}} + \frac{(15a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{1}{x^{4/3} \sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{16d \sqrt{a - a \csc(c + dx)}}$$

$$= -\frac{15a \cot(c + dx)}{8d (1 + \sqrt{3} - \sqrt[3]{\csc(c + dx)}) \sqrt{a + a \csc(c + dx)}} - \frac{3a \cos(c + dx)}{4d \sqrt[3]{\csc(c + dx)} \sqrt{a + a \csc(c + dx)}} - \frac{15a^2 \cot(c + dx)}{16d \sqrt{a - a \csc(c + dx)}}$$

Mathematica [C] time = 1.20123, size = 72, normalized size = 0.13

$$\frac{a \cos(c + dx) \left(5 \csc^{\frac{4}{3}}(c + dx) \text{Hypergeometric2F1} \left(\frac{1}{2}, \frac{4}{3}, \frac{3}{2}, 1 - \csc(c + dx) \right) + 3 \right)}{4d \sqrt[3]{\csc(c + dx)} \sqrt{a(\csc(c + dx) + 1)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + a*Csc[c + d*x]]/Csc[c + d*x]^(4/3), x]

[Out] -(a*Cos[c + d*x]*(3 + 5*Csc[c + d*x]^(4/3)*Hypergeometric2F1[1/2, 4/3, 3/2, 1 - Csc[c + d*x]]))/(4*d*Csc[c + d*x]^(1/3)*Sqrt[a*(1 + Csc[c + d*x])])

Maple [F] time = 0.323, size = 0, normalized size = 0.

$$\int \sqrt{a + a \csc(dx + c)} (\csc(dx + c))^{-\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(4/3), x)

[Out] int((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(4/3), x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(4/3), x, algorithm="maxima")

[Out] integrate(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(4/3), x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral} \left(\frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{4}{3}}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(4/3),x, algorithm="fricas")

[Out] integral(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(4/3), x)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a(\csc(c + dx) + 1)}}{\csc^{\frac{4}{3}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(d*x+c))**(1/2)/csc(d*x+c)**(4/3),x)

[Out] Integral(sqrt(a*(csc(c + d*x) + 1))/csc(c + d*x)**(4/3), x)

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sqrt{a \csc(dx + c) + a}}{\csc(dx + c)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*csc(d*x+c))^(1/2)/csc(d*x+c)^(4/3),x, algorithm="giac")

[Out] integrate(sqrt(a*csc(d*x + c) + a)/csc(d*x + c)^(4/3), x)

3.28 $\int \csc^n(c + dx) \sqrt{a + a \csc(c + dx)} dx$

Optimal. Leaf size=48

$$\frac{2a \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, 1 - \csc(c + dx)\right)}{d \sqrt{a \csc(c + dx) + a}}$$

[Out] $(-2*a*\cot[c + d*x]*\operatorname{Hypergeometric2F1}[1/2, 1 - n, 3/2, 1 - \operatorname{Csc}[c + d*x]])/(d*\sqrt{a + a*\operatorname{Csc}[c + d*x]})$

Rubi [A] time = 0.0584631, antiderivative size = 48, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {3806, 65}

$$\frac{2a \cot(c + dx) {}_2F_1\left(\frac{1}{2}, 1 - n; \frac{3}{2}; 1 - \csc(c + dx)\right)}{d \sqrt{a \csc(c + dx) + a}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[c + d*x]^n*\sqrt{a + a*\operatorname{Csc}[c + d*x]}, x]$

[Out] $(-2*a*\cot[c + d*x]*\operatorname{Hypergeometric2F1}[1/2, 1 - n, 3/2, 1 - \operatorname{Csc}[c + d*x]])/(d*\sqrt{a + a*\operatorname{Csc}[c + d*x]})$

Rule 3806

$\operatorname{Int}[(\operatorname{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*\sqrt{\operatorname{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)}, x_Symbol] \rightarrow \operatorname{Dist}[(a^2*d*\cot[e + f*x])/(f*\sqrt{a + b*\operatorname{Csc}[e + f*x]})*\sqrt{a - b*\operatorname{Csc}[e + f*x]}], \operatorname{Subst}[\operatorname{Int}[(d*x)^{(n - 1)}/\sqrt{a - b*x}], x], x, \operatorname{Csc}[e + f*x], x] /;$ $\operatorname{FreeQ}\{a, b, d, e, f, n\}, x \ \&\& \ \operatorname{EqQ}[a^2 - b^2, 0]$

Rule 65

$\operatorname{Int}[(b_.*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(c + d*x)^{(n + 1)}*\operatorname{Hypergeometric2F1}[-m, n + 1, n + 2, 1 + (d*x)/c]/(d*(n + 1)*(-(d/(b*c)))^m), x] /;$ $\operatorname{FreeQ}\{b, c, d, m, n\}, x \ \&\& \ !\operatorname{IntegerQ}[n] \ \&\& \ (\operatorname{IntegerQ}[m] \ || \ \operatorname{GtQ}[-(d/(b*c)), 0])$

Rubi steps

$$\int \csc^n(c + dx) \sqrt{a + a \csc(c + dx)} dx = \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{x^{-1+n}}{\sqrt{a-ax}} dx, x, \csc(c + dx)\right)}{d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\ = -\frac{2a \cot(c + dx) {}_2F_1\left(\frac{1}{2}, 1 - n; \frac{3}{2}; 1 - \csc(c + dx)\right)}{d \sqrt{a + a \csc(c + dx)}}$$

Mathematica [A] time = 0.169233, size = 48, normalized size = 1.

$$\frac{2a \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, 1 - \csc(c + dx)\right)}{d \sqrt{a(\csc(c + dx) + 1)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[c + d*x]^n*Sqrt[a + a*Csc[c + d*x]], x]

[Out] (-2*a*Cot[c + d*x]*Hypergeometric2F1[1/2, 1 - n, 3/2, 1 - Csc[c + d*x]])/(d*Sqrt[a*(1 + Csc[c + d*x])])

Maple [F] time = 0.401, size = 0, normalized size = 0.

$$\int (\csc(dx + c))^n \sqrt{a + a \csc(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(d*x+c)^n*(a+a*csc(d*x+c))^(1/2), x)

[Out] int(csc(d*x+c)^n*(a+a*csc(d*x+c))^(1/2), x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a} \csc(dx + c)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^n*(a+a*csc(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^n, x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\sqrt{a \csc(dx + c) + a} \csc(dx + c)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^n*(a+a*csc(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^n, x)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a(\csc(c + dx) + 1)} \csc^n(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)**n*(a+a*csc(d*x+c))**(1/2),x)

[Out] Integral(sqrt(a*(csc(c + d*x) + 1))*csc(c + d*x)**n, x)

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{a \csc(dx + c) + a} \csc(dx + c)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(d*x+c)^n*(a+a*csc(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(a*csc(d*x + c) + a)*csc(d*x + c)^n, x)

3.29 $\int \csc^n(c + dx) \sqrt{a - a \csc(c + dx)} dx$

Optimal. Leaf size=69

$$\frac{2a \cos(c + dx)(-\csc(c + dx))^{-n} \csc^{n+1}(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, \csc(c + dx) + 1\right)}{d\sqrt{a - a \csc(c + dx)}}$$

[Out] $(-2*a*\cos[c + d*x]*\operatorname{Csc}[c + d*x]^{(1 + n)}*\operatorname{Hypergeometric2F1}[1/2, 1 - n, 3/2, 1 + \operatorname{Csc}[c + d*x]])/(d*(-\operatorname{Csc}[c + d*x])^n*\operatorname{Sqrt}[a - a*\operatorname{Csc}[c + d*x]])$

Rubi [A] time = 0.0701908, antiderivative size = 69, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3806, 67, 65}

$$\frac{2a \cos(c + dx)(-\csc(c + dx))^{-n} \csc^{n+1}(c + dx) {}_2F_1\left(\frac{1}{2}, 1 - n; \frac{3}{2}; \csc(c + dx) + 1\right)}{d\sqrt{a - a \csc(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[c + d*x]^n*\operatorname{Sqrt}[a - a*\operatorname{Csc}[c + d*x]], x]$

[Out] $(-2*a*\cos[c + d*x]*\operatorname{Csc}[c + d*x]^{(1 + n)}*\operatorname{Hypergeometric2F1}[1/2, 1 - n, 3/2, 1 + \operatorname{Csc}[c + d*x]])/(d*(-\operatorname{Csc}[c + d*x])^n*\operatorname{Sqrt}[a - a*\operatorname{Csc}[c + d*x]])$

Rule 3806

$\operatorname{Int}[(\operatorname{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*\operatorname{Sqrt}[\operatorname{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] \rightarrow \operatorname{Dist}[(a^2*d*\operatorname{Cot}[e + f*x])/(f*\operatorname{Sqrt}[a + b*\operatorname{Csc}[e + f*x]]*\operatorname{Sqrt}[a - b*\operatorname{Csc}[e + f*x]]), \operatorname{Subst}[\operatorname{Int}[(d*x)^{(n-1)}/\operatorname{Sqrt}[a - b*x], x], x, \operatorname{Csc}[e + f*x]], x] /;$ $\operatorname{FreeQ}\{a, b, d, e, f, n\}, x \ \&\& \ \operatorname{EqQ}[a^2 - b^2, 0]$

Rule 67

$\operatorname{Int}[(b_.)*(x_.)^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Dist}[(b*c/d)^m*\operatorname{IntPart}[m]*(b*x)^{\operatorname{FracPart}[m]}/(-(d*x)/c)^{\operatorname{FracPart}[m]}, \operatorname{Int}[(d*x)/c]^m*(c + d*x)^n, x] /;$ $\operatorname{FreeQ}\{b, c, d, m, n\}, x \ \&\& \ !\operatorname{IntegerQ}[m] \ \&\& \ !\operatorname{IntegerQ}[n] \ \&\& \ !\operatorname{GtQ}[c, 0] \ \&\& \ !\operatorname{GtQ}[-(d/(b*c)), 0]$

Rule 65

```
Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + (d*x)/c])/(d*(n + 1)*(-(d/(b*c)))^m), x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-(d/(b*c)), 0])
```

Rubi steps

$$\begin{aligned} \int \csc^n(c + dx) \sqrt{a - a \csc(c + dx)} dx &= \frac{(a^2 \cot(c + dx)) \operatorname{Subst}\left(\int \frac{x^{-1+n}}{\sqrt{a+ax}} dx, x, \csc(c + dx)\right)}{d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\ &= -\frac{(a^2 \cos(c + dx) (-\csc(c + dx))^{-n} \csc^{1+n}(c + dx)) \operatorname{Subst}\left(\int \frac{(-x)^{-1+n}}{\sqrt{a+ax}} dx, x, \csc(c + dx)\right)}{d \sqrt{a - a \csc(c + dx)} \sqrt{a + a \csc(c + dx)}} \\ &= -\frac{2a \cos(c + dx) (-\csc(c + dx))^{-n} \csc^{1+n}(c + dx) {}_2F_1\left(\frac{1}{2}, 1 - n; \frac{3}{2}; 1 + \csc(c + dx)\right)}{d \sqrt{a - a \csc(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.913166, size = 73, normalized size = 1.06

$$\frac{2a \cos(c + dx) \csc^{2n+1}(c + dx) (-\csc^2(c + dx))^{-n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, \csc(c + dx) + 1\right)}{d \sqrt{a - a \csc(c + dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[c + d*x]^n*Sqrt[a - a*Csc[c + d*x]], x]
```

```
[Out] (-2*a*Cos[c + d*x]*Csc[c + d*x]^(1 + 2*n)*Hypergeometric2F1[1/2, 1 - n, 3/2, 1 + Csc[c + d*x]])/(d*(-Csc[c + d*x]^2)^n*Sqrt[a - a*Csc[c + d*x]])
```

Maple [F] time = 0.484, size = 0, normalized size = 0.

$$\int (\csc(dx + c))^n \sqrt{a - a \csc(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(d*x+c)^n*(a-a*csc(d*x+c))^(1/2), x)
```

[Out] `int(csc(d*x+c)^n*(a-a*csc(d*x+c))^(1/2),x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{-a \csc(dx + c) + a \csc(dx + c)^n} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(d*x+c)^n*(a-a*csc(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-a*csc(d*x + c) + a)*csc(d*x + c)^n, x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\sqrt{-a \csc(dx + c) + a \csc(dx + c)^n}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(d*x+c)^n*(a-a*csc(d*x+c))^(1/2),x, algorithm="fricas")`

[Out] `integral(sqrt(-a*csc(d*x + c) + a)*csc(d*x + c)^n, x)`

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{-a (\csc(c + dx) - 1) \csc^n(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(d*x+c)**n*(a-a*csc(d*x+c))**(1/2),x)`

[Out] `Integral(sqrt(-a*(csc(c + d*x) - 1))*csc(c + d*x)**n, x)`

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int \sqrt{-a \csc(dx + c) + a \csc(dx + c)^n} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(d*x+c)^n*(a-a*csc(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(-a*csc(d*x + c) + a)*csc(d*x + c)^n, x)
```

3.30 $\int \csc^3(e + fx)(a + a \csc(e + fx))^m dx$

Optimal. Leaf size=156

$$\frac{2^{m+\frac{1}{2}}(m^2 + m + 1) \cot(e + fx)(\csc(e + fx) + 1)^{-m-\frac{1}{2}}(a \csc(e + fx) + a)^m \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - m, \frac{3}{2}, \frac{1}{2}(1 - \csc(e + fx))\right)}{f(m+1)(m+2)}$$

[Out] (Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(2 + 3*m + m^2)) - (Cot[e + f*x]*(a + a*Csc[e + f*x])^(1 + m))/(a*f*(2 + m)) - (2^(1/2 + m)*(1 + m + m^2)*Cot[e + f*x]*(1 + Csc[e + f*x])^(-1/2 - m)*(a + a*Csc[e + f*x])^m*Hypergeometric2F1[1/2, 1/2 - m, 3/2, (1 - Csc[e + f*x])/2])/(f*(1 + m)*(2 + m))

Rubi [A] time = 0.189615, antiderivative size = 156, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3800, 4001, 3828, 3827, 69}

$$\frac{2^{m+\frac{1}{2}}(m^2 + m + 1) \cot(e + fx)(\csc(e + fx) + 1)^{-m-\frac{1}{2}}(a \csc(e + fx) + a)^m {}_2F_1\left(\frac{1}{2}, \frac{1}{2} - m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx))\right)}{f(m+1)(m+2)} + \frac{\cot(e + fx)}{f}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^3*(a + a*Csc[e + f*x])^m,x]

[Out] (Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(2 + 3*m + m^2)) - (Cot[e + f*x]*(a + a*Csc[e + f*x])^(1 + m))/(a*f*(2 + m)) - (2^(1/2 + m)*(1 + m + m^2)*Cot[e + f*x]*(1 + Csc[e + f*x])^(-1/2 - m)*(a + a*Csc[e + f*x])^m*Hypergeometric2F1[1/2, 1/2 - m, 3/2, (1 - Csc[e + f*x])/2])/(f*(1 + m)*(2 + m))

Rule 3800

Int[csc[(e_.) + (f_.)*(x_)]^3*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := -Simp[(Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1))/(b*f*(m + 2)), x] + Dist[1/(b*(m + 2)), Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(b*(m + 1) - a*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f, m}, x] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]

Rule 4001

Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := -Simp[(B*Cot[e + f*x]*(a


```
+ b*Csc[e + f*x]^m)/(f*(m + 1)), x] + Dist[(a*B*m + A*b*(m + 1))/(b*(m + 1)), Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m, x], x] /; FreeQ[{a, b, A, B, e, f, m}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && NeQ[a*B*m + A*b*(m + 1), 0] && !LtQ[m, -2^(-1)]
```

Rule 3828

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m, x_Symbol] := Dist[(a^IntPart[m]*(a + b*Csc[e + f*x])^FracPart[m])/((1 + (b*Csc[e + f*x])/a)^FracPart[m], Int[(1 + (b*Csc[e + f*x])/a]^m*(d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && !GtQ[a, 0]
```

Rule 3827

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m, x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[((d*x)^(n - 1)*(a + b*x)^(m - 1/2))/Sqrt[a - b*x], x], x, Csc[e + f*x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && GtQ[a, 0]
```

Rule 69

```
Int[((a_) + (b_.)*(x_))^m*((c_) + (d_.)*(x_))^n, x_Symbol] := Simp[((a + b*x)^(m + 1)*Hypergeometric2F1[-n, m + 1, m + 2, -((d*(a + b*x))/(b*c - a*d))]/(b*(m + 1)*(b/(b*c - a*d))^n), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-(d/(b*c - a*d)), 0]))
```

Rubi steps

$$\begin{aligned}
\int \csc^3(e+fx)(a+a\csc(e+fx))^m dx &= -\frac{\cot(e+fx)(a+a\csc(e+fx))^{1+m}}{af(2+m)} + \frac{\int \csc(e+fx)(a(1+m)-a\csc(e+fx))}{a(2+m)} \\
&= \frac{\cot(e+fx)(a+a\csc(e+fx))^m}{f(2+3m+m^2)} - \frac{\cot(e+fx)(a+a\csc(e+fx))^{1+m}}{af(2+m)} + \frac{(1+m)}{f(2+3m+m^2)} \\
&= \frac{\cot(e+fx)(a+a\csc(e+fx))^m}{f(2+3m+m^2)} - \frac{\cot(e+fx)(a+a\csc(e+fx))^{1+m}}{af(2+m)} + \frac{((1+m))}{f(2+3m+m^2)} \\
&= \frac{\cot(e+fx)(a+a\csc(e+fx))^m}{f(2+3m+m^2)} - \frac{\cot(e+fx)(a+a\csc(e+fx))^{1+m}}{af(2+m)} + \frac{((1+m))}{f(2+3m+m^2)} \\
&= \frac{\cot(e+fx)(a+a\csc(e+fx))^m}{f(2+3m+m^2)} - \frac{\cot(e+fx)(a+a\csc(e+fx))^{1+m}}{af(2+m)} - \frac{2^{\frac{1}{2}+m}}{f(2+3m+m^2)}
\end{aligned}$$

Mathematica [A] time = 0.725878, size = 199, normalized size = 1.28

$$\frac{\cot^2\left(\frac{1}{2}(e+fx)\right)\left(\tan\left(\frac{1}{2}(e+fx)\right)+1\right)^{-2m}(a(\csc(e+fx)+1))^m\left(-4\text{Hypergeometric2F1}\left(-2m-3,-m-2,-m-1,\right.\right.$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^3*(a + a*Csc[e + f*x])^m,x]

[Out] -(Cot[(e + f*x)/2]^2*(a*(1 + Csc[e + f*x]))^m*(-4*Hypergeometric2F1[-3 - 2*m, -2 - m, -1 - m, -Tan[(e + f*x)/2]] - 8*Hypergeometric2F1[-1 - 2*m, -2 - m, -1 - m, -Tan[(e + f*x)/2]] + 4*Hypergeometric2F1[-2 - m, -2*m, -1 - m, -Tan[(e + f*x)/2]] + 8*Hypergeometric2F1[-2 - m, -2*(1 + m), -1 - m, -Tan[(e + f*x)/2]] + Hypergeometric2F1[-2 - m, -2*(2 + m), -1 - m, -Tan[(e + f*x)/2]]))/(4*f*(2 + m)*(1 + Tan[(e + f*x)/2])^(2*m))

Maple [F] time = 0.609, size = 0, normalized size = 0.

$$\int (\csc(fx + e))^3 (a + a \csc(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^3*(a+a*csc(f*x+e))^m,x)`

[Out] `int(csc(f*x+e)^3*(a+a*csc(f*x+e))^m,x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \csc(fx + e)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^3*(a+a*csc(f*x+e))^m,x, algorithm="maxima")`

[Out] `integrate((a*csc(f*x + e) + a)^m*csc(f*x + e)^3, x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(a \csc(fx + e) + a\right)^m \csc(fx + e)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^3*(a+a*csc(f*x+e))^m,x, algorithm="fricas")`

[Out] `integral((a*csc(f*x + e) + a)^m*csc(f*x + e)^3, x)`

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a (\csc(e + fx) + 1))^m \csc^3(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**3*(a+a*csc(f*x+e))**m,x)`

[Out] `Integral((a*(csc(e + f*x) + 1))**m*csc(e + f*x)**3, x)`

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \csc(fx + e)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^3*(a+a*csc(f*x+e))^m,x, algorithm="giac")
```

```
[Out] integrate((a*csc(f*x + e) + a)^m*csc(f*x + e)^3, x)
```

3.31 $\int \csc^2(e + fx)(a + a \csc(e + fx))^m dx$

Optimal. Leaf size=109

$$\frac{2^{m+\frac{1}{2}} m \cot(e + fx) (\csc(e + fx) + 1)^{-m-\frac{1}{2}} (a \csc(e + fx) + a)^m \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - m, \frac{3}{2}, \frac{1}{2}(1 - \csc(e + fx))\right)}{f(m+1)}$$

[Out] -((Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(1 + m))) - (2^(1/2 + m)*m*Cot[e + f*x]*(1 + Csc[e + f*x])^(-1/2 - m)*(a + a*Csc[e + f*x])^m*Hypergeometric2F1[1/2, 1/2 - m, 3/2, (1 - Csc[e + f*x])/2])/(f*(1 + m))

Rubi [A] time = 0.103176, antiderivative size = 109, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.19$, Rules used = {3798, 3828, 3827, 69}

$$\frac{2^{m+\frac{1}{2}} m \cot(e + fx) (\csc(e + fx) + 1)^{-m-\frac{1}{2}} (a \csc(e + fx) + a)^m {}_2F_1\left(\frac{1}{2}, \frac{1}{2} - m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx))\right)}{f(m+1)} - \frac{\cot(e + fx)(a \csc(e + fx) + a)^m}{f(m)}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^2*(a + a*Csc[e + f*x])^m,x]

[Out] -((Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(1 + m))) - (2^(1/2 + m)*m*Cot[e + f*x]*(1 + Csc[e + f*x])^(-1/2 - m)*(a + a*Csc[e + f*x])^m*Hypergeometric2F1[1/2, 1/2 - m, 3/2, (1 - Csc[e + f*x])/2])/(f*(1 + m))

Rule 3798

Int[csc[(e_.) + (f_.)*(x_)]^2*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m), x_Symbol] :> -Simp[(Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(f*(m + 1)), x] + Dist[(a*m)/(b*(m + 1)), Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m, x], x] /; FreeQ[{a, b, e, f, m}, x] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]

Rule 3828

Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m), x_Symbol] :> Dist[(a^IntPart[m]*(a + b*Csc[e + f*x])^FracPart[m])/(1 + (b*Csc[e + f*x])/a)^FracPart[m], Int[(1 + (b*Csc[e + f*x])/a)^m*(d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && !GtQ[a, 0]

Rule 3827

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) +
(a_.)^(m_.), x_Symbol] :> Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*
x])*Sqrt[a - b*Csc[e + f*x]]], Subst[Int[((d*x)^(n - 1)*(a + b*x)^(m - 1/2)
)/Sqrt[a - b*x], x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, d, e, f, m, n}, x
] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && GtQ[a, 0]
```

Rule 69

```
Int[((a_) + (b_.)*(x_.))^(m_)*((c_) + (d_.)*(x_.))^(n_), x_Symbol] :> Simp[((
a + b*x)^(m + 1)*Hypergeometric2F1[-n, m + 1, m + 2, -((d*(a + b*x))/(b*c -
a*d))]/(b*(m + 1)*(b/(b*c - a*d))^(n)), x] /; FreeQ[{a, b, c, d, m, n}, x]
&& NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d)
, 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-(d/(b*c - a*d)), 0]))
```

Rubi steps

$$\begin{aligned} \int \csc^2(e + fx)(a + a \csc(e + fx))^m dx &= -\frac{\cot(e + fx)(a + a \csc(e + fx))^m}{f(1 + m)} + \frac{m \int \csc(e + fx)(a + a \csc(e + fx))^m dx}{1 + m} \\ &= -\frac{\cot(e + fx)(a + a \csc(e + fx))^m}{f(1 + m)} + \frac{(m(1 + \csc(e + fx))^{-m}(a + a \csc(e + fx))^m)}{1 + m} \\ &= -\frac{\cot(e + fx)(a + a \csc(e + fx))^m}{f(1 + m)} + \frac{(m \cot(e + fx)(1 + \csc(e + fx))^{-\frac{1}{2}-m}(a + a \csc(e + fx))^m)}{f(1 + m)} \\ &= -\frac{\cot(e + fx)(a + a \csc(e + fx))^m}{f(1 + m)} - \frac{2^{\frac{1}{2}+m} m \cot(e + fx)(1 + \csc(e + fx))^{-\frac{1}{2}-m}(a + a \csc(e + fx))^m}{f(1 + m)} \end{aligned}$$

Mathematica [A] time = 0.431278, size = 131, normalized size = 1.2

$$\frac{\cot\left(\frac{1}{2}(e + fx)\right)\left(\tan\left(\frac{1}{2}(e + fx)\right) + 1\right)^{-2m}(a(\csc(e + fx) + 1))^m\left(-2\text{Hypergeometric2F1}\left(-2m - 1, -m - 1, -m, -\tan\left(\frac{1}{2}(e + fx)\right)\right)\right)}{f(1 + m)}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]^2*(a + a*Csc[e + f*x])^m,x]
```

```
[Out] -(Cot[(e + f*x)/2]*(a*(1 + Csc[e + f*x]))^m*(-2*Hypergeometric2F1[-1 - 2*m,
-1 - m, -m, -Tan[(e + f*x)/2]] + 2*Hypergeometric2F1[-1 - m, -2*m, -m, -Ta
```

$n[(e + f*x)/2]] + \text{Hypergeometric2F1}[-1 - m, -2*(1 + m), -m, -\text{Tan}[(e + f*x)/2]])) / (2*f*(1 + m)*(1 + \text{Tan}[(e + f*x)/2])^{(2*m)})$

Maple [F] time = 0.448, size = 0, normalized size = 0.

$$\int (\csc(fx + e))^2 (a + a \csc(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^2*(a+a*csc(f*x+e))^m,x)

[Out] int(csc(f*x+e)^2*(a+a*csc(f*x+e))^m,x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \csc(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(a+a*csc(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((a*csc(f*x + e) + a)^m*csc(f*x + e)^2, x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(a \csc(fx + e) + a\right)^m \csc(fx + e)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(a+a*csc(f*x+e))^m,x, algorithm="fricas")

[Out] integral((a*csc(f*x + e) + a)^m*csc(f*x + e)^2, x)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a(\csc(e + fx) + 1))^m \csc^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**2*(a+a*csc(f*x+e))**m,x)

[Out] Integral((a*(csc(e + f*x) + 1))**m*csc(e + f*x)**2, x)

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \csc(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(a+a*csc(f*x+e))^m,x, algorithm="giac")

[Out] integrate((a*csc(f*x + e) + a)^m*csc(f*x + e)^2, x)

3.32 $\int \csc(e + fx)(a + a \csc(e + fx))^m dx$

Optimal. Leaf size=74

$$\frac{2^{m+\frac{1}{2}} \cot(e + fx)(\csc(e + fx) + 1)^{-m-\frac{1}{2}}(a \csc(e + fx) + a)^m \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - m, \frac{3}{2}, \frac{1}{2}(1 - \csc(e + fx))\right)}{f}$$

[Out] -((2^(1/2 + m)*Cot[e + f*x]*(1 + Csc[e + f*x])^(-1/2 - m)*(a + a*Csc[e + f*x])^m*Hypergeometric2F1[1/2, 1/2 - m, 3/2, (1 - Csc[e + f*x])/2])/f)

Rubi [A] time = 0.0564489, antiderivative size = 74, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3828, 3827, 69}

$$\frac{2^{m+\frac{1}{2}} \cot(e + fx)(\csc(e + fx) + 1)^{-m-\frac{1}{2}}(a \csc(e + fx) + a)^m {}_2F_1\left(\frac{1}{2}, \frac{1}{2} - m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx))\right)}{f}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]*(a + a*Csc[e + f*x])^m,x]

[Out] -((2^(1/2 + m)*Cot[e + f*x]*(1 + Csc[e + f*x])^(-1/2 - m)*(a + a*Csc[e + f*x])^m*Hypergeometric2F1[1/2, 1/2 - m, 3/2, (1 - Csc[e + f*x])/2])/f)

Rule 3828

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_Symbol] :> Dist[(a^IntPart[m]*(a + b*Csc[e + f*x])^FracPart[m])/(1 + (b*Csc[e + f*x])/a)^FracPart[m], Int[(1 + (b*Csc[e + f*x])/a)^m*(d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && !GtQ[a, 0]
```

Rule 3827

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_Symbol] :> Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[((d*x)^(n - 1)*(a + b*x)^(m - 1/2))/Sqrt[a - b*x], x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && GtQ[a, 0]
```

Rule 69

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(((
a + b*x)^(m + 1)*Hypergeometric2F1[-n, m + 1, m + 2, -((d*(a + b*x))/(b*c -
a*d))])/(b*(m + 1)*(b/(b*c - a*d))^n), x] /; FreeQ[{a, b, c, d, m, n}, x]
&& NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d)
, 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-(d/(b*c - a*d)), 0]))
```

Rubi steps

$$\int \csc(e + fx)(a + a \csc(e + fx))^m dx = \left((1 + \csc(e + fx))^{-m} (a + a \csc(e + fx))^m \right) \int \csc(e + fx)(1 + \csc(e + fx))^m dx$$

$$= \frac{\left(\cot(e + fx)(1 + \csc(e + fx))^{-\frac{1}{2}-m} (a + a \csc(e + fx))^m \right) \text{Subst} \left(\int \frac{(1+x)^{-\frac{1}{2}+m}}{\sqrt{1-x}} dx, x, \frac{1 - \csc(e + fx)}{1 + \csc(e + fx)} \right)}{f \sqrt{1 - \csc(e + fx)}}$$

$$= -\frac{2^{\frac{1}{2}+m} \cot(e + fx)(1 + \csc(e + fx))^{-\frac{1}{2}-m} (a + a \csc(e + fx))^m {}_2F_1 \left(\frac{1}{2}, \frac{1}{2} - m; \frac{3}{2}; \frac{1 - \csc(e + fx)}{1 + \csc(e + fx)} \right)}{f}$$

Mathematica [A] time = 0.202145, size = 60, normalized size = 0.81

$$\frac{\left(\tan \left(\frac{1}{2}(e + fx) \right) + 1 \right)^{-2m} (a(\csc(e + fx) + 1))^m \text{Hypergeometric2F1} \left(-2m, -m, 1 - m, -\tan \left(\frac{1}{2}(e + fx) \right) \right)}{fm}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]*(a + a*Csc[e + f*x])^m,x]
```

```
[Out] -(((a*(1 + Csc[e + f*x]))^m*Hypergeometric2F1[-2*m, -m, 1 - m, -Tan[(e + f*x)/2]])/(f*m*(1 + Tan[(e + f*x)/2])^(2*m)))
```

Maple [F] time = 0.429, size = 0, normalized size = 0.

$$\int \csc(fx + e)(a + a \csc(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(f*x+e)*(a+a*csc(f*x+e))^m,x)
```

[Out] `int(csc(f*x+e)*(a+a*csc(f*x+e))^m,x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \csc(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(a+a*csc(f*x+e))^m,x, algorithm="maxima")`

[Out] `integrate((a*csc(f*x + e) + a)^m*csc(f*x + e), x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(a \csc(fx + e) + a\right)^m \csc(fx + e), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(a+a*csc(f*x+e))^m,x, algorithm="fricas")`

[Out] `integral((a*csc(f*x + e) + a)^m*csc(f*x + e), x)`

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a(\csc(e + fx) + 1))^m \csc(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(a+a*csc(f*x+e))**m,x)`

[Out] `Integral((a*(csc(e + f*x) + 1))**m*csc(e + f*x), x)`

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \csc(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)*(a+a*csc(f*x+e))^m,x, algorithm="giac")
```

```
[Out] integrate((a*csc(f*x + e) + a)^m*csc(f*x + e), x)
```

3.33 $\int (a + a \csc(e + fx))^m dx$

Optimal. Leaf size=84

$$-\frac{\sqrt{2} \cot(e + fx)(a \csc(e + fx) + a)^m F_1\left(m + \frac{1}{2}; \frac{1}{2}, 1; m + \frac{3}{2}; \frac{1}{2}(\csc(e + fx) + 1), \csc(e + fx) + 1\right)}{f(2m + 1)\sqrt{1 - \csc(e + fx)}}$$

[Out] -((Sqrt[2]*AppellF1[1/2 + m, 1/2, 1, 3/2 + m, (1 + Csc[e + f*x])/2, 1 + Csc[e + f*x]]*Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[1 - Csc[e + f*x]]))

Rubi [A] time = 0.0574375, antiderivative size = 84, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.25$, Rules used = {3779, 3778, 136}

$$-\frac{\sqrt{2} \cot(e + fx)(a \csc(e + fx) + a)^m F_1\left(m + \frac{1}{2}; \frac{1}{2}, 1; m + \frac{3}{2}; \frac{1}{2}(\csc(e + fx) + 1), \csc(e + fx) + 1\right)}{f(2m + 1)\sqrt{1 - \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Csc[e + f*x])^m,x]

[Out] -((Sqrt[2]*AppellF1[1/2 + m, 1/2, 1, 3/2 + m, (1 + Csc[e + f*x])/2, 1 + Csc[e + f*x]]*Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[1 - Csc[e + f*x]]))

Rule 3779

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^n_, x_Symbol] :> Dist[(a^IntPart[n]*(a + b*Csc[c + d*x])^FracPart[n])/(1 + (b*Csc[c + d*x])/a)^FracPart[n], Int[(1 + (b*Csc[c + d*x])/a)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]

Rule 3778

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^n_, x_Symbol] :> Dist[(a^n*Cot[c + d*x]/(d*Sqrt[1 + Csc[c + d*x]]*Sqrt[1 - Csc[c + d*x]]), Subst[Int[(1 + (b*x)/a)^(n - 1/2)/(x*Sqrt[1 - (b*x)/a]), x], x, Csc[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]

Rule 136

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_Symbol] :> Simp[((b*e - a*f)^p*(a + b*x)^(m + 1)*AppellF1[m + 1, -
n, -p, m + 2, -((d*(a + b*x))/(b*c - a*d)), -((f*(a + b*x))/(b*e - a*f))]/
(b^(p + 1)*(m + 1)*(b/(b*c - a*d))^n), x] /; FreeQ[{a, b, c, d, e, f, m, n},
x] && !IntegerQ[m] && !IntegerQ[n] && IntegerQ[p] && GtQ[b/(b*c - a*d),
0] && !(GtQ[d/(d*a - c*b), 0] && SimplerQ[c + d*x, a + b*x])
```

Rubi steps

$$\begin{aligned} \int (a + a \csc(e + fx))^m dx &= \left((1 + \csc(e + fx))^{-m} (a + a \csc(e + fx))^m \right) \int (1 + \csc(e + fx))^m dx \\ &= \frac{\left(\cot(e + fx) (1 + \csc(e + fx))^{-\frac{1}{2}-m} (a + a \csc(e + fx))^m \right) \text{Subst} \left(\int \frac{(1+x)^{-\frac{1}{2}+m}}{\sqrt{1-xx}} dx, x, \csc(e + fx) \right)}{f \sqrt{1 - \csc(e + fx)}} \\ &= \frac{\sqrt{2} F_1 \left(\frac{1}{2} + m; \frac{1}{2}, 1; \frac{3}{2} + m; \frac{1}{2} (1 + \csc(e + fx)), 1 + \csc(e + fx) \right) \cot(e + fx) (a + a \csc(e + fx))^m}{f (1 + 2m) \sqrt{1 - \csc(e + fx)}} \end{aligned}$$

Mathematica [F] time = 0.594863, size = 0, normalized size = 0.

$$\int (a + a \csc(e + fx))^m dx$$

Verification is Not applicable to the result.

[In] Integrate[(a + a*Csc[e + f*x])^m, x]

[Out] Integrate[(a + a*Csc[e + f*x])^m, x]

Maple [F] time = 0.287, size = 0, normalized size = 0.

$$\int (a + a \csc(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+a*csc(f*x+e))^m, x)

[Out] `int((a+a*csc(f*x+e))^m,x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))^m,x, algorithm="maxima")`

[Out] `integrate((a*csc(f*x + e) + a)^m, x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(a \csc(fx + e) + a\right)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))^m,x, algorithm="fricas")`

[Out] `integral((a*csc(f*x + e) + a)^m, x)`

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(e + fx) + a)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))**m,x)`

[Out] `Integral((a*csc(e + f*x) + a)**m, x)`

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+a*csc(f*x+e))^m,x, algorithm="giac")
```

```
[Out] integrate((a*csc(f*x + e) + a)^m, x)
```


3.34 $\int (a + a \csc(e + fx))^m \sin(e + fx) dx$

Optimal. Leaf size=83

$$\frac{\sqrt{2} \cot(e + fx)(a \csc(e + fx) + a)^m F_1\left(m + \frac{1}{2}; \frac{1}{2}, 2; m + \frac{3}{2}; \frac{1}{2}(\csc(e + fx) + 1), \csc(e + fx) + 1\right)}{f(2m + 1)\sqrt{1 - \csc(e + fx)}}$$

[Out] (Sqrt[2]*AppellF1[1/2 + m, 1/2, 2, 3/2 + m, (1 + Csc[e + f*x])/2, 1 + Csc[e + f*x]]*Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[1 - Csc[e + f*x]])

Rubi [A] time = 0.0895867, antiderivative size = 83, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3828, 3827, 136}

$$\frac{\sqrt{2} \cot(e + fx)(a \csc(e + fx) + a)^m F_1\left(m + \frac{1}{2}; \frac{1}{2}, 2; m + \frac{3}{2}; \frac{1}{2}(\csc(e + fx) + 1), \csc(e + fx) + 1\right)}{f(2m + 1)\sqrt{1 - \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Csc[e + f*x])^m*Sin[e + f*x], x]

[Out] (Sqrt[2]*AppellF1[1/2 + m, 1/2, 2, 3/2 + m, (1 + Csc[e + f*x])/2, 1 + Csc[e + f*x]]*Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[1 - Csc[e + f*x]])

Rule 3828

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] := Dist[(a^IntPart[m]*(a + b*Csc[e + f*x])^FracPart[m])/(1 + (b*Csc[e + f*x])/a)^FracPart[m], Int[(1 + (b*Csc[e + f*x])/a)^m*(d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && !GtQ[a, 0]

Rule 3827

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]])*Sqrt[a - b*Csc[e + f*x]], Subst[Int[((d*x)^(n - 1)*(a + b*x)^(m - 1/2))/Sqrt[a - b*x], x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, d, e, f, m, n}, x]

] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && GtQ[a, 0]

Rule 136

Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[((b*e - a*f)^p*(a + b*x)^(m + 1)*AppellF1[m + 1, -n, -p, m + 2, -((d*(a + b*x))/(b*c - a*d)), -((f*(a + b*x))/(b*e - a*f))]/(b^(p + 1)*(m + 1)*(b/(b*c - a*d))^n), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && IntegerQ[p] && GtQ[b/(b*c - a*d), 0] && !(GtQ[d/(d*a - c*b), 0] && SimplerQ[c + d*x, a + b*x])

Rubi steps

$$\begin{aligned} \int (a + a \csc(e + fx))^m \sin(e + fx) dx &= \left((1 + \csc(e + fx))^{-m} (a + a \csc(e + fx))^m \right) \int (1 + \csc(e + fx))^m \sin(e + fx) dx \\ &= \frac{\left(\cot(e + fx) (1 + \csc(e + fx))^{-\frac{1}{2}-m} (a + a \csc(e + fx))^m \right) \text{Subst} \left(\int \frac{(1+x)^{-\frac{1}{2}+m}}{\sqrt{1-xx^2}} dx, x \right)}{f \sqrt{1 - \csc(e + fx)}} \\ &= \frac{\sqrt{2} F_1 \left(\frac{1}{2} + m; \frac{1}{2}, 2; \frac{3}{2} + m; \frac{1}{2} (1 + \csc(e + fx)), 1 + \csc(e + fx) \right) \cot(e + fx) (a + a \csc(e + fx))^m}{f (1 + 2m) \sqrt{1 - \csc(e + fx)}} \end{aligned}$$

Mathematica [F] time = 3.64719, size = 0, normalized size = 0.

$$\int (a + a \csc(e + fx))^m \sin(e + fx) dx$$

Verification is Not applicable to the result.

[In] Integrate[(a + a*Csc[e + f*x])^m*Sin[e + f*x], x]

[Out] Integrate[(a + a*Csc[e + f*x])^m*Sin[e + f*x], x]

Maple [F] time = 0.678, size = 0, normalized size = 0.

$$\int (a + a \csc(fx + e))^m \sin(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+a*csc(f*x+e))^m*sin(f*x+e),x)`

[Out] `int((a+a*csc(f*x+e))^m*sin(f*x+e),x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \sin(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))^m*sin(f*x+e),x, algorithm="maxima")`

[Out] `integrate((a*csc(f*x + e) + a)^m*sin(f*x + e), x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(a \csc(fx + e) + a\right)^m \sin(fx + e), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))^m*sin(f*x+e),x, algorithm="fricas")`

[Out] `integral((a*csc(f*x + e) + a)^m*sin(f*x + e), x)`

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a(\csc(e + fx) + 1))^m \sin(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))**m*sin(f*x+e),x)`

[Out] `Integral((a*(csc(e + f*x) + 1))**m*sin(e + f*x), x)`

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \sin(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+a*csc(f*x+e))^m*sin(f*x+e),x, algorithm="giac")
```

```
[Out] integrate((a*csc(f*x + e) + a)^m*sin(f*x + e), x)
```

3.35 $\int (a + a \csc(e + fx))^m \sin^2(e + fx) dx$

Optimal. Leaf size=84

$$\frac{\sqrt{2} \cot(e + fx) (a \csc(e + fx) + a)^m F_1\left(m + \frac{1}{2}; \frac{1}{2}, 3; m + \frac{3}{2}; \frac{1}{2}(\csc(e + fx) + 1), \csc(e + fx) + 1\right)}{f(2m + 1)\sqrt{1 - \csc(e + fx)}}$$

[Out] -((Sqrt[2]*AppellF1[1/2 + m, 1/2, 3, 3/2 + m, (1 + Csc[e + f*x])/2, 1 + Csc[e + f*x]]*Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[1 - Csc[e + f*x]]))

Rubi [A] time = 0.105052, antiderivative size = 84, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3828, 3827, 136}

$$\frac{\sqrt{2} \cot(e + fx) (a \csc(e + fx) + a)^m F_1\left(m + \frac{1}{2}; \frac{1}{2}, 3; m + \frac{3}{2}; \frac{1}{2}(\csc(e + fx) + 1), \csc(e + fx) + 1\right)}{f(2m + 1)\sqrt{1 - \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Csc[e + f*x])^m*Sin[e + f*x]^2,x]

[Out] -((Sqrt[2]*AppellF1[1/2 + m, 1/2, 3, 3/2 + m, (1 + Csc[e + f*x])/2, 1 + Csc[e + f*x]]*Cot[e + f*x]*(a + a*Csc[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[1 - Csc[e + f*x]]))

Rule 3828

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] := Dist[(a^IntPart[m]*(a + b*Csc[e + f*x])^FracPart[m])/(1 + (b*Csc[e + f*x])/a)^FracPart[m], Int[(1 + (b*Csc[e + f*x])/a)^m*(d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && !GtQ[a, 0]

Rule 3827

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] := Dist[(a^2*d*Cot[e + f*x])/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]]), Subst[Int[((d*x)^(n - 1)*(a + b*x)^(m - 1/2))/Sqrt[a - b*x], x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, d, e, f, m, n}, x]

] && EqQ[a^2 - b^2, 0] && !IntegerQ[m] && GtQ[a, 0]

Rule 136

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_Symbol] := Simp[((b*e - a*f)^p*(a + b*x)^(m + 1)*AppellF1[m + 1, -
n, -p, m + 2, -((d*(a + b*x))/(b*c - a*d)), -((f*(a + b*x))/(b*e - a*f))]/
(b^(p + 1)*(m + 1)*(b/(b*c - a*d))^n), x] /; FreeQ[{a, b, c, d, e, f, m, n},
x] && !IntegerQ[m] && !IntegerQ[n] && IntegerQ[p] && GtQ[b/(b*c - a*d),
0] && !(GtQ[d/(d*a - c*b), 0] && SimplerQ[c + d*x, a + b*x])
```

Rubi steps

$$\begin{aligned} \int (a + a \csc(e + fx))^m \sin^2(e + fx) dx &= \left((1 + \csc(e + fx))^{-m} (a + a \csc(e + fx))^m \right) \int (1 + \csc(e + fx))^m \sin^2(e + fx) dx \\ &= \frac{\left(\cot(e + fx) (1 + \csc(e + fx))^{-\frac{1}{2}-m} (a + a \csc(e + fx))^m \right) \text{Subst} \left(\int \frac{(1+x)^{\frac{1}{2}+m}}{\sqrt{1-xx^3}} dx, \right)}{f \sqrt{1 - \csc(e + fx)}} \\ &= - \frac{\sqrt{2} F_1 \left(\frac{1}{2} + m; \frac{1}{2}, 3; \frac{3}{2} + m; \frac{1}{2} (1 + \csc(e + fx)), 1 + \csc(e + fx) \right) \cot(e + fx) (a + a \csc(e + fx))^m}{f (1 + 2m) \sqrt{1 - \csc(e + fx)}} \end{aligned}$$

Mathematica [F] time = 7.64792, size = 0, normalized size = 0.

$$\int (a + a \csc(e + fx))^m \sin^2(e + fx) dx$$

Verification is Not applicable to the result.

[In] Integrate[(a + a*Csc[e + f*x])^m*Sin[e + f*x]^2,x]

[Out] Integrate[(a + a*Csc[e + f*x])^m*Sin[e + f*x]^2, x]

Maple [F] time = 0.882, size = 0, normalized size = 0.

$$\int (a + a \csc(fx + e))^m (\sin(fx + e))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+a*csc(f*x+e))^m*sin(f*x+e)^2,x)`

[Out] `int((a+a*csc(f*x+e))^m*sin(f*x+e)^2,x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \sin(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))^m*sin(f*x+e)^2,x, algorithm="maxima")`

[Out] `integrate((a*csc(f*x + e) + a)^m*sin(f*x + e)^2, x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(-\left(\cos(fx + e)^2 - 1\right)\left(a \csc(fx + e) + a\right)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))^m*sin(f*x+e)^2,x, algorithm="fricas")`

[Out] `integral(-(cos(f*x + e)^2 - 1)*(a*csc(f*x + e) + a)^m, x)`

Sympy [F(-1)] time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*csc(f*x+e))^m*sin(f*x+e)**2,x)`

[Out] Timed out

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (a \csc(fx + e) + a)^m \sin(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+a*csc(f*x+e))^m*sin(f*x+e)^2,x, algorithm="giac")
```

```
[Out] integrate((a*csc(f*x + e) + a)^m*sin(f*x + e)^2, x)
```


3.36 $\int (a + b \csc(c + dx))^4 dx$

Optimal. Leaf size=107

$$\frac{b^2 (17a^2 + 2b^2) \cot(c + dx)}{3d} - \frac{2ab (2a^2 + b^2) \tanh^{-1}(\cos(c + dx))}{d} + a^4 x - \frac{4ab^3 \cot(c + dx) \csc(c + dx)}{3d} - \frac{b^2 \cot(c + dx)}{3d}$$

[Out] $a^4 x - (2 a b (2 a^2 + b^2) \operatorname{ArcTanh}[\cos[c + d x]]) / d - (b^2 (17 a^2 + 2 b^2) \cot[c + d x]) / (3 d) - (4 a b^3 \cot[c + d x] \operatorname{Csc}[c + d x]) / (3 d) - (b^2 \cot[c + d x] (a + b \operatorname{Csc}[c + d x])^2) / (3 d)$

Rubi [A] time = 0.109771, antiderivative size = 107, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 5, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3782, 4048, 3770, 3767, 8}

$$\frac{b^2 (17a^2 + 2b^2) \cot(c + dx)}{3d} - \frac{2ab (2a^2 + b^2) \tanh^{-1}(\cos(c + dx))}{d} + a^4 x - \frac{4ab^3 \cot(c + dx) \csc(c + dx)}{3d} - \frac{b^2 \cot(c + dx)}{3d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Csc[c + d*x])^4, x]

[Out] $a^4 x - (2 a b (2 a^2 + b^2) \operatorname{ArcTanh}[\cos[c + d x]]) / d - (b^2 (17 a^2 + 2 b^2) \cot[c + d x]) / (3 d) - (4 a b^3 \cot[c + d x] \operatorname{Csc}[c + d x]) / (3 d) - (b^2 \cot[c + d x] (a + b \operatorname{Csc}[c + d x])^2) / (3 d)$

Rule 3782

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^n, x_Symbol] :> -Simp[(b^2*Cot[c + d*x]*(a + b*Csc[c + d*x])^(n - 2))/(d*(n - 1)), x] + Dist[1/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 4048

Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> -Simp[(b*C*Csc[e + f*x]*Cot[e + f*x])/(2*f), x] + Dist[1/2, Int[Simp[2*A*a + (2*B*a + b*(2*A + C))*Csc[e + f*x] + 2*(a*C + B*b)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rubi steps

$$\begin{aligned}
 \int (a + b \csc(c + dx))^4 dx &= -\frac{b^2 \cot(c + dx)(a + b \csc(c + dx))^2}{3d} + \frac{1}{3} \int (a + b \csc(c + dx)) (3a^3 + b(9a^2 + 2b^2) \csc(c + dx) \\
 &= -\frac{4ab^3 \cot(c + dx) \csc(c + dx)}{3d} - \frac{b^2 \cot(c + dx)(a + b \csc(c + dx))^2}{3d} + \frac{1}{6} \int (6a^4 + 12ab(2a^2 + \\
 &= a^4 x - \frac{4ab^3 \cot(c + dx) \csc(c + dx)}{3d} - \frac{b^2 \cot(c + dx)(a + b \csc(c + dx))^2}{3d} + (2ab(2a^2 + b^2)) \int \\
 &= a^4 x - \frac{2ab(2a^2 + b^2) \tanh^{-1}(\cos(c + dx))}{d} - \frac{4ab^3 \cot(c + dx) \csc(c + dx)}{3d} - \frac{b^2 \cot(c + dx)(a + b \csc(c + dx))^2}{3d} \\
 &= a^4 x - \frac{2ab(2a^2 + b^2) \tanh^{-1}(\cos(c + dx))}{d} - \frac{b^2(17a^2 + 2b^2) \cot(c + dx)}{3d} - \frac{4ab^3 \cot(c + dx) \csc(c + dx)}{3d}
 \end{aligned}$$

Mathematica [B] time = 6.23832, size = 568, normalized size = 5.31

$$\frac{\sin^4(c + dx) \csc\left(\frac{1}{2}(c + dx)\right) \left(b^4 \left(-\cos\left(\frac{1}{2}(c + dx)\right)\right) - 9a^2 b^2 \cos\left(\frac{1}{2}(c + dx)\right)\right) (a + b \csc(c + dx))^4}{3d(a \sin(c + dx) + b)^4} + \frac{2(2a^3 b + ab^3) \sin^4(c + dx)}{3d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Csc[c + d*x])^4, x]

[Out] (a^4*(c + d*x)*(a + b*Csc[c + d*x])^4*Sin[c + d*x]^4)/(d*(b + a*Sin[c + d*x])^4) + ((-9*a^2*b^2*Cos[(c + d*x)/2] - b^4*Cos[(c + d*x)/2])*Csc[(c + d*x)/2]*(a + b*Csc[c + d*x])^4*Sin[c + d*x]^4)/(3*d*(b + a*Sin[c + d*x])^4) - (a*b^3*Csc[(c + d*x)/2]^2*(a + b*Csc[c + d*x])^4*Sin[c + d*x]^4)/(2*d*(b + a

$$\begin{aligned} & * \sin[c + dx]^4) - (b^4 \cot[(c + dx)/2] \operatorname{Csc}[(c + dx)/2]^2 (a + b \operatorname{Csc}[c + dx])^4 \sin[c + dx]^4) / (24 d (b + a \sin[c + dx])^4) - (2 (2 a^3 b + a b^3) (a + b \operatorname{Csc}[c + dx])^4 \operatorname{Log}[\cos[(c + dx)/2]] \sin[c + dx]^4) / (d (b + a \sin[c + dx])^4) + (2 (2 a^3 b + a b^3) (a + b \operatorname{Csc}[c + dx])^4 \operatorname{Log}[\sin[(c + dx)/2]] \sin[c + dx]^4) / (d (b + a \sin[c + dx])^4) + (a b^3 (a + b \operatorname{Csc}[c + dx])^4 \operatorname{Sec}[(c + dx)/2]^2 \sin[c + dx]^4) / (2 d (b + a \sin[c + dx])^4) + ((a + b \operatorname{Csc}[c + dx])^4 \operatorname{Sec}[(c + dx)/2] (9 a^2 b^2 \sin[(c + dx)/2] + b^4 \sin[(c + dx)/2]) \sin[c + dx]^4) / (3 d (b + a \sin[c + dx])^4) + (b^4 (a + b \operatorname{Csc}[c + dx])^4 \operatorname{Sec}[(c + dx)/2]^2 \sin[c + dx]^4 \tan[(c + dx)/2]) / (24 d (b + a \sin[c + dx])^4) \end{aligned}$$

Maple [A] time = 0.05, size = 139, normalized size = 1.3

$$a^4 x + \frac{a^4 c}{d} + 4 \frac{a^3 b \ln(\csc(dx + c) - \cot(dx + c))}{d} - 6 \frac{a^2 b^2 \cot(dx + c)}{d} - 2 \frac{a b^3 \cot(dx + c) \csc(dx + c)}{d} + 2 \frac{a b^3 \ln(\csc(dx + c) - \cot(dx + c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+b*csc(d*x+c))^4,x)

[Out] a^4*x+1/d*a^4*c+4/d*a^3*b*ln(csc(d*x+c)-cot(d*x+c))-6/d*a^2*b^2*cot(d*x+c)-2*a*b^3*cot(d*x+c)*csc(d*x+c)/d+2/d*a*b^3*ln(csc(d*x+c)-cot(d*x+c))-2/3/d*b^4*cot(d*x+c)-1/3/d*b^4*cot(d*x+c)*csc(d*x+c)^2

Maxima [A] time = 1.00559, size = 169, normalized size = 1.58

$$a^4 x + \frac{a b^3 \left(\frac{2 \cos(dx+c)}{\cos(dx+c)^2-1} - \log(\cos(dx+c)+1) + \log(\cos(dx+c)-1) \right)}{d} - \frac{4 a^3 b \log(\cot(dx+c) + \csc(dx+c))}{d} - \frac{6}{d \tan(dx+c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))^4,x, algorithm="maxima")

[Out] a^4*x + a*b^3*(2*cos(d*x + c)/(cos(d*x + c)^2 - 1) - log(cos(d*x + c) + 1) + log(cos(d*x + c) - 1))/d - 4*a^3*b*log(cot(d*x + c) + csc(d*x + c))/d - 6*a^2*b^2/(d*tan(d*x + c)) - 1/3*(3*tan(d*x + c)^2 + 1)*b^4/(d*tan(d*x + c)^3)

Fricas [B] time = 0.523606, size = 524, normalized size = 4.9

$$2(9a^2b^2 + b^4)\cos(dx + c)^3 - 3(2a^3b + ab^3 - (2a^3b + ab^3)\cos(dx + c)^2)\log\left(\frac{1}{2}\cos(dx + c) + \frac{1}{2}\right)\sin(dx + c) + 3(2a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))^4,x, algorithm="fricas")

[Out]
$$-1/3*(2*(9*a^2*b^2 + b^4)*\cos(d*x + c)^3 - 3*(2*a^3*b + a*b^3 - (2*a^3*b + a*b^3)*\cos(d*x + c)^2)*\log(1/2*\cos(d*x + c) + 1/2)*\sin(d*x + c) + 3*(2*a^3*b + a*b^3 - (2*a^3*b + a*b^3)*\cos(d*x + c)^2)*\log(-1/2*\cos(d*x + c) + 1/2)*\sin(d*x + c) - 3*(6*a^2*b^2 + b^4)*\cos(d*x + c) - 3*(a^4*d*x*\cos(d*x + c)^2 - a^4*d*x + 2*a*b^3*\cos(d*x + c))*\sin(d*x + c))/((d*\cos(d*x + c)^2 - d)*\sin(d*x + c))$$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a + b \csc(c + dx))^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))**4,x)

[Out] Integral((a + b*csc(c + d*x))**4, x)

Giac [B] time = 1.59847, size = 277, normalized size = 2.59

$$b^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 12 ab^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 24 (dx + c)a^4 + 72 a^2 b^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 9 b^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 48 (2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))^4,x, algorithm="giac")

```
[Out] 1/24*(b^4*tan(1/2*d*x + 1/2*c)^3 + 12*a*b^3*tan(1/2*d*x + 1/2*c)^2 + 24*(d*x + c)*a^4 + 72*a^2*b^2*tan(1/2*d*x + 1/2*c) + 9*b^4*tan(1/2*d*x + 1/2*c) + 48*(2*a^3*b + a*b^3)*log(abs(tan(1/2*d*x + 1/2*c))) - (176*a^3*b*tan(1/2*d*x + 1/2*c)^3 + 88*a*b^3*tan(1/2*d*x + 1/2*c)^3 + 72*a^2*b^2*tan(1/2*d*x + 1/2*c)^2 + 9*b^4*tan(1/2*d*x + 1/2*c)^2 + 12*a*b^3*tan(1/2*d*x + 1/2*c) + b^4)/tan(1/2*d*x + 1/2*c)^3)/d
```

3.37 $\int (a + b \csc(c + dx))^3 dx$

Optimal. Leaf size=73

$$-\frac{b(6a^2 + b^2) \tanh^{-1}(\cos(c + dx))}{2d} + a^3x - \frac{5ab^2 \cot(c + dx)}{2d} - \frac{b^2 \cot(c + dx)(a + b \csc(c + dx))}{2d}$$

[Out] $a^3x - (b(6a^2 + b^2) \operatorname{ArcTanh}[\cos[c + dx]])/(2d) - (5ab^2 \cot[c + dx]x)/(2d) - (b^2 \cot[c + dx](a + b \csc[c + dx]))/(2d)$

Rubi [A] time = 0.0469065, antiderivative size = 73, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3782, 3770, 3767, 8}

$$-\frac{b(6a^2 + b^2) \tanh^{-1}(\cos(c + dx))}{2d} + a^3x - \frac{5ab^2 \cot(c + dx)}{2d} - \frac{b^2 \cot(c + dx)(a + b \csc(c + dx))}{2d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a + b \csc[c + dx])^3, x]$

[Out] $a^3x - (b(6a^2 + b^2) \operatorname{ArcTanh}[\cos[c + dx]])/(2d) - (5ab^2 \cot[c + dx]x)/(2d) - (b^2 \cot[c + dx](a + b \csc[c + dx]))/(2d)$

Rule 3782

$\operatorname{Int}[(\csc[(c_.) + (d_.)(x_)]*(b_.) + (a_.))^{(n_.)}, x_Symbol] \rightarrow -\operatorname{Simp}[(b^2 \cot[c + dx]*(a + b \csc[c + dx])^{(n-2)})/(d*(n-1)), x] + \operatorname{Dist}[1/(n-1), \operatorname{Int}[(a + b \csc[c + dx])^{(n-3)} * \operatorname{Simp}[a^3*(n-1) + (b*(b^2*(n-2) + 3*a^2*(n-1)))*\csc[c + dx] + (a*b^2*(3*n-4))*\csc[c + dx]^2, x], x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 3770

$\operatorname{Int}[\csc[(c_.) + (d_.)(x_)], x_Symbol] \rightarrow -\operatorname{Simp}[\operatorname{ArcTanh}[\cos[c + dx]]/d, x] /;$ FreeQ[{c, d}, x]

Rule 3767

$\operatorname{Int}[\csc[(c_.) + (d_.)(x_)]^{(n_.)}, x_Symbol] \rightarrow -\operatorname{Dist}[d^{(-1)}, \operatorname{Subst}[\operatorname{Int}[\operatorname{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \cot[c + dx]], x] /;$ FreeQ[{c,

d}, x] && IGtQ[n/2, 0]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rubi steps

$$\begin{aligned} \int (a + b \csc(c + dx))^3 dx &= -\frac{b^2 \cot(c + dx)(a + b \csc(c + dx))}{2d} + \frac{1}{2} \int (2a^3 + b(6a^2 + b^2) \csc(c + dx) + 5ab^2 \csc^2(c + dx)) dx \\ &= a^3 x - \frac{b^2 \cot(c + dx)(a + b \csc(c + dx))}{2d} + \frac{1}{2} (5ab^2) \int \csc^2(c + dx) dx + \frac{1}{2} (b(6a^2 + b^2)) \int \csc(c + dx) dx \\ &= a^3 x - \frac{b(6a^2 + b^2) \tanh^{-1}(\cos(c + dx))}{2d} - \frac{b^2 \cot(c + dx)(a + b \csc(c + dx))}{2d} - \frac{(5ab^2) \text{Subst}}{2d} \\ &= a^3 x - \frac{b(6a^2 + b^2) \tanh^{-1}(\cos(c + dx))}{2d} - \frac{5ab^2 \cot(c + dx)}{2d} - \frac{b^2 \cot(c + dx)(a + b \csc(c + dx))}{2d} \end{aligned}$$

Mathematica [B] time = 0.648397, size = 152, normalized size = 2.08

$$\frac{24a^2b \log\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) - 24a^2b \log\left(\cos\left(\frac{1}{2}(c + dx)\right)\right) + 8a^3c + 8a^3dx + 12ab^2 \tan\left(\frac{1}{2}(c + dx)\right) - 12ab^2 \cot\left(\frac{1}{2}(c + dx)\right)}{8d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Csc[c + d*x])^3,x]

[Out] (8*a^3*c + 8*a^3*d*x - 12*a*b^2*Cot[(c + d*x)/2] - b^3*Csc[(c + d*x)/2]^2 - 24*a^2*b*Log[Cos[(c + d*x)/2]] - 4*b^3*Log[Cos[(c + d*x)/2]] + 24*a^2*b*Log[Sin[(c + d*x)/2]] + 4*b^3*Log[Sin[(c + d*x)/2]] + b^3*Sec[(c + d*x)/2]^2 + 12*a*b^2*Tan[(c + d*x)/2])/(8*d)

Maple [A] time = 0.044, size = 99, normalized size = 1.4

$$a^3x + \frac{a^3c}{d} + 3 \frac{a^2b \ln(\csc(dx + c) - \cot(dx + c))}{d} - 3 \frac{ab^2 \cot(dx + c)}{d} - \frac{b^3 \csc(dx + c) \cot(dx + c)}{2d} + \frac{b^3 \ln(\csc(dx + c) - \cot(dx + c))}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*csc(d*x+c))^3,x)`

[Out] $a^3x + 1/d a^3c + 3/d a^2b \ln(\csc(dx+c) - \cot(dx+c)) - 3ab^2 \cot(dx+c)/d - 1/2/d b^3 \csc(dx+c) \cot(dx+c) + 1/2/d b^3 \ln(\csc(dx+c) - \cot(dx+c))$

Maxima [A] time = 1.00415, size = 128, normalized size = 1.75

$$a^3x + \frac{b^3 \left(\frac{2 \cos(dx+c)}{\cos(dx+c)^2 - 1} - \log(\cos(dx+c) + 1) + \log(\cos(dx+c) - 1) \right)}{4d} - \frac{3a^2b \log(\cot(dx+c) + \csc(dx+c))}{d} - \frac{3ab}{d \tan(dx+c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*csc(d*x+c))^3,x, algorithm="maxima")`

[Out] $a^3x + 1/4 b^3 (2 \cos(dx+c) / (\cos(dx+c)^2 - 1) - \log(\cos(dx+c) + 1) + \log(\cos(dx+c) - 1)) / d - 3a^2b \log(\cot(dx+c) + \csc(dx+c)) / d - 3ab^2 / (d \tan(dx+c))$

Fricas [B] time = 0.510696, size = 383, normalized size = 5.25

$$\frac{4a^3 dx \cos(dx+c)^2 - 4a^3 dx + 12ab^2 \cos(dx+c) \sin(dx+c) + 2b^3 \cos(dx+c) + (6a^2b + b^3 - (6a^2b + b^3) \cos(dx+c))}{4(d \cos(dx+c)^2 - d)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*csc(d*x+c))^3,x, algorithm="fricas")`

[Out] $1/4 (4a^3 dx \cos(dx+c)^2 - 4a^3 dx + 12a^2b \cos(dx+c) \sin(dx+c) + 2b^3 \cos(dx+c) + (6a^2b + b^3 - (6a^2b + b^3) \cos(dx+c)^2) \log(1/2 \cos(dx+c) + 1/2) - (6a^2b + b^3 - (6a^2b + b^3) \cos(dx+c)^2) \log(-1/2 \cos(dx+c) + 1/2)) / (d \cos(dx+c)^2 - d)$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a + b \csc(c + dx))^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))**3,x)

[Out] Integral((a + b*csc(c + d*x))**3, x)

Giac [B] time = 1.47343, size = 181, normalized size = 2.48

$$b^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 8(dx + c)a^3 + 12ab^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 4(6a^2b + b^3) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right|\right) - \frac{36a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{8d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))^3,x, algorithm="giac")

[Out] 1/8*(b^3*tan(1/2*d*x + 1/2*c)^2 + 8*(d*x + c)*a^3 + 12*a*b^2*tan(1/2*d*x + 1/2*c) + 4*(6*a^2*b + b^3)*log(abs(tan(1/2*d*x + 1/2*c)))) - (36*a^2*b*tan(1/2*d*x + 1/2*c)^2 + 6*b^3*tan(1/2*d*x + 1/2*c)^2 + 12*a*b^2*tan(1/2*d*x + 1/2*c) + b^3)/tan(1/2*d*x + 1/2*c)^2/d

3.38 $\int (a + b \csc(c + dx))^2 dx$

Optimal. Leaf size=34

$$a^2x - \frac{2ab \tanh^{-1}(\cos(c + dx))}{d} - \frac{b^2 \cot(c + dx)}{d}$$

[Out] $a^2x - (2ab \operatorname{ArcTanh}[\cos(c + dx)])/d - (b^2 \cot(c + dx))/d$

Rubi [A] time = 0.025757, antiderivative size = 34, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3773, 3770, 3767, 8}

$$a^2x - \frac{2ab \tanh^{-1}(\cos(c + dx))}{d} - \frac{b^2 \cot(c + dx)}{d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a + b \csc(c + dx))^2, x]$

[Out] $a^2x - (2ab \operatorname{ArcTanh}[\cos(c + dx)])/d - (b^2 \cot(c + dx))/d$

Rule 3773

$\operatorname{Int}[(\csc[(c_.) + (d_.)(x_.)]*(b_.) + (a_.))^2, x_Symbol] \rightarrow \operatorname{Simp}[a^2x, x] + (\operatorname{Dist}[2*a*b, \operatorname{Int}[\csc[c + dx], x], x] + \operatorname{Dist}[b^2, \operatorname{Int}[\csc[c + dx]^2, x], x]) /;$ $\operatorname{FreeQ}[\{a, b, c, d\}, x]$

Rule 3770

$\operatorname{Int}[\csc[(c_.) + (d_.)(x_.)], x_Symbol] \rightarrow -\operatorname{Simp}[\operatorname{ArcTanh}[\cos[c + dx]]/d, x] /;$ $\operatorname{FreeQ}[\{c, d\}, x]$

Rule 3767

$\operatorname{Int}[\csc[(c_.) + (d_.)(x_.)]^{(n_.)}, x_Symbol] \rightarrow -\operatorname{Dist}[d^{(-1)}, \operatorname{Subst}[\operatorname{Int}[\operatorname{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \cot[c + dx]], x] /;$ $\operatorname{FreeQ}[\{c, d\}, x] \ \&\& \ \operatorname{IGtQ}[n/2, 0]$

Rule 8

$\operatorname{Int}[a_, x_Symbol] \rightarrow \operatorname{Simp}[a*x, x] /;$ $\operatorname{FreeQ}[a, x]$

Rubi steps

$$\begin{aligned}
\int (a + b \csc(c + dx))^2 dx &= a^2x + (2ab) \int \csc(c + dx) dx + b^2 \int \csc^2(c + dx) dx \\
&= a^2x - \frac{2ab \tanh^{-1}(\cos(c + dx))}{d} - \frac{b^2 \operatorname{Subst}(\int 1 dx, x, \cot(c + dx))}{d} \\
&= a^2x - \frac{2ab \tanh^{-1}(\cos(c + dx))}{d} - \frac{b^2 \cot(c + dx)}{d}
\end{aligned}$$

Mathematica [B] time = 0.181533, size = 76, normalized size = 2.24

$$\frac{2a \left(ac + adx + 2b \log \left(\sin \left(\frac{1}{2}(c + dx) \right) \right) - 2b \log \left(\cos \left(\frac{1}{2}(c + dx) \right) \right) \right) + b^2 \tan \left(\frac{1}{2}(c + dx) \right) + b^2 \left(-\cot \left(\frac{1}{2}(c + dx) \right) \right)}{2d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Csc[c + d*x])^2, x]

[Out] $(-(b^2 \cot[(c + d*x)/2]) + 2*a*(a*c + a*d*x - 2*b*\log[\cos[(c + d*x)/2]] + 2*b*\log[\sin[(c + d*x)/2]]) + b^2*\tan[(c + d*x)/2])/(2*d)$

Maple [A] time = 0.029, size = 52, normalized size = 1.5

$$a^2x - \frac{b^2 \cot(dx + c)}{d} + 2 \frac{ab \ln(\csc(dx + c) - \cot(dx + c))}{d} + \frac{a^2c}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+b*csc(d*x+c))^2, x)

[Out] $a^2*x - b^2*\cot(d*x+c)/d + 2/d*a*b*\ln(\csc(d*x+c) - \cot(d*x+c)) + 1/d*a^2*c$

Maxima [A] time = 0.996058, size = 58, normalized size = 1.71

$$a^2x - \frac{2ab \log(\cot(dx + c) + \csc(dx + c))}{d} - \frac{b^2}{d \tan(dx + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))^2,x, algorithm="maxima")

[Out] $a^2x - 2ab \log(\cot(dx + c) + \csc(dx + c))/d - b^2/(d \tan(dx + c))$

Fricas [B] time = 0.503685, size = 209, normalized size = 6.15

$$\frac{a^2 dx \sin(dx + c) - ab \log\left(\frac{1}{2} \cos(dx + c) + \frac{1}{2}\right) \sin(dx + c) + ab \log\left(-\frac{1}{2} \cos(dx + c) + \frac{1}{2}\right) \sin(dx + c) - b^2 \cos(dx + c)}{d \sin(dx + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))^2,x, algorithm="fricas")

[Out] $(a^2 d x \sin(dx + c) - a b \log(1/2 \cos(dx + c) + 1/2) \sin(dx + c) + a b \log(-1/2 \cos(dx + c) + 1/2) \sin(dx + c) - b^2 \cos(dx + c)) / (d \sin(dx + c))$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a + b \csc(c + dx))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(d*x+c))**2,x)

[Out] Integral((a + b*csc(c + d*x))**2, x)

Giac [B] time = 1.42798, size = 100, normalized size = 2.94

$$\frac{2(dx + c)a^2 + 4ab \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right|\right) + b^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - \frac{4ab \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + b^2}{\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*csc(d*x+c))^2,x, algorithm="giac")
```

```
[Out] 1/2*(2*(d*x + c)*a^2 + 4*a*b*log(abs(tan(1/2*d*x + 1/2*c)))) + b^2*tan(1/2*d*x + 1/2*c) - (4*a*b*tan(1/2*d*x + 1/2*c) + b^2)/tan(1/2*d*x + 1/2*c))/d
```

$$3.39 \quad \int \frac{\csc^5(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=112

$$-\frac{(3a^2 + 2b^2) \cot(x)}{3b^3} + \frac{a(2a^2 + b^2) \tanh^{-1}(\cos(x))}{2b^4} - \frac{2a^4 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{b^4 \sqrt{a^2-b^2}} + \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b}$$

[Out] (a*(2*a^2 + b^2)*ArcTanh[Cos[x]])/(2*b^4) - (2*a^4*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(b^4*Sqrt[a^2 - b^2]) - ((3*a^2 + 2*b^2)*Cot[x])/(3*b^3) + (a*Cot[x]*Csc[x])/(2*b^2) - (Cot[x]*Csc[x]^2)/(3*b)

Rubi [A] time = 0.406685, antiderivative size = 112, normalized size of antiderivative = 1., number of steps used = 9, number of rules used = 9, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {3851, 4092, 4082, 3998, 3770, 3831, 2660, 618, 206}

$$-\frac{(3a^2 + 2b^2) \cot(x)}{3b^3} + \frac{a(2a^2 + b^2) \tanh^{-1}(\cos(x))}{2b^4} - \frac{2a^4 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{b^4 \sqrt{a^2-b^2}} + \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Csc[x]^5/(a + b*Csc[x]), x]

[Out] (a*(2*a^2 + b^2)*ArcTanh[Cos[x]])/(2*b^4) - (2*a^4*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(b^4*Sqrt[a^2 - b^2]) - ((3*a^2 + 2*b^2)*Cot[x])/(3*b^3) + (a*Cot[x]*Csc[x])/(2*b^2) - (Cot[x]*Csc[x]^2)/(3*b)

Rule 3851

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> -Simp[(d^3*Cot[e + f*x]*(d*Csc[e + f*x])^(n - 3))/(b*f*(n - 2)), x] + Dist[d^3/(b*(n - 2)), Int[((d*Csc[e + f*x])^(n - 3)*Simp[a*(n - 3) + b*(n - 3)*Csc[e + f*x] - a*(n - 2)*Csc[e + f*x]^2, x])/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 3]

Rule 4092

Int[csc[(e_.) + (f_.)*(x_.)]^2*((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^m, x

```

_Symbol] := -Simp[(C*Csc[e + f*x]*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)
)/(b*f*(m + 3)), x] + Dist[1/(b*(m + 3)), Int[Csc[e + f*x]*(a + b*Csc[e + f
*x])^m*Simp[a*C + b*(C*(m + 2) + A*(m + 3))*Csc[e + f*x] - (2*a*C - b*B*(m
+ 3))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && N
eQ[a^2 - b^2, 0] && !LtQ[m, -1]

```

Rule 4082

```

Int[csc[(e_.) + (f_.)*(x_)]*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e
_.) + (f_.)*(x_)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_S
ymbol] := -Simp[(C*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1))/(b*f*(m + 2))
, x] + Dist[1/(b*(m + 2)), Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*Simp[b*A
*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Csc[e + f*x], x], x], x] /; Fr
eeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]

```

Rule 3998

```

Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)))/(csc[(
e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Dist[B/b, Int[Csc[e + f*x],
x], x] + Dist[(A*b - a*B)/b, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x]
/; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0]

```

Rule 3770

```

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]

```

Rule 3831

```

Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbo
l] := Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f
}, x] && NeQ[a^2 - b^2, 0]

```

Rule 2660

```

Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = Fre
eFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*
e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[
a^2 - b^2, 0]

```

Rule 618

```

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},

```

x] && NeQ[b^2 - 4*a*c, 0]

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rubi steps

$$\begin{aligned}
 \int \frac{\csc^5(x)}{a + b \csc(x)} dx &= -\frac{\cot(x) \csc^2(x)}{3b} + \frac{\int \frac{\csc^2(x)(2a+2b \csc(x)-3a \csc^2(x))}{a+b \csc(x)} dx}{3b} \\
 &= \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b} + \frac{\int \frac{\csc(x)(-3a^2+ab \csc(x)+2(3a^2+2b^2) \csc^2(x))}{a+b \csc(x)} dx}{6b^2} \\
 &= -\frac{(3a^2 + 2b^2) \cot(x)}{3b^3} + \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b} + \frac{\int \frac{\csc(x)(-3a^2b-3a(2a^2+b^2) \csc(x))}{a+b \csc(x)} dx}{6b^3} \\
 &= -\frac{(3a^2 + 2b^2) \cot(x)}{3b^3} + \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b} + \frac{a^4 \int \frac{\csc(x)}{a+b \csc(x)} dx}{b^4} - \frac{(a(2a^2 + b^2)) \int \csc(x)}{2b^4} \\
 &= \frac{a(2a^2 + b^2) \tanh^{-1}(\cos(x))}{2b^4} - \frac{(3a^2 + 2b^2) \cot(x)}{3b^3} + \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b} + \frac{a^4 \int \frac{1}{1+\frac{a \sin(x)}{b}}}{b^5} \\
 &= \frac{a(2a^2 + b^2) \tanh^{-1}(\cos(x))}{2b^4} - \frac{(3a^2 + 2b^2) \cot(x)}{3b^3} + \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b} + \frac{(2a^4) \text{Subst}}{b^5} \\
 &= \frac{a(2a^2 + b^2) \tanh^{-1}(\cos(x))}{2b^4} - \frac{(3a^2 + 2b^2) \cot(x)}{3b^3} + \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b} - \frac{(4a^4) \text{Subst}}{b^5} \\
 &= \frac{a(2a^2 + b^2) \tanh^{-1}(\cos(x))}{2b^4} - \frac{2a^4 \tanh^{-1}\left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2 - b^2}}\right)}{b^4 \sqrt{a^2 - b^2}} - \frac{(3a^2 + 2b^2) \cot(x)}{3b^3} + \frac{a \cot(x) \csc(x)}{2b^2} - \frac{\cot(x) \csc^2(x)}{3b}
 \end{aligned}$$

Mathematica [A] time = 1.63056, size = 125, normalized size = 1.12

$$\frac{24a^4 \tan^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2 - a^2}}\right)}{\sqrt{b^2 - a^2}} + b(3a^2 + 2b^2) \cos(3x) \csc^3(x) - 3b \cot(x) \csc(x) \left((a^2 + 2b^2) \csc(x) - 2ab \right) + 6a(2a^2 + b^2) \left(\log(\cos(x)) \right)$$

12b⁴

Antiderivative was successfully verified.

[In] Integrate[Csc[x]^5/(a + b*Csc[x]),x]

[Out] $\left(\frac{24a^4 \operatorname{ArcTan}\left[\frac{a + b \tan(x/2)}{\sqrt{-a^2 + b^2}}\right]}{\sqrt{-a^2 + b^2}}\right) / \sqrt{-a^2 + b^2} + b(3a^2 + 2b^2) \cos(3x) \operatorname{Csc}[x]^3 - 3b \cot(x) \operatorname{Csc}[x](-2ab + (a^2 + 2b^2) \operatorname{Csc}[x]) + 6a(2a^2 + b^2)(\log(\cos(x/2)) - \log(\sin(x/2))) / (12b^4)$

Maple [A] time = 0.05, size = 162, normalized size = 1.5

$$\frac{1}{24b} \left(\tan\left(\frac{x}{2}\right)\right)^3 - \frac{a}{8b^2} \left(\tan\left(\frac{x}{2}\right)\right)^2 + \frac{a^2}{2b^3} \tan\left(\frac{x}{2}\right) + \frac{3}{8b} \tan\left(\frac{x}{2}\right) + 2 \frac{a^4}{b^4 \sqrt{-a^2 + b^2}} \arctan\left(\frac{1}{2} \frac{2b \tan(x/2) + 2a}{\sqrt{-a^2 + b^2}}\right) - \frac{1}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(x)^5/(a+b*csc(x)),x)

[Out] $\frac{1}{24} \frac{1}{b} \tan(1/2*x)^3 - \frac{1}{8} \frac{1}{b^2} a \tan(1/2*x)^2 + \frac{1}{2} \frac{1}{b^3} a^2 \tan(1/2*x) + \frac{3}{8} \frac{1}{b} \tan(1/2*x) + 2 \frac{a^4}{b^4 \sqrt{-a^2 + b^2}} \arctan\left(\frac{1}{2} \frac{2*b*\tan(1/2*x) + 2*a}{\sqrt{-a^2 + b^2}}\right) - \frac{1}{24} \frac{1}{b} \tan(1/2*x)^3 - \frac{1}{2} \frac{1}{b^3} \frac{1}{\tan(1/2*x)} * a^2 - \frac{3}{8} \frac{1}{b} \frac{1}{\tan(1/2*x)} + \frac{1}{8} \frac{a}{b^2} \frac{1}{\tan(1/2*x)^2} - \frac{1}{b^4} a^3 \ln(\tan(1/2*x)) - \frac{1}{2} \frac{1}{b^2} a \ln(\tan(1/2*x))$

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^5/(a+b*csc(x)),x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [B] time = 0.898211, size = 1378, normalized size = 12.3

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^5/(a+b*csc(x)),x, algorithm="fricas")

[Out]
$$\begin{aligned} & \left[\frac{1}{12} (4(3a^4b - a^2b^3 - 2b^5)\cos(x)^3 - 6(a^4\cos(x)^2 - a^4)\sqrt{(a^2 - b^2)\log(-((a^2 - 2b^2)\cos(x)^2 + 2ab\sin(x) + a^2 + b^2 - 2(b\cos(x)\sin(x) + a\cos(x))\sqrt{a^2 - b^2})) / (a^2\cos(x)^2 - 2ab\sin(x) - a^2 - b^2))\sin(x) + 6(a^3b^2 - ab^4)\cos(x)\sin(x) + 3(2a^5 - a^3b^2 - ab^4 - (2a^5 - a^3b^2 - ab^4)\cos(x)^2)\log(1/2\cos(x) + 1/2)\sin(x) - 3(2a^5 - a^3b^2 - ab^4 - (2a^5 - a^3b^2 - ab^4)\cos(x)^2)\log(-1/2\cos(x) + 1/2)\sin(x) - 12(a^4b - b^5)\cos(x)) / ((a^2b^4 - b^6 - (a^2b^4 - b^6)\cos(x)^2)\sin(x)), \right. \\ & \left. \frac{1}{12} (4(3a^4b - a^2b^3 - 2b^5)\cos(x)^3 + 12(a^4\cos(x)^2 - a^4)\sqrt{-a^2 + b^2}\arctan(-\sqrt{-a^2 + b^2}(b\sin(x) + a) / ((a^2 - b^2)\cos(x)))\sin(x) + 6(a^3b^2 - ab^4)\cos(x)\sin(x) + 3(2a^5 - a^3b^2 - ab^4 - (2a^5 - a^3b^2 - ab^4)\cos(x)^2)\log(1/2\cos(x) + 1/2)\sin(x) - 3(2a^5 - a^3b^2 - ab^4 - (2a^5 - a^3b^2 - ab^4)\cos(x)^2)\log(-1/2\cos(x) + 1/2)\sin(x) - 12(a^4b - b^5)\cos(x)) / ((a^2b^4 - b^6 - (a^2b^4 - b^6)\cos(x)^2)\sin(x)) \right] \end{aligned}$$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\csc^5(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)**5/(a+b*csc(x)),x)

[Out] Integral(csc(x)**5/(a + b*csc(x)), x)

Giac [A] time = 1.38737, size = 262, normalized size = 2.34

$$\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}} \right) \right) a^4}{\sqrt{-a^2 + b^2} b^4} + \frac{b^2 \tan\left(\frac{1}{2}x\right)^3 - 3ab \tan\left(\frac{1}{2}x\right)^2 + 12a^2 \tan\left(\frac{1}{2}x\right) + 9b^2 \tan\left(\frac{1}{2}x\right)}{24b^3} - \dots$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^5/(a+b*csc(x)),x, algorithm="giac")

```
[Out] 2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 +
b^2)))*a^4/(sqrt(-a^2 + b^2)*b^4) + 1/24*(b^2*tan(1/2*x)^3 - 3*a*b*tan(1/2*
x)^2 + 12*a^2*tan(1/2*x) + 9*b^2*tan(1/2*x))/b^3 - 1/2*(2*a^3 + a*b^2)*log(
abs(tan(1/2*x)))/b^4 + 1/24*(44*a^3*tan(1/2*x)^3 + 22*a*b^2*tan(1/2*x)^3 -
12*a^2*b*tan(1/2*x)^2 - 9*b^3*tan(1/2*x)^2 + 3*a*b^2*tan(1/2*x) - b^3)/(b^4
*tan(1/2*x)^3)
```

$$3.40 \quad \int \frac{\csc^4(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=84

$$-\frac{(2a^2 + b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{2a^3 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{b^3 \sqrt{a^2-b^2}} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b}$$

[Out] $-\left(\left(2a^2 + b^2\right) \operatorname{ArcTanh}[\cos[x]]\right) / \left(2b^3\right) + \left(2a^3 \operatorname{ArcTanh}\left[\left(a + b \tan\left[x/2\right]\right) / \sqrt{a^2 - b^2}\right]\right) / \left(b^3 \sqrt{a^2 - b^2}\right) + \left(a \cot[x]\right) / b^2 - \left(\cot[x] \csc[x]\right) / \left(2b\right)$

Rubi [A] time = 0.252198, antiderivative size = 84, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 8, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$, Rules used = {3851, 4082, 3998, 3770, 3831, 2660, 618, 206}

$$-\frac{(2a^2 + b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{2a^3 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{b^3 \sqrt{a^2-b^2}} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Csc[x]^4/(a + b*Csc[x]), x]

[Out] $-\left(\left(2a^2 + b^2\right) \operatorname{ArcTanh}[\cos[x]]\right) / \left(2b^3\right) + \left(2a^3 \operatorname{ArcTanh}\left[\left(a + b \tan\left[x/2\right]\right) / \sqrt{a^2 - b^2}\right]\right) / \left(b^3 \sqrt{a^2 - b^2}\right) + \left(a \cot[x]\right) / b^2 - \left(\cot[x] \csc[x]\right) / \left(2b\right)$

Rule 3851

Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n_]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> -Simp[(d^3*Cot[e + f*x]*(d*Csc[e + f*x])^(n - 3))/(b*f*(n - 2)), x] + Dist[d^3/(b*(n - 2)), Int[((d*Csc[e + f*x])^(n - 3)*Simp[a*(n - 3) + b*(n - 3)*Csc[e + f*x] - a*(n - 2)*Csc[e + f*x]^2, x])/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 3]

Rule 4082

Int[csc[(e_.) + (f_.)*(x_)]*(A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_, x_S

```

ymbol] := -Simp[(C*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1))/(b*f*(m + 2))
, x] + Dist[1/(b*(m + 2)), Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*Simp[b*A
*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Csc[e + f*x], x], x] /; Fr
eeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]

```

Rule 3998

```

Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)))/(csc[(
e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Dist[B/b, Int[Csc[e + f*x],
x], x] + Dist[(A*b - a*B)/b, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x]
/; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0]

```

Rule 3770

```

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]

```

Rule 3831

```

Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbo
l] := Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f
}, x] && NeQ[a^2 - b^2, 0]

```

Rule 2660

```

Int[((a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = Fre
eFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*
e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[
a^2 - b^2, 0]

```

Rule 618

```

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]

```

Rule 206

```

Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/
Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])

```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^4(x)}{a+b \csc(x)} dx &= -\frac{\cot(x) \csc(x)}{2b} + \frac{\int \frac{\csc(x)(a+b \csc(x)-2a \csc^2(x))}{a+b \csc(x)} dx}{2b} \\
&= \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} + \frac{\int \frac{\csc(x)(ab+(2a^2+b^2) \csc(x))}{a+b \csc(x)} dx}{2b^2} \\
&= \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} - \frac{a^3 \int \frac{\csc(x)}{a+b \csc(x)} dx}{b^3} + \frac{(2a^2+b^2) \int \csc(x) dx}{2b^3} \\
&= -\frac{(2a^2+b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} - \frac{a^3 \int \frac{1}{1+\frac{a \sin(x)}{b}} dx}{b^4} \\
&= -\frac{(2a^2+b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} - \frac{(2a^3) \text{Subst} \left(\int \frac{1}{1+\frac{2ax}{b}+x^2} dx, x, \tan\left(\frac{x}{2}\right) \right)}{b^4} \\
&= -\frac{(2a^2+b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} + \frac{(4a^3) \text{Subst} \left(\int \frac{1}{-4\left(1-\frac{a^2}{b^2}\right)-x^2} dx, x, \frac{2a}{b} + 2 \tan\left(\frac{x}{2}\right) \right)}{b^4} \\
&= -\frac{(2a^2+b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{2a^3 \tanh^{-1} \left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2-b^2}} \right)}{b^3 \sqrt{a^2-b^2}} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b}
\end{aligned}$$

Mathematica [A] time = 0.486741, size = 144, normalized size = 1.71

$$\frac{-\frac{16a^3 \tan^{-1} \left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}} \right)}{\sqrt{b^2-a^2}} + 8a^2 \log \left(\sin \left(\frac{x}{2} \right) \right) - 8a^2 \log \left(\cos \left(\frac{x}{2} \right) \right) - 4ab \tan \left(\frac{x}{2} \right) + 4ab \cot \left(\frac{x}{2} \right) - b^2 \csc^2 \left(\frac{x}{2} \right) + b^2 \sec^2 \left(\frac{x}{2} \right) + 4b^2}{8b^3}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[x]^4/(a + b*Csc[x]),x]

[Out] ((-16*a^3*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]]/Sqrt[-a^2 + b^2] + 4*a*b*Cot[x/2] - b^2*Csc[x/2]^2 - 8*a^2*Log[Cos[x/2]] - 4*b^2*Log[Cos[x/2]] + 8*a^2*Log[Sin[x/2]] + 4*b^2*Log[Sin[x/2]] + b^2*Sec[x/2]^2 - 4*a*b*Tan[x/2])/(8*b^3)

Maple [A] time = 0.045, size = 112, normalized size = 1.3

$$\frac{1}{8b} \left(\tan\left(\frac{x}{2}\right) \right)^2 - \frac{a}{2b^2} \tan\left(\frac{x}{2}\right) - 2 \frac{a^3}{b^3 \sqrt{-a^2 + b^2}} \arctan\left(\frac{1}{2} \frac{2b \tan(x/2) + 2a}{\sqrt{-a^2 + b^2}}\right) - \frac{1}{8b} \left(\tan\left(\frac{x}{2}\right) \right)^{-2} + \frac{a^2}{b^3} \ln\left(\tan\left(\frac{x}{2}\right)\right) +$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(x)^4/(a+b*csc(x)),x)

[Out] 1/8/b*tan(1/2*x)^2-1/2/b^2*a*tan(1/2*x)-2*a^3/b^3/(-a^2+b^2)^(1/2)*arctan(1/2*(2*b*tan(1/2*x)+2*a)/(-a^2+b^2)^(1/2))-1/8/b/tan(1/2*x)^2+1/b^3*ln(tan(1/2*x))*a^2+1/2/b*ln(tan(1/2*x))+1/2*a/b^2/tan(1/2*x)

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^4/(a+b*csc(x)),x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [B] time = 0.878674, size = 1146, normalized size = 13.64

$$\left[\frac{4(a^3b - ab^3) \cos(x) \sin(x) - 2(a^3 \cos(x)^2 - a^3) \sqrt{a^2 - b^2} \log\left(\frac{(a^2 - 2b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 + 2(b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2ab \sin(x) - a^2 - b^2}\right)}{\right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^4/(a+b*csc(x)),x, algorithm="fricas")

[Out] [1/4*(4*(a^3*b - a*b^3)*cos(x)*sin(x) - 2*(a^3*cos(x)^2 - a^3)*sqrt(a^2 - b^2)*log(((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) - 2*(a^2*b^2 - b^4)*cos(x) - (2*a^4 - a^2*b^2 - b^4 - (2*a^4 - a^2*b^2 - b^4)*cos(x)^2)*log(1/2*cos(x) + 1/2) + (2*a^4 - a^2*b^2 - b^4 - (2*a^4 - a^2*b^2 - b^4)*cos(x)^2)*log(1/2*cos(x) - 1/2) + (2*a^4 - a^2*b^2 - b^4 - (2*a^4 - a^2*b^2 - b^4)*cos(x)^2)*log(1/2*cos(x) + 1/2) + (2*a^4 - a^2*b^2 - b^4 - (2*a^4 - a^2*b^2 - b^4)*cos(x)^2)*log(1/2*cos(x) - 1/2)]

$$2*b^2 - b^4)*\cos(x)^2*\log(-1/2*\cos(x) + 1/2))/(a^2*b^3 - b^5 - (a^2*b^3 - b^5)*\cos(x)^2), 1/4*(4*(a^3*b - a*b^3)*\cos(x)*\sin(x) - 4*(a^3*\cos(x)^2 - a^3)*\sqrt{-a^2 + b^2}*\arctan(-\sqrt{-a^2 + b^2}*(b*\sin(x) + a)/((a^2 - b^2)*\cos(x)))) - 2*(a^2*b^2 - b^4)*\cos(x) - (2*a^4 - a^2*b^2 - b^4 - (2*a^4 - a^2*b^2 - b^4)*\cos(x)^2)*\log(1/2*\cos(x) + 1/2) + (2*a^4 - a^2*b^2 - b^4 - (2*a^4 - a^2*b^2 - b^4)*\cos(x)^2)*\log(-1/2*\cos(x) + 1/2))/(a^2*b^3 - b^5 - (a^2*b^3 - b^5)*\cos(x)^2)]$$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\csc^4(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)**4/(a+b*csc(x)),x)

[Out] Integral(csc(x)**4/(a + b*csc(x)), x)

Giac [A] time = 1.44523, size = 190, normalized size = 2.26

$$\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan \left(\frac{1}{2} x \right) + a}{\sqrt{-a^2 + b^2}} \right) \right) a^3}{\sqrt{-a^2 + b^2} b^3} + \frac{b \tan \left(\frac{1}{2} x \right)^2 - 4 a \tan \left(\frac{1}{2} x \right)}{8 b^2} + \frac{(2 a^2 + b^2) \log \left(\left| \tan \left(\frac{1}{2} x \right) \right| \right)}{2 b^3} - \frac{12 a^2 t}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^4/(a+b*csc(x)),x, algorithm="giac")

[Out] -2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))*a^3/(sqrt(-a^2 + b^2)*b^3) + 1/8*(b*tan(1/2*x)^2 - 4*a*tan(1/2*x))/b^2 + 1/2*(2*a^2 + b^2)*log(abs(tan(1/2*x)))/b^3 - 1/8*(12*a^2*tan(1/2*x)^2 + 6*b^2*tan(1/2*x)^2 - 4*a*b*tan(1/2*x) + b^2)/(b^3*tan(1/2*x)^2)

$$3.41 \quad \int \frac{\csc^3(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=62

$$-\frac{2a^2 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{b^2 \sqrt{a^2-b^2}} + \frac{a \tanh^{-1}(\cos(x))}{b^2} - \frac{\cot(x)}{b}$$

[Out] (a*ArcTanh[Cos[x]])/b^2 - (2*a^2*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(b^2*Sqrt[a^2 - b^2]) - Cot[x]/b

Rubi [A] time = 0.154549, antiderivative size = 62, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 7, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3790, 3789, 3770, 3831, 2660, 618, 206}

$$-\frac{2a^2 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{b^2 \sqrt{a^2-b^2}} + \frac{a \tanh^{-1}(\cos(x))}{b^2} - \frac{\cot(x)}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[x]^3/(a + b*Csc[x]),x]

[Out] (a*ArcTanh[Cos[x]])/b^2 - (2*a^2*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(b^2*Sqrt[a^2 - b^2]) - Cot[x]/b

Rule 3790

Int[csc[(e_.) + (f_.)*(x_)]^3/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> -Simp[Cot[e + f*x]/(b*f), x] - Dist[a/b, Int[Csc[e + f*x]^2/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x]

Rule 3789

Int[csc[(e_.) + (f_.)*(x_)]^2/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Dist[1/b, Int[Csc[e + f*x], x], x] - Dist[a/b, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x]

Rule 3770

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x]
  /; FreeQ[{c, d}, x]
```

Rule 3831

```
Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol]
:= Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f}, x]
  && NeQ[a^2 - b^2, 0]
```

Rule 2660

```
Int[((a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[
Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x,
Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

Rule 618

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[
b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 206

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]]/
(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^3(x)}{a + b \csc(x)} dx &= -\frac{\cot(x)}{b} - \frac{a \int \frac{\csc^2(x)}{a+b \csc(x)} dx}{b} \\
&= -\frac{\cot(x)}{b} - \frac{a \int \csc(x) dx}{b^2} + \frac{a^2 \int \frac{\csc(x)}{a+b \csc(x)} dx}{b^2} \\
&= \frac{a \tanh^{-1}(\cos(x))}{b^2} - \frac{\cot(x)}{b} + \frac{a^2 \int \frac{1}{1+\frac{a \sin(x)}{b}} dx}{b^3} \\
&= \frac{a \tanh^{-1}(\cos(x))}{b^2} - \frac{\cot(x)}{b} + \frac{(2a^2) \text{Subst} \left(\int \frac{1}{1+\frac{2ax}{b}+x^2} dx, x, \tan\left(\frac{x}{2}\right) \right)}{b^3} \\
&= \frac{a \tanh^{-1}(\cos(x))}{b^2} - \frac{\cot(x)}{b} - \frac{(4a^2) \text{Subst} \left(\int \frac{1}{-4\left(1-\frac{a^2}{b^2}\right)-x^2} dx, x, \frac{2a}{b} + 2 \tan\left(\frac{x}{2}\right) \right)}{b^3} \\
&= \frac{a \tanh^{-1}(\cos(x))}{b^2} - \frac{2a^2 \tanh^{-1} \left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2-b^2}} \right)}{b^2 \sqrt{a^2-b^2}} - \frac{\cot(x)}{b}
\end{aligned}$$

Mathematica [A] time = 0.203814, size = 106, normalized size = 1.71

$$\frac{\csc\left(\frac{x}{2}\right) \sec\left(\frac{x}{2}\right) \left(2a^2 \sin(x) \tan^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}}\right) + \sqrt{b^2-a^2} \left(a \sin(x) \left(\log\left(\cos\left(\frac{x}{2}\right)\right) - \log\left(\sin\left(\frac{x}{2}\right)\right) \right) - b \cos(x) \right) \right)}{2b^2 \sqrt{b^2-a^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[x]^3/(a + b*Csc[x]), x]

[Out] (Csc[x/2]*Sec[x/2]*(2*a^2*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]]*Sin[x] + Sqrt[-a^2 + b^2]*(-(b*Cos[x]) + a*(Log[Cos[x/2]] - Log[Sin[x/2]]))*Sin[x]))/(2*b^2*Sqrt[-a^2 + b^2])

Maple [A] time = 0.049, size = 77, normalized size = 1.2

$$\frac{1}{2b} \tan\left(\frac{x}{2}\right) + 2 \frac{a^2}{b^2 \sqrt{-a^2 + b^2}} \arctan\left(\frac{1}{2} \frac{2b \tan\left(\frac{x}{2}\right) + 2a}{\sqrt{-a^2 + b^2}}\right) - \frac{1}{2b} \left(\tan\left(\frac{x}{2}\right)\right)^{-1} - \frac{a}{b^2} \ln\left(\tan\left(\frac{x}{2}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(x)^3/(a+b*csc(x)),x)`

[Out] $\frac{1}{2} \frac{b \tan\left(\frac{1}{2}x\right) + 2a^2/b^2 \sqrt{-a^2+b^2} \arctan\left(\frac{1}{2} \frac{2b \tan\left(\frac{1}{2}x\right) + 2a}{-a^2+b^2}\right) - 1/b \tan\left(\frac{1}{2}x\right) - 1/b^2 a \ln\left(\tan\left(\frac{1}{2}x\right)\right)}{(-a^2+b^2)^{1/2}}$

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(x)^3/(a+b*csc(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError

Fricas [B] time = 0.623509, size = 768, normalized size = 12.39

$$\frac{\sqrt{a^2 - b^2} a^2 \log\left(-\frac{(a^2 - 2b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 - 2(b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2ab \sin(x) - a^2 - b^2}\right) \sin(x) + (a^3 - ab^2) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) \sin(x)}{2(a^2 b^2 - b^4) \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(x)^3/(a+b*csc(x)),x, algorithm="fricas")`

[Out] $\left[\frac{1}{2} \frac{(\sqrt{a^2 - b^2} a^2 \log(-((a^2 - 2b^2) \cos(x)^2 + 2a b \sin(x) + a^2 + b^2 - 2(b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2})) / (a^2 \cos(x)^2 - 2a b \sin(x) - a^2 - b^2)) \sin(x) + (a^3 - a b^2) \log(1/2 \cos(x) + 1/2) \sin(x) - (a^3 - a b^2) \log(-1/2 \cos(x) + 1/2) \sin(x) - 2(a^2 b - b^3) \cos(x)}{(a^2 b^2 - b^4) \sin(x)}, -\frac{1}{2} \frac{2 \sqrt{-a^2 + b^2} a^2 \arctan(-\sqrt{-a^2 + b^2} (b \sin(x) + a) / ((a^2 - b^2) \cos(x))) \sin(x) - (a^3 - a b^2) \log(1/2 \cos(x) + 1/2) \sin(x) + (a^3 - a b^2) \log(-1/2 \cos(x) + 1/2) \sin(x) + 2(a^2 b - b^3) \cos(x)}{(a^2 b^2 - b^4) \sin(x)}\right]$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\csc^3(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)**3/(a+b*csc(x)),x)

[Out] Integral(csc(x)**3/(a + b*csc(x)), x)

Giac [A] time = 1.28532, size = 132, normalized size = 2.13

$$\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}} \right) \right) a^2}{\sqrt{-a^2 + b^2} b^2} - \frac{a \log \left(\left| \tan \left(\frac{1}{2}x \right) \right| \right)}{b^2} + \frac{\tan \left(\frac{1}{2}x \right)}{2b} + \frac{2a \tan \left(\frac{1}{2}x \right) - b}{2b^2 \tan \left(\frac{1}{2}x \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^3/(a+b*csc(x)),x, algorithm="giac")

[Out] 2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))*a^2/(sqrt(-a^2 + b^2)*b^2) - a*log(abs(tan(1/2*x)))/b^2 + 1/2*tan(1/2*x)/b + 1/2*(2*a*tan(1/2*x) - b)/(b^2*tan(1/2*x))

$$3.42 \quad \int \frac{\csc^2(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=53

$$\frac{2a \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{b\sqrt{a^2-b^2}} - \frac{\tanh^{-1}(\cos(x))}{b}$$

[Out] -(ArcTanh[Cos[x]]/b) + (2*a*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(b*Sqrt[a^2 - b^2])

Rubi [A] time = 0.112866, antiderivative size = 53, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 6, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3789, 3770, 3831, 2660, 618, 206}

$$\frac{2a \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{b\sqrt{a^2-b^2}} - \frac{\tanh^{-1}(\cos(x))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[x]^2/(a + b*Csc[x]),x]

[Out] -(ArcTanh[Cos[x]]/b) + (2*a*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(b*Sqrt[a^2 - b^2])

Rule 3789

Int[csc[(e_.) + (f_.)*(x_)]^2/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Dist[1/b, Int[Csc[e + f*x], x], x] - Dist[a/b, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3831

Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f}

}, x] && NeQ[a^2 - b^2, 0]

Rule 2660

Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 618

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 206

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rubi steps

$$\begin{aligned}
 \int \frac{\csc^2(x)}{a + b \csc(x)} dx &= \frac{\int \csc(x) dx}{b} - \frac{a \int \frac{\csc(x)}{a + b \csc(x)} dx}{b} \\
 &= -\frac{\tanh^{-1}(\cos(x))}{b} - \frac{a \int \frac{1}{1 + \frac{a \sin(x)}{b}} dx}{b^2} \\
 &= -\frac{\tanh^{-1}(\cos(x))}{b} - \frac{(2a) \operatorname{Subst} \left(\int \frac{1}{1 + \frac{2ax}{b} + x^2} dx, x, \tan\left(\frac{x}{2}\right) \right)}{b^2} \\
 &= -\frac{\tanh^{-1}(\cos(x))}{b} + \frac{(4a) \operatorname{Subst} \left(\int \frac{1}{-4\left(1 - \frac{a^2}{b^2}\right) - x^2} dx, x, \frac{2a}{b} + 2 \tan\left(\frac{x}{2}\right) \right)}{b^2} \\
 &= -\frac{\tanh^{-1}(\cos(x))}{b} + \frac{2a \tanh^{-1} \left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2 - b^2}} \right)}{b\sqrt{a^2 - b^2}}
 \end{aligned}$$

Mathematica [A] time = 0.0607206, size = 62, normalized size = 1.17

$$\frac{2a \tan^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}}\right)}{\sqrt{b^2-a^2}} + \log\left(\sin\left(\frac{x}{2}\right)\right) - \log\left(\cos\left(\frac{x}{2}\right)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[x]^2/(a + b*Csc[x]),x]

[Out] ((-2*a*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] - Log[Cos[x/2]] + Log[Sin[x/2]])/b

Maple [A] time = 0.045, size = 53, normalized size = 1.

$$-2 \frac{a}{b\sqrt{-a^2 + b^2}} \arctan\left(\frac{1}{2} \frac{2b \tan(x/2) + 2a}{\sqrt{-a^2 + b^2}}\right) + \frac{1}{b} \ln\left(\tan\left(\frac{x}{2}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(x)^2/(a+b*csc(x)),x)

[Out] -2*a/b/(-a^2+b^2)^(1/2)*arctan(1/2*(2*b*tan(1/2*x)+2*a)/(-a^2+b^2)^(1/2))+1/b*ln(tan(1/2*x))

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^2/(a+b*csc(x)),x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [B] time = 0.612126, size = 595, normalized size = 11.23

$$\frac{\sqrt{a^2 - b^2} a \log\left(\frac{(a^2 - 2b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 + 2(b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2ab \sin(x) - a^2 - b^2}\right) - (a^2 - b^2) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + (a^2 - b^2) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)}{2(a^2 b - b^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^2/(a+b*csc(x)),x, algorithm="fricas")

[Out] [1/2*(sqrt(a^2 - b^2)*a*log(((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) - (a^2 - b^2)*log(1/2*cos(x) + 1/2) + (a^2 - b^2)*log(-1/2*cos(x) + 1/2))/(a^2*b - b^3), 1/2*(2*sqrt(-a^2 + b^2)*a*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/((a^2 - b^2)*cos(x))) - (a^2 - b^2)*log(1/2*cos(x) + 1/2) + (a^2 - b^2)*log(-1/2*cos(x) + 1/2))/(a^2*b - b^3)]

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\csc^2(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)**2/(a+b*csc(x)),x)

[Out] Integral(csc(x)**2/(a + b*csc(x)), x)

Giac [A] time = 1.5833, size = 85, normalized size = 1.6

$$-\frac{2\left(\pi\left[\frac{x}{2\pi} + \frac{1}{2}\right] \operatorname{sgn}(b) + \arctan\left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}}\right)\right)a}{\sqrt{-a^2 + b^2}b} + \frac{\log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)^2/(a+b*csc(x)),x, algorithm="giac")

```
[Out] -2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 +  
b^2)))*a/(sqrt(-a^2 + b^2)*b) + log(abs(tan(1/2*x)))/b
```

$$3.43 \quad \int \frac{\csc(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=40

$$-\frac{2 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}}$$

[Out] $(-2*\text{ArcTanh}[(a + b*\text{Tan}[x/2])/Sqrt[a^2 - b^2]])/Sqrt[a^2 - b^2]$

Rubi [A] time = 0.0702075, antiderivative size = 40, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3831, 2660, 618, 206}

$$-\frac{2 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[x]/(a + b*\text{Csc}[x]), x]$

[Out] $(-2*\text{ArcTanh}[(a + b*\text{Tan}[x/2])/Sqrt[a^2 - b^2]])/Sqrt[a^2 - b^2]$

Rule 3831

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_.)]/(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Dist}[1/b, \text{Int}[1/(1 + (a*\text{Sin}[e + f*x])/b), x], x] /;$ $\text{FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Rule 2660

$\text{Int}[(a_.) + (b_.)*\text{sin}[(c_.) + (d_.)*(x_.)]^{(-1)}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Dist}[(2*e)/d, \text{Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Rule 618

$\text{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{(-1)}, x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /;$ $\text{FreeQ}\{a, b, c\},$

`x] && NeQ[b^2 - 4*a*c, 0]`

Rule 206

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rubi steps

$$\begin{aligned} \int \frac{\csc(x)}{a + b \csc(x)} dx &= \frac{\int \frac{1}{1 + \frac{a \sin(x)}{b}} dx}{b} \\ &= \frac{2 \operatorname{Subst} \left(\int \frac{1}{1 + \frac{2ax}{b} + x^2} dx, x, \tan\left(\frac{x}{2}\right) \right)}{b} \\ &= \frac{4 \operatorname{Subst} \left(\int \frac{1}{-4\left(1 - \frac{a^2}{b^2}\right) - x^2} dx, x, \frac{2a}{b} + 2 \tan\left(\frac{x}{2}\right) \right)}{b} \\ &= \frac{2 \tanh^{-1} \left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2 - b^2}} \right)}{\sqrt{a^2 - b^2}} \end{aligned}$$

Mathematica [A] time = 0.0242993, size = 40, normalized size = 1.

$$\frac{2 \tan^{-1} \left(\frac{a + b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2 - a^2}} \right)}{\sqrt{b^2 - a^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[x]/(a + b*Csc[x]), x]`

`[Out] (2*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2]`

Maple [A] time = 0.042, size = 39, normalized size = 1.

$$2 \frac{1}{\sqrt{-a^2 + b^2}} \arctan \left(\frac{1}{2} \frac{2b \tan(x/2) + 2a}{\sqrt{-a^2 + b^2}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(x)/(a+b*csc(x)),x)`

[Out] $2/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*b*\tan(1/2*x)+2*a)/(-a^2+b^2)^{(1/2)})$

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(x)/(a+b*csc(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError

Fricas [A] time = 0.497159, size = 360, normalized size = 9.

$$\left[\frac{\log\left(-\frac{(a^2-2b^2)\cos(x)^2+2ab\sin(x)+a^2+b^2-2(b\cos(x)\sin(x)+a\cos(x))\sqrt{a^2-b^2}}{a^2\cos(x)^2-2ab\sin(x)-a^2-b^2}\right)}{2\sqrt{a^2-b^2}}, -\frac{\sqrt{-a^2+b^2}\arctan\left(-\frac{\sqrt{-a^2+b^2}(b\sin(x)+a)}{(a^2-b^2)\cos(x)}\right)}{a^2-b^2} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(x)/(a+b*csc(x)),x, algorithm="fricas")`

[Out] $[1/2*\log(-((a^2 - 2*b^2)*\cos(x)^2 + 2*a*b*\sin(x) + a^2 + b^2 - 2*(b*\cos(x))*\sin(x) + a*\cos(x))*\sqrt{a^2 - b^2})/(a^2*\cos(x)^2 - 2*a*b*\sin(x) - a^2 - b^2))/\sqrt{a^2 - b^2}, -\sqrt{-a^2 + b^2}*\arctan(-\sqrt{-a^2 + b^2}*(b*\sin(x) + a)/((a^2 - b^2)*\cos(x)))/(a^2 - b^2)]$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\csc(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)/(a+b*csc(x)),x)

[Out] Integral(csc(x)/(a + b*csc(x)), x)

Giac [A] time = 1.43235, size = 65, normalized size = 1.62

$$\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}} \right) \right)}{\sqrt{-a^2 + b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(x)/(a+b*csc(x)),x, algorithm="giac")

[Out] 2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))/sqrt(-a^2 + b^2)

$$3.44 \quad \int \frac{1}{a+b \csc(c+dx)} dx$$

Optimal. Leaf size=57

$$\frac{2b \tanh^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2-b^2}}\right)}{ad\sqrt{a^2-b^2}} + \frac{x}{a}$$

[Out] x/a + (2*b*ArcTanh[(a + b*Tan[(c + d*x)/2])/Sqrt[a^2 - b^2]])/(a*Sqrt[a^2 - b^2]*d)

Rubi [A] time = 0.064512, antiderivative size = 57, normalized size of antiderivative = 1., number of steps used = 4, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3783, 2660, 618, 206}

$$\frac{2b \tanh^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2-b^2}}\right)}{ad\sqrt{a^2-b^2}} + \frac{x}{a}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Csc[c + d*x])^(-1), x]

[Out] x/a + (2*b*ArcTanh[(a + b*Tan[(c + d*x)/2])/Sqrt[a^2 - b^2]])/(a*Sqrt[a^2 - b^2]*d)

Rule 3783

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(-1), x_Symbol] :> Simp[x/a, x] - Dist[1/a, Int[1/(1 + (a*Sin[c + d*x])/b), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 2660

Int[((a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 618

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 206

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[Rt[-b, 2]*x]/Rt[a, 2])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rubi steps

$$\begin{aligned} \int \frac{1}{a + b \csc(c + dx)} dx &= \frac{x}{a} - \frac{\int \frac{1}{1 + \frac{a \sin(c+dx)}{b}} dx}{a} \\ &= \frac{x}{a} - \frac{2 \operatorname{Subst}\left(\int \frac{1}{1 + \frac{2ax}{b} + x^2} dx, x, \tan\left(\frac{1}{2}(c + dx)\right)\right)}{ad} \\ &= \frac{x}{a} + \frac{4 \operatorname{Subst}\left(\int \frac{1}{-4\left(1 - \frac{a^2}{b^2}\right) - x^2} dx, x, \frac{2a}{b} + 2 \tan\left(\frac{1}{2}(c + dx)\right)\right)}{ad} \\ &= \frac{x}{a} + \frac{2b \tanh^{-1}\left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{1}{2}(c+dx)\right)\right)}{\sqrt{a^2 - b^2}}\right)}{a\sqrt{a^2 - b^2}d} \end{aligned}$$

Mathematica [A] time = 0.107481, size = 59, normalized size = 1.04

$$\frac{-\frac{2b \tan^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{b^2-a^2}}\right)}{d\sqrt{b^2-a^2}} + \frac{c}{d} + x}{a}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + b*Csc[c + d*x])^(-1), x]
```

```
[Out] (c/d + x - (2*b*ArcTan[(a + b*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(Sqrt[-a^2 + b^2]*d))/a
```

Maple [A] time = 0.056, size = 70, normalized size = 1.2

$$2 \frac{\arctan(\tan(1/2 dx + c/2))}{ad} - 2 \frac{b}{ad\sqrt{-a^2 + b^2}} \arctan\left(\frac{1}{2} \frac{2b \tan(1/2 dx + c/2) + 2a}{\sqrt{-a^2 + b^2}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+b*csc(d*x+c)),x)

[Out] 2/a/d*arctan(tan(1/2*d*x+1/2*c))-2/d*b/a/(-a^2+b^2)^(1/2)*arctan(1/2*(2*b*tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^(1/2))

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c)),x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [A] time = 0.521268, size = 525, normalized size = 9.21

$$\left[\frac{2(a^2 - b^2)dx + \sqrt{a^2 - b^2}b \log\left(\frac{(a^2 - 2b^2)\cos(dx+c)^2 + 2ab\sin(dx+c) + a^2 + b^2 + 2(b\cos(dx+c)\sin(dx+c) + a\cos(dx+c))\sqrt{a^2 - b^2}}{a^2\cos(dx+c)^2 - 2ab\sin(dx+c) - a^2 - b^2}\right)}{2(a^3 - ab^2)d}, \frac{(a^2 - b^2)dx + \dots}{\dots} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c)),x, algorithm="fricas")

[Out] [1/2*(2*(a^2 - b^2)*d*x + sqrt(a^2 - b^2)*b*log(((a^2 - 2*b^2)*cos(d*x + c)^2 + 2*a*b*sin(d*x + c) + a^2 + b^2 + 2*(b*cos(d*x + c)*sin(d*x + c) + a*cos(d*x + c))*sqrt(a^2 - b^2)))/(a^2*cos(d*x + c)^2 - 2*a*b*sin(d*x + c) - a^2 - b^2)))/((a^3 - a*b^2)*d), ((a^2 - b^2)*d*x + sqrt(-a^2 + b^2)*b*arctan(-

```
sqrt(-a^2 + b^2)*(b*sin(d*x + c) + a)/((a^2 - b^2)*cos(d*x + c)))/((a^3 -
a*b^2)*d)]
```

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{a + b \csc(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a+b*csc(d*x+c)),x)
```

```
[Out] Integral(1/(a + b*csc(c + d*x)), x)
```

Giac [A] time = 1.45478, size = 104, normalized size = 1.82

$$\frac{2 \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(b) + \arctan \left(\frac{b \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + a}{\sqrt{-a^2 + b^2}} \right) \right) b}{\sqrt{-a^2 + b^2} a} - \frac{dx+c}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a+b*csc(d*x+c)),x, algorithm="giac")
```

```
[Out] -(2*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*d*x + 1/2*
c) + a)/sqrt(-a^2 + b^2)))*b/(sqrt(-a^2 + b^2)*a) - (d*x + c)/a)/d
```

$$3.45 \quad \int \frac{\sin(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=61

$$-\frac{2b^2 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a^2 \sqrt{a^2-b^2}} - \frac{bx}{a^2} - \frac{\cos(x)}{a}$$

[Out] -((b*x)/a^2) - (2*b^2*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a^2*Sqrt[a^2 - b^2]) - Cos[x]/a

Rubi [A] time = 0.106608, antiderivative size = 61, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 6, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.546$, Rules used = {3853, 12, 3783, 2660, 618, 206}

$$-\frac{2b^2 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a^2 \sqrt{a^2-b^2}} - \frac{bx}{a^2} - \frac{\cos(x)}{a}$$

Antiderivative was successfully verified.

[In] Int[Sin[x]/(a + b*Csc[x]),x]

[Out] -((b*x)/a^2) - (2*b^2*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a^2*Sqrt[a^2 - b^2]) - Cos[x]/a

Rule 3853

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_)/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Simp[(Cot[e + f*x]*(d*Csc[e + f*x])^n)/(a*f*n), x] - Dist[1/(a*d*n), Int[((d*Csc[e + f*x])^(n + 1)*Simp[b*n - a*(n + 1)*Csc[e + f*x] - b*(n + 1)*Csc[e + f*x]^2, x])/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 12

Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 3783

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))(-1), x_Symbol] := Simp[x/a, x]
- Dist[1/a, Int[1/(1 + (a*Sin[c + d*x])/b), x], x] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a2 - b2, 0]
```

Rule 2660

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e2*x2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a2 - b2, 0]
```

Rule 618

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)2)(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b2 - 4*a*c - x2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b2 - 4*a*c, 0]
```

Rule 206

```
Int[((a_) + (b_.)*(x_)2)(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{\sin(x)}{a + b \csc(x)} dx &= -\frac{\cos(x)}{a} - \frac{\int \frac{b}{a+b \csc(x)} dx}{a} \\
&= -\frac{\cos(x)}{a} - \frac{b \int \frac{1}{a+b \csc(x)} dx}{a} \\
&= -\frac{bx}{a^2} - \frac{\cos(x)}{a} + \frac{b \int \frac{1}{1+\frac{a \sin(x)}{b}} dx}{a^2} \\
&= -\frac{bx}{a^2} - \frac{\cos(x)}{a} + \frac{(2b) \text{Subst} \left(\int \frac{1}{1+\frac{2ax}{b}+x^2} dx, x, \tan\left(\frac{x}{2}\right) \right)}{a^2} \\
&= -\frac{bx}{a^2} - \frac{\cos(x)}{a} - \frac{(4b) \text{Subst} \left(\int \frac{1}{-4\left(1-\frac{a^2}{b^2}\right)-x^2} dx, x, \frac{2a}{b} + 2 \tan\left(\frac{x}{2}\right) \right)}{a^2} \\
&= -\frac{bx}{a^2} - \frac{2b^2 \tanh^{-1} \left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2-b^2}} \right)}{a^2 \sqrt{a^2-b^2}} - \frac{\cos(x)}{a}
\end{aligned}$$

Mathematica [A] time = 0.0985496, size = 56, normalized size = 0.92

$$-\frac{2b^2 \tan^{-1} \left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}} \right)}{\sqrt{b^2-a^2}} + a \cos(x) + bx$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]/(a + b*Csc[x]), x]

[Out] -((b*x - (2*b^2*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + a*Cos[x])/a^2)

Maple [A] time = 0.061, size = 72, normalized size = 1.2

$$-2 \frac{1}{a \left((\tan(x/2))^2 + 1 \right)} - 2 \frac{b \arctan(\tan(x/2))}{a^2} + 2 \frac{b^2}{a^2 \sqrt{-a^2 + b^2}} \arctan \left(\frac{1}{2} \frac{2b \tan(x/2) + 2a}{\sqrt{-a^2 + b^2}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(x)/(a+b*csc(x)),x)
```

```
[Out] -2/a/(tan(1/2*x)^2+1)-2/a^2*b*arctan(tan(1/2*x))+2*b^2/a^2/(-a^2+b^2)^(1/2)
*arctan(1/2*(2*b*tan(1/2*x)+2*a)/(-a^2+b^2)^(1/2))
```

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(x)/(a+b*csc(x)),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError
```

Fricas [A] time = 0.531966, size = 520, normalized size = 8.52

$$\frac{\sqrt{a^2 - b^2} b^2 \log\left(-\frac{(a^2 - 2b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 - 2(b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2ab \sin(x) - a^2 - b^2}\right) - 2(a^2 b - b^3)x - 2(a^3 - ab^2) \cos(x) \sqrt{-a^2 - b^2}}{2(a^4 - a^2 b^2)}, -$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(x)/(a+b*csc(x)),x, algorithm="fricas")
```

```
[Out] [1/2*(sqrt(a^2 - b^2)*b^2*log(-((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2
+ b^2 - 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*
a*b*sin(x) - a^2 - b^2)) - 2*(a^2*b - b^3)*x - 2*(a^3 - a*b^2)*cos(x))/(a^4
- a^2*b^2), -(sqrt(-a^2 + b^2)*b^2*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)
/((a^2 - b^2)*cos(x))) + (a^2*b - b^3)*x + (a^3 - a*b^2)*cos(x))/(a^4 - a^2
*b^2)]
```

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sin(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)/(a+b*csc(x)),x)

[Out] Integral(sin(x)/(a + b*csc(x)), x)

Giac [A] time = 1.45592, size = 104, normalized size = 1.7

$$\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}} \right) \right) b^2}{\sqrt{-a^2 + b^2} a^2} - \frac{bx}{a^2} - \frac{2}{\left(\tan\left(\frac{1}{2}x\right)^2 + 1 \right) a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)/(a+b*csc(x)),x, algorithm="giac")

[Out] 2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))*b^2/(sqrt(-a^2 + b^2)*a^2) - b*x/a^2 - 2/((tan(1/2*x)^2 + 1)*a)

3.46 $\int \frac{\sin^2(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=82

$$\frac{x(a^2 + 2b^2)}{2a^3} + \frac{2b^3 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a^3 \sqrt{a^2-b^2}} + \frac{b \cos(x)}{a^2} - \frac{\sin(x) \cos(x)}{2a}$$

[Out] ((a^2 + 2*b^2)*x)/(2*a^3) + (2*b^3*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a^3*Sqrt[a^2 - b^2]) + (b*Cos[x])/a^2 - (Cos[x]*Sin[x])/(2*a)

Rubi [A] time = 0.261108, antiderivative size = 82, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 7, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3853, 4104, 3919, 3831, 2660, 618, 206}

$$\frac{x(a^2 + 2b^2)}{2a^3} + \frac{2b^3 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a^3 \sqrt{a^2-b^2}} + \frac{b \cos(x)}{a^2} - \frac{\sin(x) \cos(x)}{2a}$$

Antiderivative was successfully verified.

[In] Int[Sin[x]^2/(a + b*Csc[x]),x]

[Out] ((a^2 + 2*b^2)*x)/(2*a^3) + (2*b^3*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a^3*Sqrt[a^2 - b^2]) + (b*Cos[x])/a^2 - (Cos[x]*Sin[x])/(2*a)

Rule 3853

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n_]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Simp[(Cot[e + f*x]*(d*Csc[e + f*x])^n)/(a*f*n), x] - Dist[1/(a*d*n), Int[((d*Csc[e + f*x])^(n + 1)*Simp[b*n - a*(n + 1)*Csc[e + f*x] - b*(n + 1)*Csc[e + f*x]^2, x])/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 4104

Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n_]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^m_), x_Symbol] :> Simp[(A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n)/(a*f*n), x] + Dist[1/(a*d*n), Int[(a + b*Csc[e + f*x])^m*(d*Csc[e + f*x])^(n + 1)*Simp[a*B*n - A*b*(m + n + 1) + a*(A + A*n + C*n)*C

$\text{sc}[e + f*x] + A*b*(m + n + 2)*\text{Csc}[e + f*x]^2, x], x] /; \text{FreeQ}\{a, b, d, e, f, A, B, C, m\}, x\} \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LeQ}[n, -1]$

Rule 3919

$\text{Int}[(\text{csc}[e_] + (f_)*(x_)]*(d_) + (c_))/(\text{csc}[e_] + (f_)*(x_)]*(b_) + (a_)), x_Symbol] \rightarrow \text{Simp}[(c*x)/a, x] - \text{Dist}[(b*c - a*d)/a, \text{Int}[\text{Csc}[e + f*x]/(a + b*\text{Csc}[e + f*x]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x\} \&\& \text{NeQ}[b*c - a*d, 0]$

Rule 3831

$\text{Int}[\text{csc}[e_] + (f_)*(x_)]/(\text{csc}[e_] + (f_)*(x_)]*(b_) + (a_)), x_Symbol] \rightarrow \text{Dist}[1/b, \text{Int}[1/(1 + (a*\text{Sin}[e + f*x])/b), x], x] /; \text{FreeQ}\{a, b, e, f\}, x\} \&\& \text{NeQ}[a^2 - b^2, 0]$

Rule 2660

$\text{Int}[(a_) + (b_)*\text{sin}[c_] + (d_)*(x_)]^{-1}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Dist}[(2*e)/d, \text{Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x] /; \text{FreeQ}\{a, b, c, d\}, x\} \&\& \text{NeQ}[a^2 - b^2, 0]$

Rule 618

$\text{Int}[(a_) + (b_)*(x_) + (c_)*(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\}, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 206

$\text{Int}[(a_) + (b_)*(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(1*\text{ArcTanh}[(\text{Rt}[-b, 2]*x)/\text{Rt}[a, 2]])/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]), x] /; \text{FreeQ}\{a, b\}, x\} \&\& \text{NegQ}[a/b] \&\& (\text{GtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

Rubi steps

$$\begin{aligned}
\int \frac{\sin^2(x)}{a + b \csc(x)} dx &= -\frac{\cos(x) \sin(x)}{2a} + \frac{\int \frac{(-2b+a \csc(x)+b \csc^2(x)) \sin(x)}{a+b \csc(x)} dx}{2a} \\
&= \frac{b \cos(x)}{a^2} - \frac{\cos(x) \sin(x)}{2a} - \frac{\int \frac{-a^2-2b^2-ab \csc(x)}{a+b \csc(x)} dx}{2a^2} \\
&= \frac{(a^2 + 2b^2)x}{2a^3} + \frac{b \cos(x)}{a^2} - \frac{\cos(x) \sin(x)}{2a} - \frac{b^3 \int \frac{\csc(x)}{a+b \csc(x)} dx}{a^3} \\
&= \frac{(a^2 + 2b^2)x}{2a^3} + \frac{b \cos(x)}{a^2} - \frac{\cos(x) \sin(x)}{2a} - \frac{b^2 \int \frac{1}{1+\frac{a \sin(x)}{b}} dx}{a^3} \\
&= \frac{(a^2 + 2b^2)x}{2a^3} + \frac{b \cos(x)}{a^2} - \frac{\cos(x) \sin(x)}{2a} - \frac{(2b^2) \text{Subst} \left(\int \frac{1}{1+\frac{2ax}{b}+x^2} dx, x, \tan\left(\frac{x}{2}\right) \right)}{a^3} \\
&= \frac{(a^2 + 2b^2)x}{2a^3} + \frac{b \cos(x)}{a^2} - \frac{\cos(x) \sin(x)}{2a} + \frac{(4b^2) \text{Subst} \left(\int \frac{1}{-4\left(1-\frac{a^2}{b^2}\right)-x^2} dx, x, \frac{2a}{b} + 2 \tan\left(\frac{x}{2}\right) \right)}{a^3} \\
&= \frac{(a^2 + 2b^2)x}{2a^3} + \frac{2b^3 \tanh^{-1} \left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2-b^2}} \right)}{a^3 \sqrt{a^2-b^2}} + \frac{b \cos(x)}{a^2} - \frac{\cos(x) \sin(x)}{2a}
\end{aligned}$$

Mathematica [A] time = 0.118575, size = 78, normalized size = 0.95

$$\frac{8b^3 \tan^{-1} \left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}} \right)}{\sqrt{b^2-a^2}} + \frac{2a^2x - a^2 \sin(2x) + 4ab \cos(x) + 4b^2x}{4a^3}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]^2/(a + b*Csc[x]),x]

[Out] (2*a^2*x + 4*b^2*x - (8*b^3*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + 4*a*b*Cos[x] - a^2*Sin[2*x])/(4*a^3)

Maple [A] time = 0.072, size = 142, normalized size = 1.7

$$\frac{1}{a} \left(\tan\left(\frac{x}{2}\right) \right)^3 \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-2} + 2 \frac{b \left(\tan\left(\frac{x}{2}\right) \right)^2}{a^2 \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^2} - \frac{1}{a} \tan\left(\frac{x}{2}\right) \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-2} + 2 \frac{b}{a^2 \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^2} +$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(x)^2/(a+b*csc(x)),x)`

[Out] $\frac{1}{a} \frac{1}{(\tan(1/2*x)^2+1)^2} \tan(1/2*x)^3 + \frac{2}{a^2} \frac{1}{(\tan(1/2*x)^2+1)^2} b \tan(1/2*x)^2 - \frac{1}{a} \frac{1}{(\tan(1/2*x)^2+1)^2} \tan(1/2*x) + \frac{2}{a^2} \frac{1}{(\tan(1/2*x)^2+1)^2} b + \frac{2}{a^3} \arctan(\tan(1/2*x)) * b^2 - 2 * b^3 / a^3 / (-a^2+b^2)^{(1/2)} * \arctan(1/2*(2*b*\tan(1/2*x)+2*a) / (-a^2+b^2)^{(1/2)}) + 1/2*x/a$

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^2/(a+b*csc(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError

Fricas [A] time = 0.540342, size = 644, normalized size = 7.85

$$\left[\frac{\sqrt{a^2 - b^2} b^3 \log\left(\frac{(a^2 - 2b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 + 2(b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2ab \sin(x) - a^2 - b^2}\right) - (a^4 - a^2 b^2) \cos(x) \sin(x) + (a^4 + a^2 b^2 - 2b^4) x + 2(a^3 b - a b^3) \cos(x)}{2(a^5 - a^3 b^2)}, \frac{1}{2} (2 \sqrt{-a^2 + b^2}) b^3 \arctan\left(\frac{-\sqrt{-a^2 + b^2} (b \sin(x) + a)}{(a^2 - b^2) \cos(x)}\right) - (a^4 - a^2 b^2) \cos(x) \sin(x) + (a^4 + a^2 b^2 - 2b^4) x + 2(a^3 b - a b^3) \cos(x)}{(a^5 - a^3 b^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^2/(a+b*csc(x)),x, algorithm="fricas")`

[Out] $\left[\frac{1}{2} (\sqrt{a^2 - b^2}) b^3 \log\left(\frac{(a^2 - 2b^2) \cos(x)^2 + 2a*b*\sin(x) + a^2 + b^2 + 2*(b*\cos(x)*\sin(x) + a*\cos(x))*\sqrt{a^2 - b^2}}{(a^2*\cos(x))^2 - 2*a*b*\sin(x) - a^2 - b^2}\right) - (a^4 - a^2*b^2)*\cos(x)*\sin(x) + (a^4 + a^2*b^2 - 2*b^4)*x + 2*(a^3*b - a*b^3)*\cos(x)}{(a^5 - a^3*b^2)}, \frac{1}{2} (2*\sqrt{-a^2 + b^2}) * b^3 * \arctan\left(\frac{-\sqrt{-a^2 + b^2} * (b*\sin(x) + a)}{(a^2 - b^2)*\cos(x)}\right) - (a^4 - a^2*b^2)*\cos(x)*\sin(x) + (a^4 + a^2*b^2 - 2*b^4)*x + 2*(a^3*b - a*b^3)*\cos(x)}{(a^5 - a^3*b^2)} \right]$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sin^2(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)**2/(a+b*csc(x)),x)

[Out] Integral(sin(x)**2/(a + b*csc(x)), x)

Giac [A] time = 1.38052, size = 151, normalized size = 1.84

$$\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}} \right) \right) b^3}{\sqrt{-a^2 + b^2} a^3} + \frac{(a^2 + 2b^2)x}{2a^3} + \frac{a \tan\left(\frac{1}{2}x\right)^3 + 2b \tan\left(\frac{1}{2}x\right)^2 - a \tan\left(\frac{1}{2}x\right) + 2b}{\left(\tan\left(\frac{1}{2}x\right)^2 + 1\right)^2 a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^2/(a+b*csc(x)),x, algorithm="giac")

[Out] -2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))*b^3/(sqrt(-a^2 + b^2)*a^3) + 1/2*(a^2 + 2*b^2)*x/a^3 + (a*tan(1/2*x)^3 + 2*b*tan(1/2*x)^2 - a*tan(1/2*x) + 2*b)/((tan(1/2*x)^2 + 1)^2*a^2)

$$3.47 \quad \int \frac{\sin^3(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=110

$$-\frac{bx(a^2+2b^2)}{2a^4} - \frac{(2a^2+3b^2)\cos(x)}{3a^3} - \frac{2b^4 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a^4\sqrt{a^2-b^2}} + \frac{b \sin(x) \cos(x)}{2a^2} - \frac{\sin^2(x) \cos(x)}{3a}$$

[Out] $-(b*(a^2 + 2*b^2)*x)/(2*a^4) - (2*b^4*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a^4*Sqrt[a^2 - b^2]) - ((2*a^2 + 3*b^2)*Cos[x])/(3*a^3) + (b*Cos[x]*Sin[x])/(2*a^2) - (Cos[x]*Sin[x]^2)/(3*a)$

Rubi [A] time = 0.398245, antiderivative size = 110, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 7, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3853, 4104, 3919, 3831, 2660, 618, 206}

$$-\frac{bx(a^2+2b^2)}{2a^4} - \frac{(2a^2+3b^2)\cos(x)}{3a^3} - \frac{2b^4 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a^4\sqrt{a^2-b^2}} + \frac{b \sin(x) \cos(x)}{2a^2} - \frac{\sin^2(x) \cos(x)}{3a}$$

Antiderivative was successfully verified.

[In] Int[Sin[x]^3/(a + b*Csc[x]),x]

[Out] $-(b*(a^2 + 2*b^2)*x)/(2*a^4) - (2*b^4*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a^4*Sqrt[a^2 - b^2]) - ((2*a^2 + 3*b^2)*Cos[x])/(3*a^3) + (b*Cos[x]*Sin[x])/(2*a^2) - (Cos[x]*Sin[x]^2)/(3*a)$

Rule 3853

Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n_]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Simp[(Cot[e + f*x]*(d*Csc[e + f*x])^n)/(a*f*n), x] - Dist[1/(a*d*n), Int[((d*Csc[e + f*x])^(n + 1)*Simp[b*n - a*(n + 1)*Csc[e + f*x] - b*(n + 1)*Csc[e + f*x]^2, x])/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 4104

Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^n_*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_], x_Symbol] :> Simp[(A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(d

```
*Csc[e + f*x])^n)/(a*f*n), x] + Dist[1/(a*d*n), Int[(a + b*Csc[e + f*x])^m*
(d*Csc[e + f*x])^(n + 1)*Simp[a*B*n - A*b*(m + n + 1) + a*(A + A*n + C*n)*C
sc[e + f*x] + A*b*(m + n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d,
e, f, A, B, C, m}, x] && NeQ[a^2 - b^2, 0] && LeQ[n, -1]
```

Rule 3919

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))/(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.), x_Symbol] := Simp[(c*x)/a, x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]
]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c -
a*d, 0]
```

Rule 3831

```
Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.), x_Symbol]
:= Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f
}, x] && NeQ[a^2 - b^2, 0]
```

Rule 2660

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = Fre
eFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*
e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[
a^2 - b^2, 0]
```

Rule 618

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 206

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/
Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{\sin^3(x)}{a + b \csc(x)} dx &= -\frac{\cos(x) \sin^2(x)}{3a} + \frac{\int \frac{(-3b+2a \csc(x)+2b \csc^2(x)) \sin^2(x)}{a+b \csc(x)} dx}{3a} \\
&= \frac{b \cos(x) \sin(x)}{2a^2} - \frac{\cos(x) \sin^2(x)}{3a} - \frac{\int \frac{(-2(2a^2+3b^2)-ab \csc(x)+3b^2 \csc^2(x)) \sin(x)}{a+b \csc(x)} dx}{6a^2} \\
&= -\frac{(2a^2 + 3b^2) \cos(x)}{3a^3} + \frac{b \cos(x) \sin(x)}{2a^2} - \frac{\cos(x) \sin^2(x)}{3a} + \frac{\int \frac{-3b(a^2+2b^2)-3ab^2 \csc(x)}{a+b \csc(x)} dx}{6a^3} \\
&= -\frac{b(a^2 + 2b^2)x}{2a^4} - \frac{(2a^2 + 3b^2) \cos(x)}{3a^3} + \frac{b \cos(x) \sin(x)}{2a^2} - \frac{\cos(x) \sin^2(x)}{3a} + \frac{b^4 \int \frac{\csc(x)}{a+b \csc(x)} dx}{a^4} \\
&= -\frac{b(a^2 + 2b^2)x}{2a^4} - \frac{(2a^2 + 3b^2) \cos(x)}{3a^3} + \frac{b \cos(x) \sin(x)}{2a^2} - \frac{\cos(x) \sin^2(x)}{3a} + \frac{b^3 \int \frac{1}{1+\frac{a \sin(x)}{b}} dx}{a^4} \\
&= -\frac{b(a^2 + 2b^2)x}{2a^4} - \frac{(2a^2 + 3b^2) \cos(x)}{3a^3} + \frac{b \cos(x) \sin(x)}{2a^2} - \frac{\cos(x) \sin^2(x)}{3a} + \frac{(2b^3) \text{Subst} \left(\int \frac{1}{1+\frac{2ax}{b}+x} dx \right)}{a^4} \\
&= -\frac{b(a^2 + 2b^2)x}{2a^4} - \frac{(2a^2 + 3b^2) \cos(x)}{3a^3} + \frac{b \cos(x) \sin(x)}{2a^2} - \frac{\cos(x) \sin^2(x)}{3a} - \frac{(4b^3) \text{Subst} \left(\int \frac{1}{-4\left(1-\frac{a^2}{b^2}\right)} dx \right)}{a^4} \\
&= -\frac{b(a^2 + 2b^2)x}{2a^4} - \frac{2b^4 \tanh^{-1} \left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2-b^2}} \right)}{a^4 \sqrt{a^2-b^2}} - \frac{(2a^2 + 3b^2) \cos(x)}{3a^3} + \frac{b \cos(x) \sin(x)}{2a^2} - \frac{\cos(x) \sin^2(x)}{3a}
\end{aligned}$$

Mathematica [A] time = 0.237856, size = 98, normalized size = 0.89

$$\frac{-6bx(a^2 + 2b^2) - 3a(3a^2 + 4b^2) \cos(x) + \frac{24b^4 \tan^{-1} \left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}} \right)}{\sqrt{b^2-a^2}} + 3a^2b \sin(2x) + a^3 \cos(3x)}{12a^4}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]^3/(a + b*Csc[x]), x]

[Out] (-6*b*(a^2 + 2*b^2)*x + (24*b^4*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] - 3*a*(3*a^2 + 4*b^2)*Cos[x] + a^3*Cos[3*x] + 3*a^2*b*Sin[2*x])/(12*a^4)

Maple [B] time = 0.066, size = 213, normalized size = 1.9

$$-\frac{b}{a^2} \left(\tan\left(\frac{x}{2}\right) \right)^5 \left(\left(\tan\left(\frac{x}{2}\right) \right)^2 + 1 \right)^{-3} - 2 \frac{b^2 (\tan(x/2))^4}{a^3 \left((\tan(x/2))^2 + 1 \right)^3} - 4 \frac{(\tan(x/2))^2}{a \left((\tan(x/2))^2 + 1 \right)^3} - 4 \frac{(\tan(x/2))^2 b^2}{a^3 \left((\tan(x/2))^2 + 1 \right)^3} + \frac{b}{a^2} \tan$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)^3/(a+b*csc(x)),x)

[Out]
$$-1/a^2/(\tan(1/2*x)^2+1)^3*b*\tan(1/2*x)^5-2/a^3/(\tan(1/2*x)^2+1)^3*b^2*\tan(1/2*x)^4-4/a/(\tan(1/2*x)^2+1)^3*\tan(1/2*x)^2-4/a^3/(\tan(1/2*x)^2+1)^3*\tan(1/2*x)^2*b^2+1/a^2/(\tan(1/2*x)^2+1)^3*\tan(1/2*x)*b-4/3/a/(\tan(1/2*x)^2+1)^3-2/a^3/(\tan(1/2*x)^2+1)^3*b^2-1/a^2*b*\arctan(\tan(1/2*x))-2/a^4*\arctan(\tan(1/2*x))*b^3+2*b^4/a^4/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*b*\tan(1/2*x)+2*a)/(-a^2+b^2)^{(1/2)})$$

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^3/(a+b*csc(x)),x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [A] time = 0.552073, size = 744, normalized size = 6.76

$$\left[\frac{3 \sqrt{a^2 - b^2} b^4 \log\left(-\frac{(a^2 - 2b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 - 2(b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2ab \sin(x) - a^2 - b^2} \right) + 2(a^5 - a^3 b^2) \cos(x)^3 + 3(a^4 b - a^2 b^3) \cos(x)}{6(a^6 - a^4 b^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^3/(a+b*csc(x)),x, algorithm="fricas")

[Out]
$$[1/6*(3*\sqrt{a^2 - b^2}*b^4*\log(-((a^2 - 2*b^2)*\cos(x)^2 + 2*a*b*\sin(x) + a^2 + b^2 - 2*(b*\cos(x)*\sin(x) + a*\cos(x))*\sqrt{a^2 - b^2}))/a^2*\cos(x)^2 -$$

$$2*a*b*\sin(x) - a^2 - b^2)) + 2*(a^5 - a^3*b^2)*\cos(x)^3 + 3*(a^4*b - a^2*b^3)*\cos(x)*\sin(x) - 3*(a^4*b + a^2*b^3 - 2*b^5)*x - 6*(a^5 - a*b^4)*\cos(x))/(a^6 - a^4*b^2), -1/6*(6*\sqrt{-a^2 + b^2}*b^4*\arctan(-\sqrt{-a^2 + b^2}*(b*\sin(x) + a)/((a^2 - b^2)*\cos(x)))) - 2*(a^5 - a^3*b^2)*\cos(x)^3 - 3*(a^4*b - a^2*b^3)*\cos(x)*\sin(x) + 3*(a^4*b + a^2*b^3 - 2*b^5)*x + 6*(a^5 - a*b^4)*\cos(x))/(a^6 - a^4*b^2)]$$

Sympy [F(-1)] time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)**3/(a+b*csc(x)),x)

[Out] Timed out

Giac [A] time = 1.4371, size = 201, normalized size = 1.83

$$\frac{2 \left(\pi \left[\frac{x}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(b) + \arctan \left(\frac{b \tan \left(\frac{1}{2} x \right) + a}{\sqrt{-a^2 + b^2}} \right) \right) b^4}{\sqrt{-a^2 + b^2} a^4} - \frac{(a^2 b + 2 b^3) x}{2 a^4} - \frac{3 a b \tan \left(\frac{1}{2} x \right)^5 + 6 b^2 \tan \left(\frac{1}{2} x \right)^4 + 12 a^2 \tan \left(\frac{1}{2} x \right)^2}{3 \left(\tan \left(\frac{1}{2} x \right) \right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^3/(a+b*csc(x)),x, algorithm="giac")

[Out] 2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))*b^4/(sqrt(-a^2 + b^2)*a^4) - 1/2*(a^2*b + 2*b^3)*x/a^4 - 1/3*(3*a*b*tan(1/2*x)^5 + 6*b^2*tan(1/2*x)^4 + 12*a^2*tan(1/2*x)^2 + 12*b^2*tan(1/2*x)^2 - 3*a*b*tan(1/2*x) + 4*a^2 + 6*b^2)/((tan(1/2*x)^2 + 1)^3*a^3)

$$3.48 \quad \int \frac{\sin^4(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=144

$$\frac{x(4a^2b^2 + 3a^4 + 8b^4)}{8a^5} + \frac{b(2a^2 + 3b^2)\cos(x)}{3a^4} - \frac{(3a^2 + 4b^2)\sin(x)\cos(x)}{8a^3} + \frac{2b^5 \tanh^{-1}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{a^5\sqrt{a^2-b^2}} + \frac{b \sin^2(x)\cos(x)}{3a^2}$$

[Out] $((3*a^4 + 4*a^2*b^2 + 8*b^4)*x)/(8*a^5) + (2*b^5*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a^5*Sqrt[a^2 - b^2]) + (b*(2*a^2 + 3*b^2)*Cos[x])/(3*a^4) - ((3*a^2 + 4*b^2)*Cos[x]*Sin[x])/(8*a^3) + (b*Cos[x]*Sin[x]^2)/(3*a^2) - (Cos[x]*Sin[x]^3)/(4*a)$

Rubi [A] time = 0.588463, antiderivative size = 144, normalized size of antiderivative = 1., number of steps used = 9, number of rules used = 7, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3853, 4104, 3919, 3831, 2660, 618, 206}

$$\frac{x(4a^2b^2 + 3a^4 + 8b^4)}{8a^5} + \frac{b(2a^2 + 3b^2)\cos(x)}{3a^4} - \frac{(3a^2 + 4b^2)\sin(x)\cos(x)}{8a^3} + \frac{2b^5 \tanh^{-1}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{a^5\sqrt{a^2-b^2}} + \frac{b \sin^2(x)\cos(x)}{3a^2}$$

Antiderivative was successfully verified.

[In] Int[Sin[x]^4/(a + b*Csc[x]),x]

[Out] $((3*a^4 + 4*a^2*b^2 + 8*b^4)*x)/(8*a^5) + (2*b^5*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a^5*Sqrt[a^2 - b^2]) + (b*(2*a^2 + 3*b^2)*Cos[x])/(3*a^4) - ((3*a^2 + 4*b^2)*Cos[x]*Sin[x])/(8*a^3) + (b*Cos[x]*Sin[x]^2)/(3*a^2) - (Cos[x]*Sin[x]^3)/(4*a)$

Rule 3853

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[(Cot[e + f*x]*(d*Csc[e + f*x])^n)/(a*f*n), x] - Dist[1/(a*d*n), Int[((d*Csc[e + f*x])^(n + 1)*Simp[b*n - a*(n + 1)*Csc[e + f*x] - b*(n + 1)*Csc[e + f*x]^2, x])/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && LeQ[n, -1] && IntegerQ[2*n]

Rule 4104

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_)])*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))^(m_), x_Symbol] := Simp[(A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(d
*Csc[e + f*x])^n)/(a*f*n), x] + Dist[1/(a*d*n), Int[(a + b*Csc[e + f*x])^m*
(d*Csc[e + f*x])^(n + 1)*Simp[a*B*n - A*b*(m + n + 1) + a*(A + A*n + C*n)*C
sc[e + f*x] + A*b*(m + n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d,
e, f, A, B, C, m}, x] && NeQ[a^2 - b^2, 0] && LeQ[n, -1]
```

Rule 3919

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))/(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_)), x_Symbol] := Simp[(c*x)/a, x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x
]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c -
a*d, 0]
```

Rule 3831

```
Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbo
l] := Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f
}, x] && NeQ[a^2 - b^2, 0]
```

Rule 2660

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = Fre
eFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*
e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[
a^2 - b^2, 0]
```

Rule 618

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 206

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/
Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{\sin^4(x)}{a + b \csc(x)} dx &= -\frac{\cos(x) \sin^3(x)}{4a} + \frac{\int \frac{(-4b+3a \csc(x)+3b \csc^2(x)) \sin^3(x)}{a+b \csc(x)} dx}{4a} \\
&= \frac{b \cos(x) \sin^2(x)}{3a^2} - \frac{\cos(x) \sin^3(x)}{4a} - \frac{\int \frac{(-3(3a^2+4b^2)-ab \csc(x)+8b^2 \csc^2(x)) \sin^2(x)}{a+b \csc(x)} dx}{12a^2} \\
&= -\frac{(3a^2 + 4b^2) \cos(x) \sin(x)}{8a^3} + \frac{b \cos(x) \sin^2(x)}{3a^2} - \frac{\cos(x) \sin^3(x)}{4a} + \frac{\int \frac{(-8b(2a^2+3b^2)+a(9a^2-4b^2) \csc(x)+3b(3a^2+4b^2)) \sin(x)}{a+b \csc(x)} dx}{24a^3} \\
&= \frac{b(2a^2 + 3b^2) \cos(x)}{3a^4} - \frac{(3a^2 + 4b^2) \cos(x) \sin(x)}{8a^3} + \frac{b \cos(x) \sin^2(x)}{3a^2} - \frac{\cos(x) \sin^3(x)}{4a} - \frac{\int \frac{-3(3a^4+4a^2b^2+8b^4)}{a+b \csc(x)} dx}{24a^3} \\
&= \frac{(3a^4 + 4a^2b^2 + 8b^4)x}{8a^5} + \frac{b(2a^2 + 3b^2) \cos(x)}{3a^4} - \frac{(3a^2 + 4b^2) \cos(x) \sin(x)}{8a^3} + \frac{b \cos(x) \sin^2(x)}{3a^2} - \frac{\cos(x) \sin^3(x)}{4a} \\
&= \frac{(3a^4 + 4a^2b^2 + 8b^4)x}{8a^5} + \frac{b(2a^2 + 3b^2) \cos(x)}{3a^4} - \frac{(3a^2 + 4b^2) \cos(x) \sin(x)}{8a^3} + \frac{b \cos(x) \sin^2(x)}{3a^2} - \frac{\cos(x) \sin^3(x)}{4a} \\
&= \frac{(3a^4 + 4a^2b^2 + 8b^4)x}{8a^5} + \frac{b(2a^2 + 3b^2) \cos(x)}{3a^4} - \frac{(3a^2 + 4b^2) \cos(x) \sin(x)}{8a^3} + \frac{b \cos(x) \sin^2(x)}{3a^2} - \frac{\cos(x) \sin^3(x)}{4a} \\
&= \frac{(3a^4 + 4a^2b^2 + 8b^4)x}{8a^5} + \frac{b(2a^2 + 3b^2) \cos(x)}{3a^4} - \frac{(3a^2 + 4b^2) \cos(x) \sin(x)}{8a^3} + \frac{b \cos(x) \sin^2(x)}{3a^2} - \frac{\cos(x) \sin^3(x)}{4a} \\
&= \frac{(3a^4 + 4a^2b^2 + 8b^4)x}{8a^5} + \frac{2b^5 \tanh^{-1}\left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{x}{2}\right)\right)}{\sqrt{a^2 - b^2}}\right)}{a^5 \sqrt{a^2 - b^2}} + \frac{b(2a^2 + 3b^2) \cos(x)}{3a^4} - \frac{(3a^2 + 4b^2) \cos(x) \sin(x)}{8a^3}
\end{aligned}$$

Mathematica [A] time = 0.315207, size = 129, normalized size = 0.9

$$\frac{48a^2b^2x - 24a^2b^2 \sin(2x) + 24ab(3a^2 + 4b^2) \cos(x) - \frac{192b^5 \tan^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}}\right)}{\sqrt{b^2-a^2}} - 8a^3b \cos(3x) + 36a^4x - 24a^4 \sin(2x) + 3a^4 \sin^3(x)}{96a^5}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]^4/(a + b*Csc[x]),x]

[Out] (36*a^4*x + 48*a^2*b^2*x + 96*b^4*x - (192*b^5*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + 24*a*b*(3*a^2 + 4*b^2)*Cos[x] - 8*a^3*b*C

os[3*x] - 24*a^4*Sin[2*x] - 24*a^2*b^2*Sin[2*x] + 3*a^4*Sin[4*x])/(96*a^5)

Maple [B] time = 0.079, size = 405, normalized size = 2.8

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)^4/(a+b*csc(x)),x)

[Out] $\frac{3}{4} \frac{a}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^7 + \frac{1}{a^3} \frac{1}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^7 * b^2 + \frac{2}{a^4} \frac{1}{(\tan(1/2*x)^2+1)^4} b^3 \tan(1/2*x)^6 + \frac{1}{a^3} \frac{1}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^5 * b^2 + \frac{11}{4} \frac{a}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^5 + \frac{4}{a^2} \frac{1}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^4 * b + \frac{6}{a^4} \frac{1}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^4 * b^3 - \frac{1}{a^3} \frac{1}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^3 * b^2 - \frac{11}{4} \frac{a}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^3 + \frac{6}{a^4} \frac{1}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^2 * b^3 + \frac{16}{3} \frac{a^2}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x)^2 * b - \frac{3}{4} \frac{a}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x) - \frac{1}{a^3} \frac{1}{(\tan(1/2*x)^2+1)^4} \tan(1/2*x) * b^2 + \frac{4}{3} \frac{a^2}{(\tan(1/2*x)^2+1)^4} b + \frac{2}{a^4} \frac{1}{(\tan(1/2*x)^2+1)^4} b^3 + \frac{2}{a^5} \arctan(\tan(1/2*x)) * b^4 + \frac{3}{4} \frac{a}{(\tan(1/2*x)^2+1)^4} \arctan(\tan(1/2*x)) + \frac{1}{a^3} \frac{1}{(\tan(1/2*x)^2+1)^4} \arctan(\tan(1/2*x)) * b^2 - \frac{2}{a^5} \frac{1}{(-a^2+b^2)^{1/2}} \arctan(1/2*(2*b*\tan(1/2*x)+2*a)) / (-a^2+b^2)^{1/2}$

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^4/(a+b*csc(x)),x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [A] time = 0.581091, size = 909, normalized size = 6.31

$$\frac{12 \sqrt{a^2 - b^2} b^5 \log\left(\frac{(a^2 - 2b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 + 2(b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2ab \sin(x) - a^2 - b^2}\right) - 8(a^5 b - a^3 b^3) \cos(x)^3 + 3(3a^6 + a^4 b^2)}{24(a^7 - a^5 b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^4/(a+b*csc(x)),x, algorithm="fricas")

[Out] [1/24*(12*sqrt(a^2 - b^2)*b^5*log(((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) - 8*(a^5*b - a^3*b^3)*cos(x)^3 + 3*(3*a^6 + a^4*b^2 + 4*a^2*b^4 - 8*b^6)*x + 24*(a^5*b - a*b^5)*cos(x) + 3*(2*(a^6 - a^4*b^2)*cos(x)^3 - (5*a^6 - a^4*b^2 - 4*a^2*b^4)*cos(x))*sin(x))/(a^7 - a^5*b^2), 1/24*(24*sqrt(-a^2 + b^2)*b^5*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/((a^2 - b^2)*cos(x))) - 8*(a^5*b - a^3*b^3)*cos(x)^3 + 3*(3*a^6 + a^4*b^2 + 4*a^2*b^4 - 8*b^6)*x + 24*(a^5*b - a*b^5)*cos(x) + 3*(2*(a^6 - a^4*b^2)*cos(x)^3 - (5*a^6 - a^4*b^2 - 4*a^2*b^4)*cos(x))*sin(x))/(a^7 - a^5*b^2)]

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{\sin^4(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)**4/(a+b*csc(x)),x)

[Out] Integral(sin(x)**4/(a + b*csc(x)), x)

Giac [A] time = 1.38791, size = 340, normalized size = 2.36

$$\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}} \right) \right) b^5}{\sqrt{-a^2 + b^2} a^5} + \frac{(3a^4 + 4a^2b^2 + 8b^4)x}{8a^5} + \frac{9a^3 \tan\left(\frac{1}{2}x\right)^7 + 12ab^2 \tan\left(\frac{1}{2}x\right)^7 + 24b^3}{8a^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^4/(a+b*csc(x)),x, algorithm="giac")

[Out] -2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))*b^5/(sqrt(-a^2 + b^2)*a^5) + 1/8*(3*a^4 + 4*a^2*b^2 + 8*b^4)*x/a^5

$$\begin{aligned} &+ 1/12*(9*a^3*\tan(1/2*x)^7 + 12*a*b^2*\tan(1/2*x)^7 + 24*b^3*\tan(1/2*x)^6 + \\ &33*a^3*\tan(1/2*x)^5 + 12*a*b^2*\tan(1/2*x)^5 + 48*a^2*b*\tan(1/2*x)^4 + 72*b^3*\tan(1/2*x)^4 - 33*a^3*\tan(1/2*x)^3 - 12*a*b^2*\tan(1/2*x)^3 + 64*a^2*b*\tan(1/2*x)^2 + 72*b^3*\tan(1/2*x)^2 - 9*a^3*\tan(1/2*x) - 12*a*b^2*\tan(1/2*x) + 16*a^2*b + 24*b^3)/((\tan(1/2*x)^2 + 1)^4*a^4) \end{aligned}$$

$$3.49 \quad \int \frac{1}{(a+b \csc(c+dx))^2} dx$$

Optimal. Leaf size=108

$$\frac{2b(2a^2 - b^2) \tanh^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2-b^2}}\right)}{a^2 d (a^2 - b^2)^{3/2}} - \frac{b^2 \cot(c+dx)}{ad(a^2 - b^2)(a + b \csc(c+dx))} + \frac{x}{a^2}$$

[Out] x/a^2 + (2*b*(2*a^2 - b^2)*ArcTanh[(a + b*Tan[(c + d*x)/2])/Sqrt[a^2 - b^2]])/(a^2*(a^2 - b^2)^(3/2)*d) - (b^2*Cot[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Csc[c + d*x]))

Rubi [A] time = 0.171399, antiderivative size = 108, normalized size of antiderivative = 1., number of steps used = 6, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.5$, Rules used = {3785, 3919, 3831, 2660, 618, 206}

$$\frac{2b(2a^2 - b^2) \tanh^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2-b^2}}\right)}{a^2 d (a^2 - b^2)^{3/2}} - \frac{b^2 \cot(c+dx)}{ad(a^2 - b^2)(a + b \csc(c+dx))} + \frac{x}{a^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Csc[c + d*x])^(-2), x]

[Out] x/a^2 + (2*b*(2*a^2 - b^2)*ArcTanh[(a + b*Tan[(c + d*x)/2])/Sqrt[a^2 - b^2]])/(a^2*(a^2 - b^2)^(3/2)*d) - (b^2*Cot[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Csc[c + d*x]))

Rule 3785

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^(n_.), x_Symbol] := Simp[(b^2*Cot[c + d*x]*(a + b*Csc[c + d*x])^(n + 1))/(a*d*(n + 1)*(a^2 - b^2)), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3919


```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))/(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.)), x_Symbol] := Simp[(c*x)/a, x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]
]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c -
a*d, 0]
```

Rule 3831

```
Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol]
:= Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]
```

Rule 2660

```
Int[((a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

Rule 618

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 206

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a + b \csc(c + dx))^2} dx &= -\frac{b^2 \cot(c + dx)}{a(a^2 - b^2)d(a + b \csc(c + dx))} - \frac{\int \frac{-a^2 + b^2 + ab \csc(c + dx)}{a + b \csc(c + dx)} dx}{a(a^2 - b^2)} \\
&= \frac{x}{a^2} - \frac{b^2 \cot(c + dx)}{a(a^2 - b^2)d(a + b \csc(c + dx))} - \frac{(b(2a^2 - b^2)) \int \frac{\csc(c + dx)}{a + b \csc(c + dx)} dx}{a^2(a^2 - b^2)} \\
&= \frac{x}{a^2} - \frac{b^2 \cot(c + dx)}{a(a^2 - b^2)d(a + b \csc(c + dx))} - \frac{(2a^2 - b^2) \int \frac{1}{1 + \frac{a \sin(c + dx)}{b}} dx}{a^2(a^2 - b^2)} \\
&= \frac{x}{a^2} - \frac{b^2 \cot(c + dx)}{a(a^2 - b^2)d(a + b \csc(c + dx))} - \frac{(2(2a^2 - b^2)) \text{Subst} \left(\int \frac{1}{1 + \frac{2ax}{b} + x^2} dx, x, \tan \left(\frac{1}{2}(c + dx) \right) \right)}{a^2(a^2 - b^2)d} \\
&= \frac{x}{a^2} - \frac{b^2 \cot(c + dx)}{a(a^2 - b^2)d(a + b \csc(c + dx))} + \frac{(4(2a^2 - b^2)) \text{Subst} \left(\int \frac{1}{-4 \left(1 - \frac{a^2}{b^2} \right) - x^2} dx, x, \frac{2a}{b} + 2 \tan \left(\frac{1}{2}(c + dx) \right) \right)}{a^2(a^2 - b^2)d} \\
&= \frac{x}{a^2} + \frac{2b(2a^2 - b^2) \tanh^{-1} \left(\frac{b \left(\frac{a}{b} + \tan \left(\frac{1}{2}(c + dx) \right) \right)}{\sqrt{a^2 - b^2}} \right)}{a^2(a^2 - b^2)^{3/2}d} - \frac{b^2 \cot(c + dx)}{a(a^2 - b^2)d(a + b \csc(c + dx))}
\end{aligned}$$

Mathematica [A] time = 0.455729, size = 139, normalized size = 1.29

$$\frac{\csc(c + dx)(a \sin(c + dx) + b) \left(-\frac{2b(b^2 - 2a^2) \tan^{-1} \left(\frac{a + b \tan \left(\frac{1}{2}(c + dx) \right)}{\sqrt{b^2 - a^2}} \right) (a + b \csc(c + dx))}{(b^2 - a^2)^{3/2}} + \frac{ab^2 \cot(c + dx)}{(b - a)(a + b)} + (c + dx)(a + b \csc(c + dx)) \right)}{a^2 d (a + b \csc(c + dx))^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Csc[c + d*x])^(-2), x]

[Out] (Csc[c + d*x]*((a*b^2*Cot[c + d*x])/((-a + b)*(a + b)) + (c + d*x)*(a + b*Csc[c + d*x]) - (2*b*(-2*a^2 + b^2)*ArcTan[(a + b*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])*(a + b*Csc[c + d*x]))/(-a^2 + b^2)^(3/2)*(b + a*Sin[c + d*x]))/(a^2*d*(a + b*Csc[c + d*x])^2)

Maple [B] time = 0.093, size = 247, normalized size = 2.3

$$2 \frac{\arctan(\tan(1/2 dx + c/2))}{da^2} - 2 \frac{b \tan(1/2 dx + c/2)}{d((\tan(1/2 dx + c/2))^2 b + 2 a \tan(1/2 dx + c/2) + b)(a^2 - b^2)} - 2 \frac{1}{ad((\tan(1/2 dx + c/2))^2 b + 2 a \tan(1/2 dx + c/2) + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+b*csc(d*x+c))^2,x)

[Out] 2/d/a^2*arctan(tan(1/2*d*x+1/2*c))-2/d*b/(tan(1/2*d*x+1/2*c)^2*b+2*a*tan(1/2*d*x+1/2*c)+b)/(a^2-b^2)*tan(1/2*d*x+1/2*c)-2/d*b^2/a/(tan(1/2*d*x+1/2*c)^2*b+2*a*tan(1/2*d*x+1/2*c)+b)/(a^2-b^2)-4/d*b/(a^2-b^2)/(-a^2+b^2)^(1/2)*arctan(1/2*(2*b*tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^(1/2))+2/d*b^3/a^2/(a^2-b^2)/(-a^2+b^2)^(1/2)*arctan(1/2*(2*b*tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^(1/2))

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^2,x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [B] time = 0.563269, size = 1076, normalized size = 9.96

$$\left[\frac{2(a^5 - 2a^3b^2 + ab^4)dx \sin(dx + c) + 2(a^4b - 2a^2b^3 + b^5)dx + (2a^2b^2 - b^4 + (2a^3b - ab^3) \sin(dx + c))\sqrt{a^2 - b^2} \log\left(\frac{2((a^7 - 2a^5b^2 + a^3b^4)d \sin(dx + c) + (a^2 - b^2)\sqrt{a^2 - b^2})}{2(a^2b^2 - b^4 + (2a^3b - ab^3) \sin(dx + c))\sqrt{a^2 - b^2}}\right)}{2((a^7 - 2a^5b^2 + a^3b^4)d \sin(dx + c) + (a^2 - b^2)\sqrt{a^2 - b^2})} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^2,x, algorithm="fricas")

[Out] [1/2*(2*(a^5 - 2*a^3*b^2 + a*b^4)*d*x*sin(d*x + c) + 2*(a^4*b - 2*a^2*b^3 + b^5)*d*x + (2*a^2*b^2 - b^4 + (2*a^3*b - a*b^3)*sin(d*x + c))*sqrt(a^2 - b^2)]

$$\begin{aligned} &^2) * \log(((a^2 - 2*b^2) * \cos(d*x + c))^2 + 2*a*b*\sin(d*x + c) + a^2 + b^2 + 2* \\ &(b*\cos(d*x + c)*\sin(d*x + c) + a*\cos(d*x + c))*\sqrt{a^2 - b^2}) / (a^2*\cos(d* \\ &x + c)^2 - 2*a*b*\sin(d*x + c) - a^2 - b^2)) - 2*(a^3*b^2 - a*b^4)*\cos(d*x + \\ &c)) / ((a^7 - 2*a^5*b^2 + a^3*b^4)*d*\sin(d*x + c) + (a^6*b - 2*a^4*b^3 + a^2 \\ &*b^5)*d), ((a^5 - 2*a^3*b^2 + a*b^4)*d*x*\sin(d*x + c) + (a^4*b - 2*a^2*b^3 \\ &+ b^5)*d*x + (2*a^2*b^2 - b^4 + (2*a^3*b - a*b^3)*\sin(d*x + c))*\sqrt{-a^2 + \\ &b^2} * \arctan(-\sqrt{-a^2 + b^2} * (b*\sin(d*x + c) + a) / ((a^2 - b^2) * \cos(d*x + \\ &c)))) - (a^3*b^2 - a*b^4)*\cos(d*x + c)) / ((a^7 - 2*a^5*b^2 + a^3*b^4)*d*\sin(d \\ &*x + c) + (a^6*b - 2*a^4*b^3 + a^2*b^5)*d)] \end{aligned}$$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{(a + b \csc(c + dx))^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))**2,x)

[Out] Integral((a + b*csc(c + d*x))**(-2), x)

Giac [A] time = 1.38376, size = 213, normalized size = 1.97

$$\frac{2(2a^2b - b^3) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(b) + \arctan \left(\frac{b \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + a}{\sqrt{-a^2 + b^2}} \right) \right)}{(a^4 - a^2 b^2) \sqrt{-a^2 + b^2}} + \frac{2 \left(ab \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + b^2 \right)}{(a^3 - ab^2) \left(b \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 + 2a \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + b \right)} - \frac{dx+c}{a^2}$$

$$d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^2,x, algorithm="giac")

[Out] $-(2*(2*a^2*b - b^3)*(pi*\text{floor}(1/2*(d*x + c)/pi + 1/2)*\operatorname{sgn}(b) + \arctan((b*\tan(1/2*d*x + 1/2*c) + a)/\sqrt{-a^2 + b^2}))) / ((a^4 - a^2*b^2)*\sqrt{-a^2 + b^2}) + 2*(a*b*\tan(1/2*d*x + 1/2*c) + b^2) / ((a^3 - a*b^2)*(b*\tan(1/2*d*x + 1/2*c)^2 + 2*a*\tan(1/2*d*x + 1/2*c) + b)) - (d*x + c)/a^2)/d$

$$3.50 \quad \int \frac{1}{(a+b \csc(c+dx))^3} dx$$

Optimal. Leaf size=170

$$\frac{b(-5a^2b^2 + 6a^4 + 2b^4) \tanh^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2-b^2}}\right)}{a^3d(a^2-b^2)^{5/2}} - \frac{b^2(5a^2-2b^2) \cot(c+dx)}{2a^2d(a^2-b^2)^2(a+b \csc(c+dx))} - \frac{b^2 \cot(c+dx)}{2ad(a^2-b^2)(a+b \csc(c+dx))}$$

[Out] x/a^3 + (b*(6*a^4 - 5*a^2*b^2 + 2*b^4)*ArcTanh[(a + b*Tan[(c + d*x)/2])/Sqrt[a^2 - b^2]])/(a^3*(a^2 - b^2)^(5/2)*d) - (b^2*Cot[c + d*x])/(2*a*(a^2 - b^2)*d*(a + b*Csc[c + d*x])^2) - (b^2*(5*a^2 - 2*b^2)*Cot[c + d*x])/(2*a^2*(a^2 - b^2)^2*d*(a + b*Csc[c + d*x]))

Rubi [A] time = 0.318968, antiderivative size = 170, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 7, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3785, 4060, 3919, 3831, 2660, 618, 206}

$$\frac{b(-5a^2b^2 + 6a^4 + 2b^4) \tanh^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2-b^2}}\right)}{a^3d(a^2-b^2)^{5/2}} - \frac{b^2(5a^2-2b^2) \cot(c+dx)}{2a^2d(a^2-b^2)^2(a+b \csc(c+dx))} - \frac{b^2 \cot(c+dx)}{2ad(a^2-b^2)(a+b \csc(c+dx))}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Csc[c + d*x])^(-3), x]

[Out] x/a^3 + (b*(6*a^4 - 5*a^2*b^2 + 2*b^4)*ArcTanh[(a + b*Tan[(c + d*x)/2])/Sqrt[a^2 - b^2]])/(a^3*(a^2 - b^2)^(5/2)*d) - (b^2*Cot[c + d*x])/(2*a*(a^2 - b^2)*d*(a + b*Csc[c + d*x])^2) - (b^2*(5*a^2 - 2*b^2)*Cot[c + d*x])/(2*a^2*(a^2 - b^2)^2*d*(a + b*Csc[c + d*x]))

Rule 3785

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^ (n_.), x_Symbol] :> Simp[(b^2*Cot[c + d*x]*(a + b*Csc[c + d*x])^(n + 1))/(a*d*(n + 1)*(a^2 - b^2)), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]

Rule 4060

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.)
)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_Symbol] := Simp[((A*b^2 -
a*b*B + a^2*C)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1))/(a*f*(m + 1)*(a^
2 - b^2)), x] + Dist[1/(a*(m + 1)*(a^2 - b^2)), Int[(a + b*Csc[e + f*x])^(m
+ 1)*Simp[A*(a^2 - b^2)*(m + 1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x]
+ (A*b^2 - a*b*B + a^2*C)*(m + 2)*Csc[e + f*x]^2, x], x] /; FreeQ[{a,
b, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]
```

Rule 3919

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))/(csc[(e_.) + (f_.)*(x_.)]*(b_.) +
(a_.)), x_Symbol] := Simp[(c*x)/a, x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x
]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c -
a*d, 0]
```

Rule 3831

```
Int[csc[(e_.) + (f_.)*(x_.)]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbo
l] := Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f
}, x] && NeQ[a^2 - b^2, 0]
```

Rule 2660

```
Int[((a_.) + (b_.)*sin[(c_.) + (d_.)*(x_.)])^(-1), x_Symbol] := With[{e = Fre
eFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*
e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[
a^2 - b^2, 0]
```

Rule 618

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 206

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/
Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a + b \csc(c + dx))^3} dx &= -\frac{b^2 \cot(c + dx)}{2a(a^2 - b^2)d(a + b \csc(c + dx))^2} - \frac{\int \frac{-2(a^2 - b^2) + 2ab \csc(c + dx) - b^2 \csc^2(c + dx)}{(a + b \csc(c + dx))^2} dx}{2a(a^2 - b^2)} \\
&= -\frac{b^2 \cot(c + dx)}{2a(a^2 - b^2)d(a + b \csc(c + dx))^2} - \frac{b^2(5a^2 - 2b^2) \cot(c + dx)}{2a^2(a^2 - b^2)^2 d(a + b \csc(c + dx))} + \frac{\int \frac{2(a^2 - b^2)^2 - ab(4a^2 - b^2)}{a + b \csc(c + dx)} dx}{2a^2(a^2 - b^2)^2} \\
&= \frac{x}{a^3} - \frac{b^2 \cot(c + dx)}{2a(a^2 - b^2)d(a + b \csc(c + dx))^2} - \frac{b^2(5a^2 - 2b^2) \cot(c + dx)}{2a^2(a^2 - b^2)^2 d(a + b \csc(c + dx))} - \frac{(b(6a^4 - 5a^2b^2 + 2b^4) \tanh^{-1}\left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{1}{2}(c + dx)\right)\right)}{\sqrt{a^2 - b^2}}\right))}{2a^2(a^2 - b^2)^2} \\
&= \frac{x}{a^3} - \frac{b^2 \cot(c + dx)}{2a(a^2 - b^2)d(a + b \csc(c + dx))^2} - \frac{b^2(5a^2 - 2b^2) \cot(c + dx)}{2a^2(a^2 - b^2)^2 d(a + b \csc(c + dx))} - \frac{(6a^4 - 5a^2b^2 + 2b^4) \tanh^{-1}\left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{1}{2}(c + dx)\right)\right)}{\sqrt{a^2 - b^2}}\right)}{2a^2(a^2 - b^2)^2} \\
&= \frac{x}{a^3} - \frac{b^2 \cot(c + dx)}{2a(a^2 - b^2)d(a + b \csc(c + dx))^2} - \frac{b^2(5a^2 - 2b^2) \cot(c + dx)}{2a^2(a^2 - b^2)^2 d(a + b \csc(c + dx))} - \frac{(6a^4 - 5a^2b^2 + 2b^4) \tanh^{-1}\left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{1}{2}(c + dx)\right)\right)}{\sqrt{a^2 - b^2}}\right)}{2a^2(a^2 - b^2)^2} \\
&= \frac{x}{a^3} - \frac{b^2 \cot(c + dx)}{2a(a^2 - b^2)d(a + b \csc(c + dx))^2} - \frac{b^2(5a^2 - 2b^2) \cot(c + dx)}{2a^2(a^2 - b^2)^2 d(a + b \csc(c + dx))} + \frac{(2(6a^4 - 5a^2b^2 + 2b^4) \tanh^{-1}\left(\frac{b\left(\frac{a}{b} + \tan\left(\frac{1}{2}(c + dx)\right)\right)}{\sqrt{a^2 - b^2}}\right))}{2a^3(a^2 - b^2)^{5/2}d} - \frac{b^2 \cot(c + dx)}{2a(a^2 - b^2)d(a + b \csc(c + dx))^2}
\end{aligned}$$

Mathematica [A] time = 1.10303, size = 216, normalized size = 1.27

$$\frac{\csc^2(c + dx)(a \sin(c + dx) + b) \left(-\frac{3ab^2(2a^2 - b^2) \cot(c + dx)(a \sin(c + dx) + b)}{(a - b)^2(a + b)^2} - \frac{2b(-5a^2b^2 + 6a^4 + 2b^4) \csc(c + dx)(a \sin(c + dx) + b)^2 \tan^{-1}\left(\frac{a + b \tan\left(\frac{1}{2}(c + dx)\right)}{\sqrt{b^2 - a^2}}\right)}{(b^2 - a^2)^{5/2}} \right)}{2a^3d(a + b \csc(c + dx))^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Csc[c + d*x])^(-3), x]

[Out] (Csc[c + d*x]^2*(b + a*Sin[c + d*x])*((a*b^3*Cot[c + d*x])/((a - b)*(a + b) - (3*a*b^2*(2*a^2 - b^2)*Cot[c + d*x]*(b + a*Sin[c + d*x]))/((a - b)^2*(a

$$+ b)^2) + 2*(c + d*x)*Csc[c + d*x]*(b + a*\sin[c + d*x])^2 - (2*b*(6*a^4 - 5*a^2*b^2 + 2*b^4)*ArcTan[(a + b*\tan[(c + d*x)/2])/sqrt[-a^2 + b^2]]*Csc[c + d*x]*(b + a*\sin[c + d*x])^2)/(-a^2 + b^2)^(5/2))/(2*a^3*d*(a + b*Csc[c + d*x])^3)$$

Maple [B] time = 0.115, size = 796, normalized size = 4.7

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+b*csc(d*x+c))^3,x)

[Out]
$$\begin{aligned} & 2/d/a^3*\arctan(\tan(1/2*d*x+1/2*c))-4/d*a*b^2/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)*\tan(1/2*d*x+1/2*c)^3+1/d/a*b^4/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)*\tan(1/2*d*x+1/2*c)^3-10/d*a^2*b/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)*\tan(1/2*d*x+1/2*c)^2-1/d*b^3/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)*\tan(1/2*d*x+1/2*c)^2+2/d/a^2*b^5/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)*\tan(1/2*d*x+1/2*c)^2-16/d*a*b^2/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)*\tan(1/2*d*x+1/2*c)+7/d/a*b^4/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)*\tan(1/2*d*x+1/2*c)-5/d*b^3/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)+2/d/a^2*b^5/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^2/(a^4-2*a^2*b^2+b^4)-6/d*a*b/(a^4-2*a^2*b^2+b^4)/(-a^2+b^2)^(1/2)*\arctan(1/2*(2*b*\tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^(1/2))+5/d/a*b^3/(a^4-2*a^2*b^2+b^4)/(-a^2+b^2)^(1/2)*\arctan(1/2*(2*b*\tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^(1/2))-2/d/a^3*b^5/(a^4-2*a^2*b^2+b^4)/(-a^2+b^2)^(1/2)*\arctan(1/2*(2*b*\tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^(1/2)) \end{aligned}$$

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^3,x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [B] time = 0.645056, size = 1994, normalized size = 11.73

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^3,x, algorithm="fricas")

[Out]
$$\begin{aligned} & [1/4*(4*(a^8 - 3*a^6*b^2 + 3*a^4*b^4 - a^2*b^6)*d*x*cos(d*x + c)^2 - 4*(a^8 \\ & - 2*a^6*b^2 + 2*a^2*b^6 - b^8)*d*x - (6*a^6*b + a^4*b^3 - 3*a^2*b^5 + 2*b^7 \\ & - (6*a^6*b - 5*a^4*b^3 + 2*a^2*b^5)*cos(d*x + c)^2 + 2*(6*a^5*b^2 - 5*a^3 \\ & *b^4 + 2*a*b^6)*sin(d*x + c))*sqrt(a^2 - b^2)*log(((a^2 - 2*b^2)*cos(d*x + \\ & c)^2 + 2*a*b*sin(d*x + c) + a^2 + b^2 + 2*(b*cos(d*x + c)*sin(d*x + c) + a* \\ & cos(d*x + c))*sqrt(a^2 - b^2))/(a^2*cos(d*x + c)^2 - 2*a*b*sin(d*x + c) - a \\ & ^2 - b^2)) + 2*(5*a^5*b^3 - 7*a^3*b^5 + 2*a*b^7)*cos(d*x + c) - 2*(4*(a^7*b \\ & - 3*a^5*b^3 + 3*a^3*b^5 - a*b^7)*d*x - 3*(2*a^6*b^2 - 3*a^4*b^4 + a^2*b^6) \\ & *cos(d*x + c))*sin(d*x + c))/((a^11 - 3*a^9*b^2 + 3*a^7*b^4 - a^5*b^6)*d*co \\ & s(d*x + c)^2 - 2*(a^10*b - 3*a^8*b^3 + 3*a^6*b^5 - a^4*b^7)*d*sin(d*x + c) \\ & - (a^11 - 2*a^9*b^2 + 2*a^5*b^6 - a^3*b^8)*d), 1/2*(2*(a^8 - 3*a^6*b^2 + 3* \\ & a^4*b^4 - a^2*b^6)*d*x*cos(d*x + c)^2 - 2*(a^8 - 2*a^6*b^2 + 2*a^2*b^6 - b^ \\ & 8)*d*x - (6*a^6*b + a^4*b^3 - 3*a^2*b^5 + 2*b^7 - (6*a^6*b - 5*a^4*b^3 + 2* \\ & a^2*b^5)*cos(d*x + c)^2 + 2*(6*a^5*b^2 - 5*a^3*b^4 + 2*a*b^6)*sin(d*x + c)) \\ & *sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*sin(d*x + c) + a)/((a^2 - b^2) \\ &)*cos(d*x + c))) + (5*a^5*b^3 - 7*a^3*b^5 + 2*a*b^7)*cos(d*x + c) - (4*(a^7 \\ & *b - 3*a^5*b^3 + 3*a^3*b^5 - a*b^7)*d*x - 3*(2*a^6*b^2 - 3*a^4*b^4 + a^2*b^ \\ & 6)*cos(d*x + c))*sin(d*x + c))/((a^11 - 3*a^9*b^2 + 3*a^7*b^4 - a^5*b^6)*d* \\ & cos(d*x + c)^2 - 2*(a^10*b - 3*a^8*b^3 + 3*a^6*b^5 - a^4*b^7)*d*sin(d*x + c) \\ &) - (a^11 - 2*a^9*b^2 + 2*a^5*b^6 - a^3*b^8)*d]] \end{aligned}$$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{(a + b \csc(c + dx))^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))**3,x)

[Out] Integral((a + b*csc(c + d*x))**(-3), x)

Giac [A] time = 1.18064, size = 401, normalized size = 2.36

$$\frac{(6a^4b - 5a^2b^3 + 2b^5) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(b) + \arctan \left(\frac{b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + a}{\sqrt{-a^2+b^2}} \right) \right)}{(a^7 - 2a^5b^2 + a^3b^4) \sqrt{-a^2+b^2}} + \frac{4a^3b^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - ab^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 10a^4b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + a^2b^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{(a^6 - 2a^4b^2 + a^2b^4) \left(b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + a \right)} \frac{1}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^3,x, algorithm="giac")

[Out] -((6*a^4*b - 5*a^2*b^3 + 2*b^5)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*d*x + 1/2*c) + a)/sqrt(-a^2 + b^2)))/((a^7 - 2*a^5*b^2 + a^3*b^4)*sqrt(-a^2 + b^2)) + (4*a^3*b^2*tan(1/2*d*x + 1/2*c)^3 - a*b^4*tan(1/2*d*x + 1/2*c)^3 + 10*a^4*b*tan(1/2*d*x + 1/2*c)^2 + a^2*b^3*tan(1/2*d*x + 1/2*c)^2 - 2*b^5*tan(1/2*d*x + 1/2*c)^2 + 16*a^3*b^2*tan(1/2*d*x + 1/2*c) - 7*a*b^4*tan(1/2*d*x + 1/2*c) + 5*a^2*b^3 - 2*b^5)/((a^6 - 2*a^4*b^2 + a^2*b^4)*(b*tan(1/2*d*x + 1/2*c)^2 + 2*a*tan(1/2*d*x + 1/2*c) + b)^2) - (d*x + c)/a^3)/d

$$3.51 \quad \int \frac{1}{(a+b \csc(c+dx))^4} dx$$

Optimal. Leaf size=239

$$\frac{b(-8a^4b^2 + 7a^2b^4 + 8a^6 - 2b^6) \tanh^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2-b^2}}\right)}{a^4d(a^2-b^2)^{7/2}} - \frac{b^2(-17a^2b^2 + 26a^4 + 6b^4) \cot(c+dx)}{6a^3d(a^2-b^2)^3(a+b \csc(c+dx))} - \frac{b^2(8a^2-3b^2)}{6a^2d(a^2-b^2)^2(a+b \csc(c+dx))}$$

[Out] x/a^4 + (b*(8*a^6 - 8*a^4*b^2 + 7*a^2*b^4 - 2*b^6)*ArcTanh[(a + b*Tan[(c + d*x)/2])/Sqrt[a^2 - b^2]])/(a^4*(a^2 - b^2)^(7/2)*d) - (b^2*Cot[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Csc[c + d*x])^3) - (b^2*(8*a^2 - 3*b^2)*Cot[c + d*x])/(6*a^2*(a^2 - b^2)^2*d*(a + b*Csc[c + d*x])^2) - (b^2*(26*a^4 - 17*a^2*b^2 + 6*b^4)*Cot[c + d*x])/(6*a^3*(a^2 - b^2)^3*d*(a + b*Csc[c + d*x]))

Rubi [A] time = 0.50425, antiderivative size = 239, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 7, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3785, 4060, 3919, 3831, 2660, 618, 206}

$$\frac{b(-8a^4b^2 + 7a^2b^4 + 8a^6 - 2b^6) \tanh^{-1}\left(\frac{a+b \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2-b^2}}\right)}{a^4d(a^2-b^2)^{7/2}} - \frac{b^2(-17a^2b^2 + 26a^4 + 6b^4) \cot(c+dx)}{6a^3d(a^2-b^2)^3(a+b \csc(c+dx))} - \frac{b^2(8a^2-3b^2)}{6a^2d(a^2-b^2)^2(a+b \csc(c+dx))}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Csc[c + d*x])^(-4), x]

[Out] x/a^4 + (b*(8*a^6 - 8*a^4*b^2 + 7*a^2*b^4 - 2*b^6)*ArcTanh[(a + b*Tan[(c + d*x)/2])/Sqrt[a^2 - b^2]])/(a^4*(a^2 - b^2)^(7/2)*d) - (b^2*Cot[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Csc[c + d*x])^3) - (b^2*(8*a^2 - 3*b^2)*Cot[c + d*x])/(6*a^2*(a^2 - b^2)^2*d*(a + b*Csc[c + d*x])^2) - (b^2*(26*a^4 - 17*a^2*b^2 + 6*b^4)*Cot[c + d*x])/(6*a^3*(a^2 - b^2)^3*d*(a + b*Csc[c + d*x]))

Rule 3785

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^ (n_.), x_Symbol] := Simp[(b^2*Cot[c + d*x]*(a + b*Csc[c + d*x])^(n + 1))/(a*d*(n + 1)*(a^2 - b^2)), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && Integ

rQ[2*n]

Rule 4060

Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_Symbol] :> Simp[((A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1))/(a*f*(m + 1)*(a^2 - b^2)), x] + Dist[1/(a*(m + 1)*(a^2 - b^2)), Int[(a + b*Csc[e + f*x])^(m + 1)*Simp[A*(a^2 - b^2)*(m + 1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x] + (A*b^2 - a*b*B + a^2*C)*(m + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]

Rule 3919

Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Simp[(c*x)/a, x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]

Rule 3831

Int[csc[(e_.) + (f_.)*(x_.)]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]

Rule 2660

Int[((a_.) + (b_.)*sin[(c_.) + (d_.)*(x_.)])^(-1), x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 618

Int[((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 206

Int[((a_.) + (b_.)*(x_.)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a+b \csc(c+dx))^4} dx &= -\frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3} - \frac{\int \frac{-3(a^2-b^2)+3ab \csc(c+dx)-2b^2 \csc^2(c+dx)}{(a+b \csc(c+dx))^3} dx}{3a(a^2-b^2)} \\
&= -\frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3} - \frac{b^2(8a^2-3b^2)\cot(c+dx)}{6a^2(a^2-b^2)^2 d(a+b \csc(c+dx))^2} + \frac{\int \frac{6(a^2-b^2)^2-2ab}{(a+b \csc(c+dx))^3} dx}{6a^2(a^2-b^2)^2} \\
&= -\frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3} - \frac{b^2(8a^2-3b^2)\cot(c+dx)}{6a^2(a^2-b^2)^2 d(a+b \csc(c+dx))^2} - \frac{b^2(26a^4-17a^2b^2)}{6a^3(a^2-b^2)} \\
&= \frac{x}{a^4} - \frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3} - \frac{b^2(8a^2-3b^2)\cot(c+dx)}{6a^2(a^2-b^2)^2 d(a+b \csc(c+dx))^2} - \frac{b^2(26a^4-17a^2b^2)}{6a^3(a^2-b^2)} \\
&= \frac{x}{a^4} - \frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3} - \frac{b^2(8a^2-3b^2)\cot(c+dx)}{6a^2(a^2-b^2)^2 d(a+b \csc(c+dx))^2} - \frac{b^2(26a^4-17a^2b^2)}{6a^3(a^2-b^2)} \\
&= \frac{x}{a^4} - \frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3} - \frac{b^2(8a^2-3b^2)\cot(c+dx)}{6a^2(a^2-b^2)^2 d(a+b \csc(c+dx))^2} - \frac{b^2(26a^4-17a^2b^2)}{6a^3(a^2-b^2)} \\
&= \frac{x}{a^4} - \frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3} - \frac{b^2(8a^2-3b^2)\cot(c+dx)}{6a^2(a^2-b^2)^2 d(a+b \csc(c+dx))^2} - \frac{b^2(26a^4-17a^2b^2)}{6a^3(a^2-b^2)} \\
&= \frac{x}{a^4} - \frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3} - \frac{b^2(8a^2-3b^2)\cot(c+dx)}{6a^2(a^2-b^2)^2 d(a+b \csc(c+dx))^2} - \frac{b^2(26a^4-17a^2b^2)}{6a^3(a^2-b^2)} \\
&= \frac{x}{a^4} + \frac{b(8a^6-8a^4b^2+7a^2b^4-2b^6) \tanh^{-1}\left(\frac{b\left(\frac{a}{b}+\tan\left(\frac{1}{2}(c+dx)\right)\right)}{\sqrt{a^2-b^2}}\right)}{a^4(a^2-b^2)^{7/2}d} - \frac{b^2 \cot(c+dx)}{3a(a^2-b^2)d(a+b \csc(c+dx))^3}
\end{aligned}$$

Mathematica [A] time = 2.09468, size = 279, normalized size = 1.17

$$\csc^3(c+dx)(a \sin(c+dx)+b) \left(\frac{ab^3(12a^2-7b^2)\cot(c+dx)(a \sin(c+dx)+b)}{(a-b)^2(a+b)^2} - \frac{ab^2(-32a^2b^2+36a^4+11b^4)\cot(c+dx)(a \sin(c+dx)+b)^2}{(a-b)^3(a+b)^3} - \frac{6b(8a^4b^2-7a^2b^4)}{6a^4d(a+b \csc(c+dx))^3} \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Csc[c + d*x])^(-4),x]

[Out] (Csc[c + d*x]^3*(b + a*Sin[c + d*x])*((2*a*b^4*Cot[c + d*x])/((-a + b)*(a + b)) + (a*b^3*(12*a^2 - 7*b^2)*Cot[c + d*x]*(b + a*Sin[c + d*x]))/((a - b)^2*(a + b)^2) - (a*b^2*(36*a^4 - 32*a^2*b^2 + 11*b^4)*Cot[c + d*x]*(b + a*Sin[c + d*x])^2)/((a - b)^3*(a + b)^3) + 6*(c + d*x)*Csc[c + d*x]*(b + a*Sin[c + d*x])^3 - (6*b*(-8*a^6 + 8*a^4*b^2 - 7*a^2*b^4 + 2*b^6)*ArcTan[(a + b*Tan[(c + d*x)/2]]/Sqrt[-a^2 + b^2])*Csc[c + d*x]*(b + a*Sin[c + d*x])^3)/(-a^2 + b^2)^(7/2))/((6*a^4*d*(a + b*Csc[c + d*x])^4)

Maple [B] time = 0.147, size = 1912, normalized size = 8.

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+b*csc(d*x+c))^4,x)

[Out]
$$\begin{aligned} & -46/d*a^2*b^3/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)-2/d/a^3*b^8/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^4-12/d/a^2*b^7/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^3+2/d/a^4*b^7/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*b*\tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^{(1/2)})-7/d/a^2*b^5/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*b*\tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^{(1/2)})-104/3/d*a^4*b/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^3+17/3/d/a*b^6/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)-2/d/a^3*b^8/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)-8/d/a*b^6/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^2-4/d/a^3*b^8/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^2-28/d*a^3*b^2/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^4-11/d/a^2*b^7/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)+4/d*a*b^4/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^4+1/d/a*b^6/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^4-1/d/a^2*b^7/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^5-76/d*a^3*b^2/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^2 \end{aligned}$$

$$\begin{aligned}
&+2/d/a^4*\arctan(\tan(1/2*d*x+1/2*c))+38/d*a*b^4/(\tan(1/2*d*x+1/2*c)^2*b+2*a* \\
&\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^2- \\
&8/d*a^2*b/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*b*ta \\
&n(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^{(1/2)}+8/d*b^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^ \\
&6)/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*b*\tan(1/2*d*x+1/2*c)+2*a)/(-a^2+b^2)^{(1/2} \\
&))+26/d*b^5/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4* \\
&b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^3-26/3/d*a*b^4/(\tan(1/2*d*x+1/2*c)^2* \\
&b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)+2/d*b^5/(\tan(1/ \\
&2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)* \\
&\tan(1/2*d*x+1/2*c)^5+32/d*b^5/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c \\
&)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)-6/d*a^2*b^3/(\tan(1/ \\
&2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)* \\
&\tan(1/2*d*x+1/2*c)^5-88/3/d*a^2*b^3/(\tan(1/2*d*x+1/2*c)^2*b+2*a*\tan(1/2*d*x \\
&+1/2*c)+b)^3/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\tan(1/2*d*x+1/2*c)^3
\end{aligned}$$

Maxima [F(-2)] time = 0., size = 0, normalized size = 0.

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^4,x, algorithm="maxima")

[Out] Exception raised: ValueError

Fricas [B] time = 0.777519, size = 3389, normalized size = 14.18

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^4,x, algorithm="fricas")

[Out]
$$\begin{aligned}
&[1/12*(36*(a^{10}*b - 4*a^8*b^3 + 6*a^6*b^5 - 4*a^4*b^7 + a^2*b^9)*d*x*\cos(d* \\
&x + c)^2 - 2*(36*a^9*b^2 - 68*a^7*b^4 + 43*a^5*b^6 - 11*a^3*b^8)*\cos(d*x + \\
&c)^3 - 12*(3*a^{10}*b - 11*a^8*b^3 + 14*a^6*b^5 - 6*a^4*b^7 - a^2*b^9 + b^{11}) \\
&*d*x - 3*(24*a^8*b^2 - 16*a^6*b^4 + 13*a^4*b^6 + a^2*b^8 - 2*b^{10} - 3*(8*a^ \\
&8*b^2 - 8*a^6*b^4 + 7*a^4*b^6 - 2*a^2*b^8)*\cos(d*x + c)^2 + (8*a^9*b + 16*a \\
&^7*b^3 - 17*a^5*b^5 + 19*a^3*b^7 - 6*a*b^9 - (8*a^9*b - 8*a^7*b^3 + 7*a^5*b \\
&^5 - 2*a^3*b^7)*\cos(d*x + c)^2)*\sin(d*x + c))*\sqrt{a^2 - b^2}*\log(((a^2 - 2
\end{aligned}$$

```

*b^2)*cos(d*x + c)^2 + 2*a*b*sin(d*x + c) + a^2 + b^2 + 2*(b*cos(d*x + c)*s
in(d*x + c) + a*cos(d*x + c))*sqrt(a^2 - b^2))/(a^2*cos(d*x + c)^2 - 2*a*b*
sin(d*x + c) - a^2 - b^2)) + 12*(6*a^9*b^2 - 7*a^7*b^4 + 2*a^3*b^8 - a*b^10
)*cos(d*x + c) + 6*(2*(a^11 - 4*a^9*b^2 + 6*a^7*b^4 - 4*a^5*b^6 + a^3*b^8)*
d*x*cos(d*x + c)^2 - 2*(a^11 - a^9*b^2 - 6*a^7*b^4 + 14*a^5*b^6 - 11*a^3*b^
8 + 3*a*b^10)*d*x + 5*(4*a^8*b^3 - 7*a^6*b^5 + 4*a^4*b^7 - a^2*b^9)*cos(d*x
+ c))*sin(d*x + c))/(3*(a^14*b - 4*a^12*b^3 + 6*a^10*b^5 - 4*a^8*b^7 + a^6
*b^9)*d*cos(d*x + c)^2 - (3*a^14*b - 11*a^12*b^3 + 14*a^10*b^5 - 6*a^8*b^7
- a^6*b^9 + a^4*b^11)*d + ((a^15 - 4*a^13*b^2 + 6*a^11*b^4 - 4*a^9*b^6 + a^
7*b^8)*d*cos(d*x + c)^2 - (a^15 - a^13*b^2 - 6*a^11*b^4 + 14*a^9*b^6 - 11*a
^7*b^8 + 3*a^5*b^10)*d)*sin(d*x + c)), 1/6*(18*(a^10*b - 4*a^8*b^3 + 6*a^6*
b^5 - 4*a^4*b^7 + a^2*b^9)*d*x*cos(d*x + c)^2 - (36*a^9*b^2 - 68*a^7*b^4 +
43*a^5*b^6 - 11*a^3*b^8)*cos(d*x + c)^3 - 6*(3*a^10*b - 11*a^8*b^3 + 14*a^6
*b^5 - 6*a^4*b^7 - a^2*b^9 + b^11)*d*x - 3*(24*a^8*b^2 - 16*a^6*b^4 + 13*a^
4*b^6 + a^2*b^8 - 2*b^10 - 3*(8*a^8*b^2 - 8*a^6*b^4 + 7*a^4*b^6 - 2*a^2*b^8
)*cos(d*x + c)^2 + (8*a^9*b + 16*a^7*b^3 - 17*a^5*b^5 + 19*a^3*b^7 - 6*a*b^
9 - (8*a^9*b - 8*a^7*b^3 + 7*a^5*b^5 - 2*a^3*b^7)*cos(d*x + c)^2)*sin(d*x +
c))*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*sin(d*x + c) + a)/((a^2 -
b^2)*cos(d*x + c))) + 6*(6*a^9*b^2 - 7*a^7*b^4 + 2*a^3*b^8 - a*b^10)*cos(d
*x + c) + 3*(2*(a^11 - 4*a^9*b^2 + 6*a^7*b^4 - 4*a^5*b^6 + a^3*b^8)*d*x*cos
(d*x + c)^2 - 2*(a^11 - a^9*b^2 - 6*a^7*b^4 + 14*a^5*b^6 - 11*a^3*b^8 + 3*a
*b^10)*d*x + 5*(4*a^8*b^3 - 7*a^6*b^5 + 4*a^4*b^7 - a^2*b^9)*cos(d*x + c))*
sin(d*x + c))/(3*(a^14*b - 4*a^12*b^3 + 6*a^10*b^5 - 4*a^8*b^7 + a^6*b^9)*d
*cos(d*x + c)^2 - (3*a^14*b - 11*a^12*b^3 + 14*a^10*b^5 - 6*a^8*b^7 - a^6*b
^9 + a^4*b^11)*d + ((a^15 - 4*a^13*b^2 + 6*a^11*b^4 - 4*a^9*b^6 + a^7*b^8)*
d*cos(d*x + c)^2 - (a^15 - a^13*b^2 - 6*a^11*b^4 + 14*a^9*b^6 - 11*a^7*b^8
+ 3*a^5*b^10)*d)*sin(d*x + c))]

```

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{(a + b \csc(c + dx))^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))**4,x)

[Out] Integral((a + b*csc(c + d*x))**(-4), x)

Giac [B] time = 1.39036, size = 722, normalized size = 3.02

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*csc(d*x+c))^4,x, algorithm="giac")

[Out]
$$-1/3*(3*(8*a^6*b - 8*a^4*b^3 + 7*a^2*b^5 - 2*b^7)*(pi*\text{floor}(1/2*(d*x + c)/pi + 1/2)*\text{sgn}(b) + \arctan((b*\tan(1/2*d*x + 1/2*c) + a)/\sqrt{-a^2 + b^2}))/((a^{10} - 3*a^8*b^2 + 3*a^6*b^4 - a^4*b^6)*\sqrt{-a^2 + b^2}) + (18*a^5*b^3*\tan(1/2*d*x + 1/2*c)^5 - 6*a^3*b^5*\tan(1/2*d*x + 1/2*c)^5 + 3*a*b^7*\tan(1/2*d*x + 1/2*c)^5 + 84*a^6*b^2*\tan(1/2*d*x + 1/2*c)^4 - 12*a^4*b^4*\tan(1/2*d*x + 1/2*c)^4 - 3*a^2*b^6*\tan(1/2*d*x + 1/2*c)^4 + 6*b^8*\tan(1/2*d*x + 1/2*c)^4 + 104*a^7*b*\tan(1/2*d*x + 1/2*c)^3 + 88*a^5*b^3*\tan(1/2*d*x + 1/2*c)^3 - 7*8*a^3*b^5*\tan(1/2*d*x + 1/2*c)^3 + 36*a*b^7*\tan(1/2*d*x + 1/2*c)^3 + 228*a^6*b^2*\tan(1/2*d*x + 1/2*c)^2 - 114*a^4*b^4*\tan(1/2*d*x + 1/2*c)^2 + 24*a^2*b^6*\tan(1/2*d*x + 1/2*c)^2 + 12*b^8*\tan(1/2*d*x + 1/2*c)^2 + 138*a^5*b^3*\tan(1/2*d*x + 1/2*c) - 96*a^3*b^5*\tan(1/2*d*x + 1/2*c) + 33*a*b^7*\tan(1/2*d*x + 1/2*c) + 26*a^4*b^4 - 17*a^2*b^6 + 6*b^8)/((a^9 - 3*a^7*b^2 + 3*a^5*b^4 - a^3*b^6)*(b*\tan(1/2*d*x + 1/2*c)^2 + 2*a*\tan(1/2*d*x + 1/2*c) + b)^3) - 3*(d*x + c)/a^4)/d$$

$$3.52 \quad \int \frac{1}{3+5 \csc(c+dx)} dx$$

Optimal. Leaf size=31

$$-\frac{5 \tan^{-1}\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{6d} - \frac{x}{12}$$

[Out] $-x/12 - (5*\text{ArcTan}[\text{Cos}[c + d*x]/(3 + \text{Sin}[c + d*x])])/(6*d)$

Rubi [A] time = 0.029382, antiderivative size = 31, normalized size of antiderivative = 1., number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3783, 2657}

$$-\frac{5 \tan^{-1}\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{6d} - \frac{x}{12}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(3 + 5*\text{Csc}[c + d*x])^{-1}, x]$

[Out] $-x/12 - (5*\text{ArcTan}[\text{Cos}[c + d*x]/(3 + \text{Sin}[c + d*x])])/(6*d)$

Rule 3783

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^{-1}, x_Symbol] \rightarrow \text{Simp}[x/a, x] - \text{Dist}[1/a, \text{Int}[1/(1 + (a*\text{Sin}[c + d*x])/b), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

Rule 2657

$\text{Int}[(a_. + (b_.)*\text{sin}[(c_.) + (d_.)*(x_.)])^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a^2 - b^2, 2]\}, \text{Simp}[x/q, x] + \text{Simp}[(2*\text{ArcTan}[(b*\text{Cos}[c + d*x])/(a + q + b*\text{Sin}[c + d*x])])/(d*q), x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{GtQ}[a^2 - b^2, 0] \&\& \text{PosQ}[a]$

Rubi steps

$$\int \frac{1}{3 + 5 \csc(c + dx)} dx = \frac{x}{3} - \frac{1}{3} \int \frac{1}{1 + \frac{3}{5} \sin(c + dx)} dx$$

$$= -\frac{x}{12} - \frac{5 \tan^{-1}\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{6d}$$

Mathematica [B] time = 0.0484721, size = 66, normalized size = 2.13

$$\frac{2(c + dx) - 5 \tan^{-1}\left(\frac{2\left(\sin\left(\frac{1}{2}(c+dx)\right) + \cos\left(\frac{1}{2}(c+dx)\right)\right)}{\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)}\right)}{6d}$$

Antiderivative was successfully verified.

[In] Integrate[(3 + 5*Csc[c + d*x])^(-1), x]

[Out] (2*(c + d*x) - 5*ArcTan[(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])])/(6*d)

Maple [A] time = 0.039, size = 36, normalized size = 1.2

$$\frac{2}{3d} \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \frac{5}{6d} \arctan\left(\frac{5}{4} \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(3+5*csc(d*x+c)), x)

[Out] 2/3/d*arctan(tan(1/2*d*x+1/2*c))-5/6/d*arctan(5/4*tan(1/2*d*x+1/2*c)+3/4)

Maxima [A] time = 1.47446, size = 66, normalized size = 2.13

$$\frac{5 \arctan\left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} + \frac{3}{4}\right) - 4 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{6d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3+5*csc(d*x+c)),x, algorithm="maxima")

[Out] $-1/6*(5*\arctan(5/4*\sin(d*x + c)/(\cos(d*x + c) + 1) + 3/4) - 4*\arctan(\sin(d*x + c)/(\cos(d*x + c) + 1)))/d$

Fricas [A] time = 0.492249, size = 89, normalized size = 2.87

$$\frac{4 dx - 5 \arctan\left(\frac{5 \sin(dx+c)+3}{4 \cos(dx+c)}\right)}{12 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3+5*csc(d*x+c)),x, algorithm="fricas")

[Out] $1/12*(4*d*x - 5*\arctan(1/4*(5*\sin(d*x + c) + 3)/\cos(d*x + c)))/d$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{5 \csc(c + dx) + 3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3+5*csc(d*x+c)),x)

[Out] Integral(1/(5*csc(c + d*x) + 3), x)

Giac [A] time = 1.44328, size = 66, normalized size = 2.13

$$\frac{dx + c + 10 \arctan\left(-\frac{3 \cos(dx+c)+\sin(dx+c)+3}{\cos(dx+c)-3 \sin(dx+c)-9}\right)}{12 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3+5*csc(d*x+c)),x, algorithm="giac")

[Out] $-1/12*(d*x + c + 10*\arctan(-(3*\cos(d*x + c) + \sin(d*x + c) + 3)/(\cos(d*x + c) - 3*\sin(d*x + c) - 9)))/d$

3.53 $\int \frac{1}{5+3 \csc(c+dx)} dx$

Optimal. Leaf size=68

$$\frac{3 \log\left(\sin\left(\frac{1}{2}(c+dx)\right) + 3 \cos\left(\frac{1}{2}(c+dx)\right)\right)}{20d} - \frac{3 \log\left(3 \sin\left(\frac{1}{2}(c+dx)\right) + \cos\left(\frac{1}{2}(c+dx)\right)\right)}{20d} + \frac{x}{5}$$

[Out] x/5 + (3*Log[3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/(20*d) - (3*Log[Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2]])/(20*d)

Rubi [A] time = 0.0390231, antiderivative size = 68, normalized size of antiderivative = 1., number of steps used = 5, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3783, 2660, 616, 31}

$$\frac{3 \log\left(\sin\left(\frac{1}{2}(c+dx)\right) + 3 \cos\left(\frac{1}{2}(c+dx)\right)\right)}{20d} - \frac{3 \log\left(3 \sin\left(\frac{1}{2}(c+dx)\right) + \cos\left(\frac{1}{2}(c+dx)\right)\right)}{20d} + \frac{x}{5}$$

Antiderivative was successfully verified.

[In] Int[(5 + 3*Csc[c + d*x])^(-1), x]

[Out] x/5 + (3*Log[3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/(20*d) - (3*Log[Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2]])/(20*d)

Rule 3783

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(-1), x_Symbol] := Simp[x/a, x] - Dist[1/a, Int[1/(1 + (a*Sin[c + d*x])/b), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 2660

Int[((a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 616

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[c/q, Int[1/Simp[b/2 - q/2 + c*x, x], x], x] - Dist[c/q,

`Int[1/Simp[b/2 + q/2 + c*x, x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[b^2 - 4*a*c] && PerfectSquareQ[b^2 - 4*a*c]`

Rule 31

`Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]`

Rubi steps

$$\begin{aligned} \int \frac{1}{5 + 3 \csc(c + dx)} dx &= \frac{x}{5} - \frac{1}{5} \int \frac{1}{1 + \frac{5}{3} \sin(c + dx)} dx \\ &= \frac{x}{5} - \frac{2 \operatorname{Subst}\left(\int \frac{1}{1 + \frac{10x}{3} + x^2} dx, x, \tan\left(\frac{1}{2}(c + dx)\right)\right)}{5d} \\ &= \frac{x}{5} - \frac{3 \operatorname{Subst}\left(\int \frac{1}{\frac{1}{3} + x} dx, x, \tan\left(\frac{1}{2}(c + dx)\right)\right)}{20d} + \frac{3 \operatorname{Subst}\left(\int \frac{1}{3+x} dx, x, \tan\left(\frac{1}{2}(c + dx)\right)\right)}{20d} \\ &= \frac{x}{5} + \frac{3 \log\left(3 + \tan\left(\frac{1}{2}(c + dx)\right)\right)}{20d} - \frac{3 \log\left(1 + 3 \tan\left(\frac{1}{2}(c + dx)\right)\right)}{20d} \end{aligned}$$

Mathematica [A] time = 0.0468918, size = 67, normalized size = 0.99

$$\frac{4(c + dx) + 3 \log\left(\sin\left(\frac{1}{2}(c + dx)\right) + 3 \cos\left(\frac{1}{2}(c + dx)\right)\right) - 3 \log\left(3 \sin\left(\frac{1}{2}(c + dx)\right) + \cos\left(\frac{1}{2}(c + dx)\right)\right)}{20d}$$

Antiderivative was successfully verified.

[In] `Integrate[(5 + 3*Csc[c + d*x])^(-1), x]`

[Out] `(4*(c + d*x) + 3*Log[3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] - 3*Log[Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2]])/(20*d)`

Maple [A] time = 0.045, size = 53, normalized size = 0.8

$$\frac{2}{5d} \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \frac{3}{20d} \ln(3 \tan(1/2 dx + c/2) + 1) + \frac{3}{20d} \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(5+3*csc(d*x+c)),x)`

[Out] $2/5/d*\arctan(\tan(1/2*d*x+1/2*c))-3/20/d*\ln(3*\tan(1/2*d*x+1/2*c)+1)+3/20/d*\ln(\tan(1/2*d*x+1/2*c)+3)$

Maxima [A] time = 1.46285, size = 96, normalized size = 1.41

$$\frac{8 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right) - 3 \log\left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 1\right) + 3 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 3\right)}{20 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(5+3*csc(d*x+c)),x, algorithm="maxima")`

[Out] $1/20*(8*\arctan(\sin(d*x + c)/(\cos(d*x + c) + 1)) - 3*\log(3*\sin(d*x + c)/(\cos(d*x + c) + 1) + 1) + 3*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) + 3))/d$

Fricas [A] time = 0.496222, size = 144, normalized size = 2.12

$$\frac{8 dx + 3 \log(4 \cos(dx + c) + 3 \sin(dx + c) + 5) - 3 \log(-4 \cos(dx + c) + 3 \sin(dx + c) + 5)}{40 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(5+3*csc(d*x+c)),x, algorithm="fricas")`

[Out] $1/40*(8*d*x + 3*\log(4*\cos(d*x + c) + 3*\sin(d*x + c) + 5) - 3*\log(-4*\cos(d*x + c) + 3*\sin(d*x + c) + 5))/d$

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int \frac{1}{3 \csc(c + dx) + 5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(5+3*csc(d*x+c)),x)

[Out] Integral(1/(3*csc(c + d*x) + 5), x)

Giac [A] time = 1.37382, size = 61, normalized size = 0.9

$$\frac{4dx + 4c - 3 \log\left(\left|3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) + 3 \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 3\right|\right)}{20d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(5+3*csc(d*x+c)),x, algorithm="giac")

[Out] 1/20*(4*d*x + 4*c - 3*log(abs(3*tan(1/2*d*x + 1/2*c) + 1)) + 3*log(abs(tan(1/2*d*x + 1/2*c) + 3)))/d

3.54 $\int \csc^3(e + fx)(a + b \csc(e + fx))^m dx$

Optimal. Leaf size=274

$$\frac{\sqrt{2}(a^2 + b^2(m+1)) \cot(e + fx)(a + b \csc(e + fx))^m \left(\frac{a+b \csc(e+fx)}{a+b}\right)^{-m} F_1\left(\frac{1}{2}; \frac{1}{2}, -m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx)), \frac{b(1-\csc(e+fx))}{a+b}\right)}{b^2 f(m+2) \sqrt{\csc(e + fx) + 1}}$$

[Out] -((Cot[e + f*x]*(a + b*Csc[e + f*x])^(1 + m))/(b*f*(2 + m))) + (Sqrt[2]*a*(a + b)*AppellF1[1/2, 1/2, -1 - m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b^2*f*(2 + m)*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m) - (Sqrt[2]*(a^2 + b^2*(1 + m))*AppellF1[1/2, 1/2, -m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b^2*f*(2 + m)*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m)

Rubi [A] time = 0.34266, antiderivative size = 274, normalized size of antiderivative = 1., number of steps used = 8, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3840, 4007, 3834, 139, 138}

$$\frac{\sqrt{2}(a^2 + b^2(m+1)) \cot(e + fx)(a + b \csc(e + fx))^m \left(\frac{a+b \csc(e+fx)}{a+b}\right)^{-m} F_1\left(\frac{1}{2}; \frac{1}{2}, -m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx)), \frac{b(1-\csc(e+fx))}{a+b}\right)}{b^2 f(m+2) \sqrt{\csc(e + fx) + 1}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^3*(a + b*Csc[e + f*x])^m,x]

[Out] -((Cot[e + f*x]*(a + b*Csc[e + f*x])^(1 + m))/(b*f*(2 + m))) + (Sqrt[2]*a*(a + b)*AppellF1[1/2, 1/2, -1 - m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b^2*f*(2 + m)*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m) - (Sqrt[2]*(a^2 + b^2*(1 + m))*AppellF1[1/2, 1/2, -m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b^2*f*(2 + m)*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m)

Rule 3840

Int[csc[(e_.) + (f_.)*(x_)]^3*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_), x_Symbol] := -Simp[(Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1))/(b*f*(m + 2)), x] + Dist[1/(b*(m + 2)), Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(b*(m + 1) - a*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f, m}, x] && NeQ[a^2 - b^2

, 0] && !LtQ[m, -1]

Rule 4007

Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] :> Dist[(A*b - a*B)/b, Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m, x], x] + Dist[B/b, Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, A, B, e, f, m}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0]

Rule 3834

Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] :> Dist[Cot[e + f*x]/(f*Sqrt[1 + Csc[e + f*x]]*Sqrt[1 - Csc[e + f*x]]), Subst[Int[(a + b*x)^m/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*m]

Rule 139

Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_Symbol] :> Dist[(e + f*x)^FracPart[p]/((b/(b*e - a*f))^IntPart[p]*((b*(e + f*x))/(b*e - a*f))^FracPart[p]), Int[(a + b*x)^m*(c + d*x)^n*((b*e)/(b*e - a*f) + (b*f*x)/(b*e - a*f))^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[b/(b*c - a*d), 0] && !GtQ[b/(b*e - a*f), 0]

Rule 138

Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_Symbol] :> Simp[((a + b*x)^(m + 1)*AppellF1[m + 1, -n, -p, m + 2, -((d*(a + b*x))/(b*c - a*d)), -((f*(a + b*x))/(b*e - a*f))]/(b*(m + 1)*(b/(b*c - a*d))^n*(b/(b*e - a*f))^p), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[b/(b*c - a*d), 0] && GtQ[b/(b*e - a*f), 0] && !(GtQ[d/(d*a - c*b), 0] && GtQ[d/(d*e - c*f), 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[f/(f*a - e*b), 0] && GtQ[f/(f*c - e*d), 0] && SimplerQ[e + f*x, a + b*x])

Rubi steps

$$\begin{aligned}
\int \csc^3(e+fx)(a+b \csc(e+fx))^m dx &= -\frac{\cot(e+fx)(a+b \csc(e+fx))^{1+m}}{bf(2+m)} + \frac{\int \csc(e+fx)(b(1+m) - a \csc(e+fx))}{b(2+m)} \\
&= -\frac{\cot(e+fx)(a+b \csc(e+fx))^{1+m}}{bf(2+m)} - \frac{a \int \csc(e+fx)(a+b \csc(e+fx))^{1+m} dx}{b^2(2+m)} \\
&= -\frac{\cot(e+fx)(a+b \csc(e+fx))^{1+m}}{bf(2+m)} - \frac{(a \cot(e+fx)) \text{Subst}\left(\int \frac{(a+bx)^{1+m}}{\sqrt{1-x}\sqrt{1+x}} dx, x, \frac{a+b \csc(e+fx)}{b}\right)}{b^2 f(2+m) \sqrt{1-\csc(e+fx)} \sqrt{1+\csc(e+fx)}} \\
&= -\frac{\cot(e+fx)(a+b \csc(e+fx))^{1+m}}{bf(2+m)} + \frac{\left(a(-a-b) \cot(e+fx)(a+b \csc(e+fx))\right)}{b^2 f(2+m)} \\
&= -\frac{\cot(e+fx)(a+b \csc(e+fx))^{1+m}}{bf(2+m)} + \frac{\sqrt{2}a(a+b)F_1\left(\frac{1}{2}; \frac{1}{2}, -1-m; \frac{3}{2}; \frac{1}{2}(1-\csc(e+fx))\right)}{b^2 f(2+m)}
\end{aligned}$$

Mathematica [F] time = 4.181, size = 0, normalized size = 0.

$$\int \csc^3(e+fx)(a+b \csc(e+fx))^m dx$$

Verification is Not applicable to the result.

[In] Integrate[Csc[e + f*x]^3*(a + b*Csc[e + f*x])^m,x]

[Out] Integrate[Csc[e + f*x]^3*(a + b*Csc[e + f*x])^m, x]

Maple [F] time = 0.309, size = 0, normalized size = 0.

$$\int (\csc(fx+e))^3 (a+b \csc(fx+e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^3*(a+b*csc(f*x+e))^m,x)

[Out] int(csc(f*x+e)^3*(a+b*csc(f*x+e))^m,x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \csc(fx + e)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(a+b*csc(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e) + a)^m*csc(f*x + e)^3, x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(b \csc(fx + e) + a\right)^m \csc(fx + e)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(a+b*csc(f*x+e))^m,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e) + a)^m*csc(f*x + e)^3, x)

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a + b \csc(e + fx))^m \csc^3(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**3*(a+b*csc(f*x+e))**m,x)

[Out] Integral((a + b*csc(e + f*x))**m*csc(e + f*x)**3, x)

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \csc(fx + e)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^3*(a+b*csc(f*x+e))^m,x, algorithm="giac")
```

```
[Out] integrate((b*csc(f*x + e) + a)^m*csc(f*x + e)^3, x)
```

3.55 $\int \csc^2(e + fx)(a + b \csc(e + fx))^m dx$

Optimal. Leaf size=220

$$\frac{\sqrt{2}a \cot(e + fx)(a + b \csc(e + fx))^m \left(\frac{a+b \csc(e+fx)}{a+b}\right)^{-m} F_1\left(\frac{1}{2}; \frac{1}{2}, -m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx)), \frac{b(1-\csc(e+fx))}{a+b}\right)}{bf\sqrt{\csc(e + fx) + 1}} - \frac{\sqrt{2}(a + b) \cot(e + fx)}{bf\sqrt{\csc(e + fx) + 1}}$$

```
[Out] -((Sqrt[2]*(a + b)*AppellF1[1/2, 1/2, -1 - m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b*f*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m) + (Sqrt[2]*a*AppellF1[1/2, 1/2, -m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b*f*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m)
```

Rubi [A] time = 0.225604, antiderivative size = 220, normalized size of antiderivative = 1., number of steps used = 7, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.19$, Rules used = {3838, 3834, 139, 138}

$$\frac{\sqrt{2}a \cot(e + fx)(a + b \csc(e + fx))^m \left(\frac{a+b \csc(e+fx)}{a+b}\right)^{-m} F_1\left(\frac{1}{2}; \frac{1}{2}, -m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx)), \frac{b(1-\csc(e+fx))}{a+b}\right)}{bf\sqrt{\csc(e + fx) + 1}} - \frac{\sqrt{2}(a + b) \cot(e + fx)}{bf\sqrt{\csc(e + fx) + 1}}$$

Antiderivative was successfully verified.

```
[In] Int[Csc[e + f*x]^2*(a + b*Csc[e + f*x])^m,x]
```

```
[Out] -((Sqrt[2]*(a + b)*AppellF1[1/2, 1/2, -1 - m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b*f*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m) + (Sqrt[2]*a*AppellF1[1/2, 1/2, -m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(b*f*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m)
```

Rule 3838

```
Int[csc[(e_.) + (f_.)*(x_)]^2*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_, x_Symbol] :> -Dist[a/b, Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m, x], x] + Dist[1/b, Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f, m}, x] && NeQ[a^2 - b^2, 0]
```

Rule 3834

```
Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_), x_
Symbol] := Dist[Cot[e + f*x]/(f*Sqrt[1 + Csc[e + f*x]]*Sqrt[1 - Csc[e + f*x
]]), Subst[Int[(a + b*x)^m/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Csc[e + f*x]],
x] /; FreeQ[{a, b, e, f, m}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*m]
```

Rule 139

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_Symbol] := Dist[(e + f*x)^FracPart[p]/((b/(b*e - a*f))^IntPart[p]*
((b*(e + f*x))/(b*e - a*f))^FracPart[p]), Int[(a + b*x)^m*(c + d*x)^n*((b*e
)/(b*e - a*f) + (b*f*x)/(b*e - a*f))^p, x], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[b/(b
*c - a*d), 0] && !GtQ[b/(b*e - a*f), 0]
```

Rule 138

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_Symbol] := Simp[((a + b*x)^(m + 1)*AppellF1[m + 1, -n, -p, m + 2,
-((d*(a + b*x))/(b*c - a*d)), -((f*(a + b*x))/(b*e - a*f))]/(b*(m + 1)*(b/
(b*c - a*d))^n*(b/(b*e - a*f))^p), x] /; FreeQ[{a, b, c, d, e, f, m, n, p},
x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[b/(b*c - a*d)
, 0] && GtQ[b/(b*e - a*f), 0] && !(GtQ[d/(d*a - c*b), 0] && GtQ[d/(d*e - c
*f), 0] && SimplrQ[c + d*x, a + b*x]) && !(GtQ[f/(f*a - e*b), 0] && GtQ[f
/(f*c - e*d), 0] && SimplrQ[e + f*x, a + b*x])
```

Rubi steps

$$\begin{aligned} \int \csc^2(e + fx)(a + b \csc(e + fx))^m dx &= \frac{\int \csc(e + fx)(a + b \csc(e + fx))^{1+m} dx}{b} - \frac{a \int \csc(e + fx)(a + b \csc(e + fx))^m dx}{b} \\ &= \frac{\cot(e + fx) \operatorname{Subst}\left(\int \frac{(a+bx)^{1+m}}{\sqrt{1-x}\sqrt{1+x}} dx, x, \csc(e + fx)\right)}{bf\sqrt{1 - \csc(e + fx)}\sqrt{1 + \csc(e + fx)}} - \frac{(a \cot(e + fx)) \operatorname{Subst}\left(\int \frac{(a+bx)^{1+m}}{\sqrt{1-x}\sqrt{1+x}} dx, x, \csc(e + fx)\right)}{bf\sqrt{1 - \csc(e + fx)}\sqrt{1 + \csc(e + fx)}} \\ &= -\frac{\left(a \cot(e + fx)(a + b \csc(e + fx))^m \left(-\frac{a+b \csc(e+fx)}{-a-b}\right)^{-m}\right) \operatorname{Subst}\left(\int \frac{\left(-\frac{a}{-a-b} - \frac{bx}{-a-b}\right)^m}{\sqrt{1-x}\sqrt{1+x}} dx, x, \csc(e + fx)\right)}{bf\sqrt{1 - \csc(e + fx)}\sqrt{1 + \csc(e + fx)}} \\ &= -\frac{\sqrt{2}(a + b)F_1\left(\frac{1}{2}; \frac{1}{2}, -1 - m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx)), \frac{b(1 - \csc(e+fx))}{a+b}\right) \cot(e + fx)(a + b \csc(e + fx))^m}{bf\sqrt{1 + \csc(e + fx)}} \end{aligned}$$

Mathematica [F] time = 2.71815, size = 0, normalized size = 0.

$$\int \csc^2(e + fx)(a + b \csc(e + fx))^m dx$$

Verification is Not applicable to the result.

[In] Integrate[Csc[e + f*x]^2*(a + b*Csc[e + f*x])^m,x]

[Out] Integrate[Csc[e + f*x]^2*(a + b*Csc[e + f*x])^m, x]

Maple [F] time = 0.247, size = 0, normalized size = 0.

$$\int (\csc(fx + e))^2 (a + b \csc(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^2*(a+b*csc(f*x+e))^m,x)

[Out] int(csc(f*x+e)^2*(a+b*csc(f*x+e))^m,x)

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \csc(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(a+b*csc(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e) + a)^m*csc(f*x + e)^2, x)

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(b \csc(fx + e) + a\right)^m \csc(fx + e)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2*(a+b*csc(f*x+e))^m,x, algorithm="fricas")
```

```
[Out] integral((b*csc(f*x + e) + a)^m*csc(f*x + e)^2, x)
```

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a + b \csc(e + fx))^m \csc^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)**2*(a+b*csc(f*x+e))**m,x)
```

```
[Out] Integral((a + b*csc(e + f*x))**m*csc(e + f*x)**2, x)
```

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \csc(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2*(a+b*csc(f*x+e))^m,x, algorithm="giac")
```

```
[Out] integrate((b*csc(f*x + e) + a)^m*csc(f*x + e)^2, x)
```

3.56 $\int \csc(e + fx)(a + b \csc(e + fx))^m dx$

Optimal. Leaf size=104

$$\frac{\sqrt{2} \cot(e + fx)(a + b \csc(e + fx))^m \left(\frac{a+b \csc(e+fx)}{a+b} \right)^{-m} F_1 \left(\frac{1}{2}; \frac{1}{2}, -m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx)), \frac{b(1-\csc(e+fx))}{a+b} \right)}{f \sqrt{\csc(e + fx) + 1}}$$

[Out] -((Sqrt[2]*AppellF1[1/2, 1/2, -m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(f*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m))

Rubi [A] time = 0.0738236, antiderivative size = 104, normalized size of antiderivative = 1., number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3834, 139, 138}

$$\frac{\sqrt{2} \cot(e + fx)(a + b \csc(e + fx))^m \left(\frac{a+b \csc(e+fx)}{a+b} \right)^{-m} F_1 \left(\frac{1}{2}; \frac{1}{2}, -m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx)), \frac{b(1-\csc(e+fx))}{a+b} \right)}{f \sqrt{\csc(e + fx) + 1}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m,x]

[Out] -((Sqrt[2]*AppellF1[1/2, 1/2, -m, 3/2, (1 - Csc[e + f*x])/2, (b*(1 - Csc[e + f*x]))/(a + b)]*Cot[e + f*x]*(a + b*Csc[e + f*x])^m)/(f*Sqrt[1 + Csc[e + f*x]]*((a + b*Csc[e + f*x])/(a + b))^m))

Rule 3834

Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_), x_Symbol] :> Dist[Cot[e + f*x]/(f*Sqrt[1 + Csc[e + f*x]]*Sqrt[1 - Csc[e + f*x]]), Subst[Int[(a + b*x)^m/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*m]

Rule 139

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] :> Dist[(e + f*x)^FracPart[p]/((b/(b*e - a*f))^IntPart[p]*((b*(e + f*x))/(b*e - a*f))^FracPart[p]), Int[(a + b*x)^m*(c + d*x)^n*((b*e)/(b*e - a*f) + (b*f*x)/(b*e - a*f))^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[b/(b

*c - a*d), 0] && !GtQ[b/(b*e - a*f), 0]

Rule 138

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_Symbol] :> Simp[((a + b*x)^(m + 1)*AppellF1[m + 1, -n, -p, m + 2,
-((d*(a + b*x))/(b*c - a*d)), -((f*(a + b*x))/(b*e - a*f))]/(b*(m + 1)*(b/
(b*c - a*d))^n*(b/(b*e - a*f))^p), x] /; FreeQ[{a, b, c, d, e, f, m, n, p},
x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[b/(b*c - a*d)
, 0] && GtQ[b/(b*e - a*f), 0] && !(GtQ[d/(d*a - c*b), 0] && GtQ[d/(d*e - c
*f), 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[f/(f*a - e*b), 0] && GtQ[f
/(f*c - e*d), 0] && SimplerQ[e + f*x, a + b*x])
```

Rubi steps

$$\begin{aligned} \int \csc(e + fx)(a + b \csc(e + fx))^m dx &= \frac{\cot(e + fx) \operatorname{Subst}\left(\int \frac{(a+bx)^m}{\sqrt{1-x}\sqrt{1+x}} dx, x, \csc(e + fx)\right)}{f\sqrt{1 - \csc(e + fx)}\sqrt{1 + \csc(e + fx)}} \\ &= \frac{\left(\cot(e + fx)(a + b \csc(e + fx))^m \left(-\frac{a+b \csc(e+fx)}{-a-b}\right)^{-m}\right) \operatorname{Subst}\left(\int \frac{\left(-\frac{a}{-a-b} - \frac{bx}{-a-b}\right)^m}{\sqrt{1-x}\sqrt{1+x}} dx, x\right)}{f\sqrt{1 - \csc(e + fx)}\sqrt{1 + \csc(e + fx)}} \\ &= -\frac{\sqrt{2}F_1\left(\frac{1}{2}; \frac{1}{2}, -m; \frac{3}{2}; \frac{1}{2}(1 - \csc(e + fx)), \frac{b(1 - \csc(e+fx))}{a+b}\right) \cot(e + fx)(a + b \csc(e + fx))^m}{f\sqrt{1 + \csc(e + fx)}} \end{aligned}$$

Mathematica [F] time = 1.9129, size = 0, normalized size = 0.

$$\int \csc(e + fx)(a + b \csc(e + fx))^m dx$$

Verification is Not applicable to the result.

[In] Integrate[Csc[e + f*x]*(a + b*Csc[e + f*x])^m, x]

[Out] Integrate[Csc[e + f*x]*(a + b*Csc[e + f*x])^m, x]

Maple [F] time = 0.511, size = 0, normalized size = 0.

$$\int \csc(fx + e)(a + b \csc(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)*(a+b*csc(f*x+e))^m,x)`

[Out] `int(csc(f*x+e)*(a+b*csc(f*x+e))^m,x)`

Maxima [F] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \csc(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(a+b*csc(f*x+e))^m,x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e) + a)^m*csc(f*x + e), x)`

Fricas [F] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(b \csc(fx + e) + a\right)^m \csc(fx + e), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(a+b*csc(f*x+e))^m,x, algorithm="fricas")`

[Out] `integral((b*csc(f*x + e) + a)^m*csc(f*x + e), x)`

Sympy [F] time = 0., size = 0, normalized size = 0.

$$\int (a + b \csc(e + fx))^m \csc(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(a+b*csc(f*x+e))**m,x)`

[Out] `Integral((a + b*csc(e + f*x))**m*csc(e + f*x), x)`

Giac [F] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \csc(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(a+b*csc(f*x+e))^m,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e) + a)^m*csc(f*x + e), x)

$$3.57 \quad \int (a + b \csc(e + fx))^m dx$$

Optimal. Leaf size=14

Unintegrable $((a + b \csc(e + fx))^m, x)$

[Out] Unintegrable[(a + b*Csc[e + f*x])^m, x]

Rubi [A] time = 0.0095191, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.$, Rules used = {}

$$\int (a + b \csc(e + fx))^m dx$$

Verification is Not applicable to the result.

[In] Int[(a + b*Csc[e + f*x])^m,x]

[Out] Defer[Int] [(a + b*Csc[e + f*x])^m, x]

Rubi steps

$$\int (a + b \csc(e + fx))^m dx = \int (a + b \csc(e + fx))^m dx$$

Mathematica [A] time = 1.54628, size = 0, normalized size = 0.

$$\int (a + b \csc(e + fx))^m dx$$

Verification is Not applicable to the result.

[In] Integrate[(a + b*Csc[e + f*x])^m,x]

[Out] Integrate[(a + b*Csc[e + f*x])^m, x]

Maple [A] time = 0.333, size = 0, normalized size = 0.

$$\int (a + b \csc(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+b*csc(f*x+e))^m,x)

[Out] int((a+b*csc(f*x+e))^m,x)

Maxima [A] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e) + a)^m, x)

Fricas [A] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(b \csc(fx + e) + a\right)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(f*x+e))^m,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e) + a)^m, x)

Sympy [A] time = 0., size = 0, normalized size = 0.

$$\int (a + b \csc(e + fx))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.


```
[In] integrate((a+b*csc(f*x+e))**m,x)
```

```
[Out] Integral((a + b*csc(e + f*x))**m, x)
```

Giac [A] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*csc(f*x+e))^m,x, algorithm="giac")
```

```
[Out] integrate((b*csc(f*x + e) + a)^m, x)
```

$$3.58 \quad \int (a + b \csc(e + fx))^m \sin(e + fx) dx$$

Optimal. Leaf size=21

Unintegrable $(\sin(e + fx)(a + b \csc(e + fx))^m, x)$

[Out] Unintegrable $[(a + b \cdot \text{Csc}[e + f \cdot x])^m \cdot \text{Sin}[e + f \cdot x], x]$

Rubi [A] time = 0.0318065, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.$, Rules used = {}

$$\int (a + b \csc(e + fx))^m \sin(e + fx) dx$$

Verification is Not applicable to the result.

[In] Int $[(a + b \cdot \text{Csc}[e + f \cdot x])^m \cdot \text{Sin}[e + f \cdot x], x]$

[Out] Defer[Int] $[(a + b \cdot \text{Csc}[e + f \cdot x])^m \cdot \text{Sin}[e + f \cdot x], x]$

Rubi steps

$$\int (a + b \csc(e + fx))^m \sin(e + fx) dx = \int (a + b \csc(e + fx))^m \sin(e + fx) dx$$

Mathematica [A] time = 6.05511, size = 0, normalized size = 0.

$$\int (a + b \csc(e + fx))^m \sin(e + fx) dx$$

Verification is Not applicable to the result.

[In] Integrate $[(a + b \cdot \text{Csc}[e + f \cdot x])^m \cdot \text{Sin}[e + f \cdot x], x]$

[Out] Integrate $[(a + b \cdot \text{Csc}[e + f \cdot x])^m \cdot \text{Sin}[e + f \cdot x], x]$

Maple [A] time = 0.257, size = 0, normalized size = 0.

$$\int (a + b \csc(fx + e))^m \sin(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+b*csc(f*x+e))^m*sin(f*x+e),x)

[Out] int((a+b*csc(f*x+e))^m*sin(f*x+e),x)

Maxima [A] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \sin(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(f*x+e))^m*sin(f*x+e),x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e) + a)^m*sin(f*x + e), x)

Fricas [A] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(\left(b \csc(fx + e) + a\right)^m \sin(fx + e), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(f*x+e))^m*sin(f*x+e),x, algorithm="fricas")

[Out] integral((b*csc(f*x + e) + a)^m*sin(f*x + e), x)

Sympy [A] time = 0., size = 0, normalized size = 0.

$$\int (a + b \csc(e + fx))^m \sin(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*csc(f*x+e))*m*sin(f*x+e),x)
```

```
[Out] Integral((a + b*csc(e + f*x))*m*sin(e + f*x), x)
```

Giac [A] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \sin(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*csc(f*x+e))^m*sin(f*x+e),x, algorithm="giac")
```

```
[Out] integrate((b*csc(f*x + e) + a)^m*sin(f*x + e), x)
```

$$3.59 \quad \int (a + b \csc(e + fx))^m \sin^2(e + fx) dx$$

Optimal. Leaf size=23

$$\text{Unintegrable}(\sin^2(e + fx)(a + b \csc(e + fx))^m, x)$$

[Out] Unintegrable[(a + b*Csc[e + f*x])^m*Sin[e + f*x]^2, x]

Rubi [A] time = 0.040237, antiderivative size = 0, normalized size of antiderivative = 0., number of steps used = 0, number of rules used = 0, integrand size = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.$, Rules used = {}

$$\int (a + b \csc(e + fx))^m \sin^2(e + fx) dx$$

Verification is Not applicable to the result.

[In] Int[(a + b*Csc[e + f*x])^m*Sin[e + f*x]^2,x]

[Out] Defer[Int] [(a + b*Csc[e + f*x])^m*Sin[e + f*x]^2, x]

Rubi steps

$$\int (a + b \csc(e + fx))^m \sin^2(e + fx) dx = \int (a + b \csc(e + fx))^m \sin^2(e + fx) dx$$

Mathematica [A] time = 5.90403, size = 0, normalized size = 0.

$$\int (a + b \csc(e + fx))^m \sin^2(e + fx) dx$$

Verification is Not applicable to the result.

[In] Integrate[(a + b*Csc[e + f*x])^m*Sin[e + f*x]^2,x]

[Out] Integrate[(a + b*Csc[e + f*x])^m*Sin[e + f*x]^2, x]

Maple [A] time = 0.567, size = 0, normalized size = 0.

$$\int (a + b \csc(fx + e))^m (\sin(fx + e))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+b*csc(f*x+e))^m*sin(f*x+e)^2,x)

[Out] int((a+b*csc(f*x+e))^m*sin(f*x+e)^2,x)

Maxima [A] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \sin(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(f*x+e))^m*sin(f*x+e)^2,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e) + a)^m*sin(f*x + e)^2, x)

Fricas [A] time = 0., size = 0, normalized size = 0.

$$\text{integral}\left(-\left(\cos(fx + e)\right)^2 - 1\right)\left(b \csc(fx + e) + a\right)^m, x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*csc(f*x+e))^m*sin(f*x+e)^2,x, algorithm="fricas")

[Out] integral(-(cos(f*x + e)^2 - 1)*(b*csc(f*x + e) + a)^m, x)

Sympy [F(-1)] time = 0., size = 0, normalized size = 0.

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*csc(f*x+e))^m*sin(f*x+e)**2,x)
```

```
[Out] Timed out
```

Giac [A] time = 0., size = 0, normalized size = 0.

$$\int (b \csc(fx + e) + a)^m \sin(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*csc(f*x+e))^m*sin(f*x+e)^2,x, algorithm="giac")
```

```
[Out] integrate((b*csc(f*x + e) + a)^m*sin(f*x + e)^2, x)
```


Chapter 4

Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.0.1 Mathematica and Rubi grading function

```
1 (* Original version thanks to Albert Rich emailed on 03/21/2017 *)
2 (* ::Package:: *)
3
4 (* ::Subsection:: *)
5 (*GradeAntiderivative[result,optimal]*)
6
7
8 (* ::Text:: *)
9 (*If result and optimal are mathematical expressions, *)
10 (*      GradeAntiderivative[result,optimal] returns*)
11 (* "F" if the result fails to integrate an expression that*)
12 (*      is integrable*)
13 (* "C" if result involves higher level functions than necessary*)
14 (* "B" if result is more than twice the size of the optimal*)
15 (*      antiderivative*)
16 (* "A" if result can be considered optimal*)
17
18
19 GradeAntiderivative[result_,optimal_] :=
20   If[ExpnType[result]<=ExpnType[optimal],
21     If[FreeQ[result,Complex] || Not[FreeQ[optimal,Complex]],
```

```

22     If[LeafCount[result]<=2*LeafCount[optimal],
23         "A",
24         "B"],
25     "C"],
26 If[FreeQ[result,Integrate] && FreeQ[result,Int],
27     "C",
28     "F"]]
29
30
31 (* ::Text:: *)
32 (*The following summarizes the type number assigned an *)
33 (*expression based on the functions it involves*)
34 (*1 = rational function*)
35 (*2 = algebraic function*)
36 (*3 = elementary function*)
37 (*4 = special function*)
38 (*5 = hyperpergeometric function*)
39 (*6 = appell function*)
40 (*7 = rootsum function*)
41 (*8 = integrate function*)
42 (*9 = unknown function*)
43
44
45 ExpnType[expn_] :=
46     If[AtomQ[expn],
47         1,
48     If[ListQ[expn],
49         Max[Map[ExpnType,expn]],
50     If[Head[expn]===Power,
51         If[IntegerQ[expn[[2]]],
52             ExpnType[expn[[1]]],
53         If[Head[expn[[2]]]===Rational,
54             If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
55                 1,
56                 Max[ExpnType[expn[[1]],2]],
57             Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
58     If[Head[expn]===Plus || Head[expn]===Times,
59         Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
60     If[ElementaryFunctionQ[Head[expn]],
61         Max[3,ExpnType[expn[[1]]]],
62     If[SpecialFunctionQ[Head[expn]],
63         Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
64     If[HypergeometricFunctionQ[Head[expn]],
65         Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
66     If[AppellFunctionQ[Head[expn]],
67         Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
68     If[Head[expn]===RootSum,

```

```

69   Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
70   If[Head[expn]===Integrate || Head[expn]===Int,
71     Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
72   9]]]]]]]]]]
73
74
75 ElementaryFunctionQ[func_] :=
76   MemberQ[{
77     Exp, Log,
78     Sin, Cos, Tan, Cot, Sec, Csc,
79     ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
80     Sinh, Cosh, Tanh, Coth, Sech, Csch,
81     ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
82   }, func]
83
84
85 SpecialFunctionQ[func_] :=
86   MemberQ[{
87     Erf, Erfc, Erfi,
88     FresnelS, FresnelC,
89     ExpIntegralE, ExpIntegralEi, LogIntegral,
90     SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
91     Gamma, LogGamma, PolyGamma,
92     Zeta, PolyLog, ProductLog,
93     EllipticF, EllipticE, EllipticPi
94   }, func]
95
96
97 HypergeometricFunctionQ[func_] :=
98   MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
99
100
101 AppellFunctionQ[func_] :=
102   MemberQ[{AppellF1}, func]

```

4.0.2 Maple grading function

```

1 # File: GradeAntiderivative.mpl
2 # Original version thanks to Albert Rich emailed on 03/21/2017
3
4 #Nasser 03/22/2017 Use Maple leaf count instead since buildin
5 #Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
6 #Nasser 03/24/2017 corrected the check for complex result
7 #Nasser 10/27/2017 check for leafsize and do not call ExpnType()
8 #
9 #Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
10 #
11 see problem 156, file Apostol_Problems

```

```

11
12 GradeAntiderivative := proc(result,optimal)
13 local leaf_count_result, leaf_count_optimal,ExpnType_result,ExpnType_optimal,
    debug:=false;
14
15     leaf_count_result:=leafcount(result);
16     #do NOT call ExpnType() if leaf size is too large. Recursion problem
17     if leaf_count_result > 500000 then
18         return "B";
19     fi;
20
21     leaf_count_optimal:=leafcount(optimal);
22
23     ExpnType_result:=ExpnType(result);
24     ExpnType_optimal:=ExpnType(optimal);
25
26     if debug then
27         print("ExpnType_result",ExpnType_result," ExpnType_optimal=",
    ExpnType_optimal);
28     fi;
29
30 # If result and optimal are mathematical expressions,
31 # GradeAntiderivative[result,optimal] returns
32 #   "F" if the result fails to integrate an expression that
33 #     is integrable
34 #   "C" if result involves higher level functions than necessary
35 #   "B" if result is more than twice the size of the optimal
36 #     antiderivative
37 #   "A" if result can be considered optimal
38
39 #This check below actually is not needed, since I only
40 #call this grading only for passed integrals. i.e. I check
41 #for "F" before calling this. But no harm of keeping it here.
42 #just in case.
43
44
45 if not type(result,freeof('int')) then
46     return "F";
47 end if;
48
49
50 if ExpnType_result<=ExpnType_optimal then
51     if debug then
52         print("ExpnType_result<=ExpnType_optimal");
53     fi;
54     if is_contains_complex(result) then
55         if is_contains_complex(optimal) then

```

```

56     if debug then
57         print("both result and optimal complex");
58     fi;
59     #both result and optimal complex
60     if leaf_count_result<=2*leaf_count_optimal then
61         return "A";
62     else
63         return "B";
64     end if
65 else #result contains complex but optimal is not
66     if debug then
67         print("result contains complex but optimal is not");
68     fi;
69     return "C";
70 end if
71 else # result do not contain complex
72     # this assumes optimal do not as well
73     if debug then
74         print("result do not contain complex, this assumes optimal do not
as well");
75     fi;
76     if leaf_count_result<=2*leaf_count_optimal then
77         if debug then
78             print("leaf_count_result<=2*leaf_count_optimal");
79         fi;
80         return "A";
81     else
82         if debug then
83             print("leaf_count_result>2*leaf_count_optimal");
84         fi;
85         return "B";
86     end if
87 end if
88 else #ExpnType(result) > ExpnType(optimal)
89     if debug then
90         print("ExpnType(result) > ExpnType(optimal)");
91     fi;
92     return "C";
93 end if
94
95 end proc:
96
97 #
98 # is_contains_complex(result)
99 # takes expressions and returns true if it contains "I" else false
100 #
101 #Nasser 032417

```

```

102 is_contains_complex:= proc(expression)
103   return (has(expression,I));
104 end proc:
105
106 # The following summarizes the type number assigned an expression
107 # based on the functions it involves
108 # 1 = rational function
109 # 2 = algebraic function
110 # 3 = elementary function
111 # 4 = special function
112 # 5 = hyperpergeometric function
113 # 6 = appell function
114 # 7 = rootsum function
115 # 8 = integrate function
116 # 9 = unknown function
117
118 ExpnType := proc(expn)
119   if type(expn,'atomic') then
120     1
121   elif type(expn,'list') then
122     apply(max,map(ExpnType,expn))
123   elif type(expn,'sqrt') then
124     if type(op(1,expn),'rational') then
125       1
126     else
127       max(2,ExpnType(op(1,expn)))
128     end if
129   elif type(expn,'^^') then
130     if type(op(2,expn),'integer') then
131       ExpnType(op(1,expn))
132     elif type(op(2,expn),'rational') then
133       if type(op(1,expn),'rational') then
134         1
135       else
136         max(2,ExpnType(op(1,expn)))
137       end if
138     else
139       max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
140     end if
141   elif type(expn,'+'') or type(expn,'*') then
142     max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
143   elif ElementaryFunctionQ(op(0,expn)) then
144     max(3,ExpnType(op(1,expn)))
145   elif SpecialFunctionQ(op(0,expn)) then
146     max(4,apply(max,map(ExpnType,[op(expn)])))
147   elif HypergeometricFunctionQ(op(0,expn)) then
148     max(5,apply(max,map(ExpnType,[op(expn)])))

```

```

149   elif AppellFunctionQ(op(0,expn)) then
150       max(6,apply(max,map(ExpnType,[op(expn)])))
151   elif op(0,expn)='int' then
152       max(8,apply(max,map(ExpnType,[op(expn)]))) else
153       9
154   end if
155 end proc:
156
157 ElementaryFunctionQ := proc(func)
158     member(func,[
159         exp,log,ln,
160         sin,cos,tan,cot,sec,csc,
161         arcsin,arccos,arctan,arccot,arcsec,arccsc,
162         sinh,cosh,tanh,coth,sech,csch,
163         arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
164 end proc:
165
166 SpecialFunctionQ := proc(func)
167     member(func,[
168         erf,erfc,erfi,
169         FresnelS,FresnelC,
170         Ei,Ei,Li,Si,Ci,Shi,Chi,
171         GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
172         EllipticF,EllipticE,EllipticPi])
173 end proc:
174
175 HypergeometricFunctionQ := proc(func)
176     member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
177 end proc:
178
179 AppellFunctionQ := proc(func)
180     member(func,[AppellF1])
181 end proc:
182
183
184 # u is a sum or product. rest(u) returns all but the
185 # first term or factor of u.
186 rest := proc(u) local v;
187     if nops(u)=2 then
188         op(2,u)
189     else
190         apply(op(0,u),op(2..nops(u),u))
191     end if
192 end proc:
193
194 #leafcount(u) returns the number of nodes in u.
195 #Nasser 3/23/17 Replaced by build-in leafCount from package in Maple

```

```

196 leafcount := proc(u)
197     MmaTranslator[Mma][LeafCount](u);
198 end proc:

```

4.0.3 Sympy grading function

```

1 #Dec 24, 2019. Nasser M. Abbasi:
2 #           Port of original Maple grading function by
3 #           Albert Rich to use with Sympy/Python
4 #Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
5 #           added 'exp_polar'
6 from sympy import *
7
8 def leaf_count(expr):
9     #sympy do not have leaf count function. This is approximation
10    return round(1.7*count_ops(expr))
11
12 def is_sqrt(expr):
13     if isinstance(expr,Pow):
14         if expr.args[1] == Rational(1,2):
15             return True
16         else:
17             return False
18     else:
19         return False
20
21 def is_elementary_function(func):
22     return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
23                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
24                    asinh,acosh,atanh,acoth,asech,acsch
25                    ]
26
27 def is_special_function(func):
28     return func in [ erf,erfc,erfi,
29                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
30                    gamma,loggamma,digamma,zeta,polylog,LambertW,
31                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
32                    ]
33
34 def is_hypergeometric_function(func):
35     return func in [hyper]
36
37 def is_appell_function(func):
38     return func in [appellf1]
39
40 def is_atom(expn):
41     try:

```



```

42     if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
43         return True
44     else:
45         return False
46
47     except AttributeError as error:
48         return False
49
50 def expnType(expn):
51     debug=False
52     if debug:
53         print("expn=",expn,"type(expn)=",type(expn))
54
55     if is_atom(expn):
56         return 1
57     elif isinstance(expn,list):
58         return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
59     elif is_sqrt(expn):
60         if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
61             return 1
62         else:
63             return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
64     elif isinstance(expn,Pow): #type(expn,'^^')
65         if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
66             return expnType(expn.args[0]) #ExpnType(op(1,expn))
67         elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
68             if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
69                 return 1
70             else:
71                 return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)
72 ))
73     else:
74         return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,
75 ExpnType(op(1,expn)),ExpnType(op(2,expn)))
76     elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type
77 (expn,'*')
78         m1 = expnType(expn.args[0])
79         m2 = expnType(list(expn.args[1:]))
80         return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
81     elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
82         return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
83     elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
84         m1 = max(map(expnType, list(expn.args)))
85         return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
86     elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,
87 expn))
88         m1 = max(map(expnType, list(expn.args)))

```

```

85     return max(5,m1)    #max(5,apply(max,map(ExpnType,[op(expn)])))
86 elif is_appell_function(expn.func):
87     m1 = max(map(expnType, list(expn.args)))
88     return max(6,m1)    #max(5,apply(max,map(ExpnType,[op(expn)])))
89 elif isinstance(expn,RootSum):
90     m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,
Apply[List,expn]],7]],
91     return max(7,m1)
92 elif str(expn).find("Integral") != -1:
93     m1 = max(map(expnType, list(expn.args)))
94     return max(8,m1)    #max(5,apply(max,map(ExpnType,[op(expn)])))
95 else:
96     return 9
97
98 #main function
99 def grade_antiderivative(result,optimal):
100
101     leaf_count_result  = leaf_count(result)
102     leaf_count_optimal = leaf_count(optimal)
103
104     expnType_result  = expnType(result)
105     expnType_optimal = expnType(optimal)
106
107     if str(result).find("Integral") != -1:
108         return "F"
109
110     if expnType_result <= expnType_optimal:
111         if result.has(I):
112             if optimal.has(I): #both result and optimal complex
113                 if leaf_count_result <= 2*leaf_count_optimal:
114                     return "A"
115                 else:
116                     return "B"
117             else: #result contains complex but optimal is not
118                 return "C"
119         else: # result do not contain complex, this assumes optimal do not as
well
120             if leaf_count_result <= 2*leaf_count_optimal:
121                 return "A"
122             else:
123                 return "B"
124     else:
125         return "C"

```

4.0.4 SageMath grading function

1 #Dec 24, 2019. Nasser: Ported original Maple grading function by

```

2 #           Albert Rich to use with Sagemath. This is used to
3 #           grade Fracas, Giac and Maxima results.
4 #Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
5 #           'arctan2','floor','abs','log_integral'
6
7 from sage.all import *
8 from sage.symbolic.operators import add_vararg, mul_vararg
9
10 def tree(expr):
11     debug=False;
12     if debug:
13         print ("Enter tree(expr), expr=",expr)
14         print ("expr.operator()=",expr.operator())
15         print ("expr.operands()=",expr.operands())
16         print ("map(tree, expr.operands()=",map(tree, expr.operands()))
17
18     if expr.operator() is None:
19         return expr
20     else:
21         return [expr.operator()+list(map(tree, expr.operands()))
22
23 def leaf_count(anti):
24     debug=False;
25
26     if debug: print ("Enter leaf_count, anti=", anti, " len(anti)=", len(anti))
27
28     if len(anti) == 0: #special check for optimal being 0 for some test cases.
29         if debug: print ("len(anti) == 0")
30         return 1
31     else:
32         if debug: print ("round(1.35*len(flatten(tree(anti))))=",round(1.35*len(
33         flatten(tree(anti))))
34         return round(1.35*len(flatten(tree(anti)))) #fudge factor
35         #since this estimate of leaf count is bit lower than
36         #what it should be compared to Mathematica's
37
38 def is_sqrt(expr):
39     debug=False;
40     if expr.operator() == operator.pow: #isinstance(expr,Pow):
41         if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
42             if debug: print ("expr is sqrt")
43             return True
44         else:
45             return False
46     else:
47         return False

```

```

48 def is_elementary_function(func):
49     debug = False
50
51     m = func.name() in ['exp','log','ln',
52         'sin','cos','tan','cot','sec','csc',
53         'arcsin','arccos','arctan','arccot','arcsec','arccsc',
54         'sinh','cosh','tanh','coth','sech','csch',
55         'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
56         'arctan2','floor','abs'
57     ]
58     if debug:
59         if m:
60             print ("func ", func , " is elementary_function")
61         else:
62             print ("func ", func , " is NOT elementary_function")
63
64
65     return m
66
67 def is_special_function(func):
68     debug = False
69
70     if debug: print ("type(func)=", type(func))
71
72     m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
73         'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','
74     sinh_integral'
75         'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
76         'polylog','lambert_w','elliptic_f','elliptic_e',
77         'elliptic_pi','exp_integral_e','log_integral']
78
79     if debug:
80         print ("m=",m)
81         if m:
82             print ("func ", func ," is special_function")
83         else:
84             print ("func ", func ," is NOT special_function")
85
86     return m
87
88
89 def is_hypergeometric_function(func):
90     return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U
91     ']
92
93 def is_appell_function(func):

```

```

93     return func.name() in ['hypergeometric']    #[appellf1] can't find this in
          sagemath
94
95 def is_atom(expn):
96
97     #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type-in-maple/
98     try:
99         if expn.parent() is SR:
100             return expn.operator() is None
101         if expn.parent() in (ZZ, QQ, AA, QQbar):
102             return expn in expn.parent() # Should always return True
103         if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
104             return expn in expn.parent().base_ring() or expn in expn.parent().
          gens()
105         return False
106
107     except AttributeError as error:
108         return False
109
110
111 def expnType(expn):
112     debug=False
113
114     if debug:
115         print(">>>>Enter expnType, expn=", expn)
116         print(">>>>is_atom(expn)=", is_atom(expn))
117
118     if is_atom(expn):
119         return 1
120     elif type(expn)==list:    #isinstance(expn,list):
121         return max(map(expnType, expn))    #apply(max,map(ExpnType,expn))
122     elif is_sqrt(expn):
123         if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],
          Rational):
124             return 1
125         else:
126             return max(2,expnType(expn.operands()[0]))    #max(2,expnType(expn.
          args[0]))
127     elif expn.operator() == operator.pow:    #isinstance(expn,Pow)
128         if type(expn.operands()[1])==Integer:    #isinstance(expn.args[1],Integer)
129             return expnType(expn.operands()[0])    #expnType(expn.args[0])
130         elif type(expn.operands()[1])==Rational:    #isinstance(expn.args[1],
          Rational)
131             if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],
          Rational)
132                 return 1

```

```

133         else:
134             return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
135         else:
136             return max(3,expnType(expn.operands()[0]),expnType(expn.operands()
[1])) #max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))
137         elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #
isinstance(expn,Add) or isinstance(expn,Mul)
138             m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
139             m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
140             return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
141         elif is_elementary_function(expn.operator()): #is_elementary_function(expn.
func)
142             return max(3,expnType(expn.operands()[0]))
143         elif is_special_function(expn.operator()): #is_special_function(expn.func)
144             m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
145             return max(4,m1) #max(4,m1)
146         elif is_hypergeometric_function(expn.operator()): #
is_hypergeometric_function(expn.func)
147             m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
148             return max(5,m1) #max(5,m1)
149         elif is_appell_function(expn.operator()):
150             m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
151             return max(6,m1) #max(6,m1)
152         elif str(expn).find("Integral") != -1: #this will never happen, since it
153             #is checked before calling the grading function that is passed.
154             #but kept it here.
155             m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
156             return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
157         else:
158             return 9
159
160 #main function
161 def grade_antiderivative(result,optimal):
162     debug = False;
163
164     if debug: print ("Enter grade_antiderivative for sagemath")
165
166     leaf_count_result = leaf_count(result)
167     leaf_count_optimal = leaf_count(optimal)
168
169     if debug: print ("leaf_count_result=", leaf_count_result, "
leaf_count_optimal=",leaf_count_optimal)

```

```
170
171
172     expnType_result = expnType(result)
173     expnType_optimal = expnType(optimal)
174
175     if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",
176                     expnType_optimal)
177
178     if expnType_result <= expnType_optimal:
179         if result.has(I):
180             if optimal.has(I): #both result and optimal complex
181                 if leaf_count_result <= 2*leaf_count_optimal:
182                     return "A"
183             else:
184                 return "B"
185             else: #result contains complex but optimal is not
186                 return "C"
187         else: # result do not contain complex, this assumes optimal do not as
188             well
189                 if leaf_count_result <= 2*leaf_count_optimal:
190                     return "A"
191                 else:
192                     return "B"
193         else:
194             return "C"
```