

Computer algebra independent integration tests

1-Algebraic-functions/1.1-Binomial-products/1.1.2-Quadratic/1.1.2.3-
 $a+b-x^2-p-c+d-x^2-q$

Nasser M. Abbasi

June 28, 2021

Compiled on June 28, 2021 at 7:39pm

Contents

1	Introduction	3
1.1	Listing of CAS systems tested	3
1.2	Results	3
1.3	Performance	7
1.4	list of integrals that has no closed form antiderivative	8
1.5	list of integrals solved by CAS but has no known antiderivative	8
1.6	list of integrals solved by CAS but failed verification	8
1.7	Timing	9
1.8	Verification	9
1.9	Important notes about some of the results	9
1.9.1	Important note about Maxima results	9
1.9.2	Important note about FriCAS and Giac/XCAS results	10
1.9.3	Important note about finding leaf size of antiderivative	10
1.9.4	Important note about Mupad results	11
1.10	Design of the test system	11
2	detailed summary tables of results	13
2.1	List of integrals sorted by grade for each CAS	13
2.1.1	Rubi	13
2.1.2	Mathematica	13
2.1.3	Maple	14
2.1.4	Maxima	14
2.1.5	FriCAS	14
2.1.6	Sympy	15
2.1.7	Giac	15
2.1.8	Mupad	16
2.2	Detailed conclusion table per each integral for all CAS systems	17
2.3	Detailed conclusion table specific for Rubi results	75
3	Listing of integrals	87
3.1	$\int (a + bx^2)(c + dx^2)^4 dx$	87
3.2	$\int (a + bx^2)(c + dx^2)^3 dx$	90
3.3	$\int (a + bx^2)(c + dx^2)^2 dx$	92
3.4	$\int (a + bx^2)(c + dx^2) dx$	94

3.5	$\int \frac{a+bx^2}{c+dx^2} dx$	96
3.6	$\int \frac{a+bx^2}{(c+dx^2)^2} dx$	99
3.7	$\int \frac{a+bx^2}{(c+dx^2)^3} dx$	102
3.8	$\int (a+bx^2)^2 (c+dx^2)^3 dx$	105
3.9	$\int (a+bx^2)^2 (c+dx^2)^2 dx$	107
3.10	$\int (a+bx^2)^2 (c+dx^2) dx$	109
3.11	$\int \frac{(a+bx^2)^2}{c+dx^2} dx$	111
3.12	$\int \frac{(a+bx^2)^2}{(c+dx^2)^2} dx$	114
3.13	$\int \frac{(a+bx^2)^2}{(c+dx^2)^3} dx$	117
3.14	$\int (a+bx^2)^3 (c+dx^2)^3 dx$	120
3.15	$\int (a+bx^2)^3 (c+dx^2)^2 dx$	123
3.16	$\int (a+bx^2)^3 (c+dx^2) dx$	125
3.17	$\int \frac{(a+bx^2)^3}{c+dx^2} dx$	127
3.18	$\int \frac{(a+bx^2)^3}{(c+dx^2)^2} dx$	130
3.19	$\int \frac{(a+bx^2)^3}{(c+dx^2)^3} dx$	134
3.20	$\int \frac{(c+dx^2)^4}{a+bx^2} dx$	138
3.21	$\int \frac{(c+dx^2)^3}{a+bx^2} dx$	141
3.22	$\int \frac{(c+dx^2)^2}{a+bx^2} dx$	144
3.23	$\int \frac{c+dx^2}{a+bx^2} dx$	147
3.24	$\int \frac{1}{(a+bx^2)(c+dx^2)} dx$	150
3.25	$\int \frac{1}{(a+bx^2)(c+dx^2)^2} dx$	153
3.26	$\int \frac{1}{(a+bx^2)(c+dx^2)^3} dx$	158
3.27	$\int \frac{(c+dx^2)^5}{(a+bx^2)^2} dx$	164
3.28	$\int \frac{(c+dx^2)^4}{(a+bx^2)^2} dx$	168
3.29	$\int \frac{(c+dx^2)^3}{(a+bx^2)^2} dx$	172
3.30	$\int \frac{(c+dx^2)^2}{(a+bx^2)^2} dx$	176
3.31	$\int \frac{c+dx^2}{(a+bx^2)^2} dx$	179
3.32	$\int \frac{1}{(a+bx^2)^2(c+dx^2)} dx$	182
3.33	$\int \frac{1}{(a+bx^2)^2(c+dx^2)^2} dx$	187
3.34	$\int \frac{1}{(a+bx^2)^2(c+dx^2)^3} dx$	193

3.35	$\int \frac{(c+dx^2)^5}{(a+bx^2)^3} dx$	201
3.36	$\int \frac{(c+dx^2)^4}{(a+bx^2)^3} dx$	205
3.37	$\int \frac{(c+dx^2)^3}{(a+bx^2)^3} dx$	209
3.38	$\int \frac{(c+dx^2)^2}{(a+bx^2)^3} dx$	213
3.39	$\int \frac{c+dx^2}{(a+bx^2)^3} dx$	216
3.40	$\int \frac{1}{(a+bx^2)^3(c+dx^2)} dx$	219
3.41	$\int \frac{1}{(a+bx^2)^3(c+dx^2)^2} dx$	225
3.42	$\int \frac{1}{(a+bx^2)^3(c+dx^2)^3} dx$	233
3.43	$\int \frac{(-1+x^2)^3}{(1+x^2)^4} dx$	243
3.44	$\int \frac{(-1+x^2)^4}{(1+x^2)^5} dx$	246
3.45	$\int \sqrt{a+bx^2} (c+dx^2)^3 dx$	249
3.46	$\int \sqrt{a+bx^2} (c+dx^2)^2 dx$	253
3.47	$\int \sqrt{a+bx^2} (c+dx^2) dx$	257
3.48	$\int \sqrt{a+bx^2} dx$	260
3.49	$\int \frac{\sqrt{a+bx^2}}{c+dx^2} dx$	263
3.50	$\int \frac{\sqrt{a+bx^2}}{(c+dx^2)^2} dx$	267
3.51	$\int \frac{\sqrt{a+bx^2}}{(c+dx^2)^3} dx$	271
3.52	$\int \frac{\sqrt{a+bx^2}}{(c+dx^2)^4} dx$	275
3.53	$\int (a+bx^2)^{3/2} (c+dx^2)^3 dx$	279
3.54	$\int (a+bx^2)^{3/2} (c+dx^2)^2 dx$	283
3.55	$\int (a+bx^2)^{3/2} (c+dx^2) dx$	287
3.56	$\int (a+bx^2)^{3/2} dx$	290
3.57	$\int \frac{(a+bx^2)^{3/2}}{c+dx^2} dx$	293
3.58	$\int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^2} dx$	297
3.59	$\int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^3} dx$	302
3.60	$\int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^4} dx$	306
3.61	$\int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^5} dx$	310
3.62	$\int (a+bx^2)^{5/2} (c+dx^2)^3 dx$	315
3.63	$\int (a+bx^2)^{5/2} (c+dx^2)^2 dx$	320
3.64	$\int (a+bx^2)^{5/2} (c+dx^2) dx$	324

3.65	$\int (a + bx^2)^{5/2} dx$	328
3.66	$\int \frac{(a+bx^2)^{5/2}}{c+dx^2} dx$	331
3.67	$\int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^2} dx$	336
3.68	$\int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^3} dx$	340
3.69	$\int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^4} dx$	344
3.70	$\int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^5} dx$	348
3.71	$\int \frac{\sqrt{1-x^2}}{1+x^2} dx$	352
3.72	$\int \frac{\sqrt{1+x^2}}{-1+x^2} dx$	355
3.73	$\int \frac{\sqrt{1-x^2}}{-1+2x^2} dx$	358
3.74	$\int \frac{(c+dx^2)^3}{\sqrt{a+bx^2}} dx$	361
3.75	$\int \frac{(c+dx^2)^2}{\sqrt{a+bx^2}} dx$	365
3.76	$\int \frac{c+dx^2}{\sqrt{a+bx^2}} dx$	368
3.77	$\int \frac{1}{\sqrt{a+bx^2}} dx$	371
3.78	$\int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx$	373
3.79	$\int \frac{1}{\sqrt{a+bx^2}(c+dx^2)^2} dx$	376
3.80	$\int \frac{1}{\sqrt{a+bx^2}(c+dx^2)^3} dx$	380
3.81	$\int \frac{(c+dx^2)^4}{(a+bx^2)^{3/2}} dx$	384
3.82	$\int \frac{(c+dx^2)^3}{(a+bx^2)^{3/2}} dx$	388
3.83	$\int \frac{(c+dx^2)^2}{(a+bx^2)^{3/2}} dx$	392
3.84	$\int \frac{c+dx^2}{(a+bx^2)^{3/2}} dx$	395
3.85	$\int \frac{1}{(a+bx^2)^{3/2}} dx$	398
3.86	$\int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)} dx$	400
3.87	$\int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)^2} dx$	404
3.88	$\int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)^3} dx$	408
3.89	$\int \frac{(c+dx^2)^4}{(a+bx^2)^{5/2}} dx$	414
3.90	$\int \frac{(c+dx^2)^3}{(a+bx^2)^{5/2}} dx$	418
3.91	$\int \frac{(c+dx^2)^2}{(a+bx^2)^{5/2}} dx$	422
3.92	$\int \frac{c+dx^2}{(a+bx^2)^{5/2}} dx$	425

3.93	$\int \frac{1}{(a+bx^2)^{5/2}} dx$	428
3.94	$\int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)} dx$	431
3.95	$\int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)^2} dx$	435
3.96	$\int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)^3} dx$	440
3.97	$\int \frac{(a+bx^2)^3}{(c+dx^2)^{11/2}} dx$	446
3.98	$\int \frac{(a+bx^2)^2}{(c+dx^2)^{9/2}} dx$	450
3.99	$\int \frac{a+bx^2}{(c+dx^2)^{7/2}} dx$	453
3.100	$\int \frac{1}{(c+dx^2)^{5/2}} dx$	456
3.101	$\int \frac{1}{(a+bx^2)(c+dx^2)^{3/2}} dx$	459
3.102	$\int \frac{1}{(a+bx^2)^2 \sqrt{c+dx^2}} dx$	463
3.103	$\int \frac{\sqrt{c+dx^2}}{(a+bx^2)^3} dx$	467
3.104	$\int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^4} dx$	471
3.105	$\int \frac{1}{\left(\frac{bc}{d}+bx^2\right)\sqrt{c+dx^2}} dx$	475
3.106	$\int \frac{1}{\sqrt{1-x^2}(1+x^2)} dx$	477
3.107	$\int \frac{1}{(a+bx^2)\sqrt{c+dx^2}} dx$	480
3.108	$\int \frac{-1+x^2}{(1+x^2)^{3/2}} dx$	483
3.109	$\int (a-bx^2)^{2/3} (3a+bx^2)^3 dx$	485
3.110	$\int (a-bx^2)^{2/3} (3a+bx^2)^2 dx$	489
3.111	$\int (a-bx^2)^{2/3} (3a+bx^2) dx$	493
3.112	$\int \frac{(a-bx^2)^{2/3}}{3a+bx^2} dx$	497
3.113	$\int \frac{(a-bx^2)^{2/3}}{(3a+bx^2)^2} dx$	501
3.114	$\int \frac{(a-bx^2)^{2/3}}{(3a+bx^2)^3} dx$	505
3.115	$\int \frac{(a-bx^2)^{2/3}}{(3a+bx^2)^4} dx$	510
3.116	$\int (a-bx^2)^{5/3} (3a+bx^2)^3 dx$	515
3.117	$\int (a-bx^2)^{5/3} (3a+bx^2)^2 dx$	519
3.118	$\int (a-bx^2)^{5/3} (3a+bx^2) dx$	523
3.119	$\int \frac{(a-bx^2)^{5/3}}{3a+bx^2} dx$	527
3.120	$\int \frac{(a-bx^2)^{5/3}}{(3a+bx^2)^2} dx$	531
3.121	$\int \frac{(a-bx^2)^{5/3}}{(3a+bx^2)^3} dx$	535

3.122	$\int \frac{(3a+bx^2)^4}{\sqrt[3]{a-bx^2}} dx$	540
3.123	$\int \frac{(3a+bx^2)^3}{\sqrt[3]{a-bx^2}} dx$	544
3.124	$\int \frac{(3a+bx^2)^2}{\sqrt[3]{a-bx^2}} dx$	548
3.125	$\int \frac{3a+bx^2}{\sqrt[3]{a-bx^2}} dx$	552
3.126	$\int \frac{1}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx$	556
3.127	$\int \frac{1}{\sqrt[3]{a-bx^2}(3a+bx^2)^2} dx$	559
3.128	$\int \frac{1}{\sqrt[3]{a-bx^2}(3a+bx^2)^3} dx$	563
3.129	$\int \frac{(3a+bx^2)^3}{(a-bx^2)^{4/3}} dx$	568
3.130	$\int \frac{(3a+bx^2)^2}{(a-bx^2)^{4/3}} dx$	572
3.131	$\int \frac{3a+bx^2}{(a-bx^2)^{4/3}} dx$	576
3.132	$\int \frac{1}{(a-bx^2)^{4/3}(3a+bx^2)} dx$	580
3.133	$\int \frac{1}{(a-bx^2)^{4/3}(3a+bx^2)^2} dx$	584
3.134	$\int \frac{1}{(a-bx^2)^{4/3}(3a+bx^2)^3} dx$	589
3.135	$\int \frac{(3a+bx^2)^4}{(a-bx^2)^{7/3}} dx$	594
3.136	$\int \frac{(3a+bx^2)^3}{(a-bx^2)^{7/3}} dx$	598
3.137	$\int \frac{(3a+bx^2)^2}{(a-bx^2)^{7/3}} dx$	602
3.138	$\int \frac{3a+bx^2}{(a-bx^2)^{7/3}} dx$	605
3.139	$\int \frac{1}{(a-bx^2)^{7/3}(3a+bx^2)} dx$	609
3.140	$\int \frac{1}{(a-bx^2)^{7/3}(3a+bx^2)^2} dx$	614
3.141	$\int \frac{1}{(-3a-bx^2)\sqrt[3]{-a+bx^2}} dx$	619
3.142	$\int \frac{1}{(3a-bx^2)\sqrt[3]{a+bx^2}} dx$	622
3.143	$\int \frac{1}{(c-dx^2)\sqrt[3]{c+3dx^2}} dx$	625
3.144	$\int \frac{1}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx$	628
3.145	$\int \frac{1}{\sqrt[3]{c-3dx^2}(c+dx^2)} dx$	631
3.146	$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx$	634
3.147	$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx$	638
3.148	$\int \frac{3-x}{\sqrt[3]{1-x^2}(3+x^2)} dx$	641
3.149	$\int \frac{3+x}{\sqrt[3]{1-x^2}(3+x^2)} dx$	644
3.150	$\int \frac{1}{\sqrt[3]{a+bx^2}\left(\frac{9ad}{b}+dx^2\right)} dx$	647

3.151	$\int \frac{1}{\sqrt[3]{a-bx^2}\left(-\frac{9ad}{b}+dx^2\right)} dx$	650
3.152	$\int \frac{1}{\sqrt[3]{-a+bx^2}\left(-\frac{9ad}{b}+dx^2\right)} dx$	653
3.153	$\int \frac{1}{\sqrt[3]{-a-bx^2}\left(\frac{9ad}{b}+dx^2\right)} dx$	656
3.154	$\int \frac{1}{\sqrt[3]{2+bx^2}\left(\frac{18d}{b}+dx^2\right)} dx$	659
3.155	$\int \frac{1}{\sqrt[3]{-2+bx^2}\left(-\frac{18d}{b}+dx^2\right)} dx$	662
3.156	$\int \frac{1}{\sqrt[3]{2+3x^2}(6d+dx^2)} dx$	665
3.157	$\int \frac{1}{\sqrt[3]{2-3x^2}(-6d+dx^2)} dx$	668
3.158	$\int \frac{1}{\sqrt[3]{-2+3x^2}(-6d+dx^2)} dx$	671
3.159	$\int \frac{1}{\sqrt[3]{-2-3x^2}(6d+dx^2)} dx$	674
3.160	$\int \frac{1}{\sqrt[3]{1+x^2}(9+x^2)} dx$	677
3.161	$\int \frac{1}{\sqrt[3]{1+bx^2}(9+bx^2)} dx$	681
3.162	$\int \frac{1}{\sqrt[3]{1-x^2}(9-x^2)} dx$	684
3.163	$\int \frac{\sqrt{-1+c^2x^2}}{(d-c^2dx^2)^{5/2}} dx$	687
3.164	$\int \frac{1}{(-1+c^2x^2)^{3/2}\sqrt{d-c^2dx^2}} dx$	690
3.165	$\int \frac{1}{\sqrt{-1+c^2x^2}(d-c^2dx^2)^{3/2}} dx$	693
3.166	$\int (a+bx^2)^{3/2}\sqrt{c+dx^2} dx$	696
3.167	$\int \sqrt{a+bx^2}\sqrt{c+dx^2} dx$	700
3.168	$\int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx$	703
3.169	$\int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{3/2}} dx$	706
3.170	$\int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{5/2}} dx$	709
3.171	$\int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{7/2}} dx$	712
3.172	$\int (a+bx^2)^{3/2}(c+dx^2)^{3/2} dx$	716
3.173	$\int \sqrt{a+bx^2}(c+dx^2)^{3/2} dx$	720
3.174	$\int \frac{(c+dx^2)^{3/2}}{\sqrt{a+bx^2}} dx$	724
3.175	$\int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^{3/2}} dx$	728
3.176	$\int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^{5/2}} dx$	732
3.177	$\int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^{7/2}} dx$	736
3.178	$\int \sqrt{2+bx^2}\sqrt{3+dx^2} dx$	740
3.179	$\int \sqrt{3-6x^2}\sqrt{2+4x^2} dx$	743
3.180	$\int \sqrt{2+4x^2}\sqrt{3+6x^2} dx$	746
3.181	$\int \frac{\sqrt{2+bx^2}}{\sqrt{3+dx^2}} dx$	748

3.182	$\int \frac{\sqrt{4-x^2}}{\sqrt{c+dx^2}} dx$	751
3.183	$\int \frac{\sqrt{4+x^2}}{\sqrt{c+dx^2}} dx$	754
3.184	$\int \frac{\sqrt{1-x^2}}{\sqrt{2-3x^2}} dx$	757
3.185	$\int \frac{\sqrt{4-x^2}}{\sqrt{2-3x^2}} dx$	759
3.186	$\int \frac{\sqrt{1-4x^2}}{\sqrt{2-3x^2}} dx$	761
3.187	$\int \frac{\sqrt{1+x^2}}{\sqrt{1-x^2}} dx$	763
3.188	$\int \frac{\sqrt{1+x^2}}{\sqrt{2-3x^2}} dx$	765
3.189	$\int \frac{\sqrt{4+x^2}}{\sqrt{2-3x^2}} dx$	767
3.190	$\int \frac{\sqrt{1+4x^2}}{\sqrt{2-3x^2}} dx$	769
3.191	$\int \frac{\sqrt{1-x^2}}{\sqrt{1+x^2}} dx$	771
3.192	$\int \frac{\sqrt{1-x^2}}{\sqrt{2+3x^2}} dx$	774
3.193	$\int \frac{\sqrt{4-x^2}}{\sqrt{2+3x^2}} dx$	777
3.194	$\int \frac{\sqrt{1-4x^2}}{\sqrt{2+3x^2}} dx$	780
3.195	$\int \frac{\sqrt{1+x^2}}{\sqrt{2+3x^2}} dx$	783
3.196	$\int \frac{\sqrt{4+x^2}}{\sqrt{2+3x^2}} dx$	786
3.197	$\int \frac{\sqrt{1+4x^2}}{\sqrt{2+3x^2}} dx$	789
3.198	$\int \frac{\sqrt{1-x^2}}{\sqrt{-1+2x^2}} dx$	792
3.199	$\int \frac{(a+bx^2)^{7/2}}{\sqrt{c+dx^2}} dx$	794
3.200	$\int \frac{(a+bx^2)^{5/2}}{\sqrt{c+dx^2}} dx$	799
3.201	$\int \frac{(a+bx^2)^{3/2}}{\sqrt{c+dx^2}} dx$	803
3.202	$\int \frac{\sqrt{a+bx^2}}{\sqrt{c+dx^2}} dx$	807
3.203	$\int \frac{1}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx$	810
3.204	$\int \frac{1}{(a+bx^2)^{3/2} \sqrt{c+dx^2}} dx$	813
3.205	$\int \frac{1}{(a+bx^2)^{5/2} \sqrt{c+dx^2}} dx$	817
3.206	$\int \frac{1}{(a+bx^2)^{7/2} \sqrt{c+dx^2}} dx$	821
3.207	$\int \frac{(a+bx^2)^{7/2}}{(c+dx^2)^{3/2}} dx$	825
3.208	$\int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^{3/2}} dx$	830
3.209	$\int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^{3/2}} dx$	834
3.210	$\int \frac{\sqrt{a+bx^2}}{(c+dx^2)^{3/2}} dx$	838
3.211	$\int \frac{1}{\sqrt{a+bx^2} (c+dx^2)^{3/2}} dx$	841

3.212	$\int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)^{3/2}} dx$	845
3.213	$\int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)^{3/2}} dx$	848
3.214	$\int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx$	852
3.215	$\int \frac{1}{\sqrt{a-bx^2}\sqrt{c+dx^2}} dx$	855
3.216	$\int \frac{1}{\sqrt{a+bx^2}\sqrt{c-dx^2}} dx$	858
3.217	$\int \frac{1}{\sqrt{a-bx^2}\sqrt{c-dx^2}} dx$	861
3.218	$\int \frac{1}{\sqrt{1-x^2}\sqrt{2+5x^2}} dx$	864
3.219	$\int \frac{1}{\sqrt{1-x^2}\sqrt{2+4x^2}} dx$	866
3.220	$\int \frac{1}{\sqrt{1-x^2}\sqrt{2+3x^2}} dx$	868
3.221	$\int \frac{1}{\sqrt{1-x^2}\sqrt{2+2x^2}} dx$	870
3.222	$\int \frac{1}{\sqrt{1-x^2}\sqrt{2+x^2}} dx$	872
3.223	$\int \frac{1}{\sqrt{1-x^2}\sqrt{2-x^2}} dx$	874
3.224	$\int \frac{1}{\sqrt{2-2x^2}\sqrt{1-x^2}} dx$	876
3.225	$\int \frac{1}{\sqrt{2-3x^2}\sqrt{1-x^2}} dx$	878
3.226	$\int \frac{1}{\sqrt{2-4x^2}\sqrt{1-x^2}} dx$	880
3.227	$\int \frac{1}{\sqrt{2-5x^2}\sqrt{1-x^2}} dx$	882
3.228	$\int \frac{1}{\sqrt{1+x^2}\sqrt{2+5x^2}} dx$	884
3.229	$\int \frac{1}{\sqrt{1+x^2}\sqrt{2+4x^2}} dx$	886
3.230	$\int \frac{1}{\sqrt{1+x^2}\sqrt{2+3x^2}} dx$	888
3.231	$\int \frac{1}{\sqrt{1+x^2}\sqrt{2+2x^2}} dx$	890
3.232	$\int \frac{1}{\sqrt{1+x^2}\sqrt{2+x^2}} dx$	892
3.233	$\int \frac{1}{\sqrt{2-x^2}\sqrt{1+x^2}} dx$	894
3.234	$\int \frac{1}{\sqrt{2-2x^2}\sqrt{1+x^2}} dx$	896
3.235	$\int \frac{1}{\sqrt{2-3x^2}\sqrt{1+x^2}} dx$	898
3.236	$\int \frac{1}{\sqrt{2-4x^2}\sqrt{1+x^2}} dx$	900
3.237	$\int \frac{1}{\sqrt{2-5x^2}\sqrt{1+x^2}} dx$	902
3.238	$\int \frac{1}{\sqrt{-1+x^2}\sqrt{2+5x^2}} dx$	904
3.239	$\int \frac{1}{\sqrt{-1+x^2}\sqrt{2+4x^2}} dx$	906
3.240	$\int \frac{1}{\sqrt{-1+x^2}\sqrt{2+3x^2}} dx$	908
3.241	$\int \frac{1}{\sqrt{-1+x^2}\sqrt{2+2x^2}} dx$	910
3.242	$\int \frac{1}{\sqrt{-1+x^2}\sqrt{2+x^2}} dx$	913
3.243	$\int \frac{1}{\sqrt{2-x^2}\sqrt{-1+x^2}} dx$	915
3.244	$\int \frac{1}{\sqrt{2-2x^2}\sqrt{-1+x^2}} dx$	917
3.245	$\int \frac{1}{\sqrt{2-3x^2}\sqrt{-1+x^2}} dx$	919
3.246	$\int \frac{1}{\sqrt{2-4x^2}\sqrt{-1+x^2}} dx$	921
3.247	$\int \frac{1}{\sqrt{2-5x^2}\sqrt{-1+x^2}} dx$	923
3.248	$\int \frac{1}{\sqrt{-1-x^2}\sqrt{2+5x^2}} dx$	925

3.249	$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2+4x^2}} dx$	927
3.250	$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2+3x^2}} dx$	929
3.251	$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2+2x^2}} dx$	931
3.252	$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2+x^2}} dx$	933
3.253	$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2-x^2}} dx$	935
3.254	$\int \frac{1}{\sqrt{2-2x^2} \sqrt{-1-x^2}} dx$	937
3.255	$\int \frac{1}{\sqrt{2-3x^2} \sqrt{-1-x^2}} dx$	940
3.256	$\int \frac{1}{\sqrt{2-4x^2} \sqrt{-1-x^2}} dx$	942
3.257	$\int \frac{1}{\sqrt{2-5x^2} \sqrt{-1-x^2}} dx$	944
3.258	$\int \frac{\sqrt{a+bx^2}}{\sqrt{c-dx^2}} dx$	946
3.259	$\int \frac{\sqrt{-a-bx^2}}{\sqrt{c-dx^2}} dx$	949
3.260	$\int \frac{\sqrt{a+bx^2}}{\sqrt{-c+dx^2}} dx$	952
3.261	$\int \frac{\sqrt{-a-bx^2}}{\sqrt{-c+dx^2}} dx$	955
3.262	$\int \frac{\sqrt{a-bx^2}}{\sqrt{c-dx^2}} dx$	958
3.263	$\int \frac{\sqrt{-a+bx^2}}{\sqrt{c-dx^2}} dx$	961
3.264	$\int \frac{\sqrt{a-bx^2}}{\sqrt{-c+dx^2}} dx$	964
3.265	$\int \frac{\sqrt{-a+bx^2}}{\sqrt{-c+dx^2}} dx$	967
3.266	$\int \frac{\sqrt{a+bx^2}}{\sqrt{c+dx^2}} dx$	970
3.267	$\int \frac{\sqrt{-a-bx^2}}{\sqrt{c+dx^2}} dx$	973
3.268	$\int \frac{\sqrt{a+bx^2}}{\sqrt{-c-dx^2}} dx$	976
3.269	$\int \frac{\sqrt{-a-bx^2}}{\sqrt{-c-dx^2}} dx$	979
3.270	$\int \frac{\sqrt{a-bx^2}}{\sqrt{c+dx^2}} dx$	982
3.271	$\int \frac{\sqrt{-a+bx^2}}{\sqrt{c+dx^2}} dx$	985
3.272	$\int \frac{\sqrt{a-bx^2}}{\sqrt{-c-dx^2}} dx$	988
3.273	$\int \frac{\sqrt{-a+bx^2}}{\sqrt{-c-dx^2}} dx$	991
3.274	$\int \frac{\sqrt{c+dx^2}}{\sqrt{a-bx^2}} dx$	994
3.275	$\int \frac{\sqrt{-c-dx^2}}{\sqrt{a-bx^2}} dx$	997
3.276	$\int \frac{\sqrt{c+dx^2}}{\sqrt{-a+bx^2}} dx$	1000
3.277	$\int \frac{\sqrt{-c-dx^2}}{\sqrt{-a+bx^2}} dx$	1003
3.278	$\int \frac{\sqrt{c-dx^2}}{\sqrt{a-bx^2}} dx$	1006
3.279	$\int \frac{\sqrt{-c+dx^2}}{\sqrt{a-bx^2}} dx$	1009
3.280	$\int \frac{\sqrt{c-dx^2}}{\sqrt{-a+bx^2}} dx$	1012
3.281	$\int \frac{\sqrt{-c+dx^2}}{\sqrt{-a+bx^2}} dx$	1015

3.282	$\int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx$	1018
3.283	$\int \frac{\sqrt{-c-dx^2}}{\sqrt{a+bx^2}} dx$	1021
3.284	$\int \frac{\sqrt{c+dx^2}}{\sqrt{-a-bx^2}} dx$	1024
3.285	$\int \frac{\sqrt{-c-dx^2}}{\sqrt{-a-bx^2}} dx$	1027
3.286	$\int \frac{\sqrt{c-dx^2}}{\sqrt{a+bx^2}} dx$	1030
3.287	$\int \frac{\sqrt{-c+dx^2}}{\sqrt{a+bx^2}} dx$	1033
3.288	$\int \frac{\sqrt{c-dx^2}}{\sqrt{-a-bx^2}} dx$	1036
3.289	$\int \frac{\sqrt{-c+dx^2}}{\sqrt{-a-bx^2}} dx$	1039
3.290	$\int \frac{1}{\sqrt{2+bx^2} \sqrt{3+dx^2}} dx$	1042
3.291	$\int \frac{1}{\sqrt{4-x^2} \sqrt{c+dx^2}} dx$	1044
3.292	$\int \frac{1}{\sqrt{4+x^2} \sqrt{c+dx^2}} dx$	1046
3.293	$\int \frac{1}{\sqrt{1-x^2} \sqrt{-1+2x^2}} dx$	1048
3.294	$\int \frac{\sqrt{1-c^2x^2}}{\sqrt{1+c^2x^2}} dx$	1050
3.295	$\int \frac{\sqrt{2+bx^2}}{\sqrt{3+dx^2}} dx$	1053
3.296	$\int \frac{\sqrt{-1+3x^2}}{\sqrt{2-3x^2}} dx$	1056
3.297	$\int \frac{\sqrt{1+\frac{2cx^2}{b-\sqrt{b^2-4ac}}}}{\sqrt{1-\frac{2cx^2}{b+\sqrt{b^2-4ac}}}} dx$	1058
3.298	$\int \frac{\sqrt{1-\frac{2cx^2}{b-\sqrt{b^2-4ac}}}}{\sqrt{1-\frac{2cx^2}{b+\sqrt{b^2-4ac}}}} dx$	1061
3.299	$\int \frac{\sqrt{1+\frac{2cx^2}{b-\sqrt{b^2-4ac}}}}{\sqrt{1+\frac{2cx^2}{b+\sqrt{b^2-4ac}}}} dx$	1064
3.300	$\int \frac{\sqrt{1-\frac{2cx^2}{b-\sqrt{b^2-4ac}}}}{\sqrt{1+\frac{2cx^2}{b+\sqrt{b^2-4ac}}}} dx$	1068
3.301	$\int \frac{(1-2x^2)^m}{\sqrt{1-x^2}} dx$	1072
3.302	$\int \frac{1}{\sqrt{-1+x^2} \sqrt{7-4\sqrt{3}+x^2}} dx$	1074
3.303	$\int \frac{1}{\sqrt{3-3\sqrt{3}+2\sqrt{3}x^2} \sqrt{3+(-3+\sqrt{3})x^2}} dx$	1077
3.304	$\int \frac{1}{\sqrt[4]{2+3x^2} (4+3x^2)} dx$	1080
3.305	$\int \frac{1}{\sqrt[4]{2-3x^2} (4-3x^2)} dx$	1083
3.306	$\int \frac{1}{\sqrt[4]{2+bx^2} (4+bx^2)} dx$	1086
3.307	$\int \frac{1}{\sqrt[4]{2-bx^2} (4-bx^2)} dx$	1089
3.308	$\int \frac{1}{\sqrt[4]{a+3x^2} (2a+3x^2)} dx$	1092
3.309	$\int \frac{1}{\sqrt[4]{a-3x^2} (2a-3x^2)} dx$	1095

3.310	$\int \frac{1}{\sqrt[4]{a+bx^2}(2a+bx^2)} dx$	1098
3.311	$\int \frac{1}{\sqrt[4]{a-bx^2}(2a-bx^2)} dx$	1101
3.312	$\int \frac{1}{(-2+3x^2)\sqrt[4]{-1+3x^2}} dx$	1104
3.313	$\int \frac{1}{(-2-3x^2)\sqrt[4]{-1-3x^2}} dx$	1107
3.314	$\int \frac{1}{(-2+bx^2)\sqrt[4]{-1+bx^2}} dx$	1110
3.315	$\int \frac{1}{(-2-bx^2)\sqrt[4]{-1-bx^2}} dx$	1113
3.316	$\int \frac{1}{(-2a+3x^2)\sqrt[4]{-a+3x^2}} dx$	1116
3.317	$\int \frac{1}{(-2a-3x^2)\sqrt[4]{-a-3x^2}} dx$	1119
3.318	$\int \frac{1}{(-2a+bx^2)\sqrt[4]{-a+bx^2}} dx$	1122
3.319	$\int \frac{1}{(-2a-bx^2)\sqrt[4]{-a-bx^2}} dx$	1125
3.320	$\int \frac{1}{(2-x^2)\sqrt[4]{-1+x^2}} dx$	1128
3.321	$\int \frac{(a+bx^2)^{7/4}}{c+dx^2} dx$	1131
3.322	$\int \frac{(a+bx^2)^{5/4}}{c+dx^2} dx$	1135
3.323	$\int \frac{(a+bx^2)^{3/4}}{c+dx^2} dx$	1139
3.324	$\int \frac{\sqrt[4]{a+bx^2}}{c+dx^2} dx$	1142
3.325	$\int \frac{1}{\sqrt[4]{a+bx^2}(c+dx^2)} dx$	1145
3.326	$\int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx$	1148
3.327	$\int \frac{1}{(a+bx^2)^{5/4}(c+dx^2)} dx$	1151
3.328	$\int \frac{1}{(a+bx^2)^{7/4}(c+dx^2)} dx$	1154
3.329	$\int \frac{1}{(a+bx^2)^{9/4}(c+dx^2)} dx$	1158
3.330	$\int \frac{1}{(a+bx^2)^{11/4}(c+dx^2)} dx$	1162
3.331	$\int \frac{(a+bx^2)^{7/4}}{(c+dx^2)^2} dx$	1166
3.332	$\int \frac{(a+bx^2)^{5/4}}{(c+dx^2)^2} dx$	1170
3.333	$\int \frac{(a+bx^2)^{3/4}}{(c+dx^2)^2} dx$	1174
3.334	$\int \frac{\sqrt[4]{a+bx^2}}{(c+dx^2)^2} dx$	1178
3.335	$\int \frac{1}{\sqrt[4]{a+bx^2}(c+dx^2)^2} dx$	1182
3.336	$\int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)^2} dx$	1186
3.337	$\int \frac{1}{(a+bx^2)^{5/4}(c+dx^2)^2} dx$	1190
3.338	$\int \frac{1}{(a+bx^2)^{7/4}(c+dx^2)^2} dx$	1194
3.339	$\int \frac{1}{(a+bx^2)^{9/4}(c+dx^2)^2} dx$	1198

3.340	$\int \frac{1}{(a+bx^2)^{11/4}(c+dx^2)^2} dx$	1202
3.341	$\int (a+bx^2)^p (c+dx^2)^q dx$	1207
3.342	$\int (a+bx^2)^p (c+dx^2)^3 dx$	1209
3.343	$\int (a+bx^2)^p (c+dx^2)^2 dx$	1212
3.344	$\int (a+bx^2)^p (c+dx^2) dx$	1215
3.345	$\int (a+bx^2)^p dx$	1218
3.346	$\int \frac{(a+bx^2)^p}{c+dx^2} dx$	1220
3.347	$\int \frac{(a+bx^2)^p}{(c+dx^2)^2} dx$	1223
3.348	$\int \frac{(a+bx^2)^p}{(c+dx^2)^3} dx$	1226
3.349	$\int (a+bx^2)^{-1-\frac{bc}{2bc-2ad}} (c+dx^2)^{-1+\frac{ad}{2bc-2ad}} dx$	1229
4	Listing of Grading functions	1233
4.0.1	Mathematica and Rubi grading function	1233
4.0.2	Maple grading function	1235
4.0.3	Sympy grading function	1238
4.0.4	SageMath grading function	1240

Chapter 1

Introduction

This report gives the result of running the computer algebra independent integration problems. The listing of the problems are maintained by and can be downloaded from <https://rulebasedintegration.org>

The number of integrals in this report is [349]. This is test number [20].

1.1 Listing of CAS systems tested

The following systems were tested at this time.

1. Mathematica 12.3 (64 bit) on windows 10.
2. Rubi 4.16.1 in Mathematica 12.1 on windows 10.
3. Maple 2021.1 (64 bit) on windows 10.
4. Maxima 5.44 on Linux. (via sagemath 9.3)
5. Fricas 1.3.7 on Linux (via sagemath 9.3)
6. Giac/Xcas 1.7 on Linux. (via sagemath 9.3)
7. Sympy 1.8 under Python 3.8.8 using Anaconda distribution on Ubuntu.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 under windows 10 (64 bit)

Maxima, Fricas and Giac/Xcas were called from inside SageMath. This was done using SageMath integrate command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly using Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric $2F1$ functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	solved	Failed
Rubi	% 100.00 (349)	% 0.00 (0)
Mathematica	% 100.00 (349)	% 0.00 (0)
Maple	% 74.50 (260)	% 25.50 (89)
Maxima	% 22.64 (79)	% 77.36 (270)
Fricas	% 40.40 (141)	% 59.60 (208)
Sympy	% 29.51 (103)	% 70.49 (246)
Giac	% 30.95 (108)	% 69.05 (241)
Mupad	% 18.91 (66)	% 81.09 (283)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	99.71	0.00	0.29	0.00
Mathematica	55.01	2.58	42.41	0.00
Maple	52.15	16.62	5.73	25.50
Maxima	20.34	2.29	0.00	77.36
Fricas	20.34	19.77	0.29	59.60
Sympy	16.62	10.89	2.01	70.49
Giac	23.21	7.74	0.00	69.05
Mupad	0.00	18.91	0.00	81.09

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This the typical normal failure F .

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned F(-1).

The third is due to an exception generated. Assigned F(-2). This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	0	0.00 %	0.00 %	0.00 %
Maple	89	98.88 %	1.12 %	0.00 %
Maxima	270	100.00 %	0.00 %	0.00 %
Fricas	208	75.96 %	24.04 %	0.00 %
Sympy	246	89.43 %	10.57 %	0.00 %
Giac	241	97.51 %	0.00 %	2.49 %
Mupad	283	100.00 %	0.00 %	0.00 %

Table 1.4: Time and leaf size performance for each CAS


1.3 Performance


The table below summarizes the performance of each CAS system in terms of CPU time and leaf size of results.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.12	178.99	1.00	116.00	1.00
Mathematica	0.60	132.26	1.04	102.00	1.00
Maple	1.20	846.63	5.25	116.00	1.23
Maxima	2.14	157.72	1.24	116.00	1.16
Fricas	3.66	530.65	4.06	302.00	3.27
Sympy	6.67	159.68	1.89	95.00	1.74
Giac	1.04	225.66	1.66	129.00	1.14
Mupad	3.76	912.58	5.52	87.50	1.12

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used columns from the above table.





1.4 list of integrals that has no closed form antiderivative

{}

1.5 list of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {301}

Mathematica {46, 50, 54, 59, 63, 69, 75, 83, 86, 87, 88, 91, 94, 101, 102, 110, 112, 114, 115, 117, 119, 120, 121, 124, 126, 127, 128, 132, 133, 134, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 301, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 346, 347, 348}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of _int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call has completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out is not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica. Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative produced was correct.

Verification phase has 3 minutes time out. An integral whose result was not verified could still be correct. Further investigation is needed on those integrals which failed verifications. Such integrals are marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since these integrals are run in a batch mode, using an automated script, and by using `sagemath` (SageMath uses Maxima), then any integral where Maxima needs an interactive response from the user to answer a question during evaluation of the integral in order to complete the integration, will fail and is counted as failed.

The exception raised is `ValueError`. Therefore Maxima result below is lower than what could result if Maxima was run directly and each question Maxima asks was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate of the error is due to the interactive question being asked or not.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath loading of Maxima `abs_integrate` was found to cause some problem. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS and Giac/X-CAS results

There are Few integrals which failed due to SageMath not able to translate the result back to SageMath syntax and not because these CAS system were not able to do the integrations.

These will fail With error `Exception raised: NotImplementedError`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals.

Hopefully the next version of SageMath will have complete translation of FriCAS and XCAS syntax and I will re-run all the tests again when this happens.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi and Maple, the buildin system function `LeafSize` is used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special buildin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy and Giac antiderivatives is determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which is called directly from Python, the following code is used to obtain the leafsize of its result

```
try:
```

```
# 1.7 is a fudge factor since it is low side from actual leaf count
leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative, Maple was used to determine the leaf size of Mupad output by post processing.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine,'cos(x)*sin(x)')
the_variable = evalin(symengine,'x')
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
The following field present only in Rubi and Mathematica Tables
13. integer. 1 if result was verified or 0 if not verified.
The following fields present only in Rubi Tables
14. integer. Number of rules used.
15. integer. Integrand leaf size.
16. real number. Ratio of field 14 over field 15
17. integer. 1 if result was verified or 0 if not verified.
18. String of form "{n,n,...}" which is list of the rules used by Rubi

High level overview of the CAS independent integration test build system

Chapter 2

detailed summary tables of results

2.1 List of integrals sorted by grade for each CAS

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349 }

B grade: { }

C grade: { 301 }

F grade: { }

2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 47, 48, 49, 51, 52, 53, 55, 56, 57, 58, 59, 60, 61, 62, 64, 65, 66, 67, 68, 69, 70, 71, 73, 74, 76, 77, 78, 79, 80, 81, 82, 84, 85, 89, 90, 92, 93, 95, 96, 97, 98, 99, 100, 103, 104, 105, 106, 107, 108, 137, 163, 164, 165, 168, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 198, 202, 203, 204, 211, 214, 215, 216, 217, 218, 220, 221, 223, 225, 226, 227, 231, 234, 235, 236, 237, 238, 239, 240, 242, 244, 245, 246, 247, 251, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 294, 295, 296, 297, 298, 299, 300, 302, 303, 326, 342, 343, 344, 345, 349 }

B grade: { 50, 72, 224, 243, 293, 341, 346, 347, 348 }

C grade: { 46, 54, 63, 75, 83, 86, 87, 88, 91, 94, 101, 102, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 166, 167, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 191, 192, 193, 194, 195, 196, 197, 199, 200, 201, 205, 206, 207, 208, 209, 210, 212, 213, 219, 222, 228, 229, 230, 232, 233, 241,

248, 249, 250, 252, 253, 254, 292, 301, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340 }

F grade: { }

2.1.3 Maple

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 20, 21, 22, 23, 24, 25, 30, 31, 32, 33, 34, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 53, 54, 55, 56, 62, 63, 64, 65, 71, 74, 75, 76, 77, 81, 82, 83, 84, 85, 89, 90, 91, 92, 93, 97, 98, 99, 100, 105, 106, 108, 137, 163, 164, 165, 166, 167, 168, 169, 172, 173, 174, 175, 178, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 198, 199, 200, 201, 202, 203, 204, 207, 208, 209, 210, 211, 212, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 233, 234, 235, 236, 237, 238, 239, 240, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 253, 254, 255, 256, 257, 258, 261, 262, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 277, 279, 280, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 295, 296, 349 }

B grade: { 18, 19, 26, 27, 28, 29, 35, 36, 37, 40, 49, 50, 51, 52, 57, 58, 59, 60, 61, 66, 67, 68, 69, 70, 72, 73, 78, 79, 80, 86, 87, 88, 94, 95, 96, 101, 102, 103, 104, 107, 170, 171, 176, 177, 179, 205, 206, 213, 259, 260, 263, 264, 275, 276, 278, 281, 302, 303 }

C grade: { 146, 148, 149, 156, 157, 158, 159, 160, 162, 180, 197, 232, 241, 252, 294, 304, 305, 312, 313, 320 }

F grade: { 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 138, 139, 140, 141, 142, 143, 144, 145, 147, 150, 151, 152, 153, 154, 155, 161, 297, 298, 299, 300, 301, 306, 307, 308, 309, 310, 311, 314, 315, 316, 317, 318, 319, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348 }

2.1.4 Maxima

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 27, 28, 29, 30, 31, 32, 35, 36, 37, 38, 39, 40, 43, 44, 45, 46, 47, 48, 53, 54, 55, 56, 62, 63, 64, 65, 74, 75, 76, 77, 81, 82, 83, 84, 85, 89, 90, 91, 92, 93, 98, 99, 100, 108, 137, 163 }

B grade: { 26, 33, 34, 41, 42, 72, 73, 97 }

C grade: { }

F grade: { 49, 50, 51, 52, 57, 58, 59, 60, 61, 66, 67, 68, 69, 70, 71, 78, 79, 80, 86, 87, 88, 94, 95, 96, 101, 102, 103, 104, 105, 106, 107, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349 }

2.1.5 FriCAS

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 16, 17, 20, 21, 22, 23, 24, 25, 30, 31, 32, 39, 43, 44, 45, 46, 47, 48, 49, 53, 54, 55, 56, 57, 58, 62, 63, 64, 65, 66, 67, 71, 74, 75, 76, 77, 81, 82, 83, 84, 85, 89, 90, 91, 92, 93, 97, 98, 99, 100, 105, 106, 137, 163, 164, 165, 244, 349 }

B grade: { 12, 13, 18, 19, 26, 27, 28, 29, 33, 34, 35, 36, 37, 38, 40, 41, 42, 50, 51, 52, 59, 60, 61, 68, 69, 70, 72, 73, 78, 79, 80, 86, 87, 88, 94, 95, 96, 101, 102, 103, 104, 107, 108, 146, 147, 148, 149, 160,

162, 180, 224, 231, 251, 304, 305, 306, 307, 308, 309, 310, 311, 312, 314, 315, 316, 317, 318, 319, 320
}

C grade: { 313 }

F grade: { 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 138, 139, 140, 141, 142, 143, 144, 145, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 161, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 225, 226, 227, 228, 229, 230, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 245, 246, 247, 248, 249, 250, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348 }

2.1.6 Sympy

A grade: { 1, 2, 3, 4, 7, 8, 9, 10, 14, 15, 16, 39, 43, 44, 47, 48, 56, 65, 74, 75, 76, 77, 84, 85, 109, 110, 111, 116, 117, 118, 122, 123, 124, 125, 131, 138, 180, 184, 185, 186, 188, 189, 190, 218, 220, 222, 224, 225, 226, 227, 231, 235, 236, 237, 246, 254, 256, 291 }

B grade: { 5, 6, 11, 12, 13, 17, 18, 19, 20, 21, 22, 23, 24, 27, 28, 29, 30, 31, 35, 36, 37, 38, 45, 46, 53, 54, 55, 62, 63, 64, 92, 93, 99, 100, 108, 187, 221, 234 }

C grade: { 241, 245, 247, 342, 343, 344, 345 }

F grade: { 25, 26, 32, 33, 34, 40, 41, 42, 49, 50, 51, 52, 57, 58, 59, 60, 61, 66, 67, 68, 69, 70, 71, 72, 73, 78, 79, 80, 81, 82, 83, 86, 87, 88, 89, 90, 91, 94, 95, 96, 97, 98, 101, 102, 103, 104, 105, 106, 107, 112, 113, 114, 115, 119, 120, 121, 126, 127, 128, 129, 130, 132, 133, 134, 135, 136, 137, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 181, 182, 183, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 219, 223, 228, 229, 230, 232, 233, 238, 239, 240, 242, 243, 244, 248, 249, 250, 251, 252, 253, 255, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 346, 347, 348, 349 }

2.1.7 Giac

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 53, 54, 55, 56, 62, 63, 64, 65, 74, 75, 76, 77, 78, 81, 82, 83, 84, 85, 86, 89, 90, 91, 92, 93, 97, 98, 99, 100, 101, 105, 107, 108, 180 }

B grade: { 50, 51, 52, 58, 59, 60, 61, 67, 68, 69, 70, 71, 72, 73, 79, 80, 87, 88, 94, 95, 96, 102, 103, 104, 106, 224, 231 }

C grade: { }

F grade: { 49, 57, 66, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 225, 226, 227, 228, 229, 230, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294,

295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349 }

2.1.8 Mupad

A grade: { }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 48, 56, 65, 71, 72, 73, 76, 77, 84, 85, 92, 93, 97, 98, 99, 100, 105, 106, 108, 137, 345, 349 }

C grade: { }

F grade: { 45, 46, 47, 49, 50, 51, 52, 53, 54, 55, 57, 58, 59, 60, 61, 62, 63, 64, 66, 67, 68, 69, 70, 74, 75, 78, 79, 80, 81, 82, 83, 86, 87, 88, 89, 90, 91, 94, 95, 96, 101, 102, 103, 104, 107, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 346, 347, 348 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$

Problem 1	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	94	97	96	98	107	98	88
normalized size	1	1.00	1.00	1.03	1.02	1.04	1.14	1.04	0.94
time (sec)	N/A	0.064	0.021	0.002	1.355	0.514	0.085	0.567	4.775
Problem 2	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	70	73	70	73	76	73	65
normalized size	1	1.00	1.00	1.04	1.00	1.04	1.09	1.04	0.93
time (sec)	N/A	0.042	0.015	0.002	1.372	0.571	0.082	0.565	4.753
Problem 3	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	50	49	48	50	53	50	48
normalized size	1	1.00	1.00	0.98	0.96	1.00	1.06	1.00	0.96
time (sec)	N/A	0.027	0.011	0.001	1.352	0.542	0.074	0.570	0.047
Problem 4	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	28	25	24	26	26	26	25
normalized size	1	1.00	1.00	0.89	0.86	0.93	0.93	0.93	0.89
time (sec)	N/A	0.013	0.005	0.001	1.349	0.636	0.065	0.561	0.036
Problem 5	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	40	45	34	99	82	34	31
normalized size	1	1.00	1.00	1.12	0.85	2.48	2.05	0.85	0.78
time (sec)	N/A	0.020	0.023	0.007	2.998	0.789	0.280	0.571	0.060

Problem 6	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	63	68	57	182	112	57	51
normalized size	1	1.00	1.00	1.08	0.90	2.89	1.78	0.90	0.81
time (sec)	N/A	0.020	0.047	0.008	3.115	0.664	0.392	0.583	5.006
Problem 7	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	82	90	92	300	150	78	82
normalized size	1	1.00	0.89	0.98	1.00	3.26	1.63	0.85	0.89
time (sec)	N/A	0.031	0.059	0.010	2.963	0.664	0.541	0.587	5.061
Problem 8	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	122	125	124	131	136	131	116
normalized size	1	1.00	1.00	1.02	1.02	1.07	1.11	1.07	0.95
time (sec)	N/A	0.074	0.023	0.002	1.395	0.464	0.091	0.564	4.945
Problem 9	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	82	87	82	91	97	91	75
normalized size	1	1.00	1.00	1.06	1.00	1.11	1.18	1.11	0.91
time (sec)	N/A	0.046	0.016	0.000	1.327	0.416	0.081	0.572	0.047
Problem 10	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	50	49	48	50	53	50	48
normalized size	1	1.00	1.00	0.98	0.96	1.00	1.06	1.00	0.96
time (sec)	N/A	0.028	0.007	0.000	1.320	0.571	0.073	0.569	0.046
Problem 11	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	59	95	68	179	172	72	90
normalized size	1	1.00	0.94	1.51	1.08	2.84	2.73	1.14	1.43
time (sec)	N/A	0.043	0.050	0.003	2.886	0.685	0.420	0.585	0.088

Problem 12	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	89	129	96	302	236	95	124
normalized size	1	1.00	1.09	1.57	1.17	3.68	2.88	1.16	1.51
time (sec)	N/A	0.099	0.060	0.009	2.840	0.672	0.698	0.570	5.021
Problem 13	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	121	147	138	449	223	126	130
normalized size	1	1.00	1.04	1.27	1.19	3.87	1.92	1.09	1.12
time (sec)	N/A	0.072	0.097	0.009	3.084	0.686	0.984	0.585	5.030
Problem 14	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	154	154	161	177	167	187	189	187	152
normalized size	1	1.00	1.05	1.15	1.08	1.21	1.23	1.21	0.99
time (sec)	N/A	0.103	0.030	0.003	1.378	0.688	0.100	0.567	4.905
Problem 15	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	122	125	124	131	136	131	116
normalized size	1	1.00	1.00	1.02	1.02	1.07	1.11	1.07	0.95
time (sec)	N/A	0.070	0.022	0.002	1.362	0.474	0.091	0.591	4.875
Problem 16	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	70	73	70	73	76	73	65
normalized size	1	1.00	1.00	1.04	1.00	1.04	1.09	1.04	0.93
time (sec)	N/A	0.044	0.012	0.001	1.294	0.529	0.078	0.565	0.034
Problem 17	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	93	161	122	290	238	130	145
normalized size	1	1.00	0.95	1.64	1.24	2.96	2.43	1.33	1.48
time (sec)	N/A	0.064	0.061	0.004	3.025	0.545	0.573	0.577	4.870

Problem 18	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	107	205	147	444	314	152	181
normalized size	1	1.00	1.00	1.92	1.37	4.15	2.93	1.42	1.69
time (sec)	N/A	0.096	0.059	0.010	2.991	0.618	1.054	0.581	0.100
Problem 19	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	130	141	266	187	618	422	180	240
normalized size	1	1.00	1.08	2.05	1.44	4.75	3.25	1.38	1.85
time (sec)	N/A	0.165	0.081	0.011	2.979	0.691	1.813	0.587	4.956
Problem 20	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	142	136	246	187	428	326	198	216
normalized size	1	1.00	0.96	1.73	1.32	3.01	2.30	1.39	1.52
time (sec)	N/A	0.093	0.091	0.006	3.033	0.647	0.757	0.579	4.861
Problem 21	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	92	161	122	292	238	129	146
normalized size	1	1.00	0.94	1.64	1.24	2.98	2.43	1.32	1.49
time (sec)	N/A	0.059	0.067	0.004	2.911	0.750	0.591	0.576	0.076
Problem 22	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	59	95	69	181	172	72	90
normalized size	1	1.00	0.94	1.51	1.10	2.87	2.73	1.14	1.43
time (sec)	N/A	0.041	0.051	0.004	2.934	0.562	0.435	0.568	4.902
Problem 23	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	40	45	33	98	82	33	32
normalized size	1	1.00	1.03	1.15	0.85	2.51	2.10	0.85	0.82
time (sec)	N/A	0.015	0.026	0.005	2.991	0.711	0.276	0.571	0.055

Problem 24	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	61	55	54	292	712	54	135
normalized size	1	1.00	0.87	0.79	0.77	4.17	10.17	0.77	1.93
time (sec)	N/A	0.027	0.045	0.008	2.985	0.846	2.798	0.585	0.320
Problem 25	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	109	109	95	144	133	711	0	122	3637
normalized size	1	1.00	0.87	1.32	1.22	6.52	0.00	1.12	33.37
time (sec)	N/A	0.084	0.171	0.011	3.045	0.963	0.000	0.569	5.688
Problem 26	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	160	160	158	310	277	1585	0	217	6033
normalized size	1	1.00	0.99	1.94	1.73	9.91	0.00	1.36	37.71
time (sec)	N/A	0.192	0.238	0.013	3.075	2.142	0.000	0.578	6.869
Problem 27	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	192	192	192	402	294	810	502	306	386
normalized size	1	1.00	1.00	2.09	1.53	4.22	2.61	1.59	2.01
time (sec)	N/A	0.163	0.098	0.012	3.026	0.550	1.910	0.579	5.024
Problem 28	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	142	142	296	213	612	403	220	261
normalized size	1	1.00	1.00	2.08	1.50	4.31	2.84	1.55	1.84
time (sec)	N/A	0.120	0.090	0.012	2.997	0.677	1.475	0.576	5.054
Problem 29	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	106	205	147	442	314	152	182
normalized size	1	1.00	1.00	1.93	1.39	4.17	2.96	1.43	1.72
time (sec)	N/A	0.093	0.062	0.010	2.990	0.695	1.075	0.568	0.102

Problem 30	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	88	129	95	297	236	94	124
normalized size	1	1.00	1.07	1.57	1.16	3.62	2.88	1.15	1.51
time (sec)	N/A	0.104	0.063	0.009	2.839	0.642	0.723	0.584	5.062
Problem 31	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	63	68	57	181	112	57	51
normalized size	1	1.00	1.00	1.08	0.90	2.87	1.78	0.90	0.81
time (sec)	N/A	0.021	0.047	0.009	3.114	0.725	0.399	0.572	5.042
Problem 32	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	108	109	144	132	699	0	121	3649
normalized size	1	1.00	1.01	1.33	1.22	6.47	0.00	1.12	33.79
time (sec)	N/A	0.081	0.146	0.010	2.915	0.804	0.000	0.576	5.766
Problem 33	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	167	136	238	294	1681	0	232	6183
normalized size	1	1.00	0.81	1.43	1.76	10.07	0.00	1.39	37.02
time (sec)	N/A	0.201	0.319	0.015	3.140	1.644	0.000	0.583	6.875
Problem 34	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	230	230	197	403	529	3239	0	332	8649
normalized size	1	1.00	0.86	1.75	2.30	14.08	0.00	1.44	37.60
time (sec)	N/A	0.309	0.419	0.017	3.409	6.075	0.000	0.582	7.793
Problem 35	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	196	196	196	484	334	1044	615	340	409
normalized size	1	1.00	1.00	2.47	1.70	5.33	3.14	1.73	2.09
time (sec)	N/A	0.227	0.126	0.015	3.037	0.620	4.406	0.580	5.023

Problem 36	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	160	160	160	367	253	817	515	254	318
normalized size	1	1.00	1.00	2.29	1.58	5.11	3.22	1.59	1.99
time (sec)	N/A	0.197	0.097	0.013	3.073	0.633	2.811	0.583	0.135
Problem 37	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	130	139	266	185	606	422	178	240
normalized size	1	1.00	1.07	2.05	1.42	4.66	3.25	1.37	1.85
time (sec)	N/A	0.166	0.084	0.011	2.938	0.620	1.890	0.571	5.052
Problem 38	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	124	147	138	449	223	126	130
normalized size	1	1.00	1.07	1.27	1.19	3.87	1.92	1.09	1.12
time (sec)	N/A	0.077	0.096	0.010	3.010	0.532	1.049	0.569	5.023
Problem 39	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	84	89	92	301	150	78	81
normalized size	1	1.00	0.91	0.97	1.00	3.27	1.63	0.85	0.88
time (sec)	N/A	0.033	0.064	0.009	3.002	0.697	0.585	0.576	5.016
Problem 40	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	161	161	158	309	278	1587	0	218	6033
normalized size	1	1.00	0.98	1.92	1.73	9.86	0.00	1.35	37.47
time (sec)	N/A	0.197	0.281	0.012	3.161	1.475	0.000	0.578	6.892
Problem 41	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	236	236	197	403	530	3251	0	333	8635
normalized size	1	1.00	0.83	1.71	2.25	13.78	0.00	1.41	36.59
time (sec)	N/A	0.311	0.418	0.014	3.177	7.246	0.000	0.612	7.855

Problem 42	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	315	315	233	568	820	5070	0	574	11150
normalized size	1	1.00	0.74	1.80	2.60	16.10	0.00	1.82	35.40
time (sec)	N/A	0.451	0.928	0.018	3.395	19.469	0.000	0.620	8.555
Problem 43	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	24	23	33	33	31	20	31
normalized size	1	1.00	0.71	0.68	0.97	0.97	0.91	0.59	0.91
time (sec)	N/A	0.010	0.008	0.007	1.323	0.761	0.132	0.578	4.995
Problem 44	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	41	33	48	67	46	54	47
normalized size	1	1.00	0.87	0.70	1.02	1.43	0.98	1.15	1.00
time (sec)	N/A	0.016	0.013	0.007	2.994	0.808	0.172	0.580	0.042
Problem 45	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	231	231	181	310	281	398	484	201	-1
normalized size	1	1.00	0.78	1.34	1.22	1.72	2.10	0.87	-0.00
time (sec)	N/A	0.179	5.114	0.016	1.395	0.904	20.381	0.642	0.000
Problem 46	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	B	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	149	149	160	190	168	264	291	129	-1
normalized size	1	1.00	1.07	1.28	1.13	1.77	1.95	0.87	-0.01
time (sec)	N/A	0.088	2.673	0.010	1.347	0.971	11.393	0.607	0.000
Problem 47	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	85	96	81	158	144	70	-1
normalized size	1	1.00	0.98	1.10	0.93	1.82	1.66	0.80	-0.01
time (sec)	N/A	0.028	0.157	0.005	1.350	0.493	5.681	0.600	0.000

Problem 48	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	49	36	28	94	41	37	35
normalized size	1	1.00	1.07	0.78	0.61	2.04	0.89	0.80	0.76
time (sec)	N/A	0.010	0.021	0.000	1.364	0.545	1.855	0.576	4.712
Problem 49	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	84	932	0	596	0	0	-1
normalized size	1	1.00	1.02	11.37	0.00	7.27	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.044	0.045	0.000	0.696	0.000	0.000	0.000
Problem 50	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	F	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	165	2521	0	369	0	217	-1
normalized size	1	1.00	2.01	30.74	0.00	4.50	0.00	2.65	-0.01
time (sec)	N/A	0.034	0.234	0.022	0.000	0.676	0.000	1.641	0.000
Problem 51	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	149	149	176	5101	0	698	0	487	-1
normalized size	1	1.00	1.18	34.23	0.00	4.68	0.00	3.27	-0.01
time (sec)	N/A	0.094	0.566	0.025	0.000	1.409	0.000	3.746	0.000
Problem 52	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	208	208	227	7922	0	1220	0	958	-1
normalized size	1	1.00	1.09	38.09	0.00	5.87	0.00	4.61	-0.00
time (sec)	N/A	0.213	0.990	0.029	0.000	1.645	0.000	2.870	0.000
Problem 53	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	272	272	220	393	364	502	665	260	-1
normalized size	1	1.00	0.81	1.44	1.34	1.85	2.44	0.96	-0.00
time (sec)	N/A	0.218	5.129	0.016	1.496	1.210	52.753	0.662	0.000

Problem 54	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	B	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	196	196	157	249	227	344	440	175	-1
normalized size	1	1.00	0.80	1.27	1.16	1.76	2.24	0.89	-0.01
time (sec)	N/A	0.116	2.712	0.007	1.424	0.782	29.419	0.660	0.000
Problem 55	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	118	118	109	131	116	210	253	103	-1
normalized size	1	1.00	0.92	1.11	0.98	1.78	2.14	0.87	-0.01
time (sec)	N/A	0.040	0.204	0.004	1.372	0.757	14.712	0.613	0.000
Problem 56	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	65	51	43	124	70	49	37
normalized size	1	1.00	1.00	0.78	0.66	1.91	1.08	0.75	0.57
time (sec)	N/A	0.016	0.090	0.001	1.336	0.580	2.915	0.612	4.709
Problem 57	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	110	1845	0	721	0	0	-1
normalized size	1	1.00	0.97	16.33	0.00	6.38	0.00	0.00	-0.01
time (sec)	N/A	0.108	0.200	0.016	0.000	0.831	0.000	0.000	0.000
Problem 58	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	142	4621	0	907	0	317	-1
normalized size	1	1.00	1.08	35.27	0.00	6.92	0.00	2.42	-0.01
time (sec)	N/A	0.090	0.128	0.021	0.000	0.967	0.000	0.682	0.000
Problem 59	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	163	9059	0	526	0	451	-1
normalized size	1	1.00	1.44	80.17	0.00	4.65	0.00	3.99	-0.01
time (sec)	N/A	0.057	0.693	0.025	0.000	0.785	0.000	3.723	0.000

Problem 60	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	199	199	247	13766	0	972	0	919	-1
normalized size	1	1.00	1.24	69.18	0.00	4.88	0.00	4.62	-0.01
time (sec)	N/A	0.115	0.817	0.032	0.000	1.231	0.000	2.893	0.000
Problem 61	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	300	300	362	18791	0	1604	0	1557	-1
normalized size	1	1.00	1.21	62.64	0.00	5.35	0.00	5.19	-0.00
time (sec)	N/A	0.365	1.385	0.046	0.000	3.675	0.000	9.597	0.000
Problem 62	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	349	349	270	476	447	608	796	321	-1
normalized size	1	1.00	0.77	1.36	1.28	1.74	2.28	0.92	-0.00
time (sec)	N/A	0.247	5.179	0.019	1.472	1.585	102.666	0.680	0.000
Problem 63	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	B	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	241	241	158	308	286	420	537	221	-1
normalized size	1	1.00	0.66	1.28	1.19	1.74	2.23	0.92	-0.00
time (sec)	N/A	0.148	2.802	0.010	1.388	0.870	58.630	0.656	0.000
Problem 64	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	149	149	130	166	151	260	316	135	-1
normalized size	1	1.00	0.87	1.11	1.01	1.74	2.12	0.91	-0.01
time (sec)	N/A	0.052	0.228	0.006	1.401	0.843	29.528	0.632	0.000
Problem 65	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	76	66	58	146	97	63	37
normalized size	1	1.00	0.90	0.79	0.69	1.74	1.15	0.75	0.44
time (sec)	N/A	0.023	0.112	0.003	1.328	0.629	4.282	0.608	4.693

Problem 66	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	157	140	3053	0	935	0	0	-1
normalized size	1	1.00	0.89	19.45	0.00	5.96	0.00	0.00	-0.01
time (sec)	N/A	0.199	0.122	0.019	0.000	3.228	0.000	0.000	0.000
Problem 67	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	175	175	144	7345	0	1236	0	405	-1
normalized size	1	1.00	0.82	41.97	0.00	7.06	0.00	2.31	-0.01
time (sec)	N/A	0.226	0.164	0.023	0.000	2.202	0.000	0.689	0.000
Problem 68	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	194	184	14133	0	1517	0	659	-1
normalized size	1	1.00	0.95	72.85	0.00	7.82	0.00	3.40	-0.01
time (sec)	N/A	0.194	0.203	0.030	0.000	1.297	0.000	0.733	0.000
Problem 69	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	144	144	201	21220	0	706	0	846	-1
normalized size	1	1.00	1.40	147.36	0.00	4.90	0.00	5.88	-0.01
time (sec)	N/A	0.073	0.807	0.040	0.000	0.957	0.000	2.858	0.000
Problem 70	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	249	249	306	28625	0	1258	0	1448	-1
normalized size	1	1.00	1.23	114.96	0.00	5.05	0.00	5.82	-0.00
time (sec)	N/A	0.137	1.094	0.053	0.000	2.024	0.000	9.315	0.000
Problem 71	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	33	0	42	0	95	83
normalized size	1	1.00	1.00	1.10	0.00	1.40	0.00	3.17	2.77
time (sec)	N/A	0.014	0.029	0.023	0.000	0.641	0.000	0.608	0.394

Problem 72	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	64	84	59	67	0	70	59
normalized size	1	1.00	2.37	3.11	2.19	2.48	0.00	2.59	2.19
time (sec)	N/A	0.014	0.026	0.012	2.978	0.533	0.000	0.620	0.167
Problem 73	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	187	110	74	0	118	85
normalized size	1	1.00	1.00	7.48	4.40	2.96	0.00	4.72	3.40
time (sec)	N/A	0.013	0.011	0.043	3.074	0.562	0.000	0.611	5.347
Problem 74	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	169	169	140	228	199	300	400	150	-1
normalized size	1	1.00	0.83	1.35	1.18	1.78	2.37	0.89	-0.01
time (sec)	N/A	0.145	5.095	0.014	1.360	0.547	13.273	0.639	0.000
Problem 75	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	A	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	108	108	160	131	109	192	238	90	-1
normalized size	1	1.00	1.48	1.21	1.01	1.78	2.20	0.83	-0.01
time (sec)	N/A	0.056	2.489	0.008	1.469	0.566	6.943	0.619	0.000
Problem 76	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	57	62	47	113	126	49	86
normalized size	1	1.00	0.98	1.07	0.81	1.95	2.17	0.84	1.48
time (sec)	N/A	0.017	0.021	0.006	1.329	0.687	2.780	0.599	5.515
Problem 77	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	21	13	59	17	23	20
normalized size	1	1.00	1.00	0.84	0.52	2.36	0.68	0.92	0.80
time (sec)	N/A	0.006	0.004	0.002	1.284	0.557	0.997	0.596	0.121

Problem 78	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	49	300	0	241	0	70	-1
normalized size	1	1.00	1.00	6.12	0.00	4.92	0.00	1.43	-0.02
time (sec)	N/A	0.022	0.017	0.015	0.000	1.039	0.000	0.603	0.000
Problem 79	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	126	809	0	463	0	242	-1
normalized size	1	1.00	1.25	8.01	0.00	4.58	0.00	2.40	-0.01
time (sec)	N/A	0.049	0.302	0.019	0.000	1.045	0.000	0.627	0.000
Problem 80	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	192	1815	0	864	0	538	-1
normalized size	1	1.00	1.18	11.13	0.00	5.30	0.00	3.30	-0.01
time (sec)	N/A	0.119	0.668	0.021	0.000	1.016	0.000	3.505	0.000
Problem 81	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	257	257	172	340	311	584	0	235	-1
normalized size	1	1.00	0.67	1.32	1.21	2.27	0.00	0.91	-0.00
time (sec)	N/A	0.259	5.203	0.019	1.382	0.943	0.000	0.670	0.000
Problem 82	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	169	169	122	219	197	416	0	157	-1
normalized size	1	1.00	0.72	1.30	1.17	2.46	0.00	0.93	-0.01
time (sec)	N/A	0.197	5.100	0.008	1.397	0.997	0.000	0.644	0.000
Problem 83	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	90	105	160	123	108	276	0	92	-1
normalized size	1	1.17	1.78	1.37	1.20	3.07	0.00	1.02	-0.01
time (sec)	N/A	0.062	2.529	0.008	1.377	0.712	0.000	0.632	0.000

Problem 84	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	70	54	46	167	60	50	53
normalized size	1	1.00	1.30	1.00	0.85	3.09	1.11	0.93	0.98
time (sec)	N/A	0.017	0.061	0.005	1.302	0.759	5.168	0.643	5.117
Problem 85	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	23	17	14	14
normalized size	1	1.00	1.00	0.94	0.88	1.44	1.06	0.88	0.88
time (sec)	N/A	0.002	0.004	0.003	1.324	0.685	0.614	0.596	0.040
Problem 86	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	309	618	0	441	0	107	-1
normalized size	1	1.00	3.91	7.82	0.00	5.58	0.00	1.35	-0.01
time (sec)	N/A	0.039	0.720	0.017	0.000	1.071	0.000	0.619	0.000
Problem 87	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	143	143	758	1439	0	864	0	318	-1
normalized size	1	1.00	5.30	10.06	0.00	6.04	0.00	2.22	-0.01
time (sec)	N/A	0.109	2.673	0.022	0.000	1.398	0.000	1.902	0.000
Problem 88	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F(-1)	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	225	225	1392	2919	0	1482	0	643	-1
normalized size	1	1.00	6.19	12.97	0.00	6.59	0.00	2.86	-0.00
time (sec)	N/A	0.243	5.017	0.026	0.000	2.781	0.000	2.699	0.000
Problem 89	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	255	255	157	351	392	684	0	237	-1
normalized size	1	1.00	0.62	1.38	1.54	2.68	0.00	0.93	-0.00
time (sec)	N/A	0.245	5.168	0.017	1.442	1.439	0.000	0.679	0.000

Problem 90	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	172	172	125	228	254	486	0	158	-1
normalized size	1	1.00	0.73	1.33	1.48	2.83	0.00	0.92	-0.01
time (sec)	N/A	0.157	5.101	0.009	1.493	0.669	0.000	0.682	0.000
Problem 91	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	105	105	214	136	147	318	0	103	-1
normalized size	1	1.00	2.04	1.30	1.40	3.03	0.00	0.98	-0.01
time (sec)	N/A	0.050	4.146	0.007	1.389	0.473	0.000	0.632	0.000
Problem 92	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	37	34	68	54	144	40	33
normalized size	1	1.00	0.79	0.72	1.45	1.15	3.06	0.85	0.70
time (sec)	N/A	0.010	0.016	0.004	1.366	0.621	11.030	0.618	4.785
Problem 93	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	29	26	31	47	95	27	28
normalized size	1	1.00	0.74	0.67	0.79	1.21	2.44	0.69	0.72
time (sec)	N/A	0.006	0.006	0.000	1.332	0.801	0.821	0.609	4.754
Problem 94	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	775	1070	0	764	0	320	-1
normalized size	1	1.00	6.35	8.77	0.00	6.26	0.00	2.62	-0.01
time (sec)	N/A	0.103	2.750	0.020	0.000	0.979	0.000	0.639	0.000
Problem 95	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	202	202	170	2371	0	1440	0	620	-1
normalized size	1	1.00	0.84	11.74	0.00	7.13	0.00	3.07	-0.00
time (sec)	N/A	0.228	5.498	0.027	0.000	2.275	0.000	2.070	0.000

Problem 96	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	313	313	221	4495	0	2250	0	1010	-1
normalized size	1	1.00	0.71	14.36	0.00	7.19	0.00	3.23	-0.00
time (sec)	N/A	0.400	5.662	0.029	0.000	8.343	0.000	3.760	0.000
Problem 97	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	224	224	163	190	465	229	0	218	326
normalized size	1	1.00	0.73	0.85	2.08	1.02	0.00	0.97	1.46
time (sec)	N/A	0.101	0.103	0.007	1.513	1.022	0.000	0.677	5.152
Problem 98	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	174	107	115	249	151	0	138	176
normalized size	1	1.00	0.61	0.66	1.43	0.87	0.00	0.79	1.01
time (sec)	N/A	0.071	0.068	0.008	1.500	0.639	0.000	0.641	4.987
Problem 99	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	59	57	103	87	566	72	87
normalized size	1	1.00	0.65	0.63	1.13	0.96	6.22	0.79	0.96
time (sec)	N/A	0.029	0.023	0.004	1.453	0.586	27.986	0.618	4.849
Problem 100	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	29	26	31	47	95	27	28
normalized size	1	1.00	0.74	0.67	0.79	1.21	2.44	0.69	0.72
time (sec)	N/A	0.006	0.009	0.003	1.338	0.616	0.826	0.603	4.787
Problem 101	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	236	628	0	442	0	107	-1
normalized size	1	1.00	2.99	7.95	0.00	5.59	0.00	1.35	-0.01
time (sec)	N/A	0.046	2.716	0.042	0.000	0.723	0.000	0.617	0.000

Problem 102	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	405	823	0	459	0	225	-1
normalized size	1	1.00	4.05	8.23	0.00	4.59	0.00	2.25	-0.01
time (sec)	N/A	0.055	0.779	0.025	0.000	0.975	0.000	0.623	0.000
Problem 103	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	149	149	130	5177	0	698	0	487	-1
normalized size	1	1.00	0.87	34.74	0.00	4.68	0.00	3.27	-0.01
time (sec)	N/A	0.079	5.169	0.033	0.000	1.006	0.000	3.132	0.000
Problem 104	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	199	199	179	13964	0	972	0	919	-1
normalized size	1	1.00	0.90	70.17	0.00	4.88	0.00	4.62	-0.01
time (sec)	N/A	0.113	5.270	0.049	0.000	1.282	0.000	2.812	0.000
Problem 105	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	0	27	0	18	18
normalized size	1	1.00	1.00	0.95	0.00	1.35	0.00	0.90	0.90
time (sec)	N/A	0.005	0.008	0.003	0.000	0.563	0.000	0.604	4.771
Problem 106	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	28	0	23	0	51	79
normalized size	1	1.00	1.00	1.12	0.00	0.92	0.00	2.04	3.16
time (sec)	N/A	0.008	0.006	0.006	0.000	0.550	0.000	0.604	0.368
Problem 107	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	49	306	0	241	0	70	-1
normalized size	1	1.00	1.00	6.24	0.00	4.92	0.00	1.43	-0.02
time (sec)	N/A	0.020	0.015	0.010	0.000	0.689	0.000	0.602	0.000

Problem 108	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	44	31	25	27
normalized size	1	1.00	1.00	0.93	0.87	2.93	2.07	1.67	1.80
time (sec)	N/A	0.004	0.013	0.006	2.782	0.629	4.681	0.574	0.039
Problem 109	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	648	648	99	0	0	0	136	0	-1
normalized size	1	1.00	0.15	0.00	0.00	0.00	0.21	0.00	-0.00
time (sec)	N/A	0.597	5.051	0.311	0.000	0.605	4.374	0.000	0.000
Problem 110	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	617	617	176	0	0	0	99	0	-1
normalized size	1	1.00	0.29	0.00	0.00	0.00	0.16	0.00	-0.00
time (sec)	N/A	0.426	3.378	0.303	0.000	0.646	3.298	0.000	0.000
Problem 111	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	588	588	62	0	0	0	63	0	-1
normalized size	1	1.00	0.11	0.00	0.00	0.00	0.11	0.00	-0.00
time (sec)	N/A	0.369	0.077	0.296	0.000	0.615	2.428	0.000	0.000
Problem 112	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	740	740	162	0	0	0	0	0	-1
normalized size	1	1.00	0.22	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.321	0.160	0.314	0.000	0.000	0.000	0.000	0.000
Problem 113	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	584	584	86	0	0	0	0	0	-1
normalized size	1	1.00	0.15	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.374	0.061	0.316	0.000	0.865	0.000	0.000	0.000

Problem 114	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	818	818	252	0	0	0	0	0	-1
normalized size	1	1.00	0.31	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.560	0.325	0.320	0.000	0.000	0.000	0.000	0.000
Problem 115	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	849	849	265	0	0	0	0	0	-1
normalized size	1	1.00	0.31	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.651	0.192	0.326	0.000	0.000	0.000	0.000	0.000
Problem 116	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	668	668	110	0	0	0	139	0	-1
normalized size	1	1.00	0.16	0.00	0.00	0.00	0.21	0.00	-0.00
time (sec)	N/A	0.571	5.053	0.301	0.000	1.071	5.589	0.000	0.000
Problem 117	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	637	637	173	0	0	0	131	0	-1
normalized size	1	1.00	0.27	0.00	0.00	0.00	0.21	0.00	-0.00
time (sec)	N/A	0.484	3.007	0.299	0.000	0.945	4.895	0.000	0.000
Problem 118	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	608	608	68	0	0	0	100	0	-1
normalized size	1	1.00	0.11	0.00	0.00	0.00	0.16	0.00	-0.00
time (sec)	N/A	0.420	0.064	0.300	0.000	0.941	3.688	0.000	0.000
Problem 119	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	765	765	231	0	0	0	0	0	-1
normalized size	1	1.00	0.30	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.444	0.151	0.313	0.000	0.000	0.000	0.000	0.000

Problem 120	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	775	775	235	0	0	0	0	0	-1
normalized size	1	1.00	0.30	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.437	0.147	0.333	0.000	0.000	0.000	0.000	0.000
Problem 121	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	815	815	252	0	0	0	0	0	-1
normalized size	1	1.00	0.31	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.613	0.235	0.363	0.000	0.000	0.000	0.000	0.000
Problem 122	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	659	659	98	0	0	0	165	0	-1
normalized size	1	1.00	0.15	0.00	0.00	0.00	0.25	0.00	-0.00
time (sec)	N/A	0.527	5.062	0.409	0.000	1.106	5.404	0.000	0.000
Problem 123	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	628	628	88	0	0	0	129	0	-1
normalized size	1	1.00	0.14	0.00	0.00	0.00	0.21	0.00	-0.00
time (sec)	N/A	0.440	5.051	0.298	0.000	0.947	4.379	0.000	0.000
Problem 124	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	597	597	158	0	0	0	94	0	-1
normalized size	1	1.00	0.26	0.00	0.00	0.00	0.16	0.00	-0.00
time (sec)	N/A	0.381	4.495	0.300	0.000	0.599	3.347	0.000	0.000
Problem 125	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	568	568	62	0	0	0	60	0	-1
normalized size	1	1.00	0.11	0.00	0.00	0.00	0.11	0.00	-0.00
time (sec)	N/A	0.305	0.027	0.300	0.000	0.839	2.008	0.000	0.000

Problem 126	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	204	204	162	0	0	0	0	0	-1
normalized size	1	1.00	0.79	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.031	0.047	0.310	0.000	0.000	0.000	0.000	0.000
Problem 127	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	787	787	234	0	0	0	0	0	-1
normalized size	1	1.00	0.30	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.439	0.162	0.313	0.000	0.000	0.000	0.000	0.000
Problem 128	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	818	818	255	0	0	0	0	0	-1
normalized size	1	1.00	0.31	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.543	0.172	0.326	0.000	0.000	0.000	0.000	0.000
Problem 129	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	623	623	76	0	0	0	0	0	-1
normalized size	1	1.00	0.12	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.435	5.057	0.321	0.000	0.874	0.000	0.000	0.000
Problem 130	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	592	592	62	0	0	0	0	0	-1
normalized size	1	1.00	0.10	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.368	5.048	0.324	0.000	0.592	0.000	0.000	0.000
Problem 131	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	561	561	53	0	0	0	60	0	-1
normalized size	1	1.00	0.09	0.00	0.00	0.00	0.11	0.00	-0.00
time (sec)	N/A	0.304	0.020	0.312	0.000	0.735	6.869	0.000	0.000

Problem 132	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	776	776	226	0	0	0	0	0	-1
normalized size	1	1.00	0.29	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.443	0.142	0.321	0.000	0.000	0.000	0.000	0.000
Problem 133	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	807	807	236	0	0	0	0	0	-1
normalized size	1	1.00	0.29	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.534	0.180	0.330	0.000	0.000	0.000	0.000	0.000
Problem 134	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	849	849	256	0	0	0	0	0	-1
normalized size	1	1.00	0.30	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.653	0.241	0.328	0.000	0.000	0.000	0.000	0.000
Problem 135	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	653	653	96	0	0	0	0	0	-1
normalized size	1	1.00	0.15	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.512	5.079	0.316	0.000	1.130	0.000	0.000	0.000
Problem 136	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	596	596	83	0	0	0	0	0	-1
normalized size	1	1.00	0.14	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.426	5.075	0.319	0.000	1.027	0.000	0.000	0.000
Problem 137	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	24	24	33	42	0	0	27
normalized size	1	1.00	0.55	0.55	0.75	0.95	0.00	0.00	0.61
time (sec)	N/A	0.017	5.027	0.005	1.890	0.549	0.000	0.000	4.785

Problem 138	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	590	590	74	0	0	0	60	0	-1
normalized size	1	1.00	0.13	0.00	0.00	0.00	0.10	0.00	-0.00
time (sec)	N/A	0.366	0.040	0.312	0.000	0.587	12.990	0.000	0.000
Problem 139	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	796	796	248	0	0	0	0	0	-1
normalized size	1	1.00	0.31	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.537	0.210	0.329	0.000	0.000	0.000	0.000	0.000
Problem 140	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	827	827	259	0	0	0	0	0	-1
normalized size	1	1.00	0.31	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.634	0.243	0.334	0.000	0.000	0.000	0.000	0.000
Problem 141	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	252	252	163	0	0	0	0	0	-1
normalized size	1	1.00	0.65	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.073	0.138	0.322	0.000	0.000	0.000	0.000	0.000
Problem 142	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	202	202	166	0	0	0	0	0	-1
normalized size	1	1.00	0.82	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.029	0.156	0.319	0.000	0.000	0.000	0.000	0.000
Problem 143	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	204	204	153	0	0	0	0	0	-1
normalized size	1	1.00	0.75	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.042	0.151	0.324	0.000	0.000	0.000	0.000	0.000

Problem 144	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	204	204	162	0	0	0	0	0	-1
normalized size	1	1.00	0.79	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.029	0.042	0.000	0.000	0.000	0.000	0.000	0.000
Problem 145	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	204	204	156	0	0	0	0	0	-1
normalized size	1	1.00	0.76	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.029	0.156	0.333	0.000	0.000	0.000	0.000	0.000
Problem 146	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	118	938	0	1943	0	0	-1
normalized size	1	1.00	1.04	8.30	0.00	17.19	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.042	16.872	0.000	3.851	0.000	0.000	0.000
Problem 147	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F(-1)	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	109	109	124	0	0	1685	0	0	-1
normalized size	1	1.00	1.14	0.00	0.00	15.46	0.00	0.00	-0.01
time (sec)	N/A	0.014	0.039	180.000	0.000	3.201	0.000	0.000	0.000
Problem 148	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	96	96	143	1033	0	285	0	0	-1
normalized size	1	1.00	1.49	10.76	0.00	2.97	0.00	0.00	-0.01
time (sec)	N/A	0.018	0.154	7.834	0.000	10.666	0.000	0.000	0.000
Problem 149	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	143	1553	0	315	0	0	-1
normalized size	1	1.00	1.51	16.35	0.00	3.32	0.00	0.00	-0.01
time (sec)	N/A	0.017	0.109	7.621	0.000	9.413	0.000	0.000	0.000

Problem 150	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	151	151	169	0	0	0	0	0	-1
normalized size	1	1.00	1.12	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.027	0.174	0.322	0.000	0.000	0.000	0.000	0.000
Problem 151	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	153	153	167	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.026	0.178	0.321	0.000	0.000	0.000	0.000	0.000
Problem 152	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	151	151	168	0	0	0	0	0	-1
normalized size	1	1.00	1.11	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.146	0.314	0.000	0.000	0.000	0.000	0.000
Problem 153	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	153	153	172	0	0	0	0	0	-1
normalized size	1	1.00	1.12	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.164	0.325	0.000	0.000	0.000	0.000	0.000
Problem 154	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	151	151	148	0	0	0	0	0	-1
normalized size	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.150	0.317	0.000	0.000	0.000	0.000	0.000
Problem 155	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	147	147	148	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.024	0.168	0.322	0.000	0.000	0.000	0.000	0.000

Problem 156	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	136	549	0	0	0	0	-1
normalized size	1	1.00	1.11	4.46	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.020	0.120	58.096	0.000	0.000	0.000	0.000	0.000
Problem 157	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	136	1061	0	0	0	0	-1
normalized size	1	1.00	1.11	8.63	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.017	0.125	59.072	0.000	0.000	0.000	0.000	0.000
Problem 158	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	119	119	136	1063	0	0	0	0	-1
normalized size	1	1.00	1.14	8.93	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.017	0.104	58.821	0.000	0.000	0.000	0.000	0.000
Problem 159	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	119	119	136	725	0	0	0	0	-1
normalized size	1	1.00	1.14	6.09	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.017	0.109	77.110	0.000	0.000	0.000	0.000	0.000
Problem 160	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	124	512	0	1395	0	0	-1
normalized size	1	1.00	1.77	7.31	0.00	19.93	0.00	0.00	-0.01
time (sec)	N/A	0.009	0.095	10.587	0.000	5.411	0.000	0.000	0.000
Problem 161	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	104	104	137	0	0	0	0	0	-1
normalized size	1	1.00	1.32	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.016	0.111	0.316	0.000	0.000	0.000	0.000	0.000

Problem 162	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	125	539	0	269	0	0	-1
normalized size	1	1.00	1.69	7.28	0.00	3.64	0.00	0.00	-0.01
time (sec)	N/A	0.011	0.046	2.611	0.000	1.724	0.000	0.000	0.000
Problem 163	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	91	57	94	70	315	0	0	-1
normalized size	1	1.15	0.72	1.19	0.89	3.99	0.00	0.00	-0.01
time (sec)	N/A	0.019	0.027	0.026	1.493	1.001	0.000	0.000	0.000
Problem 164	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	91	54	94	0	303	0	0	-1
normalized size	1	1.23	0.73	1.27	0.00	4.09	0.00	0.00	-0.01
time (sec)	N/A	0.018	0.019	0.022	0.000	0.661	0.000	0.000	0.000
Problem 165	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	91	57	94	0	314	0	0	-1
normalized size	1	1.20	0.75	1.24	0.00	4.13	0.00	0.00	-0.01
time (sec)	N/A	0.017	0.029	0.022	0.000	1.027	0.000	0.000	0.000
Problem 166	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	328	328	243	543	0	0	0	0	-1
normalized size	1	1.00	0.74	1.66	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.316	0.464	0.045	0.000	0.851	0.000	0.000	0.000
Problem 167	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	249	249	198	328	0	0	0	0	-1
normalized size	1	1.00	0.80	1.32	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.179	0.251	0.019	0.000	0.638	0.000	0.000	0.000

Problem 168	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	204	204	86	101	0	0	0	0	-1
normalized size	1	1.00	0.42	0.50	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.093	0.053	0.023	0.000	1.277	0.000	0.000	0.000
Problem 169	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	133	181	0	0	0	0	-1
normalized size	1	1.00	1.58	2.15	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.017	0.295	0.045	0.000	0.490	0.000	0.000	0.000
Problem 170	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	237	237	243	617	0	0	0	0	-1
normalized size	1	1.00	1.03	2.60	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.120	0.466	0.051	0.000	0.570	0.000	0.000	0.000
Problem 171	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	309	309	285	1411	0	0	0	0	-1
normalized size	1	1.00	0.92	4.57	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.228	0.546	0.059	0.000	0.546	0.000	0.000	0.000
Problem 172	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	410	410	302	780	0	0	0	0	-1
normalized size	1	1.00	0.74	1.90	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.439	0.599	0.036	0.000	0.677	0.000	0.000	0.000
Problem 173	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	336	336	246	545	0	0	0	0	-1
normalized size	1	1.00	0.73	1.62	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.281	0.423	0.026	0.000	0.496	0.000	0.000	0.000

Problem 174	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	273	273	199	330	0	0	0	0	-1
normalized size	1	1.00	0.73	1.21	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.165	0.280	0.027	0.000	0.517	0.000	0.000	0.000
Problem 175	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	267	267	191	332	0	0	0	0	-1
normalized size	1	1.00	0.72	1.24	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.159	0.299	0.041	0.000	0.513	0.000	0.000	0.000
Problem 176	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	229	229	232	607	0	0	0	0	-1
normalized size	1	1.00	1.01	2.65	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.136	0.485	0.046	0.000	0.527	0.000	0.000	0.000
Problem 177	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	315	315	285	1410	0	0	0	0	-1
normalized size	1	1.00	0.90	4.48	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.282	0.603	0.051	0.000	0.568	0.000	0.000	0.000
Problem 178	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	235	235	127	303	0	0	0	0	-1
normalized size	1	1.00	0.54	1.29	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.135	0.113	0.035	0.000	0.561	0.000	0.000	0.000
Problem 179	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	22	75	0	0	0	0	-1
normalized size	1	1.00	0.58	1.97	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.009	0.012	0.036	0.000	0.591	0.000	0.000	0.000

Problem 180	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	B	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	15	38	0	38	17	17	-1
normalized size	1	1.00	0.75	1.90	0.00	1.90	0.85	0.85	-0.05
time (sec)	N/A	0.004	0.002	0.003	0.000	0.586	3.433	0.573	0.000
Problem 181	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	182	182	37	37	0	0	0	0	-1
normalized size	1	1.00	0.20	0.20	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.081	0.009	0.027	0.000	0.592	0.000	0.000	0.000
Problem 182	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	60	78	0	0	0	0	-1
normalized size	1	1.00	0.66	0.86	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.038	0.033	0.000	0.550	0.000	0.000	0.000
Problem 183	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	150	150	60	53	0	0	0	0	-1
normalized size	1	1.00	0.40	0.35	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.033	0.032	0.000	0.525	0.000	0.000	0.000
Problem 184	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	23	0	0	34	0	-1
normalized size	1	1.00	1.00	1.15	0.00	0.00	1.70	0.00	-0.05
time (sec)	N/A	0.007	0.007	0.023	0.000	0.539	3.939	0.000	0.000
Problem 185	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	18	0	0	36	0	-1
normalized size	1	1.00	1.00	0.86	0.00	0.00	1.71	0.00	-0.05
time (sec)	N/A	0.008	0.005	0.036	0.000	0.466	4.212	0.000	0.000

Problem 186	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	29	0	0	34	0	-1
normalized size	1	1.00	1.00	1.45	0.00	0.00	1.70	0.00	-0.05
time (sec)	N/A	0.007	0.005	0.028	0.000	0.562	4.105	0.000	0.000
Problem 187	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	B	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	0	0	10	0	-1
normalized size	1	1.00	1.00	1.25	0.00	0.00	2.50	0.00	-0.25
time (sec)	N/A	0.005	0.003	0.020	0.000	0.549	2.393	0.000	0.000
Problem 188	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	0	0	36	0	-1
normalized size	1	1.00	1.00	0.95	0.00	0.00	1.80	0.00	-0.05
time (sec)	N/A	0.006	0.004	0.021	0.000	0.561	3.959	0.000	0.000
Problem 189	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	19	0	0	37	0	-1
normalized size	1	1.00	1.00	0.90	0.00	0.00	1.76	0.00	-0.05
time (sec)	N/A	0.006	0.005	0.021	0.000	0.601	4.166	0.000	0.000
Problem 190	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	0	0	36	0	-1
normalized size	1	1.00	1.00	0.95	0.00	0.00	1.80	0.00	-0.05
time (sec)	N/A	0.007	0.005	0.032	0.000	0.611	4.050	0.000	0.000
Problem 191	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	12	14	0	0	0	0	-1
normalized size	1	1.00	0.92	1.08	0.00	0.00	0.00	0.00	-0.08
time (sec)	N/A	0.014	0.003	0.009	0.000	0.799	0.000	0.000	0.000

Problem 192	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	27	27	0	0	0	0	-1
normalized size	1	1.00	0.87	0.87	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.020	0.006	0.018	0.000	0.658	0.000	0.000	0.000
Problem 193	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	27	31	0	0	0	0	-1
normalized size	1	1.00	0.77	0.89	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.021	0.005	0.031	0.000	0.524	0.000	0.000	0.000
Problem 194	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	27	31	0	0	0	0	-1
normalized size	1	1.00	0.77	0.89	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.020	0.005	0.022	0.000	0.606	0.000	0.000	0.000
Problem 195	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	27	30	0	0	0	0	-1
normalized size	1	1.00	0.21	0.23	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.004	0.020	0.000	0.580	0.000	0.000	0.000
Problem 196	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	136	136	27	26	0	0	0	0	-1
normalized size	1	1.00	0.20	0.19	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.044	0.004	0.019	0.000	0.568	0.000	0.000	0.000
Problem 197	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	148	148	27	20	0	0	0	0	-1
normalized size	1	1.00	0.18	0.14	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.050	0.005	0.030	0.000	0.528	0.000	0.000	0.000

Problem 198	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	35	32	0	0	0	0	-1
normalized size	1	1.00	0.88	0.80	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.015	0.025	0.015	0.000	0.764	0.000	0.000	0.000
Problem 199	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	423	423	321	852	0	0	0	0	-1
normalized size	1	1.00	0.76	2.01	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.431	1.566	0.041	0.000	0.512	0.000	0.000	0.000
Problem 200	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	344	344	260	615	0	0	0	0	-1
normalized size	1	1.00	0.76	1.79	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.283	0.495	0.031	0.000	0.648	0.000	0.000	0.000
Problem 201	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	260	260	216	399	0	0	0	0	-1
normalized size	1	1.00	0.83	1.53	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.160	0.353	0.027	0.000	0.614	0.000	0.000	0.000
Problem 202	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	194	86	158	0	0	0	0	-1
normalized size	1	1.00	0.44	0.81	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.090	0.053	0.020	0.000	0.561	0.000	0.000	0.000
Problem 203	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	86	100	0	0	0	0	-1
normalized size	1	1.00	0.99	1.15	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.019	0.057	0.030	0.000	0.541	0.000	0.000	0.000

Problem 204	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	273	273	112	248	0	0	0	0	-1
normalized size	1	1.00	0.41	0.91	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.143	0.253	0.043	0.000	0.724	0.000	0.000	0.000
Problem 205	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	255	255	261	752	0	0	0	0	-1
normalized size	1	1.00	1.02	2.95	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.146	0.613	0.056	0.000	0.570	0.000	0.000	0.000
Problem 206	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	334	334	301	1607	0	0	0	0	-1
normalized size	1	1.00	0.90	4.81	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.278	0.642	0.069	0.000	0.534	0.000	0.000	0.000
Problem 207	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	445	445	318	755	0	0	0	0	-1
normalized size	1	1.00	0.71	1.70	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.405	1.224	0.072	0.000	0.600	0.000	0.000	0.000
Problem 208	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	346	346	256	539	0	0	0	0	-1
normalized size	1	1.00	0.74	1.56	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.266	0.475	0.047	0.000	0.742	0.000	0.000	0.000
Problem 209	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	258	258	196	345	0	0	0	0	-1
normalized size	1	1.00	0.76	1.34	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.152	0.294	0.040	0.000	0.525	0.000	0.000	0.000

Problem 210	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	136	188	0	0	0	0	-1
normalized size	1	1.00	1.62	2.24	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.017	0.319	0.031	0.000	0.514	0.000	0.000	0.000
Problem 211	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	194	112	144	0	0	0	0	-1
normalized size	1	1.00	0.58	0.74	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.152	0.239	0.043	0.000	0.765	0.000	0.000	0.000
Problem 212	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	242	242	224	354	0	0	0	0	-1
normalized size	1	1.00	0.93	1.46	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.120	0.700	0.052	0.000	0.700	0.000	0.000	0.000
Problem 213	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	323	323	337	964	0	0	0	0	-1
normalized size	1	1.00	1.04	2.98	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.264	1.066	0.066	0.000	0.598	0.000	0.000	0.000
Problem 214	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	86	100	0	0	0	0	-1
normalized size	1	1.00	0.99	1.15	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.019	0.056	0.023	0.000	0.709	0.000	0.000	0.000
Problem 215	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	87	106	0	0	0	0	-1
normalized size	1	1.00	1.00	1.22	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.063	0.044	0.000	0.655	0.000	0.000	0.000

Problem 216	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	89	106	0	0	0	0	-1
normalized size	1	1.00	1.02	1.22	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.065	0.041	0.000	0.635	0.000	0.000	0.000
Problem 217	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	88	108	0	0	0	0	-1
normalized size	1	1.00	1.00	1.23	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.063	0.041	0.000	0.688	0.000	0.000	0.000
Problem 218	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	14	0	0	19	0	-1
normalized size	1	1.00	1.00	1.17	0.00	0.00	1.58	0.00	-0.08
time (sec)	N/A	0.007	0.005	0.033	0.000	0.729	3.727	0.000	0.000
Problem 219	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	58	14	0	0	0	0	-1
normalized size	1	1.00	5.80	1.40	0.00	0.00	0.00	0.00	-0.10
time (sec)	N/A	0.007	0.031	0.033	0.000	0.554	0.000	0.000	0.000
Problem 220	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	14	0	0	19	0	-1
normalized size	1	1.00	1.00	1.17	0.00	0.00	1.58	0.00	-0.08
time (sec)	N/A	0.007	0.005	0.028	0.000	0.517	4.306	0.000	0.000
Problem 221	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	B	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	10	0	0	73	0	-1
normalized size	1	1.00	1.00	1.00	0.00	0.00	7.30	0.00	-0.10
time (sec)	N/A	0.005	0.020	0.028	0.000	0.589	5.472	0.000	0.000

Problem 222	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	18	14	0	0	19	0	-1
normalized size	1	1.00	1.50	1.17	0.00	0.00	1.58	0.00	-0.08
time (sec)	N/A	0.007	0.021	0.032	0.000	0.537	2.503	0.000	0.000
Problem 223	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	0	0	0	0	-1
normalized size	1	1.00	1.00	1.08	0.00	0.00	0.00	0.00	-0.08
time (sec)	N/A	0.007	0.005	0.029	0.000	0.518	0.000	0.000	0.000
Problem 224	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	F	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	26	8	0	68	22	19	-1
normalized size	1	1.00	3.25	1.00	0.00	8.50	2.75	2.38	-0.12
time (sec)	N/A	0.002	0.005	0.317	0.000	0.531	2.321	0.575	0.000
Problem 225	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	0	0	17	0	-1
normalized size	1	1.00	1.00	1.08	0.00	0.00	1.42	0.00	-0.08
time (sec)	N/A	0.007	0.005	0.026	0.000	0.668	3.647	0.000	0.000
Problem 226	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	0	0	39	0	-1
normalized size	1	1.00	1.00	1.10	0.00	0.00	3.90	0.00	-0.10
time (sec)	N/A	0.007	0.009	0.028	0.000	0.628	5.854	0.000	0.000
Problem 227	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	0	0	34	0	-1
normalized size	1	1.00	1.00	1.08	0.00	0.00	2.83	0.00	-0.08
time (sec)	N/A	0.007	0.005	0.031	0.000	0.506	3.761	0.000	0.000

Problem 228	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	19	17	0	0	0	0	-1
normalized size	1	1.00	0.37	0.33	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.022	0.040	0.000	0.570	0.000	0.000	0.000
Problem 229	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	17	15	0	0	0	0	-1
normalized size	1	1.00	0.35	0.31	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.029	0.028	0.000	0.599	0.000	0.000	0.000
Problem 230	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	19	17	0	0	0	0	-1
normalized size	1	1.00	0.37	0.33	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.021	0.026	0.000	0.698	0.000	0.000	0.000
Problem 231	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	8	0	34	8	26	-1
normalized size	1	1.00	1.00	1.00	0.00	4.25	1.00	3.25	-0.12
time (sec)	N/A	0.002	0.003	0.307	0.000	1.038	2.494	0.572	0.000
Problem 232	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	19	15	0	0	0	0	-1
normalized size	1	1.00	0.40	0.32	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.020	0.033	0.000	0.654	0.000	0.000	0.000
Problem 233	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	19	14	0	0	0	0	-1
normalized size	1	1.00	1.90	1.40	0.00	0.00	0.00	0.00	-0.10
time (sec)	N/A	0.006	0.019	0.033	0.000	0.729	0.000	0.000	0.000

Problem 234	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	B	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	10	0	0	76	0	-1
normalized size	1	1.00	1.00	1.00	0.00	0.00	7.60	0.00	-0.10
time (sec)	N/A	0.005	0.014	0.020	0.000	0.712	5.264	0.000	0.000
Problem 235	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	0	0	36	0	-1
normalized size	1	1.00	1.00	0.95	0.00	0.00	1.80	0.00	-0.05
time (sec)	N/A	0.007	0.004	0.028	0.000	0.641	4.087	0.000	0.000
Problem 236	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	0	0	41	0	-1
normalized size	1	1.00	1.00	0.94	0.00	0.00	2.56	0.00	-0.06
time (sec)	N/A	0.007	0.007	0.028	0.000	0.677	6.020	0.000	0.000
Problem 237	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	0	0	36	0	-1
normalized size	1	1.00	1.00	0.95	0.00	0.00	1.80	0.00	-0.05
time (sec)	N/A	0.009	0.007	0.048	0.000	0.528	4.091	0.000	0.000
Problem 238	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	37	0	0	0	0	-1
normalized size	1	1.00	1.00	1.16	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.014	0.023	0.043	0.000	0.599	0.000	0.000	0.000
Problem 239	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	34	0	0	0	0	-1
normalized size	1	1.00	1.00	1.13	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.015	0.028	0.040	0.000	0.562	0.000	0.000	0.000

Problem 240	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	37	0	0	0	0	-1
normalized size	1	1.00	1.00	1.16	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.013	0.022	0.036	0.000	0.525	0.000	0.000	0.000
Problem 241	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	46	30	0	0	75	0	-1
normalized size	1	1.00	1.84	1.20	0.00	0.00	3.00	0.00	-0.04
time (sec)	N/A	0.011	0.011	0.029	0.000	0.554	5.550	0.000	0.000
Problem 242	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	34	0	0	0	0	-1
normalized size	1	1.00	1.00	1.06	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.013	0.021	0.033	0.000	0.690	0.000	0.000	0.000
Problem 243	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	47	28	0	0	0	0	-1
normalized size	1	1.00	3.92	2.33	0.00	0.00	0.00	0.00	-0.08
time (sec)	N/A	0.007	0.024	0.029	0.000	0.618	0.000	0.000	0.000
Problem 244	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	40	24	0	34	0	0	-1
normalized size	1	1.00	1.38	0.83	0.00	1.17	0.00	0.00	-0.03
time (sec)	N/A	0.003	0.011	0.010	0.000	0.645	0.000	0.000	0.000
Problem 245	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	40	29	0	0	37	0	-1
normalized size	1	1.00	1.25	0.91	0.00	0.00	1.16	0.00	-0.03
time (sec)	N/A	0.015	0.025	0.031	0.000	0.696	4.096	0.000	0.000

Problem 246	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	36	27	0	0	42	0	-1
normalized size	1	1.00	1.20	0.90	0.00	0.00	1.40	0.00	-0.03
time (sec)	N/A	0.015	0.029	0.030	0.000	0.526	6.073	0.000	0.000
Problem 247	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	40	29	0	0	37	0	-1
normalized size	1	1.00	1.25	0.91	0.00	0.00	1.16	0.00	-0.03
time (sec)	N/A	0.014	0.025	0.034	0.000	0.500	4.147	0.000	0.000
Problem 248	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	39	36	0	0	0	0	-1
normalized size	1	1.00	0.74	0.68	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.010	0.027	0.033	0.000	0.678	0.000	0.000	0.000
Problem 249	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	37	33	0	0	0	0	-1
normalized size	1	1.00	0.73	0.65	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.010	0.030	0.031	0.000	0.614	0.000	0.000	0.000
Problem 250	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	39	36	0	0	0	0	-1
normalized size	1	1.00	0.74	0.68	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.024	0.034	0.000	0.583	0.000	0.000	0.000
Problem 251	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	26	24	0	104	0	0	-1
normalized size	1	1.00	0.93	0.86	0.00	3.71	0.00	0.00	-0.04
time (sec)	N/A	0.004	0.011	0.010	0.000	0.647	0.000	0.000	0.000

Problem 252	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	53	33	0	0	0	0	-1
normalized size	1	1.00	1.08	0.67	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.038	0.027	0.000	0.509	0.000	0.000	0.000
Problem 253	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	39	34	0	0	0	0	-1
normalized size	1	1.00	1.26	1.10	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.016	0.025	0.030	0.000	0.697	0.000	0.000	0.000
Problem 254	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	65	48	30	0	0	73	0	-1
normalized size	1	1.55	1.14	0.71	0.00	0.00	1.74	0.00	-0.02
time (sec)	N/A	0.012	0.015	0.023	0.000	0.618	5.455	0.000	0.000
Problem 255	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	40	34	0	0	0	0	-1
normalized size	1	1.00	1.00	0.85	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.016	0.025	0.033	0.000	0.857	0.000	0.000	0.000
Problem 256	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	36	34	0	0	44	0	-1
normalized size	1	1.00	1.00	0.94	0.00	0.00	1.22	0.00	-0.03
time (sec)	N/A	0.015	0.031	0.027	0.000	0.851	5.780	0.000	0.000
Problem 257	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	40	34	0	0	0	0	-1
normalized size	1	1.00	1.00	0.85	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.015	0.028	0.033	0.000	0.908	0.000	0.000	0.000

Problem 258	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	87	106	0	0	0	0	-1
normalized size	1	1.00	1.00	1.22	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.053	0.059	0.018	0.000	0.838	0.000	0.000	0.000
Problem 259	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	90	171	0	0	0	0	-1
normalized size	1	1.00	1.00	1.90	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.051	0.026	0.000	0.632	0.000	0.000	0.000
Problem 260	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	88	168	0	0	0	0	-1
normalized size	1	1.00	1.00	1.91	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.049	0.049	0.017	0.000	0.664	0.000	0.000	0.000
Problem 261	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	91	110	0	0	0	0	-1
normalized size	1	1.00	1.00	1.21	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.049	0.017	0.000	0.683	0.000	0.000	0.000
Problem 262	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	88	109	0	0	0	0	-1
normalized size	1	1.00	1.00	1.24	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.058	0.021	0.000	0.700	0.000	0.000	0.000
Problem 263	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	89	165	0	0	0	0	-1
normalized size	1	1.00	1.00	1.85	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.053	0.054	0.028	0.000	0.551	0.000	0.000	0.000

Problem 264	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	89	166	0	0	0	0	-1
normalized size	1	1.00	1.00	1.87	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.055	0.048	0.020	0.000	0.581	0.000	0.000	0.000
Problem 265	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	90	111	0	0	0	0	-1
normalized size	1	1.00	1.00	1.23	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.048	0.019	0.000	0.504	0.000	0.000	0.000
Problem 266	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	194	86	158	0	0	0	0	-1
normalized size	1	1.00	0.44	0.81	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.084	0.048	0.013	0.000	0.722	0.000	0.000	0.000
Problem 267	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	203	203	89	104	0	0	0	0	-1
normalized size	1	1.00	0.44	0.51	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.089	0.047	0.026	0.000	0.578	0.000	0.000	0.000
Problem 268	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	203	203	89	108	0	0	0	0	-1
normalized size	1	1.00	0.44	0.53	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.092	0.050	0.020	0.000	0.640	0.000	0.000	0.000
Problem 269	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	212	212	92	165	0	0	0	0	-1
normalized size	1	1.00	0.43	0.78	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.103	0.046	0.020	0.000	0.557	0.000	0.000	0.000

Problem 270	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	189	189	89	164	0	0	0	0	-1
normalized size	1	1.00	0.47	0.87	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.125	0.052	0.019	0.000	0.523	0.000	0.000	0.000
Problem 271	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	191	191	90	109	0	0	0	0	-1
normalized size	1	1.00	0.47	0.57	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.123	0.054	0.023	0.000	0.532	0.000	0.000	0.000
Problem 272	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	194	92	111	0	0	0	0	-1
normalized size	1	1.00	0.47	0.57	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.128	0.052	0.019	0.000	0.921	0.000	0.000	0.000
Problem 273	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	198	198	93	167	0	0	0	0	-1
normalized size	1	1.00	0.47	0.84	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.131	0.049	0.019	0.000	0.672	0.000	0.000	0.000
Problem 274	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	87	106	0	0	0	0	-1
normalized size	1	1.00	1.00	1.22	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.048	0.056	0.017	0.000	0.750	0.000	0.000	0.000
Problem 275	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	90	171	0	0	0	0	-1
normalized size	1	1.00	1.00	1.90	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.049	0.024	0.000	0.602	0.000	0.000	0.000

Problem 276	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	88	168	0	0	0	0	-1
normalized size	1	1.00	1.00	1.91	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.047	0.047	0.019	0.000	0.625	0.000	0.000	0.000
Problem 277	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	91	110	0	0	0	0	-1
normalized size	1	1.00	1.00	1.21	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.050	0.043	0.016	0.000	0.470	0.000	0.000	0.000
Problem 278	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	88	164	0	0	0	0	-1
normalized size	1	1.00	1.00	1.86	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.056	0.022	0.000	0.809	0.000	0.000	0.000
Problem 279	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	89	110	0	0	0	0	-1
normalized size	1	1.00	1.00	1.24	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.049	0.027	0.000	0.625	0.000	0.000	0.000
Problem 280	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	89	110	0	0	0	0	-1
normalized size	1	1.00	1.00	1.24	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.049	0.019	0.000	0.563	0.000	0.000	0.000
Problem 281	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	90	167	0	0	0	0	-1
normalized size	1	1.00	1.00	1.86	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.043	0.019	0.000	0.570	0.000	0.000	0.000

Problem 282	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	204	204	86	101	0	0	0	0	-1
normalized size	1	1.00	0.42	0.50	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.088	0.045	0.017	0.000	0.570	0.000	0.000	0.000
Problem 283	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	214	214	89	161	0	0	0	0	-1
normalized size	1	1.00	0.42	0.75	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.096	0.046	0.025	0.000	0.533	0.000	0.000	0.000
Problem 284	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	214	214	89	162	0	0	0	0	-1
normalized size	1	1.00	0.42	0.76	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.097	0.051	0.020	0.000	0.598	0.000	0.000	0.000
Problem 285	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	222	222	92	111	0	0	0	0	-1
normalized size	1	1.00	0.41	0.50	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.103	0.052	0.021	0.000	0.588	0.000	0.000	0.000
Problem 286	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	189	189	89	164	0	0	0	0	-1
normalized size	1	1.00	0.47	0.87	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.121	0.053	0.019	0.000	0.733	0.000	0.000	0.000
Problem 287	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	191	191	90	109	0	0	0	0	-1
normalized size	1	1.00	0.47	0.57	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.122	0.047	0.023	0.000	0.590	0.000	0.000	0.000

Problem 288	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	194	92	111	0	0	0	0	-1
normalized size	1	1.00	0.47	0.57	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.125	0.051	0.019	0.000	0.562	0.000	0.000	0.000
Problem 289	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	198	198	93	167	0	0	0	0	-1
normalized size	1	1.00	0.47	0.84	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.129	0.051	0.019	0.000	0.705	0.000	0.000	0.000
Problem 290	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	37	38	0	0	0	0	-1
normalized size	1	1.00	0.47	0.49	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.014	0.013	0.036	0.000	0.724	0.000	0.000	0.000
Problem 291	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	40	38	0	0	20	0	-1
normalized size	1	1.00	1.03	0.97	0.00	0.00	0.51	0.00	-0.03
time (sec)	N/A	0.020	0.040	0.033	0.000	0.543	2.379	0.000	0.000
Problem 292	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	47	53	0	0	0	0	-1
normalized size	1	1.00	0.77	0.87	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.012	0.037	0.031	0.000	0.606	0.000	0.000	0.000
Problem 293	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	6	6	27	25	0	0	0	0	-1
normalized size	1	1.00	4.50	4.17	0.00	0.00	0.00	0.00	-0.17
time (sec)	N/A	0.006	0.025	0.026	0.000	0.585	0.000	0.000	0.000

Problem 294	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	24	28	0	0	0	0	-1
normalized size	1	1.00	1.04	1.22	0.00	0.00	0.00	0.00	-0.04
time (sec)	N/A	0.025	0.009	0.023	0.000	0.503	0.000	0.000	0.000
Problem 295	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	182	182	37	37	0	0	0	0	-1
normalized size	1	1.00	0.20	0.20	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.075	0.009	0.019	0.000	0.527	0.000	0.000	0.000
Problem 296	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	35	37	0	0	0	0	-1
normalized size	1	1.00	1.84	1.95	0.00	0.00	0.00	0.00	-0.05
time (sec)	N/A	0.007	0.025	0.025	0.000	0.452	0.000	0.000	0.000
Problem 297	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	95	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.132	0.112	0.374	0.000	0.909	0.000	0.000	0.000
Problem 298	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	95	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.102	0.095	0.336	0.000	0.791	0.000	0.000	0.000
Problem 299	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	478	478	102	0	0	0	0	0	-1
normalized size	1	1.00	0.21	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.367	0.121	0.333	0.000	0.539	0.000	0.000	0.000

Problem 300	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	215	215	102	0	0	0	0	0	-1
normalized size	1	1.00	0.47	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.208	0.093	0.325	0.000	0.747	0.000	0.000	0.000
Problem 301	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	C	C	F	F	F	F	F	F
verified	N/A	NO	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	62	23	122	0	0	0	0	0	-1
normalized size	1	0.37	1.97	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.011	0.129	0.314	0.000	0.665	0.000	0.000	0.000
Problem 302	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	48	117	0	0	0	0	-1
normalized size	1	1.00	1.04	2.54	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.041	0.085	0.145	0.000	0.619	0.000	0.000	0.000
Problem 303	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	81	207	0	0	0	0	-1
normalized size	1	1.00	1.72	4.40	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.062	0.154	0.225	0.000	0.592	0.000	0.000	0.000
Problem 304	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	129	129	135	186	0	553	0	0	-1
normalized size	1	1.00	1.05	1.44	0.00	4.29	0.00	0.00	-0.01
time (sec)	N/A	0.019	0.113	1.667	0.000	3.997	0.000	0.000	0.000
Problem 305	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	135	187	0	553	0	0	-1
normalized size	1	1.00	1.12	1.56	0.00	4.61	0.00	0.00	-0.01
time (sec)	N/A	0.016	0.128	1.596	0.000	4.337	0.000	0.000	0.000

Problem 306	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	129	129	144	0	0	755	0	0	-1
normalized size	1	1.00	1.12	0.00	0.00	5.85	0.00	0.00	-0.01
time (sec)	N/A	0.023	0.138	0.331	0.000	12.454	0.000	0.000	0.000
Problem 307	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	145	0	0	776	0	0	-1
normalized size	1	1.00	1.17	0.00	0.00	6.26	0.00	0.00	-0.01
time (sec)	N/A	0.020	0.133	0.335	0.000	13.087	0.000	0.000	0.000
Problem 308	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	155	0	0	286	0	0	-1
normalized size	1	1.00	1.29	0.00	0.00	2.38	0.00	0.00	-0.01
time (sec)	N/A	0.020	0.145	0.344	0.000	9.570	0.000	0.000	0.000
Problem 309	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	155	0	0	286	0	0	-1
normalized size	1	1.00	1.29	0.00	0.00	2.38	0.00	0.00	-0.01
time (sec)	N/A	0.017	0.162	0.340	0.000	9.736	0.000	0.000	0.000
Problem 310	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	165	0	0	337	0	0	-1
normalized size	1	1.00	1.38	0.00	0.00	2.81	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.155	0.334	0.000	58.437	0.000	0.000	0.000
Problem 311	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	162	0	0	343	0	0	-1
normalized size	1	1.00	1.31	0.00	0.00	2.77	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.164	0.336	0.000	60.662	0.000	0.000	0.000

Problem 312	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	127	138	0	104	0	0	-1
normalized size	1	1.00	2.08	2.26	0.00	1.70	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.148	1.139	0.000	4.436	0.000	0.000	0.000
Problem 313	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	C	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	127	138	0	243	0	0	-1
normalized size	1	1.00	2.08	2.26	0.00	3.98	0.00	0.00	-0.02
time (sec)	N/A	0.010	0.129	1.112	0.000	4.433	0.000	0.000	0.000
Problem 314	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	132	0	0	274	0	0	-1
normalized size	1	1.00	1.71	0.00	0.00	3.56	0.00	0.00	-0.01
time (sec)	N/A	0.013	0.166	0.335	0.000	13.267	0.000	0.000	0.000
Problem 315	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	137	0	0	273	0	0	-1
normalized size	1	1.00	1.73	0.00	0.00	3.46	0.00	0.00	-0.01
time (sec)	N/A	0.014	0.148	0.334	0.000	13.266	0.000	0.000	0.000
Problem 316	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	157	0	0	276	0	0	-1
normalized size	1	1.00	1.85	0.00	0.00	3.25	0.00	0.00	-0.01
time (sec)	N/A	0.018	0.171	0.351	0.000	9.493	0.000	0.000	0.000
Problem 317	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	157	0	0	278	0	0	-1
normalized size	1	1.00	1.85	0.00	0.00	3.27	0.00	0.00	-0.01
time (sec)	N/A	0.014	0.148	0.363	0.000	9.425	0.000	0.000	0.000

Problem 318	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	163	0	0	338	0	0	-1
normalized size	1	1.00	1.61	0.00	0.00	3.35	0.00	0.00	-0.01
time (sec)	N/A	0.019	0.164	0.342	0.000	50.674	0.000	0.000	0.000
Problem 319	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	168	0	0	350	0	0	-1
normalized size	1	1.00	1.63	0.00	0.00	3.40	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.163	0.337	0.000	56.410	0.000	0.000	0.000
Problem 320	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	115	121	0	91	0	0	-1
normalized size	1	1.00	2.17	2.28	0.00	1.72	0.00	0.00	-0.02
time (sec)	N/A	0.008	0.141	1.220	0.000	3.964	0.000	0.000	0.000
Problem 321	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	362	362	346	0	0	0	0	0	-1
normalized size	1	1.00	0.96	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.265	0.461	0.344	0.000	0.000	0.000	0.000	0.000
Problem 322	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	302	302	348	0	0	0	0	0	-1
normalized size	1	1.00	1.15	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.207	0.471	0.344	0.000	0.000	0.000	0.000	0.000
Problem 323	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	244	244	161	0	0	0	0	0	-1
normalized size	1	1.00	0.66	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.150	0.163	0.336	0.000	0.000	0.000	0.000	0.000

Problem 324	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	199	199	160	0	0	0	0	0	-1
normalized size	1	1.00	0.80	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.147	0.158	0.329	0.000	0.000	0.000	0.000	0.000
Problem 325	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	167	167	160	0	0	0	0	0	-1
normalized size	1	1.00	0.96	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.108	0.056	0.333	0.000	0.000	0.000	0.000	0.000
Problem 326	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	152	152	120	0	0	0	0	0	-1
normalized size	1	1.00	0.79	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.118	0.032	0.335	0.000	0.000	0.000	0.000	0.000
Problem 327	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	233	233	327	0	0	0	0	0	-1
normalized size	1	1.00	1.40	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.160	0.233	0.339	0.000	0.000	0.000	0.000	0.000
Problem 328	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	254	254	331	0	0	0	0	0	-1
normalized size	1	1.00	1.30	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.157	0.262	0.337	0.000	0.000	0.000	0.000	0.000
Problem 329	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	274	274	419	0	0	0	0	0	-1
normalized size	1	1.00	1.53	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.382	0.660	0.349	0.000	0.000	0.000	0.000	0.000

Problem 330	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	304	304	431	0	0	0	0	0	-1
normalized size	1	1.00	1.42	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.343	0.776	0.351	0.000	0.000	0.000	0.000	0.000
Problem 331	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	340	340	340	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.294	0.338	0.351	0.000	0.000	0.000	0.000	0.000
Problem 332	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	279	279	341	0	0	0	0	0	-1
normalized size	1	1.00	1.22	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.271	0.332	0.344	0.000	0.000	0.000	0.000	0.000
Problem 333	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	309	309	232	0	0	0	0	0	-1
normalized size	1	1.00	0.75	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.222	0.203	0.347	0.000	0.000	0.000	0.000	0.000
Problem 334	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	278	278	232	0	0	0	0	0	-1
normalized size	1	1.00	0.83	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.206	0.189	0.339	0.000	0.000	0.000	0.000	0.000
Problem 335	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	336	336	392	0	0	0	0	0	-1
normalized size	1	1.00	1.17	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.255	0.253	0.344	0.000	0.000	0.000	0.000	0.000

Problem 336	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	292	292	336	0	0	0	0	0	-1
normalized size	1	1.00	1.15	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.231	0.320	0.344	0.000	0.000	0.000	0.000	0.000
Problem 337	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	314	314	380	0	0	0	0	0	-1
normalized size	1	1.00	1.21	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.395	0.491	0.356	0.000	0.000	0.000	0.000	0.000
Problem 338	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	345	345	387	0	0	0	0	0	-1
normalized size	1	1.00	1.12	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.382	0.512	0.357	0.000	0.000	0.000	0.000	0.000
Problem 339	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	371	371	536	0	0	0	0	0	-1
normalized size	1	1.00	1.44	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.552	1.048	0.370	0.000	0.000	0.000	0.000	0.000
Problem 340	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	419	419	550	0	0	0	0	0	-1
normalized size	1	1.00	1.31	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.484	0.920	0.370	0.000	0.000	0.000	0.000	0.000
Problem 341	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F(-1)	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	172	0	0	0	0	0	-1
normalized size	1	1.00	2.18	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.225	0.507	0.000	1.562	0.000	0.000	0.000

Problem 342	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	296	296	136	0	0	0	121	0	-1
normalized size	1	1.00	0.46	0.00	0.00	0.00	0.41	0.00	-0.00
time (sec)	N/A	0.276	5.067	0.361	0.000	1.519	39.511	0.000	0.000
Problem 343	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	176	176	106	0	0	0	88	0	-1
normalized size	1	1.00	0.60	0.00	0.00	0.00	0.50	0.00	-0.01
time (sec)	N/A	0.121	5.048	0.308	0.000	1.524	20.397	0.000	0.000
Problem 344	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	85	90	0	0	0	53	0	-1
normalized size	1	0.91	0.97	0.00	0.00	0.00	0.57	0.00	-0.01
time (sec)	N/A	0.039	0.032	0.304	0.000	1.333	9.832	0.000	0.000
Problem 345	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	C	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	44	0	0	0	22	0	41
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.50	0.00	0.93
time (sec)	N/A	0.009	0.004	0.000	0.000	0.875	2.326	0.000	5.567
Problem 346	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F(-1)	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	162	0	0	0	0	0	-1
normalized size	1	1.00	2.84	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.182	0.320	0.000	0.777	0.000	0.000	0.000
Problem 347	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F(-1)	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	162	0	0	0	0	0	-1
normalized size	1	1.00	2.84	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.193	0.311	0.000	1.004	0.000	0.000	0.000

Problem 348	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F(-1)	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	162	0	0	0	0	0	-1
normalized size	1	1.00	2.84	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.245	0.313	0.000	1.001	0.000	0.000	0.000

Problem 349	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	52	71	0	91	0	0	131
normalized size	1	1.00	0.98	1.34	0.00	1.72	0.00	0.00	2.47
time (sec)	N/A	0.020	0.032	0.005	0.000	1.016	0.000	0.000	5.736

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [329] had the largest ratio of [.4286]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	2	1	1.00	17	0.059
2	A	2	1	1.00	17	0.059
3	A	2	1	1.00	17	0.059
4	A	2	1	1.00	15	0.067
5	A	2	2	1.00	17	0.118
6	A	2	2	1.00	17	0.118
7	A	3	3	1.00	17	0.176
8	A	2	1	1.00	19	0.053
9	A	2	1	1.00	19	0.053
10	A	2	1	1.00	17	0.059
11	A	3	2	1.00	19	0.105
12	A	4	3	1.00	19	0.158
13	A	3	3	1.00	19	0.158

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
14	A	2	1	1.00	19	0.053
15	A	2	1	1.00	19	0.053
16	A	2	1	1.00	17	0.059
17	A	3	2	1.00	19	0.105
18	A	4	3	1.00	19	0.158
19	A	5	4	1.00	19	0.210
20	A	3	2	1.00	19	0.105
21	A	3	2	1.00	19	0.105
22	A	3	2	1.00	19	0.105
23	A	2	2	1.00	17	0.118
24	A	3	2	1.00	19	0.105
25	A	4	3	1.00	19	0.158
26	A	5	4	1.00	19	0.210
27	A	4	3	1.00	19	0.158
28	A	4	3	1.00	19	0.158
29	A	4	3	1.00	19	0.158
30	A	4	3	1.00	19	0.158
31	A	2	2	1.00	17	0.118
32	A	4	3	1.00	19	0.158
33	A	5	4	1.00	19	0.210
34	A	6	4	1.00	19	0.210
35	A	5	4	1.00	19	0.210
36	A	5	4	1.00	19	0.210
37	A	5	4	1.00	19	0.210
38	A	3	3	1.00	19	0.158
39	A	3	3	1.00	17	0.176
40	A	5	4	1.00	19	0.210
41	A	6	4	1.00	19	0.210
42	A	7	4	1.00	19	0.210
43	A	3	3	1.00	15	0.200
44	A	5	3	1.00	15	0.200
45	A	6	6	1.00	21	0.286
46	A	5	5	1.00	21	0.238
47	A	4	4	1.00	19	0.210
48	A	3	3	1.00	11	0.273
49	A	5	5	1.00	21	0.238

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
50	A	3	3	1.00	21	0.143
51	A	4	4	1.00	21	0.190
52	A	6	5	1.00	21	0.238
53	A	7	6	1.00	21	0.286
54	A	6	5	1.00	21	0.238
55	A	5	4	1.00	19	0.210
56	A	4	3	1.00	11	0.273
57	A	6	6	1.00	21	0.286
58	A	6	6	1.00	21	0.286
59	A	4	3	1.00	21	0.143
60	A	5	4	1.00	21	0.190
61	A	7	5	1.00	21	0.238
62	A	8	6	1.00	21	0.286
63	A	7	5	1.00	21	0.238
64	A	6	4	1.00	19	0.210
65	A	5	3	1.00	11	0.273
66	A	7	7	1.00	21	0.333
67	A	7	7	1.00	21	0.333
68	A	7	7	1.00	21	0.333
69	A	5	3	1.00	21	0.143
70	A	6	4	1.00	21	0.190
71	A	4	4	1.00	19	0.210
72	A	4	4	1.00	17	0.235
73	A	4	4	1.00	21	0.190
74	A	5	5	1.00	21	0.238
75	A	4	4	1.00	21	0.190
76	A	3	3	1.00	19	0.158
77	A	2	2	1.00	11	0.182
78	A	2	2	1.00	21	0.095
79	A	3	3	1.00	21	0.143
80	A	5	5	1.00	21	0.238
81	A	6	5	1.00	21	0.238
82	A	5	5	1.00	21	0.238
83	A	4	4	1.17	21	0.190
84	A	3	3	1.00	19	0.158
85	A	1	1	1.00	11	0.091

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	3	3	1.00	21	0.143
87	A	5	5	1.00	21	0.238
88	A	6	5	1.00	21	0.238
89	A	6	6	1.00	21	0.286
90	A	5	5	1.00	21	0.238
91	A	4	4	1.00	21	0.190
92	A	2	2	1.00	19	0.105
93	A	2	2	1.00	11	0.182
94	A	5	5	1.00	21	0.238
95	A	6	5	1.00	21	0.238
96	A	7	5	1.00	21	0.238
97	A	5	3	1.00	21	0.143
98	A	4	3	1.00	21	0.143
99	A	3	3	1.00	19	0.158
100	A	2	2	1.00	11	0.182
101	A	3	3	1.00	21	0.143
102	A	3	3	1.00	21	0.143
103	A	4	4	1.00	21	0.190
104	A	5	4	1.00	21	0.190
105	A	2	2	1.00	26	0.077
106	A	2	2	1.00	19	0.105
107	A	2	2	1.00	21	0.095
108	A	2	2	1.00	15	0.133
109	A	8	8	1.00	24	0.333
110	A	7	7	1.00	24	0.292
111	A	6	6	1.00	22	0.273
112	A	6	6	1.00	24	0.250
113	A	6	6	1.00	24	0.250
114	A	8	8	1.00	24	0.333
115	A	9	8	1.00	24	0.333
116	A	9	8	1.00	24	0.333
117	A	8	7	1.00	24	0.292
118	A	7	6	1.00	22	0.273
119	A	7	7	1.00	24	0.292
120	A	7	7	1.00	24	0.292
121	A	9	9	1.00	24	0.375

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
122	A	8	7	1.00	24	0.292
123	A	7	7	1.00	24	0.292
124	A	6	6	1.00	24	0.250
125	A	5	5	1.00	22	0.227
126	A	1	1	1.00	24	0.042
127	A	7	7	1.00	24	0.292
128	A	8	8	1.00	24	0.333
129	A	7	7	1.00	24	0.292
130	A	6	6	1.00	24	0.250
131	A	5	5	1.00	22	0.227
132	A	7	7	1.00	24	0.292
133	A	8	8	1.00	24	0.333
134	A	9	8	1.00	24	0.333
135	A	8	8	1.00	24	0.333
136	A	7	7	1.00	24	0.292
137	A	2	2	1.00	24	0.083
138	A	6	6	1.00	22	0.273
139	A	8	8	1.00	24	0.333
140	A	9	8	1.00	24	0.333
141	A	1	1	1.00	26	0.038
142	A	1	1	1.00	24	0.042
143	A	1	1	1.00	23	0.043
144	A	1	1	1.00	24	0.042
145	A	1	1	1.00	22	0.045
146	A	1	1	1.00	19	0.053
147	A	1	1	1.00	19	0.053
148	A	1	1	1.00	24	0.042
149	A	1	1	1.00	22	0.045
150	A	1	1	1.00	27	0.037
151	A	1	1	1.00	28	0.036
152	A	1	1	1.00	29	0.034
153	A	1	1	1.00	30	0.033
154	A	1	1	1.00	26	0.038
155	A	1	1	1.00	26	0.038
156	A	1	1	1.00	23	0.043
157	A	1	1	1.00	23	0.043

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
158	A	1	1	1.00	23	0.043
159	A	1	1	1.00	23	0.043
160	A	1	1	1.00	17	0.059
161	A	1	1	1.00	21	0.048
162	A	1	1	1.00	21	0.048
163	A	3	3	1.15	29	0.103
164	A	3	3	1.23	29	0.103
165	A	3	3	1.20	29	0.103
166	A	6	6	1.00	23	0.261
167	A	5	5	1.00	23	0.217
168	A	4	4	1.00	23	0.174
169	A	1	1	1.00	23	0.043
170	A	4	4	1.00	23	0.174
171	A	5	5	1.00	23	0.217
172	A	7	6	1.00	23	0.261
173	A	6	6	1.00	23	0.261
174	A	5	5	1.00	23	0.217
175	A	5	5	1.00	23	0.217
176	A	4	4	1.00	23	0.174
177	A	5	5	1.00	23	0.217
178	A	5	5	1.00	23	0.217
179	A	3	3	1.00	23	0.130
180	A	2	1	1.00	23	0.043
181	A	4	4	1.00	23	0.174
182	A	5	5	1.00	23	0.217
183	A	4	4	1.00	21	0.190
184	A	1	1	1.00	23	0.043
185	A	1	1	1.00	23	0.043
186	A	1	1	1.00	23	0.043
187	A	1	1	1.00	21	0.048
188	A	1	1	1.00	21	0.048
189	A	1	1	1.00	21	0.048
190	A	1	1	1.00	23	0.043
191	A	4	4	1.00	21	0.190
192	A	3	3	1.00	23	0.130
193	A	3	3	1.00	23	0.130

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
194	A	3	3	1.00	23	0.130
195	A	4	4	1.00	21	0.190
196	A	4	4	1.00	21	0.190
197	A	4	4	1.00	23	0.174
198	A	2	2	1.00	23	0.087
199	A	7	6	1.00	23	0.261
200	A	6	6	1.00	23	0.261
201	A	5	5	1.00	23	0.217
202	A	4	4	1.00	23	0.174
203	A	1	1	1.00	23	0.043
204	A	6	6	1.00	23	0.261
205	A	4	4	1.00	23	0.174
206	A	5	5	1.00	23	0.217
207	A	7	6	1.00	23	0.261
208	A	6	6	1.00	23	0.261
209	A	5	5	1.00	23	0.217
210	A	1	1	1.00	23	0.043
211	A	6	6	1.00	23	0.261
212	A	4	4	1.00	23	0.174
213	A	5	5	1.00	23	0.217
214	A	1	1	1.00	23	0.043
215	A	3	2	1.00	24	0.083
216	A	3	2	1.00	24	0.083
217	A	3	2	1.00	25	0.080
218	A	1	1	1.00	23	0.043
219	A	1	1	1.00	23	0.043
220	A	1	1	1.00	23	0.043
221	A	2	2	1.00	23	0.087
222	A	1	1	1.00	21	0.048
223	A	1	1	1.00	23	0.043
224	A	2	2	1.00	23	0.087
225	A	1	1	1.00	23	0.043
226	A	1	1	1.00	23	0.043
227	A	1	1	1.00	23	0.043
228	A	1	1	1.00	21	0.048
229	A	1	1	1.00	21	0.048

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
230	A	1	1	1.00	21	0.048
231	A	2	2	1.00	21	0.095
232	A	1	1	1.00	19	0.053
233	A	1	1	1.00	21	0.048
234	A	2	2	1.00	21	0.095
235	A	1	1	1.00	21	0.048
236	A	1	1	1.00	21	0.048
237	A	1	1	1.00	21	0.048
238	A	2	2	1.00	21	0.095
239	A	2	2	1.00	21	0.095
240	A	2	2	1.00	21	0.095
241	A	2	2	1.00	21	0.095
242	A	2	2	1.00	19	0.105
243	A	1	1	1.00	21	0.048
244	A	2	2	1.00	21	0.095
245	A	2	2	1.00	21	0.095
246	A	2	2	1.00	21	0.095
247	A	2	2	1.00	21	0.095
248	A	1	1	1.00	23	0.043
249	A	1	1	1.00	23	0.043
250	A	1	1	1.00	23	0.043
251	A	2	2	1.00	23	0.087
252	A	1	1	1.00	21	0.048
253	A	2	2	1.00	23	0.087
254	A	2	2	1.55	23	0.087
255	A	2	2	1.00	23	0.087
256	A	2	2	1.00	23	0.087
257	A	2	2	1.00	23	0.087
258	A	3	3	1.00	24	0.125
259	A	3	3	1.00	27	0.111
260	A	3	3	1.00	25	0.120
261	A	3	3	1.00	28	0.107
262	A	3	3	1.00	25	0.120
263	A	3	3	1.00	26	0.115
264	A	3	3	1.00	26	0.115
265	A	3	3	1.00	27	0.111

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
266	A	4	4	1.00	23	0.174
267	A	4	4	1.00	26	0.154
268	A	4	4	1.00	26	0.154
269	A	4	4	1.00	29	0.138
270	A	7	6	1.00	24	0.250
271	A	7	6	1.00	25	0.240
272	A	7	6	1.00	27	0.222
273	A	7	6	1.00	28	0.214
274	A	3	3	1.00	24	0.125
275	A	3	3	1.00	27	0.111
276	A	3	3	1.00	25	0.120
277	A	3	3	1.00	28	0.107
278	A	3	3	1.00	25	0.120
279	A	3	3	1.00	26	0.115
280	A	3	3	1.00	26	0.115
281	A	3	3	1.00	27	0.111
282	A	4	4	1.00	23	0.174
283	A	4	4	1.00	26	0.154
284	A	4	4	1.00	26	0.154
285	A	4	4	1.00	29	0.138
286	A	7	6	1.00	24	0.250
287	A	7	6	1.00	25	0.240
288	A	7	6	1.00	27	0.222
289	A	7	6	1.00	28	0.214
290	A	1	1	1.00	23	0.043
291	A	2	2	1.00	23	0.087
292	A	1	1	1.00	21	0.048
293	A	1	1	1.00	23	0.043
294	A	4	4	1.00	28	0.143
295	A	4	4	1.00	23	0.174
296	A	1	1	1.00	23	0.043
297	A	1	1	1.00	59	0.017
298	A	1	1	1.00	59	0.017
299	A	4	4	1.00	59	0.068
300	A	3	3	1.00	59	0.051
301	C	1	1	0.37	21	0.048

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
302	A	2	2	1.00	26	0.077
303	A	1	1	1.00	41	0.024
304	A	1	1	1.00	21	0.048
305	A	1	1	1.00	21	0.048
306	A	1	1	1.00	21	0.048
307	A	1	1	1.00	23	0.043
308	A	1	1	1.00	23	0.043
309	A	1	1	1.00	23	0.043
310	A	1	1	1.00	23	0.043
311	A	1	1	1.00	25	0.040
312	A	1	1	1.00	21	0.048
313	A	1	1	1.00	21	0.048
314	A	1	1	1.00	21	0.048
315	A	1	1	1.00	23	0.043
316	A	1	1	1.00	25	0.040
317	A	1	1	1.00	25	0.040
318	A	1	1	1.00	25	0.040
319	A	1	1	1.00	27	0.037
320	A	1	1	1.00	19	0.053
321	A	13	8	1.00	21	0.381
322	A	12	8	1.00	21	0.381
323	A	8	7	1.00	21	0.333
324	A	8	7	1.00	21	0.333
325	A	4	3	1.00	21	0.143
326	A	5	4	1.00	21	0.190
327	A	7	6	1.00	21	0.286
328	A	9	8	1.00	21	0.381
329	A	10	9	1.00	21	0.429
330	A	10	9	1.00	21	0.429
331	A	9	8	1.00	21	0.381
332	A	9	8	1.00	21	0.381
333	A	9	8	1.00	21	0.381
334	A	9	8	1.00	21	0.381
335	A	9	8	1.00	21	0.381
336	A	9	8	1.00	21	0.381
337	A	10	9	1.00	21	0.429

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
338	A	10	9	1.00	21	0.429
339	A	11	9	1.00	21	0.429
340	A	11	9	1.00	21	0.429
341	A	3	2	1.00	19	0.105
342	A	5	5	1.00	19	0.263
343	A	4	4	1.00	19	0.210
344	A	3	3	0.91	17	0.176
345	A	2	2	1.00	9	0.222
346	A	2	2	1.00	19	0.105
347	A	2	2	1.00	19	0.105
348	A	2	2	1.00	19	0.105
349	A	1	1	1.00	50	0.020

Chapter 3

Listing of integrals

3.1 $\int (a + bx^2)(c + dx^2)^4 dx$

Optimal. Leaf size=94

$$\frac{1}{3}c^3x^3(4ad + bc) + \frac{2}{5}c^2dx^5(3ad + 2bc) + \frac{1}{9}d^3x^9(ad + 4bc) + \frac{2}{7}cd^2x^7(2ad + 3bc) + ac^4x + \frac{1}{11}bd^4x^{11}$$

[Out] a*c^4*x+1/3*c^3*(4*a*d+b*c)*x^3+2/5*c^2*d*(3*a*d+2*b*c)*x^5+2/7*c*d^2*(2*a*d+3*b*c)*x^7+1/9*d^3*(a*d+4*b*c)*x^9+1/11*b*d^4*x^11

Rubi [A] time = 0.06, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {373}

$$\frac{2}{5}c^2dx^5(3ad + 2bc) + \frac{1}{3}c^3x^3(4ad + bc) + \frac{1}{9}d^3x^9(ad + 4bc) + \frac{2}{7}cd^2x^7(2ad + 3bc) + ac^4x + \frac{1}{11}bd^4x^{11}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^4,x]

[Out] a*c^4*x + (c^3*(b*c + 4*a*d)*x^3)/3 + (2*c^2*d*(2*b*c + 3*a*d)*x^5)/5 + (2*c*d^2*(3*b*c + 2*a*d)*x^7)/7 + (d^3*(4*b*c + a*d)*x^9)/9 + (b*d^4*x^11)/11

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^4 dx &= \int (ac^4 + c^3(bc + 4ad)x^2 + 2c^2d(2bc + 3ad)x^4 + 2cd^2(3bc + 2ad)x^6 + d^3(4bc + ad)x^8 + bd^4x^{10}) dx \\ &= ac^4x + \frac{1}{3}c^3(bc + 4ad)x^3 + \frac{2}{5}c^2d(2bc + 3ad)x^5 + \frac{2}{7}cd^2(3bc + 2ad)x^7 + \frac{1}{9}d^3(4bc + ad)x^9 + \frac{1}{11}bd^4x^{11} \end{aligned}$$

Mathematica [A] time = 0.02, size = 94, normalized size = 1.00

$$\frac{1}{3}c^3x^3(4ad + bc) + \frac{2}{5}c^2dx^5(3ad + 2bc) + \frac{1}{9}d^3x^9(ad + 4bc) + \frac{2}{7}cd^2x^7(2ad + 3bc) + ac^4x + \frac{1}{11}bd^4x^{11}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^4,x]

[Out] $a*c^4*x + (c^3*(b*c + 4*a*d)*x^3)/3 + (2*c^2*d*(2*b*c + 3*a*d)*x^5)/5 + (2*c*d^2*(3*b*c + 2*a*d)*x^7)/7 + (d^3*(4*b*c + a*d)*x^9)/9 + (b*d^4*x^{11})/11$

fricas [A] time = 0.51, size = 98, normalized size = 1.04

$$\frac{1}{11}x^{11}d^4b + \frac{4}{9}x^9d^3cb + \frac{1}{9}x^9d^4a + \frac{6}{7}x^7d^2c^2b + \frac{4}{7}x^7d^3ca + \frac{4}{5}x^5dc^3b + \frac{6}{5}x^5d^2c^2a + \frac{1}{3}x^3c^4b + \frac{4}{3}x^3dc^3a + xc^4a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^4,x, algorithm="fricas")

[Out] $1/11*x^{11}*d^4*b + 4/9*x^9*d^3*c*b + 1/9*x^9*d^4*a + 6/7*x^7*d^2*c^2*b + 4/7*x^7*d^3*c*a + 4/5*x^5*d*c^3*b + 6/5*x^5*d^2*c^2*a + 1/3*x^3*c^4*b + 4/3*x^3*d*c^3*a + x*c^4*a$

giac [A] time = 0.57, size = 98, normalized size = 1.04

$$\frac{1}{11}bd^4x^{11} + \frac{4}{9}bcd^3x^9 + \frac{1}{9}ad^4x^9 + \frac{6}{7}bc^2d^2x^7 + \frac{4}{7}acd^3x^7 + \frac{4}{5}bc^3dx^5 + \frac{6}{5}ac^2d^2x^5 + \frac{1}{3}bc^4x^3 + \frac{4}{3}ac^3dx^3 + ac^4x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^4,x, algorithm="giac")

[Out] $1/11*b*d^4*x^{11} + 4/9*b*c*d^3*x^9 + 1/9*a*d^4*x^9 + 6/7*b*c^2*d^2*x^7 + 4/7*a*c*d^3*x^7 + 4/5*b*c^3*d*x^5 + 6/5*a*c^2*d^2*x^5 + 1/3*b*c^4*x^3 + 4/3*a*c^3*d*x^3 + a*c^4*x$

maple [A] time = 0.00, size = 97, normalized size = 1.03

$$\frac{bd^4x^{11}}{11} + \frac{(ad^4 + 4bcd^3)x^9}{9} + \frac{(4acd^3 + 6bc^2d^2)x^7}{7} + ac^4x + \frac{(6ac^2d^2 + 4bc^3d)x^5}{5} + \frac{(4ac^3d + bc^4)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^4,x)

[Out] $1/11*b*d^4*x^{11} + 1/9*(a*d^4 + 4*b*c*d^3)*x^9 + 1/7*(4*a*c*d^3 + 6*b*c^2*d^2)*x^7 + 1/5*(6*a*c^2*d^2 + 4*b*c^3*d)*x^5 + 1/3*(4*a*c^3*d + b*c^4)*x^3 + a*c^4*x$

maxima [A] time = 1.36, size = 96, normalized size = 1.02

$$\frac{1}{11}bd^4x^{11} + \frac{1}{9}(4bcd^3 + ad^4)x^9 + \frac{2}{7}(3bc^2d^2 + 2acd^3)x^7 + ac^4x + \frac{2}{5}(2bc^3d + 3ac^2d^2)x^5 + \frac{1}{3}(bc^4 + 4ac^3d)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^4,x, algorithm="maxima")

[Out] $1/11*b*d^4*x^{11} + 1/9*(4*b*c*d^3 + a*d^4)*x^9 + 2/7*(3*b*c^2*d^2 + 2*a*c*d^3)*x^7 + a*c^4*x + 2/5*(2*b*c^3*d + 3*a*c^2*d^2)*x^5 + 1/3*(b*c^4 + 4*a*c^3*d)*x^3$

mupad [B] time = 4.77, size = 88, normalized size = 0.94

$$x^3 \left(\frac{bc^4}{3} + \frac{4adc^3}{3} \right) + x^9 \left(\frac{ad^4}{9} + \frac{4bcd^3}{9} \right) + \frac{bd^4x^{11}}{11} + ac^4x + \frac{2c^2dx^5(3ad + 2bc)}{5} + \frac{2cd^2x^7(2ad + 3bc)}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^4,x)

```
[Out] x^3*((b*c^4)/3 + (4*a*c^3*d)/3) + x^9*((a*d^4)/9 + (4*b*c*d^3)/9) + (b*d^4*x^11)/11 + a*c^4*x + (2*c^2*d*x^5*(3*a*d + 2*b*c))/5 + (2*c*d^2*x^7*(2*a*d + 3*b*c))/7
```

sympy [A] time = 0.09, size = 107, normalized size = 1.14

$$ac^4x + \frac{bd^4x^{11}}{11} + x^9\left(\frac{ad^4}{9} + \frac{4bcd^3}{9}\right) + x^7\left(\frac{4acd^3}{7} + \frac{6bc^2d^2}{7}\right) + x^5\left(\frac{6ac^2d^2}{5} + \frac{4bc^3d}{5}\right) + x^3\left(\frac{4ac^3d}{3} + \frac{bc^4}{3}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)*(d*x**2+c)**4,x)
```

```
[Out] a*c**4*x + b*d**4*x**11/11 + x**9*(a*d**4/9 + 4*b*c*d**3/9) + x**7*(4*a*c*d**3/7 + 6*b*c**2*d**2/7) + x**5*(6*a*c**2*d**2/5 + 4*b*c**3*d/5) + x**3*(4*a*c**3*d/3 + b*c**4/3)
```

3.2 $\int (a + bx^2)(c + dx^2)^3 dx$

Optimal. Leaf size=70

$$\frac{1}{3}c^2x^3(3ad + bc) + \frac{1}{7}d^2x^7(ad + 3bc) + \frac{3}{5}cdx^5(ad + bc) + ac^3x + \frac{1}{9}bd^3x^9$$

[Out] $a*c^3*x + 1/3*c^2*(3*a*d+b*c)*x^3 + 3/5*c*d*(a*d+b*c)*x^5 + 1/7*d^2*(a*d+3*b*c)*x^7 + 1/9*b*d^3*x^9$

Rubi [A] time = 0.04, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {373}

$$\frac{1}{3}c^2x^3(3ad + bc) + \frac{1}{7}d^2x^7(ad + 3bc) + \frac{3}{5}cdx^5(ad + bc) + ac^3x + \frac{1}{9}bd^3x^9$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^3,x]

[Out] $a*c^3*x + (c^2*(b*c + 3*a*d)*x^3)/3 + (3*c*d*(b*c + a*d)*x^5)/5 + (d^2*(3*b*c + a*d)*x^7)/7 + (b*d^3*x^9)/9$

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^3 dx &= \int (ac^3 + c^2(bc + 3ad)x^2 + 3cd(bc + ad)x^4 + d^2(3bc + ad)x^6 + bd^3x^8) dx \\ &= ac^3x + \frac{1}{3}c^2(bc + 3ad)x^3 + \frac{3}{5}cd(bc + ad)x^5 + \frac{1}{7}d^2(3bc + ad)x^7 + \frac{1}{9}bd^3x^9 \end{aligned}$$

Mathematica [A] time = 0.02, size = 70, normalized size = 1.00

$$\frac{1}{3}c^2x^3(3ad + bc) + \frac{1}{7}d^2x^7(ad + 3bc) + \frac{3}{5}cdx^5(ad + bc) + ac^3x + \frac{1}{9}bd^3x^9$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^3,x]

[Out] $a*c^3*x + (c^2*(b*c + 3*a*d)*x^3)/3 + (3*c*d*(b*c + a*d)*x^5)/5 + (d^2*(3*b*c + a*d)*x^7)/7 + (b*d^3*x^9)/9$

fricas [A] time = 0.57, size = 73, normalized size = 1.04

$$\frac{1}{9}x^9d^3b + \frac{3}{7}x^7d^2cb + \frac{1}{7}x^7d^3a + \frac{3}{5}x^5dc^2b + \frac{3}{5}x^5d^2ca + \frac{1}{3}x^3c^3b + x^3dc^2a + xc^3a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3,x, algorithm="fricas")

[Out] $1/9*x^9*d^3*b + 3/7*x^7*d^2*c*b + 1/7*x^7*d^3*a + 3/5*x^5*d*c^2*b + 3/5*x^5*d^2*c*a + 1/3*x^3*c^3*b + x^3*d*c^2*a + x*c^3*a$

giac [A] time = 0.56, size = 73, normalized size = 1.04

$$\frac{1}{9}bd^3x^9 + \frac{3}{7}bcd^2x^7 + \frac{1}{7}ad^3x^7 + \frac{3}{5}bc^2dx^5 + \frac{3}{5}acd^2x^5 + \frac{1}{3}bc^3x^3 + ac^2dx^3 + ac^3x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3,x, algorithm="giac")

[Out] 1/9*b*d^3*x^9 + 3/7*b*c*d^2*x^7 + 1/7*a*d^3*x^7 + 3/5*b*c^2*d*x^5 + 3/5*a*c*d^2*x^5 + 1/3*b*c^3*x^3 + a*c^2*d*x^3 + a*c^3*x

maple [A] time = 0.00, size = 73, normalized size = 1.04

$$\frac{bd^3x^9}{9} + \frac{(ad^3 + 3bcd^2)x^7}{7} + ac^3x + \frac{(3acd^2 + 3bc^2d)x^5}{5} + \frac{(3ac^2d + bc^3)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^3,x)

[Out] 1/9*b*d^3*x^9+1/7*(a*d^3+3*b*c*d^2)*x^7+1/5*(3*a*c*d^2+3*b*c^2*d)*x^5+1/3*(3*a*c^2*d+b*c^3)*x^3+a*c^3*x

maxima [A] time = 1.37, size = 70, normalized size = 1.00

$$\frac{1}{9}bd^3x^9 + \frac{1}{7}(3bcd^2 + ad^3)x^7 + \frac{3}{5}(bc^2d + acd^2)x^5 + ac^3x + \frac{1}{3}(bc^3 + 3ac^2d)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3,x, algorithm="maxima")

[Out] 1/9*b*d^3*x^9 + 1/7*(3*b*c*d^2 + a*d^3)*x^7 + 3/5*(b*c^2*d + a*c*d^2)*x^5 + a*c^3*x + 1/3*(b*c^3 + 3*a*c^2*d)*x^3

mupad [B] time = 4.75, size = 65, normalized size = 0.93

$$x^3 \left(\frac{bc^3}{3} + ad^3 \right) + x^7 \left(\frac{ad^3}{7} + \frac{3bcd^2}{7} \right) + \frac{bd^3x^9}{9} + ac^3x + \frac{3cdx^5(ad+bc)}{5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^3,x)

[Out] x^3*((b*c^3)/3 + a*c^2*d) + x^7*((a*d^3)/7 + (3*b*c*d^2)/7) + (b*d^3*x^9)/9 + a*c^3*x + (3*c*d*x^5*(a*d + b*c))/5

sympy [A] time = 0.08, size = 76, normalized size = 1.09

$$ac^3x + \frac{bd^3x^9}{9} + x^7 \left(\frac{ad^3}{7} + \frac{3bcd^2}{7} \right) + x^5 \left(\frac{3acd^2}{5} + \frac{3bc^2d}{5} \right) + x^3 \left(ac^2d + \frac{bc^3}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**3,x)

[Out] a*c**3*x + b*d**3*x**9/9 + x**7*(a*d**3/7 + 3*b*c*d**2/7) + x**5*(3*a*c*d**2/5 + 3*b*c**2*d/5) + x**3*(a*c**2*d + b*c**3/3)

3.3 $\int (a + bx^2)(c + dx^2)^2 dx$

Optimal. Leaf size=50

$$\frac{1}{5}dx^5(ad + 2bc) + \frac{1}{3}cx^3(2ad + bc) + ac^2x + \frac{1}{7}bd^2x^7$$

[Out] $a*c^2*x + 1/3*c*(2*a*d+b*c)*x^3 + 1/5*d*(a*d+2*b*c)*x^5 + 1/7*b*d^2*x^7$

Rubi [A] time = 0.03, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {373}

$$\frac{1}{5}dx^5(ad + 2bc) + \frac{1}{3}cx^3(2ad + bc) + ac^2x + \frac{1}{7}bd^2x^7$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^2,x]

[Out] $a*c^2*x + (c*(b*c + 2*a*d)*x^3)/3 + (d*(2*b*c + a*d)*x^5)/5 + (b*d^2*x^7)/7$

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^2 dx &= \int (ac^2 + c(bc + 2ad)x^2 + d(2bc + ad)x^4 + bd^2x^6) dx \\ &= ac^2x + \frac{1}{3}c(bc + 2ad)x^3 + \frac{1}{5}d(2bc + ad)x^5 + \frac{1}{7}bd^2x^7 \end{aligned}$$

Mathematica [A] time = 0.01, size = 50, normalized size = 1.00

$$\frac{1}{5}dx^5(ad + 2bc) + \frac{1}{3}cx^3(2ad + bc) + ac^2x + \frac{1}{7}bd^2x^7$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^2,x]

[Out] $a*c^2*x + (c*(b*c + 2*a*d)*x^3)/3 + (d*(2*b*c + a*d)*x^5)/5 + (b*d^2*x^7)/7$

fricas [A] time = 0.54, size = 50, normalized size = 1.00

$$\frac{1}{7}x^7d^2b + \frac{2}{5}x^5dcb + \frac{1}{5}x^5d^2a + \frac{1}{3}x^3c^2b + \frac{2}{3}x^3dca + xc^2a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2,x, algorithm="fricas")

[Out] $1/7*x^7*d^2*b + 2/5*x^5*d*c*b + 1/5*x^5*d^2*a + 1/3*x^3*c^2*b + 2/3*x^3*d*c*a + x*c^2*a$

giac [A] time = 0.57, size = 50, normalized size = 1.00

$$\frac{1}{7}bd^2x^7 + \frac{2}{5}bcdx^5 + \frac{1}{5}ad^2x^5 + \frac{1}{3}bc^2x^3 + \frac{2}{3}acdx^3 + ac^2x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2,x, algorithm="giac")

[Out] $1/7*b*d^2*x^7 + 2/5*b*c*d*x^5 + 1/5*a*d^2*x^5 + 1/3*b*c^2*x^3 + 2/3*a*c*d*x^3 + a*c^2*x$

maple [A] time = 0.00, size = 49, normalized size = 0.98

$$\frac{b d^2 x^7}{7} + \frac{(a d^2 + 2 b c d) x^5}{5} + a c^2 x + \frac{(2 a c d + b c^2) x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^2,x)

[Out] $1/7*b*d^2*x^7+1/5*(a*d^2+2*b*c*d)*x^5+1/3*(2*a*c*d+b*c^2)*x^3+a*c^2*x$

maxima [A] time = 1.35, size = 48, normalized size = 0.96

$$\frac{1}{7} b d^2 x^7 + \frac{1}{5} (2 b c d + a d^2) x^5 + a c^2 x + \frac{1}{3} (b c^2 + 2 a c d) x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2,x, algorithm="maxima")

[Out] $1/7*b*d^2*x^7 + 1/5*(2*b*c*d + a*d^2)*x^5 + a*c^2*x + 1/3*(b*c^2 + 2*a*c*d)*x^3$

mupad [B] time = 0.05, size = 48, normalized size = 0.96

$$x^3 \left(\frac{b c^2}{3} + \frac{2 a d c}{3} \right) + x^5 \left(\frac{a d^2}{5} + \frac{2 b c d}{5} \right) + \frac{b d^2 x^7}{7} + a c^2 x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^2,x)

[Out] $x^3*((b*c^2)/3 + (2*a*c*d)/3) + x^5*((a*d^2)/5 + (2*b*c*d)/5) + (b*d^2*x^7)/7 + a*c^2*x$

sympy [A] time = 0.07, size = 53, normalized size = 1.06

$$a c^2 x + \frac{b d^2 x^7}{7} + x^5 \left(\frac{a d^2}{5} + \frac{2 b c d}{5} \right) + x^3 \left(\frac{2 a c d}{3} + \frac{b c^2}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**2,x)

[Out] $a*c**2*x + b*d**2*x**7/7 + x**5*(a*d**2/5 + 2*b*c*d/5) + x**3*(2*a*c*d/3 + b*c**2/3)$

3.4 $\int (a + bx^2)(c + dx^2) dx$

Optimal. Leaf size=28

$$\frac{1}{3}x^3(ad + bc) + acx + \frac{1}{5}bdx^5$$

[Out] a*c*x+1/3*(a*d+b*c)*x^3+1/5*b*d*x^5

Rubi [A] time = 0.01, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {373}

$$\frac{1}{3}x^3(ad + bc) + acx + \frac{1}{5}bdx^5$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2), x]

[Out] a*c*x + ((b*c + a*d)*x^3)/3 + (b*d*x^5)/5

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2) dx &= \int (ac + (bc + ad)x^2 + bdx^4) dx \\ &= acx + \frac{1}{3}(bc + ad)x^3 + \frac{1}{5}bdx^5 \end{aligned}$$

Mathematica [A] time = 0.01, size = 28, normalized size = 1.00

$$\frac{1}{3}x^3(ad + bc) + acx + \frac{1}{5}bdx^5$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2), x]

[Out] a*c*x + ((b*c + a*d)*x^3)/3 + (b*d*x^5)/5

fricas [A] time = 0.64, size = 26, normalized size = 0.93

$$\frac{1}{5}x^5db + \frac{1}{3}x^3cb + \frac{1}{3}x^3da + xca$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c), x, algorithm="fricas")

[Out] 1/5*x^5*d*b + 1/3*x^3*c*b + 1/3*x^3*d*a + x*c*a

giac [A] time = 0.56, size = 26, normalized size = 0.93

$$\frac{1}{5}bdx^5 + \frac{1}{3}bcx^3 + \frac{1}{3}adx^3 + acx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c),x, algorithm="giac")

[Out] 1/5*b*d*x^5 + 1/3*b*c*x^3 + 1/3*a*d*x^3 + a*c*x

maple [A] time = 0.00, size = 25, normalized size = 0.89

$$\frac{bdx^5}{5} + acx + \frac{(ad+bc)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c),x)

[Out] a*c*x+1/3*(a*d+b*c)*x^3+1/5*b*d*x^5

maxima [A] time = 1.35, size = 24, normalized size = 0.86

$$\frac{1}{5}bdx^5 + \frac{1}{3}(bc+ad)x^3 + acx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c),x, algorithm="maxima")

[Out] 1/5*b*d*x^5 + 1/3*(b*c + a*d)*x^3 + a*c*x

mupad [B] time = 0.04, size = 25, normalized size = 0.89

$$\frac{bdx^5}{5} + \left(\frac{ad}{3} + \frac{bc}{3}\right)x^3 + acx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2),x)

[Out] x^3*((a*d)/3 + (b*c)/3) + a*c*x + (b*d*x^5)/5

sympy [A] time = 0.06, size = 26, normalized size = 0.93

$$acx + \frac{bdx^5}{5} + x^3\left(\frac{ad}{3} + \frac{bc}{3}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c),x)

[Out] a*c*x + b*d*x**5/5 + x**3*(a*d/3 + b*c/3)

3.5 $\int \frac{a+bx^2}{c+dx^2} dx$

Optimal. Leaf size=40

$$\frac{bx}{d} - \frac{(bc - ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c} d^{3/2}}$$

[Out] $b*x/d - (-a*d+b*c)*\arctan(x*d^{(1/2)}/c^{(1/2)})/d^{(3/2)}/c^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {388, 205}

$$\frac{bx}{d} - \frac{(bc - ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c} d^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)/(c + d*x^2), x]

[Out] (b*x)/d - ((b*c - a*d)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(Sqrt[c]*d^(3/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int \frac{a + bx^2}{c + dx^2} dx &= \frac{bx}{d} - \frac{(bc - ad) \int \frac{1}{c + dx^2} dx}{d} \\ &= \frac{bx}{d} - \frac{(bc - ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c} d^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 40, normalized size = 1.00

$$\frac{bx}{d} - \frac{(bc - ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c} d^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)/(c + d*x^2), x]

[Out] (b*x)/d - ((b*c - a*d)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(Sqrt[c]*d^(3/2))

fricas [A] time = 0.79, size = 99, normalized size = 2.48

$$\left[\frac{2 b c d x + (b c - a d) \sqrt{-c d} \log \left(\frac{d x^2 - 2 \sqrt{-c d} x - c}{d x^2 + c} \right)}{2 c d^2}, \frac{b c d x - (b c - a d) \sqrt{c d} \arctan \left(\frac{\sqrt{c d} x}{c} \right)}{c d^2} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c),x, algorithm="fricas")

[Out] [1/2*(2*b*c*d*x + (b*c - a*d)*sqrt(-c*d)*log((d*x^2 - 2*sqrt(-c*d)*x - c)/(d*x^2 + c)))/(c*d^2), (b*c*d*x - (b*c - a*d)*sqrt(c*d)*arctan(sqrt(c*d)*x/c))/(c*d^2)]

giac [A] time = 0.57, size = 34, normalized size = 0.85

$$\frac{b x}{d} - \frac{(b c - a d) \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c),x, algorithm="giac")

[Out] b*x/d - (b*c - a*d)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*d)

maple [A] time = 0.01, size = 45, normalized size = 1.12

$$\frac{a \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d}} - \frac{b c \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d} d} + \frac{b x}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)/(d*x^2+c),x)

[Out] b*x/d+1/(c*d)^(1/2)*arctan(x*d/(c*d)^(1/2))*a-1/d/(c*d)^(1/2)*arctan(x*d/(c*d)^(1/2))*b*c

maxima [A] time = 3.00, size = 34, normalized size = 0.85

$$\frac{b x}{d} - \frac{(b c - a d) \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c),x, algorithm="maxima")

[Out] b*x/d - (b*c - a*d)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*d)

mupad [B] time = 0.06, size = 31, normalized size = 0.78

$$\frac{b x}{d} + \frac{\operatorname{atan} \left(\frac{\sqrt{d} x}{\sqrt{c}} \right) (a d - b c)}{\sqrt{c} d^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)/(c + d*x^2),x)

[Out] (b*x)/d + (atan((d^(1/2)*x)/c^(1/2))*(a*d - b*c))/(c^(1/2)*d^(3/2))

sympy [B] time = 0.28, size = 82, normalized size = 2.05

$$\frac{bx}{d} - \frac{\sqrt{-\frac{1}{cd^3}} (ad - bc) \log\left(-cd\sqrt{-\frac{1}{cd^3}} + x\right)}{2} + \frac{\sqrt{-\frac{1}{cd^3}} (ad - bc) \log\left(cd\sqrt{-\frac{1}{cd^3}} + x\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)/(d*x**2+c),x)

[Out] b*x/d - sqrt(-1/(c*d**3))*(a*d - b*c)*log(-c*d*sqrt(-1/(c*d**3)) + x)/2 + s
qrt(-1/(c*d**3))*(a*d - b*c)*log(c*d*sqrt(-1/(c*d**3)) + x)/2

$$3.6 \quad \int \frac{a+bx^2}{(c+dx^2)^2} dx$$

Optimal. Leaf size=63

$$\frac{(ad + bc) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{3/2}} - \frac{x(bc - ad)}{2cd(c + dx^2)}$$

[Out] $-1/2*(-a*d+b*c)*x/c/d/(d*x^2+c)+1/2*(a*d+b*c)*\arctan(x*d^{(1/2)}/c^{(1/2)})/c^{(3/2)}/d^{(3/2)}$

Rubi [A] time = 0.02, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {385, 205}

$$\frac{(ad + bc) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{3/2}} - \frac{x(bc - ad)}{2cd(c + dx^2)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)/(c + d*x^2)^2, x]

[Out] $-((b*c - a*d)*x)/(2*c*d*(c + d*x^2)) + ((b*c + a*d)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(2*c^{(3/2)}*d^{(3/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rubi steps

$$\begin{aligned} \int \frac{a + bx^2}{(c + dx^2)^2} dx &= -\frac{(bc - ad)x}{2cd(c + dx^2)} + \frac{(bc + ad) \int \frac{1}{c + dx^2} dx}{2cd} \\ &= -\frac{(bc - ad)x}{2cd(c + dx^2)} + \frac{(bc + ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 63, normalized size = 1.00

$$\frac{(ad + bc) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{3/2}} - \frac{x(bc - ad)}{2cd(c + dx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)/(c + d*x^2)^2,x]

[Out] $-1/2*((b*c - a*d)*x)/(c*d*(c + d*x^2)) + ((b*c + a*d)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(2*c^{3/2}*d^{3/2})$

fricas [A] time = 0.66, size = 182, normalized size = 2.89

$$\left[\frac{(bc^2 + acd + (bcd + ad^2)x^2)\sqrt{-cd} \log\left(\frac{dx^2 - 2\sqrt{-cd}x - c}{dx^2 + c}\right) + 2(bc^2d - acd^2)x}{4(c^2d^3x^2 + c^3d^2)}, \frac{(bc^2 + acd + (bcd + ad^2)x^2)\sqrt{cd} \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2(c^2d^3x^2 + c^3d^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] $[-1/4*((b*c^2 + a*c*d + (b*c*d + a*d^2)*x^2)*\text{sqrt}(-c*d)*\log((d*x^2 - 2*\text{sqrt}(-c*d)*x - c)/(d*x^2 + c)) + 2*(b*c^2*d - a*c*d^2)*x)/(c^2*d^3*x^2 + c^3*d^2), 1/2*((b*c^2 + a*c*d + (b*c*d + a*d^2)*x^2)*\text{sqrt}(c*d)*\text{arctan}(\text{sqrt}(c*d)*x/c) - (b*c^2*d - a*c*d^2)*x)/(c^2*d^3*x^2 + c^3*d^2)]$

giac [A] time = 0.58, size = 57, normalized size = 0.90

$$\frac{(bc + ad) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}cd} - \frac{bcx - adx}{2(dx^2 + c)cd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^2,x, algorithm="giac")

[Out] $1/2*(b*c + a*d)*\text{arctan}(d*x/\text{sqrt}(c*d))/(\text{sqrt}(c*d)*c*d) - 1/2*(b*c*x - a*d*x)/((d*x^2 + c)*c*d)$

maple [A] time = 0.01, size = 68, normalized size = 1.08

$$\frac{a \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}c} + \frac{b \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}d} + \frac{(ad - bc)x}{2(dx^2 + c)cd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)/(d*x^2+c)^2,x)

[Out] $1/2*(a*d-b*c)/c/d*x/(d*x^2+c)+1/2/c/(c*d)^{(1/2)*\text{arctan}(1/(c*d)^{(1/2)*d*x})*a+1/2/d/(c*d)^{(1/2)*\text{arctan}(1/(c*d)^{(1/2)*d*x})*b}$

maxima [A] time = 3.11, size = 57, normalized size = 0.90

$$-\frac{(bc - ad)x}{2(cd^2x^2 + c^2d)} + \frac{(bc + ad) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}cd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] $-1/2*(b*c - a*d)*x/(c*d^2*x^2 + c^2*d) + 1/2*(b*c + a*d)*\text{arctan}(d*x/\text{sqrt}(c*d))/(\text{sqrt}(c*d)*c*d)$

mupad [B] time = 5.01, size = 51, normalized size = 0.81

$$\frac{\text{atan}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)(ad + bc)}{2c^{3/2}d^{3/2}} + \frac{x(ad - bc)}{2cd(dx^2 + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)/(c + d*x^2)^2,x)`

[Out] $(\operatorname{atan}((d^{1/2}x)/c^{1/2})*(a*d + b*c))/(2*c^{3/2}*d^{3/2}) + (x*(a*d - b*c))/(2*c*d*(c + d*x^2))$

sympy [B] time = 0.39, size = 112, normalized size = 1.78

$$\frac{x(ad - bc)}{2c^2d + 2cd^2x^2} - \frac{\sqrt{-\frac{1}{c^3d^3}}(ad + bc)\log\left(-c^2d\sqrt{-\frac{1}{c^3d^3}} + x\right)}{4} + \frac{\sqrt{-\frac{1}{c^3d^3}}(ad + bc)\log\left(c^2d\sqrt{-\frac{1}{c^3d^3}} + x\right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)/(d*x**2+c)**2,x)`

[Out] $x*(a*d - b*c)/(2*c**2*d + 2*c*d**2*x**2) - \operatorname{sqrt}(-1/(c**3*d**3))*(a*d + b*c)*\log(-c**2*d*\operatorname{sqrt}(-1/(c**3*d**3)) + x)/4 + \operatorname{sqrt}(-1/(c**3*d**3))*(a*d + b*c)*\log(c**2*d*\operatorname{sqrt}(-1/(c**3*d**3)) + x)/4$

$$3.7 \quad \int \frac{a+bx^2}{(c+dx^2)^3} dx$$

Optimal. Leaf size=92

$$\frac{(3ad + bc) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{3/2}} + \frac{x(3ad + bc)}{8c^2d(c + dx^2)} - \frac{x(bc - ad)}{4cd(c + dx^2)^2}$$

[Out] $-1/4*(-a*d+b*c)*x/c/d/(d*x^2+c)^2+1/8*(3*a*d+b*c)*x/c^2/d/(d*x^2+c)+1/8*(3*a*d+b*c)*\arctan(x*d^{(1/2)}/c^{(1/2)})/c^{(5/2)}/d^{(3/2)}$

Rubi [A] time = 0.03, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {385, 199, 205}

$$\frac{(3ad + bc) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{3/2}} + \frac{x(3ad + bc)}{8c^2d(c + dx^2)} - \frac{x(bc - ad)}{4cd(c + dx^2)^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)/(c + d*x^2)^3, x]

[Out] $-((b*c - a*d)*x)/(4*c*d*(c + d*x^2)^2) + ((b*c + 3*a*d)*x)/(8*c^2*d*(c + d*x^2)) + ((b*c + 3*a*d)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(8*c^{(5/2)}*d^{(3/2)})$

Rule 199

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rubi steps

$$\begin{aligned}
\int \frac{a + bx^2}{(c + dx^2)^3} dx &= -\frac{(bc - ad)x}{4cd(c + dx^2)^2} + \frac{(bc + 3ad) \int \frac{1}{(c+dx^2)^2} dx}{4cd} \\
&= -\frac{(bc - ad)x}{4cd(c + dx^2)^2} + \frac{(bc + 3ad)x}{8c^2d(c + dx^2)} + \frac{(bc + 3ad) \int \frac{1}{c+dx^2} dx}{8c^2d} \\
&= -\frac{(bc - ad)x}{4cd(c + dx^2)^2} + \frac{(bc + 3ad)x}{8c^2d(c + dx^2)} + \frac{(bc + 3ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 82, normalized size = 0.89

$$\frac{(3ad + bc) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{3/2}} + \frac{x(ad(5c + 3dx^2) + bc(dx^2 - c))}{8c^2d(c + dx^2)^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)/(c + d*x^2)^3,x]

[Out] (x*(b*c*(-c + d*x^2) + a*d*(5*c + 3*d*x^2)))/(8*c^2*d*(c + d*x^2)^2) + ((b*c + 3*a*d)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(8*c^(5/2)*d^(3/2))

fricas [A] time = 0.66, size = 300, normalized size = 3.26

$$\left[\frac{2(bc^2d^2 + 3acd^3)x^3 - ((bcd^2 + 3ad^3)x^4 + bc^3 + 3ac^2d + 2(bc^2d + 3acd^2)x^2)\sqrt{-cd} \log\left(\frac{dx^2 - 2\sqrt{-cd}x - c}{dx^2 + c}\right) - 2}{16(c^3d^4x^4 + 2c^4d^3x^2 + c^5d^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [1/16*(2*(b*c^2*d^2 + 3*a*c*d^3)*x^3 - ((b*c*d^2 + 3*a*d^3)*x^4 + b*c^3 + 3*a*c^2*d + 2*(b*c^2*d + 3*a*c*d^2)*x^2)*sqrt(-c*d)*log((d*x^2 - 2*sqrt(-c*d)*x - c)/(d*x^2 + c)) - 2*(b*c^3*d - 5*a*c^2*d^2)*x)/(c^3*d^4*x^4 + 2*c^4*d^3*x^2 + c^5*d^2), 1/8*((b*c^2*d^2 + 3*a*c*d^3)*x^3 + ((b*c*d^2 + 3*a*d^3)*x^4 + b*c^3 + 3*a*c^2*d + 2*(b*c^2*d + 3*a*c*d^2)*x^2)*sqrt(c*d)*arctan(sqrt(c*d)*x/c) - (b*c^3*d - 5*a*c^2*d^2)*x)/(c^3*d^4*x^4 + 2*c^4*d^3*x^2 + c^5*d^2)]

giac [A] time = 0.59, size = 78, normalized size = 0.85

$$\frac{(bc + 3ad) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd}c^2d} + \frac{bcdx^3 + 3ad^2x^3 - bc^2x + 5acdx}{8(dx^2 + c)^2c^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^3,x, algorithm="giac")

[Out] 1/8*(b*c + 3*a*d)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*c^2*d) + 1/8*(b*c*d*x^3 + 3*a*d^2*x^3 - b*c^2*x + 5*a*c*d*x)/((d*x^2 + c)^2*c^2*d)

maple [A] time = 0.01, size = 90, normalized size = 0.98

$$\frac{3a \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd} c^2} + \frac{b \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd} cd} + \frac{\frac{(3ad+bc)x^3}{8c^2} + \frac{(5ad-bc)x}{8cd}}{(dx^2 + c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)/(d*x^2+c)^3,x)

[Out] (1/8*(3*a*d+b*c)/c^2*x^3+1/8*(5*a*d-b*c)/c/d*x)/(d*x^2+c)^2+3/8/c^2/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a+1/8/c/d/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*b

maxima [A] time = 2.96, size = 92, normalized size = 1.00

$$\frac{(bcd + 3ad^2)x^3 - (bc^2 - 5acd)x}{8(c^2d^3x^4 + 2c^3d^2x^2 + c^4d)} + \frac{(bc + 3ad) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd} c^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^3,x, algorithm="maxima")

[Out] 1/8*((b*c*d + 3*a*d^2)*x^3 - (b*c^2 - 5*a*c*d)*x)/(c^2*d^3*x^4 + 2*c^3*d^2*x^2 + c^4*d) + 1/8*(b*c + 3*a*d)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*c^2*d)

mupad [B] time = 5.06, size = 82, normalized size = 0.89

$$\frac{\frac{x^3(3ad+bc)}{8c^2} + \frac{x(5ad-bc)}{8cd}}{c^2 + 2cdx^2 + d^2x^4} + \frac{\operatorname{atan}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)(3ad+bc)}{8c^{5/2}d^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)/(c + d*x^2)^3,x)

[Out] ((x^3*(3*a*d + b*c))/(8*c^2) + (x*(5*a*d - b*c))/(8*c*d))/(c^2 + d^2*x^4 + 2*c*d*x^2) + (atan((d^(1/2)*x)/c^(1/2))*(3*a*d + b*c))/(8*c^(5/2)*d^(3/2))

sympy [A] time = 0.54, size = 150, normalized size = 1.63

$$-\frac{\sqrt{-\frac{1}{c^5d^3}}(3ad+bc)\log\left(-c^3d\sqrt{-\frac{1}{c^5d^3}}+x\right)}{16} + \frac{\sqrt{-\frac{1}{c^5d^3}}(3ad+bc)\log\left(c^3d\sqrt{-\frac{1}{c^5d^3}}+x\right)}{16} + \frac{x^3(3ad^2+bcd)+x(5acd)}{8c^4d+16c^3d^2x^2+8c^2d^2x^2+8c^2d^2x^2+8c^2d^2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)/(d*x**2+c)**3,x)

[Out] -sqrt(-1/(c**5*d**3))*(3*a*d + b*c)*log(-c**3*d*sqrt(-1/(c**5*d**3)) + x)/16 + sqrt(-1/(c**5*d**3))*(3*a*d + b*c)*log(c**3*d*sqrt(-1/(c**5*d**3)) + x)/16 + (x**3*(3*a*d**2 + b*c*d) + x*(5*a*c*d - b*c**2))/(8*c**4*d + 16*c**3*d**2*x**2 + 8*c**2*d**3*x**4)

3.8 $\int (a + bx^2)^2 (c + dx^2)^3 dx$

Optimal. Leaf size=122

$$\frac{1}{7}dx^7 (a^2d^2 + 6abcd + 3b^2c^2) + \frac{1}{5}cx^5 (3a^2d^2 + 6abcd + b^2c^2) + a^2c^3x + \frac{1}{3}ac^2x^3(3ad+2bc) + \frac{1}{9}bd^2x^9(2ad+3bc) + \frac{1}{11}b^2d^3x^{11}$$

[Out] a²*c³*x+1/3*a*c²*(3*a*d+2*b*c)*x³+1/5*c*(3*a²*d²+6*a*b*c*d+b²*c²)*x⁵+1/7*d*(a²*d²+6*a*b*c*d+3*b²*c²)*x⁷+1/9*b*d²*(2*a*d+3*b*c)*x⁹+1/11*b²*d³*x¹¹

Rubi [A] time = 0.07, antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {373}

$$\frac{1}{7}dx^7 (a^2d^2 + 6abcd + 3b^2c^2) + \frac{1}{5}cx^5 (3a^2d^2 + 6abcd + b^2c^2) + a^2c^3x + \frac{1}{3}ac^2x^3(3ad+2bc) + \frac{1}{9}bd^2x^9(2ad+3bc) + \frac{1}{11}b^2d^3x^{11}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^2*(c + d*x^2)^3,x]

[Out] a²*c³*x + (a*c²*(2*b*c + 3*a*d)*x³)/3 + (c*(b²*c² + 6*a*b*c*d + 3*a²*d²)*x⁵)/5 + (d*(3*b²*c² + 6*a*b*c*d + a²*d²)*x⁷)/7 + (b*d²*(3*b*c + 2*a*d)*x⁹)/9 + (b²*d³*x¹¹)/11

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^2 (c + dx^2)^3 dx &= \int (a^2c^3 + ac^2(2bc + 3ad)x^2 + c(b^2c^2 + 6abcd + 3a^2d^2)x^4 + d(3b^2c^2 + 6abcd + 3a^2d^2)x^6 + dx^8(3bd^2c + 2ad^2b) + d^2x^{10}(3bd^2c + 2ad^2b) + d^3x^{12}) dx \\ &= a^2c^3x + \frac{1}{3}ac^2(2bc + 3ad)x^3 + \frac{1}{5}c(b^2c^2 + 6abcd + 3a^2d^2)x^5 + \frac{1}{7}d(3b^2c^2 + 6abcd + 3a^2d^2)x^7 + \frac{1}{9}bd^2x^9(2ad+3bc) + \frac{1}{11}b^2d^3x^{11} \end{aligned}$$

Mathematica [A] time = 0.02, size = 122, normalized size = 1.00

$$\frac{1}{7}dx^7 (a^2d^2 + 6abcd + 3b^2c^2) + \frac{1}{5}cx^5 (3a^2d^2 + 6abcd + b^2c^2) + a^2c^3x + \frac{1}{3}ac^2x^3(3ad+2bc) + \frac{1}{9}bd^2x^9(2ad+3bc) + \frac{1}{11}b^2d^3x^{11}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^2*(c + d*x^2)^3,x]

[Out] a²*c³*x + (a*c²*(2*b*c + 3*a*d)*x³)/3 + (c*(b²*c² + 6*a*b*c*d + 3*a²*d²)*x⁵)/5 + (d*(3*b²*c² + 6*a*b*c*d + a²*d²)*x⁷)/7 + (b*d²*(3*b*c + 2*a*d)*x⁹)/9 + (b²*d³*x¹¹)/11

fricas [A] time = 0.46, size = 131, normalized size = 1.07

$$\frac{1}{11}x^{11}d^3b^2 + \frac{1}{3}x^9d^2cb^2 + \frac{2}{9}x^9d^3ba + \frac{3}{7}x^7dc^2b^2 + \frac{6}{7}x^7d^2cba + \frac{1}{7}x^7d^3a^2 + \frac{1}{5}x^5c^3b^2 + \frac{6}{5}x^5dc^2ba + \frac{3}{5}x^5d^2ca^2 + \frac{2}{3}x^3c^3ba + x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c)^3,x, algorithm="fricas")

[Out] $\frac{1}{11}x^{11}d^3b^2 + \frac{1}{3}x^9d^2c^2b^2 + \frac{2}{9}x^9d^3b^2a + \frac{3}{7}x^7d^2c^2b^2 + \frac{6}{7}x^7d^2c^2b^2a + \frac{1}{7}x^7d^3a^2 + \frac{1}{5}x^5c^3b^2 + \frac{6}{5}x^5d^2c^2b^2a + \frac{3}{5}x^5d^2c^2a^2 + \frac{2}{3}x^3c^3b^2a + x^3d^2c^2a^2 + x^3c^3a^2$

giac [A] time = 0.56, size = 131, normalized size = 1.07

$$\frac{1}{11}b^2d^3x^{11} + \frac{1}{3}b^2cd^2x^9 + \frac{2}{9}abd^3x^9 + \frac{3}{7}b^2c^2dx^7 + \frac{6}{7}abcd^2x^7 + \frac{1}{7}a^2d^3x^7 + \frac{1}{5}b^2c^3x^5 + \frac{6}{5}abc^2dx^5 + \frac{3}{5}a^2cd^2x^5 + \frac{2}{3}abc^3x^3 + \frac{1}{3}a^2c^2d^2x^3 + \frac{1}{3}a^2c^3x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c)^3,x, algorithm="giac")

[Out] $\frac{1}{11}b^2d^3x^{11} + \frac{1}{3}b^2c^2d^2x^9 + \frac{2}{9}a^2b^2d^3x^9 + \frac{3}{7}b^2c^2d^2x^7 + \frac{6}{7}a^2b^2c^2d^2x^7 + \frac{1}{7}a^2b^2c^3x^5 + \frac{6}{5}a^2b^2c^2d^2x^5 + \frac{3}{5}a^2c^2d^2x^5 + \frac{2}{3}a^2b^2c^3x^3 + a^2c^2d^2x^3 + a^2c^3x^3$

maple [A] time = 0.00, size = 125, normalized size = 1.02

$$\frac{b^2d^3x^{11}}{11} + \frac{(2abd^3 + 3b^2cd^2)x^9}{9} + \frac{(a^2d^3 + 6abcd^2 + 3b^2c^2d)x^7}{7} + a^2c^3x + \frac{(3a^2cd^2 + 6abc^2d + b^2c^3)x^5}{5} + \frac{(3a^2c^2d + 2a^2c^3)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^2*(d*x^2+c)^3,x)

[Out] $\frac{1}{11}b^2d^3x^{11} + \frac{1}{9}(2abd^3 + 3b^2cd^2)x^9 + \frac{1}{7}(a^2d^3 + 6abcd^2 + 3b^2c^2d)x^7 + \frac{1}{5}(3a^2cd^2 + 6abc^2d + b^2c^3)x^5 + \frac{1}{3}(3a^2c^2d + 2a^2c^3)x^3 + a^2c^3x$

maxima [A] time = 1.39, size = 124, normalized size = 1.02

$$\frac{1}{11}b^2d^3x^{11} + \frac{1}{9}(3b^2cd^2 + 2abd^3)x^9 + \frac{1}{7}(3b^2c^2d + 6abcd^2 + a^2d^3)x^7 + a^2c^3x + \frac{1}{5}(b^2c^3 + 6abc^2d + 3a^2cd^2)x^5 + \frac{1}{3}(2a^2c^2d + a^2c^3)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c)^3,x, algorithm="maxima")

[Out] $\frac{1}{11}b^2d^3x^{11} + \frac{1}{9}(3b^2cd^2 + 2abd^3)x^9 + \frac{1}{7}(3b^2c^2d + 6abcd^2 + a^2d^3)x^7 + a^2c^3x + \frac{1}{5}(b^2c^3 + 6abc^2d + 3a^2cd^2)x^5 + \frac{1}{3}(2a^2c^2d + a^2c^3)x^3$

mupad [B] time = 4.95, size = 116, normalized size = 0.95

$$x^5 \left(\frac{3a^2cd^2}{5} + \frac{6abc^2d}{5} + \frac{b^2c^3}{5} \right) + x^7 \left(\frac{a^2d^3}{7} + \frac{6abcd^2}{7} + \frac{3b^2c^2d}{7} \right) + a^2c^3x + \frac{b^2d^3x^{11}}{11} + \frac{ac^2x^3(3ad + 2bc)}{3} + \frac{b^2c^3x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^2*(c + d*x^2)^3,x)

[Out] $x^5 \left(\frac{b^2c^3}{5} + \frac{3a^2cd^2}{5} + \frac{6abc^2d}{5} \right) + x^7 \left(\frac{a^2d^3}{7} + \frac{6abcd^2}{7} + \frac{3b^2c^2d}{7} \right) + a^2c^3x + \frac{b^2d^3x^{11}}{11} + \frac{ac^2x^3(3ad + 2bc)}{3} + \frac{b^2c^3x^3}{3}$

sympy [A] time = 0.09, size = 136, normalized size = 1.11

$$a^2c^3x + \frac{b^2d^3x^{11}}{11} + x^9 \left(\frac{2abd^3}{9} + \frac{b^2cd^2}{3} \right) + x^7 \left(\frac{a^2d^3}{7} + \frac{6abcd^2}{7} + \frac{3b^2c^2d}{7} \right) + x^5 \left(\frac{3a^2cd^2}{5} + \frac{6abc^2d}{5} + \frac{b^2c^3}{5} \right) + x^3 \left(\frac{a^2c^2d}{3} + \frac{2a^2c^3}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**2*(d*x**2+c)**3,x)

[Out] $a^2c^3x + \frac{b^2d^3x^{11}}{11} + x^9 \left(\frac{2abd^3}{9} + \frac{b^2cd^2}{3} \right) + x^7 \left(\frac{a^2d^3}{7} + \frac{6abcd^2}{7} + \frac{3b^2c^2d}{7} \right) + x^5 \left(\frac{3a^2cd^2}{5} + \frac{6abc^2d}{5} + \frac{b^2c^3}{5} \right) + x^3 \left(\frac{a^2c^2d}{3} + \frac{2a^2c^3}{3} \right)$

3.9 $\int (a + bx^2)^2 (c + dx^2)^2 dx$

Optimal. Leaf size=82

$$\frac{1}{5}x^5(a^2d^2 + 4abcd + b^2c^2) + a^2c^2x + \frac{2}{7}bdx^7(ad + bc) + \frac{2}{3}acx^3(ad + bc) + \frac{1}{9}b^2d^2x^9$$

[Out] $a^2c^2x + 2/3*a*c*(a*d+b*c)*x^3 + 1/5*(a^2*d^2+4*a*b*c*d+b^2*c^2)*x^5 + 2/7*b*d*(a*d+b*c)*x^7 + 1/9*b^2*d^2*x^9$

Rubi [A] time = 0.05, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {373}

$$\frac{1}{5}x^5(a^2d^2 + 4abcd + b^2c^2) + a^2c^2x + \frac{2}{7}bdx^7(ad + bc) + \frac{2}{3}acx^3(ad + bc) + \frac{1}{9}b^2d^2x^9$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^2*(c + d*x^2)^2,x]

[Out] $a^2c^2x + (2*a*c*(b*c + a*d)*x^3)/3 + ((b^2*c^2 + 4*a*b*c*d + a^2*d^2)*x^5)/5 + (2*b*d*(b*c + a*d)*x^7)/7 + (b^2*d^2*x^9)/9$

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^2 (c + dx^2)^2 dx &= \int (a^2c^2 + 2ac(bc + ad)x^2 + (b^2c^2 + 4abcd + a^2d^2)x^4 + 2bd(bc + ad)x^6 + b^2d^2x^8) dx \\ &= a^2c^2x + \frac{2}{3}ac(bc + ad)x^3 + \frac{1}{5}(b^2c^2 + 4abcd + a^2d^2)x^5 + \frac{2}{7}bd(bc + ad)x^7 + \frac{1}{9}b^2d^2x^9 \end{aligned}$$

Mathematica [A] time = 0.02, size = 82, normalized size = 1.00

$$\frac{1}{5}x^5(a^2d^2 + 4abcd + b^2c^2) + a^2c^2x + \frac{2}{7}bdx^7(ad + bc) + \frac{2}{3}acx^3(ad + bc) + \frac{1}{9}b^2d^2x^9$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^2*(c + d*x^2)^2,x]

[Out] $a^2c^2x + (2*a*c*(b*c + a*d)*x^3)/3 + ((b^2*c^2 + 4*a*b*c*d + a^2*d^2)*x^5)/5 + (2*b*d*(b*c + a*d)*x^7)/7 + (b^2*d^2*x^9)/9$

fricas [A] time = 0.42, size = 91, normalized size = 1.11

$$\frac{1}{9}x^9d^2b^2 + \frac{2}{7}x^7dcb^2 + \frac{2}{7}x^7d^2ba + \frac{1}{5}x^5c^2b^2 + \frac{4}{5}x^5dcba + \frac{1}{5}x^5d^2a^2 + \frac{2}{3}x^3c^2ba + \frac{2}{3}x^3dca^2 + xc^2a^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c)^2,x, algorithm="fricas")

[Out] $1/9*x^9*d^2*b^2 + 2/7*x^7*d*c*b^2 + 2/7*x^7*d^2*b*a + 1/5*x^5*c^2*b^2 + 4/5*x^5*d*c*b*a + 1/5*x^5*d^2*a^2 + 2/3*x^3*c^2*b*a + 2/3*x^3*d*c*a^2 + x*c^2*a^2$

giac [A] time = 0.57, size = 91, normalized size = 1.11

$$\frac{1}{9}b^2d^2x^9 + \frac{2}{7}b^2cdx^7 + \frac{2}{7}abd^2x^7 + \frac{1}{5}b^2c^2x^5 + \frac{4}{5}abcdx^5 + \frac{1}{5}a^2d^2x^5 + \frac{2}{3}abc^2x^3 + \frac{2}{3}a^2cdx^3 + a^2c^2x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c)^2,x, algorithm="giac")

[Out] 1/9*b^2*d^2*x^9 + 2/7*b^2*c*d*x^7 + 2/7*a*b*d^2*x^7 + 1/5*b^2*c^2*x^5 + 4/5*a*b*c*d*x^5 + 1/5*a^2*d^2*x^5 + 2/3*a*b*c^2*x^3 + 2/3*a^2*c*d*x^3 + a^2*c^2*x

maple [A] time = 0.00, size = 87, normalized size = 1.06

$$\frac{b^2d^2x^9}{9} + \frac{(2abd^2 + 2b^2cd)x^7}{7} + a^2c^2x + \frac{(a^2d^2 + 4abcd + b^2c^2)x^5}{5} + \frac{(2a^2cd + 2abc^2)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^2*(d*x^2+c)^2,x)

[Out] 1/9*b^2*d^2*x^9+1/7*(2*a*b*d^2+2*b^2*c*d)*x^7+1/5*(a^2*d^2+4*a*b*c*d+b^2*c^2)*x^5+1/3*(2*a^2*c*d+2*a*b*c^2)*x^3+a^2*c^2*x

maxima [A] time = 1.33, size = 82, normalized size = 1.00

$$\frac{1}{9}b^2d^2x^9 + \frac{2}{7}(b^2cd + abd^2)x^7 + \frac{1}{5}(b^2c^2 + 4abcd + a^2d^2)x^5 + a^2c^2x + \frac{2}{3}(abc^2 + a^2cd)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c)^2,x, algorithm="maxima")

[Out] 1/9*b^2*d^2*x^9 + 2/7*(b^2*c*d + a*b*d^2)*x^7 + 1/5*(b^2*c^2 + 4*a*b*c*d + a^2*d^2)*x^5 + a^2*c^2*x + 2/3*(a*b*c^2 + a^2*c*d)*x^3

mapad [B] time = 0.05, size = 75, normalized size = 0.91

$$x^5 \left(\frac{a^2d^2}{5} + \frac{4abcd}{5} + \frac{b^2c^2}{5} \right) + a^2c^2x + \frac{b^2d^2x^9}{9} + \frac{2acx^3(ad+bc)}{3} + \frac{2bdx^7(ad+bc)}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^2*(c + d*x^2)^2,x)

[Out] x^5*((a^2*d^2)/5 + (b^2*c^2)/5 + (4*a*b*c*d)/5) + a^2*c^2*x + (b^2*d^2*x^9)/9 + (2*a*c*x^3*(a*d + b*c))/3 + (2*b*d*x^7*(a*d + b*c))/7

sympy [A] time = 0.08, size = 97, normalized size = 1.18

$$a^2c^2x + \frac{b^2d^2x^9}{9} + x^7 \left(\frac{2abd^2}{7} + \frac{2b^2cd}{7} \right) + x^5 \left(\frac{a^2d^2}{5} + \frac{4abcd}{5} + \frac{b^2c^2}{5} \right) + x^3 \left(\frac{2a^2cd}{3} + \frac{2abc^2}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**2*(d*x**2+c)**2,x)

[Out] a**2*c**2*x + b**2*d**2*x**9/9 + x**7*(2*a*b*d**2/7 + 2*b**2*c*d/7) + x**5*(a**2*d**2/5 + 4*a*b*c*d/5 + b**2*c**2/5) + x**3*(2*a**2*c*d/3 + 2*a*b*c**2/3)

3.10 $\int (a + bx^2)^2 (c + dx^2) dx$

Optimal. Leaf size=50

$$a^2cx + \frac{1}{5}bx^5(2ad + bc) + \frac{1}{3}ax^3(ad + 2bc) + \frac{1}{7}b^2dx^7$$

[Out] $a^2c*x + 1/3*a*(a*d+2*b*c)*x^3 + 1/5*b*(2*a*d+b*c)*x^5 + 1/7*b^2*d*x^7$

Rubi [A] time = 0.03, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {373}

$$a^2cx + \frac{1}{5}bx^5(2ad + bc) + \frac{1}{3}ax^3(ad + 2bc) + \frac{1}{7}b^2dx^7$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^2*(c + d*x^2), x]

[Out] $a^2c*x + (a*(2*b*c + a*d)*x^3)/3 + (b*(b*c + 2*a*d)*x^5)/5 + (b^2*d*x^7)/7$

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^2 (c + dx^2) dx &= \int (a^2c + a(2bc + ad)x^2 + b(bc + 2ad)x^4 + b^2dx^6) dx \\ &= a^2cx + \frac{1}{3}a(2bc + ad)x^3 + \frac{1}{5}b(bc + 2ad)x^5 + \frac{1}{7}b^2dx^7 \end{aligned}$$

Mathematica [A] time = 0.01, size = 50, normalized size = 1.00

$$a^2cx + \frac{1}{5}bx^5(2ad + bc) + \frac{1}{3}ax^3(ad + 2bc) + \frac{1}{7}b^2dx^7$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^2*(c + d*x^2), x]

[Out] $a^2c*x + (a*(2*b*c + a*d)*x^3)/3 + (b*(b*c + 2*a*d)*x^5)/5 + (b^2*d*x^7)/7$

fricas [A] time = 0.57, size = 50, normalized size = 1.00

$$\frac{1}{7}x^7db^2 + \frac{1}{5}x^5cb^2 + \frac{2}{5}x^5dba + \frac{2}{3}x^3cba + \frac{1}{3}x^3da^2 + xca^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c), x, algorithm="fricas")

[Out] $1/7*x^7*d*b^2 + 1/5*x^5*c*b^2 + 2/5*x^5*d*b*a + 2/3*x^3*c*b*a + 1/3*x^3*d*a^2 + x*c*a^2$

giac [A] time = 0.57, size = 50, normalized size = 1.00

$$\frac{1}{7}b^2dx^7 + \frac{1}{5}b^2cx^5 + \frac{2}{5}abdx^5 + \frac{2}{3}abcx^3 + \frac{1}{3}a^2dx^3 + a^2cx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c),x, algorithm="giac")

[Out] $1/7*b^2*d*x^7 + 1/5*b^2*c*x^5 + 2/5*a*b*d*x^5 + 2/3*a*b*c*x^3 + 1/3*a^2*d*x^3 + a^2*c*x$

maple [A] time = 0.00, size = 49, normalized size = 0.98

$$\frac{b^2 d x^7}{7} + \frac{(2 a b d + b^2 c) x^5}{5} + a^2 c x + \frac{(a^2 d + 2 a b c) x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^2*(d*x^2+c),x)

[Out] $1/7*b^2*d*x^7+1/5*(2*a*b*d+b^2*c)*x^5+1/3*(a^2*d+2*a*b*c)*x^3+a^2*c*x$

maxima [A] time = 1.32, size = 48, normalized size = 0.96

$$\frac{1}{7} b^2 d x^7 + \frac{1}{5} (b^2 c + 2 a b d) x^5 + a^2 c x + \frac{1}{3} (2 a b c + a^2 d) x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2*(d*x^2+c),x, algorithm="maxima")

[Out] $1/7*b^2*d*x^7 + 1/5*(b^2*c + 2*a*b*d)*x^5 + a^2*c*x + 1/3*(2*a*b*c + a^2*d)*x^3$

mupad [B] time = 0.05, size = 48, normalized size = 0.96

$$x^3 \left(\frac{d a^2}{3} + \frac{2 b c a}{3} \right) + x^5 \left(\frac{c b^2}{5} + \frac{2 a d b}{5} \right) + \frac{b^2 d x^7}{7} + a^2 c x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^2*(c + d*x^2),x)

[Out] $x^3*((a^2*d)/3 + (2*a*b*c)/3) + x^5*((b^2*c)/5 + (2*a*b*d)/5) + (b^2*d*x^7)/7 + a^2*c*x$

sympy [A] time = 0.07, size = 53, normalized size = 1.06

$$a^2 c x + \frac{b^2 d x^7}{7} + x^5 \left(\frac{2 a b d}{5} + \frac{b^2 c}{5} \right) + x^3 \left(\frac{a^2 d}{3} + \frac{2 a b c}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**2*(d*x**2+c),x)

[Out] $a**2*c*x + b**2*d*x**7/7 + x**5*(2*a*b*d/5 + b**2*c/5) + x**3*(a**2*d/3 + 2*a*b*c/3)$

$$3.11 \quad \int \frac{(a+bx^2)^2}{c+dx^2} dx$$

Optimal. Leaf size=63

$$\frac{(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}d^{5/2}} - \frac{bx(bc-2ad)}{d^2} + \frac{b^2x^3}{3d}$$

[Out] $-b*(-2*a*d+b*c)*x/d^2+1/3*b^2*x^3/d+(-a*d+b*c)^2*\arctan(x*d^{(1/2)}/c^{(1/2)})/d^{(5/2)}/c^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {390, 205}

$$-\frac{bx(bc-2ad)}{d^2} + \frac{(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}d^{5/2}} + \frac{b^2x^3}{3d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^2/(c + d*x^2), x]

[Out] $-((b*(b*c - 2*a*d)*x)/d^2) + (b^2*x^3)/(3*d) + ((b*c - a*d)^2*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(\text{Sqrt}[c]*d^{(5/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned} \int \frac{(a+bx^2)^2}{c+dx^2} dx &= \int \left(-\frac{b(bc-2ad)}{d^2} + \frac{b^2x^2}{d} + \frac{b^2c^2-2abcd+a^2d^2}{d^2(c+dx^2)} \right) dx \\ &= -\frac{b(bc-2ad)x}{d^2} + \frac{b^2x^3}{3d} + \frac{(bc-ad)^2 \int \frac{1}{c+dx^2} dx}{d^2} \\ &= -\frac{b(bc-2ad)x}{d^2} + \frac{b^2x^3}{3d} + \frac{(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}d^{5/2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 59, normalized size = 0.94

$$\frac{(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}d^{5/2}} + \frac{bx(6ad-3bc+bdx^2)}{3d^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^2/(c + d*x^2), x]

[Out] (b*x*(-3*b*c + 6*a*d + b*d*x^2))/(3*d^2) + ((b*c - a*d)^2*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(Sqrt[c]*d^(5/2))

fricas [A] time = 0.68, size = 179, normalized size = 2.84

$$\left[\frac{2 b^2 c d^2 x^3 - 3 (b^2 c^2 - 2 a b c d + a^2 d^2) \sqrt{-c d} \log \left(\frac{d x^2 - 2 \sqrt{-c d} x - c}{d x^2 + c} \right) - 6 (b^2 c^2 d - 2 a b c d^2) x}{6 c d^3}, \frac{b^2 c d^2 x^3 + 3 (b^2 c^2 - 2 a b c d)}{6 c d^3} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c), x, algorithm="fricas")

[Out] [1/6*(2*b^2*c*d^2*x^3 - 3*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(-c*d)*log((d*x^2 - 2*sqrt(-c*d)*x - c)/(d*x^2 + c)) - 6*(b^2*c^2*d - 2*a*b*c*d^2)*x)/(c*d^3), 1/3*(b^2*c*d^2*x^3 + 3*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(c*d)*arctan(sqrt(c*d)*x/c) - 3*(b^2*c^2*d - 2*a*b*c*d^2)*x)/(c*d^3)]

giac [A] time = 0.59, size = 72, normalized size = 1.14

$$\frac{(b^2 c^2 - 2 a b c d + a^2 d^2) \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d} d^2} + \frac{b^2 d^2 x^3 - 3 b^2 c d x + 6 a b d^2 x}{3 d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c), x, algorithm="giac")

[Out] (b^2*c^2 - 2*a*b*c*d + a^2*d^2)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*d^2) + 1/3*(b^2*d^2*x^3 - 3*b^2*c*d*x + 6*a*b*d^2*x)/d^3

maple [A] time = 0.00, size = 95, normalized size = 1.51

$$\frac{b^2 x^3}{3 d} + \frac{a^2 \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d}} - \frac{2 a b c \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d} d} + \frac{b^2 c^2 \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d} d^2} + \frac{2 a b x}{d} - \frac{b^2 c x}{d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^2/(d*x^2+c), x)

[Out] 1/3*b^2*x^3/d+2*b/d*a*x-b^2/d^2*x*c+1/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a^2-2/d/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a*b*c+1/d^2/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*b^2*c^2

maxima [A] time = 2.89, size = 68, normalized size = 1.08

$$\frac{(b^2 c^2 - 2 a b c d + a^2 d^2) \arctan \left(\frac{d x}{\sqrt{c d}} \right)}{\sqrt{c d} d^2} + \frac{b^2 d x^3 - 3 (b^2 c - 2 a b d) x}{3 d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c), x, algorithm="maxima")

[Out] (b^2*c^2 - 2*a*b*c*d + a^2*d^2)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*d^2) + 1/3*(b^2*d*x^3 - 3*(b^2*c - 2*a*b*d)*x)/d^2

mupad [B] time = 0.09, size = 90, normalized size = 1.43

$$\frac{b^2 x^3}{3 d} - x \left(\frac{b^2 c}{d^2} - \frac{2 a b}{d} \right) + \frac{\operatorname{atan} \left(\frac{\sqrt{d} x (a d - b c)^2}{\sqrt{c} (a^2 d^2 - 2 a b c d + b^2 c^2)} \right) (a d - b c)^2}{\sqrt{c} d^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^2/(c + d*x^2), x)`

[Out] $(b^2x^3)/(3d) - x((b^2c)/d^2 - (2ab)/d) + (\operatorname{atan}((d^{1/2})x(ad - bc)^2)/(c^{1/2}(a^2d^2 + b^2c^2 - 2abc*d))) * (ad - bc)^2 / (c^{1/2}d^{5/2})$

sympy [B] time = 0.42, size = 172, normalized size = 2.73

$$\frac{b^2x^3}{3d} + x\left(\frac{2ab}{d} - \frac{b^2c}{d^2}\right) - \frac{\sqrt{-\frac{1}{cd^5}}(ad - bc)^2 \log\left(-\frac{cd^2\sqrt{-\frac{1}{cd^5}}(ad - bc)^2}{a^2d^2 - 2abcd + b^2c^2} + x\right)}{2} + \frac{\sqrt{-\frac{1}{cd^5}}(ad - bc)^2 \log\left(\frac{cd^2\sqrt{-\frac{1}{cd^5}}(ad - bc)^2}{a^2d^2 - 2abcd + b^2c^2} + x\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**2/(d*x**2+c), x)`

[Out] $b^2x^3/(3d) + x(2ab/d - b^2c/d^2) - \sqrt{-1/(cd^5)}(ad - bc)^2 \log(-cd^2\sqrt{-1/(cd^5)}(ad - bc)^2/(a^2d^2 - 2abcd + b^2c^2) + x)/2 + \sqrt{-1/(cd^5)}(ad - bc)^2 \log(cd^2\sqrt{-1/(cd^5)}(ad - bc)^2/(a^2d^2 - 2abcd + b^2c^2) + x)/2$

$$3.12 \quad \int \frac{(a+bx^2)^2}{(c+dx^2)^2} dx$$

Optimal. Leaf size=82

$$-\frac{(bc-ad)(ad+3bc)\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{5/2}} + \frac{x(bc-ad)^2}{2cd^2(c+dx^2)} + \frac{b^2x}{d^2}$$

[Out] $b^2x/d^2 + 1/2*(-a*d+b*c)^2*x/c/d^2/(d*x^2+c) - 1/2*(-a*d+b*c)*(a*d+3*b*c)*\arctan(x*d^{(1/2)}/c^{(1/2)})/c^{(3/2)}/d^{(5/2)}$

Rubi [A] time = 0.10, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {390, 385, 205}

$$-\frac{(bc-ad)(ad+3bc)\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{5/2}} + \frac{x(bc-ad)^2}{2cd^2(c+dx^2)} + \frac{b^2x}{d^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^2/(c + d*x^2)^2, x]

[Out] $(b^2*x)/d^2 + ((b*c - a*d)^2*x)/(2*c*d^2*(c + d*x^2)) - ((b*c - a*d)*(3*b*c + a*d)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(2*c^{(3/2)}*d^{(5/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p+1))/(a*b*n*(p+1)), x] - Dist[(a*d - b*c*(n*(p+1) + 1))/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)^2}{(c + dx^2)^2} dx &= \int \left(\frac{b^2}{d^2} - \frac{b^2c^2 - a^2d^2 + 2bd(bc - ad)x^2}{d^2(c + dx^2)^2} \right) dx \\
&= \frac{b^2x}{d^2} - \frac{\int \frac{b^2c^2 - a^2d^2 + 2bd(bc - ad)x^2}{(c + dx^2)^2} dx}{d^2} \\
&= \frac{b^2x}{d^2} + \frac{(bc - ad)^2x}{2cd^2(c + dx^2)} - \frac{((bc - ad)(3bc + ad)) \int \frac{1}{c + dx^2} dx}{2cd^2} \\
&= \frac{b^2x}{d^2} + \frac{(bc - ad)^2x}{2cd^2(c + dx^2)} - \frac{(bc - ad)(3bc + ad) \tan^{-1} \left(\frac{\sqrt{d}x}{\sqrt{c}} \right)}{2c^{3/2}d^{5/2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 89, normalized size = 1.09

$$-\frac{(-a^2d^2 - 2abcd + 3b^2c^2) \tan^{-1} \left(\frac{\sqrt{d}x}{\sqrt{c}} \right)}{2c^{3/2}d^{5/2}} + \frac{x(bc - ad)^2}{2cd^2(c + dx^2)} + \frac{b^2x}{d^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^2/(c + d*x^2)^2,x]

[Out] (b^2*x)/d^2 + ((b*c - a*d)^2*x)/(2*c*d^2*(c + d*x^2)) - ((3*b^2*c^2 - 2*a*b*c*d - a^2*d^2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(2*c^(3/2)*d^(5/2))

fricas [B] time = 0.67, size = 302, normalized size = 3.68

$$\frac{4b^2c^2d^2x^3 + (3b^2c^3 - 2abc^2d - a^2cd^2 + (3b^2c^2d - 2abcd^2 - a^2d^3)x^2)\sqrt{-cd} \log\left(\frac{dx^2 - 2\sqrt{-cd}x - c}{dx^2 + c}\right) + 2(3b^2c^3d)}{4(c^2d^4x^2 + c^3d^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)^2,x, algorithm="fricas")

[Out] [1/4*(4*b^2*c^2*d^2*x^3 + (3*b^2*c^3 - 2*a*b*c^2*d - a^2*c*d^2 + (3*b^2*c^2*d - 2*a*b*c*d^2 - a^2*d^3)*x^2)*sqrt(-c*d)*log((d*x^2 - 2*sqrt(-c*d)*x - c)/(d*x^2 + c)) + 2*(3*b^2*c^3*d - 2*a*b*c^2*d^2 + a^2*c*d^3)*x)/(c^2*d^4*x^2 + c^3*d^3), 1/2*(2*b^2*c^2*d^2*x^3 - (3*b^2*c^3 - 2*a*b*c^2*d - a^2*c*d^2 + (3*b^2*c^2*d - 2*a*b*c*d^2 - a^2*d^3)*x^2)*sqrt(c*d)*arctan(sqrt(c*d)*x/c) + (3*b^2*c^3*d - 2*a*b*c^2*d^2 + a^2*c*d^3)*x)/(c^2*d^4*x^2 + c^3*d^3)]

giac [A] time = 0.57, size = 95, normalized size = 1.16

$$\frac{b^2x}{d^2} - \frac{(3b^2c^2 - 2abcd - a^2d^2) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}cd^2} + \frac{b^2c^2x - 2abcdx + a^2d^2x}{2(dx^2 + c)cd^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)^2,x, algorithm="giac")

[Out] b^2*x/d^2 - 1/2*(3*b^2*c^2 - 2*a*b*c*d - a^2*d^2)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*c*d^2) + 1/2*(b^2*c^2*x - 2*a*b*c*d*x + a^2*d^2*x)/((d*x^2 + c)*c*d^2)

maple [A] time = 0.01, size = 129, normalized size = 1.57

$$\frac{a^2 x}{2(d x^2 + c)c} + \frac{a^2 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}c} - \frac{abx}{(d x^2 + c)d} + \frac{ab \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{\sqrt{cd}d} + \frac{b^2 cx}{2(d x^2 + c)d^2} - \frac{3b^2 c \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}d^2} + \frac{b^2 x}{d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^2/(d*x^2+c)^2,x)

[Out] b^2*x/d^2+1/2/c*x/(d*x^2+c)*a^2-1/d*x/(d*x^2+c)*a*b+1/2/d^2*c*x/(d*x^2+c)*b^2+1/2/c/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a^2+1/d/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a*b-3/2/d^2*c/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*b^2

maxima [A] time = 2.84, size = 96, normalized size = 1.17

$$\frac{(b^2 c^2 - 2abcd + a^2 d^2)x}{2(cd^3 x^2 + c^2 d^2)} + \frac{b^2 x}{d^2} - \frac{(3b^2 c^2 - 2abcd - a^2 d^2) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}cd^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)^2,x, algorithm="maxima")

[Out] 1/2*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*x/(c*d^3*x^2 + c^2*d^2) + b^2*x/d^2 - 1/2*(3*b^2*c^2 - 2*a*b*c*d - a^2*d^2)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*c*d^2)

mupad [B] time = 5.02, size = 124, normalized size = 1.51

$$\frac{b^2 x}{d^2} + \frac{x(a^2 d^2 - 2abcd + b^2 c^2)}{2c(d^3 x^2 + c d^2)} + \frac{\operatorname{atan}\left(\frac{\sqrt{d} x(ad-bc)(ad+3bc)}{\sqrt{c}(a^2 d^2 + 2abcd - 3b^2 c^2)}\right)(ad-bc)(ad+3bc)}{2c^{3/2}d^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^2/(c + d*x^2)^2,x)

[Out] (b^2*x)/d^2 + (x*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d))/(2*c*(c*d^2 + d^3*x^2)) + (atan((d^(1/2)*x*(a*d - b*c)*(a*d + 3*b*c))/(c^(1/2)*(a^2*d^2 - 3*b^2*c^2 + 2*a*b*c*d)))*(a*d - b*c)*(a*d + 3*b*c))/(2*c^(3/2)*d^(5/2))

sympy [B] time = 0.70, size = 236, normalized size = 2.88

$$\frac{b^2 x}{d^2} + \frac{x(a^2 d^2 - 2abcd + b^2 c^2)}{2c^2 d^2 + 2cd^3 x^2} - \frac{\sqrt{-\frac{1}{c^3 d^5}}(ad-bc)(ad+3bc) \log\left(-\frac{c^2 d^2 \sqrt{-\frac{1}{c^3 d^5}}(ad-bc)(ad+3bc)}{a^2 d^2 + 2abcd - 3b^2 c^2} + x\right)}{4} + \frac{\sqrt{-\frac{1}{c^3 d^5}}(ad-bc)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**2/(d*x**2+c)**2,x)

[Out] b**2*x/d**2 + x*(a**2*d**2 - 2*a*b*c*d + b**2*c**2)/(2*c**2*d**2 + 2*c*d**3*x**2) - sqrt(-1/(c**3*d**5))*(a*d - b*c)*(a*d + 3*b*c)*log(-c**2*d**2*sqrt(-1/(c**3*d**5))*(a*d - b*c)*(a*d + 3*b*c)/(a**2*d**2 + 2*a*b*c*d - 3*b**2*c**2) + x)/4 + sqrt(-1/(c**3*d**5))*(a*d - b*c)*(a*d + 3*b*c)*log(c**2*d**2*sqrt(-1/(c**3*d**5))*(a*d - b*c)*(a*d + 3*b*c)/(a**2*d**2 + 2*a*b*c*d - 3*b**2*c**2) + x)/4

$$3.13 \quad \int \frac{(a+bx^2)^2}{(c+dx^2)^3} dx$$

Optimal. Leaf size=116

$$\frac{3x\left(\frac{a^2}{c^2} - \frac{b^2}{d^2}\right)}{8(c+dx^2)} + \frac{(3a^2d^2 + 2abcd + 3b^2c^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{5/2}} - \frac{x(a+bx^2)(bc-ad)}{4cd(c+dx^2)^2}$$

[Out] $-1/4*(-a*d+b*c)*x*(b*x^2+a)/c/d/(d*x^2+c)^2+3/8*(a^2/c^2-b^2/d^2)*x/(d*x^2+c)+1/8*(3*a^2*d^2+2*a*b*c*d+3*b^2*c^2)*\arctan(x*d^{(1/2)}/c^{(1/2)})/c^{(5/2)}/d^{(5/2)}$

Rubi [A] time = 0.07, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {413, 385, 205}

$$\frac{3x\left(\frac{a^2}{c^2} - \frac{b^2}{d^2}\right)}{8(c+dx^2)} + \frac{(3a^2d^2 + 2abcd + 3b^2c^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{5/2}} - \frac{x(a+bx^2)(bc-ad)}{4cd(c+dx^2)^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^2/(c + d*x^2)^3,x]

[Out] $-((b*c - a*d)*x*(a + b*x^2))/(4*c*d*(c + d*x^2)^2) + (3*(a^2/c^2 - b^2/d^2)*x)/(8*(c + d*x^2)) + ((3*b^2*c^2 + 2*a*b*c*d + 3*a^2*d^2)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(8*c^{(5/2)}*d^{(5/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rubi steps

$$\begin{aligned} \int \frac{(a+bx^2)^2}{(c+dx^2)^3} dx &= -\frac{(bc-ad)x(a+bx^2)}{4cd(c+dx^2)^2} + \frac{\int \frac{a(bc+3ad)+b(3bc+ad)x^2}{(c+dx^2)^2} dx}{4cd} \\ &= -\frac{(bc-ad)x(a+bx^2)}{4cd(c+dx^2)^2} + \frac{3\left(\frac{a^2}{c^2} - \frac{b^2}{d^2}\right)x}{8(c+dx^2)} + \frac{(3b^2c^2 + 2abcd + 3a^2d^2) \int \frac{1}{c+dx^2} dx}{8c^2d^2} \\ &= -\frac{(bc-ad)x(a+bx^2)}{4cd(c+dx^2)^2} + \frac{3\left(\frac{a^2}{c^2} - \frac{b^2}{d^2}\right)x}{8(c+dx^2)} + \frac{(3b^2c^2 + 2abcd + 3a^2d^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{5/2}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 121, normalized size = 1.04

$$\frac{x(a^2d^2(5c+3dx^2) - 2abcd(c-dx^2) - b^2c^2(3c+5dx^2))}{8c^2d^2(c+dx^2)^2} + \frac{(3a^2d^2 + 2abcd + 3b^2c^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^2/(c + d*x^2)^3,x]

[Out] (x*(-2*a*b*c*d*(c - d*x^2) + a^2*d^2*(5*c + 3*d*x^2) - b^2*c^2*(3*c + 5*d*x^2)))/(8*c^2*d^2*(c + d*x^2)^2) + (((3*b^2*c^2 + 2*a*b*c*d + 3*a^2*d^2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(8*c^(5/2)*d^(5/2))

fricas [B] time = 0.69, size = 449, normalized size = 3.87

$$\left[\frac{2(5b^2c^3d^2 - 2abc^2d^3 - 3a^2cd^4)x^3 + (3b^2c^4 + 2abc^3d + 3a^2c^2d^2 + (3b^2c^2d^2 + 2abcd^3 + 3a^2d^4))x^4 + 2(3b^2c^3d^2 - 2abcd^3 - 3a^2cd^4)}{16(c^3d^5x^4 + 2c^4d^4x^2 + c^5d^3)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [-1/16*(2*(5*b^2*c^3*d^2 - 2*a*b*c^2*d^3 - 3*a^2*c*d^4)*x^3 + (3*b^2*c^4 + 2*a*b*c^3*d + 3*a^2*c^2*d^2 + (3*b^2*c^2*d^2 + 2*a*b*c*d^3 + 3*a^2*d^4))*x^4 + 2*(3*b^2*c^3*d + 2*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(-c*d)*log((d*x^2 - 2*sqrt(-c*d)*x - c)/(d*x^2 + c)) + 2*(3*b^2*c^4*d + 2*a*b*c^3*d^2 - 5*a^2*c^2*d^3)*x)/(c^3*d^5*x^4 + 2*c^4*d^4*x^2 + c^5*d^3), -1/8*((5*b^2*c^3*d^2 - 2*a*b*c^2*d^3 - 3*a^2*c*d^4)*x^3 - (3*b^2*c^4 + 2*a*b*c^3*d + 3*a^2*c^2*d^2 + (3*b^2*c^2*d^2 + 2*a*b*c*d^3 + 3*a^2*d^4))*x^4 + 2*(3*b^2*c^3*d + 2*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(c*d)*arctan(sqrt(c*d)*x/c) + (3*b^2*c^4*d + 2*a*b*c^3*d^2 - 5*a^2*c^2*d^3)*x)/(c^3*d^5*x^4 + 2*c^4*d^4*x^2 + c^5*d^3)]

giac [A] time = 0.59, size = 126, normalized size = 1.09

$$\frac{(3b^2c^2 + 2abcd + 3a^2d^2) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd}c^2d^2} - \frac{5b^2c^2dx^3 - 2abcd^2x^3 - 3a^2d^3x^3 + 3b^2c^3x + 2abc^2dx - 5a^2cd^2x}{8(dx^2 + c)^2c^2d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)^3,x, algorithm="giac")

[Out] $\frac{1}{8}(3b^2c^2 + 2ab^2cd + 3a^2d^2) \arctan(dx/\sqrt{cd}) / (\sqrt{cd} * c^2d^2) - \frac{1}{8}(5b^2c^2d^2x^3 - 2ab^2cd^2x^3 - 3a^2d^3x^3 + 3b^2c^3x + 2ab^2c^2d^2x - 5a^2c^2d^2x) / ((dx^2 + c)^2 * c^2d^2)$

maple [A] time = 0.01, size = 147, normalized size = 1.27

$$\frac{3a^2 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd} c^2} + \frac{ab \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{4\sqrt{cd} cd} + \frac{3b^2 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd} d^2} + \frac{\frac{(3a^2d^2+2abcd-5b^2c^2)x^3}{8c^2d} + \frac{(5a^2d^2-2abcd-3b^2c^2)x}{8cd^2}}{(dx^2 + c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2+a)^2/(d*x^2+c)^3,x)`

[Out] $(1/8*(3a^2d^2+2ab^2cd-5b^2c^2)/c^2/d*x^3+1/8*(5a^2d^2-2ab^2cd-3b^2c^2)/d^2/c*x)/(d*x^2+c)^2+3/8/c^2/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*a^2+1/4/c/d/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*a*b+3/8/d^2/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*b^2$

maxima [A] time = 3.08, size = 138, normalized size = 1.19

$$\frac{(5b^2c^2d - 2abcd^2 - 3a^2d^3)x^3 + (3b^2c^3 + 2abc^2d - 5a^2cd^2)x}{8(c^2d^4x^4 + 2c^3d^3x^2 + c^4d^2)} + \frac{(3b^2c^2 + 2abcd + 3a^2d^2) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd} c^2d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+a)^2/(d*x^2+c)^3,x, algorithm="maxima")`

[Out] $-1/8*((5b^2c^2d - 2ab^2cd^2 - 3a^2d^3)*x^3 + (3b^2c^3 + 2ab^2cd^2 - 5a^2cd^2)*x)/(c^2d^4x^4 + 2c^3d^3x^2 + c^4d^2) + 1/8*(3b^2c^2 + 2ab^2cd + 3a^2d^2)*\arctan(dx/\sqrt{cd})/(\sqrt{cd}*c^2d^2)$

mupad [B] time = 5.03, size = 130, normalized size = 1.12

$$\frac{\operatorname{atan}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) (3a^2d^2 + 2abcd + 3b^2c^2)}{8c^{5/2}d^{5/2}} - \frac{x(-5a^2d^2+2abcd+3b^2c^2)}{8cd^2} - \frac{x^3(3a^2d^2+2abcd-5b^2c^2)}{8c^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^2/(c + d*x^2)^3,x)`

[Out] $(\operatorname{atan}((d^{(1/2)}*x)/c^{(1/2)})*(3a^2d^2 + 3b^2c^2 + 2ab^2cd))/(8*c^{(5/2)}*d^{(5/2)}) - ((x*(3b^2c^2 - 5a^2d^2 + 2ab^2cd))/(8*c*d^2) - (x^3*(3a^2d^2 - 5b^2c^2 + 2ab^2cd))/(8*c^2*d))/(c^2 + d^2*x^4 + 2*c*d*x^2)$

sympy [B] time = 0.98, size = 223, normalized size = 1.92

$$\frac{\sqrt{-\frac{1}{c^5d^5}} (3a^2d^2 + 2abcd + 3b^2c^2) \log\left(-c^3d^2\sqrt{-\frac{1}{c^5d^5}} + x\right)}{16} + \frac{\sqrt{-\frac{1}{c^5d^5}} (3a^2d^2 + 2abcd + 3b^2c^2) \log\left(c^3d^2\sqrt{-\frac{1}{c^5d^5}}\right)}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**2/(d*x**2+c)**3,x)`

[Out] $-\sqrt{-1/(c**5*d**5)}*(3a**2*d**2 + 2a*b*c*d + 3b**2*c**2)*\log(-c**3*d**2*\sqrt{-1/(c**5*d**5)} + x)/16 + \sqrt{-1/(c**5*d**5)}*(3a**2*d**2 + 2a*b*c*d + 3b**2*c**2)*\log(c**3*d**2*\sqrt{-1/(c**5*d**5)} + x)/16 + (x**3*(3a**2*d**3 + 2a*b*c*d**2 - 5b**2*c**2*d) + x*(5a**2*c*d**2 - 2a*b*c**2*d - 3b**2*c**3))/(8*c**4*d**2 + 16*c**3*d**3*x**2 + 8*c**2*d**4*x**4)$

3.14 $\int (a + bx^2)^3 (c + dx^2)^3 dx$

Optimal. Leaf size=154

$$a^3c^3x + \frac{1}{3}bdx^9(a^2d^2 + 3abcd + b^2c^2) + \frac{1}{7}x^7(ad+bc)(a^2d^2 + 8abcd + b^2c^2) + \frac{3}{5}acx^5(a^2d^2 + 3abcd + b^2c^2) + a^2c^2x^3(ad+bc) + \frac{1}{13}b^3d^3x^{13}$$

[Out] $a^3c^3x + a^2c^2(b^3d^3 + 9a^2bcd^2 + 9ab^2cd + 3a^3d^3)x^7/7 + (b^3d^3 + 9a^2bcd^2 + 9ab^2cd + 3a^3d^3)x^9/3 + (3b^2d^2(b^3c + a^3d)x^{11})/11 + (b^3d^3x^{13})/13$

Rubi [A] time = 0.10, antiderivative size = 154, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {373}

$$\frac{1}{3}bdx^9(a^2d^2 + 3abcd + b^2c^2) + \frac{1}{7}x^7(ad+bc)(a^2d^2 + 8abcd + b^2c^2) + \frac{3}{5}acx^5(a^2d^2 + 3abcd + b^2c^2) + a^2c^2x^3(ad+bc) + \frac{1}{13}b^3d^3x^{13}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^3*(c + d*x^2)^3,x]

[Out] $a^3c^3x + a^2c^2(b^3c + a^3d)x^7/7 + (3a^3c^3 + 9a^2c^2(b^3c + a^3d))x^9/3 + (3b^2d^2(b^3c + a^3d)x^{11})/11 + (b^3d^3x^{13})/13$

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\int (a + bx^2)^3 (c + dx^2)^3 dx = \int (a^3c^3 + 3a^2c^2(bc + ad)x^2 + 3ac(b^2c^2 + 3abcd + a^2d^2)x^4 + (bc + ad)(b^2c^2 + 8abcd + a^2d^2)x^6 + a^3c^3x^8 + \frac{1}{3}bdx^{10}(a^2d^2 + 3abcd + b^2c^2) + \frac{1}{7}x^{12}(ad+bc)(a^2d^2 + 8abcd + b^2c^2) + \frac{3}{5}acx^{14}(a^2d^2 + 3abcd + b^2c^2) + a^2c^2x^{16}(ad+bc) + \frac{1}{13}b^3d^3x^{18}) dx$$

Mathematica [A] time = 0.03, size = 161, normalized size = 1.05

$$a^3c^3x + \frac{1}{3}bdx^9(a^2d^2 + 3abcd + b^2c^2) + \frac{3}{5}acx^5(a^2d^2 + 3abcd + b^2c^2) + a^2c^2x^3(ad+bc) + \frac{1}{7}x^7(a^3d^3 + 9a^2bcd^2 + 9ab^2cd + 3a^3d^3)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^3*(c + d*x^2)^3,x]

[Out] $a^3c^3x + a^2c^2(b^3c + a^3d)x^7/7 + (3a^3c^3 + 9a^2c^2(b^3c + a^3d))x^9/3 + (3b^2d^2(b^3c + a^3d)x^{11})/11 + (b^3d^3x^{13})/13$

fricas [A] time = 0.69, size = 187, normalized size = 1.21

$$\frac{1}{13}x^{13}d^3b^3 + \frac{3}{11}x^{11}d^2cb^3 + \frac{3}{11}x^{11}d^3b^2a + \frac{1}{3}x^9dc^2b^3 + x^9d^2cb^2a + \frac{1}{3}x^9d^3ba^2 + \frac{1}{7}x^7c^3b^3 + \frac{9}{7}x^7dc^2b^2a + \frac{9}{7}x^7d^2cba^2 + \frac{1}{7}x^7d^3a^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c)^3,x, algorithm="fricas")

[Out] $\frac{1}{13}x^{13}d^3b^3 + \frac{3}{11}x^{11}d^2c^2b^3 + \frac{3}{11}x^{11}d^3b^2a + \frac{1}{3}x^9d^2c^2b^3 + x^9d^2c^2b^2a + \frac{1}{3}x^9d^3b^2a^2 + \frac{1}{7}x^7c^3b^3 + \frac{9}{7}x^7d^2c^2b^2a + \frac{9}{7}x^7d^2c^2b^2a^2 + \frac{1}{7}x^7d^3a^3 + \frac{3}{5}x^5c^3b^2a + \frac{9}{5}x^5d^2c^2b^2a^2 + \frac{3}{5}x^5d^2c^2a^3 + x^3c^3b^2a^2 + x^3d^2c^2a^3 + x^3c^3a^3$

giac [A] time = 0.57, size = 187, normalized size = 1.21

$$\frac{1}{13}b^3d^3x^{13} + \frac{3}{11}b^3cd^2x^{11} + \frac{3}{11}ab^2d^3x^{11} + \frac{1}{3}b^3c^2dx^9 + ab^2cd^2x^9 + \frac{1}{3}a^2bd^3x^9 + \frac{1}{7}b^3c^3x^7 + \frac{9}{7}ab^2c^2dx^7 + \frac{9}{7}a^2bcd^2x^7 + \frac{1}{7}a^3d^3x^7 + \frac{3}{5}a^2b^2c^3x^5 + \frac{9}{5}a^2b^2c^2d^2x^5 + \frac{3}{5}a^3c^3d^2x^5 + a^2b^2c^3x^3 + a^3c^2d^2x^3 + a^3c^3x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c)^3,x, algorithm="giac")

[Out] $\frac{1}{13}b^3d^3x^{13} + \frac{3}{11}b^3c^2d^2x^{11} + \frac{3}{11}a^2b^2d^3x^{11} + \frac{1}{3}b^3c^2d^2x^9 + a^2b^2c^2d^2x^9 + \frac{1}{3}a^2b^2d^3x^9 + \frac{1}{7}b^3c^3x^7 + \frac{9}{7}a^2b^2c^2d^2x^7 + \frac{9}{7}a^2b^2c^2d^2x^7 + \frac{1}{7}a^3d^3x^7 + \frac{3}{5}a^2b^2c^3x^5 + \frac{9}{5}a^2b^2c^2d^2x^5 + \frac{3}{5}a^3c^3d^2x^5 + a^2b^2c^3x^3 + a^3c^2d^2x^3 + a^3c^3x$

maple [A] time = 0.00, size = 177, normalized size = 1.15

$$\frac{b^3d^3x^{13}}{13} + \frac{(3ab^2d^3 + 3b^3cd^2)x^{11}}{11} + \frac{(3a^2bd^3 + 9ab^2cd^2 + 3b^3c^2d)x^9}{9} + a^3c^3x + \frac{(a^3d^3 + 9a^2bcd^2 + 9ab^2c^2d + b^3c^3)x^7}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^3*(d*x^2+c)^3,x)

[Out] $\frac{1}{13}b^3d^3x^{13} + \frac{1}{11}(3a^2b^2d^3 + 3b^3cd^2)x^{11} + \frac{1}{9}(3a^3d^3 + 9a^2b^2cd^2 + 9a^2b^2c^2d + b^3c^3)x^9 + \frac{1}{7}(a^3d^3 + 9a^2b^2cd^2 + 9a^2b^2c^2d + b^3c^3)x^7 + \frac{1}{5}(3a^3c^3d^2 + 9a^2b^2c^2d + 3a^2b^2c^3)x^5 + \frac{1}{3}(3a^3c^2d + 3a^2b^2c^3)x^3 + a^3c^3x$

maxima [A] time = 1.38, size = 167, normalized size = 1.08

$$\frac{1}{13}b^3d^3x^{13} + \frac{3}{11}(b^3cd^2 + ab^2d^3)x^{11} + \frac{1}{3}(b^3c^2d + 3ab^2cd^2 + a^2bd^3)x^9 + \frac{1}{7}(b^3c^3 + 9ab^2c^2d + 9a^2bcd^2 + a^3d^3)x^7$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c)^3,x, algorithm="maxima")

[Out] $\frac{1}{13}b^3d^3x^{13} + \frac{3}{11}(b^3c^2d^2 + a^2b^2d^3)x^{11} + \frac{1}{9}(b^3c^2d + 3a^2b^2cd^2 + a^2b^2d^3)x^9 + \frac{1}{7}(b^3c^3 + 9a^2b^2cd^2 + 9a^2b^2c^2d + a^3d^3)x^7 + a^3c^3x + \frac{3}{5}(a^2b^2c^3 + 3a^2b^2c^2d + a^3c^2d^2)x^5 + (a^2b^2c^3 + a^3c^2d)x^3$

mupad [B] time = 4.90, size = 152, normalized size = 0.99

$$x^7 \left(\frac{a^3d^3}{7} + \frac{9a^2bcd^2}{7} + \frac{9ab^2c^2d}{7} + \frac{b^3c^3}{7} \right) + a^3c^3x + \frac{b^3d^3x^{13}}{13} + \frac{3acx^5(a^2d^2 + 3abcd + b^2c^2)}{5} + \frac{bdx^9(a^2d^3 + 9a^2b^2cd^2 + 9a^2b^2c^2d + b^3c^3)}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^3*(c + d*x^2)^3,x)

[Out] $x^7 \left(\frac{a^3d^3}{7} + \frac{b^3c^3}{7} + \frac{9a^2b^2c^2d}{7} + \frac{9a^2b^2c^2d}{7} \right) + a^3c^3x + \frac{b^3d^3x^{13}}{13} + \frac{3a^2c^3x^5(a^2d^2 + b^2c^2 + 3a^2b^2cd)}{5} + \frac{b^3d^3x^9(a^2d^2 + b^2c^2 + 3a^2b^2cd)}{3} + a^2c^2x^3(a^2d + b^2c) + \frac{3b^2d^2x^{11}(a^2d + b^2c)}{11}$

sympy [A] time = 0.10, size = 189, normalized size = 1.23

$$a^3c^3x + \frac{b^3d^3x^{13}}{13} + x^{11} \left(\frac{3ab^2d^3}{11} + \frac{3b^3cd^2}{11} \right) + x^9 \left(\frac{a^2bd^3}{3} + ab^2cd^2 + \frac{b^3c^2d}{3} \right) + x^7 \left(\frac{a^3d^3}{7} + \frac{9a^2bcd^2}{7} + \frac{9ab^2c^2d}{7} + \frac{b^3c^3}{7} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**3*(d*x**2+c)**3,x)

[Out] a**3*c**3*x + b**3*d**3*x**13/13 + x**11*(3*a*b**2*d**3/11 + 3*b**3*c*d**2/11) + x**9*(a**2*b*d**3/3 + a*b**2*c*d**2 + b**3*c**2*d/3) + x**7*(a**3*d**3/7 + 9*a**2*b*c*d**2/7 + 9*a*b**2*c**2*d/7 + b**3*c**3/7) + x**5*(3*a**3*c*d**2/5 + 9*a**2*b*c**2*d/5 + 3*a*b**2*c**3/5) + x**3*(a**3*c**2*d + a**2*b*c**3)

3.15 $\int (a + bx^2)^3 (c + dx^2)^2 dx$

Optimal. Leaf size=122

$$a^3c^2x + \frac{1}{7}bx^7(3a^2d^2 + 6abcd + b^2c^2) + \frac{1}{5}ax^5(a^2d^2 + 6abcd + 3b^2c^2) + \frac{1}{3}a^2cx^3(2ad + 3bc) + \frac{1}{9}b^2dx^9(3ad + 2bc) + \frac{1}{11}b^3d^2x^{11}$$

[Out] a^3*c^2*x+1/3*a^2*c*(2*a*d+3*b*c)*x^3+1/5*a*(a^2*d^2+6*a*b*c*d+3*b^2*c^2)*x^5+1/7*b*(3*a^2*d^2+6*a*b*c*d+b^2*c^2)*x^7+1/9*b^2*d*(3*a*d+2*b*c)*x^9+1/11*b^3*d^2*x^11

Rubi [A] time = 0.07, antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {373}

$$\frac{1}{7}bx^7(3a^2d^2 + 6abcd + b^2c^2) + \frac{1}{5}ax^5(a^2d^2 + 6abcd + 3b^2c^2) + \frac{1}{3}a^2cx^3(2ad + 3bc) + a^3c^2x + \frac{1}{9}b^2dx^9(3ad + 2bc) + \frac{1}{11}b^3d^2x^{11}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^3*(c + d*x^2)^2,x]

[Out] a^3*c^2*x + (a^2*c*(3*b*c + 2*a*d)*x^3)/3 + (a*(3*b^2*c^2 + 6*a*b*c*d + a^2*d^2)*x^5)/5 + (b*(b^2*c^2 + 6*a*b*c*d + 3*a^2*d^2)*x^7)/7 + (b^2*d*(2*b*c + 3*a*d)*x^9)/9 + (b^3*d^2*x^11)/11

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^3 (c + dx^2)^2 dx &= \int (a^3c^2 + a^2c(3bc + 2ad)x^2 + a(3b^2c^2 + 6abcd + a^2d^2)x^4 + b(b^2c^2 + 6abcd + a^2d^2)x^6 + b^2d(2bc + 3ad)x^8 + b^3d^2x^{10}) dx \\ &= a^3c^2x + \frac{1}{3}a^2c(3bc + 2ad)x^3 + \frac{1}{5}a(3b^2c^2 + 6abcd + a^2d^2)x^5 + \frac{1}{7}b(b^2c^2 + 6abcd + a^2d^2)x^7 + \frac{1}{9}b^2d(2bc + 3ad)x^9 + \frac{1}{11}b^3d^2x^{11} \end{aligned}$$

Mathematica [A] time = 0.02, size = 122, normalized size = 1.00

$$a^3c^2x + \frac{1}{7}bx^7(3a^2d^2 + 6abcd + b^2c^2) + \frac{1}{5}ax^5(a^2d^2 + 6abcd + 3b^2c^2) + \frac{1}{3}a^2cx^3(2ad + 3bc) + \frac{1}{9}b^2dx^9(3ad + 2bc) + \frac{1}{11}b^3d^2x^{11}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^3*(c + d*x^2)^2,x]

[Out] a^3*c^2*x + (a^2*c*(3*b*c + 2*a*d)*x^3)/3 + (a*(3*b^2*c^2 + 6*a*b*c*d + a^2*d^2)*x^5)/5 + (b*(b^2*c^2 + 6*a*b*c*d + 3*a^2*d^2)*x^7)/7 + (b^2*d*(2*b*c + 3*a*d)*x^9)/9 + (b^3*d^2*x^11)/11

fricas [A] time = 0.47, size = 131, normalized size = 1.07

$$\frac{1}{11}x^{11}d^2b^3 + \frac{2}{9}x^9dcb^3 + \frac{1}{3}x^9d^2b^2a + \frac{1}{7}x^7c^2b^3 + \frac{6}{7}x^7dcb^2a + \frac{3}{7}x^7d^2ba^2 + \frac{3}{5}x^5c^2b^2a + \frac{6}{5}x^5dcb^2a + \frac{1}{5}x^5d^2a^3 + x^3c^2ba^2 + \frac{2}{3}x^3d^2ba^2 + \frac{1}{3}x^3d^2a^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c)^2,x, algorithm="fricas")

[Out] $\frac{1}{11}x^{11}d^2b^3 + \frac{2}{9}x^9d^2c^2b^3 + \frac{1}{3}x^9d^2b^2a + \frac{1}{7}x^7c^2b^3 + \frac{6}{7}x^7d^2c^2b^2a + \frac{3}{7}x^7d^2b^2a^2 + \frac{3}{5}x^5c^2b^2a + \frac{6}{5}x^5d^2c^2b^2a + \frac{1}{5}x^5d^2a^3 + x^3c^2b^2a^2 + \frac{2}{3}x^3d^2c^2a^3 + x^2c^2a^3$

giac [A] time = 0.59, size = 131, normalized size = 1.07

$$\frac{1}{11}b^3d^2x^{11} + \frac{2}{9}b^3cdx^9 + \frac{1}{3}ab^2d^2x^9 + \frac{1}{7}b^3c^2x^7 + \frac{6}{7}ab^2cdx^7 + \frac{3}{7}a^2bd^2x^7 + \frac{3}{5}ab^2c^2x^5 + \frac{6}{5}a^2bcdx^5 + \frac{1}{5}a^3d^2x^5 + a^2bc^2x^3 + \frac{2}{3}a^3c^2x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c)^2,x, algorithm="giac")

[Out] $\frac{1}{11}b^3d^2x^{11} + \frac{2}{9}b^3cdx^9 + \frac{1}{3}ab^2d^2x^9 + \frac{1}{7}b^3c^2x^7 + \frac{6}{7}ab^2cdx^7 + \frac{3}{7}a^2bd^2x^7 + \frac{3}{5}ab^2c^2x^5 + \frac{6}{5}a^2bcdx^5 + \frac{1}{5}a^3d^2x^5 + a^2bc^2x^3 + \frac{2}{3}a^3c^2x^3$

maple [A] time = 0.00, size = 125, normalized size = 1.02

$$\frac{b^3d^2x^{11}}{11} + \frac{(3ab^2d^2 + 2b^3cd)x^9}{9} + \frac{(3a^2bd^2 + 6ab^2cd + b^3c^2)x^7}{7} + a^3c^2x + \frac{(a^3d^2 + 6a^2bcd + 3ab^2c^2)x^5}{5} + \frac{(2a^3cd + 3a^3c^2)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^3*(d*x^2+c)^2,x)

[Out] $\frac{1}{11}b^3d^2x^{11} + \frac{1}{9}(3ab^2d^2 + 2b^3cd)x^9 + \frac{1}{7}(3a^2bd^2 + 6ab^2cd + b^3c^2)x^7 + \frac{1}{5}(a^3d^2 + 6a^2bcd + 3ab^2c^2)x^5 + \frac{1}{3}(2a^3cd + 3a^3c^2)x^3 + a^3c^2x$

maxima [A] time = 1.36, size = 124, normalized size = 1.02

$$\frac{1}{11}b^3d^2x^{11} + \frac{1}{9}(2b^3cd + 3ab^2d^2)x^9 + \frac{1}{7}(b^3c^2 + 6ab^2cd + 3a^2bd^2)x^7 + a^3c^2x + \frac{1}{5}(3ab^2c^2 + 6a^2bcd + a^3d^2)x^5 + \frac{1}{3}(3a^3cd + 3a^3c^2)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c)^2,x, algorithm="maxima")

[Out] $\frac{1}{11}b^3d^2x^{11} + \frac{1}{9}(2b^3cd + 3ab^2d^2)x^9 + \frac{1}{7}(b^3c^2 + 6ab^2cd + 3a^2bd^2)x^7 + \frac{1}{5}(3ab^2c^2 + 6a^2bcd + a^3d^2)x^5 + \frac{1}{3}(3a^3cd + 3a^3c^2)x^3 + a^3c^2x$

mupad [B] time = 4.87, size = 116, normalized size = 0.95

$$x^5 \left(\frac{a^3d^2}{5} + \frac{6a^2bcd}{5} + \frac{3ab^2c^2}{5} \right) + x^7 \left(\frac{3a^2bd^2}{7} + \frac{6ab^2cd}{7} + \frac{b^3c^2}{7} \right) + a^3c^2x + \frac{b^3d^2x^{11}}{11} + \frac{a^2cx^3(2ad + 3bc)}{3} + \frac{a^3c^2x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^3*(c + d*x^2)^2,x)

[Out] $x^5 \left(\frac{a^3d^2}{5} + \frac{3a^2bd^2}{5} + \frac{6a^2bcd}{5} \right) + x^7 \left(\frac{b^3c^2}{7} + \frac{3a^2bd^2}{7} + \frac{6a^2bcd}{7} \right) + a^3c^2x + \frac{b^3d^2x^{11}}{11} + \frac{a^2c^2x^3(2ad + 3bc)}{3} + \frac{b^2d^2x^9(3ad + 2bc)}{9}$

sympy [A] time = 0.09, size = 136, normalized size = 1.11

$$a^3c^2x + \frac{b^3d^2x^{11}}{11} + x^9 \left(\frac{ab^2d^2}{3} + \frac{2b^3cd}{9} \right) + x^7 \left(\frac{3a^2bd^2}{7} + \frac{6ab^2cd}{7} + \frac{b^3c^2}{7} \right) + x^5 \left(\frac{a^3d^2}{5} + \frac{6a^2bcd}{5} + \frac{3ab^2c^2}{5} \right) + x^3 \left(\frac{2a^3cd}{3} + \frac{a^3c^2}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**3*(d*x**2+c)**2,x)

[Out] $a^3c^2x + \frac{b^3d^2x^{11}}{11} + x^9 \left(\frac{ab^2d^2}{3} + \frac{2b^3cd}{9} \right) + x^7 \left(\frac{3a^2bd^2}{7} + \frac{6ab^2cd}{7} + \frac{b^3c^2}{7} \right) + x^5 \left(\frac{a^3d^2}{5} + \frac{6a^2bcd}{5} + \frac{3ab^2c^2}{5} \right) + x^3 \left(\frac{2a^3cd}{3} + \frac{a^3c^2}{3} \right)$

3.16 $\int (a + bx^2)^3 (c + dx^2) dx$

Optimal. Leaf size=70

$$a^3cx + \frac{1}{3}a^2x^3(ad + 3bc) + \frac{1}{7}b^2x^7(3ad + bc) + \frac{3}{5}abx^5(ad + bc) + \frac{1}{9}b^3dx^9$$

[Out] $a^3cx + \frac{1}{3}a^2x^3(ad + 3bc) + \frac{1}{7}b^2x^7(3ad + bc) + \frac{3}{5}abx^5(ad + bc) + \frac{1}{9}b^3dx^9$

Rubi [A] time = 0.04, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {373}

$$\frac{1}{3}a^2x^3(ad + 3bc) + a^3cx + \frac{1}{7}b^2x^7(3ad + bc) + \frac{3}{5}abx^5(ad + bc) + \frac{1}{9}b^3dx^9$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^3*(c + d*x^2), x]

[Out] $a^3cx + (a^2(3bc + ad)x^3)/3 + (3ab(bc + ad)x^5)/5 + (b^2(bc + 3ad)x^7)/7 + (b^3dx^9)/9$

Rule 373

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^3 (c + dx^2) dx &= \int (a^3c + a^2(3bc + ad)x^2 + 3ab(bc + ad)x^4 + b^2(bc + 3ad)x^6 + b^3dx^8) dx \\ &= a^3cx + \frac{1}{3}a^2(3bc + ad)x^3 + \frac{3}{5}ab(bc + ad)x^5 + \frac{1}{7}b^2(bc + 3ad)x^7 + \frac{1}{9}b^3dx^9 \end{aligned}$$

Mathematica [A] time = 0.01, size = 70, normalized size = 1.00

$$a^3cx + \frac{1}{3}a^2x^3(ad + 3bc) + \frac{1}{7}b^2x^7(3ad + bc) + \frac{3}{5}abx^5(ad + bc) + \frac{1}{9}b^3dx^9$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^3*(c + d*x^2), x]

[Out] $a^3cx + (a^2(3bc + ad)x^3)/3 + (3ab(bc + ad)x^5)/5 + (b^2(bc + 3ad)x^7)/7 + (b^3dx^9)/9$

fricas [A] time = 0.53, size = 73, normalized size = 1.04

$$\frac{1}{9}x^9db^3 + \frac{1}{7}x^7cb^3 + \frac{3}{7}x^7db^2a + \frac{3}{5}x^5cb^2a + \frac{3}{5}x^5dba^2 + x^3cba^2 + \frac{1}{3}x^3da^3 + xca^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c), x, algorithm="fricas")

[Out] $\frac{1}{9}x^9db^3 + \frac{1}{7}x^7cb^3 + \frac{3}{7}x^7db^2a + \frac{3}{5}x^5cb^2a + \frac{3}{5}x^5dba^2 + x^3cba^2 + \frac{1}{3}x^3da^3 + xca^3$

giac [A] time = 0.56, size = 73, normalized size = 1.04

$$\frac{1}{9}b^3dx^9 + \frac{1}{7}b^3cx^7 + \frac{3}{7}ab^2dx^7 + \frac{3}{5}ab^2cx^5 + \frac{3}{5}a^2bdx^5 + a^2bcx^3 + \frac{1}{3}a^3dx^3 + a^3cx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c),x, algorithm="giac")

[Out] 1/9*b^3*d*x^9 + 1/7*b^3*c*x^7 + 3/7*a*b^2*d*x^7 + 3/5*a*b^2*c*x^5 + 3/5*a^2*b*d*x^5 + a^2*b*c*x^3 + 1/3*a^3*d*x^3 + a^3*c*x

maple [A] time = 0.00, size = 73, normalized size = 1.04

$$\frac{b^3dx^9}{9} + \frac{(3ab^2d + b^3c)x^7}{7} + a^3cx + \frac{(3a^2bd + 3ab^2c)x^5}{5} + \frac{(a^3d + 3a^2bc)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^3*(d*x^2+c),x)

[Out] 1/9*b^3*d*x^9+1/7*(3*a*b^2*d+b^3*c)*x^7+1/5*(3*a^2*b*d+3*a*b^2*c)*x^5+1/3*(a^3*d+3*a^2*b*c)*x^3+c*a^3*x

maxima [A] time = 1.29, size = 70, normalized size = 1.00

$$\frac{1}{9}b^3dx^9 + \frac{1}{7}(b^3c + 3ab^2d)x^7 + \frac{3}{5}(ab^2c + a^2bd)x^5 + a^3cx + \frac{1}{3}(3a^2bc + a^3d)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3*(d*x^2+c),x, algorithm="maxima")

[Out] 1/9*b^3*d*x^9 + 1/7*(b^3*c + 3*a*b^2*d)*x^7 + 3/5*(a*b^2*c + a^2*b*d)*x^5 + a^3*c*x + 1/3*(3*a^2*b*c + a^3*d)*x^3

mupad [B] time = 0.03, size = 65, normalized size = 0.93

$$x^7 \left(\frac{cb^3}{7} + \frac{3adb^2}{7} \right) + x^3 \left(\frac{da^3}{3} + bca^2 \right) + \frac{b^3dx^9}{9} + a^3cx + \frac{3abx^5(ad+bc)}{5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^3*(c + d*x^2),x)

[Out] x^7*((b^3*c)/7 + (3*a*b^2*d)/7) + x^3*((a^3*d)/3 + a^2*b*c) + (b^3*d*x^9)/9 + a^3*c*x + (3*a*b*x^5*(a*d + b*c))/5

sympy [A] time = 0.08, size = 76, normalized size = 1.09

$$a^3cx + \frac{b^3dx^9}{9} + x^7 \left(\frac{3ab^2d}{7} + \frac{b^3c}{7} \right) + x^5 \left(\frac{3a^2bd}{5} + \frac{3ab^2c}{5} \right) + x^3 \left(\frac{a^3d}{3} + a^2bc \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**3*(d*x**2+c),x)

[Out] a**3*c*x + b**3*d*x**9/9 + x**7*(3*a*b**2*d/7 + b**3*c/7) + x**5*(3*a**2*b*d/5 + 3*a*b**2*c/5) + x**3*(a**3*d/3 + a**2*b*c)

$$3.17 \quad \int \frac{(a+bx^2)^3}{c+dx^2} dx$$

Optimal. Leaf size=98

$$\frac{bx(3a^2d^2 - 3abcd + b^2c^2)}{d^3} - \frac{b^2x^3(bc - 3ad)}{3d^2} - \frac{(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}d^{7/2}} + \frac{b^3x^5}{5d}$$

[Out] $b(3a^2d^2 - 3a^2b^2c^2 + 3ab^2cd + b^2c^2)x/d^3 - 1/3b^2(-3ad + bc)x^3/d^2 + 1/5b^3x^5/d - (ad + bc)^3 \arctan(x\sqrt{d}/\sqrt{c})/d^{7/2}/\sqrt{c}$

Rubi [A] time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {390, 205}

$$\frac{bx(3a^2d^2 - 3abcd + b^2c^2)}{d^3} - \frac{b^2x^3(bc - 3ad)}{3d^2} - \frac{(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}d^{7/2}} + \frac{b^3x^5}{5d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^3/(c + d*x^2), x]

[Out] $(b(b^2c^2 - 3a^2b^2c^2 + 3a^2d^2)x)/d^3 - (b^2(b^2c^2 - 3a^2d^2)x^3)/(3d^2) + (b^3x^5)/(5d) - ((b^2c^2 - 3a^2d^2)^3 \text{ArcTan}[\sqrt{d}x/\sqrt{c}])/(\sqrt{c}d^{7/2})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned} \int \frac{(a+bx^2)^3}{c+dx^2} dx &= \int \left(\frac{b(b^2c^2 - 3abcd + 3a^2d^2)}{d^3} - \frac{b^2(bc - 3ad)x^2}{d^2} + \frac{b^3x^4}{d} + \frac{-b^3c^3 + 3ab^2c^2d - 3a^2bcd^2 + a^3d^3}{d^3(c+dx^2)} \right) dx \\ &= \frac{b(b^2c^2 - 3abcd + 3a^2d^2)x}{d^3} - \frac{b^2(bc - 3ad)x^3}{3d^2} + \frac{b^3x^5}{5d} - \frac{(bc - ad)^3 \int \frac{1}{c+dx^2} dx}{d^3} \\ &= \frac{b(b^2c^2 - 3abcd + 3a^2d^2)x}{d^3} - \frac{b^2(bc - 3ad)x^3}{3d^2} + \frac{b^3x^5}{5d} - \frac{(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}d^{7/2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 93, normalized size = 0.95

$$\frac{bx(45a^2d^2 + 15abd(dx^2 - 3c) + b^2(15c^2 - 5cdx^2 + 3d^2x^4))}{15d^3} - \frac{(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}d^{7/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^3/(c + d*x^2), x]

[Out] (b*x*(45*a^2*d^2 + 15*a*b*d*(-3*c + d*x^2) + b^2*(15*c^2 - 5*c*d*x^2 + 3*d^2*x^4)))/(15*d^3) - ((b*c - a*d)^3*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(Sqrt[c]*d^(7/2))

fricas [A] time = 0.54, size = 290, normalized size = 2.96

$$\frac{6b^3cd^3x^5 - 10(b^3c^2d^2 - 3ab^2cd^3)x^3 + 15(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3)\sqrt{-cd} \log\left(\frac{dx^2 - 2\sqrt{-cd}x - c}{dx^2 + c}\right) + 30(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3)\sqrt{cd} \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{30cd^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c), x, algorithm="fricas")

[Out] [1/30*(6*b^3*c*d^3*x^5 - 10*(b^3*c^2*d^2 - 3*a*b^2*c*d^3)*x^3 + 15*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(-c*d)*log((d*x^2 - 2*sqrt(-c*d)*x - c)/(d*x^2 + c)) + 30*(b^3*c^3*d - 3*a*b^2*c^2*d^2 + 3*a^2*b*c*d^3)*x)/(c*d^4), 1/15*(3*b^3*c*d^3*x^5 - 5*(b^3*c^2*d^2 - 3*a*b^2*c*d^3)*x^3 - 15*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(c*d)*arctan(sqrt(c*d)*x/c) + 15*(b^3*c^3*d - 3*a*b^2*c^2*d^2 + 3*a^2*b*c*d^3)*x)/(c*d^4)]

giac [A] time = 0.58, size = 130, normalized size = 1.33

$$\frac{(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{\sqrt{cd}d^3} + \frac{3b^3d^4x^5 - 5b^3cd^3x^3 + 15ab^2d^4x^3 + 15b^3c^2d^2x - 45ab^2cd^3x}{15d^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c), x, algorithm="giac")

[Out] -(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*arctan(d*x/sqrt(c*d))/(sqrt(c*d)*d^3) + 1/15*(3*b^3*d^4*x^5 - 5*b^3*c*d^3*x^3 + 15*a*b^2*d^4*x^3 + 15*b^3*c^2*d^2*x - 45*a*b^2*c*d^3*x + 45*a^2*b*d^4*x)/d^5

maple [A] time = 0.00, size = 161, normalized size = 1.64

$$\frac{b^3x^5}{5d} + \frac{ab^2x^3}{d} - \frac{b^3cx^3}{3d^2} + \frac{a^3 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{\sqrt{cd}} - \frac{3a^2bc \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{\sqrt{cd}d} + \frac{3ab^2c^2 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{\sqrt{cd}d^2} - \frac{b^3c^3 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{\sqrt{cd}d^3} + \frac{3a^2bx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^3/(d*x^2+c), x)

[Out] 1/5*b^3*x^5/d + b^2/d*x^3*a - 1/3*b^3/d^2*x^3*c + 3*b/d*a^2*x - 3*b^2/d^2*a*c*x + b^3/d^3*c^2*x + 1/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a^3 - 3/d/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a^2*b*c + 3/d^2/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a*b^2*c^2 - 1/d^3/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*b^3*c^3

maxima [A] time = 3.02, size = 122, normalized size = 1.24

$$\frac{(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{\sqrt{cd}d^3} + \frac{3b^3d^2x^5 - 5(b^3cd - 3ab^2d^2)x^3 + 15(b^3c^2 - 3ab^2cd + 3a^2bd^2)x - 45ab^2cd^3x}{15d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c), x, algorithm="maxima")

[Out] $-(b^3c^3 - 3ab^2c^2d + 3a^2b^2cd^2 - a^3d^3) \arctan(dx/\sqrt{cd}) / (\sqrt{cd}d^3) + 1/15(3b^3d^2x^5 - 5(b^3cd - 3ab^2d^2)x^3 + 15(b^3c^2 - 3ab^2cd + 3a^2bd^2)x) / d^3$

mupad [B] time = 4.87, size = 145, normalized size = 1.48

$$x^3 \left(\frac{ab^2}{d} - \frac{b^3c}{3d^2} \right) + x \left(\frac{3a^2b}{d} - \frac{c \left(\frac{3ab^2}{d} - \frac{b^3c}{d^2} \right)}{d} \right) + \frac{b^3x^5}{5d} + \frac{\operatorname{atan} \left(\frac{\sqrt{d} x (ad-bc)^3}{\sqrt{c} (a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3)} \right) (ad-bc)^3}{\sqrt{c} d^{7/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^3/(c + d*x^2), x)`

[Out] $x^3((ab^2)/d - (b^3c)/(3d^2)) + x((3a^2b)/d - (c((3ab^2)/d - (b^3c)/d^2))/d) + (b^3x^5)/(5d) + (\operatorname{atan}((d^{1/2})x*(ad - bc)^3)/(c^{1/2}(a^3d^3 - b^3c^3 + 3ab^2cd^2 - 3a^2b^2cd^2)))*(ad - bc)^3/(c^{1/2})d^{7/2})$

sympy [B] time = 0.57, size = 238, normalized size = 2.43

$$\frac{b^3x^5}{5d} + x^3 \left(\frac{ab^2}{d} - \frac{b^3c}{3d^2} \right) + x \left(\frac{3a^2b}{d} - \frac{3ab^2c}{d^2} + \frac{b^3c^2}{d^3} \right) - \frac{\sqrt{-\frac{1}{cd^7}} (ad - bc)^3 \log \left(-\frac{cd^3 \sqrt{-\frac{1}{cd^7}} (ad-bc)^3}{a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3} + x \right)}{2} + \frac{\sqrt{-\frac{1}{cd^7}} (ad - bc)^3 \log \left(-\frac{cd^3 \sqrt{-\frac{1}{cd^7}} (ad-bc)^3}{a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3} + x \right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**3/(d*x**2+c), x)`

[Out] $b^3x^5/(5d) + x^3(a^2b/d - b^3c/(3d^2)) + x(3a^2b/d - 3ab^2c/d^2 + b^3c^2/d^3) - \sqrt{-1/(cd^7)}(ad - bc)^3 \log(-cd^3 \sqrt{-1/(cd^7)}(ad - bc)^3 / (a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3) + x) / 2 + \sqrt{-1/(cd^7)}(ad - bc)^3 \log(cd^3 \sqrt{-1/(cd^7)}(ad - bc)^3 / (a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3) + x) / 2$

$$3.18 \quad \int \frac{(a+bx^2)^3}{(c+dx^2)^2} dx$$

Optimal. Leaf size=107

$$-\frac{b^2x(2bc-3ad)}{d^3} + \frac{(ad+5bc)(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{7/2}} - \frac{x(bc-ad)^3}{2cd^3(c+dx^2)} + \frac{b^3x^3}{3d^2}$$

[Out] $-b^2*(-3*a*d+2*b*c)*x/d^3+1/3*b^3*x^3/d^2-1/2*(-a*d+b*c)^3*x/c/d^3/(d*x^2+c)+1/2*(-a*d+b*c)^2*(a*d+5*b*c)*\arctan(x*d^{(1/2)}/c^{(1/2)})/c^{(3/2)}/d^{(7/2)}$

Rubi [A] time = 0.10, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {390, 385, 205}

$$-\frac{b^2x(2bc-3ad)}{d^3} + \frac{(ad+5bc)(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{7/2}} - \frac{x(bc-ad)^3}{2cd^3(c+dx^2)} + \frac{b^3x^3}{3d^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^3/(c + d*x^2)^2,x]

[Out] $-((b^2*(2*b*c-3*a*d)*x)/d^3) + (b^3*x^3)/(3*d^2) - ((b*c-a*d)^3*x)/(2*c*d^3*(c+d*x^2)) + ((b*c-a*d)^2*(5*b*c+a*d)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(2*c^{(3/2)}*d^{(7/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c-a*d)*x*(a+b*x^n)^(p+1))/(a*b*n*(p+1)), x] - Dist[(a*d-b*c*(n*(p+1)+1))/(a*b*n*(p+1)), Int[(a+b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c-a*d, 0] && (LtQ[p, -1] || ILtQ[1/n+p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a+b*x^n)^p, (c+d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c-a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)^3}{(c + dx^2)^2} dx &= \int \left(-\frac{b^2(2bc - 3ad)}{d^3} + \frac{b^3x^2}{d^2} + \frac{(bc - ad)^2(2bc + ad) + 3bd(bc - ad)^2x^2}{d^3(c + dx^2)^2} \right) dx \\
&= -\frac{b^2(2bc - 3ad)x}{d^3} + \frac{b^3x^3}{3d^2} + \frac{\int \frac{(bc - ad)^2(2bc + ad) + 3bd(bc - ad)^2x^2}{(c + dx^2)^2} dx}{d^3} \\
&= -\frac{b^2(2bc - 3ad)x}{d^3} + \frac{b^3x^3}{3d^2} - \frac{(bc - ad)^3x}{2cd^3(c + dx^2)} + \frac{((bc - ad)^2(5bc + ad)) \int \frac{1}{c + dx^2} dx}{2cd^3} \\
&= -\frac{b^2(2bc - 3ad)x}{d^3} + \frac{b^3x^3}{3d^2} - \frac{(bc - ad)^3x}{2cd^3(c + dx^2)} + \frac{(bc - ad)^2(5bc + ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{7/2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 107, normalized size = 1.00

$$-\frac{b^2x(2bc - 3ad)}{d^3} + \frac{(ad + 5bc)(bc - ad)^2 \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}d^{7/2}} - \frac{x(bc - ad)^3}{2cd^3(c + dx^2)} + \frac{b^3x^3}{3d^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^3/(c + d*x^2)^2,x]

[Out] -((b^2*(2*b*c - 3*a*d)*x)/d^3) + (b^3*x^3)/(3*d^2) - ((b*c - a*d)^3*x)/(2*c*d^3*(c + d*x^2)) + ((b*c - a*d)^2*(5*b*c + a*d)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(2*c^(3/2)*d^(7/2))

fricas [B] time = 0.62, size = 444, normalized size = 4.15

$$\frac{4b^3c^2d^3x^5 - 4(5b^3c^3d^2 - 9ab^2c^2d^3)x^3 - 3(5b^3c^4 - 9ab^2c^3d + 3a^2bc^2d^2 + a^3cd^3 + (5b^3c^3d - 9ab^2c^2d^2 + 3a^2bc^2d^2 - 9a^3cd^3)x^2) \sqrt{-cd} \log((dx^2 - 2\sqrt{-cd})x - c)/(dx^2 + c) - 6(5b^3c^4d - 9a^2b^2c^3d^2 + 3a^2b^2c^2d^3 - a^3cd^4)x)/(c^2d^5x^2 + c^3d^4), 1/6*(2b^3c^2d^3x^5 - 2(5b^3c^3d^2 - 9a^2b^2c^2d^3)x^3 + 3(5b^3c^4 - 9a^2b^2c^3d + 3a^2b^2c^2d^3 - a^3cd^4)x^2) \sqrt{cd} \arctan(\sqrt{cd}x/c) - 3(5b^3c^4d - 9a^2b^2c^3d^2 + 3a^2b^2c^2d^3 - a^3cd^4)x)/(c^2d^5x^2 + c^3d^4)}{12(c^2d^5x^2 + c^3d^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c)^2,x, algorithm="fricas")

[Out] [1/12*(4*b^3*c^2*d^3*x^5 - 4*(5*b^3*c^3*d^2 - 9*a*b^2*c^2*d^3)*x^3 - 3*(5*b^3*c^4 - 9*a*b^2*c^3*d + 3*a^2*b*c^2*d^2 + a^3*c*d^3 + (5*b^3*c^3*d - 9*a*b^2*c^2*d^2 + 3*a^2*b*c^2*d^2 - 9*a^3*c*d^3)*x^2)*sqrt(-c*d)*log((d*x^2 - 2*sqrt(-c*d)*x - c)/(d*x^2 + c)) - 6*(5*b^3*c^4*d - 9*a*b^2*c^3*d^2 + 3*a^2*b*c^2*d^3 - a^3*c*d^4)*x)/(c^2*d^5*x^2 + c^3*d^4), 1/6*(2*b^3*c^2*d^3*x^5 - 2*(5*b^3*c^3*d^2 - 9*a*b^2*c^2*d^3)*x^3 + 3*(5*b^3*c^4 - 9*a*b^2*c^3*d + 3*a^2*b*c^2*d^2 + a^3*c*d^3 + (5*b^3*c^3*d - 9*a*b^2*c^2*d^2 + 3*a^2*b*c^2*d^2 - 9*a^3*c*d^3)*x^2)*sqrt(c*d)*arctan(sqrt(c*d)*x/c) - 3*(5*b^3*c^4*d - 9*a*b^2*c^3*d^2 + 3*a^2*b*c^2*d^3 - a^3*c*d^4)*x)/(c^2*d^5*x^2 + c^3*d^4)]

giac [A] time = 0.58, size = 152, normalized size = 1.42

$$\frac{(5b^3c^3 - 9ab^2c^2d + 3a^2bcd^2 + a^3d^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}cd^3} - \frac{b^3c^3x - 3ab^2c^2dx + 3a^2bcd^2x - a^3d^3x}{2(dx^2 + c)cd^3} + \frac{b^3d^4x^3 - 6b^3cd^3}{3d^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c)^2,x, algorithm="giac")

[Out] $\frac{1}{2}*(5*b^3*c^3 - 9*a*b^2*c^2*d + 3*a^2*b*c*d^2 + a^3*d^3)*\arctan(d*x/\sqrt{c*d})/(\sqrt{c*d}*c*d^3) - \frac{1}{2}*(b^3*c^3*x - 3*a*b^2*c^2*d*x + 3*a^2*b*c*d^2*x - a^3*d^3*x)/((d*x^2 + c)*c*d^3) + \frac{1}{3}*(b^3*d^4*x^3 - 6*b^3*c*d^3*x + 9*a*b^2*d^4*x)/d^6$

maple [B] time = 0.01, size = 205, normalized size = 1.92

$$\frac{b^3x^3}{3d^2} + \frac{a^3x}{2(dx^2+c)c} + \frac{a^3 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}c} - \frac{3a^2bx}{2(dx^2+c)d} + \frac{3a^2b \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}d} + \frac{3ab^2cx}{2(dx^2+c)d^2} - \frac{9ab^2c \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2+a)^3/(d*x^2+c)^2,x)`

[Out] $\frac{1}{3}b^3x^3/d^2 + 3b^2/d^2ax - 2b^3/d^3x^2c + 1/2/cx/(d*x^2+c)a^3 - 3/2/dx/(d*x^2+c)a^2b + 3/2/d^2cx/(d*x^2+c)a*b^2 - 1/2/d^3c^2x/(d*x^2+c)b^3 + 1/2/c/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*a^3 + 3/2/d/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*a^2b - 9/2/d^2c/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*a*b^2 + 5/2/d^3c^2/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*b^3$

maxima [A] time = 2.99, size = 147, normalized size = 1.37

$$\frac{(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3)x}{2(cd^4x^2 + c^2d^3)} + \frac{b^3dx^3 - 3(2b^3c - 3ab^2d)x}{3d^3} + \frac{(5b^3c^3 - 9ab^2c^2d + 3a^2bcd^2 + a^3d^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2\sqrt{cd}cd^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+a)^3/(d*x^2+c)^2,x, algorithm="maxima")`

[Out] $-\frac{1}{2}*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*x/(c*d^4*x^2 + c^2*d^3) + \frac{1}{3}*(b^3*d*x^3 - 3*(2*b^3*c - 3*a*b^2*d)*x)/d^3 + \frac{1}{2}*(5*b^3*c^3 - 9*a*b^2*c^2*d + 3*a^2*b*c*d^2 + a^3*d^3)*\arctan(d*x/\sqrt{c*d})/(\sqrt{c*d}*c*d^3)$

mupad [B] time = 0.10, size = 181, normalized size = 1.69

$$x \left(\frac{3ab^2}{d^2} - \frac{2b^3c}{d^3} \right) + \frac{b^3x^3}{3d^2} + \frac{x(a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3)}{2c(d^4x^2 + cd^3)} + \frac{\operatorname{atan}\left(\frac{\sqrt{d}x(ad-bc)^2(ad+5bc)}{\sqrt{c}(a^3d^3+3a^2bcd^2-9ab^2c^2d+5b^3c^3)}\right)}{2c^{3/2}d^{7/2}} (a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^3/(c + d*x^2)^2,x)`

[Out] $x*((3*a*b^2)/d^2 - (2*b^3*c)/d^3) + (b^3*x^3)/(3*d^2) + (x*(a^3*d^3 - b^3*c^3 + 3*a*b^2*c^2*d - 3*a^2*b*c*d^2))/(2*c*(c*d^3 + d^4*x^2)) + (\operatorname{atan}((d^{1/2}*x*(a*d - b*c)^2*(a*d + 5*b*c))/(c^{1/2)*(a^3*d^3 + 5*b^3*c^3 - 9*a*b^2*c^2*d + 3*a^2*b*c*d^2)))*(a*d - b*c)^2*(a*d + 5*b*c))/(2*c^{3/2}*d^{7/2})$

sympy [B] time = 1.05, size = 314, normalized size = 2.93

$$\frac{b^3x^3}{3d^2} + x \left(\frac{3ab^2}{d^2} - \frac{2b^3c}{d^3} \right) + \frac{x(a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3)}{2c^2d^3 + 2cd^4x^2} - \frac{\sqrt{-\frac{1}{c^3d^7}}(ad-bc)^2(ad+5bc) \log\left(-\frac{c^2d^3\sqrt{-\frac{1}{c^3d^7}}(ad-bc)^2(ad+5bc)}{a^3d^3+3a^2bcd^2-9ab^2c^2d+5b^3c^3}\right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**3/(d*x**2+c)**2,x)`


```
[Out] b**3*x**3/(3*d**2) + x*(3*a*b**2/d**2 - 2*b**3*c/d**3) + x*(a**3*d**3 - 3*a
**2*b*c*d**2 + 3*a*b**2*c**2*d - b**3*c**3)/(2*c**2*d**3 + 2*c*d**4*x**2) -
sqrt(-1/(c**3*d**7))*(a*d - b*c)**2*(a*d + 5*b*c)*log(-c**2*d**3*sqrt(-1/(
c**3*d**7))*(a*d - b*c)**2*(a*d + 5*b*c)/(a**3*d**3 + 3*a**2*b*c*d**2 - 9*a
*b**2*c**2*d + 5*b**3*c**3) + x)/4 + sqrt(-1/(c**3*d**7))*(a*d - b*c)**2*(a
*d + 5*b*c)*log(c**2*d**3*sqrt(-1/(c**3*d**7))*(a*d - b*c)**2*(a*d + 5*b*c)
/(a**3*d**3 + 3*a**2*b*c*d**2 - 9*a*b**2*c**2*d + 5*b**3*c**3) + x)/4
```

$$3.19 \quad \int \frac{(a+bx^2)^3}{(c+dx^2)^3} dx$$

Optimal. Leaf size=130

$$\frac{3(bc-ad)\left((ad+bc)^2+4b^2c^2\right)\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{7/2}} + \frac{3x(bc-ad)^2(ad+3bc)}{8c^2d^3(c+dx^2)} - \frac{x(bc-ad)^3}{4cd^3(c+dx^2)^2} + \frac{b^3x}{d^3}$$

[Out] $b^3*x/d^3-1/4*(-a*d+b*c)^3*x/c/d^3/(d*x^2+c)^2+3/8*(-a*d+b*c)^2*(a*d+3*b*c)*x/c^2/d^3/(d*x^2+c)-3/8*(-a*d+b*c)*(4*b^2*c^2+(a*d+b*c)^2)*\arctan(x*d^{(1/2)})/c^{(1/2)}/c^{(5/2)}/d^{(7/2)}$

Rubi [A] time = 0.16, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {390, 1157, 385, 205}

$$\frac{3(bc-ad)\left((ad+bc)^2+4b^2c^2\right)\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{7/2}} + \frac{3x(bc-ad)^2(ad+3bc)}{8c^2d^3(c+dx^2)} - \frac{x(bc-ad)^3}{4cd^3(c+dx^2)^2} + \frac{b^3x}{d^3}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^3/(c + d*x^2)^3,x]

[Out] $(b^3*x)/d^3 - ((b*c - a*d)^3*x)/(4*c*d^3*(c + d*x^2)^2) + (3*(b*c - a*d)^2*(3*b*c + a*d)*x)/(8*c^2*d^3*(c + d*x^2)) - (3*(b*c - a*d)*(4*b^2*c^2 + (b*c + a*d)^2)*\text{ArcTan}[\text{Sqrt}[d]*x/\text{Sqrt}[c]])/(8*c^{(5/2)}*d^{(7/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p+1))/(a*b*n*(p+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 1157

Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x, 0]}, -Simp[(R*x*(d + e*x^2)^(q+1))/(2*d*(q+1)), x] + Dist[1/(2*d*(q+1)), Int[(d + e*x^2)^(q+1)*ExpandToSum[2*d*(q+1)*Qx + R*(2*q+3), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)^3}{(c + dx^2)^3} dx &= \int \left(\frac{b^3}{d^3} - \frac{b^3c^3 - a^3d^3 + 3bd(bc - ad)(bc + ad)x^2 + 3b^2d^2(bc - ad)x^4}{d^3(c + dx^2)^3} \right) dx \\
&= \frac{b^3x}{d^3} - \frac{\int \frac{b^3c^3 - a^3d^3 + 3bd(bc - ad)(bc + ad)x^2 + 3b^2d^2(bc - ad)x^4}{(c + dx^2)^3} dx}{d^3} \\
&= \frac{b^3x}{d^3} - \frac{(bc - ad)^3x}{4cd^3(c + dx^2)^2} + \frac{\int \frac{-3(bc - ad)(bc + ad)^2 - 12b^2cd(bc - ad)x^2}{(c + dx^2)^2} dx}{4cd^3} \\
&= \frac{b^3x}{d^3} - \frac{(bc - ad)^3x}{4cd^3(c + dx^2)^2} + \frac{3(bc - ad)^2(3bc + ad)x}{8c^2d^3(c + dx^2)} - \frac{(3(bc - ad)(4b^2c^2 + (bc + ad)^2)) \int \frac{1}{c + dx^2}}{8c^2d^3} \\
&= \frac{b^3x}{d^3} - \frac{(bc - ad)^3x}{4cd^3(c + dx^2)^2} + \frac{3(bc - ad)^2(3bc + ad)x}{8c^2d^3(c + dx^2)} - \frac{3(bc - ad)(4b^2c^2 + (bc + ad)^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{7/2}}
\end{aligned}$$

Mathematica [A] time = 0.08, size = 141, normalized size = 1.08

$$\frac{3(-a^3d^3 - a^2bcd^2 - 3ab^2c^2d + 5b^3c^3) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}d^{7/2}} + \frac{3x(bc - ad)^2(ad + 3bc)}{8c^2d^3(c + dx^2)} - \frac{x(bc - ad)^3}{4cd^3(c + dx^2)^2} + \frac{b^3x}{d^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^3/(c + d*x^2)^3,x]

[Out] (b^3*x)/d^3 - ((b*c - a*d)^3*x)/(4*c*d^3*(c + d*x^2)^2) + (3*(b*c - a*d)^2*(3*b*c + a*d)*x)/(8*c^2*d^3*(c + d*x^2)) - (3*(5*b^3*c^3 - 3*a*b^2*c^2*d - a^2*b*c*d^2 - a^3*d^3)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(8*c^(5/2)*d^(7/2))

fricas [B] time = 0.69, size = 618, normalized size = 4.75

$$\left[\frac{16b^3c^3d^3x^5 + 2(25b^3c^4d^2 - 15ab^2c^3d^3 + 3a^2bc^2d^4 + 3a^3cd^5)x^3 + 3(5b^3c^5 - 3ab^2c^4d - a^2bc^3d^2 - a^3c^2d^3)}{8c^{5/2}d^{7/2}} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [1/16*(16*b^3*c^3*d^3*x^5 + 2*(25*b^3*c^4*d^2 - 15*a*b^2*c^3*d^3 + 3*a^2*b*c^2*d^4 + 3*a^3*c*d^5)*x^3 + 3*(5*b^3*c^5 - 3*a*b^2*c^4*d - a^2*b*c^3*d^2 - a^3*c^2*d^3) + (5*b^3*c^3*d^2 - 3*a*b^2*c^2*d^3 - a^2*b*c*d^4 - a^3*d^5)*x^4 + 2*(5*b^3*c^4*d - 3*a*b^2*c^3*d^2 - a^2*b*c^2*d^3 - a^3*c*d^4)*x^2)*sqrt(-c*d)*log((d*x^2 - 2*sqrt(-c*d)*x - c)/(d*x^2 + c)) + 2*(15*b^3*c^5*d - 9*a*b^2*c^4*d^2 - 3*a^2*b*c^3*d^3 + 5*a^3*c^2*d^4)*x)/(c^3*d^6*x^4 + 2*c^4*d^5*x^2 + c^5*d^4), 1/8*(8*b^3*c^3*d^3*x^5 + (25*b^3*c^4*d^2 - 15*a*b^2*c^3*d^3 + 3*a^2*b*c^2*d^4 + 3*a^3*c*d^5)*x^3 - 3*(5*b^3*c^5 - 3*a*b^2*c^4*d - a^2*b*c^3*d^2 - a^3*c^2*d^3) + (5*b^3*c^3*d^2 - 3*a*b^2*c^2*d^3 - a^2*b*c*d^4 - a^3*d^5)*x^4 + 2*(5*b^3*c^4*d - 3*a*b^2*c^3*d^2 - a^2*b*c^2*d^3 - a^3*c*d^4)*x^2)*sqrt(c*d)*arctan(sqrt(c*d)*x/c) + (15*b^3*c^5*d - 9*a*b^2*c^4*d^2 - 3*a^2*b*c^3*d^3 + 5*a^3*c^2*d^4)*x)/(c^3*d^6*x^4 + 2*c^4*d^5*x^2 + c^5*d^4)]

giac [A] time = 0.59, size = 180, normalized size = 1.38

$$\frac{b^3 x}{d^3} - \frac{3(5b^3 c^3 - 3ab^2 c^2 d - a^2 b c d^2 - a^3 d^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8\sqrt{cd}c^2 d^3} + \frac{9b^3 c^3 dx^3 - 15ab^2 c^2 d^2 x^3 + 3a^2 b c d^3 x^3 + 3a^3 d^4 x^3 + 7b^3 c^3 d^3 x^3}{8(dx^2 + c)^2 c^2 d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c)^3,x, algorithm="giac")

[Out] $b^3 x/d^3 - 3/8*(5*b^3*c^3 - 3*a*b^2*c^2*d - a^2*b*c*d^2 - a^3*d^3)*\arctan(d*x/\sqrt{c*d})/(\sqrt{c*d}*c^2*d^3) + 1/8*(9*b^3*c^3*d*x^3 - 15*a*b^2*c^2*d^2*x^3 + 3*a^2*b*c*d^3*x^3 + 3*a^3*d^4*x^3 + 7*b^3*c^3*d^3*x^3 - 9*a*b^2*c^3*d*x - 3*a^2*b*c^2*d^2*x + 5*a^3*c*d^3*x)/((d*x^2 + c)^2*c^2*d^3)$

maple [B] time = 0.01, size = 266, normalized size = 2.05

$$\frac{3a^3 d x^3}{8(dx^2 + c)^2 c^2} + \frac{3a^2 b x^3}{8(dx^2 + c)^2 c} - \frac{15a b^2 x^3}{8(dx^2 + c)^2 d} + \frac{9b^3 c x^3}{8(dx^2 + c)^2 d^2} + \frac{5a^3 x}{8(dx^2 + c)^2 c} - \frac{3a^2 b x}{8(dx^2 + c)^2 d} - \frac{9a b^2 c x}{8(dx^2 + c)^2 d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^3/(d*x^2+c)^3,x)

[Out] $b^3 x/d^3 + 3/8*d/(d*x^2+c)^2/c^2*x^3*a^3 + 3/8/(d*x^2+c)^2/c*x^3*a^2*b - 15/8/d/(d*x^2+c)^2*x^3*a*b^2 + 9/8/d^2/(d*x^2+c)^2*c*x^3*b^3 + 5/8/(d*x^2+c)^2/c*x*a^3 - 3/8/d/(d*x^2+c)^2*x*a^2*b - 9/8/d^2/(d*x^2+c)^2*c*x*a*b^2 + 7/8/d^3/(d*x^2+c)^2*c^2*x*b^3 + 3/8/c^2/(c*d)^(1/2)*\arctan(1/(c*d)^(1/2)*d*x)*a^3 + 3/8/d/c/(c*d)^(1/2)*\arctan(1/(c*d)^(1/2)*d*x)*a^2*b + 9/8/d^2/(c*d)^(1/2)*\arctan(1/(c*d)^(1/2)*d*x)*a*b^2 - 15/8/d^3*c/(c*d)^(1/2)*\arctan(1/(c*d)^(1/2)*d*x)*b^3$

maxima [A] time = 2.98, size = 187, normalized size = 1.44

$$\frac{b^3 x}{d^3} + \frac{3(3b^3 c^3 d - 5ab^2 c^2 d^2 + a^2 b c d^3 + a^3 d^4)x^3 + (7b^3 c^4 - 9ab^2 c^3 d - 3a^2 b c^2 d^2 + 5a^3 c d^3)x}{8(c^2 d^5 x^4 + 2c^3 d^4 x^2 + c^4 d^3)} - \frac{3(5b^3 c^3 - 3ab^2 c^2 d)}{8c^2 d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c)^3,x, algorithm="maxima")

[Out] $b^3 x/d^3 + 1/8*(3*(3*b^3*c^3*d - 5*a*b^2*c^2*d^2 + a^2*b*c*d^3 + a^3*d^4)*x^3 + (7*b^3*c^4 - 9*a*b^2*c^3*d - 3*a^2*b*c^2*d^2 + 5*a^3*c*d^3)*x)/(c^2*d^5*x^4 + 2*c^3*d^4*x^2 + c^4*d^3) - 3/8*(5*b^3*c^3 - 3*a*b^2*c^2*d - a^2*b*c*d^2 - a^3*d^3)*\arctan(d*x/\sqrt{c*d})/(\sqrt{c*d}*c^2*d^3)$

mupad [B] time = 4.96, size = 240, normalized size = 1.85

$$\frac{x(5a^3 d^3 - 3a^2 b c d^2 - 9a b^2 c^2 d + 7b^3 c^3)}{8c} + \frac{3x^3(a^3 d^4 + a^2 b c d^3 - 5a b^2 c^2 d^2 + 3b^3 c^3 d)}{8c^2} + \frac{b^3 x}{d^3} + \frac{3 \operatorname{atan}\left(\frac{\sqrt{d} x(a d - b c)(a^2 d^2 + 2 a b c d + 5 b^2 c^2)}{\sqrt{c}(a^3 d^3 + a^2 b c d^2 + 3 a b^2 c^2 d - 5 b^3 c^3)}\right)}{8c^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^3/(c + d*x^2)^3,x)

[Out] $((x*(5*a^3*d^3 + 7*b^3*c^3 - 9*a*b^2*c^2*d - 3*a^2*b*c*d^2))/(8*c) + (3*x^3*(a^3*d^4 + 3*b^3*c^3*d - 5*a*b^2*c^2*d^2 + a^2*b*c*d^3))/(8*c^2))/(c^2*d^3 + d^5*x^4) + (b^3*x)/d^3 + (3*\operatorname{atan}((d^(1/2)*x*(a*d - b*c)*(a^2*d^2 + 5*b^2*c^2 + 2*a*b*c*d))/(c^(1/2)*(a^3*d^3 - 5*b^3*c^3 + 3*a*b^2*c^2)))$

$2*d + a^2*b*c*d^2)))*(a*d - b*c)*(a^2*d^2 + 5*b^2*c^2 + 2*a*b*c*d))/(8*c^(5/2)*d^(7/2))$

sympy [B] time = 1.81, size = 422, normalized size = 3.25

$$\frac{b^3 x}{d^3} - \frac{3\sqrt{-\frac{1}{c^5 d^7}} (ad - bc) (a^2 d^2 + 2abcd + 5b^2 c^2) \log\left(-\frac{3c^3 d^3 \sqrt{-\frac{1}{c^5 d^7}} (ad - bc) (a^2 d^2 + 2abcd + 5b^2 c^2)}{3a^3 d^3 + 3a^2 bcd^2 + 9ab^2 c^2 d - 15b^3 c^3} + x\right)}{16} + \frac{3\sqrt{-\frac{1}{c^5 d^7}} (ad - bc) (a^2 d^2 + 2abcd + 5b^2 c^2)}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**3/(d*x**2+c)**3,x)

[Out] $b**3*x/d**3 - 3*sqrt(-1/(c**5*d**7))*(a*d - b*c)*(a**2*d**2 + 2*a*b*c*d + 5*b**2*c**2)*log(-3*c**3*d**3*sqrt(-1/(c**5*d**7))*(a*d - b*c)*(a**2*d**2 + 2*a*b*c*d + 5*b**2*c**2)/(3*a**3*d**3 + 3*a**2*b*c*d**2 + 9*a*b**2*c**2*d - 15*b**3*c**3) + x)/16 + 3*sqrt(-1/(c**5*d**7))*(a*d - b*c)*(a**2*d**2 + 2*a*b*c*d + 5*b**2*c**2)*log(3*c**3*d**3*sqrt(-1/(c**5*d**7))*(a*d - b*c)*(a**2*d**2 + 2*a*b*c*d + 5*b**2*c**2)/(3*a**3*d**3 + 3*a**2*b*c*d**2 + 9*a*b**2*c**2*d - 15*b**3*c**3) + x)/16 + (x**3*(3*a**3*d**4 + 3*a**2*b*c*d**3 - 15*a*b**2*c**2*d**2 + 9*b**3*c**3*d) + x*(5*a**3*c*d**3 - 3*a**2*b*c**2*d**2 - 9*a*b**2*c**3*d + 7*b**3*c**4))/(8*c**4*d**3 + 16*c**3*d**4*x**2 + 8*c**2*d**5*x**4)$

$$3.20 \quad \int \frac{(c+dx^2)^4}{a+bx^2} dx$$

Optimal. Leaf size=142

$$\frac{dx(2bc - ad)(a^2d^2 - 2abcd + 2b^2c^2)}{b^4} + \frac{d^2x^3(a^2d^2 - 4abcd + 6b^2c^2)}{3b^3} + \frac{(bc - ad)^4 \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}b^{9/2}} + \frac{d^3x^5(4bc - ad)}{5b^2} + \frac{d^4x^7}{7b}$$

[Out] d*(-a*d+2*b*c)*(a^2*d^2-2*a*b*c*d+2*b^2*c^2)*x/b^4+1/3*d^2*(a^2*d^2-4*a*b*c*d+6*b^2*c^2)*x^3/b^3+1/5*d^3*(-a*d+4*b*c)*x^5/b^2+1/7*d^4*x^7/b+(-a*d+b*c)^4*arctan(x*b^(1/2)/a^(1/2))/b^(9/2)/a^(1/2)

Rubi [A] time = 0.09, antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {390, 205}

$$\frac{d^2x^3(a^2d^2 - 4abcd + 6b^2c^2)}{3b^3} + \frac{dx(2bc - ad)(a^2d^2 - 2abcd + 2b^2c^2)}{b^4} + \frac{d^3x^5(4bc - ad)}{5b^2} + \frac{(bc - ad)^4 \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}b^{9/2}} + \frac{d^4x^7}{7b}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^4/(a + b*x^2), x]

[Out] (d*(2*b*c - a*d)*(2*b^2*c^2 - 2*a*b*c*d + a^2*d^2)*x)/b^4 + (d^2*(6*b^2*c^2 - 4*a*b*c*d + a^2*d^2)*x^3)/(3*b^3) + (d^3*(4*b*c - a*d)*x^5)/(5*b^2) + (d^4*x^7)/(7*b) + ((b*c - a*d)^4*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[a]*b^(9/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned} \int \frac{(c+dx^2)^4}{a+bx^2} dx &= \int \left(\frac{d(2bc - ad)(2b^2c^2 - 2abcd + a^2d^2)}{b^4} + \frac{d^2(6b^2c^2 - 4abcd + a^2d^2)x^2}{b^3} + \frac{d^3(4bc - ad)x^4}{b^2} + \frac{d^4x^6}{b} \right) dx \\ &= \frac{d(2bc - ad)(2b^2c^2 - 2abcd + a^2d^2)x}{b^4} + \frac{d^2(6b^2c^2 - 4abcd + a^2d^2)x^3}{3b^3} + \frac{d^3(4bc - ad)x^5}{5b^2} + \frac{d^4x^7}{7b} \\ &= \frac{d(2bc - ad)(2b^2c^2 - 2abcd + a^2d^2)x}{b^4} + \frac{d^2(6b^2c^2 - 4abcd + a^2d^2)x^3}{3b^3} + \frac{d^3(4bc - ad)x^5}{5b^2} + \frac{d^4x^7}{7b} \end{aligned}$$

Mathematica [A] time = 0.09, size = 136, normalized size = 0.96

$$\frac{dx(-105a^3d^3 + 35a^2bd^2(12c + dx^2) - 7ab^2d(90c^2 + 20cdx^2 + 3d^2x^4) + 3b^3(140c^3 + 70c^2dx^2 + 28cd^2x^4 + 5d^3x^6))}{105b^4}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^4/(a + b*x^2), x]

[Out] (d*x*(-105*a^3*d^3 + 35*a^2*b*d^2*(12*c + d*x^2) - 7*a*b^2*d*(90*c^2 + 20*c*d*x^2 + 3*d^2*x^4) + 3*b^3*(140*c^3 + 70*c^2*d*x^2 + 28*c*d^2*x^4 + 5*d^3*x^6))/(105*b^4) + ((b*c - a*d)^4*ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(Sqrt[a]*b^(9/2))

fricas [A] time = 0.65, size = 428, normalized size = 3.01

$$\frac{30 ab^4 d^4 x^7 + 42 (4 ab^4 cd^3 - a^2 b^3 d^4) x^5 + 70 (6 ab^4 c^2 d^2 - 4 a^2 b^3 cd^3 + a^3 b^2 d^4) x^3 - 105 (b^4 c^4 - 4 ab^3 c^3 d + 6 a^2 b^2 c^2 d^2 - 4 a^3 b c^2 d^3 + a^4 d^4) \arctan\left(\frac{bx}{\sqrt{ab}}\right) + 15 b^6 d^4 x^7 + 84 b^6 cd^3 x^5 - 21 ab^5 d^4 x^5 + 210 b^6 d^4 x^7}{210 ab^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a), x, algorithm="fricas")

[Out] [1/210*(30*a*b^4*d^4*x^7 + 42*(4*a*b^4*c*d^3 - a^2*b^3*d^4)*x^5 + 70*(6*a*b^4*c^2*d^2 - 4*a^2*b^3*c*d^3 + a^3*b^2*d^4)*x^3 - 105*(b^4*c^4 - 4*a*b^3*c^3*d + 6*a^2*b^2*c^2*d^2 - 4*a^3*b*c*d^3 + a^4*d^4)*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b)*x - a)/(b*x^2 + a)) + 210*(4*a*b^4*c^3*d - 6*a^2*b^3*c^2*d^2 + 4*a^3*b^2*c*d^3 - a^4*b*d^4)*x)/(a*b^5), 1/105*(15*a*b^4*d^4*x^7 + 21*(4*a*b^4*c*d^3 - a^2*b^3*d^4)*x^5 + 35*(6*a*b^4*c^2*d^2 - 4*a^2*b^3*c*d^3 + a^3*b^2*d^4)*x^3 + 105*(b^4*c^4 - 4*a*b^3*c^3*d + 6*a^2*b^2*c^2*d^2 - 4*a^3*b*c*d^3 + a^4*d^4)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + 105*(4*a*b^4*c^3*d - 6*a^2*b^3*c^2*d^2 + 4*a^3*b^2*c*d^3 - a^4*b*d^4)*x)/(a*b^5)]

giac [A] time = 0.58, size = 198, normalized size = 1.39

$$\frac{(b^4 c^4 - 4 ab^3 c^3 d + 6 a^2 b^2 c^2 d^2 - 4 a^3 b c d^3 + a^4 d^4) \arctan\left(\frac{bx}{\sqrt{ab}}\right) + 15 b^6 d^4 x^7 + 84 b^6 cd^3 x^5 - 21 ab^5 d^4 x^5 + 210 b^6 d^4 x^7}{\sqrt{ab} b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a), x, algorithm="giac")

[Out] (b^4*c^4 - 4*a*b^3*c^3*d + 6*a^2*b^2*c^2*d^2 - 4*a^3*b*c*d^3 + a^4*d^4)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*b^4) + 1/105*(15*b^6*d^4*x^7 + 84*b^6*c*d^3*x^5 - 21*a*b^5*d^4*x^5 + 210*b^6*c^2*d^2*x^3 - 140*a*b^5*c*d^3*x^3 + 35*a^2*b^4*d^4*x^3 + 420*b^6*c^3*d*x - 630*a*b^5*c^2*d^2*x + 420*a^2*b^4*c*d^3*x - 105*a^3*b^3*d^4*x)/b^7

maple [A] time = 0.01, size = 246, normalized size = 1.73

$$\frac{d^4 x^7}{7b} - \frac{a d^4 x^5}{5b^2} + \frac{4c d^3 x^5}{5b} + \frac{a^2 d^4 x^3}{3b^3} - \frac{4ac d^3 x^3}{3b^2} + \frac{2c^2 d^2 x^3}{b} + \frac{a^4 d^4 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab} b^4} - \frac{4a^3 c d^3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab} b^3} + \frac{6a^2 c^2 d^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab} b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^4/(b*x^2+a), x)

[Out] 1/7*d^4*x^7/b-1/5*d^4/b^2*x^5*a+4/5*d^3/b*x^5*c+1/3*d^4/b^3*x^3*a^2-4/3*d^3/b^2*x^3*a*c+2*d^2/b*x^3*c^2-d^4/b^4*a^3*x+4*d^3/b^3*a^2*c*x-6*d^2/b^2*a*c^2*x+4*d/b*c^3*x+1/b^4/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a^4*d^4-4/b^3/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a^3*c*d^3+6/b^2/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a^2*c^2*d^2-4/b/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a*c^3*d+1/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c^4

maxima [A] time = 3.03, size = 187, normalized size = 1.32

$$\frac{(b^4c^4 - 4ab^3c^3d + 6a^2b^2c^2d^2 - 4a^3bcd^3 + a^4d^4) \arctan\left(\frac{bx}{\sqrt{ab}}\right) + 15b^3d^4x^7 + 21(4b^3cd^3 - ab^2d^4)x^5 + 35(6b^3c^2d^2 - a^2b^2cd^3 + a^3d^4)x^3 + 105(4b^3c^3d - 6a^2b^2c^2d^2 + 4a^2b^3cd^3 - a^3d^4)x}{\sqrt{ab}b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a),x, algorithm="maxima")

[Out] (b^4*c^4 - 4*a*b^3*c^3*d + 6*a^2*b^2*c^2*d^2 - 4*a^3*b*c*d^3 + a^4*d^4)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*b^4) + 1/105*(15*b^3*d^4*x^7 + 21*(4*b^3*c*d^3 - a*b^2*d^4)*x^5 + 35*(6*b^3*c^2*d^2 - 4*a*b^2*c*d^3 + a^2*b*d^4)*x^3 + 105*(4*b^3*c^3*d - 6*a*b^2*c^2*d^2 + 4*a^2*b*c*d^3 - a^3*d^4)*x)/b^4

mupad [B] time = 4.86, size = 216, normalized size = 1.52

$$x \left(\frac{4c^3d}{b} - \frac{a \left(\frac{a \left(\frac{ad^4}{b^2} - \frac{4cd^3}{b} \right)}{b} + \frac{6c^2d^2}{b} \right)}{b} \right) - x^5 \left(\frac{ad^4}{5b^2} - \frac{4cd^3}{5b} \right) + x^3 \left(\frac{a \left(\frac{ad^4}{b^2} - \frac{4cd^3}{b} \right)}{3b} + \frac{2c^2d^2}{b} \right) + \frac{d^4x^7}{7b} + \frac{\operatorname{atan}\left(\frac{d^4x^7}{\sqrt{a}(a^4d^4 - 4a^3b^3cd^3 - 6a^2b^2c^2d^2 + 4ab^3cd^3 - a^4d^4)}\right)}{7b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^4/(a + b*x^2),x)

[Out] x*((4*c^3*d)/b - (a*((a*((a*d^4)/b^2 - (4*c*d^3)/b))/b + (6*c^2*d^2)/b))/b - x^5*((a*d^4)/(5*b^2) - (4*c*d^3)/(5*b)) + x^3*((a*((a*d^4)/b^2 - (4*c*d^3)/b))/(3*b) + (2*c^2*d^2)/b) + (d^4*x^7)/(7*b) + (atan((b^(1/2)*x*(a*d - b*c)^4)/(a^(1/2)*(a^4*d^4 + b^4*c^4 + 6*a^2*b^2*c^2*d^2 - 4*a*b^3*c^3*d - 4*a^3*b*c*d^3)))*(a*d - b*c)^4)/(a^(1/2)*b^(9/2))

sympy [B] time = 0.76, size = 326, normalized size = 2.30

$$x^5 \left(-\frac{ad^4}{5b^2} + \frac{4cd^3}{5b} \right) + x^3 \left(\frac{a^2d^4}{3b^3} - \frac{4acd^3}{3b^2} + \frac{2c^2d^2}{b} \right) + x \left(-\frac{a^3d^4}{b^4} + \frac{4a^2cd^3}{b^3} - \frac{6ac^2d^2}{b^2} + \frac{4c^3d}{b} \right) - \frac{\sqrt{-\frac{1}{ab^9}} (ad - bc)^4 \log\left(-\frac{a^4d^4 - 4a^3b^3cd^3 - 6a^2b^2c^2d^2 + 4ab^3cd^3 - a^4d^4}{(ad - bc)^4}\right)}{\sqrt{-\frac{1}{ab^9}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**4/(b*x**2+a),x)

[Out] x**5*(-a*d**4/(5*b**2) + 4*c*d**3/(5*b)) + x**3*(a**2*d**4/(3*b**3) - 4*a*c*d**3/(3*b**2) + 2*c**2*d**2/b) + x*(-a**3*d**4/b**4 + 4*a**2*c*d**3/b**3 - 6*a*c**2*d**2/b**2 + 4*c**3*d/b) - sqrt(-1/(a*b**9))*(a*d - b*c)**4*log(-a*b**4*sqrt(-1/(a*b**9))*(a*d - b*c)**4/(a**4*d**4 - 4*a**3*b*c*d**3 + 6*a**2*b**2*c**2*d**2 - 4*a*b**3*c**3*d + b**4*c**4) + x)/2 + sqrt(-1/(a*b**9))*(a*d - b*c)**4*log(a*b**4*sqrt(-1/(a*b**9))*(a*d - b*c)**4/(a**4*d**4 - 4*a**3*b*c*d**3 + 6*a**2*b**2*c**2*d**2 - 4*a*b**3*c**3*d + b**4*c**4) + x)/2 + d**4*x**7/(7*b)

$$3.21 \quad \int \frac{(c+dx^2)^3}{a+bx^2} dx$$

Optimal. Leaf size=98

$$\frac{dx(a^2d^2 - 3abcd + 3b^2c^2)}{b^3} + \frac{(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}b^{7/2}} + \frac{d^2x^3(3bc - ad)}{3b^2} + \frac{d^3x^5}{5b}$$

[Out] $d*(a^2*d^2-3*a*b*c*d+3*b^2*c^2)*x/b^3+1/3*d^2*(-a*d+3*b*c)*x^3/b^2+1/5*d^3*x^5/b+(-a*d+b*c)^3*arctan(x*b^(1/2)/a^(1/2))/b^(7/2)/a^(1/2)$

Rubi [A] time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {390, 205}

$$\frac{dx(a^2d^2 - 3abcd + 3b^2c^2)}{b^3} + \frac{d^2x^3(3bc - ad)}{3b^2} + \frac{(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}b^{7/2}} + \frac{d^3x^5}{5b}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^3/(a + b*x^2), x]

[Out] $(d*(3*b^2*c^2 - 3*a*b*c*d + a^2*d^2)*x)/b^3 + (d^2*(3*b*c - a*d)*x^3)/(3*b^2) + (d^3*x^5)/(5*b) + ((b*c - a*d)^3*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[a]*b^(7/2))$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned} \int \frac{(c+dx^2)^3}{a+bx^2} dx &= \int \left(\frac{d(3b^2c^2 - 3abcd + a^2d^2)}{b^3} + \frac{d^2(3bc - ad)x^2}{b^2} + \frac{d^3x^4}{b} + \frac{b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3}{b^3(a+bx^2)} \right) dx \\ &= \frac{d(3b^2c^2 - 3abcd + a^2d^2)x}{b^3} + \frac{d^2(3bc - ad)x^3}{3b^2} + \frac{d^3x^5}{5b} + \frac{(bc - ad)^3 \int \frac{1}{a+bx^2} dx}{b^3} \\ &= \frac{d(3b^2c^2 - 3abcd + a^2d^2)x}{b^3} + \frac{d^2(3bc - ad)x^3}{3b^2} + \frac{d^3x^5}{5b} + \frac{(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}b^{7/2}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 92, normalized size = 0.94

$$\frac{dx(15a^2d^2 - 5abd(9c + dx^2) + 3b^2(15c^2 + 5cdx^2 + d^2x^4))}{15b^3} + \frac{(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}b^{7/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^3/(a + b*x^2), x]

[Out] (d*x*(15*a^2*d^2 - 5*a*b*d*(9*c + d*x^2) + 3*b^2*(15*c^2 + 5*c*d*x^2 + d^2*x^4)))/(15*b^3) + ((b*c - a*d)^3*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[a]*b^(7/2))

fricas [A] time = 0.75, size = 292, normalized size = 2.98

$$\frac{6ab^3d^3x^5 + 10(3ab^3cd^2 - a^2b^2d^3)x^3 + 15(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3)\sqrt{-ab} \log\left(\frac{bx^2 + 2\sqrt{-ab}x - a}{bx^2 + a}\right) + 30(3}{30ab^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a), x, algorithm="fricas")

[Out] [1/30*(6*a*b^3*d^3*x^5 + 10*(3*a*b^3*c*d^2 - a^2*b^2*d^3)*x^3 + 15*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(-a*b)*log((b*x^2 + 2*sqrt(-a*b)*x - a)/(b*x^2 + a)) + 30*(3*a*b^3*c^2*d - 3*a^2*b^2*c*d^2 + a^3*b*d^3)*x)/(a*b^4), 1/15*(3*a*b^3*d^3*x^5 + 5*(3*a*b^3*c*d^2 - a^2*b^2*d^3)*x^3 + 15*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + 15*(3*a*b^3*c^2*d - 3*a^2*b^2*c*d^2 + a^3*b*d^3)*x)/(a*b^4)]

giac [A] time = 0.58, size = 129, normalized size = 1.32

$$\frac{(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b^3} + \frac{3b^4d^3x^5 + 15b^4cd^2x^3 - 5ab^3d^3x^3 + 45b^4c^2dx - 45ab^3cd^2x + 15b^5}{15b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a), x, algorithm="giac")

[Out] (b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*b^3) + 1/15*(3*b^4*d^3*x^5 + 15*b^4*c*d^2*x^3 - 5*a*b^3*d^3*x^3 + 45*b^4*c^2*d*x - 45*a*b^3*c*d^2*x + 15*a^2*b^2*d^3*x)/b^5

maple [A] time = 0.00, size = 161, normalized size = 1.64

$$\frac{d^3x^5}{5b} - \frac{a d^3x^3}{3b^2} + \frac{c d^2x^3}{b} - \frac{a^3d^3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b^3} + \frac{3a^2c d^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b^2} - \frac{3a c^2d \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b} + \frac{c^3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}} + \frac{a^2d^3x}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^3/(b*x^2+a), x)

[Out] 1/5/b*d^3*x^5-1/3*d^3/b^2*x^3*a+d^2/b*x^3*c+d^3/b^3*a^2*x-3*d^2/b^2*a*c*x+3*d/b*c^2*x-1/b^3/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a^3*d^3+3/b^2/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a^2*c*d^2-3/b/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a*c^2*d+1/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c^3

maxima [A] time = 2.91, size = 122, normalized size = 1.24

$$\frac{(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b^3} + \frac{3b^2d^3x^5 + 5(3b^2cd^2 - abd^3)x^3 + 15(3b^2c^2d - 3abcd^2 + a^2d^3)}{15b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a), x, algorithm="maxima")

[Out] $(b^3c^3 - 3ab^2c^2d + 3a^2b^2cd^2 - a^3d^3) \arctan(bx/\sqrt{ab}) / (\sqrt{ab}b^3) + 1/15(3b^2d^3x^5 + 5(3b^2cd^2 - ab^2d^3)x^3 + 15(3b^2c^2d - 3ab^2cd^2 + a^2d^3)x) / b^3$

mupad [B] time = 0.08, size = 146, normalized size = 1.49

$$x \left(\frac{3c^2d}{b} + \frac{a \left(\frac{ad^3}{b^2} - \frac{3cd^2}{b} \right)}{b} \right) - x^3 \left(\frac{ad^3}{3b^2} - \frac{cd^2}{b} \right) + \frac{d^3x^5}{5b} - \frac{\operatorname{atan} \left(\frac{\sqrt{b}x(ad-bc)^3}{\sqrt{a}(a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3)} \right) (ad-bc)^3}{\sqrt{a}b^{7/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^3/(a + b*x^2), x)`

[Out] $x((3c^2d)/b + (a((ad^3)/b^2 - (3cd^2)/b))/b) - x^3((ad^3)/(3b^2) - (cd^2)/b) + (d^3x^5)/(5b) - (\operatorname{atan}((b^{1/2})x(ad-bc)^3)/(a^{1/2}(a^3d^3 - b^3c^3 + 3ab^2cd^2 - 3a^2b^2cd^2)))(ad-bc)^3/(a^{1/2}b^{7/2})$

sympy [B] time = 0.59, size = 238, normalized size = 2.43

$$x^3 \left(-\frac{ad^3}{3b^2} + \frac{cd^2}{b} \right) + x \left(\frac{a^2d^3}{b^3} - \frac{3acd^2}{b^2} + \frac{3c^2d}{b} \right) + \frac{\sqrt{-\frac{1}{ab^7}} (ad-bc)^3 \log \left(-\frac{ab^3 \sqrt{-\frac{1}{ab^7}} (ad-bc)^3}{a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3} + x \right)}{2} - \frac{\sqrt{-\frac{1}{ab^7}} (ad-bc)^3}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**3/(b*x**2+a), x)`

[Out] $x^3(-ad^3/(3b^2) + cd^2/b) + x(a^2d^3/b^3 - 3acd^2/b^2 + 3c^2d/b) + \sqrt{-1/(ab^7)}(ad-bc)^3 \log(-ab^3 \sqrt{-1/(ab^7)}(ad-bc)^3 + x) / 2 - \sqrt{-1/(ab^7)}(ad-bc)^3 \log(ab^3 \sqrt{-1/(ab^7)}(ad-bc)^3 + x) / 2 + d^3x^5/(5b)$

$$3.22 \quad \int \frac{(c+dx^2)^2}{a+bx^2} dx$$

Optimal. Leaf size=63

$$\frac{(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}b^{5/2}} + \frac{dx(2bc-ad)}{b^2} + \frac{d^2x^3}{3b}$$

[Out] d*(-a*d+2*b*c)*x/b^2+1/3*d^2*x^3/b+(-a*d+b*c)^2*arctan(x*b^(1/2)/a^(1/2))/b^(5/2)/a^(1/2)

Rubi [A] time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {390, 205}

$$\frac{dx(2bc-ad)}{b^2} + \frac{(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}b^{5/2}} + \frac{d^2x^3}{3b}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^2/(a + b*x^2), x]

[Out] (d*(2*b*c - a*d)*x)/b^2 + (d^2*x^3)/(3*b) + ((b*c - a*d)^2*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[a]*b^(5/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned} \int \frac{(c+dx^2)^2}{a+bx^2} dx &= \int \left(\frac{d(2bc-ad)}{b^2} + \frac{d^2x^2}{b} + \frac{b^2c^2 - 2abcd + a^2d^2}{b^2(a+bx^2)} \right) dx \\ &= \frac{d(2bc-ad)x}{b^2} + \frac{d^2x^3}{3b} + \frac{(bc-ad)^2 \int \frac{1}{a+bx^2} dx}{b^2} \\ &= \frac{d(2bc-ad)x}{b^2} + \frac{d^2x^3}{3b} + \frac{(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}b^{5/2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 59, normalized size = 0.94

$$\frac{(bc-ad)^2 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}b^{5/2}} + \frac{dx(-3ad+6bc+bdx^2)}{3b^2}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^2/(a + b*x^2), x]

[Out] (d*x*(6*b*c - 3*a*d + b*d*x^2))/(3*b^2) + ((b*c - a*d)^2*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[a]*b^(5/2))

fricas [A] time = 0.56, size = 181, normalized size = 2.87

$$\left[\frac{2ab^2d^2x^3 - 3(b^2c^2 - 2abcd + a^2d^2)\sqrt{-ab} \log\left(\frac{bx^2 - 2\sqrt{-ab}x - a}{bx^2 + a}\right) + 6(2ab^2cd - a^2bd^2)x}{6ab^3}, \frac{ab^2d^2x^3 + 3(b^2c^2 - 2abcd + a^2d^2)\sqrt{-ab} \operatorname{arctan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{ab^3} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a), x, algorithm="fricas")

[Out] [1/6*(2*a*b^2*d^2*x^3 - 3*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b)*x - a)/(b*x^2 + a)) + 6*(2*a*b^2*c*d - a^2*b*d^2)*x)/(a*b^3), 1/3*(a*b^2*d^2*x^3 + 3*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + 3*(2*a*b^2*c*d - a^2*b*d^2)*x)/(a*b^3)]

giac [A] time = 0.57, size = 72, normalized size = 1.14

$$\frac{(b^2c^2 - 2abcd + a^2d^2) \operatorname{arctan}\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b^2} + \frac{b^2d^2x^3 + 6b^2cdx - 3abd^2x}{3b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a), x, algorithm="giac")

[Out] (b^2*c^2 - 2*a*b*c*d + a^2*d^2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*b^2) + 1/3*(b^2*d^2*x^3 + 6*b^2*c*d*x - 3*a*b*d^2*x)/b^3

maple [A] time = 0.00, size = 95, normalized size = 1.51

$$\frac{d^2x^3}{3b} + \frac{a^2d^2 \operatorname{arctan}\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b^2} - \frac{2acd \operatorname{arctan}\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b} + \frac{c^2 \operatorname{arctan}\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}} - \frac{ad^2x}{b^2} + \frac{2cdx}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^2/(b*x^2+a), x)

[Out] 1/3/b*d^2*x^3-d^2/b^2*a*x+2*d/b*x*c+1/b^2/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a^2*d^2-2/b/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a*c*d+1/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c^2

maxima [A] time = 2.93, size = 69, normalized size = 1.10

$$\frac{(b^2c^2 - 2abcd + a^2d^2) \operatorname{arctan}\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b^2} + \frac{bd^2x^3 + 3(2bcd - ad^2)x}{3b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a), x, algorithm="maxima")

[Out] (b^2*c^2 - 2*a*b*c*d + a^2*d^2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*b^2) + 1/3*(b*d^2*x^3 + 3*(2*b*c*d - a*d^2)*x)/b^2

mupad [B] time = 4.90, size = 90, normalized size = 1.43

$$\frac{d^2x^3}{3b} - x \left(\frac{ad^2}{b^2} - \frac{2cd}{b} \right) + \frac{\operatorname{atan}\left(\frac{\sqrt{b}x(ad-bc)^2}{\sqrt{a}(a^2d^2-2abcd+b^2c^2)}\right)(ad-bc)^2}{\sqrt{a}b^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^2/(a + b*x^2),x)`

[Out] $(d^2*x^3)/(3*b) - x*((a*d^2)/b^2 - (2*c*d)/b) + (\operatorname{atan}((b^{1/2})*x*(a*d - b*c)^2)/(a^{1/2}*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d)))*(a*d - b*c)^2/(a^{1/2}*b^{5/2})$

sympy [B] time = 0.44, size = 172, normalized size = 2.73

$$x\left(-\frac{ad^2}{b^2} + \frac{2cd}{b}\right) - \frac{\sqrt{-\frac{1}{ab^5}}(ad-bc)^2 \log\left(-\frac{ab^2\sqrt{-\frac{1}{ab^5}}(ad-bc)^2}{a^2d^2-2abcd+b^2c^2} + x\right)}{2} + \frac{\sqrt{-\frac{1}{ab^5}}(ad-bc)^2 \log\left(\frac{ab^2\sqrt{-\frac{1}{ab^5}}(ad-bc)^2}{a^2d^2-2abcd+b^2c^2} + x\right)}{2} + \frac{d^2}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**2/(b*x**2+a),x)`

[Out] $x*(-a*d**2/b**2 + 2*c*d/b) - \operatorname{sqrt}(-1/(a*b**5))*(a*d - b*c)**2*\log(-a*b**2*\operatorname{sqrt}(-1/(a*b**5))*(a*d - b*c)**2/(a**2*d**2 - 2*a*b*c*d + b**2*c**2) + x)/2 + \operatorname{sqrt}(-1/(a*b**5))*(a*d - b*c)**2*\log(a*b**2*\operatorname{sqrt}(-1/(a*b**5))*(a*d - b*c)**2/(a**2*d**2 - 2*a*b*c*d + b**2*c**2) + x)/2 + d**2*x**3/(3*b)$

3.23 $\int \frac{c+dx^2}{a+bx^2} dx$

Optimal. Leaf size=39

$$\frac{(bc - ad) \tan^{-1} \left(\frac{\sqrt{b}x}{\sqrt{a}} \right)}{\sqrt{a} b^{3/2}} + \frac{dx}{b}$$

[Out] d*x/b+(-a*d+b*c)*arctan(x*b^(1/2)/a^(1/2))/b^(3/2)/a^(1/2)

Rubi [A] time = 0.02, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {388, 205}

$$\frac{(bc - ad) \tan^{-1} \left(\frac{\sqrt{b}x}{\sqrt{a}} \right)}{\sqrt{a} b^{3/2}} + \frac{dx}{b}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)/(a + b*x^2), x]

[Out] (d*x)/b + ((b*c - a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[a]*b^(3/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int \frac{c + dx^2}{a + bx^2} dx &= \frac{dx}{b} - \frac{(-bc + ad) \int \frac{1}{a+bx^2} dx}{b} \\ &= \frac{dx}{b} + \frac{(bc - ad) \tan^{-1} \left(\frac{\sqrt{b}x}{\sqrt{a}} \right)}{\sqrt{a} b^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 40, normalized size = 1.03

$$\frac{dx}{b} - \frac{(ad - bc) \tan^{-1} \left(\frac{\sqrt{b}x}{\sqrt{a}} \right)}{\sqrt{a} b^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)/(a + b*x^2), x]

[Out] (d*x)/b - ((-b*c) + a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(Sqrt[a]*b^(3/2))

fricas [A] time = 0.71, size = 98, normalized size = 2.51

$$\left[\frac{2 abdx + \sqrt{-ab} (bc - ad) \log\left(\frac{bx^2 + 2\sqrt{-ab}x - a}{bx^2 + a}\right)}{2 ab^2}, \frac{abdx + \sqrt{ab} (bc - ad) \arctan\left(\frac{\sqrt{ab}x}{a}\right)}{ab^2} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a),x, algorithm="fricas")

[Out] [1/2*(2*a*b*d*x + sqrt(-a*b)*(b*c - a*d)*log((b*x^2 + 2*sqrt(-a*b)*x - a)/(b*x^2 + a)))/(a*b^2), (a*b*d*x + sqrt(a*b)*(b*c - a*d)*arctan(sqrt(a*b)*x/a))/(a*b^2)]

giac [A] time = 0.57, size = 33, normalized size = 0.85

$$\frac{dx}{b} + \frac{(bc - ad) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a),x, algorithm="giac")

[Out] d*x/b + (b*c - a*d)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*b)

maple [A] time = 0.00, size = 45, normalized size = 1.15

$$-\frac{ad \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab} b} + \frac{c \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}} + \frac{dx}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)/(b*x^2+a),x)

[Out] 1/b*d*x-1/b/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*a*d+1/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c

maxima [A] time = 2.99, size = 33, normalized size = 0.85

$$\frac{dx}{b} + \frac{(bc - ad) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a),x, algorithm="maxima")

[Out] d*x/b + (b*c - a*d)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*b)

mupad [B] time = 0.06, size = 32, normalized size = 0.82

$$\frac{dx}{b} - \frac{\operatorname{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) (ad - bc)}{\sqrt{a} b^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)/(a + b*x^2),x)

[Out] (d*x)/b - (atan((b^(1/2)*x)/a^(1/2))*(a*d - b*c))/(a^(1/2)*b^(3/2))

sympy [B] time = 0.28, size = 82, normalized size = 2.10

$$\frac{\sqrt{-\frac{1}{ab^3}} (ad - bc) \log\left(-ab\sqrt{-\frac{1}{ab^3}} + x\right)}{2} - \frac{\sqrt{-\frac{1}{ab^3}} (ad - bc) \log\left(ab\sqrt{-\frac{1}{ab^3}} + x\right)}{2} + \frac{dx}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)/(b*x**2+a),x)

[Out] sqrt(-1/(a*b**3))*(a*d - b*c)*log(-a*b*sqrt(-1/(a*b**3)) + x)/2 - sqrt(-1/(a*b**3))*(a*d - b*c)*log(a*b*sqrt(-1/(a*b**3)) + x)/2 + d*x/b

$$3.24 \quad \int \frac{1}{(a+bx^2)(c+dx^2)} dx$$

Optimal. Leaf size=70

$$\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)} - \frac{\sqrt{d} \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)}$$

[Out] arctan(x*b^(1/2)/a^(1/2))*b^(1/2)/(-a*d+b*c)/a^(1/2)-arctan(x*d^(1/2)/c^(1/2))*d^(1/2)/(-a*d+b*c)/c^(1/2)

Rubi [A] time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {391, 205}

$$\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)} - \frac{\sqrt{d} \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)*(c + d*x^2)),x]

[Out] (Sqrt[b]*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[a]*(b*c - a*d)) - (Sqrt[d]*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(Sqrt[c]*(b*c - a*d))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 391

Int[1/(((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_))), x_Symbol] := Dist[b/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[d/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)(c+dx^2)} dx &= \frac{b \int \frac{1}{a+bx^2} dx}{bc-ad} - \frac{d \int \frac{1}{c+dx^2} dx}{bc-ad} \\ &= \frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)} - \frac{\sqrt{d} \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)} \end{aligned}$$

Mathematica [A] time = 0.05, size = 61, normalized size = 0.87

$$\frac{\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\sqrt{d} \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}}}{bc-ad}$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)*(c + d*x^2)),x]

[Out] $\left(\frac{\sqrt{b} \operatorname{ArcTan}\left[\frac{\sqrt{b} x}{\sqrt{a}}\right]}{\sqrt{a}} - \frac{\sqrt{d} \operatorname{ArcTan}\left[\frac{\sqrt{d} x}{\sqrt{c}}\right]}{\sqrt{c}}\right) / (b c - a d)$

fricas [A] time = 0.85, size = 292, normalized size = 4.17

$$\left[\frac{\sqrt{-\frac{b}{a}} \log\left(\frac{bx^2 - 2ax\sqrt{-\frac{b}{a}} - a}{bx^2 + a}\right) + \sqrt{-\frac{d}{c}} \log\left(\frac{dx^2 + 2cx\sqrt{-\frac{d}{c}} - c}{dx^2 + c}\right)}{2(bc - ad)}, \frac{2\sqrt{\frac{d}{c}} \arctan\left(x\sqrt{\frac{d}{c}}\right) + \sqrt{-\frac{b}{a}} \log\left(\frac{bx^2 - 2ax\sqrt{-\frac{b}{a}} - a}{bx^2 + a}\right)}{2(bc - ad)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c),x, algorithm="fricas")

[Out] $[-1/2 * (\sqrt{-b/a} * \log((b*x^2 - 2*a*x*\sqrt{-b/a} - a)/(b*x^2 + a)) + \sqrt{-d/c} * \log((d*x^2 + 2*c*x*\sqrt{-d/c} - c)/(d*x^2 + c)))/(b*c - a*d), -1/2 * (2*\sqrt{d/c} * \arctan(x*\sqrt{d/c}) + \sqrt{-b/a} * \log((b*x^2 - 2*a*x*\sqrt{-b/a} - a)/(b*x^2 + a)))/(b*c - a*d), 1/2 * (2*\sqrt{b/a} * \arctan(x*\sqrt{b/a}) - \sqrt{-d/c} * \log((d*x^2 + 2*c*x*\sqrt{-d/c} - c)/(d*x^2 + c)))/(b*c - a*d), (\sqrt{b/a} * \arctan(x*\sqrt{b/a}) - \sqrt{d/c} * \arctan(x*\sqrt{d/c})) / (b*c - a*d)]$

giac [A] time = 0.59, size = 54, normalized size = 0.77

$$\frac{b \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}(bc - ad)} - \frac{d \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{(bc - ad)\sqrt{cd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c),x, algorithm="giac")

[Out] $b * \arctan(b*x/\sqrt{a*b}) / (\sqrt{a*b} * (b*c - a*d)) - d * \arctan(d*x/\sqrt{c*d}) / ((b*c - a*d) * \sqrt{c*d})$

maple [A] time = 0.01, size = 55, normalized size = 0.79

$$-\frac{b \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{(ad - bc)\sqrt{ab}} + \frac{d \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{(ad - bc)\sqrt{cd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)/(d*x^2+c),x)

[Out] $d/(a*d - b*c) / (c*d)^{1/2} * \arctan(1/(c*d)^{1/2} * d*x) - b/(a*d - b*c) / (a*b)^{1/2} * \arctan(1/(a*b)^{1/2} * b*x)$

maxima [A] time = 2.98, size = 54, normalized size = 0.77

$$\frac{b \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}(bc - ad)} - \frac{d \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{(bc - ad)\sqrt{cd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c),x, algorithm="maxima")

[Out] $b * \arctan(b*x/\sqrt{a*b}) / (\sqrt{a*b} * (b*c - a*d)) - d * \arctan(d*x/\sqrt{c*d}) / ((b*c - a*d) * \sqrt{c*d})$

mupad [B] time = 0.32, size = 135, normalized size = 1.93

$$\frac{\ln\left(\frac{bx - \sqrt{-ab}}{2a^2d - 2abc}\right) \sqrt{-ab}}{2(bc^2 - acd)} - \frac{\ln\left(\frac{dx + \sqrt{-cd}}{2(a^2d - abc)}\right) \sqrt{-cd}}{2(bc^2 - 2acd)} + \frac{\ln\left(\frac{bx + \sqrt{-ab}}{2(a^2d - abc)}\right) \sqrt{-ab}}{2(bc^2 - 2acd)} - \frac{\ln\left(\frac{dx - \sqrt{-cd}}{2bc^2 - 2acd}\right) \sqrt{-cd}}{2bc^2 - 2acd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((a + b*x^2)*(c + d*x^2)),x)`

[Out] $(\log(b*x - (-a*b)^{(1/2)})*(-a*b)^{(1/2)})/(2*a^2*d - 2*a*b*c) - (\log(d*x + (-c*d)^{(1/2)})*(-c*d)^{(1/2)})/(2*(b*c^2 - a*c*d)) - (\log(b*x + (-a*b)^{(1/2)})*(-a*b)^{(1/2)})/(2*(a^2*d - a*b*c)) + (\log(d*x - (-c*d)^{(1/2)})*(-c*d)^{(1/2)})/(2*b*c^2 - 2*a*c*d)$

sympy [B] time = 2.80, size = 712, normalized size = 10.17

$$\frac{\sqrt{-\frac{b}{a}} \log \left(x + \frac{-\frac{a^4 c d^3 \left(-\frac{b}{a}\right)^{\frac{3}{2}}}{(ad-bc)^3} + \frac{a^3 b c^2 d^2 \left(-\frac{b}{a}\right)^{\frac{3}{2}}}{(ad-bc)^3} + \frac{a^2 b^2 c^3 d \left(-\frac{b}{a}\right)^{\frac{3}{2}}}{(ad-bc)^3} - \frac{a^2 d^2 \sqrt{-\frac{b}{a}}}{ad-bc} - \frac{a b^3 c^4 \left(-\frac{b}{a}\right)^{\frac{3}{2}}}{(ad-bc)^3} - \frac{b^2 c^2 \sqrt{-\frac{b}{a}}}{ad-bc}}{bd} \right)}{2(ad-bc)} - \frac{\sqrt{-\frac{b}{a}} \log \left(x + \frac{\frac{a^4 c d^3 \left(-\frac{b}{a}\right)^{\frac{3}{2}}}{(ad-bc)^3} - \frac{a^3 b c^2 d^2 \left(-\frac{b}{a}\right)^{\frac{3}{2}}}{(ad-bc)^3}}{2(ad-bc)} \right)}{2(ad-bc)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x**2+a)/(d*x**2+c),x)`

[Out] $\sqrt{-b/a} \log(x + (-a**4*c*d**3*(-b/a)**(3/2)/(a*d - b*c)**3 + a**3*b*c**2*d**2*(-b/a)**(3/2)/(a*d - b*c)**3 + a**2*b**2*c**3*d*(-b/a)**(3/2)/(a*d - b*c)**3 - a**2*d**2*\sqrt{-b/a}/(a*d - b*c) - a*b**3*c**4*(-b/a)**(3/2)/(a*d - b*c)**3 - b**2*c**2*\sqrt{-b/a}/(a*d - b*c))/(b*d))/(2*(a*d - b*c)) - \sqrt{-b/a} \log(x + (a**4*c*d**3*(-b/a)**(3/2)/(a*d - b*c)**3 - a**3*b*c**2*d**2*(-b/a)**(3/2)/(a*d - b*c)**3 + a**2*b**2*c**3*d*(-b/a)**(3/2)/(a*d - b*c)**3 + a**2*d**2*\sqrt{-b/a}/(a*d - b*c) + a*b**3*c**4*(-b/a)**(3/2)/(a*d - b*c)**3 + b**2*c**2*\sqrt{-b/a}/(a*d - b*c))/(b*d))/(2*(a*d - b*c)) + \sqrt{-d/c} \log(x + (-a**4*c*d**3*(-d/c)**(3/2)/(a*d - b*c)**3 + a**3*b*c**2*d**2*(-d/c)**(3/2)/(a*d - b*c)**3 + a**2*b**2*c**3*d*(-d/c)**(3/2)/(a*d - b*c)**3 - a**2*d**2*\sqrt{-d/c}/(a*d - b*c) - a*b**3*c**4*(-d/c)**(3/2)/(a*d - b*c)**3 - b**2*c**2*\sqrt{-d/c}/(a*d - b*c))/(b*d))/(2*(a*d - b*c)) - \sqrt{-d/c} \log(x + (a**4*c*d**3*(-d/c)**(3/2)/(a*d - b*c)**3 - a**3*b*c**2*d**2*(-d/c)**(3/2)/(a*d - b*c)**3 - a**2*b**2*c**3*d*(-d/c)**(3/2)/(a*d - b*c)**3 + a**2*d**2*\sqrt{-d/c}/(a*d - b*c) + a*b**3*c**4*(-d/c)**(3/2)/(a*d - b*c)**3 + b**2*c**2*\sqrt{-d/c}/(a*d - b*c))/(b*d))/(2*(a*d - b*c))$

$$3.25 \quad \int \frac{1}{(a+bx^2)(c+dx^2)^2} dx$$

Optimal. Leaf size=109

$$\frac{b^{3/2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)^2} - \frac{\sqrt{d}(3bc-ad) \tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)}{2c^{3/2}(bc-ad)^2} - \frac{dx}{2c(c+dx^2)(bc-ad)}$$

[Out] $-1/2*d*x/c/(-a*d+b*c)/(d*x^2+c)+b^{(3/2)*\arctan(x*b^{(1/2)}/a^{(1/2)})}/(-a*d+b*c)^2/a^{(1/2)}-1/2*(-a*d+3*b*c)*\arctan(x*d^{(1/2)}/c^{(1/2)})*d^{(1/2)}/c^{(3/2)}/(-a*d+b*c)^2$

Rubi [A] time = 0.08, antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {414, 522, 205}

$$\frac{b^{3/2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)^2} - \frac{\sqrt{d}(3bc-ad) \tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)}{2c^{3/2}(bc-ad)^2} - \frac{dx}{2c(c+dx^2)(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)*(c + d*x^2)^2), x]

[Out] $-(d*x)/(2*c*(b*c - a*d)*(c + d*x^2)) + (b^{(3/2)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]})/(\text{Sqrt}[a]*(b*c - a*d)^2) - (\text{Sqrt}[d]*(3*b*c - a*d)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(2*c^{(3/2)}*(b*c - a*d)^2)$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 522

Int[((e_) + (f_.)*(x_)^(n_))/(((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_))), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)(c+dx^2)^2} dx &= -\frac{dx}{2c(bc-ad)(c+dx^2)} + \frac{\int \frac{2bc-ad-bdx^2}{(a+bx^2)(c+dx^2)} dx}{2c(bc-ad)} \\ &= -\frac{dx}{2c(bc-ad)(c+dx^2)} + \frac{b^2 \int \frac{1}{a+bx^2} dx}{(bc-ad)^2} - \frac{(d(3bc-ad)) \int \frac{1}{c+dx^2} dx}{2c(bc-ad)^2} \\ &= -\frac{dx}{2c(bc-ad)(c+dx^2)} + \frac{b^{3/2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)^2} - \frac{\sqrt{d}(3bc-ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{2c^{3/2}(bc-ad)^2} \end{aligned}$$

Mathematica [A] time = 0.17, size = 95, normalized size = 0.87

$$\frac{\frac{2b^{3/2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{\sqrt{d}(ad-3bc) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{c^{3/2}} + \frac{dx(ad-bc)}{c(c+dx^2)}}{2(bc-ad)^2}$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)*(c + d*x^2)^2), x]

[Out] ((d*(-(b*c) + a*d)*x)/(c*(c + d*x^2)) + (2*b^(3/2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/Sqrt[a] + (Sqrt[d]*(-3*b*c + a*d)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/c^(3/2))/(2*(b*c - a*d)^2)

fricas [A] time = 0.96, size = 711, normalized size = 6.52

$$\frac{2(bcdx^2 + bc^2)\sqrt{-\frac{b}{a}} \log\left(\frac{bx^2+2ax\sqrt{-\frac{b}{a}}-a}{bx^2+a}\right) - (3bc^2 - acd + (3bcd - ad^2)x^2)\sqrt{-\frac{d}{c}} \log\left(\frac{dx^2+2cx\sqrt{-\frac{d}{c}}-c}{dx^2+c}\right) - 2(bcd - a^2d)}{4(b^2c^4 - 2abc^3d + a^2c^2d^2 + (b^2c^3d - 2abc^2d^2 + a^2cd^3)x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] [1/4*(2*(b*c*d*x^2 + b*c^2)*sqrt(-b/a)*log((b*x^2 + 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) - (3*b*c^2 - a*c*d + (3*b*c*d - a*d^2)*x^2)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) - 2*(b*c*d - a*d^2)*x)/(b^2*c^4 - 2*a*b*c^3*d + a^2*c^2*d^2 + (b^2*c^3*d - 2*a*b*c^2*d^2 + a^2*c*d^3)*x^2), -1/2*((3*b*c^2 - a*c*d + (3*b*c*d - a*d^2)*x^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) - (b*c*d*x^2 + b*c^2)*sqrt(-b/a)*log((b*x^2 + 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + (b*c*d - a*d^2)*x)/(b^2*c^4 - 2*a*b*c^3*d + a^2*c^2*d^2 + (b^2*c^3*d - 2*a*b*c^2*d^2 + a^2*c*d^3)*x^2), 1/4*(4*(b*c*d*x^2 + b*c^2)*sqrt(b/a)*arctan(x*sqrt(b/a)) - (3*b*c^2 - a*c*d + (3*b*c*d - a*d^2)*x^2)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) - 2*(b*c*d - a*d^2)*x)/(b^2*c^4 - 2*a*b*c^3*d + a^2*c^2*d^2 + (b^2*c^3*d - 2*a*b*c^2*d^2 + a^2*c*d^3)*x^2), 1/2*(2*(b*c*d*x^2 + b*c^2)*sqrt(b/a)*arctan(x*sqrt(b/a)) - (3*b*c^2 - a*c*d + (3*b*c*d - a*d^2)*x^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) - (b*c*d - a*d^2)*x)/(b^2*c^4 - 2*a*b*c^3*d + a^2*c^2*d^2 + (b^2*c^3*d - 2*a*b*c^2*d^2 + a^2*c*d^3)*x^2)]

giac [A] time = 0.57, size = 122, normalized size = 1.12

$$\frac{b^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{(b^2c^2 - 2abcd + a^2d^2)\sqrt{ab}} - \frac{(3bcd - ad^2) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2(b^2c^3 - 2abc^2d + a^2cd^2)\sqrt{cd}} - \frac{dx}{2(bc^2 - acd)(dx^2 + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^2,x, algorithm="giac")

[Out] $b^2 \arctan\left(\frac{bx}{\sqrt{a*b}}\right) / ((b^2*c^2 - 2*a*b*c*d + a^2*d^2)*\sqrt{a*b}) - 1/2 * (3*b*c*d - a*d^2) \arctan\left(\frac{dx}{\sqrt{c*d}}\right) / ((b^2*c^3 - 2*a*b*c^2*d + a^2*c*d^2)*\sqrt{c*d}) - 1/2*d*x / ((b*c^2 - a*c*d)*(d*x^2 + c))$

maple [A] time = 0.01, size = 144, normalized size = 1.32

$$\frac{a d^2 x}{2(ad-bc)^2(d x^2+c)c} + \frac{a d^2 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2(ad-bc)^2 \sqrt{cd} c} + \frac{b^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{(ad-bc)^2 \sqrt{ab}} - \frac{bdx}{2(ad-bc)^2(d x^2+c)} - \frac{3bd \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2(ad-bc)^2 \sqrt{cd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)/(d*x^2+c)^2,x)

[Out] $1/2*d^2/(a*d-b*c)^2/c*x/(d*x^2+c)*a-1/2*d/(a*d-b*c)^2*x/(d*x^2+c)*b+1/2*d^2/(a*d-b*c)^2/c/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*a-3/2*d/(a*d-b*c)^2/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)*b+b^2/(a*d-b*c)^2/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)$

maxima [A] time = 3.04, size = 133, normalized size = 1.22

$$\frac{b^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{(b^2c^2 - 2abcd + a^2d^2)\sqrt{ab}} - \frac{dx}{2(bc^3 - ac^2d + (bc^2d - acd^2)x^2)} - \frac{(3bcd - ad^2) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2(b^2c^3 - 2abc^2d + a^2cd^2)\sqrt{cd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] $b^2 \arctan\left(\frac{bx}{\sqrt{a*b}}\right) / ((b^2*c^2 - 2*a*b*c*d + a^2*d^2)*\sqrt{a*b}) - 1/2 * d*x / (b*c^3 - a*c^2*d + (b*c^2*d - a*c*d^2)*x^2) - 1/2 * (3*b*c*d - a*d^2) \arctan\left(\frac{dx}{\sqrt{c*d}}\right) / ((b^2*c^3 - 2*a*b*c^2*d + a^2*c*d^2)*\sqrt{c*d})$

mupad [B] time = 5.69, size = 3637, normalized size = 33.37

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)*(c + d*x^2)^2),x)

[Out] $(d*x)/(2*c*(c + d*x^2)*(a*d - b*c)) - (\operatorname{atan}(\frac{(-c^3*d)^{(1/2)}*(a*d - 3*b*c)}{(x*(a^2*b^3*d^5 + 13*b^5*c^2*d^3 - 6*a*b^4*c*d^4))/(2*(b^2*c^4 + a^2*c^2*d^2 - 2*a*b*c^3*d)) - ((4*b^7*c^6*d^2 - 18*a*b^6*c^5*d^3 - 2*a^5*b^2*c*d^7 + 32*a^2*b^5*c^4*d^4 - 28*a^3*b^4*c^3*d^5 + 12*a^4*b^3*c^2*d^6)/(b^3*c^5 - a^3*c^2*d^3 + 3*a^2*b*c^3*d^2 - 3*a*b^2*c^4*d) - (x*(-c^3*d)^{(1/2)}*(a*d - 3*b*c)*(16*b^7*c^7*d^2 - 48*a*b^6*c^6*d^3 + 32*a^2*b^5*c^5*d^4 + 32*a^3*b^4*c^4*d^5 - 48*a^4*b^3*c^3*d^6 + 16*a^5*b^2*c^2*d^7))/(8*(b^2*c^4 + a^2*c^2*d^2 - 2*a*b*c^3*d)*(b^2*c^5 + a^2*c^3*d^2 - 2*a*b*c^4*d)))*(-c^3*d)^{(1/2)}*(a*d - 3*b*c))/(4*(b^2*c^5 + a^2*c^3*d^2 - 2*a*b*c^4*d))*1i)/(4*(b^2*c^5 + a^2*c^3*d^2 - 2*a*b*c^4*d)) + ((-c^3*d)^{(1/2)}*(a*d - 3*b*c)*((x*(a^2*b^3*d^5 + 13*b^5*c^2*d^3 - 6*a*b^4*c*d^4))/(2*(b^2*c^4 + a^2*c^2*d^2 - 2*a*b*c^3*d)) + ((4*b^7*c^6*d^2 - 18*a*b^6*c^5*d^3 - 2*a^5*b^2*c*d^7 + 32*a^2*b^5*c^4*d^4 - 28*a^3*b^4*c^3*d^5 + 12*a^4*b^3*c^2*d^6)/(b^3*c^5 - a^3*c^2*d^3 + 3*a^2*b*c^3*d^2 - 3*a*b^2*c^4*d) + (x*(-c^3*d)^{(1/2)}*(a*d - 3*b*c)*(16*b^7*c^7*d^2 - 48*a*b^6*c^6*d^3 + 32*a^2*b^5*c^5*d^4 + 32*a^3*b^4*c^4*d^5 - 48*a^4*b^3*c^3*d^6 + 16*a^5*b^2*c^2*d^7))/(8*(b^2*c^4 + a^2*c^2*d^2 - 2*a*b*c^3*d))*(b^2*c^5 + a^2*c^3*d^2 - 2*a*b*c^4*d)))*(-c^3*d)^{(1/2)}*(a*d - 3*b*c))/(4*$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x**2+a)/(d*x**2+c)**2,x)
```

```
[Out] Timed out
```

$$3.26 \quad \int \frac{1}{(a+bx^2)(c+dx^2)^3} dx$$

Optimal. Leaf size=160

$$\frac{\sqrt{d} (3a^2d^2 - 10abcd + 15b^2c^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) + \frac{b^{5/2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)^3} - \frac{dx(7bc-3ad)}{8c^2(c+dx^2)(bc-ad)^2} - \frac{dx}{4c(c+dx^2)^2(bc-ad)}}{8c^{5/2}(bc-ad)^3}$$

[Out] $-1/4*d*x/c/(-a*d+b*c)/(d*x^2+c)^2-1/8*d*(-3*a*d+7*b*c)*x/c^2/(-a*d+b*c)^2/(d*x^2+c)+b^{(5/2)*\arctan(x*b^{(1/2)}/a^{(1/2)})}/(-a*d+b*c)^3/a^{(1/2)}-1/8*(3*a^2*d^2-10*a*b*c*d+15*b^2*c^2)*\arctan(x*d^{(1/2)}/c^{(1/2)})*d^{(1/2)}/c^{(5/2)}/(-a*d+b*c)^3$

Rubi [A] time = 0.19, antiderivative size = 160, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {414, 527, 522, 205}

$$\frac{\sqrt{d} (3a^2d^2 - 10abcd + 15b^2c^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) + \frac{b^{5/2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)^3} - \frac{dx(7bc-3ad)}{8c^2(c+dx^2)(bc-ad)^2} - \frac{dx}{4c(c+dx^2)^2(bc-ad)}}{8c^{5/2}(bc-ad)^3}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)*(c + d*x^2)^3), x]

[Out] $-(d*x)/(4*c*(b*c - a*d)*(c + d*x^2)^2) - (d*(7*b*c - 3*a*d)*x)/(8*c^2*(b*c - a*d)^2*(c + d*x^2)) + (b^{(5/2)*\text{ArcTan}[\text{Sqrt}[b]*x]/\text{Sqrt}[a]})/(\text{Sqrt}[a]*(b*c - a*d)^3) - (\text{Sqrt}[d]*(15*b^2*c^2 - 10*a*b*c*d + 3*a^2*d^2)*\text{ArcTan}[\text{Sqrt}[d]*x]/\text{Sqrt}[c])/(8*c^{(5/2)}*(b*c - a*d)^3)$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 522

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(b*c - a*d)*(p+1)), x] + Dist[1/(a*n*(b*c - a*d)*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2)+1)*x^n, x], x], x] /; FreeQ

[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)(c+dx^2)^3} dx &= -\frac{dx}{4c(bc-ad)(c+dx^2)^2} + \frac{\int \frac{4bc-3ad-3bdx^2}{(a+bx^2)(c+dx^2)^2} dx}{4c(bc-ad)} \\ &= -\frac{dx}{4c(bc-ad)(c+dx^2)^2} - \frac{d(7bc-3ad)x}{8c^2(bc-ad)^2(c+dx^2)} + \frac{\int \frac{8b^2c^2-7abcd+3a^2d^2-bd(7bc-3ad)}{(a+bx^2)(c+dx^2)} dx}{8c^2(bc-ad)^2} \\ &= -\frac{dx}{4c(bc-ad)(c+dx^2)^2} - \frac{d(7bc-3ad)x}{8c^2(bc-ad)^2(c+dx^2)} + \frac{b^3 \int \frac{1}{a+bx^2} dx}{(bc-ad)^3} - \frac{d(15b^2c^2-7bcd+3a^2d^2)}{8c^2(bc-ad)^2} \\ &= -\frac{dx}{4c(bc-ad)(c+dx^2)^2} - \frac{d(7bc-3ad)x}{8c^2(bc-ad)^2(c+dx^2)} + \frac{b^{5/2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}(bc-ad)^3} - \frac{\sqrt{d}(15b^2c^2-7bcd+3a^2d^2)}{8c^2(bc-ad)^2} \end{aligned}$$

Mathematica [A] time = 0.24, size = 158, normalized size = 0.99

$$\frac{1}{8} \left(\frac{\sqrt{d} (3a^2d^2 - 10abcd + 15b^2c^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{c^{5/2}(bc-ad)^3} - \frac{8b^{5/2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}(ad-bc)^3} + \frac{dx(3ad-7bc)}{c^2(c+dx^2)(bc-ad)^2} - \frac{2dx}{c(c+dx^2)^2} \right)$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)*(c + d*x^2)^3), x]

[Out] ((-2*d*x)/(c*(b*c - a*d)*(c + d*x^2)^2) + (d*(-7*b*c + 3*a*d)*x)/(c^2*(b*c - a*d)^2*(c + d*x^2)) - (8*b^(5/2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[a]*(- (b*c) + a*d)^3) - (Sqrt[d]*(15*b^2*c^2 - 10*a*b*c*d + 3*a^2*d^2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(c^(5/2)*(b*c - a*d)^3))/8

fricas [B] time = 2.14, size = 1585, normalized size = 9.91

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [-1/16*(2*(7*b^2*c^2*d^2 - 10*a*b*c*d^3 + 3*a^2*d^4)*x^3 + 8*(b^2*c^2*d^2*x^4 + 2*b^2*c^3*d*x^2 + b^2*c^4)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + (15*b^2*c^4 - 10*a*b*c^3*d + 3*a^2*c^2*d^2 + (15*b^2*c^2*d^2 - 10*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(15*b^2*c^3*d - 10*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) + 2*(9*b^2*c^3*d - 14*a*b*c^2*d^2 + 5*a^2*c*d^3)*x)/(b^3*c^7 - 3*a*b^2*c^6*d + 3*a^2*b*c^5*d^2 - a^3*c^4*d^3 + (b^3*c^5*d^2 - 3*a*b^2*c^4*d^3 + 3*a^2*b*c^3*d^4 - a^3*c^2*d^5)*x^4 + 2*(b^3*c^6*d - 3*a*b^2*c^5*d^2 + 3*a^2*b*c^4*d^3 - a^3*c^3*d^4)*x^2), -1/8*((7*b^2*c^2*d^2 - 10*a*b*c*d^3 + 3*a^2*d^4)*x^3 + (15*b^2*c^4 - 10*a*b*c^3*d + 3*a^2*c^2*d^2 + (15*b^2*c^2*d^2 - 10*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(15*b^2*c^3*d - 10*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) + 4*(b^2*c^2*d^2*x^4 + 2*b^2*c^3*d*x^2 + b^2*c^4)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + (9*b^2*c^3*d - 14*a*b*c^2*d^2 + 5*a^2*c*d^3)*x)/(b^3*c^7 - 3*a*b^2*c^6*d + 3*a^2*b*c^5*d^2 - a^3*c^4*d^3 + (b^3*c^5*d^2 - 3*a*b^2*c^4*d^3 + 3*a^2*b*c^3*d^4

- a^3*c^2*d^5)*x^4 + 2*(b^3*c^6*d - 3*a*b^2*c^5*d^2 + 3*a^2*b*c^4*d^3 - a^3*c^3*d^4)*x^2), -1/16*(2*(7*b^2*c^2*d^2 - 10*a*b*c*d^3 + 3*a^2*d^4)*x^3 - 16*(b^2*c^2*d^2*x^4 + 2*b^2*c^3*d*x^2 + b^2*c^4)*sqrt(b/a)*arctan(x*sqrt(b/a)) + (15*b^2*c^4 - 10*a*b*c^3*d + 3*a^2*c^2*d^2 + (15*b^2*c^2*d^2 - 10*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(15*b^2*c^3*d - 10*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) + 2*(9*b^2*c^3*d - 14*a*b*c^2*d^2 + 5*a^2*c*d^3)*x)/(b^3*c^7 - 3*a*b^2*c^6*d + 3*a^2*b*c^5*d^2 - a^3*c^4*d^3 + (b^3*c^5*d^2 - 3*a*b^2*c^4*d^3 + 3*a^2*b*c^3*d^4 - a^3*c^2*d^5)*x^4 + 2*(b^3*c^6*d - 3*a*b^2*c^5*d^2 + 3*a^2*b*c^4*d^3 - a^3*c^3*d^4)*x^2), -1/8*((7*b^2*c^2*d^2 - 10*a*b*c*d^3 + 3*a^2*d^4)*x^3 - 8*(b^2*c^2*d^2*x^4 + 2*b^2*c^3*d*x^2 + b^2*c^4)*sqrt(b/a)*arctan(x*sqrt(b/a)) + (15*b^2*c^4 - 10*a*b*c^3*d + 3*a^2*c^2*d^2 + (15*b^2*c^2*d^2 - 10*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(15*b^2*c^3*d - 10*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) + (9*b^2*c^3*d - 14*a*b*c^2*d^2 + 5*a^2*c*d^3)*x)/(b^3*c^7 - 3*a*b^2*c^6*d + 3*a^2*b*c^5*d^2 - a^3*c^4*d^3 + (b^3*c^5*d^2 - 3*a*b^2*c^4*d^3 + 3*a^2*b*c^3*d^4 - a^3*c^2*d^5)*x^4 + 2*(b^3*c^6*d - 3*a*b^2*c^5*d^2 + 3*a^2*b*c^4*d^3 - a^3*c^3*d^4)*x^2)]

giac [A] time = 0.58, size = 217, normalized size = 1.36

$$\frac{b^3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3)\sqrt{ab}} - \frac{(15b^2c^2d - 10abcd^2 + 3a^2d^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8(b^3c^5 - 3ab^2c^4d + 3a^2bc^3d^2 - a^3c^2d^3)\sqrt{cd}} - \frac{7bcd^2x^3 - 3ad^3x^3 + 9b^2cd^2x}{8(b^2c^4 - 2abc^3d + a^2c^2d)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^3,x, algorithm="giac")

[Out] b^3*arctan(b*x/sqrt(a*b))/((b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(a*b)) - 1/8*(15*b^2*c^2*d - 10*a*b*c*d^2 + 3*a^2*d^3)*arctan(d*x/sqrt(c*d))/((b^3*c^5 - 3*a*b^2*c^4*d + 3*a^2*b*c^3*d^2 - a^3*c^2*d^3)*sqrt(c*d)) - 1/8*(7*b*c*d^2*x^3 - 3*a*d^3*x^3 + 9*b*c^2*d*x - 5*a*c*d^2*x)/((b^2*c^4 - 2*a*b*c^3*d + a^2*c^2*d^2)*(d*x^2 + c)^2)

maple [B] time = 0.01, size = 310, normalized size = 1.94

$$\frac{3a^2d^4x^3}{8(ad - bc)^3(d x^2 + c)^2 c^2} - \frac{5abd^3x^3}{4(ad - bc)^3(d x^2 + c)^2 c} + \frac{7b^2d^2x^3}{8(ad - bc)^3(d x^2 + c)^2} + \frac{5a^2d^3x}{8(ad - bc)^3(d x^2 + c)^2 c} - \frac{7b^2cd^2x}{4(ad - bc)^3(d x^2 + c)^2 c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)/(d*x^2+c)^3,x)

[Out] 3/8*d^4/(a*d-b*c)^3/(d*x^2+c)^2/c^2*x^3*a^2-5/4*d^3/(a*d-b*c)^3/(d*x^2+c)^2/c*x^3*a*b+7/8*d^2/(a*d-b*c)^3/(d*x^2+c)^2*x^3*b^2+5/8*d^3/(a*d-b*c)^3/(d*x^2+c)^2/c*x*a^2-7/4*d^2/(a*d-b*c)^3/(d*x^2+c)^2*x*a*b+9/8*d/(a*d-b*c)^3/(d*x^2+c)^2*c*x*b^2+3/8*d^3/(a*d-b*c)^3/c^2/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a^2-5/4*d^2/(a*d-b*c)^3/c/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a*b+15/8*d/(a*d-b*c)^3/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*b^2-b^3/(a*d-b*c)^3/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)

maxima [B] time = 3.08, size = 277, normalized size = 1.73

$$\frac{b^3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3)\sqrt{ab}} - \frac{(15b^2c^2d - 10abcd^2 + 3a^2d^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8(b^3c^5 - 3ab^2c^4d + 3a^2bc^3d^2 - a^3c^2d^3)\sqrt{cd}} - \frac{8(b^2c^6 - 2abc^5d + a^2c^4d)}{8(b^2c^6 - 2abc^5d + a^2c^4d)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^3,x, algorithm="maxima")

$$\begin{aligned}
& *c^4*d^3 - 300*a*b^6*c^3*d^4 - 60*a^3*b^4*c*d^6 + 190*a^2*b^5*c^2*d^5))/((32 \\
& *(b^4*c^8 + a^4*c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^2*b^2*c^6*d^2 - 4*a*b^3*c^7 \\
& *d)) + ((-a*b^5)^{(1/2)}*((256*b^10*c^10*d^2 - 1760*a*b^9*c^9*d^3 + 5280*a^2* \\
& b^8*c^8*d^4 - 9056*a^3*b^7*c^7*d^5 + 9760*a^4*b^6*c^6*d^6 - 6816*a^5*b^5*c^5*d^7 + 3040*a^6*b^4*c^4*d^8 - 800*a^7*b^3*c^3*d^9 + 96*a^8*b^2*c^2*d^10)/(\\
& 64*(b^6*c^10 + a^6*c^4*d^6 - 6*a^5*b*c^5*d^5 + 15*a^2*b^4*c^8*d^2 - 20*a^3* \\
& b^3*c^7*d^3 + 15*a^4*b^2*c^6*d^4 - 6*a*b^5*c^9*d)) + (x*(-a*b^5)^{(1/2)}*(256 \\
& *b^9*c^11*d^2 - 1280*a*b^8*c^10*d^3 + 2304*a^2*b^7*c^9*d^4 - 1280*a^3*b^6*c^8*d^5 - 1280*a^4*b^5*c^7*d^6 + 2304*a^5*b^4*c^6*d^7 - 1280*a^6*b^3*c^5*d^8 \\
& + 256*a^7*b^2*c^4*d^9))/(64*(a^4*d^3 - a*b^3*c^3 + 3*a^2*b^2*c^2*d - 3*a^3* \\
& *b*c*d^2))*(b^4*c^8 + a^4*c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^2*b^2*c^6*d^2 - 4* \\
& a*b^3*c^7*d))))/(2*(a^4*d^3 - a*b^3*c^3 + 3*a^2*b^2*c^2*d - 3*a^3*b*c*d^2)) \\
&))/(2*(a^4*d^3 - a*b^3*c^3 + 3*a^2*b^2*c^2*d - 3*a^3*b*c*d^2)))*(-a*b^5)^{(\\
& 1/2)*1i)/(a^4*d^3 - a*b^3*c^3 + 3*a^2*b^2*c^2*d - 3*a^3*b*c*d^2) - (atan(((\\
& ((x*(9*a^4*b^3*d^7 + 289*b^7*c^4*d^3 - 300*a*b^6*c^3*d^4 - 60*a^3*b^4*c*d^6 \\
& + 190*a^2*b^5*c^2*d^5))/(32*(b^4*c^8 + a^4*c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^ \\
& ^2*b^2*c^6*d^2 - 4*a*b^3*c^7*d)) - (((256*b^10*c^10*d^2 - 1760*a*b^9*c^9*d^ \\
& 3 + 5280*a^2*b^8*c^8*d^4 - 9056*a^3*b^7*c^7*d^5 + 9760*a^4*b^6*c^6*d^6 - 68 \\
& 16*a^5*b^5*c^5*d^7 + 3040*a^6*b^4*c^4*d^8 - 800*a^7*b^3*c^3*d^9 + 96*a^8*b^ \\
& 2*c^2*d^10)/(64*(b^6*c^10 + a^6*c^4*d^6 - 6*a^5*b*c^5*d^5 + 15*a^2*b^4*c^8* \\
& d^2 - 20*a^3*b^3*c^7*d^3 + 15*a^4*b^2*c^6*d^4 - 6*a*b^5*c^9*d)) - (x*(-c^5* \\
& d)^{(1/2)}*(3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d))*(256*b^9*c^11*d^2 - 1280*a*b \\
& ^8*c^10*d^3 + 2304*a^2*b^7*c^9*d^4 - 1280*a^3*b^6*c^8*d^5 - 1280*a^4*b^5*c^ \\
& 7*d^6 + 2304*a^5*b^4*c^6*d^7 - 1280*a^6*b^3*c^5*d^8 + 256*a^7*b^2*c^4*d^9)) \\
& /((512*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d))*(b^4*c^8 + \\
& a^4*c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^2*b^2*c^6*d^2 - 4*a*b^3*c^7*d)))*(-c^5* \\
& d)^{(1/2)}*(3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d))/(16*(b^3*c^8 - a^3*c^5*d^3 \\
& + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d))*(-c^5*d)^{(1/2)}*(3*a^2*d^2 + 15*b^2*c^2 \\
& - 10*a*b*c*d)*1i)/(16*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^ \\
& ^7*d)) + (((x*(9*a^4*b^3*d^7 + 289*b^7*c^4*d^3 - 300*a*b^6*c^3*d^4 - 60*a^3 \\
& *b^4*c*d^6 + 190*a^2*b^5*c^2*d^5))/(32*(b^4*c^8 + a^4*c^4*d^4 - 4*a^3*b*c^5 \\
& *d^3 + 6*a^2*b^2*c^6*d^2 - 4*a*b^3*c^7*d)) + (((256*b^10*c^10*d^2 - 1760*a* \\
& b^9*c^9*d^3 + 5280*a^2*b^8*c^8*d^4 - 9056*a^3*b^7*c^7*d^5 + 9760*a^4*b^6*c^ \\
& 6*d^6 - 6816*a^5*b^5*c^5*d^7 + 3040*a^6*b^4*c^4*d^8 - 800*a^7*b^3*c^3*d^9 + \\
& 96*a^8*b^2*c^2*d^10)/(64*(b^6*c^10 + a^6*c^4*d^6 - 6*a^5*b*c^5*d^5 + 15*a^ \\
& 2*b^4*c^8*d^2 - 20*a^3*b^3*c^7*d^3 + 15*a^4*b^2*c^6*d^4 - 6*a*b^5*c^9*d)) + \\
& (x*(-c^5*d)^{(1/2)}*(3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d))*(256*b^9*c^11*d^2 \\
& - 1280*a*b^8*c^10*d^3 + 2304*a^2*b^7*c^9*d^4 - 1280*a^3*b^6*c^8*d^5 - 1280* \\
& a^4*b^5*c^7*d^6 + 2304*a^5*b^4*c^6*d^7 - 1280*a^6*b^3*c^5*d^8 + 256*a^7*b^2 \\
& *c^4*d^9))/(512*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d))* \\
& (b^4*c^8 + a^4*c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^2*b^2*c^6*d^2 - 4*a*b^3*c^7*d \\
&)))*(-c^5*d)^{(1/2)}*(3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d))/(16*(b^3*c^8 - a^ \\
& 3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d))*(-c^5*d)^{(1/2)}*(3*a^2*d^2 + \\
& 15*b^2*c^2 - 10*a*b*c*d)*1i)/(16*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - \\
& 3*a*b^2*c^7*d)))/((9*a^3*b^5*d^6 - 105*b^8*c^3*d^3 + 115*a*b^7*c^2*d^4 - 5 \\
& 1*a^2*b^6*c*d^5)/(32*(b^6*c^10 + a^6*c^4*d^6 - 6*a^5*b*c^5*d^5 + 15*a^2*b^4 \\
& *c^8*d^2 - 20*a^3*b^3*c^7*d^3 + 15*a^4*b^2*c^6*d^4 - 6*a*b^5*c^9*d)) + (((x \\
& *(9*a^4*b^3*d^7 + 289*b^7*c^4*d^3 - 300*a*b^6*c^3*d^4 - 60*a^3*b^4*c*d^6 + \\
& 190*a^2*b^5*c^2*d^5))/(32*(b^4*c^8 + a^4*c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^2* \\
& b^2*c^6*d^2 - 4*a*b^3*c^7*d)) - (((256*b^10*c^10*d^2 - 1760*a*b^9*c^9*d^3 + \\
& 5280*a^2*b^8*c^8*d^4 - 9056*a^3*b^7*c^7*d^5 + 9760*a^4*b^6*c^6*d^6 - 6816* \\
& a^5*b^5*c^5*d^7 + 3040*a^6*b^4*c^4*d^8 - 800*a^7*b^3*c^3*d^9 + 96*a^8*b^2*c^ \\
& ^2*d^10)/(64*(b^6*c^10 + a^6*c^4*d^6 - 6*a^5*b*c^5*d^5 + 15*a^2*b^4*c^8*d^2 \\
& - 20*a^3*b^3*c^7*d^3 + 15*a^4*b^2*c^6*d^4 - 6*a*b^5*c^9*d)) - (x*(-c^5*d)^ \\
& (1/2)**(3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d))*(256*b^9*c^11*d^2 - 1280*a*b^8* \\
& c^10*d^3 + 2304*a^2*b^7*c^9*d^4 - 1280*a^3*b^6*c^8*d^5 - 1280*a^4*b^5*c^7*d \\
& ^6 + 2304*a^5*b^4*c^6*d^7 - 1280*a^6*b^3*c^5*d^8 + 256*a^7*b^2*c^4*d^9))/(5 \\
& 12*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d))*(b^4*c^8 + a^4 \\
& *c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^2*b^2*c^6*d^2 - 4*a*b^3*c^7*d))*(-c^5*d)^
\end{aligned}$$

$$\begin{aligned} & \left(\frac{1}{2} \right) * (3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d) / (16*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d)) * (-c^5*d)^{(1/2)} * (3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d) / (16*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d)) \\ & - \left(\left(x*(9*a^4*b^3*d^7 + 289*b^7*c^4*d^3 - 300*a*b^6*c^3*d^4 - 60*a^3*b^4*c*d^6 + 190*a^2*b^5*c^2*d^5) \right) / (32*(b^4*c^8 + a^4*c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^2*b^2*c^6*d^2 - 4*a*b^3*c^7*d)) \right) + \left(\left((256*b^10*c^10*d^2 - 1760*a*b^9*c^9*d^3 + 5280*a^2*b^8*c^8*d^4 - 9056*a^3*b^7*c^7*d^5 + 9760*a^4*b^6*c^6*d^6 - 6816*a^5*b^5*c^5*d^7 + 3040*a^6*b^4*c^4*d^8 - 800*a^7*b^3*c^3*d^9 + 96*a^8*b^2*c^2*d^10) / (64*(b^6*c^10 + a^6*c^4*d^6 - 6*a^5*b*c^5*d^5 + 15*a^2*b^4*c^8*d^2 - 20*a^3*b^3*c^7*d^3 + 15*a^4*b^2*c^6*d^4 - 6*a*b^5*c^9*d) \right) \right) + (x*(-c^5*d)^{(1/2)} * (3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d) * (256*b^9*c^11*d^2 - 1280*a*b^8*c^10*d^3 + 2304*a^2*b^7*c^9*d^4 - 1280*a^3*b^6*c^8*d^5 - 1280*a^4*b^5*c^7*d^6 + 2304*a^5*b^4*c^6*d^7 - 1280*a^6*b^3*c^5*d^8 + 256*a^7*b^2*c^4*d^9)) / (512*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d)) * (b^4*c^8 + a^4*c^4*d^4 - 4*a^3*b*c^5*d^3 + 6*a^2*b^2*c^6*d^2 - 4*a*b^3*c^7*d)) * (-c^5*d)^{(1/2)} * (3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d) / (16*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d)) * (-c^5*d)^{(1/2)} * (3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d) / (16*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d)) * (-c^5*d)^{(1/2)} * (3*a^2*d^2 + 15*b^2*c^2 - 10*a*b*c*d) * 1i / (8*(b^3*c^8 - a^3*c^5*d^3 + 3*a^2*b*c^6*d^2 - 3*a*b^2*c^7*d)) \end{aligned}$$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)/(d*x**2+c)**3,x)

[Out] Timed out

$$3.27 \quad \int \frac{(c+dx^2)^5}{(a+bx^2)^2} dx$$

Optimal. Leaf size=192

$$\frac{(bc-ad)^4(9ad+bc)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}b^{11/2}} + \frac{d^3x^3(3a^2d^2-10abcd+10b^2c^2)}{3b^4} + \frac{d^2x(-4a^3d^3+15a^2bcd^2-20ab^2c^2d+10b^3c^3)}{b^5}$$

[Out] $d^2*(-4*a^3*d^3+15*a^2*b*c*d^2-20*a*b^2*c^2*d+10*b^3*c^3)*x/b^5+1/3*d^3*(3*a^2*d^2-10*a*b*c*d+10*b^2*c^2)*x^3/b^4+1/5*d^4*(-2*a*d+5*b*c)*x^5/b^3+1/7*d^5*x^7/b^2+1/2*(-a*d+b*c)^5*x/a/b^5/(b*x^2+a)+1/2*(-a*d+b*c)^4*(9*a*d+b*c)*\arctan(x*b^{(1/2)}/a^{(1/2)})/a^{(3/2)}/b^{(11/2)}$

Rubi [A] time = 0.16, antiderivative size = 192, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {390, 385, 205}

$$\frac{d^3x^3(3a^2d^2-10abcd+10b^2c^2)}{3b^4} + \frac{d^2x(15a^2bcd^2-4a^3d^3-20ab^2c^2d+10b^3c^3)}{b^5} + \frac{(bc-ad)^4(9ad+bc)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}b^{11/2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^5/(a + b*x^2)^2,x]

[Out] $(d^2*(10*b^3*c^3-20*a*b^2*c^2*d+15*a^2*b*c*d^2-4*a^3*d^3)*x)/b^5+(d^3*(10*b^2*c^2-10*a*b*c*d+3*a^2*d^2)*x^3)/(3*b^4)+(d^4*(5*b*c-2*a*d)*x^5)/(5*b^3)+(d^5*x^7)/(7*b^2)+((b*c-a*d)^5*x)/(2*a*b^5*(a+b*x^2))+((b*c-a*d)^4*(b*c+9*a*d)*\text{ArcTan}[\text{Sqrt}[b]*x/\text{Sqrt}[a]])/(2*a^{(3/2)}*b^{(11/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\int \frac{(c + dx^2)^5}{(a + bx^2)^2} dx = \int \left(\frac{d^2 (10b^3c^3 - 20ab^2c^2d + 15a^2bcd^2 - 4a^3d^3)}{b^5} + \frac{d^3 (10b^2c^2 - 10abcd + 3a^2d^2)x^2}{b^4} + \frac{d^4 (5b^2c^2 - 5abcd + 2a^2d^2)x^4}{3b^3} \right) dx$$

$$= \frac{d^2 (10b^3c^3 - 20ab^2c^2d + 15a^2bcd^2 - 4a^3d^3)x}{b^5} + \frac{d^3 (10b^2c^2 - 10abcd + 3a^2d^2)x^3}{3b^4} + \frac{d^4 (5b^2c^2 - 5abcd + 2a^2d^2)x^5}{15b^3}$$

$$= \frac{d^2 (10b^3c^3 - 20ab^2c^2d + 15a^2bcd^2 - 4a^3d^3)x}{b^5} + \frac{d^3 (10b^2c^2 - 10abcd + 3a^2d^2)x^3}{3b^4} + \frac{d^4 (5b^2c^2 - 5abcd + 2a^2d^2)x^5}{15b^3}$$

$$= \frac{d^2 (10b^3c^3 - 20ab^2c^2d + 15a^2bcd^2 - 4a^3d^3)x}{b^5} + \frac{d^3 (10b^2c^2 - 10abcd + 3a^2d^2)x^3}{3b^4} + \frac{d^4 (5b^2c^2 - 5abcd + 2a^2d^2)x^5}{15b^3}$$

Mathematica [A] time = 0.10, size = 192, normalized size = 1.00

$$\frac{(bc - ad)^4(9ad + bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{11/2}} + \frac{d^3x^3(3a^2d^2 - 10abcd + 10b^2c^2)}{3b^4} + \frac{d^2x(-4a^3d^3 + 15a^2bcd^2 - 20ab^2c^2d + 10b^3c^3)}{b^5}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^5/(a + b*x^2)^2,x]

[Out] (d^2*(10*b^3*c^3 - 20*a*b^2*c^2*d + 15*a^2*b*c*d^2 - 4*a^3*d^3)*x)/b^5 + (d^3*(10*b^2*c^2 - 10*a*b*c*d + 3*a^2*d^2)*x^3)/(3*b^4) + (d^4*(5*b^2*c^2 - 5*a*b*c*d + 2*a^2*d^2)*x^5)/(15*b^3) + (d^5*x^7)/(7*b^2) + ((b*c - a*d)^5*x)/(2*a*b^5*(a + b*x^2)) + ((b*c - a*d)^4*(b*c + 9*a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(2*a^(3/2)*b^(11/2))

fricas [B] time = 0.55, size = 810, normalized size = 4.22

$$\frac{60 a^2 b^5 d^5 x^9 + 12 (35 a^2 b^5 c d^4 - 9 a^3 b^4 d^5) x^7 + 28 (50 a^2 b^5 c^2 d^3 - 35 a^3 b^4 c d^4 + 9 a^4 b^3 d^5) x^5 + 140 (30 a^2 b^5 c^3 d^2 - 50 a^3 b^4 c^2 d^3 + 35 a^4 b^3 c d^4 - 9 a^5 b^2 d^5) x^3 - 105 (a b^5 c^5 + 5 a^2 b^4 c^4 d - 30 a^3 b^3 c^3 d^2 + 50 a^4 b^2 c^2 d^3 - 35 a^5 b c d^4 + 9 a^6 d^5 + (b^6 c^5 + 5 a b^5 c^4 d - 30 a^2 b^4 c^3 d^2 + 50 a^3 b^3 c^2 d^3 - 35 a^4 b^2 c d^4 + 9 a^5 b d^5) x^2) \sqrt{-a b} \log\left(\frac{(b x^2 - 2 \sqrt{-a b}) x - a}{(b x^2 + a)}\right) + 210 (a b^6 c^5 - 5 a^2 b^5 c^4 d + 30 a^3 b^4 c^3 d^2 - 50 a^4 b^3 c^2 d^3 + 35 a^5 b^2 c d^4 - 9 a^6 b d^5) x}{(a^2 b^7 x^2 + a^3 b^6)} + \frac{1}{210} (30 a^2 b^5 d^5 x^9 + 6 (35 a^2 b^5 c d^4 - 9 a^3 b^4 d^5) x^7 + 14 (50 a^2 b^5 c^2 d^3 - 35 a^3 b^4 c d^4 + 9 a^4 b^3 d^5) x^5 + 70 (30 a^2 b^5 c^3 d^2 - 50 a^3 b^4 c^2 d^3 + 35 a^4 b^3 c d^4 - 9 a^5 b^2 d^5) x^3 + 105 (a b^5 c^5 + 5 a^2 b^4 c^4 d - 30 a^3 b^3 c^3 d^2 + 50 a^4 b^2 c^2 d^3 - 35 a^5 b c d^4 + 9 a^6 d^5 + (b^6 c^5 + 5 a b^5 c^4 d - 30 a^2 b^4 c^3 d^2 + 50 a^3 b^3 c^2 d^3 - 35 a^4 b^2 c d^4 + 9 a^5 b d^5) x^2) \sqrt{a b} \arctan\left(\frac{\sqrt{a b} x}{a}\right) + 105 (a b^6 c^5 - 5 a^2 b^5 c^4 d + 30 a^3 b^4 c^3 d^2 - 50 a^4 b^3 c^2 d^3 + 35 a^5 b^2 c d^4 - 9 a^6 b d^5) x}{(a^2 b^7 x^2 + a^3 b^6)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^5/(b*x^2+a)^2,x, algorithm="fricas")

[Out] [1/420*(60*a^2*b^5*d^5*x^9 + 12*(35*a^2*b^5*c*d^4 - 9*a^3*b^4*d^5)*x^7 + 28*(50*a^2*b^5*c^2*d^3 - 35*a^3*b^4*c*d^4 + 9*a^4*b^3*d^5)*x^5 + 140*(30*a^2*b^5*c^3*d^2 - 50*a^3*b^4*c^2*d^3 + 35*a^4*b^3*c*d^4 - 9*a^5*b^2*d^5)*x^3 - 105*(a*b^5*c^5 + 5*a^2*b^4*c^4*d - 30*a^3*b^3*c^3*d^2 + 50*a^4*b^2*c^2*d^3 - 35*a^5*b*c*d^4 + 9*a^6*d^5 + (b^6*c^5 + 5*a*b^5*c^4*d - 30*a^2*b^4*c^3*d^2 + 50*a^3*b^3*c^2*d^3 - 35*a^4*b^2*c*d^4 + 9*a^5*b*d^5)*x^2)*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b))*x - a)/(b*x^2 + a) + 210*(a*b^6*c^5 - 5*a^2*b^5*c^4*d + 30*a^3*b^4*c^3*d^2 - 50*a^4*b^3*c^2*d^3 + 35*a^5*b^2*c*d^4 - 9*a^6*b*d^5)*x)/(a^2*b^7*x^2 + a^3*b^6), 1/210*(30*a^2*b^5*d^5*x^9 + 6*(35*a^2*b^5*c*d^4 - 9*a^3*b^4*d^5)*x^7 + 14*(50*a^2*b^5*c^2*d^3 - 35*a^3*b^4*c*d^4 + 9*a^4*b^3*d^5)*x^5 + 70*(30*a^2*b^5*c^3*d^2 - 50*a^3*b^4*c^2*d^3 + 35*a^4*b^3*c*d^4 - 9*a^5*b^2*d^5)*x^3 + 105*(a*b^5*c^5 + 5*a^2*b^4*c^4*d - 30*a^3*b^3*c^3*d^2 + 50*a^4*b^2*c^2*d^3 - 35*a^5*b*c*d^4 + 9*a^6*d^5 + (b^6*c^5 + 5*a*b^5*c^4*d - 30*a^2*b^4*c^3*d^2 + 50*a^3*b^3*c^2*d^3 - 35*a^4*b^2*c*d^4 + 9*a^5*b*d^5)*x^2)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + 105*(a*b^6*c^5 - 5*a^2*b^5*c^4*d + 30*a^3*b^4*c^3*d^2 - 50*a^4*b^3*c^2*d^3 + 35*a^5*b^2*c*d^4 - 9*a^6*b*d^5)*x)/(a^2*b^7*x^2 + a^3*b^6)]

giac [A] time = 0.58, size = 306, normalized size = 1.59

$$\frac{(b^5c^5 + 5ab^4c^4d - 30a^2b^3c^3d^2 + 50a^3b^2c^2d^3 - 35a^4bcd^4 + 9a^5d^5) \arctan\left(\frac{bx}{\sqrt{ab}}\right) + b^5c^5x - 5ab^4c^4dx + 10a^2b^3c^3d^2}{2\sqrt{ab}ab^5} + \frac{b^5c^5x - 5ab^4c^4dx + 10a^2b^3c^3d^2}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^5/(b*x^2+a)^2,x, algorithm="giac")

[Out] 1/2*(b^5*c^5 + 5*a*b^4*c^4*d - 30*a^2*b^3*c^3*d^2 + 50*a^3*b^2*c^2*d^3 - 35*a^4*b*c*d^4 + 9*a^5*d^5)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a*b^5) + 1/2*(b^5*c^5*x - 5*a*b^4*c^4*d*x + 10*a^2*b^3*c^3*d^2*x - 10*a^3*b^2*c^2*d^3*x + 5*a^4*b*c*d^4*x - a^5*d^5*x)/((b*x^2 + a)*a*b^5) + 1/105*(15*b^12*d^5*x^7 + 105*b^12*c*d^4*x^5 - 42*a*b^11*d^5*x^5 + 350*b^12*c^2*d^3*x^3 - 350*a*b^11*c*d^4*x^3 + 105*a^2*b^10*d^5*x^3 + 1050*b^12*c^3*d^2*x - 2100*a*b^11*c^2*d^3*x + 1575*a^2*b^10*c*d^4*x - 420*a^3*b^9*d^5*x)/b^14

maple [B] time = 0.01, size = 402, normalized size = 2.09

$$\frac{d^5x^7}{7b^2} - \frac{2ad^5x^5}{5b^3} + \frac{cd^4x^5}{b^2} + \frac{a^2d^5x^3}{b^4} - \frac{10acd^4x^3}{3b^3} + \frac{10c^2d^3x^3}{3b^2} - \frac{a^4d^5x}{2(bx^2+a)b^5} + \frac{9a^4d^5 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab}b^5} + \frac{5a^3cd^4x}{2(bx^2+a)b^4} - \frac{35a^5d^5}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^5/(b*x^2+a)^2,x)

[Out] 1/7*d^5*x^7/b^2-2/5*d^5/b^3*x^5*a+d^4/b^2*x^5*c+d^5/b^4*x^3*a^2-10/3*d^4/b^3*x^3*a*c+10/3*d^3/b^2*x^3*c^2-4*d^5/b^5*a^3*x+15*d^4/b^4*a^2*c*x-20*d^3/b^3*a*c^2*x+10*d^2/b^2*c^3*x-1/2/b^5*a^4*x/(b*x^2+a)*d^5+5/2/b^4*a^3*x/(b*x^2+a)*c*d^4-5/b^3*a^2*x/(b*x^2+a)*c^2*d^3+5/b^2*a*x/(b*x^2+a)*c^3*d^2-5/2/b*x/(b*x^2+a)*c^4*d+1/2/a*x/(b*x^2+a)*c^5+9/2/b^5*a^4/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*d^5-35/2/b^4*a^3/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c*d^4+25/b^3*a^2/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c^2*d^3-15/b^2*a/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c^3*d^2+5/2/b/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c^4*d+1/2/a/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c^5

maxima [A] time = 3.03, size = 294, normalized size = 1.53

$$\frac{(b^5c^5 - 5ab^4c^4d + 10a^2b^3c^3d^2 - 10a^3b^2c^2d^3 + 5a^4bcd^4 - a^5d^5)x}{2(ab^6x^2 + a^2b^5)} + \frac{15b^3d^5x^7 + 21(5b^3cd^4 - 2ab^2d^5)x^5 + 35(10a^2b^3c^3d^2 - 10a^3b^2c^2d^3 + 5a^4bcd^4 - a^5d^5)}{2(ab^6x^2 + a^2b^5)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^5/(b*x^2+a)^2,x, algorithm="maxima")

[Out] 1/2*(b^5*c^5 - 5*a*b^4*c^4*d + 10*a^2*b^3*c^3*d^2 - 10*a^3*b^2*c^2*d^3 + 5*a^4*b*c*d^4 - a^5*d^5)*x/(a*b^6*x^2 + a^2*b^5) + 1/105*(15*b^3*d^5*x^7 + 21*(5*b^3*c*d^4 - 2*a*b^2*d^5)*x^5 + 35*(10*b^3*c^2*d^3 - 10*a*b^2*c*d^4 + 3*a^2*b*d^5)*x^3 + 105*(10*b^3*c^3*d^2 - 20*a*b^2*c^2*d^3 + 15*a^2*b*c*d^4 - 4*a^3*d^5)*x)/b^5 + 1/2*(b^5*c^5 + 5*a*b^4*c^4*d - 30*a^2*b^3*c^3*d^2 + 50*a^3*b^2*c^2*d^3 - 35*a^4*b*c*d^4 + 9*a^5*d^5)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a*b^5)

mupad [B] time = 5.02, size = 386, normalized size = 2.01

$$x \left(\frac{10c^3d^2}{b^2} - \frac{2a \left(\frac{2ad^5}{b^3} - \frac{5cd^4}{b^2} \right) - \frac{a^2d^5}{b^4} + \frac{10c^2d^3}{b^2}}{b} + \frac{a^2 \left(\frac{2ad^5}{b^3} - \frac{5cd^4}{b^2} \right)}{b^2} \right) - x^5 \left(\frac{2ad^5}{5b^3} - \frac{cd^4}{b^2} \right) + x^3 \left(\frac{2a \left(\frac{2ad^5}{b^3} - \frac{5cd^4}{b^2} \right)}{3b} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^5/(a + b*x^2)^2,x)`

[Out] $x \left(\frac{10c^3d^2}{b^2} - \frac{2a \left(\frac{2a \left(\frac{2ad^5}{b^3} - \frac{5cd^4}{b^2} \right)}{b} - \frac{a^2d^5}{b^4} + \frac{10c^2d^3}{b^2} \right)}{b} + \frac{a^2 \left(\frac{2ad^5}{b^3} - \frac{5cd^4}{b^2} \right)}{b^2} \right) - x^5 \left(\frac{2ad^5}{5b^3} - \frac{cd^4}{b^2} \right) + x^3 \left(\frac{2a \left(\frac{2ad^5}{b^3} - \frac{5cd^4}{b^2} \right)}{3b} - \frac{a^2d^5}{3b^4} + \frac{10c^2d^3}{3b^2} \right) + \frac{d^5x^7}{7b^2} - \frac{(x(a^5d^5 - b^5c^5 - 10a^2b^3c^3d^2 + 10a^3b^2c^2d^3 + 5ab^4c^4d - 5a^4b^2cd^4))}{2a(ab^5 + b^6x^2)} + \frac{\operatorname{atan}\left(\frac{b^{1/2}x(ad - bc)^4(9ad + bc)}{a^{1/2}(9a^5d^5 + b^5c^5 - 30a^2b^3c^3d^2 + 50a^3b^2c^2d^3 + 5ab^4c^4d - 35a^4b^2cd^4)}\right)}{(2a^{3/2}b^{11/2})}$

sympy [B] time = 1.91, size = 502, normalized size = 2.61

$$x^5 \left(-\frac{2ad^5}{5b^3} + \frac{cd^4}{b^2} \right) + x^3 \left(\frac{a^2d^5}{b^4} - \frac{10acd^4}{3b^3} + \frac{10c^2d^3}{3b^2} \right) + x \left(-\frac{4a^3d^5}{b^5} + \frac{15a^2cd^4}{b^4} - \frac{20ac^2d^3}{b^3} + \frac{10c^3d^2}{b^2} \right) + \frac{x(-a^5d^5 + 5a^4b^2cd^4 - 10a^3b^3c^3d^2 + 10a^2b^4c^4d - 5ab^5c^5)}{2a(ab^5 + b^6x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**5/(b*x**2+a)**2,x)`

[Out] $x^5 \left(-\frac{2ad^5}{5b^3} + \frac{cd^4}{b^2} \right) + x^3 \left(\frac{a^2d^5}{b^4} - \frac{10acd^4}{3b^3} + \frac{10c^2d^3}{3b^2} \right) + x \left(-\frac{4a^3d^5}{b^5} + \frac{15a^2cd^4}{b^4} - \frac{20ac^2d^3}{b^3} + \frac{10c^3d^2}{b^2} \right) + \frac{x(-a^5d^5 + 5a^4b^2cd^4 - 10a^3b^3c^3d^2 + 10a^2b^4c^4d - 5ab^5c^5)}{(2a^2b^5 + 2ab^6x^2) - \sqrt{-1/(a^3b^{11})}} \frac{(ad - bc)^4(9ad + bc) \log(-a^2b^5\sqrt{-1/(a^3b^{11})})(ad - bc)^4(9ad + bc)}{(9a^5d^5 - 35a^4b^2cd^4 + 50a^3b^3c^3d^2 - 30a^2b^4c^4d + b^5c^5) + x}{4} + \sqrt{-1/(a^3b^{11})} \frac{(ad - bc)^4(9ad + bc) \log(a^2b^5\sqrt{-1/(a^3b^{11})})(ad - bc)^4(9ad + bc)}{(9a^5d^5 - 35a^4b^2cd^4 + 50a^3b^3c^3d^2 - 30a^2b^4c^4d + b^5c^5) + x}{4} + \frac{d^5x^7}{7b^2}$

$$3.28 \quad \int \frac{(c+dx^2)^4}{(a+bx^2)^2} dx$$

Optimal. Leaf size=142

$$\frac{(bc-ad)^3(7ad+bc)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}b^{9/2}} + \frac{d^2x(3a^2d^2-8abcd+6b^2c^2)}{b^4} + \frac{x(bc-ad)^4}{2ab^4(a+bx^2)} + \frac{2d^3x^3(2bc-ad)}{3b^3} + \frac{d^4x^5}{5b^2}$$

[Out] $d^2*(3*a^2*d^2-8*a*b*c*d+6*b^2*c^2)*x/b^4+2/3*d^3*(-a*d+2*b*c)*x^3/b^3+1/5*d^4*x^5/b^2+1/2*(-a*d+b*c)^4*x/a/b^4/(b*x^2+a)+1/2*(-a*d+b*c)^3*(7*a*d+b*c)*\arctan(x*b^(1/2)/a^(1/2))/a^(3/2)/b^(9/2)$

Rubi [A] time = 0.12, antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {390, 385, 205}

$$\frac{d^2x(3a^2d^2-8abcd+6b^2c^2)}{b^4} + \frac{(bc-ad)^3(7ad+bc)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}b^{9/2}} + \frac{2d^3x^3(2bc-ad)}{3b^3} + \frac{x(bc-ad)^4}{2ab^4(a+bx^2)} + \frac{d^4x^5}{5b^2}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^4/(a + b*x^2)^2, x]

[Out] $(d^2*(6*b^2*c^2-8*a*b*c*d+3*a^2*d^2)*x)/b^4+(2*d^3*(2*b*c-a*d)*x^3)/(3*b^3)+(d^4*x^5)/(5*b^2)+((b*c-a*d)^4*x)/(2*a*b^4*(a+b*x^2))+((b*c-a*d)^3*(b*c+7*a*d)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]])/(2*a^(3/2)*b^(9/2))$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p+1))/(a*b*n*(p+1)), x] - Dist[(a*d - b*c*(n*(p+1) + 1))/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned}
\int \frac{(c + dx^2)^4}{(a + bx^2)^2} dx &= \int \left(\frac{d^2(6b^2c^2 - 8abcd + 3a^2d^2)}{b^4} + \frac{2d^3(2bc - ad)x^2}{b^3} + \frac{d^4x^4}{b^2} + \frac{(bc - ad)^3(bc + 3ad) + 4bd(bc - ad)^2}{b^4(a + bx^2)^2} \right) dx \\
&= \frac{d^2(6b^2c^2 - 8abcd + 3a^2d^2)x}{b^4} + \frac{2d^3(2bc - ad)x^3}{3b^3} + \frac{d^4x^5}{5b^2} + \frac{\int \frac{(bc - ad)^3(bc + 3ad) + 4bd(bc - ad)^2}{(a + bx^2)^2} dx}{b^4} \\
&= \frac{d^2(6b^2c^2 - 8abcd + 3a^2d^2)x}{b^4} + \frac{2d^3(2bc - ad)x^3}{3b^3} + \frac{d^4x^5}{5b^2} + \frac{(bc - ad)^4x}{2ab^4(a + bx^2)} + \frac{(bc - ad)^3(bc + 3ad)}{2ab^4(a + bx^2)} \\
&= \frac{d^2(6b^2c^2 - 8abcd + 3a^2d^2)x}{b^4} + \frac{2d^3(2bc - ad)x^3}{3b^3} + \frac{d^4x^5}{5b^2} + \frac{(bc - ad)^4x}{2ab^4(a + bx^2)} + \frac{(bc - ad)^3(bc + 3ad)}{2ab^4(a + bx^2)}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 142, normalized size = 1.00

$$\frac{(bc - ad)^3(7ad + bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{9/2}} + \frac{d^2x(3a^2d^2 - 8abcd + 6b^2c^2)}{b^4} + \frac{x(bc - ad)^4}{2ab^4(a + bx^2)} + \frac{2d^3x^3(2bc - ad)}{3b^3} + \frac{d^4x^5}{5b^2}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^4/(a + b*x^2)^2,x]

[Out] (d^2*(6*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*x)/b^4 + (2*d^3*(2*b*c - a*d)*x^3)/(3*b^3) + (d^4*x^5)/(5*b^2) + ((b*c - a*d)^4*x)/(2*a*b^4*(a + b*x^2)) + ((b*c - a*d)^3*(b*c + 7*a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(2*a^(3/2)*b^(9/2))

fricas [B] time = 0.68, size = 612, normalized size = 4.31

$$\frac{12a^2b^4d^4x^7 + 4(20a^2b^4cd^3 - 7a^3b^3d^4)x^5 + 20(18a^2b^4c^2d^2 - 20a^3b^3cd^3 + 7a^4b^2d^4)x^3 + 15(ab^4c^4 + 4a^2b^3cd^3 - 18a^3b^2c^2d^2 + 20a^4b^2cd^3 - 7a^5b^2d^4)x + 15(ab^4c^4 + 4a^2b^3cd^3 - 18a^3b^2c^2d^2 + 20a^4b^2cd^3 - 7a^5b^2d^4)}{2\sqrt{ab}ab^4} + \frac{b^4c^4x - 4ab^3c^3dx + 6a^2b^2c^2d^2x - 4a^3bcd^3}{2(bx^2 + a)ab^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^2,x, algorithm="fricas")

[Out] [1/60*(12*a^2*b^4*d^4*x^7 + 4*(20*a^2*b^4*c*d^3 - 7*a^3*b^3*d^4)*x^5 + 20*(18*a^2*b^4*c^2*d^2 - 20*a^3*b^3*c*d^3 + 7*a^4*b^2*d^4)*x^3 + 15*(a*b^4*c^4 + 4*a^2*b^3*c^3*d - 18*a^3*b^2*c^2*d^2 + 20*a^4*b^2*c*d^3 - 7*a^5*d^4 + (b^5*c^4 + 4*a*b^4*c^3*d - 18*a^2*b^3*c^2*d^2 + 20*a^3*b^2*c*d^3 - 7*a^4*b*d^4)*x^2)*sqrt(-a*b)*log((b*x^2 + 2*sqrt(-a*b)*x - a)/(b*x^2 + a)) + 30*(a*b^5*c^4 - 4*a^2*b^4*c^3*d + 18*a^3*b^3*c^2*d^2 - 20*a^4*b^2*c*d^3 + 7*a^5*b*d^4)*x)/(a^2*b^6*x^2 + a^3*b^5), 1/30*(6*a^2*b^4*d^4*x^7 + 2*(20*a^2*b^4*c*d^3 - 7*a^3*b^3*d^4)*x^5 + 10*(18*a^2*b^4*c^2*d^2 - 20*a^3*b^3*c*d^3 + 7*a^4*b^2*d^4)*x^3 + 15*(a*b^4*c^4 + 4*a^2*b^3*c^3*d - 18*a^3*b^2*c^2*d^2 + 20*a^4*b^2*c*d^3 - 7*a^5*d^4 + (b^5*c^4 + 4*a*b^4*c^3*d - 18*a^2*b^3*c^2*d^2 + 20*a^3*b^2*c*d^3 - 7*a^4*b*d^4)*x^2)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + 15*(a*b^5*c^4 - 4*a^2*b^4*c^3*d + 18*a^3*b^3*c^2*d^2 - 20*a^4*b^2*c*d^3 + 7*a^5*b*d^4)*x)/(a^2*b^6*x^2 + a^3*b^5)]

giac [A] time = 0.58, size = 220, normalized size = 1.55

$$\frac{(b^4c^4 + 4ab^3c^3d - 18a^2b^2c^2d^2 + 20a^3bcd^3 - 7a^4d^4) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab}ab^4} + \frac{b^4c^4x - 4ab^3c^3dx + 6a^2b^2c^2d^2x - 4a^3bcd^3}{2(bx^2 + a)ab^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^2,x, algorithm="giac")

[Out] $\frac{1}{2}*(b^4*c^4 + 4*a*b^3*c^3*d - 18*a^2*b^2*c^2*d^2 + 20*a^3*b*c*d^3 - 7*a^4*d^4)*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b}*a*b^4) + \frac{1}{2}*(b^4*c^4*x - 4*a*b^3*c^3*d*x + 6*a^2*b^2*c^2*d^2*x - 4*a^3*b*c*d^3*x + a^4*d^4*x)/((b*x^2 + a)*a*b^4) + \frac{1}{15}*(3*b^8*d^4*x^5 + 20*b^8*c*d^3*x^3 - 10*a*b^7*d^4*x^3 + 90*b^8*c^2*d^2*x - 120*a*b^7*c*d^3*x + 45*a^2*b^6*d^4*x)/b^{10}$

maple [B] time = 0.01, size = 296, normalized size = 2.08

$$\frac{d^4x^5}{5b^2} - \frac{2ad^4x^3}{3b^3} + \frac{4cd^3x^3}{3b^2} + \frac{a^3d^4x}{2(bx^2+a)b^4} - \frac{7a^3d^4 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab}b^4} - \frac{2a^2cd^3x}{(bx^2+a)b^3} + \frac{10a^2cd^3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}b^3} + \frac{3a^2d^2x}{(bx^2+a)b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^4/(b*x^2+a)^2,x)

[Out] $\frac{1}{5}d^4x^5/b^2 - \frac{2}{3}d^4/b^3x^3/a + \frac{4}{3}d^3/b^2x^3/c + \frac{3d^4/b^4a^2x - 8d^3/b^3a^2cx + 6d^2/b^2c^2x + 1/2/b^4a^3x}{(b*x^2+a)*d^4} - \frac{2/b^3a^2x}{(b*x^2+a)*c*d^3} + \frac{3/b^2ax}{(b*x^2+a)*c^2*d^2} - \frac{2/b*x}{(b*x^2+a)*c^3*d} + \frac{1/2/a*x}{(b*x^2+a)*c^4} - \frac{7/2/b^4a^3}{(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*d^4} + \frac{10/b^3a^2}{(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c*d^3} - \frac{9/b^2a}{(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c^2*d^2} + \frac{2/b}{(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c^3*d} + \frac{1/2/a}{(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c^4}$

maxima [A] time = 3.00, size = 213, normalized size = 1.50

$$\frac{(b^4c^4 - 4ab^3c^3d + 6a^2b^2c^2d^2 - 4a^3bcd^3 + a^4d^4)x}{2(ab^5x^2 + a^2b^4)} + \frac{3b^2d^4x^5 + 10(2b^2cd^3 - abd^4)x^3 + 15(6b^2c^2d^2 - 8abcd^3 + 3a^2d^4)}{15b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^2,x, algorithm="maxima")

[Out] $\frac{1}{2}*(b^4*c^4 - 4*a*b^3*c^3*d + 6*a^2*b^2*c^2*d^2 - 4*a^3*b*c*d^3 + a^4*d^4)*x/(a*b^5*x^2 + a^2*b^4) + \frac{1}{15}*(3*b^8*d^4*x^5 + 10*(2*b^2*c*d^3 - a*b*d^4)*x^3 + 15*(6*b^2*c^2*d^2 - 8*a*b*c*d^3 + 3*a^2*d^4)*x)/b^4 + \frac{1}{2}*(b^4*c^4 + 4*a*b^3*c^3*d - 18*a^2*b^2*c^2*d^2 + 20*a^3*b*c*d^3 - 7*a^4*d^4)*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b}*a*b^4)$

mupad [B] time = 5.05, size = 261, normalized size = 1.84

$$x \left(\frac{2a \left(\frac{2ad^4}{b^3} - \frac{4cd^3}{b^2} \right)}{b} - \frac{a^2d^4}{b^4} + \frac{6c^2d^2}{b^2} \right) - x^3 \left(\frac{2ad^4}{3b^3} - \frac{4cd^3}{3b^2} \right) + \frac{d^4x^5}{5b^2} + \frac{x(a^4d^4 - 4a^3bcd^3 + 6a^2b^2c^2d^2 - 4ab^3c^3d)}{2a(b^5x^2 + ab^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^4/(a + b*x^2)^2,x)

[Out] $x*((2*a*((2*a*d^4)/b^3 - (4*c*d^3)/b^2))/b - (a^2*d^4)/b^4 + (6*c^2*d^2)/b^2) - x^3*((2*a*d^4)/(3*b^3) - (4*c*d^3)/(3*b^2)) + (d^4*x^5)/(5*b^2) + (x*(a^4*d^4 + b^4*c^4 + 6*a^2*b^2*c^2*d^2 - 4*a*b^3*c^3*d - 4*a^3*b*c*d^3))/(2*a*(a*b^4 + b^5*x^2)) + (atan((b^(1/2))*x*(a*d - b*c))^3*(7*a*d + b*c))/(a^(1/2)*(b^4*c^4 - 7*a^4*d^4 - 18*a^2*b^2*c^2*d^2 + 4*a*b^3*c^3*d + 20*a^3*b*c*d^3))*(a*d - b*c)^3*(7*a*d + b*c))/(2*a^(3/2)*b^(9/2))$

sympy [B] time = 1.47, size = 403, normalized size = 2.84

$$x^3 \left(-\frac{2ad^4}{3b^3} + \frac{4cd^3}{3b^2} \right) + x \left(\frac{3a^2d^4}{b^4} - \frac{8acd^3}{b^3} + \frac{6c^2d^2}{b^2} \right) + \frac{x(a^4d^4 - 4a^3bcd^3 + 6a^2b^2c^2d^2 - 4ab^3c^3d + b^4c^4)}{2a^2b^4 + 2ab^5x^2} + \frac{\sqrt{-\frac{1}{a^3b^9}}}{1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**4/(b*x**2+a)**2,x)

[Out] x**3*(-2*a*d**4/(3*b**3) + 4*c*d**3/(3*b**2)) + x*(3*a**2*d**4/b**4 - 8*a*c*d**3/b**3 + 6*c**2*d**2/b**2) + x*(a**4*d**4 - 4*a**3*b*c*d**3 + 6*a**2*b**2*c**2*d**2 - 4*a*b**3*c**3*d + b**4*c**4)/(2*a**2*b**4 + 2*a*b**5*x**2) + sqrt(-1/(a**3*b**9))*(a*d - b*c)**3*(7*a*d + b*c)*log(-a**2*b**4*sqrt(-1/(a**3*b**9))*(a*d - b*c)**3*(7*a*d + b*c)/(7*a**4*d**4 - 20*a**3*b*c*d**3 + 18*a**2*b**2*c**2*d**2 - 4*a*b**3*c**3*d - b**4*c**4) + x)/4 - sqrt(-1/(a**3*b**9))*(a*d - b*c)**3*(7*a*d + b*c)*log(a**2*b**4*sqrt(-1/(a**3*b**9))*(a*d - b*c)**3*(7*a*d + b*c)/(7*a**4*d**4 - 20*a**3*b*c*d**3 + 18*a**2*b**2*c**2*d**2 - 4*a*b**3*c**3*d - b**4*c**4) + x)/4 + d**4*x**5/(5*b**2)

$$3.29 \quad \int \frac{(c+dx^2)^3}{(a+bx^2)^2} dx$$

Optimal. Leaf size=106

$$\frac{(5ad + bc)(bc - ad)^2 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{7/2}} + \frac{d^2x(3bc - 2ad)}{b^3} + \frac{x(bc - ad)^3}{2ab^3(a + bx^2)} + \frac{d^3x^3}{3b^2}$$

[Out] $d^2*(-2*a*d+3*b*c)*x/b^3+1/3*d^3*x^3/b^2+1/2*(-a*d+b*c)^3*x/a/b^3/(b*x^2+a)+1/2*(-a*d+b*c)^2*(5*a*d+b*c)*\arctan(x*b^{(1/2)}/a^{(1/2)})/a^{(3/2)}/b^{(7/2)}$

Rubi [A] time = 0.09, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {390, 385, 205}

$$\frac{(5ad + bc)(bc - ad)^2 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{7/2}} + \frac{d^2x(3bc - 2ad)}{b^3} + \frac{x(bc - ad)^3}{2ab^3(a + bx^2)} + \frac{d^3x^3}{3b^2}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^3/(a + b*x^2)^2,x]

[Out] $(d^2*(3*b*c - 2*a*d)*x)/b^3 + (d^3*x^3)/(3*b^2) + ((b*c - a*d)^3*x)/(2*a*b^3*(a + b*x^2)) + ((b*c - a*d)^2*(b*c + 5*a*d)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]])/(2*a^{(3/2)}*b^{(7/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

[Out] $\frac{1}{2}*(b^3*c^3 + 3*a*b^2*c^2*d - 9*a^2*b*c*d^2 + 5*a^3*d^3)*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b}*a*b^3) + \frac{1}{2}*(b^3*c^3*x - 3*a*b^2*c^2*d*x + 3*a^2*b*c*d^2*x - a^3*d^3*x)/((b*x^2 + a)*a*b^3) + \frac{1}{3}*(b^4*d^3*x^3 + 9*b^4*c*d^2*x - 6*a*b^3*d^3*x)/b^6$

maple [B] time = 0.01, size = 205, normalized size = 1.93

$$\frac{d^3 x^3}{3b^2} - \frac{a^2 d^3 x}{2(bx^2 + a)b^3} + \frac{5a^2 d^3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab} b^3} + \frac{3ac d^2 x}{2(bx^2 + a)b^2} - \frac{9ac d^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab} b^2} + \frac{c^3 x}{2(bx^2 + a)a} + \frac{c^3 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab} a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+c)^3/(b*x^2+a)^2,x)`

[Out] $\frac{1}{3}b^2d^3x^3 - 2d^3/b^3ax + 3/b^2c*d^2*x - 1/2/b^3a^2*x/(b*x^2+a)*d^3 + 3/2/b^2a*x/(b*x^2+a)*c*d^2 - 3/2/b*x/(b*x^2+a)*c^2*d + 1/2/a*x/(b*x^2+a)*c^3 + 5/2/b^3a^2/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*d^3 - 9/2/b^2a/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c*d^2 + 3/2/b/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c^2*d + 1/2/a/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c^3$

maxima [A] time = 2.99, size = 147, normalized size = 1.39

$$\frac{(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3)x}{2(ab^4x^2 + a^2b^3)} + \frac{bd^3x^3 + 3(3bcd^2 - 2ad^3)x}{3b^3} + \frac{(b^3c^3 + 3ab^2c^2d - 9a^2bcd^2 + 5a^3d^3) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab}ab^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^3/(b*x^2+a)^2,x, algorithm="maxima")`

[Out] $\frac{1}{2}*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*x/(a*b^4*x^2 + a^2*b^3) + \frac{1}{3}*(b*d^3*x^3 + 3*(3*b*c*d^2 - 2*a*d^3)*x)/b^3 + \frac{1}{2}*(b^3*c^3 + 3*a*b^2*c^2*d - 9*a^2*b*c*d^2 + 5*a^3*d^3)*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b}*a*b^3)$

mupad [B] time = 0.10, size = 182, normalized size = 1.72

$$\frac{d^3 x^3}{3b^2} - x \left(\frac{2ad^3}{b^3} - \frac{3cd^2}{b^2} \right) - \frac{x(a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3)}{2a(b^4x^2 + a^2b^3)} + \frac{\operatorname{atan}\left(\frac{\sqrt{b}x(ad-bc)^2(5ad+bc)}{\sqrt{a}(5a^3d^3-9a^2bcd^2+3ab^2c^2d+b^3c^3)}\right)}{2a^{3/2}b^{7/2}} (a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^3/(a + b*x^2)^2,x)`

[Out] $\frac{d^3x^3}{3b^2} - \frac{x((2ad^3)/b^3 - (3cd^2)/b^2)}{2a(b^4x^2 + a^2b^3)} - \frac{x(a^3d^3 - b^3c^3 + 3a^2bcd^2 - 3a^2b^2c^2d - 3a^2b^2c^2d^2)}{(2a*(a*b^3 + b^4*x^2))} + \frac{\operatorname{atan}\left(\frac{b^{1/2}x(ad-bc)^2(5ad+bc)}{a^{3/2}(5a^3d^3-9a^2bcd^2+3ab^2c^2d+b^3c^3)}\right)}{(2a^{3/2}b^{7/2})} (a^3d^3 - b^3c^3 + 3a^2bcd^2 - 3a^2b^2c^2d - 3a^2b^2c^2d^2)$

sympy [B] time = 1.08, size = 314, normalized size = 2.96

$$x \left(-\frac{2ad^3}{b^3} + \frac{3cd^2}{b^2} \right) + \frac{x(-a^3d^3 + 3a^2bcd^2 - 3ab^2c^2d + b^3c^3)}{2a^2b^3 + 2ab^4x^2} - \frac{\sqrt{-\frac{1}{a^3b^7}}(ad-bc)^2(5ad+bc) \log\left(-\frac{a^2b^3\sqrt{-\frac{1}{a^3b^7}}(ad-bc)^2}{5a^3d^3-9a^2bcd^2+3ab^2c^2d+b^3c^3}\right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**3/(b*x**2+a)**2,x)`

```
[Out] x*(-2*a*d**3/b**3 + 3*c*d**2/b**2) + x*(-a**3*d**3 + 3*a**2*b*c*d**2 - 3*a*
b**2*c**2*d + b**3*c**3)/(2*a**2*b**3 + 2*a*b**4*x**2) - sqrt(-1/(a**3*b**7
))*(a*d - b*c)**2*(5*a*d + b*c)*log(-a**2*b**3*sqrt(-1/(a**3*b**7))*(a*d -
b*c)**2*(5*a*d + b*c)/(5*a**3*d**3 - 9*a**2*b*c*d**2 + 3*a*b**2*c**2*d + b*
*3*c**3) + x)/4 + sqrt(-1/(a**3*b**7))*(a*d - b*c)**2*(5*a*d + b*c)*log(a**
2*b**3*sqrt(-1/(a**3*b**7))*(a*d - b*c)**2*(5*a*d + b*c)/(5*a**3*d**3 - 9*a
**2*b*c*d**2 + 3*a*b**2*c**2*d + b**3*c**3) + x)/4 + d**3*x**3/(3*b**2)
```

$$3.30 \quad \int \frac{(c+dx^2)^2}{(a+bx^2)^2} dx$$

Optimal. Leaf size=82

$$\frac{(bc-ad)(3ad+bc)\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{5/2}} + \frac{x(bc-ad)^2}{2ab^2(a+bx^2)} + \frac{d^2x}{b^2}$$

[Out] $d^2*x/b^2+1/2*(-a*d+b*c)^2*x/a/b^2/(b*x^2+a)+1/2*(-a*d+b*c)*(3*a*d+b*c)*\arctan(x*b^{(1/2)}/a^{(1/2)})/a^{(3/2)}/b^{(5/2)}$

Rubi [A] time = 0.10, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {390, 385, 205}

$$\frac{(bc-ad)(3ad+bc)\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{5/2}} + \frac{x(bc-ad)^2}{2ab^2(a+bx^2)} + \frac{d^2x}{b^2}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^2/(a + b*x^2)^2,x]

[Out] $(d^2*x)/b^2 + ((b*c - a*d)^2*x)/(2*a*b^2*(a + b*x^2)) + ((b*c - a*d)*(b*c + 3*a*d)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]])/(2*a^{(3/2)}*b^{(5/2)})$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rubi steps

$$\begin{aligned}
\int \frac{(c + dx^2)^2}{(a + bx^2)^2} dx &= \int \left(\frac{d^2}{b^2} + \frac{b^2c^2 - a^2d^2 + 2bd(bc - ad)x^2}{b^2(a + bx^2)^2} \right) dx \\
&= \frac{d^2x}{b^2} + \frac{\int \frac{b^2c^2 - a^2d^2 + 2bd(bc - ad)x^2}{(a + bx^2)^2} dx}{b^2} \\
&= \frac{d^2x}{b^2} + \frac{(bc - ad)^2x}{2ab^2(a + bx^2)} + \frac{((bc - ad)(bc + 3ad)) \int \frac{1}{a + bx^2} dx}{2ab^2} \\
&= \frac{d^2x}{b^2} + \frac{(bc - ad)^2x}{2ab^2(a + bx^2)} + \frac{(bc - ad)(bc + 3ad) \tan^{-1} \left(\frac{\sqrt{b}x}{\sqrt{a}} \right)}{2a^{3/2}b^{5/2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 88, normalized size = 1.07

$$\frac{(-3a^2d^2 + 2abcd + b^2c^2) \tan^{-1} \left(\frac{\sqrt{b}x}{\sqrt{a}} \right)}{2a^{3/2}b^{5/2}} + \frac{x(bc - ad)^2}{2ab^2(a + bx^2)} + \frac{d^2x}{b^2}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^2/(a + b*x^2)^2,x]

[Out] (d^2*x)/b^2 + ((b*c - a*d)^2*x)/(2*a*b^2*(a + b*x^2)) + ((b^2*c^2 + 2*a*b*c*d - 3*a^2*d^2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(2*a^(3/2)*b^(5/2))

fricas [A] time = 0.64, size = 297, normalized size = 3.62

$$\frac{4a^2b^2d^2x^3 + (ab^2c^2 + 2a^2bcd - 3a^3d^2 + (b^3c^2 + 2ab^2cd - 3a^2bd^2)x^2)\sqrt{-ab} \log\left(\frac{bx^2 + 2\sqrt{-ab}x - a}{bx^2 + a}\right) + 2(ab^3c^2 - 2a^2b^2cd + 3a^3d^2)}{4(a^2b^4x^2 + a^3b^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^2,x, algorithm="fricas")

[Out] [1/4*(4*a^2*b^2*d^2*x^3 + (a*b^2*c^2 + 2*a^2*b*c*d - 3*a^3*d^2 + (b^3*c^2 + 2*a*b^2*c*d - 3*a^2*b*d^2)*x^2)*sqrt(-a*b)*log((b*x^2 + 2*sqrt(-a*b)*x - a)/(b*x^2 + a)) + 2*(a*b^3*c^2 - 2*a^2*b^2*c*d + 3*a^3*b*d^2)*x)/(a^2*b^4*x^2 + a^3*b^3), 1/2*(2*a^2*b^2*d^2*x^3 + (a*b^2*c^2 + 2*a^2*b*c*d - 3*a^3*d^2 + (b^3*c^2 + 2*a*b^2*c*d - 3*a^2*b*d^2)*x^2)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + (a*b^3*c^2 - 2*a^2*b^2*c*d + 3*a^3*b*d^2)*x)/(a^2*b^4*x^2 + a^3*b^3)]

giac [A] time = 0.58, size = 94, normalized size = 1.15

$$\frac{d^2x}{b^2} + \frac{(b^2c^2 + 2abcd - 3a^2d^2) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab}ab^2} + \frac{b^2c^2x - 2abcdx + a^2d^2x}{2(bx^2 + a)ab^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^2,x, algorithm="giac")

[Out] d^2*x/b^2 + 1/2*(b^2*c^2 + 2*a*b*c*d - 3*a^2*d^2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a*b^2) + 1/2*(b^2*c^2*x - 2*a*b*c*d*x + a^2*d^2*x)/((b*x^2 + a)*a*b^2)

maple [A] time = 0.01, size = 129, normalized size = 1.57

$$\frac{a d^2 x}{2(b x^2 + a) b^2} - \frac{3 a d^2 \arctan\left(\frac{b x}{\sqrt{a b}}\right)}{2 \sqrt{a b} b^2} + \frac{c^2 x}{2(b x^2 + a) a} + \frac{c^2 \arctan\left(\frac{b x}{\sqrt{a b}}\right)}{2 \sqrt{a b} a} - \frac{c d x}{(b x^2 + a) b} + \frac{c d \arctan\left(\frac{b x}{\sqrt{a b}}\right)}{\sqrt{a b} b} + \frac{d^2 x}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^2/(b*x^2+a)^2,x)

[Out] 1/b^2*d^2*x+1/2/b^2*a*x/(b*x^2+a)*d^2-1/b*x/(b*x^2+a)*c*d+1/2/a*x/(b*x^2+a)*c^2-3/2/b^2*a/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*d^2+1/b/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c*d+1/2/a/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c^2

maxima [A] time = 2.84, size = 95, normalized size = 1.16

$$\frac{(b^2 c^2 - 2 a b c d + a^2 d^2) x}{2(a b^3 x^2 + a^2 b^2)} + \frac{d^2 x}{b^2} + \frac{(b^2 c^2 + 2 a b c d - 3 a^2 d^2) \arctan\left(\frac{b x}{\sqrt{a b}}\right)}{2 \sqrt{a b} a b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^2,x, algorithm="maxima")

[Out] 1/2*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*x/(a*b^3*x^2 + a^2*b^2) + d^2*x/b^2 + 1/2*(b^2*c^2 + 2*a*b*c*d - 3*a^2*d^2)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a*b^2)

mupad [B] time = 5.06, size = 124, normalized size = 1.51

$$\frac{d^2 x}{b^2} + \frac{x(a^2 d^2 - 2 a b c d + b^2 c^2)}{2 a(b^3 x^2 + a b^2)} + \frac{\operatorname{atan}\left(\frac{\sqrt{b} x(a d - b c)(3 a d + b c)}{\sqrt{a}(-3 a^2 d^2 + 2 a b c d + b^2 c^2)}\right)(a d - b c)(3 a d + b c)}{2 a^{3/2} b^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^2/(a + b*x^2)^2,x)

[Out] (d^2*x)/b^2 + (x*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d))/(2*a*(a*b^2 + b^3*x^2)) + (atan((b^(1/2)*x*(a*d - b*c)*(3*a*d + b*c))/(a^(1/2)*(b^2*c^2 - 3*a^2*d^2 + 2*a*b*c*d)))*(a*d - b*c)*(3*a*d + b*c))/(2*a^(3/2)*b^(5/2))

sympy [B] time = 0.72, size = 236, normalized size = 2.88

$$\frac{x(a^2 d^2 - 2 a b c d + b^2 c^2)}{2 a^2 b^2 + 2 a b^3 x^2} + \frac{\sqrt{-\frac{1}{a^3 b^5}}(a d - b c)(3 a d + b c) \log\left(-\frac{a^2 b^2 \sqrt{-\frac{1}{a^3 b^5}}(a d - b c)(3 a d + b c)}{3 a^2 d^2 - 2 a b c d - b^2 c^2} + x\right)}{4} - \frac{\sqrt{-\frac{1}{a^3 b^5}}(a d - b c)(3 a d + b c)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**2/(b*x**2+a)**2,x)

[Out] x*(a**2*d**2 - 2*a*b*c*d + b**2*c**2)/(2*a**2*b**2 + 2*a*b**3*x**2) + sqrt(-1/(a**3*b**5))*(a*d - b*c)*(3*a*d + b*c)*log(-a**2*b**2*sqrt(-1/(a**3*b**5)))*(a*d - b*c)*(3*a*d + b*c)/(3*a**2*d**2 - 2*a*b*c*d - b**2*c**2) + x)/4 - sqrt(-1/(a**3*b**5))*(a*d - b*c)*(3*a*d + b*c)*log(a**2*b**2*sqrt(-1/(a**3*b**5)))*(a*d - b*c)*(3*a*d + b*c)/(3*a**2*d**2 - 2*a*b*c*d - b**2*c**2) + x)/4 + d**2*x/b**2

$$3.31 \quad \int \frac{c+dx^2}{(a+bx^2)^2} dx$$

Optimal. Leaf size=63

$$\frac{(ad + bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{3/2}} + \frac{x(bc - ad)}{2ab(a + bx^2)}$$

[Out] 1/2*(-a*d+b*c)*x/a/b/(b*x^2+a)+1/2*(a*d+b*c)*arctan(x*b^(1/2)/a^(1/2))/a^(3/2)/b^(3/2)

Rubi [A] time = 0.02, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {385, 205}

$$\frac{(ad + bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{3/2}} + \frac{x(bc - ad)}{2ab(a + bx^2)}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)/(a + b*x^2)^2,x]

[Out] ((b*c - a*d)*x)/(2*a*b*(a + b*x^2)) + ((b*c + a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(2*a^(3/2)*b^(3/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rubi steps

$$\begin{aligned} \int \frac{c + dx^2}{(a + bx^2)^2} dx &= \frac{(bc - ad)x}{2ab(a + bx^2)} + \frac{(bc + ad) \int \frac{1}{a+bx^2} dx}{2ab} \\ &= \frac{(bc - ad)x}{2ab(a + bx^2)} + \frac{(bc + ad) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 63, normalized size = 1.00

$$\frac{(ad + bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}b^{3/2}} - \frac{x(ad - bc)}{2ab(a + bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)/(a + b*x^2)^2,x]

[Out] $-1/2*((-(b*c) + a*d)*x)/(a*b*(a + b*x^2)) + ((b*c + a*d)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]])/(2*a^{3/2}*b^{3/2})$

fricas [A] time = 0.72, size = 181, normalized size = 2.87

$$\left[\frac{(abc + a^2d + (b^2c + abd)x^2)\sqrt{-ab} \log\left(\frac{bx^2 - 2\sqrt{-ab}x - a}{bx^2 + a}\right) - 2(ab^2c - a^2bd)x}{4(a^2b^3x^2 + a^3b^2)}, \frac{(abc + a^2d + (b^2c + abd)x^2)\sqrt{ab} \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(a^2b^3x^2 + a^3b^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^2,x, algorithm="fricas")

[Out] $[-1/4*((a*b*c + a^2*d + (b^2*c + a*b*d)*x^2)*\text{sqrt}(-a*b)*\log((b*x^2 - 2*\text{sqrt}(-a*b)*x - a)/(b*x^2 + a)) - 2*(a*b^2*c - a^2*b*d)*x)/(a^2*b^3*x^2 + a^3*b^2), 1/2*((a*b*c + a^2*d + (b^2*c + a*b*d)*x^2)*\text{sqrt}(a*b)*\arctan(\text{sqrt}(a*b)*x/a) + (a*b^2*c - a^2*b*d)*x)/(a^2*b^3*x^2 + a^3*b^2)]$

giac [A] time = 0.57, size = 57, normalized size = 0.90

$$\frac{(bc + ad) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab} ab} + \frac{bcx - adx}{2(bx^2 + a)ab}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^2,x, algorithm="giac")

[Out] $1/2*(b*c + a*d)*\arctan(b*x/\text{sqrt}(a*b))/(\text{sqrt}(a*b)*a*b) + 1/2*(b*c*x - a*d*x)/((b*x^2 + a)*a*b)$

maple [A] time = 0.01, size = 68, normalized size = 1.08

$$\frac{c \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab} a} + \frac{d \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab} b} - \frac{(ad - bc)x}{2(bx^2 + a)ab}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)/(b*x^2+a)^2,x)

[Out] $-1/2*(a*d-b*c)/a/b*x/(b*x^2+a)+1/2/b/(a*b)^{(1/2)*\arctan(1/(a*b)^{(1/2)*b*x)*d+1/2/a/(a*b)^{(1/2)*\arctan(1/(a*b)^{(1/2)*b*x)*c}$

maxima [A] time = 3.11, size = 57, normalized size = 0.90

$$\frac{(bc - ad)x}{2(ab^2x^2 + a^2b)} + \frac{(bc + ad) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{ab} ab}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^2,x, algorithm="maxima")

[Out] $1/2*(b*c - a*d)*x/(a*b^2*x^2 + a^2*b) + 1/2*(b*c + a*d)*\arctan(b*x/\text{sqrt}(a*b))/(\text{sqrt}(a*b)*a*b)$

mupad [B] time = 5.04, size = 51, normalized size = 0.81

$$\frac{\text{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)(ad + bc)}{2a^{3/2}b^{3/2}} - \frac{x(ad - bc)}{2ab(bx^2 + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)/(a + b*x^2)^2,x)`

[Out] $(\operatorname{atan}((b^{1/2}x)/a^{1/2})*(a*d + b*c))/(2*a^{3/2}*b^{3/2}) - (x*(a*d - b*c))/(2*a*b*(a + b*x^2))$

sympy [B] time = 0.40, size = 112, normalized size = 1.78

$$\frac{x(-ad + bc)}{2a^2b + 2ab^2x^2} - \frac{\sqrt{-\frac{1}{a^3b^3}}(ad + bc)\log\left(-a^2b\sqrt{-\frac{1}{a^3b^3}} + x\right)}{4} + \frac{\sqrt{-\frac{1}{a^3b^3}}(ad + bc)\log\left(a^2b\sqrt{-\frac{1}{a^3b^3}} + x\right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)/(b*x**2+a)**2,x)`

[Out] $x*(-a*d + b*c)/(2*a**2*b + 2*a*b**2*x**2) - \operatorname{sqrt}(-1/(a**3*b**3))*(a*d + b*c)*\log(-a**2*b*\operatorname{sqrt}(-1/(a**3*b**3)) + x)/4 + \operatorname{sqrt}(-1/(a**3*b**3))*(a*d + b*c)*\log(a**2*b*\operatorname{sqrt}(-1/(a**3*b**3)) + x)/4$

$$3.32 \quad \int \frac{1}{(a+bx^2)^2(c+dx^2)} dx$$

Optimal. Leaf size=108

$$\frac{\sqrt{b}(bc-3ad)\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}(bc-ad)^2} + \frac{d^{3/2}\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)^2} + \frac{bx}{2a(a+bx^2)(bc-ad)}$$

[Out] 1/2*b*x/a/(-a*d+b*c)/(b*x^2+a)+1/2*(-3*a*d+b*c)*arctan(x*b^(1/2)/a^(1/2))*b^(1/2)/a^(3/2)/(-a*d+b*c)^2+d^(3/2)*arctan(x*d^(1/2)/c^(1/2))/(-a*d+b*c)^2/c^(1/2)

Rubi [A] time = 0.08, antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {414, 522, 205}

$$\frac{\sqrt{b}(bc-3ad)\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}(bc-ad)^2} + \frac{d^{3/2}\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)^2} + \frac{bx}{2a(a+bx^2)(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^2*(c + d*x^2)),x]

[Out] (b*x)/(2*a*(b*c - a*d)*(a + b*x^2)) + (Sqrt[b]*(b*c - 3*a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(2*a^(3/2)*(b*c - a*d)^2) + (d^(3/2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(Sqrt[c]*(b*c - a*d)^2)

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 522

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)^2(c+dx^2)} dx &= \frac{bx}{2a(bc-ad)(a+bx^2)} - \frac{\int \frac{-bc+2ad-bdx^2}{(a+bx^2)(c+dx^2)} dx}{2a(bc-ad)} \\ &= \frac{bx}{2a(bc-ad)(a+bx^2)} + \frac{d^2 \int \frac{1}{c+dx^2} dx}{(bc-ad)^2} + \frac{(b(bc-3ad)) \int \frac{1}{a+bx^2} dx}{2a(bc-ad)^2} \\ &= \frac{bx}{2a(bc-ad)(a+bx^2)} + \frac{\sqrt{b}(bc-3ad) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}(bc-ad)^2} + \frac{d^{3/2} \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)^2} \end{aligned}$$

Mathematica [A] time = 0.15, size = 109, normalized size = 1.01

$$-\frac{\sqrt{b}(3ad-bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}(ad-bc)^2} + \frac{d^{3/2} \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)^2} - \frac{bx}{2a(a+bx^2)(ad-bc)}$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^2*(c + d*x^2)), x]

[Out] -1/2*(b*x)/(a*(-(b*c) + a*d)*(a + b*x^2)) - (Sqrt[b]*(-(b*c) + 3*a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(2*a^(3/2)*(-(b*c) + a*d)^2) + (d^(3/2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(Sqrt[c]*(b*c - a*d)^2)

fricas [A] time = 0.80, size = 699, normalized size = 6.47

$$\left[\frac{(abc - 3a^2d + (b^2c - 3abd)x^2) \sqrt{-\frac{b}{a}} \log\left(\frac{bx^2 - 2ax\sqrt{-\frac{b}{a}} - a}{bx^2 + a}\right) - 2(abdx^2 + a^2d) \sqrt{-\frac{d}{c}} \log\left(\frac{dx^2 + 2cx\sqrt{-\frac{d}{c}} - c}{dx^2 + c}\right) - 2(b^2c - 3abd)x^2 \sqrt{\frac{b}{a}} \arctan\left(\frac{x\sqrt{\frac{b}{a}}}{1}\right) + (abc - 3a^2d + (b^2c - 3abd)x^2) \sqrt{-\frac{b}{a}} \log\left(\frac{bx^2 - 2ax\sqrt{-\frac{b}{a}} - a}{bx^2 + a}\right) + 2(b^2c - a*b*d)*x / (a^2*b^2*c^2 - 2*a^3*b*c*d + a^4*d^2 + (a*b^3*c^2 - 2*a^2*b^2*c*d + a^3*b*d^2)*x^2), 1/2*((a*b*c - 3*a^2*d + (b^2*c - 3*a*b*d)*x^2) \sqrt{b/a} \arctan(x \sqrt{b/a}) + (a*b*d*x^2 + a^2*d) \sqrt{-d/c} \log((d*x^2 + 2*c*x \sqrt{-d/c} - c)/(d*x^2 + c)) + (b^2*c - a*b*d)*x / (a^2*b^2*c^2 - 2*a^3*b*c*d + a^4*d^2 + (a*b^3*c^2 - 2*a^2*b^2*c*d + a^3*b*d^2)*x^2), 1/2*((a*b*c - 3*a^2*d + (b^2*c - 3*a*b*d)*x^2) \sqrt{b/a} \arctan(x \sqrt{b/a}) + 2*(a*b*d*x^2 + a^2*d) \sqrt{d/c} \arctan(x \sqrt{d/c}) + (b^2*c - a*b*d)*x / (a^2*b^2*c^2 - 2*a^3*b*c*d + a^4*d^2 + (a*b^3*c^2 - 2*a^2*b^2*c*d + a^3*b*d^2)*x^2)}{4(a^2b^2c^2 - 2a^3bcd + a^4d^2 + (ab^3c^2 - 2a^2b^2cd + a^3bd^2)x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c), x, algorithm="fricas")

[Out] [-1/4*((a*b*c - 3*a^2*d + (b^2*c - 3*a*b*d)*x^2)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) - 2*(a*b*d*x^2 + a^2*d)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) - 2*(b^2*c - a*b*d)*x/(a^2*b^2*c^2 - 2*a^3*b*c*d + a^4*d^2 + (a*b^3*c^2 - 2*a^2*b^2*c*d + a^3*b*d^2)*x^2), 1/4*(4*(a*b*d*x^2 + a^2*d)*sqrt(d/c)*arctan(x*sqrt(d/c)) - (a*b*c - 3*a^2*d + (b^2*c - 3*a*b*d)*x^2)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + 2*(b^2*c - a*b*d)*x/(a^2*b^2*c^2 - 2*a^3*b*c*d + a^4*d^2 + (a*b^3*c^2 - 2*a^2*b^2*c*d + a^3*b*d^2)*x^2), 1/2*((a*b*c - 3*a^2*d + (b^2*c - 3*a*b*d)*x^2)*sqrt(b/a)*arctan(x*sqrt(b/a)) + (a*b*d*x^2 + a^2*d)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) + (b^2*c - a*b*d)*x/(a^2*b^2*c^2 - 2*a^3*b*c*d + a^4*d^2 + (a*b^3*c^2 - 2*a^2*b^2*c*d + a^3*b*d^2)*x^2), 1/2*((a*b*c - 3*a^2*d + (b^2*c - 3*a*b*d)*x^2)*sqrt(b/a)*arctan(x*sqrt(b/a)) + 2*(a*b*d*x^2 + a^2*d)*sqrt(d/c)*arctan(x*sqrt(d/c)) + (b^2*c - a*b*d)*x/(a^2*b^2*c^2 - 2*a^3*b*c*d + a^4*d^2 + (a*b^3*c^2 - 2*a^2*b^2*c*d + a^3*b*d^2)*x^2)]

giac [A] time = 0.58, size = 121, normalized size = 1.12

$$\frac{d^2 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{(b^2c^2 - 2abcd + a^2d^2)\sqrt{cd}} + \frac{(b^2c - 3abd) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ab^2c^2 - 2a^2bcd + a^3d^2)\sqrt{ab}} + \frac{bx}{2(abc - a^2d)(bx^2 + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c),x, algorithm="giac")

[Out] $d^2 \arctan(d*x/\sqrt{c*d})/((b^2*c^2 - 2*a*b*c*d + a^2*d^2)*\sqrt{c*d}) + 1/2 * (b^2*c - 3*a*b*d) \arctan(b*x/\sqrt{a*b})/((a*b^2*c^2 - 2*a^2*b*c*d + a^3*d^2)*\sqrt{a*b}) + 1/2*b*x/((a*b*c - a^2*d)*(b*x^2 + a))$

maple [A] time = 0.01, size = 144, normalized size = 1.33

$$\frac{b^2cx}{2(ad-bc)^2(bx^2+a)a} + \frac{b^2c \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ad-bc)^2\sqrt{ab}a} - \frac{bdx}{2(ad-bc)^2(bx^2+a)} - \frac{3bd \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ad-bc)^2\sqrt{ab}} + \frac{d^2 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{(ad-bc)^2\sqrt{cd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^2/(d*x^2+c),x)

[Out] $d^2/(a*d-b*c)^2/(c*d)^{(1/2)}*\arctan(1/(c*d)^{(1/2)}*d*x)-1/2*b/(a*d-b*c)^2*x/(b*x^2+a)*d+1/2*b^2/(a*d-b*c)^2/a*x/(b*x^2+a)*c-3/2*b/(a*d-b*c)^2/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*d+1/2*b^2/(a*d-b*c)^2/a/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c$

maxima [A] time = 2.92, size = 132, normalized size = 1.22

$$\frac{d^2 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{(b^2c^2 - 2abcd + a^2d^2)\sqrt{cd}} + \frac{bx}{2(a^2bc - a^3d + (ab^2c - a^2bd)x^2)} + \frac{(b^2c - 3abd) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ab^2c^2 - 2a^2bcd + a^3d^2)\sqrt{ab}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c),x, algorithm="maxima")

[Out] $d^2 \arctan(d*x/\sqrt{c*d})/((b^2*c^2 - 2*a*b*c*d + a^2*d^2)*\sqrt{c*d}) + 1/2 * b*x/(a^2*b*c - a^3*d + (a*b^2*c - a^2*b*d)*x^2) + 1/2*(b^2*c - 3*a*b*d) \arctan(b*x/\sqrt{a*b})/((a*b^2*c^2 - 2*a^2*b*c*d + a^3*d^2)*\sqrt{a*b})$

mupad [B] time = 5.77, size = 3649, normalized size = 33.79

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^2*(c + d*x^2)),x)

[Out] $(\operatorname{atan}(\frac{(-a^3b)^{(1/2)}*(3ad-bc)*((x*(13a^2b^3d^5 + b^5c^2d^3 - 6ab^4cd^4))}{(2(a^4d^2 + a^2b^2c^2 - 2a^3b^2cd)) - ((4a^6b^2d^7 - 2a^7c^5d^2 - 18a^5b^3cd^6 + 12a^2b^6c^4d^3 - 28a^3b^5c^3d^4 + 32a^4b^4c^2d^5))}{(a^5d^3 - a^2b^3c^3 + 3a^3b^2c^2d - 3a^4b^2cd^2) - (x*(-a^3b)^{(1/2)}*(3ad-bc)*(16a^7b^2d^7 - 48a^6b^3cd^6 + 16a^2b^7c^5d^2 - 48a^3b^6c^4d^3 + 32a^4b^5c^3d^4 + 32a^5b^4c^2d^5))}{(8(a^4d^2 + a^2b^2c^2 - 2a^3b^2cd))*(a^5d^2 + a^3b^2c^2 - 2a^4b^2cd))}*(-a^3b)^{(1/2)}*(3ad-bc))/((4(a^5d^2 + a^3b^2c^2 - 2a^4b^2cd)))*1i)/(4(a^5d^2 + a^3b^2c^2 - 2a^4b^2cd)) + ((-a^3b)^{(1/2)}*(3ad-bc)*((x*(13a^2b^3d^5 + b^5c^2d^3 - 6ab^4cd^4))}{(2(a^4d^2 + a^2b^2c^2 - 2a^3b^2cd))} + ((4a^6b^2d^7 - 2a^7c^5d^2 - 18a^5b^3cd^6 + 12a^2b^6c^4d^3 - 28a^3b^5c^3d^4 + 32a^4b^4c^2d^5))}{(a^5d^3 - a^2b^3c^3 + 3a^3b^2c^2d - 3a^4b^2cd^2) + (x*(-a^3b)^{(1/2)}*(3ad-bc)*(16a^7b^2d^7 - 48a^6b^3cd^6 + 16a^2b^7c^5d^2 - 48a^3b^6c^4d^3 + 32a^4b^5c^3d^4 + 32a^5b^4c^2d^5))}{(8(a^4d^2 + a^2b^2c^2 - 2a^3b^2cd))*(a^5d^2 + a^3b^2c^2 - 2a^4b^2cd))}*(-a^3b)^{(1/2)}*(3ad-bc))/((4(a^5d^2 + a^3b^2c^2 - 2a^4b^2cd))$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x**2+a)**2/(d*x**2+c),x)
```

```
[Out] Timed out
```

$$3.33 \quad \int \frac{1}{(a+bx^2)^2(c+dx^2)^2} dx$$

Optimal. Leaf size=167

$$\frac{b^{3/2}(bc-5ad)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}(bc-ad)^3} + \frac{d^{3/2}(5bc-ad)\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)}{2c^{3/2}(bc-ad)^3} + \frac{bx}{2a(a+bx^2)(c+dx^2)(bc-ad)} + \frac{dx(ad+bc)}{2ac(c+dx^2)(bc-ad)}$$

[Out] 1/2*d*(a*d+b*c)*x/a/c/(-a*d+b*c)^2/(d*x^2+c)+1/2*b*x/a/(-a*d+b*c)/(b*x^2+a)/(d*x^2+c)+1/2*b^(3/2)*(-5*a*d+b*c)*arctan(x*b^(1/2)/a^(1/2))/a^(3/2)/(-a*d+b*c)^3+1/2*d^(3/2)*(-a*d+5*b*c)*arctan(x*d^(1/2)/c^(1/2))/c^(3/2)/(-a*d+b*c)^3

Rubi [A] time = 0.20, antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {414, 527, 522, 205}

$$\frac{b^{3/2}(bc-5ad)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{2a^{3/2}(bc-ad)^3} + \frac{d^{3/2}(5bc-ad)\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)}{2c^{3/2}(bc-ad)^3} + \frac{bx}{2a(a+bx^2)(c+dx^2)(bc-ad)} + \frac{dx(ad+bc)}{2ac(c+dx^2)(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^2*(c + d*x^2)^2), x]

[Out] (d*(b*c + a*d)*x)/(2*a*c*(b*c - a*d)^2*(c + d*x^2)) + (b*x)/(2*a*(b*c - a*d)*(a + b*x^2)*(c + d*x^2)) + (b^(3/2)*(b*c - 5*a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(2*a^(3/2)*(b*c - a*d)^3) + (d^(3/2)*(5*b*c - a*d)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(2*c^(3/2)*(b*c - a*d)^3)

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 522

Int[((e_) + (f_.)*(x_)^(n_))/(((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_))), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(b*c - a*d)*(p+1)), x] + Dist[1/(a*n*(b*c - a*d)*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2)+1)*x^n, x], x], x] /; FreeQ

[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a+bx^2)^2(c+dx^2)^2} dx &= \frac{bx}{2a(bc-ad)(a+bx^2)(c+dx^2)} - \frac{\int \frac{-bc+2ad-3bdx^2}{(a+bx^2)(c+dx^2)^2} dx}{2a(bc-ad)} \\
 &= \frac{d(bc+ad)x}{2ac(bc-ad)^2(c+dx^2)} + \frac{bx}{2a(bc-ad)(a+bx^2)(c+dx^2)} - \frac{\int \frac{-2(b^2c^2-4abcd+a^2d^2)-2bdx^2}{(a+bx^2)(c+dx^2)^2} dx}{4ac(bc-ad)^2} \\
 &= \frac{d(bc+ad)x}{2ac(bc-ad)^2(c+dx^2)} + \frac{bx}{2a(bc-ad)(a+bx^2)(c+dx^2)} + \frac{(b^2(bc-5ad)) \int \frac{1}{a+bx^2} dx}{2a(bc-ad)^3} \\
 &= \frac{d(bc+ad)x}{2ac(bc-ad)^2(c+dx^2)} + \frac{bx}{2a(bc-ad)(a+bx^2)(c+dx^2)} + \frac{b^{3/2}(bc-5ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{a}}\right)}{2a^{3/2}(bc-ad)^3}
 \end{aligned}$$

Mathematica [A] time = 0.32, size = 136, normalized size = 0.81

$$\frac{1}{2} \left(\frac{b^{3/2}(5ad-bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{a^{3/2}(ad-bc)^3} + \frac{x(bc-ad) \left(\frac{b^2}{a^2+abx^2} + \frac{d^2}{c^2+cdx^2} \right) + \frac{d^{3/2}(5bc-ad) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{c^{3/2}}}{(bc-ad)^3} \right)$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^2*(c + d*x^2)^2), x]

[Out] ((b^(3/2)*(-(b*c) + 5*a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(a^(3/2)*(-(b*c) + a*d)^3) + ((b*c - a*d)*x*(b^2/(a^2 + a*b*x^2) + d^2/(c^2 + c*d*x^2)) + (d^(3/2)*(5*b*c - a*d)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/c^(3/2))/(b*c - a*d)^3/2

fricas [B] time = 1.64, size = 1681, normalized size = 10.07

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)^2,x, algorithm="fricas")

[Out] [1/4*(2*(b^3*c^2*d - a^2*b*d^3)*x^3 + (a*b^2*c^3 - 5*a^2*b*c^2*d + (b^3*c^2*d - 5*a*b^2*c*d^2)*x^4 + (b^3*c^3 - 4*a*b^2*c^2*d - 5*a^2*b*c*d^2)*x^2)*sqrt(-b/a)*log((b*x^2 + 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + (5*a^2*b*c^2*d - a^3*c*d^2 + (5*a*b^2*c*d^2 - a^2*b*d^3)*x^4 + (5*a*b^2*c^2*d + 4*a^2*b*c*d^2 - a^3*d^3)*x^2)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) + 2*(b^3*c^3 - a*b^2*c^2*d + a^2*b*c*d^2 - a^3*d^3)*x)/(a^2*b^3*c^5 - 3*a^3*b^2*c^4*d + 3*a^4*b*c^3*d^2 - a^5*c^2*d^3 + (a*b^4*c^4*d - 3*a^2*b^3*c^3*d^2 + 3*a^3*b^2*c^2*d^3 - a^4*b*c*d^4)*x^4 + (a*b^4*c^5 - 2*a^2*b^3*c^4*d + 2*a^4*b*c^2*d^3 - a^5*c*d^4)*x^2), 1/4*(2*(b^3*c^2*d - a^2*b*d^3)*x^3 + 2*(5*a^2*b*c^2*d - a^3*c*d^2 + (5*a*b^2*c*d^2 - a^2*b*d^3)*x^4 + (5*a*b^2*c^2*d + 4*a^2*b*c*d^2 - a^3*d^3)*x^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) + (a*b^2*c^3 - 5*a^2*b*c^2*d + (b^3*c^2*d - 5*a*b^2*c*d^2)*x^4 + (b^3*c^3 - 4*a*b^2*c^2*d - 5*a^2*b*c*d^2)*x^2)*sqrt(-b/a)*log((b*x^2 + 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + 2*(b^3*c^3 - a*b^2*c^2*d + a^2*b*c*d^2 - a^3*d^3)*x)/(a^2*b^3*c^5 - 3*a^3*b^2*c^4*d + 3*a^4*b*c^3*d^2 - a^5*c^2*d^3 + (a*b^4*c^4*d - 3*

$$a^2b^3c^3d^2 + 3a^3b^2c^2d^3 - a^4b^2cd^4)x^4 + (a^2b^4c^5 - 2a^2b^3c^4d + 2a^4b^2c^2d^3 - a^5c^2d^4)x^2, 1/4*(2*(b^3c^2d - a^2b^3d^3)*x^3 + 2*(a^2b^2c^3 - 5a^2b^2c^2d + (b^3c^2d - 5a^2b^2c^2d^2)*x^4 + (b^3c^3 - 4a^2b^2c^2d - 5a^2b^2c^2d^2)*x^2)*\sqrt{b/a}*\arctan(x*\sqrt{b/a}) + (5a^2b^2c^2d - a^3c^2d^2 + (5a^2b^2c^2d^2 - a^2b^2d^3)*x^4 + (5a^2b^2c^2d + 4a^2b^2c^2d^2 - a^3d^3)*x^2)*\sqrt{-d/c}*\log((d*x^2 + 2*c*x*\sqrt{-d/c}) - c)/(d*x^2 + c)) + 2*(b^3c^3 - a^2b^2c^2d + a^2b^2c^2d^2 - a^3d^3)*x)/(a^2b^3c^5 - 3a^3b^2c^4d + 3a^4b^2c^3d^2 - a^5c^2d^3 + (a^2b^4c^4d - 3a^2b^3c^3d^2 + 3a^3b^2c^2d^3 - a^4b^2cd^4)*x^4 + (a^2b^4c^5 - 2a^2b^3c^4d + 2a^4b^2c^2d^3 - a^5c^2d^4)*x^2), 1/2*((b^3c^2d - a^2b^2d^3)*x^3 + (a^2b^2c^3 - 5a^2b^2c^2d + (b^3c^2d - 5a^2b^2c^2d^2)*x^4 + (b^3c^3 - 4a^2b^2c^2d - 5a^2b^2c^2d^2)*x^2)*\sqrt{b/a}*\arctan(x*\sqrt{b/a})) + (5a^2b^2c^2d - a^3c^2d^2 + (5a^2b^2c^2d^2 - a^2b^2d^3)*x^4 + (5a^2b^2c^2d + 4a^2b^2c^2d^2 - a^3d^3)*x^2)*\sqrt{d/c}*\arctan(x*\sqrt{d/c})) + (b^3c^3 - a^2b^2c^2d + a^2b^2c^2d^2 - a^3d^3)*x)/(a^2b^3c^5 - 3a^3b^2c^4d + 3a^4b^2c^3d^2 - a^5c^2d^3 + (a^2b^4c^4d - 3a^2b^3c^3d^2 + 3a^3b^2c^2d^3 - a^4b^2cd^4)*x^4 + (a^2b^4c^5 - 2a^2b^3c^4d + 2a^4b^2c^2d^3 - a^5c^2d^4)*x^2)]$$

giac [A] time = 0.58, size = 232, normalized size = 1.39

$$\frac{(b^3c - 5ab^2d) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ab^3c^3 - 3a^2b^2c^2d + 3a^3bcd^2 - a^4d^3)\sqrt{ab}} + \frac{(5bcd^2 - ad^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2(b^3c^4 - 3ab^2c^3d + 3a^2bc^2d^2 - a^3cd^3)\sqrt{cd}} + \frac{b^2cdx^3}{2(ab^2c^3 - 2a^2bc^2d)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)^2,x, algorithm="giac")

[Out] 1/2*(b^3*c - 5*a*b^2*d)*arctan(b*x/sqrt(a*b))/((a*b^3*c^3 - 3*a^2*b^2*c^2*d + 3*a^3*b*c*d^2 - a^4*d^3)*sqrt(a*b)) + 1/2*(5*b*c*d^2 - a*d^3)*arctan(d*x/sqrt(c*d))/((b^3*c^4 - 3*a*b^2*c^3*d + 3*a^2*b*c^2*d^2 - a^3*c*d^3)*sqrt(c*d)) + 1/2*(b^2*c*d*x^3 + a*b*d^2*x^3 + b^2*c^2*x + a^2*d^2*x)/((a*b^2*c^3 - 2*a^2*b*c^2*d + a^3*c*d^2)*(b*d*x^4 + b*c*x^2 + a*d*x^2 + a*c))

maple [A] time = 0.02, size = 238, normalized size = 1.43

$$\frac{a d^3 x}{2(ad - bc)^3 (d x^2 + c) c} + \frac{a d^3 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2(ad - bc)^3 \sqrt{cd} c} - \frac{b^3 c x}{2(ad - bc)^3 (b x^2 + a) a} - \frac{b^3 c \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ad - bc)^3 \sqrt{ab} a} + \frac{b^2 dx}{2(ad - bc)^3 (b x^2 + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^2/(d*x^2+c)^2,x)

[Out] 1/2*d^3/(a*d-b*c)^3/c*x/(d*x^2+c)*a-1/2*d^2/(a*d-b*c)^3*x/(d*x^2+c)*b+1/2*d^3/(a*d-b*c)^3/c/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a-5/2*d^2/(a*d-b*c)^3/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*b+1/2*b^2/(a*d-b*c)^3*x/(b*x^2+a)*d-1/2*b^3/(a*d-b*c)^3/a*x/(b*x^2+a)*c+5/2*b^2/(a*d-b*c)^3/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*d-1/2*b^3/(a*d-b*c)^3/a/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c

maxima [B] time = 3.14, size = 294, normalized size = 1.76

$$\frac{(b^3c - 5ab^2d) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ab^3c^3 - 3a^2b^2c^2d + 3a^3bcd^2 - a^4d^3)\sqrt{ab}} + \frac{(5bcd^2 - ad^3) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{2(b^3c^4 - 3ab^2c^3d + 3a^2bc^2d^2 - a^3cd^3)\sqrt{cd}} + \frac{b^2cdx^3}{2(a^2b^2c^4 - 2a^3bc^2d)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)^2,x, algorithm="maxima")

```
[Out] 1/2*(b^3*c - 5*a*b^2*d)*arctan(b*x/sqrt(a*b))/((a*b^3*c^3 - 3*a^2*b^2*c^2*d
+ 3*a^3*b*c*d^2 - a^4*d^3)*sqrt(a*b)) + 1/2*(5*b*c*d^2 - a*d^3)*arctan(d*x
/sqrt(c*d))/((b^3*c^4 - 3*a*b^2*c^3*d + 3*a^2*b*c^2*d^2 - a^3*c*d^3)*sqrt(c
*d)) + 1/2*((b^2*c*d + a*b*d^2)*x^3 + (b^2*c^2 + a^2*d^2)*x)/(a^2*b^2*c^4 -
2*a^3*b*c^3*d + a^4*c^2*d^2 + (a*b^3*c^3*d - 2*a^2*b^2*c^2*d^2 + a^3*b*c*d
^3)*x^4 + (a*b^3*c^4 - a^2*b^2*c^3*d - a^3*b*c^2*d^2 + a^4*c*d^3)*x^2)
```

mupad [B] time = 6.87, size = 6183, normalized size = 37.02

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((a + b*x^2)^2*(c + d*x^2)^2),x)
```

```
[Out] ((x*(a^2*d^2 + b^2*c^2))/(2*a*c*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d)) + (b*d*x^3
*(a*d + b*c))/(2*a*c*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d)))/(a*c + x^2*(a*d + b*
c) + b*d*x^4) + (atan((((x*(a^4*b^3*d^7 + b^7*c^4*d^3 - 10*a*b^6*c^3*d^4 -
10*a^3*b^4*c*d^6 + 50*a^2*b^5*c^2*d^5))/(2*(a^2*b^4*c^6 + a^6*c^2*d^4 - 4*
a^3*b^3*c^5*d - 4*a^5*b*c^3*d^3 + 6*a^4*b^2*c^4*d^2))) - (((2*a*b^10*c^9*d^2
+ 2*a^9*b^2*c*d^10 - 20*a^2*b^9*c^8*d^3 + 80*a^3*b^8*c^7*d^4 - 172*a^4*b^7
*c^6*d^5 + 220*a^5*b^6*c^5*d^6 - 172*a^6*b^5*c^4*d^7 + 80*a^7*b^4*c^3*d^8 -
20*a^8*b^3*c^2*d^9)/(a^2*b^6*c^8 + a^8*c^2*d^6 - 6*a^3*b^5*c^7*d - 6*a^7*b
*c^3*d^5 + 15*a^4*b^4*c^6*d^2 - 20*a^5*b^3*c^5*d^3 + 15*a^6*b^2*c^4*d^4) -
(x*(5*a*d - b*c)*(-a^3*b^3)^(1/2)*(16*a^2*b^9*c^9*d^2 - 80*a^3*b^8*c^8*d^3
+ 144*a^4*b^7*c^7*d^4 - 80*a^5*b^6*c^6*d^5 - 80*a^6*b^5*c^5*d^6 + 144*a^7*b
^4*c^4*d^7 - 80*a^8*b^3*c^3*d^8 + 16*a^9*b^2*c^2*d^9)))/(8*(a^6*d^3 - a^3*b^
3*c^3 + 3*a^4*b^2*c^2*d - 3*a^5*b*c*d^2)*(a^2*b^4*c^6 + a^6*c^2*d^4 - 4*a^3
*b^3*c^5*d - 4*a^5*b*c^3*d^3 + 6*a^4*b^2*c^4*d^2)))*(5*a*d - b*c)*(-a^3*b^3
)^(1/2))/(4*(a^6*d^3 - a^3*b^3*c^3 + 3*a^4*b^2*c^2*d - 3*a^5*b*c*d^2)))*(5*
a*d - b*c)*(-a^3*b^3)^(1/2)*i)/(4*(a^6*d^3 - a^3*b^3*c^3 + 3*a^4*b^2*c^2*d
- 3*a^5*b*c*d^2)) + (((x*(a^4*b^3*d^7 + b^7*c^4*d^3 - 10*a*b^6*c^3*d^4 - 1
0*a^3*b^4*c*d^6 + 50*a^2*b^5*c^2*d^5))/(2*(a^2*b^4*c^6 + a^6*c^2*d^4 - 4*a^
3*b^3*c^5*d - 4*a^5*b*c^3*d^3 + 6*a^4*b^2*c^4*d^2))) + (((2*a*b^10*c^9*d^2 +
2*a^9*b^2*c*d^10 - 20*a^2*b^9*c^8*d^3 + 80*a^3*b^8*c^7*d^4 - 172*a^4*b^7*c
^6*d^5 + 220*a^5*b^6*c^5*d^6 - 172*a^6*b^5*c^4*d^7 + 80*a^7*b^4*c^3*d^8 - 2
0*a^8*b^3*c^2*d^9)/(a^2*b^6*c^8 + a^8*c^2*d^6 - 6*a^3*b^5*c^7*d - 6*a^7*b*c
^3*d^5 + 15*a^4*b^4*c^6*d^2 - 20*a^5*b^3*c^5*d^3 + 15*a^6*b^2*c^4*d^4) + (x
*(5*a*d - b*c)*(-a^3*b^3)^(1/2)*(16*a^2*b^9*c^9*d^2 - 80*a^3*b^8*c^8*d^3 +
144*a^4*b^7*c^7*d^4 - 80*a^5*b^6*c^6*d^5 - 80*a^6*b^5*c^5*d^6 + 144*a^7*b^4
*c^4*d^7 - 80*a^8*b^3*c^3*d^8 + 16*a^9*b^2*c^2*d^9)))/(8*(a^6*d^3 - a^3*b^3*
c^3 + 3*a^4*b^2*c^2*d - 3*a^5*b*c*d^2)*(a^2*b^4*c^6 + a^6*c^2*d^4 - 4*a^3*b
^3*c^5*d - 4*a^5*b*c^3*d^3 + 6*a^4*b^2*c^4*d^2)))*(5*a*d - b*c)*(-a^3*b^3)^(
1/2))/(4*(a^6*d^3 - a^3*b^3*c^3 + 3*a^4*b^2*c^2*d - 3*a^5*b*c*d^2)))*(5*a*
d - b*c)*(-a^3*b^3)^(1/2)*i)/(4*(a^6*d^3 - a^3*b^3*c^3 + 3*a^4*b^2*c^2*d -
3*a^5*b*c*d^2)))/(((5*a^3*b^4*d^7)/4 + (5*b^7*c^3*d^4)/4 - (21*a*b^6*c^2*d
^5)/4 - (21*a^2*b^5*c*d^6)/4)/(a^2*b^6*c^8 + a^8*c^2*d^6 - 6*a^3*b^5*c^7*d
- 6*a^7*b*c^3*d^5 + 15*a^4*b^4*c^6*d^2 - 20*a^5*b^3*c^5*d^3 + 15*a^6*b^2*c^
4*d^4) - (((x*(a^4*b^3*d^7 + b^7*c^4*d^3 - 10*a*b^6*c^3*d^4 - 10*a^3*b^4*c*
d^6 + 50*a^2*b^5*c^2*d^5))/(2*(a^2*b^4*c^6 + a^6*c^2*d^4 - 4*a^3*b^3*c^5*d
- 4*a^5*b*c^3*d^3 + 6*a^4*b^2*c^4*d^2))) - (((2*a*b^10*c^9*d^2 + 2*a^9*b^2*c
*d^10 - 20*a^2*b^9*c^8*d^3 + 80*a^3*b^8*c^7*d^4 - 172*a^4*b^7*c^6*d^5 + 220
*a^5*b^6*c^5*d^6 - 172*a^6*b^5*c^4*d^7 + 80*a^7*b^4*c^3*d^8 - 20*a^8*b^3*c^
2*d^9)/(a^2*b^6*c^8 + a^8*c^2*d^6 - 6*a^3*b^5*c^7*d - 6*a^7*b*c^3*d^5 + 15*
a^4*b^4*c^6*d^2 - 20*a^5*b^3*c^5*d^3 + 15*a^6*b^2*c^4*d^4) - (x*(5*a*d - b*
c)*(-a^3*b^3)^(1/2)*(16*a^2*b^9*c^9*d^2 - 80*a^3*b^8*c^8*d^3 + 144*a^4*b^7*
c^7*d^4 - 80*a^5*b^6*c^6*d^5 - 80*a^6*b^5*c^5*d^6 + 144*a^7*b^4*c^4*d^7 - 8
0*a^8*b^3*c^3*d^8 + 16*a^9*b^2*c^2*d^9)))/(8*(a^6*d^3 - a^3*b^3*c^3 + 3*a^4*
b^2*c^2*d - 3*a^5*b*c*d^2)*(a^2*b^4*c^6 + a^6*c^2*d^4 - 4*a^3*b^3*c^5*d - 4
*a^5*b*c^3*d^3 + 6*a^4*b^2*c^4*d^2)))*(5*a*d - b*c)*(-a^3*b^3)^(1/2))/(4*(a
^6*d^3 - a^3*b^3*c^3 + 3*a^4*b^2*c^2*d - 3*a^5*b*c*d^2)))*(5*a*d - b*c)*(-a
```


$$\begin{aligned} &)/(4*(b^3*c^6 - a^3*c^3*d^3 + 3*a^2*b*c^4*d^2 - 3*a*b^2*c^5*d)))*(a*d - 5*b \\ & *c)*(-c^3*d^3)^{(1/2)})/(4*(b^3*c^6 - a^3*c^3*d^3 + 3*a^2*b*c^4*d^2 - 3*a*b^2 \\ & *c^5*d)) + (((x*(a^4*b^3*d^7 + b^7*c^4*d^3 - 10*a*b^6*c^3*d^4 - 10*a^3*b^4* \\ & c*d^6 + 50*a^2*b^5*c^2*d^5))/(2*(a^2*b^4*c^6 + a^6*c^2*d^4 - 4*a^3*b^3*c^5* \\ & d - 4*a^5*b*c^3*d^3 + 6*a^4*b^2*c^4*d^2)) + (((2*a*b^10*c^9*d^2 + 2*a^9*b^2 \\ & *c*d^10 - 20*a^2*b^9*c^8*d^3 + 80*a^3*b^8*c^7*d^4 - 172*a^4*b^7*c^6*d^5 + 2 \\ & 20*a^5*b^6*c^5*d^6 - 172*a^6*b^5*c^4*d^7 + 80*a^7*b^4*c^3*d^8 - 20*a^8*b^3* \\ & c^2*d^9)/(a^2*b^6*c^8 + a^8*c^2*d^6 - 6*a^3*b^5*c^7*d - 6*a^7*b*c^3*d^5 + 1 \\ & 5*a^4*b^4*c^6*d^2 - 20*a^5*b^3*c^5*d^3 + 15*a^6*b^2*c^4*d^4) + (x*(a*d - 5* \\ & b*c)*(-c^3*d^3)^{(1/2)}*(16*a^2*b^9*c^9*d^2 - 80*a^3*b^8*c^8*d^3 + 144*a^4*b^ \\ & 7*c^7*d^4 - 80*a^5*b^6*c^6*d^5 - 80*a^6*b^5*c^5*d^6 + 144*a^7*b^4*c^4*d^7 - \\ & 80*a^8*b^3*c^3*d^8 + 16*a^9*b^2*c^2*d^9))/(8*(b^3*c^6 - a^3*c^3*d^3 + 3*a^ \\ & 2*b*c^4*d^2 - 3*a*b^2*c^5*d))*(a^2*b^4*c^6 + a^6*c^2*d^4 - 4*a^3*b^3*c^5*d - \\ & 4*a^5*b*c^3*d^3 + 6*a^4*b^2*c^4*d^2)))*(a*d - 5*b*c)*(-c^3*d^3)^{(1/2)})/(4* \\ & (b^3*c^6 - a^3*c^3*d^3 + 3*a^2*b*c^4*d^2 - 3*a*b^2*c^5*d)))*(a*d - 5*b*c)*(- \\ & c^3*d^3)^{(1/2)})/(4*(b^3*c^6 - a^3*c^3*d^3 + 3*a^2*b*c^4*d^2 - 3*a*b^2*c^5* \\ & d)))*(-c^3*d^3)^{(1/2)})/(4*(b^3*c^6 - a^3*c^3*d^3 + 3*a^2*b*c^4*d^2 - 3*a*b^2*c^5* \\ & d)))*(-c^3*d^3)^{(1/2)}*i)/(2*(b^3*c^6 - a^3*c^3*d^3 + 3*a^2*b*c^4*d^2 - 3*a*b^2*c^5*d)) \end{aligned}$$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**2/(d*x**2+c)**2,x)

[Out] Timed out

$$3.34 \quad \int \frac{1}{(a+bx^2)^2(c+dx^2)^3} dx$$

Optimal. Leaf size=230

$$\frac{b^{5/2}(bc-7ad)\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}(bc-ad)^4} + \frac{d^{3/2}(3a^2d^2-14abcd+35b^2c^2)\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}(bc-ad)^4} + \frac{dx(4bc-ad)(3ad+bc)}{8ac^2(c+dx^2)(bc-ad)^3} + \frac{1}{2a(a+bx^2)}$$

[Out] $1/4*d*(a*d+2*b*c)*x/a/c/(-a*d+b*c)^2/(d*x^2+c)^2+1/2*b*x/a/(-a*d+b*c)/(b*x^2+a)/(d*x^2+c)^2+1/8*d*(-a*d+4*b*c)*(3*a*d+b*c)*x/a/c^2/(-a*d+b*c)^3/(d*x^2+c)+1/2*b^(5/2)*(-7*a*d+b*c)*\arctan(x*b^(1/2)/a^(1/2))/a^(3/2)/(-a*d+b*c)^4+1/8*d^(3/2)*(3*a^2*d^2-14*a*b*c*d+35*b^2*c^2)*\arctan(x*d^(1/2)/c^(1/2))/c^(5/2)/(-a*d+b*c)^4$

Rubi [A] time = 0.31, antiderivative size = 230, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {414, 527, 522, 205}

$$\frac{d^{3/2}(3a^2d^2-14abcd+35b^2c^2)\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{8c^{5/2}(bc-ad)^4} + \frac{b^{5/2}(bc-7ad)\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2a^{3/2}(bc-ad)^4} + \frac{dx(4bc-ad)(3ad+bc)}{8ac^2(c+dx^2)(bc-ad)^3} + \frac{1}{2a(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^2*(c + d*x^2)^3), x]

[Out] $(d*(2*b*c + a*d)*x)/(4*a*c*(b*c - a*d)^2*(c + d*x^2)^2) + (b*x)/(2*a*(b*c - a*d)*(a + b*x^2)*(c + d*x^2)^2) + (d*(4*b*c - a*d)*(b*c + 3*a*d)*x)/(8*a*c^2*(b*c - a*d)^3*(c + d*x^2)) + (b^(5/2)*(b*c - 7*a*d)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]])/(2*a^(3/2)*(b*c - a*d)^4) + (d^(3/2)*(35*b^2*c^2 - 14*a*b*c*d + 3*a^2*d^2)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]])/(8*c^(5/2)*(b*c - a*d)^4)$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 522

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(b*c - a*d)*(p+1)), x] + Dist[1/(a*n*(b*c - a*d)*(p

+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + bx^2)^2 (c + dx^2)^3} dx &= \frac{bx}{2a(bc - ad)(a + bx^2)(c + dx^2)^2} - \frac{\int \frac{-bc + 2ad - 5bdx^2}{(a + bx^2)(c + dx^2)^3} dx}{2a(bc - ad)} \\ &= \frac{d(2bc + ad)x}{4ac(bc - ad)^2 (c + dx^2)^2} + \frac{bx}{2a(bc - ad)(a + bx^2)(c + dx^2)^2} - \frac{\int \frac{-2(2b^2c^2 - 8abcd + 3a^2d^2)}{(a + bx^2)(c + dx^2)^3} dx}{8ac(bc - ad)^3} \\ &= \frac{d(2bc + ad)x}{4ac(bc - ad)^2 (c + dx^2)^2} + \frac{bx}{2a(bc - ad)(a + bx^2)(c + dx^2)^2} + \frac{d(4bc - ad)(bc + 3ad)}{8ac^2(bc - ad)^3 (c + dx^2)^2} \\ &= \frac{d(2bc + ad)x}{4ac(bc - ad)^2 (c + dx^2)^2} + \frac{bx}{2a(bc - ad)(a + bx^2)(c + dx^2)^2} + \frac{d(4bc - ad)(bc + 3ad)}{8ac^2(bc - ad)^3 (c + dx^2)^2} \\ &= \frac{d(2bc + ad)x}{4ac(bc - ad)^2 (c + dx^2)^2} + \frac{bx}{2a(bc - ad)(a + bx^2)(c + dx^2)^2} + \frac{d(4bc - ad)(bc + 3ad)}{8ac^2(bc - ad)^3 (c + dx^2)^2} \end{aligned}$$

Mathematica [A] time = 0.42, size = 197, normalized size = 0.86

$$\frac{1}{8} \left(\frac{4b^{5/2}(bc - 7ad) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{a^{3/2}(bc - ad)^4} + \frac{d^{3/2}(3a^2d^2 - 14abcd + 35b^2c^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{c^{5/2}(bc - ad)^4} - \frac{4b^3x}{a(a + bx^2)(ad - bc)^3} + \frac{d^2x}{c^2(c + dx^2)} \right)$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^2*(c + d*x^2)^3),x]

[Out] ((-4*b^3*x)/(a*(-(b*c) + a*d)^3*(a + b*x^2)) + (2*d^2*x)/(c*(b*c - a*d)^2*(c + d*x^2)^2) + (d^2*(11*b*c - 3*a*d)*x)/(c^2*(b*c - a*d)^3*(c + d*x^2)) + (4*b^(5/2)*(b*c - 7*a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(a^(3/2)*(b*c - a*d)^4) + (d^(3/2)*(35*b^2*c^2 - 14*a*b*c*d + 3*a^2*d^2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(c^(5/2)*(b*c - a*d)^4))/8

fricas [B] time = 6.07, size = 3239, normalized size = 14.08

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [1/16*(2*(4*b^4*c^3*d^2 + 7*a*b^3*c^2*d^3 - 14*a^2*b^2*c*d^4 + 3*a^3*b*d^5)*x^5 + 2*(8*b^4*c^4*d + 5*a*b^3*c^3*d^2 - 7*a^2*b^2*c^2*d^3 - 9*a^3*b*c*d^4 + 3*a^4*d^5)*x^3 - 4*(a*b^3*c^5 - 7*a^2*b^2*c^4*d + (b^4*c^3*d^2 - 7*a*b^3*c^2*d^3)*x^6 + (2*b^4*c^4*d - 13*a*b^3*c^3*d^2 - 7*a^2*b^2*c^2*d^3)*x^4 + (b^4*c^5 - 5*a*b^3*c^4*d - 14*a^2*b^2*c^3*d^2)*x^2)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + (35*a^2*b^2*c^4*d - 14*a^3*b*c^3*d^2 + 3*a^4*c^2*d^3 + (35*a*b^3*c^2*d^3 - 14*a^2*b^2*c*d^4 + 3*a^3*b*d^5)*x^6 +

$$\begin{aligned}
& (70*a*b^3*c^3*d^2 + 7*a^2*b^2*c^2*d^3 - 8*a^3*b*c*d^4 + 3*a^4*d^5)*x^4 + (\\
& 35*a*b^3*c^4*d + 56*a^2*b^2*c^3*d^2 - 25*a^3*b*c^2*d^3 + 6*a^4*c*d^4)*x^2)* \\
& \text{sqrt}(-d/c)*\log((d*x^2 + 2*c*x*\text{sqrt}(-d/c) - c)/(d*x^2 + c)) + 2*(4*b^4*c^5 - \\
& 4*a*b^3*c^4*d + 13*a^2*b^2*c^3*d^2 - 18*a^3*b*c^2*d^3 + 5*a^4*c*d^4)*x)/(a \\
& ^2*b^4*c^8 - 4*a^3*b^3*c^7*d + 6*a^4*b^2*c^6*d^2 - 4*a^5*b*c^5*d^3 + a^6*c^4 \\
& ^4*d^4 + (a*b^5*c^6*d^2 - 4*a^2*b^4*c^5*d^3 + 6*a^3*b^3*c^4*d^4 - 4*a^4*b^2*c^3 \\
& ^3*d^5 + a^5*b*c^2*d^6)*x^6 + (2*a*b^5*c^7*d - 7*a^2*b^4*c^6*d^2 + 8*a^3*b^3 \\
& ^3*c^5*d^3 - 2*a^4*b^2*c^4*d^4 - 2*a^5*b*c^3*d^5 + a^6*c^2*d^6)*x^4 + (a*b^5 \\
& ^5*c^8 - 2*a^2*b^4*c^7*d - 2*a^3*b^3*c^6*d^2 + 8*a^4*b^2*c^5*d^3 - 7*a^5*b*c^4 \\
& ^4*d^4 + 2*a^6*c^3*d^5)*x^2), 1/8*((4*b^4*c^3*d^2 + 7*a*b^3*c^2*d^3 - 14*a^2 \\
& ^2*b^2*c*d^4 + 3*a^3*b*d^5)*x^5 + (8*b^4*c^4*d + 5*a*b^3*c^3*d^2 - 7*a^2*b^2 \\
& ^2*c^2*d^3 - 9*a^3*b*c*d^4 + 3*a^4*d^5)*x^3 + (35*a^2*b^2*c^4*d - 14*a^3*b*c^3 \\
& ^3*d^2 + 3*a^4*c^2*d^3 + (35*a*b^3*c^2*d^3 - 14*a^2*b^2*c*d^4 + 3*a^3*b*d^5) \\
& ^5)*x^6 + (70*a*b^3*c^3*d^2 + 7*a^2*b^2*c^2*d^3 - 8*a^3*b*c*d^4 + 3*a^4*d^5)*x \\
& ^4 + (35*a*b^3*c^4*d + 56*a^2*b^2*c^3*d^2 - 25*a^3*b*c^2*d^3 + 6*a^4*c*d^4) \\
& ^2)*\text{sqrt}(d/c)*\arctan(x*\text{sqrt}(d/c)) - 2*(a*b^3*c^5 - 7*a^2*b^2*c^4*d + (b^4 \\
& ^4*c^3*d^2 - 7*a*b^3*c^2*d^3)*x^6 + (2*b^4*c^4*d - 13*a*b^3*c^3*d^2 - 7*a^2*b^2 \\
& ^2*c^2*d^3)*x^4 + (b^4*c^5 - 5*a*b^3*c^4*d - 14*a^2*b^2*c^3*d^2)*x^2)*\text{sqrt}(- \\
& b/a)*\log((b*x^2 - 2*a*x*\text{sqrt}(-b/a) - a)/(b*x^2 + a)) + (4*b^4*c^5 - 4*a*b^3 \\
& ^3*c^4*d + 13*a^2*b^2*c^3*d^2 - 18*a^3*b*c^2*d^3 + 5*a^4*c*d^4)*x)/(a^2*b^4*c^8 \\
& - 4*a^3*b^3*c^7*d + 6*a^4*b^2*c^6*d^2 - 4*a^5*b*c^5*d^3 + a^6*c^4*d^4 + \\
& (a*b^5*c^6*d^2 - 4*a^2*b^4*c^5*d^3 + 6*a^3*b^3*c^4*d^4 - 4*a^4*b^2*c^3*d^5 \\
& + a^5*b*c^2*d^6)*x^6 + (2*a*b^5*c^7*d - 7*a^2*b^4*c^6*d^2 + 8*a^3*b^3*c^5 \\
& ^5*d^3 - 2*a^4*b^2*c^4*d^4 - 2*a^5*b*c^3*d^5 + a^6*c^2*d^6)*x^4 + (a*b^5*c^8 - \\
& 2*a^2*b^4*c^7*d - 2*a^3*b^3*c^6*d^2 + 8*a^4*b^2*c^5*d^3 - 7*a^5*b*c^4*d^4 \\
& + 2*a^6*c^3*d^5)*x^2), 1/16*(2*(4*b^4*c^3*d^2 + 7*a*b^3*c^2*d^3 - 14*a^2*b^2 \\
& ^2*c*d^4 + 3*a^3*b*d^5)*x^5 + 2*(8*b^4*c^4*d + 5*a*b^3*c^3*d^2 - 7*a^2*b^2*c^2 \\
& ^2*d^3 - 9*a^3*b*c*d^4 + 3*a^4*d^5)*x^3 + 8*(a*b^3*c^5 - 7*a^2*b^2*c^4*d + \\
& (b^4*c^3*d^2 - 7*a*b^3*c^2*d^3)*x^6 + (2*b^4*c^4*d - 13*a*b^3*c^3*d^2 - 7*a^2 \\
& ^2*b^2*c^2*d^3)*x^4 + (b^4*c^5 - 5*a*b^3*c^4*d - 14*a^2*b^2*c^3*d^2)*x^2)*\text{s} \\
& \text{qrt}(b/a)*\arctan(x*\text{sqrt}(b/a)) + (35*a^2*b^2*c^4*d - 14*a^3*b*c^3*d^2 + 3*a^4 \\
& ^4*c^2*d^3 + (35*a*b^3*c^2*d^3 - 14*a^2*b^2*c*d^4 + 3*a^3*b*d^5)*x^6 + (70*a \\
& ^70*a*b^3*c^3*d^2 + 7*a^2*b^2*c^2*d^3 - 8*a^3*b*c*d^4 + 3*a^4*d^5)*x^4 + (35*a*b^3 \\
& ^3*c^4*d + 56*a^2*b^2*c^3*d^2 - 25*a^3*b*c^2*d^3 + 6*a^4*c*d^4)*x^2)*\text{sqrt}(-d \\
& /c)*\log((d*x^2 + 2*c*x*\text{sqrt}(-d/c) - c)/(d*x^2 + c)) + 2*(4*b^4*c^5 - 4*a*b^3 \\
& ^3*c^4*d + 13*a^2*b^2*c^3*d^2 - 18*a^3*b*c^2*d^3 + 5*a^4*c*d^4)*x)/(a^2*b^4*c^8 \\
& - 4*a^3*b^3*c^7*d + 6*a^4*b^2*c^6*d^2 - 4*a^5*b*c^5*d^3 + a^6*c^4*d^4 + \\
& (a*b^5*c^6*d^2 - 4*a^2*b^4*c^5*d^3 + 6*a^3*b^3*c^4*d^4 - 4*a^4*b^2*c^3*d^5 \\
& + a^5*b*c^2*d^6)*x^6 + (2*a*b^5*c^7*d - 7*a^2*b^4*c^6*d^2 + 8*a^3*b^3*c^5 \\
& ^5*d^3 - 2*a^4*b^2*c^4*d^4 - 2*a^5*b*c^3*d^5 + a^6*c^2*d^6)*x^4 + (a*b^5*c^8 - \\
& 2*a^2*b^4*c^7*d - 2*a^3*b^3*c^6*d^2 + 8*a^4*b^2*c^5*d^3 - 7*a^5*b*c^4*d^4 \\
& + 2*a^6*c^3*d^5)*x^2), 1/8*((4*b^4*c^3*d^2 + 7*a*b^3*c^2*d^3 - 14*a^2*b^2*c \\
& ^2*d^4 + 3*a^3*b*d^5)*x^5 + (8*b^4*c^4*d + 5*a*b^3*c^3*d^2 - 7*a^2*b^2*c^2*d^3 \\
& ^2*d^3 - 9*a^3*b*c*d^4 + 3*a^4*d^5)*x^3 + 4*(a*b^3*c^5 - 7*a^2*b^2*c^4*d + (b^4 \\
& ^4*c^3*d^2 - 7*a*b^3*c^2*d^3)*x^6 + (2*b^4*c^4*d - 13*a*b^3*c^3*d^2 - 7*a^2*b^2 \\
& ^2*c^2*d^3)*x^4 + (b^4*c^5 - 5*a*b^3*c^4*d - 14*a^2*b^2*c^3*d^2)*x^2)*\text{sqrt}(b \\
& /a)*\arctan(x*\text{sqrt}(b/a)) + (35*a^2*b^2*c^4*d - 14*a^3*b*c^3*d^2 + 3*a^4*c^2 \\
& ^2*d^3 + (35*a*b^3*c^2*d^3 - 14*a^2*b^2*c*d^4 + 3*a^3*b*d^5)*x^6 + (70*a*b^3 \\
& ^3*c^3*d^2 + 7*a^2*b^2*c^2*d^3 - 8*a^3*b*c*d^4 + 3*a^4*d^5)*x^4 + (35*a*b^3 \\
& ^3*c^4*d + 56*a^2*b^2*c^3*d^2 - 25*a^3*b*c^2*d^3 + 6*a^4*c*d^4)*x^2)*\text{sqrt}(d/c)*\text{a} \\
& \text{rctan}(x*\text{sqrt}(d/c)) + (4*b^4*c^5 - 4*a*b^3*c^4*d + 13*a^2*b^2*c^3*d^2 - 18*a^3 \\
& ^3*b*c^2*d^3 + 5*a^4*c*d^4)*x)/(a^2*b^4*c^8 - 4*a^3*b^3*c^7*d + 6*a^4*b^2*c^6 \\
& ^6*d^2 - 4*a^5*b*c^5*d^3 + a^6*c^4*d^4 + (a*b^5*c^6*d^2 - 4*a^2*b^4*c^5*d^3 \\
& + 6*a^3*b^3*c^4*d^4 - 4*a^4*b^2*c^3*d^5 + a^5*b*c^2*d^6)*x^6 + (2*a*b^5*c^7 \\
& ^7*d - 7*a^2*b^4*c^6*d^2 + 8*a^3*b^3*c^5*d^3 - 2*a^4*b^2*c^4*d^4 - 2*a^5*b*c^3 \\
& ^3*d^5 + a^6*c^2*d^6)*x^4 + (a*b^5*c^8 - 2*a^2*b^4*c^7*d - 2*a^3*b^3*c^6*d^2 \\
& + 8*a^4*b^2*c^5*d^3 - 7*a^5*b*c^4*d^4 + 2*a^6*c^3*d^5)*x^2)]
\end{aligned}$$

giac [A] time = 0.58, size = 332, normalized size = 1.44

$$\frac{b^3x}{2(ab^3c^3 - 3a^2b^2c^2d + 3a^3bcd^2 - a^4d^3)(bx^2 + a)} + \frac{(b^4c - 7ab^3d) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ab^4c^4 - 4a^2b^3c^3d + 6a^3b^2c^2d^2 - 4a^4bcd^3 + a^5d^4)\sqrt{ab}} + \dots$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)^3,x, algorithm="giac")
```

```
[Out] 1/2*b^3*x/((a*b^3*c^3 - 3*a^2*b^2*c^2*d + 3*a^3*b*c*d^2 - a^4*d^3)*(b*x^2 + a)) + 1/2*(b^4*c - 7*a*b^3*d)*arctan(b*x/sqrt(a*b))/((a*b^4*c^4 - 4*a^2*b^3*c^3*d + 6*a^3*b^2*c^2*d^2 - 4*a^4*b*c*d^3 + a^5*d^4)*sqrt(a*b)) + 1/8*(35*b^2*c^2*d^2 - 14*a*b*c*d^3 + 3*a^2*d^4)*arctan(d*x/sqrt(c*d))/((b^4*c^6 - 4*a*b^3*c^5*d + 6*a^2*b^2*c^4*d^2 - 4*a^3*b*c^3*d^3 + a^4*c^2*d^4)*sqrt(c*d)) + 1/8*(11*b*c*d^3*x^3 - 3*a*d^4*x^3 + 13*b*c^2*d^2*x - 5*a*c*d^3*x)/((b^3*c^5 - 3*a*b^2*c^4*d + 3*a^2*b*c^3*d^2 - a^3*c^2*d^3)*(d*x^2 + c)^2)
```

maple [A] time = 0.02, size = 403, normalized size = 1.75

$$\frac{3a^2d^5x^3}{8(ad - bc)^4(dx^2 + c)^2c^2} - \frac{7abd^4x^3}{4(ad - bc)^4(dx^2 + c)^2c} + \frac{11b^2d^3x^3}{8(ad - bc)^4(dx^2 + c)^2} + \frac{5a^2d^4x}{8(ad - bc)^4(dx^2 + c)^2c} - \dots$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(b*x^2+a)^2/(d*x^2+c)^3,x)
```

```
[Out] 3/8*d^5/(a*d-b*c)^4/(d*x^2+c)^2/c^2*x^3*a^2-7/4*d^4/(a*d-b*c)^4/(d*x^2+c)^2/c*x^3*a*b+11/8*d^3/(a*d-b*c)^4/(d*x^2+c)^2*x^3*b^2+5/8*d^4/(a*d-b*c)^4/(d*x^2+c)^2/c*x*a^2-9/4*d^3/(a*d-b*c)^4/(d*x^2+c)^2*x*a*b+13/8*d^2/(a*d-b*c)^4/(d*x^2+c)^2*c*x*b^2+3/8*d^4/(a*d-b*c)^4/c^2/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a^2-7/4*d^3/(a*d-b*c)^4/c/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*a*b+35/8*d^2/(a*d-b*c)^4/(c*d)^(1/2)*arctan(1/(c*d)^(1/2)*d*x)*b^2-1/2*b^3/(a*d-b*c)^4*x/(b*x^2+a)*d+1/2*b^4/(a*d-b*c)^4/a*x/(b*x^2+a)*c-7/2*b^3/(a*d-b*c)^4/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*d+1/2*b^4/(a*d-b*c)^4/a/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c
```

maxima [B] time = 3.41, size = 529, normalized size = 2.30

$$\frac{(b^4c - 7ab^3d) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{2(ab^4c^4 - 4a^2b^3c^3d + 6a^3b^2c^2d^2 - 4a^4bcd^3 + a^5d^4)\sqrt{ab}} + \frac{(35b^2c^2d^2 - 14abcd^3 + 3a^2d^4) \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{8(b^4c^6 - 4ab^3c^5d + 6a^2b^2c^4d^2 - 4a^3bc^3d^3 + a^4c^2d^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)^3,x, algorithm="maxima")
```

```
[Out] 1/2*(b^4*c - 7*a*b^3*d)*arctan(b*x/sqrt(a*b))/((a*b^4*c^4 - 4*a^2*b^3*c^3*d + 6*a^3*b^2*c^2*d^2 - 4*a^4*b*c*d^3 + a^5*d^4)*sqrt(a*b)) + 1/8*(35*b^2*c^2*d^2 - 14*a*b*c*d^3 + 3*a^2*d^4)*arctan(d*x/sqrt(c*d))/((b^4*c^6 - 4*a*b^3*c^5*d + 6*a^2*b^2*c^4*d^2 - 4*a^3*b*c^3*d^3 + a^4*c^2*d^4)*sqrt(c*d)) + 1/8*((4*b^3*c^2*d^2 + 11*a*b^2*c*d^3 - 3*a^2*b*d^4)*x^5 + (8*b^3*c^3*d + 13*a*b^2*c^2*d^2 + 6*a^2*b*c*d^3 - 3*a^3*d^4)*x^3 + (4*b^3*c^4 + 13*a^2*b*c^2*d^2 - 5*a^3*c*d^3)*x)/(a^2*b^3*c^7 - 3*a^3*b^2*c^6*d + 3*a^4*b*c^5*d^2 - a^5*c^4*d^3 + (a*b^4*c^5*d^2 - 3*a^2*b^3*c^4*d^3 + 3*a^3*b^2*c^3*d^4 - a^4*b*c^2*d^5)*x^6 + (2*a*b^4*c^6*d - 5*a^2*b^3*c^5*d^2 + 3*a^3*b^2*c^4*d^3 + a^4*b*c^3*d^4 - a^5*c^2*d^5)*x^4 + (a*b^4*c^7 - a^2*b^3*c^6*d - 3*a^3*b^2*c^5*d^2 + 5*a^4*b*c^4*d^3 - 2*a^5*c^3*d^4)*x^2)
```

mupad [B] time = 7.79, size = 8649, normalized size = 37.60

result too large to display

$$\begin{aligned}
& 4*c^8*d^5 + 84*a^8*b^3*c^7*d^6 - 36*a^9*b^2*c^6*d^7) - (x*(-c^5*d^3)^{(1/2)}* \\
& (3*a^2*d^2 + 35*b^2*c^2 - 14*a*b*c*d)*(256*a^2*b^11*c^13*d^2 - 1792*a^3*b^1 \\
& 0*c^12*d^3 + 5120*a^4*b^9*c^11*d^4 - 7168*a^5*b^8*c^10*d^5 + 3584*a^6*b^7*c \\
& ^9*d^6 + 3584*a^7*b^6*c^8*d^7 - 7168*a^8*b^5*c^7*d^8 + 5120*a^9*b^4*c^6*d^9 \\
& - 1792*a^10*b^3*c^5*d^10 + 256*a^11*b^2*c^4*d^11))/(512*(b^4*c^9 + a^4*c^5 \\
& *d^4 - 4*a^3*b*c^6*d^3 + 6*a^2*b^2*c^7*d^2 - 4*a*b^3*c^8*d)*(a^2*b^6*c^10 + \\
& a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6*a^7*b*c^5*d^5 + 15*a^4*b^4*c^8*d^2 - 20* \\
& a^5*b^3*c^7*d^3 + 15*a^6*b^2*c^6*d^4)))*(-c^5*d^3)^{(1/2)}*(3*a^2*d^2 + 35*b^ \\
& 2*c^2 - 14*a*b*c*d))/(16*(b^4*c^9 + a^4*c^5*d^4 - 4*a^3*b*c^6*d^3 + 6*a^2*b \\
& ^2*c^7*d^2 - 4*a*b^3*c^8*d)))*(-c^5*d^3)^{(1/2)}*(3*a^2*d^2 + 35*b^2*c^2 - 14 \\
& *a*b*c*d))/(16*(b^4*c^9 + a^4*c^5*d^4 - 4*a^3*b*c^6*d^3 + 6*a^2*b^2*c^7*d^2 \\
& - 4*a*b^3*c^8*d)) + (((x*(9*a^6*b^3*d^9 + 16*b^9*c^6*d^3 - 224*a*b^8*c^5*d \\
& ^4 - 84*a^5*b^4*c*d^8 + 2009*a^2*b^7*c^4*d^5 - 980*a^3*b^6*c^3*d^6 + 406*a^ \\
& 4*b^5*c^2*d^7))/(32*(a^2*b^6*c^10 + a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6*a^7*b \\
& *c^5*d^5 + 15*a^4*b^4*c^8*d^2 - 20*a^5*b^3*c^7*d^3 + 15*a^6*b^2*c^6*d^4)) + \\
& ((2*a*b^13*c^13*d^2 - 28*a^2*b^12*c^12*d^3 + (315*a^3*b^11*c^11*d^4)/2 - \\
& (987*a^4*b^10*c^10*d^5)/2 + 978*a^5*b^9*c^9*d^6 - 1302*a^6*b^8*c^8*d^7 + 11 \\
& 97*a^7*b^7*c^7*d^8 - 765*a^8*b^6*c^6*d^9 + 336*a^9*b^5*c^5*d^10 - 98*a^10*b \\
& ^4*c^4*d^11 + (35*a^11*b^3*c^3*d^12)/2 - (3*a^12*b^2*c^2*d^13)/2)/(a^2*b^9* \\
& c^13 - a^11*c^4*d^9 - 9*a^3*b^8*c^12*d + 9*a^10*b*c^5*d^8 + 36*a^4*b^7*c^11 \\
& *d^2 - 84*a^5*b^6*c^10*d^3 + 126*a^6*b^5*c^9*d^4 - 126*a^7*b^4*c^8*d^5 + 84 \\
& *a^8*b^3*c^7*d^6 - 36*a^9*b^2*c^6*d^7) + (x*(-c^5*d^3)^{(1/2)}*(3*a^2*d^2 + 3 \\
& 5*b^2*c^2 - 14*a*b*c*d)*(256*a^2*b^11*c^13*d^2 - 1792*a^3*b^10*c^12*d^3 + 5 \\
& 120*a^4*b^9*c^11*d^4 - 7168*a^5*b^8*c^10*d^5 + 3584*a^6*b^7*c^9*d^6 + 3584* \\
& a^7*b^6*c^8*d^7 - 7168*a^8*b^5*c^7*d^8 + 5120*a^9*b^4*c^6*d^9 - 1792*a^10*b \\
& ^3*c^5*d^10 + 256*a^11*b^2*c^4*d^11))/(512*(b^4*c^9 + a^4*c^5*d^4 - 4*a^3*b \\
& *c^6*d^3 + 6*a^2*b^2*c^7*d^2 - 4*a*b^3*c^8*d)*(a^2*b^6*c^10 + a^8*c^4*d^6 - \\
& 6*a^3*b^5*c^9*d - 6*a^7*b*c^5*d^5 + 15*a^4*b^4*c^8*d^2 - 20*a^5*b^3*c^7*d^ \\
& 3 + 15*a^6*b^2*c^6*d^4)))*(-c^5*d^3)^{(1/2)}*(3*a^2*d^2 + 35*b^2*c^2 - 14*a*b \\
& *c*d))/(16*(b^4*c^9 + a^4*c^5*d^4 - 4*a^3*b*c^6*d^3 + 6*a^2*b^2*c^7*d^2 - 4 \\
& *a*b^3*c^8*d)))*(-c^5*d^3)^{(1/2)}*(3*a^2*d^2 + 35*b^2*c^2 - 14*a*b*c*d))/(16 \\
& *(b^4*c^9 + a^4*c^5*d^4 - 4*a^3*b*c^6*d^3 + 6*a^2*b^2*c^7*d^2 - 4*a*b^3*c^8 \\
& *d)))*(-c^5*d^3)^{(1/2)}*(3*a^2*d^2 + 35*b^2*c^2 - 14*a*b*c*d)*1i)/(8*(b^4*c \\
& ^9 + a^4*c^5*d^4 - 4*a^3*b*c^6*d^3 + 6*a^2*b^2*c^7*d^2 - 4*a*b^3*c^8*d)) - \\
& ((x^5*(4*b^3*c^2*d^2 - 3*a^2*b*d^4 + 11*a*b^2*c*d^3))/(8*a*c^2*(a^3*d^3 - b \\
& ^3*c^3 + 3*a*b^2*c^2*d - 3*a^2*b*c*d^2)) + (x*(4*b^3*c^3 - 5*a^3*d^3 + 13*a \\
& ^2*b*c*d^2))/(8*a*c*(a*d - b*c)*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d)) + (d*x^3*(\\
& 8*b^3*c^3 - 3*a^3*d^3 + 13*a*b^2*c^2*d + 6*a^2*b*c*d^2))/(8*a*c^2*(a*d - b* \\
& c)*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d)))/(a*c^2 + x^2*(b*c^2 + 2*a*c*d) + x^4*(\\
& a*d^2 + 2*b*c*d) + b*d^2*x^6) + (atan((((x*(9*a^6*b^3*d^9 + 16*b^9*c^6*d^3 \\
& - 224*a*b^8*c^5*d^4 - 84*a^5*b^4*c*d^8 + 2009*a^2*b^7*c^4*d^5 - 980*a^3*b^ \\
& 6*c^3*d^6 + 406*a^4*b^5*c^2*d^7))/(32*(a^2*b^6*c^10 + a^8*c^4*d^6 - 6*a^3*b \\
& ^5*c^9*d - 6*a^7*b*c^5*d^5 + 15*a^4*b^4*c^8*d^2 - 20*a^5*b^3*c^7*d^3 + 15*a \\
& ^6*b^2*c^6*d^4)) - ((7*a*d - b*c)*(-a^3*b^5)^{(1/2)}*((2*a*b^13*c^13*d^2 - 28 \\
& *a^2*b^12*c^12*d^3 + (315*a^3*b^11*c^11*d^4)/2 - (987*a^4*b^10*c^10*d^5)/2 \\
& + 978*a^5*b^9*c^9*d^6 - 1302*a^6*b^8*c^8*d^7 + 1197*a^7*b^7*c^7*d^8 - 765*a \\
& ^8*b^6*c^6*d^9 + 336*a^9*b^5*c^5*d^10 - 98*a^10*b^4*c^4*d^11 + (35*a^11*b^3 \\
& *c^3*d^12)/2 - (3*a^12*b^2*c^2*d^13)/2)/(a^2*b^9*c^13 - a^11*c^4*d^9 - 9*a^ \\
& 3*b^8*c^12*d + 9*a^10*b*c^5*d^8 + 36*a^4*b^7*c^11*d^2 - 84*a^5*b^6*c^10*d^3 \\
& + 126*a^6*b^5*c^9*d^4 - 126*a^7*b^4*c^8*d^5 + 84*a^8*b^3*c^7*d^6 - 36*a^9* \\
& b^2*c^6*d^7) - (x*(7*a*d - b*c)*(-a^3*b^5)^{(1/2)}*(256*a^2*b^11*c^13*d^2 - 1 \\
& 792*a^3*b^10*c^12*d^3 + 5120*a^4*b^9*c^11*d^4 - 7168*a^5*b^8*c^10*d^5 + 358 \\
& 4*a^6*b^7*c^9*d^6 + 3584*a^7*b^6*c^8*d^7 - 7168*a^8*b^5*c^7*d^8 + 5120*a^9* \\
& b^4*c^6*d^9 - 1792*a^10*b^3*c^5*d^10 + 256*a^11*b^2*c^4*d^11))/(128*(a^7*d^ \\
& 4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*b*c*d^3)*(a^2 \\
& *b^6*c^10 + a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6*a^7*b*c^5*d^5 + 15*a^4*b^4*c^ \\
& 8*d^2 - 20*a^5*b^3*c^7*d^3 + 15*a^6*b^2*c^6*d^4)))/(4*(a^7*d^4 + a^3*b^4*c \\
& ^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*b*c*d^3)))*(7*a*d - b*c)* \\
& (-a^3*b^5)^{(1/2)}*1i)/(4*(a^7*d^4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2
\end{aligned}$$

$$\begin{aligned}
& *c^2*d^2 - 4*a^6*b*c*d^3)) + (((x*(9*a^6*b^3*d^9 + 16*b^9*c^6*d^3 - 224*a*b^8*c^5*d^4 - 84*a^5*b^4*c*d^8 + 2009*a^2*b^7*c^4*d^5 - 980*a^3*b^6*c^3*d^6 \\
& + 406*a^4*b^5*c^2*d^7)))/(32*(a^2*b^6*c^10 + a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6*a^7*b*c^5*d^5 + 15*a^4*b^4*c^8*d^2 - 20*a^5*b^3*c^7*d^3 + 15*a^6*b^2*c^6 \\
& *d^4)) + ((7*a*d - b*c)*(-a^3*b^5)^(1/2))*((2*a*b^13*c^13*d^2 - 28*a^2*b^12*c^12*d^3 + (315*a^3*b^11*c^11*d^4)/2 - (987*a^4*b^10*c^10*d^5)/2 + 978*a^5* \\
& b^9*c^9*d^6 - 1302*a^6*b^8*c^8*d^7 + 1197*a^7*b^7*c^7*d^8 - 765*a^8*b^6*c^6*d^9 + 336*a^9*b^5*c^5*d^10 - 98*a^10*b^4*c^4*d^11 + (35*a^11*b^3*c^3*d^12) \\
& /2 - (3*a^12*b^2*c^2*d^13)/2)/(a^2*b^9*c^13 - a^11*c^4*d^9 - 9*a^3*b^8*c^12*d + 9*a^10*b*c^5*d^8 + 36*a^4*b^7*c^11*d^2 - 84*a^5*b^6*c^10*d^3 + 126*a^6 \\
& *b^5*c^9*d^4 - 126*a^7*b^4*c^8*d^5 + 84*a^8*b^3*c^7*d^6 - 36*a^9*b^2*c^6*d^7) + (x*(7*a*d - b*c)*(-a^3*b^5)^(1/2))*(256*a^2*b^11*c^13*d^2 - 1792*a^3*b^ \\
& 10*c^12*d^3 + 5120*a^4*b^9*c^11*d^4 - 7168*a^5*b^8*c^10*d^5 + 3584*a^6*b^7*c^9*d^6 + 3584*a^7*b^6*c^8*d^7 - 7168*a^8*b^5*c^7*d^8 + 5120*a^9*b^4*c^6*d^ \\
& 9 - 1792*a^10*b^3*c^5*d^10 + 256*a^11*b^2*c^4*d^11))/(128*(a^7*d^4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*b*c*d^3)*(a^2*b^6*c^10 \\
& + a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6*a^7*b*c^5*d^5 + 15*a^4*b^4*c^8*d^2 - 20*a^5*b^3*c^7*d^3 + 15*a^6*b^2*c^6*d^4)))/(4*(a^7*d^4 + a^3*b^4*c^4 - 4*a^4 \\
& *b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*b*c*d^3))*(7*a*d - b*c)*(-a^3*b^5)^(1/2)*1i)/(4*(a^7*d^4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - \\
& 4*a^6*b*c*d^3)))/(((63*a^5*b^5*d^9)/64 + (35*b^10*c^5*d^4)/16 - (651*a*b^9*c^4*d^5)/64 - (267*a^4*b^6*c*d^8)/32 - (1275*a^2*b^8*c^3*d^6)/32 + (451*a^ \\
& 3*b^7*c^2*d^7)/16)/(a^2*b^9*c^13 - a^11*c^4*d^9 - 9*a^3*b^8*c^12*d + 9*a^10*b*c^5*d^8 + 36*a^4*b^7*c^11*d^2 - 84*a^5*b^6*c^10*d^3 + 126*a^6*b^5*c^9*d^ \\
& 4 - 126*a^7*b^4*c^8*d^5 + 84*a^8*b^3*c^7*d^6 - 36*a^9*b^2*c^6*d^7) - (((x*(9*a^6*b^3*d^9 + 16*b^9*c^6*d^3 - 224*a*b^8*c^5*d^4 - 84*a^5*b^4*c*d^8 + 200 \\
& 9*a^2*b^7*c^4*d^5 - 980*a^3*b^6*c^3*d^6 + 406*a^4*b^5*c^2*d^7)))/(32*(a^2*b^6*c^10 + a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6*a^7*b*c^5*d^5 + 15*a^4*b^4*c^8*d^ \\
& 2 - 20*a^5*b^3*c^7*d^3 + 15*a^6*b^2*c^6*d^4)) - ((7*a*d - b*c)*(-a^3*b^5)^(1/2))*((2*a*b^13*c^13*d^2 - 28*a^2*b^12*c^12*d^3 + (315*a^3*b^11*c^11*d^4)/ \\
& 2 - (987*a^4*b^10*c^10*d^5)/2 + 978*a^5*b^9*c^9*d^6 - 1302*a^6*b^8*c^8*d^7 + 1197*a^7*b^7*c^7*d^8 - 765*a^8*b^6*c^6*d^9 + 336*a^9*b^5*c^5*d^10 - 98*a^ \\
& 10*b^4*c^4*d^11 + (35*a^11*b^3*c^3*d^12)/2 - (3*a^12*b^2*c^2*d^13)/2)/(a^2*b^9*c^13 - a^11*c^4*d^9 - 9*a^3*b^8*c^12*d + 9*a^10*b*c^5*d^8 + 36*a^4*b^7*c^ \\
& 11*d^2 - 84*a^5*b^6*c^10*d^3 + 126*a^6*b^5*c^9*d^4 - 126*a^7*b^4*c^8*d^5 + 84*a^8*b^3*c^7*d^6 - 36*a^9*b^2*c^6*d^7) - (x*(7*a*d - b*c)*(-a^3*b^5)^(1/ \\
& 2))*(256*a^2*b^11*c^13*d^2 - 1792*a^3*b^10*c^12*d^3 + 5120*a^4*b^9*c^11*d^4 - 7168*a^5*b^8*c^10*d^5 + 3584*a^6*b^7*c^9*d^6 + 3584*a^7*b^6*c^8*d^7 - 71 \\
& 68*a^8*b^5*c^7*d^8 + 5120*a^9*b^4*c^6*d^9 - 1792*a^10*b^3*c^5*d^10 + 256*a^ \\
& 11*b^2*c^4*d^11))/(128*(a^7*d^4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*b*c*d^3)*(a^2*b^6*c^10 + a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6 \\
& *a^7*b*c^5*d^5 + 15*a^4*b^4*c^8*d^2 - 20*a^5*b^3*c^7*d^3 + 15*a^6*b^2*c^6*d^4)))/((7*a*d - b*c)*(-a^3*b^5)^(1/2))/(4*(a^7*d^4 + a^3*b^4*c^4 - \\
& 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*b*c*d^3)) + (((x*(9*a^6*b^3*d^9 + 16*b^9*c^6*d^3 - 224*a*b^8*c^5*d^4 - 84*a^5*b^4*c*d^8 + 2009*a^2*b^7*c^4 \\
& *d^5 - 980*a^3*b^6*c^3*d^6 + 406*a^4*b^5*c^2*d^7)))/(32*(a^2*b^6*c^10 + a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6*a^7*b*c^5*d^5 + 15*a^4*b^4*c^8*d^2 - 20*a^5*b^ \\
& 3*c^7*d^3 + 15*a^6*b^2*c^6*d^4)) + ((7*a*d - b*c)*(-a^3*b^5)^(1/2))*((2*a*b^13*c^13*d^2 - 28*a^2*b^12*c^12*d^3 + (315*a^3*b^11*c^11*d^4)/2 - (987*a^4* \\
& b^10*c^10*d^5)/2 + 978*a^5*b^9*c^9*d^6 - 1302*a^6*b^8*c^8*d^7 + 1197*a^7*b^7*c^7*d^8 - 765*a^8*b^6*c^6*d^9 + 336*a^9*b^5*c^5*d^10 - 98*a^10*b^4*c^4*d^ \\
& 11 + (35*a^11*b^3*c^3*d^12)/2 - (3*a^12*b^2*c^2*d^13)/2)/(a^2*b^9*c^13 - a^ \\
& 11*c^4*d^9 - 9*a^3*b^8*c^12*d + 9*a^10*b*c^5*d^8 + 36*a^4*b^7*c^11*d^2 - 84 \\
& *a^5*b^6*c^10*d^3 + 126*a^6*b^5*c^9*d^4 - 126*a^7*b^4*c^8*d^5 + 84*a^8*b^3*c^7*d^6 - 36*a^9*b^2*c^6*d^7) + (x*(7*a*d - b*c)*(-a^3*b^5)^(1/2))*(256*a^2* \\
& b^11*c^13*d^2 - 1792*a^3*b^10*c^12*d^3 + 5120*a^4*b^9*c^11*d^4 - 7168*a^5*b^8*c^10*d^5 + 3584*a^6*b^7*c^9*d^6 + 3584*a^7*b^6*c^8*d^7 - 7168*a^8*b^5*c^ \\
& 7*d^8 + 5120*a^9*b^4*c^6*d^9 - 1792*a^10*b^3*c^5*d^10 + 256*a^11*b^2*c^4*d^
\end{aligned}$$

```

11)))/(128*(a^7*d^4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*
a^6*b*c*d^3)*(a^2*b^6*c^10 + a^8*c^4*d^6 - 6*a^3*b^5*c^9*d - 6*a^7*b*c^5*d^
5 + 15*a^4*b^4*c^8*d^2 - 20*a^5*b^3*c^7*d^3 + 15*a^6*b^2*c^6*d^4))))/(4*(a^
7*d^4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*b*c*d^3))
)*(7*a*d - b*c)*(-a^3*b^5)^(1/2))/(4*(a^7*d^4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3
*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*b*c*d^3)))*(7*a*d - b*c)*(-a^3*b^5)^(1/2)*1
i)/(2*(a^7*d^4 + a^3*b^4*c^4 - 4*a^4*b^3*c^3*d + 6*a^5*b^2*c^2*d^2 - 4*a^6*
b*c*d^3))

```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**2/(d*x**2+c)**3,x)

[Out] Timed out

$$3.35 \quad \int \frac{(c+dx^2)^5}{(a+bx^2)^3} dx$$

Optimal. Leaf size=196

$$\frac{x(17ad + 3bc)(bc - ad)^4}{8a^2b^5(a + bx^2)} + \frac{d^3x(6a^2d^2 - 15abcd + 10b^2c^2)}{b^5} + \frac{(63a^2d^2 + 14abcd + 3b^2c^2)(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}b^{11/2}}$$

[Out] $d^3*(6*a^2*d^2-15*a*b*c*d+10*b^2*c^2)*x/b^5+1/3*d^4*(-3*a*d+5*b*c)*x^3/b^4+1/5*d^5*x^5/b^3+1/4*(-a*d+b*c)^5*x/a/b^5/(b*x^2+a)^2+1/8*(-a*d+b*c)^4*(17*a*d+3*b*c)*x/a^2/b^5/(b*x^2+a)+1/8*(-a*d+b*c)^3*(63*a^2*d^2+14*a*b*c*d+3*b^2*c^2)*\arctan(x*b^(1/2)/a^(1/2))/a^(5/2)/b^(11/2)$

Rubi [A] time = 0.23, antiderivative size = 196, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {390, 1157, 385, 205}

$$\frac{d^3x(6a^2d^2 - 15abcd + 10b^2c^2)}{b^5} + \frac{(63a^2d^2 + 14abcd + 3b^2c^2)(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}b^{11/2}} + \frac{x(17ad + 3bc)(bc - ad)^4}{8a^2b^5(a + bx^2)}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^5/(a + b*x^2)^3, x]

[Out] $(d^3*(10*b^2*c^2 - 15*a*b*c*d + 6*a^2*d^2)*x)/b^5 + (d^4*(5*b*c - 3*a*d)*x^3)/(3*b^4) + (d^5*x^5)/(5*b^3) + ((b*c - a*d)^5*x)/(4*a*b^5*(a + b*x^2)^2) + ((b*c - a*d)^4*(3*b*c + 17*a*d)*x)/(8*a^2*b^5*(a + b*x^2)) + ((b*c - a*d)^3*(3*b^2*c^2 + 14*a*b*c*d + 63*a^2*d^2)*\text{ArcTan}[\text{Sqrt}[b]*x/\text{Sqrt}[a]])/(8*a^(5/2)*b^(11/2))$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 1157

Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], 0]}, -Simp[(R*x*(d + e*x^2)^(q + 1))/(2*d*(q + 1)), x] + Dist[1/(2*d*(q + 1)), Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x],

$x], x]] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{LtQ}[q, -1]$

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^5}{(a + bx^2)^3} dx &= \int \left(\frac{d^3(10b^2c^2 - 15abcd + 6a^2d^2)}{b^5} + \frac{d^4(5bc - 3ad)x^2}{b^4} + \frac{d^5x^4}{b^3} + \frac{(bc - ad)^3(b^2c^2 + 3abcd + 6a^2d^2)}{b^5} \right) dx \\ &= \frac{d^3(10b^2c^2 - 15abcd + 6a^2d^2)x}{b^5} + \frac{d^4(5bc - 3ad)x^3}{3b^4} + \frac{d^5x^5}{5b^3} + \frac{\int \frac{(bc-ad)^3(b^2c^2+3abcd+6a^2d^2)+5bd(bc-ad)}{(a+bx^2)^3} dx}{b^5} \\ &= \frac{d^3(10b^2c^2 - 15abcd + 6a^2d^2)x}{b^5} + \frac{d^4(5bc - 3ad)x^3}{3b^4} + \frac{d^5x^5}{5b^3} + \frac{(bc - ad)^5x}{4ab^5(a + bx^2)^2} - \frac{\int \frac{-(bc-ad)^3(3b^2c^2+3abcd+6a^2d^2)}{(a+bx^2)^3} dx}{b^5} \\ &= \frac{d^3(10b^2c^2 - 15abcd + 6a^2d^2)x}{b^5} + \frac{d^4(5bc - 3ad)x^3}{3b^4} + \frac{d^5x^5}{5b^3} + \frac{(bc - ad)^5x}{4ab^5(a + bx^2)^2} + \frac{(bc - ad)^4(3b^2c^2 + 3abcd + 6a^2d^2)}{8a^2b^5(a + bx^2)} \\ &= \frac{d^3(10b^2c^2 - 15abcd + 6a^2d^2)x}{b^5} + \frac{d^4(5bc - 3ad)x^3}{3b^4} + \frac{d^5x^5}{5b^3} + \frac{(bc - ad)^5x}{4ab^5(a + bx^2)^2} + \frac{(bc - ad)^4(3b^2c^2 + 3abcd + 6a^2d^2)}{8a^2b^5(a + bx^2)} \end{aligned}$$

Mathematica [A] time = 0.13, size = 196, normalized size = 1.00

$$\frac{x(17ad + 3bc)(bc - ad)^4}{8a^2b^5(a + bx^2)} + \frac{d^3x(6a^2d^2 - 15abcd + 10b^2c^2)}{b^5} + \frac{(63a^2d^2 + 14abcd + 3b^2c^2)(bc - ad)^3 \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{11/2}} + \frac{4a^2d^5}{4a^2b^5}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^5/(a + b*x^2)^3,x]

[Out] $(d^3*(10*b^2*c^2 - 15*a*b*c*d + 6*a^2*d^2)*x)/b^5 + (d^4*(5*b*c - 3*a*d)*x^3)/(3*b^4) + (d^5*x^5)/(5*b^3) + ((b*c - a*d)^5*x)/(4*a*b^5*(a + b*x^2)^2) + ((b*c - a*d)^4*(3*b*c + 17*a*d)*x)/(8*a^2*b^5*(a + b*x^2)) + ((b*c - a*d)^3*(3*b^2*c^2 + 14*a*b*c*d + 63*a^2*d^2)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]])/(8*a^{5/2}*b^{11/2})$

fricas [B] time = 0.62, size = 1044, normalized size = 5.33

$$\frac{48 a^3 b^5 d^5 x^9 + 16 (25 a^3 b^5 c d^4 - 9 a^4 b^4 d^5) x^7 + 16 (150 a^3 b^5 c^2 d^3 - 175 a^4 b^4 c d^4 + 63 a^5 b^3 d^5) x^5 + 10 (9 a b^7 c^5 + 15 a^2 b^6 c^4 d - 150 a^3 b^5 c^3 d^2 + 750 a^4 b^4 c^2 d^3 - 875 a^5 b^3 c^2 d^4 + 315 a^6 b^2 c^2 d^5) x^3 + 15 (3 a^2 b^5 c^5 + 5 a^3 b^4 c^4 d + 30 a^4 b^3 c^3 d^2 - 150 a^5 b^2 c^2 d^3 + 175 a^6 b^2 c^2 d^4 - 63 a^7 d^5 + (3 b^7 c^5 + 5 a b^6 c^4 d + 30 a^2 b^5 c^3 d^2 - 150 a^3 b^4 c^2 d^3 + 175 a^4 b^3 c^2 d^4 - 63 a^5 b^2 c^2 d^5) x^4 + 2 (3 a b^6 c^5 + 5 a^2 b^5 c^4 d + 30 a^3 b^4 c^3 d^2 - 150 a^4 b^3 c^2 d^3 + 175 a^5 b^2 c^2 d^4 - 63 a^6 b^2 c^2 d^5) x^5}{8 a^2 b^5 (a + b x^2)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^5/(b*x^2+a)^3,x, algorithm="fricas")

[Out] $[1/240*(48*a^3*b^5*d^5*x^9 + 16*(25*a^3*b^5*c*d^4 - 9*a^4*b^4*d^5)*x^7 + 16*(150*a^3*b^5*c^2*d^3 - 175*a^4*b^4*c*d^4 + 63*a^5*b^3*d^5)*x^5 + 10*(9*a*b^7*c^5 + 15*a^2*b^6*c^4*d - 150*a^3*b^5*c^3*d^2 + 750*a^4*b^4*c^2*d^3 - 875*a^5*b^3*c^2*d^4 + 315*a^6*b^2*c^2*d^5)*x^3 + 15*(3*a^2*b^5*c^5 + 5*a^3*b^4*c^4*d + 30*a^4*b^3*c^3*d^2 - 150*a^5*b^2*c^2*d^3 + 175*a^6*b^2*c^2*d^4 - 63*a^7*d^5 + (3*b^7*c^5 + 5*a*b^6*c^4*d + 30*a^2*b^5*c^3*d^2 - 150*a^3*b^4*c^2*d^3 + 175*a^4*b^3*c^2*d^4 - 63*a^5*b^2*c^2*d^5)*x^4 + 2*(3*a*b^6*c^5 + 5*a^2*b^5*c^4*d + 30*a^3*b^4*c^3*d^2 - 150*a^4*b^3*c^2*d^3 + 175*a^5*b^2*c^2*d^4 - 63*a^6*b^2*c^2*d^5)*x^5]$

$$30a^3b^4c^3d^2 - 150a^4b^3c^2d^3 + 175a^5b^2cd^4 - 63a^6b^2d^5)x^2) \sqrt{-ab} \log\left(\frac{bx^2 + 2\sqrt{-ab}x - a}{bx^2 + a}\right) + 30(5a^2b^6c^5 - 5a^3b^5c^4d - 30a^4b^4c^3d^2 + 150a^5b^3c^2d^3 - 175a^6b^2cd^4 + 63a^7b^2d^5) \sqrt{ab} \arctan\left(\frac{bx}{\sqrt{ab}}\right) + \frac{1}{120}(24a^3b^5d^5x^9 + 8(25a^3b^5cd^4 - 9a^4b^4d^5)x^7 + 8(150a^3b^5c^2d^3 - 175a^4b^4cd^4 + 63a^5b^3d^5)x^5 + 5(9a^6b^7c^5 + 15a^2b^6c^4d - 150a^3b^5c^3d^2 + 750a^4b^4c^2d^3 - 875a^5b^3cd^4 + 315a^6b^2d^5)x^3 + 15(3a^2b^5c^5 + 5a^3b^4c^4d + 30a^4b^3c^3d^2 - 150a^5b^2c^2d^3 + 175a^6b^2cd^4 - 63a^7d^5 + (3b^7c^5 + 5ab^6c^4d + 30a^2b^5c^3d^2 - 150a^3b^4c^2d^3 + 175a^4b^3cd^4 - 63a^5b^2d^5)x^4 + 2(3a^2b^6c^5 + 5a^2b^5c^4d + 30a^3b^4c^3d^2 - 150a^4b^3c^2d^3 + 175a^5b^2cd^4 - 63a^6b^2d^5)x^2) \sqrt{ab} \arctan\left(\frac{bx}{\sqrt{ab}}\right) + 15(5a^2b^6c^5 - 5a^3b^5c^4d - 30a^4b^4c^3d^2 + 150a^5b^3c^2d^3 - 175a^6b^2cd^4 + 63a^7b^2d^5)x) / (a^3b^8x^4 + 2a^4b^7x^2 + a^5b^6)]$$

giac [A] time = 0.58, size = 340, normalized size = 1.73

$$\frac{(3b^5c^5 + 5ab^4c^4d + 30a^2b^3c^3d^2 - 150a^3b^2c^2d^3 + 175a^4bcd^4 - 63a^5d^5) \arctan\left(\frac{bx}{\sqrt{ab}}\right) + 3b^6c^5x^3 + 5ab^5c^4dx^3}{8\sqrt{ab}a^2b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^5/(b*x^2+a)^3,x, algorithm="giac")

[Out] $\frac{1}{8}(3b^5c^5 + 5a^2b^4c^4d + 30a^3b^3c^3d^2 - 150a^4b^2c^2d^3 + 175a^5bcd^4 - 63a^6d^5) \arctan\left(\frac{bx}{\sqrt{ab}}\right) / (\sqrt{ab}a^2b^5) + \frac{1}{8}(3b^6c^5x^3 + 5a^2b^5c^4dx^3 - 50a^3b^4c^3d^2x^3 + 90a^4b^3c^2d^3x^3 - 65a^5b^2cd^4x^3 + 17a^6b^2d^5x^3 + 5a^2b^5c^5x - 5a^3b^4c^4d - 30a^4b^3c^3d^2 + 70a^5b^2cd^4 - 55a^6b^2d^5) / ((bx^2 + a)^2a^2b^5) + \frac{1}{15}(3b^12d^5x^5 + 25b^12cd^4x^3 - 15a^2b^11d^5x^3 + 150b^12c^2d^3x - 225a^2b^11cd^4x + 90a^2b^10d^5x) / b^15$

maple [B] time = 0.02, size = 484, normalized size = 2.47

$$\frac{17a^3d^5x^3}{8(bx^2+a)^2b^4} - \frac{65a^2cd^4x^3}{8(bx^2+a)^2b^3} + \frac{45a^2d^3x^3}{4(bx^2+a)^2b^2} + \frac{5c^4dx^3}{8(bx^2+a)^2a} + \frac{3b^5x^3}{8(bx^2+a)^2a^2} - \frac{25c^3d^2x^3}{4(bx^2+a)^2b} + \frac{d^5x^5}{5b^3} + \frac{1}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^5/(b*x^2+a)^3,x)

[Out] $\frac{1}{5}d^5x^5/b^3 - d^5/b^4x^3/a + 5/3d^4/b^3x^3c + 6d^5/b^5a^2x - 15d^4/b^4acx + 10d^3/b^3c^2x + 17/8/b^4/(bx^2+a)^2a^3x^3d^5 - 65/8/b^3/(bx^2+a)^2a^2x^3cd^4 + 45/4/b^2/(bx^2+a)^2ax^3c^2d^3 - 25/4/b/(bx^2+a)^2x^3c^3d^2 + 5/8/(bx^2+a)^2/a^2x^3c^4d + 3/8b/(bx^2+a)^2/a^2x^3c^5 + 15/8/b^5/(bx^2+a)^2xa^4d^5 - 55/8/b^4/(bx^2+a)^2xa^3cd^4 + 35/4/b^3/(bx^2+a)^2xa^2c^2d^3 - 15/4/b^2/(bx^2+a)^2xa^2c^3d^2 - 5/8/b/(bx^2+a)^2xa^2c^4d + 5/8/(bx^2+a)^2x/ac^5 - 63/8/b^5a^3/(ab)^{1/2} \arctan(1/(ab)^{1/2}bx) * d^5 + 175/8/b^4a^2/(ab)^{1/2} \arctan(1/(ab)^{1/2}bx) * cd^4 - 75/4/b^3a/(ab)^{1/2} \arctan(1/(ab)^{1/2}bx) * c^2d^3 + 15/4/b^2/(ab)^{1/2} \arctan(1/(ab)^{1/2}bx) * c^3d^2 + 5/8/b/a/(ab)^{1/2} \arctan(1/(ab)^{1/2}bx) * c^4d + 3/8/a^2/(ab)^{1/2} \arctan(1/(ab)^{1/2}bx) * c^5$

maxima [A] time = 3.04, size = 334, normalized size = 1.70

$$\frac{(3b^6c^5 + 5ab^5c^4d - 50a^2b^4c^3d^2 + 90a^3b^3c^2d^3 - 65a^4b^2cd^4 + 17a^5bd^5)x^3 + 5(ab^5c^5 - a^2b^4c^4d - 6a^3b^3c^3d^2 + 15a^4b^2cd^4 - 5a^5bd^5)}{8(a^2b^7x^4 + 2a^3b^6x^2 + a^4b^5)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^5/(b*x^2+a)^3,x, algorithm="maxima")
```

```
[Out] 1/8*((3*b^6*c^5 + 5*a*b^5*c^4*d - 50*a^2*b^4*c^3*d^2 + 90*a^3*b^3*c^2*d^3 - 65*a^4*b^2*c*d^4 + 17*a^5*b*d^5)*x^3 + 5*(a*b^5*c^5 - a^2*b^4*c^4*d - 6*a^3*b^3*c^3*d^2 + 14*a^4*b^2*c^2*d^3 - 11*a^5*b*c*d^4 + 3*a^6*d^5)*x)/(a^2*b^7*x^4 + 2*a^3*b^6*x^2 + a^4*b^5) + 1/15*(3*b^2*d^5*x^5 + 5*(5*b^2*c*d^4 - 3*a*b*d^5)*x^3 + 15*(10*b^2*c^2*d^3 - 15*a*b*c*d^4 + 6*a^2*d^5)*x)/b^5 + 1/8*(3*b^5*c^5 + 5*a*b^4*c^4*d + 30*a^2*b^3*c^3*d^2 - 150*a^3*b^2*c^2*d^3 + 175*a^4*b*c*d^4 - 63*a^5*d^5)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^2*b^5)
```

```
mupad [B] time = 5.02, size = 409, normalized size = 2.09
```

$$\frac{5x(3a^5d^5 - 11a^4bcd^4 + 14a^3b^2c^2d^3 - 6a^2b^3c^3d^2 - ab^4c^4d + b^5c^5)}{8a} + \frac{x^3(17a^5bd^5 - 65a^4b^2cd^4 + 90a^3b^3c^2d^3 - 50a^2b^4c^3d^2 + 5ab^5c^4d + 3b^6c^5)}{8a^2}$$

$$a^2b^5 + 2ab^6x^2 + b^7x^4$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c + d*x^2)^5/(a + b*x^2)^3,x)
```

```
[Out] ((5*x*(3*a^5*d^5 + b^5*c^5 - 6*a^2*b^3*c^3*d^2 + 14*a^3*b^2*c^2*d^3 - a*b^4*c^4*d - 11*a^4*b*c*d^4))/(8*a) + (x^3*(3*b^6*c^5 + 17*a^5*b*d^5 - 65*a^4*b^2*c*d^4 - 50*a^2*b^4*c^3*d^2 + 90*a^3*b^3*c^2*d^3 + 5*a*b^5*c^4*d))/(8*a^2))/(a^2*b^5 + b^7*x^4 + 2*a*b^6*x^2) - x^3*((a*d^5)/b^4 - (5*c*d^4)/(3*b^3)) + x*((3*a*((3*a*d^5)/b^4 - (5*c*d^4)/b^3))/b - (3*a^2*d^5)/b^5 + (10*c^2*d^3)/b^3) + (d^5*x^5)/(5*b^3) + (atan((b^(1/2))*x*(a*d - b*c)^3*(63*a^2*d^2 + 3*b^2*c^2 + 14*a*b*c*d))/(a^(1/2)*(3*b^5*c^5 - 63*a^5*d^5 + 30*a^2*b^3*c^3*d^2 - 150*a^3*b^2*c^2*d^3 + 5*a*b^4*c^4*d + 175*a^4*b*c*d^4)))*(a*d - b*c)^3*(63*a^2*d^2 + 3*b^2*c^2 + 14*a*b*c*d))/(8*a^(5/2)*b^(11/2))
```

```
sympy [B] time = 4.41, size = 615, normalized size = 3.14
```

$$x^3 \left(-\frac{ad^5}{b^4} + \frac{5cd^4}{3b^3} \right) + x \left(\frac{6a^2d^5}{b^5} - \frac{15acd^4}{b^4} + \frac{10c^2d^3}{b^3} \right) + \frac{\sqrt{-\frac{1}{a^5b^{11}}} (ad - bc)^3 (63a^2d^2 + 14abcd + 3b^2c^2) \log \left(-\frac{a^3b}{63a^5d^5 - 175a^4b^2c^2d^3 + 30a^3b^3c^2d^2 - 150a^2b^4c^3d^2 + 5ab^5c^4d + 3b^6c^5} \right)}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x**2+c)**5/(b*x**2+a)**3,x)
```

```
[Out] x**3*(-a*d**5/b**4 + 5*c*d**4/(3*b**3)) + x*(6*a**2*d**5/b**5 - 15*a*c*d**4/b**4 + 10*c**2*d**3/b**3) + sqrt(-1/(a**5*b**11))*(a*d - b*c)**3*(63*a**2*d**2 + 14*a*b*c*d + 3*b**2*c**2)*log(-a**3*b**5*sqrt(-1/(a**5*b**11))*(a*d - b*c)**3*(63*a**2*d**2 + 14*a*b*c*d + 3*b**2*c**2)/(63*a**5*d**5 - 175*a**4*b*c*d**4 + 150*a**3*b**2*c**2*d**3 - 30*a**2*b**3*c**3*d**2 - 5*a*b**4*c**4*d - 3*b**5*c**5) + x)/16 - sqrt(-1/(a**5*b**11))*(a*d - b*c)**3*(63*a**2*d**2 + 14*a*b*c*d + 3*b**2*c**2)*log(a**3*b**5*sqrt(-1/(a**5*b**11))*(a*d - b*c)**3*(63*a**2*d**2 + 14*a*b*c*d + 3*b**2*c**2)/(63*a**5*d**5 - 175*a**4*b*c*d**4 + 150*a**3*b**2*c**2*d**3 - 30*a**2*b**3*c**3*d**2 - 5*a*b**4*c**4*d - 3*b**5*c**5) + x)/16 + (x**3*(17*a**5*b*d**5 - 65*a**4*b**2*c*d**4 + 90*a**3*b**3*c**2*d**3 - 50*a**2*b**4*c**3*d**2 + 5*a*b**5*c**4*d + 3*b**6*c**5) + x*(15*a**6*d**5 - 55*a**5*b*c*d**4 + 70*a**4*b**2*c**2*d**3 - 30*a**3*b**3*c**3*d**2 - 5*a**2*b**4*c**4*d + 5*a*b**5*c**5))/(8*a**4*b**5 + 16*a**3*b**6*x**2 + 8*a**2*b**7*x**4) + d**5*x**5/(5*b**3)
```


$$3.36 \quad \int \frac{(c+dx^2)^4}{(a+bx^2)^3} dx$$

Optimal. Leaf size=160

$$\frac{x(bc-ad)^3(13ad+3bc)}{8a^2b^4(a+bx^2)} + \frac{(bc-ad)^2(35a^2d^2+10abcd+3b^2c^2)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}b^{9/2}} + \frac{d^3x(4bc-3ad)}{b^4} + \frac{x(bc-ad)^4}{4ab^4(a+bx^2)}$$

[Out] d^3*(-3*a*d+4*b*c)*x/b^4+1/3*d^4*x^3/b^3+1/4*(-a*d+b*c)^4*x/a/b^4/(b*x^2+a)^2+1/8*(-a*d+b*c)^3*(13*a*d+3*b*c)*x/a^2/b^4/(b*x^2+a)+1/8*(-a*d+b*c)^2*(35*a^2*d^2+10*a*b*c*d+3*b^2*c^2)*arctan(x*b^(1/2)/a^(1/2))/a^(5/2)/b^(9/2)

Rubi [A] time = 0.20, antiderivative size = 160, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {390, 1157, 385, 205}

$$\frac{(bc-ad)^2(35a^2d^2+10abcd+3b^2c^2)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}b^{9/2}} + \frac{x(bc-ad)^3(13ad+3bc)}{8a^2b^4(a+bx^2)} + \frac{d^3x(4bc-3ad)}{b^4} + \frac{x(bc-ad)^4}{4ab^4(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^4/(a + b*x^2)^3,x]

[Out] (d^3*(4*b*c - 3*a*d)*x)/b^4 + (d^4*x^3)/(3*b^3) + ((b*c - a*d)^4*x)/(4*a*b^4*(a + b*x^2)^2) + ((b*c - a*d)^3*(3*b*c + 13*a*d)*x)/(8*a^2*b^4*(a + b*x^2)) + ((b*c - a*d)^2*(3*b^2*c^2 + 10*a*b*c*d + 35*a^2*d^2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(8*a^(5/2)*b^(9/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p+1))/(a*b*n*(p+1), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(a*b*n*(p+1), Int[(a + b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 1157

Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x, 0]}, -Simp[(R*x*(d + e*x^2)^(q+1))/(2*d*(q+1), x] + Dist[1/(2*d*(q+1)), Int[(d + e*x^2)^(q+1)*ExpandToSum[2*d*(q+1)*Qx + R*(2*q+3), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 -

$$b^4cd^3 - 7a^4b^3d^4)x^5 + (9ab^6c^4 + 12a^2b^5c^3d - 90a^3b^4c^2d^2 + 300a^4b^3cd^3 - 175a^5b^2d^4)x^3 + 3(3a^2b^4c^4 + 4a^3b^3c^3d + 18a^4b^2c^2d^2 - 60a^5b^3cd^3 + 35a^6d^4 + (3b^6c^4 + 4ab^5c^3d + 18a^2b^4c^2d^2 - 60a^3b^3cd^3 + 35a^4b^2d^4)x^4 + 2(3ab^5c^4 + 4a^2b^4c^3d + 18a^3b^3c^2d^2 - 60a^4b^2cd^3 + 35a^5bd^4)x^2) \sqrt{ab} \arctan(\sqrt{ab}x/a) + 3(5a^2b^5c^4 - 4a^3b^4c^3d - 18a^4b^3c^2d^2 + 60a^5b^2cd^3 - 35a^6bd^4)x / (a^3b^7x^4 + 2a^4b^6x^2 + a^5b^5)]$$

giac [A] time = 0.58, size = 254, normalized size = 1.59

$$\frac{(3b^4c^4 + 4ab^3c^3d + 18a^2b^2c^2d^2 - 60a^3bcd^3 + 35a^4d^4) \arctan\left(\frac{bx}{\sqrt{ab}}\right) + 3b^5c^4x^3 + 4ab^4c^3dx^3 - 30a^2b^3c^2d^2x^3}{8\sqrt{ab}a^2b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^3,x, algorithm="giac")

[Out] $\frac{1}{8}(3b^4c^4 + 4a^2b^3c^3d + 18a^2b^2c^2d^2 - 60a^3b^3cd^3 + 35a^4d^4) \arctan(bx/\sqrt{ab}) / (\sqrt{ab}a^2b^4) + \frac{1}{8}(3b^5c^4x^3 + 4ab^4c^3d^3x^3 - 30a^2b^3c^2d^2x^3 + 36a^3b^2cd^3x^3 - 13a^4bd^4x^3 + 5ab^4c^4x^3 - 4a^2b^3c^3d^3x^3 - 18a^3b^2c^2d^2x^3 + 28a^4b^2cd^3x^3 - 11a^5d^4x^3) / ((bx^2 + a)^2a^2b^4) + \frac{1}{3}(b^6d^4x^3 + 12b^6cd^3x^3 - 9ab^5d^4x^3) / b^9$

maple [B] time = 0.01, size = 367, normalized size = 2.29

$$-\frac{13a^2d^4x^3}{8(bx^2+a)^2b^3} + \frac{9acd^3x^3}{2(bx^2+a)^2b^2} + \frac{c^3dx^3}{2(bx^2+a)^2a} + \frac{3bc^4x^3}{8(bx^2+a)^2a^2} - \frac{15c^2d^2x^3}{4(bx^2+a)^2b} - \frac{11a^3d^4x}{8(bx^2+a)^2b^4} + \frac{7a^2}{2(bx^2+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^4/(b*x^2+a)^3,x)

[Out] $\frac{1}{3}d^4x^3/b^3 - 3d^4/b^4 * a*x + 4d^3/b^3 * x*c - 13/8/b^3 / (b*x^2+a)^2 * a^2 * x^3 * d^4 + 9/2/b^2 / (b*x^2+a)^2 * a * x^3 * c * d^3 - 15/4/b / (b*x^2+a)^2 * x^3 * c^2 * d^2 + 1/2 / (b*x^2+a)^2 / a * x^3 * c^3 * d + 3/8 * b / (b*x^2+a)^2 / a^2 * x^3 * c^4 - 11/8/b^4 / (b*x^2+a)^2 * x * a^3 * d^4 + 7/2/b^3 / (b*x^2+a)^2 * x * a^2 * c * d^3 - 9/4/b^2 / (b*x^2+a)^2 * x * a * c^2 * d^2 - 1/2/b / (b*x^2+a)^2 * x * c^3 * d + 5/8 / (b*x^2+a)^2 * x / a * c^4 + 35/8/b^4 * a^2 / (a*b)^(1/2) * arctan(1/(a*b)^(1/2) * b*x) * d^4 - 15/2/b^3 * a / (a*b)^(1/2) * arctan(1/(a*b)^(1/2) * b*x) * c * d^3 + 9/4/b^2 / (a*b)^(1/2) * arctan(1/(a*b)^(1/2) * b*x) * c^2 * d^2 + 1/2/b/a / (a*b)^(1/2) * arctan(1/(a*b)^(1/2) * b*x) * c^3 * d + 3/8/a^2 / (a*b)^(1/2) * arctan(1/(a*b)^(1/2) * b*x) * c^4$

maxima [A] time = 3.07, size = 253, normalized size = 1.58

$$\frac{(3b^5c^4 + 4ab^4c^3d - 30a^2b^3c^2d^2 + 36a^3b^2cd^3 - 13a^4bd^4)x^3 + (5ab^4c^4 - 4a^2b^3c^3d - 18a^3b^2c^2d^2 + 28a^4bcd^3)}{8(a^2b^6x^4 + 2a^3b^5x^2 + a^4b^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^3,x, algorithm="maxima")

[Out] $\frac{1}{8}((3b^5c^4 + 4a^2b^4c^3d - 30a^2b^3c^2d^2 + 36a^3b^2cd^3 - 13a^4bd^4)x^3 + (5ab^4c^4 - 4a^2b^3c^3d - 18a^3b^2c^2d^2 + 28a^4bcd^3 - 11a^5d^4)x) / (a^2b^6x^4 + 2a^3b^5x^2 + a^4b^4) + \frac{1}{3} * (b^6d^4x^3 + 3(4b^5cd^3 - 3a^2d^4)x) / b^4 + \frac{1}{8}(3b^4c^4 + 4a^2b^3c^3d + 18a^2b^2c^2d^2 - 60a^3b^3cd^3 + 35a^4d^4) \arctan(bx/\sqrt{ab}) / (\sqrt{ab}a^2b^4)$

mupad [B] time = 0.14, size = 318, normalized size = 1.99

$$\frac{d^4 x^3}{3b^3} - x \left(\frac{3ad^4}{b^4} - \frac{4cd^3}{b^3} \right) - \frac{x(11a^4d^4 - 28a^3bcd^3 + 18a^2b^2c^2d^2 + 4ab^3c^3d - 5b^4c^4)}{8a} - \frac{x^3(-13a^4bd^4 + 36a^3b^2cd^3 - 30a^2b^3c^2d^2 + 4ab^4c^3)}{8a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^4/(a + b*x^2)^3, x)

[Out] (d^4*x^3)/(3*b^3) - x*((3*a*d^4)/b^4 - (4*c*d^3)/b^3) - ((x*(11*a^4*d^4 - 5*b^4*c^4 + 18*a^2*b^2*c^2*d^2 + 4*a*b^3*c^3*d - 28*a^3*b*c*d^3))/(8*a) - (x^3*(3*b^5*c^4 - 13*a^4*b*d^4 + 36*a^3*b^2*c*d^3 - 30*a^2*b^3*c^2*d^2 + 4*a*b^4*c^3*d))/(8*a^2))/(a^2*b^4 + b^6*x^2 + 2*a*b^5*x^2) + (atan((b^(1/2))*x*(a*d - b*c)^2*(35*a^2*d^2 + 3*b^2*c^2 + 10*a*b*c*d))/(a^(1/2)*(35*a^4*d^4 + 3*b^4*c^4 + 18*a^2*b^2*c^2*d^2 + 4*a*b^3*c^3*d - 60*a^3*b*c*d^3)))*(a*d - b*c)^2*(35*a^2*d^2 + 3*b^2*c^2 + 10*a*b*c*d))/(8*a^(5/2)*b^(9/2))

sympy [B] time = 2.81, size = 515, normalized size = 3.22

$$x \left(-\frac{3ad^4}{b^4} + \frac{4cd^3}{b^3} \right) - \frac{\sqrt{-\frac{1}{a^5b^9}} (ad - bc)^2 (35a^2d^2 + 10abcd + 3b^2c^2) \log \left(-\frac{a^3b^4 \sqrt{-\frac{1}{a^5b^9}} (ad - bc)^2 (35a^2d^2 + 10abcd + 3b^2c^2)}{35a^4d^4 - 60a^3bcd^3 + 18a^2b^2c^2d^2 + 4ab^3c^3d + 3b^4c^4} + x \right)}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**4/(b*x**2+a)**3, x)

[Out] x*(-3*a*d**4/b**4 + 4*c*d**3/b**3) - sqrt(-1/(a**5*b**9))*(a*d - b*c)**2*(35*a**2*d**2 + 10*a*b*c*d + 3*b**2*c**2)*log(-a**3*b**4*sqrt(-1/(a**5*b**9))*(a*d - b*c)**2*(35*a**2*d**2 + 10*a*b*c*d + 3*b**2*c**2)/(35*a**4*d**4 - 60*a**3*b*c*d**3 + 18*a**2*b**2*c**2*d**2 + 4*a*b**3*c**3*d + 3*b**4*c**4) + x)/16 + sqrt(-1/(a**5*b**9))*(a*d - b*c)**2*(35*a**2*d**2 + 10*a*b*c*d + 3*b**2*c**2)*log(a**3*b**4*sqrt(-1/(a**5*b**9))*(a*d - b*c)**2*(35*a**2*d**2 + 10*a*b*c*d + 3*b**2*c**2)/(35*a**4*d**4 - 60*a**3*b*c*d**3 + 18*a**2*b**2*c**2*d**2 + 4*a*b**3*c**3*d + 3*b**4*c**4) + x)/16 + (x**3*(-13*a**4*b*d**4 + 36*a**3*b**2*c*d**3 - 30*a**2*b**3*c**2*d**2 + 4*a*b**4*c**3*d + 3*b**5*c**4) + x*(-11*a**5*d**4 + 28*a**4*b*c*d**3 - 18*a**3*b**2*c**2*d**2 - 4*a**2*b**3*c**3*d + 5*a*b**4*c**4))/(8*a**4*b**4 + 16*a**3*b**5*x**2 + 8*a**2*b**6*x**4) + d**4*x**3/(3*b**3)

$$3.37 \quad \int \frac{(c+dx^2)^3}{(a+bx^2)^3} dx$$

Optimal. Leaf size=130

$$\frac{3x(bc-ad)^2(3ad+bc)}{8a^2b^3(a+bx^2)} + \frac{3(bc-ad)(4a^2d^2+(ad+bc)^2)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}b^{7/2}} + \frac{x(bc-ad)^3}{4ab^3(a+bx^2)^2} + \frac{d^3x}{b^3}$$

[Out] $d^3*x/b^3+1/4*(-a*d+b*c)^3*x/a/b^3/(b*x^2+a)^2+3/8*(-a*d+b*c)^2*(3*a*d+b*c)*x/a^2/b^3/(b*x^2+a)+3/8*(-a*d+b*c)*(4*a^2*d^2+(a*d+b*c)^2)*\arctan(x*b^(1/2)/a^(1/2))/a^(5/2)/b^(7/2)$

Rubi [A] time = 0.17, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {390, 1157, 385, 205}

$$\frac{3(bc-ad)(4a^2d^2+(ad+bc)^2)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}b^{7/2}} + \frac{3x(bc-ad)^2(3ad+bc)}{8a^2b^3(a+bx^2)} + \frac{x(bc-ad)^3}{4ab^3(a+bx^2)^2} + \frac{d^3x}{b^3}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^3/(a + b*x^2)^3,x]

[Out] $(d^3*x)/b^3 + ((b*c - a*d)^3*x)/(4*a*b^3*(a + b*x^2)^2) + (3*(b*c - a*d)^2*(b*c + 3*a*d)*x)/(8*a^2*b^3*(a + b*x^2)) + (3*(b*c - a*d)*(4*a^2*d^2 + (b*c + a*d)^2)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]])/(8*a^(5/2)*b^(7/2))$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 390

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 1157

Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] :> With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x, 0]}, -Simp[(R*x*(d + e*x^2)^(q + 1))/(2*d*(q + 1)), x] + Dist[1/(2*d*(q + 1)), Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]

Rubi steps

$$\begin{aligned}
 \int \frac{(c + dx^2)^3}{(a + bx^2)^3} dx &= \int \left(\frac{d^3}{b^3} + \frac{b^3c^3 - a^3d^3 + 3bd(bc - ad)(bc + ad)x^2 + 3b^2d^2(bc - ad)x^4}{b^3(a + bx^2)^3} \right) dx \\
 &= \frac{d^3x}{b^3} + \frac{\int \frac{b^3c^3 - a^3d^3 + 3bd(bc - ad)(bc + ad)x^2 + 3b^2d^2(bc - ad)x^4}{(a + bx^2)^3} dx}{b^3} \\
 &= \frac{d^3x}{b^3} + \frac{(bc - ad)^3x}{4ab^3(a + bx^2)^2} - \frac{\int \frac{-3(bc - ad)(bc + ad)^2 - 12abd^2(bc - ad)x^2}{(a + bx^2)^2} dx}{4ab^3} \\
 &= \frac{d^3x}{b^3} + \frac{(bc - ad)^3x}{4ab^3(a + bx^2)^2} + \frac{3(bc - ad)^2(bc + 3ad)x}{8a^2b^3(a + bx^2)} + \frac{(3(bc - ad)(4a^2d^2 + (bc + ad)^2)) \int \frac{1}{a + bx^2}}{8a^2b^3} \\
 &= \frac{d^3x}{b^3} + \frac{(bc - ad)^3x}{4ab^3(a + bx^2)^2} + \frac{3(bc - ad)^2(bc + 3ad)x}{8a^2b^3(a + bx^2)} + \frac{3(bc - ad)(4a^2d^2 + (bc + ad)^2) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{7/2}}
 \end{aligned}$$

Mathematica [A] time = 0.08, size = 139, normalized size = 1.07

$$\frac{3x(bc - ad)^2(3ad + bc)}{8a^2b^3(a + bx^2)} + \frac{3(-5a^3d^3 + 3a^2bcd^2 + ab^2c^2d + b^3c^3) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{7/2}} + \frac{x(bc - ad)^3}{4ab^3(a + bx^2)^2} + \frac{d^3x}{b^3}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^3/(a + b*x^2)^3,x]

[Out] (d^3*x)/b^3 + ((b*c - a*d)^3*x)/(4*a*b^3*(a + b*x^2)^2) + (3*(b*c - a*d)^2*(b*c + 3*a*d)*x)/(8*a^2*b^3*(a + b*x^2)) + (3*(b^3*c^3 + a*b^2*c^2*d + 3*a^2*b*c*d^2 - 5*a^3*d^3)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(8*a^(5/2)*b^(7/2))

fricas [B] time = 0.62, size = 606, normalized size = 4.66

$$\left[\frac{16a^3b^3d^3x^5 + 2(3ab^5c^3 + 3a^2b^4c^2d - 15a^3b^3cd^2 + 25a^4b^2d^3)x^3 + 3(a^2b^3c^3 + a^3b^2c^2d + 3a^4bcd^2 - 5a^5d^3 + (b^3c^3 - a^3d^3) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right))}{8a^{5/2}b^{7/2}} + \frac{x(bc - ad)^3}{4ab^3(a + bx^2)^2} + \frac{d^3x}{b^3} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^3,x, algorithm="fricas")

[Out] [1/16*(16*a^3*b^3*d^3*x^5 + 2*(3*a*b^5*c^3 + 3*a^2*b^4*c^2*d - 15*a^3*b^3*c*d^2 + 25*a^4*b^2*d^3)*x^3 + 3*(a^2*b^3*c^3 + a^3*b^2*c^2*d + 3*a^4*b*c*d^2 - 5*a^5*d^3 + (b^3*c^3 + a*b^2*c^2*d + 3*a^2*b^3*c*d^2 - 5*a^3*b^2*d^3)*x^4 + 2*(a*b^4*c^3 + a^2*b^3*c^2*d + 3*a^3*b^2*c*d^2 - 5*a^4*b*d^3)*x^2)*sqrt(-a*b)*log((b*x^2 + 2*sqrt(-a*b)*x - a)/(b*x^2 + a)) + 2*(5*a^2*b^4*c^3 - 3*a^3*b^3*c^2*d - 9*a^4*b^2*c*d^2 + 15*a^5*b*d^3)*x)/(a^3*b^6*x^4 + 2*a^4*b^5*x^2 + a^5*b^4), 1/8*(8*a^3*b^3*d^3*x^5 + (3*a*b^5*c^3 + 3*a^2*b^4*c^2*d - 15*a^3*b^3*c*d^2 + 25*a^4*b^2*d^3)*x^3 + 3*(a^2*b^3*c^3 + a^3*b^2*c^2*d + 3*a^4*b*c*d^2 - 5*a^5*d^3 + (b^3*c^3 + a*b^2*c^2*d + 3*a^2*b^3*c*d^2 - 5*a^3*b^2*d^3)*x^4 + 2*(a*b^4*c^3 + a^2*b^3*c^2*d + 3*a^3*b^2*c*d^2 - 5*a^4*b*d^3)*x^2)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + (5*a^2*b^4*c^3 - 3*a^3*b^3*c^2*d - 9*a^4*b^2*c*d^2 + 15*a^5*b*d^3)*x)/(a^3*b^6*x^4 + 2*a^4*b^5*x^2 + a^5*b^4)]

giac [A] time = 0.57, size = 178, normalized size = 1.37

$$\frac{d^3 x}{b^3} + \frac{3(b^3 c^3 + ab^2 c^2 d + 3a^2 b c d^2 - 5a^3 d^3) \arctan\left(\frac{bx}{\sqrt{ab}}\right) + 3b^4 c^3 x^3 + 3ab^3 c^2 d x^3 - 15a^2 b^2 c d^2 x^3 + 9a^3 b d^3 x^3}{8\sqrt{ab} a^2 b^3} + \frac{3b^4 c^3 x^3 + 3ab^3 c^2 d x^3 - 15a^2 b^2 c d^2 x^3 + 9a^3 b d^3 x^3}{8(bx^2 + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^3,x, algorithm="giac")

[Out] $d^3 x/b^3 + 3/8*(b^3 c^3 + a*b^2*c^2*d + 3*a^2*b*c*d^2 - 5*a^3*d^3)*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b}*a^2*b^3) + 1/8*(3*b^4*c^3*x^3 + 3*a*b^3*c^2*d*x^3 - 15*a^2*b^2*c*d^2*x^3 + 9*a^3*b*d^3*x^3 + 5*a*b^3*c^3*x - 3*a^2*b^2*c^2*d*x - 9*a^3*b*c*d^2*x + 7*a^4*d^3*x)/((b*x^2 + a)^2*a^2*b^3)$

maple [B] time = 0.01, size = 266, normalized size = 2.05

$$\frac{9a d^3 x^3}{8(bx^2 + a)^2 b^2} + \frac{3c^2 d x^3}{8(bx^2 + a)^2 a} + \frac{3b c^3 x^3}{8(bx^2 + a)^2 a^2} - \frac{15c d^2 x^3}{8(bx^2 + a)^2 b} + \frac{7a^2 d^3 x}{8(bx^2 + a)^2 b^3} - \frac{9ac d^2 x}{8(bx^2 + a)^2 b^2} + \frac{5c^3}{8(bx^2 + a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^3/(b*x^2+a)^3,x)

[Out] $1/b^3*d^3*x+9/8/b^2/(b*x^2+a)^2*a*x^3*d^3-15/8/b/(b*x^2+a)^2*x^3*c*d^2+3/8/(b*x^2+a)^2/a*x^3*c^2*d+3/8*b/(b*x^2+a)^2/a^2*x^3*c^3+7/8/b^3/(b*x^2+a)^2*x*a^2*d^3-9/8/b^2/(b*x^2+a)^2*x*a*c*d^2-3/8/b/(b*x^2+a)^2*x*c^2*d+5/8/(b*x^2+a)^2*x/a*c^3-15/8/b^3*a/(a*b)^(1/2)*\arctan(1/(a*b)^(1/2)*b*x)*d^3+9/8/b^2/(a*b)^(1/2)*\arctan(1/(a*b)^(1/2)*b*x)*c*d^2+3/8/b/a/(a*b)^(1/2)*\arctan(1/(a*b)^(1/2)*b*x)*c^2+d+3/8/a^2/(a*b)^(1/2)*\arctan(1/(a*b)^(1/2)*b*x)*c^3$

maxima [A] time = 2.94, size = 185, normalized size = 1.42

$$\frac{d^3 x}{b^3} + \frac{3(b^4 c^3 + ab^3 c^2 d - 5a^2 b^2 c d^2 + 3a^3 b d^3) x^3 + (5ab^3 c^3 - 3a^2 b^2 c^2 d - 9a^3 b c d^2 + 7a^4 d^3) x}{8(a^2 b^5 x^4 + 2a^3 b^4 x^2 + a^4 b^3)} + \frac{3(b^3 c^3 + ab^2 c^2 d)}{8(bx^2 + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^3,x, algorithm="maxima")

[Out] $d^3 x/b^3 + 1/8*(3*(b^4*c^3 + a*b^3*c^2*d - 5*a^2*b^2*c*d^2 + 3*a^3*b*d^3)*x^3 + (5*a*b^3*c^3 - 3*a^2*b^2*c^2*d - 9*a^3*b*c*d^2 + 7*a^4*d^3)*x)/(a^2*b^5*x^4 + 2*a^3*b^4*x^2 + a^4*b^3) + 3/8*(b^3*c^3 + a*b^2*c^2*d + 3*a^2*b*c*d^2 - 5*a^3*d^3)*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b}*a^2*b^3)$

mupad [B] time = 5.05, size = 240, normalized size = 1.85

$$\frac{x(7a^3 d^3 - 9a^2 b c d^2 - 3a b^2 c^2 d + 5b^3 c^3)}{8a} + \frac{3x^3(3a^3 b d^3 - 5a^2 b^2 c d^2 + a b^3 c^2 d + b^4 c^3)}{8a^2} + \frac{d^3 x}{b^3} + \frac{3 \operatorname{atan}\left(\frac{\sqrt{b} x(a-d-bc)(5a^2 d^2 + 2abcd + b^2 c^2)}{\sqrt{a}(-5a^3 d^3 + 3a^2 b c d^2 + a b^2 c^2 d + b^3 c^3)}\right)}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^3/(a + b*x^2)^3,x)

[Out] $((x*(7*a^3*d^3 + 5*b^3*c^3 - 3*a*b^2*c^2*d - 9*a^2*b*c*d^2))/(8*a) + (3*x^3*(b^4*c^3 + 3*a^3*b*d^3 - 5*a^2*b^2*c*d^2 + a*b^3*c^2*d))/(8*a^2))/(a^2*b^3 + b^5*x^4) + (d^3*x)/b^3 + (3*atan((b^(1/2))*x*(a*d - b*c)*(5*a^2*d^2 + b^2*c^2 + 2*a*b*c*d))/(a^(1/2)*(b^3*c^3 - 5*a^3*d^3 + a*b^2*c^2*d)))/8$

$d + 3a^2b^2cd^2)))(ad - bc)(5a^2d^2 + b^2c^2 + 2abc^2d)/(8a^5/2)b^{7/2})$

sympy [B] time = 1.89, size = 422, normalized size = 3.25

$$\frac{3\sqrt{-\frac{1}{a^5b^7}}(ad - bc)(5a^2d^2 + 2abcd + b^2c^2) \log\left(-\frac{3a^3b^3\sqrt{-\frac{1}{a^5b^7}}(ad - bc)(5a^2d^2 + 2abcd + b^2c^2)}{15a^3d^3 - 9a^2bcd^2 - 3ab^2c^2d - 3b^3c^3} + x\right)}{16} - 3\sqrt{-\frac{1}{a^5b^7}}(ad - bc)(5a^2d^2 + 2abcd + b^2c^2)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**3/(b*x**2+a)**3,x)

[Out] $3\sqrt{-1/(a^5b^7)}(ad - bc)(5a^2d^2 + 2abc^2d + b^2c^2) \log(-3a^3b^3\sqrt{-1/(a^5b^7)}(ad - bc)(5a^2d^2 + 2abc^2d + b^2c^2)/(15a^3d^3 - 9a^2bcd^2 - 3ab^2c^2d - 3b^3c^3) + x)/16 - 3\sqrt{-1/(a^5b^7)}(ad - bc)(5a^2d^2 + 2abc^2d + b^2c^2) \log(3a^3b^3\sqrt{-1/(a^5b^7)}(ad - bc)(5a^2d^2 + 2abc^2d + b^2c^2)/(15a^3d^3 - 9a^2bcd^2 - 3ab^2c^2d - 3b^3c^3) + x)/16 + (x^3(9a^3bd^3 - 15a^2b^2cd^2 + 3ab^3c^2d + 3b^4c^3) + x(7a^4d^3 - 9a^3b^2cd^2 - 3a^2b^2c^2d + 5ab^3c^3))/(8a^4b^3 + 16a^3b^4x^2 + 8a^2b^5x^4) + d^3x/b^3$

$$3.38 \quad \int \frac{(c+dx^2)^2}{(a+bx^2)^3} dx$$

Optimal. Leaf size=116

$$\frac{3x\left(\frac{c^2}{a^2} - \frac{d^2}{b^2}\right)}{8(a+bx^2)} + \frac{(3a^2d^2 + 2abcd + 3b^2c^2) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{5/2}} + \frac{x(c+dx^2)(bc-ad)}{4ab(a+bx^2)^2}$$

[Out] 3/8*(c^2/a^2-d^2/b^2)*x/(b*x^2+a)+1/4*(-a*d+b*c)*x*(d*x^2+c)/a/b/(b*x^2+a)^2+1/8*(3*a^2*d^2+2*a*b*c*d+3*b^2*c^2)*arctan(x*b^(1/2)/a^(1/2))/a^(5/2)/b^(5/2)

Rubi [A] time = 0.08, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, number of rules / integrand size = 0.158, Rules used = {413, 385, 205}

$$\frac{3x\left(\frac{c^2}{a^2} - \frac{d^2}{b^2}\right)}{8(a+bx^2)} + \frac{(3a^2d^2 + 2abcd + 3b^2c^2) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{5/2}} + \frac{x(c+dx^2)(bc-ad)}{4ab(a+bx^2)^2}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^2/(a + b*x^2)^3,x]

[Out] (3*(c^2/a^2 - d^2/b^2)*x)/(8*(a + b*x^2)) + ((b*c - a*d)*x*(c + d*x^2))/(4*a*b*(a + b*x^2)^2) + ((3*b^2*c^2 + 2*a*b*c*d + 3*a^2*d^2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(8*a^(5/2)*b^(5/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^2}{(a + bx^2)^3} dx &= \frac{(bc - ad)x(c + dx^2)}{4ab(a + bx^2)^2} + \frac{\int \frac{c(3bc+ad)+d(bc+3ad)x^2}{(a+bx^2)^2} dx}{4ab} \\ &= \frac{3\left(\frac{c^2}{a^2} - \frac{d^2}{b^2}\right)x}{8(a + bx^2)} + \frac{(bc - ad)x(c + dx^2)}{4ab(a + bx^2)^2} + \frac{(3b^2c^2 + 2abcd + 3a^2d^2) \int \frac{1}{a+bx^2} dx}{8a^2b^2} \\ &= \frac{3\left(\frac{c^2}{a^2} - \frac{d^2}{b^2}\right)x}{8(a + bx^2)} + \frac{(bc - ad)x(c + dx^2)}{4ab(a + bx^2)^2} + \frac{(3b^2c^2 + 2abcd + 3a^2d^2) \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}b^{5/2}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 124, normalized size = 1.07

$$\frac{(3a^2d^2 + 2abcd + 3b^2c^2) \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}b^{5/2}} + \frac{x(-3a^3d^2 - a^2bd(2c + 5dx^2) + ab^2c(5c + 2dx^2) + 3b^3c^2x^2)}{8a^2b^2(a + bx^2)^2}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^2/(a + b*x^2)^3,x]

[Out] (x*(-3*a^3*d^2 + 3*b^3*c^2*x^2 + a*b^2*c*(5*c + 2*d*x^2) - a^2*b*d*(2*c + 5*d*x^2)))/(8*a^2*b^2*(a + b*x^2)^2) + ((3*b^2*c^2 + 2*a*b*c*d + 3*a^2*d^2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(8*a^(5/2)*b^(5/2))

fricas [B] time = 0.53, size = 449, normalized size = 3.87

$$\left[\frac{2(3ab^4c^2 + 2a^2b^3cd - 5a^3b^2d^2)x^3 - (3a^2b^2c^2 + 2a^3bcd + 3a^4d^2 + (3b^4c^2 + 2ab^3cd + 3a^2b^2d^2)x^4 + 2(3ab^3c^2 - 2a^2b^2cd + 3a^3bd^2)x^5)}{16(a^3b^5x^4 + 2a^4b^4x^2 + a^5b^3)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^3,x, algorithm="fricas")

[Out] [1/16*(2*(3*a*b^4*c^2 + 2*a^2*b^3*c*d - 5*a^3*b^2*d^2)*x^3 - (3*a^2*b^2*c^2 + 2*a^3*b*c*d + 3*a^4*d^2 + (3*b^4*c^2 + 2*a*b^3*c*d + 3*a^2*b^2*d^2)*x^4 + 2*(3*a*b^3*c^2 + 2*a^2*b^2*c*d + 3*a^3*b*d^2)*x^5)*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b)*x - a)/(b*x^2 + a)) + 2*(5*a^2*b^3*c^2 - 2*a^3*b^2*c*d - 3*a^4*b*d^2)*x)/(a^3*b^5*x^4 + 2*a^4*b^4*x^2 + a^5*b^3), 1/8*((3*a*b^4*c^2 + 2*a^2*b^3*c*d - 5*a^3*b^2*d^2)*x^3 + (3*a^2*b^2*c^2 + 2*a^3*b*c*d + 3*a^4*d^2 + (3*b^4*c^2 + 2*a*b^3*c*d + 3*a^2*b^2*d^2)*x^4 + 2*(3*a*b^3*c^2 + 2*a^2*b^2*c*d + 3*a^3*b*d^2)*x^5)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + (5*a^2*b^3*c^2 - 2*a^3*b^2*c*d - 3*a^4*b*d^2)*x)/(a^3*b^5*x^4 + 2*a^4*b^4*x^2 + a^5*b^3)]

giac [A] time = 0.57, size = 126, normalized size = 1.09

$$\frac{(3b^2c^2 + 2abcd + 3a^2d^2) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{ab}a^2b^2} + \frac{3b^3c^2x^3 + 2ab^2cdx^3 - 5a^2bd^2x^3 + 5ab^2c^2x - 2a^2bcdx - 3a^3d^2x}{8(bx^2 + a)^2a^2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^3,x, algorithm="giac")

[Out] $1/8*(3*b^2*c^2 + 2*a*b*c*d + 3*a^2*d^2)*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b})*a^2*b^2) + 1/8*(3*b^3*c^2*x^3 + 2*a*b^2*c*d*x^3 - 5*a^2*b*d^2*x^3 + 5*a*b^2*c^2*x^2 - 2*a^2*b*c*d*x - 3*a^3*d^2*x)/(b*x^2 + a)^2*a^2*b^2)$

maple [A] time = 0.01, size = 147, normalized size = 1.27

$$\frac{cd \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{4\sqrt{ab} ab} + \frac{3c^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{ab} a^2} + \frac{3d^2 \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{ab} b^2} + \frac{\frac{(5a^2d^2-2abcd-3b^2c^2)x^3}{8a^2b} - \frac{(3a^2d^2+2abcd-5b^2c^2)x}{8ab^2}}{(bx^2+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+c)^2/(b*x^2+a)^3,x)`

[Out] $(-1/8*(5*a^2*d^2-2*a*b*c*d-3*b^2*c^2)/a^2/b*x^3-1/8*(3*a^2*d^2+2*a*b*c*d-5*b^2*c^2)/a/b^2*x)/(b*x^2+a)^2+3/8/b^2/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*d^2+1/4/a/b/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c*d+3/8/a^2/(a*b)^{(1/2)}*\arctan(1/(a*b)^{(1/2)}*b*x)*c^2)$

maxima [A] time = 3.01, size = 138, normalized size = 1.19

$$\frac{(3b^3c^2 + 2ab^2cd - 5a^2bd^2)x^3 + (5ab^2c^2 - 2a^2bcd - 3a^3d^2)x}{8(a^2b^4x^4 + 2a^3b^3x^2 + a^4b^2)} + \frac{(3b^2c^2 + 2abcd + 3a^2d^2) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{ab}a^2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^2/(b*x^2+a)^3,x, algorithm="maxima")`

[Out] $1/8*((3*b^3*c^2 + 2*a*b^2*c*d - 5*a^2*b*d^2)*x^3 + (5*a*b^2*c^2 - 2*a^2*b*c*d - 3*a^3*d^2)*x)/(a^2*b^4*x^4 + 2*a^3*b^3*x^2 + a^4*b^2) + 1/8*(3*b^2*c^2 + 2*a*b*c*d + 3*a^2*d^2)*\arctan(b*x/\sqrt{a*b})/(\sqrt{a*b})*a^2*b^2)$

mupad [B] time = 5.02, size = 130, normalized size = 1.12

$$\frac{\operatorname{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) (3a^2d^2 + 2abcd + 3b^2c^2)}{8a^{5/2}b^{5/2}} - \frac{\frac{x(3a^2d^2+2abcd-5b^2c^2)}{8ab^2} - \frac{x^3(-5a^2d^2+2abcd+3b^2c^2)}{8a^2b}}{a^2 + 2abx^2 + b^2x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^2/(a + b*x^2)^3,x)`

[Out] $(\operatorname{atan}((b^{(1/2)}*x)/a^{(1/2)})*(3*a^2*d^2 + 3*b^2*c^2 + 2*a*b*c*d))/(8*a^{(5/2)}*b^{(5/2)}) - ((x*(3*a^2*d^2 - 5*b^2*c^2 + 2*a*b*c*d))/(8*a*b^2) - (x^3*(3*b^2*c^2 - 5*a^2*d^2 + 2*a*b*c*d))/(8*a^2*b))/(a^2 + b^2*x^4 + 2*a*b*x^2)$

sympy [B] time = 1.05, size = 223, normalized size = 1.92

$$\frac{\sqrt{-\frac{1}{a^5b^5}} (3a^2d^2 + 2abcd + 3b^2c^2) \log\left(-a^3b^2\sqrt{-\frac{1}{a^5b^5}} + x\right)}{16} + \frac{\sqrt{-\frac{1}{a^5b^5}} (3a^2d^2 + 2abcd + 3b^2c^2) \log\left(a^3b^2\sqrt{-\frac{1}{a^5b^5}}\right)}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**2/(b*x**2+a)**3,x)`

[Out] $-\sqrt{-1/(a**5*b**5)}*(3*a**2*d**2 + 2*a*b*c*d + 3*b**2*c**2)*\log(-a**3*b**2*\sqrt{-1/(a**5*b**5)} + x)/16 + \sqrt{-1/(a**5*b**5)}*(3*a**2*d**2 + 2*a*b*c*d + 3*b**2*c**2)*\log(a**3*b**2*\sqrt{-1/(a**5*b**5)} + x)/16 + (x**3*(-5*a**2*b*d**2 + 2*a*b**2*c*d + 3*b**3*c**2) + x*(-3*a**3*d**2 - 2*a**2*b*c*d + 5*a*b**2*c**2))/(8*a**4*b**2 + 16*a**3*b**3*x**2 + 8*a**2*b**4*x**4)$

$$3.39 \quad \int \frac{c+dx^2}{(a+bx^2)^3} dx$$

Optimal. Leaf size=92

$$\frac{(ad + 3bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{3/2}} + \frac{x(ad + 3bc)}{8a^2b(a + bx^2)} + \frac{x(bc - ad)}{4ab(a + bx^2)^2}$$

[Out] $1/4*(-a*d+b*c)*x/a/b/(b*x^2+a)^2+1/8*(a*d+3*b*c)*x/a^2/b/(b*x^2+a)+1/8*(a*d+3*b*c)*\arctan(x*b^{(1/2)}/a^{(1/2)})/a^{(5/2)}/b^{(3/2)}$

Rubi [A] time = 0.03, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {385, 199, 205}

$$\frac{(ad + 3bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{3/2}} + \frac{x(ad + 3bc)}{8a^2b(a + bx^2)} + \frac{x(bc - ad)}{4ab(a + bx^2)^2}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)/(a + b*x^2)^3, x]

[Out] $((b*c - a*d)*x)/(4*a*b*(a + b*x^2)^2) + ((3*b*c + a*d)*x)/(8*a^2*b*(a + b*x^2)) + ((3*b*c + a*d)*\text{ArcTan}[\text{Sqrt}[b]*x]/\text{Sqrt}[a])/(8*a^{(5/2)}*b^{(3/2)})$

Rule 199

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rubi steps

$$\begin{aligned} \int \frac{c + dx^2}{(a + bx^2)^3} dx &= \frac{(bc - ad)x}{4ab(a + bx^2)^2} + \frac{(3bc + ad) \int \frac{1}{(a+bx^2)^2} dx}{4ab} \\ &= \frac{(bc - ad)x}{4ab(a + bx^2)^2} + \frac{(3bc + ad)x}{8a^2b(a + bx^2)} + \frac{(3bc + ad) \int \frac{1}{a+bx^2} dx}{8a^2b} \\ &= \frac{(bc - ad)x}{4ab(a + bx^2)^2} + \frac{(3bc + ad)x}{8a^2b(a + bx^2)} + \frac{(3bc + ad) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 84, normalized size = 0.91

$$\frac{(ad + 3bc) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}b^{3/2}} + \frac{x(a^2(-d) + ab(5c + dx^2) + 3b^2cx^2)}{8a^2b(a + bx^2)^2}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)/(a + b*x^2)^3,x]

[Out] (x*(-(a^2*d) + 3*b^2*c*x^2 + a*b*(5*c + d*x^2)))/(8*a^2*b*(a + b*x^2)^2) + ((3*b*c + a*d)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(8*a^(5/2)*b^(3/2))

fricas [A] time = 0.70, size = 301, normalized size = 3.27

$$\left[\frac{2(3ab^3c + a^2b^2d)x^3 - ((3b^3c + ab^2d)x^4 + 3a^2bc + a^3d + 2(3ab^2c + a^2bd)x^2)\sqrt{-ab} \log\left(\frac{bx^2 - 2\sqrt{-ab}x - a}{bx^2 + a}\right) + 2}{16(a^3b^4x^4 + 2a^4b^3x^2 + a^5b^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^3,x, algorithm="fricas")

[Out] [1/16*(2*(3*a*b^3*c + a^2*b^2*d)*x^3 - ((3*b^3*c + a*b^2*d)*x^4 + 3*a^2*b*c + a^3*d + 2*(3*a*b^2*c + a^2*b*d)*x^2)*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b)*x - a)/(b*x^2 + a)) + 2*(5*a^2*b^2*c - a^3*b*d)*x)/(a^3*b^4*x^4 + 2*a^4*b^3*x^2 + a^5*b^2), 1/8*((3*a*b^3*c + a^2*b^2*d)*x^3 + ((3*b^3*c + a*b^2*d)*x^4 + 3*a^2*b*c + a^3*d + 2*(3*a*b^2*c + a^2*b*d)*x^2)*sqrt(a*b)*arctan(sqrt(a*b)*x/a) + (5*a^2*b^2*c - a^3*b*d)*x)/(a^3*b^4*x^4 + 2*a^4*b^3*x^2 + a^5*b^2)]

giac [A] time = 0.58, size = 78, normalized size = 0.85

$$\frac{(3bc + ad) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{ab}a^2b} + \frac{3b^2cx^3 + abdx^3 + 5abcx - a^2dx}{8(bx^2 + a)^2a^2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^3,x, algorithm="giac")

[Out] 1/8*(3*b*c + a*d)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^2*b) + 1/8*(3*b^2*c*x^3 + a*b*d*x^3 + 5*a*b*c*x - a^2*d*x)/((b*x^2 + a)^2*a^2*b)

maple [A] time = 0.01, size = 89, normalized size = 0.97

$$\frac{d \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{ab} ab} + \frac{3c \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{ab} a^2} + \frac{\frac{(ad+3bc)x^3}{8a^2} - \frac{(ad-5bc)x}{8ab}}{(bx^2 + a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)/(b*x^2+a)^3,x)

[Out] (1/8*(a*d+3*b*c)/a^2*x^3-1/8*(a*d-5*b*c)/a/b*x)/(b*x^2+a)^2+1/8/a/b/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*d+3/8/a^2/(a*b)^(1/2)*arctan(1/(a*b)^(1/2)*b*x)*c

maxima [A] time = 3.00, size = 92, normalized size = 1.00

$$\frac{(3b^2c + abd)x^3 + (5abc - a^2d)x}{8(a^2b^3x^4 + 2a^3b^2x^2 + a^4b)} + \frac{(3bc + ad) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{ab} a^2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^3,x, algorithm="maxima")

[Out] 1/8*((3*b^2*c + a*b*d)*x^3 + (5*a*b*c - a^2*d)*x)/(a^2*b^3*x^4 + 2*a^3*b^2*x^2 + a^4*b) + 1/8*(3*b*c + a*d)*arctan(b*x/sqrt(a*b))/(sqrt(a*b)*a^2*b)

mupad [B] time = 5.02, size = 81, normalized size = 0.88

$$\frac{\frac{x^3(ad+3bc)}{8a^2} - \frac{x(ad-5bc)}{8ab}}{a^2 + 2abx^2 + b^2x^4} + \frac{\operatorname{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)(ad+3bc)}{8a^{5/2}b^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)/(a + b*x^2)^3,x)

[Out] ((x^3*(a*d + 3*b*c))/(8*a^2) - (x*(a*d - 5*b*c))/(8*a*b))/(a^2 + b^2*x^4 + 2*a*b*x^2) + (atan((b^(1/2)*x)/a^(1/2))*(a*d + 3*b*c))/(8*a^(5/2)*b^(3/2))

sympy [A] time = 0.58, size = 150, normalized size = 1.63

$$-\frac{\sqrt{-\frac{1}{a^5b^3}}(ad+3bc)\log\left(-a^3b\sqrt{-\frac{1}{a^5b^3}}+x\right)}{16} + \frac{\sqrt{-\frac{1}{a^5b^3}}(ad+3bc)\log\left(a^3b\sqrt{-\frac{1}{a^5b^3}}+x\right)}{16} + \frac{x^3(abd+3b^2c)+x(-a^2+3b^2x^2+8a^2bx^2+8a^2b^3x^4)}{8a^4b+16a^3b^2x^2+8a^2b^3x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)/(b*x**2+a)**3,x)

[Out] -sqrt(-1/(a**5*b**3))*(a*d + 3*b*c)*log(-a**3*b*sqrt(-1/(a**5*b**3)) + x)/16 + sqrt(-1/(a**5*b**3))*(a*d + 3*b*c)*log(a**3*b*sqrt(-1/(a**5*b**3)) + x)/16 + (x**3*(a*b*d + 3*b**2*c) + x*(-a**2*d + 5*a*b*c))/(8*a**4*b + 16*a**3*b**2*x**2 + 8*a**2*b**3*x**4)

$$3.40 \quad \int \frac{1}{(a+bx^2)^3(c+dx^2)} dx$$

Optimal. Leaf size=161

$$\frac{bx(3bc-7ad)}{8a^2(a+bx^2)(bc-ad)^2} + \frac{\sqrt{b}(15a^2d^2-10abcd+3b^2c^2)\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}(bc-ad)^3} - \frac{d^{5/2}\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)^3} + \frac{bx}{4a(a+bx^2)^2(bc-a$$

[Out] $1/4*b*x/a/(-a*d+b*c)/(b*x^2+a)^2+1/8*b*(-7*a*d+3*b*c)*x/a^2/(-a*d+b*c)^2/(b*x^2+a)+1/8*(15*a^2*d^2-10*a*b*c*d+3*b^2*c^2)*\arctan(x*b^{(1/2)}/a^{(1/2)})*b^{(1/2)}/a^{(5/2)}/(-a*d+b*c)^3-d^{(5/2)*\arctan(x*d^{(1/2)}/c^{(1/2)})}/(-a*d+b*c)^3/c^{(1/2)}$

Rubi [A] time = 0.20, antiderivative size = 161, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {414, 527, 522, 205}

$$\frac{\sqrt{b}(15a^2d^2-10abcd+3b^2c^2)\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8a^{5/2}(bc-ad)^3} + \frac{bx(3bc-7ad)}{8a^2(a+bx^2)(bc-ad)^2} - \frac{d^{5/2}\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)^3} + \frac{bx}{4a(a+bx^2)^2(bc-a$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^3*(c + d*x^2)), x]

[Out] $(b*x)/(4*a*(b*c - a*d)*(a + b*x^2)^2) + (b*(3*b*c - 7*a*d)*x)/(8*a^2*(b*c - a*d)^2*(a + b*x^2)) + (\text{Sqrt}[b]*(3*b^2*c^2 - 10*a*b*c*d + 15*a^2*d^2)*\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]])/(8*a^{(5/2)}*(b*c - a*d)^3) - (d^{(5/2)*\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]]})/(\text{Sqrt}[c]*(b*c - a*d)^3)$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 522

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(b*c - a*d)*(p+1)), x] + Dist[1/(a*n*(b*c - a*d)*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2)+1)*x^n, x], x], x] /; FreeQ

[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)^3(c+dx^2)} dx &= \frac{bx}{4a(bc-ad)(a+bx^2)^2} - \frac{\int \frac{-3bc+4ad-3bdx^2}{(a+bx^2)^2(c+dx^2)} dx}{4a(bc-ad)} \\ &= \frac{bx}{4a(bc-ad)(a+bx^2)^2} + \frac{b(3bc-7ad)x}{8a^2(bc-ad)^2(a+bx^2)} + \frac{\int \frac{3b^2c^2-7abcd+8a^2d^2+bd(3bc-7ad)x^2}{(a+bx^2)(c+dx^2)} dx}{8a^2(bc-ad)^2} \\ &= \frac{bx}{4a(bc-ad)(a+bx^2)^2} + \frac{b(3bc-7ad)x}{8a^2(bc-ad)^2(a+bx^2)} - \frac{d^3 \int \frac{1}{c+dx^2} dx}{(bc-ad)^3} + \frac{(b(3b^2c^2-10abcd+15a^2d^2))}{8a^2(bc-ad)^2} \\ &= \frac{bx}{4a(bc-ad)(a+bx^2)^2} + \frac{b(3bc-7ad)x}{8a^2(bc-ad)^2(a+bx^2)} + \frac{\sqrt{b}(3b^2c^2-10abcd+15a^2d^2)}{8a^{5/2}(bc-ad)^3} \end{aligned}$$

Mathematica [A] time = 0.28, size = 158, normalized size = 0.98

$$\frac{1}{8} \left(\frac{bx(3bc-7ad)}{a^2(a+bx^2)(bc-ad)^2} - \frac{\sqrt{b}(15a^2d^2-10abcd+3b^2c^2) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{a^{5/2}(ad-bc)^3} - \frac{8d^{5/2} \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{\sqrt{c}(bc-ad)^3} - \frac{2bx}{a(a+bx^2)^2(ad-bc)} \right)$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^3*(c + d*x^2)),x]

[Out] ((-2*b*x)/(a*(-(b*c) + a*d)*(a + b*x^2)^2) + (b*(3*b*c - 7*a*d)*x)/(a^2*(b*c - a*d)^2*(a + b*x^2)) - (Sqrt[b]*(3*b^2*c^2 - 10*a*b*c*d + 15*a^2*d^2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(a^(5/2)*(-(b*c) + a*d)^3) - (8*d^(5/2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(Sqrt[c]*(b*c - a*d)^3))/8

fricas [B] time = 1.47, size = 1587, normalized size = 9.86

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^3/(d*x^2+c),x, algorithm="fricas")

[Out] [1/16*(2*(3*b^4*c^2 - 10*a*b^3*c*d + 7*a^2*b^2*d^2)*x^3 - (3*a^2*b^2*c^2 - 10*a^3*b*c*d + 15*a^4*d^2 + (3*b^4*c^2 - 10*a*b^3*c*d + 15*a^2*b^2*d^2)*x^4 + 2*(3*a*b^3*c^2 - 10*a^2*b^2*c*d + 15*a^3*b*d^2)*x^2)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) - 8*(a^2*b^2*d^2*x^4 + 2*a^3*b*d^2*x^2 + a^4*d^2)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) + 2*(5*a*b^3*c^2 - 14*a^2*b^2*c*d + 9*a^3*b*d^2)*x)/(a^4*b^3*c^3 - 3*a^5*b^2*c^2*d + 3*a^6*b*c*d^2 - a^7*d^3 + (a^2*b^5*c^3 - 3*a^3*b^4*c^2*d + 3*a^4*b^3*c*d^2 - a^5*b^2*d^3)*x^4 + 2*(a^3*b^4*c^3 - 3*a^4*b^3*c^2*d + 3*a^5*b^2*c*d^2 - a^6*b*d^3)*x^2), 1/16*(2*(3*b^4*c^2 - 10*a*b^3*c*d + 7*a^2*b^2*d^2)*x^3 - 16*(a^2*b^2*d^2*x^4 + 2*a^3*b*d^2*x^2 + a^4*d^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) - (3*a^2*b^2*c^2 - 10*a^3*b*c*d + 15*a^4*d^2 + (3*b^4*c^2 - 10*a*b^3*c*d + 15*a^2*b^2*d^2)*x^4 + 2*(3*a*b^3*c^2 - 10*a^2*b^2*c*d + 15*a^3*b*d^2)*x^2)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + 2*(5*a*b^3*c^2 - 14*a^2*b^2*c*d + 9*a^3*b*d^2)*x)/(a^4*b^3*c^3 - 3*a^5*b^2*c^2*d + 3*a^6*b*c*d^2 - a^7*d^3 + (a^2*b^5*c^3 - 3*a^3*b^4*c^2*d + 3*a^4*b^3*c*d^2 - a^5*b^2*d^3)*x^4 + 2*(a^3*b^4*c^3 - 3*a^4*b^3*c^2*d + 3*a^5*b^2*c*d^2 - a^6*b*d^3)*x^2)

$$d^2 - a^5 b^2 d^3) x^4 + 2(a^3 b^4 c^3 - 3a^4 b^3 c^2 d + 3a^5 b^2 c d^2 - a^6 b d^3) x^2, 1/8((3b^4 c^2 - 10a b^3 c d + 7a^2 b^2 d^2) x^3 + (3a^2 b^2 c^2 - 10a^3 b c d + 15a^4 d^2 + (3b^4 c^2 - 10a b^3 c d + 15a^2 b^2 d^2) x^4 + 2(3a^3 b^3 c^2 - 10a^2 b^2 c d + 15a^3 b d^2) x^2) \sqrt{b/a} \arctan(x \sqrt{b/a}) - 4(a^2 b^2 d^2 x^4 + 2a^3 b d^2 x^2 + a^4 d^2) \sqrt{-d/c} \log((d x^2 + 2c x \sqrt{-d/c} - c)/(d x^2 + c)) + (5a^3 b^3 c^2 - 14a^2 b^2 c d + 9a^3 b d^2) x)/(a^4 b^3 c^3 - 3a^5 b^2 c^2 d + 3a^6 b c d^2 - a^7 d^3 + (a^2 b^5 c^3 - 3a^3 b^4 c^2 d + 3a^4 b^3 c d^2 - a^5 b^2 d^3) x^4 + 2(a^3 b^4 c^3 - 3a^4 b^3 c^2 d + 3a^5 b^2 c d^2 - a^6 b d^3) x^2), 1/8((3b^4 c^2 - 10a b^3 c d + 7a^2 b^2 d^2) x^3 + (3a^2 b^2 c^2 - 10a^3 b c d + 15a^4 d^2 + (3b^4 c^2 - 10a b^3 c d + 15a^2 b^2 d^2) x^4 + 2(3a^3 b^3 c^2 - 10a^2 b^2 c d + 15a^3 b d^2) x^2) \sqrt{b/a} \arctan(x \sqrt{b/a}) - 8(a^2 b^2 d^2 x^4 + 2a^3 b d^2 x^2 + a^4 d^2) \sqrt{d/c} \arctan(x \sqrt{d/c}) + (5a^3 b^3 c^2 - 14a^2 b^2 c d + 9a^3 b d^2) x)/(a^4 b^3 c^3 - 3a^5 b^2 c^2 d + 3a^6 b c d^2 - a^7 d^3 + (a^2 b^5 c^3 - 3a^3 b^4 c^2 d + 3a^4 b^3 c d^2 - a^5 b^2 d^3) x^4 + 2(a^3 b^4 c^3 - 3a^4 b^3 c^2 d + 3a^5 b^2 c d^2 - a^6 b d^3) x^2)]$$

giac [A] time = 0.58, size = 218, normalized size = 1.35

$$\frac{d^3 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{(b^3 c^3 - 3ab^2 c^2 d + 3a^2 b c d^2 - a^3 d^3) \sqrt{cd}} + \frac{(3b^3 c^2 - 10ab^2 c d + 15a^2 b d^2) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8(a^2 b^3 c^3 - 3a^3 b^2 c^2 d + 3a^4 b c d^2 - a^5 d^3) \sqrt{ab}} + \frac{3b^3 c x^3 - 7ab^2 d x^3}{8(a^2 b^2 c^2 - 2a^3 b c d)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^3/(d*x^2+c),x, algorithm="giac")

[Out] $-d^3 \arctan(d x / \sqrt{c d}) / ((b^3 c^3 - 3 a b^2 c^2 d + 3 a^2 b c d^2 - a^3 d^3) \sqrt{c d}) + 1/8(3 b^4 c^2 - 10 a b^3 c d + 15 a^2 b^2 d^2) \arctan(b x / \sqrt{a b}) / ((a^2 b^3 c^3 - 3 a^3 b^2 c^2 d + 3 a^4 b c d^2 - a^5 d^3) \sqrt{a b}) + 1/8(3 b^3 c x^3 - 7 a b^2 d x^3 + 5 a b^2 c x - 9 a^2 b d x) / ((a^2 b^2 c^2 - 2 a^3 b c d + a^4 d^2) (b x^2 + a)^2)$

maple [B] time = 0.01, size = 309, normalized size = 1.92

$$\frac{5b^3 c d x^3}{4(ad - bc)^3 (b x^2 + a)^2 a} - \frac{3b^4 c^2 x^3}{8(ad - bc)^3 (b x^2 + a)^2 a^2} - \frac{7b^2 d^2 x^3}{8(ad - bc)^3 (b x^2 + a)^2} - \frac{9ab d^2 x}{8(ad - bc)^3 (b x^2 + a)^2} - \frac{1}{8(ad - bc)^3 (b x^2 + a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^3/(d*x^2+c),x)

[Out] $d^3/(a d - b c)^3/(c d)^{(1/2)} \arctan(1/(c d)^{(1/2)} d x) - 7/8 b^2/(a d - b c)^3/(b x^2 + a)^2 x^3 d^2 + 5/4 b^3/(a d - b c)^3/(b x^2 + a)^2/a x^3 c d - 3/8 b^4/(a d - b c)^3/(b x^2 + a)^2/a^2 x^3 c^2 - 9/8 b/(a d - b c)^3/(b x^2 + a)^2 x a d^2 + 7/4 b^2/(a d - b c)^3/(b x^2 + a)^2 x c d - 5/8 b^3/(a d - b c)^3/(b x^2 + a)^2 x/a c^2 - 15/8 b/(a d - b c)^3/(a b)^{(1/2)} \arctan(1/(a b)^{(1/2)} b x) d^2 + 5/4 b^2/(a d - b c)^3/a/(a b)^{(1/2)} \arctan(1/(a b)^{(1/2)} b x) c d - 3/8 b^3/(a d - b c)^3/a^2/(a b)^{(1/2)} \arctan(1/(a b)^{(1/2)} b x) c^2$

maxima [A] time = 3.16, size = 278, normalized size = 1.73

$$\frac{d^3 \arctan\left(\frac{dx}{\sqrt{cd}}\right)}{(b^3 c^3 - 3ab^2 c^2 d + 3a^2 b c d^2 - a^3 d^3) \sqrt{cd}} + \frac{(3b^3 c^2 - 10ab^2 c d + 15a^2 b d^2) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8(a^2 b^3 c^3 - 3a^3 b^2 c^2 d + 3a^4 b c d^2 - a^5 d^3) \sqrt{ab}} + \frac{3b^3 c x^3 - 7ab^2 d x^3}{8(a^4 b^2 c^2 - 2a^5 b c d)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^3/(d*x^2+c),x, algorithm="maxima")

```
[Out] -d^3*arctan(d*x/sqrt(c*d))/((b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(c*d)) + 1/8*(3*b^3*c^2 - 10*a*b^2*c*d + 15*a^2*b*d^2)*arctan(b*x/sqrt(a*b))/((a^2*b^3*c^3 - 3*a^3*b^2*c^2*d + 3*a^4*b*c*d^2 - a^5*d^3)*sqrt(a*b)) + 1/8*((3*b^3*c - 7*a*b^2*d)*x^3 + (5*a*b^2*c - 9*a^2*b*d)*x)/(a^4*b^2*c^2 - 2*a^5*b*c*d + a^6*d^2 + (a^2*b^4*c^2 - 2*a^3*b^3*c*d + a^4*b^2*d^2)*x^4 + 2*(a^3*b^3*c^2 - 2*a^4*b^2*c*d + a^5*b*d^2)*x^2)
```

mupad [B] time = 6.89, size = 6033, normalized size = 37.47

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((a + b*x^2)^3*(c + d*x^2)),x)
```

```
[Out] ((x^3*(3*b^3*c - 7*a*b^2*d))/(8*a^2*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d)) + (x*(5*b^2*c - 9*a*b*d))/(8*a*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d)))/(a^2 + b^2*x^4 + 2*a*b*x^2) + (atan((((x*(289*a^4*b^3*d^7 + 9*b^7*c^4*d^3 - 60*a*b^6*c^3*d^4 - 300*a^3*b^4*c*d^6 + 190*a^2*b^5*c^2*d^5))/(32*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3))) - ((-c*d^5)^(1/2))*((256*a^10*b^2*d^10 - 1760*a^9*b^3*c*d^9 + 96*a^2*b^10*c^8*d^2 - 800*a^3*b^9*c^7*d^3 + 3040*a^4*b^8*c^6*d^4 - 6816*a^5*b^7*c^5*d^5 + 9760*a^6*b^6*c^4*d^6 - 9056*a^7*b^5*c^3*d^7 + 5280*a^8*b^4*c^2*d^8)/(64*(a^10*d^6 + a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^6*b^4*c^4*d^2 - 20*a^7*b^3*c^3*d^3 + 15*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5))) - (x*(-c*d^5)^(1/2))*((256*a^11*b^2*d^9 - 1280*a^10*b^3*c*d^8 + 256*a^4*b^9*c^7*d^2 - 1280*a^5*b^8*c^6*d^3 + 2304*a^6*b^7*c^5*d^4 - 1280*a^7*b^6*c^4*d^5 - 1280*a^8*b^5*c^3*d^6 + 2304*a^9*b^4*c^2*d^7)))/(64*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3)))))/(2*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d))*(-c*d^5)^(1/2)*i)/(2*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)) + (((x*(289*a^4*b^3*d^7 + 9*b^7*c^4*d^3 - 60*a*b^6*c^3*d^4 - 300*a^3*b^4*c*d^6 + 190*a^2*b^5*c^2*d^5))/(32*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3))) + ((-c*d^5)^(1/2))*((256*a^10*b^2*d^10 - 1760*a^9*b^3*c*d^9 + 96*a^2*b^10*c^8*d^2 - 800*a^3*b^9*c^7*d^3 + 3040*a^4*b^8*c^6*d^4 - 6816*a^5*b^7*c^5*d^5 + 9760*a^6*b^6*c^4*d^6 - 9056*a^7*b^5*c^3*d^7 + 5280*a^8*b^4*c^2*d^8)/(64*(a^10*d^6 + a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^6*b^4*c^4*d^2 - 20*a^7*b^3*c^3*d^3 + 15*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5))) + (x*(-c*d^5)^(1/2))*((256*a^11*b^2*d^9 - 1280*a^10*b^3*c*d^8 + 256*a^4*b^9*c^7*d^2 - 1280*a^5*b^8*c^6*d^3 + 2304*a^6*b^7*c^5*d^4 - 1280*a^7*b^6*c^4*d^5 - 1280*a^8*b^5*c^3*d^6 + 2304*a^9*b^4*c^2*d^7)))/(64*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3)))))/(2*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d))*(-c*d^5)^(1/2)*i)/(2*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)))/((105*a^3*b^3*d^8 - 9*b^6*c^3*d^5 + 51*a*b^5*c^2*d^6 - 15*a^2*b^4*c*d^7)/(32*(a^10*d^6 + a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^6*b^4*c^4*d^2 - 20*a^7*b^3*c^3*d^3 + 15*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5))) - (((x*(289*a^4*b^3*d^7 + 9*b^7*c^4*d^3 - 60*a*b^6*c^3*d^4 - 300*a^3*b^4*c*d^6 + 190*a^2*b^5*c^2*d^5))/(32*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3))) - ((-c*d^5)^(1/2))*((256*a^10*b^2*d^10 - 1760*a^9*b^3*c*d^9 + 96*a^2*b^10*c^8*d^2 - 800*a^3*b^9*c^7*d^3 + 3040*a^4*b^8*c^6*d^4 - 6816*a^5*b^7*c^5*d^5 + 9760*a^6*b^6*c^4*d^6 - 9056*a^7*b^5*c^3*d^7 + 5280*a^8*b^4*c^2*d^8)/(64*(a^10*d^6 + a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^6*b^4*c^4*d^2 - 20*a^7*b^3*c^3*d^3 + 15*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5))) - (x*(-c*d^5)^(1/2))*((256*a^11*b^2*d^9 - 1280*a^10*b^3*c*d^8 + 256*a^4*b^9*c^7*d^2 - 1280*a^5*b^8*c^6*d^3 + 2304*a^6*b^7*c^5*d^4 - 1280*a^7*b^6*c^4*d^5 - 1280*a^8*b^5*c^3*d^6 + 2304*a^9*b^4*c^2*d^7)))/(64*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3)))))/(2*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d))*(-c*d^5)^(1/2))/(2*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)) + (((x*(289*a^4*b^3*d^7 + 9*b^7*c^4*d^3 - 60*a
```

$$\begin{aligned}
& *b^6*c^3*d^4 - 300*a^3*b^4*c*d^6 + 190*a^2*b^5*c^2*d^5)) / (32*(a^8*d^4 + a^4 \\
& *b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3)) + ((-c*d^5 \\
&)^{(1/2)} * ((256*a^10*b^2*d^10 - 1760*a^9*b^3*c*d^9 + 96*a^2*b^10*c^8*d^2 - 80 \\
& 0*a^3*b^9*c^7*d^3 + 3040*a^4*b^8*c^6*d^4 - 6816*a^5*b^7*c^5*d^5 + 9760*a^6* \\
& b^6*c^4*d^6 - 9056*a^7*b^5*c^3*d^7 + 5280*a^8*b^4*c^2*d^8) / (64*(a^10*d^6 + \\
& a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^6*b^4*c^4*d^2 - 20*a^7*b^3*c^3*d^3 + 1 \\
& 5*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5)) + (x*(-c*d^5)^{(1/2)} * (256*a^11*b^2*d^9 - \\
& 1280*a^10*b^3*c*d^8 + 256*a^4*b^9*c^7*d^2 - 1280*a^5*b^8*c^6*d^3 + 2304*a^ \\
& 6*b^7*c^5*d^4 - 1280*a^7*b^6*c^4*d^5 - 1280*a^8*b^5*c^3*d^6 + 2304*a^9*b^4* \\
& c^2*d^7)) / (64*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)) * (a^8* \\
& d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3)) \\
& / (2*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)) * (-c*d^5)^{(1/2)} \\
&) / (2*(b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d)) * (-c*d^5)^{(1/2)} \\
& * i) / (b^3*c^4 - a^3*c*d^3 + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^3*d) + (\operatorname{atan}(((\\
& (x*(289*a^4*b^3*d^7 + 9*b^7*c^4*d^3 - 60*a*b^6*c^3*d^4 - 300*a^3*b^4*c*d^6 \\
& + 190*a^2*b^5*c^2*d^5)) / (32*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a \\
& ^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3)) - (((256*a^10*b^2*d^10 - 1760*a^9*b^3*c*d^ \\
& 9 + 96*a^2*b^10*c^8*d^2 - 800*a^3*b^9*c^7*d^3 + 3040*a^4*b^8*c^6*d^4 - 6816 \\
& *a^5*b^7*c^5*d^5 + 9760*a^6*b^6*c^4*d^6 - 9056*a^7*b^5*c^3*d^7 + 5280*a^8*b \\
& ^4*c^2*d^8) / (64*(a^10*d^6 + a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^6*b^4*c^4* \\
& d^2 - 20*a^7*b^3*c^3*d^3 + 15*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5)) - (x*(-a^5* \\
& b)^{(1/2)} * (15*a^2*d^2 + 3*b^2*c^2 - 10*a*b*c*d) * (256*a^11*b^2*d^9 - 1280*a^1 \\
& 0*b^3*c*d^8 + 256*a^4*b^9*c^7*d^2 - 1280*a^5*b^8*c^6*d^3 + 2304*a^6*b^7*c^5 \\
& *d^4 - 1280*a^7*b^6*c^4*d^5 - 1280*a^8*b^5*c^3*d^6 + 2304*a^9*b^4*c^2*d^7)) \\
& / (512*(a^8*d^3 - a^5*b^3*c^3 + 3*a^6*b^2*c^2*d - 3*a^7*b*c*d^2)) * (-a^5* \\
& b)^{(1/2)} * (15*a^2*d^2 + 3*b^2*c^2 - 10*a*b*c*d)) / (16*(a^8*d^3 - a^5*b^3*c^3 \\
& + 3*a^6*b^2*c^2*d - 3*a^7*b*c*d^2)) * (-a^5*b)^{(1/2)} * (15*a^2*d^2 + 3*b^2*c^2 \\
& - 10*a*b*c*d) * i) / (16*(a^8*d^3 - a^5*b^3*c^3 + 3*a^6*b^2*c^2*d - 3*a^7*b*c \\
& *d^2)) + (((x*(289*a^4*b^3*d^7 + 9*b^7*c^4*d^3 - 60*a*b^6*c^3*d^4 - 300*a^3 \\
& *b^4*c*d^6 + 190*a^2*b^5*c^2*d^5)) / (32*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^ \\
& ^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3)) + (((256*a^10*b^2*d^10 - 1760*a^ \\
& 9*b^3*c*d^9 + 96*a^2*b^10*c^8*d^2 - 800*a^3*b^9*c^7*d^3 + 3040*a^4*b^8*c^6* \\
& d^4 - 6816*a^5*b^7*c^5*d^5 + 9760*a^6*b^6*c^4*d^6 - 9056*a^7*b^5*c^3*d^7 + \\
& 5280*a^8*b^4*c^2*d^8) / (64*(a^10*d^6 + a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^ \\
& 6*b^4*c^4*d^2 - 20*a^7*b^3*c^3*d^3 + 15*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5)) + \\
& (x*(-a^5*b)^{(1/2)} * (15*a^2*d^2 + 3*b^2*c^2 - 10*a*b*c*d) * (256*a^11*b^2*d^9 \\
& - 1280*a^10*b^3*c*d^8 + 256*a^4*b^9*c^7*d^2 - 1280*a^5*b^8*c^6*d^3 + 2304*a^ \\
& 6*b^7*c^5*d^4 - 1280*a^7*b^6*c^4*d^5 - 1280*a^8*b^5*c^3*d^6 + 2304*a^9*b^4 \\
& *c^2*d^7)) / (512*(a^8*d^3 - a^5*b^3*c^3 + 3*a^6*b^2*c^2*d - 3*a^7*b*c*d^2)) * (\\
& a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3 \\
&)) * (-a^5*b)^{(1/2)} * (15*a^2*d^2 + 3*b^2*c^2 - 10*a*b*c*d)) / (16*(a^8*d^3 - a^ \\
& 5*b^3*c^3 + 3*a^6*b^2*c^2*d - 3*a^7*b*c*d^2)) * (-a^5*b)^{(1/2)} * (15*a^2*d^2 + \\
& 3*b^2*c^2 - 10*a*b*c*d) * i) / (16*(a^8*d^3 - a^5*b^3*c^3 + 3*a^6*b^2*c^2*d - \\
& 3*a^7*b*c*d^2)) / ((105*a^3*b^3*d^8 - 9*b^6*c^3*d^5 + 51*a*b^5*c^2*d^6 - 11 \\
& 5*a^2*b^4*c*d^7) / (32*(a^10*d^6 + a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^6*b^4 \\
& *c^4*d^2 - 20*a^7*b^3*c^3*d^3 + 15*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5)) - (((x \\
& *(289*a^4*b^3*d^7 + 9*b^7*c^4*d^3 - 60*a*b^6*c^3*d^4 - 300*a^3*b^4*c*d^6 + \\
& 190*a^2*b^5*c^2*d^5)) / (32*(a^8*d^4 + a^4*b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6* \\
& b^2*c^2*d^2 - 4*a^7*b*c*d^3)) - (((256*a^10*b^2*d^10 - 1760*a^9*b^3*c*d^9 + \\
& 96*a^2*b^10*c^8*d^2 - 800*a^3*b^9*c^7*d^3 + 3040*a^4*b^8*c^6*d^4 - 6816*a^ \\
& 5*b^7*c^5*d^5 + 9760*a^6*b^6*c^4*d^6 - 9056*a^7*b^5*c^3*d^7 + 5280*a^8*b^4* \\
& c^2*d^8) / (64*(a^10*d^6 + a^4*b^6*c^6 - 6*a^5*b^5*c^5*d + 15*a^6*b^4*c^4*d^2 \\
& - 20*a^7*b^3*c^3*d^3 + 15*a^8*b^2*c^2*d^4 - 6*a^9*b*c*d^5)) - (x*(-a^5*b)^ \\
& (1/2) * (15*a^2*d^2 + 3*b^2*c^2 - 10*a*b*c*d) * (256*a^11*b^2*d^9 - 1280*a^10*b \\
& ^3*c*d^8 + 256*a^4*b^9*c^7*d^2 - 1280*a^5*b^8*c^6*d^3 + 2304*a^6*b^7*c^5*d^ \\
& 4 - 1280*a^7*b^6*c^4*d^5 - 1280*a^8*b^5*c^3*d^6 + 2304*a^9*b^4*c^2*d^7)) / (5 \\
& 12*(a^8*d^3 - a^5*b^3*c^3 + 3*a^6*b^2*c^2*d - 3*a^7*b*c*d^2)) * (a^8*d^4 + a^4 \\
& *b^4*c^4 - 4*a^5*b^3*c^3*d + 6*a^6*b^2*c^2*d^2 - 4*a^7*b*c*d^3)) * (-a^5*b)^
\end{aligned}$$

$$\begin{aligned} & \left(\frac{1}{2}\right) * (15 * a^2 * d^2 + 3 * b^2 * c^2 - 10 * a * b * c * d) / (16 * (a^8 * d^3 - a^5 * b^3 * c^3 + 3 * a^6 * b^2 * c^2 * d - 3 * a^7 * b * c * d^2)) * (-a^5 * b)^{(1/2)} * (15 * a^2 * d^2 + 3 * b^2 * c^2 - 10 * a * b * c * d) / (16 * (a^8 * d^3 - a^5 * b^3 * c^3 + 3 * a^6 * b^2 * c^2 * d - 3 * a^7 * b * c * d^2)) \\ & + ((x * (289 * a^4 * b^3 * d^7 + 9 * b^7 * c^4 * d^3 - 60 * a * b^6 * c^3 * d^4 - 300 * a^3 * b^4 * c * d^6 + 190 * a^2 * b^5 * c^2 * d^5)) / (32 * (a^8 * d^4 + a^4 * b^4 * c^4 - 4 * a^5 * b^3 * c^3 * d + 6 * a^6 * b^2 * c^2 * d^2 - 4 * a^7 * b * c * d^3)) + (((256 * a^{10} * b^2 * d^{10} - 1760 * a^9 * b^3 * c * d^9 + 96 * a^2 * b^{10} * c^8 * d^2 - 800 * a^3 * b^9 * c^7 * d^3 + 3040 * a^4 * b^8 * c^6 * d^4 - 6816 * a^5 * b^7 * c^5 * d^5 + 9760 * a^6 * b^6 * c^4 * d^6 - 9056 * a^7 * b^5 * c^3 * d^7 + 5280 * a^8 * b^4 * c^2 * d^8) / (64 * (a^{10} * d^6 + a^4 * b^6 * c^6 - 6 * a^5 * b^5 * c^5 * d + 15 * a^6 * b^4 * c^4 * d^2 - 20 * a^7 * b^3 * c^3 * d^3 + 15 * a^8 * b^2 * c^2 * d^4 - 6 * a^9 * b * c * d^5)) + (x * (-a^5 * b)^{(1/2)} * (15 * a^2 * d^2 + 3 * b^2 * c^2 - 10 * a * b * c * d) * (256 * a^{11} * b^2 * d^9 - 1280 * a^{10} * b^3 * c * d^8 + 256 * a^4 * b^9 * c^7 * d^2 - 1280 * a^5 * b^8 * c^6 * d^3 + 2304 * a^6 * b^7 * c^5 * d^4 - 1280 * a^7 * b^6 * c^4 * d^5 - 1280 * a^8 * b^5 * c^3 * d^6 + 2304 * a^9 * b^4 * c^2 * d^7)) / (512 * (a^8 * d^3 - a^5 * b^3 * c^3 + 3 * a^6 * b^2 * c^2 * d - 3 * a^7 * b * c * d^2)) * (a^8 * d^4 + a^4 * b^4 * c^4 - 4 * a^5 * b^3 * c^3 * d + 6 * a^6 * b^2 * c^2 * d^2 - 4 * a^7 * b * c * d^3)) * (-a^5 * b)^{(1/2)} * (15 * a^2 * d^2 + 3 * b^2 * c^2 - 10 * a * b * c * d) / (16 * (a^8 * d^3 - a^5 * b^3 * c^3 + 3 * a^6 * b^2 * c^2 * d - 3 * a^7 * b * c * d^2)) * (-a^5 * b)^{(1/2)} * (15 * a^2 * d^2 + 3 * b^2 * c^2 - 10 * a * b * c * d) / (16 * (a^8 * d^3 - a^5 * b^3 * c^3 + 3 * a^6 * b^2 * c^2 * d - 3 * a^7 * b * c * d^2)) * (-a^5 * b)^{(1/2)} * (15 * a^2 * d^2 + 3 * b^2 * c^2 - 10 * a * b * c * d) * 1i) / (8 * (a^8 * d^3 - a^5 * b^3 * c^3 + 3 * a^6 * b^2 * c^2 * d - 3 * a^7 * b * c * d^2)) \end{aligned}$$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**3/(d*x**2+c),x)

[Out] Timed out

$$3.41 \quad \int \frac{1}{(a+bx^2)^3 (c+dx^2)^2} dx$$

Optimal. Leaf size=236

$$\frac{dx(bc-4ad)(ad+3bc)}{8a^2c(c+dx^2)(bc-ad)^3} + \frac{3bx(bc-3ad)}{8a^2(a+bx^2)(c+dx^2)(bc-ad)^2} + \frac{b^{3/2}(35a^2d^2-14abcd+3b^2c^2)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}(bc-ad)^4}$$

[Out] $1/8*d*(-4*a*d+b*c)*(a*d+3*b*c)*x/a^2/c/(-a*d+b*c)^3/(d*x^2+c)+1/4*b*x/a/(-a*d+b*c)/(b*x^2+a)^2/(d*x^2+c)+3/8*b*(-3*a*d+b*c)*x/a^2/(-a*d+b*c)^2/(b*x^2+a)/(d*x^2+c)+1/8*b^(3/2)*(35*a^2*d^2-14*a*b*c*d+3*b^2*c^2)*\arctan(x*b^(1/2)/a^(1/2))/a^(5/2)/(-a*d+b*c)^4-1/2*d^(5/2)*(-a*d+7*b*c)*\arctan(x*d^(1/2)/c^(1/2))/c^(3/2)/(-a*d+b*c)^4$

Rubi [A] time = 0.31, antiderivative size = 236, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {414, 527, 522, 205}

$$\frac{b^{3/2}(35a^2d^2-14abcd+3b^2c^2)\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{8a^{5/2}(bc-ad)^4} + \frac{dx(bc-4ad)(ad+3bc)}{8a^2c(c+dx^2)(bc-ad)^3} + \frac{3bx(bc-3ad)}{8a^2(a+bx^2)(c+dx^2)(bc-ad)^2}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^3*(c + d*x^2)^2), x]

[Out] $(d*(b*c-4*a*d)*(3*b*c+a*d)*x)/(8*a^2*c*(b*c-a*d)^3*(c+d*x^2)) + (b*x)/(4*a*(b*c-a*d)*(a+b*x^2)^2*(c+d*x^2)) + (3*b*(b*c-3*a*d)*x)/(8*a^2*(b*c-a*d)^2*(a+b*x^2)*(c+d*x^2)) + (b^(3/2)*(3*b^2*c^2-14*a*b*c*d+35*a^2*d^2)*\text{ArcTan}[\text{Sqrt}[b]*x/\text{Sqrt}[a]])/(8*a^(5/2)*(b*c-a*d)^4) - (d^(5/2)*(7*b*c-a*d)*\text{ArcTan}[\text{Sqrt}[d]*x/\text{Sqrt}[c]])/(2*c^(3/2)*(b*c-a*d)^4)$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c-a*d)), x] + Dist[1/(a*n*(p+1)*(b*c-a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c-a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c-a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 522

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(b*c - a*d)*(p+1)), x] + Dist[1/(a*n*(b*c - a*d)*(p

+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + bx^2)^3 (c + dx^2)^2} dx &= \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)} - \frac{\int \frac{-3bc + 4ad - 5bdx^2}{(a + bx^2)^2 (c + dx^2)^2} dx}{4a(bc - ad)} \\ &= \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)} + \frac{3b(bc - 3ad)x}{8a^2(bc - ad)^2 (a + bx^2)(c + dx^2)} + \frac{\int \frac{3b^2c^2 - 5abc}{(a + bx^2)^2 (c + dx^2)^2} dx}{8} \\ &= \frac{d(bc - 4ad)(3bc + ad)x}{8a^2c(bc - ad)^3 (c + dx^2)} + \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)} + \frac{3b(bc - 3ad)}{8a^2(bc - ad)^2 (a + bx^2)} \\ &= \frac{d(bc - 4ad)(3bc + ad)x}{8a^2c(bc - ad)^3 (c + dx^2)} + \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)} + \frac{3b(bc - 3ad)}{8a^2(bc - ad)^2 (a + bx^2)} \\ &= \frac{d(bc - 4ad)(3bc + ad)x}{8a^2c(bc - ad)^3 (c + dx^2)} + \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)} + \frac{3b(bc - 3ad)}{8a^2(bc - ad)^2 (a + bx^2)} \end{aligned}$$

Mathematica [A] time = 0.42, size = 197, normalized size = 0.83

$$\frac{1}{8} \left(\frac{b^2x(11ad - 3bc)}{a^2(a + bx^2)(ad - bc)^3} + \frac{b^{3/2}(35a^2d^2 - 14abcd + 3b^2c^2) \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{a^{5/2}(bc - ad)^4} + \frac{2b^2x}{a(a + bx^2)^2(bc - ad)^2} + \frac{4d^{5/2}(ad - 7b^2)}{c^{3/2}(bc - ad)^2} \right)$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^3*(c + d*x^2)^2), x]

[Out] ((2*b^2*x)/(a*(b*c - a*d)^2*(a + b*x^2)^2) + (b^2*(-3*b*c + 11*a*d)*x)/(a^2*(-(b*c) + a*d)^3*(a + b*x^2)) - (4*d^3*x)/(c*(b*c - a*d)^3*(c + d*x^2)) + (b^(3/2)*(3*b^2*c^2 - 14*a*b*c*d + 35*a^2*d^2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(a^(5/2)*(b*c - a*d)^4) + (4*d^(5/2)*(-7*b*c + a*d)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(c^(3/2)*(b*c - a*d)^4))/8

fricas [B] time = 7.25, size = 3251, normalized size = 13.78

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^3/(d*x^2+c)^2,x, algorithm="fricas")

[Out] [1/16*(2*(3*b^5*c^3*d - 14*a*b^4*c^2*d^2 + 7*a^2*b^3*c^2*d^2 + 4*a^3*b^2*d^4)*x^5 + 2*(3*b^5*c^4 - 9*a*b^4*c^3*d - 7*a^2*b^3*c^2*d^2 + 5*a^3*b^2*c*d^3 + 8*a^4*b*d^4)*x^3 + (3*a^2*b^3*c^4 - 14*a^3*b^2*c^3*d + 35*a^4*b*c^2*d^2 + (3*b^5*c^3*d - 14*a*b^4*c^2*d^2 + 35*a^2*b^3*c*d^3)*x^6 + (3*b^5*c^4 - 8*a*b^4*c^3*d + 7*a^2*b^3*c^2*d^2 + 70*a^3*b^2*c*d^3)*x^4 + (6*a*b^4*c^4 - 25*a^2*b^3*c^3*d + 56*a^3*b^2*c^2*d^2 + 35*a^4*b*c*d^3)*x^2)*sqrt(-b/a)*log((b*x^2 + 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) - 4*(7*a^4*b*c^2*d^2 - a^5*c*d^3 + (7*a^2*b^3*c*d^3 - a^3*b^2*d^4)*x^6 + (7*a^2*b^3*c^2*d^2 + 13*a^3*b^2*c*d^3 - a^4*b*d^4)*x^4 + (3*a^2*b^3*c^2*d^2 + 13*a^3*b^2*c*d^3 - a^4*b*d^4)*x^2 + (3*a^2*b^3*c^2*d^2 + 13*a^3*b^2*c*d^3 - a^4*b*d^4)*x^0)]/16

$$\begin{aligned}
& 3 - 2a^4bd^4)x^4 + (14a^3b^2c^2d^2 + 5a^4b^3cd^3 - a^5d^4)x^2) * \\
& \sqrt{-d/c} * \log((dx^2 + 2cx\sqrt{-d/c} - c)/(dx^2 + c)) + 2(5a^4b^3cd^4 \\
& - 18a^2b^3c^3d + 13a^3b^2c^2d^2 - 4a^4b^3cd^3 + 4a^5d^4)x)/(a \\
& ^4b^4c^6 - 4a^5b^3c^5d + 6a^6b^2c^4d^2 - 4a^7b^3cd^3 + a^8c^2 \\
& ^2d^4 + (a^2b^6c^5d - 4a^3b^5c^4d^2 + 6a^4b^4c^3d^3 - 4a^5b^3c^2 \\
& ^2d^4 + a^6b^2cd^5)x^6 + (a^2b^6c^6 - 2a^3b^5c^5d - 2a^4b^4c^4 \\
& ^4d^2 + 8a^5b^3c^3d^3 - 7a^6b^2c^2d^4 + 2a^7b^3cd^5)x^4 + (2a^3 \\
& ^3b^5c^6 - 7a^4b^4c^5d + 8a^5b^3c^4d^2 - 2a^6b^2c^3d^3 - 2a^7 \\
& ^7b^3c^2d^4 + a^8cd^5)x^2), 1/16*(2(3b^5c^3d - 14ab^4c^2d^2 + 7a \\
& ^2b^3cd^3 + 4a^3b^2d^4)x^5 + 2(3b^5c^4 - 9ab^4c^3d - 7a^2b^3 \\
& ^3c^2d^2 + 5a^3b^2cd^3 + 8a^4bd^4)x^3 - 8(7a^4b^3c^2d^2 - a^5c \\
& ^5d^3 + (7a^2b^3cd^3 - a^3b^2d^4)x^6 + (7a^2b^3c^2d^2 + 13a^3b^2 \\
& ^2cd^3 - 2a^4bd^4)x^4 + (14a^3b^2c^2d^2 + 5a^4b^3cd^3 - a^5d^4) \\
& ^2)x^2) * \sqrt{d/c} * \arctan(x\sqrt{d/c}) + (3a^2b^3c^4 - 14a^3b^2c^3d + 3 \\
& ^5a^4b^3cd^2 + (3b^5c^3d - 14ab^4c^2d^2 + 35a^2b^3cd^3)x^6 + \\
& (3b^5c^4 - 8ab^4c^3d + 7a^2b^3c^2d^2 + 70a^3b^2cd^3)x^4 + (6 \\
& ^6ab^4c^4 - 25a^2b^3c^3d + 56a^3b^2c^2d^2 + 35a^4b^3cd^3)x^2) * \\
& \sqrt{-b/a} * \log((bx^2 + 2ax\sqrt{-b/a} - a)/(bx^2 + a)) + 2(5a^4b^3cd^4 \\
& - 18a^2b^3c^3d + 13a^3b^2c^2d^2 - 4a^4b^3cd^3 + 4a^5d^4)x)/(a \\
& ^4b^4c^6 - 4a^5b^3c^5d + 6a^6b^2c^4d^2 - 4a^7b^3cd^3 + a^8c^2 \\
& ^2d^4 + (a^2b^6c^5d - 4a^3b^5c^4d^2 + 6a^4b^4c^3d^3 - 4a^5b^3c^2 \\
& ^2d^4 + a^6b^2cd^5)x^6 + (a^2b^6c^6 - 2a^3b^5c^5d - 2a^4b^4c^4 \\
& ^4d^2 + 8a^5b^3c^3d^3 - 7a^6b^2c^2d^4 + 2a^7b^3cd^5)x^4 + (2a^3 \\
& ^3b^5c^6 - 7a^4b^4c^5d + 8a^5b^3c^4d^2 - 2a^6b^2c^3d^3 - 2a^7 \\
& ^7b^3c^2d^4 + a^8cd^5)x^2), 1/8*((3b^5c^3d - 14ab^4c^2d^2 + 7a^2b^3 \\
& ^3cd^3 + 4a^3b^2d^4)x^5 + (3b^5c^4 - 9ab^4c^3d - 7a^2b^3c^2 \\
& ^2d^2 + 5a^3b^2cd^3 + 8a^4bd^4)x^3 + (3a^2b^3c^4 - 14a^3b^2c^3 \\
& ^3d + 35a^4b^3cd^2 + (3b^5c^3d - 14ab^4c^2d^2 + 35a^2b^3cd^3) \\
& ^2)x^6 + (3b^5c^4 - 8ab^4c^3d + 7a^2b^3c^2d^2 + 70a^3b^2cd^3)x^4 \\
& + (6ab^4c^4 - 25a^2b^3c^3d + 56a^3b^2c^2d^2 + 35a^4b^3cd^3) \\
& ^2)x^2) * \sqrt{b/a} * \arctan(x\sqrt{b/a}) - 2(7a^4b^3cd^2 - a^5cd^3 + (7a \\
& ^2b^3cd^3 - a^3b^2d^4)x^6 + (7a^2b^3c^2d^2 + 13a^3b^2cd^3 - 2 \\
& ^4ab^3d^4)x^4 + (14a^3b^2c^2d^2 + 5a^4b^3cd^3 - a^5d^4)x^2) * \sqrt{ \\
& -d/c} * \log((dx^2 + 2cx\sqrt{-d/c} - c)/(dx^2 + c)) + (5a^4b^3cd^4 - 18a \\
& ^2b^3c^3d + 13a^3b^2c^2d^2 - 4a^4b^3cd^3 + 4a^5d^4)x)/(a^4b^4c^6 - 4a^5b^3c^5d + 6a^6b^2c^4 \\
& ^4d^2 - 4a^7b^3cd^3 + a^8c^2d^4 + (a^2b^6c^5d - 4a^3b^5c^4d^2 + 6a^4b^4c^3d^3 - 4a^5b^3c^2d^4 \\
& + a^6b^2cd^5)x^6 + (a^2b^6c^6 - 2a^3b^5c^5d - 2a^4b^4c^4d^2 \\
& + 8a^5b^3c^3d^3 - 7a^6b^2c^2d^4 + 2a^7b^3cd^5)x^4 + (2a^3b^5c^6 - 7a^4b^4c^5d + 8a^5b^3c^4 \\
& ^4d^2 - 2a^6b^2c^3d^3 - 2a^7b^3c^2d^4 + a^8cd^5)x^2), 1/8*((3b^5c^3d - 14ab^4c^2d^2 + 7a^2b^3cd^3 \\
& + 4a^3b^2d^4)x^5 + (3b^5c^4 - 9ab^4c^3d - 7a^2b^3c^2d^2 + 5a^3b^2cd^3 + 8a^4bd^4)x^3 + (3a^2b^3c^4 - 14a^3b^2c^3d + 35 \\
& ^4ab^3cd^2 + (3b^5c^3d - 14ab^4c^2d^2 + 35a^2b^3cd^3)x^6 + \\
& (3b^5c^4 - 8ab^4c^3d + 7a^2b^3c^2d^2 + 70a^3b^2cd^3)x^4 + (6 \\
& ^6ab^4c^4 - 25a^2b^3c^3d + 56a^3b^2c^2d^2 + 35a^4b^3cd^3)x^2) * \sqrt{ \\
& b/a} * \arctan(x\sqrt{b/a}) - 4(7a^4b^3cd^2 - a^5cd^3 + (7a^2b^3c^2 \\
& ^2d^2 - a^3b^2d^4)x^6 + (7a^2b^3c^2d^2 + 13a^3b^2cd^3 - 2a^4b^3 \\
& ^3d^4)x^4 + (14a^3b^2c^2d^2 + 5a^4b^3cd^3 - a^5d^4)x^2) * \sqrt{d/c} * \ar \\
& \arctan(x\sqrt{d/c}) + (5a^4b^3cd^4 - 18a^2b^3c^3d + 13a^3b^2c^2d^2 - \\
& 4a^4b^3cd^3 + 4a^5d^4)x)/(a^4b^4c^6 - 4a^5b^3c^5d + 6a^6b^2c^4 \\
& ^4d^2 - 4a^7b^3cd^3 + a^8c^2d^4 + (a^2b^6c^5d - 4a^3b^5c^4d^2 \\
& + 6a^4b^4c^3d^3 - 4a^5b^3c^2d^4 + a^6b^2cd^5)x^6 + (a^2b^6c^6 - \\
& 2a^3b^5c^5d - 2a^4b^4c^4d^2 + 8a^5b^3c^3d^3 - 7a^6b^2c^2d^4 \\
& + 2a^7b^3cd^5)x^4 + (2a^3b^5c^6 - 7a^4b^4c^5d + 8a^5b^3c^4 \\
& ^4d^2 - 2a^6b^2c^3d^3 - 2a^7b^3c^2d^4 + a^8cd^5)x^2)]
\end{aligned}$$

$$\begin{aligned}
& 2*b^{13}*c^{12}*d^2)/2 + (35*a^3*b^{12}*c^{11}*d^3)/2 - 98*a^4*b^{11}*c^{10}*d^4 + 336* \\
& a^5*b^{10}*c^9*d^5 - 765*a^6*b^9*c^8*d^6 + 1197*a^7*b^8*c^7*d^7 - 1302*a^8*b^7* \\
& c^6*d^8 + 978*a^9*b^6*c^5*d^9 - (987*a^{10}*b^5*c^4*d^{10})/2 + (315*a^{11}*b^4* \\
& c^3*d^{11})/2 - 28*a^{12}*b^3*c^2*d^{12})/(a^4*b^9*c^{11} - a^{13}*c^2*d^9 - 9*a^5*b^8* \\
& c^{10}*d + 9*a^{12}*b*c^3*d^8 + 36*a^6*b^7*c^9*d^2 - 84*a^7*b^6*c^8*d^3 + 12 \\
& 6*a^8*b^5*c^7*d^4 - 126*a^9*b^4*c^6*d^5 + 84*a^{10}*b^3*c^5*d^6 - 36*a^{11}*b^2* \\
& c^4*d^7) - (x*(-a^5*b^3)^{(1/2)}*(35*a^2*d^2 + 3*b^2*c^2 - 14*a*b*c*d)*(256* \\
& a^4*b^{11}*c^{11}*d^2 - 1792*a^5*b^{10}*c^{10}*d^3 + 5120*a^6*b^9*c^9*d^4 - 7168*a^7* \\
& b^8*c^8*d^5 + 3584*a^8*b^7*c^7*d^6 + 3584*a^9*b^6*c^6*d^7 - 7168*a^{10}*b^5* \\
& c^5*d^8 + 5120*a^{11}*b^4*c^4*d^9 - 1792*a^{12}*b^3*c^3*d^{10} + 256*a^{13}*b^2*c^2* \\
& d^{11}))/((512*(a^9*d^4 + a^5*b^4*c^4 - 4*a^6*b^3*c^3*d + 6*a^7*b^2*c^2*d^2 - \\
& 4*a^8*b*c*d^3)*(a^4*b^6*c^8 + a^{10}*c^2*d^6 - 6*a^5*b^5*c^7*d - 6*a^9*b*c^3*d^5 + \\
& 15*a^6*b^4*c^6*d^2 - 20*a^7*b^3*c^5*d^3 + 15*a^8*b^2*c^4*d^4)))*(-a^5*b^3)^{(1/2)}* \\
& (35*a^2*d^2 + 3*b^2*c^2 - 14*a*b*c*d))/(16*(a^9*d^4 + a^5*b^4*c^4 - 4*a^6*b^3*c^3*d + \\
& 6*a^7*b^2*c^2*d^2 - 4*a^8*b*c*d^3)))*(-a^5*b^3)^{(1/2)}*(35*a^2*d^2 + 3*b^2*c^2 - \\
& 14*a*b*c*d))/(16*(a^9*d^4 + a^5*b^4*c^4 - 4*a^6*b^3*c^3*d + 6*a^7*b^2*c^2*d^2 - \\
& 4*a^8*b*c*d^3)) + (((x*(16*a^6*b^3*d^9 + 9*b^9*c^6*d^3 - 84*a*b^8*c^5*d^4 - 224*a^5*b^4*c*d^8 + \\
& 406*a^2*b^7*c^4*d^5 - 980*a^3*b^6*c^3*d^6 + 2009*a^4*b^5*c^2*d^7)))/(32*(a^4*b^6*c^8 + a^{10}*c^2* \\
& d^6 - 6*a^5*b^5*c^7*d - 6*a^9*b*c^3*d^5 + 15*a^6*b^4*c^6*d^2 - 20*a^7*b^3*c^5*d^3 + \\
& 15*a^8*b^2*c^4*d^4)) + (((2*a^{13}*b^2*c*d^{13} - (3*a^2*b^{13}*c^{12}*d^2)/2 + (35*a^3*b^{12}*c^{11}*d^3)/2 - \\
& 98*a^4*b^{11}*c^{10}*d^4 + 336*a^5*b^{10}*c^9*d^5 - 765*a^6*b^9*c^8*d^6 + 1197*a^7*b^8*c^7*d^7 - \\
& 1302*a^8*b^7*c^6*d^8 + 978*a^9*b^6*c^5*d^9 - (987*a^{10}*b^5*c^4*d^{10})/2 + (315*a^{11}*b^4*c^3*d^{11})/2 - \\
& 28*a^{12}*b^3*c^2*d^{12})/(a^4*b^9*c^{11} - a^{13}*c^2*d^9 - 9*a^5*b^8*c^{10}*d + 9* \\
& a^{12}*b*c^3*d^8 + 36*a^6*b^7*c^9*d^2 - 84*a^7*b^6*c^8*d^3 + 126*a^8*b^5*c^7*d^4 - \\
& 126*a^9*b^4*c^6*d^5 + 84*a^{10}*b^3*c^5*d^6 - 36*a^{11}*b^2*c^4*d^7) + (x \\
& *(-a^5*b^3)^{(1/2)}*(35*a^2*d^2 + 3*b^2*c^2 - 14*a*b*c*d)*(256*a^4*b^{11}*c^{11}* \\
& d^2 - 1792*a^5*b^{10}*c^{10}*d^3 + 5120*a^6*b^9*c^9*d^4 - 7168*a^7*b^8*c^8*d^5 + \\
& 3584*a^8*b^7*c^7*d^6 + 3584*a^9*b^6*c^6*d^7 - 7168*a^{10}*b^5*c^5*d^8 + 5120* \\
& a^{11}*b^4*c^4*d^9 - 1792*a^{12}*b^3*c^3*d^{10} + 256*a^{13}*b^2*c^2*d^{11}))/((512* \\
& (a^9*d^4 + a^5*b^4*c^4 - 4*a^6*b^3*c^3*d + 6*a^7*b^2*c^2*d^2 - 4*a^8*b*c*d^3)*(a^4*b^6*c^8 + \\
& a^{10}*c^2*d^6 - 6*a^5*b^5*c^7*d - 6*a^9*b*c^3*d^5 + 15*a^6*b^4*c^6*d^2 - 20*a^7*b^3*c^5*d^3 + \\
& 15*a^8*b^2*c^4*d^4)))*(-a^5*b^3)^{(1/2)}*(35*a^2*d^2 + 3*b^2*c^2 - 14*a*b*c*d))/(16*(a^9*d^4 + \\
& a^5*b^4*c^4 - 4*a^6*b^3*c^3*d + 6*a^7*b^2*c^2*d^2 - 4*a^8*b*c*d^3)))*(-a^5*b^3)^{(1/2)}*(35*a^2*d^2 + \\
& 3*b^2*c^2 - 14*a*b*c*d)*1i)/(8*(a^9*d^4 + a^5*b^4*c^4 - 4*a^6*b^3*c^3*d + 6*a^7* \\
& b^2*c^2*d^2 - 4*a^8*b*c*d^3)) - (atan((((x*(16*a^6*b^3*d^9 + 9*b^9*c^6*d^3 - 84*a*b^8*c^5*d^4 - \\
& 224*a^5*b^4*c*d^8 + 406*a^2*b^7*c^4*d^5 - 980*a^3*b^6*c^3*d^6 + 2009*a^4*b^5*c^2*d^7)))/(32*(a^4*b^6*c^8 + \\
& a^{10}*c^2*d^6 - 6*a^5*b^5*c^7*d - 6*a^9*b*c^3*d^5 + 15*a^6*b^4*c^6*d^2 - 20*a^7*b^3*c^5*d^3 + 15*a^8* \\
& b^2*c^4*d^4)) - ((a*d - 7*b*c)*(-c^3*d^5)^{(1/2)}*((2*a^{13}*b^2*c*d^{13} - (3* \\
& a^2*b^{13}*c^{12}*d^2)/2 + (35*a^3*b^{12}*c^{11}*d^3)/2 - 98*a^4*b^{11}*c^{10}*d^4 + 3 \\
& 36*a^5*b^{10}*c^9*d^5 - 765*a^6*b^9*c^8*d^6 + 1197*a^7*b^8*c^7*d^7 - 1302*a^8* \\
& b^7*c^6*d^8 + 978*a^9*b^6*c^5*d^9 - (987*a^{10}*b^5*c^4*d^{10})/2 + (315*a^{11}*b^4*c^3*d^{11})/2 - \\
& 28*a^{12}*b^3*c^2*d^{12})/(a^4*b^9*c^{11} - a^{13}*c^2*d^9 - 9*a^5*b^8*c^{10}*d + 9*a^ \\
& 5*b^8*c^{10}*d + 9*a^{12}*b*c^3*d^8 + 36*a^6*b^7*c^9*d^2 - 84*a^7*b^6*c^8*d^3 + \\
& 126*a^8*b^5*c^7*d^4 - 126*a^9*b^4*c^6*d^5 + 84*a^{10}*b^3*c^5*d^6 - 36*a^{11}* \\
& b^2*c^4*d^7) - (x*(a*d - 7*b*c)*(-c^3*d^5)^{(1/2)}*(256*a^4*b^{11}*c^{11}*d^2 - 1 \\
& 792*a^5*b^{10}*c^{10}*d^3 + 5120*a^6*b^9*c^9*d^4 - 7168*a^7*b^8*c^8*d^5 + 3584* \\
& a^8*b^7*c^7*d^6 + 3584*a^9*b^6*c^6*d^7 - 7168*a^{10}*b^5*c^5*d^8 + 5120*a^{11}* \\
& b^4*c^4*d^9 - 1792*a^{12}*b^3*c^3*d^{10} + 256*a^{13}*b^2*c^2*d^{11}))/((128*(b^4*c^7 + \\
& a^4*c^3*d^4 - 4*a^3*b*c^4*d^3 + 6*a^2*b^2*c^5*d^2 - 4*a*b^3*c^6*d)*(a^4* \\
& b^6*c^8 + a^{10}*c^2*d^6 - 6*a^5*b^5*c^7*d - 6*a^9*b*c^3*d^5 + 15*a^6*b^4*c^6* \\
& d^2 - 20*a^7*b^3*c^5*d^3 + 15*a^8*b^2*c^4*d^4)))/(4*(b^4*c^7 + a^4*c^3*d^4 - 4*a^3*b*c^4*d^3 + \\
& 6*a^2*b^2*c^5*d^2 - 4*a*b^3*c^6*d)))*(a*d - 7*b*c)*(-c^3*d^5)^{(1/2)}*1i)/(4*(b^4*c^7 + \\
& a^4*c^3*d^4 - 4*a^3*b*c^4*d^3 + 6*a^2*b^2*c^5*d^2 - 4*a*b^3*c^6*d)
\end{aligned}$$


```

11)))/(128*(b^4*c^7 + a^4*c^3*d^4 - 4*a^3*b*c^4*d^3 + 6*a^2*b^2*c^5*d^2 - 4*
a*b^3*c^6*d)*(a^4*b^6*c^8 + a^10*c^2*d^6 - 6*a^5*b^5*c^7*d - 6*a^9*b*c^3*d^
5 + 15*a^6*b^4*c^6*d^2 - 20*a^7*b^3*c^5*d^3 + 15*a^8*b^2*c^4*d^4)))/(4*(b^
4*c^7 + a^4*c^3*d^4 - 4*a^3*b*c^4*d^3 + 6*a^2*b^2*c^5*d^2 - 4*a*b^3*c^6*d))
)*(a*d - 7*b*c)*(-c^3*d^5)^(1/2))/(4*(b^4*c^7 + a^4*c^3*d^4 - 4*a^3*b*c^4*d
^3 + 6*a^2*b^2*c^5*d^2 - 4*a*b^3*c^6*d)))*(a*d - 7*b*c)*(-c^3*d^5)^(1/2)*1
i)/(2*(b^4*c^7 + a^4*c^3*d^4 - 4*a^3*b*c^4*d^3 + 6*a^2*b^2*c^5*d^2 - 4*a*b^
3*c^6*d))

```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**3/(d*x**2+c)**2,x)

[Out] Timed out

$$3.42 \quad \int \frac{1}{(a+bx^2)^3 (c+dx^2)^3} dx$$

Optimal. Leaf size=315

$$\frac{3dx(ad+bc)(a^2d^2-6abcd+b^2c^2)}{8a^2c^2(c+dx^2)(bc-ad)^4} + \frac{dx(-2a^2d^2-13abcd+3b^2c^2)}{8a^2c(c+dx^2)^2(bc-ad)^3} - \frac{3d^{5/2}(a^2d^2-6abcd+21b^2c^2)\tan^{-1}\left(\frac{\sqrt{d}}{\sqrt{a}}\right)}{8c^{5/2}(bc-ad)^5}$$

[Out] 1/8*d*(-2*a^2*d^2-13*a*b*c*d+3*b^2*c^2)*x/a^2/c/(-a*d+b*c)^3/(d*x^2+c)^2+1/4*b*x/a/(-a*d+b*c)/(b*x^2+a)^2/(d*x^2+c)^2+1/8*b*(-11*a*d+3*b*c)*x/a^2/(-a*d+b*c)^2/(b*x^2+a)/(d*x^2+c)^2+3/8*d*(a*d+b*c)*(a^2*d^2-6*a*b*c*d+b^2*c^2)*x/a^2/c^2/(-a*d+b*c)^4/(d*x^2+c)+3/8*b^(5/2)*(21*a^2*d^2-6*a*b*c*d+b^2*c^2)*arctan(x*b^(1/2)/a^(1/2))/a^(5/2)/(-a*d+b*c)^5-3/8*d^(5/2)*(a^2*d^2-6*a*b*c*d+21*b^2*c^2)*arctan(x*d^(1/2)/c^(1/2))/c^(5/2)/(-a*d+b*c)^5

Rubi [A] time = 0.45, antiderivative size = 315, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {414, 527, 522, 205}

$$\frac{3dx(ad+bc)(a^2d^2-6abcd+b^2c^2)}{8a^2c^2(c+dx^2)(bc-ad)^4} + \frac{dx(-2a^2d^2-13abcd+3b^2c^2)}{8a^2c(c+dx^2)^2(bc-ad)^3} + \frac{3b^{5/2}(21a^2d^2-6abcd+b^2c^2)\tan^{-1}\left(\frac{\sqrt{b}}{\sqrt{a}}\right)}{8a^{5/2}(bc-ad)^5}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^3*(c + d*x^2)^3), x]

[Out] (d*(3*b^2*c^2 - 13*a*b*c*d - 2*a^2*d^2)*x)/(8*a^2*c*(b*c - a*d)^3*(c + d*x^2)^2) + (b*x)/(4*a*(b*c - a*d)*(a + b*x^2)^2*(c + d*x^2)^2) + (b*(3*b*c - 1*a*d)*x)/(8*a^2*(b*c - a*d)^2*(a + b*x^2)*(c + d*x^2)^2) + (3*d*(b*c + a*d)*(b^2*c^2 - 6*a*b*c*d + a^2*d^2)*x)/(8*a^2*c^2*(b*c - a*d)^4*(c + d*x^2)) + (3*b^(5/2)*(b^2*c^2 - 6*a*b*c*d + 21*a^2*d^2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]])/(8*a^(5/2)*(b*c - a*d)^5) - (3*d^(5/2)*(21*b^2*c^2 - 6*a*b*c*d + a^2*d^2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/(8*c^(5/2)*(b*c - a*d)^5)

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 522

Int[((e_) + (f_.)*(x_)^(n_))/(((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_))), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\int \frac{1}{(a + bx^2)^3 (c + dx^2)^3} dx = \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)^2} - \frac{\int \frac{-3bc+4ad-7bdx^2}{(a+bx^2)^2(c+dx^2)^3} dx}{4a(bc - ad)}$$

$$= \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)^2} + \frac{b(3bc - 11ad)x}{8a^2(bc - ad)^2 (a + bx^2) (c + dx^2)^2} + \int \frac{3b^2c^2-3c}{8a^2(bc - ad)^2}$$

$$= \frac{d(3b^2c^2 - 13abcd - 2a^2d^2)x}{8a^2c(bc - ad)^3 (c + dx^2)^2} + \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)^2} + \frac{b(3b}{8a^2(bc - ad)^2}$$

$$= \frac{d(3b^2c^2 - 13abcd - 2a^2d^2)x}{8a^2c(bc - ad)^3 (c + dx^2)^2} + \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)^2} + \frac{b(3b}{8a^2(bc - ad)^2}$$

$$= \frac{d(3b^2c^2 - 13abcd - 2a^2d^2)x}{8a^2c(bc - ad)^3 (c + dx^2)^2} + \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)^2} + \frac{b(3b}{8a^2(bc - ad)^2}$$

$$= \frac{d(3b^2c^2 - 13abcd - 2a^2d^2)x}{8a^2c(bc - ad)^3 (c + dx^2)^2} + \frac{bx}{4a(bc - ad)(a + bx^2)^2 (c + dx^2)^2} + \frac{b(3b}{8a^2(bc - ad)^2}$$

Mathematica [A] time = 0.93, size = 233, normalized size = 0.74

$$\frac{1}{8} \left(\frac{x(bc - ad) \left(\frac{3b^4c}{a^2(a+bx^2)} + \frac{b^3(-17ad+2bc-15bdx^2)}{a(a+bx^2)^2} - \frac{d^3(-2ad+17bc+15bdx^2)}{c(c+dx^2)^2} + \frac{3ad^4}{c^2(c+dx^2)} \right) - \frac{3d^{5/2}(a^2d^2-6abcd+21b^2c^2) \tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)}{c^{5/2}}}{(bc - ad)^5} \right)$$

Antiderivative was successfully verified.

```
[In] Integrate[1/((a + b*x^2)^3*(c + d*x^2)^3),x]
[Out] ((-3*b^(5/2)*(b^2*c^2 - 6*a*b*c*d + 21*a^2*d^2)*ArcTan[(Sqrt[b]*x)/Sqrt[a]]
)/(a^(5/2)*(-(b*c) + a*d)^5) + ((b*c - a*d)*x*((3*b^4*c)/(a^2*(a + b*x^2))
+ (3*a*d^4)/(c^2*(c + d*x^2)) + (b^3*(2*b*c - 17*a*d - 15*b*d*x^2))/(a*(a +
b*x^2)^2) - (d^3*(17*b*c - 2*a*d + 15*b*d*x^2))/(c*(c + d*x^2)^2)) - (3*d^(
5/2)*(21*b^2*c^2 - 6*a*b*c*d + a^2*d^2)*ArcTan[(Sqrt[d]*x)/Sqrt[c]])/c^(5/
2))/(b*c - a*d)^5)/8
```

fricas [B] time = 19.47, size = 5070, normalized size = 16.10

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^3/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [1/16*(6*(b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 6*a^3*b^3*c*d^5 - a^4*b^2*d^6)*x^7 + 2*(6*b^6*c^5*d - 31*a*b^5*c^4*d^2 - 9*a^2*b^4*c^3*d^3 + 9*a^3*b^3*c^2*d^4 + 31*a^4*b^2*c*d^5 - 6*a^5*b*d^6)*x^5 + 2*(3*b^6*c^6 - 8*a*b^5*c^5*d - 29*a^2*b^4*c^4*d^2 + 29*a^4*b^2*c^2*d^4 + 8*a^5*b*c*d^5 - 3*a^6*d^6)*x^3 - 3*(a^2*b^4*c^6 - 6*a^3*b^3*c^5*d + 21*a^4*b^2*c^4*d^2 + (b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 21*a^2*b^4*c^2*d^4)*x^8 + 2*(b^6*c^5*d - 5*a*b^5*c^4*d^2 + 15*a^2*b^4*c^3*d^3 + 21*a^3*b^3*c^2*d^4)*x^6 + (b^6*c^6 - 2*a*b^5*c^5*d - 2*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 + 21*a^4*b^2*c^2*d^4)*x^4 + 2*(a*b^5*c^6 - 5*a^2*b^4*c^5*d + 15*a^3*b^3*c^4*d^2 + 21*a^4*b^2*c^3*d^3)*x^2)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) - 3*(21*a^4*b^2*c^4*d^2 - 6*a^5*b*c^3*d^3 + a^6*c^2*d^4 + (21*a^2*b^4*c^2*d^4 - 6*a^3*b^3*c*d^5 + a^4*b^2*d^6)*x^8 + 2*(21*a^2*b^4*c^3*d^3 + 15*a^3*b^3*c^2*d^4 - 5*a^4*b^2*c*d^5 + a^5*b*d^6)*x^6 + (21*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 - 2*a^4*b^2*c^2*d^4 - 2*a^5*b*c*d^5 + a^6*d^6)*x^4 + 2*(21*a^3*b^3*c^4*d^2 + 15*a^4*b^2*c^3*d^3 - 5*a^5*b*c^2*d^4 + a^6*c*d^5)*x^2)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) + 2*(5*a*b^5*c^6 - 22*a^2*b^4*c^5*d + 17*a^3*b^3*c^4*d^2 - 17*a^4*b^2*c^3*d^3 + 22*a^5*b*c^2*d^4 - 5*a^6*c*d^5)*x)/(a^4*b^5*c^9 - 5*a^5*b^4*c^8*d + 10*a^6*b^3*c^7*d^2 - 10*a^7*b^2*c^6*d^3 + 5*a^8*b*c^5*d^4 - a^9*c^4*d^5 + (a^2*b^7*c^7*d^2 - 5*a^3*b^6*c^6*d^3 + 10*a^4*b^5*c^5*d^4 - 10*a^5*b^4*c^4*d^5 + 5*a^6*b^3*c^3*d^6 - a^7*b^2*c^2*d^7)*x^8 + 2*(a^2*b^7*c^8*d - 4*a^3*b^6*c^7*d^2 + 5*a^4*b^5*c^6*d^3 - 5*a^6*b^3*c^4*d^5 + 4*a^7*b^2*c^3*d^6 - a^8*b*c^2*d^7)*x^6 + (a^2*b^7*c^9 - a^3*b^6*c^8*d - 9*a^4*b^5*c^7*d^2 + 25*a^5*b^4*c^6*d^3 - 25*a^6*b^3*c^5*d^4 + 9*a^7*b^2*c^4*d^5 + a^8*b*c^3*d^6 - a^9*c^2*d^7)*x^4 + 2*(a^3*b^6*c^9 - 4*a^4*b^5*c^8*d + 5*a^5*b^4*c^7*d^2 - 5*a^7*b^2*c^5*d^4 + 4*a^8*b*c^4*d^5 - a^9*c^3*d^6)*x^2), 1/16*(6*(b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 6*a^3*b^3*c*d^5 - a^4*b^2*d^6)*x^7 + 2*(6*b^6*c^5*d - 31*a*b^5*c^4*d^2 - 9*a^2*b^4*c^3*d^3 + 9*a^3*b^3*c^2*d^4 + 31*a^4*b^2*c*d^5 - 6*a^5*b*d^6)*x^5 + 2*(3*b^6*c^6 - 8*a*b^5*c^5*d - 29*a^2*b^4*c^4*d^2 + 29*a^4*b^2*c^2*d^4 + 8*a^5*b*c*d^5 - 3*a^6*d^6)*x^3 - 6*(21*a^4*b^2*c^4*d^2 - 6*a^5*b*c^3*d^3 + a^6*c^2*d^4 + (21*a^2*b^4*c^2*d^4 - 6*a^3*b^3*c*d^5 + a^4*b^2*d^6)*x^8 + 2*(21*a^2*b^4*c^3*d^3 + 15*a^3*b^3*c^2*d^4 - 5*a^4*b^2*c*d^5 + a^5*b*d^6)*x^6 + (21*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 - 2*a^4*b^2*c^2*d^4 - 2*a^5*b*c*d^5 + a^6*d^6)*x^4 + 2*(21*a^3*b^3*c^4*d^2 + 15*a^4*b^2*c^3*d^3 - 5*a^5*b*c^2*d^4 + a^6*c*d^5)*x^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) - 3*(a^2*b^4*c^6 - 6*a^3*b^3*c^5*d + 21*a^4*b^2*c^4*d^2 + (b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 21*a^2*b^4*c^2*d^4)*x^8 + 2*(b^6*c^5*d - 5*a*b^5*c^4*d^2 + 15*a^2*b^4*c^3*d^3 + 21*a^3*b^3*c^2*d^4)*x^6 + (b^6*c^6 - 2*a*b^5*c^5*d - 2*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 + 21*a^4*b^2*c^2*d^4)*x^4 + 2*(a*b^5*c^6 - 5*a^2*b^4*c^5*d + 15*a^3*b^3*c^4*d^2 + 21*a^4*b^2*c^3*d^3)*x^2)*sqrt(-b/a)*log((b*x^2 - 2*a*x*sqrt(-b/a) - a)/(b*x^2 + a)) + 2*(5*a*b^5*c^6 - 22*a^2*b^4*c^5*d + 17*a^3*b^3*c^4*d^2 - 17*a^4*b^2*c^3*d^3 + 22*a^5*b*c^2*d^4 - 5*a^6*c*d^5)*x)/(a^4*b^5*c^9 - 5*a^5*b^4*c^8*d + 10*a^6*b^3*c^7*d^2 - 10*a^7*b^2*c^6*d^3 + 5*a^8*b*c^5*d^4 - a^9*c^4*d^5 + (a^2*b^7*c^7*d^2 - 5*a^3*b^6*c^6*d^3 + 10*a^4*b^5*c^5*d^4 - 10*a^5*b^4*c^4*d^5 + 5*a^6*b^3*c^3*d^6 - a^7*b^2*c^2*d^7)*x^8 + 2*(a^2*b^7*c^8*d - 4*a^3*b^6*c^7*d^2 + 5*a^4*b^5*c^6*d^3 - 5*a^6*b^3*c^4*d^5 + 4*a^7*b^2*c^3*d^6 - a^8*b*c^2*d^7)*x^6 + (a^2*b^7*c^9 - a^3*b^6*c^8*d - 9*a^4*b^5*c^7*d^2 + 25*a^5*b^4*c^6*d^3 - 25*a^6*b^3*c^5*d^4 + 9*a^7*b^2*c^4*d^5 + a^8*b*c^3*d^6 - a^9*c^2*d^7)*x^4 + 2*(a^3*b^6*c^9 - 4*a^4*b^5*c^8*d + 5*a^5*b^4*c^7*d^2 - 5*a^7*b^2*c^5*d^4 + 4*a^8*b*c^4*d^5 - a^9*c^3*d^6)*x^2), 1/16*(6*(b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 6*a^3*b^3*c*d^5 - a^4*b^2*d^6)*x^7 + 2*(6*b^6*c^5*d - 31*a*b^5*c^4*d^2 - 9*a^2*b^4*c^3*d^3 + 9*a^3*b^3*c^2*d^4 + 31*a^4*b^2*c*d^5 - 6*a^5*b*d^6)*x^5 + 2*(3*b^6*c^6 - 8*a*b^5*c^5*d - 29*a^2*b^4*c^4*d^2 + 29*a^4*b^2*c^2*d^4 + 8*a^5*b*c*d^5 - 3*a^6*d^6)*x^3 + 6*(a^2*b^4*c^6 - 6*a^3*b^3*c^5*d + 21*a^4*b^2*c^4*d^2 + (b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 21*a^2*b^4*c^2*d^4)*x^8 + 2*(b^6*c^5*d - 5*a*b^5*c^4*d^2 + 15*a^2*b^4*c^3*d^3 + 21*a^3*b^3*c^2*d^4)*x^6 + (b^6*c^6 - 2*a*b^5*c^5*d - 2*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 + 21*a^4*b^2*c^2*d^4)*x^4 + 2*(a*b^5*c^6 - 5*a^2*b^4*c^5*d + 15*a^3*b^3*c^4*d^2 + 21*a^4*b^2*c^3*d^3 + 21*a^5*b*c^2*d^4 + a^6*c*d^5)*x^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) - 3*(a^2*b^4*c^6 - 6*a^3*b^3*c^5*d + 21*a^4*b^2*c^4*d^2 + (b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 21*a^2*b^4*c^2*d^4)*x^8 + 2*(b^6*c^5*d - 5*a*b^5*c^4*d^2 + 15*a^2*b^4*c^3*d^3 + 21*a^3*b^3*c^2*d^4)*x^6 + (b^6*c^6 - 2*a*b^5*c^5*d - 2*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 + 21*a^4*b^2*c^2*d^4)*x^4 + 2*(a*b^5*c^6 - 5*a^2*b^4*c^5*d

+ 15*a^3*b^3*c^4*d^2 + 21*a^4*b^2*c^3*d^3)*x^2)*sqrt(b/a)*arctan(x*sqrt(b/a)) - 3*(21*a^4*b^2*c^4*d^2 - 6*a^5*b*c^3*d^3 + a^6*c^2*d^4 + (21*a^2*b^4*c^2*d^4 - 6*a^3*b^3*c*d^5 + a^4*b^2*d^6)*x^8 + 2*(21*a^2*b^4*c^3*d^3 + 15*a^3*b^3*c^2*d^4 - 5*a^4*b^2*c*d^5 + a^5*b*d^6)*x^6 + (21*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 - 2*a^4*b^2*c^2*d^4 - 2*a^5*b*c*d^5 + a^6*d^6)*x^4 + 2*(21*a^3*b^3*c^4*d^2 + 15*a^4*b^2*c^3*d^3 - 5*a^5*b*c^2*d^4 + a^6*c*d^5)*x^2)*sqrt(-d/c)*log((d*x^2 + 2*c*x*sqrt(-d/c) - c)/(d*x^2 + c)) + 2*(5*a*b^5*c^6 - 22*a^2*b^4*c^5*d + 17*a^3*b^3*c^4*d^2 - 17*a^4*b^2*c^3*d^3 + 22*a^5*b*c^2*d^4 - 5*a^6*c*d^5)*x)/(a^4*b^5*c^9 - 5*a^5*b^4*c^8*d + 10*a^6*b^3*c^7*d^2 - 10*a^7*b^2*c^6*d^3 + 5*a^8*b*c^5*d^4 - a^9*c^4*d^5 + (a^2*b^7*c^7*d^2 - 5*a^3*b^6*c^6*d^3 + 10*a^4*b^5*c^5*d^4 - 10*a^5*b^4*c^4*d^5 + 5*a^6*b^3*c^3*d^6 - a^7*b^2*c^2*d^7)*x^8 + 2*(a^2*b^7*c^8*d - 4*a^3*b^6*c^7*d^2 + 5*a^4*b^5*c^6*d^3 - 5*a^6*b^3*c^4*d^5 + 4*a^7*b^2*c^3*d^6 - a^8*b*c^2*d^7)*x^6 + (a^2*b^7*c^9 - a^3*b^6*c^8*d - 9*a^4*b^5*c^7*d^2 + 25*a^5*b^4*c^6*d^3 - 25*a^6*b^3*c^5*d^4 + 9*a^7*b^2*c^4*d^5 + a^8*b*c^3*d^6 - a^9*c^2*d^7)*x^4 + 2*(a^3*b^6*c^9 - 4*a^4*b^5*c^8*d + 5*a^5*b^4*c^7*d^2 - 5*a^7*b^2*c^5*d^4 + 4*a^8*b*c^4*d^5 - a^9*c^3*d^6)*x^2), 1/8*(3*(b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 6*a^3*b^3*c*d^5 - a^4*b^2*d^6)*x^7 + (6*b^6*c^5*d - 31*a*b^5*c^4*d^2 - 9*a^2*b^4*c^3*d^3 + 9*a^3*b^3*c^2*d^4 + 31*a^4*b^2*c*d^5 - 6*a^5*b*d^6)*x^5 + (3*b^6*c^6 - 8*a*b^5*c^5*d - 29*a^2*b^4*c^4*d^2 + 29*a^4*b^2*c^2*d^4 + 8*a^5*b*c*d^5 - 3*a^6*d^6)*x^3 + 3*(a^2*b^4*c^6 - 6*a^3*b^3*c^5*d + 21*a^4*b^2*c^4*d^2 + (b^6*c^4*d^2 - 6*a*b^5*c^3*d^3 + 21*a^2*b^4*c^2*d^4)*x^8 + 2*(b^6*c^5*d - 5*a*b^5*c^4*d^2 + 15*a^2*b^4*c^3*d^3 + 21*a^3*b^3*c^2*d^4)*x^6 + (b^6*c^6 - 2*a*b^5*c^5*d - 2*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 + 21*a^4*b^2*c^2*d^4)*x^4 + 2*(a*b^5*c^6 - 5*a^2*b^4*c^5*d + 15*a^3*b^3*c^4*d^2 + 21*a^4*b^2*c^3*d^3)*x^2)*sqrt(b/a)*arctan(x*sqrt(b/a)) - 3*(21*a^4*b^2*c^4*d^2 - 6*a^5*b*c^3*d^3 + a^6*c^2*d^4 + (21*a^2*b^4*c^2*d^4 - 6*a^3*b^3*c*d^5 + a^4*b^2*d^6)*x^8 + 2*(21*a^2*b^4*c^3*d^3 + 15*a^3*b^3*c^2*d^4 - 5*a^4*b^2*c*d^5 + a^5*b*d^6)*x^6 + (21*a^2*b^4*c^4*d^2 + 78*a^3*b^3*c^3*d^3 - 2*a^4*b^2*c^2*d^4 - 2*a^5*b*c*d^5 + a^6*d^6)*x^4 + 2*(21*a^3*b^3*c^4*d^2 + 15*a^4*b^2*c^3*d^3 - 5*a^5*b*c^2*d^4 + a^6*c*d^5)*x^2)*sqrt(d/c)*arctan(x*sqrt(d/c)) + (5*a*b^5*c^6 - 22*a^2*b^4*c^5*d + 17*a^3*b^3*c^4*d^2 - 17*a^4*b^2*c^3*d^3 + 22*a^5*b*c^2*d^4 - 5*a^6*c*d^5)*x)/(a^4*b^5*c^9 - 5*a^5*b^4*c^8*d + 10*a^6*b^3*c^7*d^2 - 10*a^7*b^2*c^6*d^3 + 5*a^8*b*c^5*d^4 - a^9*c^4*d^5 + (a^2*b^7*c^7*d^2 - 5*a^3*b^6*c^6*d^3 + 10*a^4*b^5*c^5*d^4 - 10*a^5*b^4*c^4*d^5 + 5*a^6*b^3*c^3*d^6 - a^7*b^2*c^2*d^7)*x^8 + 2*(a^2*b^7*c^8*d - 4*a^3*b^6*c^7*d^2 + 5*a^4*b^5*c^6*d^3 - 5*a^6*b^3*c^4*d^5 + 4*a^7*b^2*c^3*d^6 - a^8*b*c^2*d^7)*x^6 + (a^2*b^7*c^9 - a^3*b^6*c^8*d - 9*a^4*b^5*c^7*d^2 + 25*a^5*b^4*c^6*d^3 - 25*a^6*b^3*c^5*d^4 + 9*a^7*b^2*c^4*d^5 + a^8*b*c^3*d^6 - a^9*c^2*d^7)*x^4 + 2*(a^3*b^6*c^9 - 4*a^4*b^5*c^8*d + 5*a^5*b^4*c^7*d^2 - 5*a^7*b^2*c^5*d^4 + 4*a^8*b*c^4*d^5 - a^9*c^3*d^6)*x^2)]

giac [A] time = 0.62, size = 574, normalized size = 1.82

$$\frac{3(b^5c^2 - 6ab^4cd + 21a^2b^3d^2) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8(a^2b^5c^5 - 5a^3b^4c^4d + 10a^4b^3c^3d^2 - 10a^5b^2c^2d^3 + 5a^6bcd^4 - a^7d^5)\sqrt{ab}} - \frac{3(21b^2c^2d^3 - 6abcd^4 - 10a^2b^3c^5d^2 - 10a^3b^2c^4d^3 + 5a^4b^3c^3d^4 - 5a^5b^2c^2d^5 + 5a^6bcd^4 - a^7d^5)\sqrt{ab}}{8(b^5c^7 - 5ab^4c^6d + 10a^2b^3c^5d^2 - 10a^3b^2c^4d^3 + 5a^4b^3c^3d^4 - a^5c^2d^5)\sqrt{cd}} + \frac{1}{8} \left(3b^5c^3d^2x^7 - 15a^2b^4c^2d^3x^7 - 15a^2b^3c^2d^4x^7 + 3a^3b^2d^5x^7 + 6b^5c^4d^4x^5 - 25a^2b^4c^3d^2x^5 - 34a^2b^3c^2d^3x^5 - 25a^3b^2c^2d^4x^5 + 6a^4b^3d^5x^5 + 3b^5c^5x^3 - 5a^2b^4c^4d^4x^3 - 34a^3b^3c^4d^4x^3 + 3a^4b^2c^4d^4x^3 - 34a^5b^2c^4d^4x^3 + 3a^6bcd^4x^3 - 34a^7d^5x^3 \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^3/(d*x^2+c)^3,x, algorithm="giac")

[Out] 3/8*(b^5*c^2 - 6*a*b^4*c*d + 21*a^2*b^3*d^2)*arctan(b*x/sqrt(a*b))/((a^2*b^5*c^5 - 5*a^3*b^4*c^4*d + 10*a^4*b^3*c^3*d^2 - 10*a^5*b^2*c^2*d^3 + 5*a^6*b*c*d^4 - a^7*d^5)*sqrt(a*b)) - 3/8*(21*b^2*c^2*d^3 - 6*a*b*c*d^4 + a^2*d^5)*arctan(d*x/sqrt(c*d))/((b^5*c^7 - 5*a*b^4*c^6*d + 10*a^2*b^3*c^5*d^2 - 10*a^3*b^2*c^4*d^3 + 5*a^4*b^3*c^3*d^4 - a^5*c^2*d^5)*sqrt(c*d)) + 1/8*(3*b^5*c^3*d^2*x^7 - 15*a^2*b^4*c^2*d^3*x^7 - 15*a^2*b^3*c^2*d^4*x^7 + 3*a^3*b^2*d^5*x^7 + 6*b^5*c^4*d^4*x^5 - 25*a^2*b^4*c^3*d^2*x^5 - 34*a^2*b^3*c^2*d^3*x^5 - 25*a^3*b^2*c^2*d^4*x^5 + 6*a^4*b^3*d^5*x^5 + 3*b^5*c^5*x^3 - 5*a^2*b^4*c^4*d^4*x^3 - 34*a^3*b^3*c^4*d^4*x^3 + 3*a^4*b^2*c^4*d^4*x^3 - 34*a^5*b^2*c^4*d^4*x^3 + 3*a^6*b*c*d^4*x^3 - 34*a^7*d^5*x^3)

$$\frac{2b^3c^3d^2x^3 - 34a^3b^2c^2d^3x^3 - 5a^4b^2cd^4x^3 + 3a^5d^5x^3 + 5a^4b^2c^4d^5x - 17a^2b^3c^4d^5x - 17a^4b^2c^2d^3x + 5a^5c^4d^4x}{(a^2b^4c^6 - 4a^3b^3c^5d + 6a^4b^2c^4d^2 - 4a^5b^2c^3d^3 + a^6c^2d^4)(b^2d^2x^2 + b^2cd^2x + a^2d^2x^2 + a^2c^2d^2)}$$

maple [A] time = 0.02, size = 568, normalized size = 1.80

$$\frac{3a^2d^6x^3}{8(ad-bc)^5(dx^2+c)^2c^2} - \frac{9abd^5x^3}{4(ad-bc)^5(dx^2+c)^2c} + \frac{9b^5cdx^3}{4(ad-bc)^5(bx^2+a)^2a} - \frac{3b^6c^2x^3}{8(ad-bc)^5(bx^2+a)^2a^2} - \dots$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^3/(d*x^2+c)^3,x)

[Out] $\frac{3}{8}d^6/(ad-bc)^5/(d^2x^2+c)^2/c^2x^3a^2-9/4d^5/(ad-bc)^5/(d^2x^2+c)^2/c^2x^3ab+15/8d^4/(ad-bc)^5/(d^2x^2+c)^2x^3b^2+5/8d^5/(ad-bc)^5/(d^2x^2+c)^2/c^2xa^2-11/4d^4/(ad-bc)^5/(d^2x^2+c)^2x^3ab+17/8d^3/(ad-bc)^5/(d^2x^2+c)^2c^2xb^2+3/8d^5/(ad-bc)^5/c^2/(cd)^{(1/2)}\arctan(1/(cd)^{(1/2)}dx) * a^2-9/4d^4/(ad-bc)^5/c/(cd)^{(1/2)}\arctan(1/(cd)^{(1/2)}dx) * ab+63/8d^3/(ad-bc)^5/(cd)^{(1/2)}\arctan(1/(cd)^{(1/2)}dx) * b^2-15/8b^4/(ad-bc)^5/(b^2x^2+a)^2x^3d^2+9/4b^5/(ad-bc)^5/(b^2x^2+a)^2/a^2x^3cd-3/8b^6/(ad-bc)^5/(b^2x^2+a)^2/a^2x^3c^2-17/8b^3/(ad-bc)^5/(b^2x^2+a)^2x^3ad^2+11/4b^4/(ad-bc)^5/(b^2x^2+a)^2x^3cd-5/8b^5/(ad-bc)^5/(b^2x^2+a)^2x^3/a^2c^2-63/8b^3/(ad-bc)^5/(ab)^{(1/2)}\arctan(1/(ab)^{(1/2)}bx) * d^2+9/4b^4/(ad-bc)^5/a/(ab)^{(1/2)}\arctan(1/(ab)^{(1/2)}bx) * cd-3/8b^5/(ad-bc)^5/a^2/(ab)^{(1/2)}\arctan(1/(ab)^{(1/2)}bx) * c^2$

maxima [B] time = 3.39, size = 820, normalized size = 2.60

$$\frac{3(b^5c^2 - 6ab^4cd + 21a^2b^3d^2) \arctan\left(\frac{bx}{\sqrt{ab}}\right)}{8(a^2b^5c^5 - 5a^3b^4c^4d + 10a^4b^3c^3d^2 - 10a^5b^2c^2d^3 + 5a^6bcd^4 - a^7d^5)\sqrt{ab}} - \frac{3(21b^2c^2d^3 - 6abcd)}{8(b^5c^7 - 5ab^4c^6d + 10a^2b^3c^5d^2 - 10a^3b^2c^4d^3 + 5a^4b^2c^3d^4 - a^5c^2d^5)\sqrt{cd}} + \frac{1}{8} \dots$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^3/(d*x^2+c)^3,x, algorithm="maxima")

[Out] $\frac{3}{8}(b^5c^2 - 6a^4b^4cd + 21a^2b^3d^2) \arctan(bx/\sqrt{ab}) / ((a^2b^5c^5 - 5a^3b^4c^4d + 10a^4b^3c^3d^2 - 10a^5b^2c^2d^3 + 5a^6bcd^4 - a^7d^5)\sqrt{ab}) - \frac{3}{8}(21b^2c^2d^3 - 6abcd + a^2d^5) \arctan(dx/\sqrt{cd}) / ((b^5c^7 - 5ab^4c^6d + 10a^2b^3c^5d^2 - 10a^3b^2c^4d^3 + 5a^4b^2c^3d^4 - a^5c^2d^5)\sqrt{cd}) + \frac{1}{8} \dots$

mupad [B] time = 8.55, size = 11150, normalized size = 35.40

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^3*(c + d*x^2)^3),x)

[Out] ((x*(5*a^4*d^4 + 5*b^4*c^4 - 17*a*b^3*c^3*d - 17*a^3*b*c*d^3))/(8*a*c*(a^4*d^4 + b^4*c^4 + 6*a^2*b^2*c^2*d^2 - 4*a*b^3*c^3*d - 4*a^3*b*c*d^3)) - (x^3*(34*a^2*b^3*c^3*d^2 - 3*b^5*c^5 - 3*a^5*d^5 + 34*a^3*b^2*c^2*d^3 + 5*a*b^4*c^4*d + 5*a^4*b*c*d^4))/(8*a^2*c^2*(a^4*d^4 + b^4*c^4 + 6*a^2*b^2*c^2*d^2 - 4*a*b^3*c^3*d - 4*a^3*b*c*d^3)) - (x^5*(25*a*b^4*c^3*d^2 - 6*b^5*c^4*d - 6*a^4*b*d^5 + 25*a^3*b^2*c*d^4 + 34*a^2*b^3*c^2*d^3))/(8*a^2*c^2*(a^4*d^4 + b^4*c^4 + 6*a^2*b^2*c^2*d^2 - 4*a*b^3*c^3*d - 4*a^3*b*c*d^3)) + (3*b*d*x^7*(a^3*b*d^4 + b^4*c^3*d - 5*a*b^3*c^2*d^2 - 5*a^2*b^2*c*d^3))/(8*a^2*c^2*(a^4*d^4 + b^4*c^4 + 6*a^2*b^2*c^2*d^2 - 4*a*b^3*c^3*d - 4*a^3*b*c*d^3)))/(x^4*(a^2*d^2 + b^2*c^2 + 4*a*b*c*d) + x^2*(2*a*b*c^2 + 2*a^2*c*d) + x^6*(2*a*b*d^2 + 2*b^2*c*d) + a^2*c^2 + b^2*d^2*x^8) - (atan((((x*(9*a^8*b^3*d^11 + 9*b^11*c^8*d^3 - 108*a*b^10*c^7*d^4 - 108*a^7*b^4*c*d^10 + 702*a^2*b^9*c^6*d^5 - 2268*a^3*b^8*c^5*d^6 + 7938*a^4*b^7*c^4*d^7 - 2268*a^5*b^6*c^3*d^8 + 702*a^6*b^5*c^2*d^9))/(32*(a^4*b^8*c^12 + a^12*c^4*d^8 - 8*a^5*b^7*c^11*d - 8*a^11*b*c^5*d^7 + 28*a^6*b^6*c^10*d^2 - 56*a^7*b^5*c^9*d^3 + 70*a^8*b^4*c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^10*b^2*c^6*d^6)) - (3*(((3*a^2*b^16*c^16*d^2)/2 - (45*a^3*b^15*c^15*d^3)/2 + (333*a^4*b^14*c^14*d^4)/2 - 765*a^5*b^13*c^13*d^5 + (4743*a^6*b^12*c^12*d^6)/2 - (10371*a^7*b^11*c^11*d^7)/2 + (16425*a^8*b^10*c^10*d^8)/2 - 9558*a^9*b^9*c^9*d^9 + (16425*a^10*b^8*c^8*d^10)/2 - (10371*a^11*b^7*c^7*d^11)/2 + (4743*a^12*b^6*c^6*d^12)/2 - 765*a^13*b^5*c^5*d^13 + (333*a^14*b^4*c^4*d^14)/2 - (45*a^15*b^3*c^3*d^15)/2 + (3*a^16*b^2*c^2*d^16)/2)/(a^4*b^12*c^16 + a^16*c^4*d^12 - 12*a^5*b^11*c^15*d - 12*a^15*b*c^5*d^11 + 66*a^6*b^10*c^14*d^2 - 220*a^7*b^9*c^13*d^3 + 495*a^8*b^8*c^12*d^4 - 792*a^9*b^7*c^11*d^5 + 924*a^10*b^6*c^10*d^6 - 792*a^11*b^5*c^9*d^7 + 495*a^12*b^4*c^8*d^8 - 220*a^13*b^3*c^7*d^9 + 66*a^14*b^2*c^6*d^10) - (3*x*(-a^5*b^5)^(1/2)*(21*a^2*d^2 + b^2*c^2 - 6*a*b*c*d)*(256*a^4*b^13*c^15*d^2 - 2304*a^5*b^12*c^14*d^3 + 8960*a^6*b^11*c^13*d^4 - 19200*a^7*b^10*c^12*d^5 + 23040*a^8*b^9*c^11*d^6 - 10752*a^9*b^8*c^10*d^7 - 10752*a^10*b^7*c^9*d^8 + 23040*a^11*b^6*c^8*d^9 - 19200*a^12*b^5*c^7*d^10 + 8960*a^13*b^4*c^6*d^11 - 2304*a^14*b^3*c^5*d^12 + 256*a^15*b^2*c^4*d^13)))/(512*(a^10*d^5 - a^5*b^5*c^5 + 5*a^6*b^4*c^4*d - 10*a^7*b^3*c^3*d^2 + 10*a^8*b^2*c^2*d^3 - 5*a^9*b*c*d^4)*(a^4*b^8*c^12 + a^12*c^4*d^8 - 8*a^5*b^7*c^11*d - 8*a^11*b*c^5*d^7 + 28*a^6*b^6*c^10*d^2 - 56*a^7*b^5*c^9*d^3 + 70*a^8*b^4*c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^10*b^2*c^6*d^6)))*(-a^5*b^5)^(1/2)*(21*a^2*d^2 + b^2*c^2 - 6*a*b*c*d))/(16*(a^10*d^5 - a^5*b^5*c^5 + 5*a^6*b^4*c^4*d - 10*a^7*b^3*c^3*d^2 + 10*a^8*b^2*c^2*d^3 - 5*a^9*b*c*d^4)))*(-a^5*b^5)^(1/2)*(21*a^2*d^2 + b^2*c^2 - 6*a*b*c*d)*3i)/(16*(a^10*d^5 - a^5*b^5*c^5 + 5*a^6*b^4*c^4*d - 10*a^7*b^3*c^3*d^2 + 10*a^8*b^2*c^2*d^3 - 5*a^9*b*c*d^4)) + (((x*(9*a^8*b^3*d^11 + 9*b^11*c^8*d^3 - 108*a*b^10*c^7*d^4 - 108*a^7*b^4*c*d^10 + 702*a^2*b^9*c^6*d^5 - 2268*a^3*b^8*c^5*d^6 + 7938*a^4*b^7*c^4*d^7 - 2268*a^5*b^6*c^3*d^8 + 702*a^6*b^5*c^2*d^9))/(32*(a^4*b^8*c^12 + a^12*c^4*d^8 - 8*a^5*b^7*c^11*d - 8*a^11*b*c^5*d^7 + 28*a^6*b^6*c^10*d^2 - 56*a^7*b^5*c^9*d^3 + 70*a^8*b^4*c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^10*b^2*c^6*d^6)) + (3*(((3*a^2*b^16*c^16*d^2)/2 - (45*a^3*b^15*c^15*d^3)/2 + (333*a^4*b^14*c^14*d^4)/2 - 765*a^5*b^13*c^13*d^5 + (4743*a^6*b^12*c^12*d^6)/2 - (10371*a^7*b^11*c^11*d^7)/2 + (16425*a^8*b^10*c^10*d^8)/2 - 9558*a^9*b^9*c^9*d^9 + (16425*a^10*b^8*c^8*d^10)/2 - (10371*a^11*b^7*c^7*d^11)/2 + (4743*a^12*b^6*c^6*d^12)/2 - 765*a^13*b^5*c^5*d^13 + (333*a^14*b^4*c^4*d^14)/2 - (45*a^15*b^3*c^3*d^15)/2 + (3*a^16*b^2*c^2*d^16)/2)/(a^4*b^12*c^16 + a^16*c^4*d^12 - 12*a^5*b^11*c^15*d - 12*a^15*b*c^5*d^11 + 66*a^6*b^10*c^14*d^2 - 220*a^7*b^9*c^13*d^3 + 495*a^8*b^8*c^12*d^4 - 792*a^9*b^7*c^11*d^5 + 924*a^10*b^6*c^10*d^6 - 792*a^11*b^5*c^9*d^7 + 495*a^12*b^4*c^8*d^8 - 220*a^13*b^3*c^7*d^9 + 66*a^14*b^2*c^6*d^10) + (3*x*(-a^5*b^5)^(1/2)*(21*a^2*d^2 + b^2*c^2 - 6*a*b*c*d)*(256*a^4*b^13*c^15*d^2 - 2304*a^5*b^12*c^14*d^3 + 8960*a^6*b^11*c^13*d^4 - 19200*a^7*b^10*c^12*d^5 + 23040*a^8*b^9*c^11*d^6 - 10752*a^9*b^8*c^10*d^7 - 10752*a^10*b^7*c^9*d^8 + 23040*a^11*b^6*c^8*d^9 - 19200*a^12*b^5*c^7*d^10 + 8960*a^13*b^4*c^6*d^11 - 2304*a^14*b^3*c^5*d^12 + 256*a^15*b^2*c^4*d^13)))/(512*(a^10*d^5 - a^5*b^5*c^5 + 5*a^6*b^4*c^4*d - 10*a^7*b^3*c^3*d^2 + 10*

$$\begin{aligned}
& 00*a^{12}*b^5*c^7*d^{10} + 8960*a^{13}*b^4*c^6*d^{11} - 2304*a^{14}*b^3*c^5*d^{12} + 25 \\
& 6*a^{15}*b^2*c^4*d^{13}))/((512*(a^{10}*d^5 - a^5*b^5*c^5 + 5*a^6*b^4*c^4*d - 10*a \\
& ^7*b^3*c^3*d^2 + 10*a^8*b^2*c^2*d^3 - 5*a^9*b*c*d^4)*(a^4*b^8*c^{12} + a^{12}*c \\
& ^4*d^8 - 8*a^5*b^7*c^{11}*d - 8*a^{11}*b*c^5*d^7 + 28*a^6*b^6*c^{10}*d^2 - 56*a^7 \\
& *b^5*c^9*d^3 + 70*a^8*b^4*c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^{10}*b^2*c^6*d^6 \\
&))*(-a^5*b^5)^{(1/2)}*(21*a^2*d^2 + b^2*c^2 - 6*a*b*c*d))/(16*(a^{10}*d^5 - a \\
& ^5*b^5*c^5 + 5*a^6*b^4*c^4*d - 10*a^7*b^3*c^3*d^2 + 10*a^8*b^2*c^2*d^3 - 5* \\
& a^9*b*c*d^4)))*(-a^5*b^5)^{(1/2)}*(21*a^2*d^2 + b^2*c^2 - 6*a*b*c*d))/(16*(a^ \\
& 10*d^5 - a^5*b^5*c^5 + 5*a^6*b^4*c^4*d - 10*a^7*b^3*c^3*d^2 + 10*a^8*b^2*c^ \\
& 2*d^3 - 5*a^9*b*c*d^4)))*(-a^5*b^5)^{(1/2)}*(21*a^2*d^2 + b^2*c^2 - 6*a*b*c* \\
& d)*3i)/(8*(a^{10}*d^5 - a^5*b^5*c^5 + 5*a^6*b^4*c^4*d - 10*a^7*b^3*c^3*d^2 + \\
& 10*a^8*b^2*c^2*d^3 - 5*a^9*b*c*d^4)) - (\operatorname{atan}(\frac{(x*(9*a^8*b^3*d^{11} + 9*b^{11} \\
& *c^8*d^3 - 108*a*b^{10}*c^7*d^4 - 108*a^7*b^4*c*d^{10} + 702*a^2*b^9*c^6*d^5 - \\
& 2268*a^3*b^8*c^5*d^6 + 7938*a^4*b^7*c^4*d^7 - 2268*a^5*b^6*c^3*d^8 + 702*a^6 \\
& *b^5*c^2*d^9))/(32*(a^4*b^8*c^{12} + a^{12}*c^4*d^8 - 8*a^5*b^7*c^{11}*d - 8*a^1 \\
& 1*b*c^5*d^7 + 28*a^6*b^6*c^{10}*d^2 - 56*a^7*b^5*c^9*d^3 + 70*a^8*b^4*c^8*d^4 \\
& - 56*a^9*b^3*c^7*d^5 + 28*a^{10}*b^2*c^6*d^6)) - (3*((3*a^2*b^{16}*c^{16}*d^2)/ \\
& 2 - (45*a^3*b^{15}*c^{15}*d^3)/2 + (333*a^4*b^{14}*c^{14}*d^4)/2 - 765*a^5*b^{13}*c^{1 \\
& 3*d^5 + (4743*a^6*b^{12}*c^{12}*d^6)/2 - (10371*a^7*b^{11}*c^{11}*d^7)/2 + (16425*a^ \\
& 8*b^{10}*c^{10}*d^8)/2 - 9558*a^9*b^9*c^9*d^9 + (16425*a^{10}*b^8*c^8*d^{10})/2 - \\
& (10371*a^{11}*b^7*c^7*d^{11})/2 + (4743*a^{12}*b^6*c^6*d^{12})/2 - 765*a^{13}*b^5*c^5 \\
& *d^{13} + (333*a^{14}*b^4*c^4*d^{14})/2 - (45*a^{15}*b^3*c^3*d^{15})/2 + (3*a^{16}*b^2* \\
& c^2*d^{16})/2)/(a^4*b^{12}*c^{16} + a^{16}*c^4*d^{12} - 12*a^5*b^{11}*c^{15}*d - 12*a^{15}* \\
& b*c^5*d^{11} + 66*a^6*b^{10}*c^{14}*d^2 - 220*a^7*b^9*c^{13}*d^3 + 495*a^8*b^8*c^{12} \\
& *d^4 - 792*a^9*b^7*c^{11}*d^5 + 924*a^{10}*b^6*c^{10}*d^6 - 792*a^{11}*b^5*c^9*d^7 \\
& + 495*a^{12}*b^4*c^8*d^8 - 220*a^{13}*b^3*c^7*d^9 + 66*a^{14}*b^2*c^6*d^{10}) - (3* \\
& x*(-c^5*d^5)^{(1/2)}*(a^2*d^2 + 21*b^2*c^2 - 6*a*b*c*d)*(256*a^4*b^{13}*c^{15}*d^2 \\
& - 2304*a^5*b^{12}*c^{14}*d^3 + 8960*a^6*b^{11}*c^{13}*d^4 - 19200*a^7*b^{10}*c^{12}*d \\
& ^5 + 23040*a^8*b^9*c^{11}*d^6 - 10752*a^9*b^8*c^{10}*d^7 - 10752*a^{10}*b^7*c^9*d \\
& ^8 + 23040*a^{11}*b^6*c^8*d^9 - 19200*a^{12}*b^5*c^7*d^{10} + 8960*a^{13}*b^4*c^6*d \\
& ^{11} - 2304*a^{14}*b^3*c^5*d^{12} + 256*a^{15}*b^2*c^4*d^{13}))/((512*(b^5*c^{10} - a^5 \\
& *c^5*d^5 + 5*a^4*b*c^6*d^4 + 10*a^2*b^3*c^8*d^2 - 10*a^3*b^2*c^7*d^3 - 5*a* \\
& b^4*c^9*d)*(a^4*b^8*c^{12} + a^{12}*c^4*d^8 - 8*a^5*b^7*c^{11}*d - 8*a^{11}*b*c^5*d \\
& ^7 + 28*a^6*b^6*c^{10}*d^2 - 56*a^7*b^5*c^9*d^3 + 70*a^8*b^4*c^8*d^4 - 56*a^9 \\
& *b^3*c^7*d^5 + 28*a^{10}*b^2*c^6*d^6)))*(-c^5*d^5)^{(1/2)}*(a^2*d^2 + 21*b^2*c^ \\
& 2 - 6*a*b*c*d))/(16*(b^5*c^{10} - a^5*c^5*d^5 + 5*a^4*b*c^6*d^4 + 10*a^2*b^3* \\
& c^8*d^2 - 10*a^3*b^2*c^7*d^3 - 5*a*b^4*c^9*d)))*(-c^5*d^5)^{(1/2)}*(a^2*d^2 + \\
& 21*b^2*c^2 - 6*a*b*c*d)*3i)/(16*(b^5*c^{10} - a^5*c^5*d^5 + 5*a^4*b*c^6*d^4 \\
& + 10*a^2*b^3*c^8*d^2 - 10*a^3*b^2*c^7*d^3 - 5*a*b^4*c^9*d)) + (((x*(9*a^8*b \\
& ^3*d^{11} + 9*b^{11}*c^8*d^3 - 108*a*b^{10}*c^7*d^4 - 108*a^7*b^4*c*d^{10} + 702*a^ \\
& 2*b^9*c^6*d^5 - 2268*a^3*b^8*c^5*d^6 + 7938*a^4*b^7*c^4*d^7 - 2268*a^5*b^6* \\
& c^3*d^8 + 702*a^6*b^5*c^2*d^9))/(32*(a^4*b^8*c^{12} + a^{12}*c^4*d^8 - 8*a^5*b^ \\
& 7*c^{11}*d - 8*a^{11}*b*c^5*d^7 + 28*a^6*b^6*c^{10}*d^2 - 56*a^7*b^5*c^9*d^3 + 70 \\
& *a^8*b^4*c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^{10}*b^2*c^6*d^6)) + (3*((3*a^2 \\
& *b^{16}*c^{16}*d^2)/2 - (45*a^3*b^{15}*c^{15}*d^3)/2 + (333*a^4*b^{14}*c^{14}*d^4)/2 - \\
& 765*a^5*b^{13}*c^{13}*d^5 + (4743*a^6*b^{12}*c^{12}*d^6)/2 - (10371*a^7*b^{11}*c^{11}*d \\
& ^7)/2 + (16425*a^8*b^{10}*c^{10}*d^8)/2 - 9558*a^9*b^9*c^9*d^9 + (16425*a^{10}*b^ \\
& 8*c^8*d^{10})/2 - (10371*a^{11}*b^7*c^7*d^{11})/2 + (4743*a^{12}*b^6*c^6*d^{12})/2 - \\
& 765*a^{13}*b^5*c^5*d^{13} + (333*a^{14}*b^4*c^4*d^{14})/2 - (45*a^{15}*b^3*c^3*d^{15})/ \\
& 2 + (3*a^{16}*b^2*c^2*d^{16})/2)/(a^4*b^{12}*c^{16} + a^{16}*c^4*d^{12} - 12*a^5*b^{11}*c \\
& ^{15}*d - 12*a^{15}*b*c^5*d^{11} + 66*a^6*b^{10}*c^{14}*d^2 - 220*a^7*b^9*c^{13}*d^3 + \\
& 495*a^8*b^8*c^{12}*d^4 - 792*a^9*b^7*c^{11}*d^5 + 924*a^{10}*b^6*c^{10}*d^6 - 792*a \\
& ^{11}*b^5*c^9*d^7 + 495*a^{12}*b^4*c^8*d^8 - 220*a^{13}*b^3*c^7*d^9 + 66*a^{14}*b^2 \\
& *c^6*d^{10}) + (3*x*(-c^5*d^5)^{(1/2)}*(a^2*d^2 + 21*b^2*c^2 - 6*a*b*c*d)*(256* \\
& a^4*b^{13}*c^{15}*d^2 - 2304*a^5*b^{12}*c^{14}*d^3 + 8960*a^6*b^{11}*c^{13}*d^4 - 19200 \\
& *a^7*b^{10}*c^{12}*d^5 + 23040*a^8*b^9*c^{11}*d^6 - 10752*a^9*b^8*c^{10}*d^7 - 1075 \\
& 2*a^{10}*b^7*c^9*d^8 + 23040*a^{11}*b^6*c^8*d^9 - 19200*a^{12}*b^5*c^7*d^{10} + 896 \\
& 0*a^{13}*b^4*c^6*d^{11} - 2304*a^{14}*b^3*c^5*d^{12} + 256*a^{15}*b^2*c^4*d^{13}))/((512 \\
& *(b^5*c^{10} - a^5*c^5*d^5 + 5*a^4*b*c^6*d^4 + 10*a^2*b^3*c^8*d^2 - 10*a^3*b^
\end{aligned}$$

$$\begin{aligned}
& 2*c^7*d^3 - 5*a*b^4*c^9*d)*(a^4*b^8*c^12 + a^12*c^4*d^8 - 8*a^5*b^7*c^11*d \\
& - 8*a^11*b*c^5*d^7 + 28*a^6*b^6*c^10*d^2 - 56*a^7*b^5*c^9*d^3 + 70*a^8*b^4* \\
& c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^10*b^2*c^6*d^6)))*(-c^5*d^5)^{(1/2)}*(a^2 \\
& *d^2 + 21*b^2*c^2 - 6*a*b*c*d))/(16*(b^5*c^10 - a^5*c^5*d^5 + 5*a^4*b*c^6*d \\
& ^4 + 10*a^2*b^3*c^8*d^2 - 10*a^3*b^2*c^7*d^3 - 5*a*b^4*c^9*d)))*(-c^5*d^5)^{(1/2)} \\
& *(a^2*d^2 + 21*b^2*c^2 - 6*a*b*c*d)*3i)/(16*(b^5*c^10 - a^5*c^5*d^5 + \\
& 5*a^4*b*c^6*d^4 + 10*a^2*b^3*c^8*d^2 - 10*a^3*b^2*c^7*d^3 - 5*a*b^4*c^9*d)) \\
&)/(((567*a^7*b^5*d^12)/256 + (567*b^12*c^7*d^5)/256 - (6399*a*b^11*c^6*d^6) \\
& /256 - (6399*a^6*b^6*c*d^11)/256 + (27891*a^2*b^10*c^5*d^7)/256 - (49707*a^ \\
& 3*b^9*c^4*d^8)/256 - (49707*a^4*b^8*c^3*d^9)/256 + (27891*a^5*b^7*c^2*d^10) \\
& /256)/(a^4*b^12*c^16 + a^16*c^4*d^12 - 12*a^5*b^11*c^15*d - 12*a^15*b*c^5*d \\
& ^11 + 66*a^6*b^10*c^14*d^2 - 220*a^7*b^9*c^13*d^3 + 495*a^8*b^8*c^12*d^4 - \\
& 792*a^9*b^7*c^11*d^5 + 924*a^10*b^6*c^10*d^6 - 792*a^11*b^5*c^9*d^7 + 495*a \\
& ^12*b^4*c^8*d^8 - 220*a^13*b^3*c^7*d^9 + 66*a^14*b^2*c^6*d^10) + (3*((x*(9* \\
& a^8*b^3*d^11 + 9*b^11*c^8*d^3 - 108*a*b^10*c^7*d^4 - 108*a^7*b^4*c*d^10 + 7 \\
& 02*a^2*b^9*c^6*d^5 - 2268*a^3*b^8*c^5*d^6 + 7938*a^4*b^7*c^4*d^7 - 2268*a^5 \\
& *b^6*c^3*d^8 + 702*a^6*b^5*c^2*d^9)))/(32*(a^4*b^8*c^12 + a^12*c^4*d^8 - 8*a \\
& ^5*b^7*c^11*d - 8*a^11*b*c^5*d^7 + 28*a^6*b^6*c^10*d^2 - 56*a^7*b^5*c^9*d^3 \\
& + 70*a^8*b^4*c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^10*b^2*c^6*d^6)) - (3*((\\
& 3*a^2*b^16*c^16*d^2)/2 - (45*a^3*b^15*c^15*d^3)/2 + (333*a^4*b^14*c^14*d^4) \\
& /2 - 765*a^5*b^13*c^13*d^5 + (4743*a^6*b^12*c^12*d^6)/2 - (10371*a^7*b^11*c \\
& ^11*d^7)/2 + (16425*a^8*b^10*c^10*d^8)/2 - 9558*a^9*b^9*c^9*d^9 + (16425*a^ \\
& 10*b^8*c^8*d^10)/2 - (10371*a^11*b^7*c^7*d^11)/2 + (4743*a^12*b^6*c^6*d^12) \\
& /2 - 765*a^13*b^5*c^5*d^13 + (333*a^14*b^4*c^4*d^14)/2 - (45*a^15*b^3*c^3*d \\
& ^15)/2 + (3*a^16*b^2*c^2*d^16)/2)/(a^4*b^12*c^16 + a^16*c^4*d^12 - 12*a^5*b \\
& ^11*c^15*d - 12*a^15*b*c^5*d^11 + 66*a^6*b^10*c^14*d^2 - 220*a^7*b^9*c^13*d \\
& ^3 + 495*a^8*b^8*c^12*d^4 - 792*a^9*b^7*c^11*d^5 + 924*a^10*b^6*c^10*d^6 - \\
& 792*a^11*b^5*c^9*d^7 + 495*a^12*b^4*c^8*d^8 - 220*a^13*b^3*c^7*d^9 + 66*a^1 \\
& 4*b^2*c^6*d^10) - (3*x*(-c^5*d^5)^{(1/2)}*(a^2*d^2 + 21*b^2*c^2 - 6*a*b*c*d)* \\
& (256*a^4*b^13*c^15*d^2 - 2304*a^5*b^12*c^14*d^3 + 8960*a^6*b^11*c^13*d^4 - \\
& 19200*a^7*b^10*c^12*d^5 + 23040*a^8*b^9*c^11*d^6 - 10752*a^9*b^8*c^10*d^7 - \\
& 10752*a^10*b^7*c^9*d^8 + 23040*a^11*b^6*c^8*d^9 - 19200*a^12*b^5*c^7*d^10 \\
& + 8960*a^13*b^4*c^6*d^11 - 2304*a^14*b^3*c^5*d^12 + 256*a^15*b^2*c^4*d^13)) \\
& /((512*(b^5*c^10 - a^5*c^5*d^5 + 5*a^4*b*c^6*d^4 + 10*a^2*b^3*c^8*d^2 - 10*a \\
& ^3*b^2*c^7*d^3 - 5*a*b^4*c^9*d))*(a^4*b^8*c^12 + a^12*c^4*d^8 - 8*a^5*b^7*c^ \\
& 11*d - 8*a^11*b*c^5*d^7 + 28*a^6*b^6*c^10*d^2 - 56*a^7*b^5*c^9*d^3 + 70*a^8 \\
& *b^4*c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^10*b^2*c^6*d^6)))*(-c^5*d^5)^{(1/2)} \\
& *(a^2*d^2 + 21*b^2*c^2 - 6*a*b*c*d))/(16*(b^5*c^10 - a^5*c^5*d^5 + 5*a^4*b* \\
& c^6*d^4 + 10*a^2*b^3*c^8*d^2 - 10*a^3*b^2*c^7*d^3 - 5*a*b^4*c^9*d)))*(-c^5*d^ \\
& 5)^{(1/2)}*(a^2*d^2 + 21*b^2*c^2 - 6*a*b*c*d))/(16*(b^5*c^10 - a^5*c^5*d^5 \\
& + 5*a^4*b*c^6*d^4 + 10*a^2*b^3*c^8*d^2 - 10*a^3*b^2*c^7*d^3 - 5*a*b^4*c^9*d \\
&)) - (3*((x*(9*a^8*b^3*d^11 + 9*b^11*c^8*d^3 - 108*a*b^10*c^7*d^4 - 108*a^7 \\
& *b^4*c*d^10 + 702*a^2*b^9*c^6*d^5 - 2268*a^3*b^8*c^5*d^6 + 7938*a^4*b^7*c^4 \\
& *d^7 - 2268*a^5*b^6*c^3*d^8 + 702*a^6*b^5*c^2*d^9)))/(32*(a^4*b^8*c^12 + a^1 \\
& 2*c^4*d^8 - 8*a^5*b^7*c^11*d - 8*a^11*b*c^5*d^7 + 28*a^6*b^6*c^10*d^2 - 56* \\
& a^7*b^5*c^9*d^3 + 70*a^8*b^4*c^8*d^4 - 56*a^9*b^3*c^7*d^5 + 28*a^10*b^2*c^6 \\
& *d^6)) + (3*((3*a^2*b^16*c^16*d^2)/2 - (45*a^3*b^15*c^15*d^3)/2 + (333*a^4 \\
& *b^14*c^14*d^4)/2 - 765*a^5*b^13*c^13*d^5 + (4743*a^6*b^12*c^12*d^6)/2 - (1 \\
& 0371*a^7*b^11*c^11*d^7)/2 + (16425*a^8*b^10*c^10*d^8)/2 - 9558*a^9*b^9*c^9* \\
& d^9 + (16425*a^10*b^8*c^8*d^10)/2 - (10371*a^11*b^7*c^7*d^11)/2 + (4743*a^1 \\
& 2*b^6*c^6*d^12)/2 - 765*a^13*b^5*c^5*d^13 + (333*a^14*b^4*c^4*d^14)/2 - (45 \\
& *a^15*b^3*c^3*d^15)/2 + (3*a^16*b^2*c^2*d^16)/2)/(a^4*b^12*c^16 + a^16*c^4* \\
& d^12 - 12*a^5*b^11*c^15*d - 12*a^15*b*c^5*d^11 + 66*a^6*b^10*c^14*d^2 - 220 \\
& *a^7*b^9*c^13*d^3 + 495*a^8*b^8*c^12*d^4 - 792*a^9*b^7*c^11*d^5 + 924*a^10* \\
& b^6*c^10*d^6 - 792*a^11*b^5*c^9*d^7 + 495*a^12*b^4*c^8*d^8 - 220*a^13*b^3*c \\
& ^7*d^9 + 66*a^14*b^2*c^6*d^10) + (3*x*(-c^5*d^5)^{(1/2)}*(a^2*d^2 + 21*b^2*c^ \\
& 2 - 6*a*b*c*d)*(256*a^4*b^13*c^15*d^2 - 2304*a^5*b^12*c^14*d^3 + 8960*a^6*b \\
& ^11*c^13*d^4 - 19200*a^7*b^10*c^12*d^5 + 23040*a^8*b^9*c^11*d^6 - 10752*a^9 \\
& *b^8*c^10*d^7 - 10752*a^10*b^7*c^9*d^8 + 23040*a^11*b^6*c^8*d^9 - 19200*a^1
\end{aligned}$$

$$\frac{2b^5c^7d^{10} + 8960a^{13}b^4c^6d^{11} - 2304a^{14}b^3c^5d^{12} + 256a^{15}b^2c^4d^{13}}{(512(b^5c^{10} - a^5c^5d^5 + 5a^4b^3c^6d^4 + 10a^2b^3c^8d^2 - 10a^3b^2c^7d^3 - 5ab^4c^9d)) \cdot (a^4b^8c^{12} + a^{12}c^4d^8 - 8a^5b^7c^{11}d - 8a^{11}b^3c^5d^7 + 28a^6b^6c^{10}d^2 - 56a^7b^5c^9d^3 + 70a^8b^4c^8d^4 - 56a^9b^3c^7d^5 + 28a^{10}b^2c^6d^6))} \cdot (-c^5d^5)^{1/2} \cdot (a^2d^2 + 21b^2c^2 - 6abc^2d)}{(16(b^5c^{10} - a^5c^5d^5 + 5a^4b^3c^6d^4 + 10a^2b^3c^8d^2 - 10a^3b^2c^7d^3 - 5ab^4c^9d))} \cdot (-c^5d^5)^{1/2} \cdot (a^2d^2 + 21b^2c^2 - 6abc^2d)}{(16(b^5c^{10} - a^5c^5d^5 + 5a^4b^3c^6d^4 + 10a^2b^3c^8d^2 - 10a^3b^2c^7d^3 - 5ab^4c^9d))} \cdot (-c^5d^5)^{1/2} \cdot (a^2d^2 + 21b^2c^2 - 6abc^2d) \cdot 3i} / (8(b^5c^{10} - a^5c^5d^5 + 5a^4b^3c^6d^4 + 10a^2b^3c^8d^2 - 10a^3b^2c^7d^3 - 5ab^4c^9d))$$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**3/(d*x**2+c)**3,x)

[Out] Timed out

$$3.43 \quad \int \frac{(-1+x^2)^3}{(1+x^2)^4} dx$$

Optimal. Leaf size=34

$$-\frac{x(1-x^2)^2}{3(x^2+1)^3} - \frac{2x}{3(x^2+1)}$$

[Out] $-1/3*x*(-x^2+1)^2/(x^2+1)^3-2/3*x/(x^2+1)$

Rubi [A] time = 0.01, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {413, 21, 383}

$$-\frac{x(1-x^2)^2}{3(x^2+1)^3} - \frac{2x}{3(x^2+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(-1 + x^2)^3/(1 + x^2)^4, x]$

[Out] $-(x*(1 - x^2)^2)/(3*(1 + x^2)^3) - (2*x)/(3*(1 + x^2))$

Rule 21

$\text{Int}[(u_.)*((a_.) + (b_.)*(v_))^{(m_.)}*((c_.) + (d_.)*(v_))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b/d)^m, \text{Int}[u*(c + d*v)^{(m+n)}, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \&\& \text{EqQ}[b*c - a*d, 0] \&\& \text{IntegerQ}[m] \&\& (!\text{IntegerQ}[n] \mid\mid \text{SimplerQ}[c + d*x, a + b*x])$

Rule 383

$\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(c*x*(a + b*x^n)^{(p+1)})/a, x] /; \text{FreeQ}\{a, b, c, d, n, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a*d - b*c*(n*(p+1) + 1), 0]$

Rule 413

$\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)})^{(q_.)}, x_Symbol] \rightarrow \text{Simp}[(a*d - c*b)*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-1)}/(a*b*n*(p+1)), x] - \text{Dist}[1/(a*b*n*(p+1)), \text{Int}[(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(a*d - c*b*(n*(p+1) + 1)) + d*(a*d*(n*(q-1) + 1) - b*c*(n*(p+q) + 1))*x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[p, -1] \&\& \text{GtQ}[q, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rubi steps

$$\begin{aligned} \int \frac{(-1+x^2)^3}{(1+x^2)^4} dx &= -\frac{x(1-x^2)^2}{3(1+x^2)^3} + \frac{1}{6} \int \frac{(-1+x^2)(4+4x^2)}{(1+x^2)^3} dx \\ &= -\frac{x(1-x^2)^2}{3(1+x^2)^3} + \frac{2}{3} \int \frac{-1+x^2}{(1+x^2)^2} dx \\ &= -\frac{x(1-x^2)^2}{3(1+x^2)^3} - \frac{2x}{3(1+x^2)} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 0.71

$$\frac{x(3x^4 + 2x^2 + 3)}{3(x^2 + 1)^3}$$

Antiderivative was successfully verified.

[In] Integrate[(-1 + x^2)^3/(1 + x^2)^4,x]

[Out] -1/3*(x*(3 + 2*x^2 + 3*x^4))/(1 + x^2)^3

fricas [A] time = 0.76, size = 33, normalized size = 0.97

$$\frac{3x^5 + 2x^3 + 3x}{3(x^6 + 3x^4 + 3x^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2-1)^3/(x^2+1)^4,x, algorithm="fricas")

[Out] -1/3*(3*x^5 + 2*x^3 + 3*x)/(x^6 + 3*x^4 + 3*x^2 + 1)

giac [A] time = 0.58, size = 20, normalized size = 0.59

$$\frac{3\left(x + \frac{1}{x}\right)^2 - 4}{3\left(x + \frac{1}{x}\right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2-1)^3/(x^2+1)^4,x, algorithm="giac")

[Out] -1/3*(3*(x + 1/x)^2 - 4)/(x + 1/x)^3

maple [A] time = 0.01, size = 23, normalized size = 0.68

$$\frac{-x^5 - \frac{2}{3}x^3 - x}{(x^2 + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2-1)^3/(x^2+1)^4,x)

[Out] (-x^5-2/3*x^3-x)/(x^2+1)^3

maxima [A] time = 1.32, size = 33, normalized size = 0.97

$$\frac{3x^5 + 2x^3 + 3x}{3(x^6 + 3x^4 + 3x^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2-1)^3/(x^2+1)^4,x, algorithm="maxima")

[Out] -1/3*(3*x^5 + 2*x^3 + 3*x)/(x^6 + 3*x^4 + 3*x^2 + 1)

mupad [B] time = 5.00, size = 31, normalized size = 0.91

$$\frac{4x}{3(x^2 + 1)^2} - \frac{x}{x^2 + 1} - \frac{4x}{3(x^2 + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2 - 1)^3/(x^2 + 1)^4,x)`

[Out] $(4*x)/(3*(x^2 + 1)^2) - x/(x^2 + 1) - (4*x)/(3*(x^2 + 1)^3)$

sympy [A] time = 0.13, size = 31, normalized size = 0.91

$$\frac{-3x^5 - 2x^3 - 3x}{3x^6 + 9x^4 + 9x^2 + 3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((x**2-1)**3/(x**2+1)**4,x)`

[Out] $(-3*x**5 - 2*x**3 - 3*x)/(3*x**6 + 9*x**4 + 9*x**2 + 3)$

$$3.44 \quad \int \frac{(-1+x^2)^4}{(1+x^2)^5} dx$$

Optimal. Leaf size=47

$$\frac{x(1-x^2)^3}{4(x^2+1)^4} + \frac{3x(1-x^2)}{8(x^2+1)^2} + \frac{3}{8}\tan^{-1}(x)$$

[Out] 1/4*x*(-x^2+1)^3/(x^2+1)^4+3/8*x*(-x^2+1)/(x^2+1)^2+3/8*arctan(x)

Rubi [A] time = 0.02, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {413, 21, 203}

$$\frac{x(1-x^2)^3}{4(x^2+1)^4} + \frac{3x(1-x^2)}{8(x^2+1)^2} + \frac{3}{8}\tan^{-1}(x)$$

Antiderivative was successfully verified.

[In] Int[(-1 + x^2)^4/(1 + x^2)^5, x]

[Out] (x*(1 - x^2)^3)/(4*(1 + x^2)^4) + (3*x*(1 - x^2))/(8*(1 + x^2)^2) + (3*ArcTan[x])/8

Rule 21

```
Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
  Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

Rule 203

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[Rt[b, 2]*x]/Rt
[a, 2])]/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 413

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p +
1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q -
2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p
+ q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,
0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{(-1+x^2)^4}{(1+x^2)^5} dx &= \frac{x(1-x^2)^3}{4(1+x^2)^4} + \frac{1}{8} \int \frac{(-1+x^2)^2(6+6x^2)}{(1+x^2)^4} dx \\
&= \frac{x(1-x^2)^3}{4(1+x^2)^4} + \frac{3}{4} \int \frac{(-1+x^2)^2}{(1+x^2)^3} dx \\
&= \frac{x(1-x^2)^3}{4(1+x^2)^4} + \frac{3x(1-x^2)}{8(1+x^2)^2} + \frac{3}{16} \int \frac{2+2x^2}{(1+x^2)^2} dx \\
&= \frac{x(1-x^2)^3}{4(1+x^2)^4} + \frac{3x(1-x^2)}{8(1+x^2)^2} + \frac{3}{8} \int \frac{1}{1+x^2} dx \\
&= \frac{x(1-x^2)^3}{4(1+x^2)^4} + \frac{3x(1-x^2)}{8(1+x^2)^2} + \frac{3}{8} \tan^{-1}(x)
\end{aligned}$$

Mathematica [A] time = 0.01, size = 41, normalized size = 0.87

$$\frac{-5x^7 + 3x^5 - 3x^3 + 3(x^2 + 1)^4 \tan^{-1}(x) + 5x}{8(x^2 + 1)^4}$$

Antiderivative was successfully verified.

[In] Integrate[(-1 + x^2)^4/(1 + x^2)^5, x]

[Out] (5*x - 3*x^3 + 3*x^5 - 5*x^7 + 3*(1 + x^2)^4*ArcTan[x])/(8*(1 + x^2)^4)

fricas [A] time = 0.81, size = 67, normalized size = 1.43

$$\frac{5x^7 - 3x^5 + 3x^3 - 3(x^8 + 4x^6 + 6x^4 + 4x^2 + 1) \arctan(x) - 5x}{8(x^8 + 4x^6 + 6x^4 + 4x^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2-1)^4/(x^2+1)^5, x, algorithm="fricas")

[Out] -1/8*(5*x^7 - 3*x^5 + 3*x^3 - 3*(x^8 + 4*x^6 + 6*x^4 + 4*x^2 + 1)*arctan(x) - 5*x)/(x^8 + 4*x^6 + 6*x^4 + 4*x^2 + 1)

giac [A] time = 0.58, size = 54, normalized size = 1.15

$$\frac{3}{32} \pi \operatorname{sgn}(x) - \frac{5\left(x - \frac{1}{x}\right)^3 + 12x - \frac{12}{x}}{8\left(\left(x - \frac{1}{x}\right)^2 + 4\right)^2} + \frac{3}{16} \arctan\left(\frac{x^2 - 1}{2x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2-1)^4/(x^2+1)^5, x, algorithm="giac")

[Out] 3/32*pi*sgn(x) - 1/8*(5*(x - 1/x)^3 + 12*x - 12/x)/((x - 1/x)^2 + 4)^2 + 3/16*arctan(1/2*(x^2 - 1)/x)

maple [A] time = 0.01, size = 33, normalized size = 0.70

$$\frac{3 \arctan(x)}{8} + \frac{-\frac{5}{8}x^7 + \frac{3}{8}x^5 - \frac{3}{8}x^3 + \frac{5}{8}x}{(x^2 + 1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2-1)^4/(x^2+1)^5,x)`

[Out] $(-5/8*x^7+3/8*x^5-3/8*x^3+5/8*x)/(x^2+1)^4+3/8*\arctan(x)$

maxima [A] time = 2.99, size = 48, normalized size = 1.02

$$-\frac{5x^7 - 3x^5 + 3x^3 - 5x}{8(x^8 + 4x^6 + 6x^4 + 4x^2 + 1)} + \frac{3}{8} \arctan(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((x^2-1)^4/(x^2+1)^5,x, algorithm="maxima")`

[Out] $-1/8*(5*x^7 - 3*x^5 + 3*x^3 - 5*x)/(x^8 + 4*x^6 + 6*x^4 + 4*x^2 + 1) + 3/8*\arctan(x)$

mupad [B] time = 0.04, size = 47, normalized size = 1.00

$$\frac{3 \operatorname{atan}(x)}{8} + \frac{-\frac{5x^7}{8} + \frac{3x^5}{8} - \frac{3x^3}{8} + \frac{5x}{8}}{x^8 + 4x^6 + 6x^4 + 4x^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2 - 1)^4/(x^2 + 1)^5,x)`

[Out] $(3*\operatorname{atan}(x))/8 + ((5*x)/8 - (3*x^3)/8 + (3*x^5)/8 - (5*x^7)/8)/(4*x^2 + 6*x^4 + 4*x^6 + x^8 + 1)$

sympy [A] time = 0.17, size = 46, normalized size = 0.98

$$\frac{-5x^7 + 3x^5 - 3x^3 + 5x}{8x^8 + 32x^6 + 48x^4 + 32x^2 + 8} + \frac{3 \operatorname{atan}(x)}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((x**2-1)**4/(x**2+1)**5,x)`

[Out] $(-5*x**7 + 3*x**5 - 3*x**3 + 5*x)/(8*x**8 + 32*x**6 + 48*x**4 + 32*x**2 + 8) + 3*\operatorname{atan}(x)/8$

3.45 $\int \sqrt{a + bx^2} (c + dx^2)^3 dx$

Optimal. Leaf size=231

$$\frac{dx (a + bx^2)^{3/2} (15a^2d^2 - 52abcd + 72b^2c^2)}{192b^3} + \frac{x\sqrt{a + bx^2} (-5a^3d^3 + 24a^2bcd^2 - 48ab^2c^2d + 64b^3c^3)}{128b^3} + \frac{a(-5a^3d^3 + 24a^2bcd^2 - 48ab^2c^2d + 64b^3c^3)}{128b^3}$$

[Out] $1/192*d*(15*a^2*d^2-52*a*b*c*d+72*b^2*c^2)*x*(b*x^2+a)^(3/2)/b^3+1/48*d*(-5*a*d+12*b*c)*x*(b*x^2+a)^(3/2)*(d*x^2+c)/b^2+1/8*d*x*(b*x^2+a)^(3/2)*(d*x^2+c)^2/b+1/128*a*(-5*a^3*d^3+24*a^2*b*c*d^2-48*a*b^2*c^2*d+64*b^3*c^3)*\operatorname{arctanh}(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(7/2)+1/128*(-5*a^3*d^3+24*a^2*b*c*d^2-48*a*b^2*c^2*d+64*b^3*c^3)*x*(b*x^2+a)^(1/2)/b^3$

Rubi [A] time = 0.18, antiderivative size = 231, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {416, 528, 388, 195, 217, 206}

$$\frac{dx (a + bx^2)^{3/2} (15a^2d^2 - 52abcd + 72b^2c^2)}{192b^3} + \frac{x\sqrt{a + bx^2} (24a^2bcd^2 - 5a^3d^3 - 48ab^2c^2d + 64b^3c^3)}{128b^3} + \frac{a(24a^2bcd^2 - 5a^3d^3 - 48ab^2c^2d + 64b^3c^3)}{128b^3}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[a + b*x^2]*(c + d*x^2)^3,x]`

[Out] $((64*b^3*c^3 - 48*a*b^2*c^2*d + 24*a^2*b*c*d^2 - 5*a^3*d^3)*x*\operatorname{Sqrt}[a + b*x^2])/(128*b^3) + (d*(72*b^2*c^2 - 52*a*b*c*d + 15*a^2*d^2)*x*(a + b*x^2)^(3/2))/(192*b^3) + (d*(12*b*c - 5*a*d)*x*(a + b*x^2)^(3/2)*(c + d*x^2))/(48*b^2) + (d*x*(a + b*x^2)^(3/2)*(c + d*x^2)^2)/(8*b) + (a*(64*b^3*c^3 - 48*a*b^2*c^2*d + 24*a^2*b*c*d^2 - 5*a^3*d^3)*\operatorname{ArcTanh}[\operatorname{Sqrt}[b]*x/\operatorname{Sqrt}[a + b*x^2]])/(128*b^(7/2))$

Rule 195

`Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])`

Rule 206

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 217

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

Rule 388

`Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]`

Rule 416

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)),
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a
, b, c, d, n, p, q, x]
```

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rubi steps

$$\begin{aligned}
\int \sqrt{a+bx^2} (c+dx^2)^3 dx &= \frac{dx (a+bx^2)^{3/2} (c+dx^2)^2}{8b} + \frac{\int \sqrt{a+bx^2} (c+dx^2) (c(8bc-ad) + d(12bc-5ad)x^2)}{8b} \\
&= \frac{d(12bc-5ad)x (a+bx^2)^{3/2} (c+dx^2)}{48b^2} + \frac{dx (a+bx^2)^{3/2} (c+dx^2)^2}{8b} + \frac{\int \sqrt{a+bx^2} (c+dx^2)}{8b} \\
&= \frac{d(72b^2c^2 - 52abcd + 15a^2d^2)x (a+bx^2)^{3/2}}{192b^3} + \frac{d(12bc-5ad)x (a+bx^2)^{3/2} (c+dx^2)}{48b^2} \\
&= \frac{(64b^3c^3 - 48ab^2c^2d + 24a^2bcd^2 - 5a^3d^3)x\sqrt{a+bx^2}}{128b^3} + \frac{d(72b^2c^2 - 52abcd + 15a^2d^2)x\sqrt{a+bx^2}}{192b^3} \\
&= \frac{(64b^3c^3 - 48ab^2c^2d + 24a^2bcd^2 - 5a^3d^3)x\sqrt{a+bx^2}}{128b^3} + \frac{d(72b^2c^2 - 52abcd + 15a^2d^2)x\sqrt{a+bx^2}}{192b^3} \\
&= \frac{(64b^3c^3 - 48ab^2c^2d + 24a^2bcd^2 - 5a^3d^3)x\sqrt{a+bx^2}}{128b^3} + \frac{d(72b^2c^2 - 52abcd + 15a^2d^2)x\sqrt{a+bx^2}}{192b^3}
\end{aligned}$$

Mathematica [A] time = 5.11, size = 181, normalized size = 0.78

$$\frac{\sqrt{b}x\sqrt{a+bx^2} (15a^3d^3 - 2a^2bd^2(36c + 5dx^2) + 8ab^2d(18c^2 + 6cdx^2 + d^2x^4) + 48b^3(4c^3 + 6c^2dx^2 + 4cd^2x^4 + d^3x^6))}{384b^{7/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[a + b*x^2]*(c + d*x^2)^3,x]
```

```
[Out] (Sqrt[b]*x*Sqrt[a + b*x^2]*(15*a^3*d^3 - 2*a^2*b*d^2*(36*c + 5*d*x^2) + 8*a
*b^2*d*(18*c^2 + 6*c*d*x^2 + d^2*x^4) + 48*b^3*(4*c^3 + 6*c^2*d*x^2 + 4*c*d
^2*x^4 + d^3*x^6)) - 3*a*(-64*b^3*c^3 + 48*a*b^2*c^2*d - 24*a^2*b*c*d^2 + 5
*a^3*d^3)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(384*b^(7/2))
```

fricas [A] time = 0.90, size = 398, normalized size = 1.72

$$\left[\frac{3(64ab^3c^3 - 48a^2b^2c^2d + 24a^3bcd^2 - 5a^4d^3)\sqrt{b} \log(-2bx^2 + 2\sqrt{bx^2+a}\sqrt{b}x - a) - 2(48b^4d^3x^7 + 8(24b^3d^3x^5 + 48b^2d^3x^3 + 48bd^3x) + 48b^3d^3)}{384b^{7/2}} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^3,x, algorithm="fricas")

[Out] [-1/768*(3*(64*a*b^3*c^3 - 48*a^2*b^2*c^2*d + 24*a^3*b*c*d^2 - 5*a^4*d^3)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(48*b^4*d^3*x^7 + 8*(24*b^4*c*d^2 + a*b^3*d^3)*x^5 + 2*(144*b^4*c^2*d + 24*a*b^3*c*d^2 - 5*a^2*b^2*d^3)*x^3 + 3*(64*b^4*c^3 + 48*a*b^3*c^2*d - 24*a^2*b^2*c*d^2 + 5*a^3*b*d^3)*x)*sqrt(b*x^2 + a))/b^4, -1/384*(3*(64*a*b^3*c^3 - 48*a^2*b^2*c^2*d + 24*a^3*b*c*d^2 - 5*a^4*d^3)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (48*b^4*d^3*x^7 + 8*(24*b^4*c*d^2 + a*b^3*d^3)*x^5 + 2*(144*b^4*c^2*d + 24*a*b^3*c*d^2 - 5*a^2*b^2*d^3)*x^3 + 3*(64*b^4*c^3 + 48*a*b^3*c^2*d - 24*a^2*b^2*c*d^2 + 5*a^3*b*d^3)*x)*sqrt(b*x^2 + a))/b^4]

giac [A] time = 0.64, size = 201, normalized size = 0.87

$$\frac{1}{384} \left(2 \left(4 \left(6d^3x^2 + \frac{24b^6cd^2 + ab^5d^3}{b^6} \right) x^2 + \frac{144b^6c^2d + 24ab^5cd^2 - 5a^2b^4d^3}{b^6} \right) x^2 + \frac{3(64b^6c^3 + 48ab^5c^2d - 24a^2b^4d^3)}{b^6} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^3,x, algorithm="giac")

[Out] 1/384*(2*(4*(6*d^3*x^2 + (24*b^6*c*d^2 + a*b^5*d^3)/b^6)*x^2 + (144*b^6*c^2*d + 24*a*b^5*c*d^2 - 5*a^2*b^4*d^3)/b^6)*x^2 + 3*(64*b^6*c^3 + 48*a*b^5*c^2*d - 24*a^2*b^4*c*d^2 + 5*a^3*b^3*d^3)/b^6)*sqrt(b*x^2 + a)*x - 1/128*(64*a*b^3*c^3 - 48*a^2*b^2*c^2*d + 24*a^3*b*c*d^2 - 5*a^4*d^3)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(7/2)

maple [A] time = 0.02, size = 310, normalized size = 1.34

$$\frac{(bx^2 + a)^{\frac{3}{2}} d^3 x^5}{8b} - \frac{5(bx^2 + a)^{\frac{3}{2}} a d^3 x^3}{48b^2} + \frac{(bx^2 + a)^{\frac{3}{2}} c d^2 x^3}{2b} - \frac{5a^4 d^3 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{128b^{\frac{7}{2}}} + \frac{3a^3 c d^2 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{16b^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)*(d*x^2+c)^3,x)

[Out] 1/8*d^3*x^5*(b*x^2+a)^(3/2)/b-5/48*d^3*a/b^2*x^3*(b*x^2+a)^(3/2)+5/64*d^3*a^2/b^3*x*(b*x^2+a)^(3/2)-5/128*d^3*a^3/b^3*x*(b*x^2+a)^(1/2)-5/128*d^3*a^4/b^(7/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+1/2*c*d^2*x^3*(b*x^2+a)^(3/2)/b-3/8*c*d^2*a/b^2*x*(b*x^2+a)^(3/2)+3/16*c*d^2*a^2/b^2*x*(b*x^2+a)^(1/2)+3/16*c*d^2*a^3/b^(5/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+3/4*c^2*d*x*(b*x^2+a)^(3/2)/b-3/8*c^2*d*a/b*x*(b*x^2+a)^(1/2)-3/8*c^2*d*a^2/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+1/2*c^3*x*(b*x^2+a)^(1/2)+1/2*c^3*a/b^(1/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))

maxima [A] time = 1.39, size = 281, normalized size = 1.22

$$\frac{(bx^2 + a)^{\frac{3}{2}} d^3 x^5}{8b} + \frac{(bx^2 + a)^{\frac{3}{2}} cd^2 x^3}{2b} - \frac{5(bx^2 + a)^{\frac{3}{2}} ad^3 x^3}{48b^2} + \frac{1}{2} \sqrt{bx^2 + a} c^3 x + \frac{3(bx^2 + a)^{\frac{3}{2}} c^2 dx}{4b} - \frac{3\sqrt{bx^2 + a} ac^2 dx}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^3,x, algorithm="maxima")

[Out] 1/8*(b*x^2 + a)^(3/2)*d^3*x^5/b + 1/2*(b*x^2 + a)^(3/2)*c*d^2*x^3/b - 5/48*(b*x^2 + a)^(3/2)*a*d^3*x^3/b^2 + 1/2*sqrt(b*x^2 + a)*c^3*x + 3/4*(b*x^2 + a)^(3/2)*c^2*d*x/b - 3/8*sqrt(b*x^2 + a)*a*c^2*d*x/b - 3/8*(b*x^2 + a)^(3/2)*a*c*d^2*x/b^2 + 3/16*sqrt(b*x^2 + a)*a^2*c*d^2*x/b^2 + 5/64*(b*x^2 + a)^(3/2)*a^2*d^3*x/b^3 - 5/128*sqrt(b*x^2 + a)*a^3*d^3*x/b^3 + 1/2*a*c^3*arcsin

$h(b*x/\sqrt{a*b})/\sqrt{b} - 3/8*a^2*c^2*d*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{(3/2)} + 3/16*a^3*c*d^2*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{(5/2)} - 5/128*a^4*d^3*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{(7/2)}$

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{bx^2 + a} (dx^2 + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^(1/2)*(c + d*x^2)^3, x)`

[Out] `int((a + b*x^2)^(1/2)*(c + d*x^2)^3, x)`

sympy [B] time = 20.38, size = 484, normalized size = 2.10

$$\frac{5a^{\frac{7}{2}}d^3x}{128b^3\sqrt{1 + \frac{bx^2}{a}}} - \frac{3a^{\frac{5}{2}}cd^2x}{16b^2\sqrt{1 + \frac{bx^2}{a}}} + \frac{5a^{\frac{5}{2}}d^3x^3}{384b^2\sqrt{1 + \frac{bx^2}{a}}} + \frac{3a^{\frac{3}{2}}c^2dx}{8b\sqrt{1 + \frac{bx^2}{a}}} - \frac{a^{\frac{3}{2}}cd^2x^3}{16b\sqrt{1 + \frac{bx^2}{a}}} - \frac{a^{\frac{3}{2}}d^3x^5}{192b\sqrt{1 + \frac{bx^2}{a}}} + \frac{\sqrt{a}c^3x\sqrt{1 + \frac{bx^2}{a}}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**(1/2)*(d*x**2+c)**3, x)`

[Out] `5*a**(7/2)*d**3*x/(128*b**3*sqrt(1 + b*x**2/a)) - 3*a**(5/2)*c*d**2*x/(16*b**2*sqrt(1 + b*x**2/a)) + 5*a**(5/2)*d**3*x**3/(384*b**2*sqrt(1 + b*x**2/a)) + 3*a**(3/2)*c**2*d*x/(8*b*sqrt(1 + b*x**2/a)) - a**(3/2)*c*d**2*x**3/(16*b*sqrt(1 + b*x**2/a)) - a**(3/2)*d**3*x**5/(192*b*sqrt(1 + b*x**2/a)) + sqrt(a)*c**3*x*sqrt(1 + b*x**2/a)/2 + 9*sqrt(a)*c**2*d*x**3/(8*sqrt(1 + b*x**2/a)) + 5*sqrt(a)*c*d**2*x**5/(8*sqrt(1 + b*x**2/a)) + 7*sqrt(a)*d**3*x**7/(48*sqrt(1 + b*x**2/a)) - 5*a**4*d**3*asinh(sqrt(b)*x/sqrt(a))/(128*b**(7/2)) + 3*a**3*c*d**2*asinh(sqrt(b)*x/sqrt(a))/(16*b**(5/2)) - 3*a**2*c**2*d*asinh(sqrt(b)*x/sqrt(a))/(8*b**(3/2)) + a*c**3*asinh(sqrt(b)*x/sqrt(a))/(2*sqrt(b)) + 3*b*c**2*d*x**5/(4*sqrt(a)*sqrt(1 + b*x**2/a)) + b*c*d**2*x**7/(2*sqrt(a)*sqrt(1 + b*x**2/a)) + b*d**3*x**9/(8*sqrt(a)*sqrt(1 + b*x**2/a))`

3.46 $\int \sqrt{a + bx^2} (c + dx^2)^2 dx$

Optimal. Leaf size=149

$$\frac{x\sqrt{a+bx^2}(a^2d^2-4abcd+8b^2c^2)}{16b^2} + \frac{a(a^2d^2-4abcd+8b^2c^2)\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16b^{5/2}} + \frac{dx(a+bx^2)^{3/2}(8bc-3ad)}{24b^2} + \dots$$

[Out] 1/24*d*(-3*a*d+8*b*c)*x*(b*x^2+a)^(3/2)/b^2+1/6*d*x*(b*x^2+a)^(3/2)*(d*x^2+c)/b+1/16*a*(a^2*d^2-4*a*b*c*d+8*b^2*c^2)*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(5/2)+1/16*(a^2*d^2-4*a*b*c*d+8*b^2*c^2)*x*(b*x^2+a)^(1/2)/b^2

Rubi [A] time = 0.09, antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {416, 388, 195, 217, 206}

$$\frac{x\sqrt{a+bx^2}(a^2d^2-4abcd+8b^2c^2)}{16b^2} + \frac{a(a^2d^2-4abcd+8b^2c^2)\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16b^{5/2}} + \frac{dx(a+bx^2)^{3/2}(8bc-3ad)}{24b^2} + \dots$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]*(c + d*x^2)^2,x]

[Out] ((8*b^2*c^2 - 4*a*b*c*d + a^2*d^2)*x*Sqrt[a + b*x^2])/(16*b^2) + (d*(8*b*c - 3*a*d)*x*(a + b*x^2)^(3/2))/(24*b^2) + (d*x*(a + b*x^2)^(3/2)*(c + d*x^2))/(6*b) + (a*(8*b^2*c^2 - 4*a*b*c*d + a^2*d^2)*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/(16*b^(5/2))

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)), x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -

1) + 1)) * x^n, x], x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rubi steps

$$\begin{aligned} \int \sqrt{a+bx^2} (c+dx^2)^2 dx &= \frac{dx (a+bx^2)^{3/2} (c+dx^2)}{6b} + \frac{\int \sqrt{a+bx^2} (c(6bc-ad) + d(8bc-3ad)x^2) dx}{6b} \\ &= \frac{d(8bc-3ad)x (a+bx^2)^{3/2}}{24b^2} + \frac{dx (a+bx^2)^{3/2} (c+dx^2)}{6b} + \frac{(8b^2c^2 - 4abcd + a^2d^2) \int}{8b^2} \\ &= \frac{(8b^2c^2 - 4abcd + a^2d^2)x\sqrt{a+bx^2}}{16b^2} + \frac{d(8bc-3ad)x (a+bx^2)^{3/2}}{24b^2} + \frac{dx (a+bx^2)^{3/2}}{6b} \\ &= \frac{(8b^2c^2 - 4abcd + a^2d^2)x\sqrt{a+bx^2}}{16b^2} + \frac{d(8bc-3ad)x (a+bx^2)^{3/2}}{24b^2} + \frac{dx (a+bx^2)^{3/2}}{6b} \\ &= \frac{(8b^2c^2 - 4abcd + a^2d^2)x\sqrt{a+bx^2}}{16b^2} + \frac{d(8bc-3ad)x (a+bx^2)^{3/2}}{24b^2} + \frac{dx (a+bx^2)^{3/2}}{6b} \end{aligned}$$

Mathematica [C] time = 2.67, size = 160, normalized size = 1.07

$$\frac{x\sqrt{a+bx^2} \left(2bx^2 (c+dx^2)^2 {}_3F_2\left(\frac{1}{2}, \frac{3}{2}, 2; 1, \frac{9}{2}; -\frac{bx^2}{a}\right) + 4bx^2 (2c^2 + 3cdx^2 + d^2x^4) {}_2F_1\left(\frac{1}{2}, \frac{3}{2}; \frac{9}{2}; -\frac{bx^2}{a}\right) + 7a (15c^2 + \dots) \right)}{105a\sqrt{\frac{bx^2}{a} + 1}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Sqrt[a + b*x^2]*(c + d*x^2)^2,x]

[Out] (x*Sqrt[a + b*x^2]*(7*a*(15*c^2 + 10*c*d*x^2 + 3*d^2*x^4)*Hypergeometric2F1[-1/2, 1/2, 7/2, -((b*x^2)/a)] + 4*b*x^2*(2*c^2 + 3*c*d*x^2 + d^2*x^4)*Hypergeometric2F1[1/2, 3/2, 9/2, -((b*x^2)/a)] + 2*b*x^2*(c + d*x^2)^2*HypergeometricPFQ[{1/2, 3/2, 2}, {1, 9/2}, -((b*x^2)/a)]))/(105*a*Sqrt[1 + (b*x^2)/a])

fricas [A] time = 0.97, size = 264, normalized size = 1.77

$$\frac{3(8ab^2c^2 - 4a^2bcd + a^3d^2)\sqrt{b} \log(-2bx^2 - 2\sqrt{bx^2+a}\sqrt{b}x - a) + 2(8b^3d^2x^5 + 2(12b^3cd + ab^2d^2)x^3 + 3(8ab^2c^2 - 4a^2bcd + a^3d^2)\sqrt{b} \log(-2bx^2 - 2\sqrt{bx^2+a}\sqrt{b}x - a))}{96b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^2,x, algorithm="fricas")

[Out] [1/96*(3*(8*a*b^2*c^2 - 4*a^2*b*c*d + a^3*d^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 2*(8*b^3*d^2*x^5 + 2*(12*b^3*c*d + a*b^2*d^2)*x^3 + 3*(8*b^3*c^2 + 4*a*b^2*c*d - a^2*b*d^2)*x)*sqrt(b*x^2 + a))/b^3, -1/48*(3*(8*a*b^2*c^2 - 4*a^2*b*c*d + a^3*d^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (8*b^3*d^2*x^5 + 2*(12*b^3*c*d + a*b^2*d^2)*x^3 + 3*(8*b^3*c^2 + 4*a*b^2*c*d - a^2*b*d^2)*x)*sqrt(b*x^2 + a))/b^3]

giac [A] time = 0.61, size = 129, normalized size = 0.87

$$\frac{1}{48} \left(2 \left(4d^2x^2 + \frac{12b^4cd + ab^3d^2}{b^4} \right) x^2 + \frac{3(8b^4c^2 + 4ab^3cd - a^2b^2d^2)}{b^4} \right) \sqrt{bx^2 + ax} - \frac{(8ab^2c^2 - 4a^2bcd + a^3d^2) \log}{16b^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^2,x, algorithm="giac")

[Out] $\frac{1}{48}*(2*(4*d^2*x^2 + (12*b^4*c*d + a*b^3*d^2)/b^4)*x^2 + 3*(8*b^4*c^2 + 4*a*b^3*c*d - a^2*b^2*d^2)/b^4)*\sqrt{b*x^2 + a}*x - \frac{1}{16}*(8*a*b^2*c^2 - 4*a^2*b*c*d + a^3*d^2)*\log(\text{abs}(-\sqrt{b}*x + \sqrt{b*x^2 + a}))/b^{5/2}$

maple [A] time = 0.01, size = 190, normalized size = 1.28

$$\frac{(bx^2 + a)^{\frac{3}{2}} d^2 x^3}{6b} + \frac{a^3 d^2 \ln(\sqrt{b} x + \sqrt{bx^2 + a})}{16b^{\frac{5}{2}}} - \frac{a^2 cd \ln(\sqrt{b} x + \sqrt{bx^2 + a})}{4b^{\frac{3}{2}}} + \frac{a c^2 \ln(\sqrt{b} x + \sqrt{bx^2 + a})}{2\sqrt{b}} + \sqrt{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)*(d*x^2+c)^2,x)

[Out] $\frac{1}{6}*d^2*x^3*(b*x^2+a)^{(3/2)}/b - \frac{1}{8}*d^2*a/b^2*x*(b*x^2+a)^{(3/2)} + \frac{1}{16}*d^2*a^2/b^2*x*(b*x^2+a)^{(1/2)} + \frac{1}{16}*d^2*a^3/b^{5/2}*\ln(b^{(1/2)*x+(b*x^2+a)^{(1/2)})} + \frac{1}{2}*c*d*x*(b*x^2+a)^{(3/2)}/b - \frac{1}{4}*c*d*a/b*x*(b*x^2+a)^{(1/2)} - \frac{1}{4}*c*d*a^2/b^{3/2}*\ln(b^{(1/2)*x+(b*x^2+a)^{(1/2)})} + \frac{1}{2}*c^2*x*(b*x^2+a)^{(1/2)} + \frac{1}{2}*c^2*a/b^{(1/2)}*\ln(b^{(1/2)*x+(b*x^2+a)^{(1/2)})}$

maxima [A] time = 1.35, size = 168, normalized size = 1.13

$$\frac{(bx^2 + a)^{\frac{3}{2}} d^2 x^3}{6b} + \frac{1}{2} \sqrt{bx^2 + a} c^2 x + \frac{(bx^2 + a)^{\frac{3}{2}} cdx}{2b} - \frac{\sqrt{bx^2 + a} acdx}{4b} - \frac{(bx^2 + a)^{\frac{3}{2}} ad^2 x}{8b^2} + \frac{\sqrt{bx^2 + a} a^2 d^2 x}{16b^2} + \frac{ac^2 \operatorname{arcsinh}\left(\frac{\sqrt{bx^2 + a}}{\sqrt{a}}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^2,x, algorithm="maxima")

[Out] $\frac{1}{6}*(b*x^2 + a)^{(3/2)}*d^2*x^3/b + \frac{1}{2}*\sqrt{b*x^2 + a}*c^2*x + \frac{1}{2}*(b*x^2 + a)^{(3/2)}*c*d*x/b - \frac{1}{4}*\sqrt{b*x^2 + a}*a*c*d*x/b - \frac{1}{8}*(b*x^2 + a)^{(3/2)}*a*d^2*x/b^2 + \frac{1}{16}*\sqrt{b*x^2 + a}*a^2*d^2*x/b^2 + \frac{1}{2}*a*c^2*\operatorname{arcsinh}(b*x/\sqrt{a*b})/\sqrt{b} - \frac{1}{4}*a^2*c*d*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{3/2} + \frac{1}{16}*a^3*d^2*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{5/2}$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{bx^2 + a} (dx^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)*(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(1/2)*(c + d*x^2)^2, x)

sympy [B] time = 11.39, size = 291, normalized size = 1.95

$$-\frac{a^{\frac{5}{2}} d^2 x}{16b^2 \sqrt{1 + \frac{bx^2}{a}}} + \frac{a^{\frac{3}{2}} cdx}{4b \sqrt{1 + \frac{bx^2}{a}}} - \frac{a^{\frac{3}{2}} d^2 x^3}{48b \sqrt{1 + \frac{bx^2}{a}}} + \frac{\sqrt{a} c^2 x \sqrt{1 + \frac{bx^2}{a}}}{2} + \frac{3\sqrt{a} cdx^3}{4 \sqrt{1 + \frac{bx^2}{a}}} + \frac{5\sqrt{a} d^2 x^5}{24 \sqrt{1 + \frac{bx^2}{a}}} + \frac{a^3 d^2 \operatorname{asinh}\left(\frac{\sqrt{bx^2 + a}}{\sqrt{a}}\right)}{16b^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)*(d*x**2+c)**2,x)

[Out] $-a^{(5/2)}*d^{**2}*x/(16*b^{**2}*\sqrt{1 + b*x^{**2}/a}) + a^{(3/2)}*c*d*x/(4*b*\sqrt{1 + b*x^{**2}/a}) - a^{(3/2)}*d^{**2}*x^{**3}/(48*b*\sqrt{1 + b*x^{**2}/a}) + \sqrt{a}*c^{**2}$

$$\begin{aligned}
& x\sqrt{1 + b*x**2/a}/2 + 3*\sqrt{a}*c*d*x**3/(4*\sqrt{1 + b*x**2/a}) + 5*\sqrt{a}*d**2*x**5/(24*\sqrt{1 + b*x**2/a}) + a**3*d**2*\operatorname{asinh}(\sqrt{b}*x/\sqrt{a})/(16*b**(5/2)) - a**2*c*d*\operatorname{asinh}(\sqrt{b}*x/\sqrt{a})/(4*b**(3/2)) + a*c**2*\operatorname{asinh}(\sqrt{b}*x/\sqrt{a})/(2*\sqrt{b}) + b*c*d*x**5/(2*\sqrt{a}*\sqrt{1 + b*x**2/a}) + b*d**2*x**7/(6*\sqrt{a}*\sqrt{1 + b*x**2/a})
\end{aligned}$$

3.47 $\int \sqrt{a + bx^2} (c + dx^2) dx$

Optimal. Leaf size=87

$$\frac{a(4bc - ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{8b^{3/2}} + \frac{x\sqrt{a+bx^2}(4bc - ad)}{8b} + \frac{dx(a+bx^2)^{3/2}}{4b}$$

[Out] $1/4*d*x*(b*x^2+a)^{(3/2)}/b+1/8*a*(-a*d+4*b*c)*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(3/2)}+1/8*(-a*d+4*b*c)*x*(b*x^2+a)^{(1/2)}/b$

Rubi [A] time = 0.03, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {388, 195, 217, 206}

$$\frac{a(4bc - ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{8b^{3/2}} + \frac{x\sqrt{a+bx^2}(4bc - ad)}{8b} + \frac{dx(a+bx^2)^{3/2}}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]*(c + d*x^2), x]

[Out] $((4*b*c - a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(8*b) + (d*x*(a + b*x^2)^{(3/2)})/(4*b) + (a*(4*b*c - a*d)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(8*b^{(3/2)})$

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rubi steps

$$\begin{aligned}
\int \sqrt{a+bx^2} (c+dx^2) dx &= \frac{dx(a+bx^2)^{3/2}}{4b} - \frac{(-4bc+ad) \int \sqrt{a+bx^2} dx}{4b} \\
&= \frac{(4bc-ad)x\sqrt{a+bx^2}}{8b} + \frac{dx(a+bx^2)^{3/2}}{4b} + \frac{(a(4bc-ad)) \int \frac{1}{\sqrt{a+bx^2}} dx}{8b} \\
&= \frac{(4bc-ad)x\sqrt{a+bx^2}}{8b} + \frac{dx(a+bx^2)^{3/2}}{4b} + \frac{(a(4bc-ad)) \operatorname{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{8b} \\
&= \frac{(4bc-ad)x\sqrt{a+bx^2}}{8b} + \frac{dx(a+bx^2)^{3/2}}{4b} + \frac{a(4bc-ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{8b^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.16, size = 85, normalized size = 0.98

$$\frac{\sqrt{a+bx^2} \left(\sqrt{b}x(ad+4bc+2bdx^2) - \frac{\sqrt{a(ad-4bc)} \sinh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{\frac{bx^2}{a}+1}} \right)}{8b^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2]*(c + d*x^2), x]

[Out] (Sqrt[a + b*x^2]*(Sqrt[b]*x*(4*b*c + a*d + 2*b*d*x^2) - (Sqrt[a]*(-4*b*c + a*d)*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/Sqrt[1 + (b*x^2)/a]))/(8*b^(3/2))

fricas [A] time = 0.49, size = 158, normalized size = 1.82

$$\left[\frac{(4abc - a^2d)\sqrt{b} \log\left(-2bx^2 + 2\sqrt{bx^2 + a}\sqrt{b}x - a\right) - 2(2b^2dx^3 + (4b^2c + abd)x)\sqrt{bx^2 + a}}{16b^2}, \dots \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c), x, algorithm="fricas")

[Out] [-1/16*((4*a*b*c - a^2*d)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(2*b^2*d*x^3 + (4*b^2*c + a*b*d)*x)*sqrt(b*x^2 + a))/b^2, -1/8*((4*a*b*c - a^2*d)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (2*b^2*d*x^3 + (4*b^2*c + a*b*d)*x)*sqrt(b*x^2 + a))/b^2]

giac [A] time = 0.60, size = 70, normalized size = 0.80

$$\frac{1}{8} \sqrt{bx^2 + a} \left(2dx^2 + \frac{4b^2c + abd}{b^2} \right) x - \frac{(4abc - a^2d) \log\left(\left| -\sqrt{b}x + \sqrt{bx^2 + a} \right| \right)}{8b^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c), x, algorithm="giac")

[Out] 1/8*sqrt(b*x^2 + a)*(2*d*x^2 + (4*b^2*c + a*b*d)/b^2)*x - 1/8*(4*a*b*c - a^2*d)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(3/2)

maple [A] time = 0.00, size = 96, normalized size = 1.10

$$-\frac{a^2d \ln\left(\sqrt{b}x + \sqrt{bx^2 + a}\right)}{8b^{3/2}} + \frac{ac \ln\left(\sqrt{b}x + \sqrt{bx^2 + a}\right)}{2\sqrt{b}} - \frac{\sqrt{bx^2 + a} adx}{8b} + \frac{\sqrt{bx^2 + a} cx}{2} + \frac{(bx^2 + a)^{3/2} dx}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2+a)^(1/2)*(d*x^2+c), x)`

[Out] $\frac{1}{4}d*x*(b*x^2+a)^{(3/2)}/b - \frac{1}{8}d*a/b*x*(b*x^2+a)^{(1/2)} - \frac{1}{8}d*a^2/b^{(3/2)}*\ln(b^{(1/2)*x+(b*x^2+a)^{(1/2)}}) + \frac{1}{2}c*x*(b*x^2+a)^{(1/2)} + \frac{1}{2}c*a/b^{(1/2)}*\ln(b^{(1/2)*x+(b*x^2+a)^{(1/2)})}$

maxima [A] time = 1.35, size = 81, normalized size = 0.93

$$\frac{1}{2}\sqrt{bx^2+a}cx + \frac{(bx^2+a)^{\frac{3}{2}}dx}{4b} - \frac{\sqrt{bx^2+a}adx}{8b} + \frac{ac \operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{b}} - \frac{a^2d \operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{8b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+a)^(1/2)*(d*x^2+c), x, algorithm="maxima")`

[Out] $\frac{1}{2}\sqrt{bx^2+a}cx + \frac{1}{4}(bx^2+a)^{(3/2)}dx/b - \frac{1}{8}\sqrt{bx^2+a}ax/b + \frac{1}{2}ac*\operatorname{arcsinh}(bx/\sqrt{a*b})/\sqrt{b} - \frac{1}{8}a^2d*\operatorname{arcsinh}(bx/\sqrt{a*b})/b^{(3/2)}$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{bx^2+a} (dx^2+c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^(1/2)*(c + d*x^2), x)`

[Out] `int((a + b*x^2)^(1/2)*(c + d*x^2), x)`

sympy [A] time = 5.68, size = 144, normalized size = 1.66

$$\frac{a^{\frac{3}{2}}dx}{8b\sqrt{1+\frac{bx^2}{a}}} + \frac{\sqrt{a}cx\sqrt{1+\frac{bx^2}{a}}}{2} + \frac{3\sqrt{a}dx^3}{8\sqrt{1+\frac{bx^2}{a}}} - \frac{a^2d \operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8b^{\frac{3}{2}}} + \frac{ac \operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2\sqrt{b}} + \frac{bdx^5}{4\sqrt{a}\sqrt{1+\frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**(1/2)*(d*x**2+c), x)`

[Out] $a^{(3/2)}*d*x/(8*b*\sqrt{1+b*x**2/a}) + \sqrt{a}*c*x*\sqrt{1+b*x**2/a}/2 + 3*\sqrt{a}*d*x**3/(8*\sqrt{1+b*x**2/a}) - a**2*d*\operatorname{asinh}(\sqrt{b}*x/\sqrt{a})/(8*b**(3/2)) + a*c*\operatorname{asinh}(\sqrt{b}*x/\sqrt{a})/(2*\sqrt{b}) + b*d*x**5/(4*\sqrt{a}*\sqrt{1+b*x**2/a})$

3.48 $\int \sqrt{a + bx^2} dx$

Optimal. Leaf size=46

$$\frac{1}{2}x\sqrt{a + bx^2} + \frac{a \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2\sqrt{b}}$$

[Out] $1/2*a*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(1/2)}+1/2*x*(b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {195, 217, 206}

$$\frac{1}{2}x\sqrt{a + bx^2} + \frac{a \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2], x]

[Out] $(x*\operatorname{Sqrt}[a + b*x^2])/2 + (a*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(2*\operatorname{Sqrt}[b])$

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[Rt[-b, 2]*x]/Rt[a, 2])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \sqrt{a + bx^2} dx &= \frac{1}{2}x\sqrt{a + bx^2} + \frac{1}{2}a \int \frac{1}{\sqrt{a + bx^2}} dx \\ &= \frac{1}{2}x\sqrt{a + bx^2} + \frac{1}{2}a \operatorname{Subst}\left(\int \frac{1}{1 - bx^2} dx, x, \frac{x}{\sqrt{a + bx^2}}\right) \\ &= \frac{1}{2}x\sqrt{a + bx^2} + \frac{a \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2\sqrt{b}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 49, normalized size = 1.07

$$\frac{1}{2}x\sqrt{a + bx^2} + \frac{a \log\left(\sqrt{b}\sqrt{a + bx^2} + bx\right)}{2\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2], x]

[Out] (x*Sqrt[a + b*x^2])/2 + (a*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(2*Sqrt[b])

fricas [A] time = 0.55, size = 94, normalized size = 2.04

$$\left[\frac{2\sqrt{bx^2+a}bx + a\sqrt{b}\log\left(-2bx^2 - 2\sqrt{bx^2+a}\sqrt{b}x - a\right)}{4b}, \frac{\sqrt{bx^2+a}bx - a\sqrt{-b}\arctan\left(\frac{\sqrt{-b}x}{\sqrt{bx^2+a}}\right)}{2b} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b*x^2 + a)*b*x + a*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a))/b, 1/2*(sqrt(b*x^2 + a)*b*x - a*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)))/b]

giac [A] time = 0.58, size = 37, normalized size = 0.80

$$\frac{1}{2}\sqrt{bx^2+ax} - \frac{a\log\left(|-\sqrt{b}x + \sqrt{bx^2+a}|\right)}{2\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2), x, algorithm="giac")

[Out] 1/2*sqrt(b*x^2 + a)*x - 1/2*a*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/sqrt(b)

maple [A] time = 0.00, size = 36, normalized size = 0.78

$$\frac{a\ln\left(\sqrt{b}x + \sqrt{bx^2+a}\right)}{2\sqrt{b}} + \frac{\sqrt{bx^2+ax}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2), x)

[Out] 1/2*a/b^(1/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+1/2*(b*x^2+a)^(1/2)*x

maxima [A] time = 1.36, size = 28, normalized size = 0.61

$$\frac{1}{2}\sqrt{bx^2+ax} + \frac{a\operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{2\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2), x, algorithm="maxima")

[Out] 1/2*sqrt(b*x^2 + a)*x + 1/2*a*arcsinh(b*x/sqrt(a*b))/sqrt(b)

mupad [B] time = 4.71, size = 35, normalized size = 0.76

$$\frac{x\sqrt{bx^2+a}}{2} + \frac{a\ln\left(\sqrt{b}x + \sqrt{bx^2+a}\right)}{2\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^(1/2),x)`

[Out] $(x*(a + b*x^2)^{(1/2)})/2 + (a*\log(b^{(1/2)}*x + (a + b*x^2)^{(1/2)}))/(2*b^{(1/2)})$

sympy [A] time = 1.86, size = 41, normalized size = 0.89

$$\frac{\sqrt{a} x \sqrt{1 + \frac{bx^2}{a}}}{2} + \frac{a \operatorname{asinh}\left(\frac{\sqrt{b} x}{\sqrt{a}}\right)}{2\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**(1/2),x)`

[Out] $\sqrt{a}*x*\sqrt{1 + b*x**2/a}/2 + a*\operatorname{asinh}(\sqrt{b}*x/\sqrt{a})/(2*\sqrt{b})$

$$3.49 \quad \int \frac{\sqrt{a+bx^2}}{c+dx^2} dx$$

Optimal. Leaf size=82

$$\frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{d} - \frac{\sqrt{bc-ad} \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d}$$

[Out] $\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})*b^{(1/2)}/d - \operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})*(-a*d+b*c)^{(1/2)}/d/c^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {402, 217, 206, 377, 208}

$$\frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{d} - \frac{\sqrt{bc-ad} \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Sqrt}[a + b*x^2]/(c + d*x^2), x]$

[Out] $(\operatorname{Sqrt}[b]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/d - (\operatorname{Sqrt}[b*c - a*d]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(\operatorname{Sqrt}[c]*d)$

Rule 206

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1*\operatorname{ArcTanh}[(\operatorname{Rt}[-b, 2]*x)/\operatorname{Rt}[a, 2]])/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]), x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 208

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-(a/b), 2]*\operatorname{ArcTanh}[x/\operatorname{Rt}[-(a/b), 2]])/a, x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b]$

Rule 217

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_)*(x_)^2)], x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 377

$\operatorname{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}/((c_ + (d_)*(x_)^{(n_)})), x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^{(1/n)}] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \ \&\& \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \operatorname{EqQ}[n*p + 1, 0] \ \&\& \operatorname{IntegerQ}[n]$

Rule 402

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{(p_)} / ((c_ + (d_)*(x_)^2)), x_Symbol] \rightarrow \operatorname{Dist}[b/d, \operatorname{Int}[(a + b*x^2)^{(p-1)}, x], x] - \operatorname{Dist}[(b*c - a*d)/d, \operatorname{Int}[(a + b*x^2)^{(p-1)}/(c + d*x^2), x], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \ \&\& \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \operatorname{GtQ}[p, 0] \ \&\& (\operatorname{EqQ}[p, 1/2] \ || \ \operatorname{EqQ}[\operatorname{Denominator}[p], 4])$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a+bx^2}}{c+dx^2} dx &= \frac{b \int \frac{1}{\sqrt{a+bx^2}} dx}{d} - \frac{(bc-ad) \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{d} \\ &= \frac{b \operatorname{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{d} - \frac{(bc-ad) \operatorname{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{d} \\ &= \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{d} - \frac{\sqrt{bc-ad} \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d} \end{aligned}$$

Mathematica [A] time = 0.04, size = 84, normalized size = 1.02

$$\frac{\sqrt{ad-bc} \tan^{-1}\left(\frac{x\sqrt{ad-bc}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d} + \frac{\sqrt{b} \log\left(\sqrt{b}\sqrt{a+bx^2} + bx\right)}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2]/(c + d*x^2), x]

[Out] (Sqrt[-(b*c) + a*d]*ArcTan[(Sqrt[-(b*c) + a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(Sqrt[c]*d) + (Sqrt[b]*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/d

fricas [A] time = 0.70, size = 596, normalized size = 7.27

$$\left[\frac{2\sqrt{b} \log\left(-2bx^2 - 2\sqrt{bx^2+a}\sqrt{b}x - a\right) + \sqrt{\frac{bc-ad}{c}} \log\left(\frac{(8b^2c^2-8abcd+a^2d^2)x^4+a^2c^2+2(4abc^2-3a^2cd)x^2-4(ac^2x+(2bc^2-acd))}{d^2x^4+2cdx^2+c^2}\right)}{4d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*sqrt(b*x^2 + a)*sqrt((b*c - a*d)/c))/(d^2*x^4 + 2*c*d*x^2 + c^2))/d, -1/4*(4*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*sqrt(b*x^2 + a)*sqrt((b*c - a*d)/c))/(d^2*x^4 + 2*c*d*x^2 + c^2))/d, 1/2*(sqrt(-(b*c - a*d)/c)*arctan(1/2*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)*sqrt(-(b*c - a*d)/c)/((b^2*c - a*b*d)*x^3 + (a*b*c - a^2*d)*x)) + sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a)/d, -1/2*(2*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - sqrt(-(b*c - a*d)/c)*arctan(1/2*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)*sqrt(-(b*c - a*d)/c)/((b^2*c - a*b*d)*x^3 + (a*b*c - a^2*d)*x)))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

sage0*x

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c), x, algorithm="giac")

[Out] sage0*x

maple [B] time = 0.04, size = 932, normalized size = 11.37

$$\frac{a \ln \left(\frac{2\sqrt{-cd} \left(x - \frac{\sqrt{-cd}}{d} \right)^b + \frac{2ad-2bc}{d} + 2\sqrt{\frac{ad-bc}{d}} \sqrt{\left(x - \frac{\sqrt{-cd}}{d} \right)^2 + \frac{2\sqrt{-cd} \left(x - \frac{\sqrt{-cd}}{d} \right)^b + \frac{ad-bc}{d}}}{x - \frac{\sqrt{-cd}}{d}} \right)}{2\sqrt{-cd} \sqrt{\frac{ad-bc}{d}}} + \frac{a \ln \left(\frac{2\sqrt{-cd} \left(x + \frac{\sqrt{-cd}}{d} \right)^b + \frac{2ad-2bc}{d} + 2\sqrt{\frac{ad-bc}{d}} \sqrt{\left(x + \frac{\sqrt{-cd}}{d} \right)^2 + \frac{2\sqrt{-cd} \left(x + \frac{\sqrt{-cd}}{d} \right)^b + \frac{ad-bc}{d}}}{x + \frac{\sqrt{-cd}}{d}} \right)}{2\sqrt{-cd} \sqrt{\frac{ad-bc}{d}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)/(d*x^2+c), x)

[Out] $\frac{1}{2}(-c*d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*b*(-c*d)^{(1/2)}/d*(x-(-c*d)^{(1/2)}/d)} + (a*d-b*c)/d)^{(1/2)} + \frac{1}{2}*b^{(1/2)}/d*\ln((b*(-c*d)^{(1/2)}/d+(x-(-c*d)^{(1/2)}/d)*b)/b^{(1/2)} + ((x-(-c*d)^{(1/2)}/d)^{2*b+2*b*(-c*d)^{(1/2)}/d*(x-(-c*d)^{(1/2)}/d)} + (a*d-b*c)/d)^{(1/2)} - \frac{1}{2}(-c*d)^{(1/2)}/((a*d-b*c)/d)^{(1/2)}*\ln((2*(a*d-b*c)/d+2*b*(-c*d)^{(1/2)}/d*(x-(-c*d)^{(1/2)}/d)+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*b*(-c*d)^{(1/2)}/d*(x-(-c*d)^{(1/2)}/d)} + (a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d)*a + \frac{1}{2}(-c*d)^{(1/2)}/d/((a*d-b*c)/d)^{(1/2)}*\ln((2*(a*d-b*c)/d+2*b*(-c*d)^{(1/2)}/d*(x-(-c*d)^{(1/2)}/d)+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*b*(-c*d)^{(1/2)}/d*(x-(-c*d)^{(1/2)}/d)} + (a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d)*b*c - \frac{1}{2}(-c*d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*b*(-c*d)^{(1/2)}/d*(x+(-c*d)^{(1/2)}/d)} + (a*d-b*c)/d)^{(1/2)} + \frac{1}{2}*b^{(1/2)}/d*\ln((-b*(-c*d)^{(1/2)}/d+(x+(-c*d)^{(1/2)}/d)*b)/b^{(1/2)} + ((x+(-c*d)^{(1/2)}/d)^{2*b-2*b*(-c*d)^{(1/2)}/d*(x+(-c*d)^{(1/2)}/d)} + (a*d-b*c)/d)^{(1/2)} + \frac{1}{2}/(-c*d)^{(1/2)}/((a*d-b*c)/d)^{(1/2)}*\ln((2*(a*d-b*c)/d-2*b*(-c*d)^{(1/2)}/d*(x+(-c*d)^{(1/2)}/d)+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*b*(-c*d)^{(1/2)}/d*(x+(-c*d)^{(1/2)}/d)} + (a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d)*a - \frac{1}{2}/(-c*d)^{(1/2)}/d/((a*d-b*c)/d)^{(1/2)}*\ln((2*(a*d-b*c)/d-2*b*(-c*d)^{(1/2)}/d*(x+(-c*d)^{(1/2)}/d)+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*b*(-c*d)^{(1/2)}/d*(x+(-c*d)^{(1/2)}/d)} + (a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d)*b*c$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{dx^2+c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c), x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2+a)/(d*x^2+c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\left\{ \begin{array}{ll} \frac{\sqrt{-b} \operatorname{asin}\left(x \sqrt{\frac{-b}{a}}\right)}{c} & \text{if } ((a+bc=0 \wedge d=-1) \vee ad=bc) \wedge b < 0 \\ \frac{\sqrt{b} \ln\left(2\sqrt{b}x+2\sqrt{bx^2+a}\right)}{d} + \frac{\operatorname{atan}\left(\frac{x\sqrt{ad-bc}}{\sqrt{c}\sqrt{bx^2+a}}\right)\sqrt{ad-bc}}{\sqrt{c}d} & \text{if } a \neq 0 \wedge (((a+bc \neq 0 \vee d \neq -1) \wedge ad \neq bc) \vee -b < 0) \\ \int \frac{\sqrt{bx^2+a}}{dx^2+c} dx & \text{if } (((a+bc=0 \wedge d=-1) \vee ad=bc) \wedge b < 0) \vee a = 0 \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+b*x^2)^(1/2)/(c+d*x^2), x)

[Out] piecewise((a+b*c == 0 & d == -1 | a*d == b*c) & b < 0, ((-b)^(1/2)*asin(x*(-b/a)^(1/2)))/c, a ~= 0 & ((a+b*c ~= 0 | d ~= -1) & a*d ~= b*c | ~b < 0

), (b^(1/2)*log(2*b^(1/2)*x + 2*(a + b*x^2)^(1/2))/d + (atan((x*(a*d - b*c)^(1/2))/(c^(1/2)*(a + b*x^2)^(1/2)))*(a*d - b*c)^(1/2))/(c^(1/2)*d), ((a + b*c == 0 & d == -1 | a*d == b*c) & b < 0 | a == 0) & ((a + b*c != 0 | d != -1) & a*d != b*c | ~b < 0), int((a + b*x^2)^(1/2)/(c + d*x^2), x))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a + bx^2}}{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)/(d*x**2+c),x)

[Out] Integral(sqrt(a + b*x**2)/(c + d*x**2), x)

$$3.50 \quad \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^2} dx$$

Optimal. Leaf size=82

$$\frac{a \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}\sqrt{bc-ad}} + \frac{x\sqrt{a+bx^2}}{2c(c+dx^2)}$$

[Out] $1/2*a*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})/c^{(3/2)}/(-a*d+b*c)^{(1/2)}+1/2*x*(b*x^2+a)^{(1/2)}/c/(d*x^2+c)$

Rubi [A] time = 0.03, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {378, 377, 208}

$$\frac{a \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}\sqrt{bc-ad}} + \frac{x\sqrt{a+bx^2}}{2c(c+dx^2)}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[a + b*x^2]/(c + d*x^2)^2, x]`

[Out] $(x*\operatorname{Sqrt}[a + b*x^2])/(2*c*(c + d*x^2)) + (a*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(2*c^{(3/2)}*\operatorname{Sqrt}[b*c - a*d])$

Rule 208

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 377

`Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]`

Rule 378

`Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] - Dist[(c*q)/(a*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+1)+1, 0] && GtQ[q, 0] && NeQ[p, -1]`

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^2} dx &= \frac{x\sqrt{a+bx^2}}{2c(c+dx^2)} + \frac{a \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{2c} \\ &= \frac{x\sqrt{a+bx^2}}{2c(c+dx^2)} + \frac{a \operatorname{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{2c} \\ &= \frac{x\sqrt{a+bx^2}}{2c(c+dx^2)} + \frac{a \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}\sqrt{bc-ad}} \end{aligned}$$

Mathematica [B] time = 0.23, size = 165, normalized size = 2.01

$$\frac{x\sqrt{a+bx^2} \left(\sqrt{x^2 \left(\frac{d}{c} - \frac{b}{a} \right)} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} + \sqrt{\frac{dx^2}{c} + 1} \sin^{-1} \left(\frac{\sqrt{x^2 \left(\frac{d}{c} - \frac{b}{a} \right)}}{\sqrt{\frac{dx^2}{c} + 1}} \right) \right)}{2c^2 \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{dx^2}{c} + 1} \sqrt{x^2 \left(\frac{d}{c} - \frac{b}{a} \right)}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Sqrt[a + b*x^2]/(c + d*x^2)^2, x]

[Out] (x*Sqrt[a + b*x^2]*(Sqrt[(-b/a) + d/c]*x^2)*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2)]) + Sqrt[1 + (d*x^2)/c]*ArcSin[Sqrt[(-b/a) + d/c]*x^2/Sqrt[1 + (d*x^2)/c]])/(2*c^2*Sqrt[(-b/a) + d/c]*x^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c])

fricas [B] time = 0.68, size = 369, normalized size = 4.50

$$\frac{4(bc^2 - acd)\sqrt{bx^2 + a}x + (adx^2 + ac)\sqrt{bc^2 - acd} \log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4 + a^2c^2 + 2(4abc^2 - 3a^2cd)x^2 + 4((2bc - ad)x^3 + acx)\sqrt{bx^2 + a}}{d^2x^4 + 2cdx^2 + c^2}\right)}{8(bc^4 - ac^3d + (bc^3d - ac^2d^2)x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^2, x, algorithm="fricas")

[Out] [1/8*(4*(b*c^2 - a*c*d)*sqrt(b*x^2 + a)*x + (a*d*x^2 + a*c)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)))/(b*c^4 - a*c^3*d + (b*c^3*d - a*c^2*d^2)*x^2), 1/4*(2*(b*c^2 - a*c*d)*sqrt(b*x^2 + a)*x - (a*d*x^2 + a*c)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)))/(b*c^4 - a*c^3*d + (b*c^3*d - a*c^2*d^2)*x^2)]

giac [B] time = 1.64, size = 217, normalized size = 2.65

$$\frac{a\sqrt{b} \arctan\left(\frac{(\sqrt{b}x - \sqrt{bx^2+a})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{2\sqrt{-b^2c^2 + abcd}c} + \frac{2\left(\sqrt{b}x - \sqrt{bx^2+a}\right)^2 b^{\frac{3}{2}}c - \left(\sqrt{b}x - \sqrt{bx^2+a}\right)^2 a\sqrt{b}d + a^2\sqrt{b}}{\left(\left(\sqrt{b}x - \sqrt{bx^2+a}\right)^4 d + 4\left(\sqrt{b}x - \sqrt{bx^2+a}\right)^2 bc - 2\left(\sqrt{b}x - \sqrt{bx^2+a}\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

$$\frac{1}{(x+(-c*d)^{(1/2)/d})} * a^{-1/4} / (-c*d)^{(1/2)/d} / ((a*d-b*c)/d)^{(1/2)} * \ln((-2*(-c*d)^{(1/2)} * (x+(-c*d)^{(1/2)/d}) * b/d + 2*(a*d-b*c)/d + 2*((a*d-b*c)/d)^{(1/2)} * ((x+(-c*d)^{(1/2)/d})^2 * b - 2*(-c*d)^{(1/2)} * (x+(-c*d)^{(1/2)/d}) * b/d + (a*d-b*c)/d)^{(1/2)}) / (x+(-c*d)^{(1/2)/d}) * b$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + a}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)/(d*x^2 + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 + a}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)/(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(1/2)/(c + d*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a + bx^2}}{(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)/(d*x**2+c)**2,x)

[Out] Integral(sqrt(a + b*x**2)/(c + d*x**2)**2, x)

$$3.51 \quad \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^3} dx$$

Optimal. Leaf size=149

$$\frac{a(4bc - 3ad) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}(bc - ad)^{3/2}} + \frac{x\sqrt{a+bx^2}(4bc - 3ad)}{8c^2(c+dx^2)(bc - ad)} - \frac{dx(a+bx^2)^{3/2}}{4c(c+dx^2)^2(bc - ad)}$$

[Out] $-1/4*d*x*(b*x^2+a)^{(3/2)}/c/(-a*d+b*c)/(d*x^2+c)^2+1/8*a*(-3*a*d+4*b*c)*\arctanh(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})/c^{(5/2)}/(-a*d+b*c)^{(3/2)}+1/8*(-3*a*d+4*b*c)*x*(b*x^2+a)^{(1/2)}/c^2/(-a*d+b*c)/(d*x^2+c)$

Rubi [A] time = 0.09, antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {382, 378, 377, 208}

$$\frac{x\sqrt{a+bx^2}(4bc - 3ad)}{8c^2(c+dx^2)(bc - ad)} + \frac{a(4bc - 3ad) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}(bc - ad)^{3/2}} - \frac{dx(a+bx^2)^{3/2}}{4c(c+dx^2)^2(bc - ad)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]/(c + d*x^2)^3, x]

[Out] $-(d*x*(a + b*x^2)^{(3/2)})/(4*c*(b*c - a*d)*(c + d*x^2)^2) + ((4*b*c - 3*a*d)*x*\text{Sqrt}[a + b*x^2])/(8*c^2*(b*c - a*d)*(c + d*x^2)) + (a*(4*b*c - 3*a*d)*\text{ArcTanh}[(\text{Sqrt}[b*c - a*d]*x)/(\text{Sqrt}[c]*\text{Sqrt}[a + b*x^2])])/(8*c^{(5/2)}*(b*c - a*d)^{(3/2)})$

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] - Dist[(c*q)/(a*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+1) + 1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+2) + 1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{a+bx^2}}{(c+dx^2)^3} dx &= -\frac{dx(a+bx^2)^{3/2}}{4c(bc-ad)(c+dx^2)^2} + \frac{(4bc-3ad) \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^2} dx}{4c(bc-ad)} \\
&= -\frac{dx(a+bx^2)^{3/2}}{4c(bc-ad)(c+dx^2)^2} + \frac{(4bc-3ad)x\sqrt{a+bx^2}}{8c^2(bc-ad)(c+dx^2)} + \frac{(a(4bc-3ad)) \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{8c^2(bc-ad)} \\
&= -\frac{dx(a+bx^2)^{3/2}}{4c(bc-ad)(c+dx^2)^2} + \frac{(4bc-3ad)x\sqrt{a+bx^2}}{8c^2(bc-ad)(c+dx^2)} + \frac{(a(4bc-3ad)) \text{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \right)}{8c^2(bc-ad)} \\
&= -\frac{dx(a+bx^2)^{3/2}}{4c(bc-ad)(c+dx^2)^2} + \frac{(4bc-3ad)x\sqrt{a+bx^2}}{8c^2(bc-ad)(c+dx^2)} + \frac{a(4bc-3ad) \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}(bc-ad)^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.57, size = 176, normalized size = 1.18

$$\frac{x \left(c \left(a^2 d (5c + 3dx^2) + ab(-4c^2 + 3cdx^2 + 3d^2x^4) - 2b^2cx^2(2c + dx^2) \right) + \frac{a(c+dx^2)^2(3ad-4bc) \tanh^{-1}\left(\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}\right)}{\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}} \right)}{8c^3\sqrt{a+bx^2}(c+dx^2)^2(ad-bc)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2]/(c + d*x^2)^3,x]

[Out] (x*(c*(-2*b^2*c*x^2*(2*c + d*x^2) + a^2*d*(5*c + 3*d*x^2) + a*b*(-4*c^2 + 3*c*d*x^2 + 3*d^2*x^4)) + (a*(-4*b*c + 3*a*d))*(c + d*x^2)^2*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2)))]/(8*c^3*(-(b*c) + a*d)*Sqrt[a + b*x^2]*(c + d*x^2)^2)

fricas [B] time = 1.41, size = 698, normalized size = 4.68

$$\left[\frac{(4abc^3 - 3a^2c^2d + (4abcd^2 - 3a^2d^3)x^4 + 2(4abc^2d - 3a^2cd^2)x^2)\sqrt{bc^2 - acd} \log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4 + a^2c^2 + 2(4abcd - 3a^2d^3)x^2 + a^2c^2}{32(b^2c^7 - 2abc^6d + a^2c^5d^2 + (b^2c^5d^2 - 2abcd^2 - a^2c^4d^2)x^2 + a^2c^4d^2)}\right)}{32(b^2c^7 - 2abc^6d + a^2c^5d^2 + (b^2c^5d^2 - 2abcd^2 - a^2c^4d^2)x^2 + a^2c^4d^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [1/32*((4*a*b*c^3 - 3*a^2*c^2*d + (4*a*b*c*d^2 - 3*a^2*d^3)*x^4 + 2*(4*a*b*c^2*d - 3*a^2*c*d^2)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*((2*b^2*c^3*d - 5*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^3 + (4*b^2*c^4 - 9*a*b*c^3*d + 5*a^2*c^2*d^2)*x)*sqrt(b*x^2 + a))/(b^2*c^7 - 2*a*b*c^6*d + a^2*c^5*d^2 + (b^2*c^5*d^2 - 2*a*b*c^4*d^3 + a^2*c^3*d^4)*x^4 + 2*(b^2*c^6*d - 2*a*b*c^5*d^2 + a^2*c^4*d^3)*x^2), -1/16*((4*a*b*c^3 - 3*a^2*c^2*d + (4*a*b*c*d^2 - 3*a^2*d^3)*x^4 + 2*(4*a*b*c^2*d - 3*a^2*c*d^2)*x^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a))/(b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x) - 2*((2*b^2*c^3*d

$$- 5*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^3 + (4*b^2*c^4 - 9*a*b*c^3*d + 5*a^2*c^2*d^2)*x)*\sqrt{b*x^2 + a})/(b^2*c^7 - 2*a*b*c^6*d + a^2*c^5*d^2 + (b^2*c^5*d^2 - 2*a*b*c^4*d^3 + a^2*c^3*d^4)*x^4 + 2*(b^2*c^6*d - 2*a*b*c^5*d^2 + a^2*c^4*d^3)*x^2)]$$

giac [B] time = 3.75, size = 487, normalized size = 3.27

$$\frac{\left(4ab^{\frac{3}{2}}c - 3a^2\sqrt{bd}\right)\arctan\left(\frac{\left(\sqrt{bx-\sqrt{bx^2+a}}\right)^{d+2bc-ad}}{2\sqrt{-b^2c^2+abcd}}\right)}{8\sqrt{-b^2c^2+abcd}(bc^3-ac^2d)} - \frac{4\left(\sqrt{bx-\sqrt{bx^2+a}}\right)^6ab^{\frac{3}{2}}cd^2 - 3\left(\sqrt{bx-\sqrt{bx^2+a}}\right)^6a^{\frac{3}{2}}d^2}{8\sqrt{-b^2c^2+abcd}(bc^3-ac^2d)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^3,x, algorithm="giac")

[Out] $-1/8*(4*a*b^{(3/2)*c} - 3*a^2*\sqrt{b}*d)*\arctan(1/2*((\sqrt{b}*x - \sqrt{b*x^2 + a})^2*d + 2*b*c - a*d)/\sqrt{-b^2*c^2 + a*b*c*d})/(\sqrt{-b^2*c^2 + a*b*c*d})*(b*c^3 - a*c^2*d) - 1/4*(4*(\sqrt{b}*x - \sqrt{b*x^2 + a})^6*a*b^{(3/2)*c*d^2} - 3*(\sqrt{b}*x - \sqrt{b*x^2 + a})^6*a^2*\sqrt{b}*d^3 - 16*(\sqrt{b}*x - \sqrt{b*x^2 + a})^4*b^{(7/2)*c^3} + 40*(\sqrt{b}*x - \sqrt{b*x^2 + a})^4*a*b^{(5/2)*c^2*d} - 30*(\sqrt{b}*x - \sqrt{b*x^2 + a})^4*a^2*b^{(3/2)*c*d^2} + 9*(\sqrt{b}*x - \sqrt{b*x^2 + a})^4*a^3*\sqrt{b}*d^3 - 16*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*a^2*b^{(5/2)*c^2*d} + 28*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*a^3*b^{(3/2)*c*d^2} - 9*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*a^4*\sqrt{b}*d^3 - 2*a^4*b^{(3/2)*c*d^2} + 3*a^5*\sqrt{b}*d^3)/((\sqrt{b}*x - \sqrt{b*x^2 + a})^4*d + 4*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*b*c - 2*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*a*d + a^2*d)^2*(b*c^3*d - a*c^2*d^2))$

maple [B] time = 0.02, size = 5101, normalized size = 34.23

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)/(d*x^2+c)^3,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + a}}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^3,x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)/(d*x^2 + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 + a}}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)/(c + d*x^2)^3,x)

[Out] int((a + b*x^2)^(1/2)/(c + d*x^2)^3, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)/(d*x**2+c)**3,x)

[Out] Timed out

$$3.52 \quad \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^4} dx$$

Optimal. Leaf size=208

$$\frac{a(5a^2d^2 - 12abcd + 8b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{16c^{7/2}(bc-ad)^{5/2}} + \frac{x\sqrt{a+bx^2}(2bc-5ad)(4bc-3ad)}{48c^3(c+dx^2)(bc-ad)^2} + \frac{x\sqrt{a+bx^2}(4bc-5ad)}{24c^2(c+dx^2)^2(bc-ad)}$$

[Out] 1/16*a*(5*a^2*d^2-12*a*b*c*d+8*b^2*c^2)*arctanh(x*(-a*d+b*c)^(1/2)/c^(1/2)/(b*x^2+a)^(1/2))/c^(7/2)/(-a*d+b*c)^(5/2)+1/6*x*(b*x^2+a)^(1/2)/c/(d*x^2+c)^(3+1/24*(-5*a*d+4*b*c))*x*(b*x^2+a)^(1/2)/c^2/(-a*d+b*c)/(d*x^2+c)^2+1/48*(-5*a*d+2*b*c)*(-3*a*d+4*b*c)*x*(b*x^2+a)^(1/2)/c^3/(-a*d+b*c)^2/(d*x^2+c)

Rubi [A] time = 0.21, antiderivative size = 208, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {412, 527, 12, 377, 208}

$$\frac{a(5a^2d^2 - 12abcd + 8b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{16c^{7/2}(bc-ad)^{5/2}} + \frac{x\sqrt{a+bx^2}(2bc-5ad)(4bc-3ad)}{48c^3(c+dx^2)(bc-ad)^2} + \frac{x\sqrt{a+bx^2}(4bc-5ad)}{24c^2(c+dx^2)^2(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]/(c + d*x^2)^4, x]

[Out] (x*Sqrt[a + b*x^2])/(6*c*(c + d*x^2)^3) + ((4*b*c - 5*a*d)*x*Sqrt[a + b*x^2])/(24*c^2*(b*c - a*d)*(c + d*x^2)^2) + ((2*b*c - 5*a*d)*(4*b*c - 3*a*d)*x*Sqrt[a + b*x^2])/(48*c^3*(b*c - a*d)^2*(c + d*x^2)) + (a*(8*b^2*c^2 - 12*a*b*c*d + 5*a^2*d^2)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(16*c^(7/2)*(b*c - a*d)^(5/2))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 412

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] + Dist[1/(a*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1)*Simp[c*(n*(p+1) + 1) + d*(n*(p+q+1) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && LtQ[0, q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\int \frac{\sqrt{a+bx^2}}{(c+dx^2)^4} dx = \frac{x\sqrt{a+bx^2}}{6c(c+dx^2)^3} - \frac{\int \frac{-5a-4bx^2}{\sqrt{a+bx^2}(c+dx^2)^3} dx}{6c}$$

$$= \frac{x\sqrt{a+bx^2}}{6c(c+dx^2)^3} + \frac{(4bc-5ad)x\sqrt{a+bx^2}}{24c^2(bc-ad)(c+dx^2)^2} - \frac{\int \frac{-a(16bc-15ad)-2b(4bc-5ad)x^2}{\sqrt{a+bx^2}(c+dx^2)^2} dx}{24c^2(bc-ad)}$$

$$= \frac{x\sqrt{a+bx^2}}{6c(c+dx^2)^3} + \frac{(4bc-5ad)x\sqrt{a+bx^2}}{24c^2(bc-ad)(c+dx^2)^2} + \frac{(2bc-5ad)(4bc-3ad)x\sqrt{a+bx^2}}{48c^3(bc-ad)^2(c+dx^2)} - \frac{\int -\frac{3a(8b^2c^2-1)}{\sqrt{a+bx^2}} dx}{48c^3(bc-ad)^2}$$

$$= \frac{x\sqrt{a+bx^2}}{6c(c+dx^2)^3} + \frac{(4bc-5ad)x\sqrt{a+bx^2}}{24c^2(bc-ad)(c+dx^2)^2} + \frac{(2bc-5ad)(4bc-3ad)x\sqrt{a+bx^2}}{48c^3(bc-ad)^2(c+dx^2)} + \frac{a(8b^2c^2-1)}{48c^3(bc-ad)^2}$$

$$= \frac{x\sqrt{a+bx^2}}{6c(c+dx^2)^3} + \frac{(4bc-5ad)x\sqrt{a+bx^2}}{24c^2(bc-ad)(c+dx^2)^2} + \frac{(2bc-5ad)(4bc-3ad)x\sqrt{a+bx^2}}{48c^3(bc-ad)^2(c+dx^2)} + \frac{a(8b^2c^2-1)}{48c^3(bc-ad)^2}$$

$$= \frac{x\sqrt{a+bx^2}}{6c(c+dx^2)^3} + \frac{(4bc-5ad)x\sqrt{a+bx^2}}{24c^2(bc-ad)(c+dx^2)^2} + \frac{(2bc-5ad)(4bc-3ad)x\sqrt{a+bx^2}}{48c^3(bc-ad)^2(c+dx^2)} + \frac{a(8b^2c^2-1)}{48c^3(bc-ad)^2}$$

Mathematica [A] time = 0.99, size = 227, normalized size = 1.09

$$x\sqrt{a+bx^2} \left(\frac{3a(c+dx^2)^3(5a^2d^2-12abcd+8b^2c^2)\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}} \tanh^{-1}\left(\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}\right)}{x^2} + (bc-ad)(a^2d^2(33c^2+40cdx^2+15d^2x^4) - 2a^2d^2) \right) / (48c^3(c+dx^2)^3(bc-ad)^3)$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[a + b*x^2]/(c + d*x^2)^4, x]
[Out] (x*Sqrt[a + b*x^2]*((b*c - a*d)*(8*b^2*c^2*(3*c^2 + 3*c*d*x^2 + d^2*x^4) -
2*a*b*c*d*(30*c^2 + 35*c*d*x^2 + 13*d^2*x^4) + a^2*d^2*(33*c^2 + 40*c*d*x^2
+ 15*d^2*x^4)) + (3*a*(8*b^2*c^2 - 12*a*b*c*d + 5*a^2*d^2)*Sqrt[((b*c - a*
d)*x^2)/(c*(a + b*x^2))])*(c + d*x^2)^3*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a
+ b*x^2))]])/x^2)/(48*c^3*(b*c - a*d)^3*(c + d*x^2)^3)
```

fricas [B] time = 1.64, size = 1220, normalized size = 5.87

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^4,x, algorithm="fricas")

[Out] [1/192*(3*(8*a*b^2*c^5 - 12*a^2*b*c^4*d + 5*a^3*c^3*d^2 + (8*a*b^2*c^2*d^3 - 12*a^2*b*c*d^4 + 5*a^3*d^5))*x^6 + 3*(8*a*b^2*c^3*d^2 - 12*a^2*b*c^2*d^3 + 5*a^3*c*d^4)*x^4 + 3*(8*a*b^2*c^4*d - 12*a^2*b*c^3*d^2 + 5*a^3*c^2*d^3)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*((8*b^3*c^4*d^2 - 34*a*b^2*c^3*d^3 + 41*a^2*b*c^2*d^4 - 15*a^3*c*d^5))*x^5 + 2*(12*b^3*c^5*d - 47*a*b^2*c^4*d^2 + 55*a^2*b*c^3*d^3 - 20*a^3*c^2*d^4)*x^3 + 3*(8*b^3*c^6 - 28*a*b^2*c^5*d + 31*a^2*b*c^4*d^2 - 11*a^3*c^3*d^3)*x)*sqrt(b*x^2 + a))/(b^3*c^10 - 3*a*b^2*c^9*d + 3*a^2*b*c^8*d^2 - a^3*c^7*d^3 + (b^3*c^7*d^3 - 3*a*b^2*c^6*d^4 + 3*a^2*b*c^5*d^5 - a^3*c^4*d^6)*x^6 + 3*(b^3*c^8*d^2 - 3*a*b^2*c^7*d^3 + 3*a^2*b*c^6*d^4 - a^3*c^5*d^5)*x^4 + 3*(b^3*c^9*d - 3*a*b^2*c^8*d^2 + 3*a^2*b*c^7*d^3 - a^3*c^6*d^4)*x^2), -1/96*(3*(8*a*b^2*c^5 - 12*a^2*b*c^4*d + 5*a^3*c^3*d^2 + (8*a*b^2*c^2*d^3 - 12*a^2*b*c*d^4 + 5*a^3*d^5))*x^6 + 3*(8*a*b^2*c^3*d^2 - 12*a^2*b*c^2*d^3 + 5*a^3*c*d^4)*x^4 + 3*(8*a*b^2*c^4*d - 12*a^2*b*c^3*d^2 + 5*a^3*c^2*d^3)*x^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)) - 2*((8*b^3*c^4*d^2 - 34*a*b^2*c^3*d^3 + 41*a^2*b*c^2*d^4 - 15*a^3*c*d^5)*x^5 + 2*(12*b^3*c^5*d - 47*a*b^2*c^4*d^2 + 55*a^2*b*c^3*d^3 - 20*a^3*c^2*d^4)*x^3 + 3*(8*b^3*c^6 - 28*a*b^2*c^5*d + 31*a^2*b*c^4*d^2 - 11*a^3*c^3*d^3)*x)*sqrt(b*x^2 + a))/(b^3*c^10 - 3*a*b^2*c^9*d + 3*a^2*b*c^8*d^2 - a^3*c^7*d^3 + (b^3*c^7*d^3 - 3*a*b^2*c^6*d^4 + 3*a^2*b*c^5*d^5 - a^3*c^4*d^6)*x^6 + 3*(b^3*c^8*d^2 - 3*a*b^2*c^7*d^3 + 3*a^2*b*c^6*d^4 - a^3*c^5*d^5)*x^4 + 3*(b^3*c^9*d - 3*a*b^2*c^8*d^2 + 3*a^2*b*c^7*d^3 - a^3*c^6*d^4)*x^2)]

giac [B] time = 2.87, size = 958, normalized size = 4.61

$$\frac{\left(8ab^{\frac{5}{2}}c^2 - 12a^2b^{\frac{3}{2}}cd + 5a^3\sqrt{b}d^2\right) \arctan\left(\frac{\left(\sqrt{bx} - \sqrt{bx^2+a}\right)^{d+2bc-ad}}{2\sqrt{-b^2c^2+abcd}}\right)}{16\left(b^2c^5 - 2abc^4d + a^2c^3d^2\right)\sqrt{-b^2c^2+abcd}} - 24\left(\sqrt{bx} - \sqrt{bx^2+a}\right)^{10}ab^{\frac{5}{2}}c^2d^3 - 36\left(\sqrt{bx} - \sqrt{bx^2+a}\right)^{10}ab^{\frac{5}{2}}c^2d^3 - 36\left(\sqrt{bx} - \sqrt{bx^2+a}\right)^{10}ab^{\frac{5}{2}}c^2d^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^4,x, algorithm="giac")

[Out] -1/16*(8*a*b^(5/2)*c^2 - 12*a^2*b^(3/2)*c*d + 5*a^3*sqrt(b)*d^2)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/((b^2*c^5 - 2*a*b*c^4*d + a^2*c^3*d^2)*sqrt(-b^2*c^2 + a*b*c*d)) - 1/24*(24*(sqrt(b)*x - sqrt(b*x^2 + a))^10*a*b^(5/2)*c^2*d^3 - 36*(sqrt(b)*x - sqrt(b*x^2 + a))^10*a^2*b^(3/2)*c*d^4 + 15*(sqrt(b)*x - sqrt(b*x^2 + a))^10*a^3*sqrt(b)*d^5 + 240*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a*b^(7/2)*c^3*d^2 - 480*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^2*b^(5/2)*c^2*d^3 + 330*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^3*b^(3/2)*c*d^4 - 75*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^4*sqrt(b)*d^5 - 256*(sqrt(b)*x - sqrt(b*x^2 + a))^6*b^(11/2)*c^5 + 1216*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a*b^(9/2)*c^4*d - 2016*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^2*b^(7/2)*c^3*d^2 + 1736*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^3*b^(5/2)*c^2*d^3 - 800*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^4*b^(3/2)*c*d^4 + 150*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^5*sqrt(b)*d^5 - 384*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^2*b^(9/2)*c^4*d + 1392*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^3*b^(7/2)*c^3*d^2 - 1608*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^4*b^(5/2)*c^2*d^3 + 780*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^5*b^(3/2)*c*d^4 - 150*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^6*sqrt(b)*d^5 - 96*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^4*b^(7/2)*c^3*d^2 + 336*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^5*b^(5/2)*c^2*d^3 - 300*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^6*b^(3/2)*c*d^4 + 75*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^7*sqrt(b)*d^5 - 8*a^6*b^(5/2)*c^2*d^3 + 26*a^7*b^(3/2)*c*d^4

- 15*a^8*sqrt(b)*d^5)/((b^2*c^5*d - 2*a*b*c^4*d^2 + a^2*c^3*d^3)*((sqrt(b)*x - sqrt(b*x^2 + a))^4*d + 4*(sqrt(b)*x - sqrt(b*x^2 + a))^2*b*c - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*d + a^2*d)^3)

maple [B] time = 0.03, size = 7922, normalized size = 38.09

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)/(d*x^2+c)^4,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + a}}{(dx^2 + c)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^4,x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)/(d*x^2 + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{bx^2 + a}}{(dx^2 + c)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)/(c + d*x^2)^4,x)

[Out] int((a + b*x^2)^(1/2)/(c + d*x^2)^4, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)/(d*x**2+c)**4,x)

[Out] Timed out

3.53 $\int (a + bx^2)^{3/2} (c + dx^2)^3 dx$

Optimal. Leaf size=272

$$\frac{3a^2(4bc - ad)(a^2d^2 - 2abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{256b^{7/2}} + \frac{dx(a + bx^2)^{5/2}(5a^2d^2 - 20abcd + 36b^2c^2)}{160b^3} + \frac{x(a + bx^2)^{3/2}(4bc - ad)(a^2d^2 - 2abcd + 8b^2c^2)}{128b^3}$$

[Out] 1/128*(-a*d+4*b*c)*(a^2*d^2-2*a*b*c*d+8*b^2*c^2)*x*(b*x^2+a)^(3/2)/b^3+1/160*d*(5*a^2*d^2-20*a*b*c*d+36*b^2*c^2)*x*(b*x^2+a)^(5/2)/b^3+1/80*d*(-5*a*d+14*b*c)*x*(b*x^2+a)^(5/2)*(d*x^2+c)/b^2+1/10*d*x*(b*x^2+a)^(5/2)*(d*x^2+c)^2/b+3/256*a^2*(-a*d+4*b*c)*(a^2*d^2-2*a*b*c*d+8*b^2*c^2)*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(7/2)+3/256*a*(-a*d+4*b*c)*(a^2*d^2-2*a*b*c*d+8*b^2*c^2)*x*(b*x^2+a)^(1/2)/b^3

Rubi [A] time = 0.22, antiderivative size = 272, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {416, 528, 388, 195, 217, 206}

$$\frac{dx(a + bx^2)^{5/2}(5a^2d^2 - 20abcd + 36b^2c^2)}{160b^3} + \frac{x(a + bx^2)^{3/2}(4bc - ad)(a^2d^2 - 2abcd + 8b^2c^2)}{128b^3} + \frac{3ax\sqrt{a + bx^2}(4bc - ad)(a^2d^2 - 2abcd + 8b^2c^2)}{128b^3}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)*(c + d*x^2)^3,x]

[Out] (3*a*(4*b*c - a*d)*(8*b^2*c^2 - 2*a*b*c*d + a^2*d^2)*x*sqrt[a + b*x^2])/(256*b^3) + ((4*b*c - a*d)*(8*b^2*c^2 - 2*a*b*c*d + a^2*d^2)*x*(a + b*x^2)^(3/2))/(128*b^3) + (d*(36*b^2*c^2 - 20*a*b*c*d + 5*a^2*d^2)*x*(a + b*x^2)^(5/2))/(160*b^3) + (d*(14*b*c - 5*a*d)*x*(a + b*x^2)^(5/2)*(c + d*x^2))/(80*b^2) + (d*x*(a + b*x^2)^(5/2)*(c + d*x^2)^2)/(10*b) + (3*a^2*(4*b*c - a*d)*(8*b^2*c^2 - 2*a*b*c*d + a^2*d^2)*ArcTanh[(sqrt[b]*x)/sqrt[a + b*x^2]])/(256*b^3)

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 416

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)),
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a,
b, c, d, n, p, q, x]
```

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rubi steps

$$\begin{aligned}
\int (a + bx^2)^{3/2} (c + dx^2)^3 dx &= \frac{dx (a + bx^2)^{5/2} (c + dx^2)^2}{10b} + \frac{\int (a + bx^2)^{3/2} (c + dx^2) (c(10bc - ad) + d(14bc - 5ad)) dx}{10b} \\
&= \frac{d(14bc - 5ad)x (a + bx^2)^{5/2} (c + dx^2)}{80b^2} + \frac{dx (a + bx^2)^{5/2} (c + dx^2)^2}{10b} + \frac{\int (a + bx^2)^{3/2} (c + dx^2) dx}{10b} \\
&= \frac{d(36b^2c^2 - 20abcd + 5a^2d^2)x (a + bx^2)^{5/2}}{160b^3} + \frac{d(14bc - 5ad)x (a + bx^2)^{5/2} (c + dx^2)}{80b^2} \\
&= \frac{(4bc - ad)(8b^2c^2 - 2abcd + a^2d^2)x (a + bx^2)^{3/2}}{128b^3} + \frac{d(36b^2c^2 - 20abcd + 5a^2d^2)x (a + bx^2)^{5/2}}{160b^3} \\
&= \frac{3a(4bc - ad)(8b^2c^2 - 2abcd + a^2d^2)x\sqrt{a + bx^2}}{256b^3} + \frac{(4bc - ad)(8b^2c^2 - 2abcd + a^2d^2)x\sqrt{a + bx^2}}{128b^3} \\
&= \frac{3a(4bc - ad)(8b^2c^2 - 2abcd + a^2d^2)x\sqrt{a + bx^2}}{256b^3} + \frac{(4bc - ad)(8b^2c^2 - 2abcd + a^2d^2)x\sqrt{a + bx^2}}{128b^3} \\
&= \frac{3a(4bc - ad)(8b^2c^2 - 2abcd + a^2d^2)x\sqrt{a + bx^2}}{256b^3} + \frac{(4bc - ad)(8b^2c^2 - 2abcd + a^2d^2)x\sqrt{a + bx^2}}{128b^3}
\end{aligned}$$

Mathematica [A] time = 5.13, size = 220, normalized size = 0.81

$$\sqrt{b}x\sqrt{a + bx^2} (15a^4d^3 - 10a^3bd^2(9c + dx^2) + 4a^2b^2d(60c^2 + 15cdx^2 + 2d^2x^4) + 16ab^3(50c^3 + 70c^2dx^2 + 45cd^2x^4)) / (1280b^{7/2})$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)*(c + d*x^2)^3,x]

[Out] (Sqrt[b]*x*Sqrt[a + b*x^2]*(15*a^4*d^3 - 10*a^3*b*d^2*(9*c + d*x^2) + 4*a^2*b^2*d*(60*c^2 + 15*c*d*x^2 + 2*d^2*x^4) + 32*b^4*x^2*(10*c^3 + 20*c^2*d*x^2 + 15*c*d^2*x^4 + 4*d^3*x^6) + 16*a*b^3*(50*c^3 + 70*c^2*d*x^2 + 45*c*d^2*x^4 + 11*d^3*x^6)) - 15*a^2*(-4*b*c + a*d)*(8*b^2*c^2 - 2*a*b*c*d + a^2*d^2)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(1280*b^(7/2))

fricas [A] time = 1.21, size = 502, normalized size = 1.85

$$\frac{15(32a^2b^3c^3 - 16a^3b^2c^2d + 6a^4bcd^2 - a^5d^3)\sqrt{b} \log(-2bx^2 + 2\sqrt{bx^2 + a}\sqrt{b}x - a) - 2(128b^5d^3x^9 + 16(30b^5c^2d^2 + 11a^2b^4cd^3)x^7 + 8(80b^5c^2d + 90a^2b^4cd^2 + a^2b^3d^3)x^5 + 10(32b^5c^3 + 112a^2b^4cd^2 + 6a^2b^3cd^2 - a^3b^2d^3)x^3 + 5(160a^2b^4cd^3 + 48a^2b^3c^2d - 18a^3b^2cd^2 + 3a^4bd^3)x)\sqrt{bx^2 + a}}{b^4} - \frac{1}{1280}(15(32a^2b^3c^3 - 16a^3b^2c^2d + 6a^4bcd^2 - a^5d^3)\sqrt{-b} \arctan(\sqrt{-b}x/\sqrt{bx^2 + a}) - (128b^5d^3x^9 + 16(30b^5c^2d^2 + 11a^2b^4cd^3)x^7 + 8(80b^5c^2d + 90a^2b^4cd^2 + a^2b^3d^3)x^5 + 10(32b^5c^3 + 112a^2b^4cd^2 + 6a^2b^3cd^2 - a^3b^2d^3)x^3 + 5(160a^2b^4cd^3 + 48a^2b^3c^2d - 18a^3b^2cd^2 + 3a^4bd^3)x)\sqrt{bx^2 + a})/b^4$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^3,x, algorithm="fricas")

[Out] [-1/2560*(15*(32*a^2*b^3*c^3 - 16*a^3*b^2*c^2*d + 6*a^4*b*c*d^2 - a^5*d^3)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(128*b^5*d^3*x^9 + 16*(30*b^5*c^2*d^2 + 11*a*b^4*d^3)*x^7 + 8*(80*b^5*c^2*d + 90*a*b^4*c*d^2 + a^2*b^3*d^3)*x^5 + 10*(32*b^5*c^3 + 112*a*b^4*c^2*d + 6*a^2*b^3*c*d^2 - a^3*b^2*d^3)*x^3 + 5*(160*a*b^4*c^3 + 48*a^2*b^3*c^2*d - 18*a^3*b^2*c*d^2 + 3*a^4*b*d^3)*x)*sqrt(b*x^2 + a))/b^4, -1/1280*(15*(32*a^2*b^3*c^3 - 16*a^3*b^2*c^2*d + 6*a^4*b*c*d^2 - a^5*d^3)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (128*b^5*d^3*x^9 + 16*(30*b^5*c^2*d^2 + 11*a*b^4*d^3)*x^7 + 8*(80*b^5*c^2*d + 90*a*b^4*c*d^2 + a^2*b^3*d^3)*x^5 + 10*(32*b^5*c^3 + 112*a*b^4*c^2*d + 6*a^2*b^3*c*d^2 - a^3*b^2*d^3)*x^3 + 5*(160*a*b^4*c^3 + 48*a^2*b^3*c^2*d - 18*a^3*b^2*c*d^2 + 3*a^4*b*d^3)*x)*sqrt(b*x^2 + a))/b^4]

giac [A] time = 0.66, size = 260, normalized size = 0.96

$$\frac{1}{1280} \left(2 \left(4 \left(2 \left(8bd^3x^2 + \frac{30b^9cd^2 + 11ab^8d^3}{b^8} \right) x^2 + \frac{80b^9c^2d + 90ab^8cd^2 + a^2b^7d^3}{b^8} \right) x^2 + \frac{5(32b^9c^3 + 112ab^8c^2)}{b^8} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^3,x, algorithm="giac")

[Out] 1/1280*(2*(4*(2*(8*b*d^3*x^2 + (30*b^9*c*d^2 + 11*a*b^8*d^3)/b^8)*x^2 + (80*b^9*c^2*d + 90*a*b^8*c*d^2 + a^2*b^7*d^3)/b^8)*x^2 + 5*(32*b^9*c^3 + 112*a*b^8*c^2*d + 6*a^2*b^7*c*d^2 - a^3*b^6*d^3)/b^8)*x^2 + 5*(160*a*b^8*c^3 + 48*a^2*b^7*c^2*d - 18*a^3*b^6*c*d^2 + 3*a^4*b^5*d^3)/b^8)*sqrt(b*x^2 + a)*x - 3/256*(32*a^2*b^3*c^3 - 16*a^3*b^2*c^2*d + 6*a^4*b*c*d^2 - a^5*d^3)*log(a*b*(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(7/2)

maple [A] time = 0.02, size = 393, normalized size = 1.44

$$\frac{(bx^2 + a)^{\frac{5}{2}} d^3 x^5}{10b} - \frac{3a^5 d^3 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{256b^{\frac{7}{2}}} + \frac{9a^4 c d^2 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{128b^{\frac{5}{2}}} - \frac{3a^3 c^2 d \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{16b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)*(d*x^2+c)^3,x)

[Out] 1/10*d^3*x^5*(b*x^2+a)^(5/2)/b-1/16*d^3*a/b^2*x^3*(b*x^2+a)^(5/2)+1/32*d^3*a^2/b^3*x*(b*x^2+a)^(5/2)-1/128*d^3*a^3/b^3*x*(b*x^2+a)^(3/2)-3/256*d^3*a^4/b^3*x*(b*x^2+a)^(1/2)-3/256*d^3*a^5/b^(7/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+3/8*c*d^2*x^3*(b*x^2+a)^(5/2)/b-3/16*c*d^2*a/b^2*x*(b*x^2+a)^(5/2)+3/64*c*d^2*a^2/b^2*x*(b*x^2+a)^(3/2)+9/128*c*d^2*a^3/b^2*x*(b*x^2+a)^(1/2)+9/128*c*d^2*a^4/b^(5/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+1/2*c^2*d*x*(b*x^2+a)^(5/2)/b-1/8*c^2*d*a/b*x*(b*x^2+a)^(3/2)-3/16*c^2*d*a^2/b*x*(b*x^2+a)^(1/2)-3/16*c^2*d*a^3/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+1/4*c^3*x*(b*x^2+a)^(3/2)+3/8*c^3*a*x*(b*x^2+a)^(1/2)+3/8*c^3*a^2/b^(1/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))

maxima [A] time = 1.50, size = 364, normalized size = 1.34

$$\frac{(bx^2 + a)^{\frac{5}{2}} d^3 x^5}{10b} + \frac{3(bx^2 + a)^{\frac{5}{2}} cd^2 x^3}{8b} - \frac{(bx^2 + a)^{\frac{5}{2}} ad^3 x^3}{16b^2} + \frac{1}{4} (bx^2 + a)^{\frac{3}{2}} c^3 x + \frac{3}{8} \sqrt{bx^2 + a} ac^3 x + \frac{(bx^2 + a)^{\frac{5}{2}} c^2 dx}{2b} - \frac{3a^5 d^3 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{256b^{\frac{7}{2}}} + \frac{9a^4 c d^2 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{128b^{\frac{5}{2}}} - \frac{3a^3 c^2 d \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{16b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^3,x, algorithm="maxima")

[Out] $1/10*(b*x^2 + a)^{(5/2)}*d^3*x^5/b + 3/8*(b*x^2 + a)^{(5/2)}*c*d^2*x^3/b - 1/16*(b*x^2 + a)^{(5/2)}*a*d^3*x^3/b^2 + 1/4*(b*x^2 + a)^{(3/2)}*c^3*x + 3/8*\sqrt{(b*x^2 + a)}*a*c^3*x + 1/2*(b*x^2 + a)^{(5/2)}*c^2*d*x/b - 1/8*(b*x^2 + a)^{(3/2)}*a*c^2*d*x/b - 3/16*\sqrt{(b*x^2 + a)}*a^2*c^2*d*x/b - 3/16*(b*x^2 + a)^{(5/2)}*a*c*d^2*x/b^2 + 3/64*(b*x^2 + a)^{(3/2)}*a^2*c*d^2*x/b^2 + 9/128*\sqrt{(b*x^2 + a)}*a^3*c*d^2*x/b^2 + 1/32*(b*x^2 + a)^{(5/2)}*a^2*d^3*x/b^3 - 1/128*(b*x^2 + a)^{(3/2)}*a^3*d^3*x/b^3 - 3/256*\sqrt{(b*x^2 + a)}*a^4*d^3*x/b^3 + 3/8*a^2*c^3*\operatorname{arcsinh}(b*x/\sqrt{a*b})/\sqrt{b} - 3/16*a^3*c^2*d*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{(3/2)} + 9/128*a^4*c*d^2*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{(5/2)} - 3/256*a^5*d^3*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{(7/2)}$

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (bx^2 + a)^{3/2} (dx^2 + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)*(c + d*x^2)^3,x)

[Out] int((a + b*x^2)^(3/2)*(c + d*x^2)^3, x)

sympy [B] time = 52.75, size = 665, normalized size = 2.44

$$\frac{3a^{\frac{9}{2}}d^3x}{256b^3\sqrt{1+\frac{bx^2}{a}}} - \frac{9a^{\frac{7}{2}}cd^2x}{128b^2\sqrt{1+\frac{bx^2}{a}}} + \frac{a^{\frac{7}{2}}d^3x^3}{256b^2\sqrt{1+\frac{bx^2}{a}}} + \frac{3a^{\frac{5}{2}}c^2dx}{16b\sqrt{1+\frac{bx^2}{a}}} - \frac{3a^{\frac{5}{2}}cd^2x^3}{128b\sqrt{1+\frac{bx^2}{a}}} - \frac{a^{\frac{5}{2}}d^3x^5}{640b\sqrt{1+\frac{bx^2}{a}}} + \frac{a^{\frac{3}{2}}c^3x\sqrt{1+\frac{bx^2}{a}}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)*(d*x**2+c)**3,x)

[Out] $3*a^{(9/2)}*d^{**3}*x/(256*b^{**3}*\sqrt{(1 + b*x^{**2}/a)}) - 9*a^{(7/2)}*c*d^{**2}*x/(128*b^{**2}*\sqrt{(1 + b*x^{**2}/a)}) + a^{(7/2)}*d^{**3}*x^{**3}/(256*b^{**2}*\sqrt{(1 + b*x^{**2}/a)}) + 3*a^{(5/2)}*c^{**2}*d*x/(16*b*\sqrt{(1 + b*x^{**2}/a)}) - 3*a^{(5/2)}*c*d^{**2}*x^{**3}/(128*b*\sqrt{(1 + b*x^{**2}/a)}) - a^{(5/2)}*d^{**3}*x^{**5}/(640*b*\sqrt{(1 + b*x^{**2}/a)}) + a^{(3/2)}*c^{**3}*x*\sqrt{(1 + b*x^{**2}/a)}/2 + a^{(3/2)}*c^{**3}*x/(8*\sqrt{(1 + b*x^{**2}/a)}) + 17*a^{(3/2)}*c^{**2}*d*x^{**3}/(16*\sqrt{(1 + b*x^{**2}/a)}) + 39*a^{(3/2)}*c*d^{**2}*x^{**5}/(64*\sqrt{(1 + b*x^{**2}/a)}) + 23*a^{(3/2)}*d^{**3}*x^{**7}/(160*\sqrt{(1 + b*x^{**2}/a)}) + 3*\sqrt{a}*b*c^{**3}*x^{**3}/(8*\sqrt{(1 + b*x^{**2}/a)}) + 11*\sqrt{a}*b*c^{**2}*d*x^{**5}/(8*\sqrt{(1 + b*x^{**2}/a)}) + 15*\sqrt{a}*b*c*d^{**2}*x^{**7}/(16*\sqrt{(1 + b*x^{**2}/a)}) + 19*\sqrt{a}*b*d^{**3}*x^{**9}/(80*\sqrt{(1 + b*x^{**2}/a)}) - 3*a^{**5}*d^{**3}*a*\operatorname{asinh}(\sqrt{(b)*x/\sqrt{a}})/(256*b^{**}(7/2)) + 9*a^{**4}*c*d^{**2}*a*\operatorname{asinh}(\sqrt{(b)*x/\sqrt{a}})/(128*b^{**}(5/2)) - 3*a^{**3}*c^{**2}*d*a*\operatorname{asinh}(\sqrt{(b)*x/\sqrt{a}})/(16*b^{**}(3/2)) + 3*a^{**2}*c^{**3}*a*\operatorname{asinh}(\sqrt{(b)*x/\sqrt{a}})/(8*\sqrt{(b)}) + b^{**2}*c^{**3}*x^{**5}/(4*\sqrt{a}*\sqrt{(1 + b*x^{**2}/a)}) + b^{**2}*c^{**2}*d*x^{**7}/(2*\sqrt{a}*\sqrt{(1 + b*x^{**2}/a)}) + 3*b^{**2}*c*d^{**2}*x^{**9}/(8*\sqrt{a}*\sqrt{(1 + b*x^{**2}/a)}) + b^{**2}*d^{**3}*x^{**11}/(10*\sqrt{a}*\sqrt{(1 + b*x^{**2}/a)})$

3.54 $\int (a + bx^2)^{3/2} (c + dx^2)^2 dx$

Optimal. Leaf size=196

$$\frac{x(a + bx^2)^{3/2} (3a^2d^2 - 16abcd + 48b^2c^2)}{192b^2} + \frac{ax\sqrt{a + bx^2} (3a^2d^2 - 16abcd + 48b^2c^2)}{128b^2} + \frac{a^2 (3a^2d^2 - 16abcd + 48b^2c^2)}{128b^5}$$

[Out] 1/192*(3*a^2*d^2-16*a*b*c*d+48*b^2*c^2)*x*(b*x^2+a)^(3/2)/b^2+1/48*d*(-3*a*d+10*b*c)*x*(b*x^2+a)^(5/2)/b^2+1/8*d*x*(b*x^2+a)^(5/2)*(d*x^2+c)/b+1/128*a^2*(3*a^2*d^2-16*a*b*c*d+48*b^2*c^2)*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(5/2)+1/128*a*(3*a^2*d^2-16*a*b*c*d+48*b^2*c^2)*x*(b*x^2+a)^(1/2)/b^2

Rubi [A] time = 0.12, antiderivative size = 196, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {416, 388, 195, 217, 206}

$$\frac{x(a + bx^2)^{3/2} (3a^2d^2 - 16abcd + 48b^2c^2)}{192b^2} + \frac{ax\sqrt{a + bx^2} (3a^2d^2 - 16abcd + 48b^2c^2)}{128b^2} + \frac{a^2 (3a^2d^2 - 16abcd + 48b^2c^2)}{128b^5}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)*(c + d*x^2)^2,x]

[Out] (a*(48*b^2*c^2 - 16*a*b*c*d + 3*a^2*d^2)*x*sqrt[a + b*x^2])/(128*b^2) + ((48*b^2*c^2 - 16*a*b*c*d + 3*a^2*d^2)*x*(a + b*x^2)^(3/2))/(192*b^2) + (d*(10*b*c - 3*a*d)*x*(a + b*x^2)^(5/2))/(48*b^2) + (d*x*(a + b*x^2)^(5/2)*(c + d*x^2))/(8*b) + (a^2*(48*b^2*c^2 - 16*a*b*c*d + 3*a^2*d^2)*ArcTanh[(sqrt[b]*x)/sqrt[a + b*x^2]])/(128*b^(5/2))

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)),

```
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a
, b, c, d, n, p, q, x]
```

Rubi steps

$$\int (a + bx^2)^{3/2} (c + dx^2)^2 dx = \frac{dx (a + bx^2)^{5/2} (c + dx^2)}{8b} + \frac{\int (a + bx^2)^{3/2} (c(8bc - ad) + d(10bc - 3ad)x^2) dx}{8b}$$

$$= \frac{d(10bc - 3ad)x (a + bx^2)^{5/2}}{48b^2} + \frac{dx (a + bx^2)^{5/2} (c + dx^2)}{8b} - \frac{(ad(10bc - 3ad) - 6bc^2)}{48b^2}$$

$$= \frac{(48b^2c^2 - 16abcd + 3a^2d^2)x (a + bx^2)^{3/2}}{192b^2} + \frac{d(10bc - 3ad)x (a + bx^2)^{5/2}}{48b^2} + \frac{dx (a + bx^2)^{3/2}}{8b}$$

$$= \frac{a(48b^2c^2 - 16abcd + 3a^2d^2)x\sqrt{a + bx^2}}{128b^2} + \frac{(48b^2c^2 - 16abcd + 3a^2d^2)x (a + bx^2)^{3/2}}{192b^2}$$

$$= \frac{a(48b^2c^2 - 16abcd + 3a^2d^2)x\sqrt{a + bx^2}}{128b^2} + \frac{(48b^2c^2 - 16abcd + 3a^2d^2)x (a + bx^2)^{3/2}}{192b^2}$$

$$= \frac{a(48b^2c^2 - 16abcd + 3a^2d^2)x\sqrt{a + bx^2}}{128b^2} + \frac{(48b^2c^2 - 16abcd + 3a^2d^2)x (a + bx^2)^{3/2}}{192b^2}$$

Mathematica [C] time = 2.71, size = 157, normalized size = 0.80

$$\frac{x\sqrt{a + bx^2} \left(6bx^2 (c + dx^2)^2 {}_3F_2\left(-\frac{1}{2}, \frac{3}{2}, 2; 1, \frac{9}{2}; -\frac{bx^2}{a}\right) + 12bx^2 (2c^2 + 3cdx^2 + d^2x^4) {}_2F_1\left(-\frac{1}{2}, \frac{3}{2}; \frac{9}{2}; -\frac{bx^2}{a}\right) + 7a (15cd^2 + 3d^3x^2) {}_2F_1\left(-\frac{1}{2}, \frac{3}{2}; \frac{9}{2}; -\frac{bx^2}{a}\right) \right)}{105\sqrt{\frac{bx^2}{a} + 1}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(3/2)*(c + d*x^2)^2,x]

[Out] (x*Sqrt[a + b*x^2]*(7*a*(15*c^2 + 10*c*d*x^2 + 3*d^2*x^4)*Hypergeometric2F1[-3/2, 1/2, 7/2, -((b*x^2)/a)] + 12*b*x^2*(2*c^2 + 3*c*d*x^2 + d^2*x^4)*Hypergeometric2F1[-1/2, 3/2, 9/2, -((b*x^2)/a)] + 6*b*x^2*(c + d*x^2)^2*HypergeometricPFQ[{-1/2, 3/2, 2}, {1, 9/2}, -((b*x^2)/a)]))/(105*Sqrt[1 + (b*x^2)/a])

fricas [A] time = 0.78, size = 344, normalized size = 1.76

$$\frac{3(48a^2b^2c^2 - 16a^3bcd + 3a^4d^2)\sqrt{b} \log(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{b}x - a) + 2(48b^4d^2x^7 + 8(16b^4cd + 9ab^3d^2)x^6 + 2(48b^4cd^2 + 8(16b^4cd + 9ab^3d^2)x^5 + 2(48b^4cd^2 + 8(16b^4cd + 9ab^3d^2)x^4 + 2(48b^4cd^2 + 8(16b^4cd + 9ab^3d^2)x^3 + 3(80a^2b^3c^2 + 16a^2b^2c*d - 3a^3b*d^2)*x)*\sqrt{b*x^2 + a})/b^3, -1/384*(3*(48a^2b^2c^2 - 16a^3b*c*d + 3a^4*d^2)*\sqrt{-b}*\arctan(\sqrt{-b}*x/s$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^2,x, algorithm="fricas")

[Out] [1/768*(3*(48*a^2*b^2*c^2 - 16*a^3*b*c*d + 3*a^4*d^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 2*(48*b^4*d^2*x^7 + 8*(16*b^4*c*d + 9*a*b^3*d^2)*x^6 + 2*(48*b^4*c*d^2 + 8*(16*b^4*c*d + 9*a*b^3*d^2)*x^5 + 2*(48*b^4*c*d^2 + 8*(16*b^4*c*d + 9*a*b^3*d^2)*x^4 + 2*(48*b^4*c*d^2 + 8*(16*b^4*c*d + 9*a*b^3*d^2)*x^3 + 3*(80*a^2*b^3*c^2 + 16*a^2*b^2*c*d - 3*a^3*b*d^2)*x)*sqrt(b*x^2 + a))/b^3, -1/384*(3*(48*a^2*b^2*c^2 - 16*a^3*b*c*d + 3*a^4*d^2)*sqrt(-b)*arctan(sqrt(-b)*x/s

$\text{qrt}(b*x^2 + a) - (48*b^4*d^2*x^7 + 8*(16*b^4*c*d + 9*a*b^3*d^2)*x^5 + 2*(48*b^4*c^2 + 112*a*b^3*c*d + 3*a^2*b^2*d^2)*x^3 + 3*(80*a*b^3*c^2 + 16*a^2*b^2*c*d - 3*a^3*b*d^2)*x)*\text{sqrt}(b*x^2 + a)/b^3]$

giac [A] time = 0.66, size = 175, normalized size = 0.89

$$\frac{1}{384} \left(2 \left(4 \left(6 b d^2 x^2 + \frac{16 b^7 c d + 9 a b^6 d^2}{b^6} \right) x^2 + \frac{48 b^7 c^2 + 112 a b^6 c d + 3 a^2 b^5 d^2}{b^6} \right) x^2 + \frac{3 (80 a b^6 c^2 + 16 a^2 b^5 c d - 3 a^3 b^4 d^2)}{b^6} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^2,x, algorithm="giac")

[Out] 1/384*(2*(4*(6*b*d^2*x^2 + (16*b^7*c*d + 9*a*b^6*d^2)/b^6)*x^2 + (48*b^7*c^2 + 112*a*b^6*c*d + 3*a^2*b^5*d^2)/b^6)*x^2 + 3*(80*a*b^6*c^2 + 16*a^2*b^5*c*d - 3*a^3*b^4*d^2)/b^6)*sqrt(b*x^2 + a)*x - 1/128*(48*a^2*b^2*c^2 - 16*a^3*b*c*d + 3*a^4*d^2)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(5/2)

maple [A] time = 0.01, size = 249, normalized size = 1.27

$$\frac{3a^4d^2 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{128b^{\frac{5}{2}}} - \frac{a^3cd \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{8b^{\frac{3}{2}}} + \frac{3a^2c^2 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{8\sqrt{b}} + \frac{3\sqrt{bx^2 + a} a^3d^2x}{128b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)*(d*x^2+c)^2,x)

[Out] 1/8*d^2*x^3*(b*x^2+a)^(5/2)/b-1/16*d^2*a/b^2*x*(b*x^2+a)^(5/2)+1/64*d^2*a^2/b^2*x*(b*x^2+a)^(3/2)+3/128*d^2*a^3/b^2*x*(b*x^2+a)^(1/2)+3/128*d^2*a^4/b^(5/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+1/3*c*d*x*(b*x^2+a)^(5/2)/b-1/12*c*d*a/b*x*(b*x^2+a)^(3/2)-1/8*c*d*a^2/b*x*(b*x^2+a)^(1/2)-1/8*c*d*a^3/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+1/4*c^2*x*(b*x^2+a)^(3/2)+3/8*c^2*a*x*(b*x^2+a)^(1/2)+3/8*c^2*a^2/b^(1/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))

maxima [A] time = 1.42, size = 227, normalized size = 1.16

$$\frac{(bx^2 + a)^{\frac{5}{2}}d^2x^3}{8b} + \frac{1}{4}(bx^2 + a)^{\frac{3}{2}}c^2x + \frac{3}{8}\sqrt{bx^2 + a}ac^2x + \frac{(bx^2 + a)^{\frac{5}{2}}cdx}{3b} - \frac{(bx^2 + a)^{\frac{3}{2}}acd}{12b} - \frac{\sqrt{bx^2 + a}a^2cdx}{8b} - \frac{(bx^2 + a)^{\frac{5}{2}}d^2x^3}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^2,x, algorithm="maxima")

[Out] 1/8*(b*x^2 + a)^(5/2)*d^2*x^3/b + 1/4*(b*x^2 + a)^(3/2)*c^2*x + 3/8*sqrt(b*x^2 + a)*a*c^2*x + 1/3*(b*x^2 + a)^(5/2)*c*d*x/b - 1/12*(b*x^2 + a)^(3/2)*a*c*d*x/b - 1/8*sqrt(b*x^2 + a)*a^2*c*d*x/b - 1/16*(b*x^2 + a)^(5/2)*a*d^2*x/b^2 + 1/64*(b*x^2 + a)^(3/2)*a^2*d^2*x/b^2 + 3/128*sqrt(b*x^2 + a)*a^3*d^2*x/b^2 + 3/8*a^2*c^2*arcsinh(b*x/sqrt(a*b))/sqrt(b) - 1/8*a^3*c*d*arcsinh(b*x/sqrt(a*b))/b^(3/2) + 3/128*a^4*d^2*arcsinh(b*x/sqrt(a*b))/b^(5/2)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (bx^2 + a)^{3/2} (dx^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)*(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(3/2)*(c + d*x^2)^2, x)

sympy [B] time = 29.42, size = 440, normalized size = 2.24

$$-\frac{3a^{\frac{7}{2}}d^2x}{128b^2\sqrt{1+\frac{bx^2}{a}}} + \frac{a^{\frac{5}{2}}cdx}{8b\sqrt{1+\frac{bx^2}{a}}} - \frac{a^{\frac{5}{2}}d^2x^3}{128b\sqrt{1+\frac{bx^2}{a}}} + \frac{a^{\frac{3}{2}}c^2x\sqrt{1+\frac{bx^2}{a}}}{2} + \frac{a^{\frac{3}{2}}c^2x}{8\sqrt{1+\frac{bx^2}{a}}} + \frac{17a^{\frac{3}{2}}cdx^3}{24\sqrt{1+\frac{bx^2}{a}}} + \frac{13a^{\frac{3}{2}}d^2x^5}{64\sqrt{1+\frac{bx^2}{a}}} + \frac{3\sqrt{a}}{8\sqrt{1+\frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)*(d*x**2+c)**2,x)

[Out] $-3*a^{7/2}*d^{2*x}/(128*b^{2*sqrt(1 + b*x^{2}/a)) + a^{5/2}*c*d*x/(8*b*sqrt(1 + b*x^{2}/a)) - a^{5/2}*d^{2*x^3}/(128*b*sqrt(1 + b*x^{2}/a)) + a^{3/2}*c^{2*x}*sqrt(1 + b*x^{2}/a)/2 + a^{3/2}*c^{2*x}/(8*sqrt(1 + b*x^{2}/a)) + 17*a^{3/2}*c*d*x^3/(24*sqrt(1 + b*x^{2}/a)) + 13*a^{3/2}*d^{2*x^5}/(64*sqrt(1 + b*x^{2}/a)) + 3*sqrt(a)*b*c^{2*x^3}/(8*sqrt(1 + b*x^{2}/a)) + 11*sqrt(a)*b*c*d*x^5/(12*sqrt(1 + b*x^{2}/a)) + 5*sqrt(a)*b*d^{2*x^7}/(16*sqrt(1 + b*x^{2}/a)) + 3*a^{4*d^{2*asinh(sqrt(b)*x/sqrt(a))}/(128*b^{5/2}) - a^{3*c*d*asinh(sqrt(b)*x/sqrt(a))}/(8*b^{3/2}) + 3*a^{2*c^{2*asinh(sqrt(b)*x/sqrt(a))}/(8*sqrt(b)) + b^{2*c^{2*x^5}/(4*sqrt(a)*sqrt(1 + b*x^{2}/a))} + b^{2*c*d*x^7/(3*sqrt(a)*sqrt(1 + b*x^{2}/a))} + b^{2*d^{2*x^9}/(8*sqrt(a)*sqrt(1 + b*x^{2}/a))}$

3.55 $\int (a + bx^2)^{3/2} (c + dx^2) dx$

Optimal. Leaf size=118

$$\frac{a^2(6bc - ad) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16b^{3/2}} + \frac{x(a + bx^2)^{3/2} (6bc - ad)}{24b} + \frac{ax\sqrt{a + bx^2} (6bc - ad)}{16b} + \frac{dx(a + bx^2)^{5/2}}{6b}$$

[Out] $1/24*(-a*d+6*b*c)*x*(b*x^2+a)^{(3/2)}/b+1/6*d*x*(b*x^2+a)^{(5/2)}/b+1/16*a^2*(-a*d+6*b*c)*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(3/2)}+1/16*a*(-a*d+6*b*c)*x*(b*x^2+a)^{(1/2)}/b$

Rubi [A] time = 0.04, antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {388, 195, 217, 206}

$$\frac{a^2(6bc - ad) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16b^{3/2}} + \frac{x(a + bx^2)^{3/2} (6bc - ad)}{24b} + \frac{ax\sqrt{a + bx^2} (6bc - ad)}{16b} + \frac{dx(a + bx^2)^{5/2}}{6b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a + b*x^2)^{(3/2)}*(c + d*x^2), x]$

[Out] $(a*(6*b*c - a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(16*b) + ((6*b*c - a*d)*x*(a + b*x^2)^{(3/2)})/(24*b) + (d*x*(a + b*x^2)^{(5/2)})/(6*b) + (a^2*(6*b*c - a*d)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(16*b^{(3/2)})$

Rule 195

$\operatorname{Int}[(a_ + (b_.)*(x_)^{(n_))^{(p_)}, x_Symbol] := \operatorname{Simp}[(x*(a + b*x^n)^p)/(n*p + 1), x] + \operatorname{Dist}[(a*n*p)/(n*p + 1), \operatorname{Int}[(a + b*x^n)^{p-1}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

$\operatorname{Int}[(a_ + (b_.)*(x_)^2)^{-1}, x_Symbol] := \operatorname{Simp}[(1*\operatorname{ArcTanh}[(\operatorname{Rt}[-b, 2]*x)/\operatorname{Rt}[a, 2]])/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]), x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_.)*(x_)^2], x_Symbol] := \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /;$ FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

$\operatorname{Int}[(a_ + (b_.)*(x_)^{(n_))^{(p_)}*((c_ + (d_.)*(x_)^{(n_))}, x_Symbol] := \operatorname{Simp}[(d*x*(a + b*x^n)^{(p+1)})/(b*(n*(p+1) + 1)), x] - \operatorname{Dist}[(a*d - b*c*(n*(p+1) + 1))/(b*(n*(p+1) + 1)), \operatorname{Int}[(a + b*x^n)^p, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1) + 1, 0]

Rubi steps

$$\begin{aligned}
\int (a + bx^2)^{3/2} (c + dx^2) dx &= \frac{dx (a + bx^2)^{5/2}}{6b} - \frac{(-6bc + ad) \int (a + bx^2)^{3/2} dx}{6b} \\
&= \frac{(6bc - ad)x (a + bx^2)^{3/2}}{24b} + \frac{dx (a + bx^2)^{5/2}}{6b} + \frac{(a(6bc - ad)) \int \sqrt{a + bx^2} dx}{8b} \\
&= \frac{a(6bc - ad)x\sqrt{a + bx^2}}{16b} + \frac{(6bc - ad)x (a + bx^2)^{3/2}}{24b} + \frac{dx (a + bx^2)^{5/2}}{6b} + \frac{(a^2(6bc - ad)) \int \sqrt{a + bx^2} dx}{8b} \\
&= \frac{a(6bc - ad)x\sqrt{a + bx^2}}{16b} + \frac{(6bc - ad)x (a + bx^2)^{3/2}}{24b} + \frac{dx (a + bx^2)^{5/2}}{6b} + \frac{(a^2(6bc - ad)) \int \sqrt{a + bx^2} dx}{8b} \\
&= \frac{a(6bc - ad)x\sqrt{a + bx^2}}{16b} + \frac{(6bc - ad)x (a + bx^2)^{3/2}}{24b} + \frac{dx (a + bx^2)^{5/2}}{6b} + \frac{a^2(6bc - ad) \int \sqrt{a + bx^2} dx}{8b}
\end{aligned}$$

Mathematica [A] time = 0.20, size = 109, normalized size = 0.92

$$\frac{\sqrt{a + bx^2} \left(\sqrt{b} x (3a^2d + 2ab(15c + 7dx^2) + 4b^2x^2(3c + 2dx^2)) - \frac{3a^{3/2}(ad - 6bc) \sinh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{\frac{bx^2}{a} + 1}} \right)}{48b^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)*(c + d*x^2), x]

[Out] (Sqrt[a + b*x^2]*(Sqrt[b]*x*(3*a^2*d + 4*b^2*x^2*(3*c + 2*d*x^2) + 2*a*b*(15*c + 7*d*x^2)) - (3*a^(3/2)*(-6*b*c + a*d)*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/Sqrt[1 + (b*x^2)/a])/ (48*b^(3/2))

fricas [A] time = 0.76, size = 210, normalized size = 1.78

$$\left[\frac{3(6a^2bc - a^3d)\sqrt{b} \log(-2bx^2 + 2\sqrt{bx^2 + a}\sqrt{b}x - a) - 2(8b^3dx^5 + 2(6b^3c + 7ab^2d)x^3 + 3(10ab^2c + a^2bd))}{96b^2} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c), x, algorithm="fricas")

[Out] [-1/96*(3*(6*a^2*b*c - a^3*d)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(8*b^3*d*x^5 + 2*(6*b^3*c + 7*a*b^2*d)*x^3 + 3*(10*a*b^2*c + a^2*b*d)*x)*sqrt(b*x^2 + a))/b^2, -1/48*(3*(6*a^2*b*c - a^3*d)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (8*b^3*d*x^5 + 2*(6*b^3*c + 7*a*b^2*d)*x^3 + 3*(10*a*b^2*c + a^2*b*d)*x)*sqrt(b*x^2 + a))/b^2]

giac [A] time = 0.61, size = 103, normalized size = 0.87

$$\frac{1}{48} \left(2 \left(4bdx^2 + \frac{6b^5c + 7ab^4d}{b^4} \right) x^2 + \frac{3(10ab^4c + a^2b^3d)}{b^4} \right) \sqrt{bx^2 + a} x - \frac{(6a^2bc - a^3d) \log\left(\left| -\sqrt{b}x + \sqrt{bx^2 + a} \right|\right)}{16b^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c), x, algorithm="giac")

[Out] 1/48*(2*(4*b*d*x^2 + (6*b^5*c + 7*a*b^4*d)/b^4)*x^2 + 3*(10*a*b^4*c + a^2*b^3*d)/b^4)*sqrt(b*x^2 + a)*x - 1/16*(6*a^2*b*c - a^3*d)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(3/2)

maple [A] time = 0.00, size = 131, normalized size = 1.11

$$\frac{a^3 d \ln(\sqrt{b} x + \sqrt{b x^2 + a})}{16 b^{\frac{3}{2}}} + \frac{3 a^2 c \ln(\sqrt{b} x + \sqrt{b x^2 + a})}{8 \sqrt{b}} - \frac{\sqrt{b x^2 + a} a^2 dx}{16 b} + \frac{3 \sqrt{b x^2 + a} a c x}{8} - \frac{(b x^2 + a)^{\frac{3}{2}} a d x}{24 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)*(d*x^2+c), x)

[Out] 1/6*d*x*(b*x^2+a)^(5/2)/b-1/24*d*a/b*x*(b*x^2+a)^(3/2)-1/16*d*a^2/b*x*(b*x^2+a)^(1/2)-1/16*d*a^3/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+1/4*c*x*(b*x^2+a)^(3/2)+3/8*c*a*x*(b*x^2+a)^(1/2)+3/8*c*a^2/b^(1/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))

maxima [A] time = 1.37, size = 116, normalized size = 0.98

$$\frac{1}{4} (b x^2 + a)^{\frac{3}{2}} c x + \frac{3}{8} \sqrt{b x^2 + a} a c x + \frac{(b x^2 + a)^{\frac{5}{2}} dx}{6 b} - \frac{(b x^2 + a)^{\frac{3}{2}} a d x}{24 b} - \frac{\sqrt{b x^2 + a} a^2 dx}{16 b} + \frac{3 a^2 c \operatorname{arsinh}\left(\frac{b x}{\sqrt{a b}}\right)}{8 \sqrt{b}} - \frac{a^3 d \operatorname{arsinh}\left(\frac{b x}{\sqrt{a b}}\right)}{16 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c), x, algorithm="maxima")

[Out] 1/4*(b*x^2 + a)^(3/2)*c*x + 3/8*sqrt(b*x^2 + a)*a*c*x + 1/6*(b*x^2 + a)^(5/2)*d*x/b - 1/24*(b*x^2 + a)^(3/2)*a*d*x/b - 1/16*sqrt(b*x^2 + a)*a^2*d*x/b + 3/8*a^2*c*arcsinh(b*x/sqrt(a*b))/sqrt(b) - 1/16*a^3*d*arcsinh(b*x/sqrt(a*b))/b^(3/2)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (b x^2 + a)^{3/2} (d x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)*(c + d*x^2), x)

[Out] int((a + b*x^2)^(3/2)*(c + d*x^2), x)

sympy [B] time = 14.71, size = 253, normalized size = 2.14

$$\frac{a^5 dx}{16 b \sqrt{1 + \frac{b x^2}{a}}} + \frac{a^3 c x \sqrt{1 + \frac{b x^2}{a}}}{2} + \frac{a^3 c x}{8 \sqrt{1 + \frac{b x^2}{a}}} + \frac{17 a^3 dx^3}{48 \sqrt{1 + \frac{b x^2}{a}}} + \frac{3 \sqrt{a} b c x^3}{8 \sqrt{1 + \frac{b x^2}{a}}} + \frac{11 \sqrt{a} b d x^5}{24 \sqrt{1 + \frac{b x^2}{a}}} - \frac{a^3 d \operatorname{asinh}\left(\frac{\sqrt{b} x}{\sqrt{a}}\right)}{16 b^{\frac{3}{2}}} + \frac{3 a^2 c \operatorname{asinh}\left(\frac{\sqrt{b} x}{\sqrt{a}}\right)}{8 \sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)*(d*x**2+c), x)

[Out] a**(5/2)*d*x/(16*b*sqrt(1 + b*x**2/a)) + a**(3/2)*c*x*sqrt(1 + b*x**2/a)/2 + a**(3/2)*c*x/(8*sqrt(1 + b*x**2/a)) + 17*a**(3/2)*d*x**3/(48*sqrt(1 + b*x**2/a)) + 3*sqrt(a)*b*c*x**3/(8*sqrt(1 + b*x**2/a)) + 11*sqrt(a)*b*d*x**5/(24*sqrt(1 + b*x**2/a)) - a**3*d*asinh(sqrt(b)*x/sqrt(a))/(16*b**(3/2)) + 3*a**2*c*asinh(sqrt(b)*x/sqrt(a))/(8*sqrt(b)) + b**2*c*x**5/(4*sqrt(a)*sqrt(1 + b*x**2/a)) + b**2*d*x**7/(6*sqrt(a)*sqrt(1 + b*x**2/a))

3.56 $\int (a + bx^2)^{3/2} dx$

Optimal. Leaf size=65

$$\frac{3a^2 \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{8\sqrt{b}} + \frac{3}{8}ax\sqrt{a+bx^2} + \frac{1}{4}x(a+bx^2)^{3/2}$$

[Out] $1/4*x*(b*x^2+a)^{(3/2)}+3/8*a^2*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(1/2)}+3/8*a*x*(b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {195, 217, 206}

$$\frac{3a^2 \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{8\sqrt{b}} + \frac{3}{8}ax\sqrt{a+bx^2} + \frac{1}{4}x(a+bx^2)^{3/2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2), x]

[Out] $(3*a*x*\operatorname{Sqrt}[a + b*x^2])/8 + (x*(a + b*x^2)^{(3/2)})/4 + (3*a^2*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(8*\operatorname{Sqrt}[b])$

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^{3/2} dx &= \frac{1}{4}x(a + bx^2)^{3/2} + \frac{1}{4}(3a) \int \sqrt{a + bx^2} dx \\ &= \frac{3}{8}ax\sqrt{a + bx^2} + \frac{1}{4}x(a + bx^2)^{3/2} + \frac{1}{8}(3a^2) \int \frac{1}{\sqrt{a + bx^2}} dx \\ &= \frac{3}{8}ax\sqrt{a + bx^2} + \frac{1}{4}x(a + bx^2)^{3/2} + \frac{1}{8}(3a^2) \operatorname{Subst}\left(\int \frac{1}{1 - bx^2} dx, x, \frac{x}{\sqrt{a + bx^2}}\right) \\ &= \frac{3}{8}ax\sqrt{a + bx^2} + \frac{1}{4}x(a + bx^2)^{3/2} + \frac{3a^2 \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{8\sqrt{b}} \end{aligned}$$

Mathematica [A] time = 0.09, size = 65, normalized size = 1.00

$$\frac{1}{8} \sqrt{a + bx^2} \left(\frac{3a^{3/2} \sinh^{-1} \left(\frac{\sqrt{b}x}{\sqrt{a}} \right)}{\sqrt{b} \sqrt{\frac{bx^2}{a} + 1}} + 5ax + 2bx^3 \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2), x]

[Out] (Sqrt[a + b*x^2]*(5*a*x + 2*b*x^3 + (3*a^(3/2)*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[b]*Sqrt[1 + (b*x^2)/a]))) / 8

fricas [A] time = 0.58, size = 124, normalized size = 1.91

$$\left[\frac{3a^2\sqrt{b} \log\left(-2bx^2 - 2\sqrt{bx^2+a}\sqrt{b}x - a\right) + 2(2b^2x^3 + 5abx)\sqrt{bx^2+a}}{16b}, -\frac{3a^2\sqrt{-b} \arctan\left(\frac{\sqrt{-b}x}{\sqrt{bx^2+a}}\right) - (2b^2x^3 + 5abx)\sqrt{bx^2+a}}{8b} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2), x, algorithm="fricas")

[Out] [1/16*(3*a^2*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 2*(2*b^2*x^3 + 5*a*b*x)*sqrt(b*x^2 + a))/b, -1/8*(3*a^2*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (2*b^2*x^3 + 5*a*b*x)*sqrt(b*x^2 + a))/b]

giac [A] time = 0.61, size = 49, normalized size = 0.75

$$\frac{1}{8} (2bx^2 + 5a)\sqrt{bx^2 + a}x - \frac{3a^2 \log\left(\left|-\sqrt{b}x + \sqrt{bx^2 + a}\right|\right)}{8\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2), x, algorithm="giac")

[Out] 1/8*(2*b*x^2 + 5*a)*sqrt(b*x^2 + a)*x - 3/8*a^2*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/sqrt(b)

maple [A] time = 0.00, size = 51, normalized size = 0.78

$$\frac{3a^2 \ln\left(\sqrt{b}x + \sqrt{bx^2 + a}\right)}{8\sqrt{b}} + \frac{3\sqrt{bx^2 + a}ax}{8} + \frac{(bx^2 + a)^{\frac{3}{2}}x}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2), x)

[Out] 1/4*(b*x^2+a)^(3/2)*x+3/8*(b*x^2+a)^(1/2)*a*x+3/8*a^2/b^(1/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))

maxima [A] time = 1.34, size = 43, normalized size = 0.66

$$\frac{1}{4} (bx^2 + a)^{\frac{3}{2}}x + \frac{3}{8} \sqrt{bx^2 + a}ax + \frac{3a^2 \operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{8\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2), x, algorithm="maxima")

[Out] $\frac{1}{4}(bx^2 + a)^{3/2}x + \frac{3}{8}\sqrt{bx^2 + a}ax + \frac{3}{8}a^2\operatorname{arcsinh}\left(\frac{bx}{\sqrt{a}}\right)/\sqrt{b}$

mupad [B] time = 4.71, size = 37, normalized size = 0.57

$$\frac{x(bx^2 + a)^{3/2} {}_2F_1\left(-\frac{3}{2}, \frac{1}{2}; \frac{3}{2}; -\frac{bx^2}{a}\right)}{\left(\frac{bx^2}{a} + 1\right)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^(3/2), x)`

[Out] $(x(a + bx^2)^{3/2}\operatorname{hypergeom}([-3/2, 1/2], 3/2, -(bx^2)/a))/((bx^2)/a + 1)^{3/2}$

sympy [A] time = 2.91, size = 70, normalized size = 1.08

$$\frac{5a^3x\sqrt{1 + \frac{bx^2}{a}}}{8} + \frac{\sqrt{a}bx^3\sqrt{1 + \frac{bx^2}{a}}}{4} + \frac{3a^2\operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**(3/2), x)`

[Out] $5a^{3/2}x\sqrt{1 + bx^2/a}/8 + \sqrt{a}bx^3\sqrt{1 + bx^2/a}/4 + 3a^2\operatorname{asinh}(\sqrt{b}x/\sqrt{a})/(8\sqrt{b})$

$$3.57 \quad \int \frac{(a+bx^2)^{3/2}}{c+dx^2} dx$$

Optimal. Leaf size=113

$$\frac{(bc-ad)^{3/2} \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d^2} - \frac{\sqrt{b}(2bc-3ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2d^2} + \frac{bx\sqrt{a+bx^2}}{2d}$$

[Out] $-1/2*(-3*a*d+2*b*c)*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})*b^{(1/2)}/d^2+(-a*d+b*c)^{(3/2)*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})}/d^2/c^{(1/2)}+1/2*b*x*(b*x^2+a)^{(1/2)}/d$

Rubi [A] time = 0.11, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {416, 523, 217, 206, 377, 208}

$$\frac{(bc-ad)^{3/2} \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d^2} - \frac{\sqrt{b}(2bc-3ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2d^2} + \frac{bx\sqrt{a+bx^2}}{2d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)/(c + d*x^2), x]

[Out] $(b*x*\operatorname{Sqrt}[a + b*x^2])/(2*d) - (\operatorname{Sqrt}[b]*(2*b*c - 3*a*d)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(2*d^2) + ((b*c - a*d)^{(3/2)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(2*d^2)$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q) + 1)), x] + Dist[1/(b*(n*(p+q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q) + 1) - a*d) + d*(b*c*(n*(p+2*q-1) + 1) - a*d*(n*(q-1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a

, b, c, d, n, p, q, x]

Rule 523

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^n)*Sqrt[(c_) + (d_.)*(x_)^n]), x_Symbol] :> Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{3/2}}{c + dx^2} dx &= \frac{bx\sqrt{a + bx^2}}{2d} + \frac{\int \frac{-a(bc-2ad)-b(2bc-3ad)x^2}{\sqrt{a+bx^2}(c+dx^2)} dx}{2d} \\ &= \frac{bx\sqrt{a + bx^2}}{2d} - \frac{(b(2bc - 3ad)) \int \frac{1}{\sqrt{a+bx^2}} dx}{2d^2} + \frac{(bc - ad)^2 \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{d^2} \\ &= \frac{bx\sqrt{a + bx^2}}{2d} - \frac{(b(2bc - 3ad)) \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{2d^2} + \frac{(bc - ad)^2 \text{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{d^2} \\ &= \frac{bx\sqrt{a + bx^2}}{2d} - \frac{\sqrt{b}(2bc - 3ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2d^2} + \frac{(bc - ad)^{3/2} \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d^2} \end{aligned}$$

Mathematica [A] time = 0.20, size = 110, normalized size = 0.97

$$\frac{\sqrt{b}(3ad - 2bc) \log\left(\sqrt{b}\sqrt{a + bx^2} + bx\right) + \frac{2(ad-bc)^{3/2} \tan^{-1}\left(\frac{x\sqrt{ad-bc}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}} + bdx\sqrt{a + bx^2}}{2d^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)/(c + d*x^2), x]

[Out] (b*d*x*Sqrt[a + b*x^2] + (2*(-(b*c) + a*d)^(3/2)*ArcTan[(Sqrt[-(b*c) + a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/Sqrt[c] + Sqrt[b]*(-2*b*c + 3*a*d)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(2*d^2)

fricas [A] time = 0.83, size = 721, normalized size = 6.38

$$\frac{2\sqrt{bx^2 + a} bdx - (2bc - 3ad)\sqrt{b} \log\left(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{b}x - a\right) - (bc - ad)\sqrt{\frac{bc-ad}{c}} \log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4}{c}\right)}{4d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b*x^2 + a)*b*d*x - (2*b*c - 3*a*d)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - (b*c - a*d)*sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*sqrt(b*x^2 + a)*sqrt((b*c - a*d)/c)))/(d^2*x^4 + 2*c*d*x^2 + c^2))/d^2, 1/4*(2*sqrt(b*x^2 + a)*b*d*x + 2*(2*b*c - 3*a*d)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (b*c - a*d)*sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c

$$+(-c*d)^{(1/2)/d}^2*b-2*(-c*d)^{(1/2)*(x+(-c*d)^{(1/2)/d})*b/d+(a*d-b*c)/d}^{(1/2)}/(x+(-c*d)^{(1/2)/d}))*a*b*c+1/2/(-c*d)^{(1/2)/d}^2/((a*d-b*c)/d)^{(1/2)*\ln((-2*(-c*d)^{(1/2)*(x+(-c*d)^{(1/2)/d})*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)*((x+(-c*d)^{(1/2)/d})^2*b-2*(-c*d)^{(1/2)*(x+(-c*d)^{(1/2)/d})*b/d+(a*d-b*c)/d}^{(1/2)))/(x+(-c*d)^{(1/2)/d}))*b^2*c^2}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/2)/(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{3/2}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)/(c + d*x^2),x)

[Out] int((a + b*x^2)^(3/2)/(c + d*x^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{3}{2}}}{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)/(d*x**2+c),x)

[Out] Integral((a + b*x**2)**(3/2)/(c + d*x**2), x)

$$3.58 \quad \int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^2} dx$$

Optimal. Leaf size=131

$$\frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{d^2} - \frac{\sqrt{bc-ad}(ad+2bc) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}d^2} - \frac{x\sqrt{a+bx^2}(bc-ad)}{2cd(c+dx^2)}$$

[Out] $b^{3/2} \operatorname{arctanh}(x \cdot b^{1/2} / (b \cdot x^2 + a)^{1/2}) / d^2 - 1/2 \cdot (a \cdot d + 2 \cdot b \cdot c) \cdot \operatorname{arctanh}(x \cdot (-a \cdot d + b \cdot c)^{1/2} / c^{1/2} / (b \cdot x^2 + a)^{1/2}) \cdot (-a \cdot d + b \cdot c)^{1/2} / c^{3/2} / d^2 - 1/2 \cdot (-a \cdot d + b \cdot c) \cdot x \cdot (b \cdot x^2 + a)^{1/2} / c \cdot d / (d \cdot x^2 + c)$

Rubi [A] time = 0.09, antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {413, 523, 217, 206, 377, 208}

$$\frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{d^2} - \frac{\sqrt{bc-ad}(ad+2bc) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}d^2} - \frac{x\sqrt{a+bx^2}(bc-ad)}{2cd(c+dx^2)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)/(c + d*x^2)^2, x]

[Out] $-\frac{(b \cdot c - a \cdot d) \cdot x \cdot \operatorname{Sqrt}[a + b \cdot x^2]}{(2 \cdot c \cdot d \cdot (c + d \cdot x^2))} + \frac{b^{3/2} \cdot \operatorname{ArcTanh}[\frac{\operatorname{Sqrt}[b] \cdot x}{\operatorname{Sqrt}[a + b \cdot x^2]}]}{d^2} - \frac{(\operatorname{Sqrt}[b \cdot c - a \cdot d] \cdot (2 \cdot b \cdot c + a \cdot d) \cdot \operatorname{ArcTanh}[\frac{\operatorname{Sqrt}[b \cdot c - a \cdot d] \cdot x}{\operatorname{Sqrt}[c] \cdot \operatorname{Sqrt}[a + b \cdot x^2]}])}{(2 \cdot c^{3/2} \cdot d^2)}$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]]/(Rt[a, 2]*Rt[-b, 2])), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+q)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,

0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 523

Int[((e_) + (f_)*(x_)^(n_))/(((a_) + (b_)*(x_)^(n_))*Sqrt[(c_) + (d_)*(x_)^(n_)], x_Symbol] :> Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{3/2}}{(c + dx^2)^2} dx &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{2cd(c + dx^2)} + \frac{\int \frac{a(bc+ad)+2b^2cx^2}{\sqrt{a+bx^2}(c+dx^2)} dx}{2cd} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{2cd(c + dx^2)} + \frac{b^2 \int \frac{1}{\sqrt{a+bx^2}} dx}{d^2} - \frac{((bc - ad)(2bc + ad)) \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{2cd^2} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{2cd(c + dx^2)} + \frac{b^2 \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{d^2} - \frac{((bc - ad)(2bc + ad)) \text{Subst}\left(\int \frac{1}{c - dx^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{2cd^2} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{2cd(c + dx^2)} + \frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{d^2} - \frac{\sqrt{bc - ad}(2bc + ad) \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}d^2} \end{aligned}$$

Mathematica [A] time = 0.13, size = 142, normalized size = 1.08

$$\frac{(a^2d^2+abcd-2b^2c^2) \tan^{-1}\left(\frac{x\sqrt{ad-bc}}{\sqrt{c}\sqrt{a+bx^2}}\right) + 2b^{3/2} \log\left(\sqrt{b}\sqrt{a+bx^2} + bx\right) - \frac{dx\sqrt{a+bx^2}(bc-ad)}{c(c+dx^2)}}{2d^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)/(c + d*x^2)^2,x]

[Out] (-((d*(b*c - a*d)*x*Sqrt[a + b*x^2])/(c*(c + d*x^2))) + ((-2*b^2*c^2 + a*b*c*d + a^2*d^2)*ArcTan[(Sqrt[-(b*c) + a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(c^(3/2)*Sqrt[-(b*c) + a*d]) + 2*b^(3/2)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(2*d^2)

fricas [A] time = 0.97, size = 907, normalized size = 6.92

$$\left[\frac{4(bcd - ad^2)\sqrt{bx^2 + a}x - 4(bcdx^2 + bc^2)\sqrt{b} \log(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{b}x - a) - (2bc^2 + acd + (2bcd + ad^2))}{8(cd^3x^2 + c^2d^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] [-1/8*(4*(b*c*d - a*d^2)*sqrt(b*x^2 + a)*x - 4*(b*c*d*x^2 + b*c^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - (2*b*c^2 + a*c*d + (2*b*c*d + a*d^2)*x^2)*sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a

$$\begin{aligned} & c*d*x^3)*\text{sqrt}(b*x^2 + a)*\text{sqrt}((b*c - a*d)/c))/(d^2*x^4 + 2*c*d*x^2 + c^2)) \\ &)/(c*d^3*x^2 + c^2*d^2), -1/8*(4*(b*c*d - a*d^2)*\text{sqrt}(b*x^2 + a)*x + 8*(b*c \\ & *d*x^2 + b*c^2)*\text{sqrt}(-b)*\text{arctan}(\text{sqrt}(-b)*x/\text{sqrt}(b*x^2 + a)) - (2*b*c^2 + a* \\ & c*d + (2*b*c*d + a*d^2)*x^2)*\text{sqrt}((b*c - a*d)/c)*\log(((8*b^2*c^2 - 8*a*b*c* \\ & d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (\\ & 2*b*c^2 - a*c*d)*x^3)*\text{sqrt}(b*x^2 + a)*\text{sqrt}((b*c - a*d)/c))/(d^2*x^4 + 2*c*d \\ & *x^2 + c^2)))/(c*d^3*x^2 + c^2*d^2), -1/4*(2*(b*c*d - a*d^2)*\text{sqrt}(b*x^2 + a) \\ &)*x - (2*b*c^2 + a*c*d + (2*b*c*d + a*d^2)*x^2)*\text{sqrt}(-(b*c - a*d)/c)*\text{arctan} \\ & (1/2*((2*b*c - a*d)*x^2 + a*c)*\text{sqrt}(b*x^2 + a)*\text{sqrt}(-(b*c - a*d)/c)/((b^2*c \\ & - a*b*d)*x^3 + (a*b*c - a^2*d)*x)) - 2*(b*c*d*x^2 + b*c^2)*\text{sqrt}(b)*\log(-2* \\ & b*x^2 - 2*\text{sqrt}(b*x^2 + a)*\text{sqrt}(b)*x - a))/(c*d^3*x^2 + c^2*d^2), -1/4*(2*(b \\ & *c*d - a*d^2)*\text{sqrt}(b*x^2 + a)*x + 4*(b*c*d*x^2 + b*c^2)*\text{sqrt}(-b)*\text{arctan}(\text{sq \\ & rt}(-b)*x/\text{sqrt}(b*x^2 + a)) - (2*b*c^2 + a*c*d + (2*b*c*d + a*d^2)*x^2)*\text{sqrt}(- \\ & (b*c - a*d)/c)*\text{arctan}(1/2*((2*b*c - a*d)*x^2 + a*c)*\text{sqrt}(b*x^2 + a)*\text{sqrt}(- \\ & (b*c - a*d)/c)/((b^2*c - a*b*d)*x^3 + (a*b*c - a^2*d)*x)))/(c*d^3*x^2 + c^2* \\ & d^2)] \end{aligned}$$

giac [B] time = 0.68, size = 317, normalized size = 2.42

$$\frac{b^{\frac{3}{2}} \log\left(\left(\sqrt{b}x - \sqrt{bx^2 + a}\right)^2\right)}{2d^2} + \frac{\left(2b^{\frac{5}{2}}c^2 - ab^{\frac{3}{2}}cd - a^2\sqrt{b}d^2\right) \arctan\left(\frac{\left(\sqrt{b}x - \sqrt{bx^2 + a}\right)^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{2\sqrt{-b^2c^2 + abcd}cd^2} - \frac{2\left(\sqrt{b}x - \sqrt{bx^2 + a}\right)}{\left(\sqrt{b}x - \sqrt{bx^2 + a}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^2,x, algorithm="giac")

[Out] $-1/2*b^{(3/2)}*\log((\text{sqrt}(b)*x - \text{sqrt}(b*x^2 + a))^2)/d^2 + 1/2*(2*b^{(5/2)}*c^2 - a*b^{(3/2)}*c*d - a^2*\text{sqrt}(b)*d^2)*\text{arctan}(1/2*((\text{sqrt}(b)*x - \text{sqrt}(b*x^2 + a))^2*d + 2*b*c - a*d)/\text{sqrt}(-b^2*c^2 + a*b*c*d))/(\text{sqrt}(-b^2*c^2 + a*b*c*d)*c*d^2) - (2*(\text{sqrt}(b)*x - \text{sqrt}(b*x^2 + a))^2*b^{(5/2)}*c^2 - 3*(\text{sqrt}(b)*x - \text{sqrt}(b*x^2 + a))^2*a*b^{(3/2)}*c*d + (\text{sqrt}(b)*x - \text{sqrt}(b*x^2 + a))^2*a^2*\text{sqrt}(b)*d^2 + a^2*b^{(3/2)}*c*d - a^3*\text{sqrt}(b)*d^2)/(((\text{sqrt}(b)*x - \text{sqrt}(b*x^2 + a))^4*d + 4*(\text{sqrt}(b)*x - \text{sqrt}(b*x^2 + a))^2*b*c - 2*(\text{sqrt}(b)*x - \text{sqrt}(b*x^2 + a))^2*a*d + a^2*d)*c*d^2)$

maple [B] time = 0.02, size = 4621, normalized size = 35.27

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)/(d*x^2+c)^2,x)

[Out] $-3/8/c*b/(a*d-b*c)*a*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x+1/2/(-c*d)^{(1/2)}/d/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d)*a*b-1/4/(-c*d)^{(1/2)}*c/d^2/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d)*b^2-1/4/c/d*(-c*d)^{(1/2)}*b/(a*d-b*c)*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}-3/8/c*b/(a*d-b*c)*a*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x-1/2/(-c*d)^{(1/2)}/d/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d)*a*b+1/4/(-c*d)^{(1/2)}*c/d^2/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})$

$$\begin{aligned}
& d^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(\\
& x+(-c*d)^{(1/2)}/d))^{*b^2+1/4/c/d*(-c*d)^{(1/2)*b/(a*d-b*c)*((x+(-c*d)^{(1/2)}/d) \\
& ^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)+1/4/c/(a*d-b* \\
& c)/(x+(-c*d)^{(1/2)}/d)*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/ \\
&)/d)*b/d+(a*d-b*c)/d)^{(5/2)-1/4/(-c*d)^{(1/2)}/c*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(- \\
& c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)*a+1/4/(-c*d)^{(1/2)}/d*(\\
& (x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(\\
& 1/2)*b+1/4/(-c*d)^{(1/2)}/c*((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/ \\
&)/d)*b/d+(a*d-b*c)/d)^{(1/2)*a-1/4/(-c*d)^{(1/2)}/d*((x+(-c*d)^{(1/2)}/d)^{2*b \\
& +2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)*b+1/4/c/(a*d-b*c) \\
&)/(x+(-c*d)^{(1/2)}/d)*((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/ \\
&)/d)*b/d+(a*d-b*c)/d)^{(5/2)-3/4/c/d*(-c*d)^{(1/2)*b/(a*d-b*c)/((a*d-b*c)/d)^{(\\
& 1/2)*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/ \\
& d)^{(1/2)*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d \\
& -b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))*a^2+3/4/c/d*(-c*d)^{(1/2)*b/(a*d-b*c)/((\\
& a*d-b*c)/d)^{(1/2)*\ln((2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2 \\
& *((a*d-b*c)/d)^{(1/2)*((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/ \\
&)/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))*a^2+3/8/d*b^2/(a*d-b*c)*((x \\
& +(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/ \\
& 2)*x+9/8/d*b^(3/2)/(a*d-b*c)*\ln(((x+(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)*b/d)/b^(\\
& 1/2)+((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c) \\
&)/d)^{(1/2))*a-1/4/d^2*b^(3/2)*\ln(((x+(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)*b/d)/b^(\\
& 1/2)+((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b* \\
& c)/d)^{(1/2))+1/12/(-c*d)^{(1/2)}/c*((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x \\
& +(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)-1/4/d^2*b^(3/2)*\ln(((x+(-c*d)^{(1/2)}/ \\
& d)*b+(-c*d)^{(1/2)*b/d)/b^(1/2)+((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+ \\
& (-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))-1/12/(-c*d)^{(1/2)}/c*((x+(-c*d)^{(1/2)}/ \\
& d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)-3/4*c/d^2*b \\
& ^{(5/2)/(a*d-b*c)*\ln(((x+(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)*b/d)/b^(1/2)+((x+(-c \\
& *d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))- \\
& 1/4/c*b/(a*d-b*c)*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d) \\
&)*b/d+(a*d-b*c)/d)^{(3/2)*x-3/8/c*b^(1/2)/(a*d-b*c)*a^2*\ln(((x+(-c*d)^{(1/2)}/d) \\
&)*b+(-c*d)^{(1/2)*b/d)/b^(1/2)+((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c \\
& *d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))+3/8/d*b^2/(a*d-b*c)*((x+(-c*d)^{(1/2)}/d) \\
&)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)*x-1/4/c*b/(a \\
& *d-b*c)*((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d- \\
& b*c)/d)^{(3/2)*x-3/8/c*b^(1/2)/(a*d-b*c)*a^2*\ln(((x+(-c*d)^{(1/2)}/d)*b+(-c*d) \\
& ^{(1/2)*b/d)/b^(1/2)+((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/ \\
& d)*b/d+(a*d-b*c)/d)^{(1/2))+1/8/c*b/d*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2) \\
&)*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)*x+3/8/c/d*b^(1/2)*\ln(((x+(-c*d)^{(1/2)}/ \\
& d)*b+(-c*d)^{(1/2)*b/d)/b^(1/2)+((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2) \\
&)*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*a+1/4/(-c*d)^{(1/2)}/c/((a*d-b*c) \\
& /d)^{(1/2)*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d- \\
& b*c)/d)^{(1/2)*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d \\
& +(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))*a^2+1/8/c*b/d*((x+(-c*d)^{(1/2)}/d)^{ \\
& 2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)*x+3/8/c/d*b^(1 \\
& /2)*\ln(((x+(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)*b/d)/b^(1/2)+((x+(-c*d)^{(1/2)}/d)^{ \\
& 2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*a-1/4/(-c*d)^{(\\
& 1/2)}/c/((a*d-b*c)/d)^{(1/2)*\ln((2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a* \\
& d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)*((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+ \\
& (-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))*a^2+9/8/d*b^(3/2) \\
& /((a*d-b*c)*\ln(((x+(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)*b/d)/b^(1/2)+((x+(-c*d)^{(1 \\
& /2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*a-3/4/ \\
& d^2*(-c*d)^{(1/2)*b^2/(a*d-b*c)*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(- \\
& c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)+3/4/d^2*(-c*d)^{(1/2)*b^2/(a*d-b*c)*((x \\
& +(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/ \\
& 2)-3/4*c/d^2*b^(5/2)/(a*d-b*c)*\ln(((x+(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)*b/d)/b \\
& ^{(1/2)+((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b \\
& *c)/d)^{(1/2))-3/4/c/d*(-c*d)^{(1/2)*b/(a*d-b*c)*((x+(-c*d)^{(1/2)}/d)^{2*b+2*(-}
\end{aligned}$$

$c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*a-3/2/d^2*(-c*d)^{(1/2)}$
 $*b^2/(a*d-b*c)/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/$
 $d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}$
 $)*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))*a+3/4*c/d^3$
 $*(-c*d)^{(1/2)}*b^3/(a*d-b*c)/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)$
 $d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^2*b$
 $+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/$
 $d))+3/4/c/d*(-c*d)^{(1/2)}*b/(a*d-b*c)*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}$
 $*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*a+3/2/d^2*(-c*d)^{(1/2)}*b^2/(a*d-$
 $b*c)/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-$
 $b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*$
 $d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))*a-3/4*c/d^3*(-c*d)^{(1/2)}$
 $*b^3/(a*d-b*c)/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-$
 $b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)$
 $)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/2)/(d*x^2 + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{3/2}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)/(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(3/2)/(c + d*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{3}{2}}}{(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)/(d*x**2+c)**2,x)

[Out] Integral((a + b*x**2)**(3/2)/(c + d*x**2)**2, x)

$$3.59 \quad \int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^3} dx$$

Optimal. Leaf size=113

$$\frac{3a^2 \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}\sqrt{bc-ad}} + \frac{3ax\sqrt{a+bx^2}}{8c^2(c+dx^2)} + \frac{x(a+bx^2)^{3/2}}{4c(c+dx^2)^2}$$

[Out] 1/4*x*(b*x^2+a)^(3/2)/c/(d*x^2+c)^2+3/8*a^2*arctanh(x*(-a*d+b*c)^(1/2)/c^(1/2)/(b*x^2+a)^(1/2))/c^(5/2)/(-a*d+b*c)^(1/2)+3/8*a*x*(b*x^2+a)^(1/2)/c^2/(d*x^2+c)

Rubi [A] time = 0.06, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {378, 377, 208}

$$\frac{3a^2 \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}\sqrt{bc-ad}} + \frac{3ax\sqrt{a+bx^2}}{8c^2(c+dx^2)} + \frac{x(a+bx^2)^{3/2}}{4c(c+dx^2)^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)/(c + d*x^2)^3,x]

[Out] (x*(a + b*x^2)^(3/2))/(4*c*(c + d*x^2)^2) + (3*a*x*Sqrt[a + b*x^2])/(8*c^2*(c + d*x^2)) + (3*a^2*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(8*c^(5/2)*Sqrt[b*c - a*d])

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] - Dist[(c*q)/(a*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+1)+1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^3} dx &= \frac{x(a+bx^2)^{3/2}}{4c(c+dx^2)^2} + \frac{(3a) \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^2} dx}{4c} \\
&= \frac{x(a+bx^2)^{3/2}}{4c(c+dx^2)^2} + \frac{3ax\sqrt{a+bx^2}}{8c^2(c+dx^2)} + \frac{(3a^2) \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{8c^2} \\
&= \frac{x(a+bx^2)^{3/2}}{4c(c+dx^2)^2} + \frac{3ax\sqrt{a+bx^2}}{8c^2(c+dx^2)} + \frac{(3a^2) \text{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{8c^2} \\
&= \frac{x(a+bx^2)^{3/2}}{4c(c+dx^2)^2} + \frac{3ax\sqrt{a+bx^2}}{8c^2(c+dx^2)} + \frac{3a^2 \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}\sqrt{bc-ad}}
\end{aligned}$$

Mathematica [A] time = 0.69, size = 163, normalized size = 1.44

$$\frac{x\sqrt{a+bx^2} \left(\frac{\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} (5ac+3adx^2+2bcx^2)}{(c+dx^2)\sqrt{\frac{dx^2}{c}+1}} + \frac{3a \sin^{-1}\left(\frac{\sqrt{x^2\left(\frac{d}{c}-\frac{b}{a}\right)}}{\sqrt{\frac{dx^2}{c}+1}}\right)}{\sqrt{\frac{x^2(ad-bc)}{ac}}}}{8c^3\sqrt{\frac{bx^2}{a}+1}} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(3/2)/(c + d*x^2)^3,x]

[Out] (x*sqrt[a + b*x^2]*((sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*(5*a*c + 2*b*c*x^2 + 3*a*d*x^2))/((c + d*x^2)*sqrt[1 + (d*x^2)/c]) + (3*a*ArcSin[sqrt[(-b/a) + d/c]*x^2]/sqrt[1 + (d*x^2)/c]]/sqrt[(-b*c) + a*d]*x^2)/(8*c^3*sqrt[1 + (b*x^2)/a])

fricas [B] time = 0.78, size = 526, normalized size = 4.65

$$\left[\frac{3(a^2d^2x^4 + 2a^2cdx^2 + a^2c^2)\sqrt{bc^2 - acd} \log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4 + a^2c^2 + 2(4abc^2 - 3a^2cd)x^2 + 4((2bc - ad)x^3 + acx)\sqrt{bc^2 - acd}}{d^2x^4 + 2cdx^2 + c^2}\right)}{32(bc^6 - ac^5d + (bc^4d^2 - ac^3d^3)x^4 + 2(bc^5d - ac^4d^2)x^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [1/32*(3*(a^2*d^2*x^4 + 2*a^2*c*d*x^2 + a^2*c^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/((d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*((2*b^2*c^3 + a*b*c^2*d - 3*a^2*c*d^2)*x^3 + 5*(a*b*c^3 - a^2*c^2*d)*x)*sqrt(b*x^2 + a))/(b*c^6 - a*c^5*d + (b*c^4*d^2 - a*c^3*d^3)*x^4 + 2*(b*c^5*d - a*c^4*d^2)*x^2), -1/16*(3*(a^2*d^2*x^4 + 2*a^2*c*d*x^2 + a^2*c^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d))*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x) - 2*((2*b^2*c^3 + a*b*c^2*d - 3*a^2*c*d^2)*x^3 + 5*(a*b*c^3 - a^2*c^2*d)*x)*sqrt(b*x^2 + a))/(b*c^6 - a*c^5*d + (b*c^4*d^2 - a*c^3*d^3)*x^4 + 2*(b*c^5*d - a*c^4*d^2)*x^2)]

giac [B] time = 3.72, size = 451, normalized size = 3.99

$$\frac{3a^2\sqrt{b}\arctan\left(\frac{(\sqrt{bx}-\sqrt{bx^2+a})^{d+2}bc-ad}{2\sqrt{-b^2c^2+abcd}}\right)}{8\sqrt{-b^2c^2+abcd}c^2} + \frac{8(\sqrt{bx}-\sqrt{bx^2+a})^6 b^{\frac{5}{2}}c^2d - 3(\sqrt{bx}-\sqrt{bx^2+a})^6 a^2\sqrt{b}d^3 + 16(\sqrt{bx}-\sqrt{bx^2+a})^4 a^2\sqrt{b}d^2 - 9(\sqrt{bx}-\sqrt{bx^2+a})^4 a^3\sqrt{b}d^3 + 8(\sqrt{bx}-\sqrt{bx^2+a})^2 a^2 b^{\frac{5}{2}}c^2d + 16(\sqrt{bx}-\sqrt{bx^2+a})^2 a^3 b^{\frac{3}{2}}c^2d^2 - 9(\sqrt{bx}-\sqrt{bx^2+a})^2 a^4\sqrt{b}d^3 + 2a^4 b^{\frac{3}{2}}c^2d^2 + 3a^5\sqrt{b}d^3}{((\sqrt{bx}-\sqrt{bx^2+a})^4 d + 4(\sqrt{bx}-\sqrt{bx^2+a})^2 bc - 2(\sqrt{bx}-\sqrt{bx^2+a})^2 ad + a^2 d)^2 c^2 d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^3,x, algorithm="giac")

[Out] -3/8*a^2*sqrt(b)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/sqrt(-b^2*c^2 + a*b*c*d)*c^2 + 1/4*(8*(sqrt(b)*x - sqrt(b*x^2 + a))^6*b^(5/2)*c^2*d - 3*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^2*sqrt(b)*d^3 + 16*(sqrt(b)*x - sqrt(b*x^2 + a))^4*b^(7/2)*c^3 + 8*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a*b^(5/2)*c^2*d - 18*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^2*b^(3/2)*c*d^2 + 9*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^3*sqrt(b)*d^3 + 8*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^2*b^(5/2)*c^2*d + 16*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^3*b^(3/2)*c*d^2 - 9*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^4*sqrt(b)*d^3 + 2*a^4*b^(3/2)*c*d^2 + 3*a^5*sqrt(b)*d^3)/(((sqrt(b)*x - sqrt(b*x^2 + a))^4*d + 4*(sqrt(b)*x - sqrt(b*x^2 + a))^2*b*c - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*d + a^2*d)^2*c^2*d^2)

maple [B] time = 0.02, size = 9059, normalized size = 80.17

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)/(d*x^2+c)^3,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^3,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/2)/(d*x^2 + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{3/2}}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)/(c + d*x^2)^3,x)

[Out] int((a + b*x^2)^(3/2)/(c + d*x^2)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{3}{2}}}{(c + dx^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)**(3/2)/(d*x**2+c)**3,x)
```

```
[Out] Integral((a + b*x**2)**(3/2)/(c + d*x**2)**3, x)
```

$$3.60 \quad \int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^4} dx$$

Optimal. Leaf size=199

$$\frac{a^2(6bc - 5ad) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{16c^{7/2}(bc - ad)^{3/2}} + \frac{ax\sqrt{a+bx^2}(6bc - 5ad)}{16c^3(c+dx^2)(bc - ad)} + \frac{x(a+bx^2)^{3/2}(6bc - 5ad)}{24c^2(c+dx^2)^2(bc - ad)} - \frac{dx(a+bx^2)^{5/2}}{6c(c+dx^2)^3(bc - ad)}$$

[Out] $-1/6*d*x*(b*x^2+a)^{(5/2)}/c/(-a*d+b*c)/(d*x^2+c)^3+1/24*(-5*a*d+6*b*c)*x*(b*x^2+a)^{(3/2)}/c^2/(-a*d+b*c)/(d*x^2+c)^2+1/16*a^2*(-5*a*d+6*b*c)*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})/c^{(7/2)}/(-a*d+b*c)^{(3/2)}+1/16*a*(-5*a*d+6*b*c)*x*(b*x^2+a)^{(1/2)}/c^3/(-a*d+b*c)/(d*x^2+c)$

Rubi [A] time = 0.12, antiderivative size = 199, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {382, 378, 377, 208}

$$\frac{a^2(6bc - 5ad) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{16c^{7/2}(bc - ad)^{3/2}} + \frac{x(a+bx^2)^{3/2}(6bc - 5ad)}{24c^2(c+dx^2)^2(bc - ad)} + \frac{ax\sqrt{a+bx^2}(6bc - 5ad)}{16c^3(c+dx^2)(bc - ad)} - \frac{dx(a+bx^2)^{5/2}}{6c(c+dx^2)^3(bc - ad)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)/(c + d*x^2)^4, x]

[Out] $-(d*x*(a + b*x^2)^{(5/2)})/(6*c*(b*c - a*d)*(c + d*x^2)^3) + ((6*b*c - 5*a*d)*x*(a + b*x^2)^{(3/2)})/(24*c^2*(b*c - a*d)*(c + d*x^2)^2) + (a*(6*b*c - 5*a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(16*c^3*(b*c - a*d)*(c + d*x^2)) + (a^2*(6*b*c - 5*a*d)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(16*c^{(7/2)}*(b*c - a*d)^{(3/2)})$

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] - Dist[(c*q)/(a*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+1)+1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+2)+1, 0] && (LtQ[p, -1] || !LtQ

[q, -1]) && NeQ[p, -1]

Rubi steps

$$\int \frac{(a + bx^2)^{3/2}}{(c + dx^2)^4} dx = -\frac{dx (a + bx^2)^{5/2}}{6c(bc - ad)(c + dx^2)^3} + \frac{(6bc - 5ad) \int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^3} dx}{6c(bc - ad)}$$

$$= -\frac{dx (a + bx^2)^{5/2}}{6c(bc - ad)(c + dx^2)^3} + \frac{(6bc - 5ad)x (a + bx^2)^{3/2}}{24c^2(bc - ad)(c + dx^2)^2} + \frac{(a(6bc - 5ad)) \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^2} dx}{8c^2(bc - ad)}$$

$$= -\frac{dx (a + bx^2)^{5/2}}{6c(bc - ad)(c + dx^2)^3} + \frac{(6bc - 5ad)x (a + bx^2)^{3/2}}{24c^2(bc - ad)(c + dx^2)^2} + \frac{a(6bc - 5ad)x\sqrt{a + bx^2}}{16c^3(bc - ad)(c + dx^2)} + \frac{(a^2(6bc - 5ad)) \int \frac{1}{c+dx^2} dx}{8c^2(bc - ad)}$$

$$= -\frac{dx (a + bx^2)^{5/2}}{6c(bc - ad)(c + dx^2)^3} + \frac{(6bc - 5ad)x (a + bx^2)^{3/2}}{24c^2(bc - ad)(c + dx^2)^2} + \frac{a(6bc - 5ad)x\sqrt{a + bx^2}}{16c^3(bc - ad)(c + dx^2)} + \frac{(a^2(6bc - 5ad)) \int \frac{1}{c+dx^2} dx}{8c^2(bc - ad)}$$

$$= -\frac{dx (a + bx^2)^{5/2}}{6c(bc - ad)(c + dx^2)^3} + \frac{(6bc - 5ad)x (a + bx^2)^{3/2}}{24c^2(bc - ad)(c + dx^2)^2} + \frac{a(6bc - 5ad)x\sqrt{a + bx^2}}{16c^3(bc - ad)(c + dx^2)} + \frac{a^2(6bc - 5ad) \arctan\left(\frac{\sqrt{a+bx^2}}{c+dx^2}\right)}{8c^2(bc - ad)}$$

Mathematica [A] time = 0.82, size = 247, normalized size = 1.24

$$ax \left(\frac{bx^2}{a} + 1 \right) \left(\frac{3a^2(c+dx^2)^3(5ad-6bc) \tanh^{-1}\left(\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}\right)}{\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}} + c(a^3d(33c^2 + 40cdx^2 + 15d^2x^4) + a^2b(-30c^3 + 11c^2dx^2 + 32cd^2x^4 + 15d^3x^6)) + (3a^2(-6*bc + 5*a*d)*(c + d*x^2)^3 * \text{ArcTanh}\left[\frac{\sqrt{(b*c - a*d)*x^2}}{\sqrt{c*(a + b*x^2)}}\right]) / \sqrt{\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}} \right) / (48c^4(-b*c + a*d)*(a + b*x^2)^{3/2} * (c + d*x^2)^3 (ad - b^2))$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)/(c + d*x^2)^4, x]

[Out] (a*x*(1 + (b*x^2)/a)*(c*(-4*b^3*c^2*x^4*(3*c + d*x^2) - 2*a*b^2*c*x^2*(21*c^2 + 13*c*d*x^2 + 4*d^2*x^4) + a^3*d*(33*c^2 + 40*c*d*x^2 + 15*d^2*x^4) + a^2*b*(-30*c^3 + 11*c^2*d*x^2 + 32*c*d^2*x^4 + 15*d^3*x^6)) + (3*a^2*(-6*b*c + 5*a*d)*(c + d*x^2)^3*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2)))/(48*c^4*(-(b*c) + a*d)*(a + b*x^2)^(3/2)*(c + d*x^2)^3)

fricas [B] time = 1.23, size = 972, normalized size = 4.88

$$\frac{3(6a^2bc^4 - 5a^3c^3d + (6a^2bcd^3 - 5a^3d^4)x^6 + 3(6a^2bc^2d^2 - 5a^3cd^3)x^4 + 3(6a^2bc^3d - 5a^3c^2d^2)x^2)\sqrt{bc^2 - a^2}}{192}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^4,x, algorithm="fricas")

[Out] [1/192*(3*(6*a^2*b*c^4 - 5*a^3*c^3*d + (6*a^2*b*c*d^3 - 5*a^3*d^4)*x^6 + 3*(6*a^2*b*c^2*d^2 - 5*a^3*c*d^3)*x^4 + 3*(6*a^2*b*c^3*d - 5*a^3*c^2*d^2)*x^2)

) $\sqrt{b^2c^2 - a^2cd}$ $\log\left(\frac{((8b^2c^2 - 8a^2b^2cd + a^2d^2)x^4 + a^2c^2 + 2(4ab^2c^2 - 3a^2cd)x^2 + 4((2b^2c - a^2d)x^3 + acx)\sqrt{b^2c^2 - a^2cd})\sqrt{bx^2 + a}}{(d^2x^4 + 2cdx^2 + c^2)} + 4\frac{(4b^3c^4d + 4a^2b^2c^3d^2 - 23a^2b^2c^2d^3 + 15a^3c^2d^4)x^5 + 2(6b^3c^5 + 5ab^2c^4d - 31a^2b^2c^3d^2 + 20a^3c^2d^3)x^3 + 3(10ab^2c^5 - 21a^2b^2c^4d + 11a^3c^3d^2)x}{(b^2c^9 - 2ab^2c^8d + a^2c^7d^2 + (b^2c^6d^3 - 2ab^2c^5d^4 + a^2c^4d^5)x^6 + 3(b^2c^7d^2 - 2ab^2c^6d^3 + a^2c^5d^4)x^4 + 3(b^2c^8d - 2ab^2c^7d^2 + a^2c^6d^3)x^2)}\right), -\frac{1}{96}\frac{(3(6a^2b^2c^4 - 5a^3c^3d + (6a^2b^2cd^3 - 5a^3d^4)x^6 + 3(6a^2b^2cd^2 - 5a^3cd^3)x^4 + 3(6a^2b^2c^3d - 5a^3c^2d^2)x^2)\sqrt{-b^2c^2 + a^2cd})\arctan\left(\frac{1}{2}\sqrt{-b^2c^2 + a^2cd}\right)\left((2b^2c - a^2d)x^2 + ac\right)\sqrt{bx^2 + a}}{(b^2c^2 - ab^2cd)x^3 + (ab^2c^2 - a^2cd)x)} - 2\frac{((4b^3c^4d + 4a^2b^2c^3d^2 - 23a^2b^2c^2d^3 + 15a^3c^2d^4)x^5 + 2(6b^3c^5 + 5ab^2c^4d - 31a^2b^2c^3d^2 + 20a^3c^2d^3)x^3 + 3(10ab^2c^5 - 21a^2b^2c^4d + 11a^3c^3d^2)x)\sqrt{bx^2 + a}}{(b^2c^9 - 2ab^2c^8d + a^2c^7d^2 + (b^2c^6d^3 - 2ab^2c^5d^4)x^6 + 3(b^2c^7d^2 - 2ab^2c^6d^3 + a^2c^5d^4)x^4 + 3(b^2c^8d - 2ab^2c^7d^2 + a^2c^6d^3)x^2)}$

giac [B] time = 2.89, size = 919, normalized size = 4.62

$$\frac{\left(6a^2b^{\frac{3}{2}}c - 5a^3\sqrt{bd}\right)\arctan\left(\frac{\left(\sqrt{bx} - \sqrt{bx^2+a}\right)^2}{2\sqrt{-b^2c^2+abcd}}\right)}{16(bc^4 - ac^3d)\sqrt{-b^2c^2 + abcd}} - 18\left(\sqrt{bx} - \sqrt{bx^2+a}\right)^{10}a^2b^{\frac{3}{2}}cd^4 - 15\left(\sqrt{bx} - \sqrt{bx^2+a}\right)^{10}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^4,x, algorithm="giac")

[Out] $-\frac{1}{16}\frac{(6a^2b^{\frac{3}{2}}c - 5a^3\sqrt{bd})\arctan\left(\frac{1}{2}\frac{(\sqrt{bx} - \sqrt{bx^2+a})}{\sqrt{-b^2c^2 + a^2cd}}\right)}{(b^2c^4 - a^2c^3d)\sqrt{-b^2c^2 + a^2cd}} - \frac{1}{24}\frac{18(\sqrt{bx} - \sqrt{bx^2+a})^{10}a^2b^{\frac{3}{2}}cd^4 - 15(\sqrt{bx} - \sqrt{bx^2+a})^{10}a^3\sqrt{bd}d^5 - 96(\sqrt{bx} - \sqrt{bx^2+a})^8b^{\frac{9}{2}}c^4d + 96(\sqrt{bx} - \sqrt{bx^2+a})^8a^2b^{\frac{7}{2}}c^3d^2 + 180(\sqrt{bx} - \sqrt{bx^2+a})^8a^2b^{\frac{5}{2}}c^2d^3 - 240(\sqrt{bx} - \sqrt{bx^2+a})^8a^3b^{\frac{3}{2}}cd^4 + 75(\sqrt{bx} - \sqrt{bx^2+a})^8a^4\sqrt{bd}d^5 - 128(\sqrt{bx} - \sqrt{bx^2+a})^6b^{\frac{11}{2}}c^5 - 64(\sqrt{bx} - \sqrt{bx^2+a})^6a^2b^{\frac{9}{2}}c^4d + 720(\sqrt{bx} - \sqrt{bx^2+a})^6a^2b^{\frac{7}{2}}c^3d^2 - 968(\sqrt{bx} - \sqrt{bx^2+a})^6a^3b^{\frac{5}{2}}c^2d^3 + 620(\sqrt{bx} - \sqrt{bx^2+a})^6a^4b^{\frac{3}{2}}cd^4 - 150(\sqrt{bx} - \sqrt{bx^2+a})^6a^5\sqrt{bd}d^5 - 96(\sqrt{bx} - \sqrt{bx^2+a})^4a^2b^{\frac{9}{2}}c^4d - 288(\sqrt{bx} - \sqrt{bx^2+a})^4a^3b^{\frac{7}{2}}c^3d^2 + 864(\sqrt{bx} - \sqrt{bx^2+a})^4a^4b^{\frac{5}{2}}c^2d^3 - 600(\sqrt{bx} - \sqrt{bx^2+a})^4a^5b^{\frac{3}{2}}cd^4 + 150(\sqrt{bx} - \sqrt{bx^2+a})^4a^6\sqrt{bd}d^5 - 48(\sqrt{bx} - \sqrt{bx^2+a})^2a^4b^{\frac{7}{2}}c^3d^2 - 72(\sqrt{bx} - \sqrt{bx^2+a})^2a^5b^{\frac{5}{2}}c^2d^3 + 210(\sqrt{bx} - \sqrt{bx^2+a})^2a^6b^{\frac{3}{2}}cd^4 - 75(\sqrt{bx} - \sqrt{bx^2+a})^2a^7\sqrt{bd}d^5 - 4a^6b^{\frac{5}{2}}c^2d^3 - 8a^7b^{\frac{3}{2}}cd^4 + 15a^8\sqrt{bd}d^5}{(b^2c^4d^2 - a^2c^3d^3)\left((\sqrt{bx} - \sqrt{bx^2+a})^4d + 4(\sqrt{bx} - \sqrt{bx^2+a})^2b^2c - 2(\sqrt{bx} - \sqrt{bx^2+a})^2ad + a^2d\right)^3}$

maple [B] time = 0.03, size = 13766, normalized size = 69.18

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)/(d*x^2+c)^4,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{(dx^2 + c)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^4,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/2)/(d*x^2 + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{(dx^2 + c)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)/(c + d*x^2)^4,x)

[Out] int((a + b*x^2)^(3/2)/(c + d*x^2)^4, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)/(d*x**2+c)**4,x)

[Out] Timed out

$$3.61 \quad \int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^5} dx$$

Optimal. Leaf size=300

$$\frac{a^2 (35a^2d^2 - 80abcd + 48b^2c^2) \tanh^{-1} \left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}} \right)}{128c^{9/2}(bc-ad)^{5/2}} + \frac{x\sqrt{a+bx^2} (-35a^2d^2 + 24abcd + 8b^2c^2)}{192c^3d(c+dx^2)^2(bc-ad)} + \frac{x\sqrt{a+bx^2} (105a^3d^3 - 170a^2b^2cd + 40ab^2c^2d + 16b^3c^3)}{384c^4d(c+dx^2)(bc-ad)^2}$$

[Out] 1/128*a^2*(35*a^2*d^2-80*a*b*c*d+48*b^2*c^2)*arctanh(x*(-a*d+b*c)^(1/2)/c^(1/2)/(b*x^2+a)^(1/2))/c^(9/2)/(-a*d+b*c)^(5/2)-1/8*(-a*d+b*c)*x*(b*x^2+a)^(1/2)/c/d/(d*x^2+c)^4+1/48*(7*a*d+2*b*c)*x*(b*x^2+a)^(1/2)/c^2/d/(d*x^2+c)^3+1/192*(-35*a^2*d^2+24*a*b*c*d+8*b^2*c^2)*x*(b*x^2+a)^(1/2)/c^3/d/(-a*d+b*c)/(d*x^2+c)^2+1/384*(105*a^3*d^3-170*a^2*b*c*d^2+40*a*b^2*c^2*d+16*b^3*c^3)*x*(b*x^2+a)^(1/2)/c^4/d/(-a*d+b*c)^2/(d*x^2+c)

Rubi [A] time = 0.37, antiderivative size = 300, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {413, 527, 12, 377, 208}

$$\frac{x\sqrt{a+bx^2} (-170a^2bcd^2 + 105a^3d^3 + 40ab^2c^2d + 16b^3c^3)}{384c^4d(c+dx^2)(bc-ad)^2} + \frac{x\sqrt{a+bx^2} (-35a^2d^2 + 24abcd + 8b^2c^2)}{192c^3d(c+dx^2)^2(bc-ad)} + \frac{a^2(35a^2d^2 - 80abcd + 48b^2c^2)}{128c^{9/2}(bc-ad)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)/(c + d*x^2)^5, x]

[Out] -((b*c - a*d)*x*sqrt[a + b*x^2])/(8*c*d*(c + d*x^2)^4) + ((2*b*c + 7*a*d)*x*sqrt[a + b*x^2])/(48*c^2*d*(c + d*x^2)^3) + ((8*b^2*c^2 + 24*a*b*c*d - 35*a^2*d^2)*x*sqrt[a + b*x^2])/(192*c^3*d*(b*c - a*d)*(c + d*x^2)^2) + ((16*b^3*c^3 + 40*a*b^2*c^2*d - 170*a^2*b*c*d^2 + 105*a^3*d^3)*x*sqrt[a + b*x^2])/(384*c^4*d*(b*c - a*d)^2*(c + d*x^2)) + (a^2*(48*b^2*c^2 - 80*a*b*c*d + 35*a^2*d^2)*ArcTanh[(sqrt[b*c - a*d]*x)/(sqrt[c]*sqrt[a + b*x^2])])/(128*c^(9/2)*(b*c - a*d)^(5/2))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+1))], x], x]

+ q) + 1)) * x^n, x], x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{3/2}}{(c + dx^2)^5} dx &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{8cd(c + dx^2)^4} + \frac{\int \frac{a(bc+7ad)+2b(bc+3ad)x^2}{\sqrt{a+bx^2}(c+dx^2)^4} dx}{8cd} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{8cd(c + dx^2)^4} + \frac{(2bc + 7ad)x\sqrt{a + bx^2}}{48c^2d(c + dx^2)^3} + \frac{\int \frac{a(bc-ad)(4bc+35ad)+4b(bc-ad)(2bc+7ad)x^2}{\sqrt{a+bx^2}(c+dx^2)^3} dx}{48c^2d(bc - ad)} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{8cd(c + dx^2)^4} + \frac{(2bc + 7ad)x\sqrt{a + bx^2}}{48c^2d(c + dx^2)^3} + \frac{(8b^2c^2 + 24abcd - 35a^2d^2)x\sqrt{a + bx^2}}{192c^3d(bc - ad)(c + dx^2)^2} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{8cd(c + dx^2)^4} + \frac{(2bc + 7ad)x\sqrt{a + bx^2}}{48c^2d(c + dx^2)^3} + \frac{(8b^2c^2 + 24abcd - 35a^2d^2)x\sqrt{a + bx^2}}{192c^3d(bc - ad)(c + dx^2)^2} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{8cd(c + dx^2)^4} + \frac{(2bc + 7ad)x\sqrt{a + bx^2}}{48c^2d(c + dx^2)^3} + \frac{(8b^2c^2 + 24abcd - 35a^2d^2)x\sqrt{a + bx^2}}{192c^3d(bc - ad)(c + dx^2)^2} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{8cd(c + dx^2)^4} + \frac{(2bc + 7ad)x\sqrt{a + bx^2}}{48c^2d(c + dx^2)^3} + \frac{(8b^2c^2 + 24abcd - 35a^2d^2)x\sqrt{a + bx^2}}{192c^3d(bc - ad)(c + dx^2)^2} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{8cd(c + dx^2)^4} + \frac{(2bc + 7ad)x\sqrt{a + bx^2}}{48c^2d(c + dx^2)^3} + \frac{(8b^2c^2 + 24abcd - 35a^2d^2)x\sqrt{a + bx^2}}{192c^3d(bc - ad)(c + dx^2)^2} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{8cd(c + dx^2)^4} + \frac{(2bc + 7ad)x\sqrt{a + bx^2}}{48c^2d(c + dx^2)^3} + \frac{(8b^2c^2 + 24abcd - 35a^2d^2)x\sqrt{a + bx^2}}{192c^3d(bc - ad)(c + dx^2)^2} \end{aligned}$$

Mathematica [A] time = 1.38, size = 362, normalized size = 1.21

$$ax \left(\frac{bx^2}{a} + 1 \right) \left(\frac{3a^2(c+dx^2)^4(35a^2d^2-80abcd+48b^2c^2) \tanh^{-1} \left(\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}} \right)}{\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}} + c(a^4d^2(279c^3 + 511c^2dx^2 + 385cd^2x^4 + 105d^3x^6) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)/(c + d*x^2)^5, x]

[Out] (a*x*(1 + (b*x^2)/a)*(c*(16*b^4*c^3*x^4*(6*c^2 + 4*c*d*x^2 + d^2*x^4) + 8*a*b^3*c^2*x^2*(42*c^3 + 34*c^2*d*x^2 + 21*c*d^2*x^4 + 5*d^3*x^6) + a^4*d^2*(

$$279*c^3 + 511*c^2*d*x^2 + 385*c*d^2*x^4 + 105*d^3*x^6) + 2*a^2*b^2*c*(120*c^4 - 160*c^3*d*x^2 - 345*c^2*d^2*x^4 - 294*c*d^3*x^6 - 85*d^4*x^8) + a^3*b*d*(-528*c^4 - 563*c^3*d*x^2 - 117*c^2*d^2*x^4 + 215*c*d^3*x^6 + 105*d^4*x^8)) + (3*a^2*(48*b^2*c^2 - 80*a*b*c*d + 35*a^2*d^2)*(c + d*x^2)^4*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]))/(384*c^5*(b*c - a*d)^2*(a + b*x^2)^(3/2)*(c + d*x^2)^4)$$

fricas [B] time = 3.68, size = 1604, normalized size = 5.35

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^5,x, algorithm="fricas")

[Out] [1/1536*(3*(48*a^2*b^2*c^6 - 80*a^3*b*c^5*d + 35*a^4*c^4*d^2 + (48*a^2*b^2*c^2*d^4 - 80*a^3*b*c*d^5 + 35*a^4*d^6)*x^8 + 4*(48*a^2*b^2*c^3*d^3 - 80*a^3*b*c^2*d^4 + 35*a^4*c*d^5)*x^6 + 6*(48*a^2*b^2*c^4*d^2 - 80*a^3*b*c^3*d^3 + 35*a^4*c^2*d^4)*x^4 + 4*(48*a^2*b^2*c^5*d - 80*a^3*b*c^4*d^2 + 35*a^4*c^3*d^3)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*((16*b^4*c^5*d^2 + 24*a*b^3*c^4*d^3 - 210*a^2*b^2*c^3*d^4 + 275*a^3*b*c^2*d^5 - 105*a^4*c*d^6)*x^7 + (64*b^4*c^6*d + 88*a*b^3*c^5*d^2 - 780*a^2*b^2*c^4*d^3 + 1013*a^3*b*c^3*d^4 - 385*a^4*c^2*d^5)*x^5 + (96*b^4*c^7 + 112*a*b^3*c^6*d - 1050*a^2*b^2*c^5*d^2 + 1353*a^3*b*c^4*d^3 - 511*a^4*c^3*d^4)*x^3 + 3*(80*a*b^3*c^7 - 256*a^2*b^2*c^6*d + 269*a^3*b*c^5*d^2 - 93*a^4*c^4*d^3)*x)*sqrt(b*x^2 + a))/(b^3*c^12 - 3*a*b^2*c^11*d + 3*a^2*b*c^10*d^2 - a^3*c^9*d^3 + (b^3*c^8*d^4 - 3*a*b^2*c^7*d^5 + 3*a^2*b*c^6*d^6 - a^3*c^5*d^7)*x^8 + 4*(b^3*c^9*d^3 - 3*a*b^2*c^8*d^4 + 3*a^2*b*c^7*d^5 - a^3*c^6*d^6)*x^6 + 6*(b^3*c^10*d^2 - 3*a*b^2*c^9*d^3 + 3*a^2*b*c^8*d^4 - a^3*c^7*d^5)*x^4 + 4*(b^3*c^11*d - 3*a*b^2*c^10*d^2 + 3*a^2*b*c^9*d^3 - a^3*c^8*d^4)*x^2), -1/768*(3*(48*a^2*b^2*c^6 - 80*a^3*b*c^5*d + 35*a^4*c^4*d^2 + (48*a^2*b^2*c^2*d^4 - 80*a^3*b*c*d^5 + 35*a^4*d^6)*x^8 + 4*(48*a^2*b^2*c^3*d^3 - 80*a^3*b*c^2*d^4 + 35*a^4*c*d^5)*x^6 + 6*(48*a^2*b^2*c^4*d^2 - 80*a^3*b*c^3*d^3 + 35*a^4*c^2*d^4)*x^4 + 4*(48*a^2*b^2*c^5*d - 80*a^3*b*c^4*d^2 + 35*a^4*c^3*d^3)*x^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d))*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x) - 2*((16*b^4*c^5*d^2 + 24*a*b^3*c^4*d^3 - 210*a^2*b^2*c^3*d^4 + 275*a^3*b*c^2*d^5 - 105*a^4*c*d^6)*x^7 + (64*b^4*c^6*d + 88*a*b^3*c^5*d^2 - 780*a^2*b^2*c^4*d^3 + 1013*a^3*b*c^3*d^4 - 385*a^4*c^2*d^5)*x^5 + (96*b^4*c^7 + 112*a*b^3*c^6*d - 1050*a^2*b^2*c^5*d^2 + 1353*a^3*b*c^4*d^3 - 511*a^4*c^3*d^4)*x^3 + 3*(80*a*b^3*c^7 - 256*a^2*b^2*c^6*d + 269*a^3*b*c^5*d^2 - 93*a^4*c^4*d^3)*x)*sqrt(b*x^2 + a))/(b^3*c^12 - 3*a*b^2*c^11*d + 3*a^2*b*c^10*d^2 - a^3*c^9*d^3 + (b^3*c^8*d^4 - 3*a*b^2*c^7*d^5 + 3*a^2*b*c^6*d^6 - a^3*c^5*d^7)*x^8 + 4*(b^3*c^9*d^3 - 3*a*b^2*c^8*d^4 + 3*a^2*b*c^7*d^5 - a^3*c^6*d^6)*x^6 + 6*(b^3*c^10*d^2 - 3*a*b^2*c^9*d^3 + 3*a^2*b*c^8*d^4 - a^3*c^7*d^5)*x^4 + 4*(b^3*c^11*d - 3*a*b^2*c^10*d^2 + 3*a^2*b*c^9*d^3 - a^3*c^8*d^4)*x^2)]

giac [B] time = 9.60, size = 1557, normalized size = 5.19

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^5,x, algorithm="giac")

[Out] -1/128*(48*a^2*b^(5/2)*c^2 - 80*a^3*b^(3/2)*c*d + 35*a^4*sqrt(b)*d^2)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/((b^2*c^6 - 2*a*b*c^5*d + a^2*c^4*d^2)*sqrt(-b^2*c^2 + a*b*c*d)) - 1/192*(144*(sqrt(b)*x - sqrt(b*x^2 + a))^14*a^2*b^(5/2)*c^2*d^5 - 240*(sqrt(b)*x - sqrt(b*x^2 + a))^14*a^3*b^(3/2)*c*d^6 + 105*(sqrt(b)*x - sqrt(b*x^2

+ a))¹⁴a⁴sqrt(b)*d⁷ + 2016*(sqrt(b)*x - sqrt(b*x² + a))¹²a²b^(7/2)*c³d⁴ - 4368*(sqrt(b)*x - sqrt(b*x² + a))¹²a³b^(5/2)*c²d⁵ + 3150*(sqrt(b)*x - sqrt(b*x² + a))¹²a⁴b^(3/2)*c*d⁶ - 735*(sqrt(b)*x - sqrt(b*x² + a))¹²a⁵sqrt(b)*d⁷ - 2048*(sqrt(b)*x - sqrt(b*x² + a))¹⁰b^(13/2)*c⁶d + 4096*(sqrt(b)*x - sqrt(b*x² + a))¹⁰a*b^(11/2)*c⁵d² + 7936*(sqrt(b)*x - sqrt(b*x² + a))¹⁰a²b^(9/2)*c⁴d³ - 26624*(sqrt(b)*x - sqrt(b*x² + a))¹⁰a³b^(7/2)*c³d⁴ + 26944*(sqrt(b)*x - sqrt(b*x² + a))¹⁰a⁴b^(5/2)*c²d⁵ - 12320*(sqrt(b)*x - sqrt(b*x² + a))¹⁰a⁵b^(3/2)*c*d⁶ + 2205*(sqrt(b)*x - sqrt(b*x² + a))¹⁰a⁶sqrt(b)*d⁷ - 2048*(sqrt(b)*x - sqrt(b*x² + a))⁸b^(15/2)*c⁷ - 1024*(sqrt(b)*x - sqrt(b*x² + a))⁸a*b^(13/2)*c⁶d + 27392*(sqrt(b)*x - sqrt(b*x² + a))⁸a²b^(11/2)*c⁵d² - 65920*(sqrt(b)*x - sqrt(b*x² + a))⁸a³b^(9/2)*c⁴d³ + 81680*(sqrt(b)*x - sqrt(b*x² + a))⁸a⁴b^(7/2)*c³d⁴ - 58840*(sqrt(b)*x - sqrt(b*x² + a))⁸a⁵b^(5/2)*c²d⁵ + 22750*(sqrt(b)*x - sqrt(b*x² + a))⁸a⁶b^(3/2)*c*d⁶ - 3675*(sqrt(b)*x - sqrt(b*x² + a))⁸a⁷sqrt(b)*d⁷ - 2048*(sqrt(b)*x - sqrt(b*x² + a))⁶a²b^(13/2)*c⁶d - 8192*(sqrt(b)*x - sqrt(b*x² + a))⁶a³b^(11/2)*c⁵d² + 47104*(sqrt(b)*x - sqrt(b*x² + a))⁶a⁴b^(9/2)*c⁴d³ - 74240*(sqrt(b)*x - sqrt(b*x² + a))⁶a⁵b^(7/2)*c³d⁴ + 56416*(sqrt(b)*x - sqrt(b*x² + a))⁶a⁶b^(5/2)*c²d⁵ - 22400*(sqrt(b)*x - sqrt(b*x² + a))⁶a⁷b^(3/2)*c*d⁶ + 3675*(sqrt(b)*x - sqrt(b*x² + a))⁶a⁸sqrt(b)*d⁷ - 1536*(sqrt(b)*x - sqrt(b*x² + a))⁴a⁴b^(11/2)*c⁵d² - 2304*(sqrt(b)*x - sqrt(b*x² + a))⁴a⁵b^(9/2)*c⁴d³ + 17696*(sqrt(b)*x - sqrt(b*x² + a))⁴a⁶b^(7/2)*c³d⁴ - 23152*(sqrt(b)*x - sqrt(b*x² + a))⁴a⁷b^(5/2)*c²d⁵ + 11690*(sqrt(b)*x - sqrt(b*x² + a))⁴a⁸b^(3/2)*c*d⁶ - 2205*(sqrt(b)*x - sqrt(b*x² + a))⁴a⁹sqrt(b)*d⁷ - 256*(sqrt(b)*x - sqrt(b*x² + a))²a⁶b^(9/2)*c⁴d³ - 512*(sqrt(b)*x - sqrt(b*x² + a))²a⁷b^(7/2)*c³d⁴ + 2896*(sqrt(b)*x - sqrt(b*x² + a))²a⁸b^(5/2)*c²d⁵ - 2800*(sqrt(b)*x - sqrt(b*x² + a))²a⁹b^(3/2)*c*d⁶ + 735*(sqrt(b)*x - sqrt(b*x² + a))²a¹⁰sqrt(b)*d⁷ - 16*a⁸b^(7/2)*c³d⁴ - 40*a⁹b^(5/2)*c²d⁵ + 170*a¹⁰b^(3/2)*c*d⁶ - 105*a¹¹sqrt(b)*d⁷)/((b²*c⁶d² - 2*a*b*c⁵d³ + a²*c⁴d⁴)*(sqrt(b)*x - sqrt(b*x² + a))⁴d + 4*(sqrt(b)*x - sqrt(b*x² + a))²b*c - 2*(sqrt(b)*x - sqrt(b*x² + a))²a*d + a²d)⁴)

maple [B] time = 0.05, size = 18791, normalized size = 62.64

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x²+a)^(3/2)/(d*x²+c)⁵,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{(dx^2 + c)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x²+a)^(3/2)/(d*x²+c)⁵,x, algorithm="maxima")

[Out] integrate((b*x² + a)^(3/2)/(d*x² + c)⁵, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{(dx^2 + c)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a + b*x^2)^(3/2)/(c + d*x^2)^5,x)
```

```
[Out] int((a + b*x^2)^(3/2)/(c + d*x^2)^5, x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)**(3/2)/(d*x**2+c)**5,x)
```

```
[Out] Timed out
```

3.62 $\int (a + bx^2)^{5/2} (c + dx^2)^3 dx$

Optimal. Leaf size=349

$$\frac{dx (a + bx^2)^{7/2} (15a^2d^2 - 68abcd + 152b^2c^2)}{960b^3} + \frac{x (a + bx^2)^{5/2} (-5a^3d^3 + 36a^2bcd^2 - 120ab^2c^2d + 320b^3c^3)}{1920b^3} + \dots$$

[Out] $1/1536*a*(-5*a^3*d^3+36*a^2*b*c*d^2-120*a*b^2*c^2*d+320*b^3*c^3)*x*(b*x^2+a)^{(3/2)}/b^3+1/1920*(-5*a^3*d^3+36*a^2*b*c*d^2-120*a*b^2*c^2*d+320*b^3*c^3)*x*(b*x^2+a)^{(5/2)}/b^3+1/960*d*(15*a^2*d^2-68*a*b*c*d+152*b^2*c^2)*x*(b*x^2+a)^{(7/2)}/b^3+1/120*d*(-5*a*d+16*b*c)*x*(b*x^2+a)^{(7/2)}*(d*x^2+c)/b^2+1/12*d*x*(b*x^2+a)^{(7/2)}*(d*x^2+c)^2/b+1/1024*a^3*(-5*a^3*d^3+36*a^2*b*c*d^2-120*a*b^2*c^2*d+320*b^3*c^3)*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(7/2)}+1/1024*a^2*(-5*a^3*d^3+36*a^2*b*c*d^2-120*a*b^2*c^2*d+320*b^3*c^3)*x*(b*x^2+a)^{(1/2)}/b^3$

Rubi [A] time = 0.25, antiderivative size = 349, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {416, 528, 388, 195, 217, 206}

$$\frac{dx (a + bx^2)^{7/2} (15a^2d^2 - 68abcd + 152b^2c^2)}{960b^3} + \frac{x (a + bx^2)^{5/2} (36a^2bcd^2 - 5a^3d^3 - 120ab^2c^2d + 320b^3c^3)}{1920b^3} + \dots$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a + b*x^2)^{(5/2)}*(c + d*x^2)^3, x]$

[Out] $(a^2*(320*b^3*c^3 - 120*a*b^2*c^2*d + 36*a^2*b*c*d^2 - 5*a^3*d^3)*x*\operatorname{Sqrt}[a + b*x^2])/(1024*b^3) + (a*(320*b^3*c^3 - 120*a*b^2*c^2*d + 36*a^2*b*c*d^2 - 5*a^3*d^3)*x*(a + b*x^2)^{(3/2)})/(1536*b^3) + ((320*b^3*c^3 - 120*a*b^2*c^2*d + 36*a^2*b*c*d^2 - 5*a^3*d^3)*x*(a + b*x^2)^{(5/2)})/(1920*b^3) + (d*(152*b^2*c^2 - 68*a*b*c*d + 15*a^2*d^2)*x*(a + b*x^2)^{(7/2)})/(960*b^3) + (d*(16*b*c - 5*a*d)*x*(a + b*x^2)^{(7/2)}*(c + d*x^2))/(120*b^2) + (d*x*(a + b*x^2)^{(7/2)}*(c + d*x^2)^2)/(12*b) + (a^3*(320*b^3*c^3 - 120*a*b^2*c^2*d + 36*a^2*b*c*d^2 - 5*a^3*d^3)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(1024*b^{(7/2)})$

Rule 195

$\operatorname{Int}[(a + b*x^n)^p, x] := \operatorname{Simp}[(x*(a + b*x^n)^p)/(n*p + 1), x] + \operatorname{Dist}[(a*n*p)/(n*p + 1), \operatorname{Int}[(a + b*x^n)^{p-1}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

$\operatorname{Int}[(a + b*x^2)^{-1}, x] := \operatorname{Simp}[(1*\operatorname{ArcTanh}[(\operatorname{Rt}[-b, 2]*x)/\operatorname{Rt}[a, 2]])/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]), x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

$\operatorname{Int}[1/\operatorname{Sqrt}[(a + b*x^2)], x] := \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /;$ FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

$\operatorname{Int}[(a + b*x^n)^p*(c + d*x^n), x] := \operatorname{Simp}[(d*x*(a + b*x^n)^{p+1})/(b*(n*(p+1) + 1)), x] - \operatorname{Dist}[(a*d - b*c*(n*($

$(p + 1) + 1) / (b * (n * (p + 1) + 1))$, Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 416

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)), x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q - 1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 528

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] :> Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^{5/2} (c + dx^2)^3 dx &= \frac{dx (a + bx^2)^{7/2} (c + dx^2)^2}{12b} + \frac{\int (a + bx^2)^{5/2} (c + dx^2) (c(12bc - ad) + d(16bc - 5ad)) dx}{12b} \\ &= \frac{d(16bc - 5ad)x (a + bx^2)^{7/2} (c + dx^2)}{120b^2} + \frac{dx (a + bx^2)^{7/2} (c + dx^2)^2}{12b} + \frac{\int (a + bx^2)^{5/2} (c + dx^2) dx}{12b} \\ &= \frac{d(152b^2c^2 - 68abcd + 15a^2d^2)x (a + bx^2)^{7/2}}{960b^3} + \frac{d(16bc - 5ad)x (a + bx^2)^{7/2} (c + dx^2)}{120b^2} \\ &= \frac{(320b^3c^3 - 120ab^2c^2d + 36a^2bcd^2 - 5a^3d^3)x (a + bx^2)^{5/2}}{1920b^3} + \frac{d(152b^2c^2 - 68abcd + 15a^2d^2)x (a + bx^2)^{3/2}}{960b^3} \\ &= \frac{a(320b^3c^3 - 120ab^2c^2d + 36a^2bcd^2 - 5a^3d^3)x (a + bx^2)^{3/2}}{1536b^3} + \frac{(320b^3c^3 - 120ab^2c^2d + 36a^2bcd^2 - 5a^3d^3)x \sqrt{a + bx^2}}{1024b^3} \\ &= \frac{a^2(320b^3c^3 - 120ab^2c^2d + 36a^2bcd^2 - 5a^3d^3)x \sqrt{a + bx^2}}{1024b^3} + \frac{a(320b^3c^3 - 120ab^2c^2d + 36a^2bcd^2 - 5a^3d^3)}{1024b^3} \\ &= \frac{a^2(320b^3c^3 - 120ab^2c^2d + 36a^2bcd^2 - 5a^3d^3)x \sqrt{a + bx^2}}{1024b^3} + \frac{a(320b^3c^3 - 120ab^2c^2d + 36a^2bcd^2 - 5a^3d^3)}{1024b^3} \end{aligned}$$

Mathematica [A] time = 5.18, size = 270, normalized size = 0.77

$$\sqrt{b} x \sqrt{a + bx^2} (75a^5d^3 - 10a^4bd^2 (54c + 5dx^2) + 40a^3b^2d (45c^2 + 9cdx^2 + d^2x^4) + 48a^2b^3 (220c^3 + 295c^2dx^2 + 100cd^2x^4 + 10d^3x^6)) + \frac{d(152b^2c^2 - 68abcd + 15a^2d^2)x \sqrt{a + bx^2}}{960b^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2)*(c + d*x^2)^3,x]


```
[Out] (Sqrt[b]*x*Sqrt[a + b*x^2]*(75*a^5*d^3 - 10*a^4*b*d^2*(54*c + 5*d*x^2) + 40
*a^3*b^2*d*(45*c^2 + 9*c*d*x^2 + d^2*x^4) + 128*b^5*x^4*(20*c^3 + 45*c^2*d*
x^2 + 36*c*d^2*x^4 + 10*d^3*x^6) + 48*a^2*b^3*(220*c^3 + 295*c^2*d*x^2 + 18
6*c*d^2*x^4 + 45*d^3*x^6) + 64*a*b^4*x^2*(130*c^3 + 255*c^2*d*x^2 + 189*c*d
^2*x^4 + 50*d^3*x^6)) - 15*a^3*(-320*b^3*c^3 + 120*a*b^2*c^2*d - 36*a^2*b*c
*d^2 + 5*a^3*d^3)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]]/(15360*b^(7/2))
```

fricas [A] time = 1.58, size = 608, normalized size = 1.74

$$\frac{15(320a^3b^3c^3 - 120a^4b^2c^2d + 36a^5bcd^2 - 5a^6d^3)\sqrt{b} \log(-2bx^2 + 2\sqrt{bx^2 + a}\sqrt{b}x - a) - 2(1280b^6d^3x^{11} + 128(36b^6cd^2 + 25ab^5d^3)x^9 + 144(40b^6c^2d + 84a^2b^5cd^2 + 15a^2b^4d^3)x^7 + 8(320b^6c^3 + 2040ab^5c^2d + 1116a^2b^4cd^2 + 5a^3b^3d^3)x^5 + 10(832ab^5c^3 + 1416a^2b^4c^2d + 36a^3b^3cd^2 - 5a^4b^2d^3)x^3 + 15(704a^2b^4c^3 + 120a^3b^3c^2d - 36a^4b^2cd^2 + 5a^5bd^3)x)\sqrt{bx^2 + a}}{b^4} - \frac{1}{15360}(15(320a^3b^3c^3 - 120a^4b^2c^2d + 36a^5bcd^2 - 5a^6d^3)*\sqrt{b} \arctan(\sqrt{-b}x/\sqrt{bx^2 + a}) - (1280b^6d^3x^{11} + 128(36b^6cd^2 + 25ab^5d^3)x^9 + 144(40b^6c^2d + 84a^2b^5cd^2 + 15a^2b^4d^3)x^7 + 8(320b^6c^3 + 2040ab^5c^2d + 1116a^2b^4cd^2 + 5a^3b^3d^3)x^5 + 10(832ab^5c^3 + 1416a^2b^4c^2d + 36a^3b^3cd^2 - 5a^4b^2d^3)x^3 + 15(704a^2b^4c^3 + 120a^3b^3c^2d - 36a^4b^2cd^2 + 5a^5bd^3)x)\sqrt{bx^2 + a})/b^4$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(5/2)*(d*x^2+c)^3,x, algorithm="fricas")
```

```
[Out] [-1/30720*(15*(320*a^3*b^3*c^3 - 120*a^4*b^2*c^2*d + 36*a^5*b*c*d^2 - 5*a^6
*d^3)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(1280*b^6
*d^3*x^11 + 128*(36*b^6*c*d^2 + 25*a*b^5*d^3)*x^9 + 144*(40*b^6*c^2*d + 84*
a*b^5*c*d^2 + 15*a^2*b^4*d^3)*x^7 + 8*(320*b^6*c^3 + 2040*a*b^5*c^2*d + 111
6*a^2*b^4*c*d^2 + 5*a^3*b^3*d^3)*x^5 + 10*(832*a*b^5*c^3 + 1416*a^2*b^4*c^2
*d + 36*a^3*b^3*c*d^2 - 5*a^4*b^2*d^3)*x^3 + 15*(704*a^2*b^4*c^3 + 120*a^3*
b^3*c^2*d - 36*a^4*b^2*c*d^2 + 5*a^5*b*d^3)*x)*sqrt(b*x^2 + a))/b^4, -1/153
60*(15*(320*a^3*b^3*c^3 - 120*a^4*b^2*c^2*d + 36*a^5*b*c*d^2 - 5*a^6*d^3)*s
qrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (1280*b^6*d^3*x^11 + 128*(36*b
^6*c*d^2 + 25*a*b^5*d^3)*x^9 + 144*(40*b^6*c^2*d + 84*a*b^5*c*d^2 + 15*a^2*
b^4*d^3)*x^7 + 8*(320*b^6*c^3 + 2040*a*b^5*c^2*d + 1116*a^2*b^4*c*d^2 + 5*a
^3*b^3*d^3)*x^5 + 10*(832*a*b^5*c^3 + 1416*a^2*b^4*c^2*d + 36*a^3*b^3*c*d^2
- 5*a^4*b^2*d^3)*x^3 + 15*(704*a^2*b^4*c^3 + 120*a^3*b^3*c^2*d - 36*a^4*b^
2*c*d^2 + 5*a^5*b*d^3)*x)*sqrt(b*x^2 + a))/b^4]
```

giac [A] time = 0.68, size = 321, normalized size = 0.92

$$\frac{1}{15360} \left(2 \left(4 \left(2 \left(8 \left(10b^2d^3x^2 + \frac{36b^{12}cd^2 + 25ab^{11}d^3}{b^{10}} \right) x^2 + \frac{9(40b^{12}c^2d + 84ab^{11}cd^2 + 15a^2b^{10}d^3)}{b^{10}} \right) x^2 + \frac{320}{b^{10}} \right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(5/2)*(d*x^2+c)^3,x, algorithm="giac")
```

```
[Out] 1/15360*(2*(4*(2*(8*(10*b^2*d^3*x^2 + (36*b^12*c*d^2 + 25*a*b^11*d^3)/b^10)
*x^2 + 9*(40*b^12*c^2*d + 84*a*b^11*c*d^2 + 15*a^2*b^10*d^3)/b^10)*x^2 + (3
20*b^12*c^3 + 2040*a*b^11*c^2*d + 1116*a^2*b^10*c*d^2 + 5*a^3*b^9*d^3)/b^10
)*x^2 + 5*(832*a*b^11*c^3 + 1416*a^2*b^10*c^2*d + 36*a^3*b^9*c*d^2 - 5*a^4*
b^8*d^3)/b^10)*x^2 + 15*(704*a^2*b^10*c^3 + 120*a^3*b^9*c^2*d - 36*a^4*b^8*
c*d^2 + 5*a^5*b^7*d^3)/b^10)*sqrt(b*x^2 + a)*x - 1/1024*(320*a^3*b^3*c^3 -
120*a^4*b^2*c^2*d + 36*a^5*b*c*d^2 - 5*a^6*d^3)*log(abs(-sqrt(b)*x + sqrt(b
*x^2 + a)))/b^(7/2)
```

maple [A] time = 0.02, size = 476, normalized size = 1.36

$$\frac{(bx^2 + a)^{\frac{7}{2}} d^3 x^5}{12b} - \frac{5a^6 d^3 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{1024b^{\frac{7}{2}}} + \frac{9a^5 c d^2 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{256b^{\frac{5}{2}}} - \frac{15a^4 c^2 d \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{128b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*x^2+a)^(5/2)*(d*x^2+c)^3,x)
```

[Out] $1/12*d^3*x^5*(b*x^2+a)^{(7/2)}/b-1/24*d^3*a/b^2*x^3*(b*x^2+a)^{(7/2)}+1/64*d^3*a^2/b^3*x*(b*x^2+a)^{(7/2)}-1/384*d^3*a^3/b^3*x*(b*x^2+a)^{(5/2)}-5/1536*d^3*a^4/b^3*x*(b*x^2+a)^{(3/2)}-5/1024*d^3*a^5/b^3*x*(b*x^2+a)^{(1/2)}-5/1024*d^3*a^6/b^{(7/2)}*\ln(b^{(1/2)}*x+(b*x^2+a)^{(1/2)})+3/10*c*d^2*x^3*(b*x^2+a)^{(7/2)}/b-9/80*c*d^2*a/b^2*x*(b*x^2+a)^{(7/2)}+3/160*c*d^2*a^2/b^2*x*(b*x^2+a)^{(5/2)}+3/128*c*d^2*a^3/b^2*x*(b*x^2+a)^{(3/2)}+9/256*c*d^2*a^4/b^2*x*(b*x^2+a)^{(1/2)}+9/256*c*d^2*a^5/b^{(5/2)}*\ln(b^{(1/2)}*x+(b*x^2+a)^{(1/2)})+3/8*c^2*d*x*(b*x^2+a)^{(7/2)}/b-1/16*c^2*d*a/b*x*(b*x^2+a)^{(5/2)}-5/64*c^2*d*a^2/b*x*(b*x^2+a)^{(3/2)}-15/128*c^2*d*a^3/b*x*(b*x^2+a)^{(1/2)}-15/128*c^2*d*a^4/b^{(3/2)}*\ln(b^{(1/2)}*x+(b*x^2+a)^{(1/2)})+1/6*c^3*x*(b*x^2+a)^{(5/2)}+5/24*c^3*a*x*(b*x^2+a)^{(3/2)}+5/16*c^3*a^2*x*(b*x^2+a)^{(1/2)}+5/16*c^3*a^3/b^{(1/2)}*\ln(b^{(1/2)}*x+(b*x^2+a)^{(1/2)})$

maxima [A] time = 1.47, size = 447, normalized size = 1.28

$$\frac{(bx^2 + a)^{\frac{7}{2}}d^3x^5}{12b} + \frac{3(bx^2 + a)^{\frac{7}{2}}cd^2x^3}{10b} - \frac{(bx^2 + a)^{\frac{7}{2}}ad^3x^3}{24b^2} + \frac{1}{6}(bx^2 + a)^{\frac{5}{2}}c^3x + \frac{5}{24}(bx^2 + a)^{\frac{3}{2}}ac^3x + \frac{5}{16}\sqrt{bx^2 + a}a^2c^3x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)*(d*x^2+c)^3,x, algorithm="maxima")

[Out] $1/12*(b*x^2 + a)^{(7/2)}*d^3*x^5/b + 3/10*(b*x^2 + a)^{(7/2)}*c*d^2*x^3/b - 1/24*(b*x^2 + a)^{(7/2)}*a*d^3*x^3/b^2 + 1/6*(b*x^2 + a)^{(5/2)}*c^3*x + 5/24*(b*x^2 + a)^{(3/2)}*a*c^3*x + 5/16*\text{sqrt}(b*x^2 + a)*a^2*c^3*x + 3/8*(b*x^2 + a)^{(7/2)}*c^2*d*x/b - 1/16*(b*x^2 + a)^{(5/2)}*a*c^2*d*x/b - 5/64*(b*x^2 + a)^{(3/2)}*a^2*c^2*d*x/b - 15/128*\text{sqrt}(b*x^2 + a)*a^3*c^2*d*x/b - 9/80*(b*x^2 + a)^{(7/2)}*a*c*d^2*x/b^2 + 3/160*(b*x^2 + a)^{(5/2)}*a^2*c*d^2*x/b^2 + 3/128*(b*x^2 + a)^{(3/2)}*a^3*c*d^2*x/b^2 + 9/256*\text{sqrt}(b*x^2 + a)*a^4*c*d^2*x/b^2 + 1/64*(b*x^2 + a)^{(7/2)}*a^2*d^3*x/b^3 - 1/384*(b*x^2 + a)^{(5/2)}*a^3*d^3*x/b^3 - 5/1536*(b*x^2 + a)^{(3/2)}*a^4*d^3*x/b^3 - 5/1024*\text{sqrt}(b*x^2 + a)*a^5*d^3*x/b^3 + 5/16*a^3*c^3*\text{arcsinh}(b*x/\text{sqrt}(a*b))/\text{sqrt}(b) - 15/128*a^4*c^2*d*\text{arcsinh}(b*x/\text{sqrt}(a*b))/b^{(3/2)} + 9/256*a^5*c*d^2*\text{arcsinh}(b*x/\text{sqrt}(a*b))/b^{(5/2)} - 5/1024*a^6*d^3*\text{arcsinh}(b*x/\text{sqrt}(a*b))/b^{(7/2)}$

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (bx^2 + a)^{5/2} (dx^2 + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)*(c + d*x^2)^3,x)

[Out] int((a + b*x^2)^(5/2)*(c + d*x^2)^3, x)

sympy [B] time = 102.67, size = 796, normalized size = 2.28

$$\frac{5a^{\frac{11}{2}}d^3x}{1024b^3\sqrt{1 + \frac{bx^2}{a}}} - \frac{9a^{\frac{9}{2}}cd^2x}{256b^2\sqrt{1 + \frac{bx^2}{a}}} + \frac{5a^{\frac{9}{2}}d^3x^3}{3072b^2\sqrt{1 + \frac{bx^2}{a}}} + \frac{15a^{\frac{7}{2}}c^2dx}{128b\sqrt{1 + \frac{bx^2}{a}}} - \frac{3a^{\frac{7}{2}}cd^2x^3}{256b\sqrt{1 + \frac{bx^2}{a}}} - \frac{a^{\frac{7}{2}}d^3x^5}{1536b\sqrt{1 + \frac{bx^2}{a}}} + \frac{a^{\frac{5}{2}}c^3x}{16\sqrt{1 + \frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)*(d*x**2+c)**3,x)

[Out] $5*a**(11/2)*d**3*x/(1024*b**3*\text{sqrt}(1 + b*x**2/a)) - 9*a**(9/2)*c*d**2*x/(256*b**2*\text{sqrt}(1 + b*x**2/a)) + 5*a**(9/2)*d**3*x**3/(3072*b**2*\text{sqrt}(1 + b*x**2/a)) + 15*a**(7/2)*c**2*d*x/(128*b*\text{sqrt}(1 + b*x**2/a)) - 3*a**(7/2)*c*d**2*x**3/(256*b*\text{sqrt}(1 + b*x**2/a)) - a**(7/2)*d**3*x**5/(1536*b*\text{sqrt}(1 + b*x**2/a)) + a**(5/2)*c**3*x*\text{sqrt}(1 + b*x**2/a)/2 + 3*a**(5/2)*c**3*x/(16*\text{sqrt}(1 + b*x**2/a))$

$$\begin{aligned}
& 1 + b*x**2/a)) + 133*a**(5/2)*c**2*d*x**3/(128*sqrt(1 + b*x**2/a)) + 387*a* \\
& *(5/2)*c*d**2*x**5/(640*sqrt(1 + b*x**2/a)) + 55*a**(5/2)*d**3*x**7/(384*sq \\
& rt(1 + b*x**2/a)) + 35*a**(3/2)*b*c**3*x**3/(48*sqrt(1 + b*x**2/a)) + 127*a \\
& **3/2)*b*c**2*d*x**5/(64*sqrt(1 + b*x**2/a)) + 219*a**(3/2)*b*c*d**2*x**7/ \\
& (160*sqrt(1 + b*x**2/a)) + 67*a**(3/2)*b*d**3*x**9/(192*sqrt(1 + b*x**2/a)) \\
& + 17*sqrt(a)*b**2*c**3*x**5/(24*sqrt(1 + b*x**2/a)) + 23*sqrt(a)*b**2*c**2 \\
& *d*x**7/(16*sqrt(1 + b*x**2/a)) + 87*sqrt(a)*b**2*c*d**2*x**9/(80*sqrt(1 + \\
& b*x**2/a)) + 7*sqrt(a)*b**2*d**3*x**11/(24*sqrt(1 + b*x**2/a)) - 5*a**6*d** \\
& 3*asinh(sqrt(b)*x/sqrt(a))/(1024*b**(7/2)) + 9*a**5*c*d**2*asinh(sqrt(b)*x/ \\
& sqrt(a))/(256*b**(5/2)) - 15*a**4*c**2*d*asinh(sqrt(b)*x/sqrt(a))/(128*b** \\
& (3/2)) + 5*a**3*c**3*asinh(sqrt(b)*x/sqrt(a))/(16*sqrt(b)) + b**3*c**3*x**7/ \\
& (6*sqrt(a)*sqrt(1 + b*x**2/a)) + 3*b**3*c**2*d*x**9/(8*sqrt(a)*sqrt(1 + b*x \\
& **2/a)) + 3*b**3*c*d**2*x**11/(10*sqrt(a)*sqrt(1 + b*x**2/a)) + b**3*d**3*x \\
& **13/(12*sqrt(a)*sqrt(1 + b*x**2/a))
\end{aligned}$$

3.63 $\int (a + bx^2)^{5/2} (c + dx^2)^2 dx$

Optimal. Leaf size=241

$$\frac{x(a + bx^2)^{5/2} (3a^2d^2 - 20abcd + 80b^2c^2)}{480b^2} + \frac{ax(a + bx^2)^{3/2} (3a^2d^2 - 20abcd + 80b^2c^2)}{384b^2} + \frac{a^2x\sqrt{a + bx^2} (3a^2d^2 - 20abcd + 80b^2c^2)}{256b^2}$$

[Out] $\frac{1}{384} a (3 a^2 d^2 - 20 a b c d + 80 b^2 c^2) x x (b x^2 + a)^{(3/2)} / b^2 + \frac{1}{480} (3 a^2 d^2 - 20 a b c d + 80 b^2 c^2) x x (b x^2 + a)^{(5/2)} / b^2 + \frac{3}{80} d (-a d + 4 b c) x x (b x^2 + a)^{(7/2)} / b^2 + \frac{1}{10} d x x (b x^2 + a)^{(7/2)} (d x^2 + c) / b + \frac{1}{256} a^3 (3 a^2 d^2 - 20 a b c d + 80 b^2 c^2) \operatorname{arctanh}(x b^{(1/2)} / (b x^2 + a)^{(1/2)}) / b^{(5/2)} + \frac{1}{256} a^2 (3 a^2 d^2 - 20 a b c d + 80 b^2 c^2) x x (b x^2 + a)^{(1/2)} / b^2$

Rubi [A] time = 0.15, antiderivative size = 241, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {416, 388, 195, 217, 206}

$$\frac{x(a + bx^2)^{5/2} (3a^2d^2 - 20abcd + 80b^2c^2)}{480b^2} + \frac{ax(a + bx^2)^{3/2} (3a^2d^2 - 20abcd + 80b^2c^2)}{384b^2} + \frac{a^2x\sqrt{a + bx^2} (3a^2d^2 - 20abcd + 80b^2c^2)}{256b^2}$$

Antiderivative was successfully verified.

[In] `Int[(a + b*x^2)^(5/2)*(c + d*x^2)^2,x]`

[Out] $(a^2(80b^2c^2 - 20abc d + 3a^2d^2) x \operatorname{Sqrt}[a + b x^2]) / (256 b^2) + (a(80b^2c^2 - 20abc d + 3a^2d^2) x (a + b x^2)^{(3/2)}) / (384 b^2) + ((80b^2c^2 - 20abc d + 3a^2d^2) x (a + b x^2)^{(5/2)}) / (480 b^2) + (3 d (4 b c - a d) x (a + b x^2)^{(7/2)}) / (80 b^2) + (d x (a + b x^2)^{(7/2)} (c + d x^2)) / (10 b) + (a^3 (80b^2c^2 - 20abc d + 3a^2d^2) \operatorname{ArcTanh}[\operatorname{Sqrt}[b x] / \operatorname{Sqrt}[a + b x^2]]) / (256 b^{(5/2)})$

Rule 195

`Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])`

Rule 206

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[Rt[-b, 2]*x]/Rt[a, 2])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 217

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

Rule 388

`Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]`

Rule 416

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)),
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a
, b, c, d, n, p, q, x]
```

Rubi steps

$$\begin{aligned}
\int (a + bx^2)^{5/2} (c + dx^2)^2 dx &= \frac{dx (a + bx^2)^{7/2} (c + dx^2)}{10b} + \frac{\int (a + bx^2)^{5/2} (c(10bc - ad) + 3d(4bc - ad)x^2) dx}{10b} \\
&= \frac{3d(4bc - ad)x (a + bx^2)^{7/2}}{80b^2} + \frac{dx (a + bx^2)^{7/2} (c + dx^2)}{10b} - \frac{(3ad(4bc - ad) - 8bc}{80b^2} \\
&= \frac{(80b^2c^2 - 20abcd + 3a^2d^2)x (a + bx^2)^{5/2}}{480b^2} + \frac{3d(4bc - ad)x (a + bx^2)^{7/2}}{80b^2} + \frac{dx (a + bx^2)^{5/2}}{10b} \\
&= \frac{a(80b^2c^2 - 20abcd + 3a^2d^2)x (a + bx^2)^{3/2}}{384b^2} + \frac{(80b^2c^2 - 20abcd + 3a^2d^2)x (a + bx^2)^{5/2}}{480b^2} + \frac{dx (a + bx^2)^{5/2}}{10b} \\
&= \frac{a^2(80b^2c^2 - 20abcd + 3a^2d^2)x\sqrt{a + bx^2}}{256b^2} + \frac{a(80b^2c^2 - 20abcd + 3a^2d^2)x (a + bx^2)^{3/2}}{384b^2} + \frac{dx (a + bx^2)^{5/2}}{10b} \\
&= \frac{a^2(80b^2c^2 - 20abcd + 3a^2d^2)x\sqrt{a + bx^2}}{256b^2} + \frac{a(80b^2c^2 - 20abcd + 3a^2d^2)x (a + bx^2)^{3/2}}{384b^2} + \frac{dx (a + bx^2)^{5/2}}{10b} \\
&= \frac{a^2(80b^2c^2 - 20abcd + 3a^2d^2)x\sqrt{a + bx^2}}{256b^2} + \frac{a(80b^2c^2 - 20abcd + 3a^2d^2)x (a + bx^2)^{3/2}}{384b^2} + \frac{dx (a + bx^2)^{5/2}}{10b}
\end{aligned}$$

Mathematica [C] time = 2.80, size = 158, normalized size = 0.66

$$\frac{ax\sqrt{a + bx^2} \left(10bx^2 (c + dx^2)^2 {}_3F_2\left(-\frac{3}{2}, \frac{3}{2}, 2; 1, \frac{9}{2}; -\frac{bx^2}{a}\right) + 20bx^2 (2c^2 + 3cdx^2 + d^2x^4) {}_2F_1\left(-\frac{3}{2}, \frac{3}{2}; \frac{9}{2}; -\frac{bx^2}{a}\right) + 7c^2\right)}{105\sqrt{\frac{bx^2}{a} + 1}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(5/2)*(c + d*x^2)^2,x]

[Out] (a*x*Sqrt[a + b*x^2]*(7*a*(15*c^2 + 10*c*d*x^2 + 3*d^2*x^4)*Hypergeometric2F1[-5/2, 1/2, 7/2, -((b*x^2)/a)] + 20*b*x^2*(2*c^2 + 3*c*d*x^2 + d^2*x^4)*Hypergeometric2F1[-3/2, 3/2, 9/2, -((b*x^2)/a)] + 10*b*x^2*(c + d*x^2)^2*HypergeometricPFQ[{-3/2, 3/2, 2}, {1, 9/2}, -((b*x^2)/a)]))/(105*Sqrt[1 + (b*x^2)/a])

fricas [A] time = 0.87, size = 420, normalized size = 1.74

$$\left[\frac{15(80a^3b^2c^2 - 20a^4bcd + 3a^5d^2)\sqrt{b} \log\left(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{bx - a}\right) + 2(384b^5d^2x^9 + 48(20b^5cd + 21b^4d^2x^7 + 12b^3d^3x^5 + 3b^2d^4x^3 + 3bd^5x) + 3d^6x)}{105\sqrt{bx^2 + a}} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)*(d*x^2+c)^2,x, algorithm="fricas")

[Out] $\left[\frac{1}{7680} (15(80a^3b^2c^2 - 20a^4b^*cd + 3a^5d^2) \sqrt{b} \log(-2bx^2 - 2\sqrt{bx^2 + a}) \sqrt{b} x - a) + 2(384b^5d^2x^9 + 48(20b^5cd + 21ab^4d^2)x^7 + 8(80b^5c^2 + 340ab^4cd + 93a^2b^3d^2)x^5 + 10(208ab^4c^2 + 236a^2b^3cd + 3a^3b^2d^2)x^3 + 15(176a^2b^3c^2 + 20a^3b^2cd - 3a^4b^*d^2)x) \sqrt{bx^2 + a}) / b^3, -1/3840(15(80a^3b^2c^2 - 20a^4b^*cd + 3a^5d^2) \sqrt{-b} \arctan(\sqrt{-b} x / \sqrt{bx^2 + a})) - (384b^5d^2x^9 + 48(20b^5cd + 21ab^4d^2)x^7 + 8(80b^5c^2 + 340ab^4cd + 93a^2b^3d^2)x^5 + 10(208ab^4c^2 + 236a^2b^3cd + 3a^3b^2d^2)x^3 + 15(176a^2b^3c^2 + 20a^3b^2cd - 3a^4b^*d^2)x) \sqrt{bx^2 + a}) / b^3 \right]$

giac [A] time = 0.66, size = 221, normalized size = 0.92

$$\frac{1}{3840} \left(2 \left(4 \left(6 \left(8b^2d^2x^2 + \frac{20b^{10}cd + 21ab^9d^2}{b^8} \right) x^2 + \frac{80b^{10}c^2 + 340ab^9cd + 93a^2b^8d^2}{b^8} \right) x^2 + \frac{5(208ab^9c^2 + 236a^2b^8cd + 3a^3b^7d^2)}{b^8} \right) x^2 + \frac{5(208ab^9c^2 + 236a^2b^8cd + 3a^3b^7d^2)}{b^8} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+a)^(5/2)*(d*x^2+c)^2,x, algorithm="giac")`

[Out] $\frac{1}{3840} (2(4(6(8b^2d^2x^2 + (20b^{10}cd + 21ab^9d^2)/b^8)x^2 + (80b^{10}c^2 + 340ab^9cd + 93a^2b^8d^2)/b^8)x^2 + 5(208ab^9c^2 + 236a^2b^8cd + 3a^3b^7d^2)/b^8)x^2 + 15(176a^2b^8c^2 + 20a^3b^7cd - 3a^4b^6d^2)/b^8) \sqrt{bx^2 + a} x - 1/256(80a^3b^2c^2 - 20a^4b^*cd + 3a^5d^2) \log(\text{abs}(-\sqrt{b}x + \sqrt{bx^2 + a}))) / b^{5/2}$

maple [A] time = 0.01, size = 308, normalized size = 1.28

$$\frac{3a^5d^2 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{256b^{\frac{5}{2}}} - \frac{5a^4cd \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{64b^{\frac{3}{2}}} + \frac{5a^3c^2 \ln(\sqrt{b}x + \sqrt{bx^2 + a})}{16\sqrt{b}} + \frac{3\sqrt{bx^2 + a} a^4d^2x}{256b^2} - \dots$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2+a)^(5/2)*(d*x^2+c)^2,x)`

[Out] $\frac{1}{10}d^2x^3(bx^2+a)^{7/2}/b - 3/80d^2a/b^2x(bx^2+a)^{7/2} + 1/160d^2a^2/b^2x(bx^2+a)^{5/2} + 1/128d^2a^3/b^2x(bx^2+a)^{3/2} + 3/256d^2a^4/b^2x(bx^2+a)^{1/2} + 3/256d^2a^5/b^{5/2} \ln(b^{1/2}x + (bx^2+a)^{1/2}) + 1/4c*d*x(bx^2+a)^{7/2}/b - 1/24c*d*a/b*x(bx^2+a)^{5/2} - 5/96c*d*a^2/b*x(bx^2+a)^{3/2} - 5/64c*d*a^3/b*x(bx^2+a)^{1/2} - 5/64c*d*a^4/b^{3/2} \ln(b^{1/2}x + (bx^2+a)^{1/2}) + 1/6c^2*x(bx^2+a)^{5/2} + 5/24c^2*a*x(bx^2+a)^{3/2} + 5/16c^2*a^2x(bx^2+a)^{1/2} + 5/16c^2*a^3/b^{1/2} \ln(b^{1/2}x + (bx^2+a)^{1/2})$

maxima [A] time = 1.39, size = 286, normalized size = 1.19

$$\frac{(bx^2 + a)^{\frac{7}{2}} d^2 x^3}{10b} + \frac{1}{6} (bx^2 + a)^{\frac{5}{2}} c^2 x + \frac{5}{24} (bx^2 + a)^{\frac{3}{2}} a c^2 x + \frac{5}{16} \sqrt{bx^2 + a} a^2 c^2 x + \frac{(bx^2 + a)^{\frac{7}{2}} c d x}{4b} - \frac{(bx^2 + a)^{\frac{5}{2}} a c d x}{24b} - \frac{5}{24} (bx^2 + a)^{\frac{3}{2}} a^2 c d x + \frac{5}{16} a^3 c^2 \operatorname{arcsinh}(bx^2 + a)^{1/2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+a)^(5/2)*(d*x^2+c)^2,x, algorithm="maxima")`

[Out] $\frac{1}{10}(bx^2 + a)^{7/2}d^2x^3/b + 1/6(bx^2 + a)^{5/2}c^2x + 5/24(bx^2 + a)^{3/2}a^2c^2x + 5/16\sqrt{bx^2 + a}a^2c^2x + 1/4(bx^2 + a)^{7/2}c*d*x/b - 1/24(bx^2 + a)^{5/2}a^2c*d*x/b - 5/96(bx^2 + a)^{3/2}a^2c*d*x/b - 5/64\sqrt{bx^2 + a}a^3c*d*x/b - 3/80(bx^2 + a)^{7/2}a*d^2*x/b^2 + 1/160(bx^2 + a)^{5/2}a^2d^2*x/b^2 + 1/128(bx^2 + a)^{3/2}a^3d^2*x/b^2 + 3/256\sqrt{bx^2 + a}a^4d^2*x/b^2 + 5/16a^3c^2\operatorname{arcsinh}(bx^2 + a)^{1/2}$

$\sqrt{a*b})/\sqrt{b} - 5/64*a^4*c*d*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{(3/2)} + 3/256*a^5*d^2*\operatorname{arcsinh}(b*x/\sqrt{a*b})/b^{(5/2)}$

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (bx^2 + a)^{5/2} (dx^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^(5/2)*(c + d*x^2)^2,x)`

[Out] `int((a + b*x^2)^(5/2)*(c + d*x^2)^2, x)`

sympy [B] time = 58.63, size = 537, normalized size = 2.23

$$-\frac{3a^{\frac{9}{2}}d^2x}{256b^2\sqrt{1+\frac{bx^2}{a}}} + \frac{5a^{\frac{7}{2}}cdx}{64b\sqrt{1+\frac{bx^2}{a}}} - \frac{a^{\frac{7}{2}}d^2x^3}{256b\sqrt{1+\frac{bx^2}{a}}} + \frac{a^{\frac{5}{2}}c^2x\sqrt{1+\frac{bx^2}{a}}}{2} + \frac{3a^{\frac{5}{2}}c^2x}{16\sqrt{1+\frac{bx^2}{a}}} + \frac{133a^{\frac{5}{2}}cdx^3}{192\sqrt{1+\frac{bx^2}{a}}} + \frac{129a^{\frac{5}{2}}d^2x^5}{640\sqrt{1+\frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**(5/2)*(d*x**2+c)**2,x)`

[Out] `-3*a**(9/2)*d**2*x/(256*b**2*sqrt(1 + b*x**2/a)) + 5*a**(7/2)*c*d*x/(64*b*sqrt(1 + b*x**2/a)) - a**(7/2)*d**2*x**3/(256*b*sqrt(1 + b*x**2/a)) + a**(5/2)*c**2*x*sqrt(1 + b*x**2/a)/2 + 3*a**(5/2)*c**2*x/(16*sqrt(1 + b*x**2/a)) + 133*a**(5/2)*c*d*x**3/(192*sqrt(1 + b*x**2/a)) + 129*a**(5/2)*d**2*x**5/(640*sqrt(1 + b*x**2/a)) + 35*a**(3/2)*b*c**2*x**3/(48*sqrt(1 + b*x**2/a)) + 127*a**(3/2)*b*c*d*x**5/(96*sqrt(1 + b*x**2/a)) + 73*a**(3/2)*b*d**2*x**7/(160*sqrt(1 + b*x**2/a)) + 17*sqrt(a)*b**2*c**2*x**5/(24*sqrt(1 + b*x**2/a)) + 23*sqrt(a)*b**2*c*d*x**7/(24*sqrt(1 + b*x**2/a)) + 29*sqrt(a)*b**2*d**2*x**9/(80*sqrt(1 + b*x**2/a)) + 3*a**5*d**2*asinh(sqrt(b)*x/sqrt(a))/(256*b**(5/2)) - 5*a**4*c*d*asinh(sqrt(b)*x/sqrt(a))/(64*b**(3/2)) + 5*a**3*c**2*asinh(sqrt(b)*x/sqrt(a))/(16*sqrt(b)) + b**3*c**2*x**7/(6*sqrt(a)*sqrt(1 + b*x**2/a)) + b**3*c*d*x**9/(4*sqrt(a)*sqrt(1 + b*x**2/a)) + b**3*d**2*x**11/(10*sqrt(a)*sqrt(1 + b*x**2/a))`

3.64 $\int (a + bx^2)^{5/2} (c + dx^2) dx$

Optimal. Leaf size=149

$$\frac{5a^3(8bc - ad) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{128b^{3/2}} + \frac{5a^2x\sqrt{a+bx^2}(8bc - ad)}{128b} + \frac{x(a+bx^2)^{5/2}(8bc - ad)}{48b} + \frac{5ax(a+bx^2)^{3/2}(8bc - ad)}{192b}$$

[Out] $5/192*a*(-a*d+8*b*c)*x*(b*x^2+a)^{(3/2)}/b+1/48*(-a*d+8*b*c)*x*(b*x^2+a)^{(5/2)}/b+1/8*d*x*(b*x^2+a)^{(7/2)}/b+5/128*a^3*(-a*d+8*b*c)*\arctanh(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(3/2)}+5/128*a^2*(-a*d+8*b*c)*x*(b*x^2+a)^{(1/2)}/b$

Rubi [A] time = 0.05, antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {388, 195, 217, 206}

$$\frac{5a^3(8bc - ad) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{128b^{3/2}} + \frac{5a^2x\sqrt{a+bx^2}(8bc - ad)}{128b} + \frac{x(a+bx^2)^{5/2}(8bc - ad)}{48b} + \frac{5ax(a+bx^2)^{3/2}(8bc - ad)}{192b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*x^2)^{(5/2)}*(c + d*x^2), x]$

[Out] $(5*a^2*(8*b*c - a*d)*x*\text{Sqrt}[a + b*x^2])/(128*b) + (5*a*(8*b*c - a*d)*x*(a + b*x^2)^{(3/2)})/(192*b) + ((8*b*c - a*d)*x*(a + b*x^2)^{(5/2)})/(48*b) + (d*x*(a + b*x^2)^{(7/2)})/(8*b) + (5*a^3*(8*b*c - a*d)*\text{ArcTanh}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a + b*x^2]])/(128*b^{(3/2)})$

Rule 195

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] := \text{Simp}[(x*(a + b*x^n)^p)/(n*p + 1), x] + \text{Dist}[(a*n*p)/(n*p + 1), \text{Int}[(a + b*x^n)^{(p-1)}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] := \text{Simp}[(1*\text{ArcTanh}[(\text{Rt}[-b, 2]*x)/\text{Rt}[a, 2]])/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]), x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

$\text{Int}[1/\text{Sqrt}[(a_ + (b_)*(x_)^2)], x_Symbol] := \text{Subst}[\text{Int}[1/(1 - b*x^2), x], x, x/\text{Sqrt}[a + b*x^2]] /;$ FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}*((c_ + (d_)*(x_)^{(n_)})], x_Symbol] := \text{Simp}[(d*x*(a + b*x^n)^{(p+1)})/(b*(n*(p+1) + 1)), x] - \text{Dist}[(a*d - b*c*(n*(p+1) + 1))/(b*(n*(p+1) + 1)), \text{Int}[(a + b*x^n)^p, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1) + 1, 0]

Rubi steps

$$\begin{aligned}
\int (a + bx^2)^{5/2} (c + dx^2) dx &= \frac{dx (a + bx^2)^{7/2}}{8b} - \frac{(-8bc + ad) \int (a + bx^2)^{5/2} dx}{8b} \\
&= \frac{(8bc - ad)x (a + bx^2)^{5/2}}{48b} + \frac{dx (a + bx^2)^{7/2}}{8b} + \frac{(5a(8bc - ad)) \int (a + bx^2)^{3/2} dx}{48b} \\
&= \frac{5a(8bc - ad)x (a + bx^2)^{3/2}}{192b} + \frac{(8bc - ad)x (a + bx^2)^{5/2}}{48b} + \frac{dx (a + bx^2)^{7/2}}{8b} + \frac{(5a^2)}{48b} \\
&= \frac{5a^2(8bc - ad)x\sqrt{a + bx^2}}{128b} + \frac{5a(8bc - ad)x (a + bx^2)^{3/2}}{192b} + \frac{(8bc - ad)x (a + bx^2)^{5/2}}{48b} \\
&= \frac{5a^2(8bc - ad)x\sqrt{a + bx^2}}{128b} + \frac{5a(8bc - ad)x (a + bx^2)^{3/2}}{192b} + \frac{(8bc - ad)x (a + bx^2)^{5/2}}{48b} \\
&= \frac{5a^2(8bc - ad)x\sqrt{a + bx^2}}{128b} + \frac{5a(8bc - ad)x (a + bx^2)^{3/2}}{192b} + \frac{(8bc - ad)x (a + bx^2)^{5/2}}{48b}
\end{aligned}$$

Mathematica [A] time = 0.23, size = 130, normalized size = 0.87

$$\frac{\sqrt{a + bx^2} \left(\sqrt{b} x (15a^3d + 2a^2b(132c + 59dx^2)) + 8ab^2x^2(26c + 17dx^2) + 16b^3x^4(4c + 3dx^2) \right) - \frac{15a^{5/2}(ad - 8bc) \operatorname{arcsinh}\left(\frac{\sqrt{bx^2 + a}}{a}\right)}{\sqrt{\frac{bx^2}{a} + 1}}}{384b^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2)*(c + d*x^2), x]

[Out] (Sqrt[a + b*x^2]*(Sqrt[b]*x*(15*a^3*d + 16*b^3*x^4*(4*c + 3*d*x^2) + 8*a*b^2*x^2*(26*c + 17*d*x^2) + 2*a^2*b*(132*c + 59*d*x^2)) - (15*a^(5/2)*(-8*b*c + a*d)*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/Sqrt[1 + (b*x^2)/a])/(384*b^(3/2))

fricas [A] time = 0.84, size = 260, normalized size = 1.74

$$\frac{15(8a^3bc - a^4d)\sqrt{b} \log(-2bx^2 + 2\sqrt{bx^2 + a}\sqrt{b}x - a) - 2(48b^4dx^7 + 8(8b^4c + 17ab^3d)x^5 + 2(104ab^3c + 59a^2b^2d)x^3 + 3(88a^2b^2c + 5a^3b^2d)x)\sqrt{bx^2 + a}}{768b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)*(d*x^2+c), x, algorithm="fricas")

[Out] [-1/768*(15*(8*a^3*b*c - a^4*d)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(48*b^4*d*x^7 + 8*(8*b^4*c + 17*a*b^3*d)*x^5 + 2*(104*a*b^3*c + 59*a^2*b^2*d)*x^3 + 3*(88*a^2*b^2*c + 5*a^3*b^2*d)*x)*sqrt(b*x^2 + a))/b^2, -1/384*(15*(8*a^3*b*c - a^4*d)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (48*b^4*d*x^7 + 8*(8*b^4*c + 17*a*b^3*d)*x^5 + 2*(104*a*b^3*c + 59*a^2*b^2*d)*x^3 + 3*(88*a^2*b^2*c + 5*a^3*b^2*d)*x)*sqrt(b*x^2 + a))/b^2]

giac [A] time = 0.63, size = 135, normalized size = 0.91

$$\frac{1}{384} \left(2 \left(4 \left(6b^2dx^2 + \frac{8b^8c + 17ab^7d}{b^6} \right) x^2 + \frac{104ab^7c + 59a^2b^6d}{b^6} \right) x^2 + \frac{3(88a^2b^6c + 5a^3b^5d)}{b^6} \right) \sqrt{bx^2 + a} x - \frac{5(8a^3bc - a^4d)\sqrt{b} \operatorname{arcsinh}\left(\frac{\sqrt{bx^2 + a}}{a}\right)}{384b^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)*(d*x^2+c),x, algorithm="giac")

[Out] $\frac{1}{384} \cdot (2 \cdot (4 \cdot (6 \cdot b^2 \cdot d \cdot x^2 + (8 \cdot b^8 \cdot c + 17 \cdot a \cdot b^7 \cdot d) / b^6) \cdot x^2 + (104 \cdot a \cdot b^7 \cdot c + 59 \cdot a^2 \cdot b^6 \cdot d) / b^6) \cdot x^2 + 3 \cdot (88 \cdot a^2 \cdot b^6 \cdot c + 5 \cdot a^3 \cdot b^5 \cdot d) / b^6) \cdot \sqrt{b \cdot x^2 + a} \cdot x - \frac{5}{128} \cdot (8 \cdot a^3 \cdot b \cdot c - a^4 \cdot d) \cdot \log(\text{abs}(-\sqrt{b} \cdot x + \sqrt{b \cdot x^2 + a})) / b^{(3/2)}$

maple [A] time = 0.01, size = 166, normalized size = 1.11

$$\frac{5a^4 d \ln\left(\sqrt{b} x + \sqrt{bx^2 + a}\right)}{128b^{\frac{3}{2}}} + \frac{5a^3 c \ln\left(\sqrt{b} x + \sqrt{bx^2 + a}\right)}{16\sqrt{b}} - \frac{5\sqrt{bx^2 + a} a^3 dx}{128b} + \frac{5\sqrt{bx^2 + a} a^2 cx}{16} - \frac{5(bx^2 + a)^{\frac{3}{2}} a}{192b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/2)*(d*x^2+c),x)

[Out] $\frac{1}{8} \cdot d \cdot x \cdot (b \cdot x^2 + a)^{(7/2)} / b - \frac{1}{48} \cdot d \cdot a / b \cdot x \cdot (b \cdot x^2 + a)^{(5/2)} - \frac{5}{192} \cdot d \cdot a^2 / b \cdot x \cdot (b \cdot x^2 + a)^{(3/2)} - \frac{5}{128} \cdot d \cdot a^3 / b \cdot x \cdot (b \cdot x^2 + a)^{(1/2)} - \frac{5}{128} \cdot d \cdot a^4 / b^{(3/2)} \cdot \ln(b^{(1/2)} \cdot x + (b \cdot x^2 + a)^{(1/2)}) + \frac{1}{6} \cdot c \cdot x \cdot (b \cdot x^2 + a)^{(5/2)} + \frac{5}{24} \cdot c \cdot a \cdot x \cdot (b \cdot x^2 + a)^{(3/2)} + \frac{5}{16} \cdot c \cdot a^2 \cdot x \cdot (b \cdot x^2 + a)^{(1/2)} + \frac{5}{16} \cdot c \cdot a^3 / b^{(1/2)} \cdot \ln(b^{(1/2)} \cdot x + (b \cdot x^2 + a)^{(1/2)})$

maxima [A] time = 1.40, size = 151, normalized size = 1.01

$$\frac{1}{6} (bx^2 + a)^{\frac{5}{2}} cx + \frac{5}{24} (bx^2 + a)^{\frac{3}{2}} acx + \frac{5}{16} \sqrt{bx^2 + a} a^2 cx + \frac{(bx^2 + a)^{\frac{7}{2}} dx}{8b} - \frac{(bx^2 + a)^{\frac{5}{2}} adx}{48b} - \frac{5(bx^2 + a)^{\frac{3}{2}} a^2 dx}{192b} - \frac{5\sqrt{bx^2 + a} a^3 dx}{192b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)*(d*x^2+c),x, algorithm="maxima")

[Out] $\frac{1}{6} \cdot (b \cdot x^2 + a)^{(5/2)} \cdot c \cdot x + \frac{5}{24} \cdot (b \cdot x^2 + a)^{(3/2)} \cdot a \cdot c \cdot x + \frac{5}{16} \cdot \sqrt{b \cdot x^2 + a} \cdot a^2 \cdot c \cdot x + \frac{1}{8} \cdot (b \cdot x^2 + a)^{(7/2)} \cdot d \cdot x / b - \frac{1}{48} \cdot (b \cdot x^2 + a)^{(5/2)} \cdot a \cdot d \cdot x / b - \frac{5}{192} \cdot (b \cdot x^2 + a)^{(3/2)} \cdot a^2 \cdot d \cdot x / b - \frac{5}{128} \cdot \sqrt{b \cdot x^2 + a} \cdot a^3 \cdot d \cdot x / b + \frac{5}{16} \cdot a^3 \cdot c \cdot \text{arcsinh}(b \cdot x / \sqrt{a \cdot b}) / \sqrt{b} - \frac{5}{128} \cdot a^4 \cdot d \cdot \text{arcsinh}(b \cdot x / \sqrt{a \cdot b}) / b^{(3/2)}$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (bx^2 + a)^{5/2} (dx^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)*(c + d*x^2),x)

[Out] int((a + b*x^2)^(5/2)*(c + d*x^2), x)

sympy [B] time = 29.53, size = 316, normalized size = 2.12

$$\frac{5a^{\frac{7}{2}} dx}{128b\sqrt{1 + \frac{bx^2}{a}}} + \frac{a^{\frac{5}{2}} cx \sqrt{1 + \frac{bx^2}{a}}}{2} + \frac{3a^{\frac{5}{2}} cx}{16\sqrt{1 + \frac{bx^2}{a}}} + \frac{133a^{\frac{5}{2}} dx^3}{384\sqrt{1 + \frac{bx^2}{a}}} + \frac{35a^{\frac{3}{2}} bcx^3}{48\sqrt{1 + \frac{bx^2}{a}}} + \frac{127a^{\frac{3}{2}} bdx^5}{192\sqrt{1 + \frac{bx^2}{a}}} + \frac{17\sqrt{a} b^2 cx^5}{24\sqrt{1 + \frac{bx^2}{a}}} + \frac{23\sqrt{a} b^2 cx^5}{48\sqrt{1 + \frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)*(d*x**2+c),x)

[Out] $5 \cdot a^{(7/2)} \cdot d \cdot x / (128 \cdot b \cdot \sqrt{1 + b \cdot x^{**2} / a}) + a^{(5/2)} \cdot c \cdot x \cdot \sqrt{1 + b \cdot x^{**2} / a} / 2 + 3 \cdot a^{(5/2)} \cdot c \cdot x / (16 \cdot \sqrt{1 + b \cdot x^{**2} / a}) + 133 \cdot a^{(5/2)} \cdot d \cdot x^{**3} / (384 \cdot \sqrt{1 + b \cdot x^{**2} / a}) + 35 \cdot a^{(3/2)} \cdot b \cdot c \cdot x^{**3} / (48 \cdot \sqrt{1 + b \cdot x^{**2} / a}) + 127 \cdot a^{(3/2)} \cdot b \cdot d \cdot x^{**5} / (192 \cdot \sqrt{1 + b \cdot x^{**2} / a}) + 17 \cdot \sqrt{a} \cdot b^{**2} \cdot c \cdot x^{**5} / (24 \cdot \sqrt{1 + b \cdot x^{**2} / a})$

$$\begin{aligned}
& b*x**2/a)) + 23*\sqrt{a}*b**2*d*x**7/(48*\sqrt{1 + b*x**2/a}) - 5*a**4*d*\operatorname{asin} \\
& \operatorname{h}(\sqrt{b}*x/\sqrt{a})/(128*b**(3/2)) + 5*a**3*c*\operatorname{asinh}(\sqrt{b}*x/\sqrt{a})/(16 \\
& *\sqrt{b}) + b**3*c*x**7/(6*\sqrt{a}*\sqrt{1 + b*x**2/a}) + b**3*d*x**9/(8*\sqrt{a}*\sqrt{1 + b*x**2/a})
\end{aligned}$$

3.65 $\int (a + bx^2)^{5/2} dx$

Optimal. Leaf size=84

$$\frac{5a^3 \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16\sqrt{b}} + \frac{5}{16}a^2x\sqrt{a+bx^2} + \frac{5}{24}ax(a+bx^2)^{3/2} + \frac{1}{6}x(a+bx^2)^{5/2}$$

[Out] $5/24*a*x*(b*x^2+a)^{(3/2)}+1/6*x*(b*x^2+a)^{(5/2)}+5/16*a^3*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(1/2)}+5/16*a^2*x*(b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {195, 217, 206}

$$\frac{5}{16}a^2x\sqrt{a+bx^2} + \frac{5a^3 \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16\sqrt{b}} + \frac{5}{24}ax(a+bx^2)^{3/2} + \frac{1}{6}x(a+bx^2)^{5/2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a + b*x^2)^{(5/2)}, x]$

[Out] $(5*a^2*x*\operatorname{Sqrt}[a + b*x^2])/16 + (5*a*x*(a + b*x^2)^{(3/2)})/24 + (x*(a + b*x^2)^{(5/2)})/6 + (5*a^3*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(16*\operatorname{Sqrt}[b])$

Rule 195

$\operatorname{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \operatorname{Simp}[(x*(a + b*x^n)^p)/(n*p + 1), x] + \operatorname{Dist}[(a*n*p)/(n*p + 1), \operatorname{Int}[(a + b*x^n)^{(p-1)}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{(-1)}, x_Symbol] \rightarrow \operatorname{Simp}[(1*\operatorname{ArcTanh}[(\operatorname{Rt}[-b, 2]*x)/\operatorname{Rt}[a, 2]])/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]), x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_)*(x_)^2)], x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /;$ FreeQ[{a, b}, x] && !GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^{5/2} dx &= \frac{1}{6}x(a + bx^2)^{5/2} + \frac{1}{6}(5a) \int (a + bx^2)^{3/2} dx \\ &= \frac{5}{24}ax(a + bx^2)^{3/2} + \frac{1}{6}x(a + bx^2)^{5/2} + \frac{1}{8}(5a^2) \int \sqrt{a + bx^2} dx \\ &= \frac{5}{16}a^2x\sqrt{a + bx^2} + \frac{5}{24}ax(a + bx^2)^{3/2} + \frac{1}{6}x(a + bx^2)^{5/2} + \frac{1}{16}(5a^3) \int \frac{1}{\sqrt{a + bx^2}} dx \\ &= \frac{5}{16}a^2x\sqrt{a + bx^2} + \frac{5}{24}ax(a + bx^2)^{3/2} + \frac{1}{6}x(a + bx^2)^{5/2} + \frac{1}{16}(5a^3) \operatorname{Subst}\left(\int \frac{1}{1 - bx^2} dx, x, \frac{x}{\sqrt{a + bx^2}}\right) \\ &= \frac{5}{16}a^2x\sqrt{a + bx^2} + \frac{5}{24}ax(a + bx^2)^{3/2} + \frac{1}{6}x(a + bx^2)^{5/2} + \frac{5a^3 \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16\sqrt{b}} \end{aligned}$$

Mathematica [A] time = 0.11, size = 76, normalized size = 0.90

$$\frac{1}{48} \sqrt{a + bx^2} \left(\frac{15a^{5/2} \sinh^{-1} \left(\frac{\sqrt{bx}}{\sqrt{a}} \right)}{\sqrt{b} \sqrt{\frac{bx^2}{a} + 1}} + 33a^2x + 26abx^3 + 8b^2x^5 \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2), x]

[Out] (Sqrt[a + b*x^2]*(33*a^2*x + 26*a*b*x^3 + 8*b^2*x^5 + (15*a^(5/2)*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/(Sqrt[b]*Sqrt[1 + (b*x^2)/a]))) / 48

fricas [A] time = 0.63, size = 146, normalized size = 1.74

$$\left[\frac{15 a^3 \sqrt{b} \log \left(-2 b x^2 - 2 \sqrt{b x^2 + a} \sqrt{b} x - a \right) + 2 \left(8 b^3 x^5 + 26 a b^2 x^3 + 33 a^2 b x \right) \sqrt{b x^2 + a}}{96 b}, - \frac{15 a^3 \sqrt{-b} \arctan \left(\frac{\sqrt{b} x}{\sqrt{b x^2 + a}} \right)}{96 b} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2), x, algorithm="fricas")

[Out] [1/96*(15*a^3*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 2*(8*b^3*x^5 + 26*a*b^2*x^3 + 33*a^2*b*x)*sqrt(b*x^2 + a))/b, -1/48*(15*a^3*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (8*b^3*x^5 + 26*a*b^2*x^3 + 33*a^2*b*x)*sqrt(b*x^2 + a))/b]

giac [A] time = 0.61, size = 63, normalized size = 0.75

$$-\frac{5 a^3 \log \left(\left| -\sqrt{b} x + \sqrt{b x^2 + a} \right| \right)}{16 \sqrt{b}} + \frac{1}{48} \left(2 \left(4 b^2 x^2 + 13 a b \right) x^2 + 33 a^2 \right) \sqrt{b x^2 + a} x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2), x, algorithm="giac")

[Out] -5/16*a^3*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/sqrt(b) + 1/48*(2*(4*b^2*x^2 + 13*a*b)*x^2 + 33*a^2)*sqrt(b*x^2 + a)*x

maple [A] time = 0.00, size = 66, normalized size = 0.79

$$\frac{5 a^3 \ln \left(\sqrt{b} x + \sqrt{b x^2 + a} \right)}{16 \sqrt{b}} + \frac{5 \sqrt{b x^2 + a} a^2 x}{16} + \frac{5 \left(b x^2 + a \right)^{\frac{3}{2}} a x}{24} + \frac{\left(b x^2 + a \right)^{\frac{5}{2}} x}{6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/2), x)

[Out] 1/6*(b*x^2+a)^(5/2)*x+5/24*(b*x^2+a)^(3/2)*a*x+5/16*(b*x^2+a)^(1/2)*a^2*x+5/16*a^3/b^(1/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))

maxima [A] time = 1.33, size = 58, normalized size = 0.69

$$\frac{1}{6} \left(b x^2 + a \right)^{\frac{5}{2}} x + \frac{5}{24} \left(b x^2 + a \right)^{\frac{3}{2}} a x + \frac{5}{16} \sqrt{b x^2 + a} a^2 x + \frac{5 a^3 \operatorname{arsinh} \left(\frac{b x}{\sqrt{a b}} \right)}{16 \sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2),x, algorithm="maxima")

[Out] $\frac{1}{6}(bx^2 + a)^{5/2}x + \frac{5}{24}(bx^2 + a)^{3/2}ax + \frac{5}{16}\sqrt{bx^2 + a}a^2x + \frac{5}{16}a^3\operatorname{arcsinh}(bx/\sqrt{a})/\sqrt{b}$

mupad [B] time = 4.69, size = 37, normalized size = 0.44

$$\frac{x(bx^2 + a)^{5/2} {}_2F_1\left(-\frac{5}{2}, \frac{1}{2}; \frac{3}{2}; -\frac{bx^2}{a}\right)}{\left(\frac{bx^2}{a} + 1\right)^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2),x)

[Out] $(x(a + bx^2)^{5/2} \operatorname{hypergeom}([-5/2, 1/2], 3/2, -(bx^2)/a)) / ((bx^2)/a + 1)^{5/2}$

sympy [A] time = 4.28, size = 97, normalized size = 1.15

$$\frac{11a^{\frac{5}{2}}x\sqrt{1 + \frac{bx^2}{a}}}{16} + \frac{13a^{\frac{3}{2}}bx^3\sqrt{1 + \frac{bx^2}{a}}}{24} + \frac{\sqrt{a}b^2x^5\sqrt{1 + \frac{bx^2}{a}}}{6} + \frac{5a^3 \operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{16\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2),x)

[Out] $11*a^{5/2}*x*\sqrt{1 + b*x**2/a}/16 + 13*a^{3/2}*b*x**3*\sqrt{1 + b*x**2/a}/24 + \sqrt{a}*b**2*x**5*\sqrt{1 + b*x**2/a}/6 + 5*a**3*\operatorname{asinh}(\sqrt{b}*x/\sqrt{a})/(16*\sqrt{b})$

$$3.66 \quad \int \frac{(a+bx^2)^{5/2}}{c+dx^2} dx$$

Optimal. Leaf size=157

$$\frac{\sqrt{b} (15a^2d^2 - 20abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{8d^3} - \frac{(bc - ad)^{5/2} \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d^3} - \frac{bx\sqrt{a+bx^2}(4bc - 7ad)}{8d^2} + \frac{b}{d}$$

[Out] $1/4*b*x*(b*x^2+a)^{(3/2)}/d+1/8*(15*a^2*d^2-20*a*b*c*d+8*b^2*c^2)*\operatorname{arctanh}(x*b^{1/2}/(b*x^2+a)^{(1/2}))*b^{1/2}/d^3-(-a*d+b*c)^{(5/2)*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{1/2}/(b*x^2+a)^{(1/2}))/d^3/c^{1/2}-1/8*b*(-7*a*d+4*b*c)*x*(b*x^2+a)^{(1/2)}/d^2$

Rubi [A] time = 0.20, antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {416, 528, 523, 217, 206, 377, 208}

$$\frac{\sqrt{b} (15a^2d^2 - 20abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{8d^3} - \frac{bx\sqrt{a+bx^2}(4bc - 7ad)}{8d^2} - \frac{(bc - ad)^{5/2} \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}d^3} + \frac{b}{d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(5/2)/(c + d*x^2), x]

[Out] $-(b*(4*b*c - 7*a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(8*d^2) + (b*x*(a + b*x^2)^{(3/2)})/(4*d) + (\operatorname{Sqrt}[b]*(8*b^2*c^2 - 20*a*b*c*d + 15*a^2*d^2)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(8*d^3) - ((b*c - a*d)^{(5/2)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(8*d^3)$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q) + 1)), x] + Dist[1/(b*(n*(p+q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q) + 1) - a*d) + d*(b*c*(n*(p+2*q-1) + 1) - a*d*(n*(q-1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,

0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 523

Int[((e_) + (f_)*(x_)^(n_))/(((a_) + (b_)*(x_)^(n_))*Sqrt[(c_) + (d_)*(x_)^(n_)]), x_Symbol] :> Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 528

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] :> Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q) + 1)), x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{5/2}}{c + dx^2} dx &= \frac{bx(a + bx^2)^{3/2}}{4d} + \frac{\int \frac{\sqrt{a+bx^2}(-a(bc-4ad)-b(4bc-7ad)x^2)}{c+dx^2} dx}{4d} \\ &= -\frac{b(4bc-7ad)x\sqrt{a+bx^2}}{8d^2} + \frac{bx(a + bx^2)^{3/2}}{4d} + \frac{\int \frac{a(4b^2c^2-9abcd+8a^2d^2)+b(8b^2c^2-20abcd+15a^2d^2)x^2}{\sqrt{a+bx^2}(c+dx^2)} dx}{8d^2} \\ &= -\frac{b(4bc-7ad)x\sqrt{a+bx^2}}{8d^2} + \frac{bx(a + bx^2)^{3/2}}{4d} - \frac{(bc-ad)^3 \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{d^3} + \frac{b(8b^2c^2-20abcd+15a^2d^2)}{8d^3} \\ &= -\frac{b(4bc-7ad)x\sqrt{a+bx^2}}{8d^2} + \frac{bx(a + bx^2)^{3/2}}{4d} - \frac{(bc-ad)^3 \text{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{d^3} \\ &= -\frac{b(4bc-7ad)x\sqrt{a+bx^2}}{8d^2} + \frac{bx(a + bx^2)^{3/2}}{4d} + \frac{\sqrt{b}(8b^2c^2-20abcd+15a^2d^2) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{8d^3} \end{aligned}$$

Mathematica [A] time = 0.12, size = 140, normalized size = 0.89

$$\frac{\sqrt{b}(15a^2d^2 - 20abcd + 8b^2c^2) \log\left(\sqrt{b}\sqrt{a+bx^2} + bx\right) + bdx\sqrt{a+bx^2}(9ad - 4bc + 2bdx^2) + \frac{8(ad-bc)^{5/2} \tan^{-1}\left(\frac{x\sqrt{b}}{\sqrt{c}}\right)}{\sqrt{c}}}{8d^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2)/(c + d*x^2), x]

[Out] (b*d*x*Sqrt[a + b*x^2]*(-4*b*c + 9*a*d + 2*b*d*x^2) + (8*(-(b*c) + a*d)^(5/2)*ArcTan[(Sqrt[-(b*c) + a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/Sqrt[c] + Sqrt[b]*(8*b^2*c^2 - 20*a*b*c*d + 15*a^2*d^2)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2])/ (8*d^3)

fricas [A] time = 3.23, size = 935, normalized size = 5.96

$$\frac{\left(8b^2c^2 - 20abcd + 15a^2d^2\right)\sqrt{b} \log\left(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{b}x - a\right) + 4\left(b^2c^2 - 2abcd + a^2d^2\right)\sqrt{\frac{bc-ad}{c}} \log\left(\frac{8}{\dots}\right)}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c),x, algorithm="fricas")

[Out] [1/16*((8*b^2*c^2 - 20*a*b*c*d + 15*a^2*d^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 4*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*sqrt(b*x^2 + a)*sqrt((b*c - a*d)/c))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 2*(2*b^2*d^2*x^3 - (4*b^2*c*d - 9*a*b*d^2)*x)*sqrt(b*x^2 + a))/d^3, -1/8*((8*b^2*c^2 - 20*a*b*c*d + 15*a^2*d^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - 2*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*sqrt(b*x^2 + a)*sqrt((b*c - a*d)/c))/(d^2*x^4 + 2*c*d*x^2 + c^2)) - (2*b^2*d^2*x^3 - (4*b^2*c*d - 9*a*b*d^2)*x)*sqrt(b*x^2 + a))/d^3, 1/16*(8*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(-(b*c - a*d)/c)*arctan(1/2*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)*sqrt(-(b*c - a*d)/c))/((b^2*c - a*b*d)*x^3 + (a*b*c - a^2*d)*x)) + (8*b^2*c^2 - 20*a*b*c*d + 15*a^2*d^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 2*(2*b^2*d^2*x^3 - (4*b^2*c*d - 9*a*b*d^2)*x)*sqrt(b*x^2 + a))/d^3, -1/8*((8*b^2*c^2 - 20*a*b*c*d + 15*a^2*d^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - 4*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(-(b*c - a*d)/c)*arctan(1/2*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)*sqrt(-(b*c - a*d)/c))/((b^2*c - a*b*d)*x^3 + (a*b*c - a^2*d)*x)) - (2*b^2*d^2*x^3 - (4*b^2*c*d - 9*a*b*d^2)*x)*sqrt(b*x^2 + a))/d^3]

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,x):;OUTPUT:

maple [B] time = 0.02, size = 3053, normalized size = 19.45

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/2)/(d*x^2+c),x)

[Out]
$$\frac{-1/(-c*d)^{(1/2)}/d*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*a*b*c+1/2/(-c*d)^{(1/2)}/d^3/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}}{(x-(-c*d)^{(1/2)}/d)*b^3*c^3+1/(-c*d)^{(1/2)}/d*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*a*b*c-1/2/(-c*d)^{(1/2)}/d^3/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))}}{(x+(-c*d)^{(1/2)}/d)*b^3*c^3+3/2/(-c*d)^{(1/2)}/d^2/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/$$

$d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}/(x+(-c*d)^{(1/2)}/d))*a*b^2*c^2+3/2/(-c*d)^{(1/2)}/d/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}/(x-(-c*d)^{(1/2)}/d))*a^2*b*c-3/2/(-c*d)^{(1/2)}/d^2/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}/(x-(-c*d)^{(1/2)}/d))*a*b^2*c^2-3/2/(-c*d)^{(1/2)}/d/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}/(x+(-c*d)^{(1/2)}/d))*a^2*b*c+1/8*b/d*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*x+1/8*b/d*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*x+15/16/d*b^(1/2)*\ln(((x-(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)}*b/d)/b^(1/2))+((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*a^2+1/2/d^3*b^(5/2)*\ln(((x-(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)}*b/d)/b^(1/2))+((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*c^2-1/2/(-c*d)^{(1/2)}/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}/(x-(-c*d)^{(1/2)}/d))*a^3+15/16/d*b^(1/2)*\ln(((x+(-c*d)^{(1/2)}/d)*b-(-c*d)^{(1/2)}*b/d)/b^(1/2))+((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*a^2+1/2/d^3*b^(5/2)*\ln(((x+(-c*d)^{(1/2)}/d)*b-(-c*d)^{(1/2)}*b/d)/b^(1/2))+((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*c^2+1/2/(-c*d)^{(1/2)}/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}/(x+(-c*d)^{(1/2)}/d))*a^3-1/10/(-c*d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(5/2)}+1/10/(-c*d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(5/2)}-1/2/(-c*d)^{(1/2)}/d^2*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*b^2*c^2-1/6/(-c*d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*a-1/2/(-c*d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*a^2+1/6/(-c*d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*a+1/2/(-c*d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*a^2-1/6/(-c*d)^{(1/2)}/d*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*b*c+7/16*b/d*a*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x-1/4/d^2*b^2*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x*c-5/4/d^2*b^(3/2)*\ln(((x-(-c*d)^{(1/2)}/d)*b+(-c*d)^{(1/2)}*b/d)/b^(1/2))+((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*c*a+1/2/(-c*d)^{(1/2)}/d^2*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*b^2*c^2+7/16*b/d*a*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x+1/6/(-c*d)^{(1/2)}/d*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*b*c-1/4/d^2*b^2*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x*c-5/4/d^2*b^(3/2)*\ln(((x+(-c*d)^{(1/2)}/d)*b-(-c*d)^{(1/2)}*b/d)/b^(1/2))+((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2))*c*a$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/2)/(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{5/2}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)/(c + d*x^2), x)

[Out] int((a + b*x^2)^(5/2)/(c + d*x^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{5/2}}{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)/(d*x**2+c), x)

[Out] Integral((a + b*x**2)**(5/2)/(c + d*x**2), x)

$$3.67 \quad \int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^2} dx$$

Optimal. Leaf size=175

$$\frac{b^{3/2}(4bc - 5ad) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{2d^3} + \frac{(bc - ad)^{3/2}(ad + 4bc) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}d^3} + \frac{bx\sqrt{a+bx^2}(2bc - ad)}{2cd^2} - \frac{x(a+bx^2)}{2cd}$$

[Out] $-1/2*(-a*d+b*c)*x*(b*x^2+a)^{(3/2)}/c/d/(d*x^2+c)-1/2*b^{(3/2)*(-5*a*d+4*b*c)*\arctanh(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/d^3+1/2*(-a*d+b*c)^{(3/2)*(a*d+4*b*c)*\arctanh(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})/c^{(3/2)}/d^3+1/2*b*(-a*d+2*b*c)*x*(b*x^2+a)^{(1/2)}/c/d^2$

Rubi [A] time = 0.23, antiderivative size = 175, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {413, 528, 523, 217, 206, 377, 208}

$$\frac{b^{3/2}(4bc - 5ad) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{2d^3} + \frac{(bc - ad)^{3/2}(ad + 4bc) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}d^3} + \frac{bx\sqrt{a+bx^2}(2bc - ad)}{2cd^2} - \frac{x(a+bx^2)}{2cd}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(5/2)/(c + d*x^2)^2, x]

[Out] $(b*(2*b*c - a*d)*x*\text{Sqrt}[a + b*x^2])/(2*c*d^2) - ((b*c - a*d)*x*(a + b*x^2)^{(3/2)})/(2*c*d*(c + d*x^2)) - (b^{(3/2)}*(4*b*c - 5*a*d)*\text{ArcTanh}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a + b*x^2]])/(2*d^3) + ((b*c - a*d)^{(3/2)}*(4*b*c + a*d)*\text{ArcTanh}[(\text{Sqrt}[b*c - a*d]*x)/(\text{Sqrt}[c]*\text{Sqrt}[a + b*x^2])])/(2*c^{(3/2)}*d^3)$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[Rt[-b, 2]*x]/Rt[a, 2])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1) + 1)) + d*(a*d*(n*(q-1) + 1) - b*c*(n*(p

+ q) + 1)) * x^n, x], x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 523

Int[((e_) + (f_)*(x_)^(n_))/(((a_) + (b_)*(x_)^(n_))*Sqrt[(c_) + (d_)*(x_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 528

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{5/2}}{(c + dx^2)^2} dx &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{2cd(c + dx^2)} + \frac{\int \frac{\sqrt{a+bx^2}(a(bc+ad)+2b(2bc-ad)x^2)}{c+dx^2} dx}{2cd} \\ &= \frac{b(2bc - ad)x\sqrt{a + bx^2}}{2cd^2} - \frac{(bc - ad)x(a + bx^2)^{3/2}}{2cd(c + dx^2)} + \frac{\int \frac{-2a(2b^2c^2 - 2abcd - a^2d^2) - 2b^2c(4bc - 5ad)x^2}{\sqrt{a+bx^2}(c+dx^2)} dx}{4cd^2} \\ &= \frac{b(2bc - ad)x\sqrt{a + bx^2}}{2cd^2} - \frac{(bc - ad)x(a + bx^2)^{3/2}}{2cd(c + dx^2)} - \frac{(b^2(4bc - 5ad)) \int \frac{1}{\sqrt{a+bx^2}} dx}{2d^3} + \frac{(bc - ad)x(a + bx^2)^{3/2}}{2cd(c + dx^2)} \\ &= \frac{b(2bc - ad)x\sqrt{a + bx^2}}{2cd^2} - \frac{(bc - ad)x(a + bx^2)^{3/2}}{2cd(c + dx^2)} - \frac{(b^2(4bc - 5ad)) \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \sqrt{a+bx^2}\right)}{2d^3} \\ &= \frac{b(2bc - ad)x\sqrt{a + bx^2}}{2cd^2} - \frac{(bc - ad)x(a + bx^2)^{3/2}}{2cd(c + dx^2)} - \frac{b^{3/2}(4bc - 5ad) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{2d^3} + \frac{(bc - ad)x(a + bx^2)^{3/2}}{2cd(c + dx^2)} \end{aligned}$$

Mathematica [A] time = 0.16, size = 144, normalized size = 0.82

$$\frac{-\left(b^{3/2}(4bc - 5ad) \log\left(\sqrt{b} \sqrt{a + bx^2} + bx\right)\right) + dx\sqrt{a + bx^2} \left(\frac{(bc - ad)^2}{c(c + dx^2)} + b^2\right) + \frac{(ad - bc)^{3/2}(ad + 4bc) \tan^{-1}\left(\frac{x\sqrt{ad - bc}}{\sqrt{c} \sqrt{a + bx^2}}\right)}{c^{3/2}}}{2d^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2)/(c + d*x^2)^2, x]

[Out] (d*x*Sqrt[a + b*x^2]*(b^2 + (b*c - a*d)^2/(c*(c + d*x^2))) + ((-(b*c) + a*d)^(3/2)*(4*b*c + a*d)*ArcTan[(Sqrt[-(b*c) + a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])]/c^(3/2) - b^(3/2)*(4*b*c - 5*a*d)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]]/(2*d^3)

fricas [A] time = 2.20, size = 1236, normalized size = 7.06

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^2,x, algorithm="fricas")

[Out]
$$\begin{aligned} & [-1/8*(2*(4*b^2*c^3 - 5*a*b*c^2*d + (4*b^2*c^2*d - 5*a*b*c*d^2)*x^2)*\sqrt{b} \\ &)*\log(-2*b*x^2 - 2*\sqrt{b*x^2 + a}*\sqrt{b}*x - a) + (4*b^2*c^3 - 3*a*b*c^2*d \\ & - a^2*c*d^2 + (4*b^2*c^2*d - 3*a*b*c*d^2 - a^2*d^3)*x^2)*\sqrt{(b*c - a*d)} \\ & /c)*\log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3 \\ & *a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*\sqrt{b*x^2 + a}*\sqrt{(b \\ & *c - a*d)/c}))/d^2*x^4 + 2*c*d*x^2 + c^2)) - 4*(b^2*c*d^2*x^3 + (2*b^2*c^2*d \\ & - 2*a*b*c*d^2 + a^2*d^3)*x)*\sqrt{b*x^2 + a}))/c*d^4*x^2 + c^2*d^3), 1/8*(\\ & 4*(4*b^2*c^3 - 5*a*b*c^2*d + (4*b^2*c^2*d - 5*a*b*c*d^2)*x^2)*\sqrt{-b}*\arctan \\ & (\sqrt{-b}*x/\sqrt{b*x^2 + a}) - (4*b^2*c^3 - 3*a*b*c^2*d - a^2*c*d^2 + (4*b^2*c^2*d \\ & - 3*a*b*c*d^2 - a^2*d^3)*x^2)*\sqrt{(b*c - a*d)/c})*\log(((8*b^2*c^2 \\ & - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(\\ & a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*\sqrt{b*x^2 + a}*\sqrt{(b*c - a*d)/c}))/d^2*x^4 \\ & + 2*c*d*x^2 + c^2)) + 4*(b^2*c*d^2*x^3 + (2*b^2*c^2*d - 2*a*b*c*d^2 + a^2*d^3)*x) \\ &)*\sqrt{b*x^2 + a}))/c*d^4*x^2 + c^2*d^3), -1/4*((4*b^2*c^3 - 3*a*b*c^2*d - a^2*c*d^2 \\ & + (4*b^2*c^2*d - 3*a*b*c*d^2 - a^2*d^3)*x^2)*\sqrt{-(b*c - a*d)/c})*\arctan(1/2*((2*b*c \\ & - a*d)*x^2 + a*c)*\sqrt{b*x^2 + a}*\sqrt{-(b*c - a*d)/c}))/((b^2*c - a*b*d)*x^3 + (a*b*c \\ & - a^2*d)*x)) + (4*b^2*c^3 - 5*a*b*c^2*d + (4*b^2*c^2*d - 5*a*b*c*d^2)*x^2)*\sqrt{b} \\ &)*\log(-2*b*x^2 - 2*\sqrt{b*x^2 + a}*\sqrt{b}*x - a) - 2*(b^2*c*d^2*x^3 + (2*b^2*c^2*d \\ & - 2*a*b*c*d^2 + a^2*d^3)*x)*\sqrt{b*x^2 + a}))/c*d^4*x^2 + c^2*d^3), 1/4*(2*(4*b^2*c^3 \\ & - 5*a*b*c^2*d + (4*b^2*c^2*d - 5*a*b*c*d^2)*x^2)*\sqrt{-b}*\arctan(\sqrt{-b}*x/\sqrt{b*x^2 \\ & + a}) - (4*b^2*c^3 - 3*a*b*c^2*d - a^2*c*d^2 + (4*b^2*c^2*d - 3*a*b*c*d^2 \\ & - a^2*d^3)*x^2)*\sqrt{-(b*c - a*d)/c})*\arctan(1/2*((2*b*c - a*d)*x^2 + a*c)* \\ & \sqrt{b*x^2 + a}*\sqrt{-(b*c - a*d)/c}))/((b^2*c - a*b*d)*x^3 + (a*b*c - a^2*d)*x)) \\ & + 2*(b^2*c*d^2*x^3 + (2*b^2*c^2*d - 2*a*b*c*d^2 + a^2*d^3)*x)*\sqrt{b*x^2 + a}))/c*d^4*x^2 \\ & + c^2*d^3)] \end{aligned}$$

giac [B] time = 0.69, size = 405, normalized size = 2.31

$$\frac{\sqrt{bx^2 + a}b^2x}{2d^2} + \frac{\left(4b^{\frac{5}{2}}c - 5ab^{\frac{3}{2}}d\right) \log\left(\left(\sqrt{b}x - \sqrt{bx^2 + a}\right)^2\right)}{4d^3} - \frac{\left(4b^{\frac{7}{2}}c^3 - 7ab^{\frac{5}{2}}c^2d + 2a^2b^{\frac{3}{2}}cd^2 + a^3\sqrt{b}d^3\right) \arctan\left(\frac{x}{\sqrt{-b^2c^2 + abcd}cd^3}\right)}{2\sqrt{-b^2c^2 + abcd}cd^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^2,x, algorithm="giac")

[Out]
$$\begin{aligned} & 1/2*\sqrt{b*x^2 + a}*b^2*x/d^2 + 1/4*(4*b^(5/2)*c - 5*a*b^(3/2)*d)*\log((\sqrt{b} \\ &)*x - \sqrt{b*x^2 + a})^2/d^3 - 1/2*(4*b^(7/2)*c^3 - 7*a*b^(5/2)*c^2*d + \\ & 2*a^2*b^(3/2)*c*d^2 + a^3*\sqrt{b}*d^3)*\arctan(1/2*((\sqrt{b}*x - \sqrt{b*x^2 \\ & + a})^2*d + 2*b*c - a*d)/\sqrt{-b^2*c^2 + a*b*c*d}))/(\sqrt{-b^2*c^2 + a*b*c*d} \\ &)*c*d^3) + (2*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*b^(7/2)*c^3 - 5*(\sqrt{b}*x - \\ & \sqrt{b*x^2 + a})^2*a*b^(5/2)*c^2*d + 4*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*a^2* \\ & b^(3/2)*c*d^2 - (\sqrt{b}*x - \sqrt{b*x^2 + a})^2*a^3*\sqrt{b}*d^3 + a^2*b^(5/2) \\ &)*c^2*d - 2*a^3*b^(3/2)*c*d^2 + a^4*\sqrt{b}*d^3)/(((\sqrt{b}*x - \sqrt{b*x^2 \\ & + a})^4*d + 4*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*b*c - 2*(\sqrt{b}*x - \sqrt{b*x^2 \\ & + a})^2*a*d + a^2*d)*c*d^3) \end{aligned}$$

maple [B] time = 0.02, size = 7345, normalized size = 41.97

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/2)/(d*x^2+c)^2,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/2)/(d*x^2 + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{5/2}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)/(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(5/2)/(c + d*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{5}{2}}}{(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)/(d*x**2+c)**2,x)

[Out] Integral((a + b*x**2)**(5/2)/(c + d*x**2)**2, x)

$$3.68 \quad \int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^3} dx$$

Optimal. Leaf size=194

$$\frac{\sqrt{bc-ad} (3a^2d^2 + 4abcd + 8b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}d^3} + \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{d^3} - \frac{x\sqrt{a+bx^2} (bc-ad)(3ad+4bc)}{8c^2d^2(c+dx^2)}$$

[Out] $-1/4*(-a*d+b*c)*x*(b*x^2+a)^{(3/2)}/c/d/(d*x^2+c)^2+b^{(5/2)*\operatorname{arctanh}(x*b^{(1/2)})/(b*x^2+a)^{(1/2)})/d^3-1/8*(3*a^2*d^2+4*a*b*c*d+8*b^2*c^2)*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)})/(b*x^2+a)^{(1/2)}*(-a*d+b*c)^{(1/2)}/c^{(5/2)}/d^3-1/8*(-a*d+b*c)*(3*a*d+4*b*c)*x*(b*x^2+a)^{(1/2)}/c^2/d^2/(d*x^2+c)$

Rubi [A] time = 0.19, antiderivative size = 194, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {413, 526, 523, 217, 206, 377, 208}

$$\frac{\sqrt{bc-ad} (3a^2d^2 + 4abcd + 8b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}d^3} + \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{d^3} - \frac{x\sqrt{a+bx^2} (bc-ad)(3ad+4bc)}{8c^2d^2(c+dx^2)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a + b*x^2)^{(5/2)}/(c + d*x^2)^3, x]$

[Out] $-((b*c - a*d)*x*(a + b*x^2)^{(3/2)})/(4*c*d*(c + d*x^2)^2) - ((b*c - a*d)*(4*b*c + 3*a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(8*c^2*d^2*(c + d*x^2)) + (b^{(5/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[b]*x/\operatorname{Sqrt}[a + b*x^2]]})/d^3 - (\operatorname{Sqrt}[b*c - a*d]*(8*b^2*c^2 + 4*a*b*c*d + 3*a^2*d^2)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(8*c^{(5/2)*d^3})$

Rule 206

$\operatorname{Int}[(a_ + (b_.)*(x_)^2)^{-1}, x_Symbol] :> \operatorname{Simp}[(1*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*x]/\operatorname{Rt}[a, 2])]/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]), x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 208

$\operatorname{Int}[(a_ + (b_.)*(x_)^2)^{-1}, x_Symbol] :> \operatorname{Simp}[(\operatorname{Rt}[-(a/b), 2]*\operatorname{ArcTanh}[x/\operatorname{Rt}[-(a/b), 2]])/a, x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b]$

Rule 217

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_.)*(x_)^2)], x_Symbol] :> \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 377

$\operatorname{Int}[(a_ + (b_.)*(x_)^{(n_)})^{(p_)}]/((c_ + (d_.)*(x_)^{(n_)}), x_Symbol] :> \operatorname{Subst}[\operatorname{Int}[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^{(1/n)}] /; \operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \operatorname{EqQ}[n*p + 1, 0] \ \&\& \operatorname{IntegerQ}[n]$

Rule 413

$\operatorname{Int}[(a_ + (b_.)*(x_)^{(n_)})^{(p_)}*((c_ + (d_.)*(x_)^{(n_)})^{(q_)}), x_Symbol] :> \operatorname{Simp}[(a*d - c*b)*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-1)}/(a*b*n*(p + 1))]$

1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 523

Int[((e_) + (f_)*(x_)^(n_))/(((a_) + (b_)*(x_)^(n_))*Sqrt[(c_) + (d_)*(x_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 526

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{5/2}}{(c + dx^2)^3} dx &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{4cd(c + dx^2)^2} + \frac{\int \frac{\sqrt{a+bx^2}(a(bc+3ad)+4b^2cx^2)}{(c+dx^2)^2} dx}{4cd} \\ &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{4cd(c + dx^2)^2} - \frac{(bc - ad)(4bc + 3ad)x\sqrt{a + bx^2}}{8c^2d^2(c + dx^2)} - \frac{\int \frac{-a(4b^2c^2+ad(bc+3ad))-8b^3c^2x^2}{\sqrt{a+bx^2}(c+dx^2)} dx}{8c^2d^2} \\ &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{4cd(c + dx^2)^2} - \frac{(bc - ad)(4bc + 3ad)x\sqrt{a + bx^2}}{8c^2d^2(c + dx^2)} + \frac{b^3 \int \frac{1}{\sqrt{a+bx^2}} dx}{d^3} - \frac{(bc - ad)}{d^3} \\ &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{4cd(c + dx^2)^2} - \frac{(bc - ad)(4bc + 3ad)x\sqrt{a + bx^2}}{8c^2d^2(c + dx^2)} + \frac{b^3 \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\sqrt{a+bx^2}}{\sqrt{a}}\right)}{d^3} \\ &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{4cd(c + dx^2)^2} - \frac{(bc - ad)(4bc + 3ad)x\sqrt{a + bx^2}}{8c^2d^2(c + dx^2)} + \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{d^3} - \frac{\sqrt{b}}{d^3} \end{aligned}$$

Mathematica [A] time = 0.20, size = 184, normalized size = 0.95

$$\frac{(3a^3d^3+a^2bcd^2+4ab^2c^2d-8b^3c^3) \tan^{-1}\left(\frac{x\sqrt{ad-bc}}{\sqrt{c}\sqrt{a+bx^2}}\right) + 8b^{5/2} \log\left(\sqrt{b}\sqrt{a+bx^2} + bx\right) + \frac{dx\sqrt{a+bx^2}(ad-bc)(ad(5c+3dx^2)+2bc(2c+3dx^2))}{c^2(c+dx^2)^2}}{8d^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2)/(c + d*x^2)^3,x]

[Out] ((d*(-(b*c) + a*d)*x*Sqrt[a + b*x^2]*(2*b*c*(2*c + 3*d*x^2) + a*d*(5*c + 3*d*x^2)))/(c^2*(c + d*x^2)^2) + ((-8*b^3*c^3 + 4*a*b^2*c^2*d + a^2*b*c*d^2 + 3*a^3*d^3)*ArcTan[(Sqrt[-(b*c) + a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(c^5

/2)*Sqrt[-(b*c) + a*d]) + 8*b^(5/2)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(8*d^3)

fricas [B] time = 1.30, size = 1517, normalized size = 7.82

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [1/32*(16*(b^2*c^2*d^2*x^4 + 2*b^2*c^3*d*x^2 + b^2*c^4)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + (8*b^2*c^4 + 4*a*b*c^3*d + 3*a^2*c^2*d^2 + (8*b^2*c^2*d^2 + 4*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(8*b^2*c^3*d + 4*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*sqrt(b*x^2 + a)*sqrt((b*c - a*d)/c)))/(d^2*x^4 + 2*c*d*x^2 + c^2)) - 4*(3*(2*b^2*c^2*d^2 - a*b*c*d^3 - a^2*d^4)*x^3 + (4*b^2*c^3*d + a*b*c^2*d^2 - 5*a^2*c*d^3)*x)*sqrt(b*x^2 + a))/(c^2*d^5*x^4 + 2*c^3*d^4*x^2 + c^4*d^3), -1/32*(32*(b^2*c^2*d^2*x^4 + 2*b^2*c^3*d*x^2 + b^2*c^4)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (8*b^2*c^4 + 4*a*b*c^3*d + 3*a^2*c^2*d^2 + (8*b^2*c^2*d^2 + 4*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(8*b^2*c^3*d + 4*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt((b*c - a*d)/c)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 - 4*(a*c^2*x + (2*b*c^2 - a*c*d)*x^3)*sqrt(b*x^2 + a)*sqrt((b*c - a*d)/c)))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*(3*(2*b^2*c^2*d^2 - a*b*c*d^3 - a^2*d^4)*x^3 + (4*b^2*c^3*d + a*b*c^2*d^2 - 5*a^2*c*d^3)*x)*sqrt(b*x^2 + a))/(c^2*d^5*x^4 + 2*c^3*d^4*x^2 + c^4*d^3), 1/16*((8*b^2*c^4 + 4*a*b*c^3*d + 3*a^2*c^2*d^2 + (8*b^2*c^2*d^2 + 4*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(8*b^2*c^3*d + 4*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(-b*c - a*d)/c)*arctan(1/2*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)*sqrt(-b*c - a*d)/c)/((b^2*c - a*b*d)*x^3 + (a*b*c - a^2*d)*x)) + 8*(b^2*c^2*d^2*x^4 + 2*b^2*c^3*d*x^2 + b^2*c^4)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(3*(2*b^2*c^2*d^2 - a*b*c*d^3 - a^2*d^4)*x^3 + (4*b^2*c^3*d + a*b*c^2*d^2 - 5*a^2*c*d^3)*x)*sqrt(b*x^2 + a))/(c^2*d^5*x^4 + 2*c^3*d^4*x^2 + c^4*d^3), -1/16*(16*(b^2*c^2*d^2*x^4 + 2*b^2*c^3*d*x^2 + b^2*c^4)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (8*b^2*c^4 + 4*a*b*c^3*d + 3*a^2*c^2*d^2 + (8*b^2*c^2*d^2 + 4*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(8*b^2*c^3*d + 4*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(-b*c - a*d)/c)*arctan(1/2*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)*sqrt(-b*c - a*d)/c)/((b^2*c - a*b*d)*x^3 + (a*b*c - a^2*d)*x)) + 2*(3*(2*b^2*c^2*d^2 - a*b*c*d^3 - a^2*d^4)*x^3 + (4*b^2*c^3*d + a*b*c^2*d^2 - 5*a^2*c*d^3)*x)*sqrt(b*x^2 + a))/(c^2*d^5*x^4 + 2*c^3*d^4*x^2 + c^4*d^3)]

giac [B] time = 0.73, size = 659, normalized size = 3.40

$$\frac{b^{\frac{5}{2}} \log\left(\left(\sqrt{b}x - \sqrt{bx^2 + a}\right)^2\right)}{2d^3} + \frac{\left(8b^{\frac{7}{2}}c^3 - 4ab^{\frac{5}{2}}c^2d - a^2b^{\frac{3}{2}}cd^2 - 3a^3\sqrt{b}d^3\right) \arctan\left(\frac{\left(\sqrt{b}x - \sqrt{bx^2 + a}\right)^2_{d+2bc-ad}}{2\sqrt{-b^2c^2+abcd}}\right)}{8\sqrt{-b^2c^2+abcd}c^2d^3} - 16\left(\dots\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^3,x, algorithm="giac")

[Out] -1/2*b^(5/2)*log((sqrt(b)*x - sqrt(b*x^2 + a))^2)/d^3 + 1/8*(8*b^(7/2)*c^3 - 4*a*b^(5/2)*c^2*d - a^2*b^(3/2)*c*d^2 - 3*a^3*sqrt(b)*d^3)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/(sqrt(-b^2*c^2 + a*b*c*d)*c^2*d^3) - 1/4*(16*(sqrt(b)*x - sqrt(b*x^2 + a))^6*b^(7/2)*c^3*d - 20*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a*b^(5/2)*c^2*d^2 + (sqrt(b)*x - sqrt(b*x^2 + a))^6*a^2*b^(3/2)*c*d^3 + 3*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^3*d^3)

$$a)^6 a^3 \sqrt{b} d^4 + 48 (\sqrt{b} x - \sqrt{b x^2 + a})^4 b^{9/2} c^4 - 72 (\sqrt{b} x - \sqrt{b x^2 + a})^4 a^2 b^{5/2} c^2 d^2 + 15 (\sqrt{b} x - \sqrt{b x^2 + a})^4 a^3 b^{3/2} c d^3 - 9 (\sqrt{b} x - \sqrt{b x^2 + a})^4 a^4 \sqrt{b} d^4 + 32 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a^2 b^{7/2} c^3 d - 28 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a^3 b^{5/2} c^2 d^2 - 13 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a^4 b^{3/2} c d^3 + 9 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a^5 \sqrt{b} d^4 + 6 a^4 b^{5/2} c^2 d^2 - 3 a^5 b^{3/2} c d^3 - 3 a^6 \sqrt{b} d^4 / (((\sqrt{b} x - \sqrt{b x^2 + a})^4 d + 4 (\sqrt{b} x - \sqrt{b x^2 + a})^2 b c - 2 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a d + a^2 d)^2 c^2 d^3)$$

maple [B] time = 0.03, size = 14133, normalized size = 72.85

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/2)/(d*x^2+c)^3,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^3,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/2)/(d*x^2 + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{5/2}}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)/(c + d*x^2)^3,x)

[Out] int((a + b*x^2)^(5/2)/(c + d*x^2)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{5}{2}}}{(c + dx^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)/(d*x**2+c)**3,x)

[Out] Integral((a + b*x**2)**(5/2)/(c + d*x**2)**3, x)

$$3.69 \quad \int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^4} dx$$

Optimal. Leaf size=144

$$\frac{5a^3 \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{16c^{7/2}\sqrt{bc-ad}} + \frac{5a^2x\sqrt{a+bx^2}}{16c^3(c+dx^2)} + \frac{5ax(a+bx^2)^{3/2}}{24c^2(c+dx^2)^2} + \frac{x(a+bx^2)^{5/2}}{6c(c+dx^2)^3}$$

[Out] $1/6*x*(b*x^2+a)^{(5/2)}/c/(d*x^2+c)^3+5/24*a*x*(b*x^2+a)^{(3/2)}/c^2/(d*x^2+c)^2+5/16*a^3*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})/c^{(7/2)}/(-a*d+b*c)^{(1/2)}+5/16*a^2*x*(b*x^2+a)^{(1/2)}/c^3/(d*x^2+c)$

Rubi [A] time = 0.07, antiderivative size = 144, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {378, 377, 208}

$$\frac{5a^2x\sqrt{a+bx^2}}{16c^3(c+dx^2)} + \frac{5a^3 \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{16c^{7/2}\sqrt{bc-ad}} + \frac{5ax(a+bx^2)^{3/2}}{24c^2(c+dx^2)^2} + \frac{x(a+bx^2)^{5/2}}{6c(c+dx^2)^3}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(5/2)/(c + d*x^2)^4, x]

[Out] $(x*(a + b*x^2)^{(5/2)})/(6*c*(c + d*x^2)^3) + (5*a*x*(a + b*x^2)^{(3/2)})/(24*c^2*(c + d*x^2)^2) + (5*a^2*x*\operatorname{Sqrt}[a + b*x^2])/(16*c^3*(c + d*x^2)) + (5*a^3*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(16*c^{(7/2)}*\operatorname{Sqrt}[b*c - a*d])$

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] - Dist[(c*q)/(a*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+1)+1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^4} dx &= \frac{x(a+bx^2)^{5/2}}{6c(c+dx^2)^3} + \frac{(5a) \int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^3} dx}{6c} \\
&= \frac{x(a+bx^2)^{5/2}}{6c(c+dx^2)^3} + \frac{5ax(a+bx^2)^{3/2}}{24c^2(c+dx^2)^2} + \frac{(5a^2) \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^2} dx}{8c^2} \\
&= \frac{x(a+bx^2)^{5/2}}{6c(c+dx^2)^3} + \frac{5ax(a+bx^2)^{3/2}}{24c^2(c+dx^2)^2} + \frac{5a^2x\sqrt{a+bx^2}}{16c^3(c+dx^2)} + \frac{(5a^3) \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{16c^3} \\
&= \frac{x(a+bx^2)^{5/2}}{6c(c+dx^2)^3} + \frac{5ax(a+bx^2)^{3/2}}{24c^2(c+dx^2)^2} + \frac{5a^2x\sqrt{a+bx^2}}{16c^3(c+dx^2)} + \frac{(5a^3) \text{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{16c^3} \\
&= \frac{x(a+bx^2)^{5/2}}{6c(c+dx^2)^3} + \frac{5ax(a+bx^2)^{3/2}}{24c^2(c+dx^2)^2} + \frac{5a^2x\sqrt{a+bx^2}}{16c^3(c+dx^2)} + \frac{5a^3 \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{16c^{7/2}\sqrt{bc-ad}}
\end{aligned}$$

Mathematica [A] time = 0.81, size = 201, normalized size = 1.40

$$\frac{x\sqrt{a+bx^2} \left(\frac{\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} (a^2(33c^2+40cdx^2+15d^2x^4)+2abcx^2(13c+5dx^2)+8b^2c^2x^4)}{(c+dx^2)^2 \sqrt{\frac{dx^2}{c}+1}} + \frac{15a^2 \sin^{-1}\left(\frac{\sqrt{x^2\left(\frac{d}{c}-\frac{b}{a}\right)}}{\sqrt{\frac{dx^2}{c}+1}}\right)}{\sqrt{\frac{x^2(ad-bc)}{ac}}} \right)}{48c^4 \sqrt{\frac{bx^2}{a}+1}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(5/2)/(c + d*x^2)^4,x]

[Out] (x*sqrt[a + b*x^2]*((sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]*(8*b^2*c^2*x^4 + 2*a*b*c*x^2*(13*c + 5*d*x^2) + a^2*(33*c^2 + 40*c*d*x^2 + 15*d^2*x^4)))/((c + d*x^2)^2*sqrt[1 + (d*x^2)/c]) + (15*a^2*ArcSin[sqrt[(-(b/a) + d/c)*x^2]/sqrt[1 + (d*x^2)/c]])/sqrt[(-(b*c) + a*d)*x^2/(a*c)]))/(48*c^4*sqrt[1 + (b*x^2)/a])

fricas [B] time = 0.96, size = 706, normalized size = 4.90

$$\frac{15 \left(a^3 d^3 x^6 + 3 a^3 c d^2 x^4 + 3 a^3 c^2 d x^2 + a^3 c^3 \right) \sqrt{bc^2 - acd} \log \left(\frac{(8 b^2 c^2 - 8 abcd + a^2 d^2) x^4 + a^2 c^2 + 2(4 abc^2 - 3 a^2 cd) x^2 + 4((2 bc - ad) x^2 + a^2)}{d^2 x^4 + 2 cd x^2 + c^2} \right)}{192 (bc^8 - ac^7 d + (bc^5 d^2 - ac^6 d) x^2 + (bc^4 d^3 - ac^5 d) x^4 + (bc^3 d^4 - ac^4 d) x^6 + (bc^2 d^5 - ac^3 d) x^8 + bc^2 d^6 x^{10} + bc^2 d^7 x^{12} + bc^2 d^8 x^{14})}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^4,x, algorithm="fricas")

[Out] [1/192*(15*(a^3*d^3*x^6 + 3*a^3*c*d^2*x^4 + 3*a^3*c^2*d*x^2 + a^3*c^3)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*((8*b^3*c^4 + 2*a*b^2*c^3*d + 5*a^2*b*c^2*d^2 - 15*a^3*c*d^3)*x^5 + 2*(13*a*b^2*c^4 + 7*a^2*b*c^3

```
*d - 20*a^3*c^2*d^2)*x^3 + 33*(a^2*b*c^4 - a^3*c^3*d)*x)*sqrt(b*x^2 + a))/(
b*c^8 - a*c^7*d + (b*c^5*d^3 - a*c^4*d^4)*x^6 + 3*(b*c^6*d^2 - a*c^5*d^3)*x
^4 + 3*(b*c^7*d - a*c^6*d^2)*x^2), -1/96*(15*(a^3*d^3*x^6 + 3*a^3*c*d^2*x^4
+ 3*a^3*c^2*d*x^2 + a^3*c^3)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 +
a*c*d))*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3
+ (a*b*c^2 - a^2*c*d)*x)) - 2*((8*b^3*c^4 + 2*a*b^2*c^3*d + 5*a^2*b*c^2*d^2
- 15*a^3*c*d^3)*x^5 + 2*(13*a*b^2*c^4 + 7*a^2*b*c^3*d - 20*a^3*c^2*d^2)*x^
3 + 33*(a^2*b*c^4 - a^3*c^3*d)*x)*sqrt(b*x^2 + a))/(b*c^8 - a*c^7*d + (b*c^
5*d^3 - a*c^4*d^4)*x^6 + 3*(b*c^6*d^2 - a*c^5*d^3)*x^4 + 3*(b*c^7*d - a*c^6
*d^2)*x^2)]
```

giac [B] time = 2.86, size = 846, normalized size = 5.88

$$\frac{5a^3\sqrt{b}\arctan\left(\frac{(\sqrt{bx}-\sqrt{bx^2+a})^{d+2}bc-ad}{2\sqrt{-b^2c^2+abcd}}\right)}{16\sqrt{-b^2c^2+abcd}c^3} + \frac{48\left(\sqrt{b}x-\sqrt{bx^2+a}\right)^{10}b^{\frac{7}{2}}c^3d^2-15\left(\sqrt{b}x-\sqrt{bx^2+a}\right)^{10}a^3\sqrt{b}d^5+19}{16\sqrt{-b^2c^2+abcd}c^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^4,x, algorithm="giac")
```

```
[Out] -5/16*a^3*sqrt(b)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a
*d)/sqrt(-b^2*c^2 + a*b*c*d))/(sqrt(-b^2*c^2 + a*b*c*d)*c^3) + 1/24*(48*(sq
rt(b)*x - sqrt(b*x^2 + a))^10*b^(7/2)*c^3*d^2 - 15*(sqrt(b)*x - sqrt(b*x^2
+ a))^10*a^3*sqrt(b)*d^5 + 192*(sqrt(b)*x - sqrt(b*x^2 + a))^8*b^(9/2)*c^4*
d + 48*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a*b^(7/2)*c^3*d^2 - 150*(sqrt(b)*x -
sqrt(b*x^2 + a))^8*a^3*b^(3/2)*c*d^4 + 75*(sqrt(b)*x - sqrt(b*x^2 + a))^8*
a^4*sqrt(b)*d^5 + 256*(sqrt(b)*x - sqrt(b*x^2 + a))^6*b^(11/2)*c^5 - 64*(sq
rt(b)*x - sqrt(b*x^2 + a))^6*a*b^(9/2)*c^4*d + 288*(sqrt(b)*x - sqrt(b*x^2
+ a))^6*a^2*b^(7/2)*c^3*d^2 - 440*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^3*b^(5/
2)*c^2*d^3 + 440*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^4*b^(3/2)*c*d^4 - 150*(s
qrt(b)*x - sqrt(b*x^2 + a))^6*a^5*sqrt(b)*d^5 + 192*(sqrt(b)*x - sqrt(b*x^2
+ a))^4*a^2*b^(9/2)*c^4*d + 48*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^3*b^(7/2)
*c^3*d^2 + 360*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^4*b^(5/2)*c^2*d^3 - 420*(s
qrt(b)*x - sqrt(b*x^2 + a))^4*a^5*b^(3/2)*c*d^4 + 150*(sqrt(b)*x - sqrt(b*x
^2 + a))^4*a^6*sqrt(b)*d^5 + 48*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^4*b^(7/2)
*c^3*d^2 + 72*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^5*b^(5/2)*c^2*d^3 + 120*(sq
rt(b)*x - sqrt(b*x^2 + a))^2*a^6*b^(3/2)*c*d^4 - 75*(sqrt(b)*x - sqrt(b*x^2
+ a))^2*a^7*sqrt(b)*d^5 + 8*a^6*b^(5/2)*c^2*d^3 + 10*a^7*b^(3/2)*c*d^4 + 1
5*a^8*sqrt(b)*d^5)/(((sqrt(b)*x - sqrt(b*x^2 + a))^4*d + 4*(sqrt(b)*x - sq
rt(b*x^2 + a))^2*b*c - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*d + a^2*d)^3*c^3*
d^3)
```

maple [B] time = 0.04, size = 21220, normalized size = 147.36

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*x^2+a)^(5/2)/(d*x^2+c)^4,x)
```

```
[Out] result too large to display
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{(dx^2 + c)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^4,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/2)/(d*x^2 + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{5/2}}{(dx^2 + c)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)/(c + d*x^2)^4,x)

[Out] int((a + b*x^2)^(5/2)/(c + d*x^2)^4, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)/(d*x**2+c)**4,x)

[Out] Timed out

$$3.70 \quad \int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^5} dx$$

Optimal. Leaf size=249

$$\frac{5a^3(8bc-7ad) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{128c^{9/2}(bc-ad)^{3/2}} + \frac{5a^2x\sqrt{a+bx^2}(8bc-7ad)}{128c^4(c+dx^2)(bc-ad)} + \frac{5ax(a+bx^2)^{3/2}(8bc-7ad)}{192c^3(c+dx^2)^2(bc-ad)} + \frac{x(a+bx^2)^{5/2}(8bc-7ad)}{48c^2(c+dx^2)^3(bc-ad)}$$

[Out] $-1/8*d*x*(b*x^2+a)^{(7/2)}/c/(-a*d+b*c)/(d*x^2+c)^4+1/48*(-7*a*d+8*b*c)*x*(b*x^2+a)^{(5/2)}/c^2/(-a*d+b*c)/(d*x^2+c)^3+5/192*a*(-7*a*d+8*b*c)*x*(b*x^2+a)^{(3/2)}/c^3/(-a*d+b*c)/(d*x^2+c)^2+5/128*a^3*(-7*a*d+8*b*c)*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)/(b*x^2+a)^{(1/2)})/c^{(9/2)/(-a*d+b*c)^{(3/2)}+5/128*a^2*(-7*a*d+8*b*c)*x*(b*x^2+a)^{(1/2)}/c^4/(-a*d+b*c)/(d*x^2+c)$

Rubi [A] time = 0.14, antiderivative size = 249, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {382, 378, 377, 208}

$$\frac{5a^2x\sqrt{a+bx^2}(8bc-7ad)}{128c^4(c+dx^2)(bc-ad)} + \frac{5a^3(8bc-7ad) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{128c^{9/2}(bc-ad)^{3/2}} + \frac{x(a+bx^2)^{5/2}(8bc-7ad)}{48c^2(c+dx^2)^3(bc-ad)} + \frac{5ax(a+bx^2)^{3/2}(8bc-7ad)}{192c^3(c+dx^2)^2(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(5/2)/(c + d*x^2)^5, x]

[Out] $-(d*x*(a+b*x^2)^{(7/2)})/(8*c*(b*c-a*d)*(c+d*x^2)^4)+((8*b*c-7*a*d)*x*(a+b*x^2)^{(5/2)})/(48*c^2*(b*c-a*d)*(c+d*x^2)^3)+(5*a*(8*b*c-7*a*d)*x*(a+b*x^2)^{(3/2)})/(192*c^3*(b*c-a*d)*(c+d*x^2)^2)+(5*a^2*(8*b*c-7*a*d)*x*\operatorname{Sqrt}[a+b*x^2])/(128*c^4*(b*c-a*d)*(c+d*x^2))+5*a^3*(8*b*c-7*a*d)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c-a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a+b*x^2])]/(128*c^{(9/2)}*(b*c-a*d)^{(3/2)})$

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] - Dist[(c*q)/(a*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+1) + 1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)), I

nt[(a + b*x^n)^(p + 1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p + q + 2) + 1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{5/2}}{(c + dx^2)^5} dx &= -\frac{dx (a + bx^2)^{7/2}}{8c(bc - ad)(c + dx^2)^4} + \frac{(8bc - 7ad) \int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^4} dx}{8c(bc - ad)} \\ &= -\frac{dx (a + bx^2)^{7/2}}{8c(bc - ad)(c + dx^2)^4} + \frac{(8bc - 7ad)x (a + bx^2)^{5/2}}{48c^2(bc - ad)(c + dx^2)^3} + \frac{(5a(8bc - 7ad)) \int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^3} dx}{48c^2(bc - ad)} \\ &= -\frac{dx (a + bx^2)^{7/2}}{8c(bc - ad)(c + dx^2)^4} + \frac{(8bc - 7ad)x (a + bx^2)^{5/2}}{48c^2(bc - ad)(c + dx^2)^3} + \frac{5a(8bc - 7ad)x (a + bx^2)^{3/2}}{192c^3(bc - ad)(c + dx^2)^2} + \dots \\ &= -\frac{dx (a + bx^2)^{7/2}}{8c(bc - ad)(c + dx^2)^4} + \frac{(8bc - 7ad)x (a + bx^2)^{5/2}}{48c^2(bc - ad)(c + dx^2)^3} + \frac{5a(8bc - 7ad)x (a + bx^2)^{3/2}}{192c^3(bc - ad)(c + dx^2)^2} + \frac{5a^2}{12} \\ &= -\frac{dx (a + bx^2)^{7/2}}{8c(bc - ad)(c + dx^2)^4} + \frac{(8bc - 7ad)x (a + bx^2)^{5/2}}{48c^2(bc - ad)(c + dx^2)^3} + \frac{5a(8bc - 7ad)x (a + bx^2)^{3/2}}{192c^3(bc - ad)(c + dx^2)^2} + \frac{5a^2}{12} \\ &= -\frac{dx (a + bx^2)^{7/2}}{8c(bc - ad)(c + dx^2)^4} + \frac{(8bc - 7ad)x (a + bx^2)^{5/2}}{48c^2(bc - ad)(c + dx^2)^3} + \frac{5a(8bc - 7ad)x (a + bx^2)^{3/2}}{192c^3(bc - ad)(c + dx^2)^2} + \frac{5a^2}{12} \end{aligned}$$

Mathematica [A] time = 1.09, size = 306, normalized size = 1.23

$$x \left(\frac{15a^3(c+dx^2)^4(7ad-8bc) \tanh^{-1}\left(\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}\right)}{\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}} - c(-a^4d(279c^3 + 511c^2dx^2 + 385cd^2x^4 + 105d^3x^6) + a^3b(264c^4 - 21c^3d^3x^6)) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2)/(c + d*x^2)^5, x]

[Out] (x*(-(c*(16*b^4*c^3*x^6*(4*c + d*x^2) + 8*a*b^3*c^2*x^4*(34*c^2 + 13*c*d*x^2 + 3*d^2*x^4) + 2*a^2*b^2*c*x^2*(236*c^3 + 173*c^2*d*x^2 + 106*c*d^2*x^4 + 25*d^3*x^6) - a^4*d*(279*c^3 + 511*c^2*d*x^2 + 385*c*d^2*x^4 + 105*d^3*x^6) + a^3*b*(264*c^4 - 21*c^3*d*x^2 - 323*c^2*d^2*x^4 - 335*c*d^3*x^6 - 105*d^4*x^8))) + (15*a^3*(-8*b*c + 7*a*d)*(c + d*x^2)^4*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2)))]/(384*c^5*(-(b*c) + a*d)*Sqrt[a + b*x^2]*(c + d*x^2)^4)

fricas [B] time = 2.02, size = 1258, normalized size = 5.05

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^5,x, algorithm="fricas")

[Out] [1/1536*(15*(8*a^3*b*c^5 - 7*a^4*c^4*d + (8*a^3*b*c*d^4 - 7*a^4*d^5)*x^8 + 4*(8*a^3*b*c^2*d^3 - 7*a^4*c*d^4)*x^6 + 6*(8*a^3*b*c^3*d^2 - 7*a^4*c^2*d^3)*x^4 + 4*(8*a^3*b*c^4*d - 7*a^4*c^3*d^2)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*((16*b^4*c^5*d + 8*a*b^3*c^4*d^2 + 26*a^2*b^2*c^3*d^3 - 155*a^3*b*c^2*d^4 + 105*a^4*c*d^5)*x^7 + (64*b^4*c^6 + 24*a*b^3*c^5*d + 100*a^2*b^2*c^4*d^2 - 573*a^3*b*c^3*d^3 + 385*a^4*c^2*d^4)*x^5 + (208*a*b^3*c^6 + 50*a^2*b^2*c^5*d - 769*a^3*b*c^4*d^2 + 511*a^4*c^3*d^3)*x^3 + 3*(88*a^2*b^2*c^6 - 181*a^3*b*c^5*d + 93*a^4*c^4*d^2)*x)*sqrt(b*x^2 + a))/(b^2*c^11 - 2*a*b*c^10*d + a^2*c^9*d^2 + (b^2*c^7*d^4 - 2*a*b*c^6*d^5 + a^2*c^5*d^6)*x^8 + 4*(b^2*c^8*d^3 - 2*a*b*c^7*d^4 + a^2*c^6*d^5)*x^6 + 6*(b^2*c^9*d^2 - 2*a*b*c^8*d^3 + a^2*c^7*d^4)*x^4 + 4*(b^2*c^10*d - 2*a*b*c^9*d^2 + a^2*c^8*d^3)*x^2), -1/768*(15*(8*a^3*b*c^5 - 7*a^4*c^4*d + (8*a^3*b*c*d^4 - 7*a^4*d^5)*x^8 + 4*(8*a^3*b*c^2*d^3 - 7*a^4*c*d^4)*x^6 + 6*(8*a^3*b*c^3*d^2 - 7*a^4*c^2*d^3)*x^4 + 4*(8*a^3*b*c^4*d - 7*a^4*c^3*d^2)*x^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d))*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)) - 2*((16*b^4*c^5*d + 8*a*b^3*c^4*d^2 + 26*a^2*b^2*c^3*d^3 - 155*a^3*b*c^2*d^4 + 105*a^4*c*d^5)*x^7 + (64*b^4*c^6 + 24*a*b^3*c^5*d + 100*a^2*b^2*c^4*d^2 - 573*a^3*b*c^3*d^3 + 385*a^4*c^2*d^4)*x^5 + (208*a*b^3*c^6 + 50*a^2*b^2*c^5*d - 769*a^3*b*c^4*d^2 + 511*a^4*c^3*d^3)*x^3 + 3*(88*a^2*b^2*c^6 - 181*a^3*b*c^5*d + 93*a^4*c^4*d^2)*x)*sqrt(b*x^2 + a))/(b^2*c^11 - 2*a*b*c^10*d + a^2*c^9*d^2 + (b^2*c^7*d^4 - 2*a*b*c^6*d^5 + a^2*c^5*d^6)*x^8 + 4*(b^2*c^8*d^3 - 2*a*b*c^7*d^4 + a^2*c^6*d^5)*x^6 + 6*(b^2*c^9*d^2 - 2*a*b*c^8*d^3 + a^2*c^7*d^4)*x^4 + 4*(b^2*c^10*d - 2*a*b*c^9*d^2 + a^2*c^8*d^3)*x^2)]

giac [B] time = 9.31, size = 1448, normalized size = 5.82

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^5,x, algorithm="giac")

[Out] -5/128*(8*a^3*b^(3/2)*c - 7*a^4*sqrt(b)*d)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/((b*c^5 - a*c^4*d)*sqrt(-b^2*c^2 + a*b*c*d)) - 1/192*(120*(sqrt(b)*x - sqrt(b*x^2 + a))^14*a^3*b^(3/2)*c*d^6 - 105*(sqrt(b)*x - sqrt(b*x^2 + a))^14*a^4*sqrt(b)*d^7 - 768*(sqrt(b)*x - sqrt(b*x^2 + a))^12*b^(11/2)*c^5*d^2 + 768*(sqrt(b)*x - sqrt(b*x^2 + a))^12*a*b^(9/2)*c^4*d^3 + 1680*(sqrt(b)*x - sqrt(b*x^2 + a))^12*a^3*b^(5/2)*c^2*d^5 - 2310*(sqrt(b)*x - sqrt(b*x^2 + a))^12*a^4*b^(3/2)*c*d^6 + 735*(sqrt(b)*x - sqrt(b*x^2 + a))^12*a^5*sqrt(b)*d^7 - 2048*(sqrt(b)*x - sqrt(b*x^2 + a))^10*b^(13/2)*c^6*d + 2048*(sqrt(b)*x - sqrt(b*x^2 + a))^10*a^2*b^(9/2)*c^4*d^3 + 8320*(sqrt(b)*x - sqrt(b*x^2 + a))^10*a^3*b^(7/2)*c^3*d^4 - 15600*(sqrt(b)*x - sqrt(b*x^2 + a))^10*a^4*b^(5/2)*c^2*d^5 + 9800*(sqrt(b)*x - sqrt(b*x^2 + a))^10*a^5*b^(3/2)*c*d^6 - 2205*(sqrt(b)*x - sqrt(b*x^2 + a))^10*a^6*sqrt(b)*d^7 - 2048*(sqrt(b)*x - sqrt(b*x^2 + a))^8*b^(15/2)*c^7 + 1024*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a*b^(13/2)*c^6*d - 4864*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^2*b^(11/2)*c^5*d^2 + 21888*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^3*b^(9/2)*c^4*d^3 - 38000*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^4*b^(7/2)*c^3*d^4 + 37400*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^5*b^(5/2)*c^2*d^5 - 18550*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^6*b^(3/2)*c*d^6 + 3675*(sqrt(b)*x - sqrt(b*x^2 + a))^8*a^7*sqrt(b)*d^7 - 2048*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^2*b^(13/2)*c^6*d - 9472*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^4*b^(9/2)*c^4*d^3 + 32896*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^5*b^(7/2)*c^3*d^4 - 35376*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^6*b^(5/2)*c^2*d^5 + 18200*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^7*b^(3/2)*c*d^6 - 3675*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^8*sqrt(b)*d^7 - 768*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^4*b^(11/2)*c^5*d^2

- 1536*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^5*b^(9/2)*c^4*d^3 - 2944*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^6*b^(7/2)*c^3*d^4 + 12528*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^7*b^(5/2)*c^2*d^5 - 9170*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^8*b^(3/2)*c*d^6 + 2205*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^9*sqrt(b)*d^7 - 256*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^6*b^(9/2)*c^4*d^3 - 256*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^7*b^(7/2)*c^3*d^4 - 608*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^8*b^(5/2)*c^2*d^5 + 1960*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^9*b^(3/2)*c*d^6 - 735*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^10*sqrt(b)*d^7 - 16*a^8*b^(7/2)*c^3*d^4 - 24*a^9*b^(5/2)*c^2*d^5 - 50*a^10*b^(3/2)*c*d^6 + 105*a^11*sqrt(b)*d^7)/(b*c^5*d^3 - a*c^4*d^4)*((sqrt(b)*x - sqrt(b*x^2 + a))^4*d + 4*(sqrt(b)*x - sqrt(b*x^2 + a))^2*b*c - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*d + a^2*d^4)

maple [B] time = 0.05, size = 28625, normalized size = 114.96

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/2)/(d*x^2+c)^5,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{(dx^2 + c)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^5,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/2)/(d*x^2 + c)^5, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{5/2}}{(dx^2 + c)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)/(c + d*x^2)^5,x)

[Out] int((a + b*x^2)^(5/2)/(c + d*x^2)^5, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)/(d*x**2+c)**5,x)

[Out] Timed out

$$3.71 \quad \int \frac{\sqrt{1-x^2}}{1+x^2} dx$$

Optimal. Leaf size=30

$$\sqrt{2} \tan^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right) - \sin^{-1}(x)$$

[Out] `-arcsin(x)+arctan(x*2^(1/2)/(-x^2+1)^(1/2))*2^(1/2)`

Rubi [A] time = 0.01, antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {402, 216, 377, 203}

$$\sqrt{2} \tan^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right) - \sin^{-1}(x)$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[1 - x^2]/(1 + x^2), x]`

[Out] `-ArcSin[x] + Sqrt[2]*ArcTan[(Sqrt[2]*x)/Sqrt[1 - x^2]]`

Rule 203

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 216

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[(Rt[-b, 2]*x)/Sqrt[a]]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Rule 377

`Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]`

Rule 402

`Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Dist[b/d, Int[(a + b*x^2)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4])`

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{1-x^2}}{1+x^2} dx &= 2 \int \frac{1}{\sqrt{1-x^2}(1+x^2)} dx - \int \frac{1}{\sqrt{1-x^2}} dx \\ &= -\sin^{-1}(x) + 2 \operatorname{Subst}\left(\int \frac{1}{1+2x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right) \\ &= -\sin^{-1}(x) + \sqrt{2} \tan^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right) \end{aligned}$$

Mathematica [A] time = 0.03, size = 30, normalized size = 1.00

$$\sqrt{2} \tan^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right) - \sin^{-1}(x)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - x^2]/(1 + x^2), x]

[Out] -ArcSin[x] + Sqrt[2]*ArcTan[(Sqrt[2]*x)/Sqrt[1 - x^2]]

fricas [A] time = 0.64, size = 42, normalized size = 1.40

$$-\sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{-x^2+1}}{2x}\right) + 2 \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(x^2+1), x, algorithm="fricas")

[Out] -sqrt(2)*arctan(1/2*sqrt(2)*sqrt(-x^2 + 1)/x) + 2*arctan((sqrt(-x^2 + 1) - 1)/x)

giac [B] time = 0.61, size = 95, normalized size = 3.17

$$-\frac{1}{2} \pi \operatorname{sgn}(x) + \frac{1}{2} \sqrt{2} \left(\pi \operatorname{sgn}(x) + 2 \arctan\left(-\frac{\sqrt{2}x\left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2}-1\right)}{4(\sqrt{-x^2+1}-1)}\right) \right) - \arctan\left(-\frac{x\left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2}-1\right)}{2(\sqrt{-x^2+1}-1)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(x^2+1), x, algorithm="giac")

[Out] -1/2*pi*sgn(x) + 1/2*sqrt(2)*(pi*sgn(x) + 2*arctan(-1/4*sqrt(2)*x*((sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1))) - arctan(-1/2*x*((sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1))

maple [A] time = 0.02, size = 33, normalized size = 1.10

$$-\arcsin(x) - \sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{-x^2+1}x}{x^2-1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x^2+1)^(1/2)/(x^2+1), x)

[Out] -arcsin(x)-2^(1/2)*arctan(2^(1/2)*(-x^2+1)^(1/2)/(x^2-1)*x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+1}}{x^2+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(x^2+1), x, algorithm="maxima")

[Out] integrate(sqrt(-x^2 + 1)/(x^2 + 1), x)

mupad [B] time = 0.39, size = 83, normalized size = 2.77

$$-\operatorname{asin}(x) + \frac{\sqrt{2} \ln\left(\frac{\frac{\sqrt{2}(-1+xi)1i}{2} - \sqrt{1-x^2} 1i}{x-i}\right) 1i}{2} - \frac{\sqrt{2} \ln\left(\frac{\frac{\sqrt{2}(1+xi)1i}{2} + \sqrt{1-x^2} 1i}{x+1i}\right) 1i}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1 - x^2)^(1/2)/(x^2 + 1),x)`

[Out] $(2^{1/2} \log(((2^{1/2} * (x * 1i - 1) * 1i) / 2 - (1 - x^2)^{1/2} * 1i) / (x - 1i)) * 1i) / 2 - \operatorname{asin}(x) - (2^{1/2} \log(((2^{1/2} * (x * 1i + 1) * 1i) / 2 + (1 - x^2)^{1/2} * 1i) / (x + 1i)) * 1i) / 2$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(x-1)(x+1)}}{x^2+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-x**2+1)**(1/2)/(x**2+1),x)`

[Out] `Integral(sqrt(-(x - 1)*(x + 1))/(x**2 + 1), x)`

$$3.72 \quad \int \frac{\sqrt{1+x^2}}{-1+x^2} dx$$

Optimal. Leaf size=27

$$\sinh^{-1}(x) - \sqrt{2} \tanh^{-1}\left(\frac{\sqrt{2}x}{\sqrt{x^2+1}}\right)$$

[Out] arcsinh(x)-arctanh(x*2^(1/2)/(x^2+1)^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {402, 215, 377, 207}

$$\sinh^{-1}(x) - \sqrt{2} \tanh^{-1}\left(\frac{\sqrt{2}x}{\sqrt{x^2+1}}\right)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 + x^2]/(-1 + x^2), x]

[Out] ArcSinh[x] - Sqrt[2]*ArcTanh[(Sqrt[2]*x)/Sqrt[1 + x^2]]

Rule 207

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTanh[(Rt[b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 215

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[(Rt[b, 2]*x)/Sqrt[a]]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 402

Int[((a_) + (b_.)*(x_)^2)^(p_)/((c_) + (d_.)*(x_)^2), x_Symbol] := Dist[b/d, Int[(a + b*x^2)^(p-1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^2)^(p-1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4])

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{1+x^2}}{-1+x^2} dx &= 2 \int \frac{1}{(-1+x^2)\sqrt{1+x^2}} dx + \int \frac{1}{\sqrt{1+x^2}} dx \\ &= \sinh^{-1}(x) + 2 \operatorname{Subst}\left(\int \frac{1}{-1+2x^2} dx, x, \frac{x}{\sqrt{1+x^2}}\right) \\ &= \sinh^{-1}(x) - \sqrt{2} \tanh^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1+x^2}}\right) \end{aligned}$$

Mathematica [B] time = 0.03, size = 64, normalized size = 2.37

$$\frac{\log\left(\sqrt{2}\sqrt{x^2+1}-x+1\right)-\log\left(\sqrt{2}\sqrt{x^2+1}+x+1\right)+\log(1-x)-\log(x+1)}{\sqrt{2}}+\sinh^{-1}(x)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 + x^2]/(-1 + x^2), x]

[Out] ArcSinh[x] + (Log[1 - x] - Log[1 + x] + Log[1 - x + Sqrt[2]*Sqrt[1 + x^2]] - Log[1 + x + Sqrt[2]*Sqrt[1 + x^2]])/Sqrt[2]

fricas [B] time = 0.53, size = 67, normalized size = 2.48

$$\frac{1}{2}\sqrt{2}\log\left(\frac{9x^2-2\sqrt{2}(3x^2+1)-2\sqrt{x^2+1}(3\sqrt{2}x-4x)+3}{x^2-1}\right)-\log(-x+\sqrt{x^2+1})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(x^2-1), x, algorithm="fricas")

[Out] 1/2*sqrt(2)*log((9*x^2 - 2*sqrt(2)*(3*x^2 + 1) - 2*sqrt(x^2 + 1)*(3*sqrt(2)*x - 4*x) + 3)/(x^2 - 1)) - log(-x + sqrt(x^2 + 1))

giac [B] time = 0.62, size = 70, normalized size = 2.59

$$-\frac{1}{2}\sqrt{2}\log\left(\frac{\left|2\left(x-\sqrt{x^2+1}\right)^2-4\sqrt{2}-6\right|}{\left|2\left(x-\sqrt{x^2+1}\right)^2+4\sqrt{2}-6\right|}\right)-\log(-x+\sqrt{x^2+1})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(x^2-1), x, algorithm="giac")

[Out] -1/2*sqrt(2)*log(abs(2*(x - sqrt(x^2 + 1))^2 - 4*sqrt(2) - 6)/abs(2*(x - sqrt(x^2 + 1))^2 + 4*sqrt(2) - 6)) - log(-x + sqrt(x^2 + 1))

maple [B] time = 0.01, size = 84, normalized size = 3.11

$$\operatorname{arcsinh}(x)+\frac{\sqrt{2}\operatorname{arctanh}\left(\frac{(-2x+2)\sqrt{2}}{4\sqrt{-2x+(x+1)^2}}\right)}{2}-\frac{\sqrt{2}\operatorname{arctanh}\left(\frac{(2x+2)\sqrt{2}}{4\sqrt{2x+(x-1)^2}}\right)}{2}+\frac{\sqrt{-2x+(x+1)^2}}{2}+\frac{\sqrt{2x+(x-1)^2}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2+1)^(1/2)/(x^2-1), x)

[Out] -1/2*((x+1)^2-2*x)^(1/2)+arcsinh(x)+1/2*2^(1/2)*arctanh(1/4*(2-2*x)*2^(1/2)/((x+1)^2-2*x)^(1/2))+1/2*((x-1)^2+2*x)^(1/2)-1/2*2^(1/2)*arctanh(1/4*(2*x+2)*2^(1/2)/((x-1)^2+2*x)^(1/2))

maxima [B] time = 2.98, size = 59, normalized size = 2.19

$$-\frac{1}{2}\sqrt{2}\operatorname{arsinh}\left(\frac{2x}{|2x+2|}-\frac{2}{|2x+2|}\right)-\frac{1}{2}\sqrt{2}\operatorname{arsinh}\left(\frac{2x}{|2x-2|}+\frac{2}{|2x-2|}\right)+\operatorname{arsinh}(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(x^2-1), x, algorithm="maxima")

[Out] $-1/2*\sqrt{2}*\operatorname{arcsinh}(2*x/\operatorname{abs}(2*x + 2) - 2/\operatorname{abs}(2*x + 2)) - 1/2*\sqrt{2}*\operatorname{arcsinh}(2*x/\operatorname{abs}(2*x - 2) + 2/\operatorname{abs}(2*x - 2)) + \operatorname{arcsinh}(x)$

mupad [B] time = 0.17, size = 59, normalized size = 2.19

$$\operatorname{asinh}(x) + \frac{\sqrt{2} \left(\ln(x-1) - \ln\left(x + \sqrt{2} \sqrt{x^2+1} + 1\right) \right)}{2} - \frac{\sqrt{2} \left(\ln(x+1) - \ln\left(\sqrt{2} \sqrt{x^2+1} - x + 1\right) \right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2 + 1)^(1/2)/(x^2 - 1), x)`

[Out] $\operatorname{asinh}(x) + (2^{1/2}*(\log(x - 1) - \log(x + 2^{1/2}*(x^2 + 1)^{1/2} + 1)))/2 - (2^{1/2}*(\log(x + 1) - \log(2^{1/2}*(x^2 + 1)^{1/2} - x + 1)))/2$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2+1}}{(x-1)(x+1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((x**2+1)**(1/2)/(x**2-1), x)`

[Out] `Integral(sqrt(x**2 + 1)/((x - 1)*(x + 1)), x)`

$$3.73 \quad \int \frac{\sqrt{1-x^2}}{-1+2x^2} dx$$

Optimal. Leaf size=25

$$-\frac{1}{2} \tanh^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right) - \frac{1}{2} \sin^{-1}(x)$$

[Out] -1/2*arcsin(x)-1/2*arctanh(x/(-x^2+1)^(1/2))

Rubi [A] time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {402, 216, 377, 207}

$$-\frac{1}{2} \tanh^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right) - \frac{1}{2} \sin^{-1}(x)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - x^2]/(-1 + 2*x^2), x]

[Out] -ArcSin[x]/2 - ArcTanh[x/Sqrt[1 - x^2]]/2

Rule 207

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> -Simp[ArcTanh[(Rt[b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 216

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[(Rt[-b, 2]*x)/Sqrt[a]]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 402

Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] :> Dist[b/d, Int[(a + b*x^2)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4])

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{1-x^2}}{-1+2x^2} dx &= -\left(\frac{1}{2} \int \frac{1}{\sqrt{1-x^2}} dx\right) + \frac{1}{2} \int \frac{1}{\sqrt{1-x^2}(-1+2x^2)} dx \\ &= -\frac{1}{2} \sin^{-1}(x) + \frac{1}{2} \text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right) \\ &= -\frac{1}{2} \sin^{-1}(x) - \frac{1}{2} \tanh^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right) \end{aligned}$$

Mathematica [A] time = 0.01, size = 25, normalized size = 1.00

$$-\frac{1}{2} \tanh^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right) - \frac{1}{2} \sin^{-1}(x)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - x^2]/(-1 + 2*x^2), x]

[Out] -1/2*ArcSin[x] - ArcTanh[x/Sqrt[1 - x^2]]/2

fricas [B] time = 0.56, size = 74, normalized size = 2.96

$$\arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right) + \frac{1}{4} \log\left(-\frac{x^2 + \sqrt{-x^2+1}(x+1) - x - 1}{x^2}\right) - \frac{1}{4} \log\left(-\frac{x^2 - \sqrt{-x^2+1}(x-1) + x - 1}{x^2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(2*x^2-1), x, algorithm="fricas")

[Out] arctan((sqrt(-x^2 + 1) - 1)/x) + 1/4*log(-(x^2 + sqrt(-x^2 + 1)*(x + 1) - x - 1)/x^2) - 1/4*log(-(x^2 - sqrt(-x^2 + 1)*(x - 1) + x - 1)/x^2)

giac [B] time = 0.61, size = 118, normalized size = 4.72

$$-\frac{1}{4} \pi \operatorname{sgn}(x) - \frac{1}{2} \arctan\left(\frac{x \left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2} - 1\right)}{2(\sqrt{-x^2+1}-1)}\right) - \frac{1}{4} \log\left(\left|-\frac{x}{\sqrt{-x^2+1}-1} + \frac{\sqrt{-x^2+1}-1}{x} + 2\right|\right) + \frac{1}{4} \log\left(\left|-\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x} + 2\right|\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(2*x^2-1), x, algorithm="giac")

[Out] -1/4*pi*sgn(x) - 1/2*arctan(-1/2*x*((sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1)) - 1/4*log(abs(-x/(sqrt(-x^2 + 1) - 1) + (sqrt(-x^2 + 1) - 1)/x + 2)) + 1/4*log(abs(-x/(sqrt(-x^2 + 1) - 1) + (sqrt(-x^2 + 1) - 1)/x - 2))

maple [B] time = 0.04, size = 187, normalized size = 7.48

$$\frac{\sqrt{2} \left(\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\left(\left(x + \frac{\sqrt{2}}{2}\right)\sqrt{2} + 1\right)\sqrt{2}}{\sqrt{-4\left(x + \frac{\sqrt{2}}{2}\right)^2 + 4\left(x + \frac{\sqrt{2}}{2}\right)\sqrt{2} + 2}}\right)}{4} + \frac{\sqrt{2} \arcsin(x)}{4} + \frac{\sqrt{-4\left(x + \frac{\sqrt{2}}{2}\right)^2 + 4\left(x + \frac{\sqrt{2}}{2}\right)\sqrt{2} + 2}}{4} \right)}{2} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\left(\left(x - \frac{\sqrt{2}}{2}\right)\sqrt{2} + 1\right)\sqrt{2}}{\sqrt{-4\left(x - \frac{\sqrt{2}}{2}\right)^2 + 4\left(x - \frac{\sqrt{2}}{2}\right)\sqrt{2} + 2}}\right)}{4} + \frac{\sqrt{2} \arcsin(x)}{4} + \frac{\sqrt{-4\left(x - \frac{\sqrt{2}}{2}\right)^2 + 4\left(x - \frac{\sqrt{2}}{2}\right)\sqrt{2} + 2}}{4} \right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x^2+1)^(1/2)/(2*x^2-1), x)

[Out] -1/2*2^(1/2)*(1/4*(-4*(x+1/2*2^(1/2))^2+4*(x+1/2*2^(1/2))*2^(1/2)+2)^(1/2)+1/4*2^(1/2)*arcsin(x)-1/4*2^(1/2)*arctanh(((x+1/2*2^(1/2))*2^(1/2)+1)*2^(1/2)/(-4*(x+1/2*2^(1/2))^2+4*(x+1/2*2^(1/2))*2^(1/2)+2)^(1/2)))+1/2*2^(1/2)*(1/4*(-4*(x-1/2*2^(1/2))^2-4*(x-1/2*2^(1/2))*2^(1/2)+2)^(1/2)-1/4*2^(1/2)*arcsin(x)-1/4*2^(1/2)*arctanh((-x-1/2*2^(1/2))*2^(1/2)+1)*2^(1/2)/(-4*(x-1/2*2^(1/2))^2-4*(x-1/2*2^(1/2))*2^(1/2)+2)^(1/2))

maxima [B] time = 3.07, size = 110, normalized size = 4.40

$$-\frac{1}{8}\sqrt{2}\left(2\sqrt{2}\arcsin(x)-\sqrt{2}\log\left(\frac{1}{4}\sqrt{2}+\frac{\sqrt{2}\sqrt{-x^2+1}}{|4x+2\sqrt{2}|}+\frac{1}{|4x+2\sqrt{2}|}\right)+\sqrt{2}\log\left(-\frac{1}{4}\sqrt{2}+\frac{\sqrt{2}\sqrt{-x^2+1}}{|4x-2\sqrt{2}|}\right)+\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(2*x^2-1),x, algorithm="maxima")

[Out] -1/8*sqrt(2)*(2*sqrt(2)*arcsin(x) - sqrt(2)*log(1/4*sqrt(2) + sqrt(2)*sqrt(-x^2 + 1)/abs(4*x + 2*sqrt(2)) + 1/abs(4*x + 2*sqrt(2))) + sqrt(2)*log(-1/4*sqrt(2) + sqrt(2)*sqrt(-x^2 + 1)/abs(4*x - 2*sqrt(2)) + 1/abs(4*x - 2*sqrt(2)))

mupad [B] time = 5.35, size = 85, normalized size = 3.40

$$-\frac{\ln\left(\frac{\sqrt{2}\left(\frac{\sqrt{2}x-1}{2}\right)^{1i-\sqrt{1-x^2}}1i}{x-\frac{\sqrt{2}}{2}}\right)}{4}+\frac{\ln\left(\frac{\sqrt{2}\left(\frac{\sqrt{2}x+1}{2}\right)^{1i+\sqrt{1-x^2}}1i}{x+\frac{\sqrt{2}}{2}}\right)}{4}-\frac{\operatorname{asin}(x)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - x^2)^(1/2)/(2*x^2 - 1),x)

[Out] log((2^(1/2)*((2^(1/2)*x)/2 + 1)*1i + (1 - x^2)^(1/2)*1i)/(x + 2^(1/2)/2))/4 - log((2^(1/2)*((2^(1/2)*x)/2 - 1)*1i - (1 - x^2)^(1/2)*1i)/(x - 2^(1/2)/2))/4 - asin(x)/2

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(x-1)(x+1)}}{2x^2-1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x**2+1)**(1/2)/(2*x**2-1),x)

[Out] Integral(sqrt(-(x - 1)*(x + 1))/(2*x**2 - 1), x)

$$3.74 \quad \int \frac{(c+dx^2)^3}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=169

$$\frac{(2bc - ad)(5a^2d^2 - 8abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16b^{7/2}} + \frac{dx\sqrt{a+bx^2}(15a^2d^2 - 44abcd + 44b^2c^2)}{48b^3} + \frac{5dx\sqrt{a+bx^2}}{16b^{7/2}}$$

[Out] $1/16*(-a*d+2*b*c)*(5*a^2*d^2-8*a*b*c*d+8*b^2*c^2)*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(7/2)}+1/48*d*(15*a^2*d^2-44*a*b*c*d+44*b^2*c^2)*x*(b*x^2+a)^{(1/2)}/b^3+5/24*d*(-a*d+2*b*c)*x*(d*x^2+c)*(b*x^2+a)^{(1/2)}/b^2+1/6*d*x*(d*x^2+c)^2*(b*x^2+a)^{(1/2)}/b$

Rubi [A] time = 0.14, antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {416, 528, 388, 217, 206}

$$\frac{dx\sqrt{a+bx^2}(15a^2d^2 - 44abcd + 44b^2c^2)}{48b^3} + \frac{(2bc - ad)(5a^2d^2 - 8abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{16b^{7/2}} + \frac{5dx\sqrt{a+bx^2}}{16b^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^3/Sqrt[a + b*x^2], x]

[Out] $(d*(44*b^2*c^2 - 44*a*b*c*d + 15*a^2*d^2)*x*\operatorname{Sqrt}[a + b*x^2])/(48*b^3) + (5*d*(2*b*c - a*d)*x*\operatorname{Sqrt}[a + b*x^2]*(c + d*x^2))/(24*b^2) + (d*x*\operatorname{Sqrt}[a + b*x^2]*(c + d*x^2)^2)/(6*b) + ((2*b*c - a*d)*(8*b^2*c^2 - 8*a*b*c*d + 5*a^2*d^2)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(16*b^{(7/2)})$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1)/(b*(n*(p+q)+1)), x] + Dist[1/(b*(n*(p+q)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q)+1) - a*d] + d*(b*c*(n*(p+2*q-1)+1) - a*d*(n*(q-1)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q)+1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^3}{\sqrt{a + bx^2}} dx &= \frac{dx\sqrt{a + bx^2} (c + dx^2)^2}{6b} + \frac{\int \frac{(c+dx^2)(c(6bc-ad)+5d(2bc-ad)x^2)}{\sqrt{a+bx^2}} dx}{6b} \\ &= \frac{5d(2bc - ad)x\sqrt{a + bx^2} (c + dx^2)}{24b^2} + \frac{dx\sqrt{a + bx^2} (c + dx^2)^2}{6b} + \frac{\int \frac{c(24b^2c^2 - 14abcd + 5a^2d^2) + d(44b^2c^2 - 44abcd + 15a^2d^2)}{\sqrt{a+bx^2}} dx}{24b^2} \\ &= \frac{d(44b^2c^2 - 44abcd + 15a^2d^2)x\sqrt{a + bx^2}}{48b^3} + \frac{5d(2bc - ad)x\sqrt{a + bx^2} (c + dx^2)}{24b^2} + \frac{dx\sqrt{a + bx^2}}{6b} \\ &= \frac{d(44b^2c^2 - 44abcd + 15a^2d^2)x\sqrt{a + bx^2}}{48b^3} + \frac{5d(2bc - ad)x\sqrt{a + bx^2} (c + dx^2)}{24b^2} + \frac{dx\sqrt{a + bx^2}}{6b} \\ &= \frac{d(44b^2c^2 - 44abcd + 15a^2d^2)x\sqrt{a + bx^2}}{48b^3} + \frac{5d(2bc - ad)x\sqrt{a + bx^2} (c + dx^2)}{24b^2} + \frac{dx\sqrt{a + bx^2}}{6b} \end{aligned}$$

Mathematica [A] time = 5.09, size = 140, normalized size = 0.83

$$\frac{\sqrt{b} dx\sqrt{a + bx^2} (15a^2d^2 - 2abd(27c + 5dx^2) + 4b^2(18c^2 + 9cdx^2 + 2d^2x^4)) + 3(-5a^3d^3 + 18a^2bcd^2 - 24ab^2c^2d)}{48b^{7/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(c + d*x^2)^3/Sqrt[a + b*x^2], x]
```

```
[Out] (Sqrt[b]*d*x*Sqrt[a + b*x^2]*(15*a^2*d^2 - 2*a*b*d*(27*c + 5*d*x^2) + 4*b^2
*(18*c^2 + 9*c*d*x^2 + 2*d^2*x^4)) + 3*(16*b^3*c^3 - 24*a*b^2*c^2*d + 18*a^
2*b*c*d^2 - 5*a^3*d^3)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(48*b^(7/2))
```

fricas [A] time = 0.55, size = 300, normalized size = 1.78

$$\left[\frac{3(16b^3c^3 - 24ab^2c^2d + 18a^2bcd^2 - 5a^3d^3)\sqrt{b} \log(-2bx^2 + 2\sqrt{bx^2 + a}\sqrt{b}x - a) - 2(8b^3d^3x^5 + 2(18b^3cd^2 - 5a^2b^2cd^2 + 5a^2b^2cd^2 - 5a^3d^3)*x)\sqrt{bx^2 + a}}{96b^4} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^3/(b*x^2+a)^(1/2), x, algorithm="fricas")
```

```
[Out] [-1/96*(3*(16*b^3*c^3 - 24*a*b^2*c^2*d + 18*a^2*b*c*d^2 - 5*a^3*d^3)*sqrt(b)
)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(8*b^3*d^3*x^5 + 2*(1
8*b^3*c*d^2 - 5*a*b^2*d^3)*x^3 + 3*(24*b^3*c^2*d - 18*a*b^2*c*d^2 + 5*a^2*b
*d^3)*x)*sqrt(b*x^2 + a))/b^4, -1/48*(3*(16*b^3*c^3 - 24*a*b^2*c^2*d + 18*a
^2*b*c*d^2 - 5*a^3*d^3)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (8*b^
3*d^3*x^5 + 2*(18*b^3*c*d^2 - 5*a*b^2*d^3)*x^3 + 3*(24*b^3*c^2*d - 18*a*b^2
*c*d^2 + 5*a^2*b*d^3)*x)*sqrt(b*x^2 + a))/b^4]
```

giac [A] time = 0.64, size = 150, normalized size = 0.89

$$\frac{1}{48} \left(2 \left(\frac{4d^3x^2}{b} + \frac{18b^4cd^2 - 5ab^3d^3}{b^5} \right) x^2 + \frac{3(24b^4c^2d - 18ab^3cd^2 + 5a^2b^2d^3)}{b^5} \right) \sqrt{bx^2 + a} x - \frac{(16b^3c^3 - 24ab^2c^2)}{b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^(1/2),x, algorithm="giac")

[Out] 1/48*(2*(4*d^3*x^2/b + (18*b^4*c*d^2 - 5*a*b^3*d^3)/b^5)*x^2 + 3*(24*b^4*c^2*d - 18*a*b^3*c*d^2 + 5*a^2*b^2*d^3)/b^5)*sqrt(b*x^2 + a)*x - 1/16*(16*b^3*c^3 - 24*a*b^2*c^2*d + 18*a^2*b*c*d^2 - 5*a^3*d^3)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(7/2)

maple [A] time = 0.01, size = 228, normalized size = 1.35

$$\frac{\sqrt{bx^2 + a} d^3 x^5}{6b} - \frac{5\sqrt{bx^2 + a} a d^3 x^3}{24b^2} + \frac{3\sqrt{bx^2 + a} c d^2 x^3}{4b} - \frac{5a^3 d^3 \ln(\sqrt{b} x + \sqrt{bx^2 + a})}{16b^{7/2}} + \frac{9a^2 c d^2 \ln(\sqrt{b} x + \sqrt{bx^2 + a})}{8b^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^3/(b*x^2+a)^(1/2),x)

[Out] 1/6*d^3*x^5/b*(b*x^2+a)^(1/2)-5/24*d^3*a/b^2*x^3*(b*x^2+a)^(1/2)+5/16*d^3*a^2/b^3*x*(b*x^2+a)^(1/2)-5/16*d^3*a^3/b^(7/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+3/4*c*d^2*x^3/b*(b*x^2+a)^(1/2)-9/8*c*d^2*a/b^2*x*(b*x^2+a)^(1/2)+9/8*c*d^2*a^2/b^(5/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+3/2*c^2*d*x/b*(b*x^2+a)^(1/2)-3/2*c^2*d*a/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+c^3*ln(b^(1/2)*x+(b*x^2+a)^(1/2))/b^(1/2)

maxima [A] time = 1.36, size = 199, normalized size = 1.18

$$\frac{\sqrt{bx^2 + a} d^3 x^5}{6b} + \frac{3\sqrt{bx^2 + a} c d^2 x^3}{4b} - \frac{5\sqrt{bx^2 + a} a d^3 x^3}{24b^2} + \frac{3\sqrt{bx^2 + a} c^2 dx}{2b} - \frac{9\sqrt{bx^2 + a} a c d^2 x}{8b^2} + \frac{5\sqrt{bx^2 + a} a^2 d^3 x}{16b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] 1/6*sqrt(b*x^2 + a)*d^3*x^5/b + 3/4*sqrt(b*x^2 + a)*c*d^2*x^3/b - 5/24*sqrt(b*x^2 + a)*a*d^3*x^3/b^2 + 3/2*sqrt(b*x^2 + a)*c^2*d*x/b - 9/8*sqrt(b*x^2 + a)*a*c*d^2*x/b^2 + 5/16*sqrt(b*x^2 + a)*a^2*d^3*x/b^3 + c^3*arcsinh(b*x/sqrt(a*b))/sqrt(b) - 3/2*a*c^2*d*arcsinh(b*x/sqrt(a*b))/b^(3/2) + 9/8*a^2*c*d^2*arcsinh(b*x/sqrt(a*b))/b^(5/2) - 5/16*a^3*d^3*arcsinh(b*x/sqrt(a*b))/b^(7/2)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(dx^2 + c)^3}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^3/(a + b*x^2)^(1/2),x)

[Out] int((c + d*x^2)^3/(a + b*x^2)^(1/2), x)

sympy [A] time = 13.27, size = 400, normalized size = 2.37

$$\frac{5a^{\frac{5}{2}}d^3x}{16b^3\sqrt{1+\frac{bx^2}{a}}} - \frac{9a^{\frac{3}{2}}cd^2x}{8b^2\sqrt{1+\frac{bx^2}{a}}} + \frac{5a^{\frac{3}{2}}d^3x^3}{48b^2\sqrt{1+\frac{bx^2}{a}}} + \frac{3\sqrt{a}c^2dx\sqrt{1+\frac{bx^2}{a}}}{2b} - \frac{3\sqrt{a}cd^2x^3}{8b\sqrt{1+\frac{bx^2}{a}}} - \frac{\sqrt{a}d^3x^5}{24b\sqrt{1+\frac{bx^2}{a}}} - \frac{5a^3d^3\operatorname{asinh}\left(\frac{\sqrt{bx^2+a}}{\sqrt{a}}\right)}{16b^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**3/(b*x**2+a)**(1/2),x)

[Out] 5*a**(5/2)*d**3*x/(16*b**3*sqrt(1 + b*x**2/a)) - 9*a**(3/2)*c*d**2*x/(8*b**2*sqrt(1 + b*x**2/a)) + 5*a**(3/2)*d**3*x**3/(48*b**2*sqrt(1 + b*x**2/a)) + 3*sqrt(a)*c**2*d*x*sqrt(1 + b*x**2/a)/(2*b) - 3*sqrt(a)*c*d**2*x**3/(8*b*sqrt(1 + b*x**2/a)) - sqrt(a)*d**3*x**5/(24*b*sqrt(1 + b*x**2/a)) - 5*a**3*d**3*asinh(sqrt(b)*x/sqrt(a))/(16*b**(7/2)) + 9*a**2*c*d**2*asinh(sqrt(b)*x/sqrt(a))/(8*b**(5/2)) - 3*a*c**2*d*asinh(sqrt(b)*x/sqrt(a))/(2*b**(3/2)) + c**3*Piecewise((sqrt(-a/b)*asin(x*sqrt(-b/a))/sqrt(a), (a > 0) & (b < 0)), (sqrt(a/b)*asinh(x*sqrt(b/a))/sqrt(a), (a > 0) & (b > 0)), (sqrt(-a/b)*acosh(x*sqrt(-b/a))/sqrt(-a), (b > 0) & (a < 0))) + 3*c*d**2*x**5/(4*sqrt(a)*sqrt(1 + b*x**2/a)) + d**3*x**7/(6*sqrt(a)*sqrt(1 + b*x**2/a))

$$3.75 \quad \int \frac{(c+dx^2)^2}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=108

$$\frac{(3a^2d^2 - 8abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{8b^{5/2}} + \frac{3dx\sqrt{a+bx^2}(2bc - ad)}{8b^2} + \frac{dx\sqrt{a+bx^2}(c + dx^2)}{4b}$$

[Out] 1/8*(3*a^2*d^2-8*a*b*c*d+8*b^2*c^2)*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(5/2)+3/8*d*(-a*d+2*b*c)*x*(b*x^2+a)^(1/2)/b^2+1/4*d*x*(d*x^2+c)*(b*x^2+a)^(1/2)/b

Rubi [A] time = 0.06, antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {416, 388, 217, 206}

$$\frac{(3a^2d^2 - 8abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{8b^{5/2}} + \frac{3dx\sqrt{a+bx^2}(2bc - ad)}{8b^2} + \frac{dx\sqrt{a+bx^2}(c + dx^2)}{4b}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^2/Sqrt[a + b*x^2], x]

[Out] (3*d*(2*b*c - a*d)*x*Sqrt[a + b*x^2])/(8*b^2) + (d*x*Sqrt[a + b*x^2]*(c + d*x^2))/(4*b) + ((8*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/(8*b^(5/2))

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)), x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q - 1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rubi steps

$$\begin{aligned}
\int \frac{(c + dx^2)^2}{\sqrt{a + bx^2}} dx &= \frac{dx\sqrt{a + bx^2} (c + dx^2)}{4b} + \frac{\int \frac{c(4bc - ad) + 3d(2bc - ad)x^2}{\sqrt{a + bx^2}} dx}{4b} \\
&= \frac{3d(2bc - ad)x\sqrt{a + bx^2}}{8b^2} + \frac{dx\sqrt{a + bx^2} (c + dx^2)}{4b} - \frac{(3ad(2bc - ad) - 2bc(4bc - ad)) \int \frac{1}{\sqrt{a + bx^2}}}{8b^2} \\
&= \frac{3d(2bc - ad)x\sqrt{a + bx^2}}{8b^2} + \frac{dx\sqrt{a + bx^2} (c + dx^2)}{4b} - \frac{(3ad(2bc - ad) - 2bc(4bc - ad)) \operatorname{Subst}\left(\int \frac{1}{\sqrt{a + bx^2}}\right)}{8b^2} \\
&= \frac{3d(2bc - ad)x\sqrt{a + bx^2}}{8b^2} + \frac{dx\sqrt{a + bx^2} (c + dx^2)}{4b} + \frac{(8b^2c^2 - 8abcd + 3a^2d^2) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a + bx^2}}\right)}{8b^{5/2}}
\end{aligned}$$

Mathematica [C] time = 2.49, size = 160, normalized size = 1.48

$$\frac{x\sqrt{\frac{bx^2}{a} + 1} \left(-2bx^2(c + dx^2)\right)^2 {}_3F_2\left(\frac{3}{2}, \frac{3}{2}, 2; 1, \frac{9}{2}; -\frac{bx^2}{a}\right) - 4bx^2(2c^2 + 3cdx^2 + d^2x^4) {}_2F_1\left(\frac{3}{2}, \frac{3}{2}; \frac{9}{2}; -\frac{bx^2}{a}\right) + 7a(15c^2 + 10cdx^2 + 3d^2x^4)}{105a\sqrt{a + bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(c + d*x^2)^2/Sqrt[a + b*x^2], x]

[Out] (x*Sqrt[1 + (b*x^2)/a]*(7*a*(15*c^2 + 10*c*d*x^2 + 3*d^2*x^4)*Hypergeometric2F1[1/2, 1/2, 7/2, -((b*x^2)/a)] - 4*b*x^2*(2*c^2 + 3*c*d*x^2 + d^2*x^4)*Hypergeometric2F1[3/2, 3/2, 9/2, -((b*x^2)/a)] - 2*b*x^2*(c + d*x^2)^2*HypergeometricPFQ[{3/2, 3/2, 2}, {1, 9/2}, -((b*x^2)/a)]))/(105*a*Sqrt[a + b*x^2])

fricas [A] time = 0.57, size = 192, normalized size = 1.78

$$\frac{\left((8b^2c^2 - 8abcd + 3a^2d^2)\sqrt{b} \log\left(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{bx - a}\right) + 2(2b^2d^2x^3 + (8b^2cd - 3abd^2)x)\sqrt{bx^2 + a}\right)}{16b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^(1/2), x, algorithm="fricas")

[Out] [1/16*((8*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 2*(2*b^2*d^2*x^3 + (8*b^2*c*d - 3*a*b*d^2)*x)*sqrt(b*x^2 + a))/b^3, -1/8*((8*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (2*b^2*d^2*x^3 + (8*b^2*c*d - 3*a*b*d^2)*x)*sqrt(b*x^2 + a))/b^3]

giac [A] time = 0.62, size = 90, normalized size = 0.83

$$\frac{1}{8}\sqrt{bx^2 + a}\left(\frac{2d^2x^2}{b} + \frac{8b^2cd - 3abd^2}{b^3}\right)x - \frac{(8b^2c^2 - 8abcd + 3a^2d^2)\log\left(\left|-\sqrt{bx} + \sqrt{bx^2 + a}\right|\right)}{8b^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^(1/2), x, algorithm="giac")

[Out] 1/8*sqrt(b*x^2 + a)*(2*d^2*x^2/b + (8*b^2*c*d - 3*a*b*d^2)/b^3)*x - 1/8*(8*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(5/2)

maple [A] time = 0.01, size = 131, normalized size = 1.21

$$\frac{\sqrt{bx^2+a} d^2 x^3}{4b} + \frac{3a^2 d^2 \ln(\sqrt{b}x + \sqrt{bx^2+a})}{8b^{\frac{5}{2}}} - \frac{acd \ln(\sqrt{b}x + \sqrt{bx^2+a})}{b^{\frac{3}{2}}} + \frac{c^2 \ln(\sqrt{b}x + \sqrt{bx^2+a})}{\sqrt{b}} - \frac{3\sqrt{bx^2+a}}{8b^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^2/(b*x^2+a)^(1/2), x)

[Out] 1/4*d^2*x^3/b*(b*x^2+a)^(1/2)-3/8*d^2*a/b^2*x*(b*x^2+a)^(1/2)+3/8*d^2*a^2/b^(5/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+c*d*x/b*(b*x^2+a)^(1/2)-c*d*a/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+c^2*ln(b^(1/2)*x+(b*x^2+a)^(1/2))/b^(1/2)

maxima [A] time = 1.47, size = 109, normalized size = 1.01

$$\frac{\sqrt{bx^2+a} d^2 x^3}{4b} + \frac{\sqrt{bx^2+a} cdx}{b} - \frac{3\sqrt{bx^2+a} ad^2 x}{8b^2} + \frac{c^2 \operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{b}} - \frac{acd \operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{b^{\frac{3}{2}}} + \frac{3a^2 d^2 \operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{8b^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^(1/2), x, algorithm="maxima")

[Out] 1/4*sqrt(b*x^2 + a)*d^2*x^3/b + sqrt(b*x^2 + a)*c*d*x/b - 3/8*sqrt(b*x^2 + a)*a*d^2*x/b^2 + c^2*arcsinh(b*x/sqrt(a*b))/sqrt(b) - a*c*d*arcsinh(b*x/sqrt(a*b))/b^(3/2) + 3/8*a^2*d^2*arcsinh(b*x/sqrt(a*b))/b^(5/2)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(dx^2 + c)^2}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^2/(a + b*x^2)^(1/2), x)

[Out] int((c + d*x^2)^2/(a + b*x^2)^(1/2), x)

sympy [A] time = 6.94, size = 238, normalized size = 2.20

$$\frac{3a^{\frac{3}{2}}d^2x}{8b^2\sqrt{1+\frac{bx^2}{a}}} + \frac{\sqrt{a}cdx\sqrt{1+\frac{bx^2}{a}}}{b} - \frac{\sqrt{a}d^2x^3}{8b\sqrt{1+\frac{bx^2}{a}}} + \frac{3a^2d^2 \operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{8b^{\frac{5}{2}}} - \frac{acd \operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{b^{\frac{3}{2}}} + c^2 \left\{ \begin{array}{l} \frac{\sqrt{-\frac{a}{b}} \operatorname{asin}\left(x\sqrt{-\frac{b}{a}}\right)}{\sqrt{a}} \\ \frac{\sqrt{\frac{a}{b}} \operatorname{asinh}\left(x\sqrt{\frac{b}{a}}\right)}{\sqrt{a}} \\ \frac{\sqrt{-\frac{a}{b}} \operatorname{acosh}\left(x\sqrt{-\frac{b}{a}}\right)}{\sqrt{-a}} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**2/(b*x**2+a)**(1/2), x)

[Out] -3*a**(3/2)*d**2*x/(8*b**2*sqrt(1 + b*x**2/a)) + sqrt(a)*c*d*x*sqrt(1 + b*x**2/a)/b - sqrt(a)*d**2*x**3/(8*b*sqrt(1 + b*x**2/a)) + 3*a**2*d**2*asinh(sqrt(b)*x/sqrt(a))/(8*b**(5/2)) - a*c*d*asinh(sqrt(b)*x/sqrt(a))/b**(3/2) + c**2*Piecewise((sqrt(-a/b)*asin(x*sqrt(-b/a))/sqrt(a), (a > 0) & (b < 0)), (sqrt(a/b)*asinh(x*sqrt(b/a))/sqrt(a), (a > 0) & (b > 0)), (sqrt(-a/b)*acosh(x*sqrt(-b/a))/sqrt(-a), (b > 0) & (a < 0))) + d**2*x**5/(4*sqrt(a)*sqrt(1 + b*x**2/a))

$$3.76 \quad \int \frac{c+dx^2}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=58

$$\frac{(2bc - ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2b^{3/2}} + \frac{dx\sqrt{a+bx^2}}{2b}$$

[Out] 1/2*(-a*d+2*b*c)*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(3/2)+1/2*d*x*(b*x^2+a)^(1/2)/b

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {388, 217, 206}

$$\frac{(2bc - ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2b^{3/2}} + \frac{dx\sqrt{a+bx^2}}{2b}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)/Sqrt[a + b*x^2], x]

[Out] (d*x*Sqrt[a + b*x^2])/(2*b) + ((2*b*c - a*d)*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/(2*b^(3/2))

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int \frac{c+dx^2}{\sqrt{a+bx^2}} dx &= \frac{dx\sqrt{a+bx^2}}{2b} - \frac{(-2bc+ad) \int \frac{1}{\sqrt{a+bx^2}} dx}{2b} \\ &= \frac{dx\sqrt{a+bx^2}}{2b} - \frac{(-2bc+ad) \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{2b} \\ &= \frac{dx\sqrt{a+bx^2}}{2b} + \frac{(2bc-ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2b^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 57, normalized size = 0.98

$$\frac{dx\sqrt{a+bx^2}}{2b} - \frac{(ad-2bc) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2b^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)/Sqrt[a + b*x^2],x]

[Out] (d*x*Sqrt[a + b*x^2])/(2*b) - ((-2*b*c + a*d)*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/(2*b^(3/2))

fricas [A] time = 0.69, size = 113, normalized size = 1.95

$$\left[\frac{2\sqrt{bx^2+a}bdx - (2bc-ad)\sqrt{b}\log\left(-2bx^2+2\sqrt{bx^2+a}\sqrt{b}x-a\right)}{4b^2}, \frac{\sqrt{bx^2+a}bdx - (2bc-ad)\sqrt{-b}\arctan\left(\frac{\sqrt{bx^2+a}}{\sqrt{-b}}\right)}{2b^2} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b*x^2 + a)*b*d*x - (2*b*c - a*d)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a))/b^2, 1/2*(sqrt(b*x^2 + a)*b*d*x - (2*b*c - a*d)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)))/b^2]

giac [A] time = 0.60, size = 49, normalized size = 0.84

$$\frac{\sqrt{bx^2+a}dx}{2b} - \frac{(2bc-ad)\log\left(\left|-\sqrt{b}x + \sqrt{bx^2+a}\right|\right)}{2b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(1/2),x, algorithm="giac")

[Out] 1/2*sqrt(b*x^2 + a)*d*x/b - 1/2*(2*b*c - a*d)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(3/2)

maple [A] time = 0.01, size = 62, normalized size = 1.07

$$-\frac{ad\ln\left(\sqrt{b}x + \sqrt{bx^2+a}\right)}{2b^{\frac{3}{2}}} + \frac{c\ln\left(\sqrt{b}x + \sqrt{bx^2+a}\right)}{\sqrt{b}} + \frac{\sqrt{bx^2+a}dx}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)/(b*x^2+a)^(1/2),x)

[Out] 1/2*d*x*(b*x^2+a)^(1/2)/b-1/2*d*a/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+c*ln(b^(1/2)*x+(b*x^2+a)^(1/2))/b^(1/2)

maxima [A] time = 1.33, size = 47, normalized size = 0.81

$$\frac{\sqrt{bx^2+a}dx}{2b} + \frac{c\operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{b}} - \frac{ad\operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{2b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] 1/2*sqrt(b*x^2 + a)*d*x/b + c*arcsinh(b*x/sqrt(a*b))/sqrt(b) - 1/2*a*d*arcsinh(b*x/sqrt(a*b))/b^(3/2)

mupad [B] time = 5.51, size = 86, normalized size = 1.48

$$\left\{ \begin{array}{ll} \frac{dx^3+3cx}{3\sqrt{a}} & \text{if } b = 0 \\ \frac{c \ln(\sqrt{b}x + \sqrt{bx^2+a})}{\sqrt{b}} - \frac{ad \ln(2\sqrt{b}x + 2\sqrt{bx^2+a})}{2b^{3/2}} + \frac{dx\sqrt{bx^2+a}}{2b} & \text{if } b \neq 0 \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)/(a + b*x^2)^(1/2),x)

[Out] piecewise(b == 0, (3*c*x + d*x^3)/(3*a^(1/2)), b != 0, (c*log(b^(1/2)*x + (a + b*x^2)^(1/2))/b^(1/2) - (a*d*log(2*b^(1/2)*x + 2*(a + b*x^2)^(1/2)))/(2*b^(3/2)) + (d*x*(a + b*x^2)^(1/2))/(2*b))

sympy [A] time = 2.78, size = 126, normalized size = 2.17

$$\frac{\sqrt{a} dx \sqrt{1 + \frac{bx^2}{a}}}{2b} - \frac{ad \operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{2b^{\frac{3}{2}}} + c \left\{ \begin{array}{ll} \frac{\sqrt{-\frac{a}{b}} \operatorname{asin}\left(x\sqrt{-\frac{b}{a}}\right)}{\sqrt{a}} & \text{for } a > 0 \wedge b < 0 \\ \frac{\sqrt{\frac{a}{b}} \operatorname{asinh}\left(x\sqrt{\frac{b}{a}}\right)}{\sqrt{a}} & \text{for } a > 0 \wedge b > 0 \\ \frac{\sqrt{-\frac{a}{b}} \operatorname{acosh}\left(x\sqrt{-\frac{b}{a}}\right)}{\sqrt{-a}} & \text{for } b > 0 \wedge a < 0 \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)/(b*x**2+a)**(1/2),x)

[Out] sqrt(a)*d*x*sqrt(1 + b*x**2/a)/(2*b) - a*d*asinh(sqrt(b)*x/sqrt(a))/(2*b**(3/2)) + c*Piecewise((sqrt(-a/b)*asin(x*sqrt(-b/a))/sqrt(a), (a > 0) & (b < 0)), (sqrt(a/b)*asinh(x*sqrt(b/a))/sqrt(a), (a > 0) & (b > 0)), (sqrt(-a/b)*acosh(x*sqrt(-b/a))/sqrt(-a), (b > 0) & (a < 0)))

$$3.77 \quad \int \frac{1}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=25

$$\frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{\sqrt{b}}$$

[Out] arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(1/2)

Rubi [A] time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {217, 206}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a + b*x^2], x]

[Out] ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]]/Sqrt[b]

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a+bx^2}} dx &= \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right) \\ &= \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{\sqrt{b}} \end{aligned}$$

Mathematica [A] time = 0.00, size = 25, normalized size = 1.00

$$\frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[a + b*x^2], x]

[Out] ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]]/Sqrt[b]

fricas [A] time = 0.56, size = 59, normalized size = 2.36

$$\left[\frac{\log\left(-2bx^2 - 2\sqrt{bx^2+a}\sqrt{b}x - a\right)}{2\sqrt{b}}, -\frac{\sqrt{-b}\arctan\left(\frac{\sqrt{-b}x}{\sqrt{bx^2+a}}\right)}{b} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] [1/2*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a)/sqrt(b), -sqrt(-b)*arc
tan(sqrt(-b)*x/sqrt(b*x^2 + a))/b]

giac [A] time = 0.60, size = 23, normalized size = 0.92

$$-\frac{\log\left(\left|-\sqrt{b}x + \sqrt{bx^2 + a}\right|\right)}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2),x, algorithm="giac")

[Out] -log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/sqrt(b)

maple [A] time = 0.00, size = 21, normalized size = 0.84

$$\frac{\ln\left(\sqrt{b}x + \sqrt{bx^2 + a}\right)}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/2),x)

[Out] ln(b^(1/2)*x+(b*x^2+a)^(1/2))/b^(1/2)

maxima [A] time = 1.28, size = 13, normalized size = 0.52

$$\frac{\operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] arcsinh(b*x/sqrt(a*b))/sqrt(b)

mupad [B] time = 0.12, size = 20, normalized size = 0.80

$$\frac{\ln\left(\sqrt{b}x + \sqrt{bx^2 + a}\right)}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + b*x^2)^(1/2),x)

[Out] log(b^(1/2)*x + (a + b*x^2)^(1/2))/b^(1/2)

sympy [A] time = 1.00, size = 17, normalized size = 0.68

$$\frac{\operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(1/2),x)

[Out] asinh(sqrt(b)*x/sqrt(a))/sqrt(b)

$$3.78 \quad \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx$$

Optimal. Leaf size=49

$$\frac{\tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}\sqrt{bc-ad}}$$

[Out] arctanh(x*(-a*d+b*c)^(1/2)/c^(1/2)/(b*x^2+a)^(1/2))/c^(1/2)/(-a*d+b*c)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {377, 208}

$$\frac{\tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}\sqrt{bc-ad}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[a + b*x^2]*(c + d*x^2)),x]

[Out] ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])]/(Sqrt[c]*Sqrt[b*c - a*d])

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx &= \text{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right) \\ &= \frac{\tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}\sqrt{bc-ad}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 49, normalized size = 1.00

$$\frac{\tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}\sqrt{bc-ad}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[a + b*x^2]*(c + d*x^2)),x]

[Out] ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])]/(Sqrt[c]*Sqrt[b*c - a*d])

fricas [B] time = 1.04, size = 241, normalized size = 4.92

$$\left[\frac{\log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4 + a^2c^2 + 2(4abc^2 - 3a^2cd)x^2 + 4((2bc - ad)x^3 + acx)\sqrt{bc^2 - acd}\sqrt{bx^2 + a}}{d^2x^4 + 2cdx^2 + c^2}\right)}{4\sqrt{bc^2 - acd}}, -\frac{\sqrt{-bc^2 + acd} \arctan\left(\frac{\sqrt{-bc^2 + acd}((2b^2c^2 - abcd)x^2 + a)}{2((b^2c^2 - abcd)x^2 + a)}\right)}{2(bc^2 - acd)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c),x, algorithm="fricas")

[Out] [1/4*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2))/sqrt(b*c^2 - a*c*d), -1/2*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x))/(b*c^2 - a*c*d)]

giac [A] time = 0.60, size = 70, normalized size = 1.43

$$\frac{\sqrt{b} \arctan\left(\frac{(\sqrt{b}x - \sqrt{bx^2 + a})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{\sqrt{-b^2c^2 + abcd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c),x, algorithm="giac")

[Out] -sqrt(b)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/sqrt(-b^2*c^2 + a*b*c*d)

maple [B] time = 0.02, size = 300, normalized size = 6.12

$$\frac{\ln\left(\frac{\frac{2\sqrt{-cd}\left(x - \frac{\sqrt{-cd}}{d}\right)b}{d} + \frac{2ad - 2bc}{d} + 2\sqrt{\frac{ad - bc}{d}}\sqrt{\left(x - \frac{\sqrt{-cd}}{d}\right)^2 b + \frac{2\sqrt{-cd}\left(x - \frac{\sqrt{-cd}}{d}\right)b}{d} + \frac{ad - bc}{d}}}{x - \frac{\sqrt{-cd}}{d}}\right)}{2\sqrt{-cd}\sqrt{\frac{ad - bc}{d}}}\right) + \frac{\ln\left(\frac{\frac{2\sqrt{-cd}\left(x + \frac{\sqrt{-cd}}{d}\right)b}{d} + \frac{2ad - 2bc}{d} + 2\sqrt{\frac{ad - bc}{d}}\sqrt{\left(x + \frac{\sqrt{-cd}}{d}\right)^2 b + \frac{2\sqrt{-cd}\left(x + \frac{\sqrt{-cd}}{d}\right)b}{d} + \frac{ad - bc}{d}}}{x + \frac{\sqrt{-cd}}{d}}\right)}{2\sqrt{-cd}\sqrt{\frac{ad - bc}{d}}}\right)}{2\sqrt{-cd}\sqrt{\frac{ad - bc}{d}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/2)/(d*x^2+c),x)

[Out] -1/2/(-c*d)^(1/2)/((a*d-b*c)/d)^(1/2)*ln((2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^(1/2)*((x-(-c*d)^(1/2)/d)^2*b+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/2))/(x-(-c*d)^(1/2)/d)+1/2/(-c*d)^(1/2)/((a*d-b*c)/d)^(1/2)*ln((-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^(1/2)*((x+(-c*d)^(1/2)/d)^2*b-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/2))/(x+(-c*d)^(1/2)/d)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2 + a}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*x^2 + a)*(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\left\{ \begin{array}{ll} \frac{\operatorname{atan}\left(\frac{x\sqrt{ad-bc}}{\sqrt{c}\sqrt{bx^2+a}}\right)}{\sqrt{c(ad-bc)}} & \text{if } 0 < ad - bc \\ \frac{\ln\left(\frac{\sqrt{c(bx^2+a)+x\sqrt{bc-ad}}}{\sqrt{c(bx^2+a)-x\sqrt{bc-ad}}}\right)}{2\sqrt{-c(ad-bc)}} & \text{if } ad - bc < 0 \\ \int \frac{1}{\sqrt{bx^2+a}(dx^2+c)} dx & \text{if } ad - bc \notin \mathbb{R} \vee ad = bc \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(1/2)*(c + d*x^2)),x)

[Out] piecewise(0 < a*d - b*c, atan((x*(a*d - b*c)^(1/2))/(c^(1/2)*(a + b*x^2)^(1/2)))/(c*(a*d - b*c)^(1/2), a*d - b*c < 0, log(((c*(a + b*x^2)^(1/2) + x*(- a*d + b*c)^(1/2))/((c*(a + b*x^2)^(1/2) - x*(- a*d + b*c)^(1/2)))/(2*(- c*(a*d - b*c)^(1/2))), ~in(a*d - b*c, 'real') | a*d == b*c, int(1/((a + b*x^2)^(1/2)*(c + d*x^2)), x))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a + bx^2} (c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(1/2)/(d*x**2+c),x)

[Out] Integral(1/(sqrt(a + b*x**2)*(c + d*x**2)), x)

$$3.79 \quad \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)^2} dx$$

Optimal. Leaf size=101

$$\frac{(2bc - ad) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}(bc - ad)^{3/2}} - \frac{dx\sqrt{a+bx^2}}{2c(c+dx^2)(bc - ad)}$$

[Out] 1/2*(-a*d+2*b*c)*arctanh(x*(-a*d+b*c)^(1/2)/c^(1/2)/(b*x^2+a)^(1/2))/c^(3/2)/(-a*d+b*c)^(3/2)-1/2*d*x*(b*x^2+a)^(1/2)/c/(-a*d+b*c)/(d*x^2+c)

Rubi [A] time = 0.05, antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {382, 377, 208}

$$\frac{(2bc - ad) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}(bc - ad)^{3/2}} - \frac{dx\sqrt{a+bx^2}}{2c(c+dx^2)(bc - ad)}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[a + b*x^2]*(c + d*x^2)^2), x]

[Out] -(d*x*Sqrt[a + b*x^2])/(2*c*(b*c - a*d)*(c + d*x^2)) + ((2*b*c - a*d)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(2*c^(3/2)*(b*c - a*d)^(3/2))

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)], Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+2)+1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)^2} dx &= -\frac{dx\sqrt{a+bx^2}}{2c(bc-ad)(c+dx^2)} + \frac{(2bc-ad) \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{2c(bc-ad)} \\ &= -\frac{dx\sqrt{a+bx^2}}{2c(bc-ad)(c+dx^2)} + \frac{(2bc-ad) \operatorname{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{2c(bc-ad)} \\ &= -\frac{dx\sqrt{a+bx^2}}{2c(bc-ad)(c+dx^2)} + \frac{(2bc-ad) \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}(bc-ad)^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.30, size = 126, normalized size = 1.25

$$\frac{x \left(\frac{(c+dx^2)(2bc-ad) \tanh^{-1}\left(\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}\right)}{c \sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}} - d(a+bx^2) \right)}{2c\sqrt{a+bx^2}(c+dx^2)(bc-ad)}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[a + b*x^2]*(c + d*x^2)^2), x]

[Out] (x*(-(d*(a + b*x^2)) + ((2*b*c - a*d)*(c + d*x^2)*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/(c*Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]))/(2*c*(b*c - a*d)*Sqrt[a + b*x^2]*(c + d*x^2))

fricas [B] time = 1.04, size = 463, normalized size = 4.58

$$\frac{4(bc^2d - acd^2)\sqrt{bx^2 + a}x - (2bc^2 - acd + (2bcd - ad^2)x^2)\sqrt{bc^2 - acd} \log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4 + a^2c^2 + 2(4abc^2 - ad^2)x^2 + a^2c^2}{d^2}\right)}{8(b^2c^5 - 2abc^4d + a^2c^3d^2 + (b^2c^4d - 2abc^3d^2 + a^2c^2d^3)x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] [-1/8*(4*(b*c^2*d - a*c*d^2)*sqrt(b*x^2 + a)*x - (2*b*c^2 - a*c*d + (2*b*c*d - a*d^2)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)))/(b^2*c^5 - 2*a*b*c^4*d + a^2*c^3*d^2 + (b^2*c^4*d - 2*a*b*c^3*d^2 + a^2*c^2*d^3)*x^2), -1/4*(2*(b*c^2*d - a*c*d^2)*sqrt(b*x^2 + a)*x + (2*b*c^2 - a*c*d + (2*b*c*d - a*d^2)*x^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)))/(b^2*c^5 - 2*a*b*c^4*d + a^2*c^3*d^2 + (b^2*c^4*d - 2*a*b*c^3*d^2 + a^2*c^2*d^3)*x^2)]

giac [B] time = 0.63, size = 242, normalized size = 2.40

$$\frac{\frac{1}{2}b^{\frac{3}{2}} \left((2bc - ad) \arctan\left(\frac{(\sqrt{b}x - \sqrt{bx^2 + a})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right) \right)}{(b^2c^2 - abcd)\sqrt{-b^2c^2 + abcd}} - \frac{2 \left(2(\sqrt{b}x - \sqrt{bx^2 + a})^2 bc - (\sqrt{b}x - \sqrt{bx^2 + a})^4 d + 4(\sqrt{b}x - \sqrt{bx^2 + a})^2 bc - 2(\sqrt{b}x - \sqrt{bx^2 + a})^4 d \right)}{\left((\sqrt{b}x - \sqrt{bx^2 + a})^4 d + 4(\sqrt{b}x - \sqrt{bx^2 + a})^2 bc - 2(\sqrt{b}x - \sqrt{bx^2 + a})^4 d \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^2,x, algorithm="giac")
```

```
[Out] 1/2*b^(3/2)*((2*b*c - a*d)*arctan(-1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d +
2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/((b^2*c^2 - a*b*c*d)*sqrt(-b^2*c^2
+ a*b*c*d)) - 2*(2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*b*c - (sqrt(b)*x - sqrt(
b*x^2 + a))^2*a*d + a^2*d)/(((sqrt(b)*x - sqrt(b*x^2 + a))^4*d + 4*(sqrt(b)
*x - sqrt(b*x^2 + a))^2*b*c - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*d + a^2*d
)*(b^2*c^2 - a*b*c*d))
```

```
maple [B] time = 0.02, size = 809, normalized size = 8.01
```

$$\frac{\sqrt{-cd} b \ln \left(\frac{\frac{2\sqrt{-cd} \left(x - \frac{\sqrt{-cd}}{d}\right) b}{d} + \frac{2ad-2bc}{d} + 2\sqrt{\frac{ad-bc}{d}} \sqrt{\left(x - \frac{\sqrt{-cd}}{d}\right)^2 + \frac{2\sqrt{-cd} \left(x - \frac{\sqrt{-cd}}{d}\right) b + \frac{ad-bc}{d}}}{x - \frac{\sqrt{-cd}}{d}} \right)}{4(ad-bc)\sqrt{\frac{ad-bc}{d}} cd} + \frac{\sqrt{-cd} b \ln \left(\frac{\frac{2\sqrt{-cd} \left(x + \frac{\sqrt{-cd}}{d}\right) b}{d} + \frac{2ad-2bc}{d}}{\dots} \right)}{4(ad-bc)\sqrt{\frac{ad-bc}{d}} cd}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(b*x^2+a)^(1/2)/(d*x^2+c)^2,x)
```

```
[Out] 1/4/c/(a*d-b*c)/(x-(-c*d)^(1/2)/d)*((x-(-c*d)^(1/2)/d)^2*b+2*(-c*d)^(1/2)*(
x-(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/2)-1/4/c/d*(-c*d)^(1/2)*b/(a*d-b*c)/(
(a*d-b*c)/d)^(1/2)*ln((2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+2*(a*d-b*c)/d+
2*((a*d-b*c)/d)^(1/2)*((x-(-c*d)^(1/2)/d)^2*b+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)
)/d)*b/d+(a*d-b*c)/d)^(1/2))/(x-(-c*d)^(1/2)/d))+1/4/c/(a*d-b*c)/(x+(-c*d)^(
1/2)/d)*((x+(-c*d)^(1/2)/d)^2*b-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)*b/d+(a*d
-b*c)/d)^(1/2)+1/4/c/d*(-c*d)^(1/2)*b/(a*d-b*c)/((a*d-b*c)/d)^(1/2)*ln((-2*
(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^(1/2)*((x
+(-c*d)^(1/2)/d)^2*b-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/
2))/(x+(-c*d)^(1/2)/d))-1/4/c/(-c*d)^(1/2)/((a*d-b*c)/d)^(1/2)*ln((2*(-c*d)
^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^(1/2)*((x-(-c*d)
)^(1/2)/d)^2*b+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/2))/(x
-(-c*d)^(1/2)/d))+1/4/c/(-c*d)^(1/2)/((a*d-b*c)/d)^(1/2)*ln((-2*(-c*d)^(1/2)
)*(x+(-c*d)^(1/2)/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^(1/2)*((x+(-c*d)^(1/
2)/d)^2*b-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/2))/(x+(-c
d)^(1/2)/d))
```

```
maxima [F] time = 0.00, size = 0, normalized size = 0.00
```

$$\int \frac{1}{\sqrt{bx^2 + a} (dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^2,x, algorithm="maxima")
```

```
[Out] integrate(1/(sqrt(b*x^2 + a)*(d*x^2 + c)^2), x)
```

```
mupad [F] time = 0.00, size = -1, normalized size = -0.01
```

$$\int \frac{1}{\sqrt{bx^2 + a} (dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((a + b*x^2)^(1/2)*(c + d*x^2)^2),x)
```

[Out] `int(1/((a + b*x^2)^(1/2)*(c + d*x^2)^2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a + bx^2} (c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x**2+a)**(1/2)/(d*x**2+c)**2,x)`

[Out] `Integral(1/(sqrt(a + b*x**2)*(c + d*x**2)**2), x)`

$$3.80 \quad \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)^3} dx$$

Optimal. Leaf size=163

$$\frac{(3a^2d^2 - 8abcd + 8b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}(bc-ad)^{5/2}} - \frac{3dx\sqrt{a+bx^2}(2bc-ad)}{8c^2(c+dx^2)(bc-ad)^2} - \frac{dx\sqrt{a+bx^2}}{4c(c+dx^2)^2(bc-ad)}$$

[Out] 1/8*(3*a^2*d^2-8*a*b*c*d+8*b^2*c^2)*arctanh(x*(-a*d+b*c)^(1/2)/c^(1/2)/(b*x^2+a)^(1/2))/c^(5/2)/(-a*d+b*c)^(5/2)-1/4*d*x*(b*x^2+a)^(1/2)/c/(-a*d+b*c)/(d*x^2+c)^2-3/8*d*(-a*d+2*b*c)*x*(b*x^2+a)^(1/2)/c^2/(-a*d+b*c)^2/(d*x^2+c)

Rubi [A] time = 0.12, antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {414, 527, 12, 377, 208}

$$\frac{(3a^2d^2 - 8abcd + 8b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}(bc-ad)^{5/2}} - \frac{3dx\sqrt{a+bx^2}(2bc-ad)}{8c^2(c+dx^2)(bc-ad)^2} - \frac{dx\sqrt{a+bx^2}}{4c(c+dx^2)^2(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[a + b*x^2]*(c + d*x^2)^3), x]

[Out] -(d*x*Sqrt[a + b*x^2])/(4*c*(b*c - a*d)*(c + d*x^2)^2) - (3*d*(2*b*c - a*d)*x*Sqrt[a + b*x^2])/(8*c^2*(b*c - a*d)^2*(c + d*x^2)) + ((8*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(8*c^(5/2)*(b*c - a*d)^(5/2))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527


```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a+bx^2} (c+dx^2)^3} dx &= -\frac{dx\sqrt{a+bx^2}}{4c(bc-ad)(c+dx^2)^2} + \frac{\int \frac{4bc-3ad-2bdx^2}{\sqrt{a+bx^2}(c+dx^2)^2} dx}{4c(bc-ad)} \\ &= -\frac{dx\sqrt{a+bx^2}}{4c(bc-ad)(c+dx^2)^2} - \frac{3d(2bc-ad)x\sqrt{a+bx^2}}{8c^2(bc-ad)^2(c+dx^2)} + \frac{\int \frac{8b^2c^2-8abcd+3a^2d^2}{\sqrt{a+bx^2}(c+dx^2)} dx}{8c^2(bc-ad)^2} \\ &= -\frac{dx\sqrt{a+bx^2}}{4c(bc-ad)(c+dx^2)^2} - \frac{3d(2bc-ad)x\sqrt{a+bx^2}}{8c^2(bc-ad)^2(c+dx^2)} + \frac{(8b^2c^2-8abcd+3a^2d^2) \int}{8c^2(bc-ad)^2} \\ &= -\frac{dx\sqrt{a+bx^2}}{4c(bc-ad)(c+dx^2)^2} - \frac{3d(2bc-ad)x\sqrt{a+bx^2}}{8c^2(bc-ad)^2(c+dx^2)} + \frac{(8b^2c^2-8abcd+3a^2d^2) S}{8c^2} \\ &= -\frac{dx\sqrt{a+bx^2}}{4c(bc-ad)(c+dx^2)^2} - \frac{3d(2bc-ad)x\sqrt{a+bx^2}}{8c^2(bc-ad)^2(c+dx^2)} + \frac{(8b^2c^2-8abcd+3a^2d^2) ta}{8c^{5/2}(bc-ad)} \end{aligned}$$

Mathematica [A] time = 0.67, size = 192, normalized size = 1.18

$$x \frac{\left((c+dx^2)^2 (3a^2d^2 - 8abcd + 8b^2c^2) \tanh^{-1} \left(\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}} \right) - cd(a^2(-d)(5c + 3dx^2) + ab(8c^2 + cdx^2 - 3d^2x^4) + 2b^2cx^2(4c + \sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}) \right)}{8c^3\sqrt{a+bx^2}(c+dx^2)^2(bc-ad)^2}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[a + b*x^2]*(c + d*x^2)^3), x]

[Out] (x*(-(c*d*(2*b^2*c*x^2*(4*c + 3*d*x^2) - a^2*d*(5*c + 3*d*x^2) + a*b*(8*c^2 + c*d*x^2 - 3*d^2*x^4))) + ((8*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*(c + d*x^2)^2*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2)))))/(8*c^3*(b*c - a*d)^2*Sqrt[a + b*x^2]*(c + d*x^2)^3)

fricas [B] time = 1.02, size = 864, normalized size = 5.30

$$\left[\frac{(8b^2c^4 - 8abc^3d + 3a^2c^2d^2 + (8b^2c^2d^2 - 8abcd^3 + 3a^2d^4)x^4 + 2(8b^2c^3d - 8abc^2d^2 + 3a^2cd^3)x^2)\sqrt{bc^2 - a^2}}{32(b^3c^8 - 3ab^2c^7d + 3a^2bc^6d^2 - a^3c^5d^3)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^3,x, algorithm="fricas")

```
[Out] [1/32*((8*b^2*c^4 - 8*a*b*c^3*d + 3*a^2*c^2*d^2 + (8*b^2*c^2*d^2 - 8*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(8*b^2*c^3*d - 8*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) - 4*(3*(2*b^2*c^3*d^2 - 3*a*b*c^2*d^3 + a^2*c*d^4)*x^3 + (8*b^2*c^4*d - 13*a*b*c^3*d^2 + 5*a^2*c^2*d^3)*x)*sqrt(b*x^2 + a))/(b^3*c^8 - 3*a*b^2*c^7*d + 3*a^2*b*c^6*d^2 - a^3*c^5*d^3 + (b^3*c^6*d^2 - 3*a*b^2*c^5*d^3 + 3*a^2*b*c^4*d^4 - a^3*c^3*d^5)*x^4 + 2*(b^3*c^7*d - 3*a*b^2*c^6*d^2 + 3*a^2*b*c^5*d^3 - a^3*c^4*d^4)*x^2), -1/16*((8*b^2*c^4 - 8*a*b*c^3*d + 3*a^2*c^2*d^2 + (8*b^2*c^2*d^2 - 8*a*b*c*d^3 + 3*a^2*d^4)*x^4 + 2*(8*b^2*c^3*d - 8*a*b*c^2*d^2 + 3*a^2*c*d^3)*x^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a))/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x) + 2*(3*(2*b^2*c^3*d^2 - 3*a*b*c^2*d^3 + a^2*c*d^4)*x^3 + (8*b^2*c^4*d - 13*a*b*c^3*d^2 + 5*a^2*c^2*d^3)*x)*sqrt(b*x^2 + a))/(b^3*c^8 - 3*a*b^2*c^7*d + 3*a^2*b*c^6*d^2 - a^3*c^5*d^3 + (b^3*c^6*d^2 - 3*a*b^2*c^5*d^3 + 3*a^2*b*c^4*d^4 - a^3*c^3*d^5)*x^4 + 2*(b^3*c^7*d - 3*a*b^2*c^6*d^2 + 3*a^2*b*c^5*d^3 - a^3*c^4*d^4)*x^2)]
```

giac [B] time = 3.51, size = 538, normalized size = 3.30

$$-\frac{1}{8} b^{\frac{5}{2}} \left(\frac{(8b^2c^2 - 8abcd + 3a^2d^2) \arctan\left(\frac{(\sqrt{bx - \sqrt{bx^2 + a}})^2 d + bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{(b^4c^4 - 2ab^3c^3d + a^2b^2c^2d^2)\sqrt{-b^2c^2 + abcd}} \right) + \frac{2\left(8(\sqrt{bx} - \sqrt{bx^2 + a})^6 b^2c^2d - 8(\sqrt{bx} - \sqrt{bx^2 + a})^4 b^2c^2d + 8(\sqrt{bx} - \sqrt{bx^2 + a})^2 b^2c^2d - 8b^2c^2d\right)}{(b^4c^4 - 2ab^3c^3d + a^2b^2c^2d^2)\sqrt{-b^2c^2 + abcd}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^3,x, algorithm="giac")
```

```
[Out] -1/8*b^(5/2)*((8*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/((b^4*c^4 - 2*a*b^3*c^3*d + a^2*b^2*c^2*d^2)*sqrt(-b^2*c^2 + a*b*c*d)) + 2*(8*(sqrt(b)*x - sqrt(b*x^2 + a))^6*b^2*c^2*d - 8*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a*b*c*d^2 + 3*(sqrt(b)*x - sqrt(b*x^2 + a))^6*a^2*d^3 + 48*(sqrt(b)*x - sqrt(b*x^2 + a))^4*b^3*c^3 - 72*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a*b^2*c^2*d + 42*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^2*b*c*d^2 - 9*(sqrt(b)*x - sqrt(b*x^2 + a))^4*a^3*d^3 + 40*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^2*b^2*c^2*d - 40*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^3*b*c*d^2 + 9*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a^4*d^3 + 6*a^4*b*c*d^2 - 3*a^5*d^3)/((b^4*c^4 - 2*a*b^3*c^3*d + a^2*b^2*c^2*d^2)*((sqrt(b)*x - sqrt(b*x^2 + a))^4*d + 4*(sqrt(b)*x - sqrt(b*x^2 + a))^2*b*c - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*d + a^2*d^2))
```

maple [B] time = 0.02, size = 1815, normalized size = 11.13

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(b*x^2+a)^(1/2)/(d*x^2+c)^3,x)
```

```
[Out] 1/16/(-c*d)^(1/2)/c/(a*d-b*c)/(x-(-c*d)^(1/2)/d)^2*((x-(-c*d)^(1/2)/d)^2*b+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/2)-3/16/c*b/(a*d-b*c)^2/(x-(-c*d)^(1/2)/d)*((x-(-c*d)^(1/2)/d)^2*b+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/2)-3/16/(-c*d)^(1/2)*b^2/(a*d-b*c)^2/((a*d-b*c)/d)^(1/2)*ln((2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^(1/2))*((x-(-c*d)^(1/2)/d)^2*b+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+(a*d-b*c)/d)^(1/2))/(x-(-c*d)^(1/2)/d)-1/16/(-c*d)^(1/2)/c*b/(a*d-b*c)/((a*d-b*c)/d)^(1/2)*ln((2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)*b/d+2*(a*d-b*c)/d+2*((a
```

$$\begin{aligned} & *d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*} \\ & b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))-1/16/(-c*d)^{(1/2)}/c/(a*d-b*c)/(\\ & x+(-c*d)^{(1/2)}/d)^{2*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)} \\ & d)*b/d+(a*d-b*c)/d)^{(1/2)}-3/16/c*b/(a*d-b*c)^{2}/(x+(-c*d)^{(1/2)}/d)*((x+(-c*d) \\ &)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}+3/1 \\ & 6/(-c*d)^{(1/2)}*b^2/(a*d-b*c)^2/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d) \\ &)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))+1/16/(-c*d)^{(1/2)}/c*b/(a*d-b*c)/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))+3/16/c^2/(a*d-b*c)/(x+(-c*d)^{(1/2)}/d)*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}+3/16/c^2/d*(-c*d)^{(1/2)}*b/(a*d-b*c)/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))+3/16/c^2/(a*d-b*c)/(x-(-c*d)^{(1/2)}/d)*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}-3/16/c^2/d*(-c*d)^{(1/2)}*b/(a*d-b*c)/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))-3/16/(-c*d)^{(1/2)}/c^2/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))+3/16/(-c*d)^{(1/2)}/c^2/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d)) \end{aligned}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2+a}(dx^2+c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^3,x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*x^2+a)*(d*x^2+c)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{bx^2+a}(dx^2+c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a+b*x^2)^(1/2)*(c+d*x^2)^3),x)

[Out] int(1/((a+b*x^2)^(1/2)*(c+d*x^2)^3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(1/2)/(d*x**2+c)**3,x)

[Out] Timed out

$$3.81 \quad \int \frac{(c+dx^2)^4}{(a+bx^2)^{3/2}} dx$$

Optimal. Leaf size=257

$$\frac{dx\sqrt{a+bx^2}(c+dx^2)(35a^2d^2-64abcd+24b^2c^2)}{24ab^3} + \frac{d(-35a^3d^3+120a^2bcd^2-144ab^2c^2d+64b^3c^3)\tanh^{-1}\left(\frac{dx\sqrt{a+bx^2}}{\sqrt{a+bx^2}}\right)}{16b^{9/2}}$$

[Out] 1/16*d*(-35*a^3*d^3+120*a^2*b*c*d^2-144*a*b^2*c^2*d+64*b^3*c^3)*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(9/2)+(-a*d+b*c)*x*(d*x^2+c)^3/a/b/(b*x^2+a)^(1/2)-1/48*d*(-105*a^3*d^3+290*a^2*b*c*d^2-248*a*b^2*c^2*d+48*b^3*c^3)*x*(b*x^2+a)^(1/2)/a/b^4-1/24*d*(35*a^2*d^2-64*a*b*c*d+24*b^2*c^2)*x*(d*x^2+c)*(b*x^2+a)^(1/2)/a/b^3-1/6*d*(-7*a*d+6*b*c)*x*(d*x^2+c)^2*(b*x^2+a)^(1/2)/a/b^2

Rubi [A] time = 0.26, antiderivative size = 257, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {413, 528, 388, 217, 206}

$$\frac{dx\sqrt{a+bx^2}(c+dx^2)(35a^2d^2-64abcd+24b^2c^2)}{24ab^3} - \frac{dx\sqrt{a+bx^2}(290a^2bcd^2-105a^3d^3-248ab^2c^2d+48b^3c^3)}{48ab^4}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^4/(a + b*x^2)^(3/2), x]

[Out] -(d*(48*b^3*c^3 - 248*a*b^2*c^2*d + 290*a^2*b*c*d^2 - 105*a^3*d^3)*x*Sqrt[a + b*x^2])/(48*a*b^4) - (d*(24*b^2*c^2 - 64*a*b*c*d + 35*a^2*d^2)*x*Sqrt[a + b*x^2]*(c + d*x^2))/(24*a*b^3) - (d*(6*b*c - 7*a*d)*x*Sqrt[a + b*x^2]*(c + d*x^2)^2)/(6*a*b^2) + ((b*c - a*d)*x*(c + d*x^2)^3)/(a*b*Sqrt[a + b*x^2]) + (d*(64*b^3*c^3 - 144*a*b^2*c^2*d + 120*a^2*b*c*d^2 - 35*a^3*d^3)*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/(16*b^(9/2))

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+q)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,

0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 528

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^4}{(a + bx^2)^{3/2}} dx &= \frac{(bc - ad)x(c + dx^2)^3}{ab\sqrt{a + bx^2}} + \frac{\int \frac{(c+dx^2)^2(acd-d(6bc-7ad)x^2)}{\sqrt{a+bx^2}} dx}{ab} \\ &= -\frac{d(6bc - 7ad)x\sqrt{a + bx^2}(c + dx^2)^2}{6ab^2} + \frac{(bc - ad)x(c + dx^2)^3}{ab\sqrt{a + bx^2}} + \frac{\int \frac{(c+dx^2)(acd(12bc-7ad)-d(24b^2c^2-64abcd+35a^2d^2))}{\sqrt{a+bx^2}} dx}{6ab^2} \\ &= -\frac{d(24b^2c^2 - 64abcd + 35a^2d^2)x\sqrt{a + bx^2}(c + dx^2)}{24ab^3} - \frac{d(6bc - 7ad)x\sqrt{a + bx^2}(c + dx^2)}{6ab^2} \\ &= -\frac{d(48b^3c^3 - 248ab^2c^2d + 290a^2bcd^2 - 105a^3d^3)x\sqrt{a + bx^2}}{48ab^4} - \frac{d(24b^2c^2 - 64abcd + 35a^2d^2)}{24ab^2} \\ &= -\frac{d(48b^3c^3 - 248ab^2c^2d + 290a^2bcd^2 - 105a^3d^3)x\sqrt{a + bx^2}}{48ab^4} - \frac{d(24b^2c^2 - 64abcd + 35a^2d^2)}{24ab^2} \\ &= -\frac{d(48b^3c^3 - 248ab^2c^2d + 290a^2bcd^2 - 105a^3d^3)x\sqrt{a + bx^2}}{48ab^4} - \frac{d(24b^2c^2 - 64abcd + 35a^2d^2)}{24ab^2} \end{aligned}$$

Mathematica [A] time = 5.20, size = 172, normalized size = 0.67

$$\frac{\sqrt{b}x\sqrt{a + bx^2} \left(3d^2(19a^2d^2 - 56abcd + 48b^2c^2) + 2bd^3x^2(24bc - 11ad) + \frac{48(bc-ad)^4}{a(ax^2)} + 8b^2d^4x^4 \right) + 3d(-35a^3d^3)}{48b^{9/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^4/(a + b*x^2)^(3/2), x]

[Out] (Sqrt[b]*x*Sqrt[a + b*x^2]*(3*d^2*(48*b^2*c^2 - 56*a*b*c*d + 19*a^2*d^2) + 2*b*d^3*(24*b*c - 11*a*d)*x^2 + 8*b^2*d^4*x^4 + (48*(b*c - a*d)^4)/(a*(a + b*x^2))) + 3*d*(64*b^3*c^3 - 144*a*b^2*c^2*d + 120*a^2*b*c*d^2 - 35*a^3*d^3)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]]/(48*b^(9/2))

fricas [A] time = 0.94, size = 584, normalized size = 2.27

$$\left[\frac{3(64a^2b^3c^3d - 144a^3b^2c^2d^2 + 120a^4bcd^3 - 35a^5d^4 + (64ab^4c^3d - 144a^2b^3c^2d^2 + 120a^3b^2cd^3 - 35a^4bd^4))}{48b^{9/2}} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] [-1/96*(3*(64*a^2*b^3*c^3*d - 144*a^3*b^2*c^2*d^2 + 120*a^4*b*c*d^3 - 35*a^5*d^4 + (64*a*b^4*c^3*d - 144*a^2*b^3*c^2*d^2 + 120*a^3*b^2*c*d^3 - 35*a^4*b*d^4)*x^2)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(8*a*b^4*d^4*x^7 + 2*(24*a*b^4*c*d^3 - 7*a^2*b^3*d^4)*x^5 + (144*a*b^4*c^2*d^2 - 120*a^2*b^3*c*d^3 + 35*a^3*b^2*d^4)*x^3 + 3*(16*b^5*c^4 - 64*a*b^4*c^3*d + 144*a^2*b^3*c^2*d^2 - 120*a^3*b^2*c*d^3 + 35*a^4*b*d^4)*x)*sqrt(b*x^2 + a))/(a*b^6*x^2 + a^2*b^5), -1/48*(3*(64*a^2*b^3*c^3*d - 144*a^3*b^2*c^2*d^2 + 120*a^4*b*c*d^3 - 35*a^5*d^4 + (64*a*b^4*c^3*d - 144*a^2*b^3*c^2*d^2 + 120*a^3*b^2*c*d^3 - 35*a^4*b*d^4)*x^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (8*a*b^4*d^4*x^7 + 2*(24*a*b^4*c*d^3 - 7*a^2*b^3*d^4)*x^5 + (144*a*b^4*c^2*d^2 - 120*a^2*b^3*c*d^3 + 35*a^3*b^2*d^4)*x^3 + 3*(16*b^5*c^4 - 64*a*b^4*c^3*d + 144*a^2*b^3*c^2*d^2 - 120*a^3*b^2*c*d^3 + 35*a^4*b*d^4)*x)*sqrt(b*x^2 + a))/(a*b^6*x^2 + a^2*b^5)]

giac [A] time = 0.67, size = 235, normalized size = 0.91

$$\frac{\left(2\left(\frac{4d^4x^2}{b} + \frac{24ab^6cd^3 - 7a^2b^5d^4}{ab^7}\right)x^2 + \frac{144ab^6c^2d^2 - 120a^2b^5cd^3 + 35a^3b^4d^4}{ab^7}\right)x^2 + \frac{3(16b^7c^4 - 64ab^6c^3d + 144a^2b^5c^2d^2 - 120a^3b^4cd^3 + 35a^4b^3d^4)}{ab^7}}{48\sqrt{bx^2 + a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^(3/2),x, algorithm="giac")

[Out] 1/48*((2*(4*d^4*x^2/b + (24*a*b^6*c*d^3 - 7*a^2*b^5*d^4)/(a*b^7))*x^2 + (14*4*a*b^6*c^2*d^2 - 120*a^2*b^5*c*d^3 + 35*a^3*b^4*d^4)/(a*b^7))*x^2 + 3*(16*b^7*c^4 - 64*a*b^6*c^3*d + 144*a^2*b^5*c^2*d^2 - 120*a^3*b^4*c*d^3 + 35*a^4*b^3*d^4)/(a*b^7))*x/sqrt(b*x^2 + a) - 1/16*(64*b^3*c^3*d - 144*a*b^2*c^2*d^2 + 120*a^2*b*c*d^3 - 35*a^3*d^4)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(9/2)

maple [A] time = 0.02, size = 340, normalized size = 1.32

$$\frac{d^4x^7}{6\sqrt{bx^2 + a}b} - \frac{7ad^4x^5}{24\sqrt{bx^2 + a}b^2} + \frac{cd^3x^5}{\sqrt{bx^2 + a}b} + \frac{35a^2d^4x^3}{48\sqrt{bx^2 + a}b^3} - \frac{5acd^3x^3}{2\sqrt{bx^2 + a}b^2} + \frac{3c^2d^2x^3}{\sqrt{bx^2 + a}b} + \frac{35a^3d^4x}{16\sqrt{bx^2 + a}b^4} - \frac{2c^4x}{\sqrt{bx^2 + a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^4/(b*x^2+a)^(3/2),x)

[Out] 1/6*d^4*x^7/b/(b*x^2+a)^(1/2)-7/24*d^4*a/b^2*x^5/(b*x^2+a)^(1/2)+35/48*d^4*a^2/b^3*x^3/(b*x^2+a)^(1/2)+35/16*d^4*a^3/b^4*x/(b*x^2+a)^(1/2)-35/16*d^4*a^3/b^(9/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+c*d^3*x^5/b/(b*x^2+a)^(1/2)-5/2*c*d^3*a/b^2*x^3/(b*x^2+a)^(1/2)-15/2*c*d^3*a^2/b^3*x/(b*x^2+a)^(1/2)+15/2*c*d^3*a^2/b^(7/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+3*c^2*d^2*x^3/b/(b*x^2+a)^(1/2)+9*c^2*d^2*a/b^2*x/(b*x^2+a)^(1/2)-9*c^2*d^2*a/b^(5/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))-4*c^3*d*x/b/(b*x^2+a)^(1/2)+4*c^3*d/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+c^4*x/a/(b*x^2+a)^(1/2)

maxima [A] time = 1.38, size = 311, normalized size = 1.21

$$\frac{d^4x^7}{6\sqrt{bx^2 + a}b} + \frac{cd^3x^5}{\sqrt{bx^2 + a}b} - \frac{7ad^4x^5}{24\sqrt{bx^2 + a}b^2} + \frac{3c^2d^2x^3}{\sqrt{bx^2 + a}b} - \frac{5acd^3x^3}{2\sqrt{bx^2 + a}b^2} + \frac{35a^2d^4x^3}{48\sqrt{bx^2 + a}b^3} + \frac{c^4x}{\sqrt{bx^2 + a}a} - \frac{4c^3dx}{\sqrt{bx^2 + a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^(3/2),x, algorithm="maxima")

```
[Out] 1/6*d^4*x^7/(sqrt(b*x^2 + a)*b) + c*d^3*x^5/(sqrt(b*x^2 + a)*b) - 7/24*a*d^4*x^5/(sqrt(b*x^2 + a)*b^2) + 3*c^2*d^2*x^3/(sqrt(b*x^2 + a)*b) - 5/2*a*c*d^3*x^3/(sqrt(b*x^2 + a)*b^2) + 35/48*a^2*d^4*x^3/(sqrt(b*x^2 + a)*b^3) + c^4*x/(sqrt(b*x^2 + a)*a) - 4*c^3*d*x/(sqrt(b*x^2 + a)*b) + 9*a*c^2*d^2*x/(sqrt(b*x^2 + a)*b^2) - 15/2*a^2*c*d^3*x/(sqrt(b*x^2 + a)*b^3) + 35/16*a^3*d^4*x/(sqrt(b*x^2 + a)*b^4) + 4*c^3*d*arcsinh(b*x/sqrt(a*b))/b^(3/2) - 9*a*c^2*d^2*arcsinh(b*x/sqrt(a*b))/b^(5/2) + 15/2*a^2*c*d^3*arcsinh(b*x/sqrt(a*b))/b^(7/2) - 35/16*a^3*d^4*arcsinh(b*x/sqrt(a*b))/b^(9/2)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(dx^2 + c)^4}{(bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c + d*x^2)^4/(a + b*x^2)^(3/2), x)
```

```
[Out] int((c + d*x^2)^4/(a + b*x^2)^(3/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + dx^2)^4}{(a + bx^2)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x**2+c)**4/(b*x**2+a)**(3/2), x)
```

```
[Out] Integral((c + d*x**2)**4/(a + b*x**2)**(3/2), x)
```

$$3.82 \quad \int \frac{(c+dx^2)^3}{(a+bx^2)^{3/2}} dx$$

Optimal. Leaf size=169

$$\frac{3d(5a^2d^2 - 12abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{8b^{7/2}} - \frac{dx\sqrt{a+bx^2}(2bc-5ad)(4bc-3ad)}{8ab^3} - \frac{dx\sqrt{a+bx^2}(c+dx^2)(4bc-5ad)}{4ab^2}$$

[Out] 3/8*d*(5*a^2*d^2-12*a*b*c*d+8*b^2*c^2)*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(7/2)+(-a*d+b*c)*x*(d*x^2+c)^2/a/b/(b*x^2+a)^(1/2)-1/8*d*(-5*a*d+2*b*c)*(-3*a*d+4*b*c)*x*(b*x^2+a)^(1/2)/a/b^3-1/4*d*(-5*a*d+4*b*c)*x*(d*x^2+c)*(b*x^2+a)^(1/2)/a/b^2

Rubi [A] time = 0.20, antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {413, 528, 388, 217, 206}

$$\frac{3d(5a^2d^2 - 12abcd + 8b^2c^2) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{8b^{7/2}} - \frac{dx\sqrt{a+bx^2}(c+dx^2)(4bc-5ad)}{4ab^2} - \frac{dx\sqrt{a+bx^2}(2bc-5ad)(4bc-5ad)}{8ab^3}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^3/(a + b*x^2)^(3/2), x]

[Out] -(d*(2*b*c - 5*a*d)*(4*b*c - 3*a*d)*x*Sqrt[a + b*x^2])/(8*a*b^3) - (d*(4*b*c - 5*a*d)*x*Sqrt[a + b*x^2]*(c + d*x^2))/(4*a*b^2) + ((b*c - a*d)*x*(c + d*x^2)^2)/(a*b*Sqrt[a + b*x^2]) + (3*d*(8*b^2*c^2 - 12*a*b*c*d + 5*a^2*d^2)*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/(8*b^(7/2))

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+q)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 528


```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
f_)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^3}{(a + bx^2)^{3/2}} dx &= \frac{(bc - ad)x(c + dx^2)^2}{ab\sqrt{a + bx^2}} + \frac{\int \frac{(c+dx^2)(acd-d(4bc-5ad)x^2)}{\sqrt{a+bx^2}} dx}{ab} \\ &= -\frac{d(4bc - 5ad)x\sqrt{a + bx^2}(c + dx^2)}{4ab^2} + \frac{(bc - ad)x(c + dx^2)^2}{ab\sqrt{a + bx^2}} + \frac{\int \frac{acd(8bc-5ad)-d(2bc-5ad)(4bc-5ad)x^2}{\sqrt{a+bx^2}} dx}{4ab^2} \\ &= -\frac{d(2bc - 5ad)(4bc - 3ad)x\sqrt{a + bx^2}}{8ab^3} - \frac{d(4bc - 5ad)x\sqrt{a + bx^2}(c + dx^2)}{4ab^2} + \frac{(bc - ad)x(c + dx^2)^2}{ab\sqrt{a + bx^2}} \\ &= -\frac{d(2bc - 5ad)(4bc - 3ad)x\sqrt{a + bx^2}}{8ab^3} - \frac{d(4bc - 5ad)x\sqrt{a + bx^2}(c + dx^2)}{4ab^2} + \frac{(bc - ad)x(c + dx^2)^2}{ab\sqrt{a + bx^2}} \\ &= -\frac{d(2bc - 5ad)(4bc - 3ad)x\sqrt{a + bx^2}}{8ab^3} - \frac{d(4bc - 5ad)x\sqrt{a + bx^2}(c + dx^2)}{4ab^2} + \frac{(bc - ad)x(c + dx^2)^2}{ab\sqrt{a + bx^2}} \end{aligned}$$

Mathematica [A] time = 5.10, size = 122, normalized size = 0.72

$$\frac{3d(5a^2d^2 - 12abcd + 8b^2c^2) \log\left(\sqrt{b}\sqrt{a + bx^2} + bx\right)}{8b^{7/2}} + \frac{x\sqrt{a + bx^2} \left(d^2(12bc - 7ad) + \frac{8(bc-ad)^3}{a(a+bx^2)} + 2bd^3x^2\right)}{8b^3}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^3/(a + b*x^2)^(3/2), x]

[Out] (x*Sqrt[a + b*x^2]*(d^2*(12*b*c - 7*a*d) + 2*b*d^3*x^2 + (8*(b*c - a*d)^3)/(a*(a + b*x^2))))/(8*b^3) + (3*d*(8*b^2*c^2 - 12*a*b*c*d + 5*a^2*d^2)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(8*b^(7/2))

fricas [A] time = 1.00, size = 416, normalized size = 2.46

$$\frac{3(8a^2b^2c^2d - 12a^3bcd^2 + 5a^4d^3 + (8ab^3c^2d - 12a^2b^2cd^2 + 5a^3bd^3)x^2)\sqrt{b} \log\left(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{b}x - a\right)}{16(ab^5x^2 + \dots)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^(3/2), x, algorithm="fricas")

[Out] [1/16*(3*(8*a^2*b^2*c^2*d - 12*a^3*b*c*d^2 + 5*a^4*d^3 + (8*a*b^3*c^2*d - 12*a^2*b^2*c*d^2 + 5*a^3*b*d^3)*x^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 2*(2*a*b^3*d^3*x^5 + (12*a*b^3*c*d^2 - 5*a^2*b^2*d^3)*x^3 + (8*b^4*c^3 - 24*a*b^3*c^2*d + 36*a^2*b^2*c*d^2 - 15*a^3*b*d^3)*x)*sqrt(b*x^2 + a))/(a*b^5*x^2 + a^2*b^4), -1/8*(3*(8*a^2*b^2*c^2*d - 12*a^3*b*c*d^2 + 5*a^4*d^3 + (8*a*b^3*c^2*d - 12*a^2*b^2*c*d^2 + 5*a^3*b*d^3)*x^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (2*a*b^3*d^3*x^5 + (12*a*b^3*c*d^2

$- 5a^2b^2d^3)x^3 + (8b^4c^3 - 24a^2b^3c^2d + 36a^2b^2c^2d^2 - 15a^3b^2d^3)x) \sqrt{bx^2 + a} / (ab^5x^2 + a^2b^4)]$

giac [A] time = 0.64, size = 157, normalized size = 0.93

$$\frac{\left(\left(\frac{2d^3x^2}{b} + \frac{12ab^4cd^2 - 5a^2b^3d^3}{ab^5}\right)x^2 + \frac{8b^5c^3 - 24ab^4c^2d + 36a^2b^3cd^2 - 15a^3b^2d^3}{ab^5}\right)x - 3(8b^2c^2d - 12abcd^2 + 5a^2d^3) \log\left(\left|-\sqrt{b}x + \sqrt{bx^2 + a}\right|\right)}{8\sqrt{bx^2 + a} \cdot 8b^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^(3/2),x, algorithm="giac")

[Out] $\frac{1}{8} \left(\frac{(2d^3x^2/b + (12a^2b^4c^2d - 5a^2b^3d^3)/(ab^5))x^2 + (8b^5c^3 - 24a^2b^4c^2d + 36a^2b^3c^2d^2 - 15a^3b^2d^3)/(ab^5)}{ab^5} \right) x / \sqrt{bx^2 + a} - \frac{3}{8} \frac{(8b^2c^2d - 12a^2b^3cd^2 + 5a^2d^3) \log(\text{abs}(-\sqrt{b}x + \sqrt{bx^2 + a}))}{b^{7/2}}$

maple [A] time = 0.01, size = 219, normalized size = 1.30

$$\frac{d^3x^5}{4\sqrt{bx^2 + a}b} - \frac{5ad^3x^3}{8\sqrt{bx^2 + a}b^2} + \frac{3cd^2x^3}{2\sqrt{bx^2 + a}b} - \frac{15a^2d^3x}{8\sqrt{bx^2 + a}b^3} + \frac{9acd^2x}{2\sqrt{bx^2 + a}b^2} + \frac{c^3x}{\sqrt{bx^2 + a}a} - \frac{3c^2dx}{\sqrt{bx^2 + a}b} + \frac{15a^2d^3}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^3/(b*x^2+a)^(3/2),x)

[Out] $\frac{1}{4}d^3x^5/b/(b^2x^2+a)^{1/2} - \frac{5}{8}d^3a/b^2x^3/(b^2x^2+a)^{1/2} - \frac{15}{8}d^3a^2/b^3x/(b^2x^2+a)^{1/2} + \frac{15}{8}d^3a^2/b^3 \ln(b^{1/2}x + (b^2x^2+a)^{1/2}) + \frac{3}{2}cd^2x^3/b/(b^2x^2+a)^{1/2} + \frac{9}{2}cd^2a/b^2x/(b^2x^2+a)^{1/2} - \frac{9}{2}cd^2a^2/b^3 \ln(b^{1/2}x + (b^2x^2+a)^{1/2}) - \frac{3}{2}c^2dx/b/(b^2x^2+a)^{1/2} + \frac{3}{2}c^2d/b^3 \ln(b^{1/2}x + (b^2x^2+a)^{1/2}) + \frac{c^3x/a}{(b^2x^2+a)^{1/2}}$

maxima [A] time = 1.40, size = 197, normalized size = 1.17

$$\frac{d^3x^5}{4\sqrt{bx^2 + a}b} + \frac{3cd^2x^3}{2\sqrt{bx^2 + a}b} - \frac{5ad^3x^3}{8\sqrt{bx^2 + a}b^2} + \frac{c^3x}{\sqrt{bx^2 + a}a} - \frac{3c^2dx}{\sqrt{bx^2 + a}b} + \frac{9acd^2x}{2\sqrt{bx^2 + a}b^2} - \frac{15a^2d^3x}{8\sqrt{bx^2 + a}b^3} + \frac{3c^2d \operatorname{arsinh}(bx/\sqrt{a})}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] $\frac{1}{4}d^3x^5/(\sqrt{bx^2 + a}b) + \frac{3}{2}cd^2x^3/(\sqrt{bx^2 + a}b) - \frac{5}{8}d^3a^2x^3/(\sqrt{bx^2 + a}b^2) + \frac{c^3x}{(\sqrt{bx^2 + a}a)} - \frac{3}{2}c^2dx/(\sqrt{bx^2 + a}b) + \frac{9}{2}acd^2x/(\sqrt{bx^2 + a}b^2) - \frac{15}{8}a^2d^3x/(\sqrt{bx^2 + a}b^3) + \frac{3}{2}c^2d \operatorname{arcsinh}(bx/\sqrt{a})/b^{3/2} - \frac{9}{2}acd^2 \operatorname{arcsinh}(bx/\sqrt{a})/b^{5/2} + \frac{15}{8}a^2d^3 \operatorname{arcsinh}(bx/\sqrt{a})/b^{7/2}$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(dx^2 + c)^3}{(bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^3/(a + b*x^2)^(3/2),x)

[Out] int((c + d*x^2)^3/(a + b*x^2)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + dx^2)^3}{(a + bx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**3/(b*x**2+a)**(3/2), x)

[Out] Integral((c + d*x**2)**3/(a + b*x**2)**(3/2), x)

$$3.83 \quad \int \frac{(c+dx^2)^2}{(a+bx^2)^{3/2}} dx$$

Optimal. Leaf size=90

$$\frac{d(4bc - 3ad) \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2b^{5/2}} + \frac{x(bc - ad)^2}{ab^2\sqrt{a+bx^2}} + \frac{d^2x\sqrt{a+bx^2}}{2b^2}$$

[Out] $1/2*d*(-3*a*d+4*b*c)*\operatorname{arctanh}(x*b^{(1/2)}/(b*x^2+a)^{(1/2)})/b^{(5/2)}+(-a*d+b*c)^2*x/a/b^{(2)}/(b*x^2+a)^{(1/2)}+1/2*d^2*x*(b*x^2+a)^{(1/2)}/b^2$

Rubi [A] time = 0.06, antiderivative size = 105, normalized size of antiderivative = 1.17, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {413, 388, 217, 206}

$$-\frac{dx\sqrt{a+bx^2}(2bc-3ad)}{2ab^2} + \frac{d(4bc-3ad)\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{2b^{5/2}} + \frac{x(c+dx^2)(bc-ad)}{ab\sqrt{a+bx^2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^2/(a + b*x^2)^(3/2), x]

[Out] $-(d*(2*b*c - 3*a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(2*a*b^2) + ((b*c - a*d)*x*(c + d*x^2))/(a*b*\operatorname{Sqrt}[a + b*x^2]) + (d*(4*b*c - 3*a*d)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a + b*x^2]])/(2*b^{(5/2)})$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+q)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rubi steps

$$\begin{aligned}
\int \frac{(c + dx^2)^2}{(a + bx^2)^{3/2}} dx &= \frac{(bc - ad)x(c + dx^2)}{ab\sqrt{a + bx^2}} + \frac{\int \frac{acd - d(2bc - 3ad)x^2}{\sqrt{a + bx^2}} dx}{ab} \\
&= -\frac{d(2bc - 3ad)x\sqrt{a + bx^2}}{2ab^2} + \frac{(bc - ad)x(c + dx^2)}{ab\sqrt{a + bx^2}} + \frac{(d(4bc - 3ad)) \int \frac{1}{\sqrt{a + bx^2}} dx}{2b^2} \\
&= -\frac{d(2bc - 3ad)x\sqrt{a + bx^2}}{2ab^2} + \frac{(bc - ad)x(c + dx^2)}{ab\sqrt{a + bx^2}} + \frac{(d(4bc - 3ad)) \operatorname{Subst}\left(\int \frac{1}{1 - bx^2} dx, x, \sqrt{a + bx^2}\right)}{2b^2} \\
&= -\frac{d(2bc - 3ad)x\sqrt{a + bx^2}}{2ab^2} + \frac{(bc - ad)x(c + dx^2)}{ab\sqrt{a + bx^2}} + \frac{d(4bc - 3ad) \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a + bx^2}}\right)}{2b^{5/2}}
\end{aligned}$$

Mathematica [C] time = 2.53, size = 160, normalized size = 1.78

$$\frac{x\sqrt{\frac{bx^2}{a} + 1} \left(-6bx^2(c + dx^2)^2 {}_3F_2\left(\frac{3}{2}, 2, \frac{5}{2}; 1, \frac{9}{2}; -\frac{bx^2}{a}\right) - 12bx^2(2c^2 + 3cdx^2 + d^2x^4) {}_2F_1\left(\frac{3}{2}, \frac{5}{2}; \frac{9}{2}; -\frac{bx^2}{a}\right) + 7a(15) \right)}{105a^2\sqrt{a + bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(c + d*x^2)^2/(a + b*x^2)^(3/2), x]

[Out] (x*sqrt[1 + (b*x^2)/a]*(7*a*(15*c^2 + 10*c*d*x^2 + 3*d^2*x^4)*Hypergeometric2F1[1/2, 3/2, 7/2, -((b*x^2)/a)] - 12*b*x^2*(2*c^2 + 3*c*d*x^2 + d^2*x^4)*Hypergeometric2F1[3/2, 5/2, 9/2, -((b*x^2)/a)] - 6*b*x^2*(c + d*x^2)^2*HypergeometricPFQ[{3/2, 2, 5/2}, {1, 9/2}, -((b*x^2)/a)]))/(105*a^2*sqrt[a + b*x^2])

fricas [A] time = 0.71, size = 276, normalized size = 3.07

$$\left[\frac{(4a^2bcd - 3a^3d^2 + (4ab^2cd - 3a^2bd^2)x^2)\sqrt{b} \log\left(-2bx^2 + 2\sqrt{bx^2 + a}\sqrt{bx} - a\right) - 2(ab^2d^2x^3 + (2b^3c^2 - 4ab^2cd + 3a^2bd^2)x)\sqrt{bx^2 + a}}{4(ab^4x^2 + a^2b^3)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^(3/2), x, algorithm="fricas")

[Out] [-1/4*((4*a^2*b*c*d - 3*a^3*d^2 + (4*a*b^2*c*d - 3*a^2*b*d^2)*x^2)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(a*b^2*d^2*x^3 + (2*b^3*c^2 - 4*a*b^2*c*d + 3*a^2*b*d^2)*x)*sqrt(b*x^2 + a))/(a*b^4*x^2 + a^2*b^3), -1/2*((4*a^2*b*c*d - 3*a^3*d^2 + (4*a*b^2*c*d - 3*a^2*b*d^2)*x^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (a*b^2*d^2*x^3 + (2*b^3*c^2 - 4*a*b^2*c*d + 3*a^2*b*d^2)*x)*sqrt(b*x^2 + a))/(a*b^4*x^2 + a^2*b^3)]

giac [A] time = 0.63, size = 92, normalized size = 1.02

$$\frac{\left(\frac{d^2x^2}{b} + \frac{2b^3c^2 - 4ab^2cd + 3a^2bd^2}{ab^3}\right)x}{2\sqrt{bx^2 + a}} - \frac{(4bcd - 3ad^2) \log\left(\left|-\sqrt{b}x + \sqrt{bx^2 + a}\right|\right)}{2b^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^(3/2), x, algorithm="giac")

[Out] $\frac{1}{2} \cdot \frac{d^2 x^2 / b + (2 \cdot b^3 c^2 - 4 \cdot a \cdot b^2 c d + 3 \cdot a^2 b d^2)}{(a \cdot b^3)} \cdot x / \sqrt{b x^2 + a} - \frac{1}{2} \cdot \frac{(4 \cdot b \cdot c \cdot d - 3 \cdot a \cdot d^2) \cdot \log(\text{abs}(-\sqrt{b} \cdot x + \sqrt{b x^2 + a}))}{b^{5/2}}$

maple [A] time = 0.01, size = 123, normalized size = 1.37

$$\frac{\frac{d^2 x^3}{2 \sqrt{b x^2 + a} b} + \frac{3 a d^2 x}{2 \sqrt{b x^2 + a} b^2} + \frac{c^2 x}{\sqrt{b x^2 + a} a} - \frac{2 c d x}{\sqrt{b x^2 + a} b} - \frac{3 a d^2 \ln(\sqrt{b} x + \sqrt{b x^2 + a})}{2 b^{\frac{5}{2}}} + \frac{2 c d \ln(\sqrt{b} x + \sqrt{b x^2 + a})}{b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+c)^2/(b*x^2+a)^(3/2),x)`

[Out] $\frac{1}{2} \cdot \frac{d^2 x^3}{b} / (b x^2 + a)^{1/2} + \frac{3}{2} \cdot \frac{d^2 a}{b^2 x} / (b x^2 + a)^{1/2} - \frac{3}{2} \cdot \frac{d^2 a}{b^{5/2}} \cdot \ln(b^{1/2} x + (b x^2 + a)^{1/2}) - \frac{2 \cdot c \cdot d \cdot x}{b} / (b x^2 + a)^{1/2} + \frac{2 \cdot c \cdot d}{b^{3/2}} \cdot \ln(b^{1/2} x + (b x^2 + a)^{1/2}) + \frac{c^2 x}{a} / (b x^2 + a)^{1/2}$

maxima [A] time = 1.38, size = 108, normalized size = 1.20

$$\frac{\frac{d^2 x^3}{2 \sqrt{b x^2 + a} b} + \frac{c^2 x}{\sqrt{b x^2 + a} a} - \frac{2 c d x}{\sqrt{b x^2 + a} b} + \frac{3 a d^2 x}{2 \sqrt{b x^2 + a} b^2} + \frac{2 c d \operatorname{arsinh}\left(\frac{b x}{\sqrt{a b}}\right)}{b^{\frac{3}{2}}} - \frac{3 a d^2 \operatorname{arsinh}\left(\frac{b x}{\sqrt{a b}}\right)}{2 b^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^2/(b*x^2+a)^(3/2),x, algorithm="maxima")`

[Out] $\frac{1}{2} \cdot \frac{d^2 x^3}{(\sqrt{b x^2 + a}) \cdot b} + \frac{c^2 x}{(\sqrt{b x^2 + a}) \cdot a} - \frac{2 \cdot c \cdot d \cdot x}{(\sqrt{b x^2 + a}) \cdot b} + \frac{3}{2} \cdot \frac{a \cdot d^2 x}{(\sqrt{b x^2 + a}) \cdot b^2} + \frac{2 \cdot c \cdot d \cdot \operatorname{arcsinh}(b x / \sqrt{a \cdot b})}{b^{3/2}} - \frac{3}{2} \cdot \frac{a \cdot d^2 \cdot \operatorname{arcsinh}(b x / \sqrt{a \cdot b})}{b^{5/2}}$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d x^2 + c)^2}{(b x^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^2/(a + b*x^2)^(3/2),x)`

[Out] `int((c + d*x^2)^2/(a + b*x^2)^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + d x^2)^2}{(a + b x^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**2/(b*x**2+a)**(3/2),x)`

[Out] `Integral((c + d*x**2)**2/(a + b*x**2)**(3/2), x)`

$$3.84 \quad \int \frac{c+dx^2}{(a+bx^2)^{3/2}} dx$$

Optimal. Leaf size=54

$$\frac{d \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{b^{3/2}} + \frac{x(bc-ad)}{ab\sqrt{a+bx^2}}$$

[Out] d*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(3/2)+(-a*d+b*c)*x/a/b/(b*x^2+a)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {385, 217, 206}

$$\frac{d \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{b^{3/2}} + \frac{x(bc-ad)}{ab\sqrt{a+bx^2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)/(a + b*x^2)^(3/2), x]

[Out] ((b*c - a*d)*x)/(a*b*sqrt[a + b*x^2]) + (d*ArcTanh[(sqrt[b]*x)/sqrt[a + b*x^2]])/b^(3/2)

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rubi steps

$$\begin{aligned} \int \frac{c+dx^2}{(a+bx^2)^{3/2}} dx &= \frac{(bc-ad)x}{ab\sqrt{a+bx^2}} + \frac{d \int \frac{1}{\sqrt{a+bx^2}} dx}{b} \\ &= \frac{(bc-ad)x}{ab\sqrt{a+bx^2}} + \frac{d \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{b} \\ &= \frac{(bc-ad)x}{ab\sqrt{a+bx^2}} + \frac{d \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{b^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 70, normalized size = 1.30

$$\frac{a^{3/2}d\sqrt{\frac{bx^2}{a}+1}\sinh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)+\sqrt{b}x(bc-ad)}{ab^{3/2}\sqrt{a+bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)/(a + b*x^2)^(3/2), x]

[Out] (Sqrt[b]*(b*c - a*d)*x + a^(3/2)*d*Sqrt[1 + (b*x^2)/a]*ArcSinh[(Sqrt[b]*x)/Sqrt[a]])/(a*b^(3/2)*Sqrt[a + b*x^2])

fricas [A] time = 0.76, size = 167, normalized size = 3.09

$$\left[\frac{2(b^2c - abd)\sqrt{bx^2 + a}x + (abdx^2 + a^2d)\sqrt{b} \log\left(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{b}x - a\right)}{2(ab^3x^2 + a^2b^2)}, \frac{(b^2c - abd)\sqrt{bx^2 + a}x - (abdx^2 + a^2d)\sqrt{b} \arctan\left(\frac{\sqrt{b}x}{\sqrt{bx^2 + a}}\right)}{ab^3x^2} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(3/2), x, algorithm="fricas")

[Out] [1/2*(2*(b^2*c - a*b*d)*sqrt(b*x^2 + a)*x + (a*b*d*x^2 + a^2*d)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a))/(a*b^3*x^2 + a^2*b^2), ((b^2*c - a*b*d)*sqrt(b*x^2 + a)*x - (a*b*d*x^2 + a^2*d)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)))/(a*b^3*x^2 + a^2*b^2)]

giac [A] time = 0.64, size = 50, normalized size = 0.93

$$-\frac{d \log\left(\left|-\sqrt{b}x + \sqrt{bx^2 + a}\right|\right)}{b^{\frac{3}{2}}} + \frac{(bc - ad)x}{\sqrt{bx^2 + a}ab}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(3/2), x, algorithm="giac")

[Out] -d*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(3/2) + (b*c - a*d)*x/(sqrt(b*x^2 + a)*a*b)

maple [A] time = 0.00, size = 54, normalized size = 1.00

$$\frac{cx}{\sqrt{bx^2 + a}a} - \frac{dx}{\sqrt{bx^2 + a}b} + \frac{d \ln\left(\sqrt{b}x + \sqrt{bx^2 + a}\right)}{b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)/(b*x^2+a)^(3/2), x)

[Out] -d*x/b/(b*x^2+a)^(1/2)+d/b^(3/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+c*x/a/(b*x^2+a)^(1/2)

maxima [A] time = 1.30, size = 46, normalized size = 0.85

$$\frac{cx}{\sqrt{bx^2 + a}a} - \frac{dx}{\sqrt{bx^2 + a}b} + \frac{d \operatorname{arsinh}\left(\frac{bx}{\sqrt{ab}}\right)}{b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(3/2), x, algorithm="maxima")

[Out] $c*x/(sqrt(b*x^2 + a)*a) - d*x/(sqrt(b*x^2 + a)*b) + d*arcsinh(b*x/sqrt(a*b))/b^(3/2)$

mupad [B] time = 5.12, size = 53, normalized size = 0.98

$$\frac{d \ln\left(\sqrt{b} x + \sqrt{b x^2 + a}\right)}{b^{3/2}} + \frac{c x}{a \sqrt{b x^2 + a}} - \frac{d x}{b \sqrt{b x^2 + a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)/(a + b*x^2)^(3/2), x)`

[Out] $(d*\log(b^{(1/2)*x} + (a + b*x^2)^{(1/2)}))/b^{(3/2)} + (c*x)/(a*(a + b*x^2)^{(1/2)}) - (d*x)/(b*(a + b*x^2)^{(1/2)})$

sympy [A] time = 5.17, size = 60, normalized size = 1.11

$$d \left(\frac{\operatorname{asinh}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{b^{\frac{3}{2}}} - \frac{x}{\sqrt{a}b\sqrt{1 + \frac{bx^2}{a}}} \right) + \frac{cx}{a^{\frac{3}{2}}\sqrt{1 + \frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)/(b*x**2+a)**(3/2), x)`

[Out] $d*(\operatorname{asinh}(\sqrt{b}*x/\sqrt{a}))/b^{(3/2)} - x/(\sqrt{a}*b*\sqrt{1 + b*x**2/a})) + c*x/(a^{(3/2)}*\sqrt{1 + b*x**2/a})$

$$3.85 \quad \int \frac{1}{(a+bx^2)^{3/2}} dx$$

Optimal. Leaf size=16

$$\frac{x}{a\sqrt{a+bx^2}}$$

[Out] x/a/(b*x^2+a)^(1/2)

Rubi [A] time = 0.00, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {191}

$$\frac{x}{a\sqrt{a+bx^2}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(-3/2), x]

[Out] x/(a*Sqrt[a + b*x^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rubi steps

$$\int \frac{1}{(a+bx^2)^{3/2}} dx = \frac{x}{a\sqrt{a+bx^2}}$$

Mathematica [A] time = 0.00, size = 16, normalized size = 1.00

$$\frac{x}{a\sqrt{a+bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(-3/2), x]

[Out] x/(a*Sqrt[a + b*x^2])

fricas [A] time = 0.68, size = 23, normalized size = 1.44

$$\frac{\sqrt{bx^2 + a} x}{abx^2 + a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2), x, algorithm="fricas")

[Out] sqrt(b*x^2 + a)*x/(a*b*x^2 + a^2)

giac [A] time = 0.60, size = 14, normalized size = 0.88

$$\frac{x}{\sqrt{bx^2 + a} a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2),x, algorithm="giac")

[Out] x/(sqrt(b*x^2 + a)*a)

maple [A] time = 0.00, size = 15, normalized size = 0.94

$$\frac{x}{\sqrt{bx^2 + a} a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(3/2),x)

[Out] 1/(b*x^2+a)^(1/2)/a*x

maxima [A] time = 1.32, size = 14, normalized size = 0.88

$$\frac{x}{\sqrt{bx^2 + a} a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] x/(sqrt(b*x^2 + a)*a)

mupad [B] time = 0.04, size = 14, normalized size = 0.88

$$\frac{x}{a \sqrt{bx^2 + a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + b*x^2)^(3/2),x)

[Out] x/(a*(a + b*x^2)^(1/2))

sympy [A] time = 0.61, size = 17, normalized size = 1.06

$$\frac{x}{a^{\frac{3}{2}} \sqrt{1 + \frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(3/2),x)

[Out] x/(a**(3/2)*sqrt(1 + b*x**2/a))

$$3.86 \quad \int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)} dx$$

Optimal. Leaf size=79

$$\frac{bx}{a\sqrt{a+bx^2}(bc-ad)} - \frac{d \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc-ad)^{3/2}}$$

[Out] $-d \operatorname{arctanh}\left(\frac{x(-a+d+bc)^{1/2}/c^{1/2}}{(b^2x^2+a)^{1/2}}\right) / (-a+d+bc)^{3/2} / c^{1/2} + bx/a / (-a+d+bc) / (b^2x^2+a)^{1/2}$

Rubi [A] time = 0.04, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {382, 377, 208}

$$\frac{bx}{a\sqrt{a+bx^2}(bc-ad)} - \frac{d \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc-ad)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(3/2)*(c + d*x^2)),x]

[Out] $(bx)/(a*(bc - a*d)*\text{Sqrt}[a + b*x^2]) - (d*\text{ArcTanh}[(\text{Sqrt}[bc - a*d]*x)/(\text{Sqrt}[c]*\text{Sqrt}[a + b*x^2])]) / (\text{Sqrt}[c]*(bc - a*d)^{3/2})$

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+2)+1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)} dx = \frac{bx}{a(bc-ad)\sqrt{a+bx^2}} - \frac{d \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{bc-ad}$$

$$= \frac{bx}{a(bc-ad)\sqrt{a+bx^2}} - \frac{d \operatorname{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x, \frac{x}{\sqrt{a+bx^2}}\right)}{bc-ad}$$

$$= \frac{bx}{a(bc-ad)\sqrt{a+bx^2}} - \frac{d \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc-ad)^{3/2}}$$

Mathematica [C] time = 0.72, size = 309, normalized size = 3.91

$$\frac{x \left(2dx^2 \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{5/2} {}_2F_1 \left(2, \frac{5}{2}; \frac{7}{2}; \frac{(bc-ad)x^2}{c(bx^2+a)} \right) + 2c \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{5/2} {}_2F_1 \left(2, \frac{5}{2}; \frac{7}{2}; \frac{(bc-ad)x^2}{c(bx^2+a)} \right) - 10dx^2 \sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}} - 15c \sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}} \right)}{5c^2 (a+bx^2)^{3/2} \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{3/2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(3/2)*(c + d*x^2)),x]

[Out] (x*(-15*c*Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]) - 10*d*x^2*Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]) + 15*c*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]) + 10*d*x^2*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]) + 2*c*((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b*c - a*d)*x^2)/(c*(a + b*x^2))] + 2*d*x^2*((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2)*Hypergeometric2F1[2, 5/2, 7/2, ((b*c - a*d)*x^2)/(c*(a + b*x^2))]) / (5*c^2*((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(3/2)*(a + b*x^2)^(3/2))

fricas [B] time = 1.07, size = 441, normalized size = 5.58

$$\frac{4(b^2c^2 - abcd)\sqrt{bx^2 + a}x - (abdx^2 + a^2d)\sqrt{bc^2 - acd} \log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4 + a^2c^2 + 2(4abc^2 - 3a^2cd)x^2 + 4((2bc-ad)d^2x^4 + 2cdx^2 + c^2)}{4(a^2b^2c^3 - 2a^3bc^2d + a^4cd^2 + (ab^3c^3 - 2a^2b^2c^2d + a^3bcd^2)x^2)}\right)}{4(a^2b^2c^3 - 2a^3bc^2d + a^4cd^2 + (ab^3c^3 - 2a^2b^2c^2d + a^3bcd^2)x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c),x, algorithm="fricas")

[Out] [1/4*(4*(b^2*c^2 - a*b*c*d)*sqrt(b*x^2 + a)*x - (a*b*d*x^2 + a^2*d)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)))/(a^2*b^2*c^3 - 2*a^3*b*c^2*d + a^4*c*d^2 + (a*b^3*c^3 - 2*a^2*b^2*c^2*d + a^3*b*c*d^2)*x^2), 1/2*(2*(b^2*c^2 - a*b*c*d)*sqrt(b*x^2 + a)*x + (a*b*d*x^2 + a^2*d)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a))/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x))/(a^2*b^2*c^3 - 2*a^3*b*c^2*d + a^4*c*d^2 + (a*b^3*c^3 - 2*a^2*b^2*c^2*d + a^3*b*c*d^2)*x^2)]

giac [A] time = 0.62, size = 107, normalized size = 1.35

$$-\frac{\sqrt{b}d \arctan\left(-\frac{(\sqrt{bx-\sqrt{bx^2+a}})^2 d+2bc-ad}{2\sqrt{-b^2c^2+abcd}}\right)}{\sqrt{-b^2c^2+abcd}(bc-ad)} + \frac{bx}{(abc-a^2d)\sqrt{bx^2+a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c),x, algorithm="giac")

[Out] $-\sqrt{b}d \arctan\left(\frac{-1/2\left(\sqrt{b}x - \sqrt{bx^2+a}\right)^2d + 2bc - ad}{\sqrt{-b^2c^2 + a^2bcd}}\right) / \left(\sqrt{-b^2c^2 + a^2bcd}\right) + bx / \left(\left(a^2cd - a^2d\right)\sqrt{bx^2+a}\right)$

maple [B] time = 0.02, size = 618, normalized size = 7.82

$$\frac{d \ln \left(\frac{2\sqrt{-cd} \left(x - \frac{\sqrt{-cd}}{d}\right)^b + \frac{2ad-2bc}{d} + 2\sqrt{\frac{ad-bc}{d}} \sqrt{\left(x - \frac{\sqrt{-cd}}{d}\right)^2 + \frac{2\sqrt{-cd} \left(x - \frac{\sqrt{-cd}}{d}\right)^b + \frac{ad-bc}{d}}}{x - \frac{\sqrt{-cd}}{d}} \right)}{2\sqrt{-cd} (ad - bc) \sqrt{\frac{ad-bc}{d}}} + \frac{d \ln \left(\frac{2\sqrt{-cd} \left(x + \frac{\sqrt{-cd}}{d}\right)^b + \frac{2ad-2bc}{d} + 2\sqrt{\frac{ad-bc}{d}} \sqrt{\left(x + \frac{\sqrt{-cd}}{d}\right)^2 + \frac{2\sqrt{-cd} \left(x + \frac{\sqrt{-cd}}{d}\right)^b + \frac{ad-bc}{d}}}{x + \frac{\sqrt{-cd}}{d}} \right)}{2\sqrt{-cd} (ad - bc) \sqrt{\frac{ad-bc}{d}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(3/2)/(d*x^2+c),x)

[Out] $\frac{1/2/(-cd)^{1/2}/(ad-bc)d/((x-(-cd)^{1/2}/d)^2b+2(-cd)^{1/2}(x-(-cd)^{1/2}/d)b/d+(ad-bc)/d)^{1/2}-1/2/(ad-bc)/a/((x-(-cd)^{1/2}/d)^2b+2(-cd)^{1/2}(x-(-cd)^{1/2}/d)b/d+(ad-bc)/d)^{1/2}*x*b-1/2/(-cd)^{1/2}/(ad-bc)d/((ad-bc)/d)^{1/2}*\ln((2(-cd)^{1/2}(x-(-cd)^{1/2}/d)b/d+2(ad-bc)/d+2((ad-bc)/d)^{1/2}*((x-(-cd)^{1/2}/d)^2b+2(-cd)^{1/2}(x-(-cd)^{1/2}/d)b/d+(ad-bc)/d)^{1/2})/(x-(-cd)^{1/2}/d))-1/2/(-cd)^{1/2}/(ad-bc)d/((x+(-cd)^{1/2}/d)^2b-2(-cd)^{1/2}(x+(-cd)^{1/2}/d)b/d+(ad-bc)/d)^{1/2}-1/2/(ad-bc)/a/((x+(-cd)^{1/2}/d)^2b-2(-cd)^{1/2}(x+(-cd)^{1/2}/d)b/d+(ad-bc)/d)^{1/2}*x*b+1/2/(-cd)^{1/2}/(ad-bc)d/((ad-bc)/d)^{1/2}*\ln((-2(-cd)^{1/2}(x+(-cd)^{1/2}/d)b/d+2(ad-bc)/d+2((ad-bc)/d)^{1/2}*((x+(-cd)^{1/2}/d)^2b-2(-cd)^{1/2}(x+(-cd)^{1/2}/d)b/d+(ad-bc)/d)^{1/2})/(x+(-cd)^{1/2}/d))}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+a)^{\frac{3}{2}}(dx^2+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/((b*x^2+a)^(3/2)*(d*x^2+c)),x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2+a)^{3/2}(dx^2+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a+b*x^2)^(3/2)*(c+d*x^2)),x)

[Out] int(1/((a+b*x^2)^(3/2)*(c+d*x^2)),x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a+bx^2)^{\frac{3}{2}}(c+dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x**2+a)**(3/2)/(d*x**2+c),x)
```

```
[Out] Integral(1/((a + b*x**2)**(3/2)*(c + d*x**2)), x)
```

$$3.87 \quad \int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)^2} dx$$

Optimal. Leaf size=143

$$-\frac{d(4bc-ad)\tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}(bc-ad)^{5/2}} + \frac{bx(ad+2bc)}{2ac\sqrt{a+bx^2}(bc-ad)^2} - \frac{dx}{2c\sqrt{a+bx^2}(c+dx^2)(bc-ad)}$$

[Out] $-1/2*d*(-a*d+4*b*c)*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})/c^{(3/2)}/(-a*d+b*c)^{(5/2)}+1/2*b*(a*d+2*b*c)*x/a/c/(-a*d+b*c)^2/(b*x^2+a)^{(1/2)}-1/2*d*x/c/(-a*d+b*c)/(d*x^2+c)/(b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.11, antiderivative size = 143, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {414, 527, 12, 377, 208}

$$-\frac{d(4bc-ad)\tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}(bc-ad)^{5/2}} + \frac{bx(ad+2bc)}{2ac\sqrt{a+bx^2}(bc-ad)^2} - \frac{dx}{2c\sqrt{a+bx^2}(c+dx^2)(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(3/2)*(c + d*x^2)^2), x]

[Out] $(b*(2*b*c + a*d)*x)/(2*a*c*(b*c - a*d)^2*\operatorname{Sqrt}[a + b*x^2]) - (d*x)/(2*c*(b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*(c + d*x^2)) - (d*(4*b*c - a*d)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(2*c^{(3/2)}*(b*c - a*d)^{(5/2)})$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c +

$d*x^n)^{(q+1)}/(a*n*(b*c - a*d)*(p+1)), x] + \text{Dist}[1/(a*n*(b*c - a*d)*(p+1)), \text{Int}[(a + b*x^n)^{(p+1)}*(c + d*x^n)^q*\text{Simp}[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2) + 1)*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, q\}, x] \&\& \text{LtQ}[p, -1]$

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + bx^2)^{3/2} (c + dx^2)^2} dx &= -\frac{dx}{2c(bc - ad)\sqrt{a + bx^2} (c + dx^2)} + \frac{\int \frac{2bc - ad - 2bdx^2}{(a + bx^2)^{3/2} (c + dx^2)} dx}{2c(bc - ad)} \\ &= \frac{b(2bc + ad)x}{2ac(bc - ad)^2\sqrt{a + bx^2}} - \frac{dx}{2c(bc - ad)\sqrt{a + bx^2} (c + dx^2)} - \frac{\int \frac{ad(4bc - ad)}{\sqrt{a + bx^2} (c + dx^2)} dx}{2ac(bc - ad)^2} \\ &= \frac{b(2bc + ad)x}{2ac(bc - ad)^2\sqrt{a + bx^2}} - \frac{dx}{2c(bc - ad)\sqrt{a + bx^2} (c + dx^2)} - \frac{(d(4bc - ad)) \int \frac{1}{\sqrt{a + bx^2}} dx}{2c(bc - ad)^2} \\ &= \frac{b(2bc + ad)x}{2ac(bc - ad)^2\sqrt{a + bx^2}} - \frac{dx}{2c(bc - ad)\sqrt{a + bx^2} (c + dx^2)} - \frac{(d(4bc - ad)) \text{Subst}\left[\int \frac{1}{\sqrt{a + bx^2}} dx\right]}{2c(bc - ad)^2} \\ &= \frac{b(2bc + ad)x}{2ac(bc - ad)^2\sqrt{a + bx^2}} - \frac{dx}{2c(bc - ad)\sqrt{a + bx^2} (c + dx^2)} - \frac{d(4bc - ad) \tanh^{-1}\left(\frac{\sqrt{a + bx^2}}{\sqrt{c + dx^2}}\right)}{2c^3/2(bc - ad)^2} \end{aligned}$$

Mathematica [C] time = 2.67, size = 758, normalized size = 5.30

$$x \left[\frac{24d^2x^4 \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{7/2} {}_3F_2\left(2, 2, \frac{5}{2}; 1, \frac{9}{2}; \frac{(bc-ad)x^2}{c(bx^2+a)}\right)}{c^2} + \frac{48dx^2 \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{7/2} {}_3F_2\left(2, 2, \frac{5}{2}; 1, \frac{9}{2}; \frac{(bc-ad)x^2}{c(bx^2+a)}\right)}{c} + 24 \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{7/2} {}_3F_2\left(2, 2, \frac{5}{2}; 1, \frac{9}{2}; \frac{(bc-ad)x^2}{c(bx^2+a)}\right) \right]$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(3/2)*(c + d*x^2)^2), x]

[Out] $(x*(-2625*\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]) - (5250*d*x^2*\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]))/c - (2310*d^2*x^4*\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}])/c^2 + 70*((b*c - a*d)*x^2)/(c*(a + b*x^2))^{3/2} + (560*d*x^2*((b*c - a*d)*x^2)/(c*(a + b*x^2))^{3/2})/c + (280*d^2*x^4*((b*c - a*d)*x^2)/(c*(a + b*x^2))^{3/2})/c^2 + 2625*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]] + (5250*d*x^2*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]])/c + (2310*d^2*x^4*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]])/c^2 - (945*(b*c - a*d)*x^2*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]])/(c*(a + b*x^2)) + (2310*d*(-(b*c) + a*d)*x^4*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]])/(c^2*(a + b*x^2)) + (1050*d^2*(-(b*c) + a*d)*x^6*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]])/(c^3*(a + b*x^2)) + 24*((b*c - a*d)*x^2)/(c*(a + b*x^2))^{7/2}*HypergeometricPFQ[{2, 2, 5/2}, {1, 9/2}, ((b*c - a*d)*x^2)/(c*(a + b*x^2))] + (48*d*x^2*((b*c - a*d)*x^2)/(c*(a + b*x^2))^{7/2}*HypergeometricPFQ[{2, 2, 5/2}, {1, 9/2}, ((b*c - a*d)*x^2)/(c*(a + b*x^2))])/c + (24*d^2*x^4*((b*c - a*d)*x^2)/(c*(a + b*x^2))^{7/2}*HypergeometricPFQ[{2, 2, 5/2}, {1, 9/2}, ((b*c - a*d)*x^2)/(c*(a + b*x^2))])/c^2)/(210*c*((b*c - a*d)*x^2)/(c*(a + b*x^2))^{5/2}*(a + b*x^2)^{3/2}*(c + d*x^2))$

fricas [B] time = 1.40, size = 864, normalized size = 6.04

$$\frac{\left(4a^2bc^2d - a^3cd^2 + (4ab^2cd^2 - a^2bd^3)x^4 + (4ab^2c^2d + 3a^2bcd^2 - a^3d^3)x^2\right)\sqrt{bc^2 - acd} \log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4}{8(a^2b^3c^6 - 3a^3b^2c^5d + 3a^4bc^4d^2 - a^5c^3d^3 + (ab^4c^5d - \dots)}\right)}{8(a^2b^3c^6 - 3a^3b^2c^5d + 3a^4bc^4d^2 - a^5c^3d^3 + (ab^4c^5d - \dots)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^2,x, algorithm="fricas")

[Out]
$$\begin{aligned} & [-1/8*((4*a^2*b*c^2*d - a^3*c*d^2 + (4*a*b^2*c*d^2 - a^2*b*d^3)*x^4 + (4*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*x^2)*\sqrt{b*c^2 - a*c*d}*\log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*\sqrt{b*c^2 - a*c*d}*\sqrt{b*x^2 + a}))/d^2*x^4 + 2*c*d*x^2 + c^2)) - 4*((2*b^3*c^3*d - a*b^2*c^2*d^2 - a^2*b*c*d^3)*x^3 + (2*b^3*c^4 - 2*a*b^2*c^3*d + a^2*b*c^2*d^2 - a^3*c*d^3)*x)*\sqrt{b*x^2 + a})/(a^2*b^3*c^6 - 3*a^3*b^2*c^5*d + 3*a^4*b*c^4*d^2 - a^5*c^3*d^3 + (a*b^4*c^5*d - 3*a^2*b^3*c^4*d^2 + 3*a^3*b^2*c^3*d^3 - a^4*b*c^2*d^4)*x^4 + (a*b^4*c^6 - 2*a^2*b^3*c^5*d + 2*a^4*b*c^3*d^3 - a^5*c^2*d^4)*x^2), 1/4*((4*a^2*b*c^2*d - a^3*c*d^2 + (4*a*b^2*c*d^2 - a^2*b*d^3)*x^4 + (4*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*x^2)*\sqrt{-b*c^2 + a*c*d}*\arctan(1/2*\sqrt{-b*c^2 + a*c*d}*((2*b*c - a*d)*x^2 + a*c)*\sqrt{b*x^2 + a}))/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)) + 2*((2*b^3*c^3*d - a*b^2*c^2*d^2 - a^2*b*c*d^3)*x^3 + (2*b^3*c^4 - 2*a*b^2*c^3*d + a^2*b*c^2*d^2 - a^3*c*d^3)*x)*\sqrt{b*x^2 + a})/(a^2*b^3*c^6 - 3*a^3*b^2*c^5*d + 3*a^4*b*c^4*d^2 - a^5*c^3*d^3 + (a*b^4*c^5*d - 3*a^2*b^3*c^4*d^2 + 3*a^3*b^2*c^3*d^3 - a^4*b*c^2*d^4)*x^4 + (a*b^4*c^6 - 2*a^2*b^3*c^5*d + 2*a^4*b*c^3*d^3 - a^5*c^2*d^4)*x^2)] \end{aligned}$$

giac [B] time = 1.90, size = 318, normalized size = 2.22

$$\frac{b^2x}{(ab^2c^2 - 2a^2bcd + a^3d^2)\sqrt{bx^2 + a}} + \frac{(4b^{\frac{3}{2}}cd - a\sqrt{b}d^2) \arctan\left(\frac{(\sqrt{b}x - \sqrt{bx^2 + a})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{2(b^2c^3 - 2abc^2d + a^2cd^2)\sqrt{-b^2c^2 + abcd}} + \frac{2(\sqrt{b}x - \sqrt{bx^2 + a})^2 d + 2bc - ad}{(b^2c^3 - 2abc^2d + a^2cd^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^2,x, algorithm="giac")

[Out]
$$\begin{aligned} & b^2*x/((a*b^2*c^2 - 2*a^2*b*c*d + a^3*d^2)*\sqrt{b*x^2 + a}) + 1/2*(4*b^{(3/2)}*c*d - a*\sqrt{b}*d^2)*\arctan(1/2*((\sqrt{b}*x - \sqrt{b*x^2 + a})^2*d + 2*b*c - a*d)/\sqrt{-b^2*c^2 + a*b*c*d}))/((b^2*c^3 - 2*a*b*c^2*d + a^2*c*d^2)*\sqrt{-b^2*c^2 + a*b*c*d}) + (2*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*b^{(3/2)}*c*d - (\sqrt{b}*x - \sqrt{b*x^2 + a})^2*a*\sqrt{b}*d^2 + a^2*\sqrt{b}*d^2)/((b^2*c^3 - 2*a*b*c^2*d + a^2*c*d^2)*((\sqrt{b}*x - \sqrt{b*x^2 + a})^4*d + 4*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*b*c - 2*(\sqrt{b}*x - \sqrt{b*x^2 + a})^2*a*d + a^2*d)) \end{aligned}$$

maple [B] time = 0.02, size = 1439, normalized size = 10.06

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(3/2)/(d*x^2+c)^2,x)

[Out]
$$\begin{aligned} & 1/4/c/(a*d-b*c)/(x+(-c*d)^{(1/2)}/d)/((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}-3/4/c*(-c*d)^{(1/2)}*b/(a*d-b*c)^2/((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}+3/4*b^2/(a*d-b*c)^2/a/((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)} \end{aligned}$$

$$\begin{aligned} & \frac{1}{d} \sqrt{\frac{a-d-bc}{d}} + \frac{(a-d-bc)}{d} \sqrt{\frac{a-d-bc}{d}} x + \frac{3}{4} \frac{1}{c} (-c*d)^{1/2} \frac{b}{(a*d-b*c)^2} \left(\frac{(a*d-b*c)}{d} \right)^{1/2} \\ & \ln \left(\frac{-2*(-c*d)^{1/2} * (x+(-c*d)^{1/2}/d) * b/d + 2*(a*d-b*c)/d + 2*\left(\frac{(a*d-b*c)}{d}\right)^{1/2} * \left(\frac{x+(-c*d)^{1/2}}{d}\right)^2 * b - 2*(-c*d)^{1/2} * (x+(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d}{(x+(-c*d)^{1/2}/d)} \right) \\ & + \frac{1}{4} \frac{1}{c} \frac{(a*d-b*c)}{a} \left(\frac{x+(-c*d)^{1/2}}{d} \right)^2 * b - 2*(-c*d)^{1/2} * (x+(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d \sqrt{\frac{a-d-bc}{d}} \\ & x + \frac{1}{4} \frac{1}{c} \frac{(a*d-b*c)}{(x-(-c*d)^{1/2}/d)} \left(\frac{x-(-c*d)^{1/2}}{d} \right)^2 * b + 2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d \sqrt{\frac{a-d-bc}{d}} \\ & + \frac{3}{4} \frac{1}{c} (-c*d)^{1/2} \frac{b}{(a*d-b*c)^2} \left(\frac{x-(-c*d)^{1/2}}{d} \right)^2 * b + 2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d \sqrt{\frac{a-d-bc}{d}} \\ & + \frac{3}{4} \frac{1}{c} \frac{(a*d-b*c)}{a} \left(\frac{x-(-c*d)^{1/2}}{d} \right)^2 * b + 2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d \sqrt{\frac{a-d-bc}{d}} \\ & x - \frac{3}{4} \frac{1}{c} (-c*d)^{1/2} \frac{b}{(a*d-b*c)^2} \left(\frac{(a*d-b*c)}{d} \right)^{1/2} \ln \left(\frac{2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + 2*(a*d-b*c)/d + 2*\left(\frac{(a*d-b*c)}{d}\right)^{1/2} * \left(\frac{x-(-c*d)^{1/2}}{d}\right)^2 * b + 2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d}{(x-(-c*d)^{1/2}/d)} \right) \\ & + \frac{1}{4} \frac{1}{c} \frac{(a*d-b*c)}{a} \left(\frac{x-(-c*d)^{1/2}}{d} \right)^2 * b + 2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d \sqrt{\frac{a-d-bc}{d}} \\ & x + \frac{1}{4} \frac{1}{c} \frac{(a*d-b*c)}{(x-(-c*d)^{1/2}/d)} \left(\frac{x-(-c*d)^{1/2}}{d} \right)^2 * b + 2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d \sqrt{\frac{a-d-bc}{d}} \\ & x - \frac{1}{4} \frac{1}{c} (-c*d)^{1/2} \frac{b}{(a*d-b*c)} \frac{d}{(x-(-c*d)^{1/2}/d)^2 * b + 2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d} \\ & \ln \left(\frac{2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + 2*(a*d-b*c)/d + 2*\left(\frac{(a*d-b*c)}{d}\right)^{1/2} * \left(\frac{x-(-c*d)^{1/2}}{d}\right)^2 * b + 2*(-c*d)^{1/2} * (x-(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d}{(x-(-c*d)^{1/2}/d)} \right) \\ & - \frac{1}{4} \frac{1}{c} (-c*d)^{1/2} \frac{b}{(a*d-b*c)} \frac{d}{(x+(-c*d)^{1/2}/d)^2 * b - 2*(-c*d)^{1/2} * (x+(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d} \\ & + \frac{1}{4} \frac{1}{c} (-c*d)^{1/2} \frac{b}{(a*d-b*c)} \frac{d}{(x+(-c*d)^{1/2}/d)^2 * b - 2*(-c*d)^{1/2} * (x+(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d} \ln \left(\frac{-2*(-c*d)^{1/2} * (x+(-c*d)^{1/2}/d) * b/d + 2*(a*d-b*c)/d + 2*\left(\frac{(a*d-b*c)}{d}\right)^{1/2} * \left(\frac{x+(-c*d)^{1/2}}{d}\right)^2 * b - 2*(-c*d)^{1/2} * (x+(-c*d)^{1/2}/d) * b/d + (a*d-b*c)/d}{(x+(-c*d)^{1/2}/d)} \right) \end{aligned}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{2}} (dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(3/2)*(d*x^2 + c)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + a)^{3/2} (dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(3/2)*(c + d*x^2)^2),x)

[Out] int(1/((a + b*x^2)^(3/2)*(c + d*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{3}{2}} (c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(3/2)/(d*x**2+c)**2,x)

[Out] Integral(1/((a + b*x**2)**(3/2)*(c + d*x**2)**2), x)

$$3.88 \quad \int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)^3} dx$$

Optimal. Leaf size=225

$$\frac{3d(a^2d^2 - 4abcd + 8b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}(bc-ad)^{7/2}} + \frac{dx\sqrt{a+bx^2}(4bc-ad)(3ad+2bc)}{8ac^2(c+dx^2)(bc-ad)^3} + \frac{bx(ad+4bc)}{4ac\sqrt{a+bx^2}(c+dx^2)(bc-ad)}$$

[Out] $-3/8*d*(a^2*d^2-4*a*b*c*d+8*b^2*c^2)*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})/c^{(5/2)}/(-a*d+b*c)^{(7/2)}-1/4*d*x/c/(-a*d+b*c)/(d*x^2+c)^2/(b*x^2+a)^{(1/2)}+1/4*b*(a*d+4*b*c)*x/a/c/(-a*d+b*c)^2/(d*x^2+c)/(b*x^2+a)^{(1/2)}+1/8*d*(-a*d+4*b*c)*(3*a*d+2*b*c)*x*(b*x^2+a)^{(1/2)}/a/c^2/(-a*d+b*c)^3/(d*x^2+c)$

Rubi [A] time = 0.24, antiderivative size = 225, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {414, 527, 12, 377, 208}

$$\frac{3d(a^2d^2 - 4abcd + 8b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}(bc-ad)^{7/2}} + \frac{dx\sqrt{a+bx^2}(4bc-ad)(3ad+2bc)}{8ac^2(c+dx^2)(bc-ad)^3} + \frac{bx(ad+4bc)}{4ac\sqrt{a+bx^2}(c+dx^2)(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(3/2)*(c + d*x^2)^3), x]

[Out] $-(d*x)/(4*c*(b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*(c + d*x^2)^2) + (b*(4*b*c + a*d)*x)/(4*a*c*(b*c - a*d)^2*\operatorname{Sqrt}[a + b*x^2]*(c + d*x^2)) + (d*(4*b*c - a*d)*(2*b*c + 3*a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(8*a*c^2*(b*c - a*d)^3*(c + d*x^2)) - (3*d*(8*b^2*c^2 - 4*a*b*c*d + a^2*d^2)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(8*c^{(5/2)}*(b*c - a*d)^{(7/2)})$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rubi steps

$$\int \frac{1}{(a + bx^2)^{3/2} (c + dx^2)^3} dx = -\frac{dx}{4c(bc - ad)\sqrt{a + bx^2} (c + dx^2)^2} + \frac{\int \frac{4bc - 3ad - 4bdx^2}{(a + bx^2)^{3/2} (c + dx^2)^2} dx}{4c(bc - ad)}$$

$$= -\frac{dx}{4c(bc - ad)\sqrt{a + bx^2} (c + dx^2)^2} + \frac{b(4bc + ad)x}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)} - \frac{\int \frac{ad(8bc - ad)}{(a + bx^2)^{3/2} (c + dx^2)^2} dx}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)}$$

$$= -\frac{dx}{4c(bc - ad)\sqrt{a + bx^2} (c + dx^2)^2} + \frac{b(4bc + ad)x}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)} + \frac{d(4bc - ad)}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)}$$

$$= -\frac{dx}{4c(bc - ad)\sqrt{a + bx^2} (c + dx^2)^2} + \frac{b(4bc + ad)x}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)} + \frac{d(4bc - ad)}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)}$$

$$= -\frac{dx}{4c(bc - ad)\sqrt{a + bx^2} (c + dx^2)^2} + \frac{b(4bc + ad)x}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)} + \frac{d(4bc - ad)}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)}$$

$$= -\frac{dx}{4c(bc - ad)\sqrt{a + bx^2} (c + dx^2)^2} + \frac{b(4bc + ad)x}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)} + \frac{d(4bc - ad)}{4ac(bc - ad)^2\sqrt{a + bx^2} (c + dx^2)}$$

Mathematica [C] time = 5.02, size = 1392, normalized size = 6.19

result too large to display

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(3/2)*(c + d*x^2)^3), x]

[Out] (x*(-108045*sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]) - (324135*d*x^2*sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))])/c - (324135*d^2*x^4*sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))])/c^2 - (103320*d^3*x^6*sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))])/c^3 + 42735*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(3/2) + (128205*d*x^2*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(3/2))/c + (139545*d^2*x^4*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(3/2))/c^2 + (46200*d^3*x^6*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(3/2))/c^3 - 3864*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2) - (4032*d*x^2*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2))/c - (4032*d^2*x^4*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2))/c^2 - (1344*d^3*x^6*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2))/c^3 + 108045*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]] + (324135*d*x^2*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/c + (324135*d^2*x^4*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/c^2 + (103320*d^3*x^6*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/c^3 + (8505*(b*c - a*d)^2*x^4*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/c^4

$$\begin{aligned} & 2))]]/(c^2*(a + b*x^2)^2) + (17955*d*(b*c - a*d)^2*x^6*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/(c^3*(a + b*x^2)^2) + (21735*d^2*(b*c - a*d)^2*x^8*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/(c^4*(a + b*x^2)^2) + (7560*d^3*(b*c - a*d)^2*x^10*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/(c^5*(a + b*x^2)^2) - (78750*(b*c - a*d)*x^2*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/(c*(a + b*x^2)) + (236250*d*(-(b*c) + a*d)*x^4*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/(c^2*(a + b*x^2)) + (247590*d^2*(-(b*c) + a*d)*x^6*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/(c^3*(a + b*x^2)) + (80640*d^3*(-(b*c) + a*d)*x^8*ArcTanh[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]])/(c^4*(a + b*x^2)) + 64*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(9/2)*HypergeometricPFQ[{2, 2, 2, 5/2}, {1, 1, 11/2}, ((b*c - a*d)*x^2)/(c*(a + b*x^2))] + (192*d*x^2*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(9/2)*HypergeometricPFQ[{2, 2, 2, 5/2}, {1, 1, 11/2}, ((b*c - a*d)*x^2)/(c*(a + b*x^2))])/c + (192*d^2*x^4*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(9/2)*HypergeometricPFQ[{2, 2, 2, 5/2}, {1, 1, 11/2}, ((b*c - a*d)*x^2)/(c*(a + b*x^2))])/c^2 + (64*d^3*x^6*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(9/2)*HypergeometricPFQ[{2, 2, 2, 5/2}, {1, 1, 11/2}, ((b*c - a*d)*x^2)/(c*(a + b*x^2))])/c^3)/(2520*c*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(7/2)*(a + b*x^2)^(3/2)*(c + d*x^2)^2) \end{aligned}$$

fricas [B] time = 2.78, size = 1482, normalized size = 6.59

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^3,x, algorithm="fricas")

[Out]
$$\begin{aligned} & [-1/32*(3*(8*a^2*b^2*c^4*d - 4*a^3*b*c^3*d^2 + a^4*c^2*d^3 + (8*a*b^3*c^2*d^3 - 4*a^2*b^2*c*d^4 + a^3*b*d^5)*x^6 + (16*a*b^3*c^3*d^2 - 2*a^3*b*c*d^4 + a^4*d^5)*x^4 + (8*a*b^3*c^4*d + 12*a^2*b^2*c^3*d^2 - 7*a^3*b*c^2*d^3 + 2*a^4*c*d^4)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) - 4*((8*b^4*c^4*d^2 + 2*a*b^3*c^3*d^3 - 13*a^2*b^2*c^2*d^4 + 3*a^3*b*c*d^5)*x^5 + (16*b^4*c^5*d - 4*a*b^3*c^4*d^2 - 7*a^2*b^2*c^3*d^3 - 8*a^3*b*c^2*d^4 + 3*a^4*c*d^5)*x^3 + (8*b^4*c^6 - 8*a*b^3*c^5*d + 12*a^2*b^2*c^4*d^2 - 17*a^3*b*c^3*d^3 + 5*a^4*c^2*d^4)*x)*sqrt(b*x^2 + a))/(a^2*b^4*c^9 - 4*a^3*b^3*c^8*d + 6*a^4*b^2*c^7*d^2 - 4*a^5*b*c^6*d^3 + a^6*c^5*d^4 + (a*b^5*c^7*d^2 - 4*a^2*b^4*c^6*d^3 + 6*a^3*b^3*c^5*d^4 - 4*a^4*b^2*c^4*d^5 + a^5*b*c^3*d^6)*x^6 + (2*a*b^5*c^8*d - 7*a^2*b^4*c^7*d^2 + 8*a^3*b^3*c^6*d^3 - 2*a^4*b^2*c^5*d^4 - 2*a^5*b*c^4*d^5 + a^6*c^3*d^6)*x^4 + (a*b^5*c^9 - 2*a^2*b^4*c^8*d - 2*a^3*b^3*c^7*d^2 + 8*a^4*b^2*c^6*d^3 - 7*a^5*b*c^5*d^4 + 2*a^6*c^4*d^5)*x^2), 1/16*(3*(8*a^2*b^2*c^4*d - 4*a^3*b*c^3*d^2 + a^4*c^2*d^3 + (8*a*b^3*c^2*d^3 - 4*a^2*b^2*c*d^4 + a^3*b*d^5)*x^6 + (16*a*b^3*c^3*d^2 - 2*a^3*b*c*d^4 + a^4*d^5)*x^4 + (8*a*b^3*c^4*d + 12*a^2*b^2*c^3*d^2 - 7*a^3*b*c^2*d^3 + 2*a^4*c*d^4)*x^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)) + 2*((8*b^4*c^4*d^2 + 2*a*b^3*c^3*d^3 - 13*a^2*b^2*c^2*d^4 + 3*a^3*b*c*d^5)*x^5 + (16*b^4*c^5*d - 4*a*b^3*c^4*d^2 - 7*a^2*b^2*c^3*d^3 - 8*a^3*b*c^2*d^4 + 3*a^4*c*d^5)*x^3 + (8*b^4*c^6 - 8*a*b^3*c^5*d + 12*a^2*b^2*c^4*d^2 - 17*a^3*b*c^3*d^3 + 5*a^4*c^2*d^4)*x)*sqrt(b*x^2 + a))/(a^2*b^4*c^9 - 4*a^3*b^3*c^8*d + 6*a^4*b^2*c^7*d^2 - 4*a^5*b*c^6*d^3 + a^6*c^5*d^4 + (a*b^5*c^7*d^2 - 4*a^2*b^4*c^6*d^3 + 6*a^3*b^3*c^5*d^4 - 4*a^4*b^2*c^4*d^5 + a^5*b*c^3*d^6)*x^6 + (2*a*b^5*c^8*d - 7*a^2*b^4*c^7*d^2 + 8*a^3*b^3*c^6*d^3 - 2*a^4*b^2*c^5*d^4 - 2*a^5*b*c^4*d^5 + a^6*c^3*d^6)*x^4 + (a*b^5*c^9 - 2*a^2*b^4*c^8*d - 2*a^3*b^3*c^7*d^2 + 8*a^4*b^2*c^6*d^3 - 7*a^5*b*c^5*d^4 + 2*a^6*c^4*d^5)*x^2)] \end{aligned}$$

giac [B] time = 2.70, size = 643, normalized size = 2.86

$$\frac{b^3 x}{(ab^3 c^3 - 3a^2 b^2 c^2 d + 3a^3 b c d^2 - a^4 d^3) \sqrt{bx^2 + a}} + \frac{3 \left(8b^{\frac{5}{2}} c^2 d - 4ab^{\frac{3}{2}} c d^2 + a^2 \sqrt{b} d^3 \right) \arctan \left(\frac{(\sqrt{bx} - \sqrt{bx^2 + a})^2 d + 2b}{2\sqrt{-b^2 c^2 + abcd}} \right)}{8(b^3 c^5 - 3ab^2 c^4 d + 3a^2 b c^3 d^2 - a^3 c^2 d^3) \sqrt{-b^2 c^2 + abcd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^3,x, algorithm="giac")

[Out] $b^3 x / ((a b^3 c^3 - 3 a^2 b^2 c^2 d + 3 a^3 b c d^2 - a^4 d^3) \sqrt{b x^2 + a}) + 3/8 (8 b^{5/2} c^2 d - 4 a b^{3/2} c d^2 + a^2 \sqrt{b} d^3) \arctan(1/2 ((\sqrt{b} x - \sqrt{b x^2 + a})^2 d + 2 b c - a d) / \sqrt{-b^2 c^2 + a b c d}) / ((b^3 c^5 - 3 a b^2 c^4 d + 3 a^2 b c^3 d^2 - a^3 c^2 d^3) \sqrt{-b^2 c^2 + a b c d}) + 1/4 (16 (\sqrt{b} x - \sqrt{b x^2 + a})^6 b^{5/2} c^2 d^2 - 12 (\sqrt{b} x - \sqrt{b x^2 + a})^6 a b^{3/2} c d^3 + 3 (\sqrt{b} x - \sqrt{b x^2 + a})^6 a^2 \sqrt{b} d^4 + 80 (\sqrt{b} x - \sqrt{b x^2 + a})^4 b^{7/2} c^3 d - 104 (\sqrt{b} x - \sqrt{b x^2 + a})^4 a b^{5/2} c^2 d^2 + 54 (\sqrt{b} x - \sqrt{b x^2 + a})^4 a^2 b^{3/2} c d^3 - 9 (\sqrt{b} x - \sqrt{b x^2 + a})^4 a^3 \sqrt{b} d^4 + 64 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a^2 b^{5/2} c^2 d^2 - 52 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a^3 b^{3/2} c d^3 + 9 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a^4 \sqrt{b} d^4 + 10 a^4 b^{3/2} c d^3 - 3 a^5 \sqrt{b} d^4) / ((b^3 c^5 - 3 a b^2 c^4 d + 3 a^2 b c^3 d^2 - a^3 c^2 d^3) ((\sqrt{b} x - \sqrt{b x^2 + a})^4 d + 4 (\sqrt{b} x - \sqrt{b x^2 + a})^2 b c - 2 (\sqrt{b} x - \sqrt{b x^2 + a})^2 a d + a^2 d^2)$

maple [B] time = 0.03, size = 2919, normalized size = 12.97

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(3/2)/(d*x^2+c)^3,x)

[Out] $3/16/c^2/(a*d-b*c)/(x-(-c*d)^{1/2}/d)/((x-(-c*d)^{1/2}/d)^2*b+2*(-c*d)^{1/2}/d)*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}+3/16/c^2/(a*d-b*c)/(x+(-c*d)^{1/2}/d)/((x+(-c*d)^{1/2}/d)^2*b-2*(-c*d)^{1/2}/d)*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}-1/16/(-c*d)^{1/2}/c/(a*d-b*c)/(x+(-c*d)^{1/2}/d)^2/((x+(-c*d)^{1/2}/d)^2*b-2*(-c*d)^{1/2}/d)*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}-5/16/c*b/(a*d-b*c)^2/(x+(-c*d)^{1/2}/d)/((x+(-c*d)^{1/2}/d)^2*b-2*(-c*d)^{1/2}/d)*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}-3/16/(-c*d)^{1/2}/c^2/(a*d-b*c)*d/((x+(-c*d)^{1/2}/d)^2*b-2*(-c*d)^{1/2}/d)*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}-9/16/c^2*(-c*d)^{1/2}*b/(a*d-b*c)^2/((x+(-c*d)^{1/2}/d)^2*b-2*(-c*d)^{1/2}/d)*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}+3/16/(-c*d)^{1/2}/c^2/(a*d-b*c)*d/((x-(-c*d)^{1/2}/d)^2*b+2*(-c*d)^{1/2}/d)*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}+15/16/(-c*d)^{1/2}*d*b^2/(a*d-b*c)^3/((x-(-c*d)^{1/2}/d)^2*b+2*(-c*d)^{1/2}/d)*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}-15/16*b^3/(a*d-b*c)^3/a/((x-(-c*d)^{1/2}/d)^2*b+2*(-c*d)^{1/2}/d)*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}+9/16/c^2*(-c*d)^{1/2}*b/(a*d-b*c)^2/((x-(-c*d)^{1/2}/d)^2*b+2*(-c*d)^{1/2}/d)*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}-15/16/(-c*d)^{1/2}*d*b^2/(a*d-b*c)^3/((x+(-c*d)^{1/2}/d)^2*b-2*(-c*d)^{1/2}/d)*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}-15/16*b^3/(a*d-b*c)^3/a/((x+(-c*d)^{1/2}/d)^2*b-2*(-c*d)^{1/2}/d)*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2}*x-3/16/(-c*d)^{1/2}/c*d*b/(a*d-b*c)^2/((a*d-b*c)/d)^{1/2}*ln((2*(-c*d)^{1/2}/d)*(x-(-c*d)^{1/2}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{1/2}*((x-(-c*d)^{1/2}/d)^2*b+2*(-c*d)^{1/2}/d)*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d^{1/2})/(x-(-c*d)^{1/2}/d))+3/16/(-c*d)^{1/2}/c*d*b/(a*d-b*c)^2/((a*d-b*c)/d)^{1/2}*ln((-2*(-c*d)^{1/2}/d)*(x+(-c*d)^{1/2}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{1/2}*((x+(-c*d)^{1/2}/d)^2*b-$

$$2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}/(x+(-c*d)^{(1/2)}/d)+1/16/(-c*d)^{(1/2)}/c/(a*d-b*c)/(x-(-c*d)^{(1/2)}/d)^2/((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}-5/16/c*b/(a*d-b*c)^2/(x-(-c*d)^{(1/2)}/d)/((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}+15/16/(-c*d)^{(1/2)}*d*b^2/(a*d-b*c)^3/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))-1/4/c*b^2/(a*d-b*c)^2/a/((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x-3/16/(-c*d)^{(1/2)}/c*d*b/(a*d-b*c)^2/((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}+9/16/c^2*(-c*d)^{(1/2)}*b/(a*d-b*c)^2/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))+3/16/c^2/(a*d-b*c)/a/((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x*b+3/16/c^2/(a*d-b*c)/a/((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x*b-3/16/(-c*d)^{(1/2)}/c^2/(a*d-b*c)*d/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))+3/16/(-c*d)^{(1/2)}/c^2/(a*d-b*c)*d/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))-9/16/c^2*(-c*d)^{(1/2)}*b/(a*d-b*c)^2/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))-15/16/(-c*d)^{(1/2)}*d*b^2/(a*d-b*c)^3/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))-1/4/c*b^2/(a*d-b*c)^2/a/((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x+3/16/(-c*d)^{(1/2)}/c*d*b/(a*d-b*c)^2/((x-(-c*d)^{(1/2)}/d)^2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{2}}(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^3,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(3/2)*(d*x^2 + c)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{3/2}(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(3/2)*(c + d*x^2)^3),x)

[Out] int(1/((a + b*x^2)^(3/2)*(c + d*x^2)^3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.


```
[In] integrate(1/(b*x**2+a)**(3/2)/(d*x**2+c)**3,x)
```

```
[Out] Timed out
```

$$3.89 \quad \int \frac{(c+dx^2)^4}{(a+bx^2)^{5/2}} dx$$

Optimal. Leaf size=255

$$\frac{x(c+dx^2)^2(bc-ad)(7ad+2bc)}{3a^2b^2\sqrt{a+bx^2}} + \frac{d^2(35a^2d^2-80abcd+48b^2c^2)\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{8b^{9/2}} - \frac{dx\sqrt{a+bx^2}(c+dx^2)(-35a^2d^2+24abcd+8b^2c^2)}{12a^2b^3}$$

[Out] 1/3*(-a*d+b*c)*x*(d*x^2+c)^3/a/b/(b*x^2+a)^(3/2)+1/8*d^2*(35*a^2*d^2-80*a*b*c*d+48*b^2*c^2)*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(9/2)+1/3*(-a*d+b*c)*(7*a*d+2*b*c)*x*(d*x^2+c)^2/a^2/b^2/(b*x^2+a)^(1/2)-1/24*d*(105*a^3*d^3-170*a^2*b*c*d^2+40*a*b^2*c^2*d+16*b^3*c^3)*x*(b*x^2+a)^(1/2)/a^2/b^4-1/12*d*(-35*a^2*d^2+24*a*b*c*d+8*b^2*c^2)*x*(d*x^2+c)*(b*x^2+a)^(1/2)/a^2/b^3

Rubi [A] time = 0.24, antiderivative size = 255, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {413, 526, 528, 388, 217, 206}

$$\frac{dx\sqrt{a+bx^2}(c+dx^2)(-35a^2d^2+24abcd+8b^2c^2)}{12a^2b^3} - \frac{dx\sqrt{a+bx^2}(-170a^2bcd^2+105a^3d^3+40ab^2c^2d+16b^3c^3)}{24a^2b^4}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^4/(a + b*x^2)^(5/2), x]

[Out] -(d*(16*b^3*c^3 + 40*a*b^2*c^2*d - 170*a^2*b*c*d^2 + 105*a^3*d^3)*x*sqrt[a + b*x^2])/(24*a^2*b^4) - (d*(8*b^2*c^2 + 24*a*b*c*d - 35*a^2*d^2)*x*sqrt[a + b*x^2]*(c + d*x^2))/(12*a^2*b^3) + ((b*c - a*d)*(2*b*c + 7*a*d)*x*(c + d*x^2)^2)/(3*a^2*b^2*sqrt[a + b*x^2]) + ((b*c - a*d)*x*(c + d*x^2)^3)/(3*a*b*(a + b*x^2)^(3/2)) + (d^2*(48*b^2*c^2 - 80*a*b*c*d + 35*a^2*d^2)*ArcTanh[(sqrt[b]*x)/sqrt[a + b*x^2]])/(8*b^(9/2))

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+q)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,

0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 526

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rule 528

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^4}{(a + bx^2)^{5/2}} dx &= \frac{(bc - ad)x(c + dx^2)^3}{3ab(a + bx^2)^{3/2}} + \frac{\int \frac{(c + dx^2)^2(c(2bc + ad) - d(4bc - 7ad)x^2)}{(a + bx^2)^{3/2}} dx}{3ab} \\ &= \frac{(bc - ad)(2bc + 7ad)x(c + dx^2)^2}{3a^2b^2\sqrt{a + bx^2}} + \frac{(bc - ad)x(c + dx^2)^3}{3ab(a + bx^2)^{3/2}} - \frac{\int \frac{(c + dx^2)(acd(4bc - 7ad) + d(8b^2c^2 + 2ad^2))}{\sqrt{a + bx^2}} dx}{3a^2b^2} \\ &= -\frac{d(8b^2c^2 + 24abcd - 35a^2d^2)x\sqrt{a + bx^2}(c + dx^2)}{12a^2b^3} + \frac{(bc - ad)(2bc + 7ad)x(c + dx^2)^2}{3a^2b^2\sqrt{a + bx^2}} \\ &= -\frac{d(16b^3c^3 + 40ab^2c^2d - 170a^2bcd^2 + 105a^3d^3)x\sqrt{a + bx^2}}{24a^2b^4} - \frac{d(8b^2c^2 + 24abcd - 35a^2d^2)}{12a^2b^2\sqrt{a + bx^2}} \\ &= -\frac{d(16b^3c^3 + 40ab^2c^2d - 170a^2bcd^2 + 105a^3d^3)x\sqrt{a + bx^2}}{24a^2b^4} - \frac{d(8b^2c^2 + 24abcd - 35a^2d^2)}{12a^2b^2\sqrt{a + bx^2}} \\ &= -\frac{d(16b^3c^3 + 40ab^2c^2d - 170a^2bcd^2 + 105a^3d^3)x\sqrt{a + bx^2}}{24a^2b^4} - \frac{d(8b^2c^2 + 24abcd - 35a^2d^2)}{12a^2b^2\sqrt{a + bx^2}} \end{aligned}$$

Mathematica [A] time = 5.17, size = 157, normalized size = 0.62

$$\frac{x\sqrt{a + bx^2} \left(\frac{16(bc - ad)^3(5ad + bc)}{a^2(a + bx^2)} + 3d^3(16bc - 11ad) + \frac{8(bc - ad)^4}{a(a + bx^2)^2} + 6bd^4x^2 \right)}{24b^4} + \frac{d^2(35a^2d^2 - 80abcd + 48b^2c^2) \log\left(\sqrt{a + bx^2}\right)}{8b^9/2}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^4/(a + b*x^2)^(5/2), x]

[Out] (x*Sqrt[a + b*x^2]*(3*d^3*(16*b*c - 11*a*d) + 6*b*d^4*x^2 + (8*(b*c - a*d)^4)/(a*(a + b*x^2)^2) + (16*(b*c - a*d)^3*(b*c + 5*a*d))/(a^2*(a + b*x^2))))

$$\frac{/(24*b^4) + (d^2*(48*b^2*c^2 - 80*a*b*c*d + 35*a^2*d^2)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(8*b^(9/2))$$

fricas [A] time = 1.44, size = 684, normalized size = 2.68

$$\frac{3(48a^4b^2c^2d^2 - 80a^5bcd^3 + 35a^6d^4 + (48a^2b^4c^2d^2 - 80a^3b^3cd^3 + 35a^4b^2d^4)x^4 + 2(48a^3b^3c^2d^2 - 80a^4b^2cd^3 - 35a^5b^2d^4)x^3 + 3(8a^4b^2c^2d^2 - 80a^5b^2cd^3 + 35a^6d^4)x^2 + 2(48a^3b^3c^2d^2 - 80a^4b^2cd^3 + 35a^5b^2d^4)x + 4(4b^8c^4 + 8ab^7c^3d - 48a^2b^6c^2d^2 + 80a^3b^5cd^3 - 35a^4b^4d^4)}{24(bx^2 + a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^(5/2),x, algorithm="fricas")

[Out] [1/48*(3*(48*a^4*b^2*c^2*d^2 - 80*a^5*b*c*d^3 + 35*a^6*d^4 + (48*a^2*b^4*c^2*d^2 - 80*a^3*b^3*c*d^3 + 35*a^4*b^2*d^4)*x^4 + 2*(48*a^3*b^3*c^2*d^2 - 80*a^4*b^2*c*d^3 + 35*a^5*b*d^4)*x^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a))*sqrt(b)*x - a) + 2*(6*a^2*b^4*d^4*x^7 + 3*(16*a^2*b^4*c*d^3 - 7*a^3*b^3*d^4)*x^5 + 4*(4*b^6*c^4 + 8*a*b^5*c^3*d - 48*a^2*b^4*c^2*d^2 + 80*a^3*b^3*c*d^3 - 35*a^4*b^2*d^4)*x^3 + 3*(8*a*b^5*c^4 - 48*a^3*b^3*c^2*d^2 + 80*a^4*b^2*c*d^3 - 35*a^5*b*d^4)*x)*sqrt(b*x^2 + a))/(a^2*b^7*x^4 + 2*a^3*b^6*x^2 + a^4*b^5), -1/24*(3*(48*a^4*b^2*c^2*d^2 - 80*a^5*b*c*d^3 + 35*a^6*d^4 + (48*a^2*b^4*c^2*d^2 - 80*a^3*b^3*c*d^3 + 35*a^4*b^2*d^4)*x^4 + 2*(48*a^3*b^3*c^2*d^2 - 80*a^4*b^2*c*d^3 + 35*a^5*b*d^4)*x^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (6*a^2*b^4*d^4*x^7 + 3*(16*a^2*b^4*c*d^3 - 7*a^3*b^3*d^4)*x^5 + 4*(4*b^6*c^4 + 8*a*b^5*c^3*d - 48*a^2*b^4*c^2*d^2 + 80*a^3*b^3*c*d^3 - 35*a^4*b^2*d^4)*x^3 + 3*(8*a*b^5*c^4 - 48*a^3*b^3*c^2*d^2 + 80*a^4*b^2*c*d^3 - 35*a^5*b*d^4)*x)*sqrt(b*x^2 + a))/(a^2*b^7*x^4 + 2*a^3*b^6*x^2 + a^4*b^5)]

giac [A] time = 0.68, size = 237, normalized size = 0.93

$$\frac{\left(\left(3\left(\frac{2d^4x^2}{b} + \frac{16a^2b^6cd^3-7a^3b^5d^4}{a^2b^7}\right)x^2 + \frac{4(4b^8c^4+8ab^7c^3d-48a^2b^6c^2d^2+80a^3b^5cd^3-35a^4b^4d^4)}{a^2b^7}\right)x^2 + \frac{3(8ab^7c^4-48a^3b^5c^2d^2+80a^4b^4cd^3-35a^5b^2d^4)}{a^2b^7}\right)}{24(bx^2 + a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^(5/2),x, algorithm="giac")

[Out] 1/24*((3*(2*d^4*x^2/b + (16*a^2*b^6*c*d^3 - 7*a^3*b^5*d^4)/(a^2*b^7))*x^2 + 4*(4*b^8*c^4 + 8*a*b^7*c^3*d - 48*a^2*b^6*c^2*d^2 + 80*a^3*b^5*c*d^3 - 35*a^4*b^4*d^4)/(a^2*b^7))*x^2 + 3*(8*a*b^7*c^4 - 48*a^3*b^5*c^2*d^2 + 80*a^4*b^4*c*d^3 - 35*a^5*b^3*d^4)/(a^2*b^7))*x/(b*x^2 + a)^(3/2) - 1/8*(48*b^2*c^2*d^2 - 80*a*b*c*d^3 + 35*a^2*d^4)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(9/2)

maple [A] time = 0.02, size = 351, normalized size = 1.38

$$\frac{d^4x^7}{4(bx^2 + a)^{\frac{3}{2}}b} - \frac{7ad^4x^5}{8(bx^2 + a)^{\frac{3}{2}}b^2} + \frac{2cd^3x^5}{(bx^2 + a)^{\frac{3}{2}}b} - \frac{35a^2d^4x^3}{24(bx^2 + a)^{\frac{3}{2}}b^3} + \frac{10acd^3x^3}{3(bx^2 + a)^{\frac{3}{2}}b^2} - \frac{2c^2d^2x^3}{(bx^2 + a)^{\frac{3}{2}}b} + \frac{c^4x}{3(bx^2 + a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^4/(b*x^2+a)^(5/2),x)

[Out] 1/4*d^4*x^7/b/(b*x^2+a)^(3/2)-7/8*d^4*a/b^2*x^5/(b*x^2+a)^(3/2)-35/24*d^4*a^2/b^3*x^3/(b*x^2+a)^(3/2)-35/8*d^4*a^2/b^4*x/(b*x^2+a)^(1/2)+35/8*d^4*a^2/b^(9/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))+2*c*d^3*x^5/b/(b*x^2+a)^(3/2)+10/3*c*d^3*a/b^2*x^3/(b*x^2+a)^(3/2)+10*c*d^3*a/b^3*x/(b*x^2+a)^(1/2)-10*c*d^3*a/b

$$\begin{aligned} & \frac{d^4 x^7}{4(bx^2 + a)^{\frac{3}{2}} b} + \frac{2cd^3 x^5}{(bx^2 + a)^{\frac{3}{2}} b} - \frac{7ad^4 x^5}{8(bx^2 + a)^{\frac{3}{2}} b^2} - 2c^2 d^2 x \left(\frac{3x^2}{(bx^2 + a)^{\frac{3}{2}} b} + \frac{2a}{(bx^2 + a)^{\frac{3}{2}} b^2} \right) + \frac{10acd^3 x \left(\frac{3x^2}{(bx^2 + a)^{\frac{3}{2}} b} + \frac{2a}{(bx^2 + a)^{\frac{3}{2}} b^2} \right)}{3b} \\ & - 6c^2 d^2 / b^2 * x / (bx^2 + a)^{(1/2)} + 6c^2 d^2 / b^{(5/2)} * \ln(b^{(1/2)} * x + (bx^2 + a)^{(1/2)}) - 4 / 3 * c^3 * d / b * x / (bx^2 + a)^{(3/2)} + 4 / 3 * c^3 * d / a / b * x / (bx^2 + a)^{(1/2)} + 1 / 3 * c^4 * x / a / (bx^2 + a)^{(3/2)} + 2 / 3 * c^4 / a^2 * x / (bx^2 + a)^{(1/2)} \end{aligned}$$

maxima [A] time = 1.44, size = 392, normalized size = 1.54

$$\frac{d^4 x^7}{4(bx^2 + a)^{\frac{3}{2}} b} + \frac{2cd^3 x^5}{(bx^2 + a)^{\frac{3}{2}} b} - \frac{7ad^4 x^5}{8(bx^2 + a)^{\frac{3}{2}} b^2} - 2c^2 d^2 x \left(\frac{3x^2}{(bx^2 + a)^{\frac{3}{2}} b} + \frac{2a}{(bx^2 + a)^{\frac{3}{2}} b^2} \right) + \frac{10acd^3 x \left(\frac{3x^2}{(bx^2 + a)^{\frac{3}{2}} b} + \frac{2a}{(bx^2 + a)^{\frac{3}{2}} b^2} \right)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^4/(b*x^2+a)^(5/2),x, algorithm="maxima")

[Out] $\frac{1}{4}d^4x^7/((bx^2 + a)^{(3/2)}b) + 2c*d^3*x^5/((bx^2 + a)^{(3/2)}b) - 7/8 * a*d^4*x^5/((bx^2 + a)^{(3/2)}*b^2) - 2*c^2*d^2*x*(3*x^2/((bx^2 + a)^{(3/2)}*b) + 2*a/((bx^2 + a)^{(3/2)}*b^2)) + 10/3*a*c*d^3*x*(3*x^2/((bx^2 + a)^{(3/2)}*b) + 2*a/((bx^2 + a)^{(3/2)}*b^2))/b - 35/24*a^2*d^4*x*(3*x^2/((bx^2 + a)^{(3/2)}*b) + 2*a/((bx^2 + a)^{(3/2)}*b^2))/b^2 + 2/3*c^4*x/(sqrt(b*x^2 + a)*a^2) + 1/3*c^4*x/((bx^2 + a)^{(3/2)}*a) - 4/3*c^3*d*x/((bx^2 + a)^{(3/2)}*b) + 4/3*c^3*d*x/(sqrt(b*x^2 + a)*a*b) - 2*c^2*d^2*x/(sqrt(b*x^2 + a)*b^2) + 10/3*a*c*d^3*x/(sqrt(b*x^2 + a)*b^3) - 35/24*a^2*d^4*x/(sqrt(b*x^2 + a)*b^4) + 6*c^2*d^2*arcsinh(b*x/sqrt(a*b))/b^(5/2) - 10*a*c*d^3*arcsinh(b*x/sqrt(a*b))/b^(7/2) + 35/8*a^2*d^4*arcsinh(b*x/sqrt(a*b))/b^(9/2)$

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(dx^2 + c)^4}{(bx^2 + a)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^4/(a + b*x^2)^(5/2), x)

[Out] int((c + d*x^2)^4/(a + b*x^2)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + dx^2)^4}{(a + bx^2)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**4/(b*x**2+a)**(5/2), x)

[Out] Integral((c + d*x**2)**4/(a + b*x**2)**(5/2), x)

$$3.90 \quad \int \frac{(c+dx^2)^3}{(a+bx^2)^{5/2}} dx$$

Optimal. Leaf size=172

$$\frac{x(c+dx^2)(bc-ad)(5ad+2bc)}{3a^2b^2\sqrt{a+bx^2}} - \frac{dx\sqrt{a+bx^2}(-15a^2d^2+8abcd+4b^2c^2)}{6a^2b^3} + \frac{d^2(6bc-5ad)\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{2b^{7/2}} + \frac{x(c+dx^2)}{3a^2b^2\sqrt{a+bx^2}}$$

[Out] 1/3*(-a*d+b*c)*x*(d*x^2+c)^2/a/b/(b*x^2+a)^(3/2)+1/2*d^2*(-5*a*d+6*b*c)*arc tanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(7/2)+1/3*(-a*d+b*c)*(5*a*d+2*b*c)*x*(d*x^2+c)/a^2/b^2/(b*x^2+a)^(1/2)-1/6*d*(-15*a^2*d^2+8*a*b*c*d+4*b^2*c^2)*x*(b*x^2+a)^(1/2)/a^2/b^3

Rubi [A] time = 0.16, antiderivative size = 172, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {413, 526, 388, 217, 206}

$$-\frac{dx\sqrt{a+bx^2}(-15a^2d^2+8abcd+4b^2c^2)}{6a^2b^3} + \frac{x(c+dx^2)(bc-ad)(5ad+2bc)}{3a^2b^2\sqrt{a+bx^2}} + \frac{d^2(6bc-5ad)\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a+bx^2}}\right)}{2b^{7/2}} + \frac{x(c+dx^2)}{3a^2b^2\sqrt{a+bx^2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^3/(a + b*x^2)^(5/2), x]

[Out] -(d*(4*b^2*c^2 + 8*a*b*c*d - 15*a^2*d^2)*x*sqrt[a + b*x^2])/(6*a^2*b^3) + ((b*c - a*d)*(2*b*c + 5*a*d)*x*(c + d*x^2))/(3*a^2*b^2*sqrt[a + b*x^2]) + ((b*c - a*d)*x*(c + d*x^2)^2)/(3*a*b*(a + b*x^2)^(3/2)) + (d^2*(6*b*c - 5*a*d)*ArcTanh[(sqrt[b]*x)/sqrt[a + b*x^2]])/(2*b^(7/2))

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+q)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 526

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^3}{(a + bx^2)^{5/2}} dx &= \frac{(bc - ad)x(c + dx^2)^2}{3ab(a + bx^2)^{3/2}} + \frac{\int \frac{(c + dx^2)(c(2bc + ad) - d(2bc - 5ad)x^2)}{(a + bx^2)^{3/2}} dx}{3ab} \\ &= \frac{(bc - ad)(2bc + 5ad)x(c + dx^2)}{3a^2b^2\sqrt{a + bx^2}} + \frac{(bc - ad)x(c + dx^2)^2}{3ab(a + bx^2)^{3/2}} - \frac{\int \frac{acd(2bc - 5ad) + d(4b^2c^2 + 8abcd - 15a^2d^2)}{\sqrt{a + bx^2}} dx}{3a^2b^2} \\ &= -\frac{d(4b^2c^2 + 8abcd - 15a^2d^2)x\sqrt{a + bx^2}}{6a^2b^3} + \frac{(bc - ad)(2bc + 5ad)x(c + dx^2)}{3a^2b^2\sqrt{a + bx^2}} + \frac{(bc - ad)x}{3ab(a + bx^2)^{3/2}} \\ &= -\frac{d(4b^2c^2 + 8abcd - 15a^2d^2)x\sqrt{a + bx^2}}{6a^2b^3} + \frac{(bc - ad)(2bc + 5ad)x(c + dx^2)}{3a^2b^2\sqrt{a + bx^2}} + \frac{(bc - ad)x}{3ab(a + bx^2)^{3/2}} \\ &= -\frac{d(4b^2c^2 + 8abcd - 15a^2d^2)x\sqrt{a + bx^2}}{6a^2b^3} + \frac{(bc - ad)(2bc + 5ad)x(c + dx^2)}{3a^2b^2\sqrt{a + bx^2}} + \frac{(bc - ad)x}{3ab(a + bx^2)^{3/2}} \end{aligned}$$

Mathematica [A] time = 5.10, size = 125, normalized size = 0.73

$$\frac{x \left(3a^2d^3(a + bx^2)^2 + 2(a + bx^2)(bc - ad)^2(7ad + 2bc) + 2a(bc - ad)^3 \right)}{6a^2b^3(a + bx^2)^{3/2}} + \frac{d^2(6bc - 5ad) \log\left(\sqrt{b}\sqrt{a + bx^2} + b\sqrt{a}\right)}{2b^{7/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(c + d*x^2)^3/(a + b*x^2)^(5/2), x]
```

```
[Out] (x*(2*a*(b*c - a*d)^3 + 2*(b*c - a*d)^2*(2*b*c + 7*a*d)*(a + b*x^2) + 3*a^2*d^3*(a + b*x^2)^2)/(6*a^2*b^3*(a + b*x^2)^(3/2)) + (d^2*(6*b*c - 5*a*d)*Log[b*x + Sqrt[b]*Sqrt[a + b*x^2]])/(2*b^(7/2))
```

fricas [A] time = 0.67, size = 486, normalized size = 2.83

$$\left[\frac{3(6a^4bcd^2 - 5a^5d^3 + (6a^2b^3cd^2 - 5a^3b^2d^3)x^4 + 2(6a^3b^2cd^2 - 5a^4bd^3)x^2)\sqrt{b} \log\left(-2bx^2 + 2\sqrt{bx^2 + a}\sqrt{b}\right)}{12(a^2b^3\sqrt{a + bx^2})} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^3/(b*x^2+a)^(5/2), x, algorithm="fricas")
```

```
[Out] [-1/12*(3*(6*a^4*b*c*d^2 - 5*a^5*d^3 + (6*a^2*b^3*c*d^2 - 5*a^3*b^2*d^3)*x^4 + 2*(6*a^3*b^2*c*d^2 - 5*a^4*b*d^3)*x^2)*sqrt(b)*log(-2*b*x^2 + 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) - 2*(3*a^2*b^3*d^3*x^5 + 2*(2*b^5*c^3 + 3*a*b^4*c^2
```

*d - 12*a^2*b^3*c*d^2 + 10*a^3*b^2*d^3)*x^3 + 3*(2*a*b^4*c^3 - 6*a^3*b^2*c*d^2 + 5*a^4*b*d^3)*x)*sqrt(b*x^2 + a))/(a^2*b^6*x^4 + 2*a^3*b^5*x^2 + a^4*b^4), -1/6*(3*(6*a^4*b*c*d^2 - 5*a^5*d^3 + (6*a^2*b^3*c*d^2 - 5*a^3*b^2*d^3)*x^4 + 2*(6*a^3*b^2*c*d^2 - 5*a^4*b*d^3)*x^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (3*a^2*b^3*d^3*x^5 + 2*(2*b^5*c^3 + 3*a*b^4*c^2*d - 12*a^2*b^3*c*d^2 + 10*a^3*b^2*d^3)*x^3 + 3*(2*a*b^4*c^3 - 6*a^3*b^2*c*d^2 + 5*a^4*b*d^3)*x)*sqrt(b*x^2 + a))/(a^2*b^6*x^4 + 2*a^3*b^5*x^2 + a^4*b^4)]

giac [A] time = 0.68, size = 158, normalized size = 0.92

$$\frac{\left(\left(\frac{3d^3x^2}{b} + \frac{2(2b^6c^3+3ab^5c^2d-12a^2b^4cd^2+10a^3b^3d^3)}{a^2b^5}\right)x^2 + \frac{3(2ab^5c^3-6a^3b^3cd^2+5a^4b^2d^3)}{a^2b^5}\right)x}{6(bx^2+a)^{\frac{3}{2}}} - \frac{(6bcd^2-5ad^3)\log\left(\left|-\sqrt{b}x + \sqrt{bx^2+a}\right|\right)}{2b^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^(5/2),x, algorithm="giac")

[Out] 1/6*((3*d^3*x^2/b + 2*(2*b^6*c^3 + 3*a*b^5*c^2*d - 12*a^2*b^4*c*d^2 + 10*a^3*b^3*d^3)/(a^2*b^5))*x^2 + 3*(2*a*b^5*c^3 - 6*a^3*b^3*c*d^2 + 5*a^4*b^2*d^3)/(a^2*b^5))*x/(b*x^2 + a)^(3/2) - 1/2*(6*b*c*d^2 - 5*a*d^3)*log(abs(-sqrt(b)*x + sqrt(b*x^2 + a)))/b^(7/2)

maple [A] time = 0.01, size = 228, normalized size = 1.33

$$\frac{d^3x^5}{2(bx^2+a)^{\frac{3}{2}}b} + \frac{5ad^3x^3}{6(bx^2+a)^{\frac{3}{2}}b^2} - \frac{cd^2x^3}{(bx^2+a)^{\frac{3}{2}}b} + \frac{c^3x}{3(bx^2+a)^{\frac{3}{2}}a} - \frac{c^2dx}{(bx^2+a)^{\frac{3}{2}}b} + \frac{5ad^3x}{2\sqrt{bx^2+a}b^3} + \frac{c^2dx}{\sqrt{bx^2+a}ab} + \dots$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^3/(b*x^2+a)^(5/2),x)

[Out] 1/2*d^3*x^5/b/(b*x^2+a)^(3/2)+5/6*d^3*a/b^2*x^3/(b*x^2+a)^(3/2)+5/2*d^3*a/b^3*x/(b*x^2+a)^(1/2)-5/2*d^3*a/b^(7/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))-c*d^2*x^3/b/(b*x^2+a)^(3/2)-3*c*d^2/b^2*x/(b*x^2+a)^(1/2)+3*c*d^2/b^(5/2)*ln(b^(1/2)*x+(b*x^2+a)^(1/2))-c^2*d/b*x/(b*x^2+a)^(3/2)+c^2*d/a/b*x/(b*x^2+a)^(1/2)+1/3*c^3*x/a/(b*x^2+a)^(3/2)+2/3*c^3/a^2*x/(b*x^2+a)^(1/2)

maxima [A] time = 1.49, size = 254, normalized size = 1.48

$$\frac{d^3x^5}{2(bx^2+a)^{\frac{3}{2}}b} - cd^2x \left(\frac{3x^2}{(bx^2+a)^{\frac{3}{2}}b} + \frac{2a}{(bx^2+a)^{\frac{3}{2}}b^2} \right) + \frac{5ad^3x \left(\frac{3x^2}{(bx^2+a)^{\frac{3}{2}}b} + \frac{2a}{(bx^2+a)^{\frac{3}{2}}b^2} \right)}{6b} + \frac{2c^3x}{3\sqrt{bx^2+a}a^2} + \frac{c^3x}{3(bx^2+a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^3/(b*x^2+a)^(5/2),x, algorithm="maxima")

[Out] 1/2*d^3*x^5/((b*x^2 + a)^(3/2)*b) - c*d^2*x*(3*x^2/((b*x^2 + a)^(3/2)*b) + 2*a/((b*x^2 + a)^(3/2)*b^2)) + 5/6*a*d^3*x*(3*x^2/((b*x^2 + a)^(3/2)*b) + 2*a/((b*x^2 + a)^(3/2)*b^2))/b + 2/3*c^3*x/(sqrt(b*x^2 + a)*a^2) + 1/3*c^3*x/((b*x^2 + a)^(3/2)*a) - c^2*d*x/((b*x^2 + a)^(3/2)*b) + c^2*d*x/(sqrt(b*x^2 + a)*a*b) - c*d^2*x/(sqrt(b*x^2 + a)*b^2) + 5/6*a*d^3*x/(sqrt(b*x^2 + a)*b^3) + 3*c*d^2*arcsinh(b*x/sqrt(a*b))/b^(5/2) - 5/2*a*d^3*arcsinh(b*x/sqrt(a*b))/b^(7/2)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(dx^2 + c)^3}{(bx^2 + a)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^3/(a + b*x^2)^(5/2), x)

[Out] int((c + d*x^2)^3/(a + b*x^2)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + dx^2)^3}{(a + bx^2)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**3/(b*x**2+a)**(5/2), x)

[Out] Integral((c + d*x**2)**3/(a + b*x**2)**(5/2), x)

$$3.91 \quad \int \frac{(c+dx^2)^2}{(a+bx^2)^{5/2}} dx$$

Optimal. Leaf size=105

$$\frac{x(bc-ad)(3ad+2bc)}{3a^2b^2\sqrt{a+bx^2}} + \frac{d^2 \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{b^{5/2}} + \frac{x(c+dx^2)(bc-ad)}{3ab(a+bx^2)^{3/2}}$$

[Out] 1/3*(-a*d+b*c)*x*(d*x^2+c)/a/b/(b*x^2+a)^(3/2)+d^2*arctanh(x*b^(1/2)/(b*x^2+a)^(1/2))/b^(5/2)+1/3*(-a*d+b*c)*(3*a*d+2*b*c)*x/a^2/b^2/(b*x^2+a)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {413, 385, 217, 206}

$$\frac{x(bc-ad)(3ad+2bc)}{3a^2b^2\sqrt{a+bx^2}} + \frac{d^2 \tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a+bx^2}}\right)}{b^{5/2}} + \frac{x(c+dx^2)(bc-ad)}{3ab(a+bx^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^2/(a + b*x^2)^(5/2), x]

[Out] ((b*c - a*d)*(2*b*c + 3*a*d)*x)/(3*a^2*b^2*Sqrt[a + b*x^2]) + ((b*c - a*d)*x*(c + d*x^2))/(3*a*b*(a + b*x^2)^(3/2)) + (d^2*ArcTanh[(Sqrt[b]*x)/Sqrt[a + b*x^2]])/b^(5/2)

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rubi steps

$$\begin{aligned}
\int \frac{(c + dx^2)^2}{(a + bx^2)^{5/2}} dx &= \frac{(bc - ad)x(c + dx^2)}{3ab(a + bx^2)^{3/2}} + \frac{\int \frac{c(2bc + ad) + 3ad^2x^2}{(a + bx^2)^{3/2}} dx}{3ab} \\
&= \frac{(bc - ad)(2bc + 3ad)x}{3a^2b^2\sqrt{a + bx^2}} + \frac{(bc - ad)x(c + dx^2)}{3ab(a + bx^2)^{3/2}} + \frac{d^2 \int \frac{1}{\sqrt{a + bx^2}} dx}{b^2} \\
&= \frac{(bc - ad)(2bc + 3ad)x}{3a^2b^2\sqrt{a + bx^2}} + \frac{(bc - ad)x(c + dx^2)}{3ab(a + bx^2)^{3/2}} + \frac{d^2 \operatorname{Subst}\left(\int \frac{1}{1 - bx^2} dx, x, \frac{x}{\sqrt{a + bx^2}}\right)}{b^2} \\
&= \frac{(bc - ad)(2bc + 3ad)x}{3a^2b^2\sqrt{a + bx^2}} + \frac{(bc - ad)x(c + dx^2)}{3ab(a + bx^2)^{3/2}} + \frac{d^2 \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a + bx^2}}\right)}{b^{5/2}}
\end{aligned}$$

Mathematica [C] time = 4.15, size = 214, normalized size = 2.04

$$\frac{\sqrt{\frac{bx^2}{a} + 1} \left(-16b^3x^6(c + dx^2)^2 {}_3F_2\left(\frac{3}{2}, 2, \frac{7}{2}; 1, \frac{9}{2}; -\frac{bx^2}{a}\right) + \frac{7a^2(15c^2 + 10cdx^2 + 3d^2x^4) \left(\sqrt{-\frac{bx^2(a + bx^2)}{a^2}} (2bx^2 - 3a) + 3a \sin^{-1}\left(\sqrt{-\frac{bx^2}{a}}\right) \right)}{\sqrt{-\frac{bx^2}{a}}} \right)}{168a^3b^2x^3\sqrt{a + bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(c + d*x^2)^2/(a + b*x^2)^(5/2), x]

[Out] (Sqrt[1 + (b*x^2)/a]*((7*a^2*(15*c^2 + 10*c*d*x^2 + 3*d^2*x^4)*(Sqrt[-((b*x^2*(a + b*x^2))/a^2)]*(-3*a + 2*b*x^2) + 3*a*ArcSin[Sqrt[-((b*x^2)/a)]])))/Sqrt[-((b*x^2)/a)] - 32*b^3*x^6*(2*c^2 + 3*c*d*x^2 + d^2*x^4)*Hypergeometric2F1[3/2, 7/2, 9/2, -((b*x^2)/a)] - 16*b^3*x^6*(c + d*x^2)^2*HypergeometricPFQ[{3/2, 2, 7/2}, {1, 9/2}, -((b*x^2)/a)])/(168*a^3*b^2*x^3*Sqrt[a + b*x^2])

fricas [A] time = 0.47, size = 318, normalized size = 3.03

$$\frac{3(a^2b^2d^2x^4 + 2a^3bd^2x^2 + a^4d^2)\sqrt{b} \log(-2bx^2 - 2\sqrt{bx^2 + a}\sqrt{b}x - a) + 2(2(b^4c^2 + ab^3cd - 2a^2b^2d^2)x^3 + \dots)}{6(a^2b^5x^4 + 2a^3b^4x^2 + a^4b^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^(5/2), x, algorithm="fricas")

[Out] [1/6*(3*(a^2*b^2*d^2*x^4 + 2*a^3*b*d^2*x^2 + a^4*d^2)*sqrt(b)*log(-2*b*x^2 - 2*sqrt(b*x^2 + a)*sqrt(b)*x - a) + 2*(2*(b^4*c^2 + a*b^3*c*d - 2*a^2*b^2*d^2)*x^3 + 3*(a*b^3*c^2 - a^3*b*d^2)*x)*sqrt(b*x^2 + a))/(a^2*b^5*x^4 + 2*a^3*b^4*x^2 + a^4*b^3), -1/3*(3*(a^2*b^2*d^2*x^4 + 2*a^3*b*d^2*x^2 + a^4*d^2)*sqrt(-b)*arctan(sqrt(-b)*x/sqrt(b*x^2 + a)) - (2*(b^4*c^2 + a*b^3*c*d - 2*a^2*b^2*d^2)*x^3 + 3*(a*b^3*c^2 - a^3*b*d^2)*x)*sqrt(b*x^2 + a))/(a^2*b^5*x^4 + 2*a^3*b^4*x^2 + a^4*b^3)]

giac [A] time = 0.63, size = 103, normalized size = 0.98

$$\frac{x \left(\frac{2(b^4c^2 + ab^3cd - 2a^2b^2d^2)x^2}{a^2b^3} + \frac{3(ab^3c^2 - a^3bd^2)}{a^2b^3} \right)}{3(bx^2 + a)^{\frac{3}{2}}} - \frac{d^2 \log\left(\left| -\sqrt{b}x + \sqrt{bx^2 + a} \right| \right)}{b^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^(5/2),x, algorithm="giac")

[Out] $\frac{1}{3}x(2(b^4c^2 + ab^3cd - 2a^2b^2d^2)x^2/(a^2b^3) + 3(ab^3c^2 - a^3bd^2)/(a^2b^3))/(bx^2 + a)^{3/2} - d^2\log(\text{abs}(-\sqrt{b}x + \sqrt{bx^2 + a}))/b^{5/2}$

maple [A] time = 0.01, size = 136, normalized size = 1.30

$$-\frac{d^2x^3}{3(bx^2+a)^{\frac{3}{2}}b} + \frac{c^2x}{3(bx^2+a)^{\frac{3}{2}}a} - \frac{2cdx}{3(bx^2+a)^{\frac{3}{2}}b} + \frac{2cdx}{3\sqrt{bx^2+a}ab} + \frac{2c^2x}{3\sqrt{bx^2+a}a^2} - \frac{d^2x}{\sqrt{bx^2+a}b^2} + \frac{d^2\ln(\sqrt{b}x + \sqrt{bx^2+a})}{b^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^2/(b*x^2+a)^(5/2),x)

[Out] $-\frac{1}{3}d^2x^3/b/(bx^2+a)^{3/2} - d^2/b^2x/(bx^2+a)^{1/2} + d^2/b^{5/2}\ln(b^{1/2}x + (bx^2+a)^{1/2}) - \frac{2}{3}cd/bx/(bx^2+a)^{3/2} + \frac{2}{3}cd/a/bx/(bx^2+a)^{1/2} + \frac{1}{3}c^2x/a/(bx^2+a)^{3/2} + \frac{2}{3}c^2/a^2x/(bx^2+a)^{1/2}$

maxima [A] time = 1.39, size = 147, normalized size = 1.40

$$-\frac{1}{3}d^2x\left(\frac{3x^2}{(bx^2+a)^{\frac{3}{2}}b} + \frac{2a}{(bx^2+a)^{\frac{3}{2}}b^2}\right) + \frac{2c^2x}{3\sqrt{bx^2+a}a^2} + \frac{c^2x}{3(bx^2+a)^{\frac{3}{2}}a} - \frac{2cdx}{3(bx^2+a)^{\frac{3}{2}}b} + \frac{2cdx}{3\sqrt{bx^2+a}ab} - \frac{d^2x}{3\sqrt{bx^2+a}b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^2/(b*x^2+a)^(5/2),x, algorithm="maxima")

[Out] $-\frac{1}{3}d^2x(3x^2/((bx^2+a)^{3/2}b) + 2a/((bx^2+a)^{3/2}b^2)) + \frac{2}{3}c^2x/(\sqrt{bx^2+a}a^2) + \frac{1}{3}c^2x/((bx^2+a)^{3/2}a) - \frac{2}{3}cdx/((bx^2+a)^{3/2}b) + \frac{2}{3}cdx/(\sqrt{bx^2+a}ab) - \frac{1}{3}d^2x/(\sqrt{bx^2+a}b^2) + d^2\text{arcsinh}(bx/\sqrt{ab})/b^{5/2}$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(dx^2+c)^2}{(bx^2+a)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^2/(a + b*x^2)^(5/2),x)

[Out] int((c + d*x^2)^2/(a + b*x^2)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + dx^2)^2}{(a + bx^2)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**2/(b*x**2+a)**(5/2),x)

[Out] Integral((c + d*x**2)**2/(a + b*x**2)**(5/2), x)

$$3.92 \quad \int \frac{c+dx^2}{(a+bx^2)^{5/2}} dx$$

Optimal. Leaf size=47

$$\frac{2cx}{3a^2\sqrt{a+bx^2}} + \frac{x(c+dx^2)}{3a(a+bx^2)^{3/2}}$$

[Out] 1/3*x*(d*x^2+c)/a/(b*x^2+a)^(3/2)+2/3*c*x/a^2/(b*x^2+a)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {378, 191}

$$\frac{2cx}{3a^2\sqrt{a+bx^2}} + \frac{x(c+dx^2)}{3a(a+bx^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)/(a + b*x^2)^(5/2), x]

[Out] (2*c*x)/(3*a^2*sqrt[a + b*x^2]) + (x*(c + d*x^2))/(3*a*(a + b*x^2)^(3/2))

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*n*(p + 1)), x] - Dist[(c*q)/(a*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p + q + 1) + 1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{c+dx^2}{(a+bx^2)^{5/2}} dx &= \frac{x(c+dx^2)}{3a(a+bx^2)^{3/2}} + \frac{(2c) \int \frac{1}{(a+bx^2)^{3/2}} dx}{3a} \\ &= \frac{2cx}{3a^2\sqrt{a+bx^2}} + \frac{x(c+dx^2)}{3a(a+bx^2)^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 37, normalized size = 0.79

$$\frac{x(3ac + adx^2 + 2bcx^2)}{3a^2(a+bx^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)/(a + b*x^2)^(5/2), x]

[Out] (x*(3*a*c + 2*b*c*x^2 + a*d*x^2))/(3*a^2*(a + b*x^2)^(3/2))

fricas [A] time = 0.62, size = 54, normalized size = 1.15

$$\frac{((2bc + ad)x^3 + 3acx)\sqrt{bx^2 + a}}{3(a^2b^2x^4 + 2a^3bx^2 + a^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(5/2),x, algorithm="fricas")

[Out] 1/3*((2*b*c + a*d)*x^3 + 3*a*c*x)*sqrt(b*x^2 + a)/(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4)

giac [A] time = 0.62, size = 40, normalized size = 0.85

$$\frac{x\left(\frac{3c}{a} + \frac{(2b^2c+abd)x^2}{a^2b}\right)}{3(bx^2 + a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(5/2),x, algorithm="giac")

[Out] 1/3*x*(3*c/a + (2*b^2*c + a*b*d)*x^2/(a^2*b))/(b*x^2 + a)^(3/2)

maple [A] time = 0.00, size = 34, normalized size = 0.72

$$\frac{(adx^2 + 2bcx^2 + 3ac)x}{3(bx^2 + a)^{\frac{3}{2}}a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)/(b*x^2+a)^(5/2),x)

[Out] 1/3*x*(a*d*x^2+2*b*c*x^2+3*a*c)/(b*x^2+a)^(3/2)/a^2

maxima [A] time = 1.37, size = 68, normalized size = 1.45

$$\frac{2cx}{3\sqrt{bx^2 + a}a^2} + \frac{cx}{3(bx^2 + a)^{\frac{3}{2}}a} - \frac{dx}{3(bx^2 + a)^{\frac{3}{2}}b} + \frac{dx}{3\sqrt{bx^2 + a}ab}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)/(b*x^2+a)^(5/2),x, algorithm="maxima")

[Out] 2/3*c*x/(sqrt(b*x^2 + a)*a^2) + 1/3*c*x/((b*x^2 + a)^(3/2)*a) - 1/3*d*x/((b*x^2 + a)^(3/2)*b) + 1/3*d*x/(sqrt(b*x^2 + a)*a*b)

mupad [B] time = 4.78, size = 33, normalized size = 0.70

$$\frac{3acx + adx^3 + 2bcx^3}{3a^2(bx^2 + a)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)/(a + b*x^2)^(5/2),x)

[Out] (3*a*c*x + a*d*x^3 + 2*b*c*x^3)/(3*a^2*(a + b*x^2)^(3/2))

sympy [B] time = 11.03, size = 144, normalized size = 3.06

$$c \left(\frac{3ax}{3a^{\frac{7}{2}}\sqrt{1 + \frac{bx^2}{a}} + 3a^{\frac{5}{2}}bx^2\sqrt{1 + \frac{bx^2}{a}}} + \frac{2bx^3}{3a^{\frac{7}{2}}\sqrt{1 + \frac{bx^2}{a}} + 3a^{\frac{5}{2}}bx^2\sqrt{1 + \frac{bx^2}{a}}} \right) + \frac{dx^3}{3a^{\frac{5}{2}}\sqrt{1 + \frac{bx^2}{a}} + 3a^{\frac{3}{2}}bx^2\sqrt{1 + \frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)/(b*x**2+a)**(5/2),x)

[Out] c*(3*a*x/(3*a**(7/2)*sqrt(1 + b*x**2/a) + 3*a**(5/2)*b*x**2*sqrt(1 + b*x**2/a)) + 2*b*x**3/(3*a**(7/2)*sqrt(1 + b*x**2/a) + 3*a**(5/2)*b*x**2*sqrt(1 + b*x**2/a))) + d*x**3/(3*a**(5/2)*sqrt(1 + b*x**2/a) + 3*a**(3/2)*b*x**2*sqrt(1 + b*x**2/a))

$$3.93 \quad \int \frac{1}{(a+bx^2)^{5/2}} dx$$

Optimal. Leaf size=39

$$\frac{2x}{3a^2\sqrt{a+bx^2}} + \frac{x}{3a(a+bx^2)^{3/2}}$$

[Out] 1/3*x/a/(b*x^2+a)^(3/2)+2/3*x/a^2/(b*x^2+a)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {192, 191}

$$\frac{2x}{3a^2\sqrt{a+bx^2}} + \frac{x}{3a(a+bx^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(-5/2), x]

[Out] x/(3*a*(a + b*x^2)^(3/2)) + (2*x)/(3*a^2*Sqrt[a + b*x^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)^{5/2}} dx &= \frac{x}{3a(a+bx^2)^{3/2}} + \frac{2 \int \frac{1}{(a+bx^2)^{3/2}} dx}{3a} \\ &= \frac{x}{3a(a+bx^2)^{3/2}} + \frac{2x}{3a^2\sqrt{a+bx^2}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 29, normalized size = 0.74

$$\frac{x(3a + 2bx^2)}{3a^2(a + bx^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(-5/2), x]

[Out] (x*(3*a + 2*b*x^2))/(3*a^2*(a + b*x^2)^(3/2))

fricas [A] time = 0.80, size = 47, normalized size = 1.21

$$\frac{(2bx^3 + 3ax)\sqrt{bx^2 + a}}{3(a^2b^2x^4 + 2a^3bx^2 + a^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2),x, algorithm="fricas")

[Out] 1/3*(2*b*x^3 + 3*a*x)*sqrt(b*x^2 + a)/(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4)

giac [A] time = 0.61, size = 27, normalized size = 0.69

$$\frac{x\left(\frac{2bx^2}{a^2} + \frac{3}{a}\right)}{3(bx^2 + a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2),x, algorithm="giac")

[Out] 1/3*x*(2*b*x^2/a^2 + 3/a)/(b*x^2 + a)^(3/2)

maple [A] time = 0.00, size = 26, normalized size = 0.67

$$\frac{(2bx^2 + 3a)x}{3(bx^2 + a)^{\frac{3}{2}}a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(5/2),x)

[Out] 1/3*(2*b*x^2+3*a)/(b*x^2+a)^(3/2)/a^2*x

maxima [A] time = 1.33, size = 31, normalized size = 0.79

$$\frac{2x}{3\sqrt{bx^2 + a}a^2} + \frac{x}{3(bx^2 + a)^{\frac{3}{2}}a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2),x, algorithm="maxima")

[Out] 2/3*x/(sqrt(b*x^2 + a)*a^2) + 1/3*x/((b*x^2 + a)^(3/2)*a)

mupad [B] time = 4.75, size = 28, normalized size = 0.72

$$\frac{2x(bx^2 + a) + ax}{3a^2(bx^2 + a)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + b*x^2)^(5/2),x)

[Out] (2*x*(a + b*x^2) + a*x)/(3*a^2*(a + b*x^2)^(3/2))

sympy [B] time = 0.82, size = 95, normalized size = 2.44

$$\frac{3ax}{3a^{\frac{7}{2}}\sqrt{1 + \frac{bx^2}{a}} + 3a^{\frac{5}{2}}bx^2\sqrt{1 + \frac{bx^2}{a}}} + \frac{2bx^3}{3a^{\frac{7}{2}}\sqrt{1 + \frac{bx^2}{a}} + 3a^{\frac{5}{2}}bx^2\sqrt{1 + \frac{bx^2}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x**2+a)**(5/2),x)
```

```
[Out] 3*a*x/(3*a**(7/2)*sqrt(1 + b*x**2/a) + 3*a**(5/2)*b*x**2*sqrt(1 + b*x**2/a)
) + 2*b*x**3/(3*a**(7/2)*sqrt(1 + b*x**2/a) + 3*a**(5/2)*b*x**2*sqrt(1 + b*
x**2/a))
```

$$3.94 \quad \int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)} dx$$

Optimal. Leaf size=122

$$\frac{bx(2bc-5ad)}{3a^2\sqrt{a+bx^2}(bc-ad)^2} + \frac{d^2 \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc-ad)^{5/2}} + \frac{bx}{3a(a+bx^2)^{3/2}(bc-ad)}$$

[Out] $1/3*b*x/a/(-a*d+b*c)/(b*x^2+a)^{(3/2)+d^2*\operatorname{arctanh}(x*(-a*d+b*c)^{(1/2)/c^{(1/2)}}/(b*x^2+a)^{(1/2)))/(-a*d+b*c)^{(5/2)/c^{(1/2)}+1/3*b*(-5*a*d+2*b*c)*x/a^2/(-a*d+b*c)^2/(b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {414, 527, 12, 377, 208}

$$\frac{bx(2bc-5ad)}{3a^2\sqrt{a+bx^2}(bc-ad)^2} + \frac{d^2 \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc-ad)^{5/2}} + \frac{bx}{3a(a+bx^2)^{3/2}(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(5/2)*(c + d*x^2)), x]

[Out] $(b*x)/(3*a*(b*c - a*d)*(a + b*x^2)^{(3/2)} + (b*(2*b*c - 5*a*d)*x)/(3*a^2*(b*c - a*d)^2*\operatorname{Sqrt}[a + b*x^2]) + (d^2*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b*c - a*d]*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2])])/(\operatorname{Sqrt}[c]*(b*c - a*d)^{(5/2)})$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c +

$d*x^n)^{(q+1)}/(a*n*(b*c - a*d)*(p+1)), x] + \text{Dist}[1/(a*n*(b*c - a*d)*(p+1)), \text{Int}[(a + b*x^n)^{(p+1)}*(c + d*x^n)^q*\text{Simp}[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2) + 1)*x^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, q\}, x] \&\& \text{LtQ}[p, -1]$

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)} dx &= \frac{bx}{3a(bc-ad)(a+bx^2)^{3/2}} - \frac{\int \frac{-2bc+3ad-2bdx^2}{(a+bx^2)^{3/2}(c+dx^2)} dx}{3a(bc-ad)} \\ &= \frac{bx}{3a(bc-ad)(a+bx^2)^{3/2}} + \frac{b(2bc-5ad)x}{3a^2(bc-ad)^2\sqrt{a+bx^2}} + \frac{\int \frac{3a^2d^2}{\sqrt{a+bx^2}(c+dx^2)} dx}{3a^2(bc-ad)^2} \\ &= \frac{bx}{3a(bc-ad)(a+bx^2)^{3/2}} + \frac{b(2bc-5ad)x}{3a^2(bc-ad)^2\sqrt{a+bx^2}} + \frac{d^2 \int \frac{1}{\sqrt{a+bx^2}(c+dx^2)} dx}{(bc-ad)^2} \\ &= \frac{bx}{3a(bc-ad)(a+bx^2)^{3/2}} + \frac{b(2bc-5ad)x}{3a^2(bc-ad)^2\sqrt{a+bx^2}} + \frac{d^2 \text{Subst}\left(\int \frac{1}{c-(bc-ad)x^2} dx, x\right)}{(bc-ad)^2} \\ &= \frac{bx}{3a(bc-ad)(a+bx^2)^{3/2}} + \frac{b(2bc-5ad)x}{3a^2(bc-ad)^2\sqrt{a+bx^2}} + \frac{d^2 \tanh^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{c}\sqrt{a+bx^2}}\right)}{\sqrt{c}(bc-ad)^{5/2}} \end{aligned}$$

Mathematica [C] time = 2.75, size = 775, normalized size = 6.35

$$x \left(12c^2 \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{7/2} {}_3F_2 \left(2, 2, \frac{7}{2}; 1, \frac{9}{2}; \frac{(bc-ad)x^2}{c(bx^2+a)} \right) + 12d^2x^4 \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{7/2} {}_3F_2 \left(2, 2, \frac{7}{2}; 1, \frac{9}{2}; \frac{(bc-ad)x^2}{c(bx^2+a)} \right) + 24cdx^2 \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{7/2} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(5/2)*(c + d*x^2)), x]

[Out] $(x*(-315*c^2*\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]) - 420*c*d*x^2*\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]) - 168*d^2*x^4*\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]) - 105*c^2*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{3/2} - 140*c*d*x^2*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{3/2} - 56*d^2*x^4*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{3/2} + 315*c^2*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]]) + 420*c*d*x^2*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]]) + 168*d^2*x^4*\text{ArcTanh}[\text{Sqrt}[\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}]]) + 48*c^2*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{7/2}*\text{Hypergeometric2F1}[2, 7/2, 9/2, \frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}] + 84*c*d*x^2*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{7/2}*\text{Hypergeometric2F1}[2, 7/2, 9/2, \frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}] + 36*d^2*x^4*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{7/2}*\text{Hypergeometric2F1}[2, 7/2, 9/2, \frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}] + 12*c^2*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{7/2}*\text{HypergeometricPFQ}[\{2, 2, 7/2\}, \{1, 9/2\}, \frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}] + 24*c*d*x^2*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{7/2}*\text{HypergeometricPFQ}[\{2, 2, 7/2\}, \{1, 9/2\}, \frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}] + 12*d^2*x^4*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{7/2}*\text{HypergeometricPFQ}[\{2, 2, 7/2\}, \{1, 9/2\}, \frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}])]/(63*c^3*\left(\frac{(b*c - a*d)*x^2}{c*(a + b*x^2)}\right)^{5/2}*(a + b*x^2)^{5/2})$

fricas [B] time = 0.98, size = 764, normalized size = 6.26

$$\frac{3(a^2b^2d^2x^4 + 2a^3bd^2x^2 + a^4d^2)\sqrt{bc^2 - acd} \log\left(\frac{(8b^2c^2 - 8abcd + a^2d^2)x^4 + a^2c^2 + 2(4abc^2 - 3a^2cd)x^2 + 4((2bc - ad)x^3 + acx)\sqrt{bc^2 - acd}}{d^2x^4 + 2cdx^2 + c^2}\right)}{12(a^4b^3c^4 - 3a^5b^2c^3d + 3a^6bc^2d^2 - a^7cd^3 + (a^2b^5c^4 - 3a^3b^4c^3d + 3a^4b^3c^2d^2 - a^5b^2c^3d^2 + a^6b^3c^2d^3 - a^7b^4c^3d^2 + a^8b^5c^4d^3 - a^9b^6c^5d^4))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c),x, algorithm="fricas")

[Out] [1/12*(3*(a^2*b^2*d^2*x^4 + 2*a^3*b*d^2*x^2 + a^4*d^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*((2*b^4*c^3 - 7*a*b^3*c^2*d + 5*a^2*b^2*c*d^2)*x^3 + 3*(a*b^3*c^3 - 3*a^2*b^2*c^2*d + 2*a^3*b*c*d^2)*x)*sqrt(b*x^2 + a))/(a^4*b^3*c^4 - 3*a^5*b^2*c^3*d + 3*a^6*b*c^2*d^2 - a^7*c*d^3 + (a^2*b^5*c^4 - 3*a^3*b^4*c^3*d + 3*a^4*b^3*c^2*d^2 - a^5*b^2*c*d^3)*x^4 + 2*(a^3*b^4*c^4 - 3*a^4*b^3*c^3*d + 3*a^5*b^2*c^2*d^2 - a^6*b*c*d^3)*x^2), -1/6*(3*(a^2*b^2*d^2*x^4 + 2*a^3*b*d^2*x^2 + a^4*d^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a)/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)) - 2*((2*b^4*c^3 - 7*a*b^3*c^2*d + 5*a^2*b^2*c*d^2)*x^3 + 3*(a*b^3*c^3 - 3*a^2*b^2*c^2*d + 2*a^3*b*c*d^2)*x)*sqrt(b*x^2 + a))/(a^4*b^3*c^4 - 3*a^5*b^2*c^3*d + 3*a^6*b*c^2*d^2 - a^7*c*d^3 + (a^2*b^5*c^4 - 3*a^3*b^4*c^3*d + 3*a^4*b^3*c^2*d^2 - a^5*b^2*c*d^3)*x^4 + 2*(a^3*b^4*c^4 - 3*a^4*b^3*c^3*d + 3*a^5*b^2*c^2*d^2 - a^6*b*c*d^3)*x^2)]

giac [B] time = 0.64, size = 320, normalized size = 2.62

$$\frac{\sqrt{b}d^2 \arctan\left(\frac{(\sqrt{b}x - \sqrt{bx^2+a})^2 d + 2bc - ad}{2\sqrt{-b^2c^2 + abcd}}\right)}{(b^2c^2 - 2abcd + a^2d^2)\sqrt{-b^2c^2 + abcd}} + \frac{\left(\frac{(2b^6c^3 - 9ab^5c^2d + 12a^2b^4cd^2 - 5a^3b^3d^3)x^2}{a^2b^5c^4 - 4a^3b^4c^3d + 6a^4b^3c^2d^2 - 4a^5b^2cd^3 + a^6bd^4} + \frac{3(ab^5c^3 - 4a^2b^4c^2d + 5a^3b^3cd^2 - 2a^4b^2c^2d^2 + a^5b^3cd^3 - 2a^6b^4c^4d^4 + a^7b^5c^5d^5)}{a^2b^5c^4 - 4a^3b^4c^3d + 6a^4b^3c^2d^2 - 4a^5b^2cd^3 + a^6bd^4}\right)}{3(bx^2 + a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c),x, algorithm="giac")

[Out] -sqrt(b)*d^2*arctan(1/2*((sqrt(b)*x - sqrt(b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/((b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(-b^2*c^2 + a*b*c*d)) + 1/3*((2*b^6*c^3 - 9*a*b^5*c^2*d + 12*a^2*b^4*c*d^2 - 5*a^3*b^3*d^3)*x^2/(a^2*b^5*c^4 - 4*a^3*b^4*c^3*d + 6*a^4*b^3*c^2*d^2 - 4*a^5*b^2*c*d^3 + a^6*b*d^4) + 3*(a*b^5*c^3 - 4*a^2*b^4*c^2*d + 5*a^3*b^3*c*d^2 - 2*a^4*b^2*c*d^3)/(a^2*b^5*c^4 - 4*a^3*b^4*c^3*d + 6*a^4*b^3*c^2*d^2 - 4*a^5*b^2*c*d^3 + a^6*b*d^4))*x/(b*x^2 + a)^(3/2)

maple [B] time = 0.02, size = 1070, normalized size = 8.77

$$\frac{d^2 \ln\left(\frac{\frac{2\sqrt{-cd}\left(x - \frac{\sqrt{-cd}}{d}\right)b}{d} + \frac{2ad-2bc}{d} + 2\sqrt{\frac{ad-bc}{d}} \sqrt{\left(x - \frac{\sqrt{-cd}}{d}\right)^2 b + \frac{2\sqrt{-cd}\left(x - \frac{\sqrt{-cd}}{d}\right)b}{d} + \frac{ad-bc}{d}}}{x - \frac{\sqrt{-cd}}{d}}}{2\sqrt{-cd} (ad - bc)^2 \sqrt{\frac{ad-bc}{d}}}\right)}{2\sqrt{-cd} (ad - bc)^2 \sqrt{\frac{ad-bc}{d}}} + \frac{d^2 \ln\left(\frac{\frac{2\sqrt{-cd}\left(x + \frac{\sqrt{-cd}}{d}\right)b}{d} + \frac{2ad-2bc}{d} + 2\sqrt{\frac{ad-bc}{d}} \sqrt{\left(x + \frac{\sqrt{-cd}}{d}\right)^2 b + \frac{2\sqrt{-cd}\left(x + \frac{\sqrt{-cd}}{d}\right)b}{d} + \frac{ad-bc}{d}}}{x + \frac{\sqrt{-cd}}{d}}}{2\sqrt{-cd} (ad - bc)^2 \sqrt{\frac{ad-bc}{d}}}\right)}{2\sqrt{-cd} (ad - bc)^2 \sqrt{\frac{ad-bc}{d}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(5/2)/(d*x^2+c),x)

[Out] $\frac{1}{6} \frac{(-c*d)^{1/2}}{(a*d-b*c)*d} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2}*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} - \frac{1}{6} \frac{b}{(a*d-b*c)} \frac{1}{a} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2}*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} * x - \frac{1}{3} \frac{b}{(a*d-b*c)} \frac{1}{a^2} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2}*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x + \frac{1}{2} \frac{1}{(-c*d)^{1/2}} \frac{d^2}{(a*d-b*c)^2} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2}*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} - \frac{1}{2} \frac{d}{(a*d-b*c)^2} \frac{1}{a} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2}*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x * b - \frac{1}{2} \frac{1}{(-c*d)^{1/2}} \frac{d^2}{(a*d-b*c)^2} \frac{1}{((a*d-b*c)/d)^{1/2}} * \ln\left(\frac{(2*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{1/2}*((x-(-c*d)^{1/2}/d)^{2*b+2}*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}}{(x-(-c*d)^{1/2}/d)}\right) - \frac{1}{6} \frac{1}{(-c*d)^{1/2}} \frac{1}{(a*d-b*c)*d} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2}*(-c*d)^{1/2}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} - \frac{1}{6} \frac{b}{(a*d-b*c)} \frac{1}{a} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2}*(-c*d)^{1/2}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} * x - \frac{1}{3} \frac{b}{(a*d-b*c)} \frac{1}{a^2} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2}*(-c*d)^{1/2}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x - \frac{1}{2} \frac{1}{(-c*d)^{1/2}} \frac{d^2}{(a*d-b*c)^2} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2}*(-c*d)^{1/2}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x * b + \frac{1}{2} \frac{1}{(-c*d)^{1/2}} \frac{d^2}{(a*d-b*c)^2} \frac{1}{((a*d-b*c)/d)^{1/2}} * \ln\left(\frac{-2*(-c*d)^{1/2}*(x+(-c*d)^{1/2}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{1/2}*((x+(-c*d)^{1/2}/d)^{2*b-2}*(-c*d)^{1/2}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}}{(x+(-c*d)^{1/2}/d)}\right)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{2}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(5/2)*(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + a)^{5/2}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)),x)

[Out] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{5}{2}}(c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(5/2)/(d*x**2+c),x)

[Out] Integral(1/((a + b*x**2)**(5/2)*(c + d*x**2)), x)

$$3.95 \quad \int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)^2} dx$$

Optimal. Leaf size=202

$$\frac{bx(-3a^2d^2 - 16abcd + 4b^2c^2)}{6a^2c\sqrt{a+bx^2}(bc-ad)^3} + \frac{d^2(6bc-ad)\tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}(bc-ad)^{7/2}} - \frac{dx}{2c(a+bx^2)^{3/2}(c+dx^2)(bc-ad)} + \frac{bx}{6ac(a+bx^2)}$$

[Out] 1/6*b*(3*a*d+2*b*c)*x/a/c/(-a*d+b*c)^2/(b*x^2+a)^(3/2)-1/2*d*x/c/(-a*d+b*c)/(b*x^2+a)^(3/2)/(d*x^2+c)+1/2*d^2*(-a*d+6*b*c)*arctanh(x*(-a*d+b*c)^(1/2)/c^(1/2)/(b*x^2+a)^(1/2))/c^(3/2)/(-a*d+b*c)^(7/2)+1/6*b*(-3*a^2*d^2-16*a*b*c*d+4*b^2*c^2)*x/a^2/c/(-a*d+b*c)^3/(b*x^2+a)^(1/2)

Rubi [A] time = 0.23, antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {414, 527, 12, 377, 208}

$$\frac{bx(-3a^2d^2 - 16abcd + 4b^2c^2)}{6a^2c\sqrt{a+bx^2}(bc-ad)^3} + \frac{d^2(6bc-ad)\tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{2c^{3/2}(bc-ad)^{7/2}} - \frac{dx}{2c(a+bx^2)^{3/2}(c+dx^2)(bc-ad)} + \frac{bx}{6ac(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(5/2)*(c + d*x^2)^2), x]

[Out] (b*(2*b*c + 3*a*d)*x)/(6*a*c*(b*c - a*d)^2*(a + b*x^2)^(3/2)) + (b*(4*b^2*c^2 - 16*a*b*c*d - 3*a^2*d^2)*x)/(6*a^2*c*(b*c - a*d)^3*Sqrt[a + b*x^2]) - (d*x)/(2*c*(b*c - a*d)*(a + b*x^2)^(3/2)*(c + d*x^2)) + (d^2*(6*b*c - a*d)*ArcTanh[(Sqrt[b*c - a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(2*c^(3/2)*(b*c - a*d)^(7/2))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + bx^2)^{5/2} (c + dx^2)^2} dx &= -\frac{dx}{2c(bc - ad)(a + bx^2)^{3/2} (c + dx^2)} + \frac{\int \frac{2bc - ad - 4bdx^2}{(a + bx^2)^{5/2} (c + dx^2)} dx}{2c(bc - ad)} \\ &= \frac{b(2bc + 3ad)x}{6ac(bc - ad)^2 (a + bx^2)^{3/2}} - \frac{dx}{2c(bc - ad)(a + bx^2)^{3/2} (c + dx^2)} - \frac{\int \frac{-4b^2c^2 + 12abcd - 3a^2d^2}{(a + bx^2)^{5/2} (c + dx^2)} dx}{6ac(bc - ad)^2} \\ &= \frac{b(2bc + 3ad)x}{6ac(bc - ad)^2 (a + bx^2)^{3/2}} + \frac{b(4b^2c^2 - 16abcd - 3a^2d^2)x}{6a^2c(bc - ad)^3 \sqrt{a + bx^2}} - \frac{dx}{2c(bc - ad)(a + bx^2)^{3/2}} \\ &= \frac{b(2bc + 3ad)x}{6ac(bc - ad)^2 (a + bx^2)^{3/2}} + \frac{b(4b^2c^2 - 16abcd - 3a^2d^2)x}{6a^2c(bc - ad)^3 \sqrt{a + bx^2}} - \frac{dx}{2c(bc - ad)(a + bx^2)^{3/2}} \\ &= \frac{b(2bc + 3ad)x}{6ac(bc - ad)^2 (a + bx^2)^{3/2}} + \frac{b(4b^2c^2 - 16abcd - 3a^2d^2)x}{6a^2c(bc - ad)^3 \sqrt{a + bx^2}} - \frac{dx}{2c(bc - ad)(a + bx^2)^{3/2}} \\ &= \frac{b(2bc + 3ad)x}{6ac(bc - ad)^2 (a + bx^2)^{3/2}} + \frac{b(4b^2c^2 - 16abcd - 3a^2d^2)x}{6a^2c(bc - ad)^3 \sqrt{a + bx^2}} - \frac{dx}{2c(bc - ad)(a + bx^2)^{3/2}} \end{aligned}$$

Mathematica [A] time = 5.50, size = 170, normalized size = 0.84

$$\frac{1}{6} \left(x \sqrt{a + bx^2} \left(\frac{4b^2(4ad - bc)}{a^2 (a + bx^2) (ad - bc)^3} + \frac{2b^2}{a (a + bx^2)^2 (bc - ad)^2} - \frac{3d^3}{c (c + dx^2) (bc - ad)^3} \right) + \frac{3d^2(ad - 6bc) \tan^{-1} \left(\frac{\sqrt{c} \sqrt{a + bx^2}}{c + dx^2} \right)}{c^{3/2}(ad - bc)} \right)$$

Antiderivative was successfully verified.

```
[In] Integrate[1/((a + b*x^2)^(5/2)*(c + d*x^2)^2), x]
```

```
[Out] (x*Sqrt[a + b*x^2]*((2*b^2)/(a*(b*c - a*d)^2*(a + b*x^2)^2) + (4*b^2*(-(b*c) + 4*a*d))/(a^2*(-(b*c) + a*d)^3*(a + b*x^2)) - (3*d^3)/(c*(b*c - a*d)^3*(c + d*x^2))) + (3*d^2*(-6*b*c + a*d)*ArcTan[(Sqrt[-(b*c) + a*d]*x)/(Sqrt[c]*Sqrt[a + b*x^2])])/(c^(3/2)*(-(b*c) + a*d)^(7/2)))/6
```

fricas [B] time = 2.28, size = 1440, normalized size = 7.13

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^2,x, algorithm="fricas")
```



```
[Out] [1/24*(3*(6*a^4*b*c^2*d^2 - a^5*c*d^3 + (6*a^2*b^3*c*d^3 - a^3*b^2*d^4)*x^6
+ (6*a^2*b^3*c^2*d^2 + 11*a^3*b^2*c*d^3 - 2*a^4*b*d^4)*x^4 + (12*a^3*b^2*c
^2*d^2 + 4*a^4*b*c*d^3 - a^5*d^4)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2
- 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((
2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2
*c*d*x^2 + c^2)) + 4*((4*b^5*c^4*d - 20*a*b^4*c^3*d^2 + 13*a^2*b^3*c^2*d^3
+ 3*a^3*b^2*c*d^4)*x^5 + 2*(2*b^5*c^5 - 7*a*b^4*c^4*d - 4*a^2*b^3*c^3*d^2 +
6*a^3*b^2*c^2*d^3 + 3*a^4*b*c*d^4)*x^3 + 3*(2*a*b^4*c^5 - 8*a^2*b^3*c^4*d
+ 6*a^3*b^2*c^3*d^2 - a^4*b*c^2*d^3 + a^5*c*d^4)*x)*sqrt(b*x^2 + a))/(a^4*b
^4*c^7 - 4*a^5*b^3*c^6*d + 6*a^6*b^2*c^5*d^2 - 4*a^7*b*c^4*d^3 + a^8*c^3*d^
4 + (a^2*b^6*c^6*d - 4*a^3*b^5*c^5*d^2 + 6*a^4*b^4*c^4*d^3 - 4*a^5*b^3*c^3*
d^4 + a^6*b^2*c^2*d^5)*x^6 + (a^2*b^6*c^7 - 2*a^3*b^5*c^6*d - 2*a^4*b^4*c^5
*d^2 + 8*a^5*b^3*c^4*d^3 - 7*a^6*b^2*c^3*d^4 + 2*a^7*b*c^2*d^5)*x^4 + (2*a^
3*b^5*c^7 - 7*a^4*b^4*c^6*d + 8*a^5*b^3*c^5*d^2 - 2*a^6*b^2*c^4*d^3 - 2*a^7
*b*c^3*d^4 + a^8*c^2*d^5)*x^2), -1/12*(3*(6*a^4*b*c^2*d^2 - a^5*c*d^3 + (6*
a^2*b^3*c*d^3 - a^3*b^2*d^4)*x^6 + (6*a^2*b^3*c^2*d^2 + 11*a^3*b^2*c*d^3 -
2*a^4*b*d^4)*x^4 + (12*a^3*b^2*c^2*d^2 + 4*a^4*b*c*d^3 - a^5*d^4)*x^2)*sqrt
(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*
sqrt(b*x^2 + a))/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)) - 2*((4*
b^5*c^4*d - 20*a*b^4*c^3*d^2 + 13*a^2*b^3*c^2*d^3 + 3*a^3*b^2*c*d^4)*x^5 +
2*(2*b^5*c^5 - 7*a*b^4*c^4*d - 4*a^2*b^3*c^3*d^2 + 6*a^3*b^2*c^2*d^3 + 3*a^
4*b*c*d^4)*x^3 + 3*(2*a*b^4*c^5 - 8*a^2*b^3*c^4*d + 6*a^3*b^2*c^3*d^2 - a^4
*b*c^2*d^3 + a^5*c*d^4)*x)*sqrt(b*x^2 + a))/(a^4*b^4*c^7 - 4*a^5*b^3*c^6*d
+ 6*a^6*b^2*c^5*d^2 - 4*a^7*b*c^4*d^3 + a^8*c^3*d^4 + (a^2*b^6*c^6*d - 4*a^
3*b^5*c^5*d^2 + 6*a^4*b^4*c^4*d^3 - 4*a^5*b^3*c^3*d^4 + a^6*b^2*c^2*d^5)*x^
6 + (a^2*b^6*c^7 - 2*a^3*b^5*c^6*d - 2*a^4*b^4*c^5*d^2 + 8*a^5*b^3*c^4*d^3
- 7*a^6*b^2*c^3*d^4 + 2*a^7*b*c^2*d^5)*x^4 + (2*a^3*b^5*c^7 - 7*a^4*b^4*c^6
*d + 8*a^5*b^3*c^5*d^2 - 2*a^6*b^2*c^4*d^3 - 2*a^7*b*c^3*d^4 + a^8*c^2*d^5)
*x^2)]
```

giac [B] time = 2.07, size = 620, normalized size = 3.07

$$\frac{\left(\frac{2(b^8c^4 - 7ab^7c^3d + 15a^2b^6c^2d^2 - 13a^3b^5cd^3 + 4a^4b^4d^4)x^2}{a^2b^7c^6 - 6a^3b^6c^5d + 15a^4b^5c^4d^2 - 20a^5b^4c^3d^3 + 15a^6b^3c^2d^4 - 6a^7b^2cd^5 + a^8bd^6} + \frac{3(ab^7c^4 - 6a^2b^6c^3d + 12a^3b^5c^2d^2 - 10a^4b^4cd^3 + 3a^5b^3d^4)}{a^2b^7c^6 - 6a^3b^6c^5d + 15a^4b^5c^4d^2 - 20a^5b^4c^3d^3 + 15a^6b^3c^2d^4 - 6a^7b^2cd^5 + a^8bd^6} \right)}{3(bx^2 + a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^2,x, algorithm="giac")
```

```
[Out] 1/3*(2*(b^8*c^4 - 7*a*b^7*c^3*d + 15*a^2*b^6*c^2*d^2 - 13*a^3*b^5*c*d^3 + 4
*a^4*b^4*d^4)*x^2/(a^2*b^7*c^6 - 6*a^3*b^6*c^5*d + 15*a^4*b^5*c^4*d^2 - 20*
a^5*b^4*c^3*d^3 + 15*a^6*b^3*c^2*d^4 - 6*a^7*b^2*c*d^5 + a^8*b*d^6) + 3*(a*
b^7*c^4 - 6*a^2*b^6*c^3*d + 12*a^3*b^5*c^2*d^2 - 10*a^4*b^4*c*d^3 + 3*a^5*b
^3*d^4)/(a^2*b^7*c^6 - 6*a^3*b^6*c^5*d + 15*a^4*b^5*c^4*d^2 - 20*a^5*b^4*c^
3*d^3 + 15*a^6*b^3*c^2*d^4 - 6*a^7*b^2*c*d^5 + a^8*b*d^6))*x/(b*x^2 + a)^(3
/2) + 1/2*(6*b^(3/2)*c*d^2 - a*sqrt(b)*d^3)*arctan(-1/2*((sqrt(b)*x - sqrt(
b*x^2 + a))^2*d + 2*b*c - a*d)/sqrt(-b^2*c^2 + a*b*c*d))/((b^3*c^4 - 3*a*b^
2*c^3*d + 3*a^2*b*c^2*d^2 - a^3*c*d^3)*sqrt(-b^2*c^2 + a*b*c*d)) - (2*(sqrt
(b)*x - sqrt(b*x^2 + a))^2*b^(3/2)*c*d^2 - (sqrt(b)*x - sqrt(b*x^2 + a))^2*
a*sqrt(b)*d^3 + a^2*sqrt(b)*d^3)/((b^3*c^4 - 3*a*b^2*c^3*d + 3*a^2*b*c^2*d^
2 - a^3*c*d^3)*((sqrt(b)*x - sqrt(b*x^2 + a))^4*d + 4*(sqrt(b)*x - sqrt(b*x
^2 + a))^2*b*c - 2*(sqrt(b)*x - sqrt(b*x^2 + a))^2*a*d + a^2*d))
```

maple [B] time = 0.03, size = 2371, normalized size = 11.74

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(5/2)/(d*x^2+c)^2,x)

[Out] $\frac{1}{4} \frac{(-c*d)^{1/2}}{c*d^2} \frac{1}{(a*d-b*c)^2} \frac{1}{((a*d-b*c)/d)^{1/2}} \ln\left(\frac{-2*(-c*d)^{1/2}*(x+(-c*d)^{1/2}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{1/2}*((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}}{(x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d}\right) + \frac{1}{4} \frac{c*b}{(a*d-b*c)} \frac{1}{a} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} * x + \frac{1}{2} \frac{c*b}{(a*d-b*c)} \frac{1}{a^2} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} * x + \frac{5}{4} \frac{c*d}{d} * (-c*d)^{1/2} * b / (a*d-b*c)^3 \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} + \frac{5}{4} \frac{d*b^2}{(a*d-b*c)^3} \frac{1}{a} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x + \frac{1}{4} \frac{c}{(a*d-b*c)} \frac{1}{(x+(-c*d)^{1/2}/d)} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} + \frac{1}{4} \frac{c}{(a*d-b*c)} \frac{1}{(x-(-c*d)^{1/2}/d)} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} - \frac{1}{4} \frac{(-c*d)^{1/2}}{c*d^2} \frac{1}{(a*d-b*c)^2} \frac{1}{((a*d-b*c)/d)^{1/2}} \ln\left(\frac{2*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{1/2}*((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}}{(x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d}\right) - \frac{5}{4} \frac{c*d}{d} * (-c*d)^{1/2} * b / (a*d-b*c)^3 \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} + \frac{5}{4} \frac{d*b^2}{(a*d-b*c)^3} \frac{1}{a} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x + \frac{1}{4} \frac{c*b}{(a*d-b*c)} \frac{1}{a} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} * x - \frac{5}{4} \frac{c*d}{d} * (-c*d)^{1/2} * b / (a*d-b*c)^3 \frac{1}{((a*d-b*c)/d)^{1/2}} \ln\left(\frac{2*(-c*d)^{1/2}*(x-(-c*d)^{1/2}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{1/2}*((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}}{(x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d}\right) - \frac{1}{4} \frac{c*d}{(a*d-b*c)^2} \frac{1}{a} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x * b - \frac{1}{4} \frac{c*d}{(a*d-b*c)^2} \frac{1}{a} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x * b + \frac{5}{4} \frac{c*d}{d} * (-c*d)^{1/2} * b / (a*d-b*c)^3 \frac{1}{((a*d-b*c)/d)^{1/2}} \ln\left(\frac{-2*(-c*d)^{1/2}*(x+(-c*d)^{1/2}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{1/2}*((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}}{(x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d}\right) - \frac{5}{12} \frac{c}{c} * (-c*d)^{1/2} * b / (a*d-b*c)^2 \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} + \frac{5}{12} \frac{b^2}{(a*d-b*c)^2} \frac{1}{a} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} * x + \frac{5}{12} \frac{c}{c} * (-c*d)^{1/2} * b / (a*d-b*c)^2 \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} + \frac{1}{12} \frac{(-c*d)^{1/2}}{c} \frac{1}{(a*d-b*c)} \frac{1}{d} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} + \frac{1}{4} \frac{(-c*d)^{1/2}}{c*d^2} \frac{1}{(a*d-b*c)^2} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} + \frac{5}{6} \frac{b^2}{(a*d-b*c)^2} \frac{1}{a^2} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x + \frac{5}{12} \frac{b^2}{(a*d-b*c)^2} \frac{1}{a} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{3/2}} * x + \frac{5}{6} \frac{b^2}{(a*d-b*c)^2} \frac{1}{a^2} \frac{1}{((x-(-c*d)^{1/2}/d)^{2*b+2*(-c*d)^{1/2}}*(x-(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x + \frac{1}{2} \frac{c*b}{(a*d-b*c)} \frac{1}{a^2} \frac{1}{((x+(-c*d)^{1/2}/d)^{2*b-2*(-c*d)^{1/2}}*(x+(-c*d)^{1/2}/d)*b/d+(a*d-b*c)/d)^{1/2}} * x$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{2}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(5/2)*(d*x^2 + c)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{5/2} (dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)^2), x)

[Out] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{5/2} (c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(5/2)/(d*x**2+c)**2, x)

[Out] Integral(1/((a + b*x**2)**(5/2)*(c + d*x**2)**2), x)

$$3.96 \quad \int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)^3} dx$$

Optimal. Leaf size=313

$$\frac{bx(-3a^2d^2 - 40abcd + 8b^2c^2)}{12a^2c\sqrt{a+bx^2}(c+dx^2)(bc-ad)^3} + \frac{d^2(3a^2d^2 - 16abcd + 48b^2c^2) \tanh^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{c}\sqrt{a+bx^2}}\right)}{8c^{5/2}(bc-ad)^{9/2}} + \frac{dx\sqrt{a+bx^2}(9a^3d^3 - 42a^2bd^2 + 16abcd - 8b^2c^2)}{24a^2c^2(c+dx^2)(bc-ad)^3}$$

[Out] $-1/4*d*x/c/(-a*d+b*c)/(b*x^2+a)^{(3/2)}/(d*x^2+c)^2+1/12*b*(3*a*d+4*b*c)*x/a/c/(-a*d+b*c)^2/(b*x^2+a)^{(3/2)}/(d*x^2+c)+1/8*d^2*(3*a^2*d^2-16*a*b*c*d+48*b^2*c^2)*\arctanh(x*(-a*d+b*c)^{(1/2)}/c^{(1/2)}/(b*x^2+a)^{(1/2)})/c^{(5/2)}/(-a*d+b*c)^{(9/2)}+1/12*b*(-3*a^2*d^2-40*a*b*c*d+8*b^2*c^2)*x/a^2/c/(-a*d+b*c)^3/(d*x^2+c)/(b*x^2+a)^{(1/2)}+1/24*d*(9*a^3*d^3-42*a^2*b*c*d^2-88*a*b^2*c^2*d+16*b^3*c^3)*x*(b*x^2+a)^{(1/2)}/a^2/c^2/(-a*d+b*c)^4/(d*x^2+c)$

Rubi [A] time = 0.40, antiderivative size = 313, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {414, 527, 12, 377, 208}

$$\frac{dx\sqrt{a+bx^2}(-42a^2bcd^2 + 9a^3d^3 - 88ab^2c^2d + 16b^3c^3)}{24a^2c^2(c+dx^2)(bc-ad)^4} + \frac{bx(-3a^2d^2 - 40abcd + 8b^2c^2)}{12a^2c\sqrt{a+bx^2}(c+dx^2)(bc-ad)^3} + \frac{d^2(3a^2d^2 - 16abcd + 48b^2c^2)}{8c^{5/2}(bc-ad)^{9/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(5/2)*(c + d*x^2)^3), x]

[Out] $-(d*x)/(4*c*(b*c - a*d)*(a + b*x^2)^{(3/2)}*(c + d*x^2)^2) + (b*(4*b*c + 3*a*d)*x)/(12*a*c*(b*c - a*d)^2*(a + b*x^2)^{(3/2)}*(c + d*x^2)) + (b*(8*b^2*c^2 - 40*a*b*c*d - 3*a^2*d^2)*x)/(12*a^2*c*(b*c - a*d)^3*\text{Sqrt}[a + b*x^2]*(c + d*x^2)) + (d*(16*b^3*c^3 - 88*a*b^2*c^2*d - 42*a^2*b*c*d^2 + 9*a^3*d^3)*x*\text{Sqrt}[a + b*x^2])/(24*a^2*c^2*(b*c - a*d)^4*(c + d*x^2)) + (d^2*(48*b^2*c^2 - 16*a*b*c*d + 3*a^2*d^2)*\text{ArcTanh}[(\text{Sqrt}[b*c - a*d]*x)/(\text{Sqrt}[c]*\text{Sqrt}[a + b*x^2])])/(8*c^{(5/2)}*(b*c - a*d)^{(9/2)})$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1]

&& !(!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + bx^2)^{5/2} (c + dx^2)^3} dx &= -\frac{dx}{4c(bc - ad) (a + bx^2)^{3/2} (c + dx^2)^2} + \frac{\int \frac{4bc - 3ad - 6bdx^2}{(a + bx^2)^{5/2} (c + dx^2)^2} dx}{4c(bc - ad)} \\ &= -\frac{dx}{4c(bc - ad) (a + bx^2)^{3/2} (c + dx^2)^2} + \frac{b(4bc + 3ad)x}{12ac(bc - ad)^2 (a + bx^2)^{3/2} (c + dx^2)} \\ &= -\frac{dx}{4c(bc - ad) (a + bx^2)^{3/2} (c + dx^2)^2} + \frac{b(4bc + 3ad)x}{12ac(bc - ad)^2 (a + bx^2)^{3/2} (c + dx^2)} + \\ &= -\frac{dx}{4c(bc - ad) (a + bx^2)^{3/2} (c + dx^2)^2} + \frac{b(4bc + 3ad)x}{12ac(bc - ad)^2 (a + bx^2)^{3/2} (c + dx^2)} + \\ &= -\frac{dx}{4c(bc - ad) (a + bx^2)^{3/2} (c + dx^2)^2} + \frac{b(4bc + 3ad)x}{12ac(bc - ad)^2 (a + bx^2)^{3/2} (c + dx^2)} + \\ &= -\frac{dx}{4c(bc - ad) (a + bx^2)^{3/2} (c + dx^2)^2} + \frac{b(4bc + 3ad)x}{12ac(bc - ad)^2 (a + bx^2)^{3/2} (c + dx^2)} + \\ &= -\frac{dx}{4c(bc - ad) (a + bx^2)^{3/2} (c + dx^2)^2} + \frac{b(4bc + 3ad)x}{12ac(bc - ad)^2 (a + bx^2)^{3/2} (c + dx^2)} + \end{aligned}$$

Mathematica [A] time = 5.66, size = 221, normalized size = 0.71

$$\frac{1}{24} \left(x \sqrt{a + bx^2} \left(\frac{8b^3(2bc - 11ad)}{a^2 (a + bx^2) (bc - ad)^4} - \frac{8b^3}{a (a + bx^2)^2 (ad - bc)^3} + \frac{3d^3(3ad - 14bc)}{c^2 (c + dx^2) (bc - ad)^4} - \frac{6d^3}{c (c + dx^2)^2 (bc - ad)^3} \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^(5/2)*(c + d*x^2)^3), x]

[Out] (x*Sqrt[a + b*x^2]*((-8*b^3)/(a*(-(b*c) + a*d)^3*(a + b*x^2)^2) + (8*b^3*(2*b*c - 11*a*d))/(a^2*(b*c - a*d)^4*(a + b*x^2)) - (6*d^3)/(c*(b*c - a*d)^3*(c + d*x^2)^2) + (3*d^3*(-14*b*c + 3*a*d))/(c^2*(b*c - a*d)^4*(c + d*x^2)))

$$+ (3*d^2*(48*b^2*c^2 - 16*a*b*c*d + 3*a^2*d^2)*ArcTan[(\text{Sqrt}[-(b*c) + a*d]*x)/(\text{Sqrt}[c]*\text{Sqrt}[a + b*x^2])])/(c^{5/2}*(-(b*c) + a*d)^{9/2}))/24$$

fricas [B] time = 8.34, size = 2250, normalized size = 7.19

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^3,x, algorithm="fricas")

[Out] [1/96*(3*(48*a^4*b^2*c^4*d^2 - 16*a^5*b*c^3*d^3 + 3*a^6*c^2*d^4 + (48*a^2*b^4*c^2*d^4 - 16*a^3*b^3*c*d^5 + 3*a^4*b^2*d^6)*x^8 + 2*(48*a^2*b^4*c^3*d^3 + 32*a^3*b^3*c^2*d^4 - 13*a^4*b^2*c*d^5 + 3*a^5*b*d^6)*x^6 + (48*a^2*b^4*c^4*d^2 + 176*a^3*b^3*c^3*d^3 - 13*a^4*b^2*c^2*d^4 - 4*a^5*b*c*d^5 + 3*a^6*d^6)*x^4 + 2*(48*a^3*b^3*c^4*d^2 + 32*a^4*b^2*c^3*d^3 - 13*a^5*b*c^2*d^4 + 3*a^6*c*d^5)*x^2)*sqrt(b*c^2 - a*c*d)*log(((8*b^2*c^2 - 8*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(4*a*b*c^2 - 3*a^2*c*d)*x^2 + 4*((2*b*c - a*d)*x^3 + a*c*x)*sqrt(b*c^2 - a*c*d)*sqrt(b*x^2 + a))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 4*((16*b^6*c^5*d^2 - 104*a*b^5*c^4*d^3 + 46*a^2*b^4*c^3*d^4 + 51*a^3*b^3*c^2*d^5 - 9*a^4*b^2*c*d^6)*x^7 + (32*b^6*c^6*d - 184*a*b^5*c^5*d^2 + 8*a^2*b^4*c^4*d^3 + 75*a^3*b^3*c^3*d^4 + 87*a^4*b^2*c^2*d^5 - 18*a^5*b*c*d^6)*x^5 + (16*b^6*c^7 - 56*a*b^5*c^6*d - 152*a^2*b^4*c^5*d^2 + 96*a^3*b^3*c^4*d^3 + 84*a^4*b^2*c^3*d^4 + 21*a^5*b*c^2*d^5 - 9*a^6*c*d^6)*x^3 + 3*(8*a*b^5*c^7 - 40*a^2*b^4*c^6*d + 32*a^3*b^3*c^5*d^2 - 16*a^4*b^2*c^4*d^3 + 21*a^5*b*c^3*d^4 - 5*a^6*c^2*d^5)*x)*sqrt(b*x^2 + a))/(a^4*b^5*c^10 - 5*a^5*b^4*c^9*d + 10*a^6*b^3*c^8*d^2 - 10*a^7*b^2*c^7*d^3 + 5*a^8*b*c^6*d^4 - a^9*c^5*d^5 + (a^2*b^7*c^8*d^2 - 5*a^3*b^6*c^7*d^3 + 10*a^4*b^5*c^6*d^4 - 10*a^5*b^4*c^5*d^5 + 5*a^6*b^3*c^4*d^6 - a^7*b^2*c^3*d^7)*x^8 + 2*(a^2*b^7*c^9*d - 4*a^3*b^6*c^8*d^2 + 5*a^4*b^5*c^7*d^3 - 5*a^6*b^3*c^5*d^5 + 4*a^7*b^2*c^4*d^6 - a^8*b*c^3*d^7)*x^6 + (a^2*b^7*c^10 - a^3*b^6*c^9*d - 9*a^4*b^5*c^8*d^2 + 25*a^5*b^4*c^7*d^3 - 25*a^6*b^3*c^6*d^4 + 9*a^7*b^2*c^5*d^5 + a^8*b*c^4*d^6 - a^9*c^3*d^7)*x^4 + 2*(a^3*b^6*c^10 - 4*a^4*b^5*c^9*d + 5*a^5*b^4*c^8*d^2 - 5*a^7*b^2*c^6*d^4 + 4*a^8*b*c^5*d^5 - a^9*c^4*d^6)*x^2), -1/48*(3*(48*a^4*b^2*c^4*d^2 - 16*a^5*b*c^3*d^3 + 3*a^6*c^2*d^4 + (48*a^2*b^4*c^2*d^4 - 16*a^3*b^3*c*d^5 + 3*a^4*b^2*d^6)*x^8 + 2*(48*a^2*b^4*c^3*d^3 + 32*a^3*b^3*c^2*d^4 - 13*a^4*b^2*c*d^5 + 3*a^5*b*d^6)*x^6 + (48*a^2*b^4*c^4*d^2 + 176*a^3*b^3*c^3*d^3 - 13*a^4*b^2*c^2*d^4 - 4*a^5*b*c*d^5 + 3*a^6*d^6)*x^4 + 2*(48*a^3*b^3*c^4*d^2 + 32*a^4*b^2*c^3*d^3 - 13*a^5*b*c^2*d^4 + 3*a^6*c*d^5)*x^2)*sqrt(-b*c^2 + a*c*d)*arctan(1/2*sqrt(-b*c^2 + a*c*d)*((2*b*c - a*d)*x^2 + a*c)*sqrt(b*x^2 + a))/((b^2*c^2 - a*b*c*d)*x^3 + (a*b*c^2 - a^2*c*d)*x)) - 2*((16*b^6*c^5*d^2 - 104*a*b^5*c^4*d^3 + 46*a^2*b^4*c^3*d^4 + 51*a^3*b^3*c^2*d^5 - 9*a^4*b^2*c*d^6)*x^7 + (32*b^6*c^6*d - 184*a*b^5*c^5*d^2 + 8*a^2*b^4*c^4*d^3 + 75*a^3*b^3*c^3*d^4 + 87*a^4*b^2*c^2*d^5 - 18*a^5*b*c*d^6)*x^5 + (16*b^6*c^7 - 56*a*b^5*c^6*d - 152*a^2*b^4*c^5*d^2 + 96*a^3*b^3*c^4*d^3 + 84*a^4*b^2*c^3*d^4 + 21*a^5*b*c^2*d^5 - 9*a^6*c*d^6)*x^3 + 3*(8*a*b^5*c^7 - 40*a^2*b^4*c^6*d + 32*a^3*b^3*c^5*d^2 - 16*a^4*b^2*c^4*d^3 + 21*a^5*b*c^3*d^4 - 5*a^6*c^2*d^5)*x)*sqrt(b*x^2 + a))/(a^4*b^5*c^10 - 5*a^5*b^4*c^9*d + 10*a^6*b^3*c^8*d^2 - 10*a^7*b^2*c^7*d^3 + 5*a^8*b*c^6*d^4 - a^9*c^5*d^5 + (a^2*b^7*c^8*d^2 - 5*a^3*b^6*c^7*d^3 + 10*a^4*b^5*c^6*d^4 - 10*a^5*b^4*c^5*d^5 + 5*a^6*b^3*c^4*d^6 - a^7*b^2*c^3*d^7)*x^8 + 2*(a^2*b^7*c^9*d - 4*a^3*b^6*c^8*d^2 + 5*a^4*b^5*c^7*d^3 - 5*a^6*b^3*c^5*d^5 + 4*a^7*b^2*c^4*d^6 - a^8*b*c^3*d^7)*x^6 + (a^2*b^7*c^10 - a^3*b^6*c^9*d - 9*a^4*b^5*c^8*d^2 + 25*a^5*b^4*c^7*d^3 - 25*a^6*b^3*c^6*d^4 + 9*a^7*b^2*c^5*d^5 + a^8*b*c^4*d^6 - a^9*c^3*d^7)*x^4 + 2*(a^3*b^6*c^10 - 4*a^4*b^5*c^9*d + 5*a^5*b^4*c^8*d^2 - 5*a^7*b^2*c^6*d^4 + 4*a^8*b*c^5*d^5 - a^9*c^4*d^6)*x^2)]

$$\begin{aligned}
& b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))-7/16/c*b \\
& / (a*d-b*c)^2/(x+(-c*d)^{(1/2)}/d)/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}-35/48/(-c*d)^{(1/2)}*d*b^2/(a*d-b*c)^3/ \\
& ((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}-35/48*b^3/(a*d-b*c)^3/a/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c \\
& *d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*x-35/24*b^3/(a*d-b*c)^3/a^2/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x-35/ \\
& 16/(-c*d)^{(1/2)}*d^2*b^2/(a*d-b*c)^4/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}+5/16/c^2*(-c*d)^{(1/2)}*b/(a*d-b*c) \\
& ^2/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}+3/16/(-c*d)^{(1/2)}/c^2*d^2/(a*d-b*c)^2/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c \\
& *d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)} \\
&))/(x+(-c*d)^{(1/2)}/d))+15/16/c^2*d*(-c*d)^{(1/2)}*b/(a*d-b*c)^3/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}-15/16/c \\
& ^2*d*(-c*d)^{(1/2)}*b/(a*d-b*c)^3/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}+3/16/c^2*b/(a*d-b*c)/a/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*x+3/8/c^ \\
& 2*b/(a*d-b*c)/a^2/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x-35/16*d*b^3/(a*d-b*c)^4/a/((x+(-c*d)^{(1/2)}/d)^{2*b \\
& -2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x+35/16/(-c*d)^{(1/2)}*d^2*b^2/(a*d-b*c)^4/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c \\
& *d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))-3/8/c*b^2/(a*d-b*c)^2/a/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*x-3/4/c*b^2/(a*d-b*c)^2/a^2/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x-5/48/(-c \\
& *d)^{(1/2)}/c*d*b/(a*d-b*c)^2/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}-5/16/(-c*d)^{(1/2)}/c*d^2*b/(a*d-b*c)^3/((x+ \\
& (-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}-35/16*d*b^3/(a*d-b*c)^4/a/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d) \\
&)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x-35/16/(-c*d)^{(1/2)}*d^2*b^2/(a*d-b*c)^4/ \\
& ((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d \\
& +2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))-3/8/c*b^2/(a*d-b*c)^2/a/ \\
& (x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*x-3/4/c*b^2/(a*d-b*c)^2/a^2/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x- \\
& (-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x+15/16/c^2*d*(-c*d)^{(1/2)}*b/(a*d-b* \\
& c)^3/((a*d-b*c)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d- \\
& b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c* \\
& d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))+5/8/c*d*b^2/(a*d-b* \\
& c)^3/a/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b \\
& *c)/d)^{(1/2)}*x+5/16/(-c*d)^{(1/2)}/c*d^2*b/(a*d-b*c)^3/((a*d-b*c)/d)^{(1/2)}*\ln \\
& ((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)} \\
&)*((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d \\
&)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))+5/8/c*d*b^2/(a*d-b*c)^3/a/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c \\
& *d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x-5/16/(-c*d)^{(1/2)}/c*d^2*b/(a*d-b*c)^3/((a*d-b*c)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+2*(a*d-b*c)/d+2*((a*d-b*c)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c \\
& *d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d)) \\
& -3/16/c^2*d/(a*d-b*c)^2/a/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x*b-3/16/c^2*d/(a*d-b*c)^2/a/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x*b+35/4 \\
& 8/(-c*d)^{(1/2)}*d*b^2/(a*d-b*c)^3/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x- \\
& (-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}-35/48*b^3/(a*d-b*c)^3/a/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}*x-35/2 \\
& 4*b^3/(a*d-b*c)^3/a^2/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)}*x+35/16/(-c*d)^{(1/2)}*d^2*b^2/(a*d-b*c)^4/((x-(-
\end{aligned}$$

$$c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)+1/16/(-c*d)^{(1/2)}/c^2/(a*d-b*c)*d/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)+3/16/(-c*d)^{(1/2)}/c^2*d^2/(a*d-b*c)^2/((x-(-c*d)^{(1/2)}/d)^{2*b+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)-1/16/(-c*d)^{(1/2)}/c^2/(a*d-b*c)*d/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)-3/16/(-c*d)^{(1/2)}/c^2*d^2/(a*d-b*c)^2/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(1/2)-5/16/c^2*(-c*d)^{(1/2)*b/(a*d-b*c)^2/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)-1/16/(-c*d)^{(1/2)}/c/(a*d-b*c)/(x+(-c*d)^{(1/2)}/d)^2/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)*b/(a*d-b*c)^2/((x+(-c*d)^{(1/2)}/d)^{2*b-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)*b/d+(a*d-b*c)/d)^{(3/2)}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{2}}(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^3,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(5/2)*(d*x^2 + c)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{5/2}(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)^3),x)

[Out] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)^3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(5/2)/(d*x**2+c)**3,x)

[Out] Timed out

$$3.97 \quad \int \frac{(a+bx^2)^3}{(c+dx^2)^{11/2}} dx$$

Optimal. Leaf size=224

$$\frac{16a^3x(9bc-8ad)}{315c^5\sqrt{c+dx^2}(bc-ad)} + \frac{8a^2x(a+bx^2)(9bc-8ad)}{315c^4(c+dx^2)^{3/2}(bc-ad)} + \frac{2ax(a+bx^2)^2(9bc-8ad)}{105c^3(c+dx^2)^{5/2}(bc-ad)} + \frac{x(a+bx^2)^3(9bc-8ad)}{63c^2(c+dx^2)^{7/2}(bc-ad)}$$

[Out] $-1/9*d*x*(b*x^2+a)^4/c/(-a*d+b*c)/(d*x^2+c)^{(9/2)}+1/63*(-8*a*d+9*b*c)*x*(b*x^2+a)^3/c^2/(-a*d+b*c)/(d*x^2+c)^{(7/2)}+2/105*a*(-8*a*d+9*b*c)*x*(b*x^2+a)^2/c^3/(-a*d+b*c)/(d*x^2+c)^{(5/2)}+8/315*a^2*(-8*a*d+9*b*c)*x*(b*x^2+a)/c^4/(-a*d+b*c)/(d*x^2+c)^{(3/2)}+16/315*a^3*(-8*a*d+9*b*c)*x/c^5/(-a*d+b*c)/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 224, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {382, 378, 191}

$$\frac{8a^2x(a+bx^2)(9bc-8ad)}{315c^4(c+dx^2)^{3/2}(bc-ad)} + \frac{16a^3x(9bc-8ad)}{315c^5\sqrt{c+dx^2}(bc-ad)} + \frac{x(a+bx^2)^3(9bc-8ad)}{63c^2(c+dx^2)^{7/2}(bc-ad)} + \frac{2ax(a+bx^2)^2(9bc-8ad)}{105c^3(c+dx^2)^{5/2}(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^3/(c + d*x^2)^(11/2), x]

[Out] $-(d*x*(a+b*x^2)^4)/(9*c*(b*c-a*d)*(c+d*x^2)^{(9/2)}) + ((9*b*c-8*a*d)*x*(a+b*x^2)^3)/(63*c^2*(b*c-a*d)*(c+d*x^2)^{(7/2)}) + (2*a*(9*b*c-8*a*d)*x*(a+b*x^2)^2)/(105*c^3*(b*c-a*d)*(c+d*x^2)^{(5/2)}) + (8*a^2*(9*b*c-8*a*d)*x*(a+b*x^2))/(315*c^4*(b*c-a*d)*(c+d*x^2)^{(3/2)}) + (16*a^3*(9*b*c-8*a*d)*x)/(315*c^5*(b*c-a*d)*\text{Sqrt}[c+d*x^2])$

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*n*(p + 1)), x] - Dist[(c*q)/(a*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p + q + 1) + 1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[(b*c + n*(p + 1)*(b*c - a*d))/(a*n*(p + 1)*(b*c - a*d)], Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p + q + 2) + 1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^3}{(c+dx^2)^{11/2}} dx &= -\frac{dx(a+bx^2)^4}{9c(bc-ad)(c+dx^2)^{9/2}} + \frac{(9bc-8ad) \int \frac{(a+bx^2)^3}{(c+dx^2)^{9/2}} dx}{9c(bc-ad)} \\
&= -\frac{dx(a+bx^2)^4}{9c(bc-ad)(c+dx^2)^{9/2}} + \frac{(9bc-8ad)x(a+bx^2)^3}{63c^2(bc-ad)(c+dx^2)^{7/2}} + \frac{(2a(9bc-8ad)) \int \frac{(a+bx^2)^2}{(c+dx^2)^{7/2}} dx}{21c^2(bc-ad)} \\
&= -\frac{dx(a+bx^2)^4}{9c(bc-ad)(c+dx^2)^{9/2}} + \frac{(9bc-8ad)x(a+bx^2)^3}{63c^2(bc-ad)(c+dx^2)^{7/2}} + \frac{2a(9bc-8ad)x(a+bx^2)^2}{105c^3(bc-ad)(c+dx^2)^{5/2}} + \dots \\
&= -\frac{dx(a+bx^2)^4}{9c(bc-ad)(c+dx^2)^{9/2}} + \frac{(9bc-8ad)x(a+bx^2)^3}{63c^2(bc-ad)(c+dx^2)^{7/2}} + \frac{2a(9bc-8ad)x(a+bx^2)^2}{105c^3(bc-ad)(c+dx^2)^{5/2}} + \dots \\
&= -\frac{dx(a+bx^2)^4}{9c(bc-ad)(c+dx^2)^{9/2}} + \frac{(9bc-8ad)x(a+bx^2)^3}{63c^2(bc-ad)(c+dx^2)^{7/2}} + \frac{2a(9bc-8ad)x(a+bx^2)^2}{105c^3(bc-ad)(c+dx^2)^{5/2}} + \dots
\end{aligned}$$

Mathematica [A] time = 0.10, size = 163, normalized size = 0.73

$$\frac{a^3 (315c^4x + 840c^3dx^3 + 1008c^2d^2x^5 + 576cd^3x^7 + 128d^4x^9) + 3a^2bcx^3 (105c^3 + 126c^2dx^2 + 72cd^2x^4 + 16d^3x^6) + a^3 (315c^4x + 840c^3dx^3 + 1008c^2d^2x^5 + 576cd^3x^7 + 128d^4x^9)}{315c^5 (c + dx^2)^{9/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^3/(c + d*x^2)^(11/2), x]

[Out] (5*b^3*c^3*x^7*(9*c + 2*d*x^2) + 3*a*b^2*c^2*x^5*(63*c^2 + 36*c*d*x^2 + 8*d^2*x^4) + 3*a^2*b*c*x^3*(105*c^3 + 126*c^2*d*x^2 + 72*c*d^2*x^4 + 16*d^3*x^6) + a^3*(315*c^4*x + 840*c^3*d*x^3 + 1008*c^2*d^2*x^5 + 576*c*d^3*x^7 + 128*d^4*x^9))/(315*c^5*(c + d*x^2)^(9/2))

fricas [A] time = 1.02, size = 229, normalized size = 1.02

$$\frac{(2(5b^3c^3d + 12ab^2c^2d^2 + 24a^2bcd^3 + 64a^3d^4)x^9 + 315a^3c^4x + 9(5b^3c^4 + 12ab^2c^3d + 24a^2bc^2d^2 + 64a^3cd^3))}{315(c^5d^5x^{10} + 5c^6d^4x^8 + 10c^7d^3x^6 + 10c^8d^2x^4 + 5c^9d^1x^2 + c^{10})}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c)^(11/2), x, algorithm="fricas")

[Out] 1/315*(2*(5*b^3*c^3*d + 12*a*b^2*c^2*d^2 + 24*a^2*b*c*d^3 + 64*a^3*d^4)*x^9 + 315*a^3*c^4*x + 9*(5*b^3*c^4 + 12*a*b^2*c^3*d + 24*a^2*b*c^2*d^2 + 64*a^3*c*d^3)*x^7 + 63*(3*a*b^2*c^4 + 6*a^2*b*c^3*d + 16*a^3*c^2*d^2)*x^5 + 105*(3*a^2*b*c^4 + 8*a^3*c^3*d)*x^3)*sqrt(d*x^2 + c)/(c^5*d^5*x^10 + 5*c^6*d^4*x^8 + 10*c^7*d^3*x^6 + 10*c^8*d^2*x^4 + 5*c^9*d*x^2 + c^10)

giac [A] time = 0.68, size = 218, normalized size = 0.97

$$\frac{\left(\left(x^2 \left(\frac{2(5b^3c^3d^5 + 12ab^2c^2d^6 + 24a^2bcd^7 + 64a^3d^8)x^2}{c^5d^4} + \frac{9(5b^3c^4d^4 + 12ab^2c^3d^5 + 24a^2bc^2d^6 + 64a^3cd^7)}{c^5d^4} \right) \right) + \frac{63(3ab^2c^4d^4 + 6a^2bc^3d^5 + 16a^3c^2d^6)}{c^5d^4} \right)}{315(dx^2 + c)^{9/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^3/(d*x^2+c)^(11/2),x, algorithm="giac")
```

```
[Out] 1/315*(((x^2*(2*(5*b^3*c^3*d^5 + 12*a*b^2*c^2*d^6 + 24*a^2*b*c*d^7 + 64*a^3*d^8))*x^2/(c^5*d^4) + 9*(5*b^3*c^4*d^4 + 12*a*b^2*c^3*d^5 + 24*a^2*b*c^2*d^6 + 64*a^3*c*d^7)/(c^5*d^4)) + 63*(3*a*b^2*c^4*d^4 + 6*a^2*b*c^3*d^5 + 16*a^3*c^2*d^6)/(c^5*d^4))*x^2 + 105*(3*a^2*b*c^4*d^4 + 8*a^3*c^3*d^5)/(c^5*d^4))*x^2 + 315*a^3/c)*x/(d*x^2 + c)^(9/2)
```

```
maple [A] time = 0.01, size = 190, normalized size = 0.85
```

$$\frac{(128a^3d^4x^8 + 48a^2bcd^3x^8 + 24ab^2c^2d^2x^8 + 10b^3c^3dx^8 + 576a^3cd^3x^6 + 216a^2b^2c^2d^2x^6 + 108ab^2c^3dx^6 + 45b^3c^4x^6)}{315(dx^2 + c)^{\frac{9}{2}}c^5}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*x^2+a)^3/(d*x^2+c)^(11/2),x)
```

```
[Out] 1/315*x*(128*a^3*d^4*x^8+48*a^2*b*c*d^3*x^8+24*a*b^2*c^2*d^2*x^8+10*b^3*c^3*d*x^8+576*a^3*c*d^3*x^6+216*a^2*b*c^2*d^2*x^6+108*a*b^2*c^3*d*x^6+45*b^3*c^4*x^6+1008*a^3*c^2*d^2*x^4+378*a^2*b*c^3*d*x^4+189*a*b^2*c^4*x^4+840*a^3*c^3*d*x^2+315*a^2*b*c^4*x^2+315*a^3*c^4)/(d*x^2+c)^(9/2)/c^5
```

```
maxima [B] time = 1.51, size = 465, normalized size = 2.08
```

$$-\frac{b^3x^5}{4(dx^2 + c)^{\frac{9}{2}}d} - \frac{5b^3cx^3}{24(dx^2 + c)^{\frac{9}{2}}d^2} - \frac{ab^2x^3}{2(dx^2 + c)^{\frac{9}{2}}d} + \frac{128a^3x}{315\sqrt{dx^2 + c}c^5} + \frac{64a^3x}{315(dx^2 + c)^{\frac{3}{2}}c^4} + \frac{16a^3x}{105(dx^2 + c)^{\frac{5}{2}}c^3} + \frac{8}{63(dx^2 + c)^{\frac{7}{2}}c^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^3/(d*x^2+c)^(11/2),x, algorithm="maxima")
```

```
[Out] -1/4*b^3*x^5/((d*x^2 + c)^(9/2)*d) - 5/24*b^3*c*x^3/((d*x^2 + c)^(9/2)*d^2) - 1/2*a*b^2*x^3/((d*x^2 + c)^(9/2)*d) + 128/315*a^3*x/(sqrt(d*x^2 + c)*c^5) + 64/315*a^3*x/((d*x^2 + c)^(3/2)*c^4) + 16/105*a^3*x/((d*x^2 + c)^(5/2)*c^3) + 8/63*a^3*x/((d*x^2 + c)^(7/2)*c^2) + 1/9*a^3*x/((d*x^2 + c)^(9/2)*c) + 1/84*b^3*x/((d*x^2 + c)^(5/2)*d^3) + 2/63*b^3*x/(sqrt(d*x^2 + c)*c^2*d^3) + 1/63*b^3*x/((d*x^2 + c)^(3/2)*c*d^3) + 5/504*b^3*c*x/((d*x^2 + c)^(7/2)*d^3) - 5/72*b^3*c^2*x/((d*x^2 + c)^(9/2)*d^3) + 1/42*a*b^2*x/((d*x^2 + c)^(7/2)*d^2) + 8/105*a*b^2*x/(sqrt(d*x^2 + c)*c^3*d^2) + 4/105*a*b^2*x/((d*x^2 + c)^(3/2)*c^2*d^2) + 1/35*a*b^2*x/((d*x^2 + c)^(5/2)*c*d^2) - 1/6*a*b^2*c*x/((d*x^2 + c)^(9/2)*d^2) - 1/3*a^2*b*x/((d*x^2 + c)^(9/2)*d) + 16/105*a^2*b*x/(sqrt(d*x^2 + c)*c^4*d) + 8/105*a^2*b*x/((d*x^2 + c)^(3/2)*c^3*d) + 2/35*a^2*b*x/((d*x^2 + c)^(5/2)*c^2*d) + 1/21*a^2*b*x/((d*x^2 + c)^(7/2)*c*d)
```

```
mupad [B] time = 5.15, size = 326, normalized size = 1.46
```

$$\frac{x \left(\frac{a^3}{9c} - \frac{c \left(\frac{b^3}{9d} - \frac{ab^2}{3c} \right) + \frac{a^2b}{3c}}{d} \right)}{(dx^2 + c)^{9/2}} - \frac{x \left(\frac{b^3}{5d^3} - \frac{16a^3d^3 + 6a^2bcd^2 + 3ab^2c^2d - 4b^3c^3}{105c^3d^3} \right)}{(dx^2 + c)^{5/2}} + \frac{x \left(\frac{c \left(\frac{b^3}{7d^2} - \frac{b^2(3ad-bc)}{7cd^2} \right) + \frac{8a^3d^3 + 3a^2bcd^2 - 3ab^2c^2d}{63c^2d^3}}{d} \right)}{(dx^2 + c)^{7/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a + b*x^2)^3/(c + d*x^2)^(11/2),x)
```

```
[Out] (x*(a^3/(9*c) - (c*((c*(b^3/(9*d) - (a*b^2)/(3*c)))/d + (a^2*b)/(3*c)))/d))
/(c + d*x^2)^(9/2) - (x*(b^3/(5*d^3) - (16*a^3*d^3 - 4*b^3*c^3 + 3*a*b^2*c^
2*d + 6*a^2*b*c*d^2)/(105*c^3*d^3)))/(c + d*x^2)^(5/2) + (x*((c*(b^3/(7*d^2
) - (b^2*(3*a*d - b*c))/(7*c*d^2)))/d + (8*a^3*d^3 + b^3*c^3 - 3*a*b^2*c^2*
d + 3*a^2*b*c*d^2)/(63*c^2*d^3)))/(c + d*x^2)^(7/2) + (x*(64*a^3*d^3 + 5*b^
3*c^3 + 12*a*b^2*c^2*d + 24*a^2*b*c*d^2))/(315*c^4*d^3*(c + d*x^2)^(3/2)) +
(x*(128*a^3*d^3 + 10*b^3*c^3 + 24*a*b^2*c^2*d + 48*a^2*b*c*d^2))/(315*c^5*
d^3*(c + d*x^2)^(1/2))
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)**3/(d*x**2+c)**(11/2),x)
```

[Out] Timed out

$$3.98 \quad \int \frac{(a+bx^2)^2}{(c+dx^2)^{9/2}} dx$$

Optimal. Leaf size=174

$$\frac{8a^2x(7bc-6ad)}{105c^4\sqrt{c+dx^2}(bc-ad)} + \frac{4ax(a+bx^2)(7bc-6ad)}{105c^3(c+dx^2)^{3/2}(bc-ad)} + \frac{x(a+bx^2)^2(7bc-6ad)}{35c^2(c+dx^2)^{5/2}(bc-ad)} - \frac{dx(a+bx^2)^3}{7c(c+dx^2)^{7/2}(bc-ad)}$$

[Out] $-1/7*d*x*(b*x^2+a)^3/c/(-a*d+b*c)/(d*x^2+c)^{(7/2)}+1/35*(-6*a*d+7*b*c)*x*(b*x^2+a)^2/c^2/(-a*d+b*c)/(d*x^2+c)^{(5/2)}+4/105*a*(-6*a*d+7*b*c)*x*(b*x^2+a)/c^3/(-a*d+b*c)/(d*x^2+c)^{(3/2)}+8/105*a^2*(-6*a*d+7*b*c)*x/c^4/(-a*d+b*c)/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 174, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {382, 378, 191}

$$\frac{8a^2x(7bc-6ad)}{105c^4\sqrt{c+dx^2}(bc-ad)} + \frac{x(a+bx^2)^2(7bc-6ad)}{35c^2(c+dx^2)^{5/2}(bc-ad)} + \frac{4ax(a+bx^2)(7bc-6ad)}{105c^3(c+dx^2)^{3/2}(bc-ad)} - \frac{dx(a+bx^2)^3}{7c(c+dx^2)^{7/2}(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^2/(c + d*x^2)^(9/2), x]

[Out] $-(d*x*(a + b*x^2)^3)/(7*c*(b*c - a*d)*(c + d*x^2)^{(7/2)}) + ((7*b*c - 6*a*d)*x*(a + b*x^2)^2)/(35*c^2*(b*c - a*d)*(c + d*x^2)^{(5/2)}) + (4*a*(7*b*c - 6*a*d)*x*(a + b*x^2))/(105*c^3*(b*c - a*d)*(c + d*x^2)^{(3/2)}) + (8*a^2*(7*b*c - 6*a*d)*x)/(105*c^4*(b*c - a*d)*\text{Sqrt}[c + d*x^2])$

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*n*(p + 1)), x] - Dist[(c*q)/(a*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p + q + 1) + 1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[(b*c + n*(p + 1)*(b*c - a*d))/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p + q + 2) + 1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^2}{(c+dx^2)^{9/2}} dx &= -\frac{dx(a+bx^2)^3}{7c(bc-ad)(c+dx^2)^{7/2}} + \frac{(7bc-6ad) \int \frac{(a+bx^2)^2}{(c+dx^2)^{7/2}} dx}{7c(bc-ad)} \\
&= -\frac{dx(a+bx^2)^3}{7c(bc-ad)(c+dx^2)^{7/2}} + \frac{(7bc-6ad)x(a+bx^2)^2}{35c^2(bc-ad)(c+dx^2)^{5/2}} + \frac{(4a(7bc-6ad)) \int \frac{a+bx^2}{(c+dx^2)^{5/2}} dx}{35c^2(bc-ad)} \\
&= -\frac{dx(a+bx^2)^3}{7c(bc-ad)(c+dx^2)^{7/2}} + \frac{(7bc-6ad)x(a+bx^2)^2}{35c^2(bc-ad)(c+dx^2)^{5/2}} + \frac{4a(7bc-6ad)x(a+bx^2)}{105c^3(bc-ad)(c+dx^2)^{3/2}} + \dots \\
&= -\frac{dx(a+bx^2)^3}{7c(bc-ad)(c+dx^2)^{7/2}} + \frac{(7bc-6ad)x(a+bx^2)^2}{35c^2(bc-ad)(c+dx^2)^{5/2}} + \frac{4a(7bc-6ad)x(a+bx^2)}{105c^3(bc-ad)(c+dx^2)^{3/2}} + \dots
\end{aligned}$$

Mathematica [A] time = 0.07, size = 107, normalized size = 0.61

$$\frac{3a^2(35c^3x + 70c^2dx^3 + 56cd^2x^5 + 16d^3x^7) + 2abcx^3(35c^2 + 28cdx^2 + 8d^2x^4) + 3b^2c^2x^5(7c + 2dx^2)}{105c^4(c+dx^2)^{7/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^2/(c + d*x^2)^(9/2), x]

[Out] (3*b^2*c^2*x^5*(7*c + 2*d*x^2) + 2*a*b*c*x^3*(35*c^2 + 28*c*d*x^2 + 8*d^2*x^4) + 3*a^2*(35*c^3*x + 70*c^2*d*x^3 + 56*c*d^2*x^5 + 16*d^3*x^7))/(105*c^4*(c + d*x^2)^(7/2))

fricas [A] time = 0.64, size = 151, normalized size = 0.87

$$\frac{(2(3b^2c^2d + 8abcd^2 + 24a^2d^3)x^7 + 105a^2c^3x + 7(3b^2c^3 + 8abc^2d + 24a^2cd^2)x^5 + 70(abc^3 + 3a^2c^2d)x^3)\sqrt{c+dx^2}}{105(c^4d^4x^8 + 4c^5d^3x^6 + 6c^6d^2x^4 + 4c^7dx^2 + c^8)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)^(9/2), x, algorithm="fricas")

[Out] 1/105*(2*(3*b^2*c^2*d + 8*a*b*c*d^2 + 24*a^2*d^3)*x^7 + 105*a^2*c^3*x + 7*(3*b^2*c^3 + 8*a*b*c^2*d + 24*a^2*c*d^2)*x^5 + 70*(a*b*c^3 + 3*a^2*c^2*d)*x^3)*sqrt(d*x^2 + c)/(c^4*d^4*x^8 + 4*c^5*d^3*x^6 + 6*c^6*d^2*x^4 + 4*c^7*d*x^2 + c^8)

giac [A] time = 0.64, size = 138, normalized size = 0.79

$$\frac{\left(x^2 \left(\frac{2(3b^2c^2d^4 + 8abcd^5 + 24a^2d^6)x^2}{c^4d^3} + \frac{7(3b^2c^3d^3 + 8abc^2d^4 + 24a^2cd^5)}{c^4d^3} \right) + \frac{70(abc^3d^3 + 3a^2c^2d^4)}{c^4d^3} \right) x^2 + \frac{105a^2}{c} x}{105(dx^2 + c)^{7/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)^(9/2), x, algorithm="giac")

[Out] 1/105*((x^2*(2*(3*b^2*c^2*d^4 + 8*a*b*c*d^5 + 24*a^2*d^6)*x^2/(c^4*d^3) + 7*(3*b^2*c^3*d^3 + 8*a*b*c^2*d^4 + 24*a^2*c*d^5)/(c^4*d^3)) + 70*(a*b*c^3*d^3 + 3*a^2*c^2*d^4)/(c^4*d^3))*x^2 + 105*a^2/c)*x/(d*x^2 + c)^(7/2)

maple [A] time = 0.01, size = 115, normalized size = 0.66

$$\frac{(48a^2d^3x^6 + 16abc d^2x^6 + 6b^2c^2d x^6 + 168a^2c d^2x^4 + 56ab c^2d x^4 + 21b^2c^3x^4 + 210a^2c^2d x^2 + 70ab c^3x^2 + 105a^2c^3)}{105(dx^2 + c)^{\frac{7}{2}}c^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^2/(d*x^2+c)^(9/2),x)

[Out] 1/105*x*(48*a^2*d^3*x^6+16*a*b*c*d^2*x^6+6*b^2*c^2*d*x^6+168*a^2*c*d^2*x^4+56*a*b*c^2*d*x^4+21*b^2*c^3*x^4+210*a^2*c^2*d*x^2+70*a*b*c^3*x^2+105*a^2*c^3)/(d*x^2+c)^(7/2)/c^4

maxima [A] time = 1.50, size = 249, normalized size = 1.43

$$-\frac{b^2x^3}{4(dx^2 + c)^{\frac{7}{2}}d} + \frac{16a^2x}{35\sqrt{dx^2 + c}c^4} + \frac{8a^2x}{35(dx^2 + c)^{\frac{3}{2}}c^3} + \frac{6a^2x}{35(dx^2 + c)^{\frac{5}{2}}c^2} + \frac{a^2x}{7(dx^2 + c)^{\frac{7}{2}}c} + \frac{3b^2x}{140(dx^2 + c)^{\frac{5}{2}}d^2} + \frac{2b^2}{35\sqrt{dx^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)^(9/2),x, algorithm="maxima")

[Out] -1/4*b^2*x^3/((d*x^2 + c)^(7/2)*d) + 16/35*a^2*x/(sqrt(d*x^2 + c)*c^4) + 8/35*a^2*x/((d*x^2 + c)^(3/2)*c^3) + 6/35*a^2*x/((d*x^2 + c)^(5/2)*c^2) + 1/7*a^2*x/((d*x^2 + c)^(7/2)*c) + 3/140*b^2*x/((d*x^2 + c)^(5/2)*d^2) + 2/35*b^2*x/(sqrt(d*x^2 + c)*c^2*d^2) + 1/35*b^2*x/((d*x^2 + c)^(3/2)*c*d^2) - 3/2*8*b^2*c*x/((d*x^2 + c)^(7/2)*d^2) - 2/7*a*b*x/((d*x^2 + c)^(7/2)*d) + 16/105*a*b*x/(sqrt(d*x^2 + c)*c^3*d) + 8/105*a*b*x/((d*x^2 + c)^(3/2)*c^2*d) + 2/35*a*b*x/((d*x^2 + c)^(5/2)*c*d)

mupad [B] time = 4.99, size = 176, normalized size = 1.01

$$\frac{x \left(\frac{a^2}{7c} + \frac{c \left(\frac{b^2}{7d} - \frac{2ab}{7c} \right)}{d} \right)}{(dx^2 + c)^{7/2}} - \frac{x \left(\frac{b^2}{5d^2} - \frac{6a^2d^2 + 2abcd - b^2c^2}{35c^2d^2} \right)}{(dx^2 + c)^{5/2}} + \frac{x (24a^2d^2 + 8abcd + 3b^2c^2)}{105c^3d^2(dx^2 + c)^{3/2}} + \frac{x (48a^2d^2 + 16abcd + 6b^2c^2)}{105c^4d^2\sqrt{dx^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^2/(c + d*x^2)^(9/2),x)

[Out] (x*(a^2/(7*c) + (c*(b^2/(7*d) - (2*a*b)/(7*c)))/d))/(c + d*x^2)^(7/2) - (x*(b^2/(5*d^2) - (6*a^2*d^2 - b^2*c^2 + 2*a*b*c*d)/(35*c^2*d^2)))/(c + d*x^2)^(5/2) + (x*(24*a^2*d^2 + 3*b^2*c^2 + 8*a*b*c*d))/(105*c^3*d^2*(c + d*x^2)^(3/2)) + (x*(48*a^2*d^2 + 6*b^2*c^2 + 16*a*b*c*d))/(105*c^4*d^2*(c + d*x^2)^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^2}{(c + dx^2)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**2/(d*x**2+c)**(9/2),x)

[Out] Integral((a + b*x**2)**2/(c + d*x**2)**(9/2), x)

$$3.99 \quad \int \frac{a+bx^2}{(c+dx^2)^{7/2}} dx$$

Optimal. Leaf size=91

$$\frac{2x(4ad+bc)}{15c^3d\sqrt{c+dx^2}} + \frac{x(4ad+bc)}{15c^2d(c+dx^2)^{3/2}} - \frac{x(bc-ad)}{5cd(c+dx^2)^{5/2}}$$

[Out] $-1/5*(-a*d+b*c)*x/c/d/(d*x^2+c)^{(5/2)}+1/15*(4*a*d+b*c)*x/c^2/d/(d*x^2+c)^{(3/2)}+2/15*(4*a*d+b*c)*x/c^3/d/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {385, 192, 191}

$$\frac{2x(4ad+bc)}{15c^3d\sqrt{c+dx^2}} + \frac{x(4ad+bc)}{15c^2d(c+dx^2)^{3/2}} - \frac{x(bc-ad)}{5cd(c+dx^2)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)/(c + d*x^2)^(7/2), x]

[Out] $-((b*c - a*d)*x)/(5*c*d*(c + d*x^2)^{(5/2)}) + ((b*c + 4*a*d)*x)/(15*c^2*d*(c + d*x^2)^{(3/2)}) + (2*(b*c + 4*a*d)*x)/(15*c^3*d*\text{Sqrt}[c + d*x^2])$

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rubi steps

$$\begin{aligned} \int \frac{a+bx^2}{(c+dx^2)^{7/2}} dx &= -\frac{(bc-ad)x}{5cd(c+dx^2)^{5/2}} + \frac{(bc+4ad) \int \frac{1}{(c+dx^2)^{5/2}} dx}{5cd} \\ &= -\frac{(bc-ad)x}{5cd(c+dx^2)^{5/2}} + \frac{(bc+4ad)x}{15c^2d(c+dx^2)^{3/2}} + \frac{(2(bc+4ad)) \int \frac{1}{(c+dx^2)^{3/2}} dx}{15c^2d} \\ &= -\frac{(bc-ad)x}{5cd(c+dx^2)^{5/2}} + \frac{(bc+4ad)x}{15c^2d(c+dx^2)^{3/2}} + \frac{2(bc+4ad)x}{15c^3d\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 59, normalized size = 0.65

$$\frac{a(15c^2x + 20cdx^3 + 8d^2x^5) + bcx^3(5c + 2dx^2)}{15c^3(c + dx^2)^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)/(c + d*x^2)^(7/2), x]

[Out] (b*c*x^3*(5*c + 2*d*x^2) + a*(15*c^2*x + 20*c*d*x^3 + 8*d^2*x^5))/(15*c^3*(c + d*x^2)^(5/2))

fricas [A] time = 0.59, size = 87, normalized size = 0.96

$$\frac{(2(bcd + 4ad^2)x^5 + 15ac^2x + 5(bc^2 + 4acd)x^3)\sqrt{dx^2 + c}}{15(c^3d^3x^6 + 3c^4d^2x^4 + 3c^5dx^2 + c^6)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^(7/2), x, algorithm="fricas")

[Out] 1/15*(2*(b*c*d + 4*a*d^2)*x^5 + 15*a*c^2*x + 5*(b*c^2 + 4*a*c*d)*x^3)*sqrt(d*x^2 + c)/(c^3*d^3*x^6 + 3*c^4*d^2*x^4 + 3*c^5*d*x^2 + c^6)

giac [A] time = 0.62, size = 72, normalized size = 0.79

$$\frac{\left(x^2\left(\frac{2(bcd^3+4ad^4)x^2}{c^3d^2} + \frac{5(bc^2d^2+4acd^3)}{c^3d^2}\right) + \frac{15a}{c}\right)x}{15(dx^2 + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^(7/2), x, algorithm="giac")

[Out] 1/15*(x^2*(2*(b*c*d^3 + 4*a*d^4)*x^2/(c^3*d^2) + 5*(b*c^2*d^2 + 4*a*c*d^3)/(c^3*d^2)) + 15*a/c)*x/(d*x^2 + c)^(5/2)

maple [A] time = 0.00, size = 57, normalized size = 0.63

$$\frac{(8ad^2x^4 + 2bcdx^4 + 20acd^2x^2 + 5bc^2x^2 + 15c^2a)x}{15(dx^2 + c)^{\frac{5}{2}}c^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)/(d*x^2+c)^(7/2), x)

[Out] 1/15*x*(8*a*d^2*x^4+2*b*c*d*x^4+20*a*c*d*x^2+5*b*c^2*x^2+15*a*c^2)/(d*x^2+c)^(5/2)/c^3

maxima [A] time = 1.45, size = 103, normalized size = 1.13

$$\frac{8ax}{15\sqrt{dx^2 + c}c^3} + \frac{4ax}{15(dx^2 + c)^{\frac{3}{2}}c^2} + \frac{ax}{5(dx^2 + c)^{\frac{5}{2}}c} - \frac{bx}{5(dx^2 + c)^{\frac{5}{2}}d} + \frac{2bx}{15\sqrt{dx^2 + c}c^2d} + \frac{bx}{15(dx^2 + c)^{\frac{3}{2}}cd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)^(7/2), x, algorithm="maxima")

[Out] $8/15*a*x/(sqrt(dx^2 + c)*c^3) + 4/15*a*x/((dx^2 + c)^{(3/2)}*c^2) + 1/5*a*x/((dx^2 + c)^{(5/2)}*c) - 1/5*b*x/((dx^2 + c)^{(5/2)}*d) + 2/15*b*x/(sqrt(dx^2 + c)*c^2*d) + 1/15*b*x/((dx^2 + c)^{(3/2)}*c*d)$

mupad [B] time = 4.85, size = 87, normalized size = 0.96

$$\frac{8adx(dx^2 + c)^2 - 3bc^3x + 2bcx(dx^2 + c)^2 + bc^2x(dx^2 + c) + 3ac^2dx + 4acdx(dx^2 + c)}{15c^3d(dx^2 + c)^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)/(c + d*x^2)^(7/2), x)`

[Out] $(8*a*d*x*(c + d*x^2)^2 - 3*b*c^3*x + 2*b*c*x*(c + d*x^2)^2 + b*c^2*x*(c + d*x^2) + 3*a*c^2*d*x + 4*a*c*d*x*(c + d*x^2))/(15*c^3*d*(c + d*x^2)^{(5/2)})$

sympy [B] time = 27.99, size = 566, normalized size = 6.22

$$a \left(\frac{15c^5x}{15c^{\frac{17}{2}}\sqrt{1 + \frac{dx^2}{c}} + 45c^{\frac{15}{2}}dx^2\sqrt{1 + \frac{dx^2}{c}} + 45c^{\frac{13}{2}}d^2x^4\sqrt{1 + \frac{dx^2}{c}} + 15c^{\frac{11}{2}}d^3x^6\sqrt{1 + \frac{dx^2}{c}}} + \frac{1}{15c^{\frac{17}{2}}\sqrt{1 + \frac{dx^2}{c}} + 45c^{\frac{15}{2}}dx^2\sqrt{1 + \frac{dx^2}{c}} + 45c^{\frac{13}{2}}d^2x^4\sqrt{1 + \frac{dx^2}{c}} + 15c^{\frac{11}{2}}d^3x^6\sqrt{1 + \frac{dx^2}{c}}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)/(d*x**2+c)**(7/2), x)`

[Out] $a*(15*c**5*x/(15*c**(17/2)*sqrt(1 + d*x**2/c) + 45*c**(15/2)*d*x**2*sqrt(1 + d*x**2/c) + 45*c**(13/2)*d**2*x**4*sqrt(1 + d*x**2/c) + 15*c**(11/2)*d**3*x**6*sqrt(1 + d*x**2/c)) + 35*c**4*d*x**3/(15*c**(17/2)*sqrt(1 + d*x**2/c) + 45*c**(15/2)*d*x**2*sqrt(1 + d*x**2/c) + 45*c**(13/2)*d**2*x**4*sqrt(1 + d*x**2/c) + 15*c**(11/2)*d**3*x**6*sqrt(1 + d*x**2/c)) + 28*c**3*d*x**5/(15*c**(17/2)*sqrt(1 + d*x**2/c) + 45*c**(15/2)*d*x**2*sqrt(1 + d*x**2/c) + 45*c**(13/2)*d**2*x**4*sqrt(1 + d*x**2/c) + 15*c**(11/2)*d**3*x**6*sqrt(1 + d*x**2/c)) + 8*c**2*d**3*x**7/(15*c**(17/2)*sqrt(1 + d*x**2/c) + 45*c**(15/2)*d*x**2*sqrt(1 + d*x**2/c) + 45*c**(13/2)*d**2*x**4*sqrt(1 + d*x**2/c) + 15*c**(11/2)*d**3*x**6*sqrt(1 + d*x**2/c)) + b*(5*c*x**3/(15*c**(9/2)*sqrt(1 + d*x**2/c) + 30*c**(7/2)*d*x**2*sqrt(1 + d*x**2/c) + 15*c**(5/2)*d**2*x**4*sqrt(1 + d*x**2/c)) + 2*d*x**5/(15*c**(9/2)*sqrt(1 + d*x**2/c) + 30*c**(7/2)*d*x**2*sqrt(1 + d*x**2/c) + 15*c**(5/2)*d**2*x**4*sqrt(1 + d*x**2/c))$

$$3.100 \quad \int \frac{1}{(c+dx^2)^{5/2}} dx$$

Optimal. Leaf size=39

$$\frac{2x}{3c^2\sqrt{c+dx^2}} + \frac{x}{3c(c+dx^2)^{3/2}}$$

[Out] 1/3*x/c/(d*x^2+c)^(3/2)+2/3*x/c^2/(d*x^2+c)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {192, 191}

$$\frac{2x}{3c^2\sqrt{c+dx^2}} + \frac{x}{3c(c+dx^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^(-5/2), x]

[Out] x/(3*c*(c + d*x^2)^(3/2)) + (2*x)/(3*c^2*Sqrt[c + d*x^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(c+dx^2)^{5/2}} dx &= \frac{x}{3c(c+dx^2)^{3/2}} + \frac{2 \int \frac{1}{(c+dx^2)^{3/2}} dx}{3c} \\ &= \frac{x}{3c(c+dx^2)^{3/2}} + \frac{2x}{3c^2\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 29, normalized size = 0.74

$$\frac{x(3c + 2dx^2)}{3c^2(c + dx^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^(-5/2), x]

[Out] (x*(3*c + 2*d*x^2))/(3*c^2*(c + d*x^2)^(3/2))

fricas [A] time = 0.62, size = 47, normalized size = 1.21

$$\frac{(2dx^3 + 3cx)\sqrt{dx^2 + c}}{3(c^2d^2x^4 + 2c^3dx^2 + c^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*x^2+c)^(5/2),x, algorithm="fricas")

[Out] 1/3*(2*d*x^3 + 3*c*x)*sqrt(d*x^2 + c)/(c^2*d^2*x^4 + 2*c^3*d*x^2 + c^4)

giac [A] time = 0.60, size = 27, normalized size = 0.69

$$\frac{x\left(\frac{2dx^2}{c^2} + \frac{3}{c}\right)}{3(dx^2 + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*x^2+c)^(5/2),x, algorithm="giac")

[Out] 1/3*x*(2*d*x^2/c^2 + 3/c)/(d*x^2 + c)^(3/2)

maple [A] time = 0.00, size = 26, normalized size = 0.67

$$\frac{(2dx^2 + 3c)x}{3(dx^2 + c)^{\frac{3}{2}}c^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*x^2+c)^(5/2),x)

[Out] 1/3*x*(2*d*x^2+3*c)/(d*x^2+c)^(3/2)/c^2

maxima [A] time = 1.34, size = 31, normalized size = 0.79

$$\frac{2x}{3\sqrt{dx^2 + c}c^2} + \frac{x}{3(dx^2 + c)^{\frac{3}{2}}c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*x^2+c)^(5/2),x, algorithm="maxima")

[Out] 2/3*x/(sqrt(d*x^2 + c)*c^2) + 1/3*x/((d*x^2 + c)^(3/2)*c)

mupad [B] time = 4.79, size = 28, normalized size = 0.72

$$\frac{2x(dx^2 + c) + cx}{3c^2(dx^2 + c)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c + d*x^2)^(5/2),x)

[Out] (2*x*(c + d*x^2) + c*x)/(3*c^2*(c + d*x^2)^(3/2))

sympy [B] time = 0.83, size = 95, normalized size = 2.44

$$\frac{3cx}{3c^{\frac{7}{2}}\sqrt{1 + \frac{dx^2}{c}} + 3c^{\frac{5}{2}}dx^2\sqrt{1 + \frac{dx^2}{c}}} + \frac{2dx^3}{3c^{\frac{7}{2}}\sqrt{1 + \frac{dx^2}{c}} + 3c^{\frac{5}{2}}dx^2\sqrt{1 + \frac{dx^2}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*x**2+c)**(5/2),x)
```

```
[Out] 3*c*x/(3*c**(7/2)*sqrt(1 + d*x**2/c) + 3*c**(5/2)*d*x**2*sqrt(1 + d*x**2/c)
) + 2*d*x**3/(3*c**(7/2)*sqrt(1 + d*x**2/c) + 3*c**(5/2)*d*x**2*sqrt(1 + d*
x**2/c))
```

$$3.101 \quad \int \frac{1}{(a+bx^2)(c+dx^2)^{3/2}} dx$$

Optimal. Leaf size=79

$$\frac{b \tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{\sqrt{a}(bc-ad)^{3/2}} - \frac{dx}{c\sqrt{c+dx^2}(bc-ad)}$$

[Out] b*arctan(x*(-a*d+b*c)^(1/2)/a^(1/2)/(d*x^2+c)^(1/2))/(-a*d+b*c)^(3/2)/a^(1/2)-d*x/c/(-a*d+b*c)/(d*x^2+c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {382, 377, 205}

$$\frac{b \tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{\sqrt{a}(bc-ad)^{3/2}} - \frac{dx}{c\sqrt{c+dx^2}(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)*(c + d*x^2)^(3/2)), x]

[Out] -((d*x)/(c*(b*c - a*d)*Sqrt[c + d*x^2])) + (b*ArcTan[(Sqrt[b*c - a*d]*x)/(Sqrt[a]*Sqrt[c + d*x^2])])/(Sqrt[a]*(b*c - a*d)^(3/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)], Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+2)+1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)(c+dx^2)^{3/2}} dx &= -\frac{dx}{c(bc-ad)\sqrt{c+dx^2}} + \frac{b \int \frac{1}{(a+bx^2)\sqrt{c+dx^2}} dx}{bc-ad} \\ &= -\frac{dx}{c(bc-ad)\sqrt{c+dx^2}} + \frac{b \operatorname{Subst}\left(\int \frac{1}{a-(-bc+ad)x^2} dx, x, \frac{x}{\sqrt{c+dx^2}}\right)}{bc-ad} \\ &= -\frac{dx}{c(bc-ad)\sqrt{c+dx^2}} + \frac{b \tan^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{a}\sqrt{c+dx^2}}\right)}{\sqrt{a}(bc-ad)^{3/2}} \end{aligned}$$

Mathematica [C] time = 2.72, size = 236, normalized size = 2.99

$$\frac{15c(3c+2dx^2) \left(c(a+bx^2) \sqrt{\frac{ax^2(c+dx^2)(bc-ad)}{c^2(a+bx^2)^2}} - a(c+dx^2) \sin^{-1}\left(\sqrt{\frac{x^2(bc-ad)}{c(a+bx^2)}}\right) \right)}{\sqrt{\frac{ax^2(c+dx^2)(bc-ad)}{c^2(a+bx^2)^2}}} + \frac{4x^4(c+dx^2)(bc-ad)^2 {}_2F_1\left(2, 2; \frac{7}{2}; \frac{(bc-ad)x^2}{c(bx^2+a)}\right)}{a+bx^2}$$

$$15c^3x(a+bx^2)\sqrt{c+dx^2}(ad-bc)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)*(c + d*x^2)^(3/2)), x]

[Out] $-1/15*((15*c*(3*c + 2*d*x^2)*(c*(a + b*x^2)*\operatorname{Sqrt}[(a*(b*c - a*d)*x^2*(c + d*x^2))/(c^2*(a + b*x^2)^2]] - a*(c + d*x^2)*\operatorname{ArcSin}[\operatorname{Sqrt}[(b*c - a*d)*x^2/(c*(a + b*x^2))]])/\operatorname{Sqrt}[(a*(b*c - a*d)*x^2*(c + d*x^2))/(c^2*(a + b*x^2)^2]] + (4*(b*c - a*d)^2*x^4*(c + d*x^2)*\operatorname{Hypergeometric2F1}[2, 2, 7/2, ((b*c - a*d)*x^2)/(c*(a + b*x^2))])/(a + b*x^2))/(c^3*(-(b*c) + a*d)*x*(a + b*x^2)*\operatorname{Sqrt}[c + d*x^2])$

fricas [B] time = 0.72, size = 442, normalized size = 5.59

$$\left[\frac{4(abcd - a^2d^2)\sqrt{dx^2 + c}x - (bcdx^2 + bc^2)\sqrt{-abc + a^2d} \log\left(\frac{(b^2c^2 - 8abcd + 8a^2d^2)x^4 + a^2c^2 - 2(3abc^2 - 4a^2cd)x^2 + 4((bc - 2ad)x^2 + a^2)}{b^2x^4 + 2abx^2 + a^2}\right)}{4(ab^2c^4 - 2a^2bc^3d + a^3c^2d^2 + (ab^2c^3d - 2a^2bc^2d^2 + a^3cd^3)x^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^(3/2), x, algorithm="fricas")

[Out] $[-1/4*(4*(a*b*c*d - a^2*d^2)*\operatorname{sqrt}(d*x^2 + c)*x - (b*c*d*x^2 + b*c^2)*\operatorname{sqrt}(-a*b*c + a^2*d)*\log(((b^2*c^2 - 8*a*b*c*d + 8*a^2*d^2)*x^4 + a^2*c^2 - 2*(3*a*b*c^2 - 4*a^2*c*d)*x^2 + 4*((b*c - 2*a*d)*x^3 - a*c*x)*\operatorname{sqrt}(-a*b*c + a^2*d)*\operatorname{sqrt}(d*x^2 + c))/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(a*b^2*c^4 - 2*a^2*b*c^3*d + a^3*c^2*d^2 + (a*b^2*c^3*d - 2*a^2*b*c^2*d^2 + a^3*c*d^3)*x^2), -1/2*(2*(a*b*c*d - a^2*d^2)*\operatorname{sqrt}(d*x^2 + c)*x - (b*c*d*x^2 + b*c^2)*\operatorname{sqrt}(a*b*c - a^2*d)*\operatorname{arctan}(1/2*\operatorname{sqrt}(a*b*c - a^2*d)*((b*c - 2*a*d)*x^2 - a*c)*\operatorname{sqrt}(d*x^2 + c))/((a*b*c*d - a^2*d^2)*x^3 + (a*b*c^2 - a^2*c*d)*x))/(a*b^2*c^4 - 2*a^2*b*c^3*d + a^3*c^2*d^2 + (a*b^2*c^3*d - 2*a^2*b*c^2*d^2 + a^3*c*d^3)*x^2)]$

giac [A] time = 0.62, size = 107, normalized size = 1.35

$$\frac{b\sqrt{d} \operatorname{arctan}\left(\frac{(\sqrt{d}x - \sqrt{dx^2+c})^2 b - bc + 2ad}{2\sqrt{abcd - a^2d^2}}\right)}{\sqrt{abcd - a^2d^2}(bc - ad)} - \frac{dx}{(bc^2 - acd)\sqrt{dx^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^(3/2),x, algorithm="giac")

[Out] $b\sqrt{d}\arctan\left(\frac{-1/2\left(\sqrt{d}x - \sqrt{d^2x^2 + c}\right)^2b - b^2c + 2ad}{\sqrt{a^2b^2cd - a^2d^2}}\right) / \left(\sqrt{a^2b^2cd - a^2d^2}\right) (b^2c - a^2d) - dx / \left(\left(b^2c^2 - a^2cd\right)\sqrt{d^2x^2 + c}\right)$

maple [B] time = 0.04, size = 628, normalized size = 7.95

$$b \ln \left(\frac{\frac{2\sqrt{-ab}\left(x - \frac{\sqrt{-ab}}{b}\right)^d}{b} - \frac{2(ad-bc)}{b} + 2\sqrt{-\frac{ad-bc}{b}} \sqrt{\left(x - \frac{\sqrt{-ab}}{b}\right)^2 + \frac{ad-bc}{b}}}{x - \frac{\sqrt{-ab}}{b}} \right) - b \ln \left(\frac{\frac{2\sqrt{-ab}\left(x + \frac{\sqrt{-ab}}{b}\right)^d}{b} - \frac{2(ad-bc)}{b} + 2\sqrt{-\frac{ad-bc}{b}} \sqrt{\left(x + \frac{\sqrt{-ab}}{b}\right)^2 + \frac{ad-bc}{b}}}{x + \frac{\sqrt{-ab}}{b}} \right)$$

$$\frac{2\sqrt{-ab} (ad - bc) \sqrt{-\frac{ad-bc}{b}}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)/(d*x^2+c)^(3/2),x)

[Out] $\frac{1}{2}(-a^2b)^{-1/2}/(ad-b^2c)*b/\left(\left(x+1/b*(-a^2b)^{-1/2}\right)^2d-2d*(-a^2b)^{-1/2}/b*\left(x+1/b*(-a^2b)^{-1/2}\right)-(ad-b^2c)/b\right)^{1/2}+1/2/(ad-b^2c)/c/\left(\left(x+1/b*(-a^2b)^{-1/2}\right)^2d-2d*(-a^2b)^{-1/2}/b*\left(x+1/b*(-a^2b)^{-1/2}\right)-(ad-b^2c)/b\right)^{1/2}*x*d-1/2/(-a^2b)^{-1/2}/(ad-b^2c)*b/\left(-\left(ad-b^2c\right)/b\right)^{1/2}*\ln\left(\frac{-2*(ad-b^2c)/b-2d*(-a^2b)^{-1/2}/b*\left(x+1/b*(-a^2b)^{-1/2}\right)+2*\left(-\left(ad-b^2c\right)/b\right)^{1/2}*\left(\left(x+1/b*(-a^2b)^{-1/2}\right)^2d-2d*(-a^2b)^{-1/2}/b*\left(x+1/b*(-a^2b)^{-1/2}\right)-(ad-b^2c)/b\right)^{1/2}}{\left(x+1/b*(-a^2b)^{-1/2}\right)}\right)-1/2/(-a^2b)^{-1/2}/(ad-b^2c)*b/\left(\left(x-1/b*(-a^2b)^{-1/2}\right)^2d+2d*(-a^2b)^{-1/2}/b*\left(x-1/b*(-a^2b)^{-1/2}\right)-(ad-b^2c)/b\right)^{1/2}+1/2/(ad-b^2c)/c/\left(\left(x-1/b*(-a^2b)^{-1/2}\right)^2d+2d*(-a^2b)^{-1/2}/b*\left(x-1/b*(-a^2b)^{-1/2}\right)-(ad-b^2c)/b\right)^{1/2}*x*d+1/2/(-a^2b)^{-1/2}/(ad-b^2c)*b/\left(-\left(ad-b^2c\right)/b\right)^{1/2}*\ln\left(\frac{-2*(ad-b^2c)/b+2d*(-a^2b)^{-1/2}/b*\left(x-1/b*(-a^2b)^{-1/2}\right)+2*\left(-\left(ad-b^2c\right)/b\right)^{1/2}*\left(\left(x-1/b*(-a^2b)^{-1/2}\right)^2d+2d*(-a^2b)^{-1/2}/b*\left(x-1/b*(-a^2b)^{-1/2}\right)-(ad-b^2c)/b\right)^{1/2}}{\left(x-1/b*(-a^2b)^{-1/2}\right)}\right)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)*(d*x^2 + c)^(3/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + a)(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)*(c + d*x^2)^(3/2)),x)

[Out] int(1/((a + b*x^2)*(c + d*x^2)^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)(c + dx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x**2+a)/(d*x**2+c)**(3/2),x)
```

```
[Out] Integral(1/((a + b*x**2)*(c + d*x**2)**(3/2)), x)
```

$$3.102 \quad \int \frac{1}{(a+bx^2)^2 \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=100

$$\frac{(bc - 2ad) \tan^{-1} \left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}} \right)}{2a^{3/2}(bc - ad)^{3/2}} + \frac{bx\sqrt{c+dx^2}}{2a(a+bx^2)(bc - ad)}$$

[Out] 1/2*(-2*a*d+b*c)*arctan(x*(-a*d+b*c)^(1/2)/a^(1/2)/(d*x^2+c)^(1/2))/a^(3/2)/(-a*d+b*c)^(3/2)+1/2*b*x*(d*x^2+c)^(1/2)/a/(-a*d+b*c)/(b*x^2+a)

Rubi [A] time = 0.05, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {382, 377, 205}

$$\frac{(bc - 2ad) \tan^{-1} \left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}} \right)}{2a^{3/2}(bc - ad)^{3/2}} + \frac{bx\sqrt{c+dx^2}}{2a(a+bx^2)(bc - ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^2*Sqrt[c + d*x^2]),x]

[Out] (b*x*Sqrt[c + d*x^2])/(2*a*(b*c - a*d)*(a + b*x^2)) + ((b*c - 2*a*d)*ArcTan[(Sqrt[b*c - a*d]*x)/(Sqrt[a]*Sqrt[c + d*x^2])])/(2*a^(3/2)*(b*c - a*d)^(3/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)], Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+2)+1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)^2 \sqrt{c+dx^2}} dx &= \frac{bx\sqrt{c+dx^2}}{2a(bc-ad)(a+bx^2)} + \frac{(bc-2ad) \int \frac{1}{(a+bx^2)\sqrt{c+dx^2}} dx}{2a(bc-ad)} \\ &= \frac{bx\sqrt{c+dx^2}}{2a(bc-ad)(a+bx^2)} + \frac{(bc-2ad) \text{Subst}\left(\int \frac{1}{a-(-bc+ad)x^2} dx, x, \frac{x}{\sqrt{c+dx^2}}\right)}{2a(bc-ad)} \\ &= \frac{bx\sqrt{c+dx^2}}{2a(bc-ad)(a+bx^2)} + \frac{(bc-2ad) \tan^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{a}\sqrt{c+dx^2}}\right)}{2a^{3/2}(bc-ad)^{3/2}} \end{aligned}$$

Mathematica [C] time = 0.78, size = 405, normalized size = 4.05

$$\frac{x\sqrt{c+dx^2} \left(-30dx^2 \sqrt{\frac{ax^2(c+dx^2)(bc-ad)}{c^2(a+bx^2)^2}} - 45c \sqrt{\frac{ax^2(c+dx^2)(bc-ad)}{c^2(a+bx^2)^2}} + 16dx^2 \sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}} \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)^{5/2} {}_2F_1\left(2, 3; \frac{7}{2}; \frac{(bc-ad)x}{c(bx^2+a)}\right) \right)}{30c^2(a+bx^2)^2 \left(\frac{x^2(bc-ad)}{c(a+bx^2)} \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^2*Sqrt[c + d*x^2]), x]

[Out] (x*Sqrt[c + d*x^2]*(-45*c*Sqrt[(a*(b*c - a*d)*x^2*(c + d*x^2))/(c^2*(a + b*x^2)^2)] - 30*d*x^2*Sqrt[(a*(b*c - a*d)*x^2*(c + d*x^2))/(c^2*(a + b*x^2)^2)]) + 45*c*ArcSin[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]] + 30*d*x^2*ArcSin[Sqrt[((b*c - a*d)*x^2)/(c*(a + b*x^2))]] + 16*c*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2)*Sqrt[(a*(c + d*x^2))/(c*(a + b*x^2))]*Hypergeometric2F1[2, 3, 7/2, ((b*c - a*d)*x^2)/(c*(a + b*x^2))] + 16*d*x^2*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(5/2)*Sqrt[(a*(c + d*x^2))/(c*(a + b*x^2))]*Hypergeometric2F1[2, 3, 7/2, ((b*c - a*d)*x^2)/(c*(a + b*x^2))]/(30*c^2*(((b*c - a*d)*x^2)/(c*(a + b*x^2)))^(3/2)*(a + b*x^2)^2*Sqrt[(a*(c + d*x^2))/(c*(a + b*x^2))])

fricas [B] time = 0.98, size = 459, normalized size = 4.59

$$\frac{4(ab^2c - a^2bd)\sqrt{dx^2 + c}x - (abc - 2a^2d + (b^2c - 2abd)x^2)\sqrt{-abc + a^2d} \log\left(\frac{(b^2c^2 - 8abcd + 8a^2d^2)x^4 + a^2c^2 - 2(3abc^2 - 4a^2bd)x^2 + a^2c^2}{b^2x^4}\right)}{8(a^3b^2c^2 - 2a^4bcd + a^5d^2 + (a^2b^3c^2 - 2a^3b^2cd + a^4bd^2)x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] [1/8*(4*(a*b^2*c - a^2*b*d)*sqrt(d*x^2 + c)*x - (a*b*c - 2*a^2*d + (b^2*c - 2*a*b*d)*x^2)*sqrt(-a*b*c + a^2*d)*log(((b^2*c^2 - 8*a*b*c*d + 8*a^2*d^2)*x^4 + a^2*c^2 - 2*(3*a*b*c^2 - 4*a^2*c*d)*x^2 - 4*((b*c - 2*a*d)*x^3 - a*c*x)*sqrt(-a*b*c + a^2*d)*sqrt(d*x^2 + c))/(b^2*x^4 + 2*a*b*x^2 + a^2)))/(a^3*b^2*c^2 - 2*a^4*b*c*d + a^5*d^2 + (a^2*b^3*c^2 - 2*a^3*b^2*c*d + a^4*b*d^2)*x^2), 1/4*(2*(a*b^2*c - a^2*b*d)*sqrt(d*x^2 + c)*x + sqrt(a*b*c - a^2*d)*(a*b*c - 2*a^2*d + (b^2*c - 2*a*b*d)*x^2)*arctan(1/2*sqrt(a*b*c - a^2*d)*((b*c - 2*a*d)*x^2 - a*c)*sqrt(d*x^2 + c)/((a*b*c*d - a^2*d^2)*x^3 + (a*b*c^2 - a^2*c*d)*x)))/(a^3*b^2*c^2 - 2*a^4*b*c*d + a^5*d^2 + (a^2*b^3*c^2 - 2*a^3*b^2*c*d + a^4*b*d^2)*x^2)]

giac [B] time = 0.62, size = 225, normalized size = 2.25

$$-\frac{1}{2}d^{\frac{3}{2}} \left(\frac{(bc - 2ad) \arctan \left(\frac{(\sqrt{d}x - \sqrt{dx^2 + c})^2 b - bc + 2ad}{2\sqrt{abcd - a^2d^2}} \right)}{(abcd - a^2d^2)^{\frac{3}{2}}} \right) + \frac{2 \left((\sqrt{d}x - \sqrt{dx^2 + c})^2 bc - 2(\sqrt{d}x - \sqrt{dx^2 + c})^2 b - 2(\sqrt{d}x - \sqrt{dx^2 + c})^2 bc + 4(\sqrt{d}x - \sqrt{dx^2 + c})^2 b^2 \right)}{\left((\sqrt{d}x - \sqrt{dx^2 + c})^4 b - 2(\sqrt{d}x - \sqrt{dx^2 + c})^2 bc + 4(\sqrt{d}x - \sqrt{dx^2 + c})^2 b^2 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] $-\frac{1}{2}d^{\frac{3}{2}} \left(\frac{(bc - 2ad) \arctan \left(\frac{(\sqrt{d}x - \sqrt{dx^2 + c})^2 b - bc + 2ad}{2\sqrt{abcd - a^2d^2}} \right)}{(abcd - a^2d^2)^{\frac{3}{2}}} \right) + \frac{2 \left((\sqrt{d}x - \sqrt{dx^2 + c})^2 bc - 2(\sqrt{d}x - \sqrt{dx^2 + c})^2 b - 2(\sqrt{d}x - \sqrt{dx^2 + c})^2 bc + 4(\sqrt{d}x - \sqrt{dx^2 + c})^2 b^2 \right)}{\left((\sqrt{d}x - \sqrt{dx^2 + c})^4 b - 2(\sqrt{d}x - \sqrt{dx^2 + c})^2 bc + 4(\sqrt{d}x - \sqrt{dx^2 + c})^2 b^2 \right)}$

maple [B] time = 0.02, size = 823, normalized size = 8.23

$$\frac{\sqrt{-ab} d \ln \left(\frac{2\sqrt{-ab} \left(x - \frac{\sqrt{-ab}}{b} \right) d - \frac{2(ad-bc)}{b} + 2\sqrt{-\frac{ad-bc}{b}} \sqrt{\left(x - \frac{\sqrt{-ab}}{b} \right)^2 d + \frac{2\sqrt{-ab} \left(x - \frac{\sqrt{-ab}}{b} \right) d - \frac{ad-bc}{b}}}{x - \frac{\sqrt{-ab}}{b}} \right)}{4(ad-bc) \sqrt{-\frac{ad-bc}{b}} ab} - \frac{\sqrt{-ab} d \ln \left(\frac{2\sqrt{-ab} \left(x + \frac{\sqrt{-ab}}{b} \right) d - \frac{2(ad-bc)}{b}}{x + \frac{\sqrt{-ab}}{b}} \right)}{4(ad-bc) \sqrt{-\frac{ad-bc}{b}} ab}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^2/(d*x^2+c)^(1/2),x)

[Out] $\frac{1}{4} \frac{a}{(-a*b)^{\frac{1}{2}}} \frac{1}{(-a*d-b*c/b)^{\frac{1}{2}}} \ln \left(\frac{-2*(-a*b)^{\frac{1}{2}} * (x + (-a*b)^{\frac{1}{2}}/b) / b * d - 2*(a*d-b*c)/b + 2*(-(a*d-b*c)/b)^{\frac{1}{2}} * ((x + (-a*b)^{\frac{1}{2}}/b)^2 * d - 2*(-a*b)^{\frac{1}{2}} * (x + (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}}{(x + (-a*b)^{\frac{1}{2}}/b) * ((x + (-a*b)^{\frac{1}{2}}/b)^2 * d - 2*(-a*b)^{\frac{1}{2}} * (x + (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}} \right) - \frac{1}{4} \frac{a}{(a*d-b*c)} \frac{1}{(x + (-a*b)^{\frac{1}{2}}/b)^{\frac{1}{2}}} \frac{1}{((x + (-a*b)^{\frac{1}{2}}/b)^2 * d - 2*(-a*b)^{\frac{1}{2}} * (x + (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}} - \frac{1}{4} \frac{a}{(-a*b)^{\frac{1}{2}}} \frac{1}{(-a*d-b*c/b)^{\frac{1}{2}}} \ln \left(\frac{-2*(-a*b)^{\frac{1}{2}} * (x + (-a*b)^{\frac{1}{2}}/b) / b * d - 2*(a*d-b*c)/b + 2*(-(a*d-b*c)/b)^{\frac{1}{2}} * ((x + (-a*b)^{\frac{1}{2}}/b)^2 * d - 2*(-a*b)^{\frac{1}{2}} * (x + (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}}{(x + (-a*b)^{\frac{1}{2}}/b) * ((x + (-a*b)^{\frac{1}{2}}/b)^2 * d - 2*(-a*b)^{\frac{1}{2}} * (x + (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}} \right) - \frac{1}{4} \frac{a}{(-a*b)^{\frac{1}{2}}} \frac{1}{(-a*d-b*c/b)^{\frac{1}{2}}} \ln \left(\frac{2*(-a*b)^{\frac{1}{2}} * (x - (-a*b)^{\frac{1}{2}}/b) / b * d - 2*(a*d-b*c)/b + 2*(-(a*d-b*c)/b)^{\frac{1}{2}} * ((x - (-a*b)^{\frac{1}{2}}/b)^2 * d + 2*(-a*b)^{\frac{1}{2}} * (x - (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}}{(x - (-a*b)^{\frac{1}{2}}/b) * ((x - (-a*b)^{\frac{1}{2}}/b)^2 * d + 2*(-a*b)^{\frac{1}{2}} * (x - (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}} \right) - \frac{1}{4} \frac{a}{(a*d-b*c)} \frac{1}{(x - (-a*b)^{\frac{1}{2}}/b)^{\frac{1}{2}}} \frac{1}{((x - (-a*b)^{\frac{1}{2}}/b)^2 * d + 2*(-a*b)^{\frac{1}{2}} * (x - (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}} + \frac{1}{4} \frac{a}{b} \frac{1}{a*d*(-a*b)^{\frac{1}{2}}} \frac{1}{(a*d-b*c)} \frac{1}{(-a*d-b*c/b)^{\frac{1}{2}}} \ln \left(\frac{2*(-a*b)^{\frac{1}{2}} * (x - (-a*b)^{\frac{1}{2}}/b) / b * d - 2*(a*d-b*c)/b + 2*(-(a*d-b*c)/b)^{\frac{1}{2}} * ((x - (-a*b)^{\frac{1}{2}}/b)^2 * d + 2*(-a*b)^{\frac{1}{2}} * (x - (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}}{(x - (-a*b)^{\frac{1}{2}}/b) * ((x - (-a*b)^{\frac{1}{2}}/b)^2 * d + 2*(-a*b)^{\frac{1}{2}} * (x - (-a*b)^{\frac{1}{2}}/b) / b * d - (a*d-b*c)/b)^{\frac{1}{2}}} \right)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^2 \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/sqrt(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^2*sqrt(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + a)^2 \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^2*(c + d*x^2)^(1/2)),x)

[Out] int(1/((a + b*x^2)^2*(c + d*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^2 \sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**2/(d*x**2+c)**(1/2),x)

[Out] Integral(1/((a + b*x**2)**2*sqrt(c + d*x**2)), x)

$$3.103 \quad \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^3} dx$$

Optimal. Leaf size=149

$$\frac{c(3bc - 4ad) \tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{8a^{5/2}(bc - ad)^{3/2}} + \frac{x\sqrt{c+dx^2}(3bc - 4ad)}{8a^2(a+bx^2)(bc - ad)} + \frac{bx(c+dx^2)^{3/2}}{4a(a+bx^2)^2(bc - ad)}$$

[Out] $1/4*b*x*(d*x^2+c)^(3/2)/a/(-a*d+b*c)/(b*x^2+a)^2+1/8*c*(-4*a*d+3*b*c)*\arctan(x*(-a*d+b*c)^(1/2)/a^(1/2)/(d*x^2+c)^(1/2))/a^(5/2)/(-a*d+b*c)^(3/2)+1/8*(-4*a*d+3*b*c)*x*(d*x^2+c)^(1/2)/a^2/(-a*d+b*c)/(b*x^2+a)$

Rubi [A] time = 0.08, antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {382, 378, 377, 205}

$$\frac{x\sqrt{c+dx^2}(3bc - 4ad)}{8a^2(a+bx^2)(bc - ad)} + \frac{c(3bc - 4ad) \tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{8a^{5/2}(bc - ad)^{3/2}} + \frac{bx(c+dx^2)^{3/2}}{4a(a+bx^2)^2(bc - ad)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/(a + b*x^2)^3, x]

[Out] $((3*b*c - 4*a*d)*x*\text{Sqrt}[c + d*x^2])/(8*a^2*(b*c - a*d)*(a + b*x^2)) + (b*x*(c + d*x^2)^(3/2))/(4*a*(b*c - a*d)*(a + b*x^2)^2) + (c*(3*b*c - 4*a*d)*\text{ArcTan}[(\text{Sqrt}[b*c - a*d]*x)/(\text{Sqrt}[a]*\text{Sqrt}[c + d*x^2])])/(8*a^(5/2)*(b*c - a*d)^(3/2))$

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] - Dist[(c*q)/(a*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+1)+1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+2)+1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{c+dx^2}}{(a+bx^2)^3} dx &= \frac{bx(c+dx^2)^{3/2}}{4a(bc-ad)(a+bx^2)^2} + \frac{(3bc-4ad) \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^2} dx}{4a(bc-ad)} \\
&= \frac{(3bc-4ad)x\sqrt{c+dx^2}}{8a^2(bc-ad)(a+bx^2)} + \frac{bx(c+dx^2)^{3/2}}{4a(bc-ad)(a+bx^2)^2} + \frac{(c(3bc-4ad)) \int \frac{1}{(a+bx^2)\sqrt{c+dx^2}} dx}{8a^2(bc-ad)} \\
&= \frac{(3bc-4ad)x\sqrt{c+dx^2}}{8a^2(bc-ad)(a+bx^2)} + \frac{bx(c+dx^2)^{3/2}}{4a(bc-ad)(a+bx^2)^2} + \frac{(c(3bc-4ad)) \text{Subst}\left(\int \frac{1}{a-(-bc+ad)x^2} dx, x, \right)}{8a^2(bc-ad)} \\
&= \frac{(3bc-4ad)x\sqrt{c+dx^2}}{8a^2(bc-ad)(a+bx^2)} + \frac{bx(c+dx^2)^{3/2}}{4a(bc-ad)(a+bx^2)^2} + \frac{c(3bc-4ad) \tan^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{a}\sqrt{c+dx^2}}\right)}{8a^{5/2}(bc-ad)^{3/2}}
\end{aligned}$$

Mathematica [A] time = 5.17, size = 130, normalized size = 0.87

$$\frac{\sqrt{a}x\sqrt{c+dx^2}(-4a^2d+ab(5c-2dx^2)+3b^2cx^2)}{(a+bx^2)^2(bc-ad)} + \frac{c(3bc-4ad) \tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{(bc-ad)^{3/2}}}{8a^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + d*x^2]/(a + b*x^2)^3, x]

[Out] ((Sqrt[a]*x*Sqrt[c + d*x^2]*(-4*a^2*d + 3*b^2*c*x^2 + a*b*(5*c - 2*d*x^2)))/(b*c - a*d)*(a + b*x^2)^2) + (c*(3*b*c - 4*a*d)*ArcTan[(Sqrt[b*c - a*d]*x)/(Sqrt[a]*Sqrt[c + d*x^2])])/(b*c - a*d)^(3/2))/(8*a^(5/2))

fricas [B] time = 1.01, size = 698, normalized size = 4.68

$$\left[\frac{(3a^2bc^2 - 4a^3cd + (3b^3c^2 - 4ab^2cd)x^4 + 2(3ab^2c^2 - 4a^2bcd)x^2)\sqrt{-abc + a^2d} \log\left(\frac{(b^2c^2 - 8abcd + 8a^2d^2)x^4 + a^2c^2 - 2}{32(a^5b^2c^2 - 2a^6bcd + a^7d^2 + (a^3b^4c^2 - \dots)}\right)}{32(a^5b^2c^2 - 2a^6bcd + a^7d^2 + (a^3b^4c^2 - \dots)} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^3, x, algorithm="fricas")

[Out] [-1/32*((3*a^2*b*c^2 - 4*a^3*c*d + (3*b^3*c^2 - 4*a*b^2*c*d)*x^4 + 2*(3*a*b^2*c^2 - 4*a^2*b*c*d)*x^2)*sqrt(-a*b*c + a^2*d)*log(((b^2*c^2 - 8*a*b*c*d + 8*a^2*d^2)*x^4 + a^2*c^2 - 2*(3*a*b*c^2 - 4*a^2*c*d)*x^2 - 4*((b*c - 2*a*d)*x^3 - a*c*x)*sqrt(-a*b*c + a^2*d)*sqrt(d*x^2 + c))/(b^2*x^4 + 2*a*b*x^2 + a^2)) - 4*((3*a*b^3*c^2 - 5*a^2*b^2*c*d + 2*a^3*b*d^2)*x^3 + (5*a^2*b^2*c^2 - 9*a^3*b*c*d + 4*a^4*d^2)*x)*sqrt(d*x^2 + c))/(a^5*b^2*c^2 - 2*a^6*b*c*d + a^7*d^2 + (a^3*b^4*c^2 - 2*a^4*b^3*c*d + a^5*b^2*d^2)*x^4 + 2*(a^4*b^3*c^2 - 2*a^5*b^2*c*d + a^6*b*d^2)*x^2), 1/16*((3*a^2*b*c^2 - 4*a^3*c*d + (3*b^3*c^2 - 4*a*b^2*c*d)*x^4 + 2*(3*a*b^2*c^2 - 4*a^2*b*c*d)*x^2)*sqrt(a*b*c - a^2*d)*arctan(1/2*sqrt(a*b*c - a^2*d)*((b*c - 2*a*d)*x^2 - a*c)*sqrt(d*x^2 + c))/((a*b*c*d - a^2*d^2)*x^3 + (a*b*c^2 - a^2*c*d)*x) + 2*((3*a*b^3*c^2 - 5*a^2*b^2*c*d + 2*a^3*b*d^2)*x^3 + (5*a^2*b^2*c^2 - 9*a^3*b*c*d + 4*a^4*d^2)*x)*sqrt(d*x^2 + c))/(a^5*b^2*c^2 - 2*a^6*b*c*d + a^7*d^2 + (a^3*b^4*c^2 - \dots)

$- 2*a^4*b^3*c*d + a^5*b^2*d^2)*x^4 + 2*(a^4*b^3*c^2 - 2*a^5*b^2*c*d + a^6*b*d^2)*x^2]$

giac [B] time = 3.13, size = 487, normalized size = 3.27

$$\frac{\left(3bc^2\sqrt{d} - 4acd^{\frac{3}{2}}\right) \arctan\left(\frac{\left(\sqrt{d}x - \sqrt{dx^2+c}\right)^2 b - bc + 2ad}{2\sqrt{abcd - a^2d^2}}\right) + 3\left(\sqrt{d}x - \sqrt{dx^2+c}\right)^6 b^3c^2\sqrt{d} - 4\left(\sqrt{d}x - \sqrt{dx^2+c}\right)^6 a}{8\left(a^2bc - a^3d\right)\sqrt{abcd - a^2d^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^3,x, algorithm="giac")

[Out] $-1/8*(3*b*c^2*\sqrt{d} - 4*a*c*d^{(3/2)})*\arctan(1/2*((\sqrt{d}*x - \sqrt{d*x^2 + c})^2*b - b*c + 2*a*d)/\sqrt{a*b*c*d - a^2*d^2})/((a^2*b*c - a^3*d)*\sqrt{a*b*c*d - a^2*d^2}) - 1/4*(3*(\sqrt{d}*x - \sqrt{d*x^2 + c})^6*b^3*c^2*\sqrt{d} - 4*(\sqrt{d}*x - \sqrt{d*x^2 + c})^6*a*b^2*c*d^{(3/2)} - 9*(\sqrt{d}*x - \sqrt{d*x^2 + c})^4*b^3*c^3*\sqrt{d} + 30*(\sqrt{d}*x - \sqrt{d*x^2 + c})^4*a*b^2*c^2*d^{(3/2)} - 40*(\sqrt{d}*x - \sqrt{d*x^2 + c})^4*a^2*b*c*d^{(5/2)} + 16*(\sqrt{d}*x - \sqrt{d*x^2 + c})^4*a^3*d^{(7/2)} + 9*(\sqrt{d}*x - \sqrt{d*x^2 + c})^2*b^3*c^4*\sqrt{d} - 28*(\sqrt{d}*x - \sqrt{d*x^2 + c})^2*a*b^2*c^3*d^{(3/2)} + 16*(\sqrt{d}*x - \sqrt{d*x^2 + c})^2*a^2*b*c^2*d^{(5/2)} - 3*b^3*c^5*\sqrt{d} + 2*a*b^2*c^4*d^{(3/2)})/(((\sqrt{d}*x - \sqrt{d*x^2 + c})^4*b - 2*(\sqrt{d}*x - \sqrt{d*x^2 + c})^2*b*c + 4*(\sqrt{d}*x - \sqrt{d*x^2 + c})^2*a*d + b*c^2)^2*(a^2*b^2*c - a^3*b*d))$

maple [B] time = 0.03, size = 5177, normalized size = 34.74

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^(1/2)/(b*x^2+a)^3,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{(bx^2+a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^3,x, algorithm="maxima")

[Out] integrate(sqrt(d*x^2+c)/(b*x^2+a)^3,x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{d x^2+c}}{(b x^2+a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c+d*x^2)^(1/2)/(a+b*x^2)^3,x)

[Out] int((c+d*x^2)^(1/2)/(a+b*x^2)^3,x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2}}{(a + bx^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**(1/2)/(b*x**2+a)**3,x)

[Out] Integral(sqrt(c + d*x**2)/(a + b*x**2)**3, x)

$$3.104 \quad \int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^4} dx$$

Optimal. Leaf size=199

$$\frac{c^2(5bc - 6ad) \tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{16a^{7/2}(bc - ad)^{3/2}} + \frac{cx\sqrt{c+dx^2}(5bc - 6ad)}{16a^3(a+bx^2)(bc - ad)} + \frac{x(c+dx^2)^{3/2}(5bc - 6ad)}{24a^2(a+bx^2)^2(bc - ad)} + \frac{bx(c+dx^2)^{5/2}}{6a(a+bx^2)^3(bc - ad)}$$

[Out] 1/24*(-6*a*d+5*b*c)*x*(d*x^2+c)^(3/2)/a^2/(-a*d+b*c)/(b*x^2+a)^2+1/6*b*x*(d*x^2+c)^(5/2)/a/(-a*d+b*c)/(b*x^2+a)^3+1/16*c^2*(-6*a*d+5*b*c)*arctan(x*(-a*d+b*c)^(1/2)/a^(1/2)/(d*x^2+c)^(1/2))/a^(7/2)/(-a*d+b*c)^(3/2)+1/16*c*(-6*a*d+5*b*c)*x*(d*x^2+c)^(1/2)/a^3/(-a*d+b*c)/(b*x^2+a)

Rubi [A] time = 0.11, antiderivative size = 199, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {382, 378, 377, 205}

$$\frac{c^2(5bc - 6ad) \tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{16a^{7/2}(bc - ad)^{3/2}} + \frac{x(c+dx^2)^{3/2}(5bc - 6ad)}{24a^2(a+bx^2)^2(bc - ad)} + \frac{cx\sqrt{c+dx^2}(5bc - 6ad)}{16a^3(a+bx^2)(bc - ad)} + \frac{bx(c+dx^2)^{5/2}}{6a(a+bx^2)^3(bc - ad)}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^(3/2)/(a + b*x^2)^4, x]

[Out] (c*(5*b*c - 6*a*d)*x*sqrt[c + d*x^2])/(16*a^3*(b*c - a*d)*(a + b*x^2)) + ((5*b*c - 6*a*d)*x*(c + d*x^2)^(3/2))/(24*a^2*(b*c - a*d)*(a + b*x^2)^2) + (b*x*(c + d*x^2)^(5/2))/(6*a*(b*c - a*d)*(a + b*x^2)^3) + (c^2*(5*b*c - 6*a*d)*ArcTan[(sqrt[b*c - a*d]*x)/(sqrt[a]*sqrt[c + d*x^2])])/(16*a^(7/2)*(b*c - a*d)^(3/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 378

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] - Dist[(c*q)/(a*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+1)+1, 0] && GtQ[q, 0] && NeQ[p, -1]

Rule 382

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[(b*c + n*(p+1)*(b*c - a*d))/(a*n*(p+1)*(b*c - a*d)], Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p+q+2)+1, 0] && (LtQ[p, -1] || !LtQ

[q, -1]) && NeQ[p, -1]

Rubi steps

$$\int \frac{(c + dx^2)^{3/2}}{(a + bx^2)^4} dx = \frac{bx(c + dx^2)^{5/2}}{6a(bc - ad)(a + bx^2)^3} + \frac{(5bc - 6ad) \int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^3} dx}{6a(bc - ad)}$$

$$= \frac{(5bc - 6ad)x(c + dx^2)^{3/2}}{24a^2(bc - ad)(a + bx^2)^2} + \frac{bx(c + dx^2)^{5/2}}{6a(bc - ad)(a + bx^2)^3} + \frac{(c(5bc - 6ad)) \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^2} dx}{8a^2(bc - ad)}$$

$$= \frac{c(5bc - 6ad)x\sqrt{c + dx^2}}{16a^3(bc - ad)(a + bx^2)} + \frac{(5bc - 6ad)x(c + dx^2)^{3/2}}{24a^2(bc - ad)(a + bx^2)^2} + \frac{bx(c + dx^2)^{5/2}}{6a(bc - ad)(a + bx^2)^3} + \frac{(c^2(5bc - 6ad)) \int \frac{1}{(a+bx^2)} dx}{16a^3(bc - ad)}$$

$$= \frac{c(5bc - 6ad)x\sqrt{c + dx^2}}{16a^3(bc - ad)(a + bx^2)} + \frac{(5bc - 6ad)x(c + dx^2)^{3/2}}{24a^2(bc - ad)(a + bx^2)^2} + \frac{bx(c + dx^2)^{5/2}}{6a(bc - ad)(a + bx^2)^3} + \frac{(c^2(5bc - 6ad)) \ln|ax + \sqrt{c+dx^2}|}{16a^3(bc - ad)}$$

Mathematica [A] time = 5.27, size = 179, normalized size = 0.90

$$\frac{3c^2(5bc-6ad) \tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right) - \sqrt{a}x\sqrt{c+dx^2}(-6a^3d(5c+2dx^2)+a^2b(33c^2-22cdx^2-4d^2x^4)+8ab^2cx^2(5c-dx^2)+15b^3c^2x^4)}{(bc-ad)^{3/2} (a+bx^2)^3(ad-bc)} \cdot \frac{1}{48a^{7/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(c + d*x^2)^(3/2)/(a + b*x^2)^4,x]
[Out] (-((Sqrt[a]*x*Sqrt[c + d*x^2]*(15*b^3*c^2*x^4 + 8*a*b^2*c*x^2*(5*c - d*x^2) - 6*a^3*d*(5*c + 2*d*x^2) + a^2*b*(33*c^2 - 22*c*d*x^2 - 4*d^2*x^4)))/((- (b*c) + a*d)*(a + b*x^2)^3)) + (3*c^2*(5*b*c - 6*a*d)*ArcTan[(Sqrt[b*c - a*d]*x)/(Sqrt[a]*Sqrt[c + d*x^2])])/(b*c - a*d)^(3/2))/(48*a^(7/2))
```

fricas [B] time = 1.28, size = 972, normalized size = 4.88

$$\left[\frac{3(5a^3bc^3 - 6a^4c^2d + (5b^4c^3 - 6ab^3c^2d)x^6 + 3(5ab^3c^3 - 6a^2b^2c^2d)x^4 + 3(5a^2b^2c^3 - 6a^3bc^2d)x^2)\sqrt{-abc + a^2}}{192} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^4,x, algorithm="fricas")
[Out] [-1/192*(3*(5*a^3*b*c^3 - 6*a^4*c^2*d + (5*b^4*c^3 - 6*a*b^3*c^2*d)*x^6 + 3*(5*a^2*b^2*c^3 - 6*a^3*b*c^2*d)*x^2)*sqrt(-a*b*c + a^2*d)*log(((b^2*c^2 - 8*a*b*c*d + 8*a^2*d^2)*x^4 + a^2*c^2 - 2*(3*a*b*c^2 - 4*a^2*c*d)*x^2 - 4*((b*c - 2*a*d)*x^3 - a*c*x)*sqrt(-a*b*c + a^2*d)*sqrt(d*x^2 + c))/(b^2*x^4 + 2*a*b*x^2 + a^2)) - 4*((15*a*b^4*c^3 - 23*a^2*b^3*c^2*d + 4*a^3*b^2*c*d^2 + 4*a^4*b*d^3)*x^5 + 2*(20*a^2*b^3*c
```

$$\begin{aligned} &^3 - 31a^3b^2c^2d + 5a^4b^2c^2d + 6a^5d^3)x^3 + 3(11a^3b^2c^3 \\ &- 21a^4b^2c^2d + 10a^5c^2d^2)x) \sqrt{dx^2 + c}) / (a^7b^2c^2 - 2a^8b \\ &^2c^2d + a^9d^2 + (a^4b^5c^2 - 2a^5b^4c^2d + a^6b^3d^2)x^6 + 3(a^5b \\ &^4c^2 - 2a^6b^3c^2d + a^7b^2d^2)x^4 + 3(a^6b^3c^2 - 2a^7b^2c^2d \\ &+ a^8b^2d^2)x^2), 1/96(3(5a^3b^2c^3 - 6a^4c^2d + (5b^4c^3 - 6a^2b^3 \\ &^3c^2d)x^6 + 3(5a^2b^3c^3 - 6a^2b^2c^2d)x^4 + 3(5a^2b^2c^3 - 6 \\ &a^3b^2c^2d)x^2) \sqrt{abc - a^2d}) \arctan(1/2 \sqrt{abc - a^2d}) ((b^2c \\ &- 2a^2d)x^2 - ac) \sqrt{dx^2 + c}) / ((a^2b^2c^2d - a^2d^2)x^3 + (a^2b^2c^2 - \\ &a^2c^2d)x) + 2((15a^2b^4c^3 - 23a^2b^3c^2d + 4a^3b^2c^2d^2 + 4a^4 \\ &^4b^2d^3)x^5 + 2(20a^2b^3c^3 - 31a^3b^2c^2d + 5a^4b^2c^2d + 6a^5 \\ &^5d^3)x^3 + 3(11a^3b^2c^3 - 21a^4b^2c^2d + 10a^5c^2d^2)x) \sqrt{dx^2 \\ &+ c}) / (a^7b^2c^2 - 2a^8b^2c^2d + a^9d^2 + (a^4b^5c^2 - 2a^5b^4c^2d \\ &+ a^6b^3d^2)x^6 + 3(a^5b^4c^2 - 2a^6b^3c^2d + a^7b^2d^2)x^4 + 3 \\ &^3(a^6b^3c^2 - 2a^7b^2c^2d + a^8b^2d^2)x^2)] \end{aligned}$$

giac [B] time = 2.81, size = 919, normalized size = 4.62

$$\frac{\left(5bc^3\sqrt{d} - 6ac^2d^{\frac{3}{2}}\right) \arctan\left(\frac{\left(\sqrt{d}x - \sqrt{dx^2+c}\right)^2 b - bc + 2ad}{2\sqrt{abcd - a^2d^2}}\right) - 15\left(\sqrt{d}x - \sqrt{dx^2+c}\right)^{10} b^5c^3\sqrt{d} - 18\left(\sqrt{d}x - \sqrt{dx^2+c}\right)^8 b^4c^3\sqrt{d}}{16(a^3bc - a^4d)\sqrt{abcd - a^2d^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((dx^2+c)^(3/2)/(bx^2+a)^4,x, algorithm="giac")

[Out]
$$\begin{aligned} &-1/16(5b^2c^3\sqrt{d} - 6a^2c^2d^{3/2}) \arctan(1/2((\sqrt{d})x - \sqrt{dx^2 \\ &^2 + c}))^2 b - bc + 2a^2d) / \sqrt{abc^2d - a^2d^2}) / ((a^3b^2c^2 - a^4d) \sqrt{ \\ &^2 abc^2d - a^2d^2}) - 1/24(15(\sqrt{d})x - \sqrt{dx^2 + c})^{10} b^5c^3 \sqrt{d} \\ &^3 \sqrt{d} - 18(\sqrt{d})x - \sqrt{dx^2 + c})^{10} a^2b^4c^2d^{3/2} - 75(\sqrt{d} \\ &^2)x - \sqrt{dx^2 + c})^8 b^5c^4 \sqrt{d} + 240(\sqrt{d})x - \sqrt{dx^2 + c} \\ &^8 a^2b^4c^3d^{3/2} - 180(\sqrt{d})x - \sqrt{dx^2 + c})^8 a^2b^3c^2d^{5/2} - 96(\sqrt{d})x \\ &^8 a^3b^2c^2d^{7/2} + 96(\sqrt{d})x \\ &- \sqrt{dx^2 + c})^8 a^4b^2d^{9/2} + 150(\sqrt{d})x - \sqrt{dx^2 + c})^6 b \\ &^5c^5 \sqrt{d} - 620(\sqrt{d})x - \sqrt{dx^2 + c})^6 a^2b^4c^4d^{3/2} + 96 \\ &^8(\sqrt{d})x - \sqrt{dx^2 + c})^6 a^2b^3c^3d^{5/2} - 720(\sqrt{d})x - \sqrt{ \\ &^2 dx^2 + c})^6 a^3b^2c^2d^{7/2} + 64(\sqrt{d})x - \sqrt{dx^2 + c})^6 a \\ &^4b^2c^2d^{9/2} + 128(\sqrt{d})x - \sqrt{dx^2 + c})^6 a^5d^{11/2} - 150(\sqrt{ \\ &^2 d})x - \sqrt{dx^2 + c})^4 b^5c^6 \sqrt{d} + 600(\sqrt{d})x - \sqrt{dx^2 \\ &^2 + c})^4 a^2b^4c^5d^{3/2} - 864(\sqrt{d})x - \sqrt{dx^2 + c})^4 a^2b^3c^4 \\ &^4d^{5/2} + 288(\sqrt{d})x - \sqrt{dx^2 + c})^4 a^3b^2c^3d^{7/2} + 96(\sqrt{ \\ &^2 d})x - \sqrt{dx^2 + c})^4 a^4b^2c^2d^{9/2} + 75(\sqrt{d})x - \sqrt{dx^2 \\ &^2 + c})^2 b^5c^7 \sqrt{d} - 210(\sqrt{d})x - \sqrt{dx^2 + c})^2 a^2b^4c^6d^{3/2} \\ &^3 + 72(\sqrt{d})x - \sqrt{dx^2 + c})^2 a^2b^3c^5d^{5/2} + 48(\sqrt{d} \\ &^2)x - \sqrt{dx^2 + c})^2 a^3b^2c^4d^{7/2} - 15b^5c^8 \sqrt{d} + 8a^2b^4 \\ &^4c^7d^{3/2} + 4a^2b^3c^6d^{5/2}) / ((a^3b^3c - a^4b^2d)((\sqrt{d})x \\ &- \sqrt{dx^2 + c})^4 b - 2(\sqrt{d})x - \sqrt{dx^2 + c})^2 b^2c + 4(\sqrt{d} \\ &^2)x - \sqrt{dx^2 + c})^2 a^2d + b^2c^2)^3 \end{aligned}$$

maple [B] time = 0.05, size = 13964, normalized size = 70.17

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((dx^2+c)^(3/2)/(bx^2+a)^4,x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^4,x, algorithm="maxima")

[Out] integrate((d*x^2 + c)^(3/2)/(b*x^2 + a)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^(3/2)/(a + b*x^2)^4,x)

[Out] int((c + d*x^2)^(3/2)/(a + b*x^2)^4, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**(3/2)/(b*x**2+a)**4,x)

[Out] Timed out

$$3.105 \quad \int \frac{1}{\left(\frac{bc}{d} + bx^2\right) \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=20

$$\frac{dx}{bc\sqrt{c+dx^2}}$$

[Out] d*x/b/c/(d*x^2+c)^(1/2)

Rubi [A] time = 0.00, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {21, 191}

$$\frac{dx}{bc\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Int[1/(((b*c)/d + b*x^2)*Sqrt[c + d*x^2]),x]

[Out] (d*x)/(b*c*Sqrt[c + d*x^2])

Rule 21

Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rubi steps

$$\int \frac{1}{\left(\frac{bc}{d} + bx^2\right) \sqrt{c+dx^2}} dx = \frac{d \int \frac{1}{(c+dx^2)^{3/2}} dx}{b} = \frac{dx}{bc\sqrt{c+dx^2}}$$

Mathematica [A] time = 0.01, size = 20, normalized size = 1.00

$$\frac{dx}{bc\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(((b*c)/d + b*x^2)*Sqrt[c + d*x^2]),x]

[Out] (d*x)/(b*c*Sqrt[c + d*x^2])

fricas [A] time = 0.56, size = 27, normalized size = 1.35

$$\frac{\sqrt{dx^2 + c} dx}{bcdx^2 + bc^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*c/d+b*x^2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] sqrt(d*x^2 + c)*d*x/(b*c*d*x^2 + b*c^2)

giac [A] time = 0.60, size = 18, normalized size = 0.90

$$\frac{dx}{\sqrt{dx^2 + c} bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*c/d+b*x^2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] d*x/(sqrt(d*x^2 + c)*b*c)

maple [A] time = 0.00, size = 19, normalized size = 0.95

$$\frac{dx}{\sqrt{d x^2 + c} bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*c/d+b*x^2)/(d*x^2+c)^(1/2),x)

[Out] d*x/b/c/(d*x^2+c)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(bx^2 + \frac{bc}{d}\right)\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*c/d+b*x^2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + b*c/d)*sqrt(d*x^2 + c)), x)

mupad [B] time = 4.77, size = 18, normalized size = 0.90

$$\frac{dx}{bc\sqrt{dx^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c + d*x^2)^(1/2)*(b*x^2 + (b*c)/d)),x)

[Out] (d*x)/(b*c*(c + d*x^2)^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{d \int \frac{1}{c\sqrt{c+dx^2}+dx^2\sqrt{c+dx^2}} dx}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*c/d+b*x**2)/(d*x**2+c)**(1/2),x)

[Out] d*Integral(1/(c*sqrt(c + d*x**2) + d*x**2*sqrt(c + d*x**2)), x)/b

$$3.106 \quad \int \frac{1}{\sqrt{1-x^2}(1+x^2)} dx$$

Optimal. Leaf size=25

$$\frac{\tan^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right)}{\sqrt{2}}$$

[Out] 1/2*arctan(x*2^(1/2)/(-x^2+1)^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {377, 203}

$$\frac{\tan^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 - x^2]*(1 + x^2)), x]

[Out] ArcTan[(Sqrt[2]*x)/Sqrt[1 - x^2]]/Sqrt[2]

Rule 203

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{1-x^2}(1+x^2)} dx &= \text{Subst}\left(\int \frac{1}{1+2x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right) \\ &= \frac{\tan^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 25, normalized size = 1.00

$$\frac{\tan^{-1}\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 - x^2]*(1 + x^2)), x]

[Out] ArcTan[(Sqrt[2]*x)/Sqrt[1 - x^2]]/Sqrt[2]

fricas [A] time = 0.55, size = 23, normalized size = 0.92

$$-\frac{1}{2}\sqrt{2}\arctan\left(\frac{\sqrt{2}\sqrt{-x^2+1}}{2x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] -1/2*sqrt(2)*arctan(1/2*sqrt(2)*sqrt(-x^2 + 1)/x)

giac [B] time = 0.60, size = 51, normalized size = 2.04

$$\frac{1}{4}\sqrt{2}\left(\pi\operatorname{sgn}(x)+2\arctan\left(-\frac{\sqrt{2}x\left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2}-1\right)}{4(\sqrt{-x^2+1}-1)}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(2)*(pi*sgn(x) + 2*arctan(-1/4*sqrt(2)*x*((sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1)))

maple [A] time = 0.01, size = 28, normalized size = 1.12

$$\frac{\sqrt{2}\arctan\left(\frac{\sqrt{2}\sqrt{-x^2+1}x}{x^2-1}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2+1)/(-x^2+1)^(1/2),x)

[Out] -1/2*2^(1/2)*arctan(2^(1/2)*(-x^2+1)^(1/2)/(x^2-1)*x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(x^2+1)\sqrt{-x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/((x^2 + 1)*sqrt(-x^2 + 1)), x)

mupad [B] time = 0.37, size = 79, normalized size = 3.16

$$\frac{\sqrt{2}\ln\left(\frac{\frac{\sqrt{2}(-1+xi)1i}{2}-\sqrt{1-x^2}1i}{x-i}\right)1i}{4}-\frac{\sqrt{2}\ln\left(\frac{\frac{\sqrt{2}(1+xi)1i}{2}+\sqrt{1-x^2}1i}{x+1i}\right)1i}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/2)*(x^2 + 1)),x)

[Out] (2^(1/2)*log(((2^(1/2)*(x*1i - 1)*1i)/2 - (1 - x^2)^(1/2)*1i)/(x - 1i))*1i)/4 - (2^(1/2)*log(((2^(1/2)*(x*1i + 1)*1i)/2 + (1 - x^2)^(1/2)*1i)/(x + 1i))*1i)/4

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-(x-1)(x+1)}(x^2+1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(x**2+1)/(-x**2+1)**(1/2), x)
```

```
[Out] Integral(1/(sqrt(-(x - 1)*(x + 1))*(x**2 + 1)), x)
```

$$3.107 \quad \int \frac{1}{(a+bx^2)\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=49

$$\frac{\tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{\sqrt{a}\sqrt{bc-ad}}$$

[Out] arctan(x*(-a*d+b*c)^(1/2)/a^(1/2)/(d*x^2+c)^(1/2))/a^(1/2)/(-a*d+b*c)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {377, 205}

$$\frac{\tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{\sqrt{a}\sqrt{bc-ad}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)*Sqrt[c + d*x^2]),x]

[Out] ArcTan[(Sqrt[b*c - a*d]*x)/(Sqrt[a]*Sqrt[c + d*x^2])]/(Sqrt[a]*Sqrt[b*c - a*d])

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)\sqrt{c+dx^2}} dx &= \text{Subst}\left(\int \frac{1}{a - (-bc+ad)x^2} dx, x, \frac{x}{\sqrt{c+dx^2}}\right) \\ &= \frac{\tan^{-1}\left(\frac{\sqrt{bc-ad}x}{\sqrt{a}\sqrt{c+dx^2}}\right)}{\sqrt{a}\sqrt{bc-ad}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 49, normalized size = 1.00

$$\frac{\tan^{-1}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{c+dx^2}}\right)}{\sqrt{a}\sqrt{bc-ad}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)*Sqrt[c + d*x^2]),x]

[Out] ArcTan[(Sqrt[b*c - a*d]*x)/(Sqrt[a]*Sqrt[c + d*x^2])]/(Sqrt[a]*Sqrt[b*c - a*d])

fricas [B] time = 0.69, size = 241, normalized size = 4.92

$$\left[\frac{\sqrt{-abc + a^2d} \log\left(\frac{(b^2c^2 - 8abcd + 8a^2d^2)x^4 + a^2c^2 - 2(3abc^2 - 4a^2cd)x^2 - 4((bc - 2ad)x^3 - acx)\sqrt{-abc + a^2d}\sqrt{dx^2 + c}}{b^2x^4 + 2abx^2 + a^2}\right)}{4(abc - a^2d)}, \arctan\left(\frac{\sqrt{abc - a^2d}}{2\sqrt{\dots}}\right) \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] [-1/4*sqrt(-a*b*c + a^2*d)*log(((b^2*c^2 - 8*a*b*c*d + 8*a^2*d^2)*x^4 + a^2*c^2 - 2*(3*a*b*c^2 - 4*a^2*c*d)*x^2 - 4*((b*c - 2*a*d)*x^3 - a*c*x)*sqrt(-a*b*c + a^2*d)*sqrt(d*x^2 + c))/(b^2*x^4 + 2*a*b*x^2 + a^2))/(a*b*c - a^2*d), 1/2*arctan(1/2*sqrt(a*b*c - a^2*d)*((b*c - 2*a*d)*x^2 - a*c)*sqrt(d*x^2 + c)/((a*b*c*d - a^2*d^2)*x^3 + (a*b*c^2 - a^2*c*d)*x))/sqrt(a*b*c - a^2*d)]

giac [A] time = 0.60, size = 70, normalized size = 1.43

$$\frac{\sqrt{d} \arctan\left(\frac{(\sqrt{d}x - \sqrt{dx^2 + c})^2 b - bc + 2ad}{2\sqrt{abcd - a^2d^2}}\right)}{\sqrt{abcd - a^2d^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] -sqrt(d)*arctan(1/2*((sqrt(d)*x - sqrt(d*x^2 + c))^2*b - b*c + 2*a*d)/sqrt(a*b*c*d - a^2*d^2))/sqrt(a*b*c*d - a^2*d^2)

maple [B] time = 0.01, size = 306, normalized size = 6.24

$$\frac{\ln\left(\frac{2\sqrt{-ab}\left(x - \frac{\sqrt{-ab}}{b}\right)d - \frac{2(ad-bc)}{b} + 2\sqrt{-\frac{ad-bc}{b}}\sqrt{\left(x - \frac{\sqrt{-ab}}{b}\right)^2 d + \frac{2\sqrt{-ab}\left(x - \frac{\sqrt{-ab}}{b}\right)d - \frac{ad-bc}{b}}}{x - \frac{\sqrt{-ab}}{b}}\right)}{2\sqrt{-ab}\sqrt{-\frac{ad-bc}{b}}}\right) + \ln\left(\frac{2\sqrt{-ab}\left(x + \frac{\sqrt{-ab}}{b}\right)d - \frac{2(ad-bc)}{b} + 2\sqrt{-\frac{ad-bc}{b}}\sqrt{\left(x + \frac{\sqrt{-ab}}{b}\right)^2 d + \frac{2\sqrt{-ab}\left(x + \frac{\sqrt{-ab}}{b}\right)d - \frac{ad-bc}{b}}}{x + \frac{\sqrt{-ab}}{b}}\right)}{2\sqrt{-ab}\sqrt{-\frac{ad-bc}{b}}}\right)}{2\sqrt{-ab}\sqrt{-\frac{ad-bc}{b}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)/(d*x^2+c)^(1/2),x)

[Out] 1/2/(-a*b)^(1/2)/(-a*d-b*c)/b)^(1/2)*ln((-2*(-a*b)^(1/2)*(x+(-a*b)^(1/2)/b)/b*d-2*(a*d-b*c)/b+2*(-(a*d-b*c)/b)^(1/2)*((x+(-a*b)^(1/2)/b)^2*d-2*(-a*b)^(1/2)*(x+(-a*b)^(1/2)/b)/b*d-(a*d-b*c)/b)^(1/2))/(x+(-a*b)^(1/2)/b))-1/2/(-a*b)^(1/2)/(-a*d-b*c)/b)^(1/2)*ln((2*(-a*b)^(1/2)*(x-(-a*b)^(1/2)/b)/b*d-2*(a*d-b*c)/b+2*(-(a*d-b*c)/b)^(1/2)*((x-(-a*b)^(1/2)/b)^2*d+2*(-a*b)^(1/2)*(x-(-a*b)^(1/2)/b)/b*d-(a*d-b*c)/b)^(1/2))/(x-(-a*b)^(1/2)/b))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)*sqrt(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\left\{ \begin{array}{ll} \frac{\operatorname{atan}\left(\frac{x\sqrt{bc-ad}}{\sqrt{a}\sqrt{dx^2+c}}\right)}{\sqrt{-a(ad-bc)}} & \text{if } 0 < bc - ad \\ \frac{\ln\left(\frac{\sqrt{a(dx^2+c)+x}\sqrt{ad-bc}}{\sqrt{a(dx^2+c)-x}\sqrt{ad-bc}}\right)}{2\sqrt{a(ad-bc)}} & \text{if } bc - ad < 0 \\ \int \frac{1}{(bx^2+a)\sqrt{dx^2+c}} dx & \text{if } bc - ad \notin \mathbb{R} \vee ad = bc \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)*(c + d*x^2)^(1/2)),x)

[Out] piecewise(0 < - a*d + b*c, atan((x*(- a*d + b*c)^(1/2))/(a^(1/2)*(c + d*x^2)^(1/2)))/(-a*(a*d - b*c))^(1/2), - a*d + b*c < 0, log(((a*(c + d*x^2))^(1/2) + x*(a*d - b*c)^(1/2))/((a*(c + d*x^2))^(1/2) - x*(a*d - b*c)^(1/2)))/(2*(a*(a*d - b*c))^(1/2)), ~in(- a*d + b*c, 'real') | a*d == b*c, int(1/((a + b*x^2)*(c + d*x^2)^(1/2)), x))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)/(d*x**2+c)**(1/2),x)

[Out] Integral(1/((a + b*x**2)*sqrt(c + d*x**2)), x)

$$3.108 \quad \int \frac{-1+x^2}{(1+x^2)^{3/2}} dx$$

Optimal. Leaf size=15

$$\sinh^{-1}(x) - \frac{2x}{\sqrt{x^2+1}}$$

[Out] arcsinh(x)-2*x/(x^2+1)^(1/2)

Rubi [A] time = 0.00, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {385, 215}

$$\sinh^{-1}(x) - \frac{2x}{\sqrt{x^2+1}}$$

Antiderivative was successfully verified.

[In] Int[(-1 + x^2)/(1 + x^2)^(3/2), x]

[Out] (-2*x)/Sqrt[1 + x^2] + ArcSinh[x]

Rule 215

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[(Rt[b, 2]*x)/Sqrt[a]]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rubi steps

$$\begin{aligned} \int \frac{-1+x^2}{(1+x^2)^{3/2}} dx &= -\frac{2x}{\sqrt{1+x^2}} + \int \frac{1}{\sqrt{1+x^2}} dx \\ &= -\frac{2x}{\sqrt{1+x^2}} + \sinh^{-1}(x) \end{aligned}$$

Mathematica [A] time = 0.01, size = 15, normalized size = 1.00

$$\sinh^{-1}(x) - \frac{2x}{\sqrt{x^2+1}}$$

Antiderivative was successfully verified.

[In] Integrate[(-1 + x^2)/(1 + x^2)^(3/2), x]

[Out] (-2*x)/Sqrt[1 + x^2] + ArcSinh[x]

fricas [B] time = 0.63, size = 44, normalized size = 2.93

$$\frac{2x^2 + (x^2 + 1) \log(-x + \sqrt{x^2 + 1}) + 2\sqrt{x^2 + 1}x + 2}{x^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2-1)/(x^2+1)^(3/2),x, algorithm="fricas")

[Out] $-(2x^2 + (x^2 + 1)\log(-x + \sqrt{x^2 + 1}) + 2\sqrt{x^2 + 1}x + 2)/(x^2 + 1)$

giac [A] time = 0.57, size = 25, normalized size = 1.67

$$-\frac{2x}{\sqrt{x^2 + 1}} - \log\left(-x + \sqrt{x^2 + 1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2-1)/(x^2+1)^(3/2),x, algorithm="giac")

[Out] $-2x/\sqrt{x^2 + 1} - \log(-x + \sqrt{x^2 + 1})$

maple [A] time = 0.01, size = 14, normalized size = 0.93

$$-\frac{2x}{\sqrt{x^2 + 1}} + \operatorname{arcsinh}(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2-1)/(x^2+1)^(3/2),x)

[Out] $\operatorname{arcsinh}(x) - 2x/(x^2+1)^{(1/2)}$

maxima [A] time = 2.78, size = 13, normalized size = 0.87

$$-\frac{2x}{\sqrt{x^2 + 1}} + \operatorname{arsinh}(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2-1)/(x^2+1)^(3/2),x, algorithm="maxima")

[Out] $-2x/\sqrt{x^2 + 1} + \operatorname{arcsinh}(x)$

mupad [B] time = 0.04, size = 27, normalized size = 1.80

$$\frac{\operatorname{asinh}(x) + x^2 \operatorname{asinh}(x) - 2x\sqrt{x^2 + 1}}{x^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2 - 1)/(x^2 + 1)^(3/2),x)

[Out] $(\operatorname{asinh}(x) + x^2 \operatorname{asinh}(x) - 2x(x^2 + 1)^{(1/2)})/(x^2 + 1)$

sympy [B] time = 4.68, size = 31, normalized size = 2.07

$$\frac{x^2 \operatorname{asinh}(x)}{x^2 + 1} - \frac{2x}{\sqrt{x^2 + 1}} + \frac{\operatorname{asinh}(x)}{x^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x**2-1)/(x**2+1)**(3/2),x)

[Out] $x**2*\operatorname{asinh}(x)/(x**2 + 1) - 2x/\sqrt{x**2 + 1} + \operatorname{asinh}(x)/(x**2 + 1)$

$$3.109 \quad \int (a - bx^2)^{2/3} (3a + bx^2)^3 dx$$

Optimal. Leaf size=648

$$\frac{24192\sqrt{2}3^{3/4}a^{13/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{1235bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] 18144/1235*a^3*x*(-b*x^2+a)^(2/3)-23544/6175*a^2*x*(-b*x^2+a)^(5/3)-378/475*a*x*(-b*x^2+a)^(5/3)*(b*x^2+3*a)-3/25*x*(-b*x^2+a)^(5/3)*(b*x^2+3*a)^2-725/76/1235*a^4*x/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))+24192/1235*3^(3/4)*a^(13/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*2^(1/2)*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)-36288/1235*3^(1/4)*a^(13/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))*2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)*((1/2*6^(1/2)+1/2*2^(1/2))/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)

Rubi [A] time = 0.60, antiderivative size = 648, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {416, 528, 388, 195, 235, 304, 219, 1879}

$$\frac{72576a^4x}{1235\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}+\frac{18144a^3x(a-bx^2)^{2/3}}{1235}-\frac{23544a^2x(a-bx^2)^{5/3}}{6175}+\frac{24192\sqrt{2}3^{3/4}a^{13/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{1235bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(2/3)*(3*a + b*x^2)^3,x]

[Out] (18144*a^3*x*(a - b*x^2)^(2/3))/1235 - (23544*a^2*x*(a - b*x^2)^(5/3))/6175 - (378*a*x*(a - b*x^2)^(5/3)*(3*a + b*x^2))/475 - (3*x*(a - b*x^2)^(5/3)*(3*a + b*x^2)^2)/25 - (72576*a^4*x)/(1235*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) - (36288*3^(1/4)*Sqrt[2 + Sqrt[3]]*a^(13/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))]^2)*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(1235*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)))^2]) + (24192*Sqrt[2]*3^(3/4)*a^(13/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))]^2)*EllipticF[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(1235*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)))^2])

Rule 195

Int[(a_) + (b_.)*(x_)^(n_)]^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; Free

$Q[\{a, b\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[p, 0] \&\& (\text{IntegerQ}[2*p] \parallel (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[4*p]) \parallel (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[3*p]) \parallel \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$

Rule 219

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] \text{ :> With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, \text{Simp}[(2*\text{Sqrt}[2 - \text{Sqrt}[3]]*(s + r*x)*\text{Sqrt}[(s^2 - r*s*x + r^2*x^2)]/((1 - \text{Sqrt}[3])*s + r*x)^2*\text{EllipticF}[\text{ArcSin}[(1 + \text{Sqrt}[3])*s + r*x]/((1 - \text{Sqrt}[3])*s + r*x)], -7 + 4*\text{Sqrt}[3])]/(3^{1/4}*r*\text{Sqrt}[a + b*x^3]*\text{Sqrt}[-((s*(s + r*x))/((1 - \text{Sqrt}[3])*s + r*x)^2)]), x]] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 235

$\text{Int}[(a_) + (b_)*(x_)^2]^{-1/3}, x_Symbol] \text{ :> Dist}[(3*\text{Sqrt}[b*x^2])/(2*b*x), \text{Subst}[\text{Int}[x/\text{Sqrt}[-a + x^3], x], x, (a + b*x^2)^{1/3}], x] /; \text{FreeQ}[\{a, b\}, x]$

Rule 304

$\text{Int}[(x_)/\text{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] \text{ :> With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\text{Sqrt}[2]*s)/(\text{Sqrt}[2 - \text{Sqrt}[3]]*r), \text{Int}[1/\text{Sqrt}[a + b*x^3], x], x] + \text{Dist}[1/r, \text{Int}[(1 + \text{Sqrt}[3])*s + r*x]/\text{Sqrt}[a + b*x^3], x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 388

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}]^{(p_)}*((c_) + (d_)*(x_)^{(n_)}), x_Symbol] \text{ :> Simp}[(d*x*(a + b*x^n)^{(p+1)})/(b*(n*(p+1) + 1)), x] - \text{Dist}[(a*d - b*c*(n*(p+1) + 1))/(b*(n*(p+1) + 1)), \text{Int}[(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[n*(p+1) + 1, 0]$

Rule 416

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}]^{(p_)}*((c_) + (d_)*(x_)^{(n_)})^{(q_)}, x_Symbol] \text{ :> Simp}[(d*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-1)})/(b*(n*(p+q) + 1)), x] + \text{Dist}[1/(b*(n*(p+q) + 1)), \text{Int}[(a + b*x^n)^p*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(b*c*(n*(p+q) + 1) - a*d) + d*(b*c*(n*(p+2*q-1) + 1) - a*d*(n*(q-1) + 1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, n, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[q, 1] \&\& \text{NeQ}[n*(p+q) + 1, 0] \&\& !\text{IGtQ}[p, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 528

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}]^{(p_)}*((c_) + (d_)*(x_)^{(n_)})^{(q_)}*((e_) + (f_)*(x_)^{(n_)}), x_Symbol] \text{ :> Simp}[(f*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^q]/(b*(n*(p+q+1) + 1)), x] + \text{Dist}[1/(b*(n*(p+q+1) + 1)), \text{Int}[(a + b*x^n)^p*(c + d*x^n)^{(q-1)}*\text{Simp}[c*(b*e - a*f + b*e*n*(p+q+1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p+q+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \&\& \text{GtQ}[q, 0] \&\& \text{NeQ}[n*(p+q+1) + 1, 0]$

Rule 1879

$\text{Int}[(c_) + (d_)*(x_)]/\text{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] \text{ :> With}[\{r = \text{Numer}[\text{Simplify}[(1 + \text{Sqrt}[3])*d]/c], s = \text{Denom}[\text{Simplify}[(1 + \text{Sqrt}[3])*d]/c]]\}, \text{Simp}[(2*d*s^3*\text{Sqrt}[a + b*x^3])/(a*r^2*((1 - \text{Sqrt}[3])*s + r*x)), x] + \text{Simp}[(3^{1/4}*\text{Sqrt}[2 + \text{Sqrt}[3]]*d*s*(s + r*x)*\text{Sqrt}[(s^2 - r*s*x + r^2*x^2)]/((1 - \text{Sqrt}[3])*s + r*x)^2*\text{EllipticE}[\text{ArcSin}[(1 + \text{Sqrt}[3])*s + r*x]/((1 - \text{Sqrt}[3])*s + r*x)]), x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{GtQ}[d, 0] \&\& \text{NeQ}[a, 0] \&\& \text{NeQ}[b, 0]$

rt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)) / ((1 - Sqrt[3])*s + r*x)^2))), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned} \int (a - bx^2)^{2/3} (3a + bx^2)^3 dx &= -\frac{3}{25}x(a - bx^2)^{5/3} (3a + bx^2)^2 - \frac{3 \int (a - bx^2)^{2/3} (3a + bx^2) (-78a^2b - 42ab^2)}{25b} \\ &= -\frac{378}{475}ax(a - bx^2)^{5/3} (3a + bx^2) - \frac{3}{25}x(a - bx^2)^{5/3} (3a + bx^2)^2 + \frac{9 \int (a - bx^2)^{2/3} (3a + bx^2)^2}{25} \\ &= -\frac{23544a^2x(a - bx^2)^{5/3}}{6175} - \frac{378}{475}ax(a - bx^2)^{5/3} (3a + bx^2) - \frac{3}{25}x(a - bx^2)^{5/3} (3a + bx^2)^2 \\ &= \frac{18144a^3x(a - bx^2)^{2/3}}{1235} - \frac{23544a^2x(a - bx^2)^{5/3}}{6175} - \frac{378}{475}ax(a - bx^2)^{5/3} (3a + bx^2) \\ &= \frac{18144a^3x(a - bx^2)^{2/3}}{1235} - \frac{23544a^2x(a - bx^2)^{5/3}}{6175} - \frac{378}{475}ax(a - bx^2)^{5/3} (3a + bx^2) \\ &= \frac{18144a^3x(a - bx^2)^{2/3}}{1235} - \frac{23544a^2x(a - bx^2)^{5/3}}{6175} - \frac{378}{475}ax(a - bx^2)^{5/3} (3a + bx^2) \\ &= \frac{18144a^3x(a - bx^2)^{2/3}}{1235} - \frac{23544a^2x(a - bx^2)^{5/3}}{6175} - \frac{378}{475}ax(a - bx^2)^{5/3} (3a + bx^2) \end{aligned}$$

Mathematica [C] time = 5.05, size = 99, normalized size = 0.15

$$\frac{3 \left(-40320a^4x \sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1 \left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a} \right) - 15255a^4x + 3390a^3bx^3 + 8992a^2b^2x^5 + 2626ab^3x^7 + 247b^4x^9 \right)}{6175 \sqrt[3]{a - bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a - b*x^2)^(2/3)*(3*a + b*x^2)^3,x]

[Out] (-3*(-15255*a^4*x + 3390*a^3*b*x^3 + 8992*a^2*b^2*x^5 + 2626*a*b^3*x^7 + 247*b^4*x^9 - 40320*a^4*x*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a]))/(6175*(a - b*x^2)^(1/3))

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral} \left((b^3x^6 + 9ab^2x^4 + 27a^2bx^2 + 27a^3)(-bx^2 + a)^{\frac{2}{3}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a)^3,x, algorithm="fricas")

[Out] integral((b^3*x^6 + 9*a*b^2*x^4 + 27*a^2*b*x^2 + 27*a^3)*(-b*x^2 + a)^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)^3 (-bx^2 + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a)^3,x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^3*(-b*x^2 + a)^(2/3), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int (-bx^2 + a)^{\frac{2}{3}} (bx^2 + 3a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(2/3)*(b*x^2+3*a)^3,x)

[Out] int((-b*x^2+a)^(2/3)*(b*x^2+3*a)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)^3 (-bx^2 + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a)^3,x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^3*(-b*x^2 + a)^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (a - bx^2)^{\frac{2}{3}} (bx^2 + 3a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(2/3)*(3*a + b*x^2)^3,x)

[Out] int((a - b*x^2)^(2/3)*(3*a + b*x^2)^3, x)

sympy [A] time = 4.37, size = 136, normalized size = 0.21

$$27a^{\frac{11}{3}} x {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{1}{2} \\ \frac{3}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) + 9a^{\frac{8}{3}} bx^3 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{3}{2} \\ \frac{5}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) + \frac{9a^{\frac{5}{3}} b^2 x^5 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{5}{2} \\ \frac{7}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{5} + \frac{a^{\frac{2}{3}} b^3 x^7 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{7}{2} \\ \frac{9}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(2/3)*(b*x**2+3*a)**3,x)

[Out] 27*a**(11/3)*x*hyper((-2/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) + 9*a**(8/3)*b*x**3*hyper((-2/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a) + 9*a**(5/3)*b**2*x**5*hyper((-2/3, 5/2), (7/2,), b*x**2*exp_polar(2*I*pi)/a)/5 + a**(2/3)*b**3*x**7*hyper((-2/3, 7/2), (9/2,), b*x**2*exp_polar(2*I*pi)/a)/7

$$3.110 \quad \int (a - bx^2)^{2/3} (3a + bx^2)^2 dx$$

Optimal. Leaf size=617

$$\frac{10368\sqrt{2}3^{3/4}a^{10/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{1729bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] 7776/1729*a^2*x*(-b*x^2+a)^(2/3)-252/247*a*x*(-b*x^2+a)^(5/3)-3/19*x*(-b*x^2+a)^(5/3)*(b*x^2+3*a)-31104/1729*a^3*x/((-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))+10368/1729*3^(3/4)*a^(10/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/((-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*2^(1/2)*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/((-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3))/((-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)-15552/1729*3^(1/4)*a^(10/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/((-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/((-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3))/((-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)

Rubi [A] time = 0.43, antiderivative size = 617, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {416, 388, 195, 235, 304, 219, 1879}

$$\frac{31104a^3x}{1729\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}+\frac{7776a^2x(a-bx^2)^{2/3}}{1729}+\frac{10368\sqrt{2}3^{3/4}a^{10/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}{1729bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(2/3)*(3*a + b*x^2)^2,x]

[Out] (7776*a^2*x*(a - b*x^2)^(2/3))/1729 - (252*a*x*(a - b*x^2)^(5/3))/247 - (3*x*(a - b*x^2)^(5/3)*(3*a + b*x^2))/19 - (31104*a^3*x)/(1729*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) - (15552*3^(1/4)*Sqrt[2 + Sqrt[3]]*a^(10/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]]]/(1729*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)]) + (10368*Sqrt[2]*3^(3/4)*a^(10/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)*EllipticF[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]]]/(1729*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)])

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],

Denominator[p]])

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 388

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 416

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)),
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a
, b, c, d, n, p, q, x]
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((
1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int (a - bx^2)^{2/3} (3a + bx^2)^2 dx &= -\frac{3}{19}x(a - bx^2)^{5/3} (3a + bx^2) - \frac{3 \int (a - bx^2)^{2/3} (-60a^2b - 28ab^2x^2) dx}{19b} \\
&= -\frac{252}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{5/3} (3a + bx^2) + \frac{1}{247} (2592a^2) \int (a - bx^2)^{2/3} dx \\
&= \frac{7776a^2x(a - bx^2)^{2/3}}{1729} - \frac{252}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{5/3} (3a + bx^2) + \frac{1}{247} (2592a^2) \int (a - bx^2)^{2/3} dx \\
&= \frac{7776a^2x(a - bx^2)^{2/3}}{1729} - \frac{252}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{5/3} (3a + bx^2) - \frac{1}{247} (2592a^2) \int (a - bx^2)^{2/3} dx \\
&= \frac{7776a^2x(a - bx^2)^{2/3}}{1729} - \frac{252}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{5/3} (3a + bx^2) + \frac{1}{247} (2592a^2) \int (a - bx^2)^{2/3} dx \\
&= \frac{7776a^2x(a - bx^2)^{2/3}}{1729} - \frac{252}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{5/3} (3a + bx^2) - \frac{1}{247} (2592a^2) \int (a - bx^2)^{2/3} dx
\end{aligned}$$

Mathematica [C] time = 3.38, size = 176, normalized size = 0.29

$$\frac{x(a - bx^2)^{2/3} \left(4b \operatorname{Gamma}\left(\frac{1}{3}\right) (3ax + bx^3)^2 {}_3F_2\left(\frac{1}{3}, \frac{3}{2}, 2; 1, \frac{9}{2}; \frac{bx^2}{a}\right) + 8bx^2 \operatorname{Gamma}\left(\frac{1}{3}\right) (18a^2 + 9abx^2 + b^2x^4) \right)}{105a \operatorname{Gamma}\left(-\frac{2}{3}\right) \left(1 - \frac{bx^2}{a}\right)^2}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a - b*x^2)^(2/3)*(3*a + b*x^2)^2,x]

[Out] (x*(a - b*x^2)^(2/3)*(21*a*(45*a^2 + 10*a*b*x^2 + b^2*x^4)*Gamma[-2/3]*Hypergeometric2F1[-2/3, 1/2, 7/2, (b*x^2)/a] + 8*b*x^2*(18*a^2 + 9*a*b*x^2 + b^2*x^4)*Gamma[1/3]*Hypergeometric2F1[1/3, 3/2, 9/2, (b*x^2)/a] + 4*b*(3*a*x + b*x^3)^2*Gamma[1/3]*HypergeometricPFQ[{1/3, 3/2, 2}, {1, 9/2}, (b*x^2)/a])/((105*a*(1 - (b*x^2)/a)^(2/3)*Gamma[-2/3])

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\left(b^2x^4 + 6abx^2 + 9a^2\right)\left(-bx^2 + a\right)^{\frac{2}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a)^2,x, algorithm="fricas")

[Out] integral((b^2*x^4 + 6*a*b*x^2 + 9*a^2)*(-b*x^2 + a)^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)^2 (-bx^2 + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a)^2,x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^2*(-b*x^2 + a)^(2/3), x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int (-bx^2 + a)^{\frac{2}{3}} (bx^2 + 3a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(2/3)*(b*x^2+3*a)^2,x)

[Out] int((-b*x^2+a)^(2/3)*(b*x^2+3*a)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)^2 (-bx^2 + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^2*(-b*x^2 + a)^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (a - bx^2)^{\frac{2}{3}} (bx^2 + 3a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(2/3)*(3*a + b*x^2)^2,x)

[Out] int((a - b*x^2)^(2/3)*(3*a + b*x^2)^2, x)

sympy [A] time = 3.30, size = 99, normalized size = 0.16

$$9a^{\frac{8}{3}}x_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{1}{2} \\ \frac{3}{2} \end{matrix} \middle| \frac{bx^2e^{2i\pi}}{a}\right) + 2a^{\frac{5}{3}}bx^3_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{3}{2} \\ \frac{5}{2} \end{matrix} \middle| \frac{bx^2e^{2i\pi}}{a}\right) + \frac{a^{\frac{2}{3}}b^2x^5_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{5}{2} \\ \frac{7}{2} \end{matrix} \middle| \frac{bx^2e^{2i\pi}}{a}\right)}{5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(2/3)*(b*x**2+3*a)**2,x)

[Out] 9*a**(8/3)*x*hyper((-2/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) + 2*a**(5/3)*b*x**3*hyper((-2/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a) + a**(2/3)*b**2*x**5*hyper((-2/3, 5/2), (7/2,), b*x**2*exp_polar(2*I*pi)/a)/5

3.111 $\int (a - bx^2)^{2/3} (3a + bx^2) dx$

Optimal. Leaf size=588

$$\frac{24\sqrt{2}3^{3/4}a^{7/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{13bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

```
[Out] 18/13*a*x*(-b*x^2+a)^(2/3)-3/13*x*(-b*x^2+a)^(5/3)-72/13*a^2*x/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))+24/13*3^(3/4)*a^(7/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*2^(1/2)*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)-36/13*3^(1/4)*a^(7/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)
```

Rubi [A] time = 0.37, antiderivative size = 588, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {388, 195, 235, 304, 219, 1879}

$$\frac{72a^2x}{13\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}+\frac{24\sqrt{2}3^{3/4}a^{7/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}F\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{13bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

```
[In] Int[(a - b*x^2)^(2/3)*(3*a + b*x^2), x]
```

```
[Out] (18*a*x*(a - b*x^2)^(2/3))/13 - (3*x*(a - b*x^2)^(5/3))/13 - (72*a^2*x)/(13*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) - (36*3^(1/4)*Sqrt[2 + Sqrt[3]]*a^(7/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))]^2)*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(13*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)))^(1/2)] + (24*Sqrt[2]*3^(3/4)*a^(7/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))]^2)*EllipticF[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(13*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)))^(1/2)] - (a - b*x^2)^(1/3))^2])
```

Rule 195

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],
```

Denominator[p]])

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 388

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/
(1 - Sqrt[3])*s + r*x]^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int (a - bx^2)^{2/3} (3a + bx^2) dx &= -\frac{3}{13}x(a - bx^2)^{5/3} + \frac{1}{13}(42a) \int (a - bx^2)^{2/3} dx \\
&= \frac{18}{13}ax(a - bx^2)^{2/3} - \frac{3}{13}x(a - bx^2)^{5/3} + \frac{1}{13}(24a^2) \int \frac{1}{\sqrt[3]{a - bx^2}} dx \\
&= \frac{18}{13}ax(a - bx^2)^{2/3} - \frac{3}{13}x(a - bx^2)^{5/3} - \frac{(36a^2\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{x}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{13bx} \\
&= \frac{18}{13}ax(a - bx^2)^{2/3} - \frac{3}{13}x(a - bx^2)^{5/3} + \frac{(36a^2\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{(1+\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx\right)}{13bx} \\
&= \frac{18}{13}ax(a - bx^2)^{2/3} - \frac{3}{13}x(a - bx^2)^{5/3} - \frac{72a^2x}{13\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} - \frac{36\sqrt[4]{3}}{13}
\end{aligned}$$

Mathematica [C] time = 0.08, size = 62, normalized size = 0.11

$$\frac{3}{13}x(a - bx^2)^{2/3} \left(\frac{14a {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a}\right)}{\left(1 - \frac{bx^2}{a}\right)^{2/3}} - a + bx^2 \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a - b*x^2)^(2/3)*(3*a + b*x^2), x]

[Out] (3*x*(a - b*x^2)^(2/3)*(-a + b*x^2 + (14*a*Hypergeometric2F1[-2/3, 1/2, 3/2, (b*x^2)/a])/(1 - (b*x^2)/a)^(2/3)))/13

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\left(bx^2 + 3a\right)\left(-bx^2 + a\right)^{\frac{2}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a), x, algorithm="fricas")

[Out] integral((b*x^2 + 3*a)*(-b*x^2 + a)^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)(-bx^2 + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a), x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)*(-b*x^2 + a)^(2/3), x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int (-bx^2 + a)^{\frac{2}{3}} (bx^2 + 3a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-b*x^2+a)^(2/3)*(b*x^2+3*a),x)`

[Out] `int((-b*x^2+a)^(2/3)*(b*x^2+3*a),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)(-bx^2 + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x^2+a)^(2/3)*(b*x^2+3*a),x, algorithm="maxima")`

[Out] `integrate((b*x^2 + 3*a)*(-b*x^2 + a)^(2/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (a - bx^2)^{\frac{2}{3}} (bx^2 + 3a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a - b*x^2)^(2/3)*(3*a + b*x^2),x)`

[Out] `int((a - b*x^2)^(2/3)*(3*a + b*x^2), x)`

sympy [A] time = 2.43, size = 63, normalized size = 0.11

$$3a^{\frac{5}{3}}x {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{1}{2} \\ \frac{3}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) + \frac{a^{\frac{2}{3}}bx^3 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{3}{2} \\ \frac{5}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x**2+a)**(2/3)*(b*x**2+3*a),x)`

[Out] `3*a**(5/3)*x*hyper((-2/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) + a**(2/3)*b*x**3*hyper((-2/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a)/3`

$$3.112 \quad \int \frac{(a-bx^2)^{2/3}}{3a+bx^2} dx$$

Optimal. Leaf size=740

$$\frac{\sqrt{2} 3^{3/4} \sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} \operatorname{EllipticF} \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right), 4\sqrt{3} - 7 \right)}{bx \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}$$

[Out] $3*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+2^{(1/3)}*a^{(1/6)}*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/(a^{(1/3)}+2^{(1/3)}*(-b*x^2+a)^{(1/3)})/b^{(1/2)}-1/3*2^{(1/3)}*a^{(1/6)}*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/b^{(1/2)}+1/3*2^{(1/3)}*a^{(1/6)}*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(-b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})/b^{(1/2)}+1/3*2^{(1/3)}*a^{(1/6)}*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})/b^{(1/2)}-3^{(3/4)}*a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}+3/2*3^{(1/4)}*a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.32, antiderivative size = 740, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {396, 235, 304, 219, 1879, 393}

$$\frac{\sqrt{2} 3^{3/4} \sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} F \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right) \middle| -7 + 4\sqrt{3} \right)}{bx \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(2/3)/(3*a + b*x^2), x]

[Out] $(3*x)/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}) + (2^{(1/3)}*a^{(1/6)}*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[b]) + (2^{(1/3)}*a^{(1/6)}*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)} - 2^{(1/3)}*(a - b*x^2)^{(1/3)})/(\operatorname{Sqrt}[b]*x)]/(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[b]) - (2^{(1/3)}*a^{(1/6)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]])/(3*\operatorname{Sqrt}[b]) + (2^{(1/3)}*a^{(1/6)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{(1/6)}*(a^{(1/3)} + 2^{(1/3)}*(a - b*x^2)^{(1/3)}))]/\operatorname{Sqrt}[b] + (3*3^{(1/4)}*\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]]*a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})], -7 + 4*\operatorname{Sqrt}[3]))/(2*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]) - (\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}), -7 + 4*\operatorname{Sqrt}[3]))/(b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])])$

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 393

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Wit
h[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^
(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2
)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(
1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)
))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]]] /; FreeQ[{a, b, c, d}
, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]
```

Rule 396

```
Int[((a_) + (b_.)*(x_)^2)^(2/3)/((c_) + (d_.)*(x_)^2), x_Symbol] := Dist[b/
d, Int[1/(a + b*x^2)^(1/3), x], x] - Dist[(b*c - a*d)/d, Int[1/((a + b*x^2)
^(1/3)*(c + d*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]
&& EqQ[b*c + 3*a*d, 0]
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/(
(1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a - bx^2)^{2/3}}{3a + bx^2} dx &= (4a) \int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx - \int \frac{1}{\sqrt[3]{a - bx^2}} dx \\
&= \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} + \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a - bx^2})}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} - \frac{\sqrt[3]{2} \sqrt[6]{a} \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{3\sqrt{b}} + \\
&= \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} + \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a - bx^2})}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} - \frac{\sqrt[3]{2} \sqrt[6]{a} \tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{3\sqrt{b}} + \\
&= \frac{3x}{(1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2}} + \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} + \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a - bx^2})}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}}
\end{aligned}$$

Mathematica [C] time = 0.16, size = 162, normalized size = 0.22

$$\frac{9ax(a - bx^2)^{2/3} F_1\left(\frac{1}{2}; -\frac{2}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{(3a + bx^2) \left(9a F_1\left(\frac{1}{2}; -\frac{2}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - 2bx^2 \left(F_1\left(\frac{3}{2}; -\frac{2}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) + 2F_1\left(\frac{3}{2}; \frac{1}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a - b*x^2)^(2/3)/(3*a + b*x^2), x]

[Out] (9*a*x*(a - b*x^2)^(2/3)*AppellF1[1/2, -2/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/((3*a + b*x^2)*(9*a*AppellF1[1/2, -2/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] - 2*b*x^2*(AppellF1[3/2, -2/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a), x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{2/3}}{bx^2 + 3a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a), x, algorithm="giac")

[Out] integrate((-b*x^2 + a)^(2/3)/(b*x^2 + 3*a), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{2/3}}{bx^2 + 3a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-b*x^2+a)^(2/3)/(b*x^2+3*a),x)`

[Out] `int((-b*x^2+a)^(2/3)/(b*x^2+3*a),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{bx^2 + 3a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a),x, algorithm="maxima")`

[Out] `integrate((-b*x^2 + a)^(2/3)/(b*x^2 + 3*a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a - bx^2)^{\frac{2}{3}}}{bx^2 + 3a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a - b*x^2)^(2/3)/(3*a + b*x^2),x)`

[Out] `int((a - b*x^2)^(2/3)/(3*a + b*x^2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a - bx^2)^{\frac{2}{3}}}{3a + bx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x**2+a)**(2/3)/(b*x**2+3*a),x)`

[Out] `Integral((a - b*x**2)**(2/3)/(3*a + b*x**2), x)`

$$3.113 \quad \int \frac{(a-bx^2)^{2/3}}{(3a+bx^2)^2} dx$$

Optimal. Leaf size=584

$$\frac{\left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}}\right), 4\sqrt{3} - 7\right) \sqrt{2 + \sqrt{3}} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{3\sqrt{2} \sqrt[4]{3} a^{2/3} bx \sqrt{-\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] $1/6*x*(-b*x^2+a)^{(2/3)}/a/(b*x^2+3*a)-1/6*x/a/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+1/18*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}(((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*3^{(3/4)}/a^{(2/3)}/b/x*2^{(1/2)}/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}-1/12*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}(((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})*3^{(1/4)}/a^{(2/3)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.37, antiderivative size = 584, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {412, 21, 235, 304, 219, 1879}

$$\frac{\left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}} F\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}}\right) \middle| -7 + 4\sqrt{3}\right) \sqrt{2 + \sqrt{3}} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{3\sqrt{2} \sqrt[4]{3} a^{2/3} bx \sqrt{-\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a - b*x^2)^{(2/3)}/(3*a + b*x^2)^2, x]$

[Out] $(x*(a - b*x^2)^{(2/3)})/(6*a*(3*a + b*x^2)) - x/(6*a*((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})) - (\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]]*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}))], -7 + 4*\operatorname{Sqrt}[3])]/(4*3^{(3/4)}*a^{(2/3)}*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]) + ((a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}))], -7 + 4*\operatorname{Sqrt}[3])]/(3*\operatorname{Sqrt}[2]*3^{(1/4)}*a^{(2/3)}*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])$

Rule 21

$\operatorname{Int}[(u_.)*((a_.) + (b_.)*(v_))^{(m_.)}*((c_.) + (d_.)*(v_))^{(n_.)}, x_Symbol] := \operatorname{Dist}[(b/d)^m, \operatorname{Int}[u*(c + d*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{a, b, c, d, n\}, x] \&\& \operatorname{EqQ}[b*c - a*d, 0] \&\& \operatorname{IntegerQ}[m] \&\& (!\operatorname{IntegerQ}[n] || \operatorname{SimplerQ}[c + d*x, a + b*x])$

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 412

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*n*(p + 1)), x] + Dist[1/(
a*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(n*(p + 1)
+ 1) + d*(n*(p + q + 1) + 1)*x^n, x], x]] /; FreeQ[{a, b, c, d, n}, x]
&& NeQ[b*c - a*d, 0] && LtQ[p, -1] && LtQ[0, q, 1] && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/(
(1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a - bx^2)^{2/3}}{(3a + bx^2)^2} dx &= \frac{x(a - bx^2)^{2/3}}{6a(3a + bx^2)} - \frac{\int \frac{-a - bx^2}{\sqrt[3]{a - bx^2}(3a + bx^2)} dx}{6a} \\
&= \frac{x(a - bx^2)^{2/3}}{6a(3a + bx^2)} + \frac{\int \frac{1}{\sqrt[3]{a - bx^2}} dx}{18a} \\
&= \frac{x(a - bx^2)^{2/3}}{6a(3a + bx^2)} - \frac{\sqrt{-bx^2} \operatorname{Subst}\left(\int \frac{x}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{12abx} \\
&= \frac{x(a - bx^2)^{2/3}}{6a(3a + bx^2)} + \frac{\sqrt{-bx^2} \operatorname{Subst}\left(\int \frac{(1+\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{12abx} - \frac{\left(\sqrt{\frac{1}{2}(2 + \sqrt{3})} \sqrt{-bx^2}\right)}{12abx} \\
&= \frac{x(a - bx^2)^{2/3}}{6a(3a + bx^2)} - \frac{x}{6a\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} - \frac{\sqrt{2 + \sqrt{3}}\left(\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)}{4 \cdot 3^{3/4} a^2} \sqrt{\frac{a^{2/3} + \sqrt[3]{a}}{(1 - \sqrt{3})}}
\end{aligned}$$

Mathematica [C] time = 0.06, size = 86, normalized size = 0.15

$$\frac{x \sqrt[3]{\frac{a - bx^2}{a}} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a}\right)}{18a \sqrt[3]{a - bx^2}} + \frac{x(a - bx^2)^{2/3}}{6a(3a + bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(a - b*x^2)^(2/3)/(3*a + b*x^2)^2, x]

[Out] (x*(a - b*x^2)^(2/3))/(6*a*(3*a + b*x^2)) + (x*((a - b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a])/(18*a*(a - b*x^2)^(1/3))

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\frac{(-bx^2 + a)^{\frac{2}{3}}}{b^2x^4 + 6abx^2 + 9a^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^2,x, algorithm="fricas")

[Out] integral((-b*x^2 + a)^(2/3)/(b^2*x^4 + 6*a*b*x^2 + 9*a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^2,x, algorithm="giac")

[Out] integrate((-b*x^2 + a)^(2/3)/(b*x^2 + 3*a)^2, x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(2/3)/(b*x^2+3*a)^2,x)

[Out] int((-b*x^2+a)^(2/3)/(b*x^2+3*a)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^2,x, algorithm="maxima")

[Out] integrate((-b*x^2 + a)^(2/3)/(b*x^2 + 3*a)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a - bx^2)^{\frac{2}{3}}}{(bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(2/3)/(3*a + b*x^2)^2,x)

[Out] int((a - b*x^2)^(2/3)/(3*a + b*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a - bx^2)^{\frac{2}{3}}}{(3a + bx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(2/3)/(b*x**2+3*a)**2,x)

[Out] Integral((a - b*x**2)**(2/3)/(3*a + b*x**2)**2, x)

$$3.114 \quad \int \frac{(a-bx^2)^{2/3}}{(3a+bx^2)^3} dx$$

Optimal. Leaf size=818

$$\frac{(a-bx^2)^{2/3} x}{36a^2 (bx^2+3a)} - \frac{x}{36a^2 \left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)} + \frac{(a-bx^2)^{2/3} x}{12a (bx^2+3a)^2} + \frac{\tan^{-1} \left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}} \right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\sqrt{bx}} \right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}}$$

[Out] 1/12*x*(-b*x^2+a)^(2/3)/a/(b*x^2+3*a)^2+1/36*x*(-b*x^2+a)^(2/3)/a^2/(b*x^2+3*a)-1/36*x/a^2/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))+1/144*arctanh(x*b^(1/2)/a^(1/6)/(a^(1/3)+2^(1/3)*(-b*x^2+a)^(1/3)))*2^(1/3)/a^(11/6)/b^(1/2)-1/432*arctanh(x*b^(1/2)/a^(1/2))*2^(1/3)/a^(11/6)/b^(1/2)+1/432*arctan(a^(1/6)*(a^(1/3)-2^(1/3)*(-b*x^2+a)^(1/3))*3^(1/2)/x/b^(1/2))*2^(1/3)/a^(11/6)*3^(1/2)/b^(1/2)+1/432*arctan(3^(1/2)*a^(1/2)/x/b^(1/2))*2^(1/3)/a^(11/6)*3^(1/2)/b^(1/2)+1/108*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)*3^(3/4)/a^(5/3)/b/x*2^(1/2)/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)-1/72*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))*3^(1/4)/a^(5/3)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)

Rubi [A] time = 0.56, antiderivative size = 818, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {412, 527, 530, 235, 304, 219, 1879, 393}

$$\frac{(a-bx^2)^{2/3} x}{36a^2 (bx^2+3a)} - \frac{x}{36a^2 \left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)} + \frac{(a-bx^2)^{2/3} x}{12a (bx^2+3a)^2} + \frac{\tan^{-1} \left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}} \right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\sqrt{bx}} \right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(2/3)/(3*a + b*x^2)^3,x]

[Out] (x*(a - b*x^2)^(2/3))/(12*a*(3*a + b*x^2)^2) + (x*(a - b*x^2)^(2/3))/(36*a^2*(3*a + b*x^2)) - x/(36*a^2*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) + ArcTan[(Sqrt[3]*Sqrt[a])/(Sqrt[b]*x)]/(72*2^(2/3)*Sqrt[3]*a^(11/6)*Sqrt[b]) + ArcTan[(Sqrt[3]*a^(1/6)*(a^(1/3) - 2^(1/3)*(a - b*x^2)^(1/3)))/(Sqrt[b]*x)]/(72*2^(2/3)*Sqrt[3]*a^(11/6)*Sqrt[b]) - ArcTanh[(Sqrt[b]*x)/Sqrt[a]]/(216*2^(2/3)*a^(11/6)*Sqrt[b]) + ArcTanh[(Sqrt[b]*x)/(a^(1/6)*(a^(1/3) + 2^(1/3)*(a - b*x^2)^(1/3)))]/(72*2^(2/3)*a^(11/6)*Sqrt[b]) - (Sqrt[2 + Sqrt[3]]*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/(1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)], -7 + 4*Sqrt[3]]]/(24*3^(3/4)*a^(5/3)*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)]) + ((a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)))

$$\int \frac{1}{\sqrt{(a + b x^3)}} \operatorname{EllipticF}\left[\operatorname{ArcSin}\left(\frac{(1 + \sqrt{3})a^{1/3} - (a - b x^2)^{1/3}}{(1 - \sqrt{3})a^{1/3} - (a - b x^2)^{1/3}}\right), -7 + 4\sqrt{3}\right] / (18\sqrt{2} \cdot 3^{1/4} a^{5/3} b x \sqrt{-(a^{1/3}(a^{1/3} - (a - b x^2)^{1/3}) / ((1 - \sqrt{3})a^{1/3} - (a - b x^2)^{1/3}))^2})}{1}$$
Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]]]/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2))], x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x),
Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b},
x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sqr
t[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 393

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Wit
h[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(
1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)
)^(1/3)])/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x]/(6*2^(2/3)*a^(
1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3))
)/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]]) /; FreeQ[{a, b, c, d},
x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]
```

Rule 412

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*n*(p + 1)), x] + Dist[1/(
a*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(n*(p + 1)
+ 1) + d*(n*(p + q + 1) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x]
&& NeQ[b*c - a*d, 0] && LtQ[p, -1] && LtQ[0, q, 1] && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 530

```
Int[(((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p}
```

, n}, x]

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2))], x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{(a - bx^2)^{2/3}}{(3a + bx^2)^3} dx &= \frac{x(a - bx^2)^{2/3}}{12a(3a + bx^2)^2} - \frac{\int \frac{-3a + \frac{5bx^2}{3}}{\sqrt[3]{a - bx^2}(3a + bx^2)^2} dx}{12a} \\ &= \frac{x(a - bx^2)^{2/3}}{12a(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{36a^2(3a + bx^2)} + \frac{\int \frac{16a^2b + \frac{8}{3}ab^2x^2}{\sqrt[3]{a - bx^2}(3a + bx^2)} dx}{288a^3b} \\ &= \frac{x(a - bx^2)^{2/3}}{12a(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{36a^2(3a + bx^2)} + \frac{\int \frac{1}{\sqrt[3]{a - bx^2}} dx}{108a^2} + \frac{\int \frac{1}{\sqrt[3]{a - bx^2}(3a + bx^2)} dx}{36a} \\ &= \frac{x(a - bx^2)^{2/3}}{12a(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{36a^2(3a + bx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{bx}}\right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} \\ &= \frac{x(a - bx^2)^{2/3}}{12a(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{36a^2(3a + bx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{bx}}\right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} \\ &= \frac{x(a - bx^2)^{2/3}}{12a(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{36a^2(3a + bx^2)} - \frac{x}{36a^2\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{72 \cdot 2^{2/3} \sqrt{3} a^{11/6}} \end{aligned}$$

Mathematica [C] time = 0.32, size = 252, normalized size = 0.31

$$\frac{bx^3 \sqrt[3]{1 - \frac{bx^2}{a}} F_1\left(\frac{3}{2}; \frac{1}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{a^3} + \frac{27x \left(-\frac{b^2x^4}{a^2} + \frac{18(3a + bx^2) F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - \frac{5bx^2}{a} + 6 \right)}{(3a + bx^2)^2}$$

$$972 \sqrt[3]{a - bx^2}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a - b*x^2)^(2/3)/(3*a + b*x^2)^3,x]

[Out] ((b*x^3*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])/a^3 + (27*x*(6 - (5*b*x^2)/a - (b^2*x^4)/a^2 + (18*(3*a + b*x^2)*

AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a]/(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))/(3*a + b*x^2)^2/(972*(a - b*x^2)^(1/3))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^3,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^3,x, algorithm="giac")

[Out] integrate((-b*x^2 + a)^(2/3)/(b*x^2 + 3*a)^3, x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(2/3)/(b*x^2+3*a)^3,x)

[Out] int((-b*x^2+a)^(2/3)/(b*x^2+3*a)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^3,x, algorithm="maxima")

[Out] integrate((-b*x^2 + a)^(2/3)/(b*x^2 + 3*a)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a - bx^2)^{\frac{2}{3}}}{(bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(2/3)/(3*a + b*x^2)^3,x)

[Out] int((a - b*x^2)^(2/3)/(3*a + b*x^2)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a - bx^2)^{\frac{2}{3}}}{(3a + bx^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(2/3)/(b*x**2+3*a)**3,x)

[Out] Integral((a - b*x**2)**(2/3)/(3*a + b*x**2)**3, x)

$$3.115 \quad \int \frac{(a-bx^2)^{2/3}}{(3a+bx^2)^4} dx$$

Optimal. Leaf size=849

$$\frac{(a-bx^2)^{2/3} x}{144a^3 (bx^2+3a)} - \frac{x}{144a^3 \left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)} + \frac{(a-bx^2)^{2/3} x}{54a^2 (bx^2+3a)^2} + \frac{(a-bx^2)^{2/3} x}{18a (bx^2+3a)^3} + \frac{7 \tan^{-1} \left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}} \right)}{1296 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}}$$

[Out] 1/18*x*(-b*x^2+a)^(2/3)/a/(b*x^2+3*a)^3+1/54*x*(-b*x^2+a)^(2/3)/a^2/(b*x^2+3*a)^2+1/144*x*(-b*x^2+a)^(2/3)/a^3/(b*x^2+3*a)-1/144*x/a^3/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))) + 7/2592*arctanh(x*b^(1/2)/a^(1/6))/(a^(1/3)+2^(1/3)*(-b*x^2+a)^(1/3))*2^(1/3)/a^(17/6)/b^(1/2)-7/7776*arctanh(x*b^(1/2)/a^(1/2))*2^(1/3)/a^(17/6)/b^(1/2)+7/7776*arctan(a^(1/6)*(a^(1/3)-2^(1/3)*(-b*x^2+a)^(1/3))*3^(1/2)/x/b^(1/2))*2^(1/3)/a^(17/6)*3^(1/2)/b^(1/2)+7/7776*arctan(3^(1/2)*a^(1/2)/x/b^(1/2))*2^(1/3)/a^(17/6)*3^(1/2)/b^(1/2)+1/432*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))), 2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)*3^(3/4)/a^(8/3)/b/x*2^(1/2)/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)-1/288*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))), 2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))*3^(1/4)/a^(8/3)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)

Rubi [A] time = 0.65, antiderivative size = 849, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {412, 527, 530, 235, 304, 219, 1879, 393}

$$\frac{(a-bx^2)^{2/3} x}{144a^3 (bx^2+3a)} - \frac{x}{144a^3 \left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)} + \frac{(a-bx^2)^{2/3} x}{54a^2 (bx^2+3a)^2} + \frac{(a-bx^2)^{2/3} x}{18a (bx^2+3a)^3} + \frac{7 \tan^{-1} \left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}} \right)}{1296 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(2/3)/(3*a + b*x^2)^4,x]

[Out] (x*(a - b*x^2)^(2/3))/(18*a*(3*a + b*x^2)^3) + (x*(a - b*x^2)^(2/3))/(54*a^2*(3*a + b*x^2)^2) + (x*(a - b*x^2)^(2/3))/(144*a^3*(3*a + b*x^2)) - x/(144*a^3*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) + (7*ArcTan[(Sqrt[3]*Sqrt[a])/(Sqrt[b]*x)])/(1296*2^(2/3)*Sqrt[3]*a^(17/6)*Sqrt[b]) + (7*ArcTan[(Sqrt[3]*a^(1/6)*(a^(1/3) - 2^(1/3)*(a - b*x^2)^(1/3)))/(Sqrt[b]*x)])/(1296*2^(2/3)*Sqrt[3]*a^(17/6)*Sqrt[b]) - (7*ArcTanh[(Sqrt[b]*x)/Sqrt[a]])/(3888*2^(2/3)*a^(17/6)*Sqrt[b]) + (7*ArcTanh[(Sqrt[b]*x)/(a^(1/6)*(a^(1/3) + 2^(1/3)*(a - b*x^2)^(1/3)))]/(1296*2^(2/3)*a^(17/6)*Sqrt[b]) - (Sqrt[2 + Sqrt[3]]*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(96*3^(3/4)*a^(8/3)*b*x*Sqrt[-(a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))])

$$\frac{1}{3})^2)] + ((a^{1/3} - (a - b*x^2)^{1/3}) * \text{Sqrt}[(a^{2/3} + a^{1/3} * (a - b*x^2)^{1/3} + (a - b*x^2)^{2/3}) / ((1 - \text{Sqrt}[3]) * a^{1/3} - (a - b*x^2)^{1/3})]^2 * \text{EllipticF}[\text{ArcSin}[\frac{(1 + \text{Sqrt}[3]) * a^{1/3} - (a - b*x^2)^{1/3}}{(1 - \text{Sqrt}[3]) * a^{1/3} - (a - b*x^2)^{1/3}}], -7 + 4 * \text{Sqrt}[3]]) / (72 * \text{Sqrt}[2] * 3^{1/4} * a^{8/3} * b * x * \text{Sqrt}[-((a^{1/3} * (a^{1/3} - (a - b*x^2)^{1/3})) / ((1 - \text{Sqrt}[3]) * a^{1/3} - (a - b*x^2)^{1/3})^2)])$$

Rule 219

$$\text{Int}[1/\text{Sqrt}[(a_) + (b_.)*(x_)^3], x_Symbol] := \text{With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, \text{Simp}[(2 * \text{Sqrt}[2 - \text{Sqrt}[3]] * (s + r*x) * \text{Sqrt}[(s^2 - r*s*x + r^2*x^2) / ((1 - \text{Sqrt}[3]) * s + r*x)]^2 * \text{EllipticF}[\text{ArcSin}[\frac{(1 + \text{Sqrt}[3]) * s + r*x}{(1 - \text{Sqrt}[3]) * s + r*x}], -7 + 4 * \text{Sqrt}[3]]) / (3^{1/4} * r * \text{Sqrt}[a + b*x^3] * \text{Sqrt}[-((s*(s + r*x)) / ((1 - \text{Sqrt}[3]) * s + r*x)^2)])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a]$$

Rule 235

$$\text{Int}[(a_) + (b_.)*(x_)^2]^{-1/3}, x_Symbol] := \text{Dist}[(3 * \text{Sqrt}[b*x^2]) / (2 * b * x), \text{Subst}[\text{Int}[x/\text{Sqrt}[-a + x^3], x], x, (a + b*x^2)^{1/3}], x] /; \text{FreeQ}[\{a, b\}, x]$$

Rule 304

$$\text{Int}[(x_)/\text{Sqrt}[(a_) + (b_.)*(x_)^3], x_Symbol] := \text{With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\text{Sqrt}[2] * s) / (\text{Sqrt}[2 - \text{Sqrt}[3]] * r), \text{Int}[1/\text{Sqrt}[a + b*x^3], x], x] + \text{Dist}[1/r, \text{Int}[\frac{(1 + \text{Sqrt}[3]) * s + r*x}{\text{Sqrt}[a + b*x^3]}, x], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a]$$

Rule 393

$$\text{Int}[1/((a_) + (b_.)*(x_)^2)^{1/3} * ((c_) + (d_.)*(x_)^2), x_Symbol] := \text{With}[\{q = \text{Rt}[-(b/a), 2]\}, \text{Simp}[(q * \text{ArcTan}[\text{Sqrt}[3] / (q*x)]) / (2 * 2^{2/3} * \text{Sqrt}[3] * a^{1/3} * d), x] + (\text{Simp}[(q * \text{ArcTanh}[(a^{1/3} * q*x) / (a^{1/3} + 2^{1/3} * (a + b*x^2)^{1/3})]) / (2 * 2^{2/3} * a^{1/3} * d), x] - \text{Simp}[(q * \text{ArcTanh}[q*x]) / (6 * 2^{2/3} * a^{1/3} * d), x] + \text{Simp}[(q * \text{ArcTan}[(\text{Sqrt}[3] * (a^{1/3} - 2^{1/3} * (a + b*x^2)^{1/3})) / (a^{1/3} * q*x)]) / (2 * 2^{2/3} * \text{Sqrt}[3] * a^{1/3} * d), x])] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[b*c + 3*a*d, 0] \&\& \text{NegQ}[b/a]$$

Rule 412

$$\text{Int}[(a_) + (b_.)*(x_)^{(n_)}]^{(p_)} * ((c_) + (d_.)*(x_)^{(n_)})^{(q_)}, x_Symbol] := -\text{Simp}[(x * (a + b*x^n)^{(p+1)} * (c + d*x^n)^q) / (a*n*(p+1)), x] + \text{Dist}[1/(a*n*(p+1)), \text{Int}[(a + b*x^n)^{(p+1)} * (c + d*x^n)^{(q-1)} * \text{Simp}[c*(n*(p+1) + 1) + d*(n*(p+q+1) + 1)*x^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[p, -1] \&\& \text{LtQ}[0, q, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$$

Rule 527

$$\text{Int}[(a_) + (b_.)*(x_)^{(n_)}]^{(p_)} * ((c_) + (d_.)*(x_)^{(n_)})^{(q_)} * ((e_) + (f_.)*(x_)^{(n_)}), x_Symbol] := -\text{Simp}[(b*e - a*f) * x * (a + b*x^n)^{(p+1)} * (c + d*x^n)^{(q+1)} / (a*n*(b*c - a*d)*(p+1)), x] + \text{Dist}[1/(a*n*(b*c - a*d)*(p+1)), \text{Int}[(a + b*x^n)^{(p+1)} * (c + d*x^n)^q * \text{Simp}[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2) + 1)*x^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, q\}, x] \&\& \text{LtQ}[p, -1]$$

Rule 530

$$\text{Int}[(a_) + (b_.)*(x_)^{(n_)}]^{(p_)} * ((e_) + (f_.)*(x_)^{(n_)})) / ((c_) + (d_.)*$$

$(x_)^{(n_)}), x_Symbol] :> \text{Dist}[f/d, \text{Int}[(a + b*x^n)^p, x], x] + \text{Dist}[(d*e - c*f)/d, \text{Int}[(a + b*x^n)^p/(c + d*x^n), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, p, n\}, x]$

Rule 1879

$\text{Int}[(c_ + (d_)*(x_))/\text{Sqrt}[(a_ + (b_)*(x_)^3], x_Symbol] :> \text{With}\{r = \text{Numerator}[\text{Simplify}[(1 + \text{Sqrt}[3])*d]/c], s = \text{Denom}[\text{Simplify}[(1 + \text{Sqrt}[3])*d]/c]\}, \text{Simp}[(2*d*s^3*\text{Sqrt}[a + b*x^3])/(a*r^2*((1 - \text{Sqrt}[3])*s + r*x)), x] + \text{Simp}[(3^{(1/4)}*\text{Sqrt}[2 + \text{Sqrt}[3])*d*s*(s + r*x)*\text{Sqrt}[(s^2 - r*s*x + r^2*x^2)/(1 - \text{Sqrt}[3])*s + r*x]^2*\text{EllipticE}[\text{ArcSin}[(1 + \text{Sqrt}[3])*s + r*x]/((1 - \text{Sqrt}[3])*s + r*x)], -7 + 4*\text{Sqrt}[3]])/(r^2*\text{Sqrt}[a + b*x^3]*\text{Sqrt}[-((s*(s + r*x))/(1 - \text{Sqrt}[3])*s + r*x)^2])), x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NegQ}[a] \&\& \text{EqQ}[b*c^3 - 2*(5 + 3*\text{Sqrt}[3])*a*d^3, 0]$

Rubi steps

$$\begin{aligned} \int \frac{(a - bx^2)^{2/3}}{(3a + bx^2)^4} dx &= \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)^3} - \frac{\int \frac{-5a + \frac{11bx^2}{3}}{\sqrt[3]{a - bx^2}(3a + bx^2)^3} dx}{18a} \\ &= \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)^3} + \frac{x(a - bx^2)^{2/3}}{54a^2(3a + bx^2)^2} + \frac{\int \frac{64a^2b - \frac{80}{3}ab^2x^2}{\sqrt[3]{a - bx^2}(3a + bx^2)^2} dx}{864a^3b} \\ &= \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)^3} + \frac{x(a - bx^2)^{2/3}}{54a^2(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{144a^3(3a + bx^2)} - \frac{\int \frac{-368a^3b^2 - 48a^2b^3x^2}{\sqrt[3]{a - bx^2}(3a + bx^2)} dx}{20736a^5b^2} \\ &= \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)^3} + \frac{x(a - bx^2)^{2/3}}{54a^2(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{144a^3(3a + bx^2)} + \frac{\int \frac{1}{\sqrt[3]{a - bx^2}} dx}{432a^3} + \frac{7 \int \frac{1}{\sqrt[3]{a - bx^2}(3a + bx^2)}}{648a^2} \\ &= \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)^3} + \frac{x(a - bx^2)^{2/3}}{54a^2(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{144a^3(3a + bx^2)} + \frac{7 \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx^2}}\right)}{1296 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{7 \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx^2}}\right)}{1296 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} \\ &= \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)^3} + \frac{x(a - bx^2)^{2/3}}{54a^2(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{144a^3(3a + bx^2)} + \frac{7 \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx^2}}\right)}{1296 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{7 \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx^2}}\right)}{1296 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} \\ &= \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)^3} + \frac{x(a - bx^2)^{2/3}}{54a^2(3a + bx^2)^2} + \frac{x(a - bx^2)^{2/3}}{144a^3(3a + bx^2)} - \frac{x}{144a^3 \left((1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2} \right)} \end{aligned}$$

Mathematica [C] time = 0.19, size = 265, normalized size = 0.31

$$\frac{x \left(\frac{621a^3 F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{(3a + bx^2) \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right)}{3888a^4 \sqrt[3]{a - bx^2}} + \frac{9a(a - bx^2)(75a^2 + 26abx^2 + 3b^2x^4)}{(3a + bx^2)^3} + bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} \right)}{3888a^4 \sqrt[3]{a - bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a - b*x^2)^(2/3)/(3*a + b*x^2)^4,x]

[Out] (x*((9*a*(a - b*x^2)*(75*a^2 + 26*a*b*x^2 + 3*b^2*x^4))/(3*a + b*x^2)^3 + b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + (621*a^3*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a]))/((3*a + b*x^2)*(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))))/(3888*a^4*(a - b*x^2)^(1/3))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^4,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^4,x, algorithm="giac")

[Out] integrate((-b*x^2 + a)^(2/3)/(b*x^2 + 3*a)^4, x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(2/3)/(b*x^2+3*a)^4,x)

[Out] int((-b*x^2+a)^(2/3)/(b*x^2+3*a)^4,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{2}{3}}}{(bx^2 + 3a)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(2/3)/(b*x^2+3*a)^4,x, algorithm="maxima")

[Out] integrate((-b*x^2 + a)^(2/3)/(b*x^2 + 3*a)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a - bx^2)^{\frac{2}{3}}}{(bx^2 + 3a)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a - b*x^2)^(2/3)/(3*a + b*x^2)^4, x)`

[Out] `int((a - b*x^2)^(2/3)/(3*a + b*x^2)^4, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a - bx^2)^{\frac{2}{3}}}{(3a + bx^2)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x**2+a)**(2/3)/(b*x**2+3*a)**4, x)`

[Out] `Integral((a - b*x**2)**(2/3)/(3*a + b*x**2)**4, x)`

$$3.116 \quad \int (a - bx^2)^{5/3} (3a + bx^2)^3 dx$$

Optimal. Leaf size=668

$$\frac{3746304\sqrt{2}3^{3/4}a^{16/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}\right)}{267995bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] 2809728/267995*a^4*x*(-b*x^2+a)^(2/3)+1404864/191425*a^3*x*(-b*x^2+a)^(5/3)-33264/14725*a^2*x*(-b*x^2+a)^(8/3)-432/775*a*x*(-b*x^2+a)^(8/3)*(b*x^2+3*a)-3/31*x*(-b*x^2+a)^(8/3)*(b*x^2+3*a)^2-11238912/267995*a^5*x/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))+3746304/267995*3^(3/4)*a^(16/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*2^(1/2)*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^(1/2)-5619456/267995*3^(1/4)*a^(16/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^(1/2)

Rubi [A] time = 0.57, antiderivative size = 668, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {416, 528, 388, 195, 235, 304, 219, 1879}

$$\frac{11238912a^5x}{267995\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}+\frac{2809728a^4x(a-bx^2)^{2/3}}{267995}+\frac{1404864a^3x(a-bx^2)^{5/3}}{191425}-\frac{33264a^2x(a-bx^2)^{8/3}}{14725}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(5/3)*(3*a + b*x^2)^3,x]

[Out] (2809728*a^4*x*(a - b*x^2)^(2/3))/267995 + (1404864*a^3*x*(a - b*x^2)^(5/3))/191425 - (33264*a^2*x*(a - b*x^2)^(8/3))/14725 - (432*a*x*(a - b*x^2)^(8/3)*(3*a + b*x^2))/775 - (3*x*(a - b*x^2)^(8/3)*(3*a + b*x^2)^2)/31 - (11238912*a^5*x)/(267995*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) - (5619456*3^(1/4)*Sqrt[2 + Sqrt[3]]*a^(16/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(267995*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2])) + (3746304*Sqrt[2]*3^(3/4)*a^(16/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]*EllipticF[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(267995*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]))

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; Free

$Q[\{a, b\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[p, 0] \&\& (\text{IntegerQ}[2*p] \parallel (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[4*p]) \parallel (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[3*p]) \parallel \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$

Rule 219

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] \text{ :> With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, \text{Simp}[(2*\text{Sqrt}[2 - \text{Sqrt}[3]]*(s + r*x)*\text{Sqrt}[(s^2 - r*s*x + r^2*x^2)/((1 - \text{Sqrt}[3])*s + r*x)^2]*\text{EllipticF}[\text{ArcSin}[(1 + \text{Sqrt}[3])*s + r*x]/((1 - \text{Sqrt}[3])*s + r*x)], -7 + 4*\text{Sqrt}[3])]/(3^(1/4)*r*\text{Sqrt}[a + b*x^3]*\text{Sqrt}[-((s*(s + r*x))/((1 - \text{Sqrt}[3])*s + r*x)^2)]), x]] \text{ /; FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 235

$\text{Int}[(a_) + (b_)*(x_)^2)^{-1/3}, x_Symbol] \text{ :> Dist}[(3*\text{Sqrt}[b*x^2])/(2*b*x), \text{Subst}[\text{Int}[x/\text{Sqrt}[-a + x^3], x], x, (a + b*x^2)^{1/3}], x] \text{ /; FreeQ}[\{a, b\}, x]$

Rule 304

$\text{Int}[(x_)/\text{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] \text{ :> With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\text{Sqrt}[2]*s)/(\text{Sqrt}[2 - \text{Sqrt}[3]]*r), \text{Int}[1/\text{Sqrt}[a + b*x^3], x], x] + \text{Dist}[1/r, \text{Int}[(1 + \text{Sqrt}[3])*s + r*x]/\text{Sqrt}[a + b*x^3], x], x]] \text{ /; FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 388

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}]^{(p_)*((c_) + (d_)*(x_)^{(n_)}), x_Symbol] \text{ :> Simp}[(d*x*(a + b*x^n)^{(p+1)})/(b*(n*(p+1) + 1)), x] - \text{Dist}[(a*d - b*c*(n*(p+1) + 1))/(b*(n*(p+1) + 1)), \text{Int}[(a + b*x^n)^p, x], x] \text{ /; FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[n*(p+1) + 1, 0]$

Rule 416

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}]^{(p_)*((c_) + (d_)*(x_)^{(n_)})^{(q_)}], x_Symbol] \text{ :> Simp}[(d*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-1)})/(b*(n*(p+q) + 1)), x] + \text{Dist}[1/(b*(n*(p+q) + 1)), \text{Int}[(a + b*x^n)^p*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(b*c*(n*(p+q) + 1) - a*d) + d*(b*c*(n*(p+2*q-1) + 1) - a*d*(n*(q-1) + 1))*x^n, x], x], x] \text{ /; FreeQ}[\{a, b, c, d, n, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[q, 1] \&\& \text{NeQ}[n*(p+q) + 1, 0] \&\& !\text{IGtQ}[p, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 528

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}]^{(p_)*((c_) + (d_)*(x_)^{(n_)})^{(q_)}*(e_) + (f_)*(x_)^{(n_)}], x_Symbol] \text{ :> Simp}[(f*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^q)/(b*(n*(p+q) + 1)), x] + \text{Dist}[1/(b*(n*(p+q) + 1)), \text{Int}[(a + b*x^n)^p*(c + d*x^n)^{(q-1)}*\text{Simp}[c*(b*e - a*f + b*e*n*(p+q) + 1) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p+q) + 1))*x^n, x], x], x] \text{ /; FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \&\& \text{GtQ}[q, 0] \&\& \text{NeQ}[n*(p+q) + 1, 0]$

Rule 1879

$\text{Int}[(c_) + (d_)*(x_)]/\text{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] \text{ :> With}[\{r = \text{Numer}[\text{Simplify}[(1 + \text{Sqrt}[3])*d]/c], s = \text{Denom}[\text{Simplify}[(1 + \text{Sqrt}[3])*d]/c]]\}, \text{Simp}[(2*d*s^3*\text{Sqrt}[a + b*x^3])/(a*r^2*((1 - \text{Sqrt}[3])*s + r*x)), x] + \text{Simp}[(3^(1/4)*\text{Sqrt}[2 + \text{Sqrt}[3]]*d*s*(s + r*x)*\text{Sqrt}[(s^2 - r*s*x + r^2*x^2)/((1 - \text{Sqrt}[3])*s + r*x)^2]*\text{EllipticE}[\text{ArcSin}[(1 + \text{Sqrt}[3])*s + r*x]/((1 - \text{Sqrt}[3])*s + r*x)]), x]] \text{ /; FreeQ}[\{a, b, c, d\}, x] \&\& \text{GtQ}[d, 0] \&\& \text{NeQ}[a, 0] \&\& \text{NeQ}[b, 0] \&\& \text{NeQ}[c, 0] \&\& \text{NeQ}[d, 0]$


```
rt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2))), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned} \int (a - bx^2)^{5/3} (3a + bx^2)^3 dx &= -\frac{3}{31}x(a - bx^2)^{8/3} (3a + bx^2)^2 - \frac{3 \int (a - bx^2)^{5/3} (3a + bx^2) (-96a^2b - 48ab^2) dx}{31b} \\ &= -\frac{432}{775}ax(a - bx^2)^{8/3} (3a + bx^2) - \frac{3}{31}x(a - bx^2)^{8/3} (3a + bx^2)^2 + \frac{9 \int (a - bx^2)^{5/3} (3a + bx^2)^2 dx}{31} \\ &= -\frac{33264a^2x(a - bx^2)^{8/3}}{14725} - \frac{432}{775}ax(a - bx^2)^{8/3} (3a + bx^2) - \frac{3}{31}x(a - bx^2)^{8/3} (3a + bx^2)^2 \\ &= \frac{1404864a^3x(a - bx^2)^{5/3}}{191425} - \frac{33264a^2x(a - bx^2)^{8/3}}{14725} - \frac{432}{775}ax(a - bx^2)^{8/3} (3a + bx^2) \\ &= \frac{2809728a^4x(a - bx^2)^{2/3}}{267995} + \frac{1404864a^3x(a - bx^2)^{5/3}}{191425} - \frac{33264a^2x(a - bx^2)^{8/3}}{14725} \\ &= \frac{2809728a^4x(a - bx^2)^{2/3}}{267995} + \frac{1404864a^3x(a - bx^2)^{5/3}}{191425} - \frac{33264a^2x(a - bx^2)^{8/3}}{14725} \\ &= \frac{2809728a^4x(a - bx^2)^{2/3}}{267995} + \frac{1404864a^3x(a - bx^2)^{5/3}}{191425} - \frac{33264a^2x(a - bx^2)^{8/3}}{14725} \\ &= \frac{2809728a^4x(a - bx^2)^{2/3}}{267995} + \frac{1404864a^3x(a - bx^2)^{5/3}}{191425} - \frac{33264a^2x(a - bx^2)^{8/3}}{14725} \end{aligned}$$

Mathematica [C] time = 5.05, size = 110, normalized size = 0.16

$$\frac{3 \left(6243840a^5x^3 \sqrt{1 - \frac{bx^2}{a}} {}_2F_1 \left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a} \right) + 5815935a^5x - 5312355a^4bx^3 - 1675114a^3b^2x^5 + 749658a^2b^3x^7 + 378651ab^4x^9 + 43225b^5x^{11} + 6243840a^5x \left(1 - \frac{bx^2}{a} \right)^{1/3} \text{Hypergeometric2F1} \left[\frac{1}{3}, \frac{1}{2}, \frac{3}{2}, \frac{bx^2}{a} \right] \right)}{1339975 \sqrt[3]{a - bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a - b*x^2)^(5/3)*(3*a + b*x^2)^3,x]

[Out] (3*(5815935*a^5*x - 5312355*a^4*b*x^3 - 1675114*a^3*b^2*x^5 + 749658*a^2*b^3*x^7 + 378651*a*b^4*x^9 + 43225*b^5*x^11 + 6243840*a^5*x*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a]))/(1339975*(a - b*x^2)^(1/3))

fricas [F] time = 1.07, size = 0, normalized size = 0.00

$$\text{integral} \left(-(b^4x^8 + 8ab^3x^6 + 18a^2b^2x^4 - 27a^4)(-bx^2 + a)^{\frac{2}{3}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a)^3,x, algorithm="fricas")

[Out] integral(-(b^4*x^8 + 8*a*b^3*x^6 + 18*a^2*b^2*x^4 - 27*a^4)*(-b*x^2 + a)^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)^3 (-bx^2 + a)^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a)^3,x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^3*(-b*x^2 + a)^(5/3), x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int (-bx^2 + a)^{\frac{5}{3}} (bx^2 + 3a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(5/3)*(b*x^2+3*a)^3,x)

[Out] int((-b*x^2+a)^(5/3)*(b*x^2+3*a)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)^3 (-bx^2 + a)^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a)^3,x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^3*(-b*x^2 + a)^(5/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (a - bx^2)^{\frac{5}{3}} (bx^2 + 3a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(5/3)*(3*a + b*x^2)^3,x)

[Out] int((a - b*x^2)^(5/3)*(3*a + b*x^2)^3, x)

sympy [A] time = 5.59, size = 139, normalized size = 0.21

$$27a^{\frac{14}{3}} x {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{1}{2} \\ \frac{3}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) - \frac{18a^{\frac{8}{3}} b^2 x^5 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{5}{2} \\ \frac{7}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{5} - \frac{8a^{\frac{5}{3}} b^3 x^7 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{7}{2} \\ \frac{9}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{7} - \frac{a^{\frac{2}{3}} b^4 x^9 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{9}{2} \\ \frac{11}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(5/3)*(b*x**2+3*a)**3,x)

[Out] 27*a**(14/3)*x*hyper((-2/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) - 18*a**(8/3)*b**2*x**5*hyper((-2/3, 5/2), (7/2,), b*x**2*exp_polar(2*I*pi)/a)/5 - 8*a**(5/3)*b**3*x**7*hyper((-2/3, 7/2), (9/2,), b*x**2*exp_polar(2*I*pi)/a)/7 - a**(2/3)*b**4*x**9*hyper((-2/3, 9/2), (11/2,), b*x**2*exp_polar(2*I*pi)/a)/9

$$3.117 \quad \int (a - bx^2)^{5/3} (3a + bx^2)^2 dx$$

Optimal. Leaf size=637

$$\frac{38016\sqrt{2}3^{3/4}a^{13/3} \left(\sqrt[3]{a} - \sqrt[3]{a - bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a - bx^2} + (a - bx^2)^{2/3}}{\left((1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2} \right)^2}} \operatorname{EllipticF} \left(\sin^{-1} \left(\frac{(1 + \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2}}{(1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2}} \right), 4\sqrt{3} - 7 \right)}{8645bx \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a - bx^2} \right)}{\left((1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2} \right)^2}}$$

[Out] 28512/8645*a^3*x*(-b*x^2+a)^(2/3)+14256/6175*a^2*x*(-b*x^2+a)^(5/3)-306/475*a*x*(-b*x^2+a)^(8/3)-3/25*x*(-b*x^2+a)^(8/3)*(b*x^2+3*a)-114048/8645*a^4*x/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))+38016/8645*3^(3/4)*a^(13/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*2^(1/2)*((a^(2/3)+a^(1/3))*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)-57024/8645*3^(1/4)*a^(13/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3))*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)

Rubi [A] time = 0.48, antiderivative size = 637, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {416, 388, 195, 235, 304, 219, 1879}

$$\frac{114048a^4x}{8645 \left((1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2} \right)} + \frac{28512a^3x(a - bx^2)^{2/3}}{8645} + \frac{14256a^2x(a - bx^2)^{5/3}}{6175} + \frac{38016\sqrt{2}3^{3/4}a^{13/3} \left(\sqrt[3]{a} - \sqrt[3]{a - bx^2} \right)}{8645 \left((1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2} \right)}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(5/3)*(3*a + b*x^2)^2,x]

[Out] (28512*a^3*x*(a - b*x^2)^(2/3))/8645 + (14256*a^2*x*(a - b*x^2)^(5/3))/6175 - (306*a*x*(a - b*x^2)^(8/3))/475 - (3*x*(a - b*x^2)^(8/3)*(3*a + b*x^2))/25 - (114048*a^4*x)/(8645*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) - (57024*3^(1/4)*Sqrt[2 + Sqrt[3]]*a^(13/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))]^2)*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(8645*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)))^2])) + (38016*Sqrt[2]*3^(3/4)*a^(13/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))]^2)*EllipticF[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(8645*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)))^2]))

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] &&

IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],
Denominator[p]])

Rule 219

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b}, x
&& NegQ[a]

Rule 235

Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x),
Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b},
x]

Rule 304

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[(d*x*(a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)),
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x]] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a,
b, c, d, n, p, q, x]

Rule 1879

Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/(
(1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned}
\int (a - bx^2)^{5/3} (3a + bx^2)^2 dx &= -\frac{3}{25}x(a - bx^2)^{8/3} (3a + bx^2) - \frac{3 \int (a - bx^2)^{5/3} (-78a^2b - 34ab^2x^2) dx}{25b} \\
&= -\frac{306}{475}ax(a - bx^2)^{8/3} - \frac{3}{25}x(a - bx^2)^{8/3} (3a + bx^2) + \frac{1}{475} (4752a^2) \int (a - bx^2)^{5/3} dx \\
&= \frac{14256a^2x(a - bx^2)^{5/3}}{6175} - \frac{306}{475}ax(a - bx^2)^{8/3} - \frac{3}{25}x(a - bx^2)^{8/3} (3a + bx^2) + \dots \\
&= \frac{28512a^3x(a - bx^2)^{2/3}}{8645} + \frac{14256a^2x(a - bx^2)^{5/3}}{6175} - \frac{306}{475}ax(a - bx^2)^{8/3} - \frac{3}{25}x(a - bx^2)^{8/3} (3a + bx^2) \\
&= \frac{28512a^3x(a - bx^2)^{2/3}}{8645} + \frac{14256a^2x(a - bx^2)^{5/3}}{6175} - \frac{306}{475}ax(a - bx^2)^{8/3} - \frac{3}{25}x(a - bx^2)^{8/3} (3a + bx^2) \\
&= \frac{28512a^3x(a - bx^2)^{2/3}}{8645} + \frac{14256a^2x(a - bx^2)^{5/3}}{6175} - \frac{306}{475}ax(a - bx^2)^{8/3} - \frac{3}{25}x(a - bx^2)^{8/3} (3a + bx^2) \\
&= \frac{28512a^3x(a - bx^2)^{2/3}}{8645} + \frac{14256a^2x(a - bx^2)^{5/3}}{6175} - \frac{306}{475}ax(a - bx^2)^{8/3} - \frac{3}{25}x(a - bx^2)^{8/3} (3a + bx^2)
\end{aligned}$$

Mathematica [C] time = 3.01, size = 173, normalized size = 0.27

$$\frac{x(a - bx^2)^{2/3} \left(4b \operatorname{Gamma}\left(-\frac{2}{3}\right) (3ax + bx^3)^2 {}_3F_2\left(-\frac{2}{3}, \frac{3}{2}, 2; 1, \frac{9}{2}; \frac{bx^2}{a}\right) + 8bx^2 \operatorname{Gamma}\left(-\frac{2}{3}\right) (18a^2 + 9abx^2 + b^2x^4) \right)}{105 \operatorname{Gamma}\left(-\frac{5}{3}\right) \left(1 - \frac{bx^2}{a}\right)^{2/3}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a - b*x^2)^(5/3)*(3*a + b*x^2)^2,x]

[Out] (x*(a - b*x^2)^(2/3)*(21*a*(45*a^2 + 10*a*b*x^2 + b^2*x^4)*Gamma[-5/3]*Hypergeometric2F1[-5/3, 1/2, 7/2, (b*x^2)/a] + 8*b*x^2*(18*a^2 + 9*a*b*x^2 + b^2*x^4)*Gamma[-2/3]*Hypergeometric2F1[-2/3, 3/2, 9/2, (b*x^2)/a] + 4*b*(3*a*x + b*x^3)^2*Gamma[-2/3]*HypergeometricPFQ[{-2/3, 3/2, 2}, {1, 9/2}, (b*x^2)/a])/(105*(1 - (b*x^2)/a)^(2/3)*Gamma[-5/3])

fricas [F] time = 0.95, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\left(b^3x^6 + 5ab^2x^4 + 3a^2bx^2 - 9a^3\right)\left(-bx^2 + a\right)^{\frac{2}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a)^2,x, algorithm="fricas")

[Out] integral(-(b^3*x^6 + 5*a*b^2*x^4 + 3*a^2*b*x^2 - 9*a^3)*(-b*x^2 + a)^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)^2 (-bx^2 + a)^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a)^2,x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^2*(-b*x^2 + a)^(5/3), x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int (-bx^2 + a)^{\frac{5}{3}} (bx^2 + 3a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(5/3)*(b*x^2+3*a)^2,x)

[Out] int((-b*x^2+a)^(5/3)*(b*x^2+3*a)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)^2 (-bx^2 + a)^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^2*(-b*x^2 + a)^(5/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (a - bx^2)^{\frac{5}{3}} (bx^2 + 3a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(5/3)*(3*a + b*x^2)^2,x)

[Out] int((a - b*x^2)^(5/3)*(3*a + b*x^2)^2, x)

sympy [A] time = 4.90, size = 131, normalized size = 0.21

$$9a^{\frac{11}{3}} x {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{1}{2} \\ \frac{3}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) - a^{\frac{8}{3}} bx^3 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{3}{2} \\ \frac{5}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) - a^{\frac{5}{3}} b^2 x^5 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{5}{2} \\ \frac{7}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) - \frac{a^{\frac{2}{3}} b^3 x^7 {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{7}{2} \\ \frac{9}{2} \end{matrix} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(5/3)*(b*x**2+3*a)**2,x)

[Out] 9*a**(11/3)*x*hyper((-2/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) - a**(8/3)*b*x**3*hyper((-2/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a) - a**(5/3)*b**2*x**5*hyper((-2/3, 5/2), (7/2,), b*x**2*exp_polar(2*I*pi)/a) - a**(2/3)*b**3*x**7*hyper((-2/3, 7/2), (9/2,), b*x**2*exp_polar(2*I*pi)/a)/7

3.118 $\int (a - bx^2)^{5/3} (3a + bx^2) dx$

Optimal. Leaf size=608

$$\frac{2400\sqrt{2}3^{3/4}a^{10/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{1729bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}$$

[Out] $1800/1729*a^2*x*(-b*x^2+a)^{(2/3)}+180/247*a*x*(-b*x^2+a)^{(5/3)}-3/19*x*(-b*x^2+a)^{(8/3)}-7200/1729*a^3*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+2400/1729*3^{(3/4)}*a^{(10/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*EllipticF((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})^2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2^{(1/2)}-3600/1729*3^{(1/4)}*a^{(10/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*EllipticE((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2^{(1/2)}$

Rubi [A] time = 0.42, antiderivative size = 608, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {388, 195, 235, 304, 219, 1879}

$$\frac{7200a^3x}{1729\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}+\frac{1800a^2x(a-bx^2)^{2/3}}{1729}+\frac{2400\sqrt{2}3^{3/4}a^{10/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}{1729bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(5/3)*(3*a + b*x^2), x]

[Out] $(1800*a^2*x*(a-b*x^2)^{(2/3)})/1729+(180*a*x*(a-b*x^2)^{(5/3)})/247-(3*x*(a-b*x^2)^{(8/3)})/19-(7200*a^3*x)/(1729*((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))-(3600*3^{(1/4)}*\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*a^{(10/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})],-7+4*\operatorname{Sqrt}[3]])/(1729*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2])+(2400*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(10/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})],-7+4*\operatorname{Sqrt}[3]])/(1729*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2])$

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],

Denominator[p]])

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 388

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/
(1 - Sqrt[3])*s + r*x]^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int (a - bx^2)^{5/3} (3a + bx^2) dx &= -\frac{3}{19}x(a - bx^2)^{8/3} + \frac{1}{19}(60a) \int (a - bx^2)^{5/3} dx \\
&= \frac{180}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{8/3} + \frac{1}{247}(600a^2) \int (a - bx^2)^{2/3} dx \\
&= \frac{1800a^2x(a - bx^2)^{2/3}}{1729} + \frac{180}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{8/3} + \frac{(2400a^3) \int \sqrt[3]{a - bx^2}}{1729} \\
&= \frac{1800a^2x(a - bx^2)^{2/3}}{1729} + \frac{180}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{8/3} - \frac{(3600a^3\sqrt{-bx^2})}{1729} \\
&= \frac{1800a^2x(a - bx^2)^{2/3}}{1729} + \frac{180}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{8/3} + \frac{(3600a^3\sqrt{-bx^2})}{1729} \\
&= \frac{1800a^2x(a - bx^2)^{2/3}}{1729} + \frac{180}{247}ax(a - bx^2)^{5/3} - \frac{3}{19}x(a - bx^2)^{8/3} - \frac{7}{1729} \left((1 - \sqrt{3}) \right)
\end{aligned}$$

Mathematica [C] time = 0.06, size = 68, normalized size = 0.11

$$\frac{3}{19}x(a - bx^2)^{2/3} \left(\frac{20a^2 {}_2F_1\left(-\frac{5}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a}\right)}{\left(1 - \frac{bx^2}{a}\right)^{2/3}} - (a - bx^2)^2 \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a - b*x^2)^(5/3)*(3*a + b*x^2), x]

[Out] (3*x*(a - b*x^2)^(2/3)*(-(a - b*x^2)^2 + (20*a^2*Hypergeometric2F1[-5/3, 1/2, 3/2, (b*x^2)/a])/(1 - (b*x^2)/a)^(2/3)))/19

fricas [F] time = 0.94, size = 0, normalized size = 0.00

$$\text{integral}\left(-\left(b^2x^4 + 2abx^2 - 3a^2\right)\left(-bx^2 + a\right)^{\frac{2}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a), x, algorithm="fricas")

[Out] integral(-(b^2*x^4 + 2*a*b*x^2 - 3*a^2)*(-b*x^2 + a)^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)(-bx^2 + a)^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a), x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)*(-b*x^2 + a)^(5/3), x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int (-bx^2 + a)^{\frac{5}{3}} (bx^2 + 3a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-b*x^2+a)^(5/3)*(b*x^2+3*a),x)`

[Out] `int((-b*x^2+a)^(5/3)*(b*x^2+3*a),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + 3a)(-bx^2 + a)^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x^2+a)^(5/3)*(b*x^2+3*a),x, algorithm="maxima")`

[Out] `integrate((b*x^2 + 3*a)*(-b*x^2 + a)^(5/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (a - bx^2)^{\frac{5}{3}} (bx^2 + 3a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a - b*x^2)^(5/3)*(3*a + b*x^2),x)`

[Out] `int((a - b*x^2)^(5/3)*(3*a + b*x^2), x)`

sympy [A] time = 3.69, size = 100, normalized size = 0.16

$$3a^{\frac{8}{3}}x^2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{1}{2} \\ \frac{3}{2} \end{matrix} \middle| \frac{bx^2e^{2i\pi}}{a}\right) - \frac{2a^{\frac{5}{3}}bx^3{}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{3}{2} \\ \frac{5}{2} \end{matrix} \middle| \frac{bx^2e^{2i\pi}}{a}\right)}{3} - \frac{a^{\frac{2}{3}}b^2x^5{}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{5}{2} \\ \frac{7}{2} \end{matrix} \middle| \frac{bx^2e^{2i\pi}}{a}\right)}{5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x**2+a)**(5/3)*(b*x**2+3*a),x)`

[Out] `3*a**(8/3)*x*hyper((-2/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) - 2*a**(5/3)*b*x**3*hyper((-2/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a)/3 - a**(2/3)*b**2*x**5*hyper((-2/3, 5/2), (7/2,), b*x**2*exp_polar(2*I*pi)/a)/5`

$$3.119 \quad \int \frac{(a-bx^2)^{5/3}}{3a+bx^2} dx$$

Optimal. Leaf size=765

$$\frac{32\sqrt{2}3^{3/4}a^{4/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{7bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] $-3/7*x*(-b*x^2+a)^{(2/3)}+96/7*a*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))+4*2^{(1/3)}*a^{(7/6)}*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/(a^{(1/3)}+2^{(1/3)}*(-b*x^2+a)^{(1/3)})/b^{(1/2)}-4/3*2^{(1/3)}*a^{(7/6)}*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/2)})/b^{(1/2)}+4/3*2^{(1/3)}*a^{(7/6)}*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(-b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})/b^{(1/2)}+4/3*2^{(1/3)}*a^{(7/6)}*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})/b^{(1/2)}-32/7*3^{(3/4)}*a^{(4/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))),2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}+48/7*3^{(1/4)}*a^{(4/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}$

Rubi [A] time = 0.44, antiderivative size = 765, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {416, 530, 235, 304, 219, 1879, 393}

$$\frac{4\sqrt[3]{2}a^{7/6}\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}}+\frac{4\sqrt[3]{2}a^{7/6}\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt[6]{a}\left(\sqrt[3]{2}\sqrt[3]{a-bx^2}+\sqrt[3]{a}\right)}\right)}{\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(5/3)/(3*a + b*x^2), x]

[Out] $(-3*x*(a-b*x^2)^{(2/3)})/7+(96*a*x)/(7*((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))+((4*2^{(1/3)}*a^{(7/6)}*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[b])+(4*2^{(1/3)}*a^{(7/6)}*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(a-b*x^2)^{(1/3)})]/(\operatorname{Sqrt}[b]*x)]/(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[b])-(4*2^{(1/3)}*a^{(7/6)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(3*\operatorname{Sqrt}[b])+(4*2^{(1/3)}*a^{(7/6)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{(1/6)}*(a^{(1/3)}+2^{(1/3)}*(a-b*x^2)^{(1/3)})])/(\operatorname{Sqrt}[b]+(48*3^{(1/4)}*\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*a^{(4/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})],-7+4*\operatorname{Sqrt}[3]])/(7*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2)]-(32*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(4/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})],-7+4*\operatorname{Sqrt}[3]])/(7*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2)]))$

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 393

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Wit
h[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^
(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2
)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^
(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)
))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]]] /; FreeQ[{a, b, c, d}
, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]
```

Rule 416

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)),
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a
, b, c, d, n, p, q, x]
```

Rule 530

```
Int[(((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p
, n}, x]
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/
(1 - Sqrt[3])*s + r*x]^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a - bx^2)^{5/3}}{3a + bx^2} dx &= -\frac{3}{7}x(a - bx^2)^{2/3} + \frac{3 \int \frac{\frac{16a^2b}{3} - \frac{32}{3}ab^2x^2}{\sqrt[3]{a - bx^2}(3a + bx^2)} dx}{7b} \\
&= -\frac{3}{7}x(a - bx^2)^{2/3} - \frac{1}{7}(32a) \int \frac{1}{\sqrt[3]{a - bx^2}} dx + (16a^2) \int \frac{1}{\sqrt[3]{a - bx^2}(3a + bx^2)} dx \\
&= -\frac{3}{7}x(a - bx^2)^{2/3} + \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}} + \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}} - \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}} \\
&= -\frac{3}{7}x(a - bx^2)^{2/3} + \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}} + \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}} - \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}} \\
&= -\frac{3}{7}x(a - bx^2)^{2/3} + \frac{96ax}{7\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} + \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}} + \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}} - \frac{4\sqrt[3]{2} a^{7/6} \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{bx}}\right)}{\sqrt{3}\sqrt{b}}
\end{aligned}$$

Mathematica [C] time = 0.15, size = 231, normalized size = 0.30

$$x \left(27 \left(\frac{48a^3 F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{(3a + bx^2) \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - a + bx^2 \right) - 32bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) \right) / (63 \sqrt[3]{a - bx^2})$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a - b*x^2)^(5/3)/(3*a + b*x^2), x]

[Out] (x*(-32*b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 27*(-a + b*x^2 + (48*a^3*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/((3*a + b*x^2)*(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))))/((63*(a - b*x^2)^(1/3)))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a), x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{5/3}}{bx^2 + 3a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a),x, algorithm="giac")

[Out] integrate((-b*x^2 + a)^(5/3)/(b*x^2 + 3*a), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{5}{3}}}{bx^2 + 3a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(5/3)/(b*x^2+3*a),x)

[Out] int((-b*x^2+a)^(5/3)/(b*x^2+3*a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{5}{3}}}{bx^2 + 3a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a),x, algorithm="maxima")

[Out] integrate((-b*x^2 + a)^(5/3)/(b*x^2 + 3*a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a - bx^2)^{\frac{5}{3}}}{bx^2 + 3a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(5/3)/(3*a + b*x^2),x)

[Out] int((a - b*x^2)^(5/3)/(3*a + b*x^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a - bx^2)^{\frac{5}{3}}}{3a + bx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(5/3)/(b*x**2+3*a),x)

[Out] Integral((a - b*x**2)**(5/3)/(3*a + b*x**2), x)

$$3.120 \quad \int \frac{(a-bx^2)^{5/3}}{(3a+bx^2)^2} dx$$

Optimal. Leaf size=775

$$\frac{11\sqrt{2} \sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} \operatorname{EllipticF} \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right), 4\sqrt{3} - 7 \right) 11\sqrt{2}}{3\sqrt[4]{3} bx \sqrt{-\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}$$

[Out] $2/3*x*(-b*x^2+a)^{(2/3)}/(b*x^2+3*a)-11/3*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))-2^{(1/3)}*a^{(1/6)}*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/(a^{(1/3)}+2^{(1/3)}*(-b*x^2+a)^{(1/3)})/b^{(1/2)}+1/3*2^{(1/3)}*a^{(1/6)}*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/2)})/b^{(1/2)}-1/3*2^{(1/3)}*a^{(1/6)}*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(-b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})/b^{(1/2)}-1/3*2^{(1/3)}*a^{(1/6)}*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})/b^{(1/2)}+11/9*3^{(3/4)}*a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}-11/6*3^{(1/4)}*a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}$

Rubi [A] time = 0.44, antiderivative size = 775, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {413, 530, 235, 304, 219, 1879, 393}

$$\frac{11\sqrt{2} \sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} F \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right) \middle| -7 + 4\sqrt{3} \right) 11\sqrt{2 + \sqrt{3}}}{3\sqrt[4]{3} bx \sqrt{-\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(5/3)/(3*a + b*x^2)^2, x]

[Out] $(2*x*(a - b*x^2)^{(2/3)})/(3*(3*a + b*x^2)) - (11*x)/(3*((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})) - (2^{(1/3)}*a^{(1/6)}*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[b]) - (2^{(1/3)}*a^{(1/6)}*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)} - 2^{(1/3)}*(a - b*x^2)^{(1/3)}))/(\operatorname{Sqrt}[b]*x)]/(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[b]) + (2^{(1/3)}*a^{(1/6)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[b]) - (2^{(1/3)}*a^{(1/6)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{(1/6)}*(a^{(1/3)} + 2^{(1/3)}*(a - b*x^2)^{(1/3)}))]/\operatorname{Sqrt}[b] - (11*\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3])*a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}]/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})], -7 + 4*\operatorname{Sqrt}[3])]/(2*3^{(3/4)}*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])) + (11*\operatorname{Sqrt}[2]*a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}]/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})], -7$

+ 4*Sqrt[3]]/(3*3^(1/4)*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/(1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2])

Rule 219

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 235

Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

Rule 304

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sqrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 393

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3))])/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x]] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 530

Int[(((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1879

Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&

EqQ[b*c^3 - 2*(5 + 3*sqrt(3))*a*d^3, 0]

Rubi steps

$$\begin{aligned}
 \int \frac{(a - bx^2)^{5/3}}{(3a + bx^2)^2} dx &= \frac{2x(a - bx^2)^{2/3}}{3(3a + bx^2)} + \frac{\int \frac{-2a^2b + \frac{22}{3}ab^2x^2}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx}{6ab} \\
 &= \frac{2x(a - bx^2)^{2/3}}{3(3a + bx^2)} + \frac{11}{9} \int \frac{1}{\sqrt[3]{a - bx^2}} dx - (4a) \int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx \\
 &= \frac{2x(a - bx^2)^{2/3}}{3(3a + bx^2)} - \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} - \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} + \frac{\sqrt[3]{2} \sqrt[6]{a}}{\sqrt{3} \sqrt{b}} \\
 &= \frac{2x(a - bx^2)^{2/3}}{3(3a + bx^2)} - \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} - \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} + \frac{\sqrt[3]{2} \sqrt[6]{a}}{\sqrt{3} \sqrt{b}} \\
 &= \frac{2x(a - bx^2)^{2/3}}{3(3a + bx^2)} - \frac{11x}{3\left((1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} - \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}} - \frac{\sqrt[3]{2} \sqrt[6]{a} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{\sqrt{3} \sqrt{b}}
 \end{aligned}$$

Mathematica [C] time = 0.15, size = 235, normalized size = 0.30

$$x \left(\frac{27 \left(\frac{9a^2 F_1\left(\frac{1}{2}, \frac{1}{3}, 1; \frac{3}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{2bx^2 \left(F_1\left(\frac{3}{2}, \frac{4}{3}, 1; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}, \frac{1}{3}, 2; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}, \frac{1}{3}, 1; \frac{3}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) + 2a - 2bx^2 \right)}{3a + bx^2} + \frac{11bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} F_1\left(\frac{3}{2}, \frac{1}{3}, 1; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{a} \right)$$

$$\frac{\hspace{10em}}{81 \sqrt[3]{a - bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a - b*x^2)^(5/3)/(3*a + b*x^2)^2,x]

[Out] (x*((11*b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])/a + (27*(2*a - 2*b*x^2 - (9*a^2*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])))/(3*a + b*x^2)))/(81*(a - b*x^2)^(1/3))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{5}{3}}}{(bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a)^2,x, algorithm="giac")

[Out] integrate((-b*x^2 + a)^(5/3)/(b*x^2 + 3*a)^2, x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{5}{3}}}{(bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(5/3)/(b*x^2+3*a)^2,x)

[Out] int((-b*x^2+a)^(5/3)/(b*x^2+3*a)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{5}{3}}}{(bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a)^2,x, algorithm="maxima")

[Out] integrate((-b*x^2 + a)^(5/3)/(b*x^2 + 3*a)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a - bx^2)^{\frac{5}{3}}}{(bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(5/3)/(3*a + b*x^2)^2,x)

[Out] int((a - b*x^2)^(5/3)/(3*a + b*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a - bx^2)^{\frac{5}{3}}}{(3a + bx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(5/3)/(b*x**2+3*a)**2,x)

[Out] Integral((a - b*x**2)**(5/3)/(3*a + b*x**2)**2, x)

$$3.121 \quad \int \frac{(a-bx^2)^{5/3}}{(3a+bx^2)^3} dx$$

Optimal. Leaf size=815

$$\frac{(a-bx^2)^{2/3} x}{18a(bx^2+3a)} + \frac{x}{18a\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{(a-bx^2)^{2/3} x}{3(bx^2+3a)^2} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a}\right)}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}}$$

[Out] $\frac{1}{3}x(-bx^2+a)^{2/3}/(bx^2+3a)^2 - \frac{1}{18}x(-bx^2+a)^{2/3}/a/(bx^2+3a) + \frac{1}{18}x/a/(-(-bx^2+a)^{1/3}+a^{1/3}(1-3^{1/2})) + \frac{1}{36}\operatorname{arctanh}(x*b^{1/2}/a^{1/6})/(a^{1/3}+2^{1/3}*(-bx^2+a)^{1/3}) * 2^{1/3}/a^{5/6}/b^{1/2} - \frac{1}{108}\operatorname{arctanh}(x*b^{1/2}/a^{1/2}) * 2^{1/3}/a^{5/6}/b^{1/2} + \frac{1}{108}\operatorname{arctan}(a^{1/6}*(a^{1/3}-2^{1/3}*(-bx^2+a)^{1/3})*3^{1/2}/x/b^{1/2}) * 2^{1/3}/a^{5/6} * 3^{1/2}/b^{1/2} + \frac{1}{108}\operatorname{arctan}(3^{1/2}*a^{1/2}/x/b^{1/2}) * 2^{1/3}/a^{5/6} * 3^{1/2}/b^{1/2} - \frac{1}{54}(a^{1/3}-(-bx^2+a)^{1/3})*\operatorname{EllipticF}((-(-bx^2+a)^{1/3}+a^{1/3}*(1+3^{1/2}))/(-(-bx^2+a)^{1/3}+a^{1/3}*(1-3^{1/2})), 2*I-I*3^{1/2})*((a^{2/3}+a^{1/3}*(-bx^2+a)^{1/3}+(-bx^2+a)^{2/3})/(-(-bx^2+a)^{1/3}+a^{1/3}*(1-3^{1/2})))^2)^{1/2} * 3^{3/4}/a^{2/3}/b/x * 2^{1/2}/(-a^{1/3}*(a^{1/3}-(-bx^2+a)^{1/3}))/(-(-bx^2+a)^{1/3}+a^{1/3}*(1-3^{1/2})))^2)^{1/2} + \frac{1}{36}(a^{1/3}-(-bx^2+a)^{1/3})*\operatorname{EllipticE}((-(-bx^2+a)^{1/3}+a^{1/3}*(1+3^{1/2}))/(-(-bx^2+a)^{1/3}+a^{1/3}*(1-3^{1/2}))), 2*I-I*3^{1/2})*((a^{2/3}+a^{1/3}*(-bx^2+a)^{1/3}+(-bx^2+a)^{2/3})/(-(-bx^2+a)^{1/3}+a^{1/3}*(1-3^{1/2})))^2)^{1/2} * (1/2 * 6^{1/2} + 1/2 * 2^{1/2}) * 3^{1/4}/a^{2/3}/b/x / (-a^{1/3}*(a^{1/3}-(-bx^2+a)^{1/3}))/(-(-bx^2+a)^{1/3}+a^{1/3}*(1-3^{1/2})))^2)^{1/2}$

Rubi [A] time = 0.61, antiderivative size = 815, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {413, 12, 471, 530, 235, 304, 219, 1879, 393}

$$\frac{(a-bx^2)^{2/3} x}{18a(bx^2+3a)} + \frac{x}{18a\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{(a-bx^2)^{2/3} x}{3(bx^2+3a)^2} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a}\right)}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[(a - b*x^2)^(5/3)/(3*a + b*x^2)^3, x]

[Out] $(x*(a-bx^2)^{2/3})/(3*(3a+bx^2)^2) - (x*(a-bx^2)^{2/3})/(18*a*(3a+bx^2)) + x/(18*a*((1-\operatorname{Sqrt}[3])*a^{1/3} - (a-bx^2)^{1/3})) + \operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(18*2^{2/3}*\operatorname{Sqrt}[3]*a^{5/6}*\operatorname{Sqrt}[b]) + \operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{1/6}*(a^{1/3}-2^{1/3}*(a-bx^2)^{1/3}))/(\operatorname{Sqrt}[b]*x)]/(18*2^{2/3}*\operatorname{Sqrt}[3]*a^{5/6}*\operatorname{Sqrt}[b]) - \operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(54*2^{2/3})*a^{5/6}*\operatorname{Sqrt}[b]) + \operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{1/6}*(a^{1/3}+2^{1/3}*(a-bx^2)^{1/3}))]/(18*2^{2/3})*a^{5/6}*\operatorname{Sqrt}[b]) + (\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*(a^{1/3} - (a-bx^2)^{1/3}))*\operatorname{Sqrt}[(a^{2/3}+a^{1/3}*(a-bx^2)^{1/3}+(a-bx^2)^{2/3})]/((1-\operatorname{Sqrt}[3])*a^{1/3} - (a-bx^2)^{1/3})^2)*\operatorname{EllipticE}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{1/3} - (a-bx^2)^{1/3}]/((1-\operatorname{Sqrt}[3])*a^{1/3} - (a-bx^2)^{1/3})], -7+4*\operatorname{Sqrt}[3]]/(12*3^{3/4})*a^{2/3}*b*x*\operatorname{Sqrt}[-((a^{1/3}*(a^{1/3} - (a-bx^2)^{1/3}))/((1-\operatorname{Sqrt}[3])*a^{1/3} - (a-bx^2)^{1/3}))^2]) - ((a^{1/3} - (a-bx^2)^{1/3}))*\operatorname{Sqrt}[(a^{2/3}+a^{1/3}*(a-bx^2)^{1/3}+(a-bx^2)^{2/3})]/((1-\operatorname{Sqrt}[3])*a^{1/3} - (a-bx^2)^{1/3})^2)*\operatorname{Elliptic}$

$$F[\text{ArcSin}[\frac{(1 + \sqrt{3})a^{1/3} - (a - bx^2)^{1/3}}{(1 - \sqrt{3})a^{1/3} - (a - bx^2)^{1/3}}], -7 + 4\sqrt{3}]/(9\sqrt{2} \cdot 3^{1/4} a^{2/3} b x \sqrt{t[-((a^{1/3})(a^{1/3}) - (a - bx^2)^{1/3})]/((1 - \sqrt{3})a^{1/3} - (a - bx^2)^{1/3})^2})]$$

Rule 12

$$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*)(v_)] /; \text{FreeQ}[b, x]$$

Rule 219

$$\text{Int}[1/\sqrt{(a_*) + (b_*)(x_)^3}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, \text{Simp}[(2\sqrt{2 - \sqrt{3}})(s + rx)\sqrt{(s^2 - r^2sx + r^2x^2)/((1 - \sqrt{3})s + rx)^2} \text{EllipticF}[\text{ArcSin}[\frac{(1 + \sqrt{3})s + rx}{(1 - \sqrt{3})s + rx}], -7 + 4\sqrt{3}]/(3^{1/4} r \sqrt{a + bx^3}) \sqrt{-(s(s + rx)/((1 - \sqrt{3})s + rx)^2)}], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a]$$

Rule 235

$$\text{Int}[(a_*) + (b_*)(x_)^2)^{-1/3}, x_Symbol] \rightarrow \text{Dist}[(3\sqrt{bx^2})/(2b^2x), \text{Subst}[\text{Int}[x/\sqrt{-a + x^3}, x], x, (a + bx^2)^{1/3}], x] /; \text{FreeQ}[\{a, b\}, x]$$

Rule 304

$$\text{Int}[(x_)/\sqrt{(a_*) + (b_*)(x_)^3}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\sqrt{2}s)/(\sqrt{2 - \sqrt{3}}r), \text{Int}[1/\sqrt{a + bx^3}, x], x] + \text{Dist}[1/r, \text{Int}[(1 + \sqrt{3})s + rx]/\sqrt{a + bx^3}, x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a]$$

Rule 393

$$\text{Int}[1/((a_*) + (b_*)(x_)^2)^{1/3}((c_*) + (d_*)(x_)^2)), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-(b/a), 2]\}, \text{Simp}[(q \text{ArcTan}[\sqrt{3}/(qx)])/(2 \cdot 2^{2/3} \sqrt{3} a^{1/3} d), x] + (\text{Simp}[(q \text{ArcTanh}[(a^{1/3}qx)/(a^{1/3} + 2^{1/3}(a + bx^2)^{1/3})])/(2 \cdot 2^{2/3} a^{1/3} d), x] - \text{Simp}[(q \text{ArcTanh}[qx])/(6 \cdot 2^{2/3} a^{1/3} d), x] + \text{Simp}[(q \text{ArcTan}[(\sqrt{3}(a^{1/3} - 2^{1/3}(a + bx^2)^{1/3})/(a^{1/3}qx)])/(2 \cdot 2^{2/3} \sqrt{3} a^{1/3} d), x])]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b^2c - a^2d, 0] \ \&\& \ \text{EqQ}[b^2c + 3a^2d, 0] \ \&\& \ \text{NegQ}[b/a]$$

Rule 413

$$\text{Int}[(a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)})^{(q_*)}, x_Symbol] \rightarrow \text{Simp}[(a^d - c^b)x^{n+1}(a + bx^n)^{p+1}(c + dx^n)^{q-1}/(a^b n (p+1)), x] - \text{Dist}[1/(a^b n (p+1)), \text{Int}[(a + bx^n)^{p+1}(c + dx^n)^{q-2} \text{Simp}[c(a^d - c^b(n(p+1) + 1)) + d(a^d(n(q-1) + 1) - b^2c(n(p+q) + 1))x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{NeQ}[b^2c - a^2d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$$

Rule 471

$$\text{Int}[(e_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)})^{(q_*)}, x_Symbol] \rightarrow \text{Simp}[(e^{n-1}(e^x)^{m-n+1}(a + bx^n)^{p+1}(c + dx^n)^{q+1})/(n(b^2c - a^2d)(p+1)), x] - \text{Dist}[e^n/(n(b^2c - a^2d)(p+1)), \text{Int}[(e^x)^{m-n}(a + bx^n)^{p+1}(c + dx^n)^q \text{Simp}[c(m-n+1) + d(m+n(p+q+1)+1)x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, q\}, x] \ \&\& \ \text{NeQ}[b^2c - a^2d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GeQ}[n, m-n+1]$$

1] && GtQ[m - n + 1, 0] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 530

Int[(((a_) + (b_)*(x_)^(n_))^(p_)*((e_) + (f_)*(x_)^(n_)))/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1879

Int[((c_) + (d_)*(x_))/Sqrt[(a_) + (b_)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned}
 \int \frac{(a - bx^2)^{5/3}}{(3a + bx^2)^3} dx &= \frac{x(a - bx^2)^{2/3}}{3(3a + bx^2)^2} + \frac{\int \frac{16ab^2x^2}{3\sqrt[3]{a-bx^2}(3a+bx^2)^2} dx}{12ab} \\
 &= \frac{x(a - bx^2)^{2/3}}{3(3a + bx^2)^2} + \frac{1}{9}(4b) \int \frac{x^2}{\sqrt[3]{a - bx^2} (3a + bx^2)^2} dx \\
 &= \frac{x(a - bx^2)^{2/3}}{3(3a + bx^2)^2} - \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)} + \frac{\int \frac{a - \frac{bx^2}{3}}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx}{18a} \\
 &= \frac{x(a - bx^2)^{2/3}}{3(3a + bx^2)^2} - \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)} + \frac{1}{9} \int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx - \frac{\int \frac{1}{\sqrt[3]{a-bx^2}} dx}{54a} \\
 &= \frac{x(a - bx^2)^{2/3}}{3(3a + bx^2)^2} - \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} - \frac{\int \frac{1}{\sqrt[3]{a-bx^2}} dx}{54a} \\
 &= \frac{x(a - bx^2)^{2/3}}{3(3a + bx^2)^2} - \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} - \frac{\int \frac{1}{\sqrt[3]{a-bx^2}} dx}{54a} \\
 &= \frac{x(a - bx^2)^{2/3}}{3(3a + bx^2)^2} - \frac{x(a - bx^2)^{2/3}}{18a(3a + bx^2)} + \frac{x}{18a\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{18 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\int \frac{1}{\sqrt[3]{a-bx^2}} dx}{54a}
 \end{aligned}$$

Mathematica [C] time = 0.24, size = 252, normalized size = 0.31

$$\frac{27x \left(\frac{9a(3a+bx^2)F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9aF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) + \frac{b^2x^4}{a} + 3a - 4bx^2}{(3a+bx^2)^2} - \frac{bx^3 \sqrt[3]{1-\frac{bx^2}{a}} F_1\left(\frac{3}{2}; \frac{1}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{a^2} \right)}{486\sqrt[3]{a-bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a - b*x^2)^(5/3)/(3*a + b*x^2)^3, x]

[Out] (-((b*x^3*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])/a^2) + (27*x*(3*a - 4*b*x^2 + (b^2*x^4)/a + (9*a*(3*a + b*x^2)*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])))/(3*a + b*x^2)^2)/(486*(a - b*x^2)^(1/3))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a)^3,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{5}{3}}}{(bx^2 + 3a)^{\frac{3}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a)^3,x, algorithm="giac")

[Out] integrate((-b*x^2 + a)^(5/3)/(b*x^2 + 3*a)^3, x)

maple [F] time = 0.36, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{5}{3}}}{(bx^2 + 3a)^{\frac{3}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(5/3)/(b*x^2+3*a)^3,x)

[Out] int((-b*x^2+a)^(5/3)/(b*x^2+3*a)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-bx^2 + a)^{\frac{5}{3}}}{(bx^2 + 3a)^{\frac{3}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(5/3)/(b*x^2+3*a)^3,x, algorithm="maxima")

[Out] integrate((-b*x^2 + a)^(5/3)/(b*x^2 + 3*a)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a - bx^2)^{5/3}}{(bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(5/3)/(3*a + b*x^2)^3, x)

[Out] int((a - b*x^2)^(5/3)/(3*a + b*x^2)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a - bx^2)^{5/3}}{(3a + bx^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(5/3)/(b*x**2+3*a)**3, x)

[Out] Integral((a - b*x**2)**(5/3)/(3*a + b*x**2)**3, x)

$$3.122 \quad \int \frac{(3a+bx^2)^4}{\sqrt[3]{a-bx^2}} dx$$

Optimal. Leaf size=659

$$\frac{1264896\sqrt{2}3^{3/4}a^{13/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{8645bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] $-1552608/43225*a^3*x*(-b*x^2+a)^{(2/3)}-36288/6175*a^2*x*(-b*x^2+a)^{(2/3)}*(b*x^2+3*a)-18/19*a*x*(-b*x^2+a)^{(2/3)}*(b*x^2+3*a)^2-3/25*x*(-b*x^2+a)^{(2/3)}*(b*x^2+3*a)^3-3794688/8645*a^4*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+1264896/8645*3^{(3/4)}*a^{(13/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}-1897344/8645*3^{(1/4)}*a^{(13/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*(a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}$

Rubi [A] time = 0.53, antiderivative size = 659, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {416, 528, 388, 235, 304, 219, 1879}

$$\frac{3794688a^4x}{8645\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}-\frac{1552608a^3x(a-bx^2)^{2/3}}{43225}-\frac{36288a^2x(a-bx^2)^{2/3}(3a+bx^2)}{6175}+\frac{1264896\sqrt{2}3^{3/4}}{8645}$$

Antiderivative was successfully verified.

[In] Int[(3*a + b*x^2)^4/(a - b*x^2)^(1/3),x]

[Out] $(-1552608*a^3*x*(a-b*x^2)^{(2/3)}/43225-(36288*a^2*x*(a-b*x^2)^{(2/3)}*(3*a+b*x^2))/6175-(18*a*x*(a-b*x^2)^{(2/3)}*(3*a+b*x^2)^2)/19-(3*x*(a-b*x^2)^{(2/3)}*(3*a+b*x^2)^3)/25-(3794688*a^4*x)/(8645*((1-\operatorname{Sqrt}[3])*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))-(1897344*3^{(1/4)}*\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*a^{(13/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})), -7+4*\operatorname{Sqrt}[3]])/(8645*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]))+(1264896*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(13/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})), -7+4*\operatorname{Sqrt}[3]])/(8645*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]))$

Rule 219

Int[1/Sqrt[(a_) + (b_)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s

$$\frac{x + r^2 x^2}{((1 - \sqrt{3})s + rx)^2} \text{EllipticF}[\text{ArcSin}[\frac{(1 + \sqrt{3})s + rx}{(1 - \sqrt{3})s + rx}], -7 + 4\sqrt{3}]/(3^{1/4} r \sqrt{a + bx^3}) \sqrt{-((s(s + rx))/((1 - \sqrt{3})s + rx)^2)}], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a]$$

Rule 235

$$\text{Int}[(a + (b \cdot x)^2)^{-1/3}, x_Symbol] \rightarrow \text{Dist}[(3\sqrt{b}x^2)/(2bx), \text{Subst}[\text{Int}[x/\sqrt{-a + x^3}], x], x, (a + bx^2)^{1/3}], x] /; \text{FreeQ}\{a, b\}, x]$$

Rule 304

$$\text{Int}[x/\sqrt{(a + (b \cdot x)^3)}, x_Symbol] \rightarrow \text{With}\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\sqrt{2}s)/(\sqrt{2 - \sqrt{3}}r), \text{Int}[1/\sqrt[3]{a + bx^3}], x], x] + \text{Dist}[1/r, \text{Int}[(1 + \sqrt{3})s + rx]/\sqrt{a + bx^3}], x], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a]$$

Rule 388

$$\text{Int}[(a + (b \cdot x)^n)^{p_1} ((c + (d \cdot x)^n)), x_Symbol] \rightarrow \text{Simp}[(d \cdot x (a + bx^n)^{p_1 + 1}) / (b(n(p_1 + 1) + 1)), x] - \text{Dist}[(a \cdot d - b \cdot c (n(p_1 + 1) + 1)) / (b(n(p_1 + 1) + 1)), \text{Int}[(a + bx^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{NeQ}[n(p_1 + 1) + 1, 0]$$

Rule 416

$$\text{Int}[(a + (b \cdot x)^n)^{p_1} ((c + (d \cdot x)^n)^{q_1}), x_Symbol] \rightarrow \text{Simp}[(d \cdot x (a + bx^n)^{p_1 + 1} (c + dx^n)^{q_1 - 1}) / (b(n(p_1 + q_1) + 1)), x] + \text{Dist}[1/(b(n(p_1 + q_1) + 1)), \text{Int}[(a + bx^n)^p (c + dx^n)^{q_1 - 2} \text{Simp}[c(b \cdot c(n(p_1 + q_1) + 1) - a \cdot d) + d(b \cdot c(n(p_1 + 2q_1 - 1) + 1) - a \cdot d(n(q_1 - 1) + 1)) \cdot x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, n, p\}, x] \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{GtQ}[q_1, 1] \&\& \text{NeQ}[n(p_1 + q_1) + 1, 0] \&\& !\text{IGtQ}[p, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$$

Rule 528

$$\text{Int}[(a + (b \cdot x)^n)^{p_1} ((c + (d \cdot x)^n)^{q_1}) ((e + (f \cdot x)^n)), x_Symbol] \rightarrow \text{Simp}[(f \cdot x (a + bx^n)^{p_1 + 1} (c + dx^n)^{q_1}) / (b(n(p_1 + q_1 + 1) + 1)), x] + \text{Dist}[1/(b(n(p_1 + q_1 + 1) + 1)), \text{Int}[(a + bx^n)^p (c + dx^n)^{q_1 - 1} \text{Simp}[c(b \cdot e - a \cdot f + b \cdot e \cdot n(p_1 + q_1 + 1)) + (d(b \cdot e - a \cdot f) + f \cdot n \cdot q_1 (b \cdot c - a \cdot d) + b \cdot d \cdot e \cdot n(p_1 + q_1 + 1)) \cdot x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \text{GtQ}[q_1, 0] \&\& \text{NeQ}[n(p_1 + q_1 + 1) + 1, 0]$$

Rule 1879

$$\text{Int}[(c + (d \cdot x))/\sqrt{(a + (b \cdot x)^3)}, x_Symbol] \rightarrow \text{With}\{r = \text{Numer}[\text{Simplify}[(1 + \sqrt{3})d/c], s = \text{Denom}[\text{Simplify}[(1 + \sqrt{3})d/c]]\}, \text{Simp}[(2 \cdot d \cdot s^3 \sqrt{a + bx^3}) / (a \cdot r^2 ((1 - \sqrt{3})s + rx)), x] + \text{Simp}[(3^{1/4} \sqrt{2 + \sqrt{3}} d \cdot s (s + rx) \sqrt{(s^2 - r \cdot s \cdot x + r^2 x^2)}) / ((1 - \sqrt{3})s + rx)^2 \text{EllipticE}[\text{ArcSin}[(1 + \sqrt{3})s + rx]/((1 - \sqrt{3})s + rx)], -7 + 4\sqrt{3}]/(r^2 \sqrt{a + bx^3}) \sqrt{-((s(s + rx))/((1 - \sqrt{3})s + rx)^2)}], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NegQ}[a] \&\& \text{EqQ}[b \cdot c^3 - 2(5 + 3\sqrt{3}) \cdot a \cdot d^3, 0]$$

Rubi steps

$$\begin{aligned}
\int \frac{(3a + bx^2)^4}{\sqrt[3]{a - bx^2}} dx &= -\frac{3}{25}x(a - bx^2)^{2/3}(3a + bx^2)^3 - \frac{3 \int \frac{(3a + bx^2)^2(-78a^2b - 50ab^2x^2)}{\sqrt[3]{a - bx^2}} dx}{25b} \\
&= -\frac{18}{19}ax(a - bx^2)^{2/3}(3a + bx^2)^2 - \frac{3}{25}x(a - bx^2)^{2/3}(3a + bx^2)^3 + \frac{9 \int \frac{(3a + bx^2)(1632a^3b^2 + 1344a^2b^3)}{\sqrt[3]{a - bx^2}}}{475b^2} \\
&= -\frac{36288a^2x(a - bx^2)^{2/3}(3a + bx^2)}{6175} - \frac{18}{19}ax(a - bx^2)^{2/3}(3a + bx^2)^2 - \frac{3}{25}x(a - bx^2)^{2/3}(3a + bx^2)^3 \\
&= -\frac{1552608a^3x(a - bx^2)^{2/3}}{43225} - \frac{36288a^2x(a - bx^2)^{2/3}(3a + bx^2)}{6175} - \frac{18}{19}ax(a - bx^2)^{2/3}(3a + bx^2)^2 \\
&= -\frac{1552608a^3x(a - bx^2)^{2/3}}{43225} - \frac{36288a^2x(a - bx^2)^{2/3}(3a + bx^2)}{6175} - \frac{18}{19}ax(a - bx^2)^{2/3}(3a + bx^2)^2 \\
&= -\frac{1552608a^3x(a - bx^2)^{2/3}}{43225} - \frac{36288a^2x(a - bx^2)^{2/3}(3a + bx^2)}{6175} - \frac{18}{19}ax(a - bx^2)^{2/3}(3a + bx^2)^2 \\
&= -\frac{1552608a^3x(a - bx^2)^{2/3}}{43225} - \frac{36288a^2x(a - bx^2)^{2/3}(3a + bx^2)}{6175} - \frac{18}{19}ax(a - bx^2)^{2/3}(3a + bx^2)^2
\end{aligned}$$

Mathematica [C] time = 5.06, size = 98, normalized size = 0.15

$$\frac{3x \left(2108160a^4 \sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1 \left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a} \right) - 941085a^4 + 727830a^3bx^2 + 184044a^2b^2x^4 + 27482ab^3x^6 + 1729b^4x^8 \right)}{43225 \sqrt[3]{a - bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)^4/(a - b*x^2)^(1/3), x]

[Out] (3*x*(-941085*a^4 + 727830*a^3*b*x^2 + 184044*a^2*b^2*x^4 + 27482*a*b^3*x^6 + 1729*b^4*x^8 + 2108160*a^4*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a]))/(43225*(a - b*x^2)^(1/3))

fricas [F] time = 1.11, size = 0, normalized size = 0.00

$$\text{integral} \left(-\frac{(b^4x^8 + 12ab^3x^6 + 54a^2b^2x^4 + 108a^3bx^2 + 81a^4)(-bx^2 + a)^{2/3}}{bx^2 - a}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^4/(-b*x^2+a)^(1/3), x, algorithm="fricas")

[Out] integral(-(b^4*x^8 + 12*a*b^3*x^6 + 54*a^2*b^2*x^4 + 108*a^3*b*x^2 + 81*a^4)*(-b*x^2 + a)^(2/3)/(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^4}{(-bx^2 + a)^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^4/(-b*x^2+a)^(1/3),x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^4/(-b*x^2 + a)^(1/3), x)

maple [F] time = 0.41, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^4}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)^4/(-b*x^2+a)^(1/3),x)

[Out] int((b*x^2+3*a)^4/(-b*x^2+a)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^4}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^4/(-b*x^2+a)^(1/3),x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^4/(-b*x^2 + a)^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + 3a)^4}{(a - bx^2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)^4/(a - b*x^2)^(1/3),x)

[Out] int((3*a + b*x^2)^4/(a - b*x^2)^(1/3), x)

sympy [A] time = 5.40, size = 165, normalized size = 0.25

$$81a^{\frac{11}{3}}x_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{bx^2e^{2i\pi}}{a}\right) + 36a^{\frac{8}{3}}bx^3_2F_1\left(\frac{1}{3}, \frac{3}{2} \middle| \frac{bx^2e^{2i\pi}}{a}\right) + \frac{54a^{\frac{5}{3}}b^2x^5_2F_1\left(\frac{1}{3}, \frac{5}{2} \middle| \frac{bx^2e^{2i\pi}}{a}\right)}{5} + \frac{12a^{\frac{2}{3}}b^3x^7_2F_1\left(\frac{1}{3}, \frac{7}{2} \middle| \frac{bx^2e^{2i\pi}}{a}\right)}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)**4/(-b*x**2+a)**(1/3),x)

[Out] 81*a**(11/3)*x*hyper((1/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) + 36*a**(8/3)*b*x**3*hyper((1/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a) + 54*a**(5/3)*b**2*x**5*hyper((1/3, 5/2), (7/2,), b*x**2*exp_polar(2*I*pi)/a)/5 + 12*a**(2/3)*b**3*x**7*hyper((1/3, 7/2), (9/2,), b*x**2*exp_polar(2*I*pi)/a)/7 + b**4*x**9*hyper((1/3, 9/2), (11/2,), b*x**2*exp_polar(2*I*pi)/a)/(9*a**(1/3))

3.123 $\int \frac{(3a+bx^2)^3}{\sqrt[3]{a-bx^2}} dx$

Optimal. Leaf size=628

$$\frac{71712\sqrt{2}3^{3/4}a^{10/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{1729bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] -15768/1729*a^2*x*(-b*x^2+a)^(2/3)-324/247*a*x*(-b*x^2+a)^(2/3)*(b*x^2+3*a)
 -3/19*x*(-b*x^2+a)^(2/3)*(b*x^2+3*a)^2-215136/1729*a^3*x/(-b*x^2+a)^(1/3)
 +a^(1/3)*(1-3^(1/2)))+71712/1729*3^(3/4)*a^(10/3)*(a^(1/3)-(-b*x^2+a)^(1/3))
)*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-b*x^2+a)^(1/3)+a^(1/3)
 *(1-3^(1/2))),2*I-I*3^(1/2))*2^(1/2)*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+
 (-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)/b/x/(-a^(1/3)
 *(a^(1/3)-(-b*x^2+a)^(1/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)
 ^2)^(1/2)-107568/1729*3^(1/4)*a^(10/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-
 (-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))
)),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-
 b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))/b/x/
 (-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))
)^2)^(1/2)

Rubi [A] time = 0.44, antiderivative size = 628, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 24, number of rules / integrand size = 0.292, Rules used = {416, 528, 388, 235, 304, 219, 1879}

$$\frac{215136a^3x}{1729\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}-\frac{15768a^2x(a-bx^2)^{2/3}}{1729}+\frac{71712\sqrt{2}3^{3/4}a^{10/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}{1729bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

[In] Int[(3*a + b*x^2)^3/(a - b*x^2)^(1/3), x]

[Out] (-15768*a^2*x*(a - b*x^2)^(2/3))/1729 - (324*a*x*(a - b*x^2)^(2/3)*(3*a + b*x^2))/247 - (3*x*(a - b*x^2)^(2/3)*(3*a + b*x^2)^2)/19 - (215136*a^3*x)/(1729*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) - (107568*3^(1/4)*Sqrt[2 + Sqrt[3]]*a^(10/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]]]/(1729*b*x*Sqrt[-(a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2])) + (71712*Sqrt[2]*3^(3/4)*a^(10/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)*EllipticF[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]]]/(1729*b*x*Sqrt[-(a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]))

Rule 219

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s

$$\frac{x + r^2 x^2}{((1 - \sqrt{3})s + rx)^2} \text{EllipticF}[\text{ArcSin}[\frac{(1 + \sqrt{3})s + rx}{(1 - \sqrt{3})s + rx}], -7 + 4\sqrt{3}]/(3^{1/4} r \sqrt{a + bx^3}) \sqrt{-((s(s + rx))/((1 - \sqrt{3})s + rx)^2)}], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a]$$

Rule 235

$$\text{Int}[(a + (b \cdot x)^2)^{-1/3}, x_Symbol] \rightarrow \text{Dist}[(3\sqrt{bx^2})/(2bx), \text{Subst}[\text{Int}[x/\sqrt{-a + x^3}], x], x, (a + bx^2)^{1/3}], x] /; \text{FreeQ}\{a, b\}, x]$$

Rule 304

$$\text{Int}[x/\sqrt{(a + (b \cdot x)^3)}, x_Symbol] \rightarrow \text{With}\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\sqrt{2}s)/(\sqrt{2 - \sqrt{3}}r), \text{Int}[1/\sqrt{a + bx^3}], x], x] + \text{Dist}[1/r, \text{Int}[(1 + \sqrt{3})s + rx]/\sqrt{a + bx^3}], x], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a]$$

Rule 388

$$\text{Int}[(a + (b \cdot x)^n)^p \cdot ((c + (d \cdot x)^n)), x_Symbol] \rightarrow \text{Simp}[(d \cdot x \cdot (a + bx^n)^{p+1})/(b \cdot (n \cdot (p+1) + 1)), x] - \text{Dist}[(a \cdot d - b \cdot c \cdot (n \cdot (p+1) + 1))/(b \cdot (n \cdot (p+1) + 1)), \text{Int}[(a + bx^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{NeQ}[n \cdot (p+1) + 1, 0]$$

Rule 416

$$\text{Int}[(a + (b \cdot x)^n)^p \cdot ((c + (d \cdot x)^n))^q, x_Symbol] \rightarrow \text{Simp}[(d \cdot x \cdot (a + bx^n)^{p+1} \cdot (c + dx^n)^{q-1})/(b \cdot (n \cdot (p+q) + 1)), x] + \text{Dist}[1/(b \cdot (n \cdot (p+q) + 1)), \text{Int}[(a + bx^n)^p \cdot (c + dx^n)^{q-2} \cdot \text{Simp}[c \cdot (b \cdot c \cdot (n \cdot (p+q) + 1) - a \cdot d) + d \cdot (b \cdot c \cdot (n \cdot (p+2q-1) + 1) - a \cdot d \cdot (n \cdot (q-1) + 1)) \cdot x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, n, p\}, x] \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{GtQ}[q, 1] \&\& \text{NeQ}[n \cdot (p+q) + 1, 0] \&\& !\text{IGtQ}[p, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$$

Rule 528

$$\text{Int}[(a + (b \cdot x)^n)^p \cdot ((c + (d \cdot x)^n))^q \cdot ((e + (f \cdot x)^n)), x_Symbol] \rightarrow \text{Simp}[(f \cdot x \cdot (a + bx^n)^{p+1} \cdot (c + dx^n)^q)/(b \cdot (n \cdot (p+q+1) + 1)), x] + \text{Dist}[1/(b \cdot (n \cdot (p+q+1) + 1)), \text{Int}[(a + bx^n)^p \cdot (c + dx^n)^{q-1} \cdot \text{Simp}[c \cdot (b \cdot e - a \cdot f + b \cdot e \cdot n \cdot (p+q+1)) + (d \cdot (b \cdot e - a \cdot f) + f \cdot n \cdot q \cdot (b \cdot c - a \cdot d) + b \cdot d \cdot e \cdot n \cdot (p+q+1)) \cdot x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \text{GtQ}[q, 0] \&\& \text{NeQ}[n \cdot (p+q+1) + 1, 0]$$

Rule 1879

$$\text{Int}[(c + (d \cdot x))/\sqrt{(a + (b \cdot x)^3)}, x_Symbol] \rightarrow \text{With}\{r = \text{Numer}[\text{Simplify}[(1 + \sqrt{3})d/c]], s = \text{Denom}[\text{Simplify}[(1 + \sqrt{3})d/c]]\}, \text{Simp}[(2 \cdot d \cdot s^3 \sqrt{a + bx^3})/(a \cdot r^2 \cdot ((1 - \sqrt{3})s + rx)), x] + \text{Simp}[(3^{1/4} \sqrt{2 + \sqrt{3}} \cdot d \cdot s \cdot (s + rx) \sqrt{(s^2 - r \cdot s \cdot x + r^2 \cdot x^2)})/((1 - \sqrt{3})s + rx)^2 \cdot \text{EllipticE}[\text{ArcSin}[\frac{(1 + \sqrt{3})s + rx}{(1 - \sqrt{3})s + rx}], -7 + 4\sqrt{3}]/(r^2 \sqrt{a + bx^3} \sqrt{-((s(s + rx))/((1 - \sqrt{3})s + rx)^2)}), x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NegQ}[a] \&\& \text{EqQ}[b \cdot c^3 - 2 \cdot (5 + 3\sqrt{3}) \cdot a \cdot d^3, 0]$$

Rubi steps

$$\begin{aligned}
\int \frac{(3a + bx^2)^3}{\sqrt[3]{a - bx^2}} dx &= -\frac{3}{19}x(a - bx^2)^{2/3}(3a + bx^2)^2 - \frac{3 \int \frac{(3a+bx^2)(-60a^2b-36ab^2x^2)}{\sqrt[3]{a-bx^2}} dx}{19b} \\
&= -\frac{324}{247}ax(a - bx^2)^{2/3}(3a + bx^2) - \frac{3}{19}x(a - bx^2)^{2/3}(3a + bx^2)^2 + \frac{9 \int \frac{888a^3b^2+584a^2b^3x^2}{\sqrt[3]{a-bx^2}} dx}{247b^2} \\
&= -\frac{15768a^2x(a - bx^2)^{2/3}}{1729} - \frac{324}{247}ax(a - bx^2)^{2/3}(3a + bx^2) - \frac{3}{19}x(a - bx^2)^{2/3}(3a + bx^2)^2 + \dots \\
&= -\frac{15768a^2x(a - bx^2)^{2/3}}{1729} - \frac{324}{247}ax(a - bx^2)^{2/3}(3a + bx^2) - \frac{3}{19}x(a - bx^2)^{2/3}(3a + bx^2)^2 - \dots \\
&= -\frac{15768a^2x(a - bx^2)^{2/3}}{1729} - \frac{324}{247}ax(a - bx^2)^{2/3}(3a + bx^2) - \frac{3}{19}x(a - bx^2)^{2/3}(3a + bx^2)^2 + \dots \\
&= -\frac{15768a^2x(a - bx^2)^{2/3}}{1729} - \frac{324}{247}ax(a - bx^2)^{2/3}(3a + bx^2) - \frac{3}{19}x(a - bx^2)^{2/3}(3a + bx^2)^2 - \dots
\end{aligned}$$

Mathematica [C] time = 5.05, size = 88, normalized size = 0.14

$$\frac{3 \left(23904a^3x \sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1 \left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a} \right) - 8343a^3x + 7041a^2bx^3 + 1211ab^2x^5 + 91b^3x^7 \right)}{1729 \sqrt[3]{a - bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)^3/(a - b*x^2)^(1/3), x]

[Out] (3*(-8343*a^3*x + 7041*a^2*b*x^3 + 1211*a*b^2*x^5 + 91*b^3*x^7 + 23904*a^3*x*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a]))/(1729*(a - b*x^2)^(1/3))

fricas [F] time = 0.95, size = 0, normalized size = 0.00

$$\text{integral} \left(-\frac{(b^3x^6 + 9ab^2x^4 + 27a^2bx^2 + 27a^3)(-bx^2 + a)^{\frac{2}{3}}}{bx^2 - a}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(1/3), x, algorithm="fricas")

[Out] integral(-(b^3*x^6 + 9*a*b^2*x^4 + 27*a^2*b*x^2 + 27*a^3)*(-b*x^2 + a)^(2/3)/(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(1/3),x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^3/(-b*x^2 + a)^(1/3), x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)^3/(-b*x^2+a)^(1/3),x)

[Out] int((b*x^2+3*a)^3/(-b*x^2+a)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(1/3),x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^3/(-b*x^2 + a)^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + 3a)^3}{(a - bx^2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)^3/(a - b*x^2)^(1/3),x)

[Out] int((3*a + b*x^2)^3/(a - b*x^2)^(1/3), x)

sympy [A] time = 4.38, size = 129, normalized size = 0.21

$$27a^{\frac{8}{3}}x {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) + 9a^{\frac{5}{3}}bx^3 {}_2F_1\left(\frac{1}{3}, \frac{3}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) + \frac{9a^{\frac{2}{3}}b^2x^5 {}_2F_1\left(\frac{1}{3}, \frac{5}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{5} + \frac{b^3x^7 {}_2F_1\left(\frac{1}{3}, \frac{7}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{7\sqrt[3]{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)**3/(-b*x**2+a)**(1/3),x)

[Out] 27*a**(8/3)*x*hyper((1/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) + 9*a**(5/3)*b*x**3*hyper((1/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a) + 9*a**(2/3)*b**2*x**5*hyper((1/3, 5/2), (7/2,), b*x**2*exp_polar(2*I*pi)/a)/5 + b**3*x**7*hyper((1/3, 7/2), (9/2,), b*x**2*exp_polar(2*I*pi)/a)/(7*a**(1/3))

$$3.124 \quad \int \frac{(3a+bx^2)^2}{\sqrt[3]{a-bx^2}} dx$$

Optimal. Leaf size=597

$$\frac{1080\sqrt{2}3^{3/4}a^{7/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{91bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] $-198/91*a*x*(-b*x^2+a)^{(2/3)}-3/13*x*(-b*x^2+a)^{(2/3)}*(b*x^2+3*a)-3240/91*a^2*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+1080/91*3^{(3/4)}*a^{(7/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}-1620/91*3^{(1/4)}*a^{(7/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))),2*I-I*3^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.38, antiderivative size = 597, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {416, 388, 235, 304, 219, 1879}

$$\frac{3240a^2x}{91\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{1080\sqrt{2}3^{3/4}a^{7/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}F\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right)\right)}{91bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(3a + b*x^2)^2/(a - b*x^2)^{(1/3)}, x]$

[Out] $(-198*a*x*(a - b*x^2)^{(2/3)})/91 - (3*x*(a - b*x^2)^{(2/3)}*(3*a + b*x^2))/13 - (3240*a^2*x)/(91*((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})) - (1620*3^{(1/4)}*\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]]*a^{(7/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}))], -7 + 4*\operatorname{Sqrt}[3])]/(91*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])) + (1080*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(7/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}))], -7 + 4*\operatorname{Sqrt}[3])]/(91*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]))$

Rule 219

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_.)*(x_)^3], x_Symbol] \rightarrow \operatorname{With}[\{r = \operatorname{Numer}[\operatorname{Rt}[b/a, 3]], s = \operatorname{Denom}[\operatorname{Rt}[b/a, 3]]\}, \operatorname{Simp}[(2*\operatorname{Sqrt}[2 - \operatorname{Sqrt}[3]]*(s + r*x)*\operatorname{Sqrt}[(s^2 - r*s*x + r^2*x^2)]/((1 - \operatorname{Sqrt}[3])*s + r*x))^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*s + r*x)/((1 - \operatorname{Sqrt}[3])*s + r*x))], -7 + 4*\operatorname{Sqrt}[3])]/(3^{(1/4)}*r*\operatorname{Sqrt}[a + b*x^3]$

] * Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)], x]] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 235

Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

Rule 304

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sqrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)), x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q - 1) + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 1879

Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(3a + bx^2)^2}{\sqrt[3]{a - bx^2}} dx &= -\frac{3}{13}x(a - bx^2)^{2/3}(3a + bx^2) - \frac{3 \int \frac{-42a^2b - 22ab^2x^2}{\sqrt[3]{a - bx^2}} dx}{13b} \\
&= -\frac{198}{91}ax(a - bx^2)^{2/3} - \frac{3}{13}x(a - bx^2)^{2/3}(3a + bx^2) + \frac{1}{91}(1080a^2) \int \frac{1}{\sqrt[3]{a - bx^2}} dx \\
&= -\frac{198}{91}ax(a - bx^2)^{2/3} - \frac{3}{13}x(a - bx^2)^{2/3}(3a + bx^2) - \frac{(1620a^2\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{x}{\sqrt{-a+x^3}} dx, x\right)}{91bx} \\
&= -\frac{198}{91}ax(a - bx^2)^{2/3} - \frac{3}{13}x(a - bx^2)^{2/3}(3a + bx^2) + \frac{(1620a^2\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{(1+\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx, x\right)}{91bx} \\
&= -\frac{198}{91}ax(a - bx^2)^{2/3} - \frac{3}{13}x(a - bx^2)^{2/3}(3a + bx^2) - \frac{3240a^2x}{91\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} - \frac{1620a^2}{91}
\end{aligned}$$

Mathematica [C] time = 4.49, size = 158, normalized size = 0.26

$$\frac{x\sqrt[3]{1 - \frac{bx^2}{a}} \left(4b(3ax + bx^3)^2 {}_3F_2\left(\frac{4}{3}, \frac{3}{2}, 2; 1, \frac{9}{2}; \frac{bx^2}{a}\right) + 8bx^2(18a^2 + 9abx^2 + b^2x^4) {}_2F_1\left(\frac{4}{3}, \frac{3}{2}; \frac{9}{2}; \frac{bx^2}{a}\right) + 63a(45a^2 + 10abx^2 + b^2x^4)\right)}{315a\sqrt[3]{a - bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(3*a + b*x^2)^2/(a - b*x^2)^(1/3), x]

[Out] (x*(1 - (b*x^2)/a)^(1/3)*(63*a*(45*a^2 + 10*a*b*x^2 + b^2*x^4)*Hypergeometric2F1[1/3, 1/2, 7/2, (b*x^2)/a] + 8*b*x^2*(18*a^2 + 9*a*b*x^2 + b^2*x^4)*Hypergeometric2F1[4/3, 3/2, 9/2, (b*x^2)/a] + 4*b*(3*a*x + b*x^3)^2*HypergeometricPFQ[{4/3, 3/2, 2}, {1, 9/2}, (b*x^2)/a]))/(315*a*(a - b*x^2)^(1/3))

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{(b^2x^4 + 6abx^2 + 9a^2)(-bx^2 + a)^{\frac{2}{3}}}{bx^2 - a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(1/3), x, algorithm="fricas")

[Out] integral(-(b^2*x^4 + 6*a*b*x^2 + 9*a^2)*(-b*x^2 + a)^(2/3)/(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^2}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(1/3), x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^2/(-b*x^2 + a)^(1/3), x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^2}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)^2/(-b*x^2+a)^(1/3), x)

[Out] int((b*x^2+3*a)^2/(-b*x^2+a)^(1/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^2}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(1/3), x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^2/(-b*x^2 + a)^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + 3a)^2}{(a - bx^2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)^2/(a - b*x^2)^(1/3), x)

[Out] int((3*a + b*x^2)^2/(a - b*x^2)^(1/3), x)

sympy [A] time = 3.35, size = 94, normalized size = 0.16

$$9a^{\frac{5}{3}}x {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) + 2a^{\frac{2}{3}}bx^3 {}_2F_1\left(\frac{1}{3}, \frac{3}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) + \frac{b^2x^5 {}_2F_1\left(\frac{1}{3}, \frac{5}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{5\sqrt[3]{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)**2/(-b*x**2+a)**(1/3), x)

[Out] 9*a**(5/3)*x*hyper((1/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) + 2*a**(2/3)*b*x**3*hyper((1/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a) + b**2*x**5*hyper((1/3, 5/2), (7/2,), b*x**2*exp_polar(2*I*pi)/a)/(5*a**(1/3))

$$3.125 \quad \int \frac{3a+bx^2}{\sqrt[3]{a-bx^2}} dx$$

Optimal. Leaf size=568

$$24\sqrt{2}3^{3/4}a^{4/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)36\sqrt[4]{3}$$

$$7bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}$$

[Out] $-3/7*x*(-b*x^2+a)^{(2/3)}-72/7*a*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+24/7*3^{(3/4)}*a^{(4/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}-36/7*3^{(1/4)}*a^{(4/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}$

Rubi [A] time = 0.30, antiderivative size = 568, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {388, 235, 304, 219, 1879}

$$24\sqrt{2}3^{3/4}a^{4/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}F\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right)\middle| -7+4\sqrt{3}\right)36\sqrt[4]{3}\sqrt{2}$$

$$7bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}$$

Antiderivative was successfully verified.

[In] Int[(3*a + b*x^2)/(a - b*x^2)^(1/3), x]

[Out] $(-3*x*(a-b*x^2)^{(2/3)})/7-(72*a*x)/(7*((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))-(36*3^{(1/4)}*\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*a^{(4/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))],-7+4*\operatorname{Sqrt}[3])/((7*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]))+(24*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(4/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))],-7+4*\operatorname{Sqrt}[3])/((7*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]))$

Rule 219

Int[1/Sqrt[(a_) + (b_)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^{(1/4)}*r*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2))], x] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 235

Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

Rule 304

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sqrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 1879

Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)]], -7 + 4*Sqrt[3])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/(1 - Sqrt[3])*s + r*x)^2]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned} \int \frac{3a + bx^2}{\sqrt[3]{a - bx^2}} dx &= -\frac{3}{7}x(a - bx^2)^{2/3} + \frac{1}{7}(24a) \int \frac{1}{\sqrt[3]{a - bx^2}} dx \\ &= -\frac{3}{7}x(a - bx^2)^{2/3} - \frac{(36a\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{x}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{7bx} \\ &= -\frac{3}{7}x(a - bx^2)^{2/3} + \frac{(36a\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{(1+\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{7bx} - \frac{(36\sqrt{2(2+\sqrt{3})})}{7bx} \\ &= -\frac{3}{7}x(a - bx^2)^{2/3} - \frac{72ax}{7\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} - \frac{36^4\sqrt{3}\sqrt{2+\sqrt{3}}a^{4/3}\left(\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)}{7\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} \end{aligned}$$

Mathematica [C] time = 0.03, size = 62, normalized size = 0.11

$$\frac{3x\left(8a\sqrt[3]{1-\frac{bx^2}{a}} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a}\right) - a + bx^2\right)}{7\sqrt[3]{a - bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)/(a - b*x^2)^(1/3),x]

[Out] (3*x*(-a + b*x^2 + 8*a*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a]))/(7*(a - b*x^2)^(1/3))

fricas [F] time = 0.84, size = 0, normalized size = 0.00

$$\text{integral} \left(-\frac{(bx^2 + 3a)(-bx^2 + a)^{\frac{2}{3}}}{bx^2 - a}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(1/3),x, algorithm="fricas")

[Out] integral(-(b*x^2 + 3*a)*(-b*x^2 + a)^(2/3)/(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(1/3),x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)/(-b*x^2 + a)^(1/3), x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)/(-b*x^2+a)^(1/3),x)

[Out] int((b*x^2+3*a)/(-b*x^2+a)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(1/3),x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)/(-b*x^2 + a)^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{bx^2 + 3a}{(a - bx^2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)/(a - b*x^2)^(1/3),x)

[Out] int((3*a + b*x^2)/(a - b*x^2)^(1/3), x)

sympy [A] time = 2.01, size = 60, normalized size = 0.11

$$3a^{\frac{2}{3}}x {}_2F_1\left(\frac{1}{3}, \frac{1}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right) + \frac{bx^3 {}_2F_1\left(\frac{1}{3}, \frac{3}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{3\sqrt[3]{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)/(-b*x**2+a)**(1/3),x)

[Out] 3*a**(2/3)*x*hyper((1/3, 1/2), (3/2,), b*x**2*exp_polar(2*I*pi)/a) + b*x**3*hyper((1/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a)/(3*a**(1/3))

$$3.126 \quad \int \frac{1}{\sqrt[3]{a-bx^2} (3a+bx^2)} dx$$

Optimal. Leaf size=204

$$\frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt[6]{a} (\sqrt[3]{2} \sqrt[3]{a-bx^2} + \sqrt[3]{a})}\right)}{2 \cdot 2^{2/3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6 \cdot 2^{2/3} a^{5/6} \sqrt{b}}$$

[Out] $1/4 \cdot \arctan(x \cdot b^{1/2} / a^{1/6} / (a^{1/3} + 2^{1/3}) \cdot (-b \cdot x^2 + a)^{1/3}) \cdot 2^{1/3} / a^{5/6} / b^{1/2} - 1/12 \cdot \arctan(x \cdot b^{1/2} / a^{1/6}) \cdot 2^{1/3} / a^{5/6} / b^{1/2} + 1/12 \cdot \arctan(a^{1/6} \cdot (a^{1/3} - 2^{1/3}) \cdot (-b \cdot x^2 + a)^{1/3}) \cdot 3^{1/2} / x / b^{1/2} \cdot 2^{1/3} / a^{5/6} \cdot 3^{1/2} / b^{1/2} + 1/12 \cdot \arctan(3^{1/2} \cdot a^{1/2} / x / b^{1/2}) \cdot 2^{1/3} / a^{5/6} \cdot 3^{1/2} / b^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 204, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {393}

$$\frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt[6]{a} (\sqrt[3]{2} \sqrt[3]{a-bx^2} + \sqrt[3]{a})}\right)}{2 \cdot 2^{2/3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6 \cdot 2^{2/3} a^{5/6} \sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(1/3)*(3*a + b*x^2)), x]

[Out] ArcTan[(Sqrt[3]*Sqrt[a])/(Sqrt[b]*x)]/(2*2^(2/3)*Sqrt[3]*a^(5/6)*Sqrt[b]) + ArcTan[(Sqrt[3]*a^(1/6)*(a^(1/3) - 2^(1/3)*(a - b*x^2)^(1/3)))/(Sqrt[b]*x)]/(2*2^(2/3)*Sqrt[3]*a^(5/6)*Sqrt[b]) - ArcTanh[(Sqrt[b]*x)/Sqrt[a]]/(6*2^(2/3)*a^(5/6)*Sqrt[b]) + ArcTanh[(Sqrt[b]*x)/(a^(1/6)*(a^(1/3) + 2^(1/3)*(a - b*x^2)^(1/3)))]/(2*2^(2/3)*a^(5/6)*Sqrt[b])

Rule 393

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{a-bx^2} (3a+bx^2)} dx = \frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6 \cdot 2^{2/3} a^{5/6} \sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt[6]{a} (\sqrt[3]{2} \sqrt[3]{a-bx^2} + \sqrt[3]{a})}\right)}{2 \cdot 2^{2/3} a^{5/6} \sqrt{b}}$$

Mathematica [C] time = 0.05, size = 162, normalized size = 0.79

$$\frac{9axF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{\sqrt[3]{a-bx^2} (3a+bx^2) \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)\right) + 9aF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(1/3)*(3*a + b*x^2)),x]

[Out] (9*a*x*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/((a - b*x^2)^(1/3)*(3*a + b*x^2)*(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 3*a)*(-b*x^2 + a)^(1/3)), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{1}{3}}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x)

[Out] int(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 3*a)*(-b*x^2 + a)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{1/3}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(1/3)*(3*a + b*x^2)),x)

[Out] `int(1/((a - b*x^2)^(1/3)*(3*a + b*x^2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-b*x**2+a)**(1/3)/(b*x**2+3*a), x)`

[Out] `Integral(1/((a - b*x**2)**(1/3)*(3*a + b*x**2)), x)`

$$3.127 \quad \int \frac{1}{\sqrt[3]{a-bx^2} (3a+bx^2)^2} dx$$

Optimal. Leaf size=787

$$\frac{\left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}}\right), 4\sqrt{3} - 7\right) \tan^{-1}\left(\frac{\sqrt{3} \sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{8 \cdot 2^{2/3} \sqrt{3} a}\right)}{12\sqrt{2} \sqrt[3]{3} a^{5/3} bx \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}} + \frac{\sqrt{3} \sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{8 \cdot 2^{2/3} \sqrt{3} a}$$

[Out] $1/24*x*(-b*x^2+a)^{(2/3)}/a^2/(b*x^2+3*a)-1/24*x/a^2/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+1/16*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/(a^{(1/3)}+2^{(1/3)}*(-b*x^2+a)^{(1/3)})^2^{(1/3)}/a^{(11/6)}/b^{(1/2)}-1/48*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/2)})^2^{(1/3)}/a^{(11/6)}/b^{(1/2)}+1/48*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(-b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})^2^{(1/3)}/a^{(11/6)}*3^{(1/2)}/b^{(1/2)}+1/48*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})^2^{(1/3)}/a^{(11/6)}*3^{(1/2)}/b^{(1/2)}+1/72*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2^{(1/2)}*3^{(3/4)}/a^{(5/3)}/b/x*2^{(1/2)}/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2^{(1/2)}-1/48*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})*3^{(1/4)}/a^{(5/3)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2^{(1/2)}$

Rubi [A] time = 0.44, antiderivative size = 787, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {414, 530, 235, 304, 219, 1879, 393}

$$\frac{x(a-bx^2)^{2/3}}{24a^2(3a+bx^2)} - \frac{x}{24a^2\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{8 \cdot 2^{2/3}\sqrt{3}a^{11/6}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt[3]{a}\left(\sqrt[3]{2}\sqrt[3]{a-bx^2} + \sqrt[3]{a}\right)}\right)}{8 \cdot 2^{2/3}a^{11/6}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(1/3)*(3*a + b*x^2)^2), x]

[Out] $(x*(a-b*x^2)^{(2/3)})/(24*a^2*(3*a+b*x^2))-x/(24*a^2*((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))+\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(8*2^{(2/3)}*\operatorname{Sqrt}[3]*a^{(11/6)}*\operatorname{Sqrt}[b])+\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(a-b*x^2)^{(1/3)}))/(\operatorname{Sqrt}[b]*x)]/(8*2^{(2/3)}*\operatorname{Sqrt}[3]*a^{(11/6)}*\operatorname{Sqrt}[b])-\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(24*2^{(2/3)}*a^{(11/6)}*\operatorname{Sqrt}[b])+\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{(1/6)}*(a^{(1/3)}+2^{(1/3)}*(a-b*x^2)^{(1/3)}))]/(8*2^{(2/3)}*a^{(11/6)}*\operatorname{Sqrt}[b])-(\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})], -7+4*\operatorname{Sqrt}[3]]/(16*3^{(3/4)}*a^{(5/3)}*b*x*\operatorname{Sqrt}[-(a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]))+(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})], -7+4*\operatorname{Sqrt}[3]]$

)/(12*Sqrt[2]*3^(1/4)*a^(5/3)*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/(1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]])

Rule 219

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]]]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2))], x] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 235

Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

Rule 304

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sqrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 393

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3))])/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 530

Int[(((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1879

Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]]]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)

)/((1 - Sqrt[3])*s + r*x)^2)), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt[3]{a-bx^2} (3a+bx^2)^2} dx &= \frac{x(a-bx^2)^{2/3}}{24a^2(3a+bx^2)} - \frac{\int \frac{-7ab-\frac{b^2x^2}{3}}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx}{24a^2b} \\ &= \frac{x(a-bx^2)^{2/3}}{24a^2(3a+bx^2)} + \frac{\int \frac{1}{\sqrt[3]{a-bx^2}} dx}{72a^2} + \frac{\int \frac{1}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx}{4a} \\ &= \frac{x(a-bx^2)^{2/3}}{24a^2(3a+bx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{24 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} \\ &= \frac{x(a-bx^2)^{2/3}}{24a^2(3a+bx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{24 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} \\ &= \frac{x(a-bx^2)^{2/3}}{24a^2(3a+bx^2)} - \frac{x}{24a^2\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{24 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} \end{aligned}$$

Mathematica [C] time = 0.16, size = 234, normalized size = 0.30

$$x \left(\frac{bx^2 \sqrt[3]{1-\frac{bx^2}{a}} F_1\left(\frac{3}{2}; \frac{1}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{a^3} + \frac{27 \left(\frac{a-bx^2}{a^2} + \frac{{}_6F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{3a+bx^2} \right) \right) / (648 \sqrt[3]{a-bx^2})$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(1/3)*(3*a + b*x^2)^2), x]

[Out] (x*((b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])/a^3 + (27*((a - b*x^2)/a^2 + (63*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])))/(3*a + b*x^2))/((648*(a - b*x^2)^(1/3))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^2 (-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^2,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 3*a)^2*(-b*x^2 + a)^(1/3)), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{1}{3}} (bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^2,x)

[Out] int(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^2 (-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 3*a)^2*(-b*x^2 + a)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{\frac{1}{3}} (bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(1/3)*(3*a + b*x^2)^2),x)

[Out] int(1/((a - b*x^2)^(1/3)*(3*a + b*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2+a)**(1/3)/(b*x**2+3*a)**2,x)

[Out] Integral(1/((a - b*x**2)**(1/3)*(3*a + b*x**2)**2), x)

$$3.128 \quad \int \frac{1}{\sqrt[3]{a-bx^2} (3a+bx^2)^3} dx$$

Optimal. Leaf size=818

$$\frac{5(a-bx^2)^{2/3}x}{288a^3(bx^2+3a)} - \frac{5x}{288a^3\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)} + \frac{(a-bx^2)^{2/3}x}{48a^2(bx^2+3a)^2} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{144 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}}{\sqrt[6]{bx}}\right)}{144 \cdot 2^{2/3}}$$

[Out] $1/48*x*(-b*x^2+a)^{(2/3)}/a^2/(b*x^2+3*a)^2+5/288*x*(-b*x^2+a)^{(2/3)}/a^3/(b*x^2+3*a)-5/288*x/a^3/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+5/288*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/(a^{(1/3)}+2^{(1/3)}*(-b*x^2+a)^{(1/3)})*2^{(1/3)}/a^{(17/6)}/b^{(1/2)}-5/864*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/2)})*2^{(1/3)}/a^{(17/6)}/b^{(1/2)}+5/864*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(-b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})*2^{(1/3)}/a^{(17/6)}*3^{(1/2)}/b^{(1/2)}+5/864*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})*2^{(1/3)}/a^{(17/6)}*3^{(1/2)}/b^{(1/2)}+5/864*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*3^{(3/4)}/a^{(8/3)}/b/x*2^{(1/2)}/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}-5/576*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})*3^{(1/4)}/a^{(8/3)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.54, antiderivative size = 818, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {414, 527, 530, 235, 304, 219, 1879, 393}

$$\frac{5(a-bx^2)^{2/3}x}{288a^3(bx^2+3a)} - \frac{5x}{288a^3\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)} + \frac{(a-bx^2)^{2/3}x}{48a^2(bx^2+3a)^2} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{144 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}}{\sqrt[6]{bx}}\right)}{144 \cdot 2^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(1/3)*(3*a + b*x^2)^3), x]

[Out] $(x*(a-b*x^2)^{(2/3)})/(48*a^2*(3*a+b*x^2)^2)+(5*x*(a-b*x^2)^{(2/3)})/(288*a^3*(3*a+b*x^2))-5*x/(288*a^3*((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))+5*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(144*2^{(2/3)}*\operatorname{Sqrt}[3]*a^{(17/6)}*\operatorname{Sqrt}[b])+5*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(a-b*x^2)^{(1/3)}))/(\operatorname{Sqrt}[b]*x)]/(144*2^{(2/3)}*\operatorname{Sqrt}[3]*a^{(17/6)}*\operatorname{Sqrt}[b])-5*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(432*2^{(2/3)}*a^{(17/6)}*\operatorname{Sqrt}[b])+5*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{(1/6)}*(a^{(1/3)}+2^{(1/3)}*(a-b*x^2)^{(1/3)}))]/(144*2^{(2/3)}*a^{(17/6)}*\operatorname{Sqrt}[b])-5*\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})), -7+4*\operatorname{Sqrt}[3]]/(19*2*3^{(3/4)}*a^{(8/3)}*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]))+(5*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})), -7+4*\operatorname{Sqrt}[3]]/(19*2*3^{(3/4)}*a^{(8/3)}*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]))]$

3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3)), -7 + 4*Sqrt[3]]/(144*Sqrt[2]*3^(1/4)*a^(8/3)*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)])

Rule 219

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)]], -7 + 4*Sqrt[3]]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2))], x] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 235

Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

Rule 304

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sqrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 393

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3))])/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]) /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 530

Int((((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p}

, n}, x]

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)]], -7 + 4*Sqrt[3])]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt[3]{a-bx^2} (3a+bx^2)^3} dx &= \frac{x(a-bx^2)^{2/3}}{48a^2(3a+bx^2)^2} - \frac{\int \frac{-15ab + \frac{5b^2x^2}{3}}{\sqrt[3]{a-bx^2}(3a+bx^2)^2} dx}{48a^2b} \\ &= \frac{x(a-bx^2)^{2/3}}{48a^2(3a+bx^2)^2} + \frac{5x(a-bx^2)^{2/3}}{288a^3(3a+bx^2)} + \frac{\int \frac{100a^2b^2 + \frac{20}{3}ab^3x^2}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx}{1152a^4b^2} \\ &= \frac{x(a-bx^2)^{2/3}}{48a^2(3a+bx^2)^2} + \frac{5x(a-bx^2)^{2/3}}{288a^3(3a+bx^2)} + \frac{5 \int \frac{1}{\sqrt[3]{a-bx^2}} dx}{864a^3} + \frac{5 \int \frac{1}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx}{72a^2} \\ &= \frac{x(a-bx^2)^{2/3}}{48a^2(3a+bx^2)^2} + \frac{5x(a-bx^2)^{2/3}}{288a^3(3a+bx^2)} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{144 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}}{\sqrt[3]{a-bx^2}}\right)}{144 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} \\ &= \frac{x(a-bx^2)^{2/3}}{48a^2(3a+bx^2)^2} + \frac{5x(a-bx^2)^{2/3}}{288a^3(3a+bx^2)} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{144 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}}{\sqrt[3]{a-bx^2}}\right)}{144 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} \\ &= \frac{x(a-bx^2)^{2/3}}{48a^2(3a+bx^2)^2} + \frac{5x(a-bx^2)^{2/3}}{288a^3(3a+bx^2)} - \frac{5x}{288a^3\left((1-\sqrt{3})\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)} + \frac{5 \tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}}{\sqrt[3]{a-bx^2}}\right)}{144 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} \end{aligned}$$

Mathematica [C] time = 0.17, size = 255, normalized size = 0.31

$$x \frac{\left(\frac{6075a^3 F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{(3a+bx^2) \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right)}{7776a^4 \sqrt[3]{a-bx^2}} + 5bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} F_1\left(\frac{3}{2}; \frac{1}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(1/3)*(3*a + b*x^2)^3), x]

[Out] (x*((27*a*(a - b*x^2)*(21*a + 5*b*x^2))/(3*a + b*x^2)^2 + 5*b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + (6075*

$a^3 \text{AppellF1}[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] / ((3*a + b*x^2) * (9*a \text{AppellF1}[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2 * (-\text{AppellF1}[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + \text{AppellF1}[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])))) / (7776*a^4*(a - b*x^2)^{(1/3)})$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^3,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^3 (-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^3,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 3*a)^3*(-b*x^2 + a)^(1/3)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{1}{3}} (bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^3,x)

[Out] int(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^3 (-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^3,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 3*a)^3*(-b*x^2 + a)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{1/3} (bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(1/3)*(3*a + b*x^2)^3),x)

[Out] int(1/((a - b*x^2)^(1/3)*(3*a + b*x^2)^3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-b*x**2+a)**(1/3)/(b*x**2+3*a)**3,x)
```

```
[Out] Integral(1/((a - b*x**2)**(1/3)*(3*a + b*x**2)**3), x)
```

$$3.129 \quad \int \frac{(3a+bx^2)^3}{(a-bx^2)^{4/3}} dx$$

Optimal. Leaf size=623

$$\frac{6696\sqrt{2}3^{3/4}a^{7/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{91bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] 2538/91*a*x*(-b*x^2+a)^(2/3)+81/13*x*(-b*x^2+a)^(2/3)*(b*x^2+3*a)+6*x*(b*x^2+3*a)^2/(-b*x^2+a)^(1/3)+20088/91*a^2*x/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))-6696/91*3^(3/4)*a^(7/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*2^(1/2)*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)+10044/91*3^(1/4)*a^(7/3)*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^(1/2)

Rubi [A] time = 0.43, antiderivative size = 623, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {413, 528, 388, 235, 304, 219, 1879}

$$\frac{20088a^2x}{91\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}\frac{6696\sqrt{2}3^{3/4}a^{7/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}F\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}}{(1-\sqrt{3})\sqrt[3]{a}}\right)\right)}{91bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

[In] Int[(3*a + b*x^2)^3/(a - b*x^2)^(4/3),x]

[Out] (2538*a*x*(a - b*x^2)^(2/3))/91 + (81*x*(a - b*x^2)^(2/3)*(3*a + b*x^2))/13 + (6*x*(3*a + b*x^2)^2)/(a - b*x^2)^(1/3) + (20088*a^2*x)/(91*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) + (10044*3^(1/4)*Sqrt[2 + Sqrt[3]]*a^(7/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(91*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)] - (6696*Sqrt[2]*3^(3/4)*a^(7/3)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]*EllipticF[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(91*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)])

Rule 219

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s

$*x + r^2*x^2)/((1 - \text{Sqrt}[3])*s + r*x)^2]*\text{EllipticF}[\text{ArcSin}[\frac{(1 + \text{Sqrt}[3])*s + r*x}{(1 - \text{Sqrt}[3])*s + r*x}], -7 + 4*\text{Sqrt}[3]]/(3^{1/4}*r*\text{Sqrt}[a + b*x^3])*\text{Sqrt}[-((s*(s + r*x))/((1 - \text{Sqrt}[3])*s + r*x)^2))], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 235

$\text{Int}[(a + (b_*)*(x_)^2)^{-1/3}, x_Symbol] := \text{Dist}[(3*\text{Sqrt}[b*x^2])/(2*b*x), \text{Subst}[\text{Int}[x/\text{Sqrt}[-a + x^3], x], x, (a + b*x^2)^{1/3}], x] /; \text{FreeQ}\{a, b\}, x]$

Rule 304

$\text{Int}[(x_)/\text{Sqrt}[(a_ + (b_*)*(x_)^3], x_Symbol] := \text{With}\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\text{Sqrt}[2]*s)/(\text{Sqrt}[2 - \text{Sqrt}[3]]*r), \text{Int}[1/\text{Sqrt}[a + b*x^3], x], x] + \text{Dist}[1/r, \text{Int}[\frac{(1 + \text{Sqrt}[3])*s + r*x}{\text{Sqrt}[a + b*x^3]}, x], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 388

$\text{Int}[(a + (b_*)*(x_)^{(n_)})^{(p_)*((c_ + (d_*)*(x_)^{(n_)})}, x_Symbol] := \text{Simp}[(d*x*(a + b*x^n)^{(p+1)})/(b*(n*(p+1)+1)), x] - \text{Dist}[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), \text{Int}[(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[n*(p+1)+1, 0]$

Rule 413

$\text{Int}[(a + (b_*)*(x_)^{(n_)})^{(p_)*((c_ + (d_*)*(x_)^{(n_)})^{(q_)}), x_Symbol] := \text{Simp}[(a*d - c*b)*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-1)}/(a*b*n*(p+1)), x] - \text{Dist}[1/(a*b*n*(p+1)), \text{Int}[(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+q)+1))*x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[p, -1] \&\& \text{GtQ}[q, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 528

$\text{Int}[(a + (b_*)*(x_)^{(n_)})^{(p_)*((c_ + (d_*)*(x_)^{(n_)})^{(q_)*((e_ + (f_*)*(x_)^{(n_)})}, x_Symbol] := \text{Simp}[(f*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^q]/(b*(n*(p+q+1)+1)), x] + \text{Dist}[1/(b*(n*(p+q+1)+1)), \text{Int}[(a + b*x^n)^p*(c + d*x^n)^{(q-1)}*\text{Simp}[c*(b*e - a*f + b*e*n*(p+q+1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p+q+1))*x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \text{GtQ}[q, 0] \&\& \text{NeQ}[n*(p+q+1)+1, 0]$

Rule 1879

$\text{Int}[(c + (d_*)*(x_))/\text{Sqrt}[(a + (b_*)*(x_)^3], x_Symbol] := \text{With}\{r = \text{Numer}[\text{Simplify}[\frac{(1 + \text{Sqrt}[3])*d}{c}], s = \text{Denom}[\text{Simplify}[\frac{(1 + \text{Sqrt}[3])*d}{c}]]\}, \text{Simp}[(2*d*s^3*\text{Sqrt}[a + b*x^3])/(a*r^2*((1 - \text{Sqrt}[3])*s + r*x)), x] + \text{Simp}[(3^{1/4}*\text{Sqrt}[2 + \text{Sqrt}[3]]*d*s*(s + r*x)*\text{Sqrt}[(s^2 - r*s*x + r^2*x^2)/((1 - \text{Sqrt}[3])*s + r*x)^2]*\text{EllipticE}[\text{ArcSin}[\frac{(1 + \text{Sqrt}[3])*s + r*x}{(1 - \text{Sqrt}[3])*s + r*x}], -7 + 4*\text{Sqrt}[3]]/(r^2*\text{Sqrt}[a + b*x^3]*\text{Sqrt}[-((s*(s + r*x))/((1 - \text{Sqrt}[3])*s + r*x)^2))], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NegQ}[a] \&\& \text{EqQ}[b*c^3 - 2*(5 + 3*\text{Sqrt}[3])*a*d^3, 0]$

Rubi steps

$$\begin{aligned}
\int \frac{(3a + bx^2)^3}{(a - bx^2)^{4/3}} dx &= \frac{6x(3a + bx^2)^2}{\sqrt[3]{a - bx^2}} - \frac{3 \int \frac{(3a + bx^2)(6a^2b + 18ab^2x^2)}{\sqrt[3]{a - bx^2}} dx}{2ab} \\
&= \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) + \frac{6x(3a + bx^2)^2}{\sqrt[3]{a - bx^2}} + \frac{9 \int \frac{-132a^3b^2 - 188a^2b^3x^2}{\sqrt[3]{a - bx^2}} dx}{26ab^2} \\
&= \frac{2538}{91}ax(a - bx^2)^{2/3} + \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) + \frac{6x(3a + bx^2)^2}{\sqrt[3]{a - bx^2}} - \frac{1}{91}(6696a^2) \int \frac{1}{\sqrt[3]{a - bx^2}} dx \\
&= \frac{2538}{91}ax(a - bx^2)^{2/3} + \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) + \frac{6x(3a + bx^2)^2}{\sqrt[3]{a - bx^2}} + \frac{(10044a^2\sqrt{-bx^2}) \operatorname{Su}}{\sqrt[3]{a - bx^2}} \\
&= \frac{2538}{91}ax(a - bx^2)^{2/3} + \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) + \frac{6x(3a + bx^2)^2}{\sqrt[3]{a - bx^2}} - \frac{(10044a^2\sqrt{-bx^2}) \operatorname{Su}}{\sqrt[3]{a - bx^2}} \\
&= \frac{2538}{91}ax(a - bx^2)^{2/3} + \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) + \frac{6x(3a + bx^2)^2}{\sqrt[3]{a - bx^2}} + \frac{20088a^2x}{91((1 - \sqrt{3})\sqrt[3]{a - bx^2})}
\end{aligned}$$

Mathematica [C] time = 5.06, size = 76, normalized size = 0.12

$$\frac{3x \left(2232a^2 \sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1 \left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a} \right) - 3051a^2 + 132abx^2 + 7b^2x^4 \right)}{91 \sqrt[3]{a - bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)^3/(a - b*x^2)^(4/3), x]

[Out] (-3*x*(-3051*a^2 + 132*a*b*x^2 + 7*b^2*x^4 + 2232*a^2*(1 - (b*x^2)/a)^(1/3)) *Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a])/(91*(a - b*x^2)^(1/3))

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\operatorname{integral} \left(\frac{(b^3x^6 + 9ab^2x^4 + 27a^2bx^2 + 27a^3)(-bx^2 + a)^{\frac{2}{3}}}{b^2x^4 - 2abx^2 + a^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(4/3), x, algorithm="fricas")

[Out] integral((b^3*x^6 + 9*a*b^2*x^4 + 27*a^2*b*x^2 + 27*a^3)*(-b*x^2 + a)^(2/3)/(b^2*x^4 - 2*a*b*x^2 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(4/3),x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^3/(-b*x^2 + a)^(4/3), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)^3/(-b*x^2+a)^(4/3),x)

[Out] int((b*x^2+3*a)^3/(-b*x^2+a)^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(4/3),x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^3/(-b*x^2 + a)^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + 3a)^3}{(a - bx^2)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)^3/(a - b*x^2)^(4/3),x)

[Out] int((3*a + b*x^2)^3/(a - b*x^2)^(4/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(3a + bx^2)^3}{(a - bx^2)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)**3/(-b*x**2+a)**(4/3),x)

[Out] Integral((3*a + b*x**2)**3/(a - b*x**2)**(4/3), x)

$$3.130 \quad \int \frac{(3a+bx^2)^2}{(a-bx^2)^{4/3}} dx$$

Optimal. Leaf size=592

$$108\sqrt{2}3^{3/4}a^{4/3} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} \operatorname{EllipticF} \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right), 4\sqrt{3} - 7 \right) - \frac{162\sqrt{3} \sqrt[3]{a} \sqrt[3]{a-bx^2}}{7bx \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}$$

[Out] $45/7*x*(-b*x^2+a)^{(2/3)}+6*x*(b*x^2+3*a)/(-b*x^2+a)^{(1/3)}+324/7*a*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))-108/7*3^{(3/4)}*a^{(4/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}+162/7*3^{(1/4)}*a^{(4/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/2)})^{(1/2)}$

Rubi [A] time = 0.37, antiderivative size = 592, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {413, 388, 235, 304, 219, 1879}

$$108\sqrt{2}3^{3/4}a^{4/3} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} F \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right) \middle| -7 + 4\sqrt{3} \right) - \frac{162\sqrt{3} \sqrt[3]{a} \sqrt[3]{a-bx^2}}{7bx \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(3*a + b*x^2)^2/(a - b*x^2)^{4/3}, x]$

[Out] $(45*x*(a - b*x^2)^{(2/3)})/7 + (6*x*(3*a + b*x^2))/(a - b*x^2)^{(1/3)} + (324*a*x)/(7*((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})) + (162*3^{(1/4)}*\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]]*a^{(4/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^{(1/2)}]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}]/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})], -7 + 4*\operatorname{Sqrt}[3]])/(7*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^{(1/2)}]) - (108*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(4/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^{(1/2)}]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}]/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})], -7 + 4*\operatorname{Sqrt}[3]])/(7*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^{(1/2)}])$

Rule 219

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] \rightarrow \operatorname{With}[\{r = \operatorname{Numer}[\operatorname{Rt}[b/a, 3]], s = \operatorname{Denom}[\operatorname{Rt}[b/a, 3]]\}, \operatorname{Simp}[(2*\operatorname{Sqrt}[2 - \operatorname{Sqrt}[3]]*(s + r*x)*\operatorname{Sqrt}[(s^2 - r*s*x + r^2*x^2)]/((1 - \operatorname{Sqrt}[3])*s + r*x)^2)*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 + \operatorname{Sqrt}[3])*s$


```
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(3^(1/4)*r*Sqrt[a + b*x^3]
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b}, x]
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sq
rt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 388

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 413

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p +
1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q -
2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p
+ q) + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,
0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/(
(1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(3a + bx^2)^2}{(a - bx^2)^{4/3}} dx &= \frac{6x(3a + bx^2)}{\sqrt[3]{a - bx^2}} - \frac{3 \int \frac{6a^2b + 10ab^2x^2}{\sqrt[3]{a - bx^2}} dx}{2ab} \\
&= \frac{45}{7}x(a - bx^2)^{2/3} + \frac{6x(3a + bx^2)}{\sqrt[3]{a - bx^2}} - \frac{1}{7}(108a) \int \frac{1}{\sqrt[3]{a - bx^2}} dx \\
&= \frac{45}{7}x(a - bx^2)^{2/3} + \frac{6x(3a + bx^2)}{\sqrt[3]{a - bx^2}} + \frac{(162a\sqrt{-bx^2}) \text{Subst}\left(\int \frac{x}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{7bx} \\
&= \frac{45}{7}x(a - bx^2)^{2/3} + \frac{6x(3a + bx^2)}{\sqrt[3]{a - bx^2}} - \frac{(162a\sqrt{-bx^2}) \text{Subst}\left(\int \frac{(1+\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{7bx} + \\
&= \frac{45}{7}x(a - bx^2)^{2/3} + \frac{6x(3a + bx^2)}{\sqrt[3]{a - bx^2}} + \frac{324ax}{7\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} + \frac{162\sqrt[4]{3}\sqrt{2 + \sqrt{3}}a^{4/3}}{7}
\end{aligned}$$

Mathematica [C] time = 5.05, size = 62, normalized size = 0.10

$$\frac{3x\left(36a\sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a}\right) - 57a + bx^2\right)}{7\sqrt[3]{a - bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)^2/(a - b*x^2)^(4/3), x]

[Out] (-3*x*(-57*a + b*x^2 + 36*a*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a]))/(7*(a - b*x^2)^(1/3))

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b^2x^4 + 6abx^2 + 9a^2)(-bx^2 + a)^{2/3}}{b^2x^4 - 2abx^2 + a^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(4/3), x, algorithm="fricas")

[Out] integral((b^2*x^4 + 6*a*b*x^2 + 9*a^2)*(-b*x^2 + a)^(2/3)/(b^2*x^4 - 2*a*b*x^2 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^2}{(-bx^2 + a)^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(4/3), x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^2/(-b*x^2 + a)^(4/3), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^2}{(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)^2/(-b*x^2+a)^(4/3), x)

[Out] int((b*x^2+3*a)^2/(-b*x^2+a)^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^2}{(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(4/3), x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^2/(-b*x^2 + a)^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + 3a)^2}{(a - bx^2)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)^2/(a - b*x^2)^(4/3), x)

[Out] int((3*a + b*x^2)^2/(a - b*x^2)^(4/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(3a + bx^2)^2}{(a - bx^2)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)**2/(-b*x**2+a)**(4/3), x)

[Out] Integral((3*a + b*x**2)**2/(a - b*x**2)**(4/3), x)

$$3.131 \quad \int \frac{3a+bx^2}{(a-bx^2)^{4/3}} dx$$

Optimal. Leaf size=561

$$\frac{3\sqrt{2}3^{3/4}\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{9\sqrt[4]{3}} + \frac{bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}{1}$$

[Out] $6*x/(-b*x^2+a)^{(1/3)}+9*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))-3*3^{(3/4)}*a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}+9/2*3^{(1/4)}*a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.30, antiderivative size = 561, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {385, 235, 304, 219, 1879}

$$\frac{3\sqrt{2}3^{3/4}\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}F\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right)\right)-7+4\sqrt{3}}{9\sqrt[4]{3}\sqrt{2}} + \frac{bx\sqrt{\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}{1}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(3*a + b*x^2)/(a - b*x^2)^{(4/3)}, x]$

[Out] $(6*x)/(a - b*x^2)^{(1/3)} + (9*x)/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}) + (9*3^{(1/4)}*\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]]*a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}))], -7 + 4*\operatorname{Sqrt}[3]])/(2*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]) - (3*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}))], -7 + 4*\operatorname{Sqrt}[3]))/(b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])$

Rule 219

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] := \operatorname{With}[\{r = \operatorname{Numer}[\operatorname{Rt}[b/a, 3]], s = \operatorname{Denom}[\operatorname{Rt}[b/a, 3]]\}, \operatorname{Simp}[(2*\operatorname{Sqrt}[2 - \operatorname{Sqrt}[3]]*(s + r*x)*\operatorname{Sqrt}[(s^2 - r*s*x + r^2*x^2)/((1 - \operatorname{Sqrt}[3])*s + r*x)^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*s + r*x)/((1 - \operatorname{Sqrt}[3])*s + r*x))], -7 + 4*\operatorname{Sqrt}[3]])/(3^{(1/4)}*r*\operatorname{Sqrt}[a + b*x^3]*\operatorname{Sqrt}[-((s*(s + r*x))/((1 - \operatorname{Sqrt}[3])*s + r*x)^2))], x]] /; \operatorname{FreeQ}[\{a, b\}, x]$

] && NegQ[a]

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x),
Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sqrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[
((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]],
s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)]], -7 + 4*Sqrt[3])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\int \frac{3a + bx^2}{(a - bx^2)^{4/3}} dx = \frac{6x}{\sqrt[3]{a - bx^2}} - 3 \int \frac{1}{\sqrt[3]{a - bx^2}} dx$$

$$= \frac{6x}{\sqrt[3]{a - bx^2}} + \frac{(9\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{x}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{2bx}$$

$$= \frac{6x}{\sqrt[3]{a - bx^2}} - \frac{(9\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{(1+\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{2bx} + \frac{(9\sqrt{\frac{1}{2}(2 + \sqrt{3})} \sqrt[3]{a} \sqrt{-bx^2})}{2bx}$$

$$= \frac{6x}{\sqrt[3]{a - bx^2}} + \frac{9x}{(1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2}} + \frac{9\sqrt[4]{3} \sqrt{2 + \sqrt{3}} \sqrt[3]{a} (\sqrt[3]{a} - \sqrt[3]{a - bx^2}) \sqrt{\frac{a^{2/3} + \sqrt[3]{a}}{(1 - \sqrt{3})}}}{2bx \sqrt{-bx^2}}$$

Mathematica [C] time = 0.02, size = 53, normalized size = 0.09

$$\frac{6x - 3x\sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a}\right)}{\sqrt[3]{a - bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)/(a - b*x^2)^(4/3), x]

[Out] (6*x - 3*x*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a])/ (a - b*x^2)^(1/3)

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(bx^2 + 3a)(-bx^2 + a)^{\frac{2}{3}}}{b^2x^4 - 2abx^2 + a^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(4/3), x, algorithm="fricas")

[Out] integral((b*x^2 + 3*a)*(-b*x^2 + a)^(2/3)/(b^2*x^4 - 2*a*b*x^2 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(4/3), x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)/(-b*x^2 + a)^(4/3), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)/(-b*x^2+a)^(4/3), x)

[Out] int((b*x^2+3*a)/(-b*x^2+a)^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(4/3), x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)/(-b*x^2 + a)^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{bx^2 + 3a}{(a - bx^2)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)/(a - b*x^2)^(4/3), x)

[Out] `int((3*a + b*x^2)/(a - b*x^2)^(4/3), x)`

sympy [A] time = 6.87, size = 60, normalized size = 0.11

$$\frac{3x {}_2F_1\left(\frac{1}{2}, \frac{4}{3} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{\sqrt[3]{a}} + \frac{bx^3 {}_2F_1\left(\frac{4}{3}, \frac{3}{2} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{3a^{\frac{4}{3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+3*a)/(-b*x**2+a)**(4/3), x)`

[Out] `3*x*hyper((1/2, 4/3), (3/2,), b*x**2*exp_polar(2*I*pi)/a)/a**(1/3) + b*x**3*hyper((4/3, 3/2), (5/2,), b*x**2*exp_polar(2*I*pi)/a)/(3*a**(4/3))`

3.132 $\int \frac{1}{(a-bx^2)^{4/3}(3+bx^2)} dx$

Optimal. Leaf size=776

$$\frac{3^{3/4} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} \operatorname{EllipticF} \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right), 4\sqrt{3} - 7 \right) \tan^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a}}{\dots} \right)}{4\sqrt{2} a^{5/3} bx \sqrt{-\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} + \frac{\dots}{8 \cdot 2^{2/3} \sqrt{\dots}}$$

[Out] 3/8*x/a^2/(-b*x^2+a)^(1/3)+3/8*x/a^2/(-(-b*x^2+a)^(1/3)+a^(1/3))*(1-3^(1/2)))+1/16*arctanh(x*b^(1/2)/a^(1/6)/(a^(1/3)+2^(1/3)*(-b*x^2+a)^(1/3))) * 2^(1/3)/a^(11/6)/b^(1/2)-1/48*arctanh(x*b^(1/2)/a^(1/2)) * 2^(1/3)/a^(11/6)/b^(1/2)+1/48*arctan(a^(1/6)*(a^(1/3)-2^(1/3)*(-b*x^2+a)^(1/3)) * 3^(1/2)/x/b^(1/2)) * 2^(1/3)/a^(11/6) * 3^(1/2)/b^(1/2)+1/48*arctan(3^(1/2)*a^(1/2)/x/b^(1/2)) * 2^(1/3)/a^(11/6) * 3^(1/2)/b^(1/2)-1/8*(a^(1/3)-(-b*x^2+a)^(1/3)) * EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))) , 2*I-I*3^(1/2)) * ((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2) * 3^(3/4)/a^(5/3)/b/x * 2^(1/2)/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)+3/16*(a^(1/3)-(-b*x^2+a)^(1/3)) * EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))) , 2*I-I*3^(1/2)) * ((a^(2/3)+a^(1/3)*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2) * (1/2*6^(1/2)+1/2*2^(1/2)) * 3^(1/4)/a^(5/3)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))))^2)^(1/2)

Rubi [A] time = 0.44, antiderivative size = 776, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 24, number of rules / integrand size = 0.292, Rules used = {414, 530, 235, 304, 219, 1879, 393}

$$\frac{3x}{8a^2 \sqrt[3]{a-bx^2}} + \frac{3x}{8a^2 \left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)} + \frac{\tan^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2} \right)}{\sqrt{bx}} \right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tanh^{-1} \left(\frac{\sqrt{bx}}{\sqrt[6]{a} \left(\sqrt[3]{2} \sqrt[3]{a-bx^2} + \sqrt[3]{a} \right)} \right)}{8 \cdot 2^{2/3} a^{11/6} \sqrt{b}} - \frac{3^{3/4}}{\dots}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(4/3)*(3*a + b*x^2)), x]

[Out] (3*x)/(8*a^2*(a - b*x^2)^(1/3)) + (3*x)/(8*a^2*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) + ArcTan[(Sqrt[3]*Sqrt[a])/(Sqrt[b]*x)]/(8*2^(2/3)*Sqrt[3]*a^(11/6)*Sqrt[b]) + ArcTan[(Sqrt[3]*a^(1/6)*(a^(1/3) - 2^(1/3)*(a - b*x^2)^(1/3)))/(Sqrt[b]*x)]/(8*2^(2/3)*Sqrt[3]*a^(11/6)*Sqrt[b]) - ArcTanh[(Sqrt[b]*x)/Sqrt[a]]/(24*2^(2/3)*a^(11/6)*Sqrt[b]) + ArcTanh[(Sqrt[b]*x)/(a^(1/6)*(a^(1/3) + 2^(1/3)*(a - b*x^2)^(1/3)))]/(8*2^(2/3)*a^(11/6)*Sqrt[b]) + (3*3^(1/4)*Sqrt[2 + Sqrt[3]]*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])/(16*a^(5/3)*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)]) - (3^(3/4)*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2]*EllipticF[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3]])

$$\frac{1}{(4\sqrt{2}a^{5/3}bx\sqrt{-(a^{1/3}(a^{1/3} - (a - bx^2)^{1/3})) / ((1 - \sqrt{3})a^{1/3} - (a - bx^2)^{1/3})^2})}$$

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x
), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 393

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Wit
h[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^
(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2
)^(1/3)])/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(
1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)
))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]]) /; FreeQ[{a, b, c, d}
, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]
```

Rule 414

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c -
a*d), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d), Int[(a + b*x^n)^(p + 1)*(c +
d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x]
, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1]
&& !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 530

```
Int((((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p
, n}, x]
```

Rule 1879

```
Int(((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((
1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
```

)/((1 - Sqrt[3])*s + r*x)^2)), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a - bx^2)^{4/3} (3a + bx^2)} dx &= \frac{3x}{8a^2 \sqrt[3]{a - bx^2}} + \frac{3 \int \frac{\frac{ab}{3} - \frac{b^2 x^2}{3}}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx}{8a^2 b} \\ &= \frac{3x}{8a^2 \sqrt[3]{a - bx^2}} - \frac{\int \frac{1}{\sqrt[3]{a - bx^2}} dx}{8a^2} + \frac{\int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx}{4a} \\ &= \frac{3x}{8a^2 \sqrt[3]{a - bx^2}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{b}x}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{b}x}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}}{\sqrt{a}}\right)}{24 \cdot 2^{2/3} a^{11/6}} \\ &= \frac{3x}{8a^2 \sqrt[3]{a - bx^2}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{b}x}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a} - \sqrt[3]{2}\sqrt[3]{a - bx^2}\right)}{\sqrt{b}x}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}}{\sqrt{a}}\right)}{24 \cdot 2^{2/3} a^{11/6}} \\ &= \frac{3x}{8a^2 \sqrt[3]{a - bx^2}} + \frac{3x}{8a^2 \left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{b}x}\right)}{8 \cdot 2^{2/3} \sqrt{3} a^{11/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{b}}{\sqrt{a}}\right)}{8 \cdot 2^{2/3} a^{11/6}} \end{aligned}$$

Mathematica [C] time = 0.14, size = 226, normalized size = 0.29

$$x \left(\frac{27 \left(\frac{1}{a^2} - \frac{3F_1\left(\frac{1}{2}, \frac{1}{3}, 1; \frac{3}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{(3a + bx^2) \left(2bx^2 \left(F_1\left(\frac{3}{2}, \frac{4}{3}, 1; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}, \frac{1}{3}, 2; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}, \frac{1}{3}, 1; \frac{3}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right)} \right)}{72 \sqrt[3]{a - bx^2}} - \frac{bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} F_1\left(\frac{3}{2}, \frac{1}{3}, 1; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{a^3} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(4/3)*(3*a + b*x^2)), x]

[Out] (x*(-((b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])/a^3) + 27*(a^(-2) - (3*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/((3*a + b*x^2)*(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))))/((72*(a - b*x^2)^(1/3)))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 3*a)*(-b*x^2 + a)^(4/3)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{4}{3}}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a),x)

[Out] int(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)(-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 3*a)*(-b*x^2 + a)^(4/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{\frac{4}{3}}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(4/3)*(3*a + b*x^2)),x)

[Out] int(1/((a - b*x^2)^(4/3)*(3*a + b*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a - bx^2)^{\frac{4}{3}}(3a + bx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2+a)**(4/3)/(b*x**2+3*a),x)

[Out] Integral(1/((a - b*x**2)**(4/3)*(3*a + b*x**2)), x)

$$3.133 \quad \int \frac{1}{(a-bx^2)^{4/3}(3+bx^2)^2} dx$$

Optimal. Leaf size=807

$$\frac{x}{12a^3\sqrt[3]{a-bx^2}} + \frac{x}{24a^2\sqrt[3]{a-bx^2}(bx^2+3a)} + \frac{x}{12a^3\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{16\cdot 2^{2/3}\sqrt{3}a^{17/6}\sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}}{\sqrt{bx}}\right)}{16\cdot 2^{2/3}\sqrt{3}a^{17/6}\sqrt{b}}$$

[Out] $\frac{1}{12}x/a^{3/2}/(-b*x^2+a)^{(1/3)}+1/24*x/a^2/(-b*x^2+a)^{(1/3)}/(b*x^2+3*a)+1/12*x/a^{3/2}/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))+1/32*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/(a^{(1/3)}+2^{(1/3)}*(-b*x^2+a)^{(1/3)}))*2^{(1/3)}/a^{(17/6)}/b^{(1/2)}-1/96*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/2)})*2^{(1/3)}/a^{(17/6)}/b^{(1/2)}+1/96*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(-b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})*2^{(1/3)}/a^{(17/6)}*3^{(1/2)}/b^{(1/2)}+1/96*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})*2^{(1/3)}/a^{(17/6)}*3^{(1/2)}/b^{(1/2)}-1/36*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)})*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*3^{(3/4)}/a^{(8/3)}/b/x*2^{(1/2)}/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}+1/24*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})*3^{(1/4)}/a^{(8/3)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.53, antiderivative size = 807, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {414, 527, 530, 235, 304, 219, 1879, 393}

$$\frac{x}{12a^3\sqrt[3]{a-bx^2}} + \frac{x}{24a^2\sqrt[3]{a-bx^2}(bx^2+3a)} + \frac{x}{12a^3\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{16\cdot 2^{2/3}\sqrt{3}a^{17/6}\sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}}{\sqrt{bx}}\right)}{16\cdot 2^{2/3}\sqrt{3}a^{17/6}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(4/3)*(3*a + b*x^2)^2), x]

[Out] $x/(12*a^3*(a-b*x^2)^{(1/3)})+x/(24*a^2*(a-b*x^2)^{(1/3)}*(3*a+b*x^2))+x/(12*a^3*((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))+\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(16*2^{(2/3)}*\operatorname{Sqrt}[3]*a^{(17/6)}*\operatorname{Sqrt}[b])+\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(a-b*x^2)^{(1/3)}))/(\operatorname{Sqrt}[b]*x)]/(16*2^{(2/3)}*\operatorname{Sqrt}[3]*a^{(17/6)}*\operatorname{Sqrt}[b])-\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(48*2^{(2/3)}*a^{(17/6)}*\operatorname{Sqrt}[b])+\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{(1/6)}*(a^{(1/3)}+2^{(1/3)}*(a-b*x^2)^{(1/3)}))]/(16*2^{(2/3)}*a^{(17/6)}*\operatorname{Sqrt}[b])+(\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})],-7+4*\operatorname{Sqrt}[3)]/(8*3^{(3/4)}*a^{(8/3)}*b*x*\operatorname{Sqrt}[-(a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2])-(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})]$

$b*x^2)^{(1/3)}], -7 + 4*\text{Sqrt}[3]]/(6*\text{Sqrt}[2]*3^{(1/4)}*a^{(8/3)}*b*x*\text{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})))/((1 - \text{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3}))^2])]$

Rule 219

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] := \text{With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, \text{Simp}[(2*\text{Sqrt}[2 - \text{Sqrt}[3]]*(s + r*x)*\text{Sqrt}[(s^2 - r*s*x + r^2*x^2)/((1 - \text{Sqrt}[3])*s + r*x)^2]*\text{EllipticF}[\text{ArcSin}[(1 + \text{Sqrt}[3])*s + r*x]/((1 - \text{Sqrt}[3])*s + r*x)], -7 + 4*\text{Sqrt}[3]]/(3^{(1/4)}*r*\text{Sqrt}[a + b*x^3]*\text{Sqrt}[-((s*(s + r*x))/((1 - \text{Sqrt}[3])*s + r*x)^2)]), x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 235

$\text{Int}[(a_) + (b_)*(x_)^2)^{-1/3}, x_Symbol] := \text{Dist}[(3*\text{Sqrt}[b*x^2])/(2*b*x), \text{Subst}[\text{Int}[x/\text{Sqrt}[-a + x^3], x], x, (a + b*x^2)^{(1/3)}], x] /; \text{FreeQ}[\{a, b\}, x]$

Rule 304

$\text{Int}[(x_)/\text{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] := \text{With}[\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\text{Sqrt}[2]*s)/(\text{Sqrt}[2 - \text{Sqrt}[3]]*r), \text{Int}[1/\text{Sqrt}[a + b*x^3], x], x] + \text{Dist}[1/r, \text{Int}[(1 + \text{Sqrt}[3])*s + r*x]/\text{Sqrt}[a + b*x^3], x], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 393

$\text{Int}[1/(((a_) + (b_)*(x_)^2)^{(1/3)}*((c_) + (d_)*(x_)^2)), x_Symbol] := \text{With}[\{q = \text{Rt}[-(b/a), 2]\}, \text{Simp}[(q*\text{ArcTan}[\text{Sqrt}[3]/(q*x)])/(2*2^{(2/3)}*\text{Sqrt}[3]*a^{(1/3)*d}), x] + (\text{Simp}[(q*\text{ArcTanh}[(a^{(1/3)}*q*x)/(a^{(1/3)} + 2^{(1/3)}*(a + b*x^2)^{(1/3)})]/(2*2^{(2/3)}*a^{(1/3)*d}), x] - \text{Simp}[(q*\text{ArcTanh}[q*x])/(6*2^{(2/3)}*a^{(1/3)*d}), x] + \text{Simp}[(q*\text{ArcTan}[(\text{Sqrt}[3]*(a^{(1/3)} - 2^{(1/3)}*(a + b*x^2)^{(1/3)})]/(a^{(1/3)}*q*x))]/(2*2^{(2/3)}*\text{Sqrt}[3]*a^{(1/3)*d}), x]]) /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[b*c + 3*a*d, 0] \&\& \text{NegQ}[b/a]$

Rule 414

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}])^{(p_)}*((c_) + (d_)*(x_)^{(n_)}])^{(q_)}, x_Symbol] := -\text{Simp}[(b*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q+1)})/(a*n*(p+1)*(b*c - a*d)), x] + \text{Dist}[1/(a*n*(p+1)*(b*c - a*d)), \text{Int}[(a + b*x^n)^{(p+1)}*(c + d*x^n)^q*\text{Simp}[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, n, q\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[p, -1] \&\& !(IntegerQ[p] \&\& IntegerQ[q] \&\& \text{LtQ}[q, -1]) \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 527

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}])^{(p_)}*((c_) + (d_)*(x_)^{(n_)}])^{(q_)}*((e_) + (f_)*(x_)^{(n_)}), x_Symbol] := -\text{Simp}[(b*e - a*f)*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q+1)})/(a*n*(b*c - a*d)*(p+1)), x] + \text{Dist}[1/(a*n*(b*c - a*d)*(p+1)), \text{Int}[(a + b*x^n)^{(p+1)}*(c + d*x^n)^q*\text{Simp}[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2)+1)*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, q\}, x] \&\& \text{LtQ}[p, -1]$

Rule 530

$\text{Int}[(a_) + (b_)*(x_)^{(n_)}])^{(p_)}*((e_) + (f_)*(x_)^{(n_)}))/((c_) + (d_)*(x_)^{(n_)}), x_Symbol] := \text{Dist}[f/d, \text{Int}[(a + b*x^n)^p, x], x] + \text{Dist}[(d*e - c*f)/d, \text{Int}[(a + b*x^n)^p/(c + d*x^n), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, p\}, x]$

, n}, x]

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)]], -7 + 4*Sqrt[3])]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/(1 - Sqrt[3])*s + r*x)^2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\int \frac{1}{(a - bx^2)^{4/3} (3a + bx^2)^2} dx = \frac{x}{24a^2 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{\int \frac{-7ab + \frac{5b^2x^2}{3}}{(a - bx^2)^{4/3} (3a + bx^2)} dx}{24a^2b}$$

$$= \frac{x}{12a^3 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{\int \frac{-\frac{8}{3}a^2b^2 + \frac{16}{9}ab^3x^2}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx}{64a^4b^2}$$

$$= \frac{x}{12a^3 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{\int \frac{1}{\sqrt[3]{a - bx^2}} dx}{36a^3} + \frac{\int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx}{8a^2}$$

$$= \frac{x}{12a^3 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 \sqrt[3]{a - bx^2} (3a + bx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{16 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{16 \cdot 2^{2/3}}$$

$$= \frac{x}{12a^3 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 \sqrt[3]{a - bx^2} (3a + bx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{16 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{16 \cdot 2^{2/3}}$$

$$= \frac{x}{12a^3 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 \sqrt[3]{a - bx^2} (3a + bx^2)} + \frac{x}{12a^3 \left((1 - \sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a - bx^2} \right)}$$

Mathematica [C] time = 0.18, size = 236, normalized size = 0.29

$$x \left(\frac{27a \left(\frac{9a^2 F_1\left(\frac{1}{2}, \frac{1}{3}, 1; \frac{3}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{2bx^2 \left(F_1\left(\frac{3}{2}, \frac{4}{3}, 1; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}, \frac{1}{3}, 2; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}, \frac{1}{3}, 1; \frac{3}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) + 7a + 2bx^2 \right)}{3a + bx^2} - 2bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} F_1\left(\frac{3}{2}, \frac{1}{3}, 1; \frac{5}{2}, \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) \frac{1}{648a^4 \sqrt[3]{a - bx^2}}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[1/((a - b*x^2)^(4/3)*(3*a + b*x^2)^2), x]
```

```
[Out] (x*(-2*b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + (27*a*(7*a + 2*b*x^2 + (9*a^2*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a]))/(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))) / (3*a + b*x^2)) / (648*a^4*(a - b*x^2)^(1/3))
```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^2,x, algorithm="fricas")
```

```
[Out] Timed out
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^2 (-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^2,x, algorithm="giac")
```

```
[Out] integrate(1/((b*x^2 + 3*a)^2*(-b*x^2 + a)^(4/3)), x)
```

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{4}{3}} (bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^2,x)
```

```
[Out] int(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^2,x)
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^2 (-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^2,x, algorithm="maxima")
```

```
[Out] integrate(1/((b*x^2 + 3*a)^2*(-b*x^2 + a)^(4/3)), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{\frac{4}{3}} (bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((a - b*x^2)^(4/3)*(3*a + b*x^2)^2),x)
```

```
[Out] int(1/((a - b*x^2)^(4/3)*(3*a + b*x^2)^2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a - bx^2)^{\frac{4}{3}} (3a + bx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2+a)**(4/3)/(b*x**2+3*a)**2,x)

[Out] Integral(1/((a - b*x**2)**(4/3)*(3*a + b*x**2)**2), x)

$$3.134 \quad \int \frac{1}{(a-bx^2)^{4/3} (3a+bx^2)^3} dx$$

Optimal. Leaf size=849

$$\frac{19(a-bx^2)^{2/3}x}{1152a^4(bx^2+3a)} + \frac{17x}{192a^3\sqrt[3]{a-bx^2}(bx^2+3a)} + \frac{19x}{1152a^4\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{x}{48a^2\sqrt[3]{a-bx^2}(bx^2+3a)}$$

[Out] 1/48*x/a^2/(-b*x^2+a)^(1/3)/(b*x^2+3*a)^2+17/192*x/a^3/(-b*x^2+a)^(1/3)/(b*x^2+3*a)-19/1152*x*(-b*x^2+a)^(2/3)/a^4/(b*x^2+3*a)+19/1152*x/a^4/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))+7/576*arctanh(x*b^(1/2)/a^(1/6))/(a^(1/3)+2^(1/3)*(-b*x^2+a)^(1/3))*2^(1/3)/a^(23/6)/b^(1/2)-7/1728*arctanh(x*b^(1/2)/a^(1/2))*2^(1/3)/a^(23/6)/b^(1/2)+7/1728*arctan(a^(1/6)*(a^(1/3)-2^(1/3))*(-b*x^2+a)^(1/3))*3^(1/2)/x/b^(1/2))*2^(1/3)/a^(23/6)*3^(1/2)/b^(1/2)+7/1728*arctan(3^(1/2)*a^(1/2)/x/b^(1/2))*2^(1/3)/a^(23/6)*3^(1/2)/b^(1/2)-19/3456*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticF((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3))*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)*3^(3/4)/a^(11/3)/b/x*2^(1/2)/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)+19/2304*(a^(1/3)-(-b*x^2+a)^(1/3))*EllipticE((-(-b*x^2+a)^(1/3)+a^(1/3)*(1+3^(1/2)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2))),2*I-I*3^(1/2))*((a^(2/3)+a^(1/3))*(-b*x^2+a)^(1/3)+(-b*x^2+a)^(2/3))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)*(1/2*6^(1/2)+1/2*2^(1/2))*3^(1/4)/a^(11/3)/b/x/(-a^(1/3)*(a^(1/3)-(-b*x^2+a)^(1/3)))/(-(-b*x^2+a)^(1/3)+a^(1/3)*(1-3^(1/2)))^2)^(1/2)

Rubi [A] time = 0.65, antiderivative size = 849, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {414, 527, 530, 235, 304, 219, 1879, 393}

$$\frac{19(a-bx^2)^{2/3}x}{1152a^4(bx^2+3a)} + \frac{17x}{192a^3\sqrt[3]{a-bx^2}(bx^2+3a)} + \frac{19x}{1152a^4\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{x}{48a^2\sqrt[3]{a-bx^2}(bx^2+3a)}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(4/3)*(3*a + b*x^2)^3),x]

[Out] x/(48*a^2*(a - b*x^2)^(1/3)*(3*a + b*x^2)^2) + (17*x)/(192*a^3*(a - b*x^2)^(1/3)*(3*a + b*x^2)) - (19*x*(a - b*x^2)^(2/3))/(1152*a^4*(3*a + b*x^2)) + (19*x)/(1152*a^4*((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))) + (7*ArcTan[(Sqrt[3]*Sqrt[a])/(Sqrt[b]*x)]/(288*2^(2/3)*Sqrt[3]*a^(23/6)*Sqrt[b]) + (7*ArcTan[(Sqrt[3]*a^(1/6)*(a^(1/3) - 2^(1/3)*(a - b*x^2)^(1/3)))/(Sqrt[b]*x)]/(288*2^(2/3)*Sqrt[3]*a^(23/6)*Sqrt[b]) - (7*ArcTanh[(Sqrt[b]*x)/Sqrt[a]]/(864*2^(2/3)*a^(23/6)*Sqrt[b]) + (7*ArcTanh[(Sqrt[b]*x)/(a^(1/6)*(a^(1/3) + 2^(1/3)*(a - b*x^2)^(1/3)))]/(288*2^(2/3)*a^(23/6)*Sqrt[b]) + (19*Sqrt[2 + Sqrt[3]]*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) + a^(1/3)*(a - b*x^2)^(1/3) + (a - b*x^2)^(2/3))]/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)*EllipticE[ArcSin[((1 + Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))], -7 + 4*Sqrt[3])]/(768*3^(3/4)*a^(11/3)*b*x*Sqrt[-((a^(1/3)*(a^(1/3) - (a - b*x^2)^(1/3)))/((1 - Sqrt[3])*a^(1/3) - (a - b*x^2)^(1/3))^2)]) - (19*(a^(1/3) - (a - b*x^2)^(1/3))*Sqrt[(a^(2/3) +

$$a^{1/3}*(a - b*x^2)^{1/3} + (a - b*x^2)^{2/3}/((1 - \text{Sqrt}[3])*a^{1/3} - (a - b*x^2)^{1/3})^2*\text{EllipticF}[\text{ArcSin}[(1 + \text{Sqrt}[3])*a^{1/3} - (a - b*x^2)^{1/3}]/((1 - \text{Sqrt}[3])*a^{1/3} - (a - b*x^2)^{1/3})], -7 + 4*\text{Sqrt}[3]]/(576*\text{Sqrt}[2]*3^{1/4}*a^{11/3}*b*x*\text{Sqrt}[-((a^{1/3}*(a^{1/3} - (a - b*x^2)^{1/3}))/((1 - \text{Sqrt}[3])*a^{1/3} - (a - b*x^2)^{1/3})^2)])$$

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]]/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sq
rt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 393

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Wit
h[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)]/(2*2^(2/3)*Sqrt[3]*a^
(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2
)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^
(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)
))/(a^(1/3)*q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]]) /; FreeQ[{a, b, c, d}
, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]
```

Rule 414

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c -
a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c +
d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x]
, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1]
&& !( !IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 530

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*
```

$(x_)^{(n_)}$, x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1879

Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a - bx^2)^{4/3} (3a + bx^2)^3} dx &= \frac{x}{48a^2 \sqrt[3]{a - bx^2} (3a + bx^2)^2} - \frac{\int \frac{-15ab + \frac{11b^2x^2}{3}}{(a - bx^2)^{4/3} (3a + bx^2)^2} dx}{48a^2b} \\
 &= \frac{x}{48a^2 \sqrt[3]{a - bx^2} (3a + bx^2)^2} + \frac{17x}{192a^3 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{\int \frac{-6a^2b^2 - \frac{170}{9}ab^3x^2}{\sqrt[3]{a - bx^2} (3a + bx^2)^2} dx}{128a^4b^2} \\
 &= \frac{x}{48a^2 \sqrt[3]{a - bx^2} (3a + bx^2)^2} + \frac{17x}{192a^3 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{19x(a - bx^2)^{2/3}}{1152a^4 (3a + bx^2)^2} \\
 &= \frac{x}{48a^2 \sqrt[3]{a - bx^2} (3a + bx^2)^2} + \frac{17x}{192a^3 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{19x(a - bx^2)^{2/3}}{1152a^4 (3a + bx^2)^2} \\
 &= \frac{x}{48a^2 \sqrt[3]{a - bx^2} (3a + bx^2)^2} + \frac{17x}{192a^3 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{19x(a - bx^2)^{2/3}}{1152a^4 (3a + bx^2)^2} \\
 &= \frac{x}{48a^2 \sqrt[3]{a - bx^2} (3a + bx^2)^2} + \frac{17x}{192a^3 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{19x(a - bx^2)^{2/3}}{1152a^4 (3a + bx^2)^2} \\
 &= \frac{x}{48a^2 \sqrt[3]{a - bx^2} (3a + bx^2)^2} + \frac{17x}{192a^3 \sqrt[3]{a - bx^2} (3a + bx^2)} - \frac{19x(a - bx^2)^{2/3}}{1152a^4 (3a + bx^2)^2}
 \end{aligned}$$

Mathematica [C] time = 0.24, size = 256, normalized size = 0.30

$$x \left(\frac{27a \left(\frac{333a^2(3a+bx^2)F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9aF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) + 273a^2 + 140abx^2 + 19b^2x^4}{(3a+bx^2)^2} \right) - 19bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} F_1\left(\frac{3}{2}; \frac{1}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right)}{31104a^5 \sqrt[3]{a - bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(4/3)*(3*a + b*x^2)^3), x]

[Out] (x*(-19*b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + (27*a*(273*a^2 + 140*a*b*x^2 + 19*b^2*x^4 + (333*a^2*(3*a + b*x^2)*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])))/(3*a + b*x^2)^2))/(31104*a^5*(a - b*x^2)^(1/3))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^3,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^3 (-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^3,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 3*a)^3*(-b*x^2 + a)^(4/3)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{4}{3}} (bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^3,x)

[Out] int(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^3 (-bx^2 + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(4/3)/(b*x^2+3*a)^3,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 3*a)^3*(-b*x^2 + a)^(4/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{4/3} (bx^2 + 3a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(4/3)*(3*a + b*x^2)^3), x)

[Out] int(1/((a - b*x^2)^(4/3)*(3*a + b*x^2)^3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a - bx^2)^{4/3} (3a + bx^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2+a)**(4/3)/(b*x**2+3*a)**3, x)

[Out] Integral(1/((a - b*x**2)**(4/3)*(3*a + b*x**2)**3), x)

3.135 $\int \frac{(3a+bx^2)^4}{(a-bx^2)^{7/3}} dx$

Optimal. Leaf size=653

$$\frac{12312\sqrt{2}3^{3/4}a^{7/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{91bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

[Out] $-3240/91*a*x*(-b*x^2+a)^{(2/3)}-81/13*x*(-b*x^2+a)^{(2/3)}*(b*x^2+3*a)-9/2*x*(b*x^2+3*a)^2/(-b*x^2+a)^{(1/3)}+3/2*x*(b*x^2+3*a)^3/(-b*x^2+a)^{(4/3)}-36936/91*a^2*x/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+12312/91*3^{(3/4)}*a^{(7/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}-18468/91*3^{(1/4)}*a^{(7/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.51, antiderivative size = 653, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {413, 526, 528, 388, 235, 304, 219, 1879}

$$\frac{36936a^2x}{91\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{12312\sqrt{2}3^{3/4}a^{7/3}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)\sqrt{\frac{a^{2/3}+\sqrt[3]{a}\sqrt[3]{a-bx^2}+(a-bx^2)^{2/3}}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}F\left(\sin^{-1}\left(\frac{(1+\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}{(1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}}\right),4\sqrt{3}-7\right)}{91bx\sqrt{-\frac{\sqrt[3]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)^2}}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(3a + b*x^2)^4/(a - b*x^2)^{(7/3)}, x]$

[Out] $(-3240*a*x*(a - b*x^2)^{(2/3)}/91 - (81*x*(a - b*x^2)^{(2/3)}*(3*a + b*x^2))/13 - (9*x*(3*a + b*x^2)^2)/(2*(a - b*x^2)^{(1/3)}) + (3*x*(3*a + b*x^2)^3)/(2*(a - b*x^2)^{(4/3)}) - (36936*a^2*x)/(91*((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})) - (18468*3^{(1/4)}*\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]]*a^{(7/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}))], -7 + 4*\operatorname{Sqrt}[3])]/(91*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])) + (12312*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(7/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)}))], -7 + 4*\operatorname{Sqrt}[3])]/(91*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]))$

Rule 219

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_)*(x_)^3], x_Symbol] := \operatorname{With}[\{r = \operatorname{Numer}[\operatorname{Rt}[b/a, 3]], s = \operatorname{Denom}[\operatorname{Rt}[b/a, 3]]\}, \operatorname{Simp}[(2*\operatorname{Sqrt}[2 - \operatorname{Sqrt}[3]]*(s + r*x)*\operatorname{Sqrt}[(s^2 - r*s$

$*x + r^2*x^2)/((1 - \text{Sqrt}[3])*s + r*x)^2]*\text{EllipticF}[\text{ArcSin}[\frac{(1 + \text{Sqrt}[3])*s + r*x}{(1 - \text{Sqrt}[3])*s + r*x}], -7 + 4*\text{Sqrt}[3]]/(3^{1/4}*r*\text{Sqrt}[a + b*x^3]*\text{Sqrt}[-((s*(s + r*x))/(1 - \text{Sqrt}[3])*s + r*x)^2])], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 235

$\text{Int}[(a + b*x^2)^{-1/3}, x_Symbol] := \text{Dist}[(3*\text{Sqrt}[b*x^2])/(2*b*x), \text{Subst}[\text{Int}[x/\text{Sqrt}[-a + x^3], x], x, (a + b*x^2)^{1/3}], x] /; \text{FreeQ}\{a, b\}, x]$

Rule 304

$\text{Int}[x/\text{Sqrt}[a + b*x^3], x_Symbol] := \text{With}\{r = \text{Numer}[\text{Rt}[b/a, 3]], s = \text{Denom}[\text{Rt}[b/a, 3]]\}, -\text{Dist}[(\text{Sqrt}[2]*s)/(\text{Sqrt}[2 - \text{Sqrt}[3]]*r), \text{Int}[1/\text{Sqrt}[a + b*x^3], x], x] + \text{Dist}[1/r, \text{Int}[(1 + \text{Sqrt}[3])*s + r*x]/\text{Sqrt}[a + b*x^3], x], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a]$

Rule 388

$\text{Int}[(a + b*x^n)^p * (c + d*x^n), x_Symbol] := \text{Simp}[(d*x*(a + b*x^n)^{p+1})/(b*(n*(p+1) + 1)), x] - \text{Dist}[(a*d - b*c*(n*(p+1) + 1))/(b*(n*(p+1) + 1)), \text{Int}[(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[n*(p+1) + 1, 0]$

Rule 413

$\text{Int}[(a + b*x^n)^p * (c + d*x^n)^q, x_Symbol] := \text{Simp}[(a*d - c*b)*x*(a + b*x^n)^{p+1}*(c + d*x^n)^{q-1})/(a*b*n*(p+1)), x] - \text{Dist}[1/(a*b*n*(p+1)), \text{Int}[(a + b*x^n)^{p+1}*(c + d*x^n)^{q-2}*\text{Simp}[c*(a*d - c*b*(n*(p+1) + 1)) + d*(a*d*(n*(q-1) + 1) - b*c*(n*(p+q) + 1))*x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[p, -1] \&\& \text{GtQ}[q, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 526

$\text{Int}[(a + b*x^n)^p * (c + d*x^n)^q * (e + f*x^n), x_Symbol] := -\text{Simp}[(b*e - a*f)*x*(a + b*x^n)^{p+1}*(c + d*x^n)^q/(a*b*n*(p+1)), x] + \text{Dist}[1/(a*b*n*(p+1)), \text{Int}[(a + b*x^n)^{p+1}*(c + d*x^n)^{q-1}*\text{Simp}[c*(b*e*n*(p+1) + b*e - a*f) + d*(b*e*n*(p+1) + (b*e - a*f)*(n*q + 1))*x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{LtQ}[p, -1] \&\& \text{GtQ}[q, 0]$

Rule 528

$\text{Int}[(a + b*x^n)^p * (c + d*x^n)^q * (e + f*x^n), x_Symbol] := \text{Simp}[(f*x*(a + b*x^n)^{p+1}*(c + d*x^n)^q)/(b*(n*(p+q+1) + 1)), x] + \text{Dist}[1/(b*(n*(p+q+1) + 1)), \text{Int}[(a + b*x^n)^p*(c + d*x^n)^{q-1}*\text{Simp}[c*(b*e - a*f + b*e*n*(p+q+1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p+q+1))*x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \text{GtQ}[q, 0] \&\& \text{NeQ}[n*(p+q+1) + 1, 0]$

Rule 1879

$\text{Int}[(c + d*x)/\text{Sqrt}[a + b*x^3], x_Symbol] := \text{With}\{r = \text{Numer}[\text{Simplify}[(1 + \text{Sqrt}[3])*d/c]], s = \text{Denom}[\text{Simplify}[(1 + \text{Sqrt}[3])*d/c]]\}, \text{Simp}[(2*d*s^3*\text{Sqrt}[a + b*x^3])/(a*r^2*((1 - \text{Sqrt}[3])*s + r*x)), x] + \text{Simp}[(3^{1/4}*\text{Sqrt}[2 + \text{Sqrt}[3]]*d*s*(s + r*x)*\text{Sqrt}[(s^2 - r*s*x + r^2*x^2)/(1 - \text{Sqrt}[3])*s + r*x]^2]*\text{EllipticE}[\text{ArcSin}[\frac{(1 + \text{Sqrt}[3])*s + r*x}{(1 - \text{Sqrt}[3])*s + r*x}], x]$

rt[3])*s + r*x)], -7 + 4*sqrt[3]]/(r^2*sqrt[a + b*x^3]*sqrt[-((s*(s + r*x)) / ((1 - sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned} \int \frac{(3a + bx^2)^4}{(a - bx^2)^{7/3}} dx &= \frac{3x(3a + bx^2)^3}{2(a - bx^2)^{4/3}} - \frac{3 \int \frac{(3a + bx^2)^2(-12a^2b + 20ab^2x^2)}{(a - bx^2)^{4/3}} dx}{8ab} \\ &= -\frac{9x(3a + bx^2)^2}{2\sqrt[3]{a - bx^2}} + \frac{3x(3a + bx^2)^3}{2(a - bx^2)^{4/3}} - \frac{9 \int \frac{(3a + bx^2)(-48a^3b^2 - 48a^2b^3x^2)}{\sqrt[3]{a - bx^2}} dx}{16a^2b^2} \\ &= -\frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) - \frac{9x(3a + bx^2)^2}{2\sqrt[3]{a - bx^2}} + \frac{3x(3a + bx^2)^3}{2(a - bx^2)^{4/3}} + \frac{27 \int \frac{768a^4b^3 + 640a^3b^4x^2}{\sqrt[3]{a - bx^2}} dx}{208a^2b^3} \\ &= -\frac{3240}{91}ax(a - bx^2)^{2/3} - \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) - \frac{9x(3a + bx^2)^2}{2\sqrt[3]{a - bx^2}} + \frac{3x(3a + bx^2)^3}{2(a - bx^2)^{4/3}} + \frac{1}{91} \\ &= -\frac{3240}{91}ax(a - bx^2)^{2/3} - \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) - \frac{9x(3a + bx^2)^2}{2\sqrt[3]{a - bx^2}} + \frac{3x(3a + bx^2)^3}{2(a - bx^2)^{4/3}} - \frac{1}{91} \\ &= -\frac{3240}{91}ax(a - bx^2)^{2/3} - \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) - \frac{9x(3a + bx^2)^2}{2\sqrt[3]{a - bx^2}} + \frac{3x(3a + bx^2)^3}{2(a - bx^2)^{4/3}} + \frac{1}{91} \\ &= -\frac{3240}{91}ax(a - bx^2)^{2/3} - \frac{81}{13}x(a - bx^2)^{2/3}(3a + bx^2) - \frac{9x(3a + bx^2)^2}{2\sqrt[3]{a - bx^2}} + \frac{3x(3a + bx^2)^3}{2(a - bx^2)^{4/3}} - \frac{1}{91} \end{aligned}$$

Mathematica [C] time = 5.08, size = 96, normalized size = 0.15

$$\frac{3 \left(1647a^3x - 4104a^2x(a - bx^2) \sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1 \left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a} \right) - 4743a^2bx^3 + 177ab^2x^5 + 7b^3x^7 \right)}{91(a - bx^2)^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)^4/(a - b*x^2)^(7/3), x]

[Out] (-3*(1647*a^3*x - 4743*a^2*b*x^3 + 177*a*b^2*x^5 + 7*b^3*x^7 - 4104*a^2*x*(a - b*x^2)*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a]))/(91*(a - b*x^2)^(4/3))

fricas [F] time = 1.13, size = 0, normalized size = 0.00

$$\text{integral} \left(-\frac{(b^4x^8 + 12ab^3x^6 + 54a^2b^2x^4 + 108a^3bx^2 + 81a^4)(-bx^2 + a)^{\frac{2}{3}}}{b^3x^6 - 3ab^2x^4 + 3a^2bx^2 - a^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^4/(-b*x^2+a)^(7/3),x, algorithm="fricas")

[Out] integral(-(b^4*x^8 + 12*a*b^3*x^6 + 54*a^2*b^2*x^4 + 108*a^3*b*x^2 + 81*a^4)*(-b*x^2 + a)^(2/3)/(b^3*x^6 - 3*a*b^2*x^4 + 3*a^2*b*x^2 - a^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^4}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^4/(-b*x^2+a)^(7/3),x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^4/(-b*x^2 + a)^(7/3), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^4}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)^4/(-b*x^2+a)^(7/3),x)

[Out] int((b*x^2+3*a)^4/(-b*x^2+a)^(7/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^4}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^4/(-b*x^2+a)^(7/3),x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^4/(-b*x^2 + a)^(7/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + 3a)^4}{(a - bx^2)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)^4/(a - b*x^2)^(7/3),x)

[Out] int((3*a + b*x^2)^4/(a - b*x^2)^(7/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(3a + bx^2)^4}{(a - bx^2)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)**4/(-b*x**2+a)**(7/3),x)

[Out] Integral((3*a + b*x**2)**4/(a - b*x**2)**(7/3), x)

$$3.136 \quad \int \frac{(3a+bx^2)^3}{(a-bx^2)^{7/3}} dx$$

Optimal. Leaf size=596

$$108\sqrt{2}3^{3/4}a^{4/3} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} \operatorname{EllipticF} \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right), 4\sqrt{3} - 7 \right) - 162 \sqrt[4]{3} \sqrt[3]{a} \sqrt[3]{a-bx^2} \\ \frac{7bx \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}{\sqrt[4]{3} \sqrt[3]{a} \sqrt[3]{a-bx^2}}$$

[Out] $-27/14*x*(-b*x^2+a)^{(2/3)}+3/2*x*(b*x^2+3*a)^2/(-b*x^2+a)^{(4/3)}-324/7*a*x/(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))+108/7*3^{(3/4)}*a^{(4/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*2^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}-162/7*3^{(1/4)}*a^{(4/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.43, antiderivative size = 596, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {413, 21, 388, 235, 304, 219, 1879}

$$108\sqrt{2}3^{3/4}a^{4/3} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} F \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right) \middle| -7 + 4\sqrt{3} \right) - 162\sqrt[4]{3} \sqrt[3]{a} \sqrt[3]{a-bx^2} \\ \frac{7bx \sqrt{\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}{\sqrt[4]{3} \sqrt[3]{a} \sqrt[3]{a-bx^2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(3*a + b*x^2)^3/(a - b*x^2)^{(7/3)}, x]$

[Out] $(-27*x*(a - b*x^2)^{(2/3)})/14 + (3*x*(3*a + b*x^2)^2)/(2*(a - b*x^2)^{(4/3)}) - (324*a*x)/(7*((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})) - (162*3^{(1/4)}*\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]]*a^{(4/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})], -7 + 4*\operatorname{Sqrt}[3]])/(7*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])) + (108*\operatorname{Sqrt}[2]*3^{(3/4)}*a^{(4/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})], -7 + 4*\operatorname{Sqrt}[3]])/(7*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]))$

Rule 21

$\operatorname{Int}[(u_*)*((a_*) + (b_*)*(v_*))^{(m_*)}*((c_*) + (d_*)*(v_*))^{(n_*)}, x_Symbol] :> \operatorname{Dist}[(b/d)^m, \operatorname{Int}[u*(c + d*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, n\}, x$ && $\operatorname{EqQ}[b*c - a*d, 0]$ && $\operatorname{IntegerQ}[m]$ && $(! \operatorname{IntegerQ}[n] || \operatorname{SimplerQ}[c + d*x,$

$a + b*x]$)

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/S
qrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 388

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 413

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p +
1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q -
2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p
+ q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,
0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((
1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(3a + bx^2)^3}{(a - bx^2)^{7/3}} dx &= \frac{3x(3a + bx^2)^2}{2(a - bx^2)^{4/3}} - \frac{3 \int \frac{(3a + bx^2)(-12a^2b + 12ab^2x^2)}{(a - bx^2)^{4/3}} dx}{8ab} \\
&= \frac{3x(3a + bx^2)^2}{2(a - bx^2)^{4/3}} + \frac{9}{2} \int \frac{3a + bx^2}{\sqrt[3]{a - bx^2}} dx \\
&= -\frac{27}{14}x(a - bx^2)^{2/3} + \frac{3x(3a + bx^2)^2}{2(a - bx^2)^{4/3}} + \frac{1}{7}(108a) \int \frac{1}{\sqrt[3]{a - bx^2}} dx \\
&= -\frac{27}{14}x(a - bx^2)^{2/3} + \frac{3x(3a + bx^2)^2}{2(a - bx^2)^{4/3}} - \frac{(162a\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{x}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{7bx} \\
&= -\frac{27}{14}x(a - bx^2)^{2/3} + \frac{3x(3a + bx^2)^2}{2(a - bx^2)^{4/3}} + \frac{(162a\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{(1+\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{7bx} \\
&= -\frac{27}{14}x(a - bx^2)^{2/3} + \frac{3x(3a + bx^2)^2}{2(a - bx^2)^{4/3}} - \frac{324ax}{7((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2})} - \frac{162\sqrt[4]{3}\sqrt{2 + \sqrt{3}} a^{4/3}}{7}
\end{aligned}$$

Mathematica [C] time = 5.07, size = 83, normalized size = 0.14

$$\frac{81a^2x + 108ax(a - bx^2) \sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a}\right) + 90abx^3 - 3b^2x^5}{7(a - bx^2)^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)^3/(a - b*x^2)^(7/3), x]

[Out] (81*a^2*x + 90*a*b*x^3 - 3*b^2*x^5 + 108*a*x*(a - b*x^2)*(1 - (b*x^2)/a)^(1/3))*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a]/(7*(a - b*x^2)^(4/3))

fricas [F] time = 1.03, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{(b^3x^6 + 9ab^2x^4 + 27a^2bx^2 + 27a^3)(-bx^2 + a)^{2/3}}{b^3x^6 - 3ab^2x^4 + 3a^2bx^2 - a^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(7/3), x, algorithm="fricas")

[Out] integral(-(b^3*x^6 + 9*a*b^2*x^4 + 27*a^2*b*x^2 + 27*a^3)*(-b*x^2 + a)^(2/3))/(b^3*x^6 - 3*a*b^2*x^4 + 3*a^2*b*x^2 - a^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{7/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(7/3),x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^3/(-b*x^2 + a)^(7/3), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)^3/(-b*x^2+a)^(7/3),x)

[Out] int((b*x^2+3*a)^3/(-b*x^2+a)^(7/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^3}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^3/(-b*x^2+a)^(7/3),x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)^3/(-b*x^2 + a)^(7/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + 3a)^3}{(a - bx^2)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)^3/(a - b*x^2)^(7/3),x)

[Out] int((3*a + b*x^2)^3/(a - b*x^2)^(7/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(3a + bx^2)^3}{(a - bx^2)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)**3/(-b*x**2+a)**(7/3),x)

[Out] Integral((3*a + b*x**2)**3/(a - b*x**2)**(7/3), x)

$$3.137 \quad \int \frac{(3a+bx^2)^2}{(a-bx^2)^{7/3}} dx$$

Optimal. Leaf size=44

$$\frac{3x(3a+bx^2)}{2(a-bx^2)^{4/3}} + \frac{9x}{2\sqrt[3]{a-bx^2}}$$

[Out] $9/2*x/(-b*x^2+a)^{(1/3)}+3/2*x*(b*x^2+3*a)/(-b*x^2+a)^{(4/3)}$

Rubi [A] time = 0.02, antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {413, 383}

$$\frac{3x(3a+bx^2)}{2(a-bx^2)^{4/3}} + \frac{9x}{2\sqrt[3]{a-bx^2}}$$

Antiderivative was successfully verified.

[In] Int[(3*a + b*x^2)^2/(a - b*x^2)^(7/3), x]

[Out] (9*x)/(2*(a - b*x^2)^(1/3)) + (3*x*(3*a + b*x^2))/(2*(a - b*x^2)^(4/3))

Rule 383

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(c*x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a*d - b*c*(n*(p + 1) + 1), 0]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rubi steps

$$\begin{aligned} \int \frac{(3a+bx^2)^2}{(a-bx^2)^{7/3}} dx &= \frac{3x(3a+bx^2)}{2(a-bx^2)^{4/3}} - \frac{3 \int \frac{-12a^2b+4ab^2x^2}{(a-bx^2)^{4/3}} dx}{8ab} \\ &= \frac{9x}{2\sqrt[3]{a-bx^2}} + \frac{3x(3a+bx^2)}{2(a-bx^2)^{4/3}} \end{aligned}$$

Mathematica [A] time = 5.03, size = 24, normalized size = 0.55

$$\frac{9ax - 3bx^3}{(a - bx^2)^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)^2/(a - b*x^2)^(7/3), x]

[Out] (9*a*x - 3*b*x^3)/(a - b*x^2)^(4/3)

fricas [A] time = 0.55, size = 42, normalized size = 0.95

$$\frac{3(bx^3 - 3ax)(-bx^2 + a)^{\frac{2}{3}}}{b^2x^4 - 2abx^2 + a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(7/3), x, algorithm="fricas")

[Out] -3*(b*x^3 - 3*a*x)*(-b*x^2 + a)^(2/3)/(b^2*x^4 - 2*a*b*x^2 + a^2)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + 3a)^2}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(7/3), x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)^2/(-b*x^2 + a)^(7/3), x)

maple [A] time = 0.00, size = 24, normalized size = 0.55

$$\frac{3(-bx^2 + 3a)x}{(-bx^2 + a)^{\frac{4}{3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)^2/(-b*x^2+a)^(7/3), x)

[Out] 3/(-b*x^2+a)^(4/3)*x*(-b*x^2+3*a)

maxima [A] time = 1.89, size = 33, normalized size = 0.75

$$\frac{3(bx^3 - 3ax)}{(bx^2 - a)(-bx^2 + a)^{\frac{1}{3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)^2/(-b*x^2+a)^(7/3), x, algorithm="maxima")

[Out] 3*(b*x^3 - 3*a*x)/((b*x^2 - a)*(-b*x^2 + a)^(1/3))

mupad [B] time = 4.78, size = 27, normalized size = 0.61

$$\frac{3x(a - bx^2) + 6ax}{(a - bx^2)^{4/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)^2/(a - b*x^2)^(7/3), x)

[Out] (3*x*(a - b*x^2) + 6*a*x)/(a - b*x^2)^(4/3)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(3a + bx^2)^2}{(a - bx^2)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)**2/(-b*x**2+a)**(7/3),x)

[Out] Integral((3*a + b*x**2)**2/(a - b*x**2)**(7/3), x)

$$3.138 \quad \int \frac{3a+bx^2}{(a-bx^2)^{7/3}} dx$$

Optimal. Leaf size=590

$$\frac{3 \cdot 3^{3/4} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} \operatorname{EllipticF} \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right), 4\sqrt{3} - 7 \right) + 9\sqrt[4]{3} \sqrt{2}}{2\sqrt{2} a^{2/3} bx \sqrt{-\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}$$

[Out] $3/2*x/(-b*x^2+a)^{(4/3)}+9/4*x/a/(-b*x^2+a)^{(1/3)}+9/4*x/a/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))-3/4*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*3^{(3/4)}/a^{(2/3)}/b/x*2^{(1/2)}/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}+9/8*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)}))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})), 2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})*3^{(1/4)}/a^{(2/3)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^2)^{(1/2)}$

Rubi [A] time = 0.37, antiderivative size = 590, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {385, 199, 235, 304, 219, 1879}

$$\frac{3 \cdot 3^{3/4} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right) \sqrt{\frac{a^{2/3} + \sqrt[3]{a} \sqrt[3]{a-bx^2} + (a-bx^2)^{2/3}}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}} F \left(\sin^{-1} \left(\frac{(1+\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}}{(1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2}} \right) \middle| -7 + 4\sqrt{3} \right) + 9\sqrt[4]{3} \sqrt{2 + \sqrt{3}}}{2\sqrt{2} a^{2/3} bx \sqrt{-\frac{\sqrt[3]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)}{\left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)^2}}}$$

Antiderivative was successfully verified.

[In] Int[(3*a + b*x^2)/(a - b*x^2)^(7/3), x]

[Out] $(3*x)/(2*(a - b*x^2)^{(4/3)}) + (9*x)/(4*a*(a - b*x^2)^{(1/3)}) + (9*x)/(4*a*((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})) + (9*3^{(1/4)}*\operatorname{Sqrt}[2 + \operatorname{Sqrt}[3]]*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})), -7 + 4*\operatorname{Sqrt}[3]])/(8*a^{(2/3)}*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]) - (3*3^{(3/4)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)} + a^{(1/3)}*(a - b*x^2)^{(1/3)} + (a - b*x^2)^{(2/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])*\operatorname{EllipticF}[\operatorname{ArcSin}(((1 + \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})), -7 + 4*\operatorname{Sqrt}[3]])/(2*\operatorname{Sqrt}[2]*a^{(2/3)}*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \operatorname{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2])])$

Rule 199

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denomin

ator[p + 1/n] < Denominator[p])

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x)
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sq
rt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -S
imp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b
*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; Fre
eQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n +
p, 0])
```

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = N
umer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c
]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + S
imp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/(
(1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sq
rt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x)
)/((1 - Sqrt[3])*s + r*x)^2)]), x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&
EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{3a + bx^2}{(a - bx^2)^{7/3}} dx &= \frac{3x}{2(a - bx^2)^{4/3}} + \frac{3}{2} \int \frac{1}{(a - bx^2)^{4/3}} dx \\
&= \frac{3x}{2(a - bx^2)^{4/3}} + \frac{9x}{4a\sqrt[3]{a - bx^2}} - \frac{3 \int \frac{1}{\sqrt[3]{a - bx^2}} dx}{4a} \\
&= \frac{3x}{2(a - bx^2)^{4/3}} + \frac{9x}{4a\sqrt[3]{a - bx^2}} + \frac{(9\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{x}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{8abx} \\
&= \frac{3x}{2(a - bx^2)^{4/3}} + \frac{9x}{4a\sqrt[3]{a - bx^2}} - \frac{(9\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{(1+\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{8abx} + \frac{(9\sqrt{-bx^2}) \operatorname{Subst}\left(\int \frac{(1-\sqrt{3})\sqrt[3]{a-x}}{\sqrt{-a+x^3}} dx, x, \sqrt[3]{a - bx^2}\right)}{8abx} \\
&= \frac{3x}{2(a - bx^2)^{4/3}} + \frac{9x}{4a\sqrt[3]{a - bx^2}} + \frac{9x}{4a\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)} + \frac{9\sqrt[4]{3}\sqrt{2 + \sqrt{3}}\left(\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)}{4a\left((1 - \sqrt{3})\sqrt[3]{a} - \sqrt[3]{a - bx^2}\right)}
\end{aligned}$$

Mathematica [C] time = 0.04, size = 74, normalized size = 0.13

$$\frac{-3x(a - bx^2)\sqrt[3]{1 - \frac{bx^2}{a}} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{3}{2}; \frac{bx^2}{a}\right) + 15ax - 9bx^3}{4a(a - bx^2)^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(3*a + b*x^2)/(a - b*x^2)^(7/3), x]

[Out] (15*a*x - 9*b*x^3 - 3*x*(a - b*x^2)*(1 - (b*x^2)/a)^(1/3)*Hypergeometric2F1[1/3, 1/2, 3/2, (b*x^2)/a])/(4*a*(a - b*x^2)^(4/3))

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{(bx^2 + 3a)(-bx^2 + a)^{\frac{2}{3}}}{b^3x^6 - 3ab^2x^4 + 3a^2bx^2 - a^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(7/3), x, algorithm="fricas")

[Out] integral(-(b*x^2 + 3*a)*(-b*x^2 + a)^(2/3)/(b^3*x^6 - 3*a*b^2*x^4 + 3*a^2*b*x^2 - a^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(7/3), x, algorithm="giac")

[Out] integrate((b*x^2 + 3*a)/(-b*x^2 + a)^(7/3), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+3*a)/(-b*x^2+a)^(7/3),x)

[Out] int((b*x^2+3*a)/(-b*x^2+a)^(7/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + 3a}{(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+3*a)/(-b*x^2+a)^(7/3),x, algorithm="maxima")

[Out] integrate((b*x^2 + 3*a)/(-b*x^2 + a)^(7/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{bx^2 + 3a}{(a - bx^2)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*a + b*x^2)/(a - b*x^2)^(7/3),x)

[Out] int((3*a + b*x^2)/(a - b*x^2)^(7/3), x)

sympy [A] time = 12.99, size = 60, normalized size = 0.10

$$\frac{3x {}_2F_1\left(\frac{1}{2}, \frac{7}{3} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{a^{\frac{4}{3}}} + \frac{bx^3 {}_2F_1\left(\frac{3}{2}, \frac{7}{3} \middle| \frac{bx^2 e^{2i\pi}}{a}\right)}{3a^{\frac{7}{3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+3*a)/(-b*x**2+a)**(7/3),x)

[Out] 3*x*hyper((1/2, 7/3), (3/2,), b*x**2*exp_polar(2*I*pi)/a)/a**(4/3) + b*x**3*hyper((3/2, 7/3), (5/2,), b*x**2*exp_polar(2*I*pi)/a)/(3*a**(7/3))

$$3.139 \quad \int \frac{1}{(a-bx^2)^{7/3}(3a+bx^2)} dx$$

Optimal. Leaf size=796

$$\frac{21x}{64a^3\sqrt[3]{a-bx^2}} + \frac{21x}{64a^3\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{3x}{32a^2(a-bx^2)^{4/3}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{32\ 2^{2/3}\sqrt{3}\ a^{17/6}\sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{32\ 2^{2/3}\sqrt{3}\ a^{17/6}\sqrt{b}}$$

[Out] $3/32*x/a^2/(-b*x^2+a)^{(4/3)}+21/64*x/a^3/(-b*x^2+a)^{(1/3)}+21/64*x/a^3/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))^{(1/3)}+1/64*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/(a^{(1/3)}+2^{(1/3)}*(-b*x^2+a)^{(1/3)})^{(1/3)}+2^{(1/3)}/a^{(17/6)}/b^{(1/2)}-1/192*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})^{(1/3)}+2^{(1/3)}/a^{(17/6)}/b^{(1/2)}+1/192*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(-b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})^{(1/3)}+2^{(1/3)}/a^{(17/6)}*3^{(1/2)}/b^{(1/2)}+1/192*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})^{(1/3)}+2^{(1/3)}/a^{(17/6)}*3^{(1/2)}/b^{(1/2)}-7/64*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}+21/128*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)}*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)})*3^{(1/4)}/a^{(8/3)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}$

Rubi [A] time = 0.54, antiderivative size = 796, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {414, 527, 530, 235, 304, 219, 1879, 393}

$$\frac{21x}{64a^3\sqrt[3]{a-bx^2}} + \frac{21x}{64a^3\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{3x}{32a^2(a-bx^2)^{4/3}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{32\ 2^{2/3}\sqrt{3}\ a^{17/6}\sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{32\ 2^{2/3}\sqrt{3}\ a^{17/6}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(7/3)*(3*a + b*x^2)),x]

[Out] $(3*x)/(32*a^2*(a-b*x^2)^{(4/3)})+(21*x)/(64*a^3*(a-b*x^2)^{(1/3)})+(21*x)/(64*a^3*((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))+\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(32*2^{(2/3)}*\operatorname{Sqrt}[3]*a^{(17/6)}*\operatorname{Sqrt}[b])+\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(a-b*x^2)^{(1/3)}))/(\operatorname{Sqrt}[b]*x)]/(32*2^{(2/3)}*\operatorname{Sqrt}[3]*a^{(17/6)}*\operatorname{Sqrt}[b])-\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(96*2^{(2/3)}*a^{(17/6)}*\operatorname{Sqrt}[b])+\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{(1/6)}*(a^{(1/3)}+2^{(1/3)}*(a-b*x^2)^{(1/3)}))]/(32*2^{(2/3)}*a^{(17/6)}*\operatorname{Sqrt}[b])+(21*3^{(1/4)}*\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})],-7+4*\operatorname{Sqrt}[3]]/(128*a^{(8/3)}*b*x*\operatorname{Sqrt}[-(a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2])-(7*3^{(3/4)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)})*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*$

$a^{(1/3)} - (a - b*x^2)^{(1/3)}$], $-7 + 4*\text{Sqrt}[3]$]/(32*Sqrt[2]* $a^{(8/3)}$ * $b*x*\text{Sqrt}[-((a^{(1/3)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/((1 - \text{Sqrt}[3])*a^{(1/3)} - (a - b*x^2)^{(1/3)})^2]$)

Rule 219

Int[1/Sqrt[(a_) + (b_)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)]], -7 + 4*Sqrt[3])/((3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2)]), x] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 235

Int[((a_) + (b_)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

Rule 304

Int[(x_)/Sqrt[(a_) + (b_)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sqrt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 393

Int[1/(((a_) + (b_)*(x_)^2)^(1/3)*((c_) + (d_)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]* $a^{(1/3)}$ *d), x] + (Simp[(q*ArcTanh[($a^{(1/3)}$ *q*x)/($a^{(1/3)}$ + 2^(1/3)*(a + b*x^2)^(1/3)))]/(2*2^(2/3)* $a^{(1/3)}$ *d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)* $a^{(1/3)}$ *d), x] + Simp[(q*ArcTan[(Sqrt[3]*($a^{(1/3)}$ - 2^(1/3)*(a + b*x^2)^(1/3)))/($a^{(1/3)}$ *q*x)]/(2*2^(2/3)*Sqrt[3]* $a^{(1/3)}$ *d), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rule 414

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 530

Int((((a_) + (b_)*(x_)^(n_))^(p_)*((e_) + (f_)*(x_)^(n_)))/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p}

, n}, x]

Rule 1879

```
Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)]], -7 + 4*Sqrt[3])]/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/((1 - Sqrt[3])*s + r*x)^2))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]
```

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a-bx^2)^{7/3}(3a+bx^2)} dx &= \frac{3x}{32a^2(a-bx^2)^{4/3}} + \frac{3 \int \frac{\frac{23ab}{3} + \frac{5b^2x^2}{3}}{(a-bx^2)^{4/3}(3a+bx^2)} dx}{32a^2b} \\
 &= \frac{3x}{32a^2(a-bx^2)^{4/3}} + \frac{21x}{64a^3\sqrt[3]{a-bx^2}} + \frac{9 \int \frac{-\frac{68}{9}a^2b^2 - \frac{28}{9}ab^3x^2}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx}{256a^4b^2} \\
 &= \frac{3x}{32a^2(a-bx^2)^{4/3}} + \frac{21x}{64a^3\sqrt[3]{a-bx^2}} - \frac{7 \int \frac{1}{\sqrt[3]{a-bx^2}} dx}{64a^3} + \frac{\int \frac{1}{\sqrt[3]{a-bx^2}(3a+bx^2)} dx}{16a^2} \\
 &= \frac{3x}{32a^2(a-bx^2)^{4/3}} + \frac{21x}{64a^3\sqrt[3]{a-bx^2}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{32 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\sqrt{b}}\right)}{32 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} \\
 &= \frac{3x}{32a^2(a-bx^2)^{4/3}} + \frac{21x}{64a^3\sqrt[3]{a-bx^2}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{32 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\sqrt{b}}\right)}{32 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}} \\
 &= \frac{3x}{32a^2(a-bx^2)^{4/3}} + \frac{21x}{64a^3\sqrt[3]{a-bx^2}} + \frac{21x}{64a^3\left((1-\sqrt{3})\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)} + \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{a-bx^2}\right)}{\sqrt{b}}\right)}{32 \cdot 2^{2/3} \sqrt{3} a^{17/6} \sqrt{b}}
 \end{aligned}$$

Mathematica [C] time = 0.21, size = 248, normalized size = 0.31

$$\frac{x \left(27a \left(\frac{9a-7bx^2}{a-bx^2} - \frac{51a^2 F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{(3a+bx^2) \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) + 9a F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right) \right) - 7bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} F_1\left(\frac{3}{2}; \frac{1}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) \right)}{576a^4 \sqrt[3]{a-bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(7/3)*(3*a + b*x^2)), x]

[Out] (x*(-7*b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 27*a*((9*a - 7*b*x^2)/(a - b*x^2) - (51*a^2*AppellF1[1/2, 1

/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/((3*a + b*x^2)*(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])))))/(576*a^4*(a - b*x^2)^(1/3))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 3*a)*(-b*x^2 + a)^(7/3)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{7}{3}}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a),x)

[Out] int(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)(-bx^2 + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 3*a)*(-b*x^2 + a)^(7/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{\frac{7}{3}}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(7/3)*(3*a + b*x^2)),x)

[Out] int(1/((a - b*x^2)^(7/3)*(3*a + b*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a - bx^2)^{\frac{7}{3}}(3a + bx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-b*x**2+a)**(7/3)/(b*x**2+3*a),x)
```

```
[Out] Integral(1/((a - b*x**2)**(7/3)*(3*a + b*x**2)), x)
```

$$3.140 \quad \int \frac{1}{(a-bx^2)^{7/3} (3a+bx^2)^2} dx$$

Optimal. Leaf size=827

$$\frac{79x}{768a^4 \sqrt[3]{a-bx^2}} + \frac{x}{24a^2 (a-bx^2)^{4/3} (bx^2+3a)} + \frac{79x}{768a^4 \left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)} + \frac{5x}{384a^3 (a-bx^2)^{4/3}} + \frac{\sqrt{3} \tan^{-1} \left(\frac{x}{\sqrt{a-bx^2}} \right)}{128 \cdot 2^{2/3}}$$

[Out] $5/384*x/a^3/(-b*x^2+a)^{(4/3)}+79/768*x/a^4/(-b*x^2+a)^{(1/3)}+1/24*x/a^2/(-b*x^2+a)^{(4/3)}/(b*x^2+3*a)+79/768*x/a^4/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)}))+3/256*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/6)})/(a^{(1/3)}+2^{(1/3)}*(-b*x^2+a)^{(1/3)}))*2^{(1/3)}/a^{(23/6)}/b^{(1/2)}-1/256*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/2)}))*2^{(1/3)}/a^{(23/6)}/b^{(1/2)}+1/256*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(-b*x^2+a)^{(1/3)}))*3^{(1/2)}/x/b^{(1/2)}))*2^{(1/3)}/a^{(23/6)}*3^{(1/2)}/b^{(1/2)}+1/256*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)}))*2^{(1/3)}/a^{(23/6)}*3^{(1/2)}/b^{(1/2)}-79/2304*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticF}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}*3^{(3/4)}/a^{(11/3)}/b/x*2^{(1/2)}/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}+79/1536*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*\operatorname{EllipticE}((-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1+3^{(1/2)})))/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})),2*I-I*3^{(1/2)})*((a^{(2/3)}+a^{(1/3)}*(-b*x^2+a)^{(1/3)}+(-b*x^2+a)^{(2/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}*(1/2*6^{(1/2)}+1/2*2^{(1/2)}))*3^{(1/4)}/a^{(11/3)}/b/x/(-a^{(1/3)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})/(-(-b*x^2+a)^{(1/3)}+a^{(1/3)}*(1-3^{(1/2)})))^{(1/2)}$

Rubi [A] time = 0.63, antiderivative size = 827, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {414, 527, 530, 235, 304, 219, 1879, 393}

$$\frac{79x}{768a^4 \sqrt[3]{a-bx^2}} + \frac{x}{24a^2 (a-bx^2)^{4/3} (bx^2+3a)} + \frac{79x}{768a^4 \left((1-\sqrt{3}) \sqrt[3]{a} - \sqrt[3]{a-bx^2} \right)} + \frac{5x}{384a^3 (a-bx^2)^{4/3}} + \frac{\sqrt{3} \tan^{-1} \left(\frac{x}{\sqrt{a-bx^2}} \right)}{128 \cdot 2^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(7/3)*(3*a + b*x^2)^2),x]

[Out] $(5*x)/(384*a^3*(a-b*x^2)^{(4/3)})+(79*x)/(768*a^4*(a-b*x^2)^{(1/3)})+x/(24*a^2*(a-b*x^2)^{(4/3)}*(3*a+b*x^2))+79*x/(768*a^4*((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}))+(\operatorname{Sqrt}[3]*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)])/(128*2^{(2/3)}*a^{(23/6)}*\operatorname{Sqrt}[b])+(\operatorname{Sqrt}[3]*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(a-b*x^2)^{(1/3)}))/(\operatorname{Sqrt}[b]*x)])/(128*2^{(2/3)}*a^{(23/6)}*\operatorname{Sqrt}[b])-\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(128*2^{(2/3)}*a^{(23/6)}*\operatorname{Sqrt}[b])+(3*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(a^{(1/6)}*(a^{(1/3)}+2^{(1/3)}*(a-b*x^2)^{(1/3)}))]/(128*2^{(2/3)}*a^{(23/6)}*\operatorname{Sqrt}[b])+(79*\operatorname{Sqrt}[2+\operatorname{Sqrt}[3]]*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2]*\operatorname{EllipticE}[\operatorname{ArcSin}[(1+\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)}]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})],-7+4*\operatorname{Sqrt}[3]))/(512*3^{(3/4)}*a^{(11/3)}*b*x*\operatorname{Sqrt}[-((a^{(1/3)}*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2])-(79*(a^{(1/3)}-(a-b*x^2)^{(1/3)}))*\operatorname{Sqrt}[(a^{(2/3)}+a^{(1/3)}*(a-b*x^2)^{(1/3)}+(a-b*x^2)^{(2/3)})]/((1-\operatorname{Sqrt}[3])*a^{(1/3)}-(a-b*x^2)^{(1/3)})^2])$

$$\frac{2/3)}{((1 - \sqrt{3})a^{1/3} - (a - bx^2)^{1/3})^2} \text{EllipticF}[\text{ArcSin}[\frac{(1 + \sqrt{3})a^{1/3} - (a - bx^2)^{1/3}}{(1 - \sqrt{3})a^{1/3} - (a - bx^2)^{1/3}}], -7 + 4\sqrt{3}]] / (384\sqrt{2} \cdot 3^{1/4} a^{11/3} b x \sqrt{-(a^{1/3}) \cdot (a^{1/3} - (a - bx^2)^{1/3})}) / ((1 - \sqrt{3})a^{1/3} - (a - bx^2)^{1/3})^2]$$

Rule 219

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[(2*Sqrt[2 - Sqrt[3]]*(s + r*x)*Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticF[ArcSin[((1 + Sqrt[3])*s
+ r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(3^(1/4)*r*Sqrt[a + b*x^3
]*Sqrt[-((s*(s + r*x))/(1 - Sqrt[3])*s + r*x)^2]), x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

Rule 235

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[(3*Sqrt[b*x^2])/(2*b*x
), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

Rule 304

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, -Dist[(Sqrt[2]*s)/(Sqrt[2 - Sqrt[3]]*r), Int[1/Sq
rt[a + b*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x
^3], x], x]] /; FreeQ[{a, b}, x] && NegQ[a]
```

Rule 393

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Wit
h[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^
(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2
)^(1/3))])/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^
(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3))
)/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]]) /; FreeQ[{a, b, c, d}
, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]
```

Rule 414

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c -
a*d), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d), Int[(a + b*x^n)^(p + 1)*(c +
d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x]
, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1]
&& !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 530

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*
```

$(x_)^{(n_)}$, x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1879

Int[((c_) + (d_.)*(x_))/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Simplify[((1 + Sqrt[3])*d)/c]], s = Denom[Simplify[((1 + Sqrt[3])*d)/c]]}, Simp[(2*d*s^3*Sqrt[a + b*x^3])/(a*r^2*((1 - Sqrt[3])*s + r*x)), x] + Simp[(3^(1/4)*Sqrt[2 + Sqrt[3]]*d*s*(s + r*x)*Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]*EllipticE[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]])/(r^2*Sqrt[a + b*x^3]*Sqrt[-((s*(s + r*x))/(1 - Sqrt[3])*s + r*x)^2])], x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b*c^3 - 2*(5 + 3*Sqrt[3])*a*d^3, 0]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a - bx^2)^{7/3} (3a + bx^2)^2} dx &= \frac{x}{24a^2 (a - bx^2)^{4/3} (3a + bx^2)} - \frac{\int \frac{-7ab + \frac{11b^2x^2}{3}}{(a - bx^2)^{7/3} (3a + bx^2)} dx}{24a^2b} \\
 &= \frac{5x}{384a^3 (a - bx^2)^{4/3}} + \frac{x}{24a^2 (a - bx^2)^{4/3} (3a + bx^2)} - \frac{\int \frac{-\frac{194}{3}a^2b^2 - \frac{50}{9}ab^3x^2}{(a - bx^2)^{4/3} (3a + bx^2)} dx}{256a^4b^2} \\
 &= \frac{5x}{384a^3 (a - bx^2)^{4/3}} + \frac{79x}{768a^4 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 (a - bx^2)^{4/3} (3a + bx^2)} - \frac{3 \int \frac{\frac{344a^3b}{9}}{\sqrt[3]{a - bx^2}} dx}{20} \\
 &= \frac{5x}{384a^3 (a - bx^2)^{4/3}} + \frac{79x}{768a^4 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 (a - bx^2)^{4/3} (3a + bx^2)} - \frac{79 \int \frac{1}{\sqrt[3]{a - bx^2}} dx}{2304} \\
 &= \frac{5x}{384a^3 (a - bx^2)^{4/3}} + \frac{79x}{768a^4 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 (a - bx^2)^{4/3} (3a + bx^2)} + \frac{\sqrt{3} \tan^{-1}\left(\frac{x}{\sqrt{a - bx^2}}\right)}{128 \cdot 2^{2/3}} \\
 &= \frac{5x}{384a^3 (a - bx^2)^{4/3}} + \frac{79x}{768a^4 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 (a - bx^2)^{4/3} (3a + bx^2)} + \frac{\sqrt{3} \tan^{-1}\left(\frac{x}{\sqrt{a - bx^2}}\right)}{128 \cdot 2^{2/3}} \\
 &= \frac{5x}{384a^3 (a - bx^2)^{4/3}} + \frac{79x}{768a^4 \sqrt[3]{a - bx^2}} + \frac{x}{24a^2 (a - bx^2)^{4/3} (3a + bx^2)} + \frac{\sqrt{3} \tan^{-1}\left(\frac{x}{\sqrt{a - bx^2}}\right)}{768a^4 \left(\frac{1}{\sqrt{a - bx^2}}\right)}
 \end{aligned}$$

Mathematica [C] time = 0.24, size = 259, normalized size = 0.31

$$x \left(\frac{27a \left(\frac{299a^2 - 148abx^2 - 79b^2x^4}{a - bx^2} - \frac{387a^2 F_1 \left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a} \right)}{2bx^2 \left(F_1 \left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a} \right) - F_1 \left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a} \right) \right) + 9a F_1 \left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a} \right)}{3a + bx^2} \right) - 79bx^2 \sqrt[3]{1 - \frac{bx^2}{a}} F_1 \left(\frac{3}{2}; \frac{1}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a} \right) \right) \frac{1}{20736a^5 \sqrt[3]{a - bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(7/3)*(3*a + b*x^2)^2), x]

[Out] (x*(-79*b*x^2*(1 - (b*x^2)/a)^(1/3)*AppellF1[3/2, 1/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + (27*a*((299*a^2 - 148*a*b*x^2 - 79*b^2*x^4)/(a - b*x^2) - (387*a^2*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a])))/(3*a + b*x^2)))/(20736*a^5*(a - b*x^2)^(1/3))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^2 (-bx^2 + a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a)^2,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 3*a)^2*(-b*x^2 + a)^(7/3)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^3 (bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a)^2,x)

[Out] int(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)^2 (-bx^2 + a)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(7/3)/(b*x^2+3*a)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 3*a)^2*(-b*x^2 + a)^(7/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{7/3} (bx^2 + 3a)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(7/3)*(3*a + b*x^2)^2), x)

[Out] int(1/((a - b*x^2)^(7/3)*(3*a + b*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a - bx^2)^{7/3} (3a + bx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2+a)**(7/3)/(b*x**2+3*a)**2, x)

[Out] Integral(1/((a - b*x**2)**(7/3)*(3*a + b*x**2)**2), x)

$$3.141 \quad \int \frac{1}{(-3a-bx^2)\sqrt[3]{-a+bx^2}} dx$$

Optimal. Leaf size=252

$$\frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}\left(\sqrt[3]{-a}-\sqrt[3]{2}\sqrt[3]{bx^2-a}\right)}{\sqrt[3]{-a}\sqrt{bx}}\right)}{2\sqrt[2]{3}\sqrt{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt[3]{-a}\sqrt{bx}}{\sqrt{a}\left(\sqrt[3]{2}\sqrt[3]{bx^2-a}+\sqrt[3]{-a}\right)}\right)}{2\sqrt[2]{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}} - \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{2\sqrt[2]{3}\sqrt{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6\sqrt[2]{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}}$$

[Out] $1/12*\operatorname{arctanh}(x*b^{(1/2)}/a^{(1/2)})*2^{(1/3)}/(-a)^{(1/3)}/a^{(1/2)}/b^{(1/2)}-1/4*\operatorname{arctanh}((-a)^{(1/3)}*x*b^{(1/2)}/((-a)^{(1/3)}+2^{(1/3)}*(b*x^2-a)^{(1/3)})/a^{(1/2)}*2^{(1/3)}/(-a)^{(1/3)}/a^{(1/2)}/b^{(1/2)}-1/12*\operatorname{arctan}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})*2^{(1/3)}/(-a)^{(1/3)}*3^{(1/2)}/a^{(1/2)}/b^{(1/2)}-1/12*\operatorname{arctan}((-a)^{(1/3)}-2^{(1/3)}*(b*x^2-a)^{(1/3)})*3^{(1/2)}*a^{(1/2)}/(-a)^{(1/3)}/x/b^{(1/2)}*2^{(1/3)}/(-a)^{(1/3)}*3^{(1/2)}/a^{(1/2)}/b^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 252, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {393}

$$\frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}\left(\sqrt[3]{-a}-\sqrt[3]{2}\sqrt[3]{bx^2-a}\right)}{\sqrt[3]{-a}\sqrt{bx}}\right)}{2\sqrt[2]{3}\sqrt{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt[3]{-a}\sqrt{bx}}{\sqrt{a}\left(\sqrt[3]{2}\sqrt[3]{bx^2-a}+\sqrt[3]{-a}\right)}\right)}{2\sqrt[2]{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}} - \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{2\sqrt[2]{3}\sqrt{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6\sqrt[2]{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((-3*a - b*x^2)*(-a + b*x^2)^(1/3)), x]

[Out] $-\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a])/(\operatorname{Sqrt}[b]*x)]/(2*2^{(2/3)}*\operatorname{Sqrt}[3]*(-a)^{(1/3)}*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b]) - \operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[a]*((-a)^{(1/3)} - 2^{(1/3)}*(-a + b*x^2)^{(1/3)}))/((-a)^{(1/3)}*\operatorname{Sqrt}[b]*x)]/(2*2^{(2/3)}*\operatorname{Sqrt}[3]*(-a)^{(1/3)}*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b]) + \operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/(6*2^{(2/3)}*(-a)^{(1/3)}*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b]) - \operatorname{ArcTanh}[((-a)^{(1/3)}*\operatorname{Sqrt}[b]*x)/(\operatorname{Sqrt}[a]*((-a)^{(1/3)} + 2^{(1/3)}*(-a + b*x^2)^{(1/3)}))]/(2*2^{(2/3)}*(-a)^{(1/3)}*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b])$

Rule 393

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rubi steps

$$\int \frac{1}{(-3a-bx^2)\sqrt[3]{-a+bx^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{bx}}\right)}{2\sqrt[2]{3}\sqrt{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}} - \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{a}\left(\sqrt[3]{-a}-\sqrt[3]{2}\sqrt[3]{-a+bx^2}\right)}{\sqrt[3]{-a}\sqrt{bx}}\right)}{2\sqrt[2]{3}\sqrt{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6\sqrt[2]{3}\sqrt[3]{-a}\sqrt{a}\sqrt{b}}$$

Mathematica [C] time = 0.14, size = 163, normalized size = 0.65

$$\frac{9axF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{\sqrt[3]{bx^2-a}(3a+bx^2)} \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)\right) + 9aF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-3*a - b*x^2)*(-a + b*x^2)^(1/3)),x]

[Out] (-9*a*x*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/((-a + b*x^2)^(1/3)*(3*a + b*x^2)*(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-3*a)/(b*x^2-a)^(1/3),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(bx^2 + 3a)(bx^2 - a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-3*a)/(b*x^2-a)^(1/3),x, algorithm="giac")

[Out] integrate(-1/((b*x^2 + 3*a)*(b*x^2 - a)^(1/3)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 - 3a)(bx^2 - a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2-3*a)/(b*x^2-a)^(1/3),x)

[Out] int(1/(-b*x^2-3*a)/(b*x^2-a)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(bx^2 + 3a)(bx^2 - a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-3*a)/(b*x^2-a)^(1/3),x, algorithm="maxima")

[Out] -integrate(1/((b*x^2 + 3*a)*(b*x^2 - a)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$-\int \frac{1}{(bx^2 - a)^{\frac{1}{3}}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((b*x^2 - a)^(1/3)*(3*a + b*x^2)),x)

[Out] `-int(1/((b*x^2 - a)^(1/3)*(3*a + b*x^2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{3a\sqrt[3]{-a + bx^2} + bx^2\sqrt[3]{-a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-b*x**2-3*a)/(b*x**2-a)**(1/3), x)`

[Out] `-Integral(1/(3*a*(-a + b*x**2)**(1/3) + b*x**2*(-a + b*x**2)**(1/3)), x)`

$$3.142 \quad \int \frac{1}{(3a-bx^2)\sqrt[3]{a+bx^2}} dx$$

Optimal. Leaf size=202

$$\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt[6]{a}\left(\sqrt[3]{2}\sqrt[3]{a+bx^2}+\sqrt[3]{a}\right)}\right)}{2^{2/3}a^{5/6}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a+bx^2}\right)}{\sqrt{b}x}\right)}{2^{2/3}\sqrt{3}a^{5/6}\sqrt{b}} - \frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{6^{2/3}a^{5/6}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{b}x}\right)}{2^{2/3}\sqrt{3}a^{5/6}\sqrt{b}}$$

[Out] $1/4*\arctan(x*b^{(1/2)}/a^{(1/6)}/(a^{(1/3)}+2^{(1/3)}*(b*x^2+a)^{(1/3)}))*2^{(1/3)}/a^{(5/6)}/b^{(1/2)}-1/12*\arctan(x*b^{(1/2)}/a^{(1/2)})*2^{(1/3)}/a^{(5/6)}/b^{(1/2)}-1/12*\arctanh(a^{(1/6)}*(a^{(1/3)}-2^{(1/3)}*(b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})*2^{(1/3)}/a^{(5/6)}*3^{(1/2)}/b^{(1/2)}-1/12*\operatorname{arctanh}(3^{(1/2)}*a^{(1/2)}/x/b^{(1/2)})*2^{(1/3)}/a^{(5/6)}*3^{(1/2)}/b^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {392}

$$\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt[6]{a}\left(\sqrt[3]{2}\sqrt[3]{a+bx^2}+\sqrt[3]{a}\right)}\right)}{2^{2/3}a^{5/6}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a+bx^2}\right)}{\sqrt{b}x}\right)}{2^{2/3}\sqrt{3}a^{5/6}\sqrt{b}} - \frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{6^{2/3}a^{5/6}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{b}x}\right)}{2^{2/3}\sqrt{3}a^{5/6}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((3*a - b*x^2)*(a + b*x^2)^(1/3)),x]

[Out] $-\operatorname{ArcTan}\left[\frac{\sqrt{b}x}{\sqrt{a}}\right]/(6*2^{(2/3)}*a^{(5/6)}*\sqrt{b}) + \operatorname{ArcTan}\left[\frac{\sqrt{b}x}{a^{(1/6)}*(a^{(1/3)} + 2^{(1/3)}*(a + b*x^2)^{(1/3)})}\right]/(2*2^{(2/3)}*a^{(5/6)}*\sqrt{b}) - \operatorname{ArcTanh}\left[\frac{\sqrt{3}*\sqrt{a}}{\sqrt{b}x}\right]/(2*2^{(2/3)}*a^{(5/6)}*\sqrt{b}) - \operatorname{ArcTanh}\left[\frac{\sqrt{3}*a^{(1/6)}*(a^{(1/3)} - 2^{(1/3)}*(a + b*x^2)^{(1/3)})}{\sqrt{b}x}\right]/(2*2^{(2/3)}*a^{(5/6)}*\sqrt{b})$

Rule 392

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[b/a, 2]}, Simp[(q*ArcTanh[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (-Simp[(q*ArcTan[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3)])]/(2*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTanh[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{(3a-bx^2)\sqrt[3]{a+bx^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{6^{2/3}a^{5/6}\sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt[6]{a}\left(\sqrt[3]{a}+\sqrt[3]{2}\sqrt[3]{a+bx^2}\right)}\right)}{2^{2/3}a^{5/6}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{a}}{\sqrt{b}x}\right)}{2^{2/3}\sqrt{3}a^{5/6}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt[6]{a}\left(\sqrt[3]{a}-\sqrt[3]{2}\sqrt[3]{a+bx^2}\right)}{\sqrt{b}x}\right)}{2^{2/3}\sqrt{3}a^{5/6}\sqrt{b}}$$

Mathematica [C] time = 0.16, size = 166, normalized size = 0.82

$$\frac{9axF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{bx^2}{a}, \frac{bx^2}{3a}\right)}{(3a-bx^2)\sqrt[3]{a+bx^2}\left(2bx^2\left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -\frac{bx^2}{a}, \frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -\frac{bx^2}{a}, \frac{bx^2}{3a}\right)\right) + 9aF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{bx^2}{a}, \frac{bx^2}{3a}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((3*a - b*x^2)*(a + b*x^2)^(1/3)),x]

[Out] (9*a*x*AppellF1[1/2, 1/3, 1, 3/2, -((b*x^2)/a), (b*x^2)/(3*a)]/((3*a - b*x^2)*(a + b*x^2)^(1/3)*(9*a*AppellF1[1/2, 1/3, 1, 3/2, -((b*x^2)/a), (b*x^2)/(3*a)]/(3*a) + 2*b*x^2*(AppellF1[3/2, 1/3, 2, 5/2, -((b*x^2)/a), (b*x^2)/(3*a)] - AppellF1[3/2, 4/3, 1, 5/2, -((b*x^2)/a), (b*x^2)/(3*a)])))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+3*a)/(b*x^2+a)^(1/3),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(bx^2 + a)^{\frac{1}{3}}(bx^2 - 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+3*a)/(b*x^2+a)^(1/3),x, algorithm="giac")

[Out] integrate(-1/((b*x^2 + a)^(1/3)*(b*x^2 - 3*a)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + 3a)(bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+3*a)/(b*x^2+a)^(1/3),x)

[Out] int(1/(-b*x^2+3*a)/(b*x^2+a)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(bx^2 + a)^{\frac{1}{3}}(bx^2 - 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+3*a)/(b*x^2+a)^(1/3),x, algorithm="maxima")

[Out] -integrate(1/((b*x^2 + a)^(1/3)*(b*x^2 - 3*a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{1}{3}}(3a - bx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(1/3)*(3*a - b*x^2)),x)

[Out] `int(1/((a + b*x^2)^(1/3)*(3*a - b*x^2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{-3a\sqrt[3]{a+bx^2} + bx^2\sqrt[3]{a+bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-b*x**2+3*a)/(b*x**2+a)**(1/3), x)`

[Out] `-Integral(1/(-3*a*(a + b*x**2)**(1/3) + b*x**2*(a + b*x**2)**(1/3)), x)`

$$3.143 \quad \int \frac{1}{(c-dx^2)\sqrt[3]{c+3dx^2}} dx$$

Optimal. Leaf size=204

$$\frac{\sqrt{3} \tan^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt[6]{c}\left(\sqrt[3]{2}\sqrt[3]{c+3dx^2}+\sqrt[3]{c}\right)}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt[6]{c}\left(\sqrt[3]{c}-\sqrt[3]{2}\sqrt[3]{c+3dx^2}\right)}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt{c}}\right)}{2^{2/3}\sqrt{3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{c}}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}}$$

[Out] $-1/4*\operatorname{arctanh}(c^{(1/6)}*(c^{(1/3)}-2^{(1/3)}*(3*d*x^2+c)^{(1/3)})/x/d^{(1/2)})*2^{(1/3)}/c^{(5/6)}/d^{(1/2)}-1/4*\operatorname{arctanh}(1/x/d^{(1/2)}*c^{(1/2)})*2^{(1/3)}/c^{(5/6)}/d^{(1/2)}-1/12*\operatorname{arctan}(x*3^{(1/2)}*d^{(1/2)}/c^{(1/2)})*2^{(1/3)}/c^{(5/6)}*3^{(1/2)}/d^{(1/2)}+1/4*\operatorname{arctan}(x*3^{(1/2)}*d^{(1/2)}/c^{(1/6)})/(c^{(1/3)}+2^{(1/3)}*(3*d*x^2+c)^{(1/3)}))*3^{(1/2)}*2^{(1/3)}/c^{(5/6)}/d^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 204, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {392}

$$\frac{\sqrt{3} \tan^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt[6]{c}\left(\sqrt[3]{2}\sqrt[3]{c+3dx^2}+\sqrt[3]{c}\right)}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt[6]{c}\left(\sqrt[3]{c}-\sqrt[3]{2}\sqrt[3]{c+3dx^2}\right)}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt{c}}\right)}{2^{2/3}\sqrt{3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{c}}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}}$$

Antiderivative was successfully verified.

[In] Int[1/((c - d*x^2)*(c + 3*d*x^2)^(1/3)), x]

[Out] $-\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]]/(2*2^{(2/3)}*\operatorname{Sqrt}[3]*c^{(5/6)}*\operatorname{Sqrt}[d]) + (\operatorname{Sqrt}[3]*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[d]*x)/(c^{(1/6)}*(c^{(1/3)} + 2^{(1/3)}*(c + 3*d*x^2)^{(1/3)}))])/(2*2^{(2/3)}*c^{(5/6)}*\operatorname{Sqrt}[d]) - \operatorname{ArcTanh}[\operatorname{Sqrt}[c]/(\operatorname{Sqrt}[d]*x)]/(2*2^{(2/3)}*c^{(5/6)}*\operatorname{Sqrt}[d]) - \operatorname{ArcTanh}[(c^{(1/6)}*(c^{(1/3)} - 2^{(1/3)}*(c + 3*d*x^2)^{(1/3)}))/(\operatorname{Sqrt}[d]*x)]/(2*2^{(2/3)}*c^{(5/6)}*\operatorname{Sqrt}[d])$

Rule 392

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[b/a, 2]}, Simp[(q*ArcTanh[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (-Simp[(q*ArcTan[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3))])/(2*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTanh[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{(c-dx^2)\sqrt[3]{c+3dx^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt{c}}\right)}{2^{2/3}\sqrt{3}c^{5/6}\sqrt{d}} + \frac{\sqrt{3} \tan^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt[6]{c}\left(\sqrt[3]{c}+\sqrt[3]{2}\sqrt[3]{c+3dx^2}\right)}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{c}}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt[6]{c}\left(\sqrt[3]{c}-\sqrt[3]{2}\sqrt[3]{c+3dx^2}\right)}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}}$$

Mathematica [C] time = 0.15, size = 153, normalized size = 0.75

$$\frac{3cxF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{3dx^2}{c}, \frac{dx^2}{c}\right)}{(c-dx^2)\sqrt[3]{c+3dx^2}} \left(2dx^2\left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -\frac{3dx^2}{c}, \frac{dx^2}{c}\right) - F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -\frac{3dx^2}{c}, \frac{dx^2}{c}\right)\right) + 3cF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{3dx^2}{c}, \frac{dx^2}{c}\right)\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((c - d*x^2)*(c + 3*d*x^2)^(1/3)),x]

[Out] (3*c*x*AppellF1[1/2, 1/3, 1, 3/2, (-3*d*x^2)/c, (d*x^2)/c])/((c - d*x^2)*(c + 3*d*x^2)^(1/3)*(3*c*AppellF1[1/2, 1/3, 1, 3/2, (-3*d*x^2)/c, (d*x^2)/c] + 2*d*x^2*(AppellF1[3/2, 1/3, 2, 5/2, (-3*d*x^2)/c, (d*x^2)/c] - AppellF1[3/2, 4/3, 1, 5/2, (-3*d*x^2)/c, (d*x^2)/c])))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-d*x^2+c)/(3*d*x^2+c)^(1/3),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(3dx^2 + c)^{\frac{1}{3}}(dx^2 - c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-d*x^2+c)/(3*d*x^2+c)^(1/3),x, algorithm="giac")

[Out] integrate(-1/((3*d*x^2 + c)^(1/3)*(d*x^2 - c)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(-dx^2 + c)(3dx^2 + c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-d*x^2+c)/(3*d*x^2+c)^(1/3),x)

[Out] int(1/(-d*x^2+c)/(3*d*x^2+c)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(3dx^2 + c)^{\frac{1}{3}}(dx^2 - c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-d*x^2+c)/(3*d*x^2+c)^(1/3),x, algorithm="maxima")

[Out] -integrate(1/((3*d*x^2 + c)^(1/3)*(d*x^2 - c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(c - dx^2)(3dx^2 + c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c - d*x^2)*(c + 3*d*x^2)^(1/3)),x)

[Out] `int(1/((c - d*x^2)*(c + 3*d*x^2)^(1/3)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{-c\sqrt[3]{c+3dx^2} + dx^2\sqrt[3]{c+3dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-d*x**2+c)/(3*d*x**2+c)**(1/3), x)`

[Out] `-Integral(1/(-c*(c + 3*d*x**2)**(1/3) + d*x**2*(c + 3*d*x**2)**(1/3)), x)`

$$3.144 \quad \int \frac{1}{\sqrt[3]{a-bx^2} (3a+bx^2)} dx$$

Optimal. Leaf size=204

$$\frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt[6]{a} (\sqrt[3]{2} \sqrt[3]{a-bx^2} + \sqrt[3]{a})}\right)}{2 \cdot 2^{2/3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6 \cdot 2^{2/3} a^{5/6} \sqrt{b}}$$

[Out] $1/4 \cdot \arctan(x \cdot b^{1/2} / a^{1/6} / (a^{1/3} + 2^{1/3}) \cdot (-b \cdot x^2 + a)^{1/3}) \cdot 2^{1/3} / a^{5/6} / b^{1/2} - 1/12 \cdot \arctan(x \cdot b^{1/2} / a^{1/6}) \cdot 2^{1/3} / a^{5/6} / b^{1/2} + 1/12 \cdot \arctan(a^{1/6} \cdot (a^{1/3} - 2^{1/3}) \cdot (-b \cdot x^2 + a)^{1/3}) \cdot 3^{1/2} / x / b^{1/2} \cdot 2^{1/3} / a^{5/6} \cdot 3^{1/2} / b^{1/2} + 1/12 \cdot \arctan(3^{1/2} \cdot a^{1/2} / x / b^{1/2}) \cdot 2^{1/3} / a^{5/6} \cdot 3^{1/2} / b^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 204, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {393}

$$\frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt[6]{a} (\sqrt[3]{2} \sqrt[3]{a-bx^2} + \sqrt[3]{a})}\right)}{2 \cdot 2^{2/3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6 \cdot 2^{2/3} a^{5/6} \sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(1/3)*(3*a + b*x^2)), x]

[Out] ArcTan[(Sqrt[3]*Sqrt[a])/(Sqrt[b]*x)]/(2*2^(2/3)*Sqrt[3]*a^(5/6)*Sqrt[b]) + ArcTan[(Sqrt[3]*a^(1/6)*(a^(1/3) - 2^(1/3)*(a - b*x^2)^(1/3)))/(Sqrt[b]*x)]/(2*2^(2/3)*Sqrt[3]*a^(5/6)*Sqrt[b]) - ArcTanh[(Sqrt[b]*x)/Sqrt[a]]/(6*2^(2/3)*a^(5/6)*Sqrt[b]) + ArcTanh[(Sqrt[b]*x)/(a^(1/6)*(a^(1/3) + 2^(1/3)*(a - b*x^2)^(1/3)))]/(2*2^(2/3)*a^(5/6)*Sqrt[b])

Rule 393

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{a-bx^2} (3a+bx^2)} dx = \frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt{a}}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} (\sqrt[3]{a} - \sqrt[3]{2} \sqrt[3]{a-bx^2})}{\sqrt{bx}}\right)}{2 \cdot 2^{2/3} \sqrt{3} a^{5/6} \sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{6 \cdot 2^{2/3} a^{5/6} \sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{bx}}{\sqrt[6]{a} (\sqrt[3]{2} \sqrt[3]{a-bx^2} + \sqrt[3]{a})}\right)}{2 \cdot 2^{2/3} a^{5/6} \sqrt{b}}$$

Mathematica [C] time = 0.04, size = 162, normalized size = 0.79

$$\frac{9axF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}{\sqrt[3]{a-bx^2} (3a+bx^2) \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)\right) + 9aF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, -\frac{bx^2}{3a}\right)}\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(1/3)*(3*a + b*x^2)),x]

[Out] (9*a*x*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a])/((a - b*x^2)^(1/3)*(3*a + b*x^2)*(9*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, -1/3*(b*x^2)/a] + 2*b*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a] + AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, -1/3*(b*x^2)/a]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 3*a)*(-b*x^2 + a)^(1/3)), x)

maple [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{1}{3}}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x)

[Out] int(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 3a)(-bx^2 + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(b*x^2+3*a),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 3*a)*(-b*x^2 + a)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a - bx^2)^{1/3}(bx^2 + 3a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(1/3)*(3*a + b*x^2)),x)

[Out] `int(1/((a - b*x^2)^(1/3)*(3*a + b*x^2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{a - bx^2} (3a + bx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-b*x**2+a)**(1/3)/(b*x**2+3*a), x)`

[Out] `Integral(1/((a - b*x**2)**(1/3)*(3*a + b*x**2)), x)`

$$3.145 \quad \int \frac{1}{\sqrt[3]{c-3dx^2} (c+dx^2)} dx$$

Optimal. Leaf size=204

$$\frac{\tan^{-1}\left(\frac{\sqrt[6]{c}\left(\sqrt[3]{c}-\sqrt[3]{2}\sqrt[3]{c-3dx^2}\right)}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} + \frac{\sqrt{3}\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt[6]{c}\left(\sqrt[3]{2}\sqrt[3]{c-3dx^2}+\sqrt[3]{c}\right)}\right)}{2^{2/3}c^{5/6}\sqrt{d}} + \frac{\tan^{-1}\left(\frac{\sqrt{c}}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt{c}}\right)}{2^{2/3}\sqrt{3}c^{5/6}\sqrt{d}}$$

[Out] $1/4*\arctan(c^{(1/6)}*(c^{(1/3)}-2^{(1/3)}*(-3*d*x^2+c)^{(1/3)})/x/d^{(1/2)})*2^{(1/3)}/c^{(5/6)}/d^{(1/2)}+1/4*\arctan(1/x/d^{(1/2)}*c^{(1/2)})*2^{(1/3)}/c^{(5/6)}/d^{(1/2)}-1/2*\operatorname{arctanh}(x*3^{(1/2)}*d^{(1/2)}/c^{(1/2)})*2^{(1/3)}/c^{(5/6)}*3^{(1/2)}/d^{(1/2)}+1/4*\operatorname{arctanh}(x*3^{(1/2)}*d^{(1/2)}/c^{(1/6)})/(c^{(1/3)}+2^{(1/3)}*(-3*d*x^2+c)^{(1/3)})*3^{(1/2)}*2^{(1/3)}/c^{(5/6)}/d^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 204, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {393}

$$\frac{\tan^{-1}\left(\frac{\sqrt[6]{c}\left(\sqrt[3]{c}-\sqrt[3]{2}\sqrt[3]{c-3dx^2}\right)}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} + \frac{\sqrt{3}\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt[6]{c}\left(\sqrt[3]{2}\sqrt[3]{c-3dx^2}+\sqrt[3]{c}\right)}\right)}{2^{2/3}c^{5/6}\sqrt{d}} + \frac{\tan^{-1}\left(\frac{\sqrt{c}}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt{c}}\right)}{2^{2/3}\sqrt{3}c^{5/6}\sqrt{d}}$$

Antiderivative was successfully verified.

[In] Int[1/((c - 3*d*x^2)^(1/3)*(c + d*x^2)), x]

[Out] ArcTan[Sqrt[c]/(Sqrt[d]*x)]/(2*2^(2/3)*c^(5/6)*Sqrt[d]) + ArcTan[(c^(1/6)*(c^(1/3) - 2^(1/3)*(c - 3*d*x^2)^(1/3)))/(Sqrt[d]*x)]/(2*2^(2/3)*c^(5/6)*Sqrt[d]) - ArcTanh[(Sqrt[3]*Sqrt[d]*x)/Sqrt[c]]/(2*2^(2/3)*Sqrt[3]*c^(5/6)*Sqrt[d]) + (Sqrt[3]*ArcTanh[(Sqrt[3]*Sqrt[d]*x)/(c^(1/6)*(c^(1/3) + 2^(1/3)*(c - 3*d*x^2)^(1/3)))]/(2*2^(2/3)*c^(5/6)*Sqrt[d]))

Rule 393

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x])/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{c-3dx^2} (c+dx^2)} dx = \frac{\tan^{-1}\left(\frac{\sqrt{c}}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} + \frac{\tan^{-1}\left(\frac{\sqrt[6]{c}\left(\sqrt[3]{c}-\sqrt[3]{2}\sqrt[3]{c-3dx^2}\right)}{\sqrt{dx}}\right)}{2^{2/3}c^{5/6}\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt{c}}\right)}{2^{2/3}\sqrt{3}c^{5/6}\sqrt{d}} + \frac{\sqrt{3}\tanh^{-1}\left(\frac{\sqrt{3}\sqrt{dx}}{\sqrt[6]{c}\left(\sqrt[3]{2}\sqrt[3]{c-3dx^2}+\sqrt[3]{c}\right)}\right)}{2^{2/3}c^{5/6}\sqrt{d}}$$

Mathematica [C] time = 0.16, size = 156, normalized size = 0.76

$$\frac{3cx F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{3dx^2}{c}, -\frac{dx^2}{c}\right)}{\sqrt[3]{c-3dx^2} (c+dx^2)} \left(2dx^2 \left(F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{3dx^2}{c}, -\frac{dx^2}{c}\right) - F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{3dx^2}{c}, -\frac{dx^2}{c}\right)\right) + 3c F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{3dx^2}{c}, -\frac{dx^2}{c}\right)\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((c - 3*d*x^2)^(1/3)*(c + d*x^2)),x]

[Out] (3*c*x*AppellF1[1/2, 1/3, 1, 3/2, (3*d*x^2)/c, -((d*x^2)/c)]/((c - 3*d*x^2)^(1/3)*(c + d*x^2)*(3*c*AppellF1[1/2, 1/3, 1, 3/2, (3*d*x^2)/c, -((d*x^2)/c)] + 2*d*x^2*(-AppellF1[3/2, 1/3, 2, 5/2, (3*d*x^2)/c, -((d*x^2)/c)] + AppellF1[3/2, 4/3, 1, 5/2, (3*d*x^2)/c, -((d*x^2)/c)]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*d*x^2+c)^(1/3)/(d*x^2+c),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 + c)(-3dx^2 + c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*d*x^2+c)^(1/3)/(d*x^2+c),x, algorithm="giac")

[Out] integrate(1/((d*x^2 + c)*(-3*d*x^2 + c)^(1/3)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(-3dx^2 + c)^{\frac{1}{3}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*d*x^2+c)^(1/3)/(d*x^2+c),x)

[Out] int(1/(-3*d*x^2+c)^(1/3)/(d*x^2+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 + c)(-3dx^2 + c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*d*x^2+c)^(1/3)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/((d*x^2 + c)*(-3*d*x^2 + c)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(dx^2 + c)(c - 3dx^2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c + d*x^2)*(c - 3*d*x^2)^(1/3)),x)

[Out] `int(1/((c + d*x^2)*(c - 3*d*x^2)^(1/3)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{c - 3dx^2} (c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-3*d*x**2+c)**(1/3)/(d*x**2+c), x)`

[Out] `Integral(1/((c - 3*d*x**2)**(1/3)*(c + d*x**2)), x)`

$$3.146 \quad \int \frac{1}{\sqrt[3]{1-x^2} (3+x^2)} dx$$

Optimal. Leaf size=113

$$\frac{\tan^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1-x^2})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\tanh^{-1}\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2}+1}\right)}{2 \cdot 2^{2/3}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\tanh^{-1}(x)}{6 \cdot 2^{2/3}}$$

[Out] $-1/12 \cdot \arctan(x) \cdot 2^{1/3} + 1/4 \cdot \arctan(x/(1+2^{1/3} \cdot (-x^2+1)^{1/3})) \cdot 2^{1/3} + 1/12 \cdot \arctan(3^{1/2}/x) \cdot 2^{1/3} \cdot 3^{1/2} + 1/12 \cdot \arctan((1-2^{1/3}) \cdot (-x^2+1)^{1/3}) \cdot 3^{1/2}/x \cdot 2^{1/3} \cdot 3^{1/2}$

Rubi [A] time = 0.02, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {393}

$$\frac{\tan^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1-x^2})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\tanh^{-1}\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2}+1}\right)}{2 \cdot 2^{2/3}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\tanh^{-1}(x)}{6 \cdot 2^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[1/((1 - x^2)^(1/3)*(3 + x^2)),x]

[Out] ArcTan[Sqrt[3]/x]/(2*2^(2/3)*Sqrt[3]) + ArcTan[(Sqrt[3]*(1 - 2^(1/3)*(1 - x^2)^(1/3)))/x]/(2*2^(2/3)*Sqrt[3]) - ArcTanh[x]/(6*2^(2/3)) + ArcTanh[x/(1 + 2^(1/3)*(1 - x^2)^(1/3))]/(2*2^(2/3))

Rule 393

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, Simp[(q*ArcTan[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (Simp[(q*ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] - Simp[(q*ArcTanh[q*x]/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{1-x^2} (3+x^2)} dx = \frac{\tan^{-1}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\tan^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1-x^2})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\tanh^{-1}(x)}{6 \cdot 2^{2/3}} + \frac{\tanh^{-1}\left(\frac{x}{1+\sqrt[3]{2}\sqrt[3]{1-x^2}}\right)}{2 \cdot 2^{2/3}}$$

Mathematica [C] time = 0.04, size = 118, normalized size = 1.04

$$\frac{9x F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; x^2, -\frac{x^2}{3}\right)}{\sqrt[3]{1-x^2} (x^2+3) \left(2x^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; x^2, -\frac{x^2}{3}\right) - F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; x^2, -\frac{x^2}{3}\right)\right) - 9F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; x^2, -\frac{x^2}{3}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((1 - x^2)^(1/3)*(3 + x^2)),x]

```
[Out] (-9*x*AppellF1[1/2, 1/3, 1, 3/2, x^2, -1/3*x^2])/((1 - x^2)^(1/3)*(3 + x^2)
*(-9*AppellF1[1/2, 1/3, 1, 3/2, x^2, -1/3*x^2] + 2*x^2*(AppellF1[3/2, 1/3,
2, 5/2, x^2, -1/3*x^2] - AppellF1[3/2, 4/3, 1, 5/2, x^2, -1/3*x^2]))
```

fricas [B] time = 3.85, size = 1943, normalized size = 17.19

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="fricas")
```

```
[Out] -1/20736*432^(5/6)*sqrt(3)*log(10368*(6*2^(2/3)*(x^6 + 225*x^4 - 189*x^2 +
27) + 144*432^(1/6)*sqrt(3)*(x^5 - x^3) + (432^(5/6)*sqrt(3)*(7*x^3 - 3*x)
+ 216*2^(1/3)*(x^4 + 3*x^2))*(-x^2 + 1)^(2/3) - 72*(x^5 + 18*x^4 + 24*x^3 -
18*x^2 - 9*x)*(-x^2 + 1)^(1/3))/(x^6 + 9*x^4 + 27*x^2 + 27)) - 1/20736*432
^(5/6)*sqrt(3)*log(2592*(6*2^(2/3)*(x^6 + 225*x^4 - 189*x^2 + 27) + 144*432
^(1/6)*sqrt(3)*(x^5 - x^3) + (432^(5/6)*sqrt(3)*(7*x^3 - 3*x) + 216*2^(1/3)
*(x^4 + 3*x^2))*(-x^2 + 1)^(2/3) - 72*(x^5 + 18*x^4 + 24*x^3 - 18*x^2 - 9*x
)*(-x^2 + 1)^(1/3))/(x^6 + 9*x^4 + 27*x^2 + 27)) + 1/20736*432^(5/6)*sqrt(3
)*log(10368*(6*2^(2/3)*(x^6 + 225*x^4 - 189*x^2 + 27) - 144*432^(1/6)*sqrt(
3)*(x^5 - x^3) - (432^(5/6)*sqrt(3)*(7*x^3 - 3*x) - 216*2^(1/3)*(x^4 + 3*x^
2))*(-x^2 + 1)^(2/3) + 72*(x^5 - 18*x^4 + 24*x^3 + 18*x^2 - 9*x)*(-x^2 + 1)
^(1/3))/(x^6 + 9*x^4 + 27*x^2 + 27)) + 1/20736*432^(5/6)*sqrt(3)*log(2592*(
6*2^(2/3)*(x^6 + 225*x^4 - 189*x^2 + 27) - 144*432^(1/6)*sqrt(3)*(x^5 - x^3
) - (432^(5/6)*sqrt(3)*(7*x^3 - 3*x) - 216*2^(1/3)*(x^4 + 3*x^2))*(-x^2 + 1
)^(2/3) + 72*(x^5 - 18*x^4 + 24*x^3 + 18*x^2 - 9*x)*(-x^2 + 1)^(1/3))/(x^6
+ 9*x^4 + 27*x^2 + 27)) - 1/1296*432^(5/6)*arctan(1/36*(432^(5/6)*(x^5 - 18
*x^3 + 9*x)*(-x^2 + 1)^(1/3) + sqrt(3)*2^(1/3)*(432^(5/6)*(x^4 + 9*x^2)*(-x
^2 + 1)^(2/3) - 288*sqrt(3)*(2*x^4 - 3*x^2)*(-x^2 + 1)^(1/3) + 6*432^(1/6)*
(x^6 + 141*x^4 - 153*x^2 + 27)) - 648*432^(1/6)*(3*x^3 - x)*(-x^2 + 1)^(2/3
) - 72*sqrt(3)*(7*x^5 + 6*x^3 - 9*x))/(x^6 - 225*x^4 + 243*x^2 - 27)) - 1/2
592*432^(5/6)*arctan(-1/18*(sqrt(2)*(18*sqrt(3)*2^(2/3)*(29*x^11 + 879*x^9
- 12078*x^7 + 10638*x^5 - 3807*x^3 + 243*x) - 2*(-x^2 + 1)^(2/3)*(432^(5/6)
*(x^10 + 153*x^8 - 1701*x^6 + 459*x^4) - 216*sqrt(3)*2^(1/3)*(31*x^9 - 297*
x^7 - 27*x^5 - 27*x^3)) - 36*(-x^2 + 1)^(1/3)*(sqrt(3)*(x^11 + 1167*x^9 - 1
3158*x^7 + 17550*x^5 - 4779*x^3 + 243*x) - 8*sqrt(3)*(13*x^10 - 6*x^8 - 140
4*x^6 + 1350*x^4 - 81*x^2)) - 3*432^(1/6)*(x^12 + 7620*x^10 - 92115*x^8 + 1
69776*x^6 - 109269*x^4 + 16524*x^2 - 729))*sqrt((6*2^(2/3)*(x^6 + 225*x^4 -
189*x^2 + 27) + 144*432^(1/6)*sqrt(3)*(x^5 - x^3) + (432^(5/6)*sqrt(3)*(7*
x^3 - 3*x) + 216*2^(1/3)*(x^4 + 3*x^2))*(-x^2 + 1)^(2/3) - 72*(x^5 + 18*x^4
+ 24*x^3 - 18*x^2 - 9*x)*(-x^2 + 1)^(1/3))/(x^6 + 9*x^4 + 27*x^2 + 27)) -
216*(sqrt(3)*2^(2/3)*(x^10 + 144*x^8 - 918*x^6 + 2808*x^4 - 243*x^2) - 3*43
2^(1/6)*(31*x^9 - 568*x^7 + 1710*x^5 - 432*x^3 + 27*x))*(-x^2 + 1)^(2/3) -
18*sqrt(3)*(x^12 - 366*x^10 + 14535*x^8 - 42660*x^6 + 58239*x^4 - 14094*x^2
+ 729) + 144*sqrt(3)*(11*x^11 - 807*x^9 + 4518*x^7 - 5238*x^5 + 3807*x^3 -
243*x) - (-x^2 + 1)^(1/3)*(432^(5/6)*(x^11 - 1215*x^9 + 11754*x^7 - 21006*
x^5 + 5589*x^3 - 243*x) - 432*sqrt(3)*2^(1/3)*(13*x^10 - 120*x^8 + 1242*x^6
- 1728*x^4 + 81*x^2)))/(x^12 - 8334*x^10 + 110727*x^8 - 301860*x^6 + 18783
9*x^4 - 21870*x^2 + 729)) - 1/2592*432^(5/6)*arctan(1/18*(sqrt(2)*(18*sqrt(
3)*2^(2/3)*(29*x^11 + 879*x^9 - 12078*x^7 + 10638*x^5 - 3807*x^3 + 243*x) +
2*(-x^2 + 1)^(2/3)*(432^(5/6)*(x^10 + 153*x^8 - 1701*x^6 + 459*x^4) + 216*
sqrt(3)*2^(1/3)*(31*x^9 - 297*x^7 - 27*x^5 - 27*x^3)) - 36*(-x^2 + 1)^(1/3)
*(sqrt(3)*(x^11 + 1167*x^9 - 13158*x^7 + 17550*x^5 - 4779*x^3 + 243*x) + 8*
sqrt(3)*(13*x^10 - 6*x^8 - 1404*x^6 + 1350*x^4 - 81*x^2)) + 3*432^(1/6)*(x^
12 + 7620*x^10 - 92115*x^8 + 169776*x^6 - 109269*x^4 + 16524*x^2 - 729))*sq
rt((6*2^(2/3)*(x^6 + 225*x^4 - 189*x^2 + 27) - 144*432^(1/6)*sqrt(3)*(x^5 -
x^3) - (432^(5/6)*sqrt(3)*(7*x^3 - 3*x) - 216*2^(1/3)*(x^4 + 3*x^2))*(-x^2
+ 1)^(2/3) + 72*(x^5 - 18*x^4 + 24*x^3 + 18*x^2 - 9*x)*(-x^2 + 1)^(1/3))/(
x^6 + 9*x^4 + 27*x^2 + 27)) - 216*(sqrt(3)*2^(2/3)*(x^10 + 144*x^8 - 918*x^
6 + 2808*x^4 - 243*x^2) + 3*432^(1/6)*(31*x^9 - 568*x^7 + 1710*x^5 - 432*x^
```

$(3 + 27x))(-x^2 + 1)^{2/3} - 18\sqrt{3}(x^{12} - 366x^{10} + 14535x^8 - 42660x^6 + 58239x^4 - 14094x^2 + 729) - 144\sqrt{3}(11x^{11} - 807x^9 + 4518x^7 - 5238x^5 + 3807x^3 - 243x) + (-x^2 + 1)^{1/3}(432^{5/6}(x^{11} - 1215x^9 + 11754x^7 - 21006x^5 + 5589x^3 - 243x) + 432\sqrt{3}2^{1/3}(13x^{10} - 120x^8 + 1242x^6 - 1728x^4 + 81x^2)))/(x^{12} - 8334x^{10} + 10727x^8 - 301860x^6 + 187839x^4 - 21870x^2 + 729))$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(x^2 + 3)(-x^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="giac")

[Out] integrate(1/((x^2 + 3)*(-x^2 + 1)^(1/3)), x)

maple [C] time = 16.87, size = 938, normalized size = 8.30

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/3)/(x^2+3),x)

[Out] $\frac{1}{36}\sqrt[3]{Z^6+108}\ln(-(-4050\sqrt[3]{Z^6+108})x^4+\sqrt[3]{Z^6+108})^4x^6+225\sqrt[3]{Z^6+108}^4x^4-72x^5\sqrt[3]{Z^6+108}^4+1296x^5\sqrt[3]{Z^6+108}+72x^3\sqrt[3]{Z^6+108}^4-189\sqrt[3]{Z^6+108}^4x^2+3402\sqrt[3]{Z^6+108}x^2-1296\sqrt[3]{Z^6+108}x^3+6\sqrt[3]{Z^6+108}^5(-x^2+1)^{1/3}x^5-108\sqrt[3]{Z^6+108}^5(-x^2+1)^{1/3}x^4+144\sqrt[3]{Z^6+108}^5(-x^2+1)^{1/3}x^3+108\sqrt[3]{Z^6+108}^5(-x^2+1)^{1/3}x^2-36\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x^5-54\sqrt[3]{Z^6+108}^5(-x^2+1)^{1/3}x+648\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x^4-864\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x^3-648\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x^2+324\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x-486\sqrt[3]{Z^6+108}+27\sqrt[3]{Z^6+108}^4-1296(-x^2+1)^{2/3}x^4+9072(-x^2+1)^{2/3}x^3-3888(-x^2+1)^{2/3}x^2-3888(-x^2+1)^{2/3}x-18\sqrt[3]{Z^6+108}x^6)/(x^2+3)^3-1/432\ln((\sqrt[3]{Z^6+108})^4x^6+72x^5\sqrt[3]{Z^6+108}^4+225\sqrt[3]{Z^6+108}^4x^4+36\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x^5-72x^3\sqrt[3]{Z^6+108}^4+648\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x^4-189\sqrt[3]{Z^6+108}^4x^2+864\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x^3+648(-x^2+1)^{2/3}x^4-648\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x^2+4536(-x^2+1)^{2/3}x^3+27\sqrt[3]{Z^6+108}^4-324\sqrt[3]{Z^6+108}^2(-x^2+1)^{1/3}x+1944(-x^2+1)^{2/3}x^2-1944(-x^2+1)^{2/3}x)/((x^2+3)^3)\sqrt[3]{Z^6+108}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(x^2 + 3)(-x^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="maxima")

[Out] integrate(1/((x^2 + 3)*(-x^2 + 1)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(1-x^2)^{1/3} (x^2+3)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/3)*(x^2 + 3)), x)

[Out] int(1/((1 - x^2)^(1/3)*(x^2 + 3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{-(x-1)(x+1)} (x^2+3)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/3)/(x**2+3), x)

[Out] Integral(1/((-x - 1)*(x + 1))**(1/3)*(x**2 + 3)), x)

$$3.147 \quad \int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx$$

Optimal. Leaf size=109

$$\frac{\tan^{-1}\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{x^2+1}}\right)}{2 \cdot 2^{2/3}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{x^2+1})}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} - \frac{\tan^{-1}(x)}{6 \cdot 2^{2/3}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}}$$

[Out] $-1/12*\arctan(x)*2^{(1/3)}+1/4*\arctan(x/(1+2^{(1/3)}*(x^2+1)^{(1/3)}))*2^{(1/3)}-1/12*\arctanh(3^{(1/2)}/x)*2^{(1/3)}*3^{(1/2)}-1/12*\arctanh((1-2^{(1/3)}*(x^2+1)^{(1/3)}))*3^{(1/2)}/x)*2^{(1/3)}*3^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {392}

$$\frac{\tan^{-1}\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{x^2+1}}\right)}{2 \cdot 2^{2/3}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{x^2+1})}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} - \frac{\tan^{-1}(x)}{6 \cdot 2^{2/3}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[1/((3 - x^2)*(1 + x^2)^(1/3)),x]

[Out] $-\text{ArcTan}[x]/(6*2^{(2/3)}) + \text{ArcTan}[x/(1 + 2^{(1/3)}*(1 + x^2)^{(1/3)})]/(2*2^{(2/3)}) - \text{ArcTanh}[\text{Sqrt}[3]/x]/(2*2^{(2/3)}*\text{Sqrt}[3]) - \text{ArcTanh}[(\text{Sqrt}[3]*(1 - 2^{(1/3)}*(1 + x^2)^{(1/3)}))/x]/(2*2^{(2/3)}*\text{Sqrt}[3])$

Rule 392

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[(q*ArcTanh[Sqrt[3]/(q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x] + (-Simp[(q*ArcTan[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3)))]/(2*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTan[q*x]/(6*2^(2/3)*a^(1/3)*d), x] + Simp[(q*ArcTanh[(Sqrt[3]*(a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)])/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx = -\frac{\tan^{-1}(x)}{6 \cdot 2^{2/3}} + \frac{\tan^{-1}\left(\frac{x}{1+\sqrt[3]{2}\sqrt[3]{1+x^2}}\right)}{2 \cdot 2^{2/3}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1+x^2})}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}}$$

Mathematica [C] time = 0.04, size = 124, normalized size = 1.14

$$\frac{9x F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -x^2, \frac{x^2}{3}\right)}{(x^2-3)\sqrt[3]{x^2+1} \left(2x^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -x^2, \frac{x^2}{3}\right) - F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -x^2, \frac{x^2}{3}\right)\right) + 9F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -x^2, \frac{x^2}{3}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((3 - x^2)*(1 + x^2)^(1/3)),x]

```
[Out] (-9*x*AppellF1[1/2, 1/3, 1, 3/2, -x^2, x^2/3])/((-3 + x^2)*(1 + x^2)^(1/3)*
(9*AppellF1[1/2, 1/3, 1, 3/2, -x^2, x^2/3] + 2*x^2*(AppellF1[3/2, 1/3, 2, 5
/2, -x^2, x^2/3] - AppellF1[3/2, 4/3, 1, 5/2, -x^2, x^2/3]))
```

fricas [B] time = 3.20, size = 1685, normalized size = 15.46

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="fricas")
```

```
[Out] 1/2592*432^(5/6)*sqrt(3)*arctan(-1/54*(2592*x^11 - 393984*x^9 - 699840*x^7
- 373248*x^5 - 69984*x^3 - sqrt(6)*(18*sqrt(3)*2^(2/3)*(19*x^11 + 111*x^9 +
6030*x^7 + 7182*x^5 + 2511*x^3 + 243*x) + 3*432^(1/6)*sqrt(3)*(x^12 + 924*
x^10 - 33363*x^8 - 60912*x^6 - 36693*x^4 - 8748*x^2 - 729) + (432^(5/6)*sqr
t(3)*(x^10 - 78*x^8 - 720*x^6 - 594*x^4 - 81*x^2) + 432*sqrt(3)*2^(1/3)*(13
*x^9 - 177*x^7 - 153*x^5 - 27*x^3))*(x^2 + 1)^(2/3) + 36*(96*x^10 - 4032*x^
8 - 2592*x^6 + sqrt(3)*(x^11 + 369*x^9 - 3654*x^7 - 5454*x^5 - 2187*x^3 - 2
43*x))*(x^2 + 1)^(1/3))*sqrt((2*2^(2/3)*(x^6 - 57*x^4 - 117*x^2 - 27) + (x^
2 + 1)^(2/3)*(432^(5/6)*(x^3 + x) + 24*2^(1/3)*(x^4 + 9*x^2)) - 8*(6*x^4 -
18*x^2 + sqrt(3)*(x^5 - 9*x))*(x^2 + 1)^(1/3) - 8*432^(1/6)*(x^5 + 18*x^3 +
9*x))/(x^6 - 9*x^4 + 27*x^2 - 27)) + 216*(sqrt(3)*2^(2/3)*(x^10 + 276*x^8
+ 1206*x^6 + 756*x^4 + 81*x^2) + 432^(1/6)*sqrt(3)*(31*x^9 - 1620*x^7 - 207
0*x^5 - 756*x^3 - 81*x))*(x^2 + 1)^(2/3) + 18*sqrt(3)*(x^12 + 1422*x^10 + 2
1447*x^8 + 27108*x^6 + 16767*x^4 + 6318*x^2 + 729) + (432^(5/6)*sqrt(3)*(x^
11 - 681*x^9 + 4338*x^7 + 6102*x^5 + 2349*x^3 + 243*x) + 3888*sqrt(3)*2^(1/
3)*(x^10 + 44*x^8 + 94*x^6 + 60*x^4 + 9*x^2))*(x^2 + 1)^(1/3))/(x^12 - 2178
*x^10 + 46791*x^8 + 83268*x^6 + 47871*x^4 + 10206*x^2 + 729)) + 1/2592*432^
(5/6)*sqrt(3)*arctan(-1/54*(2592*x^11 - 393984*x^9 - 699840*x^7 - 373248*x^
5 - 69984*x^3 + sqrt(6)*(18*sqrt(3)*2^(2/3)*(19*x^11 + 111*x^9 + 6030*x^7 +
7182*x^5 + 2511*x^3 + 243*x) - 3*432^(1/6)*sqrt(3)*(x^12 + 924*x^10 - 3336
3*x^8 - 60912*x^6 - 36693*x^4 - 8748*x^2 - 729) - (432^(5/6)*sqrt(3)*(x^10
- 78*x^8 - 720*x^6 - 594*x^4 - 81*x^2) - 432*sqrt(3)*2^(1/3)*(13*x^9 - 177*
x^7 - 153*x^5 - 27*x^3))*(x^2 + 1)^(2/3) - 36*(96*x^10 - 4032*x^8 - 2592*x^
6 - sqrt(3)*(x^11 + 369*x^9 - 3654*x^7 - 5454*x^5 - 2187*x^3 - 243*x))*(x^2
+ 1)^(1/3))*sqrt((2*2^(2/3)*(x^6 - 57*x^4 - 117*x^2 - 27) - (x^2 + 1)^(2/3
))*(432^(5/6)*(x^3 + x) - 24*2^(1/3)*(x^4 + 9*x^2)) - 8*(6*x^4 - 18*x^2 - sq
rt(3)*(x^5 - 9*x))*(x^2 + 1)^(1/3) + 8*432^(1/6)*(x^5 + 18*x^3 + 9*x))/(x^6
- 9*x^4 + 27*x^2 - 27)) - 216*(sqrt(3)*2^(2/3)*(x^10 + 276*x^8 + 1206*x^6
+ 756*x^4 + 81*x^2) - 432^(1/6)*sqrt(3)*(31*x^9 - 1620*x^7 - 2070*x^5 - 756
*x^3 - 81*x))*(x^2 + 1)^(2/3) - 18*sqrt(3)*(x^12 + 1422*x^10 + 21447*x^8 +
27108*x^6 + 16767*x^4 + 6318*x^2 + 729) + (432^(5/6)*sqrt(3)*(x^11 - 681*x^
9 + 4338*x^7 + 6102*x^5 + 2349*x^3 + 243*x) - 3888*sqrt(3)*2^(1/3)*(x^10 +
44*x^8 + 94*x^6 + 60*x^4 + 9*x^2))*(x^2 + 1)^(1/3))/(x^12 - 2178*x^10 + 467
91*x^8 + 83268*x^6 + 47871*x^4 + 10206*x^2 + 729)) + 1/5184*432^(5/6)*log(-
(432^(5/6)*(x^6 + 69*x^4 + 63*x^2 + 27) + 864*(9*x^3 + sqrt(3)*(x^4 + 9*x^2
) + 9*x)*(x^2 + 1)^(2/3) + 432*2^(1/3)*(5*x^5 + 30*x^3 + 9*x) + 432*(x^2 +
1)^(1/3)*(2^(2/3)*(x^5 + 18*x^3 + 9*x) + 4*432^(1/6)*(x^4 + 3*x^2)))/(x^6 -
9*x^4 + 27*x^2 - 27)) - 1/5184*432^(5/6)*log((432^(5/6)*(x^6 + 69*x^4 + 63
*x^2 + 27) - 864*(9*x^3 - sqrt(3)*(x^4 + 9*x^2) + 9*x)*(x^2 + 1)^(2/3) - 43
2*2^(1/3)*(5*x^5 + 30*x^3 + 9*x) - 432*(x^2 + 1)^(1/3)*(2^(2/3)*(x^5 + 18*x
^3 + 9*x) - 4*432^(1/6)*(x^4 + 3*x^2)))/(x^6 - 9*x^4 + 27*x^2 - 27)) - 1/10
368*432^(5/6)*log(31104*(2*2^(2/3)*(x^6 - 57*x^4 - 117*x^2 - 27) + (x^2 + 1
)^(2/3)*(432^(5/6)*(x^3 + x) + 24*2^(1/3)*(x^4 + 9*x^2)) - 8*(6*x^4 - 18*x^
2 + sqrt(3)*(x^5 - 9*x))*(x^2 + 1)^(1/3) - 8*432^(1/6)*(x^5 + 18*x^3 + 9*x)
)/(x^6 - 9*x^4 + 27*x^2 - 27)) + 1/10368*432^(5/6)*log(31104*(2*2^(2/3)*(x^
6 - 57*x^4 - 117*x^2 - 27) - (x^2 + 1)^(2/3)*(432^(5/6)*(x^3 + x) - 24*2^(1
/3)*(x^4 + 9*x^2)) - 8*(6*x^4 - 18*x^2 - sqrt(3)*(x^5 - 9*x))*(x^2 + 1)^(1/
3) + 8*432^(1/6)*(x^5 + 18*x^3 + 9*x))/(x^6 - 9*x^4 + 27*x^2 - 27))
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(x^2 + 1)^{\frac{1}{3}}(x^2 - 3)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="giac")

[Out] integrate(-1/((x^2 + 1)^(1/3)*(x^2 - 3)), x)

maple [F(-1)] time = 180.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-x^2 + 3)(x^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+3)/(x^2+1)^(1/3),x)

[Out] int(1/(-x^2+3)/(x^2+1)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(x^2 + 1)^{\frac{1}{3}}(x^2 - 3)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="maxima")

[Out] -integrate(1/((x^2 + 1)^(1/3)*(x^2 - 3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(x^2 + 1)^{1/3} (x^2 - 3)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((x^2 + 1)^(1/3)*(x^2 - 3)),x)

[Out] -int(1/((x^2 + 1)^(1/3)*(x^2 - 3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{x^2 \sqrt[3]{x^2 + 1} - 3 \sqrt[3]{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+3)/(x**2+1)**(1/3),x)

[Out] -Integral(1/(x**2*(x**2 + 1)**(1/3) - 3*(x**2 + 1)**(1/3)), x)

$$3.148 \quad \int \frac{3-x}{\sqrt[3]{1-x^2} (3+x^2)} dx$$

Optimal. Leaf size=96

$$-\frac{\log(x^2+3)}{2 \cdot 2^{2/3}} + \frac{3 \log((x+1)^{2/3} + \sqrt[3]{2} \sqrt[3]{1-x})}{2 \cdot 2^{2/3}} - \frac{\sqrt{3} \tan^{-1}\left(\frac{1}{\sqrt{3}} - \frac{2^{2/3}(x+1)^{2/3}}{\sqrt{3} \sqrt[3]{1-x}}\right)}{2^{2/3}}$$

[Out] $-1/4 \cdot \ln(x^2+3) \cdot 2^{(1/3)} + 3/4 \cdot \ln(2^{(1/3)} \cdot (1-x)^{(1/3)} + (1+x)^{(2/3)}) \cdot 2^{(1/3)} + 1/2 \cdot \arctan(-1/3 \cdot 3^{(1/2)} + 1/3 \cdot 2^{(2/3)} \cdot (1+x)^{(2/3)} / ((1-x)^{(1/3)} \cdot 3^{(1/2)})) \cdot 3^{(1/2)} \cdot 2^{(1/3)}$

Rubi [A] time = 0.02, antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {1008}

$$-\frac{\log(x^2+3)}{2 \cdot 2^{2/3}} + \frac{3 \log((x+1)^{2/3} + \sqrt[3]{2} \sqrt[3]{1-x})}{2 \cdot 2^{2/3}} - \frac{\sqrt{3} \tan^{-1}\left(\frac{1}{\sqrt{3}} - \frac{2^{2/3}(x+1)^{2/3}}{\sqrt{3} \sqrt[3]{1-x}}\right)}{2^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[(3 - x)/((1 - x^2)^(1/3)*(3 + x^2)), x]

[Out] $-\left(\frac{\sqrt{3} \operatorname{ArcTan}\left[\frac{1}{\sqrt{3}} - \frac{2^{(2/3)}(1+x)^{(2/3)}}{\sqrt{3}(1-x)^{(1/3)}}\right]}{2^{(2/3)}} - \frac{\operatorname{Log}[3+x^2]}{2 \cdot 2^{(2/3)}} + \frac{3 \operatorname{Log}\left[2^{(1/3)}(1-x)^{(1/3)} + (1+x)^{(2/3)}\right]}{2 \cdot 2^{(2/3)}}\right)$

Rule 1008

Int[((g_) + (h_.)*(x_))/(((a_) + (c_.)*(x_)^2)^(1/3)*((d_) + (f_.)*(x_)^2)), x_Symbol] :> Simp[(Sqrt[3]*h*ArcTan[1/Sqrt[3] - (2^(2/3)*(1 - (3*h*x)/g)^(2/3))/(Sqrt[3]*(1 + (3*h*x)/g)^(1/3))]/(2^(2/3)*a^(1/3)*f), x] + (-Simp[(3*h*Log[(1 - (3*h*x)/g)^(2/3) + 2^(1/3)*(1 + (3*h*x)/g)^(1/3)])/(2^(5/3)*a^(1/3)*f), x] + Simp[(h*Log[d + f*x^2])/(2^(5/3)*a^(1/3)*f), x]) /; FreeQ[{a, c, d, f, g, h}, x] && EqQ[c*d + 3*a*f, 0] && EqQ[c*g^2 + 9*a*h^2, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{3-x}{\sqrt[3]{1-x^2} (3+x^2)} dx = -\frac{\sqrt{3} \tan^{-1}\left(\frac{1}{\sqrt{3}} - \frac{2^{2/3}(1+x)^{2/3}}{\sqrt{3} \sqrt[3]{1-x}}\right)}{2^{2/3}} - \frac{\log(3+x^2)}{2 \cdot 2^{2/3}} + \frac{3 \log(\sqrt[3]{2} \sqrt[3]{1-x} + (1+x)^{2/3})}{2 \cdot 2^{2/3}}$$

Mathematica [C] time = 0.15, size = 143, normalized size = 1.49

$$-\frac{1}{6} x^2 F_1\left(1; \frac{1}{3}, 1, 2; x^2, -\frac{x^2}{3}\right) - \frac{27 x F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; x^2, -\frac{x^2}{3}\right)}{\sqrt[3]{1-x^2} (x^2+3)} \left(2 x^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; x^2, -\frac{x^2}{3}\right) - F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; x^2, -\frac{x^2}{3}\right)\right) - 9 F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; x^2, -\frac{x^2}{3}\right)\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[(3 - x)/((1 - x^2)^(1/3)*(3 + x^2)), x]

[Out] $-1/6 \cdot (x^2 \operatorname{AppellF1}[1, 1/3, 1, 2, x^2, -1/3 \cdot x^2]) - (27 \cdot x \operatorname{AppellF1}[1/2, 1/3, 1, 3/2, x^2, -1/3 \cdot x^2]) / ((1 - x^2)^{(1/3)} \cdot (3 + x^2) \cdot (-9 \operatorname{AppellF1}[1/2, 1/3, 1, 3/2, x^2, -1/3 \cdot x^2]))$

1, 3/2, x^2, -1/3*x^2] + 2*x^2*(AppellF1[3/2, 1/3, 2, 5/2, x^2, -1/3*x^2] - AppellF1[3/2, 4/3, 1, 5/2, x^2, -1/3*x^2]))

fricas [B] time = 10.67, size = 285, normalized size = 2.97

$$-\frac{1}{6} \cdot 4^{\frac{1}{6}} \sqrt{3} \arctan \left(\frac{4^{\frac{1}{6}} \sqrt{3} \left(12 \cdot 4^{\frac{2}{3}} (x^4 + 3x^3 + 3x^2 + 9x) (-x^2 + 1)^{\frac{2}{3}} + 4^{\frac{1}{3}} (x^6 - 18x^5 - 117x^4 - 36x^3 + 207x^2 + 54x - 27) \right)}{6 (x^6 + 54x^5 + 171x^4 + 108x^3 - 81x^2 - 162x - 27)} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3-x)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="fricas")

[Out] -1/6*4^(1/6)*sqrt(3)*arctan(1/6*4^(1/6)*sqrt(3)*(12*4^(2/3)*(x^4 + 3*x^3 + 3*x^2 + 9*x)*(-x^2 + 1)^(2/3) + 4^(1/3)*(x^6 - 18*x^5 - 117*x^4 - 36*x^3 + 207*x^2 + 54*x - 27) + 12*(x^5 + 19*x^4 + 42*x^3 + 6*x^2 - 27*x - 9)*(-x^2 + 1)^(1/3))/(x^6 + 54*x^5 + 171*x^4 + 108*x^3 - 81*x^2 - 162*x - 27)) - 1/2*4^(2/3)*log((6*4^(2/3)*(x^2 + 3*x)*(-x^2 + 1)^(2/3) + 4^(1/3)*(x^4 + 18*x^3 + 24*x^2 - 18*x - 9) - 6*(x^3 + 7*x^2 + 3*x - 3)*(-x^2 + 1)^(1/3))/(x^4 + 6*x^2 + 9)) + 1/12*4^(2/3)*log((4^(2/3)*(x^2 + 3) + 6*4^(1/3)*(-x^2 + 1)^(1/3)*(x + 1) + 12*(-x^2 + 1)^(2/3))/(x^2 + 3))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{x-3}{(x^2+3)(-x^2+1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3-x)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="giac")

[Out] integrate(-(x - 3)/((x^2 + 3)*(-x^2 + 1)^(1/3)), x)

maple [C] time = 7.83, size = 1033, normalized size = 10.76

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x+3)/(-x^2+1)^(1/3)/(x^2+3),x)

[Out] 1/2*RootOf(_Z^3-2)*ln((12*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)^2*RootOf(_Z^3-2)^2*x^2+2*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^3*x^2+36*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)^2*RootOf(_Z^3-2)^2*x+6*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^3*x-18*(-x^2+1)^(2/3)*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^2-12*(-x^2+1)^(1/3)*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)*x-9*(-x^2+1)^(1/3)*RootOf(_Z^3-2)^2*x-12*(-x^2+1)^(1/3)*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)+6*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*x^2-9*(-x^2+1)^(1/3)*RootOf(_Z^3-2)^2+RootOf(_Z^3-2)*x^2+36*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*x+6*RootOf(_Z^3-2)*x-6*(-x^2+1)^(2/3)-18*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)-3*RootOf(_Z^3-2))/(2*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^2*x+x-3)/(2*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^2*x+x+3))+RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*ln(-(4*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)^2*RootOf(_Z^3-2)^2*x^2+6*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^3*x^2+12*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)^2*RootOf(_Z^3-2)^2*x+18*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^3*x-18*(-x^2+1

)^(2/3)*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^2-6*(-x^2+1)^(1/3)*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)*x-9*(-x^2+1)^(1/3)*RootOf(_Z^3-2)^2*x-6*(-x^2+1)^(1/3)*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)+2*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*x^2-9*(-x^2+1)^(1/3)*RootOf(_Z^3-2)^2+3*RootOf(_Z^3-2)*x^2-12*(-x^2+1)^(2/3)+6*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)+9*RootOf(_Z^3-2))/(2*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^2*x+x-3)/(2*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^2*x+x+3))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{x-3}{(x^2+3)(-x^2+1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3-x)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="maxima")

[Out] -integrate((x-3)/((x^2+3)*(-x^2+1)^(1/3)),x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{x-3}{(1-x^2)^{1/3}(x^2+3)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-(x-3)/((1-x^2)^(1/3)*(x^2+3)),x)

[Out] -int((x-3)/((1-x^2)^(1/3)*(x^2+3)),x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{x}{x^2\sqrt[3]{1-x^2}+3\sqrt[3]{1-x^2}} dx - \int \left(-\frac{3}{x^2\sqrt[3]{1-x^2}+3\sqrt[3]{1-x^2}} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3-x)/(-x**2+1)**(1/3)/(x**2+3),x)

[Out] -Integral(x/(x**2*(1-x**2)**(1/3)+3*(1-x**2)**(1/3)),x) - Integral(-3/(x**2*(1-x**2)**(1/3)+3*(1-x**2)**(1/3)),x)

$$3.149 \quad \int \frac{3+x}{\sqrt[3]{1-x^2} (3+x^2)} dx$$

Optimal. Leaf size=95

$$\frac{\log(x^2+3)}{2 \cdot 2^{2/3}} - \frac{3 \log((1-x)^{2/3} + \sqrt[3]{2} \sqrt[3]{x+1})}{2 \cdot 2^{2/3}} + \frac{\sqrt{3} \tan^{-1}\left(\frac{1}{\sqrt{3}} - \frac{2^{2/3}(1-x)^{2/3}}{\sqrt{3} \sqrt[3]{x+1}}\right)}{2^{2/3}}$$

[Out] 1/4*ln(x^2+3)*2^(1/3)-3/4*ln((1-x)^(2/3)+2^(1/3)*(1+x)^(1/3))*2^(1/3)-1/2*arctan(-1/3*3^(1/2)+1/3*2^(2/3)*(1-x)^(2/3)/(1+x)^(1/3)*3^(1/2))*3^(1/2)*2^(1/3)

Rubi [A] time = 0.02, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {1008}

$$\frac{\log(x^2+3)}{2 \cdot 2^{2/3}} - \frac{3 \log((1-x)^{2/3} + \sqrt[3]{2} \sqrt[3]{x+1})}{2 \cdot 2^{2/3}} + \frac{\sqrt{3} \tan^{-1}\left(\frac{1}{\sqrt{3}} - \frac{2^{2/3}(1-x)^{2/3}}{\sqrt{3} \sqrt[3]{x+1}}\right)}{2^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[(3 + x)/((1 - x^2)^(1/3)*(3 + x^2)), x]

[Out] (Sqrt[3]*ArcTan[1/Sqrt[3] - (2^(2/3)*(1 - x)^(2/3))/(Sqrt[3]*(1 + x)^(1/3))])/2^(2/3) + Log[3 + x^2]/(2*2^(2/3)) - (3*Log[(1 - x)^(2/3) + 2^(1/3)*(1 + x)^(1/3)])/(2*2^(2/3))

Rule 1008

Int[((g_) + (h_.)*(x_))/(((a_) + (c_.)*(x_)^2)^(1/3)*((d_) + (f_.)*(x_)^2)), x_Symbol] := Simp[(Sqrt[3]*h*ArcTan[1/Sqrt[3] - (2^(2/3)*(1 - (3*h*x)/g)^(2/3))/(Sqrt[3]*(1 + (3*h*x)/g)^(1/3))]/(2^(2/3)*a^(1/3)*f), x] + (-Simp[(3*h*Log[(1 - (3*h*x)/g)^(2/3) + 2^(1/3)*(1 + (3*h*x)/g)^(1/3)])/(2^(5/3)*a^(1/3)*f), x] + Simp[(h*Log[d + f*x^2])/(2^(5/3)*a^(1/3)*f), x]) /; FreeQ[{a, c, d, f, g, h}, x] && EqQ[c*d + 3*a*f, 0] && EqQ[c*g^2 + 9*a*h^2, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{3+x}{\sqrt[3]{1-x^2} (3+x^2)} dx = \frac{\sqrt{3} \tan^{-1}\left(\frac{1}{\sqrt{3}} - \frac{2^{2/3}(1-x)^{2/3}}{\sqrt{3} \sqrt[3]{1+x}}\right)}{2^{2/3}} + \frac{\log(3+x^2)}{2 \cdot 2^{2/3}} - \frac{3 \log((1-x)^{2/3} + \sqrt[3]{2} \sqrt[3]{1+x})}{2 \cdot 2^{2/3}}$$

Mathematica [C] time = 0.11, size = 143, normalized size = 1.51

$$\frac{1}{6} x^2 F_1\left(1; \frac{1}{3}, 1; 2; x^2, -\frac{x^2}{3}\right) - \frac{27 x F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; x^2, -\frac{x^2}{3}\right)}{\sqrt[3]{1-x^2} (x^2+3) \left(2 x^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; x^2, -\frac{x^2}{3}\right) - F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; x^2, -\frac{x^2}{3}\right)\right) - 9 F_1\left(\frac{1}{2}; \frac{1}{3}, 1;\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(3 + x)/((1 - x^2)^(1/3)*(3 + x^2)), x]

[Out] (x^2*AppellF1[1, 1/3, 1, 2, x^2, -1/3*x^2])/6 - (27*x*AppellF1[1/2, 1/3, 1, 3/2, x^2, -1/3*x^2])/((1 - x^2)^(1/3)*(3 + x^2)*(-9*AppellF1[1/2, 1/3, 1,

$3/2, x^2, -1/3*x^2] + 2*x^2*(\text{AppellF1}[3/2, 1/3, 2, 5/2, x^2, -1/3*x^2] - \text{AppellF1}[3/2, 4/3, 1, 5/2, x^2, -1/3*x^2]))$

fricas [B] time = 9.41, size = 315, normalized size = 3.32

$$-\frac{1}{6} \cdot 4^{\frac{1}{6}} \sqrt{3} (-1)^{\frac{1}{3}} \arctan \left(\frac{4^{\frac{1}{6}} \sqrt{3} \left(12 \cdot 4^{\frac{2}{3}} (-1)^{\frac{2}{3}} (x^4 - 3x^3 + 3x^2 - 9x) (-x^2 + 1)^{\frac{2}{3}} + 12 (-1)^{\frac{1}{3}} (x^5 - 19x^4 + 42x^3 - 6x^2 - 27x + 9) (-x^2 + 1)^{\frac{1}{3}} + 4^{\frac{1}{3}} (x^6 + 18x^5 - 117x^4 + 36x^3 + 207x^2 - 54x - 27) \right)}{6(x^6 - 54x^5 + 171x^4 - 108x^3 - 81x^2 + 162x - 27)} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3+x)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="fricas")

[Out] $-1/6 \cdot 4^{1/6} \cdot \sqrt{3} \cdot (-1)^{1/3} \cdot \arctan\left(\frac{1/6 \cdot 4^{1/6} \cdot \sqrt{3} \cdot (12 \cdot 4^{2/3} \cdot (-1)^{2/3} \cdot (x^4 - 3x^3 + 3x^2 - 9x) \cdot (-x^2 + 1)^{2/3} + 12 \cdot (-1)^{1/3} \cdot (x^5 - 19x^4 + 42x^3 - 6x^2 - 27x + 9) \cdot (-x^2 + 1)^{1/3} + 4^{1/3} \cdot (x^6 + 18x^5 - 117x^4 + 36x^3 + 207x^2 - 54x - 27))}{x^6 - 54x^5 + 171x^4 - 108x^3 - 81x^2 + 162x - 27}\right) - 1/24 \cdot 4^{2/3} \cdot (-1)^{1/3} \cdot \log\left(\frac{-6 \cdot 4^{2/3} \cdot (-1)^{1/3} \cdot (x^2 - 3x) \cdot (-x^2 + 1)^{2/3} - 4^{1/3} \cdot (-1)^{2/3} \cdot (x^4 - 18x^3 + 24x^2 + 18x - 9) - 6 \cdot (x^3 - 7x^2 + 3x + 3) \cdot (-x^2 + 1)^{1/3}}{x^4 + 6x^2 + 9}\right) + 1/12 \cdot 4^{2/3} \cdot (-1)^{1/3} \cdot \log\left(\frac{-6 \cdot 4^{1/3} \cdot (-1)^{2/3} \cdot (-x^2 + 1)^{1/3} \cdot (x - 1) + 4^{2/3} \cdot (-1)^{1/3} \cdot (x^2 + 3) - 12 \cdot (-x^2 + 1)^{2/3}}{x^2 + 3}\right)$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x+3}{(x^2+3)(-x^2+1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3+x)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="giac")

[Out] integrate((x + 3)/((x^2 + 3)*(-x^2 + 1)^(1/3)), x)

maple [C] time = 7.62, size = 1553, normalized size = 16.35

Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x+3)/(-x^2+1)^(1/3)/(x^2+3),x)

[Out] $-1/2 \cdot \ln\left(\frac{-8 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2)^2 \cdot \text{RootOf}(_Z^3+2)^2 \cdot x^2 - 2 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2)^3 \cdot x^2 - 24 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2)^2 \cdot \text{RootOf}(_Z^3+2)^2 \cdot x + 6 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2)^3 \cdot x + 18 \cdot (-x^2+1)^{2/3} \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2)^2 - 18 \cdot (-x^2+1)^{1/3} \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2) \cdot x - 6 \cdot (-x^2+1)^{1/3} \cdot \text{RootOf}(_Z^3+2)^2 \cdot x + 18 \cdot (-x^2+1)^{1/3} \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2) - 4 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot x^2 + 6 \cdot (-x^2+1)^{1/3} \cdot \text{RootOf}(_Z^3+2)^2 + \text{RootOf}(_Z^3+2) \cdot x^2 - 12 \cdot (-x^2+1)^{2/3} - 12 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) + 3 \cdot \text{RootOf}(_Z^3+2)}{(2 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2)^2 \cdot x - x - 3)}\right) / (2 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2)^2 \cdot x - x + 3) \cdot \text{RootOf}(_Z^3+2) - \ln\left(\frac{-8 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2)^2 \cdot \text{RootOf}(_Z^3+2)^2 \cdot x^2 - 2 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2)^3 \cdot x^2 - 24 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2)^2 \cdot \text{RootOf}(_Z^3+2)^2 \cdot x + 6 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2)^3 \cdot x + 18 \cdot (-x^2+1)^{2/3} \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2) - 18 \cdot (-x^2+1)^{1/3} \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2) \cdot x - 6 \cdot (-x^2+1)^{1/3} \cdot \text{RootOf}(_Z^3+2)^2 \cdot x + 18 \cdot (-x^2+1)^{1/3} \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2) - 4 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot x^2 + 6 \cdot (-x^2+1)^{1/3} \cdot \text{RootOf}(_Z^3+2)^2 + \text{RootOf}(_Z^3+2) \cdot x^2 - 12 \cdot (-x^2+1)^{2/3} - 12 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) + 3 \cdot \text{RootOf}(_Z^3+2)}{(2 \cdot \text{RootOf}(\text{RootOf}(_Z^3+2)^2+2 \cdot _Z \cdot \text{RootOf}(_Z^3+2)+4 \cdot _Z^2) \cdot \text{RootOf}(_Z^3+2)^2 \cdot x - x + 3)}\right)$

$4*_Z^2)*\text{RootOf}(_Z^3+2)^2-18*(-x^2+1)^{(1/3)}*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)*x-6*(-x^2+1)^{(1/3)}*\text{RootOf}(_Z^3+2)^2*x+18*(-x^2+1)^{(1/3)}*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)-4*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*x^2+6*(-x^2+1)^{(1/3)}*\text{RootOf}(_Z^3+2)^2+\text{RootOf}(_Z^3+2)*x^2-12*(-x^2+1)^{(2/3)}-12*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)+3*\text{RootOf}(_Z^3+2))/ (2*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)^2*x-x-3)/ (2*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)^2*x-x+3))*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)+\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\ln(- (8*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)^2*\text{RootOf}(_Z^3+2)^2*x^2+6*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)^3*x^2-24*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)^2*\text{RootOf}(_Z^3+2)^2*x-18*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)^3*x-18*(-x^2+1)^{(2/3)}*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)^2+18*(-x^2+1)^{(1/3)}*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)*x+3*(-x^2+1)^{(1/3)}*\text{RootOf}(_Z^3+2)^2*x-18*(-x^2+1)^{(1/3)}*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)-4*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*x^2-3*(-x^2+1)^{(1/3)}*\text{RootOf}(_Z^3+2)^2-3*\text{RootOf}(_Z^3+2)*x^2+24*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*x+18*\text{RootOf}(_Z^3+2)*x+6*(-x^2+1)^{(2/3)}+12*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)+9*\text{RootOf}(_Z^3+2))/ (2*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)^2*x-x-3)/ (2*\text{RootOf}(\text{RootOf}(_Z^3+2)^2+2*_Z*\text{RootOf}(_Z^3+2)+4*_Z^2)*\text{RootOf}(_Z^3+2)^2*x-x+3))$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x+3}{(x^2+3)(-x^2+1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3+x)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="maxima")

[Out] integrate((x+3)/((x^2+3)*(-x^2+1)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x+3}{(1-x^2)^{1/3}(x^2+3)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x+3)/((1-x^2)^(1/3)*(x^2+3)),x)

[Out] int((x+3)/((1-x^2)^(1/3)*(x^2+3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x+3}{\sqrt[3]{-(x-1)(x+1)}(x^2+3)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3+x)/(-x**2+1)**(1/3)/(x**2+3),x)

[Out] Integral((x+3)/((-x-1)*(x+1)**(1/3)*(x**2+3)), x)

$$3.150 \quad \int \frac{1}{\sqrt[3]{a+bx^2} \left(\frac{9ad}{b} + dx^2 \right)} dx$$

Optimal. Leaf size=151

$$\frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{a} - \sqrt[3]{a+bx^2} \right)^2}{3 \sqrt[6]{a} \sqrt{bx}} \right)}{12a^{5/6}d} - \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{a+bx^2} \right)}{\sqrt{bx}} \right)}{4\sqrt{3} a^{5/6}d} + \frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3\sqrt[6]{a}} \right)}{12a^{5/6}d}$$

[Out] $1/12*\arctan(1/3*(a^{(1/3)}-(b*x^2+a)^{(1/3)})^2/a^{(1/6)}/x/b^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d+1/12*\arctan(1/3*x*b^{(1/2)}/a^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d-1/12*\operatorname{arctanh}(a^{(1/6)}*(a^{(1/3)}-(b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d*3^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 151, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.037$, Rules used = {394}

$$\frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{a} - \sqrt[3]{a+bx^2} \right)^2}{3 \sqrt[6]{a} \sqrt{bx}} \right)}{12a^{5/6}d} - \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{a+bx^2} \right)}{\sqrt{bx}} \right)}{4\sqrt{3} a^{5/6}d} + \frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3\sqrt[6]{a}} \right)}{12a^{5/6}d}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(1/3)*((9*a*d)/b + d*x^2)), x]

[Out] (Sqrt[b]*ArcTan[(Sqrt[b]*x)/(3*Sqrt[a])])/(12*a^(5/6)*d) + (Sqrt[b]*ArcTan[a^(1/3) - (a + b*x^2)^(1/3)]^2/(3*a^(1/6)*Sqrt[b]*x))/(12*a^(5/6)*d) - (Sqrt[b]*ArcTanh[(Sqrt[3]*a^(1/6)*(a^(1/3) - (a + b*x^2)^(1/3)))/(Sqrt[b]*x)])/(4*Sqrt[3]*a^(5/6)*d)

Rule 394

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[(q*ArcTan[(q*x)/3])/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTan[(Rt[a, 3] - (a + b*x^2)^(1/3)]^2/(3*Rt[a, 3]^2*q*x))]/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTanh[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))/(Rt[a, 3]*q*x)])/(4*Sqrt[3]*Rt[a, 3]*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{a+bx^2} \left(\frac{9ad}{b} + dx^2 \right)} dx = \frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3\sqrt[6]{a}} \right)}{12a^{5/6}d} + \frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{a} - \sqrt[3]{a+bx^2} \right)^2}{3 \sqrt[6]{a} \sqrt{bx}} \right)}{12a^{5/6}d} - \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{a+bx^2} \right)}{\sqrt{bx}} \right)}{4\sqrt{3} a^{5/6}d}$$

Mathematica [C] time = 0.17, size = 169, normalized size = 1.12

$$\frac{27abx F_1 \left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{9a} \right)}{d \sqrt[3]{a+bx^2} (9a+bx^2) \left(27a F_1 \left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{9a} \right) - 2bx^2 \left(F_1 \left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{9a} \right) + 3F_1 \left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -\frac{bx^2}{a}, - \right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(1/3)*((9*a*d)/b + d*x^2)),x]

[Out] (27*a*b*x*AppellF1[1/2, 1/3, 1, 3/2, -((b*x^2)/a), -1/9*(b*x^2)/a])/(d*(a + b*x^2)^(1/3)*(9*a + b*x^2)*(27*a*AppellF1[1/2, 1/3, 1, 3/2, -((b*x^2)/a), -1/9*(b*x^2)/a] - 2*b*x^2*(AppellF1[3/2, 1/3, 2, 5/2, -((b*x^2)/a), -1/9*(b*x^2)/a] + 3*AppellF1[3/2, 4/3, 1, 5/2, -((b*x^2)/a), -1/9*(b*x^2)/a]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/3)/(9*a*d/b+d*x^2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{1}{3}} \left(dx^2 + \frac{9ad}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/3)/(9*a*d/b+d*x^2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(1/3)*(d*x^2 + 9*a*d/b)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{1}{3}} \left(dx^2 + \frac{9ad}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/3)/(9*a*d/b+d*x^2),x)

[Out] int(1/(b*x^2+a)^(1/3)/(9*a*d/b+d*x^2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{1}{3}} \left(dx^2 + \frac{9ad}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/3)/(9*a*d/b+d*x^2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(1/3)*(d*x^2 + 9*a*d/b)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + a)^{1/3} \left(dx^2 + \frac{9ad}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(1/3)*(d*x^2 + (9*a*d)/b)),x)

[Out] int(1/((a + b*x^2)^(1/3)*(d*x^2 + (9*a*d)/b)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{b \int \frac{1}{9a \sqrt[3]{a+bx^2} + bx^2 \sqrt[3]{a+bx^2}} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(1/3)/(9*a*d/b+d*x**2), x)

[Out] b*Integral(1/(9*a*(a + b*x**2)**(1/3) + b*x**2*(a + b*x**2)**(1/3)), x)/d

$$3.151 \quad \int \frac{1}{\sqrt[3]{a-bx^2} \left(-\frac{9ad}{b} + dx^2\right)} dx$$

Optimal. Leaf size=153

$$-\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{4\sqrt{3} a^{5/6} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}{3\sqrt[6]{a} \sqrt{bx}}\right)}{12a^{5/6} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3\sqrt[6]{a}}\right)}{12a^{5/6} d}$$

[Out] $1/12*\operatorname{arctanh}(1/3*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})^2/a^{(1/6)}/x/b^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d-1/12*\operatorname{arctanh}(1/3*x*b^{(1/2)}/a^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d-1/12*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}-(-b*x^2+a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d*3^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.036$, Rules used = {395}

$$-\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{4\sqrt{3} a^{5/6} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}{3\sqrt[6]{a} \sqrt{bx}}\right)}{12a^{5/6} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3\sqrt[6]{a}}\right)}{12a^{5/6} d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[1/\left((a - b*x^2)^{(1/3)}*((-9*a*d)/b + d*x^2)\right), x\right]$

[Out] $-(\operatorname{Sqrt}[b]*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)} - (a - b*x^2)^{(1/3)}))/(\operatorname{Sqrt}[b]*x)])/(4*\operatorname{Sqrt}[3]*a^{(5/6)*d} - (\operatorname{Sqrt}[b]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(3*\operatorname{Sqrt}[a]])/(12*a^{(5/6)*d} + (\operatorname{Sqrt}[b]*\operatorname{ArcTanh}[(a^{(1/3)} - (a - b*x^2)^{(1/3)})^2/(3*a^{(1/6)}*\operatorname{Sqrt}[b]*x)])/(12*a^{(5/6)*d})$

Rule 395

$\operatorname{Int}\left[1/\left(\left((a_{-}) + (b_{-})*(x_{-})^2\right)^{(1/3)}*\left((c_{-}) + (d_{-})*(x_{-})^2\right)\right), x_{\text{Symbol}}\right] :> \operatorname{With}\left[\{q = \operatorname{Rt}[-(b/a), 2]\}, -\operatorname{Simp}\left[\left(q*\operatorname{ArcTanh}\left[\frac{q*x}{3}\right]\right)/\left(12*\operatorname{Rt}[a, 3]*d\right), x\right] + \left(\operatorname{Simp}\left[\left(q*\operatorname{ArcTanh}\left[\frac{\operatorname{Rt}[a, 3] - (a + b*x^2)^{(1/3)}^2}{3*\operatorname{Rt}[a, 3]^2*q*x}\right]\right)/\left(12*\operatorname{Rt}[a, 3]*d\right), x\right] - \operatorname{Simp}\left[\left(q*\operatorname{ArcTan}\left[\frac{\operatorname{Sqrt}[3]*\left(\operatorname{Rt}[a, 3] - (a + b*x^2)^{(1/3)}\right)}{\operatorname{Rt}[a, 3]*q*x}\right]\right)/\left(4*\operatorname{Sqrt}[3]*\operatorname{Rt}[a, 3]*d\right), x\right]\right] /; \operatorname{FreeQ}\{a, b, c, d\}, x\} \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{EqQ}[b*c - 9*a*d, 0] \&\& \operatorname{NegQ}[b/a]$

Rubi steps

$$\int \frac{1}{\sqrt[3]{a-bx^2} \left(-\frac{9ad}{b} + dx^2\right)} dx = -\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)}{\sqrt{bx}}\right)}{4\sqrt{3} a^{5/6} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3\sqrt[6]{a}}\right)}{12a^{5/6} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{a} - \sqrt[3]{a-bx^2}\right)^2}{3\sqrt[6]{a} \sqrt{bx}}\right)}{12a^{5/6} d}$$

Mathematica [C] time = 0.18, size = 167, normalized size = 1.09

$$\frac{27abx F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, \frac{bx^2}{9a}\right)}{\sqrt[3]{a-bx^2} (9a-bx^2) \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, \frac{bx^2}{9a}\right) + 3F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, \frac{bx^2}{9a}\right)\right) + 27a F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, \frac{bx^2}{9a}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(1/3)*((-9*a*d)/b + d*x^2)),x]

[Out] (-27*a*b*x*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, (b*x^2)/(9*a)]/(d*(a - b*x^2)^(1/3)*(9*a - b*x^2)*(27*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, (b*x^2)/(9*a)]/(9*a)] + 2*b*x^2*(AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, (b*x^2)/(9*a)] + 3*AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, (b*x^2)/(9*a)]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(-9*a*d/b+d*x^2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{1}{3}} \left(dx^2 - \frac{9ad}{b} \right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(-9*a*d/b+d*x^2),x, algorithm="giac")

[Out] integrate(1/((-b*x^2 + a)^(1/3)*(d*x^2 - 9*a*d/b)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{1}{3}} \left(dx^2 - \frac{9ad}{b} \right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(1/3)/(-9*a/b*d+d*x^2),x)

[Out] int(1/(-b*x^2+a)^(1/3)/(-9*a/b*d+d*x^2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{1}{3}} \left(dx^2 - \frac{9ad}{b} \right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/3)/(-9*a*d/b+d*x^2),x, algorithm="maxima")

[Out] integrate(1/((-b*x^2 + a)^(1/3)*(d*x^2 - 9*a*d/b)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(a - bx^2)^{\frac{1}{3}} \left(dx^2 - \frac{9ad}{b} \right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(1/3)*(d*x^2 - (9*a*d)/b)),x)

[Out] int(1/((a - b*x^2)^(1/3)*(d*x^2 - (9*a*d)/b)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{b \int \frac{1}{-9a \sqrt[3]{a-bx^2} + bx^2 \sqrt[3]{a-bx^2}} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2+a)**(1/3)/(-9*a*d/b+d*x**2), x)

[Out] b*Integral(1/(-9*a*(a - b*x**2)**(1/3) + b*x**2*(a - b*x**2)**(1/3)), x)/d

$$3.152 \quad \int \frac{1}{\sqrt[3]{-a+bx^2} \left(-\frac{9ad}{b} + dx^2\right)} dx$$

Optimal. Leaf size=151

$$\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{bx^2-a} + \sqrt[3]{a}\right)}{\sqrt{bx}}\right)}{4\sqrt{3} a^{5/6} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{bx^2-a} + \sqrt[3]{a}\right)^2}{3\sqrt[6]{a} \sqrt{bx}}\right)}{12a^{5/6} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3\sqrt[3]{a}}\right)}{12a^{5/6} d}$$

[Out] $-1/12*\operatorname{arctanh}(1/3*(a^{(1/3)}+(b*x^2-a)^{(1/3)})^2/a^{(1/6)}/x/b^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d+1/12*\operatorname{arctanh}(1/3*x*b^{(1/2)}/a^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d+1/12*\operatorname{arctan}(a^{(1/6)}*(a^{(1/3)}+(b*x^2-a)^{(1/3)})*3^{(1/2)}/x/b^{(1/2)})*b^{(1/2)}/a^{(5/6)}/d*3^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 151, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 29, $\frac{\text{number of rules}}{\text{integrand size}} = 0.034$, Rules used = {395}

$$\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{bx^2-a} + \sqrt[3]{a}\right)}{\sqrt{bx}}\right)}{4\sqrt{3} a^{5/6} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{bx^2-a} + \sqrt[3]{a}\right)^2}{3\sqrt[6]{a} \sqrt{bx}}\right)}{12a^{5/6} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3\sqrt[3]{a}}\right)}{12a^{5/6} d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/((-a + b*x^2)^{(1/3)}*((-9*a*d)/b + d*x^2)), x]$

[Out] $(\operatorname{Sqrt}[b]*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*a^{(1/6)}*(a^{(1/3)} + (-a + b*x^2)^{(1/3)}))/(\operatorname{Sqrt}[b]*x)])/(4*\operatorname{Sqrt}[3]*a^{(5/6)*d} + (\operatorname{Sqrt}[b]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(3*\operatorname{Sqrt}[a])]))/(12*a^{(5/6)*d} - (\operatorname{Sqrt}[b]*\operatorname{ArcTanh}[(a^{(1/3)} + (-a + b*x^2)^{(1/3)})^2/(3*a^{(1/6)}*\operatorname{Sqrt}[b]*x)))/(12*a^{(5/6)*d})$

Rule 395

$\operatorname{Int}[1/(((a_) + (b_)*(x_)^2)^{(1/3)}*((c_) + (d_)*(x_)^2)), x_Symbol] := \operatorname{With}[{q = \operatorname{Rt}[-(b/a), 2]}, -\operatorname{Simp}[(q*\operatorname{ArcTanh}[(q*x)/3])/(12*\operatorname{Rt}[a, 3]*d), x] + (\operatorname{Simp}[(q*\operatorname{ArcTanh}[(\operatorname{Rt}[a, 3] - (a + b*x^2)^{(1/3)})^2/(3*\operatorname{Rt}[a, 3]^2*q*x)])/(12*\operatorname{Rt}[a, 3]*d), x] - \operatorname{Simp}[(q*\operatorname{ArcTan}[(\operatorname{Sqrt}[3]*(\operatorname{Rt}[a, 3] - (a + b*x^2)^{(1/3)})))/(\operatorname{Rt}[a, 3]*q*x)])/(4*\operatorname{Sqrt}[3]*\operatorname{Rt}[a, 3]*d), x)] /; \operatorname{FreeQ}[{a, b, c, d}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{EqQ}[b*c - 9*a*d, 0] \&\& \operatorname{NegQ}[b/a]$

Rubi steps

$$\int \frac{1}{\sqrt[3]{-a+bx^2} \left(-\frac{9ad}{b} + dx^2\right)} dx = \frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} + \sqrt[3]{-a+bx^2}\right)}{\sqrt{bx}}\right)}{4\sqrt{3} a^{5/6} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3\sqrt[3]{a}}\right)}{12a^{5/6} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{a} + \sqrt[3]{-a+bx^2}\right)^2}{3\sqrt[6]{a} \sqrt{bx}}\right)}{12a^{5/6} d}$$

Mathematica [C] time = 0.15, size = 168, normalized size = 1.11

$$\frac{27abx F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, \frac{bx^2}{9a}\right)}{d(9a - bx^2) \sqrt[3]{bx^2 - a} \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{a}, \frac{bx^2}{9a}\right) + 3F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{a}, \frac{bx^2}{9a}\right)\right) + 27a F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{a}, \frac{bx^2}{9a}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-a + b*x^2)^(1/3)*((-9*a*d)/b + d*x^2)),x]

[Out] (-27*a*b*x*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, (b*x^2)/(9*a)]/(d*(9*a - b*x^2)*(-a + b*x^2)^(1/3)*(27*a*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/a, (b*x^2)/(9*a)] + 2*b*x^2*(AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/a, (b*x^2)/(9*a)] + 3*AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/a, (b*x^2)/(9*a)])))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-a)^(1/3)/(-9*a*d/b+d*x^2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - a)^{\frac{1}{3}} \left(dx^2 - \frac{9ad}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-a)^(1/3)/(-9*a*d/b+d*x^2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 - a)^(1/3)*(d*x^2 - 9*a*d/b)), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - a)^{\frac{1}{3}} \left(dx^2 - \frac{9ad}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2-a)^(1/3)/(d*x^2-9*a/b*d),x)

[Out] int(1/(b*x^2-a)^(1/3)/(d*x^2-9*a/b*d),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - a)^{\frac{1}{3}} \left(dx^2 - \frac{9ad}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-a)^(1/3)/(-9*a*d/b+d*x^2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 - a)^(1/3)*(d*x^2 - 9*a*d/b)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 - a)^{1/3} \left(dx^2 - \frac{9ad}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((b*x^2 - a)^(1/3)*(d*x^2 - (9*a*d)/b)),x)

[Out] int(1/((b*x^2 - a)^(1/3)*(d*x^2 - (9*a*d)/b)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{b \int \frac{1}{-9a \sqrt[3]{-a+bx^2} + bx^2 \sqrt[3]{-a+bx^2}} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2-a)**(1/3)/(-9*a*d/b+d*x**2), x)

[Out] b*Integral(1/(-9*a*(-a + b*x**2)**(1/3) + b*x**2*(-a + b*x**2)**(1/3)), x)/
d

$$3.153 \quad \int \frac{1}{\sqrt[3]{-a-bx^2} \left(\frac{9ad}{b} + dx^2 \right)} dx$$

Optimal. Leaf size=153

$$-\frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{-a-bx^2} + \sqrt[3]{a} \right)^2}{3\sqrt[6]{a} \sqrt{bx}} \right)}{12a^{5/6}d} + \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{-a-bx^2} + \sqrt[3]{a} \right)}{\sqrt{bx}} \right)}{4\sqrt{3} a^{5/6}d} - \frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3\sqrt[6]{a}} \right)}{12a^{5/6}d}$$

[Out] $-1/12 \arctan(1/3 * (a^{1/3} + (-b*x^2 - a)^{1/3})^2 / a^{1/6} / x / b^{1/2}) * b^{1/2} / a^{5/6} / d - 1/12 \arctan(1/3 * x * b^{1/2} / a^{1/2}) * b^{1/2} / a^{5/6} / d + 1/12 \operatorname{arctanh}(a^{1/6} * (a^{1/3} + (-b*x^2 - a)^{1/3}) * 3^{1/2} / x / b^{1/2}) * b^{1/2} / a^{5/6} / d * 3^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 0.033$, Rules used = {394}

$$-\frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{-a-bx^2} + \sqrt[3]{a} \right)^2}{3\sqrt[6]{a} \sqrt{bx}} \right)}{12a^{5/6}d} + \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{-a-bx^2} + \sqrt[3]{a} \right)}{\sqrt{bx}} \right)}{4\sqrt{3} a^{5/6}d} - \frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3\sqrt[6]{a}} \right)}{12a^{5/6}d}$$

Antiderivative was successfully verified.

[In] Int[1/((-a - b*x^2)^(1/3)*((9*a*d)/b + d*x^2)),x]

[Out] $-(\text{Sqrt}[b] * \text{ArcTan}[(\text{Sqrt}[b] * x) / (3 * \text{Sqrt}[a])]) / (12 * a^{5/6} * d) - (\text{Sqrt}[b] * \text{ArcTan}[(a^{1/3} + (-a - b * x^2)^{1/3})^2 / (3 * a^{1/6} * \text{Sqrt}[b] * x)]) / (12 * a^{5/6} * d) + (\text{Sqrt}[b] * \text{ArcTanh}[(\text{Sqrt}[3] * a^{1/6} * (a^{1/3} + (-a - b * x^2)^{1/3})) / (\text{Sqrt}[b] * x)]) / (4 * \text{Sqrt}[3] * a^{5/6} * d)$

Rule 394

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[b/a, 2]}, Simp[(q * ArcTan[(q * x) / 3]) / (12 * Rt[a, 3] * d), x] + (Simp[(q * ArcTan[(Rt[a, 3] - (a + b * x^2)^(1/3))^2 / (3 * Rt[a, 3]^2 * q * x)]) / (12 * Rt[a, 3] * d), x] - Simp[(q * ArcTanh[(Sqrt[3] * (Rt[a, 3] - (a + b * x^2)^(1/3)))] / (Rt[a, 3] * q * x)]) / (4 * Sqrt[3] * Rt[a, 3] * d), x)] /; FreeQ[{a, b, c, d}, x] && NeQ[b * c - a * d, 0] && EqQ[b * c - 9 * a * d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{-a-bx^2} \left(\frac{9ad}{b} + dx^2 \right)} dx = -\frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3\sqrt[6]{a}} \right)}{12a^{5/6}d} - \frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{a} + \sqrt[3]{-a-bx^2} \right)^2}{3\sqrt[6]{a} \sqrt{bx}} \right)}{12a^{5/6}d} + \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt{3} \sqrt[6]{a} \left(\sqrt[3]{a} + \sqrt[3]{-a-bx^2} \right)}{\sqrt{bx}} \right)}{4\sqrt{3} a^{5/6}d}$$

Mathematica [C] time = 0.16, size = 172, normalized size = 1.12

$$\frac{27abx F_1 \left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{9a} \right)}{d \sqrt[3]{-a-bx^2} (9a + bx^2) \left(27a F_1 \left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{9a} \right) - 2bx^2 \left(F_1 \left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{9a} \right) + 3F_1 \left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{9a} \right) \right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-a - b*x^2)^(1/3)*((9*a*d)/b + d*x^2)),x]

[Out] (27*a*b*x*AppellF1[1/2, 1/3, 1, 3/2, -((b*x^2)/a), -1/9*(b*x^2)/a])/(d*(-a - b*x^2)^(1/3)*(9*a + b*x^2)*(27*a*AppellF1[1/2, 1/3, 1, 3/2, -((b*x^2)/a), -1/9*(b*x^2)/a] - 2*b*x^2*(AppellF1[3/2, 1/3, 2, 5/2, -((b*x^2)/a), -1/9*(b*x^2)/a] + 3*AppellF1[3/2, 4/3, 1, 5/2, -((b*x^2)/a), -1/9*(b*x^2)/a]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-a)^(1/3)/(9*a*d/b+d*x^2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 - a)^{\frac{1}{3}} \left(dx^2 + \frac{9ad}{b} \right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-a)^(1/3)/(9*a*d/b+d*x^2),x, algorithm="giac")

[Out] integrate(1/((-b*x^2 - a)^(1/3)*(d*x^2 + 9*a*d/b)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 - a)^{\frac{1}{3}} \left(dx^2 + \frac{9ad}{b} \right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2-a)^(1/3)/(d*x^2+9*a/b*d),x)

[Out] int(1/(-b*x^2-a)^(1/3)/(d*x^2+9*a/b*d),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 - a)^{\frac{1}{3}} \left(dx^2 + \frac{9ad}{b} \right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-a)^(1/3)/(9*a*d/b+d*x^2),x, algorithm="maxima")

[Out] integrate(1/((-b*x^2 - a)^(1/3)*(d*x^2 + 9*a*d/b)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(-bx^2 - a)^{\frac{1}{3}} \left(dx^2 + \frac{9ad}{b} \right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((- a - b*x^2)^(1/3)*(d*x^2 + (9*a*d)/b)),x)

[Out] int(1/((- a - b*x^2)^(1/3)*(d*x^2 + (9*a*d)/b)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{b \int \frac{1}{9a \sqrt[3]{-a-bx^2} + bx^2 \sqrt[3]{-a-bx^2}} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2-a)**(1/3)/(9*a*d/b+d*x**2), x)

[Out] b*Integral(1/(9*a*(-a - b*x**2)**(1/3) + b*x**2*(-a - b*x**2)**(1/3)), x)/d

$$3.154 \quad \int \frac{1}{\sqrt[3]{2+bx^2} \left(\frac{18d}{b} + dx^2 \right)} dx$$

Optimal. Leaf size=151

$$\frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{2} - \sqrt[3]{bx^2+2} \right)^2}{3 \sqrt[6]{2} \sqrt{bx}} \right)}{12 \cdot 2^{5/6} d} - \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt[6]{2} \sqrt{3} \left(\sqrt[3]{2} - \sqrt[3]{bx^2+2} \right)}{\sqrt{bx}} \right)}{4 \cdot 2^{5/6} \sqrt{3} d} + \frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3 \sqrt[6]{2}} \right)}{12 \cdot 2^{5/6} d}$$

[Out] $1/24 \cdot \arctan(1/6 \cdot (2^{1/3} - (b \cdot x^2 + 2)^{1/3})^2 \cdot 2^{5/6} / x / b^{1/2}) \cdot b^{1/2} \cdot 2^{1/6} / d + 1/24 \cdot \arctan(1/6 \cdot x \cdot b^{1/2} \cdot 2^{1/2}) \cdot b^{1/2} \cdot 2^{1/6} / d - 1/24 \cdot \operatorname{arctanh}(2^{1/6} \cdot (2^{1/3} - (b \cdot x^2 + 2)^{1/3}) \cdot 3^{1/2} / x / b^{1/2}) \cdot b^{1/2} \cdot 2^{1/6} / d \cdot 3^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 151, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {394}

$$\frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{2} - \sqrt[3]{bx^2+2} \right)^2}{3 \sqrt[6]{2} \sqrt{bx}} \right)}{12 \cdot 2^{5/6} d} - \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt[6]{2} \sqrt{3} \left(\sqrt[3]{2} - \sqrt[3]{bx^2+2} \right)}{\sqrt{bx}} \right)}{4 \cdot 2^{5/6} \sqrt{3} d} + \frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3 \sqrt[6]{2}} \right)}{12 \cdot 2^{5/6} d}$$

Antiderivative was successfully verified.

[In] Int[1/((2 + b*x^2)^(1/3)*((18*d)/b + d*x^2)),x]

[Out] (Sqrt[b]*ArcTan[(Sqrt[b]*x)/(3*Sqrt[2])])/(12*2^(5/6)*d) + (Sqrt[b]*ArcTan[(2^(1/3) - (2 + b*x^2)^(1/3))^2/(3*2^(1/6)*Sqrt[b]*x)]/(12*2^(5/6)*d) - (Sqrt[b]*ArcTanh[(2^(1/6)*Sqrt[3]*(2^(1/3) - (2 + b*x^2)^(1/3)))/(Sqrt[b]*x)]/(4*2^(5/6)*Sqrt[3]*d)

Rule 394

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[(q*ArcTan[(q*x)/3])/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTan[(Rt[a, 3] - (a + b*x^2)^(1/3))^2/(3*Rt[a, 3]^2*q*x)]/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTanh[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))/(Rt[a, 3]*q*x)]/(4*Sqrt[3]*Rt[a, 3]*d), x)]) /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{2+bx^2} \left(\frac{18d}{b} + dx^2 \right)} dx = \frac{\sqrt{b} \tan^{-1} \left(\frac{\sqrt{bx}}{3 \sqrt[6]{2}} \right)}{12 \cdot 2^{5/6} d} + \frac{\sqrt{b} \tan^{-1} \left(\frac{\left(\sqrt[3]{2} - \sqrt[3]{2+bx^2} \right)^2}{3 \sqrt[6]{2} \sqrt{bx}} \right)}{12 \cdot 2^{5/6} d} - \frac{\sqrt{b} \tanh^{-1} \left(\frac{\sqrt[6]{2} \sqrt{3} \left(\sqrt[3]{2} - \sqrt[3]{2+bx^2} \right)}{\sqrt{bx}} \right)}{4 \cdot 2^{5/6} \sqrt{3} d}$$

Mathematica [C] time = 0.15, size = 148, normalized size = 0.98

$$\frac{27bx F_1 \left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{bx^2}{2}, -\frac{bx^2}{18} \right)}{d \sqrt[3]{bx^2+2} (bx^2+18) \left(bx^2 \left(F_1 \left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -\frac{bx^2}{2}, -\frac{bx^2}{18} \right) + 3 F_1 \left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -\frac{bx^2}{2}, -\frac{bx^2}{18} \right) \right) - 27 F_1 \left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{bx^2}{2}, -\frac{bx^2}{18} \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((2 + b*x^2)^(1/3)*((18*d)/b + d*x^2)),x]

[Out] (-27*b*x*AppellF1[1/2, 1/3, 1, 3/2, -1/2*(b*x^2), -1/18*(b*x^2)]/(d*(2 + b*x^2)^(1/3)*(18 + b*x^2)*(-27*AppellF1[1/2, 1/3, 1, 3/2, -1/2*(b*x^2), -1/18*(b*x^2)] + b*x^2*(AppellF1[3/2, 1/3, 2, 5/2, -1/2*(b*x^2), -1/18*(b*x^2)] + 3*AppellF1[3/2, 4/3, 1, 5/2, -1/2*(b*x^2), -1/18*(b*x^2)])))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/3)/(18*d/b+d*x^2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 2)^{\frac{1}{3}} \left(dx^2 + \frac{18d}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/3)/(18*d/b+d*x^2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 2)^(1/3)*(d*x^2 + 18*d/b)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 2)^{\frac{1}{3}} \left(dx^2 + \frac{18d}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+2)^(1/3)/(18/b*d+d*x^2),x)

[Out] int(1/(b*x^2+2)^(1/3)/(18/b*d+d*x^2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 2)^{\frac{1}{3}} \left(dx^2 + \frac{18d}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/3)/(18*d/b+d*x^2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 2)^(1/3)*(d*x^2 + 18*d/b)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{18d}{b} + dx^2\right) (bx^2 + 2)^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(((18*d)/b + d*x^2)*(b*x^2 + 2)^(1/3)),x)

[Out] int(1/(((18*d)/b + d*x^2)*(b*x^2 + 2)^(1/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$b \int \frac{1}{bx^2 \sqrt[3]{bx^2+2} + 18 \sqrt[3]{bx^2+2}} dx$$

$$\frac{1}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+2)**(1/3)/(18*d/b+d*x**2), x)

[Out] b*Integral(1/(b*x**2*(b*x**2 + 2)**(1/3) + 18*(b*x**2 + 2)**(1/3)), x)/d

$$3.155 \quad \int \frac{1}{\sqrt[3]{-2+bx^2} \left(-\frac{18d}{b} + dx^2\right)} dx$$

Optimal. Leaf size=147

$$\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt[6]{2} \sqrt{3} \left(\sqrt[3]{bx^2-2} + \sqrt[3]{2}\right)}{\sqrt{bx}}\right)}{4 \cdot 2^{5/6} \sqrt{3} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{bx^2-2} + \sqrt[3]{2}\right)^2}{3 \sqrt[6]{2} \sqrt{bx}}\right)}{12 \cdot 2^{5/6} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3 \sqrt[6]{2}}\right)}{12 \cdot 2^{5/6} d}$$

[Out] $-1/24 \cdot \operatorname{arctanh}\left(\frac{1}{6} \cdot \left(2^{1/3} + (bx^2-2)^{1/3}\right)^2 \cdot 2^{5/6} / x / b^{1/2}\right) \cdot b^{1/2} \cdot 2^{1/6} / d + 1/24 \cdot \operatorname{arctanh}\left(\frac{1}{6} \cdot x \cdot b^{1/2} \cdot 2^{1/2}\right) \cdot b^{1/2} \cdot 2^{1/6} / d + 1/24 \cdot \operatorname{arctan}\left(\frac{1}{6} \cdot \left(2^{1/3} + (bx^2-2)^{1/3}\right) \cdot 3^{1/2} / x / b^{1/2}\right) \cdot b^{1/2} \cdot 2^{1/6} / d \cdot 3^{1/2}$

Rubi [A] time = 0.02, antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {395}

$$\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt[6]{2} \sqrt{3} \left(\sqrt[3]{bx^2-2} + \sqrt[3]{2}\right)}{\sqrt{bx}}\right)}{4 \cdot 2^{5/6} \sqrt{3} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{bx^2-2} + \sqrt[3]{2}\right)^2}{3 \sqrt[6]{2} \sqrt{bx}}\right)}{12 \cdot 2^{5/6} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3 \sqrt[6]{2}}\right)}{12 \cdot 2^{5/6} d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[1/\left((-2 + bx^2)^{1/3} \cdot \left(-18d/b + dx^2\right)\right), x\right]$

[Out] $\left(\operatorname{Sqrt}[b] \cdot \operatorname{ArcTan}\left[\frac{2^{1/6} \operatorname{Sqrt}[3] \cdot \left(2^{1/3} + (-2 + bx^2)^{1/3}\right)}{\operatorname{Sqrt}[b] \cdot x}\right]\right) / \left(4 \cdot 2^{5/6} \cdot \operatorname{Sqrt}[3] \cdot d\right) + \left(\operatorname{Sqrt}[b] \cdot \operatorname{ArcTanh}\left[\frac{\operatorname{Sqrt}[b] \cdot x}{3 \cdot \operatorname{Sqrt}[2]}\right]\right) / \left(12 \cdot 2^{5/6} \cdot d\right) - \left(\operatorname{Sqrt}[b] \cdot \operatorname{ArcTanh}\left[\frac{2^{1/3} + (-2 + bx^2)^{1/3}}{3 \cdot 2^{1/6} \cdot \operatorname{Sqrt}[b] \cdot x}\right]\right) / \left(12 \cdot 2^{5/6} \cdot d\right)$

Rule 395

$\operatorname{Int}\left[1/\left(\left((a_) + (b_) \cdot (x_)^2\right)^{1/3} \cdot \left((c_) + (d_) \cdot (x_)^2\right)\right), x_Symbol\right] :> \operatorname{With}\left[\{q = \operatorname{Rt}\left[-(b/a), 2\right]\}, -\operatorname{Simp}\left[\frac{q \cdot \operatorname{ArcTanh}\left[(q \cdot x)/3\right]}{12 \cdot \operatorname{Rt}[a, 3] \cdot d}, x\right] + \left(\operatorname{Simp}\left[\frac{q \cdot \operatorname{ArcTanh}\left[\operatorname{Rt}[a, 3] - (a + b \cdot x^2)^{1/3}\right]}{3 \cdot \operatorname{Rt}[a, 3]^2 \cdot q \cdot x}\right] / \left(12 \cdot \operatorname{Rt}[a, 3] \cdot d\right), x\right) - \operatorname{Simp}\left[\frac{q \cdot \operatorname{ArcTan}\left[\operatorname{Sqrt}[3] \cdot \left(\operatorname{Rt}[a, 3] - (a + b \cdot x^2)^{1/3}\right)\right]}{\operatorname{Rt}[a, 3] \cdot q \cdot x}\right] / \left(4 \cdot \operatorname{Sqrt}[3] \cdot \operatorname{Rt}[a, 3] \cdot d\right), x\right]\right] /; \operatorname{FreeQ}\{a, b, c, d\}, x \&\& \operatorname{NeQ}[b \cdot c - a \cdot d, 0] \&\& \operatorname{EqQ}[b \cdot c - 9 \cdot a \cdot d, 0] \&\& \operatorname{NegQ}[b/a]$

Rubi steps

$$\int \frac{1}{\sqrt[3]{-2+bx^2} \left(-\frac{18d}{b} + dx^2\right)} dx = \frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt[6]{2} \sqrt{3} \left(\sqrt[3]{2} + \sqrt[3]{-2+bx^2}\right)}{\sqrt{bx}}\right)}{4 \cdot 2^{5/6} \sqrt{3} d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{bx}}{3 \sqrt[6]{2}}\right)}{12 \cdot 2^{5/6} d} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\left(\sqrt[3]{2} + \sqrt[3]{-2+bx^2}\right)^2}{3 \sqrt[6]{2} \sqrt{bx}}\right)}{12 \cdot 2^{5/6} d}$$

Mathematica [C] time = 0.17, size = 148, normalized size = 1.01

$$\frac{27bx F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{2}, \frac{bx^2}{18}\right)}{d \left(bx^2 - 18\right) \sqrt[3]{bx^2 - 2} \left(bx^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{bx^2}{2}, \frac{bx^2}{18}\right) + 3F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{bx^2}{2}, \frac{bx^2}{18}\right)\right) + 27F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{bx^2}{2}, \frac{bx^2}{18}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2 + b*x^2)^(1/3)*((-18*d)/b + d*x^2)),x]

[Out] (27*b*x*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/2, (b*x^2)/18])/(d*(-18 + b*x^2) * (-2 + b*x^2)^(1/3)*(27*AppellF1[1/2, 1/3, 1, 3/2, (b*x^2)/2, (b*x^2)/18] + b*x^2*(AppellF1[3/2, 1/3, 2, 5/2, (b*x^2)/2, (b*x^2)/18] + 3*AppellF1[3/2, 4/3, 1, 5/2, (b*x^2)/2, (b*x^2)/18]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2)^(1/3)/(-18*d/b+d*x^2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 2)^{\frac{1}{3}} \left(dx^2 - \frac{18d}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2)^(1/3)/(-18*d/b+d*x^2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 - 2)^(1/3)*(d*x^2 - 18*d/b)), x)

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 2)^{\frac{1}{3}} \left(dx^2 - \frac{18d}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2-2)^(1/3)/(-18/b*d+d*x^2),x)

[Out] int(1/(b*x^2-2)^(1/3)/(-18/b*d+d*x^2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 2)^{\frac{1}{3}} \left(dx^2 - \frac{18d}{b}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2)^(1/3)/(-18*d/b+d*x^2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 - 2)^(1/3)*(d*x^2 - 18*d/b)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int -\frac{1}{\left(\frac{18d}{b} - dx^2\right) (bx^2 - 2)^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/(((18*d)/b - d*x^2)*(b*x^2 - 2)^(1/3)),x)

[Out] int(-1/(((18*d)/b - d*x^2)*(b*x^2 - 2)^(1/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{b \int \frac{1}{bx^2 \sqrt[3]{bx^2-2} - 18 \sqrt[3]{bx^2-2}} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2-2)**(1/3)/(-18*d/b+d*x**2), x)

[Out] b*Integral(1/(b*x**2*(b*x**2 - 2)**(1/3) - 18*(b*x**2 - 2)**(1/3)), x)/d

$$3.156 \quad \int \frac{1}{\sqrt[3]{2+3x^2} (6d+dx^2)} dx$$

Optimal. Leaf size=123

$$\frac{\tan^{-1}\left(\frac{(\sqrt[3]{2}-\sqrt[3]{3x^2+2})^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} - \frac{\tanh^{-1}\left(\frac{\sqrt[6]{2}(\sqrt[3]{2}-\sqrt[3]{3x^2+2})}{x}\right)}{4 \cdot 2^{5/6}d} + \frac{\tan^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d}$$

[Out] $-1/8*\operatorname{arctanh}(2^{(1/6)}*(2^{(1/3)}-(3*x^2+2)^{(1/3)})/x)*2^{(1/6)}/d+1/24*\operatorname{arctan}(1/18*(2^{(1/3)}-(3*x^2+2)^{(1/3)})^2*2^{(5/6)}/x*3^{(1/2)})*2^{(1/6)}/d*3^{(1/2)}+1/24*\operatorname{arctan}(1/6*x*6^{(1/2)})*2^{(1/6)}/d*3^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {394}

$$\frac{\tan^{-1}\left(\frac{(\sqrt[3]{2}-\sqrt[3]{3x^2+2})^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} - \frac{\tanh^{-1}\left(\frac{\sqrt[6]{2}(\sqrt[3]{2}-\sqrt[3]{3x^2+2})}{x}\right)}{4 \cdot 2^{5/6}d} + \frac{\tan^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d}$$

Antiderivative was successfully verified.

[In] Int[1/((2 + 3*x^2)^(1/3)*(6*d + d*x^2)),x]

[Out] ArcTan[x/Sqrt[6]]/(4*2^(5/6)*Sqrt[3]*d) + ArcTan[(2^(1/3) - (2 + 3*x^2)^(1/3))^2/(3*2^(1/6)*Sqrt[3]*x)]/(4*2^(5/6)*Sqrt[3]*d) - ArcTanh[(2^(1/6)*(2^(1/3) - (2 + 3*x^2)^(1/3)))/x]/(4*2^(5/6)*d)

Rule 394

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[(q*ArcTan[(q*x)/3])/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTan[(Rt[a, 3] - (a + b*x^2)^(1/3))^2/(3*Rt[a, 3]^2*q*x)])/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTanh[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))/(Rt[a, 3]*q*x)])/(4*Sqrt[3]*Rt[a, 3]*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{2+3x^2} (6d+dx^2)} dx = \frac{\tan^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} + \frac{\tan^{-1}\left(\frac{(\sqrt[3]{2}-\sqrt[3]{2+3x^2})^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} - \frac{\tanh^{-1}\left(\frac{\sqrt[6]{2}(\sqrt[3]{2}-\sqrt[3]{2+3x^2})}{x}\right)}{4 \cdot 2^{5/6}d}$$

Mathematica [C] time = 0.12, size = 136, normalized size = 1.11

$$\frac{9x F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{3x^2}{2}, -\frac{x^2}{6}\right)}{d(x^2+6)\sqrt[3]{3x^2+2} \left(x^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -\frac{3x^2}{2}, -\frac{x^2}{6}\right) + 3F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -\frac{3x^2}{2}, -\frac{x^2}{6}\right)\right) - 9F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{3x^2}{2}, -\frac{x^2}{6}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((2 + 3*x^2)^(1/3)*(6*d + d*x^2)),x]

[Out] $(-9*x*AppellF1[1/2, 1/3, 1, 3/2, (-3*x^2)/2, -1/6*x^2])/(d*(6 + x^2)*(2 + 3*x^2)^{(1/3)}*(-9*AppellF1[1/2, 1/3, 1, 3/2, (-3*x^2)/2, -1/6*x^2] + x^2*(AppellF1[3/2, 1/3, 2, 5/2, (-3*x^2)/2, -1/6*x^2] + 3*AppellF1[3/2, 4/3, 1, 5/2, (-3*x^2)/2, -1/6*x^2]))$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(3*x^2+2)^(1/3)/(d*x^2+6*d),x, algorithm="fricas")`

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 + 6d)(3x^2 + 2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(3*x^2+2)^(1/3)/(d*x^2+6*d),x, algorithm="giac")`

[Out] `integrate(1/((d*x^2 + 6*d)*(3*x^2 + 2)^(1/3)), x)`

maple [C] time = 58.10, size = 549, normalized size = 4.46

$$24 \operatorname{RootOf}\left(576_Z^2 - 24_Z \operatorname{RootOf}\left(_Z^6 + 54\right) + \operatorname{RootOf}\left(_Z^6 + 54\right)^2\right) \ln\left(-\frac{192x \operatorname{RootOf}\left(576_Z^2 - 24_Z \operatorname{RootOf}\left(_Z^6 + 54\right) + \operatorname{RootOf}\left(_Z^6 + 54\right)^2\right)}{\dots}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(3*x^2+2)^(1/3)/(d*x^2+6*d),x)`

[Out] $-1/24*(24*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)*\ln(-192*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)*\operatorname{RootOf}(_Z^6+54)^6*x-4*\operatorname{RootOf}(_Z^6+54)^7*x-288*(3*x^2+2)^{(1/3)}*\operatorname{RootOf}(_Z^6+54)^4*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)*x+6*(3*x^2+2)^{(1/3)}*\operatorname{RootOf}(_Z^6+54)^5*x+9*x^2*\operatorname{RootOf}(_Z^6+54)^4-18*\operatorname{RootOf}(_Z^6+54)^4-108*\operatorname{RootOf}(_Z^6+54)^2*(3*x^2+2)^{(1/3)}+324*(3*x^2+2)^{(2/3)})/(x^2+6))- \operatorname{RootOf}(_Z^6+54)*\ln(-(768*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)^2*\operatorname{RootOf}(_Z^6+54)^5*x-16*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)*\operatorname{RootOf}(_Z^6+54)^6*x+1152*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)^2*\operatorname{RootOf}(_Z^6+54)^3*(3*x^2+2)^{(1/3)}*x-72*(3*x^2+2)^{(1/3)}*\operatorname{RootOf}(_Z^6+54)^4*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)*x+(3*x^2+2)^{(1/3)}*\operatorname{RootOf}(_Z^6+54)^5*x-36*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)*\operatorname{RootOf}(_Z^6+54)^3*x^2+72*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)*\operatorname{RootOf}(_Z^6+54)^3-432*\operatorname{RootOf}(\operatorname{RootOf}(_Z^6+54)^2-24*_Z*\operatorname{RootOf}(_Z^6+54)+576*_Z^2)*\operatorname{RootOf}(_Z^6+54)*(3*x^2+2)^{(1/3)}+18*\operatorname{RootOf}(_Z^6+54)^2*(3*x^2+2)^{(1/3)}+54*(3*x^2+2)^{(2/3)})/(x^2+6)))/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 + 6d)(3x^2 + 2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2+2)^(1/3)/(d*x^2+6*d),x, algorithm="maxima")

[Out] integrate(1/((d*x^2 + 6*d)*(3*x^2 + 2)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(3x^2 + 2)^{1/3} (dx^2 + 6d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((3*x^2 + 2)^(1/3)*(6*d + d*x^2)),x)

[Out] int(1/((3*x^2 + 2)^(1/3)*(6*d + d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^2 \sqrt[3]{3x^2+2} + 6 \sqrt[3]{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x**2+2)**(1/3)/(d*x**2+6*d),x)

[Out] Integral(1/(x**2*(3*x**2 + 2)**(1/3) + 6*(3*x**2 + 2)**(1/3)), x)/d

$$3.157 \quad \int \frac{1}{\sqrt[3]{2-3x^2}(-6d+dx^2)} dx$$

Optimal. Leaf size=123

$$-\frac{\tan^{-1}\left(\frac{\sqrt[6]{2}\left(\sqrt[3]{2}-\sqrt[3]{2-3x^2}\right)}{x}\right)}{4 \cdot 2^{5/6}d} + \frac{\tanh^{-1}\left(\frac{\left(\sqrt[3]{2}-\sqrt[3]{2-3x^2}\right)^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} - \frac{\tanh^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d}$$

[Out] $-1/8*\arctan(2^{(1/6)}*(2^{(1/3)}-(-3*x^2+2)^{(1/3)})/x)*2^{(1/6)}/d+1/24*\operatorname{arctanh}(1/18*(2^{(1/3)}-(-3*x^2+2)^{(1/3)})^2*2^{(5/6)}/x*3^{(1/2)})*2^{(1/6)}/d*3^{(1/2)}-1/24*\operatorname{arctanh}(1/6*x*6^{(1/2)})*2^{(1/6)}/d*3^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {395}

$$-\frac{\tan^{-1}\left(\frac{\sqrt[6]{2}\left(\sqrt[3]{2}-\sqrt[3]{2-3x^2}\right)}{x}\right)}{4 \cdot 2^{5/6}d} + \frac{\tanh^{-1}\left(\frac{\left(\sqrt[3]{2}-\sqrt[3]{2-3x^2}\right)^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} - \frac{\tanh^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d}$$

Antiderivative was successfully verified.

[In] Int[1/((2 - 3*x^2)^(1/3)*(-6*d + d*x^2)),x]

[Out] $-\operatorname{ArcTan}\left[\frac{2^{(1/6)}*(2^{(1/3)} - (2 - 3*x^2)^{(1/3)})}{x}\right]/(4*2^{(5/6)*d}) - \operatorname{ArcTanh}\left[\frac{x/\operatorname{Sqrt}[6]}{(4*2^{(5/6)*\operatorname{Sqrt}[3]*d})} + \operatorname{ArcTanh}\left[\frac{(2^{(1/3)} - (2 - 3*x^2)^{(1/3)})^2}{(3*2^{(1/6)*\operatorname{Sqrt}[3]*x})}\right]/(4*2^{(5/6)*\operatorname{Sqrt}[3]*d})\right]$

Rule 395

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b/a), 2]}, -Simp[(q*ArcTanh[(q*x)/3])/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTanh[(Rt[a, 3] - (a + b*x^2)^(1/3))]^2/(3*Rt[a, 3]^2*q*x)]/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTan[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))]/(Rt[a, 3]*q*x)]/(4*Sqrt[3]*Rt[a, 3]*d), x)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && NegQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{2-3x^2}(-6d+dx^2)} dx = -\frac{\tan^{-1}\left(\frac{\sqrt[6]{2}\left(\sqrt[3]{2}-\sqrt[3]{2-3x^2}\right)}{x}\right)}{4 \cdot 2^{5/6}d} - \frac{\tanh^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} + \frac{\tanh^{-1}\left(\frac{\left(\sqrt[3]{2}-\sqrt[3]{2-3x^2}\right)^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d}$$

Mathematica [C] time = 0.13, size = 136, normalized size = 1.11

$$\frac{9x F_1\left(\frac{1}{2}, \frac{1}{3}, 1; \frac{3}{2}, \frac{3x^2}{2}, \frac{x^2}{6}\right)}{d\sqrt[3]{2-3x^2}(x^2-6)\left(x^2\left(F_1\left(\frac{3}{2}, \frac{1}{3}, 2; \frac{5}{2}, \frac{3x^2}{2}, \frac{x^2}{6}\right) + 3F_1\left(\frac{3}{2}, \frac{4}{3}, 1; \frac{5}{2}, \frac{3x^2}{2}, \frac{x^2}{6}\right)\right) + 9F_1\left(\frac{1}{2}, \frac{1}{3}, 1; \frac{3}{2}, \frac{3x^2}{2}, \frac{x^2}{6}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((2 - 3*x^2)^(1/3)*(-6*d + d*x^2)),x]


```
[Out] (9*x*AppellF1[1/2, 1/3, 1, 3/2, (3*x^2)/2, x^2/6])/(d*(2 - 3*x^2)^(1/3)*(-6
+ x^2)*(9*AppellF1[1/2, 1/3, 1, 3/2, (3*x^2)/2, x^2/6] + x^2*(AppellF1[3/2
, 1/3, 2, 5/2, (3*x^2)/2, x^2/6] + 3*AppellF1[3/2, 4/3, 1, 5/2, (3*x^2)/2,
x^2/6])))
```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-3*x^2+2)^(1/3)/(d*x^2-6*d),x, algorithm="fricas")
```

```
[Out] Timed out
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 - 6d)(-3x^2 + 2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-3*x^2+2)^(1/3)/(d*x^2-6*d),x, algorithm="giac")
```

```
[Out] integrate(1/((d*x^2 - 6*d)*(-3*x^2 + 2)^(1/3)), x)
```

maple [C] time = 59.07, size = 1061, normalized size = 8.63

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(-3*x^2+2)^(1/3)/(d*x^2-6*d),x)
```

```
[Out] 1/24*(24*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*ln((-4608
*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*RootOf(_Z^6-54)
^5*x+288*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z
^6-54)^6*x-4*RootOf(_Z^6-54)^7*x+6912*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-
54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*(-3*x^2+2)^(1/3)*x-144*RootOf(_Z^6-
54)^4*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*(-3*x^2+2)^(
1/3)*x-216*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)
+576*_Z^2)*x^2+9*x^2*RootOf(_Z^6-54)^4-432*RootOf(RootOf(_Z^6-54)^2-24*_Z*R
ootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-54)^3+18*RootOf(_Z^6-54)^4-2592*RootOf
(_Z^6-54)*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*(-3*x^2+
2)^(1/3)+324*(-3*x^2+2)^(2/3))/(x^2-6))-24*ln((-768*RootOf(RootOf(_Z^6-54)^
2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*RootOf(_Z^6-54)^5*x+16*RootOf(RootOf(_Z
^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-54)^6*x+1152*RootOf(_Z
^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*(-3*x^2
+2)^(1/3)*x-72*RootOf(_Z^6-54)^4*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6
-54)+576*_Z^2)*(-3*x^2+2)^(1/3)*x+(-3*x^2+2)^(1/3)*RootOf(_Z^6-54)^5*x+36*R
ootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*x
^2+72*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-
54)^3+432*RootOf(_Z^6-54)*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+57
6*_Z^2)*(-3*x^2+2)^(1/3)-18*(-3*x^2+2)^(1/3)*RootOf(_Z^6-54)^2+54*(-3*x^2+2
)^(2/3))/(x^2-6))*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)+
ln((-768*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*RootOf(
_Z^6-54)^5*x+16*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*Ro
otOf(_Z^6-54)^6*x+1152*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*Ro
otOf(_Z^6-54)+576*_Z^2)^2*(-3*x^2+2)^(1/3)*x-72*RootOf(_Z^6-54)^4*RootOf(Ro
otOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*(-3*x^2+2)^(1/3)*x+(-3*x^2+2
)^(1/3)*RootOf(_Z^6-54)^5*x+36*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-2
```

$4*_Z*\text{RootOf}(_Z^6-54)+576*_Z^2)*x^2+72*\text{RootOf}(\text{RootOf}(_Z^6-54)^2-24*_Z*\text{RootOf}(_Z^6-54)+576*_Z^2)*\text{RootOf}(_Z^6-54)^3+432*\text{RootOf}(_Z^6-54)*\text{RootOf}(\text{RootOf}(_Z^6-54)^2-24*_Z*\text{RootOf}(_Z^6-54)+576*_Z^2)*(-3*x^2+2)^{(1/3)}-18*(-3*x^2+2)^{(1/3)})*\text{RootOf}(_Z^6-54)^2+54*(-3*x^2+2)^{(2/3)})/(x^2-6))*\text{RootOf}(_Z^6-54))/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 - 6d)(-3x^2 + 2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/3)/(d*x^2-6*d),x, algorithm="maxima")

[Out] integrate(1/((d*x^2 - 6*d)*(-3*x^2 + 2)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(2 - 3x^2)^{1/3} (6d - dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((2 - 3*x^2)^(1/3)*(6*d - d*x^2)),x)

[Out] -int(1/((2 - 3*x^2)^(1/3)*(6*d - d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{1}{x^2 \sqrt[3]{2-3x^2} - 6 \sqrt[3]{2-3x^2}} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x**2+2)**(1/3)/(d*x**2-6*d),x)

[Out] Integral(1/(x**2*(2 - 3*x**2)**(1/3) - 6*(2 - 3*x**2)**(1/3)), x)/d

$$3.158 \quad \int \frac{1}{\sqrt[3]{-2+3x^2}(-6d+dx^2)} dx$$

Optimal. Leaf size=119

$$\frac{\tan^{-1}\left(\frac{\sqrt[6]{2}(\sqrt[3]{3x^2-2}+\sqrt[3]{2})}{x}\right)}{4 \cdot 2^{5/6}d} - \frac{\tanh^{-1}\left(\frac{(\sqrt[3]{3x^2-2}+\sqrt[3]{2})^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} + \frac{\tanh^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d}$$

[Out] 1/8*arctan(2^(1/6)*(2^(1/3)+(3*x^2-2)^(1/3))/x)*2^(1/6)/d-1/24*arctanh(1/18*(2^(1/3)+(3*x^2-2)^(1/3))^2*2^(5/6)/x*3^(1/2))*2^(1/6)/d*3^(1/2)+1/24*arctanh(1/6*x*6^(1/2))*2^(1/6)/d*3^(1/2)

Rubi [A] time = 0.02, antiderivative size = 119, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {395}

$$\frac{\tan^{-1}\left(\frac{\sqrt[6]{2}(\sqrt[3]{3x^2-2}+\sqrt[3]{2})}{x}\right)}{4 \cdot 2^{5/6}d} - \frac{\tanh^{-1}\left(\frac{(\sqrt[3]{3x^2-2}+\sqrt[3]{2})^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} + \frac{\tanh^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d}$$

Antiderivative was successfully verified.

[In] Int[1/((-2 + 3*x^2)^(1/3)*(-6*d + d*x^2)),x]

[Out] ArcTan[(2^(1/6)*(2^(1/3) + (-2 + 3*x^2)^(1/3)))/x]/(4*2^(5/6)*d) + ArcTanh[x/Sqrt[6]]/(4*2^(5/6)*Sqrt[3]*d) - ArcTanh[(2^(1/3) + (-2 + 3*x^2)^(1/3))^2/(3*2^(1/6)*Sqrt[3]*x)]/(4*2^(5/6)*Sqrt[3]*d)

Rule 395

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b/a), 2]}, -Simp[(q*ArcTanh[(q*x)/3])/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTanh[(Rt[a, 3] - (a + b*x^2)^(1/3))^2/(3*Rt[a, 3]^2*q*x)])/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTan[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))/(Rt[a, 3]*q*x)])/(4*Sqrt[3]*Rt[a, 3]*d), x]) /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && NegQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{-2+3x^2}(-6d+dx^2)} dx = \frac{\tan^{-1}\left(\frac{\sqrt[6]{2}(\sqrt[3]{2}+\sqrt[3]{-2+3x^2})}{x}\right)}{4 \cdot 2^{5/6}d} + \frac{\tanh^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4 \cdot 2^{5/6}\sqrt{3}d} - \frac{\tanh^{-1}\left(\frac{(\sqrt[3]{2}+\sqrt[3]{-2+3x^2})^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4 \cdot 2^{5/6}\sqrt{3}d}$$

Mathematica [C] time = 0.10, size = 136, normalized size = 1.14

$$\frac{9x {}_2F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{3x^2}{2}, \frac{x^2}{6}\right)}{d(x^2-6)\sqrt[3]{3x^2-2}\left(x^2\left({}_2F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; \frac{3x^2}{2}, \frac{x^2}{6}\right) + 3{}_2F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; \frac{3x^2}{2}, \frac{x^2}{6}\right)\right) + 9{}_2F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; \frac{3x^2}{2}, \frac{x^2}{6}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2 + 3*x^2)^(1/3)*(-6*d + d*x^2)),x]

```
[Out] (9*x*AppellF1[1/2, 1/3, 1, 3/2, (3*x^2)/2, x^2/6])/(d*(-6 + x^2)*(-2 + 3*x^2)^(1/3)*(9*AppellF1[1/2, 1/3, 1, 3/2, (3*x^2)/2, x^2/6] + x^2*(AppellF1[3/2, 1/3, 2, 5/2, (3*x^2)/2, x^2/6] + 3*AppellF1[3/2, 4/3, 1, 5/2, (3*x^2)/2, x^2/6])))
```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(3*x^2-2)^(1/3)/(d*x^2-6*d),x, algorithm="fricas")
```

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 - 6d)(3x^2 - 2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(3*x^2-2)^(1/3)/(d*x^2-6*d),x, algorithm="giac")
```

[Out] integrate(1/((d*x^2 - 6*d)*(3*x^2 - 2)^(1/3)), x)

maple [C] time = 58.82, size = 1063, normalized size = 8.93

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(3*x^2-2)^(1/3)/(d*x^2-6*d),x)
```

```
[Out] -1/24*(24*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*ln(-(4608*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*RootOf(_Z^6-54)^5*x-288*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-54)^6*x+4*RootOf(_Z^6-54)^7*x+6912*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*x-144*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)^4*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*x+216*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*x^2-9*x^2*RootOf(_Z^6-54)^4+432*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-54)^3-18*RootOf(_Z^6-54)^4-2592*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)-324*(3*x^2-2)^(2/3))/(x^2-6))+24*ln((768*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*RootOf(_Z^6-54)^5*x-16*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-54)^6*x+1152*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*x-72*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)^4*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*x+RootOf(_Z^6-54)^5*(3*x^2-2)^(1/3)*x+36*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*x^2+72*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-54)^3-432*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)+18*RootOf(_Z^6-54)^2*(3*x^2-2)^(1/3)+54*(3*x^2-2)^(2/3))/(x^2-6))*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)-ln((768*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*RootOf(_Z^6-54)^5*x-16*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-54)^6*x+1152*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)^2*x-72*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)^4*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*x+RootOf(_Z^6-54)^5*(3*x^2-2)^(1/3)*x+36*RootOf(_Z^6-54)^3*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*x^2+72*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)*RootOf(_Z^6-54)^3-432*(3*x^2-2)^(1/3)*RootOf(_Z^6-54)*RootOf(RootOf(_Z^6-54)^2-24*_Z*RootOf(_Z^6-54)+576*_Z^2)+18*RootOf(_Z^6-54)^2*(3*x^2-2)^(1/3)+54*(3*x^2-2)^(2/3))/(x^2-6))
```

$$\sqrt[6]{-54} + 576 \sqrt[2]{Z} * x^2 + 72 * \text{RootOf}(\text{RootOf}(\sqrt[6]{-54})^2 - 24 * \sqrt[2]{Z} * \text{RootOf}(\sqrt[6]{-54}) + 576 * \sqrt[2]{Z} * \text{RootOf}(\sqrt[6]{-54})^3 - 432 * (3 * x^2 - 2)^{1/3} * \text{RootOf}(\sqrt[6]{-54}) * \text{RootOf}(\text{RootOf}(\sqrt[6]{-54})^2 - 24 * \sqrt[2]{Z} * \text{RootOf}(\sqrt[6]{-54}) + 576 * \sqrt[2]{Z} + 18 * \text{RootOf}(\sqrt[6]{-54})^2 * (3 * x^2 - 2)^{1/3} + 54 * (3 * x^2 - 2)^{2/3})) / (x^2 - 6)) * \text{RootOf}(\sqrt[6]{-54})) / d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 - 6d)(3x^2 - 2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2-2)^(1/3)/(d*x^2-6*d),x, algorithm="maxima")

[Out] integrate(1/((d*x^2 - 6*d)*(3*x^2 - 2)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$- \int \frac{1}{(3x^2 - 2)^{1/3} (6d - dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((3*x^2 - 2)^(1/3)*(6*d - d*x^2)),x)

[Out] -int(1/((3*x^2 - 2)^(1/3)*(6*d - d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{1}{x^2 \sqrt[3]{3x^2-2} - 6 \sqrt[3]{3x^2-2}} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x**2-2)**(1/3)/(d*x**2-6*d),x)

[Out] Integral(1/(x**2*(3*x**2 - 2)**(1/3) - 6*(3*x**2 - 2)**(1/3)), x)/d

$$3.159 \quad \int \frac{1}{\sqrt[3]{-2-3x^2} (6d+dx^2)} dx$$

Optimal. Leaf size=119

$$-\frac{\tan^{-1}\left(\frac{\left(\sqrt[3]{-3x^2-2}+\sqrt[3]{2}\right)^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4\ 2^{5/6}\sqrt{3}d} + \frac{\tanh^{-1}\left(\frac{\sqrt[6]{2}\left(\sqrt[3]{-3x^2-2}+\sqrt[3]{2}\right)}{x}\right)}{4\ 2^{5/6}d} - \frac{\tan^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4\ 2^{5/6}\sqrt{3}d}$$

[Out] 1/8*arctanh(2^(1/6)*(2^(1/3)+(-3*x^2-2)^(1/3))/x)*2^(1/6)/d-1/24*arctan(1/18*(2^(1/3)+(-3*x^2-2)^(1/3))^2*2^(5/6)/x*3^(1/2))*2^(1/6)/d*3^(1/2)-1/24*arctan(1/6*x*6^(1/2))*2^(1/6)/d*3^(1/2)

Rubi [A] time = 0.02, antiderivative size = 119, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {394}

$$-\frac{\tan^{-1}\left(\frac{\left(\sqrt[3]{-3x^2-2}+\sqrt[3]{2}\right)^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4\ 2^{5/6}\sqrt{3}d} + \frac{\tanh^{-1}\left(\frac{\sqrt[6]{2}\left(\sqrt[3]{-3x^2-2}+\sqrt[3]{2}\right)}{x}\right)}{4\ 2^{5/6}d} - \frac{\tan^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4\ 2^{5/6}\sqrt{3}d}$$

Antiderivative was successfully verified.

[In] Int[1/((-2 - 3*x^2)^(1/3)*(6*d + d*x^2)),x]

[Out] -ArcTan[x/Sqrt[6]]/(4*2^(5/6)*Sqrt[3]*d) - ArcTan[(2^(1/3) + (-2 - 3*x^2)^(1/3))^2/(3*2^(1/6)*Sqrt[3]*x)]/(4*2^(5/6)*Sqrt[3]*d) + ArcTanh[(2^(1/6)*(2^(1/3) + (-2 - 3*x^2)^(1/3)))/x]/(4*2^(5/6)*d)

Rule 394

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[(q*ArcTan[(q*x)/3])/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTan[(Rt[a, 3] - (a + b*x^2)^(1/3))^2/(3*Rt[a, 3]^2*q*x)])/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTanh[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))/(Rt[a, 3]*q*x)])/(4*Sqrt[3]*Rt[a, 3]*d), x]]) /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{-2-3x^2} (6d+dx^2)} dx = -\frac{\tan^{-1}\left(\frac{x}{\sqrt{6}}\right)}{4\ 2^{5/6}\sqrt{3}d} - \frac{\tan^{-1}\left(\frac{\left(\sqrt[3]{2}+\sqrt[3]{-2-3x^2}\right)^2}{3\sqrt[6]{2}\sqrt{3}x}\right)}{4\ 2^{5/6}\sqrt{3}d} + \frac{\tanh^{-1}\left(\frac{\sqrt[6]{2}\left(\sqrt[3]{2}+\sqrt[3]{-2-3x^2}\right)}{x}\right)}{4\ 2^{5/6}d}$$

Mathematica [C] time = 0.11, size = 136, normalized size = 1.14

$$\frac{9x F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{3x^2}{2}, -\frac{x^2}{6}\right)}{d\sqrt[3]{-3x^2-2} (x^2+6) \left(x^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -\frac{3x^2}{2}, -\frac{x^2}{6}\right) + 3F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -\frac{3x^2}{2}, -\frac{x^2}{6}\right)\right) - 9F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -\frac{3x^2}{2}, -\frac{x^2}{6}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2 - 3*x^2)^(1/3)*(6*d + d*x^2)),x]

```
[Out] (-9*x*AppellF1[1/2, 1/3, 1, 3/2, (-3*x^2)/2, -1/6*x^2])/(d*(-2 - 3*x^2)^(1/3)*(6 + x^2)*(-9*AppellF1[1/2, 1/3, 1, 3/2, (-3*x^2)/2, -1/6*x^2] + x^2*(AppellF1[3/2, 1/3, 2, 5/2, (-3*x^2)/2, -1/6*x^2] + 3*AppellF1[3/2, 4/3, 1, 5/2, (-3*x^2)/2, -1/6*x^2])))
```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-3*x^2-2)^(1/3)/(d*x^2+6*d),x, algorithm="fricas")
```

```
[Out] Timed out
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 + 6d)(-3x^2 - 2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-3*x^2-2)^(1/3)/(d*x^2+6*d),x, algorithm="giac")
```

```
[Out] integrate(1/((d*x^2 + 6*d)*(-3*x^2 - 2)^(1/3)), x)
```

maple [C] time = 77.11, size = 725, normalized size = 6.09

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(-3*x^2-2)^(1/3)/(d*x^2+6*d),x)
```

```
[Out] -1/24*(RootOf(_Z^6+54)*ln(-(4*RootOf(_Z^6+54)^7*x-288*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)^6*x+4608*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)^2*RootOf(_Z^6+54)^5*x+144*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)^4*(-3*x^2-2)^(1/3)*x-6912*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)^2*RootOf(_Z^6+54)^3*(-3*x^2-2)^(1/3)*x-9*x^2*RootOf(_Z^6+54)^4+216*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)^3*x^2+18*RootOf(_Z^6+54)^4-432*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)^3-2592*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)*(-3*x^2-2)^(1/3)+324*(-3*x^2-2)^(2/3))/(x^2+6))+ln(-(4*RootOf(_Z^6+54)^7*x+192*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)^6*x-6*(-3*x^2-2)^(1/3)*RootOf(_Z^6+54)^5*x+288*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)^4*(-3*x^2-2)^(1/3)*x+9*x^2*RootOf(_Z^6+54)^4-18*RootOf(_Z^6+54)^4+108*(-3*x^2-2)^(1/3)*RootOf(_Z^6+54)^2+324*(-3*x^2-2)^(2/3))/(x^2+6))*RootOf(_Z^6+54)-24*ln(-(4*RootOf(_Z^6+54)^7*x+192*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)^6*x-6*(-3*x^2-2)^(1/3)*RootOf(_Z^6+54)^5*x+288*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2)*RootOf(_Z^6+54)^4*(-3*x^2-2)^(1/3)*x+9*x^2*RootOf(_Z^6+54)^4-18*RootOf(_Z^6+54)^4+108*(-3*x^2-2)^(1/3)*RootOf(_Z^6+54)^2+324*(-3*x^2-2)^(2/3))/(x^2+6))*RootOf(RootOf(_Z^6+54)^2-24*_Z*RootOf(_Z^6+54)+576*_Z^2))/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 + 6d)(-3x^2 - 2)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2-2)^(1/3)/(d*x^2+6*d),x, algorithm="maxima")

[Out] integrate(1/((d*x^2 + 6*d)*(-3*x^2 - 2)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(-3x^2 - 2)^{1/3} (dx^2 + 6d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((- 3*x^2 - 2)^(1/3)*(6*d + d*x^2)),x)

[Out] int(1/((- 3*x^2 - 2)^(1/3)*(6*d + d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^2 \sqrt[3]{-3x^2-2} + 6 \sqrt[3]{-3x^2-2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x**2-2)**(1/3)/(d*x**2+6*d),x)

[Out] Integral(1/(x**2*(-3*x**2 - 2)**(1/3) + 6*(-3*x**2 - 2)**(1/3)), x)/d

$$3.160 \quad \int \frac{1}{\sqrt[3]{1+x^2} (9+x^2)} dx$$

Optimal. Leaf size=70

$$\frac{1}{12} \tan^{-1} \left(\frac{(1 - \sqrt[3]{x^2+1})^2}{3x} \right) - \frac{\tanh^{-1} \left(\frac{\sqrt{3}(1 - \sqrt[3]{x^2+1})}{x} \right)}{4\sqrt{3}} + \frac{1}{12} \tan^{-1} \left(\frac{x}{3} \right)$$

[Out] 1/12*arctan(1/3*x)+1/12*arctan(1/3*(1-(x^2+1)^(1/3))^2/x)-1/12*arctanh((1-(x^2+1)^(1/3))*3^(1/2)/x)*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {394}

$$\frac{1}{12} \tan^{-1} \left(\frac{(1 - \sqrt[3]{x^2+1})^2}{3x} \right) - \frac{\tanh^{-1} \left(\frac{\sqrt{3}(1 - \sqrt[3]{x^2+1})}{x} \right)}{4\sqrt{3}} + \frac{1}{12} \tan^{-1} \left(\frac{x}{3} \right)$$

Antiderivative was successfully verified.

[In] Int[1/((1 + x^2)^(1/3)*(9 + x^2)), x]

[Out] ArcTan[x/3]/12 + ArcTan[(1 - (1 + x^2)^(1/3))^2/(3*x)]/12 - ArcTanh[(Sqrt[3]*(1 - (1 + x^2)^(1/3)))/x]/(4*Sqrt[3])

Rule 394

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[(q*ArcTan[(q*x)/3])/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTan[(Rt[a, 3] - (a + b*x^2)^(1/3))^2/(3*Rt[a, 3]^2*q*x)])/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTanh[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))/(Rt[a, 3]*q*x)])/(4*Sqrt[3]*Rt[a, 3]*d), x]]) /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{1+x^2} (9+x^2)} dx = \frac{1}{12} \tan^{-1} \left(\frac{x}{3} \right) + \frac{1}{12} \tan^{-1} \left(\frac{(1 - \sqrt[3]{1+x^2})^2}{3x} \right) - \frac{\tanh^{-1} \left(\frac{\sqrt{3}(1 - \sqrt[3]{1+x^2})}{x} \right)}{4\sqrt{3}}$$

Mathematica [C] time = 0.10, size = 124, normalized size = 1.77

$$\frac{27xF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -x^2, -\frac{x^2}{9}\right)}{\sqrt[3]{x^2+1} (x^2+9) \left(2x^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -x^2, -\frac{x^2}{9}\right) + 3F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -x^2, -\frac{x^2}{9}\right)\right) - 27F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -x^2, -\frac{x^2}{9}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((1 + x^2)^(1/3)*(9 + x^2)), x]

```
[Out] (-27*x*AppellF1[1/2, 1/3, 1, 3/2, -x^2, -1/9*x^2])/((1 + x^2)^(1/3)*(9 + x^2)*(-27*AppellF1[1/2, 1/3, 1, 3/2, -x^2, -1/9*x^2] + 2*x^2*(AppellF1[3/2, 1/3, 2, 5/2, -x^2, -1/9*x^2] + 3*AppellF1[3/2, 4/3, 1, 5/2, -x^2, -1/9*x^2]))
```

fricas [B] time = 5.41, size = 1395, normalized size = 19.93

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(x^2+1)^(1/3)/(x^2+9),x, algorithm="fricas")
```

```
[Out] 1/144*sqrt(3)*log(4*(x^6 + 1647*x^4 + 891*x^2 + 18*(3*x^4 + 32*sqrt(3)*x^3 + 126*x^2 + 27)*(x^2 + 1)^(2/3) + 108*sqrt(3)*(x^5 + 10*x^3 + 9*x) + 6*(81*x^4 + 162*x^2 + sqrt(3)*(x^5 + 210*x^3 + 81*x) + 81)*(x^2 + 1)^(1/3) - 243)/(x^6 + 27*x^4 + 243*x^2 + 729)) - 1/144*sqrt(3)*log(4*(x^6 + 1647*x^4 + 891*x^2 + 18*(3*x^4 - 32*sqrt(3)*x^3 + 126*x^2 + 27)*(x^2 + 1)^(2/3) - 108*sqrt(3)*(x^5 + 10*x^3 + 9*x) + 6*(81*x^4 + 162*x^2 - sqrt(3)*(x^5 + 210*x^3 + 81*x) + 81)*(x^2 + 1)^(1/3) - 243)/(x^6 + 27*x^4 + 243*x^2 + 729)) - 1/36*arctan((384*x^11 - 130320*x^9 + 2379456*x^7 - 629856*x^5 - 1259712*x^3 + 36*(388*x^9 - 27864*x^7 + 303264*x^5 + 17496*x^3 + sqrt(3)*(x^10 + 549*x^8 - 8046*x^6 + 129762*x^4 - 19683*x^2 + 59049) - 236196*x)*(x^2 + 1)^(2/3) + sqrt(3)*(x^12 - 234*x^10 + 229311*x^8 - 1214028*x^6 + 6816879*x^4 + 6022998*x^2 + 531441) + 2*(x^12 + 50616*x^10 - 1869399*x^8 - 3773304*x^6 - 6908733*x^4 + 72*(x^10 + 1620*x^8 - 63666*x^6 - 43740*x^4 + 59049*x^2 + 12*sqrt(3)*(11*x^9 - 261*x^7 - 6075*x^5 - 2187*x^3))*(x^2 + 1)^(2/3) + 6*sqrt(3)*(43*x^11 + 14055*x^9 - 563922*x^7 - 1307826*x^5 - 898857*x^3 + 177147*x) + 6*(453*x^10 + 21141*x^8 - 1483758*x^6 - 1404054*x^4 - 885735*x^2 + sqrt(3)*(x^11 + 8985*x^9 - 349110*x^7 + 118098*x^5 + 32805*x^3 - 177147*x) + 531441)*(x^2 + 1)^(1/3) + 1594323)*sqrt((x^6 + 1647*x^4 + 891*x^2 + 18*(3*x^4 - 32*sqrt(3)*x^3 + 126*x^2 + 27)*(x^2 + 1)^(2/3) - 108*sqrt(3)*(x^5 + 10*x^3 + 9*x) + 6*(81*x^4 + 162*x^2 - sqrt(3)*(x^5 + 210*x^3 + 81*x) + 81)*(x^2 + 1)^(1/3) - 243)/(x^6 + 27*x^4 + 243*x^2 + 729)) + 12*(x^11 - 6423*x^9 + 225018*x^7 - 1106622*x^5 - 1541835*x^3 + 3*sqrt(3)*(37*x^10 - 675*x^8 + 34722*x^6 - 97686*x^4 + 59049*x^2 + 59049) - 177147*x)*(x^2 + 1)^(1/3) - 8503056*x)/(x^12 - 48978*x^10 + 2332071*x^8 - 16419996*x^6 - 24151041*x^4 - 9565938*x^2 + 4782969)) + 1/36*arctan(-(384*x^11 - 130320*x^9 + 2379456*x^7 - 629856*x^5 - 1259712*x^3 + 36*(388*x^9 - 27864*x^7 + 303264*x^5 + 17496*x^3 - sqrt(3)*(x^10 + 549*x^8 - 8046*x^6 + 129762*x^4 - 19683*x^2 + 59049) - 236196*x)*(x^2 + 1)^(2/3) - sqrt(3)*(x^12 - 234*x^10 + 229311*x^8 - 1214028*x^6 + 6816879*x^4 + 6022998*x^2 + 531441) + 2*(x^12 + 50616*x^10 - 1869399*x^8 - 3773304*x^6 - 6908733*x^4 + 72*(x^10 + 1620*x^8 - 63666*x^6 - 43740*x^4 + 59049*x^2 - 12*sqrt(3)*(11*x^9 - 261*x^7 - 6075*x^5 - 2187*x^3))*(x^2 + 1)^(2/3) - 6*sqrt(3)*(43*x^11 + 14055*x^9 - 563922*x^7 - 1307826*x^5 - 898857*x^3 + 177147*x) + 6*(453*x^10 + 21141*x^8 - 1483758*x^6 - 1404054*x^4 - 885735*x^2 - sqrt(3)*(x^11 + 8985*x^9 - 349110*x^7 + 118098*x^5 + 32805*x^3 - 177147*x) + 531441)*(x^2 + 1)^(1/3) + 1594323)*sqrt((x^6 + 1647*x^4 + 891*x^2 + 18*(3*x^4 + 32*sqrt(3)*x^3 + 126*x^2 + 27)*(x^2 + 1)^(2/3) + 108*sqrt(3)*(x^5 + 10*x^3 + 9*x) + 6*(81*x^4 + 162*x^2 + sqrt(3)*(x^5 + 210*x^3 + 81*x) + 81)*(x^2 + 1)^(1/3) - 243)/(x^6 + 27*x^4 + 243*x^2 + 729)) + 12*(x^11 - 6423*x^9 + 225018*x^7 - 1106622*x^5 - 1541835*x^3 - 3*sqrt(3)*(37*x^10 - 675*x^8 + 34722*x^6 - 97686*x^4 + 59049*x^2 + 59049) - 177147*x)*(x^2 + 1)^(1/3) - 8503056*x)/(x^12 - 48978*x^10 + 2332071*x^8 - 16419996*x^6 - 24151041*x^4 - 9565938*x^2 + 4782969)) - 1/36*arctan(6*(11*x^5 + 30*x^3 + 6*(23*x^3 + 27*x)*(x^2 + 1)^(2/3) + (x^5 - 240*x^3 - 81*x)*(x^2 + 1)^(1/3) - 81*x)/(x^6 - 1971*x^4 - 1701*x^2 - 729))
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(x^2 + 9)(x^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/3)/(x^2+9),x, algorithm="giac")

[Out] integrate(1/((x^2 + 9)*(x^2 + 1)^(1/3)), x)

maple [C] time = 10.59, size = 512, normalized size = 7.31

$$-\text{RootOf}\left(144_Z^2 + 12_Z \text{RootOf}\left(_Z^2 + 1\right) - 1\right) \ln \left(\frac{-x^2 + 48(x^2 + 1)^{\frac{1}{3}} x \text{RootOf}\left(144_Z^2 + 12_Z \text{RootOf}\left(_Z^2 + 1\right) - 1\right)}{\dots} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2+1)^(1/3)/(x^2+9),x)

[Out] 1/12*RootOf(_Z^2+1)*ln((24*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*RootOf(_Z^2+1)^2*(x^2+1)^(1/3)*x+576*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)^2*RootOf(_Z^2+1)*(x^2+1)^(1/3)*x-48*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*RootOf(_Z^2+1)^2*x-1152*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)^2*RootOf(_Z^2+1)*x-12*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*RootOf(_Z^2+1)*x^2+72*(x^2+1)^(1/3)*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*RootOf(_Z^2+1)+36*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*RootOf(_Z^2+1)+4*RootOf(_Z^2+1)*x+96*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*x-6*(x^2+1)^(2/3)+x^2-3)/(x^2+9))-ln((2*(x^2+1)^(1/3)*RootOf(_Z^2+1)*x+48*(x^2+1)^(1/3)*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*x+4*RootOf(_Z^2+1)*x+96*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*x-6*(x^2+1)^(2/3)-x^2-6*(x^2+1)^(1/3)+3)/(x^2+9))*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)-1/12*ln((2*(x^2+1)^(1/3)*RootOf(_Z^2+1)*x+48*(x^2+1)^(1/3)*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*x+4*RootOf(_Z^2+1)*x+96*RootOf(12*_Z*RootOf(_Z^2+1)+144*_Z^2-1)*x-6*(x^2+1)^(2/3)-x^2-6*(x^2+1)^(1/3)+3)/(x^2+9))*RootOf(_Z^2+1)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(x^2 + 9)(x^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/3)/(x^2+9),x, algorithm="maxima")

[Out] integrate(1/((x^2 + 9)*(x^2 + 1)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(x^2 + 1)^{\frac{1}{3}}(x^2 + 9)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/3)*(x^2 + 9)),x)

[Out] int(1/((x^2 + 1)^(1/3)*(x^2 + 9)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{x^2 + 1}(x^2 + 9)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(x**2+1)**(1/3)/(x**2+9),x)
```

```
[Out] Integral(1/((x**2 + 1)**(1/3)*(x**2 + 9)), x)
```

$$3.161 \quad \int \frac{1}{\sqrt[3]{1+bx^2} (9+bx^2)} dx$$

Optimal. Leaf size=104

$$\frac{\tan^{-1}\left(\frac{(1-\sqrt[3]{bx^2+1})^2}{3\sqrt{bx^2+1}}\right)}{12\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{bx^2+1})}{\sqrt{bx^2+1}}\right)}{4\sqrt{3}\sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{bx^2+1}}{3}\right)}{12\sqrt{b}}$$

[Out] 1/12*arctan(1/3*(1-(b*x^2+1)^(1/3))^2/x/b^(1/2))/b^(1/2)+1/12*arctan(1/3*x*b^(1/2))/b^(1/2)-1/12*arctanh((1-(b*x^2+1)^(1/3))*3^(1/2)/x/b^(1/2))*3^(1/2)/b^(1/2)

Rubi [A] time = 0.02, antiderivative size = 104, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {394}

$$\frac{\tan^{-1}\left(\frac{(1-\sqrt[3]{bx^2+1})^2}{3\sqrt{bx^2+1}}\right)}{12\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{bx^2+1})}{\sqrt{bx^2+1}}\right)}{4\sqrt{3}\sqrt{b}} + \frac{\tan^{-1}\left(\frac{\sqrt{bx^2+1}}{3}\right)}{12\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((1 + b*x^2)^(1/3)*(9 + b*x^2)),x]

[Out] ArcTan[(Sqrt[b]*x)/3]/(12*Sqrt[b]) + ArcTan[(1 - (1 + b*x^2)^(1/3))^2/(3*Sqrt[b]*x)]/(12*Sqrt[b]) - ArcTanh[(Sqrt[3]*(1 - (1 + b*x^2)^(1/3)))/(Sqrt[b]*x)]/(4*Sqrt[3]*Sqrt[b])

Rule 394

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[(q*ArcTan[(q*x)/3])/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTan[(Rt[a, 3] - (a + b*x^2)^(1/3))^2/(3*Rt[a, 3]^2*q*x)])/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTanh[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))/(Rt[a, 3]*q*x)])/(4*Sqrt[3]*Rt[a, 3]*d), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && PosQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{1+bx^2} (9+bx^2)} dx = \frac{\tan^{-1}\left(\frac{\sqrt{bx^2+1}}{3}\right)}{12\sqrt{b}} + \frac{\tan^{-1}\left(\frac{(1-\sqrt[3]{1+bx^2})^2}{3\sqrt{1+bx^2}}\right)}{12\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{1+bx^2})}{\sqrt{1+bx^2}}\right)}{4\sqrt{3}\sqrt{b}}$$

Mathematica [C] time = 0.11, size = 137, normalized size = 1.32

$$\frac{27xF_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -bx^2, -\frac{bx^2}{9}\right)}{\sqrt[3]{bx^2+1} (bx^2+9) \left(2bx^2 \left(F_1\left(\frac{3}{2}; \frac{1}{3}, 2; \frac{5}{2}; -bx^2, -\frac{bx^2}{9}\right) + 3F_1\left(\frac{3}{2}; \frac{4}{3}, 1; \frac{5}{2}; -bx^2, -\frac{bx^2}{9}\right)\right) - 27F_1\left(\frac{1}{2}; \frac{1}{3}, 1; \frac{3}{2}; -bx^2, -\frac{bx^2}{9}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((1 + b*x^2)^(1/3)*(9 + b*x^2)),x]

[Out] $(-27*x*AppellF1[1/2, 1/3, 1, 3/2, -(b*x^2), -1/9*(b*x^2)]/((1 + b*x^2)^{(1/3)}*(9 + b*x^2)*(-27*AppellF1[1/2, 1/3, 1, 3/2, -(b*x^2), -1/9*(b*x^2)] + 2*b*x^2*(AppellF1[3/2, 1/3, 2, 5/2, -(b*x^2), -1/9*(b*x^2)] + 3*AppellF1[3/2, 4/3, 1, 5/2, -(b*x^2), -1/9*(b*x^2)])))$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x^2+1)^(1/3)/(b*x^2+9),x, algorithm="fricas")`

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 9)(bx^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x^2+1)^(1/3)/(b*x^2+9),x, algorithm="giac")`

[Out] `integrate(1/((b*x^2 + 9)*(b*x^2 + 1)^(1/3)), x)`

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 1)^{\frac{1}{3}}(bx^2 + 9)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b*x^2+1)^(1/3)/(b*x^2+9),x)`

[Out] `int(1/(b*x^2+1)^(1/3)/(b*x^2+9),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + 9)(bx^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x^2+1)^(1/3)/(b*x^2+9),x, algorithm="maxima")`

[Out] `integrate(1/((b*x^2 + 9)*(b*x^2 + 1)^(1/3)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + 1)^{\frac{1}{3}}(bx^2 + 9)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((b*x^2 + 1)^(1/3)*(b*x^2 + 9)),x)`

[Out] `int(1/((b*x^2 + 1)^(1/3)*(b*x^2 + 9)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{bx^2 + 1}(bx^2 + 9)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x**2+1)**(1/3)/(b*x**2+9),x)
```

```
[Out] Integral(1/((b*x**2 + 1)**(1/3)*(b*x**2 + 9)), x)
```

$$3.162 \quad \int \frac{1}{\sqrt[3]{1-x^2} (9-x^2)} dx$$

Optimal. Leaf size=74

$$\frac{\tan^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{1-x^2})}{x}\right)}{4\sqrt{3}} - \frac{1}{12} \tanh^{-1}\left(\frac{(1-\sqrt[3]{1-x^2})^2}{3x}\right) + \frac{1}{12} \tanh^{-1}\left(\frac{x}{3}\right)$$

[Out] 1/12*arctanh(1/3*x)-1/12*arctanh(1/3*(1-(-x^2+1)^(1/3))^2/x)+1/12*arctan((1-(-x^2+1)^(1/3))*3^(1/2)/x)*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {395}

$$\frac{\tan^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{1-x^2})}{x}\right)}{4\sqrt{3}} - \frac{1}{12} \tanh^{-1}\left(\frac{(1-\sqrt[3]{1-x^2})^2}{3x}\right) + \frac{1}{12} \tanh^{-1}\left(\frac{x}{3}\right)$$

Antiderivative was successfully verified.

[In] Int[1/((1 - x^2)^(1/3)*(9 - x^2)),x]

[Out] ArcTan[(Sqrt[3]*(1 - (1 - x^2)^(1/3)))/x]/(4*Sqrt[3]) + ArcTanh[x/3]/12 - ArcTanh[(1 - (1 - x^2)^(1/3))^2/(3*x)]/12

Rule 395

Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b/a), 2]}, -Simp[(q*ArcTanh[(q*x)/3]]/(12*Rt[a, 3]*d), x] + (Simp[(q*ArcTanh[(Rt[a, 3] - (a + b*x^2)^(1/3))]^2/(3*Rt[a, 3]^2*q*x)]/(12*Rt[a, 3]*d), x] - Simp[(q*ArcTan[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))]/(Rt[a, 3]*q*x)]/(4*Sqrt[3]*Rt[a, 3]*d), x)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && NegQ[b/a]

Rubi steps

$$\int \frac{1}{\sqrt[3]{1-x^2} (9-x^2)} dx = \frac{\tan^{-1}\left(\frac{\sqrt{3}(1-\sqrt[3]{1-x^2})}{x}\right)}{4\sqrt{3}} + \frac{1}{12} \tanh^{-1}\left(\frac{x}{3}\right) - \frac{1}{12} \tanh^{-1}\left(\frac{(1-\sqrt[3]{1-x^2})^2}{3x}\right)$$

Mathematica [C] time = 0.05, size = 125, normalized size = 1.69

$$\frac{\sqrt[3]{x-1} \sqrt[3]{x+1} F_1\left(\frac{2}{3}; \frac{1}{3}, \frac{1}{3}; \frac{5}{3}; -\frac{4}{x-3}, -\frac{2}{x-3}\right) - \sqrt[3]{x-1} \sqrt[3]{x+1} F_1\left(\frac{2}{3}; \frac{1}{3}, \frac{1}{3}; \frac{5}{3}; \frac{2}{x+3}, \frac{4}{x+3}\right)}{4\sqrt[3]{1-x^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((1 - x^2)^(1/3)*(9 - x^2)),x]

[Out] (((-1 + x)/(-3 + x))^(1/3)*((1 + x)/(-3 + x))^(1/3)*AppellF1[2/3, 1/3, 1/3, 5/3, -4/(-3 + x), -2/(-3 + x)] - ((-1 + x)/(3 + x))^(1/3)*((1 + x)/(3 + x))^(1/3)*AppellF1[2/3, 1/3, 1/3, 5/3, 2/(3 + x), 4/(3 + x)])/4\sqrt[3]{1-x^2}

)^(1/3)*AppellF1[2/3, 1/3, 1/3, 5/3, 2/(3 + x), 4/(3 + x)]/(4*(1 - x^2)^(1/3))

fricas [B] time = 1.72, size = 269, normalized size = 3.64

$$-\frac{1}{36} \sqrt{3} \arctan \left(\frac{36 \sqrt{3} (x^4 - 32x^3 - 42x^2 + 9)(-x^2 + 1)^{\frac{2}{3}} + 12 \sqrt{3} (x^5 + 27x^4 - 210x^3 - 54x^2 + 81x + 27)}{3(x^6 + 108x^5 - 1647x^4 - 1080x^3 + 891x^2 + 972x + 243)} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/3)/(-x^2+9),x, algorithm="fricas")

[Out] -1/36*sqrt(3)*arctan(1/3*(36*sqrt(3)*(x^4 - 32*x^3 - 42*x^2 + 9)*(-x^2 + 1)^(2/3) + 12*sqrt(3)*(x^5 + 27*x^4 - 210*x^3 - 54*x^2 + 81*x + 27)*(-x^2 + 1)^(1/3) + sqrt(3)*(x^6 - 108*x^5 - 567*x^4 + 1080*x^3 + 459*x^2 - 972*x - 405))/(x^6 + 108*x^5 - 1647*x^4 - 1080*x^3 + 891*x^2 + 972*x + 243)) - 1/72*log((x^3 + 33*x^2 + 18*(-x^2 + 1)^(2/3)*(x + 1) - 6*(x^2 + 6*x - 3)*(-x^2 + 1)^(1/3) - 9*x - 9)/(x^3 + 9*x^2 + 27*x + 27)) + 1/36*log(-(-x^3 - 33*x^2 + 18*(-x^2 + 1)^(2/3)*(x - 1) + 6*(x^2 - 6*x - 3)*(-x^2 + 1)^(1/3) - 9*x + 9)/(x^3 + 9*x^2 + 27*x + 27))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(x^2 - 9)(-x^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/3)/(-x^2+9),x, algorithm="giac")

[Out] integrate(-1/((x^2 - 9)*(-x^2 + 1)^(1/3)), x)

maple [C] time = 2.61, size = 539, normalized size = 7.28

$$-\text{RootOf}(144_Z^2 + 12_Z + 1) \ln \left(\frac{-6x^2 \text{RootOf}(144_Z^2 + 12_Z + 1) + 288(-x^2 + 1)^{\frac{1}{3}} x \text{RootOf}(144_Z^2 + 12_Z + 1)}{\dots} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/3)/(-x^2+9),x)

[Out] -1/12*ln((288*RootOf(144*_Z^2+12*_Z+1)^2*(-x^2+1)^(1/3)*x-576*RootOf(144*_Z^2+12*_Z+1)^2*x+36*RootOf(144*_Z^2+12*_Z+1)*(-x^2+1)^(1/3)*x-6*RootOf(144*_Z^2+12*_Z+1)*x^2-36*RootOf(144*_Z^2+12*_Z+1)*(-x^2+1)^(1/3)-24*RootOf(144*_Z^2+12*_Z+1)*x+3*(-x^2+1)^(2/3)+(-x^2+1)^(1/3)*x-18*RootOf(144*_Z^2+12*_Z+1)-3*(-x^2+1)^(1/3))/(x-3)/(x+3))-ln((288*RootOf(144*_Z^2+12*_Z+1)^2*(-x^2+1)^(1/3)*x-576*RootOf(144*_Z^2+12*_Z+1)^2*x+36*RootOf(144*_Z^2+12*_Z+1)*(-x^2+1)^(1/3)*x-6*RootOf(144*_Z^2+12*_Z+1)*x^2-36*RootOf(144*_Z^2+12*_Z+1)*(-x^2+1)^(1/3)-24*RootOf(144*_Z^2+12*_Z+1)*x+3*(-x^2+1)^(2/3)+(-x^2+1)^(1/3)*x-18*RootOf(144*_Z^2+12*_Z+1)-3*(-x^2+1)^(1/3))/(x-3)/(x+3))*RootOf(144*_Z^2+12*_Z+1)+RootOf(144*_Z^2+12*_Z+1)*ln((576*RootOf(144*_Z^2+12*_Z+1)^2*(-x^2+1)^(1/3)*x-1152*RootOf(144*_Z^2+12*_Z+1)^2*x+24*RootOf(144*_Z^2+12*_Z+1)*(-x^2+1)^(1/3)*x+12*RootOf(144*_Z^2+12*_Z+1)*x^2+72*RootOf(144*_Z^2+12*_Z+1)*(-x^2+1)^(1/3)-144*RootOf(144*_Z^2+12*_Z+1)*x+6*(-x^2+1)^(2/3)+x^2+36*RootOf(144*_Z^2+12*_Z+1)-4*x+3)/(x-3)/(x+3))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(x^2 - 9)(-x^2 + 1)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/3)/(-x^2+9),x, algorithm="maxima")

[Out] -integrate(1/((x^2 - 9)*(-x^2 + 1)^(1/3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(1 - x^2)^{\frac{1}{3}}(x^2 - 9)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((1 - x^2)^(1/3)*(x^2 - 9)),x)

[Out] -int(1/((1 - x^2)^(1/3)*(x^2 - 9)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{x^2 \sqrt[3]{1-x^2} - 9 \sqrt[3]{1-x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/3)/(-x**2+9),x)

[Out] -Integral(1/(x**2*(1 - x**2)**(1/3) - 9*(1 - x**2)**(1/3)), x)

$$3.163 \quad \int \frac{\sqrt{-1+c^2x^2}}{(d-c^2dx^2)^{5/2}} dx$$

Optimal. Leaf size=79

$$\frac{\sqrt{c^2x^2-1} \tanh^{-1}(cx)}{2cd^2\sqrt{d-c^2dx^2}} + \frac{x\sqrt{c^2x^2-1}}{2d(d-c^2dx^2)^{3/2}}$$

[Out] 1/2*x*(c^2*x^2-1)^(1/2)/d/(-c^2*d*x^2+d)^(3/2)+1/2*arctanh(c*x)*(c^2*x^2-1)^(1/2)/c/d^2/(-c^2*d*x^2+d)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 91, normalized size of antiderivative = 1.15, number of steps used = 3, number of rules used = 3, integrand size = 29, $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$, Rules used = {23, 199, 208}

$$\frac{x\sqrt{c^2x^2-1}}{2d^2(1-c^2x^2)\sqrt{d-c^2dx^2}} + \frac{\sqrt{c^2x^2-1} \tanh^{-1}(cx)}{2cd^2\sqrt{d-c^2dx^2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-1 + c^2*x^2]/(d - c^2*d*x^2)^(5/2), x]

[Out] (x*Sqrt[-1 + c^2*x^2])/((2*d^2*(1 - c^2*x^2)*Sqrt[d - c^2*d*x^2]) + (Sqrt[-1 + c^2*x^2]*ArcTanh[c*x]))/(2*c*d^2*Sqrt[d - c^2*d*x^2])

Rule 23

Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :> Dist[(a + b*v)^m/(c + d*v)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[b*c - a*d, 0] && !(IntegerQ[m] || IntegerQ[n] || GtQ[b/d, 0])

Rule 199

Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])

Rule 208

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-1+c^2x^2}}{(d-c^2dx^2)^{5/2}} dx &= \frac{\sqrt{-1+c^2x^2} \int \frac{1}{(d-c^2dx^2)^2} dx}{\sqrt{d-c^2dx^2}} \\ &= \frac{x\sqrt{-1+c^2x^2}}{2d^2(1-c^2x^2)\sqrt{d-c^2dx^2}} + \frac{\sqrt{-1+c^2x^2} \int \frac{1}{d-c^2dx^2} dx}{2d\sqrt{d-c^2dx^2}} \\ &= \frac{x\sqrt{-1+c^2x^2}}{2d^2(1-c^2x^2)\sqrt{d-c^2dx^2}} + \frac{\sqrt{-1+c^2x^2} \tanh^{-1}(cx)}{2cd^2\sqrt{d-c^2dx^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 57, normalized size = 0.72

$$\frac{(c^2x^2 - 1) \tanh^{-1}(cx) - cx}{2cd^2\sqrt{c^2x^2 - 1}\sqrt{d - c^2dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-1 + c^2*x^2]/(d - c^2*d*x^2)^(5/2), x]

[Out] $(-c*x) + (-1 + c^2*x^2)*\text{ArcTanh}[c*x]/(2*c*d^2*\text{Sqrt}[-1 + c^2*x^2]*\text{Sqrt}[d - c^2*d*x^2])$

fricas [A] time = 1.00, size = 315, normalized size = 3.99

$$\frac{4\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}cx - (c^4x^4 - 2c^2x^2 + 1)\sqrt{-d} \log\left(-\frac{c^6dx^6 + 5c^4dx^4 - 5c^2dx^2 - 4(c^3x^3 + cx)\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}\sqrt{-d - d}}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1}\right)}{8(c^5d^3x^4 - 2c^3d^3x^2 + cd^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c^2*x^2-1)^(1/2)/(-c^2*d*x^2+d)^(5/2), x, algorithm="fricas")

[Out] $[1/8*(4*\text{sqrt}(-c^2*d*x^2 + d)*\text{sqrt}(c^2*x^2 - 1)*c*x - (c^4*x^4 - 2*c^2*x^2 + 1)*\text{sqrt}(-d)*\log(-(c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 - 4*(c^3*x^3 + c*x)*\text{sqrt}(-c^2*d*x^2 + d)*\text{sqrt}(c^2*x^2 - 1)*\text{sqrt}(-d) - d)/(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)))/(c^5*d^3*x^4 - 2*c^3*d^3*x^2 + c*d^3), 1/4*(2*\text{sqrt}(-c^2*d*x^2 + d)*\text{sqrt}(c^2*x^2 - 1)*c*x - (c^4*x^4 - 2*c^2*x^2 + 1)*\text{sqrt}(d)*\text{arctan}(2*\text{sqrt}(-c^2*d*x^2 + d)*\text{sqrt}(c^2*x^2 - 1)*c*\text{sqrt}(d)*x/(c^4*d*x^4 - d)))/(c^5*d^3*x^4 - 2*c^3*d^3*x^2 + c*d^3)]$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c^2x^2 - 1}}{(-c^2dx^2 + d)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c^2*x^2-1)^(1/2)/(-c^2*d*x^2+d)^(5/2), x, algorithm="giac")

[Out] integrate(sqrt(c^2*x^2 - 1)/(-c^2*d*x^2 + d)^(5/2), x)

maple [A] time = 0.03, size = 94, normalized size = 1.19

$$\frac{\sqrt{-(c^2x^2 - 1)}d(-c^2x^2 \ln(cx - 1) + c^2x^2 \ln(cx + 1) - 2cx + \ln(cx - 1) - \ln(cx + 1))}{4\sqrt{c^2x^2 - 1}(cx + 1)(cx - 1)cd^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c^2*x^2-1)^(1/2)/(-c^2*d*x^2+d)^(5/2), x)

[Out] $-1/4/(c^2*x^2-1)^(1/2)*(-(c^2*x^2-1)*d)^(1/2)*(ln(c*x+1)*x^2*c^2-\ln(c*x-1)*x^2*c^2-2*c*x-\ln(c*x+1)+\ln(c*x-1))/d^3/c/(c*x+1)/(c*x-1)$

maxima [A] time = 1.49, size = 70, normalized size = 0.89

$$-\frac{x}{2(c^2\sqrt{-d}d^2x^2 - \sqrt{-d}d^2)} - \frac{\sqrt{-d} \log(cx + 1)}{4cd^3} + \frac{\sqrt{-d} \log(cx - 1)}{4cd^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c^2*x^2-1)^(1/2)/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")

[Out] -1/2*x/(c^2*sqrt(-d)*d^2*x^2 - sqrt(-d)*d^2) - 1/4*sqrt(-d)*log(c*x + 1)/(c*d^3) + 1/4*sqrt(-d)*log(c*x - 1)/(c*d^3)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{c^2 x^2 - 1}}{(d - c^2 d x^2)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c^2*x^2 - 1)^(1/2)/(d - c^2*d*x^2)^(5/2),x)

[Out] int((c^2*x^2 - 1)^(1/2)/(d - c^2*d*x^2)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{(cx - 1)(cx + 1)}}{(-d(cx - 1)(cx + 1))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c**2*x**2-1)**(1/2)/(-c**2*d*x**2+d)**(5/2),x)

[Out] Integral(sqrt((c*x - 1)*(c*x + 1))/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)

$$3.164 \quad \int \frac{1}{(-1+c^2x^2)^{3/2} \sqrt{d-c^2dx^2}} dx$$

Optimal. Leaf size=74

$$\frac{dx\sqrt{c^2x^2-1}}{2(d-c^2dx^2)^{3/2}} + \frac{\sqrt{c^2x^2-1} \tanh^{-1}(cx)}{2c\sqrt{d-c^2dx^2}}$$

[Out] $1/2*d*x*(c^2*x^2-1)^{(1/2)/(-c^2*d*x^2+d)^{(3/2)+1/2*\arctanh(c*x)*(c^2*x^2-1)^{(1/2)/c/(-c^2*d*x^2+d)^{(1/2)}}$

Rubi [A] time = 0.02, antiderivative size = 91, normalized size of antiderivative = 1.23, number of steps used = 3, number of rules used = 3, integrand size = 29, $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$, Rules used = {23, 199, 208}

$$\frac{x(d-c^2dx^2)^{3/2}}{2d^2(1-c^2x^2)(c^2x^2-1)^{3/2}} + \frac{(d-c^2dx^2)^{3/2} \tanh^{-1}(cx)}{2cd^2(c^2x^2-1)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((-1 + c^2*x^2)^(3/2)*Sqrt[d - c^2*d*x^2]), x]

[Out] $(x*(d - c^2*d*x^2)^{(3/2)})/(2*d^2*(1 - c^2*x^2)*(-1 + c^2*x^2)^{(3/2)}) + ((d - c^2*d*x^2)^{(3/2)*ArcTanh[c*x]}/(2*c*d^2*(-1 + c^2*x^2)^{(3/2)})$

Rule 23

Int[(u_.)*((a_) + (b_.)*(v_))^(m_)*((c_) + (d_.)*(v_))^(n_), x_Symbol] :> Dist[(a + b*v)^m/(c + d*v)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[b*c - a*d, 0] && !(IntegerQ[m] || IntegerQ[n] || GtQ[b/d, 0])

Rule 199

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rubi steps

$$\begin{aligned} \int \frac{1}{(-1+c^2x^2)^{3/2} \sqrt{d-c^2dx^2}} dx &= \frac{(d-c^2dx^2)^{3/2} \int \frac{1}{(d-c^2dx^2)^2} dx}{(-1+c^2x^2)^{3/2}} \\ &= \frac{x(d-c^2dx^2)^{3/2}}{2d^2(1-c^2x^2)(-1+c^2x^2)^{3/2}} + \frac{(d-c^2dx^2)^{3/2} \int \frac{1}{d-c^2dx^2} dx}{2d(-1+c^2x^2)^{3/2}} \\ &= \frac{x(d-c^2dx^2)^{3/2}}{2d^2(1-c^2x^2)(-1+c^2x^2)^{3/2}} + \frac{(d-c^2dx^2)^{3/2} \tanh^{-1}(cx)}{2cd^2(-1+c^2x^2)^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 54, normalized size = 0.73

$$\frac{(c^2x^2 - 1) \tanh^{-1}(cx) - cx}{2c\sqrt{c^2x^2 - 1} \sqrt{d - c^2dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((-1 + c^2*x^2)^(3/2)*Sqrt[d - c^2*d*x^2]), x]

[Out] $(-(c*x) + (-1 + c^2*x^2)*ArcTanh[c*x]) / (2*c*Sqrt[-1 + c^2*x^2]*Sqrt[d - c^2*d*x^2])$

fricas [A] time = 0.66, size = 303, normalized size = 4.09

$$\frac{4\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}cx - (c^4x^4 - 2c^2x^2 + 1)\sqrt{-d} \log\left(-\frac{c^6dx^6 + 5c^4dx^4 - 5c^2dx^2 - 4(c^3x^3 + cx)\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}\sqrt{-d}}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1}\right)}{8(c^5dx^4 - 2c^3dx^2 + cd)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c^2*x^2-1)^(3/2)/(-c^2*d*x^2+d)^(1/2), x, algorithm="fricas")

[Out] $[1/8*(4*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*c*x - (c^4*x^4 - 2*c^2*x^2 + 1)*\sqrt{-d}*\log(-\frac{c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 - 4*(c^3*x^3 + c*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*\sqrt{-d} - d}{(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)})) / (c^5*d*x^4 - 2*c^3*d*x^2 + c*d), 1/4*(2*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*c*x - (c^4*x^4 - 2*c^2*x^2 + 1)*\sqrt{d}*\arctan(2*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*c*\sqrt{d}*x / (c^4*d*x^4 - d))) / (c^5*d*x^4 - 2*c^3*d*x^2 + c*d)]$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-c^2dx^2 + d}(c^2x^2 - 1)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c^2*x^2-1)^(3/2)/(-c^2*d*x^2+d)^(1/2), x, algorithm="giac")

[Out] integrate(1/(sqrt(-c^2*d*x^2 + d)*(c^2*x^2 - 1)^(3/2)), x)

maple [A] time = 0.02, size = 94, normalized size = 1.27

$$\frac{\sqrt{-(c^2x^2 - 1)d} (-c^2x^2 \ln(cx - 1) + c^2x^2 \ln(cx + 1) - 2cx + \ln(cx - 1) - \ln(cx + 1))}{4\sqrt{c^2x^2 - 1} (cx + 1)(cx - 1)cd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c^2*x^2-1)^(3/2)/(-c^2*d*x^2+d)^(1/2), x)

[Out] $-1/4/(c^2*x^2-1)^(1/2)*(-(c^2*x^2-1)*d)^(1/2)*(-c^2*x^2*\ln(c*x-1)+c^2*x^2*\ln(c*x+1)-2*c*x+\ln(c*x-1)-\ln(c*x+1))/d/c/(c*x+1)/(c*x-1)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-c^2dx^2 + d}(c^2x^2 - 1)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c^2*x^2-1)^(3/2)/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-c^2*d*x^2 + d)*(c^2*x^2 - 1)^(3/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{d - c^2 d x^2} (c^2 x^2 - 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((d - c^2*d*x^2)^(1/2)*(c^2*x^2 - 1)^(3/2)), x)

[Out] int(1/((d - c^2*d*x^2)^(1/2)*(c^2*x^2 - 1)^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{((cx - 1)(cx + 1))^{\frac{3}{2}} \sqrt{-d}(cx - 1)(cx + 1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c**2*x**2-1)**(3/2)/(-c**2*d*x**2+d)**(1/2),x)

[Out] Integral(1/(((c*x - 1)*(c*x + 1))**(3/2)*sqrt(-d*(c*x - 1)*(c*x + 1))), x)

$$3.165 \quad \int \frac{1}{\sqrt{-1+c^2x^2} (d-c^2dx^2)^{3/2}} dx$$

Optimal. Leaf size=76

$$-\frac{x\sqrt{c^2x^2-1}}{2(d-c^2dx^2)^{3/2}} - \frac{\sqrt{c^2x^2-1} \tanh^{-1}(cx)}{2cd\sqrt{d-c^2dx^2}}$$

[Out] $-1/2*x*(c^2*x^2-1)^{(1/2)/(-c^2*d*x^2+d)^{(3/2)}-1/2*\operatorname{arctanh}(c*x)*(c^2*x^2-1)^{(1/2)}/c/d/(-c^2*d*x^2+d)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 91, normalized size of antiderivative = 1.20, number of steps used = 3, number of rules used = 3, integrand size = 29, $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$, Rules used = {23, 199, 208}

$$\frac{x\sqrt{d-c^2dx^2}}{2d^2(1-c^2x^2)\sqrt{c^2x^2-1}} + \frac{\sqrt{d-c^2dx^2} \tanh^{-1}(cx)}{2cd^2\sqrt{c^2x^2-1}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(\operatorname{Sqrt}[-1+c^2*x^2]*(d-c^2*d*x^2)^{(3/2)}),x]$

[Out] $(x*\operatorname{Sqrt}[d-c^2*d*x^2])/(2*d^2*(1-c^2*x^2)*\operatorname{Sqrt}[-1+c^2*x^2]) + (\operatorname{Sqrt}[d-c^2*d*x^2]*\operatorname{ArcTanh}[c*x])/(2*c*d^2*\operatorname{Sqrt}[-1+c^2*x^2])$

Rule 23

$\operatorname{Int}[(a_+)*(b_+*(v_+))^{(m_+)}*((c_+)+(d_+)*(v_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Dist}[(a+b*v)^m/(c+d*v)^m, \operatorname{Int}[u*(c+d*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}[\{a, b, c, d, m, n\}, x] \ \&\& \ \operatorname{EqQ}[b*c-a*d, 0] \ \&\& \ !(\operatorname{IntegerQ}[m] \ || \ \operatorname{IntegerQ}[n] \ || \ \operatorname{GtQ}[b/d, 0])$

Rule 199

$\operatorname{Int}[(a_+)+(b_+*(x_+)^{(n_+))^{(p_+)}, x_Symbol] \rightarrow -\operatorname{Simp}[(x*(a+b*x^n)^{(p+1)})/(a*n*(p+1)), x] + \operatorname{Dist}[(n*(p+1)+1)/(a*n*(p+1)), \operatorname{Int}[(a+b*x^n)^{(p+1)}, x], x] /;$ $\operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{LtQ}[p, -1] \ \&\& \ (\operatorname{IntegerQ}[2*p] \ || \ (n == 2 \ \&\& \ \operatorname{IntegerQ}[4*p]) \ || \ (n == 2 \ \&\& \ \operatorname{IntegerQ}[3*p]) \ || \ \operatorname{Denominator}[p+1/n] < \operatorname{Denominator}[p])$

Rule 208

$\operatorname{Int}[(a_+)+(b_+*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-(a/b), 2]*\operatorname{ArcTanh}[x/\operatorname{Rt}[-(a/b), 2]])/a, x] /;$ $\operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{NegQ}[a/b]$

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1+c^2x^2} (d-c^2dx^2)^{3/2}} dx &= \frac{\sqrt{d-c^2dx^2} \int \frac{1}{(d-c^2dx^2)^2} dx}{\sqrt{-1+c^2x^2}} \\ &= \frac{x\sqrt{d-c^2dx^2}}{2d^2(1-c^2x^2)\sqrt{-1+c^2x^2}} + \frac{\sqrt{d-c^2dx^2} \int \frac{1}{d-c^2dx^2} dx}{2d\sqrt{-1+c^2x^2}} \\ &= \frac{x\sqrt{d-c^2dx^2}}{2d^2(1-c^2x^2)\sqrt{-1+c^2x^2}} + \frac{\sqrt{d-c^2dx^2} \tanh^{-1}(cx)}{2cd^2\sqrt{-1+c^2x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 57, normalized size = 0.75

$$\frac{(1 - c^2x^2) \tanh^{-1}(cx) + cx}{2cd\sqrt{c^2x^2 - 1}\sqrt{d - c^2dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 + c^2*x^2]*(d - c^2*d*x^2)^(3/2)), x]

[Out] (c*x + (1 - c^2*x^2)*ArcTanh[c*x])/(2*c*d*Sqrt[-1 + c^2*x^2]*Sqrt[d - c^2*d*x^2])

fricas [A] time = 1.03, size = 314, normalized size = 4.13

$$\left[\frac{4\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}cx + (c^4x^4 - 2c^2x^2 + 1)\sqrt{-d} \log\left(-\frac{c^6dx^6 + 5c^4dx^4 - 5c^2dx^2 + 4(c^3x^3 + cx)\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}\sqrt{-d}}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1}\right)}{8(c^5d^2x^4 - 2c^3d^2x^2 + cd^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c^2*x^2-1)^(1/2)/(-c^2*d*x^2+d)^(3/2), x, algorithm="fricas")

[Out] [-1/8*(4*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*c*x + (c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-d)*log(-(c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 + 4*(c^3*x^3 + c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*sqrt(-d) - d)/(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)))/(c^5*d^2*x^4 - 2*c^3*d^2*x^2 + c*d^2), -1/4*(2*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*c*x - (c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(d)*arctan(2*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*c*sqrt(d)*x/(c^4*d*x^4 - d)))/(c^5*d^2*x^4 - 2*c^3*d^2*x^2 + c*d^2)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-c^2dx^2 + d)^{\frac{3}{2}}\sqrt{c^2x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c^2*x^2-1)^(1/2)/(-c^2*d*x^2+d)^(3/2), x, algorithm="giac")

[Out] integrate(1/((-c^2*d*x^2 + d)^(3/2)*sqrt(c^2*x^2 - 1)), x)

maple [A] time = 0.02, size = 94, normalized size = 1.24

$$\frac{\sqrt{-(c^2x^2 - 1)d} (-c^2x^2 \ln(cx - 1) + c^2x^2 \ln(cx + 1) - 2cx + \ln(cx - 1) - \ln(cx + 1))}{4\sqrt{c^2x^2 - 1} (cx + 1)(cx - 1)cd^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c^2*x^2-1)^(1/2)/(-c^2*d*x^2+d)^(3/2), x)

[Out] 1/4/(c^2*x^2-1)^(1/2)*(-(c^2*x^2-1)*d)^(1/2)*(-c^2*x^2*ln(c*x-1)+c^2*x^2*ln(c*x+1)-2*c*x*ln(c*x-1)-ln(c*x+1))/d^2/c/(c*x+1)/(c*x-1)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-c^2dx^2 + d)^{\frac{3}{2}}\sqrt{c^2x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c^2*x^2-1)^(1/2)/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")

[Out] integrate(1/((-c^2*d*x^2 + d)^(3/2)*sqrt(c^2*x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(d - c^2 d x^2)^{3/2} \sqrt{c^2 x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((d - c^2*d*x^2)^(3/2)*(c^2*x^2 - 1)^(1/2)),x)

[Out] int(1/((d - c^2*d*x^2)^(3/2)*(c^2*x^2 - 1)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{(cx - 1)(cx + 1)} (-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c**2*x**2-1)**(1/2)/(-c**2*d*x**2+d)**(3/2),x)

[Out] Integral(1/(sqrt((c*x - 1)*(c*x + 1))*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)

3.166 $\int (a + bx^2)^{3/2} \sqrt{c + dx^2} dx$

Optimal. Leaf size=328

$$\frac{c^{3/2}\sqrt{a+bx^2}(bc-9ad)\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right) + \sqrt{c}\sqrt{a+bx^2}(-3a^2d^2-7abcd+2b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\right)}{15d^{3/2}\sqrt{c+dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}} + \frac{15bd^{3/2}\sqrt{c+dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}}{15d^{3/2}\sqrt{c+dx^2}}$$

[Out] $\frac{1}{5}bxx(d^2x^2+c)^{3/2}(bx^2+a)^{1/2}/d + \frac{1}{15}(7ac-2bc^2/d+3a^2d/b)xx(bx^2+a)^{1/2}/(d^2x^2+c)^{1/2} - \frac{1}{15}c^{3/2}(-9ad+bc)(1/(1+d^2x^2/c))^{1/2}(1+d^2x^2/c)^{1/2}\operatorname{EllipticF}(xd^{1/2}/c^{1/2}/(1+d^2x^2/c)^{1/2}, (1-bc/a/d)^{1/2})(bx^2+a)^{1/2}/d^{3/2}/(c(bx^2+a)/a/(d^2x^2+c))^{1/2}/(d^2x^2+c)^{1/2} + \frac{1}{15}(-3a^2d^2-7abc^2d+2b^2c^2)(1/(1+d^2x^2/c))^{1/2}(1+d^2x^2/c)^{1/2}\operatorname{EllipticE}(xd^{1/2}/c^{1/2}/(1+d^2x^2/c)^{1/2}, (1-bc/a/d)^{1/2})c^{1/2}(bx^2+a)^{1/2}/b/d^{3/2}/(c(bx^2+a)/a/(d^2x^2+c))^{1/2}/(d^2x^2+c)^{1/2} - \frac{2}{15}(-3ad+bc)xx(bx^2+a)^{1/2}(d^2x^2+c)^{1/2}/d$

Rubi [A] time = 0.32, antiderivative size = 328, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {416, 528, 531, 418, 492, 411}

$$\frac{\sqrt{c}\sqrt{a+bx^2}(-3a^2d^2-7abcd+2b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right) + x\sqrt{a+bx^2}\left(\frac{3a^2d}{b}+7ac-\frac{2bc^2}{d}\right) + c^{3/2}\sqrt{a+bx^2}(bc-9ad)\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right)}{15bd^{3/2}\sqrt{c+dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}} + \frac{15bd^{3/2}\sqrt{c+dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}}{15\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a + bx^2)^{3/2}\sqrt{c + dx^2}, x]$

[Out] $((7ac - (2bc^2)/d + (3a^2d)/b)xx\sqrt{a + bx^2})/(15\sqrt{c + dx^2}) - (2(bc - 3ad)xx\sqrt{a + bx^2}\sqrt{c + dx^2})/(15d) + (bxx\sqrt{a + bx^2}(c + dx^2)^{3/2})/(5d) + (\sqrt{c}(2b^2c^2 - 7abc^2d - 3a^2d^2)\sqrt{a + bx^2}\operatorname{EllipticE}[\operatorname{ArcTan}[(\sqrt{d}x)/\sqrt{c}], 1 - (bc)/(ad)])/(15bd^{3/2}\sqrt{c + dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}) - (c^{3/2}(bc - 9ad)\sqrt{a + bx^2}\operatorname{EllipticF}[\operatorname{ArcTan}[(\sqrt{d}x)/\sqrt{c}], 1 - (bc)/(ad)])/(15d^{3/2}\sqrt{c + dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}) + \sqrt{c}\sqrt{a + bx^2}(-3a^2d^2 - 7abcd + 2b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right) + x\sqrt{a + bx^2}\left(\frac{3a^2d}{b} + 7ac - \frac{2bc^2}{d}\right)$

Rule 411

$\operatorname{Int}[\sqrt{(a_1 + (b_1)(x_1)^2)/((c_1) + (d_1)(x_1)^2)^{3/2}}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\sqrt{a + bx^2}\operatorname{EllipticE}[\operatorname{ArcTan}[\operatorname{Rt}[d/c, 2]x], 1 - (bc)/(ad)])/(c\operatorname{Rt}[d/c, 2]\sqrt{c + dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}), x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \operatorname{PosQ}[b/a] \ \&\& \operatorname{PosQ}[d/c]$

Rule 416

$\operatorname{Int}[(a_1 + (b_1)(x_1)^{n_1})^{p_1}((c_1) + (d_1)(x_1)^{n_1})^{q_1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(d^p(a + bx^n)^{p+1}(c + dx^n)^{q-1})/(b(n(p+q)+1)), x] + \operatorname{Dist}[1/(b(n(p+q)+1)), \operatorname{Int}[(a + bx^n)^p(c + dx^n)^{q-2}\operatorname{Simp}[c(b^p(n(p+q)+1) - ad) + d(b^p(n(p+2q-1)+1) - a^p(n(q-1)+1))x^n, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, n, p\}, x \ \&\& \operatorname{NeQ}[b^p - ad, 0] \ \&\& \operatorname{GtQ}[q, 1] \ \&\& \operatorname{NeQ}[n(p+q)+1, 0] \ \&\& \operatorname{!IGtQ}[p, 1] \ \&\& \operatorname{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 418

```
Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)]/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
:= Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rule 528

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
f_)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rule 531

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
f_)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x],
x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c,
d, e, f, n, p, q}, x]
```

Rubi steps

$$\begin{aligned} \int (a + bx^2)^{3/2} \sqrt{c + dx^2} dx &= \frac{bx\sqrt{a + bx^2} (c + dx^2)^{3/2}}{5d} + \frac{\int \frac{\sqrt{c+dx^2}(-a(bc-5ad)-2b(bc-3ad)x^2)}{\sqrt{a+bx^2}} dx}{5d} \\ &= -\frac{2(bc - 3ad)x\sqrt{a + bx^2} \sqrt{c + dx^2}}{15d} + \frac{bx\sqrt{a + bx^2} (c + dx^2)^{3/2}}{5d} + \frac{\int \frac{-abc(bc-9ad)-b^2x^2}{\sqrt{a+bx^2}} dx}{5d} \\ &= -\frac{2(bc - 3ad)x\sqrt{a + bx^2} \sqrt{c + dx^2}}{15d} + \frac{bx\sqrt{a + bx^2} (c + dx^2)^{3/2}}{5d} - \frac{(ac(bc - 9ad))}{5d} \\ &= \frac{\left(7ac - \frac{2bc^2}{d} + \frac{3a^2d}{b}\right)x\sqrt{a + bx^2}}{15\sqrt{c + dx^2}} - \frac{2(bc - 3ad)x\sqrt{a + bx^2} \sqrt{c + dx^2}}{15d} + \frac{bx\sqrt{a + bx^2} (c + dx^2)^{3/2}}{5d} \\ &= \frac{\left(7ac - \frac{2bc^2}{d} + \frac{3a^2d}{b}\right)x\sqrt{a + bx^2}}{15\sqrt{c + dx^2}} - \frac{2(bc - 3ad)x\sqrt{a + bx^2} \sqrt{c + dx^2}}{15d} + \frac{bx\sqrt{a + bx^2} (c + dx^2)^{3/2}}{5d} \end{aligned}$$

Mathematica [C] time = 0.46, size = 243, normalized size = 0.74

$$\frac{-2ic\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}(3a^2d^2 - 4abcd + b^2c^2)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) - ic\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}(3a^2d^2 - 4abcd + b^2c^2)}{15d^2\sqrt{\frac{b}{a}}\sqrt{a + bx^2}\sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)*Sqrt[c + d*x^2], x]

[Out] (Sqrt[b/a]*d*x*(a + b*x^2)*(c + d*x^2)*(6*a*d + b*(c + 3*d*x^2)) - I*c*(-2*b^2*c^2 + 7*a*b*c*d + 3*a^2*d^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - (2*I)*c*(b^2*c^2 - 4*a*b*c*d + 3*a^2*d^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(15*Sqrt[b/a]*d^2*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.85, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^{\frac{3}{2}} \sqrt{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral((b*x^2 + a)^(3/2)*sqrt(d*x^2 + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^{\frac{3}{2}} \sqrt{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^(1/2), x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(3/2)*sqrt(d*x^2 + c), x)

maple [A] time = 0.04, size = 543, normalized size = 1.66

$$\sqrt{bx^2 + a} \sqrt{dx^2 + c} \left(3\sqrt{-\frac{b}{a}} b^2 d^3 x^7 + 9\sqrt{-\frac{b}{a}} ab d^3 x^5 + 4\sqrt{-\frac{b}{a}} b^2 c d^2 x^5 + 6\sqrt{-\frac{b}{a}} a^2 d^3 x^3 + 10\sqrt{-\frac{b}{a}} abc d^2 x^3 + \sqrt{\dots} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)*(d*x^2+c)^(1/2), x)

[Out] 1/15*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(3*(-1/a*b)^(1/2)*x^7*b^2*d^3+9*(-1/a*b)^(1/2)*x^5*a*b*d^3+4*(-1/a*b)^(1/2)*x^5*b^2*c*d^2+6*(-1/a*b)^(1/2)*x^3*a^2*d^3+10*(-1/a*b)^(1/2)*x^3*a*b*c*d^2+(-1/a*b)^(1/2)*x^3*b^2*c^2*d+6*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(-1/a*b)^(1/2), (a*d/b/c)^(1/2))*a^2*c*d^2-8*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(-1/a*b)^(1/2), (a*d/b/c)^(1/2))*a*b*c^2*d+2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(x*(-1/a*b)^(1/2), (a*d/b/c)^(1/2))*b^2*c^3+3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(-1/a*b)^(1/2), (a*d/b/c)^(1/2))*a^2*c*d^2+7*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(-1/a*b)^(1/2), (a*d/b/c)^(1/2))*a*b*c^2*d-2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE(x*(-1/a*b)^(1/2), (a*d/b/c)^(1/2))*b^2*c^3+6*(-1/a*b)^(1/2)*x*a^2*c*d^2+(-1/a*b)^(1/2)*x*a*b*c^2*d)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/d^2/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^{\frac{3}{2}} \sqrt{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/2)*sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (bx^2 + a)^{3/2} \sqrt{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)*(c + d*x^2)^(1/2),x)

[Out] int((a + b*x^2)^(3/2)*(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)^{3/2} \sqrt{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)*(d*x**2+c)**(1/2),x)

[Out] Integral((a + b*x**2)**(3/2)*sqrt(c + d*x**2), x)

3.167 $\int \sqrt{a + bx^2} \sqrt{c + dx^2} dx$

Optimal. Leaf size=249

$$\frac{2c^{3/2}\sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + \frac{1}{3}x\sqrt{a+bx^2}\sqrt{c+dx^2} + \frac{x\sqrt{a+bx^2}(ad+bc)}{3b\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{a+bx^2}(ad+bc)}{3b\sqrt{d}\sqrt{c+dx^2}}}{3\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $\frac{1}{3}*(a*d+b*c)*x*(b*x^2+a)^{(1/2)}/b/(d*x^2+c)^{(1/2)}+2/3*c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-1/3*(a*d+b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/b/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/3*x*(b*x^2+a)^{(1/2)}*(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.18, antiderivative size = 249, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {417, 531, 418, 492, 411}

$$\frac{2c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) + \frac{1}{3}x\sqrt{a+bx^2}\sqrt{c+dx^2} + \frac{x\sqrt{a+bx^2}(ad+bc)}{3b\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{a+bx^2}(ad+bc)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{3b\sqrt{d}\sqrt{c+dx^2}}}{3\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[a + b*x^2]*Sqrt[c + d*x^2], x]`

[Out] $((b*c + a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(3*b*\operatorname{Sqrt}[c + d*x^2]) + (x*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2])/3 - (\operatorname{Sqrt}[c]*(b*c + a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(3*b*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (2*c^{(3/2)}*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(3*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

`Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]`

Rule 417

`Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(x*(a + b*x^n)^p*(c + d*x^n)^q)/(n*(p + q) + 1), x] + Dist[n/(n*(p + q) + 1), Int[(a + b*x^n)^(p - 1)*(c + d*x^n)^(q - 1)*Simp[a*c*(p + q) + (q*(b*c - a*d) + a*d*(p + q))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 0] && GtQ[p, 0] && IntBinomialQ[a, b, c, d, n, p, q, x]`

Rule 418

`Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]`

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
  := Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
  + b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
  a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rule 531

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (
  f_.)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x],
  x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c,
  d, e, f, n, p, q}, x]
```

Rubi steps

$$\begin{aligned} \int \sqrt{a+bx^2} \sqrt{c+dx^2} dx &= \frac{1}{3}x\sqrt{a+bx^2} \sqrt{c+dx^2} + \frac{2}{3} \int \frac{ac + \frac{1}{2}(bc+ad)x^2}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx \\ &= \frac{1}{3}x\sqrt{a+bx^2} \sqrt{c+dx^2} + \frac{1}{3}(2ac) \int \frac{1}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx + \frac{1}{3}(bc+ad) \int \frac{1}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx \\ &= \frac{(bc+ad)x\sqrt{a+bx^2}}{3b\sqrt{c+dx^2}} + \frac{1}{3}x\sqrt{a+bx^2} \sqrt{c+dx^2} + \frac{2c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{c(a+bx^2)}{a(c+dx^2)}\right)}{3\sqrt{d} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} \sqrt{c+dx^2}} \\ &= \frac{(bc+ad)x\sqrt{a+bx^2}}{3b\sqrt{c+dx^2}} + \frac{1}{3}x\sqrt{a+bx^2} \sqrt{c+dx^2} - \frac{\sqrt{c}(bc+ad)\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{c(a+bx^2)}{a(c+dx^2)}\right)}{3b\sqrt{d} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} \sqrt{c+dx^2}} \end{aligned}$$

Mathematica [C] time = 0.25, size = 198, normalized size = 0.80

$$\frac{-ic\sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{dx^2}{c} + 1} (ad - bc) \operatorname{EllipticF}\left(i \sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) + dx\sqrt{\frac{b}{a}} (a + bx^2) (c + dx^2) - ic\sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{dx^2}{c} + 1}}{3d\sqrt{\frac{b}{a}} \sqrt{a + bx^2} \sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[a + b*x^2]*Sqrt[c + d*x^2], x]
```

```
[Out] (Sqrt[b/a]*d*x*(a + b*x^2)*(c + d*x^2) - I*c*(b*c + a*d)*Sqrt[1 + (b*x^2)/a]
*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*c*
(-(b*c) + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[
Sqrt[b/a]*x], (a*d)/(b*c))]/(3*Sqrt[b/a]*d*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])
```

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\sqrt{bx^2 + a} \sqrt{dx^2 + c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{bx^2 + a} \sqrt{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 + a)*sqrt(d*x^2 + c), x)

maple [A] time = 0.02, size = 328, normalized size = 1.32

$$\sqrt{bx^2 + a} \sqrt{dx^2 + c} \left(\sqrt{\frac{-b}{a}} b d^2 x^5 + \sqrt{\frac{-b}{a}} a d^2 x^3 + \sqrt{\frac{-b}{a}} bcd x^3 + \sqrt{\frac{-b}{a}} acdx + \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} acd \operatorname{EllipticE} \left(\sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2),x)

[Out] $\frac{1}{3}(bx^2+a)^{1/2}(dx^2+c)^{1/2}((-1/a*b)^{1/2}*x^5*b*d^2+(-1/a*b)^{1/2}) * x^3*a*d^2+(-1/a*b)^{1/2}*x^3*b*c*d+a*c*((b*x^2+a)/a)^{1/2}*((d*x^2+c)/c)^{1/2} * \operatorname{EllipticF}((-1/a*b)^{1/2}*x, (a/b/c*d)^{1/2}) * d - ((b*x^2+a)/a)^{1/2} * ((d*x^2+c)/c)^{1/2} * \operatorname{EllipticF}((-1/a*b)^{1/2}*x, (a/b/c*d)^{1/2}) * b*c^2 + ((b*x^2+a)/a)^{1/2} * ((d*x^2+c)/c)^{1/2} * \operatorname{EllipticE}((-1/a*b)^{1/2}*x, (a/b/c*d)^{1/2}) * a*c*d + ((b*x^2+a)/a)^{1/2} * ((d*x^2+c)/c)^{1/2} * \operatorname{EllipticE}((-1/a*b)^{1/2}*x, (a/b/c*d)^{1/2}) * b*c^2 + (-1/a*b)^{1/2} * x * a * c * d / (b*d*x^4+a*d*x^2+b*c*x^2+a*c) / (-1/a*b)^{1/2} / d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{bx^2 + a} \sqrt{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)*sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{bx^2 + a} \sqrt{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2),x)

[Out] int((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a + bx^2} \sqrt{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)*(d*x**2+c)**(1/2),x)

[Out] Integral(sqrt(a + b*x**2)*sqrt(c + d*x**2), x)

$$3.168 \quad \int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=204

$$\frac{c^{3/2}\sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + \frac{dx\sqrt{a+bx^2}}{b\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}}{a\sqrt{d}\sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $d*x*(b*x^2+a)^{(1/2)}/b/(d*x^2+c)^{(1/2)}+c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/a/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/b/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.09, antiderivative size = 204, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {422, 418, 492, 411}

$$\frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) + \frac{dx\sqrt{a+bx^2}}{b\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}}{a\sqrt{d}\sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x]

[Out] $(d*x*\operatorname{Sqrt}[a + b*x^2])/(b*\operatorname{Sqrt}[c + d*x^2]) - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2])*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)]/(b*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (c^{(3/2)}*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)]/(a*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx &= c \int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx + d \int \frac{x^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx \\ &= \frac{dx\sqrt{a+bx^2}}{b\sqrt{c+dx^2}} + \frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} - \frac{(cd) \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^{3/2}} dx}{b} \\ &= \frac{dx\sqrt{a+bx^2}}{b\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} + \frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 86, normalized size = 0.42

$$\frac{\sqrt{\frac{a+bx^2}{a}}\sqrt{c+dx^2} E\left(\sin^{-1}\left(\sqrt{-\frac{b}{a}}x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{-\frac{b}{a}}\sqrt{a+bx^2}\sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], (a*d)/(b*c)))/(Sqrt[-(b/a)]*Sqrt[a + b*x^2]*Sqrt[(c + d*x^2)/c])
```

fricas [F] time = 1.28, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{dx^2+c}}{\sqrt{bx^2+a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(d*x^2 + c)/sqrt(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(d*x^2 + c)/sqrt(b*x^2 + a), x)
```

maple [A] time = 0.02, size = 101, normalized size = 0.50

$$\frac{\sqrt{dx^2+c} \sqrt{bx^2+a} \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} c \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right)}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+c)^(1/2)/(b*x^2+a)^(1/2),x)`

[Out] `(d*x^2+c)^(1/2)*(b*x^2+a)^(1/2)*c*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*x^2+c)/sqrt(b*x^2+a),x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c+d*x^2)^(1/2)/(a+b*x^2)^(1/2),x)`

[Out] `int((c+d*x^2)^(1/2)/(a+b*x^2)^(1/2),x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**(1/2)/(b*x**2+a)**(1/2),x)`

[Out] `Integral(sqrt(c+d*x**2)/sqrt(a+b*x**2),x)`

$$3.169 \quad \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{3/2}} dx$$

Optimal. Leaf size=84

$$\frac{\sqrt{c+dx^2} E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\left|1-\frac{ad}{bc}\right.\right)}{\sqrt{a}\sqrt{b}\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

[Out] $(1/(1+b*x^2/a))^{(1/2)}*(1+b*x^2/a)^{(1/2)}*EllipticE(x*b^{(1/2)}/a^{(1/2)}/(1+b*x^2/a)^{(1/2)},(1-a*d/b/c)^{(1/2)})*(d*x^2+c)^{(1/2)}/a^{(1/2)}/b^{(1/2)}/(b*x^2+a)^{(1/2)}/(a*(d*x^2+c)/c/(b*x^2+a))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {411}

$$\frac{\sqrt{c+dx^2} E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\left|1-\frac{ad}{bc}\right.\right)}{\sqrt{a}\sqrt{b}\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/(a + b*x^2)^(3/2), x]

[Out] $(\text{Sqrt}[c + d*x^2]*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]], 1 - (a*d)/(b*c)])/(\text{Sqrt}[a]*\text{Sqrt}[b]*\text{Sqrt}[a + b*x^2]*\text{Sqrt}[(a*(c + d*x^2))/(c*(a + b*x^2))])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rubi steps

$$\int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{3/2}} dx = \frac{\sqrt{c+dx^2} E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\left|1-\frac{ad}{bc}\right.\right)}{\sqrt{a}\sqrt{b}\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

Mathematica [C] time = 0.29, size = 133, normalized size = 1.58

$$\frac{x(c+dx^2) + \frac{ic\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}\left(E\left(i\sinh^{-1}\left(\sqrt{\frac{b}{a}}x\right)\left|\frac{ad}{bc}\right.\right)-\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right),\frac{ad}{bc}\right)\right)}{\sqrt{\frac{b}{a}}}}{a\sqrt{a+bx^2}\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + d*x^2]/(a + b*x^2)^(3/2), x]

[Out] $(x*(c + d*x^2) + (I*c*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*(EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]))/Sqrt[b/a])/(a*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])$

fricas [F] time = 0.49, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2 + a}\sqrt{dx^2 + c}}{b^2x^4 + 2abx^2 + a^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(3/2),x, algorithm="fricas")`

[Out] `integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b^2*x^4 + 2*a*b*x^2 + a^2), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 + c}}{(bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(3/2),x, algorithm="giac")`

[Out] `integrate(sqrt(d*x^2 + c)/(b*x^2 + a)^(3/2), x)`

maple [A] time = 0.04, size = 181, normalized size = 2.15

$$\frac{\sqrt{dx^2 + c} \sqrt{bx^2 + a} \left(\sqrt{-\frac{b}{a}} dx^3 + \sqrt{-\frac{b}{a}} cx - \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} c \text{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right) + \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} c \right)}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{b}{a}} a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+c)^(1/2)/(b*x^2+a)^(3/2),x)`

[Out] $(d*x^2+c)^{(1/2)}*(b*x^2+a)^{(1/2)}*(x^3*d*(-1/a*b)^{(1/2)}+EllipticF((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*c*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}-EllipticE((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*c*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}+x*c*(-1/a*b)^{(1/2)})/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/a/(-1/a*b)^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 + c}}{(bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(3/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*x^2 + c)/(b*x^2 + a)^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{dx^2 + c}}{(bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^(1/2)/(a + b*x^2)^(3/2), x)`

[Out] `int((c + d*x^2)^(1/2)/(a + b*x^2)^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2}}{(a + bx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**(1/2)/(b*x**2+a)**(3/2), x)`

[Out] `Integral(sqrt(c + d*x**2)/(a + b*x**2)**(3/2), x)`

$$3.170 \quad \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{5/2}} dx$$

Optimal. Leaf size=237

$$\frac{c^{3/2}\sqrt{d}\sqrt{a+bx^2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)+\sqrt{c+dx^2}(2bc-ad)E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|1-\frac{ad}{bc}\right)+\frac{x\sqrt{c+dx^2}}{3a(a+bx^2)}}{3a^2\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}+3a^{3/2}\sqrt{b}\sqrt{a+bx^2}(bc-ad)\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

[Out] $-1/3*c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a^2/(-a*d+b*c)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/3*x*(d*x^2+c)^{(1/2)}/a/(b*x^2+a)^{(3/2)}+1/3*(-a*d+2*b*c)*(1/(1+b*x^2/a))^{(1/2)}*(1+b*x^2/a)^{(1/2)}*\operatorname{EllipticE}(x*b^{(1/2)}/a^{(1/2)}/(1+b*x^2/a)^{(1/2)},(1-a*d/b/c)^{(1/2)})*(d*x^2+c)^{(1/2)}/a^{(3/2)}/(-a*d+b*c)/b^{(1/2)}/(b*x^2+a)^{(1/2)}/(a*(d*x^2+c)/c/(b*x^2+a))^{(1/2)}$

Rubi [A] time = 0.12, antiderivative size = 237, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {412, 525, 418, 411}

$$\frac{c^{3/2}\sqrt{d}\sqrt{a+bx^2}F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right)+\sqrt{c+dx^2}(2bc-ad)E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|1-\frac{ad}{bc}\right)+\frac{x\sqrt{c+dx^2}}{3a(a+bx^2)^{3/2}}}{3a^2\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}+3a^{3/2}\sqrt{b}\sqrt{a+bx^2}(bc-ad)\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/(a + b*x^2)^(5/2), x]

[Out] $(x*\operatorname{Sqrt}[c + d*x^2])/(3*a*(a + b*x^2)^{(3/2)}) + ((2*b*c - a*d)*\operatorname{Sqrt}[c + d*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]], 1 - (a*d)/(b*c)])/(3*a^{(3/2)}*\operatorname{Sqrt}[b]*(b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[(a*(c + d*x^2))/(c*(a + b*x^2))]) - (c^{(3/2)}*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(3*a^2*(b*c - a*d)*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 412

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*n*(p + 1)), x] + Dist[1/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(n*(p + 1) + 1) + d*(n*(p + q + 1) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && LtQ[0, q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre

$eQ[\{a, b, c, d\}, x] \&\& \text{PosQ}[d/c] \&\& \text{PosQ}[b/a] \&\& !\text{SimplerSqrtQ}[b/a, d/c]$

Rule 525

$\text{Int}[(e_ + (f_ \cdot)(x_)^2)/(\text{Sqrt}[(a_ + (b_ \cdot)(x_)^2] \cdot ((c_ + (d_ \cdot)(x_)^2)^{3/2})], x_Symbol] := \text{Dist}[(b \cdot e - a \cdot f)/(b \cdot c - a \cdot d), \text{Int}[1/(\text{Sqrt}[a + b \cdot x^2] \cdot \text{Sqrt}[c + d \cdot x^2]), x], x] - \text{Dist}[(d \cdot e - c \cdot f)/(b \cdot c - a \cdot d), \text{Int}[\text{Sqrt}[a + b \cdot x^2]/(c + d \cdot x^2)^{3/2}, x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && PosQ[b/a] && PosQ[d/c]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{5/2}} dx &= \frac{x\sqrt{c+dx^2}}{3a(a+bx^2)^{3/2}} - \frac{\int \frac{-2c-dx^2}{(a+bx^2)^{3/2}\sqrt{c+dx^2}} dx}{3a} \\ &= \frac{x\sqrt{c+dx^2}}{3a(a+bx^2)^{3/2}} - \frac{(cd) \int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx}{3a(bc-ad)} + \frac{(2bc-ad) \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{3/2}} dx}{3a(bc-ad)} \\ &= \frac{x\sqrt{c+dx^2}}{3a(a+bx^2)^{3/2}} + \frac{(2bc-ad)\sqrt{c+dx^2} E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 1 - \frac{ad}{bc}\right)}{3a^{3/2}\sqrt{b}(bc-ad)\sqrt{a+bx^2} \sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}} - \frac{c^{3/2}\sqrt{d}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{c(a+bx^2)}{c(a+bx^2)}\right)}{3a^2(bc-ad)\sqrt{\frac{c(a+bx^2)}{c(a+bx^2)}}} \end{aligned}$$

Mathematica [C] time = 0.47, size = 243, normalized size = 1.03

$$\frac{-2ic(a+bx^2)\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}(ad-bc)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) + x\sqrt{\frac{b}{a}}(c+dx^2)(2a^2d+ab(dx^2-3c))}{3a^2\sqrt{\frac{b}{a}}(a+bx^2)^{3/2}\sqrt{c+dx^2}(ad-bc)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + d*x^2]/(a + b*x^2)^(5/2), x]

[Out] (Sqrt[b/a]*x*(c + d*x^2)*(2*a^2*d - 2*b^2*c*x^2 + a*b*(-3*c + d*x^2)) + I*c*(-2*b*c + a*d)*(a + b*x^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - (2*I)*c*(-(b*c) + a*d)*(a + b*x^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)])/(3*a^2*Sqrt[b/a]*(-(b*c) + a*d)*(a + b*x^2)^(3/2)*Sqrt[c + d*x^2])

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+a}\sqrt{dx^2+c}}{b^3x^6+3ab^2x^4+3a^2bx^2+a^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b^3*x^6 + 3*a*b^2*x^4 + 3*a^2*b*x^2 + a^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{(bx^2+a)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(d*x^2 + c)/(b*x^2 + a)^(5/2), x)

maple [B] time = 0.05, size = 617, normalized size = 2.60

$$\sqrt{-\frac{b}{a}} ab d^2 x^5 - 2\sqrt{-\frac{b}{a}} b^2 cd x^5 + 2\sqrt{-\frac{b}{a}} a^2 d^2 x^3 - 2\sqrt{-\frac{b}{a}} abcd x^3 - \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} abcd x^2 \text{EllipticE}\left(\sqrt{-\frac{b}{a}} x,\right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^(1/2)/(b*x^2+a)^(5/2),x)

[Out] $\frac{1}{3}(x^5 a b d^2 (-1/a b)^{1/2} - 2 x^5 b^2 c d (-1/a b)^{1/2} + 2 \text{EllipticF}((-1/a b)^{1/2} x, (a/b/c d)^{1/2}) x^2 a b c d ((b x^2 + a)/a)^{1/2} ((d x^2 + c)/c)^{1/2} - 2 \text{EllipticF}((-1/a b)^{1/2} x, (a/b/c d)^{1/2}) x^2 b^2 c^2 ((b x^2 + a)/a)^{1/2} ((d x^2 + c)/c)^{1/2} - \text{EllipticE}((-1/a b)^{1/2} x, (a/b/c d)^{1/2}) x^2 a b c d ((b x^2 + a)/a)^{1/2} ((d x^2 + c)/c)^{1/2} + 2 \text{EllipticE}((-1/a b)^{1/2} x, (a/b/c d)^{1/2}) x^2 b^2 c^2 ((b x^2 + a)/a)^{1/2} ((d x^2 + c)/c)^{1/2} + 2 x^3 a^2 d^2 (-1/a b)^{1/2} - 2 x^3 a b c d (-1/a b)^{1/2} - 2 x^3 b^2 c^2 (-1/a b)^{1/2} + 2 \text{EllipticF}((-1/a b)^{1/2} x, (a/b/c d)^{1/2}) a^2 c d ((b x^2 + a)/a)^{1/2} ((d x^2 + c)/c)^{1/2} - 2 \text{EllipticF}((-1/a b)^{1/2} x, (a/b/c d)^{1/2}) a b c^2 ((b x^2 + a)/a)^{1/2} ((d x^2 + c)/c)^{1/2} - \text{EllipticE}((-1/a b)^{1/2} x, (a/b/c d)^{1/2}) a^2 c d ((b x^2 + a)/a)^{1/2} ((d x^2 + c)/c)^{1/2} + 2 \text{EllipticE}((-1/a b)^{1/2} x, (a/b/c d)^{1/2}) a b c^2 ((b x^2 + a)/a)^{1/2} ((d x^2 + c)/c)^{1/2} + 2 x a^2 c d (-1/a b)^{1/2} - 3 x a b c^2 (-1/a b)^{1/2}) / (d x^2 + c)^{1/2} / (-1/a b)^{1/2} / (a d - b c) / a^2 / (b x^2 + a)^{3/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 + c}}{(bx^2 + a)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(5/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*x^2 + c)/(b*x^2 + a)^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{dx^2 + c}}{(bx^2 + a)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^(1/2)/(a + b*x^2)^(5/2),x)

[Out] int((c + d*x^2)^(1/2)/(a + b*x^2)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2}}{(a + bx^2)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**(1/2)/(b*x**2+a)**(5/2),x)

[Out] Integral(sqrt(c + d*x**2)/(a + b*x**2)**(5/2), x)

$$3.171 \quad \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{7/2}} dx$$

Optimal. Leaf size=309

$$\frac{2c^{3/2}\sqrt{d}\sqrt{a+bx^2}(2bc-3ad)\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)+x\sqrt{c+dx^2}(4bc-3ad)+\sqrt{c+dx^2}(3a^2d^2-15a^3\sqrt{c+dx^2}(bc-ad)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}{15a^2(a+bx^2)^{3/2}(bc-ad)+15a^5\sqrt{b}}$$

[Out] $-2/15*c^{(3/2)}*(-3*a*d+2*b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a^3/(-a*d+b*c)^2/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/5*x*(d*x^2+c)^{(1/2)}/a/(b*x^2+a)^{(5/2)}+1/15*(-3*a*d+4*b*c)*x*(d*x^2+c)^{(1/2)}/a^2/(-a*d+b*c)/(b*x^2+a)^{(3/2)}+1/15*(3*a^2*d^2-13*a*b*c*d+8*b^2*c^2)*(1/(1+b*x^2/a))^{(1/2)}*(1+b*x^2/a)^{(1/2)}*\operatorname{EllipticE}(x*b^{(1/2)}/a^{(1/2)}/(1+b*x^2/a)^{(1/2)},(1-a*d/b/c)^{(1/2)}*(d*x^2+c)^{(1/2)}/a^{(5/2)}/(-a*d+b*c)^2/b^{(1/2)}/(b*x^2+a)^{(1/2)}/(a*(d*x^2+c)/c/(b*x^2+a))^{(1/2)}$

Rubi [A] time = 0.23, antiderivative size = 309, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {412, 527, 525, 418, 411}

$$\frac{\sqrt{c+dx^2}(3a^2d^2-13abcd+8b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\left|1-\frac{ad}{bc}\right.\right)+2c^{3/2}\sqrt{d}\sqrt{a+bx^2}(2bc-3ad)F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{15a^{5/2}\sqrt{b}\sqrt{a+bx^2}(bc-ad)^2\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}-15a^3\sqrt{c+dx^2}(bc-ad)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/(a + b*x^2)^(7/2),x]

[Out] $(x*\operatorname{Sqrt}[c + d*x^2])/(5*a*(a + b*x^2)^{(5/2)}) + ((4*b*c - 3*a*d)*x*\operatorname{Sqrt}[c + d*x^2])/(15*a^2*(b*c - a*d)*(a + b*x^2)^{(3/2)}) + ((8*b^2*c^2 - 13*a*b*c*d + 3*a^2*d^2)*\operatorname{Sqrt}[c + d*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]], 1 - (a*d)/(b*c)])/(15*a^{(5/2)}*\operatorname{Sqrt}[b]*(b*c - a*d)^2*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[(a*(c + d*x^2))/(c*(a + b*x^2))]) - (2*c^{(3/2)}*\operatorname{Sqrt}[d]*(2*b*c - 3*a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(15*a^3*(b*c - a*d)^2*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 412

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*n*(p+1)), x] + Dist[1/(a*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1)*Simp[c*(n*(p+1)+1) + d*(n*(p+q+1)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && LtQ[0, q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 525

```
Int[((e_) + (f_.)*(x_)^2)/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)^(
3/2)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(Sqrt[a + b*x^2]*S
qrt[c + d*x^2]), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[Sqrt[a + b*x^2]
/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && PosQ[b/a] &&
PosQ[d/c]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c+dx^2}}{(a+bx^2)^{7/2}} dx &= \frac{x\sqrt{c+dx^2}}{5a(a+bx^2)^{5/2}} - \frac{\int \frac{-4c-3dx^2}{(a+bx^2)^{5/2}\sqrt{c+dx^2}} dx}{5a} \\ &= \frac{x\sqrt{c+dx^2}}{5a(a+bx^2)^{5/2}} + \frac{(4bc-3ad)x\sqrt{c+dx^2}}{15a^2(bc-ad)(a+bx^2)^{3/2}} + \frac{\int \frac{c(8bc-9ad)+d(4bc-3ad)x^2}{(a+bx^2)^{3/2}\sqrt{c+dx^2}} dx}{15a^2(bc-ad)} \\ &= \frac{x\sqrt{c+dx^2}}{5a(a+bx^2)^{5/2}} + \frac{(4bc-3ad)x\sqrt{c+dx^2}}{15a^2(bc-ad)(a+bx^2)^{3/2}} - \frac{(2cd(2bc-3ad)) \int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx}{15a^2(bc-ad)^2} + \frac{(8b^2c^2-13abcd+3a^2d^2)\sqrt{c+dx^2}E(\tan^{-1}(\frac{x\sqrt{c+dx^2}}{\sqrt{a+bx^2}}))}{15a^5\sqrt{b}(bc-ad)^2\sqrt{a+bx^2}} \end{aligned}$$

Mathematica [C] time = 0.55, size = 285, normalized size = 0.92

$$\frac{x\sqrt{\frac{b}{a}(c+dx^2)}\left(\left(a+bx^2\right)^2\left(3a^2d^2-13abcd+8b^2c^2\right)+3a^2(bc-ad)^2+a\left(a+bx^2\right)(ad-bc)(3ad-4bc)\right)+ic}{15a^5\sqrt{b}(bc-ad)^2\sqrt{a+bx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[c + d*x^2]/(a + b*x^2)^(7/2), x]
```

```
[Out] (Sqrt[b/a]*x*(c + d*x^2)*(3*a^2*(b*c - a*d)^2 + a*(-(b*c) + a*d)*(-4*b*c +
3*a*d)*(a + b*x^2) + (8*b^2*c^2 - 13*a*b*c*d + 3*a^2*d^2)*(a + b*x^2)^2) +
I*c*(a + b*x^2)^2*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*((8*b^2*c^2 - 13*
a*b*c*d + 3*a^2*d^2)*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + (-8*b
```

$\sqrt{2c^2 + 17ab^3cd - 9a^2d^2} \text{EllipticF}[\text{I} \cdot \text{ArcSinh}[\sqrt{b/a}x], (ad)/(bc)] / (15a^3 \sqrt{b/a} (bc - ad)^2 (a + bx^2)^{5/2} \sqrt{c + dx^2})$

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{bx^2 + a} \sqrt{dx^2 + c}}{b^4x^8 + 4ab^3x^6 + 6a^2b^2x^4 + 4a^3bx^2 + a^4}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(7/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b^4*x^8 + 4*a*b^3*x^6 + 6*a^2*b^2*x^4 + 4*a^3*b*x^2 + a^4), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 + c}}{(bx^2 + a)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(7/2),x, algorithm="giac")

[Out] integrate(sqrt(d*x^2 + c)/(b*x^2 + a)^(7/2), x)

maple [B] time = 0.06, size = 1411, normalized size = 4.57

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^(1/2)/(b*x^2+a)^(7/2),x)

[Out] $-1/15*(30*x^5*a^2*b^2*c*d^2*(-1/a*b)^{(1/2)} - 7*x^5*a*b^3*c^2*d*(-1/a*b)^{(1/2)} + 17*x^3*a^3*b*c*d^2*(-1/a*b)^{(1/2)} + 18*x^3*a^2*b^2*c^2*d*(-1/a*b)^{(1/2)} + 26*x*a^3*b*c^2*d*(-1/a*b)^{(1/2)} + 8*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*b^4*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 8*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*b^4*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 3*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^4*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 8*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^2*b^2*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 9*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^4*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 8*x^5*b^4*c^3*(-1/a*b)^{(1/2)} - 9*x^3*a^4*d^3*(-1/a*b)^{(1/2)} + 13*x^7*a*b^3*c*d^2*(-1/a*b)^{(1/2)} - 3*x^7*a^2*b^2*d^3*(-1/a*b)^{(1/2)} - 13*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*a*b^3*c^2*d*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 3*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*a^2*b^2*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 34*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a^2*b^2*c^2*d*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 6*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a^3*b*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 17*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*a*b^3*c^2*d*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 9*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*a^2*b^2*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 18*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a^3*b*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 26*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a^2*b^2*c^2*d*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 16*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a*b^3*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 16*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a*b^3*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 13*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^3*b*c^2*d*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 17*\text{Ellip$

```
ticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^3*b*c^2*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-8*x^7*b^4*c^2*d*(-1/a*b)^(1/2)-9*x^5*a^3*b*d^3*(-1/a*b)^(1/2)-20*x^3*a*b^3*c^3*(-1/a*b)^(1/2)-9*x*a^4*c*d^2*(-1/a*b)^(1/2)-15*x*a^2*b^2*c^3*(-1/a*b)^(1/2))/(d*x^2+c)^(1/2)/(-1/a*b)^(1/2)/(a*d-b*c)^2/a^3/(b*x^2+a)^(5/2)
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 + c}}{(bx^2 + a)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(7/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(d*x^2 + c)/(b*x^2 + a)^(7/2), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{dx^2 + c}}{(bx^2 + a)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c + d*x^2)^(1/2)/(a + b*x^2)^(7/2),x)
```

```
[Out] int((c + d*x^2)^(1/2)/(a + b*x^2)^(7/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2}}{(a + bx^2)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x**2+c)**(1/2)/(b*x**2+a)**(7/2),x)
```

```
[Out] Integral(sqrt(c + d*x**2)/(a + b*x**2)**(7/2), x)
```

$$3.172 \quad \int (a + bx^2)^{3/2} (c + dx^2)^{3/2} dx$$

Optimal. Leaf size=410

$$\frac{c^{3/2}\sqrt{a+bx^2} (a^2d^2 - 18abcd + b^2c^2) \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) 2x\sqrt{a+bx^2} (ad+bc) (a^2d^2 - 6abcd + b^2c^2)}{35bd^{3/2}\sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} 35b^2d\sqrt{c+dx^2}}$$

[Out] $-2/35*(a*d+b*c)*(a^2*d^2-6*a*b*c*d+b^2*c^2)*x*(b*x^2+a)^{(1/2)}/b^2/d/(d*x^2+c)^{(1/2)}-1/35*c^{(3/2)}*(a^2*d^2-18*a*b*c*d+b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/b/d^{(3/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+2/35*(a*d+b*c)*(a^2*d^2-6*a*b*c*d+b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/b^2/d^{(3/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+2/35*(-a*d+4*b*c)*x*(b*x^2+a)^{(3/2)}*(d*x^2+c)^{(1/2)}/b+1/7*d*x*(b*x^2+a)^{(5/2)}*(d*x^2+c)^{(1/2)}/b+1/35*(9*a*c+b*c^2/d-2*a^2*d/b)*x*(b*x^2+a)^{(1/2)}*(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.44, antiderivative size = 410, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {416, 528, 531, 418, 492, 411}

$$\frac{2x\sqrt{a+bx^2} (ad+bc) (a^2d^2 - 6abcd + b^2c^2)}{35b^2d\sqrt{c+dx^2}} - \frac{c^{3/2}\sqrt{a+bx^2} (a^2d^2 - 18abcd + b^2c^2) F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{35bd^{3/2}\sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{2\sqrt{c+dx^2}}{35b^2d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)*(c + d*x^2)^(3/2), x]

[Out] $(-2*(b*c + a*d)*(b^2*c^2 - 6*a*b*c*d + a^2*d^2)*x*\operatorname{Sqrt}[a + b*x^2])/(35*b^2*d*\operatorname{Sqrt}[c + d*x^2]) + ((9*a*c + (b*c^2)/d - (2*a^2*d)/b)*x*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2])/35 + (2*(4*b*c - a*d)*x*(a + b*x^2)^{(3/2)}*\operatorname{Sqrt}[c + d*x^2])/(35*b) + (d*x*(a + b*x^2)^{(5/2)}*\operatorname{Sqrt}[c + d*x^2])/(7*b) + (2*\operatorname{Sqrt}[c]*(b*c + a*d)*(b^2*c^2 - 6*a*b*c*d + a^2*d^2)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(35*b^2*d^{(3/2)}*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) - (c^{(3/2)}*(b^2*c^2 - 18*a*b*c*d + a^2*d^2)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(35*b*d^{(3/2)}*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_)*(x_)^2]/((c_) + (d_)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 416

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q)+1)), x] + Dist[1/(b*(n*(p+q)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q)+1) - a*d) + d*(b*c*(n*(p+2*q-1)+1) - a*d*(n*(q-1)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q)+1, 0] && !IGtQ[p, 1] && IntBinomialQ[a

, b, c, d, n, p, q, x]

Rule 418

```
Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
:= Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rule 528

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
f_)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rule 531

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
f_)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x],
x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c,
d, e, f, n, p, q}, x]
```

Rubi steps

$$\begin{aligned}
 \int (a + bx^2)^{3/2} (c + dx^2)^{3/2} dx &= \frac{dx (a + bx^2)^{5/2} \sqrt{c + dx^2}}{7b} + \frac{\int \frac{(a+bx^2)^{3/2} (c(7bc-ad)+2d(4bc-ad)x^2)}{\sqrt{c+dx^2}} dx}{7b} \\
 &= \frac{2(4bc - ad)x (a + bx^2)^{3/2} \sqrt{c + dx^2}}{35b} + \frac{dx (a + bx^2)^{5/2} \sqrt{c + dx^2}}{7b} + \frac{\int \frac{\sqrt{a+bx^2} (3ac + bc^2 - 2a^2d)}{\sqrt{c+dx^2}} dx}{35b} \\
 &= \frac{1}{35} \left(9ac + \frac{bc^2}{d} - \frac{2a^2d}{b} \right) x \sqrt{a + bx^2} \sqrt{c + dx^2} + \frac{2(4bc - ad)x (a + bx^2)^{3/2} \sqrt{c + dx^2}}{35b} \\
 &= \frac{1}{35} \left(9ac + \frac{bc^2}{d} - \frac{2a^2d}{b} \right) x \sqrt{a + bx^2} \sqrt{c + dx^2} + \frac{2(4bc - ad)x (a + bx^2)^{3/2} \sqrt{c + dx^2}}{35b} \\
 &= -\frac{2(bc + ad) (b^2c^2 - 6abcd + a^2d^2) x \sqrt{a + bx^2}}{35b^2d\sqrt{c + dx^2}} + \frac{1}{35} \left(9ac + \frac{bc^2}{d} - \frac{2a^2d}{b} \right) x \sqrt{a + bx^2} \sqrt{c + dx^2} \\
 &= -\frac{2(bc + ad) (b^2c^2 - 6abcd + a^2d^2) x \sqrt{a + bx^2}}{35b^2d\sqrt{c + dx^2}} + \frac{1}{35} \left(9ac + \frac{bc^2}{d} - \frac{2a^2d}{b} \right) x \sqrt{a + bx^2} \sqrt{c + dx^2}
 \end{aligned}$$

Mathematica [C] time = 0.60, size = 302, normalized size = 0.74

$$-ic\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}\left(a^3d^3+8a^2bcd^2-11ab^2c^2d+2b^3c^3\right)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right),\frac{ad}{bc}\right)+dx\sqrt{\frac{b}{a}}\left(a+bx^2\right)(c$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)*(c + d*x^2)^(3/2), x]

[Out] (Sqrt[b/a]*d*x*(a + b*x^2)*(c + d*x^2)*(a^2*d^2 + a*b*d*(17*c + 8*d*x^2) + b^2*(c^2 + 8*c*d*x^2 + 5*d^2*x^4)) + (2*I)*c*(b^3*c^3 - 5*a*b^2*c^2*d - 5*a^2*b*c*d^2 + a^3*d^3)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*c*(2*b^3*c^3 - 11*a*b^2*c^2*d + 8*a^2*b*c*d^2 + a^3*d^3)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(35*b*Sqrt[b/a]*d^2*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(bdx^4 + (bc + ad)x^2 + ac\right)\sqrt{bx^2 + a}\sqrt{dx^2 + c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^(3/2), x, algorithm="fricas")

[Out] integral((b*d*x^4 + (b*c + a*d)*x^2 + a*c)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^{\frac{3}{2}}(dx^2 + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^(3/2), x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(3/2)*(d*x^2 + c)^(3/2), x)

maple [A] time = 0.04, size = 780, normalized size = 1.90

$$\sqrt{bx^2 + a}\sqrt{dx^2 + c}\left(5\sqrt{-\frac{b}{a}}b^3d^4x^9 + 13\sqrt{-\frac{b}{a}}ab^2d^4x^7 + 13\sqrt{-\frac{b}{a}}b^3cd^3x^7 + 9\sqrt{-\frac{b}{a}}a^2bd^4x^5 + 38\sqrt{-\frac{b}{a}}ab^2cd^3x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)*(d*x^2+c)^(3/2), x)

[Out] 1/35*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(5*(-1/a*b)^(1/2)*x^9*b^3*d^4+13*(-1/a*b)^(1/2)*x^7*a*b^2*d^4+13*(-1/a*b)^(1/2)*x^7*b^3*c*d^3+9*(-1/a*b)^(1/2)*x^5*a^2*b*d^4+38*(-1/a*b)^(1/2)*x^5*a*b^2*c*d^3+9*(-1/a*b)^(1/2)*x^5*b^3*c^2*d^2+(-1/a*b)^(1/2)*x^3*a^3*d^4+26*(-1/a*b)^(1/2)*x^3*a^2*b*c*d^3+26*(-1/a*b)^(1/2)*x^3*a*b^2*c^2*d^2+(-1/a*b)^(1/2)*x^3*b^3*c^3*d+((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^3*c*d^3+8*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b*c^2*d^2-11*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b^2*c^3*d+2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b^3*c^4-2*((b*x^

$2+a)/a)^{1/2}*((d*x^2+c)/c)^{1/2}*EllipticE((-1/a*b)^{1/2}*x, (a/b/c*d)^{1/2})$
 $)^3*c*d^3+10*((b*x^2+a)/a)^{1/2}*((d*x^2+c)/c)^{1/2}*EllipticE((-1/a*b)^{1/2}$
 $^{1/2}*x, (a/b/c*d)^{1/2})^2*b*c^2*d^2+10*((b*x^2+a)/a)^{1/2}*((d*x^2+c)/c)$
 $^{1/2}*EllipticE((-1/a*b)^{1/2}*x, (a/b/c*d)^{1/2})^2*a*b^2*c^3*d-2*((b*x^2+a)$
 $/a)^{1/2}*((d*x^2+c)/c)^{1/2}*EllipticE((-1/a*b)^{1/2}*x, (a/b/c*d)^{1/2})^2*b$
 $^3*c^4+(-1/a*b)^{1/2}*x*a^3*c*d^3+17*(-1/a*b)^{1/2}*x*a^2*b*c^2*d^2+(-1/a*b)$
 $)^{1/2}*x*a*b^2*c^3*d)/b/d^2/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^{1/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^{\frac{3}{2}} (dx^2 + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)*(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/2)*(d*x^2 + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (bx^2 + a)^{3/2} (dx^2 + c)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)*(c + d*x^2)^(3/2),x)

[Out] int((a + b*x^2)^(3/2)*(c + d*x^2)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)^{\frac{3}{2}} (c + dx^2)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)*(d*x**2+c)**(3/2),x)

[Out] Integral((a + b*x**2)**(3/2)*(c + d*x**2)**(3/2), x)

3.173 $\int \sqrt{a + bx^2} (c + dx^2)^{3/2} dx$

Optimal. Leaf size=336

$$\frac{c^{3/2}\sqrt{a+bx^2}(9bc-ad)\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)+x\sqrt{a+bx^2}(-2a^2d^2+7abcd+3b^2c^2)\sqrt{c}\sqrt{a+bx^2}}{15b\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}+15b^2\sqrt{c+dx^2}}$$

[Out] $\frac{1}{15}*(-2*a^2*d^2+7*a*b*c*d+3*b^2*c^2)*x*(b*x^2+a)^{(1/2)}/b^2/(d*x^2+c)^{(1/2)}$
 $+1/15*c^{(3/2)}*(-a*d+9*b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/b/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$
 $-1/15*(-2*a^2*d^2+7*a*b*c*d+3*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/b^2/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$
 $+1/5*d*x*(b*x^2+a)^{(3/2)}*(d*x^2+c)^{(1/2)}/b+2/15*(-a*d+3*b*c)*x*(b*x^2+a)^{(1/2)}*(d*x^2+c)^{(1/2)}/b$

Rubi [A] time = 0.28, antiderivative size = 336, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {416, 528, 531, 418, 492, 411}

$$\frac{x\sqrt{a+bx^2}(-2a^2d^2+7abcd+3b^2c^2)\sqrt{c}\sqrt{a+bx^2}(-2a^2d^2+7abcd+3b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)+c^{3/2}\sqrt{a+bx^2}}{15b^2\sqrt{c+dx^2}+15b^2\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[a + b*x^2]*(c + d*x^2)^(3/2),x]`

[Out] $((3*b^2*c^2 + 7*a*b*c*d - 2*a^2*d^2)*x*\operatorname{Sqrt}[a + b*x^2])/(15*b^2*\operatorname{Sqrt}[c + d*x^2]) + (2*(3*b*c - a*d)*x*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2])/(15*b) + (d*x*(a + b*x^2)^{(3/2)}*\operatorname{Sqrt}[c + d*x^2])/(5*b) - (\operatorname{Sqrt}[c]*(3*b^2*c^2 + 7*a*b*c*d - 2*a^2*d^2)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]],1 - (b*c)/(a*d)])/(15*b^2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (c^{(3/2)}*(9*b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]],1 - (b*c)/(a*d)])/(15*b*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

`Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]`

Rule 416

`Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q)+1)), x] + Dist[1/(b*(n*(p+q)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q)+1) - a*d] + d*(b*c*(n*(p+2*q-1)+1) - a*d*(n*(q-1)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q)+1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]`

Rule 418

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)]/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:= Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rule 531

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x],
x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c,
d, e, f, n, p, q}, x]
```

Rubi steps

$$\begin{aligned} \int \sqrt{a+bx^2} (c+dx^2)^{3/2} dx &= \frac{dx (a+bx^2)^{3/2} \sqrt{c+dx^2}}{5b} + \frac{\int \frac{\sqrt{a+bx^2} (c(5bc-ad)+2d(3bc-ad)x^2)}{\sqrt{c+dx^2}} dx}{5b} \\ &= \frac{2(3bc-ad)x\sqrt{a+bx^2} \sqrt{c+dx^2}}{15b} + \frac{dx (a+bx^2)^{3/2} \sqrt{c+dx^2}}{5b} + \frac{\int \frac{acd(9bc-ad)+d(3bc-ad)x^2}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx}{5b} \\ &= \frac{2(3bc-ad)x\sqrt{a+bx^2} \sqrt{c+dx^2}}{15b} + \frac{dx (a+bx^2)^{3/2} \sqrt{c+dx^2}}{5b} + \frac{(ac(9bc-ad)) \int \frac{dx}{\sqrt{a+bx^2} \sqrt{c+dx^2}}}{5b} \\ &= \frac{(3b^2c^2 + 7abcd - 2a^2d^2) x \sqrt{a+bx^2}}{15b^2 \sqrt{c+dx^2}} + \frac{2(3bc-ad)x\sqrt{a+bx^2} \sqrt{c+dx^2}}{15b} + \frac{dx (ac(9bc-ad))}{15b} \\ &= \frac{(3b^2c^2 + 7abcd - 2a^2d^2) x \sqrt{a+bx^2}}{15b^2 \sqrt{c+dx^2}} + \frac{2(3bc-ad)x\sqrt{a+bx^2} \sqrt{c+dx^2}}{15b} + \frac{dx (ac(9bc-ad))}{15b} \end{aligned}$$

Mathematica [C] time = 0.42, size = 246, normalized size = 0.73

$$\frac{-ic\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}(a^2d^2+2abcd-3b^2c^2)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right),\frac{ad}{bc}\right)+ic\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}(2a^2d^2-15bd\sqrt{\frac{b}{a}}\sqrt{a+bx^2}\sqrt{c+dx^2})}{15bd\sqrt{\frac{b}{a}}\sqrt{a+bx^2}\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2]*(c + d*x^2)^(3/2), x]

[Out] (Sqrt[b/a]*d*x*(a + b*x^2)*(c + d*x^2)*(6*b*c + a*d + 3*b*d*x^2) + I*c*(-3*b^2*c^2 - 7*a*b*c*d + 2*a^2*d^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*c*(-3*b^2*c^2 + 2*a*b*c*d + a^2*d^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(15*b*Sqrt[b/a]*d*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{bx^2 + a}(dx^2 + c)^{\frac{3}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{bx^2 + a}(dx^2 + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2), x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2), x)

maple [A] time = 0.03, size = 545, normalized size = 1.62

$$\sqrt{bx^2 + a} \sqrt{dx^2 + c} \left(3\sqrt{-\frac{b}{a}} b^2 d^3 x^7 + 4\sqrt{-\frac{b}{a}} ab d^3 x^5 + 9\sqrt{-\frac{b}{a}} b^2 c d^2 x^5 + \sqrt{-\frac{b}{a}} a^2 d^3 x^3 + 10\sqrt{-\frac{b}{a}} abc d^2 x^3 + 6\sqrt{-\frac{b}{a}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2), x)

[Out] 1/15*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(3*(-1/a*b)^(1/2)*b^2*d^3*x^7+4*(-1/a*b)^(1/2)*a*b*d^3*x^5+9*(-1/a*b)^(1/2)*b^2*c*d^2*x^5+(-1/a*b)^(1/2)*a^2*d^3*x^3+10*(-1/a*b)^(1/2)*a*b*c*d^2*x^3+6*(-1/a*b)^(1/2)*b^2*c^2*d*x^3+((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*c*d^2+2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b*c^2*d-3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b^2*c^3-2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*c*d^2+7*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b*c^2*d+3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b^2*c^3+(-1/a*b)^(1/2)*a^2*c*d^2*x+6*(-1/a*b)^(1/2)*a*b*c^2*d*x)/d/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/b/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{bx^2 + a}(dx^2 + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)*(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{bx^2 + a} (dx^2 + c)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)*(c + d*x^2)^(3/2),x)

[Out] int((a + b*x^2)^(1/2)*(c + d*x^2)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a + bx^2} (c + dx^2)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)*(d*x**2+c)**(3/2),x)

[Out] Integral(sqrt(a + b*x**2)*(c + d*x**2)**(3/2), x)

$$3.174 \quad \int \frac{(c+dx^2)^{3/2}}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=273

$$\frac{c^{3/2}\sqrt{a+bx^2}(3bc-ad)\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right)}{3ab\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{2dx\sqrt{a+bx^2}(2bc-ad)}{3b^2\sqrt{c+dx^2}} - \frac{2\sqrt{c}\sqrt{d}\sqrt{a+bx^2}(2bc-ad)E}{3b^2\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $\frac{2}{3}d*(-a*d+2*b*c)*x*(b*x^2+a)^{(1/2)}/b^2/(d*x^2+c)^{(1/2)}+1/3*c^{(3/2)}*(-a*d+3*b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/a/b/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-2/3*(-a*d+2*b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/b^2/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/3*d*x*(b*x^2+a)^{(1/2)}*(d*x^2+c)^{(1/2)}/b$

Rubi [A] time = 0.16, antiderivative size = 273, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {416, 531, 418, 492, 411}

$$\frac{2dx\sqrt{a+bx^2}(2bc-ad)}{3b^2\sqrt{c+dx^2}} - \frac{2\sqrt{c}\sqrt{d}\sqrt{a+bx^2}(2bc-ad)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{3b^2\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{c^{3/2}\sqrt{a+bx^2}(3bc-ad)F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)}{3ab\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^(3/2)/Sqrt[a + b*x^2], x]

[Out] $\frac{(2*d*(2*b*c - a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(3*b^2*\operatorname{Sqrt}[c + d*x^2]) + (d*x*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2])/(3*b) - (2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*(2*b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(3*b^2*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (c^{(3/2)}*(3*b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(3*a*b*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q)+1)), x] + Dist[1/(b*(n*(p+q)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q)+1) - a*d) + d*(b*c*(n*(p+2*q-1)+1) - a*d*(n*(q-1)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q)+1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R

$\text{t}[d/c, 2] * \text{Sqrt}[c + d*x^2] * \text{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))], x] /;$ FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 492

$\text{Int}[(x_)^2/(\text{Sqrt}[(a_) + (b_.)*(x_)^2]*\text{Sqrt}[(c_) + (d_.)*(x_)^2]), x_Symbol]$
 $:= \text{Simp}[(x*\text{Sqrt}[a + b*x^2])/(b*\text{Sqrt}[c + d*x^2]), x] - \text{Dist}[c/b, \text{Int}[\text{Sqrt}[a + b*x^2]/(c + d*x^2)^{(3/2)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rule 531

$\text{Int}[(a_) + (b_.)*(x_)^{(n_)}]^{(p_.)}*((c_) + (d_.)*(x_)^{(n_)})^{(q_.)}*((e_) + (f_.)*(x_)^{(n_)}), x_Symbol]$:= Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x], x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, e, f, n, p, q}, x]

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^{3/2}}{\sqrt{a + bx^2}} dx &= \frac{dx\sqrt{a + bx^2}\sqrt{c + dx^2}}{3b} + \frac{\int \frac{c(3bc - ad) + 2d(2bc - ad)x^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{3b} \\ &= \frac{dx\sqrt{a + bx^2}\sqrt{c + dx^2}}{3b} + \frac{(2d(2bc - ad)) \int \frac{x^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{3b} + \frac{(c(3bc - ad)) \int \frac{1}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{3b} \\ &= \frac{2d(2bc - ad)x\sqrt{a + bx^2}}{3b^2\sqrt{c + dx^2}} + \frac{dx\sqrt{a + bx^2}\sqrt{c + dx^2}}{3b} + \frac{c^{3/2}(3bc - ad)\sqrt{a + bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}}{\sqrt{c}}\right), \frac{c(a + bx^2)}{a(c + dx^2)}\right)}{3ab\sqrt{d}\sqrt{c + dx^2}} \\ &= \frac{2d(2bc - ad)x\sqrt{a + bx^2}}{3b^2\sqrt{c + dx^2}} + \frac{dx\sqrt{a + bx^2}\sqrt{c + dx^2}}{3b} - \frac{2\sqrt{c}\sqrt{d}(2bc - ad)\sqrt{a + bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}}{\sqrt{c}}\right), \frac{c(a + bx^2)}{a(c + dx^2)}\right)}{3b^2\sqrt{c + dx^2}} \end{aligned}$$

Mathematica [C] time = 0.28, size = 199, normalized size = 0.73

$$\frac{-ic\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}(ad - bc)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) + dx\sqrt{\frac{b}{a}}(a + bx^2)(c + dx^2) + 2ic\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}}{3b\sqrt{\frac{b}{a}}\sqrt{a + bx^2}\sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^(3/2)/Sqrt[a + b*x^2], x]

[Out] (Sqrt[b/a]*d*x*(a + b*x^2)*(c + d*x^2) + (2*I)*c*(-2*b*c + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*c*(-(b*c) + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(3*b*Sqrt[b/a]*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(dx^2 + c)^{\frac{3}{2}}}{\sqrt{bx^2 + a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] integral((d*x^2 + c)^(3/2)/sqrt(b*x^2 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((d*x^2 + c)^(3/2)/sqrt(b*x^2 + a), x)

maple [A] time = 0.03, size = 330, normalized size = 1.21

$$\frac{\sqrt{dx^2 + c} \sqrt{bx^2 + a} \left(\sqrt{-\frac{b}{a}} b d^2 x^5 + \sqrt{-\frac{b}{a}} a d^2 x^3 + \sqrt{-\frac{b}{a}} bcd x^3 + \sqrt{-\frac{b}{a}} acdx - 2\sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} acd \text{EllipticE} \left(\right) \right)}{1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^(3/2)/(b*x^2+a)^(1/2),x)

[Out] 1/3*(d*x^2+c)^(1/2)*(b*x^2+a)^(1/2)*((-1/a*b)^(1/2)*b*d^2*x^5+(-1/a*b)^(1/2)*a*d^2*x^3+(-1/a*b)^(1/2)*b*c*d*x^3+a*c*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*d-((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b*c^2-2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*c*d+4*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b*c^2+(-1/a*b)^(1/2)*a*c*d*x)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/b/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((d*x^2 + c)^(3/2)/sqrt(b*x^2 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^(3/2)/(a + b*x^2)^(1/2),x)

[Out] int((c + d*x^2)^(3/2)/(a + b*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + dx^2)^{\frac{3}{2}}}{\sqrt{a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**(3/2)/(b*x**2+a)**(1/2), x)

[Out] Integral((c + d*x**2)**(3/2)/sqrt(a + b*x**2), x)

$$3.175 \quad \int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^{3/2}} dx$$

Optimal. Leaf size=267

$$\frac{c^{3/2}\sqrt{d}\sqrt{a+bx^2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{ab\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{dx\sqrt{a+bx^2}(bc-2ad)}{ab^2\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2}(bc-2ad)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)}{ab^2\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $-d*(-2*a*d+b*c)*x*(b*x^2+a)^{(1/2)}/a/b^2/(d*x^2+c)^{(1/2)+c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a/b/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+(-2*a*d+b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a/b^2/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+(-a*d+b*c)*x*(d*x^2+c)^{(1/2)}/a/b/(b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.16, antiderivative size = 267, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {413, 531, 418, 492, 411}

$$\frac{dx\sqrt{a+bx^2}(bc-2ad)}{ab^2\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2}(bc-2ad)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)\left[1-\frac{bc}{ad}\right]}{ab^2\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{c^{3/2}\sqrt{d}\sqrt{a+bx^2}F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)}{ab\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] `Int[(c + d*x^2)^(3/2)/(a + b*x^2)^(3/2), x]`

[Out] $-((d*(b*c - 2*a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(a*b^2*\operatorname{Sqrt}[c + d*x^2])) + ((b*c - a*d)*x*\operatorname{Sqrt}[c + d*x^2])/(a*b*\operatorname{Sqrt}[a + b*x^2]) + (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*(b*c - 2*a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*b^2*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (c^{(3/2)}*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*b*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

`Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]`

Rule 413

`Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]`

Rule 418

`Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R`

$\int \frac{t[d/c, 2] \sqrt{c + dx^2} \sqrt{(c + bx^2)/(a + dx^2)}}{x} dx$; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 492

$\text{Int}[(x)^2/(\sqrt{(a) + (b)(x)^2})\sqrt{(c) + (d)(x)^2}], x_Symbol] \rightarrow \text{Simp}[(x\sqrt{a + bx^2})/(b\sqrt{c + dx^2}), x] - \text{Dist}[c/b, \text{Int}[\sqrt{a + bx^2}/(c + dx^2)^{3/2}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rule 531

$\text{Int}[(a + (b)(x)^n)^{p((c) + (d)(x)^n)^q}((e) + (f)(x)^n)], x_Symbol] \rightarrow \text{Dist}[e, \text{Int}[(a + bx^n)^p(c + dx^n)^q, x], x] + \text{Dist}[f, \text{Int}[x^n(a + bx^n)^p(c + dx^n)^q, x], x] /;$ FreeQ[{a, b, c, d, e, f, n, p, q}, x]

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^{3/2}}{(a + bx^2)^{3/2}} dx &= \frac{(bc - ad)x\sqrt{c + dx^2}}{ab\sqrt{a + bx^2}} + \frac{\int \frac{acd - d(bc - 2ad)x^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{ab} \\ &= \frac{(bc - ad)x\sqrt{c + dx^2}}{ab\sqrt{a + bx^2}} + \frac{(cd) \int \frac{1}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{b} - \frac{(d(bc - 2ad)) \int \frac{x^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{ab} \\ &= -\frac{d(bc - 2ad)x\sqrt{a + bx^2}}{ab^2\sqrt{c + dx^2}} + \frac{(bc - ad)x\sqrt{c + dx^2}}{ab\sqrt{a + bx^2}} + \frac{c^{3/2}\sqrt{d}\sqrt{a + bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{ab\sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}}\sqrt{c + dx^2}} \\ &= -\frac{d(bc - 2ad)x\sqrt{a + bx^2}}{ab^2\sqrt{c + dx^2}} + \frac{(bc - ad)x\sqrt{c + dx^2}}{ab\sqrt{a + bx^2}} + \frac{\sqrt{c}\sqrt{d}(bc - 2ad)\sqrt{a + bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{ab^2\sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}}\sqrt{c + dx^2}} \end{aligned}$$

Mathematica [C] time = 0.30, size = 191, normalized size = 0.72

$$\frac{(bc - ad) \left(x\sqrt{\frac{b}{a}}(c + dx^2) - ic\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1} \text{EllipticF}\left(i \sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) \right) - ic\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}(2ad)}{a^2 \left(\frac{b}{a}\right)^{3/2} \sqrt{a + bx^2} \sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + dx^2)^(3/2)/(a + bx^2)^(3/2), x]

[Out] ((-I)*c*(-(b*c) + 2*a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + (b*c - a*d)*(Sqrt[b/a]*x*(c + d*x^2) - I*c*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)))/(a^2*(b/a)^(3/2)*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{bx^2 + a} (dx^2 + c)^{\frac{3}{2}}}{b^2x^4 + 2abx^2 + a^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)/(b^2*x^4 + 2*a*b*x^2 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((d*x^2 + c)^(3/2)/(b*x^2 + a)^(3/2), x)

maple [A] time = 0.04, size = 332, normalized size = 1.24

$$\left(-\sqrt{-\frac{b}{a}} a d^2 x^3 + \sqrt{-\frac{b}{a}} b c d x^3 - \sqrt{-\frac{b}{a}} a c d x + 2\sqrt{\frac{b x^2 + a}{a}} \sqrt{\frac{d x^2 + c}{c}} a c d \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{a d}{b c}}\right) - \sqrt{\frac{b x^2 + a}{a}} \sqrt{\frac{d x^2 + c}{c}}\right) a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^(3/2)/(b*x^2+a)^(3/2),x)

[Out] $(-(-1/a*b)^{(1/2)}*a*d^2*x^3+(-1/a*b)^{(1/2)}*b*c*d*x^3-a*c*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticF}((-1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*d+((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticF}((-1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*b*c^2+2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticE}((-1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*a*c*d-((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticE}((-1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*b*c^2-(-1/a*b)^{(1/2)}*a*c*d*x+x*b*c^2*(-1/a*b)^{(1/2)}*((d*x^2+c)^{(1/2)}*(b*x^2+a)^{(1/2)}/b/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/a/(-1/a*b)^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((d*x^2 + c)^(3/2)/(b*x^2 + a)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(d x^2 + c)^{3/2}}{(b x^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^(3/2)/(a + b*x^2)^(3/2),x)

[Out] int((c + d*x^2)^(3/2)/(a + b*x^2)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + dx^2)^{\frac{3}{2}}}{(a + bx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**(3/2)/(b*x**2+a)**(3/2), x)

[Out] Integral((c + d*x**2)**(3/2)/(a + b*x**2)**(3/2), x)

$$3.176 \quad \int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^{5/2}} dx$$

Optimal. Leaf size=229

$$\frac{c^{3/2}\sqrt{d}\sqrt{a+bx^2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)+2\sqrt{c+dx^2}(ad+bc)E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|1-\frac{ad}{bc}\right)+x\sqrt{c+dx^2}(bc-ad)}{3a^2b\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}+3a^{3/2}b^{3/2}\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}+3ab(a+bx^2)^{3/2}}$$

[Out] $-1/3*c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a^2/b/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/3*(-a*d+b*c)*x*(d*x^2+c)^{(1/2)}/a/b/(b*x^2+a)^{(3/2)}+2/3*(a*d+b*c)*(1/(1+b*x^2/a))^{(1/2)}*(1+b*x^2/a)^{(1/2)}*\operatorname{EllipticE}(x*b^{(1/2)}/a^{(1/2)}/(1+b*x^2/a)^{(1/2)},(1-a*d/b/c)^{(1/2)})*(d*x^2+c)^{(1/2)}/a^{(3/2)}/b^{(3/2)}/(b*x^2+a)^{(1/2)}/(a*(d*x^2+c)/c/(b*x^2+a))^{(1/2)}$

Rubi [A] time = 0.14, antiderivative size = 229, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {413, 525, 418, 411}

$$\frac{2\sqrt{c+dx^2}(ad+bc)E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|1-\frac{ad}{bc}\right)+c^{3/2}\sqrt{d}\sqrt{a+bx^2}F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right)+x\sqrt{c+dx^2}(bc-ad)}{3a^{3/2}b^{3/2}\sqrt{a+bx^2}\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}+3a^2b\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}+3ab(a+bx^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^(3/2)/(a + b*x^2)^(5/2), x]

[Out] $((b*c - a*d)*x*\operatorname{Sqrt}[c + d*x^2])/(3*a*b*(a + b*x^2)^{(3/2)}) + (2*(b*c + a*d)*\operatorname{Sqrt}[c + d*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]], 1 - (a*d)/(b*c)])/(3*a^{(3/2)}*b^{(3/2)}*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[(a*(c + d*x^2))/(c*(a + b*x^2))]) - (c^{(3/2)}*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(3*a^2*b*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_)*(x_)^2]/((c_) + (d_)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 413

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre

$eQ[\{a, b, c, d\}, x] \ \&\& \ \text{PosQ}[d/c] \ \&\& \ \text{PosQ}[b/a] \ \&\& \ \text{!SimplerSqrtQ}[b/a, d/c]$

Rule 525

$\text{Int}[(e_ + (f_)*(x_)^2)/(\text{Sqrt}[(a_ + (b_)*(x_)^2]*((c_ + (d_)*(x_)^2)^{3/2})], x_Symbol] \rightarrow \text{Dist}[(b*e - a*f)/(b*c - a*d), \text{Int}[1/(\text{Sqrt}[a + b*x^2]*\text{Sqrt}[c + d*x^2]), x], x] - \text{Dist}[(d*e - c*f)/(b*c - a*d), \text{Int}[\text{Sqrt}[a + b*x^2]/(c + d*x^2)^{3/2}, x], x] \ /; \ \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{PosQ}[b/a] \ \&\& \ \text{PosQ}[d/c]$

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^{3/2}}{(a + bx^2)^{5/2}} dx &= \frac{(bc - ad)x\sqrt{c + dx^2}}{3ab(a + bx^2)^{3/2}} + \frac{\int \frac{c(2bc + ad) + d(bc + 2ad)x^2}{(a + bx^2)^{3/2}\sqrt{c + dx^2}} dx}{3ab} \\ &= \frac{(bc - ad)x\sqrt{c + dx^2}}{3ab(a + bx^2)^{3/2}} - \frac{(cd) \int \frac{1}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{3ab} + \frac{(2(bc + ad)) \int \frac{\sqrt{c + dx^2}}{(a + bx^2)^{3/2}} dx}{3ab} \\ &= \frac{(bc - ad)x\sqrt{c + dx^2}}{3ab(a + bx^2)^{3/2}} + \frac{2(bc + ad)\sqrt{c + dx^2} E\left(\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| 1 - \frac{ad}{bc}\right)}{3a^{3/2}b^{3/2}\sqrt{a + bx^2}\sqrt{\frac{a(c + dx^2)}{c(a + bx^2)}}} - \frac{c^{3/2}\sqrt{d}\sqrt{a + bx^2} F\left(\frac{c(a + bx^2)}{a(c + dx^2)}\right)}{3a^2b\sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}}} \end{aligned}$$

Mathematica [C] time = 0.48, size = 232, normalized size = 1.01

$$\frac{-ic(a + bx^2)\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}(ad + 2bc)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) + x\sqrt{\frac{b}{a}}(c + dx^2)(a^2d + ab(3c + 2d))}{3a^3\left(\frac{b}{a}\right)^{3/2}(a + bx^2)^{3/2}\sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c + d*x^2)^(3/2)/(a + b*x^2)^(5/2), x]

[Out] (Sqrt[b/a]*x*(c + d*x^2)*(a^2*d + 2*b^2*c*x^2 + a*b*(3*c + 2*d*x^2)) + (2*I)*c*(b*c + a*d)*(a + b*x^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*c*(2*b*c + a*d)*(a + b*x^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)))/(3*a^3*(b/a)^(3/2)*(a + b*x^2)^(3/2)*Sqrt[c + d*x^2])

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2 + a}(dx^2 + c)^{\frac{3}{2}}}{b^3x^6 + 3ab^2x^4 + 3a^2bx^2 + a^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)/(b^3*x^6 + 3*a*b^2*x^4 + 3*a^2*b*x^2 + a^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(5/2),x, algorithm="giac")

[Out] integrate((d*x^2 + c)^(3/2)/(b*x^2 + a)^(5/2), x)

maple [B] time = 0.05, size = 607, normalized size = 2.65

$$2\sqrt{-\frac{b}{a}} ab d^2 x^5 + 2\sqrt{-\frac{b}{a}} b^2 cd x^5 + \sqrt{-\frac{b}{a}} a^2 d^2 x^3 + 5\sqrt{-\frac{b}{a}} abcd x^3 - 2\sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} abcd x^2 \text{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{bx^2+a}{a}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^(3/2)/(b*x^2+a)^(5/2),x)

[Out] $\frac{1}{3} * (2 * (-1/a*b)^{(1/2)} * a*b*d^2*x^5 + 2 * (-1/a*b)^{(1/2)} * b^2*c*d*x^5 + \text{EllipticF}((-1/a*b)^{(1/2)} * x, (a/b/c*d)^{(1/2)}) * x^2 * a*b*c*d * ((b*x^2+a)/a)^{(1/2)} * ((d*x^2+c)/c)^{(1/2)} + 2 * \text{EllipticF}((-1/a*b)^{(1/2)} * x, (a/b/c*d)^{(1/2)}) * x^2 * b^2*c^2 * ((b*x^2+a)/a)^{(1/2)} * ((d*x^2+c)/c)^{(1/2)} - 2 * \text{EllipticE}((-1/a*b)^{(1/2)} * x, (a/b/c*d)^{(1/2)}) * x^2 * a*b*c*d * ((b*x^2+a)/a)^{(1/2)} * ((d*x^2+c)/c)^{(1/2)} - 2 * \text{EllipticE}((-1/a*b)^{(1/2)} * x, (a/b/c*d)^{(1/2)}) * x^2 * b^2*c^2 * ((b*x^2+a)/a)^{(1/2)} * ((d*x^2+c)/c)^{(1/2)} + (-1/a*b)^{(1/2)} * a^2*d^2*x^3 + 5 * (-1/a*b)^{(1/2)} * a*b*c*d*x^3 + 2 * (-1/a*b)^{(1/2)} * b^2*c^2*x^3 + \text{EllipticF}((-1/a*b)^{(1/2)} * x, (a/b/c*d)^{(1/2)}) * a^2*c*d * ((b*x^2+a)/a)^{(1/2)} * ((d*x^2+c)/c)^{(1/2)} + 2 * \text{EllipticF}((-1/a*b)^{(1/2)} * x, (a/b/c*d)^{(1/2)}) * a*b*c^2 * ((b*x^2+a)/a)^{(1/2)} * ((d*x^2+c)/c)^{(1/2)} - 2 * \text{EllipticE}((-1/a*b)^{(1/2)} * x, (a/b/c*d)^{(1/2)}) * a^2*c*d * ((b*x^2+a)/a)^{(1/2)} * ((d*x^2+c)/c)^{(1/2)} - 2 * \text{EllipticE}((-1/a*b)^{(1/2)} * x, (a/b/c*d)^{(1/2)}) * a*b*c^2 * ((b*x^2+a)/a)^{(1/2)} * ((d*x^2+c)/c)^{(1/2)} + (-1/a*b)^{(1/2)} * a^2*c*d*x + 3 * (-1/a*b)^{(1/2)} * a*b*c^2*x) / (d*x^2+c)^{(1/2)} / a^2 / (-1/a*b)^{(1/2)} / (b*x^2+a)^{(3/2)} / b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(5/2),x, algorithm="maxima")

[Out] integrate((d*x^2 + c)^(3/2)/(b*x^2 + a)^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^(3/2)/(a + b*x^2)^(5/2),x)

[Out] int((c + d*x^2)^(3/2)/(a + b*x^2)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c + dx^2)^{\frac{3}{2}}}{(a + bx^2)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**(3/2)/(b*x**2+a)**(5/2), x)

[Out] Integral((c + d*x**2)**(3/2)/(a + b*x**2)**(5/2), x)

$$3.177 \quad \int \frac{(c+dx^2)^{3/2}}{(a+bx^2)^{7/2}} dx$$

Optimal. Leaf size=315

$$\frac{c^{3/2}\sqrt{d}\sqrt{a+bx^2}(4bc-ad)\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right) + 2x\sqrt{c+dx^2}(ad+2bc) + \sqrt{c+dx^2}(-2a^2d^2-3abcd)}{15a^3b\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} + 15a^2b(a+bx^2)^{3/2} + 15a^{5/2}b^{3/2}\sqrt{a+bx^2}}$$

[Out] $-1/15*c^{(3/2)}*(-a*d+4*b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a^3/b/(-a*d+b*c)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/5*(-a*d+b*c)*x*(d*x^2+c)^{(1/2)}/a/b/(b*x^2+a)^{(5/2)}+2/15*(a*d+2*b*c)*x*(d*x^2+c)^{(1/2)}/a^2/b/(b*x^2+a)^{(3/2)}+1/15*(-2*a^2*d^2-3*a*b*c*d+8*b^2*c^2)*(1/(1+b*x^2/a))^{(1/2)}*(1+b*x^2/a)^{(1/2)}*\operatorname{EllipticE}(x*b^{(1/2)}/a^{(1/2)}/(1+b*x^2/a)^{(1/2)}, (1-a*d/b/c)^{(1/2)}*(d*x^2+c)^{(1/2)}/a^{(5/2)}/b^{(3/2)}/(-a*d+b*c)/(b*x^2+a)^{(1/2)}/(a*(d*x^2+c)/c/(b*x^2+a))^{(1/2)}$

Rubi [A] time = 0.28, antiderivative size = 315, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {413, 527, 525, 418, 411}

$$\frac{\sqrt{c+dx^2}(-2a^2d^2-3abcd+8b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\left|1-\frac{ad}{bc}\right.\right) + c^{3/2}\sqrt{d}\sqrt{a+bx^2}(4bc-ad)F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{15a^{5/2}b^{3/2}\sqrt{a+bx^2}(bc-ad)\sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}} + 15a^3b\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[(c + d*x^2)^(3/2)/(a + b*x^2)^(7/2), x]

[Out] $((b*c - a*d)*x*\operatorname{Sqrt}[c + d*x^2])/(5*a*b*(a + b*x^2)^{(5/2)}) + (2*(2*b*c + a*d)*x*\operatorname{Sqrt}[c + d*x^2])/(15*a^2*b*(a + b*x^2)^{(3/2)}) + ((8*b^2*c^2 - 3*a*b*c*d - 2*a^2*d^2)*\operatorname{Sqrt}[c + d*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]], 1 - (a*d)/(b*c)])/(15*a^{(5/2)}*b^{(3/2)}*(b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[(a*(c + d*x^2))/(c*(a + b*x^2))]) - (c^{(3/2)}*\operatorname{Sqrt}[d]*(4*b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(15*a^3*b*(b*c - a*d)*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_)*(x_)^2]/((c_) + (d_)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 413

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] :> Simp[((a*d - c*b)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(a*b*n*(p+1)), x] - Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-2)*Simp[c*(a*d - c*b*(n*(p+1)+1)) + d*(a*d*(n*(q-1)+1) - b*c*(n*(p+q)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)]/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 525

```
Int[((e_) + (f_.)*(x_)^2)/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)^(
3/2)), x_Symbol] :> Dist[(b*e - a*f)/(b*c - a*d), Int[1/(Sqrt[a + b*x^2]*S
qrt[c + d*x^2]), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[Sqrt[a + b*x^2]
/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && PosQ[b/a] &&
PosQ[d/c]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned} \int \frac{(c + dx^2)^{3/2}}{(a + bx^2)^{7/2}} dx &= \frac{(bc - ad)x\sqrt{c + dx^2}}{5ab(a + bx^2)^{5/2}} + \frac{\int \frac{c(4bc + ad) + d(3bc + 2ad)x^2}{(a + bx^2)^{5/2}\sqrt{c + dx^2}} dx}{5ab} \\ &= \frac{(bc - ad)x\sqrt{c + dx^2}}{5ab(a + bx^2)^{5/2}} + \frac{2(2bc + ad)x\sqrt{c + dx^2}}{15a^2b(a + bx^2)^{3/2}} - \frac{\int \frac{-c(bc - ad)(8bc + ad) - 2d(bc - ad)(2bc + ad)x^2}{(a + bx^2)^{3/2}\sqrt{c + dx^2}} dx}{15a^2b(bc - ad)} \\ &= \frac{(bc - ad)x\sqrt{c + dx^2}}{5ab(a + bx^2)^{5/2}} + \frac{2(2bc + ad)x\sqrt{c + dx^2}}{15a^2b(a + bx^2)^{3/2}} - \frac{(cd(4bc - ad)) \int \frac{1}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{15a^2b(bc - ad)} + \frac{(8b^2c^2 - 3abcd - 2a^2d^2)\sqrt{c + dx^2} E(\tan^{-1}(\frac{\sqrt{b}x}{\sqrt{a + bx^2}}))}{15a^{5/2}b^{3/2}(bc - ad)\sqrt{a + bx^2}} \end{aligned}$$

Mathematica [C] time = 0.60, size = 285, normalized size = 0.90

$$\frac{x\sqrt{\frac{b}{a}}(c + dx^2)\left((a + bx^2)^2(-2a^2d^2 - 3abcd + 8b^2c^2) + 3a^2(bc - ad)^2 + 2a(a + bx^2)(ad + 2bc)(bc - ad)\right) - ic}{15a^4}$$

Antiderivative was successfully verified.

```
[In] Integrate[(c + d*x^2)^(3/2)/(a + b*x^2)^(7/2), x]
```

```
[Out] (Sqrt[b/a]*x*(c + d*x^2)*(3*a^2*(b*c - a*d)^2 + 2*a*(b*c - a*d)*(2*b*c + a*
d)*(a + b*x^2) + (8*b^2*c^2 - 3*a*b*c*d - 2*a^2*d^2)*(a + b*x^2)^2) - I*c*(
a + b*x^2)^2*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*((-8*b^2*c^2 + 3*a*b*c
*d + 2*a^2*d^2)*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + (8*b^2*c^2
```

- 7*a*b*c*d - a^2*d^2)*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)))/(15*a^4*(b/a)^(3/2)*(b*c - a*d)*(a + b*x^2)^(5/2)*Sqrt[c + d*x^2])

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{bx^2 + a} (dx^2 + c)^{\frac{3}{2}}}{b^4x^8 + 4ab^3x^6 + 6a^2b^2x^4 + 4a^3bx^2 + a^4}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(7/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*(d*x^2 + c)^(3/2)/(b^4*x^8 + 4*a*b^3*x^6 + 6*a^2*b^2*x^4 + 4*a^3*b*x^2 + a^4), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(7/2),x, algorithm="giac")

[Out] integrate((d*x^2 + c)^(3/2)/(b*x^2 + a)^(7/2), x)

maple [B] time = 0.05, size = 1410, normalized size = 4.48

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^(3/2)/(b*x^2+a)^(7/2),x)

[Out] 1/15*(-2*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^4*c*d^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+8*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b^2*c^3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^4*c*d^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-8*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b^2*c^3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+10*(-1/a*b)^(1/2)*a^2*b^2*c*d^2*x^5-17*(-1/a*b)^(1/2)*a*b^3*c^2*d*x^5+17*(-1/a*b)^(1/2)*a^3*b*c*d^2*x^3-7*(-1/a*b)^(1/2)*a^2*b^2*c^2*d*x^3+11*(-1/a*b)^(1/2)*a^3*b*c^2*d*x+8*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^4*b^4*c^3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-8*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^4*b^4*c^3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-8*(-1/a*b)^(1/2)*b^4*c^3*x^5+(-1/a*b)^(1/2)*a^4*d^3*x^3+3*(-1/a*b)^(1/2)*a*b^3*c*d^2*x^7-3*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^4*a*b^3*c^2*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+2*(-1/a*b)^(1/2)*a^2*b^2*d^3*x^7-2*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^4*a^2*b^2*c*d^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+14*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a^2*b^2*c^2*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-4*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a^3*b*c*d^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+7*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^4*a*b^3*c^2*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^4*a^2*b^2*c*d^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+2*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a^3*b*c*d^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-6*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a^2*b^2*c^2*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+x*a^4*c*d^2*(-1/a*b)^(1/2)+16*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a*b^3*c^3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-16*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))

/2))*x^2*a*b^3*c^3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-3*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^3*b*c^2*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+7*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^3*b*c^2*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-8*(-1/a*b)^(1/2)*b^4*c^2*d*x^7+6*(-1/a*b)^(1/2)*a^3*b*d^3*x^5-20*(-1/a*b)^(1/2)*a*b^3*c^3*x^3-15*(-1/a*b)^(1/2)*a^2*b^2*c^3*x)/(d*x^2+c)^(1/2)/a^3/(a*d-b*c)/(-1/a*b)^(1/2)/(b*x^2+a)^(5/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(dx^2 + c)^{\frac{3}{2}}}{(bx^2 + a)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(3/2)/(b*x^2+a)^(7/2),x, algorithm="maxima")

[Out] integrate((d*x^2 + c)^(3/2)/(b*x^2 + a)^(7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(dx^2 + c)^{3/2}}{(bx^2 + a)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c + d*x^2)^(3/2)/(a + b*x^2)^(7/2),x)

[Out] int((c + d*x^2)^(3/2)/(a + b*x^2)^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**(3/2)/(b*x**2+a)**(7/2),x)

[Out] Timed out

3.178 $\int \sqrt{2 + bx^2} \sqrt{3 + dx^2} dx$

Optimal. Leaf size=235

$$\frac{2\sqrt{2}\sqrt{bx^2+2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right), 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}} + \frac{1}{3}x\sqrt{bx^2+2}\sqrt{dx^2+3} + \frac{x(3b+2d)\sqrt{bx^2+2}}{3b\sqrt{dx^2+3}} - \frac{\sqrt{2}(3b+2d)\sqrt{bx^2+2}}{3b\sqrt{d}}$$

[Out] $\frac{1}{3}*(3*b+2*d)*x*(b*x^2+2)^{(1/2)}/b/(d*x^2+3)^{(1/2)} - \frac{1}{3}*(3*b+2*d)*(1/(3*d*x^2+9))^{(1/2)}*(3*d*x^2+9)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}*3^{(1/2)/(3*d*x^2+9)^{(1/2)}, 1/2*(4-6*b/d)^{(1/2)})^2^{(1/2)}*(b*x^2+2)^{(1/2)}/b/d^{(1/2)}/((b*x^2+2)/(d*x^2+3))^{(1/2)}/(d*x^2+3)^{(1/2)} + 2*(1/(3*d*x^2+9))^{(1/2)}*(3*d*x^2+9)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}*3^{(1/2)/(3*d*x^2+9)^{(1/2)}, 1/2*(4-6*b/d)^{(1/2)})^2^{(1/2)}*(b*x^2+2)^{(1/2)}/d^{(1/2)}/((b*x^2+2)/(d*x^2+3))^{(1/2)}/(d*x^2+3)^{(1/2)} + 1/3*x*(b*x^2+2)^{(1/2)}*(d*x^2+3)^{(1/2)}$

Rubi [A] time = 0.13, antiderivative size = 235, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {417, 531, 418, 492, 411}

$$\frac{1}{3}x\sqrt{bx^2+2}\sqrt{dx^2+3} + \frac{x(3b+2d)\sqrt{bx^2+2}}{3b\sqrt{dx^2+3}} + \frac{2\sqrt{2}\sqrt{bx^2+2}F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right)\middle|1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}} - \frac{\sqrt{2}(3b+2d)\sqrt{bx^2+2}E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right)\middle|1 - \frac{3b}{2d}\right)}{3b\sqrt{d}\sqrt{dx^2+3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[2 + b*x^2]*Sqrt[3 + d*x^2], x]

[Out] $((3*b + 2*d)*x*\operatorname{Sqrt}[2 + b*x^2])/(3*b*\operatorname{Sqrt}[3 + d*x^2]) + (x*\operatorname{Sqrt}[2 + b*x^2]*\operatorname{Sqrt}[3 + d*x^2])/3 - (\operatorname{Sqrt}[2]*(3*b + 2*d)*\operatorname{Sqrt}[2 + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[3]], 1 - (3*b)/(2*d)])/(3*b*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(2 + b*x^2)/(3 + d*x^2)]*\operatorname{Sqrt}[3 + d*x^2]) + (2*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[2 + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[3]], 1 - (3*b)/(2*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(2 + b*x^2)/(3 + d*x^2)]*\operatorname{Sqrt}[3 + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 417

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(x*(a + b*x^n)^p*(c + d*x^n)^q)/(n*(p + q) + 1), x] + Dist[n/(n*(p + q) + 1), Int[(a + b*x^n)^(p - 1)*(c + d*x^n)^(q - 1)*Simp[a*c*(p + q) + (q*(b*c - a*d) + a*d*(p + q))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 0] && GtQ[p, 0] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 492


```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
  := Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
  + b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
  a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rule 531

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
  f_)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x],
  x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c,
  d, e, f, n, p, q}, x]
```

Rubi steps

$$\begin{aligned} \int \sqrt{2+bx^2} \sqrt{3+dx^2} dx &= \frac{1}{3}x\sqrt{2+bx^2} \sqrt{3+dx^2} + \frac{2}{3} \int \frac{6 + \frac{1}{2}(3b+2d)x^2}{\sqrt{2+bx^2} \sqrt{3+dx^2}} dx \\ &= \frac{1}{3}x\sqrt{2+bx^2} \sqrt{3+dx^2} + 4 \int \frac{1}{\sqrt{2+bx^2} \sqrt{3+dx^2}} dx + \frac{1}{3}(3b+2d) \int \frac{x^2}{\sqrt{2+bx^2} \sqrt{3+dx^2}} dx \\ &= \frac{(3b+2d)x\sqrt{2+bx^2}}{3b\sqrt{3+dx^2}} + \frac{1}{3}x\sqrt{2+bx^2} \sqrt{3+dx^2} + \frac{2\sqrt{2}\sqrt{2+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right)\right)}{\sqrt{d}\sqrt{\frac{2+bx^2}{3+dx^2}}\sqrt{3+dx^2}} \\ &= \frac{(3b+2d)x\sqrt{2+bx^2}}{3b\sqrt{3+dx^2}} + \frac{1}{3}x\sqrt{2+bx^2} \sqrt{3+dx^2} - \frac{\sqrt{2}(3b+2d)\sqrt{2+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right)\right)}{3b\sqrt{d}\sqrt{\frac{2+bx^2}{3+dx^2}}\sqrt{3+dx^2}} \end{aligned}$$

Mathematica [C] time = 0.11, size = 127, normalized size = 0.54

$$\frac{i\sqrt{3}(3b-2d) \operatorname{EllipticF}\left(i \sinh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}}\right), \frac{2d}{3b}\right) + \sqrt{b} dx \sqrt{bx^2+2} \sqrt{dx^2+3} - i\sqrt{3}(3b+2d) E\left(i \sinh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}}\right) \middle| \frac{2d}{3b}\right)}{3\sqrt{bd}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[2 + b*x^2]*Sqrt[3 + d*x^2], x]
```

```
[Out] (Sqrt[b]*d*x*Sqrt[2 + b*x^2]*Sqrt[3 + d*x^2] - I*Sqrt[3]*(3*b + 2*d)*EllipticE[I*ArcSinh[(Sqrt[b]*x)/Sqrt[2]], (2*d)/(3*b)] + I*Sqrt[3]*(3*b - 2*d)*EllipticF[I*ArcSinh[(Sqrt[b]*x)/Sqrt[2]], (2*d)/(3*b)])/(3*Sqrt[b]*d)
```

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\sqrt{bx^2+2} \sqrt{dx^2+3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+2)^(1/2)*(d*x^2+3)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*x^2 + 2)*sqrt(d*x^2 + 3), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{bx^2+2} \sqrt{dx^2+3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+2)^(1/2)*(d*x^2+3)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 + 2)*sqrt(d*x^2 + 3), x)

maple [A] time = 0.04, size = 303, normalized size = 1.29

$$\sqrt{bx^2 + 2} \sqrt{dx^2 + 3} \left(\sqrt{-d} b^2 dx^5 + 3\sqrt{-d} b^2 x^3 + 2\sqrt{-d} b dx^3 + 6\sqrt{-d} bx + 3\sqrt{2} \sqrt{bx^2 + 2} \sqrt{dx^2 + 3} b \text{Elliptic} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+2)^(1/2)*(d*x^2+3)^(1/2),x)

[Out] 1/3*(b*x^2+2)^(1/2)*(d*x^2+3)^(1/2)*(x^5*b^2*d*(-d)^(1/2)+3*x^3*b^2*(-d)^(1/2)+2*x^3*b*d*(-d)^(1/2)+3*2^(1/2)*EllipticF(1/3*x*3^(1/2)*(-d)^(1/2),1/2*2^(1/2)*3^(1/2)*(1/d*b)^(1/2))*b*(b*x^2+2)^(1/2)*(d*x^2+3)^(1/2)-2*2^(1/2)*EllipticF(1/3*x*3^(1/2)*(-d)^(1/2),1/2*2^(1/2)*3^(1/2)*(1/d*b)^(1/2))*d*(b*x^2+2)^(1/2)*(d*x^2+3)^(1/2)+3*2^(1/2)*EllipticE(1/3*x*3^(1/2)*(-d)^(1/2),1/2*2^(1/2)*3^(1/2)*(1/d*b)^(1/2))*b*(b*x^2+2)^(1/2)*(d*x^2+3)^(1/2)+2*2^(1/2)*EllipticE(1/3*x*3^(1/2)*(-d)^(1/2),1/2*2^(1/2)*3^(1/2)*(1/d*b)^(1/2))*d*(b*x^2+2)^(1/2)*(d*x^2+3)^(1/2)+6*x*b*(-d)^(1/2))/(b*d*x^4+3*b*x^2+2*d*x^2+6)/(-d)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{bx^2 + 2} \sqrt{dx^2 + 3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+2)^(1/2)*(d*x^2+3)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + 2)*sqrt(d*x^2 + 3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{bx^2 + 2} \sqrt{dx^2 + 3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2 + 2)^(1/2)*(d*x^2 + 3)^(1/2),x)

[Out] int((b*x^2 + 2)^(1/2)*(d*x^2 + 3)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{bx^2 + 2} \sqrt{dx^2 + 3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+2)**(1/2)*(d*x**2+3)**(1/2),x)

[Out] Integral(sqrt(b*x**2 + 2)*sqrt(d*x**2 + 3), x)

$$3.179 \quad \int \sqrt{3 - 6x^2} \sqrt{2 + 4x^2} dx$$

Optimal. Leaf size=38

$$\frac{2 \operatorname{EllipticF}(\sin^{-1}(\sqrt{2}x), -1)}{\sqrt{3}} + \sqrt{\frac{2}{3}} \sqrt{1 - 4x^4} x$$

[Out] $2/3 * \operatorname{EllipticF}(x * 2^{(1/2)}, 1) * 3^{(1/2)} + 1/3 * x * 6^{(1/2)} * (-4 * x^4 + 1)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {248, 195, 221}

$$\sqrt{\frac{2}{3}} \sqrt{1 - 4x^4} x + \frac{2F(\sin^{-1}(\sqrt{2}x) | -1)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[3 - 6*x^2]*Sqrt[2 + 4*x^2], x]

[Out] Sqrt[2/3]*x*Sqrt[1 - 4*x^4] + (2*EllipticF[ArcSin[Sqrt[2]*x], -1])/Sqrt[3]

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] :> Simp[EllipticF[ArcSin[(Rt[-b, 4]*x)/Rt[a, 4]], -1]/(Rt[a, 4]*Rt[-b, 4]), x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]

Rule 248

Int[((a1_.) + (b1_.)*(x_)^(n_))^(p_.)*((a2_.) + (b2_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rubi steps

$$\begin{aligned} \int \sqrt{3 - 6x^2} \sqrt{2 + 4x^2} dx &= \int \sqrt{6 - 24x^4} dx \\ &= \sqrt{\frac{2}{3}} x \sqrt{1 - 4x^4} + 4 \int \frac{1}{\sqrt{6 - 24x^4}} dx \\ &= \sqrt{\frac{2}{3}} x \sqrt{1 - 4x^4} + \frac{2F(\sin^{-1}(\sqrt{2}x) | -1)}{\sqrt{3}} \end{aligned}$$

Mathematica [C] time = 0.01, size = 22, normalized size = 0.58

$$\sqrt{6} x {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{5}{4}; 4x^4\right)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[3 - 6*x^2]*Sqrt[2 + 4*x^2],x]

[Out] Sqrt[6]*x*Hypergeometric2F1[-1/2, 1/4, 5/4, 4*x^4]

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{4x^2+2}\sqrt{-6x^2+3},x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-6*x^2+3)^(1/2)*(4*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(4*x^2 + 2)*sqrt(-6*x^2 + 3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{4x^2+2}\sqrt{-6x^2+3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-6*x^2+3)^(1/2)*(4*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(4*x^2 + 2)*sqrt(-6*x^2 + 3), x)

maple [B] time = 0.04, size = 75, normalized size = 1.97

$$\frac{\sqrt{-6x^2+3}\sqrt{2}\sqrt{2x^2+1}\left(-12x^5+3x+\sqrt{2}\sqrt{3}\sqrt{-6x^2+3}\sqrt{2x^2+1}\text{EllipticF}\left(\sqrt{2}x,i\right)\right)}{9\left(4x^4-1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-6*x^2+3)^(1/2)*(4*x^2+2)^(1/2),x)

[Out] -1/9*(-6*x^2+3)^(1/2)*2^(1/2)*(2*x^2+1)^(1/2)*(2^(1/2)*3^(1/2)*(-6*x^2+3)^(1/2)*(2*x^2+1)^(1/2)*EllipticF(x*2^(1/2),I)-12*x^5+3*x)/(4*x^4-1)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{4x^2+2}\sqrt{-6x^2+3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-6*x^2+3)^(1/2)*(4*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(4*x^2 + 2)*sqrt(-6*x^2 + 3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \sqrt{4x^2+2}\sqrt{3-6x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4*x^2 + 2)^(1/2)*(3 - 6*x^2)^(1/2),x)

[Out] int((4*x^2 + 2)^(1/2)*(3 - 6*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\sqrt{6}\int\sqrt{1-2x^2}\sqrt{2x^2+1}dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-6*x**2+3)**(1/2)*(4*x**2+2)**(1/2),x)
```

```
[Out] sqrt(6)*Integral(sqrt(1 - 2*x**2)*sqrt(2*x**2 + 1), x)
```

$$3.180 \quad \int \sqrt{2 + 4x^2} \sqrt{3 + 6x^2} dx$$

Optimal. Leaf size=20

$$2\sqrt{\frac{2}{3}}x^3 + \sqrt{6}x$$

[Out] $2/3*x^3*6^{(1/2)}+x*6^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {22}

$$2\sqrt{\frac{2}{3}}x^3 + \sqrt{6}x$$

Antiderivative was successfully verified.

[In] Int[Sqrt[2 + 4*x^2]*Sqrt[3 + 6*x^2],x]

[Out] Sqrt[6]*x + 2*Sqrt[2/3]*x^3

Rule 22

Int[(u_.)*((a_) + (b_.)*(v_))^(m_)*((c_) + (d_.)*(v_))^(n_), x_Symbol] :> Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[b*c - a*d, 0] && GtQ[b/d, 0] && !(IntegerQ[m] || IntegerQ[n])

Rubi steps

$$\begin{aligned} \int \sqrt{2 + 4x^2} \sqrt{3 + 6x^2} dx &= \sqrt{\frac{2}{3}} \int (3 + 6x^2) dx \\ &= \sqrt{6}x + 2\sqrt{\frac{2}{3}}x^3 \end{aligned}$$

Mathematica [A] time = 0.00, size = 15, normalized size = 0.75

$$\sqrt{6} \left(\frac{2x^3}{3} + x \right)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[2 + 4*x^2]*Sqrt[3 + 6*x^2],x]

[Out] Sqrt[6]*(x + (2*x^3)/3)

fricas [B] time = 0.59, size = 38, normalized size = 1.90

$$\frac{(2x^3 + 3x)\sqrt{6x^2 + 3}\sqrt{4x^2 + 2}}{3(2x^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+2)^(1/2)*(6*x^2+3)^(1/2),x, algorithm="fricas")

[Out] 1/3*(2*x^3 + 3*x)*sqrt(6*x^2 + 3)*sqrt(4*x^2 + 2)/(2*x^2 + 1)

giac [A] time = 0.57, size = 17, normalized size = 0.85

$$\frac{1}{3} \sqrt{3} \sqrt{2} (2x^3 + 3x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+2)^(1/2)*(6*x^2+3)^(1/2),x, algorithm="giac")

[Out] 1/3*sqrt(3)*sqrt(2)*(2*x^3 + 3*x)

maple [C] time = 0.00, size = 38, normalized size = 1.90

$$\frac{(2x^2 + 3) \sqrt{4x^2 + 2} x}{\sqrt{6x^2 + 3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4*x^2+2)^(1/2)*(6*x^2+3)^(1/2),x)

[Out] 1/3*x*(2*x^2+3)*(4*x^2+2)^(1/2)*(6*x^2+3)^(1/2)/(2*x^2+1)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{6x^2 + 3} \sqrt{4x^2 + 2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+2)^(1/2)*(6*x^2+3)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(6*x^2 + 3)*sqrt(4*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \sqrt{4x^2 + 2} \sqrt{6x^2 + 3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4*x^2 + 2)^(1/2)*(6*x^2 + 3)^(1/2),x)

[Out] int((4*x^2 + 2)^(1/2)*(6*x^2 + 3)^(1/2), x)

sympy [A] time = 3.43, size = 17, normalized size = 0.85

$$\frac{2\sqrt{6}x^3}{3} + \sqrt{6}x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x**2+2)**(1/2)*(6*x**2+3)**(1/2),x)

[Out] 2*sqrt(6)*x**3/3 + sqrt(6)*x

$$3.181 \quad \int \frac{\sqrt{2+bx^2}}{\sqrt{3+dx^2}} dx$$

Optimal. Leaf size=182

$$\frac{\sqrt{2}\sqrt{bx^2+2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right), 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}} + \frac{x\sqrt{bx^2+2}}{\sqrt{dx^2+3}} - \frac{\sqrt{2}\sqrt{bx^2+2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}}$$

[Out] $x*(b*x^2+2)^{(1/2)}/(d*x^2+3)^{(1/2)} - (1/(3*d*x^2+9))^{(1/2)}*(3*d*x^2+9)^{(1/2)}*E$
 $llipticE(x*d^{(1/2)}*3^{(1/2)}/(3*d*x^2+9)^{(1/2)}, 1/2*(4-6*b/d)^{(1/2)})*2^{(1/2)}*($
 $b*x^2+2)^{(1/2)}/d^{(1/2)}/((b*x^2+2)/(d*x^2+3))^{(1/2)}/(d*x^2+3)^{(1/2)} + (1/(3*d*$
 $x^2+9))^{(1/2)}*(3*d*x^2+9)^{(1/2)}*EllipticF(x*d^{(1/2)}*3^{(1/2)}/(3*d*x^2+9)^{(1/2)}$
 $, 1/2*(4-6*b/d)^{(1/2)})*2^{(1/2)}*(b*x^2+2)^{(1/2)}/d^{(1/2)}/((b*x^2+2)/(d*x^2+3$
 $))^{(1/2)}/(d*x^2+3)^{(1/2)}$

Rubi [A] time = 0.08, antiderivative size = 182, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {422, 418, 492, 411}

$$\frac{x\sqrt{bx^2+2}}{\sqrt{dx^2+3}} + \frac{\sqrt{2}\sqrt{bx^2+2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}} - \frac{\sqrt{2}\sqrt{bx^2+2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[2 + b*x^2]/Sqrt[3 + d*x^2], x]

[Out] $(x*\operatorname{Sqrt}[2 + b*x^2])/ \operatorname{Sqrt}[3 + d*x^2] - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[2 + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[3]], 1 - (3*b)/(2*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(2 + b*x^2)/(3 + d*x^2)]*\operatorname{Sqrt}[3 + d*x^2]) + (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[2 + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[3]], 1 - (3*b)/(2*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(2 + b*x^2)/(3 + d*x^2)]*\operatorname{Sqrt}[3 + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a

$+ b*x^2)/(c + d*x^2)^{(3/2)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{PosQ}[b/a] \ \&\& \ \text{PosQ}[d/c] \ \&\& \ \text{!SimplerSqrtQ}[b/a, d/c]$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{2+bx^2}}{\sqrt{3+dx^2}} dx &= 2 \int \frac{1}{\sqrt{2+bx^2}\sqrt{3+dx^2}} dx + b \int \frac{x^2}{\sqrt{2+bx^2}\sqrt{3+dx^2}} dx \\ &= \frac{x\sqrt{2+bx^2}}{\sqrt{3+dx^2}} + \frac{\sqrt{2}\sqrt{2+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{\frac{2+bx^2}{3+dx^2}}\sqrt{3+dx^2}} - 3 \int \frac{\sqrt{2+bx^2}}{(3+dx^2)^{3/2}} dx \\ &= \frac{x\sqrt{2+bx^2}}{\sqrt{3+dx^2}} - \frac{\sqrt{2}\sqrt{2+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{\frac{2+bx^2}{3+dx^2}}\sqrt{3+dx^2}} + \frac{\sqrt{2}\sqrt{2+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{\frac{2+bx^2}{3+dx^2}}\sqrt{3+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 37, normalized size = 0.20

$$\frac{\sqrt{2} E\left(\sin^{-1}\left(\frac{\sqrt{-d}x}{\sqrt{3}}\right) \middle| \frac{3b}{2d}\right)}{\sqrt{-d}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[2 + b*x^2]/Sqrt[3 + d*x^2], x]

[Out] (Sqrt[2]*EllipticE[ArcSin[(Sqrt[-d]*x)/Sqrt[3]], (3*b)/(2*d)])/Sqrt[-d]

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+2}}{\sqrt{dx^2+3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+2)^(1/2)/(d*x^2+3)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + 2)/sqrt(d*x^2 + 3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+2}}{\sqrt{dx^2+3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+2)^(1/2)/(d*x^2+3)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 + 2)/sqrt(d*x^2 + 3), x)

maple [A] time = 0.03, size = 37, normalized size = 0.20

$$\frac{\sqrt{2} \text{EllipticE}\left(\frac{\sqrt{3}\sqrt{-d}x}{3}, \frac{\sqrt{2}\sqrt{3}\sqrt{\frac{b}{d}}}{2}\right)}{\sqrt{-d}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2+2)^(1/2)/(d*x^2+3)^(1/2),x)`

[Out] `EllipticE(1/3*3^(1/2)*(-d)^(1/2)*x,1/2*2^(1/2)*3^(1/2)*(b/d)^(1/2))*2^(1/2)/(-d)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + 2}}{\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+2)^(1/2)/(d*x^2+3)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*x^2 + 2)/sqrt(d*x^2 + 3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 + 2}}{\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2 + 2)^(1/2)/(d*x^2 + 3)^(1/2),x)`

[Out] `int((b*x^2 + 2)^(1/2)/(d*x^2 + 3)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + 2}}{\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+2)**(1/2)/(d*x**2+3)**(1/2),x)`

[Out] `Integral(sqrt(b*x**2 + 2)/sqrt(d*x**2 + 3), x)`

$$3.182 \quad \int \frac{\sqrt{4-x^2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=91

$$\frac{(c+4d)\sqrt{\frac{dx^2}{c}+1} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{x}{2}\right), -\frac{4d}{c}\right)}{d\sqrt{c+dx^2}} - \frac{\sqrt{c+dx^2} E\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -\frac{4d}{c}\right)}{d\sqrt{\frac{dx^2}{c}+1}}$$

[Out] -EllipticE(1/2*x, 2*(-d/c)^(1/2))*(d*x^2+c)^(1/2)/d/(1+d*x^2/c)^(1/2)+(c+4*d)*EllipticF(1/2*x, 2*(-d/c)^(1/2))*(1+d*x^2/c)^(1/2)/d/(d*x^2+c)^(1/2)

Rubi [A] time = 0.06, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {423, 426, 424, 421, 419}

$$\frac{(c+4d)\sqrt{\frac{dx^2}{c}+1} F\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -\frac{4d}{c}\right)}{d\sqrt{c+dx^2}} - \frac{\sqrt{c+dx^2} E\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -\frac{4d}{c}\right)}{d\sqrt{\frac{dx^2}{c}+1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[4 - x^2]/Sqrt[c + d*x^2], x]

[Out] -((Sqrt[c + d*x^2]*EllipticE[ArcSin[x/2], (-4*d)/c])/(d*Sqrt[1 + (d*x^2)/c])) + ((c + 4*d)*Sqrt[1 + (d*x^2)/c]*EllipticF[ArcSin[x/2], (-4*d)/c])/(d*Sqrt[c + d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{4-x^2}}{\sqrt{c+dx^2}} dx &= -\frac{\int \frac{\sqrt{c+dx^2}}{\sqrt{4-x^2}} dx}{d} - \frac{(-c-4d) \int \frac{1}{\sqrt{4-x^2} \sqrt{c+dx^2}} dx}{d} \\
&= -\frac{\sqrt{c+dx^2} \int \frac{\sqrt{1+\frac{dx^2}{c}}}{\sqrt{4-x^2}} dx}{d\sqrt{1+\frac{dx^2}{c}}} - \frac{\left((-c-4d)\sqrt{1+\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{4-x^2} \sqrt{1+\frac{dx^2}{c}}} dx}{d\sqrt{c+dx^2}} \\
&= -\frac{\sqrt{c+dx^2} E\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -\frac{4d}{c}\right)}{d\sqrt{1+\frac{dx^2}{c}}} + \frac{(c+4d)\sqrt{1+\frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -\frac{4d}{c}\right)}{d\sqrt{c+dx^2}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 60, normalized size = 0.66

$$\frac{2\sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{-\frac{d}{c}}x\right) \middle| -\frac{c}{4d}\right)}{\sqrt{-\frac{d}{c}} \sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[4 - x^2]/Sqrt[c + d*x^2], x]

[Out] (2*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], -1/4*c/d])/(Sqrt[-(d/c)]*Sqrt[c + d*x^2])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+4}}{\sqrt{dx^2+c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+4)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-x^2 + 4)/sqrt(d*x^2 + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+4}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+4)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-x^2 + 4)/sqrt(d*x^2 + c), x)

maple [A] time = 0.03, size = 78, normalized size = 0.86

$$\frac{\left(-c \text{EllipticE}\left(\frac{x}{2}, 2\sqrt{-\frac{d}{c}}\right) + c \text{EllipticF}\left(\frac{x}{2}, 2\sqrt{-\frac{d}{c}}\right) + 4d \text{EllipticF}\left(\frac{x}{2}, 2\sqrt{-\frac{d}{c}}\right)\right) \sqrt{\frac{dx^2+c}{c}}}{\sqrt{dx^2+c}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-x^2+4)^(1/2)/(d*x^2+c)^(1/2),x)`

[Out] `(c*EllipticF(1/2*x,2*(-d/c)^(1/2))+4*EllipticF(1/2*x,2*(-d/c)^(1/2))*d-c*EllipticE(1/2*x,2*(-d/c)^(1/2)))*((d*x^2+c)/c)^(1/2)/(d*x^2+c)^(1/2)/d`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2 + 4}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-x^2+4)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-x^2 + 4)/sqrt(d*x^2 + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{4 - x^2}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((4 - x^2)^(1/2)/(c + d*x^2)^(1/2),x)`

[Out] `int((4 - x^2)^(1/2)/(c + d*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(x-2)(x+2)}}{\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-x**2+4)**(1/2)/(d*x**2+c)**(1/2),x)`

[Out] `Integral(sqrt(-(x - 2)*(x + 2))/sqrt(c + d*x**2), x)`

$$3.183 \quad \int \frac{\sqrt{4+x^2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=150

$$\frac{4\sqrt{c+dx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{x}{2}\right), 1 - \frac{4d}{c}\right)}{c\sqrt{x^2+4} \sqrt{\frac{c+dx^2}{c(x^2+4)}}} + \frac{x\sqrt{c+dx^2}}{d\sqrt{x^2+4}} - \frac{\sqrt{c+dx^2} E\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| 1 - \frac{4d}{c}\right)}{d\sqrt{x^2+4} \sqrt{\frac{c+dx^2}{c(x^2+4)}}}$$

[Out] $x*(d*x^2+c)^{(1/2)}/d/(x^2+4)^{(1/2)}-(1/(x^2+4))^{(1/2)}*\operatorname{EllipticE}(x/(x^2+4)^{(1/2)}, (1-4*d/c)^{(1/2)})*(d*x^2+c)^{(1/2)}/d/((d*x^2+c)/c/(x^2+4))^{(1/2)}+4*(1/(x^2+4))^{(1/2)}*\operatorname{EllipticF}(x/(x^2+4)^{(1/2)}, (1-4*d/c)^{(1/2)})*(d*x^2+c)^{(1/2)}/c/((d*x^2+c)/c/(x^2+4))^{(1/2)}$

Rubi [A] time = 0.06, antiderivative size = 150, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {422, 418, 492, 411}

$$\frac{x\sqrt{c+dx^2}}{d\sqrt{x^2+4}} + \frac{4\sqrt{c+dx^2} F\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| 1 - \frac{4d}{c}\right)}{c\sqrt{x^2+4} \sqrt{\frac{c+dx^2}{c(x^2+4)}}} - \frac{\sqrt{c+dx^2} E\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| 1 - \frac{4d}{c}\right)}{d\sqrt{x^2+4} \sqrt{\frac{c+dx^2}{c(x^2+4)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[4 + x^2]/Sqrt[c + d*x^2], x]

[Out] $(x*\operatorname{Sqrt}[c + d*x^2])/(d*\operatorname{Sqrt}[4 + x^2]) - (\operatorname{Sqrt}[c + d*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[x/2], 1 - (4*d)/c])/(d*\operatorname{Sqrt}[4 + x^2]*\operatorname{Sqrt}[(c + d*x^2)/(c*(4 + x^2))]) + (4*\operatorname{Sqrt}[c + d*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[x/2], 1 - (4*d)/c])/(c*\operatorname{Sqrt}[4 + x^2]*\operatorname{Sqrt}[(c + d*x^2)/(c*(4 + x^2))])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a + b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{4+x^2}}{\sqrt{c+dx^2}} dx &= 4 \int \frac{1}{\sqrt{4+x^2} \sqrt{c+dx^2}} dx + \int \frac{x^2}{\sqrt{4+x^2} \sqrt{c+dx^2}} dx \\
&= \frac{x\sqrt{c+dx^2}}{d\sqrt{4+x^2}} + \frac{4\sqrt{c+dx^2} F\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| 1 - \frac{4d}{c}\right)}{c\sqrt{4+x^2} \sqrt{\frac{c+dx^2}{c(4+x^2)}}} - \frac{4 \int \frac{\sqrt{c+dx^2}}{(4+x^2)^{3/2}} dx}{d} \\
&= \frac{x\sqrt{c+dx^2}}{d\sqrt{4+x^2}} - \frac{\sqrt{c+dx^2} E\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| 1 - \frac{4d}{c}\right)}{d\sqrt{4+x^2} \sqrt{\frac{c+dx^2}{c(4+x^2)}}} + \frac{4\sqrt{c+dx^2} F\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| 1 - \frac{4d}{c}\right)}{c\sqrt{4+x^2} \sqrt{\frac{c+dx^2}{c(4+x^2)}}}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 60, normalized size = 0.40

$$\frac{2\sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}} x\right) \middle| \frac{c}{4d}\right)}{\sqrt{-\frac{d}{c}} \sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[4 + x^2]/Sqrt[c + d*x^2], x]

[Out] (2*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], c/(4*d)]/(Sqrt[-(d/c)]*Sqrt[c + d*x^2]))

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{x^2+4}}{\sqrt{dx^2+c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+4)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(x^2 + 4)/sqrt(d*x^2 + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2+4}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+4)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(x^2 + 4)/sqrt(d*x^2 + c), x)

maple [A] time = 0.03, size = 53, normalized size = 0.35

$$\frac{2\sqrt{\frac{dx^2+c}{c}} \text{EllipticE}\left(\sqrt{-\frac{d}{c}} x, \frac{\sqrt{\frac{c}{d}}}{2}\right)}{\sqrt{dx^2+c} \sqrt{-\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2+4)^(1/2)/(d*x^2+c)^(1/2),x)`

[Out] `2*EllipticE(x*(-1/c*d)^(1/2),1/2*(c/d)^(1/2))*((d*x^2+c)/c)^(1/2)/(d*x^2+c)^(1/2)/(-1/c*d)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2 + 4}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((x^2+4)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(x^2 + 4)/sqrt(d*x^2 + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{x^2 + 4}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2 + 4)^(1/2)/(c + d*x^2)^(1/2),x)`

[Out] `int((x^2 + 4)^(1/2)/(c + d*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2 + 4}}{\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((x**2+4)**(1/2)/(d*x**2+c)**(1/2),x)`

[Out] `Integral(sqrt(x**2 + 4)/sqrt(c + d*x**2), x)`

$$3.184 \quad \int \frac{\sqrt{1-x^2}}{\sqrt{2-3x^2}} dx$$

Optimal. Leaf size=20

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{2}{3}\right)}{\sqrt{3}}$$

[Out] 1/3*EllipticE(1/2*x*6^(1/2),1/3*6^(1/2))*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {424}

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{2}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - x^2]/Sqrt[2 - 3*x^2], x]

[Out] EllipticE[ArcSin[Sqrt[3/2]*x], 2/3]/Sqrt[3]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{1-x^2}}{\sqrt{2-3x^2}} dx = \frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{2}{3}\right)}{\sqrt{3}}$$

Mathematica [A] time = 0.01, size = 20, normalized size = 1.00

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{2}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - x^2]/Sqrt[2 - 3*x^2], x]

[Out] EllipticE[ArcSin[Sqrt[3/2]*x], 2/3]/Sqrt[3]

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-x^2+1}\sqrt{-3x^2+2}}{3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(-3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(-x^2 + 1)*sqrt(-3*x^2 + 2)/(3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2 + 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-x^2 + 1)/sqrt(-3*x^2 + 2), x)

maple [A] time = 0.02, size = 23, normalized size = 1.15

$$\frac{\sqrt{2} \left(2 \operatorname{EllipticE} \left(x, \frac{\sqrt{6}}{2} \right) + \operatorname{EllipticF} \left(x, \frac{\sqrt{6}}{2} \right) \right)}{6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x)

[Out] 1/6*2^(1/2)*(EllipticF(x,1/2*6^(1/2))+2*EllipticE(x,1/2*6^(1/2)))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2 + 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-x^2 + 1)/sqrt(-3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{\sqrt{1 - x^2}}{\sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - x^2)^(1/2)/(2 - 3*x^2)^(1/2),x)

[Out] int((1 - x^2)^(1/2)/(2 - 3*x^2)^(1/2), x)

sympy [A] time = 3.94, size = 34, normalized size = 1.70

$$\left\{ \frac{\sqrt{3} E \left(\operatorname{asin} \left(\frac{\sqrt{6}x}{2} \right) \middle| \frac{2}{3} \right)}{3} \quad \text{for } x > -\frac{\sqrt{6}}{3} \wedge x < \frac{\sqrt{6}}{3} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x**2+1)**(1/2)/(-3*x**2+2)**(1/2),x)

[Out] Piecewise((sqrt(3)*elliptic_e(asin(sqrt(6)*x/2), 2/3)/3, (x > -sqrt(6)/3) & (x < sqrt(6)/3))

$$3.185 \quad \int \frac{\sqrt{4-x^2}}{\sqrt{2-3x^2}} dx$$

Optimal. Leaf size=21

$$\frac{2E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{1}{6}\right)}{\sqrt{3}}$$

[Out] 2/3*EllipticE(1/2*x*6^(1/2),1/6*6^(1/2))*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {424}

$$\frac{2E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{1}{6}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[4 - x^2]/Sqrt[2 - 3*x^2], x]

[Out] (2*EllipticE[ArcSin[Sqrt[3/2]*x], 1/6])/Sqrt[3]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{4-x^2}}{\sqrt{2-3x^2}} dx = \frac{2E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{1}{6}\right)}{\sqrt{3}}$$

Mathematica [A] time = 0.00, size = 21, normalized size = 1.00

$$\frac{2E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{1}{6}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[4 - x^2]/Sqrt[2 - 3*x^2], x]

[Out] (2*EllipticE[ArcSin[Sqrt[3/2]*x], 1/6])/Sqrt[3]

fricas [F] time = 0.47, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-x^2+4}\sqrt{-3x^2+2}}{3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+4)^(1/2)/(-3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(-x^2 + 4)*sqrt(-3*x^2 + 2)/(3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2 + 4}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+4)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-x^2 + 4)/sqrt(-3*x^2 + 2), x)

maple [A] time = 0.04, size = 18, normalized size = 0.86

$$\frac{2\sqrt{3} \operatorname{EllipticE}\left(\frac{\sqrt{6}x}{2}, \frac{\sqrt{6}}{6}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x^2+4)^(1/2)/(-3*x^2+2)^(1/2),x)

[Out] 2/3*EllipticE(1/2*x*6^(1/2),1/6*6^(1/2))*3^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2 + 4}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+4)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-x^2 + 4)/sqrt(-3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{\sqrt{4 - x^2}}{\sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4 - x^2)^(1/2)/(2 - 3*x^2)^(1/2),x)

[Out] int((4 - x^2)^(1/2)/(2 - 3*x^2)^(1/2), x)

sympy [A] time = 4.21, size = 36, normalized size = 1.71

$$\left\{ \frac{2\sqrt{3} E\left(\operatorname{asin}\left(\frac{\sqrt{6}x}{2}\right)\middle|\frac{1}{6}\right)}{3} \quad \text{for } x > -\frac{\sqrt{6}}{3} \wedge x < \frac{\sqrt{6}}{3} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x**2+4)**(1/2)/(-3*x**2+2)**(1/2),x)

[Out] Piecewise((2*sqrt(3)*elliptic_e(asin(sqrt(6)*x/2), 1/6)/3, (x > -sqrt(6)/3) & (x < sqrt(6)/3))

$$3.186 \quad \int \frac{\sqrt{1-4x^2}}{\sqrt{2-3x^2}} dx$$

Optimal. Leaf size=20

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{8}{3}\right)}{\sqrt{3}}$$

[Out] 1/3*EllipticE(1/2*x*6^(1/2),2/3*6^(1/2))*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {424}

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{8}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - 4*x^2]/Sqrt[2 - 3*x^2],x]

[Out] EllipticE[ArcSin[Sqrt[3/2]*x], 8/3]/Sqrt[3]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{1-4x^2}}{\sqrt{2-3x^2}} dx = \frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{8}{3}\right)}{\sqrt{3}}$$

Mathematica [A] time = 0.01, size = 20, normalized size = 1.00

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{8}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - 4*x^2]/Sqrt[2 - 3*x^2],x]

[Out] EllipticE[ArcSin[Sqrt[3/2]*x], 8/3]/Sqrt[3]

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-3x^2+2}\sqrt{-4x^2+1}}{3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-4*x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(-3*x^2 + 2)*sqrt(-4*x^2 + 1)/(3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-4x^2 + 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-4*x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-4*x^2 + 1)/sqrt(-3*x^2 + 2), x)

maple [A] time = 0.03, size = 29, normalized size = 1.45

$$\frac{\sqrt{2} \left(-8 \operatorname{EllipticE} \left(2x, \frac{\sqrt{6}}{4} \right) + 5 \operatorname{EllipticF} \left(2x, \frac{\sqrt{6}}{4} \right) \right)}{12}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-4*x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x)

[Out] -1/12*2^(1/2)*(5*EllipticF(2*x,1/4*6^(1/2))-8*EllipticE(2*x,1/4*6^(1/2)))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-4x^2 + 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-4*x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-4*x^2 + 1)/sqrt(-3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{\sqrt{1 - 4x^2}}{\sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - 4*x^2)^(1/2)/(2 - 3*x^2)^(1/2),x)

[Out] int((1 - 4*x^2)^(1/2)/(2 - 3*x^2)^(1/2), x)

sympy [A] time = 4.10, size = 34, normalized size = 1.70

$$\left\{ \frac{\sqrt{3} E \left(\operatorname{asin} \left(\frac{\sqrt{6}x}{2} \right) \middle| \frac{8}{3} \right)}{3} \quad \text{for } x > -\frac{\sqrt{6}}{3} \wedge x < \frac{\sqrt{6}}{3} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-4*x**2+1)**(1/2)/(-3*x**2+2)**(1/2),x)

[Out] Piecewise((sqrt(3)*elliptic_e(asin(sqrt(6)*x/2), 8/3)/3, (x > -sqrt(6)/3) & (x < sqrt(6)/3)))

$$3.187 \quad \int \frac{\sqrt{1+x^2}}{\sqrt{1-x^2}} dx$$

Optimal. Leaf size=4

$$E(\sin^{-1}(x)|-1)$$

[Out] EllipticE(x,I)

Rubi [A] time = 0.01, antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {424}

$$E(\sin^{-1}(x)|-1)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 + x^2]/Sqrt[1 - x^2], x]

[Out] EllipticE[ArcSin[x], -1]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{1+x^2}}{\sqrt{1-x^2}} dx = E(\sin^{-1}(x)|-1)$$

Mathematica [A] time = 0.00, size = 4, normalized size = 1.00

$$E(\sin^{-1}(x)|-1)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 + x^2]/Sqrt[1 - x^2], x]

[Out] EllipticE[ArcSin[x], -1]

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2+1}\sqrt{-x^2+1}}{x^2-1}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(-x^2+1)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(x^2 + 1)*sqrt(-x^2 + 1)/(x^2 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2+1}}{\sqrt{-x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(x^2 + 1)/sqrt(-x^2 + 1), x)

maple [A] time = 0.02, size = 5, normalized size = 1.25

EllipticE(x, i)

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2+1)^(1/2)/(-x^2+1)^(1/2),x)

[Out] EllipticE(x,I)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2 + 1}}{\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(x^2 + 1)/sqrt(-x^2 + 1), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.25

$$\int \frac{\sqrt{x^2 + 1}}{\sqrt{1 - x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2 + 1)^(1/2)/(1 - x^2)^(1/2),x)

[Out] int((x^2 + 1)^(1/2)/(1 - x^2)^(1/2), x)

sympy [B] time = 2.39, size = 10, normalized size = 2.50

$$\left\{ E(\operatorname{asin}(x)|-1) \quad \text{for } x > -1 \wedge x < 1 \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x**2+1)**(1/2)/(-x**2+1)**(1/2),x)

[Out] Piecewise((elliptic_e(asin(x), -1), (x > -1) & (x < 1)))

$$3.188 \quad \int \frac{\sqrt{1+x^2}}{\sqrt{2-3x^2}} dx$$

Optimal. Leaf size=20

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{2}{3}\right)}{\sqrt{3}}$$

[Out] 1/3*EllipticE(1/2*x*6^(1/2),1/3*I*6^(1/2))*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {424}

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{2}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 + x^2]/Sqrt[2 - 3*x^2], x]

[Out] EllipticE[ArcSin[Sqrt[3/2]*x], -2/3]/Sqrt[3]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{1+x^2}}{\sqrt{2-3x^2}} dx = \frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{2}{3}\right)}{\sqrt{3}}$$

Mathematica [A] time = 0.00, size = 20, normalized size = 1.00

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{2}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 + x^2]/Sqrt[2 - 3*x^2], x]

[Out] EllipticE[ArcSin[Sqrt[3/2]*x], -2/3]/Sqrt[3]

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2+1}\sqrt{-3x^2+2}}{3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(-3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(x^2 + 1)*sqrt(-3*x^2 + 2)/(3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2 + 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(x^2 + 1)/sqrt(-3*x^2 + 2), x)

maple [A] time = 0.02, size = 19, normalized size = 0.95

$$\frac{\sqrt{3} \operatorname{EllipticE}\left(\frac{\sqrt{6}x}{2}, \frac{i\sqrt{6}}{3}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x)

[Out] 1/3*EllipticE(1/2*6^(1/2)*x,1/3*I*6^(1/2))*3^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2 + 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(x^2 + 1)/sqrt(-3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{\sqrt{x^2 + 1}}{\sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2 + 1)^(1/2)/(2 - 3*x^2)^(1/2),x)

[Out] int((x^2 + 1)^(1/2)/(2 - 3*x^2)^(1/2), x)

sympy [A] time = 3.96, size = 36, normalized size = 1.80

$$\left\{ \frac{\sqrt{3} E\left(\operatorname{asin}\left(\frac{\sqrt{6}x}{2}\right) \middle| -\frac{2}{3}\right)}{3} \quad \text{for } x > -\frac{\sqrt{6}}{3} \wedge x < \frac{\sqrt{6}}{3} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x**2+1)**(1/2)/(-3*x**2+2)**(1/2),x)

[Out] Piecewise((sqrt(3)*elliptic_e(asin(sqrt(6)*x/2), -2/3)/3, (x > -sqrt(6)/3) & (x < sqrt(6)/3))

$$3.189 \quad \int \frac{\sqrt{4+x^2}}{\sqrt{2-3x^2}} dx$$

Optimal. Leaf size=21

$$\frac{2E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{1}{6}\right)}{\sqrt{3}}$$

[Out] 2/3*EllipticE(1/2*x*6^(1/2),1/6*I*6^(1/2))*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {424}

$$\frac{2E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{1}{6}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[4 + x^2]/Sqrt[2 - 3*x^2], x]

[Out] (2*EllipticE[ArcSin[Sqrt[3/2]*x], -1/6])/Sqrt[3]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{4+x^2}}{\sqrt{2-3x^2}} dx = \frac{2E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{1}{6}\right)}{\sqrt{3}}$$

Mathematica [A] time = 0.00, size = 21, normalized size = 1.00

$$\frac{2E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{1}{6}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[4 + x^2]/Sqrt[2 - 3*x^2], x]

[Out] (2*EllipticE[ArcSin[Sqrt[3/2]*x], -1/6])/Sqrt[3]

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2+4}\sqrt{-3x^2+2}}{3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+4)^(1/2)/(-3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(x^2 + 4)*sqrt(-3*x^2 + 2)/(3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2 + 4}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+4)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(x^2 + 4)/sqrt(-3*x^2 + 2), x)

maple [A] time = 0.02, size = 19, normalized size = 0.90

$$\frac{2\sqrt{3} \operatorname{EllipticE}\left(\frac{\sqrt{6}x}{2}, \frac{i\sqrt{6}}{6}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2+4)^(1/2)/(-3*x^2+2)^(1/2),x)

[Out] 2/3*EllipticE(1/2*6^(1/2)*x,1/6*I*6^(1/2))*3^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2 + 4}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+4)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(x^2 + 4)/sqrt(-3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{\sqrt{x^2 + 4}}{\sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2 + 4)^(1/2)/(2 - 3*x^2)^(1/2),x)

[Out] int((x^2 + 4)^(1/2)/(2 - 3*x^2)^(1/2), x)

sympy [A] time = 4.17, size = 37, normalized size = 1.76

$$\left\{ \frac{2\sqrt{3}E\left(\operatorname{asin}\left(\frac{\sqrt{6}x}{2}\right)\middle|-\frac{1}{6}\right)}{3} \quad \text{for } x > -\frac{\sqrt{6}}{3} \wedge x < \frac{\sqrt{6}}{3} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x**2+4)**(1/2)/(-3*x**2+2)**(1/2),x)

[Out] Piecewise((2*sqrt(3)*elliptic_e(asin(sqrt(6)*x/2), -1/6)/3, (x > -sqrt(6)/3) & (x < sqrt(6)/3))

$$3.190 \quad \int \frac{\sqrt{1+4x^2}}{\sqrt{2-3x^2}} dx$$

Optimal. Leaf size=20

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{8}{3}\right)}{\sqrt{3}}$$

[Out] 1/3*EllipticE(1/2*x*6^(1/2),2/3*I*6^(1/2))*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {424}

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{8}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 + 4*x^2]/Sqrt[2 - 3*x^2], x]

[Out] EllipticE[ArcSin[Sqrt[3/2]*x], -8/3]/Sqrt[3]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{1+4x^2}}{\sqrt{2-3x^2}} dx = \frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{8}{3}\right)}{\sqrt{3}}$$

Mathematica [A] time = 0.00, size = 20, normalized size = 1.00

$$\frac{E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|-\frac{8}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 + 4*x^2]/Sqrt[2 - 3*x^2], x]

[Out] EllipticE[ArcSin[Sqrt[3/2]*x], -8/3]/Sqrt[3]

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{4x^2+1}\sqrt{-3x^2+2}}{3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+1)^(1/2)/(-3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(4*x^2 + 1)*sqrt(-3*x^2 + 2)/(3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{4x^2 + 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(4*x^2 + 1)/sqrt(-3*x^2 + 2), x)

maple [A] time = 0.03, size = 19, normalized size = 0.95

$$\frac{\sqrt{3} \operatorname{EllipticE}\left(\frac{\sqrt{6}x}{2}, \frac{2i\sqrt{6}}{3}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4*x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x)

[Out] 1/3*EllipticE(1/2*6^(1/2)*x,2/3*I*6^(1/2))*3^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{4x^2 + 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(4*x^2 + 1)/sqrt(-3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{\sqrt{4x^2 + 1}}{\sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4*x^2 + 1)^(1/2)/(2 - 3*x^2)^(1/2),x)

[Out] int((4*x^2 + 1)^(1/2)/(2 - 3*x^2)^(1/2), x)

sympy [A] time = 4.05, size = 36, normalized size = 1.80

$$\left\{ \frac{\sqrt{3} E\left(\operatorname{asin}\left(\frac{\sqrt{6}x}{2}\right) \middle| -\frac{8}{3}\right)}{3} \quad \text{for } x > -\frac{\sqrt{6}}{3} \wedge x < \frac{\sqrt{6}}{3} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x**2+1)**(1/2)/(-3*x**2+2)**(1/2),x)

[Out] Piecewise((sqrt(3)*elliptic_e(asin(sqrt(6)*x/2), -8/3)/3, (x > -sqrt(6)/3) & (x < sqrt(6)/3))

$$3.191 \quad \int \frac{\sqrt{1-x^2}}{\sqrt{1+x^2}} dx$$

Optimal. Leaf size=13

$$2 \operatorname{EllipticF}(\sin^{-1}(x), -1) - E(\sin^{-1}(x) | -1)$$

[Out] -EllipticE(x,I)+2*EllipticF(x,I)

Rubi [A] time = 0.01, antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {423, 424, 248, 221}

$$2F(\sin^{-1}(x) | -1) - E(\sin^{-1}(x) | -1)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - x^2]/Sqrt[1 + x^2], x]

[Out] -EllipticE[ArcSin[x], -1] + 2*EllipticF[ArcSin[x], -1]

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := Simp[EllipticF[ArcSin[(Rt[-b, 4]*x)/Rt[a, 4]], -1]/(Rt[a, 4]*Rt[-b, 4]), x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]

Rule 248

Int[((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 423

Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{1-x^2}}{\sqrt{1+x^2}} dx &= 2 \int \frac{1}{\sqrt{1-x^2}\sqrt{1+x^2}} dx - \int \frac{\sqrt{1+x^2}}{\sqrt{1-x^2}} dx \\ &= -E(\sin^{-1}(x) | -1) + 2 \int \frac{1}{\sqrt{1-x^4}} dx \\ &= -E(\sin^{-1}(x) | -1) + 2F(\sin^{-1}(x) | -1) \end{aligned}$$

Mathematica [C] time = 0.00, size = 12, normalized size = 0.92

$$-iE(i \sinh^{-1}(x) | -1)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - x^2]/Sqrt[1 + x^2],x]

[Out] (-I)*EllipticE[I*ArcSinh[x], -1]

fricas [F] time = 0.80, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+1}}{\sqrt{x^2+1}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-x^2 + 1)/sqrt(x^2 + 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+1}}{\sqrt{x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-x^2 + 1)/sqrt(x^2 + 1), x)

maple [A] time = 0.01, size = 14, normalized size = 1.08

$$- \text{EllipticE}(x, i) + 2 \text{EllipticF}(x, i)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x^2+1)^(1/2)/(x^2+1)^(1/2),x)

[Out] -EllipticE(x,I)+2*EllipticF(x,I)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+1}}{\sqrt{x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-x^2 + 1)/sqrt(x^2 + 1), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.08

$$\int \frac{\sqrt{1-x^2}}{\sqrt{x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - x^2)^(1/2)/(x^2 + 1)^(1/2),x)

[Out] int((1 - x^2)^(1/2)/(x^2 + 1)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(x-1)(x+1)}}{\sqrt{x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-x**2+1)**(1/2)/(x**2+1)**(1/2),x)
```

```
[Out] Integral(sqrt(-(x - 1)*(x + 1))/sqrt(x**2 + 1), x)
```

$$3.192 \quad \int \frac{\sqrt{1-x^2}}{\sqrt{2+3x^2}} dx$$

Optimal. Leaf size=31

$$\frac{5 \operatorname{EllipticF}\left(\sin^{-1}(x), -\frac{3}{2}\right)}{3\sqrt{2}} - \frac{1}{3}\sqrt{2} E\left(\sin^{-1}(x) \middle| -\frac{3}{2}\right)$$

[Out] 5/6*EllipticF(x,1/2*I*6^(1/2))*2^(1/2)-1/3*EllipticE(x,1/2*I*6^(1/2))*2^(1/2)

Rubi [A] time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {423, 424, 419}

$$\frac{5F\left(\sin^{-1}(x) \middle| -\frac{3}{2}\right)}{3\sqrt{2}} - \frac{1}{3}\sqrt{2} E\left(\sin^{-1}(x) \middle| -\frac{3}{2}\right)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - x^2]/Sqrt[2 + 3*x^2], x]

[Out] -(Sqrt[2]*EllipticE[ArcSin[x], -3/2])/3 + (5*EllipticF[ArcSin[x], -3/2])/(3*Sqrt[2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{1-x^2}}{\sqrt{2+3x^2}} dx &= -\left(\frac{1}{3} \int \frac{\sqrt{2+3x^2}}{\sqrt{1-x^2}} dx\right) + \frac{5}{3} \int \frac{1}{\sqrt{1-x^2} \sqrt{2+3x^2}} dx \\ &= -\frac{1}{3}\sqrt{2} E\left(\sin^{-1}(x) \middle| -\frac{3}{2}\right) + \frac{5F\left(\sin^{-1}(x) \middle| -\frac{3}{2}\right)}{3\sqrt{2}} \end{aligned}$$

Mathematica [C] time = 0.01, size = 27, normalized size = 0.87

$$-\frac{iE\left(i \sinh^{-1}\left(\sqrt{\frac{3}{2}}x\right) \middle| -\frac{2}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - x^2]/Sqrt[2 + 3*x^2], x]

[Out] ((-I)*EllipticE[I*ArcSinh[Sqrt[3/2]*x], -2/3])/Sqrt[3]

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+1}}{\sqrt{3x^2+2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-x^2 + 1)/sqrt(3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-x^2 + 1)/sqrt(3*x^2 + 2), x)

maple [A] time = 0.02, size = 27, normalized size = 0.87

$$\frac{\left(-2 \text{EllipticE}\left(x, \frac{i\sqrt{6}}{2}\right) + 5 \text{EllipticF}\left(x, \frac{i\sqrt{6}}{2}\right)\right) \sqrt{2}}{6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x^2+1)^(1/2)/(3*x^2+2)^(1/2), x)

[Out] 1/6*(5*EllipticF(x, 1/2*I*6^(1/2))-2*EllipticE(x, 1/2*I*6^(1/2)))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(-x^2 + 1)/sqrt(3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\sqrt{1-x^2}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - x^2)^(1/2)/(3*x^2 + 2)^(1/2), x)

[Out] int((1 - x^2)^(1/2)/(3*x^2 + 2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(x-1)(x+1)}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-x**2+1)**(1/2)/(3*x**2+2)**(1/2),x)
```

```
[Out] Integral(sqrt(-(x - 1)*(x + 1))/sqrt(3*x**2 + 2), x)
```

$$3.193 \quad \int \frac{\sqrt{4-x^2}}{\sqrt{2+3x^2}} dx$$

Optimal. Leaf size=35

$$\frac{7}{3}\sqrt{2} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{x}{2}\right), -6\right) - \frac{1}{3}\sqrt{2} E\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -6\right)$$

[Out] $-1/3*\operatorname{EllipticE}(1/2*x, I*6^{(1/2)})*2^{(1/2)}+7/3*\operatorname{EllipticF}(1/2*x, I*6^{(1/2)})*2^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {423, 424, 419}

$$\frac{7}{3}\sqrt{2} F\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -6\right) - \frac{1}{3}\sqrt{2} E\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -6\right)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[4 - x^2]/Sqrt[2 + 3*x^2], x]

[Out] $-(\operatorname{Sqrt}[2]*\operatorname{EllipticE}[\operatorname{ArcSin}[x/2], -6])/3 + (7*\operatorname{Sqrt}[2]*\operatorname{EllipticF}[\operatorname{ArcSin}[x/2], -6])/3$

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{4-x^2}}{\sqrt{2+3x^2}} dx &= -\left(\frac{1}{3} \int \frac{\sqrt{2+3x^2}}{\sqrt{4-x^2}} dx\right) + \frac{14}{3} \int \frac{1}{\sqrt{4-x^2}\sqrt{2+3x^2}} dx \\ &= -\frac{1}{3}\sqrt{2} E\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -6\right) + \frac{7}{3}\sqrt{2} F\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -6\right) \end{aligned}$$

Mathematica [C] time = 0.00, size = 27, normalized size = 0.77

$$-\frac{2iE\left(i \sinh^{-1}\left(\sqrt{\frac{3}{2}}x\right) \middle| -\frac{1}{6}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[4 - x^2]/Sqrt[2 + 3*x^2], x]

[Out] $((-2*I)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[3/2]*x], -1/6))/\text{Sqrt}[3]$

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+4}}{\sqrt{3x^2+2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+4)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-x^2 + 4)/sqrt(3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+4}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+4)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-x^2 + 4)/sqrt(3*x^2 + 2), x)

maple [A] time = 0.03, size = 31, normalized size = 0.89

$$\frac{(-\text{EllipticE}\left(\frac{x}{2}, i\sqrt{6}\right) + 7\text{EllipticF}\left(\frac{x}{2}, i\sqrt{6}\right))\sqrt{2}}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x^2+4)^(1/2)/(3*x^2+2)^(1/2), x)

[Out] $1/3*(7*\text{EllipticF}(1/2*x, I*6^(1/2))-\text{EllipticE}(1/2*x, I*6^(1/2)))*2^(1/2)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+4}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+4)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(-x^2 + 4)/sqrt(3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\sqrt{4-x^2}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4 - x^2)^(1/2)/(3*x^2 + 2)^(1/2), x)

[Out] int((4 - x^2)^(1/2)/(3*x^2 + 2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(x-2)(x+2)}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-x**2+4)**(1/2)/(3*x**2+2)**(1/2),x)
```

```
[Out] Integral(sqrt(-(x - 2)*(x + 2))/sqrt(3*x**2 + 2), x)
```

$$3.194 \quad \int \frac{\sqrt{1-4x^2}}{\sqrt{2+3x^2}} dx$$

Optimal. Leaf size=35

$$\frac{11 \operatorname{EllipticF}\left(\sin^{-1}(2x), -\frac{3}{8}\right)}{6\sqrt{2}} - \frac{2}{3}\sqrt{2} E\left(\sin^{-1}(2x) \middle| -\frac{3}{8}\right)$$

[Out] 11/12*EllipticF(2*x,1/4*I*6^(1/2))*2^(1/2)-2/3*EllipticE(2*x,1/4*I*6^(1/2))*2^(1/2)

Rubi [A] time = 0.02, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {423, 424, 419}

$$\frac{11F\left(\sin^{-1}(2x) \middle| -\frac{3}{8}\right)}{6\sqrt{2}} - \frac{2}{3}\sqrt{2} E\left(\sin^{-1}(2x) \middle| -\frac{3}{8}\right)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - 4*x^2]/Sqrt[2 + 3*x^2],x]

[Out] (-2*Sqrt[2]*EllipticE[ArcSin[2*x], -3/8])/3 + (11*EllipticF[ArcSin[2*x], -3/8])/(6*Sqrt[2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{1-4x^2}}{\sqrt{2+3x^2}} dx &= -\left(\frac{4}{3} \int \frac{\sqrt{2+3x^2}}{\sqrt{1-4x^2}} dx\right) + \frac{11}{3} \int \frac{1}{\sqrt{1-4x^2}\sqrt{2+3x^2}} dx \\ &= -\frac{2}{3}\sqrt{2} E\left(\sin^{-1}(2x) \middle| -\frac{3}{8}\right) + \frac{11F\left(\sin^{-1}(2x) \middle| -\frac{3}{8}\right)}{6\sqrt{2}} \end{aligned}$$

Mathematica [C] time = 0.00, size = 27, normalized size = 0.77

$$\frac{iE\left(i \sinh^{-1}\left(\sqrt{\frac{3}{2}}x\right) \middle| -\frac{8}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - 4*x^2]/Sqrt[2 + 3*x^2], x]

[Out] ((-I)*EllipticE[I*ArcSinh[Sqrt[3/2]*x], -8/3])/Sqrt[3]

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-4x^2+1}}{\sqrt{3x^2+2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-4*x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-4*x^2 + 1)/sqrt(3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-4x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-4*x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-4*x^2 + 1)/sqrt(3*x^2 + 2), x)

maple [A] time = 0.02, size = 31, normalized size = 0.89

$$\frac{\left(-8 \text{EllipticE}\left(2x, \frac{i\sqrt{6}}{4}\right) + 11 \text{EllipticF}\left(2x, \frac{i\sqrt{6}}{4}\right)\right) \sqrt{2}}{12}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-4*x^2+1)^(1/2)/(3*x^2+2)^(1/2), x)

[Out] 1/12*(11*EllipticF(2*x, 1/4*I*6^(1/2))-8*EllipticE(2*x, 1/4*I*6^(1/2)))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-4x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-4*x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(-4*x^2 + 1)/sqrt(3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\sqrt{1-4x^2}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - 4*x^2)^(1/2)/(3*x^2 + 2)^(1/2), x)

[Out] int((1 - 4*x^2)^(1/2)/(3*x^2 + 2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(2x-1)(2x+1)}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-4*x**2+1)**(1/2)/(3*x**2+2)**(1/2),x)

[Out] Integral(sqrt(-(2*x - 1)*(2*x + 1))/sqrt(3*x**2 + 2), x)

$$3.195 \quad \int \frac{\sqrt{1+x^2}}{\sqrt{2+3x^2}} dx$$

Optimal. Leaf size=131

$$\frac{\sqrt{3x^2+2} \operatorname{EllipticF}\left(\tan^{-1}(x), -\frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{3x^2+2}{x^2+1}}} + \frac{\sqrt{3x^2+2} x}{3\sqrt{x^2+1}} - \frac{\sqrt{2} \sqrt{3x^2+2} E\left(\tan^{-1}(x) \mid -\frac{1}{2}\right)}{3\sqrt{x^2+1} \sqrt{\frac{3x^2+2}{x^2+1}}}$$

[Out] 1/3*x*(3*x^2+2)^(1/2)/(x^2+1)^(1/2)+1/2*(1/(x^2+1))^(1/2)*EllipticF(x/(x^2+1)^(1/2),1/2*I*2^(1/2))*(3*x^2+2)^(1/2)*2^(1/2)/((3*x^2+2)/(x^2+1))^(1/2)-1/3*(1/(x^2+1))^(1/2)*EllipticE(x/(x^2+1)^(1/2),1/2*I*2^(1/2))*2^(1/2)*(3*x^2+2)^(1/2)/((3*x^2+2)/(x^2+1))^(1/2)

Rubi [A] time = 0.04, antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {422, 418, 492, 411}

$$\frac{\sqrt{3x^2+2} x}{3\sqrt{x^2+1}} + \frac{\sqrt{3x^2+2} F\left(\tan^{-1}(x) \mid -\frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{3x^2+2}{x^2+1}}} - \frac{\sqrt{2} \sqrt{3x^2+2} E\left(\tan^{-1}(x) \mid -\frac{1}{2}\right)}{3\sqrt{x^2+1} \sqrt{\frac{3x^2+2}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 + x^2]/Sqrt[2 + 3*x^2], x]

[Out] (x*Sqrt[2 + 3*x^2])/(3*Sqrt[1 + x^2]) - (Sqrt[2]*Sqrt[2 + 3*x^2]*EllipticE[ArcTan[x], -1/2])/(3*Sqrt[1 + x^2]*Sqrt[(2 + 3*x^2)/(1 + x^2)]) + (Sqrt[2 + 3*x^2]*EllipticF[ArcTan[x], -1/2])/(Sqrt[2]*Sqrt[1 + x^2]*Sqrt[(2 + 3*x^2)/(1 + x^2)])

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a + b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{1+x^2}}{\sqrt{2+3x^2}} dx &= \int \frac{1}{\sqrt{1+x^2} \sqrt{2+3x^2}} dx + \int \frac{x^2}{\sqrt{1+x^2} \sqrt{2+3x^2}} dx \\
&= \frac{x\sqrt{2+3x^2}}{3\sqrt{1+x^2}} + \frac{\sqrt{2+3x^2} F\left(\tan^{-1}(x) \middle| -\frac{1}{2}\right)}{\sqrt{2} \sqrt{1+x^2} \sqrt{\frac{2+3x^2}{1+x^2}}} - \frac{1}{3} \int \frac{\sqrt{2+3x^2}}{(1+x^2)^{3/2}} dx \\
&= \frac{x\sqrt{2+3x^2}}{3\sqrt{1+x^2}} - \frac{\sqrt{2} \sqrt{2+3x^2} E\left(\tan^{-1}(x) \middle| -\frac{1}{2}\right)}{3\sqrt{1+x^2} \sqrt{\frac{2+3x^2}{1+x^2}}} + \frac{\sqrt{2+3x^2} F\left(\tan^{-1}(x) \middle| -\frac{1}{2}\right)}{\sqrt{2} \sqrt{1+x^2} \sqrt{\frac{2+3x^2}{1+x^2}}}
\end{aligned}$$

Mathematica [C] time = 0.00, size = 27, normalized size = 0.21

$$-\frac{iE\left(i \sinh^{-1}\left(\sqrt{\frac{3}{2}} x\right) \middle| \frac{2}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 + x^2]/Sqrt[2 + 3*x^2], x]

[Out] ((-I)*EllipticE[I*ArcSinh[Sqrt[3/2]*x], 2/3])/Sqrt[3]

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{x^2+1}}{\sqrt{3x^2+2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(x^2 + 1)/sqrt(3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(x^2 + 1)/sqrt(3*x^2 + 2), x)

maple [A] time = 0.02, size = 30, normalized size = 0.23

$$\frac{i\left(2 \text{EllipticE}\left(ix, \frac{\sqrt{6}}{2}\right) + \text{EllipticF}\left(ix, \frac{\sqrt{6}}{2}\right)\right) \sqrt{2}}{6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2+1)^(1/2)/(3*x^2+2)^(1/2), x)

[Out] -1/6*I*(EllipticF(I*x, 1/2*6^(1/2))+2*EllipticE(I*x, 1/2*6^(1/2)))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(x^2 + 1)/sqrt(3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2 + 1)^(1/2)/(3*x^2 + 2)^(1/2), x)

[Out] int((x^2 + 1)^(1/2)/(3*x^2 + 2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x**2+1)**(1/2)/(3*x**2+2)**(1/2), x)

[Out] Integral(sqrt(x**2 + 1)/sqrt(3*x**2 + 2), x)

$$3.196 \quad \int \frac{\sqrt{4+x^2}}{\sqrt{2+3x^2}} dx$$

Optimal. Leaf size=136

$$\frac{2\sqrt{2}\sqrt{3x^2+2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{x}{2}\right), -5\right)}{\sqrt{x^2+4}\sqrt{\frac{3x^2+2}{x^2+4}}} + \frac{\sqrt{3x^2+2}x}{3\sqrt{x^2+4}} - \frac{\sqrt{2}\sqrt{3x^2+2}E\left(\tan^{-1}\left(\frac{x}{2}\right)\right) - 5}{3\sqrt{x^2+4}\sqrt{\frac{3x^2+2}{x^2+4}}}$$

[Out] $1/3*x*(3*x^2+2)^(1/2)/(x^2+4)^(1/2)-1/3*(1/(x^2+4))^(1/2)*\operatorname{EllipticE}(x/(x^2+4)^(1/2), I*5^(1/2))*2^(1/2)*(3*x^2+2)^(1/2)/((3*x^2+2)/(x^2+4))^(1/2)+2*(1/(x^2+4))^(1/2)*\operatorname{EllipticF}(x/(x^2+4)^(1/2), I*5^(1/2))*2^(1/2)*(3*x^2+2)^(1/2)/((3*x^2+2)/(x^2+4))^(1/2)$

Rubi [A] time = 0.04, antiderivative size = 136, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {422, 418, 492, 411}

$$\frac{\sqrt{3x^2+2}x}{3\sqrt{x^2+4}} + \frac{2\sqrt{2}\sqrt{3x^2+2}F\left(\tan^{-1}\left(\frac{x}{2}\right)\right) - 5}{\sqrt{x^2+4}\sqrt{\frac{3x^2+2}{x^2+4}}} - \frac{\sqrt{2}\sqrt{3x^2+2}E\left(\tan^{-1}\left(\frac{x}{2}\right)\right) - 5}{3\sqrt{x^2+4}\sqrt{\frac{3x^2+2}{x^2+4}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[4 + x^2]/Sqrt[2 + 3*x^2], x]

[Out] $(x*\operatorname{Sqrt}[2 + 3*x^2])/(3*\operatorname{Sqrt}[4 + x^2]) - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[2 + 3*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[x/2], -5])/(3*\operatorname{Sqrt}[4 + x^2]*\operatorname{Sqrt}[(2 + 3*x^2)/(4 + x^2)]) + (2*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[2 + 3*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[x/2], -5])/((\operatorname{Sqrt}[4 + x^2]*\operatorname{Sqrt}[(2 + 3*x^2)/(4 + x^2)]))$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a + b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{4+x^2}}{\sqrt{2+3x^2}} dx &= 4 \int \frac{1}{\sqrt{4+x^2} \sqrt{2+3x^2}} dx + \int \frac{x^2}{\sqrt{4+x^2} \sqrt{2+3x^2}} dx \\ &= \frac{x\sqrt{2+3x^2}}{3\sqrt{4+x^2}} + \frac{2\sqrt{2}\sqrt{2+3x^2} F\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| -5\right)}{\sqrt{4+x^2} \sqrt{\frac{2+3x^2}{4+x^2}}} - \frac{4}{3} \int \frac{\sqrt{2+3x^2}}{(4+x^2)^{3/2}} dx \\ &= \frac{x\sqrt{2+3x^2}}{3\sqrt{4+x^2}} - \frac{\sqrt{2}\sqrt{2+3x^2} E\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| -5\right)}{3\sqrt{4+x^2} \sqrt{\frac{2+3x^2}{4+x^2}}} + \frac{2\sqrt{2}\sqrt{2+3x^2} F\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| -5\right)}{\sqrt{4+x^2} \sqrt{\frac{2+3x^2}{4+x^2}}} \end{aligned}$$

Mathematica [C] time = 0.00, size = 27, normalized size = 0.20

$$-\frac{2iE\left(i \sinh^{-1}\left(\sqrt{\frac{3}{2}}x\right) \middle| \frac{1}{6}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[4 + x^2]/Sqrt[2 + 3*x^2], x]

[Out] ((-2*I)*EllipticE[I*ArcSinh[Sqrt[3/2]*x], 1/6])/Sqrt[3]

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{x^2+4}}{\sqrt{3x^2+2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+4)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(x^2 + 4)/sqrt(3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2+4}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+4)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(x^2 + 4)/sqrt(3*x^2 + 2), x)

maple [A] time = 0.02, size = 26, normalized size = 0.19

$$\frac{i\left(\text{EllipticE}\left(\frac{ix}{2}, \sqrt{6}\right) + 5 \text{EllipticF}\left(\frac{ix}{2}, \sqrt{6}\right)\right) \sqrt{2}}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2+4)^(1/2)/(3*x^2+2)^(1/2), x)

[Out] -1/3*I*(5*EllipticF(1/2*I*x, 6^(1/2))+EllipticE(1/2*I*x, 6^(1/2)))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2+4}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+4)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(x^2 + 4)/sqrt(3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{x^2 + 4}}{\sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2 + 4)^(1/2)/(3*x^2 + 2)^(1/2),x)

[Out] int((x^2 + 4)^(1/2)/(3*x^2 + 2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{x^2 + 4}}{\sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x**2+4)**(1/2)/(3*x**2+2)**(1/2),x)

[Out] Integral(sqrt(x**2 + 4)/sqrt(3*x**2 + 2), x)

$$3.197 \quad \int \frac{\sqrt{1+4x^2}}{\sqrt{2+3x^2}} dx$$

Optimal. Leaf size=148

$$\frac{\sqrt{3x^2+2} \operatorname{EllipticF}\left(\tan^{-1}(2x), \frac{5}{8}\right)}{2\sqrt{2} \sqrt{\frac{3x^2+2}{4x^2+1}} \sqrt{4x^2+1}} + \frac{4\sqrt{3x^2+2}x}{3\sqrt{4x^2+1}} - \frac{2\sqrt{2} \sqrt{3x^2+2} E\left(\tan^{-1}(2x) \middle| \frac{5}{8}\right)}{3\sqrt{\frac{3x^2+2}{4x^2+1}} \sqrt{4x^2+1}}$$

[Out] $4/3*x*(3*x^2+2)^{(1/2)}/(4*x^2+1)^{(1/2)}+1/4*(1/(4*x^2+1))^{(1/2)}*\operatorname{EllipticF}(2*x/(4*x^2+1)^{(1/2)}, 1/4*10^{(1/2)})*(3*x^2+2)^{(1/2)}*2^{(1/2)}/((3*x^2+2)/(4*x^2+1))^{(1/2)}-2/3*(1/(4*x^2+1))^{(1/2)}*\operatorname{EllipticE}(2*x/(4*x^2+1)^{(1/2)}, 1/4*10^{(1/2)})*2^{(1/2)}*(3*x^2+2)^{(1/2)}/((3*x^2+2)/(4*x^2+1))^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 148, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {422, 418, 492, 411}

$$\frac{4\sqrt{3x^2+2}x}{3\sqrt{4x^2+1}} + \frac{\sqrt{3x^2+2} F\left(\tan^{-1}(2x) \middle| \frac{5}{8}\right)}{2\sqrt{2} \sqrt{\frac{3x^2+2}{4x^2+1}} \sqrt{4x^2+1}} - \frac{2\sqrt{2} \sqrt{3x^2+2} E\left(\tan^{-1}(2x) \middle| \frac{5}{8}\right)}{3\sqrt{\frac{3x^2+2}{4x^2+1}} \sqrt{4x^2+1}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[1 + 4*x^2]/Sqrt[2 + 3*x^2], x]`

[Out] $(4*x*\operatorname{Sqrt}[2 + 3*x^2])/((3*\operatorname{Sqrt}[1 + 4*x^2]) - (2*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[2 + 3*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[2*x], 5/8])/(3*\operatorname{Sqrt}[(2 + 3*x^2)/(1 + 4*x^2)]*\operatorname{Sqrt}[1 + 4*x^2]) + (\operatorname{Sqrt}[2 + 3*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[2*x], 5/8])/(2*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[(2 + 3*x^2)/(1 + 4*x^2)]*\operatorname{Sqrt}[1 + 4*x^2]))$

Rule 411

`Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]`

Rule 418

`Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]`

Rule 422

`Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]`

Rule 492

`Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a + b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]`

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{1+4x^2}}{\sqrt{2+3x^2}} dx &= 4 \int \frac{x^2}{\sqrt{2+3x^2}\sqrt{1+4x^2}} dx + \int \frac{1}{\sqrt{2+3x^2}\sqrt{1+4x^2}} dx \\
&= \frac{4x\sqrt{2+3x^2}}{3\sqrt{1+4x^2}} + \frac{\sqrt{2+3x^2} F\left(\tan^{-1}(2x)\middle|\frac{5}{8}\right)}{2\sqrt{2}\sqrt{\frac{2+3x^2}{1+4x^2}}\sqrt{1+4x^2}} - \frac{4}{3} \int \frac{\sqrt{2+3x^2}}{(1+4x^2)^{3/2}} dx \\
&= \frac{4x\sqrt{2+3x^2}}{3\sqrt{1+4x^2}} - \frac{2\sqrt{2}\sqrt{2+3x^2} E\left(\tan^{-1}(2x)\middle|\frac{5}{8}\right)}{3\sqrt{\frac{2+3x^2}{1+4x^2}}\sqrt{1+4x^2}} + \frac{\sqrt{2+3x^2} F\left(\tan^{-1}(2x)\middle|\frac{5}{8}\right)}{2\sqrt{2}\sqrt{\frac{2+3x^2}{1+4x^2}}\sqrt{1+4x^2}}
\end{aligned}$$

Mathematica [C] time = 0.00, size = 27, normalized size = 0.18

$$-\frac{iE\left(i\sinh^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|\frac{8}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 + 4*x^2]/Sqrt[2 + 3*x^2], x]

[Out] ((-I)*EllipticE[I*ArcSinh[Sqrt[3/2]*x], 8/3])/Sqrt[3]

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{4x^2+1}}{\sqrt{3x^2+2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(4*x^2 + 1)/sqrt(3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{4x^2+1}}{\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+1)^(1/2)/(3*x^2+2)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(4*x^2 + 1)/sqrt(3*x^2 + 2), x)

maple [C] time = 0.03, size = 20, normalized size = 0.14

$$\frac{i\sqrt{3} \text{EllipticE}\left(\frac{i\sqrt{6}x}{2}, \frac{2\sqrt{6}}{3}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4*x^2+1)^(1/2)/(3*x^2+2)^(1/2), x)

[Out] -1/3*I*EllipticE(1/2*I*x*6^(1/2), 2/3*6^(1/2))*3^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{4x^2 + 1}}{\sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x^2+1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(4*x^2 + 1)/sqrt(3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{4x^2 + 1}}{\sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((4*x^2 + 1)^(1/2)/(3*x^2 + 2)^(1/2), x)

[Out] int((4*x^2 + 1)^(1/2)/(3*x^2 + 2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{4x^2 + 1}}{\sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((4*x**2+1)**(1/2)/(3*x**2+2)**(1/2), x)

[Out] Integral(sqrt(4*x**2 + 1)/sqrt(3*x**2 + 2), x)

$$3.198 \quad \int \frac{\sqrt{1-x^2}}{\sqrt{-1+2x^2}} dx$$

Optimal. Leaf size=40

$$\frac{\sqrt{1-2x^2} E\left(\sin^{-1}(\sqrt{2}x) \middle| \frac{1}{2}\right)}{\sqrt{2}\sqrt{2x^2-1}}$$

[Out] 1/2*EllipticE(x*2^(1/2),1/2*2^(1/2))*(-2*x^2+1)^(1/2)*2^(1/2)/(2*x^2-1)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {427, 424}

$$\frac{\sqrt{1-2x^2} E\left(\sin^{-1}(\sqrt{2}x) \middle| \frac{1}{2}\right)}{\sqrt{2}\sqrt{2x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - x^2]/Sqrt[-1 + 2*x^2],x]

[Out] (Sqrt[1 - 2*x^2]*EllipticE[ArcSin[Sqrt[2]*x], 1/2])/(Sqrt[2]*Sqrt[-1 + 2*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{1-x^2}}{\sqrt{-1+2x^2}} dx &= \frac{\sqrt{1-2x^2} \int \frac{\sqrt{1-x^2}}{\sqrt{1-2x^2}} dx}{\sqrt{-1+2x^2}} \\ &= \frac{\sqrt{1-2x^2} E\left(\sin^{-1}(\sqrt{2}x) \middle| \frac{1}{2}\right)}{\sqrt{2}\sqrt{-1+2x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 35, normalized size = 0.88

$$\frac{\sqrt{1-2x^2} E\left(\sin^{-1}(\sqrt{2}x) \middle| \frac{1}{2}\right)}{\sqrt{4x^2-2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - x^2]/Sqrt[-1 + 2*x^2],x]

[Out] (Sqrt[1 - 2*x^2]*EllipticE[ArcSin[Sqrt[2]*x], 1/2])/Sqrt[-2 + 4*x^2]

fricas [F] time = 0.76, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+1}}{\sqrt{2x^2-1}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(2*x^2-1)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-x^2 + 1)/sqrt(2*x^2 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+1}}{\sqrt{2x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(2*x^2-1)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-x^2 + 1)/sqrt(2*x^2 - 1), x)

maple [A] time = 0.02, size = 32, normalized size = 0.80

$$\frac{(\text{EllipticE}(x, \sqrt{2}) + \text{EllipticF}(x, \sqrt{2})) \sqrt{-2x^2+1}}{2\sqrt{2x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-x^2+1)^(1/2)/(2*x^2-1)^(1/2),x)

[Out] 1/2*(EllipticF(x, 2^(1/2))+EllipticE(x, 2^(1/2)))*(-2*x^2+1)^(1/2)/(2*x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-x^2+1}}{\sqrt{2x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x^2+1)^(1/2)/(2*x^2-1)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-x^2 + 1)/sqrt(2*x^2 - 1), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{1-x^2}}{\sqrt{2x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - x^2)^(1/2)/(2*x^2 - 1)^(1/2),x)

[Out] int((1 - x^2)^(1/2)/(2*x^2 - 1)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(x-1)(x+1)}}{\sqrt{2x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-x**2+1)**(1/2)/(2*x**2-1)**(1/2),x)

[Out] Integral(sqrt(-(x - 1)*(x + 1))/sqrt(2*x**2 - 1), x)

$$3.199 \quad \int \frac{(a+bx^2)^{7/2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=423

$$\frac{\sqrt{c} \sqrt{a+bx^2} (3bc-7ad) (15a^2d^2-11abcd+8b^2c^2) \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right) + 8\sqrt{c} \sqrt{a+bx^2} (bc-2ad)}{105d^{7/2} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $-8/105*(-2*a*d+b*c)*(11*a^2*d^2-11*a*b*c*d+6*b^2*c^2)*x*(b*x^2+a)^{(1/2)}/d^3/(d*x^2+c)^{(1/2)}+8/105*(-2*a*d+b*c)*(11*a^2*d^2-11*a*b*c*d+6*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)})/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)}*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(7/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-1/105*(-7*a*d+3*b*c)*(15*a^2*d^2-11*a*b*c*d+8*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)})/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)}*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(7/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-6/35*b*(-2*a*d+b*c)*x*(b*x^2+a)^{(3/2)}*(d*x^2+c)^{(1/2)}/d^2+1/7*b*x*(b*x^2+a)^{(5/2)}*(d*x^2+c)^{(1/2)}/d+1/105*b*(71*a^2*d^2-71*a*b*c*d+24*b^2*c^2)*x*(b*x^2+a)^{(1/2)}*(d*x^2+c)^{(1/2)}/d^3$

Rubi [A] time = 0.43, antiderivative size = 423, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {416, 528, 531, 418, 492, 411}

$$\frac{bx\sqrt{a+bx^2} \sqrt{c+dx^2} (71a^2d^2-71abcd+24b^2c^2)}{105d^3} - \frac{8x\sqrt{a+bx^2} (bc-2ad) (11a^2d^2-11abcd+6b^2c^2)}{105d^3 \sqrt{c+dx^2}} - \frac{\sqrt{c} \sqrt{a+bx^2}}{105d^3}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(7/2)/Sqrt[c + d*x^2], x]

[Out] $(-8*(b*c-2*a*d)*(6*b^2*c^2-11*a*b*c*d+11*a^2*d^2)*x*\operatorname{Sqrt}[a+b*x^2])/(105*d^3*\operatorname{Sqrt}[c+d*x^2]) + (b*(24*b^2*c^2-71*a*b*c*d+71*a^2*d^2)*x*\operatorname{Sqrt}[a+b*x^2]*\operatorname{Sqrt}[c+d*x^2])/(105*d^3) - (6*b*(b*c-2*a*d)*x*(a+b*x^2)^{(3/2)}*\operatorname{Sqrt}[c+d*x^2])/(35*d^2) + (b*x*(a+b*x^2)^{(5/2)}*\operatorname{Sqrt}[c+d*x^2])/(7*d) + (8*\operatorname{Sqrt}[c]*(b*c-2*a*d)*(6*b^2*c^2-11*a*b*c*d+11*a^2*d^2)*\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1-(b*c)/(a*d)])/(105*d^{(7/2)}*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2)])*\operatorname{Sqrt}[c+d*x^2]) - (\operatorname{Sqrt}[c]*(3*b*c-7*a*d)*(8*b^2*c^2-11*a*b*c*d+15*a^2*d^2)*\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1-(b*c)/(a*d)])/(105*d^{(7/2)}*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2)])*\operatorname{Sqrt}[c+d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q)+1)), x] + Dist[1/(b*(n*(p+q)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q)+1) - a*d] + d*(b*c*(n*(p+2*q-1)+1) - a*d*(n*(q-1)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,

0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 492

Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a + b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rule 528

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rule 531

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x], x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, e, f, n, p, q}, x]

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^{7/2}}{\sqrt{c+dx^2}} dx &= \frac{bx(a+bx^2)^{5/2}\sqrt{c+dx^2}}{7d} + \frac{\int \frac{(a+bx^2)^{3/2}(-a(bc-7ad)-6b(bc-2ad)x^2)}{\sqrt{c+dx^2}} dx}{7d} \\
&= -\frac{6b(bc-2ad)x(a+bx^2)^{3/2}\sqrt{c+dx^2}}{35d^2} + \frac{bx(a+bx^2)^{5/2}\sqrt{c+dx^2}}{7d} + \frac{\int \frac{\sqrt{a+bx^2}(a(6b^2c^2-17abcd+105a^2d^2)-6b(bc-2ad)x^2)}{\sqrt{c+dx^2}} dx}{35d^2} \\
&= \frac{b(24b^2c^2-71abcd+71a^2d^2)x\sqrt{a+bx^2}\sqrt{c+dx^2}}{105d^3} - \frac{6b(bc-2ad)x(a+bx^2)^{3/2}\sqrt{c+dx^2}}{35d^2} + \frac{\int \frac{\sqrt{a+bx^2}(a(6b^2c^2-17abcd+105a^2d^2)-6b(bc-2ad)x^2)}{\sqrt{c+dx^2}} dx}{35d^2} \\
&= \frac{b(24b^2c^2-71abcd+71a^2d^2)x\sqrt{a+bx^2}\sqrt{c+dx^2}}{105d^3} - \frac{6b(bc-2ad)x(a+bx^2)^{3/2}\sqrt{c+dx^2}}{35d^2} + \frac{\int \frac{\sqrt{a+bx^2}(a(6b^2c^2-17abcd+105a^2d^2)-6b(bc-2ad)x^2)}{\sqrt{c+dx^2}} dx}{35d^2} \\
&= -\frac{8(bc-2ad)(6b^2c^2-11abcd+11a^2d^2)x\sqrt{a+bx^2}}{105d^3\sqrt{c+dx^2}} + \frac{b(24b^2c^2-71abcd+71a^2d^2)x\sqrt{a+bx^2}}{105d^3} + \frac{\int \frac{\sqrt{a+bx^2}(a(6b^2c^2-17abcd+105a^2d^2)-6b(bc-2ad)x^2)}{\sqrt{c+dx^2}} dx}{35d^2} \\
&= -\frac{8(bc-2ad)(6b^2c^2-11abcd+11a^2d^2)x\sqrt{a+bx^2}}{105d^3\sqrt{c+dx^2}} + \frac{b(24b^2c^2-71abcd+71a^2d^2)x\sqrt{a+bx^2}}{105d^3} + \frac{\int \frac{\sqrt{a+bx^2}(a(6b^2c^2-17abcd+105a^2d^2)-6b(bc-2ad)x^2)}{\sqrt{c+dx^2}} dx}{35d^2}
\end{aligned}$$

Mathematica [C] time = 1.57, size = 321, normalized size = 0.76

$$-i\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}\left(105a^4d^4-298a^3bcd^3+353a^2b^2c^2d^2-208ab^3c^3d+48b^4c^4\right)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right),\frac{ad}{bc}\right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(7/2)/Sqrt[c + d*x^2], x]

[Out] (b*Sqrt[b/a]*d*x*(a + b*x^2)*(c + d*x^2)*(122*a^2*d^2 + a*b*d*(-89*c + 66*d*x^2) + 3*b^2*(8*c^2 - 6*c*d*x^2 + 5*d^2*x^4)) - (8*I)*b*c*(-6*b^3*c^3 + 23*a*b^2*c^2*d - 33*a^2*b*c*d^2 + 22*a^3*d^3)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*(48*b^4*c^4 - 208*a*b^3*c^3*d + 353*a^2*b^2*c^2*d^2 - 298*a^3*b*c*d^3 + 105*a^4*d^4)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(105*Sqrt[b/a]*d^4*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b^3x^6 + 3ab^2x^4 + 3a^2bx^2 + a^3)\sqrt{bx^2 + a}}{\sqrt{dx^2 + c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral((b^3*x^6 + 3*a*b^2*x^4 + 3*a^2*b*x^2 + a^3)*sqrt(b*x^2 + a)/sqrt(d*x^2 + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{7/2}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(7/2)/sqrt(d*x^2 + c), x)

maple [A] time = 0.04, size = 852, normalized size = 2.01

$$\sqrt{bx^2 + a} \sqrt{dx^2 + c} \left(15\sqrt{-\frac{b}{a}} b^4 d^4 x^9 + 81\sqrt{-\frac{b}{a}} a b^3 d^4 x^7 - 3\sqrt{-\frac{b}{a}} b^4 c d^3 x^7 + 188\sqrt{-\frac{b}{a}} a^2 b^2 d^4 x^5 - 26\sqrt{-\frac{b}{a}} a b \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(7/2)/(d*x^2+c)^(1/2),x)

[Out] 1/105*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(15*(-1/a*b)^(1/2)*x^9*b^4*d^4+81*(-1/a*b)^(1/2)*x^7*a*b^3*d^4-3*(-1/a*b)^(1/2)*x^7*b^4*c*d^3+188*(-1/a*b)^(1/2)*x^5*a^2*b^2*d^4-26*(-1/a*b)^(1/2)*x^5*a*b^3*c*d^3+6*(-1/a*b)^(1/2)*x^5*b^4*c^2*d^2+122*(-1/a*b)^(1/2)*x^3*a^3*b*d^4+99*(-1/a*b)^(1/2)*x^3*a^2*b^2*c*d^3-83*(-1/a*b)^(1/2)*x^3*a*b^3*c^2*d^2+24*(-1/a*b)^(1/2)*x^3*b^4*c^3*d+105*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^4*d^4-298*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^3*b*c*d^3+353*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^2*b^2*c^2*d^2-208*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*b^3*c^3*d+48*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b^4*c^4+176*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^3*b*c*d^3-264*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^2*b^2*c^2*d^2+184*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*b^3*c^3*d-48*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b^4*c^4+122*(-1/a*b)^(1/2)*x*a^3*b*c*d^3-89*(-1/a*b)^(1/2)*x*a^2*b^2*c^2*d^2+24*(-1/a*b)^(1/2)*x*a*b^3*c^3*d)/d^4/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{7}{2}}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(7/2)/sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{7/2}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(7/2)/(c + d*x^2)^(1/2),x)

[Out] int((a + b*x^2)^(7/2)/(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{7}{2}}}{\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)**(7/2)/(d*x**2+c)**(1/2), x)
```

```
[Out] Integral((a + b*x**2)**(7/2)/sqrt(c + d*x**2), x)
```

$$3.200 \quad \int \frac{(a+bx^2)^{5/2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=344

$$\frac{\sqrt{c} \sqrt{a+bx^2} (15a^2d^2 - 11abcd + 4b^2c^2) \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + x\sqrt{a+bx^2} (23a^2d^2 - 23abcd + 8b^2c^2)}{15d^{5/2}\sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $1/15*(23*a^2*d^2-23*a*b*c*d+8*b^2*c^2)*x*(b*x^2+a)^{(1/2)}/d^2/(d*x^2+c)^{(1/2)}$
 $-1/15*(23*a^2*d^2-23*a*b*c*d+8*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}$
 $*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}$
 $(b*x^2+a)^{(1/2)}/d^{(5/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$
 $+1/15*(15*a^2*d^2-11*a*b*c*d+4*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}$
 $*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}$
 $(b*x^2+a)^{(1/2)}/d^{(5/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$
 $+1/5*b*x*(b*x^2+a)^{(3/2)}*(d*x^2+c)^{(1/2)}/d-4/15*b*(-2*a*d+b*c)*x*(b*x^2+a)^{(1/2)}$
 $(d*x^2+c)^{(1/2)}/d^2$

Rubi [A] time = 0.28, antiderivative size = 344, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {416, 528, 531, 418, 492, 411}

$$\frac{x\sqrt{a+bx^2} (23a^2d^2 - 23abcd + 8b^2c^2)}{15d^2\sqrt{c+dx^2}} + \frac{\sqrt{c} \sqrt{a+bx^2} (15a^2d^2 - 11abcd + 4b^2c^2) F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) \sqrt{c}}{15d^{5/2}\sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a + b*x^2)^{(5/2)}/\operatorname{Sqrt}[c + d*x^2], x]$

[Out] $((8*b^2*c^2 - 23*a*b*c*d + 23*a^2*d^2)*x*\operatorname{Sqrt}[a + b*x^2])/(15*d^2*\operatorname{Sqrt}[c + d*x^2]) - (4*b*(b*c - 2*a*d)*x*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2])/(15*d^2) + (b*x*(a + b*x^2)^{(3/2)}*\operatorname{Sqrt}[c + d*x^2])/(5*d) - (\operatorname{Sqrt}[c]*(8*b^2*c^2 - 23*a*b*c*d + 23*a^2*d^2)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(15*d^{(5/2)}*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (\operatorname{Sqrt}[c]*(4*b^2*c^2 - 11*a*b*c*d + 15*a^2*d^2)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(15*d^{(5/2)}*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

$\operatorname{Int}[\operatorname{Sqrt}[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^{(3/2)}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[\operatorname{Rt}[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*\operatorname{Rt}[d/c, 2]*\operatorname{Sqrt}[c + d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{PosQ}[b/a] \&\& \operatorname{PosQ}[d/c]$

Rule 416

$\operatorname{Int}[(a_) + (b_.)*(x_)^{(n_)}]^{(p_)}*((c_) + (d_.)*(x_)^{(n_)})^{(q_)}, x_Symbol] \rightarrow \operatorname{Simp}[(d*x*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-1)})/(b*(n*(p+q) + 1)), x] + \operatorname{Dist}[1/(b*(n*(p+q) + 1)), \operatorname{Int}[(a + b*x^n)^p*(c + d*x^n)^{(q-2)}*\operatorname{Simp}[c*(b*c*(n*(p+q) + 1) - a*d) + d*(b*c*(n*(p+2*q-1) + 1) - a*d*(n*(q-1) + 1))*x^n, x], x] /; \operatorname{FreeQ}[\{a, b, c, d, n, p\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{GtQ}[q, 1] \&\& \operatorname{NeQ}[n*(p+q) + 1, 0] \&\& !\operatorname{IGtQ}[p, 1] \&\& \operatorname{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 418

```
Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
:= Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rule 528

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
f_)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rule 531

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
f_)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x],
x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c,
d, e, f, n, p, q}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{5/2}}{\sqrt{c + dx^2}} dx &= \frac{bx(a + bx^2)^{3/2} \sqrt{c + dx^2}}{5d} + \frac{\int \frac{\sqrt{a+bx^2}(-a(bc-5ad)-4b(bc-2ad)x^2)}{\sqrt{c+dx^2}} dx}{5d} \\ &= -\frac{4b(bc - 2ad)x\sqrt{a + bx^2} \sqrt{c + dx^2}}{15d^2} + \frac{bx(a + bx^2)^{3/2} \sqrt{c + dx^2}}{5d} + \frac{\int \frac{a(4b^2c^2 - 11abcd + 15a^2d^2) + b(8b^2c^2 - 23abcd + 23a^2d^2)}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx}{15d^2} \\ &= -\frac{4b(bc - 2ad)x\sqrt{a + bx^2} \sqrt{c + dx^2}}{15d^2} + \frac{bx(a + bx^2)^{3/2} \sqrt{c + dx^2}}{5d} + \frac{a(4b^2c^2 - 11abcd + 15a^2d^2)}{15d^2} \\ &= \frac{(8b^2c^2 - 23abcd + 23a^2d^2)x\sqrt{a + bx^2}}{15d^2\sqrt{c + dx^2}} - \frac{4b(bc - 2ad)x\sqrt{a + bx^2} \sqrt{c + dx^2}}{15d^2} + \frac{bx(a + bx^2)^{3/2}}{5d} \\ &= \frac{(8b^2c^2 - 23abcd + 23a^2d^2)x\sqrt{a + bx^2}}{15d^2\sqrt{c + dx^2}} - \frac{4b(bc - 2ad)x\sqrt{a + bx^2} \sqrt{c + dx^2}}{15d^2} + \frac{bx(a + bx^2)^{3/2}}{5d} \end{aligned}$$

Mathematica [C] time = 0.49, size = 260, normalized size = 0.76

$$\frac{-i\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}(15a^3d^3 - 34a^2bcd^2 + 27ab^2c^2d - 8b^3c^3)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) - ibc\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}}{15d^3\sqrt{\frac{b}{a}}\sqrt{a + bx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2)/Sqrt[c + d*x^2], x]

[Out] (b*Sqrt[b/a]*d*x*(a + b*x^2)*(c + d*x^2)*(-4*b*c + 11*a*d + 3*b*d*x^2) - I*b*c*(8*b^2*c^2 - 23*a*b*c*d + 23*a^2*d^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*(-8*b^3*c^3 + 27*a*b^2*c^2*d - 34*a^2*b*c*d^2 + 15*a^3*d^3)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(15*Sqrt[b/a]*d^3*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b^2x^4 + 2abx^2 + a^2)\sqrt{bx^2 + a}}{\sqrt{dx^2 + c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral((b^2*x^4 + 2*a*b*x^2 + a^2)*sqrt(b*x^2 + a)/sqrt(d*x^2 + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^(1/2), x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(5/2)/sqrt(d*x^2 + c), x)

maple [A] time = 0.03, size = 615, normalized size = 1.79

$$\sqrt{bx^2 + a} \sqrt{dx^2 + c} \left(3\sqrt{-\frac{b}{a}} b^3 d^3 x^7 + 14\sqrt{-\frac{b}{a}} a b^2 d^3 x^5 - \sqrt{-\frac{b}{a}} b^3 c d^2 x^5 + 11\sqrt{-\frac{b}{a}} a^2 b d^3 x^3 + 10\sqrt{-\frac{b}{a}} a b^2 c d^2 \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/2)/(d*x^2+c)^(1/2), x)

[Out] 1/15*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(3*(-1/a*b)^(1/2)*x^7*b^3*d^3+14*(-1/a*b)^(1/2)*x^5*a*b^2*d^3-(-1/a*b)^(1/2)*x^5*b^3*c*d^2+11*(-1/a*b)^(1/2)*x^3*a^2*b*d^3+10*(-1/a*b)^(1/2)*x^3*a*b^2*c*d^2-4*(-1/a*b)^(1/2)*x^3*b^3*c^2*d+15*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^3*d^3-34*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b*c*d^2+27*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b^2*c^2*d-8*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b^3*c^3+23*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b*c*d^2-23*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b^2*c^2*d+8*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b^3*c^3+11*(-1/a*b)^(1/2)*x*a^2*b*c*d^2-4*(-1/a*b)^(1/2)*x*a*b^2*c^2*d/d^3/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/2)/sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{5/2}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)/(c + d*x^2)^(1/2),x)

[Out] int((a + b*x^2)^(5/2)/(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{5/2}}{\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)/(d*x**2+c)**(1/2),x)

[Out] Integral((a + b*x**2)**(5/2)/sqrt(c + d*x**2), x)

$$3.201 \quad \int \frac{(a+bx^2)^{3/2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=260

$$\frac{\sqrt{c} \sqrt{a+bx^2} (bc-3ad) \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right) + 2\sqrt{c} \sqrt{a+bx^2} (bc-2ad) E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1-\frac{bc}{ad}\right)}{3d^{3/2} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{bx \sqrt{a+bx^2}}{3d^{3/2} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $-2/3*(-2*a*d+b*c)*x*(b*x^2+a)^{(1/2)}/d/(d*x^2+c)^{(1/2)}+2/3*(-2*a*d+b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c))^{(1/2)}, (1-b*c/a/d)^{(1/2))*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(3/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-1/3*(-3*a*d+b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(3/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/3*b*x*(b*x^2+a)^{(1/2)}*(d*x^2+c)^{(1/2)}/d$

Rubi [A] time = 0.16, antiderivative size = 260, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {416, 531, 418, 492, 411}

$$\frac{\sqrt{c} \sqrt{a+bx^2} (bc-3ad) F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1-\frac{bc}{ad}\right) + 2\sqrt{c} \sqrt{a+bx^2} (bc-2ad) E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1-\frac{bc}{ad}\right) + bx \sqrt{a+bx^2}}{3d^{3/2} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{bx \sqrt{a+bx^2}}{3d^{3/2} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)/Sqrt[c + d*x^2], x]

[Out] $(-2*(b*c - 2*a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(3*d*\operatorname{Sqrt}[c + d*x^2]) + (b*x*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2])/(3*d) + (2*\operatorname{Sqrt}[c]*(b*c - 2*a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(3*d^{(3/2)}*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) - (\operatorname{Sqrt}[c]*(b*c - 3*a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(3*d^{(3/2)}*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q)+1)), x] + Dist[1/(b*(n*(p+q)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q)+1) - a*d) + d*(b*c*(n*(p+2*q-1)+1) - a*d*(n*(q-1)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q)+1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

$\text{t}[d/c, 2] * \text{Sqrt}[c + d*x^2] * \text{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))], x] /;$ FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 492

$\text{Int}[(x_)^2/(\text{Sqrt}[(a_) + (b_.)*(x_)^2]*\text{Sqrt}[(c_) + (d_.)*(x_)^2]), x_Symbol]$
 $:= \text{Simp}[(x*\text{Sqrt}[a + b*x^2])/(b*\text{Sqrt}[c + d*x^2]), x] - \text{Dist}[c/b, \text{Int}[\text{Sqrt}[a + b*x^2]/(c + d*x^2)^{(3/2)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rule 531

$\text{Int}[(a_) + (b_.)*(x_)^{(n_)}]^{(p_.)*((c_) + (d_.)*(x_)^{(n_)})^{(q_.)*((e_) + (f_.)*(x_)^{(n_)}), x_Symbol]$:= Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x], x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, e, f, n, p, q}, x]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{3/2}}{\sqrt{c + dx^2}} dx &= \frac{bx\sqrt{a + bx^2}\sqrt{c + dx^2}}{3d} + \frac{\int \frac{-a(bc - 3ad) - 2b(bc - 2ad)x^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{3d} \\ &= \frac{bx\sqrt{a + bx^2}\sqrt{c + dx^2}}{3d} - \frac{(a(bc - 3ad)) \int \frac{1}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{3d} - \frac{(2b(bc - 2ad)) \int \frac{x^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{3d} \\ &= -\frac{2(bc - 2ad)x\sqrt{a + bx^2}}{3d\sqrt{c + dx^2}} + \frac{bx\sqrt{a + bx^2}\sqrt{c + dx^2}}{3d} - \frac{\sqrt{c}(bc - 3ad)\sqrt{a + bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\right)}{3d^{3/2}\sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}}\sqrt{c + dx^2}} \\ &= -\frac{2(bc - 2ad)x\sqrt{a + bx^2}}{3d\sqrt{c + dx^2}} + \frac{bx\sqrt{a + bx^2}\sqrt{c + dx^2}}{3d} + \frac{2\sqrt{c}(bc - 2ad)\sqrt{a + bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\right)}{3d^{3/2}\sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}}\sqrt{c + dx^2}} \end{aligned}$$

Mathematica [C] time = 0.35, size = 216, normalized size = 0.83

$$\frac{-i\sqrt{\frac{bx^2}{a} + 1}\sqrt{\frac{dx^2}{c} + 1}(3a^2d^2 - 5abcd + 2b^2c^2)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) + bdx\sqrt{\frac{b}{a}}(a + bx^2)(c + dx^2) - 2il}{3d^2\sqrt{\frac{b}{a}}\sqrt{a + bx^2}\sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)/Sqrt[c + d*x^2], x]

[Out] (b*Sqrt[b/a]*d*x*(a + b*x^2)*(c + d*x^2) - (2*I)*b*c*(-(b*c) + 2*a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*(2*b^2*c^2 - 5*a*b*c*d + 3*a^2*d^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(3*Sqrt[b/a]*d^2*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(bx^2 + a)^{\frac{3}{2}}}{\sqrt{dx^2 + c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral((b*x^2 + a)^(3/2)/sqrt(d*x^2 + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(3/2)/sqrt(d*x^2 + c), x)

maple [A] time = 0.03, size = 399, normalized size = 1.53

$$\sqrt{bx^2 + a} \sqrt{dx^2 + c} \left(\sqrt{-\frac{b}{a}} b^2 d^2 x^5 + \sqrt{-\frac{b}{a}} ab d^2 x^3 + \sqrt{-\frac{b}{a}} b^2 cd x^3 + 3\sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} a^2 d^2 \text{EllipticF}\left(\sqrt{-\frac{b}{a}} x, \frac{a}{b/c*d}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)/(d*x^2+c)^(1/2),x)

[Out] 1/3*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*((-1/a*b)^(1/2)*x^5*b^2*d^2+(-1/a*b)^(1/2)*x^3*a*b*d^2+(-1/a*b)^(1/2)*x^3*b^2*c*d+3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*d^2-5*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b*c*d+2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b^2*c^2+4*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b*c*d-2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b^2*c^2+(-1/a*b)^(1/2)*x*a*b*c*d/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/d^2/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/2)/sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{3/2}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)/(c + d*x^2)^(1/2),x)

[Out] int((a + b*x^2)^(3/2)/(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{3}{2}}}{\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)/(d*x**2+c)**(1/2), x)

[Out] Integral((a + b*x**2)**(3/2)/sqrt(c + d*x**2), x)

3.202 $\int \frac{\sqrt{a+bx^2}}{\sqrt{c+dx^2}} dx$

Optimal. Leaf size=194

$$\frac{\sqrt{c} \sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{x\sqrt{a+bx^2}}{\sqrt{c+dx^2}} - \frac{\sqrt{c} \sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $x*(b*x^2+a)^{(1/2)}/(d*x^2+c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*E$
 $llipticE(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*$
 $x^2+a)^{(1/2)}/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+(1/(1+$
 $d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*EllipticF(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{($
 $1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x$
 $^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.09, antiderivative size = 194, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {422, 418, 492, 411}

$$\frac{x\sqrt{a+bx^2}}{\sqrt{c+dx^2}} + \frac{\sqrt{c} \sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{\sqrt{c} \sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x]

[Out] $(x*\operatorname{Sqrt}[a + b*x^2])/ \operatorname{Sqrt}[c + d*x^2] - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/((c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/((a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a+bx^2}}{\sqrt{c+dx^2}} dx &= a \int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx + b \int \frac{x^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx \\ &= \frac{x\sqrt{a+bx^2}}{\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} - c \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^{3/2}} dx \\ &= \frac{x\sqrt{a+bx^2}}{\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 86, normalized size = 0.44

$$\frac{\sqrt{a+bx^2}\sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{-\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{-\frac{d}{c}}\sqrt{\frac{a+bx^2}{a}}\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x]
```

```
[Out] (Sqrt[a + b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], (b*c)/(a*d)))/(Sqrt[-(d/c)]*Sqrt[(a + b*x^2)/a]*Sqrt[c + d*x^2])
```

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+a}}{\sqrt{dx^2+c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*x^2 + a)/sqrt(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*x^2 + a)/sqrt(d*x^2 + c), x)
```

maple [A] time = 0.02, size = 158, normalized size = 0.81

$$\frac{\sqrt{bx^2+a} \sqrt{dx^2+c} \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} \left(ad \operatorname{EllipticF} \left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}} \right) + bc \operatorname{EllipticE} \left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}} \right) - bc \operatorname{EllipticF} \left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}} \right) \right)}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{b}{a}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x)

[Out] (b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*(a*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*d-b*c*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))+b*c*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2)))/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)/sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)/(c + d*x^2)^(1/2),x)

[Out] int((a + b*x^2)^(1/2)/(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a+bx^2}}{\sqrt{c+dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)/(d*x**2+c)**(1/2),x)

[Out] Integral(sqrt(a + b*x**2)/sqrt(c + d*x**2), x)

$$3.203 \quad \int \frac{1}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=87

$$\frac{\sqrt{c} \sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{a\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/a/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {418}

$$\frac{\sqrt{c} \sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]),x]

[Out] $(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx = \frac{\sqrt{c} \sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} \sqrt{c+dx^2}}$$

Mathematica [A] time = 0.06, size = 86, normalized size = 0.99

$$\frac{\sqrt{\frac{a+bx^2}{a}} \sqrt{\frac{c+dx^2}{c}} \operatorname{EllipticF}\left(\sin^{-1}\left(x\sqrt{-\frac{b}{a}}\right), \frac{ad}{bc}\right)}{\sqrt{-\frac{b}{a}} \sqrt{a+bx^2} \sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]),x]

[Out] $(\operatorname{Sqrt}[(a + b*x^2)/a]*\operatorname{Sqrt}[(c + d*x^2)/c]*\operatorname{EllipticF}[\operatorname{ArcSin}[\operatorname{Sqrt}[-(b/a)]*x], (a*d)/(b*c)])/(\operatorname{Sqrt}[-(b/a)]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2])$

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+a}\sqrt{dx^2+c}}{bdx^4+(bc+ad)x^2+ac}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b*d*x^4 + (b*c + a*d)*x^2 + a*c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2+a}\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)), x)

maple [A] time = 0.03, size = 100, normalized size = 1.15

$$\frac{\sqrt{\frac{dx^2+c}{c}} \sqrt{\frac{bx^2+a}{a}} \sqrt{bx^2+a} \sqrt{dx^2+c} \text{EllipticF}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right)}{\sqrt{-\frac{b}{a}} (bdx^4 + adx^2 + bcx^2 + ac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x)

[Out] EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/(-1/a*b)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2+a}\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{bx^2+a}\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)),x)

[Out] int(1/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x**2+a)**(1/2)/(d*x**2+c)**(1/2), x)
```

```
[Out] Integral(1/(sqrt(a + b*x**2)*sqrt(c + d*x**2)), x)
```


$$3.204 \quad \int \frac{1}{(a+bx^2)^{3/2} \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=273

$$\frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{a\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{bx\sqrt{c+dx^2}}{a\sqrt{a+bx^2}(bc-ad)} - \frac{dx\sqrt{a+bx^2}}{a\sqrt{c+dx^2}(bc-ad)} + \frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2}}{a\sqrt{c+dx^2}(bc-ad)}$$

[Out] $-d*x*(b*x^2+a)^{(1/2)}/a/(-a*d+b*c)/(d*x^2+c)^{(1/2)}+(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a/(-a*d+b*c)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a/(-a*d+b*c)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+b*x*(d*x^2+c)^{(1/2)}/a/(-a*d+b*c)/(b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.14, antiderivative size = 273, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {414, 21, 422, 418, 492, 411}

$$\frac{bx\sqrt{c+dx^2}}{a\sqrt{a+bx^2}(bc-ad)} - \frac{dx\sqrt{a+bx^2}}{a\sqrt{c+dx^2}(bc-ad)} - \frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{c+dx^2}(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(3/2)*Sqrt[c + d*x^2]),x]

[Out] $-((d*x*\operatorname{Sqrt}[a + b*x^2])/(a*(b*c - a*d)*\operatorname{Sqrt}[c + d*x^2])) + (b*x*\operatorname{Sqrt}[c + d*x^2])/(a*(b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]) + (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*(b*c - a*d)*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*(b*c - a*d)*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 21

Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])

Rule 411

Int[Sqrt[(a_) + (b_)*(x_)^2]/((c_) + (d_)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 414

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1]

&& !(!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)]/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] :> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a + b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a + bx^2)^{3/2} \sqrt{c + dx^2}} dx &= \frac{bx\sqrt{c + dx^2}}{a(bc - ad)\sqrt{a + bx^2}} - \frac{\int \frac{ad + bdx^2}{\sqrt{a + bx^2} \sqrt{c + dx^2}} dx}{a(bc - ad)} \\
 &= \frac{bx\sqrt{c + dx^2}}{a(bc - ad)\sqrt{a + bx^2}} - \frac{d \int \frac{\sqrt{a + bx^2}}{\sqrt{c + dx^2}} dx}{a(bc - ad)} \\
 &= \frac{bx\sqrt{c + dx^2}}{a(bc - ad)\sqrt{a + bx^2}} - \frac{d \int \frac{1}{\sqrt{a + bx^2} \sqrt{c + dx^2}} dx}{bc - ad} - \frac{(bd) \int \frac{x^2}{\sqrt{a + bx^2} \sqrt{c + dx^2}} dx}{a(bc - ad)} \\
 &= -\frac{dx\sqrt{a + bx^2}}{a(bc - ad)\sqrt{c + dx^2}} + \frac{bx\sqrt{c + dx^2}}{a(bc - ad)\sqrt{a + bx^2}} - \frac{\sqrt{c} \sqrt{d} \sqrt{a + bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1\right)}{a(bc - ad) \sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}} \sqrt{c + dx^2}} \\
 &= -\frac{dx\sqrt{a + bx^2}}{a(bc - ad)\sqrt{c + dx^2}} + \frac{bx\sqrt{c + dx^2}}{a(bc - ad)\sqrt{a + bx^2}} + \frac{\sqrt{c} \sqrt{d} \sqrt{a + bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1\right)}{a(bc - ad) \sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}} \sqrt{c + dx^2}}
 \end{aligned}$$

Mathematica [A] time = 0.25, size = 112, normalized size = 0.41

$$\frac{ad \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{dx^2}{c} + 1} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}} x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}}} - bx(c + dx^2)$$

$$\frac{\hspace{10em}}{a\sqrt{a + bx^2} \sqrt{c + dx^2} (ad - bc)}$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^(3/2)*Sqrt[c + d*x^2]),x]

[Out] $(-(b*x*(c + d*x^2)) + (a*d*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], (b*c)/(a*d)]/Sqrt[-(d/c)])/(a*(-(b*c) + a*d)*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])$

fricas [F] time = 0.72, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2 + a} \sqrt{dx^2 + c}}{b^2 dx^6 + (b^2 c + 2 abd)x^4 + a^2 c + (2 abc + a^2 d)x^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b^2*d*x^6 + (b^2*c + 2*a*b*d)*x^4 + a^2*c + (2*a*b*c + a^2*d)*x^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{2}} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(3/2)*sqrt(d*x^2 + c)), x)

maple [A] time = 0.04, size = 248, normalized size = 0.91

$$\frac{\left(-\sqrt{-\frac{b}{a}} bd x^3 + \sqrt{\frac{dx^2+c}{c}} \sqrt{\frac{bx^2+a}{a}} ad \text{EllipticF}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right) - \sqrt{-\frac{b}{a}} bcx + \sqrt{\frac{dx^2+c}{c}} \sqrt{\frac{bx^2+a}{a}} bc \text{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right)\right)}{\sqrt{-\frac{b}{a}} (ad - bc) (bd x^4 + ad x^2 + bc x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(3/2)/(d*x^2+c)^(1/2),x)

[Out] $(-x^3*b*d*(-1/a*b)^(1/2)+EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*d*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b*c*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b*c*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-x*b*c*(-1/a*b)^(1/2)*(d*x^2+c)^(1/2)*(b*x^2+a)^(1/2)/(-1/a*b)^(1/2)/a/(a*d-b*c)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{2}} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(3/2)*sqrt(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{3/2} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((a + b*x^2)^(3/2)*(c + d*x^2)^(1/2)), x)`

[Out] `int(1/((a + b*x^2)^(3/2)*(c + d*x^2)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{3}{2}} \sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x**2+a)**(3/2)/(d*x**2+c)**(1/2), x)`

[Out] `Integral(1/((a + b*x**2)**(3/2)*sqrt(c + d*x**2)), x)`

$$3.205 \quad \int \frac{1}{(a+bx^2)^{5/2} \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=255

$$\frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} (bc-3ad) \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right) + 2\sqrt{b} \sqrt{c+dx^2} (bc-2ad) E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 1-\frac{ad}{bc}\right)}{3a^2 \sqrt{c+dx^2} (bc-ad)^2 \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} + 3a^{3/2} \sqrt{a+bx^2} (bc-ad)^2 \sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}}}$$

[Out] $-1/3*(-3*a*d+b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a^2/(-a*d+b*c)^2/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/3*b*x*(d*x^2+c)^{(1/2)}/a/(-a*d+b*c)/(b*x^2+a)^{(3/2)}+2/3*(-2*a*d+b*c)*(1/(1+b*x^2/a))^{(1/2)}*(1+b*x^2/a)^{(1/2)}*\operatorname{EllipticE}(x*b^{(1/2)}/a^{(1/2)}/(1+b*x^2/a)^{(1/2)},(1-a*d/b/c)^{(1/2)})*b^{(1/2)}*(d*x^2+c)^{(1/2)}/a^{(3/2)}/(-a*d+b*c)^2/(b*x^2+a)^{(1/2)}/(a*(d*x^2+c)/c/(b*x^2+a))^{(1/2)}$

Rubi [A] time = 0.15, antiderivative size = 255, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {414, 525, 418, 411}

$$\frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} (bc-3ad) F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1-\frac{bc}{ad}\right) + 2\sqrt{b} \sqrt{c+dx^2} (bc-2ad) E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 1-\frac{ad}{bc}\right)}{3a^2 \sqrt{c+dx^2} (bc-ad)^2 \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} + 3a^{3/2} \sqrt{a+bx^2} (bc-ad)^2 \sqrt{\frac{a(c+dx^2)}{c(a+bx^2)}} + 3a(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(5/2)*Sqrt[c + d*x^2]),x]

[Out] $(b*x*\operatorname{Sqrt}[c+d*x^2])/(3*a*(b*c-a*d)*(a+b*x^2)^{(3/2)})+(2*\operatorname{Sqrt}[b]*(b*c-2*a*d)*\operatorname{Sqrt}[c+d*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[\operatorname{Sqrt}[b]*x]/\operatorname{Sqrt}[a]],1-(a*d)/(b*c))/(3*a^{(3/2)}*(b*c-a*d)^2*\operatorname{Sqrt}[a+b*x^2]*\operatorname{Sqrt}[(a*(c+d*x^2))/(c*(a+b*x^2))])-(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*(b*c-3*a*d)*\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[\operatorname{Sqrt}[d]*x]/\operatorname{Sqrt}[c]],1-(b*c)/(a*d))/(3*a^2*(b*c-a*d)^2*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2))]*\operatorname{Sqrt}[c+d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(p+1)*(b*c - a*d)), x] + Dist[1/(a*n*(p+1)*(b*c - a*d)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]), x]

$\text{t}[d/c, 2] * \text{Sqrt}[c + d*x^2] * \text{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{PosQ}[d/c] \&\& \text{PosQ}[b/a] \&\& !\text{SimplerSqrtQ}[b/a, d/c]$

Rule 525

$\text{Int}[(e_ + (f_)*(x_)^2)/(\text{Sqrt}[(a_ + (b_)*(x_)^2]*((c_ + (d_)*(x_)^2)^{(3/2)})), x_Symbol] := \text{Dist}[(b*e - a*f)/(b*c - a*d), \text{Int}[1/(\text{Sqrt}[a + b*x^2]*\text{Sqrt}[c + d*x^2]), x], x] - \text{Dist}[(d*e - c*f)/(b*c - a*d), \text{Int}[\text{Sqrt}[a + b*x^2]/(c + d*x^2)^{(3/2)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \text{PosQ}[b/a] \&\& \text{PosQ}[d/c]$

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + bx^2)^{5/2} \sqrt{c + dx^2}} dx &= \frac{bx\sqrt{c + dx^2}}{3a(bc - ad)(a + bx^2)^{3/2}} - \frac{\int \frac{-2bc + 3ad - bdx^2}{(a + bx^2)^{3/2} \sqrt{c + dx^2}} dx}{3a(bc - ad)} \\ &= \frac{bx\sqrt{c + dx^2}}{3a(bc - ad)(a + bx^2)^{3/2}} - \frac{(d(bc - 3ad)) \int \frac{1}{\sqrt{a + bx^2} \sqrt{c + dx^2}} dx}{3a(bc - ad)^2} + \frac{(2b(bc - 2ad)) \int \frac{\sqrt{c + dx^2}}{a + bx^2} dx}{3a(bc - ad)^2} \\ &= \frac{bx\sqrt{c + dx^2}}{3a(bc - ad)(a + bx^2)^{3/2}} + \frac{2\sqrt{b}(bc - 2ad)\sqrt{c + dx^2} E\left(\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 1 - \frac{ad}{bc}\right)}{3a^{3/2}(bc - ad)^2 \sqrt{a + bx^2} \sqrt{\frac{a(c + dx^2)}{c(a + bx^2)}}} - \frac{\sqrt{c}}{3a(bc - ad)} \end{aligned}$$

Mathematica [C] time = 0.61, size = 261, normalized size = 1.02

$$\frac{-i(a + bx^2) \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{dx^2}{c} + 1} (3a^2d^2 - 5abcd + 2b^2c^2) \text{EllipticF}\left(i \sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) + bx\sqrt{\frac{b}{a}}(c + dx^2) (-5a^2d^2 + 2b^2c^2)}{3a^2\sqrt{\frac{b}{a}}(a + bx^2)^{3/2}\sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^(5/2)*Sqrt[c + d*x^2]),x]

[Out] (b*Sqrt[b/a]*x*(c + d*x^2)*(-5*a^2*d + 2*b^2*c*x^2 + a*b*(3*c - 4*d*x^2)) - (2*I)*b*c*(-(b*c) + 2*a*d)*(a + b*x^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*(2*b^2*c^2 - 5*a*b*c*d + 3*a^2*d^2)*(a + b*x^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(3*a^2*Sqrt[b/a]*(b*c - a*d)^2*(a + b*x^2)^(3/2)*Sqrt[c + d*x^2])

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2 + a} \sqrt{dx^2 + c}}{b^3 dx^8 + (b^3 c + 3 ab^2 d)x^6 + 3(ab^2 c + a^2 bd)x^4 + a^3 c + (3a^2 bc + a^3 d)x^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b^3*d*x^8 + (b^3*c + 3*a*b^2*d)*x^6 + 3*(a*b^2*c + a^2*b*d)*x^4 + a^3*c + (3*a^2*b*c + a^3*d)*x^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{2}} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(5/2)*sqrt(d*x^2 + c)), x)

maple [B] time = 0.06, size = 752, normalized size = 2.95

$$-4\sqrt{-\frac{b}{a}} a b^2 d^2 x^5 + 2\sqrt{-\frac{b}{a}} b^3 c d x^5 - 5\sqrt{-\frac{b}{a}} a^2 b d^2 x^3 + 3\sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} a^2 b d^2 x^2 \text{EllipticF}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right) -$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(5/2)/(d*x^2+c)^(1/2),x)

[Out] 1/3*(-4*x^5*a*b^2*d^2*(-1/a*b)^(1/2)+2*x^5*b^3*c*d*(-1/a*b)^(1/2)+3*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*x^2*a^2*b*d^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-5*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*x^2*a*b^2*c*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+2*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*x^2*b^3*c^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+4*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*x^2*a*b^2*c*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-2*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*x^2*b^3*c^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-5*x^3*a^2*b*d^2*(-1/a*b)^(1/2)-x^3*a*b^2*c*d*(-1/a*b)^(1/2)+2*x^3*b^3*c^2*(-1/a*b)^(1/2)+3*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^3*d^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-5*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^2*b*c*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+2*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*b^2*c^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)+4*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^2*b*c*d*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-2*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*b^2*c^2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)-5*x*a^2*b*c*d*(-1/a*b)^(1/2)+3*x*a*b^2*c^2*(-1/a*b)^(1/2))/(d*x^2+c)^(1/2)/(a*d-b*c)^2/a^2/(-1/a*b)^(1/2)/(b*x^2+a)^(3/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{2}} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(5/2)*sqrt(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{5/2} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)^(1/2)),x)

[Out] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{5}{2}} \sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(5/2)/(d*x**2+c)**(1/2), x)

[Out] Integral(1/((a + b*x**2)**(5/2)*sqrt(c + d*x**2)), x)

$$3.206 \quad \int \frac{1}{(a+bx^2)^{7/2} \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=334

$$\frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} (15a^2d^2 - 11abcd + 4b^2c^2) \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + \frac{4bx\sqrt{c+dx^2}(bc-2ad)}{15a^2(a+bx^2)^{3/2}(bc-ad)^2}}{15a^3\sqrt{c+dx^2}(bc-ad)^3 \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $-1/15*(15*a^2*d^2-11*a*b*c*d+4*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a^3/(-a*d+b*c)^3/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/5*b*x*(d*x^2+c)^{(1/2)}/a/(-a*d+b*c)/(b*x^2+a)^{(5/2)}+4/15*b*(-2*a*d+b*c)*x*(d*x^2+c)^{(1/2)}/a^2/(-a*d+b*c)^2/(b*x^2+a)^{(3/2)}+1/15*(23*a^2*d^2-23*a*b*c*d+8*b^2*c^2)*(1/(1+b*x^2/a))^{(1/2)}*(1+b*x^2/a)^{(1/2)}*\operatorname{EllipticE}(x*b^{(1/2)}/a^{(1/2)}/(1+b*x^2/a)^{(1/2)},(1-a*d/b/c)^{(1/2)})*b^{(1/2)}*(d*x^2+c)^{(1/2)}/a^{(5/2)}/(-a*d+b*c)^3/(b*x^2+a)^{(1/2)}/(a*(d*x^2+c)/c/(b*x^2+a))^{(1/2)}$

Rubi [A] time = 0.28, antiderivative size = 334, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {414, 527, 525, 418, 411}

$$\frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} (15a^2d^2 - 11abcd + 4b^2c^2) F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) + \sqrt{b} \sqrt{c+dx^2} (23a^2d^2 - 23abcd + 8b^2c^2)}{15a^3\sqrt{c+dx^2}(bc-ad)^3 \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} + 15a^{5/2}\sqrt{a+bx^2}(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(7/2)*Sqrt[c + d*x^2]),x]

[Out] $(b*x*\operatorname{Sqrt}[c+d*x^2])/(5*a*(b*c-a*d)*(a+b*x^2)^{(5/2)})+(4*b*(b*c-2*a*d)*x*\operatorname{Sqrt}[c+d*x^2])/(15*a^2*(b*c-a*d)^2*(a+b*x^2)^{(3/2)})+(\operatorname{Sqrt}[b]*(8*b^2*c^2-23*a*b*c*d+23*a^2*d^2)*\operatorname{Sqrt}[c+d*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]],1-(a*d)/(b*c)])/(15*a^{(5/2)}*(b*c-a*d)^3*\operatorname{Sqrt}[a+b*x^2]*\operatorname{Sqrt}[(a*(c+d*x^2))/(c*(a+b*x^2))])-(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*(4*b^2*c^2-11*a*b*c*d+15*a^2*d^2)*\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]],1-(b*c)/(a*d)])/(15*a^3*(b*c-a*d)^3*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2))]*\operatorname{Sqrt}[c+d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 525

```
Int[((e_) + (f_.)*(x_)^2)/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)^(
3/2)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(Sqrt[a + b*x^2]*S
qrt[c + d*x^2]), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[Sqrt[a + b*x^2]
/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && PosQ[b/a] &&
PosQ[d/c]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\int \frac{1}{(a + bx^2)^{7/2} \sqrt{c + dx^2}} dx = \frac{bx\sqrt{c + dx^2}}{5a(bc - ad)(a + bx^2)^{5/2}} - \frac{\int \frac{-4bc+5ad-3bdx^2}{(a+bx^2)^{5/2} \sqrt{c+dx^2}} dx}{5a(bc - ad)}$$

$$= \frac{bx\sqrt{c + dx^2}}{5a(bc - ad)(a + bx^2)^{5/2}} + \frac{4b(bc - 2ad)x\sqrt{c + dx^2}}{15a^2(bc - ad)^2(a + bx^2)^{3/2}} + \frac{\int \frac{8b^2c^2-19abcd+15a^2d^2+4bd(b^2c-d^2)}{(a+bx^2)^{3/2} \sqrt{c+dx^2}} dx}{15a^2(bc - ad)^2}$$

$$= \frac{bx\sqrt{c + dx^2}}{5a(bc - ad)(a + bx^2)^{5/2}} + \frac{4b(bc - 2ad)x\sqrt{c + dx^2}}{15a^2(bc - ad)^2(a + bx^2)^{3/2}} - \frac{d(4b^2c^2 - 11abcd + 15a^2d^2)}{15a^2(bc - ad)^2}$$

$$= \frac{bx\sqrt{c + dx^2}}{5a(bc - ad)(a + bx^2)^{5/2}} + \frac{4b(bc - 2ad)x\sqrt{c + dx^2}}{15a^2(bc - ad)^2(a + bx^2)^{3/2}} + \frac{\sqrt{b}(8b^2c^2 - 23abcd + 23a^2d^2)}{15a^{5/2}(bc - ad)}$$

Mathematica [C] time = 0.64, size = 301, normalized size = 0.90

$$\frac{bx\sqrt{\frac{b}{a}}(c + dx^2)\left((a + bx^2)^2(23a^2d^2 - 23abcd + 8b^2c^2) + 3a^2(bc - ad)^2 + 4a(a + bx^2)(bc - 2ad)(bc - ad)\right) + i\sqrt{\dots}}{\dots}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/((a + b*x^2)^(7/2)*Sqrt[c + d*x^2]), x]
[Out] (b*Sqrt[b/a]*x*(c + d*x^2)*(3*a^2*(b*c - a*d)^2 + 4*a*(b*c - 2*a*d)*(b*c -
a*d)*(a + b*x^2) + (8*b^2*c^2 - 23*a*b*c*d + 23*a^2*d^2)*(a + b*x^2)^2) + I
*(a + b*x^2)^2*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*(b*c*(8*b^2*c^2 - 23
```

$*a*b*c*d + 23*a^2*d^2)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[b/a]*x], (a*d)/(b*c)] + (-8*b^3*c^3 + 27*a*b^2*c^2*d - 34*a^2*b*c*d^2 + 15*a^3*d^3)*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[b/a]*x], (a*d)/(b*c)))/(15*a^3*\text{Sqrt}[b/a]*(b*c - a*d)^3*(a + b*x^2)^{(5/2)*\text{Sqrt}[c + d*x^2]}$

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2 + a}\sqrt{dx^2 + c}}{b^4dx^{10} + (b^4c + 4ab^3d)x^8 + 2(2ab^3c + 3a^2b^2d)x^6 + a^4c + 2(3a^2b^2c + 2a^3bd)x^4 + (4a^3bc + a^4d)x^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b^4*d*x^10 + (b^4*c + 4*a*b^3*d)*x^8 + 2*(2*a*b^3*c + 3*a^2*b^2*d)*x^6 + a^4*c + 2*(3*a^2*b^2*c + 2*a^3*b*d)*x^4 + (4*a^3*b*c + a^4*d)*x^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{7}{2}}\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(7/2)*sqrt(d*x^2 + c)), x)

maple [B] time = 0.07, size = 1607, normalized size = 4.81

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(7/2)/(d*x^2+c)^(1/2),x)

[Out] $1/15*(-23*x^7*a^2*b^3*d^3*(-1/a*b)^{(1/2)} - 13*x^3*a^3*b^2*c*d^2*(-1/a*b)^{(1/2)} - 8*x^5*b^5*c^3*(-1/a*b)^{(1/2)} + 43*x^3*a^2*b^3*c^2*d*(-1/a*b)^{(1/2)} - 34*x*a^4*b*c*d^2*(-1/a*b)^{(1/2)} + 23*x^7*a*b^4*c*d^2*(-1/a*b)^{(1/2)} + 41*x*a^3*b^2*c^2*d*(-1/a*b)^{(1/2)} + 35*x^5*a^2*b^3*c*d^2*(-1/a*b)^{(1/2)} + 3*x^5*a*b^4*c^2*d*(-1/a*b)^{(1/2)} + 8*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*b^5*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 8*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*b^5*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 8*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^2*b^3*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 46*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a^3*b^2*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 8*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^2*b^3*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 8*x^7*b^5*c^2*d*(-1/a*b)^{(1/2)} - 54*x^5*a^3*b^2*d^3*(-1/a*b)^{(1/2)} - 34*x^3*a^4*b*d^3*(-1/a*b)^{(1/2)} - 20*x^3*a*b^4*c^3*(-1/a*b)^{(1/2)} - 15*x*a^2*b^3*c^3*(-1/a*b)^{(1/2)} + 15*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^5*d^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 15*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^4*a^3*b^2*d^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 16*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a*b^4*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 30*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a^4*b*d^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 16*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*x^2*a*b^4*c^3*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 23*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^4*b*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 23*\text{EllipticE}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^3*b^2*c^2*d*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} - 34*\text{EllipticF}((-1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})*a^4*b*c*d^2*((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)} + 27*\text{EllipticF}((-1/a*b)^{(1/2)}*x, ($

$$\frac{a/b/c*d)^{(1/2)}*a^3*b^2*c^2*d*((b*x^2+a)/a)^{(1/2)*((d*x^2+c)/c)^{(1/2)}-46*EllipticE((-1/a*b)^{(1/2)*x, (a/b/c*d)^{(1/2)})*x^2*a^2*b^3*c^2*d*((b*x^2+a)/a)^{(1/2)*((d*x^2+c)/c)^{(1/2)}-68*EllipticF((-1/a*b)^{(1/2)*x, (a/b/c*d)^{(1/2)})*x^2*a^3*b^2*c*d^2*((b*x^2+a)/a)^{(1/2)*((d*x^2+c)/c)^{(1/2)}+54*EllipticF((-1/a*b)^{(1/2)*x, (a/b/c*d)^{(1/2)})*x^2*a^2*b^3*c^2*d*((b*x^2+a)/a)^{(1/2)*((d*x^2+c)/c)^{(1/2)}+23*EllipticE((-1/a*b)^{(1/2)*x, (a/b/c*d)^{(1/2)})*x^4*a^2*b^3*c*d^2*((b*x^2+a)/a)^{(1/2)*((d*x^2+c)/c)^{(1/2)}-23*EllipticE((-1/a*b)^{(1/2)*x, (a/b/c*d)^{(1/2)})*x^4*a*b^4*c^2*d*((b*x^2+a)/a)^{(1/2)*((d*x^2+c)/c)^{(1/2)}-34*EllipticF((-1/a*b)^{(1/2)*x, (a/b/c*d)^{(1/2)})*x^4*a^2*b^3*c*d^2*((b*x^2+a)/a)^{(1/2)*((d*x^2+c)/c)^{(1/2)}+27*EllipticF((-1/a*b)^{(1/2)*x, (a/b/c*d)^{(1/2)})*x^4*a*b^4*c^2*d*((b*x^2+a)/a)^{(1/2)*((d*x^2+c)/c)^{(1/2)}}/(d*x^2+c)^{(1/2)/(a*d-b*c)^3/(-1/a*b)^{(1/2)/a^3/(b*x^2+a)^{(5/2)}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{7}{2}} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(7/2)*sqrt(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{7}{2}} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(7/2)*(c + d*x^2)^(1/2)),x)

[Out] int(1/((a + b*x^2)^(7/2)*(c + d*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{7}{2}} \sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(7/2)/(d*x**2+c)**(1/2),x)

[Out] Integral(1/((a + b*x**2)**(7/2)*sqrt(c + d*x**2)), x)

$$3.207 \quad \int \frac{(a+bx^2)^{7/2}}{(c+dx^2)^{3/2}} dx$$

Optimal. Leaf size=445

$$\frac{b\sqrt{c}\sqrt{a+bx^2}(45a^2d^2 - 61abcd + 24b^2c^2)\text{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + bx\sqrt{a+bx^2}\sqrt{c+dx^2}(15a^2d^2 - 43abcd + 24b^2c^2)}{15d^{7/2}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} \quad 15cd^3$$

[Out] $-(-a*d+b*c)*x*(b*x^2+a)^{(5/2)}/c/d/(d*x^2+c)^{(1/2)}+1/15*(-15*a^3*d^3+103*a^2*b*c*d^2-128*a*b^2*c^2*d+48*b^3*c^3)*x*(b*x^2+a)^{(1/2)}/c/d^3/(d*x^2+c)^{(1/2)}$
 $-1/15*(-15*a^3*d^3+103*a^2*b*c*d^2-128*a*b^2*c^2*d+48*b^3*c^3)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\text{EllipticE}(x*d^{(1/2)}/c^{(1/2)})/(1+d*x^2/c)^{(1/2)}$
 $, (1-b*c/a/d)^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(7/2)}/c^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/15*b*(45*a^2*d^2-61*a*b*c*d+24*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\text{EllipticF}(x*d^{(1/2)}/c^{(1/2)})/(1+d*x^2/c)^{(1/2)}$
 $, (1-b*c/a/d)^{(1/2)}*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(7/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/5*b*(-5*a*d+6*b*c)*x*(b*x^2+a)^{(3/2)}*(d*x^2+c)^{(1/2)}/c/d^2-1/15*b*(15*a^2*d^2-43*a*b*c*d+24*b^2*c^2)*x*(b*x^2+a)^{(1/2)}*(d*x^2+c)^{(1/2)}/c/d^3$

Rubi [A] time = 0.41, antiderivative size = 445, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {413, 528, 531, 418, 492, 411}

$$\frac{bx\sqrt{a+bx^2}\sqrt{c+dx^2}(15a^2d^2 - 43abcd + 24b^2c^2)}{15cd^3} + \frac{x\sqrt{a+bx^2}(103a^2bcd^2 - 15a^3d^3 - 128ab^2c^2d + 48b^3c^3)}{15cd^3\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(7/2)/(c + d*x^2)^(3/2), x]

[Out] $((48*b^3*c^3 - 128*a*b^2*c^2*d + 103*a^2*b*c*d^2 - 15*a^3*d^3)*x*\text{Sqrt}[a + b*x^2])/(15*c*d^3*\text{Sqrt}[c + d*x^2]) - ((b*c - a*d)*x*(a + b*x^2)^{(5/2)})/(c*d*\text{Sqrt}[c + d*x^2]) - (b*(24*b^2*c^2 - 43*a*b*c*d + 15*a^2*d^2)*x*\text{Sqrt}[a + b*x^2]*\text{Sqrt}[c + d*x^2])/(15*c*d^3) + (b*(6*b*c - 5*a*d)*x*(a + b*x^2)^{(3/2)}*\text{Sqrt}[c + d*x^2])/(5*c*d^2) - ((48*b^3*c^3 - 128*a*b^2*c^2*d + 103*a^2*b*c*d^2 - 15*a^3*d^3)*\text{Sqrt}[a + b*x^2]*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]], 1 - (b*c)/(a*d)])/(15*\text{Sqrt}[c]*d^{(7/2)}*\text{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\text{Sqrt}[c + d*x^2]) + (b*\text{Sqrt}[c]*(24*b^2*c^2 - 61*a*b*c*d + 45*a^2*d^2)*\text{Sqrt}[a + b*x^2]*\text{EllipticF}[\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]], 1 - (b*c)/(a*d)])/(15*d^{(7/2)}*\text{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\text{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1), x]]

```
2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p
+ q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,
0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rule 418

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:= Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rule 531

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x],
x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c,
d, e, f, n, p, q}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^{7/2}}{(c+dx^2)^{3/2}} dx &= -\frac{(bc-ad)x(a+bx^2)^{5/2}}{cd\sqrt{c+dx^2}} + \frac{\int \frac{(a+bx^2)^{3/2}(abc+b(6bc-5ad)x^2)}{\sqrt{c+dx^2}} dx}{cd} \\
&= -\frac{(bc-ad)x(a+bx^2)^{5/2}}{cd\sqrt{c+dx^2}} + \frac{b(6bc-5ad)x(a+bx^2)^{3/2}\sqrt{c+dx^2}}{5cd^2} + \frac{\int \frac{\sqrt{a+bx^2}(-2abc(3bc-5ad))}{\sqrt{c+dx^2}} dx}{5cd^2} \\
&= -\frac{(bc-ad)x(a+bx^2)^{5/2}}{cd\sqrt{c+dx^2}} - \frac{b(24b^2c^2-43abcd+15a^2d^2)x\sqrt{a+bx^2}\sqrt{c+dx^2}}{15cd^3} + \frac{b(6bc-5ad)}{5cd^2} \\
&= -\frac{(bc-ad)x(a+bx^2)^{5/2}}{cd\sqrt{c+dx^2}} - \frac{b(24b^2c^2-43abcd+15a^2d^2)x\sqrt{a+bx^2}\sqrt{c+dx^2}}{15cd^3} + \frac{b(6bc-5ad)}{5cd^2} \\
&= \frac{(48b^3c^3-128ab^2c^2d+103a^2bcd^2-15a^3d^3)x\sqrt{a+bx^2}}{15cd^3\sqrt{c+dx^2}} - \frac{(bc-ad)x(a+bx^2)^{5/2}}{cd\sqrt{c+dx^2}} - \frac{b(24b^2c^2-43abcd+15a^2d^2)}{15cd^2} \\
&= \frac{(48b^3c^3-128ab^2c^2d+103a^2bcd^2-15a^3d^3)x\sqrt{a+bx^2}}{15cd^3\sqrt{c+dx^2}} - \frac{(bc-ad)x(a+bx^2)^{5/2}}{cd\sqrt{c+dx^2}} - \frac{b(24b^2c^2-43abcd+15a^2d^2)}{15cd^2}
\end{aligned}$$

Mathematica [C] time = 1.22, size = 318, normalized size = 0.71

$$4ibc\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}\left(-15a^3d^3+41a^2bcd^2-38ab^2c^2d+12b^3c^3\right)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right),\frac{ad}{bc}\right)+dx\sqrt{\frac{b}{a}}\left(a\right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(7/2)/(c + d*x^2)^(3/2), x]

[Out] (Sqrt[b/a]*d*x*(a + b*x^2)*(-45*a^2*b*c*d^2 + 15*a^3*d^3 + a*b^2*c*d*(61*c + 16*d*x^2) - 3*b^3*c*(8*c^2 + 2*c*d*x^2 - d^2*x^4)) + I*b*c*(-48*b^3*c^3 + 128*a*b^2*c^2*d - 103*a^2*b*c*d^2 + 15*a^3*d^3)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + (4*I)*b*c*(12*b^3*c^3 - 38*a*b^2*c^2*d + 41*a^2*b*c*d^2 - 15*a^3*d^3)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(15*Sqrt[b/a]*c*d^4*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b^3x^6+3ab^2x^4+3a^2bx^2+a^3)\sqrt{bx^2+a}\sqrt{dx^2+c}}{d^2x^4+2cdx^2+c^2},x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/2)/(d*x^2+c)^(3/2), x, algorithm="fricas")

[Out] integral((b^3*x^6 + 3*a*b^2*x^4 + 3*a^2*b*x^2 + a^3)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(d^2*x^4 + 2*c*d*x^2 + c^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{7}{2}}}{(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/2)/(d*x^2+c)^(3/2),x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(7/2)/(d*x^2 + c)^(3/2), x)

maple [A] time = 0.07, size = 755, normalized size = 1.70

$$\sqrt{bx^2 + a} \sqrt{dx^2 + c} \left(3\sqrt{-\frac{b}{a}} b^4 c d^3 x^7 + 19\sqrt{-\frac{b}{a}} a b^3 c d^3 x^5 - 6\sqrt{-\frac{b}{a}} b^4 c^2 d^2 x^5 + 15\sqrt{-\frac{b}{a}} a^3 b d^4 x^3 - 29\sqrt{-\frac{b}{a}} a^2 b^2 c \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(7/2)/(d*x^2+c)^(3/2),x)

[Out] 1/15*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(3*(-1/a*b)^(1/2)*b^4*c*d^3*x^7+19*(-1/a*b)^(1/2)*a*b^3*c*d^3*x^5-6*(-1/a*b)^(1/2)*b^4*c^2*d^2*x^5+15*(-1/a*b)^(1/2)*a^3*b*d^4*x^3-29*(-1/a*b)^(1/2)*a^2*b^2*c*d^3*x^3+55*(-1/a*b)^(1/2)*a*b^3*c^2*d^2*x^3-24*(-1/a*b)^(1/2)*b^4*c^3*d*x^3+60*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^3*b*c*d^3-164*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^2*b^2*c^2*d^2+152*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*b^3*c^3*d-48*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b^4*c^4-15*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^3*b*c*d^3+103*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a^2*b^2*c^2*d^2-128*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*b^3*c^3*d+48*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b^4*c^4+15*x*a^4*d^4*(-1/a*b)^(1/2)-45*(-1/a*b)^(1/2)*a^3*b*c*d^3*x+61*(-1/a*b)^(1/2)*a^2*b^2*c^2*d^2*x-24*(-1/a*b)^(1/2)*a*b^3*c^3*d*x)/d^4/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)/c

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{7}{2}}}{(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/2)/(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(7/2)/(d*x^2 + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{7/2}}{(dx^2 + c)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^(7/2)/(c + d*x^2)^(3/2), x)`

[Out] `int((a + b*x^2)^(7/2)/(c + d*x^2)^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{7}{2}}}{(c + dx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**(7/2)/(d*x**2+c)**(3/2), x)`

[Out] `Integral((a + b*x**2)**(7/2)/(c + d*x**2)**(3/2), x)`

$$3.208 \quad \int \frac{(a+bx^2)^{5/2}}{(c+dx^2)^{3/2}} dx$$

Optimal. Leaf size=346

$$\frac{2b\sqrt{c}\sqrt{a+bx^2}(2bc-3ad)\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right)}{3d^{5/2}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{x\sqrt{a+bx^2}(3a^2d^2-13abcd+8b^2c^2)}{3cd^2\sqrt{c+dx^2}} + \frac{\sqrt{a+bx^2}}{bx\sqrt{a+bx^2}}$$

[Out] $-(a*d+b*c)*x*(b*x^2+a)^{(3/2)}/c/d/(d*x^2+c)^{(1/2)}-1/3*(3*a^2*d^2-13*a*b*c*d+8*b^2*c^2)*x*(b*x^2+a)^{(1/2)}/c/d^2/(d*x^2+c)^{(1/2)}+1/3*(3*a^2*d^2-13*a*b*c*d+8*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)})/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(5/2)}/c^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-2/3*b*(-3*a*d+2*b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)})/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)}*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(5/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+1/3*b*(-3*a*d+4*b*c)*x*(b*x^2+a)^{(1/2)}*(d*x^2+c)^{(1/2)}/c/d^2$

Rubi [A] time = 0.27, antiderivative size = 346, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {413, 528, 531, 418, 492, 411}

$$\frac{x\sqrt{a+bx^2}(3a^2d^2-13abcd+8b^2c^2)}{3cd^2\sqrt{c+dx^2}} + \frac{\sqrt{a+bx^2}(3a^2d^2-13abcd+8b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{3\sqrt{c}d^{5/2}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{bx\sqrt{a+bx^2}}{bx\sqrt{a+bx^2}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(5/2)/(c + d*x^2)^(3/2), x]

[Out] $-\left(\left(8*b^2*c^2-13*a*b*c*d+3*a^2*d^2\right)*x*\operatorname{Sqrt}[a+b*x^2]\right)/\left(3*c*d^2*\operatorname{Sqrt}[c+d*x^2]\right)-\left(\left(b*c-a*d\right)*x*\left(a+b*x^2\right)^{(3/2)}\right)/\left(c*d*\operatorname{Sqrt}[c+d*x^2]\right)+\left(b*(4*b*c-3*a*d)*x*\operatorname{Sqrt}[a+b*x^2]*\operatorname{Sqrt}[c+d*x^2]\right)/\left(3*c*d^2\right)+\left(\left(8*b^2*c^2-13*a*b*c*d+3*a^2*d^2\right)*\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticE}\left[\operatorname{ArcTan}\left[\frac{\operatorname{Sqrt}[d]*x}{\operatorname{Sqrt}[c]}\right], 1-\frac{b*c}{a*d}\right]\right)/\left(3*\operatorname{Sqrt}[c]*d^{(5/2)}*\operatorname{Sqrt}\left[\frac{c*(a+b*x^2)}{a*(c+d*x^2)}\right]*\operatorname{Sqrt}[c+d*x^2]\right)-\left(2*b*\operatorname{Sqrt}[c]*(2*b*c-3*a*d)*\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticF}\left[\operatorname{ArcTan}\left[\frac{\operatorname{Sqrt}[d]*x}{\operatorname{Sqrt}[c]}\right], 1-\frac{b*c}{a*d}\right]\right)/\left(3*d^{(5/2)}*\operatorname{Sqrt}\left[\frac{c*(a+b*x^2)}{a*(c+d*x^2)}\right]*\operatorname{Sqrt}[c+d*x^2]\right)$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 413

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:= Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a,
b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rule 531

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Dist[e, Int[(a + b*x^n)^p*(c + d*x^n)^q, x],
x] + Dist[f, Int[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c,
d, e, f, n, p, q}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{5/2}}{(c + dx^2)^{3/2}} dx &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{cd\sqrt{c + dx^2}} + \frac{\int \frac{\sqrt{a+bx^2}(abc+b(4bc-3ad)x^2)}{\sqrt{c+dx^2}} dx}{cd} \\ &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{cd\sqrt{c + dx^2}} + \frac{b(4bc - 3ad)x\sqrt{a + bx^2}\sqrt{c + dx^2}}{3cd^2} + \frac{\int \frac{-2abc(2bc-3ad)-b(8b^2c^2-13abcd)}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx}{3cd^2} \\ &= -\frac{(bc - ad)x(a + bx^2)^{3/2}}{cd\sqrt{c + dx^2}} + \frac{b(4bc - 3ad)x\sqrt{a + bx^2}\sqrt{c + dx^2}}{3cd^2} - \frac{(2ab(2bc - 3ad)) \int \frac{1}{\sqrt{a+bx^2}} dx}{3d^2} \\ &= -\frac{(8b^2c^2 - 13abcd + 3a^2d^2)x\sqrt{a + bx^2}}{3cd^2\sqrt{c + dx^2}} - \frac{(bc - ad)x(a + bx^2)^{3/2}}{cd\sqrt{c + dx^2}} + \frac{b(4bc - 3ad)x\sqrt{a + bx^2}\sqrt{c + dx^2}}{3cd^2} \\ &= -\frac{(8b^2c^2 - 13abcd + 3a^2d^2)x\sqrt{a + bx^2}}{3cd^2\sqrt{c + dx^2}} - \frac{(bc - ad)x(a + bx^2)^{3/2}}{cd\sqrt{c + dx^2}} + \frac{b(4bc - 3ad)x\sqrt{a + bx^2}\sqrt{c + dx^2}}{3cd^2} \end{aligned}$$

Mathematica [C] time = 0.47, size = 256, normalized size = 0.74

$$\frac{-ibc\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}(9a^2d^2-17abcd+8b^2c^2)\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right),\frac{ad}{bc}\right)+ibc\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}(3a^2d^2-17abcd+8b^2c^2)\text{EllipticE}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right),\frac{ad}{bc}\right)}{3cd^3\sqrt{\frac{b}{a}}\sqrt{a+bx^2}\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(5/2)/(c + d*x^2)^(3/2), x]

[Out] (Sqrt[b/a]*d*x*(a + b*x^2)*(-6*a*b*c*d + 3*a^2*d^2 + b^2*c*(4*c + d*x^2)) + I*b*c*(8*b^2*c^2 - 13*a*b*c*d + 3*a^2*d^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - I*b*c*(8*b^2*c^2 - 17*a*b*c*d + 9*a^2*d^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(3*Sqrt[b/a]*c*d^3*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.74, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b^2x^4 + 2abx^2 + a^2)\sqrt{bx^2 + a}\sqrt{dx^2 + c}}{d^2x^4 + 2cdx^2 + c^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^(3/2), x, algorithm="fricas")

[Out] integral((b^2*x^4 + 2*a*b*x^2 + a^2)*sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(d^2*x^4 + 2*c*d*x^2 + c^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^(3/2), x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(5/2)/(d*x^2 + c)^(3/2), x)

maple [A] time = 0.05, size = 539, normalized size = 1.56

$$\sqrt{bx^2 + a}\sqrt{dx^2 + c}\left(\sqrt{\frac{-b}{a}}b^3cd^2x^5 + 3\sqrt{\frac{-b}{a}}a^2bd^3x^3 - 5\sqrt{\frac{-b}{a}}ab^2cd^2x^3 + 4\sqrt{\frac{-b}{a}}b^3c^2dx^3 + 3\sqrt{\frac{-b}{a}}a^3d^3x - 6\sqrt{\frac{-b}{a}}a^2bd^2x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/2)/(d*x^2+c)^(3/2), x)

[Out] 1/3*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*((-1/a*b)^(1/2)*b^3*c*d^2*x^5+3*(-1/a*b)^(1/2)*a^2*b*d^3*x^3-5*(-1/a*b)^(1/2)*a*b^2*c*d^2*x^3+4*(-1/a*b)^(1/2)*b^3*c^2*d*x^3+9*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b*c*d^2-17*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b^2*c^2*d+8*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*b^3*c^3-3*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b*c*d^2+13*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*Ellipti

$cE\left(\left(-1/a*b\right)^{1/2}*x, \left(a/b/c*d\right)^{1/2}\right)*a*b^2*c^2*d-8*\left(\left(b*x^2+a\right)/a\right)^{1/2}*\left(\left(d*x^2+c\right)/c\right)^{1/2}*EllipticE\left(\left(-1/a*b\right)^{1/2}*x, \left(a/b/c*d\right)^{1/2}\right)*b^3*c^3+3*x*a^3*d^3*\left(-1/a*b\right)^{1/2}-6*\left(-1/a*b\right)^{1/2}*a^2*b*c*d^2*x+4*\left(-1/a*b\right)^{1/2}*a*b^2*c^2*d*x\right)/\left(b*d*x^4+a*d*x^2+b*c*x^2+a*c\right)/d^3/c/\left(-1/a*b\right)^{1/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{2}}}{(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/2)/(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/2)/(d*x^2 + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{5/2}}{(dx^2 + c)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/2)/(c + d*x^2)^(3/2),x)

[Out] int((a + b*x^2)^(5/2)/(c + d*x^2)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{5}{2}}}{(c + dx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/2)/(d*x**2+c)**(3/2),x)

[Out] Integral((a + b*x**2)**(5/2)/(c + d*x**2)**(3/2), x)

$$3.209 \quad \int \frac{(a+bx^2)^{3/2}}{(c+dx^2)^{3/2}} dx$$

Optimal. Leaf size=258

$$\frac{b\sqrt{c}\sqrt{a+bx^2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right)\sqrt{a+bx^2}(2bc-ad)E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)+x\sqrt{a+bx^2}(bc-ad)}{d^{3/2}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c}d^{3/2}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}cd\sqrt{c+dx^2}}$$

[Out] $-(-a*d+b*c)*x*(b*x^2+a)^{(1/2)}/c/d/(d*x^2+c)^{(1/2)}+(-a*d+2*b*c)*x*(b*x^2+a)^{(1/2)}/c/d/(d*x^2+c)^{(1/2)}-(-a*d+2*b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/d^{(3/2)}/c^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+b*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(3/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.15, antiderivative size = 258, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {413, 531, 418, 492, 411}

$$\frac{b\sqrt{c}\sqrt{a+bx^2}F\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)\sqrt{a+bx^2}(2bc-ad)E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)+x\sqrt{a+bx^2}(bc-ad)+x\sqrt{a+bx^2}}{d^{3/2}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c}d^{3/2}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}cd\sqrt{c+dx^2}+cd\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/2)/(c + d*x^2)^(3/2), x]

[Out] $-(((b*c - a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(c*d*\operatorname{Sqrt}[c + d*x^2])) + ((2*b*c - a*d)*x*\operatorname{Sqrt}[a + b*x^2])/(c*d*\operatorname{Sqrt}[c + d*x^2]) - ((2*b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[c]*d^{(3/2)}*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (b*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(d^{(3/2)}*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_)*(x_)^2]/((c_) + (d_)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 413

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p + 1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p + q) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R

$\int \frac{d/c + 2\sqrt{c + dx^2}\sqrt{(c + b^2x^2)/(a + c + dx^2)}}{x} dx$; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 492

$\text{Int}[(x_)^2/(\text{Sqrt}[(a_)+(b_)(x_)^2]*\text{Sqrt}[(c_)+(d_)(x_)^2]), x_Symbol]$
 $\rightarrow \text{Simp}[(x*\text{Sqrt}[a + b*x^2])/(b*\text{Sqrt}[c + d*x^2]), x] - \text{Dist}[c/b, \text{Int}[\text{Sqrt}[a + b*x^2]/(c + d*x^2)^{3/2}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]

Rule 531

$\text{Int}[(a_)+(b_)(x_)^{n_})^{p_}*((c_)+(d_)(x_)^{n_})^{q_}*((e_)+(f_)(x_)^{n_}), x_Symbol]$ $\rightarrow \text{Dist}[e, \text{Int}[(a + b*x^n)^p*(c + d*x^n)^q, x], x] + \text{Dist}[f, \text{Int}[x^n*(a + b*x^n)^p*(c + d*x^n)^q, x], x] /;$ FreeQ[{a, b, c, d, e, f, n, p, q}, x]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{3/2}}{(c + dx^2)^{3/2}} dx &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{cd\sqrt{c + dx^2}} + \frac{\int \frac{abc + b(2bc - ad)x^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{cd} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{cd\sqrt{c + dx^2}} + \frac{(ab) \int \frac{1}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{d} + \frac{(b(2bc - ad)) \int \frac{x^2}{\sqrt{a + bx^2}\sqrt{c + dx^2}} dx}{cd} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{cd\sqrt{c + dx^2}} + \frac{(2bc - ad)x\sqrt{a + bx^2}}{cd\sqrt{c + dx^2}} + \frac{b\sqrt{c}\sqrt{a + bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{d^{3/2} \sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}} \sqrt{c + dx^2}} \\ &= -\frac{(bc - ad)x\sqrt{a + bx^2}}{cd\sqrt{c + dx^2}} + \frac{(2bc - ad)x\sqrt{a + bx^2}}{cd\sqrt{c + dx^2}} - \frac{(2bc - ad)\sqrt{a + bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{c} d^{3/2} \sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}} \sqrt{c + dx^2}} \end{aligned}$$

Mathematica [C] time = 0.29, size = 196, normalized size = 0.76

$$\frac{(ad - bc) \left(dx \sqrt{\frac{b}{a}} (a + bx^2) - 2ibc \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{dx^2}{c} + 1} \text{EllipticF}\left(i \sinh^{-1}\left(x \sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) \right) + ibc \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{dx^2}{c} + 1}}{cd^2 \sqrt{\frac{b}{a}} \sqrt{a + bx^2} \sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^(3/2)/(c + d*x^2)^(3/2), x]

[Out] (I*b*c*(-2*b*c + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + (-b*c + a*d)*(Sqrt[b/a]*d*x*(a + b*x^2) - (2*I)*b*c*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)))/(Sqrt[b/a]*c*d^2*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(bx^2 + a)^{\frac{3}{2}} \sqrt{dx^2 + c}}{d^2 x^4 + 2cdx^2 + c^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^(3/2),x, algorithm="fricas")

[Out] integral((b*x^2 + a)^(3/2)*sqrt(d*x^2 + c)/(d^2*x^4 + 2*c*d*x^2 + c^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^(3/2),x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(3/2)/(d*x^2 + c)^(3/2), x)

maple [A] time = 0.04, size = 345, normalized size = 1.34

$$\sqrt{bx^2 + a} \sqrt{dx^2 + c} \left(\sqrt{-\frac{b}{a}} ab d^2 x^3 - \sqrt{-\frac{b}{a}} b^2 cd x^3 + \sqrt{-\frac{b}{a}} a^2 d^2 x - \sqrt{-\frac{b}{a}} abcdx - \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} abcd \text{EllipticF} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/2)/(d*x^2+c)^(3/2),x)

[Out] (b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*((-1/a*b)^(1/2)*a*b*d^2*x^3-(-1/a*b)^(1/2)*b^2*c*d*x^3+2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*b*c*d-2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b^2*c^2-((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*a*b*c*d+2*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b^2*c^2+x*a^2*d^2*(-1/a*b)^(1/2)-(-1/a*b)^(1/2)*a*b*c*d*x/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/d^2/c/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{2}}}{(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/2)/(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/2)/(d*x^2 + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{3/2}}{(dx^2 + c)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/2)/(c + d*x^2)^(3/2),x)

[Out] int((a + b*x^2)^(3/2)/(c + d*x^2)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{3}{2}}}{(c + dx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/2)/(d*x**2+c)**(3/2), x)

[Out] Integral((a + b*x**2)**(3/2)/(c + d*x**2)**(3/2), x)

$$3.210 \quad \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^{3/2}} dx$$

Optimal. Leaf size=84

$$\frac{\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{\sqrt{c}\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*EllipticE(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/c^{(1/2)}/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {411}

$$\frac{\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{\sqrt{c}\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]/(c + d*x^2)^(3/2),x]

[Out] $(\text{Sqrt}[a + b*x^2]*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[d]*x)/\text{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\text{Sqrt}[c]*\text{Sqrt}[d]*\text{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\text{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rubi steps

$$\int \frac{\sqrt{a+bx^2}}{(c+dx^2)^{3/2}} dx = \frac{\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{\sqrt{c}\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}}$$

Mathematica [C] time = 0.32, size = 136, normalized size = 1.62

$$\frac{\frac{x(a+bx^2)}{c} + \frac{ia\sqrt{\frac{b}{a}}\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}\left(E\left(i\sinh^{-1}\left(\sqrt{\frac{b}{a}}x\right)\left|\frac{ad}{bc}\right.\right)-\text{EllipticF}\left(i\sinh^{-1}\left(x\sqrt{\frac{b}{a}}\right),\frac{ad}{bc}\right)\right)}{d}}{\sqrt{a+bx^2}\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2]/(c + d*x^2)^(3/2),x]

[Out] $((x*(a + b*x^2))/c + (I*a*Sqrt[b/a]*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*(EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] - EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]))/d)/(\text{Sqrt}[a + b*x^2]*\text{Sqrt}[c + d*x^2])$

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+a}\sqrt{dx^2+c}}{d^2x^4+2cdx^2+c^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(d^2*x^4 + 2*c*d*x^2 + c^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{(dx^2+c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 + a)/(d*x^2 + c)^(3/2), x)

maple [A] time = 0.03, size = 188, normalized size = 2.24

$$\frac{\sqrt{bx^2+a}\sqrt{dx^2+c}\left(\sqrt{-\frac{b}{a}}bdx^3+\sqrt{-\frac{b}{a}}adx-\sqrt{\frac{dx^2+c}{c}}\sqrt{\frac{bx^2+a}{a}}bc\text{EllipticE}\left(\sqrt{-\frac{b}{a}}x,\sqrt{\frac{ad}{bc}}\right)+\sqrt{\frac{dx^2+c}{c}}\sqrt{\frac{bx^2}{a}}\right)}{(bdx^4+adx^2+bcx^2+ac)\sqrt{-\frac{b}{a}}cd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)/(d*x^2+c)^(3/2),x)

[Out] (b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*((-1/a*b)^(1/2)*b*d*x^3+EllipticF((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b*c*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b*c*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+x*a*d*(-1/a*b)^(1/2))/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/c/d/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{(dx^2+c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)/(d*x^2 + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2+a}}{(dx^2+c)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)/(c + d*x^2)^(3/2),x)

[Out] int((a + b*x^2)^(1/2)/(c + d*x^2)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a + bx^2}}{(c + dx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)/(d*x**2+c)**(3/2), x)

[Out] Integral(sqrt(a + b*x**2)/(c + d*x**2)**(3/2), x)

$$3.211 \quad \int \frac{1}{\sqrt{a+bx^2} (c+dx^2)^{3/2}} dx$$

Optimal. Leaf size=194

$$\frac{b\sqrt{c}\sqrt{a+bx^2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{\sqrt{d}\sqrt{a+bx^2}E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right)}{\sqrt{c}\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $b*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/a/(-a*d+b*c)/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*d^{(1/2)}*(b*x^2+a)^{(1/2)}/(-a*d+b*c)/c^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.15, antiderivative size = 194, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {414, 21, 422, 418, 492, 411}

$$\frac{b\sqrt{c}\sqrt{a+bx^2}F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{\sqrt{d}\sqrt{a+bx^2}E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\middle|1-\frac{bc}{ad}\right)}{\sqrt{c}\sqrt{c+dx^2}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[a + b*x^2]*(c + d*x^2)^(3/2)), x]

[Out] $-((\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[c]*(b*c - a*d)*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))])* \operatorname{Sqrt}[c + d*x^2])) + (b*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*\operatorname{Sqrt}[d]*(b*c - a*d)*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))])* \operatorname{Sqrt}[c + d*x^2])$

Rule 21

Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

```
Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 422

```
Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Dist[
a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[
a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c]
&& PosQ[b/a]
```

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
:= Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a+bx^2} (c+dx^2)^{3/2}} dx &= -\frac{dx\sqrt{a+bx^2}}{c(bc-ad)\sqrt{c+dx^2}} + \frac{\int \frac{bc+bdx^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx}{c(bc-ad)} \\ &= -\frac{dx\sqrt{a+bx^2}}{c(bc-ad)\sqrt{c+dx^2}} + \frac{b \int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx}{c(bc-ad)} \\ &= -\frac{dx\sqrt{a+bx^2}}{c(bc-ad)\sqrt{c+dx^2}} + \frac{b \int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx}{bc-ad} + \frac{(bd) \int \frac{x^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx}{c(bc-ad)} \\ &= \frac{b\sqrt{c}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} - \frac{d \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^{3/2}} dx}{bc-ad} \\ &= -\frac{\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{c}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} + \frac{b\sqrt{c}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}(bc-ad)\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.24, size = 112, normalized size = 0.58

$$\frac{bc\sqrt{\frac{bx^2}{a}+1}\sqrt{\frac{dx^2}{c}+1}E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}}x\right)\middle|\frac{ad}{bc}\right)-dx(a+bx^2)}{\sqrt{\frac{b}{a}}c\sqrt{a+bx^2}\sqrt{c+dx^2}(bc-ad)}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/(Sqrt[a + b*x^2]*(c + d*x^2)^(3/2)), x]
```

```
[Out] (-d*x*(a + b*x^2) + (b*c*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*Elliptic
E[ArcSin[Sqrt[-(b/a)]*x], (a*d)/(b*c)]/Sqrt[-(b/a)])/(c*(b*c - a*d)*Sqrt[a
+ b*x^2]*Sqrt[c + d*x^2])
```

fricas [F] time = 0.77, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+a}\sqrt{dx^2+c}}{bd^2x^6+(2bcd+ad^2)x^4+ac^2+(bc^2+2acd)x^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2+a)*sqrt(d*x^2+c)/(b*d^2*x^6+(2*b*c*d+a*d^2)*x^4+a*c^2+(b*c^2+2*a*c*d)*x^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2+a}(dx^2+c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*x^2+a)*(d*x^2+c)^(3/2)), x)

maple [A] time = 0.04, size = 144, normalized size = 0.74

$$\frac{\left(\sqrt{-\frac{b}{a}}bdx^3+\sqrt{-\frac{b}{a}}adx-\sqrt{\frac{dx^2+c}{c}}\sqrt{\frac{bx^2+a}{a}}bc\text{EllipticE}\left(\sqrt{-\frac{b}{a}}x,\sqrt{\frac{ad}{bc}}\right)\right)\sqrt{dx^2+c}\sqrt{bx^2+a}}{\sqrt{-\frac{b}{a}}(ad-bc)(bdx^4+adx^2+bcx^2+ac)c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2),x)

[Out] ((-1/a*b)^(1/2)*b*d*x^3-EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*b*c*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+(-1/a*b)^(1/2)*a*d*x*(d*x^2+c)^(1/2)*(b*x^2+a)^(1/2)/c/(-1/a*b)^(1/2)/(a*d-b*c)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2+a}(dx^2+c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*x^2+a)*(d*x^2+c)^(3/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{bx^2+a}(dx^2+c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a+b*x^2)^(1/2)*(c+d*x^2)^(3/2)),x)

[Out] int(1/((a+b*x^2)^(1/2)*(c+d*x^2)^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a + bx^2} (c + dx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(1/2)/(d*x**2+c)**(3/2), x)

[Out] Integral(1/(sqrt(a + b*x**2)*(c + d*x**2)**(3/2)), x)

$$3.212 \quad \int \frac{1}{(a+bx^2)^{3/2}(c+dx^2)^{3/2}} dx$$

Optimal. Leaf size=242

$$\frac{2b\sqrt{c}\sqrt{d}\sqrt{a+bx^2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right),1-\frac{bc}{ad}\right)}{a\sqrt{c+dx^2}(bc-ad)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{bx}{a\sqrt{a+bx^2}\sqrt{c+dx^2}(bc-ad)} + \frac{\sqrt{d}\sqrt{a+bx^2}(ad+bc)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)}{a\sqrt{c}\sqrt{c+dx^2}(bc-ad)}$$

[Out] $b*x/a/(-a*d+b*c)/(b*x^2+a)^{(1/2)/(d*x^2+c)^{(1/2)}+(a*d+b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a/(-a*d+b*c)^2/c^{(1/2)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)/(d*x^2+c)^{(1/2)}-2*b*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a/(-a*d+b*c)^2/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.12, antiderivative size = 242, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {414, 525, 418, 411}

$$\frac{bx}{a\sqrt{a+bx^2}\sqrt{c+dx^2}(bc-ad)} + \frac{2b\sqrt{c}\sqrt{d}\sqrt{a+bx^2}F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\left|1-\frac{bc}{ad}\right.\right)}{a\sqrt{c+dx^2}(bc-ad)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{d}\sqrt{a+bx^2}(ad+bc)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)}{a\sqrt{c}\sqrt{c+dx^2}(bc-ad)^2\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(3/2)*(c + d*x^2)^(3/2)),x]

[Out] $(b*x)/(a*(b*c - a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2]) + (\operatorname{Sqrt}[d]*(b*c + a*d)*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]],1 - (b*c)/(a*d)])/(a*\operatorname{Sqrt}[c]*(b*c - a*d)^2*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2)])*\operatorname{Sqrt}[c + d*x^2]) - (2*b*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]],1 - (b*c)/(a*d)])/(a*(b*c - a*d)^2*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2)])*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R

$\text{t}[d/c, 2] * \text{Sqrt}[c + d*x^2] * \text{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{PosQ}[d/c] \&\& \text{PosQ}[b/a] \&\& !\text{SimplerSqrtQ}[b/a, d/c]$

Rule 525

$\text{Int}[(e_ + (f_)*(x_)^2)/(\text{Sqrt}[(a_ + (b_)*(x_)^2]*((c_ + (d_)*(x_)^2)^{(3/2)})), x_Symbol] :> \text{Dist}[(b*e - a*f)/(b*c - a*d), \text{Int}[1/(\text{Sqrt}[a + b*x^2]*\text{Sqrt}[c + d*x^2]), x], x] - \text{Dist}[(d*e - c*f)/(b*c - a*d), \text{Int}[\text{Sqrt}[a + b*x^2]/(c + d*x^2)^{(3/2)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \text{PosQ}[b/a] \&\& \text{PosQ}[d/c]$

Rubi steps

$$\int \frac{1}{(a + bx^2)^{3/2} (c + dx^2)^{3/2}} dx = \frac{bx}{a(bc - ad)\sqrt{a + bx^2} \sqrt{c + dx^2}} - \frac{\int \frac{ad - bdx^2}{\sqrt{a + bx^2} (c + dx^2)^{3/2}} dx}{a(bc - ad)}$$

$$= \frac{bx}{a(bc - ad)\sqrt{a + bx^2} \sqrt{c + dx^2}} - \frac{(2bd) \int \frac{1}{\sqrt{a + bx^2} \sqrt{c + dx^2}} dx}{(bc - ad)^2} + \frac{(d(bc + ad)) \int \frac{\sqrt{a + bx^2}}{(c + dx^2)^{3/2}} dx}{a(bc - ad)^2}$$

$$= \frac{bx}{a(bc - ad)\sqrt{a + bx^2} \sqrt{c + dx^2}} + \frac{\sqrt{d} (bc + ad) \sqrt{a + bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{c} (bc - ad)^2 \sqrt{\frac{c(a + bx^2)}{a(c + dx^2)}} \sqrt{c + dx^2}}$$

Mathematica [C] time = 0.70, size = 224, normalized size = 0.93

$$\frac{\sqrt{\frac{b}{a}} \left(i b c \sqrt{\frac{b x^2}{a} + 1} \sqrt{\frac{d x^2}{c} + 1} (a d - b c) \text{EllipticF}\left(i \sinh^{-1}\left(x \sqrt{\frac{b}{a}}\right), \frac{a d}{b c}\right) + x \sqrt{\frac{b}{a}} \left(a^2 d^2 + a b d^2 x^2 + b^2 c (c + d x^2)\right) + i b \sqrt{a + b x^2} \sqrt{c + d x^2} \right)}{b c \sqrt{a + b x^2} \sqrt{c + d x^2} (b c - a d)^2}$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^(3/2)*(c + d*x^2)^(3/2)), x]

[Out] (Sqrt[b/a]*(Sqrt[b/a]*x*(a^2*d^2 + a*b*d^2*x^2 + b^2*c*(c + d*x^2)) + I*b*c*(b*c + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)] + I*b*c*(-(b*c) + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)])/(b*c*(b*c - a*d)^2*Sqrt[a + b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b x^2 + a} \sqrt{d x^2 + c}}{b^2 d^2 x^8 + 2 (b^2 c d + a b d^2) x^6 + (b^2 c^2 + 4 a b c d + a^2 d^2) x^4 + a^2 c^2 + 2 (a b c^2 + a^2 c d) x^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b^2*d^2*x^8 + 2*(b^2*c*d + a*b*d^2)*x^6 + (b^2*c^2 + 4*a*b*c*d + a^2*d^2)*x^4 + a^2*c^2 + 2*(a*b*c^2 + a^2*c*d)*x^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{2}}(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(3/2)*(d*x^2 + c)^(3/2)), x)

maple [A] time = 0.05, size = 354, normalized size = 1.46

$$\left(\sqrt{-\frac{b}{a}} ab d^2 x^3 + \sqrt{-\frac{b}{a}} b^2 cd x^3 + \sqrt{-\frac{b}{a}} a^2 d^2 x - \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} abcd \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right) - \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(3/2)/(d*x^2+c)^(3/2),x)

[Out] $((-1/a*b)^{(1/2)}*a*b*d^2*x^3+(-1/a*b)^{(1/2)}*b^2*c*d*x^3-((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticF}((-1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*a*b*c*d+((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticF}((-1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*b^2*c^2-((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticE}((-1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*a*b*c*d-((b*x^2+a)/a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*\operatorname{EllipticE}((-1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*b^2*c^2+(-1/a*b)^{(1/2)}*a^2*d^2*x+x*b^2*c^2*(-1/a*b)^{(1/2)}*((d*x^2+c)^{(1/2)}*(b*x^2+a)^{(1/2)}/a/c/(-1/a*b)^{(1/2)}/(a*d-b*c)^2/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{2}}(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/2)/(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(3/2)*(d*x^2 + c)^(3/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{3/2}(dx^2 + c)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(3/2)*(c + d*x^2)^(3/2)),x)

[Out] int(1/((a + b*x^2)^(3/2)*(c + d*x^2)^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{3}{2}}(c + dx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(3/2)/(d*x**2+c)**(3/2),x)

[Out] Integral(1/((a + b*x**2)**(3/2)*(c + d*x**2)**(3/2)), x)

$$3.213 \quad \int \frac{1}{(a+bx^2)^{5/2}(c+dx^2)^{3/2}} dx$$

Optimal. Leaf size=323

$$\frac{b\sqrt{c}\sqrt{d}\sqrt{a+bx^2}(bc-9ad)\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1-\frac{bc}{ad}\right)\sqrt{d}\sqrt{a+bx^2}(-3a^2d^2-7abcd+2b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)}{3a^2\sqrt{c+dx^2}(bc-ad)^3\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{b\sqrt{c}\sqrt{d}\sqrt{a+bx^2}(-3a^2d^2-7abcd+2b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)}{3a^2\sqrt{c+dx^2}(bc-ad)^3\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $1/3*b*x/a/(-a*d+b*c)/(b*x^2+a)^{(3/2)}/(d*x^2+c)^{(1/2)}+2/3*b*(-3*a*d+b*c)*x/a^2/(-a*d+b*c)^2/(b*x^2+a)^{(1/2)}/(d*x^2+c)^{(1/2)}+1/3*(-3*a^2*d^2-7*a*b*c*d+2*b^2*c^2)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)})/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a^2/(-a*d+b*c)^3/c^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-1/3*b*(-9*a*d+b*c)*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)})/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)}*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/a^2/(-a*d+b*c)^3/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.26, antiderivative size = 323, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {414, 527, 525, 418, 411}

$$\frac{\sqrt{d}\sqrt{a+bx^2}(-3a^2d^2-7abcd+2b^2c^2)E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right)\right)\left[1-\frac{bc}{ad}\right]}{3a^2\sqrt{c+dx^2}(bc-ad)^3\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{2bx(bc-3ad)}{3a^2\sqrt{a+bx^2}\sqrt{c+dx^2}(bc-ad)^2} - \frac{b\sqrt{c}\sqrt{d}\sqrt{a+bx^2}}{3a^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/((a+b*x^2)^{(5/2)}*(c+d*x^2)^{(3/2)}), x]$

[Out] $(b*x)/(3*a*(b*c-a*d)*(a+b*x^2)^{(3/2)}*\operatorname{Sqrt}[c+d*x^2])+(2*b*(b*c-3*a*d)*x)/(3*a^2*(b*c-a*d)^2*\operatorname{Sqrt}[a+b*x^2]*\operatorname{Sqrt}[c+d*x^2])+(\operatorname{Sqrt}[d]*(2*b^2*c^2-7*a*b*c*d-3*a^2*d^2)*\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1-(b*c)/(a*d)])/(3*a^2*\operatorname{Sqrt}[c]*(b*c-a*d)^3*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2)])*\operatorname{Sqrt}[c+d*x^2])-(b*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*(b*c-9*a*d)*\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1-(b*c)/(a*d)])/(3*a^2*(b*c-a*d)^3*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2)])*\operatorname{Sqrt}[c+d*x^2])$

Rule 411

$\operatorname{Int}[\operatorname{Sqrt}[(a_.)+(b_.)*(x_)^2]/((c_.)+(d_.)*(x_)^2)^{(3/2)}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[a+b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[\operatorname{Rt}[d/c, 2]*x], 1-(b*c)/(a*d)])/(c*\operatorname{Rt}[d/c, 2]*\operatorname{Sqrt}[c+d*x^2]*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2)]), x] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{PosQ}[b/a] \&\& \operatorname{PosQ}[d/c]$

Rule 414

$\operatorname{Int}[(a_.)+(b_.)*(x_)^{(n_)}]^{(p_)}*((c_.)+(d_.)*(x_)^{(n_)})^{(q_)}, x_Symbol] \rightarrow -\operatorname{Simp}[(b*x*(a+b*x^n)^{(p+1)}*(c+d*x^n)^{(q+1)})/(a*n*(p+1)*(b*c-a*d)), x] + \operatorname{Dist}[1/(a*n*(p+1)*(b*c-a*d)), \operatorname{Int}[(a+b*x^n)^{(p+1)}*(c+d*x^n)^q*\operatorname{Simp}[b*c+n*(p+1)*(b*c-a*d)+d*b*(n*(p+q+2)+1)*x^n, x], x] /; \operatorname{FreeQ}[\{a, b, c, d, n, q\}, x] \&\& \operatorname{NeQ}[b*c-a*d, 0] \&\& \operatorname{LtQ}[p, -1] \&\& !(IntegerQ[p] \&\& IntegerQ[q] \&\& \operatorname{LtQ}[q, -1]) \&\& \operatorname{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 418

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 525

```
Int[((e_) + (f_.)*(x_)^2)/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)^(
3/2)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(Sqrt[a + b*x^2]*S
qrt[c + d*x^2]), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[Sqrt[a + b*x^2]
/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && PosQ[b/a] &&
PosQ[d/c]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + bx^2)^{5/2} (c + dx^2)^{3/2}} dx &= \frac{bx}{3a(bc - ad)(a + bx^2)^{3/2} \sqrt{c + dx^2}} - \frac{\int \frac{-2bc + 3ad - 3bdx^2}{(a + bx^2)^{3/2} (c + dx^2)^{3/2}} dx}{3a(bc - ad)} \\ &= \frac{bx}{3a(bc - ad)(a + bx^2)^{3/2} \sqrt{c + dx^2}} + \frac{2b(bc - 3ad)x}{3a^2(bc - ad)^2 \sqrt{a + bx^2} \sqrt{c + dx^2}} + \frac{\int \frac{a}{(a + bx^2)^{3/2} \sqrt{c + dx^2}} dx}{3a^2(bc - ad)^2 \sqrt{a + bx^2} \sqrt{c + dx^2}} \\ &= \frac{bx}{3a(bc - ad)(a + bx^2)^{3/2} \sqrt{c + dx^2}} + \frac{2b(bc - 3ad)x}{3a^2(bc - ad)^2 \sqrt{a + bx^2} \sqrt{c + dx^2}} - \frac{(bd)}{3a^2(bc - ad)^2 \sqrt{a + bx^2} \sqrt{c + dx^2}} \\ &= \frac{bx}{3a(bc - ad)(a + bx^2)^{3/2} \sqrt{c + dx^2}} + \frac{2b(bc - 3ad)x}{3a^2(bc - ad)^2 \sqrt{a + bx^2} \sqrt{c + dx^2}} + \frac{\sqrt{d}}{3a^2(bc - ad)^2 \sqrt{a + bx^2} \sqrt{c + dx^2}} \end{aligned}$$

Mathematica [C] time = 1.07, size = 337, normalized size = 1.04

$$\frac{2ibc(a + bx^2) \sqrt{\frac{bx^2}{a} + 1} \sqrt{\frac{dx^2}{c} + 1} (3a^2d^2 - 4abcd + b^2c^2) \text{EllipticF}\left(i \sinh^{-1}\left(x \sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right) + ibc(a + bx^2) \sqrt{\frac{bx^2}{a}}}{3a^2(bc - ad)^2 \sqrt{a + bx^2} \sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/((a + b*x^2)^(5/2)*(c + d*x^2)^(3/2)), x]
```

```
[Out] (Sqrt[b/a]*x*(3*a^4*d^3 + 6*a^3*b*d^3*x^2 - 2*b^4*c^2*x^2*(c + d*x^2) + a^2
*b^2*d*(8*c^2 + 8*c*d*x^2 + 3*d^2*x^4) + a*b^3*c*(-3*c^2 + 4*c*d*x^2 + 7*d^
2*x^4)) + I*b*c*(-2*b^2*c^2 + 7*a*b*c*d + 3*a^2*d^2)*(a + b*x^2)*Sqrt[1 + (
b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticE[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]
```

] + (2*I)*b*c*(b^2*c^2 - 4*a*b*c*d + 3*a^2*d^2)*(a + b*x^2)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[I*ArcSinh[Sqrt[b/a]*x], (a*d)/(b*c)]/(3*a^2*Sqrt[b/a]*c*(-(b*c) + a*d)^3*(a + b*x^2)^(3/2)*Sqrt[c + d*x^2])

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{bx^2 + a} \sqrt{dx^2 + c}}{b^3 d^2 x^{10} + (2 b^3 c d + 3 a b^2 d^2) x^8 + (b^3 c^2 + 6 a b^2 c d + 3 a^2 b d^2) x^6 + a^3 c^2 + (3 a b^2 c^2 + 6 a^2 b c d + a^3 d^2) x^4 + \dots} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b^3*d^2*x^10 + (2*b^3*c*d + 3*a*b^2*d^2)*x^8 + (b^3*c^2 + 6*a*b^2*c*d + 3*a^2*b*d^2)*x^6 + a^3*c^2 + (3*a*b^2*c^2 + 6*a^2*b*c*d + a^3*d^2)*x^4 + (3*a^2*b*c*d + 2*a^3*c*d)*x^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{2}}(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(5/2)*(d*x^2 + c)^(3/2)), x)

maple [B] time = 0.07, size = 964, normalized size = 2.98

$$\frac{-3\sqrt{-\frac{b}{a}} a^2 b^2 d^3 x^5 - 7\sqrt{-\frac{b}{a}} a b^3 c d^2 x^5 + 2\sqrt{-\frac{b}{a}} b^4 c^2 d x^5 - 6\sqrt{-\frac{b}{a}} a^3 b d^3 x^3 - 8\sqrt{-\frac{b}{a}} a^2 b^2 c d^2 x^3 + 3\sqrt{\frac{d x^2 + c}{c}} \sqrt{\frac{b x^2}{a}}}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(5/2)/(d*x^2+c)^(3/2),x)

[Out] -1/3*(-3*x^5*a^2*b^2*d^3*(-1/a*b)^(1/2)-7*x^5*a*b^3*c*d^2*(-1/a*b)^(1/2)+2*x^5*b^4*c^2*d*(-1/a*b)^(1/2)+3*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a^2*b^2*c*d^2*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+7*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a*b^3*c^2*d*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-2*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*b^4*c^3*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+6*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a^2*b^2*c*d^2*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-8*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*a*b^3*c^2*d*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+2*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*x^2*b^4*c^3*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-6*x^3*a^3*b*d^3*(-1/a*b)^(1/2)-8*x^3*a^2*b^2*c*d^2*(-1/a*b)^(1/2)-4*x^3*a*b^3*c^2*d*(-1/a*b)^(1/2)+2*x^3*b^4*c^3*(-1/a*b)^(1/2)+3*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^3*b*c*d^2*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+7*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b^2*c^2*d*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-2*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b^3*c^3*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+6*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^3*b*c*d^2*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-8*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a^2*b^2*c^2*d*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)+2*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*a*b^3*c^3*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)-3*x*a^4*d^3*(-1/a*b)^(1/2)-8*x*a^2*b^2*c^2*d*(-1/a*b)^(1/2)+3*x*a*b^3*c^3*(-1/a*b)^(1/2))/(d*x^2+c)^(1/2)/(a*d-b*c)^3/(-1/a*b)^(1/2)/a^2/c/(b*x^2+a)^(3/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{2}}(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/2)/(d*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(5/2)*(d*x^2 + c)^(3/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{2}}(dx^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)^(3/2)),x)

[Out] int(1/((a + b*x^2)^(5/2)*(c + d*x^2)^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{5}{2}}(c + dx^2)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(5/2)/(d*x**2+c)**(3/2),x)

[Out] Integral(1/((a + b*x**2)**(5/2)*(c + d*x**2)**(3/2)), x)

$$3.214 \quad \int \frac{1}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=87

$$\frac{\sqrt{c} \sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{a\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/a/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {418}

$$\frac{\sqrt{c} \sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]),x]

[Out] $(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{a+bx^2} \sqrt{c+dx^2}} dx = \frac{\sqrt{c} \sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}} \sqrt{c+dx^2}}$$

Mathematica [A] time = 0.06, size = 86, normalized size = 0.99

$$\frac{\sqrt{\frac{a+bx^2}{a}} \sqrt{\frac{c+dx^2}{c}} \operatorname{EllipticF}\left(\sin^{-1}\left(x\sqrt{-\frac{b}{a}}\right), \frac{ad}{bc}\right)}{\sqrt{-\frac{b}{a}} \sqrt{a+bx^2} \sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]),x]

[Out] $(\operatorname{Sqrt}[(a + b*x^2)/a]*\operatorname{Sqrt}[(c + d*x^2)/c]*\operatorname{EllipticF}[\operatorname{ArcSin}[\operatorname{Sqrt}[-(b/a)]*x], (a*d)/(b*c)])/(\operatorname{Sqrt}[-(b/a)]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[c + d*x^2])$

fricas [F] time = 0.71, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2 + a} \sqrt{dx^2 + c}}{bdx^4 + (bc + ad)x^2 + ac}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)/(b*d*x^4 + (b*c + a*d)*x^2 + a*c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2 + a} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)), x)

maple [A] time = 0.02, size = 100, normalized size = 1.15

$$\frac{\sqrt{\frac{dx^2+c}{c}} \sqrt{\frac{bx^2+a}{a}} \sqrt{bx^2 + a} \sqrt{dx^2 + c} \text{EllipticF}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right)}{\sqrt{-\frac{b}{a}} (bdx^4 + adx^2 + bcx^2 + ac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x)

[Out] EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)*(b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/(-1/a*b)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2 + a} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*x^2 + a)*sqrt(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{bx^2 + a} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)),x)

[Out] int(1/((a + b*x^2)^(1/2)*(c + d*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a + bx^2} \sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x**2+a)**(1/2)/(d*x**2+c)**(1/2), x)
```

```
[Out] Integral(1/(sqrt(a + b*x**2)*sqrt(c + d*x**2)), x)
```

$$3.215 \quad \int \frac{1}{\sqrt{a-bx^2} \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=87

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{c + dx^2}}$$

[Out] $\operatorname{EllipticF}(x \cdot b^{(1/2)}/a^{(1/2)}, (-a \cdot d/b/c)^{(1/2)}) \cdot a^{(1/2)} \cdot (1 - b \cdot x^2/a)^{(1/2)} \cdot (1 + d \cdot x^2/c)^{(1/2)} / b^{(1/2)} / (-b \cdot x^2 + a)^{(1/2)} / (d \cdot x^2 + c)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {421, 419}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(\operatorname{Sqrt}[a - b \cdot x^2] \cdot \operatorname{Sqrt}[c + d \cdot x^2]), x]$

[Out] $(\operatorname{Sqrt}[a] \cdot \operatorname{Sqrt}[1 - (b \cdot x^2)/a] \cdot \operatorname{Sqrt}[1 + (d \cdot x^2)/c] \cdot \operatorname{EllipticF}[\operatorname{ArcSin}[(\operatorname{Sqrt}[b] \cdot x)/\operatorname{Sqrt}[a]], -(a \cdot d)/(b \cdot c)]) / (\operatorname{Sqrt}[b] \cdot \operatorname{Sqrt}[a - b \cdot x^2] \cdot \operatorname{Sqrt}[c + d \cdot x^2])$

Rule 419

$\operatorname{Int}[1/(\operatorname{Sqrt}[(a_) + (b_) \cdot (x_)^2] \cdot \operatorname{Sqrt}[(c_) + (d_) \cdot (x_)^2]), x_Symbol] \rightarrow \operatorname{Simp}[(1 \cdot \operatorname{EllipticF}[\operatorname{ArcSin}[\operatorname{Rt}[-(d/c), 2] \cdot x], (b \cdot c)/(a \cdot d)]) / (\operatorname{Sqrt}[a] \cdot \operatorname{Sqrt}[c] \cdot \operatorname{Rt}[-(d/c), 2]), x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \ \operatorname{NegQ}[d/c] \ \&\& \ \operatorname{GtQ}[c, 0] \ \&\& \ \operatorname{GtQ}[a, 0] \ \&\& \ !(\operatorname{NegQ}[b/a] \ \&\& \ \operatorname{SimplerSqrtQ}[-(b/a), -(d/c)])$

Rule 421

$\operatorname{Int}[1/(\operatorname{Sqrt}[(a_) + (b_) \cdot (x_)^2] \cdot \operatorname{Sqrt}[(c_) + (d_) \cdot (x_)^2]), x_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Sqrt}[1 + (d \cdot x^2)/c] / \operatorname{Sqrt}[c + d \cdot x^2], \operatorname{Int}[1/(\operatorname{Sqrt}[a + b \cdot x^2] \cdot \operatorname{Sqrt}[1 + (d \cdot x^2)/c]), x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \ !\operatorname{GtQ}[c, 0]$

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a-bx^2} \sqrt{c+dx^2}} dx &= \frac{\sqrt{1 + \frac{dx^2}{c}} \int \frac{1}{\sqrt{a-bx^2} \sqrt{1 + \frac{dx^2}{c}}} dx}{\sqrt{c + dx^2}} \\ &= \frac{\left(\sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}}} dx}{\sqrt{a - bx^2} \sqrt{c + dx^2}} \\ &= \frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{c + dx^2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 87, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{\frac{c+dx^2}{c}} \operatorname{EllipticF}\left(\sin^{-1}\left(x \sqrt{\frac{b}{a}}\right), -\frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{a - bx^2} \sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[a - b*x^2]*Sqrt[c + d*x^2]),x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[(c + d*x^2)/c]*EllipticF[ArcSin[Sqrt[b/a]*x], -((a*d)/(b*c)))]/(Sqrt[b/a]*Sqrt[a - b*x^2]*Sqrt[c + d*x^2])

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2+a}\sqrt{dx^2+c}}{bdx^4+(bc-ad)x^2-ac}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)/(b*d*x^4 + (b*c - a*d)*x^2 - a*c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-bx^2+a}\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)), x)

maple [A] time = 0.04, size = 106, normalized size = 1.22

$$-\frac{\sqrt{\frac{dx^2+c}{c}}\sqrt{-\frac{bx^2-a}{a}}\sqrt{-bx^2+a}\sqrt{dx^2+c}\text{EllipticF}\left(\sqrt{\frac{b}{a}}x,\sqrt{-\frac{ad}{bc}}\right)}{\sqrt{\frac{b}{a}}(bdx^4-adx^2+bcx^2-ac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x)

[Out] -EllipticF(x*(1/a*b)^(1/2), (-a/b/c*d)^(1/2))*((d*x^2+c)/c)^(1/2)*(-b*x^2-a)/a)^(1/2)*(-b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)/(1/a*b)^(1/2)/(b*d*x^4-a*d*x^2+b*c*x^2-a*c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-bx^2+a}\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{a-bx^2}\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)),x)

[Out] `int(1/((a - b*x^2)^(1/2)*(c + d*x^2)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a - bx^2} \sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-b*x**2+a)**(1/2)/(d*x**2+c)**(1/2), x)`

[Out] `Integral(1/(sqrt(a - b*x**2)*sqrt(c + d*x**2)), x)`

$$3.216 \quad \int \frac{1}{\sqrt{a+bx^2} \sqrt{c-dx^2}} dx$$

Optimal. Leaf size=87

$$\frac{\sqrt{c} \sqrt{\frac{bx^2}{a} + 1} \sqrt{1 - \frac{dx^2}{c}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{a + bx^2} \sqrt{c - dx^2}}$$

[Out] EllipticF(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(1+b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {421, 419}

$$\frac{\sqrt{c} \sqrt{\frac{bx^2}{a} + 1} \sqrt{1 - \frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{a + bx^2} \sqrt{c - dx^2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]),x]

[Out] (Sqrt[c]*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -((b*c)/(a*d))]/(Sqrt[d]*Sqrt[a + b*x^2]*Sqrt[c - d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a+bx^2} \sqrt{c-dx^2}} dx &= \frac{\sqrt{1 - \frac{dx^2}{c}} \int \frac{1}{\sqrt{a+bx^2} \sqrt{1 - \frac{dx^2}{c}}} dx}{\sqrt{c - dx^2}} \\ &= \frac{\left(\sqrt{1 + \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1 + \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}}} dx}{\sqrt{a + bx^2} \sqrt{c - dx^2}} \\ &= \frac{\sqrt{c} \sqrt{1 + \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{a + bx^2} \sqrt{c - dx^2}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 89, normalized size = 1.02

$$\frac{\sqrt{\frac{a+bx^2}{a}} \sqrt{\frac{c-dx^2}{c}} \operatorname{EllipticF}\left(\sin^{-1}\left(x\sqrt{\frac{b}{a}}\right), -\frac{ad}{bc}\right)}{\sqrt{-\frac{b}{a}} \sqrt{a + bx^2} \sqrt{c - dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[a + b*x^2]*Sqrt[c - d*x^2]),x]

[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[(c - d*x^2)/c]*EllipticF[ArcSin[Sqrt[-(b/a)]*x],
-((a*d)/(b*c))]/(Sqrt[-(b/a)]*Sqrt[a + b*x^2]*Sqrt[c - d*x^2])

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{bx^2+a}\sqrt{-dx^2+c}}{bdx^4-(bc-ad)x^2-ac},x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)/(b*d*x^4 - (b*c - a*d)*x^2 - a*c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2+a}\sqrt{-dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)), x)

maple [A] time = 0.04, size = 106, normalized size = 1.22

$$\frac{\sqrt{\frac{bx^2+a}{a}}\sqrt{\frac{-dx^2-c}{c}}\sqrt{bx^2+a}\sqrt{-dx^2+c}\text{EllipticF}\left(\sqrt{\frac{d}{c}}x,\sqrt{\frac{-bc}{ad}}\right)}{\sqrt{\frac{d}{c}}(bdx^4+adx^2-bcx^2-ac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x)

[Out] -EllipticF(x*(1/c*d)^(1/2),(-b*c/a/d)^(1/2))*((b*x^2+a)/a)^(1/2)*(-d*x^2-c)/c)^(1/2)*(b*x^2+a)^(1/2)*(-d*x^2+c)^(1/2)/(1/c*d)^(1/2)/(b*d*x^4+a*d*x^2-b*c*x^2-a*c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2+a}\sqrt{-dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{bx^2+a}\sqrt{c-dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)),x)

[Out] `int(1/((a + b*x^2)^(1/2)*(c - d*x^2)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a + bx^2} \sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2), x)`

[Out] `Integral(1/(sqrt(a + b*x**2)*sqrt(c - d*x**2)), x)`

$$3.217 \quad \int \frac{1}{\sqrt{a-bx^2} \sqrt{c-dx^2}} dx$$

Optimal. Leaf size=88

$$\frac{\sqrt{c} \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{a - bx^2} \sqrt{c - dx^2}}$$

[Out] $\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}, (b*c/a/d)^{(1/2)}) * c^{(1/2)} * (1-b*x^2/a)^{(1/2)} * (1-d*x^2/c)^{(1/2)} / d^{(1/2)} / (-b*x^2+a)^{(1/2)} / (-d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.06, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {421, 419}

$$\frac{\sqrt{c} \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{a - bx^2} \sqrt{c - dx^2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(\operatorname{Sqrt}[a - b*x^2]*\operatorname{Sqrt}[c - d*x^2]), x]$

[Out] $(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[1 - (b*x^2)/a]*\operatorname{Sqrt}[1 - (d*x^2)/c]*\operatorname{EllipticF}[\operatorname{ArcSin}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a - b*x^2]*\operatorname{Sqrt}[c - d*x^2])$

Rule 419

$\operatorname{Int}[1/(\operatorname{Sqrt}[(a_) + (b_.)*(x_)^2]*\operatorname{Sqrt}[(c_) + (d_.)*(x_)^2]), x_Symbol] \rightarrow \operatorname{Simp}[(1*\operatorname{EllipticF}[\operatorname{ArcSin}[\operatorname{Rt}[-(d/c), 2]*x], (b*c)/(a*d)])/(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[c]*\operatorname{Rt}[-(d/c), 2]), x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \ \operatorname{NegQ}[d/c] \ \&\& \ \operatorname{GtQ}[c, 0] \ \&\& \ \operatorname{GtQ}[a, 0] \ \&\& \ !(\operatorname{NegQ}[b/a] \ \&\& \ \operatorname{SimplerSqrtQ}[-(b/a), -(d/c)])$

Rule 421

$\operatorname{Int}[1/(\operatorname{Sqrt}[(a_) + (b_.)*(x_)^2]*\operatorname{Sqrt}[(c_) + (d_.)*(x_)^2]), x_Symbol] \rightarrow \operatorname{Dist}[\operatorname{Sqrt}[1 + (d*x^2)/c]/\operatorname{Sqrt}[c + d*x^2], \operatorname{Int}[1/(\operatorname{Sqrt}[a + b*x^2]*\operatorname{Sqrt}[1 + (d*x^2)/c]), x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \ !\operatorname{GtQ}[c, 0]$

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a-bx^2} \sqrt{c-dx^2}} dx &= \frac{\sqrt{1 - \frac{dx^2}{c}} \int \frac{1}{\sqrt{a-bx^2} \sqrt{1 - \frac{dx^2}{c}}} dx}{\sqrt{c - dx^2}} \\ &= \frac{\left(\sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}}} dx}{\sqrt{a - bx^2} \sqrt{c - dx^2}} \\ &= \frac{\sqrt{c} \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 - \frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{a - bx^2} \sqrt{c - dx^2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 88, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{\frac{c-dx^2}{c}} \operatorname{EllipticF}\left(\sin^{-1}\left(x\sqrt{\frac{b}{a}}\right), \frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{a - bx^2} \sqrt{c - dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[a - b*x^2]*Sqrt[c - d*x^2]),x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[(c - d*x^2)/c]*EllipticF[ArcSin[Sqrt[b/a]*x], (a*d)/(b*c)]/(Sqrt[b/a]*Sqrt[a - b*x^2]*Sqrt[c - d*x^2])

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-bx^2 + a} \sqrt{-dx^2 + c}}{bdx^4 - (bc + ad)x^2 + ac}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)/(b*d*x^4 - (b*c + a*d)*x^2 + a*c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-bx^2 + a} \sqrt{-dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)), x)

maple [A] time = 0.04, size = 108, normalized size = 1.23

$$\frac{\sqrt{-\frac{bx^2-a}{a}} \sqrt{-\frac{dx^2-c}{c}} \sqrt{-bx^2 + a} \sqrt{-dx^2 + c} \text{EllipticF}\left(\sqrt{\frac{d}{c}} x, \sqrt{\frac{bc}{ad}}\right)}{\sqrt{\frac{d}{c}} (bdx^4 - adx^2 - bcx^2 + ac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x)

[Out] EllipticF((1/c*d)^(1/2)*x, (1/a*b*c/d)^(1/2))*(-b*x^2-a)/a)^(1/2)*(-d*x^2-c)/c)^(1/2)*(-b*x^2+a)^(1/2)*(-d*x^2+c)^(1/2)/(1/c*d)^(1/2)/(b*d*x^4-a*d*x^2-b*c*x^2+a*c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-bx^2 + a} \sqrt{-dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{a - bx^2} \sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)),x)

[Out] `int(1/((a - b*x^2)^(1/2)*(c - d*x^2)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a - bx^2} \sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2), x)`

[Out] `Integral(1/(sqrt(a - b*x**2)*sqrt(c - d*x**2)), x)`

$$3.218 \quad \int \frac{1}{\sqrt{1-x^2} \sqrt{2+5x^2}} dx$$

Optimal. Leaf size=12

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), -\frac{5}{2}\right)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,1/2*I*10^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {419}

$$\frac{F\left(\sin^{-1}(x) \mid -\frac{5}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 - x^2]*Sqrt[2 + 5*x^2]),x]

[Out] EllipticF[ArcSin[x], -5/2]/Sqrt[2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S imp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{1-x^2} \sqrt{2+5x^2}} dx = \frac{F\left(\sin^{-1}(x) \mid -\frac{5}{2}\right)}{\sqrt{2}}$$

Mathematica [A] time = 0.01, size = 12, normalized size = 1.00

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), -\frac{5}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 - x^2]*Sqrt[2 + 5*x^2]),x]

[Out] EllipticF[ArcSin[x], -5/2]/Sqrt[2]

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{5x^2+2}\sqrt{-x^2+1}}{5x^4-3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(5*x^2 + 2)*sqrt(-x^2 + 1)/(5*x^4 - 3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{5x^2 + 2}\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(5*x^2 + 2)*sqrt(-x^2 + 1)), x)

maple [A] time = 0.03, size = 14, normalized size = 1.17

$$\frac{\sqrt{2} \operatorname{EllipticF}\left(x, \frac{i\sqrt{10}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/2)/(5*x^2+2)^(1/2),x)

[Out] 1/2*EllipticF(x,1/2*I*10^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{5x^2 + 2}\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(5*x^2 + 2)*sqrt(-x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.08

$$\int \frac{1}{\sqrt{1-x^2}\sqrt{5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1-x^2)^(1/2)*(5*x^2+2)^(1/2)),x)

[Out] int(1/((1-x^2)^(1/2)*(5*x^2+2)^(1/2)), x)

sympy [A] time = 3.73, size = 19, normalized size = 1.58

$$\begin{cases} \frac{\sqrt{2} F\left(\operatorname{asin}(x) \middle| -\frac{5}{2}\right)}{2} & \text{for } x > -1 \wedge x < 1 \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/2)/(5*x**2+2)**(1/2),x)

[Out] Piecewise((sqrt(2)*elliptic_f(asin(x), -5/2)/2, (x > -1) & (x < 1)))

$$3.219 \quad \int \frac{1}{\sqrt{1-x^2} \sqrt{2+4x^2}} dx$$

Optimal. Leaf size=10

$$\frac{\text{EllipticF}(\sin^{-1}(x), -2)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,I*2^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {419}

$$\frac{F(\sin^{-1}(x) | -2)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 - x^2]*Sqrt[2 + 4*x^2]),x]

[Out] EllipticF[ArcSin[x], -2]/Sqrt[2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{1-x^2} \sqrt{2+4x^2}} dx = \frac{F(\sin^{-1}(x) | -2)}{\sqrt{2}}$$

Mathematica [C] time = 0.03, size = 58, normalized size = 5.80

$$\frac{i\sqrt{1-x^2} \sqrt{2x^2+1} \text{EllipticF}\left(i \sinh^{-1}(\sqrt{2}x), -\frac{1}{2}\right)}{2\sqrt{-2x^4+x^2+1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 - x^2]*Sqrt[2 + 4*x^2]),x]

[Out] ((-1/2*I)*Sqrt[1 - x^2]*Sqrt[1 + 2*x^2]*EllipticF[I*ArcSinh[Sqrt[2]*x], -1/2])/Sqrt[1 + x^2 - 2*x^4]

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{4x^2+2}\sqrt{-x^2+1}}{2(2x^4-x^2-1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(-1/2*sqrt(4*x^2 + 2)*sqrt(-x^2 + 1)/(2*x^4 - x^2 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{4x^2 + 2} \sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(4*x^2 + 2)*sqrt(-x^2 + 1)), x)

maple [A] time = 0.03, size = 14, normalized size = 1.40

$$\frac{\sqrt{2} \operatorname{EllipticF}(x, i\sqrt{2})}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/2)/(4*x^2+2)^(1/2),x)

[Out] 1/2*EllipticF(x,I*2^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{4x^2 + 2} \sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(4*x^2 + 2)*sqrt(-x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.10

$$\int \frac{1}{\sqrt{1-x^2} \sqrt{4x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/2)*(4*x^2 + 2)^(1/2)),x)

[Out] int(1/((1 - x^2)^(1/2)*(4*x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\sqrt{2} \int \frac{1}{\sqrt{1-x^2} \sqrt{2x^2+1}} dx}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/2)/(4*x**2+2)**(1/2),x)

[Out] sqrt(2)*Integral(1/(sqrt(1 - x**2)*sqrt(2*x**2 + 1)), x)/2

$$3.220 \quad \int \frac{1}{\sqrt{1-x^2} \sqrt{2+3x^2}} dx$$

Optimal. Leaf size=12

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), -\frac{3}{2}\right)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,1/2*I*6^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {419}

$$\frac{F\left(\sin^{-1}(x) \mid -\frac{3}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 - x^2]*Sqrt[2 + 3*x^2]),x]

[Out] EllipticF[ArcSin[x], -3/2]/Sqrt[2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S imp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{1-x^2} \sqrt{2+3x^2}} dx = \frac{F\left(\sin^{-1}(x) \mid -\frac{3}{2}\right)}{\sqrt{2}}$$

Mathematica [A] time = 0.01, size = 12, normalized size = 1.00

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), -\frac{3}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 - x^2]*Sqrt[2 + 3*x^2]),x]

[Out] EllipticF[ArcSin[x], -3/2]/Sqrt[2]

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{3x^2+2}\sqrt{-x^2+1}}{3x^4-x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(3*x^2 + 2)*sqrt(-x^2 + 1)/(3*x^4 - x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{3x^2 + 2}\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(3*x^2 + 2)*sqrt(-x^2 + 1)), x)

maple [A] time = 0.03, size = 14, normalized size = 1.17

$$\frac{\sqrt{2} \operatorname{EllipticF}\left(x, \frac{i\sqrt{6}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/2)/(3*x^2+2)^(1/2),x)

[Out] 1/2*EllipticF(x,1/2*I*6^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{3x^2 + 2}\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(3*x^2 + 2)*sqrt(-x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.08

$$\int \frac{1}{\sqrt{1-x^2}\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1-x^2)^(1/2)*(3*x^2+2)^(1/2)),x)

[Out] int(1/((1-x^2)^(1/2)*(3*x^2+2)^(1/2)), x)

sympy [A] time = 4.31, size = 19, normalized size = 1.58

$$\begin{cases} \frac{\sqrt{2} F\left(\operatorname{asin}(x) \middle| -\frac{3}{2}\right)}{2} & \text{for } x > -1 \wedge x < 1 \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/2)/(3*x**2+2)**(1/2),x)

[Out] Piecewise((sqrt(2)*elliptic_f(asin(x), -3/2)/2, (x > -1) & (x < 1)))

$$3.221 \quad \int \frac{1}{\sqrt{1-x^2} \sqrt{2+2x^2}} dx$$

Optimal. Leaf size=10

$$\frac{\text{EllipticF}(\sin^{-1}(x), -1)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,I)*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {248, 221}

$$\frac{F(\sin^{-1}(x)|-1)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 - x^2]*Sqrt[2 + 2*x^2]),x]

[Out] EllipticF[ArcSin[x], -1]/Sqrt[2]

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := Simp[EllipticF[ArcSin[(Rt[-b, 4]*x)/Rt[a, 4]], -1]/(Rt[a, 4]*Rt[-b, 4]), x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]

Rule 248

Int[((a1_.) + (b1_)*(x_)^(n_))^(p_)*((a2_.) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{1-x^2} \sqrt{2+2x^2}} dx &= \int \frac{1}{\sqrt{2-2x^4}} dx \\ &= \frac{F(\sin^{-1}(x)|-1)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 10, normalized size = 1.00

$$\frac{\text{EllipticF}(\sin^{-1}(x), -1)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 - x^2]*Sqrt[2 + 2*x^2]),x]

[Out] EllipticF[ArcSin[x], -1]/Sqrt[2]

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{2x^2+2}\sqrt{-x^2+1}}{2(x^4-1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(-1/2*sqrt(2*x^2 + 2)*sqrt(-x^2 + 1)/(x^4 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2 + 2}\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(2*x^2 + 2)*sqrt(-x^2 + 1)), x)

maple [A] time = 0.03, size = 10, normalized size = 1.00

$$\frac{\sqrt{2} \operatorname{EllipticF}(x, i)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/2)/(2*x^2+2)^(1/2),x)

[Out] 1/2*EllipticF(x,I)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2 + 2}\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(2*x^2 + 2)*sqrt(-x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.10

$$\int \frac{1}{\sqrt{1-x^2}\sqrt{2x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1-x^2)^(1/2)*(2*x^2+2)^(1/2)),x)

[Out] int(1/((1-x^2)^(1/2)*(2*x^2+2)^(1/2)), x)

sympy [B] time = 5.47, size = 73, normalized size = 7.30

$$\frac{\sqrt{2} G_{6,6}^{5,3} \left(\begin{matrix} \frac{1}{2}, 1, 1 & \frac{3}{4}, \frac{3}{4}, \frac{5}{4} \\ \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1, \frac{5}{4} & 0 \end{matrix} \middle| \frac{e^{-2i\pi}}{x^4} \right)}{16\pi^{\frac{3}{2}}} + \frac{\sqrt{2} G_{6,6}^{3,5} \left(\begin{matrix} -\frac{1}{4}, 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4} & 1 \\ 0, \frac{1}{2}, 0 & -\frac{1}{4}, \frac{1}{4}, \frac{1}{4} \end{matrix} \middle| \frac{1}{x^4} \right)}{16\pi^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/2)/(2*x**2+2)**(1/2),x)

[Out] -sqrt(2)*meijerg(((1/2, 1, 1), (3/4, 3/4, 5/4)), ((1/4, 1/2, 3/4, 1, 5/4), (0,)), exp_polar(-2*I*pi)/x**4)/(16*pi**(3/2)) + sqrt(2)*meijerg(((1/4, 0, 1/4, 1/2, 3/4), (1,)), ((0, 1/2, 0), (-1/4, 1/4, 1/4)), x**(-4))/(16*pi**(3/2))

$$3.222 \quad \int \frac{1}{\sqrt{1-x^2} \sqrt{2+x^2}} dx$$

Optimal. Leaf size=12

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), -\frac{1}{2}\right)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,1/2*I*2^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {419}

$$\frac{F\left(\sin^{-1}(x) \mid -\frac{1}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 - x^2]*Sqrt[2 + x^2]),x]

[Out] EllipticF[ArcSin[x], -1/2]/Sqrt[2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S imp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{1-x^2} \sqrt{2+x^2}} dx = \frac{F\left(\sin^{-1}(x) \mid -\frac{1}{2}\right)}{\sqrt{2}}$$

Mathematica [C] time = 0.02, size = 18, normalized size = 1.50

$$-i \text{EllipticF}\left(i \sinh^{-1}\left(\frac{x}{\sqrt{2}}\right), -2\right)$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 - x^2]*Sqrt[2 + x^2]),x]

[Out] (-I)*EllipticF[I*ArcSinh[x/Sqrt[2]], -2]

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2+2}\sqrt{-x^2+1}}{x^4+x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(x^2 + 2)*sqrt(-x^2 + 1)/(x^4 + x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 2} \sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 2)*sqrt(-x^2 + 1)), x)

maple [A] time = 0.03, size = 14, normalized size = 1.17

$$\frac{\sqrt{2} \operatorname{EllipticF}\left(x, \frac{i\sqrt{2}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/2)/(x^2+2)^(1/2),x)

[Out] 1/2*EllipticF(x,1/2*I*2^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 2} \sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 2)*sqrt(-x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.08

$$\int \frac{1}{\sqrt{1 - x^2} \sqrt{x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/2)*(x^2 + 2)^(1/2)),x)

[Out] int(1/((1 - x^2)^(1/2)*(x^2 + 2)^(1/2)), x)

sympy [A] time = 2.50, size = 19, normalized size = 1.58

$$\begin{cases} \frac{\sqrt{2} F\left(\operatorname{asin}(x) \middle| -\frac{1}{2}\right)}{2} & \text{for } x > -1 \wedge x < 1 \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/2)/(x**2+2)**(1/2),x)

[Out] Piecewise((sqrt(2)*elliptic_f(asin(x), -1/2)/2, (x > -1) & (x < 1)))

$$3.223 \quad \int \frac{1}{\sqrt{1-x^2} \sqrt{2-x^2}} dx$$

Optimal. Leaf size=12

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), \frac{1}{2}\right)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,1/2*2^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {419}

$$\frac{F\left(\sin^{-1}(x)\middle|\frac{1}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 - x^2]*Sqrt[2 - x^2]),x]

[Out] EllipticF[ArcSin[x], 1/2]/Sqrt[2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S imp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{1-x^2} \sqrt{2-x^2}} dx = \frac{F\left(\sin^{-1}(x)\middle|\frac{1}{2}\right)}{\sqrt{2}}$$

Mathematica [A] time = 0.01, size = 12, normalized size = 1.00

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), \frac{1}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 - x^2]*Sqrt[2 - x^2]),x]

[Out] EllipticF[ArcSin[x], 1/2]/Sqrt[2]

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+2}\sqrt{-x^2+1}}{x^4-3x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(-x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-x^2+2)*sqrt(-x^2+1)/(x^4-3*x^2+2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2 + 2} \sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(-x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2 + 2)*sqrt(-x^2 + 1)), x)

maple [A] time = 0.03, size = 13, normalized size = 1.08

$$\frac{\sqrt{2} \operatorname{EllipticF}\left(x, \frac{\sqrt{2}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/2)/(-x^2+2)^(1/2),x)

[Out] 1/2*EllipticF(x,1/2*2^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2 + 2} \sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(-x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 + 2)*sqrt(-x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.08

$$\int \frac{1}{\sqrt{1-x^2} \sqrt{2-x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1-x^2)^(1/2)*(2-x^2)^(1/2)),x)

[Out] int(1/((1-x^2)^(1/2)*(2-x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-(x-1)(x+1)} \sqrt{2-x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/2)/(-x**2+2)**(1/2),x)

[Out] Integral(1/(sqrt(-(x-1)*(x+1))*sqrt(2-x**2)), x)

$$3.224 \quad \int \frac{1}{\sqrt{2-2x^2} \sqrt{1-x^2}} dx$$

Optimal. Leaf size=8

$$\frac{\tanh^{-1}(x)}{\sqrt{2}}$$

[Out] 1/2*arctanh(x)*2^(1/2)

Rubi [A] time = 0.00, antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {22, 206}

$$\frac{\tanh^{-1}(x)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 2*x^2]*Sqrt[1 - x^2]),x]

[Out] ArcTanh[x]/Sqrt[2]

Rule 22

Int[(u_.)*((a_) + (b_.)*(v_))^(m_)*((c_) + (d_.)*(v_))^(n_), x_Symbol] :> Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[b*c - a*d, 0] && GtQ[b/d, 0] && !(IntegerQ[m] || IntegerQ[n])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rubi steps

$$\int \frac{1}{\sqrt{2-2x^2} \sqrt{1-x^2}} dx = \frac{\int \frac{1}{1-x^2} dx}{\sqrt{2}} = \frac{\tanh^{-1}(x)}{\sqrt{2}}$$

Mathematica [B] time = 0.01, size = 26, normalized size = 3.25

$$\frac{\frac{1}{2} \log(1-x) - \frac{1}{2} \log(x+1)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 2*x^2]*Sqrt[1 - x^2]),x]

[Out] -((Log[1 - x]/2 - Log[1 + x]/2)/Sqrt[2])

fricas [B] time = 0.53, size = 68, normalized size = 8.50

$$\frac{1}{8} \sqrt{2} \log \left(-\frac{x^6 + 5x^4 - 2\sqrt{2}(x^3 + x)\sqrt{-x^2 + 1}\sqrt{-2x^2 + 2} - 5x^2 - 1}{x^6 - 3x^4 + 3x^2 - 1} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] 1/8*sqrt(2)*log(-(x^6 + 5*x^4 - 2*sqrt(2)*(x^3 + x)*sqrt(-x^2 + 1)*sqrt(-2*x^2 + 2) - 5*x^2 - 1)/(x^6 - 3*x^4 + 3*x^2 - 1))

giac [B] time = 0.58, size = 19, normalized size = 2.38

$$\frac{1}{4}\sqrt{2}\log(x+1) - \frac{1}{4}\sqrt{2}\log(x-1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(2)*log(x + 1) - 1/4*sqrt(2)*log(x - 1)

maple [A] time = 0.32, size = 8, normalized size = 1.00

$$\frac{\sqrt{2} \operatorname{arctanh}(x)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-2*x^2+2)^(1/2)/(-x^2+1)^(1/2),x)

[Out] 1/2*arctanh(x)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+1}\sqrt{-2x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 + 1)*sqrt(-2*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.12

$$\int \frac{1}{\sqrt{1-x^2}\sqrt{2-2x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/2)*(2 - 2*x^2)^(1/2)),x)

[Out] int(1/((1 - x^2)^(1/2)*(2 - 2*x^2)^(1/2)), x)

sympy [A] time = 2.32, size = 22, normalized size = 2.75

$$-\sqrt{2} \begin{cases} -\frac{\operatorname{acoth}(x)}{2} & \text{for } x^2 > 1 \\ -\frac{\operatorname{atanh}(x)}{2} & \text{for } x^2 < 1 \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x**2+2)**(1/2)/(-x**2+1)**(1/2),x)

[Out] -sqrt(2)*Piecewise((-acoth(x)/2, x**2 > 1), (-atanh(x)/2, x**2 < 1))

$$3.225 \quad \int \frac{1}{\sqrt{2-3x^2} \sqrt{1-x^2}} dx$$

Optimal. Leaf size=12

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), \frac{3}{2}\right)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,1/2*6^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {419}

$$\frac{F\left(\sin^{-1}(x)\middle|\frac{3}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 3*x^2]*Sqrt[1 - x^2]),x]

[Out] EllipticF[ArcSin[x], 3/2]/Sqrt[2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S
imp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt
[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ
[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{2-3x^2} \sqrt{1-x^2}} dx = \frac{F\left(\sin^{-1}(x)\middle|\frac{3}{2}\right)}{\sqrt{2}}$$

Mathematica [A] time = 0.01, size = 12, normalized size = 1.00

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), \frac{3}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 3*x^2]*Sqrt[1 - x^2]),x]

[Out] EllipticF[ArcSin[x], 3/2]/Sqrt[2]

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+1}\sqrt{-3x^2+2}}{3x^4-5x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-x^2 + 1)*sqrt(-3*x^2 + 2)/(3*x^4 - 5*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+1}\sqrt{-3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2 + 1)*sqrt(-3*x^2 + 2)), x)

maple [A] time = 0.03, size = 13, normalized size = 1.08

$$\frac{\sqrt{2} \operatorname{EllipticF}\left(x, \frac{\sqrt{6}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*x^2+2)^(1/2)/(-x^2+1)^(1/2),x)

[Out] 1/2*EllipticF(x,1/2*6^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+1}\sqrt{-3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 + 1)*sqrt(-3*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.08

$$\int \frac{1}{\sqrt{1-x^2}\sqrt{2-3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/2)*(2 - 3*x^2)^(1/2)),x)

[Out] int(1/((1 - x^2)^(1/2)*(2 - 3*x^2)^(1/2)), x)

sympy [A] time = 3.65, size = 17, normalized size = 1.42

$$\begin{cases} \frac{\sqrt{2}F\left(\operatorname{asin}(x)\middle|\frac{3}{2}\right)}{2} & \text{for } x > -1 \wedge x < 1 \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x**2+2)**(1/2)/(-x**2+1)**(1/2),x)

[Out] Piecewise((sqrt(2)*elliptic_f(asin(x), 3/2)/2, (x > -1) & (x < 1)))

$$3.226 \quad \int \frac{1}{\sqrt{2-4x^2} \sqrt{1-x^2}} dx$$

Optimal. Leaf size=10

$$\frac{\text{EllipticF}(\sin^{-1}(x), 2)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,2^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {419}

$$\frac{F(\sin^{-1}(x)|2)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 4*x^2]*Sqrt[1 - x^2]),x]

[Out] EllipticF[ArcSin[x], 2]/Sqrt[2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{2-4x^2} \sqrt{1-x^2}} dx = \frac{F(\sin^{-1}(x)|2)}{\sqrt{2}}$$

Mathematica [A] time = 0.01, size = 10, normalized size = 1.00

$$\frac{\text{EllipticF}(\sin^{-1}(x), 2)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 4*x^2]*Sqrt[1 - x^2]),x]

[Out] EllipticF[ArcSin[x], 2]/Sqrt[2]

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+1} \sqrt{-4x^2+2}}{2(2x^4-3x^2+1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(1/2*sqrt(-x^2 + 1)*sqrt(-4*x^2 + 2)/(2*x^4 - 3*x^2 + 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+1}\sqrt{-4x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2 + 1)*sqrt(-4*x^2 + 2)), x)

maple [A] time = 0.03, size = 11, normalized size = 1.10

$$\frac{\sqrt{2} \operatorname{EllipticF}(x, \sqrt{2})}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-4*x^2+2)^(1/2)/(-x^2+1)^(1/2),x)

[Out] 1/2*EllipticF(x,2^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+1}\sqrt{-4x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 + 1)*sqrt(-4*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.10

$$\int \frac{1}{\sqrt{1-x^2}\sqrt{2-4x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/2)*(2 - 4*x^2)^(1/2)),x)

[Out] int(1/((1 - x^2)^(1/2)*(2 - 4*x^2)^(1/2)), x)

sympy [A] time = 5.85, size = 39, normalized size = 3.90

$$\frac{\sqrt{2} \left(\left\{ \frac{\sqrt{2} F\left(\operatorname{asin}\left(\sqrt{2}x\right)\middle|\frac{1}{2}\right)}{2} \quad \text{for } x > -\frac{\sqrt{2}}{2} \wedge x < \frac{\sqrt{2}}{2} \right\} \right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x**2+2)**(1/2)/(-x**2+1)**(1/2),x)

[Out] sqrt(2)*Piecewise((sqrt(2)*elliptic_f(asin(sqrt(2)*x), 1/2)/2, (x > -sqrt(2)/2) & (x < sqrt(2)/2)))/2

$$3.227 \quad \int \frac{1}{\sqrt{2-5x^2} \sqrt{1-x^2}} dx$$

Optimal. Leaf size=12

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), \frac{5}{2}\right)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,1/2*10^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {419}

$$\frac{F\left(\sin^{-1}(x)\middle|\frac{5}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 5*x^2]*Sqrt[1 - x^2]),x]

[Out] EllipticF[ArcSin[x], 5/2]/Sqrt[2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S imp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{2-5x^2} \sqrt{1-x^2}} dx = \frac{F\left(\sin^{-1}(x)\middle|\frac{5}{2}\right)}{\sqrt{2}}$$

Mathematica [A] time = 0.01, size = 12, normalized size = 1.00

$$\frac{\text{EllipticF}\left(\sin^{-1}(x), \frac{5}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 5*x^2]*Sqrt[1 - x^2]),x]

[Out] EllipticF[ArcSin[x], 5/2]/Sqrt[2]

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+1}\sqrt{-5x^2+2}}{5x^4-7x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-x^2 + 1)*sqrt(-5*x^2 + 2)/(5*x^4 - 7*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+1}\sqrt{-5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2 + 1)*sqrt(-5*x^2 + 2)), x)

maple [A] time = 0.03, size = 13, normalized size = 1.08

$$\frac{\sqrt{2} \operatorname{EllipticF}\left(x, \frac{\sqrt{10}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-5*x^2+2)^(1/2)/(-x^2+1)^(1/2),x)

[Out] 1/2*EllipticF(x,1/2*10^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+1}\sqrt{-5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 + 1)*sqrt(-5*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.08

$$\int \frac{1}{\sqrt{1-x^2}\sqrt{2-5x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/2)*(2 - 5*x^2)^(1/2)),x)

[Out] int(1/((1 - x^2)^(1/2)*(2 - 5*x^2)^(1/2)), x)

sympy [A] time = 3.76, size = 34, normalized size = 2.83

$$\begin{cases} \frac{\sqrt{5}F\left(\operatorname{asin}\left(\frac{\sqrt{10}x}{2}\right)\middle|\frac{2}{5}\right)}{5} & \text{for } x > -\frac{\sqrt{10}}{5} \wedge x < \frac{\sqrt{10}}{5} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x**2+2)**(1/2)/(-x**2+1)**(1/2),x)

[Out] Piecewise((sqrt(5)*elliptic_f(asin(sqrt(10)*x/2), 2/5)/5, (x > -sqrt(10)/5) & (x < sqrt(10)/5)))

$$3.228 \quad \int \frac{1}{\sqrt{1+x^2} \sqrt{2+5x^2}} dx$$

Optimal. Leaf size=51

$$\frac{\sqrt{5x^2+2} \operatorname{EllipticF}\left(\tan^{-1}(x), -\frac{3}{2}\right)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{5x^2+2}{x^2+1}}}$$

[Out] $1/2*(1/(x^2+1))^{(1/2)}*\operatorname{EllipticF}(x/(x^2+1)^{(1/2)}, 1/2*I*6^{(1/2)})*(5*x^2+2)^{(1/2)}*2^{(1/2)}/((5*x^2+2)/(x^2+1))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {418}

$$\frac{\sqrt{5x^2+2} F\left(\tan^{-1}(x) \mid -\frac{3}{2}\right)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{5x^2+2}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 + x^2]*Sqrt[2 + 5*x^2]),x]

[Out] (Sqrt[2 + 5*x^2]*EllipticF[ArcTan[x], -3/2])/(Sqrt[2]*Sqrt[1 + x^2]*Sqrt[(2 + 5*x^2)/(1 + x^2)])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{1+x^2} \sqrt{2+5x^2}} dx = \frac{\sqrt{2+5x^2} F\left(\tan^{-1}(x) \mid -\frac{3}{2}\right)}{\sqrt{2} \sqrt{1+x^2} \sqrt{\frac{2+5x^2}{1+x^2}}}$$

Mathematica [C] time = 0.02, size = 19, normalized size = 0.37

$$\frac{i \operatorname{EllipticF}\left(i \sinh^{-1}(x), \frac{5}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 + x^2]*Sqrt[2 + 5*x^2]),x]

[Out] ((-I)*EllipticF[I*ArcSinh[x], 5/2])/Sqrt[2]

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\frac{\sqrt{5x^2+2} \sqrt{x^2+1}}{5x^4+7x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(5*x^2 + 2)*sqrt(x^2 + 1)/(5*x^4 + 7*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{5x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(5*x^2 + 2)*sqrt(x^2 + 1)), x)

maple [A] time = 0.04, size = 17, normalized size = 0.33

$$-\frac{i\sqrt{2} \operatorname{EllipticF}\left(ix, \frac{\sqrt{10}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2+1)^(1/2)/(5*x^2+2)^(1/2),x)

[Out] -1/2*I*EllipticF(I*x,1/2*10^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{5x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(5*x^2 + 2)*sqrt(x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{5x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(5*x^2 + 2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(5*x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{5x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2+1)**(1/2)/(5*x**2+2)**(1/2),x)

[Out] Integral(1/(sqrt(x**2 + 1)*sqrt(5*x**2 + 2)), x)

$$3.229 \quad \int \frac{1}{\sqrt{1+x^2} \sqrt{2+4x^2}} dx$$

Optimal. Leaf size=49

$$\frac{\sqrt{2x^2+1} \operatorname{EllipticF}(\tan^{-1}(x), -1)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{2x^2+1}{x^2+1}}}$$

[Out] $1/2*(1/(x^2+1))^{(1/2)}*\operatorname{EllipticF}(x/(x^2+1)^{(1/2)}, I)*(2*x^2+1)^{(1/2)}*2^{(1/2)}/((2*x^2+1)/(x^2+1))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {418}

$$\frac{\sqrt{2x^2+1} F(\tan^{-1}(x) | -1)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{2x^2+1}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 + x^2]*Sqrt[2 + 4*x^2]),x]

[Out] (Sqrt[1 + 2*x^2]*EllipticF[ArcTan[x], -1])/(Sqrt[2]*Sqrt[1 + x^2]*Sqrt[(1 + 2*x^2)/(1 + x^2)])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{1+x^2} \sqrt{2+4x^2}} dx = \frac{\sqrt{1+2x^2} F(\tan^{-1}(x) | -1)}{\sqrt{2} \sqrt{1+x^2} \sqrt{\frac{1+2x^2}{1+x^2}}}$$

Mathematica [C] time = 0.03, size = 17, normalized size = 0.35

$$-\frac{i \operatorname{EllipticF}(i \sinh^{-1}(x), 2)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 + x^2]*Sqrt[2 + 4*x^2]),x]

[Out] ((-I)*EllipticF[I*ArcSinh[x], 2])/Sqrt[2]

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\frac{\sqrt{4x^2+2}\sqrt{x^2+1}}{2(2x^4+3x^2+1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(1/2*sqrt(4*x^2 + 2)*sqrt(x^2 + 1)/(2*x^4 + 3*x^2 + 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{4x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(4*x^2 + 2)*sqrt(x^2 + 1)), x)

maple [A] time = 0.03, size = 15, normalized size = 0.31

$$\frac{i\sqrt{2} \operatorname{EllipticF}(ix, \sqrt{2})}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2+1)^(1/2)/(4*x^2+2)^(1/2),x)

[Out] -1/2*I*EllipticF(I*x, 2^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{4x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(4*x^2 + 2)*sqrt(x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{4x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(4*x^2 + 2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(4*x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\sqrt{2} \int \frac{1}{\sqrt{x^2+1} \sqrt{2x^2+1}} dx}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2+1)**(1/2)/(4*x**2+2)**(1/2),x)

[Out] sqrt(2)*Integral(1/(sqrt(x**2 + 1)*sqrt(2*x**2 + 1)), x)/2

$$3.230 \quad \int \frac{1}{\sqrt{1+x^2} \sqrt{2+3x^2}} dx$$

Optimal. Leaf size=51

$$\frac{\sqrt{3x^2+2} \operatorname{EllipticF}\left(\tan^{-1}(x), -\frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{3x^2+2}{x^2+1}}}$$

[Out] $1/2*(1/(x^2+1))^{(1/2)}*\operatorname{EllipticF}(x/(x^2+1)^{(1/2)}, 1/2*I*2^{(1/2)})*(3*x^2+2)^{(1/2)}*2^{(1/2)}/((3*x^2+2)/(x^2+1))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {418}

$$\frac{\sqrt{3x^2+2} F\left(\tan^{-1}(x) \mid -\frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{3x^2+2}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 + x^2]*Sqrt[2 + 3*x^2]),x]

[Out] (Sqrt[2 + 3*x^2]*EllipticF[ArcTan[x], -1/2])/(Sqrt[2]*Sqrt[1 + x^2]*Sqrt[(2 + 3*x^2)/(1 + x^2)])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{1+x^2} \sqrt{2+3x^2}} dx = \frac{\sqrt{2+3x^2} F\left(\tan^{-1}(x) \mid -\frac{1}{2}\right)}{\sqrt{2} \sqrt{1+x^2} \sqrt{\frac{2+3x^2}{1+x^2}}}$$

Mathematica [C] time = 0.02, size = 19, normalized size = 0.37

$$\frac{i \operatorname{EllipticF}\left(i \sinh^{-1}(x), \frac{3}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 + x^2]*Sqrt[2 + 3*x^2]),x]

[Out] ((-I)*EllipticF[I*ArcSinh[x], 3/2])/Sqrt[2]

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\frac{\sqrt{3x^2+2} \sqrt{x^2+1}}{3x^4+5x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(3*x^2 + 2)*sqrt(x^2 + 1)/(3*x^4 + 5*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{3x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(3*x^2 + 2)*sqrt(x^2 + 1)), x)

maple [A] time = 0.03, size = 17, normalized size = 0.33

$$\frac{i\sqrt{2} \operatorname{EllipticF}\left(ix, \frac{\sqrt{6}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2+1)^(1/2)/(3*x^2+2)^(1/2),x)

[Out] -1/2*I*EllipticF(I*x,1/2*6^(1/2))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{3x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(3*x^2 + 2)*sqrt(x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(3*x^2 + 2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(3*x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2+1)**(1/2)/(3*x**2+2)**(1/2),x)

[Out] Integral(1/(sqrt(x**2 + 1)*sqrt(3*x**2 + 2)), x)

$$3.231 \quad \int \frac{1}{\sqrt{1+x^2} \sqrt{2+2x^2}} dx$$

Optimal. Leaf size=8

$$\frac{\tan^{-1}(x)}{\sqrt{2}}$$

[Out] 1/2*arctan(x)*2^(1/2)

Rubi [A] time = 0.00, antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {22, 203}

$$\frac{\tan^{-1}(x)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 + x^2]*Sqrt[2 + 2*x^2]),x]

[Out] ArcTan[x]/Sqrt[2]

Rule 22

Int[(u_.)*((a_) + (b_.)*(v_))^(m_)*((c_) + (d_.)*(v_))^(n_), x_Symbol] :> Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[b*c - a*d, 0] && GtQ[b/d, 0] && !(IntegerQ[m] || IntegerQ[n])

Rule 203

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{1+x^2} \sqrt{2+2x^2}} dx &= \sqrt{2} \int \frac{1}{2+2x^2} dx \\ &= \frac{\tan^{-1}(x)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] time = 0.00, size = 8, normalized size = 1.00

$$\frac{\tan^{-1}(x)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 + x^2]*Sqrt[2 + 2*x^2]),x]

[Out] ArcTan[x]/Sqrt[2]

fricas [B] time = 1.04, size = 34, normalized size = 4.25

$$-\frac{1}{4} \sqrt{2} \arctan\left(\frac{\sqrt{2} \sqrt{2x^2 + 2} \sqrt{x^2 + 1} x}{x^4 - 1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="fricas")

[Out] -1/4*sqrt(2)*arctan(sqrt(2)*sqrt(2*x^2 + 2)*sqrt(x^2 + 1)*x/(x^4 - 1))

giac [B] time = 0.57, size = 26, normalized size = 3.25

$$\frac{1}{4} \sqrt{2} i \log(ix - 1) - \frac{1}{4} \sqrt{2} i \log(-ix - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(2)*i*log(i*x - 1) - 1/4*sqrt(2)*i*log(-i*x - 1)

maple [A] time = 0.31, size = 8, normalized size = 1.00

$$\frac{\sqrt{2} \arctan(x)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2+1)^(1/2)/(2*x^2+2)^(1/2),x)

[Out] 1/2*arctan(x)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(2*x^2 + 2)*sqrt(x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.12

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{2x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(2*x^2 + 2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(2*x^2 + 2)^(1/2)), x)

sympy [A] time = 2.49, size = 8, normalized size = 1.00

$$\frac{\sqrt{2} \operatorname{atan}(x)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2+1)**(1/2)/(2*x**2+2)**(1/2),x)

[Out] sqrt(2)*atan(x)/2

$$3.232 \quad \int \frac{1}{\sqrt{1+x^2} \sqrt{2+x^2}} dx$$

Optimal. Leaf size=47

$$\frac{\sqrt{x^2+2} \operatorname{EllipticF}\left(\tan^{-1}(x), \frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{x^2+2}{x^2+1}}}$$

[Out] $1/2*(1/(x^2+1))^{(1/2)}*\operatorname{EllipticF}(x/(x^2+1)^{(1/2)}, 1/2*2^{(1/2)})*(x^2+2)^{(1/2)}*2^{(1/2)}/((x^2+2)/(x^2+1))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {418}

$$\frac{\sqrt{x^2+2} F\left(\tan^{-1}(x) \middle| \frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2+1} \sqrt{\frac{x^2+2}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 + x^2]*Sqrt[2 + x^2]), x]

[Out] (Sqrt[2 + x^2]*EllipticF[ArcTan[x], 1/2])/(Sqrt[2]*Sqrt[1 + x^2]*Sqrt[(2 + x^2)/(1 + x^2)])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{1+x^2} \sqrt{2+x^2}} dx = \frac{\sqrt{2+x^2} F\left(\tan^{-1}(x) \middle| \frac{1}{2}\right)}{\sqrt{2} \sqrt{1+x^2} \sqrt{\frac{2+x^2}{1+x^2}}}$$

Mathematica [C] time = 0.02, size = 19, normalized size = 0.40

$$\frac{i \operatorname{EllipticF}\left(i \sinh^{-1}(x), \frac{1}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 + x^2]*Sqrt[2 + x^2]), x]

[Out] ((-I)*EllipticF[I*ArcSinh[x], 1/2])/Sqrt[2]

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\frac{\sqrt{x^2+2} \sqrt{x^2+1}}{x^4+3x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(x^2 + 2)*sqrt(x^2 + 1)/(x^4 + 3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 2)*sqrt(x^2 + 1)), x)

maple [C] time = 0.03, size = 15, normalized size = 0.32

$$-i \operatorname{EllipticF}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2+1)^(1/2)/(x^2+2)^(1/2), x)

[Out] -I*EllipticF(1/2*I*x*2^(1/2), 2^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+1)^(1/2)/(x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 2)*sqrt(x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(x^2 + 2)^(1/2)), x)

[Out] int(1/((x^2 + 1)^(1/2)*(x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2+1)**(1/2)/(x**2+2)**(1/2), x)

[Out] Integral(1/(sqrt(x**2 + 1)*sqrt(x**2 + 2)), x)

$$3.233 \quad \int \frac{1}{\sqrt{2-x^2} \sqrt{1+x^2}} dx$$

Optimal. Leaf size=10

$$\text{EllipticF}\left(\sin^{-1}\left(\frac{x}{\sqrt{2}}\right), -2\right)$$

[Out] EllipticF(1/2*x*2^(1/2), I*2^(1/2))

Rubi [A] time = 0.01, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {419}

$$F\left(\sin^{-1}\left(\frac{x}{\sqrt{2}}\right) \middle| -2\right)$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[x/Sqrt[2]], -2]

Rule 419

Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])]

Rubi steps

$$\int \frac{1}{\sqrt{2-x^2} \sqrt{1+x^2}} dx = F\left(\sin^{-1}\left(\frac{x}{\sqrt{2}}\right) \middle| -2\right)$$

Mathematica [C] time = 0.02, size = 19, normalized size = 1.90

$$\frac{i \text{EllipticF}\left(i \sinh^{-1}(x), -\frac{1}{2}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - x^2]*Sqrt[1 + x^2]),x]

[Out] ((-I)*EllipticF[I*ArcSinh[x], -1/2])/Sqrt[2]

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2+1} \sqrt{-x^2+2}}{x^4-x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(x^2 + 1)*sqrt(-x^2 + 2)/(x^4 - x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2+1} \sqrt{-x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-x^2 + 2)), x)

maple [A] time = 0.03, size = 14, normalized size = 1.40

$$\text{EllipticF}\left(\frac{\sqrt{2}x}{2}, i\sqrt{2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+2)^(1/2)/(x^2+1)^(1/2),x)

[Out] EllipticF(1/2*2^(1/2)*x,I*2^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{-x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.10

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{2 - x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(2 - x^2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(2 - x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2 - x^2} \sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+2)**(1/2)/(x**2+1)**(1/2),x)

[Out] Integral(1/(sqrt(2 - x**2)*sqrt(x**2 + 1)), x)

$$3.234 \quad \int \frac{1}{\sqrt{2-2x^2} \sqrt{1+x^2}} dx$$

Optimal. Leaf size=10

$$\frac{\text{EllipticF}(\sin^{-1}(x), -1)}{\sqrt{2}}$$

[Out] 1/2*EllipticF(x,I)*2^(1/2)

Rubi [A] time = 0.00, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {248, 221}

$$\frac{F(\sin^{-1}(x)|-1)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 2*x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[x], -1]/Sqrt[2]

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := Simp[EllipticF[ArcSin[(Rt[-b, 4]*x)/Rt[a, 4]], -1]/(Rt[a, 4]*Rt[-b, 4]), x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]

Rule 248

Int[((a1_.) + (b1_)*(x_)^(n_))^(p_)*((a2_.) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-2x^2} \sqrt{1+x^2}} dx &= \int \frac{1}{\sqrt{2-2x^4}} dx \\ &= \frac{F(\sin^{-1}(x)|-1)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 10, normalized size = 1.00

$$\frac{\text{EllipticF}(\sin^{-1}(x), -1)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 2*x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[x], -1]/Sqrt[2]

fricas [F] time = 0.71, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2+1}\sqrt{-2x^2+2}}{2(x^4-1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(-1/2*sqrt(x^2 + 1)*sqrt(-2*x^2 + 2)/(x^4 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{-2x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-2*x^2 + 2)), x)

maple [A] time = 0.02, size = 10, normalized size = 1.00

$$\frac{\sqrt{2} \operatorname{EllipticF}(x, i)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-2*x^2+2)^(1/2)/(x^2+1)^(1/2),x)

[Out] 1/2*EllipticF(x,I)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{-2x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-2*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.10

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{2 - 2x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(2 - 2*x^2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(2 - 2*x^2)^(1/2)), x)

sympy [B] time = 5.26, size = 76, normalized size = 7.60

$$\frac{\sqrt{2} i G_{6,6}^{5,3} \left(\begin{matrix} \frac{1}{2}, 1, 1 & \frac{3}{4}, \frac{3}{4}, \frac{5}{4} \\ \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1, \frac{5}{4} & 0 \end{matrix} \middle| \frac{1}{x^4} \right)}{16\pi^{\frac{3}{2}}} - \frac{\sqrt{2} i G_{6,6}^{3,5} \left(\begin{matrix} -\frac{1}{4}, 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4} & 1 \\ 0, \frac{1}{2}, 0 & -\frac{1}{4}, \frac{1}{4}, \frac{1}{4} \end{matrix} \middle| \frac{e^{-2i\pi}}{x^4} \right)}{16\pi^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x**2+2)**(1/2)/(x**2+1)**(1/2),x)

[Out] sqrt(2)*I*meijerg(((1/2, 1, 1), (3/4, 3/4, 5/4)), ((1/4, 1/2, 3/4, 1, 5/4), (0,)), x**(-4))/(16*pi**(3/2)) - sqrt(2)*I*meijerg(((1/4, 0, 1/4, 1/2, 3/4), (1,)), ((0, 1/2, 0), (-1/4, 1/4, 1/4)), exp_polar(-2*I*pi)/x**4)/(16*pi**(3/2))

$$3.235 \quad \int \frac{1}{\sqrt{2-3x^2} \sqrt{1+x^2}} dx$$

Optimal. Leaf size=20

$$\frac{\text{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right), -\frac{2}{3}\right)}{\sqrt{3}}$$

[Out] 1/3*EllipticF(1/2*x*6^(1/2),1/3*I*6^(1/2))*3^(1/2)

Rubi [A] time = 0.01, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {419}

$$\frac{F\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right) \middle| -\frac{2}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 3*x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[Sqrt[3/2]*x], -2/3]/Sqrt[3]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S imp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{2-3x^2} \sqrt{1+x^2}} dx = \frac{F\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right) \middle| -\frac{2}{3}\right)}{\sqrt{3}}$$

Mathematica [A] time = 0.00, size = 20, normalized size = 1.00

$$\frac{\text{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right), -\frac{2}{3}\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 3*x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[Sqrt[3/2]*x], -2/3]/Sqrt[3]

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2+1}\sqrt{-3x^2+2}}{3x^4+x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(x^2 + 1)*sqrt(-3*x^2 + 2)/(3*x^4 + x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-3*x^2 + 2)), x)

maple [A] time = 0.03, size = 19, normalized size = 0.95

$$\frac{\sqrt{3} \operatorname{EllipticF}\left(\frac{\sqrt{6}x}{2}, \frac{i\sqrt{6}}{3}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*x^2+2)^(1/2)/(x^2+1)^(1/2),x)

[Out] 1/3*EllipticF(1/2*6^(1/2)*x,1/3*I*6^(1/2))*3^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-3*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(2 - 3*x^2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(2 - 3*x^2)^(1/2)), x)

sympy [A] time = 4.09, size = 36, normalized size = 1.80

$$\left\{ \frac{\sqrt{3} F\left(\operatorname{asin}\left(\frac{\sqrt{6}x}{2}\right) \middle| -\frac{2}{3}\right)}{3} \quad \text{for } x > -\frac{\sqrt{6}}{3} \wedge x < \frac{\sqrt{6}}{3} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x**2+2)**(1/2)/(x**2+1)**(1/2),x)

[Out] Piecewise((sqrt(3)*elliptic_f(asin(sqrt(6)*x/2), -2/3)/3, (x > -sqrt(6)/3) & (x < sqrt(6)/3)))

$$3.236 \quad \int \frac{1}{\sqrt{2-4x^2} \sqrt{1+x^2}} dx$$

Optimal. Leaf size=16

$$\frac{1}{2} \operatorname{EllipticF}\left(\sin^{-1}(\sqrt{2}x), -\frac{1}{2}\right)$$

[Out] 1/2*EllipticF(x*2^(1/2),1/2*I*2^(1/2))

Rubi [A] time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {419}

$$\frac{1}{2} F\left(\sin^{-1}(\sqrt{2}x) \mid -\frac{1}{2}\right)$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 4*x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[Sqrt[2]*x], -1/2]/2

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{2-4x^2} \sqrt{1+x^2}} dx = \frac{1}{2} F\left(\sin^{-1}(\sqrt{2}x) \mid -\frac{1}{2}\right)$$

Mathematica [A] time = 0.01, size = 16, normalized size = 1.00

$$\frac{1}{2} \operatorname{EllipticF}\left(\sin^{-1}(\sqrt{2}x), -\frac{1}{2}\right)$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 4*x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[Sqrt[2]*x], -1/2]/2

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{\sqrt{x^2+1}\sqrt{-4x^2+2}}{2(2x^4+x^2-1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(-1/2*sqrt(x^2 + 1)*sqrt(-4*x^2 + 2)/(2*x^4 + x^2 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2+1}\sqrt{-4x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-4*x^2 + 2)), x)

maple [A] time = 0.03, size = 15, normalized size = 0.94

$$\frac{\text{EllipticF}\left(\sqrt{2} x, \frac{i\sqrt{2}}{2}\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-4*x^2+2)^(1/2)/(x^2+1)^(1/2),x)

[Out] 1/2*EllipticF(2^(1/2)*x,1/2*I*2^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{-4x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-4*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.06

$$\int \frac{1}{\sqrt{x^2 + 1} \sqrt{2 - 4x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(2 - 4*x^2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(2 - 4*x^2)^(1/2)), x)

sympy [A] time = 6.02, size = 41, normalized size = 2.56

$$\frac{\sqrt{2} \left(\left\{ \frac{\sqrt{2} F\left(\text{asin}\left(\sqrt{2} x\right) \middle| -\frac{1}{2}\right)}{2} \quad \text{for } x > -\frac{\sqrt{2}}{2} \wedge x < \frac{\sqrt{2}}{2} \right\} \right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x**2+2)**(1/2)/(x**2+1)**(1/2),x)

[Out] sqrt(2)*Piecewise((sqrt(2)*elliptic_f(asin(sqrt(2)*x), -1/2)/2, (x > -sqrt(2)/2) & (x < sqrt(2)/2)))/2

$$3.237 \quad \int \frac{1}{\sqrt{2-5x^2} \sqrt{1+x^2}} dx$$

Optimal. Leaf size=20

$$\frac{\text{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right), -\frac{2}{5}\right)}{\sqrt{5}}$$

[Out] 1/5*EllipticF(1/2*x*10^(1/2),1/5*I*10^(1/2))*5^(1/2)

Rubi [A] time = 0.01, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {419}

$$\frac{F\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right) \middle| -\frac{2}{5}\right)}{\sqrt{5}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 5*x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[Sqrt[5/2]*x], -2/5]/Sqrt[5]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> S imp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rubi steps

$$\int \frac{1}{\sqrt{2-5x^2} \sqrt{1+x^2}} dx = \frac{F\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right) \middle| -\frac{2}{5}\right)}{\sqrt{5}}$$

Mathematica [A] time = 0.01, size = 20, normalized size = 1.00

$$\frac{\text{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right), -\frac{2}{5}\right)}{\sqrt{5}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 5*x^2]*Sqrt[1 + x^2]),x]

[Out] EllipticF[ArcSin[Sqrt[5/2]*x], -2/5]/Sqrt[5]

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2+1}\sqrt{-5x^2+2}}{5x^4+3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(x^2 + 1)*sqrt(-5*x^2 + 2)/(5*x^4 + 3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2+1}\sqrt{-5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-5*x^2 + 2)), x)

maple [A] time = 0.05, size = 19, normalized size = 0.95

$$\frac{\sqrt{5} \operatorname{EllipticF}\left(\frac{\sqrt{10}x}{2}, \frac{i\sqrt{10}}{5}\right)}{5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-5*x^2+2)^(1/2)/(x^2+1)^(1/2),x)

[Out] 1/5*EllipticF(1/2*x*10^(1/2),1/5*I*10^(1/2))*5^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2+1}\sqrt{-5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 1)*sqrt(-5*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{1}{\sqrt{x^2+1}\sqrt{2-5x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 1)^(1/2)*(2 - 5*x^2)^(1/2)),x)

[Out] int(1/((x^2 + 1)^(1/2)*(2 - 5*x^2)^(1/2)), x)

sympy [A] time = 4.09, size = 36, normalized size = 1.80

$$\left\{ \frac{\sqrt{5} F\left(\operatorname{asin}\left(\frac{\sqrt{10}x}{2}\right) \middle| -\frac{2}{5}\right)}{5} \quad \text{for } x > -\frac{\sqrt{10}}{5} \wedge x < \frac{\sqrt{10}}{5} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x**2+2)**(1/2)/(x**2+1)**(1/2),x)

[Out] Piecewise((sqrt(5)*elliptic_f(asin(sqrt(10)*x/2), -2/5)/5, (x > -sqrt(10)/5) & (x < sqrt(10)/5))

$$3.238 \quad \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+5x^2}} dx$$

Optimal. Leaf size=32

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), -\frac{5}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

[Out] 1/2*EllipticF(x,1/2*I*10^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {421, 419}

$$\frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \mid -\frac{5}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 + x^2]*Sqrt[2 + 5*x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -5/2])/(Sqrt[2]*Sqrt[-1 + x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+5x^2}} dx &= \frac{\sqrt{1-x^2} \int \frac{1}{\sqrt{1-x^2} \sqrt{2+5x^2}} dx}{\sqrt{-1+x^2}} \\ &= \frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \mid -\frac{5}{2}\right)}{\sqrt{2} \sqrt{-1+x^2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), -\frac{5}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 + x^2]*Sqrt[2 + 5*x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -5/2])/(Sqrt[2]*Sqrt[-1 + x^2])

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{5x^2+2}\sqrt{x^2-1}}{5x^4-3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(5*x^2 + 2)*sqrt(x^2 - 1)/(5*x^4 - 3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{5x^2+2}\sqrt{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(5*x^2 + 2)*sqrt(x^2 - 1)), x)

maple [A] time = 0.04, size = 37, normalized size = 1.16

$$-\frac{i\sqrt{-x^2+1}\sqrt{5}\text{EllipticF}\left(\frac{i\sqrt{10}x}{2}, \frac{i\sqrt{10}}{5}\right)}{5\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2-1)^(1/2)/(5*x^2+2)^(1/2),x)

[Out] -1/5*I*EllipticF(1/2*I*x*10^(1/2), 1/5*I*10^(1/2))*(-x^2+1)^(1/2)*5^(1/2)/(x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{5x^2+2}\sqrt{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(5*x^2 + 2)*sqrt(x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(5*x^2 + 2)^(1/2)),x)

[Out] int(1/((x^2 - 1)^(1/2)*(5*x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{(x-1)(x+1)}\sqrt{5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2-1)**(1/2)/(5*x**2+2)**(1/2),x)

[Out] Integral(1/(sqrt((x - 1)*(x + 1))*sqrt(5*x**2 + 2)), x)

$$3.239 \quad \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+4x^2}} dx$$

Optimal. Leaf size=30

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}(\sin^{-1}(x), -2)}{\sqrt{2} \sqrt{x^2-1}}$$

[Out] 1/2*EllipticF(x,I*2^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {421, 419}

$$\frac{\sqrt{1-x^2} F(\sin^{-1}(x) | -2)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 + x^2]*Sqrt[2 + 4*x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -2])/(Sqrt[2]*Sqrt[-1 + x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+4x^2}} dx &= \frac{\sqrt{1-x^2} \int \frac{1}{\sqrt{1-x^2} \sqrt{2+4x^2}} dx}{\sqrt{-1+x^2}} \\ &= \frac{\sqrt{1-x^2} F(\sin^{-1}(x) | -2)}{\sqrt{2} \sqrt{-1+x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 30, normalized size = 1.00

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}(\sin^{-1}(x), -2)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 + x^2]*Sqrt[2 + 4*x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -2])/(Sqrt[2]*Sqrt[-1 + x^2])

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\frac{\sqrt{4x^2+2}\sqrt{x^2-1}}{2(2x^4-x^2-1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(x^2-1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="fricas")
[Out] integral(1/2*sqrt(4*x^2 + 2)*sqrt(x^2 - 1)/(2*x^4 - x^2 - 1), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00
```

$$\int \frac{1}{\sqrt{4x^2 + 2} \sqrt{x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(x^2-1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="giac")
[Out] integrate(1/(sqrt(4*x^2 + 2)*sqrt(x^2 - 1)), x)
maple [A] time = 0.04, size = 34, normalized size = 1.13
```

$$\frac{i\sqrt{-x^2 + 1} \operatorname{EllipticF}\left(i\sqrt{2} x, \frac{i\sqrt{2}}{2}\right)}{2\sqrt{x^2 - 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x^2-1)^(1/2)/(4*x^2+2)^(1/2),x)
[Out] -1/2*I*EllipticF(I*x*2^(1/2), 1/2*I*2^(1/2))*(-x^2+1)^(1/2)/(x^2-1)^(1/2)
maxima [F] time = 0.00, size = 0, normalized size = 0.00
```

$$\int \frac{1}{\sqrt{4x^2 + 2} \sqrt{x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(x^2-1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="maxima")
[Out] integrate(1/(sqrt(4*x^2 + 2)*sqrt(x^2 - 1)), x)
mupad [F] time = 0.00, size = -1, normalized size = -0.03
```

$$\int \frac{1}{\sqrt{x^2 - 1} \sqrt{4x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((x^2 - 1)^(1/2)*(4*x^2 + 2)^(1/2)),x)
[Out] int(1/((x^2 - 1)^(1/2)*(4*x^2 + 2)^(1/2)), x)
sympy [F] time = 0.00, size = 0, normalized size = 0.00
```

$$\frac{\sqrt{2} \int \frac{1}{\sqrt{x^2-1} \sqrt{2x^2+1}} dx}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(x**2-1)**(1/2)/(4*x**2+2)**(1/2),x)
[Out] sqrt(2)*Integral(1/(sqrt(x**2 - 1)*sqrt(2*x**2 + 1)), x)/2
```

$$3.240 \quad \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+3x^2}} dx$$

Optimal. Leaf size=32

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), -\frac{3}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

[Out] 1/2*EllipticF(x,1/2*I*6^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {421, 419}

$$\frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \middle| -\frac{3}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 + x^2]*Sqrt[2 + 3*x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -3/2])/(Sqrt[2]*Sqrt[-1 + x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+3x^2}} dx &= \frac{\sqrt{1-x^2} \int \frac{1}{\sqrt{1-x^2} \sqrt{2+3x^2}} dx}{\sqrt{-1+x^2}} \\ &= \frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \middle| -\frac{3}{2}\right)}{\sqrt{2} \sqrt{-1+x^2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), -\frac{3}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 + x^2]*Sqrt[2 + 3*x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -3/2])/(Sqrt[2]*Sqrt[-1 + x^2])

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{3x^2+2}\sqrt{x^2-1}}{3x^4-x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(3*x^2 + 2)*sqrt(x^2 - 1)/(3*x^4 - x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{3x^2+2}\sqrt{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(3*x^2 + 2)*sqrt(x^2 - 1)), x)

maple [A] time = 0.04, size = 37, normalized size = 1.16

$$-\frac{i\sqrt{-x^2+1}\sqrt{3}\text{EllipticF}\left(\frac{i\sqrt{6}x}{2}, \frac{i\sqrt{6}}{3}\right)}{3\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2-1)^(1/2)/(3*x^2+2)^(1/2),x)

[Out] -1/3*I*EllipticF(1/2*I*x*6^(1/2),1/3*I*6^(1/2))*(-x^2+1)^(1/2)*3^(1/2)/(x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{3x^2+2}\sqrt{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(3*x^2 + 2)*sqrt(x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(3*x^2 + 2)^(1/2)),x)

[Out] int(1/((x^2 - 1)^(1/2)*(3*x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{(x-1)(x+1)}\sqrt{3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2-1)**(1/2)/(3*x**2+2)**(1/2),x)

[Out] Integral(1/(sqrt((x - 1)*(x + 1))*sqrt(3*x**2 + 2)), x)

$$3.241 \quad \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+2x^2}} dx$$

Optimal. Leaf size=25

$$\frac{1}{2} \text{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{2}x}{\sqrt{x^2-1}}\right), \frac{1}{2}\right)$$

[Out] 1/2*EllipticF(x*2^(1/2)/(x^2-1)^(1/2),1/2*2^(1/2))

Rubi [A] time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {253, 222}

$$\frac{1}{2} F\left(\sin^{-1}\left(\frac{\sqrt{2}x}{\sqrt{x^2-1}}\right) \middle| \frac{1}{2}\right)$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 + x^2]*Sqrt[2 + 2*x^2]),x]

[Out] EllipticF[ArcSin[(Sqrt[2]*x)/Sqrt[-1 + x^2]], 1/2]/2

Rule 222

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[-(a*b), 2]}, Simp[(Sqrt[-a + q*x^2]*Sqrt[(a + q*x^2)/q]*EllipticF[ArcSin[x/Sqrt[(a + q*x^2)/(2*q)]], 1/2])/(Sqrt[2]*Sqrt[-a]*Sqrt[a + b*x^4]), x] /; IntegerQ[q] /; FreeQ[{a, b}, x] && LtQ[a, 0] && GtQ[b, 0]

Rule 253

Int[((a1_.) + (b1_.)*(x_)^(n_))^(p_)*((a2_.) + (b2_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[((a1 + b1*x^n)^FracPart[p]*(a2 + b2*x^n)^FracPart[p])/(a1*a2 + b1*b2*x^(2*n))^FracPart[p], Int[(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+2x^2}} dx &= \frac{\sqrt{-2+2x^4} \int \frac{1}{\sqrt{-2+2x^4}} dx}{\sqrt{-1+x^2} \sqrt{2+2x^2}} \\ &= \frac{1}{2} F\left(\sin^{-1}\left(\frac{\sqrt{2}x}{\sqrt{-1+x^2}}\right) \middle| \frac{1}{2}\right) \end{aligned}$$

Mathematica [C] time = 0.01, size = 46, normalized size = 1.84

$$\frac{x\sqrt{1-x^4} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; x^4\right)}{\sqrt{x^2-1} \sqrt{2x^2+2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 + x^2]*Sqrt[2 + 2*x^2]),x]

[Out] (x*Sqrt[1 - x^4]*Hypergeometric2F1[1/4, 1/2, 5/4, x^4])/(Sqrt[-1 + x^2]*Sqrt[2 + 2*x^2])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{2x^2+2}\sqrt{x^2-1}}{2(x^4-1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(1/2*sqrt(2*x^2 + 2)*sqrt(x^2 - 1)/(x^4 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2+2}\sqrt{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(2*x^2 + 2)*sqrt(x^2 - 1)), x)

maple [C] time = 0.03, size = 30, normalized size = 1.20

$$\frac{i\sqrt{-x^2+1}\sqrt{2}\text{EllipticF}(ix,i)}{2\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2-1)^(1/2)/(2*x^2+2)^(1/2),x)

[Out] -1/2*I*EllipticF(I*x,I)*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2+2}\sqrt{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(2*x^2 + 2)*sqrt(x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.04

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{2x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(2*x^2 + 2)^(1/2)),x)

[Out] int(1/((x^2 - 1)^(1/2)*(2*x^2 + 2)^(1/2)), x)

sympy [C] time = 5.55, size = 75, normalized size = 3.00

$$\frac{\sqrt{2}iG_{6,6}^{5,3}\left(\begin{matrix} \frac{1}{2}, 1, 1 & \frac{3}{4}, \frac{3}{4}, \frac{5}{4} \\ \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1, \frac{5}{4} & 0 \end{matrix} \middle| \frac{e^{2i\pi}}{x^4}\right)}{16\pi^{\frac{3}{2}}} - \frac{\sqrt{2}iG_{6,6}^{3,5}\left(\begin{matrix} -\frac{1}{4}, 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4} & 1 \\ 0, \frac{1}{2}, 0 & -\frac{1}{4}, \frac{1}{4}, \frac{1}{4} \end{matrix} \middle| \frac{1}{x^4}\right)}{16\pi^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(x**2-1)**(1/2)/(2*x**2+2)**(1/2),x)
```

```
[Out] sqrt(2)*I*meijerg(((1/2, 1, 1), (3/4, 3/4, 5/4)), ((1/4, 1/2, 3/4, 1, 5/4),  
  (0,)), exp_polar(2*I*pi)/x**4)/(16*pi**(3/2)) - sqrt(2)*I*meijerg((-1/4,  
  0, 1/4, 1/2, 3/4), (1,)), ((0, 1/2, 0), (-1/4, 1/4, 1/4)), x**(-4))/(16*pi*  
  *(3/2))
```

$$3.242 \quad \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+x^2}} dx$$

Optimal. Leaf size=32

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), -\frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

[Out] 1/2*EllipticF(x,1/2*I*2^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {421, 419}

$$\frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \middle| -\frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 + x^2]*Sqrt[2 + x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -1/2])/(Sqrt[2]*Sqrt[-1 + x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplersqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1+x^2} \sqrt{2+x^2}} dx &= \frac{\sqrt{1-x^2} \int \frac{1}{\sqrt{1-x^2} \sqrt{2+x^2}} dx}{\sqrt{-1+x^2}} \\ &= \frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \middle| -\frac{1}{2}\right)}{\sqrt{2} \sqrt{-1+x^2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), -\frac{1}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 + x^2]*Sqrt[2 + x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -1/2])/(Sqrt[2]*Sqrt[-1 + x^2])

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{x^2+2}\sqrt{x^2-1}}{x^4+x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(x^2 + 2)*sqrt(x^2 - 1)/(x^4 + x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2+2}\sqrt{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 2)*sqrt(x^2 - 1)), x)

maple [A] time = 0.03, size = 34, normalized size = 1.06

$$-\frac{i\sqrt{-x^2+1} \text{EllipticF}\left(\frac{i\sqrt{2}x}{2}, i\sqrt{2}\right)}{\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2-1)^(1/2)/(x^2+2)^(1/2),x)

[Out] -I*EllipticF(1/2*I*2^(1/2)*x,I*2^(1/2))*(-x^2+1)^(1/2)/(x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2+2}\sqrt{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 2)*sqrt(x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(x^2 + 2)^(1/2)),x)

[Out] int(1/((x^2 - 1)^(1/2)*(x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{(x-1)(x+1)}\sqrt{x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2-1)**(1/2)/(x**2+2)**(1/2),x)

[Out] Integral(1/(sqrt((x - 1)*(x + 1))*sqrt(x**2 + 2)), x)

$$3.243 \quad \int \frac{1}{\sqrt{2-x^2} \sqrt{-1+x^2}} dx$$

Optimal. Leaf size=12

$$-\text{EllipticF}\left(\cos^{-1}\left(\frac{x}{\sqrt{2}}\right), 2\right)$$

[Out] $-(x^2)^{(1/2)}/x*\text{EllipticF}(1/2*(-2*x^2+4)^{(1/2)}, 2^{(1/2)})$

Rubi [A] time = 0.01, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {420}

$$-F\left(\cos^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right)$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - x^2]*Sqrt[-1 + x^2]), x]

[Out] -EllipticF[ArcCos[x/Sqrt[2]], 2]

Rule 420

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> -Simp[EllipticF[ArcCos[Rt[-(d/c), 2]*x], (b*c)/(b*c - a*d)]/(Sqrt[c]*Rt[-(d/c), 2]*Sqrt[a - (b*c)/d]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a - (b*c)/d, 0]

Rubi steps

$$\int \frac{1}{\sqrt{2-x^2} \sqrt{-1+x^2}} dx = -F\left(\cos^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right)$$

Mathematica [B] time = 0.02, size = 47, normalized size = 3.92

$$\frac{\sqrt{1-x^2} \sqrt{1-\frac{x^2}{2}} \text{EllipticF}\left(\sin^{-1}(x), \frac{1}{2}\right)}{\sqrt{-x^4+3x^2-2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - x^2]*Sqrt[-1 + x^2]), x]

[Out] (Sqrt[1 - x^2]*Sqrt[1 - x^2/2]*EllipticF[ArcSin[x], 1/2])/Sqrt[-2 + 3*x^2 - x^4]

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2-1} \sqrt{-x^2+2}}{x^4-3x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)^(1/2)/(x^2-1)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(x^2 - 1)*sqrt(-x^2 + 2)/(x^4 - 3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-x^2 + 2)), x)

maple [A] time = 0.03, size = 28, normalized size = 2.33

$$\frac{\sqrt{-x^2+1} \operatorname{EllipticF}\left(\frac{\sqrt{2}x}{2}, \sqrt{2}\right)}{\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+2)^(1/2)/(x^2-1)^(1/2),x)

[Out] EllipticF(1/2*2^(1/2)*x,2^(1/2))*(-x^2+1)^(1/2)/(x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.08

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{2-x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(2 - x^2)^(1/2)),x)

[Out] int(1/((x^2 - 1)^(1/2)*(2 - x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{(x-1)(x+1)}\sqrt{2-x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+2)**(1/2)/(x**2-1)**(1/2),x)

[Out] Integral(1/(sqrt((x - 1)*(x + 1))*sqrt(2 - x**2)), x)

$$3.244 \quad \int \frac{1}{\sqrt{2-2x^2} \sqrt{-1+x^2}} dx$$

Optimal. Leaf size=29

$$-\frac{\sqrt{x^2-1} \tanh^{-1}(x)}{\sqrt{2} \sqrt{1-x^2}}$$

[Out] $-1/2*\operatorname{arctanh}(x)*(x^2-1)^{(1/2)}*2^{(1/2)/(-x^2+1)^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {23, 207}

$$-\frac{\sqrt{x^2-1} \tanh^{-1}(x)}{\sqrt{2} \sqrt{1-x^2}}$$

Antiderivative was successfully verified.

[In] `Int[1/(Sqrt[2 - 2*x^2]*Sqrt[-1 + x^2]),x]`

[Out] `-((Sqrt[-1 + x^2]*ArcTanh[x])/(Sqrt[2]*Sqrt[1 - x^2]))`

Rule 23

`Int[(u_.)*((a_) + (b_.)*(v_))^(m_)*((c_) + (d_.)*(v_))^(n_), x_Symbol] :> Dist[(a + b*v)^m/(c + d*v)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[b*c - a*d, 0] && !(IntegerQ[m] || IntegerQ[n] || GtQ[b/d, 0])`

Rule 207

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> -Simp[ArcTanh[(Rt[b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-2x^2} \sqrt{-1+x^2}} dx &= \frac{\sqrt{-1+x^2} \int \frac{1}{-1+x^2} dx}{\sqrt{2-2x^2}} \\ &= -\frac{\sqrt{-1+x^2} \tanh^{-1}(x)}{\sqrt{2} \sqrt{1-x^2}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 40, normalized size = 1.38

$$\frac{(x^2-1)(\log(1-x) - \log(x+1))}{2\sqrt{2} \sqrt{-(x^2-1)^2}}$$

Antiderivative was successfully verified.

[In] `Integrate[1/(Sqrt[2 - 2*x^2]*Sqrt[-1 + x^2]),x]`

[Out] `((-1 + x^2)*(Log[1 - x] - Log[1 + x]))/(2*Sqrt[2]*Sqrt[-(-1 + x^2)^2])`

fricas [A] time = 0.65, size = 34, normalized size = 1.17

$$\frac{1}{4} \sqrt{2} \arctan\left(\frac{\sqrt{2} \sqrt{x^2-1} \sqrt{-2x^2+2x}}{x^4-1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="fricas")

[Out] 1/4*sqrt(2)*arctan(sqrt(2)*sqrt(x^2 - 1)*sqrt(-2*x^2 + 2)*x/(x^4 - 1))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-2x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-2*x^2 + 2)), x)

maple [A] time = 0.01, size = 24, normalized size = 0.83

$$\frac{\sqrt{2} \sqrt{-x^2+1} \operatorname{arctanh}(x)}{2\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-2*x^2+2)^(1/2)/(x^2-1)^(1/2),x)

[Out] 1/2*2^(1/2)*(-x^2+1)^(1/2)/(x^2-1)^(1/2)*arctanh(x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-2x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-2*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{2-2x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(2 - 2*x^2)^(1/2)),x)

[Out] int(1/((x^2 - 1)^(1/2)*(2 - 2*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\sqrt{2} \int \frac{1}{\sqrt{1-x^2}\sqrt{x^2-1}} dx}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x**2+2)**(1/2)/(x**2-1)**(1/2),x)

[Out] sqrt(2)*Integral(1/(sqrt(1 - x**2)*sqrt(x**2 - 1)), x)/2

$$3.245 \quad \int \frac{1}{\sqrt{2-3x^2} \sqrt{-1+x^2}} dx$$

Optimal. Leaf size=32

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), \frac{3}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

[Out] $1/2 * \operatorname{EllipticF}(x, 1/2 * 6^{(1/2)}) * (-x^2+1)^{(1/2)} * 2^{(1/2)} / (x^2-1)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {421, 419}

$$\frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \middle| \frac{3}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 3*x^2]*Sqrt[-1 + x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], 3/2])/(Sqrt[2]*Sqrt[-1 + x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-3x^2} \sqrt{-1+x^2}} dx &= \frac{\sqrt{1-x^2} \int \frac{1}{\sqrt{2-3x^2} \sqrt{1-x^2}} dx}{\sqrt{-1+x^2}} \\ &= \frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \middle| \frac{3}{2}\right)}{\sqrt{2} \sqrt{-1+x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 40, normalized size = 1.25

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right), \frac{2}{3}\right)}{\sqrt{3} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 3*x^2]*Sqrt[-1 + x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[Sqrt[3/2]*x], 2/3])/(Sqrt[3]*Sqrt[-1 + x^2])

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2-1}\sqrt{-3x^2+2}}{3x^4-5x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(x^2 - 1)*sqrt(-3*x^2 + 2)/(3*x^4 - 5*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-3*x^2 + 2)), x)

maple [A] time = 0.03, size = 29, normalized size = 0.91

$$\frac{\sqrt{-x^2+1}\sqrt{2}\text{EllipticF}\left(x, \frac{\sqrt{6}}{2}\right)}{2\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*x^2+2)^(1/2)/(x^2-1)^(1/2),x)

[Out] 1/2*EllipticF(x,1/2*6^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-3*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{2-3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(2 - 3*x^2)^(1/2)),x)

[Out] int(1/((x^2 - 1)^(1/2)*(2 - 3*x^2)^(1/2)), x)

sympy [C] time = 4.10, size = 37, normalized size = 1.16

$$\begin{cases} \frac{\sqrt{3}iF\left(\text{asin}\left(\frac{\sqrt{6}x}{2}\right)\middle|\frac{2}{3}\right)}{3} & \text{for } x > -\frac{\sqrt{6}}{3} \wedge x < \frac{\sqrt{6}}{3} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x**2+2)**(1/2)/(x**2-1)**(1/2),x)

[Out] Piecewise((-sqrt(3)*I*elliptic_f(asin(sqrt(6)*x/2), 2/3)/3, (x > -sqrt(6)/3) & (x < sqrt(6)/3))

$$3.246 \quad \int \frac{1}{\sqrt{2-4x^2} \sqrt{-1+x^2}} dx$$

Optimal. Leaf size=30

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}(\sin^{-1}(x), 2)}{\sqrt{2} \sqrt{x^2-1}}$$

[Out] 1/2*EllipticF(x, 2^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {421, 419}

$$\frac{\sqrt{1-x^2} F(\sin^{-1}(x)|2)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 4*x^2]*Sqrt[-1 + x^2]), x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], 2])/(Sqrt[2]*Sqrt[-1 + x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplersqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-4x^2} \sqrt{-1+x^2}} dx &= \frac{\sqrt{1-x^2} \int \frac{1}{\sqrt{2-4x^2} \sqrt{1-x^2}} dx}{\sqrt{-1+x^2}} \\ &= \frac{\sqrt{1-x^2} F(\sin^{-1}(x)|2)}{\sqrt{2} \sqrt{-1+x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 36, normalized size = 1.20

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}(\sin^{-1}(\sqrt{2}x), \frac{1}{2})}{2\sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 4*x^2]*Sqrt[-1 + x^2]), x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[Sqrt[2]*x], 1/2])/(2*Sqrt[-1 + x^2])

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{\sqrt{x^2-1} \sqrt{-4x^2+2}}{2(2x^4-3x^2+1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="fricas")
 [Out] integral(-1/2*sqrt(x^2 - 1)*sqrt(-4*x^2 + 2)/(2*x^4 - 3*x^2 + 1), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-4x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="giac")
 [Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-4*x^2 + 2)), x)
maple [A] time = 0.03, size = 27, normalized size = 0.90

$$\frac{\sqrt{-x^2+1}\sqrt{2}\operatorname{EllipticF}(x,\sqrt{2})}{2\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-4*x^2+2)^(1/2)/(x^2-1)^(1/2),x)
 [Out] 1/2*EllipticF(x,2^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)
maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-4x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="maxima")
 [Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-4*x^2 + 2)), x)
mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{2-4x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(2 - 4*x^2)^(1/2)),x)
 [Out] int(1/((x^2 - 1)^(1/2)*(2 - 4*x^2)^(1/2)), x)
sympy [A] time = 6.07, size = 42, normalized size = 1.40

$$\frac{\sqrt{2}\left(\left\{-\frac{\sqrt{2}iF\left(\operatorname{asin}\left(\sqrt{2}x\right)\middle|\frac{1}{2}\right)}{2}\right.\right)}{2}\right)}{2}\quad \text{for } x > -\frac{\sqrt{2}}{2} \wedge x < \frac{\sqrt{2}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x**2+2)**(1/2)/(x**2-1)**(1/2),x)
 [Out] sqrt(2)*Piecewise((-sqrt(2)*I*elliptic_f(asin(sqrt(2)*x), 1/2)/2, (x > -sqrt(2)/2) & (x < sqrt(2)/2)))/2

$$3.247 \quad \int \frac{1}{\sqrt{2-5x^2} \sqrt{-1+x^2}} dx$$

Optimal. Leaf size=32

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), \frac{5}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

[Out] 1/2*EllipticF(x,1/2*10^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {421, 419}

$$\frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \middle| \frac{5}{2}\right)}{\sqrt{2} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 5*x^2]*Sqrt[-1 + x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], 5/2])/(Sqrt[2]*Sqrt[-1 + x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-5x^2} \sqrt{-1+x^2}} dx &= \frac{\sqrt{1-x^2} \int \frac{1}{\sqrt{2-5x^2} \sqrt{1-x^2}} dx}{\sqrt{-1+x^2}} \\ &= \frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \middle| \frac{5}{2}\right)}{\sqrt{2} \sqrt{-1+x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 40, normalized size = 1.25

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right), \frac{2}{5}\right)}{\sqrt{5} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 5*x^2]*Sqrt[-1 + x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[Sqrt[5/2]*x], 2/5])/(Sqrt[5]*Sqrt[-1 + x^2])

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{x^2-1}\sqrt{-5x^2+2}}{5x^4-7x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(x^2 - 1)*sqrt(-5*x^2 + 2)/(5*x^4 - 7*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-5*x^2 + 2)), x)

maple [A] time = 0.03, size = 29, normalized size = 0.91

$$\frac{\sqrt{-x^2+1}\sqrt{2}\text{EllipticF}\left(x, \frac{\sqrt{10}}{2}\right)}{2\sqrt{x^2-1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-5*x^2+2)^(1/2)/(x^2-1)^(1/2),x)

[Out] 1/2*EllipticF(x,1/2*10^(1/2))*(-x^2+1)^(1/2)*2^(1/2)/(x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{-5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(x^2-1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 - 1)*sqrt(-5*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{x^2-1}\sqrt{2-5x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 - 1)^(1/2)*(2 - 5*x^2)^(1/2)),x)

[Out] int(1/((x^2 - 1)^(1/2)*(2 - 5*x^2)^(1/2)), x)

sympy [C] time = 4.15, size = 37, normalized size = 1.16

$$\begin{cases} \frac{\sqrt{5}iF\left(\text{asin}\left(\frac{\sqrt{10}x}{2}\right)\right)\frac{2}{5}}{5} & \text{for } x > -\frac{\sqrt{10}}{5} \wedge x < \frac{\sqrt{10}}{5} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x**2+2)**(1/2)/(x**2-1)**(1/2),x)

[Out] Piecewise((-sqrt(5)*I*elliptic_f(asin(sqrt(10)*x/2), 2/5)/5, (x > -sqrt(10)/5) & (x < sqrt(10)/5)))

$$3.248 \quad \int \frac{1}{\sqrt{-1-x^2} \sqrt{2+5x^2}} dx$$

Optimal. Leaf size=53

$$\frac{\sqrt{5x^2+2} \operatorname{EllipticF}\left(\tan^{-1}(x), -\frac{3}{2}\right)}{\sqrt{2} \sqrt{-x^2-1} \sqrt{\frac{5x^2+2}{x^2+1}}}$$

[Out] $1/2*(1/(x^2+1))^{(1/2)}*(x^2+1)^{(1/2)}*\operatorname{EllipticF}(x/(x^2+1)^{(1/2)}, 1/2*I*6^{(1/2)})*(5*x^2+2)^{(1/2)}*2^{(1/2)}/(-x^2-1)^{(1/2)}/((5*x^2+2)/(x^2+1))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {418}

$$\frac{\sqrt{5x^2+2} F\left(\tan^{-1}(x) \mid -\frac{3}{2}\right)}{\sqrt{2} \sqrt{-x^2-1} \sqrt{\frac{5x^2+2}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 - x^2]*Sqrt[2 + 5*x^2]), x]

[Out] (Sqrt[2 + 5*x^2]*EllipticF[ArcTan[x], -3/2])/(Sqrt[2]*Sqrt[-1 - x^2]*Sqrt[(2 + 5*x^2)/(1 + x^2)])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2+5x^2}} dx = \frac{\sqrt{2+5x^2} F\left(\tan^{-1}(x) \mid -\frac{3}{2}\right)}{\sqrt{2} \sqrt{-1-x^2} \sqrt{\frac{2+5x^2}{1+x^2}}}$$

Mathematica [C] time = 0.03, size = 39, normalized size = 0.74

$$-\frac{i\sqrt{x^2+1} \operatorname{EllipticF}\left(i \sinh^{-1}(x), \frac{5}{2}\right)}{\sqrt{2} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 - x^2]*Sqrt[2 + 5*x^2]), x]

[Out] ((-I)*Sqrt[1 + x^2]*EllipticF[I*ArcSinh[x], 5/2])/(Sqrt[2]*Sqrt[-1 - x^2])

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{\sqrt{5x^2+2} \sqrt{-x^2-1}}{5x^4+7x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="fricas")
 [Out] integral(-sqrt(5*x^2 + 2)*sqrt(-x^2 - 1)/(5*x^4 + 7*x^2 + 2), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{5x^2 + 2}\sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="giac")
 [Out] integrate(1/(sqrt(5*x^2 + 2)*sqrt(-x^2 - 1)), x)
maple [A] time = 0.03, size = 36, normalized size = 0.68

$$\frac{i\sqrt{5}\sqrt{-x^2-1}\operatorname{EllipticF}\left(\frac{i\sqrt{10}x}{2}, \frac{\sqrt{10}}{5}\right)}{5\sqrt{x^2+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2-1)^(1/2)/(5*x^2+2)^(1/2),x)
 [Out] 1/5*I*EllipticF(1/2*I*10^(1/2)*x,1/5*10^(1/2))/(x^2+1)^(1/2)*5^(1/2)*(-x^2-1)^(1/2)
maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{5x^2 + 2}\sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(5*x^2+2)^(1/2),x, algorithm="maxima")
 [Out] integrate(1/(sqrt(5*x^2 + 2)*sqrt(-x^2 - 1)), x)
mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{-x^2 - 1}\sqrt{5x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((-x^2-1)^(1/2)*(5*x^2+2)^(1/2)),x)
 [Out] int(1/((-x^2-1)^(1/2)*(5*x^2+2)^(1/2)),x)
sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2 - 1}\sqrt{5x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2-1)**(1/2)/(5*x**2+2)**(1/2),x)
 [Out] Integral(1/(sqrt(-x**2 - 1)*sqrt(5*x**2 + 2)), x)

$$3.249 \quad \int \frac{1}{\sqrt{-1-x^2} \sqrt{2+4x^2}} dx$$

Optimal. Leaf size=51

$$\frac{\sqrt{2x^2+1} \operatorname{EllipticF}(\tan^{-1}(x), -1)}{\sqrt{2} \sqrt{-x^2-1} \sqrt{\frac{2x^2+1}{x^2+1}}}$$

[Out] $1/2*(1/(x^2+1))^{(1/2)}*(x^2+1)^{(1/2)}*\operatorname{EllipticF}(x/(x^2+1)^{(1/2)}, I)*(2*x^2+1)^{(1/2)}*2^{(1/2)}/(-x^2-1)^{(1/2)}/((2*x^2+1)/(x^2+1))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {418}

$$\frac{\sqrt{2x^2+1} F(\tan^{-1}(x) | -1)}{\sqrt{2} \sqrt{-x^2-1} \sqrt{\frac{2x^2+1}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 - x^2]*Sqrt[2 + 4*x^2]), x]

[Out] (Sqrt[1 + 2*x^2]*EllipticF[ArcTan[x], -1])/(Sqrt[2]*Sqrt[-1 - x^2]*Sqrt[(1 + 2*x^2)/(1 + x^2)])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2+4x^2}} dx = \frac{\sqrt{1+2x^2} F(\tan^{-1}(x) | -1)}{\sqrt{2} \sqrt{-1-x^2} \sqrt{\frac{1+2x^2}{1+x^2}}}$$

Mathematica [C] time = 0.03, size = 37, normalized size = 0.73

$$\frac{i\sqrt{x^2+1} \operatorname{EllipticF}(i \sinh^{-1}(x), 2)}{\sqrt{2} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 - x^2]*Sqrt[2 + 4*x^2]), x]

[Out] ((-I)*Sqrt[1 + x^2]*EllipticF[I*ArcSinh[x], 2])/(Sqrt[2]*Sqrt[-1 - x^2])

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{\sqrt{4x^2+2}\sqrt{-x^2-1}}{2(2x^4+3x^2+1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(4*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(-1/2*sqrt(4*x^2 + 2)*sqrt(-x^2 - 1)/(2*x^4 + 3*x^2 + 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{4x^2 + 2} \sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(4*x^2 + 2)*sqrt(-x^2 - 1)), x)

maple [A] time = 0.03, size = 33, normalized size = 0.65

$$\frac{i\sqrt{-x^2 - 1} \operatorname{EllipticF}\left(i\sqrt{2} x, \frac{\sqrt{2}}{2}\right)}{2\sqrt{x^2 + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2-1)^(1/2)/(4*x^2+2)^(1/2),x)

[Out] 1/2*I*EllipticF(I*2^(1/2)*x,1/2*2^(1/2))/(x^2+1)^(1/2)*(-x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{4x^2 + 2} \sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(4*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(4*x^2 + 2)*sqrt(-x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{-x^2 - 1} \sqrt{4x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((- x^2 - 1)^(1/2)*(4*x^2 + 2)^(1/2)),x)

[Out] int(1/((- x^2 - 1)^(1/2)*(4*x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\sqrt{2} \int \frac{1}{\sqrt{-x^2-1} \sqrt{2x^2+1}} dx}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2-1)**(1/2)/(4*x**2+2)**(1/2),x)

[Out] sqrt(2)*Integral(1/(sqrt(-x**2 - 1)*sqrt(2*x**2 + 1)), x)/2

$$3.250 \quad \int \frac{1}{\sqrt{-1-x^2} \sqrt{2+3x^2}} dx$$

Optimal. Leaf size=53

$$\frac{\sqrt{3x^2+2} \operatorname{EllipticF}\left(\tan^{-1}(x), -\frac{1}{2}\right)}{\sqrt{2} \sqrt{-x^2-1} \sqrt{\frac{3x^2+2}{x^2+1}}}$$

[Out] $1/2*(1/(x^2+1))^{(1/2)}*(x^2+1)^{(1/2)}*\operatorname{EllipticF}(x/(x^2+1)^{(1/2)}, 1/2*I*2^{(1/2)})*(3*x^2+2)^{(1/2)}*2^{(1/2)}/(-x^2-1)^{(1/2)}/((3*x^2+2)/(x^2+1))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {418}

$$\frac{\sqrt{3x^2+2} F\left(\tan^{-1}(x) \mid -\frac{1}{2}\right)}{\sqrt{2} \sqrt{-x^2-1} \sqrt{\frac{3x^2+2}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 - x^2]*Sqrt[2 + 3*x^2]), x]

[Out] (Sqrt[2 + 3*x^2]*EllipticF[ArcTan[x], -1/2])/(Sqrt[2]*Sqrt[-1 - x^2]*Sqrt[(2 + 3*x^2)/(1 + x^2)])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2+3x^2}} dx = \frac{\sqrt{2+3x^2} F\left(\tan^{-1}(x) \mid -\frac{1}{2}\right)}{\sqrt{2} \sqrt{-1-x^2} \sqrt{\frac{2+3x^2}{1+x^2}}}$$

Mathematica [C] time = 0.02, size = 39, normalized size = 0.74

$$-\frac{i\sqrt{x^2+1} \operatorname{EllipticF}\left(i \sinh^{-1}(x), \frac{3}{2}\right)}{\sqrt{2} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 - x^2]*Sqrt[2 + 3*x^2]), x]

[Out] ((-I)*Sqrt[1 + x^2]*EllipticF[I*ArcSinh[x], 3/2])/(Sqrt[2]*Sqrt[-1 - x^2])

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{\sqrt{3x^2+2} \sqrt{-x^2-1}}{3x^4+5x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="fricas")
 [Out] integral(-sqrt(3*x^2 + 2)*sqrt(-x^2 - 1)/(3*x^4 + 5*x^2 + 2), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{3x^2 + 2}\sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="giac")
 [Out] integrate(1/(sqrt(3*x^2 + 2)*sqrt(-x^2 - 1)), x)
maple [A] time = 0.03, size = 36, normalized size = 0.68

$$\frac{i\sqrt{3}\sqrt{-x^2-1}\operatorname{EllipticF}\left(\frac{i\sqrt{6}x}{2}, \frac{\sqrt{6}}{3}\right)}{3\sqrt{x^2+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2-1)^(1/2)/(3*x^2+2)^(1/2),x)
 [Out] 1/3*I*EllipticF(1/2*I*x*6^(1/2),1/3*6^(1/2))/(x^2+1)^(1/2)*3^(1/2)*(-x^2-1)^(1/2)
maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{3x^2 + 2}\sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(3*x^2+2)^(1/2),x, algorithm="maxima")
 [Out] integrate(1/(sqrt(3*x^2 + 2)*sqrt(-x^2 - 1)), x)
mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{-x^2 - 1}\sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((-x^2-1)^(1/2)*(3*x^2+2)^(1/2)),x)
 [Out] int(1/((-x^2-1)^(1/2)*(3*x^2+2)^(1/2)),x)
sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2 - 1}\sqrt{3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2-1)**(1/2)/(3*x**2+2)**(1/2),x)
 [Out] Integral(1/(sqrt(-x**2 - 1)*sqrt(3*x**2 + 2)), x)

$$3.251 \quad \int \frac{1}{\sqrt{-1-x^2} \sqrt{2+2x^2}} dx$$

Optimal. Leaf size=28

$$\frac{\sqrt{x^2+1} \tan^{-1}(x)}{\sqrt{2} \sqrt{-x^2-1}}$$

[Out] 1/2*arctan(x)*(x^2+1)^(1/2)*2^(1/2)/(-x^2-1)^(1/2)

Rubi [A] time = 0.00, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {23, 203}

$$\frac{\sqrt{x^2+1} \tan^{-1}(x)}{\sqrt{2} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 - x^2]*Sqrt[2 + 2*x^2]),x]

[Out] (Sqrt[1 + x^2]*ArcTan[x])/(Sqrt[2]*Sqrt[-1 - x^2])

Rule 23

Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :> Dist[(a + b*v)^m/(c + d*v)^n, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[b*c - a*d, 0] && !(IntegerQ[m] || IntegerQ[n] || GtQ[b/d, 0])

Rule 203

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1-x^2} \sqrt{2+2x^2}} dx &= \frac{\sqrt{2+2x^2} \int \frac{1}{2+2x^2} dx}{\sqrt{-1-x^2}} \\ &= \frac{\sqrt{1+x^2} \tan^{-1}(x)}{\sqrt{2} \sqrt{-1-x^2}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 26, normalized size = 0.93

$$\frac{(x^2+1) \tan^{-1}(x)}{\sqrt{2} \sqrt{-(x^2+1)^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 - x^2]*Sqrt[2 + 2*x^2]),x]

[Out] ((1 + x^2)*ArcTan[x])/(Sqrt[2]*Sqrt[-(1 + x^2)^2])

fricas [B] time = 0.65, size = 104, normalized size = 3.71

$$\frac{1}{8} \sqrt{2} \log \left(\frac{2 \left(2 \sqrt{2x^2+2} \sqrt{-x^2-1} x + \sqrt{2} (x^4-1) \right)}{x^4+2x^2+1} \right) - \frac{1}{8} \sqrt{2} \log \left(\frac{2 \left(2 \sqrt{2x^2+2} \sqrt{-x^2-1} x - \sqrt{2} (x^4-1) \right)}{x^4+2x^2+1} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="fricas")

[Out] 1/8*sqrt(2)*log(2*(2*sqrt(2*x^2 + 2)*sqrt(-x^2 - 1)*x + sqrt(2)*(x^4 - 1))/(x^4 + 2*x^2 + 1)) - 1/8*sqrt(2)*log(2*(2*sqrt(2*x^2 + 2)*sqrt(-x^2 - 1)*x - sqrt(2)*(x^4 - 1))/(x^4 + 2*x^2 + 1))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2 + 2}\sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(2*x^2 + 2)*sqrt(-x^2 - 1)), x)

maple [A] time = 0.01, size = 24, normalized size = 0.86

$$\frac{\sqrt{-x^2 - 1} \sqrt{2} \arctan(x)}{2\sqrt{x^2 + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2-1)^(1/2)/(2*x^2+2)^(1/2),x)

[Out] -1/2*(-x^2-1)^(1/2)*2^(1/2)/(x^2+1)^(1/2)*arctan(x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2 + 2}\sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(2*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(2*x^2 + 2)*sqrt(-x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.04

$$\int \frac{1}{\sqrt{-x^2 - 1}\sqrt{2x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((-x^2 - 1)^(1/2)*(2*x^2 + 2)^(1/2)),x)

[Out] int(1/((-x^2 - 1)^(1/2)*(2*x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\sqrt{2} \int \frac{1}{\sqrt{-x^2-1}\sqrt{x^2+1}} dx}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2-1)**(1/2)/(2*x**2+2)**(1/2),x)

[Out] sqrt(2)*Integral(1/(sqrt(-x**2 - 1)*sqrt(x**2 + 1)), x)/2

$$3.252 \quad \int \frac{1}{\sqrt{-1-x^2} \sqrt{2+x^2}} dx$$

Optimal. Leaf size=49

$$\frac{\sqrt{x^2+2} \operatorname{EllipticF}\left(\tan^{-1}(x), \frac{1}{2}\right)}{\sqrt{2} \sqrt{-x^2-1} \sqrt{\frac{x^2+2}{x^2+1}}}$$

[Out] $1/2*(1/(x^2+1))^{(1/2)}*(x^2+1)^{(1/2)}*\operatorname{EllipticF}(x/(x^2+1)^{(1/2)}, 1/2*2^{(1/2)})*(x^2+2)^{(1/2)}*2^{(1/2)/(-x^2-1)^{(1/2)/((x^2+2)/(x^2+1))^{(1/2)}}$

Rubi [A] time = 0.01, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {418}

$$\frac{\sqrt{x^2+2} F\left(\tan^{-1}(x)\middle|\frac{1}{2}\right)}{\sqrt{2} \sqrt{-x^2-1} \sqrt{\frac{x^2+2}{x^2+1}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 - x^2]*Sqrt[2 + x^2]), x]

[Out] (Sqrt[2 + x^2]*EllipticF[ArcTan[x], 1/2])/(Sqrt[2]*Sqrt[-1 - x^2]*Sqrt[(2 + x^2)/(1 + x^2)])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{-1-x^2} \sqrt{2+x^2}} dx = \frac{\sqrt{2+x^2} F\left(\tan^{-1}(x)\middle|\frac{1}{2}\right)}{\sqrt{2} \sqrt{-1-x^2} \sqrt{\frac{2+x^2}{1+x^2}}}$$

Mathematica [C] time = 0.04, size = 53, normalized size = 1.08

$$\frac{i\sqrt{x^2+1} \sqrt{x^2+2} \operatorname{EllipticF}\left(i \sinh^{-1}(x), \frac{1}{2}\right)}{\sqrt{2} \sqrt{-((x^2+1)(x^2+2))}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 - x^2]*Sqrt[2 + x^2]), x]

[Out] ((-I)*Sqrt[1 + x^2]*Sqrt[2 + x^2]*EllipticF[I*ArcSinh[x], 1/2])/(Sqrt[2]*Sqrt[-((1 + x^2)*(2 + x^2))])

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{\sqrt{x^2+2} \sqrt{-x^2-1}}{x^4+3x^2+2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(x^2 + 2)*sqrt(-x^2 - 1)/(x^4 + 3*x^2 + 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 2} \sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 + 2)*sqrt(-x^2 - 1)), x)

maple [C] time = 0.03, size = 33, normalized size = 0.67

$$\frac{i\sqrt{2} \sqrt{-x^2 - 1} \operatorname{EllipticF}\left(ix, \frac{\sqrt{2}}{2}\right)}{2\sqrt{x^2 + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2-1)^(1/2)/(x^2+2)^(1/2),x)

[Out] 1/2*I*EllipticF(I*x,1/2*2^(1/2))*2^(1/2)/(x^2+1)^(1/2)*(-x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 + 2} \sqrt{-x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 + 2)*sqrt(-x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{-x^2 - 1} \sqrt{x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((- x^2 - 1)^(1/2)*(x^2 + 2)^(1/2)),x)

[Out] int(1/((- x^2 - 1)^(1/2)*(x^2 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2 - 1} \sqrt{x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2-1)**(1/2)/(x**2+2)**(1/2),x)

[Out] Integral(1/(sqrt(-x**2 - 1)*sqrt(x**2 + 2)), x)

$$3.253 \quad \int \frac{1}{\sqrt{-1-x^2} \sqrt{2-x^2}} dx$$

Optimal. Leaf size=31

$$\frac{\sqrt{x^2+1} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{x}{\sqrt{2}}\right), -2\right)}{\sqrt{-x^2-1}}$$

[Out] EllipticF(1/2*x*2^(1/2), I*2^(1/2))*(x^2+1)^(1/2)/(-x^2-1)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {421, 419}

$$\frac{\sqrt{x^2+1} F\left(\sin^{-1}\left(\frac{x}{\sqrt{2}}\right) \middle| -2\right)}{\sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 - x^2]*Sqrt[2 - x^2]), x]

[Out] (Sqrt[1 + x^2]*EllipticF[ArcSin[x/Sqrt[2]], -2])/Sqrt[-1 - x^2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1-x^2} \sqrt{2-x^2}} dx &= \frac{\sqrt{1+x^2} \int \frac{1}{\sqrt{2-x^2} \sqrt{1+x^2}} dx}{\sqrt{-1-x^2}} \\ &= \frac{\sqrt{1+x^2} F\left(\sin^{-1}\left(\frac{x}{\sqrt{2}}\right) \middle| -2\right)}{\sqrt{-1-x^2}} \end{aligned}$$

Mathematica [C] time = 0.03, size = 39, normalized size = 1.26

$$\frac{i\sqrt{x^2+1} \operatorname{EllipticF}\left(i \sinh^{-1}(x), -\frac{1}{2}\right)}{\sqrt{2} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 - x^2]*Sqrt[2 - x^2]), x]

[Out] ((-I)*Sqrt[1 + x^2]*EllipticF[I*ArcSinh[x], -1/2])/(Sqrt[2]*Sqrt[-1 - x^2])

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2+2}\sqrt{-x^2-1}}{x^4-x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(-x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-x^2+2)*sqrt(-x^2-1)/(x^4-x^2-2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+2}\sqrt{-x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(-x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2+2)*sqrt(-x^2-1)), x)

maple [A] time = 0.03, size = 34, normalized size = 1.10

$$\frac{i\sqrt{2}\sqrt{-x^2-1}\text{EllipticF}\left(ix, \frac{i\sqrt{2}}{2}\right)}{2\sqrt{x^2+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2-1)^(1/2)/(-x^2+2)^(1/2),x)

[Out] 1/2*I*EllipticF(I*x,1/2*I*2^(1/2))/(x^2+1)^(1/2)*2^(1/2)*(-x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2+2}\sqrt{-x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2-1)^(1/2)/(-x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2+2)*sqrt(-x^2-1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{2-x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((-x^2-1)^(1/2)*(2-x^2)^(1/2)),x)

[Out] int(1/((-x^2-1)^(1/2)*(2-x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2-x^2}\sqrt{-x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2-1)**(1/2)/(-x**2+2)**(1/2),x)

[Out] Integral(1/(sqrt(2-x**2)*sqrt(-x**2-1)), x)

$$3.254 \quad \int \frac{1}{\sqrt{2-2x^2} \sqrt{-1-x^2}} dx$$

Optimal. Leaf size=42

$$\frac{\sqrt{1-\frac{1}{x^4}} x^2 \operatorname{EllipticF}\left(\operatorname{csc}^{-1}(x), -1\right)}{\sqrt{2-2x^2} \sqrt{-x^2-1}}$$

[Out] $-x^2 \operatorname{EllipticF}(1/x, 1) (1-1/x^4)^{1/2} / (-2x^2+2)^{1/2} / (-x^2-1)^{1/2}$

Rubi [A] time = 0.01, antiderivative size = 65, normalized size of antiderivative = 1.55, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {253, 222}

$$\frac{\sqrt{x^2-1} \sqrt{x^2+1} F\left(\sin^{-1}\left(\frac{\sqrt{2}x}{\sqrt{x^2-1}}\right) \middle| \frac{1}{2}\right)}{2\sqrt{-x^2-1} \sqrt{1-x^2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 2*x^2]*Sqrt[-1 - x^2]),x]

[Out] (Sqrt[-1 + x^2]*Sqrt[1 + x^2]*EllipticF[ArcSin[(Sqrt[2]*x)/Sqrt[-1 + x^2]], 1/2])/(2*Sqrt[-1 - x^2]*Sqrt[1 - x^2])

Rule 222

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[-(a*b), 2]}, Simp[(Sqrt[-a + q*x^2]*Sqrt[(a + q*x^2)/q]*EllipticF[ArcSin[x/Sqrt[(a + q*x^2)/(2*q)]], 1/2])/(Sqrt[2]*Sqrt[-a]*Sqrt[a + b*x^4]), x] /; IntegerQ[q] /; FreeQ[{a, b}, x] && LtQ[a, 0] && GtQ[b, 0]

Rule 253

Int[((a1_.) + (b1_.)*(x_)^(n_))^(p_)*((a2_.) + (b2_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[((a1 + b1*x^n)^FracPart[p]*(a2 + b2*x^n)^FracPart[p])/(a1*a2 + b1*b2*x^(2*n))^FracPart[p], Int[(a1*a2 + b1*b2*x^(2*n))^p, x], x] /; FreeQ[{a1, b1, a2, b2, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-2x^2} \sqrt{-1-x^2}} dx &= \frac{\sqrt{-2+2x^4} \int \frac{1}{\sqrt{-2+2x^4}} dx}{\sqrt{2-2x^2} \sqrt{-1-x^2}} \\ &= \frac{\sqrt{-1+x^2} \sqrt{1+x^2} F\left(\sin^{-1}\left(\frac{\sqrt{2}x}{\sqrt{-1+x^2}}\right) \middle| \frac{1}{2}\right)}{2\sqrt{-1-x^2} \sqrt{1-x^2}} \end{aligned}$$

Mathematica [C] time = 0.02, size = 48, normalized size = 1.14

$$\frac{x\sqrt{1-x^4} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; x^4\right)}{\sqrt{2-2x^2} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 2*x^2]*Sqrt[-1 - x^2]),x]

[Out] (x*Sqrt[1 - x^4]*Hypergeometric2F1[1/4, 1/2, 5/4, x^4])/(Sqrt[2 - 2*x^2]*Sqrt[-1 - x^2])

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2-1}\sqrt{-2x^2+2}}{2(x^4-1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(-x^2-1)^(1/2), x, algorithm="fricas")

[Out] integral(1/2*sqrt(-x^2 - 1)*sqrt(-2*x^2 + 2)/(x^4 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{-2x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(-x^2-1)^(1/2), x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2 - 1)*sqrt(-2*x^2 + 2)), x)

maple [A] time = 0.02, size = 30, normalized size = 0.71

$$\frac{i\sqrt{2}\sqrt{-x^2-1}\text{EllipticF}(ix, i)}{2\sqrt{x^2+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-2*x^2+2)^(1/2)/(-x^2-1)^(1/2), x)

[Out] 1/2*I*EllipticF(I*x, I)*2^(1/2)/(x^2+1)^(1/2)*(-x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{-2x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-2*x^2+2)^(1/2)/(-x^2-1)^(1/2), x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 - 1)*sqrt(-2*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{2-2x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((-x^2-1)^(1/2)*(2-2*x^2)^(1/2)), x)

[Out] int(1/((-x^2-1)^(1/2)*(2-2*x^2)^(1/2)), x)

sympy [A] time = 5.45, size = 73, normalized size = 1.74

$$\frac{\sqrt{2}G_{6,6}^{5,3}\left(\begin{matrix} \frac{1}{2}, 1, 1 & \frac{3}{4}, \frac{3}{4}, \frac{5}{4} \\ \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1, \frac{5}{4} & 0 \end{matrix} \middle| \frac{1}{x^4}\right)}{16\pi^{\frac{3}{2}}} - \frac{\sqrt{2}G_{6,6}^{3,5}\left(\begin{matrix} -\frac{1}{4}, 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4} & 1 \\ 0, \frac{1}{2}, 0 & -\frac{1}{4}, \frac{1}{4}, \frac{1}{4} \end{matrix} \middle| \frac{e^{-2i\pi}}{x^4}\right)}{16\pi^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(-2*x**2+2)**(1/2)/(-x**2-1)**(1/2),x)
```

```
[Out] sqrt(2)*meijerg(((1/2, 1, 1), (3/4, 3/4, 5/4)), ((1/4, 1/2, 3/4, 1, 5/4), (0,)), x**(-4))/(16*pi**(3/2)) - sqrt(2)*meijerg((-1/4, 0, 1/4, 1/2, 3/4), (1,)), ((0, 1/2, 0), (-1/4, 1/4, 1/4)), exp_polar(-2*I*pi)/x**4)/(16*pi**(3/2))
```

$$3.255 \quad \int \frac{1}{\sqrt{2-3x^2} \sqrt{-1-x^2}} dx$$

Optimal. Leaf size=40

$$\frac{\sqrt{x^2+1} \operatorname{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right), -\frac{2}{3}\right)}{\sqrt{3} \sqrt{-x^2-1}}$$

[Out] 1/3*EllipticF(1/2*x*6^(1/2), 1/3*I*6^(1/2))*(x^2+1)^(1/2)*3^(1/2)/(-x^2-1)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {421, 419}

$$\frac{\sqrt{x^2+1} F\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right) \middle| -\frac{2}{3}\right)}{\sqrt{3} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 3*x^2]*Sqrt[-1 - x^2]), x]

[Out] (Sqrt[1 + x^2]*EllipticF[ArcSin[Sqrt[3/2]*x], -2/3])/(Sqrt[3]*Sqrt[-1 - x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-3x^2} \sqrt{-1-x^2}} dx &= \frac{\sqrt{1+x^2} \int \frac{1}{\sqrt{2-3x^2} \sqrt{1+x^2}} dx}{\sqrt{-1-x^2}} \\ &= \frac{\sqrt{1+x^2} F\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right) \middle| -\frac{2}{3}\right)}{\sqrt{3} \sqrt{-1-x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 40, normalized size = 1.00

$$\frac{\sqrt{x^2+1} \operatorname{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right), -\frac{2}{3}\right)}{\sqrt{3} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 3*x^2]*Sqrt[-1 - x^2]), x]

[Out] (Sqrt[1 + x^2]*EllipticF[ArcSin[Sqrt[3/2]*x], -2/3])/(Sqrt[3]*Sqrt[-1 - x^2])

fricas [F] time = 0.86, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2-1}\sqrt{-3x^2+2}}{3x^4+x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(-x^2-1)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-x^2 - 1)*sqrt(-3*x^2 + 2)/(3*x^4 + x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{-3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(-x^2-1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2 - 1)*sqrt(-3*x^2 + 2)), x)

maple [A] time = 0.03, size = 34, normalized size = 0.85

$$\frac{i\sqrt{-x^2-1}\sqrt{2}\text{EllipticF}\left(ix, \frac{i\sqrt{6}}{2}\right)}{2\sqrt{x^2+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*x^2+2)^(1/2)/(-x^2-1)^(1/2),x)

[Out] 1/2*I*EllipticF(I*x,1/2*I*6^(1/2))/(x^2+1)^(1/2)*(-x^2-1)^(1/2)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{-3x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/2)/(-x^2-1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 - 1)*sqrt(-3*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{2-3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((-x^2 - 1)^(1/2)*(2 - 3*x^2)^(1/2)),x)

[Out] int(1/((-x^2 - 1)^(1/2)*(2 - 3*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2-3x^2}\sqrt{-x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x**2+2)**(1/2)/(-x**2-1)**(1/2),x)

[Out] Integral(1/(sqrt(2 - 3*x**2)*sqrt(-x**2 - 1)), x)

$$3.256 \quad \int \frac{1}{\sqrt{2-4x^2} \sqrt{-1-x^2}} dx$$

Optimal. Leaf size=36

$$\frac{\sqrt{x^2+1} \operatorname{EllipticF}\left(\sin^{-1}(\sqrt{2}x), -\frac{1}{2}\right)}{2\sqrt{-x^2-1}}$$

[Out] 1/2*EllipticF(x*2^(1/2),1/2*I*2^(1/2))*(x^2+1)^(1/2)/(-x^2-1)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {421, 419}

$$\frac{\sqrt{x^2+1} F\left(\sin^{-1}(\sqrt{2}x) \mid -\frac{1}{2}\right)}{2\sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 4*x^2]*Sqrt[-1 - x^2]),x]

[Out] (Sqrt[1 + x^2]*EllipticF[ArcSin[Sqrt[2]*x], -1/2])/(2*Sqrt[-1 - x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-4x^2} \sqrt{-1-x^2}} dx &= \frac{\sqrt{1+x^2} \int \frac{1}{\sqrt{2-4x^2} \sqrt{1+x^2}} dx}{\sqrt{-1-x^2}} \\ &= \frac{\sqrt{1+x^2} F\left(\sin^{-1}(\sqrt{2}x) \mid -\frac{1}{2}\right)}{2\sqrt{-1-x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 36, normalized size = 1.00

$$\frac{\sqrt{x^2+1} \operatorname{EllipticF}\left(\sin^{-1}(\sqrt{2}x), -\frac{1}{2}\right)}{2\sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 4*x^2]*Sqrt[-1 - x^2]),x]

[Out] (Sqrt[1 + x^2]*EllipticF[ArcSin[Sqrt[2]*x], -1/2])/(2*Sqrt[-1 - x^2])

fricas [F] time = 0.85, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2-1}\sqrt{-4x^2+2}}{2(2x^4+x^2-1)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(-x^2-1)^(1/2), x, algorithm="fricas")

[Out] integral(1/2*sqrt(-x^2 - 1)*sqrt(-4*x^2 + 2)/(2*x^4 + x^2 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{-4x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(-x^2-1)^(1/2), x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2 - 1)*sqrt(-4*x^2 + 2)), x)

maple [A] time = 0.03, size = 34, normalized size = 0.94

$$\frac{i\sqrt{2}\sqrt{-x^2-1}\text{EllipticF}(ix, i\sqrt{2})}{2\sqrt{x^2+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-4*x^2+2)^(1/2)/(-x^2-1)^(1/2), x)

[Out] 1/2*I*EllipticF(I*x, I*2^(1/2))*2^(1/2)/(x^2+1)^(1/2)*(-x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{-4x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x^2+2)^(1/2)/(-x^2-1)^(1/2), x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 - 1)*sqrt(-4*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{2-4x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((-x^2-1)^(1/2)*(2-4*x^2)^(1/2)), x)

[Out] int(1/((-x^2-1)^(1/2)*(2-4*x^2)^(1/2)), x)

sympy [A] time = 5.78, size = 44, normalized size = 1.22

$$\frac{\sqrt{2}\left(\left\{-\frac{\sqrt{2}iF\left(\text{asin}\left(\sqrt{2}x\right)\middle|-\frac{1}{2}\right)}{2}\right.\right)}{2}\text{ for } x > -\frac{\sqrt{2}}{2} \wedge x < \frac{\sqrt{2}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-4*x**2+2)**(1/2)/(-x**2-1)**(1/2), x)

[Out] sqrt(2)*Piecewise((-sqrt(2)*I*elliptic_f(asin(sqrt(2)*x), -1/2)/2, (x > -sqrt(2)/2) & (x < sqrt(2)/2)))/2

$$3.257 \quad \int \frac{1}{\sqrt{2-5x^2} \sqrt{-1-x^2}} dx$$

Optimal. Leaf size=40

$$\frac{\sqrt{x^2+1} \operatorname{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right), -\frac{2}{5}\right)}{\sqrt{5} \sqrt{-x^2-1}}$$

[Out] 1/5*EllipticF(1/2*x*10^(1/2),1/5*I*10^(1/2))*(x^2+1)^(1/2)*5^(1/2)/(-x^2-1)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {421, 419}

$$\frac{\sqrt{x^2+1} F\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right) \middle| -\frac{2}{5}\right)}{\sqrt{5} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 - 5*x^2]*Sqrt[-1 - x^2]),x]

[Out] (Sqrt[1 + x^2]*EllipticF[ArcSin[Sqrt[5/2]*x], -2/5])/(Sqrt[5]*Sqrt[-1 - x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{2-5x^2} \sqrt{-1-x^2}} dx &= \frac{\sqrt{1+x^2} \int \frac{1}{\sqrt{2-5x^2} \sqrt{1+x^2}} dx}{\sqrt{-1-x^2}} \\ &= \frac{\sqrt{1+x^2} F\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right) \middle| -\frac{2}{5}\right)}{\sqrt{5} \sqrt{-1-x^2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 40, normalized size = 1.00

$$\frac{\sqrt{x^2+1} \operatorname{EllipticF}\left(\sin^{-1}\left(\sqrt{\frac{5}{2}}x\right), -\frac{2}{5}\right)}{\sqrt{5} \sqrt{-x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 - 5*x^2]*Sqrt[-1 - x^2]),x]

[Out] (Sqrt[1 + x^2]*EllipticF[ArcSin[Sqrt[5/2]*x], -2/5])/(Sqrt[5]*Sqrt[-1 - x^2])

fricas [F] time = 0.91, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-x^2-1}\sqrt{-5x^2+2}}{5x^4+3x^2-2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(-x^2-1)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(-x^2 - 1)*sqrt(-5*x^2 + 2)/(5*x^4 + 3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{-5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(-x^2-1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(-x^2 - 1)*sqrt(-5*x^2 + 2)), x)

maple [A] time = 0.03, size = 34, normalized size = 0.85

$$\frac{i\sqrt{2}\sqrt{-x^2-1}\text{EllipticF}\left(ix, \frac{i\sqrt{10}}{2}\right)}{2\sqrt{x^2+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-5*x^2+2)^(1/2)/(-x^2-1)^(1/2),x)

[Out] 1/2*I*EllipticF(I*x, 1/2*I*10^(1/2))*2^(1/2)/(x^2+1)^(1/2)*(-x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{-5x^2+2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x^2+2)^(1/2)/(-x^2-1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(-x^2 - 1)*sqrt(-5*x^2 + 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{-x^2-1}\sqrt{2-5x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((-x^2 - 1)^(1/2)*(2 - 5*x^2)^(1/2)),x)

[Out] int(1/((-x^2 - 1)^(1/2)*(2 - 5*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2-5x^2}\sqrt{-x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-5*x**2+2)**(1/2)/(-x**2-1)**(1/2),x)

[Out] Integral(1/(sqrt(2 - 5*x**2)*sqrt(-x**2 - 1)), x)

$$3.258 \quad \int \frac{\sqrt{a+bx^2}}{\sqrt{c-dx^2}} dx$$

Optimal. Leaf size=87

$$\frac{\sqrt{c} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{\frac{bx^2}{a}+1} \sqrt{c-dx^2}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(b*x^2+a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(1+b*x^2/a)^(1/2)/(-d*x^2+c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {427, 426, 424}

$$\frac{\sqrt{c} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{\frac{bx^2}{a}+1} \sqrt{c-dx^2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]/Sqrt[c - d*x^2], x]

[Out] (Sqrt[c]*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d)))/(Sqrt[d]*Sqrt[1 + (b*x^2)/a]*Sqrt[c - d*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{a+bx^2}}{\sqrt{c-dx^2}} dx &= \frac{\sqrt{1-\frac{dx^2}{c}} \int \frac{\sqrt{a+bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{c-dx^2}} \\
&= \frac{\left(\sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1+\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{1+\frac{bx^2}{a}} \sqrt{c-dx^2}} \\
&= \frac{\sqrt{c} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1+\frac{bx^2}{a}} \sqrt{c-dx^2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 87, normalized size = 1.00

$$\frac{\sqrt{a+bx^2} \sqrt{\frac{c-dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a+bx^2}{a}} \sqrt{c-dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2]/Sqrt[c - d*x^2], x]

[Out] (Sqrt[a + b*x^2]*Sqrt[(c - d*x^2)/c]*EllipticE[ArcSin[Sqrt[d/c]*x], -((b*c)/(a*d)))]/(Sqrt[d/c]*Sqrt[(a + b*x^2)/a]*Sqrt[c - d*x^2])

fricas [F] time = 0.84, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{bx^2+a}\sqrt{-dx^2+c}}{dx^2-c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(b*x^2 + a)*sqrt(-d*x^2 + c)/(d*x^2 - c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{-dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 + a)/sqrt(-d*x^2 + c), x)

maple [A] time = 0.02, size = 106, normalized size = 1.22

$$\frac{\sqrt{bx^2+a} \sqrt{-dx^2+c} \sqrt{-\frac{dx^2-c}{c}} \sqrt{\frac{bx^2+a}{a}} a \text{EllipticE}\left(\sqrt{\frac{d}{c}}x, \sqrt{-\frac{bc}{ad}}\right)}{(-bdx^4 - adx^2 + bcx^2 + ac) \sqrt{\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x)`

[Out] `(b*x^2+a)^(1/2)*(-d*x^2+c)^(1/2)*a*(-(d*x^2-c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)*EllipticE((1/c*d)^(1/2)*x,(-1/a*b*c/d)^(1/2))/(-b*d*x^4-a*d*x^2+b*c*x^2+a*c)/(1/c*d)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + a}}{\sqrt{-dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*x^2 + a)/sqrt(-d*x^2 + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 + a}}{\sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^(1/2)/(c - d*x^2)^(1/2),x)`

[Out] `int((a + b*x^2)^(1/2)/(c - d*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a + bx^2}}{\sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2),x)`

[Out] `Integral(sqrt(a + b*x**2)/sqrt(c - d*x**2), x)`

$$3.259 \quad \int \frac{\sqrt{-a-bx^2}}{\sqrt{c-dx^2}} dx$$

Optimal. Leaf size=90

$$\frac{\sqrt{c} \sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{\frac{bx^2}{a}+1} \sqrt{c-dx^2}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(-b*x^2-a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(1+b*x^2/a)^(1/2)/(-d*x^2+c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {427, 426, 424}

$$\frac{\sqrt{c} \sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{\frac{bx^2}{a}+1} \sqrt{c-dx^2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-a - b*x^2]/Sqrt[c - d*x^2], x]

[Out] (Sqrt[c]*Sqrt[-a - b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d)))/(Sqrt[d]*Sqrt[1 + (b*x^2)/a]*Sqrt[c - d*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\int \frac{\sqrt{-a-bx^2}}{\sqrt{c-dx^2}} dx = \frac{\sqrt{1-\frac{dx^2}{c}} \int \frac{\sqrt{-a-bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{c-dx^2}}$$

$$= \frac{\left(\sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1+\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{1+\frac{bx^2}{a}} \sqrt{c-dx^2}}$$

$$= \frac{\sqrt{c} \sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1+\frac{bx^2}{a}} \sqrt{c-dx^2}}$$

Mathematica [A] time = 0.05, size = 90, normalized size = 1.00

$$\frac{\sqrt{-a-bx^2} \sqrt{\frac{c-dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a+bx^2}{a}} \sqrt{c-dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-a - b*x^2]/Sqrt[c - d*x^2], x]

[Out] (Sqrt[-a - b*x^2]*Sqrt[(c - d*x^2)/c]*EllipticE[ArcSin[Sqrt[d/c]*x], -(b*c)/(a*d)))/(Sqrt[d/c]*Sqrt[(a + b*x^2)/a]*Sqrt[c - d*x^2])

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2-a}\sqrt{-dx^2+c}}{dx^2-c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(-b*x^2 - a)*sqrt(-d*x^2 + c)/(d*x^2 - c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2-a}}{\sqrt{-dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-b*x^2 - a)/sqrt(-d*x^2 + c), x)

maple [B] time = 0.03, size = 171, normalized size = 1.90

$$\frac{\left(-ad \text{EllipticF}\left(\sqrt{-\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right) + bc \text{EllipticE}\left(\sqrt{-\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right) - bc \text{EllipticF}\left(\sqrt{-\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right)\right) \sqrt{-bx^2-a} \sqrt{-d}}{(bdx^4 + adx^2 - bcx^2 - ac) \sqrt{-\frac{b}{a}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2),x)`

[Out] `(-a*EllipticF((-1/a*b)^(1/2)*x,(-a/b/c*d)^(1/2))*d-b*c*EllipticF((-1/a*b)^(1/2)*x,(-a/b/c*d)^(1/2))+b*c*EllipticE((-1/a*b)^(1/2)*x,(-a/b/c*d)^(1/2)))*(-b*x^2-a)^(1/2)*(-d*x^2+c)^(1/2)*((b*x^2+a)/a)^(1/2)*(-(d*x^2-c)/c)^(1/2)/(b*d*x^4+a*d*x^2-b*c*x^2-a*c)/(-1/a*b)^(1/2)/d`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{-dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-b*x^2 - a)/sqrt(-d*x^2 + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((- a - b*x^2)^(1/2)/(c - d*x^2)^(1/2),x)`

[Out] `int((- a - b*x^2)^(1/2)/(c - d*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-a - bx^2}}{\sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x**2-a)**(1/2)/(-d*x**2+c)**(1/2),x)`

[Out] `Integral(sqrt(-a - b*x**2)/sqrt(c - d*x**2), x)`

$$3.260 \quad \int \frac{\sqrt{a+bx^2}}{\sqrt{-c+dx^2}} dx$$

Optimal. Leaf size=88

$$\frac{\sqrt{c} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{\frac{bx^2}{a}+1} \sqrt{dx^2-c}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(b*x^2+a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(1+b*x^2/a)^(1/2)/(d*x^2-c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {427, 426, 424}

$$\frac{\sqrt{c} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{\frac{bx^2}{a}+1} \sqrt{dx^2-c}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]/Sqrt[-c + d*x^2], x]

[Out] (Sqrt[c]*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d)))/(Sqrt[d]*Sqrt[1 + (b*x^2)/a]*Sqrt[-c + d*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\int \frac{\sqrt{a+bx^2}}{\sqrt{-c+dx^2}} dx = \frac{\sqrt{1-\frac{dx^2}{c}} \int \frac{\sqrt{a+bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{-c+dx^2}}$$

$$= \frac{\left(\sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1+\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{1+\frac{bx^2}{a}} \sqrt{-c+dx^2}}$$

$$= \frac{\sqrt{c} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1+\frac{bx^2}{a}} \sqrt{-c+dx^2}}$$

Mathematica [A] time = 0.05, size = 88, normalized size = 1.00

$$\frac{\sqrt{a+bx^2} \sqrt{\frac{c-dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a+bx^2}{a}} \sqrt{dx^2-c}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2]/Sqrt[-c + d*x^2], x]

[Out] (Sqrt[a + b*x^2]*Sqrt[(c - d*x^2)/c]*EllipticE[ArcSin[Sqrt[d/c]*x], -((b*c)/(a*d)))]/(Sqrt[d/c]*Sqrt[(a + b*x^2)/a]*Sqrt[-c + d*x^2])

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+a}}{\sqrt{dx^2-c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2-c)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + a)/sqrt(d*x^2 - c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2-c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 + a)/sqrt(d*x^2 - c), x)

maple [B] time = 0.02, size = 168, normalized size = 1.91

$$\frac{\sqrt{bx^2+a} \sqrt{dx^2-c} \sqrt{\frac{bx^2+a}{a}} \sqrt{-\frac{dx^2-c}{c}} \left(ad \text{EllipticF}\left(\sqrt{\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right) - bc \text{EllipticE}\left(\sqrt{\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right) + bc \text{Ell}\right)}{(bdx^4 + adx^2 - bcx^2 - ac) \sqrt{-\frac{b}{a}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2+a)^(1/2)/(d*x^2-c)^(1/2),x)`

[Out] $(b*x^2+a)^{1/2}*(d*x^2-c)^{1/2}*((b*x^2+a)/a)^{1/2}*(-(d*x^2-c)/c)^{1/2}*(a*EllipticF((-1/a*b)^{1/2}*x, (-a/b/c*d)^{1/2})*d+b*c*EllipticF((-1/a*b)^{1/2}*x, (-a/b/c*d)^{1/2}))-b*c*EllipticE((-1/a*b)^{1/2}*x, (-a/b/c*d)^{1/2})/(b*d*x^4+a*d*x^2-b*c*x^2-a*c)/(-1/a*b)^{1/2}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + a}}{\sqrt{dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+a)^(1/2)/(d*x^2-c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*x^2 + a)/sqrt(d*x^2 - c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 + a}}{\sqrt{dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^(1/2)/(d*x^2 - c)^(1/2),x)`

[Out] `int((a + b*x^2)^(1/2)/(d*x^2 - c)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a + bx^2}}{\sqrt{-c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**(1/2)/(d*x**2-c)**(1/2),x)`

[Out] `Integral(sqrt(a + b*x**2)/sqrt(-c + d*x**2), x)`

$$3.261 \quad \int \frac{\sqrt{-a-bx^2}}{\sqrt{-c+dx^2}} dx$$

Optimal. Leaf size=91

$$\frac{\sqrt{c} \sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{\frac{bx^2}{a}+1} \sqrt{dx^2-c}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(-b*x^2-a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(1+b*x^2/a)^(1/2)/(d*x^2-c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {427, 426, 424}

$$\frac{\sqrt{c} \sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{\frac{bx^2}{a}+1} \sqrt{dx^2-c}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-a - b*x^2]/Sqrt[-c + d*x^2], x]

[Out] (Sqrt[c]*Sqrt[-a - b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d)))/(Sqrt[d]*Sqrt[1 + (b*x^2)/a]*Sqrt[-c + d*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{-a-bx^2}}{\sqrt{-c+dx^2}} dx &= \frac{\sqrt{1-\frac{dx^2}{c}} \int \frac{\sqrt{-a-bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{-c+dx^2}} \\
&= \frac{\left(\sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1+\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{1+\frac{bx^2}{a}} \sqrt{-c+dx^2}} \\
&= \frac{\sqrt{c} \sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1+\frac{bx^2}{a}} \sqrt{-c+dx^2}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 91, normalized size = 1.00

$$\frac{\sqrt{-a-bx^2} \sqrt{\frac{c-dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a+bx^2}{a}} \sqrt{dx^2-c}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-a - b*x^2]/Sqrt[-c + d*x^2], x]

[Out] (Sqrt[-a - b*x^2]*Sqrt[(c - d*x^2)/c]*EllipticE[ArcSin[Sqrt[d/c]*x], -(b*c)/(a*d)))/(Sqrt[d/c]*Sqrt[(a + b*x^2)/a]*Sqrt[-c + d*x^2])

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-bx^2-a}}{\sqrt{dx^2-c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2-a)^(1/2)/(d*x^2-c)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-b*x^2 - a)/sqrt(d*x^2 - c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2-a}}{\sqrt{dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2-a)^(1/2)/(d*x^2-c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-b*x^2 - a)/sqrt(d*x^2 - c), x)

maple [A] time = 0.02, size = 110, normalized size = 1.21

$$\frac{\sqrt{-bx^2-a} \sqrt{dx^2-c} \sqrt{-\frac{dx^2-c}{c}} \sqrt{\frac{bx^2+a}{a}} a \text{EllipticE}\left(\sqrt{\frac{d}{c}}x, \sqrt{-\frac{bc}{ad}}\right)}{(bdx^4 + adx^2 - bcx^2 - ac) \sqrt{\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2-a)^(1/2)/(d*x^2-c)^(1/2),x)

[Out] (-b*x^2-a)^(1/2)*(d*x^2-c)^(1/2)*a*(-(d*x^2-c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)*EllipticE((1/c*d)^(1/2)*x,(-1/a*b*c/d)^(1/2))/(b*d*x^4+a*d*x^2-b*c*x^2-a*c)/(1/c*d)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2-a)^(1/2)/(d*x^2-c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-b*x^2 - a)/sqrt(d*x^2 - c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((- a - b*x^2)^(1/2)/(d*x^2 - c)^(1/2),x)

[Out] int((- a - b*x^2)^(1/2)/(d*x^2 - c)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-a - bx^2}}{\sqrt{-c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2-a)**(1/2)/(d*x**2-c)**(1/2),x)

[Out] Integral(sqrt(-a - b*x**2)/sqrt(-c + d*x**2), x)

$$3.262 \quad \int \frac{\sqrt{a-bx^2}}{\sqrt{c-dx^2}} dx$$

Optimal. Leaf size=88

$$\frac{\sqrt{c} \sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (b*c/a/d)^(1/2))*c^(1/2)*(-b*x^2+a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(1-b*x^2/a)^(1/2)/(-d*x^2+c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, number of rules / integrand size = 0.120, Rules used = {427, 426, 424}

$$\frac{\sqrt{c} \sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a - b*x^2]/Sqrt[c - d*x^2], x]

[Out] (Sqrt[c]*Sqrt[a - b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)]/(Sqrt[d]*Sqrt[1 - (b*x^2)/a]*Sqrt[c - d*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a-bx^2}}{\sqrt{c-dx^2}} dx &= \frac{\sqrt{1-\frac{dx^2}{c}} \int \frac{\sqrt{a-bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{c-dx^2}} \\ &= \frac{\left(\sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1-\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2}} \\ &= \frac{\sqrt{c} \sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 88, normalized size = 1.00

$$\frac{\sqrt{a-bx^2} \sqrt{\frac{c-dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a-bx^2}{a}} \sqrt{c-dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a - b*x^2]/Sqrt[c - d*x^2], x]

[Out] (Sqrt[a - b*x^2]*Sqrt[(c - d*x^2)/c]*EllipticE[ArcSin[Sqrt[d/c]*x], (b*c)/(a*d)))/(Sqrt[d/c]*Sqrt[(a - b*x^2)/a]*Sqrt[c - d*x^2])

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2+a}\sqrt{-dx^2+c}}{dx^2-c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)/(d*x^2 - c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2+a}}{\sqrt{-dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-b*x^2 + a)/sqrt(-d*x^2 + c), x)

maple [A] time = 0.02, size = 109, normalized size = 1.24

$$\frac{\sqrt{-bx^2+a} \sqrt{-dx^2+c} \sqrt{-\frac{dx^2-c}{c}} \sqrt{-\frac{bx^2-a}{a}} a \text{EllipticE}\left(\sqrt{\frac{d}{c}}x, \sqrt{\frac{bc}{ad}}\right)}{(bdx^4 - adx^2 - bcx^2 + ac) \sqrt{\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x)`

[Out] $(-b*x^2+a)^{(1/2)}*(-d*x^2+c)^{(1/2)}*a*(-(d*x^2-c)/c)^{(1/2)}*(-(b*x^2-a)/a)^{(1/2)}*EllipticE((1/c*d)^{(1/2)}*x,(1/a*b*c/d)^{(1/2)})/(b*d*x^4-a*d*x^2-b*c*x^2+a*c)/(1/c*d)^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 + a}}{\sqrt{-dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-b*x^2 + a)/sqrt(-d*x^2 + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{a - bx^2}}{\sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a - b*x^2)^(1/2)/(c - d*x^2)^(1/2),x)`

[Out] `int((a - b*x^2)^(1/2)/(c - d*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a - bx^2}}{\sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x**2+a)**(1/2)/(-d*x**2+c)**(1/2),x)`

[Out] `Integral(sqrt(a - b*x**2)/sqrt(c - d*x**2), x)`

$$3.263 \quad \int \frac{\sqrt{-a+bx^2}}{\sqrt{c-dx^2}} dx$$

Optimal. Leaf size=89

$$\frac{\sqrt{c} \sqrt{bx^2 - a} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c - dx^2}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (b*c/a/d)^(1/2))*c^(1/2)*(b*x^2-a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(1-b*x^2/a)^(1/2)/(-d*x^2+c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {427, 426, 424}

$$\frac{\sqrt{c} \sqrt{bx^2 - a} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c - dx^2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-a + b*x^2]/Sqrt[c - d*x^2], x]

[Out] (Sqrt[c]*Sqrt[-a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)])/(Sqrt[d]*Sqrt[1 - (b*x^2)/a]*Sqrt[c - d*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\int \frac{\sqrt{-a+bx^2}}{\sqrt{c-dx^2}} dx = \frac{\sqrt{1-\frac{dx^2}{c}} \int \frac{\sqrt{-a+bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{c-dx^2}}$$

$$= \frac{\left(\sqrt{-a+bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1-\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2}}$$

$$= \frac{\sqrt{c} \sqrt{-a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2}}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 1.00

$$\frac{\sqrt{bx^2-a} \sqrt{\frac{c-dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a-bx^2}{a}} \sqrt{c-dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-a + b*x^2]/Sqrt[c - d*x^2], x]

[Out] (Sqrt[-a + b*x^2]*Sqrt[(c - d*x^2)/c]*EllipticE[ArcSin[Sqrt[d/c]*x], (b*c)/(a*d)))/(Sqrt[d/c]*Sqrt[(a - b*x^2)/a]*Sqrt[c - d*x^2])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{bx^2-a} \sqrt{-dx^2+c}}{dx^2-c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(b*x^2 - a)*sqrt(-d*x^2 + c)/(d*x^2 - c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2-a}}{\sqrt{-dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 - a)/sqrt(-d*x^2 + c), x)

maple [B] time = 0.03, size = 165, normalized size = 1.85

$$\frac{\sqrt{bx^2-a} \sqrt{-dx^2+c} \sqrt{-\frac{bx^2-a}{a}} \sqrt{-\frac{dx^2-c}{c}} \left(ad \text{EllipticF}\left(\sqrt{\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right) + bc \text{EllipticE}\left(\sqrt{\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right) - bc \text{EllipticF}\left(\sqrt{\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right)\right)}{(bdx^4 - adx^2 - bcx^2 + ac) \sqrt{\frac{b}{a}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2),x)`

[Out] $(b*x^2-a)^{(1/2)}*(-d*x^2+c)^{(1/2)}*(-(b*x^2-a)/a)^{(1/2)}*(-(d*x^2-c)/c)^{(1/2)}*(a*\text{EllipticF}((1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})*d-b*c*\text{EllipticF}((1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)})+b*c*\text{EllipticE}((1/a*b)^{(1/2)}*x,(a/b/c*d)^{(1/2)}))/((b*d*x^4-a*d*x^2-b*c*x^2+a*c)/(1/a*b)^{(1/2)}/d)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{-dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*x^2 - a)/sqrt(-d*x^2 + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2 - a)^(1/2)/(c - d*x^2)^(1/2),x)`

[Out] `int((b*x^2 - a)^(1/2)/(c - d*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-a + bx^2}}{\sqrt{c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2-a)**(1/2)/(-d*x**2+c)**(1/2),x)`

[Out] `Integral(sqrt(-a + b*x**2)/sqrt(c - d*x**2), x)`

$$3.264 \quad \int \frac{\sqrt{a-bx^2}}{\sqrt{-c+dx^2}} dx$$

Optimal. Leaf size=89

$$\frac{\sqrt{c} \sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1-\frac{bx^2}{a}} \sqrt{dx^2-c}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (b*c/a/d)^(1/2))*c^(1/2)*(-b*x^2+a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(1-b*x^2/a)^(1/2)/(d*x^2-c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {427, 426, 424}

$$\frac{\sqrt{c} \sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1-\frac{bx^2}{a}} \sqrt{dx^2-c}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a - b*x^2]/Sqrt[-c + d*x^2],x]

[Out] (Sqrt[c]*Sqrt[a - b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)]/(Sqrt[d]*Sqrt[1 - (b*x^2)/a]*Sqrt[-c + d*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a-bx^2}}{\sqrt{-c+dx^2}} dx &= \frac{\sqrt{1-\frac{dx^2}{c}} \int \frac{\sqrt{a-bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{-c+dx^2}} \\ &= \frac{\left(\sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1-\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{1-\frac{bx^2}{a}} \sqrt{-c+dx^2}} \\ &= \frac{\sqrt{c} \sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1-\frac{bx^2}{a}} \sqrt{-c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 1.00

$$\frac{\sqrt{a-bx^2} \sqrt{\frac{c-dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a-bx^2}{a}} \sqrt{dx^2-c}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a - b*x^2]/Sqrt[-c + d*x^2], x]

[Out] (Sqrt[a - b*x^2]*Sqrt[(c - d*x^2)/c]*EllipticE[ArcSin[Sqrt[d/c]*x], (b*c)/(a*d)))/(Sqrt[d/c]*Sqrt[(a - b*x^2)/a]*Sqrt[-c + d*x^2])

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-bx^2+a}}{\sqrt{dx^2-c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(1/2)/(d*x^2-c)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-b*x^2 + a)/sqrt(d*x^2 - c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2+a}}{\sqrt{dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(1/2)/(d*x^2-c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-b*x^2 + a)/sqrt(d*x^2 - c), x)

maple [B] time = 0.02, size = 166, normalized size = 1.87

$$\frac{\left(-ad \text{EllipticF}\left(\sqrt{\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right) - bc \text{EllipticE}\left(\sqrt{\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right) + bc \text{EllipticF}\left(\sqrt{\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right)\right) \sqrt{-bx^2+a} \sqrt{dx^2-c}}{(bdx^4 - adx^2 - bcx^2 + ac) \sqrt{\frac{b}{a}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-b*x^2+a)^(1/2)/(d*x^2-c)^(1/2),x)`

[Out] `(-a*EllipticF((1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))*d+b*c*EllipticF((1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))-b*c*EllipticE((1/a*b)^(1/2)*x,(a/b/c*d)^(1/2)))*(-b*x^2+a)^(1/2)*(d*x^2-c)^(1/2)*(-(b*x^2-a)/a)^(1/2)*(-(d*x^2-c)/c)^(1/2)/(b*d*x^4-a*d*x^2-b*c*x^2+a*c)/(1/a*b)^(1/2)/d`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 + a}}{\sqrt{dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x^2+a)^(1/2)/(d*x^2-c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-b*x^2 + a)/sqrt(d*x^2 - c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{a - bx^2}}{\sqrt{dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a - b*x^2)^(1/2)/(d*x^2 - c)^(1/2),x)`

[Out] `int((a - b*x^2)^(1/2)/(d*x^2 - c)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a - bx^2}}{\sqrt{-c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-b*x**2+a)**(1/2)/(d*x**2-c)**(1/2),x)`

[Out] `Integral(sqrt(a - b*x**2)/sqrt(-c + d*x**2), x)`

$$3.265 \quad \int \frac{\sqrt{-a+bx^2}}{\sqrt{-c+dx^2}} dx$$

Optimal. Leaf size=90

$$\frac{\sqrt{c} \sqrt{bx^2 - a} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1 - \frac{bx^2}{a}} \sqrt{dx^2 - c}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (b*c/a/d)^(1/2))*c^(1/2)*(b*x^2-a)^(1/2)*(1-d*x^2/c)^(1/2)/d^(1/2)/(1-b*x^2/a)^(1/2)/(d*x^2-c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {427, 426, 424}

$$\frac{\sqrt{c} \sqrt{bx^2 - a} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1 - \frac{bx^2}{a}} \sqrt{dx^2 - c}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-a + b*x^2]/Sqrt[-c + d*x^2], x]

[Out] (Sqrt[c]*Sqrt[-a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], (b*c)/(a*d)])/(Sqrt[d]*Sqrt[1 - (b*x^2)/a]*Sqrt[-c + d*x^2])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{-a+bx^2}}{\sqrt{-c+dx^2}} dx &= \frac{\sqrt{1-\frac{dx^2}{c}} \int \frac{\sqrt{-a+bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{-c+dx^2}} \\
&= \frac{\left(\sqrt{-a+bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1-\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{\sqrt{1-\frac{bx^2}{a}} \sqrt{-c+dx^2}} \\
&= \frac{\sqrt{c} \sqrt{-a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{1-\frac{bx^2}{a}} \sqrt{-c+dx^2}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 90, normalized size = 1.00

$$\frac{\sqrt{bx^2-a} \sqrt{\frac{c-dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a-bx^2}{a}} \sqrt{dx^2-c}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-a + b*x^2]/Sqrt[-c + d*x^2], x]

[Out] (Sqrt[-a + b*x^2]*Sqrt[(c - d*x^2)/c]*EllipticE[ArcSin[Sqrt[d/c]*x], (b*c)/(a*d)))/(Sqrt[d/c]*Sqrt[(a - b*x^2)/a]*Sqrt[-c + d*x^2])

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2-a}}{\sqrt{dx^2-c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2-a)^(1/2)/(d*x^2-c)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 - a)/sqrt(d*x^2 - c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2-a}}{\sqrt{dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2-a)^(1/2)/(d*x^2-c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 - a)/sqrt(d*x^2 - c), x)

maple [A] time = 0.02, size = 111, normalized size = 1.23

$$\frac{\sqrt{bx^2-a} \sqrt{dx^2-c} \sqrt{-\frac{dx^2-c}{c}} \sqrt{-\frac{bx^2-a}{a}} a \text{EllipticE}\left(\sqrt{\frac{d}{c}}x, \sqrt{\frac{bc}{ad}}\right)}{(-bdx^4 + adx^2 + bcx^2 - ac) \sqrt{\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2-a)^(1/2)/(d*x^2-c)^(1/2),x)`

[Out] `1/(-b*d*x^4+a*d*x^2+b*c*x^2-a*c)/(1/c*d)^(1/2)*(b*x^2-a)^(1/2)*(d*x^2-c)^(1/2)*a*(-(d*x^2-c)/c)^(1/2)*(-(b*x^2-a)/a)^(1/2)*EllipticE((1/c*d)^(1/2)*x,(1/a*b*c/d)^(1/2))`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2-a)^(1/2)/(d*x^2-c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*x^2 - a)/sqrt(d*x^2 - c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2 - a)^(1/2)/(d*x^2 - c)^(1/2),x)`

[Out] `int((b*x^2 - a)^(1/2)/(d*x^2 - c)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-a + bx^2}}{\sqrt{-c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2-a)**(1/2)/(d*x**2-c)**(1/2),x)`

[Out] `Integral(sqrt(-a + b*x**2)/sqrt(-c + d*x**2), x)`

$$3.266 \quad \int \frac{\sqrt{a+bx^2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=194

$$\frac{\sqrt{c} \sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{x\sqrt{a+bx^2}}{\sqrt{c+dx^2}} - \frac{\sqrt{c} \sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $x*(b*x^2+a)^{(1/2)}/(d*x^2+c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*E$
 $llipticE(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*$
 $x^2+a)^{(1/2)}/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+(1/(1+$
 $d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*EllipticF(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},$
 $(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x$
 $^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.08, antiderivative size = 194, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {422, 418, 492, 411}

$$\frac{x\sqrt{a+bx^2}}{\sqrt{c+dx^2}} + \frac{\sqrt{c} \sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{\sqrt{c} \sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x]

[Out] $(x*\operatorname{Sqrt}[a + b*x^2])/ \operatorname{Sqrt}[c + d*x^2] - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a+bx^2}}{\sqrt{c+dx^2}} dx &= a \int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx + b \int \frac{x^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx \\ &= \frac{x\sqrt{a+bx^2}}{\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} - c \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^{3/2}} dx \\ &= \frac{x\sqrt{a+bx^2}}{\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 86, normalized size = 0.44

$$\frac{\sqrt{a+bx^2}\sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{-\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{-\frac{d}{c}}\sqrt{\frac{a+bx^2}{a}}\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[a + b*x^2]/Sqrt[c + d*x^2], x]
```

```
[Out] (Sqrt[a + b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], (b*c)/(a*d)))/(Sqrt[-(d/c)]*Sqrt[(a + b*x^2)/a]*Sqrt[c + d*x^2])
```

fricas [F] time = 0.72, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+a}}{\sqrt{dx^2+c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*x^2 + a)/sqrt(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*x^2 + a)/sqrt(d*x^2 + c), x)
```

maple [A] time = 0.01, size = 158, normalized size = 0.81

$$\frac{\sqrt{bx^2+a} \sqrt{dx^2+c} \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} \left(ad \operatorname{EllipticF}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right) + bc \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right) - bc \operatorname{EllipticF}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right) \right)}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{b}{a}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)/(d*x^2+c)^(1/2), x)

[Out] (b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*(a*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*d-b*c*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))+b*c*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2)))/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)/sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)/(c + d*x^2)^(1/2), x)

[Out] int((a + b*x^2)^(1/2)/(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a + bx^2}}{\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)/(d*x**2+c)**(1/2), x)

[Out] Integral(sqrt(a + b*x**2)/sqrt(c + d*x**2), x)

$$3.267 \quad \int \frac{\sqrt{-a-bx^2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=203

$$\frac{\sqrt{c} \sqrt{-a-bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + \frac{x\sqrt{-a-bx^2}}{\sqrt{c+dx^2}} - \frac{\sqrt{c} \sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $x*(-b*x^2-a)^{(1/2)}/(d*x^2+c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}* \operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(-b*x^2-a)^{(1/2)}/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(-b*x^2-a)^{(1/2)}/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.09, antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {422, 418, 492, 411}

$$\frac{x\sqrt{-a-bx^2}}{\sqrt{c+dx^2}} + \frac{\sqrt{c} \sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{\sqrt{c} \sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{c+dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-a - b*x^2]/Sqrt[c + d*x^2], x]

[Out] $(x*\operatorname{Sqrt}[-a - b*x^2])/ \operatorname{Sqrt}[c + d*x^2] - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[-a - b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)]/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[-a - b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)]/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/((c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/((a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-a-bx^2}}{\sqrt{c+dx^2}} dx &= -\left(a \int \frac{1}{\sqrt{-a-bx^2}\sqrt{c+dx^2}} dx\right) - b \int \frac{x^2}{\sqrt{-a-bx^2}\sqrt{c+dx^2}} dx \\ &= \frac{x\sqrt{-a-bx^2}}{\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} - c \int \frac{\sqrt{-a-bx^2}}{(c+dx^2)^{3/2}} dx \\ &= \frac{x\sqrt{-a-bx^2}}{\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 0.44

$$\frac{\sqrt{-a-bx^2}\sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{-\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{-\frac{d}{c}}\sqrt{\frac{a+bx^2}{a}}\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[-a - b*x^2]/Sqrt[c + d*x^2], x]
```

```
[Out] (Sqrt[-a - b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], (b*c)/(a*d)))/(Sqrt[-(d/c)]*Sqrt[(a + b*x^2)/a]*Sqrt[c + d*x^2])
```

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-bx^2 - a}}{\sqrt{dx^2 + c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-b*x^2-a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(-b*x^2 - a)/sqrt(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-b*x^2-a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(-b*x^2 - a)/sqrt(d*x^2 + c), x)
```

maple [A] time = 0.03, size = 104, normalized size = 0.51

$$\frac{\sqrt{-bx^2 - a} \sqrt{dx^2 + c} \sqrt{\frac{dx^2 + c}{c}} \sqrt{\frac{bx^2 + a}{a}} a \operatorname{EllipticE}\left(\sqrt{-\frac{d}{c}} x, \sqrt{\frac{bc}{ad}}\right)}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2-a)^(1/2)/(d*x^2+c)^(1/2), x)

[Out] (-b*x^2-a)^(1/2)*(d*x^2+c)^(1/2)*a*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)*
EllipticE(x*(-1/c*d)^(1/2), (1/a*b*c/d)^(1/2))/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)
/(-1/c*d)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2-a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(-b*x^2 - a)/sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((- a - b*x^2)^(1/2)/(c + d*x^2)^(1/2), x)

[Out] int((- a - b*x^2)^(1/2)/(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-a - bx^2}}{\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2-a)**(1/2)/(d*x**2+c)**(1/2), x)

[Out] Integral(sqrt(-a - b*x**2)/sqrt(c + d*x**2), x)

$$3.268 \quad \int \frac{\sqrt{a+bx^2}}{\sqrt{-c-dx^2}} dx$$

Optimal. Leaf size=203

$$\frac{\sqrt{c} \sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{x\sqrt{a+bx^2}}{\sqrt{-c-dx^2}} - \frac{\sqrt{c} \sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $x*(b*x^2+a)^{(1/2)/(-d*x^2-c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}* \operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(1/2)/(-d*x^2-c)^{(1/2)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}+(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(b*x^2+a)^{(1/2)}/d^{(1/2)/(-d*x^2-c)^{(1/2)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}}$

Rubi [A] time = 0.09, antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {422, 418, 492, 411}

$$\frac{x\sqrt{a+bx^2}}{\sqrt{-c-dx^2}} + \frac{\sqrt{c} \sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{\sqrt{c} \sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2]/Sqrt[-c - d*x^2], x]

[Out] $(x*\operatorname{Sqrt}[a + b*x^2])/ \operatorname{Sqrt}[-c - d*x^2] - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-c - d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]) + (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-c - d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a+bx^2}}{\sqrt{-c-dx^2}} dx &= a \int \frac{1}{\sqrt{a+bx^2}\sqrt{-c-dx^2}} dx + b \int \frac{x^2}{\sqrt{a+bx^2}\sqrt{-c-dx^2}} dx \\ &= \frac{x\sqrt{a+bx^2}}{\sqrt{-c-dx^2}} + \frac{\sqrt{c}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + c \int \frac{\sqrt{a+bx^2}}{(-c-dx^2)^{3/2}} dx \\ &= \frac{x\sqrt{a+bx^2}}{\sqrt{-c-dx^2}} - \frac{\sqrt{c}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{c}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 0.44

$$\frac{\sqrt{a+bx^2}\sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{-\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{-\frac{d}{c}}\sqrt{\frac{a+bx^2}{a}}\sqrt{-c-dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[a + b*x^2]/Sqrt[-c - d*x^2], x]
```

```
[Out] (Sqrt[a + b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], (b*c)/(a*d)))/(Sqrt[-(d/c)]*Sqrt[(a + b*x^2)/a]*Sqrt[-c - d*x^2])
```

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{bx^2+a}\sqrt{-dx^2-c}}{dx^2+c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(-sqrt(b*x^2 + a)*sqrt(-d*x^2 - c)/(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{-dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*x^2 + a)/sqrt(-d*x^2 - c), x)
```

maple [A] time = 0.02, size = 108, normalized size = 0.53

$$\frac{\sqrt{bx^2+a} \sqrt{-dx^2-c} \sqrt{\frac{dx^2+c}{c}} \sqrt{\frac{bx^2+a}{a}} a \operatorname{EllipticE}\left(\sqrt{-\frac{d}{c}} x, \sqrt{\frac{bc}{ad}}\right)}{(-bdx^4 - adx^2 - bcx^2 - ac) \sqrt{-\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2), x)

[Out] (b*x^2+a)^(1/2)*(-d*x^2-c)^(1/2)*a*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)*EllipticE((-1/c*d)^(1/2)*x, (1/a*b*c/d)^(1/2))/(-b*d*x^4-a*d*x^2-b*c*x^2-a*c)/(-1/c*d)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{-dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 + a)/sqrt(-d*x^2 - c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{bx^2+a}}{\sqrt{-dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/2)/(-c - d*x^2)^(1/2), x)

[Out] int((a + b*x^2)^(1/2)/(-c - d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a+bx^2}}{\sqrt{-c-dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/2)/(-d*x**2-c)**(1/2), x)

[Out] Integral(sqrt(a + b*x**2)/sqrt(-c - d*x**2), x)

$$3.269 \quad \int \frac{\sqrt{-a-bx^2}}{\sqrt{-c-dx^2}} dx$$

Optimal. Leaf size=212

$$\frac{\sqrt{c} \sqrt{-a-bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{x\sqrt{-a-bx^2}}{\sqrt{-c-dx^2}} - \frac{\sqrt{c} \sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $x*(-b*x^2-a)^{(1/2)/(-d*x^2-c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}$
 $*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*($
 $-b*x^2-a)^{(1/2)}/d^{(1/2)/(-d*x^2-c)^{(1/2)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}+(1$
 $/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)/(1+d*x^2/$
 $c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*(-b*x^2-a)^{(1/2)}/d^{(1/2)/(-d*x^2-c)^{(1/2)$
 $2)/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 212, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 29, $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$, Rules used = {422, 418, 492, 411}

$$\frac{x\sqrt{-a-bx^2}}{\sqrt{-c-dx^2}} + \frac{\sqrt{c} \sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{\sqrt{c} \sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d} \sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-a - b*x^2]/Sqrt[-c - d*x^2], x]

[Out] $(x*\operatorname{Sqrt}[-a - b*x^2])/(\operatorname{Sqrt}[-c - d*x^2]) - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[-a - b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-c - d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]) + (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[-a - b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-c - d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-a-bx^2}}{\sqrt{-c-dx^2}} dx &= -\left(a \int \frac{1}{\sqrt{-a-bx^2}\sqrt{-c-dx^2}} dx\right) - b \int \frac{x^2}{\sqrt{-a-bx^2}\sqrt{-c-dx^2}} dx \\ &= \frac{x\sqrt{-a-bx^2}}{\sqrt{-c-dx^2}} + \frac{\sqrt{c}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + c \int \frac{\sqrt{-a-bx^2}}{(-c-dx^2)^{3/2}} dx \\ &= \frac{x\sqrt{-a-bx^2}}{\sqrt{-c-dx^2}} - \frac{\sqrt{c}\sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{c}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{\sqrt{d}\sqrt{-c-dx^2} \sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 92, normalized size = 0.43

$$\frac{\sqrt{-a-bx^2} \sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{-\frac{d}{c}}x\right) \middle| \frac{bc}{ad}\right)}{\sqrt{-\frac{d}{c}} \sqrt{\frac{a+bx^2}{a}} \sqrt{-c-dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[-a - b*x^2]/Sqrt[-c - d*x^2], x]
```

```
[Out] (Sqrt[-a - b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], (b*c)/(a*d)))/(Sqrt[-(d/c)]*Sqrt[(a + b*x^2)/a]*Sqrt[-c - d*x^2])
```

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2-a}\sqrt{-dx^2-c}}{dx^2+c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(-sqrt(-b*x^2 - a)*sqrt(-d*x^2 - c)/(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2-a}}{\sqrt{-dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(-b*x^2 - a)/sqrt(-d*x^2 - c), x)
```


maple [A] time = 0.02, size = 165, normalized size = 0.78

$$\frac{\left(-ad \operatorname{EllipticF}\left(\sqrt{-\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right) - bc \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right) + bc \operatorname{EllipticF}\left(\sqrt{-\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right)\right) \sqrt{-bx^2 - a} \sqrt{-dx^2 - c}}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{b}{a}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2),x)

[Out] (-a*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*d+b*c*EllipticF((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))-b*c*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))*(-b*x^2-a)^(1/2)*(-d*x^2-c)^(1/2)*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{-dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-b*x^2 - a)/sqrt(-d*x^2 - c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{-bx^2 - a}}{\sqrt{-dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((- a - b*x^2)^(1/2)/(- c - d*x^2)^(1/2),x)

[Out] int((- a - b*x^2)^(1/2)/(- c - d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-a - bx^2}}{\sqrt{-c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2-a)**(1/2)/(-d*x**2-c)**(1/2),x)

[Out] Integral(sqrt(-a - b*x**2)/sqrt(-c - d*x**2), x)

$$3.270 \quad \int \frac{\sqrt{a-bx^2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=189

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} (ad + bc) \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), -\frac{ad}{bc}\right)}{\sqrt{b}d\sqrt{a-bx^2}\sqrt{c+dx^2}} - \frac{\sqrt{a}\sqrt{b}\sqrt{1 - \frac{bx^2}{a}}\sqrt{c+dx^2}E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle| -\frac{ad}{bc}\right)}{d\sqrt{a-bx^2}\sqrt{\frac{dx^2}{c} + 1}}$$

[Out] -EllipticE(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*b^(1/2)*(1-b*x^2/a)^(1/2)*(d*x^2+c)^(1/2)/d/(-b*x^2+a)^(1/2)/(1+d*x^2/c)^(1/2)+(a*d+b*c)*EllipticF(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/d/b^(1/2)/(-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2)

Rubi [A] time = 0.12, antiderivative size = 189, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {423, 427, 426, 424, 421, 419}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} (ad + bc) F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle| -\frac{ad}{bc}\right)}{\sqrt{b}d\sqrt{a-bx^2}\sqrt{c+dx^2}} - \frac{\sqrt{a}\sqrt{b}\sqrt{1 - \frac{bx^2}{a}}\sqrt{c+dx^2}E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle| -\frac{ad}{bc}\right)}{d\sqrt{a-bx^2}\sqrt{\frac{dx^2}{c} + 1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a - b*x^2]/Sqrt[c + d*x^2], x]

[Out] -((Sqrt[a]*Sqrt[b]*Sqrt[1 - (b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -(a*d)/(b*c)])/(d*Sqrt[a - b*x^2]*Sqrt[1 + (d*x^2)/c]) + (Sqrt[a]*(b*c + a*d)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -(a*d)/(b*c)])/(Sqrt[b]*d*Sqrt[a - b*x^2]*Sqrt[c + d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]
```

Rule 427

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a-bx^2}}{\sqrt{c+dx^2}} dx &= -\frac{b \int \frac{\sqrt{c+dx^2}}{\sqrt{a-bx^2}} dx}{d} + \frac{(bc+ad) \int \frac{1}{\sqrt{a-bx^2} \sqrt{c+dx^2}} dx}{d} \\ &= -\frac{\left(b\sqrt{1-\frac{bx^2}{a}}\right) \int \frac{\sqrt{c+dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{d\sqrt{a-bx^2}} + \frac{\left((bc+ad)\sqrt{1+\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{a-bx^2} \sqrt{1+\frac{dx^2}{c}}} dx}{d\sqrt{c+dx^2}} \\ &= -\frac{\left(b\sqrt{1-\frac{bx^2}{a}} \sqrt{c+dx^2}\right) \int \frac{\sqrt{1+\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{d\sqrt{a-bx^2} \sqrt{1+\frac{dx^2}{c}}} + \frac{\left((bc+ad)\sqrt{1-\frac{bx^2}{a}} \sqrt{1+\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1-\frac{bx^2}{a}} \sqrt{1+\frac{dx^2}{c}}} dx}{d\sqrt{a-bx^2} \sqrt{c+dx^2}} \\ &= -\frac{\sqrt{a} \sqrt{b} \sqrt{1-\frac{bx^2}{a}} \sqrt{c+dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{d\sqrt{a-bx^2} \sqrt{1+\frac{dx^2}{c}}} + \frac{\sqrt{a} (bc+ad) \sqrt{1-\frac{bx^2}{a}} \sqrt{1+\frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{b} d \sqrt{a-bx^2} \sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 0.47

$$\frac{\sqrt{a-bx^2} \sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a-bx^2}{a}} \sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[a - b*x^2]/Sqrt[c + d*x^2], x]
```

```
[Out] (Sqrt[a - b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], -((b*c)/(a*d))]/(Sqrt[-(d/c)]*Sqrt[(a - b*x^2)/a]*Sqrt[c + d*x^2])
```

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-bx^2+a}}{\sqrt{dx^2+c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(-b*x^2 + a)/sqrt(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2+a}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-b*x^2 + a)/sqrt(d*x^2 + c), x)

maple [A] time = 0.02, size = 164, normalized size = 0.87

$$\frac{\sqrt{-bx^2+a} \sqrt{dx^2+c} \sqrt{-\frac{bx^2-a}{a}} \sqrt{\frac{dx^2+c}{c}} \left(-ad \operatorname{EllipticF}\left(\sqrt{\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right) + bc \operatorname{EllipticE}\left(\sqrt{\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right) - bc \operatorname{EllipticF}\left(\sqrt{\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right) \right)}{(bdx^4 - adx^2 + bcx^2 - ac) \sqrt{\frac{b}{a}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x)

[Out] (-b*x^2+a)^(1/2)*(d*x^2+c)^(1/2)*(-b*x^2-a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*(b*c*EllipticE((1/a*b)^(1/2)*x, (-a/b/c*d)^(1/2))-a*EllipticF((1/a*b)^(1/2)*x, (-a/b/c*d)^(1/2))*d-b*c*EllipticF((1/a*b)^(1/2)*x, (-a/b/c*d)^(1/2)))/(b*d*x^4-a*d*x^2+b*c*x^2-a*c)/(1/a*b)^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2+a}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-b*x^2 + a)/sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{a-bx^2}}{\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(1/2)/(c + d*x^2)^(1/2),x)

[Out] int((a - b*x^2)^(1/2)/(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a-bx^2}}{\sqrt{c+dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(1/2)/(d*x**2+c)**(1/2),x)

[Out] Integral(sqrt(a - b*x**2)/sqrt(c + d*x**2), x)

$$3.271 \quad \int \frac{\sqrt{-a+bx^2}}{\sqrt{c+dx^2}} dx$$

Optimal. Leaf size=191

$$\frac{\sqrt{a} \sqrt{b} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c + dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{d\sqrt{bx^2 - a} \sqrt{\frac{dx^2}{c} + 1}} - \frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} (ad + bc) \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), -\frac{ad}{bc}\right)}{\sqrt{b} d \sqrt{bx^2 - a} \sqrt{c + dx^2}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*b^(1/2)*(1-b*x^2/a)^(1/2)*(d*x^2+c)^(1/2)/d/(b*x^2-a)^(1/2)/((1+d*x^2/c)^(1/2)-(a*d+b*c)*EllipticF(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/d/b^(1/2)/(b*x^2-a)^(1/2)/(d*x^2+c)^(1/2)

Rubi [A] time = 0.12, antiderivative size = 191, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {423, 427, 426, 424, 421, 419}

$$\frac{\sqrt{a} \sqrt{b} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c + dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{d\sqrt{bx^2 - a} \sqrt{\frac{dx^2}{c} + 1}} - \frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} (ad + bc) F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} d \sqrt{bx^2 - a} \sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-a + b*x^2]/Sqrt[c + d*x^2], x]

[Out] (Sqrt[a]*Sqrt[b]*Sqrt[1 - (b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -(a*d)/(b*c))]/(d*Sqrt[-a + b*x^2]*Sqrt[1 + (d*x^2)/c]) - (Sqrt[a]*(b*c + a*d)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -(a*d)/(b*c)])/((Sqrt[b]*d*Sqrt[-a + b*x^2]*Sqrt[c + d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0
]
```

Rule 427

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-a + bx^2}}{\sqrt{c + dx^2}} dx &= \frac{b \int \frac{\sqrt{c+dx^2}}{\sqrt{-a+bx^2}} dx}{d} - \frac{(bc + ad) \int \frac{1}{\sqrt{-a+bx^2} \sqrt{c+dx^2}} dx}{d} \\ &= \frac{\left(b\sqrt{1 - \frac{bx^2}{a}}\right) \int \frac{\sqrt{c+dx^2}}{\sqrt{1 - \frac{bx^2}{a}}} dx}{d\sqrt{-a + bx^2}} - \frac{\left((bc + ad)\sqrt{1 + \frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{-a+bx^2} \sqrt{1 + \frac{dx^2}{c}}} dx}{d\sqrt{c + dx^2}} \\ &= \frac{\left(b\sqrt{1 - \frac{bx^2}{a}} \sqrt{c + dx^2}\right) \int \frac{\sqrt{1 + \frac{dx^2}{c}}}{\sqrt{1 - \frac{bx^2}{a}}} dx}{d\sqrt{-a + bx^2} \sqrt{1 + \frac{dx^2}{c}}} - \frac{\left((bc + ad)\sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}}} dx}{d\sqrt{-a + bx^2} \sqrt{c + dx^2}} \\ &= \frac{\sqrt{a} \sqrt{b} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c + dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{d\sqrt{-a + bx^2} \sqrt{1 + \frac{dx^2}{c}}} - \frac{\sqrt{a} (bc + ad) \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} d \sqrt{-a + bx^2} \sqrt{c + dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 90, normalized size = 0.47

$$\frac{\sqrt{bx^2 - a} \sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}} x\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a-bx^2}{a}} \sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[-a + b*x^2]/Sqrt[c + d*x^2], x]
```

```
[Out] (Sqrt[-a + b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], -((
b*c)/(a*d))]/(Sqrt[-(d/c)]*Sqrt[(a - b*x^2)/a]*Sqrt[c + d*x^2])
```

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2 - a}}{\sqrt{dx^2 + c}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2-a)^(1/2)/(d*x^2+c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*x^2 - a)/sqrt(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2-a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 - a)/sqrt(d*x^2 + c), x)

maple [A] time = 0.02, size = 109, normalized size = 0.57

$$\frac{\sqrt{bx^2 - a} \sqrt{dx^2 + c} \sqrt{\frac{dx^2+c}{c}} \sqrt{\frac{bx^2-a}{a}} a \operatorname{EllipticE}\left(\sqrt{\frac{d}{c}} x, \sqrt{\frac{bc}{ad}}\right)}{(-bdx^4 + adx^2 - bcx^2 + ac) \sqrt{\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2-a)^(1/2)/(d*x^2+c)^(1/2),x)

[Out] 1/(-b*d*x^4+a*d*x^2-b*c*x^2+a*c)/(-1/c*d)^(1/2)*(b*x^2-a)^(1/2)*(d*x^2+c)^(1/2)*a*((d*x^2+c)/c)^(1/2)*(-(b*x^2-a)/a)^(1/2)*EllipticE((-1/c*d)^(1/2)*x, (-1/a*b*c/d)^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2-a)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 - a)/sqrt(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2 - a)^(1/2)/(c + d*x^2)^(1/2),x)

[Out] int((b*x^2 - a)^(1/2)/(c + d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-a + bx^2}}{\sqrt{c + dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2-a)**(1/2)/(d*x**2+c)**(1/2),x)

[Out] Integral(sqrt(-a + b*x**2)/sqrt(c + d*x**2), x)

$$3.272 \quad \int \frac{\sqrt{a-bx^2}}{\sqrt{-c-dx^2}} dx$$

Optimal. Leaf size=194

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} (ad + bc) \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), -\frac{ad}{bc}\right) + \sqrt{a} \sqrt{b} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} d \sqrt{a - bx^2} \sqrt{-c - dx^2} + d \sqrt{a - bx^2} \sqrt{\frac{dx^2}{c} + 1}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*b^(1/2)*(1-b*x^2/a)^(1/2)*(-d*x^2-c)^(1/2)/d/(-b*x^2+a)^(1/2)/(1+d*x^2/c)^(1/2)+(a*d+b*c)*EllipticF(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/d/b^(1/2)/(-b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2)

Rubi [A] time = 0.13, antiderivative size = 194, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {423, 427, 426, 424, 421, 419}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} (ad + bc) F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right) + \sqrt{a} \sqrt{b} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} d \sqrt{a - bx^2} \sqrt{-c - dx^2} + d \sqrt{a - bx^2} \sqrt{\frac{dx^2}{c} + 1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a - b*x^2]/Sqrt[-c - d*x^2], x]

[Out] (Sqrt[a]*Sqrt[b]*Sqrt[1 - (b*x^2)/a]*Sqrt[-c - d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))]/(d*Sqrt[a - b*x^2]*Sqrt[1 + (d*x^2)/c]) + (Sqrt[a]*(b*c + a*d)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))]/(Sqrt[b]*d*Sqrt[a - b*x^2]*Sqrt[-c - d*x^2]))

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])]

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426


```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]
```

Rule 427

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]
```

Rubi steps

$$\int \frac{\sqrt{a - bx^2}}{\sqrt{-c - dx^2}} dx = \frac{b \int \frac{\sqrt{-c-dx^2}}{\sqrt{a-bx^2}} dx}{d} + \frac{(bc + ad) \int \frac{1}{\sqrt{a-bx^2} \sqrt{-c-dx^2}} dx}{d}$$

$$= \frac{\left(b\sqrt{1 - \frac{bx^2}{a}}\right) \int \frac{\sqrt{-c-dx^2}}{\sqrt{1 - \frac{bx^2}{a}}} dx}{d\sqrt{a - bx^2}} + \frac{\left((bc + ad)\sqrt{1 + \frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{a-bx^2} \sqrt{1 + \frac{dx^2}{c}}} dx}{d\sqrt{-c - dx^2}}$$

$$= \frac{\left(b\sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2}\right) \int \frac{\sqrt{1 + \frac{dx^2}{c}}}{\sqrt{1 - \frac{bx^2}{a}}} dx}{d\sqrt{a - bx^2} \sqrt{1 + \frac{dx^2}{c}}} + \frac{\left((bc + ad)\sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}}} dx}{d\sqrt{a - bx^2} \sqrt{-c - dx^2}}$$

$$= \frac{\sqrt{a} \sqrt{b} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{d\sqrt{a - bx^2} \sqrt{1 + \frac{dx^2}{c}}} + \frac{\sqrt{a} (bc + ad) \sqrt{1 - \frac{bx^2}{a}} \sqrt{1 + \frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} d\sqrt{a - bx^2} \sqrt{-c - dx^2}}$$

Mathematica [A] time = 0.05, size = 92, normalized size = 0.47

$$\frac{\sqrt{a - bx^2} \sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}} x\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a-bx^2}{a}} \sqrt{-c - dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[a - b*x^2]/Sqrt[-c - d*x^2], x]
[Out] (Sqrt[a - b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], -((b*c)/(a*d))]/(Sqrt[-(d/c)]*Sqrt[(a - b*x^2)/a]*Sqrt[-c - d*x^2])
```

fricas [F] time = 0.92, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2 + a} \sqrt{-dx^2 - c}}{dx^2 + c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2), x, algorithm="fricas")
[Out] integral(-sqrt(-b*x^2 + a)*sqrt(-d*x^2 - c)/(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 + a}}{\sqrt{-dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-b*x^2 + a)/sqrt(-d*x^2 - c), x)

maple [A] time = 0.02, size = 111, normalized size = 0.57

$$\frac{\sqrt{-bx^2 + a} \sqrt{-dx^2 - c} \sqrt{\frac{dx^2+c}{c}} \sqrt{\frac{bx^2-a}{a}} a \operatorname{EllipticE}\left(\sqrt{\frac{d}{c}} x, \sqrt{\frac{bc}{ad}}\right)}{(bdx^4 - adx^2 + bcx^2 - ac) \sqrt{\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2),x)

[Out] (-b*x^2+a)^(1/2)*(-d*x^2-c)^(1/2)*a*((d*x^2+c)/c)^(1/2)*(-b*x^2-a)/a)^(1/2)*EllipticE((-1/c*d)^(1/2)*x, (-1/a*b*c/d)^(1/2))/(b*d*x^4-a*d*x^2+b*c*x^2-a*c)/(-1/c*d)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-bx^2 + a}}{\sqrt{-dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x^2+a)^(1/2)/(-d*x^2-c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-b*x^2 + a)/sqrt(-d*x^2 - c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{a - bx^2}}{\sqrt{-dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a - b*x^2)^(1/2)/(-c - d*x^2)^(1/2),x)

[Out] int((a - b*x^2)^(1/2)/(-c - d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a - bx^2}}{\sqrt{-c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-b*x**2+a)**(1/2)/(-d*x**2-c)**(1/2),x)

[Out] Integral(sqrt(a - b*x**2)/sqrt(-c - d*x**2), x)

$$3.273 \quad \int \frac{\sqrt{-a+bx^2}}{\sqrt{-c-dx^2}} dx$$

Optimal. Leaf size=198

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} (ad + bc) \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), -\frac{ad}{bc}\right)}{\sqrt{b} d \sqrt{bx^2 - a} \sqrt{-c - dx^2}} - \frac{\sqrt{a} \sqrt{b} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{d \sqrt{bx^2 - a} \sqrt{\frac{dx^2}{c} + 1}}$$

[Out] -EllipticE(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*b^(1/2)*(1-b*x^2/a)^(1/2)*(-d*x^2-c)^(1/2)/d/(b*x^2-a)^(1/2)/(1+d*x^2/c)^(1/2)-(a*d+b*c)*EllipticF(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(1+d*x^2/c)^(1/2)/d/b^(1/2)/(b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2)

Rubi [A] time = 0.13, antiderivative size = 198, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {423, 427, 426, 424, 421, 419}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{\frac{dx^2}{c} + 1} (ad + bc) F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} d \sqrt{bx^2 - a} \sqrt{-c - dx^2}} - \frac{\sqrt{a} \sqrt{b} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{d \sqrt{bx^2 - a} \sqrt{\frac{dx^2}{c} + 1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-a + b*x^2]/Sqrt[-c - d*x^2], x]

[Out] -((Sqrt[a]*Sqrt[b]*Sqrt[1 - (b*x^2)/a]*Sqrt[-c - d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(d*Sqrt[-a + b*x^2]*Sqrt[1 + (d*x^2)/c])) - (Sqrt[a]*(b*c + a*d)*Sqrt[1 - (b*x^2)/a]*Sqrt[1 + (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -((a*d)/(b*c))])/(Sqrt[b]*d*Sqrt[-a + b*x^2]*Sqrt[-c - d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0
]
```

Rule 427

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-a+bx^2}}{\sqrt{-c-dx^2}} dx &= -\frac{b \int \frac{\sqrt{-c-dx^2}}{\sqrt{-a+bx^2}} dx}{d} - \frac{(bc+ad) \int \frac{1}{\sqrt{-a+bx^2} \sqrt{-c-dx^2}} dx}{d} \\ &= -\frac{\left(b\sqrt{1-\frac{bx^2}{a}}\right) \int \frac{\sqrt{-c-dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{d\sqrt{-a+bx^2}} - \frac{\left((bc+ad)\sqrt{1+\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{-a+bx^2} \sqrt{1+\frac{dx^2}{c}}} dx}{d\sqrt{-c-dx^2}} \\ &= -\frac{\left(b\sqrt{1-\frac{bx^2}{a}} \sqrt{-c-dx^2}\right) \int \frac{\sqrt{1+\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{d\sqrt{-a+bx^2} \sqrt{1+\frac{dx^2}{c}}} - \frac{\left((bc+ad)\sqrt{1-\frac{bx^2}{a}} \sqrt{1+\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1-\frac{bx^2}{a}} \sqrt{1+\frac{dx^2}{c}}} dx}{d\sqrt{-a+bx^2} \sqrt{-c-dx^2}} \\ &= -\frac{\sqrt{a} \sqrt{b} \sqrt{1-\frac{bx^2}{a}} \sqrt{-c-dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{d\sqrt{-a+bx^2} \sqrt{1+\frac{dx^2}{c}}} - \frac{\sqrt{a}(bc+ad)\sqrt{1-\frac{bx^2}{a}} \sqrt{1+\frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b}d\sqrt{-a+bx^2} \sqrt{-c-dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 93, normalized size = 0.47

$$\frac{\sqrt{bx^2-a} \sqrt{\frac{c+dx^2}{c}} E\left(\sin^{-1}\left(\sqrt{\frac{d}{c}}x\right) \middle| -\frac{bc}{ad}\right)}{\sqrt{\frac{d}{c}} \sqrt{\frac{a-bx^2}{a}} \sqrt{-c-dx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[-a + b*x^2]/Sqrt[-c - d*x^2], x]
```

```
[Out] (Sqrt[-a + b*x^2]*Sqrt[(c + d*x^2)/c]*EllipticE[ArcSin[Sqrt[-(d/c)]*x], -(
b*c)/(a*d)))/(Sqrt[-(d/c)]*Sqrt[(a - b*x^2)/a]*Sqrt[-c - d*x^2])
```

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{bx^2-a} \sqrt{-dx^2-c}}{dx^2+c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(-sqrt(b*x^2 - a)*sqrt(-d*x^2 - c)/(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2-a}}{\sqrt{-dx^2-c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 - a)/sqrt(-d*x^2 - c), x)

maple [A] time = 0.02, size = 167, normalized size = 0.84

$$\frac{\sqrt{bx^2 - a} \sqrt{-dx^2 - c} \sqrt{-\frac{bx^2 - a}{a}} \sqrt{\frac{dx^2 + c}{c}} \left(ad \operatorname{EllipticF} \left(\sqrt{\frac{b}{a}} x, \sqrt{-\frac{ad}{bc}} \right) - bc \operatorname{EllipticE} \left(\sqrt{\frac{b}{a}} x, \sqrt{-\frac{ad}{bc}} \right) + bc \operatorname{EllipticF} \left(\sqrt{\frac{b}{a}} x, \sqrt{-\frac{ad}{bc}} \right) \right)}{(bdx^4 - adx^2 + bcx^2 - ac) \sqrt{\frac{b}{a}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2),x)

[Out] (b*x^2-a)^(1/2)*(-d*x^2-c)^(1/2)*(-(b*x^2-a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*(a*EllipticF((1/a*b)^(1/2)*x, (-a/b/c*d)^(1/2))*d+b*c*EllipticF((1/a*b)^(1/2)*x, (-a/b/c*d)^(1/2))-b*c*EllipticE((1/a*b)^(1/2)*x, (-a/b/c*d)^(1/2)))/(b*d*x^4-a*d*x^2+b*c*x^2-a*c)/(1/a*b)^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{-dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2-a)^(1/2)/(-d*x^2-c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*x^2 - a)/sqrt(-d*x^2 - c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 - a}}{\sqrt{-dx^2 - c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2 - a)^(1/2)/(-c - d*x^2)^(1/2),x)

[Out] int((b*x^2 - a)^(1/2)/(-c - d*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-a + bx^2}}{\sqrt{-c - dx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2-a)**(1/2)/(-d*x**2-c)**(1/2),x)

[Out] Integral(sqrt(-a + b*x**2)/sqrt(-c - d*x**2), x)

$$3.274 \quad \int \frac{\sqrt{c+dx^2}}{\sqrt{a-bx^2}} dx$$

Optimal. Leaf size=87

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c + dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{\frac{dx^2}{c} + 1}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(d*x^2+c)^(1/2)/b^(1/2)/(-b*x^2+a)^(1/2)/(1+d*x^2/c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {427, 426, 424}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c + dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{\frac{dx^2}{c} + 1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/Sqrt[a - b*x^2], x]

[Out] (Sqrt[a]*Sqrt[1 - (b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -(a*d)/(b*c)))/(Sqrt[b]*Sqrt[a - b*x^2]*Sqrt[1 + (d*x^2)/c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\int \frac{\sqrt{c+dx^2}}{\sqrt{a-bx^2}} dx = \frac{\sqrt{1-\frac{bx^2}{a}} \int \frac{\sqrt{c+dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{a-bx^2}}$$

$$= \frac{\left(\sqrt{1-\frac{bx^2}{a}} \sqrt{c+dx^2}\right) \int \frac{\sqrt{1+\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{a-bx^2} \sqrt{1+\frac{dx^2}{c}}}$$

$$= \frac{\sqrt{a} \sqrt{1-\frac{bx^2}{a}} \sqrt{c+dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a-bx^2} \sqrt{1+\frac{dx^2}{c}}}$$

Mathematica [A] time = 0.06, size = 87, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{c+dx^2} E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}}x\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{a-bx^2} \sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + d*x^2]/Sqrt[a - b*x^2], x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[Sqrt[b/a]*x], -((a*d)/(b*c)))]/(Sqrt[b/a]*Sqrt[a - b*x^2]*Sqrt[(c + d*x^2)/c])

fricas [F] time = 0.75, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2+a}\sqrt{dx^2+c}}{bx^2-a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(-b*x^2+a)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(-b*x^2 + a)*sqrt(d*x^2 + c)/(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{-bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(-b*x^2+a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(d*x^2 + c)/sqrt(-b*x^2 + a), x)

maple [A] time = 0.02, size = 106, normalized size = 1.22

$$\frac{\sqrt{dx^2+c} \sqrt{-bx^2+a} \sqrt{-\frac{bx^2-a}{a}} \sqrt{\frac{dx^2+c}{c}} c \text{EllipticE}\left(\sqrt{\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right)}{(-bdx^4 + adx^2 - bcx^2 + ac) \sqrt{\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+c)^(1/2)/(-b*x^2+a)^(1/2),x)`

[Out] `(d*x^2+c)^(1/2)*(-b*x^2+a)^(1/2)*c*(-(b*x^2-a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((1/a*b)^(1/2)*x,(-a/b/c*d)^(1/2))/(-b*d*x^4+a*d*x^2-b*c*x^2+a*c)/(1/a*b)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 + c}}{\sqrt{-bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^(1/2)/(-b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*x^2 + c)/sqrt(-b*x^2 + a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{dx^2 + c}}{\sqrt{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^(1/2)/(a - b*x^2)^(1/2),x)`

[Out] `int((c + d*x^2)^(1/2)/(a - b*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2}}{\sqrt{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**(1/2)/(-b*x**2+a)**(1/2),x)`

[Out] `Integral(sqrt(c + d*x**2)/sqrt(a - b*x**2), x)`

$$3.275 \quad \int \frac{\sqrt{-c-dx^2}}{\sqrt{a-bx^2}} dx$$

Optimal. Leaf size=90

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{\frac{dx^2}{c} + 1}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(-d*x^2-c)^(1/2)/b^(1/2)/(-b*x^2+a)^(1/2)/(1+d*x^2/c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {427, 426, 424}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{\frac{dx^2}{c} + 1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-c - d*x^2]/Sqrt[a - b*x^2], x]

[Out] (Sqrt[a]*Sqrt[1 - (b*x^2)/a]*Sqrt[-c - d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -(a*d)/(b*c))]/(Sqrt[b]*Sqrt[a - b*x^2]*Sqrt[1 + (d*x^2)/c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\int \frac{\sqrt{-c-dx^2}}{\sqrt{a-bx^2}} dx = \frac{\sqrt{1-\frac{bx^2}{a}} \int \frac{\sqrt{-c-dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{a-bx^2}}$$

$$= \frac{\left(\sqrt{1-\frac{bx^2}{a}} \sqrt{-c-dx^2}\right) \int \frac{\sqrt{1+\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{a-bx^2} \sqrt{1+\frac{dx^2}{c}}}$$

$$= \frac{\sqrt{a} \sqrt{1-\frac{bx^2}{a}} \sqrt{-c-dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a-bx^2} \sqrt{1+\frac{dx^2}{c}}}$$

Mathematica [A] time = 0.05, size = 90, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{-c-dx^2} E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}}x\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{a-bx^2} \sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-c - d*x^2]/Sqrt[a - b*x^2], x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[-c - d*x^2]*EllipticE[ArcSin[Sqrt[b/a]*x], -(a*d)/(b*c)))/(Sqrt[b/a]*Sqrt[a - b*x^2]*Sqrt[(c + d*x^2)/c])

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2+a}\sqrt{-dx^2-c}}{bx^2-a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2-c)^(1/2)/(-b*x^2+a)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(-b*x^2 + a)*sqrt(-d*x^2 - c)/(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2-c}}{\sqrt{-bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2-c)^(1/2)/(-b*x^2+a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-d*x^2 - c)/sqrt(-b*x^2 + a), x)

maple [B] time = 0.02, size = 171, normalized size = 1.90

$$\frac{\left(ad \text{EllipticE}\left(\sqrt{-\frac{d}{c}}x, \sqrt{-\frac{bc}{ad}}\right) - ad \text{EllipticF}\left(\sqrt{-\frac{d}{c}}x, \sqrt{-\frac{bc}{ad}}\right) - bc \text{EllipticF}\left(\sqrt{-\frac{d}{c}}x, \sqrt{-\frac{bc}{ad}}\right)\right) \sqrt{-dx^2-c} \sqrt{-bx^2+a}}{\left(bdx^4 - adx^2 + bcx^2 - ac\right) \sqrt{-\frac{d}{c}}b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-d*x^2-c)^(1/2)/(-b*x^2+a)^(1/2),x)`

[Out] `(-a*d*EllipticF((-1/c*d)^(1/2)*x,(-1/a*b*c/d)^(1/2))-c*EllipticF((-1/c*d)^(1/2)*x,(-1/a*b*c/d)^(1/2))*b+a*d*EllipticE((-1/c*d)^(1/2)*x,(-1/a*b*c/d)^(1/2)))*(-d*x^2-c)^(1/2)*(-b*x^2+a)^(1/2)*((d*x^2+c)/c)^(1/2)*(-(b*x^2-a)/a)^(1/2)/(b*d*x^4-a*d*x^2+b*c*x^2-a*c)/(-1/c*d)^(1/2)/b`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2 - c}}{\sqrt{-bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x^2-c)^(1/2)/(-b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-d*x^2 - c)/sqrt(-b*x^2 + a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{-dx^2 - c}}{\sqrt{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-c-d*x^2)^(1/2)/(a-b*x^2)^(1/2),x)`

[Out] `int((-c-d*x^2)^(1/2)/(a-b*x^2)^(1/2),x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c - dx^2}}{\sqrt{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x**2-c)**(1/2)/(-b*x**2+a)**(1/2),x)`

[Out] `Integral(sqrt(-c - d*x**2)/sqrt(a - b*x**2), x)`

$$3.276 \quad \int \frac{\sqrt{c+dx^2}}{\sqrt{-a+bx^2}} dx$$

Optimal. Leaf size=88

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c + dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{bx^2 - a} \sqrt{\frac{dx^2}{c} + 1}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(d*x^2+c)^(1/2)/b^(1/2)/(b*x^2-a)^(1/2)/(1+d*x^2/c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {427, 426, 424}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c + dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{bx^2 - a} \sqrt{\frac{dx^2}{c} + 1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/Sqrt[-a + b*x^2], x]

[Out] (Sqrt[a]*Sqrt[1 - (b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -(a*d)/(b*c)))/(Sqrt[b]*Sqrt[-a + b*x^2]*Sqrt[1 + (d*x^2)/c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\int \frac{\sqrt{c+dx^2}}{\sqrt{-a+bx^2}} dx = \frac{\sqrt{1-\frac{bx^2}{a}} \int \frac{\sqrt{c+dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{-a+bx^2}}$$

$$= \frac{\left(\sqrt{1-\frac{bx^2}{a}} \sqrt{c+dx^2}\right) \int \frac{\sqrt{1+\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{-a+bx^2} \sqrt{1+\frac{dx^2}{c}}}$$

$$= \frac{\sqrt{a} \sqrt{1-\frac{bx^2}{a}} \sqrt{c+dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{-a+bx^2} \sqrt{1+\frac{dx^2}{c}}}$$

Mathematica [A] time = 0.05, size = 88, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{c+dx^2} E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}}x\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{bx^2-a} \sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + d*x^2]/Sqrt[-a + b*x^2], x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[Sqrt[b/a]*x], -((a*d)/(b*c)))]/(Sqrt[b/a]*Sqrt[-a + b*x^2]*Sqrt[(c + d*x^2)/c])

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{dx^2+c}}{\sqrt{bx^2-a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2-a)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(d*x^2 + c)/sqrt(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(b*x^2-a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(d*x^2 + c)/sqrt(b*x^2 - a), x)

maple [B] time = 0.02, size = 168, normalized size = 1.91

$$\frac{\sqrt{dx^2+c} \sqrt{bx^2-a} \sqrt{\frac{dx^2+c}{c}} \sqrt{-\frac{bx^2-a}{a}} \left(-ad \text{EllipticE}\left(\sqrt{-\frac{d}{c}}x, \sqrt{-\frac{bc}{ad}}\right) + ad \text{EllipticF}\left(\sqrt{-\frac{d}{c}}x, \sqrt{-\frac{bc}{ad}}\right) + bc E\right)}{(bdx^4 - adx^2 + bcx^2 - ac) \sqrt{-\frac{d}{c}} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+c)^(1/2)/(b*x^2-a)^(1/2),x)`

[Out] $(d*x^2+c)^{(1/2)}*(b*x^2-a)^{(1/2)}*((d*x^2+c)/c)^{(1/2)}*(-(b*x^2-a)/a)^{(1/2)}*(a*d*\text{EllipticF}((-1/c*d)^{(1/2)}*x, (-1/a*b*c/d)^{(1/2)})+c*\text{EllipticF}((-1/c*d)^{(1/2)}*x, (-1/a*b*c/d)^{(1/2)})*b-a*d*\text{EllipticE}((-1/c*d)^{(1/2)}*x, (-1/a*b*c/d)^{(1/2)}))/((b*d*x^4-a*d*x^2+b*c*x^2-a*c)/(-1/c*d)^{(1/2)}/b)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 + c}}{\sqrt{bx^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^(1/2)/(b*x^2-a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*x^2 + c)/sqrt(b*x^2 - a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{dx^2 + c}}{\sqrt{bx^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c + d*x^2)^(1/2)/(b*x^2 - a)^(1/2),x)`

[Out] `int((c + d*x^2)^(1/2)/(b*x^2 - a)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2}}{\sqrt{-a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**(1/2)/(b*x**2-a)**(1/2),x)`

[Out] `Integral(sqrt(c + d*x**2)/sqrt(-a + b*x**2), x)`

$$3.277 \quad \int \frac{\sqrt{-c-dx^2}}{\sqrt{-a+bx^2}} dx$$

Optimal. Leaf size=91

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{bx^2 - a} \sqrt{\frac{dx^2}{c} + 1}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (-a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(-d*x^2-c)^(1/2)/b^(1/2)/(b*x^2-a)^(1/2)/(1+d*x^2/c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {427, 426, 424}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{-c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{bx^2 - a} \sqrt{\frac{dx^2}{c} + 1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-c - d*x^2]/Sqrt[-a + b*x^2], x]

[Out] (Sqrt[a]*Sqrt[1 - (b*x^2)/a]*Sqrt[-c - d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], -(a*d)/(b*c))]/(Sqrt[b]*Sqrt[-a + b*x^2]*Sqrt[1 + (d*x^2)/c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{-c-dx^2}}{\sqrt{-a+bx^2}} dx &= \frac{\sqrt{1-\frac{bx^2}{a}} \int \frac{\sqrt{-c-dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{-a+bx^2}} \\
&= \frac{\left(\sqrt{1-\frac{bx^2}{a}} \sqrt{-c-dx^2}\right) \int \frac{\sqrt{1+\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{-a+bx^2} \sqrt{1+\frac{dx^2}{c}}} \\
&= \frac{\sqrt{a} \sqrt{1-\frac{bx^2}{a}} \sqrt{-c-dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{b} \sqrt{-a+bx^2} \sqrt{1+\frac{dx^2}{c}}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 91, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{-c-dx^2} E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}}x\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{bx^2-a} \sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-c - d*x^2]/Sqrt[-a + b*x^2], x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[-c - d*x^2]*EllipticE[ArcSin[Sqrt[b/a]*x], -(a*d)/(b*c)))/(Sqrt[b/a]*Sqrt[-a + b*x^2]*Sqrt[(c + d*x^2)/c])

fricas [F] time = 0.47, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-dx^2-c}}{\sqrt{bx^2-a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2-c)^(1/2)/(b*x^2-a)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-d*x^2 - c)/sqrt(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2-c}}{\sqrt{bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2-c)^(1/2)/(b*x^2-a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-d*x^2 - c)/sqrt(b*x^2 - a), x)

maple [A] time = 0.02, size = 110, normalized size = 1.21

$$\frac{\sqrt{-dx^2-c} \sqrt{bx^2-a} \sqrt{-\frac{bx^2-a}{a}} \sqrt{\frac{dx^2+c}{c}} c \text{EllipticE}\left(\sqrt{\frac{b}{a}}x, \sqrt{-\frac{ad}{bc}}\right)}{(bdx^4 - adx^2 + bcx^2 - ac) \sqrt{\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-d*x^2-c)^(1/2)/(b*x^2-a)^(1/2),x)`

[Out] `(-d*x^2-c)^(1/2)*(b*x^2-a)^(1/2)*c*(-(b*x^2-a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((1/a*b)^(1/2)*x,(-a/b/c*d)^(1/2))/(b*d*x^4-a*d*x^2+b*c*x^2-a*c)/(1/a*b)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2 - c}}{\sqrt{bx^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x^2-c)^(1/2)/(b*x^2-a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-d*x^2 - c)/sqrt(b*x^2 - a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{-dx^2 - c}}{\sqrt{bx^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-c-d*x^2)^(1/2)/(b*x^2-a)^(1/2),x)`

[Out] `int((-c-d*x^2)^(1/2)/(b*x^2-a)^(1/2),x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c - dx^2}}{\sqrt{-a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x**2-c)**(1/2)/(b*x**2-a)**(1/2),x)`

[Out] `Integral(sqrt(-c - d*x**2)/sqrt(-a + b*x**2), x)`

$$3.278 \quad \int \frac{\sqrt{c-dx^2}}{\sqrt{a-bx^2}} dx$$

Optimal. Leaf size=88

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{1 - \frac{dx^2}{c}}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(-d*x^2+c)^(1/2)/b^(1/2)/(-b*x^2+a)^(1/2)/(1-d*x^2/c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {427, 426, 424}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{1 - \frac{dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c - d*x^2]/Sqrt[a - b*x^2], x]

[Out] (Sqrt[a]*Sqrt[1 - (b*x^2)/a]*Sqrt[c - d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], (a*d)/(b*c)])/(Sqrt[b]*Sqrt[a - b*x^2]*Sqrt[1 - (d*x^2)/c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c-dx^2}}{\sqrt{a-bx^2}} dx &= \frac{\sqrt{1-\frac{bx^2}{a}} \int \frac{\sqrt{c-dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{a-bx^2}} \\ &= \frac{\left(\sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2}\right) \int \frac{\sqrt{1-\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}}} \\ &= \frac{\sqrt{a} \sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 88, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{c-dx^2} E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}} x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{a-bx^2} \sqrt{\frac{c-dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c - d*x^2]/Sqrt[a - b*x^2], x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[c - d*x^2]*EllipticE[ArcSin[Sqrt[b/a]*x], (a*d)/(b*c)))/(Sqrt[b/a]*Sqrt[a - b*x^2]*Sqrt[(c - d*x^2)/c])

fricas [F] time = 0.81, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2+a}\sqrt{-dx^2+c}}{bx^2-a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2+c)^(1/2)/(-b*x^2+a)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(-b*x^2 + a)*sqrt(-d*x^2 + c)/(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2+c}}{\sqrt{-bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2+c)^(1/2)/(-b*x^2+a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-d*x^2 + c)/sqrt(-b*x^2 + a), x)

maple [B] time = 0.02, size = 164, normalized size = 1.86

$$\frac{\left(ad \text{EllipticE}\left(\sqrt{\frac{d}{c}} x, \sqrt{\frac{bc}{ad}}\right) - ad \text{EllipticF}\left(\sqrt{\frac{d}{c}} x, \sqrt{\frac{bc}{ad}}\right) + bc \text{EllipticF}\left(\sqrt{\frac{d}{c}} x, \sqrt{\frac{bc}{ad}}\right)\right) \sqrt{-dx^2+c} \sqrt{-bx^2+a}}{(bdx^4 - adx^2 - bcx^2 + ac) \sqrt{\frac{d}{c}} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-d*x^2+c)^(1/2)/(-b*x^2+a)^(1/2),x)`

[Out] $(-a*d*\text{EllipticF}((1/c*d)^{1/2}*x, (1/a*b*c/d)^{1/2})+c*\text{EllipticF}((1/c*d)^{1/2})*x, (1/a*b*c/d)^{1/2})*b+a*d*\text{EllipticE}((1/c*d)^{1/2}*x, (1/a*b*c/d)^{1/2}))*(-d*x^2+c)^{1/2}*(-b*x^2+a)^{1/2}*(-(d*x^2-c)/c)^{1/2}*(-(b*x^2-a)/a)^{1/2}/(b*d*x^4-a*d*x^2-b*c*x^2+a*c)/(1/c*d)^{1/2}/b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2 + c}}{\sqrt{-bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x^2+c)^(1/2)/(-b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-d*x^2 + c)/sqrt(-b*x^2 + a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{c - dx^2}}{\sqrt{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c - d*x^2)^(1/2)/(a - b*x^2)^(1/2),x)`

[Out] `int((c - d*x^2)^(1/2)/(a - b*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c - dx^2}}{\sqrt{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x**2+c)**(1/2)/(-b*x**2+a)**(1/2),x)`

[Out] `Integral(sqrt(c - d*x**2)/sqrt(a - b*x**2), x)`

$$3.279 \quad \int \frac{\sqrt{-c+dx^2}}{\sqrt{a-bx^2}} dx$$

Optimal. Leaf size=89

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{dx^2 - c} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{1 - \frac{dx^2}{c}}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(d*x^2-c)^(1/2)/b^(1/2)/(-b*x^2+a)^(1/2)/(1-d*x^2/c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {427, 426, 424}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{dx^2 - c} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a - bx^2} \sqrt{1 - \frac{dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-c + d*x^2]/Sqrt[a - b*x^2], x]

[Out] (Sqrt[a]*Sqrt[1 - (b*x^2)/a]*Sqrt[-c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], (a*d)/(b*c)])/(Sqrt[b]*Sqrt[a - b*x^2]*Sqrt[1 - (d*x^2)/c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\int \frac{\sqrt{-c+dx^2}}{\sqrt{a-bx^2}} dx = \frac{\sqrt{1-\frac{bx^2}{a}} \int \frac{\sqrt{-c+dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{a-bx^2}}$$

$$= \frac{\left(\sqrt{1-\frac{bx^2}{a}} \sqrt{-c+dx^2}\right) \int \frac{\sqrt{1-\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}}}$$

$$= \frac{\sqrt{a} \sqrt{1-\frac{bx^2}{a}} \sqrt{-c+dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{a-bx^2} \sqrt{1-\frac{dx^2}{c}}}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{dx^2-c} E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}}x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{a-bx^2} \sqrt{\frac{c-dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-c + d*x^2]/Sqrt[a - b*x^2], x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[-c + d*x^2]*EllipticE[ArcSin[Sqrt[b/a]*x], (a*d)/(b*c)))/(Sqrt[b/a]*Sqrt[a - b*x^2]*Sqrt[(c - d*x^2)/c])

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2+a}\sqrt{dx^2-c}}{bx^2-a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2-c)^(1/2)/(-b*x^2+a)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(-b*x^2 + a)*sqrt(d*x^2 - c)/(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2-c}}{\sqrt{-bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2-c)^(1/2)/(-b*x^2+a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(d*x^2 - c)/sqrt(-b*x^2 + a), x)

maple [A] time = 0.03, size = 110, normalized size = 1.24

$$\frac{\sqrt{dx^2-c} \sqrt{-bx^2+a} \sqrt{-\frac{bx^2-a}{a}} \sqrt{-\frac{dx^2-c}{c}} c \text{EllipticE}\left(\sqrt{\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right)}{(bdx^4 - adx^2 - bcx^2 + ac) \sqrt{\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2-c)^(1/2)/(-b*x^2+a)^(1/2),x)`

[Out] `(d*x^2-c)^(1/2)*(-b*x^2+a)^(1/2)*c*(-(b*x^2-a)/a)^(1/2)*(-(d*x^2-c)/c)^(1/2)*EllipticE((1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))/(b*d*x^4-a*d*x^2-b*c*x^2+a*c)/(1/a*b)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 - c}}{\sqrt{-bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2-c)^(1/2)/(-b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*x^2 - c)/sqrt(-b*x^2 + a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{dx^2 - c}}{\sqrt{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2 - c)^(1/2)/(a - b*x^2)^(1/2),x)`

[Out] `int((d*x^2 - c)^(1/2)/(a - b*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c + dx^2}}{\sqrt{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2-c)**(1/2)/(-b*x**2+a)**(1/2),x)`

[Out] `Integral(sqrt(-c + d*x**2)/sqrt(a - b*x**2), x)`

$$3.280 \quad \int \frac{\sqrt{c-dx^2}}{\sqrt{-a+bx^2}} dx$$

Optimal. Leaf size=89

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{bx^2 - a} \sqrt{1 - \frac{dx^2}{c}}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(-d*x^2+c)^(1/2)/b^(1/2)/(b*x^2-a)^(1/2)/(1-d*x^2/c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {427, 426, 424}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{c - dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{bx^2 - a} \sqrt{1 - \frac{dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c - d*x^2]/Sqrt[-a + b*x^2],x]

[Out] (Sqrt[a]*Sqrt[1 - (b*x^2)/a]*Sqrt[c - d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], (a*d)/(b*c)])/(Sqrt[b]*Sqrt[-a + b*x^2]*Sqrt[1 - (d*x^2)/c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c-dx^2}}{\sqrt{-a+bx^2}} dx &= \frac{\sqrt{1-\frac{bx^2}{a}} \int \frac{\sqrt{c-dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{-a+bx^2}} \\ &= \frac{\left(\sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2}\right) \int \frac{\sqrt{1-\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{-a+bx^2} \sqrt{1-\frac{dx^2}{c}}} \\ &= \frac{\sqrt{a} \sqrt{1-\frac{bx^2}{a}} \sqrt{c-dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{-a+bx^2} \sqrt{1-\frac{dx^2}{c}}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{c-dx^2} E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}}x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{bx^2-a} \sqrt{\frac{c-dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c - d*x^2]/Sqrt[-a + b*x^2], x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[c - d*x^2]*EllipticE[ArcSin[Sqrt[b/a]*x], (a*d)/(b*c)))/(Sqrt[b/a]*Sqrt[-a + b*x^2]*Sqrt[(c - d*x^2)/c])

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-dx^2+c}}{\sqrt{bx^2-a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2+c)^(1/2)/(b*x^2-a)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-d*x^2 + c)/sqrt(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2+c}}{\sqrt{bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2+c)^(1/2)/(b*x^2-a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-d*x^2 + c)/sqrt(b*x^2 - a), x)

maple [A] time = 0.02, size = 110, normalized size = 1.24

$$\frac{\sqrt{-dx^2+c} \sqrt{bx^2-a} \sqrt{-\frac{bx^2-a}{a}} \sqrt{-\frac{dx^2-c}{c}} c \text{EllipticE}\left(\sqrt{\frac{b}{a}}x, \sqrt{\frac{ad}{bc}}\right)}{(-bdx^4 + adx^2 + bcx^2 - ac) \sqrt{\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-d*x^2+c)^(1/2)/(b*x^2-a)^(1/2),x)`

[Out] $(-d*x^2+c)^{(1/2)}*(b*x^2-a)^{(1/2)}*c*(-(b*x^2-a)/a)^{(1/2)}*(-(d*x^2-c)/c)^{(1/2)}$
 $*\text{EllipticE}((1/a*b)^{(1/2)}*x, (a/b/c*d)^{(1/2)})/(-b*d*x^4+a*d*x^2+b*c*x^2-a*c)$
 $/(1/a*b)^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2 + c}}{\sqrt{bx^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x^2+c)^(1/2)/(b*x^2-a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(-d*x^2 + c)/sqrt(b*x^2 - a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{c - dx^2}}{\sqrt{bx^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c - d*x^2)^(1/2)/(b*x^2 - a)^(1/2),x)`

[Out] `int((c - d*x^2)^(1/2)/(b*x^2 - a)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c - dx^2}}{\sqrt{-a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x**2+c)**(1/2)/(b*x**2-a)**(1/2),x)`

[Out] `Integral(sqrt(c - d*x**2)/sqrt(-a + b*x**2), x)`

$$3.281 \quad \int \frac{\sqrt{-c+dx^2}}{\sqrt{-a+bx^2}} dx$$

Optimal. Leaf size=90

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{dx^2 - c} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{bx^2 - a} \sqrt{1 - \frac{dx^2}{c}}}$$

[Out] EllipticE(x*b^(1/2)/a^(1/2), (a*d/b/c)^(1/2))*a^(1/2)*(1-b*x^2/a)^(1/2)*(d*x^2-c)^(1/2)/b^(1/2)/(b*x^2-a)^(1/2)/(1-d*x^2/c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {427, 426, 424}

$$\frac{\sqrt{a} \sqrt{1 - \frac{bx^2}{a}} \sqrt{dx^2 - c} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{bx^2 - a} \sqrt{1 - \frac{dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-c + d*x^2]/Sqrt[-a + b*x^2], x]

[Out] (Sqrt[a]*Sqrt[1 - (b*x^2)/a]*Sqrt[-c + d*x^2]*EllipticE[ArcSin[(Sqrt[b]*x)/Sqrt[a]], (a*d)/(b*c)])/(Sqrt[b]*Sqrt[-a + b*x^2]*Sqrt[1 - (d*x^2)/c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]

Rule 427

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]

Rubi steps

$$\int \frac{\sqrt{-c+dx^2}}{\sqrt{-a+bx^2}} dx = \frac{\sqrt{1-\frac{bx^2}{a}} \int \frac{\sqrt{-c+dx^2}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{-a+bx^2}}$$

$$= \frac{\left(\sqrt{1-\frac{bx^2}{a}} \sqrt{-c+dx^2}\right) \int \frac{\sqrt{1-\frac{dx^2}{c}}}{\sqrt{1-\frac{bx^2}{a}}} dx}{\sqrt{-a+bx^2} \sqrt{1-\frac{dx^2}{c}}}$$

$$= \frac{\sqrt{a} \sqrt{1-\frac{bx^2}{a}} \sqrt{-c+dx^2} E\left(\sin^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| \frac{ad}{bc}\right)}{\sqrt{b} \sqrt{-a+bx^2} \sqrt{1-\frac{dx^2}{c}}}$$

Mathematica [A] time = 0.04, size = 90, normalized size = 1.00

$$\frac{\sqrt{\frac{a-bx^2}{a}} \sqrt{dx^2-c} E\left(\sin^{-1}\left(\sqrt{\frac{b}{a}}x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{\frac{b}{a}} \sqrt{bx^2-a} \sqrt{\frac{c-dx^2}{c}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-c + d*x^2]/Sqrt[-a + b*x^2], x]

[Out] (Sqrt[(a - b*x^2)/a]*Sqrt[-c + d*x^2]*EllipticE[ArcSin[Sqrt[b/a]*x], (a*d)/(b*c)))/(Sqrt[b/a]*Sqrt[-a + b*x^2]*Sqrt[(c - d*x^2)/c])

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{dx^2-c}}{\sqrt{bx^2-a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2-c)^(1/2)/(b*x^2-a)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(d*x^2 - c)/sqrt(b*x^2 - a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2-c}}{\sqrt{bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2-c)^(1/2)/(b*x^2-a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(d*x^2 - c)/sqrt(b*x^2 - a), x)

maple [B] time = 0.02, size = 167, normalized size = 1.86

$$\frac{\sqrt{dx^2-c} \sqrt{bx^2-a} \sqrt{-\frac{dx^2-c}{c}} \sqrt{-\frac{bx^2-a}{a}} \left(-ad \text{EllipticE}\left(\sqrt{\frac{d}{c}}x, \sqrt{\frac{bc}{ad}}\right) + ad \text{EllipticF}\left(\sqrt{\frac{d}{c}}x, \sqrt{\frac{bc}{ad}}\right) - bc \text{EllipticF}\right)}{(bdx^4 - adx^2 - bcx^2 + ac) \sqrt{\frac{d}{c}} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2-c)^(1/2)/(b*x^2-a)^(1/2),x)`

[Out] $(d*x^2-c)^{(1/2)}*(b*x^2-a)^{(1/2)}*(-(d*x^2-c)/c)^{(1/2)}*(-(b*x^2-a)/a)^{(1/2)}*(a*d*EllipticF((1/c*d)^{(1/2)}*x,(1/a*b*c/d)^{(1/2)})-c*EllipticF((1/c*d)^{(1/2)}*x,(1/a*b*c/d)^{(1/2)})*b-a*d*EllipticE((1/c*d)^{(1/2)}*x,(1/a*b*c/d)^{(1/2)}))/(b*d*x^4-a*d*x^2-b*c*x^2+a*c)/(1/c*d)^{(1/2)}/b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2 - c}}{\sqrt{bx^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2-c)^(1/2)/(b*x^2-a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*x^2 - c)/sqrt(b*x^2 - a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{d x^2 - c}}{\sqrt{b x^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2 - c)^(1/2)/(b*x^2 - a)^(1/2),x)`

[Out] `int((d*x^2 - c)^(1/2)/(b*x^2 - a)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c + dx^2}}{\sqrt{-a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2-c)**(1/2)/(b*x**2-a)**(1/2),x)`

[Out] `Integral(sqrt(-c + d*x**2)/sqrt(-a + b*x**2), x)`

$$3.282 \quad \int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=204

$$\frac{c^{3/2}\sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{dx\sqrt{a+bx^2}}{b\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $d*x*(b*x^2+a)^{(1/2)}/b/(d*x^2+c)^{(1/2)}+c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/a/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)},(1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/b/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.09, antiderivative size = 204, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {422, 418, 492, 411}

$$\frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{dx\sqrt{a+bx^2}}{b\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{c+dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x]

[Out] $(d*x*\operatorname{Sqrt}[a + b*x^2])/(b*\operatorname{Sqrt}[c + d*x^2]) - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2])*\operatorname{EllipticE}(\operatorname{ArcTan}[\operatorname{Sqrt}[d]*x]/\operatorname{Sqrt}[c], 1 - (b*c)/(a*d))/(b*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2]) + (c^{(3/2)}*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}(\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d))/(a*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]*\operatorname{Sqrt}[c + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx &= c \int \frac{1}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx + d \int \frac{x^2}{\sqrt{a+bx^2}\sqrt{c+dx^2}} dx \\ &= \frac{dx\sqrt{a+bx^2}}{b\sqrt{c+dx^2}} + \frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} - \frac{(cd) \int \frac{\sqrt{a+bx^2}}{(c+dx^2)^{3/2}} dx}{b} \\ &= \frac{dx\sqrt{a+bx^2}}{b\sqrt{c+dx^2}} - \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} + \frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 86, normalized size = 0.42

$$\frac{\sqrt{\frac{a+bx^2}{a}}\sqrt{c+dx^2} E\left(\sin^{-1}\left(\sqrt{-\frac{b}{a}}x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{-\frac{b}{a}}\sqrt{a+bx^2}\sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], (a*d)/(b*c)))/(Sqrt[-(b/a)]*Sqrt[a + b*x^2]*Sqrt[(c + d*x^2)/c])
```

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{dx^2+c}}{\sqrt{bx^2+a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(d*x^2 + c)/sqrt(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(d*x^2 + c)/sqrt(b*x^2 + a), x)
```

maple [A] time = 0.02, size = 101, normalized size = 0.50

$$\frac{\sqrt{dx^2+c} \sqrt{bx^2+a} \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} c \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right)}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+c)^(1/2)/(b*x^2+a)^(1/2),x)`

[Out] `(d*x^2+c)^(1/2)*(b*x^2+a)^(1/2)*c*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x,(a/b/c*d)^(1/2))/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+c)^(1/2)/(b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*x^2+c)/sqrt(b*x^2+a),x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c+d*x^2)^(1/2)/(a+b*x^2)^(1/2),x)`

[Out] `int((c+d*x^2)^(1/2)/(a+b*x^2)^(1/2),x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c+dx^2}}{\sqrt{a+bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x**2+c)**(1/2)/(b*x**2+a)**(1/2),x)`

[Out] `Integral(sqrt(c+d*x**2)/sqrt(a+b*x**2),x)`

$$3.283 \quad \int \frac{\sqrt{-c-dx^2}}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=214

$$\frac{c^{3/2}\sqrt{a+bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) dx\sqrt{a+bx^2}}{a\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

[Out] $-d*x*(b*x^2+a)^{(1/2)}/b/(-d*x^2-c)^{(1/2)}-c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*(b*x^2+a)^{(1/2)}/a/d^{(1/2)}/(-d*x^2-c)^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}+(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(b*x^2+a)^{(1/2)}/b/(-d*x^2-c)^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 214, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {422, 418, 492, 411}

$$\frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) dx\sqrt{a+bx^2}}{a\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-c - d*x^2]/Sqrt[a + b*x^2], x]

[Out] $-((d*x*\operatorname{Sqrt}[a + b*x^2])/(b*\operatorname{Sqrt}[-c - d*x^2])) + (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(b*\operatorname{Sqrt}[-c - d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]) - (c^{(3/2)}*\operatorname{Sqrt}[a + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-c - d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-c-dx^2}}{\sqrt{a+bx^2}} dx &= -\left(c \int \frac{1}{\sqrt{a+bx^2}\sqrt{-c-dx^2}} dx\right) - d \int \frac{x^2}{\sqrt{a+bx^2}\sqrt{-c-dx^2}} dx \\ &= -\frac{dx\sqrt{a+bx^2}}{b\sqrt{-c-dx^2}} - \frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{(cd) \int \frac{\sqrt{a+bx^2}}{(-c-dx^2)^{3/2}} dx}{b} \\ &= -\frac{dx\sqrt{a+bx^2}}{b\sqrt{-c-dx^2}} + \frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} - \frac{c^{3/2}\sqrt{a+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 0.42

$$\frac{\sqrt{\frac{a+bx^2}{a}}\sqrt{-c-dx^2} E\left(\sin^{-1}\left(\sqrt{-\frac{b}{a}}x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{-\frac{b}{a}}\sqrt{a+bx^2}\sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[-c - d*x^2]/Sqrt[a + b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[-c - d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], (a*d)/(b*c)))/(Sqrt[-(b/a)]*Sqrt[a + b*x^2]*Sqrt[(c + d*x^2)/c])
```

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-dx^2-c}}{\sqrt{bx^2+a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-d*x^2-c)^(1/2)/(b*x^2+a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(-d*x^2 - c)/sqrt(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2-c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-d*x^2-c)^(1/2)/(b*x^2+a)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(-d*x^2 - c)/sqrt(b*x^2 + a), x)
```

maple [A] time = 0.02, size = 161, normalized size = 0.75

$$\frac{\left(ad \operatorname{EllipticE}\left(\sqrt{-\frac{d}{c}}x, \sqrt{\frac{bc}{ad}}\right) - ad \operatorname{EllipticF}\left(\sqrt{-\frac{d}{c}}x, \sqrt{\frac{bc}{ad}}\right) + bc \operatorname{EllipticF}\left(\sqrt{-\frac{d}{c}}x, \sqrt{\frac{bc}{ad}}\right)\right) \sqrt{-dx^2 - c} \sqrt{bx^2 + a}}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{d}{c}}b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-d*x^2-c)^(1/2)/(b*x^2+a)^(1/2),x)

[Out] (-a*d*EllipticF((-1/c*d)^(1/2)*x, (1/a*b*c/d)^(1/2))+c*EllipticF((-1/c*d)^(1/2)*x, (1/a*b*c/d)^(1/2))*b+a*d*EllipticE((-1/c*d)^(1/2)*x, (1/a*b*c/d)^(1/2)))*(-d*x^2-c)^(1/2)*(b*x^2+a)^(1/2)*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/c*d)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2 - c}}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2-c)^(1/2)/(b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-d*x^2 - c)/sqrt(b*x^2 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{-dx^2 - c}}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-c - d*x^2)^(1/2)/(a + b*x^2)^(1/2),x)

[Out] int((-c - d*x^2)^(1/2)/(a + b*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c - dx^2}}{\sqrt{a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x**2-c)**(1/2)/(b*x**2+a)**(1/2),x)

[Out] Integral(sqrt(-c - d*x**2)/sqrt(a + b*x**2), x)

$$3.284 \quad \int \frac{\sqrt{c+dx^2}}{\sqrt{-a-bx^2}} dx$$

Optimal. Leaf size=214

$$\frac{c^{3/2}\sqrt{-a-bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) dx\sqrt{-a-bx^2}}{a\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{c}\sqrt{d}\sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{c+dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}}$$

[Out] $-d*x*(-b*x^2-a)^{(1/2)}/b/(d*x^2+c)^{(1/2)}-c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*(-b*x^2-a)^{(1/2)}/a/d^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}+(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(-b*x^2-a)^{(1/2)}/b/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}/(d*x^2+c)^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 214, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {422, 418, 492, 411}

$$\frac{c^{3/2}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) dx\sqrt{-a-bx^2}}{a\sqrt{d}\sqrt{c+dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}} + \frac{\sqrt{c}\sqrt{d}\sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{c+dx^2}\sqrt{\frac{c(ax^2+bx^2)}{a(c+dx^2)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + d*x^2]/Sqrt[-a - b*x^2], x]

[Out] $-\left(\frac{d*x*\operatorname{Sqrt}[-a-b*x^2]}{b*\operatorname{Sqrt}[c+d*x^2]}\right) + \left(\frac{\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-a-b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[\operatorname{Sqrt}[d]*x]/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)]}{b*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2))]*\operatorname{Sqrt}[c+d*x^2]} - \frac{c^{(3/2)}*\operatorname{Sqrt}[-a-b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[\operatorname{Sqrt}[d]*x]/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)]}{a*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(c*(a+b*x^2))/(a*(c+d*x^2))]*\operatorname{Sqrt}[c+d*x^2]}\right)$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c+dx^2}}{\sqrt{-a-bx^2}} dx &= c \int \frac{1}{\sqrt{-a-bx^2}\sqrt{c+dx^2}} dx + d \int \frac{x^2}{\sqrt{-a-bx^2}\sqrt{c+dx^2}} dx \\ &= -\frac{dx\sqrt{-a-bx^2}}{b\sqrt{c+dx^2}} - \frac{c^{3/2}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} + \frac{(cd) \int \frac{\sqrt{-a-bx^2}}{(c+dx^2)^{3/2}} dx}{b} \\ &= -\frac{dx\sqrt{-a-bx^2}}{b\sqrt{c+dx^2}} + \frac{\sqrt{c}\sqrt{d}\sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} - \frac{c^{3/2}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}\sqrt{c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 0.42

$$\frac{\sqrt{\frac{a+bx^2}{a}}\sqrt{c+dx^2} E\left(\sin^{-1}\left(\sqrt{-\frac{b}{a}}x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{-\frac{b}{a}}\sqrt{-a-bx^2}\sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[c + d*x^2]/Sqrt[-a - b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[c + d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], (a*d
)/(b*c)))/(Sqrt[-(b/a)]*Sqrt[-a - b*x^2]*Sqrt[(c + d*x^2)/c])
```

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2-a}\sqrt{dx^2+c}}{bx^2+a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^(1/2)/(-b*x^2-a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(-sqrt(-b*x^2 - a)*sqrt(d*x^2 + c)/(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+c)^(1/2)/(-b*x^2-a)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(d*x^2 + c)/sqrt(-b*x^2 - a), x)
```

maple [A] time = 0.02, size = 162, normalized size = 0.76

$$\frac{\sqrt{dx^2+c} \sqrt{-bx^2-a} \sqrt{\frac{dx^2+c}{c}} \sqrt{\frac{bx^2+a}{a}} \left(-ad \operatorname{EllipticE} \left(\sqrt{-\frac{d}{c}} x, \sqrt{\frac{bc}{ad}} \right) + ad \operatorname{EllipticF} \left(\sqrt{-\frac{d}{c}} x, \sqrt{\frac{bc}{ad}} \right) - bc \operatorname{EllipticE} \left(\sqrt{-\frac{d}{c}} x, \sqrt{\frac{bc}{ad}} \right) \right)}{(bdx^4 + adx^2 + bcx^2 + ac) \sqrt{-\frac{d}{c}} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+c)^(1/2)/(-b*x^2-a)^(1/2),x)

[Out] (d*x^2+c)^(1/2)*(-b*x^2-a)^(1/2)*((d*x^2+c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)*(a*d*EllipticF((-1/c*d)^(1/2)*x,(1/a*b*c/d)^(1/2))-c*EllipticF((-1/c*d)^(1/2)*x,(1/a*b*c/d)^(1/2))*b-a*d*EllipticE((-1/c*d)^(1/2)*x,(1/a*b*c/d)^(1/2)))/(b*d*x^4+a*d*x^2+b*c*x^2+a*c)/(-1/c*d)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+c)^(1/2)/(-b*x^2-a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*x^2+c)/sqrt(-b*x^2-a),x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{dx^2+c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c+d*x^2)^(1/2)/(-a-b*x^2)^(1/2),x)

[Out] int((c+d*x^2)^(1/2)/(-a-b*x^2)^(1/2),x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c+dx^2}}{\sqrt{-a-bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+c)**(1/2)/(-b*x**2-a)**(1/2),x)

[Out] Integral(sqrt(c+d*x**2)/sqrt(-a-b*x**2),x)

$$3.285 \quad \int \frac{\sqrt{-c-dx^2}}{\sqrt{-a-bx^2}} dx$$

Optimal. Leaf size=222

$$\frac{c^{3/2}\sqrt{-a-bx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), 1 - \frac{bc}{ad}\right) + dx\sqrt{-a-bx^2}}{a\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(ax^2+b)}{a(cx^2+d)}}} - \frac{\sqrt{c}\sqrt{d}\sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{-c-dx^2}\sqrt{\frac{c(ax^2+b)}{a(cx^2+d)}}}$$

[Out] $d*x*(-b*x^2-a)^{(1/2)}/b/(-d*x^2-c)^{(1/2)}+c^{(3/2)}*(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*(-b*x^2-a)^{(1/2)}/a/d^{(1/2)}/(-d*x^2-c)^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}-(1/(1+d*x^2/c))^{(1/2)}*(1+d*x^2/c)^{(1/2)}*\operatorname{EllipticE}(x*d^{(1/2)}/c^{(1/2)}/(1+d*x^2/c)^{(1/2)}, (1-b*c/a/d)^{(1/2)})*c^{(1/2)}*d^{(1/2)}*(-b*x^2-a)^{(1/2)}/b/(-d*x^2-c)^{(1/2)}/(c*(b*x^2+a)/a/(d*x^2+c))^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 222, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 29, $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$, Rules used = {422, 418, 492, 411}

$$\frac{c^{3/2}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right) + dx\sqrt{-a-bx^2}}{a\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(ax^2+b)}{a(cx^2+d)}}} - \frac{\sqrt{c}\sqrt{d}\sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{-c-dx^2}\sqrt{\frac{c(ax^2+b)}{a(cx^2+d)}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-c - d*x^2]/Sqrt[-a - b*x^2], x]

[Out] $(d*x*\operatorname{Sqrt}[-a - b*x^2])/(b*\operatorname{Sqrt}[-c - d*x^2]) - (\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-a - b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(b*\operatorname{Sqrt}[-c - d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))]) + (c^{(3/2)}*\operatorname{Sqrt}[-a - b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[c]], 1 - (b*c)/(a*d)])/(a*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-c - d*x^2]*\operatorname{Sqrt}[(c*(a + b*x^2))/(a*(c + d*x^2))])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:> Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-c-dx^2}}{\sqrt{-a-bx^2}} dx &= -\left(c \int \frac{1}{\sqrt{-a-bx^2}\sqrt{-c-dx^2}} dx\right) - d \int \frac{x^2}{\sqrt{-a-bx^2}\sqrt{-c-dx^2}} dx \\ &= \frac{dx\sqrt{-a-bx^2}}{b\sqrt{-c-dx^2}} + \frac{c^{3/2}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{(cd) \int \frac{\sqrt{-a-bx^2}}{(-c-dx^2)^{3/2}} dx}{b} \\ &= \frac{dx\sqrt{-a-bx^2}}{b\sqrt{-c-dx^2}} - \frac{\sqrt{c}\sqrt{d}\sqrt{-a-bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{b\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} + \frac{c^{3/2}\sqrt{-a-bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| 1 - \frac{bc}{ad}\right)}{a\sqrt{d}\sqrt{-c-dx^2}\sqrt{\frac{c(a+bx^2)}{a(c+dx^2)}}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 92, normalized size = 0.41

$$\frac{\sqrt{\frac{a+bx^2}{a}}\sqrt{-c-dx^2} E\left(\sin^{-1}\left(\sqrt{-\frac{b}{a}}x\right) \middle| \frac{ad}{bc}\right)}{\sqrt{-\frac{b}{a}}\sqrt{-a-bx^2}\sqrt{\frac{c+dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[-c - d*x^2]/Sqrt[-a - b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[-c - d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], (a*d)/(b*c))]/(Sqrt[-(b/a)]*Sqrt[-a - b*x^2]*Sqrt[(c + d*x^2)/c])
```

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2-a}\sqrt{-dx^2-c}}{bx^2+a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-d*x^2-c)^(1/2)/(-b*x^2-a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(-sqrt(-b*x^2 - a)*sqrt(-d*x^2 - c)/(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2-c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-d*x^2-c)^(1/2)/(-b*x^2-a)^(1/2), x, algorithm="giac")
```

```
[Out] integrate(sqrt(-d*x^2 - c)/sqrt(-b*x^2 - a), x)
```


maple [A] time = 0.02, size = 111, normalized size = 0.50

$$\frac{\sqrt{-dx^2-c} \sqrt{-bx^2-a} \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2+c}{c}} c \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{\frac{ad}{bc}}\right)}{(-bdx^4 - adx^2 - bcx^2 - ac) \sqrt{-\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-d*x^2-c)^(1/2)/(-b*x^2-a)^(1/2), x)`

[Out] `1/(-b*d*x^4-a*d*x^2-b*c*x^2-a*c)/(-1/a*b)^(1/2)*(-d*x^2-c)^(1/2)*(-b*x^2-a)^(1/2)*c*((b*x^2+a)/a)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (a/b/c*d)^(1/2))`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2-c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x^2-c)^(1/2)/(-b*x^2-a)^(1/2), x, algorithm="maxima")`

[Out] `integrate(sqrt(-d*x^2 - c)/sqrt(-b*x^2 - a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{-dx^2-c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((-c-d*x^2)^(1/2)/(-a-b*x^2)^(1/2), x)`

[Out] `int((-c-d*x^2)^(1/2)/(-a-b*x^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c-dx^2}}{\sqrt{-a-bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-d*x**2-c)**(1/2)/(-b*x**2-a)**(1/2), x)`

[Out] `Integral(sqrt(-c - d*x**2)/sqrt(-a - b*x**2), x)`

$$3.286 \quad \int \frac{\sqrt{c-dx^2}}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=189

$$\frac{\sqrt{c} \sqrt{\frac{bx^2}{a} + 1} \sqrt{1 - \frac{dx^2}{c}} (ad + bc) \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{a + bx^2} \sqrt{c - dx^2}} - \frac{\sqrt{c} \sqrt{d} \sqrt{a + bx^2} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{\frac{bx^2}{a} + 1} \sqrt{c - dx^2}}$$

[Out] -EllipticE(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*d^(1/2)*(b*x^2+a)^(1/2)*(1-d*x^2/c)^(1/2)/b/(1+b*x^2/a)^(1/2)/(-d*x^2+c)^(1/2)+(a*d+b*c)*EllipticF(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(1+b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)/b/d^(1/2)/(b*x^2+a)^(1/2)/(-d*x^2+c)^(1/2)

Rubi [A] time = 0.12, antiderivative size = 189, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {423, 427, 426, 424, 421, 419}

$$\frac{\sqrt{c} \sqrt{\frac{bx^2}{a} + 1} \sqrt{1 - \frac{dx^2}{c}} (ad + bc) F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{a + bx^2} \sqrt{c - dx^2}} - \frac{\sqrt{c} \sqrt{d} \sqrt{a + bx^2} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{\frac{bx^2}{a} + 1} \sqrt{c - dx^2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c - d*x^2]/Sqrt[a + b*x^2], x]

[Out] -((Sqrt[c]*Sqrt[d]*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d)])/(b*Sqrt[1 + (b*x^2)/a]*Sqrt[c - d*x^2]) + (Sqrt[c]*(b*c + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d)])/(b*Sqrt[d]*Sqrt[a + b*x^2]*Sqrt[c - d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]
```

Rule 427

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c-dx^2}}{\sqrt{a+bx^2}} dx &= -\frac{d \int \frac{\sqrt{a+bx^2}}{\sqrt{c-dx^2}} dx}{b} + \frac{(bc+ad) \int \frac{1}{\sqrt{a+bx^2}\sqrt{c-dx^2}} dx}{b} \\ &= -\frac{\left(d\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{a+bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{c-dx^2}} + \frac{\left((bc+ad)\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{a+bx^2}\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{c-dx^2}} \\ &= -\frac{\left(d\sqrt{a+bx^2}\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1+\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{1+\frac{bx^2}{a}}\sqrt{c-dx^2}} + \frac{\left((bc+ad)\sqrt{1+\frac{bx^2}{a}}\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1+\frac{bx^2}{a}}\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{a+bx^2}\sqrt{c-dx^2}} \\ &= -\frac{\sqrt{c}\sqrt{d}\sqrt{a+bx^2}\sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{1+\frac{bx^2}{a}}\sqrt{c-dx^2}} + \frac{\sqrt{c}(bc+ad)\sqrt{1+\frac{bx^2}{a}}\sqrt{1-\frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d}\sqrt{a+bx^2}\sqrt{c-dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 89, normalized size = 0.47

$$\frac{\sqrt{\frac{a+bx^2}{a}}\sqrt{c-dx^2} E\left(\sin^{-1}\left(\sqrt{\frac{-b}{a}}x\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{\frac{-b}{a}}\sqrt{a+bx^2}\sqrt{\frac{c-dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[c - d*x^2]/Sqrt[a + b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[c - d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], -((a
*d)/(b*c))])/(Sqrt[-(b/a)]*Sqrt[a + b*x^2]*Sqrt[(c - d*x^2)/c])
```

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-dx^2+c}}{\sqrt{bx^2+a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-d*x^2+c)^(1/2)/(b*x^2+a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(-d*x^2 + c)/sqrt(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2+c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2+c)^(1/2)/(b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-d*x^2 + c)/sqrt(b*x^2 + a), x)

maple [A] time = 0.02, size = 164, normalized size = 0.87

$$\frac{\left(ad \operatorname{EllipticE}\left(\sqrt{\frac{d}{c}} x, \sqrt{-\frac{bc}{ad}}\right) - ad \operatorname{EllipticF}\left(\sqrt{\frac{d}{c}} x, \sqrt{-\frac{bc}{ad}}\right) - bc \operatorname{EllipticF}\left(\sqrt{\frac{d}{c}} x, \sqrt{-\frac{bc}{ad}}\right)\right) \sqrt{-dx^2 + c} \sqrt{bx^2 + a}}{(bdx^4 + adx^2 - bcx^2 - ac) \sqrt{\frac{d}{c}} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-d*x^2+c)^(1/2)/(b*x^2+a)^(1/2),x)

[Out] (-a*d*EllipticF((1/c*d)^(1/2)*x, (-1/a*b*c/d)^(1/2))-c*EllipticF((1/c*d)^(1/2)*x, (-1/a*b*c/d)^(1/2))*b+a*d*EllipticE((1/c*d)^(1/2)*x, (-1/a*b*c/d)^(1/2)))*(-d*x^2+c)^(1/2)*(b*x^2+a)^(1/2)*(-(d*x^2-c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)/(b*d*x^4+a*d*x^2-b*c*x^2-a*c)/(1/c*d)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2 + c}}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2+c)^(1/2)/(b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-d*x^2 + c)/sqrt(b*x^2 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{c - dx^2}}{\sqrt{bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c - d*x^2)^(1/2)/(a + b*x^2)^(1/2),x)

[Out] int((c - d*x^2)^(1/2)/(a + b*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c - dx^2}}{\sqrt{a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x**2+c)**(1/2)/(b*x**2+a)**(1/2),x)

[Out] Integral(sqrt(c - d*x**2)/sqrt(a + b*x**2), x)

$$3.287 \quad \int \frac{\sqrt{-c+dx^2}}{\sqrt{a+bx^2}} dx$$

Optimal. Leaf size=191

$$\frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{\frac{bx^2}{a}+1} \sqrt{dx^2-c}} - \frac{\sqrt{c} \sqrt{\frac{bx^2}{a}+1} \sqrt{1-\frac{dx^2}{c}} (ad+bc) \text{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{a+bx^2} \sqrt{dx^2-c}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*d^(1/2)*(b*x^2+a)^(1/2)*(1-d*x^2/c)^(1/2)/b/(1+b*x^2/a)^(1/2)/(d*x^2-c)^(1/2)-(a*d+b*c)*EllipticF(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(1+b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)/b/d^(1/2)/(b*x^2+a)^(1/2)/(d*x^2-c)^(1/2)

Rubi [A] time = 0.12, antiderivative size = 191, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {423, 427, 426, 424, 421, 419}

$$\frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{\frac{bx^2}{a}+1} \sqrt{dx^2-c}} - \frac{\sqrt{c} \sqrt{\frac{bx^2}{a}+1} \sqrt{1-\frac{dx^2}{c}} (ad+bc) F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{a+bx^2} \sqrt{dx^2-c}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-c + d*x^2]/Sqrt[a + b*x^2], x]

[Out] (Sqrt[c]*Sqrt[d]*Sqrt[a + b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d))]/(b*Sqrt[1 + (b*x^2)/a]*Sqrt[-c + d*x^2]) - (Sqrt[c]*(b*c + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d))]/(b*Sqrt[d]*Sqrt[a + b*x^2]*Sqrt[-c + d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0
]
```

Rule 427

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-c+dx^2}}{\sqrt{a+bx^2}} dx &= \frac{d \int \frac{\sqrt{a+bx^2}}{\sqrt{-c+dx^2}} dx}{b} - \frac{(bc+ad) \int \frac{1}{\sqrt{a+bx^2} \sqrt{-c+dx^2}} dx}{b} \\ &= \frac{\left(d\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{a+bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{-c+dx^2}} - \frac{\left((bc+ad)\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{-c+dx^2}} \\ &= \frac{\left(d\sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1+\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{1+\frac{bx^2}{a}} \sqrt{-c+dx^2}} - \frac{\left((bc+ad)\sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{a+bx^2} \sqrt{-c+dx^2}} \\ &= \frac{\sqrt{c} \sqrt{d} \sqrt{a+bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{1+\frac{bx^2}{a}} \sqrt{-c+dx^2}} - \frac{\sqrt{c} (bc+ad) \sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{a+bx^2} \sqrt{-c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 90, normalized size = 0.47

$$\frac{\sqrt{\frac{a+bx^2}{a}} \sqrt{dx^2-c} E\left(\sin^{-1}\left(\sqrt{\frac{-b}{a}} x\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{\frac{-b}{a}} \sqrt{a+bx^2} \sqrt{\frac{c-dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[-c + d*x^2]/Sqrt[a + b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[-c + d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], -((
a*d)/(b*c))])/(Sqrt[-(b/a)]*Sqrt[a + b*x^2]*Sqrt[(c - d*x^2)/c])
```

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{dx^2-c}}{\sqrt{bx^2+a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2-c)^(1/2)/(b*x^2+a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(d*x^2 - c)/sqrt(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2-c}}{\sqrt{bx^2+a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2-c)^(1/2)/(b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*x^2 - c)/sqrt(b*x^2 + a), x)

maple [A] time = 0.02, size = 109, normalized size = 0.57

$$\frac{\sqrt{d x^2 - c} \sqrt{b x^2 + a} \sqrt{\frac{b x^2 + a}{a}} \sqrt{-\frac{d x^2 - c}{c}} c \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{-\frac{a d}{b c}}\right)}{\left(-b d x^4 - a d x^2 + b c x^2 + a c\right) \sqrt{-\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2-c)^(1/2)/(b*x^2+a)^(1/2),x)

[Out] 1/(-b*d*x^4-a*d*x^2+b*c*x^2+a*c)/(-1/a*b)^(1/2)*(d*x^2-c)^(1/2)*(b*x^2+a)^(1/2)*c*((b*x^2+a)/a)^(1/2)*(-(d*x^2-c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (-a/b/c*d)^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d x^2 - c}}{\sqrt{b x^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2-c)^(1/2)/(b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*x^2 - c)/sqrt(b*x^2 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{d x^2 - c}}{\sqrt{b x^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2 - c)^(1/2)/(a + b*x^2)^(1/2),x)

[Out] int((d*x^2 - c)^(1/2)/(a + b*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c + d x^2}}{\sqrt{a + b x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2-c)**(1/2)/(b*x**2+a)**(1/2),x)

[Out] Integral(sqrt(-c + d*x**2)/sqrt(a + b*x**2), x)

$$3.288 \quad \int \frac{\sqrt{c-dx^2}}{\sqrt{-a-bx^2}} dx$$

Optimal. Leaf size=194

$$\frac{\sqrt{c} \sqrt{\frac{bx^2}{a} + 1} \sqrt{1 - \frac{dx^2}{c}} (ad + bc) \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right), -\frac{bc}{ad}\right) + \sqrt{c} \sqrt{d} \sqrt{-a - bx^2} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{-a - bx^2} \sqrt{c - dx^2} + b\sqrt{\frac{bx^2}{a} + 1} \sqrt{c - dx^2}}$$

[Out] EllipticE(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*d^(1/2)*(-b*x^2-a)^(1/2)*(1-d*x^2/c)^(1/2)/b/(1+b*x^2/a)^(1/2)/(-d*x^2+c)^(1/2)+(a*d+b*c)*EllipticF(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(1+b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)/b/d^(1/2)/(-b*x^2-a)^(1/2)/(-d*x^2+c)^(1/2)

Rubi [A] time = 0.13, antiderivative size = 194, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {423, 427, 426, 424, 421, 419}

$$\frac{\sqrt{c} \sqrt{\frac{bx^2}{a} + 1} \sqrt{1 - \frac{dx^2}{c}} (ad + bc) F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right) + \sqrt{c} \sqrt{d} \sqrt{-a - bx^2} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{-a - bx^2} \sqrt{c - dx^2} + b\sqrt{\frac{bx^2}{a} + 1} \sqrt{c - dx^2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c - d*x^2]/Sqrt[-a - b*x^2], x]

[Out] (Sqrt[c]*Sqrt[d]*Sqrt[-a - b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d))]/(b*Sqrt[1 + (b*x^2)/a]*Sqrt[c - d*x^2]) + (Sqrt[c]*(b*c + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d))]/(b*Sqrt[d]*Sqrt[-a - b*x^2]*Sqrt[c - d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426


```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0]
```

Rule 427

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c-dx^2}}{\sqrt{-a-bx^2}} dx &= \frac{d \int \frac{\sqrt{-a-bx^2}}{\sqrt{c-dx^2}} dx}{b} + \frac{(bc+ad) \int \frac{1}{\sqrt{-a-bx^2} \sqrt{c-dx^2}} dx}{b} \\ &= \frac{\left(d\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{-a-bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{c-dx^2}} + \frac{\left((bc+ad)\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{c-dx^2}} \\ &= \frac{\left(d\sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1+\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{1+\frac{bx^2}{a}} \sqrt{c-dx^2}} + \frac{\left((bc+ad)\sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{-a-bx^2} \sqrt{c-dx^2}} \\ &= \frac{\sqrt{c} \sqrt{d} \sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{1+\frac{bx^2}{a}} \sqrt{c-dx^2}} + \frac{\sqrt{c} (bc+ad) \sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{-a-bx^2} \sqrt{c-dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 92, normalized size = 0.47

$$\frac{\sqrt{\frac{a+bx^2}{a}} \sqrt{c-dx^2} E\left(\sin^{-1}\left(\sqrt{\frac{-b}{a}} x\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{\frac{-b}{a}} \sqrt{-a-bx^2} \sqrt{\frac{c-dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[c - d*x^2]/Sqrt[-a - b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[c - d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], -((a*d)/(b*c))])/(Sqrt[-(b/a)]*Sqrt[-a - b*x^2]*Sqrt[(c - d*x^2)/c])
```

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2-a} \sqrt{-dx^2+c}}{bx^2+a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((-d*x^2+c)^(1/2)/(-b*x^2-a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(-sqrt(-b*x^2 - a)*sqrt(-d*x^2 + c)/(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2+c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2+c)^(1/2)/(-b*x^2-a)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-d*x^2 + c)/sqrt(-b*x^2 - a), x)

maple [A] time = 0.02, size = 111, normalized size = 0.57

$$\frac{\sqrt{-dx^2+c} \sqrt{-bx^2-a} \sqrt{\frac{bx^2+a}{a}} \sqrt{\frac{dx^2-c}{c}} c \operatorname{EllipticE}\left(\sqrt{-\frac{b}{a}} x, \sqrt{-\frac{ad}{bc}}\right)}{(bdx^4 + adx^2 - bcx^2 - ac) \sqrt{-\frac{b}{a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-d*x^2+c)^(1/2)/(-b*x^2-a)^(1/2),x)

[Out] (-d*x^2+c)^(1/2)*(-b*x^2-a)^(1/2)*c*((b*x^2+a)/a)^(1/2)*(-(d*x^2-c)/c)^(1/2)*EllipticE((-1/a*b)^(1/2)*x, (-a/b/c*d)^(1/2))/(b*d*x^4+a*d*x^2-b*c*x^2-a*c)/(-1/a*b)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-dx^2+c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x^2+c)^(1/2)/(-b*x^2-a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-d*x^2 + c)/sqrt(-b*x^2 - a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{c-dx^2}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c - d*x^2)^(1/2)/(-a - b*x^2)^(1/2),x)

[Out] int((c - d*x^2)^(1/2)/(-a - b*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c-dx^2}}{\sqrt{-a-bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-d*x**2+c)**(1/2)/(-b*x**2-a)**(1/2),x)

[Out] Integral(sqrt(c - d*x**2)/sqrt(-a - b*x**2), x)

$$3.289 \quad \int \frac{\sqrt{-c+dx^2}}{\sqrt{-a-bx^2}} dx$$

Optimal. Leaf size=198

$$\frac{\sqrt{c} \sqrt{\frac{bx^2}{a} + 1} \sqrt{1 - \frac{dx^2}{c}} (ad + bc) \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right), -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{-a - bx^2} \sqrt{dx^2 - c}} - \frac{\sqrt{c} \sqrt{d} \sqrt{-a - bx^2} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right)\right)}{b\sqrt{\frac{bx^2}{a} + 1} \sqrt{dx^2 - c}}$$

[Out] -EllipticE(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*d^(1/2)*(-b*x^2-a)^(1/2)*(1-d*x^2/c)^(1/2)/b/(1+b*x^2/a)^(1/2)/(d*x^2-c)^(1/2)-(a*d+b*c)*EllipticF(x*d^(1/2)/c^(1/2), (-b*c/a/d)^(1/2))*c^(1/2)*(1+b*x^2/a)^(1/2)*(1-d*x^2/c)^(1/2)/b/d^(1/2)/(-b*x^2-a)^(1/2)/(d*x^2-c)^(1/2)

Rubi [A] time = 0.13, antiderivative size = 198, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {423, 427, 426, 424, 421, 419}

$$\frac{\sqrt{c} \sqrt{\frac{bx^2}{a} + 1} \sqrt{1 - \frac{dx^2}{c}} (ad + bc) F\left(\sin^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{-a - bx^2} \sqrt{dx^2 - c}} - \frac{\sqrt{c} \sqrt{d} \sqrt{-a - bx^2} \sqrt{1 - \frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{dx}}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{\frac{bx^2}{a} + 1} \sqrt{dx^2 - c}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-c + d*x^2]/Sqrt[-a - b*x^2], x]

[Out] -((Sqrt[c]*Sqrt[d]*Sqrt[-a - b*x^2]*Sqrt[1 - (d*x^2)/c]*EllipticE[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d)])/(b*Sqrt[1 + (b*x^2)/a]*Sqrt[-c + d*x^2])) - (Sqrt[c]*(b*c + a*d)*Sqrt[1 + (b*x^2)/a]*Sqrt[1 - (d*x^2)/c]*EllipticF[ArcSin[(Sqrt[d]*x)/Sqrt[c]], -(b*c)/(a*d)])/(b*Sqrt[d]*Sqrt[-a - b*x^2]*Sqrt[-c + d*x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 426

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[a + b*x^2]/Sqrt[1 + (b*x^2)/a], Int[Sqrt[1 + (b*x^2)/a]/Sqrt[c + d*x^2
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && !GtQ[a, 0
]
```

Rule 427

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[Sqrt[a + b*x^2]/Sqrt[1 + (d*x^2)/c
], x], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && !GtQ[c, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{-c+dx^2}}{\sqrt{-a-bx^2}} dx &= -\frac{d \int \frac{\sqrt{-a-bx^2}}{\sqrt{-c+dx^2}} dx}{b} - \frac{(bc+ad) \int \frac{1}{\sqrt{-a-bx^2} \sqrt{-c+dx^2}} dx}{b} \\ &= -\frac{\left(d\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{-a-bx^2}}{\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{-c+dx^2}} - \frac{\left((bc+ad)\sqrt{1-\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{-c+dx^2}} \\ &= -\frac{\left(d\sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{\sqrt{1+\frac{bx^2}{a}}}{\sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{1+\frac{bx^2}{a}} \sqrt{-c+dx^2}} - \frac{\left((bc+ad)\sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}}\right) \int \frac{1}{\sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}}} dx}{b\sqrt{-a-bx^2} \sqrt{-c+dx^2}} \\ &= -\frac{\sqrt{c} \sqrt{d} \sqrt{-a-bx^2} \sqrt{1-\frac{dx^2}{c}} E\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{1+\frac{bx^2}{a}} \sqrt{-c+dx^2}} - \frac{\sqrt{c}(bc+ad)\sqrt{1+\frac{bx^2}{a}} \sqrt{1-\frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c}}\right) \middle| -\frac{bc}{ad}\right)}{b\sqrt{d} \sqrt{-a-bx^2} \sqrt{-c+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 93, normalized size = 0.47

$$\frac{\sqrt{\frac{a+bx^2}{a}} \sqrt{dx^2-c} E\left(\sin^{-1}\left(\sqrt{\frac{-b}{a}}x\right) \middle| -\frac{ad}{bc}\right)}{\sqrt{\frac{-b}{a}} \sqrt{-a-bx^2} \sqrt{\frac{c-dx^2}{c}}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[-c + d*x^2]/Sqrt[-a - b*x^2], x]
```

```
[Out] (Sqrt[(a + b*x^2)/a]*Sqrt[-c + d*x^2]*EllipticE[ArcSin[Sqrt[-(b/a)]*x], -((a*d)/(b*c))])/(Sqrt[-(b/a)]*Sqrt[-a - b*x^2]*Sqrt[(c - d*x^2)/c])
```

fricas [F] time = 0.71, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-bx^2-a} \sqrt{dx^2-c}}{bx^2+a}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2-c)^(1/2)/(-b*x^2-a)^(1/2), x, algorithm="fricas")
```

```
[Out] integral(-sqrt(-b*x^2 - a)*sqrt(d*x^2 - c)/(b*x^2 + a), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{dx^2-c}}{\sqrt{-bx^2-a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2-c)^(1/2)/(-b*x^2-a)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*x^2 - c)/sqrt(-b*x^2 - a), x)

maple [A] time = 0.02, size = 167, normalized size = 0.84

$$\frac{\sqrt{d x^2 - c} \sqrt{-b x^2 - a} \sqrt{-\frac{d x^2 - c}{c}} \sqrt{\frac{b x^2 + a}{a}} \left(-ad \operatorname{EllipticE} \left(\sqrt{\frac{d}{c}} x, \sqrt{-\frac{bc}{ad}} \right) + ad \operatorname{EllipticF} \left(\sqrt{\frac{d}{c}} x, \sqrt{-\frac{bc}{ad}} \right) + bc \operatorname{EllipticE} \left(\sqrt{\frac{d}{c}} x, \sqrt{-\frac{bc}{ad}} \right) \right)}{(bd x^4 + ad x^2 - bc x^2 - ac) \sqrt{\frac{d}{c}} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2-c)^(1/2)/(-b*x^2-a)^(1/2),x)

[Out] (d*x^2-c)^(1/2)*(-b*x^2-a)^(1/2)*(-(d*x^2-c)/c)^(1/2)*((b*x^2+a)/a)^(1/2)*(a*d*EllipticF((1/c*d)^(1/2)*x, (-1/a*b*c/d)^(1/2))+c*EllipticF((1/c*d)^(1/2)*x, (-1/a*b*c/d)^(1/2))*b-a*d*EllipticE((1/c*d)^(1/2)*x, (-1/a*b*c/d)^(1/2)))/(b*d*x^4+a*d*x^2-b*c*x^2-a*c)/(1/c*d)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d x^2 - c}}{\sqrt{-b x^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2-c)^(1/2)/(-b*x^2-a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*x^2 - c)/sqrt(-b*x^2 - a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{d x^2 - c}}{\sqrt{-b x^2 - a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2 - c)^(1/2)/(- a - b*x^2)^(1/2),x)

[Out] int((d*x^2 - c)^(1/2)/(- a - b*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c + d x^2}}{\sqrt{-a - b x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2-c)**(1/2)/(-b*x**2-a)**(1/2),x)

[Out] Integral(sqrt(-c + d*x**2)/sqrt(-a - b*x**2), x)

$$3.290 \quad \int \frac{1}{\sqrt{2+bx^2} \sqrt{3+dx^2}} dx$$

Optimal. Leaf size=78

$$\frac{\sqrt{bx^2+2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right), 1 - \frac{3b}{2d}\right)}{\sqrt{2} \sqrt{d} \sqrt{dx^2+3} \sqrt{\frac{bx^2+2}{dx^2+3}}}$$

[Out] $1/2*(1/(3*d*x^2+9))^{(1/2)}*(3*d*x^2+9)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}*3^{(1/2)}/(3*d*x^2+9)^{(1/2)}, 1/2*(4-6*b/d)^{(1/2)})*2^{(1/2)}*(b*x^2+2)^{(1/2)}/d^{(1/2)}/((b*x^2+2)/(d*x^2+3))^{(1/2)}/(d*x^2+3)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {418}

$$\frac{\sqrt{bx^2+2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{2} \sqrt{d} \sqrt{dx^2+3} \sqrt{\frac{bx^2+2}{dx^2+3}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[2 + b*x^2]*Sqrt[3 + d*x^2]), x]

[Out] (Sqrt[2 + b*x^2]*EllipticF[ArcTan[(Sqrt[d]*x)/Sqrt[3]], 1 - (3*b)/(2*d)])/(Sqrt[2]*Sqrt[d]*Sqrt[(2 + b*x^2)/(3 + d*x^2)]*Sqrt[3 + d*x^2])

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{2+bx^2} \sqrt{3+dx^2}} dx = \frac{\sqrt{2+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{2} \sqrt{d} \sqrt{\frac{2+bx^2}{3+dx^2}} \sqrt{3+dx^2}}$$

Mathematica [A] time = 0.01, size = 37, normalized size = 0.47

$$\frac{\operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{-b}x}{\sqrt{2}}\right), \frac{2d}{3b}\right)}{\sqrt{3} \sqrt{-b}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[2 + b*x^2]*Sqrt[3 + d*x^2]), x]

[Out] EllipticF[ArcSin[(Sqrt[-b]*x)/Sqrt[2]], (2*d)/(3*b)]/(Sqrt[3]*Sqrt[-b])

fricas [F] time = 0.72, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\frac{\sqrt{bx^2+2} \sqrt{dx^2+3}}{bdx^4 + (3b + 2d)x^2 + 6}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/2)/(d*x^2+3)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + 2)*sqrt(d*x^2 + 3)/(b*d*x^4 + (3*b + 2*d)*x^2 + 6), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2 + 2}\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/2)/(d*x^2+3)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*x^2 + 2)*sqrt(d*x^2 + 3)), x)

maple [A] time = 0.04, size = 38, normalized size = 0.49

$$\frac{\sqrt{2} \operatorname{EllipticF}\left(\frac{\sqrt{3} \sqrt{-d} x}{3}, \frac{\sqrt{2} \sqrt{3} \sqrt{\frac{b}{d}}}{2}\right)}{2\sqrt{-d}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+2)^(1/2)/(d*x^2+3)^(1/2),x)

[Out] 1/2*2^(1/2)*EllipticF(1/3*3^(1/2)*(-d)^(1/2)*x,1/2*2^(1/2)*3^(1/2)*(b/d)^(1/2))/(-d)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2 + 2}\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/2)/(d*x^2+3)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*x^2 + 2)*sqrt(d*x^2 + 3)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{bx^2 + 2}\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((b*x^2 + 2)^(1/2)*(d*x^2 + 3)^(1/2)),x)

[Out] int(1/((b*x^2 + 2)^(1/2)*(d*x^2 + 3)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{bx^2 + 2}\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+2)**(1/2)/(d*x**2+3)**(1/2),x)

[Out] Integral(1/(sqrt(b*x**2 + 2)*sqrt(d*x**2 + 3)), x)

$$3.291 \quad \int \frac{1}{\sqrt{4-x^2} \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=39

$$\frac{\sqrt{\frac{dx^2}{c} + 1} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{x}{2}\right), -\frac{4d}{c}\right)}{\sqrt{c + dx^2}}$$

[Out] EllipticF(1/2*x, 2*(-d/c)^(1/2))*(1+d*x^2/c)^(1/2)/(d*x^2+c)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {421, 419}

$$\frac{\sqrt{\frac{dx^2}{c} + 1} F\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -\frac{4d}{c}\right)}{\sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[4 - x^2]*Sqrt[c + d*x^2]), x]

[Out] (Sqrt[1 + (d*x^2)/c]*EllipticF[ArcSin[x/2], (-4*d)/c])/Sqrt[c + d*x^2]

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{4-x^2} \sqrt{c+dx^2}} dx &= \frac{\sqrt{1 + \frac{dx^2}{c}} \int \frac{1}{\sqrt{4-x^2} \sqrt{1 + \frac{dx^2}{c}}} dx}{\sqrt{c + dx^2}} \\ &= \frac{\sqrt{1 + \frac{dx^2}{c}} F\left(\sin^{-1}\left(\frac{x}{2}\right) \middle| -\frac{4d}{c}\right)}{\sqrt{c + dx^2}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 40, normalized size = 1.03

$$\frac{\sqrt{\frac{c+dx^2}{c}} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{x}{2}\right), -\frac{4d}{c}\right)}{\sqrt{c + dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[4 - x^2]*Sqrt[c + d*x^2]), x]

[Out] (Sqrt[(c + d*x^2)/c]*EllipticF[ArcSin[x/2], (-4*d)/c])/Sqrt[c + d*x^2]

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{dx^2+c}\sqrt{-x^2+4}}{dx^4+(c-4d)x^2-4c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+4)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(d*x^2 + c)*sqrt(-x^2 + 4)/(d*x^4 + (c - 4*d)*x^2 - 4*c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{dx^2+c}\sqrt{-x^2+4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+4)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(d*x^2 + c)*sqrt(-x^2 + 4)), x)

maple [A] time = 0.03, size = 38, normalized size = 0.97

$$\frac{\sqrt{\frac{dx^2+c}{c}} \text{EllipticF}\left(\frac{x}{2}, 2\sqrt{-\frac{d}{c}}\right)}{\sqrt{dx^2+c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+4)^(1/2)/(d*x^2+c)^(1/2),x)

[Out] 1/(d*x^2+c)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF(1/2*x, 2*(-1/c*d)^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{dx^2+c}\sqrt{-x^2+4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+4)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(d*x^2 + c)*sqrt(-x^2 + 4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{4-x^2}\sqrt{dx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((4 - x^2)^(1/2)*(c + d*x^2)^(1/2)),x)

[Out] int(1/((4 - x^2)^(1/2)*(c + d*x^2)^(1/2)), x)

sympy [A] time = 2.38, size = 20, normalized size = 0.51

$$\begin{cases} \frac{F\left(\text{asin}\left(\frac{x}{2}\right)\middle|-\frac{4d}{c}\right)}{\sqrt{c}} & \text{for } x > -2 \wedge x < 2 \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+4)**(1/2)/(d*x**2+c)**(1/2),x)

[Out] Piecewise((elliptic_f(asin(x/2), -4*d/c)/sqrt(c), (x > -2) & (x < 2)))

$$3.292 \quad \int \frac{1}{\sqrt{4+x^2} \sqrt{c+dx^2}} dx$$

Optimal. Leaf size=61

$$\frac{\sqrt{c+dx^2} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{x}{2}\right), 1 - \frac{4d}{c}\right)}{c\sqrt{x^2+4} \sqrt{\frac{c+dx^2}{c(x^2+4)}}}$$

[Out] $(1/(x^2+4))^{(1/2)} * \operatorname{EllipticF}(x/(x^2+4)^{(1/2)}, (1-4*d/c)^{(1/2)}) * (d*x^2+c)^{(1/2)} / c / ((d*x^2+c)/c/(x^2+4))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {418}

$$\frac{\sqrt{c+dx^2} F\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| 1 - \frac{4d}{c}\right)}{c\sqrt{x^2+4} \sqrt{\frac{c+dx^2}{c(x^2+4)}}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[4 + x^2]*Sqrt[c + d*x^2]),x]

[Out] (Sqrt[c + d*x^2]*EllipticF[ArcTan[x/2], 1 - (4*d)/c])/(c*Sqrt[4 + x^2]*Sqrt[(c + d*x^2)/(c*(4 + x^2))])

Rule 418

Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rubi steps

$$\int \frac{1}{\sqrt{4+x^2} \sqrt{c+dx^2}} dx = \frac{\sqrt{c+dx^2} F\left(\tan^{-1}\left(\frac{x}{2}\right) \middle| 1 - \frac{4d}{c}\right)}{c\sqrt{4+x^2} \sqrt{\frac{c+dx^2}{c(4+x^2)}}}$$

Mathematica [C] time = 0.04, size = 47, normalized size = 0.77

$$\frac{i\sqrt{\frac{c+dx^2}{c}} \operatorname{EllipticF}\left(i \sinh^{-1}\left(\frac{x}{2}\right), \frac{4d}{c}\right)}{\sqrt{c+dx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[4 + x^2]*Sqrt[c + d*x^2]),x]

[Out] ((-1)*Sqrt[(c + d*x^2)/c]*EllipticF[I*ArcSinh[x/2], (4*d)/c])/Sqrt[c + d*x^2]

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(\frac{\sqrt{dx^2+c} \sqrt{x^2+4}}{dx^4 + (c+4d)x^2 + 4c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+4)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*x^2 + c)*sqrt(x^2 + 4)/(d*x^4 + (c + 4*d)*x^2 + 4*c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{dx^2 + c} \sqrt{x^2 + 4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+4)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(d*x^2 + c)*sqrt(x^2 + 4)), x)

maple [A] time = 0.03, size = 53, normalized size = 0.87

$$\frac{\sqrt{\frac{dx^2+c}{c}} \operatorname{EllipticF}\left(\sqrt{-\frac{d}{c}} x, \frac{\sqrt{\frac{c}{d}}}{2}\right)}{2\sqrt{d} \sqrt{x^2 + c} \sqrt{-\frac{d}{c}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2+4)^(1/2)/(d*x^2+c)^(1/2),x)

[Out] 1/2/(d*x^2+c)^(1/2)*((d*x^2+c)/c)^(1/2)*EllipticF((-1/c*d)^(1/2)*x,1/2*(c/d)^(1/2))/(-1/c*d)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{dx^2 + c} \sqrt{x^2 + 4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2+4)^(1/2)/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(d*x^2 + c)*sqrt(x^2 + 4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{x^2 + 4} \sqrt{dx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((x^2 + 4)^(1/2)*(c + d*x^2)^(1/2)),x)

[Out] int(1/((x^2 + 4)^(1/2)*(c + d*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c + dx^2} \sqrt{x^2 + 4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x**2+4)**(1/2)/(d*x**2+c)**(1/2),x)

[Out] Integral(1/(sqrt(c + d*x**2)*sqrt(x**2 + 4)), x)

$$3.293 \quad \int \frac{1}{\sqrt{1-x^2} \sqrt{-1+2x^2}} dx$$

Optimal. Leaf size=6

$$-\text{EllipticF}(\cos^{-1}(x), 2)$$

[Out] $-(x^2)^{(1/2)}/x*\text{EllipticF}((-x^2+1)^{(1/2)}, 2^{(1/2)})$

Rubi [A] time = 0.01, antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {420}

$$-F(\cos^{-1}(x)|2)$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[1 - x^2]*Sqrt[-1 + 2*x^2]), x]

[Out] -EllipticF[ArcCos[x], 2]

Rule 420

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> -Simp[EllipticF[ArcCos[Rt[-(d/c), 2]*x], (b*c)/(b*c - a*d)]/(Sqrt[c]*Rt[-(d/c), 2]*Sqrt[a - (b*c)/d]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a - (b*c)/d, 0]

Rubi steps

$$\int \frac{1}{\sqrt{1-x^2} \sqrt{-1+2x^2}} dx = -F(\cos^{-1}(x)|2)$$

Mathematica [B] time = 0.03, size = 27, normalized size = 4.50

$$\frac{\sqrt{1-2x^2} \text{EllipticF}(\sin^{-1}(x), 2)}{\sqrt{2x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[1 - x^2]*Sqrt[-1 + 2*x^2]), x]

[Out] (Sqrt[1 - 2*x^2]*EllipticF[ArcSin[x], 2])/Sqrt[-1 + 2*x^2]

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{2x^2-1}\sqrt{-x^2+1}}{2x^4-3x^2+1}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((-x^2+1)^(1/2))/(2*x^2-1)^(1/2), x, algorithm="fricas")

[Out] integral(-sqrt(2*x^2 - 1)*sqrt(-x^2 + 1)/(2*x^4 - 3*x^2 + 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2-1} \sqrt{-x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(2*x^2-1)^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(2*x^2 - 1)*sqrt(-x^2 + 1)), x)

maple [A] time = 0.03, size = 25, normalized size = 4.17

$$\frac{\sqrt{-2x^2 + 1} \operatorname{EllipticF}\left(x, \sqrt{2}\right)}{\sqrt{2x^2 - 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+1)^(1/2)/(2*x^2-1)^(1/2),x)

[Out] EllipticF(x,2^(1/2))*(-2*x^2+1)^(1/2)/(2*x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{2x^2 - 1} \sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+1)^(1/2)/(2*x^2-1)^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(2*x^2 - 1)*sqrt(-x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.17

$$\int \frac{1}{\sqrt{1 - x^2} \sqrt{2x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1 - x^2)^(1/2)*(2*x^2 - 1)^(1/2)),x)

[Out] int(1/((1 - x^2)^(1/2)*(2*x^2 - 1)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-(x - 1)(x + 1)} \sqrt{2x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+1)**(1/2)/(2*x**2-1)**(1/2),x)

[Out] Integral(1/(sqrt(-(x - 1)*(x + 1))*sqrt(2*x**2 - 1)), x)

$$3.294 \quad \int \frac{\sqrt{1-c^2x^2}}{\sqrt{1+c^2x^2}} dx$$

Optimal. Leaf size=23

$$\frac{2 \operatorname{EllipticF}(\sin^{-1}(cx), -1)}{c} - \frac{E(\sin^{-1}(cx)|-1)}{c}$$

[Out] -EllipticE(c*x,I)/c+2*EllipticF(c*x,I)/c

Rubi [A] time = 0.02, antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {423, 424, 248, 221}

$$\frac{2F(\sin^{-1}(cx)|-1)}{c} - \frac{E(\sin^{-1}(cx)|-1)}{c}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - c^2*x^2]/Sqrt[1 + c^2*x^2], x]

[Out] -(EllipticE[ArcSin[c*x], -1]/c) + (2*EllipticF[ArcSin[c*x], -1])/c

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := Simp[EllipticF[ArcSin[(Rt[-b, 4]*x)/Rt[a, 4]], -1]/(Rt[a, 4]*Rt[-b, 4]), x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]

Rule 248

Int[((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 423

Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{1-c^2x^2}}{\sqrt{1+c^2x^2}} dx &= 2 \int \frac{1}{\sqrt{1-c^2x^2} \sqrt{1+c^2x^2}} dx - \int \frac{\sqrt{1+c^2x^2}}{\sqrt{1-c^2x^2}} dx \\ &= -\frac{E(\sin^{-1}(cx)|-1)}{c} + 2 \int \frac{1}{\sqrt{1-c^4x^4}} dx \\ &= -\frac{E(\sin^{-1}(cx)|-1)}{c} + \frac{2F(\sin^{-1}(cx)|-1)}{c} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 1.04

$$\frac{E\left(\sin^{-1}\left(\sqrt{-c^2}x\right)\middle| -1\right)}{\sqrt{-c^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - c^2*x^2]/Sqrt[1 + c^2*x^2], x]

[Out] EllipticE[ArcSin[Sqrt[-c^2]*x], -1]/Sqrt[-c^2]

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{-c^2x^2+1}}{\sqrt{c^2x^2+1}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-c^2*x^2+1)^(1/2)/(c^2*x^2+1)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(-c^2*x^2 + 1)/sqrt(c^2*x^2 + 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c^2x^2+1}}{\sqrt{c^2x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-c^2*x^2+1)^(1/2)/(c^2*x^2+1)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(-c^2*x^2 + 1)/sqrt(c^2*x^2 + 1), x)

maple [C] time = 0.02, size = 28, normalized size = 1.22

$$\frac{(-\text{EllipticE}(cx \text{csgn}(c), i) + 2 \text{EllipticF}(cx \text{csgn}(c), i)) \text{csgn}(c)}{c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-c^2*x^2+1)^(1/2)/(c^2*x^2+1)^(1/2), x)

[Out] (2*EllipticF(x*csgn(c)*c, I)-EllipticE(x*csgn(c)*c, I))*csgn(c)/c

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-c^2x^2+1}}{\sqrt{c^2x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-c^2*x^2+1)^(1/2)/(c^2*x^2+1)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(-c^2*x^2 + 1)/sqrt(c^2*x^2 + 1), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.04

$$\int \frac{\sqrt{1-c^2x^2}}{\sqrt{c^2x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1 - c^2*x^2)^(1/2)/(c^2*x^2 + 1)^(1/2), x)`

[Out] `int((1 - c^2*x^2)^(1/2)/(c^2*x^2 + 1)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-(cx-1)(cx+1)}}{\sqrt{c^2x^2+1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((-c**2*x**2+1)**(1/2)/(c**2*x**2+1)**(1/2), x)`

[Out] `Integral(sqrt(-(c*x - 1)*(c*x + 1))/sqrt(c**2*x**2 + 1), x)`

$$3.295 \quad \int \frac{\sqrt{2+bx^2}}{\sqrt{3+dx^2}} dx$$

Optimal. Leaf size=182

$$\frac{\sqrt{2}\sqrt{bx^2+2}\operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right),1-\frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}} + \frac{x\sqrt{bx^2+2}}{\sqrt{dx^2+3}} - \frac{\sqrt{2}\sqrt{bx^2+2}E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right)\middle|1-\frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}}$$

[Out] $x*(b*x^2+2)^{(1/2)}/(d*x^2+3)^{(1/2)}-(1/(3*d*x^2+9))^{(1/2)}*(3*d*x^2+9)^{(1/2)}*E$
 $llipticE(x*d^{(1/2)}*3^{(1/2)}/(3*d*x^2+9)^{(1/2)},1/2*(4-6*b/d)^{(1/2)})*2^{(1/2)}*($
 $b*x^2+2)^{(1/2)}/d^{(1/2)}/((b*x^2+2)/(d*x^2+3))^{(1/2)}/(d*x^2+3)^{(1/2)}+(1/(3*d*$
 $x^2+9))^{(1/2)}*(3*d*x^2+9)^{(1/2)}*\operatorname{EllipticF}(x*d^{(1/2)}*3^{(1/2)}/(3*d*x^2+9)^{(1/2)}$
 $),1/2*(4-6*b/d)^{(1/2)})*2^{(1/2)}*(b*x^2+2)^{(1/2)}/d^{(1/2)}/((b*x^2+2)/(d*x^2+3)$
 $)^{(1/2)}/(d*x^2+3)^{(1/2)}$

Rubi [A] time = 0.08, antiderivative size = 182, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {422, 418, 492, 411}

$$\frac{x\sqrt{bx^2+2}}{\sqrt{dx^2+3}} + \frac{\sqrt{2}\sqrt{bx^2+2}F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right)\middle|1-\frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}} - \frac{\sqrt{2}\sqrt{bx^2+2}E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right)\middle|1-\frac{3b}{2d}\right)}{\sqrt{d}\sqrt{dx^2+3}\sqrt{\frac{bx^2+2}{dx^2+3}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[2 + b*x^2]/Sqrt[3 + d*x^2], x]

[Out] $(x*\operatorname{Sqrt}[2 + b*x^2])/ \operatorname{Sqrt}[3 + d*x^2] - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[2 + b*x^2]*\operatorname{EllipticE}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[3]], 1 - (3*b)/(2*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(2 + b*x^2)/(3 + d*x^2)]*\operatorname{Sqrt}[3 + d*x^2]) + (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[2 + b*x^2]*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[d]*x)/\operatorname{Sqrt}[3]], 1 - (3*b)/(2*d)])/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[(2 + b*x^2)/(3 + d*x^2)]*\operatorname{Sqrt}[3 + d*x^2])$

Rule 411

Int[Sqrt[(a_) + (b_.)*(x_)^2]/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/((c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 418

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/((a*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]

Rule 422

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a]

Rule 492

Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(x*Sqrt[a + b*x^2])/((b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a

$+ b*x^2]/(c + d*x^2)^{(3/2)}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{PosQ}[b/a] \ \&\& \ \text{PosQ}[d/c] \ \&\& \ \text{!SimplerSqrtQ}[b/a, d/c]$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{2+bx^2}}{\sqrt{3+dx^2}} dx &= 2 \int \frac{1}{\sqrt{2+bx^2}\sqrt{3+dx^2}} dx + b \int \frac{x^2}{\sqrt{2+bx^2}\sqrt{3+dx^2}} dx \\ &= \frac{x\sqrt{2+bx^2}}{\sqrt{3+dx^2}} + \frac{\sqrt{2}\sqrt{2+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{\frac{2+bx^2}{3+dx^2}}\sqrt{3+dx^2}} - 3 \int \frac{\sqrt{2+bx^2}}{(3+dx^2)^{3/2}} dx \\ &= \frac{x\sqrt{2+bx^2}}{\sqrt{3+dx^2}} - \frac{\sqrt{2}\sqrt{2+bx^2} E\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{\frac{2+bx^2}{3+dx^2}}\sqrt{3+dx^2}} + \frac{\sqrt{2}\sqrt{2+bx^2} F\left(\tan^{-1}\left(\frac{\sqrt{d}x}{\sqrt{3}}\right) \middle| 1 - \frac{3b}{2d}\right)}{\sqrt{d}\sqrt{\frac{2+bx^2}{3+dx^2}}\sqrt{3+dx^2}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 37, normalized size = 0.20

$$\frac{\sqrt{2} E\left(\sin^{-1}\left(\frac{\sqrt{-d}x}{\sqrt{3}}\right) \middle| \frac{3b}{2d}\right)}{\sqrt{-d}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[2 + b*x^2]/Sqrt[3 + d*x^2], x]

[Out] (Sqrt[2]*EllipticE[ArcSin[(Sqrt[-d]*x)/Sqrt[3]], (3*b)/(2*d)])/Sqrt[-d]

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{bx^2+2}}{\sqrt{dx^2+3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+2)^(1/2)/(d*x^2+3)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*x^2 + 2)/sqrt(d*x^2 + 3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2+2}}{\sqrt{dx^2+3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+2)^(1/2)/(d*x^2+3)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*x^2 + 2)/sqrt(d*x^2 + 3), x)

maple [A] time = 0.02, size = 37, normalized size = 0.20

$$\frac{\sqrt{2} \text{EllipticE}\left(\frac{\sqrt{3}\sqrt{-d}x}{3}, \frac{\sqrt{2}\sqrt{3}\sqrt{\frac{b}{d}}}{2}\right)}{\sqrt{-d}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2+2)^(1/2)/(d*x^2+3)^(1/2),x)`

[Out] `EllipticE(1/3*3^(1/2)*(-d)^(1/2)*x,1/2*2^(1/2)*3^(1/2)*(b/d)^(1/2))*2^(1/2)/(-d)^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + 2}}{\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+2)^(1/2)/(d*x^2+3)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*x^2 + 2)/sqrt(d*x^2 + 3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{bx^2 + 2}}{\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*x^2 + 2)^(1/2)/(d*x^2 + 3)^(1/2),x)`

[Out] `int((b*x^2 + 2)^(1/2)/(d*x^2 + 3)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{bx^2 + 2}}{\sqrt{dx^2 + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+2)**(1/2)/(d*x**2+3)**(1/2),x)`

[Out] `Integral(sqrt(b*x**2 + 2)/sqrt(d*x**2 + 3), x)`

$$3.296 \quad \int \frac{\sqrt{-1+3x^2}}{\sqrt{2-3x^2}} dx$$

Optimal. Leaf size=19

$$-\frac{E\left(\cos^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|2\right)}{\sqrt{3}}$$

[Out] $-1/3*(x^2)^{(1/2)}/x*\text{EllipticE}(1/2*(-6*x^2+4)^{(1/2)},2^{(1/2)})*3^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {425}

$$-\frac{E\left(\cos^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|2\right)}{\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[-1 + 3*x^2]/Sqrt[2 - 3*x^2],x]

[Out] -(EllipticE[ArcCos[Sqrt[3/2]*x], 2]/Sqrt[3])

Rule 425

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> -Simp[(Sqrt[a - (b*c)/d]*EllipticE[ArcCos[Rt[-(d/c), 2]*x], (b*c)/(b*c - a*d)])/ (Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a - (b*c)/d, 0]

Rubi steps

$$\int \frac{\sqrt{-1+3x^2}}{\sqrt{2-3x^2}} dx = -\frac{E\left(\cos^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|2\right)}{\sqrt{3}}$$

Mathematica [A] time = 0.03, size = 35, normalized size = 1.84

$$\frac{\sqrt{3x^2-1}E\left(\sin^{-1}\left(\sqrt{\frac{3}{2}}x\right)\middle|2\right)}{\sqrt{3-9x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[-1 + 3*x^2]/Sqrt[2 - 3*x^2],x]

[Out] (Sqrt[-1 + 3*x^2]*EllipticE[ArcSin[Sqrt[3/2]*x], 2])/Sqrt[3 - 9*x^2]

fricas [F] time = 0.45, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{3x^2-1}\sqrt{-3x^2+2}}{3x^2-2},x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3*x^2-1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(3*x^2 - 1)*sqrt(-3*x^2 + 2)/(3*x^2 - 2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{3x^2 - 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3*x^2-1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(3*x^2 - 1)/sqrt(-3*x^2 + 2), x)

maple [A] time = 0.02, size = 37, normalized size = 1.95

$$\frac{\sqrt{-3x^2 + 1} \sqrt{3} \operatorname{EllipticE}\left(\frac{\sqrt{2} \sqrt{3} x}{2}, \sqrt{2}\right)}{3\sqrt{3x^2 - 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*x^2-1)^(1/2)/(-3*x^2+2)^(1/2),x)

[Out] -1/3*EllipticE(1/2*2^(1/2)*3^(1/2)*x,2^(1/2))*(-3*x^2+1)^(1/2)*3^(1/2)/(3*x^2-1)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{3x^2 - 1}}{\sqrt{-3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3*x^2-1)^(1/2)/(-3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(3*x^2 - 1)/sqrt(-3*x^2 + 2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \frac{\sqrt{3x^2 - 1}}{\sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((3*x^2 - 1)^(1/2)/(2 - 3*x^2)^(1/2),x)

[Out] int((3*x^2 - 1)^(1/2)/(2 - 3*x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{3x^2 - 1}}{\sqrt{2 - 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((3*x**2-1)**(1/2)/(-3*x**2+2)**(1/2),x)

[Out] Integral(sqrt(3*x**2 - 1)/sqrt(2 - 3*x**2), x)

$$3.297 \quad \int \frac{\sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 - \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx$$

Optimal. Leaf size=95

$$\frac{\sqrt{\sqrt{b^2 - 4ac} + b} E\left(\sin^{-1}\left(\frac{\sqrt{2}\sqrt{c}x}{\sqrt{b + \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right)}{\sqrt{2}\sqrt{c}}$$

[Out] 1/2*EllipticE(x*2^(1/2)*c^(1/2)/(b+(-4*a*c+b^2)^(1/2))^(1/2),((-b-(-4*a*c+b^2)^(1/2))/(b-(-4*a*c+b^2)^(1/2)))^(1/2))*(b+(-4*a*c+b^2)^(1/2))^(1/2)*2^(1/2)/c^(1/2)

Rubi [A] time = 0.13, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 59, $\frac{\text{number of rules}}{\text{integrand size}} = 0.017$, Rules used = {424}

$$\frac{\sqrt{\sqrt{b^2 - 4ac} + b} E\left(\sin^{-1}\left(\frac{\sqrt{2}\sqrt{c}x}{\sqrt{b + \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right)}{\sqrt{2}\sqrt{c}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]/Sqrt[1 - (2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])], x]

[Out] (Sqrt[b + Sqrt[b^2 - 4*a*c]]*EllipticE[ArcSin[(Sqrt[2]*Sqrt[c]*x)/Sqrt[b + Sqrt[b^2 - 4*a*c]]], -((b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]))]/(Sqrt[2]*Sqrt[c])

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 - \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx = \frac{\sqrt{b + \sqrt{b^2 - 4ac}} E\left(\sin^{-1}\left(\frac{\sqrt{2}\sqrt{c}x}{\sqrt{b + \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right)}{\sqrt{2}\sqrt{c}}$$

Mathematica [A] time = 0.11, size = 95, normalized size = 1.00

$$\frac{\sqrt{\sqrt{b^2 - 4ac} + b} E\left(\sin^{-1}\left(\frac{\sqrt{2}\sqrt{c}x}{\sqrt{b + \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right)}{\sqrt{2}\sqrt{c}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]/Sqrt[1 - (2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])],x]

[Out] (Sqrt[b + Sqrt[b^2 - 4*a*c]]*EllipticE[ArcSin[(Sqrt[2]*Sqrt[c]*x)/Sqrt[b + Sqrt[b^2 - 4*a*c]]], -(b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]))/(Sqrt[2]*Sqrt[c])

fricas [F] time = 0.91, size = 0, normalized size = 0.00

$$\text{integral} \left(-\frac{\left(bx^2 + \sqrt{b^2 - 4ac} x^2 - 2a \right) \sqrt{\frac{bx^2 + \sqrt{b^2 - 4ac} x^2 + 2a}{a}} \sqrt{\frac{bx^2 - \sqrt{b^2 - 4ac} x^2 - 2a}{a}}}{4(cx^4 - bx^2 + a)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1-2*c*x^2/(b+(-4*a*c+b^2)^(1/2)))^(1/2),x, algorithm="fricas")

[Out] integral(-1/4*(b*x^2 + sqrt(b^2 - 4*a*c)*x^2 - 2*a)*sqrt((b*x^2 + sqrt(b^2 - 4*a*c)*x^2 + 2*a)/a)*sqrt(-(b*x^2 - sqrt(b^2 - 4*a*c)*x^2 - 2*a)/a)/(c*x^4 - b*x^2 + a), x)

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1-2*c*x^2/(b+(-4*a*c+b^2)^(1/2)))^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,x):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen & e ,const index_m & i,const vecteur & l) Error: Bad Argument ValueWarning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%}, 0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [71.707969239,70,22]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%}, 0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [78.6493344628,0,0]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%}, 0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [50.5901726987,49,-6]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%}, 0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [91.0141688026,0,0]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%}, 0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [51.2413111906,0,0]Warning, choosing root of [1,0,%%{8,[1,1,0]%%}+%%{-4,[1,0,0]%%}+%%{-2,[0,0,2]%%}, 0,%%{16,[2,2,0]%%}+%%{16,[2,1,0]%%}+%%{4,[2,0,0]%%}+%%{-8,[1,1,2]%%}+%%{-4,[1,0,2]%%}+%%{1,[0,0,4]%%}] at parameters values [-64,2,62]Evaluation time: 11.37

maple [F] time = 0.37, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{2cx^2}{b-\sqrt{-4ac+b^2}} + 1}}{\sqrt{-\frac{2cx^2}{b+\sqrt{-4ac+b^2}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1-2*c*x^2/(b+(-4*a*c+b^2)^(1/2))))^(1/2),x)`

[Out] `int((1+2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1-2*c*x^2/(b+(-4*a*c+b^2)^(1/2))))^(1/2),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{2cx^2}{b-\sqrt{b^2-4ac}} + 1}}{\sqrt{\frac{2cx^2}{b+\sqrt{b^2-4ac}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1-2*c*x^2/(b+(-4*a*c+b^2)^(1/2))))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(2*c*x^2/(b - sqrt(b^2 - 4*a*c)) + 1)/sqrt(-2*c*x^2/(b + sqrt(b^2 - 4*a*c)) + 1), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{2cx^2}{b-\sqrt{b^2-4ac}} + 1}}{\sqrt{1 - \frac{2cx^2}{b+\sqrt{b^2-4ac}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((2*c*x^2)/(b - (b^2 - 4*a*c)^(1/2)) + 1)^(1/2)/(1 - (2*c*x^2)/(b + (b^2 - 4*a*c)^(1/2))))^(1/2),x)`

[Out] `int(((2*c*x^2)/(b - (b^2 - 4*a*c)^(1/2)) + 1)^(1/2)/(1 - (2*c*x^2)/(b + (b^2 - 4*a*c)^(1/2))))^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{b+2cx^2-\sqrt{-4ac+b^2}}{b-\sqrt{-4ac+b^2}}}}{\sqrt{\frac{-b+2cx^2-\sqrt{-4ac+b^2}}{b+\sqrt{-4ac+b^2}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+2*c*x**2/(b-(-4*a*c+b**2)**(1/2)))**1/2/(1-2*c*x**2/(b+(-4*a*c+b**2)**(1/2))))**1/2,x)`

[Out] `Integral(sqrt((b + 2*c*x**2 - sqrt(-4*a*c + b**2))/(b - sqrt(-4*a*c + b**2)))/sqrt(-(-b + 2*c*x**2 - sqrt(-4*a*c + b**2))/(b + sqrt(-4*a*c + b**2))), x)`

$$3.298 \quad \int \frac{\sqrt{1 - \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 - \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx$$

Optimal. Leaf size=94

$$\frac{\sqrt{\sqrt{b^2 - 4ac}} + b E \left(\sin^{-1} \left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b + \sqrt{b^2 - 4ac}}} \right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right)}{\sqrt{2} \sqrt{c}}$$

[Out] $1/2 * \text{EllipticE}(x * 2^{(1/2)} * c^{(1/2)} / (b + (-4 * a * c + b^2)^{(1/2)})^{(1/2)}, ((b + (-4 * a * c + b^2)^{(1/2)}) / (b - (-4 * a * c + b^2)^{(1/2)}))^{(1/2)}) * (b + (-4 * a * c + b^2)^{(1/2)})^{(1/2)} * 2^{(1/2)} / c^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 59, $\frac{\text{number of rules}}{\text{integrand size}} = 0.017$, Rules used = {424}

$$\frac{\sqrt{\sqrt{b^2 - 4ac}} + b E \left(\sin^{-1} \left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b + \sqrt{b^2 - 4ac}}} \right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right)}{\sqrt{2} \sqrt{c}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[1 - (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]/Sqrt[1 - (2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])],x]`

[Out] `(Sqrt[b + Sqrt[b^2 - 4*a*c]]*EllipticE[ArcSin[(Sqrt[2]*Sqrt[c]*x)/Sqrt[b + Sqrt[b^2 - 4*a*c]]], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]))/(Sqrt[2]*Sqrt[c])`

Rule 424

`Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]`

Rubi steps

$$\int \frac{\sqrt{1 - \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 - \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx = \frac{\sqrt{b + \sqrt{b^2 - 4ac}} E \left(\sin^{-1} \left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b + \sqrt{b^2 - 4ac}}} \right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right)}{\sqrt{2} \sqrt{c}}$$

Mathematica [A] time = 0.09, size = 95, normalized size = 1.01

$$\frac{\sqrt{\sqrt{b^2 - 4ac}} + b E \left(\sin^{-1} \left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b + \sqrt{b^2 - 4ac}}} \right) \middle| -\frac{b + \sqrt{b^2 - 4ac}}{\sqrt{b^2 - 4ac} - b} \right)}{\sqrt{2} \sqrt{c}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]/Sqrt[1 - (2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])],x]

[Out] (Sqrt[b + Sqrt[b^2 - 4*a*c]]*EllipticE[ArcSin[(Sqrt[2]*Sqrt[c]*x)/Sqrt[b + Sqrt[b^2 - 4*a*c]]], -((b + Sqrt[b^2 - 4*a*c])/(-b + Sqrt[b^2 - 4*a*c]))]/(Sqrt[2]*Sqrt[c])

fricas [F] time = 0.79, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\left(bx^2 + \sqrt{b^2 - 4ac} x^2 - 2a \right) \sqrt{-\frac{bx^2 + \sqrt{b^2 - 4ac} x^2 - 2a}{a}} \sqrt{-\frac{bx^2 - \sqrt{b^2 - 4ac} x^2 - 2a}{a}}}{4 \left(cx^4 - bx^2 + a \right)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1-2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2))/(1-2*c*x^2/(b+(-4*a*c+b^2)^(1/2))))^(1/2),x, algorithm="fricas")

[Out] integral(-1/4*(b*x^2 + sqrt(b^2 - 4*a*c)*x^2 - 2*a)*sqrt(-(b*x^2 + sqrt(b^2 - 4*a*c)*x^2 - 2*a)/a)*sqrt(-(b*x^2 - sqrt(b^2 - 4*a*c)*x^2 - 2*a)/a)/(c*x^4 - b*x^2 + a), x)

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1-2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2))/(1-2*c*x^2/(b+(-4*a*c+b^2)^(1/2))))^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,x);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen & e ,const index_m & i,const vecteur & l) Error: Bad Argument ValueWarning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [71.707969239,70,22]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [78.6493344628,0,0]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [50.5901726987,49,-6]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [91.0141688026,0,0]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [51.2413111906,0,0]Warning, choosing root of [1,0,%%{8,[1,1,0]%%}+%%{-4,[1,0,0]%%}+%%{-2,[0,0,2]%%},0,%%{16,[2,2,0]%%}+%%{16,[2,1,0]%%}+%%{4,[2,0,0]%%}+%%{-8,[1,1,2]%%}+%%{-4,[1,0,2]%%}+%%{1,[0,0,4]%%}] at parameters values [-64,2,62]Evaluation time: 19.19

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-\frac{2cx^2}{b-\sqrt{-4ac+b^2}} + 1}}{\sqrt{-\frac{2cx^2}{b+\sqrt{-4ac+b^2}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((1-2/(b-(-4*a*c+b^2)^(1/2))*c*x^2)^(1/2)/(-2/(b+(-4*a*c+b^2)^(1/2))*c*x^2+1)^(1/2),x)
```

```
[Out] int((1-2/(b-(-4*a*c+b^2)^(1/2))*c*x^2)^(1/2)/(-2/(b+(-4*a*c+b^2)^(1/2))*c*x^2+1)^(1/2),x)
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{2cx^2}{b-\sqrt{b^2-4ac}} + 1}}{\sqrt{\frac{2cx^2}{b+\sqrt{b^2-4ac}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1-2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1-2*c*x^2/(b+(-4*a*c+b^2)^(1/2)))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(-2*c*x^2/(b - sqrt(b^2 - 4*a*c)) + 1)/sqrt(-2*c*x^2/(b + sqrt(b^2 - 4*a*c)) + 1), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{1 - \frac{2cx^2}{b-\sqrt{b^2-4ac}}}}{\sqrt{1 - \frac{2cx^2}{b+\sqrt{b^2-4ac}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((1 - (2*c*x^2)/(b - (b^2 - 4*a*c)^(1/2)))^(1/2)/(1 - (2*c*x^2)/(b + (b^2 - 4*a*c)^(1/2)))^(1/2),x)
```

```
[Out] int((1 - (2*c*x^2)/(b - (b^2 - 4*a*c)^(1/2)))^(1/2)/(1 - (2*c*x^2)/(b + (b^2 - 4*a*c)^(1/2)))^(1/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{-b+2cx^2+\sqrt{-4ac+b^2}}{b-\sqrt{-4ac+b^2}}}}{\sqrt{\frac{-b+2cx^2-\sqrt{-4ac+b^2}}{b+\sqrt{-4ac+b^2}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1-2*c*x**2/(b-(-4*a*c+b**2)**(1/2)))**1/2/(1-2*c*x**2/(b+(-4*a*c+b**2)**(1/2)))**1/2,x)
```

```
[Out] Integral(sqrt(-(-b + 2*c*x**2 + sqrt(-4*a*c + b**2))/(b - sqrt(-4*a*c + b**2)))/sqrt(-(-b + 2*c*x**2 - sqrt(-4*a*c + b**2))/(b + sqrt(-4*a*c + b**2))), x)
```

$$3.299 \quad \int \frac{\sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx$$

Optimal. Leaf size=478

$$\frac{\sqrt{\sqrt{b^2 - 4ac} + b} \sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1} \operatorname{EllipticF}\left(\tan^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{\sqrt{b^2 - 4ac} + b}}\right), -\frac{2\sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right) x \sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1} \sqrt{\sqrt{b^2 - 4ac} + b}}{\sqrt{2} \sqrt{c} \sqrt{\frac{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1}{\frac{2cx^2}{\sqrt{b^2 - 4ac} + b} + 1}} \sqrt{\frac{2cx^2}{\sqrt{b^2 - 4ac} + b} + 1}}$$

[Out] $x \sqrt{1 + 2cx^2 / (b - (-4ac + b^2)^{1/2})}^{1/2} / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2}))^{1/2} - 1/2 * (1 / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2}))^{1/2}) * \operatorname{EllipticE}(x^2 / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2}))^{1/2}) * c^{1/2} / (b + (-4ac + b^2)^{1/2})^{1/2} / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2}))^{1/2})^{1/2}, (-2 * (-4ac + b^2)^{1/2} / (b - (-4ac + b^2)^{1/2}))^{1/2} * (1 + 2cx^2 / (b - (-4ac + b^2)^{1/2}))^{1/2} * (b + (-4ac + b^2)^{1/2})^{1/2} * 2^{1/2} / c^{1/2} / ((1 + 2cx^2 / (b - (-4ac + b^2)^{1/2})) / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2})))^{1/2} + 1/2 * (1 / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2}))^{1/2}) * \operatorname{EllipticF}(x^2 / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2}))^{1/2}) * c^{1/2} / (b + (-4ac + b^2)^{1/2})^{1/2} / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2}))^{1/2})^{1/2}, (-2 * (-4ac + b^2)^{1/2} / (b - (-4ac + b^2)^{1/2}))^{1/2} * (1 + 2cx^2 / (b - (-4ac + b^2)^{1/2}))^{1/2} * (b + (-4ac + b^2)^{1/2})^{1/2} * 2^{1/2} / c^{1/2} / ((1 + 2cx^2 / (b - (-4ac + b^2)^{1/2})) / (1 + 2cx^2 / (b + (-4ac + b^2)^{1/2})))^{1/2}$

Rubi [A] time = 0.37, antiderivative size = 478, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 59, $\frac{\text{number of rules}}{\text{integrand size}} = 0.068$, Rules used = {422, 418, 492, 411}

$$\frac{x \sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1} \sqrt{\sqrt{b^2 - 4ac} + b} \sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1} F\left(\tan^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b + \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{2\sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right) \sqrt{\sqrt{b^2 - 4ac} + b} \sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1}}{\sqrt{\frac{2cx^2}{\sqrt{b^2 - 4ac} + b} + 1} \sqrt{2} \sqrt{c} \sqrt{\frac{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1}{\frac{2cx^2}{\sqrt{b^2 - 4ac} + b} + 1}} \sqrt{\frac{2cx^2}{\sqrt{b^2 - 4ac} + b} + 1} \sqrt{2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[\frac{\sqrt{1 + (2cx^2)/(b - \sqrt{b^2 - 4ac})}}{\sqrt{1 + (2cx^2)/(b + \sqrt{b^2 - 4ac})}}\right], x$

[Out] $(x \sqrt{1 + (2cx^2)/(b - \sqrt{b^2 - 4ac})}) / \sqrt{1 + (2cx^2)/(b + \sqrt{b^2 - 4ac})} - (\sqrt{b + \sqrt{b^2 - 4ac}} \sqrt{1 + (2cx^2)/(b - \sqrt{b^2 - 4ac})}) * \operatorname{EllipticE}(\operatorname{ArcTan}[(\sqrt{2} \sqrt{c} x) / \sqrt{b + \sqrt{b^2 - 4ac}}]), (-2 \sqrt{b^2 - 4ac}) / (b - \sqrt{b^2 - 4ac})) / (\sqrt{2} \sqrt{c} \sqrt{1 + (2cx^2)/(b - \sqrt{b^2 - 4ac})}) * \sqrt{1 + (2cx^2)/(b + \sqrt{b^2 - 4ac})}) + (\sqrt{b + \sqrt{b^2 - 4ac}} \sqrt{1 + (2cx^2)/(b - \sqrt{b^2 - 4ac})}) * \operatorname{EllipticF}(\operatorname{ArcTan}[(\sqrt{2} \sqrt{c} x) / \sqrt{b + \sqrt{b^2 - 4ac}}]), (-2 \sqrt{b^2 - 4ac}) / (b - \sqrt{b^2 - 4ac})) / (\sqrt{2} \sqrt{c} \sqrt{1 + (2cx^2)/(b - \sqrt{b^2 - 4ac})}) * \sqrt{1 + (2cx^2)/(b + \sqrt{b^2 - 4ac})})$

Rule 411

$\operatorname{Int}[\sqrt{(a_.) + (b_.) * (x_.)^2} / ((c_.) + (d_.) * (x_.)^2)^{3/2}, x_Symbol] := \operatorname{Simp}[(\sqrt{a + bx^2} * \operatorname{EllipticE}(\operatorname{ArcTan}[\operatorname{Rt}[d/c, 2] * x], 1 - (b*c)/(a*d))] / (c * \operatorname{Rt}[d/c, 2] * \sqrt{c + dx^2} * \sqrt{(c * (a + bx^2)) / (a * (c + dx^2))}), x] /; \operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \operatorname{PosQ}[b/a] \ \&\& \operatorname{PosQ}[d/c]$

Rule 418

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(Sqrt[a + b*x^2]*EllipticF[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(a*R
t[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; Fre
eQ[{a, b, c, d}, x] && PosQ[d/c] && PosQ[b/a] && !SimplerSqrtQ[b/a, d/c]
```

Rule 422

```
Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] := Dist[
a, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] + Dist[b, Int[x^2/(Sqrt[
a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c]
&& PosQ[b/a]
```

Rule 492

```
Int[(x_)^2/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol]
:= Simp[(x*Sqrt[a + b*x^2])/(b*Sqrt[c + d*x^2]), x] - Dist[c/b, Int[Sqrt[a
+ b*x^2]/(c + d*x^2)^(3/2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && PosQ[b/a] && PosQ[d/c] && !SimplerSqrtQ[b/a, d/c]
```

Rubi steps

$$\int \frac{\sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx = \frac{(2c) \int \frac{x^2}{\sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx}{b - \sqrt{b^2 - 4ac}} + \int \frac{1}{\sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx$$

$$= \frac{x \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} + \frac{\sqrt{b + \sqrt{b^2 - 4ac}} \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} F\left(\tan^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b + \sqrt{b^2 - 4ac}}}\right)\right) - \frac{2\sqrt{c} x}{b - \sqrt{b^2 - 4ac}}}{\sqrt{2} \sqrt{c} \sqrt{\frac{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} \sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}}$$

$$= \frac{x \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} - \frac{\sqrt{b + \sqrt{b^2 - 4ac}} \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} E\left(\tan^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b + \sqrt{b^2 - 4ac}}}\right)\right) - \frac{2\sqrt{c} x}{b - \sqrt{b^2 - 4ac}}}{\sqrt{2} \sqrt{c} \sqrt{\frac{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} \sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}}$$

Mathematica [A] time = 0.12, size = 102, normalized size = 0.21

$$\frac{\sqrt{-\sqrt{b^2 - 4ac}} - b E\left(\sin^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{-b - \sqrt{b^2 - 4ac}}}\right)\right) \Big|_{\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{2} \sqrt{c}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]/Sqrt[1 + (2*c*x^2)/(b
+ Sqrt[b^2 - 4*a*c])], x]
```

```
[Out] (Sqrt[-b - Sqrt[b^2 - 4*a*c]]*EllipticE[ArcSin[(Sqrt[2]*Sqrt[c]*x)/Sqrt[-b
- Sqrt[b^2 - 4*a*c]]], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])/(S
qrt[2]*Sqrt[c])
```

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\left(bx^2 + \sqrt{b^2 - 4ac} x^2 + 2a \right) \sqrt{\frac{bx^2 + \sqrt{b^2 - 4ac} x^2 + 2a}{a}} \sqrt{\frac{bx^2 - \sqrt{b^2 - 4ac} x^2 + 2a}{a}}}{4(cx^4 + bx^2 + a)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1+2*c*x^2/(b+(-4*a*c+b^2)^(1/2)))^(1/2),x, algorithm="fricas")

[Out] integral(1/4*(b*x^2 + sqrt(b^2 - 4*a*c)*x^2 + 2*a)*sqrt((b*x^2 + sqrt(b^2 - 4*a*c)*x^2 + 2*a)/a)*sqrt((b*x^2 - sqrt(b^2 - 4*a*c)*x^2 + 2*a)/a)/(c*x^4 + b*x^2 + a), x)

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1+2*c*x^2/(b+(-4*a*c+b^2)^(1/2)))^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,x):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen & e ,const index_m & i,const vecteur & l) Error: Bad Argument ValueWarning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [71.707969239,70,22]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [78.6493344628,0,0]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [50.5901726987,49,-6]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [91.0141688026,0,0]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [94.3029591851,0,0]Warning, choosing root of [1,0,%%{8,[1,1,0]%%}+%%{-4,[1,0,0]%%}+%%{-2,[0,0,2]%%},0,%%{16,[2,2,0]%%}+%%{16,[2,1,0]%%}+%%{4,[2,0,0]%%}+%%{-8,[1,1,2]%%}+%%{-4,[1,0,2]%%}+%%{1,[0,0,4]%%}] at parameters values [-64,2,62]Evaluation time: 21.17

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{2cx^2}{b-\sqrt{-4ac+b^2}} + 1}}{\sqrt{\frac{2cx^2}{b+\sqrt{-4ac+b^2}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((2/(b-(-4*a*c+b^2)^(1/2))*c*x^2+1)^(1/2)/(1+2/(b+(-4*a*c+b^2)^(1/2))*c*x^2)^(1/2),x)

[Out] $\int \frac{(2/(b - (-4ac + b^2)^{1/2})) * cx^2 + 1)^{1/2}}{(1 + 2/(b + (-4ac + b^2)^{1/2})) * cx^2)^{1/2}} dx$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1}}{\sqrt{\frac{2cx^2}{b + \sqrt{b^2 - 4ac}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1+2*c*x^2/(b+(-4*a*c+b^2)^(1/2))))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(2*c*x^2/(b - sqrt(b^2 - 4*a*c)) + 1)/sqrt(2*c*x^2/(b + sqrt(b^2 - 4*a*c)) + 1), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}} + 1}}{\sqrt{\frac{2cx^2}{b + \sqrt{b^2 - 4ac}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((2*c*x^2)/(b - (b^2 - 4*a*c)^(1/2)) + 1)^(1/2)/((2*c*x^2)/(b + (b^2 - 4*a*c)^(1/2)) + 1)^(1/2),x)`

[Out] `int(((2*c*x^2)/(b - (b^2 - 4*a*c)^(1/2)) + 1)^(1/2)/((2*c*x^2)/(b + (b^2 - 4*a*c)^(1/2)) + 1)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{b+2cx^2-\sqrt{-4ac+b^2}}{b-\sqrt{-4ac+b^2}}}}{\sqrt{\frac{b+2cx^2+\sqrt{-4ac+b^2}}{b+\sqrt{-4ac+b^2}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+2*c*x**2/(b-(-4*a*c+b**2)**(1/2)))**1/2/(1+2*c*x**2/(b+(-4*a*c+b**2)**(1/2))))**1/2,x)`

[Out] `Integral(sqrt((b + 2*c*x**2 - sqrt(-4*a*c + b**2))/(b - sqrt(-4*a*c + b**2)))/sqrt((b + 2*c*x**2 + sqrt(-4*a*c + b**2))/(b + sqrt(-4*a*c + b**2))), x)`

$$3.300 \quad \int \frac{\sqrt{1 - \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx$$

Optimal. Leaf size=215

$$\frac{\sqrt{2} b \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b - \sqrt{b^2 - 4ac}}}\right), -\frac{b - \sqrt{b^2 - 4ac}}{\sqrt{b^2 - 4ac} + b}\right) \left(\sqrt{b^2 - 4ac} + b\right) E\left(\sin^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b - \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{b - \sqrt{b^2 - 4ac}}{b + \sqrt{b^2 - 4ac}}\right)}{\sqrt{c} \sqrt{b - \sqrt{b^2 - 4ac}} \sqrt{2} \sqrt{c} \sqrt{b - \sqrt{b^2 - 4ac}}}$$

[Out] b*EllipticF(x*2^(1/2)*c^(1/2)/(b-(-4*a*c+b^2)^(1/2))^(1/2), ((-b+(-4*a*c+b^2)^(1/2))/(b+(-4*a*c+b^2)^(1/2)))^(1/2))*2^(1/2)/c^(1/2)/(b-(-4*a*c+b^2)^(1/2))^(1/2)-1/2*EllipticE(x*2^(1/2)*c^(1/2)/(b-(-4*a*c+b^2)^(1/2))^(1/2), ((-b+(-4*a*c+b^2)^(1/2))/(b+(-4*a*c+b^2)^(1/2)))^(1/2))*2^(1/2)/c^(1/2)/(b-(-4*a*c+b^2)^(1/2))^(1/2)

Rubi [A] time = 0.21, antiderivative size = 215, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 59, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {423, 424, 419}

$$\frac{\sqrt{2} b F\left(\sin^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b - \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{b - \sqrt{b^2 - 4ac}}{b + \sqrt{b^2 - 4ac}}\right) \left(\sqrt{b^2 - 4ac} + b\right) E\left(\sin^{-1}\left(\frac{\sqrt{2} \sqrt{c} x}{\sqrt{b - \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{b - \sqrt{b^2 - 4ac}}{b + \sqrt{b^2 - 4ac}}\right)}{\sqrt{c} \sqrt{b - \sqrt{b^2 - 4ac}} \sqrt{2} \sqrt{c} \sqrt{b - \sqrt{b^2 - 4ac}}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 - (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]/Sqrt[1 + (2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])], x]

[Out] -(((b + Sqrt[b^2 - 4*a*c])*EllipticE[ArcSin[(Sqrt[2]*Sqrt[c]*x)/Sqrt[b - Sqrt[b^2 - 4*a*c]]], -((b - Sqrt[b^2 - 4*a*c])/(b + Sqrt[b^2 - 4*a*c])))]/(Sqrt[2]*Sqrt[c]*Sqrt[b - Sqrt[b^2 - 4*a*c]]) + (Sqrt[2]*b*EllipticF[ArcSin[(Sqrt[2]*Sqrt[c]*x)/Sqrt[b - Sqrt[b^2 - 4*a*c]]], -((b - Sqrt[b^2 - 4*a*c])/(b + Sqrt[b^2 - 4*a*c])))]/(Sqrt[c]*Sqrt[b - Sqrt[b^2 - 4*a*c]]))

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplifierSqrtQ[-(b/a), -(d/c)])]

Rule 423

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Dist[b/d, Int[Sqrt[c + d*x^2]/Sqrt[a + b*x^2], x], x] - Dist[(b*c - a*d)/d, Int[1/(Sqrt[a + b*x^2]*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d}, x] && PosQ[d/c] && NegQ[b/a]

Rule 424

Int[Sqrt[(a_) + (b_.)*(x_)^2]/Sqrt[(c_) + (d_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[a]*EllipticE[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)]/(Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rubi steps

$$\int \frac{\sqrt{1 - \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}}{\sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx = \frac{(2b) \int \frac{1}{\sqrt{1 - \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}} dx}{b - \sqrt{b^2 - 4ac}} - \frac{(b + \sqrt{b^2 - 4ac}) \int \frac{\sqrt{1 + \frac{2cx^2}{b + \sqrt{b^2 - 4ac}}}}{\sqrt{1 - \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}}} dx}{b - \sqrt{b^2 - 4ac}}$$

$$= -\frac{(b + \sqrt{b^2 - 4ac}) E\left(\sin^{-1}\left(\frac{\sqrt{2}\sqrt{c}x}{\sqrt{b - \sqrt{b^2 - 4ac}}}\right) \middle| -\frac{b - \sqrt{b^2 - 4ac}}{b + \sqrt{b^2 - 4ac}}\right)}{\sqrt{2}\sqrt{c}\sqrt{b - \sqrt{b^2 - 4ac}}} + \frac{\sqrt{2}bF\left(\sin^{-1}\left(\frac{\sqrt{2}\sqrt{c}x}{\sqrt{b - \sqrt{b^2 - 4ac}}}\right)\right)}{\sqrt{c}\sqrt{b - \sqrt{b^2 - 4ac}}}$$

Mathematica [A] time = 0.09, size = 102, normalized size = 0.47

$$\frac{\sqrt{-\sqrt{b^2 - 4ac} - b} E\left(\sin^{-1}\left(\frac{\sqrt{2}\sqrt{c}x}{\sqrt{-b - \sqrt{b^2 - 4ac}}}\right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{\sqrt{b^2 - 4ac} - b}\right)}{\sqrt{2}\sqrt{c}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 - (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]/Sqrt[1 + (2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])], x]

[Out] (Sqrt[-b - Sqrt[b^2 - 4*a*c]]*EllipticE[ArcSin[(Sqrt[2]*Sqrt[c]*x)/Sqrt[-b - Sqrt[b^2 - 4*a*c]]], (b + Sqrt[b^2 - 4*a*c])/(-b + Sqrt[b^2 - 4*a*c])])/(Sqrt[2]*Sqrt[c])

fricas [F] time = 0.75, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\left(bx^2 + \sqrt{b^2 - 4ac} x^2 + 2a \right) \sqrt{\frac{bx^2 + \sqrt{b^2 - 4ac} x^2 - 2a}{a}} \sqrt{\frac{bx^2 - \sqrt{b^2 - 4ac} x^2 + 2a}{a}}}{4(cx^4 + bx^2 + a)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1-2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1+2*c*x^2/(b+(-4*a*c+b^2)^(1/2)))^(1/2), x, algorithm="fricas")

[Out] integral(1/4*(b*x^2 + sqrt(b^2 - 4*a*c)*x^2 + 2*a)*sqrt(-(b*x^2 + sqrt(b^2 - 4*a*c)*x^2 - 2*a)/a)*sqrt((b*x^2 - sqrt(b^2 - 4*a*c)*x^2 + 2*a)/a)/(c*x^4 + b*x^2 + a), x)

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1-2*c*x^2/(b-(-4*a*c+b^2)^(1/2)))^(1/2)/(1+2*c*x^2/(b+(-4*a*c+b^2)^(1/2)))^(1/2), x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,x):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen & e ,const index_m & i,const vecteur & l) Error: Bad Argument ValueWarning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%}, 0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}

+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [71.707969239,70,22]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [78.6493344628,0,0]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [50.5901726987,49,-6]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [91.0141688026,0,0]Warning, choosing root of [1,0,%%{8,[1,0,1]%%}+%%{-2,[1,0,0]%%}+%%{-2,[0,2,0]%%},0,%%{16,[2,0,2]%%}+%%{8,[2,0,1]%%}+%%{1,[2,0,0]%%}+%%{-8,[1,2,1]%%}+%%{-2,[1,2,0]%%}+%%{1,[0,4,0]%%}] at parameters values [94.3029591851,0,0]Warning, choosing root of [1,0,%%{8,[1,1,0]%%}+%%{-4,[1,0,0]%%}+%%{-2,[0,0,2]%%},0,%%{16,[2,2,0]%%}+%%{16,[2,1,0]%%}+%%{4,[2,0,0]%%}+%%{-8,[1,1,2]%%}+%%{-4,[1,0,2]%%}+%%{1,[0,0,4]%%}] at parameters values [-64,2,62]Evaluation time: 28.88

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-\frac{2cx^2}{b-\sqrt{-4ac+b^2}} + 1}}{\sqrt{\frac{2cx^2}{b+\sqrt{-4ac+b^2}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-2/(b-(-4*a*c+b^2)^(1/2))*c*x^2+1)^(1/2)/(2/(b+(-4*a*c+b^2)^(1/2))*c*x^2+1)^(1/2),x)

[Out] int((-2/(b-(-4*a*c+b^2)^(1/2))*c*x^2+1)^(1/2)/(2/(b+(-4*a*c+b^2)^(1/2))*c*x^2+1)^(1/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-\frac{2cx^2}{b-\sqrt{b^2-4ac}} + 1}}{\sqrt{\frac{2cx^2}{b+\sqrt{b^2-4ac}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1-2*c*x^2/(b-(-4*a*c+b^2)^(1/2))))^(1/2)/((1+2*c*x^2/(b+(-4*a*c+b^2)^(1/2))))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-2*c*x^2/(b - sqrt(b^2 - 4*a*c)) + 1)/sqrt(2*c*x^2/(b + sqrt(b^2 - 4*a*c)) + 1), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{1 - \frac{2cx^2}{b-\sqrt{b^2-4ac}}}}{\sqrt{\frac{2cx^2}{b+\sqrt{b^2-4ac}} + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - (2*c*x^2)/(b - (b^2 - 4*a*c)^(1/2))))^(1/2)/((2*c*x^2)/(b + (b^2 - 4*a*c)^(1/2)) + 1)^(1/2),x)

```
[Out] int((1 - (2*c*x^2)/(b - (b^2 - 4*a*c)^(1/2)))^(1/2)/((2*c*x^2)/(b + (b^2 - 4*a*c)^(1/2)) + 1)^(1/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{-b+2cx^2+\sqrt{-4ac+b^2}}{b-\sqrt{-4ac+b^2}}}}{\sqrt{\frac{b+2cx^2+\sqrt{-4ac+b^2}}{b+\sqrt{-4ac+b^2}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1-2*c*x**2/(b-(-4*a*c+b**2)**(1/2)))**1/2/(1+2*c*x**2/(b+(-4*a*c+b**2)**(1/2)))**1/2,x)
```

```
[Out] Integral(sqrt(-(-b + 2*c*x**2 + sqrt(-4*a*c + b**2))/(b - sqrt(-4*a*c + b**2)))/sqrt((b + 2*c*x**2 + sqrt(-4*a*c + b**2))/(b + sqrt(-4*a*c + b**2))), x)
```

$$3.301 \quad \int \frac{(1-2x^2)^m}{\sqrt{1-x^2}} dx$$

Optimal. Leaf size=62

$$-\frac{2^{-m-2}\sqrt{x^2}(2-4x^2)^{m+1} {}_2F_1\left(\frac{1}{2}, \frac{m+1}{2}; \frac{m+3}{2}; (1-2x^2)^2\right)}{(m+1)x}$$

[Out] $-2^{-(2+m)}(-4x^2+2)^{(1+m)}\text{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2}+\frac{1}{2}m\right], \left[\frac{3}{2}+\frac{1}{2}m\right], (-2x^2+1)^2\right)(x^2)^{(1/2)}/(1+m)/x$

Rubi [C] time = 0.01, antiderivative size = 23, normalized size of antiderivative = 0.37, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {429}

$$xF_1\left(\frac{1}{2}; -m, \frac{1}{2}; \frac{3}{2}; 2x^2, x^2\right)$$

Warning: Unable to verify antiderivative.

[In] Int[(1 - 2*x^2)^m/Sqrt[1 - x^2], x]

[Out] x*AppellF1[1/2, -m, 1/2, 3/2, 2*x^2, x^2]

Rule 429

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 := Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, -((b*x^n)/a), -((d*x^n)/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rubi steps

$$\int \frac{(1-2x^2)^m}{\sqrt{1-x^2}} dx = xF_1\left(\frac{1}{2}; -m, \frac{1}{2}; \frac{3}{2}; 2x^2, x^2\right)$$

Mathematica [C] time = 0.13, size = 122, normalized size = 1.97

$$\frac{3x(1-2x^2)^m F_1\left(\frac{1}{2}; -m, \frac{1}{2}; \frac{3}{2}; 2x^2, x^2\right)}{\sqrt{1-x^2} \left(x^2 \left(F_1\left(\frac{3}{2}; -m, \frac{3}{2}; \frac{5}{2}; 2x^2, x^2\right) - 4mF_1\left(\frac{3}{2}; 1-m, \frac{1}{2}; \frac{5}{2}; 2x^2, x^2\right) \right) + 3F_1\left(\frac{1}{2}; -m, \frac{1}{2}; \frac{3}{2}; 2x^2, x^2\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(1 - 2*x^2)^m/Sqrt[1 - x^2], x]

[Out] (3*x*(1 - 2*x^2)^m*AppellF1[1/2, -m, 1/2, 3/2, 2*x^2, x^2])/(Sqrt[1 - x^2]*(3*AppellF1[1/2, -m, 1/2, 3/2, 2*x^2, x^2] + x^2*(-4*m*AppellF1[3/2, 1 - m, 1/2, 5/2, 2*x^2, x^2] + AppellF1[3/2, -m, 3/2, 5/2, 2*x^2, x^2])))

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(-\frac{\sqrt{-x^2+1}(-2x^2+1)^m}{x^2-1}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-2*x^2+1)^m/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral(-sqrt(-x^2 + 1)*(-2*x^2 + 1)^m/(x^2 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-2x^2 + 1)^m}{\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-2*x^2+1)^m/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate((-2*x^2 + 1)^m/sqrt(-x^2 + 1), x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{(-2x^2 + 1)^m}{\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-2*x^2+1)^m/(-x^2+1)^(1/2),x)

[Out] int((-2*x^2+1)^m/(-x^2+1)^(1/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(-2x^2 + 1)^m}{\sqrt{-x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-2*x^2+1)^m/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate((-2*x^2 + 1)^m/sqrt(-x^2 + 1), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(1 - 2x^2)^m}{\sqrt{1 - x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1 - 2*x^2)^m/(1 - x^2)^(1/2),x)

[Out] int((1 - 2*x^2)^m/(1 - x^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(1 - 2x^2)^m}{\sqrt{-(x - 1)(x + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-2*x**2+1)**m/(-x**2+1)**(1/2),x)

[Out] Integral((1 - 2*x**2)**m/sqrt(-(x - 1)*(x + 1)), x)

$$3.302 \quad \int \frac{1}{\sqrt{-1+x^2} \sqrt{7-4\sqrt{3}+x^2}} dx$$

Optimal. Leaf size=46

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), -7-4\sqrt{3}\right)}{\sqrt{7-4\sqrt{3}} \sqrt{x^2-1}}$$

[Out] EllipticF(x,I*3^(1/2)+2*I)*(-x^2+1)^(1/2)/(x^2-1)^(1/2)/(2-3^(1/2))

Rubi [A] time = 0.04, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {421, 419}

$$\frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \mid -7-4\sqrt{3}\right)}{\sqrt{7-4\sqrt{3}} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[-1 + x^2]*Sqrt[7 - 4*Sqrt[3] + x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], -7 - 4*Sqrt[3]])/(Sqrt[7 - 4*Sqrt[3]]*Sqrt[-1 + x^2])

Rule 419

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Simp[(1*EllipticF[ArcSin[Rt[-(d/c), 2]*x], (b*c)/(a*d)])/(Sqrt[a]*Sqrt[c]*Rt[-(d/c), 2]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0] && !(NegQ[b/a] && SimplerSqrtQ[-(b/a), -(d/c)])

Rule 421

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> Dist[Sqrt[1 + (d*x^2)/c]/Sqrt[c + d*x^2], Int[1/(Sqrt[a + b*x^2]*Sqrt[1 + (d*x^2)/c]), x], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-1+x^2} \sqrt{7-4\sqrt{3}+x^2}} dx &= \frac{\sqrt{1-x^2} \int \frac{1}{\sqrt{1-x^2} \sqrt{7-4\sqrt{3}+x^2}} dx}{\sqrt{-1+x^2}} \\ &= \frac{\sqrt{1-x^2} F\left(\sin^{-1}(x) \mid -7-4\sqrt{3}\right)}{\sqrt{7-4\sqrt{3}} \sqrt{-1+x^2}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 48, normalized size = 1.04

$$\frac{\sqrt{1-x^2} \operatorname{EllipticF}\left(\sin^{-1}(x), \frac{1}{4\sqrt{3}-7}\right)}{\sqrt{7-4\sqrt{3}} \sqrt{x^2-1}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[-1 + x^2]*Sqrt[7 - 4*Sqrt[3] + x^2]),x]

[Out] (Sqrt[1 - x^2]*EllipticF[ArcSin[x], (-7 + 4*Sqrt[3])^(-1)])/(Sqrt[7 - 4*Sqrt[3]*Sqrt[-1 + x^2])

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(x^2 + 4\sqrt{3} + 7)\sqrt{x^2 - 4\sqrt{3} + 7}\sqrt{x^2 - 1}}{x^6 + 13x^4 - 13x^2 - 1}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(7+x^2-4*3^(1/2))^(1/2),x, algorithm="fricas")

[Out] integral((x^2 + 4*sqrt(3) + 7)*sqrt(x^2 - 4*sqrt(3) + 7)*sqrt(x^2 - 1)/(x^6 + 13*x^4 - 13*x^2 - 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 - 4\sqrt{3} + 7}\sqrt{x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(7+x^2-4*3^(1/2))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(x^2 - 4*sqrt(3) + 7)*sqrt(x^2 - 1)), x)

maple [B] time = 0.14, size = 117, normalized size = 2.54

$$\frac{i\sqrt{-x^2+1} \sqrt{-(-x^2+4\sqrt{3}-7)(-4\sqrt{3}+7)} (-2+\sqrt{3}) \sqrt{x^2-1} \sqrt{x^2+7-4\sqrt{3}} \text{EllipticF}\left(\frac{ix}{-2+\sqrt{3}}, 2i-iv\right)}{(4\sqrt{3}-7)(-x^4+4\sqrt{3}x^2-6x^2-4\sqrt{3}+7)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2-1)^(1/2)/(7+x^2-4*3^(1/2))^(1/2),x)

[Out] -I*EllipticF(I*x/(-2+3^(1/2)), 2*I-I*3^(1/2))*(-x^2+1)^(1/2)*(-(-x^2+4*3^(1/2)-7)*(-4*3^(1/2)+7))^(1/2)/(4*3^(1/2)-7)*(-2+3^(1/2))*(x^2-1)^(1/2)*(7+x^2-4*3^(1/2))^(1/2)/(-x^4+4*x^2*3^(1/2)-6*x^2-4*3^(1/2)+7)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2 - 4\sqrt{3} + 7}\sqrt{x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(x^2-1)^(1/2)/(7+x^2-4*3^(1/2))^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(x^2 - 4*sqrt(3) + 7)*sqrt(x^2 - 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{x^2 - 1}\sqrt{x^2 - 4\sqrt{3} + 7}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((x^2 - 1)^(1/2)*(x^2 - 4*3^(1/2) + 7)^(1/2)),x)`

[Out] `int(1/((x^2 - 1)^(1/2)*(x^2 - 4*3^(1/2) + 7)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{(x-1)(x+1)}\sqrt{x^2 - 4\sqrt{3} + 7}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(x**2-1)**(1/2)/(7+x**2-4*3**(1/2))**(1/2),x)`

[Out] `Integral(1/(sqrt((x - 1)*(x + 1))*sqrt(x**2 - 4*sqrt(3) + 7)), x)`

$$3.303 \quad \int \frac{1}{\sqrt{3-3\sqrt{3}+2\sqrt{3}x^2} \sqrt{3+(-3+\sqrt{3})x^2}} dx$$

Optimal. Leaf size=47

$$-\frac{1}{6}\sqrt{3+\sqrt{3}} \operatorname{EllipticF}\left(\cos^{-1}\left(\sqrt{\frac{1}{3}(3-\sqrt{3})}x\right), \frac{1}{2}(1+\sqrt{3})\right)$$

[Out] $-1/6*(x^2*(9-3*3^{(1/2)}))^{(1/2)}/x/(9-3*3^{(1/2)})^{(1/2)}*\operatorname{EllipticF}(1/3*(9-x^2*(9-3*3^{(1/2)}))^{(1/2)}, 1/2*(2+2*3^{(1/2)})^{(1/2)}*(3+3^{(1/2)})^{(1/2)})$

Rubi [A] time = 0.06, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 41, $\frac{\text{number of rules}}{\text{integrand size}} = 0.024$, Rules used = {420}

$$-\frac{1}{6}\sqrt{3+\sqrt{3}} F\left(\cos^{-1}\left(\sqrt{\frac{1}{3}(3-\sqrt{3})}x\right)\middle|\frac{1}{2}(1+\sqrt{3})\right)$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[3 - 3*Sqrt[3] + 2*Sqrt[3]*x^2]*Sqrt[3 + (-3 + Sqrt[3])*x^2]), x]

[Out] $-(\operatorname{Sqrt}[3 + \operatorname{Sqrt}[3]]*\operatorname{EllipticF}[\operatorname{ArcCos}[\operatorname{Sqrt}[(3 - \operatorname{Sqrt}[3])/3]*x], (1 + \operatorname{Sqrt}[3])/2])/6$

Rule 420

Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] :> -Simp[EllipticF[ArcCos[Rt[-(d/c), 2]*x], (b*c)/(b*c - a*d)]/(Sqrt[c]*Rt[-(d/c), 2]*Sqrt[a - (b*c)/d]), x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a - (b*c)/d, 0]

Rubi steps

$$\int \frac{1}{\sqrt{3-3\sqrt{3}+2\sqrt{3}x^2} \sqrt{3+(-3+\sqrt{3})x^2}} dx = -\frac{1}{6}\sqrt{3+\sqrt{3}} F\left(\cos^{-1}\left(\sqrt{\frac{1}{3}(3-\sqrt{3})}x\right)\middle|\frac{1}{2}(1+\sqrt{3})\right)$$

Mathematica [A] time = 0.15, size = 81, normalized size = 1.72

$$\frac{\sqrt{-2\sqrt{3}x^2+3\sqrt{3}-3} \operatorname{EllipticF}\left(\sin^{-1}\left(\frac{\sqrt{1+\sqrt{3}}x}{\sqrt[4]{3}}\right), 2-\sqrt{3}\right)}{3^{3/4}\sqrt{4\sqrt{3}x^2-6\sqrt{3}+6}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(Sqrt[3 - 3*Sqrt[3] + 2*Sqrt[3]*x^2]*Sqrt[3 + (-3 + Sqrt[3])*x^2]), x]

[Out] $(\operatorname{Sqrt}[-3 + 3*\operatorname{Sqrt}[3] - 2*\operatorname{Sqrt}[3]*x^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[(\operatorname{Sqrt}[1 + \operatorname{Sqrt}[3]]*x)/3^{(1/4)}], 2 - \operatorname{Sqrt}[3]])/(3^{(3/4)}*\operatorname{Sqrt}[6 - 6*\operatorname{Sqrt}[3] + 4*\operatorname{Sqrt}[3]*x^2])$

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\operatorname{integral}\left(-\frac{\sqrt{\sqrt{3}x^2-3x^2+3}\sqrt{\sqrt{3}(2x^2-3)+3(\sqrt{3}+1)}}{6(2x^4-6x^2+3)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(3+x^2*(-3+3^(1/2)))^(1/2)/(3-3*3^(1/2)+2*3^(1/2)*x^2)^(1/2),x,
algorithm="fricas")
```

```
[Out] integral(-1/6*sqrt(sqrt(3)*x^2 - 3*x^2 + 3)*sqrt(sqrt(3)*(2*x^2 - 3) + 3)*
sqrt(3) + 1)/(2*x^4 - 6*x^2 + 3), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2(\sqrt{3}-3)+3}\sqrt{2\sqrt{3}x^2-3\sqrt{3}+3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(3+x^2*(-3+3^(1/2)))^(1/2)/(3-3*3^(1/2)+2*3^(1/2)*x^2)^(1/2),x,
algorithm="giac")
```

```
[Out] integrate(1/(sqrt(x^2*(sqrt(3) - 3) + 3)*sqrt(2*sqrt(3)*x^2 - 3*sqrt(3) + 3
)), x)
```

maple [B] time = 0.22, size = 207, normalized size = 4.40

$$\frac{\sqrt{\sqrt{3}x^2-3x^2+3}\sqrt{2\sqrt{3}x^2+3-3\sqrt{3}}\sqrt{2}\sqrt{-(4\sqrt{3}x^2-6x^2-3\sqrt{3}+3)(\sqrt{3}-1)}\sqrt{-(2\sqrt{3}x^2+3-3\sqrt{3})}}{18(\sqrt{3}-1)^2(2\sqrt{3}x^4-2x^4-6\sqrt{3}x^2+6x^2+3\sqrt{3}-3)\sqrt{\dots}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(3+x^2*(-3+3^(1/2)))^(1/2)/(3-3*3^(1/2)+2*3^(1/2)*x^2)^(1/2),x)
```

```
[Out] 1/18*(3^(1/2)*x^2-3*x^2+3)^(1/2)*(3-3*3^(1/2)+2*3^(1/2)*x^2)^(1/2)*2^(1/2)/
(3^(1/2)-1)^2*(-(4*3^(1/2)*x^2-6*x^2-3*3^(1/2)+3)*(3^(1/2)-1))^(1/2)*(-(3-3
*3^(1/2)+2*3^(1/2)*x^2)*(3^(1/2)-1))^(1/2)*EllipticF(1/3*x*2^(1/2)*3^(1/2)/
(3^(1/2)-1)*((2*3^(1/2)-3)*(3^(1/2)-1))^(1/2),1/(3^(1/2)-1)*((3^(1/2)-1)*(1
+3^(1/2)))^(1/2))*(-3+3^(1/2))/(2*x^4*3^(1/2)-2*x^4-6*3^(1/2)*x^2+6*x^2+3*3
^(1/2)-3)/((2*3^(1/2)-3)*(3^(1/2)-1))^(1/2)
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{x^2(\sqrt{3}-3)+3}\sqrt{2\sqrt{3}x^2-3\sqrt{3}+3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(3+x^2*(-3+3^(1/2)))^(1/2)/(3-3*3^(1/2)+2*3^(1/2)*x^2)^(1/2),x,
algorithm="maxima")
```

```
[Out] integrate(1/(sqrt(x^2*(sqrt(3) - 3) + 3)*sqrt(2*sqrt(3)*x^2 - 3*sqrt(3) + 3
)), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{(\sqrt{3}-3)x^2+3}\sqrt{2\sqrt{3}x^2-3\sqrt{3}+3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((x^2*(3^(1/2) - 3) + 3)^(1/2)*(2*3^(1/2)*x^2 - 3*3^(1/2) + 3)^(1/2))
,x)
```

```
[Out] int(1/((x^2*(3^(1/2) - 3) + 3)^(1/2)*(2*3^(1/2)*x^2 - 3*3^(1/2) + 3)^(1/2)), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-3x^2 + \sqrt{3}x^2 + 3} \sqrt{2\sqrt{3}x^2 - 3\sqrt{3} + 3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(3+x**2*(-3+3**(1/2)))**(1/2)/(3-3*3**(1/2)+2*3**(1/2)*x**2)**(1/2), x)
```

```
[Out] Integral(1/(sqrt(-3*x**2 + sqrt(3)*x**2 + 3)*sqrt(2*sqrt(3)*x**2 - 3*sqrt(3) + 3)), x)
```

$$3.304 \quad \int \frac{1}{\sqrt[4]{2+3x^2} (4+3x^2)} dx$$

Optimal. Leaf size=129

$$-\frac{\tan^{-1}\left(\frac{2\sqrt[4]{2}\sqrt{3x^2+2}+2\cdot 2^{3/4}}{2\sqrt{3}x\sqrt[4]{3x^2+2}}\right)}{2\cdot 2^{3/4}\sqrt{3}} - \frac{\tanh^{-1}\left(\frac{2\cdot 2^{3/4}-2\sqrt[4]{2}\sqrt{3x^2+2}}{2\sqrt{3}x\sqrt[4]{3x^2+2}}\right)}{2\cdot 2^{3/4}\sqrt{3}}$$

[Out] $-1/12*\arctan(1/6*(2*2^{(3/4)}+2*2^{(1/4)}*(3*x^2+2)^{(1/2)})/x/(3*x^2+2)^{(1/4)}*3^{(1/2)})*2^{(1/4)}*3^{(1/2)}-1/12*\operatorname{arctanh}(1/6*(2*2^{(3/4)}-2*2^{(1/4)}*(3*x^2+2)^{(1/2)})/x/(3*x^2+2)^{(1/4)}*3^{(1/2)})*2^{(1/4)}*3^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 129, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {397}

$$-\frac{\tan^{-1}\left(\frac{2\sqrt[4]{2}\sqrt{3x^2+2}+2\cdot 2^{3/4}}{2\sqrt{3}x\sqrt[4]{3x^2+2}}\right)}{2\cdot 2^{3/4}\sqrt{3}} - \frac{\tanh^{-1}\left(\frac{2\cdot 2^{3/4}-2\sqrt[4]{2}\sqrt{3x^2+2}}{2\sqrt{3}x\sqrt[4]{3x^2+2}}\right)}{2\cdot 2^{3/4}\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[1/((2 + 3*x^2)^(1/4)*(4 + 3*x^2)),x]

[Out] $-\operatorname{ArcTan}[(2*2^{(3/4)} + 2*2^{(1/4)}*\operatorname{Sqrt}[2 + 3*x^2])/(2*\operatorname{Sqrt}[3]*x*(2 + 3*x^2)^{(1/4)})]/(2*2^{(3/4)}*\operatorname{Sqrt}[3]) - \operatorname{ArcTanh}[(2*2^{(3/4)} - 2*2^{(1/4)}*\operatorname{Sqrt}[2 + 3*x^2])/(2*\operatorname{Sqrt}[3]*x*(2 + 3*x^2)^{(1/4)})]/(2*2^{(3/4)}*\operatorname{Sqrt}[3])$

Rule 397

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[b^2/a, 4]}, -Simp[(b*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])/(2*a*d*q), x] - Simp[(b*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])/(2*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]

Rubi steps

$$\int \frac{1}{\sqrt[4]{2+3x^2} (4+3x^2)} dx = -\frac{\tan^{-1}\left(\frac{2\cdot 2^{3/4}+2\sqrt[4]{2}\sqrt{2+3x^2}}{2\sqrt{3}x\sqrt[4]{2+3x^2}}\right)}{2\cdot 2^{3/4}\sqrt{3}} - \frac{\tanh^{-1}\left(\frac{2\cdot 2^{3/4}-2\sqrt[4]{2}\sqrt{2+3x^2}}{2\sqrt{3}x\sqrt[4]{2+3x^2}}\right)}{2\cdot 2^{3/4}\sqrt{3}}$$

Mathematica [C] time = 0.11, size = 135, normalized size = 1.05

$$\frac{4xF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{3x^2}{2}, -\frac{3x^2}{4}\right)}{\sqrt[4]{3x^2+2} (3x^2+4) \left(x^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{3x^2}{2}, -\frac{3x^2}{4}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{3x^2}{2}, -\frac{3x^2}{4}\right)\right) - 4F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{3x^2}{2}, -\frac{3x^2}{4}\right)}\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((2 + 3*x^2)^(1/4)*(4 + 3*x^2)),x]

[Out] $(-4*x*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, (-3*x^2)/2, (-3*x^2)/4])/((2 + 3*x^2)^{(1/4)}*(4 + 3*x^2))*(-4*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, (-3*x^2)/2, (-3*x^2)/4] + x^2*$

(2*AppellF1[3/2, 1/4, 2, 5/2, (-3*x^2)/2, (-3*x^2)/4] + AppellF1[3/2, 5/4, 1, 5/2, (-3*x^2)/2, (-3*x^2)/4]))

fricas [B] time = 4.00, size = 553, normalized size = 4.29

$$\frac{1}{72} \cdot 18^{\frac{3}{4}} \sqrt{2} \arctan \left(\frac{6 \cdot 18^{\frac{3}{4}} \sqrt{2} (3x^2 + 2)^{\frac{1}{4}} x^3 + 54x^4 + 24 \cdot 18^{\frac{1}{4}} \sqrt{2} (3x^2 + 2)^{\frac{3}{4}} x + 12 \sqrt{2} (3x^2 + 4) \sqrt{3x^2 + 2}}{\dots} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2+2)^(1/4)/(3*x^2+4),x, algorithm="fricas")

[Out] 1/72*18^(3/4)*sqrt(2)*arctan(-1/6*(6*18^(3/4)*sqrt(2)*(3*x^2 + 2)^(1/4)*x^3 + 54*x^4 + 24*18^(1/4)*sqrt(2)*(3*x^2 + 2)^(3/4)*x + 12*sqrt(2)*(3*x^2 + 4)*sqrt(3*x^2 + 2) + 72*x^2 - (18^(3/4)*sqrt(2)*(3*x^3 - 4*x)*sqrt(3*x^2 + 2) + 72*(3*x^2 + 2)^(1/4)*x^2 + 6*18^(1/4)*sqrt(2)*(3*x^3 + 4*x) + 48*sqrt(2)*(3*x^2 + 2)^(3/4))*sqrt((3*sqrt(2)*x^2 + 2*18^(1/4)*sqrt(2)*(3*x^2 + 2)^(1/4)*x + 4*sqrt(3*x^2 + 2))/(3*x^2 + 4)))/(9*x^4 - 24*x^2 - 16)) - 1/72*18^(3/4)*sqrt(2)*arctan(1/6*(6*18^(3/4)*sqrt(2)*(3*x^2 + 2)^(1/4)*x^3 - 54*x^4 + 24*18^(1/4)*sqrt(2)*(3*x^2 + 2)^(3/4)*x - 12*sqrt(2)*(3*x^2 + 4)*sqrt(3*x^2 + 2) - 72*x^2 - (18^(3/4)*sqrt(2)*(3*x^3 - 4*x)*sqrt(3*x^2 + 2) - 72*(3*x^2 + 2)^(1/4)*x^2 + 6*18^(1/4)*sqrt(2)*(3*x^3 + 4*x) - 48*sqrt(2)*(3*x^2 + 2)^(3/4))*sqrt((3*sqrt(2)*x^2 - 2*18^(1/4)*sqrt(2)*(3*x^2 + 2)^(1/4)*x + 4*sqrt(3*x^2 + 2))/(3*x^2 + 4)))/(9*x^4 - 24*x^2 - 16)) + 1/288*18^(3/4)*sqrt(2)*log(36*(3*sqrt(2)*x^2 + 2*18^(1/4)*sqrt(2)*(3*x^2 + 2)^(1/4)*x + 4*sqrt(3*x^2 + 2))/(3*x^2 + 4)) - 1/288*18^(3/4)*sqrt(2)*log(36*(3*sqrt(2)*x^2 - 2*18^(1/4)*sqrt(2)*(3*x^2 + 2)^(1/4)*x + 4*sqrt(3*x^2 + 2))/(3*x^2 + 4))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 + 4)(3x^2 + 2)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2+2)^(1/4)/(3*x^2+4),x, algorithm="giac")

[Out] integrate(1/((3*x^2 + 4)*(3*x^2 + 2)^(1/4)), x)

maple [C] time = 1.67, size = 186, normalized size = 1.44

$$\text{RootOf} \left(_Z^2 + \text{RootOf} \left(_Z^4 + 72 \right)^2 \right) \ln \left(\frac{3x \text{RootOf} \left(_Z^4 + 72 \right)^2 + (3x^2 + 2)^{\frac{1}{4}} \text{RootOf} \left(_Z^2 + \text{RootOf} \left(_Z^4 + 72 \right)^2 \right) \text{RootOf} \left(_Z^4 + 72 \right)^2}{3x^2 + 4} \right)$$

24

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(3*x^2+2)^(1/4)/(3*x^2+4),x)

[Out] 1/24*RootOf(_Z^2+RootOf(_Z^4+72)^2)*ln(-((3*x^2+2)^(1/4)*RootOf(_Z^4+72)^2*RootOf(_Z^2+RootOf(_Z^4+72)^2)-6*(3*x^2+2)^(3/4)*RootOf(_Z^2+RootOf(_Z^4+72)^2)+3*RootOf(_Z^4+72)^2*x-18*(3*x^2+2)^(1/2)*x)/(3*x^2+4))+1/24*RootOf(_Z^4+72)*ln(((3*x^2+2)^(1/4)*RootOf(_Z^4+72)^3+6*(3*x^2+2)^(3/4)*RootOf(_Z^4+72)^2+3*RootOf(_Z^4+72)^2*x+18*(3*x^2+2)^(1/2)*x)/(3*x^2+4))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 + 4)(3x^2 + 2)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2+2)^(1/4)/(3*x^2+4),x, algorithm="maxima")

[Out] integrate(1/((3*x^2 + 4)*(3*x^2 + 2)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(3x^2 + 2)^{\frac{1}{4}}(3x^2 + 4)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((3*x^2 + 2)^(1/4)*(3*x^2 + 4)),x)

[Out] int(1/((3*x^2 + 2)^(1/4)*(3*x^2 + 4)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[4]{3x^2 + 2}(3x^2 + 4)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x**2+2)**(1/4)/(3*x**2+4),x)

[Out] Integral(1/((3*x**2 + 2)**(1/4)*(3*x**2 + 4)), x)

$$3.305 \quad \int \frac{1}{\sqrt[4]{2-3x^2} (4-3x^2)} dx$$

Optimal. Leaf size=120

$$\frac{\tan^{-1}\left(\frac{2-\sqrt{2}\sqrt{2-3x^2}}{\sqrt[4]{2}\sqrt{3x}\sqrt[4]{2-3x^2}}\right)}{2^{2^{3/4}}\sqrt{3}} + \frac{\tanh^{-1}\left(\frac{\sqrt{2}\sqrt{2-3x^2}+2}{\sqrt[4]{2}\sqrt{3x}\sqrt[4]{2-3x^2}}\right)}{2^{2^{3/4}}\sqrt{3}}$$

[Out] 1/12*arctan(1/6*(2-2^(1/2))*(-3*x^2+2)^(1/2))*2^(3/4)/x/(-3*x^2+2)^(1/4)*3^(1/2))*2^(1/4)*3^(1/2)+1/12*arctanh(1/6*(2+2^(1/2))*(-3*x^2+2)^(1/2))*2^(3/4)/x/(-3*x^2+2)^(1/4)*3^(1/2))*2^(1/4)*3^(1/2)

Rubi [A] time = 0.02, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {397}

$$\frac{\tan^{-1}\left(\frac{2-\sqrt{2}\sqrt{2-3x^2}}{\sqrt[4]{2}\sqrt{3x}\sqrt[4]{2-3x^2}}\right)}{2^{2^{3/4}}\sqrt{3}} + \frac{\tanh^{-1}\left(\frac{\sqrt{2}\sqrt{2-3x^2}+2}{\sqrt[4]{2}\sqrt{3x}\sqrt[4]{2-3x^2}}\right)}{2^{2^{3/4}}\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[1/((2 - 3*x^2)^(1/4)*(4 - 3*x^2)),x]

[Out] ArcTan[(2 - Sqrt[2]*Sqrt[2 - 3*x^2])/(2^(1/4)*Sqrt[3]*x*(2 - 3*x^2)^(1/4))]/(2*2^(3/4)*Sqrt[3]) + ArcTanh[(2 + Sqrt[2]*Sqrt[2 - 3*x^2])/(2^(1/4)*Sqrt[3]*x*(2 - 3*x^2)^(1/4))]/(2*2^(3/4)*Sqrt[3])

Rule 397

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b^2/a, 4]}, -Simp[(b*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])]/(2*a*d*q), x] - Simp[(b*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])]/(2*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]

Rubi steps

$$\int \frac{1}{\sqrt[4]{2-3x^2} (4-3x^2)} dx = \frac{\tan^{-1}\left(\frac{2-\sqrt{2}\sqrt{2-3x^2}}{\sqrt[4]{2}\sqrt{3x}\sqrt[4]{2-3x^2}}\right)}{2^{2^{3/4}}\sqrt{3}} + \frac{\tanh^{-1}\left(\frac{2+\sqrt{2}\sqrt{2-3x^2}}{\sqrt[4]{2}\sqrt{3x}\sqrt[4]{2-3x^2}}\right)}{2^{2^{3/4}}\sqrt{3}}$$

Mathematica [C] time = 0.13, size = 135, normalized size = 1.12

$$\frac{4xF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{3x^2}{2}, \frac{3x^2}{4}\right)}{\sqrt[4]{2-3x^2} (3x^2-4) \left(x^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; \frac{3x^2}{2}, \frac{3x^2}{4}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; \frac{3x^2}{2}, \frac{3x^2}{4}\right)\right) + 4F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{3x^2}{2}, \frac{3x^2}{4}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((2 - 3*x^2)^(1/4)*(4 - 3*x^2)),x]

[Out] (-4*x*AppellF1[1/2, 1/4, 1, 3/2, (3*x^2)/2, (3*x^2)/4])/((2 - 3*x^2)^(1/4)*(-4 + 3*x^2)*(4*AppellF1[1/2, 1/4, 1, 3/2, (3*x^2)/2, (3*x^2)/4] + x^2*(2*A

ppellF1[3/2, 1/4, 2, 5/2, (3*x^2)/2, (3*x^2)/4] + AppellF1[3/2, 5/4, 1, 5/2, (3*x^2)/2, (3*x^2)/4]))

fricas [B] time = 4.34, size = 553, normalized size = 4.61

$$\frac{1}{72} \cdot 18^{\frac{3}{4}} \sqrt{2} \arctan \left(\frac{6 \cdot 18^{\frac{3}{4}} \sqrt{2} (-3x^2 + 2)^{\frac{1}{4}} x^3 + 54x^4 + 24 \cdot 18^{\frac{1}{4}} \sqrt{2} (-3x^2 + 2)^{\frac{3}{4}} x + 12 \sqrt{2} (3x^2 - 4) \sqrt{-3x^2 + 2}}{\dots} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/4)/(-3*x^2+4),x, algorithm="fricas")

[Out] 1/72*18^(3/4)*sqrt(2)*arctan(-1/6*(6*18^(3/4)*sqrt(2)*(-3*x^2 + 2)^(1/4)*x^3 + 54*x^4 + 24*18^(1/4)*sqrt(2)*(-3*x^2 + 2)^(3/4)*x + 12*sqrt(2)*(3*x^2 - 4)*sqrt(-3*x^2 + 2) - 72*x^2 + (18^(3/4)*sqrt(2)*(3*x^3 + 4*x)*sqrt(-3*x^2 + 2) - 72*(-3*x^2 + 2)^(1/4)*x^2 - 6*18^(1/4)*sqrt(2)*(3*x^3 - 4*x) - 48*sqrt(2)*(-3*x^2 + 2)^(3/4))*sqrt(-3*sqrt(2)*x^2 + 2*18^(1/4)*sqrt(2)*(-3*x^2 + 2)^(1/4)*x + 4*sqrt(-3*x^2 + 2))/(3*x^2 - 4))/(9*x^4 + 24*x^2 - 16)) - 1/72*18^(3/4)*sqrt(2)*arctan(1/6*(6*18^(3/4)*sqrt(2)*(-3*x^2 + 2)^(1/4)*x^3 - 54*x^4 + 24*18^(1/4)*sqrt(2)*(-3*x^2 + 2)^(3/4)*x - 12*sqrt(2)*(3*x^2 - 4)*sqrt(-3*x^2 + 2) + 72*x^2 + (18^(3/4)*sqrt(2)*(3*x^3 + 4*x)*sqrt(-3*x^2 + 2) + 72*(-3*x^2 + 2)^(1/4)*x^2 - 6*18^(1/4)*sqrt(2)*(3*x^3 - 4*x) + 48*sqrt(2)*(-3*x^2 + 2)^(3/4))*sqrt(-3*sqrt(2)*x^2 - 2*18^(1/4)*sqrt(2)*(-3*x^2 + 2)^(1/4)*x + 4*sqrt(-3*x^2 + 2))/(3*x^2 - 4))/(9*x^4 + 24*x^2 - 16)) + 1/288*18^(3/4)*sqrt(2)*log(-36*(3*sqrt(2)*x^2 + 2*18^(1/4)*sqrt(2)*(-3*x^2 + 2)^(1/4)*x + 4*sqrt(-3*x^2 + 2))/(3*x^2 - 4)) - 1/288*18^(3/4)*sqrt(2)*log(-36*(3*sqrt(2)*x^2 - 2*18^(1/4)*sqrt(2)*(-3*x^2 + 2)^(1/4)*x + 4*sqrt(-3*x^2 + 2))/(3*x^2 - 4))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(3x^2 - 4)(-3x^2 + 2)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/4)/(-3*x^2+4),x, algorithm="giac")

[Out] integrate(-1/((3*x^2 - 4)*(-3*x^2 + 2)^(1/4)), x)

maple [C] time = 1.60, size = 187, normalized size = 1.56

$$\text{RootOf}(-Z^2 + \text{RootOf}(-Z^4 + 72)^2) \ln \left(-\frac{-3x \text{RootOf}(-Z^4 + 72)^2 + (-3x^2 + 2)^{\frac{1}{4}} \text{RootOf}(-Z^2 + \text{RootOf}(-Z^4 + 72)^2) \text{RootOf}(-Z^4 + 72)^2}{3x^2 - 4} \right)$$

24

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*x^2+2)^(1/4)/(-3*x^2+4),x)

[Out] 1/24*RootOf(-Z^4+72)*ln(-((-3*x^2+2)^(1/4)*RootOf(-Z^4+72)^3-6*(-3*x^2+2)^(3/4)*RootOf(-Z^4+72)+3*RootOf(-Z^4+72)^2*x-18*(-3*x^2+2)^(1/2)*x)/(3*x^2-4))-1/24*RootOf(-Z^2+RootOf(-Z^4+72)^2)*ln(-((-3*x^2+2)^(1/4)*RootOf(-Z^4+72)^2*RootOf(-Z^2+RootOf(-Z^4+72)^2)+6*(-3*x^2+2)^(3/4)*RootOf(-Z^2+RootOf(-Z^4+72)^2)-3*RootOf(-Z^4+72)^2*x-18*(-3*x^2+2)^(1/2)*x)/(3*x^2-4))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(3x^2 - 4)(-3x^2 + 2)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+2)^(1/4)/(-3*x^2+4),x, algorithm="maxima")

[Out] -integrate(1/((3*x^2 - 4)*(-3*x^2 + 2)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(2 - 3x^2)^{\frac{1}{4}} (3x^2 - 4)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((2 - 3*x^2)^(1/4)*(3*x^2 - 4)),x)

[Out] -int(1/((2 - 3*x^2)^(1/4)*(3*x^2 - 4)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{3x^2\sqrt[4]{2-3x^2} - 4\sqrt[4]{2-3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x**2+2)**(1/4)/(-3*x**2+4),x)

[Out] -Integral(1/(3*x**2*(2 - 3*x**2)**(1/4) - 4*(2 - 3*x**2)**(1/4)), x)

$$3.306 \quad \int \frac{1}{\sqrt[4]{2+bx^2} (4+bx^2)} dx$$

Optimal. Leaf size=129

$$-\frac{\tan^{-1}\left(\frac{2\sqrt[4]{2}\sqrt{bx^2+2}+2^{3/4}}{2\sqrt{b}x\sqrt[4]{bx^2+2}}\right)}{2^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{2^{3/4}-2\sqrt[4]{2}\sqrt{bx^2+2}}{2\sqrt{b}x\sqrt[4]{bx^2+2}}\right)}{2^{3/4}\sqrt{b}}$$

[Out] $-1/4*\arctan(1/2*(2*2^{(3/4)}+2*2^{(1/4)}*(b*x^2+2)^{(1/2)})/x/(b*x^2+2)^{(1/4)}/b^{(1/2)})*2^{(1/4)}/b^{(1/2)}-1/4*\operatorname{arctanh}(1/2*(2*2^{(3/4)}-2*2^{(1/4)}*(b*x^2+2)^{(1/2)})/x/(b*x^2+2)^{(1/4)}/b^{(1/2)})*2^{(1/4)}/b^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 129, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {397}

$$-\frac{\tan^{-1}\left(\frac{2\sqrt[4]{2}\sqrt{bx^2+2}+2^{3/4}}{2\sqrt{b}x\sqrt[4]{bx^2+2}}\right)}{2^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{2^{3/4}-2\sqrt[4]{2}\sqrt{bx^2+2}}{2\sqrt{b}x\sqrt[4]{bx^2+2}}\right)}{2^{3/4}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((2 + b*x^2)^(1/4)*(4 + b*x^2)),x]

[Out] $-\operatorname{ArcTan}[(2*2^{(3/4)} + 2*2^{(1/4)}*\operatorname{Sqrt}[2 + b*x^2])/(2*\operatorname{Sqrt}[b]*x*(2 + b*x^2)^{(1/4)})]/(2*2^{(3/4)}*\operatorname{Sqrt}[b]) - \operatorname{ArcTanh}[(2*2^{(3/4)} - 2*2^{(1/4)}*\operatorname{Sqrt}[2 + b*x^2])/(2*\operatorname{Sqrt}[b]*x*(2 + b*x^2)^{(1/4)})]/(2*2^{(3/4)}*\operatorname{Sqrt}[b])$

Rule 397

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[b^2/a, 4]}, -Simp[(b*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])/(2*a*d*q), x] - Simp[(b*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])/(2*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]

Rubi steps

$$\int \frac{1}{\sqrt[4]{2+bx^2} (4+bx^2)} dx = -\frac{\tan^{-1}\left(\frac{2^{3/4}+2\sqrt[4]{2}\sqrt{2+bx^2}}{2\sqrt{b}x\sqrt[4]{2+bx^2}}\right)}{2^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{2^{3/4}-2\sqrt[4]{2}\sqrt{2+bx^2}}{2\sqrt{b}x\sqrt[4]{2+bx^2}}\right)}{2^{3/4}\sqrt{b}}$$

Mathematica [C] time = 0.14, size = 144, normalized size = 1.12

$$\frac{12xF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{2}, -\frac{bx^2}{4}\right)}{\sqrt[4]{bx^2+2} (bx^2+4) \left(bx^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{2}, -\frac{bx^2}{4}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{2}, -\frac{bx^2}{4}\right)\right) - 12F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{2}, -\frac{bx^2}{4}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((2 + b*x^2)^(1/4)*(4 + b*x^2)),x]

[Out] $(-12*x*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, -1/2*(b*x^2), -1/4*(b*x^2)])/((2 + b*x^2)^{(1/4)}*(4 + b*x^2))*(-12*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, -1/2*(b*x^2), -1/4*(b*x^2)$

2)] + b*x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, -1/2*(b*x^2), -1/4*(b*x^2)] + AppellF1[3/2, 5/4, 1, 5/2, -1/2*(b*x^2), -1/4*(b*x^2)]))

fricas [B] time = 12.45, size = 755, normalized size = 5.85

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/4)/(b*x^2+4),x, algorithm="fricas")

[Out] $\frac{1}{4}\sqrt{2}\left(\frac{1}{2}\right)^{1/4}(b^{-2})^{1/4}\arctan\left(-\frac{2\sqrt{2}\left(\frac{1}{2}\right)^{1/4}(b^2x^2+2)^{1/4}b^2(b^{-2})^{1/4}x^3+b^2x^4+8\sqrt{2}\left(\frac{1}{2}\right)^{3/4}(b^2x^2+2)^{3/4}b^2(b^{-2})^{3/4}x+4b^2x^2+4\sqrt{1/2}(b^2x^2+4b)\sqrt{b^2x^2+2}\sqrt{b^{-2}}-2\sqrt{1/2}(4(b^2x^2+2)^{1/4}b^2x^2+2\sqrt{2}\left(\frac{1}{2}\right)^{3/4}(b^3x^3+4b^2x)(b^{-2})^{3/4}+16\sqrt{1/2}(b^2x^2+2)^{3/4}b\sqrt{b^{-2}}+\sqrt{2}\left(\frac{1}{2}\right)^{1/4}(b^2x^3-4b^2x)\sqrt{b^2x^2+2}(b^{-2})^{1/4}\sqrt{(2\sqrt{2}\left(\frac{1}{2}\right)^{3/4}(b^2x^2+2)^{1/4}b^2(b^{-2})^{3/4}x+\sqrt{1/2}b^2\sqrt{b^{-2}}x^2+2\sqrt{b^2x^2+2})/(b^2x^2+4)))/(b^2x^4-8b^2x^2-16)}-1/4\sqrt{2}\left(\frac{1}{2}\right)^{1/4}(b^{-2})^{1/4}\arctan\left(\frac{2\sqrt{2}\left(\frac{1}{2}\right)^{1/4}(b^2x^2+2)^{1/4}b^2(b^{-2})^{1/4}x^3-b^2x^4+8\sqrt{2}\left(\frac{1}{2}\right)^{3/4}(b^2x^2+2)^{3/4}b^2(b^{-2})^{3/4}x-4b^2x^2-4\sqrt{1/2}(b^2x^2+4b)\sqrt{b^2x^2+2}\sqrt{b^{-2}}+2\sqrt{1/2}(4(b^2x^2+2)^{1/4}b^2x^2-2\sqrt{2}\left(\frac{1}{2}\right)^{3/4}(b^3x^3+4b^2x)(b^{-2})^{3/4}+16\sqrt{1/2}(b^2x^2+2)^{3/4}b\sqrt{b^{-2}}-\sqrt{2}\left(\frac{1}{2}\right)^{1/4}(b^2x^3-4b^2x)\sqrt{b^2x^2+2}(b^{-2})^{1/4}\sqrt{-(2\sqrt{2}\left(\frac{1}{2}\right)^{3/4}(b^2x^2+2)^{1/4}b^2(b^{-2})^{3/4}x-\sqrt{1/2}b^2\sqrt{b^{-2}}x^2-2\sqrt{b^2x^2+2})/(b^2x^2+4)))/(b^2x^4-8b^2x^2-16)}+1/16\sqrt{2}\left(\frac{1}{2}\right)^{1/4}(b^{-2})^{1/4}\log\left(\frac{1/2(2\sqrt{2}\left(\frac{1}{2}\right)^{3/4}(b^2x^2+2)^{1/4}b^2(b^{-2})^{3/4}x+\sqrt{1/2}b^2\sqrt{b^{-2}}x^2+2\sqrt{b^2x^2+2})/(b^2x^2+4)}{-1/2(2\sqrt{2}\left(\frac{1}{2}\right)^{3/4}(b^2x^2+2)^{1/4}b^2(b^{-2})^{3/4}x-\sqrt{1/2}b^2\sqrt{b^{-2}}x^2-2\sqrt{b^2x^2+2})/(b^2x^2+4)}\right)\right)$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+4)(bx^2+2)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/4)/(b*x^2+4),x, algorithm="giac")

[Out] integrate(1/((b*x^2+4)*(b*x^2+2)^(1/4)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+2)^{\frac{1}{4}}(bx^2+4)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+2)^(1/4)/(b*x^2+4), x)

[Out] int(1/(b*x^2+2)^(1/4)/(b*x^2+4), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+4)(bx^2+2)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+2)^(1/4)/(b*x^2+4),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 4)*(b*x^2 + 2)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + 2)^{1/4} (bx^2 + 4)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((b*x^2 + 2)^(1/4)*(b*x^2 + 4)),x)

[Out] int(1/((b*x^2 + 2)^(1/4)*(b*x^2 + 4)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[4]{bx^2 + 2} (bx^2 + 4)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+2)**(1/4)/(b*x**2+4),x)

[Out] Integral(1/((b*x**2 + 2)**(1/4)*(b*x**2 + 4)), x)

$$3.307 \quad \int \frac{1}{\sqrt[4]{2-bx^2} (4-bx^2)} dx$$

Optimal. Leaf size=124

$$\frac{\tan^{-1}\left(\frac{2-\sqrt{2}\sqrt{2-bx^2}}{\sqrt[4]{2}\sqrt{bx^2}\sqrt[4]{2-bx^2}}\right)}{2^{2^{3/4}}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{2}\sqrt{2-bx^2}+2}{\sqrt[4]{2}\sqrt{bx^2}\sqrt[4]{2-bx^2}}\right)}{2^{2^{3/4}}\sqrt{b}}$$

[Out] $1/4*\arctan(1/2*(2-2^{(1/2)}*(-b*x^2+2)^{(1/2)})*2^{(3/4)}/x/(-b*x^2+2)^{(1/4)}/b^{(1/2)})*2^{(1/4)}/b^{(1/2)}+1/4*\operatorname{arctanh}(1/2*(2+2^{(1/2)}*(-b*x^2+2)^{(1/2)})*2^{(3/4)}/x/(-b*x^2+2)^{(1/4)}/b^{(1/2)})*2^{(1/4)}/b^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {397}

$$\frac{\tan^{-1}\left(\frac{2-\sqrt{2}\sqrt{2-bx^2}}{\sqrt[4]{2}\sqrt{bx^2}\sqrt[4]{2-bx^2}}\right)}{2^{2^{3/4}}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{\sqrt{2}\sqrt{2-bx^2}+2}{\sqrt[4]{2}\sqrt{bx^2}\sqrt[4]{2-bx^2}}\right)}{2^{2^{3/4}}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((2 - b*x^2)^(1/4)*(4 - b*x^2)),x]

[Out] ArcTan[(2 - Sqrt[2]*Sqrt[2 - b*x^2])/(2^(1/4)*Sqrt[b]*x*(2 - b*x^2)^(1/4))]/(2*2^(3/4)*Sqrt[b]) + ArcTanh[(2 + Sqrt[2]*Sqrt[2 - b*x^2])/(2^(1/4)*Sqrt[b]*x*(2 - b*x^2)^(1/4))]/(2*2^(3/4)*Sqrt[b])

Rule 397

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b^2/a, 4]}, -Simp[(b*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])]/(2*a*d*q), x] - Simp[(b*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])]/(2*a*d*q), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]

Rubi steps

$$\int \frac{1}{\sqrt[4]{2-bx^2} (4-bx^2)} dx = \frac{\tan^{-1}\left(\frac{2-\sqrt{2}\sqrt{2-bx^2}}{\sqrt[4]{2}\sqrt{bx^2}\sqrt[4]{2-bx^2}}\right)}{2^{2^{3/4}}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{2+\sqrt{2}\sqrt{2-bx^2}}{\sqrt[4]{2}\sqrt{bx^2}\sqrt[4]{2-bx^2}}\right)}{2^{2^{3/4}}\sqrt{b}}$$

Mathematica [C] time = 0.13, size = 145, normalized size = 1.17

$$\frac{12xF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{bx^2}{2}, \frac{bx^2}{4}\right)}{\sqrt[4]{2-bx^2} (bx^2-4) \left(bx^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; \frac{bx^2}{2}, \frac{bx^2}{4}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; \frac{bx^2}{2}, \frac{bx^2}{4}\right)\right) + 12F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{bx^2}{2}, \frac{bx^2}{4}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((2 - b*x^2)^(1/4)*(4 - b*x^2)),x]

[Out] (-12*x*AppellF1[1/2, 1/4, 1, 3/2, (b*x^2)/2, (b*x^2)/4])/((2 - b*x^2)^(1/4))*(-4 + b*x^2)*(12*AppellF1[1/2, 1/4, 1, 3/2, (b*x^2)/2, (b*x^2)/4] + b*x^2*

(2*AppellF1[3/2, 1/4, 2, 5/2, (b*x^2)/2, (b*x^2)/4] + AppellF1[3/2, 5/4, 1, 5/2, (b*x^2)/2, (b*x^2)/4]))

fricas [B] time = 13.09, size = 776, normalized size = 6.26

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+2)^(1/4)/(-b*x^2+4),x, algorithm="fricas")

[Out] 1/4*sqrt(2)*(1/2)^(1/4)*(b^(-2))^(1/4)*arctan(-(2*sqrt(2)*(1/2)^(1/4)*(-b*x^2 + 2)^(1/4)*b^2*(b^(-2))^(1/4)*x^3 + b^2*x^4 + 8*sqrt(2)*(1/2)^(3/4)*(-b*x^2 + 2)^(3/4)*b^2*(b^(-2))^(3/4)*x - 4*b*x^2 + 4*sqrt(1/2)*(b^2*x^2 - 4*b)*sqrt(-b*x^2 + 2)*sqrt(b^(-2)) - 2*sqrt(1/2)*(4*(-b*x^2 + 2)^(1/4)*b*x^2 + 2*sqrt(2)*(1/2)^(3/4)*(b^3*x^3 - 4*b^2*x)*(b^(-2))^(3/4) + 16*sqrt(1/2)*(-b*x^2 + 2)^(3/4)*b*sqrt(b^(-2)) - sqrt(2)*(1/2)^(1/4)*(b^2*x^3 + 4*b*x)*sqrt(-b*x^2 + 2)*(b^(-2))^(1/4))*sqrt(-(2*sqrt(2)*(1/2)^(3/4)*(-b*x^2 + 2)^(1/4)*b^2*(b^(-2))^(3/4)*x + sqrt(1/2)*b^2*sqrt(b^(-2))*x^2 + 2*sqrt(-b*x^2 + 2))/(b*x^2 - 4)))/(b^2*x^4 + 8*b*x^2 - 16)) + 1/4*sqrt(2)*(1/2)^(1/4)*(b^(-2))^(1/4)*arctan(-(2*sqrt(2)*(1/2)^(1/4)*(-b*x^2 + 2)^(1/4)*b^2*(b^(-2))^(1/4)*x^3 - b^2*x^4 + 8*sqrt(2)*(1/2)^(3/4)*(-b*x^2 + 2)^(3/4)*b^2*(b^(-2))^(3/4)*x + 4*b*x^2 - 4*sqrt(1/2)*(b^2*x^2 - 4*b)*sqrt(-b*x^2 + 2)*sqrt(b^(-2)) + 2*sqrt(1/2)*(4*(-b*x^2 + 2)^(1/4)*b*x^2 - 2*sqrt(2)*(1/2)^(3/4)*(b^3*x^3 - 4*b^2*x)*(b^(-2))^(3/4) + 16*sqrt(1/2)*(-b*x^2 + 2)^(3/4)*b*sqrt(b^(-2)) + sqrt(2)*(1/2)^(1/4)*(b^2*x^3 + 4*b*x)*sqrt(-b*x^2 + 2)*(b^(-2))^(1/4))*sqrt((2*sqrt(2)*(1/2)^(3/4)*(-b*x^2 + 2)^(1/4)*b^2*(b^(-2))^(3/4)*x - sqrt(1/2)*b^2*sqrt(b^(-2))*x^2 - 2*sqrt(-b*x^2 + 2))/(b*x^2 - 4)))/(b^2*x^4 + 8*b*x^2 - 16)) + 1/16*sqrt(2)*(1/2)^(1/4)*(b^(-2))^(1/4)*log(-1/2*(2*sqrt(2)*(1/2)^(3/4)*(-b*x^2 + 2)^(1/4)*b^2*(b^(-2))^(3/4)*x + sqrt(1/2)*b^2*sqrt(b^(-2))*x^2 + 2*sqrt(-b*x^2 + 2))/(b*x^2 - 4)) - 1/16*sqrt(2)*(1/2)^(1/4)*(b^(-2))^(1/4)*log(1/2*(2*sqrt(2)*(1/2)^(3/4)*(-b*x^2 + 2)^(1/4)*b^2*(b^(-2))^(3/4)*x - sqrt(1/2)*b^2*sqrt(b^(-2))*x^2 - 2*sqrt(-b*x^2 + 2))/(b*x^2 - 4))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 4)(-bx^2 + 2)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+2)^(1/4)/(-b*x^2+4),x, algorithm="giac")

[Out] integrate(-1/((b*x^2 - 4)*(-b*x^2 + 2)^(1/4)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + 2)^{\frac{1}{4}}(-bx^2 + 4)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+2)^(1/4)/(-b*x^2+4),x)

[Out] int(1/(-b*x^2+2)^(1/4)/(-b*x^2+4),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$- \int \frac{1}{(bx^2 - 4)(-bx^2 + 2)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+2)^(1/4)/(-b*x^2+4),x, algorithm="maxima")

[Out] -integrate(1/((b*x^2 - 4)*(-b*x^2 + 2)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(2 - b x^2)^{1/4} (b x^2 - 4)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((2 - b*x^2)^(1/4)*(b*x^2 - 4)),x)

[Out] -int(1/((2 - b*x^2)^(1/4)*(b*x^2 - 4)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{b x^2 \sqrt[4]{-b x^2 + 2} - 4 \sqrt[4]{-b x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2+2)**(1/4)/(-b*x**2+4),x)

[Out] -Integral(1/(b*x**2*(-b*x**2 + 2)**(1/4) - 4*(-b*x**2 + 2)**(1/4)), x)

$$3.308 \quad \int \frac{1}{\sqrt[4]{a+3x^2} (2a+3x^2)} dx$$

Optimal. Leaf size=120

$$\frac{\tan^{-1}\left(\frac{a^{3/4}\left(\frac{\sqrt{a+3x^2}}{\sqrt{a}}+1\right)}{\sqrt{3}x\sqrt[4]{a+3x^2}}\right)}{2\sqrt{3}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a+3x^2}}{\sqrt{a}}\right)}{\sqrt{3}x\sqrt[4]{a+3x^2}}\right)}{2\sqrt{3}a^{3/4}}$$

[Out] $-1/6*\arctan(1/3*a^{(3/4)}*(1+(3*x^2+a)^{(1/2)}/a^{(1/2)})/x/(3*x^2+a)^{(1/4)}*3^{(1/2)})/a^{(3/4)}*3^{(1/2)}-1/6*\operatorname{arctanh}(1/3*a^{(3/4)}*(1-(3*x^2+a)^{(1/2)}/a^{(1/2)})/x/(3*x^2+a)^{(1/4)}*3^{(1/2)})/a^{(3/4)}*3^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {397}

$$\frac{\tan^{-1}\left(\frac{a^{3/4}\left(\frac{\sqrt{a+3x^2}}{\sqrt{a}}+1\right)}{\sqrt{3}x\sqrt[4]{a+3x^2}}\right)}{2\sqrt{3}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a+3x^2}}{\sqrt{a}}\right)}{\sqrt{3}x\sqrt[4]{a+3x^2}}\right)}{2\sqrt{3}a^{3/4}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + 3*x^2)^(1/4)*(2*a + 3*x^2)), x]

[Out] $-\operatorname{ArcTan}[(a^{(3/4)}*(1 + \operatorname{Sqrt}[a + 3*x^2]/\operatorname{Sqrt}[a]))/(\operatorname{Sqrt}[3]*x*(a + 3*x^2)^{(1/4)})]/(2*\operatorname{Sqrt}[3]*a^{(3/4)}) - \operatorname{ArcTanh}[(a^{(3/4)}*(1 - \operatorname{Sqrt}[a + 3*x^2]/\operatorname{Sqrt}[a]))/(\operatorname{Sqrt}[3]*x*(a + 3*x^2)^{(1/4)})]/(2*\operatorname{Sqrt}[3]*a^{(3/4)})$

Rule 397

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b^2/a, 4]}, -Simp[(b*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])/(2*a*d*q), x] - Simp[(b*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])/(2*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]

Rubi steps

$$\int \frac{1}{\sqrt[4]{a+3x^2} (2a+3x^2)} dx = -\frac{\tan^{-1}\left(\frac{a^{3/4}\left(1+\frac{\sqrt{a+3x^2}}{\sqrt{a}}\right)}{\sqrt{3}x\sqrt[4]{a+3x^2}}\right)}{2\sqrt{3}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a+3x^2}}{\sqrt{a}}\right)}{\sqrt{3}x\sqrt[4]{a+3x^2}}\right)}{2\sqrt{3}a^{3/4}}$$

Mathematica [C] time = 0.15, size = 155, normalized size = 1.29

$$\frac{2axF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{3x^2}{a}, -\frac{3x^2}{2a}\right)}{\sqrt[4]{a+3x^2} (2a+3x^2) \left(x^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{3x^2}{a}, -\frac{3x^2}{2a}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{3x^2}{a}, -\frac{3x^2}{2a}\right)\right) - 2aF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{3x^2}{a}, -\frac{3x^2}{2a}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + 3*x^2)^(1/4)*(2*a + 3*x^2)), x]

[Out] $(-2ax \operatorname{AppellF1}[1/2, 1/4, 1, 3/2, (-3x^2)/a, (-3x^2)/(2a)]) / ((a + 3x^2)^{1/4} (2a + 3x^2) (-2a \operatorname{AppellF1}[1/2, 1/4, 1, 3/2, (-3x^2)/a, (-3x^2)/(2a)] + x^2 (2 \operatorname{AppellF1}[3/2, 1/4, 2, 5/2, (-3x^2)/a, (-3x^2)/(2a)] + \operatorname{AppellF1}[3/2, 5/4, 1, 5/2, (-3x^2)/a, (-3x^2)/(2a)]))$

fricas [B] time = 9.57, size = 286, normalized size = 2.38

$$\left(\frac{1}{36}\right)^{\frac{1}{4}} \left(-\frac{1}{a^3}\right)^{\frac{1}{4}} \arctan \left(\frac{2 \left(\sqrt{\frac{1}{2}} \left(6 \left(\frac{1}{36}\right)^{\frac{3}{4}} a^3 \left(-\frac{1}{a^3}\right)^{\frac{3}{4}} + \left(\frac{1}{36}\right)^{\frac{1}{4}} \sqrt{3x^2 + a} a \left(-\frac{1}{a^3}\right)^{\frac{1}{4}} \right) \sqrt{-a \sqrt{-\frac{1}{a^3}}} - \left(\frac{1}{36}\right)^{\frac{1}{4}} (3x^2 + a)^{\frac{1}{4}}}{x} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(3*x^2+a)^(1/4)/(3*x^2+2*a),x, algorithm="fricas")`

[Out] $(1/36)^{1/4} (-1/a^3)^{1/4} \arctan(2(\sqrt{1/2}(6(1/36)^{3/4} a^3 (-1/a^3)^{3/4} + (1/36)^{1/4} \sqrt{3x^2 + a} a (-1/a^3)^{1/4}) \sqrt{-a \sqrt{-1/a^3}} - (1/36)^{1/4} (3x^2 + a)^{1/4}) / x - 1/4 (1/36)^{1/4} (-1/a^3)^{1/4} \log((18(1/36)^{3/4} \sqrt{3x^2 + a} a^2 x (-1/a^3)^{3/4} + (3x^2 + a)^{1/4} a^2 \sqrt{-1/a^3} - 3(1/36)^{1/4} a x (-1/a^3)^{1/4} + (3x^2 + a)^{3/4}) / (3x^2 + 2a)) + 1/4 (1/36)^{1/4} (-1/a^3)^{1/4} \log(-18(1/36)^{3/4} \sqrt{3x^2 + a} a^2 x (-1/a^3)^{3/4} - (3x^2 + a)^{1/4} a^2 \sqrt{-1/a^3} - 3(1/36)^{1/4} a x (-1/a^3)^{1/4} - (3x^2 + a)^{3/4}) / (3x^2 + 2a))$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 + 2a)(3x^2 + a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(3*x^2+a)^(1/4)/(3*x^2+2*a),x, algorithm="giac")`

[Out] `integrate(1/((3*x^2 + 2*a)*(3*x^2 + a)^(1/4)), x)`

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 + a)^{\frac{1}{4}} (3x^2 + 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(3*x^2+a)^(1/4)/(3*x^2+2*a), x)`

[Out] `int(1/(3*x^2+a)^(1/4)/(3*x^2+2*a), x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 + 2a)(3x^2 + a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(3*x^2+a)^(1/4)/(3*x^2+2*a),x, algorithm="maxima")`

[Out] `integrate(1/((3*x^2 + 2*a)*(3*x^2 + a)^(1/4)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(3x^2 + 2a)(3x^2 + a)^{1/4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((2*a + 3*x^2)*(a + 3*x^2)^(1/4)),x)

[Out] int(1/((2*a + 3*x^2)*(a + 3*x^2)^(1/4)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[4]{a + 3x^2} (2a + 3x^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x**2+a)**(1/4)/(3*x**2+2*a),x)

[Out] Integral(1/((a + 3*x**2)**(1/4)*(2*a + 3*x**2)), x)

$$3.309 \quad \int \frac{1}{\sqrt[4]{a-3x^2} (2a-3x^2)} dx$$

Optimal. Leaf size=120

$$\frac{\tan^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a-3x^2}}{\sqrt{a}}\right)}{\sqrt{3}x\sqrt[4]{a-3x^2}}\right)}{2\sqrt{3}a^{3/4}} + \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(\frac{\sqrt{a-3x^2}}{\sqrt{a}}+1\right)}{\sqrt{3}x\sqrt[4]{a-3x^2}}\right)}{2\sqrt{3}a^{3/4}}$$

[Out] 1/6*arctan(1/3*a^(3/4)*(1-(-3*x^2+a)^(1/2)/a^(1/2))/x/(-3*x^2+a)^(1/4)*3^(1/2))/a^(3/4)*3^(1/2)+1/6*arctanh(1/3*a^(3/4)*(1+(-3*x^2+a)^(1/2)/a^(1/2))/x/(-3*x^2+a)^(1/4)*3^(1/2))/a^(3/4)*3^(1/2)

Rubi [A] time = 0.02, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {397}

$$\frac{\tan^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a-3x^2}}{\sqrt{a}}\right)}{\sqrt{3}x\sqrt[4]{a-3x^2}}\right)}{2\sqrt{3}a^{3/4}} + \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(\frac{\sqrt{a-3x^2}}{\sqrt{a}}+1\right)}{\sqrt{3}x\sqrt[4]{a-3x^2}}\right)}{2\sqrt{3}a^{3/4}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - 3*x^2)^(1/4)*(2*a - 3*x^2)), x]

[Out] ArcTan[(a^(3/4)*(1 - Sqrt[a - 3*x^2]/Sqrt[a]))/(Sqrt[3]*x*(a - 3*x^2)^(1/4))]/(2*Sqrt[3]*a^(3/4)) + ArcTanh[(a^(3/4)*(1 + Sqrt[a - 3*x^2]/Sqrt[a]))/(Sqrt[3]*x*(a - 3*x^2)^(1/4))]/(2*Sqrt[3]*a^(3/4))

Rule 397

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b^2/a, 4]}, -Simp[(b*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])]/(2*a*d*q), x] - Simp[(b*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])]/(2*a*d*q), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]

Rubi steps

$$\int \frac{1}{\sqrt[4]{a-3x^2} (2a-3x^2)} dx = \frac{\tan^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a-3x^2}}{\sqrt{a}}\right)}{\sqrt{3}x\sqrt[4]{a-3x^2}}\right)}{2\sqrt{3}a^{3/4}} + \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(1+\frac{\sqrt{a-3x^2}}{\sqrt{a}}\right)}{\sqrt{3}x\sqrt[4]{a-3x^2}}\right)}{2\sqrt{3}a^{3/4}}$$

Mathematica [C] time = 0.16, size = 155, normalized size = 1.29

$$\frac{2axF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{3x^2}{a}, \frac{3x^2}{2a}\right)}{\sqrt[4]{a-3x^2} (3x^2-2a) \left(x^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; \frac{3x^2}{a}, \frac{3x^2}{2a}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; \frac{3x^2}{a}, \frac{3x^2}{2a}\right)\right) + 2aF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{3x^2}{a}, \frac{3x^2}{2a}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - 3*x^2)^(1/4)*(2*a - 3*x^2)), x]

[Out] $(-2*a*x*AppellF1[1/2, 1/4, 1, 3/2, (3*x^2)/a, (3*x^2)/(2*a)])/((a - 3*x^2)^{(1/4)}*(-2*a + 3*x^2)*(2*a*AppellF1[1/2, 1/4, 1, 3/2, (3*x^2)/a, (3*x^2)/(2*a)] + x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, (3*x^2)/a, (3*x^2)/(2*a)] + AppellF1[3/2, 5/4, 1, 5/2, (3*x^2)/a, (3*x^2)/(2*a)]))$

fricas [B] time = 9.74, size = 286, normalized size = 2.38

$$\left(\frac{1}{36}\right)^{\frac{1}{4}} \left(-\frac{1}{a^3}\right)^{\frac{1}{4}} \arctan \left(\frac{2 \left(\sqrt{\frac{1}{2}} \left(6 \left(\frac{1}{36}\right)^{\frac{3}{4}} a^3 \left(-\frac{1}{a^3}\right)^{\frac{3}{4}} - \left(\frac{1}{36}\right)^{\frac{1}{4}} \sqrt{-3x^2 + a} a \left(-\frac{1}{a^3}\right)^{\frac{1}{4}} \right) \sqrt{a \sqrt{-\frac{1}{a^3}} - \left(\frac{1}{36}\right)^{\frac{1}{4}} (-3x^2 + a)^{\frac{1}{4}}}}{x} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+a)^(1/4)/(-3*x^2+2*a),x, algorithm="fricas")

[Out] $(1/36)^{(1/4)}*(-1/a^3)^{(1/4)}*\arctan(2*(\sqrt{1/2}*(6*(1/36)^{(3/4)}*a^3*(-1/a^3)^{(3/4)} - (1/36)^{(1/4)}*\sqrt{-3*x^2 + a}*a*(-1/a^3)^{(1/4)})*\sqrt{a*\sqrt{-1/a^3}}) - (1/36)^{(1/4)}*\sqrt{-3*x^2 + a}*a*(-1/a^3)^{(1/4)}/x) + 1/4*(1/36)^{(1/4)}*(-1/a^3)^{(1/4)}*\log(-18*(1/36)^{(3/4)}*\sqrt{-3*x^2 + a}*a^2*x*(-1/a^3)^{(3/4)} + (-3*x^2 + a)^{(1/4)}*a^2*\sqrt{-1/a^3} + 3*(1/36)^{(1/4)}*a*x*(-1/a^3)^{(1/4)} - (-3*x^2 + a)^{(3/4)})/(3*x^2 - 2*a)) - 1/4*(1/36)^{(1/4)}*(-1/a^3)^{(1/4)}*\log((18*(1/36)^{(3/4)}*\sqrt{-3*x^2 + a}*a^2*x*(-1/a^3)^{(3/4)} - (-3*x^2 + a)^{(1/4)}*a^2*\sqrt{-1/a^3} + 3*(1/36)^{(1/4)}*a*x*(-1/a^3)^{(1/4)} + (-3*x^2 + a)^{(3/4)})/(3*x^2 - 2*a))$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 - 2a)(-3x^2 + a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+a)^(1/4)/(-3*x^2+2*a),x, algorithm="giac")

[Out] integrate(-1/((3*x^2 - 2*a)*(-3*x^2 + a)^(1/4)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(-3x^2 + a)^{\frac{1}{4}}(-3x^2 + 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*x^2+a)^(1/4)/(-3*x^2+2*a),x)

[Out] int(1/(-3*x^2+a)^(1/4)/(-3*x^2+2*a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$- \int \frac{1}{(3x^2 - 2a)(-3x^2 + a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2+a)^(1/4)/(-3*x^2+2*a),x, algorithm="maxima")

[Out] -integrate(1/((3*x^2 - 2*a)*(-3*x^2 + a)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(2a - 3x^2)(a - 3x^2)^{1/4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((2*a - 3*x^2)*(a - 3*x^2)^(1/4)), x)

[Out] int(1/((2*a - 3*x^2)*(a - 3*x^2)^(1/4)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{-2a\sqrt[4]{a-3x^2} + 3x^2\sqrt[4]{a-3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x**2+a)**(1/4)/(-3*x**2+2*a), x)

[Out] -Integral(1/(-2*a*(a - 3*x**2)**(1/4) + 3*x**2*(a - 3*x**2)**(1/4)), x)

$$3.310 \quad \int \frac{1}{\sqrt[4]{a+bx^2} (2a+bx^2)} dx$$

Optimal. Leaf size=120

$$\frac{\tan^{-1}\left(\frac{a^{3/4}\left(\frac{\sqrt{a+bx^2}}{\sqrt{a}}+1\right)}{\sqrt{b}x\sqrt[4]{a+bx^2}}\right)}{2a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a+bx^2}}{\sqrt{a}}\right)}{\sqrt{b}x\sqrt[4]{a+bx^2}}\right)}{2a^{3/4}\sqrt{b}}$$

[Out] $-1/2*\arctan(a^{(3/4)}*(1+(b*x^2+a)^{(1/2)}/a^{(1/2)})/x/(b*x^2+a)^{(1/4)}/b^{(1/2)})/a^{(3/4)}/b^{(1/2)}-1/2*\operatorname{arctanh}(a^{(3/4)}*(1-(b*x^2+a)^{(1/2)}/a^{(1/2)})/x/(b*x^2+a)^{(1/4)}/b^{(1/2)})/a^{(3/4)}/b^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {397}

$$\frac{\tan^{-1}\left(\frac{a^{3/4}\left(\frac{\sqrt{a+bx^2}}{\sqrt{a}}+1\right)}{\sqrt{b}x\sqrt[4]{a+bx^2}}\right)}{2a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a+bx^2}}{\sqrt{a}}\right)}{\sqrt{b}x\sqrt[4]{a+bx^2}}\right)}{2a^{3/4}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(1/4)*(2*a + b*x^2)), x]

[Out] $-\operatorname{ArcTan}\left[\frac{a^{(3/4)}*(1 + \operatorname{Sqrt}[a + b*x^2]/\operatorname{Sqrt}[a])}{\operatorname{Sqrt}[b]*x*(a + b*x^2)^{(1/4)}}\right]/(2*a^{(3/4)}*\operatorname{Sqrt}[b]) - \operatorname{ArcTanh}\left[\frac{a^{(3/4)}*(1 - \operatorname{Sqrt}[a + b*x^2]/\operatorname{Sqrt}[a])}{\operatorname{Sqrt}[b]*x*(a + b*x^2)^{(1/4)}}\right]/(2*a^{(3/4)}*\operatorname{Sqrt}[b])$

Rule 397

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[b^2/a, 4]}, -Simp[(b*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])/(2*a*d*q), x] - Simp[(b*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])/(2*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]

Rubi steps

$$\int \frac{1}{\sqrt[4]{a+bx^2} (2a+bx^2)} dx = -\frac{\tan^{-1}\left(\frac{a^{3/4}\left(1+\frac{\sqrt{a+bx^2}}{\sqrt{a}}\right)}{\sqrt{b}x\sqrt[4]{a+bx^2}}\right)}{2a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a+bx^2}}{\sqrt{a}}\right)}{\sqrt{b}x\sqrt[4]{a+bx^2}}\right)}{2a^{3/4}\sqrt{b}}$$

Mathematica [C] time = 0.15, size = 165, normalized size = 1.38

$$\frac{6axF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{2a}\right)}{\sqrt[4]{a+bx^2} (2a+bx^2)} \left(6aF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{2a}\right) - bx^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{2a}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{2a}\right)\right)\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(1/4)*(2*a + b*x^2)), x]

[Out] $(6*a*x*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -1/2*(b*x^2)/a])/((a + b*x^2)^{1/4}*(2*a + b*x^2)*(6*a*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -1/2*(b*x^2)/a] - b*x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, -((b*x^2)/a), -1/2*(b*x^2)/a] + AppellF1[3/2, 5/4, 1, 5/2, -((b*x^2)/a), -1/2*(b*x^2)/a]))$

fricas [B] time = 58.44, size = 337, normalized size = 2.81

$$\left(\frac{1}{4}\right)^{\frac{1}{4}} \left(-\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \arctan \left(\frac{2 \left(\sqrt{\frac{1}{2}} \left(2 \left(\frac{1}{4}\right)^{\frac{3}{4}} a^3 b \left(-\frac{1}{a^3 b^2}\right)^{\frac{3}{4}} + \left(\frac{1}{4}\right)^{\frac{1}{4}} \sqrt{b x^2 + a} a \left(-\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \right) \sqrt{-a b} \sqrt{-\frac{1}{a^3 b^2}} - \left(\frac{1}{4}\right)^{\frac{1}{4}} (b x^2 + a)^{\frac{1}{4}}}{x} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/4)/(b*x^2+2*a),x, algorithm="fricas")

[Out] $(1/4)^{1/4}*(-1/(a^3*b^2))^{1/4}*arctan(2*(sqrt(1/2)*(2*(1/4)^{3/4}*a^3*b*(-1/(a^3*b^2))^{3/4} + (1/4)^{1/4}*sqrt(b*x^2 + a)*a*(-1/(a^3*b^2))^{1/4})*sqrt(-a*b*sqrt(-1/(a^3*b^2)))) - (1/4)^{1/4}*(b*x^2 + a)^{1/4}*a*(-1/(a^3*b^2))^{1/4})/x - 1/4*(1/4)^{1/4}*(-1/(a^3*b^2))^{1/4}*log((2*(1/4)^{3/4}*sqrt(b*x^2 + a)*a^2*b^2*x*(-1/(a^3*b^2))^{3/4} + (b*x^2 + a)^{1/4}*a^2*b*sqrt(-1/(a^3*b^2)) - (1/4)^{1/4}*a*b*x*(-1/(a^3*b^2))^{1/4} + (b*x^2 + a)^{3/4}))/ (b*x^2 + 2*a)) + 1/4*(1/4)^{1/4}*(-1/(a^3*b^2))^{1/4}*log(-(2*(1/4)^{3/4}*sqrt(b*x^2 + a)*a^2*b^2*x*(-1/(a^3*b^2))^{3/4} - (b*x^2 + a)^{1/4}*a^2*b*sqrt(-1/(a^3*b^2)) - (1/4)^{1/4}*a*b*x*(-1/(a^3*b^2))^{1/4} - (b*x^2 + a)^{3/4}))/ (b*x^2 + 2*a))$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b x^2 + 2 a)(b x^2 + a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/4)/(b*x^2+2*a),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + 2*a)*(b*x^2 + a)^(1/4)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(b x^2 + a)^{\frac{1}{4}} (b x^2 + 2 a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/4)/(b*x^2+2*a),x)

[Out] int(1/(b*x^2+a)^(1/4)/(b*x^2+2*a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b x^2 + 2 a)(b x^2 + a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/4)/(b*x^2+2*a),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + 2*a)*(b*x^2 + a)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + a)^{1/4} (bx^2 + 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(1/4)*(2*a + b*x^2)), x)

[Out] int(1/((a + b*x^2)^(1/4)*(2*a + b*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[4]{a + bx^2} (2a + bx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(1/4)/(b*x**2+2*a), x)

[Out] Integral(1/((a + b*x**2)**(1/4)*(2*a + b*x**2)), x)

$$3.311 \quad \int \frac{1}{\sqrt[4]{a-bx^2} (2a-bx^2)} dx$$

Optimal. Leaf size=124

$$\frac{\tan^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a-bx^2}}{\sqrt{a}}\right)}{\sqrt{b}x\sqrt[4]{a-bx^2}}\right)}{2a^{3/4}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(\frac{\sqrt{a-bx^2}}{\sqrt{a}}+1\right)}{\sqrt{b}x\sqrt[4]{a-bx^2}}\right)}{2a^{3/4}\sqrt{b}}$$

[Out] 1/2*arctan(a^(3/4)*(1-(-b*x^2+a)^(1/2)/a^(1/2))/x/(-b*x^2+a)^(1/4)/b^(1/2))/a^(3/4)/b^(1/2)+1/2*arctanh(a^(3/4)*(1+(-b*x^2+a)^(1/2)/a^(1/2))/x/(-b*x^2+a)^(1/4)/b^(1/2))/a^(3/4)/b^(1/2)

Rubi [A] time = 0.02, antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {397}

$$\frac{\tan^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a-bx^2}}{\sqrt{a}}\right)}{\sqrt{b}x\sqrt[4]{a-bx^2}}\right)}{2a^{3/4}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(\frac{\sqrt{a-bx^2}}{\sqrt{a}}+1\right)}{\sqrt{b}x\sqrt[4]{a-bx^2}}\right)}{2a^{3/4}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((a - b*x^2)^(1/4)*(2*a - b*x^2)), x]

[Out] ArcTan[(a^(3/4)*(1 - Sqrt[a - b*x^2]/Sqrt[a]))/(Sqrt[b]*x*(a - b*x^2)^(1/4))]/(2*a^(3/4)*Sqrt[b]) + ArcTanh[(a^(3/4)*(1 + Sqrt[a - b*x^2]/Sqrt[a]))/(Sqrt[b]*x*(a - b*x^2)^(1/4))]/(2*a^(3/4)*Sqrt[b])

Rule 397

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b^2/a, 4]}, -Simp[(b*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])]/(2*a*d*q), x] - Simp[(b*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))])]/(2*a*d*q), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]

Rubi steps

$$\int \frac{1}{\sqrt[4]{a-bx^2} (2a-bx^2)} dx = \frac{\tan^{-1}\left(\frac{a^{3/4}\left(1-\frac{\sqrt{a-bx^2}}{\sqrt{a}}\right)}{\sqrt{b}x\sqrt[4]{a-bx^2}}\right)}{2a^{3/4}\sqrt{b}} + \frac{\tanh^{-1}\left(\frac{a^{3/4}\left(1+\frac{\sqrt{a-bx^2}}{\sqrt{a}}\right)}{\sqrt{b}x\sqrt[4]{a-bx^2}}\right)}{2a^{3/4}\sqrt{b}}$$

Mathematica [C] time = 0.16, size = 162, normalized size = 1.31

$$\frac{6axF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{bx^2}{a}, \frac{bx^2}{2a}\right)}{\sqrt[4]{a-bx^2} (2a-bx^2) \left(bx^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; \frac{bx^2}{a}, \frac{bx^2}{2a}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; \frac{bx^2}{a}, \frac{bx^2}{2a}\right)\right) + 6aF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{bx^2}{a}, \frac{bx^2}{2a}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a - b*x^2)^(1/4)*(2*a - b*x^2)), x]

[Out] (6*a*x*AppellF1[1/2, 1/4, 1, 3/2, (b*x^2)/a, (b*x^2)/(2*a)]/((a - b*x^2)^(1/4)*(2*a - b*x^2)*(6*a*AppellF1[1/2, 1/4, 1, 3/2, (b*x^2)/a, (b*x^2)/(2*a)] + b*x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, (b*x^2)/a, (b*x^2)/(2*a)] + AppellF1[3/2, 5/4, 1, 5/2, (b*x^2)/a, (b*x^2)/(2*a)])))

fricas [B] time = 60.66, size = 343, normalized size = 2.77

$$\left(\frac{1}{4}\right)^{\frac{1}{4}} \left(-\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \arctan \left(\frac{2 \left(\sqrt{\frac{1}{2}} \left(2 \left(\frac{1}{4}\right)^{\frac{3}{4}} a^3 b \left(-\frac{1}{a^3 b^2}\right)^{\frac{3}{4}} - \left(\frac{1}{4}\right)^{\frac{1}{4}} \sqrt{-bx^2 + a} a \left(-\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \right) \sqrt{ab} \sqrt{-\frac{1}{a^3 b^2}} - \left(\frac{1}{4}\right)^{\frac{1}{4}} (-bx^2 + a)^{\frac{1}{4}}}{x} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/4)/(-b*x^2+2*a),x, algorithm="fricas")

[Out] (1/4)^(1/4)*(-1/(a^3*b^2))^(1/4)*arctan(2*(sqrt(1/2)*(2*(1/4)^(3/4)*a^3*b*(-1/(a^3*b^2))^(3/4) - (1/4)^(1/4)*sqrt(-b*x^2 + a)*a*(-1/(a^3*b^2))^(1/4))*sqrt(a*b*sqrt(-1/(a^3*b^2))) - (1/4)^(1/4)*(-b*x^2 + a)^(1/4)*a*(-1/(a^3*b^2)^(1/4))/x) + 1/4*(1/4)^(1/4)*(-1/(a^3*b^2))^(1/4)*log(-(2*(1/4)^(3/4)*sqrt(-b*x^2 + a)*a^2*b^2*x*(-1/(a^3*b^2))^(3/4) + (-b*x^2 + a)^(1/4)*a^2*b*sqrt(-1/(a^3*b^2)) + (1/4)^(1/4)*a*b*x*(-1/(a^3*b^2))^(1/4) - (-b*x^2 + a)^(3/4))/(b*x^2 - 2*a)) - 1/4*(1/4)^(1/4)*(-1/(a^3*b^2))^(1/4)*log((2*(1/4)^(3/4)*sqrt(-b*x^2 + a)*a^2*b^2*x*(-1/(a^3*b^2))^(3/4) - (-b*x^2 + a)^(1/4)*a^2*b*sqrt(-1/(a^3*b^2)) + (1/4)^(1/4)*a*b*x*(-1/(a^3*b^2))^(1/4) + (-b*x^2 + a)^(3/4))/(b*x^2 - 2*a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(bx^2 - 2a)(-bx^2 + a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/4)/(-b*x^2+2*a),x, algorithm="giac")

[Out] integrate(-1/((b*x^2 - 2*a)*(-b*x^2 + a)^(1/4)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 + a)^{\frac{1}{4}}(-bx^2 + 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2+a)^(1/4)/(-b*x^2+2*a),x)

[Out] int(1/(-b*x^2+a)^(1/4)/(-b*x^2+2*a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(bx^2 - 2a)(-bx^2 + a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2+a)^(1/4)/(-b*x^2+2*a),x, algorithm="maxima")

[Out] -integrate(1/((b*x^2 - 2*a)*(-b*x^2 + a)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(a - bx^2)^{1/4} (2a - bx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a - b*x^2)^(1/4)*(2*a - b*x^2)), x)

[Out] int(1/((a - b*x^2)^(1/4)*(2*a - b*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{-2a\sqrt[4]{a - bx^2} + bx^2\sqrt[4]{a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2+a)**(1/4)/(-b*x**2+2*a), x)

[Out] -Integral(1/(-2*a*(a - b*x**2)**(1/4) + b*x**2*(a - b*x**2)**(1/4)), x)

$$3.312 \quad \int \frac{1}{(-2+3x^2)\sqrt[4]{-1+3x^2}} dx$$

Optimal. Leaf size=61

$$-\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{3x^2-1}}\right)}{2\sqrt{6}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{3x^2-1}}\right)}{2\sqrt{6}}$$

[Out] $-1/12*\arctan(1/2*x*6^{(1/2)}/(3*x^2-1)^{(1/4)})*6^{(1/2)}-1/12*\operatorname{arctanh}(1/2*x*6^{(1/2)}/(3*x^2-1)^{(1/4)})*6^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {398}

$$-\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{3x^2-1}}\right)}{2\sqrt{6}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{3x^2-1}}\right)}{2\sqrt{6}}$$

Antiderivative was successfully verified.

[In] Int[1/((-2 + 3*x^2)*(-1 + 3*x^2)^(1/4)),x]

[Out] $-\operatorname{ArcTan}[(\operatorname{Sqrt}[3/2]*x)/(-1 + 3*x^2)^{(1/4)}]/(2*\operatorname{Sqrt}[6]) - \operatorname{ArcTanh}[(\operatorname{Sqrt}[3/2]*x)/(-1 + 3*x^2)^{(1/4)}]/(2*\operatorname{Sqrt}[6])$

Rule 398

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(-2+3x^2)\sqrt[4]{-1+3x^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{-1+3x^2}}\right)}{2\sqrt{6}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{-1+3x^2}}\right)}{2\sqrt{6}}$$

Mathematica [C] time = 0.15, size = 127, normalized size = 2.08

$$\frac{2x F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; 3x^2, \frac{3x^2}{2}\right)}{(3x^2 - 2)\sqrt[4]{3x^2 - 1} \left(x^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; 3x^2, \frac{3x^2}{2}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; 3x^2, \frac{3x^2}{2}\right)\right) + 2F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; 3x^2, \frac{3x^2}{2}\right)}\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2 + 3*x^2)*(-1 + 3*x^2)^(1/4)),x]

[Out] $(2*x*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, 3*x^2, (3*x^2)/2])/((-2 + 3*x^2)*(-1 + 3*x^2)^{(1/4)}*(2*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, 3*x^2, (3*x^2)/2] + x^2*(2*\operatorname{AppellF1}[3/2, 1/4, 2, 5/2, 3*x^2, (3*x^2)/2] + \operatorname{AppellF1}[3/2, 5/4, 1, 5/2, 3*x^2, (3*x^2)/2])))$

fricas [B] time = 4.44, size = 104, normalized size = 1.70

$$\frac{1}{12} \sqrt{6} \arctan\left(\frac{\sqrt{6}(3x^2-1)^{\frac{1}{4}}}{3x}\right) + \frac{1}{24} \sqrt{6} \log\left(\frac{9x^4 - 6\sqrt{6}(3x^2-1)^{\frac{1}{4}}x^3 + 12\sqrt{3x^2-1}x^2 - 4\sqrt{6}(3x^2-1)^{\frac{3}{4}}x}{9x^4 - 12x^2 + 4}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2-2)/(3*x^2-1)^(1/4),x, algorithm="fricas")

[Out] 1/12*sqrt(6)*arctan(1/3*sqrt(6)*(3*x^2 - 1)^(1/4)/x) + 1/24*sqrt(6)*log(-(9*x^4 - 6*sqrt(6)*(3*x^2 - 1)^(1/4)*x^3 + 12*sqrt(3*x^2 - 1)*x^2 - 4*sqrt(6)*(3*x^2 - 1)^(3/4)*x + 12*x^2 - 4)/(9*x^4 - 12*x^2 + 4))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2-1)^{\frac{1}{4}}(3x^2-2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2-2)/(3*x^2-1)^(1/4),x, algorithm="giac")

[Out] integrate(1/((3*x^2 - 1)^(1/4)*(3*x^2 - 2)), x)

maple [C] time = 1.14, size = 138, normalized size = 2.26

$$\frac{\text{RootOf}(-Z^2-6) \ln\left(-\frac{-3\sqrt{3x^2-1}x-3x+(3x^2-1)^{\frac{3}{4}}\text{RootOf}(-Z^2-6)+(3x^2-1)^{\frac{1}{4}}\text{RootOf}(-Z^2-6)}{3x^2-2}\right) \text{RootOf}(-Z^2+6) \ln\left(\frac{3\sqrt{3x^2-1}x-3x+(3x^2-1)^{\frac{3}{4}}\text{RootOf}(-Z^2-6)+(3x^2-1)^{\frac{1}{4}}\text{RootOf}(-Z^2-6)}{3x^2-2}\right)}{12}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(3*x^2-2)/(3*x^2-1)^(1/4),x)

[Out] -1/12*RootOf(-Z^2+6)*ln((RootOf(-Z^2+6)*(3*x^2-1)^(3/4)+3*(3*x^2-1)^(1/2)*x-RootOf(-Z^2+6)*(3*x^2-1)^(1/4)-3*x)/(3*x^2-2))+1/12*RootOf(-Z^2-6)*ln(-(RootOf(-Z^2-6)*(3*x^2-1)^(3/4)-3*(3*x^2-1)^(1/2)*x+RootOf(-Z^2-6)*(3*x^2-1)^(1/4)-3*x)/(3*x^2-2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2-1)^{\frac{1}{4}}(3x^2-2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2-2)/(3*x^2-1)^(1/4),x, algorithm="maxima")

[Out] integrate(1/((3*x^2 - 1)^(1/4)*(3*x^2 - 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(3x^2-1)^{1/4}(3x^2-2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((3*x^2 - 1)^(1/4)*(3*x^2 - 2)),x)

```
[Out] int(1/((3*x^2 - 1)^(1/4)*(3*x^2 - 2)), x)
```

```
sympy [F] time = 0.00, size = 0, normalized size = 0.00
```

$$\int \frac{1}{(3x^2 - 2)\sqrt[4]{3x^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(3*x**2-2)/(3*x**2-1)**(1/4),x)
```

```
[Out] Integral(1/((3*x**2 - 2)*(3*x**2 - 1)**(1/4)), x)
```

$$3.313 \quad \int \frac{1}{(-2-3x^2)\sqrt[4]{-1-3x^2}} dx$$

Optimal. Leaf size=61

$$-\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{-3x^2-1}}\right)}{2\sqrt{6}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{-3x^2-1}}\right)}{2\sqrt{6}}$$

[Out] $-1/12*\arctan(1/2*x*6^(1/2)/(-3*x^2-1)^(1/4))*6^(1/2)-1/12*\operatorname{arctanh}(1/2*x*6^(1/2)/(-3*x^2-1)^(1/4))*6^(1/2)$

Rubi [A] time = 0.01, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {398}

$$-\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{-3x^2-1}}\right)}{2\sqrt{6}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{-3x^2-1}}\right)}{2\sqrt{6}}$$

Antiderivative was successfully verified.

[In] Int[1/((-2 - 3*x^2)*(-1 - 3*x^2)^(1/4)),x]

[Out] $-\operatorname{ArcTan}[(\operatorname{Sqrt}[3/2]*x)/(-1 - 3*x^2)^(1/4)]/(2*\operatorname{Sqrt}[6]) - \operatorname{ArcTanh}[(\operatorname{Sqrt}[3/2]*x)/(-1 - 3*x^2)^(1/4)]/(2*\operatorname{Sqrt}[6])$

Rule 398

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(-2-3x^2)\sqrt[4]{-1-3x^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{-1-3x^2}}\right)}{2\sqrt{6}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{-1-3x^2}}\right)}{2\sqrt{6}}$$

Mathematica [C] time = 0.13, size = 127, normalized size = 2.08

$$\frac{2x F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -3x^2, -\frac{3x^2}{2}\right)}{\sqrt[4]{-3x^2-1} (3x^2+2) \left(x^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -3x^2, -\frac{3x^2}{2}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -3x^2, -\frac{3x^2}{2}\right)\right) - 2F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -3x^2, -\frac{3x^2}{2}\right)}\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2 - 3*x^2)*(-1 - 3*x^2)^(1/4)),x]

[Out] $(2*x*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, -3*x^2, (-3*x^2)/2])/((-1 - 3*x^2)^(1/4)*(2 + 3*x^2)*(-2*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, -3*x^2, (-3*x^2)/2] + x^2*(2*\operatorname{AppellF1}[3/2, 1/4, 2, 5/2, -3*x^2, (-3*x^2)/2] + \operatorname{AppellF1}[3/2, 5/4, 1, 5/2, -3*x^2, (-3*x^2)/2]))$

fricas [C] time = 4.43, size = 243, normalized size = 3.98

$$-\frac{1}{24} \sqrt{6} \log \left(\frac{\sqrt{6} \sqrt{-3x^2-1} x - \sqrt{6} x + 2(-3x^2-1)^{\frac{3}{4}} - 2(-3x^2-1)^{\frac{1}{4}}}{3(3x^2+2)} \right) + \frac{1}{24} \sqrt{6} \log \left(\frac{\sqrt{6} \sqrt{-3x^2-1} x - \sqrt{6} x}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2-2)/(-3*x^2-1)^(1/4),x, algorithm="fricas")

[Out] -1/24*sqrt(6)*log(1/3*(sqrt(6)*sqrt(-3*x^2 - 1)*x - sqrt(6)*x + 2*(-3*x^2 - 1)^(3/4) - 2*(-3*x^2 - 1)^(1/4))/(3*x^2 + 2)) + 1/24*sqrt(6)*log(-1/3*(sqrt(6)*sqrt(-3*x^2 - 1)*x - sqrt(6)*x - 2*(-3*x^2 - 1)^(3/4) + 2*(-3*x^2 - 1)^(1/4))/(3*x^2 + 2)) + 1/24*I*sqrt(6)*log(1/3*(I*sqrt(6)*sqrt(-3*x^2 - 1)*x + I*sqrt(6)*x + 2*(-3*x^2 - 1)^(3/4) + 2*(-3*x^2 - 1)^(1/4))/(3*x^2 + 2)) - 1/24*I*sqrt(6)*log(1/3*(-I*sqrt(6)*sqrt(-3*x^2 - 1)*x - I*sqrt(6)*x + 2*(-3*x^2 - 1)^(3/4) + 2*(-3*x^2 - 1)^(1/4))/(3*x^2 + 2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(3x^2+2)(-3x^2-1)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2-2)/(-3*x^2-1)^(1/4),x, algorithm="giac")

[Out] integrate(-1/((3*x^2 + 2)*(-3*x^2 - 1)^(1/4)), x)

maple [C] time = 1.11, size = 138, normalized size = 2.26

$$\frac{\text{RootOf}(-Z^2-6) \ln \left(\frac{3\sqrt{-3x^2-1} x - 3x + (-3x^2-1)^{\frac{3}{4}} \text{RootOf}(-Z^2-6) - (-3x^2-1)^{\frac{1}{4}} \text{RootOf}(-Z^2-6)}{3x^2+2} \right)}{12} + \frac{\text{RootOf}(-Z^2+6) \ln \left(-\frac{3\sqrt{-3x^2-1} x - 3x + (-3x^2-1)^{\frac{3}{4}} \text{RootOf}(-Z^2-6) - (-3x^2-1)^{\frac{1}{4}} \text{RootOf}(-Z^2-6)}{3x^2+2} \right)}{12}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*x^2-2)/(-3*x^2-1)^(1/4),x)

[Out] -1/12*RootOf(-Z^2-6)*ln((RootOf(-Z^2-6)*(-3*x^2-1)^(3/4)+3*(-3*x^2-1)^(1/2)*x-RootOf(-Z^2-6)*(-3*x^2-1)^(1/4)-3*x)/(3*x^2+2))+1/12*RootOf(-Z^2+6)*ln(-(RootOf(-Z^2+6)*(-3*x^2-1)^(3/4)-3*(-3*x^2-1)^(1/2)*x+RootOf(-Z^2+6)*(-3*x^2-1)^(1/4)-3*x)/(3*x^2+2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(3x^2+2)(-3x^2-1)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2-2)/(-3*x^2-1)^(1/4),x, algorithm="maxima")

[Out] -integrate(1/((3*x^2 + 2)*(-3*x^2 - 1)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$-\int \frac{1}{(-3x^2-1)^{\frac{1}{4}}(3x^2+2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(-1/((- 3*x^2 - 1)^(1/4)*(3*x^2 + 2)),x)`

[Out] `-int(1/((- 3*x^2 - 1)^(1/4)*(3*x^2 + 2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{3x^2\sqrt[4]{-3x^2-1} + 2\sqrt[4]{-3x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-3*x**2-2)/(-3*x**2-1)**(1/4),x)`

[Out] `-Integral(1/(3*x**2*(-3*x**2 - 1)**(1/4) + 2*(-3*x**2 - 1)**(1/4)), x)`

$$3.314 \quad \int \frac{1}{(-2+bx^2)\sqrt[4]{-1+bx^2}} dx$$

Optimal. Leaf size=77

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{bx^2-1}}\right)}{2\sqrt{2}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{bx^2-1}}\right)}{2\sqrt{2}\sqrt{b}}$$

[Out] $-1/4*\arctan(1/2*x*b^{(1/2)}/(b*x^2-1)^{(1/4)}*2^{(1/2)})*2^{(1/2)}/b^{(1/2)}-1/4*\arctanh(1/2*x*b^{(1/2)}/(b*x^2-1)^{(1/4)}*2^{(1/2)})*2^{(1/2)}/b^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {398}

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{bx^2-1}}\right)}{2\sqrt{2}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{bx^2-1}}\right)}{2\sqrt{2}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((-2 + b*x^2)*(-1 + b*x^2)^(1/4)),x]

[Out] $-\text{ArcTan}[(\text{Sqrt}[b]*x)/(\text{Sqrt}[2]*(-1 + b*x^2)^{(1/4)})]/(2*\text{Sqrt}[2]*\text{Sqrt}[b]) - \text{ArcTanh}[(\text{Sqrt}[b]*x)/(\text{Sqrt}[2]*(-1 + b*x^2)^{(1/4)})]/(2*\text{Sqrt}[2]*\text{Sqrt}[b])$

Rule 398

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4))]]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4))]]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(-2+bx^2)\sqrt[4]{-1+bx^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{-1+bx^2}}\right)}{2\sqrt{2}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{-1+bx^2}}\right)}{2\sqrt{2}\sqrt{b}}$$

Mathematica [C] time = 0.17, size = 132, normalized size = 1.71

$$\frac{6x F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; bx^2, \frac{bx^2}{2}\right)}{(bx^2 - 2)\sqrt[4]{bx^2 - 1} \left(bx^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; bx^2, \frac{bx^2}{2}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; bx^2, \frac{bx^2}{2}\right) \right) + 6F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; bx^2, \frac{bx^2}{2}\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2 + b*x^2)*(-1 + b*x^2)^(1/4)),x]

[Out] $(6*x*\text{AppellF1}[1/2, 1/4, 1, 3/2, b*x^2, (b*x^2)/2])/((-2 + b*x^2)*(-1 + b*x^2)^{(1/4)}*(6*\text{AppellF1}[1/2, 1/4, 1, 3/2, b*x^2, (b*x^2)/2] + b*x^2*(2*\text{AppellF1}[3/2, 1/4, 2, 5/2, b*x^2, (b*x^2)/2] + \text{AppellF1}[3/2, 5/4, 1, 5/2, b*x^2, (b*x^2)/2])))$

fricas [B] time = 13.27, size = 274, normalized size = 3.56

$$\left[\frac{2\sqrt{2}\sqrt{b}\arctan\left(\frac{\sqrt{2}(bx^2-1)^{\frac{1}{4}}}{\sqrt{bx}}\right) + \sqrt{2}\sqrt{b}\log\left(-\frac{b^2x^4-2\sqrt{2}(bx^2-1)^{\frac{1}{4}}b^{\frac{3}{2}}x^3+4\sqrt{bx^2-1}bx^2+4bx^2-4\sqrt{2}(bx^2-1)^{\frac{3}{4}}\sqrt{bx}-4}{b^2x^4-4bx^2+4}\right)}{8b} \right], \frac{2\sqrt{2}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2)/(b*x^2-1)^(1/4),x, algorithm="fricas")

[Out] [1/8*(2*sqrt(2)*sqrt(b)*arctan(sqrt(2)*(b*x^2 - 1)^(1/4)/(sqrt(b)*x)) + sqrt(2)*sqrt(b)*log(-(b^2*x^4 - 2*sqrt(2)*(b*x^2 - 1)^(1/4)*b^(3/2)*x^3 + 4*sqrt(b*x^2 - 1)*b*x^2 + 4*b*x^2 - 4*sqrt(2)*(b*x^2 - 1)^(3/4)*sqrt(b)*x - 4)/(b^2*x^4 - 4*b*x^2 + 4)))/b, 1/8*(2*sqrt(2)*sqrt(-b)*arctan(sqrt(2)*(b*x^2 - 1)^(1/4)*sqrt(-b)/(b*x)) - sqrt(2)*sqrt(-b)*log(-(b^2*x^4 + 2*sqrt(2)*(b*x^2 - 1)^(1/4)*sqrt(-b)*b*x^3 - 4*sqrt(b*x^2 - 1)*b*x^2 + 4*b*x^2 - 4*sqrt(2)*(b*x^2 - 1)^(3/4)*sqrt(-b)*x - 4)/(b^2*x^4 - 4*b*x^2 + 4)))/b]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 1)^{\frac{1}{4}}(bx^2 - 2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2)/(b*x^2-1)^(1/4),x, algorithm="giac")

[Out] integrate(1/((b*x^2 - 1)^(1/4)*(b*x^2 - 2)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 2)(bx^2 - 1)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2-2)/(b*x^2-1)^(1/4),x)

[Out] int(1/(b*x^2-2)/(b*x^2-1)^(1/4),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 1)^{\frac{1}{4}}(bx^2 - 2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2)/(b*x^2-1)^(1/4),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 - 1)^(1/4)*(b*x^2 - 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 - 1)^{1/4}(bx^2 - 2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((b*x^2 - 1)^(1/4)*(b*x^2 - 2)),x)`

[Out] `int(1/((b*x^2 - 1)^(1/4)*(b*x^2 - 2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 2)\sqrt[4]{bx^2 - 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x**2-2)/(b*x**2-1)**(1/4),x)`

[Out] `Integral(1/((b*x**2 - 2)*(b*x**2 - 1)**(1/4)), x)`

$$3.315 \quad \int \frac{1}{(-2-bx^2)\sqrt[4]{-1-bx^2}} dx$$

Optimal. Leaf size=79

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{-bx^2-1}}\right)}{2\sqrt{2}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{-bx^2-1}}\right)}{2\sqrt{2}\sqrt{b}}$$

[Out] $-1/4*\arctan(1/2*x*b^{(1/2)/(-b*x^2-1)^{(1/4)}*2^{(1/2)})*2^{(1/2)/b^{(1/2)}-1/4*\operatorname{arctanh}(1/2*x*b^{(1/2)/(-b*x^2-1)^{(1/4)}*2^{(1/2)})*2^{(1/2)/b^{(1/2)}}$

Rubi [A] time = 0.01, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {398}

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{-bx^2-1}}\right)}{2\sqrt{2}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{-bx^2-1}}\right)}{2\sqrt{2}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((-2 - b*x^2)*(-1 - b*x^2)^(1/4)),x]

[Out] $-\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/(\operatorname{Sqrt}[2]*(-1 - b*x^2)^{(1/4)})]/(2*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]) - \operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(\operatorname{Sqrt}[2]*(-1 - b*x^2)^{(1/4)})]/(2*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b])$

Rule 398

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(-2-bx^2)\sqrt[4]{-1-bx^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{-1-bx^2}}\right)}{2\sqrt{2}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{-1-bx^2}}\right)}{2\sqrt{2}\sqrt{b}}$$

Mathematica [C] time = 0.15, size = 137, normalized size = 1.73

$$\frac{6xF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -bx^2, -\frac{bx^2}{2}\right)}{\sqrt[4]{-bx^2-1} (bx^2+2) \left(bx^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -bx^2, -\frac{bx^2}{2}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -bx^2, -\frac{bx^2}{2}\right)\right) - 6F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -bx^2, -\frac{bx^2}{2}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2 - b*x^2)*(-1 - b*x^2)^(1/4)),x]

[Out] $(6*x*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, -(b*x^2), -1/2*(b*x^2)]/((-1 - b*x^2)^{(1/4)}*(2 + b*x^2)*(-6*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, -(b*x^2), -1/2*(b*x^2)] + b*x^2*2*(2*\operatorname{AppellF1}[3/2, 1/4, 2, 5/2, -(b*x^2), -1/2*(b*x^2)] + \operatorname{AppellF1}[3/2, 5/4, 1, 5/2, -(b*x^2), -1/2*(b*x^2)]))$

fricas [B] time = 13.27, size = 273, normalized size = 3.46

$$\left[\frac{2\sqrt{2}\sqrt{b}\arctan\left(\frac{\sqrt{2}(-bx^2-1)^{\frac{1}{4}}}{\sqrt{bx}}\right) + \sqrt{2}\sqrt{b}\log\left(\frac{b^2x^4+4\sqrt{-bx^2-1}bx^2-4bx^2-2\sqrt{2}\left((-bx^2-1)^{\frac{1}{4}}bx^3+2(-bx^2-1)^{\frac{3}{4}}x\right)\sqrt{b-4}}{b^2x^4+4bx^2+4}\right)}{8b}, \frac{2\sqrt{2}\sqrt{b}\arctan\left(\frac{\sqrt{2}(-bx^2-1)^{\frac{1}{4}}}{\sqrt{bx}}\right) + \sqrt{2}\sqrt{b}\log\left(\frac{b^2x^4+4\sqrt{-bx^2-1}bx^2-4bx^2-2\sqrt{2}\left((-bx^2-1)^{\frac{1}{4}}bx^3+2(-bx^2-1)^{\frac{3}{4}}x\right)\sqrt{b-4}}{b^2x^4+4bx^2+4}\right)}{8b} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-2)/(-b*x^2-1)^(1/4),x, algorithm="fricas")

[Out] [1/8*(2*sqrt(2)*sqrt(b)*arctan(sqrt(2)*(-b*x^2 - 1)^(1/4)/(sqrt(b)*x)) + sqrt(2)*sqrt(b)*log(-(b^2*x^4 + 4*sqrt(-b*x^2 - 1)*b*x^2 - 4*b*x^2 - 2*sqrt(2)*((-b*x^2 - 1)^(1/4)*b*x^3 + 2*(-b*x^2 - 1)^(3/4)*x)*sqrt(b) - 4)/(b^2*x^4 + 4*b*x^2 + 4))/b, 1/8*(2*sqrt(2)*sqrt(-b)*arctan(sqrt(2)*(-b*x^2 - 1)^(1/4)*sqrt(-b)/(b*x)) - sqrt(2)*sqrt(-b)*log(-(b^2*x^4 - 4*sqrt(-b*x^2 - 1)*b*x^2 - 4*b*x^2 + 2*sqrt(2)*((-b*x^2 - 1)^(1/4)*b*x^3 - 2*(-b*x^2 - 1)^(3/4)*x)*sqrt(-b) - 4)/(b^2*x^4 + 4*b*x^2 + 4))/b]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(bx^2 + 2)(-bx^2 - 1)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-2)/(-b*x^2-1)^(1/4),x, algorithm="giac")

[Out] integrate(-1/((b*x^2 + 2)*(-b*x^2 - 1)^(1/4)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 - 2)(-bx^2 - 1)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2-2)/(-b*x^2-1)^(1/4),x)

[Out] int(1/(-b*x^2-2)/(-b*x^2-1)^(1/4),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(bx^2 + 2)(-bx^2 - 1)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-2)/(-b*x^2-1)^(1/4),x, algorithm="maxima")

[Out] -integrate(1/((b*x^2 + 2)*(-b*x^2 - 1)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(-bx^2 - 1)^{\frac{1}{4}}(bx^2 + 2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(-1/((- b*x^2 - 1)^(1/4)*(b*x^2 + 2)), x)`

[Out] `-int(1/((- b*x^2 - 1)^(1/4)*(b*x^2 + 2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{bx^2\sqrt[4]{-bx^2-1} + 2\sqrt[4]{-bx^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-b*x**2-2)/(-b*x**2-1)**(1/4), x)`

[Out] `-Integral(1/(b*x**2*(-b*x**2 - 1)**(1/4) + 2*(-b*x**2 - 1)**(1/4)), x)`

$$3.316 \quad \int \frac{1}{(-2a+3x^2)\sqrt[4]{-a+3x^2}} dx$$

Optimal. Leaf size=85

$$-\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{3x^2-a}}\right)}{2\sqrt{6}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{3x^2-a}}\right)}{2\sqrt{6}a^{3/4}}$$

[Out] -1/12*arctan(1/2*x*6^(1/2)/a^(1/4)/(3*x^2-a)^(1/4))/a^(3/4)*6^(1/2)-1/12*arctanh(1/2*x*6^(1/2)/a^(1/4)/(3*x^2-a)^(1/4))/a^(3/4)*6^(1/2)

Rubi [A] time = 0.02, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {398}

$$-\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{3x^2-a}}\right)}{2\sqrt{6}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{3x^2-a}}\right)}{2\sqrt{6}a^{3/4}}$$

Antiderivative was successfully verified.

[In] Int[1/((-2*a + 3*x^2)*(-a + 3*x^2)^(1/4)),x]

[Out] -ArcTan[(Sqrt[3/2]*x)/(a^(1/4)*(-a + 3*x^2)^(1/4))]/(2*Sqrt[6]*a^(3/4)) - ArcTanh[(Sqrt[3/2]*x)/(a^(1/4)*(-a + 3*x^2)^(1/4))]/(2*Sqrt[6]*a^(3/4))

Rule 398

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(-2a + 3x^2)\sqrt[4]{-a + 3x^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{-a+3x^2}}\right)}{2\sqrt{6}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{-a+3x^2}}\right)}{2\sqrt{6}a^{3/4}}$$

Mathematica [C] time = 0.17, size = 157, normalized size = 1.85

$$\frac{2axF_1\left(\frac{1}{2}, \frac{1}{4}, 1; \frac{3}{2}, \frac{3x^2}{a}, \frac{3x^2}{2a}\right)}{(3x^2 - 2a)\sqrt[4]{3x^2 - a} \left(x^2 \left(2F_1\left(\frac{3}{2}, \frac{1}{4}, 2; \frac{5}{2}, \frac{3x^2}{a}, \frac{3x^2}{2a}\right) + F_1\left(\frac{3}{2}, \frac{5}{4}, 1; \frac{5}{2}, \frac{3x^2}{a}, \frac{3x^2}{2a}\right)\right) + 2aF_1\left(\frac{1}{2}, \frac{1}{4}, 1; \frac{3}{2}, \frac{3x^2}{a}, \frac{3x^2}{2a}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2*a + 3*x^2)*(-a + 3*x^2)^(1/4)),x]

[Out] (2*a*x*AppellF1[1/2, 1/4, 1, 3/2, (3*x^2)/a, (3*x^2)/(2*a)]/((-2*a + 3*x^2)*(-a + 3*x^2)^(1/4)*(2*a*AppellF1[1/2, 1/4, 1, 3/2, (3*x^2)/a, (3*x^2)/(2*a)] + x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, (3*x^2)/a, (3*x^2)/(2*a)] + AppellF1[3/2, 5/4, 1, 5/2, (3*x^2)/a, (3*x^2)/(2*a)]))

fricas [B] time = 9.49, size = 276, normalized size = 3.25

$$-\left(\frac{1}{36}\right)^{\frac{1}{4}} \frac{1}{a^{\frac{1}{4}}} \arctan \left(\frac{2 \left(\sqrt{\frac{1}{2}} \left(6 \left(\frac{1}{36} \right)^{\frac{3}{4}} a^3 \frac{1}{a^{\frac{3}{4}}} + \left(\frac{1}{36} \right)^{\frac{1}{4}} \sqrt{3x^2 - a} a \frac{1}{a^{\frac{1}{4}}} \right) \sqrt{a} \sqrt{\frac{1}{a^3}} - \left(\frac{1}{36} \right)^{\frac{1}{4}} (3x^2 - a)^{\frac{1}{4}} a \frac{1}{a^{\frac{1}{4}}} \right)}{x} \right) - \frac{1}{4} \left(\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2-2*a)/(3*x^2-a)^(1/4),x, algorithm="fricas")

[Out] $-(1/36)^{(1/4)}*(a^{(-3)})^{(1/4)}*\arctan(2*(\sqrt{1/2}*(6*(1/36)^{(3/4)}*a^3*(a^{(-3)})^{(3/4)} + (1/36)^{(1/4)}*\sqrt{3*x^2 - a}*a*(a^{(-3)})^{(1/4)})*\sqrt{a*\sqrt{a^{(-3)}}}) - (1/36)^{(1/4)}*(3*x^2 - a)^{(1/4)}*a*(a^{(-3)})^{(1/4)})/x) - 1/4*(1/36)^{(1/4)}*(a^{(-3)})^{(1/4)}*\log((18*(1/36)^{(3/4)}*\sqrt{3*x^2 - a}*a^2*(a^{(-3)})^{(3/4)}*x + (3*x^2 - a)^{(1/4)}*a^2*\sqrt{a^{(-3)}} + 3*(1/36)^{(1/4)}*a*(a^{(-3)})^{(1/4)}*x + (3*x^2 - a)^{(3/4)})/(3*x^2 - 2*a)) + 1/4*(1/36)^{(1/4)}*(a^{(-3)})^{(1/4)}*\log(-18*(1/36)^{(3/4)}*\sqrt{3*x^2 - a}*a^2*(a^{(-3)})^{(3/4)}*x - (3*x^2 - a)^{(1/4)}*a^2*\sqrt{a^{(-3)}} + 3*(1/36)^{(1/4)}*a*(a^{(-3)})^{(1/4)}*x - (3*x^2 - a)^{(3/4)})/(3*x^2 - 2*a))$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 - a)^{\frac{1}{4}}(3x^2 - 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2-2*a)/(3*x^2-a)^(1/4),x, algorithm="giac")

[Out] integrate(1/((3*x^2 - a)^(1/4)*(3*x^2 - 2*a)), x)

maple [F] time = 0.35, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 - 2a)(3x^2 - a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(3*x^2-2*a)/(3*x^2-a)^(1/4),x)

[Out] int(1/(3*x^2-2*a)/(3*x^2-a)^(1/4),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(3x^2 - a)^{\frac{1}{4}}(3x^2 - 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(3*x^2-2*a)/(3*x^2-a)^(1/4),x, algorithm="maxima")

[Out] integrate(1/((3*x^2 - a)^(1/4)*(3*x^2 - 2*a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(2a - 3x^2)(3x^2 - a)^{1/4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(-1/((2*a - 3*x^2)*(3*x^2 - a)^(1/4)), x)`

[Out] `-int(1/((2*a - 3*x^2)*(3*x^2 - a)^(1/4)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-2a + 3x^2) \sqrt[4]{-a + 3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(3*x**2-2*a)/(3*x**2-a)**(1/4), x)`

[Out] `Integral(1/((-2*a + 3*x**2)*(-a + 3*x**2)**(1/4)), x)`

$$3.317 \quad \int \frac{1}{(-2a-3x^2)\sqrt[4]{-a-3x^2}} dx$$

Optimal. Leaf size=85

$$-\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{-a-3x^2}}\right)}{2\sqrt{6}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{-a-3x^2}}\right)}{2\sqrt{6}a^{3/4}}$$

[Out] $-1/12*\arctan(1/2*x*6^(1/2)/a^(1/4)/(-3*x^2-a)^(1/4))/a^(3/4)*6^(1/2)-1/12*arctanh(1/2*x*6^(1/2)/a^(1/4)/(-3*x^2-a)^(1/4))/a^(3/4)*6^(1/2)$

Rubi [A] time = 0.01, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {398}

$$-\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{-a-3x^2}}\right)}{2\sqrt{6}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{-a-3x^2}}\right)}{2\sqrt{6}a^{3/4}}$$

Antiderivative was successfully verified.

[In] Int[1/((-2*a - 3*x^2)*(-a - 3*x^2)^(1/4)),x]

[Out] $-\text{ArcTan}[(\text{Sqrt}[3/2]*x)/(a^(1/4)*(-a - 3*x^2)^(1/4))]/(2*\text{Sqrt}[6]*a^(3/4)) - \text{ArcTanh}[(\text{Sqrt}[3/2]*x)/(a^(1/4)*(-a - 3*x^2)^(1/4))]/(2*\text{Sqrt}[6]*a^(3/4))$

Rule 398

Int[1/(((a_) + (b_)*(x_)^2)^(1/4)*((c_) + (d_)*(x_)^2)), x_Symbol] := With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4))]]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4))]]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(-2a-3x^2)\sqrt[4]{-a-3x^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{-a-3x^2}}\right)}{2\sqrt{6}a^{3/4}} - \frac{\tanh^{-1}\left(\frac{\sqrt{\frac{3}{2}}x}{\sqrt[4]{a}\sqrt[4]{-a-3x^2}}\right)}{2\sqrt{6}a^{3/4}}$$

Mathematica [C] time = 0.15, size = 157, normalized size = 1.85

$$\frac{2axF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{3x^2}{a}, -\frac{3x^2}{2a}\right)}{\sqrt[4]{-a-3x^2}(2a+3x^2)\left(x^2\left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{3x^2}{a}, -\frac{3x^2}{2a}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{3x^2}{a}, -\frac{3x^2}{2a}\right)\right) - 2aF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{3x^2}{a}, -\frac{3x^2}{2a}\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2*a - 3*x^2)*(-a - 3*x^2)^(1/4)),x]

[Out] $(2*a*x*\text{AppellF1}[1/2, 1/4, 1, 3/2, (-3*x^2)/a, (-3*x^2)/(2*a)])/((-a - 3*x^2)^(1/4)*(2*a + 3*x^2)*(-2*a*\text{AppellF1}[1/2, 1/4, 1, 3/2, (-3*x^2)/a, (-3*x^2)/(2*a)]/(2*a)] + x^2*(2*\text{AppellF1}[3/2, 1/4, 2, 5/2, (-3*x^2)/a, (-3*x^2)/(2*a)] + \text{AppellF1}[3/2, 5/4, 1, 5/2, (-3*x^2)/a, (-3*x^2)/(2*a)]))$

fricas [B] time = 9.42, size = 278, normalized size = 3.27

$$-\left(\frac{1}{36}\right)^{\frac{1}{4}} \frac{1}{a^{\frac{3}{4}}} \arctan \left(\frac{2 \left(\sqrt{\frac{1}{2}} \left(6 \left(\frac{1}{36} \right)^{\frac{3}{4}} a^{\frac{3}{4}} - \left(\frac{1}{36} \right)^{\frac{1}{4}} \sqrt{-3x^2 - a} a^{\frac{1}{4}} \right) \sqrt{-a} \sqrt{\frac{1}{a^3}} - \left(\frac{1}{36} \right)^{\frac{1}{4}} (-3x^2 - a)^{\frac{1}{4}} a^{\frac{1}{4}} \right)}{x} \right) + \frac{1}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2-2*a)/(-3*x^2-a)^(1/4),x, algorithm="fricas")

[Out] $-(1/36)^{1/4} * (a^{-3})^{1/4} * \arctan(2 * (\sqrt{1/2}) * (6 * (1/36)^{3/4} * a^{3/4} * (a^{-3})^{1/4}) - (1/36)^{1/4} * \sqrt{-3x^2 - a} * a^{1/4}) * \sqrt{-a} * \sqrt{1/a^3} - (1/36)^{1/4} * (-3x^2 - a)^{1/4} * a^{1/4}) / x + 1/4 * (1/36)^{1/4} * (a^{-3})^{1/4} * \log(-18 * (1/36)^{3/4} * \sqrt{-3x^2 - a} * a^{2/4} * (a^{-3})^{3/4} * x + (-3x^2 - a)^{1/4} * a^{2/4} * \sqrt{a^{-3}} - 3 * (1/36)^{1/4} * a * (a^{-3})^{1/4} * x - (-3x^2 - a)^{3/4}) / (3x^2 + 2a)) - 1/4 * (1/36)^{1/4} * (a^{-3})^{1/4} * \log((18 * (1/36)^{3/4} * \sqrt{-3x^2 - a} * a^{2/4} * (a^{-3})^{3/4} * x - (-3x^2 - a)^{1/4} * a^{2/4} * \sqrt{a^{-3}} - 3 * (1/36)^{1/4} * a * (a^{-3})^{1/4} * x + (-3x^2 - a)^{3/4}) / (3x^2 + 2a))$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(3x^2 + 2a)(-3x^2 - a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2-2*a)/(-3*x^2-a)^(1/4),x, algorithm="giac")

[Out] integrate(-1/((3*x^2 + 2*a)*(-3*x^2 - a)^(1/4)), x)

maple [F] time = 0.36, size = 0, normalized size = 0.00

$$\int \frac{1}{(-3x^2 - 2a)(-3x^2 - a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-3*x^2-2*a)/(-3*x^2-a)^(1/4),x)

[Out] int(1/(-3*x^2-2*a)/(-3*x^2-a)^(1/4),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(3x^2 + 2a)(-3x^2 - a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-3*x^2-2*a)/(-3*x^2-a)^(1/4),x, algorithm="maxima")

[Out] -integrate(1/((3*x^2 + 2*a)*(-3*x^2 - a)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(3x^2 + 2a)(-3x^2 - a)^{1/4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(-1/((2*a + 3*x^2)*(- a - 3*x^2)^(1/4)), x)`

[Out] `-int(1/((2*a + 3*x^2)*(- a - 3*x^2)^(1/4)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{2a\sqrt[4]{-a-3x^2} + 3x^2\sqrt[4]{-a-3x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-3*x**2-2*a)/(-3*x**2-a)**(1/4), x)`

[Out] `-Integral(1/(2*a*(-a - 3*x**2)**(1/4) + 3*x**2*(-a - 3*x**2)**(1/4)), x)`

$$3.318 \quad \int \frac{1}{(-2a+bx^2)\sqrt[4]{-a+bx^2}} dx$$

Optimal. Leaf size=101

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{bx^2-a}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{bx^2-a}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}}$$

[Out] $-1/4*\arctan(1/2*x*b^{(1/2)}/a^{(1/4)}/(b*x^2-a)^{(1/4)}*2^{(1/2)})/a^{(3/4)}*2^{(1/2)}/b^{(1/2)}-1/4*\operatorname{arctanh}(1/2*x*b^{(1/2)}/a^{(1/4)}/(b*x^2-a)^{(1/4)}*2^{(1/2)})/a^{(3/4)}*2^{(1/2)}/b^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {398}

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{bx^2-a}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{bx^2-a}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((-2*a + b*x^2)*(-a + b*x^2)^(1/4)),x]

[Out] $-\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/(\operatorname{Sqrt}[2]*a^{(1/4)}*(-a + b*x^2)^{(1/4)})]/(2*\operatorname{Sqrt}[2]*a^{(3/4)}*\operatorname{Sqrt}[b]) - \operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(\operatorname{Sqrt}[2]*a^{(1/4)}*(-a + b*x^2)^{(1/4)})]/(2*\operatorname{Sqrt}[2]*a^{(3/4)}*\operatorname{Sqrt}[b])$

Rule 398

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(-2a + bx^2)\sqrt[4]{-a + bx^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{-a+bx^2}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{-a+bx^2}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}}$$

Mathematica [C] time = 0.16, size = 163, normalized size = 1.61

$$\frac{6axF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{bx^2}{a}, \frac{bx^2}{2a}\right)}{(2a - bx^2)\sqrt[4]{bx^2 - a} \left(bx^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; \frac{bx^2}{a}, \frac{bx^2}{2a}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; \frac{bx^2}{a}, \frac{bx^2}{2a}\right) \right) + 6aF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; \frac{bx^2}{a}, \frac{bx^2}{2a}\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2*a + b*x^2)*(-a + b*x^2)^(1/4)),x]

[Out] $(-6*a*x*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, (b*x^2)/a, (b*x^2)/(2*a)])/((2*a - b*x^2)*(-a + b*x^2)^{(1/4)}*(6*a*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, (b*x^2)/a, (b*x^2)/(2*$

a)] + b*x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, (b*x^2)/a, (b*x^2)/(2*a)] + AppellF1[3/2, 5/4, 1, 5/2, (b*x^2)/a, (b*x^2)/(2*a)]))

fricas [B] time = 50.67, size = 338, normalized size = 3.35

$$-\left(\frac{1}{4}\right)^{\frac{1}{4}} \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \arctan \left(\frac{2 \left(\sqrt{\frac{1}{2}} \left(2 \left(\frac{1}{4}\right)^{\frac{3}{4}} a^3 b \left(\frac{1}{a^3 b^2}\right)^{\frac{3}{4}} + \left(\frac{1}{4}\right)^{\frac{1}{4}} \sqrt{bx^2 - a} a \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \right) \sqrt{ab \sqrt{\frac{1}{a^3 b^2}} - \left(\frac{1}{4}\right)^{\frac{1}{4}} (bx^2 - a)^{\frac{1}{4}}}}{x} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2*a)/(b*x^2-a)^(1/4),x, algorithm="fricas")

[Out] $-(1/4)^{(1/4)} * (1/(a^3*b^2))^{(1/4)} * \arctan(2*(\sqrt{1/2} * (2*(1/4)^{(3/4)} * a^3*b * (1/(a^3*b^2))^{(3/4)} + (1/4)^{(1/4)} * \sqrt{b*x^2 - a} * a * (1/(a^3*b^2))^{(1/4)} * \sqrt{a*b*\sqrt{1/(a^3*b^2))}) - (1/4)^{(1/4)} * (b*x^2 - a)^{(1/4)} * a * (1/(a^3*b^2))^{(1/4)})/x - 1/4*(1/4)^{(1/4)} * (1/(a^3*b^2))^{(1/4)} * \log((2*(1/4)^{(3/4)} * \sqrt{b*x^2 - a} * a^2*b^2*x * (1/(a^3*b^2))^{(3/4)} + (b*x^2 - a)^{(1/4)} * a^2*b*\sqrt{1/(a^3*b^2)}) + (1/4)^{(1/4)} * a*b*x * (1/(a^3*b^2))^{(1/4)} + (b*x^2 - a)^{(3/4)})/(b*x^2 - 2*a)) + 1/4*(1/4)^{(1/4)} * (1/(a^3*b^2))^{(1/4)} * \log(-(2*(1/4)^{(3/4)} * \sqrt{b*x^2 - a} * a^2*b^2*x * (1/(a^3*b^2))^{(3/4)} - (b*x^2 - a)^{(1/4)} * a^2*b*\sqrt{1/(a^3*b^2)}) + (1/4)^{(1/4)} * a*b*x * (1/(a^3*b^2))^{(1/4)} - (b*x^2 - a)^{(3/4)})/(b*x^2 - 2*a))$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - a)^{\frac{1}{4}}(bx^2 - 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2*a)/(b*x^2-a)^(1/4),x, algorithm="giac")

[Out] integrate(1/((b*x^2 - a)^(1/4)*(b*x^2 - 2*a)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - 2a)(bx^2 - a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2-2*a)/(b*x^2-a)^(1/4),x)

[Out] int(1/(b*x^2-2*a)/(b*x^2-a)^(1/4),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 - a)^{\frac{1}{4}}(bx^2 - 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2-2*a)/(b*x^2-a)^(1/4),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 - a)^(1/4)*(b*x^2 - 2*a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(bx^2 - a)^{1/4} (2a - bx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((b*x^2 - a)^(1/4)*(2*a - b*x^2)), x)

[Out] -int(1/((b*x^2 - a)^(1/4)*(2*a - b*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-2a + bx^2) \sqrt[4]{-a + bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2-2*a)/(b*x**2-a)**(1/4), x)

[Out] Integral(1/((-2*a + b*x**2)*(-a + b*x**2)**(1/4)), x)

$$3.319 \quad \int \frac{1}{(-2a-bx^2)\sqrt[4]{-a-bx^2}} dx$$

Optimal. Leaf size=103

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{-a-bx^2}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{-a-bx^2}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}}$$

[Out] $-1/4*\arctan(1/2*x*b^{(1/2)}/a^{(1/4)}/(-b*x^2-a)^{(1/4)}*2^{(1/2)})/a^{(3/4)}*2^{(1/2)}/b^{(1/2)}-1/4*\operatorname{arctanh}(1/2*x*b^{(1/2)}/a^{(1/4)}/(-b*x^2-a)^{(1/4)}*2^{(1/2)})/a^{(3/4)}*2^{(1/2)}/b^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.037$, Rules used = {398}

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{-a-bx^2}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{-a-bx^2}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}}$$

Antiderivative was successfully verified.

[In] Int[1/((-2*a - b*x^2)*(-a - b*x^2)^(1/4)),x]

[Out] $-\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/(\operatorname{Sqrt}[2]*a^{(1/4)}*(-a - b*x^2)^{(1/4)})]/(2*\operatorname{Sqrt}[2]*a^{(3/4)}*\operatorname{Sqrt}[b]) - \operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*x)/(\operatorname{Sqrt}[2]*a^{(1/4)}*(-a - b*x^2)^{(1/4)})]/(2*\operatorname{Sqrt}[2]*a^{(3/4)}*\operatorname{Sqrt}[b])$

Rule 398

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4)))]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(-2a-bx^2)\sqrt[4]{-a-bx^2}} dx = -\frac{\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{-a-bx^2}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{b}x}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{-a-bx^2}}\right)}{2\sqrt{2}a^{3/4}\sqrt{b}}$$

Mathematica [C] time = 0.16, size = 168, normalized size = 1.63

$$\frac{6axF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{2a}\right)}{\sqrt[4]{-a-bx^2}(2a+bx^2)\left(6aF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{2a}\right) - bx^2\left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{2a}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{bx^2}{2a}\right)\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((-2*a - b*x^2)*(-a - b*x^2)^(1/4)),x]

[Out] $(-6*a*x*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -1/2*(b*x^2)/a])/((-a - b*x^2)^{(1/4)}*(2*a + b*x^2)*(6*a*\operatorname{AppellF1}[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -1/2$

$\frac{b^2 x^2}{a} - b^2 x^2 (2 \operatorname{AppellF1}[3/2, 1/4, 2, 5/2, -((b^2 x^2)/a), -1/2(b^2 x^2)/a] + \operatorname{AppellF1}[3/2, 5/4, 1, 5/2, -((b^2 x^2)/a), -1/2(b^2 x^2)/a])$

fricas [B] time = 56.41, size = 350, normalized size = 3.40

$$-\left(\frac{1}{4}\right)^{\frac{1}{4}} \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \arctan \left(\frac{2 \left(\sqrt{\frac{1}{2}} \left(2 \left(\frac{1}{4}\right)^{\frac{3}{4}} a^3 b \left(\frac{1}{a^3 b^2}\right)^{\frac{3}{4}} - \left(\frac{1}{4}\right)^{\frac{1}{4}} \sqrt{-bx^2 - a} a \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \right) \sqrt{-ab} \sqrt{\frac{1}{a^3 b^2}} - \left(\frac{1}{4}\right)^{\frac{1}{4}} (-bx^2 - a)^{\frac{1}{4}}}{x} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-2*a)/(-b*x^2-a)^(1/4),x, algorithm="fricas")

[Out] $-\left(\frac{1}{4}\right)^{\frac{1}{4}} \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \arctan\left(2 \left(\sqrt{\frac{1}{2}}\right)^{\frac{3}{4}} a^3 b \left(\frac{1}{a^3 b^2}\right)^{\frac{3}{4}} - \left(\frac{1}{4}\right)^{\frac{1}{4}} \sqrt{-bx^2 - a} a \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}}\right) \sqrt{-ab} \sqrt{\frac{1}{a^3 b^2}} - \left(\frac{1}{4}\right)^{\frac{1}{4}} (-bx^2 - a)^{\frac{1}{4}}$
 $\frac{1}{x} + \frac{1}{4} \left(\frac{1}{4}\right)^{\frac{1}{4}} \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \log\left(-2 \left(\frac{1}{4}\right)^{\frac{3}{4}} \sqrt{-bx^2 - a} a^2 b^2 x \left(\frac{1}{a^3 b^2}\right)^{\frac{3}{4}} + (-bx^2 - a)^{\frac{1}{4}} a^2 b \sqrt{1/(a^3 b^2)} - \left(\frac{1}{4}\right)^{\frac{1}{4}} a b x \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} - (-bx^2 - a)^{\frac{3}{4}}\right) / (bx^2 + 2a)$
 $- \frac{1}{4} \left(\frac{1}{4}\right)^{\frac{1}{4}} \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} \log\left(2 \left(\frac{1}{4}\right)^{\frac{3}{4}} \sqrt{-bx^2 - a} a^2 b^2 x \left(\frac{1}{a^3 b^2}\right)^{\frac{3}{4}} - (-bx^2 - a)^{\frac{1}{4}} a^2 b \sqrt{1/(a^3 b^2)} - \left(\frac{1}{4}\right)^{\frac{1}{4}} a b x \left(\frac{1}{a^3 b^2}\right)^{\frac{1}{4}} + (-bx^2 - a)^{\frac{3}{4}}\right) / (bx^2 + 2a)$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(bx^2 + 2a)(-bx^2 - a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-2*a)/(-b*x^2-a)^(1/4),x, algorithm="giac")

[Out] integrate(-1/((b*x^2 + 2*a)*(-b*x^2 - a)^(1/4)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(-bx^2 - 2a)(-bx^2 - a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-b*x^2-2*a)/(-b*x^2-a)^(1/4),x)

[Out] int(1/(-b*x^2-2*a)/(-b*x^2-a)^(1/4),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(bx^2 + 2a)(-bx^2 - a)^{\frac{1}{4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x^2-2*a)/(-b*x^2-a)^(1/4),x, algorithm="maxima")

[Out] -integrate(1/((b*x^2 + 2*a)*(-b*x^2 - a)^(1/4)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$-\int \frac{1}{(-bx^2 - a)^{1/4} (bx^2 + 2a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((- a - b*x^2)^(1/4)*(2*a + b*x^2)), x)

[Out] -int(1/((- a - b*x^2)^(1/4)*(2*a + b*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{2a\sqrt[4]{-a - bx^2} + bx^2\sqrt[4]{-a - bx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-b*x**2-2*a)/(-b*x**2-a)**(1/4), x)

[Out] -Integral(1/(2*a*(-a - b*x**2)**(1/4) + b*x**2*(-a - b*x**2)**(1/4)), x)

$$3.320 \quad \int \frac{1}{(2-x^2)\sqrt[4]{-1+x^2}} dx$$

Optimal. Leaf size=53

$$\frac{\tan^{-1}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} + \frac{\tanh^{-1}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}}$$

[Out] 1/4*arctan(1/2*x/(x^2-1)^(1/4)*2^(1/2))*2^(1/2)+1/4*arctanh(1/2*x/(x^2-1)^(1/4)*2^(1/2))*2^(1/2)

Rubi [A] time = 0.01, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {398}

$$\frac{\tan^{-1}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} + \frac{\tanh^{-1}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[1/((2 - x^2)*(-1 + x^2)^(1/4)),x]

[Out] ArcTan[x/(Sqrt[2]*(-1 + x^2)^(1/4))]/(2*Sqrt[2]) + ArcTanh[x/(Sqrt[2]*(-1 + x^2)^(1/4))]/(2*Sqrt[2])

Rule 398

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With[{q = Rt[-(b^2/a), 4]}, Simp[(b*ArcTan[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4))]]/(2*Sqrt[2]*a*d*q), x] + Simp[(b*ArcTanh[(q*x)/(Sqrt[2]*(a + b*x^2)^(1/4))]]/(2*Sqrt[2]*a*d*q), x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] & NegQ[b^2/a]

Rubi steps

$$\int \frac{1}{(2-x^2)\sqrt[4]{-1+x^2}} dx = \frac{\tan^{-1}\left(\frac{x}{\sqrt{2}\sqrt[4]{-1+x^2}}\right)}{2\sqrt{2}} + \frac{\tanh^{-1}\left(\frac{x}{\sqrt{2}\sqrt[4]{-1+x^2}}\right)}{2\sqrt{2}}$$

Mathematica [C] time = 0.14, size = 115, normalized size = 2.17

$$\frac{6x F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; x^2, \frac{x^2}{2}\right)}{(x^2 - 2)\sqrt[4]{x^2 - 1} \left(x^2 \left(2F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; x^2, \frac{x^2}{2}\right) + F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; x^2, \frac{x^2}{2}\right)\right) + 6F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; x^2, \frac{x^2}{2}\right)}\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((2 - x^2)*(-1 + x^2)^(1/4)),x]

[Out] (-6*x*AppellF1[1/2, 1/4, 1, 3/2, x^2, x^2/2])/((-2 + x^2)*(-1 + x^2)^(1/4)*(6*AppellF1[1/2, 1/4, 1, 3/2, x^2, x^2/2] + x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, x^2, x^2/2] + AppellF1[3/2, 5/4, 1, 5/2, x^2, x^2/2])))

fricas [B] time = 3.96, size = 91, normalized size = 1.72

$$-\frac{1}{4}\sqrt{2}\arctan\left(\frac{\sqrt{2}(x^2-1)^{\frac{1}{4}}}{x}\right)+\frac{1}{8}\sqrt{2}\log\left(\frac{x^4+2\sqrt{2}(x^2-1)^{\frac{1}{4}}x^3+4\sqrt{x^2-1}x^2+4\sqrt{2}(x^2-1)^{\frac{3}{4}}x+4x^2-4}{x^4-4x^2+4}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)/(x^2-1)^(1/4),x, algorithm="fricas")

[Out] -1/4*sqrt(2)*arctan(sqrt(2)*(x^2 - 1)^(1/4)/x) + 1/8*sqrt(2)*log(-(x^4 + 2*sqrt(2)*(x^2 - 1)^(1/4)*x^3 + 4*sqrt(x^2 - 1)*x^2 + 4*sqrt(2)*(x^2 - 1)^(3/4)*x + 4*x^2 - 4)/(x^4 - 4*x^2 + 4))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{1}{(x^2-1)^{\frac{1}{4}}(x^2-2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)/(x^2-1)^(1/4),x, algorithm="giac")

[Out] integrate(-1/((x^2 - 1)^(1/4)*(x^2 - 2)), x)

maple [C] time = 1.22, size = 121, normalized size = 2.28

$$\frac{\text{RootOf}(-Z^2-2)\ln\left(\frac{-\sqrt{x^2-1}x-x+(x^2-1)^{\frac{3}{4}}\text{RootOf}(-Z^2-2)+(x^2-1)^{\frac{1}{4}}\text{RootOf}(-Z^2-2)}{x^2-2}\right)}{4} - \frac{\text{RootOf}(-Z^2+2)\ln\left(\frac{-\sqrt{x^2-1}}{x^2-2}\right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-x^2+2)/(x^2-1)^(1/4),x)

[Out] -1/4*RootOf(-Z^2-2)*ln(-(RootOf(-Z^2-2)*(x^2-1)^(3/4)-(x^2-1)^(1/2)*x+RootOf(-Z^2-2)*(x^2-1)^(1/4)-x)/(x^2-2))-1/4*RootOf(-Z^2+2)*ln(-(RootOf(-Z^2+2)*(x^2-1)^(3/4)-(x^2-1)^(1/2)*x-RootOf(-Z^2+2)*(x^2-1)^(1/4)+x)/(x^2-2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{(x^2-1)^{\frac{1}{4}}(x^2-2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x^2+2)/(x^2-1)^(1/4),x, algorithm="maxima")

[Out] -integrate(1/((x^2 - 1)^(1/4)*(x^2 - 2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$-\int \frac{1}{(x^2-1)^{\frac{1}{4}}(x^2-2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(-1/((x^2 - 1)^(1/4)*(x^2 - 2)),x)

[Out] -int(1/((x^2 - 1)^(1/4)*(x^2 - 2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{x^2 \sqrt[4]{x^2-1} - 2\sqrt[4]{x^2-1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-x**2+2)/(x**2-1)**(1/4),x)

[Out] -Integral(1/(x**2*(x**2 - 1)**(1/4) - 2*(x**2 - 1)**(1/4)), x)

$$3.321 \quad \int \frac{(a+bx^2)^{7/4}}{c+dx^2} dx$$

Optimal. Leaf size=362

$$\frac{6a^{3/2}\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{5d\sqrt[4]{a+bx^2}} + \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}(ad-bc)^{3/2}\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle|-1\right)}{d^{5/2}x} - \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}}{d^{5/2}x}$$

[Out] $6/5*a*b*x/d/(b*x^2+a)^{(1/4)}-2*b*(-a*d+b*c)*x/d^2/(b*x^2+a)^{(1/4)}+2/5*b*x*(b*x^2+a)^{(3/4)}/d-6/5*a^{(3/2)}*(1+b*x^2/a)^{(1/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*b^{(1/2)}/d/(b*x^2+a)^{(1/4)}+2*(-a*d+b*c)*(1+b*x^2/a)^{(1/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*a^{(1/2)}*b^{(1/2)}/d^2/(b*x^2+a)^{(1/4)}+a^{(1/4)}*(a*d-b*c)^{(3/2)}*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/d^{(5/2)}/x-a^{(1/4)}*(a*d-b*c)^{(3/2)}*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/d^{(5/2)}/x$

Rubi [A] time = 0.27, antiderivative size = 362, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {402, 195, 229, 227, 196, 399, 490, 1218}

$$\frac{6a^{3/2}\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{5d\sqrt[4]{a+bx^2}} - \frac{2bx(bc-ad)}{d^2\sqrt[4]{a+bx^2}} + \frac{2\sqrt{a}\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}(bc-ad)E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{d^2\sqrt[4]{a+bx^2}} + \frac{\sqrt[4]{a}}{d^{5/2}x}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(7/4)/(c + d*x^2), x]

[Out] $(6*a*b*x)/(5*d*(a+b*x^2)^{(1/4)}) - (2*b*(b*c-a*d)*x)/(d^2*(a+b*x^2)^{(1/4)}) + (2*b*x*(a+b*x^2)^{(3/4)})/(5*d) - (6*a^{(3/2)}*\text{Sqrt}[b]*(1+(b*x^2)/a)^{(1/4)}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/(5*d*(a+b*x^2)^{(1/4)}) + (2*\text{Sqrt}[a]*\text{Sqrt}[b]*(b*c-a*d)*(1+(b*x^2)/a)^{(1/4)}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/(d^2*(a+b*x^2)^{(1/4)}) + (a^{(1/4)}*(-(b*c)+a*d)^{(3/2)}*\text{Sqrt}[-(b*x^2)/a]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c)+a*d])], \text{ArcSin}[(a+b*x^2)^{(1/4)}/a^{(1/4)}], -1))/(d^{(5/2)}*x) - (a^{(1/4)}*(-(b*c)+a*d)^{(3/2)}*\text{Sqrt}[-(b*x^2)/a]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c)+a*d])], \text{ArcSin}[(a+b*x^2)^{(1/4)}/a^{(1/4)}], -1))/(d^{(5/2)}*x)$

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 227

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2*x)/(a + b*x^2)^(1/4), x] - Dist[a, Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a,

, 0] && PosQ[b/a]

Rule 229

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a + b*x^2)^(1/4), Int[1/(1 + (b*x^2)/a)^(1/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 399

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 402

Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Dist[b/d, Int[(a + b*x^2)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4])

Rule 490

Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] := With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2)*Sqrt[c + d*x^4]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^{7/4}}{c + dx^2} dx &= \frac{b \int (a + bx^2)^{3/4} dx}{d} - \frac{(bc - ad) \int \frac{(a + bx^2)^{3/4}}{c + dx^2} dx}{d} \\ &= \frac{2bx(a + bx^2)^{3/4}}{5d} + \frac{(3ab) \int \frac{1}{\sqrt[4]{a + bx^2}} dx}{5d} - \frac{(b(bc - ad)) \int \frac{1}{\sqrt[4]{a + bx^2}} dx}{d^2} + \frac{(bc - ad)^2 \int \frac{1}{\sqrt[4]{a + bx^2}(c + dx^2)} dx}{d^2} \\ &= \frac{2bx(a + bx^2)^{3/4}}{5d} + \frac{\left(2(bc - ad)^2 \sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{x^2}{\sqrt{1 - \frac{x^4}{a}(bc - ad + dx^4)}} dx, x, \sqrt[4]{a + bx^2}\right)}{d^2 x} + \frac{(3ab) \int \frac{1}{\sqrt[4]{a + bx^2}} dx}{d^2} \\ &= \frac{6abx}{5d \sqrt[4]{a + bx^2}} - \frac{2b(bc - ad)x}{d^2 \sqrt[4]{a + bx^2}} + \frac{2bx(a + bx^2)^{3/4}}{5d} - \frac{\left((bc - ad)^2 \sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{(\sqrt{-bc + ad} - \sqrt{a + bx^2})} dx, x, \sqrt[4]{a + bx^2}\right)}{d^{5/2} x} \\ &= \frac{6abx}{5d \sqrt[4]{a + bx^2}} - \frac{2b(bc - ad)x}{d^2 \sqrt[4]{a + bx^2}} + \frac{2bx(a + bx^2)^{3/4}}{5d} - \frac{6a^{3/2} \sqrt{b} \sqrt[4]{1 + \frac{bx^2}{a}} E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\right) \left| 2 \right.}{5d \sqrt[4]{a + bx^2}} + \end{aligned}$$

Mathematica [C] time = 0.46, size = 346, normalized size = 0.96

$$x \frac{\left(6 \left(bx^2(a+bx^2)(c+dx^2) \left(4adF_1\left(\frac{3}{2}, \frac{1}{4}, 2; \frac{5}{2}, -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + bcF_1\left(\frac{3}{2}, \frac{5}{4}, 1; \frac{5}{2}, -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) - 3ac(5a^2d+2abdx^2+2b^2x^2(c+dx^2))F_1\left(\frac{1}{2}, \frac{1}{4}, 1; \frac{3}{2}, -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) \right)}{(c+dx^2) \left(x^2 \left(4adF_1\left(\frac{3}{2}, \frac{1}{4}, 2; \frac{5}{2}, -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + bcF_1\left(\frac{3}{2}, \frac{5}{4}, 1; \frac{5}{2}, -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) - 6acF_1\left(\frac{1}{2}, \frac{1}{4}, 1; \frac{3}{2}, -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right)}$$

$$15d\sqrt[4]{a+bx^2}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(7/4)/(c + d*x^2), x]

[Out] (x*((b*(-5*b*c + 8*a*d))*x^2*(1 + (b*x^2)/a)^(1/4)*AppellF1[3/2, 1/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])/c + (6*(-3*a*c*(5*a^2*d + 2*a*b*d*x^2 + 2*b^2*x^2*(c + d*x^2))*AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -(d*x^2)/c] + b*x^2*(a + b*x^2)*(c + d*x^2)*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -(d*x^2)/c] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])))/(c + d*x^2)*(-6*a*c*AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -(d*x^2)/c] + x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -(d*x^2)/c] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c]))))/(15*d*(a + b*x^2)^(1/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/4)/(d*x^2+c), x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{7}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/4)/(d*x^2+c), x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(7/4)/(d*x^2 + c), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{7}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(7/4)/(d*x^2+c), x)

[Out] int((b*x^2+a)^(7/4)/(d*x^2+c), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{7}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/4)/(d*x^2+c), x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(7/4)/(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{7/4}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(7/4)/(c + d*x^2), x)

[Out] int((a + b*x^2)^(7/4)/(c + d*x^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{7/4}}{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(7/4)/(d*x**2+c), x)

[Out] Integral((a + b*x**2)**(7/4)/(c + d*x**2), x)

$$3.322 \quad \int \frac{(a+bx^2)^{5/4}}{c+dx^2} dx$$

Optimal. Leaf size=302

$$\frac{2a^{3/2}\sqrt{b}\left(\frac{bx^2}{a}+1\right)^{3/4}\operatorname{EllipticF}\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right),2\right)}{3d(a+bx^2)^{3/4}} - \frac{2\sqrt{a}\sqrt{b}\left(\frac{bx^2}{a}+1\right)^{3/4}(bc-ad)\operatorname{EllipticF}\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right),2\right)}{d^2(a+bx^2)^{3/4}}$$

[Out] $2/3*b*x*(b*x^2+a)^{(1/4)}/d+2/3*a^{(3/2)}*(1+b*x^2/a)^{(3/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\operatorname{EllipticF}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*b^{(1/2)}/d/(b*x^2+a)^{(3/4)}-2*(-a*d+b*c)*(1+b*x^2/a)^{(3/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\operatorname{EllipticF}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*a^{(1/2)}*b^{(1/2)}/d^2/(b*x^2+a)^{(3/4)}+a^{(1/4)}*(-a*d+b*c)*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/d^2/x+a^{(1/4)}*(-a*d+b*c)*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/d^2/x$

Rubi [A] time = 0.21, antiderivative size = 302, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {402, 195, 233, 231, 401, 108, 409, 1218}

$$\frac{2a^{3/2}\sqrt{b}\left(\frac{bx^2}{a}+1\right)^{3/4}F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{3d(a+bx^2)^{3/4}} - \frac{2\sqrt{a}\sqrt{b}\left(\frac{bx^2}{a}+1\right)^{3/4}(bc-ad)F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{d^2(a+bx^2)^{3/4}} + \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}(bc-ad)}{d^2(a+bx^2)^{3/4}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(5/4)/(c + d*x^2), x]

[Out] $(2*b*x*(a + b*x^2)^{(1/4)})/(3*d) + (2*a^{(3/2)}*\operatorname{Sqrt}[b]*(1 + (b*x^2)/a)^{(3/4)}*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/2, 2])/(3*d*(a + b*x^2)^{(3/4)}) - (2*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b]*(b*c - a*d)*(1 + (b*x^2)/a)^{(3/4)}*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/2, 2])/(d^2*(a + b*x^2)^{(3/4)}) + (a^{(1/4)}*(b*c - a*d)*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[-((\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b*c) + a*d])], \operatorname{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(d^2*x) + (a^{(1/4)}*(b*c - a*d)*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b*c) + a*d]), \operatorname{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(d^2*x)$

Rule 108

Int[1/(((a_.) + (b_.)*(x_.))*Sqrt[(c_.) + (d_.)*(x_.)]*((e_.) + (f_.)*(x_.))^(3/4)), x_Symbol] := Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 231

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2*EllipticF[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(3/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a

, 0] && PosQ[b/a]

Rule 233

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(3/4)/(a + b*x^2)^(3/4), Int[1/(1 + (b*x^2)/a)^(3/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 401

Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)])*(a + b*x)^(3/4)*(c + d*x)], x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 402

Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Dist[b/d, Int[(a + b*x^2)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4])

Rule 409

Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^{5/4}}{c+dx^2} dx &= \frac{b \int \sqrt[4]{a+bx^2} dx}{d} - \frac{(bc-ad) \int \frac{\sqrt[4]{a+bx^2}}{c+dx^2} dx}{d} \\
&= \frac{2bx\sqrt[4]{a+bx^2}}{3d} + \frac{(ab) \int \frac{1}{(a+bx^2)^{3/4}} dx}{3d} - \frac{(b(bc-ad)) \int \frac{1}{(a+bx^2)^{3/4}} dx}{d^2} + \frac{(bc-ad)^2 \int \frac{1}{(a+bx^2)^{3/4}} dx}{d^2} \\
&= \frac{2bx\sqrt[4]{a+bx^2}}{3d} + \frac{\left((bc-ad)^2 \sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-\frac{bx}{a}}(a+bx)^{3/4}(c+dx)} dx, x, x^2\right)}{2d^2x} + \frac{\left(ab\left(1+\frac{bx^2}{a}\right)\right)}{3d} \\
&= \frac{2bx\sqrt[4]{a+bx^2}}{3d} + \frac{2a^{3/2}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{3d(a+bx^2)^{3/4}} - \frac{2\sqrt{a}\sqrt{b}(bc-ad)\left(1+\frac{bx^2}{a}\right)^3}{d^2(a+bx^2)^{3/4}} \\
&= \frac{2bx\sqrt[4]{a+bx^2}}{3d} + \frac{2a^{3/2}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{3d(a+bx^2)^{3/4}} - \frac{2\sqrt{a}\sqrt{b}(bc-ad)\left(1+\frac{bx^2}{a}\right)^3}{d^2(a+bx^2)^{3/4}} \\
&= \frac{2bx\sqrt[4]{a+bx^2}}{3d} + \frac{2a^{3/2}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{3d(a+bx^2)^{3/4}} - \frac{2\sqrt{a}\sqrt{b}(bc-ad)\left(1+\frac{bx^2}{a}\right)^3}{d^2(a+bx^2)^{3/4}}
\end{aligned}$$

Mathematica [C] time = 0.47, size = 348, normalized size = 1.15

$$\frac{x \left(6 \left(bx^2(a+bx^2)(c+dx^2) \left(4adF_1\left(\frac{3}{2}; \frac{3}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + 3bcF_1\left(\frac{3}{2}; \frac{7}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) - 3ac(3a^2d+2abd^2+2b^2x^2(c+dx^2))F_1\left(\frac{1}{2}; \frac{3}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right)}{(c+dx^2) \left(x^2 \left(4adF_1\left(\frac{3}{2}; \frac{3}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + 3bcF_1\left(\frac{3}{2}; \frac{7}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) - 6acF_1\left(\frac{1}{2}; \frac{3}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right)}{9d(a+bx^2)^{3/4}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(5/4)/(c + d*x^2), x]

[Out] (x*((b*(-3*b*c + 4*a*d))*x^2*(1 + (b*x^2)/a)^(3/4)*AppellF1[3/2, 3/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])/c + (6*(-3*a*c*(3*a^2*d + 2*a*b*d*x^2 + 2*b^2*x^2*(c + d*x^2))*AppellF1[1/2, 3/4, 1, 3/2, -(b*x^2)/a, -(d*x^2)/c] + b*x^2*(a + b*x^2)*(c + d*x^2)*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -(b*x^2)/a, -(d*x^2)/c] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])))/(c + d*x^2)*(-6*a*c*AppellF1[1/2, 3/4, 1, 3/2, -(b*x^2)/a, -(d*x^2)/c] + x^2*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -(b*x^2)/a, -(d*x^2)/c] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])))/(9*d*(a + b*x^2)^(3/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/4)/(d*x^2+c), x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/4)/(d*x^2+c),x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(5/4)/(d*x^2 + c), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/4)/(d*x^2+c),x)

[Out] int((b*x^2+a)^(5/4)/(d*x^2+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/4)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/4)/(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/4)/(c + d*x^2),x)

[Out] int((a + b*x^2)^(5/4)/(c + d*x^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{5}{4}}}{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/4)/(d*x**2+c),x)

[Out] Integral((a + b*x**2)**(5/4)/(c + d*x**2), x)

$$3.323 \quad \int \frac{(a+bx^2)^{3/4}}{c+dx^2} dx$$

Optimal. Leaf size=244

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \sqrt{ad-bc} \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{d^{3/2}x} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \sqrt{ad-bc} \Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{d^{3/2}x}$$

[Out] $2*b*x/d/(b*x^2+a)^{(1/4)}-2*(1+b*x^2/a)^{(1/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*a^{(1/2)}*b^{(1/2)}/d/(b*x^2+a)^{(1/4)}+a^{(1/4)}*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(a*d-b*c)^{(1/2)}*(-b*x^2/a)^{(1/2)}/d^{(3/2)}/x-a^{(1/4)}*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(a*d-b*c)^{(1/2)}*(-b*x^2/a)^{(1/2)}/d^{(3/2)}/x$

Rubi [A] time = 0.15, antiderivative size = 244, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {402, 229, 227, 196, 399, 490, 1218}

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \sqrt{ad-bc} \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{d^{3/2}x} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \sqrt{ad-bc} \Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{d^{3/2}x}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/4)/(c + d*x^2), x]

[Out] $(2*b*x)/(d*(a + b*x^2)^{(1/4)}) - (2*\text{Sqrt}[a]*\text{Sqrt}[b]*(1 + (b*x^2)/a)^{(1/4)}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/(d*(a + b*x^2)^{(1/4)}) + (a^{(1/4)}*\text{Sqrt}[-(b*c) + a*d]*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d])], \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(d^{(3/2)}*x) - (a^{(1/4)}*\text{Sqrt}[-(b*c) + a*d]*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(d^{(3/2)}*x)$

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 227

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2*x)/(a + b*x^2)^(1/4), x] - Dist[a, Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 229

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a + b*x^2)^(1/4), Int[1/(1 + (b*x^2)/a)^(1/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 399

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -

a*d, 0]

Rule 402

```
Int[((a_) + (b_)*(x_)^2)^(p_)/((c_) + (d_)*(x_)^2), x_Symbol] := Dist[b/d, Int[(a + b*x^2)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4])
```

Rule 490

```
Int[(x_)^2/(((a_) + (b_)*(x_)^4)*Sqrt[(c_) + (d_)*(x_)^4]), x_Symbol] := With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/((r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]
```

Rule 1218

```
Int[1/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\int \frac{(a + bx^2)^{3/4}}{c + dx^2} dx = \frac{b \int \frac{1}{\sqrt[4]{a+bx^2}} dx}{d} - \frac{(bc - ad) \int \frac{1}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{d}$$

$$= -\frac{\left(2(bc - ad)\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{x^2}{\sqrt{1-\frac{x^4}{a}}(bc-ad+dx^4)} dx, x, \sqrt[4]{a+bx^2}\right)}{dx} + \frac{\left(b\sqrt[4]{1+\frac{bx^2}{a}}\right) \int \frac{1}{\sqrt[4]{1+\frac{bx^2}{a}}} dx}{d\sqrt[4]{a+bx^2}}$$

$$= \frac{2bx}{d\sqrt[4]{a+bx^2}} + \frac{\left((bc - ad)\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{(\sqrt{-bc+ad}-\sqrt{d}x^2)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2}\right)}{d^{3/2}x} - \frac{\left((bc - ad)\sqrt{-\frac{bx^2}{a}}\right) \int \frac{1}{(\sqrt{-bc+ad}+\sqrt{d}x^2)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2}}{d^{3/2}x}$$

$$= \frac{2bx}{d\sqrt[4]{a+bx^2}} - \frac{2\sqrt{a}\sqrt{b}\sqrt[4]{1+\frac{bx^2}{a}} E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{d\sqrt[4]{a+bx^2}} + \frac{\sqrt[4]{a}\sqrt{-bc+ad}\sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{-bc+ad}}\right)}{d^{3/2}x}$$

Mathematica [C] time = 0.16, size = 161, normalized size = 0.66

$$\frac{6acx(a + bx^2)^{3/4} F_1\left(\frac{1}{2}; -\frac{3}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{(c + dx^2) \left(x^2 \left(3bcF_1\left(\frac{3}{2}; \frac{1}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - 4adF_1\left(\frac{3}{2}; -\frac{3}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) + 6acF_1\left(\frac{1}{2}; -\frac{3}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(3/4)/(c + d*x^2), x]

[Out] (6*a*c*x*(a + b*x^2)^(3/4)*AppellF1[1/2, -3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)]/((c + d*x^2)*(6*a*c*AppellF1[1/2, -3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^2*(-4*a*d*AppellF1[3/2, -3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 1/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/4)/(d*x^2+c),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/4)/(d*x^2+c),x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(3/4)/(d*x^2 + c), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/4)/(d*x^2+c),x)

[Out] int((b*x^2+a)^(3/4)/(d*x^2+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/4)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/4)/(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/4)/(c + d*x^2),x)

[Out] int((a + b*x^2)^(3/4)/(c + d*x^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{3}{4}}}{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/4)/(d*x**2+c),x)

[Out] Integral((a + b*x**2)**(3/4)/(c + d*x**2), x)

$$3.324 \quad \int \frac{\sqrt[4]{a+bx^2}}{c+dx^2} dx$$

Optimal. Leaf size=199

$$\frac{2\sqrt{a}\sqrt{b}\left(\frac{bx^2}{a}+1\right)^{3/4}\operatorname{EllipticF}\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right),2\right)}{d(a+bx^2)^{3/4}} - \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle| -1\right)}{dx} + \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle| -1\right)}{dx}$$

[Out] $2*(1+b*x^2/a)^{(3/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^{(1/2)})^2/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\operatorname{EllipticF}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*a^{(1/2)}*b^{(1/2)}/d/(b*x^2+a)^{(3/4)}-a^{(1/4)}*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/d/x-a^{(1/4)}*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/d/x$

Rubi [A] time = 0.15, antiderivative size = 199, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 21, number of rules / integrand size = 0.333, Rules used = {402, 233, 231, 401, 108, 409, 1218}

$$\frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle| -1\right)}{dx} - \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}\Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle| -1\right)}{dx} + \frac{2\sqrt{a}\sqrt{b}\left(\frac{bx^2}{a}+1\right)^{3/4}}{d(a+bx^2)^{3/4}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(1/4)/(c + d*x^2), x]

[Out] $(2*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b]*(1+(b*x^2)/a)^{(3/4)}*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/2, 2])/(d*(a+b*x^2)^{(3/4)}) - (a^{(1/4)}*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[-((\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b*c)+a*d]), \operatorname{ArcSin}[(a+b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(d*x) - (a^{(1/4)}*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b*c)+a*d]), \operatorname{ArcSin}[(a+b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(d*x)$

Rule 108

Int[1/(((a_.) + (b_.)*(x_.))*Sqrt[(c_.) + (d_.)*(x_.)]*((e_.) + (f_.)*(x_.))^(3/4)), x_Symbol] := Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 231

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2*EllipticF[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(3/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 233

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(3/4)/(a + b*x^2)^(3/4), Int[1/(1 + (b*x^2)/a)^(3/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 401

Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 402

```
Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Dist[b/d, Int[(a + b*x^2)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4])
```

Rule 409

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]
```

Rule 1218

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\int \frac{\sqrt[4]{a+bx^2}}{c+dx^2} dx = \frac{b \int \frac{1}{(a+bx^2)^{3/4}} dx}{d} - \frac{(bc-ad) \int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx}{d}$$

$$= -\frac{\left((bc-ad)\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{\frac{-bx}{a}}(a+bx)^{3/4}(c+dx)} dx, x, x^2\right)}{2dx} + \frac{\left(b\left(1+\frac{bx^2}{a}\right)^{3/4}\right) \int \frac{1}{\left(1+\frac{bx^2}{a}\right)^{3/4}} dx}{d(a+bx^2)^{3/4}}$$

$$= \frac{2\sqrt{a}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{d(a+bx^2)^{3/4}} + \frac{\left(2(bc-ad)\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^4}{a}}(-bc+ad-dx^4)} dx\right)}{dx}$$

$$= \frac{2\sqrt{a}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{d(a+bx^2)^{3/4}} - \frac{\sqrt{-\frac{bx^2}{a}} \text{Subst}\left(\int \frac{1}{\left(1-\frac{\sqrt{d}x^2}{\sqrt{-bc+ad}}\right)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2}\right)}{dx}$$

$$= \frac{2\sqrt{a}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{d(a+bx^2)^{3/4}} - \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{-bc+ad}}; \sin^{-1}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right)\middle|-1\right)}{dx}$$

Mathematica [C] time = 0.16, size = 160, normalized size = 0.80

$$\frac{6acx\sqrt[4]{a+bx^2} F_1\left(\frac{1}{2}; -\frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{(c+dx^2)\left(x^2\left(bcF_1\left(\frac{3}{2}; \frac{3}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - 4adF_1\left(\frac{3}{2}; -\frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)\right) + 6acF_1\left(\frac{1}{2}; -\frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[(a + b*x^2)^(1/4)/(c + d*x^2), x]
```

```
[Out] (6*a*c*x*(a + b*x^2)^(1/4)*AppellF1[1/2, -1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)]/((c + d*x^2)*(6*a*c*AppellF1[1/2, -1/4, 1, 3/2, -((b*x^2)/a), -((d
```

$*x^2)/c]] + x^2*(-4*a*d*AppellF1[3/2, -1/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 3/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/4)/(d*x^2+c),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{1}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/4)/(d*x^2+c),x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(1/4)/(d*x^2 + c), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{1}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/4)/(d*x^2+c),x)

[Out] int((b*x^2+a)^(1/4)/(d*x^2+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{1}{4}}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/4)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(1/4)/(d*x^2 + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^{1/4}}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/4)/(c + d*x^2),x)

[Out] int((a + b*x^2)^(1/4)/(c + d*x^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[4]{a + bx^2}}{c + dx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/4)/(d*x**2+c),x)

[Out] Integral((a + b*x**2)**(1/4)/(c + d*x**2), x)

$$3.325 \quad \int \frac{1}{\sqrt[4]{a+bx^2} (c+dx^2)} dx$$

Optimal. Leaf size=167

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{\sqrt{d} x \sqrt{ad-bc}} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{\sqrt{d} x \sqrt{ad-bc}}$$

[Out] $a^{1/4} \text{EllipticPi}((b*x^2+a)^{1/4}/a^{1/4}, -a^{1/2}*d^{1/2}/(a*d-b*c)^{1/2}, I) * (-b*x^2/a)^{1/2}/x/d^{1/2}/(a*d-b*c)^{1/2} - a^{1/4} \text{EllipticPi}((b*x^2+a)^{1/4}/a^{1/4}, a^{1/2}*d^{1/2}/(a*d-b*c)^{1/2}, I) * (-b*x^2/a)^{1/2}/x/d^{1/2}/(a*d-b*c)^{1/2}$

Rubi [A] time = 0.11, antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {399, 490, 1218}

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{\sqrt{d} x \sqrt{ad-bc}} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{\sqrt{d} x \sqrt{ad-bc}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(1/4)*(c + d*x^2)), x]

[Out] $(a^{1/4} \text{Sqrt}[-((b*x^2)/a)] \text{EllipticPi}[-((\text{Sqrt}[a] \text{Sqrt}[d])/ \text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{1/4}/a^{1/4}], -1]) / (\text{Sqrt}[d] \text{Sqrt}[-(b*c) + a*d]*x) - (a^{1/4} \text{Sqrt}[-((b*x^2)/a)] \text{EllipticPi}[(\text{Sqrt}[a] \text{Sqrt}[d])/ \text{Sqrt}[-(b*c) + a*d], \text{ArcSin}[(a + b*x^2)^{1/4}/a^{1/4}], -1]) / (\text{Sqrt}[d] \text{Sqrt}[-(b*c) + a*d]*x)$

Rule 399

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 490

Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] := With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/((r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\int \frac{1}{\sqrt[4]{a+bx^2}(c+dx^2)} dx = \frac{\left(2\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{x^2}{\sqrt{1-\frac{x^4}{a}(bc-ad+dx^4)}} dx, x, \sqrt[4]{a+bx^2}\right)}{x}$$

$$= \frac{\sqrt{-\frac{bx^2}{a}} \text{Subst}\left(\int \frac{1}{(\sqrt{-bc+ad}-\sqrt{d}x^2)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2}\right)}{\sqrt{d}x} + \frac{\sqrt{-\frac{bx^2}{a}} \text{Subst}\left(\int \frac{1}{(\sqrt{-bc+ad}+\sqrt{d}x^2)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2}\right)}{\sqrt{d}x}$$

$$= \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{-bc+ad}}; \sin^{-1}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) \middle| -1\right)}{\sqrt{d}\sqrt{-bc+ad}x} - \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}} \Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{-bc+ad}}; \sin^{-1}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) \middle| -1\right)}{\sqrt{d}\sqrt{-bc+ad}x}$$

Mathematica [C] time = 0.06, size = 160, normalized size = 0.96

$$\frac{6acx F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{\sqrt[4]{a+bx^2}(c+dx^2) \left(x^2 \left(4ad F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + bc F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)\right) - 6ac F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(1/4)*(c + d*x^2)),x]

[Out] (-6*a*c*x*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)]/((a + b*x^2)^(1/4)*(c + d*x^2)*(-6*a*c*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/4)/(d*x^2+c),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+a)^{\frac{1}{4}}(dx^2+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((b*x^2 + a)^(1/4)*(d*x^2 + c)), x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(1/4)*(d*x^2 + c)), x)

maple [F] time = 0.33, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+a)^{\frac{1}{4}}(dx^2+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b*x^2+a)^(1/4)/(d*x^2+c),x)`

[Out] `int(1/(b*x^2+a)^(1/4)/(d*x^2+c),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{1}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x^2+a)^(1/4)/(d*x^2+c),x, algorithm="maxima")`

[Out] `integrate(1/((b*x^2 + a)^(1/4)*(d*x^2 + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + a)^{1/4} (dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((a + b*x^2)^(1/4)*(c + d*x^2)),x)`

[Out] `int(1/((a + b*x^2)^(1/4)*(c + d*x^2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[4]{a + bx^2} (c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x**2+a)**(1/4)/(d*x**2+c),x)`

[Out] `Integral(1/((a + b*x**2)**(1/4)*(c + d*x**2)), x)`

$$3.326 \quad \int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx$$

Optimal. Leaf size=152

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{x(bc-ad)} + \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{x(bc-ad)}$$

[Out] $a^{1/4} \text{EllipticPi}((b*x^2+a)^{1/4}/a^{1/4}, -a^{1/2}*d^{1/2}/(a*d-b*c)^{1/2}, I)*(-b*x^2/a)^{1/2}/(-a*d+b*c)/x + a^{1/4} \text{EllipticPi}((b*x^2+a)^{1/4}/a^{1/4}, a^{1/2}*d^{1/2}/(a*d-b*c)^{1/2}, I)*(-b*x^2/a)^{1/2}/(-a*d+b*c)/x$

Rubi [A] time = 0.12, antiderivative size = 152, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {401, 108, 409, 1218}

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{x(bc-ad)} + \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{x(bc-ad)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(3/4)*(c + d*x^2)), x]

[Out] $(a^{1/4} \text{Sqrt}[-((b*x^2)/a)] \text{EllipticPi}[-((\text{Sqrt}[a] \text{Sqrt}[d])/ \text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{1/4}/a^{1/4}], -1])/((b*c - a*d)*x) + (a^{1/4} \text{Sqrt}[-((b*x^2)/a)] \text{EllipticPi}[(\text{Sqrt}[a] \text{Sqrt}[d])/ \text{Sqrt}[-(b*c) + a*d], \text{ArcSin}[(a + b*x^2)^{1/4}/a^{1/4}], -1])/((b*c - a*d)*x)$

Rule 108

Int[1/(((a_.) + (b_.)*(x_.))*Sqrt[(c_.) + (d_.)*(x_.)]*((e_.) + (f_.)*(x_.))^(3/4)), x_Symbol] := Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 401

Int[1/(((a_.) + (b_.)*(x_.)^2)^(3/4)*((c_.) + (d_.)*(x_.)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 409

Int[1/(Sqrt[(a_.) + (b_.)*(x_.)^4]*((c_.) + (d_.)*(x_.)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 1218

Int[1/(((d_.) + (e_.)*(x_.)^2)*Sqrt[(a_.) + (c_.)*(x_.)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx &= \frac{\sqrt{-\frac{bx^2}{a}} \operatorname{Subst}\left(\int \frac{1}{\sqrt{-\frac{bx}{a}}(a+bx)^{3/4}(c+dx)} dx, x, x^2\right)}{2x} \\
&= \frac{\left(2\sqrt{-\frac{bx^2}{a}}\right) \operatorname{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^4}{a}}(-bc+ad-dx^4)} dx, x, \sqrt[4]{a+bx^2}\right)}{x} \\
&= \frac{\sqrt{-\frac{bx^2}{a}} \operatorname{Subst}\left(\int \frac{1}{\left(1-\frac{\sqrt{d}x^2}{\sqrt{-bc+ad}}\right)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2}\right)}{(bc-ad)x} + \frac{\sqrt{-\frac{bx^2}{a}} \operatorname{Subst}\left(\int \frac{1}{\left(1+\frac{\sqrt{d}}{\sqrt{-bc+ad}}\right)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2}\right)}{(bc-ad)x} \\
&= \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{-bc+ad}}; \sin^{-1}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) \middle| -1\right)}{(bc-ad)x} + \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{-bc+ad}}; \sin^{-1}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right) \middle| -1\right)}{(bc-ad)x}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 120, normalized size = 0.79

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \left(\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right) + \Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right) \right)}{x(bc-ad)}$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)^(3/4)*(c + d*x^2)), x]

[Out] (a^(1/4)*Sqrt[-((b*x^2)/a)]*(EllipticPi[-((Sqrt[a]*Sqrt[d])/Sqrt[-(b*c) + a*d]), ArcSin[(a + b*x^2)^(1/4)/a^(1/4)], -1] + EllipticPi[(Sqrt[a]*Sqrt[d])/Sqrt[-(b*c) + a*d], ArcSin[(a + b*x^2)^(1/4)/a^(1/4)], -1]))/((b*c - a*d)*x)

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/4)/(d*x^2+c), x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+a)^{\frac{3}{4}}(dx^2+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/4)/(d*x^2+c), x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(3/4)*(d*x^2 + c)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+a)^{\frac{3}{4}}(dx^2+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b*x^2+a)^(3/4)/(d*x^2+c),x)`

[Out] `int(1/(b*x^2+a)^(3/4)/(d*x^2+c),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x^2+a)^(3/4)/(d*x^2+c),x, algorithm="maxima")`

[Out] `integrate(1/((b*x^2 + a)^(3/4)*(d*x^2 + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((a + b*x^2)^(3/4)*(c + d*x^2)),x)`

[Out] `int(1/((a + b*x^2)^(3/4)*(c + d*x^2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{3}{4}}(c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x**2+a)**(3/4)/(d*x**2+c),x)`

[Out] `Integral(1/((a + b*x**2)**(3/4)*(c + d*x**2)), x)`

$$3.327 \quad \int \frac{1}{(a+bx^2)^{5/4}(c+dx^2)} dx$$

Optimal. Leaf size=233

$$\frac{2\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{\sqrt{a}\sqrt[4]{a+bx^2}(bc-ad)} + \frac{\sqrt[4]{a}\sqrt{d}\sqrt{-\frac{bx^2}{a}}\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle|-1\right)}{x(ad-bc)^{3/2}} - \frac{\sqrt[4]{a}\sqrt{d}\sqrt{-\frac{bx^2}{a}}\Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle|-1\right)}{x(ad-bc)^{3/2}}$$

[Out] $2*(1+b*x^2/a)^{(1/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^{(1/2)})^2/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*b^{(1/2)}/(-a*d+b*c)/(b*x^2+a)^{(1/4)}/a^{(1/2)}+a^{(1/4)}*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*d^{(1/2)}*(-b*x^2/a)^{(1/2)}/(a*d-b*c)^{(3/2)}/x-a^{(1/4)}*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*d^{(1/2)}*(-b*x^2/a)^{(1/2)}/(a*d-b*c)^{(3/2)}/x$

Rubi [A] time = 0.16, antiderivative size = 233, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {403, 197, 196, 399, 490, 1218}

$$\frac{2\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{\sqrt{a}\sqrt[4]{a+bx^2}(bc-ad)} + \frac{\sqrt[4]{a}\sqrt{d}\sqrt{-\frac{bx^2}{a}}\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle|-1\right)}{x(ad-bc)^{3/2}} - \frac{\sqrt[4]{a}\sqrt{d}\sqrt{-\frac{bx^2}{a}}\Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle|-1\right)}{x(ad-bc)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(5/4)*(c + d*x^2)),x]

[Out] $(2*\text{Sqrt}[b]*(1 + (b*x^2)/a)^{(1/4)}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/(\text{Sqrt}[a]*(b*c - a*d)*(a + b*x^2)^{(1/4)}) + (a^{(1/4)}*\text{Sqrt}[d]*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d])], \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1))/((-b*c) + a*d)^{(3/2)*x} - (a^{(1/4)}*\text{Sqrt}[d]*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1))/((-b*c) + a*d)^{(3/2)*x}$

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 197

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a*(a + b*x^2)^(1/4)), Int[1/(1 + (b*x^2)/a)^(5/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a] && PosQ[b/a]

Rule 399

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 403

Int[((a_) + (b_.)*(x_)^2)^(p_)/((c_) + (d_.)*(x_)^2), x_Symbol] := Dist[b/(b*c - a*d), Int[(a + b*x^2)^p, x], x] - Dist[d/(b*c - a*d), Int[(a + b*x^2)^(p + 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

&& LtQ[p, -1] && EqQ[Denominator[p], 4] && (EqQ[p, -5/4] || EqQ[p, -7/4])

Rule 490

```
Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] :=
With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s
/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/
(r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c
- a*d, 0]
```

Rule 1218

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*
Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\int \frac{1}{(a + bx^2)^{5/4} (c + dx^2)} dx = \frac{b \int \frac{1}{(a+bx^2)^{5/4}} dx}{bc - ad} - \frac{d \int \frac{1}{\sqrt[4]{a+bx^2} (c+dx^2)} dx}{bc - ad}$$

$$= -\frac{\left(2d\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{x^2}{\sqrt{1-\frac{x^4}{a}(bc-ad+dx^4)}} dx, x, \sqrt[4]{a+bx^2}\right)}{(bc - ad)x} + \frac{\left(b\sqrt[4]{1+\frac{bx^2}{a}}\right) \int \frac{1}{\left(1+\frac{bx^2}{a}\right)}}{a(bc - ad)\sqrt[4]{a+bx^2}}$$

$$= \frac{2\sqrt{b} \sqrt[4]{1+\frac{bx^2}{a}} E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| 2\right)}{\sqrt{a}(bc - ad)\sqrt[4]{a+bx^2}} + \frac{\left(\sqrt{d} \sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{(\sqrt{-bc+ad}-\sqrt{d}x^2)\sqrt{1-\frac{x^4}{a}}}\right)}{(bc - ad)x}$$

$$= \frac{2\sqrt{b} \sqrt[4]{1+\frac{bx^2}{a}} E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| 2\right)}{\sqrt{a}(bc - ad)\sqrt[4]{a+bx^2}} + \frac{\sqrt[4]{a} \sqrt{d} \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{-bc+ad}}; \sin^{-1}\left(\frac{\sqrt[4]{a+bx^2}}{\sqrt[4]{a}}\right)\right)}{(-bc + ad)^{3/2}x}$$

Mathematica [C] time = 0.23, size = 327, normalized size = 1.40

$$x \frac{\left(\frac{bdx^2 \sqrt[4]{\frac{bx^2}{a}} + {}_1F_1\left(\frac{3}{2}; \frac{1}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c} + \frac{6\left(bx^2(c+dx^2)\left(4ad {}_1F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + bc {}_1F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)\right) + 3ac(ad-b(c+2dx^2)) {}_1F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{(c+dx^2)\left(6ac {}_1F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - x^2\left(4ad {}_1F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + bc {}_1F_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)\right)}\right)}{3a\sqrt[4]{a+bx^2}(ad - bc)}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[1/((a + b*x^2)^(5/4)*(c + d*x^2)), x]
[Out] (x*((b*d*x^2*(1 + (b*x^2)/a)^(1/4)*AppellF1[3/2, 1/4, 1, 5/2, -((b*x^2)/a),
-((d*x^2)/c)])/c + (6*(3*a*c*(a*d - b*(c + 2*d*x^2))*AppellF1[1/2, 1/4, 1,
3/2, -((b*x^2)/a), -((d*x^2)/c)] + b*x^2*(c + d*x^2)*(4*a*d*AppellF1[3/2,
1/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -
((b*x^2)/a), -((d*x^2)/c)])))/((c + d*x^2)*(6*a*c*AppellF1[1/2, 1/4, 1, 3/2
, -((b*x^2)/a), -((d*x^2)/c)] - x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -((b*
x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -((b*x^2)/a), -((d*
x^2)/c)])))/((3*a*(-(b*c) + a*d)*(a + b*x^2)^(1/4))
```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/4)/(d*x^2+c),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/4)/(d*x^2+c),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(5/4)*(d*x^2 + c)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(5/4)/(d*x^2+c),x)

[Out] int(1/(b*x^2+a)^(5/4)/(d*x^2+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/4)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(5/4)*(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(5/4)*(c + d*x^2)),x)

[Out] int(1/((a + b*x^2)^(5/4)*(c + d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{5}{4}}(c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(5/4)/(d*x**2+c),x)

[Out] Integral(1/((a + b*x**2)**(5/4)*(c + d*x**2)), x)

$$3.328 \quad \int \frac{1}{(a+bx^2)^{7/4}(c+dx^2)} dx$$

Optimal. Leaf size=254

$$\frac{2\sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), 2\right)}{3\sqrt{a} (a+bx^2)^{3/4} (bc-ad)} + \frac{2bx}{3a (a+bx^2)^{3/4} (bc-ad)} - \frac{\sqrt[4]{a} d \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\right)}{x(bc-ad)^2}$$

[Out] $2/3*b*x/a/(-a*d+b*c)/(b*x^2+a)^{(3/4)}+2/3*(1+b*x^2/a)^{(3/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\text{EllipticF}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})), 2^{(1/2)})*b^{(1/2)}/(-a*d+b*c)/(b*x^2+a)^{(3/4)}/a^{(1/2)}-a^{(1/4)}*d*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, -a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/(-a*d+b*c)^2/x-a^{(1/4)}*d*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/(-a*d+b*c)^2/x$

Rubi [A] time = 0.16, antiderivative size = 254, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {403, 199, 233, 231, 401, 108, 409, 1218}

$$\frac{2bx}{3a (a+bx^2)^{3/4} (bc-ad)} + \frac{2\sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{3\sqrt{a} (a+bx^2)^{3/4} (bc-ad)} - \frac{\sqrt[4]{a} d \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{x(bc-ad)^2}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(7/4)*(c + d*x^2)), x]

[Out] $(2*b*x)/(3*a*(b*c - a*d)*(a + b*x^2)^{(3/4)}) + (2*\text{Sqrt}[b]*(1 + (b*x^2)/a)^{(3/4)}*\text{EllipticF}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/(3*\text{Sqrt}[a]*(b*c - a*d)*(a + b*x^2)^{(3/4)}) - (a^{(1/4)}*d*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/((b*c - a*d)^2*x) - (a^{(1/4)}*d*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/((b*c - a*d)^2*x)$

Rule 108

Int[1/(((a_.) + (b_.)*(x_))*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(3/4)), x_Symbol] :> Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 199

Int[((a_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])

Rule 231

Int[((a_.) + (b_.)*(x_)^2)^(-3/4), x_Symbol] :> Simp[(2*EllipticF[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(3/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 233

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(3/4)/(a + b*x^2)^(3/4), Int[1/(1 + (b*x^2)/a)^(3/4), x], x] /; FreeQ[{a, b}, x] & PosQ[a]

Rule 401

Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 403

Int[((a_) + (b_.)*(x_)^2)^(p_)/((c_) + (d_.)*(x_)^2), x_Symbol] := Dist[b/(b*c - a*d), Int[(a + b*x^2)^p, x], x] - Dist[d/(b*c - a*d), Int[(a + b*x^2)^(p + 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && EqQ[Denominator[p], 4] && (EqQ[p, -5/4] || EqQ[p, -7/4])

Rule 409

Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + bx^2)^{7/4} (c + dx^2)} dx &= \frac{b \int \frac{1}{(a+bx^2)^{7/4}} dx}{bc - ad} - \frac{d \int \frac{1}{(a+bx^2)^{3/4} (c+dx^2)} dx}{bc - ad} \\ &= \frac{2bx}{3a(bc - ad)(a + bx^2)^{3/4}} + \frac{b \int \frac{1}{(a+bx^2)^{3/4}} dx}{3a(bc - ad)} - \frac{\left(d\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-\frac{bx}{a}}(a+bx)^{3/4}} dx, x, \sqrt[4]{a + bx^2}\right)}{2(bc - ad)x} \\ &= \frac{2bx}{3a(bc - ad)(a + bx^2)^{3/4}} + \frac{\left(2d\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^4}{a}}(-bc+ad-dx^4)} dx, x, \sqrt[4]{a + bx^2}\right)}{(bc - ad)x} \\ &= \frac{2bx}{3a(bc - ad)(a + bx^2)^{3/4}} + \frac{2\sqrt{b} \left(1 + \frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{3\sqrt{a}(bc - ad)(a + bx^2)^{3/4}} - \frac{\left(d\sqrt{-\frac{bx^2}{a}}\right)}{2(bc - ad)x} \\ &= \frac{2bx}{3a(bc - ad)(a + bx^2)^{3/4}} + \frac{2\sqrt{b} \left(1 + \frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{3\sqrt{a}(bc - ad)(a + bx^2)^{3/4}} - \frac{\sqrt[4]{a} d \sqrt{-\frac{bx^2}{a}}}{2(bc - ad)x} \end{aligned}$$

Mathematica [C] time = 0.26, size = 331, normalized size = 1.30

$$x \frac{\left(6 \left(b x^2 (c + d x^2) \left(4 a d F_1 \left(\frac{3}{2}, \frac{3}{4}, 2, \frac{5}{2}; -\frac{b x^2}{a}, -\frac{d x^2}{c} \right) + 3 b c F_1 \left(\frac{3}{2}, \frac{7}{4}, 1, \frac{5}{2}; -\frac{b x^2}{a}, -\frac{d x^2}{c} \right) \right) + 3 a c (3 a d - 3 b c - 2 b d x^2) F_1 \left(\frac{1}{2}, \frac{3}{4}, 1, \frac{3}{2}; -\frac{b x^2}{a}, -\frac{d x^2}{c} \right) \right) b d x^2 \left(\frac{b x^2}{a} + 1 \right)^{3/4} F_1 \left(\frac{3}{2}, \frac{3}{4}, 1, \frac{5}{2}; -\frac{b x^2}{a}, -\frac{d x^2}{c} \right)}{\left(c + d x^2 \right) \left(6 a c F_1 \left(\frac{1}{2}, \frac{3}{4}, 1, \frac{3}{2}; -\frac{b x^2}{a}, -\frac{d x^2}{c} \right) - x^2 \left(4 a d F_1 \left(\frac{3}{2}, \frac{3}{4}, 2, \frac{5}{2}; -\frac{b x^2}{a}, -\frac{d x^2}{c} \right) + 3 b c F_1 \left(\frac{3}{2}, \frac{7}{4}, 1, \frac{5}{2}; -\frac{b x^2}{a}, -\frac{d x^2}{c} \right) \right) \right)} 9 a \left(a + b x^2 \right)^{3/4} (a d - b c)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(7/4)*(c + d*x^2)),x]

[Out] (x*(-((b*d*x^2*(1 + (b*x^2)/a)^(3/4)*AppellF1[3/2, 3/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])/c) + (6*(3*a*c*(-3*b*c + 3*a*d - 2*b*d*x^2)*AppellF1[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + b*x^2*(c + d*x^2)*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/((c + d*x^2)*(6*a*c*AppellF1[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] - x^2*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/((9*a*(-(b*c) + a*d)*(a + b*x^2)^(3/4)))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/4)/(d*x^2+c),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b x^2 + a)^{\frac{7}{4}} (d x^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/4)/(d*x^2+c),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(7/4)*(d*x^2 + c)), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(b x^2 + a)^{\frac{7}{4}} (d x^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(7/4)/(d*x^2+c),x)

[Out] int(1/(b*x^2+a)^(7/4)/(d*x^2+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b x^2 + a)^{\frac{7}{4}} (d x^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/4)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(7/4)*(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{7/4} (dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(7/4)*(c + d*x^2)), x)

[Out] int(1/((a + b*x^2)^(7/4)*(c + d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{7/4} (c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(7/4)/(d*x**2+c), x)

[Out] Integral(1/((a + b*x**2)**(7/4)*(c + d*x**2)), x)

$$3.329 \quad \int \frac{1}{(a+bx^2)^{9/4}(c+dx^2)} dx$$

Optimal. Leaf size=274

$$\frac{2\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}(3bc-8ad)E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{5a^{3/2}\sqrt[4]{a+bx^2}(bc-ad)^2} + \frac{\sqrt[4]{a}d^{3/2}\sqrt{-\frac{bx^2}{a}}\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle|-1\right)}{x(ad-bc)^{5/2}} - \frac{\sqrt[4]{a}d^{3/2}\sqrt{-\frac{bx^2}{a}}}{x(ad-bc)^{5/2}}$$

[Out] $2/5*b*x/a/(-a*d+b*c)/(b*x^2+a)^{(5/4)}+2/5*(-8*a*d+3*b*c)*(1+b*x^2/a)^{(1/4)*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*b^{(1/2)}/a^{(3/2)}/(-a*d+b*c)^2/(b*x^2+a)^{(1/4)}+a^{(1/4)}*d^{(3/2)}*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/(a*d-b*c)^{(5/2)}/x-a^{(1/4)}*d^{(3/2)}*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/(a*d-b*c)^{(5/2)}/x$

Rubi [A] time = 0.38, antiderivative size = 274, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {414, 527, 530, 229, 227, 196, 399, 490, 1218}

$$\frac{2\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}(3bc-8ad)E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{5a^{3/2}\sqrt[4]{a+bx^2}(bc-ad)^2} + \frac{\sqrt[4]{a}d^{3/2}\sqrt{-\frac{bx^2}{a}}\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\middle|-1\right)}{x(ad-bc)^{5/2}} - \frac{\sqrt[4]{a}d^{3/2}\sqrt{-\frac{bx^2}{a}}}{x(ad-bc)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(9/4)*(c + d*x^2)),x]

[Out] $(2*b*x)/(5*a*(b*c - a*d)*(a + b*x^2)^{(5/4)}) + (2*\text{Sqrt}[b]*(3*b*c - 8*a*d)*(1 + (b*x^2)/a)^{(1/4)}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/ (5*a^{(3/2)}*(b*c - a*d)^2*(a + b*x^2)^{(1/4)}) + (a^{(1/4)}*d^{(3/2)}*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/ \text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1))/((-b*c) + a*d)^{(5/2)*x} - (a^{(1/4)}*d^{(3/2)}*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/ \text{Sqrt}[-(b*c) + a*d], \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1))/((-b*c) + a*d)^{(5/2)*x}$

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 227

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2*x)/(a + b*x^2)^(1/4), x] - Dist[a, Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 229

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a + b*x^2)^(1/4), Int[1/(1 + (b*x^2)/a)^(1/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 399

Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x

$\wedge 4)), x], x, (a + b*x^2)^{(1/4)}, x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 490

Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] :> With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 530

Int((((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] :> With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a+bx^2)^{9/4}(c+dx^2)} dx &= \frac{2bx}{5a(bc-ad)(a+bx^2)^{5/4}} - \frac{2 \int \frac{\frac{1}{2}(-3bc+5ad) - \frac{3}{2}bdx^2}{(a+bx^2)^{5/4}(c+dx^2)} dx}{5a(bc-ad)} \\
 &= \frac{2bx}{5a(bc-ad)(a+bx^2)^{5/4}} + \frac{2b(3bc-8ad)x}{5a^2(bc-ad)^2 \sqrt[4]{a+bx^2}} + \frac{4 \int \frac{\frac{1}{4}(-3b^2c^2+8abcd+5a^2d^2) - \frac{1}{4}bd(3bc+5ad) - \frac{3}{4}bd^2x^2}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{5a^2(bc-ad)^2} \\
 &= \frac{2bx}{5a(bc-ad)(a+bx^2)^{5/4}} + \frac{2b(3bc-8ad)x}{5a^2(bc-ad)^2 \sqrt[4]{a+bx^2}} + \frac{d^2 \int \frac{1}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{(bc-ad)^2} - \frac{bd(3bc+5ad)}{(bc-ad)^2} \\
 &= \frac{2bx}{5a(bc-ad)(a+bx^2)^{5/4}} + \frac{2b(3bc-8ad)x}{5a^2(bc-ad)^2 \sqrt[4]{a+bx^2}} + \frac{\left(2d^2 \sqrt{-\frac{bx^2}{a}}\right) \text{Subst} \left(\int \frac{1}{\sqrt{1-\frac{x^2}{a}}} dx, x, \sqrt[4]{a+bx^2} \right)}{(bc-ad)^2} \\
 &= \frac{2bx}{5a(bc-ad)(a+bx^2)^{5/4}} - \frac{\left(d^{3/2} \sqrt{-\frac{bx^2}{a}}\right) \text{Subst} \left(\int \frac{1}{(\sqrt{-bc+ad}-\sqrt{d}x^2)\sqrt{1-\frac{x^2}{a}}} dx, x, \sqrt[4]{a+bx^2} \right)}{(bc-ad)^2 x} \\
 &= \frac{2bx}{5a(bc-ad)(a+bx^2)^{5/4}} + \frac{2\sqrt{b}(3bc-8ad)\sqrt[4]{1+\frac{bx^2}{a}} E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\right) \Big|_2}{5a^{3/2}(bc-ad)^2 \sqrt[4]{a+bx^2}} + \frac{\sqrt[4]{a} d}{(bc-ad)^2}
 \end{aligned}$$

Mathematica [C] time = 0.66, size = 419, normalized size = 1.53

$$x \frac{\left(\frac{bdx^2 \sqrt{\frac{bx^2}{a}+1} (8ad-3bc) F_1\left(\frac{3}{2}; \frac{1}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c} - \frac{6\left(bx^2(c+dx^2)(9a^2d-4ab(c-2dx^2))-3b^2cx^2\right)\left(4adF_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)+bcF_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)\right)}{(a+bx^2)(c+dx^2)\left(x^2\left(4adF_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)+bcF_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)\right)\right)}}{15a^2 \sqrt[4]{a+bx^2} (bc-ad)^2}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[1/((a + b*x^2)^(9/4)*(c + d*x^2)), x]
```

```
[Out] (x*((b*d*(-3*b*c + 8*a*d)*x^2*(1 + (b*x^2)/a)^(1/4)*AppellF1[3/2, 1/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])/c - (6*(3*a*c*(5*a^3*d^2 + 3*b^3*c*x^2*(c + 2*d*x^2) - a^2*b*d*(10*c + 13*d*x^2) + a*b^2*(5*c^2 - 16*d^2*x^4))*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + b*x^2*(c + d*x^2)*(9*a^2*d - 3*b^2*c*x^2 - 4*a*b*(c - 2*d*x^2))*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/((a + b*x^2)*(c + d*x^2)*(-6*a*c*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/((15*a^2*(b*c - a*d)^2*(a + b*x^2)^(1/4))
```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*x^2+a)^(9/4)/(d*x^2+c), x, algorithm="fricas")
```

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{9}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(9/4)/(d*x^2+c),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(9/4)*(d*x^2 + c)), x)

maple [F] time = 0.35, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{9}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(9/4)/(d*x^2+c),x)

[Out] int(1/(b*x^2+a)^(9/4)/(d*x^2+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{9}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(9/4)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(9/4)*(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{9}{4}}(dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(9/4)*(c + d*x^2)),x)

[Out] int(1/((a + b*x^2)^(9/4)*(c + d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{9}{4}}(c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(9/4)/(d*x**2+c),x)

[Out] Integral(1/((a + b*x**2)**(9/4)*(c + d*x**2)), x)

$$3.330 \quad \int \frac{1}{(a+bx^2)^{11/4}(c+dx^2)} dx$$

Optimal. Leaf size=304

$$\frac{2\sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} (5bc - 12ad) \operatorname{EllipticF}\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), 2\right)}{21a^{3/2} (a + bx^2)^{3/4} (bc - ad)^2} + \frac{2bx(5bc - 12ad)}{21a^2 (a + bx^2)^{3/4} (bc - ad)^2} + \frac{\sqrt[4]{a} d^2 \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a} \sqrt{b}}{\sqrt{ad-bc}}\right)}{x(bc - ad)^3}$$

[Out] $2/7*b*x/a/(-a*d+b*c)/(b*x^2+a)^{(7/4)}+2/21*b*(-12*a*d+5*b*c)*x/a^2/(-a*d+b*c)^2/(b*x^2+a)^{(3/4)}+2/21*(-12*a*d+5*b*c)*(1+b*x^2/a)^{(3/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\operatorname{EllipticF}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})), 2^{(1/2)})*b^{(1/2)}/a^{(3/2)}/(-a*d+b*c)^2/(b*x^2+a)^{(3/4)}+a^{(1/4)}*d^2*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, -a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/(-a*d+b*c)^3/x+a^{(1/4)}*d^2*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/(-a*d+b*c)^3/x$

Rubi [A] time = 0.34, antiderivative size = 304, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {414, 527, 530, 233, 231, 401, 108, 409, 1218}

$$\frac{2bx(5bc - 12ad)}{21a^2 (a + bx^2)^{3/4} (bc - ad)^2} + \frac{2\sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} (5bc - 12ad) F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{21a^{3/2} (a + bx^2)^{3/4} (bc - ad)^2} + \frac{\sqrt[4]{a} d^2 \sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a} \sqrt{b}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt{a} \sqrt{b}}{\sqrt{ad-bc}}\right)\right)}{x(bc - ad)^3}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(11/4)*(c + d*x^2)), x]

[Out] $(2*b*x)/(7*a*(b*c - a*d)*(a + b*x^2)^{(7/4)}) + (2*b*(5*b*c - 12*a*d)*x)/(21*a^2*(b*c - a*d)^2*(a + b*x^2)^{(3/4)}) + (2*\operatorname{Sqrt}[b]*(5*b*c - 12*a*d)*(1 + (b*x^2)/a)^{(3/4)}*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/2, 2])/(21*a^{(3/2)}*(b*c - a*d)^2*(a + b*x^2)^{(3/4)}) + (a^{(1/4)}*d^2*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[-((\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b*c) + a*d])], \operatorname{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/((b*c - a*d)^3*x) + (a^{(1/4)}*d^2*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b*c) + a*d]), \operatorname{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/((b*c - a*d)^3*x)$

Rule 108

Int[1/(((a_) + (b_)*(x_))*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))^(3/4)), x_Symbol] :> Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 231

Int[((a_) + (b_)*(x_)^2)^(-3/4), x_Symbol] :> Simp[(2*EllipticF[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(3/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 233

Int[((a_) + (b_)*(x_)^2)^(-3/4), x_Symbol] :> Dist[(1 + (b*x^2)/a)^(3/4)/(a + b*x^2)^(3/4), Int[1/(1 + (b*x^2)/a)^(3/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 401

Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 409

Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 530

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a+bx^2)^{11/4}(c+dx^2)} dx &= \frac{2bx}{7a(bc-ad)(a+bx^2)^{7/4}} - \frac{2 \int \frac{\frac{1}{2}(-5bc+7ad) - \frac{5}{2}bdx^2}{(a+bx^2)^{7/4}(c+dx^2)} dx}{7a(bc-ad)} \\
&= \frac{2bx}{7a(bc-ad)(a+bx^2)^{7/4}} + \frac{2b(5bc-12ad)x}{21a^2(bc-ad)^2(a+bx^2)^{3/4}} + \frac{4 \int \frac{\frac{1}{4}(5b^2c^2-12abcd+21a^2d^2)}{(a+bx^2)^{3/4}(c+dx^2)} dx}{21a^2(bc-ad)^2} \\
&= \frac{2bx}{7a(bc-ad)(a+bx^2)^{7/4}} + \frac{2b(5bc-12ad)x}{21a^2(bc-ad)^2(a+bx^2)^{3/4}} + \frac{d^2 \int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx}{(bc-ad)^2} \\
&= \frac{2bx}{7a(bc-ad)(a+bx^2)^{7/4}} + \frac{2b(5bc-12ad)x}{21a^2(bc-ad)^2(a+bx^2)^{3/4}} + \frac{\left(d^2 \sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{c+dx^2}} dx\right)}{2(bc-ad)^2} \\
&= \frac{2bx}{7a(bc-ad)(a+bx^2)^{7/4}} + \frac{2b(5bc-12ad)x}{21a^2(bc-ad)^2(a+bx^2)^{3/4}} + \frac{2\sqrt{b}(5bc-12ad)\left(1 + \frac{bx^2}{a}\right)}{21a^{3/2}(bc-ad)^2} \\
&= \frac{2bx}{7a(bc-ad)(a+bx^2)^{7/4}} + \frac{2b(5bc-12ad)x}{21a^2(bc-ad)^2(a+bx^2)^{3/4}} + \frac{2\sqrt{b}(5bc-12ad)\left(1 + \frac{bx^2}{a}\right)}{21a^{3/2}(bc-ad)^2} \\
&= \frac{2bx}{7a(bc-ad)(a+bx^2)^{7/4}} + \frac{2b(5bc-12ad)x}{21a^2(bc-ad)^2(a+bx^2)^{3/4}} + \frac{2\sqrt{b}(5bc-12ad)\left(1 + \frac{bx^2}{a}\right)}{21a^{3/2}(bc-ad)^2}
\end{aligned}$$

Mathematica [C] time = 0.78, size = 431, normalized size = 1.42

$$x \frac{6 \left(bx^2(c+dx^2)(15a^2d+ab(12dx^2-8c)-5b^2cx^2) \left(4adF_1\left(\frac{3}{2}; \frac{3}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + 3bcF_1\left(\frac{3}{2}; \frac{7}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) + 3ac(21a^3d^2-3a^2bd(14c+3dx^2)+ab^2(21c^2-20cdx^2-24d^2x^4)) \right)}{(a+bx^2)(c+dx^2) \left(x^2 \left(4adF_1\left(\frac{3}{2}; \frac{3}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + 3bcF_1\left(\frac{3}{2}; \frac{7}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) - 6acF_1\left(\frac{1}{2}; \frac{3}{4}, 1; \frac{3}{2}; -\frac{(b*x^2)}{a}, -\frac{(d*x^2)}{c}\right) \right)}{63a^2(a+bx^2)^{3/4}(bc-ad)^2}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(11/4)*(c + d*x^2)), x]

[Out]
$$\frac{-1/63 * (x * ((b*d*(-5*b*c + 12*a*d) * x^2 * (1 + (b*x^2)/a)^{3/4} * \text{AppellF1}[3/2, 3/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]) / c + (6 * (3*a*c*(21*a^3*d^2 + 5*b^3*c*x^2*(3*c + 2*d*x^2) - 3*a^2*b*d*(14*c + 3*d*x^2) + a*b^2*(21*c^2 - 20*c*d*x^2 - 24*d^2*x^4)) * \text{AppellF1}[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + b*x^2*(c + d*x^2)*(15*a^2*d - 5*b^2*c*x^2 + a*b*(-8*c + 12*d*x^2)) * (4*a*d*\text{AppellF1}[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*\text{AppellF1}[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])) / ((a + b*x^2)*(c + d*x^2)*(-6*a*c*\text{AppellF1}[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^2*(4*a*d*\text{AppellF1}[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*\text{AppellF1}[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])) / (a^2*(b*c - a*d)^2*(a + b*x^2)^{3/4})$$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(11/4)/(d*x^2+c),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{11}{4}} (dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(11/4)/(d*x^2+c),x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(11/4)*(d*x^2 + c)), x)

maple [F] time = 0.35, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{11}{4}} (dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(11/4)/(d*x^2+c),x)

[Out] int(1/(b*x^2+a)^(11/4)/(d*x^2+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{11}{4}} (dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(11/4)/(d*x^2+c),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(11/4)*(d*x^2 + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{11/4} (dx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(11/4)*(c + d*x^2)),x)

[Out] int(1/((a + b*x^2)^(11/4)*(c + d*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{11}{4}} (c + dx^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(11/4)/(d*x**2+c),x)

[Out] Integral(1/((a + b*x**2)**(11/4)*(c + d*x**2)), x)

$$3.331 \quad \int \frac{(a+bx^2)^{7/4}}{(c+dx^2)^2} dx$$

Optimal. Leaf size=340

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \sqrt{ad-bc} (2ad+5bc) \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{4cd^{5/2}x} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} \sqrt{ad-bc} (2ad+5bc) \Pi\left(\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{4cd^{5/2}x}$$

[Out] $\frac{1}{2}bx(-ad+5bc)x/c/d^2/(bx^2+a)^{1/4} - \frac{1}{2}(-ad+bc)x*(bx^2+a)^{3/4}/c/d/(dx^2+c) - \frac{1}{2}(-ad+5bc)*(1+bx^2/a)^{1/4}*(\cos(1/2*\arctan(x*b^{1/2}/a^{1/2})))^2)^{1/2}/\cos(1/2*\arctan(x*b^{1/2}/a^{1/2}))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{1/2}/a^{1/2})), 2^{1/2})*a^{1/2}*b^{1/2}/c/d^2/(bx^2+a)^{1/4} + \frac{1}{4}a^{1/4}*(2ad+5bc)*\text{EllipticPi}((bx^2+a)^{1/4}/a^{1/4}, -a^{1/2}*d^{1/2}/(ad-bc)^{1/2}, I)*(ad-bc)^{1/2}*(-bx^2/a)^{1/2}/c/d^{5/2}/x - \frac{1}{4}a^{1/4}*(2ad+5bc)*\text{EllipticPi}((bx^2+a)^{1/4}/a^{1/4}, a^{1/2}*d^{1/2}/(ad-bc)^{1/2}, I)*(ad-bc)^{1/2}*(-bx^2/a)^{1/2}/c/d^{5/2}/x$

Rubi [A] time = 0.29, antiderivative size = 340, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {413, 530, 229, 227, 196, 399, 490, 1218}

$$\frac{bx(5bc-ad)}{2cd^2\sqrt[4]{a+bx^2}} - \frac{\sqrt{a}\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}(5bc-ad)E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{2cd^2\sqrt[4]{a+bx^2}} + \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}\sqrt{ad-bc}(2ad+5bc)\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{4cd^{5/2}x}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(7/4)/(c + d*x^2)^2, x]

[Out] $\frac{b*(5*b*c - a*d)*x}{(2*c*d^2*(a + b*x^2)^{1/4})} - \frac{((b*c - a*d)*x*(a + b*x^2)^{3/4})}{(2*c*d*(c + d*x^2))} - \frac{(\text{Sqrt}[a]*\text{Sqrt}[b]*(5*b*c - a*d)*(1 + (b*x^2)/a)^{1/4}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])}{(2*c*d^2*(a + b*x^2)^{1/4})} + \frac{(a^{1/4}*\text{Sqrt}[-(b*c) + a*d]*(5*b*c + 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d])), \text{ArcSin}[(a + b*x^2)^{1/4}/a^{1/4}], -1])}{(4*c*d^{5/2}*x)} - \frac{(a^{1/4}*\text{Sqrt}[-(b*c) + a*d]*(5*b*c + 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{1/4}/a^{1/4}], -1])}{(4*c*d^{5/2}*x)}$

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 227

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2*x)/(a + b*x^2)^(1/4), x] - Dist[a, Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 229

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a + b*x^2)^(1/4), Int[1/(1 + (b*x^2)/a)^(1/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 399

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dis
t[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x
^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0]
```

Rule 413

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p +
1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q -
2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p
+ q) + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,
0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rule 490

```
Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] :=
With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s
/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/(
(r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c
- a*d, 0]
```

Rule 530

```
Int((((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p
, n}, x]
```

Rule 1218

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*
Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^{7/4}}{(c+dx^2)^2} dx &= -\frac{(bc-ad)x(a+bx^2)^{3/4}}{2cd(c+dx^2)} + \frac{\int \frac{a(bc+ad)+\frac{1}{2}b(5bc-ad)x^2}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{2cd} \\
&= -\frac{(bc-ad)x(a+bx^2)^{3/4}}{2cd(c+dx^2)} + \frac{(b(5bc-ad)) \int \frac{1}{\sqrt[4]{a+bx^2}} dx}{4cd^2} - \frac{((bc-ad)(5bc+2ad)) \int \frac{1}{\sqrt[4]{a+bx^2}(c+dx^2)}}{4cd^2} \\
&= -\frac{(bc-ad)x(a+bx^2)^{3/4}}{2cd(c+dx^2)} - \frac{\left((bc-ad)(5bc+2ad)\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{x^2}{\sqrt{1-\frac{x^4}{a}(bc-ad+dx^4)}} dx, x, \sqrt[4]{a+bx^2}\right)}{2cd^2x} \\
&= \frac{b(5bc-ad)x}{2cd^2\sqrt[4]{a+bx^2}} - \frac{(bc-ad)x(a+bx^2)^{3/4}}{2cd(c+dx^2)} + \frac{\left((bc-ad)(5bc+2ad)\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{(\sqrt{-bc+ad}}\right)}{4cd^{5/2}x} \\
&= \frac{b(5bc-ad)x}{2cd^2\sqrt[4]{a+bx^2}} - \frac{(bc-ad)x(a+bx^2)^{3/4}}{2cd(c+dx^2)} - \frac{\sqrt{a}\sqrt{b}(5bc-ad)\sqrt[4]{1+\frac{bx^2}{a}} E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\right)}{2cd^2\sqrt[4]{a+bx^2}} \Big|_2
\end{aligned}$$

Mathematica [C] time = 0.34, size = 340, normalized size = 1.00

$$x \frac{\left(6c\left(x^2(a+bx^2)(ad-bc)\left(4adF_1\left(\frac{3}{2};\frac{1}{4},2;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)+bcF_1\left(\frac{3}{2};\frac{5}{4},1;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)\right)-6ac(2a^2d+abdx^2-b^2cx^2)F_1\left(\frac{1}{2};\frac{1}{4},1;\frac{3}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)\right)}{(c+dx^2)\left(x^2\left(4adF_1\left(\frac{3}{2};\frac{1}{4},2;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)+bcF_1\left(\frac{3}{2};\frac{5}{4},1;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)\right)-6acF_1\left(\frac{1}{2};\frac{1}{4},1;\frac{3}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)\right)} - bx^2 \sqrt[4]{\frac{bx^2}{a}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(7/4)/(c + d*x^2)^2,x]

[Out] (x*(-(b*(-5*b*c + a*d))*x^2*(1 + (b*x^2)/a)^(1/4)*AppellF1[3/2, 1/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)]) + (6*c*(-6*a*c*(2*a^2*d - b^2*c*x^2 + a*b*d*x^2)*AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -((d*x^2)/c)] + (-b*c) + a*d)*x^2*(a + b*x^2)*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)])))/((c + d*x^2)*(-6*a*c*AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -((d*x^2)/c)] + x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)])))/((12*c^2*d*(a + b*x^2)^(1/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{7/4}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(7/4)/(d*x^2 + c)^2, x)

maple [F] time = 0.35, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{7}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(7/4)/(d*x^2+c)^2,x)

[Out] int((b*x^2+a)^(7/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{7}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(7/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(7/4)/(d*x^2 + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{7/4}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(7/4)/(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(7/4)/(c + d*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{7}{4}}}{(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(7/4)/(d*x**2+c)**2,x)

[Out] Integral((a + b*x**2)**(7/4)/(c + d*x**2)**2, x)

$$3.332 \quad \int \frac{(a+bx^2)^{5/4}}{(c+dx^2)^2} dx$$

Optimal. Leaf size=279

$$\frac{\sqrt{a} \sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} (ad + 3bc) \operatorname{EllipticF}\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), 2\right) \sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} (2ad + 3bc) \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\right)}{2cd^2 (a + bx^2)^{3/4} 4cd^2 x}$$

[Out] $-1/2*(-a*d+b*c)*x*(b*x^2+a)^{(1/4)}/c/d/(d*x^2+c)+1/2*(a*d+3*b*c)*(1+b*x^2/a)^{(3/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\operatorname{EllipticF}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})), 2^{(1/2)})*a^{(1/2)}*b^{(1/2)}/c/d^2/(b*x^2+a)^{(3/4)}-1/4*a^{(1/4)}*(2*a*d+3*b*c)*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, -a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/c/d^2/x-1/4*a^{(1/4)}*(2*a*d+3*b*c)*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/c/d^2/x$

Rubi [A] time = 0.27, antiderivative size = 279, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {413, 530, 233, 231, 401, 108, 409, 1218}

$$\frac{\sqrt{a} \sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} (ad + 3bc) F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right) \sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} (2ad + 3bc) \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right) \sqrt[4]{a}}{2cd^2 (a + bx^2)^{3/4} 4cd^2 x}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(5/4)/(c + d*x^2)^2,x]

[Out] $-\left((b*c - a*d)*x*(a + b*x^2)^{(1/4)}\right)/(2*c*d*(c + d*x^2)) + (\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b]*(3*b*c + a*d)*(1 + (b*x^2)/a)^{(3/4)}*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/2, 2])/ (2*c*d^2*(a + b*x^2)^{(3/4)}) - (a^{(1/4)}*(3*b*c + 2*a*d)*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[-((\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/ \operatorname{Sqrt}[-(b*c) + a*d]), \operatorname{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/ (4*c*d^2*x) - (a^{(1/4)}*(3*b*c + 2*a*d)*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/ \operatorname{Sqrt}[-(b*c) + a*d], \operatorname{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/ (4*c*d^2*x)$

Rule 108

Int[1/(((a_.) + (b_.)*(x_)) * Sqrt[(c_.) + (d_.)*(x_)] * ((e_.) + (f_.)*(x_))^(3/4)), x_Symbol] := Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 231

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2*EllipticF[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(3/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 233

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(3/4)/(a + b*x^2)^(3/4), Int[1/(1 + (b*x^2)/a)^(3/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 401

```
Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dis
t[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(
c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]
```

Rule 409

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[
1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(
2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[b*c - a*d, 0]
```

Rule 413

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[((a*d - c*b)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(a*b*n*(p +
1)), x] - Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q -
2)*Simp[c*(a*d - c*b*(n*(p + 1) + 1)) + d*(a*d*(n*(q - 1) + 1) - b*c*(n*(p
+ q) + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d,
0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rule 530

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p,
n}, x]
```

Rule 1218

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*
Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^{5/4}}{(c+dx^2)^2} dx &= -\frac{(bc-ad)x\sqrt[4]{a+bx^2}}{2cd(c+dx^2)} + \frac{\int \frac{a(bc+ad)+\frac{1}{2}b(3bc+ad)x^2}{(a+bx^2)^{3/4}(c+dx^2)} dx}{2cd} \\
&= -\frac{(bc-ad)x\sqrt[4]{a+bx^2}}{2cd(c+dx^2)} + \frac{(b(3bc+ad)) \int \frac{1}{(a+bx^2)^{3/4}} dx}{4cd^2} - \frac{((bc-ad)(3bc+2ad)) \int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx}{4cd^2} \\
&= -\frac{(bc-ad)x\sqrt[4]{a+bx^2}}{2cd(c+dx^2)} - \frac{\left((bc-ad)(3bc+2ad)\sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-\frac{bx}{a}}(a+bx)^{3/4}(c+dx)} dx, x, x^2\right)}{8cd^2x} \\
&= -\frac{(bc-ad)x\sqrt[4]{a+bx^2}}{2cd(c+dx^2)} + \frac{\sqrt{a}\sqrt{b}(3bc+ad)\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\right)\Big|_2}{2cd^2(a+bx^2)^{3/4}} + \frac{\left((bc-ad)(3bc+2ad)\right) \int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx}{4cd^2} \\
&= -\frac{(bc-ad)x\sqrt[4]{a+bx^2}}{2cd(c+dx^2)} + \frac{\sqrt{a}\sqrt{b}(3bc+ad)\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\right)\Big|_2}{2cd^2(a+bx^2)^{3/4}} - \frac{\left((3bc+2ad)\int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx\right)}{4cd^2} \\
&= -\frac{(bc-ad)x\sqrt[4]{a+bx^2}}{2cd(c+dx^2)} + \frac{\sqrt{a}\sqrt{b}(3bc+ad)\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\right)\Big|_2}{2cd^2(a+bx^2)^{3/4}} - \frac{\sqrt[4]{a}(3bc+2ad)\int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx}{4cd^2}
\end{aligned}$$

Mathematica [C] time = 0.33, size = 341, normalized size = 1.22

$$x \frac{\left(6c\left(x^2(a+bx^2)(ad-bc)\left(4adF_1\left(\frac{3}{2};\frac{3}{4},2;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)+3bcF_1\left(\frac{3}{2};\frac{7}{4},1;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)\right)-6ac(2a^2d+abdx^2-b^2cx^2)F_1\left(\frac{1}{2};\frac{3}{4},1;\frac{3}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)\right)}{(c+dx^2)\left(x^2\left(4adF_1\left(\frac{3}{2};\frac{3}{4},2;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)+3bcF_1\left(\frac{3}{2};\frac{7}{4},1;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)\right)-6acF_1\left(\frac{1}{2};\frac{3}{4},1;\frac{3}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)\right)} + bx^2 \left(\frac{bx^2}{a}\right)$$

$$12c^2d(a+bx^2)^{3/4}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(5/4)/(c + d*x^2)^2,x]

[Out] (x*(b*(3*b*c + a*d)*x^2*(1 + (b*x^2)/a)^(3/4)*AppellF1[3/2, 3/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + (6*c*(-6*a*c*(2*a^2*d - b^2*c*x^2 + a*b*d*x^2)*AppellF1[1/2, 3/4, 1, 3/2, -(b*x^2)/a, -((d*x^2)/c)] + (-b*c) + a*d)*x^2*(a + b*x^2)*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)])))/((c + d*x^2)*(-6*a*c*AppellF1[1/2, 3/4, 1, 3/2, -(b*x^2)/a, -((d*x^2)/c)] + x^2*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)])))/((12*c^2*d*(a + b*x^2)^(3/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(5/4)/(d*x^2 + c)^2, x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(5/4)/(d*x^2+c)^2,x)

[Out] int((b*x^2+a)^(5/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(5/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(5/4)/(d*x^2 + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{\frac{5}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(5/4)/(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(5/4)/(c + d*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{5}{4}}}{(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(5/4)/(d*x**2+c)**2,x)

[Out] Integral((a + b*x**2)**(5/4)/(c + d*x**2)**2, x)

$$3.333 \quad \int \frac{(a+bx^2)^{3/4}}{(c+dx^2)^2} dx$$

Optimal. Leaf size=309

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} (2ad + bc) \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{4cd^{3/2}x\sqrt{ad-bc}} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} (2ad + bc) \Pi\left(\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{4cd^{3/2}x\sqrt{ad-bc}}$$

[Out] $-1/2*b*x/c/d/(b*x^2+a)^{(1/4)}+1/2*x*(b*x^2+a)^{(3/4)}/c/(d*x^2+c)+1/2*(1+b*x^2/a)^{(1/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})), 2^{(1/2)})*a^{(1/2)}*b^{(1/2)}/c/d/(b*x^2+a)^{(1/4)}+1/4*a^{(1/4)}*(2*a*d+b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, -a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/c/d^{(3/2)}/x/(a*d-b*c)^{(1/2)}-1/4*a^{(1/4)}*(2*a*d+b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/c/d^{(3/2)}/x/(a*d-b*c)^{(1/2)}$

Rubi [A] time = 0.22, antiderivative size = 309, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {412, 530, 229, 227, 196, 399, 490, 1218}

$$\frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} (2ad + bc) \Pi\left(-\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{4cd^{3/2}x\sqrt{ad-bc}} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} (2ad + bc) \Pi\left(\frac{\sqrt{a} \sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{4cd^{3/2}x\sqrt{ad-bc}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^(3/4)/(c + d*x^2)^2, x]

[Out] $-(b*x)/(2*c*d*(a + b*x^2)^{(1/4)}) + (x*(a + b*x^2)^{(3/4)})/(2*c*(c + d*x^2)) + (\text{Sqrt}[a]*\text{Sqrt}[b]*(1 + (b*x^2)/a)^{(1/4)}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/(2*c*d*(a + b*x^2)^{(1/4)}) + (a^{(1/4)}*(b*c + 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d])], \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*d^{(3/2)}*\text{Sqrt}[-(b*c) + a*d]*x) - (a^{(1/4)}*(b*c + 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*d^{(3/2)}*\text{Sqrt}[-(b*c) + a*d]*x)$

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 227

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2*x)/(a + b*x^2)^(1/4), x] - Dist[a, Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 229

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a + b*x^2)^(1/4), Int[1/(1 + (b*x^2)/a)^(1/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 399

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dis
t[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x
^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0]
```

Rule 412

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*n*(p + 1)), x] + Dist[1/(
a*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(n*(p + 1)
+ 1) + d*(n*(p + q + 1) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n}, x]
&& NeQ[b*c - a*d, 0] && LtQ[p, -1] && LtQ[0, q, 1] && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 490

```
Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] :=
With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s
/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/(
(r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c
- a*d, 0]
```

Rule 530

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p
, n}, x]
```

Rule 1218

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*
Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a+bx^2)^{3/4}}{(c+dx^2)^2} dx &= \frac{x(a+bx^2)^{3/4}}{2c(c+dx^2)} - \frac{\int \frac{-a+\frac{bx^2}{2}}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{2c} \\
&= \frac{x(a+bx^2)^{3/4}}{2c(c+dx^2)} - \frac{b \int \frac{1}{\sqrt[4]{a+bx^2}} dx}{4cd} + \frac{(bc+2ad) \int \frac{1}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{4cd} \\
&= \frac{x(a+bx^2)^{3/4}}{2c(c+dx^2)} + \frac{\left((bc+2ad)\sqrt{-\frac{bx^2}{a}} \right) \text{Subst} \left(\int \frac{x^2}{\sqrt{1-\frac{x^4}{a}(bc-ad+dx^4)}} dx, x, \sqrt[4]{a+bx^2} \right)}{2cdx} - \frac{\left(b\sqrt[4]{1+\frac{bx^2}{a}} \right) \text{Subst} \left(\int \frac{1}{(\sqrt{-bc+ad}-\sqrt{d}x^2)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2} \right)}{4cd^{3/2}x} \\
&= -\frac{bx}{2cd\sqrt[4]{a+bx^2}} + \frac{x(a+bx^2)^{3/4}}{2c(c+dx^2)} - \frac{\left((bc+2ad)\sqrt{-\frac{bx^2}{a}} \right) \text{Subst} \left(\int \frac{1}{(\sqrt{-bc+ad}-\sqrt{d}x^2)\sqrt{1-\frac{x^4}{a}}} dx, x, \sqrt[4]{a+bx^2} \right)}{4cd^{3/2}x} \\
&= -\frac{bx}{2cd\sqrt[4]{a+bx^2}} + \frac{x(a+bx^2)^{3/4}}{2c(c+dx^2)} + \frac{\sqrt{a}\sqrt{b}\sqrt[4]{1+\frac{bx^2}{a}} E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{2cd\sqrt[4]{a+bx^2}} + \frac{\sqrt[4]{a}(bc+2ad)\sqrt{b}}{2cd\sqrt[4]{a+bx^2}}
\end{aligned}$$

Mathematica [C] time = 0.20, size = 232, normalized size = 0.75

$$x \left(\frac{6 \left(\frac{a+bx^2}{c} \frac{{}_6a^2F_1\left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{x^2 \left(4adF_1\left(\frac{3}{2}, \frac{1}{4}, 2, \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + bcF_1\left(\frac{3}{2}, \frac{5}{4}, 1, \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) - 6acF_1\left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c+dx^2} - \frac{bx^2 \sqrt[4]{\frac{bx^2}{a}+1} F_1\left(\frac{3}{2}, \frac{1}{4}, 1, \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c^2} \right)}{12\sqrt[4]{a+bx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(3/4)/(c + d*x^2)^2,x]

[Out] (x*(-((b*x^2*(1 + (b*x^2)/a)^(1/4)*AppellF1[3/2, 1/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))/c^2) + (6*((a + b*x^2)/c - (6*a^2*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)])/(-6*a*c*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/(c + d*x^2))/(12*(a + b*x^2)^(1/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(3/4)/(d*x^2 + c)^2, x)

maple [F] time = 0.35, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(3/4)/(d*x^2+c)^2,x)

[Out] int((b*x^2+a)^(3/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(3/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(3/4)/(d*x^2 + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{\frac{3}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(3/4)/(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(3/4)/(c + d*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^{\frac{3}{4}}}{(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(3/4)/(d*x**2+c)**2,x)

[Out] Integral((a + b*x**2)**(3/4)/(c + d*x**2)**2, x)

$$3.334 \quad \int \frac{\sqrt[4]{a+bx^2}}{(c+dx^2)^2} dx$$

Optimal. Leaf size=278

$$\frac{\sqrt{a} \sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} \operatorname{EllipticF}\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), 2\right)}{2cd(a+bx^2)^{3/4}} + \frac{x\sqrt[4]{a+bx^2}}{2c(c+dx^2)} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} (bc-2ad) \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right)\right)}{4cdx(bc-ad)}$$

[Out] $\frac{1}{2}x(bx^2+a)^{1/4}/c/(dx^2+c) + \frac{1}{2}(1+bx^2/a)^{3/4}(\cos(1/2\arctan(xb^{1/2}/a^{1/2}))^2)^{1/2}/\cos(1/2\arctan(xb^{1/2}/a^{1/2}))\operatorname{EllipticF}(\sin(1/2\arctan(xb^{1/2}/a^{1/2})), 2^{1/2})a^{1/2}b^{1/2}/c/d/(bx^2+a)^{3/4} - 1/4a^{1/4}(-2ad+bc)\operatorname{EllipticPi}((bx^2+a)^{1/4}/a^{1/4}, -a^{1/2}d^{1/2}/(ad-bc)^{1/2}, I) (-bx^2/a)^{1/2}/c/d/(-ad+bc)/x - 1/4a^{1/4}(-2ad+bc)\operatorname{EllipticPi}((bx^2+a)^{1/4}/a^{1/4}, a^{1/2}d^{1/2}/(ad-bc)^{1/2}, I) (-bx^2/a)^{1/2}/c/d/(-ad+bc)/x$

Rubi [A] time = 0.21, antiderivative size = 278, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {412, 530, 233, 231, 401, 108, 409, 1218}

$$\frac{x\sqrt[4]{a+bx^2}}{2c(c+dx^2)} + \frac{\sqrt{a} \sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{2cd(a+bx^2)^{3/4}} - \frac{\sqrt[4]{a} \sqrt{-\frac{bx^2}{a}} (bc-2ad) \Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt[4]{bx^2+a}}{\sqrt[4]{a}}\right) \middle| -1\right)}{4cdx(bc-ad)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a+bx^2)^{1/4}/(c+dx^2)^2, x]$

[Out] $(x(a+bx^2)^{1/4})/(2c(c+dx^2)) + (\operatorname{Sqrt}[a]\operatorname{Sqrt}[b](1+(bx^2)/a)^{3/4}\operatorname{EllipticF}[\operatorname{ArcTan}[\operatorname{Sqrt}[b]x]/\operatorname{Sqrt}[a]/2, 2])/(2cd(a+bx^2)^{3/4}) - (a^{1/4}(bc-2ad)\operatorname{Sqrt}[-(bx^2)/a]\operatorname{EllipticPi}[-(\operatorname{Sqrt}[a]\operatorname{Sqrt}[d])/\operatorname{Sqrt}[-(bc)+ad], \operatorname{ArcSin}[(a+bx^2)^{1/4}/a^{1/4}], -1])/(4cd(bc-a*d)x) - (a^{1/4}(bc-2ad)\operatorname{Sqrt}[-(bx^2)/a]\operatorname{EllipticPi}[(\operatorname{Sqrt}[a]\operatorname{Sqrt}[d])/\operatorname{Sqrt}[-(bc)+ad], \operatorname{ArcSin}[(a+bx^2)^{1/4}/a^{1/4}], -1])/(4cd(bc-a*d)x)$

Rule 108

$\operatorname{Int}[1/(((a_.) + (b_.)(x_.))\operatorname{Sqrt}[(c_.) + (d_.)(x_.)]*((e_.) + (f_.)(x_.))^{3/4}), x_Symbol] \rightarrow \operatorname{Dist}[-4, \operatorname{Subst}[\operatorname{Int}[1/((b*e - a*f - b*x^4)\operatorname{Sqrt}[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^{1/4}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \operatorname{GtQ}[-(f/(d*e - c*f)), 0]$

Rule 231

$\operatorname{Int}[(a_.) + (b_.)(x_.)^2]^{-3/4}, x_Symbol] \rightarrow \operatorname{Simp}[(2*\operatorname{EllipticF}[(1*\operatorname{ArcTan}[\operatorname{Rt}[b/a, 2]*x])/2, 2])/(a^{3/4}*\operatorname{Rt}[b/a, 2]), x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \operatorname{GtQ}[a, 0] \ \&\& \operatorname{PosQ}[b/a]$

Rule 233

$\operatorname{Int}[(a_.) + (b_.)(x_.)^2]^{-3/4}, x_Symbol] \rightarrow \operatorname{Dist}[(1 + (bx^2)/a)^{3/4}/(a + bx^2)^{3/4}, \operatorname{Int}[1/(1 + (bx^2)/a)^{3/4}, x], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \operatorname{PosQ}[a]$

Rule 401

Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 409

Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 412

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*n*(p + 1)), x] + Dist[1/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(n*(p + 1) + 1) + d*(n*(p + q + 1) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && LtQ[0, q, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 530

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt[4]{a+bx^2}}{(c+dx^2)^2} dx &= \frac{x\sqrt[4]{a+bx^2}}{2c(c+dx^2)} - \frac{\int \frac{-a-\frac{bx^2}{2}}{(a+bx^2)^{3/4}(c+dx^2)} dx}{2c} \\
&= \frac{x\sqrt[4]{a+bx^2}}{2c(c+dx^2)} + \frac{b \int \frac{1}{(a+bx^2)^{3/4}} dx}{4cd} - \frac{\left(\frac{bc}{2} - ad\right) \int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx}{2cd} \\
&= \frac{x\sqrt[4]{a+bx^2}}{2c(c+dx^2)} - \frac{\left(\left(\frac{bc}{2} - ad\right) \sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-\frac{bx}{a}}(a+bx)^{3/4}(c+dx)} dx, x, x^2\right)}{4cdx} + \frac{\left(b\left(1 + \frac{bx^2}{a}\right)^{3/4}\right) \int \frac{1}{\sqrt{1-\frac{x}{a}}}}{4cd(a+bx^2)} \\
&= \frac{x\sqrt[4]{a+bx^2}}{2c(c+dx^2)} + \frac{\sqrt{a}\sqrt{b}\left(1 + \frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{2cd(a+bx^2)^{3/4}} + \frac{\left(\left(\frac{bc}{2} - ad\right) \sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x}{a}}}}{cdx}\right)}{cdx} \\
&= \frac{x\sqrt[4]{a+bx^2}}{2c(c+dx^2)} + \frac{\sqrt{a}\sqrt{b}\left(1 + \frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{2cd(a+bx^2)^{3/4}} - \frac{\left(\left(\frac{bc}{2} - ad\right) \sqrt{-\frac{bx^2}{a}}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{x}{a}\right)^{3/4}}\right)}{2cd(bc-a)} \\
&= \frac{x\sqrt[4]{a+bx^2}}{2c(c+dx^2)} + \frac{\sqrt{a}\sqrt{b}\left(1 + \frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{2cd(a+bx^2)^{3/4}} - \frac{\sqrt[4]{a}(bc-2ad)\sqrt{-\frac{bx^2}{a}} \Pi\left(-\frac{\sqrt{a}\sqrt{a}}{\sqrt{-bc+ad}}\right)}{4cd(bc-ad)x}
\end{aligned}$$

Mathematica [C] time = 0.19, size = 232, normalized size = 0.83

$$x \left(\frac{\left(\frac{a+bx^2}{c} - \frac{6a^2 F_1\left(\frac{1}{2}, \frac{3}{4}, 1, \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{x^2 \left(4ad F_1\left(\frac{3}{2}, \frac{3}{4}, 2, \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + 3bc F_1\left(\frac{3}{2}, \frac{7}{4}, 1, \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - 6ac F_1\left(\frac{1}{2}, \frac{3}{4}, 1, \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right)}{c+dx^2} + \frac{bx^2 \left(\frac{bx^2}{a} + 1\right)^{3/4} F_1\left(\frac{3}{2}, \frac{3}{4}, 1, \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c^2} \right)}{12(a+bx^2)^{3/4}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^(1/4)/(c + d*x^2)^2, x]

[Out] (x*((b*x^2*(1 + (b*x^2)/a)^(3/4)*AppellF1[3/2, 3/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])/c^2 + (6*((a + b*x^2)/c - (6*a^2*AppellF1[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)])/(-6*a*c*AppellF1[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^2*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/(c + d*x^2)))/(12*(a + b*x^2)^(3/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{1}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(1/4)/(d*x^2 + c)^2, x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{1}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(1/4)/(d*x^2+c)^2,x)

[Out] int((b*x^2+a)^(1/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^{\frac{1}{4}}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(1/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(1/4)/(d*x^2 + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^{1/4}}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^(1/4)/(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^(1/4)/(c + d*x^2)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[4]{a + bx^2}}{(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**(1/4)/(d*x**2+c)**2,x)

[Out] Integral((a + b*x**2)**(1/4)/(c + d*x**2)**2, x)

$$3.335 \quad \int \frac{1}{\sqrt[4]{a+bx^2} (c+dx^2)^2} dx$$

Optimal. Leaf size=336

$$\frac{bx}{2c\sqrt[4]{a+bx^2}(bc-ad)} - \frac{dx(a+bx^2)^{3/4}}{2c(c+dx^2)(bc-ad)} - \frac{\sqrt{a}\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{2c\sqrt[4]{a+bx^2}(bc-ad)} - \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}(3bc-2ad)\Pi\left(-\frac{bx}{\sqrt[4]{a+bx^2}(bc-ad)}\right)}{4c\sqrt{d}x}$$

[Out] $1/2*b*x/c/(-a*d+b*c)/(b*x^2+a)^{(1/4)}-1/2*d*x*(b*x^2+a)^{(3/4)}/c/(-a*d+b*c)/(d*x^2+c)-1/2*(1+b*x^2/a)^{(1/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*a^{(1/2)}*b^{(1/2)}/c/(-a*d+b*c)/(b*x^2+a)^{(1/4)}-1/4*a^{(1/4)}*(-2*a*d+3*b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/c/(a*d-b*c)^{(3/2)}/x/d^{(1/2)}+1/4*a^{(1/4)}*(-2*a*d+3*b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/c/(a*d-b*c)^{(3/2)}/x/d^{(1/2)}$

Rubi [A] time = 0.25, antiderivative size = 336, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {414, 530, 229, 227, 196, 399, 490, 1218}

$$\frac{bx}{2c\sqrt[4]{a+bx^2}(bc-ad)} - \frac{dx(a+bx^2)^{3/4}}{2c(c+dx^2)(bc-ad)} - \frac{\sqrt{a}\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\middle|2\right)}{2c\sqrt[4]{a+bx^2}(bc-ad)} - \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}(3bc-2ad)\Pi\left(-\frac{bx}{\sqrt[4]{a+bx^2}(bc-ad)}\right)}{4c\sqrt{d}x}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(1/4)*(c + d*x^2)^2), x]

[Out] $(b*x)/(2*c*(b*c - a*d)*(a + b*x^2)^{(1/4)}) - (d*x*(a + b*x^2)^{(3/4)})/(2*c*(b*c - a*d)*(c + d*x^2)) - (\text{Sqrt}[a]*\text{Sqrt}[b]*(1 + (b*x^2)/a)^{(1/4)}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/(2*c*(b*c - a*d)*(a + b*x^2)^{(1/4)}) - (a^{(1/4)}*(3*b*c - 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d])], \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*\text{Sqrt}[d]*(-(b*c) + a*d)^{(3/2)}*x) + (a^{(1/4)}*(3*b*c - 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*\text{Sqrt}[d]*(-(b*c) + a*d)^{(3/2)}*x)$

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 227

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2*x)/(a + b*x^2)^(1/4), x] - Dist[a, Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 229

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a + b*x^2)^(1/4), Int[1/(1 + (b*x^2)/a)^(1/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 399

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dis
t[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x
^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0]
```

Rule 414

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c -
a*d), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d), Int[(a + b*x^n)^(p + 1)*(c +
d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x]
, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1]
&& !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 490

```
Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] :=
With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s
/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/(
(r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c
- a*d, 0]
```

Rule 530

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p
, n}, x]
```

Rule 1218

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*
Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sqrt[4]{a+bx^2} (c+dx^2)^2} dx &= -\frac{dx (a+bx^2)^{3/4}}{2c(bc-ad)(c+dx^2)} + \frac{\int \frac{2bc-ad+\frac{1}{2}bdx^2}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{2c(bc-ad)} \\
&= -\frac{dx (a+bx^2)^{3/4}}{2c(bc-ad)(c+dx^2)} + \frac{b \int \frac{1}{\sqrt[4]{a+bx^2}} dx}{4c(bc-ad)} + \frac{(3bc-2ad) \int \frac{1}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{4c(bc-ad)} \\
&= -\frac{dx (a+bx^2)^{3/4}}{2c(bc-ad)(c+dx^2)} + \frac{\left((3bc-2ad)\sqrt{-\frac{bx^2}{a}} \right) \text{Subst} \left(\int \frac{x^2}{\sqrt{1-\frac{x^4}{a}(bc-ad+dx^4)}} dx, x, \sqrt[4]{a+bx^2} \right)}{2c(bc-ad)x} \\
&= \frac{bx}{2c(bc-ad)\sqrt[4]{a+bx^2}} - \frac{dx (a+bx^2)^{3/4}}{2c(bc-ad)(c+dx^2)} - \frac{\left((3bc-2ad)\sqrt{-\frac{bx^2}{a}} \right) \text{Subst} \left(\int \frac{1}{(\sqrt{-t}} \right)}{4c\sqrt{d}(bc-ad)} \\
&= \frac{bx}{2c(bc-ad)\sqrt[4]{a+bx^2}} - \frac{dx (a+bx^2)^{3/4}}{2c(bc-ad)(c+dx^2)} - \frac{\sqrt{a}\sqrt{b}\sqrt[4]{1+\frac{bx^2}{a}} E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\right)}{2c(bc-ad)\sqrt[4]{a+bx^2}}
\end{aligned}$$

Mathematica [C] time = 0.25, size = 392, normalized size = 1.17

$$\frac{-6acx F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \left(bdx^2 \sqrt{\frac{bx^2}{a}} + 1 (c+dx^2) F_1\left(\frac{3}{2}; \frac{1}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - 6c(2ad-2bc+bdx^2) \right) - dx^2}{12c^2 \sqrt[4]{a+bx^2} (c+dx^2) (bc-ad) \left(x^2 \left(4ad F_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(1/4)*(c + d*x^2)^2), x]

[Out] (-6*a*c*x*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)]*(-6*c*(-2*b*c + 2*a*d + b*d*x^2) + b*d*x^2*(1 + (b*x^2)/a)^(1/4)*(c + d*x^2)*AppellF1[3/2, 1/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]) - d*x^3*(6*c*(a + b*x^2) - b*x^2*(1 + (b*x^2)/a)^(1/4)*(c + d*x^2)*AppellF1[3/2, 1/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))/(12*c^2*(b*c - a*d)*(a + b*x^2)^(1/4)*(c + d*x^2)*(-6*a*c*AppellF1[1/2, 1/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2+a)^{\frac{1}{4}}(dx^2+c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(1/4)*(d*x^2 + c)^2), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{1}{4}} (dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(1/4)/(d*x^2+c)^2,x)

[Out] int(1/(b*x^2+a)^(1/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{1}{4}} (dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(1/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(1/4)*(d*x^2 + c)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{1/4} (dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(1/4)*(c + d*x^2)^2),x)

[Out] int(1/((a + b*x^2)^(1/4)*(c + d*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[4]{a + bx^2} (c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(1/4)/(d*x**2+c)**2,x)

[Out] Integral(1/((a + b*x**2)**(1/4)*(c + d*x**2)**2), x)

$$3.336 \quad \int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)^2} dx$$

Optimal. Leaf size=292

$$\frac{\sqrt{a}\sqrt{b}\left(\frac{bx^2}{a}+1\right)^{3/4}\text{EllipticF}\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right),2\right)}{2c(a+bx^2)^{3/4}(bc-ad)} - \frac{dx\sqrt[4]{a+bx^2}}{2c(c+dx^2)(bc-ad)} + \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}(5bc-2ad)\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2}}{\sqrt[4]{a}}\right)\right)}{4cx(bc-ad)^2}$$

[Out] $-1/2*d*x*(b*x^2+a)^{(1/4)}/c/(-a*d+b*c)/(d*x^2+c)-1/2*(1+b*x^2/a)^{(3/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})))*\text{EllipticF}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*a^{(1/2)}*b^{(1/2)}/c/(-a*d+b*c)/(b*x^2+a)^{(3/4)}+1/4*a^{(1/4)}*(-2*a*d+5*b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/c/(-a*d+b*c)^2/x+1/4*a^{(1/4)}*(-2*a*d+5*b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/c/(-a*d+b*c)^2/x$

Rubi [A] time = 0.23, antiderivative size = 292, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {414, 530, 233, 231, 401, 108, 409, 1218}

$$\frac{dx\sqrt[4]{a+bx^2}}{2c(c+dx^2)(bc-ad)} - \frac{\sqrt{a}\sqrt{b}\left(\frac{bx^2}{a}+1\right)^{3/4}F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{2c(a+bx^2)^{3/4}(bc-ad)} + \frac{\sqrt[4]{a}\sqrt{-\frac{bx^2}{a}}(5bc-2ad)\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}};\sin^{-1}\left(\frac{\sqrt[4]{bx^2}}{\sqrt[4]{a}}\right)\right)}{4cx(bc-ad)^2}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(3/4)*(c + d*x^2)^2),x]

[Out] $-(d*x*(a+b*x^2)^{(1/4)})/(2*c*(b*c-a*d)*(c+d*x^2)) - (\text{Sqrt}[a]*\text{Sqrt}[b]*(1+(b*x^2)/a)^{(3/4)}*\text{EllipticF}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2,2])/(2*c*(b*c-a*d)*(a+b*x^2)^{(3/4)}) + (a^{(1/4)}*(5*b*c-2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c)+a*d])],\text{ArcSin}[(a+b*x^2)^{(1/4)}/a^{(1/4)}],-1])/(4*c*(b*c-a*d)^2*x) + (a^{(1/4)}*(5*b*c-2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c)+a*d]),\text{ArcSin}[(a+b*x^2)^{(1/4)}/a^{(1/4)}],-1])/(4*c*(b*c-a*d)^2*x$

Rule 108

Int[1/(((a_.) + (b_.)*(x_))*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(3/4)), x_Symbol] := Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 231

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Simp[(2*EllipticF[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(3/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 233

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(3/4)/(a + b*x^2)^(3/4), Int[1/(1 + (b*x^2)/a)^(3/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 401

Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 409

Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 530

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)^2} dx &= -\frac{dx\sqrt[4]{a+bx^2}}{2c(bc-ad)(c+dx^2)} + \frac{\int \frac{2bc-ad-\frac{1}{2}bdx^2}{(a+bx^2)^{3/4}(c+dx^2)} dx}{2c(bc-ad)} \\
&= -\frac{dx\sqrt[4]{a+bx^2}}{2c(bc-ad)(c+dx^2)} - \frac{b \int \frac{1}{(a+bx^2)^{3/4}} dx}{4c(bc-ad)} + \frac{(5bc-2ad) \int \frac{1}{(a+bx^2)^{3/4}(c+dx^2)} dx}{4c(bc-ad)} \\
&= -\frac{dx\sqrt[4]{a+bx^2}}{2c(bc-ad)(c+dx^2)} + \frac{\left((5bc-2ad)\sqrt{-\frac{bx^2}{a}} \right) \text{Subst}\left(\int \frac{1}{\sqrt{-\frac{bx}{a}}(a+bx)^{3/4}(c+dx)} dx, x, \dots \right)}{8c(bc-ad)x} \\
&= -\frac{dx\sqrt[4]{a+bx^2}}{2c(bc-ad)(c+dx^2)} - \frac{\sqrt{a}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{2c(bc-ad)(a+bx^2)^{3/4}} - \frac{(5bc-2ad)}{\dots} \\
&= -\frac{dx\sqrt[4]{a+bx^2}}{2c(bc-ad)(c+dx^2)} - \frac{\sqrt{a}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{2c(bc-ad)(a+bx^2)^{3/4}} + \frac{(5bc-2ad)}{\dots} \\
&= -\frac{dx\sqrt[4]{a+bx^2}}{2c(bc-ad)(c+dx^2)} - \frac{\sqrt{a}\sqrt{b}\left(1+\frac{bx^2}{a}\right)^{3/4} F\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{2c(bc-ad)(a+bx^2)^{3/4}} + \frac{\sqrt[4]{a}(5bc-2)}{\dots}
\end{aligned}$$

Mathematica [C] time = 0.32, size = 336, normalized size = 1.15

$$x \frac{\left(bdx^2\left(\frac{bx^2}{a}+1\right)^{3/4} F_1\left(\frac{3}{2};\frac{3}{4},1;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right) + c\left(36ac(2ad-2bc+bdx^2)F_1\left(\frac{1}{2};\frac{3}{4},1;\frac{3}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)-6dx^2(a+bx^2)\left(4adF_1\left(\frac{3}{2};\frac{3}{4},2;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)+3bcF_1\left(\frac{3}{2};\frac{7}{4},1,5/2,-((b*x^2)/a),-((d*x^2)/c)\right)\right)\right)}{ad-bc} + \frac{(c+dx^2)(bc-ad)\left(x^2\left(4adF_1\left(\frac{3}{2};\frac{3}{4},2;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)+3bcF_1\left(\frac{3}{2};\frac{7}{4},1,5/2,-((b*x^2)/a),-((d*x^2)/c)\right)\right)-6acF_1\left(\frac{1}{2};\frac{3}{4},1,3/2,-((b*x^2)/a),-((d*x^2)/c)\right)\right)}{12c^2(a+bx^2)^{3/4}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(3/4)*(c + d*x^2)^2), x]

[Out] (x*((b*d*x^2*(1 + (b*x^2)/a)^(3/4)*AppellF1[3/2, 3/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])/(-b*c) + a*d) + (c*(36*a*c*(-2*b*c + 2*a*d + b*d*x^2)*AppellF1[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] - 6*d*x^2*(a + b*x^2)*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/((b*c - a*d)*(c + d*x^2)*(-6*a*c*AppellF1[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^2*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/((12*c^2*(a + b*x^2)^(3/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(3/4)*(d*x^2 + c)^2), x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(3/4)/(d*x^2+c)^2,x)

[Out] int(1/(b*x^2+a)^(3/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(3/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(3/4)*(d*x^2 + c)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{3}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(3/4)*(c + d*x^2)^2),x)

[Out] int(1/((a + b*x^2)^(3/4)*(c + d*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{3}{4}}(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(3/4)/(d*x**2+c)**2,x)

[Out] Integral(1/((a + b*x**2)**(3/4)*(c + d*x**2)**2), x)

$$3.337 \quad \int \frac{1}{(a+bx^2)^{5/4}(c+dx^2)^2} dx$$

Optimal. Leaf size=314

$$\frac{dx}{2c\sqrt[4]{a+bx^2}(c+dx^2)(bc-ad)} + \frac{\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}(ad+4bc)E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{2\sqrt{a}c\sqrt[4]{a+bx^2}(bc-ad)^2} - \frac{\sqrt[4]{a}\sqrt{d}\sqrt{-\frac{bx^2}{a}}(7bc-2ad)\Pi\left(-\frac{\sqrt{b}x}{\sqrt{a}}\right)}{4cx(ad-bc)}$$

[Out] $-1/2*d*x/c/(-a*d+b*c)/(b*x^2+a)^{(1/4)}/(d*x^2+c)+1/2*(a*d+4*b*c)*(1+b*x^2/a)^{(1/4)*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})))/a^{(1/2)})^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*b^{(1/2)}/c/(-a*d+b*c)^2/(b*x^2+a)^{(1/4)}/a^{(1/2)}-1/4*a^{(1/4)*(-2*a*d+7*b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*d^{(1/2)*(-b*x^2/a)^{(1/2)}/c/(a*d-b*c)^{(5/2)}/x+1/4*a^{(1/4)*(-2*a*d+7*b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*d^{(1/2)*(-b*x^2/a)^{(1/2)}/c/(a*d-b*c)^{(5/2)}/x$

Rubi [A] time = 0.40, antiderivative size = 314, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {414, 527, 530, 229, 227, 196, 399, 490, 1218}

$$\frac{dx}{2c\sqrt[4]{a+bx^2}(c+dx^2)(bc-ad)} + \frac{\sqrt{b}\sqrt[4]{\frac{bx^2}{a}+1}(ad+4bc)E\left(\frac{1}{2}\tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\middle|2\right)}{2\sqrt{a}c\sqrt[4]{a+bx^2}(bc-ad)^2} - \frac{\sqrt[4]{a}\sqrt{d}\sqrt{-\frac{bx^2}{a}}(7bc-2ad)\Pi\left(-\frac{\sqrt{b}x}{\sqrt{a}}\right)}{4cx(ad-bc)}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(5/4)*(c + d*x^2)^2), x]

[Out] $-(d*x)/(2*c*(b*c - a*d)*(a + b*x^2)^{(1/4)*(c + d*x^2)} + (\text{Sqrt}[b]*(4*b*c + a*d)*(1 + (b*x^2)/a)^{(1/4)*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2]})/(2*\text{Sqrt}[a]*c*(b*c - a*d)^2*(a + b*x^2)^{(1/4)}) - (a^{(1/4)*\text{Sqrt}[d]*(7*b*c - 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d])], \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*(-(b*c) + a*d)^{(5/2)*x} + (a^{(1/4)*\text{Sqrt}[d]*(7*b*c - 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*(-(b*c) + a*d)^{(5/2)*x}$

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 227

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2*x)/(a + b*x^2)^(1/4), x] - Dist[a, Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 229

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a + b*x^2)^(1/4), Int[1/(1 + (b*x^2)/a)^(1/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 399

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dis
t[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x
^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0]
```

Rule 414

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c -
a*d), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d), Int[(a + b*x^n)^(p + 1)*(c +
d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x]
, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1]
&& !( !IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 490

```
Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] :=
With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s
/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/(
(r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c
- a*d, 0]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 530

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p
, n}, x]
```

Rule 1218

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*
Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a+bx^2)^{5/4}(c+dx^2)^2} dx &= -\frac{dx}{2c(bc-ad)\sqrt[4]{a+bx^2}(c+dx^2)} + \frac{\int \frac{2bc-ad-\frac{3}{2}bdx^2}{(a+bx^2)^{5/4}(c+dx^2)} dx}{2c(bc-ad)} \\
&= \frac{b(4bc+ad)x}{2ac(bc-ad)^2\sqrt[4]{a+bx^2}} - \frac{dx}{2c(bc-ad)\sqrt[4]{a+bx^2}(c+dx^2)} - \frac{\int \frac{\frac{1}{2}(2b^2c^2+4abcd-a^2d^2)+}{\sqrt[4]{a+bx^2}(c+dx^2)} dx}{ac(bc-ad)} \\
&= \frac{b(4bc+ad)x}{2ac(bc-ad)^2\sqrt[4]{a+bx^2}} - \frac{dx}{2c(bc-ad)\sqrt[4]{a+bx^2}(c+dx^2)} - \frac{(d(7bc-2ad)) \int \frac{dx}{\sqrt[4]{a+bx^2}}}{4c(bc-ad)} \\
&= \frac{b(4bc+ad)x}{2ac(bc-ad)^2\sqrt[4]{a+bx^2}} - \frac{dx}{2c(bc-ad)\sqrt[4]{a+bx^2}(c+dx^2)} - \frac{\left(d(7bc-2ad)\sqrt{-\frac{bx^2}{a}}\right)}{4c(bc-ad)^2x} \\
&= -\frac{dx}{2c(bc-ad)\sqrt[4]{a+bx^2}(c+dx^2)} + \frac{\left(\sqrt{d}(7bc-2ad)\sqrt{-\frac{bx^2}{a}}\right) \text{Subst} \left(\int \frac{1}{(\sqrt{-bc+ad}-\sqrt{bx^2})} dx \right)}{4c(bc-ad)^2x} \\
&= -\frac{dx}{2c(bc-ad)\sqrt[4]{a+bx^2}(c+dx^2)} + \frac{\sqrt{b}(4bc+ad)\sqrt[4]{1+\frac{bx^2}{a}} E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)\right)}{2\sqrt{a}c(bc-ad)^2\sqrt[4]{a+bx^2}}
\end{aligned}$$

Mathematica [C] time = 0.49, size = 380, normalized size = 1.21

$$x \frac{c \left(36ac(2a^2d^2+abd(dx^2-4c))+2b^2c(c+2dx^2) \right) F_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - 6x^2(a^2d^2+abd^2x^2+4b^2c(c+dx^2)) \left(4adF_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + bcF_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right)}{(c+dx^2) \left(6acF_1\left(\frac{1}{2}; \frac{1}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - x^2 \left(4adF_1\left(\frac{3}{2}; \frac{1}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + bcF_1\left(\frac{3}{2}; \frac{5}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) \right)}$$

$$12ac^2\sqrt[4]{a+bx^2}(bc-ad)^2$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(5/4)*(c + d*x^2)^2), x]

[Out] (x*(-(b*d*(4*b*c + a*d)*x^2*(1 + (b*x^2)/a)^(1/4)*AppellF1[3/2, 1/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)]) + (c*(36*a*c*(2*a^2*d^2 + a*b*d*(-4*c + d*x^2) + 2*b^2*c*(c + 2*d*x^2))*AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -((d*x^2)/c)] - 6*x^2*(a^2*d^2 + a*b*d^2*x^2 + 4*b^2*c*(c + d*x^2))*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)])))/((c + d*x^2)*(6*a*c*AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -((d*x^2)/c)] - x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)])))/((12*a*c^2*(b*c - a*d)^2*(a + b*x^2)^(1/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(5/4)*(d*x^2 + c)^2), x)

maple [F] time = 0.36, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(5/4)/(d*x^2+c)^2,x)

[Out] int(1/(b*x^2+a)^(5/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(5/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(5/4)*(d*x^2 + c)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{5}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(5/4)*(c + d*x^2)^2),x)

[Out] int(1/((a + b*x^2)^(5/4)*(c + d*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{5}{4}}(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(5/4)/(d*x**2+c)**2,x)

[Out] Integral(1/((a + b*x**2)**(5/4)*(c + d*x**2)**2), x)

$$3.338 \quad \int \frac{1}{(a+bx^2)^{7/4}(c+dx^2)^2} dx$$

Optimal. Leaf size=345

$$\frac{\sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} (3ad + 4bc) \operatorname{EllipticF}\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), 2\right)}{6\sqrt{a}c(a+bx^2)^{3/4}(bc-ad)^2} + \frac{bx(3ad+4bc)}{6ac(a+bx^2)^{3/4}(bc-ad)^2} - \frac{dx}{2c(a+bx^2)^{3/4}(c+dx^2)^2}$$

[Out] $\frac{1}{6} b^* (3 a^* d + 4 b^* c) x / a / c / (-a^* d + b^* c)^2 / (b^* x^2 + a)^{3/4} - 1/2 d^* x / c / (-a^* d + b^* c) / (b^* x^2 + a)^{3/4} / (d^* x^2 + c) + 1/6 (3 a^* d + 4 b^* c) (1 + b^* x^2 / a)^{3/4} (\cos(1/2 \arctan(x^* b^{1/2} / a^{1/2}))^2)^{1/2} / \cos(1/2 \arctan(x^* b^{1/2} / a^{1/2})) * \operatorname{EllipticF}(\sin(1/2 \arctan(x^* b^{1/2} / a^{1/2})), 2^{1/2}) * b^{1/2} / c / (-a^* d + b^* c)^2 / (b^* x^2 + a)^{3/4} / a^{1/2} - 1/4 a^{1/4} d^* (-2 a^* d + 9 b^* c) * \operatorname{EllipticPi}((b^* x^2 + a)^{1/4} / a^{1/4}, -a^{1/2} d^{1/2} / (a^* d - b^* c)^{1/2}, I) * (-b^* x^2 / a)^{1/2} / c / (-a^* d + b^* c)^3 / x - 1/4 a^{1/4} d^* (-2 a^* d + 9 b^* c) * \operatorname{EllipticPi}((b^* x^2 + a)^{1/4} / a^{1/4}, a^{1/2} d^{1/2} / (a^* d - b^* c)^{1/2}, I) * (-b^* x^2 / a)^{1/2} / c / (-a^* d + b^* c)^3 / x$

Rubi [A] time = 0.38, antiderivative size = 345, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {414, 527, 530, 233, 231, 401, 108, 409, 1218}

$$\frac{bx(3ad+4bc)}{6ac(a+bx^2)^{3/4}(bc-ad)^2} - \frac{dx}{2c(a+bx^2)^{3/4}(c+dx^2)^2} + \frac{\sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} (3ad+4bc) F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{6\sqrt{a}c(a+bx^2)^{3/4}(bc-ad)^2}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(7/4)*(c + d*x^2)^2), x]

[Out] $(b^*(4*b^*c + 3*a^*d)*x)/(6*a^*c*(b^*c - a^*d)^2*(a + b^*x^2)^{3/4}) - (d*x)/(2*c*(b^*c - a^*d)*(a + b^*x^2)^{3/4}*(c + d*x^2)) + (\operatorname{Sqrt}[b]*(4*b^*c + 3*a^*d)*(1 + (b^*x^2)/a)^{3/4}*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/2, 2])/(6*\operatorname{Sqrt}[a]*c*(b^*c - a^*d)^2*(a + b^*x^2)^{3/4}) - (a^{1/4}*d*(9*b^*c - 2*a^*d)*\operatorname{Sqrt}[-((b^*x^2)/a)]*\operatorname{EllipticPi}[-((\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b^*c) + a^*d])), \operatorname{ArcSin}[(a + b^*x^2)^{1/4}/a^{1/4}], -1])/(4*c*(b^*c - a^*d)^3*x) - (a^{1/4}*d*(9*b^*c - 2*a^*d)*\operatorname{Sqrt}[-((b^*x^2)/a)]*\operatorname{EllipticPi}[(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b^*c) + a^*d]), \operatorname{ArcSin}[(a + b^*x^2)^{1/4}/a^{1/4}], -1])/(4*c*(b^*c - a^*d)^3*x)$

Rule 108

Int[1/(((a_.) + (b_.)*(x_)) * Sqrt[(c_.) + (d_.)*(x_)]) * ((e_.) + (f_.)*(x_))^(3/4)), x_Symbol] :> Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 231

Int[((a_.) + (b_.)*(x_)^2)^(-3/4), x_Symbol] :> Simp[(2*EllipticF[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(3/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 233

Int[((a_.) + (b_.)*(x_)^2)^(-3/4), x_Symbol] :> Dist[(1 + (b*x^2)/a)^(3/4)/(a + b*x^2)^(3/4), Int[1/(1 + (b*x^2)/a)^(3/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 401

Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 409

Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 530

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a+bx^2)^{7/4}(c+dx^2)^2} dx &= -\frac{dx}{2c(bc-ad)(a+bx^2)^{3/4}(c+dx^2)} + \frac{\int \frac{2bc-ad-\frac{5}{2}bdx^2}{(a+bx^2)^{7/4}(c+dx^2)} dx}{2c(bc-ad)} \\
 &= \frac{b(4bc+3ad)x}{6ac(bc-ad)^2(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{3/4}(c+dx^2)} - \frac{\int \frac{\frac{1}{2}(-2b^2c^2+12abcd)}{(a+b)}}{3ac} \\
 &= \frac{b(4bc+3ad)x}{6ac(bc-ad)^2(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{3/4}(c+dx^2)} - \frac{(d(9bc-2ad)) \int}{4c(b)} \\
 &= \frac{b(4bc+3ad)x}{6ac(bc-ad)^2(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{3/4}(c+dx^2)} - \frac{(d(9bc-2ad)\sqrt{}}{4c(b)} \\
 &= \frac{b(4bc+3ad)x}{6ac(bc-ad)^2(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{3/4}(c+dx^2)} + \frac{\sqrt{b}(4bc+3ad)}{6\sqrt{a}c} \\
 &= \frac{b(4bc+3ad)x}{6ac(bc-ad)^2(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{3/4}(c+dx^2)} + \frac{\sqrt{b}(4bc+3ad)}{6\sqrt{a}c} \\
 &= \frac{b(4bc+3ad)x}{6ac(bc-ad)^2(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{3/4}(c+dx^2)} + \frac{\sqrt{b}(4bc+3ad)}{6\sqrt{a}c}
 \end{aligned}$$

Mathematica [C] time = 0.51, size = 387, normalized size = 1.12

$$x \frac{\left(c \left(36ac(6a^2d^2+3abd(dx^2-4c)+2b^2c(3c+2dx^2)) F_1\left(\frac{1}{2}; \frac{3}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - 6x^2(3a^2d^2+3abd^2x^2+4b^2c(c+dx^2)) \left(4adF_1\left(\frac{3}{2}; \frac{3}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + 3bcF_1\left(\frac{3}{2}; \frac{3}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) \right)}{(c+dx^2) \left(6acF_1\left(\frac{1}{2}; \frac{3}{4}, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - x^2 \left(4adF_1\left(\frac{3}{2}; \frac{3}{4}, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + 3bcF_1\left(\frac{3}{2}; \frac{3}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) \right)}{36ac^2(a+bx^2)^{3/4}(bc-ad)^2}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(7/4)*(c + d*x^2)^2), x]

[Out] (x*(b*d*(4*b*c + 3*a*d)*x^2*(1 + (b*x^2)/a)^(3/4)*AppellF1[3/2, 3/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + (c*(36*a*c*(6*a^2*d^2 + 3*a*b*d*(-4*c + d*x^2) + 2*b^2*c*(3*c + 2*d*x^2))*AppellF1[1/2, 3/4, 1, 3/2, -(b*x^2)/a, -((d*x^2)/c)] - 6*x^2*(3*a^2*d^2 + 3*a*b*d^2*x^2 + 4*b^2*c*(c + d*x^2))*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)])))/((c + d*x^2)*(6*a*c*AppellF1[1/2, 3/4, 1, 3/2, -(b*x^2)/a, -((d*x^2)/c)] - x^2*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -(b*x^2)/a, -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -(b*x^2)/a, -((d*x^2)/c)])))/((36*a*c^2*(b*c - a*d)^2*(a + b*x^2)^(3/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{7}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(7/4)*(d*x^2 + c)^2), x)

maple [F] time = 0.36, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{7}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(7/4)/(d*x^2+c)^2,x)

[Out] int(1/(b*x^2+a)^(7/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{7}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(7/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(7/4)*(d*x^2 + c)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{7}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(7/4)*(c + d*x^2)^2),x)

[Out] int(1/((a + b*x^2)^(7/4)*(c + d*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{7}{4}}(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(7/4)/(d*x**2+c)**2,x)

[Out] Integral(1/((a + b*x**2)**(7/4)*(c + d*x**2)**2), x)

$$3.339 \quad \int \frac{1}{(a+bx^2)^{9/4}(c+dx^2)^2} dx$$

Optimal. Leaf size=371

$$\frac{\sqrt{b} \sqrt[4]{\frac{bx^2}{a} + 1} (-5a^2d^2 - 52abcd + 12b^2c^2) E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| 2\right)}{10a^{3/2}c\sqrt[4]{a+bx^2}(bc-ad)^3} - \frac{\sqrt[4]{a}d^{3/2}\sqrt{-\frac{bx^2}{a}}(11bc-2ad)\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\right)}{4cx(ad-bc)^{7/2}}$$

[Out] $\frac{1}{10}b*(5*a*d+4*b*c)*x/a/c/(-a*d+b*c)^2/(b*x^2+a)^{(5/4)}-1/2*d*x/c/(-a*d+b*c)/(b*x^2+a)^{(5/4)}/(d*x^2+c)+1/10*(-5*a^2*d^2-52*a*b*c*d+12*b^2*c^2)*(1+b*x^2/a)^{(1/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\text{EllipticE}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})),2^{(1/2)})*b^{(1/2)}/a^{(3/2)}/c/(-a*d+b*c)^3/(b*x^2+a)^{(1/4)}-1/4*a^{(1/4)}*d^{(3/2)}*(-2*a*d+11*b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},-a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/c/(a*d-b*c)^{(7/2)}/x+1/4*a^{(1/4)}*d^{(3/2)}*(-2*a*d+11*b*c)*\text{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)},a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)},I)*(-b*x^2/a)^{(1/2)}/c/(a*d-b*c)^{(7/2)}/x$

Rubi [A] time = 0.55, antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {414, 527, 530, 229, 227, 196, 399, 490, 1218}

$$\frac{\sqrt{b} \sqrt[4]{\frac{bx^2}{a} + 1} (-5a^2d^2 - 52abcd + 12b^2c^2) E\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right) \middle| 2\right)}{10a^{3/2}c\sqrt[4]{a+bx^2}(bc-ad)^3} - \frac{\sqrt[4]{a}d^{3/2}\sqrt{-\frac{bx^2}{a}}(11bc-2ad)\Pi\left(-\frac{\sqrt{a}\sqrt{d}}{\sqrt{ad-bc}}; \sin^{-1}\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)\right)}{4cx(ad-bc)^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(9/4)*(c + d*x^2)^2), x]

[Out] $\frac{b*(4*b*c + 5*a*d)*x}{(10*a*c*(b*c - a*d)^2*(a + b*x^2)^{(5/4)} - (d*x)/(2*c*(b*c - a*d)*(a + b*x^2)^{(5/4)*(c + d*x^2))} + (\text{Sqrt}[b]*(12*b^2*c^2 - 52*a*b*c*d - 5*a^2*d^2)*(1 + (b*x^2)/a)^{(1/4)}*\text{EllipticE}[\text{ArcTan}[(\text{Sqrt}[b]*x)/\text{Sqrt}[a]]/2, 2])/(10*a^{(3/2)}*c*(b*c - a*d)^3*(a + b*x^2)^{(1/4)} - (a^{(1/4)}*d^{(3/2)}*(11*b*c - 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[-((\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*(-(b*c) + a*d)^{(7/2)*x} + (a^{(1/4)}*d^{(3/2)}*(11*b*c - 2*a*d)*\text{Sqrt}[-((b*x^2)/a)]*\text{EllipticPi}[(\text{Sqrt}[a]*\text{Sqrt}[d])/(\text{Sqrt}[-(b*c) + a*d]), \text{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*(-(b*c) + a*d)^{(7/2)*x}}$

Rule 196

Int[((a_) + (b_.)*(x_)^2)^(-5/4), x_Symbol] := Simp[(2*EllipticE[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^(5/4)*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 227

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2*x)/(a + b*x^2)^(1/4), x] - Dist[a, Int[1/(a + b*x^2)^(5/4), x], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 229

Int[((a_) + (b_.)*(x_)^2)^(-1/4), x_Symbol] := Dist[(1 + (b*x^2)/a)^(1/4)/(a + b*x^2)^(1/4), Int[1/(1 + (b*x^2)/a)^(1/4), x], x] /; FreeQ[{a, b}, x] && PosQ[a]

Rule 399

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dis
t[(2*Sqrt[-((b*x^2)/a)])/x, Subst[Int[x^2/(Sqrt[1 - x^4/a]*(b*c - a*d + d*x
^4)), x], x, (a + b*x^2)^(1/4)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0]
```

Rule 414

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c -
a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c +
d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x]
, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1]
&& !( !IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c,
d, n, p, q, x]
```

Rule 490

```
Int[(x_)^2/(((a_) + (b_.)*(x_)^4)*Sqrt[(c_) + (d_.)*(x_)^4]), x_Symbol] :=
With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s
/(2*b), Int[1/((r + s*x^2)*Sqrt[c + d*x^4]), x], x] - Dist[s/(2*b), Int[1/(
(r - s*x^2)*Sqrt[c + d*x^4]), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c
- a*d, 0]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c +
d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p
+ 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c
- a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ
[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 530

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_))/((c_) + (d_.)*
(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e -
c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p
, n}, x]
```

Rule 1218

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*
Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a+bx^2)^{9/4}(c+dx^2)^2} dx &= -\frac{dx}{2c(bc-ad)(a+bx^2)^{5/4}(c+dx^2)} + \frac{\int \frac{2bc-ad-\frac{7}{2}bdx^2}{(a+bx^2)^{9/4}(c+dx^2)} dx}{2c(bc-ad)} \\
&= \frac{b(4bc+5ad)x}{10ac(bc-ad)^2(a+bx^2)^{5/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{5/4}(c+dx^2)} - \frac{\int \frac{\frac{1}{2}(-6b^2c^2+20abcd-5a^2d^2)}{(a+bx^2)^{9/4}(c+dx^2)} dx}{5ac(bc-ad)^2} \\
&= \frac{b(4bc+5ad)x}{10ac(bc-ad)^2(a+bx^2)^{5/4}} + \frac{b(12b^2c^2-52abcd-5a^2d^2)x}{10a^2c(bc-ad)^3\sqrt[4]{a+bx^2}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{5/4}(c+dx^2)} \\
&= \frac{b(4bc+5ad)x}{10ac(bc-ad)^2(a+bx^2)^{5/4}} + \frac{b(12b^2c^2-52abcd-5a^2d^2)x}{10a^2c(bc-ad)^3\sqrt[4]{a+bx^2}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{5/4}(c+dx^2)} \\
&= \frac{b(4bc+5ad)x}{10ac(bc-ad)^2(a+bx^2)^{5/4}} + \frac{b(12b^2c^2-52abcd-5a^2d^2)x}{10a^2c(bc-ad)^3\sqrt[4]{a+bx^2}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{5/4}(c+dx^2)} \\
&= \frac{b(4bc+5ad)x}{10ac(bc-ad)^2(a+bx^2)^{5/4}} + \frac{b(12b^2c^2-52abcd-5a^2d^2)x}{10a^2c(bc-ad)^3\sqrt[4]{a+bx^2}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{5/4}(c+dx^2)} \\
&= \frac{b(4bc+5ad)x}{10ac(bc-ad)^2(a+bx^2)^{5/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{5/4}(c+dx^2)} - \frac{d^{3/2}(11bc-2ad)}{5ac(bc-ad)^2} \\
&= \frac{b(4bc+5ad)x}{10ac(bc-ad)^2(a+bx^2)^{5/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{5/4}(c+dx^2)} + \frac{\sqrt{b}(12b^2c^2-52abcd-5a^2d^2)}{10a^2c(bc-ad)^3}
\end{aligned}$$

Mathematica [C] time = 1.05, size = 536, normalized size = 1.44

$$bdx^3 \sqrt[4]{\frac{bx^2}{a} + 1} (5a^2d^2 + 52abcd - 12b^2c^2) F_1\left(\frac{3}{2}; \frac{1}{4}, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + \frac{6c(x^3(5a^4d^3 + 10a^3bd^3x^2 + a^2b^2d(56c^2 + 56cdx^2 + 5d^2x^4) + 4a^2d^3 + 15a^3b^2d^2(-2c + dx^2) - 6b^4c^2x^2(c + 2dx^2) + a^2b^2d(30c^2 + 26cdx^2 + 5d^2x^4) + 2a^2b^3c(-5c^2 + 5cdx^2 + 26d^2x^4)) * AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -(d*x^2)/c] + x^3(5a^4d^3 + 10a^3b^2d^3x^2 - 12b^4c^2x^2(c + dx^2) + a^2b^2d(56c^2 + 56cdx^2 + 5d^2x^4) + 4a^2b^3c(-4c^2 + 9cdx^2 + 13d^2x^4)) * (4ad * AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -(d*x^2)/c] + bc * AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])}{(a + bx^2)(c + dx^2)(6ac * AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -(d*x^2)/c] - x^2(4ad * AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -(d*x^2)/c] + bc * AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])}}{(60a^2c^2(bc - ad)^3(a + bx^2)^{1/4})}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(9/4)*(c + d*x^2)^2), x]

[Out] (b*d*(-12*b^2*c^2 + 52*a*b*c*d + 5*a^2*d^2)*x^3*(1 + (b*x^2)/a)^(1/4)*AppellF1[3/2, 1/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c] + (6*c*(-6*a*c*x*(10*a^4*d^3 + 15*a^3*b*d^2*(-2*c + d*x^2) - 6*b^4*c^2*x^2*(c + 2*d*x^2) + a^2*b^2*d*(30*c^2 + 26*c*d*x^2 + 5*d^2*x^4) + 2*a*b^3*c*(-5*c^2 + 5*c*d*x^2 + 26*d^2*x^4))*AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -(d*x^2)/c] + x^3*(5*a^4*d^3 + 10*a^3*b*d^3*x^2 - 12*b^4*c^2*x^2*(c + d*x^2) + a^2*b^2*d*(56*c^2 + 56*c*d*x^2 + 5*d^2*x^4) + 4*a*b^3*c*(-4*c^2 + 9*c*d*x^2 + 13*d^2*x^4))*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -(d*x^2)/c] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])))/(a + b*x^2)*(c + d*x^2)*(6*a*c*AppellF1[1/2, 1/4, 1, 3/2, -(b*x^2)/a, -(d*x^2)/c] - x^2*(4*a*d*AppellF1[3/2, 1/4, 2, 5/2, -(b*x^2)/a, -(d*x^2)/c] + b*c*AppellF1[3/2, 5/4, 1, 5/2, -(b*x^2)/a, -(d*x^2)/c])))/(60*a^2*c^2*(b*c - a*d)^3*(a + b*x^2)^(1/4))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(9/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{9}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(9/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(9/4)*(d*x^2 + c)^2), x)

maple [F] time = 0.37, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{9}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(9/4)/(d*x^2+c)^2,x)

[Out] int(1/(b*x^2+a)^(9/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{9}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(9/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(9/4)*(d*x^2 + c)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{9/4}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(9/4)*(c + d*x^2)^2),x)

[Out] int(1/((a + b*x^2)^(9/4)*(c + d*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^{\frac{9}{4}}(c + dx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(9/4)/(d*x**2+c)**2,x)

[Out] Integral(1/((a + b*x**2)**(9/4)*(c + d*x**2)**2), x)

$$3.340 \quad \int \frac{1}{(a+bx^2)^{11/4} (c+dx^2)^2} dx$$

Optimal. Leaf size=419

$$\frac{\sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} (-21a^2d^2 - 76abcd + 20b^2c^2) \operatorname{EllipticF}\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right), 2\right) + bx(-21a^2d^2 - 76abcd + 20b^2c^2) \sqrt[4]{a} d^2 \sqrt{-}}{42a^{3/2}c(a+bx^2)^{3/4}(bc-ad)^3 + 42a^2c(a+bx^2)^{3/4}(bc-ad)^3}$$

[Out] $1/14*b*(7*a*d+4*b*c)*x/a/c/(-a*d+b*c)^2/(b*x^2+a)^{(7/4)}+1/42*b*(-21*a^2*d^2-76*a*b*c*d+20*b^2*c^2)*x/a^2/c/(-a*d+b*c)^3/(b*x^2+a)^{(3/4)}-1/2*d*x/c/(-a*d+b*c)/(b*x^2+a)^{(7/4)}/(d*x^2+c)+1/42*(-21*a^2*d^2-76*a*b*c*d+20*b^2*c^2)*(1+b*x^2/a)^{(3/4)}*(\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))^2)^{(1/2)}/\cos(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)}))*\operatorname{EllipticF}(\sin(1/2*\arctan(x*b^{(1/2)}/a^{(1/2)})), 2^{(1/2)})*b^{(1/2)}/a^{(3/2)}/c/(-a*d+b*c)^3/(b*x^2+a)^{(3/4)}+1/4*a^{(1/4)}*d^2*(-2*a*d+13*b*c)*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, -a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/c/(-a*d+b*c)^4/x+1/4*a^{(1/4)}*d^2*(-2*a*d+13*b*c)*\operatorname{EllipticPi}((b*x^2+a)^{(1/4)}/a^{(1/4)}, a^{(1/2)}*d^{(1/2)}/(a*d-b*c)^{(1/2)}, I)*(-b*x^2/a)^{(1/2)}/c/(-a*d+b*c)^4/x$

Rubi [A] time = 0.48, antiderivative size = 419, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {414, 527, 530, 233, 231, 401, 108, 409, 1218}

$$\frac{bx(-21a^2d^2 - 76abcd + 20b^2c^2)}{42a^2c(a+bx^2)^{3/4}(bc-ad)^3} + \frac{\sqrt{b} \left(\frac{bx^2}{a} + 1\right)^{3/4} (-21a^2d^2 - 76abcd + 20b^2c^2) F\left(\frac{1}{2} \tan^{-1}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right) \middle| 2\right)}{42a^{3/2}c(a+bx^2)^{3/4}(bc-ad)^3} + \frac{\sqrt[4]{a} d^2 \sqrt{-}}{42a^{3/2}c(a+bx^2)^{3/4}(bc-ad)^3}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^(11/4)*(c + d*x^2)^2), x]

[Out] $(b*(4*b*c + 7*a*d)*x)/(14*a*c*(b*c - a*d)^2*(a + b*x^2)^{(7/4)} + (b*(20*b^2*c^2 - 76*a*b*c*d - 21*a^2*d^2)*x)/(42*a^2*c*(b*c - a*d)^3*(a + b*x^2)^{(3/4)}) - (d*x)/(2*c*(b*c - a*d)*(a + b*x^2)^{(7/4)}*(c + d*x^2)) + (\operatorname{Sqrt}[b]*(20*b^2*c^2 - 76*a*b*c*d - 21*a^2*d^2)*(1 + (b*x^2)/a)^{(3/4)}*\operatorname{EllipticF}[\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*x)/\operatorname{Sqrt}[a]]/2, 2])/(42*a^{(3/2)}*c*(b*c - a*d)^3*(a + b*x^2)^{(3/4)}) + (a^{(1/4)}*d^2*(13*b*c - 2*a*d)*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[-((\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b*c) + a*d]), \operatorname{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*(b*c - a*d)^4*x) + (a^{(1/4)}*d^2*(13*b*c - 2*a*d)*\operatorname{Sqrt}[-((b*x^2)/a)]*\operatorname{EllipticPi}[(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[d])/(\operatorname{Sqrt}[-(b*c) + a*d]), \operatorname{ArcSin}[(a + b*x^2)^{(1/4)}/a^{(1/4)}], -1])/(4*c*(b*c - a*d)^4*x)$

Rule 108

Int[1/(((a_.) + (b_.)*(x_))*Sqrt[(c_.) + (d_.)*(x_)])*((e_.) + (f_.)*(x_))^(3/4)), x_Symbol] :> Dist[-4, Subst[Int[1/((b*e - a*f - b*x^4)*Sqrt[c - (d*e)/f + (d*x^4)/f]), x], x, (e + f*x)^(1/4)], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[-(f/(d*e - c*f)), 0]

Rule 231

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] :> Simp[(2*EllipticF[(1*ArcTan[Rt[b/a, 2]*x])/2, 2])/(a^{(3/4)}*Rt[b/a, 2]), x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b/a]

Rule 233

Int[((a_) + (b_.)*(x_)^2)^(-3/4), x_Symbol] :> Dist[(1 + (b*x^2)/a)^(3/4)/(a + b*x^2)^(3/4), Int[1/(1 + (b*x^2)/a)^(3/4), x], x] /; FreeQ[{a, b}, x] &

& PosQ[a]

Rule 401

Int[1/(((a_) + (b_.)*(x_)^2)^(3/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Dist[Sqrt[-((b*x^2)/a)]/(2*x), Subst[Int[1/(Sqrt[-((b*x)/a)]*(a + b*x)^(3/4)*(c + d*x)), x], x, x^2], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 409

Int[1/(Sqrt[(a_) + (b_.)*(x_)^4]*((c_) + (d_.)*(x_)^4)), x_Symbol] := Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 - Rt[-(d/c), 2]*x^2)), x], x] + Dist[1/(2*c), Int[1/(Sqrt[a + b*x^4]*(1 + Rt[-(d/c), 2]*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 414

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := -Simp[(b*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q + 1))/(a*n*(b*c - a*d)*(p + 1)), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 530

Int[(((a_) + (b_.)*(x_)^(n_))^(p_)*((e_) + (f_.)*(x_)^(n_)))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[f/d, Int[(a + b*x^n)^p, x], x] + Dist[(d*e - c*f)/d, Int[(a + b*x^n)^p/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, p, n}, x]

Rule 1218

Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[-(c/a), 4]}, Simp[(1*EllipticPi[-(e/(d*q^2)), ArcSin[q*x], -1])/(d*Sqrt[a]*q), x]] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && GtQ[a, 0]

Rubi steps

$$\int \frac{1}{(a+bx^2)^{11/4}(c+dx^2)^2} dx = -\frac{dx}{2c(bc-ad)(a+bx^2)^{7/4}(c+dx^2)} + \frac{\int \frac{2bc-ad-\frac{9}{2}bdx^2}{(a+bx^2)^{11/4}(c+dx^2)} dx}{2c(bc-ad)}$$

$$= \frac{b(4bc+7ad)x}{14ac(bc-ad)^2(a+bx^2)^{7/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{7/4}(c+dx^2)} - \int \frac{\frac{1}{2}(-10b^2c^2+28b^2cd-10c^2d^2)}{(a+bx^2)^{11/4}(c+dx^2)} dx$$

$$= \frac{b(4bc+7ad)x}{14ac(bc-ad)^2(a+bx^2)^{7/4}} + \frac{b(20b^2c^2-76abcd-21a^2d^2)x}{42a^2c(bc-ad)^3(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{7/4}(c+dx^2)}$$

$$= \frac{b(4bc+7ad)x}{14ac(bc-ad)^2(a+bx^2)^{7/4}} + \frac{b(20b^2c^2-76abcd-21a^2d^2)x}{42a^2c(bc-ad)^3(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{7/4}(c+dx^2)}$$

$$= \frac{b(4bc+7ad)x}{14ac(bc-ad)^2(a+bx^2)^{7/4}} + \frac{b(20b^2c^2-76abcd-21a^2d^2)x}{42a^2c(bc-ad)^3(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{7/4}(c+dx^2)}$$

$$= \frac{b(4bc+7ad)x}{14ac(bc-ad)^2(a+bx^2)^{7/4}} + \frac{b(20b^2c^2-76abcd-21a^2d^2)x}{42a^2c(bc-ad)^3(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{7/4}(c+dx^2)}$$

$$= \frac{b(4bc+7ad)x}{14ac(bc-ad)^2(a+bx^2)^{7/4}} + \frac{b(20b^2c^2-76abcd-21a^2d^2)x}{42a^2c(bc-ad)^3(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{7/4}(c+dx^2)}$$

$$= \frac{b(4bc+7ad)x}{14ac(bc-ad)^2(a+bx^2)^{7/4}} + \frac{b(20b^2c^2-76abcd-21a^2d^2)x}{42a^2c(bc-ad)^3(a+bx^2)^{3/4}} - \frac{dx}{2c(bc-ad)(a+bx^2)^{7/4}(c+dx^2)}$$

Mathematica [C] time = 0.92, size = 550, normalized size = 1.31

$$\frac{bdx^3\left(\frac{bx^2}{a}+1\right)^{3/4}\left(21a^2d^2+76abcd-20b^2c^2\right)F_1\left(\frac{3}{2};\frac{3}{4},1;\frac{5}{2};-\frac{bx^2}{a},-\frac{dx^2}{c}\right)}{(ad-bc)^3} + \frac{6c\left(x^3(21a^4d^3+42a^3bd^3x^2+a^2b^2d(88c^2+88cdx^2+21d^2x^4))+4ab^3c(-8c^2+11cdx^2+\right)}{(ad-bc)^3}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/((a + b*x^2)^(11/4)*(c + d*x^2)^2), x]

[Out] ((b*d*(-20*b^2*c^2 + 76*a*b*c*d + 21*a^2*d^2)*x^3*(1 + (b*x^2)/a)^(3/4)*AppellF1[3/2, 3/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)]/(-(b*c) + a*d)^3 + (6*c*(-6*a*c*x*(42*a^4*d^3 + 63*a^3*b*d^2*(-2*c + d*x^2) - 10*b^4*c^2*x^2*(3*c + 2*d*x^2) + a^2*b^2*d*(126*c^2 - 38*c*d*x^2 + 21*d^2*x^4) + 2*a*b^3*c*(-21*c^2 + 41*c*d*x^2 + 38*d^2*x^4))*AppellF1[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + x^3*(21*a^4*d^3 + 42*a^3*b*d^3*x^2 - 20*b^4*c^2*x^2*(c + d*x^2) + 4*a*b^3*c*(-8*c^2 + 11*c*d*x^2 + 19*d^2*x^4) + a^2*b^2*d*(88*c^2 + 8

$$\frac{8*c*d*x^2 + 21*d^2*x^4)*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])}{((b*c - a*d)^3*(a + b*x^2)*(c + d*x^2)*(6*a*c*AppellF1[1/2, 3/4, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)] - x^2*(4*a*d*AppellF1[3/2, 3/4, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] + 3*b*c*AppellF1[3/2, 7/4, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c)])))/(252*a^2*c^2*(a + b*x^2)^(3/4))}$$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(11/4)/(d*x^2+c)^2,x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{11}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(11/4)/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate(1/((b*x^2 + a)^(11/4)*(d*x^2 + c)^2), x)

maple [F] time = 0.37, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{11}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^(11/4)/(d*x^2+c)^2,x)

[Out] int(1/(b*x^2+a)^(11/4)/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^{\frac{11}{4}}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^(11/4)/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^(11/4)*(d*x^2 + c)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^{11/4}(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^(11/4)*(c + d*x^2)^2),x)

[Out] int(1/((a + b*x^2)^(11/4)*(c + d*x^2)^2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**(11/4)/(d*x**2+c)**2,x)

[Out] Timed out

3.341 $\int (a + bx^2)^p (c + dx^2)^q dx$

Optimal. Leaf size=79

$$x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} (c + dx^2)^q \left(\frac{dx^2}{c} + 1\right)^{-q} F_1\left(\frac{1}{2}; -p, -q; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)$$

[Out] $x*(b*x^2+a)^p*(d*x^2+c)^q*AppellF1(1/2, -p, -q, 3/2, -b*x^2/a, -d*x^2/c)/((1+b*x^2/a)^p)/((1+d*x^2/c)^q)$

Rubi [A] time = 0.04, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {430, 429}

$$x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} (c + dx^2)^q \left(\frac{dx^2}{c} + 1\right)^{-q} F_1\left(\frac{1}{2}; -p, -q; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^p*(c + d*x^2)^q,x]

[Out] $(x*(a + b*x^2)^p*(c + d*x^2)^q*AppellF1[1/2, -p, -q, 3/2, -((b*x^2)/a), -((d*x^2)/c)])/((1 + (b*x^2)/a)^p*(1 + (d*x^2)/c)^q)$

Rule 429

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 :> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, -((b*x^n)/a), -((d*x^n)/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 430

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 :> Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])

Rubi steps

$$\begin{aligned} \int (a + bx^2)^p (c + dx^2)^q dx &= \left((a + bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} \right) \int \left(1 + \frac{bx^2}{a}\right)^p (c + dx^2)^q dx \\ &= \left((a + bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} (c + dx^2)^q \left(1 + \frac{dx^2}{c}\right)^{-q} \right) \int \left(1 + \frac{bx^2}{a}\right)^p \left(1 + \frac{dx^2}{c}\right)^q dx \\ &= x(a + bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} (c + dx^2)^q \left(1 + \frac{dx^2}{c}\right)^{-q} F_1\left(\frac{1}{2}; -p, -q; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \end{aligned}$$

Mathematica [B] time = 0.23, size = 172, normalized size = 2.18

$$\frac{3acx(a + bx^2)^p (c + dx^2)^q F_1\left(\frac{1}{2}; -p, -q; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{2x^2 \left(bcpF_1\left(\frac{3}{2}; 1 - p, -q; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) + adqF_1\left(\frac{3}{2}; -p, 1 - q; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) + 3acF_1\left(\frac{1}{2}; -p, -q; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^p*(c + d*x^2)^q,x]

[Out] (3*a*c*x*(a + b*x^2)^p*(c + d*x^2)^q*AppellF1[1/2, -p, -q, 3/2, -((b*x^2)/a), -((d*x^2)/c)]/(3*a*c*AppellF1[1/2, -p, -q, 3/2, -((b*x^2)/a), -((d*x^2)/c)] + 2*x^2*(b*c*p*AppellF1[3/2, 1 - p, -q, 5/2, -((b*x^2)/a), -((d*x^2)/c)]) + a*d*q*AppellF1[3/2, -p, 1 - q, 5/2, -((b*x^2)/a), -((d*x^2)/c)])

fricas [F] time = 1.56, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(bx^2 + a\right)^p\left(dx^2 + c\right)^q, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c)^q,x, algorithm="fricas")

[Out] integral((b*x^2 + a)^p*(d*x^2 + c)^q, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^p (dx^2 + c)^q dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c)^q,x, algorithm="giac")

[Out] integrate((b*x^2 + a)^p*(d*x^2 + c)^q, x)

maple [F] time = 0.51, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^p (dx^2 + c)^q dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^p*(d*x^2+c)^q,x)

[Out] int((b*x^2+a)^p*(d*x^2+c)^q,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^p (dx^2 + c)^q dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c)^q,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^p*(d*x^2 + c)^q, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (bx^2 + a)^p (dx^2 + c)^q dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^p*(c + d*x^2)^q,x)

[Out] int((a + b*x^2)^p*(c + d*x^2)^q, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**p*(d*x**2+c)**q,x)

[Out] Timed out

3.342 $\int (a + bx^2)^p (c + dx^2)^3 dx$

Optimal. Leaf size=296

$$\frac{dx (a + bx^2)^{p+1} (15a^2d^2 - 8abcd(p+6) + b^2c^2(4p^2 + 28p + 57))}{b^3(2p+3)(2p+5)(2p+7)} x (a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} (15a^3d^3 - 9a^2bcd^2)$$

[Out] $d*(15*a^2*d^2-8*a*b*c*d*(6+p)+b^2*c^2*(4*p^2+28*p+57))*x*(b*x^2+a)^(1+p)/b^3/(8*p^3+60*p^2+142*p+105)-d*(5*a*d-b*c*(11+2*p))*x*(b*x^2+a)^(1+p)*(d*x^2+c)/b^2/(4*p^2+24*p+35)+d*x*(b*x^2+a)^(1+p)*(d*x^2+c)^2/b/(7+2*p)-(15*a^3*d^3-9*a^2*b*c*d^2*(7+2*p)+3*a*b^2*c^2*d*(4*p^2+24*p+35)-b^3*c^3*(8*p^3+60*p^2+142*p+105))*x*(b*x^2+a)^p*hypergeom([1/2, -p], [3/2], -b*x^2/a)/b^3/(8*p^3+60*p^2+142*p+105)/((1+b*x^2/a)^p)$

Rubi [A] time = 0.28, antiderivative size = 296, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {416, 528, 388, 246, 245}

$$\frac{x (a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} (-9a^2bcd^2(2p+7) + 15a^3d^3 + 3ab^2c^2d(4p^2 + 24p + 35) - b^3c^3(8p^3 + 60p^2 + 142p + 105))}{b^3(2p+3)(2p+5)(2p+7)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^p*(c + d*x^2)^3,x]

[Out] $(d*(15*a^2*d^2 - 8*a*b*c*d*(6 + p) + b^2*c^2*(57 + 28*p + 4*p^2))*x*(a + b*x^2)^(1 + p))/(b^3*c*(3 + 2*p)*(5 + 2*p)*(7 + 2*p)) - (d*(5*a*d - b*c*(11 + 2*p))*x*(a + b*x^2)^(1 + p)*(c + d*x^2))/(b^2*(5 + 2*p)*(7 + 2*p)) + (d*x*(a + b*x^2)^(1 + p)*(c + d*x^2)^2)/(b*(7 + 2*p)) - ((15*a^3*d^3 - 9*a^2*b*c*d^2*(7 + 2*p) + 3*a*b^2*c^2*d*(35 + 24*p + 4*p^2) - b^3*c^3*(105 + 142*p + 60*p^2 + 8*p^3))*x*(a + b*x^2)^p*Hypergeometric2F1[1/2, -p, 3/2, -(b*x^2)/a])/b^3*c*(3 + 2*p)*(5 + 2*p)*(7 + 2*p)*(1 + (b*x^2)/a)^p$

Rule 245

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, -(b*x^n)/a], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])

Rule 246

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q)+1)),

```
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a
, b, c, d, n, p, q, x]
```

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (
f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/
(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rubi steps

$$\begin{aligned} \int (a + bx^2)^p (c + dx^2)^3 dx &= \frac{dx (a + bx^2)^{1+p} (c + dx^2)^2}{b(7 + 2p)} + \frac{\int (a + bx^2)^p (c + dx^2) (-c(ad - bc(7 + 2p)) - d(5ad - bc(11 + 2p))) dx}{b(7 + 2p)} \\ &= -\frac{d(5ad - bc(11 + 2p))x (a + bx^2)^{1+p} (c + dx^2)}{b^2(5 + 2p)(7 + 2p)} + \frac{dx (a + bx^2)^{1+p} (c + dx^2)^2}{b(7 + 2p)} + \frac{\int (a + bx^2)^p (c + dx^2) (-c(ad - bc(7 + 2p)) - d(5ad - bc(11 + 2p))) dx}{b(7 + 2p)} \\ &= \frac{d(15a^2d^2 - 8abcd(6 + p) + b^2c^2(57 + 28p + 4p^2))x (a + bx^2)^{1+p}}{b^3(3 + 2p)(5 + 2p)(7 + 2p)} - \frac{d(5ad - bc(11 + 2p))x (a + bx^2)^{1+p}}{b^2(5 + 2p)(7 + 2p)} \\ &= \frac{d(15a^2d^2 - 8abcd(6 + p) + b^2c^2(57 + 28p + 4p^2))x (a + bx^2)^{1+p}}{b^3(3 + 2p)(5 + 2p)(7 + 2p)} - \frac{d(5ad - bc(11 + 2p))x (a + bx^2)^{1+p}}{b^2(5 + 2p)(7 + 2p)} \\ &= \frac{d(15a^2d^2 - 8abcd(6 + p) + b^2c^2(57 + 28p + 4p^2))x (a + bx^2)^{1+p}}{b^3(3 + 2p)(5 + 2p)(7 + 2p)} - \frac{d(5ad - bc(11 + 2p))x (a + bx^2)^{1+p}}{b^2(5 + 2p)(7 + 2p)} \end{aligned}$$

Mathematica [A] time = 5.07, size = 136, normalized size = 0.46

$$\frac{1}{35}x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} \left(35c^3 {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right) + dx^2 \left(35c^2 {}_2F_1\left(\frac{3}{2}, -p; \frac{5}{2}; -\frac{bx^2}{a}\right) + dx^2 \left(21c {}_2F_1\left(\frac{5}{2}, -p; \frac{7}{2}; -\frac{bx^2}{a}\right) + 5d {}_2F_1\left(\frac{7}{2}, -p; \frac{9}{2}; -\frac{bx^2}{a}\right)\right)\right)\right) / (35(1 + (bx^2)/a)^p)$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + b*x^2)^p*(c + d*x^2)^3,x]
```

```
[Out] (x*(a + b*x^2)^p*(35*c^3*Hypergeometric2F1[1/2, -p, 3/2, -((b*x^2)/a)] + d*
x^2*(35*c^2*Hypergeometric2F1[3/2, -p, 5/2, -((b*x^2)/a)] + d*x^2*(21*c*Hypergeometric2F1[5/2, -p, 7/2, -((b*x^2)/a)] + 5*d*x^2*Hypergeometric2F1[7/2, -p, 9/2, -((b*x^2)/a)])))/(35*(1 + (b*x^2)/a)^p)
```

fricas [F] time = 1.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(d^3x^6 + 3cd^2x^4 + 3c^2dx^2 + c^3\right)(bx^2 + a)^p, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^p*(d*x^2+c)^3,x, algorithm="fricas")
```

```
[Out] integral((d^3*x^6 + 3*c*d^2*x^4 + 3*c^2*d*x^2 + c^3)*(b*x^2 + a)^p, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (dx^2 + c)^3 (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c)^3,x, algorithm="giac")

[Out] integrate((d*x^2 + c)^3*(b*x^2 + a)^p, x)

maple [F] time = 0.36, size = 0, normalized size = 0.00

$$\int (dx^2 + c)^3 (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^p*(d*x^2+c)^3,x)

[Out] int((b*x^2+a)^p*(d*x^2+c)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (dx^2 + c)^3 (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c)^3,x, algorithm="maxima")

[Out] integrate((d*x^2 + c)^3*(b*x^2 + a)^p, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int (bx^2 + a)^p (dx^2 + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^p*(c + d*x^2)^3,x)

[Out] int((a + b*x^2)^p*(c + d*x^2)^3, x)

sympy [C] time = 39.51, size = 121, normalized size = 0.41

$$a^p c^3 x {}_2F_1 \left(\frac{1}{2}, -p \left| \frac{bx^2 e^{i\pi}}{a} \right. \right) + a^p c^2 dx^3 {}_2F_1 \left(\frac{3}{2}, -p \left| \frac{bx^2 e^{i\pi}}{a} \right. \right) + \frac{3a^p cd^2 x^5 {}_2F_1 \left(\frac{5}{2}, -p \left| \frac{bx^2 e^{i\pi}}{a} \right. \right)}{5} + \frac{a^p d^3 x^7 {}_2F_1 \left(\frac{7}{2}, -p \left| \frac{bx^2 e^{i\pi}}{a} \right. \right)}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**p*(d*x**2+c)**3,x)

[Out] a**p*c**3*x*hyper((1/2, -p), (3/2,), b*x**2*exp_polar(I*pi)/a) + a**p*c**2*d*x**3*hyper((3/2, -p), (5/2,), b*x**2*exp_polar(I*pi)/a) + 3*a**p*c*d**2*x**5*hyper((5/2, -p), (7/2,), b*x**2*exp_polar(I*pi)/a)/5 + a**p*d**3*x**7*hyper((7/2, -p), (9/2,), b*x**2*exp_polar(I*pi)/a)/7

3.343 $\int (a + bx^2)^p (c + dx^2)^2 dx$

Optimal. Leaf size=176

$$\frac{x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} (3a^2d^2 - 2abcd(2p + 5) + b^2c^2(4p^2 + 16p + 15)) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right) dx (a + bx^2)^{p+1}}{b^2(2p + 3)(2p + 5) b^2(2p + 3)}$$

[Out] $-d*(3*a*d-b*c*(7+2*p))*x*(b*x^2+a)^{(1+p)}/b^2/(4*p^2+16*p+15)+d*x*(b*x^2+a)^{(1+p)*(d*x^2+c)/b/(5+2*p)+(3*a^2*d^2-2*a*b*c*d*(5+2*p)+b^2*c^2*(4*p^2+16*p+15))*x*(b*x^2+a)^p*\text{hypergeom}([1/2, -p], [3/2], -b*x^2/a)/b^2/(4*p^2+16*p+15)/((1+b*x^2/a)^p)$

Rubi [A] time = 0.12, antiderivative size = 176, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {416, 388, 246, 245}

$$\frac{x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} (3a^2d^2 - 2abcd(2p + 5) + b^2c^2(4p^2 + 16p + 15)) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right) dx (a + bx^2)^{p+1}}{b^2(2p + 3)(2p + 5) b^2(2p + 3)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^p*(c + d*x^2)^2,x]

[Out] $-((d*(3*a*d - b*c*(7 + 2*p))*x*(a + b*x^2)^{(1 + p)})/(b^2*(3 + 2*p)*(5 + 2*p))) + (d*x*(a + b*x^2)^{(1 + p)*(c + d*x^2)})/(b*(5 + 2*p)) + ((3*a^2*d^2 - 2*a*b*c*d*(5 + 2*p) + b^2*c^2*(15 + 16*p + 4*p^2))*x*(a + b*x^2)^p*\text{Hypergeometric2F1}[1/2, -p, 3/2, -((b*x^2)/a)])/((b^2*(3 + 2*p)*(5 + 2*p)*(1 + (b*x^2)/a)^p)$

Rule 245

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, -((b*x^n)/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])

Rule 246

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1))/(b*(n*(p + q) + 1)), x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp[c*(b*c*(n*(p + q) + 1) - a*d] + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q - 1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a

, b, c, d, n, p, q, x]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^p (c + dx^2)^2 dx &= \frac{dx (a + bx^2)^{1+p} (c + dx^2)}{b(5 + 2p)} + \frac{\int (a + bx^2)^p (-c(ad - bc(5 + 2p)) - d(3ad - bc(7 + 2p))) dx}{b(5 + 2p)} \\ &= -\frac{d(3ad - bc(7 + 2p))x (a + bx^2)^{1+p}}{b^2(3 + 2p)(5 + 2p)} + \frac{dx (a + bx^2)^{1+p} (c + dx^2)}{b(5 + 2p)} + \frac{(3a^2d^2 - 2ad^2)}{b^2(3 + 2p)(5 + 2p)} \\ &= -\frac{d(3ad - bc(7 + 2p))x (a + bx^2)^{1+p}}{b^2(3 + 2p)(5 + 2p)} + \frac{dx (a + bx^2)^{1+p} (c + dx^2)}{b(5 + 2p)} + \frac{\left((3a^2d^2 - 2ad^2) \right)}{b^2(3 + 2p)(5 + 2p)} \\ &= -\frac{d(3ad - bc(7 + 2p))x (a + bx^2)^{1+p}}{b^2(3 + 2p)(5 + 2p)} + \frac{dx (a + bx^2)^{1+p} (c + dx^2)}{b(5 + 2p)} + \frac{(3a^2d^2 - 2ad^2)}{b^2(3 + 2p)(5 + 2p)} \end{aligned}$$

Mathematica [A] time = 5.05, size = 106, normalized size = 0.60

$$\frac{1}{15}x(a + bx^2)^p \left(\frac{bx^2}{a} + 1 \right)^{-p} \left(15c^2 {}_2F_1 \left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a} \right) + dx^2 \left(10c {}_2F_1 \left(\frac{3}{2}, -p; \frac{5}{2}; -\frac{bx^2}{a} \right) + 3dx^2 {}_2F_1 \left(\frac{5}{2}, -p; \frac{7}{2}; -\frac{bx^2}{a} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^p*(c + d*x^2)^2,x]

[Out] (x*(a + b*x^2)^p*(15*c^2*Hypergeometric2F1[1/2, -p, 3/2, -((b*x^2)/a)] + d*x^2*(10*c*Hypergeometric2F1[3/2, -p, 5/2, -((b*x^2)/a)] + 3*d*x^2*Hypergeometric2F1[5/2, -p, 7/2, -((b*x^2)/a)]))/(15*(1 + (b*x^2)/a)^p)

fricas [F] time = 1.52, size = 0, normalized size = 0.00

$$\text{integral} \left((d^2x^4 + 2cdx^2 + c^2)(bx^2 + a)^p, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c)^2,x, algorithm="fricas")

[Out] integral((d^2*x^4 + 2*c*d*x^2 + c^2)*(b*x^2 + a)^p, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (dx^2 + c)^2 (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate((d*x^2 + c)^2*(b*x^2 + a)^p, x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int (dx^2 + c)^2 (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^p*(d*x^2+c)^2,x)

[Out] int((b*x^2+a)^p*(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (dx^2 + c)^2 (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((d*x^2 + c)^2*(b*x^2 + a)^p, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (bx^2 + a)^p (dx^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^p*(c + d*x^2)^2,x)

[Out] int((a + b*x^2)^p*(c + d*x^2)^2, x)

sympy [C] time = 20.40, size = 88, normalized size = 0.50

$$a^p c^2 x {}_2F_1 \left(\begin{matrix} \frac{1}{2}, -p \\ \frac{3}{2} \end{matrix} \middle| \frac{bx^2 e^{i\pi}}{a} \right) + \frac{2a^p c d x^3 {}_2F_1 \left(\begin{matrix} \frac{3}{2}, -p \\ \frac{5}{2} \end{matrix} \middle| \frac{bx^2 e^{i\pi}}{a} \right)}{3} + \frac{a^p d^2 x^5 {}_2F_1 \left(\begin{matrix} \frac{5}{2}, -p \\ \frac{7}{2} \end{matrix} \middle| \frac{bx^2 e^{i\pi}}{a} \right)}{5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**p*(d*x**2+c)**2,x)

[Out] a**p*c**2*x*hyper((1/2, -p), (3/2,), b*x**2*exp_polar(I*pi)/a) + 2*a**p*c*d*x**3*hyper((3/2, -p), (5/2,), b*x**2*exp_polar(I*pi)/a)/3 + a**p*d**2*x**5*hyper((5/2, -p), (7/2,), b*x**2*exp_polar(I*pi)/a)/5

3.344 $\int (a + bx^2)^p (c + dx^2) dx$

Optimal. Leaf size=93

$$\frac{dx (a + bx^2)^{p+1}}{b(2p + 3)} - \frac{x (a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} (ad - bc(2p + 3)) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right)}{b(2p + 3)}$$

[Out] d*x*(b*x^2+a)^(1+p)/b/(3+2*p)-(a*d-b*c*(3+2*p))*x*(b*x^2+a)^p*hypergeom([1/2, -p], [3/2], -b*x^2/a)/b/(3+2*p)/((1+b*x^2/a)^p)

Rubi [A] time = 0.04, antiderivative size = 85, normalized size of antiderivative = 0.91, number of steps used = 3, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {388, 246, 245}

$$x (a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} \left(c - \frac{ad}{2bp + 3b}\right) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right) + \frac{dx (a + bx^2)^{p+1}}{b(2p + 3)}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^p*(c + d*x^2), x]

[Out] (d*x*(a + b*x^2)^(1 + p))/(b*(3 + 2*p)) + ((c - (a*d)/(3*b + 2*b*p))*x*(a + b*x^2)^p*Hypergeometric2F1[1/2, -p, 3/2, -((b*x^2)/a)])/(1 + (b*x^2)/a)^p

Rule 245

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, -((b*x^n)/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])

Rule 246

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)^p (c + dx^2) dx &= \frac{dx (a + bx^2)^{1+p}}{b(3 + 2p)} - \left(-c + \frac{ad}{3b + 2bp}\right) \int (a + bx^2)^p dx \\ &= \frac{dx (a + bx^2)^{1+p}}{b(3 + 2p)} - \left(\left(-c + \frac{ad}{3b + 2bp}\right) (a + bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p}\right) \int \left(1 + \frac{bx^2}{a}\right)^p dx \\ &= \frac{dx (a + bx^2)^{1+p}}{b(3 + 2p)} + \left(c - \frac{ad}{3b + 2bp}\right) x (a + bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right) \end{aligned}$$

Mathematica [A] time = 0.03, size = 90, normalized size = 0.97

$$\frac{x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} \left((bc(2p + 3) - ad) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right) + d(a + bx^2) \left(\frac{bx^2}{a} + 1\right)^p\right)}{b(2p + 3)}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^p*(c + d*x^2), x]

[Out] (x*(a + b*x^2)^p*(d*(a + b*x^2)*(1 + (b*x^2)/a)^p + (-a*d) + b*c*(3 + 2*p))*Hypergeometric2F1[1/2, -p, 3/2, -((b*x^2)/a)]/(b*(3 + 2*p)*(1 + (b*x^2)/a)^p)

fricas [F] time = 1.33, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(dx^2 + c\right)\left(bx^2 + a\right)^p, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c), x, algorithm="fricas")

[Out] integral((d*x^2 + c)*(b*x^2 + a)^p, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (dx^2 + c)(bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c), x, algorithm="giac")

[Out] integrate((d*x^2 + c)*(b*x^2 + a)^p, x)

maple [F] time = 0.30, size = 0, normalized size = 0.00

$$\int (dx^2 + c)(bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^p*(d*x^2+c), x)

[Out] int((b*x^2+a)^p*(d*x^2+c), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (dx^2 + c)(bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p*(d*x^2+c), x, algorithm="maxima")

[Out] integrate((d*x^2 + c)*(b*x^2 + a)^p, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (bx^2 + a)^p (dx^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^p*(c + d*x^2), x)`

[Out] `int((a + b*x^2)^p*(c + d*x^2), x)`

sympy [C] time = 9.83, size = 53, normalized size = 0.57

$$a^p c x {}_2F_1\left(\frac{1}{2}, -p \mid \frac{bx^2 e^{i\pi}}{a}\right) + \frac{a^p dx^3 {}_2F_1\left(\frac{3}{2}, -p \mid \frac{bx^2 e^{i\pi}}{a}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**p*(d*x**2+c), x)`

[Out] `a**p*c*x*hyper((1/2, -p), (3/2,), b*x**2*exp_polar(I*pi)/a) + a**p*d*x**3*hyper((3/2, -p), (5/2,), b*x**2*exp_polar(I*pi)/a)/3`

3.345 $\int (a + bx^2)^p dx$

Optimal. Leaf size=44

$$x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right)$$

[Out] $x*(b*x^2+a)^p*\text{hypergeom}([1/2, -p], [3/2], -b*x^2/a)/((1+b*x^2/a)^p)$

Rubi [A] time = 0.01, antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {246, 245}

$$x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right)$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^p, x]

[Out] $(x*(a + b*x^2)^p*\text{Hypergeometric2F1}[1/2, -p, 3/2, -((b*x^2)/a)])/(1 + (b*x^2)/a)^p$

Rule 245

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, -((b*x^n)/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])

Rule 246

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p, x], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])

Rubi steps

$$\begin{aligned} \int (a + bx^2)^p dx &= \left((a + bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} \right) \int \left(1 + \frac{bx^2}{a}\right)^p dx \\ &= x(a + bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right) \end{aligned}$$

Mathematica [A] time = 0.00, size = 44, normalized size = 1.00

$$x(a + bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right)$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^p, x]

[Out] $(x*(a + b*x^2)^p*\text{Hypergeometric2F1}[1/2, -p, 3/2, -((b*x^2)/a)])/(1 + (b*x^2)/a)^p$

fricas [F] time = 0.88, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(bx^2 + a\right)^p, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p,x, algorithm="fricas")

[Out] integral((b*x^2 + a)^p, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p,x, algorithm="giac")

[Out] integrate((b*x^2 + a)^p, x)

maple [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^p,x)

[Out] int((b*x^2+a)^p,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^p, x)

mupad [B] time = 5.57, size = 41, normalized size = 0.93

$$\frac{x (bx^2 + a)^p {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{bx^2}{a}\right)}{\left(\frac{bx^2}{a} + 1\right)^p}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^p,x)

[Out] (x*(a + b*x^2)^p*hypergeom([1/2, -p], 3/2, -(b*x^2)/a))/((b*x^2)/a + 1)^p

sympy [C] time = 2.33, size = 22, normalized size = 0.50

$$a^p x {}_2F_1\left(\frac{1}{2}, -p \left| \frac{bx^2 e^{i\pi}}{a} \right. \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**p,x)

[Out] a**p*x*hyper((1/2, -p), (3/2,), b*x**2*exp_polar(I*pi)/a)

$$3.346 \quad \int \frac{(a+bx^2)^p}{c+dx^2} dx$$

Optimal. Leaf size=57

$$\frac{x(a+bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} F_1\left(\frac{1}{2}; -p, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c}$$

[Out] x*(b*x^2+a)^p*AppellF1(1/2,-p,1,3/2,-b*x^2/a,-d*x^2/c)/c/((1+b*x^2/a)^p)

Rubi [A] time = 0.03, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {430, 429}

$$\frac{x(a+bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} F_1\left(\frac{1}{2}; -p, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^p/(c + d*x^2), x]

[Out] (x*(a + b*x^2)^p*AppellF1[1/2, -p, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c)])/(c*(1 + (b*x^2)/a)^p)

Rule 429

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 :> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, -((b*x^n)/a), -((d*x^n)/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 430

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 :> Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])

Rubi steps

$$\begin{aligned} \int \frac{(a+bx^2)^p}{c+dx^2} dx &= \left((a+bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} \right) \int \frac{\left(1 + \frac{bx^2}{a}\right)^p}{c+dx^2} dx \\ &= \frac{x(a+bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} F_1\left(\frac{1}{2}; -p, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c} \end{aligned}$$

Mathematica [B] time = 0.18, size = 162, normalized size = 2.84

$$\frac{3acx(a+bx^2)^p F_1\left(\frac{1}{2}; -p, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{(c+dx^2) \left(2x^2 \left(adF_1\left(\frac{3}{2}; -p, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - bcpF_1\left(\frac{3}{2}; 1-p, 1; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) \right) - 3acF_1\left(\frac{1}{2}; -p, 1; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) }$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^p/(c + d*x^2), x]


```
[Out] (-3*a*c*x*(a + b*x^2)^p*AppellF1[1/2, -p, 1, 3/2, -((b*x^2)/a), -((d*x^2)/c
)])/((c + d*x^2)*(-3*a*c*AppellF1[1/2, -p, 1, 3/2, -((b*x^2)/a), -((d*x^2)/
c)] + 2*x^2*(-(b*c*p*AppellF1[3/2, 1 - p, 1, 5/2, -((b*x^2)/a), -((d*x^2)/c
)]) + a*d*AppellF1[3/2, -p, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))
```

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(bx^2 + a)^p}{dx^2 + c}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^p/(d*x^2+c),x, algorithm="fricas")
```

```
[Out] integral((b*x^2 + a)^p/(d*x^2 + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^p/(d*x^2+c),x, algorithm="giac")
```

```
[Out] integrate((b*x^2 + a)^p/(d*x^2 + c), x)
```

maple [F] time = 0.32, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*x^2+a)^p/(d*x^2+c),x)
```

```
[Out] int((b*x^2+a)^p/(d*x^2+c),x)
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^p/(d*x^2+c),x, algorithm="maxima")
```

```
[Out] integrate((b*x^2 + a)^p/(d*x^2 + c), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(bx^2 + a)^p}{dx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a + b*x^2)^p/(c + d*x^2),x)
```

```
[Out] int((a + b*x^2)^p/(c + d*x^2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)**p/(d*x**2+c),x)
```

```
[Out] Timed out
```

$$3.347 \quad \int \frac{(a+bx^2)^p}{(c+dx^2)^2} dx$$

Optimal. Leaf size=57

$$\frac{x(a+bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} F_1\left(\frac{1}{2}; -p, 2; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c^2}$$

[Out] $x*(b*x^2+a)^p*AppellF1(1/2,-p,2,3/2,-b*x^2/a,-d*x^2/c)/c^2/((1+b*x^2/a)^p)$

Rubi [A] time = 0.03, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {430, 429}

$$\frac{x(a+bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} F_1\left(\frac{1}{2}; -p, 2; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^p/(c + d*x^2)^2,x]

[Out] $(x*(a + b*x^2)^p*AppellF1[1/2, -p, 2, 3/2, -((b*x^2)/a), -((d*x^2)/c)])/(c^2*(1 + (b*x^2)/a)^p)$

Rule 429

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 :> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, -((b*x^n)/a), -((d*x^n)/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 430

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 :> Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])

Rubi steps

$$\begin{aligned} \int \frac{(a+bx^2)^p}{(c+dx^2)^2} dx &= \left((a+bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} \right) \int \frac{\left(1 + \frac{bx^2}{a}\right)^p}{(c+dx^2)^2} dx \\ &= \frac{x(a+bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} F_1\left(\frac{1}{2}; -p, 2; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c^2} \end{aligned}$$

Mathematica [B] time = 0.19, size = 162, normalized size = 2.84

$$\frac{3acx(a+bx^2)^p F_1\left(\frac{1}{2}; -p, 2; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{(c+dx^2)^2 \left(-2x^2 \left(bcp F_1\left(\frac{3}{2}; 1-p, 2; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - 2ad F_1\left(\frac{3}{2}; -p, 3; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)\right) - 3ac F_1\left(\frac{1}{2}; -p, 2; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^p/(c + d*x^2)^2,x]

[Out] (-3*a*c*x*(a + b*x^2)^p*AppellF1[1/2, -p, 2, 3/2, -((b*x^2)/a), -((d*x^2)/c)])/((c + d*x^2)^2*(-3*a*c*AppellF1[1/2, -p, 2, 3/2, -((b*x^2)/a), -((d*x^2)/c)] - 2*x^2*(b*c*p*AppellF1[3/2, 1 - p, 2, 5/2, -((b*x^2)/a), -((d*x^2)/c)] - 2*a*d*AppellF1[3/2, -p, 3, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))

fricas [F] time = 1.00, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(bx^2 + a)^p}{d^2x^4 + 2cdx^2 + c^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p/(d*x^2+c)^2,x, algorithm="fricas")

[Out] integral((b*x^2 + a)^p/(d^2*x^4 + 2*c*d*x^2 + c^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p/(d*x^2+c)^2,x, algorithm="giac")

[Out] integrate((b*x^2 + a)^p/(d*x^2 + c)^2, x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^p/(d*x^2+c)^2,x)

[Out] int((b*x^2+a)^p/(d*x^2+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p/(d*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^p/(d*x^2 + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(bx^2 + a)^p}{(dx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^p/(c + d*x^2)^2,x)

```
[Out] int((a + b*x^2)^p/(c + d*x^2)^2, x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)**p/(d*x**2+c)**2,x)
```

```
[Out] Timed out
```

$$3.348 \quad \int \frac{(a+bx^2)^p}{(c+dx^2)^3} dx$$

Optimal. Leaf size=57

$$\frac{x(a+bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} F_1\left(\frac{1}{2}; -p, 3; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c^3}$$

[Out] $x*(b*x^2+a)^p*AppellF1(1/2, -p, 3, 3/2, -b*x^2/a, -d*x^2/c)/c^3/((1+b*x^2/a)^p)$

Rubi [A] time = 0.03, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {430, 429}

$$\frac{x(a+bx^2)^p \left(\frac{bx^2}{a} + 1\right)^{-p} F_1\left(\frac{1}{2}; -p, 3; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c^3}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^p/(c + d*x^2)^3,x]

[Out] $(x*(a + b*x^2)^p*AppellF1[1/2, -p, 3, 3/2, -((b*x^2)/a), -((d*x^2)/c)])/(c^3*(1 + (b*x^2)/a)^p)$

Rule 429

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 := Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, -((b*x^n)/a), -((d*x^n)/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 430

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 := Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])

Rubi steps

$$\begin{aligned} \int \frac{(a+bx^2)^p}{(c+dx^2)^3} dx &= \left((a+bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} \right) \int \frac{\left(1 + \frac{bx^2}{a}\right)^p}{(c+dx^2)^3} dx \\ &= \frac{x(a+bx^2)^p \left(1 + \frac{bx^2}{a}\right)^{-p} F_1\left(\frac{1}{2}; -p, 3; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{c^3} \end{aligned}$$

Mathematica [B] time = 0.25, size = 162, normalized size = 2.84

$$\frac{3acx(a+bx^2)^p F_1\left(\frac{1}{2}; -p, 3; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}{(c+dx^2)^3 \left(-2x^2 \left(bcpF_1\left(\frac{3}{2}; 1-p, 3; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right) - 3adF_1\left(\frac{3}{2}; -p, 4; \frac{5}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)\right) - 3acF_1\left(\frac{1}{2}; -p, 3; \frac{3}{2}; -\frac{bx^2}{a}, -\frac{dx^2}{c}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a + b*x^2)^p/(c + d*x^2)^3,x]

[Out] (-3*a*c*x*(a + b*x^2)^p*AppellF1[1/2, -p, 3, 3/2, -((b*x^2)/a), -((d*x^2)/c)])/((c + d*x^2)^3*(-3*a*c*AppellF1[1/2, -p, 3, 3/2, -((b*x^2)/a), -((d*x^2)/c)] - 2*x^2*(b*c*p*AppellF1[3/2, 1 - p, 3, 5/2, -((b*x^2)/a), -((d*x^2)/c)] - 3*a*d*AppellF1[3/2, -p, 4, 5/2, -((b*x^2)/a), -((d*x^2)/c)]))

fricas [F] time = 1.00, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(bx^2 + a)^p}{d^3x^6 + 3cd^2x^4 + 3c^2dx^2 + c^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p/(d*x^2+c)^3,x, algorithm="fricas")

[Out] integral((b*x^2 + a)^p/(d^3*x^6 + 3*c*d^2*x^4 + 3*c^2*d*x^2 + c^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p/(d*x^2+c)^3,x, algorithm="giac")

[Out] integrate((b*x^2 + a)^p/(d*x^2 + c)^3, x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^p/(d*x^2+c)^3,x)

[Out] int((b*x^2+a)^p/(d*x^2+c)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^p}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^p/(d*x^2+c)^3,x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^p/(d*x^2 + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(bx^2 + a)^p}{(dx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^p/(c + d*x^2)^3,x)

```
[Out] int((a + b*x^2)^p/(c + d*x^2)^3, x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)**p/(d*x**2+c)**3,x)
```

```
[Out] Timed out
```


$$3.349 \quad \int (a + bx^2)^{-1 - \frac{bc}{2bc-2ad}} (c + dx^2)^{-1 + \frac{ad}{2bc-2ad}} dx$$

Optimal. Leaf size=53

$$\frac{x (a + bx^2)^{-\frac{bc}{2bc-2ad}} (c + dx^2)^{\frac{ad}{2bc-2ad}}}{ac}$$

[Out] $x*(d*x^2+c)^{(a*d/(-2*a*d+2*b*c))}/a/c/((b*x^2+a)^{(b*c/(-2*a*d+2*b*c))})$

Rubi [A] time = 0.02, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 50, $\frac{\text{number of rules}}{\text{integrand size}} = 0.020$, Rules used = {381}

$$\frac{x (a + bx^2)^{-\frac{bc}{2bc-2ad}} (c + dx^2)^{\frac{ad}{2bc-2ad}}}{ac}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*x^2)^{-1 - (b*c)/(2*b*c - 2*a*d)}*(c + d*x^2)^{-1 + (a*d)/(2*b*c - 2*a*d)}, x]$

[Out] $(x*(c + d*x^2)^{((a*d)/(2*b*c - 2*a*d))})/(a*c*(a + b*x^2)^{((b*c)/(2*b*c - 2*a*d))})$

Rule 381

$\text{Int}[(a + b*x^2)^{-1 - \frac{bc}{2bc-2ad}} (c + dx^2)^{-1 + \frac{ad}{2bc-2ad}} dx = \frac{x (a + bx^2)^{-\frac{bc}{2bc-2ad}} (c + dx^2)^{\frac{ad}{2bc-2ad}}}{ac}$

Rubi steps

Mathematica [A] time = 0.03, size = 52, normalized size = 0.98

$$\frac{x (a + bx^2)^{\frac{bc}{2ad-2bc}} (c + dx^2)^{\frac{ad}{2bc-2ad}}}{ac}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[(a + b*x^2)^{-1 - (b*c)/(2*b*c - 2*a*d)}*(c + d*x^2)^{-1 + (a*d)/(2*b*c - 2*a*d)}, x]$

[Out] $(x*(a + b*x^2)^{((b*c)/(-2*b*c + 2*a*d))}*(c + d*x^2)^{((a*d)/(2*b*c - 2*a*d))})/(a*c)$

fricas [A] time = 1.02, size = 91, normalized size = 1.72

$$\frac{bdx^5 + (bc + ad)x^3 + acx}{(bx^2 + a)^{\frac{3bc-2ad}{2(bc-ad)}} (dx^2 + c)^{\frac{2bc-3ad}{2(bc-ad)}} ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(-1-b*c/(-2*a*d+2*b*c))*(d*x^2+c)^(-1+a*d/(-2*a*d+2*b*c)),x, algorithm="fricas")

[Out] (b*d*x^5 + (b*c + a*d)*x^3 + a*c*x)/((b*x^2 + a)^(1/2*(3*b*c - 2*a*d)/(b*c - a*d))*(d*x^2 + c)^(1/2*(2*b*c - 3*a*d)/(b*c - a*d))*a*c)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^{-\frac{bc}{2(bc-ad)}-1} (dx^2 + c)^{\frac{ad}{2(bc-ad)}-1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(-1-b*c/(-2*a*d+2*b*c))*(d*x^2+c)^(-1+a*d/(-2*a*d+2*b*c)),x, algorithm="giac")

[Out] integrate((b*x^2 + a)^(-1/2*b*c/(b*c - a*d) - 1)*(d*x^2 + c)^(1/2*a*d/(b*c - a*d) - 1), x)

maple [A] time = 0.00, size = 71, normalized size = 1.34

$$\frac{x (bx^2 + a)^{1-\frac{2ad-3bc}{2(ad-bc)}} (dx^2 + c)^{1-\frac{3ad-2bc}{2(ad-bc)}}}{ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)^(-1-b*c/(-2*a*d+2*b*c))*(d*x^2+c)^(-1+a*d/(-2*a*d+2*b*c)),x)

[Out] (b*x^2+a)^(1-1/2*(2*a*d-3*b*c)/(a*d-b*c))*(d*x^2+c)^(1-1/2*(3*a*d-2*b*c)/(a*d-b*c))/a/c*x

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (bx^2 + a)^{-\frac{bc}{2(bc-ad)}-1} (dx^2 + c)^{\frac{ad}{2(bc-ad)}-1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^(-1-b*c/(-2*a*d+2*b*c))*(d*x^2+c)^(-1+a*d/(-2*a*d+2*b*c)),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^(-1/2*b*c/(b*c - a*d) - 1)*(d*x^2 + c)^(1/2*a*d/(b*c - a*d) - 1), x)

mupad [B] time = 5.74, size = 131, normalized size = 2.47

$$\frac{x (bx^2 + a)^{\frac{bc}{2ad-2bc}-1} + \frac{x^3 (bx^2+a)^{\frac{bc}{2ad-2bc}-1} (ad+bc)}{ac} + \frac{bdx^5 (bx^2+a)^{\frac{bc}{2ad-2bc}-1}}{ac}}{(dx^2 + c)^{\frac{ad}{2ad-2bc}+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^((b*c)/(2*a*d - 2*b*c) - 1)/(c + d*x^2)^((a*d)/(2*a*d - 2*b*c) + 1),x)

[Out] (x*(a + b*x^2)^((b*c)/(2*a*d - 2*b*c) - 1) + (x^3*(a + b*x^2)^((b*c)/(2*a*d - 2*b*c) - 1)*(a*d + b*c))/(a*c) + (b*d*x^5*(a + b*x^2)^((b*c)/(2*a*d - 2*b*c) - 1))/(a*c)/(c + d*x^2)^((a*d)/(2*a*d - 2*b*c) + 1)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)**(-1-b*c/(-2*a*d+2*b*c))*(d*x**2+c)**(-1+a*d/(-2*a*d+2*b*c)),x)
```

```
[Out] Timed out
```


Chapter 4

Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.0.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] :=
  If[ExpnType[result]<=ExpnType[optimal],
    If[FreeQ[result,Complex] || Not[FreeQ[optimal,Complex]],
      If[LeafCount[result]<=2*LeafCount[optimal],
        "A",
        "B"],
      "C"],
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      "C",
      "F"]]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
```

```

(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3, ExpnType[expn[[1]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
          If[HypergeometricFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
          If[AppellFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
          If[Head[expn]===RootSum,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
          If[Head[expn]===Integrate || Head[expn]===Int,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
          9]]]]]]]]]]

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```
AppellFunctionQ[func_] :=
  MemberQ[{AppellF1},func]
```

4.0.2 Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems

GradeAntiderivative := proc(result,optimal)
local leaf_count_result, leaf_count_optimal,ExpnType_result,ExpnType_optimal,
  debug:=false;

  leaf_count_result:=leafcount(result);
  #do NOT call ExpnType() if leaf size is too large. Recursion problem
  if leaf_count_result > 500000 then
    return "B";
  fi;

  leaf_count_optimal:=leafcount(optimal);

  ExpnType_result:=ExpnType(result);
  ExpnType_optimal:=ExpnType(optimal);

  if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",
  ExpnType_optimal);
  fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
  return "F";
end if;

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;

```

```

if is_contains_complex(result) then
  if is_contains_complex(optimal) then
    if debug then
      print("both result and optimal complex");
    fi;
    #both result and optimal complex
    if leaf_count_result<=2*leaf_count_optimal then
      return "A";
    else
      return "B";
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C";
  end if
else # result do not contain complex
  # this assumes optimal do not as well
  if debug then
    print("result do not contain complex, this assumes optimal do
not as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B";
  end if
end if
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C";
end if

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function

```



```

# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+' or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

```

```

AppellFunctionQ := proc(func)
  member(func,[AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.0.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]

def is_hypergeometric_function(func):
  return func in [hyper]

def is_appell_function(func):
  return func in [appellf1]

```

```

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+`' or
    type(expn,'*`)
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,
    expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType
    ,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:
        m1 = max(map(expnType, list(expn.args)))
        return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    else:
        return 9

```

```

#main function
def grade_antiderivative(result,optimal):

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        return "F"

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
    else:
        return "C"

```

4.0.4 SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#           Albert Rich to use with Sagemath. This is used to
#           grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#           'arctan2','floor','abs','log_integral'

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True

```

```

        else:
            return False
    else:
        return False

def is_elementary_function(func):
    debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    debug=False
    if debug: print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','
sinh_integral'
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M',
        hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in
sagemath

def is_atom(expn):

    debug=False
    if debug: print ("Enter is_atom")

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-
sagemath-equivalent-to-atomic-type-in-maple/
    try:
        if expn.parent() is SR:

```

```

        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens")
:
        return expn in expn.parent().base_ring() or expn in expn.parent().
gens()
    return False

except AttributeError as error:
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],
Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer
)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],
Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],
Rational)
                return 1
            else:
                return max(2,expnType(expn.operands()[0])) #max(2,expnType(
expn.args[0]))
        else:
            return max(3,expnType(expn.operands()[0]),expnType(expn.operands(
[1])) #max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))
    elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #
isinstance(expn,Add) or isinstance(expn,Mul)
        m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
        m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn)))
    elif is_elementary_function(expn.operator()): #is_elementary_function(expn
.func)
        return max(3,expnType(expn.operands()[0]))
    elif is_special_function(expn.operator()): #is_special_function(expn.func)
        m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
        return max(4,m1) #max(4,m1)
    elif is_hypergeometric_function(expn.operator()): #
is_hypergeometric_function(expn.func)
        m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
        return max(5,m1) #max(5,m1)
    elif is_appell_function(expn.operator()):

```

```

        m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(
expn.args)))
        return max(6,m1)      #max(6,m1)
    elif str(expn).find("Integral") != -1: #this will never happen, since it
        #is checked before calling the grading function that is passed.
        #but kept it here.
        m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(
expn.args)))
        return max(8,m1)      #max(5,apply(max,map(ExpnType,[op(expn)])))
    else:
        return 9

#main function
def grade_antiderivative(result,optimal):

    if debug: print ("Enter grade_antiderivative for sagemath")

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    if debug: print ("leaf_count_result=", leaf_count_result, "
leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",
expnType_optimal)

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
    else:
        return "C"

```