

Computer algebra independent integration tests

1-Algebraic-functions/1.2-Trinomial-products/1.2.2-Quartic/1.2.2.7-P-x-d+e-x^2-^q-a+b-x^2+c-x^4-^p

Nasser M. Abbasi

June 29, 2021

Compiled on June 29, 2021 at 10:42am

Contents

1	Introduction	5
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Performance	10
1.4	list of integrals that has no closed form antiderivative	11
1.5	list of integrals solved by CAS but has no known antiderivative	12
1.6	list of integrals solved by CAS but failed verification	12
1.7	Timing	13
1.8	Verification	13
1.9	Important notes about some of the results	13
1.9.1	Important note about Maxima results	13
1.9.2	Important note about FriCAS and Giac/XCAS results	14
1.9.3	Important note about finding leaf size of antiderivative	14
1.9.4	Important note about Mupad results	15
1.10	Design of the test system	16
2	detailed summary tables of results	17
2.1	List of integrals sorted by grade for each CAS	17
2.1.1	Rubi	17
2.1.2	Mathematica	17
2.1.3	Maple	17

2.1.4	Maxima	18
2.1.5	FriCAS	18
2.1.6	Sympy	18
2.1.7	Giac	18
2.1.8	Mupad	19
2.2	Detailed conclusion table per each integral for all CAS systems	20
2.3	Detailed conclusion table specific for Rubi results	28
3	Listing of integrals	31
3.1	$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$	31
3.2	$\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$	37
3.3	$\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+cx^4}} dx$	42
3.4	$\int \frac{A+Bx^2}{\sqrt{a+cx^4}} dx$	47
3.5	$\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx$	51
3.6	$\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+cx^4}} dx$	55
3.7	$\int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+cx^4}} dx$	61
3.8	$\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$	68
3.9	$\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$	74
3.10	$\int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$	79
3.11	$\int \frac{A+Bx^2}{(a+cx^4)^{3/2}} dx$	84
3.12	$\int \frac{A+Bx^2}{(d+ex^2)(a+cx^4)^{3/2}} dx$	88
3.13	$\int \frac{A+Bx^2}{(d+ex^2)^2(a+cx^4)^{3/2}} dx$	94
3.14	$\int \frac{A+Bx^2}{(d+ex^2)^3(a+cx^4)^{3/2}} dx$	102
3.15	$\int \frac{(A+Bx^2)(d+ex^2)^\eta}{a+cx^4} dx$	111
3.16	$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx$	115
3.17	$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$	118

3.18	$\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$	126
3.19	$\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+bx^2+cx^4}} dx$	132
3.20	$\int \frac{A+Bx^2}{\sqrt{a+bx^2+cx^4}} dx$	137
3.21	$\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$	141
3.22	$\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+bx^2+cx^4}} dx$	145
3.23	$\int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+bx^2+cx^4}} dx$	152
3.24	$\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$	160
3.25	$\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$	166
3.26	$\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$	172
3.27	$\int \frac{A+Bx^2}{(a+bx^2+cx^4)^{3/2}} dx$	177
3.28	$\int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$	182
3.29	$\int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx$	190
3.30	$\int \frac{\sqrt{a} + \sqrt{c}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$	197
3.31	$\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$	201
3.32	$\int \frac{946+315x^2}{(7+5x^2)\sqrt{2+3x^2+x^4}} dx$	205
3.33	$\int \frac{(A+Bx^2)(d+ex^2)^q}{a+bx^2+cx^4} dx$	209
3.34	$\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx$	213
3.35	$\int \frac{\sqrt{a+bx^2+cx^4}}{ad-cdx^4} dx$	218
3.36	$\int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx$	222
3.37	$\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$	226
3.38	$\int \sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$	231
3.39	$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx$	236

3.40	$\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$	242
3.41	$\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$	248
3.42	$\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$	254
4	Listing of Grading functions	261
4.0.1	Mathematica and Rubi grading function	261
4.0.2	Maple grading function	263
4.0.3	Sympy grading function	268
4.0.4	SageMath grading function	271

Chapter 1

Introduction

This report gives the result of running the computer algebra independent integration problems. The listing of the problems are maintained by and can be downloaded from <https://rulebasedintegration.org>

The number of integrals in this report is [42]. This is test number [44].

1.1 Listing of CAS systems tested

The following systems were tested at this time.

1. Mathematica 12.3 (64 bit) on windows 10.
2. Rubi 4.16.1 in Mathematica 12.1 on windows 10.
3. Maple 2021.1 (64 bit) on windows 10.
4. Maxima 5.44 on Linux. (via sagemath 9.3)
5. Fricas 1.3.7 on Linux (via sagemath 9.3)
6. Giac/Xcas 1.7 on Linux. (via sagemath 9.3)
7. Sympy 1.8 under Python 3.8.8 using Anaconda distribution on Ubuntu.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 under windows 10 (64 bit)

Maxima, Fricas and Giac/Xcas were called from inside SageMath. This was done using SageMath integrate command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly using Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	solved	Failed
Rubi	% 100.00 (42)	% 0.00 (0)
Mathematica	% 90.48 (38)	% 9.52 (4)
Maple	% 95.24 (40)	% 4.76 (2)
Maxima	% 0.00 (0)	% 100.00 (42)
Fricas	% 21.43 (9)	% 78.57 (33)
Sympy	% 14.29 (6)	% 85.71 (36)
Giac	% 11.90 (5)	% 88.10 (37)
Mupad	% 2.38 (1)	% 97.62 (41)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	14.29	0.00	76.19	9.52
Maple	23.81	33.33	38.10	4.76
Maxima	0.00	0.00	0.00	100.00
Fricas	14.29	7.14	0.00	78.57
Sympy	0.00	0.00	14.29	85.71
Giac	9.52	2.38	0.00	88.10
Mupad	0.00	2.38	0.00	97.62

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	4	50.00 %	50.00 %	0.00 %
Maple	2	100.00 %	0.00 %	0.00 %
Maxima	42	83.33 %	0.00 %	16.67 %
Fricas	33	63.64 %	36.36 %	0.00 %
Sympy	36	69.44 %	30.56 %	0.00 %
Giac	37	91.89 %	2.70 %	5.41 %
Mupad	41	100.00 %	0.00 %	0.00 %

Table 1.4: Time and leaf size performance for each CAS



1.3 Performance

The table below summarizes the performance of each CAS system in terms of CPU time and leaf size of results.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.81	576.24	1.08	396.50	1.00
Mathematica	1.62	824.05	1.30	297.50	0.66
Maple	0.03	1058.30	1.70	450.00	1.33
Maxima	0.00	0.00	0.00	0.00	0.00
Fricas	3.02	599.44	2.53	669.00	2.60
Sympy	7.04	186.00	0.53	167.00	0.51
Giac	0.82	359.00	1.32	360.00	1.17
Mupad	1.85	397.00	3.75	397.00	3.75

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used columns from the above table.



1.4 list of integrals that has no closed form antiderivative

{

1.5 list of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {29}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of _int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call has completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out is not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica. Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative produced was correct.

Verification phase has 3 minutes time out. An integral whose result was not verified could still be correct. Further investigation is needed on those integrals which failed verifications. Such integrals are marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since these integrals are run in a batch mode, using an automated script, and by using `sagemath` (SageMath uses Maxima), then any integral where Maxima needs an interactive response from the user to answer a question during evaluation of the integral in order to complete the integration, will fail and is counted as failed.

The exception raised is `ValueError`. Therefore Maxima result below is lower than what could result if Maxima was run directly and each question Maxima asks was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be indentified by looking at the output of the integration in each section for Maxima. The exception message will indicate of the error is due to the interactive question being asked or not.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath loading of Maxima `abs_integrate` was found to cause some problem. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS and Giac/X-CAS results

There are Few integrals which failed due to SageMath not able to translate the result back to SageMath syntax and not because these CAS system were not able to do the integrations.

These will fail With error `Exception raised: NotImplementedError`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals.

Hopefully the next version of SageMath will have complete translation of FriCAS and XCAS syntax and I will re-run all the tests again when this happens.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi and Maple, the builtin system function `LeafSize` is used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy and Giac antiderivatives is determined using the following function, thanks to user `slelievre` at <https://>

ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which is called directly from Python, the following code is used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative, Maple was used to determine the leaf size of Mupad output by post processing.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
 2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
 3. integer. Leaf size of result.
 4. integer. Leaf size of the optimal antiderivative.
 5. number. CPU time used to solve this integral. 0 if failed.
 6. string. The integral in Latex format
 7. string. The input used in CAS own syntax.
 8. string. The result (antiderivative) produced by CAS in Latex format
 9. string. The optimal antiderivative in Latex format.
 10. integer. 0 or 1. Indicates if problem has known antiderivative or not
 11. String. The result (antiderivative) in CAS own syntax.
 12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
- The following field present only in Rubi and Mathematica Tables*
13. integer. 1 if result was verified or 0 if not verified.
- The following fields present only in Rubi Tables*
14. integer. Number of rules used.
 15. integer. Integrand leaf size.
 16. real number. Ratio of field 14 over field 15
 17. integer. 1 if result was verified or 0 if not verified.
 18. String of form "{n,n,...}" which is list of the rules used by Rubi

High level overview of the CAS independent integration test build system

Chapter 2

detailed summary tables of results

2.1 List of integrals sorted by grade for each CAS

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 37, 38, 39, 40, 41, 42 }

B grade: { }

C grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 28, 29, 30, 31, 32, 34, 35, 36 }

F grade: { 15, 26, 27, 33 }

2.1.3 Maple

A grade: { 20, 21, 30, 31, 37, 38, 39, 40, 41, 42 }

B grade: { 17, 18, 19, 22, 23, 24, 25, 26, 27, 28, 29, 34, 35, 36 }

C grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 32 }

F grade: { 15,33 }

2.1.4 Maxima

A grade: { }

B grade: { }

C grade: { }

F grade: { 1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42 }

2.1.5 FriCAS

A grade: { 37,38,39,40,41,42 }

B grade: { 34,35,36 }

C grade: { }

F grade: { 1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33 }

2.1.6 Sympy

A grade: { }

B grade: { }

C grade: { 1,2,3,4,10,11 }

F grade: { 5,6,7,8,9,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42 }

2.1.7 Giac

A grade: { 34,37,38,41 }

B grade: { 42 }

C grade: { }

F grade: { 1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,35,36,39,40 }

2.1.8 Mupad

A grade: { }

B grade: { 34 }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 37, 38, 39, 40, 41, 42 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$

Problem 1	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	453	878	217	533	0	0	364	0	-1
normalized size	1	1.94	0.48	1.18	0.00	0.00	0.80	0.00	-0.00
time (sec)	N/A	0.509	0.266	0.030	0.000	0.874	9.014	0.000	0.000
Problem 2	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	367	706	159	403	0	0	262	0	-1
normalized size	1	1.92	0.43	1.10	0.00	0.00	0.71	0.00	-0.00
time (sec)	N/A	0.396	0.168	0.009	0.000	1.069	6.609	0.000	0.000
Problem 3	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	277	432	121	269	0	0	167	0	-1
normalized size	1	1.56	0.44	0.97	0.00	0.00	0.60	0.00	-0.00
time (sec)	N/A	0.225	0.097	0.008	0.000	1.007	4.510	0.000	0.000

Problem 4	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	226	226	77	169	0	0	78	0	-1
normalized size	1	1.00	0.34	0.75	0.00	0.00	0.35	0.00	-0.00
time (sec)	N/A	0.066	0.030	0.005	0.000	1.099	1.971	0.000	0.000

Problem 5	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	369	369	138	192	0	0	0	0	-1
normalized size	1	1.00	0.37	0.52	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.360	0.300	0.039	0.000	60.111	0.000	0.000	0.000

Problem 6	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	641	641	297	679	0	0	0	0	-1
normalized size	1	1.00	0.46	1.06	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.189	1.151	0.035	0.000	0.000	0.000	0.000	0.000

Problem 7	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	875	875	453	1591	0	0	0	0	-1
normalized size	1	1.00	0.52	1.82	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.784	2.133	0.036	0.000	0.000	0.000	0.000	0.000

Problem 8	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	912	912	222	588	0	0	0	0	-1
normalized size	1	1.00	0.24	0.64	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.726	0.281	0.041	0.000	1.142	0.000	0.000	0.000

Problem 9	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	694	694	166	458	0	0	0	0	-1
normalized size	1	1.00	0.24	0.66	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.532	0.172	0.010	0.000	1.401	0.000	0.000	0.000

Problem 10	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	395	395	126	320	0	0	167	0	-1
normalized size	1	1.00	0.32	0.81	0.00	0.00	0.42	0.00	-0.00
time (sec)	N/A	0.327	0.106	0.004	0.000	1.160	12.950	0.000	0.000

Problem 11	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	262	262	99	217	0	0	78	0	-1
normalized size	1	1.00	0.38	0.83	0.00	0.00	0.30	0.00	-0.00
time (sec)	N/A	0.096	0.044	0.005	0.000	0.914	7.186	0.000	0.000

Problem 12	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	732	732	432	564	0	0	0	0	-1
normalized size	1	1.00	0.59	0.77	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.781	0.791	0.034	0.000	0.000	0.000	0.000	0.000

Problem 13	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	1494	1494	427	1384	0	0	0	0	-1
normalized size	1	1.00	0.29	0.93	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	2.007	1.572	0.043	0.000	0.000	0.000	0.000	0.000

Problem 14	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2452	2452	630	2326	0	0	0	0	-1
normalized size	1	1.00	0.26	0.95	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	3.971	2.745	0.048	0.000	0.000	0.000	0.000	0.000

Problem 15	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-1)	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	169	169	0	0	0	0	0	0	-1
normalized size	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.219	0.376	0.105	0.000	1.207	0.000	0.000	0.000

Problem 16	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	94	81	0	0	0	0	-1
normalized size	1	1.00	1.96	1.69	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.029	0.127	0.027	0.000	1.097	0.000	0.000	0.000

Problem 17	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	755	755	4473	1708	0	0	0	0	-1
normalized size	1	1.00	5.92	2.26	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.357	6.477	0.019	0.000	0.898	0.000	0.000	0.000

Problem 18	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	528	528	674	1201	0	0	0	0	-1
normalized size	1	1.00	1.28	2.27	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.726	4.414	0.011	0.000	1.258	0.000	0.000	0.000

Problem 19	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	368	368	521	759	0	0	0	0	-1
normalized size	1	1.00	1.42	2.06	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.254	2.160	0.007	0.000	1.178	0.000	0.000	0.000

Problem 20	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	283	283	302	362	0	0	0	0	-1
normalized size	1	1.00	1.07	1.28	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.081	0.245	0.000	0.000	1.334	0.000	0.000	0.000

Problem 21	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	436	436	298	359	0	0	0	0	-1
normalized size	1	1.00	0.68	0.82	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.409	0.485	0.036	0.000	0.000	0.000	0.000	0.000

Problem 22	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	782	782	1853	1495	0	0	0	0	-1
normalized size	1	1.00	2.37	1.91	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.467	3.470	0.042	0.000	0.000	0.000	0.000	0.000

Problem 23	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	1125	1125	781	4476	0	0	0	0	-1
normalized size	1	1.00	0.69	3.98	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	3.594	5.732	0.048	0.000	0.000	0.000	0.000	0.000

Problem 24	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	859	859	5432	2445	0	0	0	0	-1
normalized size	1	1.00	6.32	2.85	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.380	6.653	0.037	0.000	1.558	0.000	0.000	0.000

Problem 25	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	628	633	766	1891	0	0	0	0	-1
normalized size	1	1.01	1.22	3.01	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.584	4.720	0.012	0.000	1.510	0.000	0.000	0.000

Problem 26	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F(-1)	B	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	481	481	0	1390	0	0	0	0	-1
normalized size	1	1.00	0.00	2.89	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.387	0.000	0.009	0.000	1.277	0.000	0.000	0.000

Problem 27	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F(-1)	B	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	398	398	0	931	0	0	0	0	-1
normalized size	1	1.00	0.00	2.34	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.195	0.000	0.006	0.000	1.310	0.000	0.000	0.000

Problem 28	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	867	1045	1736	3241	0	0	0	0	-1
normalized size	1	1.21	2.00	3.74	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.424	5.057	0.036	0.000	0.000	0.000	0.000	0.000

Problem 29	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	1301	2112	8031	8276	0	0	0	0	-1
normalized size	1	1.62	6.17	6.36	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	3.386	7.801	0.059	0.000	0.000	0.000	0.000	0.000

Problem 30	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F(-2)	F(-1)	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	273	273	310	365	0	0	0	0	-1
normalized size	1	1.00	1.14	1.34	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.136	0.504	0.102	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	271	271	312	369	0	0	0	0	-1
normalized size	1	1.00	1.15	1.36	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.187	0.427	0.081	0.000	0.000	0.000	0.000	0.000

Problem 32	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	74	93	0	0	0	0	-1
normalized size	1	1.00	0.70	0.88	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.105	0.182	0.034	0.000	1.580	0.000	0.000	0.000

Problem 33	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-1)	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	218	218	0	0	0	0	0	0	-1
normalized size	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.463	0.225	0.084	0.000	1.696	0.000	0.000	0.000

Problem 34	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	125	296	0	279	0	80	397
normalized size	1	1.00	1.18	2.79	0.00	2.63	0.00	0.75	3.75
time (sec)	N/A	0.129	0.201	0.030	0.000	1.634	0.000	0.652	1.845

Problem 35	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	145	441	238	0	603	0	0	-1
normalized size	1	1.00	3.04	1.64	0.00	4.16	0.00	0.00	-0.01
time (sec)	N/A	0.207	0.608	0.042	0.000	6.803	0.000	0.000	0.000

Problem 36	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	239	239	432	568	0	669	0	0	-1
normalized size	1	1.00	1.81	2.38	0.00	2.80	0.00	0.00	-0.00
time (sec)	N/A	0.184	0.685	0.108	0.000	6.822	0.000	0.000	0.000

Problem 37	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	309	309	214	442	0	469	0	360	-1
normalized size	1	1.00	0.69	1.43	0.00	1.52	0.00	1.17	-0.00
time (sec)	N/A	0.625	0.327	0.014	0.000	1.194	0.000	0.596	0.000

Problem 38	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	283	283	168	373	0	361	0	265	-1
normalized size	1	1.00	0.59	1.32	0.00	1.28	0.00	0.94	-0.00
time (sec)	N/A	0.341	0.223	0.010	0.000	1.156	0.000	0.642	0.000

Problem 39	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	286	286	176	251	0	743	0	0	-1
normalized size	1	1.00	0.62	0.88	0.00	2.60	0.00	0.00	-0.00
time (sec)	N/A	0.754	0.310	0.011	0.000	2.910	0.000	0.000	0.000

Problem 40	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	294	294	171	288	0	731	0	0	-1
normalized size	1	1.00	0.58	0.98	0.00	2.49	0.00	0.00	-0.00
time (sec)	N/A	0.767	0.251	0.010	0.000	2.440	0.000	0.000	0.000

Problem 41	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	288	288	177	329	0	749	0	370	-1
normalized size	1	1.00	0.61	1.14	0.00	2.60	0.00	1.28	-0.00
time (sec)	N/A	0.784	0.339	0.010	0.000	2.212	0.000	0.894	0.000

Problem 42	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	294	294	189	412	0	791	0	720	-1
normalized size	1	1.00	0.64	1.40	0.00	2.69	0.00	2.45	-0.00
time (sec)	N/A	0.787	0.287	0.013	0.000	2.049	0.000	1.339	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules**

column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [14] had the largest ratio of [.3929]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	15	5	1.94	28	0.179
2	A	12	5	1.92	28	0.179
3	A	8	5	1.56	26	0.192
4	A	3	3	1.00	19	0.158
5	A	3	3	1.00	28	0.107
6	A	6	6	1.00	28	0.214
7	A	7	6	1.00	28	0.214
8	A	12	7	1.00	28	0.250
9	A	10	6	1.00	28	0.214
10	A	7	5	1.00	26	0.192
11	A	4	4	1.00	19	0.210
12	A	9	7	1.00	28	0.250
13	A	15	10	1.00	28	0.357
14	A	22	11	1.00	28	0.393
15	A	6	3	1.00	26	0.115
16	A	2	2	1.00	27	0.074
17	A	6	4	1.00	33	0.121
18	A	5	4	1.00	33	0.121
19	A	4	4	1.00	31	0.129
20	A	3	3	1.00	24	0.125
21	A	3	3	1.00	33	0.091
22	A	6	6	1.00	33	0.182

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	7	6	1.00	33	0.182
24	A	5	5	1.00	33	0.152
25	A	4	4	1.01	33	0.121
26	A	4	4	1.00	31	0.129
27	A	4	4	1.00	24	0.167
28	A	9	7	1.21	33	0.212
29	A	15	10	1.62	33	0.303
30	A	1	1	1.00	41	0.024
31	A	1	1	1.00	41	0.024
32	A	4	4	1.00	31	0.129
33	A	6	3	1.00	31	0.097
34	A	11	7	1.00	28	0.250
35	A	4	3	1.00	30	0.100
36	A	1	1	1.00	30	0.033
37	A	6	6	1.00	38	0.158
38	A	6	6	1.00	37	0.162
39	A	8	7	1.00	40	0.175
40	A	8	7	1.00	40	0.175
41	A	8	7	1.00	40	0.175
42	A	8	7	1.00	40	0.175

Chapter 3

Listing of integrals

3.1
$$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$$

Optimal. Leaf size=453

$$(\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) (-63a^{3/2}\sqrt{c}e^2(Ae + 3Bd) + 25a^2Be^3 + 105\sqrt{a}c^{3/2}d^2(3Ae + Bd))$$

$$210\sqrt[4]{a}c^{9/4}\sqrt{a+cx^4}$$

[Out] $\frac{1}{21}e(21Acde - 5Bae^2 + 21Bcd^2)xx(c^4x+a)^{1/2}/c^2 + 1/5e^2(Ae + 3Bd)xx^3(c^4x+a)^{1/2}/c + 1/7Be^3x^5(c^4x+a)^{1/2}/c + 1/5(-3Aae^3 + 15Ac d^2e - 9Bade^2 + 5Bcd^3)xx(c^4x+a)^{1/2}/c^{3/2}/(a^{1/2} + x^2c^{1/2}) - 1/5a^{1/4}(-3Aae^3 + 15Ac d^2e - 9Bade^2 + 5Bcd^3)(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4})) * \text{EllipticE}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2, 2^{1/2}) * (a^{1/2} + x^2c^{1/2}) * ((c^4x+a)/(a^{1/2} + x^2c^{1/2}))^2)^{1/2}/c^{7/4}/(c^4x+a)^{1/2} + 1/210(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4})) * \text{EllipticF}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2, 2^{1/2}) * (105Ac^2d^3 + 25a^2Be^3 - 105acde(Ae+Bd) + 105c^{3/2}d^2(3Ae+Bd)a^{1/2} - 63a^{3/2})e^2(Ae+3Bd)c^{1/2})(a^{1/2} + x^2c^{1/2}) * ((c^4x+a)/(a^{1/2} + x^2c^{1/2}))^2)^{1/2}/a^{1/4}/c^{9/4}/(c^4x+a)^{1/2}$

Rubi [A] time = 0.51, antiderivative size = 878, normalized size of antiderivative = 1.94, number of steps used = 15, number of rules used = 5, integrand size = 28,

$\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {1721, 220, 305, 1196, 321}

$$\frac{Be^3\sqrt{cx^4+ax^5}}{7c} + \frac{e^2(3Bd+ Ae)\sqrt{cx^4+ax^3}}{5c} - \frac{5aBe^3\sqrt{cx^4+ax}}{21c^2} + \frac{de(Bd+ Ae)\sqrt{cx^4+ax}}{c} - \frac{3ae^2(3Bd+ Ae)\sqrt{cx^4+ax}}{5c^{3/2}(\sqrt{c}x^2 + \sqrt{a})}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + c*x^4], x]

[Out] (-5*a*B*e^3*x*Sqrt[a + c*x^4])/(21*c^2) + (d*e*(B*d + A*e)*x*Sqrt[a + c*x^4])/c + (e^2*(3*B*d + A*e)*x^3*Sqrt[a + c*x^4])/(5*c) + (B*e^3*x^5*Sqrt[a + c*x^4])/(7*c) - (3*a*e^2*(3*B*d + A*e)*x*Sqrt[a + c*x^4])/(5*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) + (d^2*(B*d + 3*A*e)*x*Sqrt[a + c*x^4])/(Sqrt[c]*(Sqrt[a] + Sqrt[c]*x^2)) + (3*a^(5/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(5*c^(7/4)*Sqrt[a + c*x^4]) - (a^(1/4)*d^2*(B*d + 3*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(3/4)*Sqrt[a + c*x^4]) + (A*d^3*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(1/4)*Sqrt[a + c*x^4]) + (5*a^(7/4)*B*e^3*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(42*c^(9/4)*Sqrt[a + c*x^4]) - (a^(3/4)*d*e*(B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(5/4)*Sqrt[a + c*x^4]) - (3*a^(5/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(10*c^(7/4)*Sqrt[a + c*x^4]) + (a^(1/4)*d^2*(B*d + 3*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(3/4)*Sqrt[a + c*x^4])

Rule 220

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2]]*EllipticF[2*ArcTan[q*x], 1/2])/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 305

Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Dist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 321


```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c^(
n - 1)*(c*x)^(m - n + 1)*(a + b*x^n)^(p + 1))/(b*(m + n*p + 1)), x] - Dist[
(a*c^n*(m - n + 1))/(b*(m + n*p + 1)), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 1196

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(
1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x],
1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0]] /; FreeQ[{a, c, d, e},
x] && PosQ[c/a]
```

Rule 1721

```
Int[(Px_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned} \int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx &= \int \left(\frac{Ad^3}{\sqrt{a + cx^4}} + \frac{d^2(Bd + 3Ae)x^2}{\sqrt{a + cx^4}} + \frac{3de(Bd + Ae)x^4}{\sqrt{a + cx^4}} + \frac{e^2(3Bd + Ae)x^6}{\sqrt{a + cx^4}} + \frac{Be^3x^8}{\sqrt{a + cx^4}} \right) dx \\ &= (Ad^3) \int \frac{1}{\sqrt{a + cx^4}} dx + (Be^3) \int \frac{x^8}{\sqrt{a + cx^4}} dx + (3de(Bd + Ae)) \int \frac{x^4}{\sqrt{a + cx^4}} dx \\ &= \frac{de(Bd + Ae)x\sqrt{a + cx^4}}{c} + \frac{e^2(3Bd + Ae)x^3\sqrt{a + cx^4}}{5c} + \frac{Be^3x^5\sqrt{a + cx^4}}{7c} + \frac{Ad^3}{c} \int \frac{1}{\sqrt{a + cx^4}} dx \\ &= -\frac{5aBe^3x\sqrt{a + cx^4}}{21c^2} + \frac{de(Bd + Ae)x\sqrt{a + cx^4}}{c} + \frac{e^2(3Bd + Ae)x^3\sqrt{a + cx^4}}{5c} + \frac{Be^3x^5\sqrt{a + cx^4}}{7c} + \frac{Ad^3}{c} \int \frac{1}{\sqrt{a + cx^4}} dx \\ &= -\frac{5aBe^3x\sqrt{a + cx^4}}{21c^2} + \frac{de(Bd + Ae)x\sqrt{a + cx^4}}{c} + \frac{e^2(3Bd + Ae)x^3\sqrt{a + cx^4}}{5c} + \frac{Be^3x^5\sqrt{a + cx^4}}{7c} + \frac{Ad^3}{c} \int \frac{1}{\sqrt{a + cx^4}} dx \end{aligned}$$

Mathematica [C] time = 0.27, size = 217, normalized size = 0.48

$$5x\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) (21Acd(cd^2 - ae^2) + aBe(5ae^2 - 21cd^2)) + ex(a + cx^4)(-25aBe^2 + 21Ace(5d + \dots))$$

$$105c^2\sqrt{a}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + c*x^4], x]

[Out] (e*x*(a + c*x^4)*(-25*a*B*e^2 + 21*A*c*e*(5*d + e*x^2) + 3*B*c*(35*d^2 + 21*d*e*x^2 + 5*e^2*x^4)) + 5*(21*A*c*d*(c*d^2 - a*e^2) + a*B*e*(-21*c*d^2 + 5*a*e^2))*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 7*c*(5*B*c*d^3 + 15*A*c*d^2*e - 9*a*B*d*e^2 - 3*a*A*e^3)*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)]/(105*c^2*Sqrt[a + c*x^4])

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{Be^3x^8 + (3Bde^2 + Ae^3)x^6 + 3(Bd^2e + Ade^2)x^4 + Ad^3 + (Bd^3 + 3Ad^2e)x^2}{\sqrt{cx^4 + a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2), x, algorithm="fricas")

[Out] integral((B*e^3*x^8 + (3*B*d*e^2 + A*e^3)*x^6 + 3*(B*d^2*e + A*d*e^2)*x^4 + A*d^3 + (B*d^3 + 3*A*d^2*e)*x^2)/sqrt(c*x^4 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2), x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^3/sqrt(c*x^4 + a), x)

maple [C] time = 0.03, size = 533, normalized size = 1.18

$$\frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} A d^3 \text{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a}} + \left(\frac{\sqrt{cx^4 + a} x^5}{7c} + \frac{5\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} a^2 \text{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x, i\right)}{21\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} c^2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x)`

[Out] $B*e^3*(1/7/c*x^5*(c*x^4+a)^{(1/2)}-5/21*a/c^2*x*(c*x^4+a)^{(1/2)}+5/21*a^2/c^2/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I))+(A*e^3+3*B*d*e^2)*(1/5/c*x^3*(c*x^4+a)^{(1/2)}-3/5*I*a^{(3/2)}/c^{(3/2)}/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*(EllipticF(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I)-EllipticE(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I)))+(3*A*d*e^2+3*B*d^2*e)*(1/3/c*x*(c*x^4+a)^{(1/2)}-1/3*a/c/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I))+I*(3*A*d^2*e+B*d^3)*a^{(1/2)}/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}/c^{(1/2)}*(EllipticF(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I)-EllipticE(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I))+A*d^3/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)*(e*x^2 + d)^3/sqrt(c*x^4 + a), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(1/2),x)`

[Out] `int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(1/2), x)`

sympy [C] time = 9.01, size = 364, normalized size = 0.80

$$\frac{Ad^3x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{5}{4}\right)} + \frac{3Ad^2ex^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{7}{4}\right)} + \frac{3Ade^2x^5\Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{9}{4}\right)} + \frac{Ae^3x^7\Gamma\left(\frac{7}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{7}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{11}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+a)**(1/2),x)

[Out] A*d**3*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + 3*A*d**2*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + 3*A*d*e**2*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4)) + A*e**3*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(11/4)) + B*d**3*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + 3*B*d**2*e*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4)) + 3*B*d*e**2*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(11/4)) + B*e**3*x**9*gamma(9/4)*hyper((1/2, 9/4), (13/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(13/4))

$$3.2 \quad \int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$$

Optimal. Leaf size=367

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) (-9a^{3/2}Be^2 + 15\sqrt{a}cd(2Ae + Bd) - 5a\sqrt{c}e(Ae + 2Bd) + 15Ac^3)}{30\sqrt[4]{a}c^{7/4}\sqrt{a+cx^4}}$$

[Out] $1/3*e*(A*e+2*B*d)*x*(c*x^4+a)^{(1/2)}/c+1/5*B*e^2*x^3*(c*x^4+a)^{(1/2)}/c+1/5*(10*A*c*d*e-3*B*a*e^2+5*B*c*d^2)*x*(c*x^4+a)^{(1/2)}/c^{(3/2)}/(a^{(1/2)}+x^2*c^{(1/2)})-1/5*a^{(1/4)}*(10*A*c*d*e-3*B*a*e^2+5*B*c*d^2)*(cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))*EllipticE(sin(2*arctan(c^{(1/4)}*x/a^{(1/4)})),1/2)*2^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/c^{(7/4)}/(c*x^4+a)^{(1/2)}+1/30*(cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))*EllipticF(sin(2*arctan(c^{(1/4)}*x/a^{(1/4)})),1/2)*2^{(1/2)}*(15*A*c^{(3/2)}*d^2-9*a^{(3/2)}*B*e^2+15*c*d*(2*A*e+B*d)*a^{(1/2)}-5*a*e*(A*e+2*B*d)*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(1/4)}/c^{(7/4)}/(c*x^4+a)^{(1/2)}$

Rubi [A] time = 0.40, antiderivative size = 706, normalized size of antiderivative = 1.92, number of steps used = 12, number of rules used = 5, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {1721, 220, 305, 1196, 321}

$$\frac{a^{3/4}e(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} (Ae + 2Bd)F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{6c^{5/4}\sqrt{a+cx^4}} - \frac{3a^{5/4}Be^2(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{10c^{7/4}\sqrt{a+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + c*x^4],x]

[Out] $(e*(2*B*d + A*e)*x*\text{Sqrt}[a + c*x^4])/(3*c) + (B*e^2*x^3*\text{Sqrt}[a + c*x^4])/(5*c) - (3*a*B*e^2*x*\text{Sqrt}[a + c*x^4])/(5*c^{(3/2)}*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) + (d*(B*d + 2*A*e)*x*\text{Sqrt}[a + c*x^4])/(\text{Sqrt}[c]*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) + (3*a^{(5/4)}*B*e^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)]^2)*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2]/(5*c^{(7/4)}*\text{Sqrt}[a + c*x^4]) - (a^{(1/4)}*d*(B*d + 2*A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)]^2)*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2]/(c^{(3/4)}*\text{Sqrt}[a + c*x^4]) + (A*d^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)]^2)*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2]/(2*a^{(1/4)}*c^{(1/4)}*\text{Sqrt}[a + c*x^4]) - (3*a^{(5/4)}*B*e^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)$

```
*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)
/a^(1/4)], 1/2]]/(10*c^(7/4)*Sqrt[a + c*x^4]) - (a^(3/4)*e*(2*B*d + A*e)*(S
qrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF
[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]]/(6*c^(5/4)*Sqrt[a + c*x^4]) + (a^(1/4)
)*d*(B*d + 2*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[
c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]]/(2*c^(3/4)*Sqrt[a
+ c*x^4])
```

Rule 220

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x]
, 1/2]]/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 305

```
Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, D
ist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a +
b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 321

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c^(
n - 1)*(c*x)^(m - n + 1)*(a + b*x^n)^(p + 1))/(b*(m + n*p + 1)), x] - Dist[
(a*c^n*(m - n + 1))/(b*(m + n*p + 1)), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 1196

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(
1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x],
1/2]]/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e},
x] && PosQ[c/a]
```

Rule 1721

```
Int[(Px_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx &= \int \left(\frac{Ad^2}{\sqrt{a + cx^4}} + \frac{d(Bd + 2Ae)x^2}{\sqrt{a + cx^4}} + \frac{e(2Bd + Ae)x^4}{\sqrt{a + cx^4}} + \frac{Be^2x^6}{\sqrt{a + cx^4}} \right) dx \\
&= (Ad^2) \int \frac{1}{\sqrt{a + cx^4}} dx + (Be^2) \int \frac{x^6}{\sqrt{a + cx^4}} dx + (e(2Bd + Ae)) \int \frac{x^4}{\sqrt{a + cx^4}} dx \\
&= \frac{e(2Bd + Ae)x\sqrt{a + cx^4}}{3c} + \frac{Be^2x^3\sqrt{a + cx^4}}{5c} + \frac{Ad^2(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(\frac{4\sqrt{a}d(Bd + 2Ae)x\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)}\right)}{2^4\sqrt{a}^4\sqrt{c}\sqrt{a + cx^4}} \\
&= \frac{e(2Bd + Ae)x\sqrt{a + cx^4}}{3c} + \frac{Be^2x^3\sqrt{a + cx^4}}{5c} + \frac{d(Bd + 2Ae)x\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} - \frac{4\sqrt{a}d(Bd + 2Ae)x\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} \\
&= \frac{e(2Bd + Ae)x\sqrt{a + cx^4}}{3c} + \frac{Be^2x^3\sqrt{a + cx^4}}{5c} - \frac{3aBe^2x\sqrt{a + cx^4}}{5c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} + \frac{d(Bd + 2Ae)x\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)}
\end{aligned}$$

Mathematica [C] time = 0.17, size = 159, normalized size = 0.43

$$\frac{-5x\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) (aAe^2 + 2aBde - 3Acd^2) + x^3\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{7}{4}; -\frac{cx^4}{a}\right) (-3aBe^2 + 10Acde + 5Ae^2)}{15c\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + c*x^4], x]

[Out] (e*x*(10*B*d + 5*A*e + 3*B*e*x^2)*(a + c*x^4) - 5*(-3*A*c*d^2 + 2*a*B*d*e + a*A*e^2)*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + (5*B*c*d^2 + 10*A*c*d*e - 3*a*B*e^2)*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)])/(15*c*Sqrt[a + c*x^4])

fricas [F] time = 1.07, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{Be^2x^6 + (2Bde + Ae^2)x^4 + Ad^2 + (Bd^2 + 2Ade)x^2}{\sqrt{cx^4 + a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2), x, algorithm="fricas")

[Out] integral((B*e^2*x^6 + (2*B*d*e + A*e^2)*x^4 + A*d^2 + (B*d^2 + 2*A*d*e)*x^2)/sqrt(c*x^4 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^2/sqrt(c*x^4 + a), x)

maple [C] time = 0.01, size = 403, normalized size = 1.10

$$\frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} A d^2 \operatorname{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a}} + \left(\frac{\sqrt{cx^4 + a} x^3}{5c} - \frac{3i\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \left(-\operatorname{EllipticE}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x, i\right)\right)}{5\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x)

[Out] B*e^2*(1/5*(c*x^4+a)^(1/2)/c*x^3-3/5*I*a^(3/2)/c^(3/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I)))+(A*e^2+2*B*d*e)*(1/3*(c*x^4+a)^(1/2)/c*x-1/3*a/c/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I))+I*(2*A*d*e+B*d^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I))+A*d^2/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^2/sqrt(c*x^4 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(1/2), x)

[Out] int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(1/2), x)

sympy [C] time = 6.61, size = 262, normalized size = 0.71

$$\frac{Ad^2x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{5}{4}\right)} + \frac{Adex^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{2\sqrt{a}\Gamma\left(\frac{7}{4}\right)} + \frac{Ae^2x^5\Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{9}{4}\right)} + \frac{Bd^2x^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{7}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+a)**(1/2), x)

[Out] A*d**2*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + A*d*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(2*sqrt(a)*gamma(7/4)) + A*e**2*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4)) + B*d**2*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*d*e*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(2*sqrt(a)*gamma(9/4)) + B*e**2*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(11/4))

$$3.3 \quad \int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+cx^4}} dx$$

Optimal. Leaf size=277

$$\frac{\sqrt[4]{a} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) \left(\frac{3Ac d - aBe}{\sqrt{a}} + 3\sqrt{c}(Ae + Bd)\right)}{6c^{5/4}\sqrt{a+cx^4}} - \frac{\sqrt[4]{a} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{c^{3/4}\sqrt{a+cx^4}}$$

[Out] $\frac{1}{3} B e x x (c x^4 + a)^{1/2} / c + (A e + B d) x x (c x^4 + a)^{1/2} / c^{1/2} / (a^{1/2} + x^2 c^{1/2}) - a^{1/4} (A e + B d) (\cos(2 \arctan(c^{1/4} x / a^{1/4}))^2)^{1/2} / \cos(2 \arctan(c^{1/4} x / a^{1/4})) * \text{EllipticE}(\sin(2 \arctan(c^{1/4} x / a^{1/4})), 1/2) * 2^{1/2} * (a^{1/2} + x^2 c^{1/2}) * ((c x^4 + a) / (a^{1/2} + x^2 c^{1/2}))^{1/2} / c^{3/4} / (c x^4 + a)^{1/2} + 1/6 a^{1/4} (\cos(2 \arctan(c^{1/4} x / a^{1/4}))^2)^{1/2} / \cos(2 \arctan(c^{1/4} x / a^{1/4})) * \text{EllipticF}(\sin(2 \arctan(c^{1/4} x / a^{1/4})), 1/2) * 2^{1/2} * ((3 A c d - B a e) / a^{1/2} + 3 (A e + B d) c^{1/2}) * (a^{1/2} + x^2 c^{1/2}) * ((c x^4 + a) / (a^{1/2} + x^2 c^{1/2}))^{1/2} / c^{5/4} / (c x^4 + a)^{1/2}$

Rubi [A] time = 0.22, antiderivative size = 432, normalized size of antiderivative = 1.56, number of steps used = 8, number of rules used = 5, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {1721, 220, 305, 1196, 321}

$$\frac{a^{3/4} B e (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{6c^{5/4}\sqrt{a+cx^4}} + \frac{\sqrt[4]{a} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} (Ae + Bd) F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2c^{3/4}\sqrt{a+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2))/Sqrt[a + c*x^4], x]

[Out] $\frac{B e x x \text{Sqrt}[a + c x^4]}{3 c} + \frac{(B d + A e) x x \text{Sqrt}[a + c x^4]}{(\text{Sqrt}[c] * (\text{Sqrt}[a] + \text{Sqrt}[c] x^2))} - \frac{a^{1/4} (B d + A e) (\text{Sqrt}[a] + \text{Sqrt}[c] x^2) \text{Sqrt}[(a + c x^4) / (\text{Sqrt}[a] + \text{Sqrt}[c] x^2)^2] * \text{EllipticE}[2 * \text{ArcTan}[(c^{1/4} x) / a^{1/4}], 1/2]}{(c^{3/4} * \text{Sqrt}[a + c x^4])} + \frac{A d * (\text{Sqrt}[a] + \text{Sqrt}[c] x^2) \text{Sqrt}[(a + c x^4) / (\text{Sqrt}[a] + \text{Sqrt}[c] x^2)^2] * \text{EllipticF}[2 * \text{ArcTan}[(c^{1/4} x) / a^{1/4}], 1/2]}{(2 * a^{1/4} * c^{1/4} * \text{Sqrt}[a + c x^4])} - \frac{a^{3/4} B e * (\text{Sqrt}[a] + \text{Sqrt}[c] x^2) \text{Sqrt}[(a + c x^4) / (\text{Sqrt}[a] + \text{Sqrt}[c] x^2)^2] * \text{EllipticF}[2 * \text{ArcTan}[(c^{1/4} x) / a^{1/4}], 1/2]}{(6 * c^{5/4} * \text{Sqrt}[a + c x^4])} + \frac{a^{1/4} (B d + A e) * (\text{Sqrt}[a] + \text{Sqrt}[c] x^2) \text{Sqrt}[(a + c x^4) / (\text{Sqrt}[a] + \text{Sqrt}[c] x^2)^2] * \text{EllipticF}[2 * \text{ArcTan}[(c^{1/4} x) / a^{1/4}], 1/2]}{(2 * c^{3/4} * \text{Sqrt}[a + c x^4])}$

Rule 220

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2]]/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 305

Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Dist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 321

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c^(n - 1)*(c*x)^(m - n + 1)*(a + b*x^n)^(p + 1))/(b*(m + n*p + 1)), x] - Dist[(a*c^n*(m - n + 1))/(b*(m + n*p + 1)), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 1196

Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]

Rule 1721

Int[(Px_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx &= \int \left(\frac{Ad}{\sqrt{a + cx^4}} + \frac{(Bd + Ae)x^2}{\sqrt{a + cx^4}} + \frac{Bex^4}{\sqrt{a + cx^4}} \right) dx \\
&= (Ad) \int \frac{1}{\sqrt{a + cx^4}} dx + (Be) \int \frac{x^4}{\sqrt{a + cx^4}} dx + (Bd + Ae) \int \frac{x^2}{\sqrt{a + cx^4}} dx \\
&= \frac{Bex\sqrt{a + cx^4}}{3c} + \frac{Ad(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a + cx^4}} - \frac{(aBe) \int \frac{1}{\sqrt{a + cx^4}} dx}{3c} \\
&= \frac{Bex\sqrt{a + cx^4}}{3c} + \frac{(Bd + Ae)x\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}(Bd + Ae)(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}}}{c^{3/4}\sqrt{a + cx^4}}
\end{aligned}$$

Mathematica [C] time = 0.10, size = 121, normalized size = 0.44

$$\frac{x\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) (3Acd - aBe) + cx^3\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{7}{4}; -\frac{cx^4}{a}\right) (Ae + Bd) + Bex(a + cx^4)}{3c\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2))/Sqrt[a + c*x^4], x]

[Out] (B*e*x*(a + c*x^4) + (3*A*c*d - a*B*e)*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + c*(B*d + A*e)*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)])/(3*c*Sqrt[a + c*x^4])

fricas [F] time = 1.01, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{Bex^4 + (Bd + Ae)x^2 + Ad}{\sqrt{cx^4 + a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2), x, algorithm="fricas")

[Out] integral((B*e*x^4 + (B*d + A*e)*x^2 + A*d)/sqrt(c*x^4 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)/sqrt(c*x^4 + a), x)

maple [C] time = 0.01, size = 269, normalized size = 0.97

$$\frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1}\sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1}Ad\text{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x,i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}+\left(-\frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1}\sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1}a\text{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x,i\right)}{3\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}c}\right)+$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x)

[Out] B*e*(1/3*(c*x^4+a)^(1/2)/c*x-1/3*a/c/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I))+I*(A*e+B*d)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I))+A*d/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)/sqrt(c*x^4 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(1/2),x)

[Out] `int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(1/2), x)`

sympy [C] time = 4.51, size = 167, normalized size = 0.60

$$\frac{A dx \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{5}{4}\right)} + \frac{A e x^3 \Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{7}{4}\right)} + \frac{B dx^3 \Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{7}{4}\right)} + \frac{B e x^5 \Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{9}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+a)**(1/2), x)`

[Out] `A*d*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + A*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*d*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*e*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4))`

3.4 $\int \frac{A+Bx^2}{\sqrt{a+cx^4}} dx$

Optimal. Leaf size=226

$$\frac{\sqrt[4]{a} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \left(\frac{A\sqrt{c}}{\sqrt{a}} + B \right) F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) + \sqrt[4]{a} B (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2c^{3/4}\sqrt{a+cx^4}}$$

[Out] $B*x*(c*x^4+a)^{(1/2)}/c^{(1/2)}/(a^{(1/2)}+x^2*c^{(1/2)})-a^{(1/4)}*B*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(3/4)}/(c*x^4+a)^{(1/2)}+1/2*a^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(B+A*c^{(1/2)}/a^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(3/4)}/(c*x^4+a)^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 226, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {1198, 220, 1196}

$$\frac{\sqrt[4]{a} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \left(\frac{A\sqrt{c}}{\sqrt{a}} + B \right) F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) + \sqrt[4]{a} B (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2c^{3/4}\sqrt{a+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/Sqrt[a + c*x^4], x]

[Out] $(B*x*\text{Sqrt}[a + c*x^4])/(\text{Sqrt}[c]*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (a^{(1/4)}*B*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/c^{(3/4)}*\text{Sqrt}[a + c*x^4] + (a^{(1/4)}*(B + (A*\text{Sqrt}[c])/ \text{Sqrt}[a])*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (2*c^{(3/4)}*\text{Sqrt}[a + c*x^4])$

Rule 220

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] :> With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2]]*EllipticF[2*ArcTan[q*x], 1/2])/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 1196

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(
  1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x],
  1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0]] /; FreeQ[{a, c, d, e},
  x] && PosQ[c/a]
```

Rule 1198

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
  nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0]] /; FreeQ[{a, c,
  d, e}, x] && PosQ[c/a]
```

Rubi steps

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx = \left(A + \frac{\sqrt{a}B}{\sqrt{c}} \right) \int \frac{1}{\sqrt{a + cx^4}} dx - \frac{(\sqrt{a}B) \int \frac{1 - \frac{\sqrt{c}x^2}{\sqrt{a}}}{\sqrt{a + cx^4}} dx}{\sqrt{c}}$$

$$= \frac{Bx\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}B(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) (\sqrt{a}B + A\sqrt{c})}{c^{3/4}\sqrt{a + cx^4}} + \dots$$

Mathematica [C] time = 0.03, size = 77, normalized size = 0.34

$$\frac{\sqrt{\frac{cx^4}{a} + 1} \left(3Ax {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) + Bx^3 {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{7}{4}; -\frac{cx^4}{a}\right) \right)}{3\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(A + B*x^2)/Sqrt[a + c*x^4], x]
```

```
[Out] (Sqrt[1 + (c*x^4)/a]*(3*A*x*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)]
+ B*x^3*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)]))/(3*Sqrt[a + c*x^4]
)
```

fricas [F] time = 1.10, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{Bx^2 + A}{\sqrt{cx^4 + a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="fricas")

[Out] integral((B*x^2 + A)/sqrt(c*x^4 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/sqrt(c*x^4 + a), x)

maple [C] time = 0.00, size = 169, normalized size = 0.75

$$\frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} A \operatorname{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right) + i\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \left(-\operatorname{EllipticE}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right) + \operatorname{EllipticE}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\right)\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} \sqrt{c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(c*x^4+a)^(1/2),x)

[Out] I*B*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x, I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x, I))+A/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x, I)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/sqrt(c*x^4 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((A + B*x^2)/(a + c*x^4)^(1/2), x)`

[Out] `int((A + B*x^2)/(a + c*x^4)^(1/2), x)`

sympy [C] time = 1.97, size = 78, normalized size = 0.35

$$\frac{Ax\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{5}{4}\right)} + \frac{Bx^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{7}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x**2+A)/(c*x**4+a)**(1/2), x)`

[Out] `A*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + B*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4))`

$$3.5 \quad \int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx$$

Optimal. Leaf size=369

$$\frac{a^{3/4} (\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \left(\frac{\sqrt{cd}}{\sqrt{a}} + e\right)^2 (Bd - Ae) \Pi\left(-\frac{(\sqrt{cd}-\sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right)\middle|\frac{1}{2}\right) (Bd - Ae) \tan^{-1}\left(\frac{x\sqrt{a}}{\sqrt{d}}\right)}{4\sqrt[4]{c}de\sqrt{a+cx^4} (cd^2 - ae^2)} \quad 2\sqrt{d}\sqrt{e}\sqrt{ae^2 +}$$

[Out] $-1/2*(-A*e+B*d)*\arctan(x*(a*e^2+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+a)^{(1/2)})/d^{(1/2)}/e^{(1/2)}/(a*e^2+c*d^2)^{(1/2)}-1/2*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(B*a^{(1/2)}-A*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}+1/4*a^{(3/4)}*(-A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(e+d*c^{(1/2)}/a^{(1/2)})^2*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/c^{(1/4)}/d/e/(-a*e^2+c*d^2)/(c*x^4+a)^{(1/2)}$

Rubi [A] time = 0.36, antiderivative size = 369, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {1709, 220, 1707}

$$\frac{a^{3/4} (\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \left(\frac{\sqrt{cd}}{\sqrt{a}} + e\right)^2 (Bd - Ae) \Pi\left(-\frac{(\sqrt{cd}-\sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right)\middle|\frac{1}{2}\right) (Bd - Ae) \tan^{-1}\left(\frac{x\sqrt{a}}{\sqrt{d}}\right)}{4\sqrt[4]{c}de\sqrt{a+cx^4} (cd^2 - ae^2)} \quad 2\sqrt{d}\sqrt{e}\sqrt{ae^2 +}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]),x]

[Out] $-((B*d - A*e)*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + c*x^4])])/(2*\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[c*d^2 + a*e^2]) - ((\text{Sqrt}[a]*B - A*\text{Sqrt}[c])*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/(2*a^{(1/4)}*c^{(1/4)}*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*\text{Sqrt}[a + c*x^4]) + (a^{(3/4)}*((\text{Sqrt}[c]*d)/\text{Sqrt}[a] + e)^2*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(4*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/(4*c^{(1/4)}*d*e*(c*d^2 - a*e^2)*\text{Sqrt}[a + c*x^4])$

Rule 220

`Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2]]/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

Rule 1707

`Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[(Rt[(c*d)/e + (a*e)/d, 2]*x)/Sqrt[a + c*x^4]])/(2*d*e*Rt[(c*d)/e + (a*e)/d, 2]), x] + Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a + c*x^4))/(a*(A + B*x^2)^2)]*EllipticPi[Cancel[-((B*d - A*e)^2/(4*d*e*A*B))], 2*ArcTan[q*x], 1/2])/(4*d*e*A*q*Sqrt[a + c*x^4]), x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]`

Rule 1709

`Int[((A_.) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4], x], x] + Dist[(a*(B*d - A*e)*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x], x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]`

Rubi steps

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + cx^4}} dx = -\frac{(\sqrt{a}B - A\sqrt{c}) \int \frac{1}{\sqrt{a+cx^4}} dx}{\sqrt{c}d - \sqrt{a}e} + \frac{(\sqrt{a}(Bd - Ae)) \int \frac{1 + \frac{\sqrt{c}x^2}{\sqrt{a}}}{(d+ex^2)\sqrt{a+cx^4}} dx}{\sqrt{c}d - \sqrt{a}e}$$

$$= -\frac{(Bd - Ae) \tan^{-1}\left(\frac{\sqrt{cd^2+ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+cx^4}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{cd^2+ae^2}} - \frac{(\sqrt{a}B - A\sqrt{c})(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\right)}{2^4 \sqrt{a}^4 \sqrt{c} (\sqrt{c}d - \sqrt{a}e) \sqrt{a + cx^4}}$$

Mathematica [C] time = 0.30, size = 138, normalized size = 0.37

$$\frac{i\sqrt{\frac{cx^4}{a} + 1} \left((Ae - Bd) \Pi\left(-\frac{i\sqrt{ae}}{\sqrt{cd}}; i \sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x\right)\right) - 1 \right) + Bd F\left(i \sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x\right)\right) - 1 \right)}{de \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]),x]

[Out] ((-I)*Sqrt[1 + (c*x^4)/a]*(B*d*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + (-B*d) + A*e)*EllipticPi[((-I)*Sqrt[a]*e)/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1))/((Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*e*Sqrt[a + c*x^4])

fricas [F] time = 60.11, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{cx^4 + a} (Bx^2 + A)}{cex^6 + cdx^4 + aex^2 + ad}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*x^4 + a)*(B*x^2 + A)/(c*e*x^6 + c*d*x^4 + a*e*x^2 + a*d), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a} (ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)), x)

maple [C] time = 0.04, size = 192, normalized size = 0.52

$$\frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} B \text{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right) (Ae - Bd) \sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \text{EllipticPi}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} e + \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} de}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x)

[Out] B/e/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x, I

$$\frac{(A* e - B*d)/e/d/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(-I/a^{(1/2)}*c^{(1/2)}*x^2+1)^{(1/2)}*(I/a^{(1/2)}*c^{(1/2)}*x^2+1)^{(1/2)}}{(c*x^4+a)^{(1/2)}*EllipticPi((I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*x, I*a^{(1/2)}/c^{(1/2)}*e/d, (-I/a^{(1/2)}*c^{(1/2)})^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)})}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2), x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)), x)

[Out] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}(d + ex^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+a)**(1/2), x)

[Out] Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)), x)

$$3.6 \quad \int \frac{A+Bx^2}{(d+ex^2)^2 \sqrt{a+cx^4}} dx$$

Optimal. Leaf size=641

$$\frac{\sqrt{c}x\sqrt{a+cx^4}(Bd-Ae)}{2d(\sqrt{a}+\sqrt{c}x^2)(ae^2+cd^2)} - \frac{ex\sqrt{a+cx^4}(Bd-Ae)}{2d(d+ex^2)(ae^2+cd^2)} - \frac{\sqrt[4]{a}\sqrt[4]{c}(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}(Bd-Ae)E\left(2\tan^{-1}\right)}{2d\sqrt{a+cx^4}(ae^2+cd^2)}$$

[Out] $-1/4*(-A*a*e^3-3*A*c*d^2*e-B*a*d*e^2+B*c*d^3)*\arctan(x*(a*e^2+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+a)^{(1/2)})/d^{(3/2)}/(a*e^2+c*d^2)^{(3/2)}/e^{(1/2)}-1/2*e*(-A*e+B*d)*x*(c*x^4+a)^{(1/2)}/d/(a*e^2+c*d^2)/(e*x^2+d)+1/2*(-A*e+B*d)*x*c^{(1/2)}*(c*x^4+a)^{(1/2)}/d/(a*e^2+c*d^2)/(a^{(1/2)}+x^2*c^{(1/2)})-1/2*a^{(1/4)}*c^{(1/4)}*(-A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/d/(a*e^2+c*d^2)/(c*x^4+a)^{(1/2)}+1/2*A*c^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/d/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}+1/8*(-A*a*e^3-3*A*c*d^2*e-B*a*d*e^2+B*c*d^3)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),-1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*2^{(1/2)})*(e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^2/e/(a*e^2+c*d^2)/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}$

Rubi [A] time = 1.19, antiderivative size = 641, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1697, 1715, 1196, 1709, 220, 1707}

$$\frac{\sqrt{c}x\sqrt{a+cx^4}(Bd-Ae)}{2d(\sqrt{a}+\sqrt{c}x^2)(ae^2+cd^2)} - \frac{ex\sqrt{a+cx^4}(Bd-Ae)}{2d(d+ex^2)(ae^2+cd^2)} - \frac{(-aAe^3 - aBde^2 - 3Acd^2e + Bcd^3)\tan^{-1}\left(\frac{x\sqrt{ae^2+cd^2}}{\sqrt{d}\sqrt{e}\sqrt{a+cx^4}}\right)}{4d^{3/2}\sqrt{e}(ae^2+cd^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + c*x^4]),x]

[Out] $(\text{Sqrt}[c]*(B*d - A*e)*x*\text{Sqrt}[a + c*x^4])/((2*d*(c*d^2 + a*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (e*(B*d - A*e)*x*\text{Sqrt}[a + c*x^4])/((2*d*(c*d^2 + a*e^2)*(d + e*x^2)) - ((B*c*d^3 - 3*A*c*d^2*e - a*B*d*e^2 - a*A*e^3)*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + c*x^4])]))/(4*d^{(3/2)}*\text{Sqrt}[e]*(c*d^2 +$

$$a^2 e^2)^{3/2} - (a^{1/4} c^{1/4} (Bd - A^2 e) (\sqrt{a} + \sqrt{c} x^2) \sqrt{(a + c x^4) / (\sqrt{a} + \sqrt{c} x^2)^2}) \text{EllipticE}[2 \text{ArcTan}[c^{1/4} x / a^{1/4}], 1/2]) / (2 d (c d^2 + a^2 e^2) \sqrt{a + c x^4}) + (A^2 c^{1/4} (\sqrt{a} + \sqrt{c} x^2) \sqrt{(a + c x^4) / (\sqrt{a} + \sqrt{c} x^2)^2}) \text{EllipticF}[2 \text{ArcTan}[c^{1/4} x / a^{1/4}], 1/2]) / (2 a^{1/4} d (\sqrt{c} d - \sqrt{a} e) \sqrt{a + c x^4}) + ((\sqrt{c} d + \sqrt{a} e) (B^2 c d^3 - 3 A^2 c d^2 e - a B^2 d e^2 - a^2 A e^3) (\sqrt{a} + \sqrt{c} x^2) \sqrt{(a + c x^4) / (\sqrt{a} + \sqrt{c} x^2)^2}) \text{EllipticPi}[-(\sqrt{c} d - \sqrt{a} e)^2 / (4 \sqrt{a} \sqrt{c} d e), 2 \text{ArcTan}[c^{1/4} x / a^{1/4}], 1/2]) / (8 a^{1/4} c^{1/4} d^2 e (\sqrt{c} d - \sqrt{a} e) (c d^2 + a^2 e^2) \sqrt{a + c x^4})$$

Rule 220

$$\text{Int}[1/\sqrt{(a_1) + (b_1)(x_1)^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2 x^2) \sqrt{(a + b x^4) / (a(1 + q^2 x^2)^2)}] \text{EllipticF}[2 \text{ArcTan}[q x], 1/2]) / (2 q \sqrt{a + b x^4}), x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$$

Rule 1196

$$\text{Int}(((d_1) + (e_1)(x_1)^2) / \sqrt{(a_1) + (c_1)(x_1)^4}, x_Symbol) \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, -\text{Simp}[(d x \sqrt{a + c x^4}) / (a(1 + q^2 x^2)), x] + \text{Simp}[(d(1 + q^2 x^2) \sqrt{(a + c x^4) / (a(1 + q^2 x^2)^2)}] \text{EllipticE}[2 \text{ArcTan}[q x], 1/2]) / (q \sqrt{a + c x^4}), x] /; \text{EqQ}[e + d q^2, 0] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$$

Rule 1697

$$\text{Int}(((P4x_1) * ((d_1) + (e_1)(x_1)^2)^{(q_1)}) / \sqrt{(a_1) + (c_1)(x_1)^4}, x_Symbol) \rightarrow \text{With}[\{A = \text{Coeff}[P4x, x, 0], B = \text{Coeff}[P4x, x, 2], C = \text{Coeff}[P4x, x, 4]\}, -\text{Simp}[(C d^2 - B d e + A^2 e^2) x (d + e x^2)^{(q+1)} \sqrt{a + c x^4}) / (2 d (q+1) (c d^2 + a^2 e^2)), x] + \text{Dist}[1 / (2 d (q+1) (c d^2 + a^2 e^2)), \text{Int}[(d + e x^2)^{(q+1)} \text{Simp}[a d (C d - B e) + A (a^2 e^2 (2 q + 3) + 2 c d^2 (q+1)) + 2 d (B c d - A^2 c e + a^2 C e) (q+1) x^2 + c (C d^2 - B d e + A^2 e^2) (2 q + 5) x^4, x] / \sqrt{a + c x^4}, x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{PolyQ}[P4x, x^2] \ \&\& \ \text{LeQ}[\text{Expon}[P4x, x], 4] \ \&\& \ \text{NeQ}[c d^2 + a^2 e^2, 0] \ \&\& \ \text{ILtQ}[q, -1]$$

Rule 1707

$$\text{Int}(((A_1) + (B_1)(x_1)^2) / (((d_1) + (e_1)(x_1)^2) \sqrt{(a_1) + (c_1)(x_1)^4}), x_Symbol) \rightarrow \text{With}[\{q = \text{Rt}[B/A, 2]\}, -\text{Simp}[(B d - A^2 e) \text{ArcTan}[(\text{Rt}[(c d) / e + (a e) / d, 2] x) / \sqrt{a + c x^4}]] / (2 d e \text{Rt}[(c d) / e + (a e) / d, 2]), x] + \text{Simp}[(B d + A^2 e) (A + B x^2) \sqrt{(A^2 (a + c x^4) / (a (A + B x^2)^2))} \text{EllipticPi}[\text{Cancel}[-(B d - A^2 e)^2 / (4 d e A B)], 2 \text{ArcTan}[q x], 1/2]) / (4 d e A q \sqrt{a + c x^4}), x] /; \text{FreeQ}[\{a, c, d, e, A, B\}, x] \ \&\& \ \text{NeQ}[c d^2 + a^2 e^2, 0]$$

$\wedge 2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a] \&\& \text{EqQ}[c*A^2 - a*B^2, 0]$

Rule 1709

$\text{Int}[\frac{(A_.) + (B_.)*(x_)^2}{((d_) + (e_.)*(x_)^2)*\text{Sqrt}[(a_) + (c_.)*(x_)^4]}, x_Symbol] :> \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2), \text{Int}[1/\text{Sqrt}[a + c*x^4], x], x] + \text{Dist}[(a*(B*d - A*e)*(e + d*q))/(c*d^2 - a*e^2), \text{Int}[(1 + q*x^2)/((d + e*x^2)*\text{Sqrt}[a + c*x^4]), x], x]] /;$ $\text{FreeQ}[\{a, c, d, e, A, B\}, x] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a] \&\& \text{NeQ}[c*A^2 - a*B^2, 0]$

Rule 1715

$\text{Int}[(P4x)/(((d_) + (e_.)*(x_)^2)*\text{Sqrt}[(a_) + (c_.)*(x_)^4]), x_Symbol] :> \text{With}[\{q = \text{Rt}[c/a, 2], A = \text{Coeff}[P4x, x, 0], B = \text{Coeff}[P4x, x, 2], C = \text{Coeff}[P4x, x, 4]\}, -\text{Dist}[C/(e*q), \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + c*x^4], x], x] + \text{Dist}[1/(c*e), \text{Int}[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2]/((d + e*x^2)*\text{Sqrt}[a + c*x^4]), x], x]] /;$ $\text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PolyQ}[P4x, x^2, 2] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a]$

Rubi steps

$$\begin{aligned} \int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx &= -\frac{e(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(d + ex^2)} - \frac{\int \frac{-2Acd^2 - aBde - aAe^2 - 2cd(Bd - Ae)x^2 - ce(Bd - Ae)x^4}{(d + ex^2)\sqrt{a + cx^4}} dx}{2d(cd^2 + ae^2)} \\ &= -\frac{e(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(d + ex^2)} - \frac{\int \frac{-\sqrt{a}c^{3/2}de(Bd - Ae) + ce(-2Acd^2 - aBde - aAe^2) + (-2c^2de(Bd - Ae) + ce^2)}{(d + ex^2)\sqrt{a + cx^4}}}{2cde(cd^2 + ae^2)} \\ &= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(d + ex^2)} - \frac{\sqrt[4]{a}\sqrt[4]{c}(Bd - Ae)(\sqrt{a} - \sqrt{c}x^2)}{2d} \\ &= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(d + ex^2)} - \frac{(Bcd^3 - 3Acd^2e - aBde)}{4d^{3/2}\sqrt{c}} \end{aligned}$$

Mathematica [C] time = 1.15, size = 297, normalized size = 0.46

$$\frac{d \operatorname{erfc}\left(\frac{a+cx^4}{d+ex^2}\right)(Ae-Bd)}{(d+ex^2)(ae^2+cd^2)} - \frac{i\sqrt{\frac{cx^4}{a}+1}\left((aAe^3+aBde^2+3Acd^2e-Bcd^3)\Pi\left(-\frac{i\sqrt{ae}}{\sqrt{cd}}; i\sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x\right)\right)-1\right)+\sqrt{c}d(\sqrt{c}d-i\sqrt{ae})(Bd-Ae)F\left(i\sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x\right)\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}(ae^3+cd^2e)}$$

$$2d^2\sqrt{a+cx^4}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + c*x^4]), x]

[Out] ((d*e*(-(B*d) + A*e)*x*(a + c*x^4))/((c*d^2 + a*e^2)*(d + e*x^2)) - (I*Sqrt[1 + (c*x^4)/a]*(I*Sqrt[a]*Sqrt[c]*d*e*(B*d - A*e)*EllipticE[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + Sqrt[c]*d*(Sqrt[c]*d - I*Sqrt[a]*e)*(B*d - A*e)*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + (-B*c*d^3 + 3*A*c*d^2*e + a*B*d*e^2 + a*A*e^3)*EllipticPi[(-I)*Sqrt[a]*e/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1]))/(Sqrt[(I*Sqrt[c])/Sqrt[a]]*(c*d^2*e + a*e^3)))/(2*d^2*Sqrt[a + c*x^4])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2), x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2), x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)^2), x)

maple [C] time = 0.04, size = 679, normalized size = 1.06

$$\frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} B \operatorname{EllipticPi}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, x, \frac{i\sqrt{a}e}{\sqrt{c}d}, \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\right) (Ae - Bd)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} de} + \frac{\left(\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} a e^2 \operatorname{EllipticPi}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, \sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1}\right) + \dots\right)}{2(ae^2 + cd^2) \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2), x)`

[Out] $(Ae - Bd) / e^{1/2} e^2 / (ae^2 + cd^2) / d * x * (cx^4 + a)^{1/2} / (e * x^2 + d)^{-1/2} * c / (ae^2 + cd^2) / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (cx^4 + a)^{1/2} * \operatorname{EllipticF}((I/a^{1/2} * c^{1/2})^{1/2} * x, I) - 1/2 * I * c^{1/2} * e / d / (ae^2 + cd^2) * a^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (cx^4 + a)^{1/2} * \operatorname{EllipticF}((I/a^{1/2} * c^{1/2})^{1/2} * x, I) + 1/2 * I * c^{1/2} * e / d / (ae^2 + cd^2) * a^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (cx^4 + a)^{1/2} * \operatorname{EllipticE}((I/a^{1/2} * c^{1/2})^{1/2} * x, I) + 1/2 / d^2 / (ae^2 + cd^2) * e^2 / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (cx^4 + a)^{1/2} * \operatorname{EllipticPi}((I/a^{1/2} * c^{1/2})^{1/2} * x, I * a^{1/2} / c^{1/2} / d * e, (-I/a^{1/2} * c^{1/2})^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2}) * a + 3/2 / (ae^2 + cd^2) / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (cx^4 + a)^{1/2} * \operatorname{EllipticPi}((I/a^{1/2} * c^{1/2})^{1/2} * x, I * a^{1/2} / c^{1/2} / d * e, (-I/a^{1/2} * c^{1/2})^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2}) * c + B / e / d / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (cx^4 + a)^{1/2} * \operatorname{EllipticPi}((I/a^{1/2} * c^{1/2})^{1/2} * x, I * a^{1/2} / c^{1/2} / d * e, (-I/a^{1/2} * c^{1/2})^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2})$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a} (ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2), x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)^2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a} (ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^2), x)

[Out] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+a)**(1/2), x)

[Out] Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)**2), x)

$$3.7 \quad \int \frac{A+Bx^2}{(d+ex^2)^3 \sqrt{a+cx^4}} dx$$

Optimal. Leaf size=875

$$\frac{\sqrt{c} (5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3) \sqrt{cx^4 + a} x}{8d^2 (cd^2 + ae^2)^2 (\sqrt{c} x^2 + \sqrt{a})} - \frac{e (5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3) \sqrt{cx^4 + a} x}{8d^2 (cd^2 + ae^2)^2 (ex^2 + d)} - \frac{e(Bd - 3c^2d^5)}{4d (cd^2 - 3c^2d^5)}$$

[Out] 1/16*(3*A*e*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)-B*(-a^2*d*e^4-10*a*c*d^3*e^2+3*c^2*d^5))*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(5/2)/(a*e^2+c*d^2)^(5/2)/e^(1/2)-1/4*e*(-A*e+B*d)*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)/(e*x^2+d)^2-1/8*e*(-3*A*a*e^3-9*A*c*d^2*e-B*a*d*e^2+5*B*c*d^3)*x*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^2/(e*x^2+d)+1/8*(-3*A*a*e^3-9*A*c*d^2*e-B*a*d*e^2+5*B*c*d^3)*x*c^(1/2)*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^2/(a^(1/2)+x^2*c^(1/2))-1/8*a^(1/4)*c^(1/4)*(-3*A*a*e^3-9*A*c*d^2*e-B*a*d*e^2+5*B*c*d^3)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/d^2/(a*e^2+c*d^2)^2/(c*x^4+a)^(1/2)-1/32*(3*A*e*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)-B*(-a^2*d*e^4-10*a*c*d^3*e^2+3*c^2*d^5))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*2^(1/2))*(e*a^(1/2)+d*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(1/4)/c^(1/4)/d^3/e/(a*e^2+c*d^2)^2/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+a)^(1/2)+1/8*c^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*(4*A*c*d^2+a*e*(3*A*e+B*d)+d*(-A*e+B*d)*a^(1/2)*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(1/4)/d^2/(a*e^2+c*d^2)/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+a)^(1/2)

Rubi [A] time = 1.78, antiderivative size = 875, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1697, 1715, 1196, 1709, 220, 1707}

$$\frac{\sqrt{c} (5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3) \sqrt{cx^4 + a} x}{8d^2 (cd^2 + ae^2)^2 (\sqrt{c} x^2 + \sqrt{a})} - \frac{e (5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3) \sqrt{cx^4 + a} x}{8d^2 (cd^2 + ae^2)^2 (ex^2 + d)} - \frac{e(Bd - 3c^2d^5)}{4d (cd^2 - 3c^2d^5)}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + c*x^4]),x]

```
[Out] (Sqrt[c]*(5*B*c*d^3 - 9*A*c*d^2*e - a*B*d*e^2 - 3*a*A*e^3)*x*Sqrt[a + c*x^4
])/((8*d^2*(c*d^2 + a*e^2)^2*(Sqrt[a] + Sqrt[c]*x^2)) - (e*(B*d - A*e)*x*Sqr
t[a + c*x^4]))/(4*d*(c*d^2 + a*e^2)*(d + e*x^2)^2) - (e*(5*B*c*d^3 - 9*A*c*d
^2*e - a*B*d*e^2 - 3*a*A*e^3)*x*Sqrt[a + c*x^4])/((8*d^2*(c*d^2 + a*e^2)^2*(
d + e*x^2)) + ((3*A*e*(5*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4) - B*(3*c^2*d^5
- 10*a*c*d^3*e^2 - a^2*d*e^4))*ArcTan[(Sqrt[c*d^2 + a*e^2]*x)/(Sqrt[d]*Sqrt
[e]*Sqrt[a + c*x^4])])/(16*d^(5/2)*Sqrt[e]*(c*d^2 + a*e^2)^(5/2)) - (a^(1/4
)*c^(1/4)*(5*B*c*d^3 - 9*A*c*d^2*e - a*B*d*e^2 - 3*a*A*e^3)*(Sqrt[a] + Sqrt
[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^
(1/4)*x)/a^(1/4)], 1/2])/((8*d^2*(c*d^2 + a*e^2)^2*Sqrt[a + c*x^4]) + (c^(1/
4)*(4*A*c*d^2 + Sqrt[a]*Sqrt[c]*d*(B*d - A*e) + a*e*(B*d + 3*A*e))*(Sqrt[a]
+ Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*Arc
Tan[(c^(1/4)*x)/a^(1/4)], 1/2])/((8*a^(1/4)*d^2*(Sqrt[c]*d - Sqrt[a]*e)*(c*d
^2 + a*e^2)*Sqrt[a + c*x^4]) - ((Sqrt[c]*d + Sqrt[a]*e)*(3*A*e*(5*c^2*d^4 +
2*a*c*d^2*e^2 + a^2*e^4) - B*(3*c^2*d^5 - 10*a*c*d^3*e^2 - a^2*d*e^4))*(Sqr
t[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi
[-(Sqrt[c]*d - Sqrt[a]*e)^2/(4*Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a
^(1/4)], 1/2])/((32*a^(1/4)*c^(1/4)*d^3*e*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 +
a*e^2)^2*Sqrt[a + c*x^4])
```

Rule 220

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x]
, 1/2])/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 1196

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(
1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x]
, 1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e},
x] && PosQ[c/a]
```

Rule 1697

```
Int[((P4x_)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol
] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]
}, -Simp[((C*d^2 - B*d*e + A*e^2)*x*(d + e*x^2)^(q + 1)*Sqrt[a + c*x^4])/(2
*d*(q + 1)*(c*d^2 + a*e^2)), x] + Dist[1/(2*d*(q + 1)*(c*d^2 + a*e^2)), Int
[((d + e*x^2)^(q + 1)*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*c*d^2*(
q + 1)) + 2*d*(B*c*d - A*c*e + a*C*e)*(q + 1)*x^2 + c*(C*d^2 - B*d*e + A*e^
2)*(2*q + 5)*x^4, x])/Sqrt[a + c*x^4], x], x] /; FreeQ[{a, c, d, e}, x] &&
PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[c*d^2 + a*e^2, 0] && ILtQ[
q, -1]
```

Rule 1707

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[(Rt[(c*d)/e
+ (a*e)/d, 2]*x)/Sqrt[a + c*x^4]])/(2*d*e*Rt[(c*d)/e + (a*e)/d, 2]), x] +
Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a + c*x^4))/(a*(A + B*x^2)^2)]*Ell
ipticPi[Cancel[-((B*d - A*e)^2/(4*d*e*A*B))], 2*ArcTan[q*x], 1/2])/(4*d*e*A
*q*Sqrt[a + c*x^4]), x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e
^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1709

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q)
)/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4], x], x] + Dist[(a*(B*d - A*e)*(e
+ d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x],
x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2
- a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rule 1715

```
Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] :=
With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff
[P4x, x, 4]}, -Dist[C/(e*q), Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] + Dist
[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x^2)
*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2, 2]
&& NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + cx^4}} dx &= -\frac{e(Bd - Ae)x\sqrt{a + cx^4}}{4d(cd^2 + ae^2)(d + ex^2)^2} - \frac{\int \frac{-4Acd^2 - aBde - 3aAe^2 - 4cd(Bd - Ae)x^2 + ce(Bd - Ae)x^4}{(d + ex^2)^2 \sqrt{a + cx^4}} dx}{4d(cd^2 + ae^2)} \\
&= -\frac{e(Bd - Ae)x\sqrt{a + cx^4}}{4d(cd^2 + ae^2)(d + ex^2)^2} - \frac{e(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a + cx^4}}{8d^2(cd^2 + ae^2)^2(d + ex^2)} + \int \frac{e(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a + cx^4}}{8d^2(cd^2 + ae^2)^2(d + ex^2)} dx \\
&= -\frac{e(Bd - Ae)x\sqrt{a + cx^4}}{4d(cd^2 + ae^2)(d + ex^2)^2} - \frac{e(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a + cx^4}}{8d^2(cd^2 + ae^2)^2(d + ex^2)} + \int \frac{e(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a + cx^4}}{8d^2(cd^2 + ae^2)^2(d + ex^2)} dx \\
&= \frac{\sqrt{c}(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a + cx^4}}{8d^2(cd^2 + ae^2)^2(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + cx^4}}{4d(cd^2 + ae^2)(d + ex^2)^2} - \int \frac{e(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a + cx^4}}{8d^2(cd^2 + ae^2)^2(d + ex^2)} dx \\
&= \frac{\sqrt{c}(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a + cx^4}}{8d^2(cd^2 + ae^2)^2(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + cx^4}}{4d(cd^2 + ae^2)(d + ex^2)^2} - \int \frac{e(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a + cx^4}}{8d^2(cd^2 + ae^2)^2(d + ex^2)} dx
\end{aligned}$$

Mathematica [C] time = 2.13, size = 453, normalized size = 0.52

$$\frac{de^2x(a+cx^4)(2d(ae^2+cd^2)(Bd-Ae)+(d+ex^2)(-3aAe^3-aBde^2-9Acd^2e+5Bcd^3))}{(d+ex^2)^2} - i\sqrt{\frac{cx^4}{a}+1}\left(\left(3Ae(a^2e^4+2acd^2e^2+5c^2d^4)+B(a^2de^4+10acd^3e^2-3c^2d^4)\right)\sqrt{a+cx^4}\right)$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + c*x^4]),x]

[Out] (-((d*e^2*x*(a + c*x^4)*(2*d*(B*d - A*e)*(c*d^2 + a*e^2) + (5*B*c*d^3 - 9*A*c*d^2*e - a*B*d*e^2 - 3*a*A*e^3)*(d + e*x^2)))/(d + e*x^2)^2) - (I*Sqrt[1 + (c*x^4)/a]*((-I)*Sqrt[a]*Sqrt[c]*d*e*(-5*B*c*d^3 + 9*A*c*d^2*e + a*B*d*e^2 + 3*a*A*e^3)*EllipticE[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + Sqrt[c]*d*(Sqrt[c]*d - I*Sqrt[a]*e)*(A*e*(-7*c*d^2 + (2*I)*Sqrt[a]*Sqrt[c]*d*e - 3*a*e^2) + B*d*(3*c*d^2 - (2*I)*Sqrt[a]*Sqrt[c]*d*e - a*e^2))*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + (3*A*e*(5*c^2*d^4 + 2*a*c*d^2*e

$e^2 + a^2e^4) + B*(-3c^2d^5 + 10a*c*d^3e^2 + a^2*d*e^4))*\text{EllipticPi}[\left(\frac{-I*\sqrt{a}*e}{\sqrt{c}*d}, I*\text{ArcSinh}[\frac{\sqrt{I*\sqrt{c}}}{\sqrt{a}}*x], -1\right)]/\sqrt{\frac{I*\sqrt{c}}{\sqrt{a}}}/(8*d^3*e*(c*d^2 + a*e^2)^2*\sqrt{a + c*x^4})$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)^3), x)

maple [C] time = 0.04, size = 1591, normalized size = 1.82

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x)

[Out] $B/e*(1/2/(a*e^2+c*d^2)*(c*x^4+a)^(1/2)/(e*x^2+d)/d*e^2*x-1/2*c/(a*e^2+c*d^2))/((I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*\text{EllipticF}((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-1/2*I*c^(1/2)*e/d/(a*e^2+c*d^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*\text{EllipticF}((I/a^(1/2)*c^(1/2))^(1/2)*x,I)+1/2*I*c^(1/2)*e/d/(a*e^2+c*d^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*\text{EllipticE}((I/a^(1/2)*c^(1/2))^(1/2)*x,I)+1/2/d^2/(a*e^2+c*d^2)*e^2/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*\text{EllipticPi}((I/a^(1/2)*c^(1/2))^(1/2)*x,I*a^(1/2)/c^(1/2)/d*e,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))*a+3/2/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*\text{EllipticPi}((I/a^(1/2)*c^(1/2))^(1/2)*x,I*a^(1/2)/c^(1/2)/d*e,(-I/a^(1/2)*c$

[In] `int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^3), x)`

[Out] `int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+a)**(1/2), x)`

[Out] `Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)**3), x)`

$$3.8 \quad \int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=912

$$\frac{a^{3/4}B(\sqrt{c}x^2 + \sqrt{a})\sqrt{\frac{cx^4+a}{(\sqrt{c}x^2+\sqrt{a})^2}}F\left(2\tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)e^3}{6c^{9/4}\sqrt{cx^4+a}} + \frac{Bx\sqrt{cx^4+a}e^3}{3c^2} - \frac{\sqrt[4]{a}(3Bd+ Ae)(\sqrt{c}x^2 + \sqrt{a})\sqrt{\frac{cx^4+a}{(\sqrt{c}x^2+\sqrt{a})^2}}}{c^{7/4}\sqrt{cx^4+a}}$$

[Out] $\frac{1}{2}xx(Acd(-3ae^2+cd^2)-aBee(-ae^2+3cd^2)+c(-Aae^3+3Acd^2e-3Baad^2e^2+Bcd^3)x^2)/a/c^2/(cx^4+a)^{(1/2)}+1/3Bee^3xx(cx^4+a)^{(1/2)}/c^2+e^2(Ae+3Bd)xx(cx^4+a)^{(1/2)}/c^{(3/2)}/(a^{(1/2)}+x^2c^{(1/2)})-1/2(-Aae^3+3Acd^2e-3Baad^2e^2+Bcd^3)xx(cx^4+a)^{(1/2)}/a/c^{(3/2)}/(a^{(1/2)}+x^2c^{(1/2)})-a^{(1/4)}e^2(Ae+3Bd)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))*\text{EllipticE}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2c^{(1/2)})*((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/c^{(7/4)}/(cx^4+a)^{(1/2)}+1/2*(-Aae^3+3Acd^2e-3Baad^2e^2+Bcd^3)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))*\text{EllipticE}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2c^{(1/2)})*((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/a^{(3/4)}/c^{(7/4)}/(cx^4+a)^{(1/2)}-1/6a^{(3/4)}Bee^3(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))*\text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2c^{(1/2)})*((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/c^{(9/4)}/(cx^4+a)^{(1/2)}+1/2a^{(1/4)}e^2(Ae+3Bd)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))*\text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2c^{(1/2)})*((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/a^{(1/4)}/c^{(9/4)}/(cx^4+a)^{(1/2)}+1/4(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))*\text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2*2^{(1/2)})*(A*c^2*d^3+a^2*B*ee^3-3a*c*d*ee(Ae+Bd)-c^{(3/2)}*d^2*(3Ae+Bd)*a^{(1/2)}+a^{(3/2)}*e^2*(Ae+3Bd)*c^{(1/2)})*(a^{(1/2)}+x^2c^{(1/2)})*((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/a^{(5/4)}/c^{(9/4)}/(cx^4+a)^{(1/2)}$

Rubi [A] time = 0.73, antiderivative size = 912, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 7, integrand size = 28,

$\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1721, 1179, 1198, 220, 1196, 305, 321}

$$\frac{a^{3/4}B(\sqrt{c}x^2 + \sqrt{a})\sqrt{\frac{cx^4+a}{(\sqrt{c}x^2+\sqrt{a})^2}}F\left(2\tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)e^3}{6c^{9/4}\sqrt{cx^4+a}} + \frac{Bx\sqrt{cx^4+a}e^3}{3c^2} - \frac{\sqrt[4]{a}(3Bd+ Ae)(\sqrt{c}x^2 + \sqrt{a})\sqrt{\frac{cx^4+a}{(\sqrt{c}x^2+\sqrt{a})^2}}}{c^{7/4}\sqrt{cx^4+a}}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2), x]

[Out] (x*(A*c*d*(c*d^2 - 3*a*e^2) - a*B*e*(3*c*d^2 - a*e^2) + c*(B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*x^2))/(2*a*c^2*Sqrt[a + c*x^4]) + (B*e^3*x*Sqrt[a + c*x^4])/(3*c^2) + (e^2*(3*B*d + A*e)*x*Sqrt[a + c*x^4])/(c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - ((B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*x*Sqrt[a + c*x^4])/(2*a*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - (a^(1/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(7/4)*Sqrt[a + c*x^4]) + ((B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*c^(7/4)*Sqrt[a + c*x^4]) - (a^(3/4)*B*e^3*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(6*c^(9/4)*Sqrt[a + c*x^4]) + (a^(1/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(7/4)*Sqrt[a + c*x^4]) + (e*(3*B*c*d^2 + 3*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(9/4)*Sqrt[a + c*x^4]) + ((A*c^2*d^3 + a^2*B*e^3 - 3*a*c*d*e*(B*d + A*e) + a^(3/2)*Sqrt[c]*e^2*(3*B*d + A*e) - Sqrt[a]*c^(3/2)*d^2*(B*d + 3*A*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(5/4)*c^(9/4)*Sqrt[a + c*x^4])

Rule 220

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2])/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 305

Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Dist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 321

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c^(
n - 1)*(c*x)^(m - n + 1)*(a + b*x^n)^(p + 1))/(b*(m + n*p + 1)), x] - Dist[
(a*c^n*(m - n + 1))/(b*(m + n*p + 1)), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := -Simp[(x
*(d + e*x^2)*(a + c*x^4)^(p + 1))/(4*a*(p + 1)), x] + Dist[1/(4*a*(p + 1)),
Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /;
FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2
*p]
```

Rule 1196

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(
1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x],
1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0]] /; FreeQ[{a, c, d, e},
x] && PosQ[c/a]
```

Rule 1198

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0]] /; FreeQ[{a, c,
d, e}, x] && PosQ[c/a]
```

Rule 1721

```
Int[(Px_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx &= \int \left(\frac{Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)x^2}{c^2(a + cx^4)^{3/2}} \right) dx \\
&= \frac{\int \frac{Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)x^2}{(a + cx^4)^{3/2}} dx}{c^2} + \frac{(Be^3) \int \frac{x^4}{\sqrt{a + cx^4}} dx}{c} + \dots \\
&= \frac{x(Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)x^2)}{2ac^2\sqrt{a + cx^4}} \\
&= \frac{x(Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)x^2)}{2ac^2\sqrt{a + cx^4}} \\
&= \frac{x(Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)x^2)}{2ac^2\sqrt{a + cx^4}}
\end{aligned}$$

Mathematica [C] time = 0.28, size = 222, normalized size = 0.24

$$\frac{x\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) (3Acd(3ae^2 + cd^2) + aBe(9cd^2 - 5ae^2)) + 2cx^3\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{3}{4}, \frac{3}{2}; \frac{7}{4}; -\frac{cx^4}{a}\right) (-3aAe^3 + 3Bde^2 + Ae^3)}{6ac^2\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2), x]

[Out] (3*A*c*x*(c*d^3 + a*e^2*(-3*d + 2*e*x^2)) + a*B*e*x*(5*a*e^2 + c*(-9*d^2 + 18*d*e*x^2 + 2*e^2*x^4)) + (a*B*e*(9*c*d^2 - 5*a*e^2) + 3*A*c*d*(c*d^2 + 3*a*e^2))*x*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*c*(B*c*d^3 + 3*A*c*d^2*e - 9*a*B*d*e^2 - 3*a*A*e^3)*x^3*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c^2*sqrt[a + c*x^4])

fricas [F] time = 1.14, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(Be^3x^8 + (3Bde^2 + Ae^3)x^6 + 3(Bd^2e + Ade^2)x^4 + Ad^3 + (Bd^3 + 3Ad^2e)x^2)\sqrt{cx^4 + a}}{c^2x^8 + 2acx^4 + a^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] integral((B*e^3*x^8 + (3*B*d*e^2 + A*e^3)*x^6 + 3*(B*d^2*e + A*d*e^2)*x^4 + A*d^3 + (B*d^3 + 3*A*d^2*e)*x^2)*sqrt(c*x^4 + a)/(c^2*x^8 + 2*a*c*x^4 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^3/(c*x^4 + a)^(3/2), x)

maple [C] time = 0.04, size = 588, normalized size = 0.64

$$\left(\frac{x}{2\sqrt{(x^4 + \frac{a}{c})c} a} + \frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \operatorname{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x, i\right)}{2\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} a} \right) A d^3 + \left(\frac{ax}{2\sqrt{(x^4 + \frac{a}{c})c} c^2} - \frac{5\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1}}{\sqrt{a}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x)

[Out] B*e^3*(1/2/c^2*a*x/((x^4+a/c)*c)^(1/2)+1/3/c^2*x*(c*x^4+a)^(1/2)-5/6*a/c^2/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I))+ (A*e^3+3*B*d*e^2)*(-1/2/c*x^3/((x^4+a/c)*c)^(1/2)+3/2*I/c^(3/2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I)))+(3*A*d*e^2+3*B*d^2*e)*(-1/2/c*x/((x^4+a/c)*c)^(1/2)+1/2/c/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I))+(3*A*d^2*e+B*d^3)*(1/2/a*x^3/((x^4+a/c)*c)^(1/2)-1/2*I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I)))+A*d^3*(1/2/a*x/((x^4+a/c)*c)^(1/2)+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1

$)^{1/2} * (I/a^{1/2} * c^{1/2} * x^{2+1})^{1/2} / (c*x^4+a)^{1/2} * \text{EllipticF}((I/a^{1/2} * c^{1/2})^{1/2} * x, I)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^3/(c*x^4 + a)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+a)**(3/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)**3/(a + c*x**4)**(3/2), x)

[In] Int[((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2), x]

[Out] (x*(A*c*d^2 - 2*a*B*d*e - a*A*e^2 + (B*c*d^2 + 2*A*c*d*e - a*B*e^2)*x^2))/(2*a*c*Sqrt[a + c*x^4]) + (B*e^2*x*Sqrt[a + c*x^4])/(c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - ((B*c*d^2 + 2*A*c*d*e - a*B*e^2)*x*Sqrt[a + c*x^4])/(2*a*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - (a^(1/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(7/4)*Sqrt[a + c*x^4]) + ((B*c*d^2 + 2*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*c^(7/4)*Sqrt[a + c*x^4]) + (a^(1/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(7/4)*Sqrt[a + c*x^4]) + (e*(2*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(5/4)*Sqrt[a + c*x^4]) - ((B*c*d^2 + 2*A*c*d*e - a*B*e^2 - (Sqrt[c]*(A*c*d^2 - 2*a*B*d*e - a*A*e^2)))/Sqrt[a])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(3/4)*c^(7/4)*Sqrt[a + c*x^4])

Rule 220

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2])/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 305

Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Dist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 1179

Int[((d_) + (e_.)*(x_)^2)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := -Simp[(x*(d + e*x^2)*(a + c*x^4)^(p + 1))/(4*a*(p + 1)), x] + Dist[1/(4*a*(p + 1)), Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2*p]

Rule 1196

Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]

Rule 1198

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0]] /; FreeQ[{a, c,
d, e}, x] && PosQ[c/a]
```

Rule 1721

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned} \int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx &= \int \left(\frac{Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2}{c(a + cx^4)^{3/2}} + \frac{e(2Bd + Ae)}{c\sqrt{a + cx^4}} + \frac{Be^2x^2}{c\sqrt{a + cx^4}} \right) dx \\ &= \frac{\int \frac{Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2}{(a + cx^4)^{3/2}} dx}{c} + \frac{(Be^2) \int \frac{x^2}{\sqrt{a + cx^4}} dx}{c} + \frac{(e(2Bd + Ae)) \int \frac{1}{\sqrt{a + cx^4}} dx}{c} \\ &= \frac{x(Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2)}{2ac\sqrt{a + cx^4}} + \frac{e(2Bd + Ae)(\sqrt{a} + \sqrt{c}x^2)}{2\sqrt{a + cx^4}} \\ &= \frac{x(Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2)}{2ac\sqrt{a + cx^4}} + \frac{Be^2x\sqrt{a + cx^4}}{c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} - \frac{4\sqrt{a}e(2Bd + Ae)}{c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} \\ &= \frac{x(Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2)}{2ac\sqrt{a + cx^4}} + \frac{Be^2x\sqrt{a + cx^4}}{c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} - \frac{(Bcd^2 + 2Acde - aBe^2)x}{c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} \end{aligned}$$

Mathematica [C] time = 0.17, size = 166, normalized size = 0.24

$$\frac{3x\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) (aAe^2 + 2aBde + Acd^2) + 2x^3\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{3}{4}, \frac{3}{2}; \frac{7}{4}; -\frac{cx^4}{a}\right) (-3aBe^2 + 2Acde + Bcd^2)}{6ac\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2),x]

[Out] (3*A*(c*d^2 - a*e^2)*x + 6*a*B*e*x*(-d + e*x^2) + 3*(A*c*d^2 + 2*a*B*d*e + a*A*e^2)*x*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*(B*c*d^2 + 2*A*c*d*e - 3*a*B*e^2)*x^3*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c*sqrt[a + c*x^4])

fricas [F] time = 1.40, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(Be^2x^6 + (2Bde + Ae^2)x^4 + Ad^2 + (Bd^2 + 2Ade)x^2)\sqrt{cx^4 + a}}{c^2x^8 + 2acx^4 + a^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] integral((B*e^2*x^6 + (2*B*d*e + A*e^2)*x^4 + A*d^2 + (B*d^2 + 2*A*d*e)*x^2)*sqrt(c*x^4 + a)/(c^2*x^8 + 2*a*c*x^4 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^2/(c*x^4 + a)^(3/2), x)

maple [C] time = 0.01, size = 458, normalized size = 0.66

$$\left(\frac{x}{2\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \text{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right)}{2\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} \right) A d^2 + \left(-\frac{x^3}{2\sqrt{(x^4 + \frac{a}{c})c}} + \frac{3i\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1}}{\sqrt{a}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x)

[Out] B*e^2*(-1/2/((x^4+a/c)*c)^(1/2)/c*x^3+3/2*I/c^(3/2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)

```
/(c*x^4+a)^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I)))+(A*e^2+2*B*d*e)*(-1/2/((x^4+a/c)*c)^(1/2)/c*x+1/2/c/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)))+(2*A*d*e+B*d^2)*(1/2/((x^4+a/c)*c)^(1/2)/a*x^3-1/2*I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I)))+A*d^2*(1/2/((x^4+a/c)*c)^(1/2)/a*x+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I))
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((B*x^2 + A)*(e*x^2 + d)^2/(c*x^4 + a)^(3/2), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2),x)
```

```
[Out] int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+a)**(3/2),x)
```

```
[Out] Integral((A + B*x**2)*(d + e*x**2)**2/(a + c*x**4)**(3/2), x)
```

$$3.10 \quad \int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=395

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{2}\right) (-\sqrt{a}\sqrt{c}(Ae+Bd) - aBe + Acd)}{4a^{5/4}c^{5/4}\sqrt{a+cx^4}} + \frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{2a^{3/4}c^{5/4}}$$

[Out] $\frac{1}{2}x(Acd - aBe + c(Ae + Bd)x^2)/a/c/(cx^4+a)^{1/2} - \frac{1}{2}(Ae + Bd)x(c^2x^4+a)^{1/2}/a/c^{1/2}/(a^{1/2}+x^2c^{1/2}) + \frac{1}{2}(Ae + Bd)(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4}))\text{EllipticE}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2, 2^{1/2}) * (a^{1/2}+x^2c^{1/2}) * ((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^{1/2}/a^{3/4}/c^{3/4}/(cx^4+a)^{1/2} + \frac{1}{2}BBe(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4}))\text{EllipticF}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2, 2^{1/2}) * (a^{1/2}+x^2c^{1/2}) * ((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^{1/2}/a^{1/4}/c^{5/4}/(cx^4+a)^{1/2} + \frac{1}{4}(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4}))\text{EllipticF}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2, 2^{1/2}) * (a^{1/2}+x^2c^{1/2}) * (Acd - aBe - (Ae + Bd)a^{1/2}c^{1/2}) * ((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^{1/2}/a^{5/4}/c^{5/4}/(cx^4+a)^{1/2}$

Rubi [A] time = 0.33, antiderivative size = 395, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {1721, 1179, 1198, 220, 1196}

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{2}\right) (-\sqrt{a}\sqrt{c}(Ae+Bd) - aBe + Acd)}{4a^{5/4}c^{5/4}\sqrt{a+cx^4}} + \frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{2a^{3/4}c^{5/4}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(A + Bx^2)(d + ex^2)/(a + cx^4)^{3/2}, x]$

[Out] $(x(Acd - aBe + c(Bd + Ae)x^2))/(2a*c*\text{Sqrt}[a + cx^4]) - ((Bd + Ae)*x*\text{Sqrt}[a + cx^4])/(2*a*\text{Sqrt}[c]*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) + ((Bd + Ae)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + cx^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)]^2*\text{EllipticE}[2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], 1/2])/(2*a^{3/4}*c^{3/4}*\text{Sqrt}[a + cx^4]) + (Bee*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + cx^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)]^2*\text{EllipticF}[2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], 1/2])/(2*a^{1/4}*c^{5/4}*\text{Sqrt}[a + cx^4]) + ((Acd - aBe - \text{Sqrt}[a]*\text{Sqrt}[c]*(Bd + Ae))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2))/a^{5/4}/c^{5/4}/(cx^4+a)^{1/2}$

$c*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], 1/2]/(4*a^{5/4}*c^{5/4}*\text{Sqrt}[a + c*x^4])$

Rule 220

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*\text{Sqrt}[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2]]/(2*q*\text{Sqrt}[a + b*x^4]), x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[b/a]$

Rule 1179

$\text{Int}[(d_) + (e_)*(x_)^2]*((a_) + (c_)*(x_)^4)^{(p_)}, x_Symbol] := -\text{Simp}[(x*(d + e*x^2)*(a + c*x^4)^{(p + 1)})/(4*a*(p + 1)), x] + \text{Dist}[1/(4*a*(p + 1)), \text{Int}[\text{Simp}[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{LtQ}[p, -1] \&\& \text{IntegerQ}[2*p]$

Rule 1196

$\text{Int}[(d_) + (e_)*(x_)^2]/\text{Sqrt}[(a_) + (c_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 4]\}, -\text{Simp}[(d*x*\text{Sqrt}[a + c*x^4])/(a*(1 + q^2*x^2)), x] + \text{Simp}[(d*(1 + q^2*x^2)*\text{Sqrt}[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2])/(q*\text{Sqrt}[a + c*x^4]), x] /; \text{EqQ}[e + d*q^2, 0] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PosQ}[c/a]$

Rule 1198

$\text{Int}[(d_) + (e_)*(x_)^2]/\text{Sqrt}[(a_) + (c_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d*q)/q, \text{Int}[1/\text{Sqrt}[a + c*x^4], x], x] - \text{Dist}[e/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + c*x^4], x], x] /; \text{NeQ}[e + d*q, 0] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PosQ}[c/a]$

Rule 1721

$\text{Int}[(P_x)*((d_) + (e_)*(x_)^2)^{(q_)}*((a_) + (c_)*(x_)^4)^{(p_)}, x_Symbol] := \text{Int}[\text{ExpandIntegrand}[1/\text{Sqrt}[a + c*x^4], P_x*(d + e*x^2)^q*(a + c*x^4)^{(p + 1/2)}, x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PolyQ}[P_x, x^2] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{IntegerQ}[p + 1/2] \&\& \text{IntegerQ}[q]$

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx &= \int \left(\frac{Acd - aBe + c(Bd + Ae)x^2}{c(a + cx^4)^{3/2}} + \frac{Be}{c\sqrt{a + cx^4}} \right) dx \\
&= \frac{\int \frac{Acd - aBe + c(Bd + Ae)x^2}{(a + cx^4)^{3/2}} dx}{c} + \frac{(Be) \int \frac{1}{\sqrt{a + cx^4}} dx}{c} \\
&= \frac{x(Acd - aBe + c(Bd + Ae)x^2)}{2ac\sqrt{a + cx^4}} + \frac{Be(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\right)}{2\sqrt[4]{a}c^{5/4}\sqrt{a + cx^4}} \\
&= \frac{x(Acd - aBe + c(Bd + Ae)x^2)}{2ac\sqrt{a + cx^4}} + \frac{Be(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\right)}{2\sqrt[4]{a}c^{5/4}\sqrt{a + cx^4}} \\
&= \frac{x(Acd - aBe + c(Bd + Ae)x^2)}{2ac\sqrt{a + cx^4}} - \frac{(Bd + Ae)x\sqrt{a + cx^4}}{2a\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} + \frac{(Bd + Ae)(\sqrt{a} + \sqrt{c}x^2)}{2a^{3/4}\sqrt{c}}
\end{aligned}$$

Mathematica [C] time = 0.11, size = 126, normalized size = 0.32

$$\frac{3x\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right)(aBe + Acd) + 2cx^3\sqrt{\frac{cx^4}{a} + 1} {}_2F_1\left(\frac{3}{4}, \frac{3}{2}; \frac{7}{4}; -\frac{cx^4}{a}\right)(Ae + Bd) + 3x(Acd - aBe)}{6ac\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2), x]

[Out] (3*(A*c*d - a*B*e)*x + 3*(A*c*d + a*B*e)*x*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*c*(B*d + A*e)*x^3*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c*sqrt[a + c*x^4])

fricas [F] time = 1.16, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(Bex^4 + (Bd + Ae)x^2 + Ad)\sqrt{cx^4 + a}}{c^2x^8 + 2acx^4 + a^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2), x, algorithm="fricas")

[Out] integral((B*e*x^4 + (B*d + A*e)*x^2 + A*d)*sqrt(c*x^4 + a)/(c^2*x^8 + 2*a*c*x^4 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)/(c*x^4 + a)^(3/2), x)

maple [C] time = 0.00, size = 320, normalized size = 0.81

$$\left(\frac{x}{2\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \operatorname{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right)}{2\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} \right) Ad + \left(-\frac{x}{2\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \operatorname{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right)}{2\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x)

[Out] B*e*(-1/2/((x^4+a/c)*c)^(1/2)/c*x+1/2/c/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)+(A*e+B*d)*(1/2/((x^4+a/c)*c)^(1/2)/a*x^3-1/2*I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I))+A*d*(1/2/((x^4+a/c)*c)^(1/2)/a*x+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)/(c*x^4 + a)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2), x)

[Out] int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2), x)

sympy [C] time = 12.95, size = 167, normalized size = 0.42

$$\frac{A dx \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}} \Gamma\left(\frac{5}{4}\right)} + \frac{A e x^3 \Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{3}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}} \Gamma\left(\frac{7}{4}\right)} + \frac{B dx^3 \Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{3}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}} \Gamma\left(\frac{7}{4}\right)} + \frac{B e x^5 \Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{5}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}} \Gamma\left(\frac{9}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+a)**(3/2), x)

[Out] A*d*x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(5/4)) + A*e*x**3*gamma(3/4)*hyper((3/4, 3/2), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(7/4)) + B*d*x**3*gamma(3/4)*hyper((3/4, 3/2), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(7/4)) + B*e*x**5*gamma(5/4)*hyper((5/4, 3/2), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(9/4))

$$3.11 \quad \int \frac{A+Bx^2}{(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=262

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} (\sqrt{a}B - A\sqrt{c}) F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{4a^{5/4}c^{3/4}\sqrt{a+cx^4}} + \frac{B(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2a^{3/4}c^{3/4}\sqrt{a+cx^4}}$$

[Out] $\frac{1}{2}x(Bx^2+A)/a/(cx^4+a)^{1/2} - \frac{1}{2}Bx^2/(cx^4+a)^{1/2}/a/c^{1/2}/(a^{1/2}+x^2c^{1/2}) + \frac{1}{2}B(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4})) * \text{EllipticE}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2, 2^{1/2}) * (a^{1/2}+x^2c^{1/2}) * ((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^{1/2}/a^{3/4}/c^{3/4}/(cx^4+a)^{1/2} - \frac{1}{4}(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4})) * \text{EllipticF}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2, 2^{1/2}) * (Ba^{1/2}-Ac^{1/2}) * (a^{1/2}+x^2c^{1/2}) * ((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^{1/2}/a^{5/4}/c^{3/4}/(cx^4+a)^{1/2}$

Rubi [A] time = 0.10, antiderivative size = 262, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {1179, 1198, 220, 1196}

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} (\sqrt{a}B - A\sqrt{c}) F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{4a^{5/4}c^{3/4}\sqrt{a+cx^4}} + \frac{B(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2a^{3/4}c^{3/4}\sqrt{a+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/(a + c*x^4)^(3/2), x]

[Out] $(x(A + Bx^2))/(2a\sqrt{a + cx^4}) - (Bx\sqrt{a + cx^4})/(2a\sqrt{c} * (\sqrt{a} + \sqrt{c}x^2)) + (B(\sqrt{a} + \sqrt{c}x^2)\sqrt{(a + cx^4)/(\sqrt{a} + \sqrt{c}x^2)^2}) * \text{EllipticE}[2\text{ArcTan}[(c^{1/4}x)/a^{1/4}], 1/2]/(2a^{3/4}c^{3/4}\sqrt{a + cx^4}) - ((\sqrt{a}B - A\sqrt{c}) * (\sqrt{a} + \sqrt{c}x^2)\sqrt{(a + cx^4)/(\sqrt{a} + \sqrt{c}x^2)^2}) * \text{EllipticF}[2\text{ArcTan}[(c^{1/4}x)/a^{1/4}], 1/2]/(4a^{5/4}c^{3/4}\sqrt{a + cx^4})$

Rule 220

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2]]*EllipticF[2*ArcTan[q*x], 1/2]/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := -Simp[(x
*(d + e*x^2)*(a + c*x^4)^(p + 1))/(4*a*(p + 1)), x] + Dist[1/(4*a*(p + 1)),
  Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /;
FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2
*p]
```

Rule 1196

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(
  1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2]]*EllipticE[2*ArcTan[q*x],
  1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e},
  x] && PosQ[c/a]
```

Rule 1198

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c,
d, e}, x] && PosQ[c/a]
```

Rubi steps

$$\begin{aligned} \int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx &= \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} - \frac{\int \frac{-A+Bx^2}{\sqrt{a+cx^4}} dx}{2a} \\ &= \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} + \frac{\left(A - \frac{\sqrt{a}B}{\sqrt{c}}\right) \int \frac{1}{\sqrt{a+cx^4}} dx}{2a} + \frac{B \int \frac{1 - \frac{\sqrt{c}x^2}{\sqrt{a}}}{\sqrt{a+cx^4}} dx}{2\sqrt{a}\sqrt{c}} \\ &= \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} - \frac{Bx\sqrt{a + cx^4}}{2a\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} + \frac{B(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2a^{3/4}c^{3/4}\sqrt{a + cx^4}} \end{aligned}$$

Mathematica [C] time = 0.04, size = 99, normalized size = 0.38

$$\frac{3Ax\sqrt{\frac{cx^4}{a}} + 1 {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) + 2Bx^3\sqrt{\frac{cx^4}{a}} + 1 {}_2F_1\left(\frac{3}{4}, \frac{3}{2}; \frac{7}{4}; -\frac{cx^4}{a}\right) + 3Ax}{6a\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/(a + c*x^4)^(3/2),x]

[Out] (3*A*x + 3*A*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*B*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)])/(6*a*Sqrt[a + c*x^4])

fricas [F] time = 0.91, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{cx^4 + a} (Bx^2 + A)}{c^2x^8 + 2acx^4 + a^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(c*x^4 + a)*(B*x^2 + A)/(c^2*x^8 + 2*a*c*x^4 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(c*x^4 + a)^(3/2), x)

maple [C] time = 0.00, size = 217, normalized size = 0.83

$$\left(\frac{x}{2\sqrt{(x^4 + \frac{a}{c})ca}} + \frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \text{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right)}{2\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} a} \right) A + \left(\frac{x^3}{2\sqrt{(x^4 + \frac{a}{c})ca}} - \frac{i\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}} + 1}}{2\sqrt{(x^4 + \frac{a}{c})ca}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(c*x^4+a)^(3/2),x)

[Out] B*(1/2/((x^4+a/c)*c)^(1/2)/a*x^3-1/2*I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)-EllipticE((I/a^(1/2)*c^(1/2))^(1/2)*x,I))+A*(1/2/((x^4+a/c)*c)^(1/2)/a*x+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2))

$2))^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * \text{EllipticF}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(c*x^4 + a)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/(a + c*x^4)^(3/2),x)

[Out] int((A + B*x^2)/(a + c*x^4)^(3/2), x)

sympy [C] time = 7.19, size = 78, normalized size = 0.30

$$\frac{Ax\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{5}{4}\right)} + \frac{Bx^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{3}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{7}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(c*x**4+a)**(3/2),x)

[Out] A*x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(5/4)) + B*x**3*gamma(3/4)*hyper((3/4, 3/2), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(7/4))

$$3.12 \quad \int \frac{A+Bx^2}{(d+ex^2)(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=732

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{2}\right) (-\sqrt{a}\sqrt{c}(Bd - Ae) + aBe + Acd) \sqrt[4]{c} (\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{4a^{5/4} \sqrt[4]{c} \sqrt{a+cx^4} (ae^2 + cd^2)} + \frac{\sqrt[4]{c} (\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{2a^{3/4} \sqrt{a+cx^4}}$$

[Out] $-1/2 * e^{(3/2)} * (-A * e + B * d) * \arctan(x * (a * e^2 + c * d^2)^{(1/2)} / d^{(1/2)} / e^{(1/2)} / (c * x^4 + a)^{(1/2)}) / (a * e^2 + c * d^2)^{(3/2)} / d^{(1/2)} + 1/2 * x * (A * c * d + a * B * e + c * (-A * e + B * d) * x^2) / a / (a * e^2 + c * d^2) / (c * x^4 + a)^{(1/2)} - 1/2 * (-A * e + B * d) * x * c^{(1/2)} * (c * x^4 + a)^{(1/2)} / a / (a * e^2 + c * d^2) / (a^{(1/2)} + x^2 * c^{(1/2)}) + 1/2 * c^{(1/4)} * (-A * e + B * d) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})))^2)^{(1/2)} / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticE}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^2)^{(1/2)} / a^{(3/4)} / (a * e^2 + c * d^2) / (c * x^4 + a)^{(1/2)} - 1/2 * c^{(1/4)} * e * (-A * e + B * d) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})))^2)^{(1/2)} / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticF}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^2)^{(1/2)} / a^{(1/4)} / (a * e^2 + c * d^2) / (-e * a^{(1/2)} + d * c^{(1/2)}) / (c * x^4 + a)^{(1/2)} + 1/4 * a^{(3/4)} * e * (-A * e + B * d) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})))^2)^{(1/2)} / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticPi}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), -1/4 * (-e * a^{(1/2)} + d * c^{(1/2)})^2 / d / e / a^{(1/2)} / c^{(1/2)}, 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * (e + d * c^{(1/2)} / a^{(1/2)})^2 * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^2)^{(1/2)} / c^{(1/4)} / d / (-a^2 * e^4 + c^2 * d^4) / (c * x^4 + a)^{(1/2)} + 1/4 * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})))^2)^{(1/2)} / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticF}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * (A * c * d + a * B * e - (-A * e + B * d) * a^{(1/2)} * c^{(1/2)}) * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^2)^{(1/2)} / a^{(5/4)} / c^{(1/4)} / (a * e^2 + c * d^2) / (c * x^4 + a)^{(1/2)}$

Rubi [A] time = 0.78, antiderivative size = 732, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1721, 1179, 1198, 220, 1196, 1217, 1707}

$$\frac{a^{3/4} e (\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \left(\frac{\sqrt{c}d}{\sqrt{a}} + e\right)^2 (Bd - Ae) \Pi\left(-\frac{(\sqrt{c}d - \sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{2}\right) (\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{4\sqrt[4]{c} d \sqrt{a+cx^4} (c^2 d^4 - a^2 e^4)} + \frac{\sqrt[4]{c} (\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{2a^{3/4} \sqrt{a+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)*(a + c*x^4)^(3/2)), x]


```
[Out] (x*(A*c*d + a*B*e + c*(B*d - A*e)*x^2))/(2*a*(c*d^2 + a*e^2)*Sqrt[a + c*x^4]
) - (Sqrt[c]*(B*d - A*e)*x*Sqrt[a + c*x^4])/(2*a*(c*d^2 + a*e^2)*(Sqrt[a
+ Sqrt[c]*x^2)) - (e^(3/2)*(B*d - A*e)*ArcTan[(Sqrt[c*d^2 + a*e^2]*x)/(Sqrt
[d]*Sqrt[e]*Sqrt[a + c*x^4])])/(2*Sqrt[d]*(c*d^2 + a*e^2)^(3/2)) + (c^(1/4)
*(B*d - A*e)*(Sqrt[a + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^
2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*(c*d^2 + a*
e^2)*Sqrt[a + c*x^4]) - (c^(1/4)*e*(B*d - A*e)*(Sqrt[a + Sqrt[c]*x^2)*Sqrt
[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1
/4)], 1/2])/(2*a^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 + a*e^2)*Sqrt[a + c*x
^4]) + ((A*c*d + a*B*e - Sqrt[a]*Sqrt[c]*(B*d - A*e))*(Sqrt[a + Sqrt[c]*x^
2)*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*
x)/a^(1/4)], 1/2])/(4*a^(5/4)*c^(1/4)*(c*d^2 + a*e^2)*Sqrt[a + c*x^4]) + (a
^(3/4)*e*((Sqrt[c]*d)/Sqrt[a + e]^2*(B*d - A*e)*(Sqrt[a + Sqrt[c]*x^2)*Sq
rt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2)^2]*EllipticPi[-(Sqrt[c]*d - Sqrt[a]*
e)^2/(4*Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*c^(1/
4)*d*(c^2*d^4 - a^2*e^4)*Sqrt[a + c*x^4])
```

Rule 220

```
Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])*EllipticF[2*ArcTan[q*x]
, 1/2])/(2*q*Sqrt[a + b*x^4]), x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := -Simp[(x
*(d + e*x^2)*(a + c*x^4)^(p + 1))/(4*a*(p + 1)), x] + Dist[1/(4*a*(p + 1)),
Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /;
FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2
*p]
```

Rule 1196

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(
1 + q^2*x^2)*Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])*EllipticE[2*ArcTan[q*x],
1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e},
x] && PosQ[c/a]
```

Rule 1198

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c,
d, e}, x] && PosQ[c/a]
```

Rule 1217

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[c/a, 2]}, Dist[(c*d + a*e*q)/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4]
, x], x] - Dist[(a*e*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^
2)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^
2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
```

Rule 1707

```
Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4])
, x_Symbol] := With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[(Rt[(c*d)/e
+ (a*e)/d, 2]*x)/Sqrt[a + c*x^4]])/(2*d*e*Rt[(c*d)/e + (a*e)/d, 2]), x] +
Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a + c*x^4))/(a*(A + B*x^2)^2)]*Ell
ipticPi[Cancel[-((B*d - A*e)^2/(4*d*e*A*B))], 2*ArcTan[q*x], 1/2])/(4*d*e*A
*q*Sqrt[a + c*x^4]), x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e
^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1721

```
Int[(Px_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)(a + cx^4)^{3/2}} dx &= \int \left(\frac{Acd + aBe + c(Bd - Ae)x^2}{(cd^2 + ae^2)(a + cx^4)^{3/2}} + \frac{e(-Bd + Ae)}{(cd^2 + ae^2)(d + ex^2)\sqrt{a + cx^4}} \right) dx \\
&= \frac{\int \frac{Acd + aBe + c(Bd - Ae)x^2}{(a + cx^4)^{3/2}} dx}{cd^2 + ae^2} - \frac{(e(Bd - Ae)) \int \frac{1}{(d + ex^2)\sqrt{a + cx^4}} dx}{cd^2 + ae^2} \\
&= \frac{x(Acd + aBe + c(Bd - Ae)x^2)}{2a(cd^2 + ae^2)\sqrt{a + cx^4}} - \frac{\int \frac{-Acd - aBe + c(Bd - Ae)x^2}{\sqrt{a + cx^4}} dx}{2a(cd^2 + ae^2)} - \frac{(\sqrt{c}e(Bd - Ae)) \int \frac{1}{\sqrt{d + ex^2}} dx}{(\sqrt{c}d - \sqrt{a}e)(cd^2 + ae^2)} \\
&= \frac{x(Acd + aBe + c(Bd - Ae)x^2)}{2a(cd^2 + ae^2)\sqrt{a + cx^4}} - \frac{e^{3/2}(Bd - Ae) \tan^{-1}\left(\frac{\sqrt{cd^2 + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a + cx^4}}\right)}{2\sqrt{d}(cd^2 + ae^2)^{3/2}} - \frac{\sqrt[4]{c}e(Bd - Ae)}{2\sqrt{d}(cd^2 + ae^2)} \\
&= \frac{x(Acd + aBe + c(Bd - Ae)x^2)}{2a(cd^2 + ae^2)\sqrt{a + cx^4}} - \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + cx^4}}{2a(cd^2 + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e^{3/2}(Bd - Ae) \tan^{-1}\left(\frac{\sqrt{cd^2 + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a + cx^4}}\right)}{2\sqrt{d}(cd^2 + ae^2)^{3/2}}
\end{aligned}$$

Mathematica [C] time = 0.79, size = 432, normalized size = 0.59

$$d\sqrt{\frac{cx^4}{a} + 1} (\sqrt{a}B - iA\sqrt{c})(\sqrt{c}d - i\sqrt{a}e) F\left(i \sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x\right) \middle| -1\right) - \sqrt{a}\sqrt{c}d\sqrt{\frac{cx^4}{a} + 1} (Bd - Ae) E\left(i \sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x\right) \middle| -1\right)$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)*(a + c*x^4)^(3/2)),x]

[Out] (A*Sqrt[(I*Sqrt[c])/Sqrt[a]]*c*d^2*x + a*B*Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*e*x + B*Sqrt[(I*Sqrt[c])/Sqrt[a]]*c*d^2*x^3 - A*Sqrt[(I*Sqrt[c])/Sqrt[a]]*c*d*e*x^3 - Sqrt[a]*Sqrt[c]*d*(B*d - A*e)*Sqrt[1 + (c*x^4)/a]*EllipticE[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + (Sqrt[a]*B - I*A*Sqrt[c])*d*(Sqrt[c]*d - I*Sqrt[a]*e)*Sqrt[1 + (c*x^4)/a]*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + (2*I)*a*B*d*e*Sqrt[1 + (c*x^4)/a]*EllipticPi[(-I)*Sqrt[a]*e)/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] - (2*I)*a*A*e^2*Sqrt[1 + (c*x^4)/a]*EllipticPi[(-I)*Sqrt[a]*e)/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1)/(2*a*Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*(c*d^2 + a*e^2)*Sqrt[a + c*x^4])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)), x)

maple [C] time = 0.03, size = 564, normalized size = 0.77

$$\frac{\left(\frac{x}{2\sqrt{(x^4 + \frac{a}{c})c}a} + \frac{\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \operatorname{EllipticF}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right)}{2\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}a} \right) B}{e} + \frac{(Ae - Bd) \left(-\frac{i\sqrt{-\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{\frac{i\sqrt{c}x^2}{\sqrt{a}}+1} \sqrt{c} e \operatorname{EllipticE}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x, i\right)}{2(ae^2 + cd^2)\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}\sqrt{a}} \right)}{e}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x)

[Out] B/e*(1/2/((x^4+a/c)*c)^(1/2)/a*x+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x, I)+(A*e-B*d)/e*(-2*c*(1/4/a*e/(a*e^2+c*d^2)*x^3-1/4/a*d/(a*e^2+c*d^2)*x)/((x^4+a/c)*c)^(1/2)+1/2/a*c*d/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x, I)+1/2*I/a^(1/2)*c^(1/2)*e/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x, I)-1/2*I/a^(1/2)*c^(1/2)*e/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2))^(1/2)*x, I)

) $x^2+1)^{1/2}/(cx^4+a)^{1/2}$ *EllipticE((I/a $^{1/2}$ *c $^{1/2}$) $^{1/2}$ *x,I)+1/(a*e $^2+c*d^2$)*e $^2/d/(I/a^{1/2}$ *c $^{1/2}$) $^{1/2}$ *(-I/a $^{1/2}$ *c $^{1/2}$ *x $^2+1)^{1/2}$)*(I/a $^{1/2}$ *c $^{1/2}$ *x $^2+1)^{1/2}/(cx^4+a)^{1/2}$ *EllipticPi((I/a $^{1/2}$ *c $^{1/2}$) $^{1/2}$ *x,I*a $^{1/2}/c^{1/2}/d$ *e,(-I/a $^{1/2}$ *c $^{1/2}$) $^{1/2}/(I/a^{1/2}$ *c $^{1/2}$) $^{1/2}$))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)),x)

[Out] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(a + cx^4)^{\frac{3}{2}}(d + ex^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+a)**(3/2),x)

[Out] Integral((A + B*x**2)/((a + c*x**4)**(3/2)*(d + e*x**2)), x)

$$3.13 \quad \int \frac{A+Bx^2}{(d+ex^2)^2(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=1494

$$\frac{(Bd - Ae)x\sqrt{cx^4 + a} e^3}{2d(cd^2 + ae^2)^2 (ex^2 + d)} - \frac{\sqrt[4]{a} \sqrt[4]{c} (Bd - Ae) (\sqrt{c} x^2 + \sqrt{a}) \sqrt{\frac{cx^4 + a}{(\sqrt{c} x^2 + \sqrt{a})^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c} x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) e^2}{2d(cd^2 + ae^2)^2 \sqrt{cx^4 + a}} + \frac{\sqrt{c} (Bd - Ae)}{2d(cd^2 + ae^2)}$$

[Out] $-1/4 * e^{(3/2)} * (-A * e + B * d) * (a * e^2 + 3 * c * d^2) * \arctan(x * (a * e^2 + c * d^2)^{(1/2)} / d^{(1/2)}) / e^{(1/2)} / (c * x^4 + a)^{(1/2)} / d^{(3/2)} / (a * e^2 + c * d^2)^{(5/2)} - 1/2 * e^{(3/2)} * (-2 * A * c * d * e - B * a * e^2 + B * c * d^2) * \arctan(x * (a * e^2 + c * d^2)^{(1/2)} / d^{(1/2)}) / e^{(1/2)} / (c * x^4 + a)^{(1/2)} / (a * e^2 + c * d^2)^{(5/2)} / d^{(1/2)} + 1/2 * c * x * (A * c * d^2 + 2 * a * B * d * e - a * A * e^2 + (-2 * A * c * d * e - B * a * e^2 + B * c * d^2) * x^2) / a / (a * e^2 + c * d^2)^2 / (c * x^4 + a)^{(1/2)} - 1/2 * e^3 * (-A * e + B * d) * x * (c * x^4 + a)^{(1/2)} / d / (a * e^2 + c * d^2)^2 / (e * x^2 + d) + 1/2 * e^2 * (-A * e + B * d) * x * c^{(1/2)} * (c * x^4 + a)^{(1/2)} / d / (a * e^2 + c * d^2)^2 / (a^{(1/2)} + x^2 * c^{(1/2)}) - 1/2 * (-2 * A * c * d * e - B * a * e^2 + B * c * d^2) * x * c^{(1/2)} * (c * x^4 + a)^{(1/2)} / a / (a * e^2 + c * d^2)^2 / (a^{(1/2)} + x^2 * c^{(1/2)}) - 1/2 * a^{(1/4)} * c^{(1/4)} * e^2 * (-A * e + B * d) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)}))^{(1/2)}) / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticE}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^{(1/2)} / d / (a * e^2 + c * d^2)^2 / (c * x^4 + a)^{(1/2)} + 1/2 * c^{(1/4)} * (-2 * A * c * d * e - B * a * e^2 + B * c * d^2) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)}))^{(1/2)}) / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticE}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^{(1/2)} / a^{(3/4)} / (a * e^2 + c * d^2)^2 / (c * x^4 + a)^{(1/2)} - 1/2 * c^{(1/4)} * e * (-A * e + B * d) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)}))^{(1/2)}) / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticF}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^{(1/2)} / a^{(1/4)} / d / (a * e^2 + c * d^2) / (-e * a^{(1/2)} + d * c^{(1/2)}) / (c * x^4 + a)^{(1/2)} - 1/2 * c^{(1/4)} * e * (-2 * A * c * d * e - B * a * e^2 + B * c * d^2) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)}))^{(1/2)}) / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticF}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^{(1/2)} / a^{(1/4)} / (a * e^2 + c * d^2)^2 / (-e * a^{(1/2)} + d * c^{(1/2)}) / (c * x^4 + a)^{(1/2)} + 1/8 * e * (-A * e + B * d) * (a * e^2 + 3 * c * d^2) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)}))^{(1/2)}) / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticPi}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), -1/4 * (-e * a^{(1/2)} + d * c^{(1/2)})^2 / d / e / a^{(1/2)} / c^{(1/2)}, 1/2 * 2^{(1/2)}) * (e * a^{(1/2)} + d * c^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^{(1/2)} / a^{(1/4)} / c^{(1/4)} / d^2 / (a * e^2 + c * d^2)^2 / (-e * a^{(1/2)} + d * c^{(1/2)}) / (c * x^4 + a)^{(1/2)} + 1/4 * a^{(3/4)} * e * (-2 * A * c * d * e - B * a * e^2 + B * c * d^2) * (\cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)}))^{(1/2)}) / \cos(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})) * \text{EllipticPi}(\sin(2 * \arctan(c^{(1/4)} * x / a^{(1/4)})), -1/4 * (-e * a^{(1/2)} + d * c^{(1/2)})^2 / d / e / a^{(1/2)} / c^{(1/2)}, 1/2 * 2^{(1/2)}) * (a^{(1/2)} + x^2 * c^{(1/2)}) * (e + d * c^{(1/2)} / a^{(1/2)})^2 * ((c * x^4 + a) / (a^{(1/2)} + x^2 * c^{(1/2)}))^{(1/2)} / c^{(1/4)} / d / (-a * e^2 + c * d^2) / (a * e^2 + c * d^2)^2 / (c$

$a)e^2/(4\sqrt{a}\sqrt{c}de), 2\text{ArcTan}[(c^{1/4}x)/a^{1/4}], 1/2]/(4c^{1/4}d(c^2d^2 - a^2e^2)(c^2d^2 + a^2e^2)^2\sqrt{a + cx^4})$

Rule 220

$\text{Int}[1/\sqrt{(a_+) + (b_+)(x_+)^4}, x_Symbol] := \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2x^2)\sqrt{(a + bx^4)/(a(1 + q^2x^2)^2)}]\text{EllipticF}[2\text{ArcTan}[qx], 1/2]/(2q\sqrt{a + bx^4}), x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[b/a]$

Rule 1179

$\text{Int}[(d_+) + (e_+)(x_+)^2)((a_+) + (c_+)(x_+)^4)^{p_+}, x_Symbol] := -\text{Simp}[(x(d + ex^2)(a + cx^4)^{p+1})/(4a^{p+1}), x] + \text{Dist}[1/(4a^{p+1}), \text{Int}[\text{Simp}[d(4p + 5) + e(4p + 7)x^2, x](a + cx^4)^{p+1}, x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[c^2d^2 + a^2e^2, 0] \&\& \text{LtQ}[p, -1] \&\& \text{IntegerQ}[2p]$

Rule 1196

$\text{Int}[(d_+) + (e_+)(x_+)^2]/\sqrt{(a_+) + (c_+)(x_+)^4}, x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 4]\}, -\text{Simp}[(d*x\sqrt{a + cx^4})/(a(1 + q^2x^2)), x] + \text{Simp}[(d(1 + q^2x^2)\sqrt{(a + cx^4)/(a(1 + q^2x^2)^2)}]\text{EllipticE}[2\text{ArcTan}[qx], 1/2]/(q\sqrt{a + cx^4}), x] /; \text{EqQ}[e + dq^2, 0] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PosQ}[c/a]$

Rule 1198

$\text{Int}[(d_+) + (e_+)(x_+)^2]/\sqrt{(a_+) + (c_+)(x_+)^4}, x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + dq)/q, \text{Int}[1/\sqrt{a + cx^4}, x], x] - \text{Dist}[e/q, \text{Int}[(1 - qx^2)/\sqrt{a + cx^4}, x], x] /; \text{NeQ}[e + dq, 0] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PosQ}[c/a]$

Rule 1217

$\text{Int}[1/(((d_+) + (e_+)(x_+)^2)\sqrt{(a_+) + (c_+)(x_+)^4}), x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(cd + aeq)/(c^2d^2 - a^2e^2), \text{Int}[1/\sqrt{a + cx^4}, x], x] - \text{Dist}[(a^2e(e + dq))/(c^2d^2 - a^2e^2), \text{Int}[(1 + qx^2)/((d + ex^2)\sqrt{a + cx^4}), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[c^2d^2 + a^2e^2, 0] \&\& \text{NeQ}[c^2d^2 - a^2e^2, 0] \&\& \text{PosQ}[c/a]$

Rule 1224

$\text{Int}[(d_+) + (e_+)(x_+)^2]^{q_+}/\sqrt{(a_+) + (c_+)(x_+)^4}, x_Symbol] := -\text{Simp}[(e^2x(d + ex^2)^{q+1}\sqrt{a + cx^4})/(2d(q + 1)(c^2d^2 + a^2e^2)), x] + \text{Dist}[1/(2d(q + 1)(c^2d^2 + a^2e^2)), \text{Int}[(d + ex^2)^{q+1}\text{Simp}[a^2e^2(2q + 3) + 2c^2d^2(q + 1) - 2e^2cd(q + 1)x^2 + c^2e^2(2q + 5)x$

$\wedge 4, x])/\text{Sqrt}[a + c*x^4], x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{ILtQ}[q, -1]$

Rule 1707

$\text{Int}[(A_) + (B_)*(x_)^2]/(((d_) + (e_)*(x_)^2)*\text{Sqrt}[(a_) + (c_)*(x_)^4])$
 $, x_Symbol] :> \text{With}[\{q = \text{Rt}[B/A, 2]\}, -\text{Simp}[(B*d - A*e)*\text{ArcTan}[(\text{Rt}[(c*d)/e$
 $+ (a*e)/d, 2]*x)/\text{Sqrt}[a + c*x^4]]/(2*d*e*\text{Rt}[(c*d)/e + (a*e)/d, 2]), x] +$
 $\text{Simp}[(B*d + A*e)*(A + B*x^2)*\text{Sqrt}[(A^2*(a + c*x^4))/(a*(A + B*x^2)^2)]*\text{Ell}$
 $\text{ipticPi}[\text{Cancel}[-((B*d - A*e)^2/(4*d*e*A*B))], 2*\text{ArcTan}[q*x], 1/2)]/(4*d*e*A$
 $*q*\text{Sqrt}[a + c*x^4]), x] /; \text{FreeQ}[\{a, c, d, e, A, B\}, x] \&\& \text{NeQ}[c*d^2 + a*e$
 $^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a] \&\& \text{EqQ}[c*A^2 - a*B^2, 0]$

Rule 1709

$\text{Int}[(A_) + (B_)*(x_)^2]/(((d_) + (e_)*(x_)^2)*\text{Sqrt}[(a_) + (c_)*(x_)^4])$
 $, x_Symbol] :> \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(A*(c*d + a*e*q) - a*B*(e + d*q)$
 $)/(c*d^2 - a*e^2), \text{Int}[1/\text{Sqrt}[a + c*x^4], x], x] + \text{Dist}[(a*(B*d - A*e)*(e$
 $+ d*q))/(c*d^2 - a*e^2), \text{Int}[(1 + q*x^2)/((d + e*x^2)*\text{Sqrt}[a + c*x^4]), x],$
 $x]] /; \text{FreeQ}[\{a, c, d, e, A, B\}, x] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{NeQ}[c*d^2$
 $- a*e^2, 0] \&\& \text{PosQ}[c/a] \&\& \text{NeQ}[c*A^2 - a*B^2, 0]$

Rule 1715

$\text{Int}[(P4x_)/(((d_) + (e_)*(x_)^2)*\text{Sqrt}[(a_) + (c_)*(x_)^4]), x_Symbol] :>$
 $\text{With}[\{q = \text{Rt}[c/a, 2], A = \text{Coeff}[P4x, x, 0], B = \text{Coeff}[P4x, x, 2], C = \text{Coeff}$
 $[P4x, x, 4]\}, -\text{Dist}[C/(e*q), \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + c*x^4], x], x] + \text{Dist}$
 $[1/(c*e), \text{Int}[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x^2)$
 $)*\text{Sqrt}[a + c*x^4]), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PolyQ}[P4x, x^2, 2]$
 $\&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a]$

Rule 1721

$\text{Int}[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]$
 $:> \text{Int}[\text{ExpandIntegrand}[1/\text{Sqrt}[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p$
 $+ 1/2), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PolyQ}[Px, x^2] \&\& \text{NeQ}[c*d^2 + a$
 $*e^2, 0] \&\& \text{IntegerQ}[p + 1/2] \&\& \text{IntegerQ}[q]$

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^2 (a + cx^4)^{3/2}} dx &= \int \left(\frac{c (Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{(cd^2 + ae^2)^2 (a + cx^4)^{3/2}} + \frac{e(-Bd + A)}{(cd^2 + ae^2)(d + ex^2)} \right) dx \\
&= \frac{c \int \frac{Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2}{(a + cx^4)^{3/2}} dx}{(cd^2 + ae^2)^2} - \frac{(e(Bd - Ae)) \int \frac{1}{(d + ex^2)^2 \sqrt{a + cx^4}} dx}{cd^2 + ae^2} \\
&= \frac{cx (Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{2a (cd^2 + ae^2)^2 \sqrt{a + cx^4}} - \frac{e^3(Bd - Ae)x\sqrt{a + cx^4}}{2d (cd^2 + ae^2)^2 (d + ex^2)} \\
&= \frac{cx (Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{2a (cd^2 + ae^2)^2 \sqrt{a + cx^4}} - \frac{e^3(Bd - Ae)x\sqrt{a + cx^4}}{2d (cd^2 + ae^2)^2 (d + ex^2)} \\
&= \frac{cx (Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{2a (cd^2 + ae^2)^2 \sqrt{a + cx^4}} + \frac{\sqrt{c} e^2 (Bd - Ae)x\sqrt{a + cx^4}}{2d (cd^2 + ae^2)^2 (\sqrt{a + cx^4} + \sqrt{a + cx^4})} \\
&= \frac{cx (Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{2a (cd^2 + ae^2)^2 \sqrt{a + cx^4}} + \frac{\sqrt{c} e^2 (Bd - Ae)x\sqrt{a + cx^4}}{2d (cd^2 + ae^2)^2 (\sqrt{a + cx^4} + \sqrt{a + cx^4})}
\end{aligned}$$

Mathematica [C] time = 1.57, size = 427, normalized size = 0.29

$$d \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \left(cdx(d + ex^2)(-aAe^2 + aBe(2d - ex^2) + Acd(d - 2ex^2) + Bcd^2x^2) + ae^3x(a + cx^4)(Ae - Bd) \right) - \sqrt{\frac{cx^4}{a}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^2*(a + c*x^4)^(3/2)),x]

[Out] (Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*(a*e^3*(-(B*d) + A*e)*x*(a + c*x^4) + c*d*x*(d + e*x^2)*(-(a*A*e^2) + B*c*d^2*x^2 + A*c*d*(d - 2*e*x^2) + a*B*e*(2*d - e*x^2))) - (d + e*x^2)*Sqrt[1 + (c*x^4)/a]*(-(Sqrt[a]*Sqrt[c]*d*(-(B*c*d^3) + 2*A*c*d^2*e + 2*a*B*d*e^2 - a*A*e^3)*EllipticE[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]]]))

```
Sqrt[a]]*x], -1]) + I*(Sqrt[c]*d*(Sqrt[c]*d - I*Sqrt[a]*e)*(A*c*d^2 + I*Sqr
t[a]*Sqrt[c]*d*(B*d - A*e) + a*e*(2*B*d - A*e))*EllipticF[I*ArcSinh[Sqrt[(I
*Sqrt[c])/Sqrt[a]]*x], -1] + a*e*(-5*B*c*d^3 + 7*A*c*d^2*e + a*B*d*e^2 + a*
A*e^3)*EllipticPi[(-I)*Sqrt[a]*e)/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/
Sqrt[a]]*x], -1]))/(2*a*Sqrt[(I*Sqrt[c])/Sqrt[a]]*(c*d^3 + a*d*e^2)^2*(d +
e*x^2)*Sqrt[a + c*x^4])
```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="fricas")
```

```
[Out] Timed out
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}(ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="giac")
```

```
[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)^2), x)
```

maple [C] time = 0.04, size = 1384, normalized size = 0.93

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x)
```

```
[Out] (A*e-B*d)/e*(1/2*e^4/(a*e^2+c*d^2)^2/d*x*(c*x^4+a)^(1/2)/(e*x^2+d)-2*c*(1/2
/a*c*d*e/(a*e^2+c*d^2)^2*x^3+1/4/a*(a*e^2-c*d^2)/(a*e^2+c*d^2)^2*x)/((x^4+a
/c)*c)^(1/2)-1/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(
I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*EllipticF((I/a^(1/2)*c^(1/2)
)^(1/2)*x,I)*e^2*c/(a*e^2+c*d^2)^2+1/2/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)
)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*Elli
pticF((I/a^(1/2)*c^(1/2))^(1/2)*x,I)/a*c^2/(a*e^2+c*d^2)^2*d^2-1/2*I*a^(1/2)
)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a^(1/2)*c^(1/2)*x^2+1)^(1/2)*(I/a^(1/2)*c^(
1/2)*x^2+1)^(1/2)/(c*x^4+a)^(1/2)*c^(1/2)*e^3/d/(a*e^2+c*d^2)^2*EllipticF((
I/a^(1/2)*c^(1/2))^(1/2)*x,I)+1/2*I*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(-I/a
```



```
[In] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^2), x)
```

```
[Out] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+a)**(3/2), x)
```

```
[Out] Timed out
```

$$3.14 \quad \int \frac{A+Bx^2}{(d+ex^2)^3 (a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=2452

result too large to display

```
[Out] -1/4*e^(3/2)*(a*e^2+3*c*d^2)*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(3/2)/(a*e^2+c*d^2)^(7/2)-3/16*e^(3/2)*(-A*e+B*d)*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(5/2)/(a*e^2+c*d^2)^(7/2)-1/2*c*e^(3/2)*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/(a*e^2+c*d^2)^(7/2)/d^(1/2)+1/2*c*x*(A*c*d*(-3*a*e^2+c*d^2)+a*B*e*(-a*e^2+3*c*d^2)+c*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3))*x^2/a/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)-1/4*e^3*(-A*e+B*d)*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^2/(e*x^2+d)^2-3/8*e^3*(-A*e+B*d)*(a*e^2+3*c*d^2)*x*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^3/(e*x^2+d)-1/2*e^3*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^3/(e*x^2+d)-1/2*c^(3/2)*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*x*(c*x^4+a)^(1/2)/a/(a*e^2+c*d^2)^3/(a^(1/2)+x^2*c^(1/2))+3/8*e^2*(-A*e+B*d)*(a*e^2+3*c*d^2)*x*c^(1/2)*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^3/(a^(1/2)+x^2*c^(1/2))+1/2*e^2*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*x*c^(1/2)*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^3/(a^(1/2)+x^2*c^(1/2))-3/8*a^(1/4)*c^(1/4)*e^2*(-A*e+B*d)*(a*e^2+3*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/d^2/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)-1/2*a^(1/4)*c^(1/4)*e^2*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/d/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)+1/2*c^(5/4)*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(3/4)/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)-1/2*c^(1/4)*e*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(1/4)/d/(a*e^2+c*d^2)^2/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+a)^(1/2)-1/2*c^(5/4)*e*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(1/4)/(a*e^2+c*d^2)^3/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+a)^(1/2)+1/8*e*(a*e^2+3*c*d^2)*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*2^(1/2))*(e*a^(1/2)
```

$$\begin{aligned}
& +d*c^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)} \\
& /a^{(1/4)}/c^{(1/4)}/d^2/(a*e^2+c*d^2)^3/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)} \\
& +3/32*e*(-A*e+B*d)*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)*(\cos(2*\arctan(c^{(1/4)}* \\
& x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan \\
& (c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)}, 1/ \\
& 2*2^{(1/2)}*(e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+ \\
& x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^3/(a*e^2+c*d^2)^3/(-e*a^{(1/2)}+d*c^{(\\
& 1/2)})/(c*x^4+a)^{(1/2)}+1/4*c^{(3/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)} \\
&)/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)} \\
&)), 1/2*2^{(1/2)}*(A*c^2*d^3-a^2*B*e^3+3*a*c*d*e*(-A*e+B*d)-c^{(3/2)}*d^2*(-3*A \\
& *e+B*d)*a^{(1/2)}+a^{(3/2)}*e^2*(-A*e+3*B*d)*c^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)}))*((c \\
& *x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(5/4)}/(a*e^2+c*d^2)^3/(c*x^4+a)^{(1 \\
& /2)}+1/4*a^{(3/4)}*c^{(3/4)}*e*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*(\cos(2* \\
& \arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{Elliptic} \\
& \text{Pi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(\\
& 1/2)}/c^{(1/2)}, 1/2*2^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*(e+d*c^{(1/2)}/a^{(1/2)})^2*((c \\
& *x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/d/(-a*e^2+c*d^2)/(a*e^2+c*d^2)^3/(c* \\
& x^4+a)^{(1/2)}-1/8*c^{(1/4)}*e*(-A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(\\
& 1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(\\
& 1/4)})), 1/2*2^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*(4*c*d^2+3*a*e^2-d*e*a^{(1/2)}*c^{(1 \\
& /2)}*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/d^2/(a*e^2+c*d^2)^2/ \\
& (-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}
\end{aligned}$$

Rubi [A] time = 3.97, antiderivative size = 2452, normalized size of antiderivative = 1.00, number of steps used = 22, number of rules used = 11, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.393$, Rules used = {1721, 1179, 1198, 220, 1196, 1224, 1697, 1715, 1709, 1707, 1217}

result too large to display

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^3*(a + c*x^4)^(3/2)), x]

[Out] (c*x*(A*c*d*(c*d^2 - 3*a*e^2) + a*B*e*(3*c*d^2 - a*e^2) + c*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*x^2))/((2*a*(c*d^2 + a*e^2)^3*sqrt[a + c*x^4]) + (3*sqrt[c]*e^2*(B*d - A*e)*(3*c*d^2 + a*e^2)*x*sqrt[a + c*x^4]))/(8*d^2*(c*d^2 + a*e^2)^3*(sqrt[a] + sqrt[c]*x^2)) + (sqrt[c]*e^2*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*x*sqrt[a + c*x^4])/((2*d*(c*d^2 + a*e^2)^3*(sqrt[a] + sqrt[c]*x^2)) - (c^(3/2)*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*x*sqrt[a + c*x^4])/((2*a*(c*d^2 + a*e^2)^3*(sqrt[a] + sqrt[c]*x^2)) - (e^3*(B*d - A*e)*x*sqrt[a + c*x^4]))/(4*d*(c*d^2 + a*e^2)^2*(d + e*x^2)^2) - (3*e^3*(B*d - A*e)*(3*c*d^2 + a*e^2)*x*sqrt[a + c*x^4])/((8*d^2*(c*d^2 + a*e^2)^3*(d + e*x^2)) - (e^3*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*x*sqrt[a + c*x^4]))/(2*d*(c*d^2 + a*e^2)^3*(d + e*x^2)) - (e^(3/2)*(3*c*d^2 + a*e^2)*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*ArcTan[(sqrt[c*d^2 + a*e^2]*x)/(sqrt[d]*sqrt[e]*sqrt[a + c*x^4])])/((4*d^(3/2)*(c*d^2 + a*e^2)^(7/2)) - (c*e^(3/2)*(B*c*d^3 - 3*A*c*d^2*e -

$$\begin{aligned}
& 3*a*B*d*e^2 + a*A*e^3)*ArcTan[(Sqrt[c*d^2 + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + c*x^4]))/(2*Sqrt[d]*(c*d^2 + a*e^2)^(7/2)) - (3*e^(3/2)*(B*d - A*e)* \\
& (5*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4)*ArcTan[(Sqrt[c*d^2 + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + c*x^4]))/(16*d^(5/2)*(c*d^2 + a*e^2)^(7/2)) - (3*a^(1/4)*c^(1/4)*e^2*(B*d - A*e)*(3*c*d^2 + a*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(8*d^2*(c*d^2 + a*e^2)^3*Sqrt[a + c*x^4]) - (a^(1/4)*c^(1/4)*e^2*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*d*(c*d^2 + a*e^2)^3*Sqrt[a + c*x^4]) + (c^(5/4)*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*(c*d^2 + a*e^2)^3*Sqrt[a + c*x^4]) - (c^(1/4)*e*(B*d - A*e)*(4*c*d^2 - Sqrt[a]*Sqrt[c]*d*e + 3*a*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(8*a^(1/4)*d^2*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 + a*e^2)^2*Sqrt[a + c*x^4]) - (c^(1/4)*e*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*d*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 + a*e^2)^2*Sqrt[a + c*x^4]) - (c^(5/4)*e*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 + a*e^2)^3*Sqrt[a + c*x^4]) + (c^(3/4)*(A*c^2*d^3 - a^2*B*e^3 - Sqrt[a]*c^(3/2)*d^2*(B*d - 3*A*e) + 3*a*c*d*e*(B*d - A*e) + a^(3/2)*Sqrt[c]*e^2*(3*B*d - A*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(5/4)*(c*d^2 + a*e^2)^3*Sqrt[a + c*x^4]) + (e*(Sqrt[c]*d + Sqrt[a]*e)*(3*c*d^2 + a*e^2)*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-(Sqrt[c]*d - Sqrt[a]*e)^2/(4*Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(8*a^(1/4)*c^(1/4)*d^2*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 + a*e^2)^3*Sqrt[a + c*x^4]) + (a^(3/4)*c^(3/4)*e*((Sqrt[c]*d)/Sqrt[a] + e)^2*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-(Sqrt[c]*d - Sqrt[a]*e)^2/(4*Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*d*(c*d^2 - a*e^2)*(c*d^2 + a*e^2)^3*Sqrt[a + c*x^4]) + (3*e*(Sqrt[c]*d + Sqrt[a]*e)*(B*d - A*e)*(5*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-(Sqrt[c]*d - Sqrt[a]*e)^2/(4*Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(32*a^(1/4)*c^(1/4)*d^3*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 + a*e^2)^3*Sqrt[a + c*x^4])
\end{aligned}$$

Rule 220

```

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[
(1 + q^2*x^2)*Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x
, 1/2]]/(2*q*Sqrt[a + b*x^4]), x]] /; FreeQ[{a, b}, x] && PosQ[b/a]

```


Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := -Simp[(x
*(d + e*x^2)*(a + c*x^4)^(p + 1))/(4*a*(p + 1)), x] + Dist[1/(4*a*(p + 1)),
  Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /;
FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2
*p]
```

Rule 1196

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(
  1 + q^2*x^2)*Sqrt[a + c*x^4]/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x],
  1/2])/(q*Sqrt[a + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e},
  x] && PosQ[c/a]
```

Rule 1198

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c,
  d, e}, x] && PosQ[c/a]
```

Rule 1217

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[c/a, 2]}, Dist[(c*d + a*e*q)/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4]
, x], x] - Dist[(a*e*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^
2)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2
, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
```

Rule 1224

```
Int[((d_) + (e_.)*(x_)^2)^(q_)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := -Sim
p[(e^2*x*(d + e*x^2)^(q + 1)*Sqrt[a + c*x^4])/(2*d*(q + 1)*(c*d^2 + a*e^2))
, x] + Dist[1/(2*d*(q + 1)*(c*d^2 + a*e^2)), Int[((d + e*x^2)^(q + 1)*Simp[
a*e^2*(2*q + 3) + 2*c*d^2*(q + 1) - 2*e*c*d*(q + 1)*x^2 + c*e^2*(2*q + 5)*x
^4, x])/Sqrt[a + c*x^4], x], x] /; FreeQ[{a, c, d, e}, x] && ILtQ[q, -1]
```

Rule 1697

```
Int[((P4x_)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol
] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]
}, -Simp[((C*d^2 - B*d*e + A*e^2)*x*(d + e*x^2)^(q + 1)*Sqrt[a + c*x^4])/(2
```

```
*d*(q + 1)*(c*d^2 + a*e^2)), x] + Dist[1/(2*d*(q + 1)*(c*d^2 + a*e^2)), Int
[((d + e*x^2)^(q + 1)*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*c*d^2*(
q + 1)) + 2*d*(B*c*d - A*c*e + a*C*e)*(q + 1)*x^2 + c*(C*d^2 - B*d*e + A*e^
2)*(2*q + 5)*x^4, x])/Sqrt[a + c*x^4], x], x] /; FreeQ[{a, c, d, e}, x] &&
PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[c*d^2 + a*e^2, 0] && ILtQ[
q, -1]
```

Rule 1707

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[(Rt[(c*d)/e
+ (a*e)/d, 2]*x)/Sqrt[a + c*x^4]])/(2*d*e*Rt[(c*d)/e + (a*e)/d, 2]), x] +
Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a + c*x^4))/(a*(A + B*x^2)^2)]*Ell
ipticPi[Cancel[-((B*d - A*e)^2/(4*d*e*A*B))], 2*ArcTan[q*x], 1/2])/(4*d*e*A
*q*Sqrt[a + c*x^4]), x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e
^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1709

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q)
)/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4], x], x] + Dist[(a*(B*d - A*e)*(e
+ d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x],
x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2
- a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rule 1715

```
Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] :=
With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff
[P4x, x, 4]}, -Dist[C/(e*q), Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] + Dist
[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x^2)
)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2, 2]
&& NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
```

Rule 1721

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^3 (a + cx^4)^{3/2}} dx &= \int \left(\frac{c (Acd (cd^2 - 3ae^2) + aBe (3cd^2 - ae^2) + c (Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{(cd^2 + ae^2)^3 (a + cx^4)^{3/2}} \right. \\
&= \frac{c \int \frac{Acd(cd^2-3ae^2)+aBe(3cd^2-ae^2)+c(Bcd^3-3Acd^2e-3aBde^2+aAe^3)x^2}{(a+cx^4)^{3/2}} dx}{(cd^2 + ae^2)^3} - \frac{(e(Bd - Ae)) \int \frac{1}{(d+ex^2)^3} dx}{cd^2 + ae^2} \\
&= \frac{cx (Acd (cd^2 - 3ae^2) + aBe (3cd^2 - ae^2) + c (Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a (cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx (Acd (cd^2 - 3ae^2) + aBe (3cd^2 - ae^2) + c (Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a (cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx (Acd (cd^2 - 3ae^2) + aBe (3cd^2 - ae^2) + c (Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a (cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx (Acd (cd^2 - 3ae^2) + aBe (3cd^2 - ae^2) + c (Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a (cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx (Acd (cd^2 - 3ae^2) + aBe (3cd^2 - ae^2) + c (Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a (cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx (Acd (cd^2 - 3ae^2) + aBe (3cd^2 - ae^2) + c (Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a (cd^2 + ae^2)^3 \sqrt{a + cx^4}}
\end{aligned}$$

Mathematica [C] time = 2.74, size = 630, normalized size = 0.26

$$\frac{dx \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \left(4cd^2 (d + ex^2)^2 \left(B(-a^2e^3 + 3acde(d - ex^2) + c^2d^3x^2) + Ac(ae^2(ex^2 - 3d) + cd^2(d - 3ex^2)) \right) - 2ade \right)}{2a (cd^2 + ae^2)^3 \sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^3*(a + c*x^4)^(3/2)), x]

```
[Out] (Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*x*(-2*a*d*e^3*(B*d - A*e)*(c*d^2 + a*e^2)*(a +
c*x^4) + a*e^3*(-13*B*c*d^3 + 17*A*c*d^2*e + a*B*d*e^2 + 3*a*A*e^3)*(d + e
*x^2)*(a + c*x^4) + 4*c*d^2*(d + e*x^2)^2*(B*(-(a^2*e^3) + c^2*d^3*x^2 + 3*
a*c*d*e*(d - e*x^2)) + A*c*(c*d^2*(d - 3*e*x^2) + a*e^2*(-3*d + e*x^2)))) -
(d + e*x^2)^2*Sqrt[1 + (c*x^4)/a]*(Sqrt[a]*Sqrt[c]*d*(3*A*e*(-4*c^2*d^4 +
7*a*c*d^2*e^2 + a^2*e^4) + B*(4*c^2*d^5 - 25*a*c*d^3*e^2 + a^2*d*e^4))*Elli
pticE[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + I*(Sqrt[c]*d*(Sqrt[c]*d
- I*Sqrt[a]*e)*(4*A*c^2*d^4 + (4*I)*Sqrt[a]*c^(3/2)*d^3*(B*d - 2*A*e) + 19
*a*c*d^2*e*(B*d - A*e) - (2*I)*a^(3/2)*Sqrt[c]*d*e^2*(3*B*d - A*e) - a^2*e^
3*(B*d + 3*A*e))*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + a*
e*(3*A*e*(21*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4) + B*(-35*c^2*d^5 + 26*a*c*d
^3*e^2 + a^2*d*e^4))*EllipticPi[(-I)*Sqrt[a]*e/(Sqrt[c]*d), I*ArcSinh[Sqr
t[(I*Sqrt[c])/Sqrt[a]]*x], -1)])/(8*a*Sqrt[(I*Sqrt[c])/Sqrt[a]]*(c*d^3 + a
*d*e^2)^3*(d + e*x^2)^2*Sqrt[a + c*x^4])
```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="fricas")
```

```
[Out] Timed out
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}(ex^2 + d)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="giac")
```

```
[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)^3), x)
```

maple [C] time = 0.05, size = 2326, normalized size = 0.95

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x)
```

```
[Out] B/e*(1/2/(a*e^2+c*d^2)^2*(c*x^4+a)^(1/2)/(e*x^2+d)/d*e^4*x-2*(1/2/(a*e^2+c*
d^2)^2/a*c*d*e*x^3+1/4*(a*e^2-c*d^2)/(a*e^2+c*d^2)^2/a*x)/((x^4+a/c)*c)^(1/
```

$$\begin{aligned}
& 2) * c - 1 / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} \\
&) * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * \text{EllipticF}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * \\
& x, I) * e^2 * c / (a * e^2 + c * d^2)^2 + 1/2 / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} \\
&) * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * \text{EllipticF}((I \\
& / a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) / a * c^2 / (a * e^2 + c * d^2)^2 * d^2 - 1/2 * I * a^{(1/2)} / (I/a^{(1/2)} \\
&) * c^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 \\
& + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * c^{(1/2)} * e^3 / d / (a * e^2 + c * d^2)^2 * \text{EllipticF}((I/a^{(1/2)} \\
&) * c^{(1/2)})^{(1/2)} * x, I) + 1/2 * I * a^{(1/2)} / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c \\
& ^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * c^{(1/2)} \\
& * e^3 / d / (a * e^2 + c * d^2)^2 * \text{EllipticE}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) + I/a^{(1/2)} / (\\
& I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} \\
&) * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * c^{(3/2)} * d * e / (a * e^2 + c * d^2)^2 * \text{EllipticF}((I/a^{(1/2)} \\
&) * c^{(1/2)})^{(1/2)} * x, I) - I/a^{(1/2)} / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * c^{(1/2)} \\
&) * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * c^{(3/2)} * \\
& d * e / (a * e^2 + c * d^2)^2 * \text{EllipticE}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) + 1/2 * e^4 / d^2 / (a \\
& * e^2 + c * d^2)^2 / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I \\
& / a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * \text{EllipticPi}((I/a^{(1/2)} * c^{(1/2)} \\
&)^{(1/2)} * x, I * a^{(1/2)} / c^{(1/2)} / d * e, (-I/a^{(1/2)} * c^{(1/2)})^{(1/2)} / (I/a^{(1/2)} * c^{(1/2)} \\
&)^{(1/2)}) * a + 7/2 * e^2 / (a * e^2 + c * d^2)^2 / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * \\
& c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * \text{Elliptic} \\
& \text{Pi}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I * a^{(1/2)} / c^{(1/2)} / d * e, (-I/a^{(1/2)} * c^{(1/2)}) \\
& ^{(1/2)} / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)}) * c) + (A * e - B * d) / e * (1/4 * e^4 / (a * e^2 + c * d^2)^2 / d \\
& * x * (c * x^4 + a)^{(1/2)} / (e * x^2 + d)^2 + 1/8 * e^4 * (3 * a * e^2 + 17 * c * d^2) / d^2 / (a * e^2 + c * d^2) \\
& ^3 * x * (c * x^4 + a)^{(1/2)} / (e * x^2 + d) - 2 * c * (-1/4 * c * e * (a * e^2 - 3 * c * d^2) / a / (a * e^2 + c * d^2) \\
&)^3 * x^3 + 1/4 * c * d * (3 * a * e^2 - c * d^2) / a / (a * e^2 + c * d^2)^3 * x) / ((x^4 + a/c) * c)^{(1/2)} - 1/ \\
& 8 / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * c^{(1/2)} \\
&) * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * \text{EllipticF}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) * \\
& c * e^4 / d / (a * e^2 + c * d^2)^3 * a - 27/8 / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} \\
&) * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * \text{EllipticF}((I \\
& / a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) * c^2 * e^2 * d / (a * e^2 + c * d^2)^3 + 1/2 / (I/a^{(1/2)} * c^{(1/2)} \\
&)^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / \\
& (c * x^4 + a)^{(1/2)} * \text{EllipticF}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) * c^3 * d^3 / a / (a * e^2 + c \\
& * d^2)^3 + 3/2 * I/a^{(1/2)} / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} \\
&) * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * c^{(5/2)} * e / (a * e^2 + c * d^2) \\
& ^3 * d^2 * \text{EllipticF}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) + 21/8 * I * a^{(1/2)} / (I/a^{(1/2)} \\
&) * c^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} \\
&) / (c * x^4 + a)^{(1/2)} * c^{(3/2)} * e^3 / (a * e^2 + c * d^2)^3 * \text{EllipticE}((I/a^{(1/2)} * c^{(1/2)} \\
&)^{(1/2)} * x, I) + 3/8 * I * a^{(3/2)} / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} \\
&) * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * c^{(1/2)} * e^5 / d \\
& ^2 / (a * e^2 + c * d^2)^3 * \text{EllipticE}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) - 21/8 * I * a^{(1/2)} / \\
& (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * c^{(1/2)} \\
&) * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * c^{(3/2)} * e^3 / (a * e^2 + c * d^2)^3 * \text{EllipticF}((I/a^{(1/2)} \\
&) * c^{(1/2)})^{(1/2)} * x, I) - 3/2 * I/a^{(1/2)} / (I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * (-I/a^{(1/2)} (1/ \\
& 2) * c^{(1/2)} * x^2 + 1)^{(1/2)} * (I/a^{(1/2)} * c^{(1/2)} * x^2 + 1)^{(1/2)} / (c * x^4 + a)^{(1/2)} * c^{(5/2)} * e / (a * e^2 + c * d^2)^3 * d^2 * \text{EllipticE}((I/a^{(1/2)} * c^{(1/2)})^{(1/2)} * x, I) - 3/8 * I * a
\end{aligned}$$

$$\begin{aligned} & \frac{1}{c^{3/2}} \frac{1}{(I/a^{1/2} * c^{1/2})^{1/2}} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (c * x^4 + a)^{1/2} * c^{1/2} * e^{5/d^2} / (a * e^2 + c * d^2)^3 * \text{EllipticF}((I/a^{1/2} * c^{1/2})^{1/2} * x, I) + 3/8 * e^6/d^3 / (a * e^2 + c * d^2)^3 / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (c * x^4 + a)^{1/2} * \text{EllipticPi}((I/a^{1/2} * c^{1/2})^{1/2} * x, I * a^{1/2} / c^{1/2} / d * e, (-I/a^{1/2} * c^{1/2})^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2}) * a^2 + 3/4 * e^4 / (a * e^2 + c * d^2)^3 / d / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (c * x^4 + a)^{1/2} * \text{EllipticPi}((I/a^{1/2} * c^{1/2})^{1/2} * x, I * a^{1/2} / c^{1/2} / d * e, (-I/a^{1/2} * c^{1/2})^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2}) * a * c + 63/8 * e^2 * d / (a * e^2 + c * d^2)^3 / (I/a^{1/2} * c^{1/2})^{1/2} * (-I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} * (I/a^{1/2} * c^{1/2} * x^2 + 1)^{1/2} / (c * x^4 + a)^{1/2} * \text{EllipticPi}((I/a^{1/2} * c^{1/2})^{1/2} * x, I * a^{1/2} / c^{1/2} / d * e, (-I/a^{1/2} * c^{1/2})^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2}) * c^2 \end{aligned}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{3/2} (ex^2 + d)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{3/2} (ex^2 + d)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^3),x)

[Out] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+a)**(3/2),x)

[Out] Timed out

$$3.15 \quad \int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx$$

Optimal. Leaf size=169

$$\frac{x \left(A - \frac{\sqrt{-a}B}{\sqrt{c}} \right) (d + ex^2)^q \left(\frac{ex^2}{d} + 1 \right)^{-q} F_1 \left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a} + \frac{x \left(\frac{\sqrt{-a}B}{\sqrt{c}} + A \right) (d + ex^2)^q \left(\frac{ex^2}{d} + 1 \right)^{-q} F_1 \left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a}$$

[Out] $1/2*x*(e*x^2+d)^q*AppellF1(1/2, -q, 1, 3/2, -e*x^2/d, -x^2*c^(1/2)/(-a)^(1/2))* (A-B*(-a)^(1/2)/c^(1/2))/a/((1+e*x^2/d)^q)+1/2*x*(e*x^2+d)^q*AppellF1(1/2, 1, -q, 3/2, x^2*c^(1/2)/(-a)^(1/2), -e*x^2/d)*(A+B*(-a)^(1/2)/c^(1/2))/a/((1+e*x^2/d)^q)$

Rubi [A] time = 0.22, antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {1693, 430, 429}

$$\frac{x \left(A - \frac{\sqrt{-a}B}{\sqrt{c}} \right) (d + ex^2)^q \left(\frac{ex^2}{d} + 1 \right)^{-q} F_1 \left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a} + \frac{x \left(\frac{\sqrt{-a}B}{\sqrt{c}} + A \right) (d + ex^2)^q \left(\frac{ex^2}{d} + 1 \right)^{-q} F_1 \left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x]

[Out] $((A - (\text{Sqrt}[-a]*B)/\text{Sqrt}[c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, -((\text{Sqrt}[c]*x^2)/\text{Sqrt}[-a]), -((e*x^2)/d)])/(2*a*(1 + (e*x^2)/d)^q) + ((A + (\text{Sqrt}[-a]*B)/\text{Sqrt}[c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (\text{Sqrt}[c]*x^2)/\text{Sqrt}[-a], -((e*x^2)/d)])/(2*a*(1 + (e*x^2)/d)^q)$

Rule 429

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, -((b*x^n)/a), -((d*x^n)/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 430

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])

Rule 1693

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx &= \int \left(-\frac{(\sqrt{-a}B + A\sqrt{c})(d + ex^2)^q}{2\sqrt{-a}\sqrt{c}(\sqrt{-a} - \sqrt{c}x^2)} + \frac{(\sqrt{-a}B - A\sqrt{c})(d + ex^2)^q}{2\sqrt{-a}\sqrt{c}(\sqrt{-a} + \sqrt{c}x^2)} \right) dx \\ &= -\left(\frac{1}{2} \left(\frac{A}{\sqrt{-a}} + \frac{B}{\sqrt{c}} \right) \int \frac{(d + ex^2)^q}{\sqrt{-a} - \sqrt{c}x^2} dx \right) + \frac{1}{2} \left(\frac{aA}{(-a)^{3/2}} + \frac{B}{\sqrt{c}} \right) \int \frac{(d + ex^2)^q}{\sqrt{-a} + \sqrt{c}x^2} dx \\ &= -\left(\frac{1}{2} \left(\left(\frac{A}{\sqrt{-a}} + \frac{B}{\sqrt{c}} \right) (d + ex^2)^q \left(1 + \frac{ex^2}{d} \right)^{-q} \right) \int \frac{\left(1 + \frac{ex^2}{d} \right)^q}{\sqrt{-a} - \sqrt{c}x^2} dx \right) + \frac{1}{2} \left(\left(\frac{aA}{(-a)^{3/2}} + \right. \right. \\ &= \frac{\left(A - \frac{\sqrt{-a}B}{\sqrt{c}} \right) x (d + ex^2)^q \left(1 + \frac{ex^2}{d} \right)^{-q} F_1 \left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a} - \frac{\left(\frac{A}{\sqrt{-a}} + \frac{B}{\sqrt{c}} \right) x (d + ex^2)^q \left(1 + \frac{ex^2}{d} \right)^{-q} F_1 \left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a} \end{aligned}$$

Mathematica [F] time = 0.38, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx$$

Verification is Not applicable to the result.

```
[In] Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x]
```

```
[Out] Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x]
```

fricas [F] time = 1.21, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a), x, algorithm="fricas")
```

```
[Out] integral((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + a), x)
```


giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + a), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x)

[Out] int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)*(e*x**2+d)**q/(c*x**4+a),x)
```

```
[Out] Timed out
```

$$3.16 \quad \int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx$$

Optimal. Leaf size=48

$$\frac{\sqrt{2} (x^2 + 2) E\left(\tan^{-1}(x) \middle| \frac{1}{2}\right)}{\sqrt{\frac{x^2+2}{x^2+1}} \sqrt{x^4 + 3x^2 + 2}}$$

[Out] $(x^2+2)*(1/(x^2+1))^{(1/2)}*(x^2+1)^{(1/2)}*EllipticE(x/(x^2+1)^{(1/2)}, 1/2*2^{(1/2)})*2^{(1/2)}/((x^2+2)/(x^2+1))^{(1/2)}/(x^4+3*x^2+2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$, Rules used = {1456, 411}

$$\frac{\sqrt{2} (x^2 + 2) E\left(\tan^{-1}(x) \middle| \frac{1}{2}\right)}{\sqrt{\frac{x^2+2}{x^2+1}} \sqrt{x^4 + 3x^2 + 2}}$$

Antiderivative was successfully verified.

[In] Int[(2 + x^2)/((1 + x^2)*Sqrt[2 + 3*x^2 + x^4]), x]

[Out] (Sqrt[2]*(2 + x^2)*EllipticE[ArcTan[x], 1/2])/(Sqrt[(2 + x^2)/(1 + x^2)]*Sqrt[2 + 3*x^2 + x^4])

Rule 411

Int[Sqrt[(a_) + (b_)*(x_)^2]/((c_) + (d_)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - (b*c)/(a*d)])/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(a + b*x^2))/(a*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 1456

Int[((d_) + (e_)*(x_)^(n_))^(q_)*((f_) + (g_)*(x_)^(n_))^(r_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Dist[(a + b*x^n + c*x^(2*n))^FracPart[p]/((d + e*x^n)^FracPart[p]*(a/d + (c*x^n)/e)^FracPart[p]), Int[(d + e*x^n)^(p + q)*(f + g*x^n)^r*(a/d + (c*x^n)/e)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q, r}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && !IntegerQ[p]

Rubi steps

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx = \frac{(\sqrt{1+x^2}\sqrt{2+x^2}) \int \frac{\sqrt{2+x^2}}{(1+x^2)^{3/2}} dx}{\sqrt{2+3x^2+x^4}}$$

$$= \frac{\sqrt{2} (2+x^2) E\left(\tan^{-1}(x)\middle|\frac{1}{2}\right)}{\sqrt{\frac{2+x^2}{1+x^2}} \sqrt{2+3x^2+x^4}}$$

Mathematica [C] time = 0.13, size = 94, normalized size = 1.96

$$\frac{x^3 - i\sqrt{x^2+1}\sqrt{x^2+2}F\left(i\sinh^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right) + i\sqrt{x^2+1}\sqrt{x^2+2}E\left(i\sinh^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right) + 2x}{\sqrt{x^4+3x^2+2}}$$

Antiderivative was successfully verified.

[In] Integrate[(2 + x^2)/((1 + x^2)*Sqrt[2 + 3*x^2 + x^4]), x]

[Out] (2*x + x^3 + I*Sqrt[1 + x^2]*Sqrt[2 + x^2]*EllipticE[I*ArcSinh[x/Sqrt[2]], 2] - I*Sqrt[1 + x^2]*Sqrt[2 + x^2]*EllipticF[I*ArcSinh[x/Sqrt[2]], 2])/Sqrt[2 + 3*x^2 + x^4]

fricas [F] time = 1.10, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{x^4+3x^2+2}}{x^4+2x^2+1}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(x^4 + 3*x^2 + 2)/(x^4 + 2*x^2 + 1), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2+2}{\sqrt{x^4+3x^2+2}(x^2+1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+2)/(sqrt(x^4+3*x^2+2)*(x^2+1)), x, algorithm="giac")

[Out] integrate((x^2 + 2)/(sqrt(x^4 + 3*x^2 + 2)*(x^2 + 1)), x)

maple [C] time = 0.03, size = 81, normalized size = 1.69

$$\frac{(x^2 + 2)x}{\sqrt{(x^2 + 1)(x^2 + 2)}} - \frac{i\sqrt{2} \sqrt{2x^2 + 4} \sqrt{x^2 + 1} \left(-\text{EllipticE}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right) + \text{EllipticF}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right) \right)}{2\sqrt{x^4 + 3x^2 + 2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2), x)

[Out] (x^2+2)*x/((x^2+1)*(x^2+2))^(1/2)-1/2*I*2^(1/2)*(2*x^2+4)^(1/2)*(x^2+1)^(1/2)/(x^4+3*x^2+2)^(1/2)*(EllipticF(1/2*I*x*2^(1/2), 2^(1/2))-EllipticE(1/2*I*x*2^(1/2), 2^(1/2)))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2 + 2}{\sqrt{x^4 + 3x^2 + 2}(x^2 + 1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2), x, algorithm="maxima")

[Out] integrate((x^2 + 2)/(sqrt(x^4 + 3*x^2 + 2)*(x^2 + 1)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{x^2 + 2}{(x^2 + 1) \sqrt{x^4 + 3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2 + 2)/((x^2 + 1)*(3*x^2 + x^4 + 2)^(1/2)), x)

[Out] int((x^2 + 2)/((x^2 + 1)*(3*x^2 + x^4 + 2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2 + 2}{\sqrt{(x^2 + 1)(x^2 + 2)}(x^2 + 1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((x**2+2)/(x**2+1)/(x**4+3*x**2+2)**(1/2), x)

[Out] Integral((x**2 + 2)/(sqrt((x**2 + 1)*(x**2 + 2))*(x**2 + 1)), x)

$$3.17 \quad \int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=755

$$\frac{ex\sqrt{a+bx^2+cx^4} \left(B(-ce(25ae+84bd)+24b^2e^2+105c^2d^2) + 7Ace(15cd-4be) \right)}{105c^3} + \frac{x\sqrt{a+bx^2+cx^4} \left(7Ace(-3c$$

[Out] 1/105*e*(7*A*c*e*(-4*b*e+15*c*d)+B*(105*c^2*d^2+24*b^2*e^2-c*e*(25*a*e+84*b*d)))*x*(c*x^4+b*x^2+a)^(1/2)/c^3+1/35*e^2*(7*A*c*e-6*B*b*e+21*B*c*d)*x^3*(c*x^4+b*x^2+a)^(1/2)/c^2+1/7*B*e^3*x^5*(c*x^4+b*x^2+a)^(1/2)/c+1/105*(7*A*c*e*(45*c^2*d^2+8*b^2*e^2-3*c*e*(3*a*e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^3-21*c^2*d*e*(9*a*e+10*b*d)+8*b*c*e^2*(13*a*e+21*b*d)))*x*(c*x^4+b*x^2+a)^(1/2)/c^(7/2)/(a^(1/2)+x^2*c^(1/2))-1/105*a^(1/4)*(7*A*c*e*(45*c^2*d^2+8*b^2*e^2-3*c*e*(3*a*e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^3-21*c^2*d*e*(9*a*e+10*b*d)+8*b*c*e^2*(13*a*e+21*b*d)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(15/4)/(c*x^4+b*x^2+a)^(1/2)+1/210*a^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*(7*A*c*e*(45*c^2*d^2+8*b^2*e^2-3*c*e*(3*a*e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^3-21*c^2*d*e*(9*a*e+10*b*d)+8*b*c*e^2*(13*a*e+21*b*d)))+(7*A*c*(4*a*b*e^3-15*a*c*d*e^2+15*c^2*d^3)-a*B*e*(105*c^2*d^2+24*b^2*e^2-c*e*(25*a*e+84*b*d)))*c^(1/2)/a^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(15/4)/(c*x^4+b*x^2+a)^(1/2)

Rubi [A] time = 1.36, antiderivative size = 755, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$, Rules used = {1679, 1197, 1103, 1195}

$$\frac{ex\sqrt{a+bx^2+cx^4} \left(B(-ce(25ae+84bd)+24b^2e^2+105c^2d^2) + 7Ace(15cd-4be) \right)}{105c^3} + \frac{x\sqrt{a+bx^2+cx^4} \left(7Ace(-3c$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + b*x^2 + c*x^4], x]

[Out] (e*(7*A*c*e*(15*c*d - 4*b*e) + B*(105*c^2*d^2 + 24*b^2*e^2 - c*e*(84*b*d + 25*a*e)))*x*Sqrt[a + b*x^2 + c*x^4])/(105*c^3) + (e^2*(21*B*c*d - 6*b*B*e +

$$\begin{aligned}
& 7*A*c*e)*x^3*\text{Sqrt}[a + b*x^2 + c*x^4])/(35*c^2) + (B*e^3*x^5*\text{Sqrt}[a + b*x^2 \\
& + c*x^4])/(7*c) + ((7*A*c*e*(45*c^2*d^2 + 8*b^2*e^2 - 3*c*e*(10*b*d + 3*a* \\
& e)) + B*(105*c^3*d^3 - 48*b^3*e^3 - 21*c^2*d*e*(10*b*d + 9*a*e) + 8*b*c*e^2 \\
& *(21*b*d + 13*a*e)))*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(105*c^(7/2)*(\text{Sqrt}[a] + \text{Sqr} \\
& \text{t}[c]*x^2)) - (a^(1/4)*(7*A*c*e*(45*c^2*d^2 + 8*b^2*e^2 - 3*c*e*(10*b*d + 3* \\
& a*e)) + B*(105*c^3*d^3 - 48*b^3*e^3 - 21*c^2*d*e*(10*b*d + 9*a*e) + 8*b*c*e \\
& ^2*(21*b*d + 13*a*e)))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sq} \\
& \text{rt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sq} \\
& \text{rt}[a]*\text{Sqrt}[c]))/4])/(105*c^(15/4)*\text{Sqrt}[a + b*x^2 + c*x^4]) + (a^(1/4)*(7*A* \\
& c*e*(45*c^2*d^2 + 8*b^2*e^2 - 3*c*e*(10*b*d + 3*a*e)) + B*(105*c^3*d^3 - 48 \\
& *b^3*e^3 - 21*c^2*d*e*(10*b*d + 9*a*e) + 8*b*c*e^2*(21*b*d + 13*a*e)) + (\text{Sq} \\
& \text{rt}[c]*(7*A*c*(15*c^2*d^3 - 15*a*c*d*e^2 + 4*a*b*e^3) - a*B*e*(105*c^2*d^2 + \\
& 24*b^2*e^2 - c*e*(84*b*d + 25*a*e))))/\text{Sqrt}[a]*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqr} \\
& \text{t}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^(1/4) \\
&)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(210*c^(15/4)*\text{Sqrt}[a + b*x^2 + \\
& c*x^4])
\end{aligned}$$

Rule 1103

$$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[\{(1 + q^2*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2 - (b*q^2)/(4*c)]\}/(2*q*\text{Sqrt}[a + b*x^2 + c*x^4]), x]] \text{ /; FreeQ}[\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{PosQ}[c/a]$$

Rule 1195

$$\text{Int}[\{(d_) + (e_)*(x_)^2\}/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 4]\}, -\text{Simp}[(d*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(a*(1 + q^2*x^2)), x] + \text{Simp}[(d*(1 + q^2*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2 - (b*q^2)/(4*c)]\}/(q*\text{Sqrt}[a + b*x^2 + c*x^4]), x] \text{ /; EqQ}[e + d*q^2, 0] \text{ /; FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{PosQ}[c/a]$$

Rule 1197

$$\text{Int}[\{(d_) + (e_)*(x_)^2\}/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d*q)/q, \text{Int}[1/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] - \text{Dist}[e/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] \text{ /; NeQ}[e + d*q, 0] \text{ /; FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{PosQ}[c/a]$$

Rule 1679

$$\text{Int}[(Pq_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] \text{ :> With}[\{q = \text{Expon}[Pq, x^2], e = \text{Coeff}[Pq, x^2, \text{Expon}[Pq, x^2]]\}, \text{Simp}[(e*x^(2*q - 3))*(a + b*x^2 + c*x^4)^(p + 1)]/(c*(2*q + 4*p + 1)), x] + \text{Dist}[1/(c*(2*q + 4*p$$

+ 1)), Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*q + 4*p + 1)*Pq - a*e*(2*q - 3)*x^(2*q - 4) - b*e*(2*q + 2*p - 1)*x^(2*q - 2) - c*e*(2*q + 4*p + 1)*x^(2*q), x], x] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pq, x^2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]

Rubi steps

$$\begin{aligned}
 \int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx &= \frac{Be^3x^5\sqrt{a + bx^2 + cx^4}}{7c} + \frac{\int \frac{7Acd^3 + 7cd^2(Bd + 3Ae)x^2 + e(21Bcd^2 + 21Acde - 5aBe^2)x^4 + e^2(21Bcd - 6bBe + 7Ace)x^6}{\sqrt{a + bx^2 + cx^4}} dx}{7c} \\
 &= \frac{e^2(21Bcd - 6bBe + 7Ace)x^3\sqrt{a + bx^2 + cx^4}}{35c^2} + \frac{Be^3x^5\sqrt{a + bx^2 + cx^4}}{7c} + \frac{\int \frac{35Ac^2d^3 + (7Acd^2(Bd + 3Ae) + e(21Bcd^2 + 21Acde - 5aBe^2))x^2 + e^2(21Bcd - 6bBe + 7Ace)x^4}{\sqrt{a + bx^2 + cx^4}} dx}{105c^3} \\
 &= \frac{e(7Ace(15cd - 4be) + B(105c^2d^2 + 24b^2e^2 - ce(84bd + 25ae)))x\sqrt{a + bx^2 + cx^4}}{105c^3} \\
 &= \frac{e(7Ace(15cd - 4be) + B(105c^2d^2 + 24b^2e^2 - ce(84bd + 25ae)))x\sqrt{a + bx^2 + cx^4}}{105c^3} \\
 &= \frac{e(7Ace(15cd - 4be) + B(105c^2d^2 + 24b^2e^2 - ce(84bd + 25ae)))x\sqrt{a + bx^2 + cx^4}}{105c^3}
 \end{aligned}$$

Mathematica [C] time = 6.48, size = 4473, normalized size = 5.92

Result too large to show

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + b*x^2 + c*x^4], x]

[Out] Sqrt[a + b*x^2 + c*x^4]*(-1/105*(e*(-105*B*c^2*d^2 + 84*b*B*c*d*e - 105*A*c^2*d*e - 24*b^2*B*e^2 + 28*A*b*c*e^2 + 25*a*B*c*e^2)*x)/c^3 + (e^2*(21*B*c*d - 6*b*B*e + 7*A*c*e)*x^3)/(35*c^2) + (B*e^3*x^5)/(7*c)) + (((((105*I)/2)*B*c^2*(-b + Sqrt[b^2 - 4*a*c])*d^3*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b^2 - 4*a*c])])*Sqrt[1 - (2*c*x^2)/(-b + Sqrt[b^2 - 4*a*c])])*(EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c])])]*x], (-b - Sqrt[b^2 - 4*a*c])/(-b + Sqrt[b^2 - 4*a*c])) - EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c])])]*x], (-b - Sqrt[b^2 - 4*a*c])/(-b + Sqrt[b^2 - 4*a*c])))/(Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c])])*Sqrt[a + b*x^2 + c*x^4]) - ((105*I)*b*


```

*a*c))/(-b + Sqrt[b^2 - 4*a*c])) - EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[-(c/(-b
- Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c))/(-b + Sqrt[b^2 - 4*a*c
])))/(c*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a + b*x^2 + c*x^4]) - (((
63*I)/2)*a*A*c*(-b + Sqrt[b^2 - 4*a*c])*e^3*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b
^2 - 4*a*c]])*Sqrt[1 - (2*c*x^2)/(-b + Sqrt[b^2 - 4*a*c]])*(EllipticE[I*Arc
Sinh[Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c
])/(-b + Sqrt[b^2 - 4*a*c])) - EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[-(c/(-b - S
qrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c))/(-b + Sqrt[b^2 - 4*a*c]))
)/(Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a + b*x^2 + c*x^4]) - (
(105*I)*A*c^3*d^3*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b^2 - 4*a*c]])*Sqrt[1 - (2*
c*x^2)/(-b + Sqrt[b^2 - 4*a*c]])*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[-(c/(-b -
Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c))/(-b + Sqrt[b^2 - 4*a*c])
)]/(Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a + b*x^2 + c*x^4]) +
((105*I)*a*B*c^2*d^2*e*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b^2 - 4*a*c]])*Sqrt[1
- (2*c*x^2)/(-b + Sqrt[b^2 - 4*a*c]])*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[-(c/
(-b - Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c))/(-b + Sqrt[b^2 - 4*
a*c])))/(Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a + b*x^2 + c*x^4
]) - ((42*I)*Sqrt[2]*a*b*B*c*d*e^2*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b^2 - 4*a*
c]])*Sqrt[1 - (2*c*x^2)/(-b + Sqrt[b^2 - 4*a*c]])*EllipticF[I*ArcSinh[Sqrt[
2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c))/(-b + S
qrt[b^2 - 4*a*c])))/(Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a + b*x^2 + c
*x^4]) + ((105*I)*a*A*c^2*d*e^2*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b^2 - 4*a*c])
]*Sqrt[1 - (2*c*x^2)/(-b + Sqrt[b^2 - 4*a*c]])*EllipticF[I*ArcSinh[Sqrt[2]*
Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c))/(-b + Sqrt
[b^2 - 4*a*c])))/(Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a + b*x^
2 + c*x^4]) + ((12*I)*Sqrt[2]*a*b^2*B*e^3*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b^2
- 4*a*c]])*Sqrt[1 - (2*c*x^2)/(-b + Sqrt[b^2 - 4*a*c]])*EllipticF[I*ArcSin
h[Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c))/
(-b + Sqrt[b^2 - 4*a*c])))/(Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a + b*
x^2 + c*x^4]) - ((14*I)*Sqrt[2]*a*A*b*c*e^3*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b
^2 - 4*a*c]])*Sqrt[1 - (2*c*x^2)/(-b + Sqrt[b^2 - 4*a*c]])*EllipticF[I*ArcS
inh[Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c]
))/(-b + Sqrt[b^2 - 4*a*c])))/(Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a +
b*x^2 + c*x^4]) - ((25*I)*a^2*B*c*e^3*Sqrt[1 - (2*c*x^2)/(-b - Sqrt[b^2 - 4
*a*c]])*Sqrt[1 - (2*c*x^2)/(-b + Sqrt[b^2 - 4*a*c]])*EllipticF[I*ArcSinh[Sq
rt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*x], (-b - Sqrt[b^2 - 4*a*c))/(-b
+ Sqrt[b^2 - 4*a*c])))/(Sqrt[2]*Sqrt[-(c/(-b - Sqrt[b^2 - 4*a*c]))]*Sqrt[a
+ b*x^2 + c*x^4]))/(105*c^3)

```

fricas [F] time = 0.90, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{Be^3x^8 + (3Bde^2 + Ae^3)x^6 + 3(Bd^2e + Ade^2)x^4 + Ad^3 + (Bd^3 + 3Ad^2e)x^2}{\sqrt{cx^4 + bx^2 + a}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] integral((B*e^3*x^8 + (3*B*d*e^2 + A*e^3)*x^6 + 3*(B*d^2*e + A*d*e^2)*x^4 + A*d^3 + (B*d^3 + 3*A*d^2*e)*x^2)/sqrt(c*x^4 + b*x^2 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^3/sqrt(c*x^4 + b*x^2 + a), x)

maple [B] time = 0.02, size = 1708, normalized size = 2.26

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x)

[Out] B*e^3*(1/7/c*x^5*(c*x^4+b*x^2+a)^(1/2)-6/35*b/c^2*x^3*(c*x^4+b*x^2+a)^(1/2)+1/3*(-5/7*a/c+24/35*b^2/c^2)/c*x*(c*x^4+b*x^2+a)^(1/2)-1/12*(-5/7*a/c+24/35*b^2/c^2)/c*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-1/2*(18/35*b/c^2*a-2/3*(-5/7*a/c+24/35*b^2/c^2)/c*b)*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2)))+(A*e^3+3*B*d*e^2)*(1/5*(c*x^4+b*x^2+a)^(1/2)/c*x^3-4/15*(c*x^4+b*x^2+a)^(1/2)*b/c^2*x+1/15*b/c^2*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-1/2*(-3/5*a/c+8/15*b^2/c^2)*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2

```

*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))))+(3*A*d*e^2+3*B*d^2*e)*(1/3*(c*x^4
+b*x^2+a)^(1/2)/c*x-1/12/c*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*
(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(
1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a
)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))+1/3*b/c*a^2^(1/2)/(
(-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)
*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+
b^2)^(1/2))*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*
(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*
c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))))-1/
2*(3*A*d^2*e+B*d^3)*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4
*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c
*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticF(1/2*2^(1/2)*((-b+(-4*a
*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-El
lipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b
^2)^(1/2))/a*b/c-4)^(1/2))+1/4*A*d^3*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(
1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a
*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)
^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))

```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate((B*x^2 + A)*(e*x^2 + d)^3/sqrt(c*x^4 + b*x^2 + a), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(1/2),x)
```

```
[Out] int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(1/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+b*x**2+a)**(1/2),x)
```

```
[Out] Integral((A + B*x**2)*(d + e*x**2)**3/sqrt(a + b*x**2 + c*x**4), x)
```

$$3.18 \quad \int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=528

$$\frac{x\sqrt{a+bx^2+cx^4} \left(B(-ce(9ae+20bd)+8b^2e^2+15c^2d^2)+10Ace(3cd-be) \right)}{15c^{5/2}(\sqrt{a}+\sqrt{c}x^2)} + \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(\dots\right)}{15c^{5/2}(\sqrt{a}+\sqrt{c}x^2)}$$

[Out] 1/15*e*(5*A*c*e-4*B*b*e+10*B*c*d)*x*(c*x^4+b*x^2+a)^(1/2)/c^2+1/5*B*e^2*x^3*(c*x^4+b*x^2+a)^(1/2)/c+1/15*(10*A*c*e*(-b*e+3*c*d)+B*(15*c^2*d^2+8*b^2*e^2-2*c*e*(9*a*e+20*b*d)))*x*(c*x^4+b*x^2+a)^(1/2)/c^(5/2)/(a^(1/2)+x^2*c^(1/2))-1/15*a^(1/4)*(10*A*c*e*(-b*e+3*c*d)+B*(15*c^2*d^2+8*b^2*e^2-2*c*e*(9*a*e+20*b*d)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2)))^(1/2)*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(11/4)/(c*x^4+b*x^2+a)^(1/2)+1/30*a^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2)))^(1/2)*(a^(1/2)+x^2*c^(1/2))*(10*A*c*e*(-b*e+3*c*d)+B*(15*c^2*d^2+8*b^2*e^2-2*c*e*(9*a*e+20*b*d))-(2*a*B*e*(-2*b*e+5*c*d)-5*A*c*(-a*e^2+3*c*d^2))*c^(1/2)/a^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(11/4)/(c*x^4+b*x^2+a)^(1/2)

Rubi [A] time = 0.73, antiderivative size = 528, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$, Rules used = {1679, 1197, 1103, 1195}

$$\frac{x\sqrt{a+bx^2+cx^4} \left(B(-ce(9ae+20bd)+8b^2e^2+15c^2d^2)+10Ace(3cd-be) \right)}{15c^{5/2}(\sqrt{a}+\sqrt{c}x^2)} + \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(\dots\right)}{15c^{5/2}(\sqrt{a}+\sqrt{c}x^2)}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + b*x^2 + c*x^4],x]

[Out] (e*(10*B*c*d - 4*b*B*e + 5*A*c*e)*x*Sqrt[a + b*x^2 + c*x^4])/(15*c^2) + (B*e^2*x^3*Sqrt[a + b*x^2 + c*x^4])/(5*c) + ((10*A*c*e*(3*c*d - b*e) + B*(15*c^2*d^2 + 8*b^2*e^2 - c*e*(20*b*d + 9*a*e)))*x*Sqrt[a + b*x^2 + c*x^4])/(15*c^(5/2)*(Sqrt[a] + Sqrt[c]*x^2)) - (a^(1/4)*(10*A*c*e*(3*c*d - b*e) + B*(15*c^2*d^2 + 8*b^2*e^2 - c*e*(20*b*d + 9*a*e)))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(15*c^(11/4)*Sqrt[a + b*x^2 + c*x^4])

```
x^4]) + (a^(1/4)*(10*A*c*e*(3*c*d - b*e) + B*(15*c^2*d^2 + 8*b^2*e^2 - c*e*
(20*b*d + 9*a*e)) - (Sqrt[c]*(2*a*B*e*(5*c*d - 2*b*e) - 5*A*c*(3*c*d^2 - a*
e^2)))/Sqrt[a])*Sqrt[a + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] +
Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*S
qrt[c]))/4]/(30*c^(11/4)*Sqrt[a + b*x^2 + c*x^4])
```

Rule 1103

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c
/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*
EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/ (2*q*Sqrt[a + b*x^2 + c*x^4]
), x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1195

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbo
l] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^
2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^
2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/ (q*Sqrt[a + b*x^2 + c
*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 -
4*a*c, 0] && PosQ[c/a]
```

Rule 1197

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbo
l] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4]
], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; Ne
Q[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[
c/a]
```

Rule 1679

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{q =
Expon[Pq, x^2], e = Coeff[Pq, x^2, Expon[Pq, x^2]]}, Simp[(e*x^(2*q - 3)*(
a + b*x^2 + c*x^4)^(p + 1))/(c*(2*q + 4*p + 1)), x] + Dist[1/(c*(2*q + 4*p
+ 1)), Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*q + 4*p + 1)*Pq - a*e*(2*
q - 3)*x^(2*q - 4) - b*e*(2*q + 2*p - 1)*x^(2*q - 2) - c*e*(2*q + 4*p + 1)*
x^(2*q), x], x], x]] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pq, x^2] && Expon[P
q, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]
```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx &= \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} + \frac{\int \frac{5Acd^2 + (5Bcd^2 + 10Acde - 3aBe^2)x^2 + e(10Bcd - 4bBe + 5Ace)x^4}{\sqrt{a + bx^2 + cx^4}} dx}{5c} \\
&= \frac{e(10Bcd - 4bBe + 5Ace)x\sqrt{a + bx^2 + cx^4}}{15c^2} + \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} + \frac{\int \frac{-2aBe(5cd - 2b)}{\sqrt{a + bx^2 + cx^4}} dx}{5c} \\
&= \frac{e(10Bcd - 4bBe + 5Ace)x\sqrt{a + bx^2 + cx^4}}{15c^2} + \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} - \frac{(\sqrt{a}(10Ace(3cd - 2b)))}{5c} \\
&= \frac{e(10Bcd - 4bBe + 5Ace)x\sqrt{a + bx^2 + cx^4}}{15c^2} + \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} + \frac{(10Ace(3cd - 2b))}{5c}
\end{aligned}$$

Mathematica [C] time = 4.41, size = 674, normalized size = 1.28

$$i \left(\sqrt{b^2 - 4ac} - b \right) \sqrt{\frac{\sqrt{b^2 - 4ac} + b + 2cx^2}{\sqrt{b^2 - 4ac} + b}} \sqrt{\frac{-2\sqrt{b^2 - 4ac} + 2b + 4cx^2}{b - \sqrt{b^2 - 4ac}}} E \left(i \sinh^{-1} \left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right) (B(-ce(9ae + 10cd - 4be + 3ce^2x^2)) + I(-b + \sqrt{b^2 - 4ac}))(10Ace(3cd - 2b) + B(15c^2d^2 + 8b^2e^2 - ce(20bd + 9ae))) \sqrt{\frac{(b + \sqrt{b^2 - 4ac} + 2cx^2)(b - \sqrt{b^2 - 4ac} + 4cx^2)}{(b + \sqrt{b^2 - 4ac})(b - \sqrt{b^2 - 4ac})}} \text{EllipticE} \left[I \text{ArcSinh} \left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right], \frac{(b + \sqrt{b^2 - 4ac})}{(b - \sqrt{b^2 - 4ac})} \right] - I(-8b^3Be^2 + 2b^2e(10Bcd + 5Ace + 4B\sqrt{b^2 - 4ac}e) - b^2c(15Bcd^2 + B(20\sqrt{b^2 - 4ac}d - 17ae) + 10Ae(3cd + \sqrt{b^2 - 4ac}e)) + c(B(-9a\sqrt{b^2 - 4ac}e^2 + 5cd(3\sqrt{b^2 - 4ac}d - 4ae)) + 10Ace(3cd^2 + e(3\sqrt{b^2 - 4ac}d - ae)))) \sqrt{\frac{(b + \sqrt{b^2 - 4ac} + 2cx^2)(b - \sqrt{b^2 - 4ac} + 4cx^2)}{(b + \sqrt{b^2 - 4ac})(b - \sqrt{b^2 - 4ac})}} \text{EllipticF} \left[I \text{ArcSinh} \left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right], \frac{(b + \sqrt{b^2 - 4ac})}{(b - \sqrt{b^2 - 4ac})} \right] \right) / (60c^3\sqrt{c/(b + \sqrt{b^2 - 4ac})}) \sqrt{a + bx^2 + cx^4}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + b*x^2 + c*x^4], x]

[Out] (4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*e*x*(a + b*x^2 + c*x^4)*(5*A*c*e + B*(10*c*d - 4*b*e + 3*c*e*x^2)) + I*(-b + Sqrt[b^2 - 4*a*c])*(10*A*c*e*(3*c*d - b*e) + B*(15*c^2*d^2 + 8*b^2*e^2 - c*e*(20*b*d + 9*a*e)))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - I*(-8*b^3*B*e^2 + 2*b^2*e*(10*B*c*d + 5*A*c*e + 4*B*Sqrt[b^2 - 4*a*c]*e) - b*c*(15*B*c*d^2 + B*e*(20*Sqrt[b^2 - 4*a*c]*d - 17*a*e) + 10*A*e*(3*c*d + Sqrt[b^2 - 4*a*c]*e)) + c*(B*(-9*a*Sqrt[b^2 - 4*a*c]*e^2 + 5*c*d*(3*Sqrt[b^2 - 4*a*c]*d - 4*a*e)) + 10*A*c*(3*c*d^2 + e*(3*Sqrt[b^2 - 4*a*c]*d - a*e))))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])/(60*c^3*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*Sqrt[a + b*x^2 + c*x^4])

fricas [F] time = 1.26, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{Be^2x^6 + (2Bde + Ae^2)x^4 + Ad^2 + (Bd^2 + 2Ade)x^2}{\sqrt{cx^4 + bx^2 + a}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] integral((B*e^2*x^6 + (2*B*d*e + A*e^2)*x^4 + A*d^2 + (B*d^2 + 2*A*d*e)*x^2)/sqrt(c*x^4 + b*x^2 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^2/sqrt(c*x^4 + b*x^2 + a), x)

maple [B] time = 0.01, size = 1201, normalized size = 2.27

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x)

[Out] B*e^2*(1/5*(c*x^4+b*x^2+a)^(1/2)/c*x^3-4/15*(c*x^4+b*x^2+a)^(1/2)*b/c^2*x+1/15*b/c^2*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-1/2*(-3/5*a/c+8/15*b^2/c^2)*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))) + (A*e^2+2*B*d*e)*(1/3*(c*x^4+b*x^2+a)^(1/2)/c*x-1/12/c*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)

$$\begin{aligned} &) * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * x, 1/2 * (2 * (b + (-4ac + b^2)^{1/2}) / a * b / c - 4 \\ &)^{1/2} + 1/3 * b / c * a^{1/2} / ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * (-2 * (-b + (-4ac \\ & + b^2)^{1/2}) / a * x^2 + 4)^{1/2} * (2 * (b + (-4ac + b^2)^{1/2}) / a * x^2 + 4)^{1/2} / (c * x^4 \\ & + b * x^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) * (\text{EllipticF}(1/2 * 2^{1/2} * ((-b + (-4ac \\ & + b^2)^{1/2}) / a)^{1/2} * x, 1/2 * (2 * (b + (-4ac + b^2)^{1/2}) / a * b / c - 4)^{1/2}) - \text{Ellip} \\ & \text{ticE}(1/2 * 2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * x, 1/2 * (2 * (b + (-4ac + b^2)^ \\ & ^{1/2}) / a * b / c - 4)^{1/2})) - 1/2 * (2 * A * d * e + B * d^2) * a^{1/2} / ((-b + (-4ac + b^2)^{1/2}) \\ & / a)^{1/2} * (-2 * (-b + (-4ac + b^2)^{1/2}) / a * x^2 + 4)^{1/2} * (2 * (b + (-4ac + b^2)^{1/2}) \\ &)^{1/2} / a * x^2 + 4)^{1/2} / (c * x^4 + b * x^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) * (\text{Ellip} \\ & \text{ticF}(1/2 * 2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * x, 1/2 * (2 * (b + (-4ac + b^2)^{1/2}) \\ & ^{1/2}) / a * b / c - 4)^{1/2}) - \text{EllipticE}(1/2 * 2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} \\ & * x, 1/2 * (2 * (b + (-4ac + b^2)^{1/2}) / a * b / c - 4)^{1/2})) + 1/4 * A * d^2 * 2^{1/2} / ((- \\ & b + (-4ac + b^2)^{1/2}) / a)^{1/2} * (-2 * (-b + (-4ac + b^2)^{1/2}) / a * x^2 + 4)^{1/2} * (\\ & 2 * (b + (-4ac + b^2)^{1/2}) / a * x^2 + 4)^{1/2} / (c * x^4 + b * x^2 + a)^{1/2} * \text{EllipticF}(1/2 \\ & * 2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * x, 1/2 * (2 * (b + (-4ac + b^2)^{1/2}) / \\ & a * b / c - 4)^{1/2}) \end{aligned}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^2/sqrt(c*x^4 + b*x^2 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(1/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+b*x**2+a)**(1/2),x)
```

```
[Out] Integral((A + B*x**2)*(d + e*x**2)**2/sqrt(a + b*x**2 + c*x**4), x)
```



```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)]/(2*q*Sqrt[a + b*x^2 + c*x^4]), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1195

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)]/(q*Sqrt[a + b*x^2 + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1197

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1679

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{q = Expon[Pq, x^2], e = Coeff[Pq, x^2, Expon[Pq, x^2]]}, Simp[(e*x^(2*q - 3)*(a + b*x^2 + c*x^4)^(p + 1))/(c*(2*q + 4*p + 1)), x] + Dist[1/(c*(2*q + 4*p + 1)), Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*q + 4*p + 1)*Pq - a*e*(2*q - 3)*x^(2*q - 4) - b*e*(2*q + 2*p - 1)*x^(2*q - 2) - c*e*(2*q + 4*p + 1)*x^(2*q), x], x], x] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pq, x^2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]
```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx &= \frac{Bex\sqrt{a + bx^2 + cx^4}}{3c} + \frac{\int \frac{3Ac d - aBe + (3Bcd - 2bBe + 3Ace)x^2}{\sqrt{a + bx^2 + cx^4}} dx}{3c} \\
&= \frac{Bex\sqrt{a + bx^2 + cx^4}}{3c} - \frac{(\sqrt{a}(3Bcd - 2bBe + 3Ace)) \int \frac{1 - \frac{\sqrt{c}x^2}{\sqrt{a}}}{\sqrt{a + bx^2 + cx^4}} dx}{3c^{3/2}} + \frac{(\sqrt{a}(3Bcd - 2bBe + 3Ace))}{3c^{3/2}} \\
&= \frac{Bex\sqrt{a + bx^2 + cx^4}}{3c} + \frac{(3Bcd - 2bBe + 3Ace)x\sqrt{a + bx^2 + cx^4}}{3c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}(3Bcd - 2bBe + 3Ace)}{3c^{3/2}}
\end{aligned}$$

Mathematica [C] time = 2.16, size = 521, normalized size = 1.42

$$\frac{i\sqrt{\frac{\sqrt{b^2-4ac}+b+2cx^2}{\sqrt{b^2-4ac}+b}} \sqrt{\frac{-2\sqrt{b^2-4ac}+2b+4cx^2}{b-\sqrt{b^2-4ac}}} F\left(i \sinh^{-1}\left(\sqrt{2} \sqrt{\frac{c}{b+\sqrt{b^2-4ac}}} x\right) \Big|_{\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}}\right) \left(b \left(2Be\sqrt{b^2-4ac} + 3Ace + 3Bcd\right)\right)}{3c^{3/2}(\sqrt{a} + \sqrt{c}x^2)}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2))/Sqrt[a + b*x^2 + c*x^4], x]

[Out] (4*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*e*x*(a + b*x^2 + c*x^4) - I*(-b + Sqrt[b^2 - 4*a*c])*(-3*B*c*d + 2*b*B*e - 3*A*c*e)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + I*(-2*b^2*B*e - c*(6*A*c*d + 3*B*Sqrt[b^2 - 4*a*c]*d - 2*a*B*e + 3*A*Sqrt[b^2 - 4*a*c]*e) + b*(3*B*c*d + 3*A*c*e + 2*B*Sqrt[b^2 - 4*a*c]*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))/(12*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[a + b*x^2 + c*x^4])

fricas [F] time = 1.18, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{Bex^4 + (Bd + Ae)x^2 + Ad}{\sqrt{cx^4 + bx^2 + a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2), x, algorithm="fricas")

[Out] integral((B*e*x^4 + (B*d + A*e)*x^2 + A*d)/sqrt(c*x^4 + b*x^2 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)/sqrt(c*x^4 + b*x^2 + a), x)

maple [B] time = 0.01, size = 759, normalized size = 2.06

$$\frac{\sqrt{2} \sqrt{-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}} + 4 \sqrt{\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}} + 4 Ad \operatorname{EllipticF}\left(\frac{\sqrt{2} \sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} x}{2}, \sqrt{\frac{2(b+\sqrt{-4ac+b^2})b}{ac}}\right)}{4 \sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} \sqrt{cx^4 + bx^2 + a}} + \left(\sqrt{2} \sqrt{\dots} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x)

[Out] B*e*(1/3*(c*x^4+b*x^2+a)^(1/2)/c*x-1/12/c*a*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))+1/3*b/c*a*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))))-1/2*(A*e+B*d)*a*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))))+1/4*A*d*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)/sqrt(c*x^4 + b*x^2 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(1/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)/sqrt(a + b*x**2 + c*x**4), x)

$$3.20 \quad \int \frac{A+Bx^2}{\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=283

$$\frac{\sqrt[4]{a} (\sqrt{a} + \sqrt{c} x^2) \left(\frac{A\sqrt{c}}{\sqrt{a}} + B \right) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}} - \frac{\sqrt[4]{a} B (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{c^{3/4}\sqrt{a+bx^2+cx^4}}$$

[Out] B*x*(c*x^4+b*x^2+a)^(1/2)/c^(1/2)/(a^(1/2)+x^2*c^(1/2))-a^(1/4)*B*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(3/4)/(c*x^4+b*x^2+a)^(1/2)+1/2*a^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*(B+A*c^(1/2)/a^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(3/4)/(c*x^4+b*x^2+a)^(1/2)

Rubi [A] time = 0.08, antiderivative size = 283, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {1197, 1103, 1195}

$$\frac{\sqrt[4]{a} (\sqrt{a} + \sqrt{c} x^2) \left(\frac{A\sqrt{c}}{\sqrt{a}} + B \right) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}} - \frac{\sqrt[4]{a} B (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{c^{3/4}\sqrt{a+bx^2+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/Sqrt[a + b*x^2 + c*x^4], x]

[Out] (B*x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[c]*(Sqrt[a] + Sqrt[c]*x^2)) - (a^(1/4)*B*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(3/4)*Sqrt[a + b*x^2 + c*x^4]) + (a^(1/4)*(B + (A*Sqrt[c])/Sqrt[a])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*c^(3/4)*Sqrt[a + b*x^2 + c*x^4])

Rule 1103

Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] :> With[{q = Rt[c/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2] + EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(2*q*Sqrt[a + b*x^2 + c*x^4])

), x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1195

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)]/(q*Sqrt[a + b*x^2 + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1197

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rubi steps

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx = \left(A + \frac{\sqrt{a}B}{\sqrt{c}} \right) \int \frac{1}{\sqrt{a + bx^2 + cx^4}} dx - \frac{(\sqrt{a}B) \int \frac{1 - \frac{\sqrt{c}x^2}{\sqrt{a}}}{\sqrt{a + bx^2 + cx^4}} dx}{\sqrt{c}}$$

$$= \frac{Bx\sqrt{a + bx^2 + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}B(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \Big| \frac{1}{4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{c^{3/4}\sqrt{a + bx^2 + cx^4}}$$

Mathematica [C] time = 0.24, size = 302, normalized size = 1.07

$$\frac{i \sqrt{\frac{\sqrt{b^2 - 4ac} + b + 2cx^2}{\sqrt{b^2 - 4ac} + b}} \sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} + 1 \left((-B\sqrt{b^2 - 4ac} - 2Ac + bB) F\left(i \sinh^{-1}\left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x\right) \Big| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right) + B \right)}{2\sqrt{2}c \sqrt{\frac{c}{\sqrt{b^2 - 4ac} + b}} \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/Sqrt[a + b*x^2 + c*x^4], x]

[Out] ((I/2)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(B*(-b + Sqrt[b^2 - 4*a*c])*Ellipti

`cE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + (b*B - 2*A*c - B*Sqrt[b^2 - 4*a*c])*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])]/(Sqrt[2]*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[a + b*x^2 + c*x^4])`

fricas [F] time = 1.33, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")`

[Out] `integral((B*x^2 + A)/sqrt(c*x^4 + b*x^2 + a), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

[Out] `integrate((B*x^2 + A)/sqrt(c*x^4 + b*x^2 + a), x)`

maple [A] time = 0.00, size = 362, normalized size = 1.28

$$\frac{\sqrt{2} \sqrt{-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}} + 4 \sqrt{\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}} + 4 A \text{EllipticF}\left(\frac{\sqrt{2} \sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} x}{2}, \frac{\sqrt{\frac{2(b+\sqrt{-4ac+b^2})b}{ac}-4}}{2}\right) \sqrt{2} \sqrt{-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}}{4 \sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} \sqrt{cx^4 + bx^2 + a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x)`

[Out] `-1/2*B*a^2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*`

$b/c-4)^{(1/2)})+1/4*A*2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)*(-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2+4)^{(1/2)*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2+4)^{(1/2)/(c*x^4+b*x^2+a)^{(1/2)*EllipticF(1/2*2^{(1/2)*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*b/c-4)^{(1/2))}}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/sqrt(c*x^4 + b*x^2 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/(a + b*x^2 + c*x^4)^(1/2),x)

[Out] int((A + B*x^2)/(a + b*x^2 + c*x^4)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((A + B*x**2)/sqrt(a + b*x**2 + c*x**4), x)

$$3.21 \quad \int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=436

$$\frac{a^{3/4}(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \left(\frac{\sqrt{c}d}{\sqrt{a}} + e\right)^2 (Bd - Ae) \Pi\left(-\frac{(\sqrt{c}d-\sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\right) \Big|_{1/4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)}{4\sqrt[4]{c}de\sqrt{a+bx^2+cx^4}(cd^2-ae^2)} \quad 2\sqrt{d}$$

[Out] $-1/2*(-A*e+B*d)*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})/d^{(1/2)}/e^{(1/2)}/(a*e^2-b*d*e+c*d^2)^{(1/2)}-1/2*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), 1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(B*a^{(1/2)}-A*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}+1/4*a^{(3/4)}*(-A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)}, 1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(e+d*c^{(1/2)}/a^{(1/2)})^2*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(1/4)}/d/e/(-a*e^2+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.41, antiderivative size = 436, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {1708, 1103, 1706}

$$\frac{a^{3/4}(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \left(\frac{\sqrt{c}d}{\sqrt{a}} + e\right)^2 (Bd - Ae) \Pi\left(-\frac{(\sqrt{c}d-\sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\right) \Big|_{1/4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)}{4\sqrt[4]{c}de\sqrt{a+bx^2+cx^4}(cd^2-ae^2)} \quad 2\sqrt{d}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x]

[Out] $-\left(\frac{(B*d - A*e)*\text{ArcTan}\left[\frac{\sqrt{c*d^2 - b*d*e + a*e^2}*x}{\sqrt{d}*\sqrt{e}*\sqrt{a + b*x^2 + c*x^4}}\right]}{2*\sqrt{d}*\sqrt{e}*\sqrt{c*d^2 - b*d*e + a*e^2}} - \left(\frac{(\sqrt{a}*B - A*\sqrt{c})*(\sqrt{a} + \sqrt{c}*x^2)*\sqrt{(a + b*x^2 + c*x^4)/(\sqrt{a} + \sqrt{c}*x^2)^2}}{2*\sqrt{a}*\sqrt{c}}*\text{EllipticF}\left[2*\text{ArcTan}\left[\frac{c^{(1/4)}*x}{a^{(1/4)}}\right], \frac{2 - b/(\sqrt{a}*\sqrt{c})}{4}\right]}{2*a^{(1/4)}*c^{(1/4)}*(\sqrt{c}*d - \sqrt{a}*e)*\sqrt{a + b*x^2 + c*x^4}} + a^{(3/4)}*\left(\frac{\sqrt{c}*d}{\sqrt{a}} + e\right)^2*(B*d - A*e)*(\sqrt{a} + \sqrt{c}*x^2)*\sqrt{(a + b*x^2 + c*x^4)/(\sqrt{a} + \sqrt{c}*x^2)^2}}{4*c^{(1/4)}*d*e*(c*d^2 - a*e^2)*\sqrt{a + b*x^2 + c*x^4}}\right)*\text{EllipticPi}\left[-\frac{(\sqrt{c}*d - \sqrt{a}*e)^2}{4*\sqrt{a}*\sqrt{c}*d*e}, 2*\text{ArcTan}\left[\frac{c^{(1/4)}*x}{a^{(1/4)}}\right], \frac{2 - b/(\sqrt{a}*\sqrt{c})}{4}\right]\right)$

Rule 1103

Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(2*q*Sqrt[a + b*x^2 + c*x^4]), x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1706

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[(Rt[-b + (c*d)/e + (a*e)/d, 2]*x)/Sqrt[a + b*x^2 + c*x^4]])/(2*d*e*Rt[-b + (c*d)/e + (a*e)/d, 2]), x] + Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a + b*x^2 + c*x^4))/(a*(A + B*x^2)^2)]*EllipticPi[Cancel[-((B*d - A*e)^2/(4*d*e*A*B))], 2*ArcTan[q*x], 1/2 - (b*A)/(4*a*B)]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4]), x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

Rule 1708

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] + Dist[(a*(B*d - A*e)*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]

Rubi steps

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = -\frac{(\sqrt{a}B - A\sqrt{c}) \int \frac{1}{\sqrt{a+bx^2+cx^4}} dx}{\sqrt{c}d - \sqrt{a}e} + \frac{(\sqrt{a}(Bd - Ae)) \int \frac{1 + \frac{\sqrt{c}x^2}{\sqrt{a}}}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx}{\sqrt{c}d - \sqrt{a}e}$$

$$= -\frac{(Bd - Ae) \tan^{-1}\left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d}\sqrt{e}\sqrt{a+bx^2+cx^4}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{cd^2 - bde + ae^2}} - \frac{(\sqrt{a}B - A\sqrt{c})(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2}{(\sqrt{a} + \sqrt{c}x^2)^2}}}{2^4 \sqrt{a} \sqrt{c} (\sqrt{c}d - \sqrt{a})}$$

Mathematica [C] time = 0.49, size = 298, normalized size = 0.68

$$\frac{i\sqrt{\frac{\sqrt{b^2-4ac}+b+2cx^2}{\sqrt{b^2-4ac}+b}}\sqrt{\frac{2cx^2}{b-\sqrt{b^2-4ac}}}+1\left((Ae-Bd)\Pi\left(\frac{(b+\sqrt{b^2-4ac})e}{2cd};i\sinh^{-1}\left(\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\right)x\right)\Big|_{\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}}\right)+BdF\left(\frac{\sqrt{2}de\sqrt{\frac{c}{\sqrt{b^2-4ac}+b}}\sqrt{a+bx^2+cx^4}}{\sqrt{b^2-4ac}+b}\right)}{\sqrt{2}de\sqrt{\frac{c}{\sqrt{b^2-4ac}+b}}\sqrt{a+bx^2+cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-I)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(B*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + (-B*d) + A*e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*Sqrt[a + b*x^2 + c*x^4])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)

maple [A] time = 0.04, size = 359, normalized size = 0.82

$$\frac{\sqrt{2}\sqrt{-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}+4\sqrt{\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}+4B\text{EllipticF}\left(\frac{\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}x}{2},\frac{\sqrt{\frac{2(b+\sqrt{-4ac+b^2})b}{ac}-4}}{2}\right)(Ae-Bd)}{4\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}e}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2), x)`

[Out] $\frac{1}{4} \frac{B}{e} \frac{1}{2^{1/2}} \left(\frac{-b + (-4ac + b^2)^{1/2}}{a} \right)^{1/2} \frac{(-2(-b + (-4ac + b^2)^{1/2}))}{a x^2 + 4} \frac{(2(b + (-4ac + b^2)^{1/2}))}{a x^2 + 4} \frac{1}{(c x^4 + b x^2 + a)^{1/2}} \text{EllipticF}\left(\frac{1}{2} \frac{1}{2^{1/2}} \left(\frac{-b + (-4ac + b^2)^{1/2}}{a} \right)^{1/2} x, \frac{1}{2} \frac{(2(b + (-4ac + b^2)^{1/2}))}{a b / c - 4} \right)^{1/2} + \frac{A e - B d}{e} \frac{1}{d} \frac{1}{2^{1/2}} \frac{(-1/a b + 1/a (-4ac + b^2)^{1/2})^{1/2}}{(1 + 1/2 a x^2 b - 1/2 a x^2 (-4ac + b^2)^{1/2})^{1/2}} \frac{(1 + 1/2 a x^2 b + 1/2 a x^2 (-4ac + b^2)^{1/2})^{1/2}}{(c x^4 + b x^2 + a)^{1/2}} \text{EllipticPi}\left(\frac{1}{2} \frac{1}{2^{1/2}} \left(\frac{-b + (-4ac + b^2)^{1/2}}{a} \right)^{1/2} x, -\frac{2}{(-b + (-4ac + b^2)^{1/2})} \frac{a e / d}{(-1/2 (b + (-4ac + b^2)^{1/2}) / a)^{1/2}} \frac{1}{2^{1/2}} \left(\frac{-b + (-4ac + b^2)^{1/2}}{a} \right)^{1/2} \right)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2), x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(ex^2 + d) \sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)`

[Out] `int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2), x)`

[Out] `Integral((A + B*x**2)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)`

$$3.22 \quad \int \frac{A+Bx^2}{(d+ex^2)^2 \sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=782

$$\frac{\sqrt{c}x\sqrt{a+bx^2+cx^4}(Bd-Ae)}{2d(\sqrt{a}+\sqrt{c}x^2)(ae^2-bde+cd^2)} - \frac{ex\sqrt{a+bx^2+cx^4}(Bd-Ae)}{2d(d+ex^2)(ae^2-bde+cd^2)} - \frac{\sqrt[4]{a}\sqrt[4]{c}(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}(Bd-Ae)}{2d\sqrt{a+bx^2+cx^4}(ae^2-bde+cd^2)}$$

[Out] $-1/4*(B*(-a*d*e^2+c*d^3)-A*e*(3*c*d^2-e*(-a*e+2*b*d)))*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})/d^{(3/2)}/(a*e^2-b*d*e+c*d^2)^{(3/2)}/e^{(1/2)}-1/2*e*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^{(1/2)}/d/(a*e^2-b*d*e+c*d^2)/(e*x^2+d)+1/2*(-A*e+B*d)*x*c^{(1/2)}*(c*x^4+b*x^2+a)^{(1/2)}/d/(a*e^2-b*d*e+c*d^2)/(a^{(1/2)}+x^2*c^{(1/2)})-1/2*a^{(1/4)}*c^{(1/4)}*(-A*e+B*d)*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/d/(a*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}+1/2*A*c^{(1/4)}*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/d/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}+1/8*(B*(-a*d*e^2+c*d^3)-A*e*(3*c*d^2-e*(-a*e+2*b*d)))*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),-1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^2/e/(a*e^2-b*d*e+c*d^2)/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A] time = 1.47, antiderivative size = 782, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {1696, 1714, 1195, 1708, 1103, 1706}

$$\frac{\sqrt{c}x\sqrt{a+bx^2+cx^4}(Bd-Ae)}{2d(\sqrt{a}+\sqrt{c}x^2)(ae^2-bde+cd^2)} - \frac{ex\sqrt{a+bx^2+cx^4}(Bd-Ae)}{2d(d+ex^2)(ae^2-bde+cd^2)} - \frac{(B(cd^3-ade^2)-Ae(3cd^2-e(2bd-ae)))}{4d^{3/2}\sqrt{e}(ae^2-bde+cd^2)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(A+B*x^2)/((d+e*x^2)^2*\text{Sqrt}[a+b*x^2+c*x^4]),x]$

[Out] $(\text{Sqrt}[c]*(B*d-A*e)*x*\text{Sqrt}[a+b*x^2+c*x^4])/(2*d*(c*d^2-b*d*e+a*e^2)*(\text{Sqrt}[a]+\text{Sqrt}[c]*x^2))-(e*(B*d-A*e)*x*\text{Sqrt}[a+b*x^2+c*x^4])/(2*d$

$$\begin{aligned} & * (c*d^2 - b*d*e + a*e^2)*(d + e*x^2) - ((B*(c*d^3 - a*d*e^2) - A*e*(3*c*d^2 - e*(2*b*d - a*e))) * \text{ArcTan}[\text{Sqrt}[c*d^2 - b*d*e + a*e^2]*x] / (\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4])) / (4*d^{3/2}*\text{Sqrt}[e]*(c*d^2 - b*d*e + a*e^2)^{3/2}) - (a^{1/4}*c^{1/4}*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2] * \text{EllipticE}[2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4]) / (2*d*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) + (A*c^{1/4}*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2] * \text{EllipticF}[2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4]) / (2*a^{1/4}*d*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*\text{Sqrt}[a + b*x^2 + c*x^4]) + ((\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(B*(c*d^3 - a*d*e^2) - A*e*(3*c*d^2 - e*(2*b*d - a*e))) * (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2] * \text{EllipticPi}[-(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(4*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4]) / (8*a^{1/4}*c^{1/4}*d^2*e*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) \end{aligned}$$

Rule 1103

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(2*q*Sqrt[a + b*x^2 + c*x^4]), x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1195

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(q*Sqrt[a + b*x^2 + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1696

```
Int[((P4x_)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, -Simp[((C*d^2 - B*d*e + A*e^2)*x*(d + e*x^2)^(q + 1)*Sqrt[a + b*x^2 + c*x^4])/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2)), Int[((d + e*x^2)^(q + 1)*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*d*(c*d - b*e)*(q + 1)) - 2*((B*d - A*e)*(b*e*(q + 2) - c*d*(q + 1)) - C*d*(b*d + a*e*(q + 1)))*x^2 + c*(C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x])/Sqrt[a + b*x^2 + c*x^4], x]] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && ILtQ[q, -1]
```

Rule 1706

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 +
(c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*Arc
Tan[(Rt[-b + (c*d)/e + (a*e)/d, 2]*x)/Sqrt[a + b*x^2 + c*x^4]]/(2*d*e*Rt[-
b + (c*d)/e + (a*e)/d, 2]), x] + Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a
+ b*x^2 + c*x^4))/(a*(A + B*x^2)^2)]*EllipticPi[Cancel[-((B*d - A*e)^2/(4*
d*e*A*B))], 2*ArcTan[q*x], 1/2 - (b*A)/(4*a*B)]/(4*d*e*A*q*Sqrt[a + b*x^2
+ c*x^4]), x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] &&
NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ
[c*A^2 - a*B^2, 0]
```

Rule 1708

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2
+ (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q)
- a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] +
Dist[(a*(B*d - A*e)*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2
)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && N
eQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0
] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rule 1714

```
Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x
, 2], C = Coeff[P4x, x, 4]}, -Dist[C/(e*q), Int[(1 - q*x^2)/Sqrt[a + b*x^2
+ c*x^4], x], x] + Dist[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a
*e*q))*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b,
c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*
d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && !GtQ[b^2 - 4*a*c,
0]
```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx &= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)} - \frac{\int \frac{-aBde - A(2cd^2 - e(2bd - ae)) - 2cd(Bd - Ae)x^2 - ce(Bd - Ae)}{(d + ex^2)\sqrt{a + bx^2 + cx^4}}}{2d(cd^2 - bde + ae^2)} \\
&= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)} - \frac{\int \frac{-\sqrt{a}c^{3/2}de(Bd - Ae) + ce(-aBde - A(2cd^2 - e(2bd - ae)))}{(d + ex^2)\sqrt{a + bx^2 + cx^4}}}{2cde(cd^2 - bde + ae^2)} \\
&= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)} - \frac{\sqrt[4]{a}\sqrt[4]{c}}{2cde(cd^2 - bde + ae^2)} \\
&= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)} - \frac{B(cd^2 - bde + ae^2)}{2cde(cd^2 - bde + ae^2)}
\end{aligned}$$

Mathematica [C] time = 3.47, size = 1853, normalized size = 2.37

result too large to display

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-1/8*I)*((-4*I)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*d*e^2*(B*d - A*e)*x*(a + b*x^2 + c*x^4) + Sqrt[2]*B*(b - Sqrt[b^2 - 4*a*c])*d^2*e*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])])*(d + e*x^2)*(EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) - EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + Sqrt[2]*A*(-b + Sqrt[b^2 - 4*a*c])*d*e^2*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])])*(d + e*x^2)*(EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) - EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + 2*Sqrt[2]*B*c*d^3*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])])*(d + e*x^2)*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) - 2*Sqrt[2]*A*c*d^2*e*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b

```

^2 - 4*a*c]))*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(d + e*x^2)*Ellip
ticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 -
4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - 2*Sqrt[2]*B*c*d^3*Sqrt[(b + Sqrt[b^2 - 4
*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2
- 4*a*c])]*(d + e*x^2)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*Ar
cSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(
b - Sqrt[b^2 - 4*a*c])] + 6*Sqrt[2]*A*c*d^2*e*Sqrt[(b + Sqrt[b^2 - 4*a*c] +
2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c
])]*(d + e*x^2)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[S
qrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqr
t[b^2 - 4*a*c])] - 4*Sqrt[2]*A*b*d*e^2*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^
2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(d
+ e*x^2)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*
Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 -
4*a*c])] + 2*Sqrt[2]*a*B*d*e^2*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b +
Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(d + e*x^2
)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/
(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c
])] + 2*Sqrt[2]*a*A*e^3*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2
- 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(d + e*x^2)*Ellipti
cPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt
[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])]))/(Sqr
t[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*(c*d^3 + d*e*(-(b*d) + a*e))*(d + e*x^2)*S
qrt[a + b*x^2 + c*x^4])

```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")
```

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a} (ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)^2), x)
```


Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)^2), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(ex^2 + d)^2 \sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(1/2)),x)
```

```
[Out] int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(1/2)), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+b*x**2+a)**(1/2),x)
```

```
[Out] Integral((A + B*x**2)/((d + e*x**2)**2*sqrt(a + b*x**2 + c*x**4)), x)
```

$$3.23 \quad \int \frac{A+Bx^2}{(d+ex^2)^3 \sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=1125

$$\frac{\sqrt{c} \left(3Ae \left(3cd^2 - e(2bd - ae) \right) - Bd \left(5cd^2 - e(2bd + ae) \right) \right) \sqrt{cx^4 + bx^2 + a} x}{8d^2 \left(cd^2 - bed + ae^2 \right)^2 \left(\sqrt{c} x^2 + \sqrt{a} \right)} + \frac{e \left(3Ae \left(3cd^2 - e(2bd - ae) \right) - Bd \left(5cd^2 - e(2bd + ae) \right) \right)}{8d^2 \left(cd^2 - bed + ae^2 \right)^2}$$

[Out] $-1/16*(B*d*(3*c^2*d^4-10*a*c*d^2*e^2+a*e^3*(-a*e+4*b*d))-A*e*(15*c^2*d^4-2*c*d^2*e*(-3*a*e+10*b*d)+e^2*(3*a^2*e^2-8*a*b*d*e+8*b^2*d^2)))*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/d^{(5/2)}/(a*e^2-b*d*e+c*d^2)^{(5/2)}/e^{(1/2)}-1/4*e*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^{(1/2)}/d/(a*e^2-b*d*e+c*d^2)/(e*x^2+d)^2+1/8*e*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d)))*x*(c*x^4+b*x^2+a)^{(1/2)}/d^2/(a*e^2-b*d*e+c*d^2)^2/(e*x^2+d)-1/8*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d)))*x*c^{(1/2)}*(c*x^4+b*x^2+a)^{(1/2)}/d^2/(a*e^2-b*d*e+c*d^2)^2/(a^{(1/2)}+x^2*c^{(1/2)})+1/8*a^{(1/4)}*c^{(1/4)}*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d)))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/d^2/(a*e^2-b*d*e+c*d^2)^2/(c*x^4+b*x^2+a)^{(1/2)}+1/32*(B*d*(3*c^2*d^4-10*a*c*d^2*e^2+a*e^3*(-a*e+4*b*d))-A*e*(15*c^2*d^4-2*c*d^2*e*(-3*a*e+10*b*d)+e^2*(3*a^2*e^2-8*a*b*d*e+8*b^2*d^2)))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^3/e/(a*e^2-b*d*e+c*d^2)^2/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}+1/8*c^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*(a*e*(3*A*e+B*d)+4*A*d*(-b*e+c*d)+d*(-A*e+B*d)*a^{(1/2)}*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/d^2/(a*e^2-b*d*e+c*d^2)/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A] time = 3.59, antiderivative size = 1125, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {1696, 1714, 1195, 1708, 1103, 1706}

$$\frac{\sqrt{c} \left(3Ae \left(3cd^2 - e(2bd - ae) \right) - B \left(5cd^3 - de(2bd + ae) \right) \right) \sqrt{cx^4 + bx^2 + a} x}{8d^2 \left(cd^2 - bed + ae^2 \right)^2 \left(\sqrt{c} x^2 + \sqrt{a} \right)} + \frac{e \left(3Ae \left(3cd^2 - e(2bd - ae) \right) - B \left(5cd^3 - de(2bd + ae) \right) \right)}{8d^2 \left(cd^2 - bed + ae^2 \right)^2}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + b*x^2 + c*x^4]),x]

[Out]
$$-(\text{Sqrt}[c]*(3*A*e*(3*c*d^2 - e*(2*b*d - a*e)) - B*(5*c*d^3 - d*e*(2*b*d + a*e))) * x * \text{Sqrt}[a + b*x^2 + c*x^4] / (8*d^2*(c*d^2 - b*d*e + a*e^2)^2 * (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (e*(B*d - A*e) * x * \text{Sqrt}[a + b*x^2 + c*x^4]) / (4*d*(c*d^2 - b*d*e + a*e^2)*(d + e*x^2)^2) + (e*(3*A*e*(3*c*d^2 - e*(2*b*d - a*e)) - B*(5*c*d^3 - d*e*(2*b*d + a*e))) * x * \text{Sqrt}[a + b*x^2 + c*x^4] / (8*d^2*(c*d^2 - b*d*e + a*e^2)^2*(d + e*x^2)) - ((B*(3*c^2*d^5 - 10*a*c*d^3*e^2 + a*d*e^3*(4*b*d - a*e)) - A*e*(15*c^2*d^4 - 2*c*d^2*e*(10*b*d - 3*a*e) + e^2*(8*b^2*d^2 - 8*a*b*d*e + 3*a^2*e^2))) * \text{ArcTan}[(\text{Sqrt}[c*d^2 - b*d*e + a*e^2]*x) / (\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4])] / (16*d^(5/2)*\text{Sqrt}[e]*(c*d^2 - b*d*e + a*e^2)^(5/2)) + (a^(1/4)*c^(1/4)*(3*A*e*(3*c*d^2 - e*(2*b*d - a*e)) - B*(5*c*d^3 - d*e*(2*b*d + a*e))) * (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2) * \text{Sqrt}[(a + b*x^2 + c*x^4) / (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2] * \text{EllipticE}[2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4] / (8*d^2*(c*d^2 - b*d*e + a*e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4]) + (c^(1/4)*(\text{Sqrt}[a]*\text{Sqrt}[c]*d*(B*d - A*e) + a*e*(B*d + 3*A*e) + 4*A*d*(c*d - b*e)) * (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2) * \text{Sqrt}[(a + b*x^2 + c*x^4) / (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2] * \text{EllipticF}[2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4] / (8*a^(1/4)*d^2*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) + ((\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(B*(3*c^2*d^5 - 10*a*c*d^3*e^2 + a*d*e^3*(4*b*d - a*e)) - A*e*(15*c^2*d^4 - 2*c*d^2*e*(10*b*d - 3*a*e) + e^2*(8*b^2*d^2 - 8*a*b*d*e + 3*a^2*e^2))) * (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2) * \text{Sqrt}[(a + b*x^2 + c*x^4) / (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2] * \text{EllipticPi}[-(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2 / (4*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4] / (32*a^(1/4)*c^(1/4)*d^3*e*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 - b*d*e + a*e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4])$$

Rule 1103

Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)]/(2*q*Sqrt[a + b*x^2 + c*x^4]), x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1195

Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)]/(q*Sqrt[a + b*x^2 + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1696

```

Int[((P4x_)*((d_) + (e_)*(x_)^2)^(q_))/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, -Simp[((C*d^2 - B*d*e + A*e^2)*x*(d + e*x^2)^(q + 1)*Sqrt[a + b*x^2 + c*x^4]/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2)), Int[((d + e*x^2)^(q + 1)*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*d*(c*d - b*e)*(q + 1)) - 2*((B*d - A*e)*(b*e*(q + 2) - c*d*(q + 1)) - C*d*(b*d + a*e*(q + 1)))*x^2 + c*(C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x])/Sqrt[a + b*x^2 + c*x^4], x]] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && ILtQ[q, -1]

```

Rule 1706

```

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[(Rt[-b + (c*d)/e + (a*e)/d, 2]*x)/Sqrt[a + b*x^2 + c*x^4]]/(2*d*e*Rt[-b + (c*d)/e + (a*e)/d, 2]), x] + Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a + b*x^2 + c*x^4))/(a*(A + B*x^2)^2)]*EllipticPi[Cancel[-((B*d - A*e)^2/(4*d*e*A*B))], 2*ArcTan[q*x], 1/2 - (b*A)/(4*a*B)]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4]), x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

```

Rule 1708

```

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] + Dist[(a*(B*d - A*e)*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]

```

Rule 1714

```

Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, -Dist[C/(e*q), Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] + Dist[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && !GtQ[b^2 - 4*a*c, 0]

```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx &= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{4d(cd^2 - bde + ae^2)(d + ex^2)^2} - \frac{\int \frac{-4Acd^2 - aBde + Ae(4bd - 3ae) - 2(Bd - Ae)(2cd - be)x^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}}}{4d(cd^2 - bde + ae^2)} \\
&= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{4d(cd^2 - bde + ae^2)(d + ex^2)^2} + \frac{e(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))}{8d^2(cd^2 - bde + ae^2)} \\
&= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{4d(cd^2 - bde + ae^2)(d + ex^2)^2} + \frac{e(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))}{8d^2(cd^2 - bde + ae^2)} \\
&= -\frac{\sqrt{c}(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))x\sqrt{a + bx^2 + cx^4}}{8d^2(cd^2 - bde + ae^2)^2(\sqrt{a} + \sqrt{c}x^2)} \\
&= -\frac{\sqrt{c}(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))x\sqrt{a + bx^2 + cx^4}}{8d^2(cd^2 - bde + ae^2)^2(\sqrt{a} + \sqrt{c}x^2)}
\end{aligned}$$

Mathematica [C] time = 5.73, size = 781, normalized size = 0.69

$$-\frac{4de^2x(a+bx^2+cx^4)(2d(Bd-Ae)(e(ae-bd)+cd^2)+(d+ex^2)(B(5cd^3-de(ae+2bd))-3Ae(e(ae-2bd)+3cd^2)))}{(d+ex^2)^2} - \frac{i\sqrt{2}\sqrt{\frac{\sqrt{b^2-4ac}+b+2cx^2}{\sqrt{b^2-4ac}+b}}\sqrt{\frac{2cx^2}{b-\sqrt{b^2-4ac}}+1}}{1}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-4*d*e^2*x*(a + b*x^2 + c*x^4)*(2*d*(B*d - A*e)*(c*d^2 + e*(-(b*d) + a*e)) + (-3*A*e*(3*c*d^2 + e*(-2*b*d + a*e)) + B*(5*c*d^3 - d*e*(2*b*d + a*e)))*(d + e*x^2))/(d + e*x^2)^2 - (I*Sqrt[2]*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])]*((-b + Sqrt[b^2 - 4*a*c])*d*e*(3*A*e*(3*c*d^2 + e*(-2*b*d + a*e)) + B*(-5*c*d^3 + d*e*(2*b*d + a*e)))*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])],x])

```

- 4*a*c))]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + d*(B*d*(
6*c^2*d^3 + c*d*e*(-5*b*d + 5*Sqrt[b^2 - 4*a*c]*d - 6*a*e) - (-b + Sqrt[b^2
- 4*a*c])*e^2*(2*b*d + a*e)) - A*e*(14*c^2*d^3 - 3*(-b + Sqrt[b^2 - 4*a*c]
)*e^2*(2*b*d - a*e) + c*d*e*(-17*b*d + 9*Sqrt[b^2 - 4*a*c]*d + 2*a*e)))*Ell
ipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2
- 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + 2*(B*(-3*c^2*d^5 + 10*a*c*d^3*e^2 + a
d*e^3*(-4*b*d + a*e)) + A*e*(15*c^2*d^4 + 2*c*d^2*e*(-10*b*d + 3*a*e) + e^2
*(8*b^2*d^2 - 8*a*b*d*e + 3*a^2*e^2)))*EllipticPi[((b + Sqrt[b^2 - 4*a*c])
e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt
[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))]/Sqrt[c/(b + Sqrt[b^2 - 4*a*c]))]/
(32*d^3*e*(c*d^2 + e*(-(b*d) + a*e))^2*Sqrt[a + b*x^2 + c*x^4])

```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")
)
```

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a} (ex^2 + d)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)^3), x)
```

maple [B] time = 0.05, size = 4476, normalized size = 3.98

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x)
```

```
[Out] B/e*(1/2/(a*e^2-b*d*e+c*d^2)*(c*x^4+b*x^2+a)^(1/2)/(e*x^2+d)/d*e^2*x-1/8*c/
(a*e^2-b*d*e+c*d^2)*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(2/a*b*x^2-
2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)*(2/a*b*x^2+2*(-4*a*c+b^2)^(1/2)/a*x^2+4
)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2)
)/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))+1/4*c*e/(a*e^2-b

```

$$\begin{aligned}
& *d*e+c*d^2)/d*a^2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}*(2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticF(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)}/a*b/c-4)^{(1/2)}))-1/4*c*e/(a*e^2-b*d*e+c*d^2)/d*a^2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}*(2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticE(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)}/a*b/c-4)^{(1/2)}))+1/2/(a*e^2-b*d*e+c*d^2)/d^2*e^2*2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)}*(1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, -2/(-b+(-4*a*c+b^2)^{(1/2)}/a)*d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}))*a-1/(a*e^2-b*d*e+c*d^2)/d*e*2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)}*(1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, -2/(-b+(-4*a*c+b^2)^{(1/2)}/a)*d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}))*b+3/2/(a*e^2-b*d*e+c*d^2)*2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)}*(1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, -2/(-b+(-4*a*c+b^2)^{(1/2)}/a)*d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}))*c)+(A*e-B*d)/e*(1/4*e^2/(a*e^2-b*d*e+c*d^2)/d*x*(c*x^4+b*x^2+a)^{(1/2)}/(e*x^2+d)^2+3/8*e^2*(a*e^2-2*b*d*e+3*c*d^2)/(a*e^2-b*d*e+c*d^2)^2/d^2*x*(c*x^4+b*x^2+a)^{(1/2)}/(e*x^2+d)+3/16*e^3*c/(a*e^2-b*d*e+c*d^2)^2/d^2*a^2*2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}*(2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticF(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)}/a*b/c-4)^{(1/2)}))+9/16*e*c^2/(a*e^2-b*d*e+c*d^2)^2*a^2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}*(2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticF(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)}/a*b/c-4)^{(1/2)}))-3/16*e^3*c/(a*e^2-b*d*e+c*d^2)^2/d^2*a^2*2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}*(2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticE(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)}/a*b/c-4)^{(1/2)}))-1/(a*e^2-b*d*e+c*d^2)^2/d^2*e^3*2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)}*(1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*x, -2/(-b+(-4*a*c+b^2)^{(1/2)}/a)*d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}))*a*b+3/4/(a*e^2-b*d*e+c*d^2)^2/d*e^2*2^{(1/2)}/(-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x
\end{aligned}$$

$$\begin{aligned}
& ^2+1)^{(1/2)} * (1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} * \text{EllipticPi}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, -2/(-b+(-4*a*c+b^2)^{(1/2)}) * a/d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * 2^{(1/2)} / ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * a*c-1/32*c/(a*e^2-b*d*e+c*d^2)^2/d*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} * (2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} * \text{EllipticF}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*b/c-4)^{(1/2)} * a*e^2+1/8*c/(a*e^2-b*d*e+c*d^2)^2*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} * (2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} * \text{EllipticF}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*b/c-4)^{(1/2)} * e*b+1/(a*e^2-b*d*e+c*d^2)^2/d*e^2*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} * (1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} * \text{EllipticPi}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, -2/(-b+(-4*a*c+b^2)^{(1/2)}) * a/d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * 2^{(1/2)} / ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * b^2-5/2/(a*e^2-b*d*e+c*d^2)^2*e*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} * (1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} * \text{EllipticPi}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, -2/(-b+(-4*a*c+b^2)^{(1/2)}) * a/d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * 2^{(1/2)} / ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * b*c+3/8/(a*e^2-b*d*e+c*d^2)^2/d^3*e^4*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} * (1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} * \text{EllipticPi}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, -2/(-b+(-4*a*c+b^2)^{(1/2)}) * a/d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * 2^{(1/2)} / ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * a^2-3/8*e^2*c/(a*e^2-b*d*e+c*d^2)^2/d*a*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} * (2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} / (b+(-4*a*c+b^2)^{(1/2)}) * \text{EllipticF}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*b/c-4)^{(1/2)} * b+3/8*e^2*c/(a*e^2-b*d*e+c*d^2)^2/d*a*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} * (2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} / (b+(-4*a*c+b^2)^{(1/2)}) * \text{EllipticE}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*b/c-4)^{(1/2)} * b-9/16*e*c^2/(a*e^2-b*d*e+c*d^2)^2*a*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} * (2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} / (b+(-4*a*c+b^2)^{(1/2)}) * \text{EllipticE}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*b/c-4)^{(1/2)} + 15/8/(a*e^2-b*d*e+c*d^2)^2*d*2^{(1/2)} / (-1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)} * (1/2/a*b*x^2-1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} * (1/2/a*b*x^2+1/2*(-4*a*c+b^2)^{(1/2)}/a*x^2+1)^{(1/2)} / (c*x^4+b*x^2+a)^{(1/2)} * \text{EllipticPi}(1/2*2^{(1/2)} * ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * x, -2/(-b+(-4*a*c+b^2)^{(1/2)}) * a/d*e, (-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * 2^{(1/2)} / ((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} * c^2-7/32*c^2/(a*e^2-b*d*e+c*d^2)^2*d*2^{(1/2)} / (-
\end{aligned}$$

$1/a*b+(-4*a*c+b^2)^{(1/2)}/a)^{(1/2)}*(2/a*b*x^2-2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}*(2/a*b*x^2+2*(-4*a*c+b^2)^{(1/2)}/a*x^2+4)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*x,1/2*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*b/c-4)^{(1/2))}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a} (ex^2 + d)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(ex^2 + d)^3 \sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((d + e*x^2)^3*(a + b*x^2 + c*x^4)^(1/2)),x)

[Out] int((A + B*x^2)/((d + e*x^2)^3*(a + b*x^2 + c*x^4)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((A + B*x**2)/((d + e*x**2)**3*sqrt(a + b*x**2 + c*x**4)), x)

$$3.24 \quad \int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=859

$$\frac{Bx\sqrt{cx^4 + bx^2 + a}e^3}{3c^2} \frac{(aB(6c^3d^3 - 9c^2e(bd + 6ae)d - 8b^3e^3 + bce^2(18bd + 29ae)) + 3Ac(2ab^2e^3 + 6ac(cd^2 - ae^2))}{3a^{3/4}c^{11/4}(b^2 - 4ac)}$$

[Out] $x*(A*c*(b^2*c*d^3-2*a*c*d*(-3*a*e^2+c*d^2))-a*b*e*(a*e^2+3*c*d^2))+a*B*(a*b^2*e^3+2*a*c*e*(-a*e^2+3*c*d^2)-b*c*d*(3*a*e^2+c*d^2))-(a*B*(-b*e+2*c*d)*(c^2*d^2+b^2*e^2-c*e*(3*a*e+b*d))+A*c*(a*b^2*e^3+2*a*c*e*(-a*e^2+3*c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*x^2)/a/c^2/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^{(1/2)}+1/3*B*e^3*x*(c*x^4+b*x^2+a)^{(1/2)}/c^2+1/3*(a*B*(6*c^3*d^3-8*b^3*e^3-9*c^2*d*e*(6*a*e+b*d)+b*c*e^2*(29*a*e+18*b*d))+3*A*c*(2*a*b^2*e^3+6*a*c*e*(-a*e^2+c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*x*(c*x^4+b*x^2+a)^{(1/2)}/a/c^{(5/2)}/(-4*a*c+b^2)/(a^{(1/2)}+x^2*c^{(1/2)})-1/3*(a*B*(6*c^3*d^3-8*b^3*e^3-9*c^2*d*e*(6*a*e+b*d)+b*c*e^2*(29*a*e+18*b*d))+3*A*c*(2*a*b^2*e^3+6*a*c*e*(-a*e^2+c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*(cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))*EllipticE(sin(2*arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(11/4)}/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^{(1/2)}-1/6*(cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))*EllipticF(sin(2*arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(3*A*c^3*d^3-5*a^2*B*c*e^3+a*e*(-2*b*e+3*c*d)*(3*A*c*e-4*B*b*e+3*B*c*d)-3*c^{(5/2)}*d^2*(3*A*e+B*d)*a^{(1/2)}+3*a^{(3/2)}*e^2*(3*A*c*e-4*B*b*e+9*B*c*d)*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(11/4)}/(-2*a^{(1/2)}*c^{(1/2)}+b)/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A] time = 1.38, antiderivative size = 859, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$, Rules used = {1678, 1679, 1197, 1103, 1195}

$$\frac{Bx\sqrt{cx^4 + bx^2 + a}e^3}{3c^2} \frac{(aB(6c^3d^3 - 9c^2e(bd + 6ae)d - 8b^3e^3 + bce^2(18bd + 29ae)) + 3Ac(2ab^2e^3 + 6ac(cd^2 - ae^2))}{3a^{3/4}c^{11/4}(b^2 - 4ac)}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x]


```
[Out] (x*(A*c*(b^2*c*d^3 - 2*a*c*d*(c*d^2 - 3*a*e^2) - a*b*e*(3*c*d^2 + a*e^2)) +
a*B*(a*b^2*e^3 + 2*a*c*e*(3*c*d^2 - a*e^2) - b*c*d*(c*d^2 + 3*a*e^2)) - (a
*B*(2*c*d - b*e)*(c^2*d^2 + b^2*e^2 - c*e*(b*d + 3*a*e)) + A*c*(a*b^2*e^3 +
2*a*c*e*(3*c*d^2 - a*e^2) - b*c*d*(c*d^2 + 3*a*e^2)))*x^2)/(a*c^2*(b^2 -
4*a*c)*Sqrt[a + b*x^2 + c*x^4]) + (B*e^3*x*Sqrt[a + b*x^2 + c*x^4])/(3*c^2)
+ ((a*B*(6*c^3*d^3 - 8*b^3*e^3 - 9*c^2*d*e*(b*d + 6*a*e) + b*c*e^2*(18*b*d
+ 29*a*e)) + 3*A*c*(2*a*b^2*e^3 + 6*a*c*e*(c*d^2 - a*e^2) - b*c*d*(c*d^2 +
3*a*e^2)))*x*Sqrt[a + b*x^2 + c*x^4])/(3*a*c^(5/2)*(b^2 - 4*a*c)*(Sqrt[a
+ Sqrt[c]*x^2)) - ((a*B*(6*c^3*d^3 - 8*b^3*e^3 - 9*c^2*d*e*(b*d + 6*a*e) +
b*c*e^2*(18*b*d + 29*a*e)) + 3*A*c*(2*a*b^2*e^3 + 6*a*c*e*(c*d^2 - a*e^2) -
b*c*d*(c*d^2 + 3*a*e^2)))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)
/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b
/(Sqrt[a]*Sqrt[c]))/4])/(3*a^(3/4)*c^(11/4)*(b^2 - 4*a*c)*Sqrt[a + b*x^2 +
c*x^4]) - ((3*A*c^3*d^3 - 5*a^2*B*c*e^3 - 3*Sqrt[a]*c^(5/2)*d^2*(B*d + 3*A*
e) + a*e*(3*c*d - 2*b*e)*(3*B*c*d - 4*b*B*e + 3*A*c*e) + 3*a^(3/2)*Sqrt[c]*
e^2*(9*B*c*d - 4*b*B*e + 3*A*c*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2
+ c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)]
, (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(6*a^(3/4)*(b - 2*Sqrt[a]*Sqrt[c])*c^(11/4)
*Sqrt[a + b*x^2 + c*x^4])
```

Rule 1103

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c
/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*
EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(2*q*Sqrt[a + b*x^2 + c*x^4]
), x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1195

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol]
:= With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^
2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^
2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(q*Sqrt[a + b*x^2 + c
*x^4]), x] /; EqQ[e + d*q^2, 0]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 -
4*a*c, 0] && PosQ[c/a]
```

Rule 1197

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol]
:= With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4]
], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; Ne
Q[e + d*q, 0]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[
c/a]
```

Rule 1678

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{d =
  Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 2]}, Simp[(x*(a + b*x^2 + c*x^4)^(p + 1)*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2))/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Dist[1/(2*a*(p + 1)*(b^2 - 4*a*c)), Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Pq, a + b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p + 7)*(b*d - 2*a*e)*x^2, x], x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x^2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]
```

Rule 1679

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{q =
  Expon[Pq, x^2], e = Coeff[Pq, x^2, Expon[Pq, x^2]]}, Simp[(e*x^(2*q - 3)*(a + b*x^2 + c*x^4)^(p + 1))/(c*(2*q + 4*p + 1)), x] + Dist[1/(c*(2*q + 4*p + 1)), Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*q + 4*p + 1)*Pq - a*e*(2*q - 3)*x^(2*q - 4) - b*e*(2*q + 2*p - 1)*x^(2*q - 2) - c*e*(2*q + 4*p + 1)*x^(2*q), x], x], x]] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pq, x^2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]
```

Rubi steps

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx = \frac{x \left(Ac \left(b^2 cd^3 - 2acd \left(cd^2 - 3ae^2 \right) - abe \left(3cd^2 + ae^2 \right) \right) + aB \left(ab^2 e^3 + 2ace \left(3cd^2 - ae^2 \right) \right) \right)}{(a + bx^2 + cx^4)^{3/2}}$$

$$= \frac{x \left(Ac \left(b^2 cd^3 - 2acd \left(cd^2 - 3ae^2 \right) - abe \left(3cd^2 + ae^2 \right) \right) + aB \left(ab^2 e^3 + 2ace \left(3cd^2 - ae^2 \right) \right) \right)}{(a + bx^2 + cx^4)^{3/2}}$$

$$= \frac{x \left(Ac \left(b^2 cd^3 - 2acd \left(cd^2 - 3ae^2 \right) - abe \left(3cd^2 + ae^2 \right) \right) + aB \left(ab^2 e^3 + 2ace \left(3cd^2 - ae^2 \right) \right) \right)}{(a + bx^2 + cx^4)^{3/2}}$$

Mathematica [C] time = 6.65, size = 5432, normalized size = 6.32

Result too large to show

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2),x]

[Out] Result too large to show

fricas [F] time = 1.56, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(Be^3x^8 + (3Bde^2 + Ae^3)x^6 + 3(Bd^2e + Ade^2)x^4 + Ad^3 + (Bd^3 + 3Ad^2e)x^2)\sqrt{cx^4 + bx^2 + a}}{c^2x^8 + 2bcx^6 + (b^2 + 2ac)x^4 + 2abx^2 + a^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] integral((B*e^3*x^8 + (3*B*d*e^2 + A*e^3)*x^6 + 3*(B*d^2*e + A*d*e^2)*x^4 + A*d^3 + (B*d^3 + 3*A*d^2*e)*x^2)*sqrt(c*x^4 + b*x^2 + a)/(c^2*x^8 + 2*b*c*x^6 + (b^2 + 2*a*c)*x^4 + 2*a*b*x^2 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^3/(c*x^4 + b*x^2 + a)^(3/2), x)

maple [B] time = 0.04, size = 2445, normalized size = 2.85

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x)

[Out] B*e^3*(-2*c*(-1/2*b/c^3*(3*a*c-b^2)/(4*a*c-b^2)*x^3-1/2*(2*a*c-b^2)*a/c^3/(4*a*c-b^2)*x)/((x^4+b/c*x^2+a/c)*c)^(1/2)+1/3/c^2*x*(c*x^4+b*x^2+a)^(1/2)+1/4*(-1/c^2*(2*a*c-b^2)*a/(4*a*c-b^2)-1/3*a/c^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-1/2*(-5/3*b/c^2-b/c^2*(3*a*c-b^2)/(4*a*c-b^2))*a*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)

$c-4)^{(1/2)} - \text{EllipticE}(1/2*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2}))/a)^{(1/2)}*x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2}))/a*b/c-4)^{(1/2})))$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^3/(c*x^4 + b*x^2 + a)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x)

[Out] int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)**3/(a + b*x**2 + c*x**4)**(3/2), x)

$$3.25 \quad \int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=628

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) \left(Ac(abe^2 - 4acde + bcd^2) - 2aB(-ce(3ae + bd) + b\right)}{a^{3/4}c^{7/4}(b^2 - 4ac) \sqrt{a + bx^2 + cx^4}}$$

[Out] $-x*(a*B*(a*b*e^2-4*a*c*d*e+b*c*d^2)-A*c*(b^2*d^2-2*a*b*d*e-2*a*(-a*e^2+c*d^2))-(A*c*(a*b*e^2-4*a*c*d*e+b*c*d^2)-a*B*(2*c^2*d^2+b^2*e^2-2*c*e*(a*e+b*d)))*x^2)/a/c/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^{(1/2)}-(A*c*(a*b*e^2-4*a*c*d*e+b*c*d^2)-2*a*B*(c^2*d^2+b^2*e^2-c*e*(3*a*e+b*d)))*x*(c*x^4+b*x^2+a)^{(1/2)}/a/c^{(3/2)}/(-4*a*c+b^2)/(a^{(1/2)}+x^2*c^{(1/2)})+(A*c*(a*b*e^2-4*a*c*d*e+b*c*d^2)-2*a*B*(c^2*d^2+b^2*e^2-c*e*(3*a*e+b*d)))*(cos(2*arctan(c^{(1/4)}*x/a^{(1/4)})))^2)^{(1/2)}/cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))*EllipticE(sin(2*arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(7/4)}/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^{(1/2)}-1/2*(cos(2*arctan(c^{(1/4)}*x/a^{(1/4)})))^2)^{(1/2)}/cos(2*arctan(c^{(1/4)}*x/a^{(1/4)}))*EllipticF(sin(2*arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(A*c^2*d^2+a*e*(A*c*e-2*B*b*e+2*B*c*d)-c^{(3/2)}*d*(2*A*e+B*d)*a^{(1/2)}+3*a^{(3/2)}*B*e^2*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(7/4)}/(-2*a^{(1/2)}*c^{(1/2)}+b)/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.58, antiderivative size = 633, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 4, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$, Rules used = {1678, 1197, 1103, 1195}

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) \left(Ac(abe^2 - 4acde + bcd^2) - 2aB(-ce(3ae + bd) + b\right)}{a^{3/4}c^{7/4}(b^2 - 4ac) \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] $-((x*(c*((a*B*(b*c*d^2 - 4*a*c*d*e + a*b*e^2))/c - A*(b^2*d^2 - 2*a*b*d*e - 2*a*(c*d^2 - a*e^2))) - (A*c*(b*c*d^2 - 4*a*c*d*e + a*b*e^2) - a*B*(2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)))*x^2))/(a*c*(b^2 - 4*a*c)*Sqrt[a + b*x^2 + c*x^4])) - ((A*c*(b*c*d^2 - 4*a*c*d*e + a*b*e^2) - 2*a*B*(c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)))*x^2)/(a*c*(b^2 - 4*a*c)*Sqrt[a + b*x^2 + c*x^4]))$

$$e^2 - c*e*(b*d + 3*a*e))*x*\text{Sqrt}[a + b*x^2 + c*x^4]/(a*c^{(3/2)}*(b^2 - 4*a*c)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2) + ((A*c*(b*c*d^2 - 4*a*c*d*e + a*b*e^2) - 2*a*B*(c^2*d^2 + b^2*e^2 - c*e*(b*d + 3*a*e)))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(a^{(3/4)}*c^{(7/4)}*(b^2 - 4*a*c)*\text{Sqrt}[a + b*x^2 + c*x^4]) - ((A*c^2*d^2 + 3*a^{(3/2)}*B*\text{Sqrt}[c]*e^2 - \text{Sqrt}[a]*c^{(3/2)}*d*(B*d + 2*A*e) + a*e*(2*B*c*d - 2*b*B*e + A*c*e))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^{(3/4)}*(b - 2*\text{Sqrt}[a]*\text{Sqrt}[c])*c^{(7/4)}*\text{Sqrt}[a + b*x^2 + c*x^4])$$

Rule 1103

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(2*q*Sqrt[a + b*x^2 + c*x^4]), x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1195

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(q*Sqrt[a + b*x^2 + c*x^4]), x] /; EqQ[e + d*q^2, 0]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1197

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1678

```
Int[(Pq_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{d = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 2]}, Simp[(x*(a + b*x^2 + c*x^4)^(p + 1)*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2))/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Dist[1/(2*a*(p + 1)*(b^2 - 4*a*c)), Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Pq, a + b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p + 7)*(b*d - 2*a*e)*x^2, x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x^
```

2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]

Rubi steps

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx = -\frac{x \left(c \left(\frac{aB(bcd^2 - 4acde + abe^2)}{c} - A(b^2d^2 - 2abde - 2a(cd^2 - ae^2)) \right) - (Ac(bcd^2 - 4acde) \right)}{ac(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}}$$

$$= -\frac{x \left(c \left(\frac{aB(bcd^2 - 4acde + abe^2)}{c} - A(b^2d^2 - 2abde - 2a(cd^2 - ae^2)) \right) - (Ac(bcd^2 - 4acde) \right)}{ac(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}}$$

$$= -\frac{x \left(c \left(\frac{aB(bcd^2 - 4acde + abe^2)}{c} - A(b^2d^2 - 2abde - 2a(cd^2 - ae^2)) \right) - (Ac(bcd^2 - 4acde) \right)}{ac(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}}$$

Mathematica [C] time = 4.72, size = 766, normalized size = 1.22

$$\frac{-4cx\sqrt{\frac{c}{\sqrt{b^2-4ac}+b}} \left(Ac(2a^2e^2 + abe(ex^2 - 2d) - 2acd(d + 2ex^2) + b^2d^2 + bcd^2x^2) - aB(abe^2 - 2ace(2d + ex^2) + \right)}{\dots}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2),x]

[Out] (-4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x*(-(a*B*(a*b*e^2 + 2*c^2*d^2*x^2 + b^2*e^2*x^2 + b*c*d*(d - 2*e*x^2) - 2*a*c*e*(2*d + e*x^2))) + A*c*(b^2*d^2 + 2*a^2*e^2 + b*c*d^2*x^2 + a*b*e*(-2*d + e*x^2) - 2*a*c*d*(d + 2*e*x^2))) - I*(-b + Sqrt[b^2 - 4*a*c])*(-(A*c*(b*c*d^2 - 4*a*c*d*e + a*b*e^2)) + 2*a*B*(c^2*d^2 + b^2*e^2 - c*e*(b*d + 3*a*e)))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]) + I*(2*a*B*(b^2*(-b + Sqrt[b^2 - 4*a*c])*e^2 + c^2*d*(Sqrt[b^2 - 4*a*c]*d - 4*a*e) + c*e*(b^2*d - b*Sqrt[b^2 - 4*a*c]*d + 4*a*b*e - 3*a*Sqrt[b^2 - 4*a*c]*e)) + A*c*(b^2*(c*d^2 + a*e^2) - b*Sqrt[b^2 - 4*a*c]*(c*d^2 + a*e^2) - 4*a*c*(c*d^2 - Sqrt[b^2 - 4*a*c]*d*e + a*e^2))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqr

$t[b^2 - 4ac]) * \text{EllipticF}[I * \text{ArcSinh}[\text{Sqrt}[2] * \text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4ac])] * x], (b + \text{Sqrt}[b^2 - 4ac]) / (b - \text{Sqrt}[b^2 - 4ac])] / (4ac^2(-b^2 + 4ac) * \text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4ac])] * \text{Sqrt}[a + bx^2 + cx^4])$

fricas [F] time = 1.51, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(Be^2x^6 + (2Bde + Ae^2)x^4 + Ad^2 + (Bd^2 + 2Ade)x^2)\sqrt{cx^4 + bx^2 + a}}{c^2x^8 + 2bcx^6 + (b^2 + 2ac)x^4 + 2abx^2 + a^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] integral((B*e^2*x^6 + (2*B*d*e + A*e^2)*x^4 + A*d^2 + (B*d^2 + 2*A*d*e)*x^2)*sqrt(c*x^4 + b*x^2 + a)/(c^2*x^8 + 2*b*c*x^6 + (b^2 + 2*a*c)*x^4 + 2*a*b*x^2 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^2/(c*x^4 + b*x^2 + a)^(3/2), x)

maple [B] time = 0.01, size = 1891, normalized size = 3.01

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x)

[Out] $B*e^2*(-2*(-1/2/(4*a*c-b^2)*a*b/c^2*x+1/2*(2*a*c-b^2)/(4*a*c-b^2)/c^2*x^3)/((x^4+b/c*x^2+a/c)*c)^{(1/2)*c-1/4/c*a*b/(4*a*c-b^2)*2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)*(-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2+4)^{(1/2)*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2+4)^{(1/2)/(c*x^4+b*x^2+a)^{(1/2)*\text{EllipticF}(1/2*2^{(1/2)*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*b/c-4)^{(1/2))-1/2*(1/c+(2*a*c-b^2)/(4*a*c-b^2)/c)*a*2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)*(-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2+4)^{(1/2)*(2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2+4)^{(1/2)/(c*x^4+b*x^2+a)^{(1/2)/(b+(-4*a*c+b^2)^{(1/2)})*(\text{EllipticF}(1$

```

/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))
)/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x
,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))))+(A*e^2+2*B*d*e)*(-2*(1/2/(
4*a*c-b^2)*b/c*x^3+1/(4*a*c-b^2)*a/c*x)/((x^4+b/c*x^2+a/c)*c)^(1/2)*c+1/2*a
/(4*a*c-b^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)
^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^
2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2
*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-1/2/(4*a*c-b^2)*b*a*2^(1/2)/((-b+(-
4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b
+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(
1/2))*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+
(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)
^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))))+(2*A*d*e
+B*d^2)*(-2*(-1/(4*a*c-b^2)*x^3-1/2/(4*a*c-b^2)*b/c*x)/((x^4+b/c*x^2+a/c)*c
)^(1/2)*c-1/4/(4*a*c-b^2)*b*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(
-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1
/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)
^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))+c/(4*a*c-b^2)*a*2^(1
/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(
1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4
*a*c+b^2)^(1/2))*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x
,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-
4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2)
))+A*d^2*(-2*(1/2/(4*a*c-b^2)/a*b*x^3-1/2*(2*a*c-b^2)/(4*a*c-b^2)/a/c*x)/((
x^4+b/c*x^2+a/c)*c)^(1/2)*c+1/4*(1/a-(2*a*c-b^2)/(4*a*c-b^2)/a)*2^(1/2)/((-
b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(
2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2
*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2)))/
a*b/c-4)^(1/2))-1/2*b/(4*a*c-b^2)*c*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/
2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x
^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticF(1/2*2^(
1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/
c-4)^(1/2))-EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x,1/2*(
2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))))

```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")
```

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^2/(c*x^4 + b*x^2 + a)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2), x)

[Out] int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+b*x**2+a)**(3/2), x)

[Out] Integral((A + B*x**2)*(d + e*x**2)**2/(a + b*x**2 + c*x**4)**(3/2), x)

$$3.26 \quad \int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=481

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (Ac(bd - 2ae) - aB(2cd - be)) (\sqrt{a} + \sqrt{c}x^2) (\sqrt{a}E$$

$$a^{3/4}c^{3/4} (b^2 - 4ac) \sqrt{a + bx^2 + cx^4} +$$

[Out] $-x*(a*B*(-2*a*e+b*d)-A*(-a*b*e-2*a*c*d+b^2*d)-(A*c*(-2*a*e+b*d)-a*B*(-b*e+2*c*d))*x^2)/a/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^{(1/2)}-(A*c*(-2*a*e+b*d)-a*B*(-b*e+2*c*d))*x*(c*x^4+b*x^2+a)^{(1/2)}/a/(-4*a*c+b^2)/c^{(1/2)}/(a^{(1/2)}+x^2*c^{(1/2)})+(A*c*(-2*a*e+b*d)-a*B*(-b*e+2*c*d))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^{(1/2)})/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)}))*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(3/4)}/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^{(1/2)}+1/2*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^{(1/2)})/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(B*a^{(1/2)}-A*c^{(1/2)})*(-e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)}))*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(3/4)}/(-2*a^{(1/2)}*c^{(1/2)}+b)/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A] time = 0.39, antiderivative size = 481, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 31, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used = {1678, 1197, 1103, 1195}

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (Ac(bd - 2ae) - aB(2cd - be)) (\sqrt{a} + \sqrt{c}x^2) (\sqrt{a}E$$

$$a^{3/4}c^{3/4} (b^2 - 4ac) \sqrt{a + bx^2 + cx^4} +$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] $-((x*(a*B*(b*d - 2*a*e) - A*(b^2*d - 2*a*c*d - a*b*e) - (A*c*(b*d - 2*a*e) - a*B*(2*c*d - b*e))*x^2))/(a*(b^2 - 4*a*c)*\text{Sqrt}[a + b*x^2 + c*x^4]) - ((A*c*(b*d - 2*a*e) - a*B*(2*c*d - b*e))*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(a*\text{Sqrt}[c]*(b^2 - 4*a*c)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) + ((A*c*(b*d - 2*a*e) - a*B*(2*c*d - b*e))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(a^{(3/4)}*c^{(3/4)}*(b^2 - 4*a*c)*\text{Sqrt}[a + b*x^2 + c*x^4]) + ((\text{Sqrt}[a]*B - A*\text{Sqrt}[c])*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)])$

$$\frac{2 + c*x^4}{(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2} * \text{EllipticF}[2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4]/(2*a^{3/4}*(b - 2*\text{Sqrt}[a]*\text{Sqrt}[c])*c^{3/4})*\text{Sqrt}[a + b*x^2 + c*x^4]$$

Rule 1103

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(2*q*Sqrt[a + b*x^2 + c*x^4]), x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1195

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(q*Sqrt[a + b*x^2 + c*x^4]), x] /; EqQ[e + d*q^2, 0]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1197

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1678

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{d = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 2]}, Simp[(x*(a + b*x^2 + c*x^4)^(p + 1)*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2))/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Dist[1/(2*a*(p + 1)*(b^2 - 4*a*c)), Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Pq, a + b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p + 7)*(b*d - 2*a*e)*x^2, x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x^2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]
```

Rubi steps

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{3/2}} dx = -\frac{x(aB(bd - 2ae) - A(b^2d - 2acd - abe) - (Ac(bd - 2ae) - aB(2cd - be))x^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} - \frac{\int \frac{dx}{\sqrt{a + bx^2 + cx^4}}}{a(b^2 - 4ac)}$$

$$= -\frac{x(aB(bd - 2ae) - A(b^2d - 2acd - abe) - (Ac(bd - 2ae) - aB(2cd - be))x^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} + \frac{\left(\int \frac{dx}{\sqrt{a + bx^2 + cx^4}}\right)}{a(b^2 - 4ac)}$$

$$= -\frac{x(aB(bd - 2ae) - A(b^2d - 2acd - abe) - (Ac(bd - 2ae) - aB(2cd - be))x^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} - \frac{(Ac(bd - 2ae) - aB(2cd - be))}{a(b^2 - 4ac)}$$

Mathematica [F] time = 0.00, size = 0, normalized size = 0.00

\$Aborted

Verification is Not applicable to the result.

[In] Integrate[((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] \$Aborted

fricas [F] time = 1.28, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(Bex^4 + (Bd + Ae)x^2 + Ad)\sqrt{cx^4 + bx^2 + a}}{c^2x^8 + 2bcx^6 + (b^2 + 2ac)x^4 + 2abx^2 + a^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2), x, algorithm="fricas")

[Out] integral((B*e*x^4 + (B*d + A*e)*x^2 + A*d)*sqrt(c*x^4 + b*x^2 + a)/(c^2*x^8 + 2*b*c*x^6 + (b^2 + 2*a*c)*x^4 + 2*a*b*x^2 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)/(c*x^4 + b*x^2 + a)^(3/2), x)

maple [B] time = 0.01, size = 1390, normalized size = 2.89

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x)

[Out]
$$B * e * (-2 * (1/2 / (4 * a * c - b^2)) * b / c * x^3 + 1 / (4 * a * c - b^2) * a / c * x) / ((x^4 + b / c * x^2 + a / c) * c)^{(1/2)} * c + 1/2 * a / (4 * a * c - b^2) * 2^{(1/2)} / ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * (-2 * (-b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} / (c * x^4 + b * x^2 + a)^{(1/2)} * \text{EllipticF}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)} - 1/2 / (4 * a * c - b^2) * b * a * 2^{(1/2)} / ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * (-2 * (-b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} / (c * x^4 + b * x^2 + a)^{(1/2)} / (b + (-4 * a * c + b^2)^{(1/2)}) * (\text{EllipticF}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)} - \text{EllipticE}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)})) + (A * e + B * d) * (-2 * (-1 / (4 * a * c - b^2)) * x^3 - 1/2 / (4 * a * c - b^2) * b / c * x) / ((x^4 + b / c * x^2 + a / c) * c)^{(1/2)} * c - 1/4 / (4 * a * c - b^2) * b * 2^{(1/2)} / ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * (-2 * (-b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} / (c * x^4 + b * x^2 + a)^{(1/2)} * \text{EllipticF}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)} + c / (4 * a * c - b^2) * a * 2^{(1/2)} / ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * (-2 * (-b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} / (c * x^4 + b * x^2 + a)^{(1/2)} / (b + (-4 * a * c + b^2)^{(1/2)}) * (\text{EllipticF}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)} - \text{EllipticE}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)})) + A * d * (-2 * (1/2 / (4 * a * c - b^2)) / a * b * x^3 - 1/2 * (2 * a * c - b^2) / (4 * a * c - b^2) / a / c * x) / ((x^4 + b / c * x^2 + a / c) * c)^{(1/2)} * c + 1/4 * (1 / a - (2 * a * c - b^2) / (4 * a * c - b^2) / a) * 2^{(1/2)} / ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * (-2 * (-b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} / (c * x^4 + b * x^2 + a)^{(1/2)} * \text{EllipticF}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)} - 1/2 * b / (4 * a * c - b^2) * c * 2^{(1/2)} / ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * (-2 * (-b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * x^2 + 4)^{(1/2)} / (c * x^4 + b * x^2 + a)^{(1/2)} / (b + (-4 * a * c + b^2)^{(1/2)}) * (\text{EllipticF}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)} - \text{EllipticE}(1/2 * 2^{(1/2)} * ((-b + (-4 * a * c + b^2)^{(1/2)}) / a)^{(1/2)} * x, 1/2 * (2 * (b + (-4 * a * c + b^2)^{(1/2)}) / a * b / c - 4)^{(1/2)}))$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)/(c*x^4 + b*x^2 + a)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)/(a + b*x**2 + c*x**4)**(3/2), x)

$$3.27 \quad \int \frac{A+Bx^2}{(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=398

$$\frac{\sqrt[4]{c} (\sqrt{a} + \sqrt{c} x^2) (Ab - 2aB) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (\sqrt{a} + \sqrt{c} x^2) (\sqrt{a} B - A\sqrt{c}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{a^{3/4} (b^2 - 4ac) \sqrt{a + bx^2 + cx^4}} + \frac{(\sqrt{a} + \sqrt{c} x^2) (\sqrt{a} B - A\sqrt{c}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{2a^{3/4} \sqrt[4]{c} (b - 2)}$$

[Out] $x*(A*b^2-a*b*B-2*a*A*c+(A*b-2*B*a)*c*x^2)/a/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)-(A*b-2*B*a)*x*c^(1/2)*(c*x^4+b*x^2+a)^(1/2)/a/(-4*a*c+b^2)/(a^(1/2)+x^2*c^(1/2))+(A*b-2*B*a)*c^(1/4)*(\cos(2*\arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/\cos(2*\arctan(c^(1/4)*x/a^(1/4)))*\text{EllipticE}(\sin(2*\arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)+1/2*(\cos(2*\arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/\cos(2*\arctan(c^(1/4)*x/a^(1/4)))*\text{EllipticF}(\sin(2*\arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(B*a^(1/2)-A*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/c^(1/4)/(-2*a^(1/2)*c^(1/2)+b)/(c*x^4+b*x^2+a)^(1/2)$

Rubi [A] time = 0.20, antiderivative size = 398, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1178, 1197, 1103, 1195}

$$\frac{\sqrt[4]{c} (\sqrt{a} + \sqrt{c} x^2) (Ab - 2aB) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (\sqrt{a} + \sqrt{c} x^2) (\sqrt{a} B - A\sqrt{c}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{a^{3/4} (b^2 - 4ac) \sqrt{a + bx^2 + cx^4}} + \frac{(\sqrt{a} + \sqrt{c} x^2) (\sqrt{a} B - A\sqrt{c}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}}{2a^{3/4} \sqrt[4]{c} (b - 2)}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] $(x*(A*b^2 - a*b*B - 2*a*A*c + (A*b - 2*a*B)*c*x^2))/(a*(b^2 - 4*a*c)*\text{Sqrt}[a + b*x^2 + c*x^4]) - ((A*b - 2*a*B)*\text{Sqrt}[c]*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(a*(b^2 - 4*a*c)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) + ((A*b - 2*a*B)*c^(1/4)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(a^(3/4)*(b^2 - 4*a*c)*\text{Sqrt}[a + b*x^2 + c*x^4]) + ((\text{Sqrt}[a]*B - A*\text{Sqrt}[c])*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^(3/4)*(b - 2*\text{Sqrt}[a]*\text{Sqrt}[c])*c^(1/4)*\text{Sqrt}[a + b*x^2 + c*x^4])$

Rule 1103

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(2*q*Sqrt[a + b*x^2 + c*x^4]), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1178

```
Int[((d_) + (e_.)*(x_)^2)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[(x*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)*(a + b*x^2 + c*x^4)^(p + 1))/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Dist[1/(2*a*(p + 1)*(b^2 - 4*a*c)), Int[Simp[(2*p + 3)*d*b^2 - a*b*e - 2*a*c*d*(4*p + 5) + (4*p + 7)*(d*b - 2*a*e)*c*x^2, x]*(a + b*x^2 + c*x^4)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2*p]
```

Rule 1195

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/(a*(1 + q^2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)])/(q*Sqrt[a + b*x^2 + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1197

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx &= \frac{x(Ab^2 - abB - 2aAc + (Ab - 2aB)cx^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} - \frac{\int \frac{-a(bB - 2Ac) + (Ab - 2aB)cx^2}{\sqrt{a + bx^2 + cx^4}} dx}{a(b^2 - 4ac)} \\
&= \frac{x(Ab^2 - abB - 2aAc + (Ab - 2aB)cx^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} + \frac{((Ab - 2aB)\sqrt{c}) \int \frac{1 - \frac{\sqrt{c}x^2}{\sqrt{a}}}{\sqrt{a + bx^2 + cx^4}} dx}{\sqrt{a}(b^2 - 4ac)} - \frac{((Ab - 2aB)\sqrt{c}) \int \frac{1 + \frac{\sqrt{c}x^2}{\sqrt{a}}}{\sqrt{a + bx^2 + cx^4}} dx}{\sqrt{a}(b^2 - 4ac)} \\
&= \frac{x(Ab^2 - abB - 2aAc + (Ab - 2aB)cx^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} - \frac{(Ab - 2aB)\sqrt{c}x\sqrt{a + bx^2 + cx^4}}{a(b^2 - 4ac)(\sqrt{a} + \sqrt{c}x^2)} + \frac{(Ab - 2aB)\sqrt{c}}{a(b^2 - 4ac)(\sqrt{a} - \sqrt{c}x^2)}
\end{aligned}$$

Mathematica [F] time = 0.00, size = 0, normalized size = 0.00

\$Aborted

Verification is Not applicable to the result.

[In] Integrate[(A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] \$Aborted

fricas [F] time = 1.31, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{cx^4 + bx^2 + a}(Bx^2 + A)}{c^2x^8 + 2bcx^6 + (b^2 + 2ac)x^4 + 2abx^2 + a^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(c*x^4 + b*x^2 + a)*(B*x^2 + A)/(c^2*x^8 + 2*b*c*x^6 + (b^2 + 2*a*c)*x^4 + 2*a*b*x^2 + a^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(c*x^4 + b*x^2 + a)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2),x)

[Out] int((A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Integral((A + B*x**2)/(a + b*x**2 + c*x**4)**(3/2), x)

$$3.28 \quad \int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=867

$$\frac{a^{3/4}e(Bd - Ae) \left(\sqrt{c}x^2 + \sqrt{a} \right) \sqrt{\frac{cx^4+bx^2+a}{(\sqrt{c}x^2+\sqrt{a})^2}} \Pi \left(-\frac{(\sqrt{c}d-\sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1} \left(\frac{\sqrt[4]{c}x}{\sqrt{a}} \right) \middle| \frac{1}{4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right) \left(\frac{\sqrt{c}d}{\sqrt{a}} + e \right)^2 e^{3/2}(Bd - A)}{4\sqrt[4]{c}d(cd^2 - ae^2)(cd^2 - bed + ae^2)\sqrt{cx^4 + bx^2 + a}} \quad 2\sqrt{a}$$

[Out] $-1/2*e^{(3/2)}*(-A*e+B*d)*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)})/(a*e^2-b*d*e+c*d^2)^{(3/2)}/d^{(1/2)}-x*(a*b*c*(-A*e+B*d)-(-2*a*c+b^2)*(-A*b*e+A*c*d+B*a*e)+c*(a*B*(-b*e+2*c*d)-A*(2*a*c*e-b^2*e+b*c*d))*x^2)/a/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}+(a*B*(-b*e+2*c*d)-A*(2*a*c*e-b^2*e+b*c*d))*x*c^{(1/2)}*(c*x^4+b*x^2+a)^{(1/2)}/a/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(a^{(1/2)}+x^2*c^{(1/2)})-c^{(1/4)}*(a*B*(-b*e+2*c*d)-A*(2*a*c*e-b^2*e+b*c*d))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}+1/4*a^{(3/4)}*e*(-A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)}, 1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(e+d*c^{(1/2)}/a^{(1/2)})^2*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(1/4)}/d/(-a*e^2+c*d^2)/(a*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}+1/2*c^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), 1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(B*a^{(1/2)}-A*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/(-e*a^{(1/2)}+d*c^{(1/2)})/(-2*a^{(1/2)}*c^{(1/2)}+b)/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A] time = 1.42, antiderivative size = 1045, normalized size of antiderivative = 1.21, number of steps used = 9, number of rules used = 7, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {1720, 1178, 1197, 1103, 1195, 1216, 1706}

$$\frac{a^{3/4}e(Bd - Ae) \left(\sqrt{c}x^2 + \sqrt{a} \right) \sqrt{\frac{cx^4+bx^2+a}{(\sqrt{c}x^2+\sqrt{a})^2}} \Pi \left(-\frac{(\sqrt{c}d-\sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1} \left(\frac{\sqrt[4]{c}x}{\sqrt{a}} \right) \middle| \frac{1}{4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right) \left(\frac{\sqrt{c}d}{\sqrt{a}} + e \right)^2 e^{3/2}(Bd - A)}{4\sqrt[4]{c}d(cd^2 - ae^2)(cd^2 - bed + ae^2)\sqrt{cx^4 + bx^2 + a}} \quad 2\sqrt{a}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)),x]

```
[Out] -((x*(a*b*c*(B*d - A*e) - (b^2 - 2*a*c)*(A*c*d - A*b*e + a*B*e) + c*(a*B*(2
*c*d - b*e) - A*(b*c*d - b^2*e + 2*a*c*e))*x^2))/(a*(b^2 - 4*a*c)*(c*d^2 -
b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*x^4])) + (Sqrt[c]*(a*B*(2*c*d - b*e) - A*
(b*c*d - b^2*e + 2*a*c*e))*x*Sqrt[a + b*x^2 + c*x^4))/(a*(b^2 - 4*a*c)*(c*d
^2 - b*d*e + a*e^2)*(Sqrt[a] + Sqrt[c]*x^2)) - (e^(3/2)*(B*d - A*e)*ArcTan[
(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4]))]
/(2*Sqrt[d]*(c*d^2 - b*d*e + a*e^2)^(3/2)) - (c^(1/4)*(a*B*(2*c*d - b*e) -
A*(b*c*d - b^2*e + 2*a*c*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^
4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 -
b/(Sqrt[a]*Sqrt[c]))/4])/(a^(3/4)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)*Sq
rt[a + b*x^2 + c*x^4]) - (c^(1/4)*e*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqr
t[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4
)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^(1/4)*(Sqrt[c]*d - Sqrt[a
]*e)*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*x^4]) - (c^(1/4)*(a*B*e - S
qrt[a]*Sqrt[c]*(B*d - A*e) + A*(c*d - b*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a
+ b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)
/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^(3/4)*(b - 2*Sqrt[a]*Sqrt[c])
*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*x^4]) + (a^(3/4)*e*((Sqrt[c]*d)
/Sqrt[a] + e)^2*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4
)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-(Sqrt[c]*d - Sqrt[a]*e)^2/(4*Sqrt[
a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4
])/(4*c^(1/4)*d*(c*d^2 - a*e^2)*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*
x^4])
```

Rule 1103

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c
/a, 4]}, Simp[((1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]*
EllipticF[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c))]/(2*q*Sqrt[a + b*x^2 + c*x^4]
), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1178

```
Int[((d_) + (e_.)*(x_)^2)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symb
ol] := Simp[(x*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)*(a + b*x^2 +
c*x^4)^(p + 1))/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Dist[1/(2*a*(p + 1)*(b^2
- 4*a*c)), Int[Simp[(2*p + 3)*d*b^2 - a*b*e - 2*a*c*d*(4*p + 5) + (4*p + 7
)*(d*b - 2*a*e)*c*x^2, x]*(a + b*x^2 + c*x^4)^(p + 1), x], x] /; FreeQ[{a,
b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] &&
LtQ[p, -1] && IntegerQ[2*p]
```

Rule 1195

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol]
:= With[{q = Rt[c/a, 4]}, -Simp[(d*x*Sqrt[a + b*x^2 + c*x^4])/a*(1 + q^
2*x^2)), x] + Simp[(d*(1 + q^2*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^
2))], x]
```

2)^2))*EllipticE[2*ArcTan[q*x], 1/2 - (b*q^2)/(4*c)]/(q*Sqrt[a + b*x^2 + c*x^4]), x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1197

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] :> With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1216

Int[1/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] :> With[{q = Rt[c/a, 2]}, Dist[(c*d + a*e*q)/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[(a*e*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]

Rule 1706

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] :> With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[(Rt[-b + (c*d)/e + (a*e)/d, 2]*x)/Sqrt[a + b*x^2 + c*x^4]]/(2*d*e*Rt[-b + (c*d)/e + (a*e)/d, 2]), x] + Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a + b*x^2 + c*x^4))/(a*(A + B*x^2)^2)]*EllipticPi[Cancel[-((B*d - A*e)^2/(4*d*e*A*B))], 2*ArcTan[q*x], 1/2 - (b*A)/(4*a*B)]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4]), x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

Rule 1720

Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] :> Int[ExpandIntegrand[1/Sqrt[a + b*x^2 + c*x^4], Px*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx &= \int \left(\frac{Acd - Abe + aBe + c(Bd - Ae)x^2}{(cd^2 - bde + ae^2)(a + bx^2 + cx^4)^{3/2}} + \frac{e(-Bd + Ae)}{(cd^2 - bde + ae^2)(d + ex^2)\sqrt{a + bx^2 + cx^4}} \right) dx \\
&= \frac{\int \frac{Acd - Abe + aBe + c(Bd - Ae)x^2}{(a + bx^2 + cx^4)^{3/2}} dx}{cd^2 - bde + ae^2} - \frac{(e(Bd - Ae)) \int \frac{1}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx}{cd^2 - bde + ae^2} \\
&= -\frac{x(abc(Bd - Ae) - (b^2 - 2ac)(Acd - Abe + aBe) + c(aB(2cd - be) - A(b^2 - 4ac)))}{a(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a + bx^2 + cx^4}} \\
&= -\frac{x(abc(Bd - Ae) - (b^2 - 2ac)(Acd - Abe + aBe) + c(aB(2cd - be) - A(b^2 - 4ac)))}{a(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a + bx^2 + cx^4}} \\
&= -\frac{x(abc(Bd - Ae) - (b^2 - 2ac)(Acd - Abe + aBe) + c(aB(2cd - be) - A(b^2 - 4ac)))}{a(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a + bx^2 + cx^4}}
\end{aligned}$$

Mathematica [C] time = 5.06, size = 1736, normalized size = 2.00

$$4A\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}dexb^3 + 4Ac\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}dex^3b^2 - 4Ac\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}d^2xb^2 - 4aB\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}dexb^2 + 2iaAe^2\sqrt{\frac{2cx}{b}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)),x]

[Out] (-4*A*b^2*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x + 4*a*b*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x + 8*a*A*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x + 4*A*b^3*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x - 4*a*b^2*B*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x - 12*a*A*b*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x + 8*a^2*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x - 4*A*b*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x^3 + 8*a*B*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x^3 + 4*A*b^2*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x^3 - 4*a*b*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x^3 - 8*a*A*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x^3 - I*(-b + Sqrt[b^2 - 4*a*c])*d*(a*B*(2*c*d - b*e) + A*(-(b*c*d) + b^2*e - 2*a*c*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]

```

a*c)))*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*
EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b
^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + I*d*(a*B*(b*(b - Sqrt[b^2 - 4*a*c])
*e + 2*c*(Sqrt[b^2 - 4*a*c]*d - 2*a*e)) + A*(-(b^3*e) + b*c*(-(Sqrt[b^2 - 4
*a*c]*d) + 4*a*e) + b^2*(c*d + Sqrt[b^2 - 4*a*c]*e) - 2*a*c*(2*c*d + Sqrt[b
^2 - 4*a*c]*e)))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a
*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*E
llipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^
2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - (2*I)*a*b^2*B*d*e*Sqrt[(b + Sqrt[b^2
- 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*
c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticPi[((b + Sqrt[b^2 - 4*a*c])
*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt
[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + (8*I)*a^2*B*c*d*e*Sqrt[(b + Sqrt[
b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4
*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticPi[((b + Sqrt[b^2 - 4*a*c
])e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + S
qrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + (2*I)*a*A*b^2*e^2*Sqrt[(b + Sq
rt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2
- 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticPi[((b + Sqrt[b^2 - 4*
a*c])e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b
+ Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - (8*I)*a^2*A*c*e^2*Sqrt[(b +
Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b
^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticPi[((b + Sqrt[b^2 -
4*a*c])e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x],
(b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])]/(4*a*(-b^2 + 4*a*c)*Sqrt[
c/(b + Sqrt[b^2 - 4*a*c])]*d*(c*d^2 + e*(-(b*d) + a*e))*Sqrt[a + b*x^2 + c
x^4])

```

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

$(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))/a/(4*a*c-b^2)/(a*e^2-b*d*e+c*d^2)*b^2*c*d-c^2/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(2/a*b*x^2-2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)*(2/a*b*x^2+2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))*a*e+1/2*c/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(2/a*b*x^2-2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)*(2/a*b*x^2+2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))*b^2*2*e-1/2*c^2/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(2/a*b*x^2-2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)*(2/a*b*x^2+2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))*b*d+c^2/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(2/a*b*x^2-2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)*(2/a*b*x^2+2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))*a*e-1/2*c/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(2/a*b*x^2-2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)*(2/a*b*x^2+2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))*b^2*e+1/2*c^2/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(2/a*b*x^2-2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)*(2/a*b*x^2+2*(-4*a*c+b^2)^(1/2)/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticE(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))*b*d+1/(a*e^2-b*d*e+c*d^2)*e^2/d*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(1/2/a*b*x^2-1/2*(-4*a*c+b^2)^(1/2)/a*x^2+1)^(1/2)*(1/2/a*b*x^2+1/2*(-4*a*c+b^2)^(1/2)/a*x^2+1)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticPi(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2))*x, -2/(-b+(-4*a*c+b^2)^(1/2))*a/d*e, (-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)))$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(e*x^2 + d)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{B x^2 + A}{(e x^2 + d) (c x^4 + b x^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)), x)

[Out] int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+b*x**2+a)**(3/2), x)

[Out] Timed out

$$3.29 \quad \int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=1301

$$\frac{(Bd - Ae)x\sqrt{cx^4 + bx^2 + a}e^3}{2d(cd^2 - bed + ae^2)^2(ex^2 + d)} + \frac{(Ae(7cd^2 - e(4bd - ae)) - Bd(5cd^2 - e(2bd + ae))) \tan^{-1}\left(\frac{\sqrt{cd^2 - bed + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{cx^4 + bx^2 + a}}\right)e^3}{4d^{3/2}(cd^2 - bed + ae^2)^{5/2}}$$

[Out] $1/4e^{(3/2)}*(A*e*(7*c*d^2-e*(-a*e+4*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d)))*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})/d^{(3/2)}/(a*e^2-b*d*e+c*d^2)^{(5/2)}+x*(a*b*c*(A*e*(-b*e+2*c*d)-B*(-a*e^2+c*d^2))+(-2*a*c+b^2)*(a*B*e*(-b*e+2*c*d)+A*(c^2*d^2+b^2*e^2-c*e*(a*e+2*b*d)))-c*(a*B*(2*c^2*d^2+b^2*e^2-2*c*e*(a*e+b*d))+A*(2*b^2*c*d*e-4*a*c^2*d*e-b^3*e^2-b*c*(-3*a*e^2+c*d^2)))*x^2)/a/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)^2/(c*x^4+b*x^2+a)^{(1/2)}-1/2*e^3*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^{(1/2)}/d/(a*e^2-b*d*e+c*d^2)^2/(e*x^2+d)+1/2*(a*B*d*(-4*c^2*d^2-3*b^2*e^2+4*c*e*(2*a*e+b*d))+A*(2*b^3*d*e^2+2*b*c*d*(-3*a*e^2+c*d^2)-4*a*c*e*(a*e^2-2*c*d^2)+b^2*(a*e^3-4*c*d^2*e)))*x*c^{(1/2)}*(c*x^4+b*x^2+a)^{(1/2)}/a/(4*a*c-b^2)/d/(c*d^2+e*(a*e-b*d))^2/(a^{(1/2)}+x^2*c^{(1/2)})-1/2*c^{(1/4)}*(a*B*d*(4*c^2*d^2+3*b^2*e^2-4*c*e*(2*a*e+b*d))-A*(2*b^3*d*e^2+2*b*c*d*(-3*a*e^2+c*d^2)-4*a*c*e*(a*e^2-2*c*d^2)+b^2*(a*e^3-4*c*d^2*e)))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^2)^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(3/4)}/(-4*a*c+b^2)/d/(c*d^2+e*(a*e-b*d))^2/(c*x^4+b*x^2+a)^{(1/2)}-1/8*e*(A*e*(7*c*d^2-e*(-a*e+4*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d)))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),-1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*(2-b/a^{(1/2)}/c^{(1/2)})^2)^{(1/2)}*(e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^2/(a*e^2-b*d*e+c*d^2)^2/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}+1/2*c^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^2)^{(1/2)}*((-A*e+B*d)*(-b*e+c*d)*a^{(1/2)}+a*e*(-2*A*e+B*d)*c^{(1/2)}+A*d*(b*e-c*d)*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(3/4)}/d/(-c*d^2+e*(-a*e+b*d))/(e*a^{(1/2)}-d*c^{(1/2)})/(-2*a^{(1/2)}*c^{(1/2)}+b)/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A] time = 3.39, antiderivative size = 2112, normalized size of antiderivative = 1.62, number of steps used = 15, number of rules used = 10, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {1720, 1178, 1197, 1103, 1195, 1223, 1714, 1708, 1706, 1216}

result too large to display

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x]

[Out]
$$\begin{aligned} & (x*(a*b*c*(A*e*(2*c*d - b*e) - B*(c*d^2 - a*e^2)) + (b^2 - 2*a*c)*(a*B*e*(2 \\ & *c*d - b*e) + A*(c^2*d^2 + b^2*e^2 - c*e*(2*b*d + a*e))) - c*(a*B*(2*c^2*d^2 \\ & + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d*e - 4*a*c^2*d*e - b^3*e^2 - \\ & b*c*(c*d^2 - 3*a*e^2)))*x^2)/(a*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)^2*S \\ & \text{qrt}[a + b*x^2 + c*x^4]) + (\text{Sqrt}[c]*e^2*(B*d - A*e)*x*\text{Sqrt}[a + b*x^2 + c*x^4 \\ &])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) + (\text{Sqrt}[c]*(a*B* \\ & (2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d*e - 4*a*c^2*d*e - \\ & b^3*e^2 - b*c*(c*d^2 - 3*a*e^2)))*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(a*(b^2 - 4*a* \\ & c)*(c*d^2 - b*d*e + a*e^2)^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (e^{3/2}*(B*d - A*e)*x* \\ & \text{Sqrt}[a + b*x^2 + c*x^4])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*(d + e*x^2)) - (e^{(\\ & 3/2)}*(B*d - A*e)*(3*c*d^2 - e*(2*b*d - a*e))*\text{ArcTan}[(\text{Sqrt}[c*d^2 - b*d*e + a \\ & *e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4])])/(4*d^{(3/2)}*(c*d^2 - b* \\ & d*e + a*e^2)^{(5/2)}) + (e^{(3/2)}*(A*e*(2*c*d - b*e) - B*(c*d^2 - a*e^2))*\text{ArcT} \\ & \text{an}[(\text{Sqrt}[c*d^2 - b*d*e + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4 \\ &])]/(2*\text{Sqrt}[d]*(c*d^2 - b*d*e + a*e^2)^{(5/2)}) - (a^{(1/4)}*c^{(1/4)}*e^2*(B*d - \\ & A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x \\ & ^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4 \\ &])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4]) - (c^{(1/4)}*(a*B* \\ & (2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d*e - 4*a*c^2*d*e - \\ & b^3*e^2 - b*c*(c*d^2 - 3*a*e^2)))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + \\ & c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], \\ & (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(a^{(3/4)}*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^ \\ & 2)^2*\text{Sqrt}[a + b*x^2 + c*x^4]) - (c^{(1/4)}*e*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x \\ & ^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan} \\ & [(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^{(1/4)}*d*(\text{Sqrt}[c]*d \\ & - \text{Sqrt}[a]*e)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) + (c^{(1/4)}*e \\ & *(A*e*(2*c*d - b*e) - B*(c*d^2 - a*e^2))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + \\ & b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(\\ & 1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^{(1/4)}*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c \\ & *d^2 - b*d*e + a*e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4]) - (c^{(1/4)}*(a^{(3/2)}*B*\text{Sqrt} \\ & [c]*e^2 + A*(c*d - b*e)^2 + a*e*(2*B*c*d - b*B*e - A*c*e) - \text{Sqrt}[a]*\text{Sqrt}[c] \\ & *(B*c*d^2 - A*e*(2*c*d - b*e)))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c \\ & *x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (\\ & 2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^{(3/4)}*(b - 2*\text{Sqrt}[a]*\text{Sqrt}[c])*(c*d^2 - b* \\ & d*e + a*e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4]) + (e*(\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(B*d - \\ & A*e)*(3*c*d^2 - e*(2*b*d - a*e))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + \\ & c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(4 \\ & *\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[\\ & c]))/4])/(8*a^{(1/4)}*c^{(1/4)}*d^2*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 - b*d*e + a* \\ & e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4]) - (a^{(3/4)}*e*((\text{Sqrt}[c]*d)/\text{Sqrt}[a] + e)^2*(A \\ & *e*(2*c*d - b*e) - B*(c*d^2 - a*e^2))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x \\ & ^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^ \\ & \end{aligned}$$

$2/(4*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4)/(4*c^{(1/4)}*d*(c*d^2 - a*e^2)*(c*d^2 - b*d*e + a*e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4])$

Rule 1103

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[\frac{(1 + q^2*x^2)*\text{Sqrt}[a + b*x^2 + c*x^4]}{a*(1 + q^2*x^2)^2}] * \text{EllipticF}[2*\text{ArcTan}[q*x], 1/2 - (b*q^2)/(4*c)] / (2*q*\text{Sqrt}[a + b*x^2 + c*x^4]), x] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{PosQ}[c/a]$

Rule 1178

$\text{Int}[\frac{(d_) + (e_)*(x_)^2}{(a_) + (b_)*(x_)^2 + (c_)*(x_)^4}^{(p_)}, x_Symbol] := \text{Simp}[\frac{x*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2}{(2*a*(p + 1)*(b^2 - 4*a*c))}, x] + \text{Dist}[1/(2*a*(p + 1)*(b^2 - 4*a*c)), \text{Int}[\text{Simp}[(2*p + 3)*d*b^2 - a*b*e - 2*a*c*d*(4*p + 5) + (4*p + 7)*(d*b - 2*a*e)*c*x^2, x]*(a + b*x^2 + c*x^4)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&\& \text{LtQ}[p, -1] \&\& \text{IntegerQ}[2*p]$

Rule 1195

$\text{Int}[\frac{(d_) + (e_)*(x_)^2}{\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]}, x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 4]\}, -\text{Simp}[\frac{d*x*\text{Sqrt}[a + b*x^2 + c*x^4]}{a*(1 + q^2*x^2)}, x] + \text{Simp}[\frac{d*(1 + q^2*x^2)*\text{Sqrt}[a + b*x^2 + c*x^4]}{a*(1 + q^2*x^2)^2}] * \text{EllipticE}[2*\text{ArcTan}[q*x], 1/2 - (b*q^2)/(4*c)] / (q*\text{Sqrt}[a + b*x^2 + c*x^4]), x] /; \text{EqQ}[e + d*q^2, 0] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{PosQ}[c/a]$

Rule 1197

$\text{Int}[\frac{(d_) + (e_)*(x_)^2}{\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]}, x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d*q)/q, \text{Int}[1/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] - \text{Dist}[e/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] /; \text{NeQ}[e + d*q, 0] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{PosQ}[c/a]$

Rule 1216

$\text{Int}[1/((d_) + (e_)*(x_)^2)*\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(c*d + a*e*q)/(c*d^2 - a*e^2), \text{Int}[1/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] - \text{Dist}[(a*e*(e + d*q))/(c*d^2 - a*e^2), \text{Int}[(1 + q*x^2)/((d + e*x^2)*\text{Sqrt}[a + b*x^2 + c*x^4]), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&\& \text{NeQ}[c/a, 0]$

$Q[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a]$

Rule 1223

$\text{Int}[\frac{(d_+ + (e_-)(x_-)^2)^{q_+}}{\sqrt{(a_+ + (b_-)(x_-)^2 + (c_-)(x_-)^4)}}, x_Symbol] \rightarrow -\text{Simp}[(e^{2x}(d + ex^2)^{q+1})\sqrt{a + bx^2 + cx^4}]/(2d*(q+1)(c*d^2 - b*d*e + a*e^2)), x] + \text{Dist}[1/(2*d*(q+1)*(c*d^2 - b*d*e + a*e^2)), \text{Int}[(d + ex^2)^{q+1} \text{Simp}[a*e^2*(2*q+3) + 2*d*(c*d - b*e)*(q+1) - 2*e*(c*d*(q+1) - b*e*(q+2))*x^2 + c*e^2*(2*q+5)*x^4, x]]/\sqrt{a + bx^2 + cx^4}, x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{ILtQ}[q, -1]$

Rule 1706

$\text{Int}[\frac{(A_+ + (B_-)(x_-)^2)}{((d_+ + (e_-)(x_-)^2)\sqrt{(a_+ + (b_-)(x_-)^2 + (c_-)(x_-)^4)}), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[B/A, 2]\}, -\text{Simp}[(B*d - A*e)*\text{ArcTan}[\text{Rt}[-b + (c*d)/e + (a*e)/d, 2]*x]/\sqrt{a + bx^2 + cx^4}]/(2*d*e*\text{Rt}[-b + (c*d)/e + (a*e)/d, 2]), x] + \text{Simp}[(B*d + A*e)*(A + B*x^2)*\sqrt{(A^2*(a + bx^2 + cx^4))/(a*(A + B*x^2)^2)}]*\text{EllipticPi}[\text{Cancel}[-(B*d - A*e)^2/(4*d*e*A*B)], 2*\text{ArcTan}[q*x], 1/2 - (b*A)/(4*a*B)]/(4*d*e*A*q*\sqrt{a + bx^2 + cx^4}), x] /; \text{FreeQ}\{a, b, c, d, e, A, B\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a] \&\& \text{EqQ}[c*A^2 - a*B^2, 0]$

Rule 1708

$\text{Int}[\frac{(A_+ + (B_-)(x_-)^2)}{((d_+ + (e_-)(x_-)^2)\sqrt{(a_+ + (b_-)(x_-)^2 + (c_-)(x_-)^4)}), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2), \text{Int}[1/\sqrt{a + bx^2 + cx^4}, x], x] + \text{Dist}[(a*(B*d - A*e)*(e + d*q))/(c*d^2 - a*e^2), \text{Int}[(1 + q*x^2)/((d + ex^2)*\sqrt{a + bx^2 + cx^4}), x], x]] /; \text{FreeQ}\{a, b, c, d, e, A, B\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a] \&\& \text{NeQ}[c*A^2 - a*B^2, 0]$

Rule 1714

$\text{Int}[P4x_+]/((d_+ + (e_-)(x_-)^2)\sqrt{(a_+ + (b_-)(x_-)^2 + (c_-)(x_-)^4)}, x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[c/a, 2], A = \text{Coeff}[P4x, x, 0], B = \text{Coeff}[P4x, x, 2], C = \text{Coeff}[P4x, x, 4]\}, -\text{Dist}[C/(e*q), \text{Int}[(1 - q*x^2)/\sqrt{a + bx^2 + cx^4}, x], x] + \text{Dist}[1/(c*e), \text{Int}[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2]/((d + ex^2)*\sqrt{a + bx^2 + cx^4}), x], x]] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{PolyQ}[P4x, x^2, 2] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[c/a] \&\& !\text{GtQ}[b^2 - 4*a*c, 0]$

Rule 1720

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] :> Int[ExpandIntegrand[1/Sqrt[a + b*x^2 + c*x^4], Px*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] & IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx = \int \left(\frac{aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae)) - c(Ae(2cd - be) - B(c^2d^2 - ae^2))}{(cd^2 - bde + ae^2)^2 (a + bx^2 + cx^4)^{3/2}} \right) dx$$

$$= \frac{\int \frac{aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae)) - c(Ae(2cd - be) - B(c^2d^2 - ae^2))x^2}{(a + bx^2 + cx^4)^{3/2}} dx}{(cd^2 - bde + ae^2)^2} - \frac{(e(Bd - Ae))}{cd^2}$$

$$= \frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 - ae^2)))}{a(b^2 - 2ac)(cd^2 - bde + ae^2)^2} - \frac{(e(Bd - Ae))}{cd^2}$$

$$= \frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 - ae^2)))}{a(b^2 - 2ac)(cd^2 - bde + ae^2)^2} - \frac{(e(Bd - Ae))}{cd^2}$$

$$= \frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 - ae^2)))}{a(b^2 - 2ac)(cd^2 - bde + ae^2)^2} - \frac{(e(Bd - Ae))}{cd^2}$$

$$= \frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 - ae^2)))}{a(b^2 - 2ac)(cd^2 - bde + ae^2)^2} - \frac{(e(Bd - Ae))}{cd^2}$$

Mathematica [C] time = 7.80, size = 8031, normalized size = 6.17

Result too large to show

Warning: Unable to verify antiderivative.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x]

[Out] Result too large to show

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}(ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(e*x^2 + d)^2), x)

maple [B] time = 0.06, size = 8276, normalized size = 6.36

output too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x)

[Out] result too large to display

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}(ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(e*x^2 + d)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(ex^2 + d)^2 (cx^4 + bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x)

[Out] int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Timed out

$$3.30 \quad \int \frac{\sqrt{a} + \sqrt{c}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=273

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} (\sqrt{a}e + \sqrt{c}d) \Pi\left(-\frac{(\sqrt{c}d-\sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \Big|_{\frac{1}{4}}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (\sqrt{c}d - \sqrt{a}e) \tan^{-1}}{4\sqrt[4]{a}\sqrt[4]{c}de\sqrt{a+bx^2+cx^4} \quad 2\sqrt{d}\sqrt{e}\sqrt{ae^2}}$$

[Out] $-1/2*\arctan(x*(a*e^2-b*d*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))*(-e*a^(1/2)+d*c^(1/2))/d^(1/2)/e^(1/2)/(a*e^2-b*d*e+c*d^2)^(1/2)+1/4*(\cos(2*\arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/\cos(2*\arctan(c^(1/4)*x/a^(1/4)))*\text{EllipticPi}(\sin(2*\arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(e*a^(1/2)+d*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(1/4)/c^(1/4)/d/e/(c*x^4+b*x^2+a)^(1/2)$

Rubi [A] time = 0.14, antiderivative size = 273, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 41, $\frac{\text{number of rules}}{\text{integrand size}} = 0.024$, Rules used = {1706}

$$\frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} (\sqrt{a}e + \sqrt{c}d) \Pi\left(-\frac{(\sqrt{c}d-\sqrt{a}e)^2}{4\sqrt{a}\sqrt{c}de}; 2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \Big|_{\frac{1}{4}}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (\sqrt{c}d - \sqrt{a}e) \tan^{-1}}{4\sqrt[4]{a}\sqrt[4]{c}de\sqrt{a+bx^2+cx^4} \quad 2\sqrt{d}\sqrt{e}\sqrt{ae^2}}$$

Antiderivative was successfully verified.

[In] Int[(Sqrt[a] + Sqrt[c]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] $-((\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*\text{ArcTan}[(\text{Sqrt}[c*d^2 - b*d*e + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4])])/(2*\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[c*d^2 - b*d*e + a*e^2]) + ((\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(4*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(4*a^(1/4)*c^(1/4)*d*e*\text{Sqrt}[a + b*x^2 + c*x^4])$

Rule 1706

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] :> With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[Rt[-b + (c*d)/e + (a*e)/d, 2]*x]/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + (c*d)/e + (a*e)/d, 2]), x] + Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a + b*x^2 + c*x^4))/(a*(A + B*x^2)^2)]*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)]], 2*ArcTan[q*x], 1/2 - (b*A)/(4*a*B)]/(4*d*e*A*q*Sqrt[a + b*x^2

+ c*x^4)), x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

Rubi steps

$$\int \frac{\sqrt{a} + \sqrt{c} x^2}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx = -\frac{(\sqrt{c}d - \sqrt{a}e) \tan^{-1}\left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}}\right)}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}} + \frac{(\sqrt{c}d + \sqrt{a}e)(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+b}{\sqrt{a}}}}{4\sqrt{d}}$$

Mathematica [C] time = 0.50, size = 310, normalized size = 1.14

$$\frac{i \sqrt{\frac{\sqrt{b^2 - 4ac} + b + 2cx^2}{\sqrt{b^2 - 4ac} + b}} \sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} + 1 \left((\sqrt{a}e - \sqrt{c}d) \Pi\left(\frac{(b + \sqrt{b^2 - 4ac})e}{2cd}; i \sinh^{-1}\left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x\right) \Big|_{\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}}\right) + \sqrt{c} \right)}{\sqrt{2} de \sqrt{\frac{c}{\sqrt{b^2 - 4ac} + b}} \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(Sqrt[a] + Sqrt[c]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-I)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])*(Sqrt[c]*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + ((-Sqrt[c]*d) + Sqrt[a]*e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]))]/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])*d*e*Sqrt[a + b*x^2 + c*x^4])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm m="giac")

[Out] Timed out

maple [A] time = 0.10, size = 365, normalized size = 1.34

$$\frac{\sqrt{2} \sqrt{-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}} + 4 \sqrt{\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}} + 4 \sqrt{c} \operatorname{EllipticF}\left(\frac{\sqrt{2} \sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} x}{2}, \sqrt{\frac{2(b+\sqrt{-4ac+b^2})b}{ac}-4}\right) (\sqrt{a} e - \dots)}{4 \sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} \sqrt{c x^4 + b x^2 + a} e}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x)

[Out] $\frac{1}{4} c^{1/2} / e^{2^{1/2}} / ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * (-2 * (-b + (-4ac + b^2)^{1/2})^{1/2}) / a * x^2 + 4)^{1/2} * (2 * (b + (-4ac + b^2)^{1/2}) / a * x^2 + 4)^{1/2} / (c * x^4 + b * x^2 + a)^{1/2} * \operatorname{EllipticF}(1/2 * 2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * x, 1/2 * (2 * (b + (-4ac + b^2)^{1/2}) / a * b / c - 4)^{1/2}) + (e * a^{1/2} - d * c^{1/2}) / e / d * 2^{1/2} / (-1/a * b + (-4ac + b^2)^{1/2} / a)^{1/2} * (1/2 / a * b * x^2 - 1/2 * (-4ac + b^2)^{1/2} / a * x^2 + 1)^{1/2} * (1/2 / a * b * x^2 + 1/2 * (-4ac + b^2)^{1/2} / a * x^2 + 1)^{1/2} / (c * x^4 + b * x^2 + a)^{1/2} * \operatorname{EllipticPi}(1/2 * 2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * x, -2 / (-b + (-4ac + b^2)^{1/2}) * a / d * e, (-1/2 * (b + (-4ac + b^2)^{1/2}) / a)^{1/2} * 2^{1/2} / ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2})$

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm m="maxima")

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{a} + \sqrt{c} x^2}{(e x^2 + d) \sqrt{c x^4 + b x^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a^(1/2) + c^(1/2)*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)),x)`

[Out] `int((a^(1/2) + c^(1/2)*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a} + \sqrt{c}x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a**(1/2)+x**2*c**(1/2))/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)`

[Out] `Integral((sqrt(a) + sqrt(c)*x**2)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)`

$$3.31 \quad \int \frac{1 + \sqrt{\frac{c}{a}} x^2}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx$$

Optimal. Leaf size=271

$$\frac{\left(x^2 \sqrt{\frac{c}{a}} + 1\right) \sqrt{\frac{a + bx^2 + cx^4}{a \left(x^2 \sqrt{\frac{c}{a}} + 1\right)^2}} \left(d \sqrt{\frac{c}{a}} + e\right) \Pi\left(-\frac{\left(\sqrt{\frac{c}{a}} d - e\right)^2}{4 \sqrt{\frac{c}{a}} de}; 2 \tan^{-1}\left(\sqrt[4]{\frac{c}{a}} x\right) \Big|_{\frac{1}{4}}\left(2 - \frac{b \sqrt{\frac{c}{a}}}{c}\right)\right)}{4 d e \sqrt[4]{\frac{c}{a}} \sqrt{a + bx^2 + cx^4}} \frac{\left(d \sqrt{\frac{c}{a}} - e\right) \tan^{-1}\left(\frac{x \sqrt{ae^2 - b}}{\sqrt{d} \sqrt{e} \sqrt{a}}\right)}{2 \sqrt{d} \sqrt{e} \sqrt{ae^2 - bde} +}$$

[Out] $-1/2 * \arctan(x * (a * e^2 - b * d * e + c * d^2)^{(1/2)} / d^{(1/2)} / e^{(1/2)} / (c * x^4 + b * x^2 + a)^{(1/2)}) * (-e + d * (c/a)^{(1/2)}) / d^{(1/2)} / e^{(1/2)} / (a * e^2 - b * d * e + c * d^2)^{(1/2)} + 1/4 * (\cos(2 * \arctan((c/a)^{(1/4)} * x))^{(1/2)} / \cos(2 * \arctan((c/a)^{(1/4)} * x))) * \text{EllipticPi}(\sin(2 * \arctan((c/a)^{(1/4)} * x)), -1/4 * (-e + d * (c/a)^{(1/2)})^2 / d / e / (c/a)^{(1/2)}, 1/2 * (2 - b * (c/a)^{(1/2)} / c)^{(1/2)}) * (e + d * (c/a)^{(1/2)}) * (1 + x^2 * (c/a)^{(1/2)}) * ((c * x^4 + b * x^2 + a) / a / (1 + x^2 * (c/a)^{(1/2)})^2)^{(1/2)} / (c/a)^{(1/4)} / d / e / (c * x^4 + b * x^2 + a)^{(1/2)}$

Rubi [A] time = 0.19, antiderivative size = 271, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 41, $\frac{\text{number of rules}}{\text{integrand size}} = 0.024$, Rules used = {1706}

$$\frac{\left(x^2 \sqrt{\frac{c}{a}} + 1\right) \sqrt{\frac{a + bx^2 + cx^4}{a \left(x^2 \sqrt{\frac{c}{a}} + 1\right)^2}} \left(d \sqrt{\frac{c}{a}} + e\right) \Pi\left(-\frac{\left(\sqrt{\frac{c}{a}} d - e\right)^2}{4 \sqrt{\frac{c}{a}} de}; 2 \tan^{-1}\left(\sqrt[4]{\frac{c}{a}} x\right) \Big|_{\frac{1}{4}}\left(2 - \frac{b \sqrt{\frac{c}{a}}}{c}\right)\right)}{4 d e \sqrt[4]{\frac{c}{a}} \sqrt{a + bx^2 + cx^4}} \frac{\left(d \sqrt{\frac{c}{a}} - e\right) \tan^{-1}\left(\frac{x \sqrt{ae^2 - b}}{\sqrt{d} \sqrt{e} \sqrt{a}}\right)}{2 \sqrt{d} \sqrt{e} \sqrt{ae^2 - bde} +}$$

Antiderivative was successfully verified.

[In] Int[(1 + Sqrt[c/a]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x]

[Out] $-((\text{Sqrt}[c/a] * d - e) * \text{ArcTan}[(\text{Sqrt}[c * d^2 - b * d * e + a * e^2] * x) / (\text{Sqrt}[d] * \text{Sqrt}[e] * \text{Sqrt}[a + b * x^2 + c * x^4])]) / (2 * \text{Sqrt}[d] * \text{Sqrt}[e] * \text{Sqrt}[c * d^2 - b * d * e + a * e^2]) + ((\text{Sqrt}[c/a] * d + e) * (1 + \text{Sqrt}[c/a] * x^2) * \text{Sqrt}[(a + b * x^2 + c * x^4) / (a * (1 + \text{Sqrt}[c/a] * x^2)^2)]) * \text{EllipticPi}[-(\text{Sqrt}[c/a] * d - e)^2 / (4 * \text{Sqrt}[c/a] * d * e), 2 * \text{ArcTan}[(c/a)^{(1/4)} * x], (2 - (b * \text{Sqrt}[c/a]) / c) / 4]) / (4 * (c/a)^{(1/4)} * d * e * \text{Sqrt}[a + b * x^2 + c * x^4])$

Rule 1706

Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]), x_Symbol] :> With[{q = Rt[B/A, 2]}, -Simp[((B*d - A*e)*ArcTan[(Rt[-b + (c*d)/e + (a*e)/d, 2]*x)/Sqrt[a + b*x^2 + c*x^4]])/(2*d*e*Rt[-b + (c*d)/e + (a*e)/d, 2]), x] + Simp[((B*d + A*e)*(A + B*x^2)*Sqrt[(A^2*(a

+ b*x^2 + c*x^4))/(a*(A + B*x^2)^2)]*EllipticPi[Cancel[-((B*d - A*e)^2/(4*d*e*A*B))], 2*ArcTan[q*x], 1/2 - (b*A)/(4*a*B)]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4]), x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

Rubi steps

$$\int \frac{1 + \sqrt{\frac{c}{a}} x^2}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx = -\frac{\left(\sqrt{\frac{c}{a}} d - e\right) \tan^{-1}\left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}}\right)}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}} + \frac{\left(\sqrt{\frac{c}{a}} d + e\right) \left(1 + \sqrt{\frac{c}{a}} x^2\right) \sqrt{\frac{a + bx^2 + cx^4}{a\left(1 + \sqrt{\frac{c}{a}} x^2\right)^2}}}{4\sqrt{\frac{c}{a}} d e}$$

Mathematica [C] time = 0.43, size = 312, normalized size = 1.15

$$\frac{i \sqrt{\frac{\sqrt{b^2 - 4ac} + b + 2cx^2}{\sqrt{b^2 - 4ac} + b}} \sqrt{\frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} + 1 \left(\left(e - d \sqrt{\frac{c}{a}} \right) \Pi \left(\frac{(b + \sqrt{b^2 - 4ac})e}{2cd}; i \sinh^{-1} \left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right) \Big|_{\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}} \right) + d \sqrt{\frac{c}{a}} \right)}{\sqrt{2} d e \sqrt{\frac{c}{\sqrt{b^2 - 4ac} + b}} \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(1 + Sqrt[c/a]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-1)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])*(Sqrt[c/a]*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + (-Sqrt[c/a]*d + e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*d*e*Sqrt[a + b*x^2 + c*x^4])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2 \sqrt{\frac{c}{a} + 1}}{\sqrt{cx^4 + bx^2 + a} (ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((x^2*sqrt(c/a) + 1)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)

maple [A] time = 0.08, size = 369, normalized size = 1.36

$$\frac{\sqrt{\frac{c}{a}} \sqrt{2} \sqrt{-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a} + 4} \sqrt{\frac{2(b+\sqrt{-4ac+b^2})x^2}{a} + 4} \operatorname{EllipticF}\left(\frac{\sqrt{2} \sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} x}{2}, \frac{\sqrt{\frac{2(b+\sqrt{-4ac+b^2})b}{ac} - 4}}{2}\right) \left(-\sqrt{\frac{c}{a}}\right)}{4 \sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} \sqrt{cx^4 + bx^2 + a} e}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+x^2*(1/a*c)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x)

[Out] 1/4*(1/a*c)^(1/2)/e*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)*(2*(b+(-4*a*c+b^2)^(1/2))/a*x^2+4)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x, 1/2*(2*(b+(-4*a*c+b^2)^(1/2))/a*b/c-4)^(1/2))+1/e*(-d*(1/a*c)^(1/2)+e)/d*2^(1/2)/(-1/a*b+(-4*a*c+b^2)^(1/2)/a)^(1/2)*(1/2/a*b*x^2-1/2*(-4*a*c+b^2)^(1/2)/a*x^2+1)^(1/2)*(1/2/a*b*x^2+1/2*(-4*a*c+b^2)^(1/2)/a*x^2+1)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticPi(1/2*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*x, -2/(-b+(-4*a*c+b^2)^(1/2))*a/d*e, (-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2 \sqrt{\frac{c}{a} + 1}}{\sqrt{cx^4 + bx^2 + a} (ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((x^2*sqrt(c/a) + 1)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2 \sqrt{\frac{c}{a}} + 1}{(e x^2 + d) \sqrt{c x^4 + b x^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(c/a)^(1/2) + 1)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)

[Out] int((x^2*(c/a)^(1/2) + 1)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2 \sqrt{\frac{c}{a}} + 1}{(d + e x^2) \sqrt{a + b x^2 + c x^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+x**2*(c/a)**(1/2))/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2), x)

[Out] Integral((x**2*sqrt(c/a) + 1)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)

$$3.32 \quad \int \frac{946+315x^2}{(7+5x^2)\sqrt{2+3x^2+x^4}} dx$$

Optimal. Leaf size=106

$$\frac{631(x^2+1)\sqrt{\frac{x^2+2}{x^2+1}}F\left(\tan^{-1}(x)\middle|\frac{1}{2}\right)}{2\sqrt{2}\sqrt{x^4+3x^2+2}} - \frac{2525(x^2+2)\Pi\left(\frac{2}{7};\tan^{-1}(x)\middle|\frac{1}{2}\right)}{14\sqrt{2}\sqrt{\frac{x^2+2}{x^2+1}}\sqrt{x^4+3x^2+2}}$$

[Out] -2525/28*(x^2+2)*(1/(x^2+1))^(1/2)*(x^2+1)^(1/2)*EllipticPi(x/(x^2+1)^(1/2), 2/7, 1/2*2^(1/2))*2^(1/2)/((x^2+2)/(x^2+1))^(1/2)/(x^4+3*x^2+2)^(1/2)+631/4*(x^2+1)^(3/2)*(1/(x^2+1))^(1/2)*EllipticF(x/(x^2+1)^(1/2), 1/2*2^(1/2))*((x^2+2)/(x^2+1))^(1/2)*2^(1/2)/(x^4+3*x^2+2)^(1/2)

Rubi [A] time = 0.11, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 31, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used = {1704, 1099, 1456, 539}

$$\frac{631(x^2+1)\sqrt{\frac{x^2+2}{x^2+1}}F\left(\tan^{-1}(x)\middle|\frac{1}{2}\right)}{2\sqrt{2}\sqrt{x^4+3x^2+2}} - \frac{2525(x^2+2)\Pi\left(\frac{2}{7};\tan^{-1}(x)\middle|\frac{1}{2}\right)}{14\sqrt{2}\sqrt{\frac{x^2+2}{x^2+1}}\sqrt{x^4+3x^2+2}}$$

Antiderivative was successfully verified.

[In] Int[(946 + 315*x^2)/((7 + 5*x^2)*Sqrt[2 + 3*x^2 + x^4]),x]

[Out] (631*(1 + x^2)*Sqrt[(2 + x^2)/(1 + x^2)]*EllipticF[ArcTan[x], 1/2])/(2*Sqrt[2]*Sqrt[2 + 3*x^2 + x^4]) - (2525*(2 + x^2)*EllipticPi[2/7, ArcTan[x], 1/2])/(14*Sqrt[2]*Sqrt[(2 + x^2)/(1 + x^2)]*Sqrt[2 + 3*x^2 + x^4])

Rule 539

Int[Sqrt[(c_) + (d_.)*(x_)^2]/(((a_) + (b_.)*(x_)^2)*Sqrt[(e_) + (f_.)*(x_)^2]), x_Symbol] :> Simp[(c*Sqrt[e + f*x^2]*EllipticPi[1 - (b*c)/(a*d), ArcTan[Rt[d/c, 2]*x], 1 - (c*f)/(d*e)])/(a*e*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[(c*(e + f*x^2))/(e*(c + d*x^2))]), x] /; FreeQ[{a, b, c, d, e, f}, x] && PosQ[d/c]

Rule 1099

Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[((2*a + (b + q)*x^2)*Sqrt[(2*a + (b - q)*x^2)/(2*a + (b + q)*x^2)]*EllipticF[ArcTan[Rt[(b + q)/(2*a), 2]*x], (2*q)/(b + q)])/(2*a*Rt[(b + q)/(2*a), 2]*Sqrt[a + b*x^2 + c*x^4]), x] /; PosQ[(b + q)/a] &&

!(PosQ[(b - q)/a] && SimplerSqrtQ[(b - q)/(2*a), (b + q)/(2*a)])] /; FreeQ[{a, b, c}, x] && GtQ[b^2 - 4*a*c, 0]

Rule 1456

Int[((d_) + (e_)*(x_)^(n_))^(q_)*((f_) + (g_)*(x_)^(n_))^(r_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Dist[(a + b*x^n + c*x^(2*n))^FracPart[p]/((d + e*x^n)^FracPart[p]*(a/d + (c*x^n)/e)^FracPart[p]), Int[(d + e*x^n)^(p + q)*(f + g*x^n)^r*(a/d + (c*x^n)/e)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q, r}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && !IntegerQ[p]

Rule 1704

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Sqrt[b^2 - 4*a*c]}, Dist[(2*a*B - A*(b + q))/(2*a*e - d*(b + q)), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[(B*d - A*e)/(2*a*e - d*(b + q)), Int[(2*a + (b + q)*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x] /; RationalQ[q]] /; FreeQ[{a, b, c, d, e, A, B}, x] && GtQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*A^2 - b*A*B + a*B^2, 0]

Rubi steps

$$\begin{aligned} \int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx &= \frac{631}{2} \int \frac{1}{\sqrt{2 + 3x^2 + x^4}} dx - \frac{2525}{8} \int \frac{4 + 4x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx \\ &= \frac{631(1 + x^2)\sqrt{\frac{2+x^2}{1+x^2}} F\left(\tan^{-1}(x)\middle|\frac{1}{2}\right)}{2\sqrt{2}\sqrt{2 + 3x^2 + x^4}} - \frac{\left(2525\sqrt{\frac{1}{2} + \frac{x^2}{4}}\sqrt{4 + 4x^2}\right) \int \frac{\sqrt{4+4x^2}}{\sqrt{\frac{1}{2} + \frac{x^2}{4}}(7+5x^2)} dx}{8\sqrt{2 + 3x^2 + x^4}} \\ &= \frac{631(1 + x^2)\sqrt{\frac{2+x^2}{1+x^2}} F\left(\tan^{-1}(x)\middle|\frac{1}{2}\right)}{2\sqrt{2}\sqrt{2 + 3x^2 + x^4}} - \frac{2525(2 + x^2)\Pi\left(\frac{2}{7}; \tan^{-1}(x)\middle|\frac{1}{2}\right)}{14\sqrt{2}\sqrt{\frac{2+x^2}{1+x^2}}\sqrt{2 + 3x^2 + x^4}} \end{aligned}$$

Mathematica [C] time = 0.18, size = 74, normalized size = 0.70

$$\frac{i\sqrt{x^2 + 1}\sqrt{x^2 + 2}\left(441F\left(i\sinh^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right) + 505\Pi\left(\frac{10}{7}; i\sinh^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right)\right)}{7\sqrt{x^4 + 3x^2 + 2}}$$

Antiderivative was successfully verified.

[In] Integrate[(946 + 315*x^2)/((7 + 5*x^2)*Sqrt[2 + 3*x^2 + x^4]),x]

[Out] ((-1/7*I)*Sqrt[1 + x^2]*Sqrt[2 + x^2]*(441*EllipticF[I*ArcSinh[x/Sqrt[2]], 2] + 505*EllipticPi[10/7, I*ArcSinh[x/Sqrt[2]], 2])/Sqrt[2 + 3*x^2 + x^4]

fricas [F] time = 1.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{x^4 + 3x^2 + 2}(315x^2 + 946)}{5x^6 + 22x^4 + 31x^2 + 14}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(x^4 + 3*x^2 + 2)*(315*x^2 + 946)/(5*x^6 + 22*x^4 + 31*x^2 + 14), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{315x^2 + 946}{\sqrt{x^4 + 3x^2 + 2}(5x^2 + 7)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate((315*x^2 + 946)/(sqrt(x^4 + 3*x^2 + 2)*(5*x^2 + 7)), x)

maple [C] time = 0.03, size = 93, normalized size = 0.88

$$\frac{63i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\text{EllipticF}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right)}{2\sqrt{x^4+3x^2+2}} - \frac{505i\sqrt{2}\sqrt{\frac{x^2}{2}+1}\sqrt{x^2+1}\text{EllipticPi}\left(\frac{i\sqrt{2}x}{2}, \frac{10}{7}, \sqrt{2}\right)}{7\sqrt{x^4+3x^2+2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x)

[Out] -63/2*I*2^(1/2)*(2*x^2+4)^(1/2)*(x^2+1)^(1/2)/(x^4+3*x^2+2)^(1/2)*EllipticF(1/2*I*2^(1/2)*x, 2^(1/2))-505/7*I*2^(1/2)*(1+1/2*x^2)^(1/2)*(x^2+1)^(1/2)/(x^4+3*x^2+2)^(1/2)*EllipticPi(1/2*I*2^(1/2)*x, 10/7, 2^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{315x^2 + 946}{\sqrt{x^4 + 3x^2 + 2}(5x^2 + 7)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate((315*x^2 + 946)/(sqrt(x^4 + 3*x^2 + 2)*(5*x^2 + 7)), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{315x^2 + 946}{(5x^2 + 7)\sqrt{x^4 + 3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((315*x^2 + 946)/((5*x^2 + 7)*(3*x^2 + x^4 + 2)^(1/2)),x)
```

```
[Out] int((315*x^2 + 946)/((5*x^2 + 7)*(3*x^2 + x^4 + 2)^(1/2)), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{315x^2 + 946}{\sqrt{(x^2 + 1)(x^2 + 2)}(5x^2 + 7)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((315*x**2+946)/(5*x**2+7)/(x**4+3*x**2+2)**(1/2),x)
```

```
[Out] Integral((315*x**2 + 946)/(sqrt((x**2 + 1)*(x**2 + 2))*(5*x**2 + 7)), x)
```


$$3.33 \quad \int \frac{(A+Bx^2)(d+ex^2)^q}{a+bx^2+cx^4} dx$$

Optimal. Leaf size=218

$$\frac{x(d+ex^2)^q \left(\frac{ex^2}{d} + 1\right)^{-q} \left(B - \frac{bB-2Ac}{\sqrt{b^2-4ac}}\right) F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b-\sqrt{b^2-4ac}}, -\frac{ex^2}{d}\right)}{b - \sqrt{b^2 - 4ac}} + \frac{x(d+ex^2)^q \left(\frac{ex^2}{d} + 1\right)^{-q} \left(\frac{bB-2Ac}{\sqrt{b^2-4ac}} + B\right)}{\sqrt{b^2 - 4ac}}$$

[Out] $x*(e*x^2+d)^q*AppellF1(1/2, 1, -q, 3/2, -2*c*x^2/(b-(-4*a*c+b^2)^(1/2)), -e*x^2/d)*(B+(2*A*c-B*b)/(-4*a*c+b^2)^(1/2))/((1+e*x^2/d)^q)/(b-(-4*a*c+b^2)^(1/2))+x*(e*x^2+d)^q*AppellF1(1/2, 1, -q, 3/2, -2*c*x^2/(b+(-4*a*c+b^2)^(1/2)), -e*x^2/d)*(B+(-2*A*c+B*b)/(-4*a*c+b^2)^(1/2))/((1+e*x^2/d)^q)/(b+(-4*a*c+b^2)^(1/2))$

Rubi [A] time = 0.46, antiderivative size = 218, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 31, $\frac{\text{number of rules}}{\text{integrand size}} = 0.097$, Rules used = {1692, 430, 429}

$$\frac{x(d+ex^2)^q \left(\frac{ex^2}{d} + 1\right)^{-q} \left(B - \frac{bB-2Ac}{\sqrt{b^2-4ac}}\right) F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b-\sqrt{b^2-4ac}}, -\frac{ex^2}{d}\right)}{b - \sqrt{b^2 - 4ac}} + \frac{x(d+ex^2)^q \left(\frac{ex^2}{d} + 1\right)^{-q} \left(\frac{bB-2Ac}{\sqrt{b^2-4ac}} + B\right)}{\sqrt{b^2 - 4ac}}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]

[Out] $((B - (b*B - 2*A*c)/\text{Sqrt}[b^2 - 4*a*c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (-2*c*x^2)/(b - \text{Sqrt}[b^2 - 4*a*c]), -((e*x^2)/d)])/((b - \text{Sqrt}[b^2 - 4*a*c])*(1 + (e*x^2)/d)^q) + ((B + (b*B - 2*A*c)/\text{Sqrt}[b^2 - 4*a*c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (-2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c]), -((e*x^2)/d)])/((b + \text{Sqrt}[b^2 - 4*a*c])*(1 + (e*x^2)/d)^q)$

Rule 429

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 :> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, -((b*x^n)/a), -((d*x^n)/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 430

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
 :> Dist[(a^IntPart[p]*(a + b*x^n)^FracPart[p])/(1 + (b*x^n)/a)^FracPart[p], Int[(1 + (b*x^n)/a)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}

, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])

Rule 1692

Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] :> Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && PolyQ[Px, x^2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IntegerQ[p]

Rubi steps

$$\begin{aligned} \int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx &= \int \left(\frac{\left(B + \frac{-bB+2Ac}{\sqrt{b^2-4ac}}\right)(d + ex^2)^q}{b - \sqrt{b^2 - 4ac} + 2cx^2} + \frac{\left(B - \frac{-bB+2Ac}{\sqrt{b^2-4ac}}\right)(d + ex^2)^q}{b + \sqrt{b^2 - 4ac} + 2cx^2} \right) dx \\ &= \left(B - \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right) \int \frac{(d + ex^2)^q}{b - \sqrt{b^2 - 4ac} + 2cx^2} dx + \left(B + \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right) \int \frac{(d + ex^2)^q}{b + \sqrt{b^2 - 4ac} + 2cx^2} dx \\ &= \left(\left(B - \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right)(d + ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q}\right) \int \frac{\left(1 + \frac{ex^2}{d}\right)^q}{b - \sqrt{b^2 - 4ac} + 2cx^2} dx + \left(\left(B + \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right)(d + ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q}\right) \int \frac{\left(1 + \frac{ex^2}{d}\right)^q}{b + \sqrt{b^2 - 4ac} + 2cx^2} dx \\ &= \frac{\left(B - \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right)x(d + ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b - \sqrt{b^2 - 4ac}}, -\frac{ex^2}{d}\right)}{b - \sqrt{b^2 - 4ac}} + \frac{\left(B + \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right)x(d + ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b + \sqrt{b^2 - 4ac}}, -\frac{ex^2}{d}\right)}{b + \sqrt{b^2 - 4ac}} \end{aligned}$$

Mathematica [F] time = 0.22, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx$$

Verification is Not applicable to the result.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]

[Out] Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]

fricas [F] time = 1.70, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="fricas")

[Out] integral((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + b*x^2 + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + b*x^2 + a), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x)

[Out] int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + b*x^2 + a), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**q/(c*x**4+b*x**2+a), x)

[Out] Timed out

$$3.34 \quad \int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx$$

Optimal. Leaf size=106

$$\frac{1}{4}\sqrt{3} \log\left(x^2 - \sqrt{3}\sqrt{x^2+1} + 2\right) - \frac{1}{4}\sqrt{3} \log\left(x^2 + \sqrt{3}\sqrt{x^2+1} + 2\right) - \frac{1}{2} \tan^{-1}\left(\sqrt{3} - 2\sqrt{x^2+1}\right) + \frac{1}{2} \tan^{-1}\left(2\sqrt{x^2+1}\right)$$

[Out] 1/2*arctan(-3^(1/2)+2*(x^2+1)^(1/2))+1/2*arctan(3^(1/2)+2*(x^2+1)^(1/2))+1/4*ln(2+x^2-3^(1/2)*(x^2+1)^(1/2))*3^(1/2)-1/4*ln(2+x^2+3^(1/2)*(x^2+1)^(1/2))*3^(1/2)

Rubi [A] time = 0.13, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 7, integrand size = 28, number of rules / integrand size = 0.250, Rules used = {1685, 826, 1169, 634, 618, 204, 628}

$$\frac{1}{4}\sqrt{3} \log\left(x^2 - \sqrt{3}\sqrt{x^2+1} + 2\right) - \frac{1}{4}\sqrt{3} \log\left(x^2 + \sqrt{3}\sqrt{x^2+1} + 2\right) - \frac{1}{2} \tan^{-1}\left(\sqrt{3} - 2\sqrt{x^2+1}\right) + \frac{1}{2} \tan^{-1}\left(2\sqrt{x^2+1}\right)$$

Antiderivative was successfully verified.

[In] Int[(x*(1 + 2*x^2))/(Sqrt[1 + x^2]*(1 + x^2 + x^4)),x]

[Out] -ArcTan[Sqrt[3] - 2*Sqrt[1 + x^2]]/2 + ArcTan[Sqrt[3] + 2*Sqrt[1 + x^2]]/2 + (Sqrt[3]*Log[2 + x^2 - Sqrt[3]*Sqrt[1 + x^2]])/4 - (Sqrt[3]*Log[2 + x^2 + Sqrt[3]*Sqrt[1 + x^2]])/4

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 618

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 628

Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 634

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] :> Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 826

```
Int[((f_.) + (g_.)*(x_))/(Sqrt[(d_.) + (e_.)*(x_)]*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)), x_Symbol] :> Dist[2, Subst[Int[(e*f - d*g + g*x^2)/(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4), x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0]
```

Rule 1169

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] :> With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]
```

Rule 1685

```
Int[(Px_)*(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] :> Dist[1/2, Subst[Int[(Px /. x -> Sqrt[x])*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && PolyQ[Px, x^2]
```

Rubi steps

$$\begin{aligned}
\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx &= \frac{1}{2} \text{Subst} \left(\int \frac{1+2x}{\sqrt{1+x}(1+x+x^2)} dx, x, x^2 \right) \\
&= \text{Subst} \left(\int \frac{-1+2x^2}{1-x^2+x^4} dx, x, \sqrt{1+x^2} \right) \\
&= \frac{\text{Subst} \left(\int \frac{-\sqrt{3}+3x}{1-\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2} \right)}{2\sqrt{3}} + \frac{\text{Subst} \left(\int \frac{-\sqrt{3}-3x}{1+\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2} \right)}{2\sqrt{3}} \\
&= \frac{1}{4} \text{Subst} \left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2} \right) + \frac{1}{4} \text{Subst} \left(\int \frac{1}{1+\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2} \right) \\
&= \frac{1}{4} \sqrt{3} \log \left(2+x^2 - \sqrt{3} \sqrt{1+x^2} \right) - \frac{1}{4} \sqrt{3} \log \left(2+x^2 + \sqrt{3} \sqrt{1+x^2} \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \sqrt{1+x^2} \right) \\
&= -\frac{1}{2} \tan^{-1} \left(\sqrt{3} - 2\sqrt{1+x^2} \right) + \frac{1}{2} \tan^{-1} \left(\sqrt{3} + 2\sqrt{1+x^2} \right) + \frac{1}{4} \sqrt{3} \log \left(2+x^2 - \sqrt{3} \sqrt{1+x^2} \right) - \frac{1}{4} \sqrt{3} \log \left(2+x^2 + \sqrt{3} \sqrt{1+x^2} \right)
\end{aligned}$$

Mathematica [C] time = 0.20, size = 125, normalized size = 1.18

$$\frac{\sqrt{2(1-i\sqrt{3})} \tanh^{-1} \left(\frac{\sqrt{2}\sqrt{x^2+1}}{\sqrt{1-i\sqrt{3}}} \right)}{-1+i\sqrt{3}} + \frac{\sqrt{2(1+i\sqrt{3})} \tanh^{-1} \left(\frac{\sqrt{2}\sqrt{x^2+1}}{\sqrt{1+i\sqrt{3}}} \right)}{-1-i\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Integrate[(x*(1+2*x^2))/(Sqrt[1+x^2]*(1+x^2+x^4)),x]

[Out] (Sqrt[2*(1-I*Sqrt[3])]*ArcTanh[(Sqrt[2]*Sqrt[1+x^2])/Sqrt[1-I*Sqrt[3]]])/(-1+I*Sqrt[3]) + (Sqrt[2*(1+I*Sqrt[3])]*ArcTanh[(Sqrt[2]*Sqrt[1+x^2])/Sqrt[1+I*Sqrt[3]]])/(-1-I*Sqrt[3])

fricas [B] time = 1.63, size = 279, normalized size = 2.63

$$\frac{1}{4} \sqrt{3} \log \left(32x^4 + 80x^2 + 32\sqrt{3}(x^3+x) - 16(2x^3 + \sqrt{3}(2x^2+1) + 4x)\sqrt{x^2+1} + 32 \right) - \frac{1}{4} \sqrt{3} \log \left(32x^4 + 80x^2 + 32\sqrt{3}(x^3+x) + 16(2x^3 + \sqrt{3}(2x^2+1) + 4x)\sqrt{x^2+1} + 32 \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] 1/4*sqrt(3)*log(32*x^4 + 80*x^2 + 32*sqrt(3)*(x^3 + x) - 16*(2*x^3 + sqrt(3)*(2*x^2 + 1) + 4*x)*sqrt(x^2 + 1) + 32) - 1/4*sqrt(3)*log(32*x^4 + 80*x^2 + 32*sqrt(3)*(x^3 + x) + 16*(2*x^3 + sqrt(3)*(2*x^2 + 1) + 4*x)*sqrt(x^2 + 1) + 32)

1) + 32) - arctan(2*sqrt(2*x^4 + 5*x^2 + 2*sqrt(3)*(x^3 + x) - (2*x^3 + sqrt(3)*(2*x^2 + 1) + 4*x)*sqrt(x^2 + 1) + 2)*(x + sqrt(x^2 + 1)) + sqrt(3) - 2*sqrt(x^2 + 1)) - arctan(2*sqrt(2*x^4 + 5*x^2 - 2*sqrt(3)*(x^3 + x) - (2*x^3 - sqrt(3)*(2*x^2 + 1) + 4*x)*sqrt(x^2 + 1) + 2)*(x + sqrt(x^2 + 1)) - sqrt(3) - 2*sqrt(x^2 + 1))

giac [A] time = 0.65, size = 80, normalized size = 0.75

$$-\frac{1}{4}\sqrt{3}\log\left(x^2 + \sqrt{3}\sqrt{x^2+1} + 2\right) + \frac{1}{4}\sqrt{3}\log\left(x^2 - \sqrt{3}\sqrt{x^2+1} + 2\right) + \frac{1}{2}\arctan\left(\sqrt{3} + 2\sqrt{x^2+1}\right) + \frac{1}{2}\arctan\left(-\sqrt{3} + 2\sqrt{x^2+1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] -1/4*sqrt(3)*log(x^2 + sqrt(3)*sqrt(x^2 + 1) + 2) + 1/4*sqrt(3)*log(x^2 - sqrt(3)*sqrt(x^2 + 1) + 2) + 1/2*arctan(sqrt(3) + 2*sqrt(x^2 + 1)) + 1/2*arctan(-sqrt(3) + 2*sqrt(x^2 + 1))

maple [B] time = 0.03, size = 296, normalized size = 2.79

$$\frac{\sqrt{2}\sqrt{\frac{2(x+1)^2}{(-x+1)^2}+2}\left(\sqrt{3}\operatorname{arctanh}\left(\frac{\sqrt{\frac{2(x+1)^2}{(-x+1)^2}+2}\sqrt{3}}{2}\right)+\operatorname{arctan}\left(\frac{\sqrt{\frac{2(x+1)^2}{(-x+1)^2}+2}(x+1)}{\left(\frac{(x+1)^2}{(-x+1)^2}+1\right)(-x+1)}\right)\right)}{4\sqrt{\frac{\frac{(x+1)^2}{(-x+1)^2}+1}{\left(\frac{x+1}{-x+1}+1\right)^2}}}\frac{\sqrt{2}\sqrt{\frac{2(x-1)^2}{(-x-1)^2}+2}\left(\sqrt{3}\operatorname{arctanh}\left(\frac{\sqrt{\frac{2(x-1)^2}{(-x-1)^2}+2}\sqrt{3}}{2}\right)+\operatorname{arctan}\left(\frac{\sqrt{\frac{2(x-1)^2}{(-x-1)^2}+2}(x-1)}{\left(\frac{(x-1)^2}{(-x-1)^2}+1\right)(-x-1)}\right)\right)}{4\sqrt{\frac{\frac{(x-1)^2}{(-x-1)^2}+1}{\left(\frac{x-1}{-x-1}+1\right)^2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x)

[Out] -1/4*2^(1/2)*(2*(x+1)^2/(1-x)^2+2)^(1/2)*(3^(1/2)*arctanh(1/2*(2*(x+1)^2/(1-x)^2+2)^(1/2)*3^(1/2))+arctan(1/((x+1)^2/(1-x)^2+1)*(2*(x+1)^2/(1-x)^2+2)^(1/2)*(x+1)/(1-x)))/(((x+1)^2/(1-x)^2+1)/((x+1)/(1-x)+1)^2)^(1/2)/((x+1)/(1-x)+1)-1/4*2^(1/2)*(2*(x-1)^2/(-1-x)^2+2)^(1/2)*(3^(1/2)*arctanh(1/2*(2*(x-1)^2/(-1-x)^2+2)^(1/2)*3^(1/2))+arctan(1/((x-1)^2/(-1-x)^2+1)*(2*(x-1)^2/(-1-x)^2+2)^(1/2)*(x-1)/(-1-x)))/(((x-1)^2/(-1-x)^2+1)/((x-1)/(-1-x)+1)^2)^(1/2)/((x-1)/(-1-x)+1)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(2x^2 + 1)x}{(x^4 + x^2 + 1)\sqrt{x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate((2*x^2 + 1)*x/((x^4 + x^2 + 1)*sqrt(x^2 + 1)), x)

mupad [B] time = 1.85, size = 397, normalized size = 3.75

$$\frac{\left(\ln\left(x - \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right) - \ln\left(\frac{x}{2} + \left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right)\sqrt{x^2+1} + 1 + \frac{\sqrt{3}x1i}{2}\right)\right)\left(2\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 + \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)\left(\ln\left(x + \frac{1}{2} - \frac{\sqrt{3}}{2}\right)\right)}{\sqrt{\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^2 + 1}\left(4\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 + 1 + \sqrt{3}1i\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*(2*x^2 + 1))/((x^2 + 1)^(1/2)*(x^2 + x^4 + 1)),x)

[Out] ((log(x - (3^(1/2)*1i)/2 - 1/2) - log(x/2 + (3^(1/2)/2 + 1i/2)*(x^2 + 1)^(1/2) + (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 + 1/2)^3 + 1/2))/((((3^(1/2)*1i)/2 + 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 + 1/2)^3 + 1)) + ((log(x - (3^(1/2)*1i)/2 + 1/2) - log((3^(1/2)/2 - 1i/2)*(x^2 + 1)^(1/2) - x/2 + (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 - 1/2)^3 - 1/2))/((((3^(1/2)*1i)/2 - 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 - 1/2)^3 - 1)) + ((log(x + (3^(1/2)*1i)/2 - 1/2) - log(x/2 + (3^(1/2)/2 - 1i/2)*(x^2 + 1)^(1/2) - (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 - 1/2)^3 - 1/2))/((((3^(1/2)*1i)/2 - 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 - 1/2)^3 - 1)) + ((log(x + (3^(1/2)*1i)/2 + 1/2) - log((3^(1/2)/2 + 1i/2)*(x^2 + 1)^(1/2) - x/2 - (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 + 1/2)^3 + 1/2))/((((3^(1/2)*1i)/2 + 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 + 1/2)^3 + 1))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x(2x^2+1)}{\sqrt{x^2+1}(x^2-x+1)(x^2+x+1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(2*x**2+1)/(x**4+x**2+1)/(x**2+1)**(1/2),x)

[Out] Integral(x*(2*x**2 + 1)/(sqrt(x**2 + 1)*(x**2 - x + 1)*(x**2 + x + 1)), x)

$$3.35 \quad \int \frac{\sqrt{a+bx^2+cx^4}}{ad-cdx^4} dx$$

Optimal. Leaf size=145

$$\frac{\sqrt{2\sqrt{a}\sqrt{c}+b} \tanh^{-1}\left(\frac{x\sqrt{2\sqrt{a}\sqrt{c}+b}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} - \frac{\sqrt{b-2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{x\sqrt{b-2\sqrt{a}\sqrt{c}}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}d}$$

[Out] $-1/4*\arctanh(x*(-2*a^(1/2)*c^(1/2)+b)^(1/2)/(c*x^4+b*x^2+a)^(1/2))*(-2*a^(1/2)*c^(1/2)+b)^(1/2)/d/a^(1/2)/c^(1/2)+1/4*\arctanh(x*(b+2*a^(1/2)*c^(1/2))^(1/2)/(c*x^4+b*x^2+a)^(1/2))*(b+2*a^(1/2)*c^(1/2))^(1/2)/d/a^(1/2)/c^(1/2)$

Rubi [A] time = 0.21, antiderivative size = 145, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2071, 1093, 208}

$$\frac{\sqrt{2\sqrt{a}\sqrt{c}+b} \tanh^{-1}\left(\frac{x\sqrt{2\sqrt{a}\sqrt{c}+b}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} - \frac{\sqrt{b-2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{x\sqrt{b-2\sqrt{a}\sqrt{c}}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}d}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2 + c*x^4]/(a*d - c*d*x^4), x]

[Out] $-(\text{Sqrt}[b - 2*\text{Sqrt}[a]*\text{Sqrt}[c]]*\text{ArcTanh}[(\text{Sqrt}[b - 2*\text{Sqrt}[a]*\text{Sqrt}[c]]*x)/\text{Sqrt}[a + b*x^2 + c*x^4]])/(4*\text{Sqrt}[a]*\text{Sqrt}[c]*d) + (\text{Sqrt}[b + 2*\text{Sqrt}[a]*\text{Sqrt}[c]]*\text{ArcTanh}[(\text{Sqrt}[b + 2*\text{Sqrt}[a]*\text{Sqrt}[c]]*x)/\text{Sqrt}[a + b*x^2 + c*x^4]])/(4*\text{Sqrt}[a]*\text{Sqrt}[c]*d)$

Rule 208

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 1093

Int[((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[c/q, Int[1/(b/2 - q/2 + c*x^2), x], x] - Dist[c/q, Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[b^2 - 4*a*c]

Rule 2071

```
Int[Sqrt[v_]/((d_) + (e_.)*(x_)^4), x_Symbol] := With[{a = Coeff[v, x, 0],
b = Coeff[v, x, 2], c = Coeff[v, x, 4]}, Dist[a/d, Subst[Int[1/(1 - 2*b*x^2
+ (b^2 - 4*a*c)*x^4), x], x, x/Sqrt[v]], x] /; EqQ[c*d + a*e, 0] && PosQ[a
*c]] /; FreeQ[{d, e}, x] && PolyQ[v, x^2, 2]
```

Rubi steps

$$\int \frac{\sqrt{a + bx^2 + cx^4}}{ad - cdx^4} dx = \frac{\text{Subst}\left(\int \frac{1}{1-2bx^2+(b^2-4ac)x^4} dx, x, \frac{x}{\sqrt{a+bx^2+cx^4}}\right)}{d}$$

$$= \frac{(b^2 - 4ac) \text{Subst}\left(\int \frac{1}{-b-2\sqrt{a}\sqrt{c}+(b^2-4ac)x^2} dx, x, \frac{x}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} - \frac{(b^2 - 4ac) \text{Subst}\left(\int \frac{1}{-b+2\sqrt{a}\sqrt{c}+(b^2-4ac)x^2} dx, x, \frac{x}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}d}$$

$$= -\frac{\sqrt{b-2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{\sqrt{b-2\sqrt{a}\sqrt{c}}x}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} + \frac{\sqrt{b+2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{\sqrt{b+2\sqrt{a}\sqrt{c}}x}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}d}$$

Mathematica [C] time = 0.61, size = 441, normalized size = 3.04

$$i\sqrt{\frac{\sqrt{b^2-4ac}+b+2cx^2}{\sqrt{b^2-4ac}+b}} \sqrt{\frac{2cx^2}{b-\sqrt{b^2-4ac}}} + 1 \left(2\sqrt{a}\sqrt{c} F\left(i \sinh^{-1}\left(\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}x\right) \middle| \frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right) - (2\sqrt{a}\sqrt{c} + b) \Pi\left(\frac{-b-\sqrt{b^2-4ac}}{2\sqrt{a}\sqrt{c}} \middle| \frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right) \right) / (2\sqrt{2}\sqrt{a}\sqrt{c}d\sqrt{\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}})$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2 + c*x^4]/(a*d - c*d*x^4), x]

[Out] ((I/2)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])*(2*Sqrt[a]*Sqrt[c]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) - (b + 2*Sqrt[a]*Sqrt[c])*EllipticPi[(-b - Sqrt[b^2 - 4*a*c])/(2*Sqrt[a]*Sqrt[c]), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + (b - 2*Sqrt[a]*Sqrt[c])*EllipticPi[(b + Sqrt[b^2 - 4*a*c])/(2*Sqrt[a]*Sqrt[c]), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))/(Sqrt[2]*Sqrt[a]*Sqrt[c]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*d*Sqrt[a + b*x^2 + c*x^4])

fricas [B] time = 6.80, size = 603, normalized size = 4.16

$$\frac{1}{8} \sqrt{\frac{2acd^2\sqrt{\frac{1}{acd^4}} + b}{acd^2}} \log \left(\frac{\sqrt{cx^4 + bx^2 + a} \left(ad^2\sqrt{\frac{1}{acd^4}} + x^2 \right) + \left(acd^3x^3\sqrt{\frac{1}{acd^4}} + adx \right) \sqrt{\frac{2acd^2\sqrt{\frac{1}{acd^4}} + b}{acd^2}}}{cx^4 - a} \right) - \frac{1}{8} \sqrt{\frac{2acd^2\sqrt{\frac{1}{acd^4}} + b}{acd^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d),x, algorithm="fricas")

[Out] 1/8*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2))*log((sqrt(c*x^4 + b*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) + x^2) + (a*c*d^3*x^3*sqrt(1/(a*c*d^4)) + a*d*x)*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2)))/(c*x^4 - a)) - 1/8*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2))*log((sqrt(c*x^4 + b*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) + x^2) - (a*c*d^3*x^3*sqrt(1/(a*c*d^4)) + a*d*x)*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2)))/(c*x^4 - a)) + 1/8*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2))*log(-(sqrt(c*x^4 + b*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) - x^2) + (a*c*d^3*x^3*sqrt(1/(a*c*d^4)) - a*d*x)*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2)))/(c*x^4 - a)) - 1/8*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2))*log(-(sqrt(c*x^4 + b*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) - x^2) - (a*c*d^3*x^3*sqrt(1/(a*c*d^4)) - a*d*x)*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2)))/(c*x^4 - a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int -\frac{\sqrt{cx^4 + bx^2 + a}}{cdx^4 - ad} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d),x, algorithm="giac")

[Out] integrate(-sqrt(c*x^4 + b*x^2 + a)/(c*d*x^4 - a*d), x)

maple [B] time = 0.04, size = 238, normalized size = 1.64

$$\frac{\sqrt{2} b \arctan\left(\frac{\sqrt{cx^4+bx^2+a} \sqrt{2}}{\sqrt{-2b-4\sqrt{ac}} x}\right)}{4\sqrt{ac} \sqrt{-2b-4\sqrt{ac}} d} + \frac{\sqrt{2} b \arctan\left(\frac{\sqrt{cx^4+bx^2+a} \sqrt{2}}{\sqrt{-2b+4\sqrt{ac}} x}\right)}{4\sqrt{ac} \sqrt{-2b+4\sqrt{ac}} d} - \frac{\sqrt{2} \arctan\left(\frac{\sqrt{cx^4+bx^2+a} \sqrt{2}}{\sqrt{-2b-4\sqrt{ac}} x}\right)}{2\sqrt{-2b-4\sqrt{ac}} d} - \frac{\sqrt{2} \arctan\left(\frac{\sqrt{cx^4+bx^2+a} \sqrt{2}}{\sqrt{-2b+4\sqrt{ac}} x}\right)}{2\sqrt{-2b+4\sqrt{ac}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d), x)`

[Out]
$$-1/2/d*2^{(1/2)}/(-4*(a*c)^{(1/2)}-2*b)^{(1/2)}*\arctan((c*x^4+b*x^2+a)^{(1/2)}*2^{(1/2)}/x/(-4*(a*c)^{(1/2)}-2*b)^{(1/2)})-1/4/d*2^{(1/2)}/(a*c)^{(1/2)}/(-4*(a*c)^{(1/2)}-2*b)^{(1/2)}*\arctan((c*x^4+b*x^2+a)^{(1/2)}*2^{(1/2)}/x/(-4*(a*c)^{(1/2)}-2*b)^{(1/2)})+1/4/d*2^{(1/2)}/(a*c)^{(1/2)}/(4*(a*c)^{(1/2)}-2*b)^{(1/2)}*\arctan((c*x^4+b*x^2+a)^{(1/2)}*2^{(1/2)}/x/(4*(a*c)^{(1/2)}-2*b)^{(1/2)})+1/4/d*2^{(1/2)}/(a*c)^{(1/2)}/(4*(a*c)^{(1/2)}-2*b)^{(1/2)}*\arctan((c*x^4+b*x^2+a)^{(1/2)}*2^{(1/2)}/x/(4*(a*c)^{(1/2)}-2*b)^{(1/2)})*b$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{\sqrt{cx^4 + bx^2 + a}}{cdx^4 - ad} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d), x, algorithm="maxima")`

[Out] `-integrate(sqrt(c*x^4 + b*x^2 + a)/(c*d*x^4 - a*d), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{cx^4 + bx^2 + a}}{ad - cdx^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2 + c*x^4)^(1/2)/(a*d - c*d*x^4), x)`

[Out] `int((a + b*x^2 + c*x^4)^(1/2)/(a*d - c*d*x^4), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$-\frac{\int \frac{\sqrt{a+bx^2+cx^4}}{-a+cx^4} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*x**4+b*x**2+a)**(1/2)/(-c*d*x**4+a*d), x)`

[Out] `-Integral(sqrt(a + b*x**2 + c*x**4)/(-a + c*x**4), x)/d`

$$3.36 \quad \int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx$$

Optimal. Leaf size=239

$$\frac{\sqrt{\sqrt{4ac+b^2}-b} \tanh^{-1}\left(\frac{x\sqrt{\sqrt{4ac+b^2}-b}(\sqrt{4ac+b^2}+b-2cx^2)}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a+bx^2-cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{c}d} - \frac{\sqrt{\sqrt{4ac+b^2}+b} \tan^{-1}\left(\frac{x\sqrt{\sqrt{4ac+b^2}+b}(-\sqrt{4ac+b^2}+b-2cx^2)}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a+bx^2-cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{c}d}$$

[Out] $\frac{1}{4} \arctanh\left(\frac{1}{4} x (-2cx^2 + (4ac + b^2)^{1/2} + b) (-b + (4ac + b^2)^{1/2})^{1/2}\right) \frac{2^{1/2}}{a^{1/2} c^{1/2}} \frac{1}{(-cx^4 + bx^2 + a)^{1/2}} (-b + (4ac + b^2)^{1/2})^{1/2} \frac{1}{d} \frac{2^{1/2}}{a^{1/2} c^{1/2}} - \frac{1}{4} \arctan\left(\frac{1}{4} x (-2cx^2 - (4ac + b^2)^{1/2} + b) (b + (4ac + b^2)^{1/2})^{1/2}\right) \frac{2^{1/2}}{a^{1/2} c^{1/2}} \frac{1}{(-cx^4 + bx^2 + a)^{1/2}} (b + (4ac + b^2)^{1/2})^{1/2} \frac{1}{d} \frac{2^{1/2}}{a^{1/2} c^{1/2}}$

Rubi [A] time = 0.18, antiderivative size = 239, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 0.033$, Rules used = {2072}

$$\frac{\sqrt{\sqrt{4ac+b^2}-b} \tanh^{-1}\left(\frac{x\sqrt{\sqrt{4ac+b^2}-b}(\sqrt{4ac+b^2}+b-2cx^2)}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a+bx^2-cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{c}d} - \frac{\sqrt{\sqrt{4ac+b^2}+b} \tan^{-1}\left(\frac{x\sqrt{\sqrt{4ac+b^2}+b}(-\sqrt{4ac+b^2}+b-2cx^2)}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a+bx^2-cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{c}d}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a + b*x^2 - c*x^4]/(a*d + c*d*x^4), x]

[Out] $-\frac{(\text{Sqrt}[b + \text{Sqrt}[b^2 + 4ac]] \text{ArcTan}[(\text{Sqrt}[b + \text{Sqrt}[b^2 + 4ac]]) x (b - \text{Sqrt}[b^2 + 4ac] - 2cx^2)) / (2\sqrt{2} \text{Sqrt}[a] \text{Sqrt}[c] \text{Sqrt}[a + bx^2 - cx^4])}{(2\sqrt{2} \text{Sqrt}[2] \text{Sqrt}[a] \text{Sqrt}[c] d)} + \frac{(\text{Sqrt}[-b + \text{Sqrt}[b^2 + 4ac]] \text{ArcTan}[(\text{Sqrt}[-b + \text{Sqrt}[b^2 + 4ac]]) x (b + \text{Sqrt}[b^2 + 4ac] - 2cx^2)) / (2\sqrt{2} \text{Sqrt}[2] \text{Sqrt}[a] \text{Sqrt}[c] \text{Sqrt}[a + bx^2 - cx^4])}{(2\sqrt{2} \text{Sqrt}[2] \text{Sqrt}[a] \text{Sqrt}[c] d)}$

Rule 2072

Int[Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]/((d_) + (e_.)*(x_)^4), x_Symbol] := With[{q = Sqrt[b^2 - 4ac]}, -Simp[(a*Sqrt[b + q]*ArcTan[(Sqrt[b + q])*x*(b - q + 2cx^2)]/(2*Sqrt[2]*Rt[-(a*c), 2]*Sqrt[a + bx^2 + cx^4])]/(2*Sqrt[2]*Rt[-(a*c), 2]*d), x] + Simp[(a*Sqrt[-b + q]*ArcTanh[(Sqrt[-b + q])*x*(b + q + 2cx^2)]/(2*Sqrt[2]*Rt[-(a*c), 2]*Sqrt[a + bx^2 + cx^4])]/(2*Sqrt[2]*Rt[-(a*c), 2]*d), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c*d + a*e, 0] && NegQ[a*c]

Rubi steps

$$\int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx = -\frac{\sqrt{b+\sqrt{b^2+4ac}} \tan^{-1}\left(\frac{\sqrt{b+\sqrt{b^2+4ac}} x(b-\sqrt{b^2+4ac}-2cx^2)}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a+bx^2-cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{c}d} + \frac{\sqrt{-b+\sqrt{b^2+4ac}} \tanh^{-1}\left(\frac{\sqrt{-b+\sqrt{b^2+4ac}} x(b-\sqrt{b^2+4ac}-2cx^2)}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a+bx^2-cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{c}d}$$

Mathematica [C] time = 0.68, size = 432, normalized size = 1.81

$$\frac{\sqrt{\frac{4cx^2}{\sqrt{4ac+b^2}-b} + 2\sqrt{1-\frac{2cx^2}{\sqrt{4ac+b^2}+b}} \left(2i\sqrt{a}\sqrt{c} F\left(i\sinh^{-1}\left(\sqrt{2}\sqrt{-\frac{c}{b+\sqrt{b^2+4ac}}}\right)x\right)\right)^{\frac{b+\sqrt{b^2+4ac}}{b-\sqrt{b^2+4ac}}} + (b-2i\sqrt{a}\sqrt{c})\Pi\left(-i\sqrt{\frac{b+\sqrt{b^2+4ac}}{b-\sqrt{b^2+4ac}}}\right)}{4\sqrt{a}\sqrt{c}d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2 - c*x^4]/(a*d + c*d*x^4), x]

[Out] (Sqrt[2 + (4*c*x^2)/(-b + Sqrt[b^2 + 4*a*c])]*Sqrt[1 - (2*c*x^2)/(b + Sqrt[b^2 + 4*a*c])])*((2*I)*Sqrt[a]*Sqrt[c]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[-(c/(b + Sqrt[b^2 + 4*a*c])])]*x], (b + Sqrt[b^2 + 4*a*c])/(b - Sqrt[b^2 + 4*a*c])) + (b - (2*I)*Sqrt[a]*Sqrt[c])*EllipticPi[(-1/2*I)*(b + Sqrt[b^2 + 4*a*c])]/(Sqrt[a]*Sqrt[c]), I*ArcSinh[Sqrt[2]*Sqrt[-(c/(b + Sqrt[b^2 + 4*a*c])])]*x], (b + Sqrt[b^2 + 4*a*c])/(b - Sqrt[b^2 + 4*a*c]) - (b + (2*I)*Sqrt[a]*Sqrt[c])*EllipticPi[((I/2)*(b + Sqrt[b^2 + 4*a*c])]/(Sqrt[a]*Sqrt[c]), I*ArcSinh[Sqrt[2]*Sqrt[-(c/(b + Sqrt[b^2 + 4*a*c])])]*x], (b + Sqrt[b^2 + 4*a*c])/(b - Sqrt[b^2 + 4*a*c]))]/(4*Sqrt[a]*Sqrt[c]*Sqrt[-(c/(b + Sqrt[b^2 + 4*a*c]))])*d*Sqrt[a + b*x^2 - c*x^4])

fricas [B] time = 6.82, size = 669, normalized size = 2.80

$$-\frac{1}{8}\sqrt{\frac{2acd^2\sqrt{-\frac{1}{acd^4}}-b}{acd^2}}\log\left(\frac{\sqrt{-cx^4+bx^2+a}ad^2\sqrt{-\frac{1}{acd^4}}+\sqrt{-cx^4+bx^2+a}x^2+(acd^3x^3\sqrt{-\frac{1}{acd^4}}-adx)\sqrt{\frac{2acd^2\sqrt{-\frac{1}{acd^4}}-b}{acd^2}}}{cx^4+a}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-c*x^4+b*x^2+a)^(1/2)/(c*d*x^4+a*d), x, algorithm="fricas")

[Out] -1/8*sqrt((2*a*c*d^2*sqrt(-1/(a*c*d^4)) - b)/(a*c*d^2))*log(-(sqrt(-c*x^4 + b*x^2 + a)*a*d^2*sqrt(-1/(a*c*d^4)) + sqrt(-c*x^4 + b*x^2 + a)*x^2 + (a*c*d^3*x^3*sqrt(-1/(a*c*d^4)) - a*d*x)*sqrt(-1/(a*c*d^4)))/(c*d*x^4 + a*d))

$$d^3x^3\sqrt{-1/(acd^4)} - adx)\sqrt{(2acd^2\sqrt{-1/(acd^4)} - b)/(acd^2))}/(cx^4 + a) + 1/8\sqrt{(2acd^2\sqrt{-1/(acd^4)} - b)/(acd^2))*\log(-(\sqrt{-cx^4 + bx^2 + a})ad^2\sqrt{-1/(acd^4)} + \sqrt{-cx^4 + bx^2 + a})x^2 - (acd^3x^3\sqrt{-1/(acd^4)} - adx)\sqrt{(2acd^2\sqrt{-1/(acd^4)} - b)/(acd^2))}/(cx^4 + a) - 1/8\sqrt{-(2acd^2\sqrt{-1/(acd^4)} + b)/(acd^2))*\log((\sqrt{-cx^4 + bx^2 + a})ad^2\sqrt{-1/(acd^4)} - \sqrt{-cx^4 + bx^2 + a})x^2 + (acd^3x^3\sqrt{-1/(acd^4)} + adx)\sqrt{-(2acd^2\sqrt{-1/(acd^4)} + b)/(acd^2))}/(cx^4 + a) + 1/8\sqrt{-(2acd^2\sqrt{-1/(acd^4)} + b)/(acd^2))*\log((\sqrt{-cx^4 + bx^2 + a})ad^2\sqrt{-1/(acd^4)} - \sqrt{-cx^4 + bx^2 + a})x^2 - (acd^3x^3\sqrt{-1/(acd^4)} + adx)\sqrt{-(2acd^2\sqrt{-1/(acd^4)} + b)/(acd^2))}/(cx^4 + a)$$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-cx^4 + bx^2 + a}}{cdx^4 + ad} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-cx^4+bx^2+a)^(1/2)/(c*d*x^4+a*d),x, algorithm="giac")

[Out] integrate(sqrt(-cx^4 + bx^2 + a)/(c*d*x^4 + a*d), x)

maple [B] time = 0.11, size = 568, normalized size = 2.38

$$\frac{\sqrt{2} \arctan\left(\frac{-\frac{2\sqrt{-cx^4+bx^2+a}}{x}\sqrt{2}+2\sqrt{b+\sqrt{4ac+b^2}}}{2\sqrt{-b+\sqrt{4ac+b^2}}}\right)}{4\sqrt{-b+\sqrt{4ac+b^2}}d} - \frac{\sqrt{2} \arctan\left(\frac{\frac{2\sqrt{-cx^4+bx^2+a}}{x}\sqrt{2}+2\sqrt{b+\sqrt{4ac+b^2}}}{2\sqrt{-b+\sqrt{4ac+b^2}}}\right)}{4\sqrt{-b+\sqrt{4ac+b^2}}d} - \sqrt{2} \sqrt{b+\sqrt{4ac+b^2}} b l$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-cx^4+bx^2+a)^(1/2)/(c*d*x^4+a*d),x)

[Out] 1/32/d*2^(1/2)*(b+(4*a*c+b^2)^(1/2))^(1/2)/a/c*b*ln((-cx^4+bx^2+a)^(1/2)*2^(1/2)/x*(b+(4*a*c+b^2)^(1/2))^(1/2)-(-cx^4+bx^2+a)/x^2-(4*a*c+b^2)^(1/2))-1/32/d*2^(1/2)*(b+(4*a*c+b^2)^(1/2))^(1/2)/a/c*(4*a*c+b^2)^(1/2)*ln((-cx^4+bx^2+a)^(1/2)*2^(1/2)/x*(b+(4*a*c+b^2)^(1/2))^(1/2)-(-cx^4+bx^2+a)/x^2-(4*a*c+b^2)^(1/2))+1/4/d*2^(1/2)/(-b+(4*a*c+b^2)^(1/2))^(1/2)*arctan(1/2*(2*(b+(4*a*c+b^2)^(1/2))^(1/2)-2*(-cx^4+bx^2+a)^(1/2)*2^(1/2)/x)/(-b+(4*a*c+b^2)^(1/2))^(1/2))-1/32/d*2^(1/2)*(b+(4*a*c+b^2)^(1/2))^(1/2)/a/c*b*ln((-cx^4+bx^2+a)/x^2+(-cx^4+bx^2+a)^(1/2)*2^(1/2)/x*(b+(4*a*c+b^2)^(1/2))^(1/2)+(4*a*c+b^2)^(1/2))+1/32/d*2^(1/2)*(b+(4*a*c+b^2)^(1/2))^(1/2)/a/c*(4*a*c+b^2)^(1/2)*ln((-cx^4+bx^2+a)/x^2+(-cx^4+bx^2+a)^(1/2)*2^(1/2)/x*(b+(4*a*c+b^2)^(1/2))^(1/2)+(4*a*c+b^2)^(1/2))+1/4/d*2^(1/2)/(-b+(4*a*c+b^2)^(1/2))^(1/2)

$(1/2)^{(1/2)} \cdot \arctan(1/2 \cdot (2 \cdot (-c \cdot x^4 + b \cdot x^2 + a))^{(1/2)} \cdot 2^{(1/2)} / x + 2 \cdot (b + (4 \cdot a \cdot c + b^2)^{(1/2)})^{(1/2)}) / (-b + (4 \cdot a \cdot c + b^2)^{(1/2)})^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{-cx^4 + bx^2 + a}}{cdx^4 + ad} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-c*x^4+b*x^2+a)^(1/2)/(c*d*x^4+a*d),x, algorithm="maxima")

[Out] integrate(sqrt(-c*x^4 + b*x^2 + a)/(c*d*x^4 + a*d), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{-cx^4 + bx^2 + a}}{cdx^4 + ad} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2 - c*x^4)^(1/2)/(a*d + c*d*x^4),x)

[Out] int((a + b*x^2 - c*x^4)^(1/2)/(a*d + c*d*x^4), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\sqrt{a+bx^2-cx^4}}{a+cx^4} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-c*x**4+b*x**2+a)**(1/2)/(c*d*x**4+a*d),x)

[Out] Integral(sqrt(a + b*x**2 - c*x**4)/(a + c*x**4), x)/d

3.37 $\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}dx$

Optimal. Leaf size=309

$$\frac{e\sqrt{a^2+2abx^2+b^2x^4}(4cd-e^2)(-16ad^2+12bcd-7be^2)\tanh^{-1}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)+e\sqrt{a^2+2abx^2+b^2x^4}(2dx+e)}{256d^{9/2}(a+bx^2)} + \frac{12}{128d^4}$$

[Out] $1/5*b*x^2*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/d/(b*x^2+a)-1/240*(42*b*d*e*x-80*a*d^2+32*b*c*d-35*b*e^2)*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/d^3/(b*x^2+a)+1/256*e*(4*c*d-e^2)*(-16*a*d^2+12*b*c*d-7*b*e^2)*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/d^{(9/2)}/(b*x^2+a)+1/128*e*(-16*a*d^2+12*b*c*d-7*b*e^2)*(2*d*x+e)*(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/d^4/(b*x^2+a)$

Rubi [A] time = 0.63, antiderivative size = 309, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 38, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {6744, 1653, 779, 612, 621, 206}

$$\frac{\sqrt{a^2+2abx^2+b^2x^4}(c+dx^2+ex)^{3/2}(-80ad^2+32bcd+42bdex-35be^2)}{240d^3(a+bx^2)} + \frac{e\sqrt{a^2+2abx^2+b^2x^4}(2dx+e)\sqrt{c+dx^2+ex}}{128d^4}$$

Antiderivative was successfully verified.

[In] `Int[x*Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4], x]`

[Out] $(e*(12*b*c*d - 16*a*d^2 - 7*b*e^2)*(e + 2*d*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(128*d^4*(a + b*x^2)) + (b*x^2*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(5*d*(a + b*x^2)) - ((32*b*c*d - 80*a*d^2 - 35*b*e^2 + 42*b*d*e*x)*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(240*d^3*(a + b*x^2)) + (e*(4*c*d - e^2)*(12*b*c*d - 16*a*d^2 - 7*b*e^2)*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(256*d^{(9/2)}*(a + b*x^2))$

Rule 206

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 612

`Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[((b + 2*c*x)*(a + b*x + c*x^2)^p)/(2*c*(2*p + 1)), x] - Dist[(p*(b^2 - 4*a*c))/(2*c*(2*p + 1)), Int[(a + b*x + c*x^2)^(p-1), x]`

*p + 1)), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]

Rule 621

Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 779

Int[((d_) + (e_)*(x_))*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := -Simp[((b*e*g*(p + 2) - c*(e*f + d*g))*(2*p + 3) - 2*c*e*g*(p + 1)*x)*(a + b*x + c*x^2)^(p + 1))/(2*c^2*(p + 1)*(2*p + 3)), x] + Dist[(b^2*e*g*(p + 2) - 2*a*c*e*g + c*(2*c*d*f - b*(e*f + d*g))*(2*p + 3))/(2*c^2*(2*p + 3)), Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b^2 - 4*a*c, 0] && !LeQ[p, -1]

Rule 1653

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[(f*(d + e*x)^(m + q - 1)*(a + b*x + c*x^2)^(p + 1))/(c*e^(q - 1)*(m + q + 2*p + 1)), x] + Dist[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + b*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, d, e, m, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGtQ[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))

Rule 6744

Int[(u_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]

Rubi steps

$$\begin{aligned}
\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int x(2ab+2b^2x^2)\sqrt{c+ex+dx^2}dx}{2ab+2b^2x^2} \\
&= \frac{bx^2(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{5d(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4} \int}{5d(a+bx^2)} \\
&= \frac{bx^2(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{5d(a+bx^2)} - \frac{(32bcd-80ad^2-35be^2)}{5d(a+bx^2)} \\
&= \frac{e(12bcd-16ad^2-7be^2)(e+2dx)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{128d^4(a+bx^2)} \\
&= \frac{e(12bcd-16ad^2-7be^2)(e+2dx)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{128d^4(a+bx^2)} \\
&= \frac{e(12bcd-16ad^2-7be^2)(e+2dx)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{128d^4(a+bx^2)}
\end{aligned}$$

Mathematica [A] time = 0.33, size = 214, normalized size = 0.69

$$\frac{\sqrt{(a+bx^2)^2} \left(2\sqrt{d}\sqrt{c+x(dx+e)} (80ad^2(8cd+8d^2x^2+2dex-3e^2) + b(-256c^2d^2+4cd(32d^2x^2-58dex+11d^2))) \right)}{3840d^{9/2}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[x*Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4], x]

[Out] (Sqrt[(a + b*x^2)^2]*(2*Sqrt[d]*Sqrt[c + x*(e + d*x)]*(80*a*d^2*(8*c*d - 3*e^2 + 2*d*e*x + 8*d^2*x^2) + b*(-256*c^2*d^2 - 105*e^4 + 70*d*e^3*x - 56*d^2*e^2*x^2 + 48*d^3*e*x^3 + 384*d^4*x^4 + 4*c*d*(115*e^2 - 58*d*e*x + 32*d^2*x^2))) + 15*e*(-4*c*d + e^2)*(-12*b*c*d + 16*a*d^2 + 7*b*e^2)*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + x*(e + d*x)])))/(3840*d^(9/2)*(a + b*x^2))

fricas [A] time = 1.19, size = 469, normalized size = 1.52

$$\left[\frac{15(7be^5 - 8(5bcd - 2ad^2)e^3 + 16(3bc^2d^2 - 4acd^3)e)\sqrt{d} \log(8d^2x^2 + 8dex + 4\sqrt{dx^2 + ex + c}(2dx + e)\sqrt{d} + \dots}{\dots} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="fricas")

[Out] [1/7680*(15*(7*b*e^5 - 8*(5*b*c*d - 2*a*d^2)*e^3 + 16*(3*b*c^2*d^2 - 4*a*c*d^3)*e)*sqrt(d)*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(384*b*d^5*x^4 + 48*b*d^4*e*x^3 - 256*b*c^2*d^3 + 640*a*c*d^4 - 105*b*d*e^4 + 20*(23*b*c*d^2 - 12*a*d^3)*e^2 + 8*(16*b*c*d^4 + 80*a*d^5 - 7*b*d^3*e^2)*x^2 + 2*(35*b*d^2*e^3 - 4*(29*b*c*d^3 - 20*a*d^4)*e)*x)*sqrt(d*x^2 + e*x + c))/d^5, -1/3840*(15*(7*b*e^5 - 8*(5*b*c*d - 2*a*d^2)*e^3 + 16*(3*b*c^2*d^2 - 4*a*c*d^3)*e)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) - 2*(384*b*d^5*x^4 + 48*b*d^4*e*x^3 - 256*b*c^2*d^3 + 640*a*c*d^4 - 105*b*d*e^4 + 20*(23*b*c*d^2 - 12*a*d^3)*e^2 + 8*(16*b*c*d^4 + 80*a*d^5 - 7*b*d^3*e^2)*x^2 + 2*(35*b*d^2*e^3 - 4*(29*b*c*d^3 - 20*a*d^4)*e)*x)*sqrt(d*x^2 + e*x + c))/d^5]

giac [A] time = 0.60, size = 360, normalized size = 1.17

$$\frac{1}{1920} \sqrt{dx^2 + xe + c} \left(2 \left(4 \left(6 \left(8bx \operatorname{sgn}(bx^2 + a) + \frac{b \operatorname{sgn}(bx^2 + a)}{d} \right) \right) \right) x + \frac{16bcd^3 \operatorname{sgn}(bx^2 + a) + 80ad^4 \operatorname{sgn}(bx^2 + a)}{d^4} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="giac")

[Out] 1/1920*sqrt(d*x^2 + x*e + c)*(2*(4*(6*(8*b*x*sgn(b*x^2 + a) + b*e*sgn(b*x^2 + a)/d)*x + (16*b*c*d^3*sgn(b*x^2 + a) + 80*a*d^4*sgn(b*x^2 + a) - 7*b*d^2*e^2*sgn(b*x^2 + a))/d^4)*x - (116*b*c*d^2*e*sgn(b*x^2 + a) - 80*a*d^3*e*sgn(b*x^2 + a) - 35*b*d*e^3*sgn(b*x^2 + a))/d^4)*x - (256*b*c^2*d^2*sgn(b*x^2 + a) - 640*a*c*d^3*sgn(b*x^2 + a) - 460*b*c*d*e^2*sgn(b*x^2 + a) + 240*a*d^2*e^2*sgn(b*x^2 + a) + 105*b*e^4*sgn(b*x^2 + a))/d^4) - 1/256*(48*b*c^2*d^2*e*sgn(b*x^2 + a) - 64*a*c*d^3*e*sgn(b*x^2 + a) - 40*b*c*d*e^3*sgn(b*x^2 + a) + 16*a*d^2*e^3*sgn(b*x^2 + a) + 7*b*e^5*sgn(b*x^2 + a))*log(abs(-2*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))*sqrt(d) - e))/d^(9/2)

maple [A] time = 0.01, size = 442, normalized size = 1.43

$$\frac{\sqrt{(bx^2 + a)^2} \left(-960ac d^4 e \ln \left(\frac{2dx+e+2\sqrt{dx^2+ex+c} \sqrt{d}}{2\sqrt{d}} \right) + 240a d^3 e^3 \ln \left(\frac{2dx+e+2\sqrt{dx^2+ex+c} \sqrt{d}}{2\sqrt{d}} \right) + 720b c^2 d^3 e \ln \left(\frac{2dx+e+2\sqrt{dx^2+ex+c} \sqrt{d}}{2\sqrt{d}} \right) \right)}{d^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x)

```
[Out] 1/3840*((b*x^2+a)^2)^(1/2)*(768*(d*x^2+e*x+c)^(3/2)*d^(9/2)*x^2*b-672*(d*x^2+e*x+c)^(3/2)*d^(7/2)*x*b*e+1280*(d*x^2+e*x+c)^(3/2)*d^(9/2)*a-512*(d*x^2+e*x+c)^(3/2)*d^(7/2)*b*c+560*(d*x^2+e*x+c)^(3/2)*d^(5/2)*b*e^2-960*(d*x^2+e*x+c)^(1/2)*d^(9/2)*x*a*e+720*(d*x^2+e*x+c)^(1/2)*d^(7/2)*x*b*c*e-420*(d*x^2+e*x+c)^(1/2)*d^(5/2)*x*b*e^3-480*(d*x^2+e*x+c)^(1/2)*d^(7/2)*a*e^2+360*(d*x^2+e*x+c)^(1/2)*d^(5/2)*b*c*e^2-210*(d*x^2+e*x+c)^(1/2)*d^(3/2)*b*e^4-960*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*a*c*d^4*e+240*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*a*d^3*e^3+720*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*c^2*d^3*e-600*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*c*d^2*e^3+105*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*d*e^5)/(b*x^2+a)/d^(11/2)
```

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e^2-4*c*d>0)', see `assume?` for more details)Is e^2-4*c*d positive, negative or zero?
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int x \sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x*((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2),x)
```

```
[Out] int(x*((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x*(d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2),x)
```

```
[Out] Timed out
```

3.38 $\int \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4} dx$

Optimal. Leaf size=283

$$\frac{5be\sqrt{a^2 + 2abx^2 + b^2x^4} (c + dx^2 + ex)^{3/2} \sqrt{a^2 + 2abx^2 + b^2x^4} (4cd - e^2) (-16ad^2 + 4bcd - 5be^2) \tanh^{-1}\left(\frac{\sqrt{c + ex + dx^2}}{\sqrt{a^2 + 2abx^2 + b^2x^4}}\right)}{24d^2 (a + bx^2) 128d^{7/2} (a + bx^2)}$$

[Out] $-5/24*b*e*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/d^2/(b*x^2+a)+1/4*b*x*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/d/(b*x^2+a)-1/128*(4*c*d-e^2)*(-16*a*d^2+4*b*c*d-5*b*e^2)*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/d^{(7/2)}/(b*x^2+a)-1/64*(-16*a*d^2+4*b*c*d-5*b*e^2)*(2*d*x+e)*(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/d^3/(b*x^2+a)$

Rubi [A] time = 0.34, antiderivative size = 283, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 37, $\frac{\text{number of rules}}{\text{integrand size}} = 0.162$, Rules used = {6744, 1661, 640, 612, 621, 206}

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} (2dx + e) \sqrt{c + dx^2 + ex} (-16ad^2 + 4bcd - 5be^2) \sqrt{a^2 + 2abx^2 + b^2x^4} (4cd - e^2) (-16ad^2 + 4bcd - 5be^2)}{64d^3 (a + bx^2) 128d^{7/2} (a + bx^2)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4], x]$

[Out] $-((4*b*c*d - 16*a*d^2 - 5*b*e^2)*(e + 2*d*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(64*d^3*(a + b*x^2)) - (5*b*e*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(24*d^2*(a + b*x^2)) + (b*x*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(4*d*(a + b*x^2)) - ((4*c*d - e^2)*(4*b*c*d - 16*a*d^2 - 5*b*e^2)*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(128*d^{(7/2)}*(a + b*x^2))$

Rule 206

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1*\operatorname{ArcTanh}[(\operatorname{Rt}[-b, 2]*x)/\operatorname{Rt}[a, 2]])/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]), x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 612

$\operatorname{Int}[(a + (b \cdot x) + (c \cdot x)^2)^{p}, x_Symbol] \rightarrow \operatorname{Simp}[(b + 2*c*x)*(a + b*x + c*x^2)^p/(2*c*(2*p + 1)), x] - \operatorname{Dist}[(p*(b^2 - 4*a*c))/(2*c*(2*p + 1)), \operatorname{Int}[(a + b*x + c*x^2)^{p-1}, x], x] /; \operatorname{FreeQ}\{a, b, c\}, x] \ \&\& \ \operatorname{N}$

$eQ[b^2 - 4ac, 0] \ \&\& \ GtQ[p, 0] \ \&\& \ IntegerQ[4p]$

Rule 621

$Int[1/\sqrt{(a_.) + (b_.)x + (c_.)x^2}, x_Symbol] \ :> \ Dist[2, Subst[Int[1/(4c - x^2), x], x, (b + 2cx)/\sqrt{a + bx + cx^2}], x] \ /; \ FreeQ[\{a, b, c\}, x] \ \&\& \ NeQ[b^2 - 4ac, 0]$

Rule 640

$Int[((d_.) + (e_.)x) * ((a_.) + (b_.)x + (c_.)x^2)^{p_}, x_Symbol] \ :> \ Simp[(e(a + bx + cx^2)^{p+1}) / (2c(p+1)), x] + Dist[(2cd - be) / (2c), Int[(a + bx + cx^2)^p, x], x] \ /; \ FreeQ[\{a, b, c, d, e, p\}, x] \ \&\& \ NeQ[2cd - be, 0] \ \&\& \ NeQ[p, -1]$

Rule 1661

$Int[(Pq_)*((a_.) + (b_.)x + (c_.)x^2)^{p_}, x_Symbol] \ :> \ With[\{q = Expon[Pq, x], e = Coeff[Pq, x, Expon[Pq, x]]\}, Simp[(e*x^{q-1}*(a + bx + cx^2)^{p+1}) / (c*(q + 2p + 1)), x] + Dist[1/(c*(q + 2p + 1)), Int[(a + bx + cx^2)^p * ExpandToSum[c*(q + 2p + 1)*Pq - a*e*(q - 1)*x^{q-2} - b*e*(q + p)*x^{q-1} - c*e*(q + 2p + 1)*x^q, x], x], x] \ /; \ FreeQ[\{a, b, c, p\}, x] \ \&\& \ PolyQ[Pq, x] \ \&\& \ NeQ[b^2 - 4ac, 0] \ \&\& \ !LeQ[p, -1]$

Rule 6744

$Int[(u_)*((a_.) + (b_.)x^{n_}) + (c_.)x^{n2_})^{p_}, x_Symbol] \ :> \ Dist[\sqrt{a + bx^n + cx^{2n}} / ((4c)^{p-1/2} * (b + 2cx^n)), Int[u*(b + 2cx^n)^{2p}, x], x] \ /; \ FreeQ[\{a, b, c, n, p\}, x] \ \&\& \ EqQ[n2, 2n] \ \&\& \ EqQ[b^2 - 4ac, 0] \ \&\& \ IntegerQ[p - 1/2]$

Rubi steps

$$\begin{aligned}
\int \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int (2ab+2b^2x^2) \sqrt{c+ex+dx^2} dx}{2ab+2b^2x^2} \\
&= \frac{bx(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{4d(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4} \int (2ab+2b^2x^2) \sqrt{c+ex+dx^2} dx}{4d(a+bx^2)} \\
&= -\frac{5be(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{24d^2(a+bx^2)} + \frac{bx(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{4d(a+bx^2)} \\
&= -\frac{(4bcd-16ad^2-5be^2)(e+2dx)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{64d^3(a+bx^2)} \\
&= -\frac{(4bcd-16ad^2-5be^2)(e+2dx)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{64d^3(a+bx^2)} \\
&= -\frac{(4bcd-16ad^2-5be^2)(e+2dx)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{64d^3(a+bx^2)}
\end{aligned}$$

Mathematica [A] time = 0.22, size = 168, normalized size = 0.59

$$\frac{\sqrt{(a+bx^2)^2} \left(2\sqrt{d} \sqrt{c+x(dx+e)} (48ad^2(2dx+e) + b(4cd(6dx-13e) + 48d^3x^3 + 8d^2ex^2 - 10de^2x + 15e^3)) - \right)}{384d^{7/2}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4], x]

[Out] (Sqrt[(a + b*x^2)^2]*(2*Sqrt[d]*Sqrt[c + x*(e + d*x)]*(48*a*d^2*(e + 2*d*x) + b*(15*e^3 - 10*d*e^2*x + 8*d^2*e*x^2 + 48*d^3*x^3 + 4*c*d*(-13*e + 6*d*x))) - 3*(4*c*d - e^2)*(4*b*c*d - 16*a*d^2 - 5*b*e^2)*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + x*(e + d*x)])])/(384*d^(7/2)*(a + b*x^2))

fricas [A] time = 1.16, size = 361, normalized size = 1.28

$$\left[\frac{3(16bc^2d^2 - 64acd^3 + 5be^4 - 8(3bcd - 2ad^2)e^2)\sqrt{d} \log\left(8d^2x^2 + 8dex - 4\sqrt{dx^2 + ex + c}(2dx + e)\sqrt{d} + 4\right)}{384d^{7/2}(a + bx^2)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="fricas")

[Out] [1/768*(3*(16*b*c^2*d^2 - 64*a*c*d^3 + 5*b*e^4 - 8*(3*b*c*d - 2*a*d^2)*e^2)*sqrt(d)*log(8*d^2*x^2 + 8*d*e*x - 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(48*b*d^4*x^3 + 8*b*d^3*e*x^2 + 15*b*d*e^3 - 4*(13*b*c*d^2 - 12*a*d^3)*e + 2*(12*b*c*d^3 + 48*a*d^4 - 5*b*d^2*e^2)*x)*sqrt(d*x^2 + e*x + c))/d^4, 1/384*(3*(16*b*c^2*d^2 - 64*a*c*d^3 + 5*b*e^4 - 8*(3*b*c*d - 2*a*d^2)*e^2)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(48*b*d^4*x^3 + 8*b*d^3*e*x^2 + 15*b*d*e^3 - 4*(13*b*c*d^2 - 12*a*d^3)*e + 2*(12*b*c*d^3 + 48*a*d^4 - 5*b*d^2*e^2)*x)*sqrt(d*x^2 + e*x + c))/d^4]

giac [A] time = 0.64, size = 265, normalized size = 0.94

$$\frac{1}{192} \sqrt{dx^2 + xe + c} \left(2 \left(4 \left(6 b x \operatorname{sgn}(bx^2 + a) + \frac{b \operatorname{sgn}(bx^2 + a)}{d} \right) x + \frac{12 b c d^2 \operatorname{sgn}(bx^2 + a) + 48 a d^3 \operatorname{sgn}(bx^2 + a)}{d^3} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="giac")

[Out] 1/192*sqrt(d*x^2 + x*e + c)*(2*(4*(6*b*x*sgn(b*x^2 + a) + b*e*sgn(b*x^2 + a))/d)*x + (12*b*c*d^2*sgn(b*x^2 + a) + 48*a*d^3*sgn(b*x^2 + a) - 5*b*d*e^2*sgn(b*x^2 + a))/d^3)*x - (52*b*c*d*e*sgn(b*x^2 + a) - 48*a*d^2*e*sgn(b*x^2 + a) - 15*b*e^3*sgn(b*x^2 + a))/d^3 + 1/128*(16*b*c^2*d^2*sgn(b*x^2 + a) - 64*a*c*d^3*sgn(b*x^2 + a) - 24*b*c*d*e^2*sgn(b*x^2 + a) + 16*a*d^2*e^2*sgn(b*x^2 + a) + 5*b*e^4*sgn(b*x^2 + a))*log(abs(-2*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))*sqrt(d) - e))/d^(7/2)

maple [A] time = 0.01, size = 373, normalized size = 1.32

$$\frac{\sqrt{(bx^2 + a)^2} \left(192ac d^4 \ln \left(\frac{2dx+e+2\sqrt{dx^2+ex+c} \sqrt{d}}{2\sqrt{d}} \right) - 48a d^3 e^2 \ln \left(\frac{2dx+e+2\sqrt{dx^2+ex+c} \sqrt{d}}{2\sqrt{d}} \right) - 48b c^2 d^3 \ln \left(\frac{2dx+e+2\sqrt{dx^2+ex+c}}{2\sqrt{d}} \right) \right)}{1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x)

[Out] 1/384*((b*x^2+a)^2)^(1/2)*(96*d^(7/2)*(d*x^2+e*x+c)^(3/2)*x*b-80*d^(5/2)*(d*x^2+e*x+c)^(3/2)*b*e+192*d^(9/2)*(d*x^2+e*x+c)^(1/2)*x*a-48*d^(7/2)*(d*x^2+e*x+c)^(1/2)*x*b*c+60*d^(5/2)*(d*x^2+e*x+c)^(1/2)*x*b*e^2+96*d^(7/2)*(d*x^2+e*x+c)^(1/2)*a*e-24*d^(5/2)*(d*x^2+e*x+c)^(1/2)*b*c*e+30*d^(3/2)*(d*x^2+e*x+c)^(1/2)*b*e^3+192*ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^(1/2)*d^(1/2))/d^(1/2))*a*c*d^4-48*ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^(1/2)*d^(1/2))/d^(1/2))*a*d^3

```
*e^2-48*ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^(1/2)*d^(1/2))/d^(1/2))*b*c^2*d^3+7
2*ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^(1/2)*d^(1/2))/d^(1/2))*b*c*d^2*e^2-15*ln
(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^(1/2)*d^(1/2))/d^(1/2))*b*d*e^4)/(b*x^2+a)/d^
(9/2)
```

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested a
dditional constraints; using the 'assume' command before evaluation *may* h
elp (example of legal syntax is 'assume(e^2-4*c*d>0)', see `assume?` for mo
re details)Is e^2-4*c*d positive, negative or zero?
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2),x)
```

```
[Out] int(((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2),x)
```

```
[Out] Timed out
```

$$3.39 \quad \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x} dx$$

Optimal. Leaf size=286

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \sqrt{c+dx^2+ex} (8ad^2-2bdex-be^2)}{8d^2(a+bx^2)} + \frac{e\sqrt{a^2+2abx^2+b^2x^4} (8ad^2-b(4cd-e^2)) \tanh^{-1}\left(\frac{\sqrt{c+dx^2+ex}}{\sqrt{a+bx^2}}\right)}{16d^{5/2}(a+bx^2)}$$

[Out] 1/3*b*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/d/(b*x^2+a)+1/16*e*(8*a*d^2-b*(4*c*d-e^2))*arctanh(1/2*(2*d*x+e)/d^(1/2)/(d*x^2+e*x+c)^(1/2))*((b*x^2+a)^2)^(1/2)/d^(5/2)/(b*x^2+a)-a*arctanh(1/2*(e*x+2*c)/c^(1/2)/(d*x^2+e*x+c)^(1/2))*c^(1/2)*((b*x^2+a)^2)^(1/2)/(b*x^2+a)+1/8*(-2*b*d*e*x+8*a*d^2-b*e^2)*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/d^2/(b*x^2+a)

Rubi [A] time = 0.75, antiderivative size = 286, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 40, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {6744, 1653, 814, 843, 621, 206, 724}

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \sqrt{c+dx^2+ex} (8ad^2-2bdex-be^2)}{8d^2(a+bx^2)} + \frac{e\sqrt{a^2+2abx^2+b^2x^4} (8ad^2-b(4cd-e^2)) \tanh^{-1}\left(\frac{\sqrt{c+dx^2+ex}}{\sqrt{a+bx^2}}\right)}{16d^{5/2}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x,x]

[Out] ((8*a*d^2 - b*e^2 - 2*b*d*e*x)*Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/(8*d^2*(a + b*x^2)) + (b*(c + e*x + d*x^2)^(3/2)*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/(3*d*(a + b*x^2)) + (e*(8*a*d^2 - b*(4*c*d - e^2))*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4]*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + e*x + d*x^2])])/(16*d^(5/2)*(a + b*x^2)) - (a*Sqrt[c]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4]*ArcTanh[(2*c + e*x)/(2*Sqrt[c]*Sqrt[c + e*x + d*x^2])])/(a + b*x^2)

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 621

Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,

b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 724

Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 814

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[((d + e*x)^(m + 1)*(c*e*f*(m + 2*p + 2) - g*(c*d + 2*c*d*p - b*e*p) + g*c*e*(m + 2*p + 1)*x)*(a + b*x + c*x^2)^p)/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2)), x] - Dist[p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2)), Int[(d + e*x)^m*(a + b*x + c*x^2)^(p - 1)*Simp[c*e*f*(b*d - 2*a*e)*(m + 2*p + 2) + g*(a*e*(b*e - 2*c*d*m + b*e*m) + b*d*(b*e*p - c*d - 2*c*d*p)) + (c*e*f*(2*c*d - b*e)*(m + 2*p + 2) + g*(b^2*e^2*(p + m + 1) - 2*c^2*d^2*(1 + 2*p) - c*e*(b*d*(m - 2*p) + 2*a*e*(m + 2*p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[p, 0] && (IntegerQ[p] || !RationalQ[m] || (GeQ[m, -1] && LtQ[m, 0])) && !ILtQ[m + 2*p, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])

Rule 843

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1653

Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[(f*(d + e*x)^(m + q - 1)*(a + b*x + c*x^2)^(p + 1))/(c*e^(q - 1)*(m + q + 2*p + 1)), x] + Dist[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + b*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, d, e, m, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGtQ[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))

Rule 6744

Int[(u_)*((a_) + (b_)*(x_)^(n_)) + (c_)*(x_)^(n2_)]^(p_), x_Symbol] :> Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]

Rubi steps

$$\begin{aligned}
 \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(2ab+2b^2x^2)\sqrt{c+ex+dx^2}}{x} dx}{2ab+2b^2x^2} \\
 &= \frac{b(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{3d(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(6ad^2-2bdex)\sqrt{c+ex+dx^2}}{a+bx^2} dx}{3d(2ab+2b^2x^2)} \\
 &= \frac{(8ad^2-be^2-2bdex)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)} + \frac{b(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{3d(2ab+2b^2x^2)} \\
 &= \frac{(8ad^2-be^2-2bdex)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)} + \frac{b(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{3d(2ab+2b^2x^2)} \\
 &= \frac{(8ad^2-be^2-2bdex)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)} + \frac{b(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{3d(2ab+2b^2x^2)} \\
 &= \frac{(8ad^2-be^2-2bdex)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)} + \frac{b(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{3d(2ab+2b^2x^2)}
 \end{aligned}$$

Mathematica [A] time = 0.31, size = 176, normalized size = 0.62

$$\frac{\sqrt{(a+bx^2)^2} \left(2\sqrt{d} \left(\sqrt{c+x(dx+e)} (24ad^2 + b(8cd + 8d^2x^2 + 2dex - 3e^2)) - 24a\sqrt{c} d^2 \tanh^{-1} \left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+x(dx+e)}} \right) \right) \right)}{48d^{5/2}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x,x]

[Out] (Sqrt[(a + b*x^2)^2]*(3*e*(8*a*d^2 + b*(-4*c*d + e^2))*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + x*(e + d*x)])] + 2*Sqrt[d]*(Sqrt[c + x*(e + d*x)]*(24*a

$*d^2 + b*(8*c*d - 3*e^2 + 2*d*e*x + 8*d^2*x^2)) - 24*a*\text{Sqrt}[c]*d^2*\text{ArcTanh}[(2*c + e*x)/(2*\text{Sqrt}[c]*\text{Sqrt}[c + x*(e + d*x)])])]/(48*d^{(5/2)}*(a + b*x^2))$

fricas [A] time = 2.91, size = 743, normalized size = 2.60

$$\frac{48 a \sqrt{c} d^3 \log\left(\frac{8 c e x + (4 c d + e^2) x^2 - 4 \sqrt{d x^2 + e x + c} (e x + 2 c) \sqrt{c + 8 c^2}}{x^2}\right) + 3 (b e^3 - 4 (b c d - 2 a d^2) e) \sqrt{d} \log\left(8 d^2 x^2 + 8 d e x + 4\right)}{96 d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="fricas")

[Out] [1/96*(48*a*sqrt(c)*d^3*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) + 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(d)*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(8*b*d^3*x^2 + 2*b*d^2*e*x + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + e*x + c))/d^3, 1/48*(24*a*sqrt(c)*d^3*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) - 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(8*b*d^3*x^2 + 2*b*d^2*e*x + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + e*x + c))/d^3, 1/96*(96*a*sqrt(-c)*d^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) + 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(d)*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(8*b*d^3*x^2 + 2*b*d^2*e*x + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + e*x + c))/d^3, 1/48*(48*a*sqrt(-c)*d^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(8*b*d^3*x^2 + 2*b*d^2*e*x + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + e*x + c))/d^3]

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,x);;OUTPUT:Warning, integration of abs or sign assumes constant sign by intervals (correct if the argument is real):Check [abs(b*x^2+a)]Warning, replacing 0 by `u`, a substitution variable should perhaps be purged.Warning, replacing 0 by `u`, a substitution variable should perh

aps be purged.Warning, replacing 0 by `u`, a substitution variable should perhaps be purged.index.cc index_m operator + Error: Bad Argument Value

maple [A] time = 0.01, size = 251, normalized size = 0.88

$$\sqrt{(bx^2 + a)^2} \left(48a\sqrt{c} d^{\frac{7}{2}} \ln\left(\frac{ex+2c+2\sqrt{dx^2+ex+c}\sqrt{c}}{x}\right) - 24ad^3e \ln\left(\frac{2dx+e+2\sqrt{dx^2+ex+c}\sqrt{d}}{2\sqrt{d}}\right) + 12bc d^2e \ln\left(\frac{2dx+e+2\sqrt{dx^2+ex+c}\sqrt{d}}{2\sqrt{d}}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x)

[Out] $-1/48*((b*x^2+a)^2)^{(1/2)}*(48*c^{(1/2)}*d^{(7/2)}*\ln((2*c+e*x+2*c^{(1/2)}*(d*x^2+e*x+c)^{(1/2)})/x)*a-16*d^{(5/2)}*(d*x^2+e*x+c)^{(3/2)}*b+12*d^{(5/2)}*(d*x^2+e*x+c)^{(1/2)}*x*b*e-48*d^{(7/2)}*(d*x^2+e*x+c)^{(1/2)}*a+6*d^{(3/2)}*(d*x^2+e*x+c)^{(1/2)}*b*e^2-24*d^3*\ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^{(1/2)}*d^{(1/2)})/d^{(1/2)})*a*e+12*\ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^{(1/2)}*d^{(1/2)})/d^{(1/2)})*b*c*d^2*e-3*\ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^{(1/2)}*d^{(1/2)})/d^{(1/2)})*b*d*e^3)/(b*x^2+a)/d^{(7/2)}$

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e^2-4*c*d>0)', see `assume?` for more details)Is e^2-4*c*d positive, negative or zero?

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x,x)

[Out] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2 + ex} \sqrt{(a + bx^2)^2}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x,x)

[Out] Integral(sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2)/x, x)

$$3.40 \quad \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$$

Optimal. Leaf size=294

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} (8ad^2 + 4bcd - be^2) \tanh^{-1} \left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}} \right) a\sqrt{a^2 + 2abx^2 + b^2x^4} (c + dx^2 + ex)^{3/2}}{8d^{3/2} (a + bx^2) cx (a + bx^2)} + \frac{\sqrt{a^2 + 2abx^2 + b^2x^4}}{cx (a + bx^2)}$$

[Out] $-a*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/c/x/(b*x^2+a)+1/8*(8*a*d^2+4*b*c*d-b*e^2)*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/d^{(3/2)}/(b*x^2+a)-1/2*a*e*\operatorname{arctanh}(1/2*(e*x+2*c)/c^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/(b*x^2+a)/c^{(1/2)}+1/4*((4*a*d+b*c)*e+2*d*(2*a*d+b*c)*x)*(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/c/d/(b*x^2+a)$

Rubi [A] time = 0.77, antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 40, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {6744, 1650, 814, 843, 621, 206, 724}

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} (8ad^2 + 4bcd - be^2) \tanh^{-1} \left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}} \right) a\sqrt{a^2 + 2abx^2 + b^2x^4} (c + dx^2 + ex)^{3/2}}{8d^{3/2} (a + bx^2) cx (a + bx^2)} + \frac{\sqrt{a^2 + 2abx^2 + b^2x^4}}{cx (a + bx^2)}$$

Antiderivative was successfully verified.

[In] Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^2, x]

[Out] $((b*c + 4*a*d)*e + 2*d*(b*c + 2*a*d)*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]/(4*c*d*(a + b*x^2)) - (a*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]/(c*x*(a + b*x^2)) + ((4*b*c*d + 8*a*d^2 - b*e^2)*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(8*d^{(3/2)}*(a + b*x^2)) - (a*e*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(2*c + e*x)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(2*\operatorname{Sqrt}[c]*(a + b*x^2))$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 621

Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,

b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 724

Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 814

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[((d + e*x)^(m + 1)*(c*e*f*(m + 2*p + 2) - g*(c*d + 2*c*d*p - b*e*p) + g*c*e*(m + 2*p + 1)*x)*(a + b*x + c*x^2)^p)/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2)), x] - Dist[p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2)), Int[(d + e*x)^m*(a + b*x + c*x^2)^(p - 1)*Simp[c*e*f*(b*d - 2*a*e)*(m + 2*p + 2) + g*(a*e*(b*e - 2*c*d*m + b*e*m) + b*d*(b*e*p - c*d - 2*c*d*p)) + (c*e*f*(2*c*d - b*e)*(m + 2*p + 2) + g*(b^2*e^2*(p + m + 1) - 2*c^2*d^2*(1 + 2*p) - c*e*(b*d*(m - 2*p) + 2*a*e*(m + 2*p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[p, 0] && (IntegerQ[p] || !RationalQ[m] || (GeQ[m, -1] && LtQ[m, 0])) && !ILtQ[m + 2*p, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])

Rule 843

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1650

Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, d + e*x, x], R = PolynomialRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m + 1)*(c*d^2 - b*d*e + a*e^2)*Q + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R*(m + 2*p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]

Rule 6744

```
Int[(u_)*((a_) + (b_)*(x_)^(n_)) + (c_)*(x_)^(n2_))^(p_), x_Symbol] :> Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(2ab+2b^2x^2)\sqrt{c+ex+dx^2}}{x^2} dx}{2ab+2b^2x^2} \\ &= -\frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{cx(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{2ab+2b^2x^2}{x^2} dx}{c(2ab+2b^2x^2)} \\ &= \frac{((bc+4ad)e+2d(bc+2ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)} - \frac{2ab}{c} \\ &= \frac{((bc+4ad)e+2d(bc+2ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)} - \frac{2ab}{c} \\ &= \frac{((bc+4ad)e+2d(bc+2ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)} - \frac{2ab}{c} \\ &= \frac{((bc+4ad)e+2d(bc+2ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)} - \frac{2ab}{c} \end{aligned}$$

Mathematica [A] time = 0.25, size = 171, normalized size = 0.58

$$\frac{\sqrt{(a+bx^2)^2} \left(\sqrt{c} x (8ad^2 + 4bcd - be^2) \tanh^{-1} \left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+x(dx+e)}} \right) + 2\sqrt{d} \left(\sqrt{c} \sqrt{c+x(dx+e)} (bx(2dx+e) - 4ad) - \sqrt{c} \sqrt{c+x(dx+e)} \right) \right)}{8\sqrt{c} d^{3/2} x (a+bx^2)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^2,x]
```

```
[Out] (Sqrt[(a + b*x^2)^2]*(Sqrt[c]*(4*b*c*d + 8*a*d^2 - b*e^2)*x*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + x*(e + d*x)])] + 2*Sqrt[d]*(Sqrt[c]*Sqrt[c + x*(e + d*x)]*(-4*a*d + b*x*(e + 2*d*x)) - 2*a*d*e*x*ArcTanh[(2*c + e*x)/(2*Sqrt[c]*Sqrt[c + x*(e + d*x)])])))/(8*Sqrt[c]*d^(3/2)*x*(a + b*x^2))
```

fricas [A] time = 2.44, size = 731, normalized size = 2.49

$$\left[\frac{4 a \sqrt{c} d^2 e x \log \left(\frac{8 c e x + (4 c d + e^2) x^2 - 4 \sqrt{d x^2 + e x + c} (e x + 2 c) \sqrt{c} + 8 c^2}{x^2} \right) - (4 b c^2 d + 8 a c d^2 - b c e^2) \sqrt{d} x \log \left(8 d^2 x^2 + 8 d e x - 4 \right)}{16 c d^2 x} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="fricas")

[Out] [1/16*(4*a*sqrt(c)*d^2*e*x*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) - (4*b*c^2*d + 8*a*c*d^2 - b*c*e^2)*sqrt(d)*x*log(8*d^2*x^2 + 8*d*e*x - 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(2*b*c*d^2*x^2 + b*c*d*e*x - 4*a*c*d^2)*sqrt(d*x^2 + e*x + c))/(c*d^2*x), 1/8*(2*a*sqrt(c)*d^2*e*x*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) - (4*b*c^2*d + 8*a*c*d^2 - b*c*e^2)*sqrt(-d)*x*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(2*b*c*d^2*x^2 + b*c*d*e*x - 4*a*c*d^2)*sqrt(d*x^2 + e*x + c))/(c*d^2*x), 1/16*(8*a*sqrt(-c)*d^2*e*x*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - (4*b*c^2*d + 8*a*c*d^2 - b*c*e^2)*sqrt(d)*x*log(8*d^2*x^2 + 8*d*e*x - 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(2*b*c*d^2*x^2 + b*c*d*e*x - 4*a*c*d^2)*sqrt(d*x^2 + e*x + c))/(c*d^2*x), 1/8*(4*a*sqrt(-c)*d^2*e*x*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - (4*b*c^2*d + 8*a*c*d^2 - b*c*e^2)*sqrt(-d)*x*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(2*b*c*d^2*x^2 + b*c*d*e*x - 4*a*c*d^2)*sqrt(d*x^2 + e*x + c))/(c*d^2*x)]

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,x);OUTPUT:Warning, integration of abs or sign assumes constant sign by intervals (correct if the argument is real):Check [abs(b*x^2+a)]Unable to divide, perhaps due to rounding error%%{%%{[1,0]:[1,0,%%{-1,[1]%%}}]%%}, [4,0,0,0,0,0]%%}+%%{%%{[-2,0]:[1,0,%%{-1,[1]%%}}]%%}, [2,0,0,0,1,0]%%}+%%{%%{[1,0]:[1,0,%%{-1,[1]%%}}]%%}, [0,0,0,0,2,0]%%} / %%{%%{1,[1]%%}, [4,0,0,0,0,0]%%}+%%{%%{[-2,[1]%%}, [2,0,0,0,1,0]%%}+%%{%%{1,[1]%%}, [0,0,0,0,2,0]%%}} Error: Bad Argument Value

maple [A] time = 0.01, size = 288, normalized size = 0.98

$$\sqrt{(bx^2 + a)^2} \left(-8ac d^3 x \ln \left(\frac{2dx+e+2\sqrt{dx^2+ex+c} \sqrt{d}}{2\sqrt{d}} \right) + 4a\sqrt{c} d^{\frac{5}{2}} ex \ln \left(\frac{ex+2c+2\sqrt{dx^2+ex+c} \sqrt{c}}{x} \right) - 4b c^2 d^2 x \ln \left(\frac{2dx+e+2\sqrt{dx^2+ex+c}}{2\sqrt{d}} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x)

[Out] $-1/8*((b*x^2+a)^2)^{(1/2)}*(-8*(d*x^2+e*x+c)^{(1/2)}*d^{(7/2)}*x^2*a-4*(d*x^2+e*x+c)^{(1/2)}*d^{(5/2)}*x^2*b*c+4*c^{(1/2)}*\ln((e*x+2*c+2*(d*x^2+e*x+c)^{(1/2)}*c^{(1/2)})/x)*d^{(5/2)}*x*a*e+8*(d*x^2+e*x+c)^{(3/2)}*d^{(5/2)}*a-8*(d*x^2+e*x+c)^{(1/2)}*d^{(5/2)}*x*a*e-2*(d*x^2+e*x+c)^{(1/2)}*d^{(3/2)}*x*b*c*e-8*\ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^{(1/2)}*d^{(1/2)})/d^{(1/2)})*x*a*c*d^3-4*\ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^{(1/2)}*d^{(1/2)})/d^{(1/2)})*x*b*c^2*d^2+\ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^{(1/2)}*d^{(1/2)})/d^{(1/2)})*x*b*c*d*e^2)/(b*x^2+a)/c/x/d^{(5/2)}$

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e^2-4*c*d>0)', see `assume?` for more details)Is e^2-4*c*d positive, negative or zero?

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^2,x)

[Out] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^2, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**2,x)
```

```
[Out] Timed out
```

$$3.41 \quad \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$$

Optimal. Leaf size=288

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} (4acd - ae^2 + 8bc^2) \tanh^{-1}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+dx^2+ex}}\right)}{8c^{3/2}(a+bx^2)} - \frac{a\sqrt{a^2+2abx^2+b^2x^4} (c+dx^2+ex)^{3/2}}{2cx^2(a+bx^2)} + \frac{\sqrt{a^2}}{2c}$$

[Out] $-1/2*a*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/c/x^2/(b*x^2+a)-1/8*(4*a*c*d - a*e^2+8*b*c^2)*\arctanh(1/2*(e*x+2*c)/c^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/c^{(3/2)}/(b*x^2+a)+1/2*b*e*\arctanh(1/2*(2*d*x+e)/d^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/(b*x^2+a)/d^{(1/2)}+1/4*(a*e+2*(a*d+2*b*c)*x)*(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/c/x/(b*x^2+a)$

Rubi [A] time = 0.78, antiderivative size = 288, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 40, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {6744, 1650, 812, 843, 621, 206, 724}

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} (4acd - ae^2 + 8bc^2) \tanh^{-1}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+dx^2+ex}}\right)}{8c^{3/2}(a+bx^2)} - \frac{a\sqrt{a^2+2abx^2+b^2x^4} (c+dx^2+ex)^{3/2}}{2cx^2(a+bx^2)} + \frac{\sqrt{a^2}}{2c}$$

Antiderivative was successfully verified.

[In] Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^3, x]

[Out] $((a*e + 2*(2*b*c + a*d)*x)*\text{Sqrt}[c + e*x + d*x^2]*\text{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(4*c*x*(a + b*x^2)) - (a*(c + e*x + d*x^2)^{(3/2)}*\text{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(2*c*x^2*(a + b*x^2)) + (b*e*\text{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\text{ArcTanh}[(e + 2*d*x)/(2*\text{Sqrt}[d]*\text{Sqrt}[c + e*x + d*x^2])])/(2*\text{Sqrt}[d]*(a + b*x^2)) - ((8*b*c^2 + 4*a*c*d - a*e^2)*\text{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\text{ArcTanh}[(2*c + e*x)/(2*\text{Sqrt}[c]*\text{Sqrt}[c + e*x + d*x^2])])/(8*c^{(3/2)}*(a + b*x^2))$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 621

Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,

b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 724

Int[1/(((d_.) + (e_.)*(x_.))*Sqrt[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 812

Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(p_.), x_Symbol] := Simp[((d + e*x)^(m + 1)*(e*f*(m + 2*p + 2) - d*g*(2*p + 1) + e*g*(m + 1)*x)*(a + b*x + c*x^2)^p)/(e^2*(m + 1)*(m + 2*p + 2)), x] + Dist[p/(e^2*(m + 1)*(m + 2*p + 2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p - 1)*Simp[g*(b*d + 2*a*e + 2*a*e*m + 2*b*d*p) - f*b*e*(m + 2*p + 2) + (g*(2*c*d + b*e + b*e*m + 4*c*d*p) - 2*c*e*f*(m + 2*p + 2))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && RationalQ[p] && p > 0 && (LtQ[m, -1] || EqQ[p, 1] || (IntegerQ[p] && !RationalQ[m])) && NeQ[m, -1] && !ILtQ[m + 2*p + 1, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])

Rule 843

Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(p_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1650

Int[(Pq_)*((d_.) + (e_.)*(x_.))^(m_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(p_.), x_Symbol] := With[{Q = PolynomialQuotient[Pq, d + e*x, x], R = PolynomialRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m + 1)*(c*d^2 - b*d*e + a*e^2)*Q + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R*(m + 2*p + 3)*x, x], x]] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]

Rule 6744

Int[(u_.)*((a_.) + (b_.)*(x_.)^(n_.) + (c_.)*(x_.)^(2*n_.))^(p_.), x_Symbol] := Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b +

$2*c*x^n)^{(2*p)}, x], x] /; \text{FreeQ}[\{a, b, c, n, p\}, x] \ \&\& \ \text{EqQ}[n2, 2*n] \ \&\& \ \text{EqQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{IntegerQ}[p - 1/2]$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(2ab+2b^2x^2)\sqrt{c+ex+dx^2}}{x^3} dx}{2ab+2b^2x^2} \\ &= -\frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{a}{2c(2ab+2b^2x^2)} dx}{2c(2ab+2b^2x^2)} \\ &= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} \\ &= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} \\ &= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} \\ &= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} \end{aligned}$$

Mathematica [A] time = 0.34, size = 177, normalized size = 0.61

$$\frac{\sqrt{(a+bx^2)^2} \left(4bc^{3/2}ex^2 \tanh^{-1}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+x(dx+e)}}\right) - \sqrt{d} \left(x^2(4acd - ae^2 + 8bc^2) \tanh^{-1}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+x(dx+e)}}\right) + 2\sqrt{c}\sqrt{c+x(dx+e)} \right) \right)}{8c^{3/2}\sqrt{d}x^2(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^3,x]

[Out] (Sqrt[(a + b*x^2)^2]*(4*b*c^(3/2)*e*x^2*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + x*(e + d*x)])]) - Sqrt[d]*(2*Sqrt[c]*(2*a*c + a*e*x - 4*b*c*x^2)*Sqrt[c + x*(e + d*x)] + (8*b*c^2 + 4*a*c*d - a*e^2)*x^2*ArcTanh[(2*c + e*x)/(2*Sqrt[c]*Sqrt[c + x*(e + d*x)])]))/(8*c^(3/2)*Sqrt[d]*x^2*(a + b*x^2))

fricas [A] time = 2.21, size = 749, normalized size = 2.60

$$\left[\frac{4bc^2\sqrt{d}ex^2 \log\left(8d^2x^2 + 8dex + 4\sqrt{dx^2 + ex + c}(2dx + e)\sqrt{d} + 4cd + e^2\right) - (8bc^2d + 4acd^2 - ade^2)\sqrt{c}x^2 \log\left(\frac{8c^2d + 4acd^2 - ade^2}{16c^2dx^2}\right)}{16c^2dx^2} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="fricas")

[Out] [1/16*(4*b*c^2*sqrt(d)*e*x^2*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) - (8*b*c^2*d + 4*a*c*d^2 - a*d*e^2)*sqrt(c)*x^2*log((8*c*e*x + (4*c*d + e^2)*x^2 + 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) + 4*(4*b*c^2*d*x^2 - a*c*d*e*x - 2*a*c^2*d)*sqrt(d*x^2 + e*x + c)/(c^2*d*x^2), -1/16*(8*b*c^2*sqrt(-d)*e*x^2*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + (8*b*c^2*d + 4*a*c*d^2 - a*d*e^2)*sqrt(c)*x^2*log((8*c*e*x + (4*c*d + e^2)*x^2 + 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) - 4*(4*b*c^2*d*x^2 - a*c*d*e*x - 2*a*c^2*d)*sqrt(d*x^2 + e*x + c)/(c^2*d*x^2), 1/8*(2*b*c^2*sqrt(d)*e*x^2*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + (8*b*c^2*d + 4*a*c*d^2 - a*d*e^2)*sqrt(-c)*x^2*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) + 2*(4*b*c^2*d*x^2 - a*c*d*e*x - 2*a*c^2*d)*sqrt(d*x^2 + e*x + c)/(c^2*d*x^2), -1/8*(4*b*c^2*sqrt(-d)*e*x^2*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) - (8*b*c^2*d + 4*a*c*d^2 - a*d*e^2)*sqrt(-c)*x^2*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - 2*(4*b*c^2*d*x^2 - a*c*d*e*x - 2*a*c^2*d)*sqrt(d*x^2 + e*x + c)/(c^2*d*x^2)]

giac [A] time = 0.89, size = 370, normalized size = 1.28

$$\frac{be \log\left(\left|-2\left(\sqrt{d}x - \sqrt{dx^2 + xe + c}\right)d - \sqrt{d}e\right|\right) \operatorname{sgn}(bx^2 + a)}{2\sqrt{d}} + \sqrt{dx^2 + xe + c} b \operatorname{sgn}(bx^2 + a) + \frac{(8bc^2 \operatorname{sgn}(bx^2 + a))}{16c^2dx^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="giac")

[Out] -1/2*b*e*log(abs(-2*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))*d - sqrt(d)*e))*sgn(b*x^2 + a)/sqrt(d) + sqrt(d*x^2 + x*e + c)*b*sgn(b*x^2 + a) + 1/4*(8*b*c^2*sgn(b*x^2 + a) + 4*a*c*d*sgn(b*x^2 + a) - a*e^2*sgn(b*x^2 + a))*arctan(-(s

$$\frac{\sqrt{d}x - \sqrt{dx^2 + x^e + c}}{\sqrt{-c}} / (\sqrt{-c} * c) + \frac{1}{4} * (4 * (\sqrt{d} * x - \sqrt{dx^2 + x^e + c})^3 * a * c * d * \text{sgn}(b * x^2 + a) + 8 * (\sqrt{d} * x - \sqrt{dx^2 + x^e + c})^2 * a * c * \sqrt{d} * e * \text{sgn}(b * x^2 + a) + 4 * (\sqrt{d} * x - \sqrt{dx^2 + x^e + c}) * a * c^2 * d * \text{sgn}(b * x^2 + a) + (\sqrt{d} * x - \sqrt{dx^2 + x^e + c})^3 * a * e^2 * \text{sgn}(b * x^2 + a) + (\sqrt{d} * x - \sqrt{dx^2 + x^e + c}) * a * c * e^2 * \text{sgn}(b * x^2 + a)) / (((\sqrt{d} * x - \sqrt{dx^2 + x^e + c})^2 - c)^2 * c)$$

maple [A] time = 0.01, size = 329, normalized size = 1.14

$$\frac{\sqrt{(bx^2 + a)^2} \left(-4ac^{\frac{3}{2}}d^{\frac{5}{2}}x^2 \ln\left(\frac{ex+2c+2\sqrt{dx^2+ex+c}\sqrt{c}}{x}\right) + a\sqrt{c}d^{\frac{3}{2}}e^2x^2 \ln\left(\frac{ex+2c+2\sqrt{dx^2+ex+c}\sqrt{c}}{x}\right) - 8bc^{\frac{5}{2}}d^{\frac{3}{2}}x^2 \ln\left(\frac{ex+2c}{x}\right) \right)}{((\sqrt{d}x - \sqrt{dx^2 + x^e + c})^2 - c)^2 c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x)

[Out] $\frac{1}{8} * ((b * x^2 + a)^2)^{(1/2)} * (-4 * d^{(5/2)} * c^{(3/2)} * \ln((e * x + 2 * c + 2 * (d * x^2 + e * x + c)^{(1/2)} * c^{(1/2)}) / x) * x^2 * a - 8 * d^{(3/2)} * c^{(5/2)} * \ln((e * x + 2 * c + 2 * (d * x^2 + e * x + c)^{(1/2)} * c^{(1/2)}) / x) * x^2 * b - 2 * (d * x^2 + e * x + c)^{(1/2)} * d^{(5/2)} * x^3 * a * e + 4 * (d * x^2 + e * x + c)^{(1/2)} * d^{(5/2)} * x^2 * a * c + d^{(3/2)} * c^{(1/2)} * \ln((e * x + 2 * c + 2 * (d * x^2 + e * x + c)^{(1/2)} * c^{(1/2)}) / x) * x^2 * a * e^2 + 2 * (d * x^2 + e * x + c)^{(3/2)} * d^{(3/2)} * x * a * e - 2 * (d * x^2 + e * x + c)^{(1/2)} * d^{(3/2)} * x^2 * a * e^2 + 8 * (d * x^2 + e * x + c)^{(1/2)} * d^{(3/2)} * x^2 * b * c^2 - 4 * (d * x^2 + e * x + c)^{(3/2)} * d^{(3/2)} * a * c + 4 * \ln(1/2 * (2 * d * x + e + 2 * (d * x^2 + e * x + c)^{(1/2)} * d^{(1/2)}) / d^{(1/2)}) * d * x^2 * b * c^2 * e) / d^{(3/2)} / x^2 / c^2 / (b * x^2 + a)$

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e^2-4*c*d>0)', see `assume?` for more details)Is e^2-4*c*d positive, negative or zero?

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^3,x)
```

```
[Out] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^3, x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**3,x)
```

```
[Out] Timed out
```

$$3.42 \quad \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$$

Optimal. Leaf size=294

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \sqrt{c+dx^2+ex} (2ace - x(8bc^2 - ae^2))}{8c^2x^2(a+bx^2)} - \frac{e\sqrt{a^2+2abx^2+b^2x^4} (8bc^2 - a(4cd - e^2)) \tanh^{-1}\left(\frac{\sqrt{c+dx^2+ex}}{\sqrt{a+bx^2}}\right)}{16c^{5/2}(a+bx^2)}$$

[Out] $-1/3*a*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/c/x^3/(b*x^2+a)-1/16*e*(8*b*c^2-a*(4*c*d-e^2))*\operatorname{arctanh}(1/2*(e*x+2*c)/c^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/c^{(5/2)}/(b*x^2+a)+b*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*d^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/(b*x^2+a)+1/8*(2*a*c*e-(-a*e^2+8*b*c^2)*x)*(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/c^2/x^2/(b*x^2+a)$

Rubi [A] time = 0.79, antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 40, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {6744, 1650, 810, 843, 621, 206, 724}

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \sqrt{c+dx^2+ex} (2ace - x(8bc^2 - ae^2))}{8c^2x^2(a+bx^2)} - \frac{e\sqrt{a^2+2abx^2+b^2x^4} (8bc^2 - a(4cd - e^2)) \tanh^{-1}\left(\frac{\sqrt{c+dx^2+ex}}{\sqrt{a+bx^2}}\right)}{16c^{5/2}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^4, x]

[Out] $((2*a*c*e - (8*b*c^2 - a*e^2)*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(8*c^2*x^2*(a + b*x^2)) - (a*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(3*c*x^3*(a + b*x^2)) + (b*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(a + b*x^2) - (e*(8*b*c^2 - a*(4*c*d - e^2))*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(2*c + e*x)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(16*c^{(5/2)}*(a + b*x^2))$

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 621

Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,

b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 724

Int[1/(((d_.) + (e_.)*(x_.))*Sqrt[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 810

Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(p_.), x_Symbol] := -Simp[((d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*((d*g - e*f*(m + 2))*(c*d^2 - b*d*e + a*e^2) - d*p*(2*c*d - b*e)*(e*f - d*g) - e*(g*(m + 1)*(c*d^2 - b*d*e + a*e^2) + p*(2*c*d - b*e)*(e*f - d*g))*x)/(e^2*(m + 1)*(m + 2)*(c*d^2 - b*d*e + a*e^2)), x] - Dist[p/(e^2*(m + 1)*(m + 2)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 2)*(a + b*x + c*x^2)^(p - 1)*Simp[2*a*c*e*(e*f - d*g)*(m + 2) + b^2*e*(d*g*(p + 1) - e*f*(m + p + 2)) + b*(a*e^2*g*(m + 1) - c*d*(d*g*(2*p + 1) - e*f*(m + 2*p + 2))) - c*(2*c*d*(d*g*(2*p + 1) - e*f*(m + 2*p + 2)) - e*(2*a*e*g*(m + 1) - b*(d*g*(m - 2*p) + e*f*(m + 2*p + 2)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[p, 0] && LtQ[m, -2] && LtQ[m + 2*p, 0] && !ILtQ[m + 2*p + 3, 0]

Rule 843

Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(p_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1650

Int[(Pq_)*((d_.) + (e_.)*(x_.))^(m_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(p_.), x_Symbol] := With[{Q = PolynomialQuotient[Pq, d + e*x, x], R = PolynomialRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m + 1)*(c*d^2 - b*d*e + a*e^2)*Q + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R*(m + 2*p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]

Rule 6744

```
Int[(u_)*((a_) + (b_)*(x_)^(n_)) + (c_)*(x_)^(n2_)]^(p_), x_Symbol] :> Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(2ab+2b^2x^2)\sqrt{c+ex+dx^2}}{x^4} dx}{2ab+2b^2x^2} \\ &= -\frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{3cx^3(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{3}{3c(2ab+2b^2x^2)} dx}{3c(2ab+2b^2x^2)} \\ &= \frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{3cx^3(a+bx^2)} \\ &= \frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{3cx^3(a+bx^2)} \\ &= \frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{3cx^3(a+bx^2)} \\ &= \frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{3cx^3(a+bx^2)} \end{aligned}$$

Mathematica [A] time = 0.29, size = 189, normalized size = 0.64

$$\frac{\sqrt{(a+bx^2)^2} \left(-3ex^3 (a(e^2 - 4cd) + 8bc^2) \tanh^{-1} \left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+x(dx+e)}} \right) - 2\sqrt{c} \sqrt{c+x(dx+e)} (a(8c^2 + 2cx(4dx+e) - 3e^2) + 2c^2) \right)}{48c^{5/2}x^3(a+bx^2)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^4,x]
```

```
[Out] (Sqrt[(a + b*x^2)^2]*(-2*Sqrt[c]*Sqrt[c + x*(e + d*x)]*(24*b*c^2*x^2 + a*(8*c^2 - 3*e^2*x^2 + 2*c*x*(e + 4*d*x))) + 48*b*c^(5/2)*Sqrt[d]*x^3*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + x*(e + d*x)])] - 3*e*(8*b*c^2 + a*(-4*c*d +
```


$e^2)) * x^3 * \text{ArcTanh}[(2*c + e*x)/(2*\text{Sqrt}[c]*\text{Sqrt}[c + x*(e + d*x)])])]/(48*c^{(5/2)}*x^3*(a + b*x^2))$

fricas [A] time = 2.05, size = 791, normalized size = 2.69

$$\frac{48bc^3\sqrt{d}x^3\log\left(8d^2x^2 + 8dex + 4\sqrt{dx^2 + ex + c}(2dx + e)\sqrt{d} + 4cd + e^2\right) + 3\left(ae^3 + 4(2bc^2 - acd)e\right)\sqrt{c}x^3}{96c^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x, algorithm="fricas")

[Out] [1/96*(48*b*c^3*sqrt(d)*x^3*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 3*(a*e^3 + 4*(2*b*c^2 - a*c*d)*e)*sqrt(c)*x^3*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) - 4*(2*a*c^2*e*x + 8*a*c^3 + (24*b*c^3 + 8*a*c^2*d - 3*a*c*e^2)*x^2)*sqrt(d*x^2 + e*x + c))/(c^3*x^3), -1/96*(96*b*c^3*sqrt(-d)*x^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) - 3*(a*e^3 + 4*(2*b*c^2 - a*c*d)*e)*sqrt(c)*x^3*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) + 4*(2*a*c^2*e*x + 8*a*c^3 + (24*b*c^3 + 8*a*c^2*d - 3*a*c*e^2)*x^2)*sqrt(d*x^2 + e*x + c))/(c^3*x^3), 1/48*(24*b*c^3*sqrt(d)*x^3*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 3*(a*e^3 + 4*(2*b*c^2 - a*c*d)*e)*sqrt(-c)*x^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - 2*(2*a*c^2*e*x + 8*a*c^3 + (24*b*c^3 + 8*a*c^2*d - 3*a*c*e^2)*x^2)*sqrt(d*x^2 + e*x + c))/(c^3*x^3), -1/48*(48*b*c^3*sqrt(-d)*x^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) - 3*(a*e^3 + 4*(2*b*c^2 - a*c*d)*e)*sqrt(-c)*x^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) + 2*(2*a*c^2*e*x + 8*a*c^3 + (24*b*c^3 + 8*a*c^2*d - 3*a*c*e^2)*x^2)*sqrt(d*x^2 + e*x + c))/(c^3*x^3)]

giac [B] time = 1.34, size = 720, normalized size = 2.45

$$-b\sqrt{d}\log\left(\left|2\left(\sqrt{d}x - \sqrt{dx^2 + xe + c}\right)\sqrt{d} + e\right|\right)\text{sgn}(bx^2 + a) + \frac{(8bc^2\text{esgn}(bx^2 + a) - 4ac\text{desgn}(bx^2 + a) + ae^3)}{8\sqrt{-c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x, algorithm="giac")

```
[Out] -b*sqrt(d)*log(abs(2*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))*sqrt(d) + e))*sgn(
b*x^2 + a) + 1/8*(8*b*c^2*e*sgn(b*x^2 + a) - 4*a*c*d*e*sgn(b*x^2 + a) + a*e
^3*sgn(b*x^2 + a))*arctan(-(sqrt(d)*x - sqrt(d*x^2 + x*e + c))/sqrt(-c))/(s
qrt(-c)*c^2) + 1/24*(24*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^5*b*c^2*sqrt(d)
*e*sgn(b*x^2 + a) + 12*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^5*a*c*d^(3/2)*e*
sgn(b*x^2 + a) + 48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^4*b*c^3*d*sgn(b*x^2
+ a) + 48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^4*a*c^2*d^2*sgn(b*x^2 + a) -
48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^3*b*c^3*sqrt(d)*e*sgn(b*x^2 + a) +
48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^3*a*c^2*d^(3/2)*e*sgn(b*x^2 + a) - 9
6*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^2*b*c^4*d*sgn(b*x^2 + a) - 3*(sqrt(d)
*x - sqrt(d*x^2 + x*e + c))^5*a*sqrt(d)*e^3*sgn(b*x^2 + a) + 24*(sqrt(d)*x
- sqrt(d*x^2 + x*e + c))*b*c^4*sqrt(d)*e*sgn(b*x^2 + a) + 36*(sqrt(d)*x - s
qrt(d*x^2 + x*e + c))*a*c^3*d^(3/2)*e*sgn(b*x^2 + a) + 48*b*c^5*d*sgn(b*x^2
+ a) + 16*a*c^4*d^2*sgn(b*x^2 + a) + 48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c)
)^2*a*c^2*d*e^2*sgn(b*x^2 + a) + 8*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^3*a*
c*sqrt(d)*e^3*sgn(b*x^2 + a) + 3*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))*a*c^2*
sqrt(d)*e^3*sgn(b*x^2 + a))/(((sqrt(d)*x - sqrt(d*x^2 + x*e + c))^2 - c)^3*
c^2*sqrt(d))
```

maple [A] time = 0.01, size = 412, normalized size = 1.40

$$\sqrt{(bx^2 + a)^2} \left(-12ac^3d^5ex^3 \ln\left(\frac{ex+2c+2\sqrt{dx^2+ex+c}\sqrt{c}}{x}\right) + 3a\sqrt{c}d^3e^3x^3 \ln\left(\frac{ex+2c+2\sqrt{dx^2+ex+c}\sqrt{c}}{x}\right) - 48bc^3d^2x^3 \ln\left(\frac{ex+2c+2\sqrt{dx^2+ex+c}\sqrt{c}}{x}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x)
```

```
[Out] -1/48*((b*x^2+a)^2)^(1/2)*(-12*d^(5/2)*c^(3/2)*ln((e*x+2*c+2*(d*x^2+e*x+c)^(1/2)*c^(1/2))/x)*x^3*a*e+24*d^(3/2)*c^(5/2)*ln((e*x+2*c+2*(d*x^2+e*x+c)^(1/2)*c^(1/2))/x)*x^3*b*e-6*(d*x^2+e*x+c)^(1/2)*d^(5/2)*x^4*a*e^2-48*(d*x^2+e*x+c)^(1/2)*d^(5/2)*x^4*b*c^2+12*(d*x^2+e*x+c)^(1/2)*d^(5/2)*x^3*a*c*e+3*d^(3/2)*c^(1/2)*ln((e*x+2*c+2*(d*x^2+e*x+c)^(1/2)*c^(1/2))/x)*x^3*a*e^3+6*(d*x^2+e*x+c)^(3/2)*d^(3/2)*x^2*a*e^2+48*(d*x^2+e*x+c)^(3/2)*d^(3/2)*x^2*b*c^2-6*(d*x^2+e*x+c)^(1/2)*d^(3/2)*x^3*a*e^3-48*(d*x^2+e*x+c)^(1/2)*d^(3/2)*x^3*b*c^2*e-12*(d*x^2+e*x+c)^(3/2)*d^(3/2)*x*a*c*e-48*ln(1/2*(2*d*x+e+2*(d*x^2+e*x+c)^(1/2)*d^(1/2))/d^(1/2))*x^3*b*c^3*d^2+16*(d*x^2+e*x+c)^(3/2)*d^(3/2)*a*c^2)/d^(3/2)/x^3/c^3/(b*x^2+a)
```

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e^2-4*c*d>0)', see `assume?` for more details)Is e^2-4*c*d positive, negative or zero?
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^4,x)
```

```
[Out] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^4, x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**4,x)
```

```
[Out] Timed out
```


Chapter 4

Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.0.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] :=
  If[ExpnType[result]<=ExpnType[optimal],
```

```

If[FreeQ[result,Complex] || Not[FreeQ[optimal,Complex]],
  If[LeafCount[result]<=2*LeafCount[optimal],
    "A",
    "B"],
  "C"],
If[FreeQ[result,Integrate] && FreeQ[result,Int],
  "C",
"F"]]

```

```
(* ::Text:: *)
```

```
(*The following summarizes the type number assigned an *)
```

```
(*expression based on the functions it involves*)
```

```
(*1 = rational function*)
```

```
(*2 = algebraic function*)
```

```
(*3 = elementary function*)
```

```
(*4 = special function*)
```

```
(*5 = hyperpergeometric function*)
```

```
(*6 = appell function*)
```

```
(*7 = rootsum function*)
```

```
(*8 = integrate function*)
```

```
(*9 = unknown function*)
```

```
ExpnType[expn_] :=
```

```
  If[AtomQ[expn],
```

```
    1,
```

```
  If[ListQ[expn],
```

```
    Max[Map[ExpnType,expn]],
```

```
  If[Head[expn]===Power,
```

```
    If[IntegerQ[expn[[2]]],
```

```
      ExpnType[expn[[1]],
```

```
    If[Head[expn[[2]]]===Rational,
```

```
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
```

```
        1,
```

```
        Max[ExpnType[expn[[1]],2]],
```

```
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
```

```
  If[Head[expn]===Plus || Head[expn]===Times,
```

```
    Max[ExpnType[First[expn],ExpnType[Rest[expn]]],
```

```
  If[ElementaryFunctionQ[Head[expn]],
```

```
    Max[3,ExpnType[expn[[1]]],
```

```
  If[SpecialFunctionQ[Head[expn]],
```

```
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
```

```
  If[HypergeometricFunctionQ[Head[expn]],
```

```
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
```

```
  If[AppellFunctionQ[Head[expn]],
```

```
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
```

```

If[Head[expn]===RootSum,
  Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
If[Head[expn]===Integrate || Head[expn]===Int,
  Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
9]]]]]]]]]]

ElementaryFunctionQ[func_] :=
MemberQ[{
  Exp,Log,
  Sin,Cos,Tan,Cot,Sec,Csc,
  ArcSin,ArcCos,ArcTan,ArcCot,ArcSec,ArcCsc,
  Sinh,Cosh,Tanh,Coth,Sech,Csch,
  ArcSinh,ArcCosh,ArcTanh,ArcCoth,ArcSech,ArcCsch
},func]

SpecialFunctionQ[func_] :=
MemberQ[{
  Erf, Erfc, Erfi,
  FresnelS, FresnelC,
  ExpIntegralE, ExpIntegralEi, LogIntegral,
  SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
  Gamma, LogGamma, PolyGamma,
  Zeta, PolyLog, ProductLog,
  EllipticF, EllipticE, EllipticPi
},func]

HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1,Hypergeometric2F1,HypergeometricPFQ},func]

AppellFunctionQ[func_] :=
MemberQ[{AppellF1},func]

```

4.0.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems

GradeAntiderivative := proc(result,optimal)
local leaf_count_result, leaf_count_optimal,ExpnType_result,ExpnType_optimal,
  debug:=false;

  leaf_count_result:=leafcount(result);
  #do NOT call ExpnType() if leaf size is too large. Recursion problem
  if leaf_count_result > 500000 then
    return "B";
  fi;

  leaf_count_optimal:=leafcount(optimal);

  ExpnType_result:=ExpnType(result);
  ExpnType_optimal:=ExpnType(optimal);

  if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",
ExpnType_optimal);
  fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
  return "F";
end if;

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;

```



```

if is_contains_complex(result) then
  if is_contains_complex(optimal) then
    if debug then
      print("both result and optimal complex");
    fi;
    #both result and optimal complex
    if leaf_count_result<=2*leaf_count_optimal then
      return "A";
    else
      return "B";
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C";
  end if
else # result do not contain complex
  # this assumes optimal do not as well
  if debug then
    print("result do not contain complex, this assumes optimal do not
as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B";
  end if
end if
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C";
end if

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false

```

```

#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+`') or type(expn,'`*`') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  end if
end proc:

```

```

elif HypergeometricFunctionQ(op(0,expn)) then
  max(5,apply(max,map(ExpnType,[op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6,apply(max,map(ExpnType,[op(expn)])))
elif op(0,expn)='int' then
  max(8,apply(max,map(ExpnType,[op(expn)]))) else
9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func,[AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

```

```
#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma][LeafCount](u);
end proc:
```

4.0.3 Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
        ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
        ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]
```

```

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)
))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,
ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+`') or type
(expn,'`*`')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))

```

```

elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,
expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,
Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        return "F"

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
    else:
        return "C"

```

4.0.4 SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#           Albert Rich to use with Sagemath. This is used to
#           grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#           'arctan2','floor','abs','log_integral'

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:

```

```

        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    debug=False
    if debug: print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','
sinh_integral'
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U
']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in
sagemath

def is_atom(expn):

    debug=False
    if debug: print ("Enter is_atom")

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type-in-maple/
    try:
        if expn.parent() is SR:

```



```

        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().
gens()
    return False

except AttributeError as error:
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],
Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #instance(expn.args[1],
Rational)
            if type(expn.operands()[0])==Rational: #instance(expn.args[0],
Rational)
                return 1
            else:
                return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
        else:
            return max(3,expnType(expn.operands()[0]),expnType(expn.operands()
[1])) #max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))
    elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #
instance(expn,Add) or instance(expn,Mul)
        m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
        m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))

```

```

    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.
func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #
is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
#is checked before calling the grading function that is passed.
#but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

```
#main function
```

```
def grade_antiderivative(result,optimal):
```

```

    if debug: print ("Enter grade_antiderivative for sagemath")

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    if debug: print ("leaf_count_result=", leaf_count_result, "
leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",
expnType_optimal)

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex

```

```
        if leaf_count_result <= 2*leaf_count_optimal:
            return "A"
        else:
            return "B"
    else: #result contains complex but optimal is not
        return "C"
else: # result do not contain complex, this assumes optimal do not as
well
    if leaf_count_result <= 2*leaf_count_optimal:
        return "A"
    else:
        return "B"
else:
    return "C"
```