

Computer algebra independent integration tests

4-Trig-functions/4.2-Cosine/4.2.0-a-cos-^m-b-trg-ⁿ

Nasser M. Abbasi

July 17, 2021

Compiled on July 17, 2021 at 6:09am

Contents

1	Introduction	3
1.1	Listing of CAS systems tested	3
1.2	Results	3
1.3	Performance	7
1.4	list of integrals that has no closed form antiderivative	8
1.5	list of integrals solved by CAS but has no known antiderivative	8
1.6	list of integrals solved by CAS but failed verification	8
1.7	Timing	9
1.8	Verification	9
1.9	Important notes about some of the results	9
1.9.1	Important note about Maxima results	9
1.9.2	Important note about FriCAS and Giac/XCAS results	10
1.9.3	Important note about finding leaf size of antiderivative	10
1.9.4	Important note about Mupad results	11
1.10	Design of the test system	11
2	detailed summary tables of results	13
2.1	List of integrals sorted by grade for each CAS	13
2.1.1	Rubi	13
2.1.2	Mathematica	13
2.1.3	Maple	14
2.1.4	Maxima	14
2.1.5	FriCAS	14
2.1.6	Sympy	15
2.1.7	Giac	15
2.1.8	Mupad	15
2.2	Detailed conclusion table per each integral for all CAS systems	17
2.3	Detailed conclusion table specific for Rubi results	66
3	Listing of integrals	75
3.1	$\int \cos(a + bx) dx$	75
3.2	$\int \cos^2(a + bx) dx$	77
3.3	$\int \cos^3(a + bx) dx$	79
3.4	$\int \cos^4(a + bx) dx$	81
3.5	$\int \cos^5(a + bx) dx$	83
3.6	$\int \cos^6(a + bx) dx$	85

3.7	$\int \cos^7(a + bx) dx$	88
3.8	$\int \cos^8(a + bx) dx$	90
3.9	$\int \cos^{\frac{7}{2}}(a + bx) dx$	93
3.10	$\int \cos^{\frac{5}{2}}(a + bx) dx$	96
3.11	$\int \cos^{\frac{3}{2}}(a + bx) dx$	99
3.12	$\int \sqrt{\cos(a + bx)} dx$	102
3.13	$\int \frac{1}{\sqrt{\cos(a+bx)}} dx$	104
3.14	$\int \frac{1}{\cos^{\frac{3}{2}}(a+bx)} dx$	106
3.15	$\int \frac{1}{\cos^{\frac{5}{2}}(a+bx)} dx$	109
3.16	$\int \frac{1}{\cos^{\frac{7}{2}}(a+bx)} dx$	112
3.17	$\int (c \cos(a + bx))^{7/2} dx$	115
3.18	$\int (c \cos(a + bx))^{5/2} dx$	118
3.19	$\int (c \cos(a + bx))^{3/2} dx$	121
3.20	$\int \sqrt{c \cos(a + bx)} dx$	124
3.21	$\int \frac{1}{\sqrt{c \cos(a+bx)}} dx$	126
3.22	$\int \frac{1}{(c \cos(a+bx))^{3/2}} dx$	128
3.23	$\int \frac{1}{(c \cos(a+bx))^{5/2}} dx$	131
3.24	$\int \frac{1}{(c \cos(a+bx))^{7/2}} dx$	134
3.25	$\int \cos^{\frac{4}{3}}(a + bx) dx$	137
3.26	$\int \cos^{\frac{2}{3}}(a + bx) dx$	139
3.27	$\int \sqrt[3]{\cos(a + bx)} dx$	141
3.28	$\int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx$	143
3.29	$\int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx$	145
3.30	$\int \frac{1}{\cos^{\frac{4}{3}}(a+bx)} dx$	147
3.31	$\int (c \cos(a + bx))^{4/3} dx$	149
3.32	$\int (c \cos(a + bx))^{2/3} dx$	151
3.33	$\int \sqrt[3]{c \cos(a + bx)} dx$	153
3.34	$\int \frac{1}{\sqrt[3]{c \cos(a+bx)}} dx$	155
3.35	$\int \frac{1}{(c \cos(a+bx))^{2/3}} dx$	157
3.36	$\int \frac{1}{(c \cos(a+bx))^{4/3}} dx$	159
3.37	$\int \cos^n(a + bx) dx$	161
3.38	$\int (c \cos(a + bx))^n dx$	163
3.39	$\int (a \cos^2(x))^{5/2} dx$	165
3.40	$\int (a \cos^2(x))^{3/2} dx$	168
3.41	$\int \sqrt{a \cos^2(x)} dx$	170
3.42	$\int \frac{1}{\sqrt{a \cos^2(x)}} dx$	172
3.43	$\int \frac{1}{(a \cos^2(x))^{3/2}} dx$	175
3.44	$\int \frac{1}{(a \cos^2(x))^{5/2}} dx$	178
3.45	$\int (a \cos^3(x))^{5/2} dx$	182

3.46	$\int (a \cos^3(x))^{3/2} dx$	185
3.47	$\int \sqrt{a \cos^3(x)} dx$	188
3.48	$\int \frac{1}{\sqrt{a \cos^3(x)}} dx$	191
3.49	$\int \frac{1}{(a \cos^3(x))^{3/2}} dx$	194
3.50	$\int \frac{1}{(a \cos^3(x))^{5/2}} dx$	197
3.51	$\int (a \cos^4(x))^{5/2} dx$	200
3.52	$\int (a \cos^4(x))^{3/2} dx$	203
3.53	$\int \sqrt{a \cos^4(x)} dx$	206
3.54	$\int \frac{1}{\sqrt{a \cos^4(x)}} dx$	208
3.55	$\int \frac{1}{(a \cos^4(x))^{3/2}} dx$	210
3.56	$\int \frac{1}{(a \cos^4(x))^{5/2}} dx$	213
3.57	$\int (b \cos^m(c + dx))^n dx$	216
3.58	$\int (c \cos^m(a + bx))^{5/2} dx$	218
3.59	$\int (c \cos^m(a + bx))^{3/2} dx$	220
3.60	$\int \sqrt{c \cos^m(a + bx)} dx$	222
3.61	$\int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx$	224
3.62	$\int \frac{1}{(c \cos^m(a + bx))^{3/2}} dx$	227
3.63	$\int \frac{1}{(c \cos^m(a + bx))^{5/2}} dx$	230
3.64	$\int (c \cos^m(a + bx))^{\frac{1}{m}} dx$	233
3.65	$\int (a(b \cos(c + dx))^p)^n dx$	236
3.66	$\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx$	238
3.67	$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx$	241
3.68	$\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx$	244
3.69	$\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx$	247
3.70	$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx$	250
3.71	$\int \sqrt{b \cos(c + dx)} dx$	253
3.72	$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx$	255
3.73	$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx$	258
3.74	$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx$	261
3.75	$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx$	264
3.76	$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx$	267
3.77	$\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx$	270
3.78	$\int \cos^4(c + dx) (b \cos(c + dx))^{3/2} dx$	273
3.79	$\int \cos^3(c + dx) (b \cos(c + dx))^{3/2} dx$	276
3.80	$\int \cos^2(c + dx) (b \cos(c + dx))^{3/2} dx$	279
3.81	$\int \cos(c + dx) (b \cos(c + dx))^{3/2} dx$	282
3.82	$\int (b \cos(c + dx))^{3/2} dx$	285
3.83	$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx$	288
3.84	$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx$	291
3.85	$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx$	294
3.86	$\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx$	297
3.87	$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx$	300
3.88	$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx$	303
3.89	$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx$	306

3.90	$\int \cos^3(c + dx)(b \cos(c + dx))^{5/2} dx$	309
3.91	$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx$	312
3.92	$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx$	315
3.93	$\int (b \cos(c + dx))^{5/2} dx$	318
3.94	$\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx$	321
3.95	$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx$	324
3.96	$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx$	327
3.97	$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx$	330
3.98	$\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx$	333
3.99	$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx$	336
3.100	$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx$	339
3.101	$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx$	342
3.102	$\int (b \cos(c + dx))^{7/2} dx$	345
3.103	$\int \frac{\cos^6(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	348
3.104	$\int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	351
3.105	$\int \frac{\cos^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	354
3.106	$\int \frac{\cos^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	357
3.107	$\int \frac{\cos^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	360
3.108	$\int \frac{\cos(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	363
3.109	$\int \frac{1}{\sqrt{b \cos(c+dx)}} dx$	366
3.110	$\int \frac{\sec(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	368
3.111	$\int \frac{\sec^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	371
3.112	$\int \frac{\sec^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	374
3.113	$\int \frac{\sec^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	377
3.114	$\int \frac{\sec^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	380
3.115	$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	383
3.116	$\int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	386
3.117	$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	389
3.118	$\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	392
3.119	$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	395
3.120	$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	398
3.121	$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	401
3.122	$\int \frac{1}{(b \cos(c+dx))^{3/2}} dx$	404
3.123	$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	407
3.124	$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	410
3.125	$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	413
3.126	$\int \frac{\sec^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	416
3.127	$\int \frac{\cos^8(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	419
3.128	$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	422

3.129	$\int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	425
3.130	$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	428
3.131	$\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	431
3.132	$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	434
3.133	$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	437
3.134	$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	440
3.135	$\int \frac{1}{(b \cos(c+dx))^{5/2}} dx$	443
3.136	$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	446
3.137	$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	449
3.138	$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	452
3.139	$\int \frac{1}{(b \cos(c+dx))^{7/2}} dx$	455
3.140	$\int \cos^2(c+dx) \sqrt{b \cos(c+dx)} dx$	458
3.141	$\int \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)} dx$	466
3.142	$\int \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)} dx$	468
3.143	$\int \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} dx$	473
3.144	$\int \frac{\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} dx$	475
3.145	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)} dx$	477
3.146	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{5}{2}}(c+dx)} dx$	479
3.147	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{7}{2}}(c+dx)} dx$	482
3.148	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{9}{2}}(c+dx)} dx$	485
3.149	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{11}{2}}(c+dx)} dx$	488
3.150	$\int \cos^{\frac{5}{2}}(c+dx) (b \cos(c+dx))^{3/2} dx$	492
3.151	$\int \cos^{\frac{3}{2}}(c+dx) (b \cos(c+dx))^{3/2} dx$	500
3.152	$\int \sqrt{\cos(c+dx)} (b \cos(c+dx))^{3/2} dx$	503
3.153	$\int \frac{(b \cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx$	508
3.154	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{3}{2}}(c+dx)} dx$	510
3.155	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{5}{2}}(c+dx)} dx$	512
3.156	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{7}{2}}(c+dx)} dx$	514
3.157	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{9}{2}}(c+dx)} dx$	517
3.158	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{11}{2}}(c+dx)} dx$	520
3.159	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{13}{2}}(c+dx)} dx$	523
3.160	$\int \cos^{\frac{5}{2}}(c+dx) (b \cos(c+dx))^{5/2} dx$	527
3.161	$\int \cos^{\frac{3}{2}}(c+dx) (b \cos(c+dx))^{5/2} dx$	530
3.162	$\int \sqrt{\cos(c+dx)} (b \cos(c+dx))^{5/2} dx$	539

3.163	$\int \frac{(b \cos(c+dx))^{5/2}}{\sqrt{\cos(c+dx)}} dx$	541
3.164	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^3(c+dx)} dx$	544
3.165	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^5(c+dx)} dx$	546
3.166	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^7(c+dx)} dx$	548
3.167	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^9(c+dx)} dx$	550
3.168	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{11}(c+dx)} dx$	553
3.169	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{13}(c+dx)} dx$	556
3.170	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{15}(c+dx)} dx$	559
3.171	$\int \frac{\cos^{11}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	563
3.172	$\int \frac{\cos^7(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	566
3.173	$\int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	569
3.174	$\int \frac{\cos^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	572
3.175	$\int \frac{\cos(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	575
3.176	$\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx$	577
3.177	$\int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} dx$	579
3.178	$\int \frac{1}{\cos^3(c+dx) \sqrt{b \cos(c+dx)}} dx$	581
3.179	$\int \frac{1}{\cos^5(c+dx) \sqrt{b \cos(c+dx)}} dx$	584
3.180	$\int \frac{1}{\cos^7(c+dx) \sqrt{b \cos(c+dx)}} dx$	587
3.181	$\int \frac{1}{\cos^9(c+dx) \sqrt{b \cos(c+dx)}} dx$	590
3.182	$\int \frac{\cos^{11}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	594
3.183	$\int \frac{\cos^9(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	597
3.184	$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	600
3.185	$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	603
3.186	$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	605
3.187	$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx$	607
3.188	$\int \frac{1}{\sqrt{\cos(c+dx)} (b \cos(c+dx))^{3/2}} dx$	609
3.189	$\int \frac{1}{\cos^3(c+dx) (b \cos(c+dx))^{3/2}} dx$	612
3.190	$\int \frac{1}{\cos^5(c+dx) (b \cos(c+dx))^{3/2}} dx$	615
3.191	$\int \frac{1}{\cos^7(c+dx) (b \cos(c+dx))^{3/2}} dx$	618

3.192	$\int \frac{\cos^{\frac{13}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	622
3.193	$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	625
3.194	$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	628
3.195	$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	631
3.196	$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	633
3.197	$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	635
3.198	$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx$	637
3.199	$\int \frac{1}{\sqrt{\cos(c+dx)}(b \cos(c+dx))^{5/2}} dx$	640
3.200	$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx$	643
3.201	$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx$	646
3.202	$\int \cos^m(c+dx) \sqrt[3]{b \cos(c+dx)} dx$	650
3.203	$\int \cos^2(c+dx) \sqrt[3]{b \cos(c+dx)} dx$	652
3.204	$\int \cos(c+dx) \sqrt[3]{b \cos(c+dx)} dx$	654
3.205	$\int \sqrt[3]{b \cos(c+dx)} dx$	656
3.206	$\int \sqrt[3]{b \cos(c+dx)} \sec(c+dx) dx$	658
3.207	$\int \sqrt[3]{b \cos(c+dx)} \sec^2(c+dx) dx$	660
3.208	$\int \sqrt[3]{b \cos(c+dx)} \sec^3(c+dx) dx$	662
3.209	$\int \cos^m(c+dx)(b \cos(c+dx))^{2/3} dx$	664
3.210	$\int \cos^2(c+dx)(b \cos(c+dx))^{2/3} dx$	666
3.211	$\int \cos(c+dx)(b \cos(c+dx))^{2/3} dx$	668
3.212	$\int (b \cos(c+dx))^{2/3} dx$	670
3.213	$\int (b \cos(c+dx))^{2/3} \sec(c+dx) dx$	672
3.214	$\int (b \cos(c+dx))^{2/3} \sec^2(c+dx) dx$	674
3.215	$\int (b \cos(c+dx))^{2/3} \sec^3(c+dx) dx$	676
3.216	$\int \cos^m(c+dx)(b \cos(c+dx))^{4/3} dx$	678
3.217	$\int \cos^2(c+dx)(b \cos(c+dx))^{4/3} dx$	680
3.218	$\int \cos(c+dx)(b \cos(c+dx))^{4/3} dx$	682
3.219	$\int (b \cos(c+dx))^{4/3} dx$	684
3.220	$\int (b \cos(c+dx))^{4/3} \sec(c+dx) dx$	686
3.221	$\int (b \cos(c+dx))^{4/3} \sec^2(c+dx) dx$	688
3.222	$\int (b \cos(c+dx))^{4/3} \sec^3(c+dx) dx$	690
3.223	$\int \frac{\cos^m(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	692
3.224	$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	695
3.225	$\int \frac{\cos(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	698
3.226	$\int \frac{1}{\sqrt[3]{b \cos(c+dx)}} dx$	700
3.227	$\int \frac{\sec(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	702
3.228	$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	704
3.229	$\int \frac{\sec^3(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	707
3.230	$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	710
3.231	$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	713

3.232	$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	716
3.233	$\int \frac{1}{(b \cos(c+dx))^{2/3}} dx$	718
3.234	$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	720
3.235	$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	722
3.236	$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	724
3.237	$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	726
3.238	$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	729
3.239	$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	731
3.240	$\int \frac{1}{(b \cos(c+dx))^{4/3}} dx$	733
3.241	$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	735
3.242	$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	737
3.243	$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	739
3.244	$\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx$	741
3.245	$\int \cos^2(c + dx) (b \cos(c + dx))^n dx$	743
3.246	$\int \cos(c + dx) (b \cos(c + dx))^n dx$	745
3.247	$\int (b \cos(c + dx))^n dx$	747
3.248	$\int (b \cos(c + dx))^n \sec(c + dx) dx$	749
3.249	$\int (b \cos(c + dx))^n \sec^2(c + dx) dx$	751
3.250	$\int (b \cos(c + dx))^n \sec^3(c + dx) dx$	753
3.251	$\int (b \cos(c + dx))^n \sec^4(c + dx) dx$	755
3.252	$\int \cos^{\frac{5}{2}}(c + dx) (b \cos(c + dx))^n dx$	757
3.253	$\int \cos^{\frac{3}{2}}(c + dx) (b \cos(c + dx))^n dx$	759
3.254	$\int \sqrt{\cos(c + dx)} (b \cos(c + dx))^n dx$	761
3.255	$\int \frac{(b \cos(c+dx))^n}{\sqrt{\cos(c+dx)}} dx$	763
3.256	$\int \frac{(b \cos(c+dx))^n}{\sqrt[3]{\cos(c+dx)}} dx$	766
3.257	$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^n} dx$	769
3.258	$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^n} dx$	772
3.259	$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^n} dx$	775
3.260	$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx$	778
3.261	$\int \cos(a + bx) \sqrt{\csc(a + bx)} dx$	780
3.262	$\int \frac{\cos(a+bx)}{\sqrt{\csc(a+bx)}} dx$	782
3.263	$\int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx$	784
3.264	$\int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$	787
3.265	$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx$	790
3.266	$\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx$	792
3.267	$\int \frac{\cos^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$	794
3.268	$\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx$	796
3.269	$\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$	799
3.270	$\int \cos(x) \csc^{\frac{7}{3}}(x) dx$	802
3.271	$\int \sqrt{\csc(a + bx)} \sec(a + bx) dx$	804

3.272	$\int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx$	807
3.273	$\int \sqrt{\csc(a+bx)} \sec^2(a+bx) dx$	810
3.274	$\int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$	813
3.275	$\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx$	816
3.276	$\int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$	819
3.277	$\int \sqrt{\csc(a+bx)} \sec^4(a+bx) dx$	822
3.278	$\int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$	825
3.279	$\int (d \cos(a+bx))^{3/2} \csc^p(a+bx) dx$	828
3.280	$\int \sqrt{d \cos(a+bx)} \csc^p(a+bx) dx$	830
3.281	$\int \frac{\csc^p(a+bx)}{\sqrt{d \cos(a+bx)}} dx$	832
3.282	$\int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{3/2}} dx$	834
3.283	$\int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{5/2}} dx$	837
3.284	$\int \cos^m(e+fx) \csc^n(e+fx) dx$	839
3.285	$\int (a \cos(e+fx))^m \csc^n(e+fx) dx$	842
3.286	$\int \cos^m(e+fx) (b \csc(e+fx))^n dx$	845
3.287	$\int (a \cos(e+fx))^m (b \csc(e+fx))^n dx$	848
3.288	$\int (a \cos(e+fx))^m (b \csc(e+fx))^{7/2} dx$	851
3.289	$\int (a \cos(e+fx))^m (b \csc(e+fx))^{5/2} dx$	853
3.290	$\int (a \cos(e+fx))^m (b \csc(e+fx))^{3/2} dx$	855
3.291	$\int (a \cos(e+fx))^m \sqrt{b \csc(e+fx)} dx$	857
3.292	$\int \frac{(a \cos(e+fx))^m}{\sqrt{b \csc(e+fx)}} dx$	859
3.293	$\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{3/2}} dx$	862
3.294	$\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{5/2}} dx$	865

4	Listing of Grading functions	869
4.0.1	Mathematica and Rubi grading function	869
4.0.2	Maple grading function	871
4.0.3	Sympy grading function	874
4.0.4	SageMath grading function	876

Chapter 1

Introduction

This report gives the result of running the computer algebra independent integration problems. The listing of the problems are maintained by and can be downloaded from <https://rulebasedintegration.org>

The number of integrals in this report is [294]. This is test number [82].

1.1 Listing of CAS systems tested

The following systems were tested at this time.

1. Mathematica 12.3 (64 bit) on windows 10.
2. Rubi 4.16.1 in Mathematica 12.1 on windows 10.
3. Maple 2021.1 (64 bit) on windows 10.
4. Maxima 5.44 on Linux. (via sagemath 9.3)
5. Fricas 1.3.7 on Linux (via sagemath 9.3)
6. Giac/Xcas 1.7 on Linux. (via sagemath 9.3)
7. Sympy 1.8 under Python 3.8.8 using Anaconda distribution on Ubuntu.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 under windows 10 (64 bit)

Maxima, Fricas and Giac/Xcas were called from inside SageMath. This was done using SageMath integrate command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly using Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric $2F1$ functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	solved	Failed
Rubi	% 100.00 (294)	% 0.00 (0)
Mathematica	% 100.00 (294)	% 0.00 (0)
Maple	% 66.67 (196)	% 33.33 (98)
Maxima	% 31.29 (92)	% 68.71 (202)
Fricas	% 31.63 (93)	% 68.37 (201)
Sympy	% 5.10 (15)	% 94.90 (279)
Giac	% 8.84 (26)	% 91.16 (268)
Mupad	% 27.21 (80)	% 72.79 (214)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	96.94	1.02	2.04	0.00
Maple	41.50	22.11	3.06	33.33
Maxima	21.09	10.20	0.00	68.71
Fricas	29.93	1.70	0.00	68.37
Sympy	5.10	0.00	0.00	94.90
Giac	8.16	0.68	0.00	91.16
Mupad	0.00	27.21	0.00	72.79

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input

within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	0	0.00 %	0.00 %	0.00 %
Maple	98	100.00 %	0.00 %	0.00 %
Maxima	202	99.01 %	0.50 %	0.50 %
Fricas	201	97.01 %	0.00 %	2.99 %
Sympy	279	41.58 %	58.42 %	0.00 %
Giac	268	95.90 %	1.87 %	2.24 %
Mupad	214	100.00 %	0.00 %	0.00 %

Table 1.4: Time and leaf size performance for each CAS


1.3 Performance


The table below summarizes the performance of each CAS system in terms of CPU time and leaf size of results.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.04	68.33	1.00	69.00	1.00
Mathematica	0.18	59.70	0.91	55.50	0.91
Maple	0.14	140.38	1.98	121.00	1.66
Maxima	1.11	221.09	2.81	48.50	0.86
Fricas	0.66	88.82	1.62	47.00	0.97
Sympy	19.47	52.07	1.29	36.00	1.41
Giac	0.66	40.77	1.19	25.00	0.70
Mupad	0.54	54.98	1.09	43.50	0.87

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used columns from the above table.





1.4 list of integrals that has no closed form antiderivative

{}

1.5 list of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {284, 285, 286, 287, 292}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of _int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call has completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out is not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica. Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative produced was correct.

Verification phase has 3 minutes time out. An integral whose result was not verified could still be correct. Further investigation is needed on those integrals which failed verifications. Such integrals are marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since these integrals are run in a batch mode, using an automated script, and by using `sagemath` (SageMath uses Maxima), then any integral where Maxima needs an interactive response from the user to answer a question during evaluation of the integral in order to complete the integration, will fail and is counted as failed.

The exception raised is `ValueError`. Therefore Maxima result below is lower than what could result if Maxima was run directly and each question Maxima asks was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate of the error is due to the interactive question being asked or not.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
```

```
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath loading of Maxima `abs_integrate` was found to cause some problem. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS and Giac/X-CAS results

There are Few integrals which failed due to SageMath not able to translate the result back to SageMath syntax and not because these CAS system were not able to do the integrations.

These will fail With error `Exception raised: NotImplementedError`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals.

Hopefully the next version of SageMath will have complete translation of FriCAS and XCAS syntax and I will re-run all the tests again when this happens.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi and Maple, the builtin system function `LeafSize` is used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy and Giac antiderivatives is determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which is called directly from Python, the following code is used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative, Maple was used to determine the leaf size of Mupad output by post processing.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
- The following field present only in Rubi and Mathematica Tables*
13. integer. 1 if result was verified or 0 if not verified.
- The following fields present only in Rubi Tables*
14. integer. Number of rules used.
15. integer. Integrand leaf size.
16. real number. Ratio of field 14 over field 15
17. integer. 1 if result was verified or 0 if not verified.
18. String of form "{n,n,..}" which is list of the rules used by Rubi

High level overview of the CAS independent integration test build system

Chapter 2

detailed summary tables of results

2.1 List of integrals sorted by grade for each CAS

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 277, 278, 279, 280, 281, 282, 283, 288, 289, 290, 291, 293, 294 }

B grade: { 1, 42, 43 }

C grade: { 276, 284, 285, 286, 287, 292 }

F grade: { }

2.1.3 Maple

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 14, 17, 22, 39, 40, 41, 44, 51, 52, 53, 54, 55, 56, 66, 68, 73, 78, 80, 85, 90, 92, 97, 102, 103, 104, 105, 110, 115, 116, 117, 122, 127, 128, 129, 134, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 275, 276, 277, 278 }

B grade: { 9, 10, 11, 12, 15, 16, 18, 19, 20, 23, 24, 42, 43, 67, 69, 70, 71, 72, 74, 75, 76, 77, 79, 81, 82, 83, 84, 86, 87, 88, 89, 91, 93, 94, 95, 96, 98, 99, 100, 101, 106, 107, 108, 111, 112, 113, 114, 118, 119, 120, 121, 123, 124, 125, 126, 130, 131, 132, 133, 135, 136, 137, 138, 139, 274 }

C grade: { 13, 21, 45, 46, 47, 48, 49, 50, 109 }

F grade: { 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 57, 58, 59, 60, 61, 62, 63, 64, 65, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

2.1.4 Maxima

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 39, 40, 41, 51, 52, 53, 54, 55, 56, 140, 141, 142, 143, 144, 146, 150, 151, 152, 153, 154, 156, 160, 161, 162, 163, 164, 165, 167, 171, 172, 173, 174, 175, 176, 182, 183, 184, 185, 186, 192, 193, 194, 195, 196, 261, 262, 265, 266, 267, 270, 271, 272, 275, 276 }

B grade: { 42, 43, 44, 145, 147, 148, 149, 155, 157, 158, 159, 166, 168, 169, 170, 177, 178, 179, 180, 181, 187, 188, 189, 190, 191, 197, 198, 199, 200, 201 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 45, 46, 47, 48, 49, 50, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 263, 264, 268, 269, 273, 274, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

2.1.5 FriCAS

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 39, 40, 41, 43, 44, 51, 52, 53, 54, 55, 56, 64, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 261, 262, 265, 266, 267, 270 }

B grade: { 42, 271, 272, 275, 276 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 45, 46, 47, 48, 49, 50, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 263, 264, 268, 269, 273, 274, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

2.1.6 Sympy

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 41, 64, 143, 144, 154, 176, 186 }

B grade: { }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 145, 146, 147, 148, 149, 150, 151, 152, 153, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 177, 178, 179, 180, 181, 182, 183, 184, 185, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

2.1.7 Giac

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 39, 40, 41, 43, 51, 52, 53, 54, 56, 143, 261, 262, 265, 266, 267, 270 }

B grade: { 44, 64 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 42, 45, 46, 47, 48, 49, 50, 55, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 263, 264, 268, 269, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

2.1.8 Mupad

A grade: { }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 21, 25, 26, 27, 28, 29, 30, 37, 41, 54, 55, 56, 64, 107, 108, 109, 140, 141, 142, 143, 144, 146, 148, 150, 151, 152, 153, 154, 156, 158, 160, 161, 162, 163, 164, 165, 167, 169, 171, 172, 173, 174, 175, 176, 178, 180, 182, 183, 184, 185, 186, 188, 190, 192, 193, 194, 195, 196, 198, 200, 261, 262, 265, 270 }

C grade: { }

F grade: { 17, 18, 19, 20, 22, 23, 24, 31, 32, 33, 34, 35, 36, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 145, 147, 149, 155, 157, 159, 166, 168, 170, 177, 179, 181, 187, 189, 191, 197, 199, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258,

259, 260, 263, 264, 266, 267, 268, 269, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$

Problem 1	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	21	11	10	10	12	10	10
normalized size	1	1.00	2.10	1.10	1.00	1.00	1.20	1.00	1.00
time (sec)	N/A	0.004	0.081	0.053	0.469	0.402	0.115	0.163	0.249
Problem 2	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	23	27	22	22	46	18	18
normalized size	1	1.00	0.92	1.08	0.88	0.88	1.84	0.72	0.72
time (sec)	N/A	0.009	0.023	0.047	0.301	0.428	0.191	0.191	0.228
Problem 3	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	22	22	21	36	22	24
normalized size	1	1.00	1.00	0.85	0.85	0.81	1.38	0.85	0.92
time (sec)	N/A	0.011	0.008	0.117	0.298	0.421	0.410	0.159	0.071
Problem 4	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	33	38	33	36	95	32	31
normalized size	1	1.00	0.72	0.83	0.72	0.78	2.07	0.70	0.67
time (sec)	N/A	0.021	0.037	0.080	0.485	0.413	0.859	0.156	0.188
Problem 5	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	32	34	33	58	34	31
normalized size	1	1.00	1.00	0.78	0.83	0.80	1.41	0.83	0.76
time (sec)	N/A	0.013	0.013	0.099	0.308	0.399	1.607	0.239	0.095

Problem 6	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	43	48	48	46	139	46	42
normalized size	1	1.00	0.64	0.72	0.72	0.69	2.07	0.69	0.63
time (sec)	N/A	0.033	0.035	0.095	0.330	0.423	3.113	0.190	0.222
Problem 7	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	54	42	44	43	78	44	43
normalized size	1	1.00	1.00	0.78	0.81	0.80	1.44	0.81	0.80
time (sec)	N/A	0.016	0.012	0.096	0.691	0.412	5.154	0.208	0.090
Problem 8	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	55	58	59	56	184	60	53
normalized size	1	1.00	0.62	0.66	0.67	0.64	2.09	0.68	0.60
time (sec)	N/A	0.047	0.051	0.102	0.316	0.405	8.671	0.210	0.279
Problem 9	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	51	199	0	0	0	0	42
normalized size	1	1.00	0.78	3.06	0.00	0.00	0.00	0.00	0.65
time (sec)	N/A	0.032	0.265	0.195	0.000	0.414	0.000	0.000	0.294
Problem 10	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	40	202	0	0	0	0	42
normalized size	1	1.00	0.95	4.81	0.00	0.00	0.00	0.00	1.00
time (sec)	N/A	0.018	0.045	0.076	0.000	0.405	0.000	0.000	0.166
Problem 11	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	36	179	0	0	0	0	35
normalized size	1	1.00	0.86	4.26	0.00	0.00	0.00	0.00	0.83
time (sec)	N/A	0.018	0.039	0.107	0.000	0.413	0.000	0.000	0.078

Problem 12	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	133	0	0	0	0	15
normalized size	1	1.00	1.00	8.31	0.00	0.00	0.00	0.00	0.94
time (sec)	N/A	0.009	0.025	0.060	0.000	0.418	0.000	0.000	0.116
Problem 13	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	18	0	0	0	0	15
normalized size	1	1.00	1.00	1.12	0.00	0.00	0.00	0.00	0.94
time (sec)	N/A	0.009	0.028	0.127	0.000	0.391	0.000	0.000	0.097
Problem 14	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	101	0	0	0	0	42
normalized size	1	1.00	1.00	2.66	0.00	0.00	0.00	0.00	1.11
time (sec)	N/A	0.018	0.060	0.085	0.000	0.421	0.000	0.000	0.246
Problem 15	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	36	213	0	0	0	0	42
normalized size	1	1.00	0.86	5.07	0.00	0.00	0.00	0.00	1.00
time (sec)	N/A	0.018	0.058	0.116	0.000	0.411	0.000	0.000	0.265
Problem 16	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	59	358	0	0	0	0	42
normalized size	1	1.00	0.91	5.51	0.00	0.00	0.00	0.00	0.65
time (sec)	N/A	0.029	0.093	0.134	0.000	0.405	0.000	0.000	0.309
Problem 17	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	76	210	0	0	0	0	-1
normalized size	1	1.00	0.78	2.14	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.090	0.127	0.000	0.414	0.000	0.000	0.000

Problem 18	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	62	213	0	0	0	0	-1
normalized size	1	1.00	0.89	3.04	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.036	0.084	0.072	0.000	0.416	0.000	0.000	0.000
Problem 19	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	58	190	0	0	0	0	-1
normalized size	1	1.00	0.83	2.71	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.043	0.075	0.000	0.453	0.000	0.000	0.000
Problem 20	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	142	0	0	0	0	-1
normalized size	1	1.00	1.00	3.74	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.022	0.018	0.069	0.000	0.467	0.000	0.000	0.000
Problem 21	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	54	0	0	0	0	33
normalized size	1	1.00	1.00	1.42	0.00	0.00	0.00	0.00	0.87
time (sec)	N/A	0.023	0.025	0.054	0.000	0.486	0.000	0.000	0.146
Problem 22	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	50	168	0	0	0	0	-1
normalized size	1	1.00	0.74	2.47	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.036	0.028	0.111	0.000	0.520	0.000	0.000	0.000
Problem 23	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	51	241	0	0	0	0	-1
normalized size	1	1.00	0.71	3.35	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.035	0.059	0.208	0.000	0.547	0.000	0.000	0.000

Problem 24	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	68	366	0	0	0	0	-1
normalized size	1	1.00	0.68	3.66	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.091	0.195	0.000	0.495	0.000	0.000	0.000
Problem 25	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	42
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.79
time (sec)	N/A	0.013	0.061	0.184	0.000	0.415	0.000	0.000	0.193
Problem 26	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	42
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.79
time (sec)	N/A	0.012	0.030	0.227	0.000	0.424	0.000	0.000	0.191
Problem 27	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	42
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.79
time (sec)	N/A	0.012	0.028	0.120	0.000	0.480	0.000	0.000	0.182
Problem 28	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	42
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.79
time (sec)	N/A	0.012	0.025	0.088	0.000	0.411	0.000	0.000	0.210
Problem 29	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	51	0	0	0	0	0	42
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.82
time (sec)	N/A	0.012	0.024	0.091	0.000	0.414	0.000	0.000	0.203

Problem 30	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	51	0	0	0	0	0	42
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.82
time (sec)	N/A	0.012	0.025	0.050	0.000	0.422	0.000	0.000	0.227

Problem 31	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.014	0.048	0.059	0.000	0.409	0.000	0.000	0.000

Problem 32	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.015	0.039	0.139	0.000	0.394	0.000	0.000	0.000

Problem 33	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.015	0.035	0.128	0.000	0.406	0.000	0.000	0.000

Problem 34	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.015	0.039	0.105	0.000	0.398	0.000	0.000	0.000

Problem 35	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	53	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.015	0.039	0.088	0.000	0.403	0.000	0.000	0.000

Problem 36	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	53	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.039	0.058	0.000	0.392	0.000	0.000	0.000
Problem 37	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	64	0	0	0	0	0	57
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.89
time (sec)	N/A	0.017	0.044	0.388	0.000	0.414	0.000	0.000	0.535
Problem 38	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	64	0	0	0	0	0	-1
normalized size	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.020	0.041	0.313	0.000	0.434	0.000	0.000	0.000
Problem 39	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	36	32	31	40	0	34	-1
normalized size	1	1.00	0.68	0.60	0.58	0.75	0.00	0.64	-0.02
time (sec)	N/A	0.039	0.018	0.090	0.683	0.417	0.000	0.536	0.000
Problem 40	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	26	24	17	26	0	17	-1
normalized size	1	1.00	0.76	0.71	0.50	0.76	0.00	0.50	-0.03
time (sec)	N/A	0.027	0.009	0.082	0.893	0.418	0.000	0.294	0.000
Problem 41	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	15	6	15	19	9	46
normalized size	1	1.00	1.00	1.15	0.46	1.15	1.46	0.69	3.54
time (sec)	N/A	0.011	0.004	0.055	0.676	0.407	0.543	0.393	0.214

Problem 42	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	B	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	46	48	38	65	0	0	-1
normalized size	1	1.00	2.88	3.00	2.38	4.06	0.00	0.00	-0.06
time (sec)	N/A	0.014	0.025	0.082	0.678	0.452	0.000	0.000	0.000
Problem 43	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	91	70	304	40	0	47	-1
normalized size	1	1.00	2.17	1.67	7.24	0.95	0.00	1.12	-0.02
time (sec)	N/A	0.022	0.059	0.088	0.676	0.418	0.000	0.415	0.000
Problem 44	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	72	89	933	49	0	126	-1
normalized size	1	1.00	1.18	1.46	15.30	0.80	0.00	2.07	-0.02
time (sec)	N/A	0.044	0.139	0.095	1.375	0.473	0.000	0.562	0.000
Problem 45	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	61	114	0	0	0	0	-1
normalized size	1	1.00	0.52	0.97	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.066	0.114	0.387	0.000	0.609	0.000	0.000	0.000
Problem 46	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	50	198	0	0	0	0	-1
normalized size	1	1.00	0.75	2.96	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.071	0.159	0.000	0.546	0.000	0.000	0.000
Problem 47	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	37	76	0	0	0	0	-1
normalized size	1	1.00	0.84	1.73	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.024	0.180	0.000	0.645	0.000	0.000	0.000

Problem 48	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	31	191	0	0	0	0	-1
normalized size	1	1.00	0.74	4.55	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	0.019	0.306	0.000	0.629	0.000	0.000	0.000
Problem 49	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	44	87	0	0	0	0	-1
normalized size	1	1.00	0.62	1.23	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.030	0.057	0.309	0.000	0.470	0.000	0.000	0.000
Problem 50	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	57	223	0	0	0	0	-1
normalized size	1	1.00	0.49	1.91	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.097	0.379	0.000	0.920	0.000	0.000	0.000
Problem 51	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	53	57	85	68	0	57	-1
normalized size	1	1.00	0.40	0.43	0.64	0.52	0.00	0.43	-0.01
time (sec)	N/A	0.051	0.116	0.412	0.903	0.529	0.000	0.596	0.000
Problem 52	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	38	41	55	42	0	25	-1
normalized size	1	1.00	0.49	0.53	0.71	0.54	0.00	0.32	-0.01
time (sec)	N/A	0.032	0.068	0.171	0.669	0.517	0.000	0.549	0.000
Problem 53	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	25	22	22	21	0	13	-1
normalized size	1	1.00	0.69	0.61	0.61	0.58	0.00	0.36	-0.03
time (sec)	N/A	0.015	0.014	0.091	1.073	0.505	0.000	0.270	0.000

Problem 54	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	6	18	0	6	6
normalized size	1	1.00	1.00	0.93	0.40	1.20	0.00	0.40	0.40
time (sec)	N/A	0.014	0.005	0.082	0.997	0.486	0.000	0.282	0.231
Problem 55	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	30	29	22	33	0	0	36
normalized size	1	1.00	0.45	0.43	0.33	0.49	0.00	0.00	0.54
time (sec)	N/A	0.021	0.026	0.097	0.870	0.729	0.000	0.000	0.541
Problem 56	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	47	41	34	45	0	34	306
normalized size	1	1.00	0.40	0.35	0.29	0.38	0.00	0.29	2.62
time (sec)	N/A	0.032	0.045	0.149	0.520	0.505	0.000	0.526	3.744
Problem 57	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	72	0	0	0	0	0	-1
normalized size	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.035	0.064	0.191	0.000	0.451	0.000	0.000	0.000
Problem 58	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F(-2)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	74	0	0	0	0	0	-1
normalized size	1	1.00	0.83	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.158	0.413	0.000	0.000	0.000	0.000	0.000
Problem 59	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F(-2)	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	72	0	0	0	0	0	-1
normalized size	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.113	0.096	0.000	0.000	0.000	0.000	0.000

Problem 60	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	68	0	0	0	0	0	-1
normalized size	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.068	0.081	0.000	0.000	0.000	0.000	0.000
Problem 61	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	72	0	0	0	0	0	-1
normalized size	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.047	0.073	0.085	0.000	0.000	0.000	0.000	0.000
Problem 62	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	72	0	0	0	0	0	-1
normalized size	1	1.00	0.81	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.047	0.103	0.068	0.000	0.000	0.000	0.000	0.000
Problem 63	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	74	0	0	0	0	0	-1
normalized size	1	1.00	0.83	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.045	0.103	0.073	0.000	0.000	0.000	0.000	0.000
Problem 64	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	0	0	15	65	300	40
normalized size	1	1.00	1.00	0.00	0.00	0.62	2.71	12.50	1.67
time (sec)	N/A	0.020	0.026	0.143	0.000	0.536	1.582	5.144	0.404
Problem 65	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	74	0	0	0	0	0	-1
normalized size	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.053	0.202	0.000	0.515	0.000	0.000	0.000

Problem 66	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	83	234	0	0	0	0	-1
normalized size	1	1.00	0.67	1.90	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.090	0.196	0.156	0.000	0.486	0.000	0.000	0.000
Problem 67	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	75	221	0	0	0	0	-1
normalized size	1	1.00	0.77	2.28	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.069	0.145	0.129	0.000	0.661	0.000	0.000	0.000
Problem 68	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	73	208	0	0	0	0	-1
normalized size	1	1.00	0.77	2.19	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.066	0.137	0.124	0.000	0.638	0.000	0.000	0.000
Problem 69	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	62	211	0	0	0	0	-1
normalized size	1	1.00	0.90	3.06	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.060	0.113	0.000	0.561	0.000	0.000	0.000
Problem 70	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	61	188	0	0	0	0	-1
normalized size	1	1.00	0.91	2.81	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.058	0.108	0.000	0.566	0.000	0.000	0.000
Problem 71	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	142	0	0	0	0	-1
normalized size	1	1.00	1.00	3.74	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.020	0.022	0.089	0.000	0.832	0.000	0.000	0.000

Problem 72	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	39	142	0	0	0	0	-1
normalized size	1	1.00	1.00	3.64	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.029	0.029	0.105	0.000	0.476	0.000	0.000	0.000
Problem 73	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	48	166	0	0	0	0	-1
normalized size	1	1.00	0.76	2.63	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.055	0.054	0.145	0.000	0.487	0.000	0.000	0.000
Problem 74	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	49	239	0	0	0	0	-1
normalized size	1	1.00	0.70	3.41	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.086	0.138	0.000	0.617	0.000	0.000	0.000
Problem 75	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	69	363	0	0	0	0	-1
normalized size	1	1.00	0.73	3.82	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.220	0.197	0.000	0.545	0.000	0.000	0.000
Problem 76	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	69	396	0	0	0	0	-1
normalized size	1	1.00	0.70	4.04	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.180	0.162	0.000	0.828	0.000	0.000	0.000
Problem 77	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	79	412	0	0	0	0	-1
normalized size	1	1.00	0.64	3.35	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.097	0.204	0.196	0.000	0.628	0.000	0.000	0.000

Problem 78	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	83	236	0	0	0	0	-1
normalized size	1	1.00	0.66	1.87	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.086	0.098	0.125	0.000	0.456	0.000	0.000	0.000
Problem 79	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	75	223	0	0	0	0	-1
normalized size	1	1.00	0.79	2.35	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.060	0.099	0.119	0.000	1.049	0.000	0.000	0.000
Problem 80	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	73	210	0	0	0	0	-1
normalized size	1	1.00	0.74	2.14	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.061	0.097	0.125	0.000	0.605	0.000	0.000	0.000
Problem 81	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	65	213	0	0	0	0	-1
normalized size	1	1.00	0.97	3.18	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.084	0.122	0.000	0.546	0.000	0.000	0.000
Problem 82	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	58	190	0	0	0	0	-1
normalized size	1	1.00	0.83	2.71	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.015	0.119	0.000	0.522	0.000	0.000	0.000
Problem 83	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	39	144	0	0	0	0	-1
normalized size	1	1.00	1.00	3.69	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.029	0.014	0.104	0.000	0.492	0.000	0.000	0.000

Problem 84	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	0	0	0	-1
normalized size	1	1.00	1.00	3.51	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.038	0.020	0.115	0.000	0.663	0.000	0.000	0.000
Problem 85	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	50	168	0	0	0	0	-1
normalized size	1	1.00	0.76	2.55	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.054	0.053	0.132	0.000	0.580	0.000	0.000	0.000
Problem 86	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	51	241	0	0	0	0	-1
normalized size	1	1.00	0.71	3.35	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.064	0.131	0.000	0.518	0.000	0.000	0.000
Problem 87	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	69	364	0	0	0	0	-1
normalized size	1	1.00	0.70	3.71	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.074	0.113	0.182	0.000	0.665	0.000	0.000	0.000
Problem 88	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	69	398	0	0	0	0	-1
normalized size	1	1.00	0.69	3.98	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.124	0.151	0.000	0.613	0.000	0.000	0.000
Problem 89	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	79	414	0	0	0	0	-1
normalized size	1	1.00	0.63	3.29	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.095	0.173	0.196	0.000	0.485	0.000	0.000	0.000

Problem 90	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	125	83	236	0	0	0	0	-1
normalized size	1	1.00	0.66	1.89	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.080	0.133	0.143	0.000	0.686	0.000	0.000	0.000
Problem 91	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	75	223	0	0	0	0	-1
normalized size	1	1.00	0.77	2.28	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.100	0.128	0.000	0.534	0.000	0.000	0.000
Problem 92	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	76	210	0	0	0	0	-1
normalized size	1	1.00	0.78	2.16	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.093	0.104	0.000	0.514	0.000	0.000	0.000
Problem 93	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	62	213	0	0	0	0	-1
normalized size	1	1.00	0.89	3.04	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.015	0.126	0.000	0.498	0.000	0.000	0.000
Problem 94	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	59	190	0	0	0	0	-1
normalized size	1	1.00	0.82	2.64	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.048	0.022	0.135	0.000	0.732	0.000	0.000	0.000
Problem 95	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	0	0	0	-1
normalized size	1	1.00	1.00	3.51	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.039	0.014	0.096	0.000	0.597	0.000	0.000	0.000

Problem 96	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	38	144	0	0	0	0	-1
normalized size	1	1.00	0.93	3.51	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.038	0.044	0.132	0.000	0.569	0.000	0.000	0.000
Problem 97	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	50	168	0	0	0	0	-1
normalized size	1	1.00	0.74	2.47	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.055	0.055	0.128	0.000	0.877	0.000	0.000	0.000
Problem 98	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	51	241	0	0	0	0	-1
normalized size	1	1.00	0.71	3.35	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.064	0.139	0.000	0.475	0.000	0.000	0.000
Problem 99	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	69	366	0	0	0	0	-1
normalized size	1	1.00	0.69	3.66	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.151	0.205	0.000	0.687	0.000	0.000	0.000
Problem 100	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	69	398	0	0	0	0	-1
normalized size	1	1.00	0.69	3.98	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.074	0.177	0.150	0.000	0.522	0.000	0.000	0.000
Problem 101	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	79	414	0	0	0	0	-1
normalized size	1	1.00	0.62	3.23	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.095	0.241	0.231	0.000	0.641	0.000	0.000	0.000

Problem 102	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	76	210	0	0	0	0	-1
normalized size	1	1.00	0.78	2.14	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.018	0.136	0.000	0.646	0.000	0.000	0.000
Problem 103	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	125	73	233	0	0	0	0	-1
normalized size	1	1.00	0.58	1.86	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.091	0.132	0.150	0.000	0.542	0.000	0.000	0.000
Problem 104	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	71	220	0	0	0	0	-1
normalized size	1	1.00	0.71	2.20	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.116	0.143	0.000	0.606	0.000	0.000	0.000
Problem 105	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	63	207	0	0	0	0	-1
normalized size	1	1.00	0.65	2.13	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.102	0.152	0.000	0.565	0.000	0.000	0.000
Problem 106	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	58	210	0	0	0	0	-1
normalized size	1	1.00	0.81	2.92	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.056	0.146	0.000	0.931	0.000	0.000	0.000
Problem 107	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	51	187	0	0	0	0	58
normalized size	1	1.00	0.74	2.71	0.00	0.00	0.00	0.00	0.84
time (sec)	N/A	0.039	0.046	0.169	0.000	0.730	0.000	0.000	0.140

Problem 108	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	141	0	0	0	0	33
normalized size	1	1.00	1.00	3.44	0.00	0.00	0.00	0.00	0.80
time (sec)	N/A	0.023	0.016	0.127	0.000	0.827	0.000	0.000	0.148
Problem 109	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	54	0	0	0	0	33
normalized size	1	1.00	1.00	1.42	0.00	0.00	0.00	0.00	0.87
time (sec)	N/A	0.019	0.014	0.053	0.000	0.663	0.000	0.000	0.192
Problem 110	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	47	165	0	0	0	0	-1
normalized size	1	1.00	0.72	2.54	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.044	0.038	0.161	0.000	0.546	0.000	0.000	0.000
Problem 111	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	48	238	0	0	0	0	-1
normalized size	1	1.00	0.72	3.55	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.060	0.159	0.000	0.810	0.000	0.000	0.000
Problem 112	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	65	366	0	0	0	0	-1
normalized size	1	1.00	0.67	3.77	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.089	0.247	0.000	0.524	0.000	0.000	0.000
Problem 113	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	63	395	0	0	0	0	-1
normalized size	1	1.00	0.66	4.16	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.136	0.185	0.000	0.634	0.000	0.000	0.000

Problem 114	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	125	77	411	0	0	0	0	-1
normalized size	1	1.00	0.62	3.29	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.094	0.299	0.250	0.000	0.593	0.000	0.000	0.000
Problem 115	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	76	236	0	0	0	0	-1
normalized size	1	1.00	0.59	1.84	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.083	0.088	0.154	0.000	0.789	0.000	0.000	0.000
Problem 116	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	74	223	0	0	0	0	-1
normalized size	1	1.00	0.74	2.23	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.051	0.155	0.000	0.707	0.000	0.000	0.000
Problem 117	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	66	210	0	0	0	0	-1
normalized size	1	1.00	0.66	2.10	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.060	0.049	0.203	0.000	0.587	0.000	0.000	0.000
Problem 118	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	61	213	0	0	0	0	-1
normalized size	1	1.00	0.85	2.96	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.036	0.158	0.000	0.910	0.000	0.000	0.000
Problem 119	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	54	190	0	0	0	0	-1
normalized size	1	1.00	0.75	2.64	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.034	0.238	0.000	0.585	0.000	0.000	0.000

Problem 120	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	0	0	0	-1
normalized size	1	1.00	1.00	3.51	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	0.015	0.190	0.000	0.548	0.000	0.000	0.000
Problem 121	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	0	0	0	-1
normalized size	1	1.00	1.00	3.51	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.024	0.018	0.132	0.000	0.741	0.000	0.000	0.000
Problem 122	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	50	168	0	0	0	0	-1
normalized size	1	1.00	0.74	2.47	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.015	0.158	0.000	0.978	0.000	0.000	0.000
Problem 123	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	51	241	0	0	0	0	-1
normalized size	1	1.00	0.74	3.49	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.048	0.025	0.165	0.000	0.522	0.000	0.000	0.000
Problem 124	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	68	366	0	0	0	0	-1
normalized size	1	1.00	0.69	3.73	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.076	0.051	0.256	0.000	0.764	0.000	0.000	0.000
Problem 125	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	66	398	0	0	0	0	-1
normalized size	1	1.00	0.68	4.10	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.075	0.044	0.182	0.000	0.726	0.000	0.000	0.000

Problem 126	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	80	414	0	0	0	0	-1
normalized size	1	1.00	0.63	3.29	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.096	0.040	0.263	0.000	0.547	0.000	0.000	0.000
Problem 127	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	76	236	0	0	0	0	-1
normalized size	1	1.00	0.59	1.84	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.082	0.052	0.194	0.000	0.511	0.000	0.000	0.000
Problem 128	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	74	223	0	0	0	0	-1
normalized size	1	1.00	0.74	2.23	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.060	0.057	0.181	0.000	0.473	0.000	0.000	0.000
Problem 129	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	66	210	0	0	0	0	-1
normalized size	1	1.00	0.66	2.10	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.060	0.041	0.185	0.000	0.580	0.000	0.000	0.000
Problem 130	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	61	213	0	0	0	0	-1
normalized size	1	1.00	0.85	2.96	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.033	0.162	0.000	0.575	0.000	0.000	0.000
Problem 131	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	54	190	0	0	0	0	-1
normalized size	1	1.00	0.75	2.64	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.029	0.159	0.000	0.601	0.000	0.000	0.000

Problem 132	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	0	0	0	-1
normalized size	1	1.00	1.00	3.51	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	0.013	0.147	0.000	0.592	0.000	0.000	0.000
Problem 133	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	38	144	0	0	0	0	-1
normalized size	1	1.00	0.93	3.51	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.050	0.135	0.000	0.554	0.000	0.000	0.000
Problem 134	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	50	168	0	0	0	0	-1
normalized size	1	1.00	0.74	2.47	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.042	0.021	0.185	0.000	0.645	0.000	0.000	0.000
Problem 135	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	51	241	0	0	0	0	-1
normalized size	1	1.00	0.71	3.35	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.035	0.016	0.166	0.000	0.860	0.000	0.000	0.000
Problem 136	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	68	366	0	0	0	0	-1
normalized size	1	1.00	0.70	3.77	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.070	0.026	0.266	0.000	0.646	0.000	0.000	0.000
Problem 137	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	66	398	0	0	0	0	-1
normalized size	1	1.00	0.67	4.06	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.079	0.062	0.193	0.000	0.522	0.000	0.000	0.000

Problem 138	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	125	80	414	0	0	0	0	-1
normalized size	1	1.00	0.64	3.31	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.098	0.041	0.286	0.000	0.549	0.000	0.000	0.000
Problem 139	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	68	366	0	0	0	0	-1
normalized size	1	1.00	0.68	3.66	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.053	0.016	0.260	0.000	0.813	0.000	0.000	0.000
Problem 140	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	55	62	49	176	0	0	75
normalized size	1	1.00	0.56	0.63	0.50	1.80	0.00	0.00	0.77
time (sec)	N/A	0.027	0.084	0.180	0.771	0.679	0.000	0.000	1.257
Problem 141	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	45	40	42	39	0	0	57
normalized size	1	1.00	0.64	0.57	0.60	0.56	0.00	0.00	0.81
time (sec)	N/A	0.017	0.085	0.112	1.119	0.869	0.000	0.000	0.739
Problem 142	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	45	42	25	150	0	0	62
normalized size	1	1.00	0.71	0.67	0.40	2.38	0.00	0.00	0.98
time (sec)	N/A	0.014	0.053	0.118	1.099	0.642	0.000	0.000	0.689
Problem 143	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	13	28	29	31	44
normalized size	1	1.00	1.00	0.91	0.41	0.88	0.91	0.97	1.38
time (sec)	N/A	0.007	0.024	0.112	0.867	0.731	10.761	2.564	0.389

Problem 144	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	28	26	94	5	0	20
normalized size	1	1.00	1.00	1.17	1.08	3.92	0.21	0.00	0.83
time (sec)	N/A	0.003	0.010	0.083	0.879	0.701	1.751	0.000	0.096
Problem 145	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	42	65	113	0	0	-1
normalized size	1	1.00	1.00	1.27	1.97	3.42	0.00	0.00	-0.03
time (sec)	N/A	0.008	0.011	0.107	1.268	0.597	0.000	0.000	0.000
Problem 146	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	54	28	0	0	59
normalized size	1	1.00	1.00	0.91	1.69	0.88	0.00	0.00	1.84
time (sec)	N/A	0.012	0.017	0.115	1.102	0.607	0.000	0.000	0.702
Problem 147	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	52	104	661	201	0	0	-1
normalized size	1	1.00	0.72	1.44	9.18	2.79	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.042	0.141	1.129	0.833	0.000	0.000	0.000
Problem 148	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	45	42	294	41	0	0	128
normalized size	1	1.00	0.64	0.60	4.20	0.59	0.00	0.00	1.83
time (sec)	N/A	0.018	0.082	0.124	1.127	0.570	0.000	0.000	1.855
Problem 149	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	66	121	1656	227	0	0	-1
normalized size	1	1.00	0.62	1.13	15.48	2.12	0.00	0.00	-0.01
time (sec)	N/A	0.036	0.100	0.197	1.230	0.603	0.000	0.000	0.000

Problem 150	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	55	62	53	183	0	0	76
normalized size	1	1.00	0.54	0.61	0.52	1.81	0.00	0.00	0.75
time (sec)	N/A	0.029	0.078	0.138	1.047	0.596	0.000	0.000	1.048
Problem 151	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	45	40	45	43	0	0	58
normalized size	1	1.00	0.62	0.56	0.62	0.60	0.00	0.00	0.81
time (sec)	N/A	0.018	0.109	0.084	1.248	0.552	0.000	0.000	0.704
Problem 152	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	45	42	28	153	0	0	63
normalized size	1	1.00	0.69	0.65	0.43	2.35	0.00	0.00	0.97
time (sec)	N/A	0.015	0.051	0.096	1.544	0.699	0.000	0.000	0.528
Problem 153	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	32	29	13	29	0	0	29
normalized size	1	1.00	0.97	0.88	0.39	0.88	0.00	0.00	0.88
time (sec)	N/A	0.007	0.037	0.097	1.094	0.514	0.000	0.000	0.240
Problem 154	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	24	28	26	95	5	0	21
normalized size	1	1.00	0.96	1.12	1.04	3.80	0.20	0.00	0.84
time (sec)	N/A	0.003	0.015	0.058	0.828	0.593	125.707	0.000	0.090
Problem 155	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	33	42	68	114	0	0	-1
normalized size	1	1.00	0.97	1.24	2.00	3.35	0.00	0.00	-0.03
time (sec)	N/A	0.008	0.014	0.081	1.106	0.570	0.000	0.000	0.000

Problem 156	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	32	29	54	29	0	0	60
normalized size	1	1.00	0.97	0.88	1.64	0.88	0.00	0.00	1.82
time (sec)	N/A	0.012	0.016	0.096	0.831	0.791	0.000	0.000	0.504
Problem 157	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	52	104	691	204	0	0	-1
normalized size	1	1.00	0.70	1.41	9.34	2.76	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.047	0.111	1.205	0.548	0.000	0.000	0.000
Problem 158	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	45	42	299	42	0	0	129
normalized size	1	1.00	0.62	0.58	4.15	0.58	0.00	0.00	1.79
time (sec)	N/A	0.018	0.049	0.086	1.697	0.544	0.000	0.000	1.140
Problem 159	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	110	67	121	1742	234	0	0	-1
normalized size	1	1.00	0.61	1.10	15.84	2.13	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.082	0.140	1.232	0.687	0.000	0.000	0.000
Problem 160	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	57	52	77	61	0	0	73
normalized size	1	1.00	0.49	0.45	0.66	0.53	0.00	0.00	0.63
time (sec)	N/A	0.026	0.116	0.121	1.015	0.603	0.000	0.000	1.355
Problem 161	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	55	62	59	193	0	0	78
normalized size	1	1.00	0.51	0.58	0.55	1.80	0.00	0.00	0.73
time (sec)	N/A	0.030	0.081	0.121	1.099	0.676	0.000	0.000	1.026

Problem 162	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	45	40	49	47	0	0	60
normalized size	1	1.00	0.59	0.53	0.64	0.62	0.00	0.00	0.79
time (sec)	N/A	0.018	0.126	0.106	1.226	0.731	0.000	0.000	0.621
Problem 163	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	45	42	32	159	0	0	40
normalized size	1	1.00	0.65	0.61	0.46	2.30	0.00	0.00	0.58
time (sec)	N/A	0.016	0.067	0.116	1.089	1.036	0.000	0.000	0.444
Problem 164	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	29	13	31	0	0	31
normalized size	1	1.00	0.91	0.83	0.37	0.89	0.00	0.00	0.89
time (sec)	N/A	0.008	0.049	0.079	1.482	0.534	0.000	0.000	0.323
Problem 165	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	24	28	26	97	0	0	23
normalized size	1	1.00	0.89	1.04	0.96	3.59	0.00	0.00	0.85
time (sec)	N/A	0.003	0.012	0.063	1.204	0.649	0.000	0.000	0.090
Problem 166	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	33	42	72	116	0	0	-1
normalized size	1	1.00	0.92	1.17	2.00	3.22	0.00	0.00	-0.03
time (sec)	N/A	0.009	0.021	0.082	1.357	0.547	0.000	0.000	0.000
Problem 167	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	29	54	31	0	0	62
normalized size	1	1.00	0.91	0.83	1.54	0.89	0.00	0.00	1.77
time (sec)	N/A	0.012	0.018	0.084	1.120	0.624	0.000	0.000	0.424

Problem 168	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	52	104	747	210	0	0	-1
normalized size	1	1.00	0.67	1.33	9.58	2.69	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.052	0.110	1.831	0.538	0.000	0.000	0.000
Problem 169	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	45	42	311	46	0	0	131
normalized size	1	1.00	0.59	0.55	4.09	0.61	0.00	0.00	1.72
time (sec)	N/A	0.019	0.052	0.104	1.265	0.592	0.000	0.000	1.178
Problem 170	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	66	121	1914	244	0	0	-1
normalized size	1	1.00	0.57	1.04	16.50	2.10	0.00	0.00	-0.01
time (sec)	N/A	0.036	0.101	0.161	1.584	0.652	0.000	0.000	0.000
Problem 171	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	57	52	68	54	0	0	73
normalized size	1	1.00	0.53	0.49	0.64	0.50	0.00	0.00	0.68
time (sec)	N/A	0.024	0.103	0.120	1.427	0.670	0.000	0.000	1.178
Problem 172	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	55	62	49	182	0	0	78
normalized size	1	1.00	0.56	0.63	0.50	1.86	0.00	0.00	0.80
time (sec)	N/A	0.027	0.065	0.150	1.342	0.784	0.000	0.000	1.061
Problem 173	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	45	40	42	42	0	0	60
normalized size	1	1.00	0.64	0.57	0.60	0.60	0.00	0.00	0.86
time (sec)	N/A	0.017	0.066	0.104	0.956	0.793	0.000	0.000	0.638

Problem 174	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	45	42	25	157	0	0	65
normalized size	1	1.00	0.71	0.67	0.40	2.49	0.00	0.00	1.03
time (sec)	N/A	0.015	0.052	0.130	1.257	0.629	0.000	0.000	0.653
Problem 175	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	13	31	0	0	47
normalized size	1	1.00	1.00	0.91	0.41	0.97	0.00	0.00	1.47
time (sec)	N/A	0.007	0.025	0.113	1.257	0.649	0.000	0.000	0.363
Problem 176	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	28	26	97	5	0	37
normalized size	1	1.00	1.00	1.17	1.08	4.04	0.21	0.00	1.54
time (sec)	N/A	0.003	0.011	0.078	0.937	0.744	2.430	0.000	0.263
Problem 177	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	42	65	116	0	0	-1
normalized size	1	1.00	1.00	1.27	1.97	3.52	0.00	0.00	-0.03
time (sec)	N/A	0.007	0.011	0.101	1.574	0.627	0.000	0.000	0.000
Problem 178	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	59	31	0	0	62
normalized size	1	1.00	1.00	0.91	1.84	0.97	0.00	0.00	1.94
time (sec)	N/A	0.012	0.021	0.115	1.592	0.656	0.000	0.000	0.542
Problem 179	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	52	104	661	207	0	0	-1
normalized size	1	1.00	0.72	1.44	9.18	2.88	0.00	0.00	-0.01
time (sec)	N/A	0.022	0.035	0.136	1.247	0.995	0.000	0.000	0.000

Problem 180	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	45	42	294	44	0	0	131
normalized size	1	1.00	0.64	0.60	4.20	0.63	0.00	0.00	1.87
time (sec)	N/A	0.018	0.058	0.097	1.600	0.702	0.000	0.000	1.374
Problem 181	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	66	121	1656	233	0	0	-1
normalized size	1	1.00	0.62	1.13	15.48	2.18	0.00	0.00	-0.01
time (sec)	N/A	0.035	0.056	0.151	1.301	0.748	0.000	0.000	0.000
Problem 182	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	55	62	49	182	0	0	78
normalized size	1	1.00	0.51	0.58	0.46	1.70	0.00	0.00	0.73
time (sec)	N/A	0.029	0.050	0.137	1.252	0.934	0.000	0.000	0.984
Problem 183	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	45	40	42	42	0	0	60
normalized size	1	1.00	0.59	0.53	0.55	0.55	0.00	0.00	0.79
time (sec)	N/A	0.019	0.048	0.086	1.002	0.846	0.000	0.000	0.695
Problem 184	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	45	42	25	157	0	0	65
normalized size	1	1.00	0.65	0.61	0.36	2.28	0.00	0.00	0.94
time (sec)	N/A	0.016	0.037	0.107	1.500	1.154	0.000	0.000	0.594
Problem 185	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	29	13	31	0	0	47
normalized size	1	1.00	0.91	0.83	0.37	0.89	0.00	0.00	1.34
time (sec)	N/A	0.007	0.035	0.080	1.480	0.915	0.000	0.000	0.323

Problem 186	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	24	28	26	97	5	0	37
normalized size	1	1.00	0.89	1.04	0.96	3.59	0.19	0.00	1.37
time (sec)	N/A	0.003	0.012	0.080	1.047	0.755	129.109	0.000	0.277
Problem 187	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	33	42	65	116	0	0	-1
normalized size	1	1.00	0.92	1.17	1.81	3.22	0.00	0.00	-0.03
time (sec)	N/A	0.008	0.012	0.110	1.159	0.848	0.000	0.000	0.000
Problem 188	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	29	67	31	0	0	62
normalized size	1	1.00	0.91	0.83	1.91	0.89	0.00	0.00	1.77
time (sec)	N/A	0.013	0.012	0.116	0.999	0.624	0.000	0.000	0.511
Problem 189	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	52	102	670	207	0	0	-1
normalized size	1	1.00	0.67	1.31	8.59	2.65	0.00	0.00	-0.01
time (sec)	N/A	0.022	0.028	0.105	1.572	1.390	0.000	0.000	0.000
Problem 190	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	45	42	311	44	0	0	131
normalized size	1	1.00	0.59	0.55	4.09	0.58	0.00	0.00	1.72
time (sec)	N/A	0.019	0.023	0.107	1.768	0.635	0.000	0.000	1.264
Problem 191	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	66	121	1679	233	0	0	-1
normalized size	1	1.00	0.57	1.04	14.47	2.01	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.039	0.167	2.001	0.781	0.000	0.000	0.000

Problem 192	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	58	62	49	182	0	0	78
normalized size	1	1.00	0.54	0.58	0.46	1.70	0.00	0.00	0.73
time (sec)	N/A	0.028	0.069	0.151	1.187	0.854	0.000	0.000	0.999
Problem 193	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	48	40	42	42	0	0	60
normalized size	1	1.00	0.63	0.53	0.55	0.55	0.00	0.00	0.79
time (sec)	N/A	0.018	0.043	0.097	1.579	0.976	0.000	0.000	0.602
Problem 194	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	48	42	25	157	0	0	65
normalized size	1	1.00	0.70	0.61	0.36	2.28	0.00	0.00	0.94
time (sec)	N/A	0.015	0.040	0.109	1.437	0.940	0.000	0.000	0.606
Problem 195	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	35	29	13	31	0	0	47
normalized size	1	1.00	1.00	0.83	0.37	0.89	0.00	0.00	1.34
time (sec)	N/A	0.007	0.021	0.097	1.628	0.678	0.000	0.000	0.410
Problem 196	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	24	28	26	97	0	0	37
normalized size	1	1.00	0.89	1.04	0.96	3.59	0.00	0.00	1.37
time (sec)	N/A	0.003	0.011	0.077	1.271	0.820	0.000	0.000	0.283
Problem 197	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	33	42	65	116	0	0	-1
normalized size	1	1.00	0.92	1.17	1.81	3.22	0.00	0.00	-0.03
time (sec)	N/A	0.008	0.019	0.084	1.431	0.917	0.000	0.000	0.000

Problem 198	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	29	67	31	0	0	87
normalized size	1	1.00	0.91	0.83	1.91	0.89	0.00	0.00	2.49
time (sec)	N/A	0.013	0.013	0.119	1.347	0.954	0.000	0.000	1.073
Problem 199	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	55	104	688	207	0	0	-1
normalized size	1	1.00	0.71	1.33	8.82	2.65	0.00	0.00	-0.01
time (sec)	N/A	0.022	0.042	0.115	1.563	0.690	0.000	0.000	0.000
Problem 200	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	45	42	343	44	0	0	131
normalized size	1	1.00	0.59	0.55	4.51	0.58	0.00	0.00	1.72
time (sec)	N/A	0.018	0.025	0.111	1.508	0.950	0.000	0.000	1.292
Problem 201	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	66	121	1729	233	0	0	-1
normalized size	1	1.00	0.57	1.04	14.91	2.01	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.043	0.155	1.945	0.831	0.000	0.000	0.000
Problem 202	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	82	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.026	0.099	0.134	0.000	0.658	0.000	0.000	0.000
Problem 203	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	63	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.058	0.187	0.000	0.938	0.000	0.000	0.000

Problem 204	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.051	0.078	0.000	0.769	0.000	0.000	0.000
Problem 205	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.014	0.036	0.131	0.000	1.022	0.000	0.000	0.000
Problem 206	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	54	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.024	0.046	0.129	0.000	0.739	0.000	0.000	0.000
Problem 207	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	56	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.058	0.108	0.000	0.631	0.000	0.000	0.000
Problem 208	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	63	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.032	0.043	0.118	0.000	0.870	0.000	0.000	0.000
Problem 209	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	82	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.112	0.090	0.000	0.921	0.000	0.000	0.000

Problem 210	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	63	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.057	0.157	0.000	0.888	0.000	0.000	0.000
Problem 211	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.051	0.071	0.000	0.945	0.000	0.000	0.000
Problem 212	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.014	0.035	0.091	0.000	0.780	0.000	0.000	0.000
Problem 213	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	56	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.027	0.045	0.131	0.000	0.956	0.000	0.000	0.000
Problem 214	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	54	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.036	0.055	0.111	0.000	0.990	0.000	0.000	0.000
Problem 215	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	63	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.036	0.042	0.121	0.000	0.986	0.000	0.000	0.000

Problem 216	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	82	0	0	0	0	0	-1
normalized size	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.026	0.166	0.101	0.000	0.729	0.000	0.000	0.000
Problem 217	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	63	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.073	0.163	0.000	0.688	0.000	0.000	0.000
Problem 218	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.068	0.076	0.000	0.917	0.000	0.000	0.000
Problem 219	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.014	0.003	0.022	0.000	0.790	0.000	0.000	0.000
Problem 220	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	56	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.005	0.103	0.000	0.943	0.000	0.000	0.000
Problem 221	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	56	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.036	0.009	0.123	0.000	0.814	0.000	0.000	0.000

Problem 222	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.037	0.056	0.102	0.000	0.670	0.000	0.000	0.000
Problem 223	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	82	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.026	0.108	0.069	0.000	0.770	0.000	0.000	0.000
Problem 224	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	63	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.068	0.128	0.000	1.048	0.000	0.000	0.000
Problem 225	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.005	0.092	0.000	1.018	0.000	0.000	0.000
Problem 226	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.014	0.003	0.043	0.000	0.956	0.000	0.000	0.000
Problem 227	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	54	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.044	0.084	0.000	0.872	0.000	0.000	0.000

Problem 228	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	58	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.034	0.048	0.087	0.000	0.673	0.000	0.000	0.000
Problem 229	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.032	0.101	0.128	0.000	0.488	0.000	0.000	0.000
Problem 230	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	82	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.025	0.103	0.066	0.000	0.602	0.000	0.000	0.000
Problem 231	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	63	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.069	0.135	0.000	0.635	0.000	0.000	0.000
Problem 232	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.004	0.072	0.000	0.569	0.000	0.000	0.000
Problem 233	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	53	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.015	0.003	0.033	0.000	0.500	0.000	0.000	0.000

Problem 234	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	56	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.044	0.078	0.000	0.784	0.000	0.000	0.000
Problem 235	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	58	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.037	0.052	0.086	0.000	0.648	0.000	0.000	0.000
Problem 236	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.045	0.097	0.123	0.000	0.958	0.000	0.000	0.000
Problem 237	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	82	0	0	0	0	0	-1
normalized size	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.031	0.103	0.072	0.000	0.641	0.000	0.000	0.000
Problem 238	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.021	0.004	0.130	0.000	0.680	0.000	0.000	0.000
Problem 239	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.021	0.008	0.092	0.000	0.522	0.000	0.000	0.000

Problem 240	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	53	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.016	0.003	0.023	0.000	0.588	0.000	0.000	0.000
Problem 241	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	56	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.027	0.008	0.082	0.000	0.659	0.000	0.000	0.000
Problem 242	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	58	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.059	0.058	0.099	0.000	0.566	0.000	0.000	0.000
Problem 243	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.042	0.104	0.110	0.000	0.684	0.000	0.000	0.000
Problem 244	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	77	0	0	0	0	0	-1
normalized size	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.084	0.891	0.000	0.666	0.000	0.000	0.000
Problem 245	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	72	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.072	1.330	0.000	1.013	0.000	0.000	0.000

Problem 246	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	70	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.049	0.051	1.147	0.000	0.622	0.000	0.000	0.000
Problem 247	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	64	0	0	0	0	0	-1
normalized size	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.024	0.042	0.231	0.000	0.520	0.000	0.000	0.000
Problem 248	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	63	0	0	0	0	0	-1
normalized size	1	1.00	1.05	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.030	0.055	0.479	0.000	0.597	0.000	0.000	0.000
Problem 249	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	67	0	0	0	0	0	-1
normalized size	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.045	0.063	0.402	0.000	0.892	0.000	0.000	0.000
Problem 250	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	70	0	0	0	0	0	-1
normalized size	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.042	0.046	0.568	0.000	0.545	0.000	0.000	0.000
Problem 251	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	72	0	0	0	0	0	-1
normalized size	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.042	0.053	0.341	0.000	1.042	0.000	0.000	0.000

Problem 252	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.027	0.130	0.132	0.000	0.545	0.000	0.000	0.000
Problem 253	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.110	0.104	0.000	0.636	0.000	0.000	0.000
Problem 254	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.027	0.084	0.124	0.000	0.596	0.000	0.000	0.000
Problem 255	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.027	0.054	0.099	0.000	0.785	0.000	0.000	0.000
Problem 256	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.078	0.096	0.000	0.438	0.000	0.000	0.000
Problem 257	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.082	0.096	0.000	0.659	0.000	0.000	0.000

Problem 258	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.080	0.092	0.000	0.618	0.000	0.000	0.000
Problem 259	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.083	0.085	0.000	0.509	0.000	0.000	0.000
Problem 260	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	89	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.070	10.906	0.842	0.000	0.668	0.000	0.000	0.000
Problem 261	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	0	13	15
normalized size	1	1.00	1.00	0.93	0.87	0.87	0.00	0.87	1.00
time (sec)	N/A	0.025	0.020	0.038	0.639	0.638	0.000	0.463	0.304
Problem 262	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	23	0	13	15
normalized size	1	1.00	1.00	0.82	0.76	1.35	0.00	0.76	0.88
time (sec)	N/A	0.025	0.023	0.023	0.839	0.539	0.000	0.625	0.213
Problem 263	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	53	88	0	0	0	0	-1
normalized size	1	1.00	0.79	1.31	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.097	0.160	0.000	0.488	0.000	0.000	0.000

Problem 264	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	61	142	0	0	0	0	-1
normalized size	1	1.00	0.91	2.12	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.048	0.145	0.140	0.000	0.502	0.000	0.000	0.000
Problem 265	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	18	14	13	22	0	14	16
normalized size	1	1.00	0.86	0.67	0.62	1.05	0.00	0.67	0.76
time (sec)	N/A	0.024	0.025	0.096	0.601	0.493	0.000	0.579	0.335
Problem 266	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	27	26	25	23	0	24	-1
normalized size	1	1.00	0.82	0.79	0.76	0.70	0.00	0.73	-0.03
time (sec)	N/A	0.033	0.062	0.093	0.697	0.577	0.000	1.091	0.000
Problem 267	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	29	26	25	33	0	25	-1
normalized size	1	1.00	0.83	0.74	0.71	0.94	0.00	0.71	-0.03
time (sec)	N/A	0.033	0.065	0.076	0.517	0.547	0.000	0.506	0.000
Problem 268	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	63	100	0	0	0	0	-1
normalized size	1	1.00	0.68	1.09	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.080	0.136	0.159	0.000	0.622	0.000	0.000	0.000
Problem 269	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	63	152	0	0	0	0	-1
normalized size	1	1.00	0.68	1.65	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.079	0.406	0.140	0.000	0.656	0.000	0.000	0.000

Problem 270	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	7	6	6	0	6	8
normalized size	1	1.00	1.00	0.70	0.60	0.60	0.00	0.60	0.80
time (sec)	N/A	0.017	0.008	0.020	1.131	0.573	0.000	0.370	0.203
Problem 271	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	47	28	41	95	0	0	-1
normalized size	1	1.00	1.47	0.88	1.28	2.97	0.00	0.00	-0.03
time (sec)	N/A	0.028	0.029	0.175	1.349	0.821	0.000	0.000	0.000
Problem 272	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	50	48	41	97	0	0	-1
normalized size	1	1.00	1.61	1.55	1.32	3.13	0.00	0.00	-0.03
time (sec)	N/A	0.029	0.033	0.188	0.704	0.649	0.000	0.000	0.000
Problem 273	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	49	123	0	0	0	0	-1
normalized size	1	1.00	0.80	2.02	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.048	0.142	0.188	0.000	0.534	0.000	0.000	0.000
Problem 274	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	54	177	0	0	0	0	-1
normalized size	1	1.00	0.87	2.85	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.048	0.148	0.192	0.000	0.828	0.000	0.000	0.000
Problem 275	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	73	73	65	131	0	0	-1
normalized size	1	1.00	1.18	1.18	1.05	2.11	0.00	0.00	-0.02
time (sec)	N/A	0.053	0.112	0.189	1.328	0.676	0.000	0.000	0.000

Problem 276	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	33	71	63	141	0	0	-1
normalized size	1	1.00	0.53	1.15	1.02	2.27	0.00	0.00	-0.02
time (sec)	N/A	0.046	0.028	0.173	1.443	0.749	0.000	0.000	0.000
Problem 277	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-1)	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	64	168	0	0	0	0	-1
normalized size	1	1.00	0.70	1.83	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.081	0.378	0.201	0.000	0.598	0.000	0.000	0.000
Problem 278	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	76	160	0	0	0	0	-1
normalized size	1	1.00	0.83	1.74	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.080	0.232	0.323	0.000	0.743	0.000	0.000	0.000
Problem 279	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	105	0	0	0	0	0	-1
normalized size	1	1.00	1.38	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.105	0.680	0.219	0.000	0.560	0.000	0.000	0.000
Problem 280	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	70	0	0	0	0	0	-1
normalized size	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.092	0.170	0.141	0.000	0.543	0.000	0.000	0.000
Problem 281	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	68	0	0	0	0	0	-1
normalized size	1	1.00	0.89	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.097	0.220	0.132	0.000	0.702	0.000	0.000	0.000

Problem 282	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	68	0	0	0	0	0	-1
normalized size	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.111	0.260	0.128	0.000	0.588	0.000	0.000	0.000
Problem 283	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	70	0	0	0	0	0	-1
normalized size	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.110	0.280	0.148	0.000	0.628	0.000	0.000	0.000
Problem 284	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	312	0	0	0	0	0	-1
normalized size	1	1.00	3.67	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.083	1.966	0.653	0.000	0.657	0.000	0.000	0.000
Problem 285	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	314	0	0	0	0	0	-1
normalized size	1	1.00	3.57	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.091	0.227	0.555	0.000	0.679	0.000	0.000	0.000
Problem 286	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	314	0	0	0	0	0	-1
normalized size	1	1.00	3.57	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.092	0.237	0.627	0.000	0.688	0.000	0.000	0.000
Problem 287	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	316	0	0	0	0	0	-1
normalized size	1	1.00	3.47	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.099	0.227	0.661	0.000	0.854	0.000	0.000	0.000

Problem 288	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	76	94	0	0	0	0	0	-1
normalized size	1	0.97	1.21	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.118	7.909	0.171	0.000	0.582	0.000	0.000	0.000
Problem 289	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	94	0	0	0	0	0	-1
normalized size	1	1.00	1.24	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.116	2.386	0.170	0.000	0.652	0.000	0.000	0.000
Problem 290	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	94	0	0	0	0	0	-1
normalized size	1	1.00	1.24	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.112	1.240	0.156	0.000	0.645	0.000	0.000	0.000
Problem 291	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	96	0	0	0	0	0	-1
normalized size	1	1.00	1.23	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.098	1.050	0.157	0.000	0.873	0.000	0.000	0.000
Problem 292	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	225	0	0	0	0	0	-1
normalized size	1	1.00	2.88	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.098	1.639	0.140	0.000	0.842	0.000	0.000	0.000
Problem 293	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	116	0	0	0	0	0	-1
normalized size	1	1.00	1.49	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.112	6.449	0.136	0.000	0.694	0.000	0.000	0.000

Problem 294	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	125	0	0	0	0	0	-1
normalized size	1	1.00	1.60	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.111	1.060	0.132	0.000	0.972	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [275] had the largest ratio of [.3158]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	6	0.167
2	A	2	2	1.00	8	0.250
3	A	2	1	1.00	8	0.125
4	A	3	2	1.00	8	0.250
5	A	2	1	1.00	8	0.125
6	A	4	2	1.00	8	0.250
7	A	2	1	1.00	8	0.125
8	A	5	2	1.00	8	0.250
9	A	3	2	1.00	10	0.200
10	A	2	2	1.00	10	0.200
11	A	2	2	1.00	10	0.200
12	A	1	1	1.00	10	0.100
13	A	1	1	1.00	10	0.100
14	A	2	2	1.00	10	0.200
15	A	2	2	1.00	10	0.200
16	A	3	2	1.00	10	0.200
17	A	4	3	1.00	12	0.250
18	A	3	3	1.00	12	0.250
19	A	3	3	1.00	12	0.250
20	A	2	2	1.00	12	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
21	A	2	2	1.00	12	0.167
22	A	3	3	1.00	12	0.250
23	A	3	3	1.00	12	0.250
24	A	4	3	1.00	12	0.250
25	A	1	1	1.00	10	0.100
26	A	1	1	1.00	10	0.100
27	A	1	1	1.00	10	0.100
28	A	1	1	1.00	10	0.100
29	A	1	1	1.00	10	0.100
30	A	1	1	1.00	10	0.100
31	A	1	1	1.00	12	0.083
32	A	1	1	1.00	12	0.083
33	A	1	1	1.00	12	0.083
34	A	1	1	1.00	12	0.083
35	A	1	1	1.00	12	0.083
36	A	1	1	1.00	12	0.083
37	A	1	1	1.00	8	0.125
38	A	1	1	1.00	10	0.100
39	A	4	3	1.00	10	0.300
40	A	3	3	1.00	10	0.300
41	A	2	2	1.00	10	0.200
42	A	2	2	1.00	10	0.200
43	A	3	3	1.00	10	0.300
44	A	4	3	1.00	10	0.300
45	A	6	3	1.00	10	0.300
46	A	4	3	1.00	10	0.300
47	A	3	3	1.00	10	0.300
48	A	3	3	1.00	10	0.300
49	A	4	3	1.00	10	0.300
50	A	6	3	1.00	10	0.300
51	A	7	3	1.00	10	0.300
52	A	5	3	1.00	10	0.300
53	A	3	3	1.00	10	0.300
54	A	3	3	1.00	10	0.300
55	A	3	2	1.00	10	0.200
56	A	3	2	1.00	10	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	2	2	1.00	12	0.167
58	A	2	2	1.00	14	0.143
59	A	2	2	1.00	14	0.143
60	A	2	2	1.00	14	0.143
61	A	2	2	1.00	14	0.143
62	A	2	2	1.00	14	0.143
63	A	2	2	1.00	14	0.143
64	A	2	2	1.00	14	0.143
65	A	2	2	1.00	14	0.143
66	A	6	4	1.00	21	0.190
67	A	5	4	1.00	21	0.190
68	A	5	4	1.00	21	0.190
69	A	4	4	1.00	21	0.190
70	A	4	4	1.00	19	0.210
71	A	2	2	1.00	12	0.167
72	A	3	3	1.00	19	0.158
73	A	4	4	1.00	21	0.190
74	A	4	4	1.00	21	0.190
75	A	5	4	1.00	21	0.190
76	A	5	4	1.00	21	0.190
77	A	6	4	1.00	21	0.190
78	A	6	4	1.00	21	0.190
79	A	5	4	1.00	21	0.190
80	A	5	4	1.00	21	0.190
81	A	4	4	1.00	19	0.210
82	A	3	3	1.00	12	0.250
83	A	3	3	1.00	19	0.158
84	A	3	3	1.00	21	0.143
85	A	4	4	1.00	21	0.190
86	A	4	4	1.00	21	0.190
87	A	5	4	1.00	21	0.190
88	A	5	4	1.00	21	0.190
89	A	6	4	1.00	21	0.190
90	A	6	4	1.00	21	0.190
91	A	5	4	1.00	21	0.190
92	A	5	4	1.00	19	0.210

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
93	A	3	3	1.00	12	0.250
94	A	4	4	1.00	19	0.210
95	A	3	3	1.00	21	0.143
96	A	3	3	1.00	21	0.143
97	A	4	4	1.00	21	0.190
98	A	4	4	1.00	21	0.190
99	A	5	4	1.00	21	0.190
100	A	5	4	1.00	21	0.190
101	A	6	4	1.00	21	0.190
102	A	4	3	1.00	12	0.250
103	A	6	4	1.00	21	0.190
104	A	5	4	1.00	21	0.190
105	A	5	4	1.00	21	0.190
106	A	4	4	1.00	21	0.190
107	A	4	4	1.00	21	0.190
108	A	3	3	1.00	19	0.158
109	A	2	2	1.00	12	0.167
110	A	4	4	1.00	19	0.210
111	A	4	4	1.00	21	0.190
112	A	5	4	1.00	21	0.190
113	A	5	4	1.00	21	0.190
114	A	6	4	1.00	21	0.190
115	A	6	4	1.00	21	0.190
116	A	5	4	1.00	21	0.190
117	A	5	4	1.00	21	0.190
118	A	4	4	1.00	21	0.190
119	A	4	4	1.00	21	0.190
120	A	3	3	1.00	21	0.143
121	A	3	3	1.00	19	0.158
122	A	3	3	1.00	12	0.250
123	A	4	4	1.00	19	0.210
124	A	5	4	1.00	21	0.190
125	A	5	4	1.00	21	0.190
126	A	6	4	1.00	21	0.190
127	A	6	4	1.00	21	0.190
128	A	5	4	1.00	21	0.190

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
129	A	5	4	1.00	21	0.190
130	A	4	4	1.00	21	0.190
131	A	4	4	1.00	21	0.190
132	A	3	3	1.00	21	0.143
133	A	3	3	1.00	21	0.143
134	A	4	4	1.00	19	0.210
135	A	3	3	1.00	12	0.250
136	A	5	4	1.00	19	0.210
137	A	5	4	1.00	21	0.190
138	A	6	4	1.00	21	0.190
139	A	4	3	1.00	12	0.250
140	A	4	3	1.00	23	0.130
141	A	3	2	1.00	23	0.087
142	A	3	3	1.00	23	0.130
143	A	2	2	1.00	23	0.087
144	A	2	2	1.00	23	0.087
145	A	2	2	1.00	23	0.087
146	A	3	3	1.00	23	0.130
147	A	3	3	1.00	23	0.130
148	A	3	2	1.00	23	0.087
149	A	4	3	1.00	23	0.130
150	A	4	3	1.00	23	0.130
151	A	3	2	1.00	23	0.087
152	A	3	3	1.00	23	0.130
153	A	2	2	1.00	23	0.087
154	A	2	2	1.00	23	0.087
155	A	2	2	1.00	23	0.087
156	A	3	3	1.00	23	0.130
157	A	3	3	1.00	23	0.130
158	A	3	2	1.00	23	0.087
159	A	4	3	1.00	23	0.130
160	A	3	2	1.00	23	0.087
161	A	4	3	1.00	23	0.130
162	A	3	2	1.00	23	0.087
163	A	3	3	1.00	23	0.130
164	A	2	2	1.00	23	0.087

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
165	A	2	2	1.00	23	0.087
166	A	2	2	1.00	23	0.087
167	A	3	3	1.00	23	0.130
168	A	3	3	1.00	23	0.130
169	A	3	2	1.00	23	0.087
170	A	4	3	1.00	23	0.130
171	A	3	2	1.00	23	0.087
172	A	4	3	1.00	23	0.130
173	A	3	2	1.00	23	0.087
174	A	3	3	1.00	23	0.130
175	A	2	2	1.00	23	0.087
176	A	2	2	1.00	23	0.087
177	A	2	2	1.00	23	0.087
178	A	3	3	1.00	23	0.130
179	A	3	3	1.00	23	0.130
180	A	3	2	1.00	23	0.087
181	A	4	3	1.00	23	0.130
182	A	4	3	1.00	23	0.130
183	A	3	2	1.00	23	0.087
184	A	3	3	1.00	23	0.130
185	A	2	2	1.00	23	0.087
186	A	2	2	1.00	23	0.087
187	A	2	2	1.00	23	0.087
188	A	3	3	1.00	23	0.130
189	A	3	3	1.00	23	0.130
190	A	3	2	1.00	23	0.087
191	A	4	3	1.00	23	0.130
192	A	4	3	1.00	23	0.130
193	A	3	2	1.00	23	0.087
194	A	3	3	1.00	23	0.130
195	A	2	2	1.00	23	0.087
196	A	2	2	1.00	23	0.087
197	A	2	2	1.00	23	0.087
198	A	3	3	1.00	23	0.130
199	A	3	3	1.00	23	0.130
200	A	3	2	1.00	23	0.087

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
201	A	4	3	1.00	23	0.130
202	A	2	2	1.00	21	0.095
203	A	2	2	1.00	21	0.095
204	A	2	2	1.00	19	0.105
205	A	1	1	1.00	12	0.083
206	A	2	2	1.00	19	0.105
207	A	2	2	1.00	21	0.095
208	A	2	2	1.00	21	0.095
209	A	2	2	1.00	21	0.095
210	A	2	2	1.00	21	0.095
211	A	2	2	1.00	19	0.105
212	A	1	1	1.00	12	0.083
213	A	2	2	1.00	19	0.105
214	A	2	2	1.00	21	0.095
215	A	2	2	1.00	21	0.095
216	A	2	2	1.00	21	0.095
217	A	2	2	1.00	21	0.095
218	A	2	2	1.00	19	0.105
219	A	1	1	1.00	12	0.083
220	A	2	2	1.00	19	0.105
221	A	2	2	1.00	21	0.095
222	A	2	2	1.00	21	0.095
223	A	2	2	1.00	21	0.095
224	A	2	2	1.00	21	0.095
225	A	2	2	1.00	19	0.105
226	A	1	1	1.00	12	0.083
227	A	2	2	1.00	19	0.105
228	A	2	2	1.00	21	0.095
229	A	2	2	1.00	21	0.095
230	A	2	2	1.00	21	0.095
231	A	2	2	1.00	21	0.095
232	A	2	2	1.00	19	0.105
233	A	1	1	1.00	12	0.083
234	A	2	2	1.00	19	0.105
235	A	2	2	1.00	21	0.095
236	A	2	2	1.00	21	0.095

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
237	A	2	2	1.00	21	0.095
238	A	2	2	1.00	21	0.095
239	A	2	2	1.00	19	0.105
240	A	1	1	1.00	12	0.083
241	A	2	2	1.00	19	0.105
242	A	2	2	1.00	21	0.095
243	A	2	2	1.00	21	0.095
244	A	2	2	1.00	21	0.095
245	A	2	2	1.00	19	0.105
246	A	2	2	1.00	17	0.118
247	A	1	1	1.00	10	0.100
248	A	2	2	1.00	17	0.118
249	A	2	2	1.00	19	0.105
250	A	2	2	1.00	19	0.105
251	A	2	2	1.00	19	0.105
252	A	2	2	1.00	21	0.095
253	A	2	2	1.00	21	0.095
254	A	2	2	1.00	21	0.095
255	A	2	2	1.00	21	0.095
256	A	2	2	1.00	21	0.095
257	A	2	2	1.00	21	0.095
258	A	2	2	1.00	21	0.095
259	A	2	2	1.00	21	0.095
260	A	2	2	1.00	21	0.095
261	A	2	2	1.00	17	0.118
262	A	2	2	1.00	17	0.118
263	A	3	3	1.00	19	0.158
264	A	3	3	1.00	19	0.158
265	A	3	2	1.00	11	0.182
266	A	3	2	1.00	19	0.105
267	A	3	2	1.00	19	0.105
268	A	4	3	1.00	19	0.158
269	A	4	3	1.00	19	0.158
270	A	2	2	1.00	9	0.222
271	A	5	5	1.00	17	0.294
272	A	5	5	1.00	17	0.294

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
273	A	3	3	1.00	19	0.158
274	A	3	3	1.00	19	0.158
275	A	6	6	1.00	19	0.316
276	A	6	6	1.00	19	0.316
277	A	4	3	1.00	19	0.158
278	A	4	3	1.00	19	0.158
279	A	2	2	1.00	21	0.095
280	A	2	2	1.00	21	0.095
281	A	2	2	1.00	21	0.095
282	A	2	2	1.00	21	0.095
283	A	2	2	1.00	21	0.095
284	A	2	2	1.00	17	0.118
285	A	2	2	1.00	19	0.105
286	A	2	2	1.00	19	0.105
287	A	2	2	1.00	21	0.095
288	A	2	2	0.97	23	0.087
289	A	2	2	1.00	23	0.087
290	A	2	2	1.00	23	0.087
291	A	2	2	1.00	23	0.087
292	A	2	2	1.00	23	0.087
293	A	2	2	1.00	23	0.087
294	A	2	2	1.00	23	0.087

Chapter 3

Listing of integrals

3.1 $\int \cos(a + bx) dx$

Optimal. Leaf size=10

$$\frac{\sin(a + bx)}{b}$$

[Out] sin(b*x+a)/b

Rubi [A] time = 0.00, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2637}

$$\frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x],x]

[Out] Sin[a + b*x]/b

Rule 2637

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] :> Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]

Rubi steps

$$\int \cos(a + bx) dx = \frac{\sin(a + bx)}{b}$$

Mathematica [B] time = 0.08, size = 21, normalized size = 2.10

$$\frac{\sin(a) \cos(bx)}{b} + \frac{\cos(a) \sin(bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x],x]

[Out] (Cos[b*x]*Sin[a])/b + (Cos[a]*Sin[b*x])/b

fricas [A] time = 0.40, size = 10, normalized size = 1.00

$$\frac{\sin(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a),x, algorithm="fricas")

[Out] sin(b*x + a)/b

giac [A] time = 0.16, size = 10, normalized size = 1.00

$$\frac{\sin(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a),x, algorithm="giac")

[Out] sin(b*x + a)/b

maple [A] time = 0.05, size = 11, normalized size = 1.10

$$\frac{\sin(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a),x)

[Out] sin(b*x+a)/b

maxima [A] time = 0.47, size = 10, normalized size = 1.00

$$\frac{\sin(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a),x, algorithm="maxima")

[Out] sin(b*x + a)/b

mupad [B] time = 0.25, size = 10, normalized size = 1.00

$$\frac{\sin(a + bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x),x)

[Out] sin(a + b*x)/b

sympy [A] time = 0.12, size = 12, normalized size = 1.20

$$\begin{cases} \frac{\sin(a+bx)}{b} & \text{for } b \neq 0 \\ x \cos(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a),x)

[Out] Piecewise((sin(a + b*x)/b, Ne(b, 0)), (x*cos(a), True))

3.2 $\int \cos^2(a + bx) dx$

Optimal. Leaf size=25

$$\frac{\sin(a + bx) \cos(a + bx)}{2b} + \frac{x}{2}$$

[Out] 1/2*x+1/2*cos(b*x+a)*sin(b*x+a)/b

Rubi [A] time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 8}

$$\frac{\sin(a + bx) \cos(a + bx)}{2b} + \frac{x}{2}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2,x]

[Out] x/2 + (Cos[a + b*x]*Sin[a + b*x])/(2*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x] * (b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^2(a + bx) dx &= \frac{\cos(a + bx) \sin(a + bx)}{2b} + \frac{\int 1 dx}{2} \\ &= \frac{x}{2} + \frac{\cos(a + bx) \sin(a + bx)}{2b} \end{aligned}$$

Mathematica [A] time = 0.02, size = 23, normalized size = 0.92

$$\frac{2(a + bx) + \sin(2(a + bx))}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^2,x]

[Out] (2*(a + b*x) + Sin[2*(a + b*x)])/(4*b)

fricas [A] time = 0.43, size = 22, normalized size = 0.88

$$\frac{bx + \cos(bx + a) \sin(bx + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2,x, algorithm="fricas")

[Out] 1/2*(b*x + cos(b*x + a)*sin(b*x + a))/b

giac [A] time = 0.19, size = 18, normalized size = 0.72

$$\frac{1}{2}x + \frac{\sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2,x, algorithm="giac")

[Out] 1/2*x + 1/4*sin(2*b*x + 2*a)/b

maple [A] time = 0.05, size = 27, normalized size = 1.08

$$\frac{\frac{\cos(bx+a)\sin(bx+a)}{2} + \frac{bx}{2} + \frac{a}{2}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2,x)

[Out] 1/b*(1/2*cos(b*x+a)*sin(b*x+a)+1/2*b*x+1/2*a)

maxima [A] time = 0.30, size = 22, normalized size = 0.88

$$\frac{2bx + 2a + \sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2,x, algorithm="maxima")

[Out] 1/4*(2*b*x + 2*a + sin(2*b*x + 2*a))/b

mupad [B] time = 0.23, size = 18, normalized size = 0.72

$$\frac{x}{2} + \frac{\sin(2a + 2bx)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2,x)

[Out] x/2 + sin(2*a + 2*b*x)/(4*b)

sympy [A] time = 0.19, size = 46, normalized size = 1.84

$$\begin{cases} \frac{x \sin^2(a+bx)}{2} + \frac{x \cos^2(a+bx)}{2} + \frac{\sin(a+bx) \cos(a+bx)}{2b} & \text{for } b \neq 0 \\ x \cos^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2,x)

[Out] Piecewise((x*sin(a + b*x)**2/2 + x*cos(a + b*x)**2/2 + sin(a + b*x)*cos(a + b*x)/(2*b), Ne(b, 0)), (x*cos(a)**2, True))

3.3 $\int \cos^3(a + bx) dx$

Optimal. Leaf size=26

$$\frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

[Out] $\sin(b*x+a)/b-1/3*\sin(b*x+a)^3/b$

Rubi [A] time = 0.01, antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2633}

$$\frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3,x]

[Out] Sin[a + b*x]/b - Sin[a + b*x]^3/(3*b)

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :- Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) dx &= -\frac{\text{Subst}\left(\int (1 - x^2) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A] time = 0.01, size = 26, normalized size = 1.00

$$\frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^3,x]

[Out] Sin[a + b*x]/b - Sin[a + b*x]^3/(3*b)

fricas [A] time = 0.42, size = 21, normalized size = 0.81

$$\frac{(\cos(bx + a)^2 + 2) \sin(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3,x, algorithm="fricas")

[Out] 1/3*(cos(b*x + a)^2 + 2)*sin(b*x + a)/b

giac [A] time = 0.16, size = 22, normalized size = 0.85

$$-\frac{\sin(bx + a)^3 - 3 \sin(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3,x, algorithm="giac")

[Out] -1/3*(sin(b*x + a)^3 - 3*sin(b*x + a))/b

maple [A] time = 0.12, size = 22, normalized size = 0.85

$$\frac{(2 + \cos^2(bx + a)) \sin(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^3,x)

[Out] 1/3/b*(2+cos(b*x+a)^2)*sin(b*x+a)

maxima [A] time = 0.30, size = 22, normalized size = 0.85

$$\frac{\sin(bx + a)^3 - 3 \sin(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3,x, algorithm="maxima")

[Out] -1/3*(sin(b*x + a)^3 - 3*sin(b*x + a))/b

mupad [B] time = 0.07, size = 24, normalized size = 0.92

$$\frac{3 \sin(a + bx) - \sin(a + bx)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3,x)

[Out] (3*sin(a + b*x) - sin(a + b*x)^3)/(3*b)

sympy [A] time = 0.41, size = 36, normalized size = 1.38

$$\begin{cases} \frac{2 \sin^3(a+bx)}{3b} + \frac{\sin(a+bx) \cos^2(a+bx)}{b} & \text{for } b \neq 0 \\ x \cos^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3,x)

[Out] Piecewise((2*sin(a + b*x)**3/(3*b) + sin(a + b*x)*cos(a + b*x)**2/b, Ne(b, 0)), (x*cos(a)**3, True))

3.4 $\int \cos^4(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sin(a + bx) \cos^3(a + bx)}{4b} + \frac{3 \sin(a + bx) \cos(a + bx)}{8b} + \frac{3x}{8}$$

[Out] 3/8*x+3/8*cos(b*x+a)*sin(b*x+a)/b+1/4*cos(b*x+a)^3*sin(b*x+a)/b

Rubi [A] time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 8}

$$\frac{\sin(a + bx) \cos^3(a + bx)}{4b} + \frac{3 \sin(a + bx) \cos(a + bx)}{8b} + \frac{3x}{8}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4,x]

[Out] (3*x)/8 + (3*Cos[a + b*x]*Sin[a + b*x])/(8*b) + (Cos[a + b*x]^3*Sin[a + b*x])/b

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^4(a + bx) dx &= \frac{\cos^3(a + bx) \sin(a + bx)}{4b} + \frac{3}{4} \int \cos^2(a + bx) dx \\ &= \frac{3 \cos(a + bx) \sin(a + bx)}{8b} + \frac{\cos^3(a + bx) \sin(a + bx)}{4b} + \frac{3 \int 1 dx}{8} \\ &= \frac{3x}{8} + \frac{3 \cos(a + bx) \sin(a + bx)}{8b} + \frac{\cos^3(a + bx) \sin(a + bx)}{4b} \end{aligned}$$

Mathematica [A] time = 0.04, size = 33, normalized size = 0.72

$$\frac{12(a + bx) + 8 \sin(2(a + bx)) + \sin(4(a + bx))}{32b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^4,x]

[Out] (12*(a + b*x) + 8*Sin[2*(a + b*x)] + Sin[4*(a + b*x)])/(32*b)

fricas [A] time = 0.41, size = 36, normalized size = 0.78

$$\frac{3bx + (2 \cos(bx + a))^3 + 3 \cos(bx + a) \sin(bx + a)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4,x, algorithm="fricas")

[Out] 1/8*(3*b*x + (2*cos(b*x + a)^3 + 3*cos(b*x + a))*sin(b*x + a))/b

giac [A] time = 0.16, size = 32, normalized size = 0.70

$$\frac{3}{8}x + \frac{\sin(4bx + 4a)}{32b} + \frac{\sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4,x, algorithm="giac")

[Out] 3/8*x + 1/32*sin(4*b*x + 4*a)/b + 1/4*sin(2*b*x + 2*a)/b

maple [A] time = 0.08, size = 38, normalized size = 0.83

$$\frac{\left(\cos^3(bx+a) + \frac{3\cos(bx+a)}{2}\right)\sin(bx+a)}{4} + \frac{3bx}{8} + \frac{3a}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^4,x)

[Out] 1/b*(1/4*(cos(b*x+a)^3+3/2*cos(b*x+a))*sin(b*x+a)+3/8*b*x+3/8*a)

maxima [A] time = 0.49, size = 33, normalized size = 0.72

$$\frac{12bx + 12a + \sin(4bx + 4a) + 8\sin(2bx + 2a)}{32b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4,x, algorithm="maxima")

[Out] 1/32*(12*b*x + 12*a + sin(4*b*x + 4*a) + 8*sin(2*b*x + 2*a))/b

mupad [B] time = 0.19, size = 31, normalized size = 0.67

$$\frac{3x}{8} + \frac{\frac{\sin(2a+2bx)}{4} + \frac{\sin(4a+4bx)}{32}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^4,x)

[Out] (3*x)/8 + (sin(2*a + 2*b*x)/4 + sin(4*a + 4*b*x)/32)/b

sympy [A] time = 0.86, size = 95, normalized size = 2.07

$$\begin{cases} \frac{3x \sin^4(a+bx)}{8} + \frac{3x \sin^2(a+bx) \cos^2(a+bx)}{4} + \frac{3x \cos^4(a+bx)}{8} + \frac{3 \sin^3(a+bx) \cos(a+bx)}{8b} + \frac{5 \sin(a+bx) \cos^3(a+bx)}{8b} & \text{for } b \neq 0 \\ x \cos^4(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**4,x)

[Out] Piecewise(((3*x*sin(a + b*x)**4/8 + 3*x*sin(a + b*x)**2*cos(a + b*x)**2/4 + 3*x*cos(a + b*x)**4/8 + 3*sin(a + b*x)**3*cos(a + b*x)/(8*b) + 5*sin(a + b*x)*cos(a + b*x)**3/(8*b), Ne(b, 0)), (x*cos(a)**4, True))

3.5 $\int \cos^5(a + bx) dx$

Optimal. Leaf size=41

$$\frac{\sin^5(a + bx)}{5b} - \frac{2 \sin^3(a + bx)}{3b} + \frac{\sin(a + bx)}{b}$$

[Out] $\sin(b*x+a)/b-2/3*\sin(b*x+a)^3/b+1/5*\sin(b*x+a)^5/b$

Rubi [A] time = 0.01, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2633}

$$\frac{\sin^5(a + bx)}{5b} - \frac{2 \sin^3(a + bx)}{3b} + \frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^5,x]

[Out] Sin[a + b*x]/b - (2*Sin[a + b*x]^3)/(3*b) + Sin[a + b*x]^5/(5*b)

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \cos^5(a + bx) dx &= -\frac{\text{Subst}\left(\int (1 - 2x^2 + x^4) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{\sin(a + bx)}{b} - \frac{2 \sin^3(a + bx)}{3b} + \frac{\sin^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A] time = 0.01, size = 41, normalized size = 1.00

$$\frac{\sin^5(a + bx)}{5b} - \frac{2 \sin^3(a + bx)}{3b} + \frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^5,x]

[Out] Sin[a + b*x]/b - (2*Sin[a + b*x]^3)/(3*b) + Sin[a + b*x]^5/(5*b)

fricas [A] time = 0.40, size = 33, normalized size = 0.80

$$\frac{(3 \cos(bx + a)^4 + 4 \cos(bx + a)^2 + 8) \sin(bx + a)}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5,x, algorithm="fricas")

[Out] 1/15*(3*cos(b*x + a)^4 + 4*cos(b*x + a)^2 + 8)*sin(b*x + a)/b

giac [A] time = 0.24, size = 34, normalized size = 0.83

$$\frac{3 \sin(bx + a)^5 - 10 \sin(bx + a)^3 + 15 \sin(bx + a)}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5,x, algorithm="giac")

[Out] 1/15*(3*sin(b*x + a)^5 - 10*sin(b*x + a)^3 + 15*sin(b*x + a))/b

maple [A] time = 0.10, size = 32, normalized size = 0.78

$$\frac{\left(\frac{8}{3} + \cos^4(bx + a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx + a)}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^5,x)

[Out] 1/5/b*(8/3+cos(b*x+a)^4+4/3*cos(b*x+a)^2)*sin(b*x+a)

maxima [A] time = 0.31, size = 34, normalized size = 0.83

$$\frac{3 \sin(bx + a)^5 - 10 \sin(bx + a)^3 + 15 \sin(bx + a)}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5,x, algorithm="maxima")

[Out] 1/15*(3*sin(b*x + a)^5 - 10*sin(b*x + a)^3 + 15*sin(b*x + a))/b

mupad [B] time = 0.10, size = 31, normalized size = 0.76

$$\frac{\frac{\sin(a+bx)^5}{5} - \frac{2\sin(a+bx)^3}{3} + \sin(a+bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^5,x)

[Out] (sin(a + b*x) - (2*sin(a + b*x)^3)/3 + sin(a + b*x)^5/5)/b

sympy [A] time = 1.61, size = 58, normalized size = 1.41

$$\begin{cases} \frac{8 \sin^5(a+bx)}{15b} + \frac{4 \sin^3(a+bx) \cos^2(a+bx)}{3b} + \frac{\sin(a+bx) \cos^4(a+bx)}{b} & \text{for } b \neq 0 \\ x \cos^5(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**5,x)

[Out] Piecewise((8*sin(a + b*x)**5/(15*b) + 4*sin(a + b*x)**3*cos(a + b*x)**2/(3*b) + sin(a + b*x)*cos(a + b*x)**4/b, Ne(b, 0)), (x*cos(a)**5, True))

3.6 $\int \cos^6(a + bx) dx$

Optimal. Leaf size=67

$$\frac{\sin(a + bx) \cos^5(a + bx)}{6b} + \frac{5 \sin(a + bx) \cos^3(a + bx)}{24b} + \frac{5 \sin(a + bx) \cos(a + bx)}{16b} + \frac{5x}{16}$$

[Out] 5/16*x+5/16*cos(b*x+a)*sin(b*x+a)/b+5/24*cos(b*x+a)^3*sin(b*x+a)/b+1/6*cos(b*x+a)^5*sin(b*x+a)/b

Rubi [A] time = 0.03, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 8}

$$\frac{\sin(a + bx) \cos^5(a + bx)}{6b} + \frac{5 \sin(a + bx) \cos^3(a + bx)}{24b} + \frac{5 \sin(a + bx) \cos(a + bx)}{16b} + \frac{5x}{16}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^6,x]

[Out] (5*x)/16 + (5*Cos[a + b*x]*Sin[a + b*x])/(16*b) + (5*Cos[a + b*x]^3*Sin[a + b*x])/(24*b) + (Cos[a + b*x]^5*Sin[a + b*x])/(6*b)

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*Sine[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^6(a + bx) dx &= \frac{\cos^5(a + bx) \sin(a + bx)}{6b} + \frac{5}{6} \int \cos^4(a + bx) dx \\ &= \frac{5 \cos^3(a + bx) \sin(a + bx)}{24b} + \frac{\cos^5(a + bx) \sin(a + bx)}{6b} + \frac{5}{8} \int \cos^2(a + bx) dx \\ &= \frac{5 \cos(a + bx) \sin(a + bx)}{16b} + \frac{5 \cos^3(a + bx) \sin(a + bx)}{24b} + \frac{\cos^5(a + bx) \sin(a + bx)}{6b} + \frac{5}{16} \int 1 dx \\ &= \frac{5x}{16} + \frac{5 \cos(a + bx) \sin(a + bx)}{16b} + \frac{5 \cos^3(a + bx) \sin(a + bx)}{24b} + \frac{\cos^5(a + bx) \sin(a + bx)}{6b} \end{aligned}$$

Mathematica [A] time = 0.03, size = 43, normalized size = 0.64

$$\frac{45 \sin(2(a + bx)) + 9 \sin(4(a + bx)) + \sin(6(a + bx)) + 60a + 60bx}{192b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^6,x]

[Out] (60*a + 60*b*x + 45*Sin[2*(a + b*x)] + 9*Sin[4*(a + b*x)] + Sin[6*(a + b*x)])/(192*b)

fricas [A] time = 0.42, size = 46, normalized size = 0.69

$$\frac{15bx + (8 \cos(bx + a)^5 + 10 \cos(bx + a)^3 + 15 \cos(bx + a)) \sin(bx + a)}{48b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6,x, algorithm="fricas")

[Out] 1/48*(15*b*x + (8*cos(b*x + a)^5 + 10*cos(b*x + a)^3 + 15*cos(b*x + a))*sin(b*x + a))/b

giac [A] time = 0.19, size = 46, normalized size = 0.69

$$\frac{5}{16}x + \frac{\sin(6bx + 6a)}{192b} + \frac{3 \sin(4bx + 4a)}{64b} + \frac{15 \sin(2bx + 2a)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6,x, algorithm="giac")

[Out] 5/16*x + 1/192*sin(6*b*x + 6*a)/b + 3/64*sin(4*b*x + 4*a)/b + 15/64*sin(2*b*x + 2*a)/b

maple [A] time = 0.10, size = 48, normalized size = 0.72

$$\frac{\left(\cos^5(bx+a) + \frac{5\cos^3(bx+a)}{4} + \frac{15\cos(bx+a)}{8}\right)\sin(bx+a)}{6} + \frac{5bx}{16} + \frac{5a}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^6,x)

[Out] 1/b*(1/6*(cos(b*x+a)^5+5/4*cos(b*x+a)^3+15/8*cos(b*x+a))*sin(b*x+a)+5/16*b*x+5/16*a)

maxima [A] time = 0.33, size = 48, normalized size = 0.72

$$-\frac{4 \sin(2bx + 2a)^3 - 60bx - 60a - 9 \sin(4bx + 4a) - 48 \sin(2bx + 2a)}{192b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6,x, algorithm="maxima")

[Out] -1/192*(4*sin(2*b*x + 2*a)^3 - 60*b*x - 60*a - 9*sin(4*b*x + 4*a) - 48*sin(2*b*x + 2*a))/b

mupad [B] time = 0.22, size = 42, normalized size = 0.63

$$\frac{5x}{16} + \frac{\frac{15 \sin(2a+2bx)}{64} + \frac{3 \sin(4a+4bx)}{64} + \frac{\sin(6a+6bx)}{192}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6,x)

[Out] (5*x)/16 + ((15*sin(2*a + 2*b*x))/64 + (3*sin(4*a + 4*b*x))/64 + sin(6*a + 6*b*x)/192)/b

sympy [A] time = 3.11, size = 139, normalized size = 2.07

$$\left\{ \begin{array}{l} \frac{5x \sin^6(a+bx)}{16} + \frac{15x \sin^4(a+bx) \cos^2(a+bx)}{16} + \frac{15x \sin^2(a+bx) \cos^4(a+bx)}{16} + \frac{5x \cos^6(a+bx)}{16} + \frac{5 \sin^5(a+bx) \cos(a+bx)}{16b} + \frac{5 \sin^3(a+bx) \cos^3(a+bx)}{6b} \\ x \cos^6(a) \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**6,x)

[Out] Piecewise((5*x*sin(a + b*x)**6/16 + 15*x*sin(a + b*x)**4*cos(a + b*x)**2/16 + 15*x*sin(a + b*x)**2*cos(a + b*x)**4/16 + 5*x*cos(a + b*x)**6/16 + 5*sin(a + b*x)**5*cos(a + b*x)/(16*b) + 5*sin(a + b*x)**3*cos(a + b*x)**3/(6*b) + 11*sin(a + b*x)*cos(a + b*x)**5/(16*b), Ne(b, 0)), (x*cos(a)**6, True))

3.7 $\int \cos^7(a + bx) dx$

Optimal. Leaf size=54

$$-\frac{\sin^7(a + bx)}{7b} + \frac{3 \sin^5(a + bx)}{5b} - \frac{\sin^3(a + bx)}{b} + \frac{\sin(a + bx)}{b}$$

[Out] $\sin(b*x+a)/b - \sin(b*x+a)^3/b + 3/5*\sin(b*x+a)^5/b - 1/7*\sin(b*x+a)^7/b$

Rubi [A] time = 0.02, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2633}

$$-\frac{\sin^7(a + bx)}{7b} + \frac{3 \sin^5(a + bx)}{5b} - \frac{\sin^3(a + bx)}{b} + \frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^7, x]

[Out] Sin[a + b*x]/b - Sin[a + b*x]^3/b + (3*Sin[a + b*x]^5)/(5*b) - Sin[a + b*x]^7/(7*b)

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \cos^7(a + bx) dx &= -\frac{\text{Subst}\left(\int (1 - 3x^2 + 3x^4 - x^6) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{b} + \frac{3 \sin^5(a + bx)}{5b} - \frac{\sin^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A] time = 0.01, size = 54, normalized size = 1.00

$$-\frac{\sin^7(a + bx)}{7b} + \frac{3 \sin^5(a + bx)}{5b} - \frac{\sin^3(a + bx)}{b} + \frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^7, x]

[Out] Sin[a + b*x]/b - Sin[a + b*x]^3/b + (3*Sin[a + b*x]^5)/(5*b) - Sin[a + b*x]^7/(7*b)

fricas [A] time = 0.41, size = 43, normalized size = 0.80

$$\frac{(5 \cos(bx + a)^6 + 6 \cos(bx + a)^4 + 8 \cos(bx + a)^2 + 16) \sin(bx + a)}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7,x, algorithm="fricas")

[Out] 1/35*(5*cos(b*x + a)^6 + 6*cos(b*x + a)^4 + 8*cos(b*x + a)^2 + 16)*sin(b*x + a)/b

giac [A] time = 0.21, size = 44, normalized size = 0.81

$$\frac{5 \sin (bx+a)^7 - 21 \sin (bx+a)^5 + 35 \sin (bx+a)^3 - 35 \sin (bx+a)}{35 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7,x, algorithm="giac")

[Out] -1/35*(5*sin(b*x + a)^7 - 21*sin(b*x + a)^5 + 35*sin(b*x + a)^3 - 35*sin(b*x + a))/b

maple [A] time = 0.10, size = 42, normalized size = 0.78

$$\frac{\left(\frac{16}{5} + \cos^6 (bx+a) + \frac{6(\cos^4 (bx+a))}{5} + \frac{8(\cos^2 (bx+a))}{5}\right) \sin (bx+a)}{7b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^7,x)

[Out] 1/7/b*(16/5*cos(b*x+a)^6+6/5*cos(b*x+a)^4+8/5*cos(b*x+a)^2)*sin(b*x+a)

maxima [A] time = 0.69, size = 44, normalized size = 0.81

$$\frac{5 \sin (bx+a)^7 - 21 \sin (bx+a)^5 + 35 \sin (bx+a)^3 - 35 \sin (bx+a)}{35 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7,x, algorithm="maxima")

[Out] -1/35*(5*sin(b*x + a)^7 - 21*sin(b*x + a)^5 + 35*sin(b*x + a)^3 - 35*sin(b*x + a))/b

mupad [B] time = 0.09, size = 43, normalized size = 0.80

$$\frac{\sin (a+bx) \left(5 \sin (a+bx)^6 - 21 \sin (a+bx)^4 + 35 \sin (a+bx)^2 - 35\right)}{35 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^7,x)

[Out] -(sin(a + b*x)*(35*sin(a + b*x)^2 - 21*sin(a + b*x)^4 + 5*sin(a + b*x)^6 - 35))/(35*b)

sympy [A] time = 5.15, size = 78, normalized size = 1.44

$$\begin{cases} \frac{16 \sin^7 (a+bx)}{35 b} + \frac{8 \sin^5 (a+bx) \cos^2 (a+bx)}{5 b} + \frac{2 \sin^3 (a+bx) \cos^4 (a+bx)}{b} + \frac{\sin (a+bx) \cos^6 (a+bx)}{b} & \text{for } b \neq 0 \\ x \cos^7 (a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**7,x)

[Out] Piecewise((16*sin(a + b*x)**7/(35*b) + 8*sin(a + b*x)**5*cos(a + b*x)**2/(5*b) + 2*sin(a + b*x)**3*cos(a + b*x)**4/b + sin(a + b*x)*cos(a + b*x)**6/b, Ne(b, 0)), (x*cos(a)**7, True))

3.8 $\int \cos^8(a + bx) dx$

Optimal. Leaf size=88

$$\frac{\sin(a + bx) \cos^7(a + bx)}{8b} + \frac{7 \sin(a + bx) \cos^5(a + bx)}{48b} + \frac{35 \sin(a + bx) \cos^3(a + bx)}{192b} + \frac{35 \sin(a + bx) \cos(a + bx)}{128b} + \frac{35x}{128}$$

[Out] 35/128*x+35/128*cos(b*x+a)*sin(b*x+a)/b+35/192*cos(b*x+a)^3*sin(b*x+a)/b+7/48*cos(b*x+a)^5*sin(b*x+a)/b+1/8*cos(b*x+a)^7*sin(b*x+a)/b

Rubi [A] time = 0.05, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 8}

$$\frac{\sin(a + bx) \cos^7(a + bx)}{8b} + \frac{7 \sin(a + bx) \cos^5(a + bx)}{48b} + \frac{35 \sin(a + bx) \cos^3(a + bx)}{192b} + \frac{35 \sin(a + bx) \cos(a + bx)}{128b} + \frac{35x}{128}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^8, x]

[Out] (35*x)/128 + (35*Cos[a + b*x]*Sin[a + b*x])/(128*b) + (35*Cos[a + b*x]^3*Sin[a + b*x])/(192*b) + (7*Cos[a + b*x]^5*Sin[a + b*x])/(48*b) + (Cos[a + b*x]^7*Sin[a + b*x])/(8*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^8(a + bx) dx &= \frac{\cos^7(a + bx) \sin(a + bx)}{8b} + \frac{7}{8} \int \cos^6(a + bx) dx \\ &= \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} + \frac{\cos^7(a + bx) \sin(a + bx)}{8b} + \frac{35}{48} \int \cos^4(a + bx) dx \\ &= \frac{35 \cos^3(a + bx) \sin(a + bx)}{192b} + \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} + \frac{\cos^7(a + bx) \sin(a + bx)}{8b} + \frac{35x}{64} \\ &= \frac{35 \cos(a + bx) \sin(a + bx)}{128b} + \frac{35 \cos^3(a + bx) \sin(a + bx)}{192b} + \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} + \frac{35x}{128} \\ &= \frac{35x}{128} + \frac{35 \cos(a + bx) \sin(a + bx)}{128b} + \frac{35 \cos^3(a + bx) \sin(a + bx)}{192b} + \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} \end{aligned}$$

Mathematica [A] time = 0.05, size = 55, normalized size = 0.62

$$\frac{672 \sin(2(a + bx)) + 168 \sin(4(a + bx)) + 32 \sin(6(a + bx)) + 3 \sin(8(a + bx)) + 840a + 840bx}{3072b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^8, x]

[Out] $(840*a + 840*b*x + 672*\sin[2*(a + b*x)] + 168*\sin[4*(a + b*x)] + 32*\sin[6*(a + b*x)] + 3*\sin[8*(a + b*x)])/(3072*b)$

fricas [A] time = 0.41, size = 56, normalized size = 0.64

$$\frac{105 bx + (48 \cos (bx + a)^7 + 56 \cos (bx + a)^5 + 70 \cos (bx + a)^3 + 105 \cos (bx + a)) \sin (bx + a)}{384 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^8,x, algorithm="fricas")`

[Out] $1/384*(105*b*x + (48*\cos(b*x + a)^7 + 56*\cos(b*x + a)^5 + 70*\cos(b*x + a)^3 + 105*\cos(b*x + a))*\sin(b*x + a))/b$

giac [A] time = 0.21, size = 60, normalized size = 0.68

$$\frac{35}{128} x + \frac{\sin(8bx + 8a)}{1024b} + \frac{\sin(6bx + 6a)}{96b} + \frac{7 \sin(4bx + 4a)}{128b} + \frac{7 \sin(2bx + 2a)}{32b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^8,x, algorithm="giac")`

[Out] $35/128*x + 1/1024*\sin(8*b*x + 8*a)/b + 1/96*\sin(6*b*x + 6*a)/b + 7/128*\sin(4*b*x + 4*a)/b + 7/32*\sin(2*b*x + 2*a)/b$

maple [A] time = 0.10, size = 58, normalized size = 0.66

$$\frac{\left(\cos^7(bx+a) + \frac{7(\cos^5(bx+a))}{6} + \frac{35(\cos^3(bx+a))}{24} + \frac{35\cos(bx+a)}{16}\right) \sin(bx+a)}{8} + \frac{35bx}{128} + \frac{35a}{128}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^8,x)`

[Out] $1/b*(1/8*(\cos(b*x+a)^7+7/6*\cos(b*x+a)^5+35/24*\cos(b*x+a)^3+35/16*\cos(b*x+a))*\sin(b*x+a)+35/128*b*x+35/128*a)$

maxima [A] time = 0.32, size = 59, normalized size = 0.67

$$\frac{128 \sin(2bx + 2a)^3 - 840bx - 840a - 3 \sin(8bx + 8a) - 168 \sin(4bx + 4a) - 768 \sin(2bx + 2a)}{3072b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^8,x, algorithm="maxima")`

[Out] $-1/3072*(128*\sin(2*b*x + 2*a)^3 - 840*b*x - 840*a - 3*\sin(8*b*x + 8*a) - 168*\sin(4*b*x + 4*a) - 768*\sin(2*b*x + 2*a))/b$

mupad [B] time = 0.28, size = 53, normalized size = 0.60

$$\frac{35x}{128} + \frac{\frac{7 \sin(2a+2bx)}{32} + \frac{7 \sin(4a+4bx)}{128} + \frac{\sin(6a+6bx)}{96} + \frac{\sin(8a+8bx)}{1024}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^8,x)`

[Out] $(35*x)/128 + ((7*\sin(2*a + 2*b*x))/32 + (7*\sin(4*a + 4*b*x))/128 + \sin(6*a + 6*b*x)/96 + \sin(8*a + 8*b*x)/1024)/b$

sympy [A] time = 8.67, size = 184, normalized size = 2.09

$$\left\{ \begin{array}{l} \frac{35x \sin^8(a+bx)}{128} + \frac{35x \sin^6(a+bx) \cos^2(a+bx)}{32} + \frac{105x \sin^4(a+bx) \cos^4(a+bx)}{64} + \frac{35x \sin^2(a+bx) \cos^6(a+bx)}{32} + \frac{35x \cos^8(a+bx)}{128} + \frac{35 \sin^7(a+bx)}{128} \\ x \cos^8(a) \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**8,x)

[Out] Piecewise((35*x*sin(a + b*x)**8/128 + 35*x*sin(a + b*x)**6*cos(a + b*x)**2/32 + 105*x*sin(a + b*x)**4*cos(a + b*x)**4/64 + 35*x*sin(a + b*x)**2*cos(a + b*x)**6/32 + 35*x*cos(a + b*x)**8/128 + 35*sin(a + b*x)**7*cos(a + b*x)/(128*b) + 385*sin(a + b*x)**5*cos(a + b*x)**3/(384*b) + 511*sin(a + b*x)**3*cos(a + b*x)**5/(384*b) + 93*sin(a + b*x)*cos(a + b*x)**7/(128*b), Ne(b, 0)), (x*cos(a)**8, True))

3.9 $\int \cos^{\frac{7}{2}}(a + bx) dx$

Optimal. Leaf size=65

$$\frac{10F\left(\frac{1}{2}(a + bx)\middle|2\right)}{21b} + \frac{2 \sin(a + bx) \cos^{\frac{5}{2}}(a + bx)}{7b} + \frac{10 \sin(a + bx) \sqrt{\cos(a + bx)}}{21b}$$

[Out] 10/21*(cos(1/2*b*x+1/2*a)^2)^(1/2)/cos(1/2*b*x+1/2*a)*EllipticF(sin(1/2*b*x+1/2*a),2^(1/2))/b+2/7*cos(b*x+a)^(5/2)*sin(b*x+a)/b+10/21*sin(b*x+a)*cos(b*x+a)^(1/2)/b

Rubi [A] time = 0.03, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2635, 2641}

$$\frac{10F\left(\frac{1}{2}(a + bx)\middle|2\right)}{21b} + \frac{2 \sin(a + bx) \cos^{\frac{5}{2}}(a + bx)}{7b} + \frac{10 \sin(a + bx) \sqrt{\cos(a + bx)}}{21b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(7/2), x]

[Out] (10*EllipticF[(a + b*x)/2, 2])/(21*b) + (10*Sqrt[Cos[a + b*x]]*Sin[a + b*x])/(21*b) + (2*Cos[a + b*x]^(5/2)*Sin[a + b*x])/(7*b)

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x] * (b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{7}{2}}(a + bx) dx &= \frac{2 \cos^{\frac{5}{2}}(a + bx) \sin(a + bx)}{7b} + \frac{5}{7} \int \cos^{\frac{3}{2}}(a + bx) dx \\ &= \frac{10 \sqrt{\cos(a + bx)} \sin(a + bx)}{21b} + \frac{2 \cos^{\frac{5}{2}}(a + bx) \sin(a + bx)}{7b} + \frac{5}{21} \int \frac{1}{\sqrt{\cos(a + bx)}} dx \\ &= \frac{10F\left(\frac{1}{2}(a + bx)\middle|2\right)}{21b} + \frac{10 \sqrt{\cos(a + bx)} \sin(a + bx)}{21b} + \frac{2 \cos^{\frac{5}{2}}(a + bx) \sin(a + bx)}{7b} \end{aligned}$$

Mathematica [A] time = 0.27, size = 51, normalized size = 0.78

$$\frac{20F\left(\frac{1}{2}(a + bx)\middle|2\right) + (23 \sin(a + bx) + 3 \sin(3(a + bx))) \sqrt{\cos(a + bx)}}{42b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(7/2), x]

[Out] (20*EllipticF[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*(23*Sin[a + b*x] + 3*Sin[3*(a + b*x)]))/(42*b)

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left(\cos(bx + a)^{\frac{7}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(7/2), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(7/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx + a)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(7/2), x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(7/2), x)

maple [B] time = 0.20, size = 199, normalized size = 3.06

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} \left(48\left(\cos^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 120\left(\cos^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 128\left(\cos^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 21\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}\right)}{21\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(7/2), x)

[Out] -2/21*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(48*cos(1/2*b*x+1/2*a)^9-120*cos(1/2*b*x+1/2*a)^7+128*cos(1/2*b*x+1/2*a)^5-72*cos(1/2*b*x+1/2*a)^3+5*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))+16*cos(1/2*b*x+1/2*a))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx + a)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(7/2), x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(7/2), x)

mupad [B] time = 0.29, size = 42, normalized size = 0.65

$$\frac{2\cos(a + bx)^{9/2}\sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(a + bx)^2\right)}{9b\sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)^(7/2),x)
```

```
[Out] -(2*cos(a + b*x)^(9/2)*sin(a + b*x)*hypergeom([1/2, 9/4], 13/4, cos(a + b*x)  
)^2))/(9*b*(sin(a + b*x)^2)^(1/2))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)**(7/2),x)
```

```
[Out] Timed out
```

3.10 $\int \cos^{\frac{5}{2}}(a + bx) dx$

Optimal. Leaf size=42

$$\frac{6E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{5b} + \frac{2 \sin(a + bx) \cos^{\frac{3}{2}}(a + bx)}{5b}$$

[Out] $6/5 * (\cos(1/2 * b * x + 1/2 * a) \wedge 2) \wedge (1/2) / \cos(1/2 * b * x + 1/2 * a) * \text{EllipticE}(\sin(1/2 * b * x + 1/2 * a), 2 \wedge (1/2)) / b + 2/5 * \cos(b * x + a) \wedge (3/2) * \sin(b * x + a) / b$

Rubi [A] time = 0.02, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2635, 2639}

$$\frac{6E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{5b} + \frac{2 \sin(a + bx) \cos^{\frac{3}{2}}(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(5/2), x]

[Out] $(6 * \text{EllipticE}[(a + b * x) / 2, 2]) / (5 * b) + (2 * \text{Cos}[a + b * x] \wedge (3/2) * \text{Sin}[a + b * x]) / (5 * b)$

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x] * (b*SIN[c + d*x])^(n - 1)) / (d*n), x] + Dist[(b^2*(n - 1)) / n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2]) / d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{5}{2}}(a + bx) dx &= \frac{2 \cos^{\frac{3}{2}}(a + bx) \sin(a + bx)}{5b} + \frac{3}{5} \int \sqrt{\cos(a + bx)} dx \\ &= \frac{6E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{5b} + \frac{2 \cos^{\frac{3}{2}}(a + bx) \sin(a + bx)}{5b} \end{aligned}$$

Mathematica [A] time = 0.05, size = 40, normalized size = 0.95

$$\frac{6E\left(\frac{1}{2}(a + bx) \middle| 2\right) + \sin(2(a + bx)) \sqrt{\cos(a + bx)}}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(5/2), x]

[Out] $(6 * \text{EllipticE}[(a + b * x) / 2, 2] + \text{Sqrt}[\text{Cos}[a + b * x]] * \text{Sin}[2 * (a + b * x)]) / (5 * b)$

fricas [F] time = 0.40, size = 0, normalized size = 0.00

$$\text{integral}\left(\cos(bx+a)^{\frac{5}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(5/2),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(5/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx+a)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(5/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(5/2), x)

maple [B] time = 0.08, size = 202, normalized size = 4.81

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} \left(-8\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\left(\sin^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 8\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{5\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(5/2),x)

[Out] $-2/5*((2*\cos(1/2*b*x+1/2*a)^2-1)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(-8*\cos(1/2*b*x+1/2*a)*\sin(1/2*b*x+1/2*a)^6+8*\sin(1/2*b*x+1/2*a)^4*\cos(1/2*b*x+1/2*a)-3*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*b*x+1/2*a),2^{(1/2)})-2*\sin(1/2*b*x+1/2*a)^2*\cos(1/2*b*x+1/2*a))/(-2*\sin(1/2*b*x+1/2*a)^4+\sin(1/2*b*x+1/2*a)^2)^{(1/2)}/\sin(1/2*b*x+1/2*a)/(2*\cos(1/2*b*x+1/2*a)^2-1)^{(1/2)}/b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx+a)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(5/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(5/2), x)

mupad [B] time = 0.17, size = 42, normalized size = 1.00

$$\frac{2\cos(a+bx)^{7/2}\sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(a+bx)^2\right)}{7b\sqrt{\sin(a+bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(5/2),x)

[Out] $-(2*\cos(a + b*x)^{(7/2)}*\sin(a + b*x)*\text{hypergeom}([1/2, 7/4], 11/4, \cos(a + b*x)^2))/ (7*b*(\sin(a + b*x)^2)^{(1/2)})$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(5/2),x)

[Out] Timed out

3.11 $\int \cos^{\frac{3}{2}}(a + bx) dx$

Optimal. Leaf size=42

$$\frac{2F\left(\frac{1}{2}(a + bx)\middle|2\right)}{3b} + \frac{2 \sin(a + bx)\sqrt{\cos(a + bx)}}{3b}$$

[Out] $2/3*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})/b+2/3*\sin(b*x+a)*\cos(b*x+a)^{(1/2)}/b$

Rubi [A] time = 0.02, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2635, 2641}

$$\frac{2F\left(\frac{1}{2}(a + bx)\middle|2\right)}{3b} + \frac{2 \sin(a + bx)\sqrt{\cos(a + bx)}}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(3/2), x]

[Out] $(2*\text{EllipticF}[(a + b*x)/2, 2])/(3*b) + (2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/(3*b)$

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x] * (b*Ssin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Ssin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{3}{2}}(a + bx) dx &= \frac{2\sqrt{\cos(a + bx)} \sin(a + bx)}{3b} + \frac{1}{3} \int \frac{1}{\sqrt{\cos(a + bx)}} dx \\ &= \frac{2F\left(\frac{1}{2}(a + bx)\middle|2\right)}{3b} + \frac{2\sqrt{\cos(a + bx)} \sin(a + bx)}{3b} \end{aligned}$$

Mathematica [A] time = 0.04, size = 36, normalized size = 0.86

$$\frac{2\left(F\left(\frac{1}{2}(a + bx)\middle|2\right) + \sin(a + bx)\sqrt{\cos(a + bx)}\right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(3/2), x]

[Out] $(2*(\text{EllipticF}[(a + b*x)/2, 2] + \text{Sqrt}[\text{Cos}[a + b*x]]*\text{Sin}[a + b*x]))/(3*b)$

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left(\cos(bx+a)^{\frac{3}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(3/2),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx+a)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(3/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(3/2), x)

maple [B] time = 0.11, size = 179, normalized size = 4.26

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\left(4\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\cos\left(\frac{bx}{2} + \frac{a}{2}\right) + \sqrt{\frac{1 - \cos(bx+a)}{2}}\sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\right)}{3\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{2\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(3/2),x)

[Out]
$$-2/3*((2*\cos(1/2*b*x+1/2*a)^2-1)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(4*\sin(1/2*b*x+1/2*a)^4*\cos(1/2*b*x+1/2*a)+(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*\text{EllipticF}(\cos(1/2*b*x+1/2*a),2^{(1/2)})-2*\sin(1/2*b*x+1/2*a)^2*\cos(1/2*b*x+1/2*a))/(-2*\sin(1/2*b*x+1/2*a)^4+\sin(1/2*b*x+1/2*a)^2)^{(1/2)}/\sin(1/2*b*x+1/2*a)/(2*\cos(1/2*b*x+1/2*a)^2-1)^{(1/2)}/b$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx+a)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(3/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(3/2), x)

mupad [B] time = 0.08, size = 35, normalized size = 0.83

$$\frac{2F\left(\frac{a}{2} + \frac{bx}{2} \middle| 2\right)}{3b} + \frac{2\sqrt{\cos(a+bx)}\sin(a+bx)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(3/2),x)

[Out]
$$(2*\text{ellipticF}(a/2 + (b*x)/2, 2))/(3*b) + (2*\cos(a + b*x)^{(1/2)}*\sin(a + b*x))/(3*b)$$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos^{\frac{3}{2}}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(3/2), x)

[Out] Integral(cos(a + b*x)**(3/2), x)

3.12 $\int \sqrt{\cos(a + bx)} dx$

Optimal. Leaf size=16

$$\frac{2E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b}$$

[Out] $2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})/b$

Rubi [A] time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2639}

$$\frac{2E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Cos[a + b*x]], x]

[Out] (2*EllipticE[(a + b*x)/2, 2])/b

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \sqrt{\cos(a + bx)} dx = \frac{2E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b}$$

Mathematica [A] time = 0.03, size = 16, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Cos[a + b*x]], x]

[Out] (2*EllipticE[(a + b*x)/2, 2])/b

fricas [F] time = 0.42, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{\cos(bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(cos(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\cos(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(cos(b*x + a)), x)

maple [B] time = 0.06, size = 133, normalized size = 8.31

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\sqrt{-2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1}\operatorname{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(1/2),x)

[Out] 2*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\cos(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(cos(b*x + a)), x)

mupad [B] time = 0.12, size = 15, normalized size = 0.94

$$\frac{2E\left(\frac{a}{2} + \frac{bx}{2} \middle| 2\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(1/2),x)

[Out] (2*ellipticE(a/2 + (b*x)/2, 2))/b

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(1/2),x)

[Out] Integral(sqrt(cos(a + b*x)), x)

$$3.13 \quad \int \frac{1}{\sqrt{\cos(a+bx)}} dx$$

Optimal. Leaf size=16

$$\frac{2F\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

[Out] $2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})/b$

Rubi [A] time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2641}

$$\frac{2F\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[Cos[a + b*x]], x]

[Out] (2*EllipticF[(a + b*x)/2, 2])/b

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \frac{1}{\sqrt{\cos(a+bx)}} dx = \frac{2F\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

Mathematica [A] time = 0.03, size = 16, normalized size = 1.00

$$\frac{2F\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[Cos[a + b*x]], x]

[Out] (2*EllipticF[(a + b*x)/2, 2])/b

fricas [F] time = 0.39, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\sqrt{\cos(bx+a)}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(1/sqrt(cos(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\cos(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(cos(b*x + a)), x)

maple [C] time = 0.13, size = 18, normalized size = 1.12

$$\frac{2 \operatorname{am}^{-1}\left(\frac{bx}{2} + \frac{a}{2} \mid \sqrt{2}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(b*x+a)^(1/2),x)

[Out] 2/b*InverseJacobiAM(1/2*b*x+1/2*a,2^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\cos(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(cos(b*x + a)), x)

mupad [B] time = 0.10, size = 15, normalized size = 0.94

$$\frac{2 F\left(\frac{a}{2} + \frac{bx}{2} \mid 2\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^(1/2),x)

[Out] (2*ellipticF(a/2 + (b*x)/2, 2))/b

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)**(1/2),x)

[Out] Integral(1/sqrt(cos(a + b*x)), x)

$$3.14 \quad \int \frac{1}{\cos^{\frac{3}{2}}(a+bx)} dx$$

Optimal. Leaf size=38

$$\frac{2 \sin(a+bx)}{b\sqrt{\cos(a+bx)}} - \frac{2E\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

[Out] $-2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a),2^{(1/2)})/b+2*\sin(b*x+a)/b/\cos(b*x+a)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2636, 2639}

$$\frac{2 \sin(a+bx)}{b\sqrt{\cos(a+bx)}} - \frac{2E\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(-3/2), x]

[Out] $(-2*\text{EllipticE}[(a + b*x)/2, 2])/b + (2*\text{Sin}[a + b*x])/(b*\text{Sqrt}[\text{Cos}[a + b*x]])$

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{3}{2}}(a+bx)} dx &= \frac{2 \sin(a+bx)}{b\sqrt{\cos(a+bx)}} - \int \sqrt{\cos(a+bx)} dx \\ &= -\frac{2E\left(\frac{1}{2}(a+bx)\middle|2\right)}{b} + \frac{2 \sin(a+bx)}{b\sqrt{\cos(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 38, normalized size = 1.00

$$\frac{2 \sin(a+bx)}{b\sqrt{\cos(a+bx)}} - \frac{2E\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(-3/2), x]

[Out] $(-2*\text{EllipticE}[(a + b*x)/2, 2])/b + (2*\text{Sin}[a + b*x])/(b*\text{Sqrt}[\text{Cos}[a + b*x]])$

fricas [F] time = 0.42, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\cos(bx+a)^{\frac{3}{2}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(3/2),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(-3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx+a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(3/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-3/2), x)

maple [A] time = 0.08, size = 101, normalized size = 2.66

$$\frac{2\left(\sqrt{\frac{1}{2}-\frac{\cos(bx+a)}{2}}\sqrt{2\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-1}\text{EllipticE}\left(\cos\left(\frac{bx}{2}+\frac{a}{2}\right),\sqrt{2}\right)-2\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\cos\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{\sin\left(\frac{bx}{2}+\frac{a}{2}\right)\sqrt{2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-1}b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(b*x+a)^(3/2),x)

[Out] -2*((sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx+a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(3/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-3/2), x)

mupad [B] time = 0.25, size = 42, normalized size = 1.11

$$\frac{2\sin(a+bx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(a+bx)^2\right)}{b\sqrt{\cos(a+bx)}\sqrt{\sin(a+bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^(3/2),x)

[Out] (2*sin(a + b*x)*hypergeom([-1/4, 1/2], 3/4, cos(a + b*x)^2))/(b*cos(a + b*x)^(1/2)*(sin(a + b*x)^2)^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)**(3/2), x)

[Out] Integral(cos(a + b*x)**(-3/2), x)

$$3.15 \quad \int \frac{1}{\cos^{\frac{5}{2}}(a+bx)} dx$$

Optimal. Leaf size=42

$$\frac{2F\left(\frac{1}{2}(a+bx)\middle|2\right)}{3b} + \frac{2\sin(a+bx)}{3b\cos^{\frac{3}{2}}(a+bx)}$$

[Out] $2/3*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a),2^{(1/2)})/b+2/3*\sin(b*x+a)/b/\cos(b*x+a)^{(3/2)}$

Rubi [A] time = 0.02, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2636, 2641}

$$\frac{2F\left(\frac{1}{2}(a+bx)\middle|2\right)}{3b} + \frac{2\sin(a+bx)}{3b\cos^{\frac{3}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(-5/2), x]

[Out] $(2*\text{EllipticF}[(a+b*x)/2, 2])/(3*b) + (2*\text{Sin}[a+b*x])/(3*b*\text{Cos}[a+b*x]^{(3/2)})$

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{5}{2}}(a+bx)} dx &= \frac{2\sin(a+bx)}{3b\cos^{\frac{3}{2}}(a+bx)} + \frac{1}{3} \int \frac{1}{\sqrt{\cos(a+bx)}} dx \\ &= \frac{2F\left(\frac{1}{2}(a+bx)\middle|2\right)}{3b} + \frac{2\sin(a+bx)}{3b\cos^{\frac{3}{2}}(a+bx)} \end{aligned}$$

Mathematica [A] time = 0.06, size = 36, normalized size = 0.86

$$\frac{2\left(F\left(\frac{1}{2}(a+bx)\middle|2\right) + \frac{\sin(a+bx)}{\cos^{\frac{3}{2}}(a+bx)}\right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(-5/2), x]

[Out] $(2*(\text{EllipticF}[(a + b*x)/2, 2] + \text{Sin}[a + b*x]/\text{Cos}[a + b*x]^{(3/2)}))/ (3*b)$
fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\cos(bx + a)^{5/2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/cos(b*x+a)^(5/2),x, algorithm="fricas")`

[Out] `integral(cos(b*x + a)^(-5/2), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/cos(b*x+a)^(5/2),x, algorithm="giac")`

[Out] `integrate(cos(b*x + a)^(-5/2), x)`

maple [B] time = 0.12, size = 213, normalized size = 5.07

$$\frac{2\left(-2\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1} \text{EllipticF}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}\right)}{3\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/cos(b*x+a)^(5/2),x)`

[Out] $-2/3*(-2*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*\text{EllipticF}(\cos(1/2*b*x+1/2*a), 2^{(1/2)})*\sin(1/2*b*x+1/2*a)^2+(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*\text{EllipticF}(\cos(1/2*b*x+1/2*a), 2^{(1/2)})-2*\sin(1/2*b*x+1/2*a)^2*\cos(1/2*b*x+1/2*a))*((2*\cos(1/2*b*x+1/2*a)^2-1)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}/(-2*\sin(1/2*b*x+1/2*a)^4+\sin(1/2*b*x+1/2*a)^2)^{(1/2)}/(2*\cos(1/2*b*x+1/2*a)^2-1)^{(3/2)}/\sin(1/2*b*x+1/2*a)/b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/cos(b*x+a)^(5/2),x, algorithm="maxima")`

[Out] `integrate(cos(b*x + a)^(-5/2), x)`

mupad [B] time = 0.27, size = 42, normalized size = 1.00

$$\frac{2 \sin(a + bx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(a + bx)^2\right)}{3b \cos(a + bx)^{3/2} \sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/cos(a + b*x)^(5/2),x)`

[Out] `(2*sin(a + b*x)*hypergeom([-3/4, 1/2], 1/4, cos(a + b*x)^2))/(3*b*cos(a + b*x)^(3/2)*(sin(a + b*x)^2)^(1/2))`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/cos(b*x+a)**(5/2),x)`

[Out] `Integral(cos(a + b*x)**(-5/2), x)`

$$3.16 \quad \int \frac{1}{\cos^{\frac{7}{2}}(a+bx)} dx$$

Optimal. Leaf size=65

$$-\frac{6E\left(\frac{1}{2}(a+bx)\middle|2\right)}{5b} + \frac{2\sin(a+bx)}{5b\cos^{\frac{5}{2}}(a+bx)} + \frac{6\sin(a+bx)}{5b\sqrt{\cos(a+bx)}}$$

[Out] $-6/5*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a),2^{(1/2)})/b+2/5*\sin(b*x+a)/b/\cos(b*x+a)^{(5/2)}+6/5*\sin(b*x+a)/b/\cos(b*x+a)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2636, 2639}

$$-\frac{6E\left(\frac{1}{2}(a+bx)\middle|2\right)}{5b} + \frac{2\sin(a+bx)}{5b\cos^{\frac{5}{2}}(a+bx)} + \frac{6\sin(a+bx)}{5b\sqrt{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(-7/2), x]

[Out] $(-6*\text{EllipticE}[(a+b*x)/2, 2])/(5*b) + (2*\text{Sin}[a+b*x])/(5*b*\text{Cos}[a+b*x]^{(5/2)}) + (6*\text{Sin}[a+b*x])/(5*b*\text{Sqrt}[\text{Cos}[a+b*x]])$

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{7}{2}}(a+bx)} dx &= \frac{2\sin(a+bx)}{5b\cos^{\frac{5}{2}}(a+bx)} + \frac{3}{5} \int \frac{1}{\cos^{\frac{3}{2}}(a+bx)} dx \\ &= \frac{2\sin(a+bx)}{5b\cos^{\frac{5}{2}}(a+bx)} + \frac{6\sin(a+bx)}{5b\sqrt{\cos(a+bx)}} - \frac{3}{5} \int \sqrt{\cos(a+bx)} dx \\ &= -\frac{6E\left(\frac{1}{2}(a+bx)\middle|2\right)}{5b} + \frac{2\sin(a+bx)}{5b\cos^{\frac{5}{2}}(a+bx)} + \frac{6\sin(a+bx)}{5b\sqrt{\cos(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.09, size = 59, normalized size = 0.91

$$\frac{3\sin(2(a+bx)) + 2\tan(a+bx) - 6\cos^{\frac{3}{2}}(a+bx)E\left(\frac{1}{2}(a+bx)\middle|2\right)}{5b\cos^{\frac{3}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(-7/2), x]

[Out] $(-6*\text{Cos}[a + b*x]^{(3/2)}*\text{EllipticE}[(a + b*x)/2, 2] + 3*\text{Sin}[2*(a + b*x)] + 2*\text{Tan}[a + b*x])/(5*b*\text{Cos}[a + b*x]^{(3/2)})$

fricas [F] time = 0.40, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\cos(bx + a)^{\frac{7}{2}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(7/2), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(-7/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(7/2), x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-7/2), x)

maple [B] time = 0.13, size = 358, normalized size = 5.51

$$2\sqrt{-\left(-2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} \left(12 \text{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right) \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2\left(\sin^2\left(\frac{bx}{2}\right)}\right)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(b*x+a)^(7/2), x)

[Out] $\frac{2}{5}*(-(-2*\text{cos}(1/2*b*x+1/2*a)^2+1)*\text{sin}(1/2*b*x+1/2*a)^2)^{(1/2)}/(8*\text{sin}(1/2*b*x+1/2*a)^6-12*\text{sin}(1/2*b*x+1/2*a)^4+6*\text{sin}(1/2*b*x+1/2*a)^2-1)/\text{sin}(1/2*b*x+1/2*a)^3*(12*\text{EllipticE}(\text{cos}(1/2*b*x+1/2*a), 2^{(1/2)})*(\text{sin}(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\text{sin}(1/2*b*x+1/2*a)^2-1)^{(1/2)}*\text{sin}(1/2*b*x+1/2*a)^4-24*\text{cos}(1/2*b*x+1/2*a)*\text{sin}(1/2*b*x+1/2*a)^6-12*\text{EllipticE}(\text{cos}(1/2*b*x+1/2*a), 2^{(1/2)})*(\text{sin}(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\text{sin}(1/2*b*x+1/2*a)^2-1)^{(1/2)}*\text{sin}(1/2*b*x+1/2*a)^2+24*\text{sin}(1/2*b*x+1/2*a)^4*\text{cos}(1/2*b*x+1/2*a)+3*(\text{sin}(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\text{sin}(1/2*b*x+1/2*a)^2-1)^{(1/2)}*\text{EllipticE}(\text{cos}(1/2*b*x+1/2*a), 2^{(1/2)})-8*\text{sin}(1/2*b*x+1/2*a)^2*\text{cos}(1/2*b*x+1/2*a))*(-2*\text{sin}(1/2*b*x+1/2*a)^4+\text{sin}(1/2*b*x+1/2*a)^2)^{(1/2)}/(2*\text{cos}(1/2*b*x+1/2*a)^2-1)^{(1/2)}/b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(7/2), x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-7/2), x)

mupad [B] time = 0.31, size = 42, normalized size = 0.65

$$\frac{2 \sin(a + bx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(a + bx)^2\right)}{5b \cos(a + bx)^{5/2} \sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^(7/2), x)

[Out] (2*sin(a + b*x)*hypergeom([-5/4, 1/2], -1/4, cos(a + b*x)^2))/(5*b*cos(a + b*x)^(5/2)*(sin(a + b*x)^2)^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)**(7/2), x)

[Out] Timed out

3.17 $\int (c \cos(a + bx))^{7/2} dx$

Optimal. Leaf size=98

$$\frac{10c^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{21b \sqrt{c \cos(a + bx)}} + \frac{10c^3 \sin(a + bx) \sqrt{c \cos(a + bx)}}{21b} + \frac{2c \sin(a + bx) (c \cos(a + bx))^{5/2}}{7b}$$

[Out] $2/7*c*(c*\cos(b*x+a))^{(5/2)}*\sin(b*x+a)/b+10/21*c^4*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a),2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(c*\cos(b*x+a))^{(1/2)}+10/21*c^3*\sin(b*x+a)*(c*\cos(b*x+a))^{(1/2)}/b$

Rubi [A] time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 2642, 2641}

$$\frac{10c^3 \sin(a + bx) \sqrt{c \cos(a + bx)}}{21b} + \frac{10c^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{21b \sqrt{c \cos(a + bx)}} + \frac{2c \sin(a + bx) (c \cos(a + bx))^{5/2}}{7b}$$

Antiderivative was successfully verified.

[In] Int[(c*Cos[a + b*x])^(7/2), x]

[Out] $(10*c^4*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(21*b*\text{Sqrt}[c*\text{Cos}[a + b*x]]) + (10*c^3*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/(21*b) + (2*c*(c*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x])/(7*b)$

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int (c \cos(a + bx))^{7/2} dx &= \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b} + \frac{1}{7} (5c^2) \int (c \cos(a + bx))^{3/2} dx \\ &= \frac{10c^3 \sqrt{c \cos(a + bx)} \sin(a + bx)}{21b} + \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b} + \frac{1}{21} (5c^4) \int \frac{1}{\sqrt{c \cos(a + bx)}} dx \\ &= \frac{10c^3 \sqrt{c \cos(a + bx)} \sin(a + bx)}{21b} + \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b} + \frac{(5c^4 \sqrt{\cos(a + bx)})}{21 \sqrt{c \cos(a + bx)}} \\ &= \frac{10c^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{21b \sqrt{c \cos(a + bx)}} + \frac{10c^3 \sqrt{c \cos(a + bx)} \sin(a + bx)}{21b} + \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b} \end{aligned}$$

Mathematica [A] time = 0.09, size = 76, normalized size = 0.78

$$\frac{c^3 \sqrt{\cos(a+bx)} \left(20F\left(\frac{1}{2}(a+bx) \middle| 2\right) + (23 \sin(a+bx) + 3 \sin(3(a+bx))) \sqrt{\cos(a+bx)} \right)}{42b \sqrt{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Cos[a + b*x])^(7/2), x]

[Out] (c^3*Sqrt[c*Cos[a + b*x]]*(20*EllipticF[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*(23*Sin[a + b*x] + 3*Sin[3*(a + b*x)])))/(42*b*Sqrt[Cos[a + b*x]])

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \cos(bx + a)} c^3 \cos(bx + a)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(7/2), x, algorithm="fricas")

[Out] integral(sqrt(c*cos(b*x + a))*c^3*cos(b*x + a)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a))^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(7/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(7/2), x)

maple [A] time = 0.13, size = 210, normalized size = 2.14

$$\frac{2\sqrt{c\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) c^4\left(48\left(\cos^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 120\left(\cos^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 128\left(\cos^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}{21\sqrt{-c\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a))^(7/2), x)

[Out] -2/21*(c*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*c^4*(48*cos(1/2*b*x+1/2*a)^9-120*cos(1/2*b*x+1/2*a)^7+128*cos(1/2*b*x+1/2*a)^5-72*cos(1/2*b*x+1/2*a)^3+5*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))+16*cos(1/2*b*x+1/2*a))/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(c*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a))^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(7/2), x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (c \cos(a + b x))^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x))^(7/2), x)

[Out] int((c*cos(a + b*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))**(7/2), x)

[Out] Timed out

3.18 $\int (c \cos(a + bx))^{5/2} dx$

Optimal. Leaf size=70

$$\frac{6c^2 E\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \cos(a + bx)}}{5b \sqrt{\cos(a + bx)}} + \frac{2c \sin(a + bx)(c \cos(a + bx))^{3/2}}{5b}$$

[Out] $2/5*c*(c*\cos(b*x+a))^{(3/2)*\sin(b*x+a)/b+6/5*c^2*(\cos(1/2*b*x+1/2*a))^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a),2^{(1/2)})*(c*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 2640, 2639}

$$\frac{6c^2 E\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \cos(a + bx)}}{5b \sqrt{\cos(a + bx)}} + \frac{2c \sin(a + bx)(c \cos(a + bx))^{3/2}}{5b}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x])^(5/2), x]

[Out] $(6*c^2*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*\text{Sqrt}[\text{Cos}[a + b*x]]) + (2*c*(c*\text{Cos}[a + b*x])^{(3/2)*\text{Sin}[a + b*x]})/(5*b)$

Rule 2635

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*cos[c + d*x])*(b*sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[b*sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int (c \cos(a + bx))^{5/2} dx &= \frac{2c(c \cos(a + bx))^{3/2} \sin(a + bx)}{5b} + \frac{1}{5} (3c^2) \int \sqrt{c \cos(a + bx)} dx \\ &= \frac{2c(c \cos(a + bx))^{3/2} \sin(a + bx)}{5b} + \frac{(3c^2 \sqrt{c \cos(a + bx)}) \int \sqrt{\cos(a + bx)} dx}{5\sqrt{\cos(a + bx)}} \\ &= \frac{6c^2 \sqrt{c \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{5b \sqrt{\cos(a + bx)}} + \frac{2c(c \cos(a + bx))^{3/2} \sin(a + bx)}{5b} \end{aligned}$$

Mathematica [A] time = 0.08, size = 62, normalized size = 0.89

$$\frac{(c \cos(a + bx))^{5/2} \left(6E\left(\frac{1}{2}(a + bx) \middle| 2\right) + \sin(2(a + bx))\sqrt{\cos(a + bx)}\right)}{5b \cos^{\frac{5}{2}}(a + bx)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x])^(5/2), x]

[Out] ((c*cos[a + b*x])^(5/2)*(6*EllipticE[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*Sin[2*(a + b*x)]))/(5*b*cos[a + b*x]^(5/2))

fricas [F] time = 0.42, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \cos(bx + a)} c^2 \cos(bx + a)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(c*cos(b*x + a))*c^2*cos(b*x + a)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(5/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(5/2), x)

maple [B] time = 0.07, size = 213, normalized size = 3.04

$$\frac{2\sqrt{c}\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) c^3\left(-8\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\left(\sin^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 8\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\cos\left(\frac{bx}{2}\right)\right)}{5\sqrt{-c}\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a))^(5/2), x)

[Out] -2/5*(c*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*c^3*(-8*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^6+8*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)-3*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(c*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(5/2), x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (c \cos(a + bx))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*cos(a + b*x))^(5/2),x)
```

```
[Out] int((c*cos(a + b*x))^(5/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*cos(b*x+a))**(5/2),x)
```

```
[Out] Timed out
```


3.19 $\int (c \cos(a + bx))^{3/2} dx$

Optimal. Leaf size=70

$$\frac{2c^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{3b \sqrt{c \cos(a + bx)}} + \frac{2c \sin(a + bx) \sqrt{c \cos(a + bx)}}{3b}$$

[Out] $2/3*c^2*(\cos(1/2*b*x+1/2*a))^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(c*\cos(b*x+a))^{(1/2)}+2/3*c*\sin(b*x+a)*(c*\cos(b*x+a))^{(1/2)}/b$

Rubi [A] time = 0.04, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 2642, 2641}

$$\frac{2c^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{3b \sqrt{c \cos(a + bx)}} + \frac{2c \sin(a + bx) \sqrt{c \cos(a + bx)}}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Cos}[a + b*x])^{(3/2)}, x]$

[Out] $(2*c^2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(3*b*\text{Sqrt}[c*\text{Cos}[a + b*x]]) + (2*c*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/(3*b)$

Rule 2635

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c + d*x])*(b*\text{Sin}[c + d*x])^{(n - 1)}]/(d*n), x] + \text{Dist}[(b^2*(n - 1))/n, \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /;$ $\text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /;$ $\text{FreeQ}\{c, d, x\}$

Rule 2642

$\text{Int}[1/\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[c + d*x]]/\text{Sqrt}[b*\text{Sin}[c + d*x]], \text{Int}[1/\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ $\text{FreeQ}\{b, c, d, x\}$

Rubi steps

$$\begin{aligned} \int (c \cos(a + bx))^{3/2} dx &= \frac{2c \sqrt{c \cos(a + bx)} \sin(a + bx)}{3b} + \frac{1}{3} c^2 \int \frac{1}{\sqrt{c \cos(a + bx)}} dx \\ &= \frac{2c \sqrt{c \cos(a + bx)} \sin(a + bx)}{3b} + \frac{(c^2 \sqrt{\cos(a + bx)}) \int \frac{1}{\sqrt{\cos(a + bx)}} dx}{3 \sqrt{c \cos(a + bx)}} \\ &= \frac{2c^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{3b \sqrt{c \cos(a + bx)}} + \frac{2c \sqrt{c \cos(a + bx)} \sin(a + bx)}{3b} \end{aligned}$$

Mathematica [A] time = 0.04, size = 58, normalized size = 0.83

$$\frac{2(c \cos(a + bx))^{3/2} \left(F\left(\frac{1}{2}(a + bx) \middle| 2\right) + \sin(a + bx) \sqrt{\cos(a + bx)} \right)}{3b \cos^{\frac{3}{2}}(a + bx)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x])^(3/2), x]

[Out] (2*(c*cos[a + b*x])^(3/2)*(EllipticF[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*Sin[a + b*x]))/(3*b*cos[a + b*x]^(3/2))

fricas [F] time = 0.45, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \cos(bx + a)} c \cos(bx + a), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(c*cos(b*x + a))*c*cos(b*x + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(3/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(3/2), x)

maple [B] time = 0.08, size = 190, normalized size = 2.71

$$\frac{2\sqrt{c}\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) c^2\left(4\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \cos\left(\frac{bx}{2} + \frac{a}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\right)}{3\sqrt{-c}\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right) \sin\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a))^(3/2), x)

[Out] -2/3*(c*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*c^2*(4*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(c*(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(3/2), x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (c \cos(a + b x))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x))^(3/2), x)

[Out] int((c*cos(a + b*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))**(3/2), x)

[Out] Timed out

3.20 $\int \sqrt{c \cos(a + bx)} dx$

Optimal. Leaf size=38

$$\frac{2E\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \cos(a + bx)}}{b\sqrt{\cos(a + bx)}}$$

[Out] $2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*(c*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2640, 2639}

$$\frac{2E\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \cos(a + bx)}}{b\sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Cos[a + b*x]], x]

[Out] $(2*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(b*\text{Sqrt}[\text{Cos}[a + b*x]])$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x, x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \sqrt{c \cos(a + bx)} dx &= \frac{\sqrt{c \cos(a + bx)} \int \sqrt{\cos(a + bx)} dx}{\sqrt{\cos(a + bx)}} \\ &= \frac{2\sqrt{c \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b\sqrt{\cos(a + bx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 38, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \cos(a + bx)}}{b\sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Cos[a + b*x]], x]

[Out] $(2*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(b*\text{Sqrt}[\text{Cos}[a + b*x]])$

fricas [F] time = 0.47, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \cos(bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*cos(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \cos(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(c*cos(b*x + a)), x)

maple [B] time = 0.07, size = 142, normalized size = 3.74

$$\frac{2\sqrt{c\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) c\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{-2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1} \operatorname{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{\sqrt{-c\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)} \sin\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{c\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)} b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a))^(1/2),x)

[Out] 2*(c*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*c*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(c*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \cos(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*cos(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \sqrt{c \cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x))^(1/2),x)

[Out] int((c*cos(a + b*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))**(1/2),x)

[Out] Integral(sqrt(c*cos(a + b*x)), x)

$$3.21 \quad \int \frac{1}{\sqrt{c \cos(a+bx)}} dx$$

Optimal. Leaf size=38

$$\frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b\sqrt{c \cos(a+bx)}}$$

[Out] $2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(c*\cos(b*x+a))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2642, 2641}

$$\frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b\sqrt{c \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[c*Cos[a + b*x]], x]

[Out] (2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(b*Sqrt[c*Cos[a + b*x]])

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{c \cos(a+bx)}} dx &= \frac{\sqrt{\cos(a+bx)} \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{\sqrt{c \cos(a+bx)}} \\ &= \frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b\sqrt{c \cos(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 38, normalized size = 1.00

$$\frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b\sqrt{c \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[c*Cos[a + b*x]], x]

[Out] (2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(b*Sqrt[c*Cos[a + b*x]])

fricas [F] time = 0.49, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{c \cos(bx+a)}}{c \cos(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*cos(b*x + a))/(c*cos(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \cos(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(c*cos(b*x + a)), x)

maple [C] time = 0.05, size = 54, normalized size = 1.42

$$\frac{2\sqrt{2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1} \operatorname{am}^{-1}\left(\frac{bx}{2} + \frac{a}{2} \mid \sqrt{2}\right)}{b\sqrt{c\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a))^(1/2),x)

[Out] 2/b/(c*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)*(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)*InverseJacobiAM(1/2*b*x+1/2*a,2^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \cos(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(c*cos(b*x + a)), x)

mupad [B] time = 0.15, size = 33, normalized size = 0.87

$$\frac{2\sqrt{\cos(a + bx)} F\left(\frac{a}{2} + \frac{bx}{2} \mid 2\right)}{b\sqrt{c \cos(a + bx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x))^(1/2),x)

[Out] (2*cos(a + b*x)^(1/2)*ellipticF(a/2 + (b*x)/2, 2))/(b*(c*cos(a + b*x))^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(1/2),x)

[Out] Integral(1/sqrt(c*cos(a + b*x)), x)

$$3.22 \quad \int \frac{1}{(c \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=68

$$\frac{2 \sin(a+bx)}{bc\sqrt{c \cos(a+bx)}} - \frac{2E\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{c \cos(a+bx)}}{bc^2\sqrt{\cos(a+bx)}}$$

[Out] $2*\sin(b*x+a)/b/c/(c*\cos(b*x+a))^{(1/2)}-2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*(c*\cos(b*x+a))^{(1/2)}/b/c^{(1/2)}/\cos(b*x+a)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2636, 2640, 2639}

$$\frac{2 \sin(a+bx)}{bc\sqrt{c \cos(a+bx)}} - \frac{2E\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{c \cos(a+bx)}}{bc^2\sqrt{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Cos[a + b*x])^(-3/2), x]

[Out] $(-2*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(b*c^2*\text{Sqrt}[\text{Cos}[a + b*x]]) + (2*\text{Sin}[a + b*x])/(b*c*\text{Sqrt}[c*\text{Cos}[a + b*x]])$

Rule 2636

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(c \cos(a+bx))^{3/2}} dx &= \frac{2 \sin(a+bx)}{bc\sqrt{c \cos(a+bx)}} - \frac{\int \sqrt{c \cos(a+bx)} dx}{c^2} \\ &= \frac{2 \sin(a+bx)}{bc\sqrt{c \cos(a+bx)}} - \frac{\sqrt{c \cos(a+bx)} \int \sqrt{\cos(a+bx)} dx}{c^2\sqrt{\cos(a+bx)}} \\ &= -\frac{2\sqrt{c \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{bc^2\sqrt{\cos(a+bx)}} + \frac{2 \sin(a+bx)}{bc\sqrt{c \cos(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 50, normalized size = 0.74

$$\frac{2 \left(\sin(a + bx) - \sqrt{\cos(a + bx)} E \left(\frac{1}{2}(a + bx) \middle| 2 \right) \right)}{bc\sqrt{c \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Cos[a + b*x])^(-3/2), x]

[Out] (2*(-(Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]) + Sin[a + b*x]))/(b*c*Sqrt[c*Cos[a + b*x]])

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{c \cos(bx + a)}}{c^2 \cos(bx + a)^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(c*cos(b*x + a))/(c^2*cos(b*x + a)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(3/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(3/2), x)

maple [A] time = 0.11, size = 168, normalized size = 2.47

$$\frac{2\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)c + c\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\left(\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}\text{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}{c\sqrt{-c\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{c\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a))^(3/2), x)

[Out] -2/c*(-2*sin(1/2*b*x+1/2*a)^4*c+c*sin(1/2*b*x+1/2*a)^2)^(1/2)*((sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/sin(1/2*b*x+1/2*a)/(c*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(3/2), x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(c \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x))^(3/2), x)

[Out] int(1/(c*cos(a + b*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))**(3/2), x)

[Out] Integral((c*cos(a + b*x))**(-3/2), x)

$$3.23 \quad \int \frac{1}{(c \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{3bc^2\sqrt{c \cos(a+bx)}} + \frac{2 \sin(a+bx)}{3bc(c \cos(a+bx))^{3/2}}$$

[Out] $2/3*\sin(b*x+a)/b/c/(c*\cos(b*x+a))^{(3/2)}+2/3*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/c^{(1/2)}/(c*\cos(b*x+a))^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2636, 2642, 2641}

$$\frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{3bc^2\sqrt{c \cos(a+bx)}} + \frac{2 \sin(a+bx)}{3bc(c \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Cos}[a + b*x])^{(-5/2)}, x]$

[Out] $(2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(3*b*c^2*\text{Sqrt}[c*\text{Cos}[a + b*x]]) + (2*\text{Sin}[a + b*x])/(3*b*c*(c*\text{Cos}[a + b*x])^{(3/2)})$

Rule 2636

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x])*(b*\text{Sin}[c + d*x])^{(n + 1)}]/(b*d*(n + 1)), x] + \text{Dist}[(n + 2)/(b^2*(n + 1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n + 2)}, x], x] /;$ $\text{FreeQ}\{b, c, d\}, x \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /;$ $\text{FreeQ}\{c, d\}, x]$

Rule 2642

$\text{Int}[1/\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[c + d*x]]/\text{Sqrt}[b*\text{Sin}[c + d*x]], \text{Int}[1/\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ $\text{FreeQ}\{b, c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{1}{(c \cos(a+bx))^{5/2}} dx &= \frac{2 \sin(a+bx)}{3bc(c \cos(a+bx))^{3/2}} + \frac{\int \frac{1}{\sqrt{c \cos(a+bx)}} dx}{3c^2} \\ &= \frac{2 \sin(a+bx)}{3bc(c \cos(a+bx))^{3/2}} + \frac{\sqrt{\cos(a+bx)} \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{3c^2\sqrt{c \cos(a+bx)}} \\ &= \frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{3bc^2\sqrt{c \cos(a+bx)}} + \frac{2 \sin(a+bx)}{3bc(c \cos(a+bx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 51, normalized size = 0.71

$$\frac{2 \left(\tan(a + bx) + \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \right)}{3bc^2 \sqrt{c \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Cos[a + b*x])^(-5/2), x]

[Out] (2*(Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + Tan[a + b*x]))/(3*b*c^2*Sqrt[c*Cos[a + b*x]])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{c \cos(bx + a)}}{c^3 \cos(bx + a)^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(c*cos(b*x + a))/(c^3*cos(b*x + a)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(5/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(5/2), x)

maple [B] time = 0.21, size = 241, normalized size = 3.35

$$\frac{2 \left(-2 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right), \sqrt{2} \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \right)}{3c^2 \sqrt{-c \left(2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a))^(5/2), x)

[Out] -2/3*(-2*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))*sin(1/2*b*x+1/2*a)^2+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/c^2*(c*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/(2*cos(1/2*b*x+1/2*a)^2-1)/sin(1/2*b*x+1/2*a)/(c*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(5/2), x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(-5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(c \cos(a + b x))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x))^(5/2), x)

[Out] int(1/(c*cos(a + b*x))^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(a + b x))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(5/2), x)

[Out] Integral((c*cos(a + b*x))^(-5/2), x)

$$3.24 \quad \int \frac{1}{(c \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=100

$$-\frac{6E\left(\frac{1}{2}(a+bx)\middle|2\right)\sqrt{c\cos(a+bx)}}{5bc^4\sqrt{\cos(a+bx)}} + \frac{6\sin(a+bx)}{5bc^3\sqrt{c\cos(a+bx)}} + \frac{2\sin(a+bx)}{5bc(c\cos(a+bx))^{5/2}}$$

[Out] $2/5*\sin(b*x+a)/b/c/(c*\cos(b*x+a))^{(5/2)}+6/5*\sin(b*x+a)/b/c^3/(c*\cos(b*x+a))^{(1/2)}-6/5*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a),2^{(1/2)})*(c*\cos(b*x+a))^{(1/2)}/b/c^4/\cos(b*x+a)^{(1/2)}$

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2636, 2640, 2639}

$$\frac{6\sin(a+bx)}{5bc^3\sqrt{c\cos(a+bx)}} - \frac{6E\left(\frac{1}{2}(a+bx)\middle|2\right)\sqrt{c\cos(a+bx)}}{5bc^4\sqrt{\cos(a+bx)}} + \frac{2\sin(a+bx)}{5bc(c\cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x])^(-7/2), x]

[Out] $(-6*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*c^4*\text{Sqrt}[\text{Cos}[a + b*x]]) + (2*\text{Sin}[a + b*x])/(5*b*c*(c*\text{Cos}[a + b*x])^{(5/2)}) + (6*\text{Sin}[a + b*x])/(5*b*c^3*\text{Sqrt}[c*\text{Cos}[a + b*x]])$

Rule 2636

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(c \cos(a+bx))^{7/2}} dx &= \frac{2\sin(a+bx)}{5bc(c\cos(a+bx))^{5/2}} + \frac{3 \int \frac{1}{(c\cos(a+bx))^{3/2}} dx}{5c^2} \\ &= \frac{2\sin(a+bx)}{5bc(c\cos(a+bx))^{5/2}} + \frac{6\sin(a+bx)}{5bc^3\sqrt{c\cos(a+bx)}} - \frac{3 \int \sqrt{c\cos(a+bx)} dx}{5c^4} \\ &= \frac{2\sin(a+bx)}{5bc(c\cos(a+bx))^{5/2}} + \frac{6\sin(a+bx)}{5bc^3\sqrt{c\cos(a+bx)}} - \frac{(3\sqrt{c\cos(a+bx)}) \int \sqrt{\cos(a+bx)} dx}{5c^4\sqrt{\cos(a+bx)}} \\ &= -\frac{6\sqrt{c\cos(a+bx)} E\left(\frac{1}{2}(a+bx)\middle|2\right)}{5bc^4\sqrt{\cos(a+bx)}} + \frac{2\sin(a+bx)}{5bc(c\cos(a+bx))^{5/2}} + \frac{6\sin(a+bx)}{5bc^3\sqrt{c\cos(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.09, size = 68, normalized size = 0.68

$$\frac{6 \sin(a + bx) - 6\sqrt{\cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right) + 2 \tan(a + bx) \sec(a + bx)}{5bc^3\sqrt{c \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Cos[a + b*x])^(-7/2), x]

[Out] (-6*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2] + 6*Sin[a + b*x] + 2*Sec[a + b*x]*Tan[a + b*x])/(5*b*c^3*Sqrt[c*Cos[a + b*x]])

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{c \cos(bx + a)}}{c^4 \cos(bx + a)^4}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(7/2), x, algorithm="fricas")

[Out] integral(sqrt(c*cos(b*x + a))/(c^4*cos(b*x + a)^4), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(7/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(7/2), x)

maple [B] time = 0.20, size = 366, normalized size = 3.66

$$2\sqrt{c \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} \left(12 \text{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right) \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a))^(7/2), x)

[Out] 2/5*(c*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/c^4/sin(1/2*b*x+1/2*a)^3/(8*sin(1/2*b*x+1/2*a)^6-12*sin(1/2*b*x+1/2*a)^4+6*sin(1/2*b*x+1/2*a)^2-1)*(12*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*sin(1/2*b*x+1/2*a)^4-24*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^6-12*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*sin(1/2*b*x+1/2*a)^2+24*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)+3*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))-8*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))*(-2*sin(1/2*b*x+1/2*a)^4*c+c*sin(1/2*b*x+1/2*a)^2)^(1/2)/(c*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(7/2),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(-7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(c \cos(a + bx))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x))^(7/2),x)

[Out] int(1/(c*cos(a + b*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))**(7/2),x)

[Out] Timed out

3.25 $\int \cos^{\frac{4}{3}}(a + bx) dx$

Optimal. Leaf size=53

$$\frac{3 \sin(a + bx) \cos^{\frac{7}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right)}{7b\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/7*\cos(b*x+a)^{(7/3)}*hypergeom([1/2, 7/6], [13/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2643}

$$\frac{3 \sin(a + bx) \cos^{\frac{7}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right)}{7b\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(4/3), x]

[Out] $(-3*\cos[a + b*x]^{(7/3)}*Hypergeometric2F1[1/2, 7/6, 13/6, \cos[a + b*x]^2]*\sin[a + b*x])/(7*b*\sqrt{[\sin[a + b*x]^2]})$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \cos^{\frac{4}{3}}(a + bx) dx = -\frac{3 \cos^{\frac{7}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right) \sin(a + bx)}{7b\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.06, size = 53, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(a + bx)} \cos^{\frac{7}{3}}(a + bx) \csc(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right)}{7b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(4/3), x]

[Out] $(-3*\cos[a + b*x]^{(7/3)}*Csc[a + b*x]*Hypergeometric2F1[1/2, 7/6, 13/6, \cos[a + b*x]^2]*\sqrt{[\sin[a + b*x]^2]})/(7*b)$

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left(\cos(bx + a)^{\frac{4}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(4/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(4/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx + a)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(4/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(4/3), x)

maple [F] time = 0.18, size = 0, normalized size = 0.00

$$\int \cos^{\frac{4}{3}}(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(4/3),x)

[Out] int(cos(b*x+a)^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx + a)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(4/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(4/3), x)

mupad [B] time = 0.19, size = 42, normalized size = 0.79

$$\frac{3 \cos(a + bx)^{7/3} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos(a + bx)^2\right)}{7b \sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(4/3),x)

[Out] -(3*cos(a + b*x)^(7/3)*sin(a + b*x)*hypergeom([1/2, 7/6], 13/6, cos(a + b*x)^2))/(7*b*(sin(a + b*x)^2)^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(4/3),x)

[Out] Timed out

3.26 $\int \cos^{\frac{2}{3}}(a + bx) dx$

Optimal. Leaf size=53

$$\frac{3 \sin(a + bx) \cos^{\frac{5}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right)}{5b\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/5*\cos(b*x+a)^{(5/3)}*\text{hypergeom}([1/2, 5/6], [11/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2643}

$$\frac{3 \sin(a + bx) \cos^{\frac{5}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right)}{5b\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(2/3), x]

[Out] $(-3*\text{Cos}[a + b*x]^{(5/3)}*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(5*b*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \cos^{\frac{2}{3}}(a + bx) dx = -\frac{3 \cos^{\frac{5}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right) \sin(a + bx)}{5b\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.03, size = 53, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(a + bx)} \cos^{\frac{5}{3}}(a + bx) \csc(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(2/3), x]

[Out] $(-3*\text{Cos}[a + b*x]^{(5/3)}*\text{Csc}[a + b*x]*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[a + b*x]^2]*\text{Sqrt}[\text{Sin}[a + b*x]^2])/(5*b)$

fricas [F] time = 0.42, size = 0, normalized size = 0.00

$$\text{integral}\left(\cos(bx + a)^{\frac{2}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(2/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(2/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(2/3), x)

maple [F] time = 0.23, size = 0, normalized size = 0.00

$$\int \cos^{\frac{2}{3}}(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(2/3),x)

[Out] int(cos(b*x+a)^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(2/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(2/3), x)

mupad [B] time = 0.19, size = 42, normalized size = 0.79

$$\frac{3 \cos(a + bx)^{5/3} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos(a + bx)^2\right)}{5b \sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(2/3),x)

[Out] -(3*cos(a + b*x)^(5/3)*sin(a + b*x)*hypergeom([1/2, 5/6], 11/6, cos(a + b*x)^2))/(5*b*(sin(a + b*x)^2)^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos^{\frac{2}{3}}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(2/3),x)

[Out] Integral(cos(a + b*x)**(2/3), x)

3.27 $\int \sqrt[3]{\cos(a + bx)} dx$

Optimal. Leaf size=53

$$\frac{3 \sin(a + bx) \cos^{\frac{4}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a + bx)\right)}{4b\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/4*\cos(b*x+a)^{(4/3)}*hypergeom([1/2, 2/3], [5/3], \cos(b*x+a)^2)*\sin(b*x+a)/b$
 $/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2643}

$$\frac{3 \sin(a + bx) \cos^{\frac{4}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a + bx)\right)}{4b\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(1/3), x]

[Out] $(-3*\cos[a + b*x]^{(4/3)}*Hypergeometric2F1[1/2, 2/3, 5/3, \cos[a + b*x]^2]*\sin[a + b*x])/(4*b*\sqrt{\sin[a + b*x]^2})$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \sqrt[3]{\cos(a + bx)} dx = -\frac{3 \cos^{\frac{4}{3}}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a + bx)\right) \sin(a + bx)}{4b\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.03, size = 53, normalized size = 1.00

$$-\frac{3\sqrt{\sin^2(a + bx)} \cos^{\frac{4}{3}}(a + bx) \csc(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a + bx)\right)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(1/3), x]

[Out] $(-3*\cos[a + b*x]^{(4/3)}*Csc[a + b*x]*Hypergeometric2F1[1/2, 2/3, 5/3, \cos[a + b*x]^2]*\sqrt{\sin[a + b*x]^2})/(4*b)$

fricas [F] time = 0.48, size = 0, normalized size = 0.00

$$\text{integral}\left(\cos(bx + a)^{\frac{1}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(1/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(1/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx + a)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(1/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(1/3), x)

maple [F] time = 0.12, size = 0, normalized size = 0.00

$$\int \cos^{\frac{1}{3}}(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(1/3),x)

[Out] int(cos(b*x+a)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx + a)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(1/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(1/3), x)

mupad [B] time = 0.18, size = 42, normalized size = 0.79

$$\frac{3 \cos(a + bx)^{4/3} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos(a + bx)^2\right)}{4b \sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(1/3),x)

[Out] -(3*cos(a + b*x)^(4/3)*sin(a + b*x)*hypergeom([1/2, 2/3], 5/3, cos(a + b*x)^2))/(4*b*(sin(a + b*x)^2)^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{\cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(1/3),x)

[Out] Integral(cos(a + b*x)**(1/3), x)

$$3.28 \quad \int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx$$

Optimal. Leaf size=53

$$-\frac{3 \sin(a+bx) \cos^{\frac{2}{3}}(a+bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right)}{2b\sqrt{\sin^2(a+bx)}}$$

[Out] $-3/2*\cos(b*x+a)^{(2/3)}*\text{hypergeom}([1/3, 1/2], [4/3], \cos(b*x+a)^2)*\sin(b*x+a)/b$
 $/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2643}

$$-\frac{3 \sin(a+bx) \cos^{\frac{2}{3}}(a+bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right)}{2b\sqrt{\sin^2(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(-1/3), x]

[Out] $(-3*\text{Cos}[a + b*x]^{(2/3)}*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(2*b*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx = -\frac{3 \cos^{\frac{2}{3}}(a+bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right) \sin(a+bx)}{2b\sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.02, size = 53, normalized size = 1.00

$$-\frac{3\sqrt{\sin^2(a+bx)} \cos^{\frac{2}{3}}(a+bx) \csc(a+bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(-1/3), x]

[Out] $(-3*\text{Cos}[a + b*x]^{(2/3)}*\text{Csc}[a + b*x]*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[a + b*x]^2]*\text{Sqrt}[\text{Sin}[a + b*x]^2])/(2*b)$

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\cos(bx+a)^{\frac{1}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(1/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(-1/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(1/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-1/3), x)

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(b*x+a)^(1/3),x)

[Out] int(1/cos(b*x+a)^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(1/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-1/3), x)

mupad [B] time = 0.21, size = 42, normalized size = 0.79

$$\frac{3 \cos(a + bx)^{2/3} \sin(a + bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos(a + bx)^2\right)}{2b \sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^(1/3),x)

[Out] -(3*cos(a + b*x)^(2/3)*sin(a + b*x)*hypergeom([1/3, 1/2], 4/3, cos(a + b*x)^2))/(2*b*(sin(a + b*x)^2)^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{\cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)**(1/3),x)

[Out] Integral(cos(a + b*x)**(-1/3), x)

$$3.29 \quad \int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx$$

Optimal. Leaf size=51

$$\frac{3 \sin(a+bx) \sqrt[3]{\cos(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a+bx)\right)}{b \sqrt{\sin^2(a+bx)}}$$

[Out] $-3 \cos(b*x+a)^{(1/3)} * \text{hypergeom}([1/6, 1/2], [7/6], \cos(b*x+a)^2) * \sin(b*x+a) / b / (\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2643}

$$\frac{3 \sin(a+bx) \sqrt[3]{\cos(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a+bx)\right)}{b \sqrt{\sin^2(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(-2/3), x]

[Out] $(-3 * \text{Cos}[a + b*x]^{(1/3)} * \text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[a + b*x]^2] * \text{Sin}[a + b*x]) / (b * \text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2]) / (b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx = - \frac{3 \sqrt[3]{\cos(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a+bx)\right) \sin(a+bx)}{b \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.02, size = 51, normalized size = 1.00

$$- \frac{3 \sqrt{\sin^2(a+bx)} \sqrt[3]{\cos(a+bx)} \csc(a+bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a+bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(-2/3), x]

[Out] $(-3 * \text{Cos}[a + b*x]^{(1/3)} * \text{Csc}[a + b*x] * \text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[a + b*x]^2] * \text{Sqrt}[\text{Sin}[a + b*x]^2]) / b$

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\cos^{\frac{2}{3}}(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(2/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(-2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(2/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-2/3), x)

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(b*x+a)^(2/3),x)

[Out] int(1/cos(b*x+a)^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(2/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-2/3), x)

mupad [B] time = 0.20, size = 42, normalized size = 0.82

$$\frac{3 \cos(a + bx)^{1/3} \sin(a + bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos(a + bx)^2\right)}{b \sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^(2/3),x)

[Out] -(3*cos(a + b*x)^(1/3)*sin(a + b*x)*hypergeom([1/6, 1/2], 7/6, cos(a + b*x)^2))/(b*(sin(a + b*x)^2)^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos^{\frac{2}{3}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)**(2/3),x)

[Out] Integral(cos(a + b*x)**(-2/3), x)

$$3.30 \quad \int \frac{1}{\cos^{\frac{4}{3}}(a+bx)} dx$$

Optimal. Leaf size=51

$$\frac{3 \sin(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a+bx)\right)}{b \sqrt{\sin^2(a+bx)} \sqrt[3]{\cos(a+bx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(b*x+a)^2)*sin(b*x+a)/b/cos(b*x+a)^(1/3)/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2643}

$$\frac{3 \sin(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a+bx)\right)}{b \sqrt{\sin^2(a+bx)} \sqrt[3]{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(-4/3), x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sin[a + b*x])/(b*Cos[a + b*x]^(1/3)*Sqrt[Sin[a + b*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{\cos^{\frac{4}{3}}(a+bx)} dx = \frac{{}_3F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a+bx)\right) \sin(a+bx)}{b \sqrt[3]{\cos(a+bx)} \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.02, size = 51, normalized size = 1.00

$$\frac{3 \sqrt{\sin^2(a+bx)} \csc(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a+bx)\right)}{b \sqrt[3]{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(-4/3), x]

[Out] (3*Csc[a + b*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*Cos[a + b*x]^(1/3))

fricas [F] time = 0.42, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\cos(bx+a)^{\frac{4}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(4/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(-4/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(4/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-4/3), x)

maple [F] time = 0.05, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(b*x+a)^(4/3),x)

[Out] int(1/cos(b*x+a)^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos(bx + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)^(4/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-4/3), x)

mupad [B] time = 0.23, size = 42, normalized size = 0.82

$$\frac{3 \sin(a + bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos(a + bx)^2\right)}{b \cos(a + bx)^{1/3} \sqrt{\sin(a + bx)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^(4/3),x)

[Out] (3*sin(a + b*x)*hypergeom([-1/6, 1/2], 5/6, cos(a + b*x)^2))/(b*cos(a + b*x)^(1/3)*(sin(a + b*x)^2)^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(b*x+a)**(4/3),x)

[Out] Integral(cos(a + b*x)**(-4/3), x)

3.31 $\int (c \cos(a + bx))^{4/3} dx$

Optimal. Leaf size=58

$$-\frac{3 \sin(a + bx)(c \cos(a + bx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right)}{7bc\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/7*(c*\cos(b*x+a))^{(7/3)}*\text{hypergeom}([1/2, 7/6], [13/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$-\frac{3 \sin(a + bx)(c \cos(a + bx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right)}{7bc\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Cos[a + b*x])^(4/3), x]

[Out] $(-3*(c*\text{Cos}[a + b*x])^{(7/3)}*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(7*b*c*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (c \cos(a + bx))^{4/3} dx = -\frac{3(c \cos(a + bx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right) \sin(a + bx)}{7bc\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.05, size = 55, normalized size = 0.95

$$-\frac{3\sqrt{\sin^2(a + bx)} \cot(a + bx)(c \cos(a + bx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right)}{7b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Cos[a + b*x])^(4/3), x]

[Out] $(-3*(c*\text{Cos}[a + b*x])^{(4/3)}*\text{Cot}[a + b*x]*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[a + b*x]^2]*\text{Sqrt}[\text{Sin}[a + b*x]^2])/(7*b)$

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left((c \cos(bx + a))^{1/3} c \cos(bx + a), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(4/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(1/3)*c*cos(b*x + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(4/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(4/3), x)

maple [F] time = 0.06, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a))^(4/3),x)

[Out] int((c*cos(b*x+a))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(4/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int (c \cos (a + bx))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x))^(4/3),x)

[Out] int((c*cos(a + b*x))^(4/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))**(4/3),x)

[Out] Timed out

3.32 $\int (c \cos(a + bx))^{2/3} dx$

Optimal. Leaf size=58

$$-\frac{3 \sin(a + bx)(c \cos(a + bx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right)}{5bc\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/5*(c*\cos(b*x+a))^{(5/3)}*\text{hypergeom}([1/2, 5/6], [11/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$-\frac{3 \sin(a + bx)(c \cos(a + bx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right)}{5bc\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x])^(2/3), x]

[Out] $(-3*(c*\text{Cos}[a + b*x])^{(5/3)}*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(5*b*c*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (c \cos(a + bx))^{2/3} dx = -\frac{3(c \cos(a + bx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right) \sin(a + bx)}{5bc\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.04, size = 55, normalized size = 0.95

$$-\frac{3\sqrt{\sin^2(a + bx)} \cot(a + bx)(c \cos(a + bx))^{2/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x])^(2/3), x]

[Out] $(-3*(c*\text{Cos}[a + b*x])^{(2/3)}*\text{Cot}[a + b*x]*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[a + b*x]^2]*\text{Sqrt}[\text{Sin}[a + b*x]^2])/(5*b)$

fricas [F] time = 0.39, size = 0, normalized size = 0.00

$$\text{integral}\left((c \cos(bx + a))^{2/3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(2/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(2/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(2/3), x)

maple [F] time = 0.14, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a))^(2/3),x)

[Out] int((c*cos(b*x+a))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(2/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int (c \cos (a + bx))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x))^(2/3),x)

[Out] int((c*cos(a + b*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (a + bx))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))**(2/3),x)

[Out] Integral((c*cos(a + b*x))**(2/3), x)

3.33 $\int \sqrt[3]{c \cos(a + bx)} dx$

Optimal. Leaf size=58

$$\frac{3 \sin(a + bx)(c \cos(a + bx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a + bx)\right)}{4bc\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/4*(c*\cos(b*x+a))^{(4/3)}*\text{hypergeom}([1/2, 2/3], [5/3], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$\frac{3 \sin(a + bx)(c \cos(a + bx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a + bx)\right)}{4bc\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x])^(1/3), x]

[Out] $(-3*(c*\cos[a + b*x])^{(4/3)}*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \cos[a + b*x]^2]*\sin[a + b*x])/(4*b*c*\text{Sqrt}[\sin[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \sqrt[3]{c \cos(a + bx)} dx = -\frac{3(c \cos(a + bx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a + bx)\right) \sin(a + bx)}{4bc\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.03, size = 55, normalized size = 0.95

$$-\frac{3\sqrt{\sin^2(a + bx)} \cot(a + bx) \sqrt[3]{c \cos(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a + bx)\right)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x])^(1/3), x]

[Out] $(-3*(c*\cos[a + b*x])^{(1/3)}*\text{Cot}[a + b*x]*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \cos[a + b*x]^2]*\text{Sqrt}[\sin[a + b*x]^2])/(4*b)$

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}\left((c \cos(bx + a))^{\frac{1}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(1/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(1/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(1/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(1/3), x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a))^(1/3),x)

[Out] int((c*cos(b*x+a))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^(1/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int (c \cos (a + bx))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x))^(1/3),x)

[Out] int((c*cos(a + b*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{c \cos (a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))**(1/3),x)

[Out] Integral((c*cos(a + b*x))**(1/3), x)

$$3.34 \quad \int \frac{1}{\sqrt[3]{c \cos(a+bx)}} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(a+bx)(c \cos(a+bx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right)}{2bc\sqrt{\sin^2(a+bx)}}$$

[Out] -3/2*(c*cos(b*x+a))^(2/3)*hypergeom([1/3, 1/2], [4/3], cos(b*x+a)^2)*sin(b*x+a)/b/c/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$\frac{3 \sin(a+bx)(c \cos(a+bx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right)}{2bc\sqrt{\sin^2(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x])^(-1/3), x]

[Out] (-3*(c*cos[a + b*x])^(2/3)*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[a + b*x]^2]*Sin[a + b*x])/(2*b*c*Sqrt[Sin[a + b*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{\sqrt[3]{c \cos(a+bx)}} dx = -\frac{3(c \cos(a+bx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right) \sin(a+bx)}{2bc\sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.04, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\sin^2(a+bx)} \cot(a+bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right)}{2b\sqrt[3]{c \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x])^(-1/3), x]

[Out] (-3*Cot[a + b*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(2*b*(c*cos[a + b*x])^(1/3))

fricas [F] time = 0.40, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(c \cos(bx + a))^{2/3}}{c \cos(bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(1/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(2/3)/(c*cos(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(1/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(-1/3), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a))^(1/3),x)

[Out] int(1/(c*cos(b*x+a))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(1/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(-1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(c \cos(a + bx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x))^(1/3),x)

[Out] int(1/(c*cos(a + b*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))**(1/3),x)

[Out] Integral((c*cos(a + b*x))**(-1/3), x)

$$3.35 \quad \int \frac{1}{(c \cos(a+bx))^{2/3}} dx$$

Optimal. Leaf size=56

$$-\frac{3 \sin(a+bx) \sqrt[3]{c \cos(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a+bx)\right)}{bc \sqrt{\sin^2(a+bx)}}$$

[Out] $-3*(c*\cos(b*x+a))^{(1/3)}*\text{hypergeom}([1/6, 1/2], [7/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$-\frac{3 \sin(a+bx) \sqrt[3]{c \cos(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a+bx)\right)}{bc \sqrt{\sin^2(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x])^(-2/3), x]

[Out] $(-3*(c*\cos[a + b*x])^{(1/3)}*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \cos[a + b*x]^2]*\sin[a + b*x])/(b*c*\text{Sqrt}[\sin[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{(c \cos(a+bx))^{2/3}} dx = -\frac{3 \sqrt[3]{c \cos(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a+bx)\right) \sin(a+bx)}{bc \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.04, size = 53, normalized size = 0.95

$$-\frac{3 \sqrt{\sin^2(a+bx)} \cot(a+bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a+bx)\right)}{b(c \cos(a+bx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x])^(-2/3), x]

[Out] $(-3*\text{Cot}[a + b*x]*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \cos[a + b*x]^2]*\text{Sqrt}[\sin[a + b*x]^2])/(b*(c*\cos[a + b*x])^{(2/3)})$

fricas [F] time = 0.40, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(c \cos(bx + a))^{1/3}}{c \cos(bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(2/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(1/3)/(c*cos(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(2/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(-2/3), x)

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a))^(2/3),x)

[Out] int(1/(c*cos(b*x+a))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(2/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(-2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(c \cos(a + bx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x))^(2/3),x)

[Out] int(1/(c*cos(a + b*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(a + bx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))**(2/3),x)

[Out] Integral((c*cos(a + b*x))**(-2/3), x)

$$3.36 \quad \int \frac{1}{(c \cos(a+bx))^{4/3}} dx$$

Optimal. Leaf size=56

$$\frac{3 \sin(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a+bx)\right)}{bc \sqrt{\sin^2(a+bx)} \sqrt[3]{c \cos(a+bx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(b*x+a)^2)*sin(b*x+a)/b/c/(c*cos(b*x+a))^(1/3)/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$\frac{3 \sin(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a+bx)\right)}{bc \sqrt{\sin^2(a+bx)} \sqrt[3]{c \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x])^(-4/3), x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sin[a + b*x])/(b*c*(c*cos[a + b*x])^(1/3)*Sqrt[Sin[a + b*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{(c \cos(a+bx))^{4/3}} dx = \frac{3 {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a+bx)\right) \sin(a+bx)}{bc \sqrt[3]{c \cos(a+bx)} \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.04, size = 53, normalized size = 0.95

$$\frac{3 \sqrt{\sin^2(a+bx)} \cot(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a+bx)\right)}{b(c \cos(a+bx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x])^(-4/3), x]

[Out] (3*Cot[a + b*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*(c*cos[a + b*x])^(4/3))

fricas [F] time = 0.39, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(c \cos(bx + a))^{2/3}}{c^2 \cos(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(4/3),x, algorithm="fricas")
 [Out] integral((c*cos(b*x + a))^(2/3)/(c^2*cos(b*x + a)^2), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(4/3),x, algorithm="giac")
 [Out] integrate((c*cos(b*x + a))^(-4/3), x)
maple [F] time = 0.06, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a))^(4/3),x)
 [Out] int(1/(c*cos(b*x+a))^(4/3),x)
maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))^(4/3),x, algorithm="maxima")
 [Out] integrate((c*cos(b*x + a))^(-4/3), x)
mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(c \cos(a + bx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x))^(4/3),x)
 [Out] int(1/(c*cos(a + b*x))^(4/3), x)
sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(a + bx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a))**(4/3),x)
 [Out] Integral((c*cos(a + b*x))**(-4/3), x)

3.37 $\int \cos^n(a + bx) dx$

Optimal. Leaf size=64

$$-\frac{\sin(a + bx) \cos^{n+1}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{b(n+1)\sqrt{\sin^2(a + bx)}}$$

[Out] $-\cos(b*x+a)^{(1+n)}*\text{hypergeom}([1/2, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*\sin(b*x+a)/b/(1+n)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2643}

$$-\frac{\sin(a + bx) \cos^{n+1}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{b(n+1)\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^n, x]

[Out] $-\left(\cos[a + b*x]^{(1+n)}*\text{Hypergeometric2F1}[1/2, (1+n)/2, (3+n)/2, \cos[a + b*x]^2]*\sin[a + b*x]\right)/(b*(1+n)*\text{Sqrt}[\sin[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \cos^n(a + bx) dx = -\frac{\cos^{1+n}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx)}{b(1+n)\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.04, size = 64, normalized size = 1.00

$$-\frac{\sqrt{\sin^2(a + bx)} \csc(a + bx) \cos^{n+1}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{b(n+1)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^n, x]

[Out] $-\left(\cos[a + b*x]^{(1+n)}*\text{Csc}[a + b*x]*\text{Hypergeometric2F1}[1/2, (1+n)/2, (3+n)/2, \cos[a + b*x]^2]*\text{Sqrt}[\sin[a + b*x]^2]\right)/(b*(1+n))$

fricas [F] time = 0.41, size = 0, normalized size = 0.00

$$\text{integral}(\cos(bx + a)^n, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^n,x, algorithm="fricas")

[Out] integral(cos(b*x + a)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos (bx + a)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^n,x, algorithm="giac")

[Out] integrate(cos(b*x + a)^n, x)

maple [F] time = 0.39, size = 0, normalized size = 0.00

$$\int \cos^n (bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^n,x)

[Out] int(cos(b*x+a)^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos (bx + a)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^n,x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^n, x)

mupad [B] time = 0.53, size = 57, normalized size = 0.89

$$\frac{\cos (a + bx)^{n+1} \sin (a + bx) {}_2F_1\left(\frac{1}{2}, \frac{n}{2} + \frac{1}{2}; \frac{n}{2} + \frac{3}{2}; \cos (a + bx)^2\right)}{b \sqrt{\sin (a + bx)^2} (n + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^n,x)

[Out] -(cos(a + b*x)^(n + 1)*sin(a + b*x)*hypergeom([1/2, n/2 + 1/2], n/2 + 3/2, cos(a + b*x)^2))/(b*(sin(a + b*x)^2)^(1/2)*(n + 1))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos^n (a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**n,x)

[Out] Integral(cos(a + b*x)**n, x)

3.38 $\int (c \cos(a + bx))^n dx$

Optimal. Leaf size=69

$$\frac{\sin(a + bx)(c \cos(a + bx))^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bc(n+1)\sqrt{\sin^2(a + bx)}}$$

[Out] $-(c \cos(b*x+a))^{(1+n)} \text{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2}+1/2*n\right], \left[\frac{3}{2}+1/2*n\right], \cos(b*x+a)^2\right) * \sin(b*x+a) / b / c / (1+n) / (\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2643}

$$\frac{\sin(a + bx)(c \cos(a + bx))^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bc(n+1)\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Cos[a + b*x])^n, x]

[Out] $-\left(\left(\left(c \cos[a + b*x]\right)^{(1+n)} \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1+n)}{2}, \frac{(3+n)}{2}, \cos[a + b*x]^2\right] * \sin[a + b*x]\right) / (b*c*(1+n)*\text{Sqrt}[\sin[a + b*x]^2])\right)$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2]) / (b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (c \cos(a + bx))^n dx = -\frac{(c \cos(a + bx))^{1+n} {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx)}{bc(1+n)\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.04, size = 64, normalized size = 0.93

$$\frac{\sqrt{\sin^2(a + bx)} \cot(a + bx)(c \cos(a + bx))^n {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{b(n+1)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Cos[a + b*x])^n, x]

[Out] $-\left(\left(\left(c \cos[a + b*x]\right)^n \cot[a + b*x] \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1+n)}{2}, \frac{(3+n)}{2}, \cos[a + b*x]^2\right] * \text{Sqrt}[\sin[a + b*x]^2]\right) / (b*(1+n))\right)$

fricas [F] time = 0.43, size = 0, normalized size = 0.00

$$\text{integral}\left((c \cos(bx + a))^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^n, x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^n,x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^n, x)

maple [F] time = 0.31, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a))^n,x)

[Out] int((c*cos(b*x+a))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))^n,x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (c \cos (a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x))^n,x)

[Out] int((c*cos(a + b*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a))**n,x)

[Out] Integral((c*cos(a + b*x))**n, x)

3.39 $\int (a \cos^2(x))^{5/2} dx$

Optimal. Leaf size=53

$$\frac{8}{15}a^2 \tan(x)\sqrt{a \cos^2(x)} + \frac{1}{5} \tan(x) (a \cos^2(x))^{5/2} + \frac{4}{15}a \tan(x) (a \cos^2(x))^{3/2}$$

[Out] $4/15*a*(a*\cos(x)^2)^{(3/2)}*\tan(x)+1/5*(a*\cos(x)^2)^{(5/2)}*\tan(x)+8/15*a^2*(a*\cos(x)^2)^{(1/2)}*\tan(x)$

Rubi [A] time = 0.04, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3203, 3207, 2637}

$$\frac{8}{15}a^2 \tan(x)\sqrt{a \cos^2(x)} + \frac{1}{5} \tan(x) (a \cos^2(x))^{5/2} + \frac{4}{15}a \tan(x) (a \cos^2(x))^{3/2}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[x]^2)^(5/2), x]

[Out] $(8*a^2*\text{Sqrt}[a*\text{Cos}[x]^2]*\text{Tan}[x])/15 + (4*a*(a*\text{Cos}[x]^2)^{(3/2)}*\text{Tan}[x])/15 + (a*\text{Cos}[x]^2)^{(5/2)}*\text{Tan}[x])/5$

Rule 2637

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] :> Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3203

Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> -Simp[(Cot[e + f*x] * (b*Sin[e + f*x]^2)^p)/(2*f*p), x] + Dist[(b*(2*p - 1))/(2*p), Int[(b*Sin[e + f*x]^2)^(p - 1), x], x] /; FreeQ[{b, e, f}, x] && !IntegerQ[p] && GtQ[p, 1]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int (a \cos^2(x))^{5/2} dx &= \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x) + \frac{1}{5} (4a) \int (a \cos^2(x))^{3/2} dx \\ &= \frac{4}{15} a (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x) + \frac{1}{15} (8a^2) \int \sqrt{a \cos^2(x)} dx \\ &= \frac{4}{15} a (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x) + \frac{1}{15} (8a^2 \sqrt{a \cos^2(x)} \sec(x)) \int \cos(x) dx \\ &= \frac{8}{15} a^2 \sqrt{a \cos^2(x)} \tan(x) + \frac{4}{15} a (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x) \end{aligned}$$

Mathematica [A] time = 0.02, size = 36, normalized size = 0.68

$$\frac{1}{240} a^2 (150 \sin(x) + 25 \sin(3x) + 3 \sin(5x)) \sec(x) \sqrt{a \cos^2(x)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^2)^(5/2),x]

[Out] (a^2*Sqrt[a*cos[x]^2]*Sec[x]*(150*Sin[x] + 25*Sin[3*x] + 3*Sin[5*x]))/240

fricas [A] time = 0.42, size = 40, normalized size = 0.75

$$\frac{(3a^2 \cos(x)^4 + 4a^2 \cos(x)^2 + 8a^2)\sqrt{a \cos(x)^2} \sin(x)}{15 \cos(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^2)^(5/2),x, algorithm="fricas")

[Out] 1/15*(3*a^2*cos(x)^4 + 4*a^2*cos(x)^2 + 8*a^2)*sqrt(a*cos(x)^2)*sin(x)/cos(x)

giac [A] time = 0.54, size = 34, normalized size = 0.64

$$\frac{1}{15} (3a^2 \sin(x)^5 - 10a^2 \sin(x)^3 + 15a^2 \sin(x))\sqrt{a} \operatorname{sgn}(\cos(x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^2)^(5/2),x, algorithm="giac")

[Out] 1/15*(3*a^2*sin(x)^5 - 10*a^2*sin(x)^3 + 15*a^2*sin(x))*sqrt(a)*sgn(cos(x))

maple [A] time = 0.09, size = 32, normalized size = 0.60

$$\frac{a^3 \cos(x) \sin(x) (3(\cos^4(x)) + 4(\cos^2(x)) + 8)}{15\sqrt{a}(\cos^2(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^2)^(5/2),x)

[Out] 1/15*a^3*cos(x)*sin(x)*(3*cos(x)^4+4*cos(x)^2+8)/(a*cos(x)^2)^(1/2)

maxima [A] time = 0.68, size = 31, normalized size = 0.58

$$\frac{1}{240} (3a^2 \sin(5x) + 25a^2 \sin(3x) + 150a^2 \sin(x))\sqrt{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^2)^(5/2),x, algorithm="maxima")

[Out] 1/240*(3*a^2*sin(5*x) + 25*a^2*sin(3*x) + 150*a^2*sin(x))*sqrt(a)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int (a \cos(x)^2)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^2)^(5/2),x)

[Out] int((a*cos(x)^2)^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*cos(x)**2)**(5/2),x)
```

```
[Out] Timed out
```

3.40 $\int (a \cos^2(x))^{3/2} dx$

Optimal. Leaf size=34

$$\frac{1}{3} \tan(x) (a \cos^2(x))^{3/2} + \frac{2}{3} a \tan(x) \sqrt{a \cos^2(x)}$$

[Out] 1/3*(a*cos(x)^2)^(3/2)*tan(x)+2/3*a*(a*cos(x)^2)^(1/2)*tan(x)

Rubi [A] time = 0.03, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3203, 3207, 2637}

$$\frac{1}{3} \tan(x) (a \cos^2(x))^{3/2} + \frac{2}{3} a \tan(x) \sqrt{a \cos^2(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^2)^(3/2), x]

[Out] (2*a*Sqrt[a*cos[x]^2]*Tan[x])/3 + ((a*cos[x]^2)^(3/2)*Tan[x])/3

Rule 2637

Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3203

Int[((b_.)*sin[(e_.) + (f_.)*(x_.)]^2)^(p_), x_Symbol] := -Simp[(Cot[e + f*x] * (b*Ssin[e + f*x]^2)^p)/(2*f*p), x] + Dist[(b*(2*p - 1))/(2*p), Int[(b*Ssin[e + f*x]^2)^(p - 1), x], x] /; FreeQ[{b, e, f}, x] && !IntegerQ[p] && GtQ[p, 1]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p] * (b*Ssin[e + f*x]^n)^FracPart[p]) / (Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u] * (Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int (a \cos^2(x))^{3/2} dx &= \frac{1}{3} (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{3} (2a) \int \sqrt{a \cos^2(x)} dx \\ &= \frac{1}{3} (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{3} (2a \sqrt{a \cos^2(x)} \sec(x)) \int \cos(x) dx \\ &= \frac{2}{3} a \sqrt{a \cos^2(x)} \tan(x) + \frac{1}{3} (a \cos^2(x))^{3/2} \tan(x) \end{aligned}$$

Mathematica [A] time = 0.01, size = 26, normalized size = 0.76

$$\frac{1}{12} a (9 \sin(x) + \sin(3x)) \sec(x) \sqrt{a \cos^2(x)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^2)^(3/2),x]

[Out] (a*Sqrt[a*cos[x]^2]*Sec[x]*(9*Sin[x] + Sin[3*x]))/12

fricas [A] time = 0.42, size = 26, normalized size = 0.76

$$\frac{(a \cos(x)^2 + 2a)\sqrt{a \cos(x)^2} \sin(x)}{3 \cos(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^2)^(3/2),x, algorithm="fricas")

[Out] 1/3*(a*cos(x)^2 + 2*a)*sqrt(a*cos(x)^2)*sin(x)/cos(x)

giac [A] time = 0.29, size = 17, normalized size = 0.50

$$-\frac{1}{3}(\sin(x)^3 - 3 \sin(x))a^{\frac{3}{2}}\operatorname{sgn}(\cos(x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^2)^(3/2),x, algorithm="giac")

[Out] -1/3*(sin(x)^3 - 3*sin(x))*a^(3/2)*sgn(cos(x))

maple [A] time = 0.08, size = 24, normalized size = 0.71

$$\frac{a^2 \cos(x) \sin(x) (2 + \cos^2(x))}{3\sqrt{a}(\cos^2(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^2)^(3/2),x)

[Out] 1/3*a^2*cos(x)*sin(x)*(2+cos(x)^2)/(a*cos(x)^2)^(1/2)

maxima [A] time = 0.89, size = 17, normalized size = 0.50

$$\frac{1}{12}(a \sin(3x) + 9a \sin(x))\sqrt{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^2)^(3/2),x, algorithm="maxima")

[Out] 1/12*(a*sin(3*x) + 9*a*sin(x))*sqrt(a)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int (a \cos(x)^2)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^2)^(3/2),x)

[Out] int((a*cos(x)^2)^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)**2)**(3/2),x)

[Out] Timed out

3.41 $\int \sqrt{a \cos^2(x)} dx$

Optimal. Leaf size=13

$$\tan(x)\sqrt{a \cos^2(x)}$$

[Out] (a*cos(x)^2)^(1/2)*tan(x)

Rubi [A] time = 0.01, antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3207, 2637}

$$\tan(x)\sqrt{a \cos^2(x)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a*Cos[x]^2],x]

[Out] Sqrt[a*Cos[x]^2]*Tan[x]

Rule 2637

Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int \sqrt{a \cos^2(x)} dx &= \left(\sqrt{a \cos^2(x)} \sec(x) \right) \int \cos(x) dx \\ &= \sqrt{a \cos^2(x)} \tan(x) \end{aligned}$$

Mathematica [A] time = 0.00, size = 13, normalized size = 1.00

$$\tan(x)\sqrt{a \cos^2(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a*Cos[x]^2],x]

[Out] Sqrt[a*Cos[x]^2]*Tan[x]

fricas [A] time = 0.41, size = 15, normalized size = 1.15

$$\frac{\sqrt{a \cos(x)^2} \sin(x)}{\cos(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^2)^(1/2),x, algorithm="fricas")

[Out] $\sqrt{a \cos(x)^2} \sin(x) / \cos(x)$

giac [A] time = 0.39, size = 9, normalized size = 0.69

$$\sqrt{a} \operatorname{sgn}(\cos(x)) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(x)^2)^(1/2),x, algorithm="giac")`

[Out] $\sqrt{a} \operatorname{sgn}(\cos(x)) \sin(x)$

maple [A] time = 0.06, size = 15, normalized size = 1.15

$$\frac{a \cos(x) \sin(x)}{\sqrt{a (\cos^2(x))}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*cos(x)^2)^(1/2),x)`

[Out] $1/(a \cos(x)^2)^{1/2} * a \cos(x) \sin(x)$

maxima [A] time = 0.68, size = 6, normalized size = 0.46

$$\sqrt{a} \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(x)^2)^(1/2),x, algorithm="maxima")`

[Out] $\sqrt{a} \sin(x)$

mupad [B] time = 0.21, size = 46, normalized size = 3.54

$$\frac{\sqrt{2} \sqrt{a} \sqrt{\cos(2x) + 1} (\cos(2x) - 1 + \sin(2x) 1i)}{2 (\cos(2x) 1i - \sin(2x) + 1i)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*cos(x)^2)^(1/2),x)`

[Out] $(2^{1/2} * a^{1/2} * (\cos(2*x) + 1)^{1/2} * (\cos(2*x) + \sin(2*x) * 1i - 1)) / (2 * (\cos(2*x) * 1i - \sin(2*x) + 1i))$

sympy [A] time = 0.54, size = 19, normalized size = 1.46

$$\frac{\sqrt{a} \sqrt{\cos^2(x)} \sin(x)}{\cos(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(x)**2)**(1/2),x)`

[Out] $\sqrt{a} \sqrt{\cos(x)^2} \sin(x) / \cos(x)$

$$3.42 \quad \int \frac{1}{\sqrt{a \cos^2(x)}} dx$$

Optimal. Leaf size=16

$$\frac{\cos(x) \tanh^{-1}(\sin(x))}{\sqrt{a \cos^2(x)}}$$

[Out] arctanh(sin(x))*cos(x)/(a*cos(x)^2)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3207, 3770}

$$\frac{\cos(x) \tanh^{-1}(\sin(x))}{\sqrt{a \cos^2(x)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a*Cos[x]^2], x]

[Out] (ArcTanh[Sin[x]]*Cos[x])/Sqrt[a*Cos[x]^2]

Rule 3207

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

Rule 3770

```
Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a \cos^2(x)}} dx &= \frac{\cos(x) \int \sec(x) dx}{\sqrt{a \cos^2(x)}} \\ &= \frac{\tanh^{-1}(\sin(x)) \cos(x)}{\sqrt{a \cos^2(x)}} \end{aligned}$$

Mathematica [B] time = 0.02, size = 46, normalized size = 2.88

$$\frac{\cos(x) \left(\log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) - \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) \right)}{\sqrt{a \cos^2(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[a*Cos[x]^2], x]

[Out] (Cos[x]*(-Log[Cos[x/2] - Sin[x/2]] + Log[Cos[x/2] + Sin[x/2]]))/Sqrt[a*Cos[x]^2]

fricas [B] time = 0.45, size = 65, normalized size = 4.06

$$\left[\frac{\sqrt{a \cos(x)^2} \log\left(\frac{\sin(x)-1}{\sin(x)+1}\right)}{2 a \cos(x)}, \frac{\sqrt{-a} \arctan\left(\frac{\sqrt{a \cos(x)^2} \sqrt{-a} \sin(x)}{a \cos(x)}\right)}{a} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(1/2), x, algorithm="fricas")

[Out] [-1/2*sqrt(a*cos(x)^2)*log(-(sin(x) - 1)/(sin(x) + 1))/(a*cos(x)), -sqrt(-a)*arctan(sqrt(a*cos(x)^2)*sqrt(-a)*sin(x)/(a*cos(x)))/a]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \cos(x)^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(1/2), x, algorithm="giac")

[Out] integrate(1/sqrt(a*cos(x)^2), x)

maple [B] time = 0.08, size = 48, normalized size = 3.00

$$\frac{\cos(x) \sqrt{a (\sin^2(x))} \ln\left(\frac{2\sqrt{a} \sqrt{a(\sin^2(x))+2a}}{\cos(x)}\right)}{\sqrt{a} \sin(x) \sqrt{a (\cos^2(x))}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^2)^(1/2), x)

[Out] cos(x)*(a*sin(x)^2)^(1/2)/a^(1/2)*ln(2*(a^(1/2)*(a*sin(x)^2)^(1/2)+a)/cos(x))/sin(x)/(a*cos(x)^2)^(1/2)

maxima [B] time = 0.68, size = 38, normalized size = 2.38

$$\frac{\log(\cos(x)^2 + \sin(x)^2 + 2 \sin(x) + 1) - \log(\cos(x)^2 + \sin(x)^2 - 2 \sin(x) + 1)}{2 \sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(1/2), x, algorithm="maxima")

[Out] 1/2*(log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) - log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1))/sqrt(a)

mupad [F] time = 0.00, size = -1, normalized size = -0.06

$$\int \frac{1}{\sqrt{a \cos(x)^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^2)^(1/2), x)

[Out] int(1/(a*cos(x)^2)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)**2)**(1/2),x)

[Out] Integral(1/sqrt(a*cos(x)**2), x)

$$3.43 \quad \int \frac{1}{(a \cos^2(x))^{3/2}} dx$$

Optimal. Leaf size=42

$$\frac{\tan(x)}{2a\sqrt{a \cos^2(x)}} + \frac{\cos(x) \tanh^{-1}(\sin(x))}{2a\sqrt{a \cos^2(x)}}$$

[Out] 1/2*arctanh(sin(x))*cos(x)/a/(a*cos(x)^2)^(1/2)+1/2*tan(x)/a/(a*cos(x)^2)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3204, 3207, 3770}

$$\frac{\tan(x)}{2a\sqrt{a \cos^2(x)}} + \frac{\cos(x) \tanh^{-1}(\sin(x))}{2a\sqrt{a \cos^2(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^2)^(-3/2), x]

[Out] (ArcTanh[Sin[x]]*Cos[x])/(2*a*Sqrt[a*cos[x]^2]) + Tan[x]/(2*a*Sqrt[a*cos[x]^2])

Rule 3204

Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Simp[(Cot[e + f*x] * (b*Sin[e + f*x]^2)^(p + 1))/(b*f*(2*p + 1)), x] + Dist[(2*(p + 1))/(b*(2*p + 1)), Int[(b*Sin[e + f*x]^2)^(p + 1), x], x] /; FreeQ[{b, e, f}, x] && ! IntegerQ[p] && LtQ[p, -1]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && ! IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a \cos^2(x))^{3/2}} dx &= \frac{\tan(x)}{2a\sqrt{a \cos^2(x)}} + \frac{\int \frac{1}{\sqrt{a \cos^2(x)}} dx}{2a} \\ &= \frac{\tan(x)}{2a\sqrt{a \cos^2(x)}} + \frac{\cos(x) \int \sec(x) dx}{2a\sqrt{a \cos^2(x)}} \\ &= \frac{\tanh^{-1}(\sin(x)) \cos(x)}{2a\sqrt{a \cos^2(x)}} + \frac{\tan(x)}{2a\sqrt{a \cos^2(x)}} \end{aligned}$$

Mathematica [B] time = 0.06, size = 91, normalized size = 2.17

$$\frac{\cos(x) \left(-2 \sin(x) + \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + \cos(2x) \left(\log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) - \log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) \right) - \log \left(\cos \left(\frac{x}{2} \right) + \sin \left(\frac{x}{2} \right) \right)}{4 \left(a \cos^2(x) \right)^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Cos[x]^2)^(-3/2),x]

[Out] -1/4*(Cos[x]*(Log[Cos[x/2] - Sin[x/2]] + Cos[2*x]*(Log[Cos[x/2] - Sin[x/2]] - Log[Cos[x/2] + Sin[x/2]])) - Log[Cos[x/2] + Sin[x/2]] - 2*Sin[x])/(a*Cos[x]^2)^(3/2)

fricas [A] time = 0.42, size = 40, normalized size = 0.95

$$\frac{\sqrt{a \cos(x)^2} \left(\cos(x)^2 \log \left(-\frac{\sin(x)-1}{\sin(x)+1} \right) - 2 \sin(x) \right)}{4 a^2 \cos(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(3/2),x, algorithm="fricas")

[Out] -1/4*sqrt(a*cos(x)^2)*(cos(x)^2*log(-(sin(x) - 1)/(sin(x) + 1)) - 2*sin(x))/(a^2*cos(x)^3)

giac [A] time = 0.42, size = 47, normalized size = 1.12

$$\frac{\frac{\log \left(\left| -\sqrt{a} \tan(x) + \sqrt{a \tan(x)^2 + a} \right| \right)}{\sqrt{a}} - \frac{\sqrt{a \tan(x)^2 + a} \tan(x)}{a}}{2 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(3/2),x, algorithm="giac")

[Out] -1/2*(log(abs(-sqrt(a)*tan(x) + sqrt(a*tan(x)^2 + a)))/sqrt(a) - sqrt(a*tan(x)^2 + a)*tan(x)/a)/a

maple [B] time = 0.09, size = 70, normalized size = 1.67

$$\frac{\sqrt{a \left(\sin^2(x) \right)} \left(\ln \left(\frac{2 \sqrt{a} \sqrt{a \left(\sin^2(x) \right) + 2a}}{\cos(x)} \right) a \left(\cos^2(x) \right) + \sqrt{a} \sqrt{a \left(\sin^2(x) \right)} \right)}{2 a^{\frac{5}{2}} \cos(x) \sin(x) \sqrt{a \left(\cos^2(x) \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^2)^(3/2),x)

[Out] 1/2/a^(5/2)/cos(x)*(a*sin(x)^2)^(1/2)*(ln(2*(a^(1/2)*(a*sin(x)^2)^(1/2)+a)/cos(x))*a*cos(x)^2+a^(1/2)*(a*sin(x)^2)^(1/2))/sin(x)/(a*cos(x)^2)^(1/2)

maxima [B] time = 0.68, size = 304, normalized size = 7.24

$$\frac{4 \left(\sin(3x) - \sin(x) \right) \cos(4x) + \left(2 \left(2 \cos(2x) + 1 \right) \cos(4x) + \cos(4x)^2 + 4 \cos(2x)^2 + \sin(4x)^2 + 4 \sin(4x) \right)}{4 a^2 \cos(x) \sin(x) \sqrt{a \left(\cos^2(x) \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(3/2),x, algorithm="maxima")


```
[Out] 1/4*(4*(sin(3*x) - sin(x))*cos(4*x) + (2*(2*cos(2*x) + 1))*cos(4*x) + cos(4*x)^2 + 4*cos(2*x)^2 + sin(4*x)^2 + 4*sin(4*x)*sin(2*x) + 4*sin(2*x)^2 + 4*cos(2*x) + 1)*log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) - (2*(2*cos(2*x) + 1))*cos(4*x) + cos(4*x)^2 + 4*cos(2*x)^2 + sin(4*x)^2 + 4*sin(4*x)*sin(2*x) + 4*sin(2*x)^2 + 4*cos(2*x) + 1)*log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1) - 4*(cos(3*x) - cos(x))*sin(4*x) + 4*(2*cos(2*x) + 1)*sin(3*x) - 8*cos(3*x)*sin(2*x) + 8*cos(x)*sin(2*x) - 8*cos(2*x)*sin(x) - 4*sin(x))/((a*cos(4*x)^2 + 4*a*cos(2*x)^2 + a*sin(4*x)^2 + 4*a*sin(4*x)*sin(2*x) + 4*a*sin(2*x)^2 + 2*(2*a*cos(2*x) + a)*cos(4*x) + 4*a*cos(2*x) + a)*sqrt(a))
```

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(a \cos(x)^2)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(a*cos(x)^2)^(3/2), x)
```

```
[Out] int(1/(a*cos(x)^2)^(3/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \cos^2(x))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a*cos(x)**2)**(3/2), x)
```

```
[Out] Integral((a*cos(x)**2)**(-3/2), x)
```

$$3.44 \quad \int \frac{1}{(a \cos^2(x))^{5/2}} dx$$

Optimal. Leaf size=61

$$\frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{3 \cos(x) \tanh^{-1}(\sin(x))}{8a^2 \sqrt{a \cos^2(x)}} + \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}}$$

[Out] 3/8*arctanh(sin(x))*cos(x)/a^2/(a*cos(x)^2)^(1/2)+1/4*tan(x)/a/(a*cos(x)^2)^(3/2)+3/8*tan(x)/a^2/(a*cos(x)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3204, 3207, 3770}

$$\frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{3 \cos(x) \tanh^{-1}(\sin(x))}{8a^2 \sqrt{a \cos^2(x)}} + \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^2)^(-5/2),x]

[Out] (3*ArcTanh[Sin[x]]*Cos[x])/(8*a^2*Sqrt[a*cos[x]^2]) + Tan[x]/(4*a*(a*cos[x]^2)^(3/2)) + (3*Tan[x])/(8*a^2*Sqrt[a*cos[x]^2])

Rule 3204

Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Simp[(Cot[e + f*x] * (b*Sin[e + f*x]^2)^(p + 1))/(b*f*(2*p + 1)), x] + Dist[(2*(p + 1))/(b*(2*p + 1)), Int[(b*Sin[e + f*x]^2)^(p + 1), x], x] /; FreeQ[{b, e, f}, x] && ! IntegerQ[p] && LtQ[p, -1]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && ! IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a \cos^2(x))^{5/2}} dx &= \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \int \frac{1}{(a \cos^2(x))^{3/2}} dx}{4a} \\
&= \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{3 \int \frac{1}{\sqrt{a \cos^2(x)}} dx}{8a^2} \\
&= \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{(3 \cos(x)) \int \sec(x) dx}{8a^2 \sqrt{a \cos^2(x)}} \\
&= \frac{3 \tanh^{-1}(\sin(x)) \cos(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}}
\end{aligned}$$

Mathematica [A] time = 0.14, size = 72, normalized size = 1.18

$$\frac{\cos^5(x) \left(\frac{1}{2} (11 \sin(x) + 3 \sin(3x)) \sec^4(x) - 6 \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + 6 \log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) \right)}{16 (a \cos^2(x))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Cos[x]^2)^(-5/2), x]

[Out] (Cos[x]^5*(-6*Log[Cos[x/2] - Sin[x/2]] + 6*Log[Cos[x/2] + Sin[x/2]] + (Sec[x]^4*(11*Sin[x] + 3*Sin[3*x]))/2))/(16*(a*Cos[x]^2)^(5/2))

fricas [A] time = 0.47, size = 49, normalized size = 0.80

$$\frac{\left(3 \cos(x)^4 \log \left(-\frac{\sin(x)-1}{\sin(x)+1} \right) - 2 \left(3 \cos(x)^2 + 2 \right) \sin(x) \right) \sqrt{a \cos(x)^2}}{16 a^3 \cos(x)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(5/2), x, algorithm="fricas")

[Out] -1/16*(3*cos(x)^4*log(-(sin(x) - 1)/(sin(x) + 1)) - 2*(3*cos(x)^2 + 2)*sin(x))*sqrt(a*cos(x)^2)/(a^3*cos(x)^5)

giac [B] time = 0.56, size = 126, normalized size = 2.07

$$\frac{3 \log \left(\left| \frac{1}{\tan \left(\frac{1}{2} x \right)} + \tan \left(\frac{1}{2} x \right) + 2 \right| \right)}{\operatorname{sgn} \left(-\tan \left(\frac{1}{2} x \right)^2 + 1 \right)} - \frac{3 \log \left(\left| \frac{1}{\tan \left(\frac{1}{2} x \right)} + \tan \left(\frac{1}{2} x \right) - 2 \right| \right)}{\operatorname{sgn} \left(-\tan \left(\frac{1}{2} x \right)^2 + 1 \right)} + \frac{4 \left(5 \left(\frac{1}{\tan \left(\frac{1}{2} x \right)} + \tan \left(\frac{1}{2} x \right) \right)^3 - \frac{12}{\tan \left(\frac{1}{2} x \right)} - 12 \tan \left(\frac{1}{2} x \right) \right)}{\left(\left(\frac{1}{\tan \left(\frac{1}{2} x \right)} + \tan \left(\frac{1}{2} x \right) \right)^2 - 4 \right)^2 \operatorname{sgn} \left(-\tan \left(\frac{1}{2} x \right)^2 + 1 \right)}$$

$$\frac{5}{16 a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(5/2), x, algorithm="giac")

[Out] 1/16*(3*log(abs(1/tan(1/2*x) + tan(1/2*x) + 2))/sgn(-tan(1/2*x)^2 + 1) - 3*log(abs(1/tan(1/2*x) + tan(1/2*x) - 2))/sgn(-tan(1/2*x)^2 + 1) + 4*(5*(1/tan(1/2*x) + tan(1/2*x))^3 - 12/tan(1/2*x) - 12*tan(1/2*x))/(((1/tan(1/2*x) + tan(1/2*x))^2 - 4)^2*sgn(-tan(1/2*x)^2 + 1))/a^(5/2)

maple [A] time = 0.10, size = 89, normalized size = 1.46

$$\frac{\sqrt{a(\sin^2(x))} \left(3 \ln \left(\frac{2\sqrt{a} \sqrt{a(\sin^2(x))+2a}}{\cos(x)} \right) a (\cos^4(x)) + 3\sqrt{a(\sin^2(x))} (\cos^2(x)) \sqrt{a} + 2\sqrt{a} \sqrt{a(\sin^2(x))} \right)}{8a^{\frac{7}{2}} \cos(x)^3 \sin(x) \sqrt{a(\cos^2(x))}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^2)^(5/2),x)

[Out] 1/8/a^(7/2)/cos(x)^3*(a*sin(x)^2)^(1/2)*(3*ln(2*(a^(1/2)*(a*sin(x)^2)^(1/2)+a)/cos(x))*a*cos(x)^4+3*(a*sin(x)^2)^(1/2)*cos(x)^2*a^(1/2)+2*a^(1/2)*(a*sin(x)^2)^(1/2))/sin(x)/(a*cos(x)^2)^(1/2)

maxima [B] time = 1.38, size = 933, normalized size = 15.30

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^2)^(5/2),x, algorithm="maxima")

[Out] 1/16*(4*(3*sin(7*x) + 11*sin(5*x) - 11*sin(3*x) - 3*sin(x))*cos(8*x) - 24*(2*sin(6*x) + 3*sin(4*x) + 2*sin(2*x))*cos(7*x) + 16*(11*sin(5*x) - 11*sin(3*x) - 3*sin(x))*cos(6*x) - 88*(3*sin(4*x) + 2*sin(2*x))*cos(5*x) - 24*(11*sin(3*x) + 3*sin(x))*cos(4*x) + 3*(2*(4*cos(6*x) + 6*cos(4*x) + 4*cos(2*x) + 1)*cos(8*x) + cos(8*x)^2 + 8*(6*cos(4*x) + 4*cos(2*x) + 1)*cos(6*x) + 16*cos(6*x)^2 + 12*(4*cos(2*x) + 1)*cos(4*x) + 36*cos(4*x)^2 + 16*cos(2*x)^2 + 4*(2*sin(6*x) + 3*sin(4*x) + 2*sin(2*x))*sin(8*x) + sin(8*x)^2 + 16*(3*sin(4*x) + 2*sin(2*x))*sin(6*x) + 16*sin(6*x)^2 + 36*sin(4*x)^2 + 48*sin(4*x)*sin(2*x) + 16*sin(2*x)^2 + 8*cos(2*x) + 1)*log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) - 3*(2*(4*cos(6*x) + 6*cos(4*x) + 4*cos(2*x) + 1)*cos(8*x) + cos(8*x)^2 + 8*(6*cos(4*x) + 4*cos(2*x) + 1)*cos(6*x) + 16*cos(6*x)^2 + 12*(4*cos(2*x) + 1)*cos(4*x) + 36*cos(4*x)^2 + 16*cos(2*x)^2 + 4*(2*sin(6*x) + 3*sin(4*x) + 2*sin(2*x))*sin(8*x) + sin(8*x)^2 + 16*(3*sin(4*x) + 2*sin(2*x))*sin(6*x) + 16*sin(6*x)^2 + 36*sin(4*x)^2 + 48*sin(4*x)*sin(2*x) + 16*sin(2*x)^2 + 8*cos(2*x) + 1)*log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1) - 4*(3*cos(7*x) + 11*cos(5*x) - 11*cos(3*x) - 3*cos(x))*sin(8*x) + 12*(4*cos(6*x) + 6*cos(4*x) + 4*cos(2*x) + 1)*sin(7*x) - 16*(11*cos(5*x) - 11*cos(3*x) - 3*cos(x))*sin(6*x) + 44*(6*cos(4*x) + 4*cos(2*x) + 1)*sin(5*x) + 24*(11*cos(3*x) + 3*cos(x))*sin(4*x) - 44*(4*cos(2*x) + 1)*sin(3*x) + 176*cos(3*x)*sin(2*x) + 48*cos(x)*sin(2*x) - 48*cos(2*x)*sin(x) - 12*sin(x))/((a^2*cos(8*x)^2 + 16*a^2*cos(6*x)^2 + 36*a^2*cos(4*x)^2 + 16*a^2*cos(2*x)^2 + a^2*sin(8*x)^2 + 16*a^2*sin(6*x)^2 + 36*a^2*sin(4*x)^2 + 48*a^2*sin(4*x)*sin(2*x) + 16*a^2*sin(2*x)^2 + 8*a^2*cos(2*x) + a^2 + 2*(4*a^2*cos(6*x) + 6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(8*x) + 8*(6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(6*x) + 12*(4*a^2*cos(2*x) + a^2)*cos(4*x) + 4*(2*a^2*sin(6*x) + 3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(8*x) + 16*(3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(6*x))*sqrt(a))

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(a \cos(x)^2)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^2)^(5/2),x)

[Out] int(1/(a*cos(x)^2)^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)**2)**(5/2), x)

[Out] Timed out

3.45 $\int (a \cos^3(x))^{5/2} dx$

Optimal. Leaf size=117

$$\frac{26}{165}a^2 \sin(x) \cos^3(x) \sqrt{a \cos^3(x)} + \frac{78}{385}a^2 \sin(x) \cos(x) \sqrt{a \cos^3(x)} + \frac{26}{77}a^2 \tan(x) \sqrt{a \cos^3(x)} + \frac{26a^2 F\left(\frac{x}{2} \middle| 2\right) \sqrt{a \cos^3(x)}}{77 \cos^2(x)}$$

[Out] 26/77*a^2*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticF(sin(1/2*x),2^(1/2))*(a*cos(x)^3)^(1/2)/cos(x)^(3/2)+78/385*a^2*cos(x)*sin(x)*(a*cos(x)^3)^(1/2)+26/165*a^2*cos(x)^3*sin(x)*(a*cos(x)^3)^(1/2)+2/15*a^2*cos(x)^5*sin(x)*(a*cos(x)^3)^(1/2)+26/77*a^2*(a*cos(x)^3)^(1/2)*tan(x)

Rubi [A] time = 0.07, antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2635, 2641}

$$\frac{2}{15}a^2 \sin(x) \cos^5(x) \sqrt{a \cos^3(x)} + \frac{26}{165}a^2 \sin(x) \cos^3(x) \sqrt{a \cos^3(x)} + \frac{78}{385}a^2 \sin(x) \cos(x) \sqrt{a \cos^3(x)} + \frac{26}{77}a^2 \tan(x) \sqrt{a \cos^3(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^3)^(5/2),x]

[Out] (26*a^2*Sqrt[a*cos[x]^3]*EllipticF[x/2, 2])/(77*cos[x]^(3/2)) + (78*a^2*cos[x]*Sqrt[a*cos[x]^3]*Sin[x])/385 + (26*a^2*cos[x]^3*Sqrt[a*cos[x]^3]*Sin[x])/165 + (2*a^2*cos[x]^5*Sqrt[a*cos[x]^3]*Sin[x])/15 + (26*a^2*Sqrt[a*cos[x]^3]*Tan[x])/77

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*cos[c + d*x])*(b*sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*sin[e + f*x])^(n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
\int (a \cos^3(x))^{5/2} dx &= \frac{(a^2 \sqrt{a \cos^3(x)}) \int \cos^{\frac{15}{2}}(x) dx}{\cos^{\frac{3}{2}}(x)} \\
&= \frac{2}{15} a^2 \cos^5(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{(13a^2 \sqrt{a \cos^3(x)}) \int \cos^{\frac{11}{2}}(x) dx}{15 \cos^{\frac{3}{2}}(x)} \\
&= \frac{26}{165} a^2 \cos^3(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{2}{15} a^2 \cos^5(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{(39a^2 \sqrt{a \cos^3(x)})}{55 \cos^{\frac{3}{2}}(x)} \\
&= \frac{78}{385} a^2 \cos(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{26}{165} a^2 \cos^3(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{2}{15} a^2 \cos^5(x) \sqrt{a \cos^3(x)} \\
&= \frac{78}{385} a^2 \cos(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{26}{165} a^2 \cos^3(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{2}{15} a^2 \cos^5(x) \sqrt{a \cos^3(x)} \\
&= \frac{26a^2 \sqrt{a \cos^3(x)} F\left(\frac{x}{2} \middle| 2\right)}{77 \cos^{\frac{3}{2}}(x)} + \frac{78}{385} a^2 \cos(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{26}{165} a^2 \cos^3(x) \sqrt{a \cos^3(x)}
\end{aligned}$$

Mathematica [A] time = 0.11, size = 61, normalized size = 0.52

$$\frac{a (a \cos^3(x))^{3/2} \left(12480 F\left(\frac{x}{2} \middle| 2\right) + (15465 \sin(x) + 3657 \sin(3x) + 749 \sin(5x) + 77 \sin(7x)) \sqrt{\cos(x)}\right)}{36960 \cos^{\frac{9}{2}}(x)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^3)^(5/2), x]

[Out] (a*(a*cos[x]^3)^(3/2)*(12480*EllipticF[x/2, 2] + Sqrt[Cos[x]]*(15465*Sin[x] + 3657*Sin[3*x] + 749*Sin[5*x] + 77*Sin[7*x])))/(36960*Cos[x]^(9/2))

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{a \cos(x)^3} a^2 \cos(x)^6, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^3)^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(a*cos(x)^3)*a^2*cos(x)^6, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(x)^3)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^3)^(5/2), x, algorithm="giac")

[Out] integrate((a*cos(x)^3)^(5/2), x)

maple [C] time = 0.39, size = 114, normalized size = 0.97

$$\frac{2(-1 + \cos(x)) \left(-77 (\cos^8(x)) + 77 (\cos^7(x)) - 91 (\cos^6(x)) + 195i \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} \text{EllipticF}\left(\frac{i(-1+\cos(x))}{\sin(x)}\right)\right)}{1155}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*cos(x)^3)^(5/2),x)`

[Out] $-2/1155*(-1+\cos(x))*(-77*\cos(x)^8+77*\cos(x)^7-91*\cos(x)^6+195*I*(1/(\cos(x)+1))^{1/2}*(\cos(x)/(\cos(x)+1))^{1/2}*EllipticF(I*(-1+\cos(x))/\sin(x),I)*\sin(x)+91*\cos(x)^5-117*\cos(x)^4+117*\cos(x)^3-195*\cos(x)^2+195*\cos(x))*(\cos(x)+1)^2*(a*\cos(x)^3)^{5/2}/\sin(x)^3/\cos(x)^8$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(x)^3)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(x)^3)^(5/2),x, algorithm="maxima")`

[Out] `integrate((a*cos(x)^3)^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(x)^3)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*cos(x)^3)^(5/2),x)`

[Out] `int((a*cos(x)^3)^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(x)**3)**(5/2),x)`

[Out] Timed out

3.46 $\int (a \cos^3(x))^{3/2} dx$

Optimal. Leaf size=67

$$\frac{14}{45} a \sin(x) \sqrt{a \cos^3(x)} + \frac{14aE\left(\frac{x}{2} \middle| 2\right) \sqrt{a \cos^3(x)}}{15 \cos^2(x)} + \frac{2}{9} a \sin(x) \cos^2(x) \sqrt{a \cos^3(x)}$$

[Out] 14/15*a*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticE(sin(1/2*x),2^(1/2))*(a*cos(x)^3)^(1/2)/cos(x)^(3/2)+14/45*a*sin(x)*(a*cos(x)^3)^(1/2)+2/9*a*cos(x)^2*sin(x)*(a*cos(x)^3)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2635, 2639}

$$\frac{2}{9} a \sin(x) \cos^2(x) \sqrt{a \cos^3(x)} + \frac{14}{45} a \sin(x) \sqrt{a \cos^3(x)} + \frac{14aE\left(\frac{x}{2} \middle| 2\right) \sqrt{a \cos^3(x)}}{15 \cos^2(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^3)^(3/2),x]

[Out] (14*a*Sqrt[a*cos[x]^3]*EllipticE[x/2, 2])/(15*cos[x]^(3/2)) + (14*a*Sqrt[a*cos[x]^3]*Sin[x])/45 + (2*a*cos[x]^2*Sqrt[a*cos[x]^3]*Sin[x])/9

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := -Simp[(b*cos[c + d*x])*(b*sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
\int (a \cos^3(x))^{3/2} dx &= \frac{(a\sqrt{a \cos^3(x)}) \int \cos^{\frac{9}{2}}(x) dx}{\cos^{\frac{3}{2}}(x)} \\
&= \frac{2}{9} a \cos^2(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{(7a\sqrt{a \cos^3(x)}) \int \cos^{\frac{5}{2}}(x) dx}{9 \cos^{\frac{3}{2}}(x)} \\
&= \frac{14}{45} a \sqrt{a \cos^3(x)} \sin(x) + \frac{2}{9} a \cos^2(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{(7a\sqrt{a \cos^3(x)}) \int \sqrt{\cos(x)} dx}{15 \cos^{\frac{3}{2}}(x)} \\
&= \frac{14a\sqrt{a \cos^3(x)} E\left(\frac{x}{2} \middle| 2\right)}{15 \cos^{\frac{3}{2}}(x)} + \frac{14}{45} a \sqrt{a \cos^3(x)} \sin(x) + \frac{2}{9} a \cos^2(x) \sqrt{a \cos^3(x)} \sin(x)
\end{aligned}$$

Mathematica [A] time = 0.07, size = 50, normalized size = 0.75

$$\frac{(a \cos^3(x))^{3/2} \left(168 E\left(\frac{x}{2} \middle| 2\right) + (38 \sin(2x) + 5 \sin(4x)) \sqrt{\cos(x)}\right)}{180 \cos^{\frac{9}{2}}(x)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^3)^(3/2), x]

[Out] ((a*cos[x]^3)^(3/2)*(168*EllipticE[x/2, 2] + Sqrt[Cos[x]]*(38*Sin[2*x] + 5*Sin[4*x]))) / (180*cos[x]^(9/2))

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{a \cos(x)^3} a \cos(x)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^3)^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(a*cos(x)^3)*a*cos(x)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(x)^3)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^3)^(3/2), x, algorithm="giac")

[Out] integrate((a*cos(x)^3)^(3/2), x)

maple [C] time = 0.16, size = 198, normalized size = 2.96

$$\frac{2 \left(5 (\cos^6(x)) - 21i \cos(x) \sin(x) \text{EllipticF}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} + 21i \cos(x) \sin(x) \text{EllipticE}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right) \right)}{15 \cos^{\frac{3}{2}}(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^3)^(3/2), x)

```
[Out] -2/45*(5*cos(x)^6-21*I*cos(x)*sin(x)*EllipticF(I*(-1+cos(x))/sin(x),I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)+21*I*cos(x)*sin(x)*EllipticE(I*(-1+cos(x))/sin(x),I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)-21*I*sin(x)*EllipticF(I*(-1+cos(x))/sin(x),I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)+21*I*sin(x)*EllipticE(I*(-1+cos(x))/sin(x),I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)+2*cos(x)^4+14*cos(x)^2-21*cos(x))*(a*cos(x)^3)^(3/2)/cos(x)^5/sin(x)
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(x)^3)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*cos(x)^3)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((a*cos(x)^3)^(3/2), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(x)^3)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*cos(x)^3)^(3/2),x)
```

```
[Out] int((a*cos(x)^3)^(3/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*cos(x)**3)**(3/2),x)
```

```
[Out] Timed out
```

3.47 $\int \sqrt{a \cos^3(x)} dx$

Optimal. Leaf size=44

$$\frac{2}{3} \tan(x) \sqrt{a \cos^3(x)} + \frac{2F\left(\frac{x}{2} \middle| 2\right) \sqrt{a \cos^3(x)}}{3 \cos^{\frac{3}{2}}(x)}$$

[Out] $\frac{2}{3} (\cos(1/2*x)^2)^{(1/2)} / \cos(1/2*x) * \text{EllipticF}(\sin(1/2*x), 2^{(1/2)}) * (a * \cos(x)^3)^{(1/2)} / \cos(x)^{(3/2)} + 2/3 * (a * \cos(x)^3)^{(1/2)} * \tan(x)$

Rubi [A] time = 0.03, antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2635, 2641}

$$\frac{2}{3} \tan(x) \sqrt{a \cos^3(x)} + \frac{2F\left(\frac{x}{2} \middle| 2\right) \sqrt{a \cos^3(x)}}{3 \cos^{\frac{3}{2}}(x)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a*Cos[x]^3], x]

[Out] $(2 * \text{Sqrt}[a * \text{Cos}[x]^3] * \text{EllipticF}[x/2, 2]) / (3 * \text{Cos}[x]^{(3/2)}) + (2 * \text{Sqrt}[a * \text{Cos}[x]^3] * \text{Tan}[x]) / 3$

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x] * (b*SIN[c + d*x])^(n - 1)) / (d*n), x] + Dist[(b^2*(n - 1)) / n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2]) / d, x] /; FreeQ[{c, d}, x]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[SIN[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p] * (b*SIN[e + f*x])^(n - IntPart[p]) / (SIN[e + f*x] / ff)^(n*FracPart[p])), Int[ActivateTrig[u] * (SIN[e + f*x] / ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int \sqrt{a \cos^3(x)} dx &= \frac{\sqrt{a \cos^3(x)} \int \cos^{\frac{3}{2}}(x) dx}{\cos^{\frac{3}{2}}(x)} \\ &= \frac{2}{3} \sqrt{a \cos^3(x)} \tan(x) + \frac{\sqrt{a \cos^3(x)} \int \frac{1}{\sqrt{\cos(x)}} dx}{3 \cos^{\frac{3}{2}}(x)} \\ &= \frac{2\sqrt{a \cos^3(x)} F\left(\frac{x}{2} \middle| 2\right)}{3 \cos^{\frac{3}{2}}(x)} + \frac{2}{3} \sqrt{a \cos^3(x)} \tan(x) \end{aligned}$$

Mathematica [A] time = 0.02, size = 37, normalized size = 0.84

$$\frac{2\sqrt{a \cos^3(x)} \left(F\left(\frac{x}{2} \middle| 2\right) + \sin(x)\sqrt{\cos(x)} \right)}{3 \cos^{\frac{3}{2}}(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a*Cos[x]^3], x]

[Out] (2*Sqrt[a*Cos[x]^3]*(EllipticF[x/2, 2] + Sqrt[Cos[x]]*Sin[x]))/(3*Cos[x]^(3/2))

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{a \cos(x)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^3)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(a*cos(x)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \cos(x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^3)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(a*cos(x)^3), x)

maple [C] time = 0.18, size = 76, normalized size = 1.73

$$\frac{2(-1 + \cos(x)) \left(i \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} \text{EllipticF}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right) \sin(x) - (\cos^2(x) + \cos(x)) (\cos(x) + 1)^2 \sqrt{a} \right)}{3 \cos(x)^2 \sin(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^3)^(1/2), x)

[Out] -2/3*(-1+cos(x))*(I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x), I)*sin(x)-cos(x)^2+cos(x))*(cos(x)+1)^2*(a*cos(x)^3)^(1/2)/cos(x)^2/sin(x)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \cos(x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^3)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(a*cos(x)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \sqrt{a \cos(x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*cos(x)^3)^(1/2),x)
```

```
[Out] int((a*cos(x)^3)^(1/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*cos(x)**3)**(1/2),x)
```

```
[Out] Timed out
```

$$3.48 \quad \int \frac{1}{\sqrt{a \cos^3(x)}} dx$$

Optimal. Leaf size=42

$$\frac{2 \sin(x) \cos(x)}{\sqrt{a \cos^3(x)}} - \frac{2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right)}{\sqrt{a \cos^3(x)}}$$

[Out] $-2 \cos(x)^{(3/2)} (\cos(1/2*x)^2)^{(1/2)} / \cos(1/2*x) * \text{EllipticE}(\sin(1/2*x), 2^{(1/2)}) / (a * \cos(x)^3)^{(1/2)} + 2 * \cos(x) * \sin(x) / (a * \cos(x)^3)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2636, 2639}

$$\frac{2 \sin(x) \cos(x)}{\sqrt{a \cos^3(x)}} - \frac{2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right)}{\sqrt{a \cos^3(x)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a*Cos[x]^3], x]

[Out] $(-2 * \text{Cos}[x]^{(3/2)} * \text{EllipticE}[x/2, 2]) / \text{Sqrt}[a * \text{Cos}[x]^3] + (2 * \text{Cos}[x] * \text{Sin}[x]) / \text{Sqrt}[a * \text{Cos}[x]^3]$

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a \cos^3(x)}} dx &= \frac{\cos^{\frac{3}{2}}(x) \int \frac{1}{\cos^{\frac{3}{2}}(x)} dx}{\sqrt{a \cos^3(x)}} \\ &= \frac{2 \cos(x) \sin(x)}{\sqrt{a \cos^3(x)}} - \frac{\cos^{\frac{3}{2}}(x) \int \sqrt{\cos(x)} dx}{\sqrt{a \cos^3(x)}} \\ &= -\frac{2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right)}{\sqrt{a \cos^3(x)}} + \frac{2 \cos(x) \sin(x)}{\sqrt{a \cos^3(x)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 31, normalized size = 0.74

$$\frac{\sin(2x) - 2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right)}{\sqrt{a \cos^3(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[a*Cos[x]^3], x]

[Out] (-2*Cos[x]^(3/2)*EllipticE[x/2, 2] + Sin[2*x])/Sqrt[a*Cos[x]^3]

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{a \cos(x)^3}}{a \cos(x)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^3)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(a*cos(x)^3)/(a*cos(x)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \cos(x)^3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^3)^(1/2), x, algorithm="giac")

[Out] integrate(1/sqrt(a*cos(x)^3), x)

maple [C] time = 0.31, size = 191, normalized size = 4.55

$$2(\cos(x)+1)^2(-1+\cos(x))^2 \left(i \cos(x) \sin(x) \text{EllipticE}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} - i \cos(x) \sin(x) \text{Ellip} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^3)^(1/2), x)

[Out] 2*(cos(x)+1)^2*(-1+cos(x))^2*(I*cos(x)*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(-1+cos(x))/sin(x), I)-I*cos(x)*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x), I)+I*sin(x)*EllipticE(I*(-1+cos(x))/sin(x), I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)-I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x), I)*sin(x)-cos(x)+1)*cos(x)/(a*cos(x)^3)^(1/2)/sin(x)^5

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \cos(x)^3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^3)^(1/2), x, algorithm="maxima")

[Out] integrate(1/sqrt(a*cos(x)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{a \cos(x)^3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^3)^(1/2), x)

[Out] int(1/(a*cos(x)^3)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)**3)**(1/2), x)

[Out] Integral(1/sqrt(a*cos(x)**3), x)

$$3.49 \quad \int \frac{1}{(a \cos^3(x))^{3/2}} dx$$

Optimal. Leaf size=71

$$\frac{10 \sin(x)}{21a\sqrt{a \cos^3(x)}} + \frac{10 \cos^2(x) F\left(\frac{x}{2} \middle| 2\right)}{21a\sqrt{a \cos^3(x)}} + \frac{2 \tan(x) \sec(x)}{7a\sqrt{a \cos^3(x)}}$$

[Out] 10/21*cos(x)^(3/2)*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticF(sin(1/2*x),2^(1/2))/a/(a*cos(x)^3)^(1/2)+10/21*sin(x)/a/(a*cos(x)^3)^(1/2)+2/7*sec(x)*tan(x)/a/(a*cos(x)^3)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2636, 2641}

$$\frac{10 \sin(x)}{21a\sqrt{a \cos^3(x)}} + \frac{10 \cos^2(x) F\left(\frac{x}{2} \middle| 2\right)}{21a\sqrt{a \cos^3(x)}} + \frac{2 \tan(x) \sec(x)}{7a\sqrt{a \cos^3(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^3)^(-3/2),x]

[Out] (10*cos[x]^(3/2)*EllipticF[x/2, 2])/(21*a*Sqrt[a*cos[x]^3]) + (10*sin[x])/(21*a*Sqrt[a*cos[x]^3]) + (2*Sec[x]*Tan[x])/(7*a*Sqrt[a*cos[x]^3])

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a \cos^3(x))^{3/2}} dx &= \frac{\cos^{\frac{3}{2}}(x) \int \frac{1}{\cos^{\frac{3}{2}}(x)} dx}{a \sqrt{a \cos^3(x)}} \\
&= \frac{2 \sec(x) \tan(x)}{7a \sqrt{a \cos^3(x)}} + \frac{\left(5 \cos^{\frac{3}{2}}(x)\right) \int \frac{1}{\cos^{\frac{3}{2}}(x)} dx}{7a \sqrt{a \cos^3(x)}} \\
&= \frac{10 \sin(x)}{21a \sqrt{a \cos^3(x)}} + \frac{2 \sec(x) \tan(x)}{7a \sqrt{a \cos^3(x)}} + \frac{\left(5 \cos^{\frac{3}{2}}(x)\right) \int \frac{1}{\sqrt{\cos(x)}} dx}{21a \sqrt{a \cos^3(x)}} \\
&= \frac{10 \cos^{\frac{3}{2}}(x) F\left(\frac{x}{2} \middle| 2\right)}{21a \sqrt{a \cos^3(x)}} + \frac{10 \sin(x)}{21a \sqrt{a \cos^3(x)}} + \frac{2 \sec(x) \tan(x)}{7a \sqrt{a \cos^3(x)}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 44, normalized size = 0.62

$$\frac{2 \cos^2(x) \left(3 \tan(x) + 5 \cos^{\frac{5}{2}}(x) F\left(\frac{x}{2} \middle| 2\right) + 5 \sin(x) \cos(x)\right)}{21 (a \cos^3(x))^{\frac{3}{2}}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Cos[x]^3)^(-3/2), x]

[Out] (2*Cos[x]^2*(5*Cos[x]^(5/2)*EllipticF[x/2, 2] + 5*Cos[x]*Sin[x] + 3*Tan[x]) / (21*(a*Cos[x]^3)^(3/2))

fricas [F] time = 0.47, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{a \cos(x)^3}}{a^2 \cos(x)^6}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^3)^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(a*cos(x)^3)/(a^2*cos(x)^6), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \cos(x)^3)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^3)^(3/2), x, algorithm="giac")

[Out] integrate((a*cos(x)^3)^(-3/2), x)

maple [C] time = 0.31, size = 87, normalized size = 1.23

$$\frac{2 (\cos(x) + 1)^2 (-1 + \cos(x)) \left(5i (\cos^3(x)) \sin(x) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} \text{EllipticF}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right) - 5 (\cos^3(x))\right)}{21 \sin(x)^3 (a (\cos^3(x)))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a*cos(x)^3)^(3/2),x)`

[Out] `-2/21*(cos(x)+1)^2*(-1+cos(x))*(5*I*cos(x)^3*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x),I)-5*cos(x)^3+5*cos(x)^2-3*cos(x)+3)*cos(x)/sin(x)^3/(a*cos(x)^3)^(3/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \cos(x)^3)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*cos(x)^3)^(3/2),x, algorithm="maxima")`

[Out] `integrate((a*cos(x)^3)^(-3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(a \cos(x)^3)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a*cos(x)^3)^(3/2),x)`

[Out] `int(1/(a*cos(x)^3)^(3/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*cos(x)**3)**(3/2),x)`

[Out] Timed out

$$3.50 \quad \int \frac{1}{(a \cos^3(x))^{5/2}} dx$$

Optimal. Leaf size=117

$$\frac{154 \sin(x) \cos(x)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} - \frac{154 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{2 \tan(x) \sec^4(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{22 \tan(x) \sec^2(x)}{117a^2 \sqrt{a \cos^3(x)}}$$

[Out] $-154/195 \cos(x)^{(3/2)} (\cos(1/2*x)^2)^{(1/2)} / \cos(1/2*x) * \text{EllipticE}(\sin(1/2*x), 2^{(1/2)}) / a^2 / (a \cos(x)^3)^{(1/2)} + 154/195 \cos(x) \sin(x) / a^2 / (a \cos(x)^3)^{(1/2)} + 154/585 \tan(x) / a^2 / (a \cos(x)^3)^{(1/2)} + 22/117 \sec(x)^2 \tan(x) / a^2 / (a \cos(x)^3)^{(1/2)} + 2/13 \sec(x)^4 \tan(x) / a^2 / (a \cos(x)^3)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2636, 2639}

$$\frac{154 \sin(x) \cos(x)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} - \frac{154 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{2 \tan(x) \sec^4(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{22 \tan(x) \sec^2(x)}{117a^2 \sqrt{a \cos^3(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^3)^(-5/2), x]

[Out] $(-154 \cos[x]^{(3/2)} \text{EllipticE}[x/2, 2]) / (195 a^2 \sqrt{a \cos[x]^3}) + (154 \cos[x] \sin[x]) / (195 a^2 \sqrt{a \cos[x]^3}) + (154 \tan[x]) / (585 a^2 \sqrt{a \cos[x]^3}) + (22 \sec[x]^2 \tan[x]) / (117 a^2 \sqrt{a \cos[x]^3}) + (2 \sec[x]^4 \tan[x]) / (13 a^2 \sqrt{a \cos[x]^3})$

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*SIN[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[SIN[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*SIN[e + f*x]^n)^FracPart[p])/(SIN[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(SIN[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a \cos^3(x))^{5/2}} dx &= \frac{\cos^{3/2}(x) \int \frac{1}{\cos^2(x)} dx}{a^2 \sqrt{a \cos^3(x)}} \\
&= \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{\left(11 \cos^{3/2}(x)\right) \int \frac{1}{\cos^2(x)} dx}{13a^2 \sqrt{a \cos^3(x)}} \\
&= \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{\left(77 \cos^{3/2}(x)\right) \int \frac{1}{\cos^2(x)} dx}{117a^2 \sqrt{a \cos^3(x)}} \\
&= \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} + \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{\left(77 \cos^{3/2}(x)\right) \int \frac{1}{\cos^2(x)} dx}{195a^2 \sqrt{a \cos^3(x)}} \\
&= \frac{154 \cos(x) \sin(x)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} + \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}} - \frac{\left(77 \cos^{3/2}(x)\right) \int \frac{1}{\cos^2(x)} dx}{195a^2 \sqrt{a \cos^3(x)}} \\
&= -\frac{154 \cos^{3/2}(x) E\left(\frac{x}{2} \middle| 2\right)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \cos(x) \sin(x)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} + \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 57, normalized size = 0.49

$$\frac{-462 \cos^{3/2}(x) E\left(\frac{x}{2} \middle| 2\right) + 462 \sin(x) \cos(x) + 2 \tan(x) (45 \sec^4(x) + 55 \sec^2(x) + 77)}{585a^2 \sqrt{a \cos^3(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^3)^(-5/2), x]

[Out] (-462*cos[x]^(3/2)*EllipticE[x/2, 2] + 462*cos[x]*Sin[x] + 2*(77 + 55*Sec[x]^2 + 45*Sec[x]^4)*Tan[x])/(585*a^2*Sqrt[a*cos[x]^3])

fricas [F] time = 0.92, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{a \cos(x)^3}}{a^3 \cos(x)^9}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^3)^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(a*cos(x)^3)/(a^3*cos(x)^9), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \cos(x)^3)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^3)^(5/2), x, algorithm="giac")

[Out] integrate((a*cos(x)^3)^(-5/2), x)

maple [C] time = 0.38, size = 223, normalized size = 1.91

$$2(\cos(x)+1)^2(-1+\cos(x))^2\left(231i(\cos^7(x))\sin(x)\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticF}\left(\frac{i(-1+\cos(x))}{\sin(x)},i\right)-231i(\cos(x)+1)^2(-1+\cos(x))^2\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticE}\left(\frac{i(-1+\cos(x))}{\sin(x)},i\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^3)^(5/2),x)

[Out] -2/585*(cos(x)+1)^2*(-1+cos(x))^2*(231*I*cos(x)^7*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x),I)-231*I*cos(x)^7*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(-1+cos(x))/sin(x),I)+231*I*cos(x)^6*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x),I)-231*I*cos(x)^6*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(-1+cos(x))/sin(x),I)+231*cos(x)^7-154*cos(x)^6-22*cos(x)^4-10*cos(x)^2-45)*cos(x)/sin(x)^5/(a*cos(x)^3)^(5/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \cos(x)^3)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^3)^(5/2),x, algorithm="maxima")

[Out] integrate((a*cos(x)^3)^(-5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(a \cos(x)^3)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^3)^(5/2),x)

[Out] int(1/(a*cos(x)^3)^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)**3)**(5/2),x)

[Out] Timed out

3.51 $\int (a \cos^4(x))^{5/2} dx$

Optimal. Leaf size=132

$$\frac{21}{128}a^2 \sin(x) \cos(x) \sqrt{a \cos^4(x)} + \frac{63}{256}a^2 \tan(x) \sqrt{a \cos^4(x)} + \frac{63}{256}a^2 x \sec^2(x) \sqrt{a \cos^4(x)} + \frac{1}{10}a^2 \sin(x) \cos^7(x) \sqrt{a \cos^4(x)}$$

[Out] 63/256*a^2*x*sec(x)^2*(a*cos(x)^4)^(1/2)+21/128*a^2*cos(x)*sin(x)*(a*cos(x)^4)^(1/2)+21/160*a^2*cos(x)^3*sin(x)*(a*cos(x)^4)^(1/2)+9/80*a^2*cos(x)^5*sin(x)*(a*cos(x)^4)^(1/2)+1/10*a^2*cos(x)^7*sin(x)*(a*cos(x)^4)^(1/2)+63/256*a^2*(a*cos(x)^4)^(1/2)*tan(x)

Rubi [A] time = 0.05, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2635, 8}

$$\frac{1}{10}a^2 \sin(x) \cos^7(x) \sqrt{a \cos^4(x)} + \frac{9}{80}a^2 \sin(x) \cos^5(x) \sqrt{a \cos^4(x)} + \frac{21}{160}a^2 \sin(x) \cos^3(x) \sqrt{a \cos^4(x)} + \frac{21}{128}a^2 \sin(x) \cos(x) \sqrt{a \cos^4(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^4)^(5/2), x]

[Out] (63*a^2*x*sqrt[a*cos[x]^4]*Sec[x]^2)/256 + (21*a^2*cos[x]*sqrt[a*cos[x]^4]*Sin[x])/128 + (21*a^2*cos[x]^3*sqrt[a*cos[x]^4]*Sin[x])/160 + (9*a^2*cos[x]^5*sqrt[a*cos[x]^4]*Sin[x])/80 + (a^2*cos[x]^7*sqrt[a*cos[x]^4]*Sin[x])/10 + (63*a^2*sqrt[a*cos[x]^4]*Tan[x])/256

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*cos[c+d*x])*(b*sin[c+d*x])^(n-1)/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*sin[c+d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3207

Int[(u_)*((b_)*sin[(e_)+(f_)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e+f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*sin[e+f*x])^FracPart[p])/(Sin[e+f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e+f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_)*(trig_)[e+f*x])^(m_) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
\int (a \cos^4(x))^{5/2} dx &= \left(a^2 \sqrt{a \cos^4(x)} \sec^2(x) \right) \int \cos^{10}(x) dx \\
&= \frac{1}{10} a^2 \cos^7(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{10} \left(9a^2 \sqrt{a \cos^4(x)} \sec^2(x) \right) \int \cos^8(x) dx \\
&= \frac{9}{80} a^2 \cos^5(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{10} a^2 \cos^7(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{80} \left(63a^2 \sqrt{a \cos^4(x)} \sec^2(x) \right) \int \cos^6(x) dx \\
&= \frac{21}{160} a^2 \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{9}{80} a^2 \cos^5(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{10} a^2 \cos^7(x) \sqrt{a \cos^4(x)} \sin(x) \\
&= \frac{21}{128} a^2 \cos(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{21}{160} a^2 \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{9}{80} a^2 \cos^5(x) \sqrt{a \cos^4(x)} \sin(x) \\
&= \frac{21}{128} a^2 \cos(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{21}{160} a^2 \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{9}{80} a^2 \cos^5(x) \sqrt{a \cos^4(x)} \sin(x) \\
&= \frac{63}{256} a^2 x \sqrt{a \cos^4(x)} \sec^2(x) + \frac{21}{128} a^2 \cos(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{21}{160} a^2 \cos^3(x) \sqrt{a \cos^4(x)} \sin(x)
\end{aligned}$$

Mathematica [A] time = 0.12, size = 53, normalized size = 0.40

$$\frac{a(2520x + 2100 \sin(2x) + 600 \sin(4x) + 150 \sin(6x) + 25 \sin(8x) + 2 \sin(10x)) \sec^6(x) (a \cos^4(x))^{3/2}}{10240}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^4)^(5/2),x]

[Out] (a*(a*cos[x]^4)^(3/2)*Sec[x]^6*(2520*x + 2100*Sin[2*x] + 600*Sin[4*x] + 150*Sin[6*x] + 25*Sin[8*x] + 2*Sin[10*x]))/10240

fricas [A] time = 0.53, size = 68, normalized size = 0.52

$$\frac{\sqrt{a \cos(x)^4} \left(315 a^2 x + (128 a^2 \cos(x)^9 + 144 a^2 \cos(x)^7 + 168 a^2 \cos(x)^5 + 210 a^2 \cos(x)^3 + 315 a^2 \cos(x)) \sin(x) \right)}{1280 \cos(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(5/2),x, algorithm="fricas")

[Out] 1/1280*sqrt(a*cos(x)^4)*(315*a^2*x + (128*a^2*cos(x)^9 + 144*a^2*cos(x)^7 + 168*a^2*cos(x)^5 + 210*a^2*cos(x)^3 + 315*a^2*cos(x))*sin(x))/cos(x)^2

giac [A] time = 0.60, size = 57, normalized size = 0.43

$$\frac{1}{10240} \left(2520 a^2 x + 2 a^2 \sin(10x) + 25 a^2 \sin(8x) + 150 a^2 \sin(6x) + 600 a^2 \sin(4x) + 2100 a^2 \sin(2x) \right) \sqrt{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(5/2),x, algorithm="giac")

[Out] 1/10240*(2520*a^2*x + 2*a^2*sin(10*x) + 25*a^2*sin(8*x) + 150*a^2*sin(6*x) + 600*a^2*sin(4*x) + 2100*a^2*sin(2*x))*sqrt(a)

maple [A] time = 0.41, size = 57, normalized size = 0.43

$$\frac{\left(a \left(\cos^4(x) \right) \right)^{5/2} \left(128 \sin(x) \left(\cos^9(x) \right) + 144 \sin(x) \left(\cos^7(x) \right) + 168 \sin(x) \left(\cos^5(x) \right) + 210 \left(\cos^3(x) \right) \sin(x) + 315 a \right)}{1280 \cos(x)^{10}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^4)^(5/2),x)

[Out] 1/1280*(a*cos(x)^4)^(5/2)*(128*sin(x)*cos(x)^9+144*sin(x)*cos(x)^7+168*sin(x)*cos(x)^5+210*cos(x)^3*sin(x)+315*cos(x)*sin(x)+315*x)/cos(x)^10

maxima [A] time = 0.90, size = 85, normalized size = 0.64

$$\frac{63}{256} a^{\frac{5}{2}} x + \frac{315 a^{\frac{5}{2}} \tan(x)^9 + 1470 a^{\frac{5}{2}} \tan(x)^7 + 2688 a^{\frac{5}{2}} \tan(x)^5 + 2370 a^{\frac{5}{2}} \tan(x)^3 + 965 a^{\frac{5}{2}} \tan(x)}{1280 (\tan(x)^{10} + 5 \tan(x)^8 + 10 \tan(x)^6 + 10 \tan(x)^4 + 5 \tan(x)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(5/2),x, algorithm="maxima")

[Out] 63/256*a^(5/2)*x + 1/1280*(315*a^(5/2)*tan(x)^9 + 1470*a^(5/2)*tan(x)^7 + 2688*a^(5/2)*tan(x)^5 + 2370*a^(5/2)*tan(x)^3 + 965*a^(5/2)*tan(x))/(tan(x)^10 + 5*tan(x)^8 + 10*tan(x)^6 + 10*tan(x)^4 + 5*tan(x)^2 + 1)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(x)^4)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^4)^(5/2),x)

[Out] int((a*cos(x)^4)^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)**4)**(5/2),x)

[Out] Timed out

3.52 $\int (a \cos^4(x))^{3/2} dx$

Optimal. Leaf size=78

$$\frac{5}{24}a \sin(x) \cos(x) \sqrt{a \cos^4(x)} + \frac{5}{16}a \tan(x) \sqrt{a \cos^4(x)} + \frac{5}{16}ax \sec^2(x) \sqrt{a \cos^4(x)} + \frac{1}{6}a \sin(x) \cos^3(x) \sqrt{a \cos^4(x)}$$

[Out] 5/16*a*x*sec(x)^2*(a*cos(x)^4)^(1/2)+5/24*a*cos(x)*sin(x)*(a*cos(x)^4)^(1/2)+1/6*a*cos(x)^3*sin(x)*(a*cos(x)^4)^(1/2)+5/16*a*(a*cos(x)^4)^(1/2)*tan(x)

Rubi [A] time = 0.03, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2635, 8}

$$\frac{1}{6}a \sin(x) \cos^3(x) \sqrt{a \cos^4(x)} + \frac{5}{24}a \sin(x) \cos(x) \sqrt{a \cos^4(x)} + \frac{5}{16}a \tan(x) \sqrt{a \cos^4(x)} + \frac{5}{16}ax \sec^2(x) \sqrt{a \cos^4(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^4)^(3/2), x]

[Out] (5*a*x*Sqrt[a*cos[x]^4]*Sec[x]^2)/16 + (5*a*cos[x]*Sqrt[a*cos[x]^4]*Sin[x])/24 + (a*cos[x]^3*Sqrt[a*cos[x]^4]*Sin[x])/6 + (5*a*Sqrt[a*cos[x]^4]*Tan[x])/16

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*cos[c + d*x])*(b*sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*sin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int (a \cos^4(x))^{3/2} dx &= \left(a \sqrt{a \cos^4(x)} \sec^2(x) \right) \int \cos^6(x) dx \\ &= \frac{1}{6}a \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{6} \left(5a \sqrt{a \cos^4(x)} \sec^2(x) \right) \int \cos^4(x) dx \\ &= \frac{5}{24}a \cos(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{6}a \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{8} \left(5a \sqrt{a \cos^4(x)} \sec^2(x) \right) \int \cos^2(x) dx \\ &= \frac{5}{24}a \cos(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{6}a \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{5}{16}a \sqrt{a \cos^4(x)} \tan(x) + \frac{5}{16}ax \sec^2(x) \sqrt{a \cos^4(x)} \\ &= \frac{5}{16}ax \sqrt{a \cos^4(x)} \sec^2(x) + \frac{5}{24}a \cos(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{1}{6}a \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) \end{aligned}$$

Mathematica [A] time = 0.07, size = 38, normalized size = 0.49

$$\frac{1}{192}(60x + 45 \sin(2x) + 9 \sin(4x) + \sin(6x)) \sec^6(x) (a \cos^4(x))^{3/2}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^4)^(3/2),x]

[Out] ((a*cos[x]^4)^(3/2)*Sec[x]^6*(60*x + 45*Sin[2*x] + 9*Sin[4*x] + Sin[6*x]))/192

fricas [A] time = 0.52, size = 42, normalized size = 0.54

$$\frac{\sqrt{a \cos(x)^4} (15 a x + (8 a \cos(x)^5 + 10 a \cos(x)^3 + 15 a \cos(x)) \sin(x))}{48 \cos(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(3/2),x, algorithm="fricas")

[Out] 1/48*sqrt(a*cos(x)^4)*(15*a*x + (8*a*cos(x)^5 + 10*a*cos(x)^3 + 15*a*cos(x))*sin(x))/cos(x)^2

giac [A] time = 0.55, size = 25, normalized size = 0.32

$$\frac{1}{192} a^{\frac{3}{2}} (60x + \sin(6x) + 9 \sin(4x) + 45 \sin(2x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(3/2),x, algorithm="giac")

[Out] 1/192*a^(3/2)*(60*x + sin(6*x) + 9*sin(4*x) + 45*sin(2*x))

maple [A] time = 0.17, size = 41, normalized size = 0.53

$$\frac{(a (\cos^4(x)))^{\frac{3}{2}} (8 \sin(x) (\cos^5(x)) + 10 (\cos^3(x)) \sin(x) + 15 \cos(x) \sin(x) + 15x)}{48 \cos(x)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^4)^(3/2),x)

[Out] 1/48*(a*cos(x)^4)^(3/2)*(8*sin(x)*cos(x)^5+10*cos(x)^3*sin(x)+15*cos(x)*sin(x)+15*x)/cos(x)^6

maxima [A] time = 0.67, size = 55, normalized size = 0.71

$$\frac{5}{16} a^{\frac{3}{2}} x + \frac{15 a^{\frac{3}{2}} \tan(x)^5 + 40 a^{\frac{3}{2}} \tan(x)^3 + 33 a^{\frac{3}{2}} \tan(x)}{48 (\tan(x)^6 + 3 \tan(x)^4 + 3 \tan(x)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(3/2),x, algorithm="maxima")

[Out] 5/16*a^(3/2)*x + 1/48*(15*a^(3/2)*tan(x)^5 + 40*a^(3/2)*tan(x)^3 + 33*a^(3/2)*tan(x))/(tan(x)^6 + 3*tan(x)^4 + 3*tan(x)^2 + 1)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(x)^4)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*cos(x)^4)^(3/2),x)
```

```
[Out] int((a*cos(x)^4)^(3/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*cos(x)**4)**(3/2),x)
```

```
[Out] Timed out
```

3.53 $\int \sqrt{a \cos^4(x)} dx$

Optimal. Leaf size=36

$$\frac{1}{2} \tan(x) \sqrt{a \cos^4(x)} + \frac{1}{2} x \sec^2(x) \sqrt{a \cos^4(x)}$$

[Out] $1/2*x*\sec(x)^2*(a*\cos(x)^4)^{(1/2)}+1/2*(a*\cos(x)^4)^{(1/2)}*\tan(x)$

Rubi [A] time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 2635, 8}

$$\frac{1}{2} \tan(x) \sqrt{a \cos^4(x)} + \frac{1}{2} x \sec^2(x) \sqrt{a \cos^4(x)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a*Cos[x]^4], x]

[Out] (x*Sqrt[a*Cos[x]^4]*Sec[x]^2)/2 + (Sqrt[a*Cos[x]^4]*Tan[x])/2

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x] * (b*SIN[c + d*x])^(n - 1)) / (d*n), x] + Dist[(b^2*(n - 1)) / n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] := With[{ff = FreeFactors[SIN[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p] * (b*SIN[e + f*x]^n)^FracPart[p]) / (SIN[e + f*x] / ff)^(n*FracPart[p]), Int[ActivateTrig[u] * (SIN[e + f*x] / ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int \sqrt{a \cos^4(x)} dx &= \left(\sqrt{a \cos^4(x)} \sec^2(x) \right) \int \cos^2(x) dx \\ &= \frac{1}{2} \sqrt{a \cos^4(x)} \tan(x) + \frac{1}{2} \left(\sqrt{a \cos^4(x)} \sec^2(x) \right) \int 1 dx \\ &= \frac{1}{2} x \sqrt{a \cos^4(x)} \sec^2(x) + \frac{1}{2} \sqrt{a \cos^4(x)} \tan(x) \end{aligned}$$

Mathematica [A] time = 0.01, size = 25, normalized size = 0.69

$$\frac{1}{2} \sec^2(x) \sqrt{a \cos^4(x)} (x + \sin(x) \cos(x))$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a*Cos[x]^4], x]

[Out] (Sqrt[a*Cos[x]^4]*Sec[x]^2*(x + Cos[x]*Sin[x]))/2

fricas [A] time = 0.51, size = 21, normalized size = 0.58

$$\frac{\sqrt{a \cos(x)^4} (\cos(x) \sin(x) + x)}{2 \cos(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(1/2),x, algorithm="fricas")

[Out] 1/2*sqrt(a*cos(x)^4)*(cos(x)*sin(x) + x)/cos(x)^2

giac [A] time = 0.27, size = 13, normalized size = 0.36

$$\frac{1}{4} \sqrt{a} (2x + \sin(2x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(a)*(2*x + sin(2*x))

maple [A] time = 0.09, size = 22, normalized size = 0.61

$$\frac{\sqrt{a (\cos^4(x))} (\cos(x) \sin(x) + x)}{2 \cos(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^4)^(1/2),x)

[Out] 1/2*(a*cos(x)^4)^(1/2)*(cos(x)*sin(x)+x)/cos(x)^2

maxima [A] time = 1.07, size = 22, normalized size = 0.61

$$\frac{1}{2} \sqrt{a} x + \frac{\sqrt{a} \tan(x)}{2 (\tan(x)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)^4)^(1/2),x, algorithm="maxima")

[Out] 1/2*sqrt(a)*x + 1/2*sqrt(a)*tan(x)/(tan(x)^2 + 1)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \sqrt{a \cos(x)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(x)^4)^(1/2),x)

[Out] int((a*cos(x)^4)^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(x)**4)**(1/2),x)

[Out] Timed out

$$3.54 \quad \int \frac{1}{\sqrt{a \cos^4(x)}} dx$$

Optimal. Leaf size=15

$$\frac{\sin(x) \cos(x)}{\sqrt{a \cos^4(x)}}$$

[Out] cos(x)*sin(x)/(a*cos(x)^4)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3207, 3767, 8}

$$\frac{\sin(x) \cos(x)}{\sqrt{a \cos^4(x)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a*Cos[x]^4],x]

[Out] (Cos[x]*Sin[x])/Sqrt[a*Cos[x]^4]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Ssin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a \cos^4(x)}} dx &= \frac{\cos^2(x) \int \sec^2(x) dx}{\sqrt{a \cos^4(x)}} \\ &= -\frac{\cos^2(x) \text{Subst}(\int 1 dx, x, -\tan(x))}{\sqrt{a \cos^4(x)}} \\ &= \frac{\cos(x) \sin(x)}{\sqrt{a \cos^4(x)}} \end{aligned}$$

Mathematica [A] time = 0.00, size = 15, normalized size = 1.00

$$\frac{\sin(x) \cos(x)}{\sqrt{a \cos^4(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[a*Cos[x]^4],x]

[Out] (Cos[x]*Sin[x])/Sqrt[a*Cos[x]^4]

fricas [A] time = 0.49, size = 18, normalized size = 1.20

$$\frac{\sqrt{a \cos(x)^4} \sin(x)}{a \cos(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(1/2),x, algorithm="fricas")

[Out] sqrt(a*cos(x)^4)*sin(x)/(a*cos(x)^3)

giac [A] time = 0.28, size = 6, normalized size = 0.40

$$\frac{\tan(x)}{\sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(1/2),x, algorithm="giac")

[Out] tan(x)/sqrt(a)

maple [A] time = 0.08, size = 14, normalized size = 0.93

$$\frac{\cos(x) \sin(x)}{\sqrt{a (\cos^4(x))}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^4)^(1/2),x)

[Out] cos(x)*sin(x)/(a*cos(x)^4)^(1/2)

maxima [A] time = 1.00, size = 6, normalized size = 0.40

$$\frac{\tan(x)}{\sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(1/2),x, algorithm="maxima")

[Out] tan(x)/sqrt(a)

mupad [B] time = 0.23, size = 6, normalized size = 0.40

$$\frac{\tan(x)}{\sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^4)^(1/2),x)

[Out] tan(x)/a^(1/2)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)**4)**(1/2),x)

[Out] Timed out

$$3.55 \quad \int \frac{1}{(a \cos^4(x))^{3/2}} dx$$

Optimal. Leaf size=67

$$\frac{\sin(x) \cos(x)}{a\sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^3(x)}{5a\sqrt{a \cos^4(x)}} + \frac{2 \sin^2(x) \tan(x)}{3a\sqrt{a \cos^4(x)}}$$

[Out] $\cos(x) \sin(x) / a / (a \cos(x)^4)^{(1/2)} + 2/3 \sin(x)^2 \tan(x) / a / (a \cos(x)^4)^{(1/2)} + 1/5 \sin(x)^2 \tan(x)^3 / a / (a \cos(x)^4)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3207, 3767}

$$\frac{\sin(x) \cos(x)}{a\sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^3(x)}{5a\sqrt{a \cos^4(x)}} + \frac{2 \sin^2(x) \tan(x)}{3a\sqrt{a \cos^4(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*cos[x]^4)^(-3/2), x]

[Out] (Cos[x]*Sin[x])/(a*Sqrt[a*Cos[x]^4]) + (2*Sin[x]^2*Tan[x])/(3*a*Sqrt[a*Cos[x]^4]) + (Sin[x]^2*Tan[x]^3)/(5*a*Sqrt[a*Cos[x]^4])

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*ff^n)^IntPart[p]*(b*Ssin[e + f*x]^n)^FracPart[p])/(Sin[e + f*x]/ff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a \cos^4(x))^{3/2}} dx &= \frac{\cos^2(x) \int \sec^6(x) dx}{a\sqrt{a \cos^4(x)}} \\ &= -\frac{\cos^2(x) \text{Subst}\left(\int (1 + 2x^2 + x^4) dx, x, -\tan(x)\right)}{a\sqrt{a \cos^4(x)}} \\ &= \frac{\cos(x) \sin(x)}{a\sqrt{a \cos^4(x)}} + \frac{2 \sin^2(x) \tan(x)}{3a\sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^3(x)}{5a\sqrt{a \cos^4(x)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 30, normalized size = 0.45

$$\frac{\sin(x) \cos(x) (6 \cos(2x) + \cos(4x) + 8)}{15 (a \cos^4(x))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^4)^(-3/2), x]

[Out] (Cos[x]*(8 + 6*cos[2*x] + Cos[4*x])*Sin[x])/(15*(a*cos[x]^4)^(3/2))

fricas [A] time = 0.73, size = 33, normalized size = 0.49

$$\frac{\sqrt{a} \cos(x)^4 (8 \cos(x)^4 + 4 \cos(x)^2 + 3) \sin(x)}{15 a^2 \cos(x)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(3/2), x, algorithm="fricas")

[Out] 1/15*sqrt(a*cos(x)^4)*(8*cos(x)^4 + 4*cos(x)^2 + 3)*sin(x)/(a^2*cos(x)^7)

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(3/2), x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,x):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value

maple [A] time = 0.10, size = 29, normalized size = 0.43

$$\frac{\sin(x) (8 (\cos^4(x)) + 4 (\cos^2(x)) + 3) \cos(x)}{15 (a (\cos^4(x)))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^4)^(3/2), x)

[Out] 1/15*sin(x)*(8*cos(x)^4+4*cos(x)^2+3)*cos(x)/(a*cos(x)^4)^(3/2)

maxima [A] time = 0.87, size = 22, normalized size = 0.33

$$\frac{3 \tan(x)^5 + 10 \tan(x)^3 + 15 \tan(x)}{15 a^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(3/2), x, algorithm="maxima")

[Out] 1/15*(3*tan(x)^5 + 10*tan(x)^3 + 15*tan(x))/a^(3/2)

mupad [B] time = 0.54, size = 36, normalized size = 0.54

$$\frac{4 \sin(x)}{5 a^{3/2} \cos(x)^3} + \frac{\sin(x)}{5 a^{3/2} \cos(x)^5} - \frac{8 \sin(x)^3}{15 a^{3/2} \cos(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^4)^(3/2), x)

[Out] (4*sin(x))/(5*a^(3/2)*cos(x)^3) + sin(x)/(5*a^(3/2)*cos(x)^5) - (8*sin(x)^3)/(15*a^(3/2)*cos(x)^3)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)**4)**(3/2),x)

[Out] Timed out

$$3.56 \quad \int \frac{1}{(a \cos^4(x))^{5/2}} dx$$

Optimal. Leaf size=117

$$\frac{\sin(x) \cos(x)}{a^2 \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^7(x)}{9a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan^5(x)}{7a^2 \sqrt{a \cos^4(x)}} + \frac{6 \sin^2(x) \tan^3(x)}{5a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan(x)}{3a^2 \sqrt{a \cos^4(x)}}$$

[Out] $\cos(x) \sin(x) / a^2 / (a \cos(x)^4)^{(1/2)} + 4/3 \sin(x)^2 \tan(x) / a^2 / (a \cos(x)^4)^{(1/2)} + 6/5 \sin(x)^2 \tan(x)^3 / a^2 / (a \cos(x)^4)^{(1/2)} + 4/7 \sin(x)^2 \tan(x)^5 / a^2 / (a \cos(x)^4)^{(1/2)} + 1/9 \sin(x)^2 \tan(x)^7 / a^2 / (a \cos(x)^4)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3207, 3767}

$$\frac{\sin(x) \cos(x)}{a^2 \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^7(x)}{9a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan^5(x)}{7a^2 \sqrt{a \cos^4(x)}} + \frac{6 \sin^2(x) \tan^3(x)}{5a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan(x)}{3a^2 \sqrt{a \cos^4(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[x]^4)^(-5/2), x]

[Out] $(\cos[x] \sin[x]) / (a^2 \sqrt{a \cos[x]^4}) + (4 \sin[x]^2 \tan[x]) / (3 a^2 \sqrt{a \cos[x]^4}) + (6 \sin[x]^2 \tan[x]^3) / (5 a^2 \sqrt{a \cos[x]^4}) + (4 \sin[x]^2 \tan[x]^5) / (7 a^2 \sqrt{a \cos[x]^4}) + (\sin[x]^2 \tan[x]^7) / (9 a^2 \sqrt{a \cos[x]^4})$

Rule 3207

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] :> With[{fff = FreeFactors[Sin[e + f*x], x]}, Dist[((b*fff^n)^IntPart[p]*(b*Ssin[e + f*x]^n)^FracPart[p]) / (Sin[e + f*x]/fff)^(n*FracPart[p]), Int[ActivateTrig[u]*(Sin[e + f*x]/fff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a \cos^4(x))^{5/2}} dx &= \frac{\cos^2(x) \int \sec^{10}(x) dx}{a^2 \sqrt{a \cos^4(x)}} \\ &= -\frac{\cos^2(x) \text{Subst}\left(\int (1 + 4x^2 + 6x^4 + 4x^6 + x^8) dx, x, -\tan(x)\right)}{a^2 \sqrt{a \cos^4(x)}} \\ &= \frac{\cos(x) \sin(x)}{a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan(x)}{3a^2 \sqrt{a \cos^4(x)}} + \frac{6 \sin^2(x) \tan^3(x)}{5a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan^5(x)}{7a^2 \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^7(x)}{9a^2 \sqrt{a \cos^4(x)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 47, normalized size = 0.40

$$\frac{(130 \cos(2x) + 46 \cos(4x) + 10 \cos(6x) + \cos(8x) + 128) \tan(x) \sec^6(x)}{315a^2 \sqrt{a \cos^4(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[x]^4)^(-5/2), x]

[Out] ((128 + 130*cos[2*x] + 46*cos[4*x] + 10*cos[6*x] + Cos[8*x])*Sec[x]^6*Tan[x])/ (315*a^2*Sqrt[a*cos[x]^4])

fricas [A] time = 0.51, size = 45, normalized size = 0.38

$$\frac{(128 \cos(x)^8 + 64 \cos(x)^6 + 48 \cos(x)^4 + 40 \cos(x)^2 + 35)\sqrt{a \cos(x)^4} \sin(x)}{315 a^3 \cos(x)^{11}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(5/2), x, algorithm="fricas")

[Out] 1/315*(128*cos(x)^8 + 64*cos(x)^6 + 48*cos(x)^4 + 40*cos(x)^2 + 35)*sqrt(a*cos(x)^4)*sin(x)/(a^3*cos(x)^11)

giac [A] time = 0.53, size = 34, normalized size = 0.29

$$\frac{35 \tan(x)^9 + 180 \tan(x)^7 + 378 \tan(x)^5 + 420 \tan(x)^3 + 315 \tan(x)}{315 a^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(5/2), x, algorithm="giac")

[Out] 1/315*(35*tan(x)^9 + 180*tan(x)^7 + 378*tan(x)^5 + 420*tan(x)^3 + 315*tan(x))/a^(5/2)

maple [A] time = 0.15, size = 41, normalized size = 0.35

$$\frac{\sin(x) (128 (\cos^8(x)) + 64 (\cos^6(x)) + 48 (\cos^4(x)) + 40 (\cos^2(x)) + 35) \cos(x)}{315 (a (\cos^4(x)))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*cos(x)^4)^(5/2), x)

[Out] 1/315*sin(x)*(128*cos(x)^8+64*cos(x)^6+48*cos(x)^4+40*cos(x)^2+35)*cos(x)/(a*cos(x)^4)^(5/2)

maxima [A] time = 0.52, size = 34, normalized size = 0.29

$$\frac{35 \tan(x)^9 + 180 \tan(x)^7 + 378 \tan(x)^5 + 420 \tan(x)^3 + 315 \tan(x)}{315 a^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*cos(x)^4)^(5/2), x, algorithm="maxima")

[Out] 1/315*(35*tan(x)^9 + 180*tan(x)^7 + 378*tan(x)^5 + 420*tan(x)^3 + 315*tan(x))/a^(5/2)

mupad [B] time = 3.74, size = 306, normalized size = 2.62

$$\frac{e^{x4i} \sqrt{a \left(\frac{e^{-x1i}}{2} + \frac{e^{x1i}}{2} \right)^4} 2048i}{5 a^3 (e^{x2i} + 1)^5 (e^{x2i} + 2 e^{x4i} + e^{x6i})} - \frac{e^{x4i} \sqrt{a \left(\frac{e^{-x1i}}{2} + \frac{e^{x1i}}{2} \right)^4} 4096i}{3 a^3 (e^{x2i} + 1)^6 (e^{x2i} + 2 e^{x4i} + e^{x6i})} + \frac{e^{x4i} \sqrt{a \left(\frac{e^{-x1i}}{2} + \frac{e^{x1i}}{2} \right)^4} 12288i}{7 a^3 (e^{x2i} + 1)^7 (e^{x2i} + 2 e^{x4i} + e^{x6i})}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(a*cos(x)^4)^(5/2),x)
```

```
[Out] (exp(x*4i)*(a*(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*2048i)/(5*a^3*(exp(x*2i) + 1)^5*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (exp(x*4i)*(a*(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*4096i)/(3*a^3*(exp(x*2i) + 1)^6*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) + (exp(x*4i)*(a*(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*12288i)/(7*a^3*(exp(x*2i) + 1)^7*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (exp(x*4i)*(a*(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*1024i)/(a^3*(exp(x*2i) + 1)^8*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) + (exp(x*4i)*(a*(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*2048i)/(9*a^3*(exp(x*2i) + 1)^9*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i)))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a*cos(x)**4)**(5/2),x)
```

```
[Out] Timed out
```

3.57 $\int (b \cos^m(c + dx))^n dx$

Optimal. Leaf size=78

$$\frac{\sin(c + dx) \cos(c + dx) (b \cos^m(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(mn + 1); \frac{1}{2}(mn + 3); \cos^2(c + dx)\right)}{d(mn + 1)\sqrt{\sin^2(c + dx)}}$$

[Out] $-\cos(d*x+c)*(b*\cos(d*x+c)^m)^n*\text{hypergeom}([1/2, 1/2*m*n+1/2], [1/2*m*n+3/2], \cos(d*x+c)^2)*\sin(d*x+c)/d/(m*n+1)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3208, 2643}

$$\frac{\sin(c + dx) \cos(c + dx) (b \cos^m(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(mn + 1); \frac{1}{2}(mn + 3); \cos^2(c + dx)\right)}{d(mn + 1)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x]^m)^n, x]$

[Out] $-\left(\text{Cos}[c + d*x]*(b*\text{Cos}[c + d*x]^m)^n*\text{Hypergeometric2F1}[1/2, (1 + m*n)/2, (3 + m*n)/2, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x]\right)/(d*(1 + m*n)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x$ && $!\text{IntegerQ}[2*n]$

Rule 3208

$\text{Int}[(u_*)*((b_*)*((c_*)*\sin[(e_*) + (f_*)*(x_)]^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Dist}[(b^{\text{IntPart}[p]}*(b*(c*\text{Sin}[e + f*x])^n)^{\text{FracPart}[p]})/(c*\text{Sin}[e + f*x])^{(n*\text{FracPart}[p])}, \text{Int}[\text{ActivateTrig}[u]*(c*\text{Sin}[e + f*x])^{(n*p)}, x], x] /;$ $\text{FreeQ}\{b, c, e, f, n, p\}, x$ && $!\text{IntegerQ}[p]$ && $!\text{IntegerQ}[n]$ && $(\text{EqQ}[u, 1] \parallel \text{MatchQ}[u, ((d_*)*(\text{trig}_)[e + f*x])^{(m_*)}) /;$ $\text{FreeQ}\{d, m\}, x$ && $\text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}\}$)

Rubi steps

$$\begin{aligned} \int (b \cos^m(c + dx))^n dx &= (\cos^{-mn}(c + dx) (b \cos^m(c + dx))^n) \int \cos^{mn}(c + dx) dx \\ &= \frac{\cos(c + dx) (b \cos^m(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + mn); \frac{1}{2}(3 + mn); \cos^2(c + dx)\right) \sin(c + dx)}{d(1 + mn)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 72, normalized size = 0.92

$$\frac{\sqrt{\sin^2(c + dx)} \cot(c + dx) (b \cos^m(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(mn + 1); \frac{1}{2}(mn + 3); \cos^2(c + dx)\right)}{d(mn + 1)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x]^m)^n,x]

[Out] -(((b*cos[c + d*x]^m)^n*cot[c + d*x]*Hypergeometric2F1[1/2, (1 + m*n)/2, (3 + m*n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2]))/(d*(1 + m*n))

fricas [F] time = 0.45, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)^m\right)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c)^m)^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c)^m)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c)^m)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c)^m)^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c)^m)^n, x)

maple [F] time = 0.19, size = 0, normalized size = 0.00

$$\int (b (\cos^m(dx + c)))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c)^m)^n,x)

[Out] int((b*cos(d*x+c)^m)^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c)^m)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c)^m)^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c)^m)^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (b \cos(c + dx)^m)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x)^m)^n,x)

[Out] int((b*cos(c + d*x)^m)^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos^m(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c)**m)**n,x)

[Out] Integral((b*cos(c + d*x)**m)**n, x)

3.58 $\int (c \cos^m(a + bx))^{5/2} dx$

Optimal. Leaf size=89

$$\frac{2c^2 \sin(a + bx) \cos^{2m+1}(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(5m + 2); \frac{1}{4}(5m + 6); \cos^2(a + bx)\right)}{b(5m + 2) \sqrt{\sin^2(a + bx)}}$$

[Out] $-2*c^2*\cos(b*x+a)^{(1+2*m)}*hypergeom([1/2, 1/2+5/4*m], [3/2+5/4*m], \cos(b*x+a)^2)*\sin(b*x+a)*(c*\cos(b*x+a)^m)^{(1/2)}/b/(2+5*m)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3208, 2643}

$$\frac{2c^2 \sin(a + bx) \cos^{2m+1}(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(5m + 2); \frac{1}{4}(5m + 6); \cos^2(a + bx)\right)}{b(5m + 2) \sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Cos}[a + b*x]^m)^{(5/2)}, x]$

[Out] $(-2*c^2*\text{Cos}[a + b*x]^{(1 + 2*m)}*\text{Sqrt}[c*\text{Cos}[a + b*x]^m]*\text{Hypergeometric2F1}[1/2, (2 + 5*m)/4, (6 + 5*m)/4, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(b*(2 + 5*m)*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3208

$\text{Int}[(u_*)*((b_*)*((c_*)*\sin[(e_*) + (f_*)*(x_)]^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Dist}[(b^{\text{IntPart}[p]}*(b*(c*\text{Sin}[e + f*x])^n)^{\text{FracPart}[p]})/(c*\text{Sin}[e + f*x])^{(n*\text{FracPart}[p])}, \text{Int}[\text{ActivateTrig}[u]*(c*\text{Sin}[e + f*x])^{(n*p)}, x], x] /;$ FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_*)*(trig_)[e + f*x])^{(m_*)} /]; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])

Rubi steps

$$\int (c \cos^m(a + bx))^{5/2} dx = \left(c^2 \cos^{-\frac{m}{2}}(a + bx) \sqrt{c \cos^m(a + bx)} \right) \int \cos^{\frac{5m}{2}}(a + bx) dx$$

$$= \frac{2c^2 \cos^{1+2m}(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2 + 5m); \frac{1}{4}(6 + 5m); \cos^2(a + bx)\right) \sin(a + bx)}{b(2 + 5m) \sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.16, size = 74, normalized size = 0.83

$$\frac{2\sqrt{\sin^2(a + bx)} \cot(a + bx) (c \cos^m(a + bx))^{5/2} {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(5m + 2); \frac{1}{4}(5m + 6); \cos^2(a + bx)\right)}{b(5m + 2)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x]^m)^(5/2), x]

[Out] $(-2*(c*\cos[a + b*x]^m)^{(5/2)}*\cot[a + b*x]*\text{Hypergeometric2F1}[1/2, (2 + 5*m)/4, (6 + 5*m)/4, \cos[a + b*x]^2]*\sqrt{\sin[a + b*x]^2})/(b*(2 + 5*m))$

fricas [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(5/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a)^m)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(5/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a)^m)^(5/2), x)

maple [F] time = 0.41, size = 0, normalized size = 0.00

$$\int (c (\cos^m(bx + a)))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a)^m)^(5/2), x)

[Out] int((c*cos(b*x+a)^m)^(5/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a)^m)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(5/2), x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a)^m)^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (c \cos(a + bx)^m)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x)^m)^(5/2), x)

[Out] int((c*cos(a + b*x)^m)^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)**m)**(5/2), x)

[Out] Timed out

3.59 $\int (c \cos^m(a + bx))^{3/2} dx$

Optimal. Leaf size=83

$$\frac{2c \sin(a + bx) \cos^{m+1}(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(3m + 2); \frac{3(m+2)}{4}; \cos^2(a + bx)\right)}{b(3m + 2) \sqrt{\sin^2(a + bx)}}$$

[Out] $-2*c*\cos(b*x+a)^{(1+m)}*\text{hypergeom}([1/2, 1/2+3/4*m], [3/2+3/4*m], \cos(b*x+a)^2)*\sin(b*x+a)*(c*\cos(b*x+a)^m)^{(1/2)}/b/(2+3*m)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3208, 2643}

$$\frac{2c \sin(a + bx) \cos^{m+1}(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(3m + 2); \frac{3(m+2)}{4}; \cos^2(a + bx)\right)}{b(3m + 2) \sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Cos}[a + b*x]^m)^{(3/2)}, x]$

[Out] $(-2*c*\text{Cos}[a + b*x]^{(1 + m)}*\text{Sqrt}[c*\text{Cos}[a + b*x]^m]*\text{Hypergeometric2F1}[1/2, (2 + 3*m)/4, (3*(2 + m))/4, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(b*(2 + 3*m)*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x$ && $\text{IntegerQ}[2*n]$

Rule 3208

$\text{Int}[(u_*)*((b_*)*((c_*)*\sin[(e_*) + (f_*)*(x_)]^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Dist}[(b^{\text{IntPart}[p]}*(b*(c*\text{Sin}[e + f*x])^n)^{\text{FracPart}[p]})/(c*\text{Sin}[e + f*x])^{(n*\text{FracPart}[p])}, \text{Int}[\text{ActivateTrig}[u]*(c*\text{Sin}[e + f*x])^{(n*p)}, x], x] /;$ $\text{FreeQ}\{b, c, e, f, n, p\}, x$ && $\text{IntegerQ}[p]$ && $\text{IntegerQ}[n]$ && $(\text{EqQ}[u, 1] \parallel \text{MatchQ}[u, ((d_*)*(\text{trig}_)[e + f*x])^{(m_*)}) /;$ $\text{FreeQ}\{d, m\}, x$ && $\text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}\}$)

Rubi steps

$$\int (c \cos^m(a + bx))^{3/2} dx = \left(c \cos^{-\frac{m}{2}}(a + bx) \sqrt{c \cos^m(a + bx)} \right) \int \cos^{\frac{3m}{2}}(a + bx) dx$$

$$= \frac{2c \cos^{1+m}(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2 + 3m); \frac{3(2+m)}{4}; \cos^2(a + bx)\right) \sin(a + bx)}{b(2 + 3m) \sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.11, size = 72, normalized size = 0.87

$$\frac{2\sqrt{\sin^2(a + bx)} \cot(a + bx) (c \cos^m(a + bx))^{3/2} {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(3m + 2); \frac{3(m+2)}{4}; \cos^2(a + bx)\right)}{b(3m + 2)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x]^m)^(3/2), x]

[Out] $(-2*(c*\cos[a + b*x]^m)^{(3/2)}*\cot[a + b*x]*\text{Hypergeometric2F1}[1/2, (2 + 3*m)/4, (3*(2 + m))/4, \cos[a + b*x]^2]*\sqrt{\sin[a + b*x]^2})/(b*(2 + 3*m))$

fricas [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(3/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos(bx + a)^m)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(3/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a)^m)^(3/2), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int (c (\cos^m(bx + a)))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a)^m)^(3/2), x)

[Out] int((c*cos(b*x+a)^m)^(3/2), x)

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(3/2), x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: BINDING-STACK overflow at size 10240. Stack can probably be resized. Proceed with caution.

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (c \cos(a + bx)^m)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x)^m)^(3/2), x)

[Out] int((c*cos(a + b*x)^m)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos^m(a + bx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)**m)**(3/2), x)

[Out] Integral((c*cos(a + b*x)**m)**(3/2), x)

3.60 $\int \sqrt{c \cos^m(a + bx)} dx$

Optimal. Leaf size=74

$$\frac{2 \sin(a + bx) \cos(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{m+2}{4}; \frac{m+6}{4}; \cos^2(a + bx)\right)}{b(m+2) \sqrt{\sin^2(a + bx)}}$$

[Out] $-2*\cos(b*x+a)*\text{hypergeom}([1/2, 1/2+1/4*m], [3/2+1/4*m], \cos(b*x+a)^2)*\sin(b*x+a)*(c*\cos(b*x+a)^m)^{(1/2)}/b/(2+m)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3208, 2643}

$$\frac{2 \sin(a + bx) \cos(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{m+2}{4}; \frac{m+6}{4}; \cos^2(a + bx)\right)}{b(m+2) \sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Cos[a + b*x]^m], x]

[Out] $(-2*\text{Cos}[a + b*x]*\text{Sqrt}[c*\text{Cos}[a + b*x]^m]*\text{Hypergeometric2F1}[1/2, (2 + m)/4, (6 + m)/4, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(b*(2 + m)*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3208

Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :> Dist[(b^IntPart[p]*(b*(c*Sin[e + f*x])^n)^FracPart[p])/(c*Sin[e + f*x])^(n*FracPart[p]), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])

Rubi steps

$$\begin{aligned} \int \sqrt{c \cos^m(a + bx)} dx &= \left(\cos^{-\frac{m}{2}}(a + bx) \sqrt{c \cos^m(a + bx)} \right) \int \cos^{\frac{m}{2}}(a + bx) dx \\ &= \frac{2 \cos(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{4}; \frac{6+m}{4}; \cos^2(a + bx)\right) \sin(a + bx)}{b(2 + m) \sqrt{\sin^2(a + bx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 68, normalized size = 0.92

$$\frac{2 \sqrt{\sin^2(a + bx)} \cot(a + bx) \sqrt{c \cos^m(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{m+2}{4}; \frac{m+6}{4}; \cos^2(a + bx)\right)}{b(m+2)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Cos[a + b*x]^m], x]

[Out] $(-2\sqrt{c\cos[a + b*x]^m}*\cot[a + b*x]*\text{Hypergeometric2F1}[1/2, (2 + m)/4, (6 + m)/4, \cos[a + b*x]^2]*\sqrt{\sin[a + b*x]^2})/(b*(2 + m))$

fricas [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(1/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \cos(bx + a)^m} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(c*cos(b*x + a)^m), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \sqrt{c (\cos^m(bx + a))} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a)^m)^(1/2), x)

[Out] int((c*cos(b*x+a)^m)^(1/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \cos(bx + a)^m} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(c*cos(b*x + a)^m), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{c \cos(a + bx)^m} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x)^m)^(1/2), x)

[Out] int((c*cos(a + b*x)^m)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \cos^m(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)**m)**(1/2), x)

[Out] Integral(sqrt(c*cos(a + b*x)**m), x)

3.61 $\int \frac{1}{\sqrt{c \cos^m(a+bx)}} dx$

Optimal. Leaf size=80

$$\frac{2 \sin(a+bx) \cos(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2-m}{4}; \frac{6-m}{4}; \cos^2(a+bx)\right)}{b(2-m)\sqrt{\sin^2(a+bx)}\sqrt{c \cos^m(a+bx)}}$$

[Out] $-2*\cos(b*x+a)*\text{hypergeom}([1/2, 1/2-1/4*m], [3/2-1/4*m], \cos(b*x+a)^2)*\sin(b*x+a)/b/(2-m)/(c*\cos(b*x+a)^m)^{(1/2)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3208, 2643}

$$\frac{2 \sin(a+bx) \cos(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2-m}{4}; \frac{6-m}{4}; \cos^2(a+bx)\right)}{b(2-m)\sqrt{\sin^2(a+bx)}\sqrt{c \cos^m(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[c*Cos[a + b*x]^m], x]

[Out] $(-2*\text{Cos}[a + b*x]*\text{Hypergeometric2F1}[1/2, (2 - m)/4, (6 - m)/4, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(b*(2 - m)*\text{Sqrt}[c*\text{Cos}[a + b*x]^m]*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3208

Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :> Dist[(b^IntPart[p]*(b*(c*Sin[e + f*x])^n)^FracPart[p])/(c*Sin[e + f*x])^(n*FracPart[p]), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{c \cos^m(a+bx)}} dx &= \frac{\cos^{\frac{m}{2}}(a+bx) \int \cos^{-\frac{m}{2}}(a+bx) dx}{\sqrt{c \cos^m(a+bx)}} \\ &= -\frac{2 \cos(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2-m}{4}; \frac{6-m}{4}; \cos^2(a+bx)\right) \sin(a+bx)}{b(2-m)\sqrt{c \cos^m(a+bx)}\sqrt{\sin^2(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 72, normalized size = 0.90

$$\frac{2\sqrt{\sin^2(a+bx)} \cot(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2-m}{4}; \frac{6-m}{4}; \cos^2(a+bx)\right)}{b(m-2)\sqrt{c \cos^m(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[c*Cos[a + b*x]^m], x]

[Out] (2*Cot[a + b*x]*Hypergeometric2F1[1/2, (2 - m)/4, (6 - m)/4, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*(-2 + m)*Sqrt[c*Cos[a + b*x]^m])

fricas [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(1/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \cos(bx + a)^m}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(1/2), x, algorithm="giac")

[Out] integrate(1/sqrt(c*cos(b*x + a)^m), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c (\cos^m(bx + a))}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a)^m)^(1/2), x)

[Out] int(1/(c*cos(b*x+a)^m)^(1/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \cos(bx + a)^m}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(1/2), x, algorithm="maxima")

[Out] integrate(1/sqrt(c*cos(b*x + a)^m), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{c \cos(a + bx)^m}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x)^m)^(1/2), x)

[Out] int(1/(c*cos(a + b*x)^m)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(c*cos(b*x+a)**m)**(1/2),x)
```

```
[Out] Integral(1/sqrt(c*cos(a + b*x)**m), x)
```

$$3.62 \quad \int \frac{1}{(c \cos^m(a+bx))^{3/2}} dx$$

Optimal. Leaf size=89

$$\frac{2 \sin(a+bx) \cos^{1-m}(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2-3m); \frac{3(2-m)}{4}; \cos^2(a+bx)\right)}{bc(2-3m)\sqrt{\sin^2(a+bx)}\sqrt{c \cos^m(a+bx)}}$$

[Out] $-2*\cos(b*x+a)^{(1-m)}*\text{hypergeom}([1/2, 1/2-3/4*m], [3/2-3/4*m], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(2-3*m)/(c*\cos(b*x+a)^m)^{(1/2)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3208, 2643}

$$\frac{2 \sin(a+bx) \cos^{1-m}(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2-3m); \frac{3(2-m)}{4}; \cos^2(a+bx)\right)}{bc(2-3m)\sqrt{\sin^2(a+bx)}\sqrt{c \cos^m(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x]^m)^(-3/2), x]

[Out] $(-2*\cos[a + b*x]^{(1-m)}*\text{Hypergeometric2F1}[1/2, (2-3*m)/4, (3*(2-m))/4, \cos[a + b*x]^2]*\sin[a + b*x])/(b*c*(2-3*m)*\text{Sqrt}[c*\cos[a + b*x]^m]*\text{Sqrt}[\sin[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3208

Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :> Dist[(b^IntPart[p]*(b*(c*Sin[e + f*x])^n)^FracPart[p])/(c*Sin[e + f*x])^(n*FracPart[p]), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int \frac{1}{(c \cos^m(a+bx))^{3/2}} dx &= \frac{\cos^{\frac{m}{2}}(a+bx) \int \cos^{-\frac{3m}{2}}(a+bx) dx}{c\sqrt{c \cos^m(a+bx)}} \\ &= -\frac{2 \cos^{1-m}(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2-3m); \frac{3(2-m)}{4}; \cos^2(a+bx)\right) \sin(a+bx)}{bc(2-3m)\sqrt{c \cos^m(a+bx)}\sqrt{\sin^2(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 72, normalized size = 0.81

$$\frac{\sqrt{\sin^2(a+bx)} \cot(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2-3m); -\frac{3}{4}(m-2); \cos^2(a+bx)\right)}{\left(b - \frac{3bm}{2}\right) (c \cos^m(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x]^m)^(-3/2),x]

[Out] -((Cot[a + b*x]*Hypergeometric2F1[1/2, (2 - 3*m)/4, (-3*(-2 + m))/4, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/((b - (3*b*m)/2)*(c*cos[a + b*x]^m)^(3/2)))

fricas [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(3/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(3/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a)^m)^(-3/2), x)

maple [F] time = 0.07, size = 0, normalized size = 0.00

$$\int \frac{1}{(c(\cos^m(bx + a)))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a)^m)^(3/2),x)

[Out] int(1/(c*cos(b*x+a)^m)^(3/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(3/2),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a)^m)^(-3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(c \cos(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x)^m)^(3/2),x)

[Out] int(1/(c*cos(a + b*x)^m)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos^m(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(c*cos(b*x+a)**m)**(3/2), x)
```

```
[Out] Integral((c*cos(a + b*x)**m)**(-3/2), x)
```

$$3.63 \quad \int \frac{1}{(c \cos^m(a+bx))^{5/2}} dx$$

Optimal. Leaf size=89

$$\frac{2 \sin(a+bx) \cos^{1-2m}(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2-5m); \frac{1}{4}(6-5m); \cos^2(a+bx)\right)}{bc^2(2-5m)\sqrt{\sin^2(a+bx)}\sqrt{c \cos^m(a+bx)}}$$

[Out] $-2*\cos(b*x+a)^{(1-2*m)}*\text{hypergeom}([1/2, 1/2-5/4*m], [3/2-5/4*m], \cos(b*x+a)^2)*\sin(b*x+a)/b/c^2/(2-5*m)/(c*\cos(b*x+a)^m)^{(1/2)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3208, 2643}

$$\frac{2 \sin(a+bx) \cos^{1-2m}(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2-5m); \frac{1}{4}(6-5m); \cos^2(a+bx)\right)}{bc^2(2-5m)\sqrt{\sin^2(a+bx)}\sqrt{c \cos^m(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*cos[a + b*x]^m)^(-5/2), x]

[Out] $(-2*\cos[a + b*x]^{(1 - 2*m)}*\text{Hypergeometric2F1}[1/2, (2 - 5*m)/4, (6 - 5*m)/4, \cos[a + b*x]^2]*\sin[a + b*x])/(b*c^2*(2 - 5*m)*\text{Sqrt}[c*\cos[a + b*x]^m]*\text{Sqrt}[\sin[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3208

Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_.))^(p_), x_Symbol] :> Dist[(b^IntPart[p]*(b*(c*Sin[e + f*x])^n)^FracPart[p])/(c*Sin[e + f*x])^(n*FracPart[p]), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])

Rubi steps

$$\begin{aligned} \int \frac{1}{(c \cos^m(a+bx))^{5/2}} dx &= \frac{\cos^{\frac{m}{2}}(a+bx) \int \cos^{-\frac{5m}{2}}(a+bx) dx}{c^2 \sqrt{c \cos^m(a+bx)}} \\ &= \frac{2 \cos^{1-2m}(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2-5m); \frac{1}{4}(6-5m); \cos^2(a+bx)\right) \sin(a+bx)}{bc^2(2-5m)\sqrt{c \cos^m(a+bx)}\sqrt{\sin^2(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 74, normalized size = 0.83

$$\frac{\sqrt{\sin^2(a+bx)} \cot(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2-5m); \frac{1}{4}(6-5m); \cos^2(a+bx)\right)}{\left(b - \frac{5bm}{2}\right) (c \cos^m(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*cos[a + b*x]^m)^(-5/2), x]

[Out] -((Cot[a + b*x]*Hypergeometric2F1[1/2, (2 - 5*m)/4, (6 - 5*m)/4, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/((b - (5*b*m)/2)*(c*cos[a + b*x]^m)^(5/2)))

fricas [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(5/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(5/2), x, algorithm="giac")

[Out] integrate((c*cos(b*x + a)^m)^(-5/2), x)

maple [F] time = 0.07, size = 0, normalized size = 0.00

$$\int \frac{1}{(c (\cos^m(bx + a)))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(b*x+a)^m)^(5/2), x)

[Out] int(1/(c*cos(b*x+a)^m)^(5/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)^m)^(5/2), x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a)^m)^(-5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(c \cos(a + bx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*cos(a + b*x)^m)^(5/2), x)

[Out] int(1/(c*cos(a + b*x)^m)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \cos^m(a + bx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*cos(b*x+a)**m)**(5/2), x)

[Out] Integral((c*cos(a + b*x)**m)**(-5/2), x)

3.64 $\int (c \cos^m(a + bx))^{\frac{1}{m}} dx$

Optimal. Leaf size=24

$$\frac{\tan(a + bx) (c \cos^m(a + bx))^{\frac{1}{m}}}{b}$$

[Out] (c*cos(b*x+a)^m)^(1/m)*tan(b*x+a)/b

Rubi [A] time = 0.02, antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3208, 2637}

$$\frac{\tan(a + bx) (c \cos^m(a + bx))^{\frac{1}{m}}}{b}$$

Antiderivative was successfully verified.

[In] Int[(c*Cos[a + b*x]^m)^m^(-1), x]

[Out] ((c*Cos[a + b*x]^m)^m^(-1)*Tan[a + b*x])/b

Rule 2637

Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3208

Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.))^(p_.), x_Symbol] := Dist[(b^IntPart[p]*(b*(c*SIN[e + f*x])^n)^FracPart[p])/(c*SIN[e + f*x])^(n*FracPart[p]), Int[ActivateTrig[u]*(c*SIN[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])

Rubi steps

$$\begin{aligned} \int (c \cos^m(a + bx))^{\frac{1}{m}} dx &= \left((c \cos^m(a + bx))^{\frac{1}{m}} \sec(a + bx) \right) \int \cos(a + bx) dx \\ &= \frac{(c \cos^m(a + bx))^{\frac{1}{m}} \tan(a + bx)}{b} \end{aligned}$$

Mathematica [A] time = 0.03, size = 24, normalized size = 1.00

$$\frac{\tan(a + bx) (c \cos^m(a + bx))^{\frac{1}{m}}}{b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Cos[a + b*x]^m)^m^(-1), x]

[Out] ((c*Cos[a + b*x]^m)^m^(-1)*Tan[a + b*x])/b

fricas [A] time = 0.54, size = 15, normalized size = 0.62

$$\frac{c^{\left(\frac{1}{m}\right)} \sin(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(1/m),x, algorithm="fricas")

[Out] c^(1/m)*sin(b*x + a)/b

giac [B] time = 5.14, size = 300, normalized size = 12.50

$$\frac{2 \left(|c|^{\left(\frac{1}{m}\right)} \tan\left(\frac{1}{2}bx + \frac{1}{2}a + \frac{\pi \operatorname{sgn}(c)}{4m} - \frac{\pi}{4m}\right)^2 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^3 - |c|^{\left(\frac{1}{m}\right)} \tan\left(\frac{1}{2}bx + \frac{1}{2}a + \frac{\pi \operatorname{sgn}(c)}{4m} - \frac{\pi}{4m}\right)^2 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 \right)}{b \tan\left(\frac{1}{2}bx + \frac{1}{2}a + \frac{\pi \operatorname{sgn}(c)}{4m} - \frac{\pi}{4m}\right)^2 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 + 2b \tan\left(\frac{1}{2}bx + \frac{1}{2}a + \frac{\pi \operatorname{sgn}(c)}{4m} - \frac{\pi}{4m}\right)^2 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 + b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(1/m),x, algorithm="giac")

[Out] 2*(abs(c)^(1/m)*tan(1/2*b*x + 1/2*a + 1/4*pi*sgn(c)/m - 1/4*pi/m)^2*tan(1/2*b*x + 1/2*a)^3 - abs(c)^(1/m)*tan(1/2*b*x + 1/2*a + 1/4*pi*sgn(c)/m - 1/4*pi/m)^2*tan(1/2*b*x + 1/2*a) + 4*abs(c)^(1/m)*tan(1/2*b*x + 1/2*a + 1/4*pi*sgn(c)/m - 1/4*pi/m)*tan(1/2*b*x + 1/2*a)^2 - abs(c)^(1/m)*tan(1/2*b*x + 1/2*a)^3 + abs(c)^(1/m)*tan(1/2*b*x + 1/2*a))/(b*tan(1/2*b*x + 1/2*a + 1/4*pi*sgn(c)/m - 1/4*pi/m)^2*tan(1/2*b*x + 1/2*a)^4 + 2*b*tan(1/2*b*x + 1/2*a + 1/4*pi*sgn(c)/m - 1/4*pi/m)^2*tan(1/2*b*x + 1/2*a)^2 + b*tan(1/2*b*x + 1/2*a)^4 + b*tan(1/2*b*x + 1/2*a + 1/4*pi*sgn(c)/m - 1/4*pi/m)^2 + 2*b*tan(1/2*b*x + 1/2*a)^2 + b)

maple [F] time = 0.14, size = 0, normalized size = 0.00

$$\int (c (\cos^m (bx + a)))^{\frac{1}{m}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(b*x+a)^m)^(1/m),x)

[Out] int((c*cos(b*x+a)^m)^(1/m),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \cos (bx + a)^m)^{\left(\frac{1}{m}\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)^m)^(1/m),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a)^m)^(1/m), x)

mupad [B] time = 0.40, size = 40, normalized size = 1.67

$$\frac{\sin(2a + 2bx) (c \cos(a + bx)^m)^{1/m}}{b (\cos(2a + 2bx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*cos(a + b*x)^m)^(1/m),x)

[Out] (sin(2*a + 2*b*x)*(c*cos(a + b*x)^m)^(1/m))/(b*(cos(2*a + 2*b*x) + 1))

sympy [A] time = 1.58, size = 65, normalized size = 2.71

$$\left\{ \begin{array}{ll} x (c \cos^m(a))^{\frac{1}{m}} & \text{for } b = 0 \\ x (0^m c)^{\frac{1}{m}} & \text{for } a = -bx + \frac{\pi}{2} \vee a = -bx + \frac{3\pi}{2} \\ \frac{c^{\frac{1}{m}} (\cos^m(a+bx))^{\frac{1}{m}} \sin(a+bx)}{b \cos(a+bx)} & \text{otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*cos(b*x+a)**m)**(1/m),x)

[Out] Piecewise((x*(c*cos(a)**m)**(1/m), Eq(b, 0)), (x*(0**m*c)**(1/m), Eq(a, -b*x + pi/2) | Eq(a, -b*x + 3*pi/2)), (c**(1/m)*(cos(a + b*x)**m)**(1/m)*sin(a + b*x)/(b*cos(a + b*x)), True))

3.65 $\int (a(b \cos(c + dx))^p)^n dx$

Optimal. Leaf size=80

$$\frac{\sin(c + dx) \cos(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(np + 1); \frac{1}{2}(np + 3); \cos^2(c + dx)\right) (a(b \cos(c + dx))^p)^n}{d(np + 1)\sqrt{\sin^2(c + dx)}}$$

[Out] $-\cos(d*x+c)*(a*(b*\cos(d*x+c))^p)^n*\text{hypergeom}([1/2, 1/2*n*p+1/2], [1/2*n*p+3/2], \cos(d*x+c)^2)*\sin(d*x+c)/d/(n*p+1)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3208, 2643}

$$\frac{\sin(c + dx) \cos(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(np + 1); \frac{1}{2}(np + 3); \cos^2(c + dx)\right) (a(b \cos(c + dx))^p)^n}{d(np + 1)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a*(b*\text{Cos}[c + d*x])^p)^n, x]$

[Out] $-\left(\left(\text{Cos}[c + d*x]*(a*(b*\text{Cos}[c + d*x])^p)^n*\text{Hypergeometric2F1}[1/2, (1 + n*p)/2, (3 + n*p)/2, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x]\right)/\left(d*(1 + n*p)*\text{Sqrt}[\text{Sin}[c + d*x]^2]\right)\right)$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3208

$\text{Int}[(u_*)*((b_*)*((c_*)*\sin[(e_*) + (f_*)*(x_*)]^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Dist}[(b^{\text{IntPart}[p]}*(b*(c*\text{Sin}[e + f*x])^n)^{\text{FracPart}[p]})/(c*\text{Sin}[e + f*x])^{(n*\text{FracPart}[p])}, \text{Int}[\text{ActivateTrig}[u]*(c*\text{Sin}[e + f*x])^{(n*p)}, x], x] /;$ FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_*)*(trig_)[e + f*x])^{(m_*)} /]; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]]

Rubi steps

$$\int (a(b \cos(c + dx))^p)^n dx = \left((b \cos(c + dx))^{-np} (a(b \cos(c + dx))^p)^n \right) \int (b \cos(c + dx))^{np} dx$$

$$= \frac{\cos(c + dx) (a(b \cos(c + dx))^p)^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + np); \frac{1}{2}(3 + np); \cos^2(c + dx)\right) \sin(c + dx)}{d(1 + np)\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.05, size = 74, normalized size = 0.92

$$\frac{\sqrt{\sin^2(c + dx)} \cot(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(np + 1); \frac{1}{2}(np + 3); \cos^2(c + dx)\right) (a(b \cos(c + dx))^p)^n}{d(np + 1)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*(b*cos[c + d*x])^p)^n,x]

[Out] -(((a*(b*cos[c + d*x])^p)^n*Cot[c + d*x]*Hypergeometric2F1[1/2, (1 + n*p)/2, (3 + n*p)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(1 + n*p)))

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral}\left(\left((b \cos(dx + c))^p a\right)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*(b*cos(d*x+c))^p)^n,x, algorithm="fricas")

[Out] integral(((b*cos(d*x + c))^p*a)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left((b \cos(dx + c))^p a \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*(b*cos(d*x+c))^p)^n,x, algorithm="giac")

[Out] integrate(((b*cos(d*x + c))^p*a)^n, x)

maple [F] time = 0.20, size = 0, normalized size = 0.00

$$\int \left(a (b \cos(dx + c))^p \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*(b*cos(d*x+c))^p)^n,x)

[Out] int((a*(b*cos(d*x+c))^p)^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left((b \cos(dx + c))^p a \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*(b*cos(d*x+c))^p)^n,x, algorithm="maxima")

[Out] integrate(((b*cos(d*x + c))^p*a)^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(a (b \cos(c + dx))^p \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*(b*cos(c + d*x))^p)^n,x)

[Out] int((a*(b*cos(c + d*x))^p)^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left(a (b \cos(c + dx))^p \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*(b*cos(d*x+c))**p)**n,x)

[Out] Integral((a*(b*cos(c + d*x))**p)**n, x)

3.66 $\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal. Leaf size=123

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{9/2}}{11b^4d} + \frac{18 \sin(c + dx)(b \cos(c + dx))^{5/2}}{77b^2d} + \frac{30 \sin(c + dx)\sqrt{b \cos(c + dx)}}{77d} + \frac{30b\sqrt{\cos(c + dx)}}{77d\sqrt{bc}}$$

[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b^2/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^4/d+30/77*b*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+30/77*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d

Rubi [A] time = 0.09, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{9/2}}{11b^4d} + \frac{18 \sin(c + dx)(b \cos(c + dx))^{5/2}}{77b^2d} + \frac{30 \sin(c + dx)\sqrt{b \cos(c + dx)}}{77d} + \frac{30b\sqrt{\cos(c + dx)}}{77d\sqrt{bc}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^5*Sqrt[b*Cos[c + d*x]],x]

[Out] (30*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*d*Sqrt[b*Cos[c + d*x]]) + (30*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b^2*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^4*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Ssin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Ssin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Ssin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos^5(c+dx)\sqrt{b\cos(c+dx)} dx &= \frac{\int (b\cos(c+dx))^{11/2} dx}{b^5} \\
&= \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^4d} + \frac{9 \int (b\cos(c+dx))^{7/2} dx}{11b^3} \\
&= \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^2d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^4d} + \frac{45 \int (b\cos(c+dx))^{3/2} dx}{11b^3} \\
&= \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^2d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^4d} \\
&= \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^2d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^4d} \\
&= \frac{30b\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx)\middle|2\right)}{77d\sqrt{b\cos(c+dx)}} + \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^2d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^4d}
\end{aligned}$$

Mathematica [A] time = 0.20, size = 83, normalized size = 0.67

$$\frac{\sqrt{b\cos(c+dx)} \left(240F\left(\frac{1}{2}(c+dx)\middle|2\right) + (290\sin(c+dx) + 57\sin(3(c+dx)) + 7\sin(5(c+dx)))\sqrt{\cos(c+dx)} \right)}{616d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^5*Sqrt[b*Cos[c + d*x]], x]

[Out] (Sqrt[b*Cos[c + d*x]]*(240*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(290*Sin[c + d*x] + 57*Sin[3*(c + d*x)] + 7*Sin[5*(c + d*x)])))/(616*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.49, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b\cos(dx+c)}\cos(dx+c)^5, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5*(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^5, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b\cos(dx+c)}\cos(dx+c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5*(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^5, x)

maple [A] time = 0.16, size = 234, normalized size = 1.90

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b \left(448\left(\cos^{13}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1568\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 2384\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1280\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 448\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 64\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right)}{77\sqrt{-b}\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^5*(b*cos(d*x+c))^(1/2),x)`

[Out] $-2/77*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*(448*\cos(1/2*d*x+1/2*c)^{13}-1568*\cos(1/2*d*x+1/2*c)^{11}+2384*\cos(1/2*d*x+1/2*c)^9-2040*\cos(1/2*d*x+1/2*c)^7+1084*\cos(1/2*d*x+1/2*c)^5-370*\cos(1/2*d*x+1/2*c)^3+15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+62*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \cos(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^5, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^5 \sqrt{b \cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^5*(b*cos(c + d*x))^(1/2),x)`

[Out] `int(cos(c + d*x)^5*(b*cos(c + d*x))^(1/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**5*(b*cos(d*x+c))**(1/2),x)`

[Out] Timed out

3.67 $\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal. Leaf size=97

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{7/2}}{9b^3d} + \frac{14 \sin(c + dx)(b \cos(c + dx))^{3/2}}{45bd} + \frac{14E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[Out] 14/45*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b/d+2/9*(b*cos(d*x+c))^(7/2)*sin(d*x+c)/b^3/d+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.07, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{7/2}}{9b^3d} + \frac{14 \sin(c + dx)(b \cos(c + dx))^{3/2}}{45bd} + \frac{14E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^4*Sqrt[b*Cos[c + d*x]],x]

[Out] (14*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*d*Sqrt[Cos[c + d*x]]) + (14*(b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(45*b*d) + (2*(b*Cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b^3*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x])*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[b*SIN[c + d*x]]/Sqrt[SIN[c + d*x]], Int[Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^4} \\
&= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3 d} + \frac{7 \int (b \cos(c + dx))^{5/2} dx}{9b^2} \\
&= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45bd} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3 d} + \frac{7}{15} \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45bd} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3 d} + \frac{(7\sqrt{b \cos(c + dx)}) E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)}} \\
&= \frac{14\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45bd} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3 d}
\end{aligned}$$

Mathematica [A] time = 0.15, size = 75, normalized size = 0.77

$$\frac{\sqrt{b \cos(c + dx)} \left(168E\left(\frac{1}{2}(c + dx) \middle| 2\right) + (38 \sin(2(c + dx)) + 5 \sin(4(c + dx)))\sqrt{\cos(c + dx)} \right)}{180d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^4*Sqrt[b*Cos[c + d*x]], x]

[Out] (Sqrt[b*Cos[c + d*x]]*(168*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)])))/(180*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} \cos(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^4, x)

maple [B] time = 0.13, size = 221, normalized size = 2.28

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b\left(160\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 480\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 616\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 45\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \dots\right)}\right)}{45\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \dots\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^4*(b*cos(d*x+c))^(1/2), x)

```
[Out] -2/45*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(160*cos(
1/2*d*x+1/2*c)^11-480*cos(1/2*d*x+1/2*c)^9+616*cos(1/2*d*x+1/2*c)^7-432*cos
(1/2*d*x+1/2*c)^5+160*cos(1/2*d*x+1/2*c)^3-21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-24*
cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2
)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^4, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^4 \sqrt{b \cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^4*(b*cos(c + d*x))^(1/2),x)
```

```
[Out] int(cos(c + d*x)^4*(b*cos(c + d*x))^(1/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**4*(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

3.68 $\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal. Leaf size=95

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{5/2}}{7b^2d} + \frac{10 \sin(c + dx)\sqrt{b \cos(c + dx)}}{21d} + \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b^2/d+10/21*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+10/21*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{5/2}}{7b^2d} + \frac{10 \sin(c + dx)\sqrt{b \cos(c + dx)}}{21d} + \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^3*Sqrt[b*Cos[c + d*x]], x]

[Out] $(10*b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (10*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(21*d) + (2*(b*\text{Cos}[c + d*x])^{(5/2)}*\text{Sin}[c + d*x])/(7*b^2*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos^3(c+dx)\sqrt{b\cos(c+dx)} dx &= \frac{\int (b\cos(c+dx))^{7/2} dx}{b^3} \\
&= \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^2d} + \frac{5 \int (b\cos(c+dx))^{3/2} dx}{7b} \\
&= \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21d} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^2d} + \frac{1}{21} (5b) \\
&= \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21d} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^2d} + \frac{(5b\sqrt{c})}{21} \\
&= \frac{10b\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21d\sqrt{b\cos(c+dx)}} + \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21d} + \frac{2(b)}{21}
\end{aligned}$$

Mathematica [A] time = 0.14, size = 73, normalized size = 0.77

$$\frac{\sqrt{b\cos(c+dx)} \left(20F\left(\frac{1}{2}(c+dx) \middle| 2\right) + (23\sin(c+dx) + 3\sin(3(c+dx)))\sqrt{\cos(c+dx)} \right)}{42d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3*Sqrt[b*Cos[c + d*x]], x]

[Out] (Sqrt[b*Cos[c + d*x]]*(20*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(2*3*Sin[c + d*x] + 3*Sin[3*(c + d*x)])))/(42*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b\cos(dx+c)} \cos(dx+c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b\cos(dx+c)} \cos(dx+c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^3, x)

maple [A] time = 0.12, size = 208, normalized size = 2.19

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b\left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 64\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 16\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^3*(b*cos(d*x+c))^(1/2), x)

```
[Out] -2/21*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^3, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^3 \sqrt{b \cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^3*(b*cos(c + d*x))^(1/2),x)
```

```
[Out] int(cos(c + d*x)^3*(b*cos(c + d*x))^(1/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**3*(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

3.69 $\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal. Leaf size=69

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{3/2}}{5bd} + \frac{6E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[Out] $2/5*(b*\cos(d*x+c))^{(3/2)*\sin(d*x+c)/b/d+6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{3/2}}{5bd} + \frac{6E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2*Sqrt[b*Cos[c + d*x]],x]

[Out] $(6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*(b*\text{Cos}[c + d*x])^{(3/2)*\text{Sin}[c + d*x]})/(5*b*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[b*SIN[c + d*x]]/Sqrt[SIN[c + d*x]], Int[Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx &= \frac{\int (b \cos(c + dx))^{5/2} dx}{b^2} \\
&= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5bd} + \frac{3}{5} \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5bd} + \frac{(3\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\
&= \frac{6\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5bd}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 62, normalized size = 0.90

$$\frac{\sqrt{b \cos(c + dx)} \left(6E\left(\frac{1}{2}(c + dx) \middle| 2\right) + \sin(2(c + dx))\sqrt{\cos(c + dx)} \right)}{5d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*Sqrt[b*Cos[c + d*x]], x]

[Out] (Sqrt[b*Cos[c + d*x]]*(6*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[2*(c + d*x)]))/(5*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^2, x)

maple [B] time = 0.11, size = 211, normalized size = 3.06

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b\left(-8\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2*(b*cos(d*x+c))^(1/2), x)

[Out] -2/5*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(-8*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+8*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(

$$\cos(1/2*d*x+1/2*c), 2^{(1/2)})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^2 \sqrt{b \cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(1/2), x)

[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(1/2), x)

[Out] Timed out

3.70 $\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal. Leaf size=67

$$\frac{2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{3d} + \frac{2b \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d \sqrt{b \cos(c + dx)}}$$

[Out] $2/3*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+2/3*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.04, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{3d} + \frac{2b \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]*Sqrt[b*Cos[c + d*x]], x]

[Out] $(2*b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(3*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos(c+dx)\sqrt{b\cos(c+dx)} dx &= \frac{\int (b\cos(c+dx))^{3/2} dx}{b} \\
&= \frac{2\sqrt{b\cos(c+dx)} \sin(c+dx)}{3d} + \frac{1}{3}b \int \frac{1}{\sqrt{b\cos(c+dx)}} dx \\
&= \frac{2\sqrt{b\cos(c+dx)} \sin(c+dx)}{3d} + \frac{(b\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3\sqrt{b\cos(c+dx)}} \\
&= \frac{2b\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3d\sqrt{b\cos(c+dx)}} + \frac{2\sqrt{b\cos(c+dx)} \sin(c+dx)}{3d}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 61, normalized size = 0.91

$$\frac{2(b\cos(c+dx))^{3/2} \left(F\left(\frac{1}{2}(c+dx) \middle| 2\right) + \sin(c+dx)\sqrt{\cos(c+dx)} \right)}{3bd\cos^2(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*Sqrt[b*Cos[c + d*x]],x]

[Out] (2*(b*Cos[c + d*x])^(3/2)*(EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[c + d*x]))/(3*b*d*Cos[c + d*x]^(3/2))

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b\cos(dx+c)}\cos(dx+c),x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b\cos(dx+c)}\cos(dx+c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c), x)

maple [B] time = 0.11, size = 188, normalized size = 2.81

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}b\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)*(b*cos(d*x+c))^(1/2),x)

[Out] -2/3*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(4*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*

$x+1/2*c)^{2-1})^{1/2}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{1/2})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2)^{1/2}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^{2-1}))^{1/2}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)*(b*cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(c + dx)} \cos(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))*cos(c + d*x), x)

3.71 $\int \sqrt{b \cos(c + dx)} dx$

Optimal. Leaf size=38

$$\frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2640, 2639}

$$\frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]], x]

[Out] $(2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \sqrt{b \cos(c + dx)} dx &= \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 38, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]], x]

[Out] $(2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]])$

fricas [F] time = 0.83, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c)), x)

maple [B] time = 0.09, size = 142, normalized size = 3.74

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \operatorname{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right),}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/2),x)

[Out] 2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \sqrt{b \cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/2),x)

[Out] int((b*cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*cos(c + d*x)), x)

3.72 $\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx$

Optimal. Leaf size=39

$$\frac{2b\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{d\sqrt{b\cos(c+dx)}}$$

[Out] $2*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 2642, 2641}

$$\frac{2b\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x],x]

[Out] $(2*b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \sqrt{b \cos(c + dx)} \sec(c + dx) dx &= b \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\ &= \frac{(b\sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{\sqrt{b \cos(c + dx)}} \\ &= \frac{2b\sqrt{\cos(c + dx)}F\left(\frac{1}{2}(c + dx)\middle|2\right)}{d\sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 39, normalized size = 1.00

$$\frac{2b\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x],x]

[Out] (2*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.48, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c), x)

maple [B] time = 0.10, size = 142, normalized size = 3.64

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \text{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)*(b*cos(d*x+c))^(1/2),x)

[Out] -2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x),x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)*(b*cos(d*x+c))**(1/2), x)
```

```
[Out] Integral(sqrt(b*cos(c + d*x))*sec(c + d*x), x)
```

3.73 $\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx$

Optimal. Leaf size=63

$$\frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[Out] $2*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^2,x]

[Out] $(-2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx &= b^2 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\
&= -\frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)}} + \frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 48, normalized size = 0.76

$$\frac{2b \left(\sin(c + dx) - \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^2, x]

[Out] (2*b*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.49, size = 0, normalized size = 0.00

$$\text{integral}(\sqrt{b \cos(dx + c)} \sec(dx + c)^2, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^2, x)

maple [A] time = 0.14, size = 166, normalized size = 2.63

$$\frac{2b \sqrt{-2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) b + \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) b \left(\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticE} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{\sqrt{-b \left(2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \right) \sin \left(\frac{dx}{2} + \frac{c}{2} \right) \sqrt{b \left(2 \left(\cos^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2*(b*cos(d*x+c))^(1/2), x)

[Out] -2*b*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)*((sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+

$\frac{1}{2}c)^4 - \sin(\frac{1}{2}d*x + \frac{1}{2}c)^2)^{(1/2)} / \sin(\frac{1}{2}d*x + \frac{1}{2}c) / (b * (2 * \cos(\frac{1}{2}d*x + \frac{1}{2}c)^2 - 1))^{(1/2)} / d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^2,x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**2*(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))*sec(c + d*x)**2, x)

3.74 $\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx$

Optimal. Leaf size=70

$$\frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}}$$

[Out] $2/3*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^(3/2)+2/3*b*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^(1/2))*\cos(d*x+c)^(1/2)/d/(b*\cos(d*x+c))^(1/2)$

Rubi [A] time = 0.06, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^3,x]

[Out] $(2*b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b^2*\text{Sin}[c + d*x])/(3*d*(b*\text{Cos}[c + d*x])^(3/2))$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx &= b^3 \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
&= \frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{1}{3} b \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{(b \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3 \sqrt{b \cos(c + dx)}} \\
&= \frac{2b \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 49, normalized size = 0.70

$$\frac{2b \left(\tan(c + dx) + \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^3,x]

[Out] (2*b*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}(\sqrt{b \cos(dx + c)} \sec(dx + c)^3, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^3, x)

maple [B] time = 0.14, size = 239, normalized size = 3.41

$$\frac{2 \left(-2 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \right)}{3 \sqrt{-b \left(2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^3*(b*cos(d*x+c))^(1/2),x)

[Out] -2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2-1)^(1/2))

$2)^{(1/2)} * (2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) ^{(1/2)} * \text{EllipticF}(\cos(1/2 * d * x + 1/2 * c), 2 ^{(1/2)}) - 2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 * \cos(1/2 * d * x + 1/2 * c) * b * (b * (2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1) * \sin(1/2 * d * x + 1/2 * c) ^ 2) ^{(1/2)} / (-b * (2 * \sin(1/2 * d * x + 1/2 * c) ^ 4 - \sin(1/2 * d * x + 1/2 * c) ^ 2)) ^{(1/2)} / (2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1) / \sin(1/2 * d * x + 1/2 * c) / (b * (2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1)) ^{(1/2)} / d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**3*(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))*sec(c + d*x)**3, x)

3.75 $\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx$

Optimal. Leaf size=95

$$\frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{6E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[Out] $2/5*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{6E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^4,x]

[Out] $(-6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^3*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*b*\text{Sin}[c + d*x])/(5*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx &= b^4 \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
&= \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{1}{5} (3b^2) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{3}{5} \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{(3\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\
&= -\frac{6\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.22, size = 69, normalized size = 0.73

$$\frac{2 \sec^2(c + dx) \sqrt{b \cos(c + dx)} \left(\frac{3}{2} \sin(2(c + dx)) + \tan(c + dx) - 3 \cos^{\frac{3}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{5d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^4, x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^2*(-3*Cos[c + d*x]^(3/2)*EllipticE[(c + d*x)/2, 2] + (3*Sin[2*(c + d*x)]/2 + Tan[c + d*x]))/(5*d)

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}(\sqrt{b \cos(dx + c)} \sec(dx + c)^4, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4*(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4*(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^4, x)

maple [B] time = 0.20, size = 363, normalized size = 3.82

$$\frac{2\sqrt{b} \left(2 \left(\cos^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \left(12 \text{EllipticE} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \sqrt{\frac{1}{2} - \cos \left(\frac{dx}{2} + \frac{c}{2} \right)} \right)}{5d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^4*(b*cos(d*x+c))^(1/2), x)

```
[Out] 2/5*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)^3/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)*(12*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^4-24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^2+24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^4*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^4, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^4,x)
```

```
[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^4, x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**4*(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

3.76 $\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx$

Optimal. Leaf size=98

$$\frac{2b^4 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^2 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

[Out] $2/7*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b^4 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^2 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^5,x]

[Out] $(10*b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b^4*\text{Sin}[c + d*x])/(7*d*(b*\text{Cos}[c + d*x])^{(7/2)}) + (10*b^2*\text{Sin}[c + d*x])/(21*d*(b*\text{Cos}[c + d*x])^{(3/2)})$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sqrt{b \cos(c+dx)} \sec^5(c+dx) dx &= b^5 \int \frac{1}{(b \cos(c+dx))^{9/2}} dx \\
&= \frac{2b^4 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{1}{7} (5b^3) \int \frac{1}{(b \cos(c+dx))^{5/2}} dx \\
&= \frac{2b^4 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b^2 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{1}{21} (5b) \int \frac{1}{\sqrt{b \cos(c+dx)}} dx \\
&= \frac{2b^4 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b^2 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{(5b\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21\sqrt{b \cos(c+dx)}} \\
&= \frac{10b\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21d\sqrt{b \cos(c+dx)}} + \frac{2b^4 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b^2 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.18, size = 69, normalized size = 0.70

$$\frac{\sec^3(c+dx)\sqrt{b \cos(c+dx)} \left(5 \sin(2(c+dx)) + 6 \tan(c+dx) + 10 \cos^{\frac{5}{2}}(c+dx) F\left(\frac{1}{2}(c+dx) \middle| 2\right)\right)}{21d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^5,x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^3*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d)

fricas [F] time = 0.83, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx+c)} \sec(dx+c)^5, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^5, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx+c)} \sec(dx+c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^5, x)

maple [B] time = 0.16, size = 396, normalized size = 4.04

$$2 \left(-40 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 60 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^5*(b*cos(d*x+c))^(1/2),x)

```
[Out] -2/21*(-40*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^6+60*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-40*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-30*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+40*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-16*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))*b*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^3/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^5, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^5,x)
```

```
[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^5, x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**5*(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

3.77 $\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx$

Optimal. Leaf size=123

$$\frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{14E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[Out] $2/9*b^5*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-14/15*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] time = 0.10, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{14E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^6,x]`

[Out] `(-14*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*d*Sqrt[Cos[c + d*x]]) + (2*b^5*Sin[c + d*x])/(9*d*(b*Cos[c + d*x])^(9/2)) + (14*b^3*Sin[c + d*x])/(45*d*(b*Cos[c + d*x])^(5/2)) + (14*b*Sin[c + d*x])/(15*d*Sqrt[b*Cos[c + d*x]])`

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2636

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2639

`Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 2640

`Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx &= b^6 \int \frac{1}{(b \cos(c + dx))^{11/2}} dx \\
&= \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{1}{9} (7b^4) \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
&= \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{1}{15} (7b^2) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{7}{15} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{7\sqrt{b \cos(c + dx)}}{15d} \\
&= -\frac{14\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}}
\end{aligned}$$

Mathematica [A] time = 0.20, size = 79, normalized size = 0.64

$$\frac{\sec^5(c + dx)\sqrt{b \cos(c + dx)} \left(150 \sin(c + dx) + 91 \sin(3(c + dx)) + 21 \sin(5(c + dx)) - 336 \cos^2(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right)\right)}{360d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^6,x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^5*(-336*Cos[c + d*x]^(9/2)*EllipticE[(c + d*x)/2, 2] + 150*Sin[c + d*x] + 91*Sin[3*(c + d*x)] + 21*Sin[5*(c + d*x)])/(360*d)

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} \sec(dx + c)^6, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^6*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^6, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^6*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^6, x)

maple [B] time = 0.20, size = 412, normalized size = 3.35

$$2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} b \left(-\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{144b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} - \frac{7\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{180b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^6*(b*cos(d*x+c))^(1/2),x)`

[Out]
$$-2*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*(-1/144*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^5-7/180*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^3-14/15*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*(\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} \sec(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^6*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^6, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^6,x)`

[Out] `int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^6, x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**6*(b*cos(d*x+c))**(1/2),x)`

[Out] Timed out

3.78 $\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal. Leaf size=126

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{9/2}}{11b^3d} + \frac{30b^2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{77d\sqrt{b \cos(c + dx)}} + \frac{18 \sin(c + dx)(b \cos(c + dx))^{5/2}}{77bd} + \frac{30b \sin(c + dx)(b \cos(c + dx))^{3/2}}{11b^2d}$$

[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^3/d+30/77*b^2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+30/77*b*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d

Rubi [A] time = 0.09, antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{9/2}}{11b^3d} + \frac{30b^2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{77d\sqrt{b \cos(c + dx)}} + \frac{18 \sin(c + dx)(b \cos(c + dx))^{5/2}}{77bd} + \frac{30b \sin(c + dx)(b \cos(c + dx))^{3/2}}{11b^2d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^4*(b*Cos[c + d*x])^(3/2), x]

[Out] (30*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*d*Sqrt[b*Cos[c + d*x]]) + (30*b*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^3*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Sine[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sine[c + d*x]]/Sqrt[b*Sine[c + d*x]], Int[1/Sqrt[Sine[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos^4(c+dx)(b \cos(c+dx))^{3/2} dx &= \frac{\int (b \cos(c+dx))^{11/2} dx}{b^4} \\
&= \frac{2(b \cos(c+dx))^{9/2} \sin(c+dx)}{11b^3d} + \frac{9 \int (b \cos(c+dx))^{7/2} dx}{11b^2} \\
&= \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77bd} + \frac{2(b \cos(c+dx))^{9/2} \sin(c+dx)}{11b^3d} + \frac{45}{77} \int (b \cos(c+dx))^{3/2} dx \\
&= \frac{30b\sqrt{b \cos(c+dx)} \sin(c+dx)}{77d} + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77bd} + \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{77d} \\
&= \frac{30b\sqrt{b \cos(c+dx)} \sin(c+dx)}{77d} + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77bd} + \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{77d} \\
&= \frac{30b^2 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{77d\sqrt{b \cos(c+dx)}} + \frac{30b\sqrt{b \cos(c+dx)} \sin(c+dx)}{77d} + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77bd}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 83, normalized size = 0.66

$$\frac{(b \cos(c+dx))^{3/2} \left(240F\left(\frac{1}{2}(c+dx) \middle| 2\right) + (290 \sin(c+dx) + 57 \sin(3(c+dx)) + 7 \sin(5(c+dx)))\sqrt{\cos(c+dx)} \right)}{616d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^4*(b*Cos[c + d*x])^(3/2), x]

[Out] ((b*Cos[c + d*x])^(3/2)*(240*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(290*Sin[c + d*x] + 57*Sin[3*(c + d*x)] + 7*Sin[5*(c + d*x)])))/(616*d*Cos[c + d*x]^(3/2))

fricas [F] time = 0.46, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx+c)} b \cos(dx+c)^5, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)^5, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx+c))^{\frac{3}{2}} \cos(dx+c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^4, x)

maple [A] time = 0.12, size = 236, normalized size = 1.87

$$\frac{2\sqrt{b} \left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) b^2 \left(448 \left(\cos^{13}\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1568 \left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 2384 \left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1120 \left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 224 \left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 16 \left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 2 \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right)}{77\sqrt{-b} \left(2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^4*(b*cos(d*x+c))^(3/2),x)`

[Out]
$$-2/77*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b^2*(448*\cos(1/2*d*x+1/2*c)^{13}-1568*\cos(1/2*d*x+1/2*c)^{11}+2384*\cos(1/2*d*x+1/2*c)^9-2040*\cos(1/2*d*x+1/2*c)^7+1084*\cos(1/2*d*x+1/2*c)^5-370*\cos(1/2*d*x+1/2*c)^3+15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+62*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^4, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^4 (b \cos(c + dx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^4*(b*cos(c + d*x))^(3/2),x)`

[Out] `int(cos(c + d*x)^4*(b*cos(c + d*x))^(3/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**4*(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

3.79 $\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal. Leaf size=95

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{7/2}}{9b^2d} + \frac{14 \sin(c + dx)(b \cos(c + dx))^{3/2}}{45d} + \frac{14bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[Out] $14/45*(b*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/d+2/9*(b*\cos(d*x+c))^{(7/2)}*\sin(d*x+c)/b^2/d+14/15*b*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.06, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{7/2}}{9b^2d} + \frac{14 \sin(c + dx)(b \cos(c + dx))^{3/2}}{45d} + \frac{14bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^3*(b*Cos[c + d*x])^(3/2), x]`

[Out] $(14*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (14*(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(45*d) + (2*(b*\text{Cos}[c + d*x])^{(7/2)}*\text{Sin}[c + d*x])/(9*b^2*d)$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2635

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2639

`Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 2640

`Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*SIN[c + d*x]]/Sqrt[SIN[c + d*x]], Int[Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^3} \\
&= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^2d} + \frac{7 \int (b \cos(c + dx))^{5/2} dx}{9b} \\
&= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^2d} + \frac{1}{15} \int (b \cos(c + dx))^{3/2} dx \\
&= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^2d} + \frac{(7b)}{15} \int (b \cos(c + dx))^{3/2} dx \\
&= \frac{14b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \dots
\end{aligned}$$

Mathematica [A] time = 0.10, size = 75, normalized size = 0.79

$$\frac{(b \cos(c + dx))^{3/2} \left(168E\left(\frac{1}{2}(c + dx) \middle| 2\right) + (38 \sin(2(c + dx)) + 5 \sin(4(c + dx)))\sqrt{\cos(c + dx)}\right)}{180d \cos^{\frac{3}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3*(b*Cos[c + d*x])^(3/2), x]

[Out] ((b*Cos[c + d*x])^(3/2)*(168*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)])))/(180*d*Cos[c + d*x]^(3/2))

fricas [F] time = 1.05, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^3, x)

maple [B] time = 0.12, size = 223, normalized size = 2.35

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \left(160\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 480\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 616\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 320\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 160\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 40\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{45\sqrt{-b}\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 2\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^3*(b*cos(d*x+c))^(3/2), x)

```
[Out] -2/45*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(160*cos(1/2*d*x+1/2*c)^11-480*cos(1/2*d*x+1/2*c)^9+616*cos(1/2*d*x+1/2*c)^7-432*cos(1/2*d*x+1/2*c)^5+160*cos(1/2*d*x+1/2*c)^3-21*(sin(1/2*d*x+1/2*c)^2)^(1/2))*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-24*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^3, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^3 (b \cos(c + dx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^3*(b*cos(c + d*x))^(3/2),x)
```

```
[Out] int(cos(c + d*x)^3*(b*cos(c + d*x))^(3/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**3*(b*cos(d*x+c))**(3/2),x)
```

```
[Out] Timed out
```

3.80 $\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal. Leaf size=98

$$\frac{10b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{21d\sqrt{b\cos(c+dx)}} + \frac{2\sin(c+dx)(b\cos(c+dx))^{5/2}}{7bd} + \frac{10b\sin(c+dx)\sqrt{b\cos(c+dx)}}{21d}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)*\sin(d*x+c)/b/d+10/21*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)+10/21*b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{10b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{21d\sqrt{b\cos(c+dx)}} + \frac{2\sin(c+dx)(b\cos(c+dx))^{5/2}}{7bd} + \frac{10b\sin(c+dx)\sqrt{b\cos(c+dx)}}{21d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2*(b*Cos[c + d*x])^(3/2),x]

[Out] $(10*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (10*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(21*d) + (2*(b*\text{Cos}[c + d*x])^{(5/2)*\text{Sin}[c + d*x]})/(7*b*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*SIN[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx &= \frac{\int (b \cos(c + dx))^{7/2} dx}{b^2} \\
&= \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd} + \frac{5}{7} \int (b \cos(c + dx))^{3/2} dx \\
&= \frac{10b\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd} + \frac{1}{21} (5b^2 \sqrt{b \cos(c + dx)}) \\
&= \frac{10b\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd} + \frac{(5b^2 \sqrt{b \cos(c + dx)})}{21} \\
&= \frac{10b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 73, normalized size = 0.74

$$\frac{(b \cos(c + dx))^{3/2} \left(20F\left(\frac{1}{2}(c + dx) \middle| 2\right) + (23 \sin(c + dx) + 3 \sin(3(c + dx))) \sqrt{\cos(c + dx)} \right)}{42d \cos^{\frac{3}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(3/2), x]

[Out] ((b*Cos[c + d*x])^(3/2)*(20*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(23*Sin[c + d*x] + 3*Sin[3*(c + d*x)])))/(42*d*Cos[c + d*x]^(3/2))

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^2, x)

maple [A] time = 0.12, size = 210, normalized size = 2.14

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2*(b*cos(d*x+c))^(3/2), x)


```
[Out] -2/21*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^2, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^2 (b \cos(c + dx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(3/2),x)
```

```
[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(3/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(3/2),x)
```

```
[Out] Timed out
```

3.81 $\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal. Leaf size=67

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{3/2}}{5d} + \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[Out] $2/5*(b*\cos(d*x+c))^{3/2}*\sin(d*x+c)/d+6/5*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{3/2}}{5d} + \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]*(b*Cos[c + d*x])^(3/2), x]

[Out] $(6*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*(b*\text{Cos}[c + d*x])^{3/2}*\text{Sin}[c + d*x])/(5*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Sin[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos(c+dx)(b \cos(c+dx))^{3/2} dx &= \frac{\int (b \cos(c+dx))^{5/2} dx}{b} \\
&= \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5d} + \frac{1}{5}(3b) \int \sqrt{b \cos(c+dx)} dx \\
&= \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5d} + \frac{(3b\sqrt{b \cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5\sqrt{\cos(c+dx)}} \\
&= \frac{6b\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5d\sqrt{\cos(c+dx)}} + \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5d}
\end{aligned}$$

Mathematica [A] time = 0.08, size = 65, normalized size = 0.97

$$\frac{(b \cos(c+dx))^{5/2} \left(6E\left(\frac{1}{2}(c+dx) \middle| 2\right) + \sin(2(c+dx))\sqrt{\cos(c+dx)}\right)}{5bd \cos^2(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(3/2), x]

[Out] ((b*Cos[c + d*x])^(5/2)*(6*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[2*(c + d*x)]))/(5*b*d*Cos[c + d*x]^(5/2))

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx+c)} b \cos(dx+c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx+c))^{\frac{3}{2}} \cos(dx+c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c), x)

maple [B] time = 0.12, size = 213, normalized size = 3.18

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \left(-8 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)*(b*cos(d*x+c))^(3/2), x)

[Out] -2/5*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(-8*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+8*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c))-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Elliptic

$E(\cos(1/2*d*x+1/2*c), 2^{(1/2)}) - 2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c) / (-b*(2*\sin(1/2*d*x+1/2*c)^4 - \sin(1/2*d*x+1/2*c)^2))^{(1/2)} / \sin(1/2*d*x+1/2*c) / (b*(2*\cos(1/2*d*x+1/2*c)^2 - 1))^{(1/2)} / d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx) (b \cos(c + dx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)*(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

3.82 $\int (b \cos(c + dx))^{3/2} dx$

Optimal. Leaf size=70

$$\frac{2b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{3d\sqrt{b\cos(c+dx)}} + \frac{2b\sin(c+dx)\sqrt{b\cos(c+dx)}}{3d}$$

[Out] $2/3*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+2/3*b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 2642, 2641}

$$\frac{2b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{3d\sqrt{b\cos(c+dx)}} + \frac{2b\sin(c+dx)\sqrt{b\cos(c+dx)}}{3d}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(3/2),x]

[Out] $(2*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(3*d)$

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*cos[c + d*x]*(b*sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{3/2} dx &= \frac{2b\sqrt{b\cos(c+dx)}\sin(c+dx)}{3d} + \frac{1}{3}b^2 \int \frac{1}{\sqrt{b\cos(c+dx)}} dx \\ &= \frac{2b\sqrt{b\cos(c+dx)}\sin(c+dx)}{3d} + \frac{(b^2\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3\sqrt{b\cos(c+dx)}} \\ &= \frac{2b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{3d\sqrt{b\cos(c+dx)}} + \frac{2b\sqrt{b\cos(c+dx)}\sin(c+dx)}{3d} \end{aligned}$$

Mathematica [A] time = 0.01, size = 58, normalized size = 0.83

$$\frac{2(b \cos(c + dx))^{3/2} \left(F\left(\frac{1}{2}(c + dx) \middle| 2\right) + \sin(c + dx) \sqrt{\cos(c + dx)} \right)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2),x]

[Out] (2*(b*Cos[c + d*x])^(3/2)*(EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[c + d*x]))/(3*d*Cos[c + d*x]^(3/2))

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.12, size = 190, normalized size = 2.71

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \left(4\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2),x)

[Out] -2/3*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(4*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (b \cos(c + dx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*cos(c + d*x))^(3/2), x)`

[Out] `int((b*cos(c + d*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*cos(d*x+c))**(3/2), x)`

[Out] `Integral((b*cos(c + d*x))**(3/2), x)`

3.83 $\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx$

Optimal. Leaf size=39

$$\frac{2bE\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

[Out] $2*b*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 2640, 2639}

$$\frac{2bE\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sec}[c + d*x], x]$

[Out] $(2*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /;$ FreeQ[{c, d}, x]

Rule 2640

$\text{Int}[\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[b*\text{Sin}[c + d*x]]/\text{Sqrt}[\text{Sin}[c + d*x]], \text{Int}[\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{3/2} \sec(c + dx) dx &= b \int \sqrt{b \cos(c + dx)} dx \\ &= \frac{(b\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{2b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx)\middle|2\right)}{d\sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 39, normalized size = 1.00

$$\frac{2bE\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(3/2)*Sec[c + d*x],x]

[Out] (2*b*Sqrt[b*cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.49, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c) \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c), x)

maple [B] time = 0.10, size = 144, normalized size = 3.69

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), 2\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)*sec(d*x+c),x)

[Out] 2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{(b \cos(c + dx))^{\frac{3}{2}}}{\cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x),x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c),x)

[Out] Timed out

3.84 $\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx$

Optimal. Leaf size=41

$$\frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[Out] $2*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2642, 2641}

$$\frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sec}[c + d*x]^2, x]$

[Out] $(2*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_)*(v_))^{(n_*)}, x_Symbol] :> \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] :> \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 2642

$\text{Int}[1/\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] :> \text{Dist}[\text{Sqrt}[\text{Sin}[c + d*x]]/\text{Sqrt}[b*\text{Sin}[c + d*x]], \text{Int}[1/\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /; \text{FreeQ}[\{b, c, d\}, x]$

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx &= b^2 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\ &= \frac{(b^2 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{\sqrt{b \cos(c + dx)}} \\ &= \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 41, normalized size = 1.00

$$\frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(3/2)*Sec[c + d*x]^2,x]

[Out] (2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*cos[c + d*x]])

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\int \sqrt{b \cos(dx + c)} b \cos(dx + c) \sec(dx + c)^2, x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^2, x)

maple [B] time = 0.12, size = 144, normalized size = 3.51

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \operatorname{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)*sec(d*x+c)^2,x)

[Out] -2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^2,x)

```
[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^2, x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**2, x)
```

```
[Out] Timed out
```

3.85 $\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx$

Optimal. Leaf size=66

$$\frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[Out] $2*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-2*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sec}[c + d*x]^3, x]$

[Out] $(-2*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^2*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] := \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2636

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] := \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /;$ $\text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_)]], x_Symbol] := \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /;$ $\text{FreeQ}\{c, d\}, x$

Rule 2640

$\text{Int}[\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_)]], x_Symbol] := \text{Dist}[\text{Sqrt}[b*\text{Sin}[c + d*x]]/\text{Sqrt}[\text{Sin}[c + d*x]], \text{Int}[\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ $\text{FreeQ}\{b, c, d\}, x$

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx &= b^3 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - b \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{(b\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\
&= -\frac{2b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 50, normalized size = 0.76

$$\frac{2b^2 \left(\sin(c + dx) - \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^3,x]

[Out] (2*b^2*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c) \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec^3(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^3, x)

maple [A] time = 0.13, size = 168, normalized size = 2.55

$$\frac{2b^2 \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) b + \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) b \left(\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1} \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \right)} \right)}{\sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b \left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1 \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)*sec(d*x+c)^3,x)

[Out] -2*b^2*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)*((sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*sin(1/2*d*x+1/2*c)*sqrt(b*(2*cos(1/2*d*x+1/2*c)^2-1)))

$x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^3, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**3,x)

[Out] Timed out

3.86 $\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx$

Optimal. Leaf size=72

$$\frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}}$$

[Out] $2/3*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(3/2)+2/3*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2))*\cos(d*x+c)^(1/2)/d/(b*\cos(d*x+c))^(1/2)$

Rubi [A] time = 0.05, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(3/2)*Sec[c + d*x]^4,x]

[Out] $(2*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b^3*\text{Sin}[c + d*x])/(3*d*(b*\text{Cos}[c + d*x])^(3/2))$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*SIN[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx &= b^4 \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
&= \frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{1}{3} b^2 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{(b^2 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
&= \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 51, normalized size = 0.71

$$\frac{2b^2 \left(\tan(c + dx) + \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^4,x]

[Out] (2*b^2*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c) \sec(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^4,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)*sec(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{3/2} \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^4,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^4, x)

maple [B] time = 0.13, size = 241, normalized size = 3.35

$$\frac{2 \left(-2 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \right)}{3 \sqrt{-b \left(2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)*sec(d*x+c)^4,x)

[Out] -2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2-1)^(1/2))

$2)^{(1/2)} * (2 * \sin(1/2 * d * x + 1/2 * c)^2 - 1)^{(1/2)} * \text{EllipticF}(\cos(1/2 * d * x + 1/2 * c), 2^{(1/2)}) - 2 * \sin(1/2 * d * x + 1/2 * c)^2 * \cos(1/2 * d * x + 1/2 * c) * b^2 * (b * (2 * \cos(1/2 * d * x + 1/2 * c)^2 - 1) * \sin(1/2 * d * x + 1/2 * c)^2)^{(1/2)} / (-b * (2 * \sin(1/2 * d * x + 1/2 * c)^4 - \sin(1/2 * d * x + 1/2 * c)^2))^{(1/2)} / (2 * \cos(1/2 * d * x + 1/2 * c)^2 - 1) / \sin(1/2 * d * x + 1/2 * c) / (b * (2 * \cos(1/2 * d * x + 1/2 * c)^2 - 1))^{(1/2)} / d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^4,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^4,x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^4, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**4,x)

[Out] Timed out

3.87 $\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx$

Optimal. Leaf size=98

$$\frac{2b^4 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^2 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[Out] $2/5*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-6/5*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^4 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^2 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sec}[c + d*x]^5, x]$

[Out] $(-6*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^4*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*b^2*\text{Sin}[c + d*x])/(5*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /;$ FreeQ[{c, d}, x]

Rule 2640

$\text{Int}[\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[b*\text{Sin}[c + d*x]]/\text{Sqrt}[\text{Sin}[c + d*x]], \text{Int}[\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx &= b^5 \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
&= \frac{2b^4 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{1}{5} (3b^3) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^4 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^2 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{1}{5} (3b) \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b^4 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^2 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{(3b\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\
&= -\frac{6b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2b^4 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^2 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.11, size = 69, normalized size = 0.70

$$\frac{\sec^4(c + dx)(b \cos(c + dx))^{3/2} \left(7 \sin(c + dx) + 3 \sin(3(c + dx)) - 12 \cos^{\frac{5}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right)\right)}{10d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^5,x]

[Out] ((b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^4*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)]))/(10*d)

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c) \sec(dx + c)^5, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^5,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)*sec(d*x + c)^5, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^5,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^5, x)

maple [B] time = 0.18, size = 364, normalized size = 3.71

$$2\sqrt{b \left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b \left(12 \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{\frac{1}{2}} - \dots\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)*sec(d*x+c)^5,x)

```
[Out] 2/5*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b/sin(1/2*d*x+1/2*c)^3/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)*(12*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^4-24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^2+24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^5,x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^5, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{\frac{3}{2}}}{\cos(c + dx)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^5,x)
```

```
[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^5, x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**5,x)
```

```
[Out] Timed out
```

3.88 $\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx$

Optimal. Leaf size=100

$$\frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

[Out] $2/7*b^5*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sec}[c + d*x]^6, x]$

[Out] $(10*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b^5*\text{Sin}[c + d*x])/(7*d*(b*\text{Cos}[c + d*x])^{(7/2)}) + (10*b^3*\text{Sin}[c + d*x])/(21*d*(b*\text{Cos}[c + d*x])^{(3/2)})$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /;$ FreeQ[{c, d}, x]

Rule 2642

$\text{Int}[1/\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[c + d*x]]/\text{Sqrt}[b*\text{Sin}[c + d*x]], \text{Int}[1/\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx &= b^6 \int \frac{1}{(b \cos(c + dx))^{9/2}} dx \\
&= \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{1}{7} (5b^4) \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
&= \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{1}{21} (5b^2) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{(5b^2 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21 \sqrt{b \cos(c + dx)}} \\
&= \frac{10b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.12, size = 69, normalized size = 0.69

$$\frac{\sec^4(c + dx)(b \cos(c + dx))^{3/2} \left(5 \sin(2(c + dx)) + 6 \tan(c + dx) + 10 \cos^{\frac{5}{2}}(c + dx) F\left(\frac{1}{2}(c + dx) \middle| 2\right)\right)}{21d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^6,x]

[Out] ((b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^4*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d)

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c) \sec(dx + c)^6, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^6,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)*sec(d*x + c)^6, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^6,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^6, x)

maple [B] time = 0.15, size = 398, normalized size = 3.98

$$\frac{2 \left(-40 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 60 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right)}{21d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)*sec(d*x+c)^6,x)


```
[Out] -2/21*(-40*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^6+60*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-40*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-30*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+40*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-16*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))*b^2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^3/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^6,x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^6, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{\frac{3}{2}}}{\cos(c + dx)^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^6,x)
```

```
[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^6, x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**6,x)
```

```
[Out] Timed out
```

3.89 $\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx$

Optimal. Leaf size=126

$$\frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^2 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{14bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[Out] $2/9*b^6*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-14/15*b*(\cos(1/2*d*x+1/2*c))^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] time = 0.09, antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^2 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{14bE\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^(3/2)*\text{Sec}[c + d*x]^7, x]$

[Out] $(-14*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/((15*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^6*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^(9/2)) + (14*b^4*\text{Sin}[c + d*x])/(45*d*(b*\text{Cos}[c + d*x])^(5/2)) + (14*b^2*\text{Sin}[c + d*x])/(15*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]))$

Rule 16

$\text{Int}[(u_*)*(v_)^(m_*)*((b_)*(v_))^(n_), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^(m + n), x], x] /;$ $\text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2636

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + \text{Dist}[(n + 2)/(b^2*(n + 1)), \text{Int}[(b*\text{Sin}[c + d*x])^(n + 2), x], x] /;$ $\text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /;$ $\text{FreeQ}[\{c, d\}, x]$

Rule 2640

$\text{Int}[\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[b*\text{Sin}[c + d*x]]/\text{Sqrt}[\text{Sin}[c + d*x]], \text{Int}[\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ $\text{FreeQ}[\{b, c, d\}, x]$

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx &= b^7 \int \frac{1}{(b \cos(c + dx))^{11/2}} dx \\
&= \frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{1}{9} (7b^5) \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
&= \frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{1}{15} (7b^3) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^2 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{1}{15} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^2 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{7}{15} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= -\frac{14b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}}
\end{aligned}$$

Mathematica [A] time = 0.17, size = 79, normalized size = 0.63

$$\frac{\sec^6(c + dx)(b \cos(c + dx))^{3/2} \left(150 \sin(c + dx) + 91 \sin(3(c + dx)) + 21 \sin(5(c + dx)) - 336 \cos^{\frac{9}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right)\right)}{360d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^7, x]

[Out] ((b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^6*(-336*Cos[c + d*x]^(9/2)*EllipticE[(c + d*x)/2, 2] + 150*Sin[c + d*x] + 91*Sin[3*(c + d*x)] + 21*Sin[5*(c + d*x)]))/ (360*d)

fricas [F] time = 0.48, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b \cos(dx + c) \sec(dx + c)^7, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^7,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b*cos(d*x + c)*sec(d*x + c)^7, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^7 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^7,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^7, x)

maple [B] time = 0.20, size = 414, normalized size = 3.29

$$2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \left(-\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{144b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} - \frac{7\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{180b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*cos(d*x+c))^(3/2)*sec(d*x+c)^7,x)`

[Out]
$$-2*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b^2*(-1/144*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^5-7/180*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^3-14/15*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*E\text{llipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*(E\text{llipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-E\text{llipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^7 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^7,x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^7, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^7} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^7,x)`

[Out] `int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^7, x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**7,x)`

[Out] Timed out

3.90 $\int \cos^3(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal. Leaf size=125

$$\frac{30b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{77d \sqrt{b \cos(c + dx)}} + \frac{2 \sin(c + dx)(b \cos(c + dx))^{9/2}}{11b^2 d} + \frac{30b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{77d} + \frac{18 \sin(c + dx)}{77d}$$

[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^2/d+30/77*b^3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+30/77*b^2*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d

Rubi [A] time = 0.08, antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c + dx)(b \cos(c + dx))^{9/2}}{11b^2 d} + \frac{30b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{77d} + \frac{30b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{77d \sqrt{b \cos(c + dx)}} + \frac{18 \sin(c + dx)}{77d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^3*(b*Cos[c + d*x])^(5/2), x]

[Out] (30*b^3*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*d*Sqrt[b*Cos[c + d*x]]) + (30*b^2*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^2*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Ssin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Ssin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Ssin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos^3(c + dx)(b \cos(c + dx))^{5/2} dx &= \frac{\int (b \cos(c + dx))^{11/2} dx}{b^3} \\
&= \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^2d} + \frac{9 \int (b \cos(c + dx))^{7/2} dx}{11b} \\
&= \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^2d} + \frac{1}{77}(45b \cos(c + dx))^{3/2} \sin(c + dx) \\
&= \frac{30b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77d} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{77d} \\
&= \frac{30b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77d} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{77d} \\
&= \frac{30b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{77d \sqrt{b \cos(c + dx)}} + \frac{30b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77d} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{77d}
\end{aligned}$$

Mathematica [A] time = 0.13, size = 83, normalized size = 0.66

$$\frac{(b \cos(c + dx))^{5/2} \left(240F\left(\frac{1}{2}(c + dx) \middle| 2\right) + (290 \sin(c + dx) + 57 \sin(3(c + dx)) + 7 \sin(5(c + dx))) \sqrt{\cos(c + dx)} \right)}{616d \cos^2(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3*(b*Cos[c + d*x])^(5/2), x]

[Out] ((b*Cos[c + d*x])^(5/2)*(240*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(290*Sin[c + d*x] + 57*Sin[3*(c + d*x)] + 7*Sin[5*(c + d*x)])))/(616*d*Cos[c + d*x]^(5/2))

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^5, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^5, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{5/2} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c)^3, x)

maple [A] time = 0.14, size = 236, normalized size = 1.89

$$\frac{2\sqrt{b} \left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) b^3 \left(448 \left(\cos^{13}\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1568 \left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 2384 \left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1120 \left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 224 \left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 16 \left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right) + 77 \sqrt{-b} \left(2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right)}{77 \sqrt{-b} \left(2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^3*(b*cos(d*x+c))^(5/2),x)`

[Out]
$$-2/77*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b^3*(448*\cos(1/2*d*x+1/2*c)^{13}-1568*\cos(1/2*d*x+1/2*c)^{11}+2384*\cos(1/2*d*x+1/2*c)^9-2040*\cos(1/2*d*x+1/2*c)^7+1084*\cos(1/2*d*x+1/2*c)^5-370*\cos(1/2*d*x+1/2*c)^3+15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+62*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c)^3, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^3 (b \cos(c + dx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^3*(b*cos(c + d*x))^(5/2),x)`

[Out] `int(cos(c + d*x)^3*(b*cos(c + d*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**3*(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

3.91 $\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal. Leaf size=98

$$\frac{14b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d \sqrt{\cos(c + dx)}} + \frac{2 \sin(c + dx)(b \cos(c + dx))^{7/2}}{9bd} + \frac{14b \sin(c + dx)(b \cos(c + dx))^{3/2}}{45d}$$

[Out] $14/45*b*(b*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/d+2/9*(b*\cos(d*x+c))^{(7/2)}*\sin(d*x+c)/b/d+14/15*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{14b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d \sqrt{\cos(c + dx)}} + \frac{2 \sin(c + dx)(b \cos(c + dx))^{7/2}}{9bd} + \frac{14b \sin(c + dx)(b \cos(c + dx))^{3/2}}{45d}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^2*(b*Cos[c + d*x])^(5/2), x]`

[Out] $(14*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (14*b*(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(45*d) + (2*(b*\text{Cos}[c + d*x])^{(7/2)}*\text{Sin}[c + d*x])/(9*b*d)$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2635

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2639

`Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 2640

`Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*SIN[c + d*x]]/Sqrt[SIN[c + d*x]], Int[Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^2} \\
&= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9bd} + \frac{7}{9} \int (b \cos(c + dx))^{5/2} dx \\
&= \frac{14b(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9bd} + \frac{1}{15} \int (b \cos(c + dx))^{1/2} dx \\
&= \frac{14b(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9bd} + \frac{7}{15} \int (b \cos(c + dx))^{1/2} dx \\
&= \frac{14b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d \sqrt{\cos(c + dx)}} + \frac{14b(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 75, normalized size = 0.77

$$\frac{(b \cos(c + dx))^{5/2} \left(168 E\left(\frac{1}{2}(c + dx) \middle| 2\right) + (38 \sin(2(c + dx)) + 5 \sin(4(c + dx))) \sqrt{\cos(c + dx)}\right)}{180d \cos^2(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(5/2), x]

[Out] ((b*Cos[c + d*x])^(5/2)*(168*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)])))/(180*d*Cos[c + d*x]^(5/2))

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{5/2} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c)^2, x)

maple [B] time = 0.13, size = 223, normalized size = 2.28

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^3 \left(160\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 480\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 616\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 448\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 224\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 56\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{45\sqrt{-b}\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2*(b*cos(d*x+c))^(5/2), x)

[Out]
$$-2/45*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b^3*(160*\cos(1/2*d*x+1/2*c)^{11}-480*\cos(1/2*d*x+1/2*c)^9+616*\cos(1/2*d*x+1/2*c)^7-432*\cos(1/2*d*x+1/2*c)^5+160*\cos(1/2*d*x+1/2*c)^3-21*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-24*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^2 (b \cos(c + dx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2*(b*cos(c + d*x))^(5/2),x)`

[Out] `int(cos(c + d*x)^2*(b*cos(c + d*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

3.92 $\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal. Leaf size=97

$$\frac{10b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{21d} + \frac{2 \sin(c + dx) (b \cos(c + dx))^{5/2}}{7d}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)*\sin(d*x+c)/d+10/21*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)+10/21*b^2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.06, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 2635, 2642, 2641}

$$\frac{10b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{21d} + \frac{10b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2 \sin(c + dx) (b \cos(c + dx))^{5/2}}{7d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]*(b*Cos[c + d*x])^(5/2), x]

[Out] $(10*b^3*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (10*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(21*d) + (2*(b*\text{Cos}[c + d*x])^{(5/2)*\text{Sin}[c + d*x]})/(7*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx &= \frac{\int (b \cos(c + dx))^{7/2} dx}{b} \\
&= \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} + \frac{1}{7}(5b) \int (b \cos(c + dx))^{3/2} dx \\
&= \frac{10b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} + \frac{1}{21} (5b^3 \sqrt{b \cos(c + dx)}) \\
&= \frac{10b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} + \frac{(5b^3 \sqrt{b \cos(c + dx)})}{21} \\
&= \frac{10b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 76, normalized size = 0.78

$$\frac{b^2 \sqrt{b \cos(c + dx)} \left(20F\left(\frac{1}{2}(c + dx) \middle| 2\right) + (23 \sin(c + dx) + 3 \sin(3(c + dx))) \sqrt{\cos(c + dx)} \right)}{42d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(5/2), x]

[Out] (b^2*Sqrt[b*Cos[c + d*x]]*(20*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(23*Sin[c + d*x] + 3*Sin[3*(c + d*x)])))/(42*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^2 \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c), x)

maple [A] time = 0.10, size = 210, normalized size = 2.16

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^3 \left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 64\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 32\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)*(b*cos(d*x+c))^(5/2), x)

```
[Out] -2/21*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx) (b \cos(c + dx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)*(b*cos(c + d*x))^(5/2),x)
```

```
[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(5/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

3.93 $\int (b \cos(c + dx))^{5/2} dx$

Optimal. Leaf size=70

$$\frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d \sqrt{\cos(c + dx)}} + \frac{2b \sin(c + dx) (b \cos(c + dx))^{3/2}}{5d}$$

[Out] $2/5*b*(b*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/d+6/5*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 2640, 2639}

$$\frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d \sqrt{\cos(c + dx)}} + \frac{2b \sin(c + dx) (b \cos(c + dx))^{3/2}}{5d}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(5/2), x]

[Out] $(6*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b*(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(5*d)$

Rule 2635

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*cos[c + d*x])*(b*sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[b*sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{5/2} dx &= \frac{2b(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d} + \frac{1}{5} (3b^2) \int \sqrt{b \cos(c + dx)} dx \\ &= \frac{2b(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d} + \frac{(3b^2 \sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{5 \sqrt{\cos(c + dx)}} \\ &= \frac{6b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d \sqrt{\cos(c + dx)}} + \frac{2b(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d} \end{aligned}$$

Mathematica [A] time = 0.01, size = 62, normalized size = 0.89

$$\frac{(b \cos(c + dx))^{5/2} \left(6E\left(\frac{1}{2}(c + dx) \middle| 2\right) + \sin(2(c + dx)) \sqrt{\cos(c + dx)}\right)}{5d \cos^{5/2}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(5/2), x]

[Out] ((b*cos[c + d*x])^(5/2)*(6*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[2*(c + d*x)]))/(5*d*cos[c + d*x]^(5/2))

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.13, size = 213, normalized size = 3.04

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^3 \left(-8\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}\right)}{5\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2), x)

[Out] -2/5*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(-8*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+8*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (b \cos(c + dx))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(5/2),x)
```

```
[Out] int((b*cos(c + d*x))^(5/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```


3.94 $\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx$

Optimal. Leaf size=72

$$\frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{3d}$$

[Out] $2/3*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+2/3*b^2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.05, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{3d} + \frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(5/2)*Sec[c + d*x],x]

[Out] $(2*b^3*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(3*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*cos[c + d*x]*(b*sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx &= b \int (b \cos(c + dx))^{3/2} dx \\
&= \frac{2b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d} + \frac{1}{3} b^3 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d} + \frac{(b^3 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
&= \frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 59, normalized size = 0.82

$$\frac{2b(b \cos(c + dx))^{3/2} \left(F\left(\frac{1}{2}(c + dx) \middle| 2\right) + \sin(c + dx) \sqrt{\cos(c + dx)} \right)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x],x]

[Out] (2*b*(b*Cos[c + d*x])^(3/2)*(EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[c + d*x]))/(3*d*Cos[c + d*x]^(3/2))

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c), x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c), x)

maple [B] time = 0.14, size = 190, normalized size = 2.64

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^3 \left(4\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)*sec(d*x+c),x)

[Out] -2/3*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(4*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*

$d*x+1/2*c)^{2-1})^{1/2}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{1/2})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{1/2}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^{2-1}))^{1/2}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{\frac{5}{2}}}{\cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x),x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c),x)

[Out] Timed out

3.95 $\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx$

Optimal. Leaf size=41

$$\frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[Out] $2*b^2*(\cos(1/2*d*x+1/2*c))^{5/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{1/2})*(b*\cos(d*x+c))^{1/2}/d/\cos(d*x+c)^{1/2}$

Rubi [A] time = 0.04, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2640, 2639}

$$\frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{5/2}*\text{Sec}[c + d*x]^2, x]$

[Out] $(2*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /;$ $\text{FreeQ}\{c, d, x\}$

Rule 2640

$\text{Int}[\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[b*\text{Sin}[c + d*x]]/\text{Sqrt}[\text{Sin}[c + d*x]], \text{Int}[\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ $\text{FreeQ}\{b, c, d, x\}$

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx &= b^2 \int \sqrt{b \cos(c + dx)} dx \\ &= \frac{(b^2 \sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{2b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 41, normalized size = 1.00

$$\frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(5/2)*Sec[c + d*x]^2,x]

[Out] (2*b^2*Sqrt[b*cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^2, x)

maple [B] time = 0.10, size = 144, normalized size = 3.51

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^3 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)*sec(d*x+c)^2,x)

[Out] 2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^2,x)

```
[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^2, x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**2,x)
```

```
[Out] Timed out
```

3.96 $\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx$

Optimal. Leaf size=41

$$\frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[Out] $2*b^3*(\cos(1/2*d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2642, 2641}

$$\frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^3,x]

[Out] (2*b^3*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx &= b^3 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\ &= \frac{(b^3 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{\sqrt{b \cos(c + dx)}} \\ &= \frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 38, normalized size = 0.93

$$\frac{2F\left(\frac{1}{2}(c + dx) \middle| 2\right) (b \cos(c + dx))^{5/2}}{d \cos^{\frac{5}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(5/2)*Sec[c + d*x]^3,x]

[Out] (2*(b*cos[c + d*x])^(5/2)*EllipticF[(c + d*x)/2, 2])/(d*cos[c + d*x]^(5/2))

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^3, x)

maple [B] time = 0.13, size = 144, normalized size = 3.51

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^3 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \text{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)*sec(d*x+c)^3,x)

[Out] -2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{\frac{5}{2}}}{\cos(c + dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^3, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**3,x)

[Out] Timed out

3.97 $\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx$

Optimal. Leaf size=68

$$\frac{2b^3 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[Out] $2*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-2*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.06, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^3 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^4,x]

[Out] $(-2*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^3*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*SIN[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*SIN[c + d*x]]/Sqrt[SIN[c + d*x]], Int[Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx &= b^4 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^3 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - b^2 \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b^3 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{(b^2 \sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\
&= -\frac{2b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 50, normalized size = 0.74

$$\frac{2b^3 \left(\sin(c + dx) - \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^4,x]

[Out] (2*b^3*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.88, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^4,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^4,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^4, x)

maple [A] time = 0.13, size = 168, normalized size = 2.47

$$\frac{2b^3 \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b + \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b \left(\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right)}{\sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b \left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1 \right)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)*sec(d*x+c)^4,x)

[Out] -2*b^3*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)*((sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*sin(1/2*d*x+1/2*c)*sqrt(b*(2*cos(1/2*d*x+1/2*c)^2-1)))

$x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^4,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{\frac{5}{2}}}{\cos(c + dx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^4,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^4, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**4,x)

[Out] Timed out

3.98 $\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx$

Optimal. Leaf size=72

$$\frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}}$$

[Out] $2/3*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^(3/2)+2/3*b^3*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2))*\cos(d*x+c)^(1/2)/d/(b*\cos(d*x+c))^(1/2)$

Rubi [A] time = 0.06, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^(5/2)*\text{Sec}[c + d*x]^5, x]$

[Out] $(2*b^3*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b^4*\text{Sin}[c + d*x])/(3*d*(b*\text{Cos}[c + d*x])^(3/2))$

Rule 16

$\text{Int}[(u_*)*(v_)^(m_*)*((b_)*(v_))^(n_), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^(m+n), x], x] /;$ $\text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2636

$\text{Int}[(b_)*\sin[(c_*) + (d_)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^(n+1))/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^(n+2), x], x] /;$ $\text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /;$ $\text{FreeQ}\{c, d, x\}$

Rule 2642

$\text{Int}[1/\text{Sqrt}[(b_)*\sin[(c_*) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[c + d*x]]/\text{Sqrt}[b*\text{Sin}[c + d*x]], \text{Int}[1/\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ $\text{FreeQ}\{b, c, d, x\}$

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx &= b^5 \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
&= \frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{1}{3} b^3 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{(b^3 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
&= \frac{2b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 51, normalized size = 0.71

$$\frac{2b^3 \left(\tan(c + dx) + \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^5,x]

[Out] (2*b^3*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.47, size = 0, normalized size = 0.00

$$\text{integral}(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c)^5, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^5,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c)^5, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{5/2} \sec(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^5,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^5, x)

maple [B] time = 0.14, size = 241, normalized size = 3.35

$$\frac{2 \left(-2 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \right)}{3 \sqrt{-b \left(2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)*sec(d*x+c)^5,x)

[Out] -2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2-1)^(1/2)*sqrt(2*(sin^2(1/2*d*x+1/2*c)-1)))/3*sqrt(-b*(2*(sin^4(1/2*d*x+1/2*c))-sin^2(1/2*d*x+1/2*c)))

$$2)^{(1/2)} * (2 * \sin(1/2 * d * x + 1/2 * c)^2 - 1)^{(1/2)} * \text{EllipticF}(\cos(1/2 * d * x + 1/2 * c), 2^{(1/2)}) - 2 * \sin(1/2 * d * x + 1/2 * c)^2 * \cos(1/2 * d * x + 1/2 * c) * b^3 * (b * (2 * \cos(1/2 * d * x + 1/2 * c)^2 - 1) * \sin(1/2 * d * x + 1/2 * c)^2)^{(1/2)} / (-b * (2 * \sin(1/2 * d * x + 1/2 * c)^4 - \sin(1/2 * d * x + 1/2 * c)^2))^{(1/2)} / (2 * \cos(1/2 * d * x + 1/2 * c)^2 - 1) / \sin(1/2 * d * x + 1/2 * c) / (b * (2 * \cos(1/2 * d * x + 1/2 * c)^2 - 1))^{(1/2)} / d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^5,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^5, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^5,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^5, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**5,x)

[Out] Timed out

3.99 $\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx$

Optimal. Leaf size=100

$$\frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[Out] $2/5*b^5*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-6/5*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(5/2)}*\text{Sec}[c + d*x]^6, x]$

[Out] $(-6*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^5*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*b^3*\text{Sin}[c + d*x])/(5*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /;$ FreeQ[{c, d}, x]

Rule 2640

$\text{Int}[\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[b*\text{Sin}[c + d*x]]/\text{Sqrt}[\text{Sin}[c + d*x]], \text{Int}[\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx &= b^6 \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
&= \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{1}{5} (3b^4) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{1}{5} (3b^2) \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{(3b^2 \sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\
&= -\frac{6b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.15, size = 69, normalized size = 0.69

$$\frac{\sec^5(c + dx)(b \cos(c + dx))^{5/2} \left(7 \sin(c + dx) + 3 \sin(3(c + dx)) - 12 \cos^2(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right)\right)}{10d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^6,x]

[Out] ((b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^5*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)]))/(10*d)

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c)^6, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^6,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c)^6, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{5/2} \sec(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^6,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^6, x)

maple [B] time = 0.20, size = 366, normalized size = 3.66

$$2\sqrt{b \left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \left(12 \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{\frac{1}{2}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)*sec(d*x+c)^6,x)

```
[Out] 2/5*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2/sin(1/2*d*x+1/2*c)^3/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)*(12*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^4-24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^2+24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^6,x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^6, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{\frac{5}{2}}}{\cos(c + dx)^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^6,x)
```

```
[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^6, x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**6,x)
```

```
[Out] Timed out
```

3.100 $\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx$

Optimal. Leaf size=100

$$\frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

[Out] $2/7*b^6*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^7,x]

[Out] $(10*b^3*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b^6*\text{Sin}[c + d*x])/(7*d*(b*\text{Cos}[c + d*x])^{(7/2)}) + (10*b^4*\text{Sin}[c + d*x])/(21*d*(b*\text{Cos}[c + d*x])^{(3/2)})$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*SIN[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx &= b^7 \int \frac{1}{(b \cos(c + dx))^{9/2}} dx \\
&= \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{1}{7} (5b^5) \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
&= \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{1}{21} (5b^3) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{(5b^3 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21 \sqrt{b \cos(c + dx)}} \\
&= \frac{10b^3 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.18, size = 69, normalized size = 0.69

$$\frac{\sec^5(c + dx)(b \cos(c + dx))^{5/2} \left(5 \sin(2(c + dx)) + 6 \tan(c + dx) + 10 \cos^{\frac{5}{2}}(c + dx) F\left(\frac{1}{2}(c + dx) \middle| 2\right)\right)}{21d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^7,x]

[Out] ((b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^5*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d)

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c)^7, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^7,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c)^7, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^7 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^7,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^7, x)

maple [B] time = 0.15, size = 398, normalized size = 3.98

$$\frac{2 \left(-40 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 60 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right)}{21d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)*sec(d*x+c)^7,x)

```
[Out] -2/21*(-40*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^6+60*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-40*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-30*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+40*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-16*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))*b^3*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^3/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{5/2} \sec(dx + c)^7 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^7,x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^7, x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^7} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^7,x)
```

```
[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^7, x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**7,x)
```

```
[Out] Timed out
```

3.101 $\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx$

Optimal. Leaf size=128

$$\frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^5 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^3 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{14b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[Out] $2/9*b^7*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*b^5*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-14/15*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*EllipticE(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] time = 0.09, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^5 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^3 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{14b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^(5/2)*\text{Sec}[c + d*x]^8,x]$

[Out] $(-14*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/((15*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^7*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^(9/2)) + (14*b^5*\text{Sin}[c + d*x])/(45*d*(b*\text{Cos}[c + d*x])^(5/2)) + (14*b^3*\text{Sin}[c + d*x])/(15*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]))$

Rule 16

$\text{Int}[(u_*)*(v_)^(m_*)*((b_)*(v_))^(n_), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^(m + n), x], x] /;$ $\text{FreeQ}\{b, n, x\} \ \&\amp; \ \text{IntegerQ}[m]$

Rule 2636

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + \text{Dist}[(n + 2)/(b^2*(n + 1)), \text{Int}[(b*\text{Sin}[c + d*x])^(n + 2), x], x] /;$ $\text{FreeQ}\{b, c, d, x\} \ \&\amp; \ \text{LtQ}[n, -1] \ \&\amp; \ \text{IntegerQ}[2*n]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /;$ $\text{FreeQ}\{c, d, x\}$

Rule 2640

$\text{Int}[\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[b*\text{Sin}[c + d*x]]/\text{Sqrt}[\text{Sin}[c + d*x]], \text{Int}[\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ $\text{FreeQ}\{b, c, d, x\}$

Rubi steps

$$\begin{aligned}
\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx &= b^8 \int \frac{1}{(b \cos(c + dx))^{11/2}} dx \\
&= \frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{1}{9} (7b^6) \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
&= \frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^5 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{1}{15} (7b^4) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^5 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^3 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{1}{15} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^5 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^3 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{7}{15} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= -\frac{14b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^5 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}}
\end{aligned}$$

Mathematica [A] time = 0.24, size = 79, normalized size = 0.62

$$\frac{\sec^7(c + dx)(b \cos(c + dx))^{5/2} \left(150 \sin(c + dx) + 91 \sin(3(c + dx)) + 21 \sin(5(c + dx)) - 336 \cos^2(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right)\right)}{360d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^8,x]

[Out] ((b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^7*(-336*Cos[c + d*x]^(9/2)*EllipticE[(c + d*x)/2, 2] + 150*Sin[c + d*x] + 91*Sin[3*(c + d*x)] + 21*Sin[5*(c + d*x)]))/ (360*d)

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c)^8, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^8,x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c)^8, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{5/2} \sec(dx + c)^8 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^8,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^8, x)

maple [B] time = 0.23, size = 414, normalized size = 3.23

$$2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^3 \left(-\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{144b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} - \frac{7\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{180b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*cos(d*x+c))^(5/2)*sec(d*x+c)^8,x)`

[Out]
$$-2*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b^3*(-1/144*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^5-7/180*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^3-14/15*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*E\ellipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*(E\ellipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-E\ellipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{5}{2}} \sec(dx + c)^8 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^8,x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^8, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{\frac{5}{2}}}{\cos(c + dx)^8} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^8,x)`

[Out] `int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^8, x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**8,x)`

[Out] Timed out

3.102 $\int (b \cos(c + dx))^{7/2} dx$

Optimal. Leaf size=98

$$\frac{10b^4 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b^3 \sin(c + dx) \sqrt{b \cos(c + dx)}}{21d} + \frac{2b \sin(c + dx) (b \cos(c + dx))^{5/2}}{7d}$$

[Out] $2/7*b*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/d+10/21*b^4*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+10/21*b^3*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.05, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2635, 2642, 2641}

$$\frac{10b^3 \sin(c + dx) \sqrt{b \cos(c + dx)}}{21d} + \frac{10b^4 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2b \sin(c + dx) (b \cos(c + dx))^{5/2}}{7d}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(7/2), x]

[Out] $(10*b^4*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (10*b^3*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(21*d) + (2*b*(b*\text{Cos}[c + d*x])^{(5/2)}*\text{Sin}[c + d*x])/(7*d)$

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*cos[c + d*x]*(b*sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{7/2} dx &= \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} + \frac{1}{7} (5b^2) \int (b \cos(c + dx))^{3/2} dx \\ &= \frac{10b^3 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} + \frac{1}{21} (5b^4) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\ &= \frac{10b^3 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} + \frac{(5b^4 \sqrt{\cos(c + dx)})}{21 \sqrt{b \cos(c + dx)}} \\ &= \frac{10b^4 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b^3 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} \end{aligned}$$

Mathematica [A] time = 0.02, size = 76, normalized size = 0.78

$$\frac{b^3 \sqrt{b \cos(c + dx)} \left(20F\left(\frac{1}{2}(c + dx) \middle| 2\right) + (23 \sin(c + dx) + 3 \sin(3(c + dx))) \sqrt{\cos(c + dx)} \right)}{42d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(7/2), x]

[Out] (b^3*Sqrt[b*cos[c + d*x]]*(20*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(23*Sin[c + d*x] + 3*Sin[3*(c + d*x)])))/(42*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \cos(dx + c)} b^3 \cos(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(7/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*b^3*cos(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(7/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(7/2), x)

maple [A] time = 0.14, size = 210, normalized size = 2.14

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^4 \left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(7/2), x)

[Out] -2/21*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^4*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(7/2), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (b \cos(c + dx))^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(7/2), x)

[Out] int((b*cos(c + d*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(7/2), x)

[Out] Timed out

3.103 $\int \frac{\cos^6(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal. Leaf size=125

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^5d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^3d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77bd} + \frac{30\sqrt{\cos(c+dx)}}{77d\sqrt{b \cos(c+dx)}}$$

[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b^3/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^5/d+30/77*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+30/77*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b/d

Rubi [A] time = 0.09, antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^5d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^3d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77bd} + \frac{30\sqrt{\cos(c+dx)}}{77d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^6/Sqrt[b*Cos[c + d*x]],x]

[Out] (30*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*d*Sqrt[b*Cos[c + d*x]]) + (30*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*b*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b^3*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^5*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n-1)/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^6(c+dx)}{\sqrt{b\cos(c+dx)}} dx &= \frac{\int (b\cos(c+dx))^{11/2} dx}{b^6} \\
&= \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^5d} + \frac{9 \int (b\cos(c+dx))^{7/2} dx}{11b^4} \\
&= \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^3d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^5d} + \frac{45 \int (b\cos(c+dx))^{5/2} dx}{77b^2} \\
&= \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77bd} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^3d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^5d} \\
&= \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77bd} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^3d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^5d} \\
&= \frac{30\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{77d\sqrt{b\cos(c+dx)}} + \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77bd} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^3d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^5d}
\end{aligned}$$

Mathematica [A] time = 0.13, size = 73, normalized size = 0.58

$$\frac{347 \sin(2(c+dx)) + 64 \sin(4(c+dx)) + 7 \sin(6(c+dx)) + 480\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{1232d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^6/Sqrt[b*Cos[c + d*x]], x]

[Out] (480*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 347*Sin[2*(c + d*x)] + 64*Sin[4*(c + d*x)] + 7*Sin[6*(c + d*x)])/(1232*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^5}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^5/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^6}{\sqrt{b\cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^6/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.15, size = 233, normalized size = 1.86

$$\frac{2\sqrt{b}\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(448\left(\cos^{13}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1568\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 2384\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1280\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 448\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 64\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{77\sqrt{-b}\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^6/(b*cos(d*x+c))^(1/2),x)`

[Out]
$$-2/77*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(448*\cos(1/2*d*x+1/2*c)^{13}-1568*\cos(1/2*d*x+1/2*c)^{11}+2384*\cos(1/2*d*x+1/2*c)^9-2040*\cos(1/2*d*x+1/2*c)^7+1084*\cos(1/2*d*x+1/2*c)^5-370*\cos(1/2*d*x+1/2*c)^3+15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)}+62*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^6}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^6/sqrt(b*cos(d*x + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^6}{\sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)^6/(b*cos(c+d*x))^(1/2),x)`

[Out] `int(cos(c+d*x)^6/(b*cos(c+d*x))^(1/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**6/(b*cos(d*x+c))**(1/2),x)`

[Out] Timed out

$$3.104 \quad \int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^4d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^2d} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15bd\sqrt{\cos(c+dx)}}$$

[Out] 14/45*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b^2/d+2/9*(b*cos(d*x+c))^(7/2)*sin(d*x+c)/b^4/d+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^4d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^2d} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15bd\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^5/Sqrt[b*Cos[c + d*x]],x]

[Out] (14*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*b*d*Sqrt[Cos[c + d*x]]) + (14*(b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(45*b^2*d) + (2*(b*Cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b^4*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n-1)/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^5(c+dx)}{\sqrt{b\cos(c+dx)}} dx &= \frac{\int (b\cos(c+dx))^{9/2} dx}{b^5} \\
&= \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^4d} + \frac{7 \int (b\cos(c+dx))^{5/2} dx}{9b^3} \\
&= \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^2d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^4d} + \frac{7 \int \sqrt{b\cos(c+dx)} dx}{15b} \\
&= \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^2d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^4d} + \frac{(7\sqrt{b\cos(c+dx)}) \int \sqrt{b\cos(c+dx)} dx}{15b\sqrt{\cos(c+dx)}} \\
&= \frac{14\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15bd\sqrt{\cos(c+dx)}} + \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^2d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^4d}
\end{aligned}$$

Mathematica [A] time = 0.12, size = 71, normalized size = 0.71

$$\frac{(38 \sin(2(c+dx)) + 5 \sin(4(c+dx))) \cos(c+dx) + 168 \sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{180d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^5/Sqrt[b*Cos[c + d*x]], x]

[Out] (168*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)]))/(180*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)}\cos(dx+c)^4}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^4/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^5}{\sqrt{b\cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^5/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.14, size = 220, normalized size = 2.20

$$\frac{2\sqrt{b}\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(160\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 480\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 616\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 280\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 112\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 14\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{45\sqrt{-b}\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^5/(b*cos(d*x+c))^(1/2),x)`

[Out]
$$-2/45*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(160*\cos(1/2*d*x+1/2*c)^{11}-480*\cos(1/2*d*x+1/2*c)^9+616*\cos(1/2*d*x+1/2*c)^7-432*\cos(1/2*d*x+1/2*c)^5+160*\cos(1/2*d*x+1/2*c)^3-21*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-24*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^5}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^5/sqrt(b*cos(d*x + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^5}{\sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)^5/(b*cos(c+d*x))^(1/2),x)`

[Out] `int(cos(c+d*x)^5/(b*cos(c+d*x))^(1/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**5/(b*cos(d*x+c))**(1/2),x)`

[Out] Timed out

$$3.105 \quad \int \frac{\cos^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=97

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^3d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21bd} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21d\sqrt{b \cos(c+dx)}}$$

[Out] 2/7*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b^3/d+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+10/21*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b/d

Rubi [A] time = 0.06, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^3d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21bd} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^4/Sqrt[b*Cos[c + d*x]],x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(21*d*Sqrt[b*Cos[c + d*x]]) + (10*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(21*b*d) + (2*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*b^3*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*SIN[c + d*x])^(n-1)/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*SIN[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^4(c+dx)}{\sqrt{b\cos(c+dx)}} dx &= \frac{\int (b\cos(c+dx))^{7/2} dx}{b^4} \\
&= \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^3d} + \frac{5 \int (b\cos(c+dx))^{3/2} dx}{7b^2} \\
&= \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21bd} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^3d} + \frac{5}{21} \int \frac{1}{\sqrt{b\cos(c+dx)}} dx \\
&= \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21bd} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^3d} + \frac{(5\sqrt{\cos(c+dx)}) \int}{21\sqrt{b\cos(c+dx)}} \\
&= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21d\sqrt{b\cos(c+dx)}} + \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21bd} + \frac{2(b\cos(c+dx))^{5/2}}{7b^3d}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 63, normalized size = 0.65

$$\frac{26 \sin(2(c+dx)) + 3 \sin(4(c+dx)) + 40\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{84d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^4/Sqrt[b*Cos[c + d*x]], x]

[Out] (40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)])/(84*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^3}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^3/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^4}{\sqrt{b\cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^4/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.15, size = 207, normalized size = 2.13

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^4/(b*cos(d*x+c))^(1/2),x)`

[Out]
$$-2/21*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(48*\cos(1/2*d*x+1/2*c)^9-120*\cos(1/2*d*x+1/2*c)^7+128*\cos(1/2*d*x+1/2*c)^5-72*\cos(1/2*d*x+1/2*c)^3+5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})+16*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^4}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^4/sqrt(b*cos(d*x + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^4}{\sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)^4/(b*cos(c+d*x))^(1/2),x)`

[Out] `int(cos(c+d*x)^4/(b*cos(c+d*x))^(1/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**4/(b*cos(d*x+c))**(1/2),x)`

[Out] Timed out

$$3.106 \quad \int \frac{\cos^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=72

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^2d} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5bd\sqrt{\cos(c+dx)}}$$

[Out] 2/5*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b^2/d+6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^2d} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5bd\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^3/Sqrt[b*Cos[c + d*x]],x]

[Out] (6*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(5*b*d*Sqrt[Cos[c + d*x]]) + (2*(b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*b^2*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Sin[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx &= \frac{\int (b \cos(c+dx))^{5/2} dx}{b^3} \\
&= \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5b^2d} + \frac{3 \int \sqrt{b \cos(c+dx)} dx}{5b} \\
&= \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5b^2d} + \frac{(3\sqrt{b \cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5b\sqrt{\cos(c+dx)}} \\
&= \frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5bd\sqrt{\cos(c+dx)}} + \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5b^2d}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 58, normalized size = 0.81

$$\frac{\sin(2(c+dx)) \cos(c+dx) + 6\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3/Sqrt[b*Cos[c + d*x]], x]

[Out] (6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*Sin[2*(c + d*x)])/(5*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.93, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \cos(dx+c)^2}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^2/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^3}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^3/sqrt(b*cos(d*x + c)), x)

maple [B] time = 0.15, size = 210, normalized size = 2.92

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(-8\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^3/(b*cos(d*x+c))^(1/2), x)

```
[Out] -2/5*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-8*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+8*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^3}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(cos(d*x + c)^3/sqrt(b*cos(d*x + c)), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^3}{\sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^3/(b*cos(c + d*x))^(1/2),x)
```

```
[Out] int(cos(c + d*x)^3/(b*cos(c + d*x))^(1/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**3/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

$$3.107 \quad \int \frac{\cos^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=69

$$\frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3bd} + \frac{2 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3d \sqrt{b \cos(c+dx)}}$$

[Out] 2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+2/3*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b/d

Rubi [A] time = 0.04, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3bd} + \frac{2 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/Sqrt[b*Cos[c + d*x]],x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(3*d*Sqrt[b*Cos[c + d*x]]) + (2*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(3*b*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Sin[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(c+dx)}{\sqrt{b\cos(c+dx)}} dx &= \frac{\int (b\cos(c+dx))^{3/2} dx}{b^2} \\
&= \frac{2\sqrt{b\cos(c+dx)} \sin(c+dx)}{3bd} + \frac{1}{3} \int \frac{1}{\sqrt{b\cos(c+dx)}} dx \\
&= \frac{2\sqrt{b\cos(c+dx)} \sin(c+dx)}{3bd} + \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3\sqrt{b\cos(c+dx)}} \\
&= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3d\sqrt{b\cos(c+dx)}} + \frac{2\sqrt{b\cos(c+dx)} \sin(c+dx)}{3bd}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 51, normalized size = 0.74

$$\frac{\sin(2(c+dx)) + 2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/Sqrt[b*Cos[c + d*x]], x]

[Out] (2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)])/(3*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)}\cos(dx+c)}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^2}{\sqrt{b\cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/sqrt(b*cos(d*x + c)), x)

maple [B] time = 0.17, size = 187, normalized size = 2.71

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(4\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)}\sqrt{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2/(b*cos(d*x+c))^(1/2), x)

[Out]
$$-2/3*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(4*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^2}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/2), x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^2/sqrt(b*cos(d*x + c)), x)`

mupad [B] time = 0.14, size = 58, normalized size = 0.84

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3d\sqrt{b\cos(c+dx)}} + \frac{2\sin(c+dx)\sqrt{b\cos(c+dx)}}{3bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2/(b*cos(c + d*x))^(1/2), x)`

[Out]
$$(2*\cos(c + d*x)^{(1/2)}*\text{ellipticF}(c/2 + (d*x)/2, 2))/(3*d*(b*\cos(c + d*x))^{(1/2)}) + (2*\sin(c + d*x)*(b*\cos(c + d*x))^{(1/2)})/(3*b*d)$$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(1/2), x)`

[Out] Timed out

$$3.108 \quad \int \frac{\cos(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=41

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 2640, 2639}

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/Sqrt[b*Cos[c + d*x]], x]

[Out] $(2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(b*d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_)^(n_)), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\cos(c+dx)}{\sqrt{b \cos(c+dx)}} dx &= \frac{\int \sqrt{b \cos(c+dx)} dx}{b} \\ &= \frac{\sqrt{b \cos(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b\sqrt{\cos(c+dx)}} \\ &= \frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\middle|2\right)}{bd\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 41, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]/Sqrt[b*Cos[c + d*x]],x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.83, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx + c)}}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{\sqrt{b \cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/sqrt(b*cos(d*x + c)), x)

maple [B] time = 0.13, size = 141, normalized size = 3.44

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}\text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*cos(d*x+c))^(1/2),x)

[Out] 2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{\sqrt{b \cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/sqrt(b*cos(d*x + c)), x)

mupad [B] time = 0.15, size = 33, normalized size = 0.80

$$\frac{2\sqrt{\cos(c + dx)}E\left(\frac{c}{2} + \frac{dx}{2}\middle|2\right)}{d\sqrt{b \cos(c + dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)/(b*cos(c + d*x))^(1/2),x)

[Out] $(2\cos(c + dx)^{1/2}\text{ellipticE}(c/2 + (dx)/2, 2))/(d(b\cos(c + dx))^{1/2})$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))**(1/2), x)

[Out] Integral(cos(c + d*x)/sqrt(b*cos(c + d*x)), x)

$$3.109 \quad \int \frac{1}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=38

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d\sqrt{b \cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2642, 2641}

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[b*Cos[c + d*x]], x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{b \cos(c+dx)}} dx &= \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 38, normalized size = 1.00

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[b*Cos[c + d*x]], x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)}}{b \cos(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(b*cos(d*x + c)), x)

maple [C] time = 0.05, size = 54, normalized size = 1.42

$$\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \operatorname{am}^{-1}\left(\frac{dx}{2} + \frac{c}{2} \mid \sqrt{2}\right)}{d\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(d*x+c))^(1/2),x)

[Out] 2/d/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)*(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)*InverseJacobiAM(1/2*d*x+1/2*c,2^(1/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(b*cos(d*x + c)), x)

mupad [B] time = 0.19, size = 33, normalized size = 0.87

$$\frac{2\sqrt{\cos(c + dx)} F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d\sqrt{b \cos(c + dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(c + d*x))^(1/2),x)

[Out] (2*cos(c + d*x)^(1/2)*ellipticF(c/2 + (d*x)/2, 2))/(d*(b*cos(c + d*x))^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(1/sqrt(b*cos(c + d*x)), x)

$$3.110 \quad \int \frac{\sec(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=65

$$\frac{2 \sin(c+dx)}{d \sqrt{b \cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}}$$

[Out] 2*sin(d*x+c)/d/(b*cos(d*x+c))^(1/2)-2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2 \sin(c+dx)}{d \sqrt{b \cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/Sqrt[b*Cos[c + d*x]],x]

[Out] (-2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]]) + (2*Sin[c + d*x])/(d*Sqrt[b*Cos[c + d*x]])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec(c+dx)}{\sqrt{b\cos(c+dx)}} dx &= b \int \frac{1}{(b\cos(c+dx))^{3/2}} dx \\
&= \frac{2\sin(c+dx)}{d\sqrt{b\cos(c+dx)}} - \frac{\int \sqrt{b\cos(c+dx)} dx}{b} \\
&= \frac{2\sin(c+dx)}{d\sqrt{b\cos(c+dx)}} - \frac{\sqrt{b\cos(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b\sqrt{\cos(c+dx)}} \\
&= -\frac{2\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx)\middle|2\right)}{bd\sqrt{\cos(c+dx)}} + \frac{2\sin(c+dx)}{d\sqrt{b\cos(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 47, normalized size = 0.72

$$\frac{2\left(\sin(c+dx) - \sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx)\middle|2\right)\right)}{d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/Sqrt[b*Cos[c + d*x]], x]

[Out] (2*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)}\sec(dx+c)}{b\cos(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{\sqrt{b\cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.16, size = 165, normalized size = 2.54

$$\frac{2\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b\left(\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\right) \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*cos(d*x+c))^(1/2), x)

[Out] -2*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)*((sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c

$), 2^{(1/2)} - 2 \sin(1/2 dx + 1/2 c)^2 \cos(1/2 dx + 1/2 c) / (-b(2 \sin(1/2 dx + 1/2 c)^4 - \sin(1/2 dx + 1/2 c)^2))^{(1/2)} / \sin(1/2 dx + 1/2 c) / (b(2 \cos(1/2 dx + 1/2 c)^2 - 1))^{(1/2)} / d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{\sqrt{b \cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/sqrt(b*cos(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx) \sqrt{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sec(c + d*x)/sqrt(b*cos(c + d*x)), x)

$$3.111 \quad \int \frac{\sec^2(c+dx)}{\sqrt{b} \cos(c+dx)} dx$$

Optimal. Leaf size=67

$$\frac{2b \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3d\sqrt{b} \cos(c+dx)}$$

[Out] $2/3*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3d\sqrt{b} \cos(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/Sqrt[b*Cos[c + d*x]],x]

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b*\text{Sin}[c + d*x])/(3*d*(b*\text{Cos}[c + d*x])^{(3/2)})$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*SIN[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx &= b^2 \int \frac{1}{(b \cos(c+dx))^{5/2}} dx \\
&= \frac{2b \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}} + \frac{1}{3} \int \frac{1}{\sqrt{b \cos(c+dx)}} dx \\
&= \frac{2b \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}} + \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3\sqrt{b \cos(c+dx)}} \\
&= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3d\sqrt{b \cos(c+dx)}} + \frac{2b \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 48, normalized size = 0.72

$$\frac{2 \left(\tan(c+dx) + \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{3d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/Sqrt[b*Cos[c + d*x]], x]

[Out] (2*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.81, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \sec(dx+c)^2}{b \cos(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^2/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/sqrt(b*cos(d*x + c)), x)

maple [B] time = 0.16, size = 238, normalized size = 3.55

$$\frac{2 \left(-2\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)} \text{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\dots} \right)}{3\sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2/(b*cos(d*x+c))^(1/2), x)

[Out] $-2/3*(-2*(\sin(1/2*d*x+1/2*c))^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c), 2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c), 2^{(1/2)})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^2-1)/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/2), x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^2/sqrt(b*cos(d*x + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^2 \sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(1/2)), x)`

[Out] `int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(1/2), x)`

[Out] `Integral(sec(c + d*x)**2/sqrt(b*cos(c + d*x)), x)`

$$3.112 \quad \int \frac{\sec^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=97

$$\frac{2b^2 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5d\sqrt{b \cos(c+dx)}} - \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5bd\sqrt{\cos(c+dx)}}$$

[Out] $2/5*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^2 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5d\sqrt{b \cos(c+dx)}} - \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5bd\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3/Sqrt[b*Cos[c + d*x]], x]

[Out] $(-6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^2*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*\text{Sin}[c + d*x])/(5*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^3(c+dx)}{\sqrt{b\cos(c+dx)}} dx &= b^3 \int \frac{1}{(b\cos(c+dx))^{7/2}} dx \\
&= \frac{2b^2 \sin(c+dx)}{5d(b\cos(c+dx))^{5/2}} + \frac{1}{5}(3b) \int \frac{1}{(b\cos(c+dx))^{3/2}} dx \\
&= \frac{2b^2 \sin(c+dx)}{5d(b\cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5d\sqrt{b\cos(c+dx)}} - \frac{3 \int \sqrt{b\cos(c+dx)} dx}{5b} \\
&= \frac{2b^2 \sin(c+dx)}{5d(b\cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5d\sqrt{b\cos(c+dx)}} - \frac{(3\sqrt{b\cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5b\sqrt{\cos(c+dx)}} \\
&= -\frac{6\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5bd\sqrt{\cos(c+dx)}} + \frac{2b^2 \sin(c+dx)}{5d(b\cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5d\sqrt{b\cos(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 65, normalized size = 0.67

$$\frac{6 \sin(c+dx) - 6\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2 \tan(c+dx) \sec(c+dx)}{5d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3/Sqrt[b*Cos[c + d*x]], x]

[Out] (-6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 6*Sin[c + d*x] + 2*Sec[c + d*x]*Tan[c + d*x])/(5*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \sec(dx+c)^3}{b\cos(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^3/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{\sqrt{b\cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/sqrt(b*cos(d*x + c)), x)

maple [B] time = 0.25, size = 366, normalized size = 3.77

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(12 \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{\frac{1}{2}} - \right)}{5bd\sqrt{\cos(c+dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^3/(b*cos(d*x+c))^(1/2),x)`

[Out] $2/5*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b/\sin(1/2*d*x+1/2*c)^3/(8*\sin(1/2*d*x+1/2*c)^6-12*\sin(1/2*d*x+1/2*c)^4+6*\sin(1/2*d*x+1/2*c)^2-1)*(12*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*\sin(1/2*d*x+1/2*c)^4-24*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6-12*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*\sin(1/2*d*x+1/2*c)^2+24*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-8*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))*(-2*\sin(1/2*d*x+1/2*c)^4*b+\sin(1/2*d*x+1/2*c)^2*b)^{(1/2)}/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^3/sqrt(b*cos(d*x + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^3 \sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(1/2)),x)`

[Out] `int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(1/2),x)`

[Out] `Integral(sec(c + d*x)**3/sqrt(b*cos(c + d*x)), x)`

$$3.113 \quad \int \frac{\sec^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=95

$$\frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21d\sqrt{b \cos(c+dx)}}$$

[Out] 2/7*b^3*sin(d*x+c)/d/(b*cos(d*x+c))^(7/2)+10/21*b*sin(d*x+c)/d/(b*cos(d*x+c))^(3/2)+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^4/Sqrt[b*Cos[c + d*x]],x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(21*d*Sqrt[b*Cos[c + d*x]]) + (2*b^3*Sin[c + d*x])/(7*d*(b*Cos[c + d*x])^(7/2)) + (10*b*Sin[c + d*x])/(21*d*(b*Cos[c + d*x])^(3/2))

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx &= b^4 \int \frac{1}{(b \cos(c+dx))^{9/2}} dx \\
&= \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{1}{7} (5b^2) \int \frac{1}{(b \cos(c+dx))^{5/2}} dx \\
&= \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{5}{21} \int \frac{1}{\sqrt{b \cos(c+dx)}} dx \\
&= \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{(5\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21\sqrt{b \cos(c+dx)}} \\
&= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21d\sqrt{b \cos(c+dx)}} + \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.14, size = 63, normalized size = 0.66

$$\frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2 \tan(c+dx) (3 \sec^2(c+dx) + 5)}{21d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^4/Sqrt[b*Cos[c + d*x]], x]

[Out] (10*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 2*(5 + 3*Sec[c + d*x]^2)*Tan[c + d*x])/(21*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \sec(dx+c)^4}{b \cos(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^4/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^4}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^4/sqrt(b*cos(d*x + c)), x)

maple [B] time = 0.18, size = 395, normalized size = 4.16

$$\frac{2\left(-40\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \text{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 60\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\right)}{21d\sqrt{b \cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^4/(b*cos(d*x+c))^(1/2),x)

[Out] $-2/21*(-40*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^6+60*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^4-40*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6-30*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2+40*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-16*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^2-1)^3/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^4}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^4/sqrt(b*cos(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^4 \sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^4*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^4*(b*cos(c + d*x))^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**4/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sec(c + d*x)**4/sqrt(b*cos(c + d*x)), x)

$$3.114 \quad \int \frac{\sec^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=125

$$\frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^2 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15bd\sqrt{\cos(c+dx)}}$$

[Out] $2/9*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(9/2)}+14/45*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+14/15*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-14/15*(\cos(1/2*d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.09, antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^2 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15bd\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^5/Sqrt[b*Cos[c + d*x]],x]

[Out] $(-14*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*b*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^4*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^{(9/2)}) + (14*b^2*\text{Sin}[c + d*x])/(45*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (14*\text{Sin}[c + d*x])/(15*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx &= b^5 \int \frac{1}{(b \cos(c+dx))^{11/2}} dx \\
&= \frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{1}{9} (7b^3) \int \frac{1}{(b \cos(c+dx))^{7/2}} dx \\
&= \frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^2 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{1}{15} (7b) \int \frac{1}{(b \cos(c+dx))^{3/2}} dx \\
&= \frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^2 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{7 \int \sqrt{b \cos(c+dx)}}{15b} \\
&= \frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^2 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{(7\sqrt{b \cos(c+dx)})}{15b\sqrt{b \cos(c+dx)}} \\
&= -\frac{14\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15bd\sqrt{\cos(c+dx)}} + \frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^2 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} +
\end{aligned}$$

Mathematica [A] time = 0.30, size = 77, normalized size = 0.62

$$\frac{42 \sin(c+dx) - 42\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2 \tan(c+dx) \sec(c+dx) (5 \sec^2(c+dx) + 7)}{45d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^5/Sqrt[b*Cos[c + d*x]], x]

[Out] (-42*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 42*Sin[c + d*x] + 2*Sec[c + d*x]*(7 + 5*Sec[c + d*x]^2)*Tan[c + d*x])/(45*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \sec(dx+c)^5}{b \cos(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^5/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^5}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^5/sqrt(b*cos(d*x + c)), x)

maple [B] time = 0.25, size = 411, normalized size = 3.29

$$\sqrt{b \left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{72b \left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} - \frac{7 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{90b \left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^5/(b*cos(d*x+c))^(1/2),x)`

[Out]
$$-(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-1/72*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^5-7/90*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^3-28/15*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+14/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-14/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*(EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^5}{\sqrt{b}\cos(dx+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^5/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^5/sqrt(b*cos(d*x + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^5 \sqrt{b}\cos(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c+d*x)^5*(b*cos(c+d*x))^(1/2)),x)`

[Out] `int(1/(cos(c+d*x)^5*(b*cos(c+d*x))^(1/2)),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**5/(b*cos(d*x+c))**(1/2),x)`

[Out] Timed out

$$3.115 \quad \int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=128

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^6d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^4d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77b^2d} + \frac{30\sqrt{\cos(c+dx)}}{77bd}$$

[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b^4/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^6/d+30/77*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)+30/77*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b^2/d

Rubi [A] time = 0.08, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^6d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^4d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77b^2d} + \frac{30\sqrt{\cos(c+dx)}}{77bd}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^7/(b*Cos[c + d*x])^(3/2), x]

[Out] (30*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*b*d*Sqrt[b*Cos[c + d*x]]) + (30*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*b^2*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b^4*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^6*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Ssin[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*Ssin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Ssin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{3/2}} dx &= \frac{\int (b\cos(c+dx))^{11/2} dx}{b^7} \\
&= \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^6d} + \frac{9 \int (b\cos(c+dx))^{7/2} dx}{11b^5} \\
&= \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^4d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^6d} + \frac{45 \int (b\cos(c+dx))^{3/2} dx}{77b^3} \\
&= \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77b^2d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^4d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^6d} \\
&= \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77b^2d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^4d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^6d} \\
&= \frac{30\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{77bd\sqrt{b\cos(c+dx)}} + \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77b^2d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^4d}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 76, normalized size = 0.59

$$\frac{347 \sin(2(c+dx)) + 64 \sin(4(c+dx)) + 7 \sin(6(c+dx)) + 480\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{1232bd\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^7/(b*Cos[c + d*x])^(3/2), x]

[Out] (480*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 347*Sin[2*(c + d*x)] + 64*Sin[4*(c + d*x)] + 7*Sin[6*(c + d*x)])/(1232*b*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.79, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^5}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^5/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^7}{(b\cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^7/(b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.15, size = 236, normalized size = 1.84

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(448\left(\cos^{13}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1568\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 2384\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1280\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 256\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 16\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 2\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{77b\sqrt{-b}\left(2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^7/(b*cos(d*x+c))^(3/2),x)`

[Out]
$$-2/77*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b*(448*\cos(1/2*d*x+1/2*c)^{13}-1568*\cos(1/2*d*x+1/2*c)^{11}+2384*\cos(1/2*d*x+1/2*c)^9-2040*\cos(1/2*d*x+1/2*c)^7+1084*\cos(1/2*d*x+1/2*c)^5-370*\cos(1/2*d*x+1/2*c)^3+15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+62*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^7}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^7/(b*cos(d*x + c))^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^7}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)^7/(b*cos(c+d*x))^(3/2),x)`

[Out] `int(cos(c+d*x)^7/(b*cos(c+d*x))^(3/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**7/(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

$$3.116 \quad \int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^5d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^3d} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15b^2d\sqrt{\cos(c+dx)}}$$

[Out] 14/45*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b^3/d+2/9*(b*cos(d*x+c))^(7/2)*sin(d*x+c)/b^5/d+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^2/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^5d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^3d} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15b^2d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^6/(b*Cos[c + d*x])^(3/2), x]

[Out] (14*sqrt[b*cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*b^2*d*sqrt[Cos[c + d*x]]) + (14*(b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(45*b^3*d) + (2*(b*cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b^5*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*cos[c + d*x])*(b*sin[c + d*x])^(n-1)/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*sin[c + d*x]]/sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^6(c+dx)}{(b\cos(c+dx))^{3/2}} dx &= \frac{\int (b\cos(c+dx))^{9/2} dx}{b^6} \\
&= \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^5d} + \frac{7 \int (b\cos(c+dx))^{5/2} dx}{9b^4} \\
&= \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^3d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^5d} + \frac{7 \int \sqrt{b\cos(c+dx)} dx}{15b^2} \\
&= \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^3d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^5d} + \frac{(7\sqrt{b\cos(c+dx)})^2}{15b^2} \\
&= \frac{14\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15b^2d\sqrt{\cos(c+dx)}} + \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^3d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^5d}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 74, normalized size = 0.74

$$\frac{(38 \sin(2(c+dx)) + 5 \sin(4(c+dx))) \cos(c+dx) + 168 \sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{180bd\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^6/(b*Cos[c + d*x])^(3/2), x]

[Out] (168*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)]))/(180*b*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.71, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)}\cos(dx+c)^4}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^4/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^6}{(b\cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^6/(b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.16, size = 223, normalized size = 2.23

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(160\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 480\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 616\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 280\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 56\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 8\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{45b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^6/(b*cos(d*x+c))^(3/2),x)`

[Out]
$$-2/45*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{1/2}/b*(160*\cos(1/2*d*x+1/2*c)^{11}-480*\cos(1/2*d*x+1/2*c)^9+616*\cos(1/2*d*x+1/2*c)^7-432*\cos(1/2*d*x+1/2*c)^5+160*\cos(1/2*d*x+1/2*c)^3-21*(\sin(1/2*d*x+1/2*c)^2)^{1/2}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{1/2}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{1/2})-24*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{1/2}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{1/2}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^6}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^6/(b*cos(d*x + c))^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^6}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)^6/(b*cos(c+d*x))^(3/2),x)`

[Out] `int(cos(c+d*x)^6/(b*cos(c+d*x))^(3/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**6/(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

$$3.117 \quad \int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^4d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21b^2d} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21bd\sqrt{b \cos(c+dx)}}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b^4/d+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}+10/21*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/b^2/d$

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^4d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21b^2d} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^5/(b*Cos[c + d*x])^(3/2), x]`

[Out] `(10*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(21*b*d*sqrt[b*Cos[c + d*x]]) + (10*sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(21*b^2*d) + (2*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*b^4*d)`

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2635

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*SIN[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*SIN[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2641

`Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 2642

`Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[SIN[c + d*x]]/sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \frac{\cos^5(c+dx)}{(b\cos(c+dx))^{3/2}} dx &= \frac{\int (b\cos(c+dx))^{7/2} dx}{b^5} \\
&= \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^4d} + \frac{5 \int (b\cos(c+dx))^{3/2} dx}{7b^3} \\
&= \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21b^2d} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^4d} + \frac{5 \int \frac{1}{\sqrt{b\cos(c+dx)}} dx}{21b} \\
&= \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21b^2d} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^4d} + \frac{(5\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21b\sqrt{b\cos(c+dx)}} \\
&= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21bd\sqrt{b\cos(c+dx)}} + \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21b^2d} + \frac{2(b\cos(c+dx))^{5/2}}{7b^4d}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 66, normalized size = 0.66

$$\frac{26 \sin(2(c+dx)) + 3 \sin(4(c+dx)) + 40\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{84bd\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^5/(b*Cos[c + d*x])^(3/2), x]

[Out] (40*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)])/(84*b*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^3}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^3/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^5}{(b\cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^5/(b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.20, size = 210, normalized size = 2.10

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{21b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^5/(b*cos(d*x+c))^(3/2),x)

[Out]
$$-2/21*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b*(48*\cos(1/2*d*x+1/2*c)^9-120*\cos(1/2*d*x+1/2*c)^7+128*\cos(1/2*d*x+1/2*c)^5-72*\cos(1/2*d*x+1/2*c)^3+5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)}+16*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^5}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^5/(b*cos(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^5}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c+d*x)^5/(b*cos(c+d*x))^(3/2),x)

[Out] int(cos(c+d*x)^5/(b*cos(c+d*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**5/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

$$3.118 \quad \int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=72

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^3d} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5b^2d\sqrt{\cos(c+dx)}}$$

[Out] $2/5*(b*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/b^3/d+6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^2/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^3d} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5b^2d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^4/(b*Cos[c + d*x])^(3/2), x]

[Out] $(6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(5*b^3*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*SIN[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*SIN[c + d*x]]/Sqrt[SIN[c + d*x]], Int[Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^4(c+dx)}{(b\cos(c+dx))^{3/2}} dx &= \frac{\int (b\cos(c+dx))^{5/2} dx}{b^4} \\
&= \frac{2(b\cos(c+dx))^{3/2} \sin(c+dx)}{5b^3d} + \frac{3 \int \sqrt{b\cos(c+dx)} dx}{5b^2} \\
&= \frac{2(b\cos(c+dx))^{3/2} \sin(c+dx)}{5b^3d} + \frac{(3\sqrt{b\cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5b^2\sqrt{\cos(c+dx)}} \\
&= \frac{6\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5b^2d\sqrt{\cos(c+dx)}} + \frac{2(b\cos(c+dx))^{3/2} \sin(c+dx)}{5b^3d}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 61, normalized size = 0.85

$$\frac{\sin(2(c+dx)) \cos(c+dx) + 6\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5bd\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^4/(b*Cos[c + d*x])^(3/2), x]

[Out] (6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*Sin[2*(c + d*x)])/(5*b*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.91, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^2}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^2/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^4}{(b\cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^4/(b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.16, size = 213, normalized size = 2.96

$$\frac{2\sqrt{b}\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(-8\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5b\sqrt{-b}\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^4/(b*cos(d*x+c))^(3/2), x)

[Out]
$$-2/5*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b*(-8*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+8*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^4}{(b \cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^4/(b*cos(d*x + c))^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^4}{(b \cos(c+dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^4/(b*cos(c + d*x))^(3/2),x)`

[Out] `int(cos(c + d*x)^4/(b*cos(c + d*x))^(3/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**4/(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

$$3.119 \quad \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=72

$$\frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3b^2d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3bd\sqrt{b \cos(c+dx)}}$$

[Out] 2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)+2/3*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b^2/d

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3b^2d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^3/(b*Cos[c + d*x])^(3/2),x]

[Out] (2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(3*b*d*sqrt[b*Cos[c + d*x]]) + (2*sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(3*b^2*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*SIN[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= \frac{\int (b \cos(c+dx))^{3/2} dx}{b^3} \\
&= \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3b^2d} + \frac{\int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{3b} \\
&= \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3b^2d} + \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b\sqrt{b \cos(c+dx)}} \\
&= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3bd\sqrt{b \cos(c+dx)}} + \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3b^2d}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 54, normalized size = 0.75

$$\frac{\sin(2(c+dx)) + 2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3/(b*Cos[c + d*x])^(3/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)])/(3*b*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \cos(dx+c)}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^3}{(b \cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^3/(b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.24, size = 190, normalized size = 2.64

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(4\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) + \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{3b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^3/(b*cos(d*x+c))^(3/2), x)

```
[Out] -2/3*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/b*(4*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^3}{(b \cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(cos(d*x + c)^3/(b*cos(d*x + c))^(3/2), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^3}{(b \cos(c+dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^3/(b*cos(c + d*x))^(3/2),x)
```

```
[Out] int(cos(c + d*x)^3/(b*cos(c + d*x))^(3/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**3/(b*cos(d*x+c))**(3/2),x)
```

```
[Out] Timed out
```

$$3.120 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=41

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b \cos(c+dx)}}{b^2 d \sqrt{\cos(c+dx)}}$$

[Out] 2*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^2/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2640, 2639}

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b \cos(c+dx)}}{b^2 d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/(b*Cos[c + d*x])^(3/2),x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b^2*d*Sqrt[Cos[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= \frac{\int \sqrt{b \cos(c+dx)} dx}{b^2} \\ &= \frac{\sqrt{b \cos(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b^2 \sqrt{\cos(c+dx)}} \\ &= \frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\middle|2\right)}{b^2 d \sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 41, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b \cos(c+dx)}}{b^2 d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(3/2), x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b^2*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx + c)}}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.19, size = 144, normalized size = 3.51

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}\text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2/(b*cos(d*x+c))^(3/2), x)

[Out] 2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))/b/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(3/2), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(3/2),x)
```

```
[Out] int(cos(c + d*x)^2/(b*cos(c + d*x))^(3/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(3/2),x)
```

```
[Out] Timed out
```


$$3.121 \quad \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=41

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd\sqrt{b \cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 2642, 2641}

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/(b*Cos[c + d*x])^(3/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(b*d*Sqrt[b*Cos[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= \frac{\int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{b} \\ &= \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b\sqrt{b \cos(c+dx)}} \\ &= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 41, normalized size = 1.00

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(3/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(b*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.74, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx + c)}}{b^2 \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/(b^2*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.13, size = 144, normalized size = 3.51

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}\text{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*cos(d*x+c))^(3/2), x)

[Out] -2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(3/2), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)/(b*cos(c + d*x))^(3/2), x)`

[Out] `int(cos(c + d*x)/(b*cos(c + d*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))**(3/2), x)`

[Out] `Integral(cos(c + d*x)/(b*cos(c + d*x))**(3/2), x)`

$$3.122 \quad \int \frac{1}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=68

$$\frac{2 \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^2 d \sqrt{\cos(c+dx)}}$$

[Out] $2*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}-2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^2/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2636, 2640, 2639}

$$\frac{2 \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^2 d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(-3/2), x]

[Out] $(-2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*\text{Sin}[c + d*x])/(b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 2636

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(b \cos(c+dx))^{3/2}} dx &= \frac{2 \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}} - \frac{\int \sqrt{b \cos(c+dx)} dx}{b^2} \\ &= \frac{2 \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}} - \frac{\sqrt{b \cos(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b^2 \sqrt{\cos(c+dx)}} \\ &= -\frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^2 d \sqrt{\cos(c+dx)}} + \frac{2 \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 50, normalized size = 0.74

$$\frac{2 \left(\sin(c + dx) - \sqrt{\cos(c + dx)} E \left(\frac{1}{2}(c + dx) \middle| 2 \right) \right)}{bd\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(-3/2), x]

[Out] (2*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(b*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.98, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \cos(dx + c)}}{b^2 \cos(dx + c)^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.16, size = 168, normalized size = 2.47

$$\frac{2\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b} \left(\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1 \right)}{b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(d*x+c))^(3/2), x)

[Out] -2/b*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)*((sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(3/2), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(b \cos(c + dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(c + d*x))^(3/2), x)

[Out] int(1/(b*cos(c + d*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))**(3/2), x)

[Out] Integral((b*cos(c + d*x))**(-3/2), x)

$$3.123 \quad \int \frac{\sec(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=69

$$\frac{2 \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3bd\sqrt{b \cos(c+dx)}}$$

[Out] 2/3*sin(d*x+c)/d/(b*cos(d*x+c))^(3/2)+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.05, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2 \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(3/2),x]

[Out] (2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(3*b*d*sqrt[b*Cos[c + d*x]]) + (2*Sin[c + d*x])/(3*d*(b*Cos[c + d*x])^(3/2))

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec(c+dx)}{(b\cos(c+dx))^{3/2}} dx &= b \int \frac{1}{(b\cos(c+dx))^{5/2}} dx \\
&= \frac{2\sin(c+dx)}{3d(b\cos(c+dx))^{3/2}} + \frac{\int \frac{1}{\sqrt{b\cos(c+dx)}} dx}{3b} \\
&= \frac{2\sin(c+dx)}{3d(b\cos(c+dx))^{3/2}} + \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b\sqrt{b\cos(c+dx)}} \\
&= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3bd\sqrt{b\cos(c+dx)}} + \frac{2\sin(c+dx)}{3d(b\cos(c+dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 51, normalized size = 0.74

$$\frac{2\left(\tan(c+dx) + \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)\right)}{3bd\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(3/2), x]

[Out] (2*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*b*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)}\sec(dx+c)}{b^2\cos(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b\cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.16, size = 241, normalized size = 3.49

$$\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\text{EllipticF}\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2}\right)}{3b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*cos(d*x+c))^(3/2), x)


```
[0ut] -2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/b*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")
```

```
[0ut] integrate(sec(d*x + c)/(b*cos(d*x + c))^(3/2), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx) (b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(3/2)),x)
```

```
[0ut] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(3/2)), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(3/2),x)
```

```
[0ut] Integral(sec(c + d*x)/(b*cos(c + d*x))**(3/2), x)
```

$$3.124 \quad \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=98

$$-\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{5b^2d\sqrt{\cos(c+dx)}} + \frac{6\sin(c+dx)}{5bd\sqrt{b\cos(c+dx)}} + \frac{2b\sin(c+dx)}{5d(b\cos(c+dx))^{5/2}}$$

[Out] $2/5*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^2/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.08, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$-\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{5b^2d\sqrt{\cos(c+dx)}} + \frac{6\sin(c+dx)}{5bd\sqrt{b\cos(c+dx)}} + \frac{2b\sin(c+dx)}{5d(b\cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Cos[c + d*x])^(3/2), x]

[Out] $(-6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*\text{Sin}[c + d*x])/(5*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= b^2 \int \frac{1}{(b \cos(c+dx))^{7/2}} dx \\
&= \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{3}{5} \int \frac{1}{(b \cos(c+dx))^{3/2}} dx \\
&= \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5bd\sqrt{b \cos(c+dx)}} - \frac{3 \int \sqrt{b \cos(c+dx)} dx}{5b^2} \\
&= \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5bd\sqrt{b \cos(c+dx)}} - \frac{(3\sqrt{b \cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5b^2\sqrt{\cos(c+dx)}} \\
&= -\frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5b^2d\sqrt{\cos(c+dx)}} + \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5bd\sqrt{b \cos(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 68, normalized size = 0.69

$$\frac{6 \sin(c+dx) - 6\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2 \tan(c+dx) \sec(c+dx)}{5bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(3/2), x]

[Out] (-6*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 6*Sin[c + d*x] + 2*Sec[c + d*x]*Tan[c + d*x])/(5*b*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.76, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \sec(dx+c)^2}{b^2 \cos(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^2/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \cos(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.26, size = 366, normalized size = 3.73

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(12 \text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{\frac{1}{2} - \dots}}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2/(b*cos(d*x+c))^(3/2),x)

[Out] $2/5*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2/\sin(1/2*d*x+1/2*c)^3/(8*\sin(1/2*d*x+1/2*c)^6-12*\sin(1/2*d*x+1/2*c)^4+6*\sin(1/2*d*x+1/2*c)^2-1)*(12*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*\sin(1/2*d*x+1/2*c)^4-24*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6-12*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*\sin(1/2*d*x+1/2*c)^2+24*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-8*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))*(-2*\sin(1/2*d*x+1/2*c)^4*b+\sin(1/2*d*x+1/2*c)^2*b)^{(1/2)}/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^2 (b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(3/2)),x)

[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(sec(c + d*x)**2/(b*cos(c + d*x))**(3/2), x)

$$3.125 \quad \int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=97

$$\frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21bd\sqrt{b \cos(c+dx)}}$$

[Out] $2/7*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3/(b*Cos[c + d*x])^(3/2), x]

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b^2*\text{Sin}[c + d*x])/(7*d*(b*\text{Cos}[c + d*x])^{(7/2)}) + (10*\text{Sin}[c + d*x])/(21*d*(b*\text{Cos}[c + d*x])^{(3/2)})$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*SIN[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= b^3 \int \frac{1}{(b \cos(c+dx))^{9/2}} dx \\
&= \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{1}{7}(5b) \int \frac{1}{(b \cos(c+dx))^{5/2}} dx \\
&= \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{5 \int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{21b} \\
&= \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{(5\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21b\sqrt{b \cos(c+dx)}} \\
&= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21bd\sqrt{b \cos(c+dx)}} + \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 66, normalized size = 0.68

$$\frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2 \tan(c+dx) (3 \sec^2(c+dx) + 5)}{21bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(3/2), x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 2*(5 + 3*Sec[c + d*x]^2)*Tan[c + d*x])/(21*b*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \sec(dx+c)^3}{b^2 \cos(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^3/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.18, size = 398, normalized size = 4.10

$$\frac{2\left(-40\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \text{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 60\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\right)}{21bd\sqrt{b \cos(c+dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^3/(b*cos(d*x+c))^(3/2),x)

[Out]
$$-2/21*(-40*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^6+60*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^4-40*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6-30*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2+40*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-16*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/b*(b*(2*\cos(1/2*d*x+1/2*c)^{2-1})*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^{2-1})^3/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^{2-1}))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^3 (b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(3/2)),x)

[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(sec(c + d*x)**3/(b*cos(c + d*x))**(3/2), x)

$$3.126 \quad \int \frac{\sec^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=126

$$\frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} - \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15b^2 d \sqrt{\cos(c+dx)}} + \frac{14b \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15bd \sqrt{b \cos(c+dx)}}$$

[Out] $2/9*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^(1/2)-14/15*(\cos(1/2*d*x+1/2*c))^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^(1/2))*(b*\cos(d*x+c))^(1/2)/b^2/d/\cos(d*x+c)^(1/2)$

Rubi [A] time = 0.10, antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} - \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15b^2 d \sqrt{\cos(c+dx)}} + \frac{14b \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15bd \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^4/(b*Cos[c + d*x])^(3/2), x]`

[Out] $(-14*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^3*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^(9/2)) + (14*b*\text{Sin}[c + d*x])/(45*d*(b*\text{Cos}[c + d*x])^(5/2)) + (14*\text{Sin}[c + d*x])/(15*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2636

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*SIN[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2639

`Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 2640

`Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*SIN[c + d*x]]/Sqrt[SIN[c + d*x]], Int[Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \frac{\sec^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= b^4 \int \frac{1}{(b \cos(c+dx))^{11/2}} dx \\
&= \frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{1}{9} (7b^2) \int \frac{1}{(b \cos(c+dx))^{7/2}} dx \\
&= \frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{7}{15} \int \frac{1}{(b \cos(c+dx))^{3/2}} dx \\
&= \frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15bd\sqrt{b \cos(c+dx)}} - \frac{7 \int \sqrt{b \cos(c+dx)}}{15b^2} \\
&= \frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15bd\sqrt{b \cos(c+dx)}} - \frac{(7\sqrt{b \cos(c+dx)})}{15b^2} \\
&= -\frac{14\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15b^2 d \sqrt{\cos(c+dx)}} + \frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 80, normalized size = 0.63

$$\frac{42 \sin(c+dx) - 42\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2 \tan(c+dx) \sec(c+dx) (5 \sec^2(c+dx) + 7)}{45bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^4/(b*Cos[c + d*x])^(3/2), x]

[Out] (-42*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 42*Sin[c + d*x] + 2*Sec[c + d*x]*(7 + 5*Sec[c + d*x]^2)*Tan[c + d*x])/(45*b*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \sec(dx+c)^4}{b^2 \cos(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^4/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^4}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^4/(b*cos(d*x + c))^(3/2), x)

maple [B] time = 0.26, size = 414, normalized size = 3.29

$$2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \left(-\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{144b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} - \frac{7\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{180b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^4/(b*cos(d*x+c))^(3/2),x)`

[Out]
$$-2*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b*(-1/144*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^5-7/180*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^3-14/15*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*(\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^4}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^4/(b*cos(d*x + c))^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^4 (b \cos(c+dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c+d*x)^4*(b*cos(c+d*x))^(3/2)),x)`

[Out] `int(1/(cos(c+d*x)^4*(b*cos(c+d*x))^(3/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**4/(b*cos(d*x+c))**(3/2),x)`

[Out] `Integral(sec(c+d*x)**4/(b*cos(c+d*x))**(3/2), x)`

$$3.127 \quad \int \frac{\cos^8(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=128

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^7d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^5d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77b^3d} + \frac{30\sqrt{\cos(c+dx)}}{77b^2d}$$

[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b^5/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^7/d+30/77*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/b^2/d/(b*cos(d*x+c))^(1/2)+30/77*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b^3/d

Rubi [A] time = 0.08, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^7d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^5d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77b^3d} + \frac{30\sqrt{\cos(c+dx)}}{77b^2d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^8/(b*Cos[c + d*x])^(5/2), x]

[Out] (30*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*b^2*d*sqrt[b*Cos[c + d*x]]) + (30*sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*b^3*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b^5*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^7*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Ssin[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*Ssin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[Sin[c + d*x]]/sqrt[b*Ssin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^8(c+dx)}{(b\cos(c+dx))^{5/2}} dx &= \frac{\int (b\cos(c+dx))^{11/2} dx}{b^8} \\
&= \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^7d} + \frac{9 \int (b\cos(c+dx))^{7/2} dx}{11b^6} \\
&= \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^5d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^7d} + \frac{45 \int (b\cos(c+dx))^{3/2} dx}{77b^4} \\
&= \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77b^3d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^5d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^7d} \\
&= \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77b^3d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^5d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^7d} \\
&= \frac{30\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{77b^2d\sqrt{b\cos(c+dx)}} + \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77b^3d} + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^5d}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 76, normalized size = 0.59

$$\frac{347 \sin(2(c+dx)) + 64 \sin(4(c+dx)) + 7 \sin(6(c+dx)) + 480\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{1232b^2d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^8/(b*Cos[c + d*x])^(5/2), x]

[Out] (480*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 347*Sin[2*(c + d*x)] + 64*Sin[4*(c + d*x)] + 7*Sin[6*(c + d*x)])/(1232*b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^5}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^8/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^5/b^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^8}{(b\cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^8/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^8/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.19, size = 236, normalized size = 1.84

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(448\left(\cos^{13}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1568\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 2384\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 224\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 16\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 2\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{77b^2\sqrt{-b}\left(2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^8/(b*cos(d*x+c))^(5/2),x)`

[Out]
$$-2/77*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(448*\cos(1/2*d*x+1/2*c)^{13}-1568*\cos(1/2*d*x+1/2*c)^{11}+2384*\cos(1/2*d*x+1/2*c)^9-2040*\cos(1/2*d*x+1/2*c)^7+1084*\cos(1/2*d*x+1/2*c)^5-370*\cos(1/2*d*x+1/2*c)^3+15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+62*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^8}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^8/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^8/(b*cos(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^8}{(b \cos(c+dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)^8/(b*cos(c+d*x))^(5/2),x)`

[Out] `int(cos(c+d*x)^8/(b*cos(c+d*x))^(5/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**8/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

$$3.128 \quad \int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^6d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^4d} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15b^3d\sqrt{\cos(c+dx)}}$$

[Out] 14/45*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b^4/d+2/9*(b*cos(d*x+c))^(7/2)*sin(d*x+c)/b^6/d+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^3/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^6d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^4d} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{15b^3d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^7/(b*Cos[c + d*x])^(5/2), x]

[Out] (14*sqrt[b*cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*b^3*d*sqrt[Cos[c + d*x]]) + (14*(b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(45*b^4*d) + (2*(b*cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b^6*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*cos[c + d*x])*(b*sin[c + d*x])^(n-1)/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_)+(d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*sin[c + d*x]]/sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{5/2}} dx &= \frac{\int (b\cos(c+dx))^{9/2} dx}{b^7} \\
&= \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^6d} + \frac{7 \int (b\cos(c+dx))^{5/2} dx}{9b^5} \\
&= \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^4d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^6d} + \frac{7 \int \sqrt{b\cos(c+dx)} dx}{15b^3} \\
&= \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^4d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^6d} + \frac{(7\sqrt{b\cos(c+dx)}) E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15b^3} \\
&= \frac{14\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15b^3d\sqrt{\cos(c+dx)}} + \frac{14(b\cos(c+dx))^{3/2} \sin(c+dx)}{45b^4d} + \frac{2(b\cos(c+dx))^{7/2} \sin(c+dx)}{9b^6d}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 74, normalized size = 0.74

$$\frac{(38 \sin(2(c+dx)) + 5 \sin(4(c+dx))) \cos(c+dx) + 168 \sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{180b^2d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^7/(b*Cos[c + d*x])^(5/2), x]

[Out] (168*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)]))/(180*b^2*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.47, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^4}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^4/b^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^7}{(b\cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^7/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.18, size = 223, normalized size = 2.23

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(160\left(\cos^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 480\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 616\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 280\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 80\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 8\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{45b^2\sqrt{-b}\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^7/(b*cos(d*x+c))^(5/2),x)`

[Out]
$$-2/45*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(160*\cos(1/2*d*x+1/2*c)^{11}-480*\cos(1/2*d*x+1/2*c)^9+616*\cos(1/2*d*x+1/2*c)^7-432*\cos(1/2*d*x+1/2*c)^5+160*\cos(1/2*d*x+1/2*c)^3-21*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-24*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^7}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^7/(b*cos(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^7}{(b \cos(c+dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)^7/(b*cos(c+d*x))^(5/2),x)`

[Out] `int(cos(c+d*x)^7/(b*cos(c+d*x))^(5/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**7/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

$$3.129 \quad \int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^5d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21b^3d} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx)\middle|2\right)}{21b^2d\sqrt{b \cos(c+dx)}}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b^5/d+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}+10/21*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/b^3/d$

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^5d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21b^3d} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx)\middle|2\right)}{21b^2d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^6/(b*Cos[c + d*x])^(5/2),x]

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (10*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(21*b^3*d) + (2*(b*\text{Cos}[c + d*x])^{(5/2)}*\text{Sin}[c + d*x])/(7*b^5*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*SIN[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^6(c+dx)}{(b\cos(c+dx))^{5/2}} dx &= \frac{\int (b\cos(c+dx))^{7/2} dx}{b^6} \\
&= \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^5d} + \frac{5 \int (b\cos(c+dx))^{3/2} dx}{7b^4} \\
&= \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21b^3d} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^5d} + \frac{5 \int \frac{1}{\sqrt{b\cos(c+dx)}} dx}{21b^2} \\
&= \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21b^3d} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^5d} + \frac{(5\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21b^2\sqrt{b\cos(c+dx)}} \\
&= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21b^2d\sqrt{b\cos(c+dx)}} + \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21b^3d} + \frac{2(b\cos(c+dx))^{5/2}}{7b^5d}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 66, normalized size = 0.66

$$\frac{26 \sin(2(c+dx)) + 3 \sin(4(c+dx)) + 40\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{84b^2d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^6/(b*Cos[c + d*x])^(5/2), x]

[Out] (40*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)])/(84*b^2*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^3}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^3/b^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^6}{(b\cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^6/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.18, size = 210, normalized size = 2.10

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{21b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^6/(b*cos(d*x+c))^(5/2),x)`

[Out]
$$-2/21*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(48*\cos(1/2*d*x+1/2*c)^9-120*\cos(1/2*d*x+1/2*c)^7+128*\cos(1/2*d*x+1/2*c)^5-72*\cos(1/2*d*x+1/2*c)^3+5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+16*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^6}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^6/(b*cos(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^6}{(b \cos(c+dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)^6/(b*cos(c+d*x))^(5/2),x)`

[Out] `int(cos(c+d*x)^6/(b*cos(c+d*x))^(5/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**6/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

$$3.130 \quad \int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^4d} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5b^3d\sqrt{\cos(c+dx)}}$$

[Out] 2/5*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b^4/d+6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^3/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2640, 2639}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^4d} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5b^3d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^5/(b*Cos[c + d*x])^(5/2), x]

[Out] (6*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(5*b^3*d*Sqrt[Cos[c + d*x]]) + (2*(b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*b^4*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*SIN[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*SIN[c + d*x]]/Sqrt[SIN[c + d*x]], Int[Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^5(c+dx)}{(b\cos(c+dx))^{5/2}} dx &= \frac{\int (b\cos(c+dx))^{5/2} dx}{b^5} \\
&= \frac{2(b\cos(c+dx))^{3/2} \sin(c+dx)}{5b^4d} + \frac{3 \int \sqrt{b\cos(c+dx)} dx}{5b^3} \\
&= \frac{2(b\cos(c+dx))^{3/2} \sin(c+dx)}{5b^4d} + \frac{(3\sqrt{b\cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5b^3\sqrt{\cos(c+dx)}} \\
&= \frac{6\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5b^3d\sqrt{\cos(c+dx)}} + \frac{2(b\cos(c+dx))^{3/2} \sin(c+dx)}{5b^4d}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 61, normalized size = 0.85

$$\frac{\sin(2(c+dx)) \cos(c+dx) + 6\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5b^2d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^5/(b*Cos[c + d*x])^(5/2), x]

[Out] (6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*Sin[2*(c + d*x)])/(5*b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)} \cos(dx+c)^2}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)^2/b^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^5}{(b\cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^5/(b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.16, size = 213, normalized size = 2.96

$$\frac{2\sqrt{b}\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(-8\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5b^2\sqrt{-b}\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^5/(b*cos(d*x+c))^(5/2), x)

[Out]
$$-2/5*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(-8*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+8*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{Elliptic E}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^5}{(b \cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^5/(b*cos(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^5}{(b \cos(c+dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^5/(b*cos(c + d*x))^(5/2),x)`

[Out] `int(cos(c + d*x)^5/(b*cos(c + d*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**5/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

$$3.131 \quad \int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3b^3d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3b^2d\sqrt{b \cos(c+dx)}}$$

[Out] 2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/b^2/d/(b*cos(d*x+c))^(1/2)+2/3*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b^3/d

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2635, 2642, 2641}

$$\frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3b^3d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3b^2d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^4/(b*Cos[c + d*x])^(5/2),x]

[Out] (2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(3*b^2*d*sqrt[b*Cos[c + d*x]]) + (2*sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(3*b^3*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n-1))/(d*n), x] + Dist[(b^2*(n-1))/n, Int[(b*SIN[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= \frac{\int (b \cos(c+dx))^{3/2} dx}{b^4} \\
&= \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3b^3 d} + \frac{\int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{3b^2} \\
&= \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3b^3 d} + \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2 \sqrt{b \cos(c+dx)}} \\
&= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}} + \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3b^3 d}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 54, normalized size = 0.75

$$\frac{\sin(2(c+dx)) + 2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^4/(b*Cos[c + d*x])^(5/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)])/(3*b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \cos(dx+c)}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*cos(d*x + c)/b^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^4}{(b \cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^4/(b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.16, size = 190, normalized size = 2.64

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(4\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) + \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{3b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^4/(b*cos(d*x+c))^(5/2), x)

[Out]
$$-2/3*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(4*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^4}{(b \cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^4/(b*cos(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)^4}{(b \cos(c+dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^4/(b*cos(c + d*x))^(5/2),x)`

[Out] `int(cos(c + d*x)^4/(b*cos(c + d*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**4/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

$$3.132 \quad \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=41

$$\frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^3 d \sqrt{\cos(c+dx)}}$$

[Out] 2*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))*(b*cos(d*x+c))^(1/2)/b^3/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2640, 2639}

$$\frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^3 d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^3/(b*Cos[c + d*x])^(5/2), x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b^3*d*Sqrt[Cos[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= \frac{\int \sqrt{b \cos(c+dx)} dx}{b^3} \\ &= \frac{\sqrt{b \cos(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b^3 \sqrt{\cos(c+dx)}} \\ &= \frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^3 d \sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 41, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^3 d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3/(b*Cos[c + d*x])^(5/2), x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b^3*d*Sqrt[Cos[c + d*x]])

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx + c)}}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/b^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^3}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^3/(b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.15, size = 144, normalized size = 3.51

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}\text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^3/(b*cos(d*x+c))^(5/2), x)

[Out] 2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))/b^2/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^3}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(5/2), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^3/(b*cos(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)^3}{(b \cos(c + dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^3/(b*cos(c + d*x))^(5/2),x)
```

```
[Out] int(cos(c + d*x)^3/(b*cos(c + d*x))^(5/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**3/(b*cos(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

$$3.133 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=41

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2642, 2641}

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/(b*Cos[c + d*x])^(5/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(b^2*d*Sqrt[b*Cos[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[Sin[c + d*x]]/Sqrt[b*Sin[c + d*x]], Int[1/Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= \frac{\int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{b^2} \\ &= \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^2 d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 38, normalized size = 0.93

$$\frac{2 \cos^{\frac{5}{2}}(c+dx) F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d(b \cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(5/2), x]

[Out] (2*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2])/(d*(b*Cos[c + d*x])^(5/2))

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx + c)}}{b^3 \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/(b^3*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.14, size = 144, normalized size = 3.51

$$\frac{2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}\text{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2/(b*cos(d*x+c))^(5/2), x)

[Out] -2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(5/2), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(5/2), x)
```

```
[Out] int(cos(c + d*x)^2/(b*cos(c + d*x))^(5/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(5/2), x)
```

```
[Out] Timed out
```

$$3.134 \quad \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=68

$$\frac{2 \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^3 d \sqrt{\cos(c+dx)}}$$

[Out] $2*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}-2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^3/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2 \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^3 d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/(b*Cos[c + d*x])^(5/2), x]

[Out] $(-2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(b^3*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\cos(c+dx)}{(b\cos(c+dx))^{5/2}} dx &= \frac{\int \frac{1}{(b\cos(c+dx))^{3/2}} dx}{b} \\
&= \frac{2\sin(c+dx)}{b^2 d \sqrt{b\cos(c+dx)}} - \frac{\int \sqrt{b\cos(c+dx)} dx}{b^3} \\
&= \frac{2\sin(c+dx)}{b^2 d \sqrt{b\cos(c+dx)}} - \frac{\sqrt{b\cos(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b^3 \sqrt{\cos(c+dx)}} \\
&= -\frac{2\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^3 d \sqrt{\cos(c+dx)}} + \frac{2\sin(c+dx)}{b^2 d \sqrt{b\cos(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 50, normalized size = 0.74

$$\frac{2\left(\sin(c+dx) - \sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)\right)}{b^2 d \sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(5/2), x]

[Out] (2*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)}}{b^3\cos(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/(b^3*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)}{(b\cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.18, size = 168, normalized size = 2.47

$$\frac{2\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b\left(\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1\right)\text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*cos(d*x+c))^(5/2), x)

[Out]
$$-2/b^2*(-2*\sin(1/2*d*x+1/2*c)^4*b+\sin(1/2*d*x+1/2*c)^2*b)^{(1/2)}*((\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2)^{(1/2)})-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)}{(b \cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)/(b*cos(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c+d*x)/(b*cos(c+d*x))^(5/2),x)`

[Out] `int(cos(c+d*x)/(b*cos(c+d*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

$$3.135 \quad \int \frac{1}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}} + \frac{2 \sin(c+dx)}{3bd(b \cos(c+dx))^{3/2}}$$

[Out] $2/3*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(3/2)}+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2636, 2642, 2641}

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}} + \frac{2 \sin(c+dx)}{3bd(b \cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(-5/2)}, x]$

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(3*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*\text{Sin}[c + d*x])/(3*b*d*(b*\text{Cos}[c + d*x])^{(3/2)})$

Rule 2636

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x])*(b*\text{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)), x] + \text{Dist}[(n + 2)/(b^2*(n + 1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n + 2)}, x], x] /;$ $\text{FreeQ}\{b, c, d\}, x$ && $\text{LtQ}[n, -1]$ && $\text{IntegerQ}[2*n]$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /;$ $\text{FreeQ}\{c, d\}, x$

Rule 2642

$\text{Int}[1/\text{Sqrt}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[c + d*x]]/\text{Sqrt}[b*\text{Sin}[c + d*x]], \text{Int}[1/\text{Sqrt}[\text{Sin}[c + d*x]], x], x] /;$ $\text{FreeQ}\{b, c, d\}, x$

Rubi steps

$$\begin{aligned} \int \frac{1}{(b \cos(c+dx))^{5/2}} dx &= \frac{2 \sin(c+dx)}{3bd(b \cos(c+dx))^{3/2}} + \frac{\int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{3b^2} \\ &= \frac{2 \sin(c+dx)}{3bd(b \cos(c+dx))^{3/2}} + \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}} + \frac{2 \sin(c+dx)}{3bd(b \cos(c+dx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 51, normalized size = 0.71

$$\frac{2 \left(\tan(c + dx) + \sqrt{\cos(c + dx)} F \left(\frac{1}{2}(c + dx) \middle| 2 \right) \right)}{3b^2 d \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(-5/2), x]

[Out] (2*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.86, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \cos(dx + c)}}{b^3 \cos(dx + c)^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/(b^3*cos(d*x + c)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.17, size = 241, normalized size = 3.35

$$\frac{2 \left(-2 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \right)}{3b^2 \sqrt{-b \left(2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(d*x+c))^(5/2), x)

[Out] -2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))/b^2*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(5/2), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(-5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(b \cos(c + dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(c + d*x))^(5/2), x)

[Out] int(1/(b*cos(c + d*x))^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(c + dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(5/2), x)

[Out] Integral((b*cos(c + d*x))^(-5/2), x)

$$3.136 \quad \int \frac{\sec(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=97

$$-\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{5b^3d\sqrt{\cos(c+dx)}} + \frac{6\sin(c+dx)}{5b^2d\sqrt{b\cos(c+dx)}} + \frac{2\sin(c+dx)}{5d(b\cos(c+dx))^{5/2}}$$

[Out] $2/5*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^3/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 2636, 2640, 2639}

$$\frac{6\sin(c+dx)}{5b^2d\sqrt{b\cos(c+dx)}} - \frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{5b^3d\sqrt{\cos(c+dx)}} + \frac{2\sin(c+dx)}{5d(b\cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(5/2), x]

[Out] $(-6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b^3*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*\text{Sin}[c + d*x])/(5*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_)*sin[(c_)+(d_)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec(c+dx)}{(b\cos(c+dx))^{5/2}} dx &= b \int \frac{1}{(b\cos(c+dx))^{7/2}} dx \\
&= \frac{2\sin(c+dx)}{5d(b\cos(c+dx))^{5/2}} + \frac{3 \int \frac{1}{(b\cos(c+dx))^{3/2}} dx}{5b} \\
&= \frac{2\sin(c+dx)}{5d(b\cos(c+dx))^{5/2}} + \frac{6\sin(c+dx)}{5b^2d\sqrt{b\cos(c+dx)}} - \frac{3 \int \sqrt{b\cos(c+dx)} dx}{5b^3} \\
&= \frac{2\sin(c+dx)}{5d(b\cos(c+dx))^{5/2}} + \frac{6\sin(c+dx)}{5b^2d\sqrt{b\cos(c+dx)}} - \frac{(3\sqrt{b\cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5b^3\sqrt{\cos(c+dx)}} \\
&= -\frac{6\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5b^3d\sqrt{\cos(c+dx)}} + \frac{2\sin(c+dx)}{5d(b\cos(c+dx))^{5/2}} + \frac{6\sin(c+dx)}{5b^2d\sqrt{b\cos(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 68, normalized size = 0.70

$$\frac{6\sin(c+dx) - 6\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2\tan(c+dx)\sec(c+dx)}{5b^2d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(5/2), x]

[Out] (-6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 6*Sin[c + d*x] + 2*Sec[c + d*x]*Tan[c + d*x])/(5*b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\cos(dx+c)}\sec(dx+c)}{b^3\cos(dx+c)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)/(b^3*cos(d*x + c)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b\cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.27, size = 366, normalized size = 3.77

$$2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(12\text{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\sqrt{\frac{1}{2}} - \dots\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*cos(d*x+c))^(5/2),x)

[Out] $\frac{2}{5} \cdot (b \cdot (2 \cdot \cos(1/2 \cdot d \cdot x + 1/2 \cdot c))^2 - 1) \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2)^{1/2} / b^3 / \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 / (8 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^6 - 12 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^4 + 6 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 - 1) \cdot (12 \cdot \text{EllipticE}(\cos(1/2 \cdot d \cdot x + 1/2 \cdot c), 2^{1/2})) \cdot (2 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 - 1)^{1/2} \cdot (\sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2)^{1/2} \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^4 - 24 \cdot \cos(1/2 \cdot d \cdot x + 1/2 \cdot c) \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^6 - 12 \cdot \text{EllipticE}(\cos(1/2 \cdot d \cdot x + 1/2 \cdot c), 2^{1/2})) \cdot (2 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 - 1)^{1/2} \cdot (\sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2)^{1/2} \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 + 24 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^4 \cdot \cos(1/2 \cdot d \cdot x + 1/2 \cdot c) + 3 \cdot (\sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2)^{1/2} \cdot (2 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 - 1)^{1/2} \cdot \text{EllipticE}(\cos(1/2 \cdot d \cdot x + 1/2 \cdot c), 2^{1/2})) - 8 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 \cdot \cos(1/2 \cdot d \cdot x + 1/2 \cdot c) \cdot (-2 \cdot \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^4 \cdot b + \sin(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 \cdot b)^{1/2} / (b \cdot (2 \cdot \cos(1/2 \cdot d \cdot x + 1/2 \cdot c))^2 - 1)^{1/2} / d$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx) (b \cos(c+dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c+d*x)*(b*cos(c+d*x))^(5/2)),x)

[Out] int(1/(cos(c+d*x)*(b*cos(c+d*x))^(5/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

$$3.137 \quad \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=98

$$\frac{10\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{21b^2d\sqrt{b\cos(c+dx)}} + \frac{10\sin(c+dx)}{21bd(b\cos(c+dx))^{3/2}} + \frac{2b\sin(c+dx)}{7d(b\cos(c+dx))^{7/2}}$$

[Out] $2/7*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(3/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.08, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2642, 2641}

$$\frac{10\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)}{21b^2d\sqrt{b\cos(c+dx)}} + \frac{10\sin(c+dx)}{21bd(b\cos(c+dx))^{3/2}} + \frac{2b\sin(c+dx)}{7d(b\cos(c+dx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Cos[c + d*x])^(5/2), x]

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(21*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (2*b*\text{Sin}[c + d*x])/(7*d*(b*\text{Cos}[c + d*x])^{(7/2)}) + (10*\text{Sin}[c + d*x])/(21*b*d*(b*\text{Cos}[c + d*x])^{(3/2)})$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1))/(b*d*(n+1)), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*SIN[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2642

Int[1/Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[Sqrt[SIN[c + d*x]]/Sqrt[b*SIN[c + d*x]], Int[1/Sqrt[SIN[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= b^2 \int \frac{1}{(b \cos(c+dx))^{9/2}} dx \\
&= \frac{2b \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{5}{7} \int \frac{1}{(b \cos(c+dx))^{5/2}} dx \\
&= \frac{2b \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21bd(b \cos(c+dx))^{3/2}} + \frac{5 \int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{21b^2} \\
&= \frac{2b \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21bd(b \cos(c+dx))^{3/2}} + \frac{(5\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21b^2 \sqrt{b \cos(c+dx)}} \\
&= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right)}{21b^2 d \sqrt{b \cos(c+dx)}} + \frac{2b \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21bd(b \cos(c+dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 66, normalized size = 0.67

$$\frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2 \tan(c+dx) (3 \sec^2(c+dx) + 5)}{21b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(5/2), x]

[Out] (10*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 2*(5 + 3*Sec[c + d*x]^2)*Tan[c + d*x])/(21*b^2*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \sec(dx+c)^2}{b^3 \cos(dx+c)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^2/(b^3*cos(d*x + c)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.19, size = 398, normalized size = 4.06

$$\frac{2 \left(-40 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} \text{EllipticF} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 60 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right)}{21b^2 d \sqrt{b \cos(c+dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2/(b*cos(d*x+c))^(5/2),x)

[Out]
$$-2/21*(-40*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^6+60*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^4-40*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6-30*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2+40*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-16*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c))/b^2*(b*(2*\cos(1/2*d*x+1/2*c)^{2-1})*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^{2-1})^3/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^{2-1}))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^2 (b \cos(c+dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(5/2)),x)

[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(5/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

$$3.138 \quad \int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=125

$$-\frac{14E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{15b^3d\sqrt{\cos(c+dx)}} + \frac{2b^2\sin(c+dx)}{9d(b\cos(c+dx))^{9/2}} + \frac{14\sin(c+dx)}{15b^2d\sqrt{b\cos(c+dx)}} + \frac{14\sin(c+dx)}{45d(b\cos(c+dx))^{5/2}}$$

[Out] $2/9*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^(1/2)-14/15*(\cos(1/2*d*x+1/2*c))^2^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/b^3/d/\cos(d*x+c)^(1/2)$

Rubi [A] time = 0.10, antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2636, 2640, 2639}

$$\frac{2b^2\sin(c+dx)}{9d(b\cos(c+dx))^{9/2}} + \frac{14\sin(c+dx)}{15b^2d\sqrt{b\cos(c+dx)}} - \frac{14E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{15b^3d\sqrt{\cos(c+dx)}} + \frac{14\sin(c+dx)}{45d(b\cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^3/(b*Cos[c + d*x])^(5/2),x]`

[Out] $(-14*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*b^3*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^2*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^(9/2)) + (14*\text{Sin}[c + d*x])/(45*d*(b*\text{Cos}[c + d*x])^(5/2)) + (14*\text{Sin}[c + d*x])/(15*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2636

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2639

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 2640

`Int[Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= b^3 \int \frac{1}{(b \cos(c+dx))^{11/2}} dx \\
&= \frac{2b^2 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{1}{9}(7b) \int \frac{1}{(b \cos(c+dx))^{7/2}} dx \\
&= \frac{2b^2 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{7 \int \frac{1}{(b \cos(c+dx))^{3/2}} dx}{15b} \\
&= \frac{2b^2 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15b^2 d \sqrt{b \cos(c+dx)}} - \frac{7 \int \sqrt{b \cos(c+dx)} dx}{15b} \\
&= \frac{2b^2 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15b^2 d \sqrt{b \cos(c+dx)}} - \frac{(7 \sqrt{b \cos(c+dx)})}{15b} \\
&= -\frac{14 \sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15b^3 d \sqrt{\cos(c+dx)}} + \frac{2b^2 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 80, normalized size = 0.64

$$\frac{42 \sin(c+dx) - 42 \sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) + 2 \tan(c+dx) \sec(c+dx) (5 \sec^2(c+dx) + 7)}{45b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(5/2), x]

[Out] (-42*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 42*Sin[c + d*x] + 2*Sec[c + d*x]*(7 + 5*Sec[c + d*x]^2)*Tan[c + d*x])/(45*b^2*d*sqrt[b*Cos[c + d*x]])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx+c)} \sec(dx+c)^3}{b^3 \cos(dx+c)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))*sec(d*x + c)^3/(b^3*cos(d*x + c)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{(b \cos(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(5/2), x)

maple [B] time = 0.29, size = 414, normalized size = 3.31

$$2\sqrt{b\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \left(-\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{144b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} - \frac{7\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{180b\left(-\frac{1}{2} + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^3/(b*cos(d*x+c))^(5/2),x)`

[Out]
$$-2*(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(-1/144*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^5-7/180*\cos(1/2*d*x+1/2*c)/b*(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(-1/2+\cos(1/2*d*x+1/2*c)^2)^3-14/15*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-7/15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*(EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c)/(b*(2*\cos(1/2*d*x+1/2*c)^2-1))^{(1/2)}/d$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^3 (b \cos(c+dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(5/2)),x)`

[Out] `int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(5/2)), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

$$3.139 \quad \int \frac{1}{(b \cos(c+dx))^{7/2}} dx$$

Optimal. Leaf size=100

$$-\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{5b^4d\sqrt{\cos(c+dx)}} + \frac{6\sin(c+dx)}{5b^3d\sqrt{b\cos(c+dx)}} + \frac{2\sin(c+dx)}{5bd(b\cos(c+dx))^{5/2}}$$

[Out] 2/5*sin(d*x+c)/b/d/(b*cos(d*x+c))^(5/2)+6/5*sin(d*x+c)/b^3/d/(b*cos(d*x+c))^(1/2)-6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^4/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2636, 2640, 2639}

$$\frac{6\sin(c+dx)}{5b^3d\sqrt{b\cos(c+dx)}} - \frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\cos(c+dx)}}{5b^4d\sqrt{\cos(c+dx)}} + \frac{2\sin(c+dx)}{5bd(b\cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(-7/2), x]

[Out] (-6*Sqrt[b*cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(5*b^4*d*Sqrt[Cos[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*cos[c + d*x])^(5/2)) + (6*Sin[c + d*x])/(5*b^3*d*Sqrt[b*cos[c + d*x]])

Rule 2636

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1))/(b*d*(n + 1)), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 2640

Int[Sqrt[(b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[Sqrt[b*Sin[c + d*x]]/Sqrt[Sin[c + d*x]], Int[Sqrt[Sin[c + d*x]], x], x] /; FreeQ[{b, c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(b \cos(c+dx))^{7/2}} dx &= \frac{2\sin(c+dx)}{5bd(b\cos(c+dx))^{5/2}} + \frac{3 \int \frac{1}{(b \cos(c+dx))^{3/2}} dx}{5b^2} \\ &= \frac{2\sin(c+dx)}{5bd(b\cos(c+dx))^{5/2}} + \frac{6\sin(c+dx)}{5b^3d\sqrt{b\cos(c+dx)}} - \frac{3 \int \sqrt{b\cos(c+dx)} dx}{5b^4} \\ &= \frac{2\sin(c+dx)}{5bd(b\cos(c+dx))^{5/2}} + \frac{6\sin(c+dx)}{5b^3d\sqrt{b\cos(c+dx)}} - \frac{(3\sqrt{b\cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5b^4\sqrt{\cos(c+dx)}} \\ &= -\frac{6\sqrt{b\cos(c+dx)} E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5b^4d\sqrt{\cos(c+dx)}} + \frac{2\sin(c+dx)}{5bd(b\cos(c+dx))^{5/2}} + \frac{6\sin(c+dx)}{5b^3d\sqrt{b\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 68, normalized size = 0.68

$$\frac{6 \sin(c + dx) - 6\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) + 2 \tan(c + dx) \sec(c + dx)}{5b^3 d \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(-7/2), x]

[Out] (-6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 6*Sin[c + d*x] + 2*Sec[c + d*x]*Tan[c + d*x])/(5*b^3*d*Sqrt[b*cos[c + d*x]])

fricas [F] time = 0.81, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \cos(dx + c)}}{b^4 \cos(dx + c)^4}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(7/2), x, algorithm="fricas")

[Out] integral(sqrt(b*cos(d*x + c))/(b^4*cos(d*x + c)^4), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(7/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(7/2), x)

maple [B] time = 0.26, size = 366, normalized size = 3.66

$$2\sqrt{b} \left(2 \left(\cos^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \left(12 \text{EllipticE} \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \sqrt{2} \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1 \right) \sqrt{\frac{1}{2} - \cos \left(\frac{dx}{2} + \frac{c}{2} \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(d*x+c))^(7/2), x)

[Out] 2/5*(b*(2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/b^4/sin(1/2*d*x+1/2*c)^3/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)*(12*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^4-24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^2+24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))-8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c))*(-2*sin(1/2*d*x+1/2*c)^4*b+sin(1/2*d*x+1/2*c)^2*b)^(1/2)/(b*(2*cos(1/2*d*x+1/2*c)^2-1))^(1/2)/d

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(7/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(b \cos(c + dx))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(c + d*x))^(7/2),x)

[Out] int(1/(b*cos(c + d*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))**(7/2),x)

[Out] Timed out

3.140 $\int \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal. Leaf size=98

$$\frac{3x\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{\sin(c + dx) \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)}}{4d} + \frac{3 \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)}}{8d}$$

[Out] $1/4*\cos(d*x+c)^{(5/2)}*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d+3/8*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+3/8*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.03, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{3x\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{\sin(c + dx) \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)}}{4d} + \frac{3 \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)}}{8d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^{(7/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]], x]$

[Out] $(3*x*\text{Sqrt}[b*\text{Cos}[c + d*x]])/(8*\text{Sqrt}[\text{Cos}[c + d*x]]) + (3*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(8*d) + (\text{Cos}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(4*d)$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_)*((a_)*(v_))^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/\text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}\{a, b, m\}, x \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 2635

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c + d*x])*(b*\text{Sin}[c + d*x])^{(n-1)})/(d*n), x] + \text{Dist}[(b^{(n-1)})/n, \text{Int}[(b*\text{Sin}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned} \int \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx &= \frac{\sqrt{b \cos(c + dx)} \int \cos^4(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{\cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d} + \frac{(3\sqrt{b \cos(c + dx)}) \int \cos^2(c + dx) dx}{4\sqrt{\cos(c + dx)}} \\ &= \frac{3\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d} + \frac{\cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d} \\ &= \frac{3x\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{3\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d} + \frac{\cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d} \end{aligned}$$

[In] integrate(cos(d*x+c)^(7/2)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d

mupad [B] time = 1.26, size = 75, normalized size = 0.77

$$\frac{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} (8 \sin(c+dx) + 9 \sin(3c+3dx) + \sin(5c+5dx) + 24dx \cos(c+dx))}{32d (\cos(2c+2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(7/2)*(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(7/2)*(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

$$3.141 \quad \int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx$$

Optimal. Leaf size=70

$$\frac{\sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{\sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}}$$

[Out] $\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}-1/3*\sin(d*x+c)^3*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{\sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{\sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) - (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x]^3)/(3*d*Sqrt[Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx &= \frac{\sqrt{b \cos(c + dx)} \int \cos^3(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{\sqrt{b \cos(c + dx)} \text{Subst}\left(\int (1 - x^2) dx, x, -\sin(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{\sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.09, size = 45, normalized size = 0.64

$$\frac{\sin(c + dx)(\cos(2(c + dx)) + 5)\sqrt{b \cos(c + dx)}}{6d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[b*Cos[c + d*x]]*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*d*Sqrt[Cos[c + d*x]])

fricas [A] time = 0.87, size = 39, normalized size = 0.56

$$\frac{\sqrt{b \cos(dx+c)} (\cos(dx+c)^2 + 2) \sin(dx+c)}{3d\sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(cos(d*x + c)^2 + 2)*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] Timed out

maple [A] time = 0.11, size = 40, normalized size = 0.57

$$\frac{(2 + \cos^2(dx+c)) \sin(dx+c) \sqrt{b \cos(dx+c)}}{3d\sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(1/2),x)

[Out] 1/3/d*(2+cos(d*x+c)^2)*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2)

maxima [A] time = 1.12, size = 42, normalized size = 0.60

$$\frac{\sqrt{b} \left(\sin(3dx+3c) + 9 \sin\left(\frac{1}{3} \arctan(\sin(3dx+3c), \cos(3dx+3c))\right) \right)}{12d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/12*sqrt(b)*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/d

mupad [B] time = 0.74, size = 57, normalized size = 0.81

$$\frac{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} (10 \sin(2c+2dx) + \sin(4c+4dx))}{12d (\cos(2c+2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(5/2)*(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

$$3.142 \quad \int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx$$

Optimal. Leaf size=63

$$\frac{x\sqrt{b \cos(c + dx)}}{2\sqrt{\cos(c + dx)}} + \frac{\sin(c + dx)\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}}{2d}$$

[Out] 1/2*x*(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2)+1/2*sin(d*x+c)*cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(1/2)/d

Rubi [A] time = 0.01, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{x\sqrt{b \cos(c + dx)}}{2\sqrt{\cos(c + dx)}} + \frac{\sin(c + dx)\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}}{2d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]], x]

[Out] (x*Sqrt[b*Cos[c + d*x]])/(2*Sqrt[Cos[c + d*x]]) + (Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(2*d)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx &= \frac{\sqrt{b \cos(c + dx)} \int \cos^2(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d} + \frac{\sqrt{b \cos(c + dx)} \int 1 dx}{2\sqrt{\cos(c + dx)}} \\ &= \frac{x\sqrt{b \cos(c + dx)}}{2\sqrt{\cos(c + dx)}} + \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d} \end{aligned}$$

Mathematica [A] time = 0.05, size = 45, normalized size = 0.71

$$\frac{(2(c + dx) + \sin(2(c + dx)))\sqrt{b \cos(c + dx)}}{4d\sqrt{\cos(c + dx)}}$$

2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)(sqrt(b)*d*x*tan((c+d*x)/2)^4+2*sqrt(b)*d*x*tan((c+d*x)/2)^2+sqrt(b)*d*x+(-2*sqrt(b))*tan((c+d*x)/2)^3+2*sqrt(b)*tan((c+d*x)/2))/(2*d*tan((c+d*x)/2)^4+4*d*tan((c+d*x)/2)^2+2*d)

maple [A] time = 0.12, size = 42, normalized size = 0.67

$$\frac{\sqrt{b \cos(dx + c)} (\cos(dx + c) \sin(dx + c) + dx + c)}{2d\sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(1/2),x)

[Out] 1/2/d*(b*cos(d*x+c))^(1/2)*(cos(d*x+c)*sin(d*x+c)+d*x+c)/cos(d*x+c)^(1/2)

maxima [A] time = 1.10, size = 25, normalized size = 0.40

$$\frac{(2dx + 2c + \sin(2dx + 2c))\sqrt{b}}{4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))*sqrt(b)/d

mupad [B] time = 0.69, size = 62, normalized size = 0.98

$$\frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (\sin(c + dx) + \sin(3c + 3dx) + 4dx \cos(c + dx))}{4d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(4*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(3/2)*(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

3.143 $\int \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} dx$

Optimal. Leaf size=32

$$\frac{\sin(c + dx)\sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[Out] $\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{\sin(c + dx)\sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2637

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} dx &= \frac{\sqrt{b \cos(c + dx)} \int \cos(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sin(c + dx)\sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])

fricas [A] time = 0.73, size = 28, normalized size = 0.88

$$\frac{\sqrt{b \cos(dx + c)} \sin(dx + c)}{d\sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [A] time = 2.56, size = 31, normalized size = 0.97

$$\frac{2\sqrt{b}\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{d\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] 2*sqrt(b)*tan(1/2*d*x + 1/2*c)/(d*tan(1/2*d*x + 1/2*c)^2 + d)

maple [A] time = 0.11, size = 29, normalized size = 0.91

$$\frac{\sin(dx + c)\sqrt{b\cos(dx + c)}}{d\sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(1/2),x)

[Out] sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

maxima [A] time = 0.87, size = 13, normalized size = 0.41

$$\frac{\sqrt{b}\sin(dx + c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] sqrt(b)*sin(d*x + c)/d

mupad [B] time = 0.39, size = 44, normalized size = 1.38

$$\frac{\sqrt{\cos(c + dx)}\sin(2c + 2dx)\sqrt{b\cos(c + dx)}}{d(\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*sin(2*c + 2*d*x)*(b*cos(c + d*x))^(1/2))/(d*(cos(2*c + 2*d*x) + 1))

sympy [A] time = 10.76, size = 29, normalized size = 0.91

$$\begin{cases} \frac{\sqrt{b}\sin(c+dx)}{d} & \text{for } d \neq 0 \\ x\sqrt{b\cos(c)}\sqrt{\cos(c)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(1/2)*(b*cos(d*x+c))**(1/2),x)

[Out] Piecewise((sqrt(b)*sin(c + d*x)/d, Ne(d, 0)), (x*sqrt(b*cos(c))*sqrt(cos(c)), True))

$$3.144 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} dx$$

Optimal. Leaf size=24

$$\frac{x\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[Out] $x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{x\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]/Sqrt[Cos[c + d*x]],x]

[Out] (x*Sqrt[b*Cos[c + d*x]])/Sqrt[Cos[c + d*x]]

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} dx &= \frac{\sqrt{b \cos(c+dx)} \int 1 dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{x\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 1.00

$$\frac{x\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Sqrt[Cos[c + d*x]],x]

[Out] (x*Sqrt[b*Cos[c + d*x]])/Sqrt[Cos[c + d*x]]

fricas [A] time = 0.70, size = 94, normalized size = 3.92

$$\left[\frac{\sqrt{-b} \log(2b \cos(dx+c)^2 - 2\sqrt{b \cos(dx+c)} \sqrt{-b} \sqrt{\cos(dx+c)} \sin(dx+c) - b)}{2d}, \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{\sqrt{b \cos(dx+c)}}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),x, algorithm="fricas")

[Out] [1/2*sqrt(-b)*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b)/d, sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \cos(dx + c)}}{\sqrt{\cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/sqrt(cos(d*x + c)), x)

maple [A] time = 0.08, size = 28, normalized size = 1.17

$$\frac{\sqrt{b \cos(dx + c)} (dx + c)}{d \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),x)

[Out] 1/d*(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2)*(d*x+c)

maxima [A] time = 0.88, size = 26, normalized size = 1.08

$$\frac{2 \sqrt{b} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d

mupad [B] time = 0.10, size = 20, normalized size = 0.83

$$\frac{x \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(1/2),x)

[Out] (x*(b*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2)

sympy [A] time = 1.75, size = 5, normalized size = 0.21

$$\sqrt{b} x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(1/2),x)

[Out] sqrt(b)*x

$$3.145 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)} dx$$

Optimal. Leaf size=33

$$\frac{\sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{\cos(c+dx)}}$$

[Out] arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{\sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(3/2),x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(d*Sqrt[Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)} dx &= \frac{\sqrt{b \cos(c+dx)} \int \sec(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 33, normalized size = 1.00

$$\frac{\sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(3/2),x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(d*Sqrt[Cos[c + d*x]])

fricas [A] time = 0.60, size = 113, normalized size = 3.42

$$\left[\frac{\sqrt{b} \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right)}{2d}, -\frac{\sqrt{-b} \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b} \sin(dx+c)}{b \sqrt{\cos(dx+c)}}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(3/2),x, algorithm="fricas")

[Out] [1/2*sqrt(b)*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3)/d, -sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \cos(dx + c)}}{\cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(3/2), x)

maple [A] time = 0.11, size = 42, normalized size = 1.27

$$\frac{2 \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right) \sqrt{b \cos(dx+c)}}{d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(3/2),x)

[Out] -2/d*arctanh((-1+cos(d*x+c))/sin(d*x+c))*(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2)

maxima [B] time = 1.27, size = 65, normalized size = 1.97

$$\frac{\sqrt{b} \left(\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2 \sin(dx+c) + 1) \right)}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(3/2),x, algorithm="maxima")

[Out] 1/2*sqrt(b)*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/d

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(3/2),x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(3/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))/cos(c + d*x)**(3/2), x)

$$3.146 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^2(c+dx)} dx$$

Optimal. Leaf size=32

$$\frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[Out] $\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(3/2)}$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[b*\text{Cos}[c + d*x]]/\text{Cos}[c + d*x]^{(5/2)}, x]$

[Out] $(\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Cos}[c + d*x]^{(3/2)})$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_.))^{(m_.)*((b_.)*(v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\amp; \text{!IntegerQ}[m] \&\amp; \text{IGtQ}[n+1/2, 0] \&\amp; \text{IntegerQ}[m+n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1+x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c+d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \&\amp; \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{5}{2}}(c+dx)} dx &= \frac{\sqrt{b \cos(c+dx)} \int \sec^2(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= -\frac{\sqrt{b \cos(c+dx)} \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{d\sqrt{\cos(c+dx)}} \\ &= \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(5/2),x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2))

fricas [A] time = 0.61, size = 28, normalized size = 0.88

$$\frac{\sqrt{b \cos(dx + c)} \sin(dx + c)}{d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(3/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \cos(dx + c)}}{\cos(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(5/2), x)

maple [A] time = 0.12, size = 29, normalized size = 0.91

$$\frac{\sin(dx + c) \sqrt{b \cos(dx + c)}}{d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2),x)

[Out] sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)

maxima [A] time = 1.10, size = 54, normalized size = 1.69

$$\frac{2 \sqrt{b} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c)^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 0.70, size = 59, normalized size = 1.84

$$\frac{\sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 1i + \sin(2c + 2dx) + 1i)}{d \sqrt{\cos(c + dx)} (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(5/2),x)

[Out] ((b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/(d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(5/2), x)

[Out] Integral(sqrt(b*cos(c + d*x))/cos(c + d*x)**(5/2), x)

$$3.147 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{7}{2}}(c+dx)} dx$$

Optimal. Leaf size=72

$$\frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{\sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2d\sqrt{\cos(c+dx)}}$$

[Out] $1/2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(5/2)}+1/2*\operatorname{arctanh}(\sin(d*x+c))*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{\sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(7/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(2*d*Sqrt[Cos[c + d*x]]) + (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(2*d*Cos[c + d*x]^(5/2))

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{7}{2}}(c+dx)} dx &= \frac{\sqrt{b \cos(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{\sqrt{b \cos(c+dx)} \int \sec(c+dx) dx}{2\sqrt{\cos(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{2d\sqrt{\cos(c+dx)}} + \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.04, size = 52, normalized size = 0.72

$$\frac{\sqrt{b \cos(c + dx)} (\sin(c + dx) + \cos^2(c + dx) \tanh^{-1}(\sin(c + dx)))}{2d \cos^{\frac{5}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(7/2), x]

[Out] (Sqrt[b*Cos[c + d*x]]*(ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x])/ (2*d*Cos[c + d*x]^(5/2)))

fricas [A] time = 0.83, size = 201, normalized size = 2.79

$$\frac{\sqrt{b} \cos(dx + c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b} \cos(dx+c) \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b} \cos(dx + c) \sqrt{\cos(dx + c)}}{4d \cos(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(7/2), x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3), -1/2*(sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \cos(dx + c)}}{\cos(dx + c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(7/2), x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(7/2), x)

maple [A] time = 0.14, size = 104, normalized size = 1.44

$$\frac{\left(\cos^2(dx + c) \ln\left(-\frac{-1 + \cos(dx+c) + \sin(dx+c)}{\sin(dx+c)}\right) - \cos^2(dx + c) \ln\left(-\frac{-\sin(dx+c) - 1 + \cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx + c)\right) \sqrt{b \cos(dx + c)}}{2d \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(7/2), x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c))*(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2)

maxima [B] time = 1.13, size = 661, normalized size = 9.18

$$\frac{\left(4(\sin(4dx + 4c) + 2\sin(2dx + 2c)) \cos\left(\frac{3}{2} \arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) - 4(\sin(4dx + 4c) + 2\sin(2dx + 2c))\right) \sqrt{b \cos(dx + c)}}{2d \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(7/2),x, algorithm="maxima")

[Out]
$$-1/4*(4*(\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\cos(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 4*(\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - (2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) + (2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 - 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) - 4*(\cos(4*d*x + 4*c) + 2*\cos(2*d*x + 2*c) + 1)*\sin(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 4*(\cos(4*d*x + 4*c) + 2*\cos(2*d*x + 2*c) + 1)*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))*\sqrt{b}/((2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*d)$$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(7/2),x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(7/2),x)

[Out] Timed out

$$3.148 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^2(c+dx)} dx$$

Optimal. Leaf size=70

$$\frac{\sin^3(c+dx)\sqrt{b \cos(c+dx)}}{3d \cos^{\frac{7}{2}}(c+dx)} + \frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[Out] $\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(3/2)}+1/3*\sin(d*x+c)^3*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(7/2)}$

Rubi [A] time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3767}

$$\frac{\sin^3(c+dx)\sqrt{b \cos(c+dx)}}{3d \cos^{\frac{7}{2}}(c+dx)} + \frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(9/2), x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)) + (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x]^3)/(3*d*Cos[c + d*x]^(7/2))

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \cos(c+dx)}}{\cos^2(c+dx)} dx &= \frac{\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} \int \sec^4(c+dx) dx \\ &= -\frac{\sqrt{b \cos(c+dx)} \operatorname{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d\sqrt{\cos(c+dx)}} \\ &= \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} + \frac{\sqrt{b \cos(c+dx)} \sin^3(c+dx)}{3d \cos^{\frac{7}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.08, size = 45, normalized size = 0.64

$$\frac{\left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx)\right) \sqrt{b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(9/2),x]

[Out] (Sqrt[b*Cos[c + d*x]]*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sqrt[Cos[c + d*x]])

fricas [A] time = 0.57, size = 41, normalized size = 0.59

$$\frac{\sqrt{b \cos(dx + c)} (2 \cos(dx + c)^2 + 1) \sin(dx + c)}{3 d \cos(dx + c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(9/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 1)*sin(d*x + c)/(d*cos(d*x + c)^(7/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \cos(dx + c)}}{\cos(dx + c)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(9/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(9/2), x)

maple [A] time = 0.12, size = 42, normalized size = 0.60

$$\frac{(2(\cos^2(dx + c)) + 1) \sqrt{b \cos(dx + c)} \sin(dx + c)}{3d \cos(dx + c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(9/2),x)

[Out] 1/3/d*(2*cos(d*x+c)^2+1)*(b*cos(d*x+c))^(1/2)*sin(d*x+c)/cos(d*x+c)^(7/2)

maxima [B] time = 1.13, size = 294, normalized size = 4.20

$$\frac{4((3 \cos(2dx + 2c) + 1) \sin(6dx + 6c) + 3(3 \cos(2dx + 2c) + 1) \sin(4dx + 4c) - 3 \cos(6dx + 6c) \sin(2dx + 2c) - 9 \cos(4dx + 4c) \sin(2dx + 2c)) \sqrt{b}}{3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \cos(4dx + 4c) + 9 \cos(4dx + 4c)^2 + 9 \cos(2dx + 2c)^2 + 6(\sin(4dx + 4c) + \sin(2dx + 2c)) \sin(6dx + 6c) + \sin(6dx + 6c)^2 + 9 \sin(4dx + 4c)^2 + 18 \sin(4dx + 4c) \sin(2dx + 2c) + 9 \sin(2dx + 2c)^2 + 6 \cos(2dx + 2c) + 1) d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(9/2),x, algorithm="maxima")

[Out] 4/3*((3*cos(2*d*x + 2*c) + 1)*sin(6*d*x + 6*c) + 3*(3*cos(2*d*x + 2*c) + 1)*sin(4*d*x + 4*c) - 3*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) - 9*cos(4*d*x + 4*c)*sin(2*d*x + 2*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 1.86, size = 128, normalized size = 1.83

$$\frac{2 \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 15i + \cos(4c + 4dx) 6i + \cos(6c + 6dx) 1i + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}{3 d \sqrt{\cos(c + dx)} (15 \cos(2c + 2dx) + 6 \cos(4c + 4dx) + \cos(6c + 6dx) + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(9/2),x)
```

```
[Out] (2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(9/2),x)
```

```
[Out] Timed out
```

$$3.149 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{11}{2}}(c+dx)} dx$$

Optimal. Leaf size=107

$$\frac{3 \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{\frac{5}{2}}(c+dx)} + \frac{\sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3 \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8d \sqrt{\cos(c+dx)}}$$

[Out] 1/4*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(9/2)+3/8*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+3/8*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{3 \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{\frac{5}{2}}(c+dx)} + \frac{\sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3 \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(11/2),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(8*d*Sqrt[Cos[c + d*x]]) + (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(4*d*Cos[c + d*x]^(9/2)) + (3*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(8*d*Cos[c + d*x]^(5/2))

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{11}{2}}(c+dx)} dx &= \frac{\sqrt{b \cos(c+dx)} \int \sec^5(c+dx) dx}{\sqrt{\cos(c+dx)}} \\
&= \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{(3\sqrt{b \cos(c+dx)}) \int \sec^3(c+dx) dx}{4\sqrt{\cos(c+dx)}} \\
&= \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3\sqrt{b \cos(c+dx)} \sin(c+dx)}{8d \cos^{\frac{5}{2}}(c+dx)} + \frac{(3\sqrt{b \cos(c+dx)}) \int \sec(c+dx) dx}{8\sqrt{\cos(c+dx)}} \\
&= \frac{3 \tanh^{-1}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{8d\sqrt{\cos(c+dx)}} + \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3\sqrt{b \cos(c+dx)}}{8d \cos^{\frac{5}{2}}(c+dx)}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 66, normalized size = 0.62

$$\frac{\sqrt{b \cos(c+dx)} (\sin(c+dx) (3 \cos^2(c+dx) + 2) + 3 \cos^4(c+dx) \tanh^{-1}(\sin(c+dx)))}{8d \cos^{\frac{9}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(11/2), x]

[Out] (Sqrt[b*Cos[c + d*x]]*(3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x]))/(8*d*Cos[c + d*x]^(9/2))

fricas [A] time = 0.60, size = 227, normalized size = 2.12

$$\left[\frac{3 \sqrt{b} \cos(dx+c)^5 \log\left(\frac{-b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2 \sqrt{b \cos(dx+c)} (3 \cos(dx+c)^2 + 2) \sin(dx+c)}{16 d \cos(dx+c)^5} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(11/2), x, algorithm="fricas")

[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5), -1/8*(3*sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^5 - sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \cos(dx+c)}}{\cos(dx+c)^{\frac{11}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(11/2), x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(11/2), x)

maple [A] time = 0.20, size = 121, normalized size = 1.13

$$\frac{\left(3 \left(\cos^4(dx+c)\right) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) - 3 \left(\cos^4(dx+c)\right) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - 3 \left(\cos^2(dx+c)\right) \ln\left(\frac{\cos(dx+c)}{\sin(dx+c)}\right)\right)}{8d \cos(dx+c)^{\frac{9}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(11/2),x)
```

```
[Out] -1/8/d*(3*cos(d*x+c)^4*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^4*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^2*sin(d*x+c)-2*sin(d*x+c))*(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(9/2)
```

maxima [B] time = 1.23, size = 1656, normalized size = 15.48

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(11/2),x, algorithm="maxima")
```

```
[Out] -1/16*(12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*d)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^{11/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(11/2), x)`

[Out] `int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(11/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(11/2), x)`

[Out] Timed out

3.150 $\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal. Leaf size=101

$$\frac{3bx\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{b \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)\sqrt{b \cos(c + dx)}}{4d} + \frac{3b \sin(c + dx)\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}}{8d}$$

[Out] $\frac{1}{4}bx\cos(dx+c)^{5/2}\sin(dx+c)(b\cos(dx+c))^{1/2}/d + \frac{3}{8}bx(b\cos(dx+c))^{1/2}/\cos(dx+c)^{1/2} + \frac{3}{8}b\sin(dx+c)\cos(dx+c)^{1/2}(b\cos(dx+c))^{1/2}/d$

Rubi [A] time = 0.03, antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{3bx\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{b \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)\sqrt{b \cos(c + dx)}}{4d} + \frac{3b \sin(c + dx)\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}}{8d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(3/2), x]

[Out] $(3bx\sqrt{b\cos[c + dx]})/(8\sqrt{\cos[c + dx]}) + (3b\sqrt{\cos[c + dx]}\sqrt{b\cos[c + dx]}\sin[c + dx])/(8d) + (b\cos[c + dx]^{5/2}\sqrt{b\cos[c + dx]}\sin[c + dx])/(4d)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Sine[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx &= \frac{(b\sqrt{b \cos(c + dx)}) \int \cos^4(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{b \cos^{\frac{5}{2}}(c + dx)\sqrt{b \cos(c + dx)} \sin(c + dx)}{4d} + \frac{(3b\sqrt{b \cos(c + dx)}) \int \cos^2(c + dx) dx}{4\sqrt{\cos(c + dx)}} \\ &= \frac{3b\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)} \sin(c + dx)}{8d} + \frac{b \cos^{\frac{5}{2}}(c + dx)\sqrt{b \cos(c + dx)}}{4d} \\ &= \frac{3bx\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{3b\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)} \sin(c + dx)}{8d} + \frac{b \cos^{\frac{5}{2}}(c + dx)\sqrt{b \cos(c + dx)}}{4d} \end{aligned}$$

maple [A] time = 0.14, size = 62, normalized size = 0.61

$$\frac{(b \cos(dx + c))^{\frac{3}{2}} \left(2 \left(\cos^3(dx + c) \right) \sin(dx + c) + 3 \cos(dx + c) \sin(dx + c) + 3dx + 3c \right)}{8d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(3/2),x)

[Out] 1/8/d*(b*cos(d*x+c))^(3/2)*(2*cos(d*x+c)^3*sin(d*x+c)+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)/cos(d*x+c)^(3/2)

maxima [A] time = 1.05, size = 53, normalized size = 0.52

$$\frac{\left(12(dx + c)b + b \sin(4dx + 4c) + 8b \sin\left(\frac{1}{2} \arctan(\sin(4dx + 4c), \cos(4dx + 4c))\right) \right) \sqrt{b}}{32d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/32*(12*(d*x + c)*b + b*sin(4*d*x + 4*c) + 8*b*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d

mupad [B] time = 1.05, size = 76, normalized size = 0.75

$$\frac{b \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx) + 24dx \cos(c + dx))}{32d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(3/2),x)

[Out] (b*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(5/2)*(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

$$3.151 \quad \int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx$$

Optimal. Leaf size=72

$$\frac{b \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{b \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}}$$

[Out] $b \sin(d*x+c) * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)} - 1/3 * b \sin(d*x+c)^3 * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{b \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{b \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(3/2), x]

[Out] $(b \sqrt{b \cos(c + dx)} \sin(c + dx)) / (d \sqrt{\cos(c + dx)}) - (b \sqrt{b \cos(c + dx)} \sin^3(c + dx)) / (3d \sqrt{\cos(c + dx)})$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_.), x_Symbol] := -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx &= \frac{(b \sqrt{b \cos(c + dx)}) \int \cos^3(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{(b \sqrt{b \cos(c + dx)}) \text{Subst}\left(\int (1 - x^2) dx, x, -\sin(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\ &= \frac{b \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{b \sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.11, size = 45, normalized size = 0.62

$$\frac{\sin(c + dx)(\cos(2(c + dx)) + 5)(b \cos(c + dx))^{3/2}}{6d \cos^{\frac{3}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(3/2), x]

[Out] $((b \cos[c + d*x])^{(3/2)} * (5 + \cos[2*(c + d*x)]) * \sin[c + d*x]) / (6*d*\cos[c + d*x]^{(3/2)})$

fricas [A] time = 0.55, size = 43, normalized size = 0.60

$$\frac{(b \cos(dx + c)^2 + 2b) \sqrt{b \cos(dx + c)} \sin(dx + c)}{3d \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out] $1/3*(b*\cos(d*x + c)^2 + 2*b)*\sqrt{b*\cos(d*x + c)}*\sin(d*x + c)/(d*\sqrt{\cos(d*x + c)})$

giac [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(3/2),x, algorithm="giac")`

[Out] Timed out

maple [A] time = 0.08, size = 40, normalized size = 0.56

$$\frac{(2 + \cos^2(dx + c)) \sin(dx + c) (b \cos(dx + c))^{\frac{3}{2}}}{3d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(3/2),x)`

[Out] $1/3/d*(2+\cos(d*x+c)^2)*\sin(d*x+c)*(b*\cos(d*x+c))^{(3/2)}/\cos(d*x+c)^{(3/2)}$

maxima [A] time = 1.25, size = 45, normalized size = 0.62

$$\frac{\left(b \sin(3dx + 3c) + 9b \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)\right) \sqrt{b}}{12d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] $1/12*(b*\sin(3*d*x + 3*c) + 9*b*\sin(1/3*\arctan2(\sin(3*d*x + 3*c), \cos(3*d*x + 3*c))))*\sqrt{b}/d$

mupad [B] time = 0.70, size = 58, normalized size = 0.81

$$\frac{b \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(3/2),x)`

[Out] $(b*\cos(c + d*x)^{(1/2)}*(b*\cos(c + d*x))^{(1/2)}*(10*\sin(2*c + 2*d*x) + \sin(4*c + 4*d*x)))/(12*d*(\cos(2*c + 2*d*x) + 1))$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**(3/2)*(b*cos(d*x+c))**(3/2),x)
```

```
[Out] Timed out
```

3.152 $\int \sqrt{\cos(c + dx)} (b \cos(c + dx))^{3/2} dx$

Optimal. Leaf size=65

$$\frac{bx\sqrt{b\cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}{2d}$$

[Out] $1/2*b*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+1/2*b*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.01, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{bx\sqrt{b\cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}{2d}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(3/2), x]

[Out] $(b*x*\text{Sqrt}[b*\text{Cos}[c + d*x]])/(2*\text{Sqrt}[\text{Cos}[c + d*x]]) + (b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(2*d)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sqrt{\cos(c + dx)} (b \cos(c + dx))^{3/2} dx &= \frac{(b\sqrt{b\cos(c+dx)}) \int \cos^2(c + dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{2d} + \frac{(b\sqrt{b\cos(c+dx)}) \int 1 dx}{2\sqrt{\cos(c+dx)}} \\ &= \frac{bx\sqrt{b\cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{2d} \end{aligned}$$

Mathematica [A] time = 0.05, size = 45, normalized size = 0.69

$$\frac{(2(c + dx) + \sin(2(c + dx)))(b \cos(c + dx))^{3/2}}{4d \cos^{\frac{3}{2}}(c + dx)}$$

2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)b*(sqrt(b)*d*x*tan((c+d*x)/2)^4+2*sqrt(b)*d*x*tan((c+d*x)/2)^2+sqrt(b)*d*x+(-2*sqrt(b))*tan((c+d*x)/2)^3+2*sqrt(b)*tan((c+d*x)/2))/(2*d*tan((c+d*x)/2)^4+4*d*tan((c+d*x)/2)^2+2*d)

maple [A] time = 0.10, size = 42, normalized size = 0.65

$$\frac{(b \cos(dx + c))^{\frac{3}{2}} (\cos(dx + c) \sin(dx + c) + dx + c)}{2d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(3/2), x)

[Out] 1/2/d*(b*cos(d*x+c))^(3/2)*(cos(d*x+c)*sin(d*x+c)+d*x+c)/cos(d*x+c)^(3/2)

maxima [A] time = 1.54, size = 28, normalized size = 0.43

$$\frac{(2(dx + c)b + b \sin(2dx + 2c))\sqrt{b}}{4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(3/2), x, algorithm="maxima")

[Out] 1/4*(2*(d*x + c)*b + b*sin(2*d*x + 2*c))*sqrt(b)/d

mupad [B] time = 0.53, size = 63, normalized size = 0.97

$$\frac{b \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (\sin(c + dx) + \sin(3c + 3dx) + 4dx \cos(c + dx))}{4d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(3/2), x)

[Out] (b*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(4*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(1/2)*(b*cos(d*x+c))**(3/2), x)

[Out] Timed out

$$3.153 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx$$

Optimal. Leaf size=33

$$\frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}}$$

[Out] b*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(3/2)/Sqrt[Cos[c + d*x]],x]

[Out] (b*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2637

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] :> Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx &= \frac{(b \sqrt{b \cos(c+dx)}) \int \cos(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 32, normalized size = 0.97

$$\frac{\sin(c+dx)(b \cos(c+dx))^{3/2}}{d \cos^2(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Sqrt[Cos[c + d*x]],x]

[Out] ((b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2))

fricas [A] time = 0.51, size = 29, normalized size = 0.88

$$\frac{\sqrt{b \cos(dx+c)} b \sin(dx+c)}{d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(1/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*b*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{3}{2}}}{\sqrt{\cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/sqrt(cos(d*x + c)), x)

maple [A] time = 0.10, size = 29, normalized size = 0.88

$$\frac{(b \cos(dx + c))^{\frac{3}{2}} \sin(dx + c)}{d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(1/2),x)

[Out] 1/d*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/cos(d*x+c)^(3/2)

maxima [A] time = 1.09, size = 13, normalized size = 0.39

$$\frac{b^{\frac{3}{2}} \sin(dx + c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(1/2),x, algorithm="maxima")

[Out] b^(3/2)*sin(d*x + c)/d

mupad [B] time = 0.24, size = 29, normalized size = 0.88

$$\frac{b \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(1/2),x)

[Out] (b*sin(c + d*x)*(b*cos(c + d*x))^(1/2))/(d*cos(c + d*x)^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(1/2),x)

[Out] Timed out

$$3.154 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{3}{2}}(c+dx)} dx$$

Optimal. Leaf size=25

$$\frac{bx\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[Out] b*x*(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2)

Rubi [A] time = 0.00, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{bx\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(3/2), x]

[Out] (b*x*Sqrt[b*Cos[c + d*x]])/Sqrt[Cos[c + d*x]]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{3}{2}}(c+dx)} dx &= \frac{(b\sqrt{b \cos(c+dx)}) \int 1 dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{bx\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 0.96

$$\frac{x(b \cos(c+dx))^{3/2}}{\cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(3/2), x]

[Out] (x*(b*Cos[c + d*x])^(3/2))/Cos[c + d*x]^(3/2)

fricas [A] time = 0.59, size = 95, normalized size = 3.80

$$\left[\frac{\sqrt{-b} b \log \left(2 b \cos (d x + c)^2 - 2 \sqrt{b \cos (d x + c)} \sqrt{-b} \sqrt{\cos (d x + c)} \sin (d x + c) - b \right)}{2 d}, \frac{b^{\frac{3}{2}} \arctan \left(\frac{\sqrt{b \cos (d x + c)} \sin (d x + c)}{\sqrt{b \cos (d x + c)}^{\frac{3}{2}}} \right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2),x, algorithm="fricas")

[Out] [1/2*sqrt(-b)*b*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b)/d, b^(3/2)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{3}{2}}}{\cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(3/2), x)

maple [A] time = 0.06, size = 28, normalized size = 1.12

$$\frac{(b \cos(dx + c))^{\frac{3}{2}} (dx + c)}{d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2),x)

[Out] 1/d*(b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2)*(d*x+c)

maxima [A] time = 0.83, size = 26, normalized size = 1.04

$$\frac{2 b^{\frac{3}{2}} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2),x, algorithm="maxima")

[Out] 2*b^(3/2)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d

mupad [B] time = 0.09, size = 21, normalized size = 0.84

$$\frac{b x \sqrt{b \cos(c + d x)}}{\sqrt{\cos(c + d x)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(3/2),x)

[Out] (b*x*(b*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2)

sympy [A] time = 125.71, size = 5, normalized size = 0.20

$$b^{\frac{3}{2}} x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(3/2),x)

[Out] b**(3/2)*x

$$3.155 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{5}{2}}(c+dx)} dx$$

Optimal. Leaf size=34

$$\frac{b\sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{\cos(c+dx)}}$$

[Out] b*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{b\sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(5/2), x]

[Out] (b*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(d*Sqrt[Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{5}{2}}(c+dx)} dx &= \frac{(b\sqrt{b \cos(c+dx)}) \int \sec(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b \tanh^{-1}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 33, normalized size = 0.97

$$\frac{(b \cos(c+dx))^{3/2} \tanh^{-1}(\sin(c+dx))}{d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(5/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*(b*Cos[c + d*x])^(3/2))/(d*Cos[c + d*x]^(3/2))

fricas [A] time = 0.57, size = 114, normalized size = 3.35

$$\left[\frac{b^{\frac{3}{2}} \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right)}{2d}, -\frac{\sqrt{-b} b \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b} \sin(dx+c)}{b \sqrt{\cos(dx+c)}}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(5/2),x, algorithm="fricas")

[Out] [1/2*b^(3/2)*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3)/d, -sqrt(-b)*b*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{3}{2}}}{\cos(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(5/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(5/2), x)

maple [A] time = 0.08, size = 42, normalized size = 1.24

$$\frac{2 \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right) (b \cos(dx + c))^{\frac{3}{2}}}{d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(5/2),x)

[Out] -2/d*arctanh((-1+cos(d*x+c))/sin(d*x+c))*(b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2)

maxima [B] time = 1.11, size = 68, normalized size = 2.00

$$\frac{(b \log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - b \log(\cos(dx + c)^2 + \sin(dx + c)^2 - 2 \sin(dx + c) + 1)) \sqrt{b}}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(5/2),x, algorithm="maxima")

[Out] 1/2*(b*log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - b*log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))*sqrt(b)/d

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{(b \cos(c + dx))^{\frac{3}{2}}}{\cos(c + dx)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(5/2),x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(5/2),x)

[Out] Timed out

$$3.156 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^2(c+dx)} dx$$

Optimal. Leaf size=33

$$\frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[Out] b*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)

Rubi [A] time = 0.01, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(7/2), x]

[Out] (b*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_.), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{3/2}}{\cos^2(c+dx)} dx &= \frac{(b\sqrt{b \cos(c+dx)}) \int \sec^2(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= -\frac{(b\sqrt{b \cos(c+dx)}) \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{d\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 0.97

$$\frac{\sin(c+dx)(b \cos(c+dx))^{3/2}}{d \cos^{\frac{5}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(3/2)/cos[c + d*x]^(7/2),x]

[Out] ((b*cos[c + d*x])^(3/2)*sin[c + d*x])/(d*cos[c + d*x]^(5/2))

fricas [A] time = 0.79, size = 29, normalized size = 0.88

$$\frac{\sqrt{b \cos(dx + c)} b \sin(dx + c)}{d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(7/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*b*sin(d*x + c)/(d*cos(d*x + c)^(3/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{3}{2}}}{\cos(dx + c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(7/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(7/2), x)

maple [A] time = 0.10, size = 29, normalized size = 0.88

$$\frac{(b \cos(dx + c))^{\frac{3}{2}} \sin(dx + c)}{d \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(7/2),x)

[Out] 1/d*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/cos(d*x+c)^(5/2)

maxima [A] time = 0.83, size = 54, normalized size = 1.64

$$\frac{2 b^{\frac{3}{2}} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c)^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1) d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(7/2),x, algorithm="maxima")

[Out] 2*b^(3/2)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 0.50, size = 60, normalized size = 1.82

$$\frac{b \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) \operatorname{li} + \sin(2c + 2dx) + \operatorname{li})}{d \sqrt{\cos(c + dx)} (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(7/2),x)

[Out] (b*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*li + sin(2*c + 2*d*x) + li))/(d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(7/2), x)

[Out] Timed out

$$3.157 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^2(c+dx)} dx$$

Optimal. Leaf size=74

$$\frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{b \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2d \sqrt{\cos(c+dx)}}$$

[Out] $1/2*b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(5/2)}+1/2*b*\operatorname{arctanh}(\sin(d*x+c))*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{b \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c+d*x])^{(3/2)}/\text{Cos}[c+d*x]^{(9/2)},x]$

[Out] $(b*\text{ArcTanh}[\text{Sin}[c+d*x]]*\text{Sqrt}[b*\text{Cos}[c+d*x]])/(2*d*\text{Sqrt}[\text{Cos}[c+d*x]]) + (b*\text{Sqrt}[b*\text{Cos}[c+d*x]]*\text{Sin}[c+d*x])/(2*d*\text{Cos}[c+d*x]^{(5/2)})$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_.))^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/\text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n+1/2, 0] && IntegerQ[m+n]

Rule 3768

$\text{Int}[(\text{csc}[c_.] + (d_.)*(x_.))*(b_.))^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c+d*x])*(b*\text{Csc}[c+d*x])^{(n-1)})/(d*(n-1)), x] + \text{Dist}[(b^{2*(n-2)})/(n-1), \text{Int}[(b*\text{Csc}[c+d*x])^{(n-2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow -\text{Simp}[\text{ArcTanh}[\text{Cos}[c+d*x]]/d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{3/2}}{\cos^2(c+dx)} dx &= \frac{(b \sqrt{b \cos(c+dx)}) \int \sec^3(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b \sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{(b \sqrt{b \cos(c+dx)}) \int \sec(c+dx) dx}{2 \sqrt{\cos(c+dx)}} \\ &= \frac{b \tanh^{-1}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{2d \sqrt{\cos(c+dx)}} + \frac{b \sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.05, size = 52, normalized size = 0.70

$$\frac{(b \cos(c + dx))^{3/2} (\sin(c + dx) + \cos^2(c + dx) \tanh^{-1}(\sin(c + dx)))}{2d \cos^2(c + dx)^{7/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(9/2), x]

[Out] ((b*Cos[c + d*x])^(3/2)*(ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x]))/(2*d*Cos[c + d*x]^(7/2))

fricas [A] time = 0.55, size = 204, normalized size = 2.76

$$\left[\frac{b^2 \cos(dx + c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2 \sqrt{b \cos(dx+c)} b \sqrt{\cos(dx+c)}}{4 d \cos(dx+c)^3} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(9/2), x, algorithm="fricas")

[Out] [1/4*(b^(3/2)*cos(d*x + c)^3*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*b*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3), -1/2*(sqrt(-b)*b*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*b*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{3/2}}{\cos(dx + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(9/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(9/2), x)

maple [A] time = 0.11, size = 104, normalized size = 1.41

$$\frac{\left((\cos^2(dx + c)) \ln\left(-\frac{-1 + \cos(dx+c) + \sin(dx+c)}{\sin(dx+c)}\right) - (\cos^2(dx + c)) \ln\left(-\frac{-\sin(dx+c) - 1 + \cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx + c) \right) (b \cos(dx + c))^{3/2}}{2d \cos(dx + c)^{7/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(9/2), x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c))*(b*cos(d*x+c))^(3/2)/cos(d*x+c)^(7/2)

maxima [B] time = 1.21, size = 691, normalized size = 9.34

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(9/2), x, algorithm="maxima")

```
[Out] -1/4*(4*(b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (b*cos(4*d*x + 4*c)^2 + 4*b*cos(2*d*x + 2*c)^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b*sin(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + (b*cos(4*d*x + 4*c)^2 + 4*b*cos(2*d*x + 2*c)^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b*sin(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 4*(b*cos(4*d*x + 4*c) + 2*b*cos(2*d*x + 2*c) + b)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(b*cos(4*d*x + 4*c) + 2*b*cos(2*d*x + 2*c) + b)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^{9/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(9/2), x)
```

```
[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(9/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(9/2), x)
```

```
[Out] Timed out
```

$$3.158 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{11}{2}}(c+dx)} dx$$

Optimal. Leaf size=72

$$\frac{b \sin^3(c+dx) \sqrt{b \cos(c+dx)}}{3d \cos^{\frac{7}{2}}(c+dx)} + \frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[Out] b*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)+1/3*b*sin(d*x+c)^3*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(7/2)

Rubi [A] time = 0.02, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3767}

$$\frac{b \sin^3(c+dx) \sqrt{b \cos(c+dx)}}{3d \cos^{\frac{7}{2}}(c+dx)} + \frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(3/2)/Cos[c + d*x]^(11/2), x]

[Out] (b*Sqrt[b*cos[c + d*x]]*Sin[c + d*x])/(d*cos[c + d*x]^(3/2)) + (b*Sqrt[b*cos[c + d*x]]*Sin[c + d*x]^3)/(3*d*cos[c + d*x]^(7/2))

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{11}{2}}(c+dx)} dx &= \frac{(b\sqrt{b \cos(c+dx)}) \int \sec^4(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= -\frac{(b\sqrt{b \cos(c+dx)}) \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} + \frac{b\sqrt{b \cos(c+dx)} \sin^3(c+dx)}{3d \cos^{\frac{7}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.05, size = 45, normalized size = 0.62

$$\frac{\left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx)\right) (b \cos(c+dx))^{3/2}}{d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(3/2)/cos[c + d*x]^(11/2), x]

[Out] ((b*cos[c + d*x])^(3/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*cos[c + d*x]^(3/2))

fricas [A] time = 0.54, size = 42, normalized size = 0.58

$$\frac{(2b \cos(dx + c)^2 + b)\sqrt{b \cos(dx + c)} \sin(dx + c)}{3d \cos(dx + c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(11/2), x, algorithm="fricas")

[Out] 1/3*(2*b*cos(d*x + c)^2 + b)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(7/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{3}{2}}}{\cos(dx + c)^{\frac{11}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(11/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(11/2), x)

maple [A] time = 0.09, size = 42, normalized size = 0.58

$$\frac{(2(\cos^2(dx + c) + 1)(b \cos(dx + c))^{\frac{3}{2}} \sin(dx + c)}{3d \cos(dx + c)^{\frac{9}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(11/2), x)

[Out] 1/3/d*(2*cos(d*x+c)^2+1)*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/cos(d*x+c)^(9/2)

maxima [B] time = 1.70, size = 299, normalized size = 4.15

$$\frac{4(3b \cos(6dx + 6c) + \dots)}{3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \dots)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(11/2), x, algorithm="maxima")

[Out] -4/3*(3*b*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) + 9*b*cos(4*d*x + 4*c)*sin(2*d*x + 2*c) - (3*b*cos(2*d*x + 2*c) + b)*sin(6*d*x + 6*c) - 3*(3*b*cos(2*d*x + 2*c) + b)*sin(4*d*x + 4*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 1.14, size = 129, normalized size = 1.79

$$\frac{2b\sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 15i + \cos(4c + 4dx) 6i + \cos(6c + 6dx) 1i + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}{3d \sqrt{\cos(c + dx)} (15 \cos(2c + 2dx) + 6 \cos(4c + 4dx) + \cos(6c + 6dx) + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(11/2),x)
```

```
[Out] (2*b*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + c
os(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*
d*x) + 10i))/(3*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d
*x) + cos(6*c + 6*d*x) + 10))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(11/2),x)
```

```
[Out] Timed out
```

$$3.159 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{13}{2}}(c+dx)} dx$$

Optimal. Leaf size=110

$$\frac{3b \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{\frac{5}{2}}(c+dx)} + \frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3b \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8d \sqrt{\cos(c+dx)}}$$

[Out] 1/4*b*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(9/2)+3/8*b*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+3/8*b*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{3b \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{\frac{5}{2}}(c+dx)} + \frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3b \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(13/2), x]

[Out] (3*b*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(8*d*Sqrt[Cos[c + d*x]]) + (b*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(4*d*Cos[c + d*x]^(9/2)) + (3*b*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(8*d*Cos[c + d*x]^(5/2))

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{13/2}(c + dx)} dx &= \frac{(b\sqrt{b \cos(c + dx)}) \int \sec^5(c + dx) dx}{\sqrt{\cos(c + dx)}} \\
&= \frac{b\sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{9/2}(c + dx)} + \frac{(3b\sqrt{b \cos(c + dx)}) \int \sec^3(c + dx) dx}{4\sqrt{\cos(c + dx)}} \\
&= \frac{b\sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{9/2}(c + dx)} + \frac{3b\sqrt{b \cos(c + dx)} \sin(c + dx)}{8d \cos^{5/2}(c + dx)} + \frac{(3b\sqrt{b \cos(c + dx)}) \int \sec(c + dx) dx}{8\sqrt{\cos(c + dx)}} \\
&= \frac{3b \tanh^{-1}(\sin(c + dx))\sqrt{b \cos(c + dx)}}{8d\sqrt{\cos(c + dx)}} + \frac{b\sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{9/2}(c + dx)} + \frac{3b\sqrt{b \cos(c + dx)}}{8d \cos^{5/2}(c + dx)}
\end{aligned}$$

Mathematica [A] time = 0.08, size = 67, normalized size = 0.61

$$\frac{b\sqrt{b \cos(c + dx)} (\sin(c + dx) (3 \cos^2(c + dx) + 2) + 3 \cos^4(c + dx) \tanh^{-1}(\sin(c + dx)))}{8d \cos^{9/2}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(13/2), x]

[Out] (b*Sqrt[b*Cos[c + d*x]]*(3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x]))/(8*d*Cos[c + d*x]^(9/2))

fricas [A] time = 0.69, size = 234, normalized size = 2.13

$$\left[\frac{3 b^{\frac{3}{2}} \cos(dx + c)^5 \log\left(-\frac{b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2 b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2 (3 b \cos(dx + c)^2 + 2 b) \sqrt{b \cos(dx + c)}}{16 d \cos(dx + c)^5} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(13/2), x, algorithm="fricas")

[Out] [1/16*(3*b^(3/2)*cos(d*x + c)^5*log(-(b*cos(d*x + c)^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*(3*b*cos(d*x + c)^2 + 2*b)*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5), -1/8*(3*sqrt(-b)*b*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^5 - (3*b*cos(d*x + c)^2 + 2*b)*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{3}{2}}}{\cos(dx + c)^{\frac{13}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(13/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(13/2), x)

maple [A] time = 0.14, size = 121, normalized size = 1.10

$$\frac{3 (\cos^4(dx + c)) \ln\left(-\frac{\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - 3 (\cos^4(dx + c)) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) + 3 (\cos^2(dx + c)) \sin(dx + c)}{8d \cos(dx + c)^{\frac{11}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(13/2),x)
```

```
[Out] 1/8/d*(3*cos(d*x+c)^4*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^4*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))*(b*cos(d*x+c))^(3/2)/cos(d*x+c)^(11/2)
```

maxima [B] time = 1.23, size = 1742, normalized size = 15.84

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(13/2),x, algorithm="maxima")
```

```
[Out] -1/16*(12*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c) + 4*b*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c) + 4*b*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c) + 4*b*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c) + 4*b*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(b*cos(8*d*x + 8*c)^2 + 16*b*cos(6*d*x + 6*c)^2 + 36*b*cos(4*d*x + 4*c)^2 + 16*b*cos(2*d*x + 2*c)^2 + b*sin(8*d*x + 8*c)^2 + 16*b*sin(6*d*x + 6*c)^2 + 36*b*sin(4*d*x + 4*c)^2 + 48*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b*sin(2*d*x + 2*c)^2 + 2*(4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*cos(8*d*x + 8*c) + 8*(6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*cos(6*d*x + 6*c) + 12*(4*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 8*b*cos(2*d*x + 2*c) + 4*(2*b*sin(6*d*x + 6*c) + 3*b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + 3*(b*cos(8*d*x + 8*c)^2 + 16*b*cos(6*d*x + 6*c)^2 + 36*b*cos(4*d*x + 4*c)^2 + 16*b*cos(2*d*x + 2*c)^2 + b*sin(8*d*x + 8*c)^2 + 16*b*sin(6*d*x + 6*c)^2 + 36*b*sin(4*d*x + 4*c)^2 + 48*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b*sin(2*d*x + 2*c)^2 + 2*(4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*cos(8*d*x + 8*c) + 8*(6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*cos(6*d*x + 6*c) + 12*(4*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 8*b*cos(2*d*x + 2*c) + 4*(2*b*sin(6*d*x + 6*c) + 3*b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 12*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 12*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 +
```

$48*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 16*\sin(2*d*x + 2*c)^2 + 8*\cos(2*d*x + 2*c) + 1)*d$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + d x))^{3/2}}{\cos(c + d x)^{13/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(13/2), x)`

[Out] `int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(13/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(13/2), x)`

[Out] Timed out

3.160 $\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal. Leaf size=116

$$\frac{b^2 \sin^5(c + dx) \sqrt{b \cos(c + dx)}}{5d \sqrt{\cos(c + dx)}} - \frac{2b^2 \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}} + \frac{b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[Out] $b^2 \sin(d*x+c) * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)} - 2/3 * b^2 \sin(d*x+c)^3 * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)} + 1/5 * b^2 \sin(d*x+c)^5 * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{b^2 \sin^5(c + dx) \sqrt{b \cos(c + dx)}}{5d \sqrt{\cos(c + dx)}} - \frac{2b^2 \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}} + \frac{b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(5/2), x]

[Out] $(b^2 \sqrt{b \cos[c + d*x]} \sin[c + d*x]) / (d \sqrt{\cos[c + d*x]}) - (2 * b^2 \sqrt{b \cos[c + d*x]} \sin[c + d*x]^3) / (3 * d \sqrt{\cos[c + d*x]}) + (b^2 \sqrt{b \cos[c + d*x]} \sin[c + d*x]^5) / (5 * d \sqrt{\cos[c + d*x]})$

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx &= \frac{(b^2 \sqrt{b \cos(c + dx)}) \int \cos^5(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{(b^2 \sqrt{b \cos(c + dx)}) \text{Subst}\left(\int (1 - 2x^2 + x^4) dx, x, -\sin(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2b^2 \sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{\cos(c + dx)}} + \frac{b^2 \sqrt{b \cos(c + dx)} \sin^5(c + dx)}{5d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.12, size = 57, normalized size = 0.49

$$\frac{\sin(c + dx) \left(3 \sin^4(c + dx) - 10 \sin^2(c + dx) + 15\right) (b \cos(c + dx))^{5/2}}{15d \cos^{\frac{5}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(5/2),x]

[Out] ((b*Cos[c + d*x])^(5/2)*Sin[c + d*x]*(15 - 10*Sin[c + d*x]^2 + 3*Sin[c + d*x]^4))/(15*d*Cos[c + d*x]^(5/2))

fricas [A] time = 0.60, size = 61, normalized size = 0.53

$$\frac{(3b^2 \cos(dx + c)^4 + 4b^2 \cos(dx + c)^2 + 8b^2) \sqrt{b \cos(dx + c)} \sin(dx + c)}{15d \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/15*(3*b^2*cos(d*x + c)^4 + 4*b^2*cos(d*x + c)^2 + 8*b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] Timed out

maple [A] time = 0.12, size = 52, normalized size = 0.45

$$\frac{(3(\cos^4(dx + c)) + 4(\cos^2(dx + c)) + 8) \sin(dx + c) (b \cos(dx + c))^{\frac{5}{2}}}{15d \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(5/2),x)

[Out] 1/15/d*(3*cos(d*x+c)^4+4*cos(d*x+c)^2+8)*sin(d*x+c)*(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2)

maxima [A] time = 1.02, size = 77, normalized size = 0.66

$$\frac{(3b^2 \sin(5dx + 5c) + 25b^2 \sin\left(\frac{3}{5} \arctan(\sin(5dx + 5c), \cos(5dx + 5c))\right) + 150b^2 \sin\left(\frac{1}{5} \arctan(\sin(5dx + 5c), \cos(5dx + 5c))\right))}{240d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/240*(3*b^2*sin(5*d*x + 5*c) + 25*b^2*sin(3/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 150*b^2*sin(1/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))))*sqrt(b)/d

mupad [B] time = 1.35, size = 73, normalized size = 0.63

$$\frac{b^2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (175 \sin(2c + 2dx) + 28 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}{240d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(5/2),x)

[Out] (b^2*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(175*sin(2*c + 2*d*x) + 28*sin(4*c + 4*d*x) + 3*sin(6*c + 6*d*x)))/(240*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(5/2)*(b*cos(d*x+c))**(5/2), x)

[Out] Timed out

3.161 $\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal. Leaf size=107

$$\frac{3b^2x\sqrt{b\cos(c+dx)}}{8\sqrt{\cos(c+dx)}} + \frac{b^2\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}}{4d} + \frac{3b^2\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}{8d}$$

[Out] $\frac{1}{4}b^2\cos(dx+c)^{(5/2)}\sin(dx+c)*(b\cos(dx+c))^{(1/2)}/d + \frac{3}{8}b^2x*(b\cos(dx+c))^{(1/2)}/\cos(dx+c)^{(1/2)} + \frac{3}{8}b^2\sin(dx+c)*\cos(dx+c)^{(1/2)}*(b\cos(dx+c))^{(1/2)}/d$

Rubi [A] time = 0.03, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{3b^2x\sqrt{b\cos(c+dx)}}{8\sqrt{\cos(c+dx)}} + \frac{b^2\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}}{4d} + \frac{3b^2\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}{8d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(5/2), x]

[Out] $(3*b^2*x*\text{Sqrt}[b*\text{Cos}[c + d*x]])/(8*\text{Sqrt}[\text{Cos}[c + d*x]]) + (3*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(8*d) + (b^2*\text{Cos}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(4*d)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx &= \frac{(b^2\sqrt{b\cos(c+dx)}) \int \cos^4(c + dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}\sin(c+dx)}{4d} + \frac{(3b^2\sqrt{b\cos(c+dx)}) \int \cos^2(c+dx) dx}{4\sqrt{\cos(c+dx)}} \\ &= \frac{3b^2\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{8d} + \frac{b^2\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}}{4d} \\ &= \frac{3b^2x\sqrt{b\cos(c+dx)}}{8\sqrt{\cos(c+dx)}} + \frac{3b^2\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{8d} + \frac{b^2\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}}{4d} \end{aligned}$$

pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2
*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to c
heck sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/
2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x
/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check
sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Una
ble to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(
-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign:
(2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable t
o check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi
/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*
pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to che
ck sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)
Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2
)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check si
gn: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unabl
e to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2
*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (
2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to
check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x
/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/
x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check
sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Un
able to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>
(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign
: (2*pi/x/2)>(-2*pi/x/2)Unable to check sign: (2*pi/x/2)>(-2*pi/x/2)(3*b^2*
sqrt(b)*d*x*tan((c+d*x)/2)^8+12*b^2*sqrt(b)*d*x*tan((c+d*x)/2)^6+18*b^2*sqrt
(b)*d*x*tan((c+d*x)/2)^4+12*b^2*sqrt(b)*d*x*tan((c+d*x)/2)^2+3*b^2*sqrt(b)
*d*x+(-10*b^2*sqrt(b))*tan((c+d*x)/2)^7+6*b^2*sqrt(b)*tan((c+d*x)/2)^5+(-6*
b^2*sqrt(b))*tan((c+d*x)/2)^3+10*b^2*sqrt(b)*tan((c+d*x)/2))/(8*d*tan((c+d*
x)/2)^8+32*d*tan((c+d*x)/2)^6+48*d*tan((c+d*x)/2)^4+32*d*tan((c+d*x)/2)^2+8
*d)

maple [A] time = 0.12, size = 62, normalized size = 0.58

$$\frac{(b \cos(dx + c))^{\frac{5}{2}} \left(2 \left(\cos^3(dx + c) \right) \sin(dx + c) + 3 \cos(dx + c) \sin(dx + c) + 3dx + 3c \right)}{8d \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(5/2),x)

[Out] 1/8/d*(b*cos(d*x+c))^(5/2)*(2*cos(d*x+c)^3*sin(d*x+c)+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)/cos(d*x+c)^(5/2)

maxima [A] time = 1.10, size = 59, normalized size = 0.55

$$\frac{\left(12(dx + c)b^2 + b^2 \sin(4dx + 4c) + 8b^2 \sin\left(\frac{1}{2} \arctan(\sin(4dx + 4c), \cos(4dx + 4c))\right) \right) \sqrt{b}}{32d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/32*(12*(d*x + c)*b^2 + b^2*sin(4*d*x + 4*c) + 8*b^2*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d

mupad [B] time = 1.03, size = 78, normalized size = 0.73

$$\frac{b^2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx) + 24dx \cos(c + dx))}{32d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(5/2), x)
```

```
[Out] (b^2*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c
+ 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*d*(cos(2*c + 2*d*x)
+ 1))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**(3/2)*(b*cos(d*x+c))**(5/2), x)
```

```
[Out] Timed out
```

3.162 $\int \sqrt{\cos(c + dx)} (b \cos(c + dx))^{5/2} dx$

Optimal. Leaf size=76

$$\frac{b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{b^2 \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}}$$

[Out] $b^2 \sin(dx+c) * (b \cos(dx+c))^{(1/2)} / d / \cos(dx+c)^{(1/2)} - 1/3 * b^2 \sin(dx+c)^3 * (b \cos(dx+c))^{(1/2)} / d / \cos(dx+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{b^2 \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(5/2),x]

[Out] $(b^2 \text{Sqrt}[b \text{Cos}[c + d*x]] * \text{Sin}[c + d*x]) / (d \text{Sqrt}[\text{Cos}[c + d*x]]) - (b^2 \text{Sqrt}[b \text{Cos}[c + d*x]] * \text{Sin}[c + d*x]^3) / (3 * d \text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \sqrt{\cos(c + dx)} (b \cos(c + dx))^{5/2} dx &= \frac{(b^2 \sqrt{b \cos(c + dx)}) \int \cos^3(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{(b^2 \sqrt{b \cos(c + dx)}) \text{Subst}\left(\int (1 - x^2) dx, x, -\sin(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{b^2 \sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.13, size = 45, normalized size = 0.59

$$\frac{\sin(c + dx)(\cos(2(c + dx)) + 5)(b \cos(c + dx))^{5/2}}{6d \cos^2(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(5/2),x]

[Out] $((b \text{Cos}[c + d*x])^{(5/2)} * (5 + \text{Cos}[2*(c + d*x)]) * \text{Sin}[c + d*x]) / (6 * d \text{Cos}[c + d*x]^{(5/2)})$

fricas [A] time = 0.73, size = 47, normalized size = 0.62

$$\frac{(b^2 \cos(dx + c)^2 + 2b^2) \sqrt{b \cos(dx + c)} \sin(dx + c)}{3d \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/3*(b^2*cos(d*x + c)^2 + 2*b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] Timed out

maple [A] time = 0.11, size = 40, normalized size = 0.53

$$\frac{(2 + \cos^2(dx + c)) \sin(dx + c) (b \cos(dx + c))^{\frac{5}{2}}}{3d \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(5/2),x)

[Out] 1/3/d*(2+cos(d*x+c)^2)*sin(d*x+c)*(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2)

maxima [A] time = 1.23, size = 49, normalized size = 0.64

$$\frac{\left(b^2 \sin(3dx + 3c) + 9b^2 \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)\right) \sqrt{b}}{12d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/12*(b^2*sin(3*d*x + 3*c) + 9*b^2*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*sqrt(b)/d

mupad [B] time = 0.62, size = 60, normalized size = 0.79

$$\frac{b^2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(5/2),x)

[Out] (b^2*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(1/2)*(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

$$3.163 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\sqrt{\cos(c+dx)}} dx$$

Optimal. Leaf size=69

$$\frac{b^2 x \sqrt{b \cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b^2 \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}{2d}$$

[Out] $1/2*b^2*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+1/2*b^2*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.02, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{b^2 x \sqrt{b \cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b^2 \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}{2d}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(5/2)/Sqrt[Cos[c + d*x]], x]

[Out] $(b^2*x*\text{Sqrt}[b*\text{Cos}[c + d*x]])/(2*\text{Sqrt}[\text{Cos}[c + d*x]]) + (b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(2*d)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{5/2}}{\sqrt{\cos(c+dx)}} dx &= \frac{(b^2 \sqrt{b \cos(c+dx)}) \int \cos^2(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} \sin(c+dx)}{2d} + \frac{(b^2 \sqrt{b \cos(c+dx)}) \int 1 dx}{2\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 x \sqrt{b \cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b^2 \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} \sin(c+dx)}{2d} \end{aligned}$$

Mathematica [A] time = 0.07, size = 45, normalized size = 0.65

$$\frac{(2(c+dx) + \sin(2(c+dx)))(b \cos(c+dx))^{5/2}}{4d \cos^2(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(5/2)/Sqrt[Cos[c + d*x]],x]

[Out] ((b*cos[c + d*x])^(5/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*cos[c + d*x]^(5/2))

fricas [A] time = 1.04, size = 159, normalized size = 2.30

$$\frac{2\sqrt{b\cos(dx+c)}b^2\sqrt{\cos(dx+c)}\sin(dx+c) + \sqrt{-b}b^2\log\left(2b\cos(dx+c)^2 - 2\sqrt{b\cos(dx+c)}\sqrt{-b}\sqrt{\cos(dx+c)}\right)}{4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(1/2),x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b*cos(d*x + c))*b^2*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(-b)*b^2*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/d, 1/2*(sqrt(b*cos(d*x + c))*b^2*sqrt(cos(d*x + c))*sin(d*x + c) + b^(5/2)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{5}{2}}}{\sqrt{\cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/sqrt(cos(d*x + c)), x)

maple [A] time = 0.12, size = 42, normalized size = 0.61

$$\frac{(\cos(dx + c)\sin(dx + c) + dx + c)(b\cos(dx + c))^{\frac{5}{2}}}{2d\cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(1/2),x)

[Out] 1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)*(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2)

maxima [A] time = 1.09, size = 32, normalized size = 0.46

$$\frac{(2(dx + c)b^2 + b^2\sin(2dx + 2c))\sqrt{b}}{4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(1/2),x, algorithm="maxima")

[Out] 1/4*(2*(d*x + c)*b^2 + b^2*sin(2*d*x + 2*c))*sqrt(b)/d

mupad [B] time = 0.44, size = 40, normalized size = 0.58

$$\frac{b^2\sqrt{b\cos(c + dx)}(\sin(2c + 2dx) + 2dx)}{4d\sqrt{\cos(c + dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(1/2),x)
```

```
[Out] (b^2*(b*cos(c + d*x))^(1/2)*(sin(2*c + 2*d*x) + 2*d*x))/(4*d*cos(c + d*x)^(1/2))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(1/2),x)
```

```
[Out] Timed out
```

$$3.164 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{3}{2}}(c+dx)} dx$$

Optimal. Leaf size=35

$$\frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}}$$

[Out] $b^2 \sin(d*x+c) * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b \cos[c + d*x])^{(5/2)} / \cos[c + d*x]^{(3/2)}, x]$

[Out] $(b^2 \sqrt{b \cos[c + d*x]} * \sin[c + d*x]) / (d \sqrt{\cos[c + d*x]})$

Rule 17

$\text{Int}[(u_.) * ((a_.) * (v_))^{(m_)} * ((b_.) * (v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)} * b^{(n-1/2)} * \sqrt{b*v}) / \sqrt{a*v}, \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.) * (x_)], x_Symbol] \rightarrow \text{Simp}[\sin[c + d*x] / d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{3}{2}}(c+dx)} dx &= \frac{(b^2 \sqrt{b \cos(c+dx)}) \int \cos(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 32, normalized size = 0.91

$$\frac{\sin(c+dx)(b \cos(c+dx))^{5/2}}{d \cos^{\frac{5}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[(b \cos[c + d*x])^{(5/2)} / \cos[c + d*x]^{(3/2)}, x]$

[Out] $((b \cos[c + d*x])^{(5/2)} * \sin[c + d*x]) / (d * \cos[c + d*x]^{(5/2)})$

fricas [A] time = 0.53, size = 31, normalized size = 0.89

$$\frac{\sqrt{b \cos(dx+c)} b^2 \sin(dx+c)}{d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(3/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*b^2*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{5}{2}}}{\cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(3/2), x)

maple [A] time = 0.08, size = 29, normalized size = 0.83

$$\frac{(b \cos(dx + c))^{\frac{5}{2}} \sin(dx + c)}{d \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(3/2),x)

[Out] 1/d*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/cos(d*x+c)^(5/2)

maxima [A] time = 1.48, size = 13, normalized size = 0.37

$$\frac{b^{\frac{5}{2}} \sin(dx + c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(3/2),x, algorithm="maxima")

[Out] b^(5/2)*sin(d*x + c)/d

mupad [B] time = 0.32, size = 31, normalized size = 0.89

$$\frac{b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(3/2),x)

[Out] (b^2*sin(c + d*x)*(b*cos(c + d*x))^(1/2))/(d*cos(c + d*x)^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(3/2),x)

[Out] Timed out

$$3.165 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx$$

Optimal. Leaf size=27

$$\frac{b^2 x \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[Out] $b^2 x (b \cos(d x + c))^{1/2} / \cos(d x + c)^{1/2}$

Rubi [A] time = 0.00, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{b^2 x \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(5/2), x]

[Out] (b^2*x*Sqrt[b*Cos[c + d*x]])/Sqrt[Cos[c + d*x]]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx &= \frac{(b^2 \sqrt{b \cos(c+dx)}) \int 1 dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 x \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 0.89

$$\frac{x(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(5/2), x]

[Out] (x*(b*Cos[c + d*x])^(5/2))/Cos[c + d*x]^(5/2)

fricas [A] time = 0.65, size = 97, normalized size = 3.59

$$\left[\frac{\sqrt{-b} b^2 \log(2 b \cos(dx+c)^2 - 2 \sqrt{b \cos(dx+c)} \sqrt{-b} \sqrt{\cos(dx+c)} \sin(dx+c) - b)}{2 d}, \frac{b^{5/2} \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{\sqrt{b} \cos(dx+c)}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2),x, algorithm="fricas")

[Out] [1/2*sqrt(-b)*b^2*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b)/d, b^(5/2)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{5}{2}}}{\cos(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(5/2), x)

maple [A] time = 0.06, size = 28, normalized size = 1.04

$$\frac{(b \cos(dx + c))^{\frac{5}{2}} (dx + c)}{d \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2),x)

[Out] 1/d*(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2)*(d*x+c)

maxima [A] time = 1.20, size = 26, normalized size = 0.96

$$\frac{2 b^{\frac{5}{2}} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2),x, algorithm="maxima")

[Out] 2*b^(5/2)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d

mupad [B] time = 0.09, size = 23, normalized size = 0.85

$$\frac{b^2 x \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(5/2),x)

[Out] (b^2*x*(b*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(5/2),x)

[Out] Timed out

$$3.166 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx$$

Optimal. Leaf size=36

$$\frac{b^2 \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d \sqrt{\cos(c+dx)}}$$

[Out] $b^2 \operatorname{arctanh}(\sin(dx+c)) * (b \cos(dx+c))^{1/2} / d / \cos(dx+c)^{1/2}$

Rubi [A] time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{b^2 \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b \cos[c + dx])^{5/2} / \cos[c + dx]^{7/2}, x]$

[Out] $(b^2 \operatorname{ArcTanh}[\sin[c + dx]] * \operatorname{Sqrt}[b \cos[c + dx]]) / (d * \operatorname{Sqrt}[\cos[c + dx]])$

Rule 17

$\text{Int}[(u_*) * ((a_*) * (v_*))^{(m_*)} * ((b_*) * (v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)} * b^{(n-1/2)} * \operatorname{Sqrt}[b*v]) / \operatorname{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ $\text{FreeQ}[\{a, b, m\}, x] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n + 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

Rule 3770

$\text{Int}[\csc[(c_*) + (d_*) * (x_*)], x_Symbol] \rightarrow -\text{Simp}[\operatorname{ArcTanh}[\cos[c + dx]] / d, x] /;$ $\text{FreeQ}[\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx &= \frac{(b^2 \sqrt{b \cos(c+dx)}) \int \sec(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \tanh^{-1}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 33, normalized size = 0.92

$$\frac{(b \cos(c+dx))^{5/2} \tanh^{-1}(\sin(c+dx))}{d \cos^{5/2}(c+dx)}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[(b \cos[c + dx])^{5/2} / \cos[c + dx]^{7/2}, x]$

[Out] $(\operatorname{ArcTanh}[\sin[c + dx]] * (b \cos[c + dx])^{5/2}) / (d \cos[c + dx]^{5/2})$

fricas [A] time = 0.55, size = 116, normalized size = 3.22

$$\left[\frac{b^{5/2} \log\left(-\frac{b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2 b \cos(dx+c)}{\cos(dx+c)^3}\right)}{2d}, -\frac{\sqrt{-b} b^2 \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b} \sin(dx+c)}{b \sqrt{\cos(dx+c)}}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(7/2),x, algorithm="fricas")

[Out] [1/2*b^(5/2)*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3)/d, -sqrt(-b)*b^2*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{5/2}}{\cos(dx + c)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(7/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(7/2), x)

maple [A] time = 0.08, size = 42, normalized size = 1.17

$$\frac{2 \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right) (b \cos(dx + c))^{5/2}}{d \cos(dx + c)^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(7/2),x)

[Out] -2/d*arctanh((-1+cos(d*x+c))/sin(d*x+c))*(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2)

maxima [B] time = 1.36, size = 72, normalized size = 2.00

$$\frac{(b^2 \log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - b^2 \log(\cos(dx + c)^2 + \sin(dx + c)^2 - 2 \sin(dx + c) + 1)) \sqrt{b}}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(7/2),x, algorithm="maxima")

[Out] 1/2*(b^2*log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - b^2*log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))*sqrt(b)/d

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(7/2),x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(7/2),x)

[Out] Timed out

$$3.167 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx$$

Optimal. Leaf size=35

$$\frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[Out] $b^2 \sin(dx+c) (b \cos(dx+c))^{1/2} / d / \cos(dx+c)^{3/2}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b \cos[c + d*x])^{5/2} / \cos[c + d*x]^{9/2}, x]$

[Out] $(b^2 \sqrt{b \cos[c + d*x]} \sin[c + d*x]) / (d \cos[c + d*x]^{3/2})$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.) * ((a_.) * (v_.))^{(m_.)} * ((b_.) * (v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)} * b^{(n-1/2)} * \text{Sqrt}[b*v]) / \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}\{a, b, m\}, x \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.) * (x_.)]^{(n_.)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1+x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c+d*x]], x] /; \text{FreeQ}\{c, d\}, x \&\& \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx &= \frac{(b^2 \sqrt{b \cos(c+dx)}) \int \sec^2(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{(b^2 \sqrt{b \cos(c+dx)}) \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{d \sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 0.91

$$\frac{\sin(c+dx)(b \cos(c+dx))^{5/2}}{d \cos^{\frac{7}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(5/2)/cos[c + d*x]^(9/2),x]

[Out] ((b*cos[c + d*x])^(5/2)*sin[c + d*x])/(d*cos[c + d*x]^(7/2))

fricas [A] time = 0.62, size = 31, normalized size = 0.89

$$\frac{\sqrt{b \cos(dx + c)} b^2 \sin(dx + c)}{d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(9/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*b^2*sin(d*x + c)/(d*cos(d*x + c)^(3/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{5}{2}}}{\cos(dx + c)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(9/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(9/2), x)

maple [A] time = 0.08, size = 29, normalized size = 0.83

$$\frac{(b \cos(dx + c))^{\frac{5}{2}} \sin(dx + c)}{d \cos(dx + c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(9/2),x)

[Out] 1/d*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/cos(d*x+c)^(7/2)

maxima [A] time = 1.12, size = 54, normalized size = 1.54

$$\frac{2 b^{\frac{5}{2}} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c)^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1) d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(9/2),x, algorithm="maxima")

[Out] 2*b^(5/2)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 0.42, size = 62, normalized size = 1.77

$$\frac{b^2 \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 1i + \sin(2c + 2dx) + 1i)}{d \sqrt{\cos(c + dx)} (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(9/2),x)

[Out] (b^2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/(d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(9/2),x)

[Out] Timed out

$$3.168 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{11}{2}}(c+dx)} dx$$

Optimal. Leaf size=78

$$\frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{b^2 \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2d \sqrt{\cos(c+dx)}}$$

[Out] $1/2*b^2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(5/2)}+1/2*b^2*\operatorname{arctanh}(\sin(d*x+c))*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{b^2 \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Cos}[c+d*x])^{(5/2)}/\text{Cos}[c+d*x]^{(11/2)},x]$

[Out] $(b^2*\text{ArcTanh}[\text{Sin}[c+d*x]]*\text{Sqrt}[b*\text{Cos}[c+d*x]])/(2*d*\text{Sqrt}[\text{Cos}[c+d*x]]) + (b^2*\text{Sqrt}[b*\text{Cos}[c+d*x]]*\text{Sin}[c+d*x])/(2*d*\text{Cos}[c+d*x]^{(5/2)})$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/\text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n+1/2, 0] && IntegerQ[m+n]

Rule 3768

$\text{Int}[(\text{csc}[c_.) + (d_.)*(x_)]*(b_.)^{(n_)}, x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c+dx])*(b*\text{Csc}[c+dx])^{(n-1)})/(d*(n-1)), x] + \text{Dist}[(b^{(n-2)})/(n-1), \text{Int}[(b*\text{Csc}[c+dx])^{(n-2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

$\text{Int}[\text{csc}[c_.) + (d_.)*(x_)], x_Symbol] \rightarrow -\text{Simp}[\text{ArcTanh}[\text{Cos}[c+dx]]/d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{11}{2}}(c+dx)} dx &= \frac{(b^2 \sqrt{b \cos(c+dx)}) \int \sec^3(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx)} + \frac{(b^2 \sqrt{b \cos(c+dx)}) \int \sec(c+dx) dx}{2\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \tanh^{-1}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{2d \sqrt{\cos(c+dx)}} + \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.05, size = 52, normalized size = 0.67

$$\frac{(b \cos(c + dx))^{5/2} (\sin(c + dx) + \cos^2(c + dx) \tanh^{-1}(\sin(c + dx)))}{2d \cos^{\frac{9}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(11/2), x]

[Out] ((b*Cos[c + d*x])^(5/2)*(ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x]))/(2*d*Cos[c + d*x]^(9/2))

fricas [A] time = 0.54, size = 210, normalized size = 2.69

$$\left[\frac{b^{\frac{5}{2}} \cos(dx + c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2 \sqrt{b \cos(dx+c)} b^2 \sqrt{\cos(dx+c)}}{4 d \cos(dx+c)^3} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(11/2), x, algorithm="fricas")

[Out] [1/4*(b^(5/2)*cos(d*x + c)^3*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*b^2*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3), -1/2*(sqrt(-b)*b^2*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*b^2*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{5}{2}}}{\cos(dx + c)^{\frac{11}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(11/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(11/2), x)

maple [A] time = 0.11, size = 104, normalized size = 1.33

$$\frac{\left((\cos^2(dx + c)) \ln\left(-\frac{-1 + \cos(dx+c) + \sin(dx+c)}{\sin(dx+c)}\right) - (\cos^2(dx + c)) \ln\left(-\frac{-\sin(dx+c) - 1 + \cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx + c) \right) (b \cos(dx + c))^{5/2}}{2d \cos(dx + c)^{\frac{9}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(11/2), x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c))*(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(9/2)

maxima [B] time = 1.83, size = 747, normalized size = 9.58

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(11/2), x, algorithm="maxima")

```
[Out] -1/4*(4*(b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + (b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 4*(b^2*cos(4*d*x + 4*c) + 2*b^2*cos(2*d*x + 2*c) + b^2)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(b^2*cos(4*d*x + 4*c) + 2*b^2*cos(2*d*x + 2*c) + b^2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^{11/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(11/2), x)
```

```
[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(11/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(11/2), x)
```

```
[Out] Timed out
```

$$3.169 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{13/2}(c+dx)} dx$$

Optimal. Leaf size=76

$$\frac{b^2 \sin^3(c+dx) \sqrt{b \cos(c+dx)}}{3d \cos^{7/2}(c+dx)} + \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{3/2}(c+dx)}$$

[Out] $b^2 \sin(d*x+c) * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(3/2)} + 1/3 * b^2 \sin(d*x+c)^3 * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(7/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3767}

$$\frac{b^2 \sin^3(c+dx) \sqrt{b \cos(c+dx)}}{3d \cos^{7/2}(c+dx)} + \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{3/2}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(5/2)/Cos[c + d*x]^(13/2), x]

[Out] $(b^2 \sqrt{b \cos[c + d*x]} \sin[c + d*x]) / (d \cos[c + d*x]^{(3/2)}) + (b^2 \sqrt{b \cos[c + d*x]} \sin^3[c + d*x]) / (3d \cos[c + d*x]^{(7/2)})$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{13/2}(c+dx)} dx &= \frac{(b^2 \sqrt{b \cos(c+dx)}) \int \sec^4(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= -\frac{(b^2 \sqrt{b \cos(c+dx)}) \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d \sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{3/2}(c+dx)} + \frac{b^2 \sqrt{b \cos(c+dx)} \sin^3(c+dx)}{3d \cos^{7/2}(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.05, size = 45, normalized size = 0.59

$$\frac{\left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx)\right) (b \cos(c+dx))^{5/2}}{d \cos^{5/2}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(5/2)/cos[c + d*x]^(13/2), x]

[Out] ((b*cos[c + d*x])^(5/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*cos[c + d*x]^(5/2))

fricas [A] time = 0.59, size = 46, normalized size = 0.61

$$\frac{(2b^2 \cos(dx + c)^2 + b^2) \sqrt{b \cos(dx + c)} \sin(dx + c)}{3d \cos(dx + c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(13/2), x, algorithm="fricas")

[Out] 1/3*(2*b^2*cos(d*x + c)^2 + b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(7/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{\frac{5}{2}}}{\cos(dx + c)^{\frac{13}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(13/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(13/2), x)

maple [A] time = 0.10, size = 42, normalized size = 0.55

$$\frac{(2(\cos^2(dx + c)) + 1)(b \cos(dx + c))^{\frac{5}{2}} \sin(dx + c)}{3d \cos(dx + c)^{\frac{11}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(13/2), x)

[Out] 1/3/d*(2*cos(d*x+c)^2+1)*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/cos(d*x+c)^(11/2)

maxima [B] time = 1.26, size = 311, normalized size = 4.09

$$\frac{4(3b^2 \cos(6dx + 6c) + \dots)}{3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \dots)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(13/2), x, algorithm="maxima")

[Out] -4/3*(3*b^2*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) + 9*b^2*cos(4*d*x + 4*c)*sin(2*d*x + 2*c) - (3*b^2*cos(2*d*x + 2*c) + b^2)*sin(6*d*x + 6*c) - 3*(3*b^2*cos(2*d*x + 2*c) + b^2)*sin(4*d*x + 4*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 1.18, size = 131, normalized size = 1.72

$$\frac{2b^2 \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 15i + \cos(4c + 4dx) 6i + \cos(6c + 6dx) 1i + 9 \sin(2c + 2dx) + 6 \dots)}{3d \sqrt{\cos(c + dx)} (15 \cos(2c + 2dx) + 6 \cos(4c + 4dx) + \cos(6c + 6dx) + \dots)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(13/2),x)
```

```
[Out] (2*b^2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i +
cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c +
6*d*x) + 10i))/(3*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4
*d*x) + cos(6*c + 6*d*x) + 10))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(13/2),x)
```

```
[Out] Timed out
```

$$3.170 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{15}{2}}(c+dx)} dx$$

Optimal. Leaf size=116

$$\frac{3b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{\frac{5}{2}}(c+dx)} + \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3b^2 \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8d \sqrt{\cos(c+dx)}}$$

[Out] 1/4*b^2*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(9/2)+3/8*b^2*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+3/8*b^2*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{3b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{\frac{5}{2}}(c+dx)} + \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3b^2 \sqrt{b \cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8d \sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(15/2), x]

[Out] (3*b^2*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(8*d*Sqrt[Cos[c + d*x]]) + (b^2*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(4*d*Cos[c + d*x]^(9/2)) + (3*b^2*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(8*d*Cos[c + d*x]^(5/2))

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{15/2}(c + dx)} dx &= \frac{(b^2 \sqrt{b \cos(c + dx)}) \int \sec^5(c + dx) dx}{\sqrt{\cos(c + dx)}} \\
&= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{9/2}(c + dx)} + \frac{(3b^2 \sqrt{b \cos(c + dx)}) \int \sec^3(c + dx) dx}{4\sqrt{\cos(c + dx)}} \\
&= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{9/2}(c + dx)} + \frac{3b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d \cos^{5/2}(c + dx)} + \frac{(3b^2 \sqrt{b \cos(c + dx)})}{8\sqrt{\cos(c + dx)}} \\
&= \frac{3b^2 \tanh^{-1}(\sin(c + dx)) \sqrt{b \cos(c + dx)}}{8d \sqrt{\cos(c + dx)}} + \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{9/2}(c + dx)} + \frac{3b^2 \sqrt{b \cos(c + dx)}}{8d \cos^{5/2}(c + dx)}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 66, normalized size = 0.57

$$\frac{(b \cos(c + dx))^{5/2} (\sin(c + dx) (3 \cos^2(c + dx) + 2) + 3 \cos^4(c + dx) \tanh^{-1}(\sin(c + dx)))}{8d \cos^{13/2}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(15/2), x]

[Out] ((b*Cos[c + d*x])^(5/2)*(3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x]))/(8*d*Cos[c + d*x]^(13/2))

fricas [A] time = 0.65, size = 244, normalized size = 2.10

$$\left[\frac{3 b^2 \cos(dx + c)^5 \log\left(-\frac{b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2 b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2 (3 b^2 \cos(dx + c)^2 + 2 b^2) \sqrt{b \cos(dx+c)}}{16 d \cos(dx + c)^5} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(15/2), x, algorithm="fricas")

[Out] [1/16*(3*b^(5/2)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*(3*b^2*cos(d*x + c)^2 + 2*b^2)*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^5), -1/8*(3*sqrt(-b)*b^2*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^5 - (3*b^2*cos(d*x + c)^2 + 2*b^2)*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^{5/2}}{\cos(dx + c)^{15/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(15/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(15/2), x)

maple [A] time = 0.16, size = 121, normalized size = 1.04

$$\frac{3 (\cos^4(dx + c)) \ln\left(-\frac{\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - 3 (\cos^4(dx + c)) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) + 3 (\cos^2(dx + c)) \sin(dx+c)}{8d \cos(dx + c)^{13/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(15/2),x)
```

```
[Out] 1/8/d*(3*cos(d*x+c)^4*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^4*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))*(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(13/2)
```

maxima [B] time = 1.58, size = 1914, normalized size = 16.50

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(15/2),x, algorithm="maxima")
```

```
[Out] -1/16*(12*(b^2*sin(8*d*x + 8*c) + 4*b^2*sin(6*d*x + 6*c) + 6*b^2*sin(4*d*x + 4*c) + 4*b^2*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(b^2*sin(8*d*x + 8*c) + 4*b^2*sin(6*d*x + 6*c) + 6*b^2*sin(4*d*x + 4*c) + 4*b^2*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(b^2*sin(8*d*x + 8*c) + 4*b^2*sin(6*d*x + 6*c) + 6*b^2*sin(4*d*x + 4*c) + 4*b^2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(b^2*sin(8*d*x + 8*c) + 4*b^2*sin(6*d*x + 6*c) + 6*b^2*sin(4*d*x + 4*c) + 4*b^2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(b^2*cos(8*d*x + 8*c)^2 + 16*b^2*cos(6*d*x + 6*c)^2 + 36*b^2*cos(4*d*x + 4*c)^2 + 16*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(8*d*x + 8*c)^2 + 16*b^2*sin(6*d*x + 6*c)^2 + 36*b^2*sin(4*d*x + 4*c)^2 + 48*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b^2*sin(2*d*x + 2*c)^2 + 8*b^2*cos(2*d*x + 2*c) + b^2 + 2*(4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*cos(8*d*x + 8*c) + 8*(6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*cos(6*d*x + 6*c) + 12*(4*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c) + 4*(2*b^2*sin(6*d*x + 6*c) + 3*b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + 3*(b^2*cos(8*d*x + 8*c)^2 + 16*b^2*cos(6*d*x + 6*c)^2 + 36*b^2*cos(4*d*x + 4*c)^2 + 16*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(8*d*x + 8*c)^2 + 16*b^2*sin(6*d*x + 6*c)^2 + 36*b^2*sin(4*d*x + 4*c)^2 + 48*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b^2*sin(2*d*x + 2*c)^2 + 8*b^2*cos(2*d*x + 2*c) + b^2 + 2*(4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*cos(8*d*x + 8*c) + 8*(6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*cos(6*d*x + 6*c) + 12*(4*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c) + 4*(2*b^2*sin(6*d*x + 6*c) + 3*b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 12*(b^2*cos(8*d*x + 8*c) + 4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(b^2*cos(8*d*x + 8*c) + 4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(b^2*cos(8*d*x + 8*c) + 4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 12*(b^2*cos(8*d*x + 8*c) + 4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) +
```

```

2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*
x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 3
6*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x
+ 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*d)

```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^{15/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(15/2), x)
```

```
[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(15/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(15/2), x)
```

```
[Out] Timed out
```

$$3.171 \quad \int \frac{\cos^{\frac{11}{2}}(c+dx)}{\sqrt{b} \cos(c+dx)} dx$$

Optimal. Leaf size=107

$$\frac{\sin^5(c+dx)\sqrt{\cos(c+dx)}}{5d\sqrt{b}\cos(c+dx)} - \frac{2\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3d\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b}\cos(c+dx)}$$

[Out] sin(d*x+c)*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)-2/3*sin(d*x+c)^3*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+1/5*sin(d*x+c)^5*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.02, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{\sin^5(c+dx)\sqrt{\cos(c+dx)}}{5d\sqrt{b}\cos(c+dx)} - \frac{2\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3d\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(11/2)/Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[b*Cos[c + d*x]]) - (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x]^3)/(3*d*Sqrt[b*Cos[c + d*x]]) + (Sqrt[Cos[c + d*x]]*Sin[c + d*x]^5)/(5*d*Sqrt[b*Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\cos^{\frac{11}{2}}(c+dx)}{\sqrt{b} \cos(c+dx)} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^5(c+dx) dx}{\sqrt{b} \cos(c+dx)} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1-2x^2+x^4) dx, x, -\sin(c+dx)\right)}{d\sqrt{b}\cos(c+dx)} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{b}\cos(c+dx)} - \frac{2\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3d\sqrt{b}\cos(c+dx)} + \frac{\sqrt{\cos(c+dx)} \sin^5(c+dx)}{5d\sqrt{b}\cos(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.10, size = 57, normalized size = 0.53

$$\frac{\sin(c+dx) \left(3 \sin^4(c+dx) - 10 \sin^2(c+dx) + 15\right) \sqrt{\cos(c+dx)}}{15d\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(11/2)/Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x]*(15 - 10*Sin[c + d*x]^2 + 3*Sin[c + d*x]^4))/(15*d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.67, size = 54, normalized size = 0.50

$$\frac{(3 \cos(dx + c)^4 + 4 \cos(dx + c)^2 + 8)\sqrt{b \cos(dx + c)} \sin(dx + c)}{15bd\sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/15*(3*cos(d*x + c)^4 + 4*cos(d*x + c)^2 + 8)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{11}{2}}}{\sqrt{b \cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(11/2)/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.12, size = 52, normalized size = 0.49

$$\frac{(3(\cos^4(dx + c)) + 4(\cos^2(dx + c)) + 8) \sin(dx + c) (\sqrt{\cos(dx + c)})}{15d\sqrt{b \cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(1/2),x)

[Out] 1/15/d*(3*cos(d*x+c)^4+4*cos(d*x+c)^2+8)*sin(d*x+c)*cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2)

maxima [A] time = 1.43, size = 68, normalized size = 0.64

$$\frac{3 \sin(5dx + 5c) + 25 \sin\left(\frac{3}{5} \arctan(\sin(5dx + 5c), \cos(5dx + 5c))\right) + 150 \sin\left(\frac{1}{5} \arctan(\sin(5dx + 5c), \cos(5dx + 5c))\right)}{240 \sqrt{bd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/240*(3*sin(5*d*x + 5*c) + 25*sin(3/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 150*sin(1/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))))/(sqrt(b)*d)

mupad [B] time = 1.18, size = 73, normalized size = 0.68

$$\frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (175 \sin(2c + 2dx) + 28 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}{240bd(\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(11/2)/(b*cos(c + d*x))^(1/2),x)

```
[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(175*sin(2*c + 2*d*x) + 28*sin(4*c + 4*d*x) + 3*sin(6*c + 6*d*x)))/(240*b*d*(cos(2*c + 2*d*x) + 1))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**(11/2)/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

$$3.172 \quad \int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b} \cos(c+dx)} dx$$

Optimal. Leaf size=98

$$\frac{3x\sqrt{\cos(c+dx)}}{8\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{4d\sqrt{b}\cos(c+dx)} + \frac{3\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{8d\sqrt{b}\cos(c+dx)}$$

[Out] $3/8*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}+1/4*\cos(d*x+c)^{(7/2)}*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}+3/8*x*\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{3x\sqrt{\cos(c+dx)}}{8\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{4d\sqrt{b}\cos(c+dx)} + \frac{3\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{8d\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(9/2)/Sqrt[b*Cos[c + d*x]], x]

[Out] $(3*x*\text{Sqrt}[\text{Cos}[c + d*x]])/(8*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (3*\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(8*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(7/2)}*\text{Sin}[c + d*x])/(4*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b} \cos(c+dx)} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^4(c+dx) dx}{\sqrt{b} \cos(c+dx)} \\ &= \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4d\sqrt{b}\cos(c+dx)} + \frac{(3\sqrt{\cos(c+dx)}) \int \cos^2(c+dx) dx}{4\sqrt{b}\cos(c+dx)} \\ &= \frac{3\cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8d\sqrt{b}\cos(c+dx)} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4d\sqrt{b}\cos(c+dx)} + \frac{(3\sqrt{\cos(c+dx)}) \int 1 dx}{8\sqrt{b}\cos(c+dx)} \\ &= \frac{3x\sqrt{\cos(c+dx)}}{8\sqrt{b}\cos(c+dx)} + \frac{3\cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8d\sqrt{b}\cos(c+dx)} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4d\sqrt{b}\cos(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.07, size = 55, normalized size = 0.56

$$\frac{(12(c + dx) + 8 \sin(2(c + dx)) + \sin(4(c + dx)))\sqrt{\cos(c + dx)}}{32d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(9/2)/Sqrt[b*Cos[c + d*x]], x]

[Out] (Sqrt[Cos[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])) / (32*d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.78, size = 182, normalized size = 1.86

$$\left[\frac{2\sqrt{b \cos(dx + c)}(2 \cos(dx + c)^2 + 3)\sqrt{\cos(dx + c)} \sin(dx + c) - 3\sqrt{-b} \log(2b \cos(dx + c)^2 + 2\sqrt{b \cos(dx + c)})}{16bd} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [1/16*(2*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 3)*sqrt(cos(d*x + c))*sin(d*x + c) - 3*sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b*d), 1/8*(sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 3)*sqrt(cos(d*x + c))*sin(d*x + c) + 3*sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))]/(b*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{9}{2}}}{\sqrt{b \cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(9/2)/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.15, size = 62, normalized size = 0.63

$$\frac{(2(\cos^3(dx + c)) \sin(dx + c) + 3 \cos(dx + c) \sin(dx + c) + 3dx + 3c)(\sqrt{\cos(dx + c)})}{8d\sqrt{b \cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2), x)

[Out] 1/8/d*(2*cos(d*x+c)^3*sin(d*x+c)+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)*cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2)

maxima [A] time = 1.34, size = 49, normalized size = 0.50

$$\frac{12dx + 12c + \sin(4dx + 4c) + 8 \sin\left(\frac{1}{2} \arctan(\sin(4dx + 4c), \cos(4dx + 4c))\right)}{32\sqrt{bd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2), x, algorithm="maxima")

[Out] $\frac{1}{32} \cdot (12 \cdot d \cdot x + 12 \cdot c + \sin(4 \cdot d \cdot x + 4 \cdot c) + 8 \cdot \sin(\frac{1}{2} \cdot \arctan2(\sin(4 \cdot d \cdot x + 4 \cdot c), \cos(4 \cdot d \cdot x + 4 \cdot c)))) / (\sqrt{b} \cdot d)$

mupad [B] time = 1.06, size = 78, normalized size = 0.80

$$\frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx) + 24 dx \cos(c + dx))}{32bd (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^(9/2)/(b*cos(c + d*x))^(1/2), x)`

[Out] $(\cos(c + d \cdot x)^{(1/2)} \cdot (b \cdot \cos(c + d \cdot x))^{(1/2)} \cdot (8 \cdot \sin(c + d \cdot x) + 9 \cdot \sin(3 \cdot c + 3 \cdot d \cdot x) + \sin(5 \cdot c + 5 \cdot d \cdot x) + 24 \cdot d \cdot x \cdot \cos(c + d \cdot x))) / (32 \cdot b \cdot d \cdot (\cos(2 \cdot c + 2 \cdot d \cdot x) + 1))$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**(9/2)/(b*cos(d*x+c))**(1/2), x)`

[Out] Timed out

$$3.173 \quad \int \frac{\cos^7(c+dx)}{\sqrt{b} \cos(c+dx)} dx$$

Optimal. Leaf size=70

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b}\cos(c+dx)} - \frac{\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3d\sqrt{b}\cos(c+dx)}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b}\cos(c+dx)} - \frac{\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3d\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(7/2)/Sqrt[b*Cos[c + d*x]], x]

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) - (\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\cos^7(c+dx)}{\sqrt{b} \cos(c+dx)} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^3(c+dx) dx}{\sqrt{b} \cos(c+dx)} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1-x^2) dx, x, -\sin(c+dx)\right)}{d\sqrt{b} \cos(c+dx)} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{b} \cos(c+dx)} - \frac{\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3d\sqrt{b} \cos(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.07, size = 45, normalized size = 0.64

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}(\cos(2(c+dx))+5)}{6d\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(7/2)/Sqrt[b*Cos[c + d*x]], x]

[Out] (Sqrt[Cos[c + d*x]]*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.79, size = 42, normalized size = 0.60

$$\frac{\sqrt{b \cos(dx + c)} (\cos(dx + c)^2 + 2) \sin(dx + c)}{3bd\sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(cos(d*x + c)^2 + 2)*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{7}{2}}}{\sqrt{b \cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(7/2)/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.10, size = 40, normalized size = 0.57

$$\frac{(2 + \cos^2(dx + c)) \sin(dx + c) (\sqrt{\cos(dx + c)})}{3d\sqrt{b \cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x)

[Out] 1/3/d*(2+cos(d*x+c)^2)*sin(d*x+c)*cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2)

maxima [A] time = 0.96, size = 42, normalized size = 0.60

$$\frac{\sin(3dx + 3c) + 9 \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)}{12\sqrt{bd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(sqrt(b)*d)

mupad [B] time = 0.64, size = 60, normalized size = 0.86

$$\frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12bd(\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(7/2)/(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*b*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(1/2), x)

[Out] Timed out

$$3.174 \quad \int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=63

$$\frac{x\sqrt{\cos(c+dx)}}{2\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}+1/2*x*\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{x\sqrt{\cos(c+dx)}}{2\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(5/2)/Sqrt[b*Cos[c + d*x]], x]

[Out] $(x*\text{Sqrt}[\text{Cos}[c + d*x]])/(2*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^2(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{\cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{2d\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \int 1 dx}{2\sqrt{b \cos(c+dx)}} \\ &= \frac{x\sqrt{\cos(c+dx)}}{2\sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{2d\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 45, normalized size = 0.71

$$\frac{(2(c+dx) + \sin(2(c+dx)))\sqrt{\cos(c+dx)}}{4d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(5/2)/Sqrt[b*Cos[c + d*x]], x]

[Out] (Sqrt[Cos[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.63, size = 157, normalized size = 2.49

$$\frac{2\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}\sin(dx+c) - \sqrt{-b}\log\left(2b\cos(dx+c)^2 + 2\sqrt{b\cos(dx+c)}\sqrt{-b}\sqrt{\cos(dx+c)}\right)}{4bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b*d), 1/2*(sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2))))/(b*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^{\frac{5}{2}}}{\sqrt{b\cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(5/2)/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.13, size = 42, normalized size = 0.67

$$\frac{(\cos(dx+c)\sin(dx+c) + dx+c)\left(\sqrt{\cos(dx+c)}\right)}{2d\sqrt{b\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2), x)

[Out] 1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)*cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2)

maxima [A] time = 1.26, size = 25, normalized size = 0.40

$$\frac{2dx + 2c + \sin(2dx + 2c)}{4\sqrt{b}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2), x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(sqrt(b)*d)

mupad [B] time = 0.65, size = 65, normalized size = 1.03

$$\frac{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}(\sin(c+dx) + \sin(3c+3dx) + 4dx\cos(c+dx))}{4bd(\cos(2c+2dx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^(5/2)/(b*cos(c + d*x))^(1/2), x)
```

```
[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(4*b*d*(cos(2*c + 2*d*x) + 1))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(1/2), x)
```

```
[Out] Timed out
```

$$3.175 \quad \int \frac{\cos^3(c+dx)}{\sqrt{b} \cos(c+dx)} dx$$

Optimal. Leaf size=32

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b}\cos(c+dx)}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^{(3/2)}/\text{Sqrt}[b*\text{Cos}[c + d*x]], x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ $\text{FreeQ}\{a, b, m\}, x] \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n + 1/2, 0] \&\& \text{IntegerQ}[m + n]$

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ $\text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{\cos^3(c+dx)}{\sqrt{b} \cos(c+dx)} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos(c+dx) dx}{\sqrt{b} \cos(c+dx)} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{b} \cos(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Cos}[c + d*x]^{(3/2)}/\text{Sqrt}[b*\text{Cos}[c + d*x]], x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

fricas [A] time = 0.65, size = 31, normalized size = 0.97

$$\frac{\sqrt{b} \cos(dx+c) \sin(dx+c)}{bd\sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^{\frac{3}{2}}}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(3/2)/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.11, size = 29, normalized size = 0.91

$$\frac{\sin(dx+c) \left(\sqrt{\cos(dx+c)} \right)}{d \sqrt{b \cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x)

[Out] sin(d*x+c)*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

maxima [A] time = 1.26, size = 13, normalized size = 0.41

$$\frac{\sin(dx+c)}{\sqrt{b}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] sin(d*x + c)/(sqrt(b)*d)

mupad [B] time = 0.36, size = 47, normalized size = 1.47

$$\frac{\sqrt{\cos(c+dx)} \sin(2c+2dx) \sqrt{b \cos(c+dx)}}{bd (\cos(2c+2dx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(3/2)/(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*sin(2*c + 2*d*x)*(b*cos(c + d*x))^(1/2))/(b*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

$$3.176 \quad \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=24

$$\frac{x\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}}$$

[Out] $x*\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{x\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Cos[c + d*x]]/Sqrt[b*Cos[c + d*x]],x]

[Out] (x*Sqrt[Cos[c + d*x]])/Sqrt[b*Cos[c + d*x]]

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx &= \frac{\sqrt{\cos(c+dx)} \int 1 dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{x\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 1.00

$$\frac{x\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Cos[c + d*x]]/Sqrt[b*Cos[c + d*x]],x]

[Out] (x*Sqrt[Cos[c + d*x]])/Sqrt[b*Cos[c + d*x]]

fricas [A] time = 0.74, size = 97, normalized size = 4.04

$$\left[\frac{\sqrt{-b} \log(2b \cos(dx+c)^2 + 2\sqrt{b \cos(dx+c)} \sqrt{-b} \sqrt{\cos(dx+c)} \sin(dx+c) - b)}{2bd}, \frac{\arctan\left(\frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{\sqrt{b \cos(dx+c)}}\right)}{\sqrt{b}d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] [-1/2*sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b)/(b*d), arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/(sqrt(b)*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\cos(dx+c)}}{\sqrt{b \cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(cos(d*x + c))/sqrt(b*cos(d*x + c)), x)

maple [A] time = 0.08, size = 28, normalized size = 1.17

$$\frac{(\sqrt{\cos(dx+c)})(dx+c)}{d\sqrt{b \cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x)

[Out] 1/d*cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2)*(d*x+c)

maxima [A] time = 0.94, size = 26, normalized size = 1.08

$$\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{\sqrt{b} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(sqrt(b)*d)

mupad [B] time = 0.26, size = 37, normalized size = 1.54

$$\frac{2x \cos(c+dx)^{3/2} \sqrt{b \cos(c+dx)}}{b (\cos(2c+2dx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(1/2)/(b*cos(c + d*x))^(1/2),x)

[Out] (2*x*cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2))/(b*(cos(2*c + 2*d*x) + 1))

sympy [A] time = 2.43, size = 5, normalized size = 0.21

$$\frac{x}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(1/2),x)

[Out] x/sqrt(b)

$$3.177 \quad \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=33

$$\frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{b \cos(c+dx)}}$$

[Out] arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.01, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 3770}

$$\frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]),x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(d*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\cos(c+dx)}}{d\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 33, normalized size = 1.00

$$\frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]),x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.63, size = 116, normalized size = 3.52

$$\left[\frac{\log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right)}{2\sqrt{b}d}, -\frac{\sqrt{-b} \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b} \sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)}{bd} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3)/(sqrt(b)*d), -sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/(b*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(dx + c)} \sqrt{\cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))), x)

maple [A] time = 0.10, size = 42, normalized size = 1.27

$$\frac{2 \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right) \left(\sqrt{\cos(dx+c)}\right)}{d\sqrt{b \cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x)

[Out] -2/d*arctanh((-1+cos(d*x+c))/sin(d*x+c))*cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2)

maxima [B] time = 1.57, size = 65, normalized size = 1.97

$$\frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2 \sin(dx+c) + 1)}{2\sqrt{b}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(sqrt(b)*d)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(c + dx)} \sqrt{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(1/(sqrt(b*cos(c + d*x))*sqrt(cos(c + d*x))), x)

$$3.178 \quad \int \frac{1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=32

$$\frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}$$

[Out] $\sin(d*x+c)/d/\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3767, 8}

$$\frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(\text{Cos}[c+d*x]^{(3/2)}*\text{Sqrt}[b*\text{Cos}[c+d*x]]),x]$

[Out] $\text{Sin}[c+d*x]/(d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Cos}[c+d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_.))^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m-1/2)}*b^{(n+1/2)}*\text{Sqrt}[a*v])/ \text{Sqrt}[b*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& \text{IntegerQ}[m] \&\& \text{ILtQ}[n-1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.)+(d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1+x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c+d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \&\& \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^2(c+dx) dx}{\sqrt{b\cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{d\sqrt{b\cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[1/(\text{Cos}[c+d*x]^{(3/2)}*\text{Sqrt}[b*\text{Cos}[c+d*x]]),x]$

[Out] $\text{Sin}[c + d*x]/(d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

fricas [A] time = 0.66, size = 31, normalized size = 0.97

$$\frac{\sqrt{b \cos(dx + c)} \sin(dx + c)}{bd \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out] `sqrt(b*cos(d*x + c))*sin(d*x + c)/(b*d*cos(d*x + c)^(3/2))`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(dx + c)} \cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")`

[Out] `integrate(1/(sqrt(b*cos(d*x + c))*cos(d*x + c)^(3/2)), x)`

maple [A] time = 0.12, size = 29, normalized size = 0.91

$$\frac{\sin(dx + c)}{d\sqrt{\cos(dx + c)} \sqrt{b \cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x)`

[Out] `sin(d*x+c)/d/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2)`

maxima [B] time = 1.59, size = 59, normalized size = 1.84

$$\frac{2\sqrt{b} \sin(2dx + 2c)}{(b \cos(2dx + 2c)^2 + b \sin(2dx + 2c)^2 + 2b \cos(2dx + 2c) + b)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `2*sqrt(b)*sin(2*d*x + 2*c)/((b*cos(2*d*x + 2*c)^2 + b*sin(2*d*x + 2*c)^2 + 2*b*cos(2*d*x + 2*c) + b)*d)`

mupad [B] time = 0.54, size = 62, normalized size = 1.94

$$\frac{\sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 1i + \sin(2c + 2dx) + 1i)}{bd \sqrt{\cos(c + dx)} (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2)),x)`

[Out] `((b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/(b*d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Integral(1/(sqrt(b*cos(c + d*x))*cos(c + d*x)**(3/2)), x)
```

$$3.179 \quad \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

Optimal. Leaf size=72

$$\frac{\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)}\tanh^{-1}(\sin(c+dx))}{2d\sqrt{b\cos(c+dx)}}$$

[Out] 1/2*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2)+1/2*arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.02, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, number of rules / integrand size = 0.130, Rules used = {18, 3768, 3770}

$$\frac{\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)}\tanh^{-1}(\sin(c+dx))}{2d\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]]),x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]]/(2*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(2*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{b\cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{2\sqrt{b\cos(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\cos(c+dx)}}{2d\sqrt{b\cos(c+dx)}} + \frac{\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 52, normalized size = 0.72

$$\frac{\sin(c + dx) + \cos^2(c + dx) \tanh^{-1}(\sin(c + dx))}{2d \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]]),x]

[Out] (ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x])/(2*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 1.00, size = 207, normalized size = 2.88

$$\frac{\sqrt{b} \cos(dx + c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b \cos(dx+c)} \sqrt{\cos(dx+c)}}{4bd \cos(dx+c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c)^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^3), -1/2*(sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^3)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(dx + c)} \cos(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*cos(d*x + c)^(5/2)), x)

maple [A] time = 0.14, size = 104, normalized size = 1.44

$$\frac{(\cos^2(dx + c)) \ln\left(-\frac{-1 + \cos(dx+c) + \sin(dx+c)}{\sin(dx+c)}\right) - (\cos^2(dx + c)) \ln\left(-\frac{-\sin(dx+c) - 1 + \cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx + c)}{2d\sqrt{b \cos(dx + c)} \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c))/(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(3/2)

maxima [B] time = 1.25, size = 661, normalized size = 9.18

$$\frac{4(\sin(4dx + 4c) + 2\sin(2dx + 2c)) \cos\left(\frac{3}{2} \arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) - 4(\sin(4dx + 4c))}{2d\sqrt{b \cos(dx + c)} \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out]
$$-1/4*(4*(\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\cos(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 4*(\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - (2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) + (2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 - 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) - 4*(\cos(4*d*x + 4*c) + 2*\cos(2*d*x + 2*c) + 1)*\sin(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 4*(\cos(4*d*x + 4*c) + 2*\cos(2*d*x + 2*c) + 1)*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))/((2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*\sqrt{b}*d)$$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^{5/2} \sqrt{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(1/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

$$3.180 \quad \int \frac{1}{\cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=70

$$\frac{\sin^3(c+dx)}{3d \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

[Out] $1/3 * \sin(d*x+c)^3 / d / \cos(d*x+c)^{(5/2)} / (b * \cos(d*x+c))^{(1/2)} + \sin(d*x+c) / d / \cos(d*x+c)^{(1/2)} / (b * \cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 3767}

$$\frac{\sin^3(c+dx)}{3d \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]]),x]

[Out] Sin[c + d*x]/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]^3/(3*d*Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^4(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d \sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} + \frac{\sin^3(c+dx)}{3d \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 45, normalized size = 0.64

$$\frac{\sqrt{\cos(c+dx)} \left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx) \right)}{d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]]),x]

[Out] (Sqrt[Cos[c + d*x]]*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.70, size = 44, normalized size = 0.63

$$\frac{\sqrt{b \cos(dx + c)} (2 \cos(dx + c)^2 + 1) \sin(dx + c)}{3bd \cos(dx + c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 1)*sin(d*x + c)/(b*d*cos(d*x + c)^(7/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \cos(dx + c)} \cos(dx + c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*cos(d*x + c)^(7/2)), x)

maple [A] time = 0.10, size = 42, normalized size = 0.60

$$\frac{\sin(dx + c) (2 (\cos^2(dx + c)) + 1)}{3d \sqrt{b \cos(dx + c)} \cos(dx + c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x)

[Out] 1/3/d*sin(d*x+c)*(2*cos(d*x+c)^2+1)/(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2)

maxima [B] time = 1.60, size = 294, normalized size = 4.20

$$\frac{4((3 \cos(2dx + 2c) + 1) \sin(6dx + 6c) + 3(3 \cos(2dx + 2c) + 1) \sin(4dx + 4c) - 3 \cos(6dx + 6c) \sin(2dx + 2c) - 9 \cos(4dx + 4c) \sin(2dx + 2c))}{3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \cos(4dx + 4c) + 9 \cos(4dx + 4c)^2 + 9 \cos(2dx + 2c)^2 + 6(\sin(4dx + 4c) + \sin(2dx + 2c)) \sin(6dx + 6c) + \sin(6dx + 6c)^2 + 9 \sin(4dx + 4c)^2 + 18 \sin(4dx + 4c) \sin(2dx + 2c) + 9 \sin(2dx + 2c)^2 + 6 \cos(2dx + 2c) + 1) \sqrt{b} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 4/3*((3*cos(2*d*x + 2*c) + 1)*sin(6*d*x + 6*c) + 3*(3*cos(2*d*x + 2*c) + 1)*sin(4*d*x + 4*c) - 3*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) - 9*cos(4*d*x + 4*c)*sin(2*d*x + 2*c))/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*sqrt(b)*d)

mupad [B] time = 1.37, size = 131, normalized size = 1.87

$$\frac{2 \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 15i + \cos(4c + 4dx) 6i + \cos(6c + 6dx) 1i + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}{3bd \sqrt{\cos(c + dx)} (15 \cos(2c + 2dx) + 6 \cos(4c + 4dx) + \cos(6c + 6dx) + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cos(c + d*x)^(7/2)*(b*cos(c + d*x))^(1/2)),x)
```

```
[Out] (2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*b*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

$$3.181 \quad \int \frac{1}{\cos^2(c+dx) \sqrt{b \cos(c+dx)}} dx$$

Optimal. Leaf size=107

$$\frac{3 \sin(c+dx)}{8d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8d \sqrt{b \cos(c+dx)}}$$

[Out] 1/4*sin(d*x+c)/d/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2)+3/8*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2)+3/8*arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.03, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3768, 3770}

$$\frac{3 \sin(c+dx)}{8d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Cos[c + d*x]^(9/2)*Sqrt[b*Cos[c + d*x]]), x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(8*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(4*d*Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]]) + (3*Sin[c + d*x])/(8*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \frac{1}{\cos^{\frac{9}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)} \int \sec^5(c+dx) dx}{\sqrt{b\cos(c+dx)}}$$

$$= \frac{\sin(c+dx)}{4d \cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{(3\sqrt{\cos(c+dx)}) \int \sec^3(c+dx) dx}{4\sqrt{b\cos(c+dx)}}$$

$$= \frac{\sin(c+dx)}{4d \cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{3 \sin(c+dx)}{8d \cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{(3\sqrt{\cos(c+dx)}) \int \sec dx}{4\sqrt{b\cos(c+dx)}}$$

$$= \frac{3 \tanh^{-1}(\sin(c+dx))\sqrt{\cos(c+dx)}}{8d\sqrt{b\cos(c+dx)}} + \frac{\sin(c+dx)}{4d \cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \dots$$

Mathematica [A] time = 0.06, size = 66, normalized size = 0.62

$$\frac{\sin(c+dx)(3\cos^2(c+dx)+2)+3\cos^4(c+dx)\tanh^{-1}(\sin(c+dx))}{8d\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Cos[c + d*x]^(9/2)*Sqrt[b*Cos[c + d*x]]), x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x])/(8*d*Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.75, size = 233, normalized size = 2.18

$$\left[\frac{3\sqrt{b}\cos(dx+c)^5 \log\left(-\frac{b\cos(dx+c)^3 - 2\sqrt{b\cos(dx+c)}\sqrt{b}\sqrt{\cos(dx+c)}\sin(dx+c) - 2b\cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b\cos(dx+c)}(3\cos(dx+c)^2 + 2)}{16bd\cos(dx+c)^5} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c)/(b*d*cos(d*x + c)^5), -1/8*(3*sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^5 - sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^5)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b\cos(dx+c)} \cos(dx+c)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*cos(d*x + c)^(9/2)), x)

maple [A] time = 0.15, size = 121, normalized size = 1.13

$$\frac{3(\cos^4(dx+c)) \ln\left(-\frac{\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - 3(\cos^4(dx+c)) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) + 3(\cos^2(dx+c)) \sqrt{\cos(dx+c)}}{8d\sqrt{b\cos(dx+c)} \cos(dx+c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2),x)
```

```
[Out] 1/8/d*(3*cos(d*x+c)^4*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^4*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))/(b*cos(d*x+c))^(1/2)/cos(d*x+c)^(7/2)
```

maxima [B] time = 1.30, size = 1656, normalized size = 15.48

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] -1/16*(12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))))/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*sqrt(b)*d)
```


mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^{9/2} \sqrt{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^(9/2)*(b*cos(c + d*x))^(1/2)), x)

[Out] int(1/(cos(c + d*x)^(9/2)*(b*cos(c + d*x))^(1/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)**(9/2)/(b*cos(d*x+c))**(1/2), x)

[Out] Timed out

$$3.182 \quad \int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=107

$$\frac{3x\sqrt{\cos(c+dx)}}{8b\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{4bd\sqrt{b}\cos(c+dx)} + \frac{3\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{8bd\sqrt{b}\cos(c+dx)}$$

[Out] $3/8*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}+1/4*\cos(d*x+c)^{(7/2)}*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}+3/8*x*\cos(d*x+c)^{(1/2)}/b/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{3x\sqrt{\cos(c+dx)}}{8b\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{4bd\sqrt{b}\cos(c+dx)} + \frac{3\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{8bd\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(11/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] $(3*x*\text{Sqrt}[\text{Cos}[c + d*x]])/(8*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (3*\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(8*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(7/2)}*\text{Sin}[c + d*x])/(4*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^4(c+dx) dx}{b\sqrt{b}\cos(c+dx)} \\ &= \frac{\cos^{\frac{7}{2}}(c+dx)\sin(c+dx)}{4bd\sqrt{b}\cos(c+dx)} + \frac{(3\sqrt{\cos(c+dx)}) \int \cos^2(c+dx) dx}{4b\sqrt{b}\cos(c+dx)} \\ &= \frac{3\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{8bd\sqrt{b}\cos(c+dx)} + \frac{\cos^{\frac{7}{2}}(c+dx)\sin(c+dx)}{4bd\sqrt{b}\cos(c+dx)} + \frac{(3\sqrt{\cos(c+dx)}) \int 1 dx}{8b\sqrt{b}\cos(c+dx)} \\ &= \frac{3x\sqrt{\cos(c+dx)}}{8b\sqrt{b}\cos(c+dx)} + \frac{3\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{8bd\sqrt{b}\cos(c+dx)} + \frac{\cos^{\frac{7}{2}}(c+dx)\sin(c+dx)}{4bd\sqrt{b}\cos(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.05, size = 55, normalized size = 0.51

$$\frac{(12(c + dx) + 8 \sin(2(c + dx)) + \sin(4(c + dx))) \cos^{\frac{3}{2}}(c + dx)}{32d(b \cos(c + dx))^{\frac{3}{2}}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(11/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] (Cos[c + d*x]^(3/2)*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])) / (32*d*(b*Cos[c + d*x])^(3/2))

fricas [A] time = 0.93, size = 182, normalized size = 1.70

$$\left[\frac{2 \sqrt{b \cos(dx + c)} (2 \cos(dx + c)^2 + 3) \sqrt{\cos(dx + c)} \sin(dx + c) - 3 \sqrt{-b} \log(2 b \cos(dx + c)^2 + 2 \sqrt{b \cos(dx + c)})}{16 b^2 d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] [1/16*(2*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 3)*sqrt(cos(d*x + c))*sin(d*x + c) - 3*sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b^2*d), 1/8*(sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 3)*sqrt(cos(d*x + c))*sin(d*x + c) + 3*sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/(b^2*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{11}{2}}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(11/2)/(b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.14, size = 62, normalized size = 0.58

$$\frac{\left(\cos^{\frac{3}{2}}(dx + c)\right) \left(2 \left(\cos^3(dx + c)\right) \sin(dx + c) + 3 \cos(dx + c) \sin(dx + c) + 3dx + 3c\right)}{8d(b \cos(dx + c))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(3/2), x)

[Out] 1/8/d*cos(d*x+c)^(3/2)*(2*cos(d*x+c)^3*sin(d*x+c)+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)/(b*cos(d*x+c))^(3/2)

maxima [A] time = 1.25, size = 49, normalized size = 0.46

$$\frac{12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin\left(\frac{1}{2} \arctan(\sin(4 dx + 4 c), \cos(4 dx + 4 c))\right)}{32 b^{\frac{3}{2}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(b^(3/2)*d)

mupad [B] time = 0.98, size = 78, normalized size = 0.73

$$\frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx) + 24dx \cos(c + dx))}{32 b^2 d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(11/2)/(b*cos(c + d*x))^(3/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*b^2*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(11/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

$$3.183 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=76

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{bd\sqrt{b \cos(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3bd\sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{bd\sqrt{b \cos(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(9/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[b*Cos[c + d*x]]) - (Sqrt[Cos[c + d*x]]*Sin[c + d*x]^3)/(3*b*d*Sqrt[b*Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^3(c+dx) dx}{b\sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1-x^2) dx, x, -\sin(c+dx)\right)}{bd\sqrt{b \cos(c+dx)}} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}} - \frac{\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3bd\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 45, normalized size = 0.59

$$\frac{\sin(c+dx) \cos^2(c+dx) (\cos(2(c+dx)) + 5)}{6d(b \cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(9/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] $(\cos[c + d*x]^{(3/2)}*(5 + \cos[2*(c + d*x)])*\sin[c + d*x])/(6*d*(b*\cos[c + d*x])^{(3/2)})$

fricas [A] time = 0.85, size = 42, normalized size = 0.55

$$\frac{\sqrt{b \cos(dx + c)} (\cos(dx + c)^2 + 2) \sin(dx + c)}{3 b^2 d \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out] $1/3*\sqrt{b*\cos(d*x + c)}*(\cos(d*x + c)^2 + 2)*\sin(d*x + c)/(b^2*d*\sqrt{\cos(d*x + c)})$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{9}{2}}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")`

[Out] `integrate(cos(d*x + c)^(9/2)/(b*cos(d*x + c))^(3/2), x)`

maple [A] time = 0.09, size = 40, normalized size = 0.53

$$\frac{(2 + \cos^2(dx + c)) \left(\cos^{\frac{3}{2}}(dx + c) \right) \sin(dx + c)}{3d (b \cos(dx + c))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(3/2),x)`

[Out] $1/3/d*(2+\cos(d*x+c)^2)*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/(b*\cos(d*x+c))^{(3/2)}$

maxima [A] time = 1.00, size = 42, normalized size = 0.55

$$\frac{\sin(3dx + 3c) + 9 \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)}{12 b^{\frac{3}{2}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] $1/12*(\sin(3*d*x + 3*c) + 9*\sin(1/3*\arctan2(\sin(3*d*x + 3*c), \cos(3*d*x + 3*c))))/(b^{(3/2)}*d)$

mupad [B] time = 0.70, size = 60, normalized size = 0.79

$$\frac{\sqrt{\cos(c + d*x)} \sqrt{b \cos(c + d*x)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12 b^2 d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^(9/2)/(b*cos(c + d*x))^(3/2),x)`

[Out] $(\cos(c + d*x)^{(1/2)}*(b*\cos(c + d*x))^{(1/2)}*(10*\sin(2*c + 2*d*x) + \sin(4*c + 4*d*x)))/(12*b^2*d*(\cos(2*c + 2*d*x) + 1))$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(9/2)/(b*cos(d*x+c))**(3/2), x)

[Out] Timed out

$$3.184 \quad \int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=69

$$\frac{x\sqrt{\cos(c+dx)}}{2b\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^3(c+dx)}{2bd\sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}+1/2*x*\cos(d*x+c)^{(1/2)}/b/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{x\sqrt{\cos(c+dx)}}{2b\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^3(c+dx)}{2bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(7/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] $(x*\text{Sqrt}[\text{Cos}[c + d*x]])/(2*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(2*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] :> -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^2(c+dx) dx}{b\sqrt{b \cos(c+dx)}} \\ &= \frac{\cos^3(c+dx) \sin(c+dx)}{2bd\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \int 1 dx}{2b\sqrt{b \cos(c+dx)}} \\ &= \frac{x\sqrt{\cos(c+dx)}}{2b\sqrt{b \cos(c+dx)}} + \frac{\cos^3(c+dx) \sin(c+dx)}{2bd\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 45, normalized size = 0.65

$$\frac{(2(c+dx) + \sin(2(c+dx))) \cos^3(c+dx)}{4d(b \cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(7/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] (Cos[c + d*x]^(3/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*(b*Cos[c + d*x])^(3/2))

fricas [A] time = 1.15, size = 157, normalized size = 2.28

$$\left[\frac{2 \sqrt{b \cos(dx + c)} \sqrt{\cos(dx + c)} \sin(dx + c) - \sqrt{-b} \log(2b \cos(dx + c)^2 + 2 \sqrt{b \cos(dx + c)} \sqrt{-b} \sqrt{\cos(dx + c)})}{4b^2d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b^2*d), 1/2*(sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c))^(3/2)))/(b^2*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{7}{2}}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(7/2)/(b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.11, size = 42, normalized size = 0.61

$$\frac{\left(\cos^{\frac{3}{2}}(dx + c)\right) (\cos(dx + c) \sin(dx + c) + dx + c)}{2d (b \cos(dx + c))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2), x)

[Out] 1/2/d*cos(d*x+c)^(3/2)*(cos(d*x+c)*sin(d*x+c)+d*x+c)/(b*cos(d*x+c))^(3/2)

maxima [A] time = 1.50, size = 25, normalized size = 0.36

$$\frac{2dx + 2c + \sin(2dx + 2c)}{4b^{\frac{3}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2), x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(b^(3/2)*d)

mupad [B] time = 0.59, size = 65, normalized size = 0.94

$$\frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (\sin(c + dx) + \sin(3c + 3dx) + 4dx \cos(c + dx))}{4b^2d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^(7/2)/(b*cos(c + d*x))^(3/2), x)
```

```
[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x)
+ 4*d*x*cos(c + d*x)))/(4*b^2*d*(cos(2*c + 2*d*x) + 1))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(3/2), x)
```

```
[Out] Timed out
```

$$3.185 \quad \int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=35

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{bd\sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^{(5/2)}/(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos(c+dx) dx}{b\sqrt{b \cos(c+dx)}} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 32, normalized size = 0.91

$$\frac{\sin(c+dx) \cos^3(c+dx)}{d(b \cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Cos}[c + d*x]^{(5/2)}/(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(d*(b*\text{Cos}[c + d*x])^{(3/2)})$

fricas [A] time = 0.92, size = 31, normalized size = 0.89

$$\frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{b^2 d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b^2*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^{\frac{5}{2}}}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(5/2)/(b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.08, size = 29, normalized size = 0.83

$$\frac{\sin(dx+c) \left(\cos^{\frac{3}{2}}(dx+c) \right)}{d (b \cos(dx+c))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x)

[Out] 1/d*sin(d*x+c)*cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2)

maxima [A] time = 1.48, size = 13, normalized size = 0.37

$$\frac{\sin(dx+c)}{b^{\frac{3}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] sin(d*x + c)/(b^(3/2)*d)

mupad [B] time = 0.32, size = 47, normalized size = 1.34

$$\frac{\sqrt{\cos(c+dx)} \sin(2c+2dx) \sqrt{b \cos(c+dx)}}{b^2 d (\cos(2c+2dx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c+d*x)^(5/2)/(b*cos(c+d*x))^(3/2),x)

[Out] (cos(c+d*x)^(1/2)*sin(2*c+2*d*x)*(b*cos(c+d*x))^(1/2))/(b^2*d*(cos(2*c+2*d*x)+1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

$$3.186 \quad \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=27

$$\frac{x\sqrt{\cos(c+dx)}}{b\sqrt{b \cos(c+dx)}}$$

[Out] $x \cos(dx+c)^{1/2} / b / (b \cos(dx+c))^{1/2}$

Rubi [A] time = 0.00, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{x\sqrt{\cos(c+dx)}}{b\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(3/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] (x*Sqrt[Cos[c + d*x]])/(b*Sqrt[b*Cos[c + d*x]])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int 1 dx}{b\sqrt{b \cos(c+dx)}} \\ &= \frac{x\sqrt{\cos(c+dx)}}{b\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 0.89

$$\frac{x \cos^3(c+dx)}{(b \cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(3/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] (x*Cos[c + d*x]^(3/2))/(b*Cos[c + d*x])^(3/2)

fricas [A] time = 0.76, size = 97, normalized size = 3.59

$$\left[\frac{\sqrt{-b} \log(2b \cos(dx+c)^2 + 2\sqrt{b \cos(dx+c)} \sqrt{-b} \sqrt{\cos(dx+c)} \sin(dx+c) - b)}{2b^2d}, \frac{\arctan\left(\frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{\sqrt{b \cos(dx+c)}^3}\right)}{b^{\frac{3}{2}}d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [-1/2*sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b)/(b^2*d), arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/(b^(3/2)*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^{\frac{3}{2}}}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(3/2)/(b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.08, size = 28, normalized size = 1.04

$$\frac{\left(\cos^{\frac{3}{2}}(dx+c)\right)(dx+c)}{d(b \cos(dx+c))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x)

[Out] 1/d*cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2)*(d*x+c)

maxima [A] time = 1.05, size = 26, normalized size = 0.96

$$\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{b^{\frac{3}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(b^(3/2)*d)

mupad [B] time = 0.28, size = 37, normalized size = 1.37

$$\frac{2x \cos(c+dx)^{3/2} \sqrt{b \cos(c+dx)}}{b^2 (\cos(2c+2dx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(3/2)/(b*cos(c + d*x))^(3/2),x)

[Out] (2*x*cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2))/(b^2*(cos(2*c + 2*d*x) + 1))

sympy [A] time = 129.11, size = 5, normalized size = 0.19

$$\frac{x}{b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(3/2),x)

[Out] x/b**(3/2)

$$3.187 \quad \int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=36

$$\frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{bd\sqrt{b \cos(c+dx)}}$$

[Out] arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Cos[c + d*x]]/(b*Cos[c + d*x])^(3/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(b*d*Sqrt[b*Cos[c + d*x]])

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{b\sqrt{b \cos(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\cos(c+dx)}}{bd\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 33, normalized size = 0.92

$$\frac{\cos^{\frac{3}{2}}(c+dx) \tanh^{-1}(\sin(c+dx))}{d(b \cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Cos[c + d*x]]/(b*Cos[c + d*x])^(3/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^(3/2))/(d*(b*Cos[c + d*x])^(3/2))

fricas [A] time = 0.85, size = 116, normalized size = 3.22

$$\left[\frac{\log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right)}{2b^{\frac{3}{2}}d}, -\frac{\sqrt{-b} \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b} \sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)}{b^2d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3)/(b^(3/2)*d), -sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/(b^2*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\cos(dx+c)}}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(cos(d*x + c))/(b*cos(d*x + c))^(3/2), x)

maple [A] time = 0.11, size = 42, normalized size = 1.17

$$\frac{2 \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right) \left(\cos^{\frac{3}{2}}(dx+c)\right)}{d (b \cos(dx+c))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x)

[Out] -2/d*arctanh((-1+cos(d*x+c))/sin(d*x+c))*cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2)

maxima [B] time = 1.16, size = 65, normalized size = 1.81

$$\frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2 \sin(dx+c) + 1)}{2 b^{\frac{3}{2}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(b^(3/2)*d)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(1/2)/(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^(1/2)/(b*cos(c + d*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(sqrt(cos(c + d*x))/(b*cos(c + d*x))**(3/2), x)

$$3.188 \quad \int \frac{1}{\sqrt{\cos(c+dx)} (b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=35

$$\frac{\sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}$$

[Out] sin(d*x+c)/b/d/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3767, 8}

$$\frac{\sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(3/2)),x]

[Out] Sin[c + d*x]/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]])

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 18

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{\cos(c+dx)} (b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^2(c+dx) dx}{b\sqrt{b\cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{bd\sqrt{b\cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 32, normalized size = 0.91

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d(b\cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(3/2)),x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(b*Cos[c + d*x])^(3/2))

fricas [A] time = 0.62, size = 31, normalized size = 0.89

$$\frac{\sqrt{b \cos(dx + c)} \sin(dx + c)}{b^2 d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b^2*d*cos(d*x + c)^(3/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{3}{2}} \sqrt{\cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(3/2)*sqrt(cos(d*x + c))), x)

maple [A] time = 0.12, size = 29, normalized size = 0.83

$$\frac{(\sqrt{\cos(dx + c)} \sin(dx + c))}{d (b \cos(dx + c))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x)

[Out] 1/d*cos(d*x+c)^(1/2)*sin(d*x+c)/(b*cos(d*x+c))^(3/2)

maxima [B] time = 1.00, size = 67, normalized size = 1.91

$$\frac{2 \sqrt{b} \sin(2 dx + 2 c)}{(b^2 \cos(2 dx + 2 c)^2 + b^2 \sin(2 dx + 2 c)^2 + 2 b^2 \cos(2 dx + 2 c) + b^2) d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*sin(2*d*x + 2*c)/((b^2*cos(2*d*x + 2*c)^2 + b^2*sin(2*d*x + 2*c)^2 + 2*b^2*cos(2*d*x + 2*c) + b^2)*d)

mupad [B] time = 0.51, size = 62, normalized size = 1.77

$$\frac{\sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 1i + \sin(2c + 2dx) + 1i)}{b^2 d \sqrt{\cos(c + dx)} (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(3/2)),x)

[Out] ((b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/(b^2*d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(c + dx))^{\frac{3}{2}} \sqrt{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(3/2),x)
```

```
[Out] Integral(1/((b*cos(c + d*x))**(3/2)*sqrt(cos(c + d*x))), x)
```

$$3.189 \quad \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=78

$$\frac{\sin(c+dx)}{2bd \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2bd\sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\sin(d*x+c)/b/d/\cos(d*x+c)^{(3/2)}/(b*\cos(d*x+c))^{(1/2)}+1/2*\operatorname{arctanh}(\sin(d*x+c))*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3768, 3770}

$$\frac{\sin(c+dx)}{2bd \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2bd\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] `Int[1/(Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(3/2)), x]`

[Out] `(ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(2*b*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(2*b*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])`

Rule 18

`Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3770

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^3(c+dx) dx}{b\sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{2bd \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{2b\sqrt{b \cos(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\cos(c+dx)}}{2bd\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2bd \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 52, normalized size = 0.67

$$\frac{\sin(c+dx) + \cos^2(c+dx) \tanh^{-1}(\sin(c+dx))}{2d\sqrt{\cos(c+dx)}(b \cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(3/2)),x]

[Out] (ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x])/(2*d*Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(3/2))

fricas [A] time = 1.39, size = 207, normalized size = 2.65

$$\left[\frac{\sqrt{b} \cos(dx + c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b \cos(dx+c)} \sqrt{\cos(dx+c)}}{4b^2 d \cos(dx+c)^3} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b^2*d*cos(d*x + c)^3), -1/2*(sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(b^2*d*cos(d*x + c)^3)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(3/2)*cos(d*x + c)^(3/2)), x)

maple [A] time = 0.10, size = 102, normalized size = 1.31

$$\frac{(\cos^2(dx + c)) \ln\left(-\frac{\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - (\cos^2(dx + c)) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) + \sin(dx + c)}{2d(b \cos(dx + c))^{\frac{3}{2}} \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x)

[Out] 1/2/d*(cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))+sin(d*x+c))/(b*cos(d*x+c))^(3/2)/cos(d*x+c)^(1/2)

maxima [B] time = 1.57, size = 670, normalized size = 8.59

$$4(\sin(4dx + 4c) + 2\sin(2dx + 2c)) \cos\left(\frac{3}{2} \arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) - 4(\sin(4dx + 4c) + 2\sin(2dx + 2c))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] -1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(1

```

/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c) +
1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x
+ 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*
cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c
)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*
arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x + 2*c)
+ 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d
*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 +
4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2
*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/
2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4*d*x + 4*c) +
2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
)) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*
x + 2*c), cos(2*d*x + 2*c))))/((b*cos(4*d*x + 4*c)^2 + 4*b*cos(2*d*x + 2*c)
^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b*sin
(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 4*b*cos(2
*d*x + 2*c) + b)*sqrt(b)*d)

```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^{3/2} (b \cos(c + dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(3/2)),x)

[Out] int(1/(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(3/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

$$3.190 \quad \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=76

$$\frac{\sin^3(c+dx)}{3bd \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{bd \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

[Out] $1/3*\sin(d*x+c)^3/b/d/\cos(d*x+c)^{(5/2)}/(b*\cos(d*x+c))^{(1/2)}+\sin(d*x+c)/b/d/c$
 $os(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 3767}

$$\frac{\sin^3(c+dx)}{3bd \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{bd \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(3/2)), x]

[Out] Sin[c + d*x]/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]^3/(3*b*d*Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^4(c+dx) dx}{b \sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{bd \sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{bd \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} + \frac{\sin^3(c+dx)}{3bd \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 45, normalized size = 0.59

$$\frac{\cos^{\frac{3}{2}}(c+dx) \left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx) \right)}{d(b \cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(3/2)),x]

[Out] (Cos[c + d*x]^(3/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*(b*Cos[c + d*x])^(3/2))

fricas [A] time = 0.63, size = 44, normalized size = 0.58

$$\frac{\sqrt{b \cos(dx + c)} (2 \cos(dx + c)^2 + 1) \sin(dx + c)}{3 b^2 d \cos(dx + c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 1)*sin(d*x + c)/(b^2*d*cos(d*x + c)^(7/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(3/2)*cos(d*x + c)^(5/2)), x)

maple [A] time = 0.11, size = 42, normalized size = 0.55

$$\frac{\sin(dx + c) (2 (\cos^2(dx + c)) + 1)}{3d (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x)

[Out] 1/3/d*sin(d*x+c)*(2*cos(d*x+c)^2+1)/(b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2)

maxima [B] time = 1.77, size = 311, normalized size = 4.09

$$\frac{4((3 \cos(6 dx + 6 c))^2 + 9 b \cos(4 dx + 4 c)^2 + 9 b \cos(2 dx + 2 c)^2 + b \sin(6 dx + 6 c)^2 + 9 b \sin(4 dx + 4 c)^2 + 18 b \sin(2 dx + 2 c)^2)}{3 b^2 d \sqrt{\cos(c + d x)} (15 \cos(2 c + 2 d x) + 6 \cos(4 c + 4 d x) + \cos(6 c + 6 d x) + 9 \sin(2 c + 2 d x) + 6 \sin(4 c + 4 d x) + \sin(6 c + 6 d x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 4/3*((3*cos(2*d*x + 2*c) + 1)*sin(6*d*x + 6*c) + 3*(3*cos(2*d*x + 2*c) + 1)*sin(4*d*x + 4*c) - 3*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) - 9*cos(4*d*x + 4*c)*sin(2*d*x + 2*c))/((b*cos(6*d*x + 6*c)^2 + 9*b*cos(4*d*x + 4*c)^2 + 9*b*cos(2*d*x + 2*c)^2 + b*sin(6*d*x + 6*c)^2 + 9*b*sin(4*d*x + 4*c)^2 + 18*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*b*sin(2*d*x + 2*c)^2 + 2*(3*b*cos(4*d*x + 4*c) + 3*b*cos(2*d*x + 2*c) + b)*cos(6*d*x + 6*c) + 6*(3*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 6*b*cos(2*d*x + 2*c) + 6*(b*sin(4*d*x + 4*c) + b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b)*sqrt(b)*d)

mupad [B] time = 1.26, size = 131, normalized size = 1.72

$$\frac{2 \sqrt{b \cos(c + d x)} (\cos(2 c + 2 d x) 15 i + \cos(4 c + 4 d x) 6 i + \cos(6 c + 6 d x) 1 i + 9 \sin(2 c + 2 d x) + 6 \sin(4 c + 4 d x) + \sin(6 c + 6 d x))}{3 b^2 d \sqrt{\cos(c + d x)} (15 \cos(2 c + 2 d x) + 6 \cos(4 c + 4 d x) + \cos(6 c + 6 d x) + 9 \sin(2 c + 2 d x) + 6 \sin(4 c + 4 d x) + \sin(6 c + 6 d x))}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(3/2)),x)
```

```
[Out] (2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*b^2*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(3/2),x)
```

```
[Out] Timed out
```

$$3.191 \quad \int \frac{1}{\cos^{\frac{7}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx$$

Optimal. Leaf size=116

$$\frac{3 \sin(c+dx)}{8bd \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4bd \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8bd \sqrt{b \cos(c+dx)}}$$

[Out] 1/4*sin(d*x+c)/b/d/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2)+3/8*sin(d*x+c)/b/d/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2)+3/8*arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.04, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3768, 3770}

$$\frac{3 \sin(c+dx)}{8bd \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4bd \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8bd \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Cos[c + d*x]^(7/2)*(b*Cos[c + d*x])^(3/2)), x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]]/(8*b*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(4*b*d*Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]]) + (3*Sin[c + d*x])/(8*b*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \frac{\sqrt{\cos(c+dx)} \int \sec^5(c+dx) dx}{b\sqrt{b\cos(c+dx)}}$$

$$= \frac{\sin(c+dx)}{4bd \cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{(3\sqrt{\cos(c+dx)}) \int \sec^3(c+dx) dx}{4b\sqrt{b\cos(c+dx)}}$$

$$= \frac{\sin(c+dx)}{4bd \cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{3 \sin(c+dx)}{8bd \cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \dots$$

$$= \frac{3 \tanh^{-1}(\sin(c+dx))\sqrt{\cos(c+dx)}}{8bd\sqrt{b\cos(c+dx)}} + \frac{\sin(c+dx)}{4bd \cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}}$$

Mathematica [A] time = 0.04, size = 66, normalized size = 0.57

$$\frac{\sin(c+dx)(3\cos^2(c+dx)+2)+3\cos^4(c+dx)\tanh^{-1}(\sin(c+dx))}{8d\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Cos[c + d*x]^(7/2)*(b*Cos[c + d*x])^(3/2)),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x])/(8*d*Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(3/2))

fricas [A] time = 0.78, size = 233, normalized size = 2.01

$$\left[\frac{3\sqrt{b}\cos(dx+c)^5 \log\left(-\frac{b\cos(dx+c)^3 - 2\sqrt{b\cos(dx+c)}\sqrt{b\cos(dx+c)}\sin(dx+c) - 2b\cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b\cos(dx+c)}(3\cos(dx+c)^2 + 2)}{16b^2d\cos(dx+c)^5} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3 + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(b^2*d*cos(d*x + c)^5), -1/8*(3*sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^5 - sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(b^2*d*cos(d*x + c)^5)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b\cos(dx+c))^{\frac{3}{2}}\cos(dx+c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(3/2)*cos(d*x + c)^(7/2)), x)

maple [A] time = 0.17, size = 121, normalized size = 1.04

$$\frac{3(\cos^4(dx+c)\ln\left(-\frac{\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - 3(\cos^4(dx+c)\ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) + 3(\cos^2(dx+c)\ln\left(-\frac{1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right)))}{8d(b\cos(dx+c))^{\frac{3}{2}}\cos(dx+c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x)
```

```
[Out] 1/8/d*(3*cos(d*x+c)^4*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^4*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))/(b*cos(d*x+c))^(3/2)/cos(d*x+c)^(5/2)
```

maxima [B] time = 2.00, size = 1679, normalized size = 14.47

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")
```

```
[Out] -1/16*(12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))/((b*cos(8*d*x + 8*c)^2 + 16*b*cos(6*d*x + 6*c)^2 + 36*b*cos(4*d*x + 4*c)^2 + 16*b*cos(2*d*x + 2*c)^2 + b*sin(8*d*x + 8*c)^2 + 16*b*sin(6*d*x + 6*c)^2 + 36*b*sin(4*d*x + 4*c)^2 + 48*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b*sin(2*d*x + 2*c)^2 + 2*(4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*cos(8*d*x + 8*c) + 8*(6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*cos(6*d*x + 6*c) + 12*(4*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 8*b*cos(2*d*x + 2*c) + 4*(2*b*sin(6*d*x + 6*c) + 3*b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b)*sqrt(b)*d)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^{7/2} (b \cos(c + dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^(7/2)*(b*cos(c + d*x))^(3/2)), x)

[Out] int(1/(cos(c + d*x)^(7/2)*(b*cos(c + d*x))^(3/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(3/2), x)

[Out] Timed out

$$3.192 \quad \int \frac{\cos^{13/2}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=107

$$\frac{3x\sqrt{\cos(c+dx)}}{8b^2\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)\cos^{7/2}(c+dx)}{4b^2d\sqrt{b}\cos(c+dx)} + \frac{3\sin(c+dx)\cos^{3/2}(c+dx)}{8b^2d\sqrt{b}\cos(c+dx)}$$

[Out] $3/8*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}+1/4*\cos(d*x+c)^{(7/2)}*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}+3/8*x*\cos(d*x+c)^{(1/2)}/b^2/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{3x\sqrt{\cos(c+dx)}}{8b^2\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)\cos^{7/2}(c+dx)}{4b^2d\sqrt{b}\cos(c+dx)} + \frac{3\sin(c+dx)\cos^{3/2}(c+dx)}{8b^2d\sqrt{b}\cos(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(13/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] $(3*x*\text{Sqrt}[\text{Cos}[c + d*x]])/(8*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (3*\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(8*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(7/2)}*\text{Sin}[c + d*x])/(4*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^{13/2}(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^4(c+dx) dx}{b^2\sqrt{b}\cos(c+dx)} \\ &= \frac{\cos^{7/2}(c+dx)\sin(c+dx)}{4b^2d\sqrt{b}\cos(c+dx)} + \frac{(3\sqrt{\cos(c+dx)}) \int \cos^2(c+dx) dx}{4b^2\sqrt{b}\cos(c+dx)} \\ &= \frac{3\cos^{3/2}(c+dx)\sin(c+dx)}{8b^2d\sqrt{b}\cos(c+dx)} + \frac{\cos^{7/2}(c+dx)\sin(c+dx)}{4b^2d\sqrt{b}\cos(c+dx)} + \frac{(3\sqrt{\cos(c+dx)}) \int 1 dx}{8b^2\sqrt{b}\cos(c+dx)} \\ &= \frac{3x\sqrt{\cos(c+dx)}}{8b^2\sqrt{b}\cos(c+dx)} + \frac{3\cos^{3/2}(c+dx)\sin(c+dx)}{8b^2d\sqrt{b}\cos(c+dx)} + \frac{\cos^{7/2}(c+dx)\sin(c+dx)}{4b^2d\sqrt{b}\cos(c+dx)} \end{aligned}$$

Mathematica [A] time = 0.07, size = 58, normalized size = 0.54

$$\frac{(12(c + dx) + 8 \sin(2(c + dx)) + \sin(4(c + dx)))\sqrt{\cos(c + dx)}}{32b^2d\sqrt{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(13/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (Sqrt[Cos[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])) / (32*b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.85, size = 182, normalized size = 1.70

$$\left[\frac{2\sqrt{b \cos(dx + c)}(2 \cos(dx + c)^2 + 3)\sqrt{\cos(dx + c)} \sin(dx + c) - 3\sqrt{-b} \log(2b \cos(dx + c)^2 + 2\sqrt{b \cos(dx + c)})}{16b^3d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(13/2)/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] [1/16*(2*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 3)*sqrt(cos(d*x + c))*sin(d*x + c) - 3*sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b^3*d), 1/8*(sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 3)*sqrt(cos(d*x + c))*sin(d*x + c) + 3*sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/(b^3*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{13}{2}}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(13/2)/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(13/2)/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.15, size = 62, normalized size = 0.58

$$\frac{\left(\cos^{\frac{5}{2}}(dx + c)\right)\left(2\left(\cos^3(dx + c)\right)\sin(dx + c) + 3\cos(dx + c)\sin(dx + c) + 3dx + 3c\right)}{8d(b \cos(dx + c))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(13/2)/(b*cos(d*x+c))^(5/2), x)

[Out] 1/8/d*cos(d*x+c)^(5/2)*(2*cos(d*x+c)^3*sin(d*x+c)+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)/(b*cos(d*x+c))^(5/2)

maxima [A] time = 1.19, size = 49, normalized size = 0.46

$$\frac{12dx + 12c + \sin(4dx + 4c) + 8 \sin\left(\frac{1}{2} \arctan(\sin(4dx + 4c), \cos(4dx + 4c))\right)}{32b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(13/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(b^(5/2)*d)

mupad [B] time = 1.00, size = 78, normalized size = 0.73

$$\frac{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} (8 \sin(c+dx) + 9 \sin(3c+3dx) + \sin(5c+5dx) + 24dx \cos(c+dx))}{32 b^3 d (\cos(2c+2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(13/2)/(b*cos(c + d*x))^(5/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*b^3*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(13/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

$$3.193 \quad \int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx$$

Optimal. Leaf size=76

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(11/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(b^2*d*Sqrt[b*Cos[c + d*x]]) - (Sqrt[Cos[c + d*x]]*Sin[c + d*x]^3)/(3*b^2*d*Sqrt[b*Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^3(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1-x^2) dx, x, -\sin(c+dx)\right)}{b^2 d \sqrt{b \cos(c+dx)}} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3b^2 d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 48, normalized size = 0.63

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}(\cos(2(c+dx))+5)}{6b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(11/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (Sqrt[Cos[c + d*x]]*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.98, size = 42, normalized size = 0.55

$$\frac{\sqrt{b \cos(dx + c)} (\cos(dx + c)^2 + 2) \sin(dx + c)}{3 b^3 d \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(cos(d*x + c)^2 + 2)*sin(d*x + c)/(b^3*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{11}{2}}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(11/2)/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.10, size = 40, normalized size = 0.53

$$\frac{(2 + \cos^2(dx + c)) \sin(dx + c) \left(\cos^{\frac{5}{2}}(dx + c)\right)}{3d (b \cos(dx + c))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(5/2),x)

[Out] 1/3/d*(2+cos(d*x+c)^2)*sin(d*x+c)*cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2)

maxima [A] time = 1.58, size = 42, normalized size = 0.55

$$\frac{\sin(3dx + 3c) + 9 \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)}{12 b^{\frac{5}{2}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(b^(5/2)*d)

mupad [B] time = 0.60, size = 60, normalized size = 0.79

$$\frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12 b^3 d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(11/2)/(b*cos(c + d*x))^(5/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*b^3*d*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(11/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

$$3.194 \quad \int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx$$

Optimal. Leaf size=69

$$\frac{x\sqrt{\cos(c+dx)}}{2b^2\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2b^2d\sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}+1/2*x*\cos(d*x+c)^{(1/2)}/b^2/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{x\sqrt{\cos(c+dx)}}{2b^2\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2b^2d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^(9/2)/(b*Cos[c + d*x])^(5/2), x]`

[Out] `(x*Sqrt[Cos[c + d*x]])/(2*b^2*Sqrt[b*Cos[c + d*x]]) + (Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*b^2*d*Sqrt[b*Cos[c + d*x]])`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 2635

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*SIN[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned} \int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos^2(c+dx) dx}{b^2\sqrt{b \cos(c+dx)}} \\ &= \frac{\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{2b^2d\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \int 1 dx}{2b^2\sqrt{b \cos(c+dx)}} \\ &= \frac{x\sqrt{\cos(c+dx)}}{2b^2\sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{2b^2d\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 48, normalized size = 0.70

$$\frac{(2(c+dx) + \sin(2(c+dx)))\sqrt{\cos(c+dx)}}{4b^2d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(9/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (Sqrt[Cos[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*b^2*d*Sqrt[b*Cos[c + d*x]])

fricas [A] time = 0.94, size = 157, normalized size = 2.28

$$\left[\frac{2 \sqrt{b \cos(dx + c)} \sqrt{\cos(dx + c)} \sin(dx + c) - \sqrt{-b} \log(2b \cos(dx + c)^2 + 2 \sqrt{b \cos(dx + c)} \sqrt{-b} \sqrt{\cos(dx + c)})}{4b^3d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b^3*d), 1/2*(sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))]/(b^3*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^{\frac{9}{2}}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(9/2)/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.11, size = 42, normalized size = 0.61

$$\frac{\left(\cos^{\frac{5}{2}}(dx + c)\right) (\cos(dx + c) \sin(dx + c) + dx + c)}{2d (b \cos(dx + c))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(5/2), x)

[Out] 1/2/d*cos(d*x+c)^(5/2)*(cos(d*x+c)*sin(d*x+c)+d*x+c)/(b*cos(d*x+c))^(5/2)

maxima [A] time = 1.44, size = 25, normalized size = 0.36

$$\frac{2dx + 2c + \sin(2dx + 2c)}{4b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(5/2), x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(b^(5/2)*d)

mupad [B] time = 0.61, size = 65, normalized size = 0.94

$$\frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (\sin(c + dx) + \sin(3c + 3dx) + 4dx \cos(c + dx))}{4b^3 d (\cos(2c + 2dx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^(9/2)/(b*cos(c + d*x))^(5/2),x)
```

```
[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x)
+ 4*d*x*cos(c + d*x)))/(4*b^3*d*(cos(2*c + 2*d*x) + 1))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**(9/2)/(b*cos(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

$$3.195 \quad \int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=35

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^{(7/2)}/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ $\text{FreeQ}\{a, b, m\}, x] \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ $\text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \cos(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 35, normalized size = 1.00

$$\frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Cos}[c + d*x]^{(7/2)}/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

fricas [A] time = 0.68, size = 31, normalized size = 0.89

$$\frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{b^3 d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b^3*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^{\frac{7}{2}}}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(7/2)/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.10, size = 29, normalized size = 0.83

$$\frac{\sin(dx+c) \left(\cos^{\frac{5}{2}}(dx+c) \right)}{d (b \cos(dx+c))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(5/2),x)

[Out] 1/d*sin(d*x+c)*cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2)

maxima [A] time = 1.63, size = 13, normalized size = 0.37

$$\frac{\sin(dx+c)}{b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] sin(d*x + c)/(b^(5/2)*d)

mupad [B] time = 0.41, size = 47, normalized size = 1.34

$$\frac{\sqrt{\cos(c+dx)} \sin(2c+2dx) \sqrt{b \cos(c+dx)}}{b^3 d (\cos(2c+2dx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c+d*x)^(7/2)/(b*cos(c+d*x))^(5/2),x)

[Out] (cos(c+d*x)^(1/2)*sin(2*c+2*d*x)*(b*cos(c+d*x))^(1/2))/(b^3*d*(cos(2*c+2*d*x)+1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

$$3.196 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=27

$$\frac{x\sqrt{\cos(c+dx)}}{b^2\sqrt{b \cos(c+dx)}}$$

[Out] $x \cos(d*x+c)^{(1/2)}/b^2/(b \cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{x\sqrt{\cos(c+dx)}}{b^2\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(5/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (x*Sqrt[Cos[c + d*x]])/(b^2*Sqrt[b*Cos[c + d*x]])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int 1 dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{x\sqrt{\cos(c+dx)}}{b^2 \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 0.89

$$\frac{x \cos^2(c+dx)}{(b \cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(5/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (x*Cos[c + d*x]^(5/2))/(b*Cos[c + d*x])^(5/2)

fricas [A] time = 0.82, size = 97, normalized size = 3.59

$$\left[\frac{\sqrt{-b} \log(2b \cos(dx+c)^2 + 2\sqrt{b \cos(dx+c)} \sqrt{-b} \sqrt{\cos(dx+c)} \sin(dx+c) - b)}{2b^3d}, \frac{\arctan\left(\frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{\sqrt{b \cos(dx+c)}^3}\right)}{b^2d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [-1/2*sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b)/(b^3*d), arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/(b^(5/2)*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^{\frac{5}{2}}}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(5/2)/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.08, size = 28, normalized size = 1.04

$$\frac{\left(\cos^{\frac{5}{2}}(dx+c)\right)(dx+c)}{d(b \cos(dx+c))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x)

[Out] 1/d*cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2)*(d*x+c)

maxima [A] time = 1.27, size = 26, normalized size = 0.96

$$\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(b^(5/2)*d)

mupad [B] time = 0.28, size = 37, normalized size = 1.37

$$\frac{2x \cos(c+dx)^{\frac{3}{2}} \sqrt{b \cos(c+dx)}}{b^3 (\cos(2c+2dx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(5/2)/(b*cos(c + d*x))^(5/2),x)

[Out] (2*x*cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2))/(b^3*(cos(2*c + 2*d*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

$$3.197 \quad \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=36

$$\frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/b^2/d/(b*cos(d*x+c))^(1/2)

Rubi [A] time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(3/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(b^2*d*Sqrt[b*Cos[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 33, normalized size = 0.92

$$\frac{\cos^5(c+dx) \tanh^{-1}(\sin(c+dx))}{d(b \cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(3/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^(5/2))/(d*(b*Cos[c + d*x])^(5/2))

fricas [A] time = 0.92, size = 116, normalized size = 3.22

$$\left[\frac{\log\left(\frac{-b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right)}{2 b^2 d}, \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b} \sin(dx+c)}{b \sqrt{\cos(dx+c)}}\right)}{b^3 d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3)/(b^(5/2)*d), -sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/(b^3*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^{\frac{3}{2}}}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(3/2)/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.08, size = 42, normalized size = 1.17

$$\frac{2 \left(\cos^{\frac{5}{2}}(dx+c) \right) \operatorname{arctanh} \left(\frac{-1+\cos(dx+c)}{\sin(dx+c)} \right)}{d (b \cos(dx+c))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x)

[Out] -2/d*cos(d*x+c)^(5/2)*arctanh((-1+cos(d*x+c))/sin(d*x+c))/(b*cos(d*x+c))^(5/2)

maxima [B] time = 1.43, size = 65, normalized size = 1.81

$$\frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2 \sin(dx+c) + 1)}{2 b^{\frac{5}{2}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(b^(5/2)*d)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\cos(c+dx)^{\frac{3}{2}}}{(b \cos(c+dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(3/2)/(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^(3/2)/(b*cos(c + d*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

$$3.198 \quad \int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=35

$$\frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[\text{Cos}[c + d*x]]/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $\text{Sin}[c + d*x]/(b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1+x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c+d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \&\& \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^2(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{b^2 d \sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 32, normalized size = 0.91

$$\frac{\sin(c+dx) \cos^3(c+dx)}{d(b \cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Sqrt}[\text{Cos}[c + d*x]]/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(d*(b*\text{Cos}[c + d*x])^{(5/2)})$

fricas [A] time = 0.95, size = 31, normalized size = 0.89

$$\frac{\sqrt{b \cos(dx + c)} \sin(dx + c)}{b^3 d \cos(dx + c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b^3*d*cos(d*x + c)^(3/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\cos(dx + c)}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(cos(d*x + c))/(b*cos(d*x + c))^(5/2), x)

maple [A] time = 0.12, size = 29, normalized size = 0.83

$$\frac{\left(\cos^{\frac{3}{2}}(dx + c)\right) \sin(dx + c)}{d (b \cos(dx + c))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x)

[Out] 1/d*cos(d*x+c)^(3/2)*sin(d*x+c)/(b*cos(d*x+c))^(5/2)

maxima [B] time = 1.35, size = 67, normalized size = 1.91

$$\frac{2 \sqrt{b} \sin(2 dx + 2 c)}{(b^3 \cos(2 dx + 2 c)^2 + b^3 \sin(2 dx + 2 c)^2 + 2 b^3 \cos(2 dx + 2 c) + b^3) d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*sin(2*d*x + 2*c)/((b^3*cos(2*d*x + 2*c)^2 + b^3*sin(2*d*x + 2*c)^2 + 2*b^3*cos(2*d*x + 2*c) + b^3)*d)

mupad [B] time = 1.07, size = 87, normalized size = 2.49

$$\frac{2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (\sin(c + dx) + \sin(3c + 3dx) + \cos(c + dx) 3i + \cos(3c + 3dx) 1i)}{b^3 d (4 \cos(2c + 2dx) + \cos(4c + 4dx) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(1/2)/(b*cos(c + d*x))^(5/2),x)

[Out] (2*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(cos(c + d*x)*3i + sin(c + d*x) + cos(3*c + 3*d*x)*1i + sin(3*c + 3*d*x)))/(b^3*d*(4*cos(2*c + 2*d*x) + cos(4*c + 4*d*x) + 3))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

$$3.199 \quad \int \frac{1}{\sqrt{\cos(c+dx)} (b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=78

$$\frac{\sin(c+dx)}{2b^2d \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2b^2d\sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\sin(d*x+c)/b^2/d/\cos(d*x+c)^{(3/2)}/(b*\cos(d*x+c))^{(1/2)}+1/2*\operatorname{arctanh}(\sin(d*x+c))*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3768, 3770}

$$\frac{\sin(c+dx)}{2b^2d \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{2b^2d\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] `Int[1/(Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(5/2)),x]`

[Out] `(ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(2*b^2*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(2*b^2*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])`

Rule 18

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3770

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{\cos(c+dx)} (b \cos(c+dx))^{5/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^3(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{2b^2d \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{2b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\cos(c+dx)}}{2b^2d\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2b^2d \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 55, normalized size = 0.71

$$\frac{\sqrt{b \cos(c+dx)} (\sin(c+dx) + \cos^2(c+dx) \tanh^{-1}(\sin(c+dx)))}{2b^3d \cos^{\frac{5}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(5/2)),x]

[Out] (Sqrt[b*Cos[c + d*x]]*(ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x])/(2*b^3*d*Cos[c + d*x]^(5/2)))

fricas [A] time = 0.69, size = 207, normalized size = 2.65

$$\frac{\sqrt{b} \cos(dx+c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b \cos(dx+c)} \sqrt{\cos(dx+c)}}{4b^3 d \cos(dx+c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(b^3*d*cos(d*x + c)^3), -1/2*(sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(b^3*d*cos(d*x + c)^3)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx+c))^{\frac{5}{2}} \sqrt{\cos(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(5/2)*sqrt(cos(d*x + c))), x)

maple [A] time = 0.12, size = 104, normalized size = 1.33

$$\frac{\left(\cos^2(dx+c) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) - \left(\cos^2(dx+c) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx+c)\right) \left(\sqrt{\cos(dx+c)}\right)}{2d(b \cos(dx+c))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c))*cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2)

maxima [B] time = 1.56, size = 688, normalized size = 8.82

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] -1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*

```

cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*
arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x + 2*c)
+ 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d
*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 +
4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2
*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/
2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4*d*x + 4*c) +
2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
)) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*
x + 2*c), cos(2*d*x + 2*c))))/((b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x +
2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) +
4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d
*x + 2*c) + b^2)*cos(4*d*x + 4*c))*sqrt(b)*d

```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{\cos(c + dx)} (b \cos(c + dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(5/2)), x)

[Out] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(5/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(5/2), x)

[Out] Timed out

$$3.200 \quad \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=76

$$\frac{\sin^3(c+dx)}{3b^2d \cos^{\frac{5}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{b^2d\sqrt{\cos(c+dx)}\sqrt{b \cos(c+dx)}}$$

[Out] $1/3*\sin(d*x+c)^3/b^2/d/\cos(d*x+c)^{(5/2)}/(b*\cos(d*x+c))^{(1/2)}+\sin(d*x+c)/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 3767}

$$\frac{\sin^3(c+dx)}{3b^2d \cos^{\frac{5}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{b^2d\sqrt{\cos(c+dx)}\sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(5/2)),x]

[Out] Sin[c + d*x]/(b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]^3/(3*b^2*d*Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx &= \frac{\sqrt{\cos(c+dx)} \int \sec^4(c+dx) dx}{b^2\sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{b^2d\sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{b^2d\sqrt{\cos(c+dx)}\sqrt{b \cos(c+dx)}} + \frac{\sin^3(c+dx)}{3b^2d \cos^{\frac{5}{2}}(c+dx)\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 45, normalized size = 0.59

$$\frac{\cos^{\frac{5}{2}}(c+dx) \left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx) \right)}{d(b \cos(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(5/2)),x]

[Out] (Cos[c + d*x]^(5/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*(b*Cos[c + d*x])^(5/2))

fricas [A] time = 0.95, size = 44, normalized size = 0.58

$$\frac{\sqrt{b \cos(dx + c)} (2 \cos(dx + c)^2 + 1) \sin(dx + c)}{3 b^3 d \cos(dx + c)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 1)*sin(d*x + c)/(b^3*d*cos(d*x + c)^(7/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(5/2)*cos(d*x + c)^(3/2)), x)

maple [A] time = 0.11, size = 42, normalized size = 0.55

$$\frac{\sin(dx + c) (2 (\cos^2(dx + c) + 1))}{3d (b \cos(dx + c))^{\frac{5}{2}} \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x)

[Out] 1/3/d*sin(d*x+c)*(2*cos(d*x+c)^2+1)/(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(1/2)

maxima [B] time = 1.51, size = 343, normalized size = 4.51

$$3 (b^2 \cos(6 dx + 6 c))^2 + 9 b^2 \cos(4 dx + 4 c)^2 + 9 b^2 \cos(2 dx + 2 c)^2 + b^2 \sin(6 dx + 6 c)^2 + 9 b^2 \sin(4 dx + 4 c)^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 4/3*((3*cos(2*d*x + 2*c) + 1)*sin(6*d*x + 6*c) + 3*(3*cos(2*d*x + 2*c) + 1)*sin(4*d*x + 4*c) - 3*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) - 9*cos(4*d*x + 4*c)*sin(2*d*x + 2*c))/((b^2*cos(6*d*x + 6*c)^2 + 9*b^2*cos(4*d*x + 4*c)^2 + 9*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(6*d*x + 6*c)^2 + 9*b^2*sin(4*d*x + 4*c)^2 + 18*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*b^2*sin(2*d*x + 2*c)^2 + 6*b^2*cos(2*d*x + 2*c) + b^2 + 2*(3*b^2*cos(4*d*x + 4*c) + 3*b^2*cos(2*d*x + 2*c) + b^2)*cos(6*d*x + 6*c) + 6*(3*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c) + 6*(b^2*sin(4*d*x + 4*c) + b^2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c))*sqrt(b)*d

mupad [B] time = 1.29, size = 131, normalized size = 1.72

$$\frac{2 \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 15i + \cos(4c + 4dx) 6i + \cos(6c + 6dx) 1i + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}{3 b^3 d \sqrt{\cos(c + dx)} (15 \cos(2c + 2dx) + 6 \cos(4c + 4dx) + \cos(6c + 6dx) + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(5/2)),x)
```

```
[Out] (2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*b^3*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

$$3.201 \quad \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx$$

Optimal. Leaf size=116

$$\frac{3 \sin(c+dx)}{8b^2 d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $1/4 * \sin(d*x+c) / b^2 / d / \cos(d*x+c)^{(7/2)} / (b * \cos(d*x+c))^{(1/2)} + 3/8 * \sin(d*x+c) / b^2 / d / \cos(d*x+c)^{(3/2)} / (b * \cos(d*x+c))^{(1/2)} + 3/8 * \operatorname{arctanh}(\sin(d*x+c)) * \cos(d*x+c)^{(1/2)} / b^2 / d / (b * \cos(d*x+c))^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3768, 3770}

$$\frac{3 \sin(c+dx)}{8b^2 d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sqrt{\cos(c+dx)} \tanh^{-1}(\sin(c+dx))}{8b^2 d \sqrt{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(\operatorname{Cos}[c+d*x]^{(5/2)} * (b * \operatorname{Cos}[c+d*x])^{(5/2)}), x]$

[Out] $(3 * \operatorname{ArcTanh}[\operatorname{Sin}[c+d*x]] * \operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]) / (8 * b^2 * d * \operatorname{Sqrt}[b * \operatorname{Cos}[c+d*x]]) + \operatorname{Sin}[c+d*x] / (4 * b^2 * d * \operatorname{Cos}[c+d*x]^{(7/2)} * \operatorname{Sqrt}[b * \operatorname{Cos}[c+d*x]]) + (3 * \operatorname{Sin}[c+d*x]) / (8 * b^2 * d * \operatorname{Cos}[c+d*x]^{(3/2)} * \operatorname{Sqrt}[b * \operatorname{Cos}[c+d*x]])$

Rule 18

$\operatorname{Int}[(u_*) * ((a_*) * (v_*))^{(m_*)} * ((b_*) * (v_*))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[(a^{(m-1/2)} * b^{(n+1/2)} * \operatorname{Sqrt}[a*v]) / \operatorname{Sqrt}[b*v], \operatorname{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3768

$\operatorname{Int}[(\operatorname{csc}[(c_*) + (d_*) * (x_*)] * (b_*))^{(n_*)}, x_Symbol] \rightarrow -\operatorname{Simp}[(b * \operatorname{Cos}[c + d*x] * (b * \operatorname{Csc}[c + d*x])^{(n-1)}) / (d * (n-1)), x] + \operatorname{Dist}[(b^2 * (n-2)) / (n-1), \operatorname{Int}[(b * \operatorname{Csc}[c + d*x])^{(n-2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

$\operatorname{Int}[\operatorname{csc}[(c_*) + (d_*) * (x_*)], x_Symbol] \rightarrow -\operatorname{Simp}[\operatorname{ArcTanh}[\operatorname{Cos}[c + d*x]] / d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{\frac{5}{2}}} dx = \frac{\sqrt{\cos(c+dx)} \int \sec^5(c+dx) dx}{b^2 \sqrt{b\cos(c+dx)}}$$

$$= \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b\cos(c+dx)}} + \frac{(3\sqrt{\cos(c+dx)}) \int \sec^3(c+dx) dx}{4b^2 \sqrt{b\cos(c+dx)}}$$

$$= \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b\cos(c+dx)}} + \frac{3 \sin(c+dx)}{8b^2 d \cos^{\frac{3}{2}}(c+dx) \sqrt{b\cos(c+dx)}}$$

$$= \frac{3 \tanh^{-1}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{8b^2 d \sqrt{b\cos(c+dx)}} + \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b\cos(c+dx)}}$$

Mathematica [A] time = 0.04, size = 66, normalized size = 0.57

$$\frac{\sin(c+dx) (3 \cos^2(c+dx) + 2) + 3 \cos^4(c+dx) \tanh^{-1}(\sin(c+dx))}{8d \cos^{\frac{3}{2}}(c+dx) (b \cos(c+dx))^{\frac{5}{2}}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(5/2)),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x])/(8*d*Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(5/2))

fricas [A] time = 0.83, size = 233, normalized size = 2.01

$$\left[\frac{3 \sqrt{b} \cos(dx+c)^5 \log\left(-\frac{b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2 \sqrt{b \cos(dx+c)} (3 \cos(dx+c)^2 + 2)}{16 b^3 d \cos(dx+c)^5} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c)/(b^3*d*cos(d*x + c)^5), -1/8*(3*sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^5 - sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(b^3*d*cos(d*x + c)^5)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx+c))^{\frac{5}{2}} \cos(dx+c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(5/2)*cos(d*x + c)^(5/2)), x)

maple [A] time = 0.16, size = 121, normalized size = 1.04

$$\frac{3 (\cos^4(dx+c) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) - 3 (\cos^4(dx+c) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - 3 (\cos^2(dx+c) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) + 3 (\cos^2(dx+c) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right))}{8d (b \cos(dx+c))^{\frac{5}{2}} \cos(dx+c)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x)
```

```
[Out] -1/8/d*(3*cos(d*x+c)^4*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^4*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^2*sin(d*x+c)-2*sin(d*x+c))/(b*cos(d*x+c))^(5/2)/cos(d*x+c)^(3/2)
```

maxima [B] time = 1.94, size = 1729, normalized size = 14.91

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")
```

```
[Out] -1/16*(12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))/(b^2*cos(8*d*x + 8*c)^2 + 16*b^2*cos(6*d*x + 6*c)^2 + 36*b^2*cos(4*d*x + 4*c)^2 + 16*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(8*d*x + 8*c)^2 + 16*b^2*sin(6*d*x + 6*c)^2 + 36*b^2*sin(4*d*x + 4*c)^2 + 48*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b^2*sin(2*d*x + 2*c)^2 + 8*b^2*cos(2*d*x + 2*c) + b^2 + 2*(4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*cos(8*d*x + 8*c) + 8*(6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*cos(6*d*x + 6*c) + 12*(4*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c) + 4*(2*b^2*sin(6*d*x + 6*c) + 3*b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c))*sqrt(b)*d)
```


mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^{5/2} (b \cos(c + dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(5/2)), x)

[Out] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(5/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(5/2), x)

[Out] Timed out

3.202 $\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx$

Optimal. Leaf size=82

$$\frac{3 \sin(c + dx) \sqrt[3]{b \cos(c + dx)} \cos^{m+1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m + 4); \frac{1}{6}(3m + 10); \cos^2(c + dx)\right)}{d(3m + 4)\sqrt{\sin^2(c + dx)}}$$

[Out] $-3*\cos(d*x+c)^{(1+m)}*(b*\cos(d*x+c))^{(1/3)}*\text{hypergeom}([1/2, 2/3+1/2*m], [5/3+1/2*m], \cos(d*x+c)^2)*\sin(d*x+c)/d/(4+3*m)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{3 \sin(c + dx) \sqrt[3]{b \cos(c + dx)} \cos^{m+1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m + 4); \frac{1}{6}(3m + 10); \cos^2(c + dx)\right)}{d(3m + 4)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^m*(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*\text{Cos}[c + d*x]^{(1 + m)}*(b*\text{Cos}[c + d*x])^{(1/3)}*\text{Hypergeometric2F1}[1/2, (4 + 3*m)/6, (10 + 3*m)/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*(4 + 3*m)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx &= \frac{\sqrt[3]{b \cos(c + dx)} \int \cos^{\frac{1}{3}+m}(c + dx) dx}{\sqrt[3]{\cos(c + dx)}} \\ &= -\frac{3 \cos^{1+m}(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(4 + 3m); \frac{1}{6}(10 + 3m); \cos^2(c + dx)\right)}{d(4 + 3m)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 82, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c + dx)} \csc(c + dx) \sqrt[3]{b \cos(c + dx)} \cos^{m+1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m + \frac{4}{3}\right); \frac{1}{2}\left(m + \frac{10}{3}\right); \cos^2(c + dx)\right)}{d\left(m + \frac{4}{3}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^m*(b*Cos[c + d*x])^(1/3), x]

[Out] -((Cos[c + d*x]^(1 + m)*(b*Cos[c + d*x])^(1/3)*Csc[c + d*x]*Hypergeometric2F1[1/2, (4/3 + m)/2, (10/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(4/3 + m))

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} \cos(dx + c)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*cos(d*x + c)^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(1/3), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c)^m, x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int (\cos^m(dx + c)) (b \cos(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^m*(b*cos(d*x+c))^(1/3), x)

[Out] int(cos(d*x+c)^m*(b*cos(d*x+c))^(1/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(1/3), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^m (b \cos(c + dx))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^m*(b*cos(c + d*x))^(1/3), x)

[Out] int(cos(c + d*x)^m*(b*cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \cos(c + dx)} \cos^m(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**m*(b*cos(d*x+c))**(1/3), x)

[Out] Integral((b*cos(c + d*x))**(1/3)*cos(c + d*x)**m, x)

3.203 $\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx) (b \cos(c + dx))^{10/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c + dx)\right)}{10b^3 d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/10*(b*\cos(d*x+c))^{(10/3)*\text{hypergeom}([1/2, 5/3], [8/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3 \sin(c + dx) (b \cos(c + dx))^{10/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c + dx)\right)}{10b^3 d \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2*(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(10/3)*\text{Hypergeometric2F1}[1/2, 5/3, 8/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(10*b^3*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2)]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx &= \frac{\int (b \cos(c + dx))^{7/3} dx}{b^2} \\ &= \frac{3(b \cos(c + dx))^{10/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{10b^3 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 63, normalized size = 1.09

$$\frac{3\sqrt{\sin^2(c + dx)} \cos^2(c + dx) \cot(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c + dx)\right)}{10d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*\text{Cos}[c + d*x]^2*(b*\text{Cos}[c + d*x])^{(1/3)*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 5/3, 8/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(10*d)$

fricas [F] time = 0.94, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c)^2, x)

maple [F] time = 0.19, size = 0, normalized size = 0.00

$$\int \left(\cos^2(dx + c)\right) (b \cos(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2*(b*cos(d*x+c))^(1/3),x)

[Out] int(cos(d*x+c)^2*(b*cos(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx)^2 (b \cos(c + dx))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(1/3),x)

[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(1/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(1/3),x)

[Out] Timed out

3.204 $\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right)}{7b^2d\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/7*(b*\cos(d*x+c))^{(7/3)}*\text{hypergeom}([1/2, 7/6], [13/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right)}{7b^2d\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]*(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(7/3)}*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(7*b^2*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx &= \frac{\int (b \cos(c + dx))^{4/3} dx}{b} \\ &= \frac{3(b \cos(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{7b^2d\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 58, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(c + dx)} \cot(c + dx)(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right)}{7bd}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(4/3)}*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(7*b*d)$

fricas [F] time = 0.77, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \cos(dx + c) (b \cos(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)*(b*cos(d*x+c))^(1/3),x)

[Out] int(cos(d*x+c)*(b*cos(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx) (b \cos(c + dx))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)*(b*cos(c + d*x))^(1/3),x)

[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \cos(c + dx)} \cos(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(1/3),x)

[Out] Integral((b*cos(c + d*x))**(1/3)*cos(c + d*x), x)

3.205 $\int \sqrt[3]{b \cos(c + dx)} dx$

Optimal. Leaf size=58

$$-\frac{3 \sin(c + dx)(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c + dx)\right)}{4bd\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/4*(b*\cos(d*x+c))^{(4/3)*\text{hypergeom}([1/2, 2/3], [5/3], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$-\frac{3 \sin(c + dx)(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c + dx)\right)}{4bd\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(4/3)*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(4*b*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2)]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \sqrt[3]{b \cos(c + dx)} dx = -\frac{3(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{4bd\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.04, size = 55, normalized size = 0.95

$$-\frac{3\sqrt{\sin^2(c + dx)} \cot(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c + dx)\right)}{4d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(1/3)*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(4*d)$

fricas [F] time = 1.02, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3), x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/3),x)

[Out] int((b*cos(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int (b \cos(c + dx))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/3),x)

[Out] int((b*cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/3),x)

[Out] Integral((b*cos(c + d*x))**(1/3), x)

3.206 $\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx$

Optimal. Leaf size=53

$$\frac{3 \sin(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3*(b*\cos(d*x+c))^{(1/3)}*\text{hypergeom}([1/6, 1/2], [7/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x], x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(1/3)}*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_)^(n_)), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx &= b \int \frac{1}{(b \cos(c + dx))^{2/3}} dx \\ &= \frac{3 \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 54, normalized size = 1.02

$$\frac{3b \sqrt{\sin^2(c + dx)} \cot(c + dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c + dx)\right)}{d(b \cos(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x], x]

[Out] $(-3*b*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(d*(b*\text{Cos}[c + d*x])^{(2/3)})$

fricas [F] time = 0.74, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c), x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/3)*sec(d*x+c),x)

[Out] int((b*cos(d*x+c))^(1/3)*sec(d*x+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{1/3}}{\cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/3)/cos(c + d*x),x)

[Out] int((b*cos(c + d*x))^(1/3)/cos(c + d*x), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/3)*sec(d*x+c),x)

[Out] Integral((b*cos(c + d*x))**(1/3)*sec(c + d*x), x)

3.207 $\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx$

Optimal. Leaf size=56

$$\frac{3b \sin(c + dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)} (b \cos(c + dx))^{2/3}}$$

[Out] $3/2*b*hypergeom([-1/3, 1/2], [2/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{2/3}/(\sin(d*x+c)^2)^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b \sin(c + dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)} (b \cos(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x]^2,x]

[Out] $(3*b*Hypergeometric2F1[-1/3, 1/2, 2/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/ (2*d*(b*\text{Cos}[c + d*x])^{2/3}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2]) / (b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx &= b^2 \int \frac{1}{(b \cos(c + dx))^{5/3}} dx \\ &= \frac{3b {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \cos(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 56, normalized size = 1.00

$$\frac{3b\sqrt{\sin^2(c + dx)} \csc(c + dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c + dx)\right)}{2d(b \cos(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x]^2,x]

[Out] $(3*b*Csc[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2]) / (2*d*(b*\text{Cos}[c + d*x])^{2/3})$

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c)^2, x)

maple [F] time = 0.11, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} (\sec^2(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/3)*sec(d*x+c)^2,x)

[Out] int((b*cos(d*x+c))^(1/3)*sec(d*x+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{1/3}}{\cos(c + dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/3)/cos(c + d*x)^2,x)

[Out] int((b*cos(c + d*x))^(1/3)/cos(c + d*x)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/3)*sec(d*x+c)**2,x)

[Out] Integral((b*cos(c + d*x))**(1/3)*sec(c + d*x)**2, x)

3.208 $\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \cos^2(c + dx)\right)}{5d\sqrt{\sin^2(c + dx)} (b \cos(c + dx))^{5/3}}$$

[Out] $3/5*b^2*\text{hypergeom}([-5/6, 1/2], [1/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \cos^2(c + dx)\right)}{5d\sqrt{\sin^2(c + dx)} (b \cos(c + dx))^{5/3}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x]^3,x]

[Out] $(3*b^2*\text{Hypergeometric2F1}[-5/6, 1/2, 1/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx &= b^3 \int \frac{1}{(b \cos(c + dx))^{8/3}} dx \\ &= \frac{3b^2 {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{5d(b \cos(c + dx))^{5/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 63, normalized size = 1.09

$$\frac{3\sqrt{\sin^2(c + dx)} \csc(c + dx) \sec^2(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \cos^2(c + dx)\right)}{5d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x]^3,x]

[Out] $(3*(b*\text{Cos}[c + d*x])^{(1/3)}*\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[-5/6, 1/2, 1/6, \text{Cos}[c + d*x]^2]*\text{Sec}[c + d*x]^2*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(5*d)$

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c)^3, x)

maple [F] time = 0.12, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} (\sec^3(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(1/3)*sec(d*x+c)^3,x)

[Out] int((b*cos(d*x+c))^(1/3)*sec(d*x+c)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{\frac{1}{3}}}{\cos(c + dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(1/3)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(1/3)/cos(c + d*x)^3, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(1/3)*sec(d*x+c)**3,x)

[Out] Timed out

3.209 $\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx$

Optimal. Leaf size=82

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{2/3} \cos^{m+1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m + 5); \frac{1}{6}(3m + 11); \cos^2(c + dx)\right)}{d(3m + 5)\sqrt{\sin^2(c + dx)}}$$

[Out] $-3*\cos(d*x+c)^{(1+m)}*(b*\cos(d*x+c))^{(2/3)}*\text{hypergeom}([1/2, 5/6+1/2*m], [11/6+1/2*m], \cos(d*x+c)^2)*\sin(d*x+c)/d/(5+3*m)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{2/3} \cos^{m+1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m + 5); \frac{1}{6}(3m + 11); \cos^2(c + dx)\right)}{d(3m + 5)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^m*(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*\text{Cos}[c + d*x]^{(1 + m)}*(b*\text{Cos}[c + d*x])^{(2/3)}*\text{Hypergeometric2F1}[1/2, (5 + 3*m)/6, (11 + 3*m)/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*(5 + 3*m)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx &= \frac{(b \cos(c + dx))^{2/3} \int \cos^{\frac{2}{3}+m}(c + dx) dx}{\cos^{\frac{2}{3}}(c + dx)} \\ &= \frac{3 \cos^{1+m}(c + dx)(b \cos(c + dx))^{2/3} {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(5 + 3m); \frac{1}{6}(11 + 3m); \cos^2(c + dx)\right)}{d(5 + 3m)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.11, size = 82, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c + dx)} \csc(c + dx)(b \cos(c + dx))^{2/3} \cos^{m+1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m + \frac{5}{3}\right); \frac{1}{2}\left(m + \frac{11}{3}\right); \cos^2(c + dx)\right)}{d\left(m + \frac{5}{3}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^m*(b*Cos[c + d*x])^(2/3), x]

[Out] -((Cos[c + d*x]^(1 + m)*(b*Cos[c + d*x])^(2/3)*Csc[c + d*x]*Hypergeometric2F1[1/2, (5/3 + m)/2, (11/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(5/3 + m))

fricas [F] time = 0.92, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{2}{3}} \cos(dx + c)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(2/3), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(2/3), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m, x)

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int (\cos^m(dx + c)) (b \cos(dx + c))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^m*(b*cos(d*x+c))^(2/3), x)

[Out] int(cos(d*x+c)^m*(b*cos(d*x+c))^(2/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(2/3), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^m (b \cos(c + dx))^{2/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^m*(b*cos(c + d*x))^(2/3), x)

[Out] int(cos(c + d*x)^m*(b*cos(c + d*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^{\frac{2}{3}} \cos^m(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**m*(b*cos(d*x+c))**(2/3), x)

[Out] Integral((b*cos(c + d*x))**(2/3)*cos(c + d*x)**m, x)

3.210 $\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{11/3} {}_2F_1\left(\frac{1}{2}, \frac{11}{6}; \frac{17}{6}; \cos^2(c + dx)\right)}{11b^3 d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/11*(b*\cos(d*x+c))^{(11/3)}*\text{hypergeom}([1/2, 11/6], [17/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{11/3} {}_2F_1\left(\frac{1}{2}, \frac{11}{6}; \frac{17}{6}; \cos^2(c + dx)\right)}{11b^3 d \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2*(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(11/3)}*\text{Hypergeometric2F1}[1/2, 11/6, 17/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/((11*b^3*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2)]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx &= \frac{\int (b \cos(c + dx))^{8/3} dx}{b^2} \\ &= \frac{3(b \cos(c + dx))^{11/3} {}_2F_1\left(\frac{1}{2}, \frac{11}{6}; \frac{17}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{11b^3 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 63, normalized size = 1.09

$$\frac{3\sqrt{\sin^2(c + dx)} \cos^2(c + dx) \cot(c + dx)(b \cos(c + dx))^{2/3} {}_2F_1\left(\frac{1}{2}, \frac{11}{6}; \frac{17}{6}; \cos^2(c + dx)\right)}{11d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*\text{Cos}[c + d*x]^2*(b*\text{Cos}[c + d*x])^{(2/3)}*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 11/6, 17/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/((11*d)$

fricas [F] time = 0.89, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{2}{3}} \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c)^2, x)

maple [F] time = 0.16, size = 0, normalized size = 0.00

$$\int (\cos^2(dx + c)) (b \cos(dx + c))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2*(b*cos(d*x+c))^(2/3),x)

[Out] int(cos(d*x+c)^2*(b*cos(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx)^2 (b \cos(c + dx))^{2/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(2/3),x)

[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(2/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(2/3),x)

[Out] Timed out

3.211 $\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{8/3} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c + dx)\right)}{8b^2d\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/8*(b*\cos(d*x+c))^{(8/3)*\text{hypergeom}([1/2, 4/3], [7/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{8/3} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c + dx)\right)}{8b^2d\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]*(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(8/3)*\text{Hypergeometric2F1}[1/2, 4/3, 7/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(8*b^2*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_)^(n_.), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos(c + dx)(b \cos(c + dx))^{2/3} dx &= \frac{\int (b \cos(c + dx))^{5/3} dx}{b} \\ &= \frac{3(b \cos(c + dx))^{8/3} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{8b^2d\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 58, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(c + dx)} \cot(c + dx)(b \cos(c + dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c + dx)\right)}{8bd}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(5/3)*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 4/3, 7/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(8*b*d)$

fricas [F] time = 0.94, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{2}{3}} \cos(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c), x)

maple [F] time = 0.07, size = 0, normalized size = 0.00

$$\int \cos(dx + c) (b \cos(dx + c))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)*(b*cos(d*x+c))^(2/3),x)

[Out] int(cos(d*x+c)*(b*cos(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx) (b \cos(c + dx))^{2/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)*(b*cos(c + d*x))^(2/3),x)

[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(2/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(2/3),x)

[Out] Timed out

3.212 $\int (b \cos(c + dx))^{2/3} dx$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right)}{5bd\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/5*(b*\cos(d*x+c))^{(5/3)}*\text{hypergeom}([1/2, 5/6], [11/6], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right)}{5bd\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(5/3)}*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(5*b*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2)]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (b \cos(c + dx))^{2/3} dx = -\frac{3(b \cos(c + dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{5bd\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.04, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\sin^2(c + dx)} \cot(c + dx)(b \cos(c + dx))^{2/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right)}{5d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(2/3)}*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(5*d)$

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^{2/3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3), x)

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(2/3),x)

[Out] int((b*cos(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int (b \cos(c + dx))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(2/3),x)

[Out] int((b*cos(c + d*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(2/3),x)

[Out] Integral((b*cos(c + d*x))**(2/3), x)

3.213 $\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx$

Optimal. Leaf size=55

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/2*(b*\cos(d*x+c))^{(2/3)*\text{hypergeom}([1/3, 1/2], [4/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x], x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(2/3)*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(2*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2)]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{2/3} \sec(c + dx) dx &= b \int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx \\ &= -\frac{3(b \cos(c + dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{2d\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 56, normalized size = 1.02

$$\frac{3b\sqrt{\sin^2(c + dx)} \cot(c + dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right)}{2d\sqrt[3]{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x], x]

[Out] $(-3*b*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(2*d*(b*\text{Cos}[c + d*x])^{(1/3)})$

fricas [F] time = 0.96, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c), x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(2/3)*sec(d*x+c),x)

[Out] int((b*cos(d*x+c))^(2/3)*sec(d*x+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{\frac{2}{3}}}{\cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(2/3)/cos(c + d*x),x)

[Out] int((b*cos(c + d*x))^(2/3)/cos(c + d*x), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(2/3)*sec(d*x+c),x)

[Out] Timed out

3.214 $\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx$

Optimal. Leaf size=54

$$\frac{3b \sin(c + dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)} \sqrt[3]{b \cos(c + dx)}}$$

[Out] 3*b*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b \sin(c + dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)} \sqrt[3]{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x]^2,x]

[Out] (3*b*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(b*Cos[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx &= b^2 \int \frac{1}{(b \cos(c + dx))^{4/3}} dx \\ &= \frac{3b {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{d \sqrt[3]{b \cos(c + dx)} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 54, normalized size = 1.00

$$\frac{3b \sqrt{\sin^2(c + dx)} \csc(c + dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right)}{d \sqrt[3]{b \cos(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x]^2,x]

[Out] (3*b*Csc[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sqrt[SIN[c + d*x]^2])/(d*(b*Cos[c + d*x])^(1/3))

fricas [F] time = 0.99, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{2}{3}} \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2, x)

maple [F] time = 0.11, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} (\sec^2(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(2/3)*sec(d*x+c)^2,x)

[Out] int((b*cos(d*x+c))^(2/3)*sec(d*x+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{\frac{2}{3}}}{\cos(c + dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(2/3)/cos(c + d*x)^2,x)

[Out] int((b*cos(c + d*x))^(2/3)/cos(c + d*x)^2, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(2/3)*sec(d*x+c)**2,x)

[Out] Timed out

3.215 $\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right)}{4d\sqrt{\sin^2(c + dx)} (b \cos(c + dx))^{4/3}}$$

[Out] $3/4*b^2*\text{hypergeom}([-2/3, 1/2], [1/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{\wedge}(4/3)/(\sin(d*x+c)^2)^{\wedge}(1/2)$

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right)}{4d\sqrt{\sin^2(c + dx)} (b \cos(c + dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x]^3,x]

[Out] $(3*b^2*\text{Hypergeometric2F1}[-2/3, 1/2, 1/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(4*d*(b*\text{Cos}[c + d*x])^{\wedge}(4/3)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx &= b^3 \int \frac{1}{(b \cos(c + dx))^{7/3}} dx \\ &= \frac{3b^2 {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{4d(b \cos(c + dx))^{4/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 63, normalized size = 1.09

$$\frac{3\sqrt{\sin^2(c + dx)} \csc(c + dx) \sec^2(c + dx) (b \cos(c + dx))^{2/3} {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right)}{4d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(2/3)*Csc[c + d*x]^3,x]

[Out] $(3*(b*\text{Cos}[c + d*x])^{\wedge}(2/3)*\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[-2/3, 1/2, 1/3, \text{Cos}[c + d*x]^2]*\text{Sec}[c + d*x]^2*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(4*d)$

fricas [F] time = 0.99, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{2}{3}} \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3, x)

maple [F] time = 0.12, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} (\sec^3(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(2/3)*sec(d*x+c)^3,x)

[Out] int((b*cos(d*x+c))^(2/3)*sec(d*x+c)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{\frac{2}{3}}}{\cos(c + dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(2/3)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(2/3)/cos(c + d*x)^3, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(2/3)*sec(d*x+c)**3,x)

[Out] Timed out

3.216 $\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx$

Optimal. Leaf size=83

$$\frac{3b \sin(c + dx) \sqrt[3]{b \cos(c + dx)} \cos^{m+2}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m + 7); \frac{1}{6}(3m + 13); \cos^2(c + dx)\right)}{d(3m + 7)\sqrt{\sin^2(c + dx)}}$$

[Out] $-3*b*\cos(d*x+c)^{(2+m)}*(b*\cos(d*x+c))^{(1/3)}*\text{hypergeom}([1/2, 7/6+1/2*m], [13/6+1/2*m], \cos(d*x+c)^2)*\sin(d*x+c)/d/(7+3*m)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{3b \sin(c + dx) \sqrt[3]{b \cos(c + dx)} \cos^{m+2}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m + 7); \frac{1}{6}(3m + 13); \cos^2(c + dx)\right)}{d(3m + 7)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^m*(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*b*\text{Cos}[c + d*x]^{(2 + m)}*(b*\text{Cos}[c + d*x])^{(1/3)}*\text{Hypergeometric2F1}[1/2, (7 + 3*m)/6, (13 + 3*m)/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*(7 + 3*m)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx &= \frac{(b \sqrt[3]{b \cos(c + dx)}) \int \cos^{4/3+m}(c + dx) dx}{\sqrt[3]{\cos(c + dx)}} \\ &= \frac{3b \cos^{2+m}(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(7 + 3m); \frac{1}{6}(13 + 3m); \cos^2(c + dx)\right)}{d(7 + 3m)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.17, size = 82, normalized size = 0.99

$$\frac{\sqrt{\sin^2(c + dx)} \csc(c + dx)(b \cos(c + dx))^{4/3} \cos^{m+1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m + \frac{7}{3}\right); \frac{1}{2}\left(m + \frac{13}{3}\right); \cos^2(c + dx)\right)}{d\left(m + \frac{7}{3}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^m*(b*Cos[c + d*x])^(4/3), x]

[Out] -((Cos[c + d*x]^(1 + m)*(b*Cos[c + d*x])^(4/3)*Csc[c + d*x]*Hypergeometric2F1[1/2, (7/3 + m)/2, (13/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(7/3 + m))

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} b \cos(dx + c)^m \cos(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)^m*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c)^m, x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int (\cos^m(dx + c)) (b \cos(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^m*(b*cos(d*x+c))^(4/3), x)

[Out] int(cos(d*x+c)^m*(b*cos(d*x+c))^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(4/3), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^m (b \cos(c + dx))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^m*(b*cos(c + d*x))^(4/3), x)

[Out] int(cos(c + d*x)^m*(b*cos(c + d*x))^(4/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**m*(b*cos(d*x+c))**(4/3), x)

[Out] Timed out

$$3.217 \quad \int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{13/3} {}_2F_1\left(\frac{1}{2}, \frac{13}{6}; \frac{19}{6}; \cos^2(c + dx)\right)}{13b^3d\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/13*(b*\cos(d*x+c))^{(13/3)}*\text{hypergeom}([1/2, 13/6], [19/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{13/3} {}_2F_1\left(\frac{1}{2}, \frac{13}{6}; \frac{19}{6}; \cos^2(c + dx)\right)}{13b^3d\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2*(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(13/3)}*\text{Hypergeometric2F1}[1/2, 13/6, 19/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(13*b^3*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx &= \frac{\int (b \cos(c + dx))^{10/3} dx}{b^2} \\ &= \frac{3(b \cos(c + dx))^{13/3} {}_2F_1\left(\frac{1}{2}, \frac{13}{6}; \frac{19}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{13b^3d\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 63, normalized size = 1.09

$$\frac{3\sqrt{\sin^2(c + dx)} \cos^2(c + dx) \cot(c + dx)(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{13}{6}; \frac{19}{6}; \cos^2(c + dx)\right)}{13d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*\text{Cos}[c + d*x]^2*(b*\text{Cos}[c + d*x])^{(4/3)}*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 13/6, 19/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(13*d)$

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} b \cos(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c)^2, x)

maple [F] time = 0.16, size = 0, normalized size = 0.00

$$\int \left(\cos^2(dx + c)\right) (b \cos(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2*(b*cos(d*x+c))^(4/3),x)

[Out] int(cos(d*x+c)^2*(b*cos(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx)^2 (b \cos(c + dx))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(4/3),x)

[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(4/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(4/3),x)

[Out] Timed out

$$3.218 \quad \int \cos(c + dx)(b \cos(c + dx))^{4/3} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{10/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c + dx)\right)}{10b^2d\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/10*(b*\cos(d*x+c))^{(10/3)*\text{hypergeom}([1/2, 5/3], [8/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{10/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c + dx)\right)}{10b^2d\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]*(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(10/3)*\text{Hypergeometric2F1}[1/2, 5/3, 8/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(10*b^2*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos(c + dx)(b \cos(c + dx))^{4/3} dx &= \frac{\int (b \cos(c + dx))^{7/3} dx}{b} \\ &= \frac{3(b \cos(c + dx))^{10/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{10b^2d\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 58, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(c + dx)} \cot(c + dx)(b \cos(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c + dx)\right)}{10bd}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(7/3)*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 5/3, 8/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(10*b*d)$

fricas [F] time = 0.92, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} b \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \cos(dx + c) (b \cos(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)*(b*cos(d*x+c))^(4/3),x)

[Out] int(cos(d*x+c)*(b*cos(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx) (b \cos(c + dx))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)*(b*cos(c + d*x))^(4/3),x)

[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(4/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(4/3),x)

[Out] Timed out

3.219 $\int (b \cos(c + dx))^{4/3} dx$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right)}{7bd\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/7*(b*\cos(d*x+c))^{(7/3)}*\text{hypergeom}([1/2, 7/6], [13/6], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$\frac{3 \sin(c + dx)(b \cos(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right)}{7bd\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(7/3)}*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(7*b*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2)]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (b \cos(c + dx))^{4/3} dx = -\frac{3(b \cos(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{7bd\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.00, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\sin^2(c + dx)} \cot(c + dx)(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right)}{7d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(4/3)}*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(7*d)$

fricas [F] time = 0.79, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} b \cos(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3), x)

maple [F] time = 0.02, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(4/3),x)

[Out] int((b*cos(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int (b \cos(c + dx))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(4/3),x)

[Out] int((b*cos(c + d*x))^(4/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(4/3),x)

[Out] Timed out

3.220 $\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx$

Optimal. Leaf size=55

$$-\frac{3 \sin(c + dx)(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c + dx)\right)}{4d\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/4*(b*\cos(d*x+c))^{(4/3)*\text{hypergeom}([1/2, 2/3], [5/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$-\frac{3 \sin(c + dx)(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c + dx)\right)}{4d\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x], x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(4/3)*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(4*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2)]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{4/3} \sec(c + dx) dx &= b \int \sqrt[3]{b \cos(c + dx)} dx \\ &= -\frac{3(b \cos(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{4d\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.00, size = 56, normalized size = 1.02

$$-\frac{3b\sqrt{\sin^2(c + dx)} \cot(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c + dx)\right)}{4d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x], x]

[Out] $(-3*b*(b*\text{Cos}[c + d*x])^{(1/3)*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(4*d)$

fricas [F] time = 0.94, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \cos(dx + c)\right)^{\frac{1}{3}} b \cos(dx + c) \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(4/3)*sec(d*x+c),x)

[Out] int((b*cos(d*x+c))^(4/3)*sec(d*x+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{\frac{4}{3}}}{\cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(4/3)/cos(c + d*x),x)

[Out] int((b*cos(c + d*x))^(4/3)/cos(c + d*x), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(4/3)*sec(d*x+c),x)

[Out] Timed out

3.221 $\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx$

Optimal. Leaf size=54

$$\frac{3b \sin(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c + dx)\right)}{d\sqrt{\sin^2(c + dx)}}$$

[Out] $-3*b*(b*\cos(d*x+c))^{(1/3)}*\text{hypergeom}([1/6, 1/2], [7/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b \sin(c + dx) \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c + dx)\right)}{d\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x]^2,x]

[Out] $(-3*b*(b*\text{Cos}[c + d*x])^{(1/3)}*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx &= b^2 \int \frac{1}{(b \cos(c + dx))^{2/3}} dx \\ &= -\frac{3b \sqrt[3]{b \cos(c + dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c + dx)\right) \sin(c + dx)}{d\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 56, normalized size = 1.04

$$\frac{3b^2 \sqrt{\sin^2(c + dx)} \cot(c + dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c + dx)\right)}{d(b \cos(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x]^2,x]

[Out] $(-3*b^2*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(d*(b*\text{Cos}[c + d*x])^{(2/3)})$

fricas [F] time = 0.81, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^{\frac{1}{3}} b \cos(dx + c) \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c)^2, x)

maple [F] time = 0.12, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} (\sec^2(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(4/3)*sec(d*x+c)^2,x)

[Out] int((b*cos(d*x+c))^(4/3)*sec(d*x+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{\frac{4}{3}}}{\cos(c + dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(4/3)/cos(c + d*x)^2,x)

[Out] int((b*cos(c + d*x))^(4/3)/cos(c + d*x)^2, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(4/3)*sec(d*x+c)**2,x)

[Out] Timed out

3.222 $\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)} (b \cos(c + dx))^{2/3}}$$

[Out] $3/2*b^2*\text{hypergeom}([-1/3, 1/2], [2/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(2/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)} (b \cos(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x]^3,x]

[Out] $(3*b^2*\text{Hypergeometric2F1}[-1/3, 1/2, 2/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/ (2*d*(b*\text{Cos}[c + d*x])^{(2/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx &= b^3 \int \frac{1}{(b \cos(c + dx))^{5/3}} dx \\ &= \frac{3b^2 {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \cos(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 58, normalized size = 1.00

$$\frac{3b^2 \sqrt{\sin^2(c + dx)} \csc(c + dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c + dx)\right)}{2d(b \cos(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x]^3,x]

[Out] $(3*b^2*\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[-1/3, 1/2, 2/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/ (2*d*(b*\text{Cos}[c + d*x])^{(2/3)})$

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^{\frac{1}{3}} b \cos(dx + c) \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c)^3, x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} (\sec^3(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^(4/3)*sec(d*x+c)^3,x)

[Out] int((b*cos(d*x+c))^(4/3)*sec(d*x+c)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^{\frac{4}{3}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^{\frac{4}{3}}}{\cos(c + dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^(4/3)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(4/3)/cos(c + d*x)^3, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**(4/3)*sec(d*x+c)**3,x)

[Out] Timed out

$$3.223 \quad \int \frac{\cos^m(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal. Leaf size=82

$$\frac{3 \sin(c+dx) \cos^{m+1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m+2); \frac{1}{6}(3m+8); \cos^2(c+dx)\right)}{d(3m+2) \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

[Out] $-3 \cos(d*x+c)^{(1+m)} \text{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{3}+\frac{1}{2}*m\right], \left[\frac{4}{3}+\frac{1}{2}*m\right], \cos(d*x+c)^2\right) \sin(d*x+c)/d/(2+3*m)/(b*\cos(d*x+c))^{(1/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{3 \sin(c+dx) \cos^{m+1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m+2); \frac{1}{6}(3m+8); \cos^2(c+dx)\right)}{d(3m+2) \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^m/(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3 \cos[c + d*x]^{(1+m)} \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(2+3*m)}{6}, \frac{(8+3*m)}{6}, \cos[c + d*x]^2\right] \sin[c + d*x]) / (d*(2+3*m)*(b*\cos[c + d*x])^{(1/3)} \sqrt{\sin[c + d*x]^2})$

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^m(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx &= \frac{\sqrt[3]{\cos(c+dx)} \int \cos^{-\frac{1}{3}+m}(c+dx) dx}{\sqrt[3]{b \cos(c+dx)}} \\ &= -\frac{3 \cos^{1+m}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(2+3m); \frac{1}{6}(8+3m); \cos^2(c+dx)\right) \sin(c+dx)}{d(2+3m) \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.11, size = 82, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c+dx)} \csc(c+dx) \cos^{m+1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m+\frac{2}{3}\right); \frac{1}{2}\left(m+\frac{8}{3}\right); \cos^2(c+dx)\right)}{d\left(m+\frac{2}{3}\right) \sqrt[3]{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^m/(b*Cos[c + d*x])^(1/3), x]

[Out] -((Cos[c + d*x]^(1 + m)*Csc[c + d*x]*Hypergeometric2F1[1/2, (2/3 + m)/2, (8/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(2/3 + m)*(b*Cos[c + d*x])^(1/3))

fricas [F] time = 0.77, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^m}{b \cos(dx + c)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(1/3), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(1/3), x)

maple [F] time = 0.07, size = 0, normalized size = 0.00

$$\int \frac{\cos^m(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^m/(b*cos(d*x+c))^(1/3), x)

[Out] int(cos(d*x+c)^m/(b*cos(d*x+c))^(1/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(1/3), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^m}{(b \cos(c + dx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^m/(b*cos(c + d*x))^(1/3), x)

[Out] int(cos(c + d*x)^m/(b*cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^m(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**m/(b*cos(d*x+c))**(1/3), x)

[Out] Integral(cos(c + d*x)**m/(b*cos(c + d*x))**(1/3), x)

$$3.224 \quad \int \frac{\cos^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{8/3} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c+dx)\right)}{8b^3 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/8*(b*\cos(d*x+c))^{(8/3)}*\text{hypergeom}([1/2, 4/3], [7/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{8/3} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c+dx)\right)}{8b^3 d \sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(8/3)}*\text{Hypergeometric2F1}[1/2, 4/3, 7/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(8*b^3*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx &= \frac{\int (b \cos(c+dx))^{5/3} dx}{b^2} \\ &= -\frac{3(b \cos(c+dx))^{8/3} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{8b^3 d \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 63, normalized size = 1.09

$$\frac{3\sqrt{\sin^2(c+dx)} \cos^2(c+dx) \cot(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c+dx)\right)}{8d\sqrt[3]{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*\cos[c + d*x]^2*\cot[c + d*x]*\text{Hypergeometric2F1}[1/2, 4/3, 7/3, \cos[c + d*x]^2]*\text{sqrt}[\sin[c + d*x]^2])/(8*d*(b*\cos[c + d*x])^{(1/3)})$

fricas [F] time = 1.05, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")`

[Out] `integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)/b, x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="giac")`

[Out] `integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(1/3), x)`

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^2/(b*cos(d*x+c))^(1/3),x)`

[Out] `int(cos(d*x+c)^2/(b*cos(d*x+c))^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(1/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2/(b*cos(c + d*x))^(1/3),x)`

[Out] `int(cos(c + d*x)^2/(b*cos(c + d*x))^(1/3), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(1/3),x)
```

```
[Out] Timed out
```

$$3.225 \quad \int \frac{\cos(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right)}{5b^2 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/5*(b*\cos(d*x+c))^{5/3}*hypergeom([1/2, 5/6], [11/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{1/2}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right)}{5b^2 d \sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*(b*\cos[c + d*x])^{5/3}*Hypergeometric2F1[1/2, 5/6, 11/6, \cos[c + d*x]^2]*\sin[c + d*x])/(5*b^2*d*\sqrt{\sin[c + d*x]^2})$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx &= \frac{\int (b \cos(c+dx))^{2/3} dx}{b} \\ &= \frac{3(b \cos(c+dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{5b^2 d \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.00, size = 58, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(c+dx)} \cot(c+dx)(b \cos(c+dx))^{2/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right)}{5bd}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(1/3), x]

[Out] $(-3*(b*\cos[c + d*x])^{2/3}*Cot[c + d*x]*Hypergeometric2F1[1/2, 5/6, 11/6, \cos[c + d*x]^2]*\sqrt{\sin[c + d*x]^2})/(5*b*d)$

fricas [F] time = 1.02, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}}}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(1/3), x)

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*cos(d*x+c))^(1/3),x)

[Out] int(cos(d*x+c)/(b*cos(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)/(b*cos(c + d*x))^(1/3),x)

[Out] int(cos(c + d*x)/(b*cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))**(1/3),x)

[Out] Integral(cos(c + d*x)/(b*cos(c + d*x))**(1/3), x)

$$3.226 \quad \int \frac{1}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal. Leaf size=58

$$-\frac{3 \sin(c+dx)(b \cos(c+dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c+dx)\right)}{2bd\sqrt{\sin^2(c+dx)}}$$

[Out] $-3/2*(b*\cos(d*x+c))^{(2/3)}*\text{hypergeom}([1/3, 1/2], [4/3], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$-\frac{3 \sin(c+dx)(b \cos(c+dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c+dx)\right)}{2bd\sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(-1/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(2/3)}*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(2*b*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{\sqrt[3]{b \cos(c+dx)}} dx = -\frac{3(b \cos(c+dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{2bd\sqrt{\sin^2(c+dx)}}$$

Mathematica [A] time = 0.00, size = 55, normalized size = 0.95

$$-\frac{3\sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c+dx)\right)}{2d\sqrt[3]{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(-1/3), x]

[Out] $(-3*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(2*d*(b*\text{Cos}[c + d*x])^{(1/3)})$

fricas [F] time = 0.96, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{2/3}}{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3), x)

maple [F] time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(d*x+c))^(1/3),x)

[Out] int(1/(b*cos(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(b \cos(c + dx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(c + d*x))^(1/3),x)

[Out] int(1/(b*cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))**(1/3),x)

[Out] Integral((b*cos(c + d*x))**(-1/3), x)

$$3.227 \quad \int \frac{\sec(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal. Leaf size=53

$$\frac{3 \sin(c+dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c+dx)\right)}{d \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c+dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c+dx)\right)}{d \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(1/3), x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(b*Cos[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx &= b \int \frac{1}{(b \cos(c+dx))^{4/3}} dx \\ &= \frac{3 {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{d \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 54, normalized size = 1.02

$$\frac{3b \sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c+dx)\right)}{d(b \cos(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(1/3), x]

[Out] (3*b*Cot[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(b*Cos[c + d*x])^(4/3))

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)}{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(1/3), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*cos(d*x+c))^(1/3),x)

[Out] int(sec(d*x+c)/(b*cos(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx) (b \cos(c + dx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(1/3)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(1/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(1/3),x)

[Out] Integral(sec(c + d*x)/(b*cos(c + d*x))**(1/3), x)

$$3.228 \quad \int \frac{\sec^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal. Leaf size=56

$$\frac{3b \sin(c+dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{4/3}}$$

[Out] $3/4*b*hypergeom([-2/3, 1/2], [1/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{4/3}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b \sin(c+dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Cos[c + d*x])^(1/3), x]

[Out] $(3*b*Hypergeometric2F1[-2/3, 1/2, 1/3, \cos[c + d*x]^2]*\sin[c + d*x])/(4*d*(b*\cos[c + d*x])^{4/3}*Sqrt[\sin[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx &= b^2 \int \frac{1}{(b \cos(c+dx))^{7/3}} dx \\ &= \frac{3b {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{4d (b \cos(c+dx))^{4/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 58, normalized size = 1.04

$$\frac{3b^2 \sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c+dx)\right)}{4d (b \cos(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(1/3), x]

[0ut] $(3*b^2*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[-2/3, 1/2, 1/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(4*d*(b*\text{Cos}[c + d*x])^{(7/3)})$

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)^2}{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")`

[0ut] `integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2/(b*cos(d*x + c)), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="giac")`

[0ut] `integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(1/3), x)`

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^2/(b*cos(d*x+c))^(1/3),x)`

[0ut] `int(sec(d*x+c)^2/(b*cos(d*x+c))^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")`

[0ut] `integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(1/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^2 (b \cos(c + dx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(1/3)),x)`

[0ut] `int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(1/3)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(1/3), x)
```

```
[Out] Integral(sec(c + d*x)**2/(b*cos(c + d*x))**(1/3), x)
```

$$3.229 \quad \int \frac{\sec^3(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c+dx)\right)}{7d\sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{7/3}}$$

[Out] 3/7*b^2*hypergeom([-7/6, 1/2], [-1/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(7/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c+dx)\right)}{7d\sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3/(b*Cos[c + d*x])^(1/3), x]

[Out] (3*b^2*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*Sin[c + d*x])/(7*d*(b*Cos[c + d*x])^(7/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx &= b^3 \int \frac{1}{(b \cos(c+dx))^{10/3}} dx \\ &= \frac{3b^2 {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{7d(b \cos(c+dx))^{7/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 58, normalized size = 1.00

$$\frac{3b^2 \sqrt{\sin^2(c+dx)} \csc(c+dx) {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c+dx)\right)}{7d(b \cos(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(1/3), x]

[Out] $(3*b^2*\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[-7/6, 1/2, -1/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(7*d*(b*\text{Cos}[c + d*x])^{(7/3)})$

fricas [F] time = 0.49, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)^3}{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")`

[Out] `integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3/(b*cos(d*x + c)), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/3),x, algorithm="giac")`

[Out] `integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(1/3), x)`

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^3/(b*cos(d*x+c))^(1/3),x)`

[Out] `int(sec(d*x+c)^3/(b*cos(d*x+c))^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(1/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^3 (b \cos(c + dx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(1/3)),x)`

[Out] `int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(1/3)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(1/3), x)
```

```
[Out] Integral(sec(c + d*x)**3/(b*cos(c + d*x))**(1/3), x)
```

$$3.230 \quad \int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal. Leaf size=82

$$\frac{3 \sin(c+dx) \cos^{m+1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m+1); \frac{1}{6}(3m+7); \cos^2(c+dx)\right)}{d(3m+1)\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{2/3}}$$

[Out] $-3*\cos(d*x+c)^{(1+m)}*\text{hypergeom}([1/2, 1/6+1/2*m], [7/6+1/2*m], \cos(d*x+c)^2)*\sin(d*x+c)/d/(1+3*m)/(b*\cos(d*x+c))^{(2/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{3 \sin(c+dx) \cos^{m+1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m+1); \frac{1}{6}(3m+7); \cos^2(c+dx)\right)}{d(3m+1)\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^m/(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*\text{Cos}[c + d*x]^{(1 + m)}*\text{Hypergeometric2F1}[1/2, (1 + 3*m)/6, (7 + 3*m)/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*(1 + 3*m)*(b*\text{Cos}[c + d*x])^{(2/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{2/3}} dx &= \frac{\cos^{2/3}(c+dx) \int \cos^{-2/3+m}(c+dx) dx}{(b \cos(c+dx))^{2/3}} \\ &= -\frac{3 \cos^{1+m}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(1+3m); \frac{1}{6}(7+3m); \cos^2(c+dx)\right) \sin(c+dx)}{d(1+3m)(b \cos(c+dx))^{2/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 82, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c+dx)} \csc(c+dx) \cos^{m+1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m + \frac{1}{3}\right); \frac{1}{2}\left(m + \frac{7}{3}\right); \cos^2(c+dx)\right)}{d\left(m + \frac{1}{3}\right)(b \cos(c+dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^m/(b*Cos[c + d*x])^(2/3), x]

[Out] -((Cos[c + d*x]^(1 + m)*Csc[c + d*x]*Hypergeometric2F1[1/2, (1/3 + m)/2, (7/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(1/3 + m)*(b*Cos[c + d*x])^(2/3))

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^m}{b \cos(dx + c)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(2/3), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*cos(d*x + c)^m/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(2/3), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(2/3), x)

maple [F] time = 0.07, size = 0, normalized size = 0.00

$$\int \frac{\cos^m(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^m/(b*cos(d*x+c))^(2/3), x)

[Out] int(cos(d*x+c)^m/(b*cos(d*x+c))^(2/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(2/3), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^m}{(b \cos(c + dx))^{2/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^m/(b*cos(c + d*x))^(2/3), x)

[Out] int(cos(c + d*x)^m/(b*cos(c + d*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**m/(b*cos(d*x+c))**(2/3), x)

[Out] Integral(cos(c + d*x)**m/(b*cos(c + d*x))**(2/3), x)

$$3.231 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal. Leaf size=58

$$-\frac{3 \sin(c+dx)(b \cos(c+dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right)}{7b^3 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/7*(b*\cos(d*x+c))^{(7/3)}*\text{hypergeom}([1/2, 7/6], [13/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$-\frac{3 \sin(c+dx)(b \cos(c+dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right)}{7b^3 d \sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(7/3)}*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(7*b^3*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx &= \frac{\int (b \cos(c+dx))^{4/3} dx}{b^2} \\ &= -\frac{3(b \cos(c+dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{7b^3 d \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 63, normalized size = 1.09

$$-\frac{3\sqrt{\sin^2(c+dx)} \cos^2(c+dx) \cot(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right)}{7d(b \cos(c+dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*\text{Cos}[c + d*x]^2*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(7*d*(b*\text{Cos}[c + d*x])^{(2/3)})$

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")`

[Out] `integral((b*cos(d*x + c))^(1/3)*cos(d*x + c)/b, x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="giac")`

[Out] `integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(2/3), x)`

maple [F] time = 0.14, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^2/(b*cos(d*x+c))^(2/3),x)`

[Out] `int(cos(d*x+c)^2/(b*cos(d*x+c))^(2/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(2/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{2/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2/(b*cos(c + d*x))^(2/3),x)`

[Out] `int(cos(c + d*x)^2/(b*cos(c + d*x))^(2/3), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(2/3),x)
```

```
[Out] Timed out
```

$$3.232 \quad \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right)}{4b^2 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/4*(b*\cos(d*x+c))^{(4/3)}*\text{hypergeom}([1/2, 2/3], [5/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right)}{4b^2 d \sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(4/3)}*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(4*b^2*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{2/3}} dx &= \frac{\int \sqrt[3]{b \cos(c+dx)} dx}{b} \\ &= \frac{3(b \cos(c+dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{4b^2 d \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.00, size = 58, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(c+dx)} \cot(c+dx) \sqrt[3]{b \cos(c+dx)} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right)}{4bd}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(1/3)}*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(4*b*d)$

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{1}{3}}}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(2/3), x)

maple [F] time = 0.07, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*cos(d*x+c))^(2/3),x)

[Out] int(cos(d*x+c)/(b*cos(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)/(b*cos(c + d*x))^(2/3),x)

[Out] int(cos(c + d*x)/(b*cos(c + d*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))**(2/3),x)

[Out] Integral(cos(c + d*x)/(b*cos(c + d*x))**(2/3), x)

$$3.233 \quad \int \frac{1}{(b \cos(c+dx))^{2/3}} dx$$

Optimal. Leaf size=56

$$\frac{3 \sin(c+dx) \sqrt[3]{b \cos(c+dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right)}{bd \sqrt{\sin^2(c+dx)}}$$

[Out] $-3*(b*\cos(d*x+c))^{(1/3)}*\text{hypergeom}([1/6, 1/2], [7/6], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$\frac{3 \sin(c+dx) \sqrt[3]{b \cos(c+dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right)}{bd \sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^(-2/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(1/3)}*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(b*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{(b \cos(c+dx))^{2/3}} dx = -\frac{3 \sqrt[3]{b \cos(c+dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{bd \sqrt{\sin^2(c+dx)}}$$

Mathematica [A] time = 0.00, size = 53, normalized size = 0.95

$$\frac{3 \sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right)}{d(b \cos(c+dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^(-2/3), x]

[Out] $(-3*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(d*(b*\text{Cos}[c + d*x])^{(2/3)})$

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{1/3}}{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)/(b*cos(d*x + c)), x)

maple [F] time = 0.03, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(d*x+c))^(2/3),x)

[Out] int(1/(b*cos(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)/(b*cos(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(c + d*x))^(2/3),x)

[Out] int(1/(b*cos(c + d*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))**(2/3),x)

[Out] Integral((b*cos(c + d*x))**(-2/3), x)

$$3.234 \quad \int \frac{\sec(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal. Leaf size=55

$$\frac{3 \sin(c+dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c+dx)\right)}{2d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{2/3}}$$

[Out] 3/2*hypergeom([-1/3, 1/2], [2/3], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(2/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c+dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c+dx)\right)}{2d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(2/3), x]

[Out] (3*Hypergeometric2F1[-1/3, 1/2, 2/3, Cos[c + d*x]^2]*Sin[c + d*x])/(2*d*(b*Cos[c + d*x])^(2/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec(c+dx)}{(b \cos(c+dx))^{2/3}} dx &= b \int \frac{1}{(b \cos(c+dx))^{5/3}} dx \\ &= \frac{3 {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{2d (b \cos(c+dx))^{2/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 56, normalized size = 1.02

$$\frac{3b \sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c+dx)\right)}{2d (b \cos(c+dx))^{5/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(2/3), x]

[Out] (3*b*Cot[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(2*d*(b*Cos[c + d*x])^(5/3))

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)}{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(2/3), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*cos(d*x+c))^(2/3),x)

[Out] int(sec(d*x+c)/(b*cos(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx) (b \cos(c + dx))^{2/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(2/3)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(2/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(2/3),x)

[Out] Integral(sec(c + d*x)/(b*cos(c + d*x))**(2/3), x)

$$3.235 \quad \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal. Leaf size=56

$$\frac{3b \sin(c+dx) {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \cos^2(c+dx)\right)}{5d\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{5/3}}$$

[Out] 3/5*b*hypergeom([-5/6, 1/2], [1/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(5/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b \sin(c+dx) {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \cos^2(c+dx)\right)}{5d\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{5/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Cos[c + d*x])^(2/3), x]

[Out] (3*b*Hypergeometric2F1[-5/6, 1/2, 1/6, Cos[c + d*x]^2]*Sin[c + d*x])/(5*d*(b*Cos[c + d*x])^(5/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx &= b^2 \int \frac{1}{(b \cos(c+dx))^{8/3}} dx \\ &= \frac{3b {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{5d(b \cos(c+dx))^{5/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 58, normalized size = 1.04

$$\frac{3b^2 \sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \cos^2(c+dx)\right)}{5d(b \cos(c+dx))^{8/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(2/3), x]

[Out] (3*b^2*Cot[c + d*x]*Hypergeometric2F1[-5/6, 1/2, 1/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(5*d*(b*Cos[c + d*x])^(8/3))

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^2}{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)^2/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(2/3), x)

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2/(b*cos(d*x+c))^(2/3),x)

[Out] int(sec(d*x+c)^2/(b*cos(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^2 (b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(2/3)),x)

[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(2/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(2/3),x)

[Out] Integral(sec(c + d*x)**2/(b*cos(c + d*x))**(2/3), x)

$$3.236 \quad \int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(-\frac{4}{3}, \frac{1}{2}; -\frac{1}{3}; \cos^2(c+dx)\right)}{8d\sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{8/3}}$$

[Out] $3/8*b^2*\text{hypergeom}([-4/3, 1/2], [-1/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(8/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(-\frac{4}{3}, \frac{1}{2}; -\frac{1}{3}; \cos^2(c+dx)\right)}{8d\sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{8/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3/(b*Cos[c + d*x])^(2/3), x]

[Out] $(3*b^2*\text{Hypergeometric2F1}[-4/3, 1/2, -1/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(8*d*(b*\text{Cos}[c + d*x])^{(8/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{2/3}} dx &= b^3 \int \frac{1}{(b \cos(c+dx))^{11/3}} dx \\ &= \frac{3b^2 {}_2F_1\left(-\frac{4}{3}, \frac{1}{2}; -\frac{1}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{8d(b \cos(c+dx))^{8/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 58, normalized size = 1.00

$$\frac{3b^2 \sqrt{\sin^2(c+dx)} \csc(c+dx) {}_2F_1\left(-\frac{4}{3}, \frac{1}{2}; -\frac{1}{3}; \cos^2(c+dx)\right)}{8d(b \cos(c+dx))^{8/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(2/3), x]

[Out] $(3*b^2*\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[-4/3, 1/2, -1/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(8*d*(b*\text{Cos}[c + d*x])^{(8/3)})$

fricas [F] time = 0.96, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^3}{b \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)^3/(b*cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(2/3), x)

maple [F] time = 0.12, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(dx + c)}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^3/(b*cos(d*x+c))^(2/3),x)

[Out] int(sec(d*x+c)^3/(b*cos(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^3 (b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(2/3)),x)

[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(2/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(2/3),x)

[Out] Integral(sec(c + d*x)**3/(b*cos(c + d*x))**(2/3), x)

$$3.237 \quad \int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal. Leaf size=83

$$\frac{3 \sin(c+dx) \cos^m(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m-1); \frac{1}{6}(3m+5); \cos^2(c+dx)\right)}{bd(1-3m)\sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

[Out] 3*cos(d*x+c)^m*hypergeom([1/2, -1/6+1/2*m], [5/6+1/2*m], cos(d*x+c)^2)*sin(d*x+c)/b/d/(1-3*m)/(b*cos(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{3 \sin(c+dx) \cos^m(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(3m-1); \frac{1}{6}(3m+5); \cos^2(c+dx)\right)}{bd(1-3m)\sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^m/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*Cos[c + d*x]^m*Hypergeometric2F1[1/2, (-1 + 3*m)/6, (5 + 3*m)/6, Cos[c + d*x]^2]*Sin[c + d*x])/(b*d*(1 - 3*m)*(b*Cos[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{4/3}} dx &= \frac{\sqrt[3]{\cos(c+dx)} \int \cos^{-\frac{4}{3}+m}(c+dx) dx}{b \sqrt[3]{b \cos(c+dx)}} \\ &= \frac{3 \cos^m(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(-1+3m); \frac{1}{6}(5+3m); \cos^2(c+dx)\right) \sin(c+dx)}{bd(1-3m)\sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 82, normalized size = 0.99

$$\frac{\sqrt{\sin^2(c+dx)} \csc(c+dx) \cos^{m+1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m-\frac{1}{3}\right); \frac{1}{2}\left(m+\frac{5}{3}\right); \cos^2(c+dx)\right)}{d\left(m-\frac{1}{3}\right)(b \cos(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^m/(b*Cos[c + d*x])^(4/3), x]

[Out] -((Cos[c + d*x]^(1 + m)*Csc[c + d*x]*Hypergeometric2F1[1/2, (-1/3 + m)/2, (5/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-1/3 + m)*(b*Cos[c + d*x])^(4/3))

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^m}{b^2 \cos(dx + c)^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(4/3), x)

maple [F] time = 0.07, size = 0, normalized size = 0.00

$$\int \frac{\cos^m(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^m/(b*cos(d*x+c))^(4/3), x)

[Out] int(cos(d*x+c)^m/(b*cos(d*x+c))^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(4/3), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^m}{(b \cos(c + dx))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^m/(b*cos(c + d*x))^(4/3), x)

[Out] int(cos(c + d*x)^m/(b*cos(c + d*x))^(4/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**m/(b*cos(d*x+c))**(4/3), x)

[Out] Integral(cos(c + d*x)**m/(b*cos(c + d*x))**(4/3), x)

$$3.238 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right)}{5b^3 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/5*(b*\cos(d*x+c))^{(5/3)}*\text{hypergeom}([1/2, 5/6], [11/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^{3/d}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right)}{5b^3 d \sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(5/3)}*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(5*b^{3*d}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx &= \frac{\int (b \cos(c+dx))^{2/3} dx}{b^2} \\ &= \frac{3(b \cos(c+dx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{5b^3 d \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.00, size = 58, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(c+dx)} \cot(c+dx)(b \cos(c+dx))^{2/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right)}{5b^2 d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*(b*\text{Cos}[c + d*x])^{(2/3)}*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(5*b^2*d)$

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}}}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(4/3), x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2/(b*cos(d*x+c))^(4/3),x)

[Out] int(cos(d*x+c)^2/(b*cos(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(4/3),x)

[Out] int(cos(c + d*x)^2/(b*cos(c + d*x))^(4/3), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(4/3),x)

[Out] Timed out

$$3.239 \quad \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c+dx)\right)}{2b^2 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/2*(b*\cos(d*x+c))^{(2/3)}*\text{hypergeom}([1/3, 1/2], [4/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c+dx)(b \cos(c+dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c+dx)\right)}{2b^2 d \sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*(b*\cos[c + d*x])^{(2/3)}*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \cos[c + d*x]^2]*\sin[c + d*x])/(2*b^2*d*\text{Sqrt}[\sin[c + d*x]^2])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{4/3}} dx &= \frac{\int \frac{1}{\sqrt[3]{b \cos(c+dx)}} dx}{b} \\ &= -\frac{3(b \cos(c+dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{2b^2 d \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 58, normalized size = 1.00

$$\frac{3\sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c+dx)\right)}{2bd\sqrt[3]{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(4/3), x]

[Out] $(-3*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[c + d*x]^2]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(2*b*d*(b*\text{Cos}[c + d*x])^{(1/3)})$

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}}}{b^2 \cos(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")`

[Out] `integral((b*cos(d*x + c))^(2/3)/(b^2*cos(d*x + c)), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="giac")`

[Out] `integrate(cos(d*x + c)/(b*cos(d*x + c))^(4/3), x)`

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)/(b*cos(d*x+c))^(4/3),x)`

[Out] `int(cos(d*x+c)/(b*cos(d*x+c))^(4/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)/(b*cos(d*x + c))^(4/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)/(b*cos(c + d*x))^(4/3),x)`

[Out] `int(cos(c + d*x)/(b*cos(c + d*x))^(4/3), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))**(4/3),x)`

[Out] Timed out

$$3.240 \quad \int \frac{1}{(b \cos(c+dx))^{4/3}} dx$$

Optimal. Leaf size=56

$$\frac{3 \sin(c+dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c+dx)\right)}{bd \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*sin(d*x+c)/b/d/(b*cos(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2643}

$$\frac{3 \sin(c+dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c+dx)\right)}{bd \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^(-4/3), x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sin[c + d*x])/(b*d*(b*cos[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{(b \cos(c+dx))^{4/3}} dx = \frac{3 {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{bd \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}}$$

Mathematica [A] time = 0.00, size = 53, normalized size = 0.95

$$\frac{3 \sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c+dx)\right)}{d(b \cos(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^(-4/3), x]

[Out] (3*Cot[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(b*cos[c + d*x])^(4/3))

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{2/3}}{b^2 \cos(dx + c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")
 [Out] integral((b*cos(d*x + c))^(2/3)/(b^2*cos(d*x + c)^2), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(4/3),x, algorithm="giac")
 [Out] integrate((b*cos(d*x + c))^(-4/3), x)
maple [F] time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(d*x+c))^(4/3),x)
 [Out] int(1/(b*cos(d*x+c))^(4/3),x)
maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")
 [Out] integrate((b*cos(d*x + c))^(-4/3), x)
mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(b \cos(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*cos(c + d*x))^(4/3),x)
 [Out] int(1/(b*cos(c + d*x))^(4/3), x)
sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \cos(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*cos(d*x+c))**(4/3),x)
 [Out] Integral((b*cos(c + d*x))**(-4/3), x)

$$3.241 \quad \int \frac{\sec(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal. Leaf size=55

$$\frac{3 \sin(c+dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c+dx)\right)}{4d\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{4/3}}$$

[Out] 3/4*hypergeom([-2/3, 1/2], [1/3], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(4/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{3 \sin(c+dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c+dx)\right)}{4d\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*Sin[c + d*x])/(4*d*(b*Cos[c + d*x])^(4/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec(c+dx)}{(b \cos(c+dx))^{4/3}} dx &= b \int \frac{1}{(b \cos(c+dx))^{7/3}} dx \\ &= \frac{3 {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{4d(b \cos(c+dx))^{4/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 56, normalized size = 1.02

$$\frac{3b\sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c+dx)\right)}{4d(b \cos(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*b*Cot[c + d*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(4*d*(b*Cos[c + d*x])^(7/3))

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)}{b^2 \cos(dx + c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(4/3), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*cos(d*x+c))^(4/3),x)

[Out] int(sec(d*x+c)/(b*cos(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx) (b \cos(c + dx))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(4/3)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(4/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(4/3),x)

[Out] Integral(sec(c + d*x)/(b*cos(c + d*x))**(4/3), x)

$$3.242 \quad \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal. Leaf size=56

$$\frac{3b \sin(c+dx) {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c+dx)\right)}{7d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{7/3}}$$

[Out] 3/7*b*hypergeom([-7/6, 1/2], [-1/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c)^(7/3)/(sin(d*x+c)^2)^(1/2))

Rubi [A] time = 0.06, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b \sin(c+dx) {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c+dx)\right)}{7d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*b*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*Sin[c + d*x])/(7*d*(b*Cos[c + d*x])^(7/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx &= b^2 \int \frac{1}{(b \cos(c+dx))^{10/3}} dx \\ &= \frac{3b {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c+dx)\right) \sin(c+dx)}{7d (b \cos(c+dx))^{7/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 58, normalized size = 1.04

$$\frac{3b^2 \sqrt{\sin^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c+dx)\right)}{7d (b \cos(c+dx))^{10/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*b^2*Cot[c + d*x]*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(7*d*(b*Cos[c + d*x])^(10/3))

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)^2}{b^2 \cos(dx + c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(4/3), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2/(b*cos(d*x+c))^(4/3),x)

[Out] int(sec(d*x+c)^2/(b*cos(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^2 (b \cos(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(4/3)),x)

[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(4/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(4/3),x)

[Out] Integral(sec(c + d*x)**2/(b*cos(c + d*x))**(4/3), x)

$$3.243 \quad \int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(-\frac{5}{3}, \frac{1}{2}; -\frac{2}{3}; \cos^2(c+dx)\right)}{10d\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{10/3}}$$

[Out] 3/10*b^2*hypergeom([-5/3, 1/2], [-2/3], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(10/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2643}

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(-\frac{5}{3}, \frac{1}{2}; -\frac{2}{3}; \cos^2(c+dx)\right)}{10d\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{10/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*b^2*Hypergeometric2F1[-5/3, 1/2, -2/3, Cos[c + d*x]^2]*Sin[c + d*x])/(10*d*(b*Cos[c + d*x])^(10/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{4/3}} dx &= b^3 \int \frac{1}{(b \cos(c+dx))^{13/3}} dx \\ &= \frac{3b^2 {}_2F_1\left(-\frac{5}{3}, \frac{1}{2}; -\frac{2}{3}; \cos^2(c+dx)\right) \sin(c+dx)}{10d(b \cos(c+dx))^{10/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 58, normalized size = 1.00

$$\frac{3b^2 \sqrt{\sin^2(c+dx)} \csc(c+dx) {}_2F_1\left(-\frac{5}{3}, \frac{1}{2}; -\frac{2}{3}; \cos^2(c+dx)\right)}{10d(b \cos(c+dx))^{10/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*b^2*Csc[c + d*x]*Hypergeometric2F1[-5/3, 1/2, -2/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(10*d*(b*Cos[c + d*x])^(10/3))

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^{\frac{2}{3}} \sec(dx + c)^3}{b^2 \cos(dx + c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3/(b^2*cos(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(4/3), x)

maple [F] time = 0.11, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(dx + c)}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^3/(b*cos(d*x+c))^(4/3),x)

[Out] int(sec(d*x+c)^3/(b*cos(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^3 (b \cos(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(4/3)),x)

[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(4/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(4/3),x)

[Out] Integral(sec(c + d*x)**3/(b*cos(c + d*x))**(4/3), x)

3.244 $\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx$

Optimal. Leaf size=82

$$\frac{\sin(e + fx)(a \cos(e + fx))^{m+1}(b \cos(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(m + n + 1); \frac{1}{2}(m + n + 3); \cos^2(e + fx)\right)}{af(m + n + 1)\sqrt{\sin^2(e + fx)}}$$

[Out] $-(a*\cos(f*x+e))^{(1+m)}*(b*\cos(f*x+e))^n*\text{hypergeom}([1/2, 1/2+1/2*m+1/2*n], [3/2+1/2*m+1/2*n], \cos(f*x+e)^2)*\sin(f*x+e)/a/f/(1+m+n)/(\sin(f*x+e)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{\sin(e + fx)(a \cos(e + fx))^{m+1}(b \cos(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(m + n + 1); \frac{1}{2}(m + n + 3); \cos^2(e + fx)\right)}{af(m + n + 1)\sqrt{\sin^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[e + f*x])^m*(b*Cos[e + f*x])^n,x]

[Out] $-\left(\left(\left(a*\cos[e + f*x]\right)^{(1 + m)}*\left(b*\cos[e + f*x]\right)^n*\text{Hypergeometric2F1}\left[\frac{1}{2}, \left(1 + m + n\right)/2, \left(3 + m + n\right)/2, \cos[e + f*x]^2*\sin[e + f*x]\right]/\left(a*f*(1 + m + n)*\text{Sqrt}[\sin[e + f*x]^2]\right)\right)$

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m + n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (a \cos(e + fx))^m (b \cos(e + fx))^n dx &= \left((a \cos(e + fx))^{-n} (b \cos(e + fx))^n \right) \int (a \cos(e + fx))^{m+n} dx \\ &= \frac{(a \cos(e + fx))^{1+m} (b \cos(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + m + n); \frac{1}{2}(3 + m + n); \cos^2(e + fx)\right)}{af(1 + m + n)\sqrt{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 77, normalized size = 0.94

$$\frac{\sqrt{\sin^2(e + fx)} \cot(e + fx)(a \cos(e + fx))^m (b \cos(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(m + n + 1); \frac{1}{2}(m + n + 3); \cos^2(e + fx)\right)}{f(m + n + 1)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[e + f*x])^m*(b*cos[e + f*x])^n,x]

[Out] -(((a*cos[e + f*x])^m*(b*cos[e + f*x])^n*Cot[e + f*x]*Hypergeometric2F1[1/2, (1 + m + n)/2, (3 + m + n)/2, Cos[e + f*x]^2]*Sqrt[Sin[e + f*x]^2])/(f*(1 + m + n)))

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(a \cos (f x+e)\right)^m\left(b \cos (f x+e)\right)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*cos(f*x+e))^n,x, algorithm="fricas")

[Out] integral((a*cos(f*x + e))^m*(b*cos(f*x + e))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int\left(a \cos (f x+e)\right)^m\left(b \cos (f x+e)\right)^n d x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*cos(f*x+e))^n,x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m*(b*cos(f*x + e))^n, x)

maple [F] time = 0.89, size = 0, normalized size = 0.00

$$\int\left(a \cos (f x+e)\right)^m\left(b \cos (f x+e)\right)^n d x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m*(b*cos(f*x+e))^n,x)

[Out] int((a*cos(f*x+e))^m*(b*cos(f*x+e))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int\left(a \cos (f x+e)\right)^m\left(b \cos (f x+e)\right)^n d x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*cos(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m*(b*cos(f*x + e))^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int\left(a \cos (e+f x)\right)^m\left(b \cos (e+f x)\right)^n d x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(e + f*x))^m*(b*cos(e + f*x))^n,x)

[Out] int((a*cos(e + f*x))^m*(b*cos(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int\left(a \cos (e+f x)\right)^m\left(b \cos (e+f x)\right)^n d x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))**m*(b*cos(f*x+e))**n,x)

[Out] Integral((a*cos(e + f*x))**m*(b*cos(e + f*x))**n, x)

3.245 $\int \cos^2(c + dx)(b \cos(c + dx))^n dx$

Optimal. Leaf size=69

$$\frac{\sin(c + dx)(b \cos(c + dx))^{n+3} {}_2F_1\left(\frac{1}{2}, \frac{n+3}{2}; \frac{n+5}{2}; \cos^2(c + dx)\right)}{b^3 d(n+3) \sqrt{\sin^2(c + dx)}}$$

[Out] $-(b \cos(dx+c))^{(3+n)} \text{hypergeom}\left(\left[\frac{1}{2}, \frac{3}{2}+\frac{1}{2}n\right], \left[\frac{5}{2}+\frac{1}{2}n\right], \cos(dx+c)^2\right) \sin(dx+c) / b^3 d / (3+n) / (\sin(dx+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{\sin(c + dx)(b \cos(c + dx))^{n+3} {}_2F_1\left(\frac{1}{2}, \frac{n+3}{2}; \frac{n+5}{2}; \cos^2(c + dx)\right)}{b^3 d(n+3) \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2*(b*Cos[c + d*x])^n,x]

[Out] $-\left(\left(b \cos[c + d*x]\right)^{(3+n)} \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(3+n)}{2}, \frac{(5+n)}{2}, \cos[c + d*x]^2\right] \sin[c + d*x]\right) / \left(b^3 d (3+n) \sqrt{\sin[c + d*x]^2}\right)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2]) / (b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^2(c + dx)(b \cos(c + dx))^n dx &= \frac{\int (b \cos(c + dx))^{2+n} dx}{b^2} \\ &= \frac{(b \cos(c + dx))^{3+n} {}_2F_1\left(\frac{1}{2}, \frac{3+n}{2}; \frac{5+n}{2}; \cos^2(c + dx)\right) \sin(c + dx)}{b^3 d(3+n) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 72, normalized size = 1.04

$$\frac{\sqrt{\sin^2(c + dx)} \cos^2(c + dx) \cot(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n+3}{2}; \frac{n+5}{2}; \cos^2(c + dx)\right)}{d(n+3)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^n,x]

[Out] $-\left(\cos[c + d*x]^2 (b \cos[c + d*x])^n \cot[c + d*x] \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(3+n)}{2}, \frac{(5+n)}{2}, \cos[c + d*x]^2\right] \sqrt{\sin[c + d*x]^2}\right) / (d(3+n))$

fricas [F] time = 1.01, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^n \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^2, x)

maple [F] time = 1.33, size = 0, normalized size = 0.00

$$\int (\cos^2(dx + c)) (b \cos(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2*(b*cos(d*x+c))^n,x)

[Out] int(cos(d*x+c)^2*(b*cos(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^2 (b \cos(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^n,x)

[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^n \cos^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n*cos(c + d*x)**2, x)

3.246 $\int \cos(c + dx)(b \cos(c + dx))^n dx$

Optimal. Leaf size=69

$$\frac{\sin(c + dx)(b \cos(c + dx))^{n+2} {}_2F_1\left(\frac{1}{2}, \frac{n+2}{2}; \frac{n+4}{2}; \cos^2(c + dx)\right)}{b^2 d(n+2) \sqrt{\sin^2(c + dx)}}$$

[Out] $-(b \cos(dx+c))^{(2+n)} \text{hypergeom}\left(\left[\frac{1}{2}, 1+1/2*n\right], [2+1/2*n], \cos(dx+c)^2\right) \sin(dx+c)/b^2/d/(2+n)/(\sin(dx+c)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {16, 2643}

$$\frac{\sin(c + dx)(b \cos(c + dx))^{n+2} {}_2F_1\left(\frac{1}{2}, \frac{n+2}{2}; \frac{n+4}{2}; \cos^2(c + dx)\right)}{b^2 d(n+2) \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]*(b*Cos[c + d*x])^n,x]

[Out] $-(((b \cos[c + d*x])^{(2 + n)} \text{Hypergeometric2F1}[1/2, (2 + n)/2, (4 + n)/2, \cos[c + d*x]^2] \sin[c + d*x]) / (b^2 * d * (2 + n) * \text{Sqrt}[\sin[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2]) / (b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos(c + dx)(b \cos(c + dx))^n dx &= \frac{\int (b \cos(c + dx))^{1+n} dx}{b} \\ &= \frac{(b \cos(c + dx))^{2+n} {}_2F_1\left(\frac{1}{2}, \frac{2+n}{2}; \frac{4+n}{2}; \cos^2(c + dx)\right) \sin(c + dx)}{b^2 d(2+n) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 70, normalized size = 1.01

$$\frac{\sqrt{\sin^2(c + dx)} \cos(c + dx) \cot(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n+2}{2}; \frac{n+4}{2}; \cos^2(c + dx)\right)}{d(n+2)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^n,x]

[Out] $-((\cos[c + d*x] * (b \cos[c + d*x])^n \cot[c + d*x] \text{Hypergeometric2F1}[1/2, (2 + n)/2, (4 + n)/2, \cos[c + d*x]^2] \text{Sqrt}[\sin[c + d*x]^2]) / (d * (2 + n)))$

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}((b \cos(dx + c))^n \cos(dx + c), x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c), x)

maple [F] time = 1.15, size = 0, normalized size = 0.00

$$\int \cos(dx + c) (b \cos(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)*(b*cos(d*x+c))^n,x)

[Out] int(cos(d*x+c)*(b*cos(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx) (b \cos(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)*(b*cos(c + d*x))^n,x)

[Out] int(cos(c + d*x)*(b*cos(c + d*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^n \cos(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n*cos(c + d*x), x)

3.247 $\int (b \cos(c + dx))^n dx$

Optimal. Leaf size=69

$$\frac{\sin(c + dx)(b \cos(c + dx))^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(c + dx)\right)}{bd(n+1)\sqrt{\sin^2(c + dx)}}$$

[Out] $-(b*\cos(d*x+c))^{(1+n)}*\text{hypergeom}([1/2, 1/2+1/2*n], [3/2+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(1+n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2643}

$$\frac{\sin(c + dx)(b \cos(c + dx))^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(c + dx)\right)}{bd(n+1)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^n, x]

[Out] $-\left(\frac{(b*\cos[c + d*x])^{(1+n)}*\text{Hypergeometric2F1}[1/2, (1+n)/2, (3+n)/2, \cos[c + d*x]^2]*\sin[c + d*x]}{b*d*(1+n)*\text{Sqrt}[\sin[c + d*x]^2]}\right)$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (b \cos(c + dx))^n dx = -\frac{(b \cos(c + dx))^{1+n} {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(c + dx)\right) \sin(c + dx)}{bd(1+n)\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.04, size = 64, normalized size = 0.93

$$\frac{\sqrt{\sin^2(c + dx)} \cot(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(c + dx)\right)}{d(n+1)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^n, x]

[Out] $-\left(\frac{(b*\cos[c + d*x])^n*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[1/2, (1+n)/2, (3+n)/2, \cos[c + d*x]^2]*\text{Sqrt}[\sin[c + d*x]^2]}{d*(1+n)}\right)$

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n, x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n, x)

maple [F] time = 0.23, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n,x)

[Out] int((b*cos(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (b \cos(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n,x)

[Out] int((b*cos(c + d*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n, x)

3.248 $\int (b \cos(c + dx))^n \sec(c + dx) dx$

Optimal. Leaf size=60

$$\frac{\sin(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n}{2}; \frac{n+2}{2}; \cos^2(c + dx)\right)}{dn\sqrt{\sin^2(c + dx)}}$$

[Out] $-(b*\cos(d*x+c))^n*\text{hypergeom}([1/2, 1/2*n], [1+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/n/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {16, 2643}

$$\frac{\sin(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n}{2}; \frac{n+2}{2}; \cos^2(c + dx)\right)}{dn\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^n*Sec[c + d*x], x]

[Out] $-\left(\frac{(b*\cos[c + d*x])^n*\text{Hypergeometric2F1}[1/2, n/2, (2 + n)/2, \cos[c + d*x]^2]*\sin[c + d*x]}{(d*n*\text{Sqrt}[\sin[c + d*x]^2])}\right)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^n \sec(c + dx) dx &= b \int (b \cos(c + dx))^{-1+n} dx \\ &= \frac{(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n}{2}; \frac{2+n}{2}; \cos^2(c + dx)\right) \sin(c + dx)}{dn\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 63, normalized size = 1.05

$$\frac{b\sqrt{\sin^2(c + dx)} \cot(c + dx)(b \cos(c + dx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{n}{2}; \frac{n+2}{2}; \cos^2(c + dx)\right)}{dn}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Cos[c + d*x])^n*Sec[c + d*x], x]

[Out] $-\left(\frac{(b*(b*\cos[c + d*x])^{(-1 + n)}*\cot[c + d*x]*\text{Hypergeometric2F1}[1/2, n/2, (2 + n)/2, \cos[c + d*x]^2]*\text{Sqrt}[\sin[c + d*x]^2])}{(d*n)}\right)$

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}((b \cos(dx + c))^n \sec(dx + c), x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c), x)

maple [F] time = 0.48, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n*sec(d*x+c),x)

[Out] int((b*cos(d*x+c))^n*sec(d*x+c),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \cos(c + dx))^n}{\cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x),x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^n \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n*sec(d*x+c),x)

[Out] Integral((b*cos(c + d*x))**n*sec(c + d*x), x)

3.249 $\int (b \cos(c + dx))^n \sec^2(c + dx) dx$

Optimal. Leaf size=68

$$\frac{b \sin(c + dx)(b \cos(c + dx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{n-1}{2}; \frac{n+1}{2}; \cos^2(c + dx)\right)}{d(1-n)\sqrt{\sin^2(c + dx)}}$$

[Out] b*(b*cos(d*x+c))⁽⁻¹⁺ⁿ⁾*hypergeom([1/2, -1/2+1/2*n], [1/2+1/2*n], cos(d*x+c)²)*sin(d*x+c)/d/(1-n)/(sin(d*x+c)²)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{b \sin(c + dx)(b \cos(c + dx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{n-1}{2}; \frac{n+1}{2}; \cos^2(c + dx)\right)}{d(1-n)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])ⁿ*Sec[c + d*x]², x]

[Out] (b*(b*cos[c + d*x])^(-1 + n)*Hypergeometric2F1[1/2, (-1 + n)/2, (1 + n)/2, Cos[c + d*x]²*Sin[c + d*x])/(d*(1 - n)*Sqrt[Sin[c + d*x]²])

Rule 16

Int[(u_.)*(v_)^{(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]}

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] :> Simp[(Cos[c + d*x]*(b*sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]²]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]²]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^n \sec^2(c + dx) dx &= b^2 \int (b \cos(c + dx))^{-2+n} dx \\ &= \frac{b(b \cos(c + dx))^{-1+n} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(-1 + n); \frac{n+1}{2}; \cos^2(c + dx)\right) \sin(c + dx)}{d(1-n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 67, normalized size = 0.99

$$\frac{b\sqrt{\sin^2(c + dx)} \csc(c + dx)(b \cos(c + dx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{n-1}{2}; \frac{n+1}{2}; \cos^2(c + dx)\right)}{d(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])ⁿ*Sec[c + d*x]², x]

[Out] -((b*(b*cos[c + d*x])^(-1 + n)*Csc[c + d*x]*Hypergeometric2F1[1/2, (-1 + n)/2, (1 + n)/2, Cos[c + d*x]²*Sqrt[Sin[c + d*x]²])/(d*(-1 + n)))

fricas [F] time = 0.89, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^n \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^2, x)

maple [F] time = 0.40, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n (\sec^2(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n*sec(d*x+c)^2,x)

[Out] int((b*cos(d*x+c))^n*sec(d*x+c)^2,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x)^2,x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^n \sec^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n*sec(d*x+c)**2,x)

[Out] Integral((b*cos(c + d*x))**n*sec(c + d*x)**2, x)

3.250 $\int (b \cos(c + dx))^n \sec^3(c + dx) dx$

Optimal. Leaf size=68

$$\frac{b^2 \sin(c + dx)(b \cos(c + dx))^{n-2} {}_2F_1\left(\frac{1}{2}, \frac{n-2}{2}; \frac{n}{2}; \cos^2(c + dx)\right)}{d(2-n)\sqrt{\sin^2(c + dx)}}$$

[Out] $b^2*(b*\cos(d*x+c))^{(-2+n)}*\text{hypergeom}([1/2, -1+1/2*n], [1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(2-n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{b^2 \sin(c + dx)(b \cos(c + dx))^{n-2} {}_2F_1\left(\frac{1}{2}, \frac{n-2}{2}; \frac{n}{2}; \cos^2(c + dx)\right)}{d(2-n)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^n*Sec[c + d*x]^3,x]

[Out] $(b^2*(b*\cos[c + d*x])^{(-2 + n)}*\text{Hypergeometric2F1}[1/2, (-2 + n)/2, n/2, \cos[c + d*x]^2]*\sin[c + d*x])/(d*(2 - n)*\text{Sqrt}[\sin[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^n \sec^3(c + dx) dx &= b^3 \int (b \cos(c + dx))^{-3+n} dx \\ &= \frac{b^2 (b \cos(c + dx))^{-2+n} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(-2 + n); \frac{n}{2}; \cos^2(c + dx)\right) \sin(c + dx)}{d(2-n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 70, normalized size = 1.03

$$\frac{\sqrt{\sin^2(c + dx)} \csc(c + dx) \sec^2(c + dx) (b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n-2}{2}; \frac{n}{2}; \cos^2(c + dx)\right)}{d(n-2)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^n*Sec[c + d*x]^3,x]

[Out] $-(((b*\cos[c + d*x])^n*\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[1/2, (-2 + n)/2, n/2, \cos[c + d*x]^2]*\text{Sec}[c + d*x]^2*\text{Sqrt}[\sin[c + d*x]^2]))/(d*(-2 + n))$

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral} \left((b \cos(dx + c))^n \sec(dx + c)^3, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^3, x)

maple [F] time = 0.57, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n (\sec^3(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n*sec(d*x+c)^3,x)

[Out] int((b*cos(d*x+c))^n*sec(d*x+c)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^n \sec^3(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n*sec(d*x+c)**3,x)

[Out] Integral((b*cos(c + d*x))**n*sec(c + d*x)**3, x)

3.251 $\int (b \cos(c + dx))^n \sec^4(c + dx) dx$

Optimal. Leaf size=70

$$\frac{b^3 \sin(c + dx)(b \cos(c + dx))^{n-3} {}_2F_1\left(\frac{1}{2}, \frac{n-3}{2}; \frac{n-1}{2}; \cos^2(c + dx)\right)}{d(3-n)\sqrt{\sin^2(c + dx)}}$$

[Out] b^3*(b*cos(d*x+c))^(n-3)*hypergeom([1/2, -3/2+1/2*n], [-1/2+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(3-n)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2643}

$$\frac{b^3 \sin(c + dx)(b \cos(c + dx))^{n-3} {}_2F_1\left(\frac{1}{2}, \frac{n-3}{2}; \frac{n-1}{2}; \cos^2(c + dx)\right)}{d(3-n)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^n*Sec[c + d*x]^4,x]

[Out] (b^3*(b*cos[c + d*x])^(n-3)*Hypergeometric2F1[1/2, (-3 + n)/2, (-1 + n)/2, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(3 - n)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (b \cos(c + dx))^n \sec^4(c + dx) dx &= b^4 \int (b \cos(c + dx))^{-4+n} dx \\ &= \frac{b^3 (b \cos(c + dx))^{-3+n} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(-3 + n); \frac{1}{2}(-1 + n); \cos^2(c + dx)\right) \sin(c + dx)}{d(3-n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 72, normalized size = 1.03

$$\frac{\sqrt{\sin^2(c + dx)} \csc(c + dx) \sec^3(c + dx) (b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n-3}{2}; \frac{n-1}{2}; \cos^2(c + dx)\right)}{d(n-3)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^n*Sec[c + d*x]^4,x]

[Out] -(((b*cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-3 + n)/2, (-1 + n)/2, Cos[c + d*x]^2]*Sec[c + d*x]^3*Sqrt[Sin[c + d*x]^2])/(d*(-3 + n)))

fricas [F] time = 1.04, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^n \sec(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^4,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*sec(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^4,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^4, x)

maple [F] time = 0.34, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n (\sec^4(dx + c)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n*sec(d*x+c)^4,x)

[Out] int((b*cos(d*x+c))^n*sec(d*x+c)^4,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^4,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x)^4,x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^4, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n*sec(d*x+c)**4,x)

[Out] Timed out

3.252 $\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx$

Optimal. Leaf size=80

$$\frac{2 \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n + 7); \frac{1}{4}(2n + 11); \cos^2(c + dx)\right)}{d(2n + 7)\sqrt{\sin^2(c + dx)}}$$

[Out] $-2*\cos(d*x+c)^{(7/2)}*(b*\cos(d*x+c))^n*\text{hypergeom}([1/2, 7/4+1/2*n], [11/4+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(7+2*n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{2 \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n + 7); \frac{1}{4}(2n + 11); \cos^2(c + dx)\right)}{d(2n + 7)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^n,x]

[Out] $(-2*\text{Cos}[c + d*x]^{(7/2)}*(b*\text{Cos}[c + d*x])^n*\text{Hypergeometric2F1}[1/2, (7 + 2*n)/4, (11 + 2*n)/4, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*(7 + 2*n)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{\frac{5}{2}+n}(c + dx) dx \\ &= -\frac{2 \cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(7 + 2n); \frac{1}{4}(11 + 2n); \cos^2(c + dx)\right)}{d(7 + 2n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.13, size = 80, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c + dx)} \cos^{\frac{7}{2}}(c + dx) \csc(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n + \frac{7}{2}\right); \frac{1}{2}\left(n + \frac{11}{2}\right); \cos^2(c + dx)\right)}{d\left(n + \frac{7}{2}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^n,x]

[Out] -((Cos[c + d*x]^(7/2)*(b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (7/2 + n)/2, (11/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(7/2 + n)))

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^n \cos(dx + c)^{\frac{5}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*cos(d*x + c)^(5/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \cos(dx + c)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^(5/2), x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int \left(\cos^{\frac{5}{2}}(dx + c)\right) (b \cos(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^n,x)

[Out] int(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \cos(dx + c)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^{5/2} (b \cos(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^n,x)

[Out] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^n, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(5/2)*(b*cos(d*x+c))**n,x)

[Out] Timed out

3.253 $\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx$

Optimal. Leaf size=80

$$\frac{2 \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n + 5); \frac{1}{4}(2n + 9); \cos^2(c + dx)\right)}{d(2n + 5)\sqrt{\sin^2(c + dx)}}$$

[Out] $-2*\cos(d*x+c)^{(5/2)}*(b*\cos(d*x+c))^n*\text{hypergeom}([1/2, 5/4+1/2*n], [9/4+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(5+2*n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{2 \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n + 5); \frac{1}{4}(2n + 9); \cos^2(c + dx)\right)}{d(2n + 5)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^n,x]

[Out] $(-2*\text{Cos}[c + d*x]^{(5/2)}*(b*\text{Cos}[c + d*x])^n*\text{Hypergeometric2F1}[1/2, (5 + 2*n)/4, (9 + 2*n)/4, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*(5 + 2*n)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{\frac{3}{2}+n}(c + dx) dx \\ &= -\frac{2 \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(5 + 2n); \frac{1}{4}(9 + 2n); \cos^2(c + dx)\right)}{d(5 + 2n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.11, size = 80, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c + dx)} \cos^{\frac{5}{2}}(c + dx) \csc(c + dx)(b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n + \frac{5}{2}\right); \frac{1}{2}\left(n + \frac{9}{2}\right); \cos^2(c + dx)\right)}{d\left(n + \frac{5}{2}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^n,x]

[Out] -((Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (5/2 + n)/2, (9/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(5/2 + n)))

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left((b \cos(dx + c))^n \cos(dx + c)^{\frac{3}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*cos(d*x + c)^(3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \cos(dx + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^(3/2), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int \left(\cos^{\frac{3}{2}}(dx + c)\right) (b \cos(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^n,x)

[Out] int(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \cos(dx + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^{3/2} (b \cos(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^n,x)

[Out] int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^n, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(3/2)*(b*cos(d*x+c))**n,x)

[Out] Timed out

3.254 $\int \sqrt{\cos(c + dx)} (b \cos(c + dx))^n dx$

Optimal. Leaf size=80

$$\frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n + 3); \frac{1}{4}(2n + 7); \cos^2(c + dx)\right)}{d(2n + 3)\sqrt{\sin^2(c + dx)}}$$

[Out] $-2*\cos(d*x+c)^{(3/2)}*(b*\cos(d*x+c))^n*\text{hypergeom}([1/2, 3/4+1/2*n], [7/4+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(3+2*n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n + 3); \frac{1}{4}(2n + 7); \cos^2(c + dx)\right)}{d(2n + 3)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^n,x]

[Out] $(-2*\text{Cos}[c + d*x]^{(3/2)}*(b*\text{Cos}[c + d*x])^n*\text{Hypergeometric2F1}[1/2, (3 + 2*n)/4, (7 + 2*n)/4, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*(3 + 2*n)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sqrt{\cos(c + dx)} (b \cos(c + dx))^n dx &= (\cos^{-n}(c + dx) (b \cos(c + dx))^n) \int \cos^{\frac{1}{2}+n}(c + dx) dx \\ &= -\frac{2 \cos^{\frac{3}{2}}(c + dx) (b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(3 + 2n); \frac{1}{4}(7 + 2n); \cos^2(c + dx)\right)}{d(3 + 2n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 80, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c + dx)} \cos^{\frac{3}{2}}(c + dx) \csc(c + dx) (b \cos(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n + \frac{3}{2}\right); \frac{1}{2}\left(n + \frac{7}{2}\right); \cos^2(c + dx)\right)}{d\left(n + \frac{3}{2}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^n,x]

[Out] -((Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (3/2 + n)/2, (7/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(3/2 + n)))

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}((b \cos(dx + c))^n \sqrt{\cos(dx + c)}, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*sqrt(cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sqrt{\cos(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*sqrt(cos(d*x + c)), x)

maple [F] time = 0.12, size = 0, normalized size = 0.00

$$\int (\sqrt{\cos(dx + c)} (b \cos(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^n,x)

[Out] int(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(dx + c))^n \sqrt{\cos(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*sqrt(cos(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{\cos(c + dx)} (b \cos(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^n,x)

[Out] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \cos(c + dx))^n \sqrt{\cos(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**(1/2)*(b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n*sqrt(cos(c + d*x)), x)

$$3.255 \quad \int \frac{(b \cos(c+dx))^n}{\sqrt{\cos(c+dx)}} dx$$

Optimal. Leaf size=80

$$\frac{2 \sin(c+dx) \sqrt{\cos(c+dx)} (b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n+1); \frac{1}{4}(2n+5); \cos^2(c+dx)\right)}{d(2n+1) \sqrt{\sin^2(c+dx)}}$$

[Out] $-2*(b*\cos(d*x+c))^n*\text{hypergeom}([1/2, 1/4+1/2*n], [5/4+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/d/(1+2*n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{2 \sin(c+dx) \sqrt{\cos(c+dx)} (b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n+1); \frac{1}{4}(2n+5); \cos^2(c+dx)\right)}{d(2n+1) \sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^n/Sqrt[Cos[c + d*x]], x]

[Out] $(-2*\text{Sqrt}[\text{Cos}[c + d*x]]*(b*\text{Cos}[c + d*x])^n*\text{Hypergeometric2F1}[1/2, (1 + 2*n)/4, (5 + 2*n)/4, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(d*(1 + 2*n)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{(b \cos(c+dx))^n}{\sqrt{\cos(c+dx)}} dx = (\cos^{-n}(c+dx)(b \cos(c+dx))^n) \int \cos^{-\frac{1}{2}+n}(c+dx) dx$$

$$= \frac{2\sqrt{\cos(c+dx)} (b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(1+2n); \frac{1}{4}(5+2n); \cos^2(c+dx)\right) \sin(c+dx)}{d(1+2n) \sqrt{\sin^2(c+dx)}}$$

Mathematica [A] time = 0.05, size = 80, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c+dx)} \sqrt{\cos(c+dx)} \csc(c+dx) (b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n + \frac{1}{2}\right); \frac{1}{2}\left(n + \frac{5}{2}\right); \cos^2(c+dx)\right)}{d\left(n + \frac{1}{2}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^n/Sqrt[Cos[c + d*x]],x]

[Out] -((Sqrt[Cos[c + d*x]]*(b*cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (1/2 + n)/2, (5/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(1/2 + n)))

fricas [F] time = 0.79, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^n}{\sqrt{\cos(dx + c)}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(1/2),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/sqrt(cos(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\sqrt{\cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/sqrt(cos(d*x + c)), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\sqrt{\cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n/cos(d*x+c)^(1/2),x)

[Out] int((b*cos(d*x+c))^n/cos(d*x+c)^(1/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\sqrt{\cos(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/sqrt(cos(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(1/2),x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(1/2), x)
```

```
[Out] Integral((b*cos(c + d*x))**n/sqrt(cos(c + d*x)), x)
```

$$3.256 \quad \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{3}{2}}(c+dx)} dx$$

Optimal. Leaf size=80

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n-1); \frac{1}{4}(2n+3); \cos^2(c+dx)\right)}{d(1-2n)\sqrt{\sin^2(c+dx)}\sqrt{\cos(c+dx)}}$$

[Out] 2*(b*cos(d*x+c))^n*hypergeom([1/2, -1/4+1/2*n], [3/4+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(1-2*n)/cos(d*x+c)^(1/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n-1); \frac{1}{4}(2n+3); \cos^2(c+dx)\right)}{d(1-2n)\sqrt{\sin^2(c+dx)}\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^n/Cos[c + d*x]^(3/2), x]

[Out] (2*(b*cos[c + d*x])^n*Hypergeometric2F1[1/2, (-1 + 2*n)/4, (3 + 2*n)/4, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(1 - 2*n)*Sqrt[Cos[c + d*x]]*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] :> Simp[(Cos[c + d*x]*(b*sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{3}{2}}(c+dx)} dx &= (\cos^{-n}(c+dx)(b \cos(c+dx))^n) \int \cos^{-\frac{3}{2}+n}(c+dx) dx \\ &= \frac{2(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-1+2n); \frac{1}{4}(3+2n); \cos^2(c+dx)\right) \sin(c+dx)}{d(1-2n)\sqrt{\cos(c+dx)}\sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 80, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c+dx)} \csc(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n - \frac{1}{2}\right); \frac{1}{2}\left(n + \frac{3}{2}\right); \cos^2(c+dx)\right)}{d\left(n - \frac{1}{2}\right)\sqrt{\cos(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^n/Cos[c + d*x]^(3/2), x]

[Out] -(((b*cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-1/2 + n)/2, (3/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-1/2 + n)*Sqrt[Cos[c + d*x]]))

fricas [F] time = 0.44, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{3}{2}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(3/2), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/cos(d*x + c)^(3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(3/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(3/2), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n/cos(d*x+c)^(3/2), x)

[Out] int((b*cos(d*x+c))^n/cos(d*x+c)^(3/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(3/2), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(3/2), x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{3}{2}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(3/2), x)

[Out] Integral((b*cos(c + d*x))**n/cos(c + d*x)**(3/2), x)

$$3.257 \quad \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{5}{2}}(c+dx)} dx$$

Optimal. Leaf size=80

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n-3); \frac{1}{4}(2n+1); \cos^2(c+dx)\right)}{d(3-2n)\sqrt{\sin^2(c+dx)} \cos^{\frac{3}{2}}(c+dx)}$$

[Out] 2*(b*cos(d*x+c))^n*hypergeom([1/2, -3/4+1/2*n], [1/4+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(3-2*n)/cos(d*x+c)^(3/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n-3); \frac{1}{4}(2n+1); \cos^2(c+dx)\right)}{d(3-2n)\sqrt{\sin^2(c+dx)} \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^n/Cos[c + d*x]^(5/2), x]

[Out] (2*(b*cos[c + d*x])^n*Hypergeometric2F1[1/2, (-3 + 2*n)/4, (1 + 2*n)/4, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(3 - 2*n)*Cos[c + d*x]^(3/2)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] :> Simp[(Cos[c + d*x]*(b*sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{5}{2}}(c+dx)} dx &= (\cos^{-n}(c+dx)(b \cos(c+dx))^n) \int \cos^{-\frac{5}{2}+n}(c+dx) dx \\ &= \frac{2(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-3+2n); \frac{1}{4}(1+2n); \cos^2(c+dx)\right) \sin(c+dx)}{d(3-2n) \cos^{\frac{3}{2}}(c+dx) \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 80, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c+dx)} \csc(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n-\frac{3}{2}\right); \frac{1}{2}\left(n+\frac{1}{2}\right); \cos^2(c+dx)\right)}{d\left(n-\frac{3}{2}\right) \cos^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^n/cos[c + d*x]^(5/2), x]

[Out] -(((b*cos[c + d*x])^n*csc[c + d*x]*Hypergeometric2F1[1/2, (-3/2 + n)/2, (1/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-3/2 + n)*Cos[c + d*x]^(3/2)))

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{5}{2}}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(5/2), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/cos(d*x + c)^(5/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(5/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(5/2), x)

maple [F] time = 0.10, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n/cos(d*x+c)^(5/2), x)

[Out] int((b*cos(d*x+c))^n/cos(d*x+c)^(5/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(5/2), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(5/2), x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{5}{2}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(5/2), x)

[Out] Integral((b*cos(c + d*x))**n/cos(c + d*x)**(5/2), x)

$$3.258 \quad \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{7}{2}}(c+dx)} dx$$

Optimal. Leaf size=80

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n-5); \frac{1}{4}(2n-1); \cos^2(c+dx)\right)}{d(5-2n)\sqrt{\sin^2(c+dx)} \cos^{\frac{5}{2}}(c+dx)}$$

[Out] 2*(b*cos(d*x+c))^n*hypergeom([1/2, -5/4+1/2*n], [-1/4+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(5-2*n)/cos(d*x+c)^(5/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n-5); \frac{1}{4}(2n-1); \cos^2(c+dx)\right)}{d(5-2n)\sqrt{\sin^2(c+dx)} \cos^{\frac{5}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*cos[c + d*x])^n/Cos[c + d*x]^(7/2), x]

[Out] (2*(b*cos[c + d*x])^n*Hypergeometric2F1[1/2, (-5 + 2*n)/4, (-1 + 2*n)/4, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(5 - 2*n)*Cos[c + d*x]^(5/2)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{7}{2}}(c+dx)} dx &= (\cos^{-n}(c+dx)(b \cos(c+dx))^n) \int \cos^{-\frac{7}{2}+n}(c+dx) dx \\ &= \frac{2(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-5+2n); \frac{1}{4}(-1+2n); \cos^2(c+dx)\right) \sin(c+dx)}{d(5-2n) \cos^{\frac{5}{2}}(c+dx) \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 80, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c+dx)} \csc(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n - \frac{5}{2}\right); \frac{1}{2}\left(n - \frac{1}{2}\right); \cos^2(c+dx)\right)}{d\left(n - \frac{5}{2}\right) \cos^{\frac{5}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^n/Cos[c + d*x]^(7/2), x]

[Out] -(((b*cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-5/2 + n)/2, (-1/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-5/2 + n)*Cos[c + d*x]^(5/2)))

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{7}{2}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(7/2), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/cos(d*x + c)^(7/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(7/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(7/2), x)

maple [F] time = 0.09, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n/cos(d*x+c)^(7/2), x)

[Out] int((b*cos(d*x+c))^n/cos(d*x+c)^(7/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(7/2), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(7/2), x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(7/2),x)

[Out] Timed out

$$3.259 \quad \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{9}{2}}(c+dx)} dx$$

Optimal. Leaf size=80

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n-7); \frac{1}{4}(2n-3); \cos^2(c+dx)\right)}{d(7-2n)\sqrt{\sin^2(c+dx)} \cos^{\frac{7}{2}}(c+dx)}$$

[Out] 2*(b*cos(d*x+c))^n*hypergeom([1/2, -7/4+1/2*n], [-3/4+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(7-2*n)/cos(d*x+c)^(7/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2643}

$$\frac{2 \sin(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(2n-7); \frac{1}{4}(2n-3); \cos^2(c+dx)\right)}{d(7-2n)\sqrt{\sin^2(c+dx)} \cos^{\frac{7}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*Cos[c + d*x])^n/Cos[c + d*x]^(9/2), x]

[Out] (2*(b*Cos[c + d*x])^n*Hypergeometric2F1[1/2, (-7 + 2*n)/4, (-3 + 2*n)/4, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(7 - 2*n)*Cos[c + d*x]^(7/2)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] :> Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{9}{2}}(c+dx)} dx &= (\cos^{-n}(c+dx)(b \cos(c+dx))^n) \int \cos^{-\frac{9}{2}+n}(c+dx) dx \\ &= \frac{2(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-7+2n); \frac{1}{4}(-3+2n); \cos^2(c+dx)\right) \sin(c+dx)}{d(7-2n) \cos^{\frac{7}{2}}(c+dx) \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 80, normalized size = 1.00

$$\frac{\sqrt{\sin^2(c+dx)} \csc(c+dx)(b \cos(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n-\frac{7}{2}\right); \frac{1}{2}\left(n-\frac{3}{2}\right); \cos^2(c+dx)\right)}{d\left(n-\frac{7}{2}\right) \cos^{\frac{7}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*cos[c + d*x])^n/Cos[c + d*x]^(9/2), x]

[Out] -(((b*cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-7/2 + n)/2, (-3/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-7/2 + n)*Cos[c + d*x]^(7/2)))

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{9}{2}}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(9/2), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/cos(d*x + c)^(9/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(9/2), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(9/2), x)

maple [F] time = 0.08, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(d*x+c))^n/cos(d*x+c)^(9/2), x)

[Out] int((b*cos(d*x+c))^n/cos(d*x+c)^(9/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(9/2), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(9/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(9/2), x)

[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(9/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(9/2), x)

[Out] Timed out

3.260 $\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx$

Optimal. Leaf size=88

$$\frac{\sin(e + fx)(a \cos(e + fx))^{m+1}(b \sec(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(m - n + 1); \frac{1}{2}(m - n + 3); \cos^2(e + fx)\right)}{af(m - n + 1)\sqrt{\sin^2(e + fx)}}$$

[Out] $-(a*\cos(f*x+e))^{(1+m)}*\text{hypergeom}([1/2, 1/2+1/2*m-1/2*n], [3/2+1/2*m-1/2*n], \cos(f*x+e)^2)*(b*\sec(f*x+e))^n*\sin(f*x+e)/a/f/(1+m-n)/(\sin(f*x+e)^2)^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2588, 2643}

$$\frac{\sin(e + fx)(a \cos(e + fx))^{m+1}(b \sec(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(m - n + 1); \frac{1}{2}(m - n + 3); \cos^2(e + fx)\right)}{af(m - n + 1)\sqrt{\sin^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a*\text{Cos}[e + f*x])^m*(b*\text{Sec}[e + f*x])^n, x]$

[Out] $-\left(\left((a*\text{Cos}[e + f*x])^{(1 + m)}*\text{Hypergeometric2F1}[1/2, (1 + m - n)/2, (3 + m - n)/2, \text{Cos}[e + f*x]^2]*(b*\text{Sec}[e + f*x])^n*\text{Sin}[e + f*x]\right)/(a*f*(1 + m - n)*\text{Sqrt}[\text{Sin}[e + f*x]^2])\right)$

Rule 2588

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Dist}[(a*b)^{\text{IntPart}[n]}*(a*\text{Sin}[e + f*x])^{\text{FracPart}[n]}*(b*\text{Csc}[e + f*x])^{\text{FracPart}[n]}, \text{Int}[(a*\text{Sin}[e + f*x])^{(m - n)}, x], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{!IntegerQ}[m] \&\& \text{!IntegerQ}[n]$

Rule 2643

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned} \int (a \cos(e + fx))^m (b \sec(e + fx))^n dx &= \left((a \cos(e + fx))^n (b \sec(e + fx))^n \right) \int (a \cos(e + fx))^{m-n} dx \\ &= -\frac{(a \cos(e + fx))^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + m - n); \frac{1}{2}(3 + m - n); \cos^2(e + fx)\right) (b)}{af(1 + m - n)\sqrt{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [A] time = 10.91, size = 89, normalized size = 1.01

$$\frac{\sin(e + fx) \cos(e + fx)(a \cos(e + fx))^m (b \sec(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(m - n + 1); \frac{1}{2}(m - n + 3); \cos^2(e + fx)\right)}{f(m - n + 1)\sqrt{\sin^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[e + f*x])^m*(b*Sec[e + f*x])^n,x]

[Out] -((Cos[e + f*x]*(a*cos[e + f*x])^m*Hypergeometric2F1[1/2, (1 + m - n)/2, (3 + m - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^n*Sin[e + f*x])/(f*(1 + m - n)*Sqrt[Sin[e + f*x]^2]))

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(a \cos(fx + e)\right)^m \left(b \sec(fx + e)\right)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*sec(f*x+e))^n,x, algorithm="fricas")

[Out] integral((a*cos(f*x + e))^m*(b*sec(f*x + e))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \sec(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*sec(f*x+e))^n,x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m*(b*sec(f*x + e))^n, x)

maple [F] time = 0.84, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \sec(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m*(b*sec(f*x+e))^n,x)

[Out] int((a*cos(f*x+e))^m*(b*sec(f*x+e))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \sec(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*sec(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m*(b*sec(f*x + e))^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(e + fx))^m \left(\frac{b}{\cos(e + fx)}\right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(e + f*x))^m*(b/cos(e + f*x))^n,x)

[Out] int((a*cos(e + f*x))^m*(b/cos(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))**m*(b*sec(f*x+e))**n,x)

[Out] Integral((a*cos(e + f*x))**m*(b*sec(e + f*x))**n, x)

3.261 $\int \cos(a + bx) \sqrt{\csc(a + bx)} dx$

Optimal. Leaf size=15

$$\frac{2}{b\sqrt{\csc(a + bx)}}$$

[Out] 2/b/csc(b*x+a)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2621, 30}

$$\frac{2}{b\sqrt{\csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Sqrt[Csc[a + b*x]],x]

[Out] 2/(b*Sqrt[Csc[a + b*x]])

Rule 30

Int[(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \sqrt{\csc(a + bx)} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{3/2}} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{2}{b\sqrt{\csc(a + bx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 15, normalized size = 1.00

$$\frac{2}{b\sqrt{\csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*Sqrt[Csc[a + b*x]],x]

[Out] 2/(b*Sqrt[Csc[a + b*x]])

fricas [A] time = 0.64, size = 13, normalized size = 0.87

$$\frac{2\sqrt{\sin(bx + a)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] 2*sqrt(sin(b*x + a))/b

giac [A] time = 0.46, size = 13, normalized size = 0.87

$$\frac{2\sqrt{\sin(bx+a)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] 2*sqrt(sin(b*x + a))/b

maple [A] time = 0.04, size = 14, normalized size = 0.93

$$\frac{2}{b\sqrt{\csc(bx+a)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)*csc(b*x+a)^(1/2),x)

[Out] 2/b/csc(b*x+a)^(1/2)

maxima [A] time = 0.64, size = 13, normalized size = 0.87

$$\frac{2\sqrt{\sin(bx+a)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 2*sqrt(sin(b*x + a))/b

mupad [B] time = 0.30, size = 15, normalized size = 1.00

$$\frac{2}{b\sqrt{\frac{1}{\sin(a+bx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)*(1/sin(a + b*x))^(1/2),x)

[Out] 2/(b*(1/sin(a + b*x))^(1/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(a + bx)\sqrt{\csc(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*csc(b*x+a)**(1/2),x)

[Out] Integral(cos(a + b*x)*sqrt(csc(a + b*x)), x)

$$3.262 \quad \int \frac{\cos(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal. Leaf size=17

$$\frac{2}{3b \csc^{\frac{3}{2}}(a+bx)}$$

[Out] 2/3/b/csc(b*x+a)^(3/2)

Rubi [A] time = 0.02, antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2621, 30}

$$\frac{2}{3b \csc^{\frac{3}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]/Sqrt[Csc[a + b*x]],x]

[Out] 2/(3*b*Csc[a + b*x]^(3/2))

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2621

Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\cos(a+bx)}{\sqrt{\csc(a+bx)}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{5/2}} dx, x, \csc(a+bx)\right)}{b} \\ &= \frac{2}{3b \csc^{\frac{3}{2}}(a+bx)} \end{aligned}$$

Mathematica [A] time = 0.02, size = 17, normalized size = 1.00

$$\frac{2}{3b \csc^{\frac{3}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]/Sqrt[Csc[a + b*x]],x]

[Out] 2/(3*b*Csc[a + b*x]^(3/2))

fricas [A] time = 0.54, size = 23, normalized size = 1.35

$$\frac{2(\cos(bx+a)^2 - 1)}{3b\sqrt{\sin(bx+a)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] -2/3*(cos(b*x + a)^2 - 1)/(b*sqrt(sin(b*x + a)))

giac [A] time = 0.63, size = 13, normalized size = 0.76

$$\frac{2 \sin (bx + a)^{\frac{3}{2}}}{3 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] 2/3*sin(b*x + a)^(3/2)/b

maple [A] time = 0.02, size = 14, normalized size = 0.82

$$\frac{2}{3 b \csc (bx + a)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)/csc(b*x+a)^(1/2),x)

[Out] 2/3/b/csc(b*x+a)^(3/2)

maxima [A] time = 0.84, size = 13, normalized size = 0.76

$$\frac{2 \sin (bx + a)^{\frac{3}{2}}}{3 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 2/3*sin(b*x + a)^(3/2)/b

mupad [B] time = 0.21, size = 15, normalized size = 0.88

$$\frac{2}{3 b \left(\frac{1}{\sin (a + b x)} \right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)/(1/sin(a + b*x))^(1/2),x)

[Out] 2/(3*b*(1/sin(a + b*x))^(3/2))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos (a + b x)}{\sqrt{\csc (a + b x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/csc(b*x+a)**(1/2),x)

[Out] Integral(cos(a + b*x)/sqrt(csc(a + b*x)), x)

3.263 $\int \cos^2(a + bx)\sqrt{\csc(a + bx)} dx$

Optimal. Leaf size=67

$$\frac{2 \cos(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{4\sqrt{\sin(a + bx)}\sqrt{\csc(a + bx)}F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right)\middle|2\right)}{3b}$$

[Out] 2/3*cos(b*x+a)/b/csc(b*x+a)^(1/2)-4/3*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticF(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] time = 0.05, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2628, 3771, 2641}

$$\frac{2 \cos(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{4\sqrt{\sin(a + bx)}\sqrt{\csc(a + bx)}F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right)\middle|2\right)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Sqrt[Csc[a + b*x]],x]

[Out] (2*Cos[a + b*x])/(3*b*Sqrt[Csc[a + b*x]]) + (4*Sqrt[Csc[a + b*x]]*EllipticF[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/(3*b)

Rule 2628

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \cos^2(a + bx)\sqrt{\csc(a + bx)} dx &= \frac{2 \cos(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{2}{3} \int \sqrt{\csc(a + bx)} dx \\ &= \frac{2 \cos(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{1}{3} \left(2\sqrt{\csc(a + bx)}\sqrt{\sin(a + bx)}\right) \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{2 \cos(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{4\sqrt{\csc(a + bx)}F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right)\middle|2\right)\sqrt{\sin(a + bx)}}{3b} \end{aligned}$$

Mathematica [A] time = 0.10, size = 53, normalized size = 0.79

$$\frac{\sqrt{\csc(a+bx)} \left(\sin(2(a+bx)) - 4\sqrt{\sin(a+bx)} F\left(\frac{1}{4}(-2a-2bx+\pi) \middle| 2\right) \right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^2*Sqrt[Csc[a + b*x]], x]

[Out] (Sqrt[Csc[a + b*x]]*(-4*EllipticF[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] + Sin[2*(a + b*x)]))/(3*b)

fricas [F] time = 0.49, size = 0, normalized size = 0.00

$$\text{integral}(\cos(bx+a)^2 \sqrt{\csc(bx+a)}, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*csc(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^2*sqrt(csc(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx+a)^2 \sqrt{\csc(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*csc(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(cos(b*x + a)^2*sqrt(csc(b*x + a)), x)

maple [A] time = 0.16, size = 88, normalized size = 1.31

$$\frac{2\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \text{EllipticF}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) + \frac{2(\cos^2(bx+a))\sin(bx+a)}{3}}{\cos(bx+a) \sqrt{\sin(bx+a)} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2*csc(b*x+a)^(1/2), x)

[Out] (2/3*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))+2/3*cos(b*x+a)^2*sin(b*x+a))/cos(b*x+a)/sin(b*x+a)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx+a)^2 \sqrt{\csc(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*csc(b*x+a)^(1/2), x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^2*sqrt(csc(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(a+bx)^2 \sqrt{\frac{1}{\sin(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)^2*(1/sin(a + b*x))^(1/2), x)
```

```
[Out] int(cos(a + b*x)^2*(1/sin(a + b*x))^(1/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)**2*csc(b*x+a)**(1/2), x)
```

```
[Out] Integral(cos(a + b*x)**2*sqrt(csc(a + b*x)), x)
```

$$3.264 \quad \int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal. Leaf size=67

$$\frac{2 \cos(a+bx)}{5b \csc^{\frac{3}{2}}(a+bx)} + \frac{4\sqrt{\sin(a+bx)}\sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a+bx-\frac{\pi}{2}\right)\middle|2\right)}{5b}$$

[Out] 2/5*cos(b*x+a)/b/csc(b*x+a)^(3/2)-4/5*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticE(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] time = 0.05, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2628, 3771, 2639}

$$\frac{2 \cos(a+bx)}{5b \csc^{\frac{3}{2}}(a+bx)} + \frac{4\sqrt{\sin(a+bx)}\sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a+bx-\frac{\pi}{2}\right)\middle|2\right)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2/Sqrt[Csc[a + b*x]],x]

[Out] (2*cos[a + b*x])/(5*b*Csc[a + b*x]^(3/2)) + (4*Sqrt[Csc[a + b*x]]*EllipticE[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/(5*b)

Rule 2628

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx &= \frac{2 \cos(a+bx)}{5b \csc^{\frac{3}{2}}(a+bx)} + \frac{2}{5} \int \frac{1}{\sqrt{\csc(a+bx)}} dx \\ &= \frac{2 \cos(a+bx)}{5b \csc^{\frac{3}{2}}(a+bx)} + \frac{1}{5} \left(2\sqrt{\csc(a+bx)}\sqrt{\sin(a+bx)}\right) \int \sqrt{\sin(a+bx)} dx \\ &= \frac{2 \cos(a+bx)}{5b \csc^{\frac{3}{2}}(a+bx)} + \frac{4\sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a-\frac{\pi}{2}+bx\right)\middle|2\right) \sqrt{\sin(a+bx)}}{5b} \end{aligned}$$

Mathematica [A] time = 0.15, size = 61, normalized size = 0.91

$$\frac{2\sqrt{\csc(a+bx)} \left(2\sqrt{\sin(a+bx)} E\left(\frac{1}{4}(-2a-2bx+\pi) \middle| 2\right) - \sin^2(a+bx) \cos(a+bx) \right)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^2/Sqrt[Csc[a + b*x]], x]

[Out] (-2*Sqrt[Csc[a + b*x]]*(2*EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] - Cos[a + b*x]*Sin[a + b*x]^2))/(5*b)

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\cos(bx+a)^2}{\sqrt{\csc(bx+a)}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/csc(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^2/sqrt(csc(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(bx+a)^2}{\sqrt{\csc(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/csc(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(cos(b*x + a)^2/sqrt(csc(b*x + a)), x)

maple [A] time = 0.14, size = 142, normalized size = 2.12

$$\frac{-\frac{2(\sin^4(bx+a))}{5} + \frac{2(\sin^2(bx+a))}{5} - \frac{4\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \text{EllipticE}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) + \frac{2\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)}}{5}}{\cos(bx+a) \sqrt{\sin(bx+a)} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2/csc(b*x+a)^(1/2), x)

[Out] (-2/5*sin(b*x+a)^4+2/5*sin(b*x+a)^2-4/5*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticE((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))+2/5*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2)))/cos(b*x+a)/sin(b*x+a)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(bx+a)^2}{\sqrt{\csc(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/csc(b*x+a)^(1/2), x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^2/sqrt(csc(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(a + bx)^2}{\sqrt{\frac{1}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2/(1/sin(a + b*x))^(1/2), x)

[Out] int(cos(a + b*x)^2/(1/sin(a + b*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(a + bx)}{\sqrt{\csc(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2/csc(b*x+a)**(1/2), x)

[Out] Integral(cos(a + b*x)**2/sqrt(csc(a + b*x)), x)

3.265 $\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx$

Optimal. Leaf size=21

$$\frac{2}{3} \csc^{\frac{3}{2}}(x) - \frac{2}{7} \csc^{\frac{7}{2}}(x)$$

[Out] $2/3*\csc(x)^{(3/2)}-2/7*\csc(x)^{(7/2)}$

Rubi [A] time = 0.02, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2621, 14}

$$\frac{2}{3} \csc^{\frac{3}{2}}(x) - \frac{2}{7} \csc^{\frac{7}{2}}(x)$$

Antiderivative was successfully verified.

[In] Int[Cos[x]^3*Csc[x]^(9/2),x]

[Out] (2*Csc[x]^(3/2))/3 - (2*Csc[x]^(7/2))/7

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.)^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> -Dist[(f*a^n)^(-1), Subst[Int[x^(m+n-1)/(-1+x^2/a^2)^(n+1)/2], x], x, a*Csc[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n+1)/2] && !(IntegerQ[(m+1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^3(x) \csc^{\frac{9}{2}}(x) dx &= -\text{Subst}\left(\int \sqrt{x} (-1+x^2) dx, x, \csc(x)\right) \\ &= -\text{Subst}\left(\int (-\sqrt{x} + x^{5/2}) dx, x, \csc(x)\right) \\ &= \frac{2}{3} \csc^{\frac{3}{2}}(x) - \frac{2}{7} \csc^{\frac{7}{2}}(x) \end{aligned}$$

Mathematica [A] time = 0.03, size = 18, normalized size = 0.86

$$\frac{2}{21} \csc^{\frac{3}{2}}(x) (7 - 3 \csc^2(x))$$

Antiderivative was successfully verified.

[In] Integrate[Cos[x]^3*Csc[x]^(9/2),x]

[Out] (2*Csc[x]^(3/2)*(7 - 3*Csc[x]^2))/21

fricas [A] time = 0.49, size = 22, normalized size = 1.05

$$\frac{2(7 \cos(x)^2 - 4)}{21(\cos(x)^2 - 1) \sin(x)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3*csc(x)^(9/2),x, algorithm="fricas")

[Out] 2/21*(7*cos(x)^2 - 4)/((cos(x)^2 - 1)*sin(x)^(3/2))

giac [A] time = 0.58, size = 14, normalized size = 0.67

$$\frac{2(7 \sin(x)^2 - 3)}{21 \sin(x)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3*csc(x)^(9/2),x, algorithm="giac")

[Out] 2/21*(7*sin(x)^2 - 3)/sin(x)^(7/2)

maple [A] time = 0.10, size = 14, normalized size = 0.67

$$-\frac{2}{7 \sin(x)^{\frac{7}{2}}} + \frac{2}{3 \sin(x)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x)^3*csc(x)^(9/2),x)

[Out] -2/7/sin(x)^(7/2)+2/3/sin(x)^(3/2)

maxima [A] time = 0.60, size = 13, normalized size = 0.62

$$\frac{2}{3 \sin(x)^{\frac{3}{2}}} - \frac{2}{7 \sin(x)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3*csc(x)^(9/2),x, algorithm="maxima")

[Out] 2/3/sin(x)^(3/2) - 2/7/sin(x)^(7/2)

mupad [B] time = 0.33, size = 16, normalized size = 0.76

$$\frac{2(7 \sin(x)^2 - 3) \left(\frac{1}{\sin(x)}\right)^{7/2}}{21}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x)^3*(1/sin(x))^(9/2),x)

[Out] (2*(7*sin(x)^2 - 3)*(1/sin(x))^(7/2))/21

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)**3*csc(x)**(9/2),x)

[Out] Timed out

3.266 $\int \cos^3(a + bx)\sqrt{\csc(a + bx)} dx$

Optimal. Leaf size=33

$$\frac{2}{b\sqrt{\csc(a + bx)}} - \frac{2}{5b \csc^{\frac{5}{2}}(a + bx)}$$

[Out] $-2/5/b/\csc(b*x+a)^{(5/2)}+2/b/\csc(b*x+a)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2621, 14}

$$\frac{2}{b\sqrt{\csc(a + bx)}} - \frac{2}{5b \csc^{\frac{5}{2}}(a + bx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3*Sqrt[Csc[a + b*x]],x]

[Out] $-2/(5*b*\csc[a + b*x]^{(5/2)}) + 2/(b*\sqrt{\csc[a + b*x]})$

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx)\sqrt{\csc(a + bx)} dx &= -\frac{\text{Subst}\left(\int \frac{-1+x^2}{x^{7/2}} dx, x, \csc(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-\frac{1}{x^{7/2}} + \frac{1}{x^{3/2}}\right) dx, x, \csc(a + bx)\right)}{b} \\ &= -\frac{2}{5b \csc^{\frac{5}{2}}(a + bx)} + \frac{2}{b\sqrt{\csc(a + bx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 27, normalized size = 0.82

$$\frac{\cos(2(a + bx)) + 9}{5b\sqrt{\csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^3*Sqrt[Csc[a + b*x]],x]

[Out] $(9 + \text{Cos}[2*(a + b*x)])/(5*b*\sqrt{\csc[a + b*x]})$

fricas [A] time = 0.58, size = 23, normalized size = 0.70

$$\frac{2 \left(\cos (bx + a)^2 + 4 \right) \sqrt{\sin (bx + a)}}{5 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] 2/5*(cos(b*x + a)^2 + 4)*sqrt(sin(b*x + a))/b

giac [A] time = 1.09, size = 24, normalized size = 0.73

$$-\frac{2 \left(\sin (bx + a)^{\frac{5}{2}} - 5 \sqrt{\sin (bx + a)} \right)}{5 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] -2/5*(sin(b*x + a)^(5/2) - 5*sqrt(sin(b*x + a)))/b

maple [A] time = 0.09, size = 26, normalized size = 0.79

$$\frac{-\frac{2 \left(\sin^{\frac{5}{2}}(bx+a) \right)}{5} + 2 \left(\sqrt{\sin (bx + a)} \right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^3*csc(b*x+a)^(1/2),x)

[Out] (-2/5*sin(b*x+a)^(5/2)+2*sin(b*x+a)^(1/2))/b

maxima [A] time = 0.70, size = 25, normalized size = 0.76

$$\frac{2 \left(\frac{5}{\sin (bx+a)^2} - 1 \right) \sin (bx + a)^{\frac{5}{2}}}{5 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 2/5*(5/sin(b*x + a)^2 - 1)*sin(b*x + a)^(5/2)/b

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \cos (a + bx)^3 \sqrt{\frac{1}{\sin (a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3*(1/sin(a + b*x))^(1/2),x)

[Out] int(cos(a + b*x)^3*(1/sin(a + b*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3*csc(b*x+a)**(1/2),x)

[Out] Timed out

$$3.267 \quad \int \frac{\cos^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal. Leaf size=35

$$\frac{2}{3b \csc^{\frac{3}{2}}(a+bx)} - \frac{2}{7b \csc^{\frac{7}{2}}(a+bx)}$$

[Out] $-2/7/b/\csc(b*x+a)^{(7/2)}+2/3/b/\csc(b*x+a)^{(3/2)}$

Rubi [A] time = 0.03, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2621, 14}

$$\frac{2}{3b \csc^{\frac{3}{2}}(a+bx)} - \frac{2}{7b \csc^{\frac{7}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3/Sqrt[Csc[a + b*x]],x]

[Out] $-2/(7*b*\csc[a + b*x]^{(7/2)}) + 2/(3*b*\csc[a + b*x]^{(3/2)})$

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.)^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := -Dist[(f*a^n)^(-1), Subst[Int[x^(m+n-1)/(-1+x^2/a^2)^((n+1)/2), x], x, a*Csc[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n+1)/2] && !(IntegerQ[(m+1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\cos^3(a+bx)}{\sqrt{\csc(a+bx)}} dx &= -\frac{\text{Subst}\left(\int \frac{-1+x^2}{x^{9/2}} dx, x, \csc(a+bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-\frac{1}{x^{9/2}} + \frac{1}{x^{5/2}}\right) dx, x, \csc(a+bx)\right)}{b} \\ &= -\frac{2}{7b \csc^{\frac{7}{2}}(a+bx)} + \frac{2}{3b \csc^{\frac{3}{2}}(a+bx)} \end{aligned}$$

Mathematica [A] time = 0.06, size = 29, normalized size = 0.83

$$\frac{2(7 \csc^2(a+bx) - 3)}{21b \csc^{\frac{7}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^3/Sqrt[Csc[a + b*x]],x]

[Out] $(2*(-3 + 7*\csc[a + b*x]^2))/(21*b*\csc[a + b*x]^{(7/2)})$

fricas [A] time = 0.55, size = 33, normalized size = 0.94

$$\frac{2(3 \cos(bx + a)^4 + \cos(bx + a)^2 - 4)}{21 b \sqrt{\sin(bx + a)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] -2/21*(3*cos(b*x + a)^4 + cos(b*x + a)^2 - 4)/(b*sqrt(sin(b*x + a)))

giac [A] time = 0.51, size = 25, normalized size = 0.71

$$\frac{2\left(\frac{7}{\sin(bx+a)^2} - 3\right) \sin(bx + a)^{\frac{7}{2}}}{21 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] 2/21*(7/sin(b*x + a)^2 - 3)*sin(b*x + a)^(7/2)/b

maple [A] time = 0.08, size = 26, normalized size = 0.74

$$\frac{-\frac{2\left(\sin^{\frac{7}{2}}(bx+a)\right)}{7} + \frac{2\left(\sin^{\frac{3}{2}}(bx+a)\right)}{3}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^3/csc(b*x+a)^(1/2),x)

[Out] (-2/7*sin(b*x+a)^(7/2)+2/3*sin(b*x+a)^(3/2))/b

maxima [A] time = 0.52, size = 25, normalized size = 0.71

$$\frac{2\left(\frac{7}{\sin(bx+a)^2} - 3\right) \sin(bx + a)^{\frac{7}{2}}}{21 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 2/21*(7/sin(b*x + a)^2 - 3)*sin(b*x + a)^(7/2)/b

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\cos(a + bx)^3}{\sqrt{\frac{1}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3/(1/sin(a + b*x))^(1/2),x)

[Out] int(cos(a + b*x)^3/(1/sin(a + b*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3/csc(b*x+a)**(1/2),x)

[Out] Timed out

3.268 $\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx$

Optimal. Leaf size=92

$$\frac{2 \cos^3(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{4 \cos(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{8 \sqrt{\sin(a + bx)} \sqrt{\csc(a + bx)} F\left(\frac{1}{2} \left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{7b}$$

[Out] $4/7 * \cos(b*x+a)/b / \csc(b*x+a)^{(1/2)} + 2/7 * \cos(b*x+a)^3 / b / \csc(b*x+a)^{(1/2)} - 8/7 * (\sin(1/2*a + 1/4*Pi + 1/2*b*x)^2)^{(1/2)} / \sin(1/2*a + 1/4*Pi + 1/2*b*x) * \text{EllipticF}(\cos(1/2*a + 1/4*Pi + 1/2*b*x), 2^{(1/2)}) * \csc(b*x+a)^{(1/2)} * \sin(b*x+a)^{(1/2)} / b$

Rubi [A] time = 0.08, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2628, 3771, 2641}

$$\frac{2 \cos^3(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{4 \cos(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{8 \sqrt{\sin(a + bx)} \sqrt{\csc(a + bx)} F\left(\frac{1}{2} \left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{7b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4*Sqrt[Csc[a + b*x]],x]

[Out] $(4 * \text{Cos}[a + b*x]) / (7 * b * \text{Sqrt}[\text{Csc}[a + b*x]]) + (2 * \text{Cos}[a + b*x]^3) / (7 * b * \text{Sqrt}[\text{Csc}[a + b*x]]) + (8 * \text{Sqrt}[\text{Csc}[a + b*x]] * \text{EllipticF}[(a - \text{Pi}/2 + b*x)/2, 2] * \text{Sqrt}[\text{Sin}[a + b*x]]) / (7 * b)$

Rule 2628

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx &= \frac{2 \cos^3(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{6}{7} \int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx \\ &= \frac{4 \cos(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{2 \cos^3(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{4}{7} \int \sqrt{\csc(a + bx)} dx \\ &= \frac{4 \cos(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{2 \cos^3(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{1}{7} \left(4 \sqrt{\csc(a + bx)} \sqrt{\sin(a + bx)} \right) \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{4 \cos(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{2 \cos^3(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{8 \sqrt{\csc(a + bx)} F\left(\frac{1}{2} \left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a + bx)}}{7b} \end{aligned}$$

Mathematica [A] time = 0.14, size = 63, normalized size = 0.68

$$\frac{\sqrt{\csc(a+bx)} \left(10 \sin(2(a+bx)) + \sin(4(a+bx)) - 32 \sqrt{\sin(a+bx)} F\left(\frac{1}{4}(-2a-2bx+\pi) \middle| 2\right) \right)}{28b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^4*Sqrt[Csc[a + b*x]], x]

[Out] (Sqrt[Csc[a + b*x]]*(-32*EllipticF[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] + 10*Sin[2*(a + b*x)] + Sin[4*(a + b*x)])/(28*b)

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}(\cos(bx+a)^4 \sqrt{\csc(bx+a)}, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*csc(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^4*sqrt(csc(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx+a)^4 \sqrt{\csc(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*csc(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(cos(b*x + a)^4*sqrt(csc(b*x + a)), x)

maple [A] time = 0.16, size = 100, normalized size = 1.09

$$\frac{\frac{2(\sin^5(bx+a))}{7} - \frac{8(\sin^3(bx+a))}{7} + \frac{6 \sin(bx+a)}{7} + \frac{4 \sqrt{\sin(bx+a)+1} \sqrt{-2 \sin(bx+a)+2} \sqrt{-\sin(bx+a)} \text{EllipticF}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right)}{\cos(bx+a) \sqrt{\sin(bx+a)} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^4*csc(b*x+a)^(1/2), x)

[Out] (2/7*sin(b*x+a)^5-8/7*sin(b*x+a)^3+6/7*sin(b*x+a)+4/7*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2)))/cos(b*x+a)/sin(b*x+a)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(bx+a)^4 \sqrt{\csc(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*csc(b*x+a)^(1/2), x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^4*sqrt(csc(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(a+bx)^4 \sqrt{\frac{1}{\sin(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)^4*(1/sin(a + b*x))^(1/2), x)
```

```
[Out] int(cos(a + b*x)^4*(1/sin(a + b*x))^(1/2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)**4*csc(b*x+a)**(1/2), x)
```

```
[Out] Integral(cos(a + b*x)**4*sqrt(csc(a + b*x)), x)
```

$$3.269 \quad \int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal. Leaf size=92

$$\frac{2 \cos^3(a+bx)}{9b \csc^2(a+bx)} + \frac{4 \cos(a+bx)}{15b \csc^2(a+bx)} + \frac{8 \sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a+bx - \frac{\pi}{2}\right) \middle| 2\right)}{15b}$$

[Out] 4/15*cos(b*x+a)/b/csc(b*x+a)^(3/2)+2/9*cos(b*x+a)^3/b/csc(b*x+a)^(3/2)-8/15*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticE(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] time = 0.08, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2628, 3771, 2639}

$$\frac{2 \cos^3(a+bx)}{9b \csc^2(a+bx)} + \frac{4 \cos(a+bx)}{15b \csc^2(a+bx)} + \frac{8 \sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a+bx - \frac{\pi}{2}\right) \middle| 2\right)}{15b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4/Sqrt[Csc[a + b*x]],x]

[Out] (4*Cos[a + b*x])/(15*b*Csc[a + b*x]^(3/2)) + (2*Cos[a + b*x]^3)/(9*b*Csc[a + b*x]^(3/2)) + (8*Sqrt[Csc[a + b*x]]*EllipticE[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/(15*b)

Rule 2628

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx &= \frac{2 \cos^3(a+bx)}{9b \csc^{\frac{3}{2}}(a+bx)} + \frac{2}{3} \int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx \\
&= \frac{4 \cos(a+bx)}{15b \csc^{\frac{3}{2}}(a+bx)} + \frac{2 \cos^3(a+bx)}{9b \csc^{\frac{3}{2}}(a+bx)} + \frac{4}{15} \int \frac{1}{\sqrt{\csc(a+bx)}} dx \\
&= \frac{4 \cos(a+bx)}{15b \csc^{\frac{3}{2}}(a+bx)} + \frac{2 \cos^3(a+bx)}{9b \csc^{\frac{3}{2}}(a+bx)} + \frac{1}{15} (4\sqrt{\csc(a+bx)} \sqrt{\sin(a+bx)}) \int \sqrt{\sin(a+bx)} \\
&= \frac{4 \cos(a+bx)}{15b \csc^{\frac{3}{2}}(a+bx)} + \frac{2 \cos^3(a+bx)}{9b \csc^{\frac{3}{2}}(a+bx)} + \frac{8\sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a+bx)}}{15b}
\end{aligned}$$

Mathematica [A] time = 0.41, size = 63, normalized size = 0.68

$$\frac{39 \cos(a+bx) + 5 \cos(3(a+bx)) - \frac{48 E\left(\frac{1}{4}(-2a-2bx+\pi) \middle| 2\right)}{\sin^{\frac{3}{2}}(a+bx)}}{90b \csc^{\frac{3}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^4/Sqrt[Csc[a + b*x]], x]

[Out] (39*Cos[a + b*x] + 5*Cos[3*(a + b*x)] - (48*EllipticE[(-2*a + Pi - 2*b*x)/4, 2])/Sin[a + b*x]^(3/2))/(90*b*Csc[a + b*x]^(3/2))

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\cos(bx+a)^4}{\sqrt{\csc(bx+a)}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/csc(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^4/sqrt(csc(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(bx+a)^4}{\sqrt{\csc(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/csc(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(cos(b*x + a)^4/sqrt(csc(b*x + a)), x)

maple [A] time = 0.14, size = 152, normalized size = 1.65

$$\frac{-\frac{2(\cos^6(bx+a))}{9} - \frac{8\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \text{EllipticE}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right)}{15} + \frac{4\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)}}{15}}{\cos(bx+a) \sqrt{\sin(bx+a)} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^4/csc(b*x+a)^(1/2), x)

[Out] $(-2/9*\cos(b*x+a)^6-8/15*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)}*\text{EllipticE}((\sin(b*x+a)+1)^{(1/2)},1/2*2^{(1/2)})+4/15*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)}*\text{EllipticF}((\sin(b*x+a)+1)^{(1/2)},1/2*2^{(1/2)})-2/45*\cos(b*x+a)^4+4/15*\cos(b*x+a)^2)/\cos(b*x+a)/\sin(b*x+a)^{(1/2)}/b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(bx+a)^4}{\sqrt{\csc(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^4/csc(b*x+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(cos(b*x + a)^4/sqrt(csc(b*x + a)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(a+bx)^4}{\sqrt{\frac{1}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^4/(1/sin(a + b*x))^(1/2),x)`

[Out] `int(cos(a + b*x)^4/(1/sin(a + b*x))^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**4/csc(b*x+a)**(1/2),x)`

[Out] `Integral(cos(a + b*x)**4/sqrt(csc(a + b*x)), x)`

$$3.270 \quad \int \cos(x) \csc^{\frac{7}{3}}(x) dx$$

Optimal. Leaf size=10

$$-\frac{3}{4} \csc^{\frac{4}{3}}(x)$$

[Out] $-3/4*\csc(x)^{(4/3)}$

Rubi [A] time = 0.02, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2621, 30}

$$-\frac{3}{4} \csc^{\frac{4}{3}}(x)$$

Antiderivative was successfully verified.

[In] Int[Cos[x]*Csc[x]^(7/3),x]

[Out] $(-3*\text{Csc}[x]^{(4/3)})/4$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2621

Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(x) \csc^{\frac{7}{3}}(x) dx &= -\text{Subst}\left(\int \sqrt[3]{x} dx, x, \csc(x)\right) \\ &= -\frac{3}{4} \csc^{\frac{4}{3}}(x) \end{aligned}$$

Mathematica [A] time = 0.01, size = 10, normalized size = 1.00

$$-\frac{3}{4} \csc^{\frac{4}{3}}(x)$$

Antiderivative was successfully verified.

[In] Integrate[Cos[x]*Csc[x]^(7/3),x]

[Out] $(-3*\text{Csc}[x]^{(4/3)})/4$

fricas [A] time = 0.57, size = 6, normalized size = 0.60

$$-\frac{3}{4 \sin(x)^{\frac{4}{3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)*csc(x)^(7/3),x, algorithm="fricas")

[Out] $-3/4/\sin(x)^{4/3}$

giac [A] time = 0.37, size = 6, normalized size = 0.60

$$-\frac{3}{4 \sin(x)^{\frac{4}{3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)*csc(x)^(7/3),x, algorithm="giac")`

[Out] $-3/4/\sin(x)^{4/3}$

maple [A] time = 0.02, size = 7, normalized size = 0.70

$$-\frac{3 \left(\csc^{\frac{4}{3}}(x) \right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(x)*csc(x)^(7/3),x)`

[Out] $-3/4*\csc(x)^{4/3}$

maxima [A] time = 1.13, size = 6, normalized size = 0.60

$$-\frac{3}{4 \sin(x)^{\frac{4}{3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)*csc(x)^(7/3),x, algorithm="maxima")`

[Out] $-3/4/\sin(x)^{4/3}$

mupad [B] time = 0.20, size = 8, normalized size = 0.80

$$-\frac{3 \left(\frac{1}{\sin(x)} \right)^{4/3}}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(x)*(1/sin(x))^(7/3),x)`

[Out] $-(3*(1/\sin(x))^{4/3})/4$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)*csc(x)**(7/3),x)`

[Out] Timed out

3.271 $\int \sqrt{\csc(a + bx)} \sec(a + bx) dx$

Optimal. Leaf size=32

$$\frac{\tanh^{-1}\left(\sqrt{\csc(a + bx)}\right)}{b} - \frac{\tan^{-1}\left(\sqrt{\csc(a + bx)}\right)}{b}$$

[Out] $-\arctan(\csc(b*x+a)^{(1/2)})/b + \operatorname{arctanh}(\csc(b*x+a)^{(1/2)})/b$

Rubi [A] time = 0.03, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2621, 329, 298, 203, 206}

$$\frac{\tanh^{-1}\left(\sqrt{\csc(a + bx)}\right)}{b} - \frac{\tan^{-1}\left(\sqrt{\csc(a + bx)}\right)}{b}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[Csc[a + b*x]]*Sec[a + b*x],x]`

[Out] `-(ArcTan[Sqrt[Csc[a + b*x]])/b) + ArcTanh[Sqrt[Csc[a + b*x]])/b`

Rule 203

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 206

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 298

`Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 329

`Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^n]^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 2621

`Int[(csc[(e_.) + (f_.)*(x_)])*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\begin{aligned}
\int \sqrt{\csc(a+bx)} \sec(a+bx) dx &= -\frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-1+x^2} dx, x, \csc(a+bx)\right)}{b} \\
&= -\frac{2 \text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \sqrt{\csc(a+bx)}\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{b} - \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{b} \\
&= -\frac{\tan^{-1}\left(\sqrt{\csc(a+bx)}\right)}{b} + \frac{\tanh^{-1}\left(\sqrt{\csc(a+bx)}\right)}{b}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 47, normalized size = 1.47

$$\frac{\sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} \left(\tan^{-1}\left(\sqrt{\sin(a+bx)}\right) + \tanh^{-1}\left(\sqrt{\sin(a+bx)}\right) \right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Csc[a + b*x]]*Sec[a + b*x], x]

[Out] ((ArcTan[Sqrt[Sin[a + b*x]]] + ArcTanh[Sqrt[Sin[a + b*x]]])*Sqrt[Csc[a + b*x]]*Sqrt[Sin[a + b*x]])/b

fricas [B] time = 0.82, size = 95, normalized size = 2.97

$$\frac{2 \arctan\left(\frac{\sin(bx+a)-1}{2\sqrt{\sin(bx+a)}}\right) + \log\left(\frac{\cos(bx+a)^2 + \frac{4(\cos(bx+a)^2 - \sin(bx+a)-1)}{\sqrt{\sin(bx+a)}} - 6\sin(bx+a)-2}{\cos(bx+a)^2 + 2\sin(bx+a)-2}\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a), x, algorithm="fricas")

[Out] 1/4*(2*arctan(1/2*(sin(b*x + a) - 1)/sqrt(sin(b*x + a))) + log((cos(b*x + a)^2 + 4*(cos(b*x + a)^2 - sin(b*x + a) - 1)/sqrt(sin(b*x + a)) - 6*sin(b*x + a) - 2)/(cos(b*x + a)^2 + 2*sin(b*x + a) - 2)))/b

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(bx+a)} \sec(bx+a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a), x, algorithm="giac")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a), x)

maple [A] time = 0.18, size = 28, normalized size = 0.88

$$\frac{\operatorname{arctanh}\left(\sqrt{\sin(bx+a)}\right)}{b} + \frac{\operatorname{arctan}\left(\sqrt{\sin(bx+a)}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)^(1/2)*sec(b*x+a), x)

[Out] $1/b \cdot \operatorname{arctanh}(\sin(bx+a)^{1/2}) + 1/b \cdot \operatorname{arctan}(\sin(bx+a)^{1/2})$

maxima [A] time = 1.35, size = 41, normalized size = 1.28

$$\frac{2 \arctan\left(\frac{1}{\sqrt{\sin(bx+a)}}\right) - \log\left(\frac{1}{\sqrt{\sin(bx+a)}} + 1\right) + \log\left(\frac{1}{\sqrt{\sin(bx+a)}} - 1\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^(1/2)*sec(b*x+a),x, algorithm="maxima")`

[Out] $-1/2 * (2 * \arctan(1/\sqrt{\sin(bx+a)}) - \log(1/\sqrt{\sin(bx+a)} + 1) + \log(1/\sqrt{\sin(bx+a)} - 1)) / b$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\sqrt{\frac{1}{\sin(a+bx)}}}{\cos(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/sin(a + b*x))^(1/2)/cos(a + b*x),x)`

[Out] `int((1/sin(a + b*x))^(1/2)/cos(a + b*x), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(a+bx)} \sec(a+bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)**(1/2)*sec(b*x+a),x)`

[Out] `Integral(sqrt(csc(a + b*x))*sec(a + b*x), x)`

$$3.272 \quad \int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal. Leaf size=31

$$\frac{\tan^{-1}\left(\sqrt{\csc(a+bx)}\right)}{b} + \frac{\tanh^{-1}\left(\sqrt{\csc(a+bx)}\right)}{b}$$

[Out] arctan(csc(b*x+a)^(1/2))/b+arctanh(csc(b*x+a)^(1/2))/b

Rubi [A] time = 0.03, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2621, 329, 212, 206, 203}

$$\frac{\tan^{-1}\left(\sqrt{\csc(a+bx)}\right)}{b} + \frac{\tanh^{-1}\left(\sqrt{\csc(a+bx)}\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]/Sqrt[Csc[a + b*x]], x]

[Out] ArcTan[Sqrt[Csc[a + b*x]]]/b + ArcTanh[Sqrt[Csc[a + b*x]]]/b

Rule 203

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 329

Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^n]^(1/k), x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*sec[(e_.) + (f_.)*(x_)^(n_.)], x_Symbol] := -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned}
\int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}(-1+x^2)} dx, x, \csc(a+bx)\right)}{b} \\
&= -\frac{2\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\csc(a+bx)}\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{b} + \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{b} \\
&= \frac{\tan^{-1}\left(\sqrt{\csc(a+bx)}\right)}{b} + \frac{\tanh^{-1}\left(\sqrt{\csc(a+bx)}\right)}{b}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 50, normalized size = 1.61

$$-\frac{\sqrt{\sin(a+bx)}\sqrt{\csc(a+bx)}\left(\tan^{-1}\left(\sqrt{\sin(a+bx)}\right) - \tanh^{-1}\left(\sqrt{\sin(a+bx)}\right)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]/Sqrt[Csc[a + b*x]], x]

[Out] -(((ArcTan[Sqrt[Sin[a + b*x]]] - ArcTanh[Sqrt[Sin[a + b*x]]])*Sqrt[Csc[a + b*x]]*Sqrt[Sin[a + b*x]])/b)

fricas [B] time = 0.65, size = 97, normalized size = 3.13

$$-\frac{2 \arctan\left(\frac{\sin(bx+a)-1}{2\sqrt{\sin(bx+a)}}\right) - \log\left(\frac{\cos(bx+a)^2 + \frac{4(\cos(bx+a)^2 - \sin(bx+a)-1)}{\sqrt{\sin(bx+a)}} - 6\sin(bx+a) - 2}{\cos(bx+a)^2 + 2\sin(bx+a) - 2}\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/csc(b*x+a)^(1/2), x, algorithm="fricas")

[Out] -1/4*(2*arctan(1/2*(sin(b*x + a) - 1)/sqrt(sin(b*x + a))) - log((cos(b*x + a)^2 + 4*(cos(b*x + a)^2 - sin(b*x + a) - 1)/sqrt(sin(b*x + a)) - 6*sin(b*x + a) - 2)/(cos(b*x + a)^2 + 2*sin(b*x + a) - 2)))/b

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(bx+a)}{\sqrt{\csc(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/csc(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(sec(b*x + a)/sqrt(csc(b*x + a)), x)

maple [A] time = 0.19, size = 48, normalized size = 1.55

$$-\frac{\ln\left(\sqrt{\sin(bx+a)}-1\right)}{2b} + \frac{\ln\left(\sqrt{\sin(bx+a)}+1\right)}{2b} - \frac{\arctan\left(\sqrt{\sin(bx+a)}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)/csc(b*x+a)^(1/2), x)

[Out] $-1/2/b*\ln(\sin(b*x+a)^{(1/2)}-1)+1/2/b*\ln(\sin(b*x+a)^{(1/2)}+1)-1/b*\arctan(\sin(b*x+a)^{(1/2)})$

maxima [A] time = 0.70, size = 41, normalized size = 1.32

$$\frac{2 \arctan\left(\frac{1}{\sqrt{\sin(bx+a)}}\right) + \log\left(\frac{1}{\sqrt{\sin(bx+a)}} + 1\right) - \log\left(\frac{1}{\sqrt{\sin(bx+a)}} - 1\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="maxima")`

[Out] $1/2*(2*\arctan(1/\sqrt{\sin(b*x + a)})) + \log(1/\sqrt{\sin(b*x + a)} + 1) - \log(1/\sqrt{\sin(b*x + a)} - 1))/b$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\cos(a + bx) \sqrt{\frac{1}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(a + b*x)*(1/sin(a + b*x))^(1/2)),x)`

[Out] `int(1/(cos(a + b*x)*(1/sin(a + b*x))^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(a + bx)}{\sqrt{\csc(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)/csc(b*x+a)**(1/2),x)`

[Out] `Integral(sec(a + b*x)/sqrt(csc(a + b*x)), x)`

3.273 $\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx$

Optimal. Leaf size=61

$$\frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{\sqrt{\sin(a + bx)} \sqrt{\csc(a + bx)} F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{b}$$

[Out] sec(b*x+a)/b/csc(b*x+a)^(1/2)-(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticF(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] time = 0.05, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, number of rules / integrand size = 0.158, Rules used = {2626, 3771, 2641}

$$\frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{\sqrt{\sin(a + bx)} \sqrt{\csc(a + bx)} F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^2,x]

[Out] Sec[a + b*x]/(b*Sqrt[Csc[a + b*x]]) + (Sqrt[Csc[a + b*x]]*EllipticF[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/b

Rule 2626

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx &= \frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{1}{2} \int \sqrt{\csc(a + bx)} dx \\ &= \frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{1}{2} \left(\sqrt{\csc(a + bx)} \sqrt{\sin(a + bx)} \right) \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{\sqrt{\csc(a + bx)} F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a + bx)}}{b} \end{aligned}$$

Mathematica [A] time = 0.14, size = 49, normalized size = 0.80

$$\frac{\sec(a + bx) + \frac{F\left(\frac{1}{4}(2a+2bx-\pi)\middle|2\right)}{\sqrt{\sin(a+bx)}}}{b\sqrt{\csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^2,x]

[Out] (Sec[a + b*x] + EllipticF[(2*a - Pi + 2*b*x)/4, 2]/Sqrt[Sin[a + b*x]])/(b*Sqrt[Csc[a + b*x]])

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{\csc(bx + a)} \sec(bx + a)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^2,x, algorithm="fricas")

[Out] integral(sqrt(csc(b*x + a))*sec(b*x + a)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(bx + a)} \sec(bx + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^2,x, algorithm="giac")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a)^2, x)

maple [A] time = 0.19, size = 123, normalized size = 2.02

$$\frac{\sqrt{(\cos^2(bx + a) \sin(bx + a))} \left(\sqrt{\sin(bx + a) + 1} \sqrt{-2 \sin(bx + a) + 2} \sqrt{-\sin(bx + a)} \text{EllipticF}\left(\sqrt{\sin(bx + a)}, \frac{1}{2}\right) \right)}{2\sqrt{-\sin(bx + a)} (\sin(bx + a) - 1) (\sin(bx + a) + 1) \cos(bx + a) \sqrt{\sin(bx + a)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)^(1/2)*sec(b*x+a)^2,x)

[Out] 1/2*(cos(b*x+a)^2*sin(b*x+a))^(1/2)*((sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))+2*sin(b*x+a))/(-sin(b*x+a)*(sin(b*x+a)-1)*(sin(b*x+a)+1))^(1/2)/cos(b*x+a)/sin(b*x+a)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(bx + a)} \sec(bx + a)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{\frac{1}{\sin(a+bx)}}}{\cos(a + bx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^2,x)`

[Out] `int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^2, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)**(1/2)*sec(b*x+a)**2,x)`

[Out] `Integral(sqrt(csc(a + b*x))*sec(a + b*x)**2, x)`

$$3.274 \quad \int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal. Leaf size=62

$$\frac{\sec(a+bx)}{b \csc^{\frac{3}{2}}(a+bx)} - \frac{\sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a+bx - \frac{\pi}{2}\right) \middle| 2\right)}{b}$$

[Out] sec(b*x+a)/b/csc(b*x+a)^(3/2)+(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticE(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] time = 0.05, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2626, 3771, 2639}

$$\frac{\sec(a+bx)}{b \csc^{\frac{3}{2}}(a+bx)} - \frac{\sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a+bx - \frac{\pi}{2}\right) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^2/Sqrt[Csc[a + b*x]],x]

[Out] Sec[a + b*x]/(b*Csc[a + b*x]^(3/2)) - (Sqrt[Csc[a + b*x]]*EllipticE[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/b

Rule 2626

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx &= \frac{\sec(a+bx)}{b \csc^{\frac{3}{2}}(a+bx)} - \frac{1}{2} \int \frac{1}{\sqrt{\csc(a+bx)}} dx \\ &= \frac{\sec(a+bx)}{b \csc^{\frac{3}{2}}(a+bx)} - \frac{1}{2} \left(\sqrt{\csc(a+bx)} \sqrt{\sin(a+bx)} \right) \int \sqrt{\sin(a+bx)} dx \\ &= \frac{\sec(a+bx)}{b \csc^{\frac{3}{2}}(a+bx)} - \frac{\sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a+bx)}}{b} \end{aligned}$$

Mathematica [A] time = 0.15, size = 54, normalized size = 0.87

$$\frac{\sqrt{\csc(a+bx)} \left(\sin(a+bx) \tan(a+bx) + \sqrt{\sin(a+bx)} E\left(\frac{1}{4}(-2a-2bx+\pi) \middle| 2\right) \right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^2/Sqrt[Csc[a + b*x]], x]

[Out] (Sqrt[Csc[a + b*x]]*(EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] + Sin[a + b*x]*Tan[a + b*x]))/b

fricas [F] time = 0.83, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sec(bx+a)^2}{\sqrt{\csc(bx+a)}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/csc(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^2/sqrt(csc(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(bx+a)^2}{\sqrt{\csc(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/csc(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^2/sqrt(csc(b*x + a)), x)

maple [B] time = 0.19, size = 177, normalized size = 2.85

$$\frac{\sqrt{(\cos^2(bx+a)) \sin(bx+a)} \left(2\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \text{EllipticE}\left(\sqrt{\sin(bx+a)} \right) + \dots \right)}{2\sqrt{-\sin(bx+a)} (\sin(bx+a) - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^2/csc(b*x+a)^(1/2), x)

[Out] 1/2*(cos(b*x+a)^2*sin(b*x+a))^(1/2)*(2*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticE((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))-sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))-2*cos(b*x+a)^2+2)/(-sin(b*x+a)*(sin(b*x+a)-1)*(sin(b*x+a)+1))^(1/2)/cos(b*x+a)/sin(b*x+a)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(bx+a)^2}{\sqrt{\csc(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/csc(b*x+a)^(1/2), x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^2/sqrt(csc(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(a+bx)^2 \sqrt{\frac{1}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^2*(1/sin(a + b*x))^(1/2)), x)

[Out] int(1/(cos(a + b*x)^2*(1/sin(a + b*x))^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**2/csc(b*x+a)**(1/2), x)

[Out] Integral(sec(a + b*x)**2/sqrt(csc(a + b*x)), x)

3.275 $\int \sqrt{\csc(a + bx)} \sec^3(a + bx) dx$

Optimal. Leaf size=62

$$-\frac{3 \tan^{-1}(\sqrt{\csc(a + bx)})}{4b} + \frac{\sec^2(a + bx)}{2b\sqrt{\csc(a + bx)}} + \frac{3 \tanh^{-1}(\sqrt{\csc(a + bx)})}{4b}$$

[Out] $-3/4*\arctan(\csc(b*x+a)^{(1/2)})/b+3/4*\operatorname{arctanh}(\csc(b*x+a)^{(1/2)})/b+1/2*\sec(b*x+a)^2/b/\csc(b*x+a)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2621, 288, 329, 298, 203, 206}

$$-\frac{3 \tan^{-1}(\sqrt{\csc(a + bx)})}{4b} + \frac{\sec^2(a + bx)}{2b\sqrt{\csc(a + bx)}} + \frac{3 \tanh^{-1}(\sqrt{\csc(a + bx)})}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^3,x]

[Out] $(-3*\operatorname{ArcTan}[\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]])/(4*b) + (3*\operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]])/(4*b) + \operatorname{Sec}[a + b*x]^2/(2*b*\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]])$

Rule 203

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 288

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c^(n - 1)*(c*x)^(m - n + 1)*(a + b*x^n)^(p + 1))/(b*n*(p + 1)), x] - Dist[(c^n*(m - n + 1))/(b*n*(p + 1)), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 298

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 329

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^n]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2621


```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^{5/2}}{(-1+x^2)^2} dx, x, \csc(a+bx)\right)}{b} \\ &= \frac{\sec^2(a+bx)}{2b\sqrt{\csc(a+bx)}} - \frac{3 \text{Subst}\left(\int \frac{\sqrt{x}}{-1+x^2} dx, x, \csc(a+bx)\right)}{4b} \\ &= \frac{\sec^2(a+bx)}{2b\sqrt{\csc(a+bx)}} - \frac{3 \text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \sqrt{\csc(a+bx)}\right)}{2b} \\ &= \frac{\sec^2(a+bx)}{2b\sqrt{\csc(a+bx)}} + \frac{3 \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{4b} - \frac{3 \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{4b} \\ &= -\frac{3 \tan^{-1}\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{3 \tanh^{-1}\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{\sec^2(a+bx)}{2b\sqrt{\csc(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.11, size = 73, normalized size = 1.18

$$\frac{\sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} \left(2\sqrt{\sin(a+bx)} \sec^2(a+bx) + 3 \left(\tan^{-1}\left(\sqrt{\sin(a+bx)}\right) + \tanh^{-1}\left(\sqrt{\sin(a+bx)}\right)\right)\right)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^3,x]

[Out] (Sqrt[Csc[a + b*x]]*(3*(ArcTan[Sqrt[Sin[a + b*x]]] + ArcTanh[Sqrt[Sin[a + b*x]]]) + 2*Sec[a + b*x]^2*Sqrt[Sin[a + b*x]]*Sqrt[Sin[a + b*x]])/(4*b)

fricas [B] time = 0.68, size = 131, normalized size = 2.11

$$\frac{6 \arctan\left(\frac{\sin(bx+a)-1}{2\sqrt{\sin(bx+a)}}\right) \cos(bx+a)^2 + 3 \cos(bx+a)^2 \log\left(\frac{\cos(bx+a)^2 + \frac{4(\cos(bx+a)^2 - \sin(bx+a)-1)}{\sqrt{\sin(bx+a)}} - 6 \sin(bx+a) - 2}{\cos(bx+a)^2 + 2 \sin(bx+a) - 2}\right) + 8 \sqrt{\sin(bx+a)}}{16 b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^3,x, algorithm="fricas")

[Out] 1/16*(6*arctan(1/2*(sin(b*x + a) - 1)/sqrt(sin(b*x + a)))*cos(b*x + a)^2 + 3*cos(b*x + a)^2*log((cos(b*x + a)^2 + 4*(cos(b*x + a)^2 - sin(b*x + a) - 1)/sqrt(sin(b*x + a)) - 6*sin(b*x + a) - 2)/(cos(b*x + a)^2 + 2*sin(b*x + a) - 2)) + 8*sqrt(sin(b*x + a)))/(b*cos(b*x + a)^2)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(bx+a)} \sec(bx+a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^3,x, algorithm="giac")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a)^3, x)

maple [A] time = 0.19, size = 73, normalized size = 1.18

$$\frac{-\left(3 \ln\left(\sqrt{\sin}(bx+a)-1\right)-3 \ln\left(\sqrt{\sin}(bx+a)+1\right)-6 \arctan\left(\sqrt{\sin}(bx+a)\right)\right)\left(\cos^2(bx+a)\right)+4\left(\sqrt{\sin}(bx+a)\right)}{8 \cos(bx+a)^2 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)^(1/2)*sec(b*x+a)^3,x)

[Out] 1/8*(-(3*ln(sin(b*x+a)^(1/2)-1)-3*ln(sin(b*x+a)^(1/2)+1)-6*arctan(sin(b*x+a)^(1/2)))*cos(b*x+a)^2+4*sin(b*x+a)^(1/2))/cos(b*x+a)^2/b

maxima [A] time = 1.33, size = 65, normalized size = 1.05

$$\frac{\frac{4}{\left(\frac{1}{\sin(bx+a)^2}-1\right)\sin(bx+a)^{\frac{3}{2}}}-6 \arctan\left(\frac{1}{\sqrt{\sin(bx+a)}}\right)+3 \log\left(\frac{1}{\sqrt{\sin(bx+a)}}+1\right)-3 \log\left(\frac{1}{\sqrt{\sin(bx+a)}}-1\right)}{8 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^3,x, algorithm="maxima")

[Out] 1/8*(4/(((1/sin(b*x + a))^2 - 1)*sin(b*x + a)^(3/2)) - 6*arctan(1/sqrt(sin(b*x + a)))) + 3*log(1/sqrt(sin(b*x + a)) + 1) - 3*log(1/sqrt(sin(b*x + a)) - 1))/b

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{\frac{1}{\sin(a+bx)}}}{\cos(a+bx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^3,x)

[Out] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)**(1/2)*sec(b*x+a)**3,x)

[Out] Integral(sqrt(csc(a + b*x))*sec(a + b*x)**3, x)

$$3.276 \quad \int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal. Leaf size=62

$$\frac{\sec^2(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} + \frac{\tan^{-1}(\sqrt{\csc(a+bx)})}{4b} + \frac{\tanh^{-1}(\sqrt{\csc(a+bx)})}{4b}$$

[Out] $1/4*\arctan(\csc(b*x+a)^{(1/2)})/b+1/4*\operatorname{arctanh}(\csc(b*x+a)^{(1/2)})/b+1/2*\sec(b*x+a)^2/b/\csc(b*x+a)^{(3/2)}$

Rubi [A] time = 0.05, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2621, 288, 329, 212, 206, 203}

$$\frac{\sec^2(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} + \frac{\tan^{-1}(\sqrt{\csc(a+bx)})}{4b} + \frac{\tanh^{-1}(\sqrt{\csc(a+bx)})}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^3/Sqrt[Csc[a + b*x]], x]

[Out] ArcTan[Sqrt[Csc[a + b*x]]]/(4*b) + ArcTanh[Sqrt[Csc[a + b*x]]]/(4*b) + Sec[a + b*x]^2/(2*b*Csc[a + b*x]^(3/2))

Rule 203

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTan[(Rt[b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] :> With[{r = Numerator[Rt[-(a/b), 2]], s = Denominator[Rt[-(a/b), 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 288

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(c^(n-1)*(c*x)^(m-n+1)*(a+b*x^n)^(p+1))/(b*n*(p+1)), x] - Dist[(c^n*(m-n+1))/(b*n*(p+1)), Int[(c*x)^(m-n)*(a+b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m+1, n] && !LtQ[(m+n*(p+1)+1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 329

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m+1)-1)*(a+(b*x^(k*n)))/c^n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*sec[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] :> -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(a + bx)}{\sqrt{\csc(a + bx)}} dx &= -\frac{\text{Subst}\left(\int \frac{x^{3/2}}{(-1+x^2)^2} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{\sec^2(a + bx)}{2b \csc^{\frac{3}{2}}(a + bx)} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}(-1+x^2)} dx, x, \csc(a + bx)\right)}{4b} \\ &= \frac{\sec^2(a + bx)}{2b \csc^{\frac{3}{2}}(a + bx)} - \frac{\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\csc(a + bx)}\right)}{2b} \\ &= \frac{\sec^2(a + bx)}{2b \csc^{\frac{3}{2}}(a + bx)} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\csc(a + bx)}\right)}{4b} + \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\csc(a + bx)}\right)}{4b} \\ &= \frac{\tan^{-1}\left(\sqrt{\csc(a + bx)}\right)}{4b} + \frac{\tanh^{-1}\left(\sqrt{\csc(a + bx)}\right)}{4b} + \frac{\sec^2(a + bx)}{2b \csc^{\frac{3}{2}}(a + bx)} \end{aligned}$$

Mathematica [C] time = 0.03, size = 33, normalized size = 0.53

$$\frac{{}_2F_1\left(\frac{3}{4}, 2; \frac{7}{4}; \sin^2(a + bx)\right)}{3b \csc^{\frac{3}{2}}(a + bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^3/Sqrt[Csc[a + b*x]], x]

[Out] (2*Hypergeometric2F1[3/4, 2, 7/4, Sin[a + b*x]^2])/(3*b*Csc[a + b*x]^(3/2))

fricas [B] time = 0.75, size = 141, normalized size = 2.27

$$\frac{2 \arctan\left(\frac{\sin(bx+a)-1}{2\sqrt{\sin(bx+a)}}\right) \cos(bx+a)^2 - \cos(bx+a)^2 \log\left(\frac{\cos(bx+a)^2 + \frac{4(\cos(bx+a)^2 - \sin(bx+a)-1)}{\sqrt{\sin(bx+a)}} - 6 \sin(bx+a) - 2}{\cos(bx+a)^2 + 2 \sin(bx+a) - 2}\right) + \frac{8(\cos(bx+a))}{\sqrt{\sin(bx+a)}}}{16b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/csc(b*x+a)^(1/2), x, algorithm="fricas")

[Out] -1/16*(2*arctan(1/2*(sin(b*x + a) - 1)/sqrt(sin(b*x + a)))*cos(b*x + a)^2 - cos(b*x + a)^2*log((cos(b*x + a)^2 + 4*(cos(b*x + a)^2 - sin(b*x + a) - 1)/sqrt(sin(b*x + a)) - 6*sin(b*x + a) - 2)/(cos(b*x + a)^2 + 2*sin(b*x + a) - 2)) + 8*(cos(b*x + a)^2 - 1)/sqrt(sin(b*x + a)))/(b*cos(b*x + a)^2)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(bx + a)^3}{\sqrt{\csc(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(sec(b*x + a)^3/sqrt(csc(b*x + a)), x)

maple [A] time = 0.17, size = 71, normalized size = 1.15

$$\frac{-\left(\ln\left(\sqrt{\sin}(bx+a)-1\right)-\ln\left(\sqrt{\sin}(bx+a)+1\right)+2\arctan\left(\sqrt{\sin}(bx+a)\right)\right)\left(\cos^2(bx+a)\right)+4\left(\sin^{\frac{3}{2}}(bx+a)\right)}{8\cos(bx+a)^2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^3/csc(b*x+a)^(1/2),x)

[Out] 1/8*(-(ln(sin(b*x+a)^(1/2)-1)-ln(sin(b*x+a)^(1/2)+1)+2*arctan(sin(b*x+a)^(1/2)))*cos(b*x+a)^2+4*sin(b*x+a)^(3/2))/cos(b*x+a)^2/b

maxima [A] time = 1.44, size = 63, normalized size = 1.02

$$\frac{\frac{4}{\left(\frac{1}{\sin(bx+a)^2}-1\right)\sqrt{\sin(bx+a)}}+2\arctan\left(\frac{1}{\sqrt{\sin(bx+a)}}\right)+\log\left(\frac{1}{\sqrt{\sin(bx+a)}}+1\right)-\log\left(\frac{1}{\sqrt{\sin(bx+a)}}-1\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 1/8*(4/((1/sin(b*x + a)^2 - 1)*sqrt(sin(b*x + a))) + 2*arctan(1/sqrt(sin(b*x + a))) + log(1/sqrt(sin(b*x + a)) + 1) - log(1/sqrt(sin(b*x + a)) - 1))/b

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(a+bx)^3 \sqrt{\frac{1}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^3*(1/sin(a + b*x))^(1/2)),x)

[Out] int(1/(cos(a + b*x)^3*(1/sin(a + b*x))^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**3/csc(b*x+a)**(1/2),x)

[Out] Integral(sec(a + b*x)**3/sqrt(csc(a + b*x)), x)

3.277 $\int \sqrt{\csc(a+bx)} \sec^4(a+bx) dx$

Optimal. Leaf size=92

$$\frac{\sec^3(a+bx)}{3b\sqrt{\csc(a+bx)}} + \frac{5\sec(a+bx)}{6b\sqrt{\csc(a+bx)}} + \frac{5\sqrt{\sin(a+bx)}\sqrt{\csc(a+bx)}F\left(\frac{1}{2}\left(a+bx-\frac{\pi}{2}\right)\middle|2\right)}{6b}$$

[Out] 5/6*sec(b*x+a)/b/csc(b*x+a)^(1/2)+1/3*sec(b*x+a)^3/b/csc(b*x+a)^(1/2)-5/6*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticF(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] time = 0.08, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2626, 3771, 2641}

$$\frac{\sec^3(a+bx)}{3b\sqrt{\csc(a+bx)}} + \frac{5\sec(a+bx)}{6b\sqrt{\csc(a+bx)}} + \frac{5\sqrt{\sin(a+bx)}\sqrt{\csc(a+bx)}F\left(\frac{1}{2}\left(a+bx-\frac{\pi}{2}\right)\middle|2\right)}{6b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^4,x]

[Out] (5*Sec[a + b*x])/(6*b*Sqrt[Csc[a + b*x]]) + Sec[a + b*x]^3/(3*b*Sqrt[Csc[a + b*x]]) + (5*Sqrt[Csc[a + b*x]]*EllipticF[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/(6*b)

Rule 2626

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sqrt{\csc(a+bx)} \sec^4(a+bx) dx &= \frac{\sec^3(a+bx)}{3b\sqrt{\csc(a+bx)}} + \frac{5}{6} \int \sqrt{\csc(a+bx)} \sec^2(a+bx) dx \\ &= \frac{5\sec(a+bx)}{6b\sqrt{\csc(a+bx)}} + \frac{\sec^3(a+bx)}{3b\sqrt{\csc(a+bx)}} + \frac{5}{12} \int \sqrt{\csc(a+bx)} dx \\ &= \frac{5\sec(a+bx)}{6b\sqrt{\csc(a+bx)}} + \frac{\sec^3(a+bx)}{3b\sqrt{\csc(a+bx)}} + \frac{1}{12} (5\sqrt{\csc(a+bx)}\sqrt{\sin(a+bx)}) \int \frac{1}{\sqrt{\csc(a+bx)}} dx \\ &= \frac{5\sec(a+bx)}{6b\sqrt{\csc(a+bx)}} + \frac{\sec^3(a+bx)}{3b\sqrt{\csc(a+bx)}} + \frac{5\sqrt{\csc(a+bx)}F\left(\frac{1}{2}\left(a-\frac{\pi}{2}+bx\right)\middle|2\right)}{6b} \sqrt{\csc(a+bx)} \end{aligned}$$

Mathematica [A] time = 0.38, size = 64, normalized size = 0.70

$$\frac{\sqrt{\csc(a+bx)} \left(\tan(a+bx) (2 \sec^2(a+bx) + 5) - 5 \sqrt{\sin(a+bx)} F\left(\frac{1}{4}(-2a-2bx+\pi) \middle| 2\right) \right)}{6b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^4,x]

[Out] (Sqrt[Csc[a + b*x]]*(-5*EllipticF[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] + (5 + 2*Sec[a + b*x]^2)*Tan[a + b*x]))/(6*b)

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{\csc(bx+a)} \sec(bx+a)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^4,x, algorithm="fricas")

[Out] integral(sqrt(csc(b*x + a))*sec(b*x + a)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(bx+a)} \sec(bx+a)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^4,x, algorithm="giac")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a)^4, x)

maple [A] time = 0.20, size = 168, normalized size = 1.83

$$\frac{\sqrt{(\cos^2(bx+a)) \sin(bx+a)} \left(5\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \text{EllipticF}\left(\sqrt{\sin(bx+a)}, \frac{1}{2}\right) \right)}{12(\sin(bx+a)+1)\sqrt{-\sin(bx+a)}(\sin(bx+a)-1)(\sin(bx+a)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)^(1/2)*sec(b*x+a)^4,x)

[Out] -1/12*(cos(b*x+a)^2*sin(b*x+a))^(1/2)*(5*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))*cos(b*x+a)^2+10*cos(b*x+a)^2*sin(b*x+a)+4*sin(b*x+a))/(sin(b*x+a)+1)/(-sin(b*x+a)*(sin(b*x+a)-1)*(sin(b*x+a)+1))^(1/2)/(sin(b*x+a)-1)/cos(b*x+a)/sin(b*x+a)^(1/2)/b

maxima [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^4,x, algorithm="maxima")

[Out] Timed out

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{1}{\sin(a+bx)}}}{\cos(a+bx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^4, x)
```

```
[Out] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^4, x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(b*x+a)**(1/2)*sec(b*x+a)**4, x)
```

```
[Out] Integral(sqrt(csc(a + b*x))*sec(a + b*x)**4, x)
```


$$3.278 \quad \int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal. Leaf size=92

$$\frac{\sec^3(a+bx)}{3b \csc^2(a+bx)} + \frac{\sec(a+bx)}{2b \csc^2(a+bx)} - \frac{\sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a+bx - \frac{\pi}{2}\right) \middle| 2\right)}{2b}$$

[Out] 1/2*sec(b*x+a)/b/csc(b*x+a)^(3/2)+1/3*sec(b*x+a)^3/b/csc(b*x+a)^(3/2)+1/2*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticE(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] time = 0.08, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2626, 3771, 2639}

$$\frac{\sec^3(a+bx)}{3b \csc^2(a+bx)} + \frac{\sec(a+bx)}{2b \csc^2(a+bx)} - \frac{\sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a+bx - \frac{\pi}{2}\right) \middle| 2\right)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^4/Sqrt[Csc[a + b*x]],x]

[Out] Sec[a + b*x]/(2*b*Csc[a + b*x]^(3/2)) + Sec[a + b*x]^3/(3*b*Csc[a + b*x]^(3/2)) - (Sqrt[Csc[a + b*x]]*EllipticE[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/(2*b)

Rule 2626

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx &= \frac{\sec^3(a+bx)}{3b \csc^{\frac{3}{2}}(a+bx)} + \frac{1}{2} \int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx \\
&= \frac{\sec(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} + \frac{\sec^3(a+bx)}{3b \csc^{\frac{3}{2}}(a+bx)} - \frac{1}{4} \int \frac{1}{\sqrt{\csc(a+bx)}} dx \\
&= \frac{\sec(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} + \frac{\sec^3(a+bx)}{3b \csc^{\frac{3}{2}}(a+bx)} - \frac{1}{4} \left(\sqrt{\csc(a+bx)} \sqrt{\sin(a+bx)} \right) \int \sqrt{\sin(a+bx)} dx \\
&= \frac{\sec(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} + \frac{\sec^3(a+bx)}{3b \csc^{\frac{3}{2}}(a+bx)} - \frac{\sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a+bx)}}{2b}
\end{aligned}$$

Mathematica [A] time = 0.23, size = 76, normalized size = 0.83

$$\frac{\cos(a+bx)\sqrt{\csc(a+bx)} \left(2\sec^4(a+bx) + \sec^2(a+bx) + 3\sqrt{\sin(a+bx)} \sec(a+bx) E\left(\frac{1}{4}(-2a-2bx+\pi) \middle| 2\right) - 3 \right)}{6b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^4/Sqrt[Csc[a + b*x]], x]

[Out] (Cos[a + b*x]*Sqrt[Csc[a + b*x]]*(-3 + Sec[a + b*x]^2 + 2*Sec[a + b*x]^4 + 3*EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sec[a + b*x]*Sqrt[Sin[a + b*x]]))/(6*b)

fricas [F] time = 0.74, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sec(bx+a)^4}{\sqrt{\csc(bx+a)}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/csc(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^4/sqrt(csc(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(bx+a)^4}{\sqrt{\csc(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/csc(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^4/sqrt(csc(b*x + a)), x)

maple [A] time = 0.32, size = 160, normalized size = 1.74

$$\frac{6\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \text{EllipticE}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) (\cos^2(bx+a)) - 3\sqrt{\sin(bx+a)+1}}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^4/csc(b*x+a)^(1/2), x)

[Out] $1/12/\sin(b*x+a)^{(1/2)}/\cos(b*x+a)^3*(6*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)}*\text{EllipticE}((\sin(b*x+a)+1)^{(1/2)},1/2*2^{(1/2)})*\cos(b*x+a)^2-3*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)})*\text{EllipticF}((\sin(b*x+a)+1)^{(1/2)},1/2*2^{(1/2)})*\cos(b*x+a)^2-6*\cos(b*x+a)^4+2*\cos(b*x+a)^2+4)/b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(bx + a)}{\sqrt{\csc(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4/csc(b*x+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sec(b*x + a)^4/sqrt(csc(b*x + a)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(a + bx)^4 \sqrt{\frac{1}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(a + b*x)^4*(1/sin(a + b*x))^(1/2)),x)`

[Out] `int(1/(cos(a + b*x)^4*(1/sin(a + b*x))^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(a + bx)}{\sqrt{\csc(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**4/csc(b*x+a)**(1/2),x)`

[Out] `Integral(sec(a + b*x)**4/sqrt(csc(a + b*x)), x)`

3.279 $\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx$

Optimal. Leaf size=76

$$\frac{d\sqrt{d \cos(a + bx)} \csc^{p-1}(a + bx) {}_2F_1\left(-\frac{1}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a + bx)\right)}{b(1-p)\sqrt[4]{\cos^2(a + bx)}}$$

[Out] d*csc(b*x+a)^(-1+p)*hypergeom([-1/4, 1/2-1/2*p], [3/2-1/2*p], sin(b*x+a)^2)*(d*cos(b*x+a))^(1/2)/b/(1-p)/(cos(b*x+a)^2)^(1/4)

Rubi [A] time = 0.11, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2587, 2577}

$$\frac{d\sqrt{d \cos(a + bx)} \csc^{p-1}(a + bx) {}_2F_1\left(-\frac{1}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a + bx)\right)}{b(1-p)\sqrt[4]{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(d*Cos[a + b*x])^(3/2)*Csc[a + b*x]^p,x]

[Out] (d*Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[-1/4, (1 - p)/2, (3 - p)/2, Sin[a + b*x]^2])/(b*(1 - p)*(Cos[a + b*x]^2)^(1/4))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2])/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx &= (\csc^p(a + bx) \sin^p(a + bx)) \int (d \cos(a + bx))^{3/2} \sin^{-p}(a + bx) dx \\ &= \frac{d\sqrt{d \cos(a + bx)} \csc^{-1+p}(a + bx) {}_2F_1\left(-\frac{1}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a + bx)\right)}{b(1-p)\sqrt[4]{\cos^2(a + bx)}} \end{aligned}$$

Mathematica [A] time = 0.68, size = 105, normalized size = 1.38

$$\frac{2(d \cos(a + bx))^{5/2} \sin^2(a + bx)^{\frac{p-1}{2}} \csc^{p-1}(a + bx) \left(5 \cos^2(a + bx) {}_2F_1\left(\frac{9}{4}, \frac{p+1}{2}; \frac{13}{4}; \cos^2(a + bx)\right) + 9 {}_2F_1\left(\frac{5}{4}, \frac{p-1}{2}; \frac{9}{4}; \cos^2(a + bx)\right)\right)}{45bd}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(3/2)*Csc[a + b*x]^p,x]

[Out] $(-2*(d*\cos[a + b*x])^{5/2}*Csc[a + b*x]^{-1 + p}*(9*Hypergeometric2F1[5/4, (-1 + p)/2, 9/4, \cos[a + b*x]^2] + 5*\cos[a + b*x]^2*Hypergeometric2F1[9/4, (1 + p)/2, 13/4, \cos[a + b*x]^2])*(\sin[a + b*x]^2)^{((-1 + p)/2)})/(45*b*d)$

fricas [F] time = 0.56, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{d \cos(bx + a)} d \csc(bx + a)^p \cos(bx + a), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x, algorithm="fricas")`

[Out] `integral(sqrt(d*cos(b*x + a))*d*csc(b*x + a)^p*cos(b*x + a), x)`

giac [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x, algorithm="giac")`

[Out] Timed out

maple [F] time = 0.22, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^{\frac{3}{2}} (\csc^p(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x)`

[Out] `int((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^{\frac{3}{2}} \csc^p(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x, algorithm="maxima")`

[Out] `integrate((d*cos(b*x + a))^(3/2)*csc(b*x + a)^p, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + bx))^{3/2} \left(\frac{1}{\sin(a + bx)}\right)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(a + b*x))^(3/2)*(1/sin(a + b*x))^p,x)`

[Out] `int((d*cos(a + b*x))^(3/2)*(1/sin(a + b*x))^p, x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(3/2)*csc(b*x+a)**p,x)`

[Out] Timed out

3.280 $\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx$

Optimal. Leaf size=76

$$\frac{d\sqrt[4]{\cos^2(a + bx)} \csc^{p-1}(a + bx) {}_2F_1\left(\frac{1}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a + bx)\right)}{b(1-p)\sqrt{d \cos(a + bx)}}$$

[Out] d*(cos(b*x+a)^2)^(1/4)*csc(b*x+a)^(-1+p)*hypergeom([1/4, 1/2-1/2*p], [3/2-1/2*p], sin(b*x+a)^2)/b/(1-p)/(d*cos(b*x+a))^(1/2)

Rubi [A] time = 0.09, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2587, 2577}

$$\frac{d\sqrt[4]{\cos^2(a + bx)} \csc^{p-1}(a + bx) {}_2F_1\left(\frac{1}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a + bx)\right)}{b(1-p)\sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^p,x]

[Out] (d*(Cos[a + b*x]^2)^(1/4)*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[1/4, (1 - p)/2, (3 - p)/2, Sin[a + b*x]^2])/(b*(1 - p)*Sqrt[d*Cos[a + b*x]])

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2])/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx &= (\csc^p(a + bx) \sin^p(a + bx)) \int \sqrt{d \cos(a + bx)} \sin^{-p}(a + bx) dx \\ &= \frac{d\sqrt[4]{\cos^2(a + bx)} \csc^{-1+p}(a + bx) {}_2F_1\left(\frac{1}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a + bx)\right)}{b(1-p)\sqrt{d \cos(a + bx)}} \end{aligned}$$

Mathematica [A] time = 0.17, size = 70, normalized size = 0.92

$$\frac{2(d \cos(a + bx))^{3/2} \sin^2(a + bx)^{\frac{p-1}{2}} \csc^{p-1}(a + bx) {}_2F_1\left(\frac{3}{4}, \frac{p+1}{2}; \frac{7}{4}; \cos^2(a + bx)\right)}{3bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^p,x]

[Out] $(-2*(d*\cos[a + b*x])^{(3/2)}*\csc[a + b*x]^{(-1 + p)}*\text{Hypergeometric2F1}[3/4, (1 + p)/2, 7/4, \cos[a + b*x]^2]*(\sin[a + b*x]^2)^{((-1 + p)/2)})/(3*b*d)$

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}(\sqrt{d \cos(bx + a)} \csc(bx + a)^p, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x, algorithm="fricas")`

[Out] `integral(sqrt(d*cos(b*x + a))*csc(b*x + a)^p, x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(bx + a)} \csc(bx + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x, algorithm="giac")`

[Out] `integrate(sqrt(d*cos(b*x + a))*csc(b*x + a)^p, x)`

maple [F] time = 0.14, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(bx + a)} (\csc^p(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x)`

[Out] `int((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(bx + a)} \csc(bx + a)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x, algorithm="maxima")`

[Out] `integrate(sqrt(d*cos(b*x + a))*csc(b*x + a)^p, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{d \cos(a + bx)} \left(\frac{1}{\sin(a + bx)} \right)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(a + b*x))^(1/2)*(1/sin(a + b*x))^p,x)`

[Out] `int((d*cos(a + b*x))^(1/2)*(1/sin(a + b*x))^p, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(1/2)*csc(b*x+a)**p,x)`

[Out] `Integral(sqrt(d*cos(a + b*x))*csc(a + b*x)**p, x)`

$$3.281 \quad \int \frac{\csc^p(a+bx)}{\sqrt{d} \cos(a+bx)} dx$$

Optimal. Leaf size=76

$$\frac{d \cos^2(a+bx)^{3/4} \csc^{p-1}(a+bx) {}_2F_1\left(\frac{3}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{b(1-p)(d \cos(a+bx))^{3/2}}$$

[Out] d*(cos(b*x+a)^2)^(3/4)*csc(b*x+a)^(-1+p)*hypergeom([3/4, 1/2-1/2*p], [3/2-1/2*p], sin(b*x+a)^2)/b/(1-p)/(d*cos(b*x+a))^(3/2)

Rubi [A] time = 0.10, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2587, 2577}

$$\frac{d \cos^2(a+bx)^{3/4} \csc^{p-1}(a+bx) {}_2F_1\left(\frac{3}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{b(1-p)(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^p/Sqrt[d*Cos[a + b*x]], x]

[Out] (d*(Cos[a + b*x]^2)^(3/4)*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[3/4, (1 - p)/2, (3 - p)/2, Sin[a + b*x]^2])/(b*(1 - p)*(d*Cos[a + b*x])^(3/2))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n]*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2])/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^n]*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{\csc^p(a+bx)}{\sqrt{d} \cos(a+bx)} dx &= (\csc^p(a+bx) \sin^p(a+bx)) \int \frac{\sin^{-p}(a+bx)}{\sqrt{d} \cos(a+bx)} dx \\ &= \frac{d \cos^2(a+bx)^{3/4} \csc^{-1+p}(a+bx) {}_2F_1\left(\frac{3}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{b(1-p)(d \cos(a+bx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.22, size = 68, normalized size = 0.89

$$\frac{2\sqrt{d} \cos(a+bx) \sin^2(a+bx)^{\frac{p+1}{2}} \csc^{p+1}(a+bx) {}_2F_1\left(\frac{1}{4}, \frac{p+1}{2}; \frac{5}{4}; \cos^2(a+bx)\right)}{bd}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^p/Sqrt[d*Cos[a + b*x]], x]

[Out] $(-2\sqrt{d\cos[a + b*x]}) * \text{Csc}[a + b*x]^{(1 + p)} * \text{Hypergeometric2F1}[1/4, (1 + p)/2, 5/4, \cos[a + b*x]^2 * (\sin[a + b*x]^2)^{((1 + p)/2)}] / (b*d)$

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \cos (bx + a)} \csc (bx + a)^p}{d \cos (bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")`

[Out] `integral(sqrt(d*cos(b*x + a))*csc(b*x + a)^p/(d*cos(b*x + a)), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc (bx + a)^p}{\sqrt{d \cos (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2),x, algorithm="giac")`

[Out] `integrate(csc(b*x + a)^p/sqrt(d*cos(b*x + a)), x)`

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int \frac{\csc^p (bx + a)}{\sqrt{d \cos (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2),x)`

[Out] `int(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc (bx + a)^p}{\sqrt{d \cos (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")`

[Out] `integrate(csc(b*x + a)^p/sqrt(d*cos(b*x + a)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\sin(a+bx)}\right)^p}{\sqrt{d \cos (a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(1/2),x)`

[Out] `int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(1/2),x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^p (a + bx)}{\sqrt{d \cos (a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)**p/(d*cos(b*x+a))**(1/2),x)`

[Out] `Integral(csc(a + b*x)**p/sqrt(d*cos(a + b*x)), x)`

$$3.282 \quad \int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=78

$$\frac{\sqrt[4]{\cos^2(a+bx)} \csc^{p-1}(a+bx) {}_2F_1\left(\frac{5}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{bd(1-p)\sqrt{d \cos(a+bx)}}$$

[Out] (cos(b*x+a)^2)^(1/4)*csc(b*x+a)^(-1+p)*hypergeom([5/4, 1/2-1/2*p], [3/2-1/2*p], sin(b*x+a)^2)/b/d/(1-p)/(d*cos(b*x+a))^(1/2)

Rubi [A] time = 0.11, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2587, 2577}

$$\frac{\sqrt[4]{\cos^2(a+bx)} \csc^{p-1}(a+bx) {}_2F_1\left(\frac{5}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{bd(1-p)\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^p/(d*Cos[a + b*x])^(3/2), x]

[Out] ((Cos[a + b*x]^2)^(1/4)*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[5/4, (1 - p)/2, (3 - p)/2, Sin[a + b*x]^2])/(b*d*(1 - p)*Sqrt[d*Cos[a + b*x]])

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2])/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{3/2}} dx &= (\csc^p(a+bx) \sin^p(a+bx)) \int \frac{\sin^{-p}(a+bx)}{(d \cos(a+bx))^{3/2}} dx \\ &= \frac{\sqrt[4]{\cos^2(a+bx)} \csc^{-1+p}(a+bx) {}_2F_1\left(\frac{5}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{bd(1-p)\sqrt{d \cos(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.26, size = 68, normalized size = 0.87

$$\frac{2 \sin^2(a+bx)^{\frac{p-1}{2}} \csc^{p-1}(a+bx) {}_2F_1\left(-\frac{1}{4}, \frac{p+1}{2}; \frac{3}{4}; \cos^2(a+bx)\right)}{bd\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^p/(d*Cos[a + b*x])^(3/2), x]

[Out] (2*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[-1/4, (1 + p)/2, 3/4, Cos[a + b*x]^2]*(Sin[a + b*x]^2)^((-1 + p)/2))/(b*d*Sqrt[d*Cos[a + b*x]])

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \cos(bx + a)} \csc(bx + a)^p}{d^2 \cos(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*csc(b*x + a)^p/(d^2*cos(b*x + a)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(bx + a)^p}{(d \cos(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2), x, algorithm="giac")

[Out] integrate(csc(b*x + a)^p/(d*cos(b*x + a))^(3/2), x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int \frac{\csc^p(bx + a)}{(d \cos(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2), x)

[Out] int(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(bx + a)^p}{(d \cos(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2), x, algorithm="maxima")

[Out] integrate(csc(b*x + a)^p/(d*cos(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\sin(a+bx)}\right)^p}{(d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(3/2), x)

[Out] int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)**p/(d*cos(b*x+a))**(3/2), x)

[Out] Integral(csc(a + b*x)**p/(d*cos(a + b*x))**(3/2), x)

$$3.283 \quad \int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=78

$$\frac{\cos^2(a+bx)^{3/4} \csc^{p-1}(a+bx) {}_2F_1\left(\frac{7}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{bd(1-p)(d \cos(a+bx))^{3/2}}$$

[Out] $(\cos(b*x+a)^2)^{(3/4)} * \csc(b*x+a)^{(-1+p)} * \text{hypergeom}([7/4, 1/2-1/2*p], [3/2-1/2*p], \sin(b*x+a)^2) / b/d / (1-p) / (d*\cos(b*x+a))^{(3/2)}$

Rubi [A] time = 0.11, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2587, 2577}

$$\frac{\cos^2(a+bx)^{3/4} \csc^{p-1}(a+bx) {}_2F_1\left(\frac{7}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{bd(1-p)(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^p/(d*Cos[a + b*x])^(5/2), x]

[Out] $((\cos[a + b*x]^2)^{(3/4)} * \csc[a + b*x]^{(-1 + p)} * \text{Hypergeometric2F1}[7/4, (1 - p)/2, (3 - p)/2, \sin[a + b*x]^2]) / (b*d*(1 - p)*(d*\cos[a + b*x])^{(3/2)})$

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2]))*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2)]/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{5/2}} dx &= (\csc^p(a+bx) \sin^p(a+bx)) \int \frac{\sin^{-p}(a+bx)}{(d \cos(a+bx))^{5/2}} dx \\ &= \frac{\cos^2(a+bx)^{3/4} \csc^{-1+p}(a+bx) {}_2F_1\left(\frac{7}{4}, \frac{1-p}{2}; \frac{3-p}{2}; \sin^2(a+bx)\right)}{bd(1-p)(d \cos(a+bx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.28, size = 70, normalized size = 0.90

$$\frac{2 \sin^2(a+bx)^{\frac{p-1}{2}} \csc^{p-1}(a+bx) {}_2F_1\left(-\frac{3}{4}, \frac{p+1}{2}; \frac{1}{4}; \cos^2(a+bx)\right)}{3bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^p/(d*Cos[a + b*x])^(5/2), x]

[Out] $(2*\text{Csc}[a + b*x]^{(-1 + p)}*\text{Hypergeometric2F1}[-3/4, (1 + p)/2, 1/4, \text{Cos}[a + b*x]^2]*(\text{Sin}[a + b*x]^2)^{((-1 + p)/2)})/(3*b*d*(d*\text{Cos}[a + b*x])^{(3/2)})$

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \cos(bx + a)} \csc(bx + a)^p}{d^3 \cos(bx + a)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x, algorithm="fricas")`

[Out] `integral(sqrt(d*cos(b*x + a))*csc(b*x + a)^p/(d^3*cos(b*x + a)^3), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(bx + a)^p}{(d \cos(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x, algorithm="giac")`

[Out] `integrate(csc(b*x + a)^p/(d*cos(b*x + a))^(5/2), x)`

maple [F] time = 0.15, size = 0, normalized size = 0.00

$$\int \frac{\csc^p(bx + a)}{(d \cos(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x)`

[Out] `int(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(bx + a)^p}{(d \cos(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x, algorithm="maxima")`

[Out] `integrate(csc(b*x + a)^p/(d*cos(b*x + a))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\sin(a+bx)}\right)^p}{(d \cos(a + bx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(5/2), x)`

[Out] `int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)**p/(d*cos(b*x+a))**(5/2), x)`

[Out] Timed out

3.284 $\int \cos^m(e + fx) \csc^n(e + fx) dx$

Optimal. Leaf size=85

$$\frac{\cos^{m-1}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{n-1}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] $\cos(f*x+e)^{-1+m}*(\cos(f*x+e)^2)^{(1/2-1/2*m)}*\csc(f*x+e)^{-1+n}*\text{hypergeom}([1/2-1/2*m, 1/2-1/2*n], [3/2-1/2*n], \sin(f*x+e)^2)/f/(1-n)$

Rubi [A] time = 0.08, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2587, 2577}

$$\frac{\cos^{m-1}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{n-1}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^m*Csc[e + f*x]^n,x]

[Out] $(\text{Cos}[e + f*x]^{-1 + m}*(\text{Cos}[e + f*x]^2)^{((1 - m)/2)}*\text{Csc}[e + f*x]^{-1 + n}*\text{Hypergeometric2F1}[(1 - m)/2, (1 - n)/2, (3 - n)/2, \text{Sin}[e + f*x]^2])/f*(1 - n)$

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2]))*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2)]/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \cos^m(e + fx) \csc^n(e + fx) dx &= \left(\csc^n(e + fx) \sin^n(e + fx) \right) \int \cos^m(e + fx) \sin^{-n}(e + fx) dx \\ &= \frac{\cos^{-1+m}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{-1+n}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)} \end{aligned}$$

Mathematica [C] time = 1.97, size = 312, normalized size = 3.67

$$\frac{2(n-3) \sin\left(\frac{1}{2}(e + fx)\right) \cos^3\left(\frac{1}{2}(e + fx)\right)}{f(n-1) \left(2 \sin^2\left(\frac{1}{2}(e + fx)\right) \left(m F_1\left(\frac{3}{2} - \frac{n}{2}; 1 - m, m - n + 1; \frac{5}{2} - \frac{n}{2}; \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) + m\right)}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Cos[e + f*x]^m*Csc[e + f*x]^n,x]

[Out] $(-2*(-3 + n)*\text{AppellF1}[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]*\text{Cos}[(e + f*x)/2]^3*\text{Cos}[e + f*x]^m*\text{Csc}[e + f*x]^n*\text{Sin}[(e + f*x)/2]) / (f*(-1 + n)*((-3 + n)*\text{AppellF1}[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]*\text{Cos}[(e + f*x)/2]^2 + 2*(m*\text{AppellF1}[3/2 - n/2, 1 - m, 1 + m - n, 5/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + (1 + m - n)*\text{AppellF1}[3/2 - n/2, -m, 2 + m - n, 5/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]))*\text{Sin}[(e + f*x)/2]^2)$

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\cos(fx + e)^m \csc(fx + e)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^m*csc(f*x+e)^n,x, algorithm="fricas")

[Out] integral(cos(f*x + e)^m*csc(f*x + e)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(fx + e)^m \csc(fx + e)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^m*csc(f*x+e)^n,x, algorithm="giac")

[Out] integrate(cos(f*x + e)^m*csc(f*x + e)^n, x)

maple [F] time = 0.65, size = 0, normalized size = 0.00

$$\int (\cos^m(fx + e)) (\csc^n(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^m*csc(f*x+e)^n,x)

[Out] int(cos(f*x+e)^m*csc(f*x+e)^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(fx + e)^m \csc(fx + e)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^m*csc(f*x+e)^n,x, algorithm="maxima")

[Out] integrate(cos(f*x + e)^m*csc(f*x + e)^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(e + fx)^m \left(\frac{1}{\sin(e + fx)}\right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^m*(1/sin(e + f*x))^n,x)

[Out] int(cos(e + f*x)^m*(1/sin(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \cos^m(e + fx) \csc^n(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)**m*csc(f*x+e)**n,x)
```

```
[Out] Integral(cos(e + f*x)**m*csc(e + f*x)**n, x)
```

3.285 $\int (a \cos(e + fx))^m \csc^n(e + fx) dx$

Optimal. Leaf size=88

$$\frac{a \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{n-1}(e + fx) (a \cos(e + fx))^{m-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] a*(a*cos(f*x+e))^(-1+m)*(cos(f*x+e)²)^(1/2-1/2*m)*csc(f*x+e)⁽⁻¹⁺ⁿ⁾*hypergeometric([1/2-1/2*m, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)²)/f/(1-n)

Rubi [A] time = 0.09, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2587, 2577}

$$\frac{a \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{n-1}(e + fx) (a \cos(e + fx))^{m-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[e + f*x])^m*Csc[e + f*x]ⁿ, x]

[Out] (a*(a*Cos[e + f*x])^(-1 + m)*(Cos[e + f*x]²)^{((1 - m)/2)}*Csc[e + f*x]^(-1 + n)*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]²)]/(f*(1 - n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(b^{(2*IntPart[(n - 1)/2] + 1)}*(b*Cos[e + f*x])^{(2*FracPart[(n - 1)/2])}*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]²)]/(a*f*(m + 1)*(Cos[e + f*x]²)^{FracPart[(n - 1)/2]}), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Dist[b²*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])ⁿ, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx = \left(\csc^n(e + fx) \sin^n(e + fx) \right) \int (a \cos(e + fx))^m \sin^{-n}(e + fx) dx$$

$$= \frac{a(a \cos(e + fx))^{-1+m} \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{-1+n}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Mathematica [C] time = 0.23, size = 314, normalized size = 3.57

$$\frac{2(n-3) \sin\left(\frac{1}{2}(e + fx)\right) \cos^3\left(\frac{1}{2}(e + fx)\right)}{f(n-1) \left(2 \sin^2\left(\frac{1}{2}(e + fx)\right) \left({}_mF_1\left(\frac{3}{2} - \frac{n}{2}; 1 - m, m - n + 1; \frac{5}{2} - \frac{n}{2}; \tan^2\left(\frac{1}{2}(e + fx)\right)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) + (m - 3)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a*cos[e + f*x])^m*csc[e + f*x]^n,x]

[Out] $(-2*(-3 + n)*\text{AppellF1}[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]*\text{Cos}[(e + f*x)/2]^3*(a*\text{Cos}[e + f*x])^m*\text{Csc}[e + f*x]^n*\text{Sin}[(e + f*x)/2])/(f*(-1 + n)*((-3 + n)*\text{AppellF1}[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]*\text{Cos}[(e + f*x)/2]^2 + 2*(m*\text{AppellF1}[3/2 - n/2, 1 - m, 1 + m - n, 5/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + (1 + m - n)*\text{AppellF1}[3/2 - n/2, -m, 2 + m - n, 5/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2))*\text{Sin}[(e + f*x)/2]^2)$

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(a \cos (f x+e)\right)^m \csc (f x+e)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*csc(f*x+e)^n,x, algorithm="fricas")

[Out] integral((a*cos(f*x + e))^m*csc(f*x + e)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos (f x+e))^m \csc (f x+e)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*csc(f*x+e)^n,x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m*csc(f*x + e)^n, x)

maple [F] time = 0.56, size = 0, normalized size = 0.00

$$\int (a \cos (f x+e))^m \left(\csc ^n (f x+e)\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m*csc(f*x+e)^n,x)

[Out] int((a*cos(f*x+e))^m*csc(f*x+e)^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos (f x+e))^m \csc (f x+e)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*csc(f*x+e)^n,x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m*csc(f*x + e)^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos (e+f x))^m \left(\frac{1}{\sin (e+f x)}\right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(e + f*x))^m*(1/sin(e + f*x))^n,x)

[Out] int((a*cos(e + f*x))^m*(1/sin(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))**m*csc(f*x+e)**n,x)

[Out] Integral((a*cos(e + f*x))**m*csc(e + f*x)**n, x)

3.286 $\int \cos^m(e + fx)(b \csc(e + fx))^n dx$

Optimal. Leaf size=88

$$\frac{b \cos^{m-1}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] b*cos(f*x+e)^(-1+m)*(cos(f*x+e)^2)^(1/2-1/2*m)*(b*csc(f*x+e))^(-1+n)*hypergeom([1/2-1/2*m, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)/f/(1-n)

Rubi [A] time = 0.09, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2587, 2577}

$$\frac{b \cos^{m-1}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^m*(b*Csc[e + f*x])^n,x]

[Out] (b*Cos[e + f*x]^(-1 + m)*(Cos[e + f*x]^2)^((1 - m)/2)*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(f*(1 - n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n_*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2])/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^n_*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \cos^m(e + fx)(b \csc(e + fx))^n dx &= \left(b^2(b \csc(e + fx))^{-1+n}(b \sin(e + fx))^{-1+n}\right) \int \cos^m(e + fx)(b \sin(e + fx))^{-1+n} dx \\ &= \frac{b \cos^{-1+m}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)} \end{aligned}$$

Mathematica [C] time = 0.24, size = 314, normalized size = 3.57

$$\frac{2(n-3) \sin\left(\frac{1}{2}(e + fx)\right) \cos^3\left(\frac{1}{2}(e + fx)\right)}{f(n-1) \left(2 \sin^2\left(\frac{1}{2}(e + fx)\right) \left(m F_1\left(\frac{3}{2} - \frac{n}{2}; 1 - m, m - n + 1; \frac{5}{2} - \frac{n}{2}; \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) + m\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Cos[e + f*x]^m*(b*Csc[e + f*x])^n,x]

[Out] $(-2*(-3 + n)*\text{AppellF1}[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]*\text{Cos}[(e + f*x)/2]^3*\text{Cos}[e + f*x]^m*(b*\text{Csc}[e + f*x])^n*\text{Sin}[(e + f*x)/2]) / (f*(-1 + n)*((-3 + n)*\text{AppellF1}[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]*\text{Cos}[(e + f*x)/2]^2 + 2*(m*\text{AppellF1}[3/2 - n/2, 1 - m, 1 + m - n, 5/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + (1 + m - n)*\text{AppellF1}[3/2 - n/2, -m, 2 + m - n, 5/2 - n/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2))*\text{Sin}[(e + f*x)/2]^2)$

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \cos(fx + e)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^m*(b*csc(f*x+e))^n,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*cos(f*x + e)^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^m*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*cos(f*x + e)^m, x)

maple [F] time = 0.63, size = 0, normalized size = 0.00

$$\int (\cos^m(fx + e))(b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^m*(b*csc(f*x+e))^n,x)

[Out] int(cos(f*x+e)^m*(b*csc(f*x+e))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^m*(b*csc(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^n*cos(f*x + e)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(e + fx)^m \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^m*(b/sin(e + f*x))^n,x)

[Out] int(cos(e + f*x)^m*(b/sin(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \cos^m(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**m*(b*csc(f*x+e))**n,x)

[Out] Integral((b*csc(e + f*x))**n*cos(e + f*x)**m, x)

3.287 $\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx$

Optimal. Leaf size=91

$$\frac{ab \cos^2(e + fx)^{\frac{1-m}{2}} (a \cos(e + fx))^{m-1} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] a*b*(a*cos(f*x+e))^(-1+m)*(cos(f*x+e)^2)^(1/2-1/2*m)*(b*csc(f*x+e))^(-1+n)*
hypergeom([1/2-1/2*m, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)/f/(1-n)

Rubi [A] time = 0.10, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2587, 2577}

$$\frac{ab \cos^2(e + fx)^{\frac{1-m}{2}} (a \cos(e + fx))^{m-1} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[e + f*x])^m*(b*Csc[e + f*x])^n,x]

[Out] (a*b*(a*Cos[e + f*x])^(-1 + m)*(Cos[e + f*x]^2)^((1 - m)/2)*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(f*(1 - n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2])/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^n*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx = (b^2 (b \csc(e + fx))^{-1+n} (b \sin(e + fx))^{-1+n}) \int (a \cos(e + fx))^m (b \sin(e + fx))^{-1+n} dx$$

$$= \frac{ab (a \cos(e + fx))^{-1+m} \cos^2(e + fx)^{\frac{1-m}{2}} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Mathematica [C] time = 0.23, size = 316, normalized size = 3.47

$$\frac{2(n-3) \sin\left(\frac{1}{2}(e+fx)\right) \cos^3\left(\frac{1}{2}(e+fx)\right)}{f(n-1) \left(2 \sin^2\left(\frac{1}{2}(e+fx)\right) \left({}_mF_1\left(\frac{3}{2} - \frac{n}{2}; 1-m, m-n+1; \frac{5}{2} - \frac{n}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) + (m-1) \tan^2\left(\frac{1}{2}(e+fx)\right)\right)} + (m-1) \tan^2\left(\frac{1}{2}(e+fx)\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a*cos[e + f*x])^m*(b*csc[e + f*x])^n,x]

[Out] (-2*(-3 + n)*AppellF1[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*(a*cos[e + f*x])^m*(b*csc[e + f*x])^n*Sin[(e + f*x)/2])/(f*(-1 + n)*((-3 + n)*AppellF1[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2 + 2*(m*AppellF1[3/2 - n/2, 1 - m, 1 + m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + (1 + m - n)*AppellF1[3/2 - n/2, -m, 2 + m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Sin[(e + f*x)/2]^2))

fricas [F] time = 0.85, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(a \cos(fx + e)\right)^m \left(b \csc(fx + e)\right)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^n,x, algorithm="fricas")

[Out] integral((a*cos(f*x + e))^m*(b*csc(f*x + e))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m*(b*csc(f*x + e))^n, x)

maple [F] time = 0.66, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m*(b*csc(f*x+e))^n,x)

[Out] int((a*cos(f*x+e))^m*(b*csc(f*x+e))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m*(b*csc(f*x + e))^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(e + fx))^m \left(\frac{b}{\sin(e + fx)}\right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^n,x)

[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))**m*(b*csc(f*x+e))**n,x)

[Out] Integral((a*cos(e + f*x))**m*(b*csc(e + f*x))**n, x)

3.288 $\int (a \cos(e + fx))^m (b \csc(e + fx))^{7/2} dx$

Optimal. Leaf size=78

$$\frac{b^3 \sqrt[4]{\sin^2(e + fx)} \sqrt{b \csc(e + fx)} (a \cos(e + fx))^{m+1} {}_2F_1\left(\frac{9}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e + fx)\right)}{af(m+1)}$$

[Out] $-b^3(a \cos(fx+e))^{(1+m)} \text{hypergeom}\left(\left[\frac{9}{4}, \frac{1}{2} + \frac{1}{2}m\right], \left[\frac{3}{2} + \frac{1}{2}m\right], \cos(fx+e)^2\right) (\sin(fx+e)^2)^{(1/4)} (b \csc(fx+e))^{(1/2)} / a/f/(1+m)$

Rubi [A] time = 0.12, antiderivative size = 76, normalized size of antiderivative = 0.97, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2587, 2576}

$$\frac{b \sin^2(e + fx)^{5/4} (b \csc(e + fx))^{5/2} (a \cos(e + fx))^{m+1} {}_2F_1\left(\frac{9}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e + fx)\right)}{af(m+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a \cos[e + fx])^m (b \csc[e + fx])^{(7/2)}, x]$

[Out] $-\left((b (a \cos[e + fx])^{(1+m)} (b \csc[e + fx])^{(5/2)} \text{Hypergeometric2F1}\left[\frac{9}{4}, \frac{(1+m)}{2}, \frac{(3+m)}{2}, \cos[e + fx]^2\right] (\sin[e + fx]^2)^{(5/4)}\right) / (a f (1+m))\right)$

Rule 2576

$\text{Int}[(\cos[(e_.) + (f_.) (x_.)]) (a_.)^{(m_.)} ((b_.) \sin[(e_.) + (f_.) (x_.)])^{(n_.)}, x_Symbol] :> -\text{Simp}[b^{(2 \text{IntPart}[(n-1)/2] + 1)} (b \sin[e + fx])^{(2 \text{FracPart}[(n-1)/2])} (a \cos[e + fx])^{(m+1)} \text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \cos[e + fx]^2)] / (a f (m+1) (\sin[e + fx]^2)^{\text{FracPart}[(n-1)/2]}), x] /;$ FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2587

$\text{Int}[(b_.) \sec[(e_.) + (f_.) (x_.)])^{(n_.)} (a_.) \sin[(e_.) + (f_.) (x_.)])^{(m_.)}, x_Symbol] :> \text{Dist}[b^{2(n-1)} (b \cos[e + fx])^{(n-1)} (b \sec[e + fx])^{(n-1)}, \text{Int}[(a \sin[e + fx])^m / (b \cos[e + fx])^n, x], x] /;$ FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (a \cos(e + fx))^m (b \csc(e + fx))^{7/2} dx &= (b^2 (b \csc(e + fx))^{5/2} (b \sin(e + fx))^{5/2}) \int \frac{(a \cos(e + fx))^m}{(b \sin(e + fx))^{7/2}} dx \\ &= -\frac{b (a \cos(e + fx))^{1+m} (b \csc(e + fx))^{5/2} {}_2F_1\left(\frac{9}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \cos^2(e + fx)\right)}{af(1+m)} \end{aligned}$$

Mathematica [A] time = 7.91, size = 94, normalized size = 1.21

$$\frac{2ab (b \csc(e + fx))^{5/2} (-\cot^2(e + fx))^{\frac{1-m}{2}} (a \cos(e + fx))^{m-1} {}_2F_1\left(\frac{1}{4}(7-2m), \frac{1-m}{2}; \frac{1}{4}(11-2m); \csc^2(e + fx)\right)}{f(2m-7)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[e + f*x])^m*(b*Csc[e + f*x])^(7/2),x]

[Out] (2*a*b*(a*cos[e + f*x])^(-1 + m)*(-Cot[e + f*x]^2)^((1 - m)/2)*(b*Csc[e + f*x])^(5/2)*Hypergeometric2F1[(7 - 2*m)/4, (1 - m)/2, (11 - 2*m)/4, Csc[e + f*x]^2])/(f*(-7 + 2*m))

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \csc(fx + e)} (a \cos(fx + e))^m b^3 \csc(fx + e)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m*b^3*csc(f*x + e)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^{\frac{7}{2}} (a \cos(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^(7/2)*(a*cos(f*x + e))^m, x)

maple [F] time = 0.17, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \csc(fx + e))^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x)

[Out] int((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^{\frac{7}{2}} (a \cos(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^(7/2)*(a*cos(f*x + e))^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(e + fx))^m \left(\frac{b}{\sin(e + fx)} \right)^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(7/2),x)

[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x)

[Out] Timed out

3.289 $\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx$

Optimal. Leaf size=76

$$\frac{b \sin^2(e + fx)^{3/4} (b \csc(e + fx))^{3/2} (a \cos(e + fx))^{m+1} {}_2F_1\left(\frac{7}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e + fx)\right)}{af(m+1)}$$

[Out] -b*(a*cos(f*x+e))^(1+m)*(b*csc(f*x+e))^(3/2)*hypergeom([7/4, 1/2+1/2*m], [3/2+1/2*m], cos(f*x+e)^2)*(sin(f*x+e)^2)^(3/4)/a/f/(1+m)

Rubi [A] time = 0.12, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2587, 2576}

$$\frac{b \sin^2(e + fx)^{3/4} (b \csc(e + fx))^{3/2} (a \cos(e + fx))^{m+1} {}_2F_1\left(\frac{7}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e + fx)\right)}{af(m+1)}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[e + f*x])^m*(b*Csc[e + f*x])^(5/2), x]

[Out] -((b*(a*Cos[e + f*x])^(1 + m)*(b*Csc[e + f*x])^(3/2)*Hypergeometric2F1[7/4, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2]*(Sin[e + f*x]^2)^(3/4))/(a*f*(1 + m))

Rule 2576

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> -Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Cos[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2])/(a*f*(m + 1)*(Sin[e + f*x]^2)^(FracPart[(n - 1)/2])), x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx &= (b^2 (b \csc(e + fx))^{3/2} (b \sin(e + fx))^{3/2}) \int \frac{(a \cos(e + fx))^m}{(b \sin(e + fx))^{5/2}} dx \\ &= -\frac{b(a \cos(e + fx))^{1+m} (b \csc(e + fx))^{3/2} {}_2F_1\left(\frac{7}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \cos^2(e + fx)\right)}{af(1+m)} \end{aligned}$$

Mathematica [A] time = 2.39, size = 94, normalized size = 1.24

$$\frac{2ab(b \csc(e + fx))^{3/2} \left(-\cot^2(e + fx)\right)^{\frac{1-m}{2}} (a \cos(e + fx))^{m-1} {}_2F_1\left(\frac{1}{4}(5 - 2m), \frac{1-m}{2}; \frac{1}{4}(9 - 2m); \csc^2(e + fx)\right)}{f(2m - 5)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[e + f*x])^m*(b*csc[e + f*x])^(5/2),x]

[Out] (2*a*b*(a*cos[e + f*x])^(-1 + m)*(-Cot[e + f*x]^2)^((1 - m)/2)*(b*csc[e + f*x])^(3/2)*Hypergeometric2F1[(5 - 2*m)/4, (1 - m)/2, (9 - 2*m)/4, Csc[e + f*x]^2])/(f*(-5 + 2*m))

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \csc(fx + e)} (a \cos(fx + e))^m b^2 \csc(fx + e)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m*b^2*csc(f*x + e)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^{\frac{5}{2}} (a \cos(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^(5/2)*(a*cos(f*x + e))^m, x)

maple [F] time = 0.17, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \csc(fx + e))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x)

[Out] int((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^{\frac{5}{2}} (a \cos(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^(5/2)*(a*cos(f*x + e))^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(e + fx))^m \left(\frac{b}{\sin(e + fx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(5/2),x)

[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x)

[Out] Timed out

3.290 $\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx$

Optimal. Leaf size=76

$$\frac{b^4 \sqrt{\sin^2(e + fx)} \sqrt{b \csc(e + fx)} (a \cos(e + fx))^{m+1} {}_2F_1\left(\frac{5}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e + fx)\right)}{af(m+1)}$$

[Out] $-b*(a*\cos(f*x+e))^{(1+m)}*\text{hypergeom}\left(\left[\frac{5}{4}, \frac{1}{2}+\frac{1}{2}*m\right], \left[\frac{3}{2}+\frac{1}{2}*m\right], \cos(f*x+e)^2\right)*(\sin(f*x+e)^2)^{(1/4)}*(b*\csc(f*x+e))^{(1/2)}/a/f/(1+m)$

Rubi [A] time = 0.11, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2587, 2576}

$$\frac{b^4 \sqrt{\sin^2(e + fx)} \sqrt{b \csc(e + fx)} (a \cos(e + fx))^{m+1} {}_2F_1\left(\frac{5}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e + fx)\right)}{af(m+1)}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[e + f*x])^m*(b*Csc[e + f*x])^(3/2), x]

[Out] $-((b*(a*\cos[e + f*x])^{(1 + m)}*\text{Sqrt}[b*\csc[e + f*x]]*\text{Hypergeometric2F1}\left[\frac{5}{4}, (1 + m)/2, (3 + m)/2, \cos[e + f*x]^2\right]*(\sin[e + f*x]^2)^{(1/4)}))/(a*f*(1 + m))$

Rule 2576

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> -Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Sine[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Cos[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2])/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2587

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx &= (b^2 \sqrt{b \csc(e + fx)} \sqrt{b \sin(e + fx)}) \int \frac{(a \cos(e + fx))^m}{(b \sin(e + fx))^{3/2}} dx \\ &= -\frac{b(a \cos(e + fx))^{1+m} \sqrt{b \csc(e + fx)} {}_2F_1\left(\frac{5}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \cos^2(e + fx)\right)}{af(1+m)} \end{aligned}$$

Mathematica [A] time = 1.24, size = 94, normalized size = 1.24

$$\frac{2ab \sqrt{b \csc(e + fx)} (-\cot^2(e + fx))^{\frac{1-m}{2}} (a \cos(e + fx))^{m-1} {}_2F_1\left(\frac{1}{4}(3-2m), \frac{1-m}{2}; \frac{1}{4}(7-2m); \csc^2(e + fx)\right)}{f(2m-3)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Cos[e + f*x])^m*(b*Csc[e + f*x])^(3/2), x]

[Out] $(2*a*b*(a*\cos[e + f*x])^{-1 + m}*(-\cot[e + f*x]^2)^{((1 - m)/2)*\sqrt{b*\csc[e + f*x]}}*\text{Hypergeometric2F1}[(3 - 2*m)/4, (1 - m)/2, (7 - 2*m)/4, \csc[e + f*x]^2])/(f*(-3 + 2*m))$

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \csc(fx + e)} (a \cos(fx + e))^m b \csc(fx + e), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(3/2),x, algorithm="fricas")`

[Out] `integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m*b*csc(f*x + e), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^{\frac{3}{2}} (a \cos(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(3/2),x, algorithm="giac")`

[Out] `integrate((b*csc(f*x + e))^(3/2)*(a*cos(f*x + e))^m, x)`

maple [F] time = 0.16, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m (b \csc(fx + e))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*cos(f*x+e))^m*(b*csc(f*x+e))^(3/2),x)`

[Out] `int((a*cos(f*x+e))^m*(b*csc(f*x+e))^(3/2),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^{\frac{3}{2}} (a \cos(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(3/2),x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^(3/2)*(a*cos(f*x + e))^m, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(e + fx))^m \left(\frac{b}{\sin(e + fx)}\right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(3/2),x)`

[Out] `int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(3/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(f*x+e))**m*(b*csc(f*x+e))**(3/2),x)`

[Out] Timed out

3.291 $\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx$

Optimal. Leaf size=78

$$\frac{\sin^2(e + fx)^{3/4} (b \csc(e + fx))^{3/2} (a \cos(e + fx))^{m+1} {}_2F_1\left(\frac{3}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e + fx)\right)}{abf(m+1)}$$

[Out] $-(a \cos(fx+e))^{(1+m)} (b \csc(fx+e))^{(3/2)} \text{hypergeom}([3/4, 1/2+1/2*m], [3/2+1/2*m], \cos(fx+e)^2) * (\sin(fx+e)^2)^{(3/4)} / a/b/f/(1+m)$

Rubi [A] time = 0.10, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2586, 2576}

$$\frac{\sin^2(e + fx)^{3/4} (b \csc(e + fx))^{3/2} (a \cos(e + fx))^{m+1} {}_2F_1\left(\frac{3}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e + fx)\right)}{abf(m+1)}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[e + f*x])^m*Sqrt[b*Csc[e + f*x]],x]

[Out] $-\left(\left((a \cos[e + f*x])^{(1+m)} (b \csc[e + f*x])^{(3/2)} \text{Hypergeometric2F1}\left[\frac{3}{4}, (1+m)/2, (3+m)/2, \cos[e + f*x]^2\right] * (\sin[e + f*x]^2)^{(3/4)}\right) / (a*b*f*(1+m))\right)$

Rule 2576

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> -Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Sine[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Cos[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2])/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2586

Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[(1*(b*Cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1))/b^2, Int[(a*Sine[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]

Rubi steps

$$\begin{aligned} \int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx &= \frac{\left((b \csc(e + fx))^{3/2} (b \sin(e + fx))^{3/2}\right) \int \frac{(a \cos(e + fx))^m}{\sqrt{b \sin(e + fx)}} dx}{b^2} \\ &= -\frac{(a \cos(e + fx))^{1+m} (b \csc(e + fx))^{3/2} {}_2F_1\left(\frac{3}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \cos^2(e + fx)\right)}{abf(1+m)} \end{aligned}$$

Mathematica [A] time = 1.05, size = 96, normalized size = 1.23

$$\frac{2 \tan(e + fx) \sqrt{b \csc(e + fx)} \left(-\cot^2(e + fx)\right)^{\frac{1-m}{2}} (a \cos(e + fx))^m {}_2F_1\left(\frac{1}{4}(1 - 2m), \frac{1-m}{2}; \frac{1}{4}(5 - 2m); \csc^2(e + fx)\right)}{f(2m - 1)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[e + f*x])^m*Sqrt[b*Csc[e + f*x]],x]

[Out] (2*(a*cos[e + f*x])^m*(-Cot[e + f*x]^2)^((1 - m)/2)*Sqrt[b*Csc[e + f*x]]*Hypergeometric2F1[(1 - 2*m)/4, (1 - m)/2, (5 - 2*m)/4, Csc[e + f*x]^2]*Tan[e + f*x])/(f*(-1 + 2*m))

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \csc(fx + e)} (a \cos(fx + e))^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \csc(fx + e)} (a \cos(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m, x)

maple [F] time = 0.16, size = 0, normalized size = 0.00

$$\int (a \cos(fx + e))^m \sqrt{b \csc(fx + e)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x)

[Out] int((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \csc(fx + e)} (a \cos(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int (a \cos(e + fx))^m \sqrt{\frac{b}{\sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(1/2),x)

[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x)

[Out] Integral((a*cos(e + f*x))^m*sqrt(b*csc(e + f*x)), x)

$$3.292 \quad \int \frac{(a \cos(e+fx))^m}{\sqrt{b \csc(e+fx)}} dx$$

Optimal. Leaf size=78

$$\frac{\sqrt[4]{\sin^2(e+fx)} \sqrt{b \csc(e+fx)} (a \cos(e+fx))^{m+1} {}_2F_1\left(\frac{1}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e+fx)\right)}{abf(m+1)}$$

[Out] $-(a \cos(fx+e))^{(1+m)} \operatorname{hypergeom}\left(\left[\frac{1}{4}, \frac{1}{2} + \frac{1}{2}m\right], \left[\frac{3}{2} + \frac{1}{2}m\right], \cos(fx+e)^2\right) (\sin(fx+e)^2)^{(1/4)} (b \csc(fx+e))^{(1/2)} / a/b/f/(1+m)$

Rubi [A] time = 0.10, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2586, 2576}

$$\frac{\sqrt[4]{\sin^2(e+fx)} \sqrt{b \csc(e+fx)} (a \cos(e+fx))^{m+1} {}_2F_1\left(\frac{1}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e+fx)\right)}{abf(m+1)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a \cos[e+fx])^m / \operatorname{Sqrt}[b \csc[e+fx]], x]$

[Out] $-\left(\left(a \cos[e+fx]\right)^{(1+m)} \operatorname{Sqrt}[b \csc[e+fx]] \operatorname{Hypergeometric2F1}\left[\frac{1}{4}, (1+m)/2, (3+m)/2, \cos[e+fx]^2\right] \left(\sin[e+fx]^2\right)^{(1/4)}\right) / (a b f (1+m))$

Rule 2576

$\operatorname{Int}[(\cos[e_.] + (f_.) (x_.) (a_.)^m) ((b_.) \sin[e_.] + (f_.) (x_.)^n), x_Symbol] \rightarrow -\operatorname{Simp}[(b^{(2 \operatorname{IntPart}[(n-1)/2] + 1)} (b \sin[e+fx])^{(2 \operatorname{FracPart}[(n-1)/2])} (a \cos[e+fx])^{(m+1)} \operatorname{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \cos[e+fx]^2]) / (a f (m+1) (\sin[e+fx]^2)^{\operatorname{FracPart}[(n-1)/2]}), x] /; \operatorname{FreeQ}\{a, b, e, f, m, n\}, x\} \&\& \operatorname{SimplerQ}[n, m]$

Rule 2586

$\operatorname{Int}[(b_.) \sec[e_.] + (f_.) (x_.)^n) ((a_.) \sin[e_.] + (f_.) (x_.)^m), x_Symbol] \rightarrow \operatorname{Dist}[(1 (b \cos[e+fx])^{(n+1)} (b \sec[e+fx])^{(n+1)}) / b^2, \operatorname{Int}[(a \sin[e+fx])^m / (b \cos[e+fx])^n, x], x] /; \operatorname{FreeQ}\{a, b, e, f, m, n\}, x\} \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{LtQ}[n, 1]$

Rubi steps

$$\begin{aligned} \int \frac{(a \cos(e+fx))^m}{\sqrt{b \csc(e+fx)}} dx &= \frac{(\sqrt{b \csc(e+fx)} \sqrt{b \sin(e+fx)})}{b^2} \int (a \cos(e+fx))^m \sqrt{b \sin(e+fx)} dx \\ &= -\frac{(a \cos(e+fx))^{1+m} \sqrt{b \csc(e+fx)} {}_2F_1\left(\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \cos^2(e+fx)\right) \sqrt[4]{\sin^2(e+fx)}}{abf(1+m)} \end{aligned}$$

Mathematica [C] time = 1.64, size = 225, normalized size = 2.88

$$\frac{14b(a \cos(e+fx))^m {}_2F_1\left(\frac{3}{4}; -m\right)}{3f(b \csc(e+fx))^{3/2} \left(7 {}_2F_1\left(\frac{3}{4}; -m, m + \frac{3}{2}; \frac{7}{4}; \tan^2\left(\frac{1}{2}(e+fx)\right)\right) - \tan^2\left(\frac{1}{2}(e+fx)\right)\right) - 2 \tan^2\left(\frac{1}{2}(e+fx)\right) \left(2m {}_2F_1\left(\frac{3}{4}; -m\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(a*cos[e + f*x])^m/Sqrt[b*Csc[e + f*x]],x]

[Out] (14*b*AppellF1[3/4, -m, 3/2 + m, 7/4, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(a*cos[e + f*x])^m)/(3*f*(b*Csc[e + f*x])^(3/2)*(7*AppellF1[3/4, -m, 3/2 + m, 7/4, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - 2*(2*m*AppellF1[7/4, 1 - m, 3/2 + m, 11/4, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + (3 + 2*m)*AppellF1[7/4, -m, 5/2 + m, 11/4, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))

fricas [F] time = 0.84, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \csc(fx + e)} (a \cos(fx + e))^m}{b \csc(fx + e)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m/(b*csc(f*x + e)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{\sqrt{b \csc(fx + e)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m/sqrt(b*csc(f*x + e)), x)

maple [F] time = 0.14, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{\sqrt{b \csc(fx + e)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m/(b*csc(f*x+e))^(1/2),x)

[Out] int((a*cos(f*x+e))^m/(b*csc(f*x+e))^(1/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{\sqrt{b \csc(fx + e)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m/sqrt(b*csc(f*x + e)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(a \cos(e + fx))^m}{\sqrt{\frac{b}{\sin(e+fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(1/2), x)`

[Out] `int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(e + fx))^m}{\sqrt{b \csc(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(f*x+e))**m/(b*csc(f*x+e))**(1/2), x)`

[Out] `Integral((a*cos(e + f*x))**m/sqrt(b*csc(e + f*x)), x)`

$$3.293 \quad \int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=78

$$\frac{(a \cos(e+fx))^{m+1} {}_2F_1\left(-\frac{1}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e+fx)\right)}{abf(m+1) \sqrt[4]{\sin^2(e+fx)} \sqrt{b \csc(e+fx)}}$$

[Out] $-(a \cos(fx+e))^{(1+m)} \text{hypergeom}\left(\left[-\frac{1}{4}, \frac{1}{2}+\frac{1}{2}m\right], \left[\frac{3}{2}+\frac{1}{2}m\right], \cos(fx+e)^2\right) / a/b/f/(1+m)/(\sin(fx+e)^2)^{(1/4)}/(b \csc(fx+e))^{(1/2)}$

Rubi [A] time = 0.11, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2586, 2576}

$$\frac{(a \cos(e+fx))^{m+1} {}_2F_1\left(-\frac{1}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e+fx)\right)}{abf(m+1) \sqrt[4]{\sin^2(e+fx)} \sqrt{b \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[e + f*x])^m/(b*Csc[e + f*x])^(3/2),x]

[Out] $-\left(\left(\left(a \cos[e + f x]\right)^{(1+m)} \text{Hypergeometric2F1}\left[-\frac{1}{4}, \frac{(1+m)}{2}, \frac{(3+m)}{2}, \cos[e + f x]^2\right]\right) / \left(a b f (1+m) \sqrt{b \csc[e + f x]} \left(\sin[e + f x]^2\right)^{(1/4)}\right)\right)$

Rule 2576

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> -Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Cos[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2])/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2586

Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[(1*(b*Cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1))/b^2, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{3/2}} dx &= \frac{\int (a \cos(e+fx))^m (b \sin(e+fx))^{3/2} dx}{b^2 \sqrt{b \csc(e+fx)} \sqrt{b \sin(e+fx)}} \\ &= -\frac{(a \cos(e+fx))^{1+m} {}_2F_1\left(-\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \cos^2(e+fx)\right)}{abf(1+m) \sqrt{b \csc(e+fx)} \sqrt[4]{\sin^2(e+fx)}} \end{aligned}$$

Mathematica [A] time = 6.45, size = 116, normalized size = 1.49

$$\frac{2a \cos(2(e+fx)) \left(-\cot^2(e+fx)\right)^{\frac{1-m}{2}} (a \cos(e+fx))^{m-1} {}_2F_1\left(\frac{1}{4}(-2m-3), \frac{1-m}{2}; \frac{1}{4}(1-2m); \csc^2(e+fx)\right)}{bf(2m+3) \left(\csc^2(e+fx) - 2\right) \sqrt{b \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[e + f*x])^m/(b*csc[e + f*x])^(3/2),x]

[Out] (2*a*(a*cos[e + f*x])^(-1 + m)*Cos[2*(e + f*x)]*(-Cot[e + f*x]^2)^((1 - m)/2)*Hypergeometric2F1[(-3 - 2*m)/4, (1 - m)/2, (1 - 2*m)/4, Csc[e + f*x]^2])/(b*f*(3 + 2*m)*Sqrt[b*csc[e + f*x]]*(-2 + Csc[e + f*x]^2))

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \csc(fx + e)} (a \cos(fx + e))^m}{b^2 \csc(fx + e)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m/(b^2*csc(f*x + e)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m/(b*csc(f*x + e))^(3/2), x)

maple [F] time = 0.14, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m/(b*csc(f*x+e))^(3/2),x)

[Out] int((a*cos(f*x+e))^m/(b*csc(f*x+e))^(3/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m/(b*csc(f*x + e))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(a \cos(e + fx))^m}{\left(\frac{b}{\sin(e+fx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(3/2), x)`

[Out] `int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*cos(f*x+e))**m/(b*csc(f*x+e))**(3/2), x)`

[Out] `Integral((a*cos(e + f*x))**m/(b*csc(e + f*x))**(3/2), x)`

$$3.294 \quad \int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{5/2}} dx$$

Optimal. Leaf size=78

$$\frac{\sqrt[4]{\sin^2(e+fx)} \sqrt{b \csc(e+fx)} (a \cos(e+fx))^{m+1} {}_2F_1\left(-\frac{3}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e+fx)\right)}{ab^3 f(m+1)}$$

[Out] $-(a \cos(f*x+e))^{(1+m)} \text{hypergeom}\left(\left[-\frac{3}{4}, \frac{1}{2}+\frac{1}{2}*m\right], \left[\frac{3}{2}+\frac{1}{2}*m\right], \cos(f*x+e)^2\right) * (\sin(f*x+e)^2)^{(1/4)} * (b * \csc(f*x+e))^{(1/2)} / a / b^3 / f / (1+m)$

Rubi [A] time = 0.11, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2586, 2576}

$$\frac{(a \cos(e+fx))^{m+1} {}_2F_1\left(-\frac{3}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \cos^2(e+fx)\right)}{abf(m+1) \sin^2(e+fx)^{3/4} (b \csc(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(a*Cos[e + f*x])^m/(b*Csc[e + f*x])^(5/2),x]

[Out] $-\left(\frac{(a \cos[e + f*x])^{(1+m)} \text{Hypergeometric2F1}\left[-\frac{3}{4}, \frac{(1+m)}{2}, \frac{(3+m)}{2}, \cos[e + f*x]^2\right]}{(a*b*f*(1+m)*(b*Csc[e + f*x])^{(3/2)}*(\sin[e + f*x]^2)^{(3/4)})}\right)$

Rule 2576

Int[(cos[(e_) + (f_)*(x_)])*(a_)^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> -Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Cos[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2])/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2586

Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[(1*(b*Cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1))/b^2, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{5/2}} dx &= \frac{\int (a \cos(e+fx))^m (b \sin(e+fx))^{5/2} dx}{b^2 (b \csc(e+fx))^{3/2} (b \sin(e+fx))^{3/2}} \\ &= \frac{(a \cos(e+fx))^{1+m} {}_2F_1\left(-\frac{3}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \cos^2(e+fx)\right)}{abf(1+m)(b \csc(e+fx))^{3/2} \sin^2(e+fx)^{3/4}} \end{aligned}$$

Mathematica [A] time = 1.06, size = 125, normalized size = 1.60

$$\frac{2(2 \cos(2(e+fx)) + 1) \tan(e+fx) \left(-\cot^2(e+fx)\right)^{\frac{1-m}{2}} (a \cos(e+fx))^m {}_2F_1\left(\frac{1}{4}(-2m-5), \frac{1-m}{2}; \frac{1}{4}(-2m-1); \cos^2(e+fx)\right)}{b^2 f(2m+5) (3 \csc^2(e+fx) - 4) \sqrt{b \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*cos[e + f*x])^m/(b*csc[e + f*x])^(5/2),x]

[Out] (2*(a*cos[e + f*x])^m*(1 + 2*cos[2*(e + f*x)])*(-cot[e + f*x]^2)^((1 - m)/2)*Hypergeometric2F1[(-5 - 2*m)/4, (1 - m)/2, (-1 - 2*m)/4, Csc[e + f*x]^2]*Tan[e + f*x])/(b^2*f*(5 + 2*m)*sqrt[b*csc[e + f*x]]*(-4 + 3*Csc[e + f*x]^2))

fricas [F] time = 0.97, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \csc(fx + e)} (a \cos(fx + e))^m}{b^3 \csc(fx + e)^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m/(b^3*csc(f*x + e)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m/(b*csc(f*x + e))^(5/2), x)

maple [F] time = 0.13, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x)

[Out] int((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m/(b*csc(f*x + e))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(a \cos(e + fx))^m}{\left(\frac{b}{\sin(e+fx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(5/2),x)
```

```
[Out] int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(5/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*cos(f*x+e))**m/(b*csc(f*x+e))**(5/2),x)
```

```
[Out] Timed out
```


Chapter 4

Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.0.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] :=
  If[ExpnType[result]<=ExpnType[optimal],
    If[FreeQ[result,Complex] || Not[FreeQ[optimal,Complex]],
      If[LeafCount[result]<=2*LeafCount[optimal],
        "A",
        "B"],
      "C"],
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      "C",
      "F"]]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
```

```

(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]], 2]],
      Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3, ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
    If[HypergeometricFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
    If[AppellFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
    If[Head[expn]===RootSum,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
    If[Head[expn]===Integrate || Head[expn]===Int,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
    9]]]]]]]]]]

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```
AppellFunctionQ[func_] :=
  MemberQ[{AppellF1},func]
```

4.0.2 Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems

GradeAntiderivative := proc(result,optimal)
local leaf_count_result, leaf_count_optimal,ExpnType_result,ExpnType_optimal,
  debug:=false;

  leaf_count_result:=leafcount(result);
  #do NOT call ExpnType() if leaf size is too large. Recursion problem
  if leaf_count_result > 500000 then
    return "B";
  fi;

  leaf_count_optimal:=leafcount(optimal);

  ExpnType_result:=ExpnType(result);
  ExpnType_optimal:=ExpnType(optimal);

  if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",
  ExpnType_optimal);
  fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
  return "F";
end if;

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
fi;
```

```

if is_contains_complex(result) then
  if is_contains_complex(optimal) then
    if debug then
      print("both result and optimal complex");
    fi;
    #both result and optimal complex
    if leaf_count_result<=2*leaf_count_optimal then
      return "A";
    else
      return "B";
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C";
  end if
else # result do not contain complex
  # this assumes optimal do not as well
  if debug then
    print("result do not contain complex, this assumes optimal do
not as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B";
  end if
end if
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C";
end if

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function

```



```

# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+' or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

```

```

AppellFunctionQ := proc(func)
  member(func,[AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.0.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]

def is_hypergeometric_function(func):
  return func in [hyper]

def is_appell_function(func):
  return func in [appellf1]

```

```

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+`' or
    type(expn,'*`)
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,
    expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType
    ,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:
        m1 = max(map(expnType, list(expn.args)))
        return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    else:
        return 9

```

```

#main function
def grade_antiderivative(result,optimal):

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        return "F"

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
    else:
        return "C"

```

4.0.4 SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#           Albert Rich to use with Sagemath. This is used to
#           grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#           'arctan2','floor','abs','log_integral'

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True

```

```

        else:
            return False
    else:
        return False

def is_elementary_function(func):
    debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    debug=False
    if debug: print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','
sinh_integral'
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M',
hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in
sagemath

def is_atom(expn):

    debug=False
    if debug: print ("Enter is_atom")

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type-in-maple/
    try:
        if expn.parent() is SR:

```

```

        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens")
:
        return expn in expn.parent().base_ring() or expn in expn.parent().
gens()
    return False

except AttributeError as error:
    return False

def expnType(expn):

    if debug:
        print(">>>>Enter expnType, expn=", expn)
        print(">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],
Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer
)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],
Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],
Rational)
                return 1
            else:
                return max(2,expnType(expn.operands()[0])) #max(2,expnType(
expn.args[0]))
        else:
            return max(3,expnType(expn.operands()[0]),expnType(expn.operands(
[1])) #max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))
    elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #
isinstance(expn,Add) or isinstance(expn,Mul)
        m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
        m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn)))
    elif is_elementary_function(expn.operator()): #is_elementary_function(expn
.func)
        return max(3,expnType(expn.operands()[0]))
    elif is_special_function(expn.operator()): #is_special_function(expn.func)
        m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
        return max(4,m1) #max(4,m1)
    elif is_hypergeometric_function(expn.operator()): #
is_hypergeometric_function(expn.func)
        m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
        return max(5,m1) #max(5,m1)
    elif is_appell_function(expn.operator()):

```

```

        m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(
expn.args)))
        return max(6,m1)      #max(6,m1)
    elif str(expn).find("Integral") != -1: #this will never happen, since it
        #is checked before calling the grading function that is passed.
        #but kept it here.
        m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(
expn.args)))
        return max(8,m1)      #max(5,apply(max,map(ExpnType,[op(expn)])))
    else:
        return 9

#main function
def grade_antiderivative(result,optimal):

    if debug: print ("Enter grade_antiderivative for sagemath")

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    if debug: print ("leaf_count_result=", leaf_count_result, "
leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",
expnType_optimal)

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
    else:
        return "C"

```